

MATH 2352 Solution Sheet 03

BY XIAOYU WEI

Created on March 6, 2015

[Problems] 2.7: 16; 3.1: 4, 11; 3.3: 14, 19; 3.4: 6, 11; 3.2: 9, 10, 14, 25; 3.5: 6, 20;

2.7 - 16. Consider the initial value problem

$$y' = t^2 + y^2, \quad y(0) = 1.$$

Use Euler's method with $h = 0.1, 0.05, 0.025$, and 0.01 to explore the solution of this problem

for $0 \leq t \leq 1$. What is your best estimate of the value of the solution at $t = 0.8$? At $t = 1$? Are your results consistent with the direction field in Problem 9?

Solution.

The following Matlab code solves the problem:

```
matlab>>hh = [ 0.1, 0.05, 0.025, 0.01 ];
          f = @(t,y)(t.^2+y.^2);
          N1 = floor(0.8./hh);
          N2 = floor(1./hh);
          v = zeros(4, 2);
          for i = 1:4
              y = 1; h = hh(i);
              for j = 1:N2(i)
                  y = y + f(j*h-h,y)*h;
                  if j == N1(i)
                      v(i,1) = y;
                  end;
              end;
              v(i,2) = y;
          end;
matlab>>disp(v);
          3.5078    7.1895
          4.2013   12.3209
          4.8004   23.9260
          5.3428   90.7555
```

3.1 - 4. Find the general solution of the given differential equation:

$$3y'' - 4y' + y = 0.$$

Solution. Since the equation can be written as

$$\begin{aligned} 3y'' - 3y' - y' + y &= 0 \\ 3(y'' - y') - (y' - y) &= 0 \\ (y' - y)' &= (y' - y)/3, \end{aligned}$$

we have

$$y' - y = C_0 e^{x/3}.$$

One solution is

$$y(x) = -\frac{3}{2}C_0 e^{x/3}.$$

Let $C_1 = -\frac{3}{2}C_0$, and notice that the solution can differ by $C_2 e^x$,

$$y(x) = C_1 e^{x/3} + C_2 e^x.$$

3.1 - 11. Find the solution of the given initial value problem. Sketch the graph of the solution and describe its behavior as t increases.

$$\begin{aligned} 12y'' - 7y' + y &= 0, \\ y(0) &= 4, \\ y'(0) &= 0. \end{aligned}$$

Solution. The equation can be written as

$$\begin{aligned} 12y'' - 3y' - 4y' + y &= 0 \\ (4y' - y)' &= (4y' - y)/3. \end{aligned}$$

From this equation $4y' - y$ can be solved.

On the other hand,

$$\begin{aligned} 12y'' - 4y' - 3y' + y &= 0 \\ (3y' - y)' &= (3y' - y)/4. \end{aligned}$$

So the general solution is


$$y(x) = C_1 e^{x/3} + C_2 e^{x/4}.$$

The solution to the I.V.P. is

$$y(x) = 16e^{x/4} - 12e^{x/3}.$$

The solution is shown in below:

GNUplot] `plot [0:10] 16*exp(x/4)-12*exp(x/3)`



3.3 - 14. Find the general solution of given differential equation:

$$9y'' + 3y' - 2y = 0.$$

Solution. Assume λ is a real constant, then

$$9y'' + \lambda y + (3 - \lambda)y - 2y = 0.$$

Let

$$\frac{9}{3 - \lambda} = \frac{\lambda}{-2}.$$

This equation has two roots $\lambda_1 = -3$, $\lambda_2 = 6$.

Then we can solve the two 1-st order equations and get the general solution

$$y(x) = C_1 e^{-2/3} + C_2 e^{1/3}.$$


3.3 - 19. Find the solution of the given initial value problem. Sketch the graph of the solution and describe its behavior as t increases.

$$\begin{aligned} y'' - 2y' + 5y &= 0, \\ y\left(\frac{\pi}{2}\right) &= 0, \\ y'\left(\frac{\pi}{2}\right) &= 4. \end{aligned}$$

Solution. The solution is

$$y(x) = -2e^{x - \frac{\pi}{2}} \sin(2x).$$

GNUplot] `plot [pi/2:11.5] -2*exp(x-(pi/2))*sin(2*x)`



3.4 - 6. Find the general solution of given differential equation:

$$y'' - 10y' + 25y = 0.$$

Solution. This is the degenerate case, meaning only one reduced equation can be written

$$(y' - 5y)' = 5(y' - 5y).$$

And the solution is $y' - 5y = C_0 e^{5x}$.

Solving this yields

$$y(x) = C_1 e^{5x} + C_2 x e^{5x}.$$

3.4 - 11. Find the solution of the given initial value problem. Sketch the graph of the solution and describe its behavior as t increases.


$$\begin{aligned} 9y'' - 12y' + 4y &= 0, \\ y(0) &= 2, \\ y'(0) &= -2. \end{aligned}$$

Solution. The solution is

$$y(x) = -\frac{2}{3}e^{2x/3}(5x-3).$$

Plot the solution:

GNUplot `plot [0:2] -2.0/3.0*exp(2*x/3)*(5*x-3)`



3.2 - 9. Determine the longest interval in which the given initial value problem is certain to have a unique twice-differentiable solution. Do not attempt to find the solution.

$$\begin{aligned} t(t-4)y'' + 3ty' + 5y &= 2, \\ y(3) &= 0, \\ y'(3) &= -1. \end{aligned}$$

Solution. $(3, 4)$.

3.2 - 10. Determine the longest interval in which the given initial value problem is certain to have a unique twice-differentiable solution. Do not attempt to find the solution.

$$\begin{aligned} y'' + (\sin t)y' + 3(\ln |t|)y &= 0, \\ y(1) &= 3, \\ y'(1) &= 1. \end{aligned}$$

Solution. $(1, +\infty)$.

3.2 - 14. Verify that $y_1(t) = 1$ and $y_2(t) = t^{1/2}$ are solution of the differential equation $y y'' + (y')^2 = 0$ for $t > 0$. Then show that $y = c_1 + c_2 t^{1/2}$ is not, in general, a solution of this equation. Explain why this result does not contradict Theorem 3.2.2.

Solution. Nonlinearity.

3.2 - 25. Verify that the functions y_1 and y_2 are solutions of the given differential equation. Do they constitute a fundamental set of solutions?

$$\begin{aligned} y'' - 4y' + 4y &= 0; \\ y_1(t) &= e^{2t}, \\ y_2(t) &= t e^{2t}. \end{aligned}$$

Solution. Yes.

3.5 - 6. Find the general solution of given differential equation:

$$y'' + 2y' = 5 + 4 \sin 2t.$$

Solution. Let $p = y'$,

$$p' = -2p + 5 + 4 \sin 2t.$$

Then $p = C_1 e^{-2t} + \sin(2t) - \cos(2t) + \frac{5}{2}$.

And $y = p + C_2$.

3.5 - 20. Find the solution of the given initial value problem:

$$\begin{aligned} y'' + 2y' + 5y &= 4e^{-t} \cos 2t, \\ y(0) &= 0, \\ y'(0) &= 0. \end{aligned}$$

Solution. $y = e^{-t} t \sin(2t)$.

Remark 1. The general solution for the homogenous problem is $y(t) = C e^{-t} \sin(2t)$. (Only real solution is considered here).

Remark 2. By assuming the solution is $y(t) = C(t) e^{-t} \sin(2t)$, one can derive the initial value problem for C as

$$\begin{aligned} (\tan 2t) C'' + 4C' - 4 &= 0, \\ C(0) &= 0. \end{aligned}$$

To solve this, let $z(t) = C'$, and solve the separable equation

$$(\tan 2t) z' + 4z - 4 = 0$$

$$(\tan 2t) \frac{dz}{dt} = 4 - 4z$$

$$\frac{dz}{4 - 4z} = \frac{dt}{\tan 2t},$$

which gives $z(t) = C_1 \csc^2(2x) + 1$. Therefore,

$$C(t) = C_1 \cot 2t + t + C_2.$$

Since $C(0)$ is well-defined and equals to zero, $C_1 = C_2 = 0$, we have

$$C(t) = t.$$