


A classical oil painting of Isaac Newton, showing him from the chest up. He has long, wavy, light-colored hair and is wearing a dark, heavy robe over a white shirt. He is looking slightly to the right with a serious expression. His right hand is resting on a surface in the lower left corner.

Calculus Through Physics

Xiao Yaozi



Contents

Contents	iii
Preface I: Calculus, the language of the universe	v
Preface II: How to read this book	vii
I Precalculus	3
1 Functions	5
1.1 Function basics	5
1.2 Function composition and inverses	10
1.3 Injection, surjection and bijection	14
1.4 An aside: A more rigorous definition of a function	19
2 Linear Algebra	21
2.1 Vector Basics	21
2.2 Scaling and dot product	26
2.3 Matrices	30
2.4 Determinants, inverses and Cramer’s rule	34
2.5 Cross products and triple products	39
2.6 An aside: Bra-ket and Hilbert space	46
2.7 An aside: Eigenfunctions and eigenvalues	46
II Calculus I	47
3 Derivatives	49
3.1 Differentiation basics	49

Preface I: Calculus, the language of the universe

As you are seated right here, on your seat, reading this book, you must realise that you are made of the same fundamental particles as everything in the universe. These fundamental particles are governed by the Schrodinger equation, which is written with notation from calculus ¹:

¹ We will assume that you are not fast enough for relativistic effects to apply.

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \Psi + V\Psi \quad (0.0.1)$$

Do not worry if you do not understand this equation. In fact, this book will teach some quantum mechanics at the end because it just shows the absolute manifestation of calculus in physics. Even if you know absolutely nothing about quantum mechanics, I have a much more subtle, yet interesting question for you: why is the area of a circle of radius r is πr^2 . You might turn to the rearrangement proof, or the “pizza” explanation as I like to call it. Take a circle, and divide it into congruent segments like cutting pizza slices. Then, take the slices and rearrange them in such way such that they form a sort of parallelogram.




Figure 0.1 Figure of the rearrangement proof.

Something that you notice in Figure 0.1 is that if you make each slice smaller, the parallelogram will become more like a rectangle. This, right here, is a very subtle yet profound fact that makes many problems in the real world much easier. As the famous saying goes, “calculus turns one big problem into infinitely smaller ones.” This little trick is the reason for developing the calculus method. Everything in calculus stems from the proposition that dividing something into infinitesimal parts reduces the complexity of a problem. So far, this has held up well. No matter if you are looking at classical mechanics or quantum mechanics, all the equations are written with calculus. While this little fact may seem very deep or even philosophical, and we will explore on the paradoxes of infinitesimals, I believe that one cannot meaningfully know what calculus means before knowing what it does. The purpose of this book is to teach you how to do calculus through the lens of physics. There are many schools of thought on the pedagogy of calculus and there are great texts on calculus such as Stewart, J. *Calculus: Early Transcendentals, 8th Ed.* and Strang, G. *Calculus*. Furthermore, if you already know the basics of calculus, I would also recommend Arfken, G. *Mathematical Methods for Physicists* and Boas, M.L. *Mathematical Methods in the Physical Sciences* is also a classic. While these books are fantastic, they either already assume a decent knowledge of calculus or does not put its focus on physics. This book seeks to fill this special niche.

I wish every reader good luck on their journey on learning calculus. It is not trivial and rereading confusing parts of the book is highly recommended. More than just knowing how to apply calculus however, I also encourage the reader to appreciate the beauty of calculus and its unreasonable effectiveness in describing our physical world. To end off, thanks for choosing this book. Happy reading.

XIAO YAOZI



Preface II: How to read this book

This book is structured into four parts. The first part deals with precalculus or the content that is needed for the later calculus chapters. Be warned, however, that the first part is rather short and sweeps over the content very quickly. The second part corresponds to the standard curriculum of Calculus I and is the beginning of the actual calculus portion. Similarly, the third and fourth part corresponds to Calculus II and multivariable calculus respectively. As with any textbook, problems are included at the end of each chapter for the reader to practice and familiarise with the topic. To aid the reader, we have included the following notation for denoting the type of problem:

1. * means that the problem is an *essential* problem, usually something that is very good to know or a theorem not derived in the text.
2. ** means that the problem is more *peripheral* and the solution might involve using a clever identity or trick.
3. *** means that the problem is very *difficult* and might take the reader very long to solve.
4. Nothing means that the problem is not very important and can be skipped.

Furthermore, to prevent confusion on sidenotes² and indices, sidenote markers are **bolded** while the latter is not. The notation for the mathematics is mostly standard, but a few notations are defined in the text. If you are confused about any notation, please refer to the table on the next page.

Finally, if you have any questions or errata, please contact me and submit an issue on GitHub. The GitHub link is <https://github.com/xyzdev-code/Calculus-through-Physics>.

² *This is a sidenote. It is italicized to prevent confusion with the main text.*

Description	Meaning	Example	Notes
Letters	Scalars, constants, matrices	a, b, c, X, Y, Z	Except for constants such as e, π, c, g . Functions may also be written as single letters, however the context would be clear. I is the identity matrix.
Bolded, upright letters	Vectors	$\mathbf{v}, \hat{\mathbf{i}}, \hat{\mathbf{j}}, \hat{\mathbf{k}}$	$\mathbf{0}$ is the zero vector.
Square brackets	Matrices	$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$	Nil
Vertical bars	Determinant	$\begin{vmatrix} 1 & 2 \\ 3 & 4 \end{vmatrix} = -2$	Nil
Functions with dots over them	Derivatives with respect to some variable (usually time)	$\dot{x}, \ddot{x}$	Nil

Table 0.1 Table of notation.

Part I

Precalculus

Functions


1.1 Function basics

Before we start on the calculus topics, this part of the book will acquaint you with the mathematics you would need to understand in order to read this book. If you are familiar with these topics, you may skip to the Part II on Calculus I. For the rest of you, the pace of this part will be very fast, it is recommended that you read up on the concepts you do not understand very well. We will start with the idea of a *function*. By now, I am sure that many of you may have seen equations like these:

$$x^2 - 5x + 6 = 0 \quad (1.1.1)$$

$$\sin(x) = 0.5, 0 \leq x \leq \frac{\pi}{2} \quad (1.1.2)$$

$$y = mx + c \quad (1.1.3)$$



Out of the three equations, notice that Equation (1.1.3) is special. There is not some fixed x values that you must plug in. Instead, the y value varies with the x value. Given a x value, the equation “produces” a y value. This brings us to the idea of a function. A function is simply something that takes in a value, and “produces” another value based on the input. You may have come across functions before, such as the trigonometric functions like $\sin$ and $\cos$. The value of $\sin(x)$ varies with different values of x . Of course, it is time for me to give you the definition of a function.

Figure 1.1 Graph of $y = x$.

Definition 1.1 Function

A **function** f is a mathematical object that assigns its inputs, or *domain*, to its outputs, or *range*. The notation $f : A \rightarrow B$ means that the function has a domain A and range B ³. The notation $f(a)$ where $a \in A$ denotes the element of B to which a is assigned to.



³ Note that A and B are sets.

Example 1.1 Evaluating functions

Here are some examples of functions:

$$f(x) = x + 1 \quad (1.1.4)$$

$$g(x) = \sin(x) \quad (1.1.5)$$

$$h(x) = x^2 - 5x + 6 \quad (1.1.6)$$

To evaluate the output of a function, you simply substitute the input to the expression that defines the function. Notice that for Equation (1.1.4), it is simply the line $y = x + 1$. Each value of $f(x)$ for some $x \in \mathbb{R}$ is just y . Take $x = 1$ for example, $f(1) = 1 + 1 = 2$ and for the line $y = x + 1$, notice the y value here is also 2. The function assigns each x value to $x + 1$. Hence, in this case, $f : \mathbb{R} \rightarrow \mathbb{R}$. For Equation (1.1.5), the function takes any real x value and maps it to $\sin(x)$. As a result, its domain would be $\mathbb{R}$ and its range would be $[-1, 1]$. Again, it is just like the graph $y = \sin(x)$. Lastly, the quadratic in Equation (1.1.6) is another real function (i.e. it has a real domain and range). Think about what happens when $h(x) = 0$, what value of x must you plug in to make $h(x) = 0$. Well, isn't it just the roots of the quadratic. Hence, the roots of a function are just the values x such that $h(x) = 0$. In this case, it would be 2 and 3. Thus, $h(2) = h(3) = 0$.

△

Actually, functions can be more than the usual $f(x) = \text{something}$. Let's say you want to have a function that outputs "1" if the input is rational and "0" if it is not, you can define it as such: ⁴

⁴ This is actually the famous Dirichlet function that shatters certain intuitions on functions and will be discussed in later chapters.

$$f(x) = \begin{cases} 1, & \text{if } x \in \mathbb{Q} \\ 0, & \text{if } x \notin \mathbb{Q} \end{cases} \quad (1.1.7)$$

Here, $f(\frac{1}{2})$ would be 1 but $f(\sqrt{2})$ would be 0. (You would probably get a migraine from graphing it.) This is an example of having conditionals in a function. Lastly, functions may take more than one input and produce more than one output, examples include:

$$f(x, y) = x^2 + y^2 \quad (1.1.8)$$

$$g(t) = (\cos(t), \sin(t)) \quad (1.1.9)$$

⁵ As you will see in Part IV of the book, this is the parametric equation of a circle.

Equation (1.1.8)⁵ is an example of a *multivariate* function, in other words, it takes in multiple inputs. If $x = 2$ and $y = 2$, then $f(2, 2) = 8$. On the other hand, Equation (1.1.9) does not have a single output. Instead, it has a tuple ⁶ as its output. Evaluating it at $t = \pi$, notice that the output is $(-1, 0)$. What is interesting is that if you trace out its outputs on a plane for all $t \in \mathbb{R}$, you will get Figure 1.2.

⁶ A tuple is basically a set where order matters. You can treat it as if it is a coordinate.

Application: Simple pendulum

In physics, we often deal with functions which enables us to describe many things whose behaviour is dependent on certain inputs. Take a pendulum for example. We can describe the angle of the pendulum to the normal line using the following formula⁷:

⁷ While not completely accurate, this formula is good enough for an example.

$$\theta(t) = \theta_0 \cos\left(\sqrt{\frac{g}{l}}t\right) \quad (1.1.10)$$




Figure 1.2 Plot of Equation (1.1.9) for all $t \in \mathbb{R}$

Here, $\theta(t)$ is the angle of the pendulum at time t , θ_0 is the initial angle of the pendulum, T is the period of the pendulum, l is the length of the rod. (g is the gravitational field strength.) While this function may look slightly complicated, we can deduce a lot about it without computing much. Firstly, at $t = 0$, $\theta(0) = \theta_0$, which makes sense because at the start, the pendulum is at its initial angle.




Figure 1.3 Figure of pendulum.

The periodic nature of the pendulum is also expressed in the cosine part of the function. Moreover, since $\cos$ can only attain values between -1 and 1 , the pendulum is guaranteed to be oscillating between $-\theta_0$ and θ_0 . This satisfies the conservation of energy because if this was not guaranteed and no energy is introduced into the system, energy must have been created or destroyed.

Challenge 1.1 A real world pendulum

This being said, in the real world, is energy lost to the surroundings? How could energy be lost to the surroundings? How would this affect the motion of the pendulum?

?

Let us calculate the period of the pendulum.

Example 1.2 Period of a pendulum

We know the formula for the angle of the pendulum for any time t . The period of a function $\cos(ax)$ is simply $\frac{2\pi}{a}$, hence the period of the pendulum is $2\pi\sqrt{\frac{l}{g}}$. This results shows us that the period of the pendulum is **independent** of the initial angle. This property is called “isochronism” that Galileo discovered.

△

In the real world, we often add a *damping factor* $d(t)$ to $\theta(t)$. A damped pendulum has the function describing its angle as $\theta_d(t) = \theta(t)d(t)$. The damping factor is to reduce the amplitude of the pendulum as time passes, mainly because air resistance opposes the motion of the pendulum.

Challenge 1.2 A suitable damping factor

From the following damping factor $d(t)$, choose the most suitable one (you may want to use a graphing calculator to assist you):

$$e^t$$

$$e^{-t}$$

$$-t$$

$$a, a \in \mathbb{R}$$

?

Problem 1.1

Given that $x = 2$ and $y = \pi$, evaluate $f(x)$ for:

a) $f(x) = x^3 + 1$

b) $f(x) = 2^x$

c) $f(x) = \begin{cases} x^4 + 1 & \text{if } x < 2 \\ 3^x & \text{otherwise} \end{cases}$

d) $f(x, y) = \frac{\sin(y)}{x}$

Problem 1.2

Given the following functions, write down $f : A \rightarrow B$ where A is the domain and B is the range (do not consider complex inputs and outputs, we shall be ignorant to it for the time being to simplify things):

a) $f(x) = x^8$

⁸ This function has a special name, the **identity** function. Notice how its input and output are the same.

- b) $f(x) = \frac{1}{x}$
c) $f(x) = \sqrt{x}$
-

Problem 1.3

Find the root(s) of the following functions:

- a) $f(x) = x^2$
b) $f(x) = 2^x - 1$
c) $f(x) = \sin(x)$
-

*** Problem 1.4**

A *recursive* function is one where the function uses itself in its definition. For example, the Fibonacci sequence can be defined like this for any Fibonacci number n :

$$f(n) = \begin{cases} f(n-1) + f(n-2) & n > 1 \\ 1 & n = 1 \\ 0 & n = 0 \end{cases}$$

- a) Compute $f(3)$ explicitly using the definition of f .
b) Define the factorial of a positive integer in a similar way using recursion, define it as $g(a)$.
c) Sketch the graph of $g(a)$.
-

*** Problem 1.5**

(*Polynomial remainder theorem*) Let $f(x)$ be a polynomial. If $f(a) = 0$, prove that $x - a$ is a factor of $f(x)$.

**** Problem 1.6**

Find $f : \mathbb{R} \setminus \{0, 1\} \rightarrow \mathbb{R}$ such that $f(x) + f\left(\frac{x-1}{x}\right) = 1 + x$ for any $x \in \mathbb{R} \setminus \{0, 1\}$.

1.2 Function composition and inverses

In this chapter, we will look at composition of functions and inverse functions. You can use the normal arithmetic operations on functions like how you would to numbers:

Example 1.3 Arithmetic operations on functions

If $x = 2$ and $f(x) = x^2 + 4$:

$$f(x) + f(x) = 2^2 + 4 + 2^2 + 4 = 16$$

$$\frac{f(x)}{2} = \frac{2^2 + 4}{2} = 4$$

△

What is more interesting, perhaps, is taking the *output* of a function and use it as the *input* to another function. You may also think of it as “piping” one function into another. This is called *function composition*.




Figure 1.4 Visual representation of function composition. ⁹

⁹ Note that the rectangles are functions and circles are values.

Another notation for composing functions together (which we will not really use except for this chapter) is as such:

$$f(g(x)) = (f \circ g)(x)$$

When you compose the same function n -times, the notation for it is¹⁰:

$$f^n(x) = \underbrace{f(f(\dots f(x)))}_{n\text{-times}}$$

Example 1.4 Composing functions

Here are some examples of composing functions together: If $x = 3$ and $f(x) = \pi x$, $g(x) = \sin(x) + 1$:

$$g(f(x)) = \sin(\pi \times 3) + 1 = 1 \quad (1.2.1)$$

$$f^3(x) = \pi^3 x \quad (1.2.2)$$

There is nothing much here, really, just evaluate the function from the innermost function to the outermost.

△

Theorem 1.1 Composition of functions is associative

Given functions $f : X \rightarrow Y$, $g : Y \rightarrow Z$ and $h : Z \rightarrow W$, prove that the composition of the three functions is *associative*, in other words:

$$h \circ (g \circ f(x)) = (h \circ g) \circ f(x)$$

QED

¹⁰ However, this does not apply to trigonometric functions. $\sin^2(x)$ is $\sin(x) \times \sin(x)$ instead of $\sin(\sin(x))$. Annoying mathematical notation strikes again. What is even better, is that $f^{-1}(x)$ denotes the inverse of f .

Proof 1.1 Composition of functions is associative

Let $x \in X$, we have:

$$h \circ (g \circ f(x)) = h(g \circ f(x)) = h(g(f(x))) = h \circ g(f(x)) = (h \circ g)f(x)$$

This proves the associativity of function composition.

QED

Remark 1.1

This is just a fun little exercise for the reader and ultimately is not that important to the entire book, just a good identity to keep in the back of your mind.



Next, we will delve into inverse functions. Let us start with one I believe you may have seen before, look at the following figure:




Figure 1.5 Figure of right triangle ABC.

If I ask you what is $\arcsin(B)$, you may say it is $\frac{b}{c}$. But, what is $\arcsin(\sin(B))$, well, that is just B , isn't it. Hence, the arcsin sort of "cancels" the sin. We call arcsin the *inverse function* or just inverse of sin.¹¹ Generally, given a function $f : X \rightarrow Y$, f^{-1} denotes the inverse function that satisfies:

¹¹ We will use arcsin as the inverse of sin instead of $\sin^{-1}$ as it is clearer.

$$f \circ f^{-1}(y) = y, y \in Y \text{ and } f^{-1} \circ f(x) = x, x \in X \quad (1.2.3)$$

Note 1.1 Cancelling functions and their "inverse"

One common misconception is that $f(f^{-1}(x)) = f^{-1}(f(x))$ for any function. While this is technically true, you may apply this fact wrongly. For example, take the following function:

$$f(x) = \sin(x), f^{-1} : \{y : -1 \leq y \leq 1\} \rightarrow \mathbb{R}$$

This looks deceptively simple, when asked to evaluate $f^{-1}(f(\pi))$, it is π (You may want to verify it yourself). But what is $f(f^{-1}(\pi))$? Well, according to our definition of f , $f^{-1}(\pi)$ is simply not defined. In actuality, to find $f^{-1}(\pi)$, we need complex numbers, which is not allowed in the definition. Hence, you must be extra careful when you are dealing with function inverses and pay attention to the domain and range.



To find the inverse function of a function f , we have to find the function that maps the outputs of the f back to its inputs. To solve algebraically, we set $f(x)$ for some arbitrary x to y , and make x the subject of the equation.

Example 1.5 Finding the inverse of a function

Given that $f(x) = \frac{x+7}{3}$:

$$\begin{aligned}f(x) &= y \\ \frac{x+7}{3} &= y \\ x+7 &= 3y \\ x &= 3y-7\end{aligned}$$

Hence, the inverse function is $f^{-1}(y) = 3y - 7$.

**Example 1.6 Common inverse functions**

Some common examples of inverse functions:

1. Trigonometry:

arcsin arccos arctan ...

2. Logarithms (Inverses of exponential functions):

ln log_b



Problem 1.7

Given $f(x) = \frac{x^2+3}{2}$ and $g(x) = \tan(x) + 1$, evaluate:

- a) $f(g(\frac{\pi}{4}))$
- b) $f^3(1)$
- c) $f(g(\arcsin(0)))$

Problem 1.8

Given that $h(x) = (x-7)^2$, $x \geq 7$, find the inverse of h . Hence, evaluate:

- a) $h^{-1}(1)$
- b) $h(h^{-1}(69))$ As Peter Lorre would have said it, “Do it ze kveek vay, Johnny”.

*** Problem 1.9**

The *dirac delta function* is defined as

$$\delta(x) = \begin{cases} \infty & x = 0 \\ 0 & \text{otherwise} \end{cases} \quad (1.2.4)$$

It also has an area of 1 “underneath” it.

- a) Show that $f(x)\delta(x - a) = f(a)\delta(x - a)$.
- b) Hence, simplify $(3x^2 - 2x - 1)\delta(x - 3)$.

*** Problem 1.10**

Given that the function $f(x)$ is defined as $f(x) > f(x - 1) + f(x - 2)$, and when $x < 3$, $f(x) = x$, show that $f(20) > 1000$. (*Adapted from the 2024 Chinese College Entrance Examination*)

1.3 Injection, surjection and bijection

You may have noticed in the previous exercise that question 2 is particularly interesting. There is a restriction on the domain of $h(x)$. Why is that? This is what this chapter is all about. However, before we explain that, I will go over some definitions of properties of functions. Firstly, we have *injection*.

Definition 1.2 Injective functions

An **injective** function, f , or a *one-to-one* function, is a function such that every distinct output of the function corresponds to only *one* distinct input, that is ¹²:

$$f(x_1) = f(x_2) \text{ implies } x_1 = x_2$$

¹² By *contraposition*, $x_1 \neq x_2$ implies $f(x_1) \neq f(x_2)$. The *contraposition* of a statement “ p then q ” is “not q then p ”.

Of course, this explanation might not have been very clear. Here is a visualization:




Figure 1.6 Illustration of an injective function with domain X and range Y .

As you can see in Figure 1.6, every output in Y of the function is associated with only one input in X , hence the function is *injective*. (Kind of like how marriages *should* be.) To prove that a function f is injective, you can either show that if $x_1 \neq x_2$, then $f(x_1) \neq f(x_2)$, or if $f(x_1) = f(x_2)$, then $x_1 = x_2$.




Figure 1.7 Illustration of a non-injective function with domain X and range Y .

Here, the function is *not* injective. As you can see in Figure 1.7, the output y is associated with more than one input in X . With this in mind, it is time to talk about *surjection*.

Definition 1.3 Surjective function

A **surjective** function f is a function such that for every output y , there exists at least one input x such that $f(x) = y$.

From Figure 1.6 and Figure 1.7, both functions are surjective as every output is associated with at least one input. Do take note that it does not matter how many inputs associate to one output, as long as every output is associated with at least one input, the function is surjective. To prove that a function is surjective, you must show that there exists an element in the domain of the function for any arbitrary output in the range.




Figure 1.8 Illustration of non-surjective function with domain X and range Y .

From Figure 1.8, we can see that y has no input associated with it. As a result, the function here is not surjective. Lastly, *bijective* functions can't be any easier.

Definition 1.4 Bijective function

A **bijective** function is a function that is both injective and surjective.

However, bijective functions have a superpower. A function is only bijective *if and only if* it has an inverse.¹³

Theorem 1.2 Bijection and inverses

A function, $f : A \rightarrow B$, has an inverse if and only if it is bijective.

¹³ P iff Q , or “ P if and only if Q ”, means that “if P then Q ” and “if Q then P ”.

QED

Proof 1.2 Bijection and inverses

To begin such proofs where there is a “if and only if” clause, we will prove both cases separately. First, we claim that if the function has an inverse, then it is bijective. To prove this, we simply prove that f is injective and surjective. Then, we claim that if a function is bijective, then it has an inverse. With both proofs below, we show that f has an inverse if and only if it is a bijection.

QED**Lemma 1.1** Inverse function has bijections

If a function, $f : A \rightarrow B$, has an inverse, then it is bijective.

QED**Proof 1.3** Inverse functions has bijections

Suppose f^{-1} is the inverse of f . Notice that $f^{-1}(f(x))$ is injective as it is simply $f(x) = x$. Suppose $a, a' \in A$ and $f(a) = f(a')$, then $f^{-1}(f(a)) = f^{-1}(f(a'))$. Notice that by injectivity of $f^{-1}(f(x))$, $a = a'$. Hence, f is injective. Then, suppose $b \in B$, we wish to show that the function is surjective. Also, $f(f^{-1}(x))$ is surjective(it is the identity function). Let $f^{-1}(b) = x$ and $f(f^{-1}(b)) = c$ for some $c \in B$. By a simple substitution, $f(f^{-1}(b)) = f(x) = c$, thereby showing there exists a $x \in A$ for every $c \in B$. The two cases proves that f is bijective if it has an inverse.

QED**Lemma 1.2** Bijective function has inverses

If a function, $f : A \rightarrow B$, is a bijection, then it has an inverse.

QED**Proof 1.4** Bijective function has inverses

Let us define f^{-1} as $f^{-1} : B \rightarrow A$. Let $b \in B$. Since f is surjective, there exists $a \in A$ such that $f(a) = b$. Let $f^{-1}(b) = a$. Since f is injective, f^{-1} is well-defined, or a function that outputs the same value given the same input. Next, we will show that f^{-1} is indeed the inverse of f by showing the composition of the two is an identity function. Again, let $a \in A$ and $b = f(a)$ which implies, by definition, $f^{-1}(b) = a$. Composing the two functions, $f(f^{-1}(b)) = f(a) = b$, which shows that f^{-1} is indeed the inverse function. The other case of $f^{-1} \circ f$ is similar. Hence, if f is a bijection, then it has an inverse.

QED**Remark 1.2**

This serves as a very good example showing how the reader should prove the injectivity, surjectivity, and bijectivity of functions. Also, this explains why in the previous exercise, the function $h(x) = (x - 7)^2$ is bounded, only taking x values larger than 7. The function is not bijective in $\mathbb{R}$, as evident in the graph as it is not injective. Hence, the bounds are needed for h to have an inverse.

△

Problem 1.11

Determine if the following functions are injective or surjective or bijective in its domain:

- a) $f(x) = \sqrt{x}, f: \mathbb{R}^+ \rightarrow \mathbb{R}^+$
- b) $f(x) = \frac{1}{x}, x > 0$
- c) $f(x) = (x - 7)^2, x \geq 5$




Figure 1.9 Plot of $(x - 7)^2$.

*** Problem 1.12**

Prove that:

- a) The composition of two injective functions is *injective*.
- b) The inverse of a function is *injective*.
- c) The inverse of a function is *surjective*.

*** Problem 1.13**

Given a quadratic function $q(x) = ax^2 + bx + c$, and $a \neq 0$, show that q cannot be bijective.

Further problems*** Problem 1.14**

We shall define an area function, $A(f, x_1, x_2)$ which gives us the area under a given function $f(x)$ bound by $x_1 \leq x \leq x_2$.

- a) Using an example, show that $A(f, x_1, x_2)$ cannot be bijective.
- b) Also using an example, disprove that $A(fg, x_1, x_2) = A(f, x_1, x_2)A(g, x_1, x_2)$.
- c) From the examples, verify geometrically that $A(f + g, x_1, x_2) = A(f, x_1, x_2) + A(g, x_1, x_2)$:
 - a) $f(x) = x + 1$ and $g(x) = x + 2$
 - b) $f(x) = x$ and $g(x) = 0$
 - c) $f(x) = 1$ and $g(x) = 2$

- d) If δ is the dirac delta function found in Problem 1.9, show that $A((x^2 + x + 1)\delta(x - 1), -\infty, \infty) = 3$. It is given that $A(cf(x), \dots) = cA(f(x), \dots)$ for some constant c .

**** Problem 1.15**

The *wavefunction* Ψ of a particle can be thought of as the quantum state of a particle. In this problem, you will solve for the wavefunction of an one-dimensional particle (that lies on the x -axis) in an infinite square well. It is given that $|\psi(x)|^2$ gives the probability of the particle existing at x and $|\Psi(x, t)|^2$ is the probability

14 If you are skeptical about the area function provides the area for any function, you are absolutely right. This property that a function can be integrated is called *integrability*. If you are interested, there are many types, such as the physicist favourite “square-integrability”. However, the point of the problem remains.

of a particle existing at x at time t . (Hence, note that $\Psi(x) \neq 0$ and $\psi(x) \neq 0$. That would imply that the particle cannot be found anywhere.) The potential function of the infinite square well is

$$V(x) = \begin{cases} 0 & 0 \leq x \leq a \\ \infty & \text{otherwise} \end{cases} \quad (1.3.1)$$

- a) Can the particle exist outside of $0 \leq x \leq a$, or more specifically, what is $|\psi(x)|^2$ (and therefore $\psi(x)$) for $x < 0$ or $x > a$?
 - b) To ensure that ψ is continuous, $\psi(0)$ and $\psi(a)$ are equal to your answer of $\psi(x)$ in (a). Also, it is known that the form of general solution inside the well (to wit: $0 \leq x \leq a$) to ψ is $\psi(x) = A \sin(kx) + B \cos(kx)$. Show that $\psi(x) = A \sin(kx)$.
 - c) Using your answer in (b), solve for k . (Hint: k has multiple solutions, generalize all the solutions by writing down k_n for $n \in \mathbb{Z} \setminus \{0\}$.)
 - d) Show that we can discard all negative solutions of k_n , hence $n \in \mathbb{N}$.
 - e) If $E(n) = \frac{\hbar^2 k_n^2}{2m}$, find $E(1)$ or the energy of the *ground state* of the particle. $\hbar$ is the reduced plank's constant. What happens as n increases?
 - f) The relation between Ψ and ψ is $\Psi(x, t) = \psi(x)e^{-\frac{iE(n)t}{\hbar}}$. Find $|\Psi|^2$ for $t = 1$ and $t = 2$, what happens to the graph of $|\Psi|^2$ as time passes? Take A, m, a as positive real numbers and choose some natural n .
-

1.4 An aside: A more rigorous definition of a function

TBD

Linear Algebra

2.1 Vector Basics

Stepping into this chapter, you may feel overwhelmed by the name, *Linear Algebra*. What is that? Please take note though, the knowledge in this chapter will not be used until Calculus III to IV in part IV of the book. If you are just willing to learn Calc I and II, you do not need the knowledge here. With that said, let's delve into Vectors. A concept that is very ubiquitous in this world is a coordinate, like the GPS coordinate in the maps app. A coordinate simply describes the position of a point in space. Generally, it is denoted (x, y) in 2 dimensions and (x, y, z) in 3 dimensions. Likewise, vectors also behave like coordinates and are written down in a *similar* fashion. Think of a vector as a coordinate.

Definition 2.1 Vector

A **vector** is a mathematical object with a magnitude, or length, and a direction. A vector, $\mathbf{a}$ is a point (x, y) in space drawn with an arrow. It is denoted as¹⁵:

$$\mathbf{a} = \begin{bmatrix} x \\ y \end{bmatrix}$$

Of course, vectors by themselves are not very useful. Hence, we shall define the vector space. For the purposes of this text, you may think of vector spaces as the 2 and 3 dimensional euclidean spaces you are very familiar with. In a vector space, we can finally meaningfully manipulate them. Take a look at the vector $\mathbf{a} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ and the vector $\mathbf{b} = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$ in the two dimensional euclidean space:

You may have noticed that the coordinate axes in Figure 2.1 are the same as the ones you are familiar with. However, in linear algebra, instead of the usual xy axes to describe the components of a vector, you have $\hat{\mathbf{i}}$ and $\hat{\mathbf{j}}$ to describe the components of a vector. (So instead of saying the “ x ” component of a vector, you would say the “ $\hat{\mathbf{i}}$ ” component of a vector.)¹⁶ This is a convention in linear algebra. $\hat{\mathbf{i}}$ translates to the x axis and $\hat{\mathbf{j}}$ translates to the y axis. Likewise, in the third dimension, $\hat{\mathbf{k}}$ represents the z axis. Another thing to note are the *basis vectors*:

¹⁵ Likewise, a vector at (x, y, z) is denoted $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$

¹⁶ Some authors use $\hat{\mathbf{x}}, \hat{\mathbf{y}}, \hat{\mathbf{z}}$ instead of the usual $\hat{\mathbf{i}}, \hat{\mathbf{j}}, \hat{\mathbf{k}}$. We shall use the latter.




Figure 2.1 Figure of vectors **a** and **b**.

Definition 2.2 Basis vectors

The set of vectors in a vector space are called **basis** if every vector in the vector space can be written uniquely written as a finite *linear combination*¹⁷ of the basis vectors. An example would be these vectors $\hat{\mathbf{i}} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\hat{\mathbf{j}} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ in the 2 dimensional euclidean space which can be thought of one unit of distance in the x and y direction respectively.



¹⁷ A linear combination of x and y is $ax + by$ where a, b are constants.

¹⁸ You can think of $\hat{\mathbf{i}}$ and $\hat{\mathbf{j}}$ as one unit of direction in the x and y axis respectively.

¹⁹ This can be extrapolated into any dimensions:

$$\begin{bmatrix} a \\ b \\ c \end{bmatrix} + \begin{bmatrix} d \\ e \\ f \end{bmatrix} = \begin{bmatrix} a+d \\ b+e \\ c+f \end{bmatrix}.$$

²⁰ Translation means moving something without rotation or any scaling.

With this in mind, another way of writing a vector $\mathbf{a} = \begin{bmatrix} x \\ y \end{bmatrix}$ is $\mathbf{a} = x\hat{\mathbf{i}} + y\hat{\mathbf{j}}$.¹⁸ Of course $\hat{\mathbf{i}}$ and $\hat{\mathbf{j}}$ are not the only basis vectors in 2d, if you have learned polar coordinates before, there is the equivalent $\hat{\mathbf{r}}$ and $\hat{\boldsymbol{\theta}}$, but this is a topic for a later chapter. Next, we shall look at the *addition* of vectors. A vector $\mathbf{a} = \begin{bmatrix} x_a \\ y_a \end{bmatrix}$ added to a vector $\mathbf{b} = \begin{bmatrix} x_b \\ y_b \end{bmatrix}$ is the vector $\begin{bmatrix} x_a+x_b \\ y_a+y_b \end{bmatrix}$. As you can see, the addition of two vectors results in another vector. Here is a illustration to help clear up things:¹⁹

Let us analyze Figure 2.2. Firstly, notice that $\mathbf{a} = 1\hat{\mathbf{i}} + 2\hat{\mathbf{j}}$ and $\mathbf{b} = 1\hat{\mathbf{i}} + 1\hat{\mathbf{j}}$. Thus, $\mathbf{a} + \mathbf{b} = 2\hat{\mathbf{i}} + 3\hat{\mathbf{j}}$, exactly like in our figure. One way to understand this is to “attach” the vector $\mathbf{a}$ ’s tail to $\mathbf{b}$ ’s head by translating $\mathbf{a}$.²⁰




Figure 2.2 Illustration of vector addition.

You may have seen this in your physics class before. Similarly, the subtraction of vectors $\mathbf{x} = \begin{bmatrix} a \\ b \end{bmatrix}$ and $\mathbf{y} = \begin{bmatrix} c \\ d \end{bmatrix}$ is $\mathbf{x} - \mathbf{y} = \begin{bmatrix} a-c \\ b-d \end{bmatrix}$. Of course, we can use the same “head to tail” model to explain vector subtraction. Subtracting $\mathbf{y}$ from $\mathbf{x}$ is simply finding the vector whose tail is on the head of $\mathbf{y}$ and whose head is on the head of $\mathbf{x}$. However, some of you may be scratching your heads right now, what if a vector does not start from the *origin* (or the coordinate $(0,0)$) but its tail lands on, say $(1,1)$. How do we express such a vector? The answer may be surprising, both vectors have the same representation. However, unless specifically stated, *all* vectors start from the origin. This is also why the “head to tail” model works, while the “tail” of the vector is not at the origin, translating it to the origin results in an *identical* vector.




Figure 2.3
Illustration of vector “head to tail” addition.

Example 2.1 Addition of vectors

Here are some examples of vectors:

$$\mathbf{a} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad \mathbf{b} = \begin{bmatrix} -3 \\ 4 \end{bmatrix} \quad \mathbf{c} = \begin{bmatrix} 2 \\ -2 \end{bmatrix}$$

Here are some examples of adding and subtracting vectors. Note that addition are *commutative*, in other words the order does not matter:

$$\mathbf{a} + \mathbf{b} = \begin{bmatrix} 1 - 3 \\ 2 + 4 \end{bmatrix} = \begin{bmatrix} -2 \\ 6 \end{bmatrix}$$

$$\mathbf{a} - \mathbf{c} = \begin{bmatrix} -1 \\ 4 \end{bmatrix}$$

$$\mathbf{a} + \mathbf{b} - \mathbf{c} = \begin{bmatrix} -2 \\ 6 \end{bmatrix} - \mathbf{c} = \begin{bmatrix} -4 \\ 8 \end{bmatrix}$$






Figure 2.4 Three identical vectors.

Given a vector $\mathbf{a}$, $-\mathbf{a}$ can be thought of the same vector with the same magnitude but pointing in the *opposite* direction. To find it, simply flip the signs of all the components in $\mathbf{a}$. If you add $\mathbf{a}$ to $-\mathbf{a}$, you get what is called the *zero vector*, or the vector $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$. This is a vector without direction has zero magnitude. We shall denote it $\mathbf{0}$. Lastly, I want to briefly touch on the idea of *magnitude* which is basically the length of a vector. The derivation of the following formula is simply the Pythagorean theorem.

Definition 2.3 Magnitude of vector

The **magnitude** of a vector $\mathbf{v}$ in n -dimensional space is as such:

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}$$

A **unit** vector is a vector with magnitude 1. A **zero** vector has magnitude 0 and is denoted $\mathbf{0}$.




Figure 2.5
Visualisation of
“head to tail” addition
of $\mathbf{a}$ and $\mathbf{b}$.

Try to attempt the following exercise:

Problem 2.1

Given that $\mathbf{a} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$ and $\mathbf{c} = \begin{bmatrix} -3 \\ -2 \end{bmatrix}$, calculate:

- $\mathbf{a} + \mathbf{b}$
- $\mathbf{a} - \mathbf{c}$
- $\mathbf{a} - \mathbf{b} - \mathbf{c}$

Problem 2.2

Are the following statements true or false:

- a) The subtraction of two vectors *always* results in a vector.
- b) When any vector is added to the zero vector, the result *always* is not a zero vector.

Problem 2.3

Find the magnitude of $\begin{bmatrix} 3 \\ 4 \end{bmatrix}$ and $\begin{bmatrix} 3 \\ 4 \\ 5 \end{bmatrix}$.

*** Problem 2.4**

Prove that the addition of vectors is *associative* given that the addition in $\mathbb{R}$ is associative.²¹

*** Problem 2.5**

Consider the 3 dimensional vector space where all vectors are described by $a_x \hat{\mathbf{i}} + a_y \hat{\mathbf{j}} + a_z \hat{\mathbf{k}}$.

- a) Consider the vector space which is the subset of vectors where $a_z = 0$. What is the dimension of this vector space?
- b) Prove that the components of a vector with respect to this basis $(\hat{\mathbf{i}}, \hat{\mathbf{j}}, \hat{\mathbf{k}})$ are unique.

²¹ An operation $\circ$ is associative if order in which it is applied does not matter, in this case $(\mathbf{x} + \mathbf{y}) + \mathbf{z} = \mathbf{x} + (\mathbf{y} + \mathbf{z})$.

2.2 Scaling and dot product

In this chapter we will deal two kinds of multiplication with vectors. By far the easiest would be scaling a vector. As we have previously discussed, the magnitude of a vector is the length of a vector. But what if I want to “elongate” the vector by some amount to make it longer or shorter. This is called *scaling*.




Figure 2.6 Two vectors, one twice as long as the other.

Take a look at Figure 2.6, $\mathbf{b}$ is twice as long as $\mathbf{a}$. What should we multiply to $\mathbf{a}$ to make as long as $\mathbf{b}$? If you guessed 2, then you are right. (Not sure what else you could have guessed.) To multiply a vector by a constant, we multiply each individual component of the vector by the constant. From the figure, we can identify that $\mathbf{a} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$ so $2\mathbf{a} = \begin{bmatrix} 2 \times 2 \\ 2 \times 2 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \end{bmatrix}$ which is equal to $\mathbf{b}$. You may extrapolate this to higher dimensions.

Definition 2.4 Scaling a vector

Scaling a n -dimensional vector $\mathbf{v}$ by a constant n is defined as such:

$$n\mathbf{v} = n \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix} = \begin{bmatrix} n \times v_1 \\ \vdots \\ n \times v_n \end{bmatrix}$$




Figure 2.7 Two vectors, one is scaled by -0.5.

Let us try something fun, what if we multiplied the vector by a negative number? Well, the computation remains the same.

As you can see from Figure 2.7, $\mathbf{a} = (-0.5)\mathbf{b}$. You may wish to compute this yourself to verify. One thing to note here is that the vector gets flipped across the origin when the constant the vector is multiplied to is negative.

Challenge 2.1 Scaled vector magnitude

Also, when scaling a vector by some constant, the reader may want to verify that the magnitude of the vector is also scaled by that constant. This should be a trivial proof.



You may have noticed that I have abstained from using either the $\cdot$ or $\times$ operators. The reason is bad notation which is sprinkled throughout mathematics. These two operators have special meanings in the realm of linear algebra. We shall first explore the easier one, the *dot product*.

Dot products use the $\cdot$ operator and only acts on vectors, do not abuse notation by using the $\cdot$ operator in the context of scaling a vector. Dot products are actually fairly straightforward to compute:

Definition 2.5 Dot product

The dot product of two vectors in n -dimensional space $\mathbf{a}$ and $\mathbf{b}$ is:

$$\mathbf{a} \cdot \mathbf{b} = a_1 b_1 + a_2 b_2 + \cdots + a_n b_n = (||\mathbf{a}||)(||\mathbf{b}||) \cos \theta \quad (2.2.1)$$

where θ is the angle between the two vectors. We shall refer to the formula between the first and second equal signs the “first definition of dot products” and the formula after the second equal sign the “second definition of dot products”.



From Equation (2.2.1), we can tell that the dot product tells us something about the relationship between the two vectors. In essence, if the dot product between two vectors is large and positive, then the vectors are pointing in roughly the same direction. Else, if the product is very negative, then the dot product is very negative.

Firstly, note down the components of each vector in Figure 2.8. Then, let us compute $\mathbf{a} \cdot \mathbf{b}$. By Equation (2.2.1), it is $1(-2) + 2(1) = 0$. Also, it is important to take note that these two vectors are orthogonal (at right angles with each other). This brings us to an important property of dot products, if two vectors are orthogonal, then their dot product is 0. This can be seen from the fact that $\cos\left(\frac{\pi}{2}\right) = 0$ and from the second definition of dot products in Equation (2.2.1). In a sense, the dot product tells you how aligned are the directions of two vectors. Another example would be taking the dot product of two identical vectors, such as $\mathbf{a} \cdot \mathbf{a}$. The dot product there is $1(1) + 2(2) = 5$. In addition, we know $\cos 0 = 1$ so according to the second definition, $\mathbf{a} \cdot \mathbf{a} = (||\mathbf{a}||)(||\mathbf{a}||) = (||\mathbf{a}||)^2$. If we take the square root on both sides, $||\mathbf{a}|| = \sqrt{\mathbf{a} \cdot \mathbf{a}}$ which can be rewritten as $||\mathbf{a}|| = \sqrt{a_1^2 + \cdots + a_n^2}$. Drum roll please. This is exactly the formula we got for the magnitude of a vector. This is a pretty nice result.

Example 2.2 Calculating dot products

From Figure 2.8, we can see that:

$$\begin{aligned} \mathbf{a} \cdot \mathbf{c} &= \begin{bmatrix} 1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix} = 1(2) + 2(1) = 4 \\ \mathbf{a} \cdot \mathbf{d} &= \begin{bmatrix} 1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} -1 \\ -2 \end{bmatrix} = 1(-1) + 2(-2) = -5 \end{aligned}$$






Figure 2.8 Four vectors pointing in different directions.

Remark 2.1

Generally, when finding the dot product, we use the first form in Equation (2.2.1) as it is typically easier to compute. It is also important to know that the dot product of two vectors results in a number, not a vector like in scaling.




Figure 2.9 Figure of a triangle formed by three vectors.

Problem 2.6

If $\mathbf{x} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$, compute the following:

- a) $2\mathbf{x}$
- b) $\mathbf{x} \cdot \mathbf{y}$
- c) $\mathbf{y} \cdot \mathbf{x}$

*** Problem 2.7**

Prove or disprove that the dot product *commutes*.²²

*** Problem 2.8**

Newton's second law states that $\mathbf{F} = m\mathbf{a}$. Consider an object with a mass 10kg in the 2d plane.

²² An operation $\circ$ is commutative if $x \circ y = y \circ x$.

- a) If the object is accelerating in the x-axis with a magnitude of 3m/s^2 and accelerating in the y-axis with a magnitude of 4m/s^2 , what is the x component of the force?
- b) What is the y-component of the force in (a). Hence, what is the magnitude of the force.
-

2.3 Matrices




Figure 2.10 Two vectors, one rotated anti-clockwise by 90° .

In this chapter, we will introduce a new mathematical object that you have probably heard before but not sure what it is, *linear transformation*. A great place to start is the *basis vectors*. All vectors can be written as a unique linear combination of the basis vectors. But $\hat{i}$ and $\hat{j}$ are not the only basis vectors out there, what if I wanted to describe vectors using another set of basis vectors? Perhaps that would be more convenient for my special use case. Well, it is actually pretty easy to do so. For the sake of this argument, let us try to rotate all of the vectors in a normal 2d vector field anti-clockwise by 90° .

Notice how we could not describe $\mathbf{b}$ as some constant multiplied to $\mathbf{a}$ in Figure 2.10. We need some other tool to help us. If we change the basis vectors of $\mathbf{a}$ to another pair of basis vectors, can we somehow get $\mathbf{b}$? Here is a thought: replace the $\hat{i}$ with $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ and $\hat{j}$ with $\begin{bmatrix} -1 \\ 0 \end{bmatrix}$. So, $\mathbf{a} = 1\hat{i} + 2\hat{j}$ will become $\mathbf{a} = \begin{bmatrix} 0 \\ 1 \end{bmatrix} + 2\begin{bmatrix} -1 \\ 0 \end{bmatrix} = \begin{bmatrix} -2 \\ 1 \end{bmatrix}$. This is pretty intriguing, by changing the basis vectors, we have basically “morphed” $\mathbf{a}$ into $\mathbf{b}$. This “morphing” is called a *linear transformation* or *linear map*. In some of the visualisations of the linear transformations I might keep the original grid in the background in a lighter color.

Example 2.3 Linear transformations

Here are some examples of linear transformations, the blue grid is the vector space after the transformation. Take note where the basis vectors are after the transformations.




Figure 2.11 A transformation in the form of an anti-clockwise *rotation*.




Figure 2.12 A transformation in the form of a *shear*.

23

²³ In Figure 2.12, the $\hat{i}$ vector remains at the same location.

To describe these transformations, we actually only need to describe where the basis vectors land after the transformation. No other information required. We present the location of the new basis vectors in the form of a *matrix*:

Definition 2.6 Matrices

In linear algebra, matrices generally represent linear transformations. A $n \times m$ matrix is one with n rows and m columns. The n th column of the matrix denotes the representation of the corresponding basis vector after the linear transformation.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

represents a matrix that moves the $\hat{i}$ basis vector to $\begin{bmatrix} a \\ c \end{bmatrix}$ and moves the $\hat{j}$ basis vector to $\begin{bmatrix} b \\ d \end{bmatrix}$.

Remark 2.2

Matrices can also represent a variety of things apart from linear transformations. It is just a mathematical object representing a bunch of things arranged in rows and columns.

It should go without saying that any vector in the original vector space will get moved after a transformation. To calculate where a vector ends up after a transformation, we multiply the matrix by the vector. This is where many textbooks make linear algebra feel laborious. In order to make the process more intuitive, recall that a vector has a $\hat{i}$ and $\hat{j}$ component. Now, we swap the basis vectors for the new basis vectors described in our matrix. This is just multiplying a vector by a constant, which you have done before.

Example 2.4 Deriving a linearly transformed vector

Take the transformation in Figure 2.12 which is represented by the matrix $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. Let us say in the original vector space, there is a vector $\mathbf{a} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$. Then, the vector's representation after the transformation:

$$\begin{aligned} \begin{bmatrix} 1 \\ 2 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} &= 1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} 3 \\ 2 \end{bmatrix} \end{aligned}$$




Figure 2.13 Figure of a vector before and after a shear transformation like in Figure 2.12.

If you have come from a linear algebra course, this may feel very refreshing. Instead of memorising what needs to be multiplied, you just swap the basis vectors for some other basis vectors. While we get the representation of $\mathbf{a}$ from Figure 2.13, it also important to take note that in the new vector space after the transformation, $\mathbf{a}$ has one unit of direction in the horizontal direction and two units of direction in the “vertical” direction. This is just like the original representation of $\mathbf{a}$, just that the vector space has been sheared. This change of basis vector is important enough to have its own name, *change of basis*.

With this knowledge, let us proceed to one of the most hated parts of linear algebra, *matrix multiplication*.

These few words can bring out so much frustration out of so many students. It is quite a laborious and boring process if you do not grasp the meaning and motivation behind it. In actuality, the idea is simple. When you multiply two matrices together, you are effectively applying two linear transformations, one after another. The result is also a matrix, which is a linear transformation, hence the operation is *closed*.²⁴

²⁴ This is only true if the matrix is $n \times n$, which are the ones we are dealing with.

Example 2.5 Deriving matrix multiplication

Say you have two matrices, a 90° anti-clockwise rotation $x = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ and a horizontal shear $y = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. xy is simply applying the shear first before the rotation. (The right to left reading is similar to that of function composition)




Figure 2.14 Figure of applying two linear transformations.

It should not be difficult to see that the end result of applying the two linear transformations in the previous examples resulted in another linear transformation (the red grid). We wish to find the representation of this linear transformation as a matrix. Let us go through this step by step. First of all, the basis vectors of the identity matrix²⁵, gets moved to some new location. In other words, given a matrix $a = \begin{bmatrix} a_1 & a_2 \\ a_3 & a_4 \end{bmatrix}$, we know that the basis vectors gets moved to $\begin{bmatrix} a_1 \\ a_3 \end{bmatrix}$ and $\begin{bmatrix} a_2 \\ a_4 \end{bmatrix}$ respectively for the $\hat{i}$ and $\hat{j}$ vectors. Now, we have another transformation $b = \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix}$. To know where the $\hat{i}$ vector lands after both transformations, notice that it is simply multiplying the intermediate position of it by b , which is $\begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix} \begin{bmatrix} a_1 \\ a_3 \end{bmatrix} = \begin{bmatrix} a_1 b_1 + a_3 b_2 \\ a_1 b_3 + a_3 b_4 \end{bmatrix}$. A similar line of reasoning may be used for $\hat{j}$. Hence, the result is:

²⁵ The identity matrix is simply the transformation that leaves the vector space unchanged.

$$ba = \begin{bmatrix} a_1 b_1 + a_3 b_2 & a_2 b_1 + a_4 b_2 \\ a_1 b_3 + a_3 b_4 & a_2 b_3 + a_4 b_4 \end{bmatrix}$$

This is the reason many students find linear algebra tedious. Most courses make students memorise the

formula which is extremely rote. I wish that you understand what is going on behind the scenes on this formula. With that said, it is best to get some experience with these multiplications:

Example 2.6 Matrix multiplication

Given that $x = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and $y = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$ and $z = \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix}$,

$$xy = \begin{bmatrix} 5 & 11 \\ 11 & 25 \end{bmatrix}$$

$$yx = \begin{bmatrix} 10 & 14 \\ 14 & 20 \end{bmatrix}$$

$$x(yz) = \begin{bmatrix} 102 & 118 \\ 230 & 266 \end{bmatrix}$$

$$(xy)z = \begin{bmatrix} 102 & 118 \\ 230 & 266 \end{bmatrix}$$

This examples shows that matrix multiplication does not commute but it is associative.



Problem 2.9

If $x = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, $a = \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix}$ and $b = \begin{bmatrix} 6 & 7 \\ 8 & 9 \end{bmatrix}$, compute:

- a) ax
- b) ab
- c) abx

* Problem 2.10

Prove that matrix multiplication is *associative* for all 2×2 matrices.

* Problem 2.11

The *trace* of an $n \times n$ matrix A is defined as

$$\text{tr}(A) = \sum_{i=1}^n a_{ii} \quad (2.3.1)$$

where a_{ii} denotes the entry for the i -th column and i -th row of A .

- a) Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$, what is the trace of A .
- b) Prove that, for square matrices A and B , $\text{tr}(A + B) = \text{tr}(A) + \text{tr}(B)$.
- c) Hence, if $B = \begin{bmatrix} 3 & 1 & 4 \\ 1 & 5 & 9 \\ 2 & 6 & 5 \end{bmatrix}$, verify that $\text{tr}(A + B) = \text{tr}(A) + \text{tr}(B)$.

2.4 Determinants, inverses and Cramer's rule

At this juncture, you should be familiar with the concept that a matrix represents a linear transformation and it morphs the vector space. Also, if you imagine the basis vectors of any vector space as lengths of a parallelogram, then you might want to find the area of that parallelogram. For example, the identity matrix's basis vectors forms a square with area 1. Likewise, given a matrix $x = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, we can display it as:




Figure 2.15 Figure of area A formed by a pair of basis vectors.

To find the area of A in Figure 2.15, we may use the sine rule (let $\mathbf{u} = \begin{bmatrix} a \\ c \end{bmatrix}$ and $\mathbf{v} = \begin{bmatrix} b \\ d \end{bmatrix}$):

$$\begin{aligned}
 A &= (||\mathbf{u}||)(||\mathbf{v}||) \sin(\theta) \quad \text{where } \theta \text{ is the angle between the vectors} \\
 &= (||\mathbf{u}'||)(||\mathbf{v}||) \cos(\theta') \quad \text{where } \mathbf{u}' \text{ is } \begin{bmatrix} a \\ -c \end{bmatrix}, \text{ perpendicular to } \mathbf{v} \text{ and } \theta' \text{ is } 90^\circ \\
 &= \begin{bmatrix} -c \\ a \end{bmatrix} \cdot \begin{bmatrix} b \\ d \end{bmatrix} \quad \text{by Equation (2.2.1), definition of dot product} \\
 &= ad - bc
 \end{aligned} \tag{2.4.1}$$

The resultant area is also known as the *determinant* of a matrix.

Definition 2.7 Determinant of 2×2 matrix

The determinant of a 2×2 matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ is represented as:

$$\det(A) = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

and it represents the signed area of a parallelogram with sides of the basis vectors described in A .

Meanwhile, a 3×3 matrix $B = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$ has a determinant of:

$$\begin{aligned} \det(B) &= \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} \\ &= a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix} \end{aligned}$$

and it represents the volume of the parallelepiped formed by the three basis vectors in B .



Example 2.7 Finding determinants

Given that $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 5 \\ 2 & 5 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$:

$$\begin{aligned} \det(A) &= 1 \times 4 - 2 \times 3 \\ &= -2 \end{aligned}$$

$$\det(B) = 0 \quad (2.4.2)$$

26

$$\begin{aligned} \det(C) &= 1 \begin{vmatrix} 5 & 6 \\ 8 & 9 \end{vmatrix} - 2 \begin{vmatrix} 4 & 6 \\ 7 & 9 \end{vmatrix} + 3 \begin{vmatrix} 4 & 5 \\ 7 & 8 \end{vmatrix} \\ &= (5 \times 9 - 8 \times 6) - 2(4 \times 9 - 6 \times 7) + 3(4 \times 8 - 7 \times 5) \\ &= 0 \end{aligned}$$



Next, we would move onto the *inverse* of a matrix. Think of a number x , $x^{-1} \times x = 1$, 1 is also called the multiplicative identity. Likewise, given a matrix Y , $YY^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ which is the identity matrix. If you think of Y as a transformation, then Y^{-1} “undo” the transformation of Y .

Definition 2.8 Inverse of a matrix

The (multiplicative) inverse of a matrix X is denoted X^{-1} where:

$$XX^{-1} = I$$

where I is the identity matrix. If X has an inverse, it is called an *invertible* matrix. X has an inverse if and only if $\det(X) \neq 0$. To find the inverse of a matrix $Y = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, we use the following formula:

$$Y^{-1} = \frac{1}{\det(Y)} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \quad (2.4.3)$$

²⁶ Equation (2.4.2) shows a more general property, given that $X = \begin{bmatrix} a & b \\ a & b \end{bmatrix}$, $\det(X) = 0$. The proof is trivial and shall be left as a verification to the reader.



27 When multiplying a matrix by a constant, you multiply each entry in matrix by the constant.

Example 2.8 G

Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 5 \\ 2 & 5 \end{bmatrix}$:

$$\begin{aligned} A^{-1} &= \frac{1}{-2} \begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix} \\ &= \begin{bmatrix} -2 & 1 \\ 1.5 & -0.5 \end{bmatrix} \end{aligned}$$

$\det(B) = 0$, hence B is not invertible.



Until now, this chapter has been focusing mainly on computation. The reader may feel that these computations are excessively dull and boring. That being said, here is an application of determinants, *Cramer's rule*. Cramer's rule may be used to solve a system of linear equations. Firstly, we will represent a simultaneous equation as a product of vectors and matrices. Given a simultaneous equation:

$$\begin{aligned} x + y &= 10 \\ 2x - y &= 14 \end{aligned} \tag{2.4.4}$$

We write it as:

$$\begin{bmatrix} 1 & 1 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 10 \\ 14 \end{bmatrix} \tag{2.4.5}$$

This is also called the “matrix form” of a simultaneous equation. It is also important to recognize that Equation (2.4.5) is in the form of $A\mathbf{v} = B$, it is easy to see that $\mathbf{v} = BA^{-1}$. From Equation (2.4.3), we can find BA^{-1}

$$\begin{aligned} BA^{-1} &= \begin{bmatrix} 10 \\ 14 \end{bmatrix} \begin{bmatrix} \frac{1}{3} & \frac{1}{3} \\ \frac{2}{3} & -\frac{1}{3} \end{bmatrix} \\ &= \begin{bmatrix} 8 \\ 2 \end{bmatrix} \end{aligned}$$

This is also the solution for Equation (2.4.5), x is indeed 8 and y is indeed 2. However, this method might not be the best. If there were three unknowns and three equations, how could you solve the equations? We may apply Cramer's rule. Consider a system of n linear equations with n unknowns $(x_1, x_2, \dots, x_n)$, it can be represented as $Ax = B$ as we have seen before. The solutions for the equations are:

$$x_i = \frac{\det(A_i)}{\det(A)} \quad i = 1, \dots, n$$

Where x_i is the i -th solution and A_i is the matrix formed by replacing the i -th column of A by B .

Example 2.9 Solving linear equations with Cramer's rule

Given the following system of linear equations of three unknowns:

$$x_1 + x_2 + x_3 = 12$$

$$x_1 - x_2 + x_3 = 4$$

$$x_1 - x_2 - x_3 = -6$$

and its matrix form is:

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & -1 & -1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 12 \\ 4 \\ -6 \end{bmatrix}$$

The solutions are:

$$\begin{aligned} x_1 &= \frac{\det(A_1)}{\det(A)} \\ &= \frac{\det \left(\begin{bmatrix} 12 & 1 & 1 \\ 4 & -1 & 1 \\ -6 & -1 & -1 \end{bmatrix} \right)}{\det \left(\begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & -1 & -1 \end{bmatrix} \right)} \\ &= \frac{12}{4} \\ &= 3 \end{aligned}$$

$$\begin{aligned} x_2 &= \frac{\det(A_2)}{\det(A)} \\ &= \frac{\det \left(\begin{bmatrix} 1 & 12 & 1 \\ 1 & 4 & 1 \\ 1 & -6 & -1 \end{bmatrix} \right)}{\det \left(\begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & -1 & -1 \end{bmatrix} \right)} \\ &= \frac{16}{4} \\ &= 4 \end{aligned}$$

$$x_3 = \frac{\det(A_3)}{\det(A)}$$

$$\begin{aligned} & \det \begin{pmatrix} 1 & 1 & 12 \\ 1 & -1 & 4 \\ 1 & -1 & -6 \end{pmatrix} \\ &= \frac{\det \begin{pmatrix} 1 & 1 & 12 \\ 1 & -1 & 4 \\ 1 & -1 & -6 \end{pmatrix}}{\det \begin{pmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & -1 & -1 \end{pmatrix}} \\ &= \frac{20}{4} \\ &= 5 \end{aligned}$$



As you can tell, Cramer's rule is very powerful. In the real world, however, *gaussian elimination* is a much faster alternative than Cramer's rule. It is also unlikely for the reader to be computing large matrices determinants by hand, we have computers for that. This also applies to matrix multiplication. This book, and more generally calculus courses only need knowledge on computing determinants of up to 3×3 matrices.

Problem 2.12

Find the determinants and inverses, if any, for the following matrices:

a) $\begin{bmatrix} 1 & 9 \\ 2 & 10 \end{bmatrix}$

b) $\begin{bmatrix} 3 & 2 \\ 6 & 4 \end{bmatrix}$

c) $\begin{bmatrix} 3 & 4 \\ 7 & 9 \end{bmatrix}$

*** Problem 2.13**

Find the determinant of the following matrix: $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 2 & 3 & 6 \end{bmatrix}$

*** Problem 2.14**

Solve the following simultaneous equations using matrices:

$$\begin{aligned} x_1 + x_2 + x_3 &= 3 \\ x_1 + 2x_2 + x_3 &= 4 \\ x_1 + 2x_2 + 2x_3 &= 6 \end{aligned}$$

2.5 Cross products and triple products

The previous section perfectly segues into this section. *Cross products* is an operation on vectors that happens solely in the third dimension. Do not confuse it with the dot product, both uses different operators and should not be used interchangeably. Firstly, we must recall that one of the basis vectors is $\hat{\mathbf{k}}$ and it is one unit in the z direction.




Figure 2.16 Figure of the three basis vectors.

The cross product is a vector, whose magnitude is the area of a parallelogram formed by the two operands (which are vectors).²⁸ More formally, the cross product is defined as:

Definition 2.9 Cross product

The **cross product** is an operation on two vectors and the result is another vector which is perpendicular to both vectors. It is defined by the formula:

$$\mathbf{a} \times \mathbf{b} = (||\mathbf{a}||)(||\mathbf{b}||) \sin(\theta) \mathbf{n}$$

where θ is the angle between $\mathbf{a}$ and $\mathbf{b}$ and $\mathbf{n}$ is the unit vector perpendicular to both of the operands such that all three vectors obey the “right-hand rule”²⁹. The common way to compute the cross product is using the following formula:

$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ a_x & a_y & a_z \\ b_x & b_y & b_z \end{vmatrix} \quad (2.5.1)$$

Lastly, do not confuse $\mathbf{n}$ with $\hat{\mathbf{k}}$ as both vectors are sometimes not the same.

²⁸ The operands of $\mathbf{a} \times \mathbf{b}$ are $\mathbf{a}$ and $\mathbf{b}$.

²⁹ The right hand rule states that if you use your right hand and point the index finger in the direction of $\mathbf{a}$, your middle finger in the direction of $\mathbf{b}$, your thumb should be pointing in the direction of $\mathbf{n}$.

The cross product is very ubiquitous in physics and vector calculus alike.

Example 2.10 Calculating cross products

iven that $\mathbf{a} = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}$, then $\mathbf{a} \times \mathbf{b}$ would be:

$$\begin{aligned} \mathbf{a} \times \mathbf{b} &= \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ 1 & 2 & 0 \\ -2 & 1 & 0 \end{vmatrix} \\ &= 0\hat{\mathbf{i}} - 0\hat{\mathbf{j}} + 5\hat{\mathbf{k}} \\ &= 5\hat{\mathbf{k}} \end{aligned}$$




Figure 2.17 Figure of two vectors and their cross product.

An important fact to note is that the operation is not commutative. In fact, the operation is called *anti-commutative*. This means that $\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a}$. In addition, if the angle between the operands is any integer multiple of π , then the result is $\mathbf{0}$. This should be trivially seen from the formula for the cross product. The derivation for the cross product is also easy, given two vectors $\mathbf{a}$ and $\mathbf{b}$ and an angle θ between them, simply use the sine rule and the result should be obvious. The cross product has many real world applications.

Application: Torque

Example 2.11 Moment arm formula

Torque is a force that is common in our daily lives, present in the turning of a doorknob or the turning of a wrench. The formula for torque is $\boldsymbol{\tau} = \mathbf{r} \times \mathbf{F}$, where $\boldsymbol{\tau}$ is the torque, $\mathbf{r}$ is the distance from the pivot point to the point where the force is applied and $\mathbf{F}$ is the force applied. The direction of the torque can be determined by the right-hand rule. Torque is also known as the moment of force, and we actually derive the infamous moment arm formula for moment of *moment = (moment arm)(force)*. We shall restrict ourselves to only two dimensions and use a Cartesian coordinate system. Set the origin such that the pivot point is at the origin. Then, we can represent $\mathbf{r}$ as the distance from the origin to where the force is applied. Let $\mathbf{F}$ be the force vector. Since there is no more z component, all of the z components of the vectors are 0. We can write down the cross product of $\mathbf{r}$ and $\mathbf{F}$ (we resolve each into its x and y components):




Figure 2.18 Figure of torque in two dimensions.

$$\begin{aligned}
 \boldsymbol{\tau} &= \mathbf{r} \times \mathbf{F} \\
 &= \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ r_x & r_y & 0 \\ F_x & F_y & 0 \end{vmatrix} \\
 &= (r_x F_y - r_y F_x) \hat{\mathbf{k}}
 \end{aligned}$$

We can consider the magnitude of the torque:

$$||\boldsymbol{\tau}|| = r_x F_y - r_y F_x \quad (2.5.2)$$

Next, we turn to the moment arm formula.

$$\begin{aligned}
 \text{moment} &= (\text{moment arm})(\text{force}) \\
 &= \sqrt{r_x^2 + r_y^2} \sqrt{F_x^2 + F_y^2}
 \end{aligned}$$

We are able to write this as I have assumed that the force and distance are perpendicular to each other (See Problem 2.18). Then, because the force and radius are perpendicular, notice that $\mathbf{r} \cdot \mathbf{F} = r_x F_x + r_y F_y = 0$. We can utilize this fact by writing down:

$$(r_x F_y - r_y F_x)^2 + (r_x F_x + r_y F_y)^2 = (r_x^2 + r_y^2)(F_x^2 + F_y^2) \quad (2.5.3)$$

Notice that the left hand side of Equation (2.5.3) is simply $||\boldsymbol{\tau}||^2$ and the right hand side is simply $(\text{moment arm})^2(\text{force})^2$. Hence, we have shown that $||\boldsymbol{\tau}|| = (\text{moment arm})(\text{force})$ which is the moment arm formula. An immediate difference between torque and moment is that moment is a scalar while torque is a vector. Another fact about torque that it has a $\hat{\mathbf{k}}$ component, which means that it is perpendicular to the plane formed by $\mathbf{r}$ and $\mathbf{F}$.



Next, we will move on to triple products which happens to also be a three dimensional topic. Firstly, take a look at the following three vectors:




Figure 2.19 Figure of triple vector multiplication.

Figure 2.19 is the example of triple vector multiplication. You would represent it as $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$. The result here is simply the volume of the parallelepiped formed in Figure 2.19.

Definition 2.10 Triple vector product

The **triple vector product** is a product of three 3-dimensional vectors. The most common of which, is the scalar triple product which is Figure 2.19. The formula for which is $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$, which yields the volume of the parallelepiped with the sides being the three vectors. Another type of triple vector multiplication is vector triple product. It is simply $\mathbf{a} \times (\mathbf{b} \times \mathbf{c})$. We can use **Lagrange's formula** to compute it easily: $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{c}(\mathbf{a} \cdot \mathbf{b})$.



Example 2.12 Volume of parallelepiped

We shall derive the formula for the volume of a parallelepiped given its height, breadth and length. We shall denote them h , b and l respectively. Notice that we define the sides of the parallelepiped like this:

$$\mathbf{a} = \begin{bmatrix} l \\ 0 \\ 0 \end{bmatrix}$$

$$\mathbf{b} = \begin{bmatrix} 0 \\ b \\ 0 \end{bmatrix}$$

$$\mathbf{c} = \begin{bmatrix} 0 \\ 0 \\ h \end{bmatrix}$$

Now, we use the triple vector multiplication formula. However, take note of the brackets and the order

of operations:

$$\begin{aligned}
 \mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) &= \mathbf{a} \cdot \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ 0 & b & 0 \\ 0 & 0 & h \end{vmatrix} \\
 &= \begin{bmatrix} l \\ 0 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} bh \\ 0 \\ 0 \end{bmatrix} \\
 &= lbh
 \end{aligned} \tag{2.5.4}$$

The result in Equation (2.5.4) is trivial and should come as no surprise to the reader.



As for vector triple product, the process of computation is very similar. We first evaluate the cross product of the two vectors in parentheses, then take the cross product of the first and resultant vector from the previous computation. We shall give a simple proof for Lagrange's formula.

Theorem 2.1 Lagrange's formula

Given $\mathbf{a}, \mathbf{b}, \mathbf{c}$, show that $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{c}(\mathbf{a} \cdot \mathbf{b})$.

QED

30

Proof 2.1 Lagrange's formula

We will start by evaluating $\mathbf{a} \times (\mathbf{b} \times \mathbf{c})$:

$$\begin{aligned}
 \mathbf{a} \times (\mathbf{b} \times \mathbf{c}) &= \mathbf{a} \times \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ b_x & b_y & b_z \\ c_x & c_y & c_z \end{vmatrix} \\
 &= \mathbf{a} \times \begin{bmatrix} b_y c_z - b_z c_y \\ b_z c_x - b_x c_z \\ b_x c_y - b_y c_x \end{bmatrix} \\
 &= \begin{vmatrix} \hat{\mathbf{i}} & \hat{\mathbf{j}} & \hat{\mathbf{k}} \\ a_x & a_y & a_z \\ b_y c_z - b_z c_y & b_z c_x - b_x c_z & b_x c_y - b_y c_x \end{vmatrix} \\
 &= \begin{bmatrix} a_y(b_x c_y - b_y c_x) - a_z(b_z c_x - b_x c_z) \\ \dots \\ a_y b_x c_y + a_z b_x c_z - a_y b_y c_x - a_z b_z c_x \end{bmatrix} \\
 &= \begin{bmatrix} a_y b_x c_y + a_z b_x c_z - a_y b_y c_x - a_z b_z c_x \\ \dots \end{bmatrix}
 \end{aligned} \tag{2.5.5}$$

³⁰ A sidenote here, when it is written $\mathbf{ab}$, this is not the cross or dot product, instead it is equal to $\begin{bmatrix} a_x b_x \\ a_y b_y \\ \dots \end{bmatrix}$.

We will examine the $\hat{\mathbf{i}}$ component of 2.5.5, the other two components are similar. Then, notice that:

$$\begin{aligned} \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{c}(\mathbf{a} \cdot \mathbf{b}) &= \mathbf{b}(a_x c_x + a_y c_y + a_z c_z) - \mathbf{c}(a_x b_x + a_y b_y + a_z b_z) \\ &= \begin{bmatrix} b_x(a_x c_x + a_y c_y + a_z c_z) \\ b_y(a_x c_x + a_y c_y + a_z c_z) \\ b_z(a_x c_x + a_y c_y + a_z c_z) \end{bmatrix} - \begin{bmatrix} c_x(a_x b_x + a_y b_y + a_z b_z) \\ c_y(a_x b_x + a_y b_y + a_z b_z) \\ c_z(a_x b_x + a_y b_y + a_z b_z) \end{bmatrix} \\ &= \begin{bmatrix} a_y b_x c_y + a_z b_x c_z - a_y b_y c_x - a_z b_z c_x \\ \dots \end{bmatrix} \end{aligned} \quad (2.5.6)$$

As you can see, the $\hat{\mathbf{i}}$ components of 2.5.5 and 2.5.6 are identical. Hence, we have shown Lagrange's formula.

QED

Remark 2.3

The above proof is a tedious pedagogical exercise mainly created to drill in the concepts of the cross products.



Problem 2.15

Find the cross products of the following vectors:

a) $\begin{bmatrix} 1 \\ 2 \\ 5 \end{bmatrix} \times \begin{bmatrix} 1 \\ 2 \\ 7 \end{bmatrix}$

b) $\begin{bmatrix} 0 \\ 2 \\ 4 \end{bmatrix} \times \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix}$

* Problem 2.16

Prove or disprove:

a) $\mathbf{a} \times \mathbf{b} = \mathbf{b} \times \mathbf{a}$

b) $\mathbf{a} \times \mathbf{a} = \mathbf{0}$

* Problem 2.17

Prove Lagrange's formula for the other two components not shown in the previous proof.

Further problems

* Problem 2.18

In this problem, you will derive the formula for the moment arm when the force and distance are not perpendicular. Refer to Figure 2.18 for when the force is perpendicular to the moment arm. Let the angle between $\mathbf{r}$ and $\mathbf{F}$ be θ .

- Find the expression for the force that is perpendicular to $\mathbf{r}$ in terms of F and θ .
- Thus, find the expression for the moment arm when the force and distance are not perpendicular.

Problem 2.19

Do not compute the cross products but just use the right-hand rule to find:

- $\hat{\mathbf{i}} \times \hat{\mathbf{j}}$
- $\hat{\mathbf{j}} \times \hat{\mathbf{k}}$
- $\hat{\mathbf{k}} \times \hat{\mathbf{i}}$
- $\hat{\mathbf{j}} \times \hat{\mathbf{i}}$
- $\hat{\mathbf{k}} \times \hat{\mathbf{j}}$
- $\hat{\mathbf{i}} \times \hat{\mathbf{k}}$

** Problem 2.20

Prove that the following determinant gives the area of a triangle with vertices at (x_1, y_1) , (x_2, y_2) and (x_3, y_3) :

$$Area = \left| \frac{1}{2} \det \begin{pmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{pmatrix} \right| \quad (2.5.7)$$

Therefore, if the determinant is 0, what does this imply about the three points?

Remark 2.4

While the linear algebra chapter in this book is ending, this is a very small subset of a standard linear algebra course. Many concepts, such as eigenvalues, are not covered due to their not so important nature. Readers are still highly encouraged to read up some of the other concepts.



2.6 An aside: Bra-ket and Hilbert space

TBD

2.7 An aside: Eigenfunctions and eigenvalues

TBD

Part II

Calculus I

Derivatives

3.1 Differentiation basics

Onto the calculus portion that is very much anticipated. No more the quick skimming over precalculus topics. We shall enter this discussion with a very simple question that will yield many interesting results. Take a look at the graph below which describes the displacement of particle x over time t ³¹:




Figure 3.1 Graph of $f(t) = t^3 - 2t + 1$.

³¹ x starts at 0, because if I were to plot the graph with negative values of x , it does not make sense physically. Unfortunately, this makes the cubic nature of the graph less obvious.

From Figure 3.1, we see that this is a cubic graph. It should not be hard to see that the graph is always changing value. This corresponds to the fact that the velocity of the particle is not constant. Also, it should not be difficult to see that the rate of change of the graph is not constant. This, meanwhile, shows that the acceleration of the particle is not constant. We can use the gradient formula to verify this:

Definition 3.1 Gradient

The gradient between two points $(x, f(x))$ and $(x + a, f(x + a))$ is:

$$\text{Gradient} = \frac{\Delta y}{\Delta x} = \frac{f(x + a) - f(x)}{a} \quad (3.1.1)$$



In case you have forgotten, the rate of change of displacement with respect to time is velocity, and the rate of change of velocity with respect to time is acceleration. Hence, we can use the gradient formula to find the average velocity of the particle over a certain time interval. For example, if we want to find the average velocity between $t = 1$ and $t = 2$, we can use Equation (3.1.1):

$$\begin{aligned} \text{Average velocity} &= \frac{f(2) - f(1)}{2 - 1} \\ &= \frac{(2^3 - 2 \cdot 2 + 1) - (1^3 - 2 \cdot 1 + 1)}{1} \\ &= \frac{5}{1} \\ &= 5 \end{aligned}$$

Using Equation (3.1.1) again, we can pick some points on the graph. You would notice that if you plug in different points then the gradient is different. The main, central question is: if the rate of change is not constant, then what is the rate of change at any single point along the function? This is where the gradient formula fails. Our current formula can only give us the average velocity for some time period, however, we want something that gives us the gradient of the graph at every single point along the graph. In physics, this is analogous to the velocity of the particle at every single point along its journey, or its *instantaneous velocity*. The forefathers of calculus were all brilliant and they noticed a special fact that applies to many functions. Let us zoom in to the graph where $t = 2$, we want to zoom in as closely as we can.




Figure 3.2 Zoomed in graph of Figure 3.1.

In Figure 3.2, we realize that the graph, when zoomed in sufficiently, essentially is a straight line. This is exciting, we can use our gradient formula which works on straight lines. In addition, on the x-axis, we mark $2 + h$, a small distance away from 2. We can infer that h is a really tiny number, almost going to 0 as we keep zooming in. We use the limit notation to denote this. Hence, we can denote it as $\lim_{h \rightarrow 0}$.³² You may be itching to use the gradient formula now. We can substitute in the values:

$$\lim_{h \rightarrow 0} \frac{f(2 + h) - f(2)}{h}$$

³² The limit notation and its concepts will be discussed in a later chapter, for now, just keep in mind that this means h is a really tiny number, almost 0.

However, notice that we have to divide throughout by h , and h is very small, infinitesimally small. Some careless manipulation and you would say the expression blows up to infinity. What the formula does tell us, though, is the gradient practically at 2. This gives us an idea. Can we just replace 2 with any value of x ? This brings us to the definition of the *derivative*.

Definition 3.2 Derivative

The *derivative* of a function $f(x)$ is the instantaneous rate of change of that function. It can be written as $f'(x)$, $\frac{df(x)}{dx}$ or $\dot{f}$.³³

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h} \quad (3.1.2)$$

In the definition of the derivative, we can see that the derivative is also a function. Also, the definition states that it is the “instantaneous rate of change”. This simply means that we are looking at the rate of change of the function over a very, very tiny range of x . Of course, the definition may feel contradictory. How can some value change “instantaneously”? Well, this is resolved by not exactly by substituting $\Delta y = 0$, instead we let it “approach” 0. As a result, using this clever little mathematical trick, we avoided this problem. However, some of you may have noticed we have not evaluated any derivatives, we just defined them. It is no good to not answer our original question. To illustrate a derivation of a derivative, we shall use a much simpler graph, $f(x) = x^2$.


³³ Sometimes, to save writing a few letters, the inputs to the functions may be omitted, such as f instead of $f(x)$. This is more common when the function takes in multiple inputs.

From Figure 3.3, we can consider what happens at $x = 2$. We shall use the definition of the derivatives, which is also more commonly referred to as “first principles”.

Example 3.1 Derivative of $f(x) = x^2$

To find the derivative at $x = 2$, we use the first principles as such³⁴:

$$\begin{aligned} f'(2) &= \lim_{h \rightarrow 0} \frac{f(2 + h) - f(2)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(2 + h)^2 - 4}{h} \\ &= \lim_{h \rightarrow 0} \frac{4 + 4h + h^2 - 4}{h} \\ &= \lim_{h \rightarrow 0} 4 + h \\ &= 4 \end{aligned}$$

Figure 3.3 Figure of $y = x^2$.

To find the derivative of f , we use a similar manipulation:

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - x^2}{h} \\ &= \lim_{h \rightarrow 0} 2x + h \\ &= 2x \end{aligned}$$

△

³⁴ We can ignore h in the second last step because as a general rule of thumb, as long as substituting $h = 0$ does not create an expression of $\frac{a}{0}$, we can let $h = 0$. Of course, this is very not rigorous and there are other exceptions that will be very evident after the reader has read the later chapter of limits.

In other words, we have also found the instantaneous rate of change of $f(x) = x^2$. We shall solve the problem we posed at the start of the chapter, what is the instantaneous rate of change or velocity for Figure 3.1.

Example 3.2 Derivative of $f(t) = t^3 - 2t + 1$

Using the first principles, we have:

$$\begin{aligned} f'(t) &= \lim_{h \rightarrow 0} \frac{f(t+h) - f(t)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(t+h)^3 - 2(t+h) + 1 - (t^3 - 2t + 1)}{h} \\ &= \lim_{h \rightarrow 0} \frac{t^3 + 3t^2h + 3th^2 + h^3 - 2t - 2h + 1 - t^3 + 2t - 1}{h} \\ &= \lim_{h \rightarrow 0} 3t^2 + 3th + h^2 - 2 \end{aligned}$$

$$= 3t^2 - 2$$

Therefore, the instantaneous velocity of the particle at any time t is $3t^2 - 2$. For example, at $t = 1$, the instantaneous velocity is simply 1, and at $t = 2$, the instantaneous velocity is 10. Remember how we found the average velocity between $t = 1$ and $t = 2$ to be 5? Notice how the instantaneous velocity at $t = 1$ is less than 5, and at $t = 2$ is greater than 5. This makes sense because the graph is constantly increasing in this interval, hence the average velocity must be between the two instantaneous velocities.



Lastly, I would like to comment that normally, if not explicitly stated in the question to use “first principles”, you can use the derivative rules you will learn in the next chapter to solve derivatives much more quickly and efficiently. In all honesty, using the “first principles” is very tedious and generally would not be used.

*** Problem 3.1**

Using first principles, find the derivatives of the following functions:

- a) $f(x) = x^2 + 1$
- b) $g(x) = x^3 + x$
- c) $h(x) = \frac{1}{x}$

*** Problem 3.2**

Let $f(x) = (x - 1)(x - 2) \cdots (x - 1729)$, find $f'(1729)$. (*Adapted from SMO 2025 Senior Section Round 1*).

*** Problem 3.3**

Given a differentiable function $f(x)$, find the equation of the tangent line at $x = a$.
