Logistics

GOV 2001 / 1002 / E-200 Section 9 Causal Inference and Estimation¹

Anton Strezhnev

Harvard University

March 23, 2016

¹These section notes are heavily indebted to past Gov 2001 TFs for slides and R code.

LOGISTICS

Reading Assignment- Ho et. al. (2007), King et. al. (2015), and Maertens and Swinnen (2015) **Problem Set 7-** Last Problem Set!

Due by 6pm Wednesday, March 13th on Canvas.

► In this section you will...

- ► In this section you will...
 - ► learn how to define a causal effect and what assumptions are needed to estimate it.

- ► In this section you will...
 - ► learn how to define a causal effect and what assumptions are needed to estimate it.
 - ▶ learn how to estimate a causal effect using data.

- ► In this section you will...
 - learn how to define a causal effect and what assumptions are needed to estimate it.
 - ▶ learn how to estimate a causal effect using data.
 - learn the benefits and drawbacks of different methods of estimating causal effects.

OUTLINE

Causal Inference

Identifying Causal Effects

Causal Effects in Observational Data

Matching

► A causal effect is a comparison of counterfactuals.

- ► A causal effect is a comparison of counterfactuals.
- One way of thinking about these counterfactuals is through the concept of "potential outcomes"

- ► A causal effect is a comparison of counterfactuals.
- One way of thinking about these counterfactuals is through the concept of "potential outcomes"
- ► For some person i, there exist two unobserved potential outcomes $Y_i(1)$ and $Y_i(0)$.

- ► A causal effect is a comparison of counterfactuals.
- ► One way of thinking about these counterfactuals is through the concept of "potential outcomes"
- For some person i, there exist two unobserved potential outcomes $Y_i(1)$ and $Y_i(0)$.
 - $Y_i(1)$ is the outcome if the person were to receive treatment T = 1.

- ► A causal effect is a comparison of counterfactuals.
- One way of thinking about these counterfactuals is through the concept of "potential outcomes"
- For some person i, there exist two unobserved potential outcomes $Y_i(1)$ and $Y_i(0)$.
 - $Y_i(1)$ is the outcome if the person were to receive treatment T = 1.
 - $Y_i(0)$ is the outcome if the person were to receive treatment T = 0.

- ► A causal effect is a comparison of counterfactuals.
- One way of thinking about these counterfactuals is through the concept of "potential outcomes"
- ► For some person i, there exist two unobserved potential outcomes $Y_i(1)$ and $Y_i(0)$.
 - $Y_i(1)$ is the outcome if the person were to receive treatment T = 1.
 - $Y_i(0)$ is the outcome if the person were to receive treatment T = 0.
- ▶ The causal effect τ_i of treatment for that individual is

$$\tau_i = Y_i(1) - Y_i(0)$$





"Everybody wanted to know what would happen if I didn't win...



"Everybody wanted to know what would happen if I didn't win...I guess we'll never know."

Logistics Overview Causal Inference Identifying Causal Effects Causal Effects in Observational Data Matching



"Everybody wanted to know what would happen if I didn't win...I guess we'll never know."

Kanye West, 2005

• We cannot estimate τ_i from data!

▶ We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.

- We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ▶ What do we do?

- ▶ We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ▶ What do we do? Change the question!

- We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ▶ What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

- ▶ We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ▶ What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

$$ATE = E[\tau_i] = E[Y_i(1)] - E[Y_i(0)]$$

- ▶ We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ▶ What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

$$ATE = E[\tau_i] = E[Y_i(1)] - E[Y_i(0)]$$

► **Important:** What causal question does the ATE answer?

- We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ▶ What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

$$ATE = E[\tau_i] = E[Y_i(1)] - E[Y_i(0)]$$

- ► **Important:** What causal question does the ATE answer?
 - ▶ Does it tell us what the effect will be if *you* get treatment?

- We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ▶ What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

$$ATE = E[\tau_i] = E[Y_i(1)] - E[Y_i(0)]$$

- ► **Important:** What causal question does the ATE answer?
 - Does it tell us what the effect will be if you get treatment? No!

- We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ▶ What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

$$ATE = E[\tau_i] = E[Y_i(1)] - E[Y_i(0)]$$

- ► **Important:** What causal question does the ATE answer?
 - Does it tell us what the effect will be if you get treatment? No!
 - ► It tells us the average of effects in the population of units.

- We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ▶ What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

$$ATE = E[\tau_i] = E[Y_i(1)] - E[Y_i(0)]$$

- ► **Important:** What causal question does the ATE answer?
 - Does it tell us what the effect will be if you get treatment? No!
 - ► It tells us the average of effects in the population of units.
 - ► This isn't always what we care about!

- ▶ We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ► What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

$$ATE = E[\tau_i] = E[Y_i(1)] - E[Y_i(0)]$$

- ► **Important:** What causal question does the ATE answer?
 - Does it tell us what the effect will be if you get treatment? No!
 - ► It tells us the average of effects in the population of units.
 - This isn't always what we care about! Sometimes the distribution matters.

- We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ► What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

$$ATE = E[\tau_i] = E[Y_i(1)] - E[Y_i(0)]$$

- ► **Important:** What causal question does the ATE answer?
 - Does it tell us what the effect will be if you get treatment? No!
 - ▶ It tells us the average of effects in the population of units.
 - This isn't always what we care about! Sometimes the distribution matters.
 - ► Example: Two medications have the same ATE,

- ▶ We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ► What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

$$ATE = E[\tau_i] = E[Y_i(1)] - E[Y_i(0)]$$

- ► **Important:** What causal question does the ATE answer?
 - Does it tell us what the effect will be if you get treatment? No!
 - ▶ It tells us the average of effects in the population of units.
 - This isn't always what we care about! Sometimes the distribution matters.
 - Example: Two medications have the same ATE, One has a constant small effect.

THE FUNDAMENTAL PROBLEM OF CAUSAL Inference

- We cannot estimate τ_i from data! For an individual, Y_i is only one of their potential outcomes.
- ▶ What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

$$ATE = E[\tau_i] = E[Y_i(1)] - E[Y_i(0)]$$

- ► **Important:** What causal question does the ATE answer?
 - ▶ Does it tell us what the effect will be if *you* get treatment? No!
 - ► It tells us the average of effects in the population of units.
 - ► This isn't always what we care about! Sometimes the distribution matters.
 - Example: Two medications have the same ATE, One has a constant small effect. The other has huge benefits but kills 1 in 1000 patients.

- We cannot estimate τ_i from data! For an individual, Y_i is only *one* of their potential outcomes.
- ▶ What do we do? Change the question!
- ► Most intuitive quantity: the Average Treatment Effect (ATE).

ATE =
$$E[\tau_i] = E[Y_i(1)] - E[Y_i(0)]$$

- ► **Important:** What causal question does the ATE answer?
 - ▶ Does it tell us what the effect will be if *you* get treatment? No!
 - ► It tells us the average of effects in the population of units.
 - ► This isn't always what we care about! Sometimes the distribution matters.
 - ► Example: Two medications have the same ATE, One has a constant small effect. The other has huge benefits but kills 1 in 1000 patients. Which would you prescribe?

CAUSAL VS. ASSOCIATIONAL CLAIMS

► What's the difference between

$$E[Y_i(1)] - E[Y_i(0)]$$

▶ and

$$E[Y_i|T_i = 1] - E[Y_i|T_i = 0]$$

?

CAUSAL VS. ASSOCIATIONAL CLAIMS

► What's the difference between

$$E[Y_i(1)] - E[Y_i(0)]$$

▶ and

Logistics

$$E[Y_i|T_i = 1] - E[Y_i|T_i = 0]$$

?

► First is a causal quantity (counterfactual statement). The second is an association (not counterfactual!)

Logistics Overview Causal Inference Identifying Causal Effects Causal Effects in Observational Data Matching

CAUSAL VS. ASSOCIATIONAL CLAIMS




Figure: Difference betweeen causal and association quantities (Hernan and Robins)

▶ Your difference-in-means is not a causal effect.

➤ Your difference-in-means is not a causal effect. Your regression coefficient is not a causal effect.

► Your difference-in-means is not a causal effect. Your regression coefficient is not a causal effect. Your post-matching linear model is not a causal effect.

➤ Your difference-in-means is not a causal effect. Your regression coefficient is not a causal effect. Your post-matching linear model is not a causal effect. Your inverse-probability of treatment weighted marginal structural model estimator is not a causal effect.

- ➤ Your difference-in-means is not a causal effect. Your regression coefficient is not a causal effect. Your post-matching linear model is not a causal effect. Your inverse-probability of treatment weighted marginal structural model estimator is not a causal effect.
- ...without additional assumptions.

Logistics

► There is *no such thing* as a causal effect derived purely from data alone.

quantities (like differences in means) to causal effects.

▶ We need *assumptions* to connect observed associational

- ▶ Basically: When does correlation actually imply causation?
- Þ
- ► This is often known as **identification** of a causal effect.

OUTLINE

Causal Inference

Identifying Causal Effects

Causal Effects in Observational Data

Matching

IDENTIFYING CAUSAL EFFECTS

Logistics

► How do we go from something we can estimate

$$E[Y_i|T_i=1] - E[Y_i|T_i=0]$$

IDENTIFYING CAUSAL EFFECTS

► How do we go from something we can estimate

$$E[Y_i|T_i = 1] - E[Y_i|T_i = 0]$$

to something we don't observe

$$E[Y_i(1)] - E[Y_i(0)]$$

► Assumption 1: Stable Unit Treatment Value Assumption (SUTVA)

- ► Assumption 1: Stable Unit Treatment Value Assumption (SUTVA)
- ▶ If a subject i receives treatment $T_i = t$, then the observed outcome Y_i is equal to the potential outcome $Y_i(t)$.

- ► Assumption 1: Stable Unit Treatment Value Assumption (SUTVA)
- ▶ If a subject i receives treatment $T_i = t$, then the observed outcome Y_i is equal to the potential outcome $Y_i(t)$.
- ► Seems trivial, but not...

- ► Assumption 1: Stable Unit Treatment Value Assumption (SUTVA)
- ▶ If a subject i receives treatment $T_i = t$, then the observed outcome Y_i is equal to the potential outcome $Y_i(t)$.
- Seems trivial, but not...
 - ► Implies that there aren't multiple versions of treatment.

- ► Assumption 1: Stable Unit Treatment Value Assumption (SUTVA)
- ▶ If a subject i receives treatment $T_i = t$, then the observed outcome Y_i is equal to the potential outcome $Y_i(t)$.
- Seems trivial, but not...
 - ► Implies that there aren't multiple versions of treatment.
 - Also implies no interference between units the potential outcome for an individual does not depend on treatments assigned to *other* individuals.

- ► Assumption 1: Stable Unit Treatment Value Assumption (SUTVA)
- ▶ If a subject i receives treatment $T_i = t$, then the observed outcome Y_i is equal to the potential outcome $Y_i(t)$.
- Seems trivial, but not...
 - ► Implies that there aren't multiple versions of treatment.
 - Also implies no interference between units the potential outcome for an individual does not depend on treatments assigned to *other* individuals.
- ► *Implication:* We can reduce the number of potential outcomes for each unit to two that depend only on the treatment it receives: $Y_i(1)$ and $Y_i(0)$.

► Assumption 2: *Ignorability/Unconfoundedness*

- ► Assumption 2: *Ignorability/Unconfoundedness*
- ▶ $\{Y_i(1), Y_i(0)\}$ is independent of T_i

- ► Assumption 2: *Ignorability/Unconfoundedness*
- $\{Y_i(1), Y_i(0)\}$ is independent of T_i
- ► This is the big assumption...
 - ► Implies that had the group that received treatment been instead assigned control

- ► Assumption 2: *Ignorability/Unconfoundedness*
- $\{Y_i(1), Y_i(0)\}$ is independent of T_i
- ► This is the big assumption...
 - ► Implies that had the group that received treatment been instead assigned control, the observed outcome in that group would have been the same as the outcome in the group that *did* receive control (and vice-versa).
 - ▶ Violated if some third factor affects probability of receiving treatment and $\{Y_i(1), Y_i(0)\}$

- ► Assumption 2: *Ignorability/Unconfoundedness*
- $\{Y_i(1), Y_i(0)\}$ is independent of T_i
- ► This is the big assumption...
 - ► Implies that had the group that received treatment been instead assigned control, the observed outcome in that group would have been the same as the outcome in the group that *did* receive control (and vice-versa).
 - ▶ Violated if some third factor affects probability of receiving treatment and $\{Y_i(1), Y_i(0)\}$
- ► Randomization is valued because it *gives us* ignorability.

► Under these assumptions, the causal effect **is identified** by the observed association between treatment/control.

- ▶ Under these assumptions, the causal effect **is identified** by the observed association between treatment/control.
- ► By SUTVA

$$E[Y_i|T_i = 1] - E[Y_i|T_i = 0] = E[Y_i(1)|T_i = 1] - E[Y_i(0)|T_i = 0]$$

- ▶ Under these assumptions, the causal effect **is identified** by the observed association between treatment/control.
- ► By SUTVA

$$E[Y_i|T_i=1] - E[Y_i|T_i=0] = E[Y_i(1)|T_i=1] - E[Y_i(0)|T_i=0]$$

► By ignorability

$$E[Y_i(1)|T_i=1] - E[Y_i(0)|T_i=0] = E[Y_i(1)] - E[Y_i(0)]$$

OUTLINE

Causal Inference

Identifying Causal Effect

Causal Effects in Observational Data

Matching

 In observational data, unconditional ignorability rarely holds.

- ► In observational data, unconditional ignorability rarely holds.
 - ► Sicker people are more likely to take medicine.

- In observational data, unconditional ignorability rarely holds.
 - Sicker people are more likely to take medicine.
 - ► Wealthier people are more likely to turn out and vote (in the U.S.)

- In observational data, unconditional ignorability rarely holds.
 - ► Sicker people are more likely to take medicine.
 - Wealthier people are more likely to turn out and vote (in the U.S.)
 - ▶ Paris of democratic countries are more likely to trade.

- In observational data, unconditional ignorability rarely holds.
 - ► Sicker people are more likely to take medicine.
 - Wealthier people are more likely to turn out and vote (in the U.S.)
 - ▶ Paris of democratic countries are more likely to trade.
- ▶ What to do?

- In observational data, unconditional ignorability rarely holds.
 - ► Sicker people are more likely to take medicine.
 - Wealthier people are more likely to turn out and vote (in the U.S.)
 - ▶ Paris of democratic countries are more likely to trade.
- What to do? Make a weaker assumption conditional ignorability.

► Assumption: Conditional Ignorability

- ► Assumption: Conditional Ignorability
- ▶ $\{Y_i(1), Y_i(0)\}$ are independent of T_i conditional on some set of covariates **X**

- ► Assumption: Conditional Ignorability
- $\{Y_i(1), Y_i(0)\}$ are independent of T_i conditional on some set of covariates **X**
 - ► Often referred to as "no unobserved confounding"

- ► Assumption: Conditional Ignorability
- $\{Y_i(1), Y_i(0)\}$ are independent of T_i conditional on some set of covariates **X**
 - ▶ Often referred to as "no unobserved confounding"
 - "Theory-driven" assumption no way of verifying from data alone that it holds!

IDENTIFICATION IN OBSERVATIONAL DATA

- Assumption: Conditional Ignorability
- $\{Y_i(1), Y_i(0)\}$ are independent of T_i conditional on some set of covariates **X**
 - ▶ Often referred to as "no unobserved confounding"
 - ► "Theory-driven" assumption no way of verifying from data alone that it holds!

Under SUTVA and conditional ignorability,

$$E[Y_i|T_i = 1, X_i = x] - E[Y_i|T_i = 0, X_i = x]$$

= $E[Y_i(1)|X_i = x] - E[Y_i(0)|X_i = x]$

 \blacktriangleright When *X* is low-dimensional, can *stratify* by *X*,

Under SUTVA and conditional ignorability,

$$E[Y_i|T_i = 1, X_i = x] - E[Y_i|T_i = 0, X_i = x]$$

= $E[Y_i(1)|X_i = x] - E[Y_i(0)|X_i = x]$

▶ When *X* is low-dimensional, can *stratify* by *X*, take differences in means within strata,

► Under SUTVA and conditional ignorability,

$$E[Y_i|T_i = 1, X_i = x] - E[Y_i|T_i = 0, X_i = x]$$

= $E[Y_i(1)|X_i = x] - E[Y_i(0)|X_i = x]$

▶ When *X* is low-dimensional, can *stratify* by *X*, take differences in means within strata, and then average those estimates to estimate ATE.

ESTIMATION IN OBSERVATIONAL DATA

Under SUTVA and conditional ignorability,

$$E[Y_i|T_i = 1, X_i = x] - E[Y_i|T_i = 0, X_i = x]$$

= $E[Y_i(1)|X_i = x] - E[Y_i(0)|X_i = x]$

▶ When *X* is low-dimensional, can *stratify* by *X*, take differences in means within strata, and then average those estimates to estimate ATE. No additional modeling assumptions necessary!

▶ When *X* is high-dimensional or continuous, what happens to the stratification estimator?

► When *X* is high-dimensional or continuous, what happens to the stratification estimator? Too high variance!

- ▶ When *X* is high-dimensional or continuous, what happens to the stratification estimator? Too high variance!
- ► Some strata might only have a single observation!
- ► First approach: Make more modeling assumptions.

Logistics

- ▶ When *X* is high-dimensional or continuous, what happens to the stratification estimator? Too high variance!
- ► Some strata might only have a single observation!
- ► First approach: Make more modeling assumptions. Regression!
- Assume not only conditional ignorability, but also a model for E[Y|T,X].

$$E[Y_i|T_i,X_i] = \beta_0 + \beta_1T_i + \beta_2X_{1i} + \beta_3X_{2i}, \dots$$

► Now we can estimate using OLS...

► Under these assumptions, what's the estimate of causal effect?

ESTIMATION IN OBSERVATIONAL DATA

Under these assumptions, what's the estimate of causal effect?

$$E[Y_i|T_i = 1, X_i = x] - E[Y_i|T_i = 0, X_i = x]$$

$$= \beta_0 + \beta_1 \times 1 + \beta_2 x_1 + \beta_3 x_2 - \beta_0 + \beta_1 \times 0 + \beta_2 x_1 + \beta_3 x_2,$$

$$= \beta_1$$

▶ Note we've assumed a constant effect!

► Under these assumptions, what's the estimate of causal effect?

$$E[Y_i|T_i = 1, X_i = x] - E[Y_i|T_i = 0, X_i = x]$$

$$= \beta_0 + \beta_1 \times 1 + \beta_2 x_1 + \beta_3 x_2 - \beta_0 + \beta_1 \times 0 + \beta_2 x_1 + \beta_3 x_2,$$

$$= \beta_1$$

Note we've assumed a constant effect! If that doesn't hold, β_1 is still an average of individual treatment effects...

► Under these assumptions, what's the estimate of causal effect?

$$E[Y_i|T_i = 1, X_i = x] - E[Y_i|T_i = 0, X_i = x]$$

$$= \beta_0 + \beta_1 \times 1 + \beta_2 x_1 + \beta_3 x_2 - \beta_0 + \beta_1 \times 0 + \beta_2 x_1 + \beta_3 x_2,$$

$$= \beta_1$$

▶ Note we've assumed a constant effect! If that doesn't hold, β_1 is still an average of individual treatment effects... but it's a weighted average and may not be representative of the ATE (see Aronow and Samii (2015))

▶ Under these assumptions, what's the estimate of causal effect?

Identifying Causal Effects

$$E[Y_i|T_i = 1, X_i = x] - E[Y_i|T_i = 0, X_i = x]$$

$$= \beta_0 + \beta_1 \times 1 + \beta_2 x_1 + \beta_3 x_2 - \beta_0 + \beta_1 \times 0 + \beta_2 x_1 + \beta_3 x_2,$$

$$= \beta_1$$

- ▶ Note we've assumed a constant effect! If that doesn't hold, β_1 is still an average of individual treatment effects... but it's a weighted average and may not be representative of the ATE (see Aronow and Samii (2015))
- ▶ If we don't like OLS, can use GLMs, GAMs, or whatever to get conditional expectations.

IMATION IN OBSERVATIONAL DATA

► Under these assumptions, what's the estimate of causal effect?

$$E[Y_i|T_i = 1, X_i = x] - E[Y_i|T_i = 0, X_i = x]$$

$$= \beta_0 + \beta_1 \times 1 + \beta_2 x_1 + \beta_3 x_2 - \beta_0 + \beta_1 \times 0 + \beta_2 x_1 + \beta_3 x_2,$$

$$= \beta_1$$

- Note we've assumed a constant effect! If that doesn't hold, β_1 is still an average of individual treatment effects... but it's a weighted average and may not be representative of the ATE (see Aronow and Samii (2015))
- ► If we don't like OLS, can use GLMs, GAMs, or whatever to get conditional expectations. How we estimate the CEF is separate from whether it's causally interpretable (identification).

► What's the right regression then?

ESTIMATION IN OBSERVATIONAL DATA

► What's the right regression then?

► What's the right regression then?

► If we pick the wrong model, we introduce bias in *estimation*.

ESTIMATION IN OBSERVATIONAL DATA

► What's the right regression then?

▶ If we pick the wrong model, we introduce bias in *estimation*. How do we avoid *model dependence*?

OUTLINE

Logistics

Causal Inference

Identifying Causal Effects

Causal Effects in Observational Data

Matching

MATCHING ESTIMATORS

► How do we estimate $E[Y_i(1)]$ and $E[Y_i(0)]$ when we have lots of covariates but don't want to assume a model for the outcome?

MATCHING ESTIMATORS

- ► How do we estimate $E[Y_i(1)]$ and $E[Y_i(0)]$ when we have lots of covariates but don't want to assume a model for the outcome?
- ▶ For each observation, we observe either $Y_i(1)$ or $Y_i(0)$.

- ► How do we estimate $E[Y_i(1)]$ and $E[Y_i(0)]$ when we have lots of covariates but don't want to assume a model for the outcome?
- ▶ For each observation, we observe either $Y_i(1)$ or $Y_i(0)$.
- ▶ If we observed everything, then our estimated ATE $\hat{\tau}$ would be

$$\hat{\tau} = \frac{1}{N} \sum_{i=1}^{N} Y_i(1) - \frac{1}{N} \sum_{i=1}^{N} Y_i(0)$$

MATCHING ESTIMATORS

• We could instead imagine coming up with estimates $Y_i(1)$ such that

► We could instead imagine coming up with estimates $\hat{Y}_i(1)$ such that

$$Y_{\hat{i}}(1) = \begin{cases} Y_i & \text{if } T_i = 1\\ Y_j^* & \text{if } T_i = 0 \end{cases}$$

▶ What's Y_j^* ?

► We could instead imagine coming up with estimates $\hat{Y_i(1)}$ such that

$$Y_{i}(1) = \begin{cases} Y_{i} & \text{if } T_{i} = 1\\ Y_{j}^{*} & \text{if } T_{i} = 0 \end{cases}$$

▶ What's Y_j^* ? It's the value of Y from some other observation j that has treatment $T_i = 0$ and covariates X_j that are "close" to covariates X_i .

• We could instead imagine coming up with estimates $Y_i(1)$ such that

$$Y_{i}(1) = \begin{cases} Y_{i} & \text{if } T_{i} = 1\\ Y_{j}^{*} & \text{if } T_{i} = 0 \end{cases}$$

- ▶ What's Y_i^* ? It's the value of Y from some other observation *j* that has treatment $T_i = 0$ and covariates X_i that are "close" to covariates X_i .
- ▶ That is, it's the outcome from a "matched" control observation (or observations).

MATCHING ESTIMATORS

► We could instead imagine coming up with estimates $\hat{Y_i(1)}$ such that

$$Y_i(1) = \begin{cases} Y_i & \text{if } T_i = 1\\ Y_j^* & \text{if } T_i = 0 \end{cases}$$

- ▶ What's Y_j^* ? It's the value of Y from some other observation j that has treatment $T_i = 0$ and covariates X_j that are "close" to covariates X_i .
- ► That is, it's the outcome from a "matched" control observation (or observations).
- ► We could do the same for $Y_i(0)$ and use a matching estimator for the ATE

$$\hat{\tau} = \frac{1}{N} \sum_{i=1}^{N} Y_i(1) - \frac{1}{N} \sum_{i=1}^{N} Y_i(0)$$

Matching is not an identification strategy.

- ► Matching is not an identification strategy.
- ► Matching is not an identification strategy.

- ► Matching is not an identification strategy.
- ► Matching is not an identification strategy.
- ► Matching is not an identification strategy.

- ► Matching is not an identification strategy.
- ► Matching is not an identification strategy.
- Matching is not an identification strategy.
- ► It does **not** eliminate bias from unobserved confounding.

- ► Matching is not an identification strategy.
- Matching is not an identification strategy.
- Matching is not an identification strategy.
- ► It does **not** eliminate bias from unobserved confounding. If the identification assumptions do not hold, then matching won't magically help.

- ► Matching is not an identification strategy.
- ► Matching is not an identification strategy.
- Matching is not an identification strategy.
- ► It does **not** eliminate bias from unobserved confounding. If the identification assumptions do not hold, then matching won't magically help.
- ► However, it *does* ameliorate bias from arbitrary choices of models for E[Y|T, X].

MATCHING IS AN ESTIMATION STRATEGY

- ► Matching is not an identification strategy.
- Matching is not an identification strategy.
- Matching is not an identification strategy.
- ► It does **not** eliminate bias from unobserved confounding. If the identification assumptions do not hold, then matching won't magically help.
- ► However, it *does* ameliorate bias from arbitrary choices of models for E[Y|T,X].
- ► And gives us a way of assessing how "well" we've adjusted for the observed confounders imbalance.

▶ Ideal case: **Exact Matching**. For each *i* find a matched observation *j* with $X_j = X_i$ but different T_i .

▶ Ideal case: **Exact Matching**. For each i find a matched observation j with $X_j = X_i$ but different T_i . Basically impossible in any normal situation.

- ▶ Ideal case: **Exact Matching**. For each i find a matched observation j with $X_j = X_i$ but different T_i . Basically impossible in any normal situation.
- ▶ Need to allow for some differences between X_j and X_i .

- ▶ Ideal case: **Exact Matching**. For each i find a matched observation j with $X_j = X_i$ but different T_i . Basically impossible in any normal situation.
- ▶ Need to allow for some differences between X_j and X_i . One approach is to define a distance metric and match the closest j ("Nearest-neighbor").

- ▶ Ideal case: **Exact Matching**. For each i find a matched observation j with $X_j = X_i$ but different T_i . Basically impossible in any normal situation.
- Need to allow for some differences between X_j and X_i. One approach is to define a distance metric and match the closest j ("Nearest-neighbor").
 - ► Mahalanobis Distance:

$$M(X_i, X_j) = [(X_i - X_j)^T S^{-1} (X_i - X_j)]^{-1/2}$$

- ► Ideal case: **Exact Matching**. For each *i* find a matched observation j with $X_i = X_i$ but different T_i . Basically impossible in any normal situation.
- ▶ Need to allow for some differences between X_i and X_i . One approach is to define a distance metric and match the closest *j* ("Nearest-neighbor").
 - Mahalanobis Distance:

$$M(X_i, X_j) = [(X_i - X_j)^T S^{-1} (X_i - X_j)]^{-1/2}$$

Propensity Score Difference:

$$\hat{\pi}_i - \hat{\pi}_i = Pr(T_i = 1|X_i) - Pr(T_i = 1|X_i)$$

Matching

- ▶ Ideal case: **Exact Matching**. For each i find a matched observation j with $X_j = X_i$ but different T_i . Basically impossible in any normal situation.
- Need to allow for some differences between X_j and X_i. One approach is to define a distance metric and match the closest j ("Nearest-neighbor").
 - ► Mahalanobis Distance:

$$M(X_i, X_j) = [(X_i - X_j)^T S^{-1} (X_i - X_j)]^{-1/2}$$

► Propensity Score Difference:

$$\hat{\pi}_i - \hat{\pi}_j = Pr(T_i = 1|X_i) - Pr(T_i = 1|X_j)$$

► Coarsened Exact Matching:

$$D(X_i, X_j) = \begin{cases} 0 & \text{if } X_i \approx X_j \\ \infty & \text{if } X_i \not\approx X_j \end{cases}$$

Goal of matching is to minimize imbalance (and thereby bias).

► Goal of matching is to minimize imbalance (and thereby bias). Often we look at differences in means across controls variables.


Logistics

► Goal of matching is to minimize imbalance (and thereby bias). Often we look at differences in means across controls variables.

Alester to the block of the	Control	Civic Duty Mean	Hawthorne Mean	Self Mean	Neighbors Mean
Household size	1.91	1.91	1.91	1.91	1.91
Nov 2002	.83	.84	.84	.84	.84
Nov 2000	.87	.87	.87	.86	.87
Aug 2004	.42	.42	.42	.42	.42
Aug 2002	.41	.41	.41	.41	.41
Aug 2000	.26	.27	.26	.26	.26
Female	.50	.50	.50	.50	.50
Age (in years)	51.98	51.85	51.87	51.91	52.01
N =	99,999	20,001	20,002	20,000	20,000

Figure: Balance table from actual experiment by Gerber, Green and Larimer (2008)

► However, means can be misleading:



► Also, don't ignore variance!

► Also, don't ignore variance! We can get perfect balance...

► Also, don't ignore variance! We can get perfect balance...by dropping all observations except for one identically matched pair.

▶ Also, don't ignore variance! We can get perfect balance...by dropping all observations except for one identically matched pair. Who wants to publish with n = 2?

- ▶ Also, don't ignore variance! We can get perfect balance...by dropping all observations except for one identically matched pair. Who wants to publish with n = 2?
- ► One advantage of propensity score matching is that it has an implicit variable selection method so we're not matching on irrelevant covariates (that are not predictive of treatment).

- ► Also, don't ignore variance! We can get perfect balance...by dropping all observations except for one identically matched pair. Who wants to publish with n = 2?
- ► One advantage of propensity score matching is that it has an implicit variable selection method so we're not matching on irrelevant covariates (that are not predictive of treatment).
- ► However, it still relies on untestable modelling assumptions about the propensity score.

- ▶ Also, don't ignore variance! We can get perfect balance...by dropping all observations except for one identically matched pair. Who wants to publish with n = 2?
- ► One advantage of propensity score matching is that it has an implicit variable selection method so we're not matching on irrelevant covariates (that are not predictive of treatment).
- ► However, it still relies on untestable modelling assumptions about the propensity score.
- ► And only approximates complete randomization rather than blocking (and so suffers in efficiency/bias reduction).

Logistics

- ▶ Also, don't ignore variance! We can get perfect balance...by dropping all observations except for one identically matched pair. Who wants to publish with n = 2?
- ► One advantage of propensity score matching is that it has an implicit variable selection method so we're not matching on irrelevant covariates (that are not predictive of treatment).
- ► However, it still relies on untestable modelling assumptions about the propensity score.
- ► And only approximates complete randomization rather than blocking (and so suffers in efficiency/bias reduction).
- ► Open research question to find best methods of variable selection in matching without relying on models like propensity score.

Matching

► When do we match versus throw away observations?

► When do we match versus throw away observations? How bad of a match are we willing to tolerate before we change the quantity of interest?

- ► When do we match versus throw away observations? How bad of a match are we willing to tolerate before we change the quantity of interest?
- ► How many matches do we match to each observation?

- ► When do we match versus throw away observations? How bad of a match are we willing to tolerate before we change the quantity of interest?
- ► How many matches do we match to each observation?

 Matching more increases bias (because we're willing to tolerate greater distance) but reduces variance (more data).
- ▶ Do we match with or without replacement?

- ► When do we match versus throw away observations? How bad of a match are we willing to tolerate before we change the quantity of interest?
- ► How many matches do we match to each observation?

 Matching more increases bias (because we're willing to tolerate greater distance) but reduces variance (more data).
- ➤ Do we match with or without replacement? Matching with replacement reduces bias (ensures that closest observations always get matched).

- ► When do we match versus throw away observations? How bad of a match are we willing to tolerate before we change the quantity of interest?
- ► How many matches do we match to each observation?

 Matching more increases bias (because we're willing to tolerate greater distance) but reduces variance (more data).
- ➤ Do we match with or without replacement? Matching with replacement reduces bias (ensures that closest observations always get matched). However, it increases variance (we're reusing observations in the matched data, reducing effective sample size).

► Which observations are you going to find matches for and which ones will you drop?


- ► Which observations are you going to find matches for and which ones will you drop?
- ► Dropping observations changes the quantity of interest.

- ► Which observations are you going to find matches for and which ones will you drop?
- ► Dropping observations changes the quantity of interest.
 - ▶ Often, we have few treateds and lots of controls.

- Which observations are you going to find matches for and which ones will you drop?
- ▶ Dropping observations changes the quantity of interest.
 - ▶ Often, we have few treateds and lots of controls.
 - Matching every control would lead to a lot of bias, so we narrow QOI to "ATT" - Average Treatment Effect on the Treated.

- Which observations are you going to find matches for and which ones will you drop?
- ▶ Dropping observations changes the quantity of interest.
 - ▶ Often, we have few treateds and lots of controls.
 - Matching every control would lead to a lot of bias, so we narrow QOI to "ATT" - Average Treatment Effect on the Treated.

$$ATT = E[Y_i(1)|A_i = 1] - E[Y_i(0)|A_i = 1]$$



► But what if finding matches for some treateds is too unreasonable.

- ► But what if finding matches for some treateds is too unreasonable.
- ► Can narrow estimand even further by dropping treateds...


- ▶ But what if finding matches for some treateds is too unreasonable.
- Can narrow estimand even further by dropping treateds... to the Feasible Sample Average Treatment Effect on the Treated (FSATT)!
- Careful this improves internal validity (reduces bias) but might limit external validity (by changing the population of interest).
- ► How do we drop?

- ▶ But what if finding matches for some treateds is too unreasonable.
- Can narrow estimand even further by dropping treateds... to the Feasible Sample Average Treatment Effect on the Treated (FSATT)!
- Careful this improves internal validity (reduces bias) but might limit external validity (by changing the population of interest).
- ► How do we drop? Common support threshold.

MORE ON DROPPING OBS AND CHANGING QOI

- ▶ But what if finding matches for some treateds is too unreasonable.
- Can narrow estimand even further by dropping treateds... to the Feasible Sample Average Treatment Effect on the Treated (FSATT)!
- Careful this improves internal validity (reduces bias) but might limit external validity (by changing the population of interest).
- ► How do we drop? Common support threshold. Caliper/maximum match distance.

MORE ON DROPPING OBS AND CHANGING QOI



► Causal effects are counterfactual comparisons - need assumptions to connect them to quantities we can estimate.

- Causal effects are counterfactual comparisons need assumptions to connect them to quantities we can estimate.
- ► Causal inference is composed of two sets of assumptions:

- Causal effects are counterfactual comparisons need assumptions to connect them to quantities we can estimate.
- ► Causal inference is composed of two sets of assumptions:
 - ► Identification: What needs to be true in order for differences in conditional expectations to be interpretable as causal quantities?

- ► Causal effects are counterfactual comparisons need assumptions to connect them to quantities we can estimate.
- ► Causal inference is composed of two sets of assumptions:
 - ► Identification: What needs to be true in order for differences in conditional expectations to be interpretable as causal quantities?
 - ► Estimation: How do we estimate the conditional expectation?

- ► Causal effects are counterfactual comparisons need assumptions to connect them to quantities we can estimate.
- ► Causal inference is composed of two sets of assumptions:
 - ► Identification: What needs to be true in order for differences in conditional expectations to be interpretable as causal quantities?
 - ► Estimation: How do we estimate the conditional expectation?
- ► Matching is a tool for **estimation** not **identification**.

- ► Causal effects are counterfactual comparisons need assumptions to connect them to quantities we can estimate.
- ► Causal inference is composed of two sets of assumptions:
 - ► Identification: What needs to be true in order for differences in conditional expectations to be interpretable as causal quantities?
 - ► Estimation: How do we estimate the conditional expectation?
- ► Matching is a tool for **estimation** not **identification**.

- Causal effects are counterfactual comparisons need assumptions to connect them to quantities we can estimate.
- ► Causal inference is composed of two sets of assumptions:
 - ► Identification: What needs to be true in order for differences in conditional expectations to be interpretable as causal quantities?
 - ► Estimation: How do we estimate the conditional expectation?
- ► Matching is a tool for **estimation** not **identification**.Relies on the same "no unobserved confounders" assumption as a regression estimator.

- Causal effects are counterfactual comparisons need assumptions to connect them to quantities we can estimate.
- ► Causal inference is composed of two sets of assumptions:
 - ► Identification: What needs to be true in order for differences in conditional expectations to be interpretable as causal quantities?
 - ► Estimation: How do we estimate the conditional expectation?
- ► Matching is a tool for **estimation** not **identification**.Relies on the same "no unobserved confounders" assumption as a regression estimator.
- ► However, it allows us to directly assess the amount of estimation bias we introduce when we adjust for observed confounders (in contrast to regression).
- ▶ But if you have strong unobserved confounding, no amount of matching will save you.

- Causal effects are counterfactual comparisons need assumptions to connect them to quantities we can estimate.
- ► Causal inference is composed of two sets of assumptions:
 - ► Identification: What needs to be true in order for differences in conditional expectations to be interpretable as causal quantities?
 - ► Estimation: How do we estimate the conditional expectation?
- ► Matching is a tool for **estimation** not **identification**.Relies on the same "no unobserved confounders" assumption as a regression estimator.
- ► However, it allows us to directly assess the amount of estimation bias we introduce when we adjust for observed confounders (in contrast to regression).
- ► But if you have strong unobserved confounding, no amount of matching will save you. Change your research design!

QUESTIONS

Logistics

Questions?