

# Advanced Quantitative Research Methodology, Lecture Notes: Research Designs for Causal Inference<sup>1</sup>

Gary King

GaryKing.org

April 10, 2016

---

<sup>1</sup>©Copyright 2016 Gary King, All Rights Reserved.

- Kosuke Imai, Gary King, and Elizabeth Stuart. **Misunderstandings among Experimentalists and Observationalists: Balance Test Fallacies in Causal Inference** *Journal of the Royal Statistical Society, Series A* Vol. 171, Part 2 (2008): Pp. 1-22  
<http://gking.harvard.edu/files/abs/matchse-abs.shtml>

# Notation

# Notation

- **Sample** of  $n$  units from a finite population of  $N$  units (typically  $N \gg n$ )

# Notation

- **Sample** of  $n$  units from a finite population of  $N$  units (typically  $N \gg n$ )
- **Sample selection:**  $I_i$  is 1 for units selected, 0 otherwise

# Notation

- **Sample** of  $n$  units from a finite population of  $N$  units (typically  $N \gg n$ )
- **Sample selection:**  $I_i$  is 1 for units selected, 0 otherwise
- **Treatment assignment:**  $T_i$  is 1 for treated group, 0 control group

# Notation

- **Sample** of  $n$  units from a finite population of  $N$  units (typically  $N \gg n$ )
- **Sample selection:**  $I_i$  is 1 for units selected, 0 otherwise
- **Treatment assignment:**  $T_i$  is 1 for treated group, 0 control group
- (**Assume:** treated and control groups are each of size  $n/2$ )

# Notation

- **Sample** of  $n$  units from a finite population of  $N$  units (typically  $N \gg n$ )
- **Sample selection:**  $I_i$  is 1 for units selected, 0 otherwise
- **Treatment assignment:**  $T_i$  is 1 for treated group, 0 control group
- (**Assume:** treated and control groups are each of size  $n/2$ )
- **Observed outcome variable:**  $Y_i$



# Notation

- **Sample** of  $n$  units from a finite population of  $N$  units (typically  $N \gg n$ )
- **Sample selection:**  $I_i$  is 1 for units selected, 0 otherwise
- **Treatment assignment:**  $T_i$  is 1 for treated group, 0 control group
- (**Assume:** treated and control groups are each of size  $n/2$ )
- **Observed outcome variable:**  $Y_i$
- **Potential outcomes:**  $Y_i(1)$  and  $Y_i(0)$ ,  $Y_i$  potential values when  $T_i$  is 1 or 0 respectively.

- **Sample** of  $n$  units from a finite population of  $N$  units (typically  $N \gg n$ )
- **Sample selection:**  $I_i$  is 1 for units selected, 0 otherwise
- **Treatment assignment:**  $T_i$  is 1 for treated group, 0 control group
- (**Assume:** treated and control groups are each of size  $n/2$ )
- **Observed outcome variable:**  $Y_i$
- **Potential outcomes:**  $Y_i(1)$  and  $Y_i(0)$ ,  $Y_i$  potential values when  $T_i$  is 1 or 0 respectively.
- **Fundamental problem of causal inference.** Only one potential outcome is ever observed:  
If  $T_i = 0$ ,  $Y_i(0) = Y_i$      $Y_i(1) = ?$   
If  $T_i = 1$ ,  $Y_i(0) = ?$      $Y_i(1) = Y_i$

- **Sample** of  $n$  units from a finite population of  $N$  units (typically  $N \gg n$ )
- **Sample selection:**  $I_i$  is 1 for units selected, 0 otherwise
- **Treatment assignment:**  $T_i$  is 1 for treated group, 0 control group
- (**Assume:** treated and control groups are each of size  $n/2$ )
- **Observed outcome variable:**  $Y_i$
- **Potential outcomes:**  $Y_i(1)$  and  $Y_i(0)$ ,  $Y_i$  potential values when  $T_i$  is 1 or 0 respectively.
- **Fundamental problem of causal inference.** Only one potential outcome is ever observed:  
If  $T_i = 0$ ,  $Y_i(0) = Y_i$      $Y_i(1) = ?$   
If  $T_i = 1$ ,  $Y_i(0) = ?$      $Y_i(1) = Y_i$
- $(I_i, T_i, Y_i)$  are random;  $Y_i(1)$  and  $Y_i(0)$  are fixed.

# Quantities of Interest

# Quantities of Interest

- Treatment Effect (for unit  $i$ ):

$$TE_i \equiv Y_i(1) - Y_i(0)$$

# Quantities of Interest

- Treatment Effect (for unit  $i$ ):

$$\text{TE}_i \equiv Y_i(1) - Y_i(0)$$

- Population Average Treatment Effect:

$$\text{PATE} \equiv \frac{1}{N} \sum_{i=1}^N \text{TE}_i$$

# Quantities of Interest

- Treatment Effect (for unit  $i$ ):

$$TE_i \equiv Y_i(1) - Y_i(0)$$

- Population Average Treatment Effect:

$$PATE \equiv \frac{1}{N} \sum_{i=1}^N TE_i$$

- Sample Average Treatment Effect:

$$SATE \equiv \frac{1}{n} \sum_{i \in \{I_i=1\}} TE_i$$

# Decomposition of Causal Effect Estimation Error



# Decomposition of Causal Effect Estimation Error

- Difference in means estimator:

$$D \equiv \left( \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=1\}} Y_i \right) - \left( \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=0\}} Y_i \right).$$

# Decomposition of Causal Effect Estimation Error

- Difference in means estimator:

$$D \equiv \left( \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=1\}} Y_i \right) - \left( \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=0\}} Y_i \right).$$

- Estimation Error:

$$\Delta \equiv \text{PATE} - D$$

# Decomposition of Causal Effect Estimation Error

- Difference in means estimator:

$$D \equiv \left( \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=1\}} Y_i \right) - \left( \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=0\}} Y_i \right).$$

- Estimation Error:

$$\Delta \equiv \text{PATE} - D$$

- Pretreatment confounders:  $X$  are observed and  $U$  are unobserved

# Decomposition of Causal Effect Estimation Error

- Difference in means estimator:

$$D \equiv \left( \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=1\}} Y_i \right) - \left( \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=0\}} Y_i \right).$$

- Estimation Error:

$$\Delta \equiv \text{PATE} - D$$

- Pretreatment confounders:  $X$  are observed and  $U$  are unobserved
- Decomposition:

$$\begin{aligned} \Delta &= \Delta_S + \Delta_T \\ &= (\Delta_{S_X} + \Delta_{S_U}) + (\Delta_{T_X} + \Delta_{T_U}) \end{aligned}$$

Error due to  $\Delta_S$  (sample selection),  $\Delta_T$  (treatment imbalance), and each due to observed ( $X_i$ ) and unobserved ( $U_i$ ) covariates

# Selection Error

- Definition:

$$\begin{aligned}\Delta_S &\equiv \text{PATE} - \text{SATE} \\ &= \frac{N - n}{N}(\text{NATE} - \text{SATE}),\end{aligned}$$

where NATE is the nonsample average treatment effect.

- Definition:

$$\begin{aligned}\Delta_S &\equiv \text{PATE} - \text{SATE} \\ &= \frac{N - n}{N}(\text{NATE} - \text{SATE}),\end{aligned}$$

where NATE is the nonsample average treatment effect.

- $\Delta_S$  vanishes if:

- Definition:

$$\begin{aligned}\Delta_S &\equiv \text{PATE} - \text{SATE} \\ &= \frac{N - n}{N}(\text{NATE} - \text{SATE}),\end{aligned}$$

where NATE is the nonsample average treatment effect.

- $\Delta_S$  vanishes if:
  - ① The sample is a census ( $I_i = 1$  for all observations and  $n = N$ );



- Definition:

$$\begin{aligned}\Delta_S &\equiv \text{PATE} - \text{SATE} \\ &= \frac{N - n}{N}(\text{NATE} - \text{SATE}),\end{aligned}$$

where NATE is the nonsample average treatment effect.

- $\Delta_S$  vanishes if:
  - 1 The sample is a census ( $I_i = 1$  for all observations and  $n = N$ );
  - 2  $\text{SATE} = \text{NATE}$ ; or

- Definition:

$$\begin{aligned}\Delta_S &\equiv \text{PATE} - \text{SATE} \\ &= \frac{N - n}{N}(\text{NATE} - \text{SATE}),\end{aligned}$$

where NATE is the nonsample average treatment effect.

- $\Delta_S$  vanishes if:
  - 1 The sample is a census ( $I_i = 1$  for all observations and  $n = N$ );
  - 2  $\text{SATE} = \text{NATE}$ ; or
  - 3 Switch quantity of interest from PATE to SATE (recommended!)

# Decomposing Selection Error

# Decomposing Selection Error

- Decomposition:

$$\Delta_S = \Delta_{S_x} + \Delta_{S_u}$$

# Decomposing Selection Error

- Decomposition:

$$\Delta_S = \Delta_{S_X} + \Delta_{S_U}$$

- $\Delta_{S_X} = 0$  when empirical distribution of (observed)  $X$  is identical in population and sample:  $\tilde{F}(X \mid I = 0) = \tilde{F}(X \mid I = 1)$ .

# Decomposing Selection Error

- Decomposition:

$$\Delta_S = \Delta_{S_X} + \Delta_{S_U}$$

- $\Delta_{S_X} = 0$  when empirical distribution of (observed)  $X$  is identical in population and sample:  $\tilde{F}(X | I = 0) = \tilde{F}(X | I = 1)$ .
- $\Delta_{S_U} = 0$  when empirical distribution of (unobserved)  $U$  is identical in population and sample:  $\tilde{F}(U | I = 0) = \tilde{F}(U | I = 1)$ .

# Decomposing Selection Error

- Decomposition:

$$\Delta_S = \Delta_{S_X} + \Delta_{S_U}$$

- $\Delta_{S_X} = 0$  when empirical distribution of (observed)  $X$  is identical in population and sample:  $\tilde{F}(X | I = 0) = \tilde{F}(X | I = 1)$ .
- $\Delta_{S_U} = 0$  when empirical distribution of (unobserved)  $U$  is identical in population and sample:  $\tilde{F}(U | I = 0) = \tilde{F}(U | I = 1)$ .
- conditions are unverifiable:  $X$  is observed only in sample and  $U$  is not observed at all.

# Decomposing Selection Error

- Decomposition:

$$\Delta_S = \Delta_{S_X} + \Delta_{S_U}$$

- $\Delta_{S_X} = 0$  when empirical distribution of (observed)  $X$  is identical in population and sample:  $\tilde{F}(X | I = 0) = \tilde{F}(X | I = 1)$ .
- $\Delta_{S_U} = 0$  when empirical distribution of (unobserved)  $U$  is identical in population and sample:  $\tilde{F}(U | I = 0) = \tilde{F}(U | I = 1)$ .
- conditions are unverifiable:  $X$  is observed only in sample and  $U$  is not observed at all.
- $\Delta_{S_X}$  vanishes if weighting on  $X$



# Decomposing Selection Error

- Decomposition:

$$\Delta_S = \Delta_{S_X} + \Delta_{S_U}$$

- $\Delta_{S_X} = 0$  when empirical distribution of (observed)  $X$  is identical in population and sample:  $\tilde{F}(X | I = 0) = \tilde{F}(X | I = 1)$ .
- $\Delta_{S_U} = 0$  when empirical distribution of (unobserved)  $U$  is identical in population and sample:  $\tilde{F}(U | I = 0) = \tilde{F}(U | I = 1)$ .
- conditions are unverifiable:  $X$  is observed only in sample and  $U$  is not observed at all.
- $\Delta_{S_X}$  vanishes if weighting on  $X$
- $\Delta_{S_U}$  cannot be corrected after the fact

# Decomposing Treatment Imbalance

# Decomposing Treatment Imbalance

- Decomposition:

$$\Delta_T = \Delta_{T_X} + \Delta_{T_U}$$

# Decomposing Treatment Imbalance

- Decomposition:

$$\Delta_T = \Delta_{T_X} + \Delta_{T_U}$$

- $\Delta_{T_X} = 0$  when  $X$  is balanced between treateds and controls:

$$\tilde{F}(X \mid T = 1, I = 1) = \tilde{F}(X \mid T = 0, I = 1).$$

Verifiable from data; can be generated ex ante by blocking or enforced ex post via matching or parametric adjustment

# Decomposing Treatment Imbalance

- Decomposition:

$$\Delta_T = \Delta_{T_X} + \Delta_{T_U}$$

- $\Delta_{T_X} = 0$  when  $X$  is balanced between treateds and controls:

$$\tilde{F}(X \mid T = 1, I = 1) = \tilde{F}(X \mid T = 0, I = 1).$$

Verifiable from data; can be generated ex ante by blocking or enforced ex post via matching or parametric adjustment

- $\Delta_{T_U} = 0$  when  $U$  is balanced between treateds and controls:

$$\tilde{F}(U \mid T = 1, I = 1) = \tilde{F}(U \mid T = 0, I = 1).$$

Unverifiable. Achieved only by assumption or, on average, by random treatment assignment

# Alternative Quantities of Interest

# Alternative Quantities of Interest

- Sample average treatment effect on the treated:

$$\text{SATT} \equiv \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=1\}} \text{TE}_i$$

# Alternative Quantities of Interest

- Sample average treatment effect on the treated:

$$\text{SATT} \equiv \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=1\}} \text{TE}_i$$

- Population average treatment effect on the treated:

$$\text{PATT} \equiv \frac{1}{N^*} \sum_{i \in \{T_i=1\}} \text{TE}_i$$

(where  $N^* = \sum_{i=1}^N T_i$  is the number of treated units in the population)



# Alternative Quantities of Interest

- Sample average treatment effect on the treated:

$$\text{SATT} \equiv \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=1\}} \text{TE}_i$$

- Population average treatment effect on the treated:

$$\text{PATT} \equiv \frac{1}{N^*} \sum_{i \in \{T_i=1\}} \text{TE}_i$$

(where  $N^* = \sum_{i=1}^N T_i$  is the number of treated units in the population)

- When are these of more interest than PATE and SATE? Why never for randomized experiments? Why usually for matching?

# Alternative Quantities of Interest

- Sample average treatment effect on the treated:

$$\text{SATT} \equiv \frac{1}{n/2} \sum_{i \in \{I_i=1, T_i=1\}} \text{TE}_i$$

- Population average treatment effect on the treated:

$$\text{PATT} \equiv \frac{1}{N^*} \sum_{i \in \{T_i=1\}} \text{TE}_i$$

(where  $N^* = \sum_{i=1}^N T_i$  is the number of treated units in the population)

- When are these of more interest than PATE and SATE? Why never for randomized experiments? Why usually for matching?
- Analogous estimation error decomposition:  $\Delta' = \text{PATT} - D$ , holds:

$$\Delta' = (\Delta'_{S_X} + \Delta'_{S_U}) + (\Delta'_{T_X} + \Delta'_{T_U})$$

# Effects of Design Components on Estimation Error

**Design Choice**

$$\Delta_{S_X}$$

$$\Delta_{S_U}$$

$$\Delta_{T_X}$$

$$\Delta_{T_U}$$

# Effects of Design Components on Estimation Error

**Design Choice**

$$\Delta_{S_X}$$

$$\Delta_{S_U}$$

$$\Delta_{T_X}$$

$$\Delta_{T_U}$$

# Effects of Design Components on Estimation Error

## Design Choice

Random sampling

$$\begin{array}{cccc} \Delta_{S_X} & \Delta_{S_U} & \Delta_{T_X} & \Delta_{T_U} \\ \stackrel{\text{avg}}{=} 0 & \stackrel{\text{avg}}{=} 0 & & \end{array}$$

# Effects of Design Components on Estimation Error

## Design Choice

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Random sampling	$\underset{\text{avg}}{=} 0$	$\underset{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\underset{\text{avg}}{=} 0$		

# Effects of Design Components on Estimation Error

## Design Choice

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Random sampling	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\stackrel{\text{avg}}{=} 0$		
Focus on SATE rather than PATE	$= 0$	$= 0$		

# Effects of Design Components on Estimation Error

## Design Choice

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Random sampling	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\stackrel{\text{avg}}{=} 0$		
Focus on SATE rather than PATE	$= 0$	$= 0$		
Weighting for nonrandom sampling	$= 0$	$= ?$		



# Effects of Design Components on Estimation Error

## Design Choice

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Random sampling	$\overset{\text{avg}}{=} 0$	$\overset{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\overset{\text{avg}}{=} 0$		
Focus on SATE rather than PATE	$= 0$	$= 0$		
Weighting for nonrandom sampling	$= 0$	$= ?$		
Large sample size	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$

# Effects of Design Components on Estimation Error

## Design Choice

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Random sampling	$\underset{\text{avg}}{=} 0$	$\underset{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\underset{\text{avg}}{=} 0$		
Focus on SATE rather than PATE	$= 0$	$= 0$		
Weighting for nonrandom sampling	$= 0$	$= ?$		
Large sample size	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$
Random treatment assignment			$\underset{\text{avg}}{=} 0$	$\underset{\text{avg}}{=} 0$

# Effects of Design Components on Estimation Error

## Design Choice

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Random sampling	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\stackrel{\text{avg}}{=} 0$		
Focus on SATE rather than PATE	$= 0$	$= 0$		
Weighting for nonrandom sampling	$= 0$	$= ?$		
Large sample size	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$
Random treatment assignment			$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$
Complete blocking			$= 0$	$= ?$

# Effects of Design Components on Estimation Error

## Design Choice

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Random sampling	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\stackrel{\text{avg}}{=} 0$		
Focus on SATE rather than PATE	$= 0$	$= 0$		
Weighting for nonrandom sampling	$= 0$	$= ?$		
Large sample size	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$
Random treatment assignment			$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$
Complete blocking			$= 0$	$= ?$
Exact matching			$= 0$	$= ?$

# Effects of Design Components on Estimation Error

## Design Choice

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Random sampling	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\stackrel{\text{avg}}{=} 0$		
Focus on SATE rather than PATE	$= 0$	$= 0$		
Weighting for nonrandom sampling	$= 0$	$= ?$		
Large sample size	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$
Random treatment assignment			$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$
Complete blocking			$= 0$	$= ?$
Exact matching			$= 0$	$= ?$

## Assumption

# Effects of Design Components on Estimation Error

## Design Choice

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Random sampling	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\stackrel{\text{avg}}{=} 0$		
Focus on SATE rather than PATE	$= 0$	$= 0$		
Weighting for nonrandom sampling	$= 0$	$= ?$		
Large sample size	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$
Random treatment assignment			$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$
Complete blocking			$= 0$	$= ?$
Exact matching			$= 0$	$= ?$

## Assumption

No selection bias	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$
-------------------	------------------------------	------------------------------

# Effects of Design Components on Estimation Error

## Design Choice

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Random sampling	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\stackrel{\text{avg}}{=} 0$		
Focus on SATE rather than PATE	$= 0$	$= 0$		
Weighting for nonrandom sampling	$= 0$	$= ?$		
Large sample size	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$
Random treatment assignment			$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$
Complete blocking			$= 0$	$= ?$
Exact matching			$= 0$	$= ?$

## Assumption

No selection bias	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Ignorability				$\stackrel{\text{avg}}{=} 0$

# Effects of Design Components on Estimation Error

## Design Choice

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Random sampling	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Complete stratified random sampling	$= 0$	$\stackrel{\text{avg}}{=} 0$		
Focus on SATE rather than PATE	$= 0$	$= 0$		
Weighting for nonrandom sampling	$= 0$	$= ?$		
Large sample size	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$	$\rightarrow ?$
Random treatment assignment			$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$
Complete blocking			$= 0$	$= ?$
Exact matching			$= 0$	$= ?$

## Assumption

No selection bias	$\stackrel{\text{avg}}{=} 0$	$\stackrel{\text{avg}}{=} 0$		
Ignorability				$\stackrel{\text{avg}}{=} 0$
No omitted variables				$= 0$



# Comparing Blocking and Matching

# Comparing Blocking and Matching

- Adding blocking to random assignment is always as or more efficient, and never biased (blocking on pretreatment variables related to the outcome is more efficient)

# Comparing Blocking and Matching

- Adding blocking to random assignment is always as or more efficient, and never biased (blocking on pretreatment variables related to the outcome is more efficient)
- Blocking is like regression adjustment, where the functional form and the parameter values are known

# Comparing Blocking and Matching

- Adding blocking to random assignment is always as or more efficient, and never biased (blocking on pretreatment variables related to the outcome is more efficient)
- Blocking is like regression adjustment, where the functional form and the parameter values are known
- Matching is like blocking, except:

# Comparing Blocking and Matching

- Adding blocking to random assignment is always as or more efficient, and never biased (blocking on pretreatment variables related to the outcome is more efficient)
- Blocking is like regression adjustment, where the functional form and the parameter values are known
- Matching is like blocking, except:
  - 1 to avoid selection error, must change quantity of interest from PATE to PATT or SATT,

# Comparing Blocking and Matching

- Adding blocking to random assignment is always as or more efficient, and never biased (blocking on pretreatment variables related to the outcome is more efficient)
- Blocking is like regression adjustment, where the functional form and the parameter values are known
- Matching is like blocking, except:
  - 1 to avoid selection error, must change quantity of interest from PATE to PATT or SATT,
  - 2 random treatment assignment following matching is impossible

# Comparing Blocking and Matching

- Adding blocking to random assignment is always as or more efficient, and never biased (blocking on pretreatment variables related to the outcome is more efficient)
- Blocking is like regression adjustment, where the functional form and the parameter values are known
- Matching is like blocking, except:
  - 1 to avoid selection error, must change quantity of interest from PATE to PATT or SATT,
  - 2 random treatment assignment following matching is impossible
  - 3 Exact matching, unlike blocking, is dependent on the already-collected data happening to contain sufficiently good matches.

# Comparing Blocking and Matching

- Adding blocking to random assignment is always as or more efficient, and never biased (blocking on pretreatment variables related to the outcome is more efficient)
- Blocking is like regression adjustment, where the functional form and the parameter values are known
- Matching is like blocking, except:
  - 1 to avoid selection error, must change quantity of interest from PATE to PATT or SATT,
  - 2 random treatment assignment following matching is impossible
  - 3 Exact matching, unlike blocking, is dependent on the already-collected data happening to contain sufficiently good matches.
  - 4 In the worst case scenerio, matching (just like regression adjustment) can increase bias (this cannot occur with blocking plus random assignment)



# Comparing Blocking and Matching

- Adding blocking to random assignment is always as or more efficient, and never biased (blocking on pretreatment variables related to the outcome is more efficient)
- Blocking is like regression adjustment, where the functional form and the parameter values are known
- Matching is like blocking, except:
  - ① to avoid selection error, must change quantity of interest from PATE to PATT or SATT,
  - ② random treatment assignment following matching is impossible
  - ③ Exact matching, unlike blocking, is dependent on the already-collected data happening to contain sufficiently good matches.
  - ④ In the worst case scenerio, matching (just like regression adjustment) can increase bias (this cannot occur with blocking plus random assignment)
- Adding matching to a parametric model almost always reduces model dependence and bias, and sometimes variance too

# The Benefits of Major Research Designs: Overview

	$\Delta_{S_X}$	$\Delta_{S_U}$	$\Delta_{T_X}$	$\Delta_{T_U}$
Ideal experiment	$\rightarrow 0$	$\rightarrow 0$	$= 0$	$\rightarrow 0$
Randomized clinical trials (Limited or no blocking)	$\neq 0$	$\neq 0$	$\text{avg} \equiv 0$	$\text{avg} \equiv 0$
Randomized clinical trials (Full blocking)	$\neq 0$	$\neq 0$	$= 0$	$\text{avg} \equiv 0$
Social Science Field Experiment (Limited or no blocking)	$\neq 0$	$\neq 0$	$\rightarrow 0$	$\rightarrow 0$
Survey Experiment (Limited or no blocking)	$\rightarrow 0$	$\rightarrow 0$	$\rightarrow 0$	$\rightarrow 0$
Observational Study (Representative data set, Well-matched)	$\approx 0$	$\approx 0$	$\approx 0$	$\neq 0$
Observational Study (Unrepresentative but partially, correctable data, well-matched)	$\approx 0$	$\neq 0$	$\approx 0$	$\neq 0$
Observational Study (Unrepresentative data set, Well-matched)	$\neq 0$	$\neq 0$	$\approx 0$	$\neq 0$

For column  $Q$ , “ $\rightarrow 0$ ” denotes  $E(Q) = 0$  and  $\lim_{n \rightarrow \infty} \text{Var}(Q) = 0$ , whereas “ $\text{avg} \equiv 0$ ” indicates zero on average, or  $E(Q) = 0$ , for a design with a small  $n$ .  $\Delta_S$  can be set to zero if we switch from PATE to SATE.

# The Ideal Experiment (according to the paper)

# The Ideal Experiment (according to the paper)

- Random selection from well-defined population

# The Ideal Experiment (according to the paper)

- Random selection from well-defined population
- large  $n$

# The Ideal Experiment (according to the paper)

- Random selection from well-defined population
- large  $n$
- blocking on all known confounders

# The Ideal Experiment (according to the paper)

- Random selection from well-defined population
- large  $n$
- blocking on all known confounders
- random treatment assignment within blocks

# The Ideal Experiment (according to the paper)

- Random selection from well-defined population
- large  $n$
- blocking on all known confounders
- random treatment assignment within blocks
- $E(\Delta_{S_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_X}) = 0$



# The Ideal Experiment (according to the paper)

- Random selection from well-defined population
- large  $n$
- blocking on all known confounders
- random treatment assignment within blocks
- $E(\Delta_{S_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_X}) = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$

# The Ideal Experiment (according to the paper)

- Random selection from well-defined population
- large  $n$
- blocking on all known confounders
- random treatment assignment within blocks
- $E(\Delta_{S_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_X}) = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$
- $\Delta_{T_X} = 0$

# The Ideal Experiment (according to the paper)

- Random selection from well-defined population
- large  $n$
- blocking on all known confounders
- random treatment assignment within blocks
- $E(\Delta_{S_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_X}) = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$
- $\Delta_{T_X} = 0$
- $E(\Delta_{T_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_U}) = 0$

# An Even More Ideal Experiment

# An Even More Ideal Experiment

- Begin with a well-defined population

# An Even More Ideal Experiment

- Begin with a well-defined population
- Define sampling strata based on cross-classification of all known confounders

# An Even More Ideal Experiment

- Begin with a well-defined population
- Define sampling strata based on cross-classification of all known confounders
- Random sampling within strata (if strata sample is proportional to population fraction, no weights are needed)

# An Even More Ideal Experiment

- Begin with a well-defined population
- Define sampling strata based on cross-classification of all known confounders
- Random sampling within strata (if strata sample is proportional to population fraction, no weights are needed)
- large  $n$



# An Even More Ideal Experiment

- Begin with a well-defined population
- Define sampling strata based on cross-classification of all known confounders
- Random sampling within strata (if strata sample is proportional to population fraction, no weights are needed)
- large  $n$
- blocking on all known confounders

# An Even More Ideal Experiment

- Begin with a well-defined population
- Define sampling strata based on cross-classification of all known confounders
- Random sampling within strata (if strata sample is proportional to population fraction, no weights are needed)
- large  $n$
- blocking on all known confounders
- random treatment assignment within blocks

# An Even More Ideal Experiment

- Begin with a well-defined population
- Define sampling strata based on cross-classification of all known confounders
- Random sampling within strata (if strata sample is proportional to population fraction, no weights are needed)
- large  $n$
- blocking on all known confounders
- random treatment assignment within blocks
- $\Delta_{S_x} = 0$

# An Even More Ideal Experiment

- Begin with a well-defined population
- Define sampling strata based on cross-classification of all known confounders
- Random sampling within strata (if strata sample is proportional to population fraction, no weights are needed)
- large  $n$
- blocking on all known confounders
- random treatment assignment within blocks
- $\Delta_{S_X} = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$

# An Even More Ideal Experiment

- Begin with a well-defined population
- Define sampling strata based on cross-classification of all known confounders
- Random sampling within strata (if strata sample is proportional to population fraction, no weights are needed)
- large  $n$
- blocking on all known confounders
- random treatment assignment within blocks
- $\Delta_{S_X} = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$
- $\Delta_{T_X} = 0$

# An Even More Ideal Experiment

- Begin with a well-defined population
- Define sampling strata based on cross-classification of all known confounders
- Random sampling within strata (if strata sample is proportional to population fraction, no weights are needed)
- large  $n$
- blocking on all known confounders
- random treatment assignment within blocks
- $\Delta_{S_X} = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$
- $\Delta_{T_X} = 0$
- $E(\Delta_{T_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_U}) = 0$

# Randomized Clinical Trials (Little or no Blocking)

# Randomized Clinical Trials (Little or no Blocking)

- nonrandom selection



# Randomized Clinical Trials (Little or no Blocking)

- nonrandom selection
- small  $n$

# Randomized Clinical Trials (Little or no Blocking)

- nonrandom selection
- small  $n$
- little or no blocking

# Randomized Clinical Trials (Little or no Blocking)

- nonrandom selection
- small  $n$
- little or no blocking
- random treatment assignment

# Randomized Clinical Trials (Little or no Blocking)

- nonrandom selection
- small  $n$
- little or no blocking
- random treatment assignment
- $\Delta_{S_x} \neq 0$

# Randomized Clinical Trials (Little or no Blocking)

- nonrandom selection
- small  $n$
- little or no blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$
- $\Delta_{S_U} \neq 0$

# Randomized Clinical Trials (Little or no Blocking)

- nonrandom selection
- small  $n$
- little or no blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$
- $\Delta_{S_U} \neq 0$
- $E(\Delta_{T_X}) = 0$

# Randomized Clinical Trials (Little or no Blocking)

- nonrandom selection
- small  $n$
- little or no blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$
- $\Delta_{S_U} \neq 0$
- $E(\Delta_{T_X}) = 0$
- $E(\Delta_{T_U}) = 0$

# Randomized Clinical Trials (Full Blocking)



# Randomized Clinical Trials (Full Blocking)

- nonrandom selection

# Randomized Clinical Trials (Full Blocking)

- nonrandom selection
- small  $n$

# Randomized Clinical Trials (Full Blocking)

- nonrandom selection
- small  $n$
- Full blocking

# Randomized Clinical Trials (Full Blocking)

- nonrandom selection
- small  $n$
- Full blocking
- random treatment assignment

# Randomized Clinical Trials (Full Blocking)

- nonrandom selection
- small  $n$
- Full blocking
- random treatment assignment
- $\Delta_{S_x} \neq 0$

# Randomized Clinical Trials (Full Blocking)

- nonrandom selection
- small  $n$
- Full blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$
- $\Delta_{S_U} \neq 0$

# Randomized Clinical Trials (Full Blocking)

- nonrandom selection
- small  $n$
- Full blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$
- $\Delta_{S_U} \neq 0$
- $\Delta_{T_X} = 0$

# Randomized Clinical Trials (Full Blocking)

- nonrandom selection
- small  $n$
- Full blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$
- $\Delta_{S_U} \neq 0$
- $\Delta_{T_X} = 0$
- $E(\Delta_{T_U}) = 0$



# Social Science Field Experiment

# Social Science Field Experiment

- nonrandom selection

# Social Science Field Experiment

- nonrandom selection
- large  $n$

# Social Science Field Experiment

- nonrandom selection
- large  $n$
- limited or no blocking

# Social Science Field Experiment

- nonrandom selection
- large  $n$
- limited or no blocking
- random treatment assignment

# Social Science Field Experiment

- nonrandom selection
- large  $n$
- limited or no blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$  or change PATE to SATE and  $\Delta_{S_X} = 0$

# Social Science Field Experiment

- nonrandom selection
- large  $n$
- limited or no blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$  or change PATE to SATE and  $\Delta_{S_X} = 0$
- $\Delta_{S_U} \neq 0$  or change PATE to SATE and  $\Delta_{S_U} = 0$

# Social Science Field Experiment

- nonrandom selection
- large  $n$
- limited or no blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$  or change PATE to SATE and  $\Delta_{S_X} = 0$
- $\Delta_{S_U} \neq 0$  or change PATE to SATE and  $\Delta_{S_U} = 0$
- $E(\Delta_{T_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_X}) = 0$



# Social Science Field Experiment

- nonrandom selection
- large  $n$
- limited or no blocking
- random treatment assignment
- $\Delta_{S_X} \neq 0$  or change PATE to SATE and  $\Delta_{S_X} = 0$
- $\Delta_{S_U} \neq 0$  or change PATE to SATE and  $\Delta_{S_U} = 0$
- $E(\Delta_{T_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_X}) = 0$
- $E(\Delta_{T_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_U}) = 0$

# Survey Experiment

# Survey Experiment

- random selection

# Survey Experiment

- random selection
- large  $n$

# Survey Experiment

- random selection
- large  $n$
- limited or no blocking

# Survey Experiment

- random selection
- large  $n$
- limited or no blocking
- random treatment assignment (only question wording treatments)

# Survey Experiment

- random selection
- large  $n$
- limited or no blocking
- random treatment assignment (only question wording treatments)
- $E(\Delta_{S_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_X}) = 0$

# Survey Experiment

- random selection
- large  $n$
- limited or no blocking
- random treatment assignment (only question wording treatments)
- $E(\Delta_{S_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_X}) = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$



# Survey Experiment

- random selection
- large  $n$
- limited or no blocking
- random treatment assignment (only question wording treatments)
- $E(\Delta_{S_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_X}) = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$
- $E(\Delta_{T_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_X}) = 0$

# Survey Experiment

- random selection
- large  $n$
- limited or no blocking
- random treatment assignment (only question wording treatments)
- $E(\Delta_{S_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_X}) = 0$
- $E(\Delta_{S_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{S_U}) = 0$
- $E(\Delta_{T_X}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_X}) = 0$
- $E(\Delta_{T_U}) = 0, \lim_{n \rightarrow \infty} V(\Delta_{T_U}) = 0$

# Observational Study, well-matched

# Observational Study, well-matched

- nonrandom selection

# Observational Study, well-matched

- nonrandom selection
- large  $n$

# Observational Study, well-matched

- nonrandom selection
- large  $n$
- no blocking

# Observational Study, well-matched

- nonrandom selection
- large  $n$
- no blocking
- nonrandom treatment assignment

# Observational Study, well-matched

- nonrandom selection
- large  $n$
- no blocking
- nonrandom treatment assignment
- $\Delta_{S_X} \approx 0$  if representative, corrected by weighting, or for estimating SATE; or  $\neq 0$  otherwise



# Observational Study, well-matched

- nonrandom selection
- large  $n$
- no blocking
- nonrandom treatment assignment
- $\Delta_{S_X} \approx 0$  if representative, corrected by weighting, or for estimating SATE; or  $\neq 0$  otherwise
- $\Delta_{S_U} \neq 0$

# Observational Study, well-matched

- nonrandom selection
- large  $n$
- no blocking
- nonrandom treatment assignment
- $\Delta_{S_X} \approx 0$  if representative, corrected by weighting, or for estimating SATE; or  $\neq 0$  otherwise
- $\Delta_{S_U} \neq 0$
- $\Delta_{T_X} \approx 0$  (due to matching well)

# Observational Study, well-matched

- nonrandom selection
- large  $n$
- no blocking
- nonrandom treatment assignment
- $\Delta_{S_X} \approx 0$  if representative, corrected by weighting, or for estimating SATE; or  $\neq 0$  otherwise
- $\Delta_{S_U} \neq 0$
- $\Delta_{T_X} \approx 0$  (due to matching well)
- $\Delta_{T_U} \neq 0$  except by assumption

# What is the Best Design?

# What is the Best Design?

- The ideal design is rarely feasible

# What is the Best Design?

- The ideal design is rarely feasible
- Effort in experimental studies: random assignment

# What is the Best Design?

- The ideal design is rarely feasible
- Effort in experimental studies: random assignment
- Effort in observational studies: measuring and adjusting for  $X$  (via matching or modeling)

# What is the Best Design?

- The ideal design is rarely feasible
- Effort in experimental studies: random assignment
- Effort in observational studies: measuring and adjusting for  $X$  (via matching or modeling)
- the Achilles heal of experimental studies:  $\Delta_S$ , small  $n$ , definition of  $T$



# What is the Best Design?

- The ideal design is rarely feasible
- Effort in experimental studies: random assignment
- Effort in observational studies: measuring and adjusting for  $X$  (via matching or modeling)
- the Achilles heal of experimental studies:  $\Delta_S$ , small  $n$ , definition of  $T$
- the Achilles heal of observational studies:  $\Delta_U$

# What is the Best Design?

- The ideal design is rarely feasible
- Effort in experimental studies: random assignment
- Effort in observational studies: measuring and adjusting for  $X$  (via matching or modeling)
- the Achilles heal of experimental studies:  $\Delta_S$ , small  $n$ , definition of  $T$
- the Achilles heal of observational studies:  $\Delta_U$
- Each design accomodates best to the applications for which it was designed

# Fallacies in Experimental Research

# Fallacies in Experimental Research

- Failure to block on all known covariates

# Fallacies in Experimental Research

- Failure to block on all known covariates
  - seen incorrectly as requiring fewer assumptions (about what to block on)

# Fallacies in Experimental Research

- Failure to block on all known covariates
  - seen incorrectly as requiring fewer assumptions (about what to block on)
  - In fact, blocking helps (except in strange situations)

# Fallacies in Experimental Research

- Failure to block on all known covariates
  - seen incorrectly as requiring fewer assumptions (about what to block on)
  - In fact, blocking helps (except in strange situations)
  - Blocking on relevant covariates is better, so choose carefully.

# Fallacies in Experimental Research

- Failure to block on all known covariates
  - seen incorrectly as requiring fewer assumptions (about what to block on)
  - In fact, blocking helps (except in strange situations)
  - Blocking on relevant covariates is better, so choose carefully.
  - “Block what you can and randomize what you cannot” (Box et al.)



# Fallacies in Experimental Research

- Failure to block on all known covariates
  - seen incorrectly as requiring fewer assumptions (about what to block on)
  - In fact, blocking helps (except in strange situations)
  - Blocking on relevant covariates is better, so choose carefully.
  - “Block what you can and randomize what you cannot” (Box et al.)
- Conducting t-tests to check for balance after random treatment assignment

# Fallacies in Experimental Research

- Failure to block on all known covariates
  - seen incorrectly as requiring fewer assumptions (about what to block on)
  - In fact, blocking helps (except in strange situations)
  - Blocking on relevant covariates is better, so choose carefully.
  - “Block what you can and randomize what you cannot” (Box et al.)
- Conducting t-tests to check for balance after random treatment assignment
  - variables blocked on balance exactly after treatment assignment; so if you’re checking, you missed an opportunity to increase efficiency


# Fallacies in Experimental Research

- Failure to block on all known covariates
  - seen incorrectly as requiring fewer assumptions (about what to block on)
  - In fact, blocking helps (except in strange situations)
  - Blocking on relevant covariates is better, so choose carefully.
  - “Block what you can and randomize what you cannot” (Box et al.)
- Conducting t-tests to check for balance after random treatment assignment
  - variables blocked on balance exactly after treatment assignment; so if you're checking, you missed an opportunity to increase efficiency
  - if variables became available after treatment assignment, a t-test only checks for whether randomization was done appropriately

# Fallacies in Experimental Research

- Failure to block on all known covariates
  - seen incorrectly as requiring fewer assumptions (about what to block on)
  - In fact, blocking helps (except in strange situations)
  - Blocking on relevant covariates is better, so choose carefully.
  - “Block what you can and randomize what you cannot” (Box et al.)
- Conducting t-tests to check for balance after random treatment assignment
  - variables blocked on balance exactly after treatment assignment; so if you’re checking, you missed an opportunity to increase efficiency
  - if variables became available after treatment assignment, a t-test only checks for whether randomization was done appropriately
  - randomization balances on average; any one random assignment is not balanced exactly (which is why its better to block)

# The Balance Test Fallacy in Matching Research



Randomly dropping observations “reduces” imbalance???

# The Balance Test Fallacy: Explanation

# The Balance Test Fallacy: Explanation

- hypothesis tests are a function of balance AND power; we only want to measure balance

# The Balance Test Fallacy: Explanation

- hypothesis tests are a function of balance AND power; we only want to measure balance
- for SATE, PATE, or SPATE, balance is a characteristic of the sample, not some superpopulation; so no need to make an inference



# The Balance Test Fallacy: Explanation

- hypothesis tests are a function of balance AND power; we only want to measure balance
- for SATE, PATE, or SPATE, balance is a characteristic of the sample, not some superpopulation; so no need to make an inference
- Simple linear model (for intuition):

# The Balance Test Fallacy: Explanation

- hypothesis tests are a function of balance AND power; we only want to measure balance
- for SATE, PATE, or SPATE, balance is a characteristic of the sample, not some superpopulation; so no need to make an inference
- Simple linear model (for intuition):
  - Suppose  $E(Y \mid T, X) = \theta + T\beta + X\gamma$

# The Balance Test Fallacy: Explanation

- hypothesis tests are a function of balance AND power; we only want to measure balance
- for SATE, PATE, or SPATE, balance is a characteristic of the sample, not some superpopulation; so no need to make an inference
- Simple linear model (for intuition):
  - Suppose  $E(Y \mid T, X) = \theta + T\beta + X\gamma$
  - Bias in coefficient on  $T$  from regressing  $Y$  on  $T$  (without  $X$ ):  
 $E(\hat{\beta} - \beta \mid T, X) = G\gamma$  (where  $G$  are coefficients from a regression  $X$  on a constant and  $T$ )

# The Balance Test Fallacy: Explanation

- hypothesis tests are a function of balance AND power; we only want to measure balance
- for SATE, PATE, or SPATE, balance is a characteristic of the sample, not some superpopulation; so no need to make an inference
- Simple linear model (for intuition):
  - Suppose  $E(Y \mid T, X) = \theta + T\beta + X\gamma$
  - Bias in coefficient on  $T$  from regressing  $Y$  on  $T$  (without  $X$ ):  
 $E(\hat{\beta} - \beta \mid T, X) = G\gamma$  (where  $G$  are coefficients from a regression  $X$  on a constant and  $T$ )
  - Imbalance is due to  $G$  only.

# The Balance Test Fallacy: Explanation

- hypothesis tests are a function of balance AND power; we only want to measure balance
- for SATE, PATE, or SPATE, balance is a characteristic of the sample, not some superpopulation; so no need to make an inference
- Simple linear model (for intuition):
  - Suppose  $E(Y \mid T, X) = \theta + T\beta + X\gamma$
  - Bias in coefficient on  $T$  from regressing  $Y$  on  $T$  (without  $X$ ):  
 $E(\hat{\beta} - \beta \mid T, X) = G\gamma$  (where  $G$  are coefficients from a regression  $X$  on a constant and  $T$ )
  - Imbalance is due to  $G$  only.
  - If  $G = 0$ , bias=0

# The Balance Test Fallacy: Explanation

- hypothesis tests are a function of balance AND power; we only want to measure balance
- for SATE, PATE, or SPATE, balance is a characteristic of the sample, not some superpopulation; so no need to make an inference
- Simple linear model (for intuition):
  - Suppose  $E(Y | T, X) = \theta + T\beta + X\gamma$
  - Bias in coefficient on  $T$  from regressing  $Y$  on  $T$  (without  $X$ ):  
 $E(\hat{\beta} - \beta | T, X) = G\gamma$  (where  $G$  are coefficients from a regression  $X$  on a constant and  $T$ )
  - Imbalance is due to  $G$  only.
  - If  $G = 0$ , bias=0
  - If  $G \neq 0$ , bias can be any size due to  $\gamma$

# The Balance Test Fallacy: Explanation

- hypothesis tests are a function of balance AND power; we only want to measure balance
- for SATE, PATE, or SPATE, balance is a characteristic of the sample, not some superpopulation; so no need to make an inference
- Simple linear model (for intuition):
  - Suppose  $E(Y \mid T, X) = \theta + T\beta + X\gamma$
  - Bias in coefficient on  $T$  from regressing  $Y$  on  $T$  (without  $X$ ):  
 $E(\hat{\beta} - \beta \mid T, X) = G\gamma$  (where  $G$  are coefficients from a regression  $X$  on a constant and  $T$ )
  - Imbalance is due to  $G$  only.
  - If  $G = 0$ , bias=0
  - If  $G \neq 0$ , bias can be any size due to  $\gamma$
  - If you cannot control  $G$  (and we don't usually try to minimize  $\gamma$  so we don't cook the books), then  $G$  must be reduced without limit.

# The Balance Test Fallacy: Explanation

- hypothesis tests are a function of balance AND power; we only want to measure balance
- for SATE, PATE, or SPATE, balance is a characteristic of the sample, not some superpopulation; so no need to make an inference
- Simple linear model (for intuition):
  - Suppose  $E(Y \mid T, X) = \theta + T\beta + X\gamma$
  - Bias in coefficient on  $T$  from regressing  $Y$  on  $T$  (without  $X$ ):  
 $E(\hat{\beta} - \beta \mid T, X) = G\gamma$  (where  $G$  are coefficients from a regression  $X$  on a constant and  $T$ )
  - Imbalance is due to  $G$  only.
  - If  $G = 0$ , bias=0
  - If  $G \neq 0$ , bias can be any size due to  $\gamma$
  - If you cannot control  $G$  (and we don't usually try to minimize  $\gamma$  so we don't cook the books), then  $G$  must be reduced without limit.
- No threshold level is safe.