

✓ **Congratulations! You passed!**

Next Item



1 / 1  
points

1.

What is the variance of the distribution of the average an IID draw of  $n$  observations from a population with mean  $\mu$  and variance  $\sigma^2$ .



$\frac{\sigma^2}{n}$



**Correct**

$\text{Var}(\bar{X}) = \sigma^2/n$



$\sigma/n$



$\sigma^2$



$2\sigma/\sqrt{n}$



1 / 1  
points

2.

Suppose that diastolic blood pressures (DBPs) for men aged 35-44 are normally distributed with a mean of 80 (mm Hg) and a standard deviation of 10. About what is the probability that a random 35-44 year old has a DBP less than 70?



16%



**Correct**

```
1 pnorm(70, mean = 80, sd = 10)
```

```
1 ## [1] 0.1587
```



8%



## Quiz 2

Quiz, 8 questions

22%

32%

8/8 points (100%)



1 / 1  
points

3.

Brain volume for adult women is normally distributed with a mean of about 1,100 cc for women with a standard deviation of 75 cc. What brain volume represents the 95th percentile?

- ☐ approximately 977
- ☐ approximately 1175
- ☐ approximately 1247
- ☒ approximately 1223

Correct

```
1 qnorm(0.95, mean = 1100, sd = 75)
```

```
1 ## [1] 1223
```



1 / 1  
points

4.

Refer to the previous question. Brain volume for adult women is about 1,100 cc for women with a standard deviation of 75 cc. Consider the sample mean of 100 random adult women from this population. What is the 95th percentile of the distribution of that sample mean?

- ☐ approximately 1110 cc
- ☒ approximately 1112 cc

Correct

```
1 qnorm(0.95, mean = 1100, sd = 75/sqrt(100))
```

```
1 [1] 1112
```

- ☐ approximately 1115 cc



1 / 1  
points

5.

You flip a fair coin 5 times, about what's the probability of getting 4 or 5 heads?

- ☐ 3%
- ☐ 6%
- ☐ 12%
- ☒ 19%

**Correct**

$$\binom{5}{4}2^{-5} + \binom{5}{5}2^{-5} \approx 19\%$$

```
1 pbinom(3, size = 5, prob = 0.5, lower.tail = FALSE)
```

```
1 ## [1] 0.1875
```



1 / 1  
points

6.

The respiratory disturbance index (RDI), a measure of sleep disturbance, for a specific population has a mean of 15 (sleep events per hour) and a standard deviation of 10. They are not normally distributed. Give your best estimate of the probability that a sample mean RDI of 100 people is between 14 and 16 events per hour?

- ☐ 95%
- ☒ 68%

**Correct**

The standard error of the mean is  $10/\sqrt{100} = 1$ . Thus between 14 and 16 is with one standard deviation of the mean of the distribution of the sample mean. Thus it should be about 68%.

```
1 pnorm(16, mean = 15, sd = 1) - pnorm(14, mean = 15, sd = 1)
```

## Quiz 2

Quiz, 8 questions

1 ## [1] 0.6827

8/8 points (100%)

☐ 34%

☐ 47.5%



1 / 1  
points

7.

Consider a standard uniform density. The mean for this density is .5 and the variance is  $1/12$ . You sample 1,000 observations from this distribution and take the sample mean, what value would you expect it to be near?

☐ 0.75

☒ 0.5



**Correct**

Via the LLN it should be near .5.

☐ 0.10

☐ 0.25



1 / 1  
points

8.

The number of people showing up at a bus stop is assumed to be

Poisson with a mean of 5 people per hour. You watch the bus

stop for 3 hours. About what's the probability of viewing 10 or fewer people?

☐ 0.06

☐ 0.08

☐ 0.03

☒ 0.12



**Correct**

```
1 ppois(10, lambda = 15)
```