

Potential FN constructs with C-terminal receptor binding (n=79 constructs / 40 genes)

Grouped by classification method · subset of the 446-pFN OASIS pool

Gene	n constr	Method	Confidence	Tiles / peptides (representative)
ADM2	1	Amidation	high	Adrenomedullin-2
ADML	1	Amidation	high	Adrenomedullin
CALCA	1	Amidation	high	Calcitonin gene-related peptide 1
CALCB	1	Amidation	high	Calcitonin gene-related peptide 2
CCKN	1	Amidation	high	Cholecystokinin
COLI	11	Amidation	high	NPP, Lipotropin beta, Met-enkephalin (+8 more)
GAST	5	Amidation	high	Gastrin-14, Gastrin, Big gastrin (+2 more)
GLUC	7	Amidation	high	Glucagon, Glicentin-related polypeptide, Glucagon-like peptide 1(7-...
GRP	2	Amidation	high	Neuromedin-C, Gastrin-releasing peptide
IAPP	1	Amidation	high	Islet amyloid polypeptide
MCH	3	Amidation	high	Neuropeptide-glutamic acid-isoleucine, Melanin-concentrating hormon...
NEU1	2	Amidation	high	Neurophysin 1, Oxytocin
NEU2	3	Amidation	high	Neurophysin 2, Arg-vasopressin, Copeptin
NMB	1	Amidation	high	Neuromedin-B-32
NPFF	3	Amidation	high	Neuropeptide FF, Neuropeptide SF, Neuropeptide AF
NPY	1	Amidation	high	Neuropeptide Y
OREX	2	Amidation	high	Orexin-B, Orexin-A
OX26	1	Amidation	high	QRF-amide
PACA	3	Amidation	high	Pituitary adenylate cyclase-activating polypeptide 27, Pituitary ad...
PAHO	1	Amidation	high	Pancreatic polypeptide
PYY	2	Amidation	high	Peptide YY(3-36), Peptide YY
SECR	1	Amidation	high	Secretin
SLIB	1	Amidation	high	Somatoliberin
SMS	3	Amidation	high	Neuronostatin, Somatostatin-14, Somatostatin-28
TKN4	3	Amidation	high	Endokinin-A/B, Endokinin-C, Endokinin-A
UCN3	1	Amidation	high	Urocortin-3
VIP	2	Amidation	high	Vasoactive intestinal peptide, Intestinal peptide PHV-42
CD40L	2	TNF family	moderate	CD40 ligand, soluble form, CD40 ligand (CD40-L) (T-cell antigen Gp3...
EDA	2	TNF family	moderate	Ectodysplasin-A, secreted form, Ectodysplasin-A (Ectodermal dysplas...
TNF11	1	TNF family	moderate	Tumor necrosis factor ligand superfamily member 11, soluble form
TNF12	1	TNF family	moderate	Tumor necrosis factor ligand superfamily member 12, secreted form
TNF13	1	TNF family	moderate	Tumor necrosis factor ligand superfamily member 13
TNF15	1	TNF family	moderate	Tumor necrosis factor ligand superfamily member 15, secreted form
TNFA	1	TNF family	moderate	Tumor necrosis factor, soluble form
TNFB	1	TNF family	moderate	Lymphotoxin-alpha
TNFC	1	TNF family	moderate	Lymphotoxin-beta (LT-beta) (Tumor necrosis factor C) (TNF-C) (Tumor...
TNFL4	1	TNF family	moderate	Tumor necrosis factor ligand superfamily member 4 (Glycoprotein Gp3...
TNFL8	1	TNF family	moderate	Tumor necrosis factor ligand superfamily member 8 (CD30 ligand) (CD...
TNFL9	1	TNF family	moderate	Tumor necrosis factor ligand superfamily member 9 (4-1BB ligand) (4...
IL15	1	PDB burial	high	Interleukin-15

Method legend: Amidation = UniProt-annotated C-terminal amidation pharmacophore (neuropeptides); TNF family = family-based (TNF superfamily C-term ectodomain, type II TM); PDB burial = PDB co-crystal C-term burial >50%.