Class14: RNASeq Mini-project

Chelsea A16871799

The authors report on differential analysis of lung fibroblasts in response to loss of the developmental transcription factor HOXA1. Their results and others indicate that HOXA1 is required for lung fibroblast and HeLa cell cycle progression. In particular their analysis show that "loss of HOXA1 results in significant expression level changes in thousands of individual transcripts, along with isoform switching events in key regulators of the cell cycle". For our session we have used their Sailfish gene-level estimated counts and hence are restricted to protein-coding genes only.

##Data Import

```
metaFile <- "GSE37704_metadata.csv"
countFile <- "GSE37704_featurecounts.csv"

# Import metadata and take a peak
colData = read.csv(metaFile, row.names=1)
head(colData)
```

```
            condition
SRR493366 control_sirna
SRR493367 control_sirna
SRR493368 control_sirna
SRR493369      hoxa1_kd
SRR493370      hoxa1_kd
SRR493371      hoxa1_kd
```

```
countData=read.csv(countFile, row.names=1)
head(countData)
```

```
                length SRR493366 SRR493367 SRR493368 SRR493369 SRR493370
ENSG00000186092    918         0         0         0         0         0
```

ENSG00000279928	718	0	0	0	0	0
ENSG00000279457	1982	23	28	29	29	28
ENSG00000278566	939	0	0	0	0	0
ENSG00000273547	939	0	0	0	0	0
ENSG00000187634	3214	124	123	205	207	212

	SRR493371
ENSG00000186092	0
ENSG00000279928	0
ENSG00000279457	46
ENSG00000278566	0
ENSG00000273547	0
ENSG00000187634	258

we need the countData and colData files to match up so we will need to remove that odd first column in countData namely contData$length

```
countData <- as.matrix(countData[,-1])
head(countData)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000186092	0	0	0	0	0	0
ENSG00000279928	0	0	0	0	0	0
ENSG00000279457	23	28	29	29	28	46
ENSG00000278566	0	0	0	0	0	0
ENSG00000273547	0	0	0	0	0	0
ENSG00000187634	124	123	205	207	212	258

Q. Complete the code below to filter countData to exclude genes (i.e. rows) where we have 0 read count across all samples (i.e. columns).

```
to.rm.inds <- rowSums(countData)==0
```

```
countData =countData[!to.rm.inds,]
head(countData)
```

	SRR493366	SRR493367	SRR493368	SRR493369	SRR493370	SRR493371
ENSG00000279457	23	28	29	29	28	46
ENSG00000187634	124	123	205	207	212	258
ENSG00000188976	1637	1831	2383	1226	1326	1504
ENSG00000187961	120	153	180	236	255	357

```
ENSG00000187583        24        48        65        44        48        64
ENSG00000187642         4         9        16        14        16        16
```

```
nrow(countData)
```

```
[1] 15975
```

##DESeq Setup and analysis

```
library(DESeq2)
```

```
Loading required package: S4Vectors
```

```
Loading required package: stats4
```

```
Loading required package: BiocGenerics
```

```

Attaching package: 'BiocGenerics'
```

```
The following objects are masked from 'package:stats':

    IQR, mad, sd, var, xtabs
```

```
The following objects are masked from 'package:base':

    anyDuplicated, aperm, append, as.data.frame, basename, cbind,
    colnames, dirname, do.call, duplicated, eval, evalq, Filter, Find,
    get, grep, grepl, intersect, is.unsorted, lapply, Map, mapply,
    match, mget, order, paste, pmax, pmax.int, pmin, pmin.int,
    Position, rank, rbind, Reduce, rownames, sapply, setdiff, sort,
    table, tapply, union, unique, unsplit, which.max, which.min
```

```

Attaching package: 'S4Vectors'
```

```
The following object is masked from 'package:utils':

    findMatches
```

```
The following objects are masked from 'package:base':

    expand.grid, I, unname

Loading required package: IRanges

Loading required package: GenomicRanges

Loading required package: GenomeInfoDb

Loading required package: SummarizedExperiment

Warning: package 'SummarizedExperiment' was built under R version 4.3.2

Loading required package: MatrixGenerics

Loading required package: matrixStats


Attaching package: 'MatrixGenerics'

The following objects are masked from 'package:matrixStats':

    colAlls, colAnyNAs, colAnys, colAvgsPerRowSet, colCollapse,
    colCounts, colCummaxs, colCummins, colCumprods, colCumsums,
    colDiffs, colIQRDiffs, colIQRs, colLogSumExps, colMadDiffs,
    colMads, colMaxs, colMeans2, colMedians, colMins, colOrderStats,
    colProds, colQuantiles, colRanges, colRanks, colSdDiffs, colSds,
    colSums2, colTabulates, colVarDiffs, colVars, colWeightedMads,
    colWeightedMeans, colWeightedMedians, colWeightedSds,
    colWeightedVars, rowAlls, rowAnyNAs, rowAnys, rowAvgsPerColSet,
    rowCollapse, rowCounts, rowCummaxs, rowCummins, rowCumprods,
    rowCumsums, rowDiffs, rowIQRDiffs, rowIQRs, rowLogSumExps,
    rowMadDiffs, rowMads, rowMaxs, rowMeans2, rowMedians, rowMins,
    rowOrderStats, rowProds, rowQuantiles, rowRanges, rowRanks,
    rowSdDiffs, rowSds, rowSums2, rowTabulates, rowVarDiffs, rowVars,
    rowWeightedMads, rowWeightedMeans, rowWeightedMedians,
    rowWeightedSds, rowWeightedVars

Loading required package: Biobase
```

```
Welcome to Bioconductor

    Vignettes contain introductory material; view with
    'browseVignettes()'. To cite Bioconductor, see
    'citation("Biobase")', and for packages 'citation("pkgname")'.


Attaching package: 'Biobase'

The following object is masked from 'package:MatrixGenerics':

    rowMedians

The following objects are masked from 'package:matrixStats':

    anyMissing, rowMedians
```

```
dds = DESeqDataSetFromMatrix(countData=countData,
                             colData=colData,
                             design=~condition)
```

```
Warning in DESeqDataSet(se, design = design, ignoreRank): some variables in
design formula are characters, converting to factors
```

```
dds = DESeq(dds)
```

```
estimating size factors

estimating dispersions

gene-wise dispersion estimates

mean-dispersion relationship

final dispersion estimates

fitting model and testing
```

```
dds
```

```
class: DESeqDataSet
dim: 15975 6
metadata(1): version
assays(4): counts mu H cooks
rownames(15975): ENSG00000279457 ENSG00000187634 ... ENSG00000276345
  ENSG00000271254
rowData names(22): baseMean baseVar ... deviance maxCooks
colnames(6): SRR493366 SRR493367 ... SRR493370 SRR493371
colData names(2): condition sizeFactor
```

```
res = results(dds, contrast=c("condition", "hoxa1_kd", "control_sirna"))
```

Q. Call the summary() function on your results to get a sense of how many genes are up or down-regulated at the default 0.1 p-value cutoff.

```
summary(res)
```

```
out of 15975 with nonzero total read count
adjusted p-value < 0.1
LFC > 0 (up)       : 4349, 27%
LFC < 0 (down)     : 4396, 28%
outliers [1]       : 0, 0%
low counts [2]     : 1237, 7.7%
(mean count < 0)
[1] see 'cooksCutoff' argument of ?results
[2] see 'independentFiltering' argument of ?results
```

volcano plot

```
plot( res$log2FoldChange, -log(res$padj) )
```



Q. Improve this plot by completing the below code, which adds color and axis labels

```
# Make a color vector for all genes
mycols <- rep("gray", nrow(res) )

# Color red the genes with absolute fold change above 2
mycols[ abs(res$log2FoldChange) > 2 ] <- "red"

# Color blue those with adjusted p-value less than 0.01
#  and absolute fold change more than 2
inds <- (res$padj<0.01) & (abs(res$log2FoldChange) > 2 )
mycols[ inds ] <- "blue"

plot( res$log2FoldChange, -log(res$padj), col=mycols, xlab="Log2(FoldChange)", ylab="-Log(
```



Q. Use the mapIDs() function multiple times to add SYMBOL, ENTREZID and GENENAME annotation to our results by completing the code below.

```
library("AnnotationDbi")
library("org.Hs.eg.db")
```

```
columns(org.Hs.eg.db)
```

```
 [1] "ACCNUM"       "ALIAS"        "ENSEMBL"      "ENSEMBLPROT"  "ENSEMBLTRANS"
 [6] "ENTREZID"     "ENZYME"       "EVIDENCE"     "EVIDENCEALL"  "GENENAME"
[11] "GENETYPE"     "GO"           "GOALL"        "IPI"          "MAP"
[16] "OMIM"         "ONTOLOGY"     "ONTOLOGYALL"  "PATH"         "PFAM"
[21] "PMID"         "PROSITE"      "REFSEQ"       "SYMBOL"       "UCSCKG"
[26] "UNIPROT"
```

```
res$symbol = mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
```

```
                    column="SYMBOL",
                    multiVals="first")
```

```
'select()' returned 1:many mapping between keys and columns
```

```
  res$entrez = mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="ENTREZID",
                    multiVals="first")
```

```
'select()' returned 1:many mapping between keys and columns
```

```
  res$name =   mapIds(org.Hs.eg.db,
                    keys=row.names(res),
                    keytype="ENSEMBL",
                    column="GENENAME",
                    multiVals="first")
```

```
'select()' returned 1:many mapping between keys and columns
```

```
  head(res, 10)
```

```
log2 fold change (MLE): condition hoxa1_kd vs control_sirna
Wald test p-value: condition hoxa1 kd vs control sirna
DataFrame with 10 rows and 9 columns
                  baseMean log2FoldChange     lfcSE       stat      pvalue
                 <numeric>      <numeric> <numeric>  <numeric>   <numeric>
ENSG00000279457   29.913579      0.1792571 0.3248216   0.551863 5.81042e-01
ENSG00000187634  183.229650      0.4264571 0.1402658   3.040350 2.36304e-03
ENSG00000188976 1651.188076     -0.6927205 0.0548465 -12.630158 1.43989e-36
ENSG00000187961  209.637938      0.7297556 0.1318599   5.534326 3.12428e-08
ENSG00000187583   47.255123      0.0405765 0.2718928   0.149237 8.81366e-01
ENSG00000187642   11.979750      0.5428105 0.5215599   1.040744 2.97994e-01
ENSG00000188290  108.922128      2.0570638 0.1969053  10.446970 1.51282e-25
ENSG00000187608  350.716868      0.2573837 0.1027266   2.505522 1.22271e-02
ENSG00000188157 9128.439422      0.3899088 0.0467163   8.346304 7.04321e-17
```

```
ENSG00000237330    0.158192      0.7859552 4.0804729   0.192614 8.47261e-01
                       padj      symbol      entrez                   name
                  <numeric> <character> <character>            <character>
ENSG00000279457 6.86555e-01          NA          NA                     NA
ENSG00000187634 5.15718e-03      SAMD11      148398 sterile alpha motif ..
ENSG00000188976 1.76549e-35       NOC2L       26155 NOC2 like nucleolar ..
ENSG00000187961 1.13413e-07      KLHL17      339451 kelch like family me..
ENSG00000187583 9.19031e-01     PLEKHN1       84069 pleckstrin homology ..
ENSG00000187642 4.03379e-01       PERM1       84808 PPARGC1 and ESRR ind..
ENSG00000188290 1.30538e-24        HES4       57801 hes family bHLH tran..
ENSG00000187608 2.37452e-02       ISG15        9636 ISG15 ubiquitin like..
ENSG00000188157 4.21963e-16        AGRN      375790                  agrin
ENSG00000237330          NA      RNF223      401934 ring finger protein ..
```

Save results

Q. Finally for this section let's reorder these results by adjusted p-value and save them to a CSV file in your current project directory.

```
res = res[order(res$pvalue),]
write.csv(res, file="deseq_results.csv")
```

##Geneset enrichment Now we can load the packages and setup the KEGG data-sets we need

```
library(pathview)
```

```
#############################################################################
Pathview is an open source software package distributed under GNU General
Public License version 3 (GPLv3). Details of GPLv3 is available at
http://www.gnu.org/licenses/gpl-3.0.html. Particullary, users are required to
formally cite the original Pathview paper (not just mention it) in publications
or products. For details, do citation("pathview") within R.

The pathview downloads and uses KEGG data. Non-academic uses may require a KEGG
license agreement (details at http://www.kegg.jp/kegg/legal.html).
#############################################################################
```

```
library(gage)
```

```
library(gageData)

data(kegg.sets.hs)
data(sigmet.idx.hs)

# Focus on signaling and metabolic pathways only
kegg.sets.hs = kegg.sets.hs[sigmet.idx.hs]

# Examine the first 3 pathways
head(kegg.sets.hs, 3)
```

```
$`hsa00232 Caffeine metabolism`
[1] "10"   "1544" "1548" "1549" "1553" "7498" "9"

$`hsa00983 Drug metabolism - other enzymes`
 [1] "10"     "1066"   "10720"  "10941"  "151531" "1548"   "1549"   "1551"
 [9] "1553"   "1576"   "1577"   "1806"   "1807"   "1890"   "221223" "2990"
[17] "3251"   "3614"   "3615"   "3704"   "51733"  "54490"  "54575"  "54576"
[25] "54577"  "54578"  "54579"  "54600"  "54657"  "54658"  "54659"  "54963"
[33] "574537" "64816"  "7083"   "7084"   "7172"   "7363"   "7364"   "7365"
[41] "7366"   "7367"   "7371"   "7372"   "7378"   "7498"   "79799"  "83549"
[49] "8824"   "8833"   "9"      "978"

$`hsa00230 Purine metabolism`
  [1] "100"    "10201"  "10606"  "10621"  "10622"  "10623"  "107"    "10714"
  [9] "108"    "10846"  "109"    "111"    "11128"  "11164"  "112"    "113"
 [17] "114"    "115"    "122481" "122622" "124583" "132"    "158"    "159"
 [25] "1633"   "171568" "1716"   "196883" "203"    "204"    "205"    "221823"
 [33] "2272"   "22978"  "23649"  "246721" "25885"  "2618"   "26289"  "270"
 [41] "271"    "27115"  "272"    "2766"   "2977"   "2982"   "2983"   "2984"
 [49] "2986"   "2987"   "29922"  "3000"   "30833"  "30834"  "318"    "3251"
 [57] "353"    "3614"   "3615"   "3704"   "377841" "471"    "4830"   "4831"
 [65] "4832"   "4833"   "4860"   "4881"   "4882"   "4907"   "50484"  "50940"
 [73] "51082"  "51251"  "51292"  "5136"   "5137"   "5138"   "5139"   "5140"
 [81] "5141"   "5142"   "5143"   "5144"   "5145"   "5146"   "5147"   "5148"
 [89] "5149"   "5150"   "5151"   "5152"   "5153"   "5158"   "5167"   "5169"
 [97] "51728"  "5198"   "5236"   "5313"   "5315"   "53343"  "54107"  "5422"
[105] "5424"   "5425"   "5426"   "5427"   "5430"   "5431"   "5432"   "5433"
[113] "5434"   "5435"   "5436"   "5437"   "5438"   "5439"   "5440"   "5441"
```

```
[121] "5471"   "548644" "55276"  "5557"   "5558"   "55703"  "55811"  "55821"
[129] "5631"   "5634"   "56655"  "56953"  "56985"  "57804"  "58497"  "6240"
[137] "6241"   "64425"  "646625" "654364" "661"    "7498"   "8382"   "84172"
[145] "84265"  "84284"  "84618"  "8622"   "8654"   "87178"  "8833"   "9060"
[153] "9061"   "93034"  "953"    "9533"   "954"    "955"    "956"    "957"
[161] "9583"   "9615"
```

```
foldchanges = res$log2FoldChange
names(foldchanges) = res$entrez
head(foldchanges)
```

```
     1266     54855      1465     51232      2034      2317
-2.422719  3.201955 -2.313738 -2.059631 -1.888019 -1.649792
```

```
# Get the results
keggres = gage(foldchanges, gsets=kegg.sets.hs)
```

```
attributes(keggres)
```

```
$names
[1] "greater" "less"    "stats"
```

```
# Look at the first few down (less) pathways
head(keggres$less)
```

```
                                      p.geomean stat.mean        p.val
hsa04110 Cell cycle                8.995727e-06 -4.378644 8.995727e-06
hsa03030 DNA replication           9.424076e-05 -3.951803 9.424076e-05
hsa03013 RNA transport             1.375901e-03 -3.028500 1.375901e-03
hsa03440 Homologous recombination  3.066756e-03 -2.852899 3.066756e-03
hsa04114 Oocyte meiosis            3.784520e-03 -2.698128 3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis 8.961413e-03 -2.405398 8.961413e-03
                                         q.val set.size         exp1
hsa04110 Cell cycle                0.001448312      121 8.995727e-06
hsa03030 DNA replication           0.007586381       36 9.424076e-05
hsa03013 RNA transport             0.073840037      144 1.375901e-03
hsa03440 Homologous recombination  0.121861535       28 3.066756e-03
hsa04114 Oocyte meiosis            0.121861535      102 3.784520e-03
hsa00010 Glycolysis / Gluconeogenesis 0.212222694       53 8.961413e-03
```

The has04110 and hsa03030 seems to be the one pick out.

```
pathview(gene.data=foldchanges, pathway.id="hsa04110")
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/chelseazhong/Desktop/bimm143/class14
```

```
Info: Writing image file hsa04110.pathview.png
```




```
pathview(gene.data=foldchanges, pathway.id="hsa03030")
```

```
'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/chelseazhong/Desktop/bimm143/class14

Info: Writing image file hsa03030.pathview.png
```




```
## Focus on top 5 upregulated pathways here for demo purposes only
keggrespathways <- rownames(keggres$greater)[1:5]
```

```
# Extract the 8 character long IDs part of each string
keggresids = substr(keggrespathways, start=1, stop=8)
keggresids
```

```
[1] "hsa04640" "hsa04630" "hsa00140" "hsa04142" "hsa04330"
```

```
pathview(gene.data=foldchanges, pathway.id=keggresids, species="hsa")
```

```
'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/chelseazhong/Desktop/bimm143/class14

Info: Writing image file hsa04640.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/chelseazhong/Desktop/bimm143/class14

Info: Writing image file hsa04630.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/chelseazhong/Desktop/bimm143/class14

Info: Writing image file hsa00140.pathview.png

'select()' returned 1:1 mapping between keys and columns

Info: Working in directory /Users/chelseazhong/Desktop/bimm143/class14

Info: Writing image file hsa04142.pathview.png

Info: some node width is different from others, and hence adjusted!

'select()' returned 1:1 mapping between keys and columns
```

```
Info: Working in directory /Users/chelseazhong/Desktop/bimm143/class14

Info: Writing image file hsa04330.pathview.png
```




JAK-STAT SIGNALING PATHWAY
-1
0
1
ECS complex
Ubiquitin mediated proteolysis
Cytokine-cytokine receptor interaction
TC-PTP
SHP1
STAM
STAT dimerization
Interleukins
IL2family
IL3family
IL6family
IL10family
IL12/23
Hormone
Interferons
IFN-I/III
IFN-II
Growth factors
GF
Receptor
JAK1/3
JAK2
JAK1/2
TYK2
JAK1
STAT
STAT1/3/5/6
STAT5
STAT1/3/6
STAT3
STAT3/4
STAT3/5
STAT1/2
STAT1/3
IRF9
TC-PTP
PIAS
CBP/P300
SLIM
DNA
Proteasome
CIS
SOCS
Bcl-2
MCL1
Bcl-XL
PIM1
Apoptosis
Anti-apoptosis
Cell cycle
c-Myc
CycD
Cell-cycle progression
p21
Cell-cycle inhibition
AOX
Lipid metabolism
GFAP
Differentiation
MAPK signaling pathway
SHP2
GRB
SOS
Ras
Raf
Proliferation
Differentiation
PI3K
AKT
mTOR
Cell cycle
Cell survival
PI3K-AKT signaling pathway
Data on KEGG graph
Rendered by Pathview


STEROID HORMONE BIOSYNTHESIS
-1
0
1
Steroid biosynthesis
Cholesterol
Cholesterol sulfate
20α-Hydroxy-cholesterol
22β-Hydroxy-cholesterol
20α,22β-Dihydroxy-cholesterol
17α,20α-Dihydroxy-cholesterol
4-Methylpentanal
Pregnenolone
Pregnenolone-sulfate
21-Hydroxy-pregnenolone
17α-Hydroxy-pregnenolone
11-Deoxy-corticosterone
Progesterone
11α-Hydroxy-progesterone
20α-Hydroxy-progesterone
7α-Hydroxy-pregnenolone
17α-Hydroxy-progesterone
17α,20α-Dihydroxy-pregn-4-en-3-one
11-Deoxycortisol
17α,21-Dihydroxy-pregnenolone
11β,17α,21-Trihydroxy-pregnenolone
Cortisol
Cortisone
11β-Hydroxyandrost-4-ene-3,17-dione
Adrenosterone
Dehydroepiandro-sterone
Dehydroepiandro-steron sulfate
7α-Hydroxydehydro-epiandrosterone
16α-Hydroxyandrost-4-ene-3,17-dione
16α-Hydroxydehydro-epiandrosterone
3β,17β-Dihydroxy-androst-5-ene
Androst-4-ene-3,17-dione
7α-Hydroxy-androstenedione
7α-Hydroxy-testosterone
Testosterone
19-Hydroxy-testosterone
19-Oxotestosterone
5β-Androstane-3,17-dione
Androsterone
Androsterone-glucuronide
5α-Dihydro-testosterone
Androstan-3alpha,17beta-diol
5β-Androstane-3α-ol-17-one
Etiocholan-3α-ol-17-one
Etiocholan-3α-ol-17-one 3-glucuronide
3-Oxo-13,17-secoandrost-4-ene-17,13α-lactone
19-Hydroxyandrost-4-ene-3,17-dione
19-Oxoandrost-4-ene-3,17-dione
5α-Dihydro-deoxycorticosterone
Allotetrahydro-deoxycorticosterone
Aldosterone hemiacetal
Aldosterone
18-Hydroxy-corticosterone
11β,21-Dihydroxy-3,20-oxo-5β-pregnan-18-al
3α,11β,21-Trihydroxy-20-oxo-5β-pregnan-18-al
11-Dehydro-corticosterone
21-Hydroxy-5β-pregnane-3,11,20-trione
Corticosterone
11β,21-Dihydroxy-5β-pregnane-3,20-dione
Tetrahydro-corticosterone
3α,20α,21-Trihydroxy-5β-pregnane-11-one
3α,21-Dihydroxy-5β-pregnane-11,20-dione
5β-Pregnane-3,20-dione
3α-Hydroxy-5β-pregnane-20-one
Pregnanediol
5α-Pregnane-3,20-dione
3α-Hydroxy-5α-pregnan-20-one
5α-Pregnane-3α,20α-diol
5α-Pregnan-20α-ol-3-one
4-Androsten-11beta-ol-3,17-dione
11β-Hydroxytestosterone
Urocortisol
Cortol
11β,17α,21-Trihydroxy-5β-pregnane-3,20-dione
17α,21-Dihydroxy-5β-pregnane-3,11,20-trione
Cortolone
Urocortisone
Estrone 3-sulfate
Estrone glucuronide
Estradiol-17α
2-Methoxyestrone-3-glucuronide
2-Methoxyestrone
2-Hydroxyestrone
2-Methoxyestrone-3-sulfate
Estrone
16-α-Hydroxyestrone
Estriol
16-Glucuronide-estriol
2-Methoxy-estradiol-17β-3-glucuronide
2-Methoxy-estradiol-17β
2-Hydroxy-estradiol-17β
2-Methoxy-estradiol-17β-3-sulfate
6β-Hydroxy-estradiol-17β
Estradiol-17β
Estradiol-17β-3-glucuronide
Estradiol-17β-3-sulfate
5β-Dihydro-testosterone
Testosterone glucuronide
C21-Steroids
C18-Steroids
C19-Steroids
Data on KEGG graph
Rendered by Pathview





gene oncology we can do more analysis

```
data(go.sets.hs)
data(go.subs.hs)

# Focus on Biological Process subset of GO
gobpsets = go.sets.hs[go.subs.hs$BP]

gobpres = gage(foldchanges, gsets=gobpsets, same.dir=TRUE)

lapply(gobpres, head)
```

```
$greater
                                           p.geomean stat.mean        p.val
GO:0007156 homophilic cell adhesion       8.519724e-05  3.824205 8.519724e-05
GO:0002009 morphogenesis of an epithelium 1.396681e-04  3.653886 1.396681e-04
GO:0048729 tissue morphogenesis           1.432451e-04  3.643242 1.432451e-04
GO:0007610 behavior                       1.925222e-04  3.565432 1.925222e-04
GO:0060562 epithelial tube morphogenesis  5.932837e-04  3.261376 5.932837e-04
GO:0035295 tube development               5.953254e-04  3.253665 5.953254e-04
                                               q.val set.size         exp1
GO:0007156 homophilic cell adhesion       0.1952430      113 8.519724e-05
GO:0002009 morphogenesis of an epithelium 0.1952430      339 1.396681e-04
GO:0048729 tissue morphogenesis           0.1952430      424 1.432451e-04
GO:0007610 behavior                       0.1968058      426 1.925222e-04
GO:0060562 epithelial tube morphogenesis  0.3566193      257 5.932837e-04
GO:0035295 tube development               0.3566193      391 5.953254e-04

$less
                                             p.geomean stat.mean        p.val
GO:0048285 organelle fission              1.536227e-15 -8.063910 1.536227e-15
GO:0000280 nuclear division               4.286961e-15 -7.939217 4.286961e-15
GO:0007067 mitosis                        4.286961e-15 -7.939217 4.286961e-15
GO:0000087 M phase of mitotic cell cycle  1.169934e-14 -7.797496 1.169934e-14
GO:0007059 chromosome segregation         2.028624e-11 -6.878340 2.028624e-11
GO:0000236 mitotic prometaphase           1.729553e-10 -6.695966 1.729553e-10
                                                 q.val set.size         exp1
GO:0048285 organelle fission              5.843127e-12      376 1.536227e-15
GO:0000280 nuclear division               5.843127e-12      352 4.286961e-15
GO:0007067 mitosis                        5.843127e-12      352 4.286961e-15
GO:0000087 M phase of mitotic cell cycle  1.195965e-11      362 1.169934e-14
GO:0007059 chromosome segregation         1.659009e-08      142 2.028624e-11
GO:0000236 mitotic prometaphase           1.178690e-07       84 1.729553e-10
```

```
$stats
                                          stat.mean     exp1
GO:0007156 homophilic cell adhesion        3.824205 3.824205
GO:0002009 morphogenesis of an epithelium  3.653886 3.653886
GO:0048729 tissue morphogenesis            3.643242 3.643242
GO:0007610 behavior                        3.565432 3.565432
GO:0060562 epithelial tube morphogenesis   3.261376 3.261376
GO:0035295 tube development                3.253665 3.253665
```

```
sig_genes <- res[res$padj <= 0.05 & !is.na(res$padj), "symbol"]
print(paste("Total number of significant genes:", length(sig_genes)))
```

```
[1] "Total number of significant genes: 8147"
```

```
write.table(sig_genes, file="significant_genes.txt", row.names=FALSE, col.names=FALSE, quo
```

#perform pathway analysis online on significant_gene.txt go to the Reactome website (https://reactome.org/PathwayBrowser/#TOOL=AT). Q: What pathway has the most significant "Entities p-value"? Do the most significant pathways listed match your previous KEGG results? What factors could cause differences between the two methods?

Cell cycle, mitotic has the most significant "Entities p-value", which matches with my previous KEGG results.