

Week 13 Tutorial

Course Review

Joe Scott / Jan Gorzny



Prepared based off of the notes of CS245 Instructors, past and present.

31 March 2017

Plan

1 Course Review

- Hoare Triples
- Conversion to CNF
- Resolution
- Natural Deduction
- Tautological Implication
- Key Theorems

2 The End

1 Course Review

- Hoare Triples
- Conversion to CNF
- Resolution
- Natural Deduction
- Tautological Implication
- Key Theorems

2 The End

Problem 1

- 1 $\llbracket (((a = a_0) \wedge (a \geq 0)) \wedge (b > 0)) \rrbracket$
- 2 while $(a - b \geq 0)$ {
- 3 $a = a - b;$
- 4 }
- 5 $r = a;$
- 6 $\llbracket (\exists q((r = a - qb) \wedge (0 \leq r < b))) \rrbracket$

Problem 1

- 1 $\langle\langle\langle\langle a = a_0 \rangle \wedge (a \geq 0) \rangle \wedge (b > 0) \rangle\rangle\rangle$
- 2 while $(a - b \geq 0)$ {
- 3 $a = a - b;$
- 4 }
- 5 $r = a;$
- 6 $\langle\langle\langle\langle \exists q((r = a - qb) \wedge (0 \leq r < b)) \rangle\rangle\rangle\rangle$

Try $I := \langle a \geq 0 \rangle$

Problem 1 - cont.

- | | | |
|---|---|---------------|
| ① | $\llbracket (((a = a_0) \wedge (a \geq 0)) \wedge (b > 0)) \rrbracket$ | assumption |
| ② | $\llbracket (a \geq 0) \rrbracket$ | Implied(a) |
| ③ | while $(a - b \geq 0)$ { | |
| ④ | $\llbracket ((a \geq 0) \wedge (a - b \geq 0)) \rrbracket$ | partial-while |
| ⑤ | $\llbracket (a - b \geq 0) \rrbracket$ | implied(b) |
| ⑥ | $a = a - b;$ | |
| ⑦ | $\llbracket a \geq 0 \rrbracket$ | assignment |
| ⑧ | } | |
| ⑨ | $\llbracket ((a \geq 0) \wedge (\neg(a - b \geq 0))) \rrbracket$ | partial while |
| ⑩ | $\llbracket (\exists q((a = a - qb) \wedge (0 \geq a < b))) \rrbracket$ | implied(c) |
| ⑪ | $r = a;$ | |
| ⑫ | $\llbracket (\exists q((r = a - qb) \wedge (0 \leq r < b))) \rrbracket$ | assignment |

Problem 1 - impliesds

For implied(a) we want to show that

$$((((a = a_0) \wedge (a \geq 0)) \wedge (b > 0)) \rightarrow (a \geq 0))$$

This is obvious from \wedge_e .
for implied(b)

$$(((a \geq 0) \wedge (a - b \geq 0)) \rightarrow (a - b \geq 0))$$

This is obvious from \wedge_e .

Problem 1 - cont.

- | | | |
|---|---|---------------|
| ① | $\llbracket (((a = a_0) \wedge (a \geq 0)) \wedge (b > 0)) \rrbracket$ | assumption |
| ② | $\llbracket (a \geq 0) \rrbracket$ | Implied(a) |
| ③ | while $(a - b \geq 0)$ { | |
| ④ | $\llbracket ((a \geq 0) \wedge (a - b \geq 0)) \rrbracket$ | partial-while |
| ⑤ | $\llbracket (a - b \geq 0) \rrbracket$ | implied(b) |
| ⑥ | $a = a - b;$ | |
| ⑦ | $\llbracket a \geq 0 \rrbracket$ | assignment |
| ⑧ | } | |
| ⑨ | $\llbracket ((a \geq 0) \wedge (\neg(a - b \geq 0))) \rrbracket$ | partial while |
| ⑩ | $\llbracket (\exists q((a = a - qb) \wedge (0 \geq a < b))) \rrbracket$ | implied(c) |
| ⑪ | $r = a;$ | |
| ⑫ | $\llbracket (\exists q((r = a - qb) \wedge (0 \leq r < b))) \rrbracket$ | assignment |

Implied c says

$$(((a \geq 0) \wedge (\neg(a - b \geq 0)) \rightarrow (\exists q((a = a - qb) \wedge (0 \leq a < b))))))$$

Firstly

$$(\neg(a - b \geq 0))$$

$$\iff a - b < 0$$

$$\iff a < b$$

As $0 \leq a$, $0 \leq a < b$. Clearly then given $q = 0$, $a = a - qb$, as needed.

Termination

We need to find a loop variant.

- ① $\langle\langle((a = a_0) \wedge (a \geq 0)) \wedge (b > 0))\rangle\rangle$
- ② while $(a - b \geq 0)$ {
- ③ $a = a - b;$
- ④ }
- ⑤ $r = a;$
- ⑥ $\langle\langle(\exists q((r = a - qb) \wedge (0 \leq r < b)))\rangle\rangle$

Termination

We need to find a loop variant.

- ① $\langle(((a = a_0) \wedge (a \geq 0)) \wedge (b > 0))\rangle$
- ② while $(a - b \geq 0)$ {
- ③ $a = a - b;$
- ④ }
- ⑤ $r = a;$
- ⑥ $\langle(\exists q((r = a - qb) \wedge (0 \leq r < b)))\rangle$

Consider the following $v = \lfloor \frac{a}{b} \rfloor - (\frac{a_0 - a}{b})$.

On initialization $v = \lfloor \frac{a}{b} \rfloor$ as $a = a_0$.

For each iteration, $\frac{a_0 - a}{b}$ takes the values 1, 2, 3, 4.. as required. The loop terminates on $v = 0$.

1 Course Review

- Hoare Triples
- **Conversion to CNF**
- Resolution
- Natural Deduction
- Tautological Implication
- Key Theorems

2 The End

CNF Conversion

Convert the following to CNF:

$$p \leftrightarrow (r \wedge s)$$

CNF Conversion

Convert the following to CNF:

$$p \leftrightarrow (r \wedge s)$$

$$\iff (p \rightarrow (r \wedge s)) \wedge ((r \wedge s) \rightarrow p) \leftrightarrow\text{-equiv}$$

CNF Conversion

Convert the following to CNF:

$$p \leftrightarrow (r \wedge s)$$

$$\iff (p \rightarrow (r \wedge s)) \wedge ((r \wedge s) \rightarrow p) \leftrightarrow\text{-equiv}$$

$$\iff (\neg p \vee (r \wedge s)) \wedge (\neg(r \wedge s) \vee p) \rightarrow\text{-equiv}$$

CNF Conversion

Convert the following to CNF:

$$p \leftrightarrow (r \wedge s)$$

$$\iff (p \rightarrow (r \wedge s)) \wedge ((r \wedge s) \rightarrow p) \leftrightarrow\text{-equiv}$$

$$\iff (\neg p \vee (r \wedge s)) \wedge (\neg(r \wedge s) \vee p) \rightarrow\text{-equiv}$$

$$\iff (\neg p \vee r) \wedge (\neg p \vee s) \wedge (\neg r \vee \neg s \vee p) \text{ Distributivity, DeMorgan's}$$

1 Course Review

- Hoare Triples
- Conversion to CNF
- **Resolution**
- Natural Deduction
- Tautological Implication
- Key Theorems

2 The End

Problem

Does $\{a, b, c, d\} \vdash_{\text{Res}} a \wedge b$?

Problem

Does $\{a, b, c, d\} \vdash_{Res} a \wedge b$?

Step 1: Negate conclusion: $a \wedge B$ becomes $\neg(a \wedge b)$.

Problem

Does $\{a, b, c, d\} \vdash_{\text{Res}} a \wedge b$?

Step 1: Negate conclusion: $a \wedge B$ becomes $\neg(a \wedge b)$.

Step 2: Ensure everything is CNF. Need to convert negated conclusion:
 $\neg(a \wedge b) \iff \neg a \vee \neg b$ by DeMorgan's

Problem

Does $\{a, b, c, d\} \vdash_{Res} a \wedge b$?

Step 1: Negate conclusion: $a \wedge B$ becomes $\neg(a \wedge b)$.

Step 2: Ensure everything is CNF. Need to convert negated conclusion:

$\neg(a \wedge b) \iff \neg a \vee \neg b$ by DeMorgan's

Step 3: Resolution

- a premise
- b premise
- c premise
- d premise
- $\neg a \vee \neg b$ negated conclusion
- $\neg b$ resolution 1,5
- \perp resolution 2,6

Done - the conclusion is implied by the premise.

1 Course Review

- Hoare Triples
- Conversion to CNF
- Resolution
- **Natural Deduction**
- Tautological Implication
- Key Theorems

2 The End

Example

Problem

Prove $\neg(A \wedge B) \vdash A \rightarrow \neg B$

Example

Problem

Prove $\neg(A \wedge B) \vdash A \rightarrow \neg B$

- ① $A, B \vdash A$ by ϵ
- ② $A, B \vdash B$ by ϵ
- ③ $A, B \vdash A \wedge B$ by $\wedge +$ 1,2
- ④ $\neg(A \wedge B), A, B \vdash A \wedge B$ by $(+)$ 3
- ⑤ $\neg(A \wedge B)A, B \vdash \neg(A \wedge B)$ by ϵ
- ⑥ $\neg(A \wedge B), A \vdash \neg B$ by $\neg +$ 4,5
- ⑦ $\neg(A \wedge B) \vdash A \rightarrow \neg B$ by $\rightarrow +$ 6

1 Course Review

- Hoare Triples
- Conversion to CNF
- Resolution
- Natural Deduction
- **Tautological Implication**
- Key Theorems

2 The End

Semantic Entailment

Do the premises semantically entail (logically imply) the conclusion?
Answer this question using a truth table. Explain.

$$\{(\neg p \wedge (q \vee \neg r))\} \models \neg r \vee p$$

Semantic Entailment

Do the premises semantically entail (logically imply) the conclusion?
Answer this question using a truth table. Explain.

$$\{(\neg p \wedge (q \vee \neg r))\} \models \neg r \vee p$$

| p | q | r | $\neg p \wedge (q \vee \neg r)$ | $\neg r \vee p$ |
|-----|-----|-----|---------------------------------|-----------------|
| 1 | 1 | 1 | | |
| 1 | 1 | 0 | | |
| 1 | 0 | 1 | | |
| 1 | 0 | 0 | | |
| 0 | 1 | 1 | | |
| 0 | 1 | 0 | | |
| 0 | 0 | 1 | | |
| 0 | 0 | 0 | | |

Semantic Entailment

Do the premises semantically entail (logically imply) the conclusion?
Answer this question using a truth table. Explain.

$$\{(\neg p \wedge (q \vee \neg r))\} \models \neg r \vee p$$

| p | q | r | $\neg p \wedge (q \vee \neg r)$ | $\neg r \vee p$ |
|-----|-----|-----|---------------------------------|-----------------|
| 1 | 1 | 1 | 0 | |
| 1 | 1 | 0 | 0 | |
| 1 | 0 | 1 | 0 | |
| 1 | 0 | 0 | 0 | |
| 0 | 1 | 1 | 1 | |
| 0 | 1 | 0 | 1 | |
| 0 | 0 | 1 | 0 | |
| 0 | 0 | 0 | 1 | |

Semantic Entailment

Do the premises semantically entail (logically imply) the conclusion?
Answer this question using a truth table. Explain.

$$\{(\neg p \wedge (q \vee \neg r))\} \models \neg r \vee p$$

| p | q | r | $\neg p \wedge (q \vee \neg r)$ | $\neg r \vee p$ |
|-----|-----|-----|---------------------------------|-----------------|
| 1 | 1 | 1 | 0 | 1 |
| 1 | 1 | 0 | 0 | 1 |
| 1 | 0 | 1 | 0 | 1 |
| 1 | 0 | 0 | 0 | 1 |
| 0 | 1 | 1 | 1 | 0 |
| 0 | 1 | 0 | 1 | 1 |
| 0 | 0 | 1 | 0 | 0 |
| 0 | 0 | 0 | 1 | 1 |

1 Course Review

- Hoare Triples
- Conversion to CNF
- Resolution
- Natural Deduction
- Tautological Implication
- Key Theorems

2 The End

Definition

A sentence ψ is true in an interpretation \mathcal{I} , denoted $\mathcal{I} \models \psi$, if for every possible sequence of elements in the interpretation, substituting these elements into the variables present in ψ yields a true sentence. Such an interpretation \mathcal{I} is called a satisfying interpretation.

Theorem (Godel's Completeness Theorem)

Let Σ be a set of formulas. If Σ is consistent, then it has a satisfying interpretation.

Theorem (Godel's Completeness Theorem)

Let Σ be a set of formulas. If Σ is consistent, then it has a satisfying interpretation.

Theorem (Godel's Completeness Theorem - Contrapositive)

If Σ does not have a satisfying interpretation, then Σ is not consistent.

Key Theorems

Theorem (Compactness Theorem)

A set of sentences Σ has a satisfying interpretation if and only if every finite set of Σ has a satisfying interpretation.

Theorem

Let A be a sentence of first-order logic such that for any $n \in \mathbb{N}$, $n \geq 1$, there is a domain D and valuation v with at least n elements in D , such that $A^v = 1$. Then A has a domain D' and a valuation v' such that $A^{v'} = 1$ and D' has an infinite number of elements.

Plan

1 Course Review

- Hoare Triples
- Conversion to CNF
- Resolution
- Natural Deduction
- Tautological Implication
- Key Theorems

2 The End

The end

That's it folks. Feel free to hang out and ask questions.

These slides are based off of the tutorial notes and lecture slides provided to you online.

If you want a copy feel free to email me. The are also available on my personal website joe-scott.net

| IA Email: | IA | email |
|-----------|-------------------------|---------------------|
| | Jan Gorzny Joe Scott | jgorzny j29scott |

Jan and Joe have an office hour **Mondays** at **3pm** in the Tutorial Center in MC.

Instructor Office Hours:

| Instructor | Time | Room | Email |
|------------|------------------|----------|---------------|
| Trefler | Tue, Thur 4:00pm | DC 2336 | trefler |
| Rahkooy | Tue, Thur 4:00pm | DC 2302B | hamid.rahkooy |