# Exercise Paper 2

Francesco Palandra, Stefano Ramagnano

May 27, 2023

### 1 Power series

**Exercise .1.** Given the following series, find:

- the center of the serie  $x_0$ ;
- the convergence radius R;
- the convergence set E.

$$\sum_{n=1}^{\infty} \log(\frac{1}{n^2+1}+1), \quad \sum_{n=1}^{\infty} \frac{9^n-8^n}{(3n^2+1)5^n} (4x+1)^n, \quad \sum_{n=1}^{\infty} (-1)^n \frac{9^n-5^n}{(4n+7)8^n} (x-\frac{1}{2})$$

$$\sum_{n=1}^{\infty} \frac{e^{\frac{1}{n}}-1}{3log(n+1)} (x-5)^n, \quad \sum_{n=1}^{\infty} \frac{n3^n+1}{2^{2n+1}-9} (7x-e)^n, \quad \sum_{n=1}^{\infty} \frac{n!(2^n+5)}{(n+3)!} (x+2)^n.$$

$$\sum_{n=1}^{\infty} \frac{(3n)^n-2n!}{2^{n+1}n^{n-1}+5} (x+\frac{1}{2})^n, \quad \sum_{n=1}^{\infty} \frac{17^n}{14^n(n+4)} (x+6)^n, \quad \sum_{n=1}^{\infty} \frac{7^n}{17^n(n+8)} (x-12)^n.$$

#### 2 Numerical series

Exercise .2. Given the following numerical series, analyse their convergence:

$$\begin{split} &\sum_{n=1}^{\infty} \sqrt[7]{\frac{n^6+n^3}{n^15+1}}, \quad \sum_{n=1}^{\infty} \sqrt[5]{\frac{n^4+n^3}{n^9+1}}, \quad \sum_{n=1}^{\infty} n^3 (\cos(\frac{1}{n})-1)^2 \\ &\sum_{n=1}^{\infty} \frac{2^n}{3^n} (n^2+\sin(e^n)), \quad \sum_{n=1}^{\infty} \sin(\frac{n^2+5}{n^4+3n^3-6n+2}), \quad \sum_{n=1}^{\infty} \sqrt{\frac{n^2+7n+13}{(n+4)^6}} \end{split}$$

#### 3 ODE first order

**Exercise .3.** Given the following ODEs (or Cauchy's problems), find the solution: (Note that if you have two different starting values you have to solve two different problems)

$$y' = y^2 - 2y - 3$$
,  $y(0) = 0$ ;  $xy' = -y + 4x^2$ ,  $y(0) = 0$ ,  $y(0) = 3$ ;  $y' = y + x$ ;  $y' = x \ln(x^2 + 1)(1 + y^2)$ ,  $y(0) = 2$ ;  $y' = \frac{x^3 + y^3}{xy^2}$ ;  $y' = \frac{1}{1 + x^2}$ ,  $y(0) = 4$ .

## 4 ODE second order

Exercise .4. Given the following ODEs or Cauchy's problems find a set of solutions or the exact one:

$$y'' + 3y = t - 3, y(0) = y'(0) = 0; \quad y'' + 2y' + y = \sin(2t), y(0) = \frac{1}{2}, y'(0) = 1;$$

$$y'' - 2y' - 3y = e^{4t}, y(0) = y'(0) = 0y'' + 2y' + 2y = \sin(t) + \cos(t);$$

$$y'' + 4y = t - \cos(3t); \quad y'' + 4y = t + \cos(t), y(0) = 0, y(0) = 1;$$

$$y'' - 2y' + y = 6te^{7t}, y(0) = y'(0) = 0.$$