Module 3: Resampling and Simulation-Based Statistical Inference

MA 563 – Statistical Computation and Simulation

Dr. Rishikesh Yadav

rishikesh@iitmandi.ac.in — TA: Vedant Vibhor (D24028@students.iitmandi.ac.in)
School of Mathematics and Statistical Sciences, IIT Mandi

$\Rightarrow$ $\mu, \sigma^2 \rightarrow F \rightarrow X_1 \cdots X_n \Rightarrow$ sample

$\mu = \rightarrow \hat{\theta}_1 = \bar{X} \rightarrow$ $\hookrightarrow se(\bar{X}) \Rightarrow (\bar{X} \pm 2\, se(\bar{X}))$

$\hat{\theta}_2 = \text{med}(X_1 \cdots X_n)$

$\hat{\theta}_3 = \text{mod}(X_1 \cdots X_n)$

↳ First way → find analytical sampling dist^n of estimator $\hat{\theta}$

↳ → $X_1^1 \cdots X_n^1 \rightarrow \hat{\theta}_1^1 \rightarrow \bar{x}_1$

we don't know this → $X_1^2 \cdots X_n^2 \rightarrow \hat{\theta}^2 \rightarrow \bar{x}_2$ → dist^n → Histogram

$X_1^N \cdots X_n^N \rightarrow \hat{\theta}^N \rightarrow \bar{x}_N$

sampling dist^n of $\bar{X}$, $\hat{\theta}$ → $\hat{\theta}$

Assumption $X_1 \cdots X_n \sim N(\mu, \sigma^2) \Rightarrow \mu, \sigma^2$

$\hat{\theta}_1 = \bar{X}_n \sim N(\mu, \frac{\sigma^2}{n})$

Sampling dist^n of $\bar{X}_n$

↳ $\bar{X}$, $\frac{S^2}{n}$, $S^2 = \frac{1}{n}\sum_{i=1}^{n}(X_i - \bar{X})^2$

↳ estimate $\bar{X}$

95% CI: $\bar{X} \pm 2S/\sqrt{n} \Rightarrow \frac{1}{n}\sum_{i=1}^{n} X_i + 2\frac{\sqrt{\frac{1}{n}\sum_{i=1}^{n}(X_i - \bar{X})^2}}{n}$

$\sigma^2 < \infty$

$X_1 \cdots X_n \sim F(\mu, \sigma^2)$.

$\Rightarrow \bar{X}_n \sim N(\mu, \sigma^2/n) \Rightarrow n \rightarrow \infty$

→ Sample median → , $X_1 - - X_n \sim F(\mu, \sigma^2)$

$$\hat{\theta}_2 = \text{med}(X_1 - - X_n)$$

$$\hat{\theta}_2 \sim N\left[\mu, \frac{1}{4[f'(\mu)]^2 n}\right]$$

f : pdf

→ sample mode → $\hat{\theta}_3 = \text{mod}(X_1 - - X_n)$

$\hat{\theta}_3 \sim$?

Table of Contents

1. Motivation: Inference When Formulas Run Out
   - Every Estimate Needs an Error Bar
   - Real Problems with No Standard-Error Formula
   - The Resampling Idea
2. The Bootstrap Principle
   - Empirical Distribution and the Plug-in Principle
   - The Nonparametric Bootstrap Algorithm
   - A Toy Example by Hand
   - Bootstrap Standard Error and Bias
   - Why It Works and How Many Resamples
3. Parametric vs Nonparametric Bootstrap
   - The Parametric Bootstrap
   - Example: Exponential Lifetimes
   - Bootstrapping Regression Models
   - Dependent Data: The Block Bootstrap
   - Choosing Between Them
4. Bootstrap Confidence Intervals
   - Normal and Percentile Intervals
   - The Basic (Pivotal) Interval
   - The Bootstrap-t Interval
   - The BCa Interval
   - Worked Example and Coverage Study
5. The Jackknife (Brief)
   - Leave-One-Out Estimates
   - Jackknife Bias and Standard Error
   - Limitations and Relation to the Bootstrap
6. Cross-Validation
   - Training Error Is Optimistic
   - Validation Set, LOOCV and K-Fold
   - Choosing K: The Bias–Variance Trade-off
   - Cross-Validation for Model Selection
   - Common Pitfalls
7. Monte Carlo Hypothesis Testing and Permutation Tests
   - Monte Carlo p-Values
   - Parametric Bootstrap Tests and Simulated Power
   - Permutation Tests
   - Example: A Two-Sample Permutation Test
   - Exchangeability: Permutation vs Bootstrap Tests
8. Likelihood-Based Inference via Simulation
   - The Sampling Distribution of the MLE by Simulation
   - Parametric Bootstrap Likelihood-Ratio Tests
   - When the Likelihood Itself Needs Simulation
   - Designing a Simulation Study
9. Simulation vs Asymptotic Inference
   - Two Routes to a Sampling Distribution
   - When the Bootstrap Fails
   - Computational Trade-offs
   - A Practical Decision Guide
10. Unit Summary

Module 3 at a Glance

Module 3: Resampling and Simulation-Based Statistical Inference (10 Lectures)

Bootstrap principles, parametric vs nonparametric bootstrap, cross-validation, jackknife (brief), simulation-based uncertainty quantification, computational trade-offs, likelihood-based inference via simulation, parametric simulation for inference, confidence intervals via simulation, hypothesis testing using Monte Carlo, permutation tests, bootstrap-based inference (deep dive), simulation vs asymptotic inference.

Module 3 at a Glance

Module 3: Resampling and Simulation-Based Statistical Inference (10 Lectures)

Bootstrap principles, parametric vs nonparametric bootstrap, cross-validation, jackknife (brief), simulation-based uncertainty quantification, computational trade-offs, likelihood-based inference via simulation, parametric simulation for inference, confidence intervals via simulation, hypothesis testing using Monte Carlo, permutation tests, bootstrap-based inference (deep dive), simulation vs asymptotic inference.

How Module 3 builds on Modules 1 and 2

- **Module 1** gave us generators: we can draw from any F we can write down.
- **Module 2** gave us Monte Carlo: any expectation can be approximated by an average.
- **Module 3** applies both to *statistical inference itself*: standard errors, confidence intervals, p-values and prediction error – computed by simulation instead of by formula.

Outline

1. Motivation: Inference When Formulas Run Out
   - Every Estimate Needs an Error Bar
   - Real Problems with No Standard-Error Formula
   - The Resampling Idea
2. The Bootstrap Principle
3. Parametric vs Nonparametric Bootstrap
4. Bootstrap Confidence Intervals
5. The Jackknife (Brief)
6. Cross-Validation
7. Monte Carlo Hypothesis Testing and Permutation Tests
8. Likelihood-Based Inference via Simulation
9. Simulation vs Asymptotic Inference
10. Unit Summary

Every Estimate Needs an Error Bar

- We observe data $x_1, \ldots, x_n \overset{\text{iid}}{\sim} F$ and compute a statistic $\hat{\theta} = s(x_1, \ldots, x_n)$ to estimate a parameter $\theta = T(F)$.

Every Estimate Needs an Error Bar

- We observe data $x_1, \ldots, x_n \overset{\text{iid}}{\sim} F$ and compute a statistic $\hat{\theta} = s(x_1, \ldots, x_n)$ to estimate a parameter $\theta = T(F)$.
- A point estimate alone is never enough. We also need:

Every Estimate Needs an Error Bar

- We observe data $x_1, \ldots, x_n \overset{\text{iid}}{\sim} F$ and compute a statistic $\hat{\theta} = s(x_1, \ldots, x_n)$ to estimate a parameter $\theta = T(F)$.
- A point estimate alone is never enough. We also need:
  - its **standard error** $\text{se}_F(\hat{\theta}) = \sqrt{\text{Var}_F(\hat{\theta})}$,

Every Estimate Needs an Error Bar

- We observe data $x_1, \ldots, x_n \overset{\text{iid}}{\sim} F$ and compute a statistic $\hat{\theta} = s(x_1, \ldots, x_n)$ to estimate a parameter $\theta = T(F)$.
- A point estimate alone is never enough. We also need:
  - its **standard error** $\text{se}_F(\hat{\theta}) = \sqrt{\text{Var}_F(\hat{\theta})}$,
  - its **bias** $\mathbb{E}_F[\hat{\theta}] - \theta$,

Every Estimate Needs an Error Bar

- We observe data $x_1, \ldots, x_n \overset{\text{iid}}{\sim} F$ and compute a statistic $\hat{\theta} = s(x_1, \ldots, x_n)$ to estimate a parameter $\theta = T(F)$.
- A point estimate alone is never enough. We also need:
  - its **standard error** $\text{se}_F(\hat{\theta}) = \sqrt{\text{Var}_F(\hat{\theta})}$,
  - its **bias** $\mathbb{E}_F[\hat{\theta}] - \theta$,
  - a **confidence interval** for θ, and/or a ***p*-value** for a hypothesis about θ.

Every Estimate Needs an Error Bar

- We observe data $x_1, \ldots, x_n \overset{\text{iid}}{\sim} F$ and compute a statistic $\hat{\theta} = s(x_1, \ldots, x_n)$ to estimate a parameter $\theta = T(F)$.
- A point estimate alone is never enough. We also need:
  - its **standard error** $\text{se}_F(\hat{\theta}) = \sqrt{\text{Var}_F(\hat{\theta})}$,
  - its **bias** $\mathbb{E}_F[\hat{\theta}] - \theta$,
  - a **confidence interval** for θ, and/or a ***p*-value** for a hypothesis about θ.
- All four are properties of the **sampling distribution** of $\hat{\theta}$ – the distribution of $\hat{\theta}$ over hypothetical repetitions of the experiment.

Every Estimate Needs an Error Bar

- We observe data $x_1, \ldots, x_n \overset{\text{iid}}{\sim} F$ and compute a statistic $\hat{\theta} = s(x_1, \ldots, x_n)$ to estimate a parameter $\theta = T(F)$.
- A point estimate alone is never enough. We also need:
  - its **standard error** $\text{se}_F(\hat{\theta}) = \sqrt{\text{Var}_F(\hat{\theta})}$,
  - its **bias** $\mathbb{E}_F[\hat{\theta}] - \theta$,
  - a **confidence interval** for θ, and/or a ***p*-value** for a hypothesis about θ.
- All four are properties of the **sampling distribution** of $\hat{\theta}$ – the distribution of $\hat{\theta}$ over hypothetical repetitions of the experiment.

The central question of Module 3

The sampling distribution depends on the *unknown* F, and we see only *one* sample. Can we approximate it anyway – for *any* statistic, without deriving a formula?

The Textbook Case – and Why It Is Special

- For the sample mean $\bar{X}$, classical theory delivers everything:

$$\mathrm{se}(\bar{X}) = \frac{\sigma}{\sqrt{n}} \approx \frac{s}{\sqrt{n}}, \qquad \bar{X} \pm t_{n-1,\,\alpha/2}\,\frac{s}{\sqrt{n}}.$$

The Textbook Case – and Why It Is Special

- For the sample mean $\bar{X}$, classical theory delivers everything:

$$\mathrm{se}(\bar{X}) = \frac{\sigma}{\sqrt{n}} \approx \frac{s}{\sqrt{n}}, \qquad \bar{X} \pm t_{n-1,\,\alpha/2}\,\frac{s}{\sqrt{n}}.$$

- This works because $\bar{X}$ is **linear** in the data and, for Normal data, its distribution is known *exactly*; otherwise the CLT gives a good approximation.

The Textbook Case – and Why It Is Special

- For the sample mean $\bar{X}$, classical theory delivers everything:

$$\mathrm{se}(\bar{X}) = \frac{\sigma}{\sqrt{n}} \approx \frac{s}{\sqrt{n}}, \qquad \bar{X} \pm t_{n-1,\,\alpha/2}\,\frac{s}{\sqrt{n}}.$$

- This works because $\bar{X}$ is **linear** in the data and, for Normal data, its distribution is known *exactly*; otherwise the CLT gives a good approximation.
- Change the statistic, and the comfort disappears:

$\theta_{\text{med}} = \text{med}(X_1, \ldots, X_n)$

The Textbook Case – and Why It Is Special

- For the sample mean $\bar{X}$, classical theory delivers everything:

$$\mathrm{se}(\bar{X}) = \frac{\sigma}{\sqrt{n}} \approx \frac{s}{\sqrt{n}}, \qquad \bar{X} \pm t_{n-1,\,\alpha/2}\,\frac{s}{\sqrt{n}}.$$

- This works because $\bar{X}$ is **linear** in the data and, for Normal data, its distribution is known *exactly*; otherwise the CLT gives a good approximation.
- Change the statistic, and the comfort disappears:
  - **Median:** $\mathrm{se} \approx 1/\{2f(m)\sqrt{n}\}$ – needs the unknown density f at the unknown median m.

The Textbook Case – and Why It Is Special

- For the sample mean $\bar{X}$, classical theory delivers everything:

$$\mathrm{se}(\bar{X}) = \frac{\sigma}{\sqrt{n}} \approx \frac{s}{\sqrt{n}}, \qquad \bar{X} \pm t_{n-1,\,\alpha/2}\,\frac{s}{\sqrt{n}}.$$

- This works because $\bar{X}$ is **linear** in the data and, for Normal data, its distribution is known *exactly*; otherwise the CLT gives a good approximation.
- Change the statistic, and the comfort disappears:
  - **Median:** se $\approx 1/\{2f(m)\sqrt{n}\}$ – needs the unknown density f at the unknown median m.
  - **Correlation r:** the classical formula assumes bivariate Normality.

The Textbook Case – and Why It Is Special

- For the sample mean $\bar{X}$, classical theory delivers everything:

$$\mathrm{se}(\bar{X}) = \frac{\sigma}{\sqrt{n}} \approx \frac{s}{\sqrt{n}}, \qquad \bar{X} \pm t_{n-1,\,\alpha/2}\,\frac{s}{\sqrt{n}}.$$

- This works because $\bar{X}$ is **linear** in the data and, for Normal data, its distribution is known *exactly*; otherwise the CLT gives a good approximation.
- Change the statistic, and the comfort disappears:
  - **Median:** se $\approx 1/\{2f(m)\sqrt{n}\}$ – needs the unknown density f at the unknown median m.
  - **Correlation** r**:** the classical formula assumes bivariate Normality.
  - **Ratio, trimmed mean, eigenvalue, a classifier's error rate:** typically no usable formula at all.

The Textbook Case – and Why It Is Special

- For the sample mean $\bar{X}$, classical theory delivers everything:

$$\mathrm{se}(\bar{X}) = \frac{\sigma}{\sqrt{n}} \approx \frac{s}{\sqrt{n}}, \qquad \bar{X} \pm t_{n-1,\,\alpha/2}\,\frac{s}{\sqrt{n}}.$$

- This works because $\bar{X}$ is **linear** in the data and, for Normal data, its distribution is known *exactly*; otherwise the CLT gives a good approximation.
- Change the statistic, and the comfort disappears:
  - **Median:** se $\approx 1/\{2f(m)\sqrt{n}\}$ – needs the unknown density f at the unknown median m.
  - **Correlation r:** the classical formula assumes bivariate Normality.
  - **Ratio, trimmed mean, eigenvalue, a classifier's error rate:** typically no usable formula at all.

The honest picture

Closed-form standard errors exist for a small, special family of statistics. Real analyses use statistics that were chosen for their *scientific* meaning, not for having a tidy variance formula.

Real Problems with No Standard-Error Formula

Medicine: median survival

A small trial ($n = 25$) reports *median* time to recovery. Patients are expensive; the data are skewed. How uncertain is the median?

$\text{med}(X_1, \ldots, X_n)$ = median time of recovery

= 5 days

$\Rightarrow$ (4, 6) $\rightarrow$ 95% CI

Real Problems with No Standard-Error Formula

Medicine: median survival

A small trial ($n = 25$) reports *median* time to recovery. Patients are expensive; the data are skewed. How uncertain is the median?

Finance: the Sharpe ratio

Sharpe ratio $= \bar{r}/s_r$ of a fund's monthly returns – a *ratio* of two random quantities, from heavy-tailed returns. What is its standard error?

Real Problems with No Standard-Error Formula

Medicine: median survival

A small trial ($n = 25$) reports *median* time to recovery. Patients are expensive; the data are skewed. How uncertain is the median?

Finance: the Sharpe ratio

Sharpe ratio $= \bar{r}/s_r$ of a fund's monthly returns – a *ratio* of two random quantities, from heavy-tailed returns. What is its standard error?

Climate: a 100-year rainfall level

From 40 years of annual-maximum rainfall in the Himalaya, estimate the level exceeded once per century – a non-linear function of fitted extreme-value parameters. How wide is its interval?

Real Problems with No Standard-Error Formula

Medicine: median survival

A small trial ($n = 25$) reports *median* time to recovery. Patients are expensive; the data are skewed. How uncertain is the median?

Finance: the Sharpe ratio

Sharpe ratio $= \bar{r}/s_r$ of a fund's monthly returns – a *ratio* of two random quantities, from heavy-tailed returns. What is its standard error?

Climate: a 100-year rainfall level

From 40 years of annual-maximum rainfall in the Himalaya, estimate the level exceeded once per century – a non-linear function of fitted extreme-value parameters. How wide is its interval?

Machine learning: test error

A classifier trained on 500 patients. What error rate will it have on *new* patients – and which of ten candidate models generalizes best?

Real Problems with No Standard-Error Formula

Medicine: median survival

A small trial ($n = 25$) reports *median* time to recovery. Patients are expensive; the data are skewed. How uncertain is the median?

Finance: the Sharpe ratio

Sharpe ratio $= \bar{r}/s_r$ of a fund's monthly returns – a *ratio* of two random quantities, from heavy-tailed returns. What is its standard error?

Climate: a 100-year rainfall level

From 40 years of annual-maximum rainfall in the Himalaya, estimate the level exceeded once per century – a non-linear function of fitted extreme-value parameters. How wide is its interval?

Machine learning: test error


A classifier trained on 500 patients. What error rate will it have on *new* patients – and which of ten candidate models generalizes best?

The common thread

Each asks for a sampling distribution that classical theory cannot supply cleanly. Each is answered in practice by **resampling** or **simulation**.

What We Would Do If We Could Repeat the Experiment

- If we could draw *fresh* samples from F again and again, we would simply compute $\hat{\theta}$ on each and look at the histogram.



What We Would Do If We Could Repeat the Experiment


- If we could draw *fresh* samples from F again and again, we would simply compute $\hat{\theta}$ on each and look at the histogram.
- This is exactly Monte Carlo (Module 2), with $h(\cdot)$ replaced by the whole estimation procedure.

What We Would Do If We Could Repeat the Experiment

- If we could draw *fresh* samples from F again and again, we would simply compute $\hat{\theta}$ on each and look at the histogram.
- This is exactly Monte Carlo (Module 2), with $h(\cdot)$ replaced by the whole estimation procedure.
- The obstacle: F is unknown, and nature lets us draw from it only **once**.

What We Would Do If We Could Repeat the Experiment

- If we could draw *fresh* samples from F again and again, we would simply compute $\hat{\theta}$ on each and look at the histogram.
- This is exactly Monte Carlo (Module 2), with $h(\cdot)$ replaced by the whole estimation procedure.
- The obstacle: F is unknown, and nature lets us draw from it only **once**.



Parametric Bootstrap

Assumption on F → parametric case → F: $N(\mu, \sigma^2)$, Gamma(α, β) e.g.

Sample $(X_1 \dots X_n)$

estimate F

estimating F $\Leftrightarrow$ estimating unknown parameter

$\Leftrightarrow$ μ, σ^2 estimating for $F = N(\mu, \sigma^2)$

↓

$\hat{\mu}_{MLE}, \hat{\sigma}^2_{MLE}$

estimate F → $\hat{F}(\hat{\mu}_{MLE}, \hat{\sigma}^2_{MLE})$

↓ Exactly know what is F

use rnorm 1 → to simulate for Normal distribution with mean $\hat{\mu}_{MLE}$, $\hat{\sigma}^2_{MLE}$

→ Sample from $\hat{F}(\hat{\mu}, \hat{\sigma}^2)$

→ $X_1^1 \dots X_n^1 \to \hat{\theta}_1$

$X_1^2 \dots X_n^2 \to \hat{\theta}_2$ ⟷ Histogram of sampling distr of $\hat{\theta}$

$\vdots$

$X_1^B \dots X_n^B \to \hat{\theta}_B$

F: Non parametric

→ can't be written in term of finite no of parameter

estimate F. by empirical cdf →

$$P(X \le t) = \hat{F}_n(t) = \frac{1}{n} \cdot \sum_{i=1}^{n} I(X_i \le t), \quad X_1 \dots X_n \text{ sample}$$

↳ estimators

$$\Rightarrow \hat{F}_n(-\infty) = \frac{1}{n} \cdot \sum_{i=1}^{n} I(X_i \le -\infty) = 0$$

$$\hat{F}_n(\infty) = \frac{1}{n} \sum_{i=1}^{n} I(X_i \le \infty) = \frac{n}{n} = 1$$

$$\text{if } t_1 < t_2 \Rightarrow \hat{F}_n(t_1) \le \hat{F}_n(t_2)$$

$$\frac{1}{n} \sum_{i=1}^{n} I(X_i \le t_1) \le \frac{1}{n} \cdot \sum_{i=1}^{n} I(X_i \le t_2)$$

$$X_i \le t_1 \Rightarrow X_i \le t_2$$

$$\frac{\sum_{i=1}^{n} I(X_i \le t_1)}{n} \le \frac{\sum_{i=1}^{n} I(X_i \le t_1)}{n}$$

$$\hat{F}_n(t_1) \le \hat{F}_n(t_2)$$

non-parametric

Empirical CDF ⇒

$$\hat{F}_n(t) = \frac{1}{n} \cdot \sum_{i=1}^{n} I(X_i \le t)$$

↳ an estimate of F in case of non-parametric

Now we can simulate sample from $\hat{F}_n(t)$


$\Rightarrow$ $X_1^1 \; \cdots \; X_n^1 \sim \hat{F}_n(t) \longrightarrow \hat{\theta}_n^1$

$\vdots$

$X_1^B \; \cdots \; X_n^B \sim \hat{F}_n(t) \longrightarrow \hat{\theta}_n^B$

Histogram of it $\rightarrow$ sampling dist^n of $\hat{\theta}$.


The Resampling Idea: Let the Sample Stand In for the Population




Efron (1979): the bootstrap

Everything we cannot compute in the real world (because F is unknown) we **can** compute in the bootstrap world (because $\hat{F}_n$ is known) – by simulation.

The Resampling Idea: Let the Sample Stand In for the Population



The Resampling Idea: Let the Sample Stand In for the Population



Efron (1979): the bootstrap

Everything we cannot compute in the real world (because F is unknown) we **can** compute in the bootstrap world (because $\hat{F}_n$ is known) – by simulation.

Outline

1. Motivation: Inference When Formulas Run Out
2. The Bootstrap Principle
   - Empirical Distribution and the Plug-in Principle
   - The Nonparametric Bootstrap Algorithm
   - A Toy Example by Hand
   - Bootstrap Standard Error and Bias
   - Why It Works and How Many Resamples
3. Parametric vs Nonparametric Bootstrap
4. Bootstrap Confidence Intervals
5. The Jackknife (Brief)
6. Cross-Validation
7. Monte Carlo Hypothesis Testing and Permutation Tests
8. Likelihood-Based Inference via Simulation
9. Simulation vs Asymptotic Inference
10. Unit Summary

The Empirical Distribution Function

Non parametric approach.

Definition

The empirical distribution $\hat{F}_n$ puts probability mass $1/n$ on each observation:

$$\hat{F}_n(t) = \frac{1}{n} \sum_{i=1}^{n} \mathbf{1}\{x_i \leq t\}.$$

indicator fn

The Empirical Distribution Function

Definition

The empirical distribution $\hat{F}_n$ puts probability mass $1/n$ on each observation:

$$\hat{F}_n(t) = \frac{1}{n} \sum_{i=1}^{n} \mathbf{1}\{x_i \leq t\}.$$

- **Glivenko–Cantelli:** $\sup_t |\hat{F}_n(t) - F(t)| \xrightarrow{\text{a.s.}} 0.$

The Empirical Distribution Function

Definition

The empirical distribution $\hat{F}_n$ puts probability mass $1/n$ on each observation:

$$\hat{F}_n(t) = \frac{1}{n} \sum_{i=1}^{n} \mathbf{1}\{x_i \le t\}.$$

- **Glivenko–Cantelli:** $\sup_t |\hat{F}_n(t) - F(t)| \xrightarrow{\text{a.s.}} 0$.
- This is the same D_n used in the Kolmogorov–Smirnov test of Module 1.

The Empirical Distribution Function

Definition

The empirical distribution $\hat{F}_n$ puts probability mass $1/n$ on each observation:

$$\hat{F}_n(t) = \frac{1}{n} \sum_{i=1}^{n} \mathbf{1}\{x_i \le t\}.$$

- **Glivenko–Cantelli:** $\sup_t |\hat{F}_n(t) - F(t)| \xrightarrow{\text{a.s.}} 0$.
- This is the same D_n used in the Kolmogorov–Smirnov test of Module 1.
- $\hat{F}_n$ is the nonparametric MLE of F: the best guess of F when we assume nothing.


The Empirical Distribution Function

Definition

The empirical distribution $\hat{F}_n$ puts probability mass $1/n$ on each observation:


$$\hat{F}_n(t) = \frac{1}{n} \sum_{i=1}^{n} \mathbf{1}\{x_i \le t\}.$$

- **Glivenko–Cantelli:** $\sup_t |\hat{F}_n(t) - F(t)| \xrightarrow{\text{a.s.}} 0$.
- This is the same D_n used in the Kolmogorov–Smirnov test of Module 1.
- $\hat{F}_n$ is the nonparametric MLE of F: the best guess of F when we assume nothing.



The Plug-in Principle

- Many parameters are *functionals* of F: $\theta = T(F)$. The **plug-in estimator** is $\hat{\theta} = T(\hat{F}_n)$.



The Plug-in Principle

- Many parameters are *functionals* of F: $\theta = T(F)$. The **plug-in estimator** is $\hat{\theta} = T(\hat{F}_n)$.

Parameter	$T(F)$	Plug-in $T(\hat{F}_n)$
Mean	$\int x\, dF(x)$	$\bar{x}$
Variance	$\int (x-\mu)^2 dF(x)$	$\hat{\sigma}^2 = \frac{1}{n}\sum (x_i - \bar{x})^2$ (divisor n!)
Median	$F^{-1}(1/2)$	sample median
Correlation	$\mathrm{Cov}(X,Y)/(\sigma_X \sigma_Y)$	sample correlation r

CDF

$\frac{dF(x)}{dx} = f(x)$ → PDF → $E(X) = \int x f(x)\, dx$

$dF(x) = f(x)\,dx$

$f(x)dx = dF(x)$ — CDF

$V(X) = \int (x-\mu)^2 f(x)\,dx$ — $dF(x)$

The Plug-in Principle

- Many parameters are *functionals* of F: $\theta = T(F)$. The **plug-in estimator** is $\hat{\theta} = T(\hat{F}_n)$.

Parameter	$T(F)$	Plug-in $T(\hat{F}_n)$
Mean	$\int x\, dF(x)$	$\bar{x}$
Variance	$\int (x-\mu)^2 dF(x)$	$\hat{\sigma}^2 = \frac{1}{n}\sum (x_i - \bar{x})^2$ (divisor n!)
Median	$F^{-1}(1/2)$	sample median
Correlation	$\mathrm{Cov}(X,Y)/(\sigma_X \sigma_Y)$	sample correlation r

The bootstrap is the plug-in principle applied one level up

The standard error is *also* a function of F: $\mathrm{se}_F(\hat{\theta})$. Its plug-in estimate is

$$\mathrm{se}_{\hat{F}_n}(\hat{\theta}^*) = \text{standard deviation of } s(\boldsymbol{x}^*) \text{ when } \boldsymbol{x}^* \overset{\text{iid}}{\sim} \hat{F}_n.$$


This is the **ideal bootstrap estimate** of the standard error.

$$\widehat{se}_{\hat{F}_n}(\theta^*) = \sqrt{\frac{1}{B}\sum_{i=1}^{B} (\hat{\theta}_i^* - \bar{\theta}^*)^2}$$

Sampling from $\hat{F}_n$ = Resampling With Replacement


- $\hat{F}_n$ is a **discrete** distribution with $P(X^* = x_i) = 1/n$, $i = 1, \ldots, n$.

$p(x = x_i) = p_i$
$= \frac{1}{n}$



Sampling from $\hat{F}_n$ = Resampling With Replacement

- $\hat{F}_n$ is a **discrete** distribution with $P(X^* = x_i) = 1/n$, $i = 1, \ldots, n$.
- By method of Module 1: draw $U \sim \text{Unif}(0,1)$, set $I = \lceil nU \rceil$, and return x_I.



Sampling from $\hat{F}_n$ = Resampling With Replacement


- $\hat{F}_n$ is a **discrete** distribution with $P(X^* = x_i) = 1/n$, $i = 1, \ldots, n$.
- By method of Module 1: draw $U \sim \text{Unif}(0, 1)$, set $I = \lceil nU \rceil$, and return x_I.
- Repeating n times gives a bootstrap sample $\boldsymbol{x}^* = (x_{I_1}, \ldots, x_{I_n})$ – some observations appear several times, some not at all.

$x_1 \cdots x_n \rightarrow$ distinct

$\boldsymbol{x}_1^* = (x_{11} \cdots x_{1n})$ —

Sampling from $\hat{F}_n$ = Resampling With Replacement

- $\hat{F}_n$ is a **discrete** distribution with $P(X^* = x_i) = 1/n$, $i = 1, \ldots, n$.
- By method of Module 1: draw $U \sim \text{Unif}(0, 1)$, set $I = \lceil nU \rceil$, and return x_I.
- Repeating n times gives a bootstrap sample $\boldsymbol{x}^* = (x_{I_1}, \ldots, x_{I_n})$ – some observations appear several times, some not at all.



The Nonparametric Bootstrap: Algorithm

Algorithm: Nonparametric Bootstrap

Given data $\boldsymbol{x} = (x_1, \ldots, x_n)$ and a statistic $s(\cdot)$, with $\hat{\theta} = s(\boldsymbol{x})$:

1. For $b = 1, \ldots, B$:
   (a) Draw $\boldsymbol{x}^{*b} = (x_1^{*b}, \ldots, x_n^{*b})$ by sampling n values **with replacement** from $\boldsymbol{x}$.
   (b) Compute the bootstrap replication $\hat{\theta}_b^* = s(\boldsymbol{x}^{*b})$.
2. Use the empirical distribution of $\hat{\theta}_1^*, \ldots, \hat{\theta}_B^*$ as an approximation to the sampling distribution of $\hat{\theta}$.