

Function and Limits

- $\lim_{x \rightarrow \pm \infty} \frac{Ax^\alpha}{Bx^\beta} \begin{cases} 0 & \text{if } \alpha < \beta \\ \frac{A}{B} & \text{if } \alpha = \beta \\ \pm \infty & \text{if } \alpha > \beta \end{cases}$
- $\lim_{x \rightarrow c} \frac{\sin(g(x))}{g(x)} = 1 \quad (\lim_{x \rightarrow c} g(x) = 0)$
- $\lim_{x \rightarrow c} \frac{\tan(g(x))}{g(x)} = 1$
- $\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$
- $\lim_{x \rightarrow 0} \frac{\tan(x)}{x} = 1$

Differentiation

parametric differentiaton: $\frac{d^2y}{dx^2} = \frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{\frac{d}{dt} \left(\frac{dy}{dx} \right)}{\frac{dx}{dt}}$

$f(x)$	$f'(x)$
$\tan x$	$\sec^2 x$
$\csc x$	$-\csc x \cot x$
$\sec x$	$\sec x \tan x$
$\cot x$	$-\csc^2 x$
$a^{f(x)}$	$\ln a \cdot f'(x) a^{f(x)}$
$\log_a f(x)$	$\log_a e \cdot \frac{f'(x)}{f(x)}$
$\sin^{-1} f(x)$	$\frac{f'(x)}{\sqrt{1-[f(x)]^2}}, \quad f(x) < 1$
$\cos^{-1} f(x)$	$-\frac{f'(x)}{\sqrt{1-[f(x)]^2}}, \quad f(x) < 1$
$\tan^{-1} f(x)$	$\frac{f'(x)}{1+[f(x)]^2}$
$\cot^{-1} f(x)$	$-\frac{f'(x)}{1+[f(x)]^2}$
$\sec^{-1} f(x)$	$\frac{f'(x)}{ f(x) \sqrt{[f(x)]^2-1}}$
$\csc^{-1} f(x)$	$-\frac{f'(x)}{ f(x) \sqrt{[f(x)]^2-1}}$

Second Derivative Test: $f'(c) = 0, f''(c) < 0$ then local max, $f''(c) > 0$ local min.

L'Hopital's Rule: Given $\lim_{x \rightarrow c} f(x)$ and $g(x) = 0 / \pm \infty$

$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}$

- Use for $\frac{0}{0}$ or $\frac{\infty}{\infty}$

Trigo Identities

- $\sec^2 x - 1 = \tan^2 x$
- $\csc^2 x - 1 = \cot^2 x$
- $\sin A \cos A = \frac{1}{2} \sin 2A$
- $\cos^2 A = \frac{1}{2} (1 + \cos 2A)$
- $\sin^2 A = \frac{1}{2} (1 - \cos 2A)$
- $\sin A \cos B = \frac{1}{2} (\sin(A+B) + \sin(A-B))$
- $\cos A \sin B = \frac{1}{2} (\sin(A+B) - \sin(A-B))$
- $\cos A \cos B = \frac{1}{2} (\cos(A+B) + \cos(A-B))$
- $\sin A \sin B = \frac{1}{2} (\cos(A+B) - \cos(A-B))$

Integration

$f(x)$	$\int f(x)$
$\tan ax$	$\frac{1}{a} \ln \sec(ax) $
$\cot ax$	$\frac{1}{a} \ln \cot(ax) $
$\sec ax$	$\frac{1}{a} \ln \sec(ax) + \tan(ax) $
$\csc ax$	$\frac{1}{a} \ln \csc(ax) + \cot(ax) $
$\frac{1}{a^2 + (x+b)^2}$	$\frac{1}{a} \tan^{-1} \left(\frac{x+b}{a} \right)$
$\frac{1}{\sqrt{a^2 - (x+b)^2}}$	$\sin^{-1} \left(\frac{x+b}{a} \right)$
$\frac{1}{a^2 - (x+b)^2}$	$\frac{1}{2a} \ln \left \frac{x+b+a}{x+b-a} \right $
$\frac{1}{(x+b)^2 - a^2}$	$\frac{1}{2a} \ln \left \frac{x+b-a}{x+b+a} \right $

Substitution $\int f(g(x)) \cdot g'(x) dx = \int f(u) du, u = g(x)$

By Parts $\int uv' dx = uv - \int u'v dx$, order: LIATE:

Differentiate to integrate

Application of Integration

about x axis


- Vol Disk: $V = \pi \int_a^b f(x)^2 - g(x)^2 dx$
- Vol Shell: $V = 2\pi \int_a^b x|f(x) - g(x)| dx$ (absolute!!)
- Length of curve: $\int_a^b \sqrt{1 + f'(x)^2} dx$

Series

TODO!!

Vectors

unit vector: $\hat{p} = \frac{p}{|p|}, \overrightarrow{AB} = \overrightarrow{OB} - \overrightarrow{OA}$



Dot Product

- $\vec{a} \cdot \vec{b} = a_1b_1 + a_2b_2 + a_3b_3 = |a||b| \cos \theta$
- $a \perp b \Rightarrow a \cdot b = 0$
- $a \parallel b \Rightarrow a \cdot b = |a||b|$

Cross Product

$a \times b = \begin{vmatrix} i & j & k \\ a_i & a_2 & a_3 \\ b_i & b_2 & b_3 \end{vmatrix} = \begin{pmatrix} (a_2b_3 - a_3b_2) \\ -(a_1b_3 - a_3b_1) \\ (a_1b_2 - a_2b_1) \end{pmatrix}$

$|a \times b| = |a||b| \sin \theta$
 $a \perp b \Rightarrow a \times b = |a||b|$
 $a \parallel b \Rightarrow a \times b = 0$
Parallelogram = $|a \times b|$

Projection



Lines

$r = r_0 + tv = \langle x, y, z \rangle$
 $\langle x_0, y_0, z_0 \rangle + t \langle a, b, c \rangle$
 $\begin{pmatrix} x_0 + at \\ y_0 + bt \\ z_0 + ct \end{pmatrix}$

Planes

$n = \langle a, b, c \rangle, r = \langle x, y, z \rangle, r_0 \langle x_0, y_0, z_0 \rangle$

Scalar: $n \cdot r = n \cdot r_0$

Cartesian: $ax + by + cz = d$

Distance from Point to Plane

$\frac{|ax_0 + by_0 + cz_0 - d|}{\sqrt{a^2 + b^2 + c^2}}$

Partial Derivatives

Chain Rule

For $z(t) = f(x(t), y(t))$,

$\frac{dz}{dt} = \frac{\partial z}{\partial x} \frac{dx}{dt} + \frac{\partial z}{\partial y} \frac{dy}{dt}$

For $z(s, t) = f(x(s, t), y(s, t))$,

$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial t}$

$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial s}$

Arc Length of $r(t)$: $\int_a^b |r'(t)| dt$

Implicit Differentiation

$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z}, \frac{\partial z}{\partial y} = -\frac{F_y}{F_z}$

Directional Derivative

Gradient vector at $f(x, y) : \Delta f = f_x i + f_y j$

$D_u f(x, y) = \langle f_x, f_y \rangle \cdot \langle a, b \rangle = \langle f_x, f_y \rangle \cdot \hat{u} = \Delta f \cdot \hat{u}$
(Unit Vector)

Tangent Plane: $\Delta f \cdot \langle x - x_0, y - y_0, z - z_0 \rangle = 0$


Critical Points

$D = f_{xx}(a, b)f_{yy}(a, b) - (f_{xy}(a, b))^2$


D	$f_{xx}(a, b)$	local
+	+	min
+	-	max
-	any	saddle point
0	any	no conclusion

Double Integrals


Type I



Type II



Polar Coordinates



$x = r \cos \theta$

$y = r \sin \theta$

$R = \{(r, \theta) : 0 \leq a \leq r \leq b, \alpha \leq \theta \leq \beta\}$

$\int_\alpha^\beta \int_a^b f(r \cos \theta, r \sin \theta) r dr d\theta$

Surface Area

$S = \iint_R \sqrt{f_x^2 + f_y^2 + 1} dA$

ODE

form	change of variable
$\frac{dy}{dx} = f(x)g(y)$	$\int \frac{1}{g(y)} dy = \int f(x) dx + C$
$y' = g(\frac{y}{x})$	Set $v = \frac{y}{x}$ $\Rightarrow y' = v + xv'$
$y' = f(ax + by + c)$ $\Rightarrow y' = \frac{ax + by + c}{\alpha x + \beta y + \gamma}$	Set $v = ax + by$
$y' + P(x)y = Q(x)$	$R = e^{\int P(x) dx}$ $\Rightarrow y \cdot R = \int Q \cdot R dx$ $z = y^{1-n}$ \Rightarrow sub in Z solve linear
$y' + P(x)y = Q(x)y^n$	