

Medications

Commertial Name	Generic Name	Active Ingiridiant	Main Usage	Main Side Effects	Allergens and Restrictions	Menner of Cunsumption	Recommended Dosage	Menner of Storage
Acamol	Paracetamol	Paracetamol	Reliev pain and fever	dizziness	keep away from kids. If you are pregnant or nursing please cunsolt your gynecologist.	Oral	one pill every 4-6 hours	keep in dark and cool place at its original package
Adizem	Diltiazem Hydrochloride	Diltiazem Hydrochloride	Diltiazem belongs to the group of drugs that block calcium channels. It is used to treat high blood pressure and angina.	Headaches, fatigue, weakness, edema of the limbs.	Not allowed during breastfeeding, not allowed during pregnancy and not allowed with alcohol. The drug should not be used in case of very low blood pressure or in case of obstructive arrhythmias.	Oral	3 times a day.	The medicine should be stored in a cool and dry place, out of the reach of children. It must be protected from exposure to light.
Advil	Ibuprofen	Ibuprofen	Reliev pain and fever	dizziness	keep away from kids. If you are pregnant or nursing please cunsolt your gynecologist.	Oral	one pill every 4-6 hours	keep in dark and cool place at its original package
Agispor shampoo	Bifonazole	Bifonazole	A broad-spectrum antifungal drug from the imidazole family intended for local treatment.	Preparations containing bifonazole usually do not have significant side effects.	keep away from children. Do not use the product without consulting a doctor before starting treatment if you are pregnant or breastfeeding.	Dermal	Wash your hair 3 times a week for 4 weeks	The preparation must be kept in the original packaging at a temperature not higher than 25 degrees Celsius. After opening for the first time, the preparation must be used within 6 months.
Agisten V 0.2	Imidazole derivatives	Imidazole derivatives	Broad spectrum antimycotic with fungicidal and trichomonacidal action (for single dose treatment).	itching, redness, rash, swelling, itching, burning. Local vagal irritation or peeling, bleeding, pain in the abdomen or pelvic area. If one of the side effects persists or worsens, see a doctor	Do not use the preparation without consulting a doctor before starting the treatment. If you are pregnant or breastfeeding.	Vaginal	one a day for 3 days per treatment	keep in a cool and dark place
Agisten cream	Clotrimazole	Clotrimazole	For the treatment of fungal infections of the skin and mucosa caused be dermatophytes, yeasts, molds and other fungi such as Malassezia furfur as well as skin infections caused by Corynebacterium minutissimum.Indicated also for Candida Vulvitis and Candida balanitis.	After using the medicine you may experience one of the following symptoms: Itching, rash, blisters, burning, discomfort, swelling, irritation, redness or peeling of the skin, feeling warm and tingling in the treated area. If one of the side effects worsens, or when you suffer from a side effect that was not mentioned in the leaflet, you should consult the doctor.	Do not use the product without consulting a doctor before starting treatment if you are pregnant or breastfeeding. Do not use if you are known to be sensitive to Clotrimazole.	Vaginal	three times a day	keep in a cool dark place
Alfuzosin	Alfuzosin Hydrochloride	Alfuzosin Hydrochloride	Lefusin is intended to treat benign enlargement of the prostate gland.	Dizziness, weakness, headaches, abdominal pain, constipation, heartburn, nausea, impotence, drop in blood pressure when getting up from lying down to the point of fainting, chest pain, accelerated pulse.	The drug is not intended for women. In case of liver disease or if there is a history of orthostatism, the attending physician should be informed.	Oral	10 milligrams once a day or 5 milligrams twice a day.	The medicine should be kept in a cool and dry place, out of the reach of children.
Anaesthetic	Phenazone+Amethocaine	Phenazone, Tetracaine hydrochloride, Glycerin	For the relief of pain in acute otitis media and the external, auditory canal.	earache	keep away from childrens.	Through the Ears	twice a day, morning and evning	keep in a cool dark place
Carbosylane	Charcoal activated	Charcoal activated,Simethicone.■	The medicine helps in cases of abdominal pain or minor disturbances in the digestive system caused by gas accumulation.	Constipation, nausea, vomiting, change in stool color (black stool).	keep awa from children	Oral	up to 3 times a day, before the meal.	keep in cool and dark place
Paxlovid	Ritonavir+Nirmatrelvir	Ritonavir,Nirmatrelvir	Paxolid is used to treat corona patients who are at high risk of deterioration in their health condition.	Paxlobide may cause liver problems. You should immediately report to your doctor if you suffer from one or more of the following symptoms that can indicate liver problems: loss of appetite, yellowing of the skin and the whites of the eyes (jaundice), dark urine, light-colored stools, itchy skin, abdominal pain.	keep away from children. A woman should inform her doctor if she is pregnant or breastfeeding before starting to use paxlobide. She must also report if she is planning a pregnancy. If you suffer from kidney failure, liver disease, you must report this to your doctor.	Oral	Once a day, Paxalobide should not be taken for more than 5 days. Consult the doctor if the feeling does not improve after 5 days.	Paxolide tablets should be stored at room temperature. If at the end of the treatment there are excess tablets left, they must be returned to the pharmacy for destruction.
Prenatal	Prenatal	Beta-Carotene,Vitamin A,Vitamin E,Vitamin C,Folic Acid,Vitamin B1,Vitamin B2,Niacinamide,Vitamin B6,Vitamin B12,Vitamin D3, Biotin,Pantothenic Acid,Calcium,Magnesium,IodineIron,Copper,Zinc,Chromium,Manganes,Molybdenum,Selenium.	Supplementation of vitamins and minerals for pregnant and lactating women.	Constipation, gas, nausea, belching.	This medicine is not intended for children and babies.	Oral	once a day	Keep at room temperature, close tightly and prevent air and moisture penetration
Rokacet	Rokacet	Caffeine,Codeine phosphate,Paracetamol.	Pain relief, to relieve cough and reduce fever accompanied by pain.	Constipation, abdominal pain, nausea, changes in alertness. The drug contains caffeine, which is a stimulant, and codeine, which has an antiaging effect. Therefore, there may be side effects such as sleepiness, impaired alertness or concentration along with hyperactivity.	keep away from children. The use of the drug should be short-term. Prolonged use may cause dependence and addiction. It is forbidden to use Rokacet during pregnancy. Pregnant women who suffer from pain should consult with the attending physician regarding other treatment that is appropriate during pregnancy. It is forbidden to use Rokacet during breastfeeding. Alcohol should be avoided while being treated with Roxacet	Oral	every 6 hours no longer then 3 days	keep in cool and drak place
Stilla	Tetrahydrozoline hydrochloride	Tetrahydrozoline hydrochloride	Irritation, burning, reddening, itching of the eye due to allergic or physical causes such as dust, smoke, plants,sun rays.	None	keep away from childrens.	Through the Eyes	up to 10 time a day as needed	keep in a cool and dark place
Turns ex sugar free	Calcium carbonate	Calcium carbonate	To relieve heartburn and acidity in the stomach.	Constipation, gas, nausea, belching.	Do not use the medicine without consulting a doctor before starting in children under 12 years old. Do not use the product without consulting a doctor before starting treatment if you are pregnant or breastfeeding. Do not use if you are known to be sensitive to Clotrimazole. The medicine contains sunset (110E (yellow), which may cause allergic reactions. The medicine contains sorbitol. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine. The medicine contains aspartame, a source of phenylalanine, which may be harmful to people with phenylketonuria.	Oral	up to 10 tablets a day as need	keep in a drak and cool place at its original package
Vitamin B12	Sodium	Ascorbic acid, Biotin,Cyanocobalamin,Folic acid,Nicotinamide,Pantothenic acid,Pyridoxine,Riboflavin,Thiamine.	Multivaitamin	Constipation and gas.	keep away from childrens.	Sublingual	For adult patients and children weighing 10 kg or more the recommended daily dosage is the contents of one vial.	keep in cool and dark plase