





Associations by Trait Category

Associations by Category



	HEMATOLOGICAL		CARDIOVASCULAR		STROKE		RENAL
	ANTHROPOMETRIC		LIPIDS		MUSCULOSKELETAL		GLYCEMIC
	HEPATIC		REPRODUCTIVE TRAITS		LUNG		METABOLITE
	AGING AND LONGEVITY		SLEEP AND CIRCADIAN		PSYCHIATRIC		IMMUNOLOGICAL
	OCULAR		OTHER		NEUROLOGICAL		NUTRITIONAL
	DIGESTIVE		SKIN CONDITIONS		CANCER		ATRIAL FIBRILLATION
	Uncategorized		ECG TRAITS		INFLAMMATORY BOWEL		NON-ADDITIVE MODELS
	COGNITIVE		DIABETIC COMPLICATIONS		TYPE 1 DIABETES		PHYSICAL ACTIVITY
	COVID-19		CEREBROVASCULAR MRI TRAITS		SULFONYLUREA RESPONSE		METFORMIN RESPONSE

This genetic concentration varies significantly depending on the specific trait class. Certain phenotypes exhibit broad support profiles, while others remain highly concentrated. For instance, metabolic conditions such as lipid-related traits systematically demonstrate higher polygenicity, averaging more than [XX] high-probability genes per trait (Fig. 4a).