

Lecture 15: Orders & Decimal Expansions

Tom Roby
University of Connecticut

23 October 2012

Outline

- Orders of elements in \mathbf{Z}_m^* ;
- Orders and powers of $a \in \mathbf{Z}_m^*$;
- Periods of decimal expansions;

Example & Definitions

- Consider \mathbf{Z}_9^* . How many units? What are they? What are their powers?

Example & Definitions

- Consider \mathbf{Z}_9^* . How many units? What are they? What are their powers?
- DEF: Given $a \in \mathbf{Z}_m^*$, the least $n \in \mathbf{Z}^+$ s.t. $a^n = 1$ is called the **order** of $a \bmod m$. Sometimes we write $ord(a)$, $ord_m(a)$, $o(a)$ or $o_m(a)$.

Example & Definitions

- Consider \mathbf{Z}_9^* . How many units? What are they? What are their powers?
- DEF: Given $a \in \mathbf{Z}_m^*$, the least $n \in \mathbf{Z}^+$ s.t. $a^n = 1$ is called the **order** of $a \bmod m$. Sometimes we write $ord(a)$, $ord_m(a)$, $o(a)$ or $o_m(a)$.
- EG: What is the order of 2 mod 9? Of 7? What do you notice about the order of elements?

Example & Definitions

- Consider \mathbf{Z}_9^* . How many units? What are they? What are their powers?
- DEF: Given $a \in \mathbf{Z}_m^*$, the least $n \in \mathbf{Z}^+$ s.t. $a^n = 1$ is called the **order** of $a \bmod m$. Sometimes we write $ord(a)$, $ord_m(a)$, $o(a)$ or $o_m(a)$.
- EG: What is the order of 2 mod 9? Of 7? What do you notice about the order of elements?
- THM: If $a \bmod m$ has order n , then for $k \in \mathbf{Z}^+$
 $a^k = 1 \iff o(a) \mid k$.

Example & Definitions

- Consider \mathbf{Z}_9^* . How many units? What are they? What are their powers?
- DEF: Given $a \in \mathbf{Z}_m^*$, the least $n \in \mathbf{Z}^+$ s.t. $a^n = 1$ is called the **order** of $a \bmod m$. Sometimes we write $ord(a)$, $ord_m(a)$, $o(a)$ or $o_m(a)$.
- EG: What is the order of 2 mod 9? Of 7? What do you notice about the order of elements?
- THM: If $a \bmod m$ has order n , then for $k \in \mathbf{Z}^+$
 $a^k = 1 \iff o(a) \mid k$.
- PF: (\Leftarrow) is easy from the definitions.
 (\Rightarrow) : Use division algorithm.

Big Theorem on Orders

Here is the THE big theorem about orders in \mathbf{Z}_m^* .

Theorem

Let $a \bmod m$ have order n . THEN

- 1 *For all $k \in \mathbf{Z}^+$, $a^k \equiv 1 \pmod{m} \iff o(a) \mid k$. In particular, $o(a) \mid \phi(m)$, and $a^k \equiv a^l \iff k \equiv l \pmod{o(a)}$.*

Big Theorem on Orders

Here is the THE big theorem about orders in \mathbf{Z}_m^* .

Theorem

Let $a \bmod m$ have order n . THEN

- ① *For all $k \in \mathbf{Z}^+$, $a^k \equiv 1 \pmod{m} \iff o(a) \mid k$. In particular, $o(a) \mid \phi(m)$, and $a^k \equiv a^l \iff k \equiv l \pmod{o(a)}$.*
- ② *If $(k, o(a)) = 1$, then a^k has the same order as a .*

Big Theorem on Orders

Here is the THE big theorem about orders in \mathbf{Z}_m^* .

Theorem

Let $a \bmod m$ have order n . THEN

- ① *For all $k \in \mathbf{Z}^+$, $a^k \equiv 1 \pmod{m} \iff o(a) \mid k$. In particular, $o(a) \mid \phi(m)$, and $a^k \equiv a^l \iff k \equiv l \pmod{o(a)}$.*
- ② *If $(k, o(a)) = 1$, then a^k has the same order as a .*
- ③ *If $d \mid o(a)$, then a^d has order $\frac{o(a)}{d}$.*

Big Theorem on Orders

Here is the THE big theorem about orders in \mathbf{Z}_m^* .

Theorem

Let $a \bmod m$ have order n . THEN

- ① *For all $k \in \mathbf{Z}^+$, $a^k \equiv 1 \pmod{m} \iff o(a) \mid k$. In particular, $o(a) \mid \phi(m)$, and $a^k \equiv a^l \iff k \equiv l \pmod{o(a)}$.*
- ② *If $(k, o(a)) = 1$, then a^k has the same order as a .*
- ③ *If $d \mid o(a)$, then a^d has order $\frac{o(a)}{d}$.*
- ④ *If a_1 has order n_1 and a_2 has order n_2 in \mathbf{Z}_m^* , with $(n_1, n_2) = 1$, then $a_1 a_2$ has order $n_1 n_2$.*

Big Theorem on Orders

Here is the THE big theorem about orders in \mathbf{Z}_m^* .

Theorem

Let $a \bmod m$ have order n . THEN

- ❶ *For all $k \in \mathbf{Z}^+$, $a^k \equiv 1 \pmod{m} \iff o(a) \mid k$. In particular, $o(a) \mid \phi(m)$, and $a^k \equiv a^l \iff k \equiv l \pmod{o(a)}$.*
- ❷ *If $(k, o(a)) = 1$, then a^k has the same order as a .*
- ❸ *If $d \mid o(a)$, then a^d has order $\frac{o(a)}{d}$.*
- ❹ *If a_1 has order n_1 and a_2 has order n_2 in \mathbf{Z}_m^* , with $(n_1, n_2) = 1$, then $a_1 a_2$ has order $n_1 n_2$.*

EG: Let's look at a power table for \mathbf{Z}_{19}^* .

Decimal expansions

Consider the decimal expansions of the following fractions:

$$\frac{1}{3}, \frac{1}{7}, \frac{7}{22}, \frac{15}{37}.$$

Decimal expansions

Consider the decimal expansions of the following fractions:

$\frac{1}{3}, \frac{1}{7}, \frac{7}{22}, \frac{15}{37}$.

$$\frac{1}{3} = .33333 \dots = .\overline{3} \quad \text{Period} = 1$$

$$\frac{1}{7} = .1428571428 \dots = .\overline{142857} \quad \text{Period} = 6$$

$$\frac{7}{22} = .3181818 \dots = 3.\overline{18} \quad \text{Period} = 2$$

$$\frac{15}{37} = .405405405 \dots = .\overline{405} \quad \text{Period} = 3$$

Decimal expansions

Consider the decimal expansions of the following fractions:

$\frac{1}{3}, \frac{1}{7}, \frac{7}{22}, \frac{15}{37}$.

$$\frac{1}{3} = .33333\ldots = .\overline{3} \quad \text{Period} = 1$$

$$\frac{1}{7} = .1428571428\ldots = .\overline{142857} \quad \text{Period} = 6$$

$$\frac{7}{22} = .3181818\ldots = 3.\overline{18} \quad \text{Period} = 2$$

$$\frac{15}{37} = .405405405\ldots = .\overline{405} \quad \text{Period} = 3$$

Any Conjectures from here or the HW?

True/False: Every rational number has a repeating decimal expansion?

Decimal expansions

Consider the decimal expansions of the following fractions:

$\frac{1}{3}, \frac{1}{7}, \frac{7}{22}, \frac{15}{37}$.

$$\frac{1}{3} = .33333\ldots = .\overline{3} \quad \text{Period} = 1$$

$$\frac{1}{7} = .1428571428\ldots = .\overline{142857} \quad \text{Period} = 6$$

$$\frac{7}{22} = .3181818\ldots = 3.\overline{18} \quad \text{Period} = 2$$

$$\frac{15}{37} = .405405405\ldots = .\overline{405} \quad \text{Period} = 3$$

Any Conjectures from here or the HW?

True/False: Every rational number has a repeating decimal expansion? ANS: **TRUE!**

Decimal Expansion Conjectures

Prove or Disprove & Salvage if Possible:

- 1 The period length of $\frac{1}{p}$ divides $p - 1$;
- 2 For each $b \geq 2$, all $\frac{a}{b}$ with $(a, b) = 1$ have same period length.
- 3 Expansion of $\frac{1}{b}$ is purely periodic when $(10, b) = 1$.
- 4 Shifting digits (cyclically) gives another fraction with same denominator.

Converting between decimals and fractions

- To go from fractions to decimal expansions is easy. Why?

Converting between decimals and fractions

- To go from fractions to decimal expansions is easy. Why?
- Conversely, how do we go from fractions to decimals? EG, what fraction is represented by $x = .\overline{15}$?

Converting between decimals and fractions

- To go from fractions to decimal expansions is easy. Why?
- Conversely, how do we go from fractions to decimals? EG, what fraction is represented by $x = .\overline{15}$?
- How about $x = .\overline{405}$

Converting between decimals and fractions

- To go from fractions to decimal expansions is easy. Why?
- Conversely, how do we go from fractions to decimals? EG, what fraction is represented by $x = .\overline{15}$?
- How about $x = .\overline{405}$
- What happens in general with $x = .\overline{c_1 c_2 \cdots c_d}$?

Converting between decimals and fractions

- To go from fractions to decimal expansions is easy. Why?
- Conversely, how do we go from fractions to decimals? EG, what fraction is represented by $x = .\overline{15}$?
- How about $x = .\overline{405}$
- What happens in general with $x = .\overline{c_1 c_2 \cdots c_d}$?
- Some algebra shows that $\frac{a}{b}$ has a *purely periodic* decimal expansion $0.\overline{c_1 \cdots c_d} \iff$ some representation of $\frac{a}{b}$ has denominator $10^d - 1$ for some $d \in \mathbf{Z}^+$.

Converting between decimals and fractions

- To go from fractions to decimal expansions is easy. Why?
- Conversely, how do we go from fractions to decimals? EG, what fraction is represented by $x = .\overline{15}$?
- How about $x = .\overline{405}$
- What happens in general with $x = .\overline{c_1 c_2 \cdots c_d}$?
- Some algebra shows that $\frac{a}{b}$ has a *purely periodic* decimal expansion $0.\overline{c_1 \cdots c_d} \iff$ some representation of $\frac{a}{b}$ has denominator $10^d - 1$ for some $d \in \mathbf{Z}^+$.
- **NB:** The denominator may be something other than $10^d - 1$ when the fraction is simplified to lowest terms.

Converting between decimals and fractions

- To go from fractions to decimal expansions is easy. Why?
- Conversely, how do we go from fractions to decimals? EG, what fraction is represented by $x = .\overline{15}$?
- How about $x = .\overline{405}$
- What happens in general with $x = .\overline{c_1 c_2 \cdots c_d}$?
- Some algebra shows that $\frac{a}{b}$ has a *purely periodic* decimal expansion $0.\overline{c_1 \cdots c_d} \iff$ some representation of $\frac{a}{b}$ has denominator $10^d - 1$ for some $d \in \mathbf{Z}^+$.
- **NB:** The denominator may be something other than $10^d - 1$ when the fraction is simplified to lowest terms.
EG: $\frac{5}{33} = \frac{15}{99} = .\overline{15}$; $\frac{1}{3} = \frac{3}{9} = 0.\overline{3}$.

Theorem

Let $x = \frac{a}{b} \in \mathbf{Q}^+$ with $(b, 10) = 1$. Then the decimal period of x is $\text{ord}_b(10)$, the order of 10 in \mathbf{Z}_b^ .*

Theorem

Let $x = \frac{a}{b} \in \mathbf{Q}^+$ with $(b, 10) = 1$. Then the decimal period of x is $\text{ord}_b(10)$, the order of 10 in \mathbf{Z}_b^ .*

Proof.

decimal period of x = least $d \geq 1$ s.t. x has denom. $10^d - 1$



Theorem

Let $x = \frac{a}{b} \in \mathbf{Q}^+$ with $(b, 10) = 1$. Then the decimal period of x is $\text{ord}_b(10)$, the order of 10 in \mathbf{Z}_b^ .*

Proof.

decimal period of x = least $d \geq 1$ s.t. x has denom. $10^d - 1$
= least $d \geq 1$ s.t. $b \mid 10^d - 1$



Theorem

Let $x = \frac{a}{b} \in \mathbf{Q}^+$ with $(b, 10) = 1$. Then the decimal period of x is $\text{ord}_b(10)$, the order of 10 in \mathbf{Z}_b^ .*

Proof.

decimal period of x = least $d \geq 1$ s.t. x has denom. $10^d - 1$
= least $d \geq 1$ s.t. $b \mid 10^d - 1$
= least $d \geq 1$ s.t. $10^d \equiv 1 \pmod{b}$



Theorem

Let $x = \frac{a}{b} \in \mathbf{Q}^+$ with $(b, 10) = 1$. Then the decimal period of x is $\text{ord}_b(10)$, the order of 10 in \mathbf{Z}_b^ .*

Proof.

decimal period of x = least $d \geq 1$ s.t. x has denom. $10^d - 1$
= least $d \geq 1$ s.t. $b \mid 10^d - 1$
= least $d \geq 1$ s.t. $10^d \equiv 1 \pmod{b}$
= order of 10 mod b



Orders & Decimals

Theorem

Let $x = \frac{a}{b} \in \mathbf{Q}^+$ with $(b, 10) = 1$. Then the decimal period of x is $\text{ord}_b(10)$, the order of 10 in \mathbf{Z}_b^* .

Proof.

decimal period of x = least $d \geq 1$ s.t. x has denom. $10^d - 1$
= least $d \geq 1$ s.t. $b \mid 10^d - 1$
= least $d \geq 1$ s.t. $10^d \equiv 1 \pmod{b}$
= order of 10 mod b



p	3	7	9	11	13	17	19	21
Period $\frac{1}{p}$	1	6	1	2	6	16	18	6
$\text{ord}_{10}(p)$	1	6	1	2	6	16	18	6