

Symmetric Chain Decomposition

Junyu Cao

School of Mathematics and Statistics
University of Connecticut

December 5, 2013

Outline

- 1 Symmetric chain decompositions in the Boolean lattice
- 2 Necklace Poset
- 3 Other quotient posets
 - Transposition
 - Group generated by transposition
 - $B_n/(1..n-1)$
 - i-cycle
- 4 Summary

Definition

A chain in a poset, $(P, <)$, is a **totally ordered subset** of P .

Definition

A chain in a poset, $(P, <)$, is a **totally ordered subset** of P .

Definition

A chain in a poset, $(P, <)$, is a **totally ordered subset** of P .

A **saturated chain** is a chain $x_1 < \dots < x_k$ such that x_i covers x_{i-1} for each $i > 1$.

Definition

A chain in a poset, $(P, <)$, is a **totally ordered subset** of P .

A **saturated chain** is a chain $x_1 < \dots < x_k$ such that x_i covers x_{i-1} for each $i > 1$.

A poset is **ranked** if it satisfies: for any $x < y$, all saturated chains from x to y have the same length. Denote the rank of x as $r(x)$.

Definition

A chain in a poset, $(P, <)$, is a **totally ordered subset** of P .

A **saturated chain** is a chain $x_1 < \dots < x_k$ such that x_i covers x_{i-1} for each $i > 1$.

A poset is **ranked** if it satisfies: for any $x < y$, all saturated chains from x to y have the same length. Denote the rank of x as $r(x)$.

The saturated chain $x_1 < x_2 < \dots < x_k$ is a **symmetric chain** in P if $r(x_1) + r(x_k) = r(P)$.

Example

$A = \{1, 3, 4, 6, 8, 9\} \in B_{10}$, then $S_A = 1011010110$ and the bracket representation is $)()()()()$

Example

$A = \{1, 3, 4, 6, 8, 9\} \in B_{10}$, then $S_A = 1011010110$ and the bracket representation is $)()()()()$

Example

$A = \{1, 3, 4, 6, 8, 9\} \in B_{10}$, then $S_A = 1011010110$ and the bracket representation is $)()()()()$
 $\tau(S_A) = 1011010111$

Example

$A = \{1, 3, 4, 6, 8, 9\} \in B_{10}$, then $S_A = 1011010110$ and the bracket representation is $)()()()()$

$$\tau(S_A) = 1011010111$$

$$\tau^{-1}(S_A) = 1011010100$$

Example

$A = \{1, 3, 4, 6, 8, 9\} \in B_{10}$, then $S_A = 1011010110$ and the bracket representation is $)()()()()$

$$\tau(S_A) = 1011010111$$

$$\tau^{-1}(S_A) = 1011010100$$

Definition

U_0 denotes the set of positions of unmatched zeros

Definition

U_0 denotes the set of positions of unmatched zeros

Definition

U_0 denotes the set of positions of unmatched zeros

U_1 denotes the set of positions of unmatched ones

Definition

U_0 denotes the set of positions of unmatched zeros

U_1 denotes the set of positions of unmatched ones

$M(x) := \{(a, b) : \text{a zero in position } a \text{ is matched to a one in position } b\}$

Definition

U_0 denotes the set of positions of unmatched zeros

U_1 denotes the set of positions of unmatched ones

$M(x) := \{(a, b) : \text{a zero in position } a \text{ is matched to a one in position } b\}$

Theorem


For a x in B_n with $|U_0(x)| = k$, let

$C_x = \{x, \tau(x), \tau^2(x), \dots, \tau^k(x)\}$. The following is a symmetric chain decomposition of B_n :

$$S = \{C_x | x \in B_n, U_1(x) = \emptyset\}.$$

—Greene and Kleitman

Example



Example

Symmetric Chain:

0000-1000-1100-1110-1111

0100-0110-0111

0010-1010-1011

0001-1001-1101

00010

00001

Theorem

For all positive integers n , $B_n/(1\dots n)$ has a symmetric chain decomposition.

–K. K. Jordan

Outline

- 1 Symmetric chain decompositions in the Boolean lattice
- 2 Necklace Poset
- 3 Other quotient posets
 - Transposition
 - Group generated by transposition
 - $B_n/(1..n-1)$
 - i-cycle
- 4 Summary

Theorem

If $G = B_n/(i, j)$, G has a symmetric chain decomposition.

Theorem

If $G = B_n/(i, j)$, G has a symmetric chain decomposition.

Theorem

If $G = B_n/(i, j)$, G has a symmetric chain decomposition.

Proof

Define the anti-sequence of $10a_3a_4\dots a_n$ is $01a_3a_4\dots a_n$ (and vice-versa).

If we remove all chains with anti-sequence of elements of the form $10a_3a_4\dots a_n$, the remain is the symmetric chain decomposition of G .

Proof

For any $s_1 = 10a_3a_4\dots a_n$ and $s_2 = 01a_3a_4\dots a_n$

Proof

For any $s_1 = 10a_3a_4\dots a_n$ and $s_2 = 01a_3a_4\dots a_n$


- case1 $\tau(s_1) = 11a_3a_4\dots a_n$

Proof

For any $s_1 = 10a_3a_4\dots a_n$ and $s_2 = 01a_3a_4\dots a_n$

- case1 $\tau(s_1) = 11a_3a_4\dots a_n$
- case2 $\tau(s_1) = 10b_3b_4\dots b_n$

Example



Example

symmetric chain decomposition:

0000-1000-1100-1110-1111

0010-1010-1011

0001-1001-1101

0101

Outline


- 1 Symmetric chain decompositions in the Boolean lattice
- 2 Necklace Poset
- 3 Other quotient posets
 - Transposition
 - Group generated by transposition
 - $B_n/(1..n-1)$
 - i-cycle
- 4 Summary

Corollary

If G is generated by $(a_1, b_1), (a_2, b_2), \dots, (a_i, b_i)$, B_n/G has a symmetric chain decomposition.

Example

$B_4/(12)(34)$



Example

Symmetric chain decomposition:

0000-1000-1100-1110-1111

0010-1010-1011

0101

Outline

- 1 Symmetric chain decompositions in the Boolean lattice
- 2 Necklace Poset
- 3 Other quotient posets
 - Transposition
 - Group generated by transposition
 - $B_n/(1..n-1)$
 - i-cycle
- 4 Summary

Theorem

$B_n/(1..n-1)$ is two copies of $B_{n-1}/(1..n-1)$. There is an edge between two parts if and only if the vertices are $a_1 a_2 \dots a_k$ and $a_1 a_2 \dots a_k n$ where $a_1, \dots, a_k < n$.

Theorem

We use the following way to generate symmetric chains:

Theorem

We use the following way to generate symmetric chains:

- 1 For two parts of $B_n/(1..n-1)$, use the same way in $B_{n-1}/(1..n-1)$ to generate chains.

Theorem

We use the following way to generate symmetric chains:

- 1 For two parts of $B_n/(1..n-1)$, use the same way in $B_{n-1}/(1..n-1)$ to generate chains.
- 2 Eliminate the top edge of every chain in the second part of $B_n/(1..n-1)$.

Theorem

We use the following way to generate symmetric chains:

- 1 For two parts of $B_n/(1..n-1)$, use the same way in $B_{n-1}/(1..n-1)$ to generate chains.
- 2 Eliminate the top edge of every chain in the second part of $B_n/(1..n-1)$.
- 3 Add the edge between $a_1 a_2 \dots a_k$ and $a_1 a_2 \dots a_k n$ where $a_1 a_2 \dots a_k$ is the top vertex of chains in the first part.

Proof

$$X_1 = B_{n-1}/(1..n-1), X_2 = B_n/(1..n-1) - X_1$$

Proof

$$X_1 = B_{n-1}/(1..n-1), X_2 = B_n/(1..n-1) - X_1$$

Proof

$$X_1 = B_{n-1}/(1..n-1), X_2 = B_n/(1..n-1) - X_1$$

$$f : X_1 \rightarrow X_2$$

$$f(a_1 a_2 \dots a_k) = a_1 a_2 \dots a_k n, f \text{ is bijective}$$

Proof

$$X_1 = B_{n-1}/(1..n-1), X_2 = B_n/(1..n-1) - X_1$$

$$f : X_1 \rightarrow X_2$$

$$f(a_1 a_2 \dots a_k) = a_1 a_2 \dots a_k n, f \text{ is bijective}$$

Based on previous theorem, we can generate chains in X_1 and X_2 separately.

Proof

$$X_1 = B_{n-1}/(1..n-1), X_2 = B_n/(1..n-1) - X_1$$

$$f : X_1 \rightarrow X_2$$

$$f(a_1 a_2 \dots a_k) = a_1 a_2 \dots a_k n, f \text{ is bijective}$$

Based on previous theorem, we can generate chains in X_1 and X_2 separately.

guarantee the symmetric property

Proof

$$X_1 = B_{n-1}/(1..n-1), X_2 = B_n/(1..n-1) - X_1$$

$$f : X_1 \rightarrow X_2$$

$$f(a_1 a_2 \dots a_k) = a_1 a_2 \dots a_k n, f \text{ is bijective}$$

Based on previous theorem, we can generate chains in X_1 and X_2 separately.

guarantee the symmetric property

→ increase one edge or decrease one edge of every chain

Proof

$$X_1 = B_{n-1}/(1..n-1), X_2 = B_n/(1..n-1) - X_1$$

$$f : X_1 \rightarrow X_2$$

$$f(a_1 a_2 \dots a_k) = a_1 a_2 \dots a_k n, f \text{ is bijective}$$

Based on previous theorem, we can generate chains in X_1 and X_2 separately.

guarantee the symmetric property

→ increase one edge or decrease one edge of every chain


eliminate the top edge of every chain of X_2

connect every top element s of X_1 with $f(s)$


Theorem

$B_n/(1..n-1)$ is two copies of $B_{n-1}/(1..n-1)$. There is an edge between two parts if and only if the vertices are $a_1 a_2 \dots a_k$ and $a_1 a_2 \dots a_k n$ where $a_1, \dots, a_k < n$.

Example



Example



Outline

- 1 Symmetric chain decompositions in the Boolean lattice
- 2 Necklace Poset
- 3 Other quotient posets
 - Transposition
 - Group generated by transposition
 - $B_n/(1..n-1)$
 - i-cycle
- 4 Summary

Theorem

For any $i \geq 2$, the poset $P = B_n/(123\dots i)$ has a symmetric chain decomposition. We can use the same way mentioned in the former theorem to generate this chains.

Theorem

For any $i \geq 2$, the poset $P = B_n/(123\dots i)$ has a symmetric chain decomposition. We can use the same way mentioned in the former theorem to generate this chains.

Proof

Use induction to prove.

Summary

$B_n/(1..i)$ has a symmetric chain decomposition

Summary

$B_n/(1..i)$ has a symmetric chain decomposition

Summary

$B_n/(1..i)$ has a symmetric chain decomposition

unproved: B_n/G has a symmetric chain decomposition in
general case

Summary

$B_n/(1..i)$ has a symmetric chain decomposition

unproved: B_n/G has a symmetric chain decomposition in
general case

key idea: the structure of chains cannot be changed

Thanks for listening