

BASE EXTENSION AND EXTERIOR POWERS

KEITH CONRAD

Let $f: R \rightarrow S$ be a ring homomorphism, through which we can base extend R -modules to S -modules: $M \rightsquigarrow S \otimes_R M$ and $\Lambda^k(M) \rightsquigarrow S \otimes_R \Lambda^k(M)$. We will write $\Lambda^k(M)$ as $\Lambda_R^k(M)$ to emphasize the dependence on the scalar ring, since we will be working with exterior powers of both R -modules and S -modules.

Theorem 1. *Base extension commutes with formation of exterior powers: for any R -module M and integer $k \geq 1$, there is a unique S -module isomorphism $S \otimes_R \Lambda_R^k(M) \cong \Lambda_S^k(S \otimes_R M)$ where*

$$s \otimes (m_1 \wedge_R \cdots \wedge_R m_k) \mapsto s((1 \otimes m_1) \wedge_S \cdots \wedge_S (1 \otimes m_k)).$$

Moreover, if $M \xrightarrow{\varphi} N$ is a linear map of R -modules then the diagram

$$\begin{array}{ccc} S \otimes_R \Lambda_R^k(M) & \xrightarrow{1 \otimes \wedge^k(\varphi)} & S \otimes_R \Lambda_R^k(N) \\ \downarrow & & \downarrow \\ \Lambda_S^k(S \otimes_R M) & \xrightarrow{\wedge^k(1 \otimes \varphi)} & \Lambda_S^k(S \otimes_R N) \end{array}$$

commutes.

For $k = 0$ the S -modules $S \otimes_R \Lambda_R^0(M) = S \otimes_R R$ and $\Lambda_S^0(S \otimes_R M) = S$ are naturally isomorphic, but the formula in Theorem 1 makes no sense for $k = 0$ so we omitted that case from the theorem.

Proof. For $k = 1$ the indicated function is $S \otimes_R M \rightarrow S \otimes_R M$ by $s \otimes m \mapsto s(1 \otimes m) = s \otimes m$, so it is just the identity map. From now on let $k \geq 2$. We want to write down S -linear maps $S \otimes_R \Lambda_R^k(M) \rightarrow \Lambda_S^k(S \otimes_R M)$ and $S \otimes_R \Lambda_R^k(N) \rightarrow \Lambda_S^k(S \otimes_R N)$ that are inverses of each other.

To write down an S -linear map

$$S \otimes_R \Lambda_R^k(M) \rightarrow \Lambda_S^k(S \otimes_R M),$$

we first ignore the S -part on the left and concentrate on getting an R -linear map

$$(1) \quad \Lambda_R^k(M) \rightarrow \Lambda_S^k(S \otimes_R M),$$

which we will then turn into an S -linear map out of $S \otimes_R \Lambda_R^k(M)$ using extension of scalars. The function we want to use is this:

$$(2) \quad m_1 \wedge_R \cdots \wedge_R m_k \mapsto (1 \otimes m_1) \wedge_S \cdots \wedge_S (1 \otimes m_k).$$

Strictly speaking, the rule (2) on its own does not *define* an R -linear map out of $\Lambda_R^k(M)$ because the elements of $\Lambda_R^k(M)$ are not all elementary wedge products, although there is at most one R -linear map with the effect (2) since elementary wedge products additively span

$\Lambda_R^k(M)$. To construct an R -linear map (1), we write down an alternating R -multilinear map out of M^k . Define $M^k \rightarrow \Lambda_S^k(S \otimes_R M)$ by

$$(m_1, \dots, m_k) \mapsto (1 \otimes_R m_1) \wedge_S \cdots \wedge_S (1 \otimes_R m_k).$$

This function is alternating and R -multilinear (treat the target $\Lambda_S^k(S \otimes_R M)$ as an R -module in the way every S -module is an R -module: restriction of scalars). Therefore the universal mapping property of $\Lambda_R^k(M)$ gives us an R -linear map $\Lambda_R^k(M) \rightarrow \Lambda_S^k(S \otimes_R M)$ satisfying (2) on elementary wedge products.

Now we use the universal mapping property of extension of scalars: from the R -linear map $\Lambda_R^k(M) \rightarrow \Lambda_S^k(S \otimes_R M)$ in (1) there is an S -linear map $\varphi: S \otimes_R \Lambda_R^k(M) \rightarrow \Lambda_S^k(S \otimes_R M)$ such that

$$(3) \quad \varphi(s \otimes_R (m_1 \wedge_R \cdots \wedge_R m_k)) = s((1 \otimes_R m_1) \wedge_S \cdots \wedge_S (1 \otimes_R m_k)).$$

This is the formula we want in the theorem.

To get an S -linear map in the other direction, $\Lambda_S^k(S \otimes_R M) \rightarrow S \otimes_R \Lambda_R^k(M)$, we want to write down an appropriate alternating S -multilinear map $(S \otimes_R M)^k \rightarrow S \otimes_R \Lambda_R^k(M)$ and then use the universal mapping property of $\Lambda_S^k(S \otimes_R M)$. Consider the composite map

$$(4) \quad (S \otimes_R M)^k \xrightarrow{\otimes_S} (S \otimes_R M)^{\otimes_S k} \cong S \otimes_R (M^{\otimes_R k}) \xrightarrow{1 \otimes \wedge_R} S \otimes_R \Lambda_R^k(M).$$

The first map is the canonical multilinear map defining tensor powers (of S -modules), the isomorphism in the middle is the inverse of the canonical one for base extensions of tensor products (recall for two R -modules M and N that $S \otimes_R (M \otimes_R N) \cong (S \otimes_R M) \otimes_S (S \otimes_R N)$ by $s \otimes (m \otimes n) \mapsto s(1 \otimes m) \otimes (1 \otimes n)$), and the third map is the base extension to S -modules of the canonical R -linear map $\wedge_R: M^{\otimes_R k} \rightarrow \Lambda_R^k(M)$. The first map is S -multilinear and the second and third are S -linear, so the composite map is S -multilinear. Its effect on a k -tuple of elementary tensors is

$$\begin{aligned} (s_1 \otimes_R m_1, \dots, s_k \otimes_R m_k) &\mapsto (s_1 \otimes_R m_1) \otimes_S \cdots \otimes_S (s_k \otimes_R m_k) \\ &\mapsto s_1 \cdots s_k \otimes_R (m_1 \otimes_R \cdots \otimes_R m_k) \\ &\mapsto s_1 \cdots s_k \otimes_R (m_1 \wedge_R \cdots \wedge_R m_k). \end{aligned}$$

Let $\alpha: (S \otimes_R M)^k \rightarrow S \otimes_R \Lambda_R^k(M)$ denote the composite map (4). We want to show α is alternating: $\alpha(t_1, t_2, \dots, t_k) = 0$ when $t_i = t_j$ for some $i \neq j$. Because α is multilinear, it suffices to check such vanishing occurs when two coordinates are equal tensors and the other coordinates are all elementary tensors. To keep the notation simple, we focus on the case $t_1 = t_2$ (other equal coordinates run in the same way):

$$\alpha(t, t, s_3 \otimes_R m_3, \dots, s_k \otimes_R m_k) \stackrel{?}{=} 0.$$

Writing $t = s'_1 \otimes_R m'_1 + \cdots + s'_n \otimes_R m'_n$, we expand the left side as

$$\begin{aligned} \alpha(t, t, s_3 \otimes_R m_3, \dots, s_k \otimes_R m_k) &= \sum_{i,j=1}^n \alpha(s'_i \otimes_R m'_i, s'_j \otimes_R m'_j, s_3 \otimes_R m_3, \dots, s_k \otimes_R m_k) \\ &= \sum_{i,j=1}^n s'_i s'_j s_3 \cdots s_k \otimes_R (m'_i \wedge_R m'_j \wedge_R m_3 \wedge_R \cdots \wedge_R m_k). \end{aligned}$$

In this sum, a term where $i = j$ is 0 because m'_i is repeated in the elementary wedge product. For $i \neq j$, the two terms with i and j in both orders are negatives of each other and thus cancel when added, so $\alpha(t, t, s_3 \otimes_R m_3, \dots, s_k \otimes_R m_k) = 0$.

Since α is alternating multilinear, there is an S -linear map $\psi: \Lambda_S^k(S \otimes_R M) \rightarrow S \otimes_R \Lambda_R^k(M)$ such that

$$\psi((s_1 \otimes m_1) \wedge_S \cdots \wedge_S (s_k \otimes m_k)) = \alpha(s_1 \otimes m_1, \dots, s_k \otimes m_k) = s_1 \cdots s_k \otimes (m_1 \wedge_R \cdots \wedge_R m_k).$$

It is left to the reader to show φ and ψ are inverses of each other. \square

Example 2. For any ideal \mathfrak{a} , $R/\mathfrak{a} \otimes_R \Lambda_R^k(M) \cong \Lambda_{R/\mathfrak{a}}^k(R/\mathfrak{a} \otimes_R M)$ as R/\mathfrak{a} -modules, and for any prime ideal \mathfrak{p} , $R_{\mathfrak{p}} \otimes_R \Lambda_R^k(M) \cong \Lambda_{R_{\mathfrak{p}}}^k(R_{\mathfrak{p}} \otimes_R M)$ as $R_{\mathfrak{p}}$ -modules.