

POLYHEXANIDE AND NORWEGIAN SPRUCE RESIN ACIDS FOR SKIN BURNS OF INTERMEDIATE THICKNESS OF THE FACE IN CHILDREN

JAROSŁAW OLSZYNA

Centrum Leczenia Oparzeń, Samodzielny Publiczny Specjalistyczny Zakład Opieki Zdrowotnej „Zdroje”, ul. Mączna 4, 70-780 Szczecin, e-mail: j.olszyna@zoz-zdroje.pl Wpłynęło: 20.08.2019
Zaakceptowano: 10.09.2019 DOI: dx.doi.org/10.15374/ChPiO2019026

ABSTRACT: Introduction Burn injuries are a significant problem for healthcare systems as they lead to permanent damage of the body, its deformation, disability and sometimes may even cause death. Burns occur twice as often in children compared to adults and usually affect small patients up to the age of four. Often, burns not only cause local effects in the form of loss of skin function, but also burn disease. Modern preparations and multi-faceted management of the burn wound have significantly improved wound healing and reduced mortality. Aim The aim of this study was to apply polyhexanide and resinous Norwegian spruce acids in pediatric patients (N=19) who suffered burns of the facial skin of intermediate depth and to determine the effect of these preparations on the wound healing processes, as well as to determine the degree of pain reduction. Material and methods This study uses the case study research method. The following materials were analyzed: medical records of patients treated at the CLO in Szczecin, photographic records, medical interviews with patients and their family, observation of wounds. The FLACC behavioral and VAS visual-analogue scale was used to assess the severity of pain. Conclusions The use of polyhexanide and resinous Norwegian spruce acids resulted in maintaining adequate moisture of the wound and affected the process of granulation and epithelialization. It also reduced pain during hospitalization. These preparations can be used in the TIME treatment strategy. **KEY WORDS:** burns, burns in children, polyhexanide, resinous acids of Norwegian spruce