

补充题: 1. 令 $\varphi(R) = \frac{1}{2\pi R} \int_{L_R} u(x, y) ds$ $\begin{matrix} x = x_0 + R \cos \theta \\ y = y_0 + R \sin \theta \end{matrix}$ $\frac{1}{2\pi} \int_0^{2\pi} u(x_0 + R \cos \theta, y_0 + R \sin \theta) d\theta \Rightarrow \varphi'(R) = \frac{1}{2\pi} \int_0^{2\pi} u_x(x_0 + R \cos \theta, y_0 + R \sin \theta) (-\sin \theta) + u_y(x_0 + R \cos \theta, y_0 + R \sin \theta) (\cos \theta) d\theta = \frac{1}{2\pi R} \int_{L_R} u_x(x, y) dy + u_y(x, y) dx$

格林 $\frac{1}{2\pi R} \int_{B((x_0, y_0), R)} u_{xx} + u_{yy} dx dy = 0 \Rightarrow \varphi(R)$ 为常值

从而 $\varphi(R) = \lim_{R \rightarrow 0^+} \varphi(R) = u(x_0, y_0)$

2. 若 $u(x, y)$ 在 (x_0, y_0) 内取到最大值, 则 $\forall 0 < R < \text{dist}((x_0, y_0), \partial D)$

$$u(x_0, y_0) = \frac{1}{2\pi R} \int_{L_R} u(x, y) ds \leq \frac{1}{2\pi R} \int_{L_R} u(x_0, y_0) ds = u(x_0, y_0) \quad (*)$$

故 $u(x, y) \equiv u(x_0, y_0), (x, y) \in L_R$, 由 D 的连通性知 $\forall (x, y) \neq (x_0, y_0)$

存在一条落于 D 内的曲线 l 连结 (x_0, y_0) 与 (x, y) , 记 $d = \text{dist}(l, \partial D)$

则 $\forall (x, y) \in l, B((x, y), \frac{d}{5}) \subset D \Rightarrow B((x, y), d) \cap l$ 上 $u \equiv u(x_0, y_0)$
 $0 < s < d$

故又 $\{B((x, y), \frac{s d}{2})\}_{(x, y) \in l}$ 为紧集 l 的一个开覆盖, 故存在 l 上有限个

点 $(x_1, y_1), \dots, (x_n, y_n)$ 使得 $l \subset \bigcup_{i=1}^n B((x_i, y_i), \frac{d}{2})$ 且不妨设

$(x_{i+1}, y_{i+1}) \in B((x_i, y_i), d)$ 不断对 (x_i, y_i) 重复 $(*)$ 中讨论可得

$\forall (x, y) \in l, u(x, y) \equiv u(x_0, y_0)$, 从而 $u(x, y) \equiv u(x_0, y_0), (x, y) \in D$

这与 $u(x, y)$ 非常数矛盾! 而 $-u$ 也是调和函数不可在 D 内取最大值从而 u 不可在 D 内取最小值.

11.3.6

(1) 原式 $\begin{matrix} x = a \cos \theta \\ y = a \sin \theta \end{matrix} \int_{\pi}^0 \frac{-a \sin \theta (-a \sin \theta) + a \cos \theta (a \cos \theta)}{a^2} d\theta = -\pi$

(2) 记 L_1 为从 B 到 $(\frac{1}{2}, 0)$ 的直线段, L_2 为从 $(\frac{1}{2}, 0)$ 沿圆周 $y = \sqrt{\frac{1}{4} - x^2}$ 到 $(-\frac{1}{2}, 0)$ 的圆弧, L_3 为从 $(-\frac{1}{2}, 0)$ 到 A 的直线段

$$\text{故 } \int_{L_1+L_2+L_3} \frac{-ydx+x dy}{x^2+y^2} \stackrel{\text{Green}}{\underset{\text{公式}}{=}} \iint_D \left(\frac{x}{x^2+y^2} \right)_x - \left(\frac{-y}{x^2+y^2} \right)_y dx dy = 0$$

$$\Rightarrow \int_{L_1} \frac{-ydx+x dy}{x^2+y^2} = \int_{L_3} \frac{-ydx+x dy}{x^2+y^2} = 0, \int_{L_2} \frac{-ydx+x dy}{x^2+y^2} \stackrel{(1)}{=} \pi$$

$$\text{故 } \int_L \frac{-ydx+x dy}{x^2+y^2} = - \int_{L_1+L_2+L_3} \frac{-ydx+x dy}{x^2+y^2} = -\pi$$

11.3.7 (1) $\oint_L \frac{\partial f}{\partial n} ds = \oint_L f_1 \vec{n}_1 + f_2 n_2 ds = \oint_L f_1 dy - f_2 dx$

$$\stackrel{\text{Green}}{\underset{\text{公式}}{=}} \oint \iint_D f_{11} - (-f_2)_2 dx dy = \iint_D \Delta f dx dy$$

$$(2) \oint_L \cos(a, n) ds = \oint_L \vec{a} \cdot \vec{n} ds = \oint_L \frac{\partial(a_1 x + a_2 y)}{\partial \vec{n}} ds = 0$$

$$(3) \text{同(1)证明有 } \oint_L \vec{F} \cdot \vec{n} ds = \oint_L -f_2 dx + f_1 dy = \iint_D \operatorname{div} \vec{F} dx dy$$

$$\text{故 } \oint_L (v \frac{\partial u}{\partial n} - u \frac{\partial v}{\partial n}) ds = \oint_L (v \nabla u - u \nabla v) \cdot \vec{n} ds = \iint_D \operatorname{div}(v \nabla u - u \nabla v) dx dy \\ = \iint_D (v \Delta u - u \Delta v) dx dy$$

11.4.1

$$(2) r(\theta, y) = (R \cos \theta, y, R \sin \theta), r_\theta = (-R \sin \theta, 0, R \cos \theta), r_y = (0, 1, 0)$$

$$(-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}, 0 \leq y \leq h), \text{ 则原式 } = \iint_D \begin{vmatrix} 0 & 0 & y R^2 \sin \theta \cos \theta \\ -R \sin \theta & 0 & R \cos \theta \\ 0 & 1 & 0 \end{vmatrix} d\theta dy \\ = -R^3 \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \sin^2 \theta \cos \theta d\theta \cdot \int_0^h y dy = \frac{1}{3} R^3 h^2$$

$$(4) r(\theta, \varphi) = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta) \quad (0 \leq \theta \leq \frac{\pi}{2}, 0 \leq \varphi \leq 2\pi)$$

$$\frac{\partial(z, x)}{\partial(\theta, \varphi)} = \begin{vmatrix} -\sin \theta & 0 \\ \cos \theta \cos \varphi & -\sin \theta \sin \varphi \end{vmatrix} = \sin^2 \theta \sin \varphi$$


$$\text{故原式} = \int_0^{\frac{\pi}{2}} d\theta \int_0^{2\pi} \sin^3 \theta \cos \theta \sin^2 \varphi d\varphi = \frac{\pi}{4}$$

$$(6) \gamma(\theta, z) = (z \cos \theta, z \sin \theta, z) \quad (0 \leq \theta \leq 2\pi, 0 \leq z \leq 1)$$

$$\begin{aligned} \text{原式} &= \int_0^{2\pi} d\theta \int_0^1 \begin{vmatrix} z \sin \theta - z & z - z \cos \theta & z \cos \theta - z \sin \theta \\ -z \sin \theta & z \cos \theta & 0 \\ \cos \theta & \sin \theta & 1 \end{vmatrix} dz \\ &= \int_0^{2\pi} (\cos \theta - \sin \theta) d\theta \cdot \int_0^1 2z^2 dz = 0 \end{aligned}$$

$$\begin{aligned} (7) \text{ 记 } D &= \{(x, y, 0) \mid x^2 + y^2 \leq a^2\}, \text{ 则 } \iint_S = \int_{S+D} x^2 z^2 dy dz + x^2 y dz dx \\ &+ y^2 z dx dy \xrightarrow{\text{Gauss}} \iiint_{\Omega} (z^2 + x^2 + y^2) dx dy dz = \frac{1}{2} \iiint_{B(0,a)} x^2 + y^2 + z^2 dx dy dz \\ &= \frac{1}{2} \int_0^a ds \int_{\partial B(0,s)} s^2 d\sigma = \frac{1}{2} \int_0^a 4\pi s^4 ds = \frac{2\pi}{5} a^5 \end{aligned}$$

11.5.1 (2) $\iint_{S+S_1+S_2+S_3} \xrightarrow{\text{Gauss}} \iiint_V (y+z+x) dx dy dz = \int_0^1 ds \int_{x+y+z=s} s d\sigma = \frac{\sqrt{3}}{16} \cdot \frac{1}{8}$



$\Rightarrow \iint_{S+S_1+S_2+S_3} = 0 \Rightarrow \text{原式} = \frac{\sqrt{3}}{16} \cdot \frac{1}{8}$ (以下“—”均代表题目中的被积式)

$$\begin{aligned} (5) \text{ 记 } D &= \{(x, y, 1) \mid x^2 + y^2 \leq 1\}, \text{ 则 } \iint_{S+D} \xrightarrow{\text{Gauss}} \iiint_V 3 dx dy dz \quad (D \text{ 取上侧}) \\ &= \int_0^1 dz \iint_{x^2+y^2 \leq z} 3 dx dy = \int_0^1 3\pi z dz = \frac{3}{2}\pi \end{aligned}$$

$$\begin{aligned} \text{而 } \iint_D \text{ — } &= \iint_{x^2+y^2 \leq 1} (x^2 + y^2 - y) dx dy \xrightarrow{\substack{x=r\cos\theta \\ y=r\sin\theta}} \int_0^1 dr \int_0^{2\pi} (r^2 - r \sin \theta) d\theta = \pi \\ \text{故原式} &= \iint_{S+D} \text{ — } - \iint_D \text{ — } = \frac{3\pi}{2} - \pi = \frac{\pi}{2}. \end{aligned}$$

11.5.3 设 ~~V 为 B~~ S_1 为 $B(0, \frac{1}{2})$ 表面, 取指向原点心一侧 ($r = \sqrt{x^2 + y^2 + z^2}$)

$$\text{则 } \iint_{S_1} \text{ — } = \iint_{S_1} \frac{1}{r^3} (x, y, z) \cdot -\frac{1}{r} (x, y, z) dS = \iint_{S_1} -4 dS = -4\pi$$

$$\text{而 } \iint_{S+S_1} \xrightarrow{\text{Gauss}} \iiint_{\Omega} \left(\frac{x}{r^3} \right)_x + \left(\frac{y}{r^3} \right)_y + \left(\frac{z}{r^3} \right)_z dx dy dz = 0$$

$$\text{故原式} = \iint_{S+S_1} \text{ — } - \int_{S_1} = 4\pi$$

$$11.5.4 \quad 0 = \oint_S \text{---} \stackrel{\text{Gauss}}{=} \iiint_V (xf)_x - (xyf)_y - (e^{2x}z)_z \, dx \, dy \, dz$$

$$= \iiint_V [xf'(x) + (1-x)f(x) - e^{2x}] \, dx \, dy \, dz \quad \text{在 } x > 0 \text{ 中任一区域 } V \text{ 成立}$$

故 $xf'(x) + (1-x)f(x) - e^{2x} = 0, \forall x > 0 \Rightarrow f(x) = \frac{e^x(e^x-1)}{x} + \frac{(c+1)e^x}{x}$

由 $\lim_{x \rightarrow 0^+} \frac{e^x}{x}$ 不存在知 $c = -1$, $f(x) = \frac{e^x(e^x-1)}{x}$

$$11.5.6 \quad \iint_S \cos \langle \vec{c}, \vec{n} \rangle \, ds = \frac{1}{|\vec{c}|} \iint_S \vec{c} \cdot \vec{n} \, ds = \frac{1}{|\vec{c}|} \iiint_V \operatorname{div} \vec{c} \, dx \, dy \, dz = 0$$

11.5.9 (2) 设 $x = a \cos t, y = a \sin t, z = h - h \cos t$ (从 $(a, 0, 0)$ 回到原处时 t 从 0 变化到 2π), 故原式 $= \int_0^{2\pi} (a \sin t - h + h \cos t)(-a \sin t) + (h - h \cos t - a \cos t)(a \cos t) + (a \cos t - a \sin t)(h \sin t) \, dt$

$$= \int_0^{2\pi} [-a^2 - ha + ha(\sin t + \cos t)] \, dt = -2\pi a(h+a)$$

(4) 设 Σ 为 $y = z$ 被 $x^2 + y^2 \leq 2y$ 所截部分, 则由 Stokes 公式, $\vec{n} = (0, \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$

$$\text{原式} = \iint_{\Sigma} \begin{vmatrix} 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y^2 & xy & xz \end{vmatrix} \, dS = \frac{1}{\sqrt{2}} \iint_{\Sigma} (y-z) \, dS = 0$$

11.5.10. (1) 圆面法向量 $\vec{n} = (0, 0, 1)$, 则 $\oint_L - = \iint_S \begin{vmatrix} 0 & 0 & 1 \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^2y^2 & 1 & z \end{vmatrix} \, d\sigma$

$$= \iint_S -3x^2y^2 \, d\sigma \quad \begin{matrix} x = r \cos \theta \\ y = r \sin \theta \end{matrix} \int_0^R dr \int_0^{2\pi} -3r^4 \cos^2 \theta \sin^2 \theta \cdot r \, d\theta = -\frac{\pi}{8} R^6$$

(2) 半球面法向量 $\vec{n} = \frac{1}{R}(x, y, z)$, 同理有 $\oint_L - = \iint_S \operatorname{rot} \vec{v} \cdot \vec{n} \, d\sigma$

$$= \frac{1}{R} \iint_S (x - 3x^2y^2z) \, d\sigma \quad \begin{matrix} x = R \sin \theta \cos \varphi \\ y = R \sin \theta \sin \varphi \\ z = R \cos \theta \end{matrix} \int_0^{\frac{\pi}{2}} d\theta \int_0^{2\pi} \frac{1}{R} (R \sin \theta \cos \varphi - 3R^5 \sin^6 \theta \cos \theta \cos^2 \varphi \sin^2 \varphi) R^2 \sin \theta \, d\theta \, d\varphi = -\frac{\pi}{8} R^6 \quad \text{结果相同}$$

11.7.3 (3) 由 $\nabla \times \vec{v} = (R_y - Q_z)\vec{i} + (P_z - R_x)\vec{j} + (Q_x - P_y)\vec{k} = 0$

知 v 为有势场, $\varphi(r, \theta, \phi) = \int_{(0,0,0)}^{(r,\theta,\phi)} r^2 \vec{r} \, d\vec{r} + C = \frac{1}{4} r^4 + C$

$$11.7.4. \nabla \times \vec{F} = (2-2a)x, (1-a)y + (3a-3)z + 5-5a = 0 \Rightarrow a=1$$

$$\text{故 } \varphi(x, y, z) = \int_{(0,0,0)}^{(x,y,z)} \vec{F} \cdot d\vec{r} + C = \int_{(0,0,0)}^{(x,y,z)} (x^2 + 5y + 3yz) dx + (5x + 3xz - 2) dy + (3xy - 4z) dz \\ = \int_0^x x^2 dx + \int_0^y (5x - 2) dy + \int_0^z (3xy - 4z) dz + C \\ = \frac{1}{3} x^3 + (5x - 2)y + (3xyz - 2z^2) + C$$

$$11.7.5 (1) \begin{cases} u_x = 3x^2 + 6xy^2 \\ u_y = 6x^2y - 4y^3 \end{cases} \Rightarrow u(x, y) = x^3 + 3x^2y^2 - y^4 + C$$

$$11.7.6 (1) P = (x-y), Q = y-x \Rightarrow Q_x = P_y \Rightarrow \text{与路径无关}$$

$$\text{故 } \int_{(0,0)}^{(1,1)} = \int_{(0,0)}^{(0,1)} + \int_{(0,1)}^{(1,1)} = \int_0^1 y dy + \int_0^1 (x-1) dx = 0$$

$$(3) \text{ 当 } (x, y) \neq (0, 0) \text{ 时, } \int_{(1,0)}^{(6,3)} \frac{1}{\sqrt{x^2+y^2}} = \int_{(1,0)}^{(6,3)} d(\sqrt{x^2+y^2}) = \sqrt{x^2+y^2} \Big|_{(1,0)}^{(6,3)} = 3\sqrt{5} - 1$$

$$11.7.7 (2) \text{ 原式 } \frac{\sqrt{x^2+y^2+z^2}=t}{dt = \frac{1}{t}(xdx+ydy+zdz)} \int_{t_0}^t t f(t) dt = 0 \quad t_0 = \sqrt{x_0^2+y_0^2+z_0^2}$$

$$11.7.11 Q_x = (2xy)_y = 2x \Rightarrow Q(x, y) = x^2 + f(y)$$

$$\text{且 } \int_{(0,0)}^{(x,y)} = \int_{(0,0)}^{(x,0)} + \int_{(x,0)}^{(x,y)} = \int_0^y Q(x^2 + f(\bar{y})) d\bar{y} = x^2 y + \int_0^y f(\bar{y}) d\bar{y}$$

$$\Rightarrow \int t^2 + \int_0^1 f(y) dy = t + \int_0^t f(y) dy \xrightarrow{\text{求导}} 2t = 1 + f(t) \Rightarrow Q(x, y) = x^2 + 2y - 1$$

$$11.7.13 (e^x \sin y + x^2 y + f(x) y)_y = (f'(x) + e^x \cos y + 2x)_x$$

$$\Rightarrow x^2 + f(x) + e^x \cos y = f'(x) + e^x \cos y + 2 \Rightarrow f'(x) - f(x) = x^2 - 2$$

$$\Rightarrow f(x) = -x^2 + e^x - e^{-x} \quad \text{由 } \begin{cases} u_x = e^x \sin y + x^2 y + (-x^2 + e^x - e^{-x}) y \\ u_y = -2x + e^x + e^{-x} + e^x \cos y + 2x \end{cases}$$

$$\text{知 } u(x, y) = e^x \sin y + (e^x + e^{-x}) y + C \Rightarrow du = 0 \Rightarrow u = C$$

$$\text{从而方程解为 } e^x \sin y + (e^x + e^{-x}) y = C.$$