


AMI on statin	• At least 2 current AMI Codes (410.* or 411.*) within a 5-Days Window • Confirmed Lab within the same time window defined by ○ Troponin-I$\geq$0.10 ng/ml ○ or Troponin-T$\geq$0.10 ng/ml, ○ or CK-MB/CK ratio$\geq$3.0 and CK-MB $\geq$10.0 ng/mL • Statin prescribed prior to the AMI event $\geq$180 days
1st AMI on statin	• AMI on statin • No any AMI codes (410.*, 411.*, or 412.*) assigned before the AMI event • No AMI mentioned in previous problem list by NLP
Revascularization while on statin	• Any CPT code for angioplasty or stent (see below) • statin prescribed prior to the procedure $\geq$180 days
1st Revascularization while on statin	• Revascularization while on statin • No exclusion codes before starting statin (see below) • No revascularization mentioned in previous problem list by NLP
Statin exposure	• The number of days that the earliest statin prescription prior to the event (AMI/Revascularization)
All-cause mortality	• EMR or social security death index (SSDI) defined mortality (date)


The timeline of AMI on statin



AMI on statins: Case definition (all three conditions required):

- At least two ICD9 code for AMI or other acute and subacute forms of ischemic heart disease within a five-day window (see AMI Case ICD9 codes below)
- Confirmed lab within the same time window (see AMI Case lab names/values below)
- Statin prescribed prior to the AMI event in medical records at least 180 days

The timeline of 1st AMI on statin



1st AMI on statins: Case definition (all four conditions required):

- Meet the above criteria for AMI on statins
- No diagnosis code for AMI, other acute and subacute forms of ischemic heart , or historical AMI assigned previously (see Case exclusion ICD 9 codes below)
- No revascularization CPT codes assigned previously (see Revascularization CPT codes below)
- No MACE found in previous problem list by NLP (see MACE Exclusion keywords listed below)

Revascularization while on statin: Case definition (both conditions required):

- At least one revascularization CPT code (see Revascularization CPT codes below)
- Statin prescribed prior to the revascularization_event in medical records at least 180 days

1st Revascularization while on statin: Case definition (all four conditions required):

- Meet the above criteria for revascularization while on statin
- No diagnosis code for AMI, other acute and subacute forms of ischemic heart , or historical AMI assigned previously (see Case exclusion ICD 9 codes below)
- No revascularization CPT codes assigned previously (see Revascularization CPT codes below)
- No MACE found in previous problem list by NLP (see MACE Exclusion keywords listed below)

Control definition

- Statin prescribed
- No diagnosis code for AMI, other acute and subacute forms of ischemic heart , or historical AMI assigned previously (see Case exclusion ICD 9 codes below)
- No revascularization CPT codes assigned previously (see Revascularization CPT codes below)
- No MACE found in previous problem list by NLP (see MACE Exclusion keywords listed below)