

Incorporation of evolutionary constraint improves genomic prediction of inbred and hybrid phenotype

Jinliang Yang^{*, 1, 2}, Sofiane Mezouk^{*, 2, 3}, Andy Baumgarten[§], Edward S. Buckler[†], Katherine E. Guille[‡], Michael D. McMullen^{‡, **, 1}, Rita H. Mumm^{§§} and Jeffrey Ross-Ibarra^{*, ††, 1}

^{*}Department of Plant Sciences, University of California, Davis, CA 95616, USA, [§]DuPont Pioneer, Johnston, IA 50131, USA, [†]US Department of Agriculture, Agricultural Research Service, Ithaca, NY 14853, USA, [‡]US Department of Agriculture, Agricultural Research Service, Columbia, MO 65211, USA, ^{**}Division of Plant Sciences, University of Missouri, Columbia, MO 65211, USA, ^{§§}Department of Crop Sciences, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA, ^{††}Center for Population Biology and Genome Center, University of California, Davis, CA 95616, USA

ABSTRACT Complementation of deleterious alleles has long been proposed as a major contributor to the hybrid vigor observed in offspring of inbred parents. We test this hypothesis using evolutionary measures of sequence conservation to ask whether incorporating information about putatively deleterious alleles can inform genomic selection (GS) models and improve phenotypic prediction. We measured a number of agronomic traits in both the inbred parents and hybrids of an elite maize partial diallel population. We resequenced the parents of the population, using genomic evolutionary rate profiling (GERP) to identify constrained sites across more than 86 Mb of the genome. We identified haplotype blocks using an identity-by-descent analysis and scored these blocks on the basis of segregating putatively deleterious variants. Incorporating sequence conservation improves prediction accuracies in a five-fold cross-validation experiment for several traits *per se* as well as heterosis for those traits. These results provide strong empirical support for the simple complementation model of heterosis, and demonstrates the utility of incorporating functional annotation and its potential usage in phenotypic prediction and plant breeding.

KEYWORDS heterosis; deleterious; genomic selection; diallel; GERP; maize

The phenomenon of heterosis or hybrid vigor has been observed across many species, from yeast (Shapira *et al.* 2014) to plants (Shull 1908) and vertebrates (Gama *et al.* 2013). A number of hypotheses have been put forth to explain the phenomenon, including gene dosage (Birchler *et al.* 2003), overdominance (East 1936; Schwartz 1973; Krieger *et al.* 2010) or pseudo-overdominance (Graham *et al.* 1997; McMullen *et al.* 2009), and epistasis (Minvielle 1987; Schnell and Cockerham 1992). Complementation of recessive deleterious alleles, however, remains the simplest genetic explanation (Charlesworth and Willis 2009), and one that is supported by considerable empirical evidences (Xiao *et al.* 1995; Frascaroli *et al.* 2007; Huang *et al.* 2015).

~~Deleterious alleles were arisen from new mutations during~~

~~meiosis. In maize, about 90 new mutations were generated per meiosis (Clark *et al.* 2005), majority of which were deleterious according to empirical estimates (Joseph and Hall 2004). In a natural outcross population, the negative effects on fitness of these deleterious alleles make them subject to be selection against, which lead the deleterious alleles to be maintained in a low frequency (Eyre-Walker and Keightley 2007). But the deleterious alleles could not be completely purged. In maize, the total number of mildly deleterious mutations is substantial because of the exponential growth of population size after domestication. The modern breeding probably aims to remove these deleterious mutations and pyramiding beneficial alleles for agronomical purposes. In practice, the relatively homogeneous maize germplasm pool was artificially divided into different heterotic groups (van Heerwaarden *et al.* 2012). It enabled the improvement of germplasm pools to be conducted in a parallel fashion, and therefore, facilitated the breeding efficiency. Using this hybrid breeding approach, One of the maize yield has been steadily improved since~~

Copyright © 2015 by the Genetics Society of America
doi: 10.1534/genetics.XXX.XXXXXX

Manuscript compiled: Monday 29th June, 2015%

¹Corresponding authors: Department of Plant Sciences, University of California, Davis, CA 95616, USA. Email: rossibarra@ucdavis.edu and jolyang@ucdavis.edu

²These authors contributed equally to this work

³Current address: KWS SAAT AG, Grimsehlstr. 31, 37555 Einbeck, Germany

the early 20th century (Duvick 2001). However, removing deleterious mutations in low recombination regions or in tightly linked regions become less effective. Studies indicated that residual heterozygosity correlates negatively with recombination (Gore et al. 2009; McMullen et al. 2009) and the low recombination is effective over long period of time (Haddrill et al. 2007). As a consequence, the deleterious alleles would be accumulated in the low recombination regions, such as the pericentromeric regions in maize, and the vigorous performance could be realized by combining two sets of non-deleterious or beneficial alleles in repulsion state, thus lead to pseudo-overdominance. A recent QTL study identified loci controlling for heterosis are enriched in centromeric regions (Laripe et al. 2012), which partly support this pseudo-overdominance hypothesis. best studied examples of hybrid vigor is that of maize. Although the benefits of cross-fertilization or hybridization had been observed frequently in the past, it was not until the work of Shull, East, and others that it's significance for agriculture was well appreciated (CITE). Now, hybrid seed makes up the vast majority in progress, still reading

Despite the importance of deleterious alleles in contributing to heterosis, they have not been systematically investigated probably because of their low frequencies in the population and mostly exhibiting minor effects. Here, we employed a genomic selection (GS) approach to simultaneously estimate genome-wide deleterious variants in a half diallel population. The diallel population was composed of a set of hybrids, which enabled us to explore different modes of inheritance of the deleterious variants. And the study can be conducted with millions of variants but using relative little sequencing efforts. In our previous study, deleterious SNPs were found to be enriched in a SNP set identified by GWAS (Mezmouk and Ross-Ibarra 2014). The deleterious variants in the study were defined as non-synonymous mutations in the coding regions. Clearly, deleterious variants are not limited to coding regions. Here, we expanded the characterization of deleterious variants to genome-wide using genomic evolutionary rate profiling (GERP) (Cooper et al. 2005). By incorporating GERP information in GS models, we demonstrated the prediction accuracies were significantly improved not only for some traits *per se*, but also for some heterosis transformations (especially for traits exhibiting high levels of hereosis). Further studies indicated that joint effects of deleterious alleles with additive and dominant modes of inheritance may contribute to heterosis.

Materials and Methods

Plant materials and phenotypic data

We selected 12 maize inbred lines, broadly representative of corn belt maize germplasm (Mikel and Dudley 2006), as parents of a partial diallel population. Each parent in a cross was used as both male and female and the resulting seed was bulked (Figure S1). We evaluated the 66 F1 hybrids, 12 inbred parents and two current commercial check hybrids in the field in Urbana, IL over three years (2009-2011) in an incomplete block design with three replicates each year. Plots consisted of four rows, with all observations taken from the inside two rows to minimize effects of shading and maturity differences from adjacent plots. We measured plant height (PHT, in cm), ear height (EHT, in cm), days to 50% silking (DTS), days to 50% pollen shed (DTP), anthesis-silking interval (ASI, in days), grain yield adjusted to 15.5% moisture (adj GY, in bu/A), and test weight (TW, in

pounds). Overall mean phenotypic values for each cross can be found at Table S1(S1).

We estimated Best Linear Unbiased Estimates (BLUES) of the genetic effects in ASReml-R (Gilmour et al. 2009) with the following linear model:

$$Y_{ijkl} = \mu + \zeta_i + \delta_{ij} + \beta_{jk} + \alpha_l + \zeta_i \cdot \alpha_l + \varepsilon$$

where Y_{ijkl} is the phenotypic value of the l^{th} genotype evaluated in the k^{th} block of the j^{th} replicate within the i^{th} year; μ , the overall mean; ζ_i , the fixed effect of the i^{th} year; δ_{ij} , the fixed effect of the j^{th} replicate nested in the i^{th} year; β_{jk} , the random effect of the k^{th} block nested in the j^{th} replicate; α_l , the the fixed genetic effect of the l^{th} individual; $\zeta_i \cdot \alpha_l$, the interaction effect of the l^{th} individual with the i^{th} year; ε , the model residuals.

We estimated best-parent heterosis (BPH) as:

$$BPH_{min,ij} = \hat{G}_{ij} - \min(\hat{G}_i, \hat{G}_j)$$

$$BPH_{max,ij} = \hat{G}_{ij} - \max(\hat{G}_i, \hat{G}_j)$$

where \hat{G}_{ij} , \hat{G}_i and \hat{G}_j are the genetic values of the hybrid and its two parents i and j . BPH_{min} was used instead of BPH_{max} for ASI. what about ear height and DTS? Did you mean plant height? We need to discuss about this. no plant height should be max, ear height maybe should be min? though it is correlated with plant height... happy to discuss

Sequencing and Genotyping

We extracted DNA from the 12 inbred lines following Doyle and Doyle (1987) and sheared the DNA on a Covaris (Woburn, Massachusetts) for library preparation. Libraries were then sequenced. Libraries were prepared using an Illumina paired end library protocol with 180 bp fragments. Libraries were then sequenced at Cornell. Kate Guill. I'll ask. text changed.

We trimmed raw sequence reads for adapter contamination with Scythe (<https://github.com/vsbuffalo/scythe>) and for quality Sofiane: what qual score? do we need to say anything about overlapping reads? and sequence length (≥ 20 nucleotides) with Sickle (<https://github.com/najoshi/sickle>). We mapped filtered reads to the maize B73 reference genome (AGPv2) with bwa-mem (Li and Durbin 2009), keeping reads with mapping quality (MAPQ) higher than 10 and with a best alignment score higher than the second best one for further analyses. We called single nucleotide polymorphisms (SNPs) using the *mpileup* function from the samtools utilities (Li et al. 2009). To deal with known issues with paralogy in maize (Chia et al. 2012), SNPs were filtered to be heterozygote in less than 3 inbred lines, have a mean minor allele depth of at least 4, have a mean depth over all individuals lower than 30 and have missing/heterozygote alleles in fewer than 6 inbred lines.

We used the fastIBD method implemented in BEAGLE (Browning and Browning 2009) to impute missing data and identify regions of identity by descent (IBD) between the 12 inbred lines. We then defined haplotype blocks as contiguous regions within which there were no IBD break points across all pairwise comparisons of the parental lines (Figure S2). IBD blocks at least 1 Kb in size were kept for further analysis.

Genomic selection using IBD blocks incorporated with GERP scores

We used genome-wide estimates of evolutionary constraint (GERP Davydov et al. 2010) estimated by Rodgers-Melnick et al.

(2015). Haplotype blocks were weighted by the summed GERP scores of all deleterious (GERP score > 0) SNPs; blocks with no deleterious SNPs were excluded from further analysis. This estimation was calculated under both additive and dominant modes of inheritance using a custom python script available at (<https://github.com/yangji/zmSNPtools>). For a particular SNP with a GERP score g , the non-reference homozygote was assigned a value of $2g$, the heterozygote a value of g , and the reference homozygote a value of 0. Under the dominant model, both the heterozygote and the non-reference homozygote were assigned a value of g , with the reference homozygote again assigned a value of 0. To conduct prediction, a 5-fold cross-validation method was used, dividing the diallel population randomly into training (80%) and validation sets (20%) 10 times. The BayesC option from GenSel4 (Habier *et al.* 2011) was used for model training, using 41,000 iterations and removing the first 1,000 as burn-in. After model training, prediction accuracies were obtained by comparing the predicted breeding values with the observed phenotypes in the corresponding validation sets. For comparison, GERP scores were circularly shuffled by permuted using 50k SNPs windows SNP (> 100Mb) windows which were circularly shuffled 10 times to estimate a null conservation score for each IBD blocks. Cross-validation experiments using the circularly shuffled permuted data were conducted on the same training and validation sets.

Results

Genetic values, heritability and heterosis

A partial diallel population was created using 12 maize inbred lines (Figure ??a). Two of them are important public inbreds, B73 and Mo17. And the other ten of proprietary inbreds (LH1, LH123HT, LH82, PH207, 4676A, PHG39, PHG47, PHG84, PHJ40, and PHZ51) that have expired from Plant Variety Protection (PVP) and represent much of the lineage of key heterotic germplasm pools used in present-day commercial corn hybrids. From this population, phenotypic data were collected for seven traits of interest during 2009-2011: anthesis-silking interval (ASI, in days), days to 50% pollen shed (DTP), days to 50% silking (DTS), ear height (EHT, in cm), grain yield adjusted to 15.5% moisture (GY, in bu/A), plant height (PHT, in cm), and test weight (TW, in pounds).

Best linear unbiased estimators (BLUEs) for genotypes of the seven traits were derived from mixed linear models (Table S1S1). In the models, all fixed effects were significant (Wald test P value < 0.05) for all traits except ASI, for which the effect of replicates within environments were not significant. As shown in Figure ??b1, BLUE values were normally distributed (normality test P values > 0.05). Broad sense heritability for these traits ranged from 0.65 for ASI to 0.95 for PHT. Using the parental phenotypic data, we then estimated best-parent heterosis (BPH) for each trait. Because the selected inbred lines are commercially relevant and fairly elite in performance, hybrids in this population exhibit relatively low hybrid vigor (overall mean percent BPH = $0.3\% \pm 0.4\%$) for most traits except GY (mean percent BPH = $95\% \pm 16\%$, Figure S3). Finally, general and specific combining ability (GCA and SCA) were estimated following (Falconer and Mackay 1996). GCA and SCA varied among traits (Table S2), but B73, PHG47 and PHG39 showed the greatest GCA for grain yield.




Figure 1 Density plots of the BLUE values for the seven phenotypic traits. On the x-axis, the phenotypic values were normalized.

Evolutionary—Sequence variation and evolutionary constraint information for genomic variants

All twelve inbreds were sequenced to an average depth of $\sim 10\times$, resulting in a filtered set of 13.8 million SNPs. We estimated the allelic error rate using three independent data sets: for all individuals using 41,292 overlapping SNPs on the maize SNP50 bead chip (van Heerwaarden *et al.* 2012); for all individuals using 180,313 overlapping SNPs identified through genotyping by sequencing (GBS) (Romay *et al.* 2013); and for B73 and Mo17 using the 10,426,715 SNP from the HapMap2 project (Chia *et al.* 2012). Compared to corresponding SNPs identified by previous studies, a concordance rate of 99.1% was observed. *Sofiane: can we separate those numbers out by study? or just report for one study and mention that similar rates were seen in other studies? either way it would be nice to know what rate went with what data. also is concordance mean identical genotype? do we have minor allele rate (which is a bit more informative)? if not, skip it.*

More than 86 million bp of the genome were annotated as conserved, with GERP scores > 0. Nonetheless, 506,898 of these sites were found to segregate among the 12 inbred parents of our diallel (Figure 2A and S4). The minor allele frequency of SNPs at conserved sites was negatively correlated with GERP score (Figure 2B; P value < 0.05, $r = -0.8$), consistent with the idea that variants at sites with more positive GERP scores are more deleterious and more strongly impacted by purifying selection.

Incorporation of GERP information improved Phenotypic prediction accuracies

Genomic variants occurred at the evolutionary constraint sites were potentially deleterious. The phenotypic effects of these genetic loads and their contributions to heterosis become an interesting area to explore. However, the population size in this study is relative small and SNPs detected at sites containing high GERP scores are generally in low frequencies. The statistical power to detect the separate effects of these putative deleterious alleles becomes very low. The small sample size of our diallel and the general low frequency of deleterious SNPs precludes association-based approaches to evaluate the impact of variants on phenotypic variation. To alleviate the statistical limitations this limitation, we conceived a haplotype-based approach for GS, which could add up genomic selection approach in which we use estimates of evolutionary constraint




Figure 2 Distribution of GERP scores and relationship between GERP scores and MAFs. (A) Histogram of GERP scores extracted from at ~1.2-1.3 million sites where the GERP scores were available SNPs. (B) Plot of average GERP scores in bins (bin size = 0.01) of minor allele frequency (MAF). The red and grey lines define the regression and its 95% confidence interval.

across the genome (Rodgers-Melnick *et al.* 2015) to sum the individual effects of deleterious alleles in IBD blocks and estimate these IBD blocks simultaneously. To conduct the analysis, first of all, 55,000 IBD blocks were identified, which had an average size of 44,980 bp (ranged from 36 to 10,320,000 bp). IBD blocks having > 1-kb in size and containing > 1 deleterious alleles (SNPs at sites with GERP scores > 0) were kept for further analysis. Secondly, the GERP scores of SNPs in an IBD block were summed under both within IBD blocks under both an additive and dominant models. Those summed GERP scores on IBD blocks were considered as the measurements of the conservation of the haplotypes. More details of this procedure were illustrated in Figure S5 model (see Methods and Figure S5).

A Bayesian-based statistical method (BayesC) (Habier *et al.* 2011) using a 5-fold cross-validation approach was employed for model training. Using a 5-fold cross-validation approach, the prediction accuracies of the real data and circularly shuffled data were compared. As shown in Figure 3, In general, average prediction accuracies were higher using the additive model (mean $r = 0.81$ and 0.49 for traits *per se* ; prediction accuracies were significantly ($FDR < 0.05$) improved and BPH) than the dominant model (mean $r = 0.70$ and 0.42), and accuracies for heterosis traits were lower than for traits *per se* (Table S3). is there a table we can cite that has all these? see edits Incorporating evolutionary constraint information improved prediction accuracy for ASI and PHT when incorporating GERP information in the IBD blocks under the additive model ; prediction accuracy was significantly improved *per se* under an additive model and for ASI under the dominant model ; For heterosis transformation traits (BPH), incorporation of GERP scores improved BPH of a dominant model ($FDR < 0.05$, Figure 3 A and B). GERP scores also improved prediction accuracies of heterosis (BPH) for GY under the additive model and improved BPH of DTP, DTS and TW under the dominant model ($FDR < 0.05$, Figure 3 C and D, Supplementary table 3). In general, the average prediction accuracies were higher using the additive model (mean $r = 0.81$ and 0.49 for traits *per se* and BPH) than the dominant model (mean $r = 0.70$ and 0.42). And the prediction accuracies decreased for predicting heterosis transformations (BPH) as compared to the

predictions for traits *per se*). It was argued that SNPs in genic regions might have higher GERP scores than those in non-genic regions. The circular shuffling permutations may shift the ref/link to supp table 3 please. also, do we need to include FDR here? what are the multiple tests the FDR is correcting for here? move table S3 to above. added FDR. FDR is for correcting multiple traits and multiple transformations. To rule out the possible confounding of high GERP scores to non-genic regions. If that is the case, the approach tended to weigh more on genic SNPs. To rule out this possibility, we elected SNPs with GERP scores > 0 in genic regions only and did the circular shuffling to assign GERP scores to the same set of the selected SNPs. By doing this, the method will not take advantage of genomic positional information any more. Noted that in this study less number of SNPs was selected ($N =$ and genic annotations, we re-permuted the data using only deleterious (GERP > 0) genic SNPs. Though this resulted in fewer SNPs (316, 983). Nevertheless, the model prediction accuracies were remained significantly improved for traits GY *per se* of GY under the additive model. For heterosis transformations, prediction accuracies were significantly improved and for BPH of GY and PHT under the additive model and the prediction accuracy was significantly improved for pBPH of GY (Figure S6 and Table S4S4).




Figure 3 Beanplots of cross-validation accuracies using SNPs with positive GERP scores. Cross-validation experiments were conducted using selected SNPs and circularly shuffled permuted data from the same set of SNPs for traits *per se* (A, B) and BPH (C, D) and pBPH (E, F) under additive (A, C, E and C) and dominant (B, D, F and D) models. Accuracies from the real data were plotted on the left side of the bean (blue) and permutation results plotted on the right (grey). Horizontal bars on beans indicate mean accuracies. The for each trait and the grey dashed lines indicate the overall average accuracies mean accuracy. Stars indicate significantly improved cross-validation accuracies with ($FDR < 0.05$) higher cross-validation accuracies.

Posterior phenotypic variance explained and model comparisons

To learn why the prediction performance varied among traits *per se* and heterosis transformations, we obtained the posterior variance explained by our models using the complete set of data. As shown in Figure 4, additive models explained more phenotypic variance for traits *per se* of DTP, DTS, EHT and PHT; but explained less phenotypic variance for heterosis transformations (BPH) of ASI, GY and TW. On the contrary, larger proportions in contrast, a larger proportion of the phenotypic variance could be explained by the dominant models for heterosis transformations (BPH) of ASI, GY and TW. For the GY in particular, 50% This difference was particularly striking for grain yield under the dominant model, where only 3% of the heterosis variance in trait *per se* could be explained by dominant model but 61% of the variance in BPH was explained.

Heterosis transformations were largely determined by the accuracies of the parental phenotypes. To take the uncertainty of the parental phenotypes into consideration, we estimated the combining abilities combining ability from the hybrid population itself to investigate which modes of inheritance perform better than the null models. We extracted the breeding values estimated with under both additive and dominant models using the genome-wide IBD blocks incorporated with the our haplotype blocks and incorporating GERP scores. Consistent with above analysis, IBD blocks coded with dominant mode of inheritance significantly (equation 1 vs. equation 2, ANOVA P value < 0.05) improved model fitting for ASI and GY. We also compared model 3 and model 4, ANOVA results indicated that 4 performed almost as good as 3, indicating that specific combining ability captured most of the parental interactions and the current method could not detect higher order of interactions. We then applied the following models:

$$Y_{ij} = \mu + GCA_i + GCA_j + \varepsilon \quad (1)$$

$$Y_{ij} = \mu + GCA_i + GCA_j + G_{ij} + \varepsilon \quad (2)$$

$$Y_{ij} = \mu + GCA_i + GCA_j + SCA_{ij} + \varepsilon \quad (3)$$

$$Y_{ij} = \mu + GCA_i + GCA_j + SCA_{ij} + G_{ij} + \varepsilon \quad (4)$$

where Y_{ij} is the BLUE value of the hybrid crossed between the i^{th} inbred and j^{th} inbred; μ , the overall mean; GCA_i , the general combining ability of the i^{th} inbred; GCA_j , the general combining ability of the j^{th} inbred; SCA_{ij} , the specific combining ability of between the i^{th} and j^{th} inbreds; G_{ij} , breeding values estimated by our GS model for hybrid crossed between the i^{th} inbred and j^{th} inbred; ε , the model residuals.

Consistent with the previous analysis, haplotype blocks coded with the dominant mode of inheritance significantly improved model fitting for ASI and GY (equation 1 vs. equation 2, ANOVA P value < 0.05 , Table S5). Comparison of model 3 and model 4 indicated that model 4 performed almost as well as 3 (ANOVA P value = X), indicating that specific combining ability captured most of the parental interactions and the current method could not detect higher order of interactions. does model 2 outperform model 1? does 3 outperform 2? 4 should outperform 3 because it has extra parameters, but the text says "almost as well"? We are comparing [1 and 2], [3 and 4] only by adding breeding value estimated from our GERP GS model. I do not think this result add anything to our story. maybe we will ditch it?




Figure 4 Barplots of the Posterior phenotypic variance explained by deleterious genic SNPs in IBD blocks incorporated with GERP scores using additive and dominant models. Dark color indicates trait *per se* and grey color indicates BPH.

Discussion

- do we support deleterious model of Mezouk et al.? **yes**
- how do results match with heritability and heterosis? **and mode of inheritance**
- Indication for breeding? genomic selection using GERP and other annotation information.

In this study, we have identified more than 500,000 deleterious SNPs were identified in in panel of elite maize lines. On average, each elite inbred line carries about 100,000 deleterious SNPs (GERP > 0). In the population cool, this should be mentioned here but also in the results. is there any big variation? some lines more than others? is mo17 worse than the pops? given the comment below i added about linkage, can we calculate mean deleterious variants per cM? that would be sweet to know and to point out that there are tons around the centromere Across lines, however, the majority of these deleterious mutations were maintained in a at low frequency, which consistent with previous observation observations (Rodgers-Melnick et al. 2015). The large number of deleterious alleles linked deleterious alleles present would likely make it hard to be completely purged completely purge all such alleles through breeding efforts. In practice, part of the genetic loads could be released in F1 hybrids by combining appropriate inbred lines (from pre-defined heterotic groups) to complement deleterious alleles. Instead, breeders have devised a strategy — hybrid breeding — that may circumvent much of this problem via complementation. Indeed, results in this study shown show that prediction accuracies were higher for yield heterosis using deleterious SNPs than permuted ones. Because we weighted SNPs with their conservation score in GS model, the improved accuracies could be attributed to deleteriousness of SNP variants. Therefore, it provided evidence of complementation of deleterious alleles for heterosis (CITE) — Note that thousands of deleterious alleles may be involved in the when information about deleterious SNPs was incorporated into the prediction model. Because there are likely thousands of deleterious alleles involved in complementation, and most of which may have minor effects. Their effects could not be easily detected by traditional QTL or GWAS (CITE) — many with relatively small effects, traditional GWAS approaches with genome-wide thresholds of multiple testing are likely unable to detect such effects. Using a liberal significance threshold, however, our previous efforts nonetheless showed an enrichment for deleterious genic SNPs among associated markers for a number

of traits (Mezmouk and Ross-Ibarra 2014).

~~To test~~ Our models did not increase the prediction accuracies equally well for traits *per se* and their heterosis transformations. This is not surprising, however, given that the variation in genetic architectures of different phenotypic traits — flowering time, for example, appears primarily determined by many loci of small additive effect (Buckler *et al.* 2009), while ~~X~~ *better to use another maize example if we can rather than rice...*. Consistent with this, we observed that the additive model increased prediction accuracies for ASI as a trait *per se* and *ideally above use a phenotype we can compare in this way...* dominant model increased the prediction accuracies for trait *per se* of GY. One limitation of our current models is that we assume phenotypic traits are determined by complete additive or complete dominant effects. Traits with a mixture of additive and dominant casual loci may thus fail to be predicted. Another limitaiton is likely the population size used; we may simply not have enough power to predict traits with low heritability.

To test the simple complementation hypothesis, we developed a GS pipeline, which incorporated evolutionary conservation information in the GS model. As the genotyping cost keeps declining, GS ~~tends to replace marker-assistant selection (MAS) (CITE)~~ gains its popularity in plant breeding (CITE). ~~Researchers developed a series of~~ (Desta and Ortiz 2014). Previous studies mainly focused on developing statistical approaches to improve the efficiency of GS model (CITE), however, none of the existing ~~approaches studies~~ have taken prior biological information into consideration. It is known that SNP variations would have varied impacts depending on their genomic position (e.g.) (CITE), biological function (e.g. transcription binding sites) and evolutionary conservation. The incorporation of GERP information under the Bayesian framework developed in this study is the first step towards more meaningful models for further GS.

~~It was not surprised that the current models did not increase the prediction accuacies equally well for traits *per se* and their heterosis transformations. Genetic arachitectures of different phenotypic traits varied in maize and other crop species; e.g., flowering time of maize was determined by many small effect additive loci (CITE) and rice yield was determined by many dominant loci (CITE). We observed additive model increased the prediction accuacies for trait *per se* of ASI and dominant model increased the prediction accuacies for trait *per se* of GY. Because under current models, we simply assume phenotypic traits were determined by complete additive or complete dominant effects. Traits with mixture effects of additive and dominant loci may fail to be predicted. In addition, the population size was relative small in the model training, we may not have enough power to predict traits with low heritability.~~

Acknowledgements

Financial support for this work came from NSF (grants IOS-0820619 and IOS-1238014), USDA (grants 2009-65300-05668 and X), DuPont Pioneer, and N2 Genetics. We'd like to thank Graham Coop, James Holland, X, and X reviewers for helpful discussion.

Literature Cited

- Birchler, J. A., D. L. Auger, and N. C. Riddle, 2003 In search of the molecular basis of heterosis. *The Plant Cell* **15**: 2236–2239.
- Browning, B. L. and S. R. Browning, 2009 A unified approach to genotype imputation and haplotype-phase inference for large data sets of trios and unrelated individuals. *Am J Hum Genet* **84**: 210–23.
- Buckler, E. S., J. B. Holland, P. J. Bradbury, C. B. Acharya, P. J. Brown, C. Browne, E. Ersoz, S. Flint-Garcia, A. Garcia, J. C. Glaubitz, *et al.*, 2009 The genetic architecture of maize flowering time. *Science* **325**: 714–718.
- Charlesworth, D. and J. H. Willis, 2009 The genetics of inbreeding depression. *Nature reviews. Genetics* **10**: 783–96.
- Chia, J.-M., C. Song, P. J. Bradbury, D. Costich, N. de Leon, J. Doebley, R. J. Elshire, B. Gaut, L. Geller, J. C. Glaubitz, M. Gore, K. E. Guill, J. Holland, M. B. Hufford, J. Lai, M. Li, X. Liu, Y. Lu, R. McCombie, R. Nelson, J. Poland, B. M. Prasanna, T. Pyhäjärvi, T. Rong, R. S. Sekhon, Q. Sun, M. I. Tenaillon, F. Tian, J. Wang, X. Xu, Z. Zhang, S. M. Kaeppler, J. Ross-Ibarra, M. D. McMullen, E. S. Buckler, G. Zhang, Y. Xu, and D. Ware, 2012 Maize hapmap2 identifies extant variation from a genome in flux. *Nat Genet* **44**: 803–7.
- Clark, R. M., S. Tavaré, and J. Doebley, 2005 Estimating a nucleotide substitution rate for maize from polymorphism at a major domestication locus. *Molecular Biology and Evolution* **22**: 2304–2312.
- Cooper, G. M., E. a. Stone, G. Asimenos, E. D. Green, S. Batzoglou, and A. Sidow, 2005 Distribution and intensity of constraint in mammalian genomic sequence. *Genome research* **15**: 901–13.
- Davydov, E. V., D. L. Goode, M. Sirota, G. M. Cooper, A. Sidow, and S. Batzoglou, 2010 Identifying a high fraction of the human genome to be under selective constraint using GERP++. *PLoS computational biology* **6**: e1001025.
- Desta, Z. A. and R. Ortiz, 2014 Genomic selection: genome-wide prediction in plant improvement. *Trends in plant science* **19**: 592–601.
- Doyle, J. J. and J. Doyle, 1987 Genomic plant dna preparation from fresh tissue-ctab method. *Phytochem Bull* **19**: 11–15.
- Duvick, D. N., 2001 Biotechnology in the 1930s: the development of hybrid maize. *Nature Reviews Genetics* **2**: 69–74.
- East, E. M., 1936 Heterosis. *Genetics* **21**: 375.
- Eyre-Walker, A. and P. D. Keightley, 2007 The distribution of fitness effects of new mutations. *Nature reviews. Genetics* **8**: 610–618.
- Falconer, D. and T. Mackay, 1996 *Introduction to Quantitative Genetics*. Longman.
- Frascaroli, E., M. A. Cane, P. Landi, G. Pea, L. Gianfranceschi, M. Villa, M. Morgante, and M. E. Pè, 2007 Classical genetic and quantitative trait loci analyses of heterosis in a maize hybrid between two elite inbred lines. *Genetics* **176**: 625–644.
- Gama, L. T., M. C. Bressan, E. C. Rodrigues, L. V. Rossato, O. C. Moreira, S. P. Alves, and R. J. B. Bessa, 2013 Heterosis for meat quality and fatty acid profiles in crosses among *Bos indicus* and *Bos taurus* finished on pasture or grain. *Meat Science* **93**: 98–104.
- Gilmour, A. R., B. Gogel, B. Cullis, and R. Thompson, 2009 *Asreml user guide release 3.0*. VSN International Ltd, Hemel Hempstead, UK.
- Gore, M. a., J.-M. Chia, R. J. Elshire, Q. Sun, E. S. Ersoz, B. L. Hurwitz, J. a. Peiffer, M. D. McMullen, G. S. Grills, J. Ross-Ibarra, D. H. Ware, and E. S. Buckler, 2009 A first-generation haplotype map of maize. *Science (New York, N.Y.)* **326**: 1115–7.
- Graham, G. I., D. W. Wolff, and C. W. Stuber, 1997 Characterization of a yield quantitative trait locus on chromosome five of maize by fine mapping. *Crop Science* **37**: 1601–1610.
- Habier, D., R. L. Fernando, K. Kizilkaya, and D. J. Garrick, 2011 Extension of the bayesian alphabet for genomic selection. *BMC bioinformatics* **12**: 186.
- Haddrill, P. R., D. L. Halligan, D. Tomaras, and B. Charlesworth, 2007 Reduced efficacy of selection in regions of the *Drosophila* genome that lack crossing over. *Genome biology* **8**: R18.
- Huang, X., S. Yang, J. Gong, Y. Zhao, Q. Feng, H. Gong, W. Li, Q. Zhan, B. Cheng, J. Xia, *et al.*, 2015 Genomic analysis of hybrid rice varieties reveals numerous superior alleles that contribute to heterosis. *Nature communications* **6**.
- Joseph, S. B. and D. W. Hall, 2004 Spontaneous mutations in diploid *Saccharomyces cerevisiae*: More beneficial than expected. *Genetics* **168**: 1817–1825.
- Krieger, U., Z. B. Lippman, and D. Zamir, 2010 The flowering gene single flower truss drives heterosis for yield in tomato. *Nature genetics* **42**: 459–463.
- Larièpe, a., B. Mangin, S. Jasson, V. Combes, F. Dumas, P. Jamin, C. Lariagon, D. Jolivot, D. Madur, J. Fiévet, A. Gallais, P. Dubreuil, A. Charcosset, and L. Moreau, 2012 The genetic basis of heterosis: multiparental quantitative trait loci mapping reveals contrasted levels of apparent overdominance among traits of agronomical interest in maize (*Zea mays* L.). *Genetics* **190**: 795–811.
- Li, H. and R. Durbin, 2009 Fast and accurate short read alignment with burrows-wheeler transform. *Bioinformatics* **25**: 1754–60.
- Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R. Durbin, and 1000 Genome Project Data Processing Subgroup, 2009 The sequence alignment/map format and samtools. *Bioinformatics* **25**: 2078–9.
- McMullen, M. D., S. Kresovich, H. S. Villeda, P. Bradbury, H. Li, Q. Sun, S. Flint-Garcia, J. Thornsberry, C. Acharya, C. Bottoms, P. Brown, C. Browne, M. Eller, K. Guill, C. Harjes, D. Kroon, N. Lepak, S. E. Mitchell, B. Peterson, G. Pressoir, S. Romero, M. Oropeza Rosas, S. Salvo, H. Yates, M. Hanson, E. Jones, S. Smith, J. C. Glaubitz, M. Goodman, D. Ware, J. B. Holland, and E. S. Buckler, 2009 Genetic properties of the maize nested association mapping population. *Science (New York, N.Y.)* **325**: 737–40.
- Mezmouk, S. and J. Ross-Ibarra, 2014 The pattern and distribution of deleterious mutations in maize. *G3 (Bethesda, Md.)* **4**: 163–71.
- Mikel, M. A. and J. W. Dudley, 2006 Evolution of north american dent corn from public to proprietary germplasm. *Crop science* **46**: 1193–1205.
- Minvielle, F., 1987 Dominance is not necessary for heterosis: a two-locus model. *Genetical research* **49**: 245–247.
- Rodgers-Melnick, E., P. J. Bradbury, R. J. Elshire, J. C. Glaubitz, C. B. Acharya, S. E. Mitchell, C. Li, Y. Li, and E. S. Buckler, 2015 Recombination in diverse maize is stable, predictable, and associated with genetic load. *Proceedings of the National Academy of Sciences* **112**: 3823–3828.
- Romay, M. C., M. J. Millard, J. C. Glaubitz, J. A. Peiffer, K. L. Swarts, T. M. Casstevens, R. J. Elshire, C. B. Acharya, S. E. Mitchell, S. A. Flint-Garcia, M. D. McMullen, J. B. Holland, E. S. Buckler, and C. A. Gardner, 2013 Comprehensive genotyping of the usa national maize inbred seed bank. *Genome Biol* **14**:

R55.

- Schnell, F. and C. Cockerham, 1992 Multiplicative vs. arbitrary gene action in heterosis. *Genetics* **131**: 461–469.
- Schwartz, D., 1973 Single gene heterosis for alcohol dehydrogenase in maize: the nature of the subunit interaction. *Theoretical and Applied Genetics* **43**: 117–120.
- Shapira, R., T. Levy, S. Shaked, E. Fridman, and L. David, 2014 Extensive heterosis in growth of yeast hybrids is explained by a combination of genetic models. *Heredity* **113**: 1–11.
- Shull, G. H., 1908 The composition of a field of maize. *Journal of Heredity* pp. 296–301.
- van Heerwaarden, J., M. B. Hufford, and J. Ross-Ibarra, 2012 Historical genomics of north american maize. *Proc Natl Acad Sci U S A* **109**: 12420–5.
- Xiao, J., J. Li, L. Yuan, and S. D. Tanksley, 1995 Dominance is the major genetic basis of heterosis in rice as revealed by qtl analysis using molecular markers. *Genetics* **140**: 745–754.

Supporting Information

Table S1 [BLUE values of the seven phenotypic traits. \(https://github.com/RILAB/pvpDiallel/blob/master/manuscript/Figure_Table/Table_S1.trait_matrix.csv\)](https://github.com/RILAB/pvpDiallel/blob/master/manuscript/Figure_Table/Table_S1.trait_matrix.csv)

Diallel experimental design and distribution of phenotypic data. (A) Twelve maize inbred lines were selected and crossed in a partial diallel. (B) Density plots of the phenotypic distributions.

Table S2 [General combining ability and specific combining ability of the seven phenotypic traits. \(https://github.com/RILAB/pvpDiallel/blob/master/manuscript/Figure_Table/Table_S2.CA.csv\)](https://github.com/RILAB/pvpDiallel/blob/master/manuscript/Figure_Table/Table_S2.CA.csv)

Table S3 [Cross-validation results using all genome-wide deleterious SNPs. \(https://github.com/RILAB/pvpDiallel/blob/master/manuscript/Figure_Table/Table_S3_allsnps_FDR.csv\)](https://github.com/RILAB/pvpDiallel/blob/master/manuscript/Figure_Table/Table_S3_allsnps_FDR.csv)

Table S4 [Cross-validation results using deleterious genic SNPs.](https://github.com/RILAB/pvpDiallel/blob/master/manuscript/Figure_Table/Table_S4_genicsnps_FDR.csv) (https://github.com/RILAB/pvpDiallel/blob/master/manuscript/Figure_Table/Table_S4_genicsnps_FDR.csv)

Table S5 [ANOVA P values of model comparisons.](https://github.com/RILAB/pvpDiallel/blob/master/manuscript/Figure_Table/Table_S5_model_comp.csv) (https://github.com/RILAB/pvpDiallel/blob/master/manuscript/Figure_Table/Table_S5_model_comp.csv)





Figure S1 [Twelve maize inbred lines were selected and crossed in a partial diallel fashion. Each inbred lines was used as both male and female and the resulting F1s were bulked.](#) *do we need to modify this diagram now that we now reciprocal crosses were pooled? any other idea to re-design the figure? or ditch it? this isn't bad, but what deleting the bottom triangle completely rather than leave the cells blank?*



	HB1	HB2	HB3	HB4	HB5	HB6	HB7	HB8	HB9	HB10	HB11	HB12	HB13	HB14
Hyb1	0	0	0	1	1	1	1	1	0	0	1	1	1	0
Hyb2	1	1	0	0	0	1	1	1	1	1	1	0	0	0
Hyb3	0	1	1	1	0	1	1	1	1	1	1	0	0	0
Hyb4	0	0	1	1	1	1	0	1	1	0	0	0	1	1

Figure S2 Haplotype block identification using an IBD approach. In the upper panel, regions in red are IBD blocks identified by pairwise comparison of the two parental lines of a hybrid. The vertical dashed lines define haplotype blocks. In the lower panel, hybrid genotypes at each block are coded as heterozygotes (0) or homozygotes (1).

Percentage of Better Parental Heterosis




Figure S3 Boxplot of percent best parent heterosis (pBPH). In the plot, ASI was calculated using pBPHmin and the other six traits were calculated using pBPHmax.




Figure S4 GERP score distribution across the genome. Shown are mean GERP scores in a 1-Mb bin region.




Figure S5 Incorporation of conservation information into IBD blocks. Regions of the genome that are identical by descent (IBD) among the 12 inbreds were identified using Beagle (Browning and Browning 2009). The GERP scores of SNPs in an IBD block were summed under both additive and dominant models. For a particular SNP with GERP score g , the homozygous non-reference genotype was assigned a value of $2g$, the heterozygote assigned a value of g , and the reference homozygote a value of 0. Under the dominant model, both the heterozygote and the non-reference homozygote were assigned a value of g , with the reference homozygote again assigned a value of 0.




Figure S6 Cross-validation accuracies using genic SNPs. Cross-validation experiments were conducted using genic SNPs and compared to circular-shuffled data for traits *per se* (**A, B**) and pBPH (**C, D**) under additive (**A, C**) and dominant (**B, D**) models. Distributions show accuracy of prediction from real data (blue) and permutations (grey), with horizontal bars to indicate mean accuracy. Stars indicate significantly higher cross-validation accuracy for the real data. The average accuracy across all traits is shown with the grey dotted line.