CYB003-101 Production Run
Tables

Date of Package: 02 September 2026

Date of Database Lock: 07 April 2026

Protocol Number: CYB003-101

A Phase 1, Open-label Study to Evaluate the Effects of CYB003 on the Pharmacokinetics of Caffeine, Bupropion, Losartan, Omeprazole, Metoprolol, and Midazolam in Healthy Adult Male Participants

TABLE OF CONTENTS

Disposition, Demographic, and Other Baseline Assessment Tables

Disposition

Table 14.1.1.1 Disposition of Participants Enrolled Population....5
Table 14.1.1.2 Important Protocol Deviations Safety Population....7
Table 14.1.1.3 Analysis Populations Enrolled Population....8

Demographic Data

Table 14.1.2.1 Demographics Safety Population....9

Medical History

Table 14.1.3.1 Medical History by MedDRA System Organ Class and Preferred Term Safety Population....12
Table 14.1.3.2 Psychedelic Use History Safety Population....14

Prior and Concomitant Medications

Table 14.1.4.1 Prior Medications Safety Population....15
Table 14.1.4.2 Concomitant Medications Safety Population....17

Extent of Exposure

Table 14.1.5.1 Summary of Exposure to Study Intervention Safety Population....18

Pharmacokinetic Data

PK Analysis Tables

Table 14.2.1.1.1 Summary of Plasma CYB003 and Metabolite Concentrations (ng/mL) PK Population....19
Table 14.2.1.1.2 Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003 PK Population....26
Table 14.2.1.2.1 Summary of Plasma PK Parameters of CYB003 and Metabolites PK Population....68
Table 14.2.1.2.2 Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003 PK Population....75

Table 14.2.1.3 Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003 DDI PK Population 117
Table 14.2.1.4 Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003 PK Population 129
Table 14.2.1.5 Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003 DDI PK Population 141
Table 14.2.1.6 Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003 PK Population 153
Table 14.2.1.7 Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003 DDI PK Population 165
Table 14.2.1.8 Summary of Percent Total Drug-Related Exposures for CYB003 Metabolites PK Population 177

Safety Data

Adverse Events

Table 14.3.1.1 Treatment-Emergent Adverse Event Overview Safety Population 178
Table 14.3.1.2 Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term Safety Population 180
Table 14.3.1.3 Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity Safety Population 186
Table 14.3.1.4 Treatment-Emergent Adverse Events with an Onset Date within 48 Hours from CYB003 Administration on Day 8 by MedDRA System Organ Class and Preferred Term Safety Population 214
Table 14.3.1.5 Related Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term Safety Population 219
Table 14.3.1.6 Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term Safety Population 225
Table 14.3.1.7 Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity Safety Population 230
Table 14.3.1.8 Related Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term Safety Population 254
Table 14.3.1.9 Potential Abuse-Related Adverse Events by MedDRA System Organ Class and Preferred Term Safety Population 259
Table 14.3.1.10 Potential Abuse-Related Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity Safety Population 261
Table 14.3.1.11 Related Potential Abuse-Related Adverse Events by MedDRA System Organ Class and Preferred Term Safety Population 270
Table 14.3.1.12 Non-SMRE Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term Safety Population 272
Table 14.3.1.13 Non-SMRE Related Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term Safety Population 274
Table 14.3.1.14 Treatment-Emergent Adverse Events by MedDRA Preferred Term Safety Population 276
Table 14.3.1.15 Treatment-Emergent Adverse Events by MedDRA High Level Term Safety Population 280

Serious Adverse Events and Discontinuations Due to Adverse Events

Table 14.3.2.1 Serious Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term Safety Population 283
Table 14.3.2.2 Serious Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity Safety Population 284
Table 14.3.2.3 Related Serious Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term Safety Population 285
Table 14.3.2.4 Treatment-Emergent Adverse Events Leading to Study Discontinuation by MedDRA System Organ Class and Preferred Term Safety Population 286
Table 14.3.2.5 Treatment-Emergent Adverse Events Resulting in Death by MedDRA System Organ Class and Preferred Term Safety Population 287

Clinical Laboratory Tests

Table 14.3.4.1 Chemistry Test Results by Visit Safety Population 288
Table 14.3.4.2 Hematology Test Results by Visit Safety Population 306
Table 14.3.4.3 Urinalysis Test Results by Visit Safety Population 323
Table 14.3.4.4 Thyroid-Stimulating Hormone Test Results by Visit Safety Population 328
Table 14.3.4.5 Chemistry Results: Shift Table from Baseline to Post-Baseline Safety Population 329
Table 14.3.4.6 Hematology Results: Shift Table from Baseline to Post-Baseline Safety Population 347
Table 14.3.4.7 Thyroid-Stimulating Hormone Results: Shift Table from Baseline to Post-Baseline Safety Population 364

Vitals Signs

Table 14.3.5.1 Vital Signs by Timepoint Safety Population 365
Table 14.3.5.2 Potentially Clinically Significant Vital Signs by Timepoint Safety Population 397

Electrocardiogram

Table 14.3.6.1 Electrocardiogram Parameter Results by Timepoint Safety Population 405
Table 14.3.6.2 Potentially Clinically Significant QTcF Findings by Timepoint Safety Population 423
Table 14.3.6.3 Electrocardiogram Investigator Assessment: Shift Table by Timepoint Safety Population 426

Columbia Suicide Severity Rating Scale (C-SSRS)

Table 14.3.7.1 Columbia Suicide Severity Rating Scale (C-SSRS) by Visit Safety Population 428

Table 14.1.1.1
Disposition of Participants
Enrolled Population

	Total N=82 n (%)
Enrolled	82
Screen Failure	45 (54.9)
Reason for Screen Failure	
Adverse Event	0
Death	0
Lost to Follow Up	0
Did not meet Inclusion/Exclusion criteria	34 (41.5)
Study terminated by Sponsor	0
Subject withdrew consent	4 (4.9)
Other [a]	7 (8.5)
Safety Population, N1 [b]	36
Completed Treatment CYP Probe alone on Day 1	36 (100)
Completed Treatment (CYB003 + CYP Probe) on Day 8	34 (94.4)

n is the number of participants with the measurement available.
The number of participants in Enrolled Population, N, is used as the denominator for percentages under Enrolled category.
a. See Listing 16.2.1.1 for other screen failure reasons and other study discontinuation reasons.
b. N1 is the number of participants who receive at least one dose of any study intervention and N1 is used as the denominator for percentages for the following categories.

Source: Listing 16.2.1.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010101.sas FINAL (2026-08-28,12:53)

Table 14.1.1.1
Disposition of Participants
Enrolled Population

	Total N=82 n (%)
Completed Study	34 (94.4)
Discontinuation from Study	2 (5.6)
Reason for Discontinuation	
Participant Withdrawal of Consent	1 (2.8)
Participant is not compliant with study procedures	0
Adverse Event	0
Protocol Violation	0
Lost to Follow-Up	0
Sponsor Request for Early Termination of Site/Study	0
Positive Pregnancy Test	0
Death	0
Lack of Efficacy	0
Investigator's Decision	0
Other [a]	1 (2.8)

n is the number of participants with the measurement available.
The number of participants in Enrolled Population, N, is used as the denominator for percentages under Enrolled category.
a. See Listing 16.2.1.1 for other screen failure reasons and other study discontinuation reasons.
b. N1 is the number of participants who receive at least one dose of any study intervention and N1 is used as the denominator for percentages for the following categories.

Source: Listing 16.2.1.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010101.sas FINAL (2026-08-28,12:53)

Table 14.1.1.2
Important Protocol Deviations
Safety Population

	CYP Probe N=36 n (%)	CYB003 + CYP Probe N=34 n (%)	Total N=36 n (%)
Participants with important protocol deviations	1 (2.8)	1 (2.9)	2 (5.6)
Assessment Missed	1 (2.8)	0	1 (2.8)
Other	0	1 (2.9)	1 (2.8)
Inclusion/Exclusion	0	0	0
Study Drug Administration	0	0	0
Visit Out of Window	0	0	0
Visit Missed	0	0	0
Assessment Out of Window	0	0	0
Concomitant Medication	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Participants with multiple important deviations are included once per category of deviation.
n is the number of participants with the measurement available.
The number of participants in Safety Population, N, is used as the denominator for percentages.

Source: Listing 16.2.2.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010102.sas FINAL (2026-08-28,12:53)

Table 14.1.1.3
Analysis Populations
Enrolled Population

	Total N=82 n (%)
Enrolled Population	82 (100)
Safety Population	36 (43.9)
Pharmacokinetic (PK) Population	35 (42.7)
DDI PK Population	33 (40.2)

Enrolled Population includes all participants who signed informed consent, regardless of whether they took any study intervention.
Safety Population includes all participants who receive at least one dose of any study intervention.
Pharmacokinetic (PK) Population includes all participants in the Safety Population who have at least one quantifiable post-dose concentration data.
DDI PK Population includes all participants who receive the complete CYP probe cocktail on Days 1 and 8 and CYB003 on Day 8, do not vomit within 4 hours of receiving study medication both on Days 1 and 8, and have sufficient number of samples to calculate the area under the curve parameters and the Cmax.
n is the number of participants with the measurement available.
The number of participants in Enrolled Population, N, is used as the denominator for percentages.

Source: Listing 16.2.3.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010103.sas FINAL (2026-08-28,12:53)

Table 14.1.2.1
Demographics
Safety Population

	Total N=36
Age (years), n	36
Mean (SD)	40.0 (12.43)
Median	39.5
Q1, Q3	29.0, 52.0
Min, Max	20, 64
Sex, n (%)	
Male	36 (100)
Race, n (%)	
American Indian or Alaska Native	0
Asian	3 (8.3)
Black or African American	19 (52.8)
Native Hawaiian or Other Pacific Islander	0
White	14 (38.9)
Other [a]	0
Ethnicity, n (%)	
Hispanic or Latino	9 (25.0)
Not Hispanic or Latino	27 (75.0)
Not reported	0
Unknown	0

n is the number of participants with the measurement available.
The number of participants in Safety Population, N, is used as the denominator for percentages.
a. See Listing 16.2.4.1 for a complete list of all other races.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.4.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010201.sas FINAL (2026-08-28,12:53)

Table 14.1.2.1
Demographics
Safety Population

	Total N=36
Weight (kg), n	36
Mean (SD)	84.94 (13.239)
Median	85.05
Q1, Q3	73.25, 95.40
Min, Max	53.5, 109.2
Height (cm), n	36
Mean (SD)	176.57 (7.663)
Median	176.70
Q1, Q3	172.50, 182.70
Min, Max	156.5, 189.4
BMI (kg/m^2), n	36
Mean (SD)	27.153 (3.2367)
Median	27.590
Q1, Q3	24.625, 30.160
Min, Max	20.37, 31.91
Creatinine Clearance (mL/min), n	36
Mean (SD)	117.832 (33.3975)
Median	111.224
Q1, Q3	91.516, 135.211
Min, Max	79.40, 220.53

n is the number of participants with the measurement available.
The number of participants in Safety Population, N, is used as the denominator for percentages.
a. See Listing 16.2.4.1 for a complete list of all other races.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.4.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010201.sas FINAL (2026-08-28,12:53)

Table 14.1.2.1
Demographics
Safety Population

	Total N=36
Psychedelic Use History, n (%)	
Yes	3 (8.3)
No	33 (91.7)

n is the number of participants with the measurement available.
The number of participants in Safety Population, N, is used as the denominator for percentages.
a. See Listing 16.2.4.1 for a complete list of all other races.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.4.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010201.sas FINAL (2026-08-28,12:53)

Table 14.1.3.1
Medical History by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	Total N=36 n (%)
Participants with any medical history	8 (22.2)
Congenital, familial and genetic disorders	1 (2.8)
Hydrocele	1 (2.8)
Hepatobiliary disorders	1 (2.8)
Cholecystitis acute	1 (2.8)
Immune system disorders	2 (5.6)
Seasonal allergy	2 (5.6)
Injury, poisoning and procedural complications	3 (8.3)
Limb injury	1 (2.8)
Shoulder fracture	1 (2.8)
Tendon rupture	1 (2.8)
Investigations	1 (2.8)
Colonoscopy	1 (2.8)
Nervous system disorders	1 (2.8)
Tension headache	1 (2.8)
Skin and subcutaneous tissue disorders	1 (2.8)
Eczema	1 (2.8)

n is the number of participants with the measurement available.
The number of participants in Safety Population, N, is used as the denominator for percentages.
MedDRA version 27.1.

Source: Listing 16.2.4.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010301.sas FINAL (2026-08-28,12:53)

Table 14.1.3.1
Medical History by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	Total N=36 n (%)
Surgical and medical procedures	5 (13.9)
Cholecystectomy	1 (2.8)
Hydrocele operation	1 (2.8)
Internal fixation of fracture	1 (2.8)
Shoulder operation	1 (2.8)
Tenoplasty	1 (2.8)

n is the number of participants with the measurement available.
The number of participants in Safety Population, N, is used as the denominator for percentages.
MedDRA version 27.1.

Source: Listing 16.2.4.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010301.sas FINAL (2026-08-28,12:53)

Table 14.1.3.2
Psychedelic Use History
Safety Population

Substance Used	Total N=36 n (%)
Participants with a history of Psychedelic drug use	3 (8.3)
Psilocybin (mushrooms)	2 (5.6)
3,4-methylenedioxymethamphetamine (MDMA)/ecstasy	1 (2.8)
Lysergic acid diethylamide (LSD/acid)	1 (2.8)
Mescaline/peyote/San Pedro	1 (2.8)

n is the number of participants with the measurement available.
The number of participants in Safety Population, N, is used as the denominator for percentages.

Source: Listing 16.2.4.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010302.sas FINAL (2026-08-28,12:53)

Table 14.1.4.1
Prior Medications
Safety Population

Therapeutic Class (ATC level 2) [a]/ Pharmacological Class (ATC level 4) Preferred Name	Total N = 36 n (%)
Participants with any prior medication	5 (13.9)
ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS	1 (2.8)
PROPIONIC ACID DERIVATIVES	1 (2.8)
IBUPROFEN	1 (2.8)
CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS	1 (2.8)
CORTICOSTEROIDS, POTENT (GROUP III)	1 (2.8)
MOMETASONE	1 (2.8)
OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS	1 (2.8)
VARIOUS ALIMENTARY TRACT AND METABOLISM PRODUCTS	1 (2.8)
FIBRE, DIETARY	1 (2.8)
PSYCHOLEPTICS	1 (2.8)
MELATONIN RECEPTOR AGONISTS	1 (2.8)
MELATONIN	1 (2.8)
UNSPECIFIED HERBAL AND TRADITIONAL MEDICINE	1 (2.8)
UNCODED	1 (2.8)
MATRICARIA CHAMOMILLA	1 (2.8)

Prior medications are medications with start and end dates prior to the date of the dose of CYP Probe Substrates on Day 1.
a. Counts, n, reflect numbers of participants reporting one or more prior medications that map to the WHO Drug Dictionary (Version Sep2024). Participants are counted once within each level of summarization (ATC Level 2, ATC Level 4 or Preferred Name).
The number of participants in Safety Population, N, is used as the denominator for percentages.
ATC Level 2 = anatomical therapeutic chemical level 2 (therapeutic class); ATC Level 4 = anatomical therapeutic chemical level 4 (chemical class).

Source: Listing 16.2.4.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010401.sas FINAL (2026-08-28,12:53)

Table 14.1.4.1
Prior Medications
Safety Population

Therapeutic Class (ATC level 2) [a]/ Pharmacological Class (ATC level 4) Preferred Name	Total N = 36 n (%)
VITAMINS	1 (2.8)
ASCORBIC ACID (VITAMIN C), PLAIN	1 (2.8)
ASCORBIC ACID	1 (2.8)
VITAMIN D AND ANALOGUES	1 (2.8)
COLECALCIFEROL	1 (2.8)

Prior medications are medications with start and end dates prior to the date of the dose of CYP Probe Substrates on Day 1.
a. Counts, n, reflect numbers of participants reporting one or more prior medications that map to the WHO Drug Dictionary (Version Sep2024). Participants are counted once within each level of summarization (ATC Level 2, ATC Level 4 or Preferred Name).
The number of participants in Safety Population, N, is used as the denominator for percentages.
ATC Level 2 = anatomical therapeutic chemical level 2 (therapeutic class); ATC Level 4 = anatomical therapeutic chemical level 4 (chemical class).

Source: Listing 16.2.4.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010401.sas FINAL (2026-08-28,12:53)

Table 14.1.4.2
Concomitant Medications
Safety Population

Therapeutic Class (ATC level 2) [a]/ Pharmacological Class (ATC level 4) Preferred Name	CYP Probe N = 36 n (%)	CYB003 + CYP Probe N = 34 n (%)	Total N = 36 n (%)
Participants with any concomitant medication	1 (2.8)	3 (8.8)	3 (8.3)
ANTIBACTERIALS FOR SYSTEMIC USE	0	1 (2.9)	1 (2.8)
TETRACYCLINES	0	1 (2.9)	1 (2.8)
DOXYCYCLINE	0	1 (2.9)	1 (2.8)
CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS	1 (2.8)	1 (2.9)	1 (2.8)
CORTICOSTEROIDS, POTENT (GROUP III)	1 (2.8)	1 (2.9)	1 (2.8)
MOMETASONE	1 (2.8)	1 (2.9)	1 (2.8)
DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES	0	1 (2.9)	1 (2.8)
SELECTIVE BETA-2-ADRENORECEPTOR AGONISTS	0	1 (2.9)	1 (2.8)
SALBUTAMOL SULFATE	0	1 (2.9)	1 (2.8)

Concomitant medications are medications with an end date after the date of the dosing for each treatment or have a missing end date.
CYP Probe refers to Period 1 and Period 1 is defined as the time of administration of CYP Probe alone (Day 1) up to and including the time prior to the administration of CYB003 + CYP Probe (Day 8).
CYB003 + CYP Probe refers to Period 2 and Period 2 is defined as the time of administration of CYB003 + CYP Probe (Day 8) through the end of follow-up or last participation in the study.
a. Counts, n, reflect numbers of participants reporting one or more prior medications that map to the WHO Drug Dictionary (Version Sep2024). Participants are counted once within each level of summarization (ATC Level 2, ATC Level 4 or Preferred Name).
The number of participants in Safety Population, N, is used as the denominator for percentages.
ATC Level 2 = anatomical therapeutic chemical level 2 (therapeutic class); ATC Level 4 = anatomical therapeutic chemical level 4 (chemical class).

Source: Listing 16.2.4.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010402.sas FINAL (2026-08-28,12:53)

Table 14.1.5.1
Summary of Exposure to Study Intervention
Safety Population

	Total N=36 n (%)
Received CYP Probe in Period 1	36 (100)
Received CYP Probe + CYB003 in Period 2	34 (94.4)
Received interventions in both study periods	34 (94.4)
Duration of Study Exposure (days) [a]	
N	36
Mean (SD)	16.1 (4.44)
Median	15.5
Q1, Q3	15.0, 17.0
Min, Max	7, 33

n is the number of participants with the measurement available.
The number of participants in Safety Population, N, is used as the denominator for percentages.
a. Duration of Exposure = the end date of participant participation - the date of first dose of CYP Probe alone + 1.

Source: Listing 16.2.5.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14010501.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.1
Summary of Plasma CYB003 and Metabolite Concentrations (ng/mL)
PK Population

Timepoint Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
Pre-dose			
n	33	33	33
nBLQ (%)	33 (100.0)	33 (100.0)	33 (100.0)
Mean (SD)	0.000 (0.000)	0.000 (0.000)	0.000 (0.000)
CV%	NC	NC	NC
Geometric Mean (CV%)	NC	NC	NC
Median	0.000	0.000	0.000
Min, Max	0.000, 0.000	0.000, 0.000	0.000, 0.000
0.5 hours post-dose			
n	33	33	33
nBLQ (%)	9 (27.3)	27 (81.8)	11 (33.3)
Mean (SD)	6.53 (6.37)	3.32 (7.94)	4.03 (4.10)
CV%	97.5	239.5	101.7
Geometric Mean (CV%)	NC	NC	NC
Median	4.79	0.000	2.97
Min, Max	0.000, 21.4	0.000, 33.1	0.000, 15.7

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; NC = not calculated; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2.

Source: Listing 16.2.6.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010101.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.1
Summary of Plasma CYB003 and Metabolite Concentrations (ng/mL)
PK Population

Timepoint Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
1 hour post-dose			
n	33	33	33
nBLQ (%)	2 (6.1)	4 (12.1)	2 (6.1)
Mean (SD)	13.4 (7.39)	41.8 (34.4)	10.9 (6.21)
CV%	55.0	82.3	57.1
Geometric Mean (CV%)	NC	NC	NC
Median	13.6	31.9	10.3
Min, Max	0.000, 26.8	0.000, 139	0.000, 25.2
1.5 hours post-dose			
n	33	33	33
nBLQ (%)	0	1 (3.0)	0
Mean (SD)	18.9 (8.05)	80.7 (47.2)	11.8 (6.09)
CV%	42.5	58.5	51.7
Geometric Mean (CV%)	17.3 (47.0)	NC	10.5 (49.8)
Median	16.3	69.3	10.1
Min, Max	5.72, 37.8	0.000, 238	3.35, 34.0

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; NC = not calculated; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2.

Source: Listing 16.2.6.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010101.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.1
Summary of Plasma CYB003 and Metabolite Concentrations (ng/mL)
PK Population

Timepoint Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
2 hours post-dose			
n	33	33	33
nBLQ (%)	0	0	0
Mean (SD)	21.7 (7.62)	110 (53.8)	11.8 (4.48)
CV%	35.2	48.9	37.9
Geometric Mean (CV%)	20.3 (39.9)	98.6 (51.1)	11.1 (37.6)
Median	21.1	95.0	11.4
Min, Max	5.96, 43.0	29.1, 259	5.83, 24.3
3 hours post-dose			
n	32	32	32
nBLQ (%)	0	0	0
Mean (SD)	25.8 (6.74)	152 (52.1)	10.7 (3.42)
CV%	26.1	34.4	31.9
Geometric Mean (CV%)	25.0 (27.6)	143 (34.9)	10.2 (30.4)
Median	26.0	138	9.70
Min, Max	13.1, 40.2	75.1, 280	6.42, 21.6

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; NC = not calculated; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2.

Source: Listing 16.2.6.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010101.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.1
Summary of Plasma CYB003 and Metabolite Concentrations (ng/mL)
PK Population

Timepoint Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
4 hours post-dose			
n	30	30	30
nBLQ (%)	0	0	0
Mean (SD)	22.7 (6.18)	167 (47.0)	7.54 (3.29)
CV%	27.2	28.2	43.6
Geometric Mean (CV%)	21.9 (27.7)	160 (28.6)	7.01 (38.3)
Median	23.1	164	6.25
Min, Max	12.8, 39.3	95.6, 278	3.57, 18.5
6 hours post-dose			
n	32	32	32
nBLQ (%)	0	0	0
Mean (SD)	16.4 (4.94)	141 (40.7)	4.07 (1.75)
CV%	30.0	28.9	43.0
Geometric Mean (CV%)	15.7 (32.4)	135 (30.9)	3.73 (44.8)
Median	16.2	141	3.88
Min, Max	7.72, 26.8	72.4, 219	1.24, 9.20

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; NC = not calculated; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2.

Source: Listing 16.2.6.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010101.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.1
Summary of Plasma CYB003 and Metabolite Concentrations (ng/mL)
PK Population

Timepoint Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
8 hours post-dose			
n	33	33	33
nBLQ (%)	0	0	3 (9.1)
Mean (SD)	11.3 (3.36)	106 (29.7)	2.71 (0.979)
CV%	29.7	28.0	36.1
Geometric Mean (CV%)	10.8 (31.4)	102 (28.4)	NC
Median	11.8	105	2.53
Min, Max	5.50, 19.0	59.3, 194	1.29, 5.09
12 hours post-dose			
n	33	33	33
nBLQ (%)	0	0	14 (42.4)
Mean (SD)	5.55 (1.85)	60.3 (21.5)	1.77 (0.505)
CV%	33.4	35.7	28.5
Geometric Mean (CV%)	5.22 (38.8)	57.0 (34.8)	NC
Median	5.45	57.6	1.54
Min, Max	1.93, 8.43	27.0, 139	1.17, 2.87

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; NC = not calculated; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2.

Source: Listing 16.2.6.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010101.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.1
Summary of Plasma CYB003 and Metabolite Concentrations (ng/mL)
PK Population

Timepoint Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
16 hours post-dose			
n	32	32	32
nBLQ (%)	0	0	25 (78.1)
Mean (SD)	2.95 (1.07)	34.5 (14.0)	1.41 (0.279)
CV%	36.4	40.6	19.8
Geometric Mean (CV%)	2.76 (39.4)	32.1 (40.1)	NC
Median	2.83	31.8	1.26
Min, Max	1.34, 5.67	15.3, 79.6	1.17, 1.96
24 hours post-dose			
n	33	33	33
nBLQ (%)	13 (39.4)	13 (39.4)	33 (100.0)
Mean (SD)	1.60 (0.330)	16.2 (5.71)	NC
CV%	20.7	35.2	NC
Geometric Mean (CV%)	NC	NC	NC
Median	1.60	14.4	NC
Min, Max	1.11, 2.46	11.0, 34.8	NC, NC

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; NC = not calculated; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2.

Source: Listing 16.2.6.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010101.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.1
Summary of Plasma CYB003 and Metabolite Concentrations (ng/mL)
PK Population

Timepoint Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
48 hours post-dose			
n	33	33	33
nBLQ (%)	33 (100.0)	33 (100.0)	33 (100.0)
Mean (SD)	NC	NC	NC
CV%	NC	NC	NC
Geometric Mean (CV%)	NC	NC	NC
Median	NC	NC	NC
Min, Max	NC, NC	NC, NC	NC, NC

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; NC = not calculated; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2.

Source: Listing 16.2.6.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010101.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

Timepoint Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
Pre-dose				
n	32	31	22	22
nBLQ (%)	22 (68.8)	21 (67.7)	22 (100.0)	17 (77.3)
Mean (SD)	15.9 (30.3)	9.61 (16.4)	0.000 (0.000)	3.50 (7.15)
CV%	190.2	170.7	NC	204.6
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.000	0.000	0.000	0.000
Min, Max	0.000, 108	0.000, 54.4	0.000, 0.000	0.000, 26.3
0.5 hours post-dose				
n	32	31	22	22
nBLQ (%)	0	0	0	0
Mean (SD)	1330 (851)	1050 (643)	133 (84.2)	109 (67.6)
CV%	63.8	61.3	63.4	61.9
Geometric Mean (CV%)	1090 (74.0)	815 (97.2)	114 (57.8)	87.4 (88.0)
Median	987	964	111	102
Min, Max	345, 3240	117, 2750	44.8, 406	11.7, 297

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

Timepoint Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
1 hour post-dose				
n	32	31	22	22
nBLQ (%)	0	0	0	0
Mean (SD)	1770 (408)	1540 (453)	262 (97.8)	212 (85.5)
CV%	23.0	29.4	37.3	40.2
Geometric Mean (CV%)	1730 (23.6)	1470 (33.5)	244 (40.7)	194 (49.4)
Median	1770	1540	265	210
Min, Max	1110, 2540	678, 2410	108, 504	73.5, 373
1.5 hours post-dose				
n	32	31	22	22
nBLQ (%)	0	0	0	0
Mean (SD)	1690 (332)	1680 (330)	319 (107)	290 (94.7)
CV%	19.7	19.7	33.5	32.7
Geometric Mean (CV%)	1660 (19.5)	1640 (22.2)	302 (34.6)	274 (37.3)
Median	1620	1730	308	286
Min, Max	1210, 2360	945, 2240	163, 554	134, 481

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

Timepoint Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
2 hours post-dose				
n	31	31	21	22
nBLQ (%)	0	0	0	0
Mean (SD)	1620 (227)	1760 (450)	351 (106)	357 (109)
CV%	14.0	25.5	30.2	30.5
Geometric Mean (CV%)	1600 (14.1)	1720 (23.3)	335 (31.7)	340 (33.7)
Median	1610	1790	353	351
Min, Max	1130, 2120	966, 3640	209, 559	185, 546
3 hours post-dose				
n	32	30	22	21
nBLQ (%)	0	0	0	0
Mean (SD)	1490 (254)	1600 (251)	419 (110)	426 (110)
CV%	17.0	15.6	26.2	25.9
Geometric Mean (CV%)	1470 (18.6)	1580 (17.0)	404 (28.9)	412 (28.0)
Median	1530	1600	423	438
Min, Max	836, 1920	937, 2190	213, 597	229, 661

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

Timepoint Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
4 hours post-dose				
n	32	28	22	20
nBLQ (%)	0	0	0	0
Mean (SD)	1390 (265)	1470 (237)	465 (120)	509 (93.5)
CV%	19.0	16.2	25.9	18.4
Geometric Mean (CV%)	1370 (20.0)	1440 (18.6)	450 (27.1)	501 (19.8)
Median	1400	1490	439	504
Min, Max	836, 1940	754, 1800	271, 676	294, 709
6 hours post-dose				
n	32	30	22	22
nBLQ (%)	0	0	0	0
Mean (SD)	1130 (225)	1240 (302)	513 (106)	543 (115)
CV%	19.9	24.4	20.6	21.3
Geometric Mean (CV%)	1110 (21.5)	1200 (29.1)	501 (23.8)	531 (22.4)
Median	1140	1270	499	544
Min, Max	573, 1540	438, 2110	239, 675	330, 811

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

Timepoint Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
8 hours post-dose				
n	32	31	22	22
nBLQ (%)	0	0	0	0
Mean (SD)	920 (230)	1050 (414)	563 (111)	582 (192)
CV%	25.0	39.4	19.8	33.0
Geometric Mean (CV%)	889 (28.8)	979 (40.6)	554 (19.0)	559 (28.2)
Median	912	1020	536	546
Min, Max	348, 1340	284, 2680	386, 849	350, 1280
12 hours post-dose				
n	32	31	22	22
nBLQ (%)	0	0	0	0
Mean (SD)	547 (208)	652 (254)	469 (93.7)	477 (107)
CV%	38.0	39.0	20.0	22.4
Geometric Mean (CV%)	504 (45.3)	591 (53.5)	458 (22.6)	465 (25.0)
Median	530	687	483	493
Min, Max	135, 952	114, 1200	241, 614	243, 765

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

Timepoint Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
16 hours post-dose				
n	31	30	22	21
nBLQ (%)	0	0	0	0
Mean (SD)	336 (159)	405 (208)	367 (86.2)	377 (113)
CV%	47.4	51.4	23.5	29.9
Geometric Mean (CV%)	296 (58.6)	342 (74.4)	356 (28.6)	356 (38.6)
Median	311	361	373	367
Min, Max	56.0, 646	47.1, 876	139, 527	131, 572
24 hours post-dose				
n	32	31	22	22
nBLQ (%)	0	0	0	0
Mean (SD)	146 (99.7)	195 (134)	204 (83.9)	237 (101)
CV%	68.4	68.9	41.1	42.6
Geometric Mean (CV%)	113 (89.8)	143 (114.6)	186 (49.8)	207 (67.6)
Median	114	178	173	205
Min, Max	12.3, 373	12.7, 499	43.3, 365	35.1, 406

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

Timepoint Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
48 hours post-dose				
n	32	31	22	22
nBLQ (%)	19 (59.4)	18 (58.1)	9 (40.9)	5 (22.7)
Mean (SD)	30.0 (13.9)	41.8 (24.1)	38.1 (26.3)	43.8 (35.4)
CV%	46.5	57.6	69.0	80.9
Geometric Mean (CV%)	NC	NC	NC	NC
Median	23.9	35.1	39.0	26.5
Min, Max	14.3, 58.6	11.0, 82.9	10.4, 90.7	10.7, 135

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

Timepoint Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
Pre-dose				
n	31	29	31	29
nBLQ (%)	30 (96.8)	29 (100.0)	30 (96.8)	23 (79.3)
Mean (SD)	0.000 (0.000)	0.000 (0.000)	0.000 (0.000)	0.377 (0.940)
CV%	NC	NC	NC	249.7
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.000	0.000	0.000	0.000
Min, Max	0.000, 0.000	0.000, 0.000	0.000, 0.000	0.000, 4.43
0.5 hours post-dose				
n	31	29	31	29
nBLQ (%)	15 (48.4)	13 (44.8)	12 (38.7)	7 (24.1)
Mean (SD)	1.40 (2.39)	1.62 (2.47)	3.13 (4.61)	3.42 (3.69)
CV%	170.8	152.8	147.2	108.0
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.283	0.779	1.30	2.54
Min, Max	0.000, 10.3	0.000, 8.81	0.000, 21.6	0.000, 14.5

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

Timepoint Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
1 hour post-dose				
n	31	29	31	29
nBLQ (%)	0	1 (3.4)	0	1 (3.4)
Mean (SD)	15.3 (10.4)	10.7 (8.08)	28.4 (17.8)	20.1 (13.0)
CV%	67.6	75.7	62.9	64.7
Geometric Mean (CV%)	11.4 (109.9)	NC	22.8 (83.8)	NC
Median	13.7	9.08	21.3	18.6
Min, Max	0.960, 37.8	0.000, 27.2	3.47, 73.4	0.000, 53.2
1.5 hours post-dose				
n	31	27	31	27
nBLQ (%)	0	0	0	0
Mean (SD)	21.0 (10.1)	17.2 (9.98)	46.2 (20.9)	35.0 (21.6)
CV%	48.1	58.1	45.2	61.6
Geometric Mean (CV%)	19.0 (48.1)	13.3 (105.0)	42.3 (44.0)	27.6 (97.5)
Median	17.8	17.8	41.3	29.8
Min, Max	7.20, 54.3	1.19, 39.1	20.6, 95.0	2.11, 98.2

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

Timepoint Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
2 hours post-dose				
n	30	29	30	29
nBLQ (%)	0	0	0	0
Mean (SD)	21.2 (7.44)	18.8 (8.56)	59.9 (26.5)	46.0 (23.2)
CV%	35.1	45.4	44.3	50.5
Geometric Mean (CV%)	19.9 (38.8)	16.4 (64.8)	54.9 (43.6)	39.7 (66.9)
Median	18.6	19.1	53.8	46.6
Min, Max	8.08, 40.9	4.14, 36.8	25.3, 123	6.41, 115
3 hours post-dose				
n	31	28	31	28
nBLQ (%)	0	0	0	0
Mean (SD)	17.7 (5.67)	19.6 (5.17)	68.2 (30.4)	58.3 (28.6)
CV%	32.0	26.4	44.6	49.0
Geometric Mean (CV%)	16.9 (32.1)	18.9 (27.8)	62.8 (41.9)	53.1 (44.5)
Median	16.3	19.4	60.5	52.2
Min, Max	9.31, 30.0	10.5, 31.0	29.3, 148	23.4, 163

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

Timepoint Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
4 hours post-dose				
n	31	26	31	26
nBLQ (%)	0	0	0	0
Mean (SD)	13.8 (4.01)	17.2 (5.21)	70.2 (33.0)	70.9 (33.6)
CV%	29.0	30.4	47.0	47.4
Geometric Mean (CV%)	13.3 (29.6)	16.4 (32.0)	63.9 (45.1)	65.5 (39.7)
Median	12.7	15.9	60.4	60.3
Min, Max	7.52, 23.9	8.39, 27.6	32.4, 157	38.8, 196
6 hours post-dose				
n	31	28	31	28
nBLQ (%)	0	0	0	0
Mean (SD)	8.14 (2.09)	10.5 (4.14)	66.4 (30.1)	68.7 (31.8)
CV%	25.7	39.5	45.3	46.3
Geometric Mean (CV%)	7.88 (27.3)	9.75 (40.7)	60.6 (45.0)	63.5 (40.0)
Median	7.92	9.88	60.1	61.4
Min, Max	4.00, 13.3	4.67, 22.1	30.9, 138	36.3, 190

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

Timepoint Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
8 hours post-dose				
n	30	29	30	29
nBLQ (%)	0	0	0	0
Mean (SD)	5.46 (1.76)	6.64 (2.60)	62.9 (27.4)	66.8 (30.5)
CV%	32.3	39.1	43.6	45.6
Geometric Mean (CV%)	5.17 (35.6)	6.19 (40.1)	57.8 (43.4)	61.5 (41.7)
Median	5.80	5.87	52.2	60.2
Min, Max	2.66, 9.11	2.50, 13.7	27.5, 133	30.5, 176
12 hours post-dose				
n	31	29	31	29
nBLQ (%)	0	0	0	0
Mean (SD)	2.61 (0.732)	3.43 (1.19)	51.2 (23.7)	54.6 (24.6)
CV%	28.0	34.7	46.3	45.0
Geometric Mean (CV%)	2.51 (29.3)	3.22 (38.4)	46.7 (44.3)	50.2 (42.3)
Median	2.51	3.25	43.1	51.8
Min, Max	1.41, 4.29	1.26, 6.40	23.7, 129	26.5, 140

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

Timepoint Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
16 hours post-dose				
n	31	27	31	27
nBLQ (%)	0	0	0	0
Mean (SD)	1.77 (0.497)	2.16 (0.683)	45.1 (21.2)	44.6 (17.1)
CV%	28.1	31.7	47.0	38.4
Geometric Mean (CV%)	1.69 (30.8)	2.05 (33.6)	41.0 (46.3)	41.7 (38.0)
Median	1.73	1.96	39.5	42.6
Min, Max	0.983, 2.57	0.988, 3.66	19.2, 114	20.3, 98.9
24 hours post-dose				
n	31	29	31	29
nBLQ (%)	0	0	0	0
Mean (SD)	1.04 (0.327)	1.36 (0.447)	40.6 (20.2)	40.9 (19.8)
CV%	31.4	33.0	49.9	48.5
Geometric Mean (CV%)	0.992 (32.6)	1.28 (34.7)	36.9 (44.6)	37.5 (42.6)
Median	0.974	1.28	36.3	38.8
Min, Max	0.506, 1.81	0.689, 2.10	17.2, 119	19.5, 123

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

Timepoint Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
48 hours post-dose				
n	31	28	31	28
nBLQ (%)	23 (74.2)	14 (50.0)	0	0
Mean (SD)	0.619 (0.156)	0.669 (0.145)	19.6 (12.1)	20.7 (11.9)
CV%	25.2	21.7	61.8	57.6
Geometric Mean (CV%)	NC	NC	17.2 (52.5)	18.6 (46.4)
Median	0.570	0.684	16.5	17.8
Min, Max	0.522, 0.993	0.510, 1.06	5.34, 73.5	9.93, 71.0

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

Timepoint Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
Pre-dose				
n	34	33	32	30
nBLQ (%)	33 (97.1)	33 (100.0)	32 (100.0)	30 (100.0)
Mean (SD)	0.042 (0.244)	0.000 (0.000)	0.000 (0.000)	0.000 (0.000)
CV%	583.1	NC	NC	NC
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.000	0.000	0.000	0.000
Min, Max	0.000, 1.42	0.000, 0.000	0.000, 0.000	0.000, 0.000
0.5 hours post-dose				
n	34	33	32	30
nBLQ (%)	3 (8.8)	6 (18.2)	30 (93.8)	27 (90.0)
Mean (SD)	25.9 (28.3)	15.8 (16.4)	0.096 (0.393)	0.142 (0.437)
CV%	109.3	103.5	409.3	307.7
Geometric Mean (CV%)	NC	NC	NC	NC
Median	14.5	10.8	0.000	0.000
Min, Max	0.000, 123	0.000, 75.7	0.000, 1.96	0.000, 1.60

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

Timepoint Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
1 hour post-dose				
n	34	33	32	30
nBLQ (%)	0	3 (9.1)	6 (18.8)	10 (33.3)
Mean (SD)	27.3 (12.8)	12.1 (8.03)	5.14 (5.19)	3.52 (4.15)
CV%	46.8	66.6	100.8	117.8
Geometric Mean (CV%)	23.1 (84.1)	NC	NC	NC
Median	26.6	10.2	3.26	1.79
Min, Max	1.11, 57.6	0.000, 36.7	0.000, 17.6	0.000, 12.9
1.5 hours post-dose				
n	34	33	32	30
nBLQ (%)	0	2 (6.1)	3 (9.4)	4 (13.3)
Mean (SD)	21.9 (9.56)	14.3 (9.49)	18.7 (15.4)	12.5 (12.8)
CV%	43.7	66.4	82.4	102.8
Geometric Mean (CV%)	19.7 (54.3)	NC	NC	NC
Median	20.1	13.1	14.8	6.39
Min, Max	3.15, 45.1	0.000, 42.2	0.000, 56.3	0.000, 43.9

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

Timepoint Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
2 hours post-dose				
n	33	33	31	30
nBLQ (%)	0	1 (3.0)	0	4 (13.3)
Mean (SD)	20.8 (8.77)	19.5 (11.1)	38.7 (27.2)	25.2 (23.2)
CV%	42.2	56.8	70.2	91.9
Geometric Mean (CV%)	19.1 (43.3)	NC	25.9 (162.4)	NC
Median	19.6	18.0	32.5	15.6
Min, Max	7.85, 47.6	0.000, 52.1	1.25, 108	0.000, 72.7
3 hours post-dose				
n	33	31	31	28
nBLQ (%)	0	0	0	0
Mean (SD)	16.1 (6.04)	23.9 (15.9)	75.2 (39.0)	57.5 (43.9)
CV%	37.5	66.4	51.9	76.4
Geometric Mean (CV%)	15.0 (42.2)	20.4 (59.0)	60.1 (102.0)	39.5 (150.3)
Median	15.5	19.5	65.7	48.1
Min, Max	6.33, 28.8	9.18, 84.9	3.45, 160	1.23, 204

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

Timepoint Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
4 hours post-dose				
n	34	30	32	27
nBLQ (%)	0	0	0	0
Mean (SD)	12.1 (5.20)	20.3 (9.70)	90.2 (41.9)	90.6 (61.3)
CV%	42.9	47.9	46.5	67.6
Geometric Mean (CV%)	11.1 (43.5)	17.9 (58.3)	75.9 (81.5)	64.6 (132.1)
Median	10.9	16.5	88.1	89.3
Min, Max	5.09, 23.7	3.44, 41.3	8.36, 193	4.52, 256
6 hours post-dose				
n	34	32	32	29
nBLQ (%)	0	0	0	0
Mean (SD)	11.5 (5.45)	12.0 (6.83)	100 (43.4)	105 (50.6)
CV%	47.4	56.9	43.2	48.2
Geometric Mean (CV%)	10.4 (48.5)	10.5 (55.9)	90.0 (55.3)	90.7 (68.9)
Median	9.98	10.6	98.2	100
Min, Max	4.04, 31.0	3.58, 36.2	19.3, 217	16.7, 235

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

Timepoint Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
8 hours post-dose				
n	34	32	32	29
nBLQ (%)	0	0	0	0
Mean (SD)	6.61 (3.71)	7.28 (4.25)	76.2 (24.4)	89.3 (41.2)
CV%	56.2	58.3	32.0	46.2
Geometric Mean (CV%)	5.89 (49.7)	6.15 (66.8)	72.2 (35.8)	81.5 (46.3)
Median	5.41	6.19	76.9	83.9
Min, Max	2.29, 21.3	1.80, 17.0	33.1, 142	30.2, 240
12 hours post-dose				
n	31	33	29	30
nBLQ (%)	2 (6.5)	1 (3.0)	0	0
Mean (SD)	3.39 (2.84)	3.24 (2.16)	41.8 (15.0)	51.8 (22.9)
CV%	83.6	66.6	35.9	44.1
Geometric Mean (CV%)	NC	NC	39.2 (38.3)	47.7 (42.3)
Median	2.79	2.51	38.5	48.0
Min, Max	1.04, 15.7	1.04, 11.1	17.4, 71.7	23.8, 120

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

Timepoint Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
16 hours post-dose				
n	34	32	32	29
nBLQ (%)	20 (58.8)	17 (53.1)	0	0
Mean (SD)	2.38 (1.61)	2.37 (1.81)	21.2 (8.33)	25.5 (14.1)
CV%	67.5	76.4	39.4	55.3
Geometric Mean (CV%)	NC	NC	19.4 (45.7)	22.4 (54.8)
Median	1.89	2.14	19.8	20.3
Min, Max	1.18, 7.12	1.01, 8.39	6.95, 39.6	6.81, 58.5
24 hours post-dose				
n	34	32	32	29
nBLQ (%)	30 (88.2)	28 (87.5)	0	0
Mean (SD)	2.26 (1.64)	1.87 (1.63)	8.66 (4.01)	10.4 (5.51)
CV%	72.6	87.3	46.3	53.2
Geometric Mean (CV%)	NC	NC	7.79 (50.4)	9.11 (57.5)
Median	1.50	1.06	7.73	9.68
Min, Max	1.32, 4.70	1.04, 4.32	3.14, 18.4	1.69, 27.0

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

Timepoint Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
48 hours post-dose				
n	34	32	32	30
nBLQ (%)	33 (97.1)	31 (96.9)	14 (43.8)	9 (30.0)
Mean (SD)	1.53 (NC)	1.13 (NC)	1.50 (0.268)	1.60 (0.469)
CV%	NC	NC	17.9	29.4
Geometric Mean (CV%)	NC	NC	NC	NC
Median	1.53	1.13	1.51	1.62
Min, Max	1.53, 1.53	1.13, 1.13	1.00, 1.93	1.04, 2.76

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

Timepoint Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
Pre-dose				
n	35	33	24	28
nBLQ (%)	35 (100.0)	33 (100.0)	23 (95.8)	28 (100.0)
Mean (SD)	0.000 (0.000)	0.000 (0.000)	0.000 (0.000)	0.000 (0.000)
CV%	NC	NC	NC	NC
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.000	0.000	0.000	0.000
Min, Max	0.000, 0.000	0.000, 0.000	0.000, 0.000	0.000, 0.000
0.5 hours post-dose				
n	35	33	25	28
nBLQ (%)	35 (100.0)	33 (100.0)	24 (96.0)	28 (100.0)
Mean (SD)	0.000 (0.000)	0.000 (0.000)	0.000 (0.000)	0.000 (0.000)
CV%	NC	NC	NC	NC
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.000	0.000	0.000	0.000
Min, Max	0.000, 0.000	0.000, 0.000	0.000, 0.000	0.000, 0.000

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

Timepoint Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
1 hour post-dose				
n	35	33	25	28
nBLQ (%)	17 (48.6)	27 (81.8)	11 (44.0)	23 (82.1)
Mean (SD)	30.7 (62.4)	6.06 (24.4)	16.8 (25.2)	4.69 (19.4)
CV%	203.6	402.4	150.6	413.2
Geometric Mean (CV%)	NC	NC	NC	NC
Median	2.21	0.000	5.62	0.000
Min, Max	0.000, 312	0.000, 139	0.000, 89.2	0.000, 102
1.5 hours post-dose				
n	35	33	25	28
nBLQ (%)	9 (25.7)	24 (72.7)	5 (20.0)	19 (67.9)
Mean (SD)	128 (152)	8.21 (19.1)	81.7 (76.7)	7.40 (19.7)
CV%	119.0	232.5	93.9	266.2
Geometric Mean (CV%)	NC	NC	NC	NC
Median	68.4	0.000	56.5	0.000
Min, Max	0.000, 525	0.000, 87.2	0.000, 243	0.000, 75.4

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

Timepoint Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
2 hours post-dose				
n	34	33	24	28
nBLQ (%)	5 (14.7)	17 (51.5)	3 (12.5)	15 (53.6)
Mean (SD)	160 (157)	64.5 (147)	96.2 (60.3)	39.0 (73.0)
CV%	98.4	227.5	62.6	187.0
Geometric Mean (CV%)	NC	NC	NC	NC
Median	139	0.000	104	0.000
Min, Max	0.000, 755	0.000, 616	0.000, 192	0.000, 273
3 hours post-dose				
n	35	32	25	27
nBLQ (%)	0	6 (18.8)	1 (4.0)	6 (22.2)
Mean (SD)	171 (149)	89.5 (153)	84.9 (52.7)	50.4 (72.6)
CV%	87.1	170.5	62.1	144.0
Geometric Mean (CV%)	101 (200.4)	NC	NC	NC
Median	107	10.4	81.6	14.5
Min, Max	2.52, 619	0.000, 613	0.000, 232	0.000, 323

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

Timepoint Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
4 hours post-dose				
n	35	30	25	25
nBLQ (%)	0	6 (20.0)	1 (4.0)	5 (20.0)
Mean (SD)	92.3 (87.1)	94.5 (127)	59.5 (34.2)	53.5 (46.8)
CV%	94.4	134.3	57.4	87.6
Geometric Mean (CV%)	57.8 (154.2)	NC	NC	NC
Median	55.0	41.3	53.2	53.0
Min, Max	3.14, 436	0.000, 559	0.000, 145	0.000, 168
6 hours post-dose				
n	35	31	25	27
nBLQ (%)	3 (8.6)	1 (3.2)	0	0
Mean (SD)	49.3 (94.2)	72.2 (77.5)	29.0 (26.6)	47.4 (34.9)
CV%	191.2	107.3	91.8	73.7
Geometric Mean (CV%)	NC	NC	21.4 (89.7)	31.9 (153.8)
Median	19.5	44.5	18.3	35.3
Min, Max	2.50, 448	0.000, 253	4.23, 116	1.62, 133

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

Timepoint Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
8 hours post-dose				
n	35	32	25	28
nBLQ (%)	12 (34.3)	3 (9.4)	0	0
Mean (SD)	27.6 (60.6)	54.3 (89.8)	11.2 (12.6)	22.9 (11.1)
CV%	219.5	165.4	112.6	48.3
Geometric Mean (CV%)	NC	NC	7.45 (108.1)	20.0 (61.3)
Median	10.3	29.7	6.72	22.7
Min, Max	2.07, 253	2.68, 395	1.24, 57.9	6.19, 48.3
12 hours post-dose				
n	35	33	25	28
nBLQ (%)	29 (82.9)	12 (36.4)	12 (48.0)	0
Mean (SD)	47.9 (85.4)	16.0 (20.8)	3.23 (3.87)	6.30 (4.81)
CV%	178.2	129.8	119.6	76.5
Geometric Mean (CV%)	NC	NC	NC	4.65 (98.5)
Median	3.68	6.70	1.77	5.04
Min, Max	2.03, 216	2.10, 73.8	1.04, 15.3	1.05, 17.1

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

Timepoint Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
16 hours post-dose				
n	35	32	25	27
nBLQ (%)	33 (94.3)	27 (84.4)	24 (96.0)	15 (55.6)
Mean (SD)	21.8 (14.4)	21.6 (31.0)	4.16 (NC)	2.49 (2.07)
CV%	66.2	143.8	NC	83.2
Geometric Mean (CV%)	NC	NC	NC	NC
Median	21.8	4.87	4.16	1.48
Min, Max	11.6, 32.0	2.91, 75.5	4.16, 4.16	1.02, 7.05
24 hours post-dose				
n	35	33	25	28
nBLQ (%)	34 (97.1)	31 (93.9)	25 (100.0)	26 (92.9)
Mean (SD)	6.54 (NC)	11.9 (11.6)	NC	1.63 (0.636)
CV%	NC	96.8	NC	39.0
Geometric Mean (CV%)	NC	NC	NC	NC
Median	6.54	11.9	NC	1.63
Min, Max	6.54, 6.54	3.76, 20.1	NC, NC	1.18, 2.08

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

Timepoint Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
48 hours post-dose				
n	35	33	25	28
nBLQ (%)	35 (100.0)	33 (100.0)	25 (100.0)	28 (100.0)
Mean (SD)	NC	NC	NC	NC
CV%	NC	NC	NC	NC
Geometric Mean (CV%)	NC	NC	NC	NC
Median	NC	NC	NC	NC
Min, Max	NC, NC	NC, NC	NC, NC	NC, NC

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

Timepoint Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
Pre-dose				
n	34	33	34	33
nBLQ (%)	34 (100.0)	33 (100.0)	34 (100.0)	33 (100.0)
Mean (SD)	0.000 (0.000)	0.000 (0.000)	0.000 (0.000)	0.000 (0.000)
CV%	NC	NC	NC	NC
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.000	0.000	0.000	0.000
Min, Max	0.000, 0.000	0.000, 0.000	0.000, 0.000	0.000, 0.000
0.5 hours post-dose				
n	34	33	34	33
nBLQ (%)	9 (26.5)	6 (18.2)	5 (14.7)	5 (15.2)
Mean (SD)	3.51 (6.66)	4.15 (6.17)	3.39 (2.99)	4.94 (5.90)
CV%	189.9	148.7	88.4	119.4
Geometric Mean (CV%)	NC	NC	NC	NC
Median	1.12	2.06	2.69	3.34
Min, Max	0.000, 35.4	0.000, 24.4	0.000, 13.7	0.000, 29.3

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

Timepoint Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
1 hour post-dose				
n	34	33	34	33
nBLQ (%)	0	1 (3.0)	0	1 (3.0)
Mean (SD)	19.1 (15.9)	12.3 (9.79)	11.2 (6.84)	8.25 (6.16)
CV%	83.4	79.6	61.0	74.7
Geometric Mean (CV%)	14.2 (94.2)	NC	9.05 (88.1)	NC
Median	14.2	9.73	10.7	7.10
Min, Max	2.52, 72.7	0.000, 37.8	0.937, 32.0	0.000, 25.2
1.5 hours post-dose				
n	34	33	34	33
nBLQ (%)	0	0	0	0
Mean (SD)	24.5 (14.5)	18.0 (11.5)	11.9 (5.60)	9.44 (6.01)
CV%	59.0	63.6	47.0	63.7
Geometric Mean (CV%)	20.8 (64.5)	14.1 (91.8)	10.6 (57.2)	7.39 (91.5)
Median	21.2	17.7	11.2	7.95
Min, Max	6.00, 68.7	2.03, 51.8	2.62, 26.6	1.13, 26.8

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

Timepoint Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
2 hours post-dose				
n	33	33	33	33
nBLQ (%)	0	0	0	0
Mean (SD)	24.1 (12.0)	19.7 (10.9)	11.2 (5.00)	9.85 (5.27)
CV%	49.6	55.1	44.8	53.5
Geometric Mean (CV%)	21.3 (55.9)	16.8 (65.9)	10.2 (47.7)	8.21 (76.5)
Median	22.2	19.1	9.85	9.26
Min, Max	6.92, 53.7	4.52, 50.7	3.25, 22.6	1.30, 19.8
3 hours post-dose				
n	34	32	34	32
nBLQ (%)	0	0	0	0
Mean (SD)	23.1 (13.0)	23.5 (12.1)	11.1 (4.10)	10.4 (4.21)
CV%	56.2	51.3	36.8	40.5
Geometric Mean (CV%)	19.9 (61.6)	20.4 (61.2)	10.4 (41.2)	9.45 (50.4)
Median	21.1	22.3	10.9	9.80
Min, Max	6.32, 67.2	5.79, 49.9	3.77, 19.8	2.60, 19.5

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

Timepoint Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
4 hours post-dose				
n	34	30	34	30
nBLQ (%)	0	0	0	0
Mean (SD)	21.2 (12.3)	23.1 (11.8)	10.5 (3.68)	9.85 (3.37)
CV%	58.2	50.9	35.1	34.3
Geometric Mean (CV%)	17.9 (67.3)	20.3 (57.5)	9.86 (37.1)	9.16 (44.0)
Median	18.3	20.7	10.2	10.1
Min, Max	5.44, 61.6	6.94, 51.1	4.11, 20.3	2.71, 17.0
6 hours post-dose				
n	34	32	34	32
nBLQ (%)	0	0	0	0
Mean (SD)	15.4 (9.17)	17.5 (11.2)	9.08 (2.35)	9.61 (3.08)
CV%	59.5	64.1	25.9	32.0
Geometric Mean (CV%)	12.8 (74.0)	13.9 (83.5)	8.76 (28.6)	9.04 (39.6)
Median	14.9	15.2	8.90	9.83
Min, Max	2.87, 44.3	2.91, 43.8	3.76, 14.8	2.76, 18.0

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

Timepoint Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
8 hours post-dose				
n	34	33	34	33
nBLQ (%)	0	0	0	0
Mean (SD)	11.5 (8.04)	13.5 (9.82)	8.05 (2.11)	8.42 (2.43)
CV%	69.7	72.8	26.2	28.9
Geometric Mean (CV%)	9.00 (88.3)	10.1 (100.2)	7.76 (28.9)	8.02 (34.5)
Median	10.0	11.3	7.95	8.32
Min, Max	1.56, 37.4	1.80, 38.8	3.85, 12.7	2.59, 13.9
12 hours post-dose				
n	34	32	34	32
nBLQ (%)	1 (2.9)	0	0	0
Mean (SD)	5.65 (4.24)	6.81 (5.46)	5.60 (1.50)	6.00 (1.61)
CV%	75.1	80.2	26.8	26.9
Geometric Mean (CV%)	NC	4.72 (120.5)	5.41 (28.1)	5.76 (30.8)
Median	5.06	6.25	5.50	6.19
Min, Max	0.605, 20.5	0.668, 21.8	3.01, 8.77	2.27, 8.99

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

Timepoint Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
16 hours post-dose				
n	34	32	34	32
nBLQ (%)	4 (11.8)	3 (9.4)	0	0
Mean (SD)	3.33 (2.63)	3.80 (3.30)	3.97 (1.13)	3.96 (1.26)
CV%	78.9	86.7	28.6	31.7
Geometric Mean (CV%)	NC	NC	3.81 (30.0)	3.77 (34.1)
Median	2.69	3.05	3.64	3.86
Min, Max	0.779, 11.2	0.575, 14.3	1.92, 6.35	1.68, 7.60
24 hours post-dose				
n	34	33	34	33
nBLQ (%)	14 (41.2)	11 (33.3)	0	0
Mean (SD)	1.74 (1.29)	2.01 (1.67)	2.04 (0.630)	2.10 (0.794)
CV%	74.3	82.9	30.9	37.8
Geometric Mean (CV%)	NC	NC	1.94 (34.8)	1.95 (43.3)
Median	1.39	1.33	1.84	2.04
Min, Max	0.592, 5.49	0.508, 7.23	0.703, 3.15	0.565, 4.48

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

Timepoint Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
48 hours post-dose				
n	34	33	34	33
nBLQ (%)	32 (94.1)	31 (93.9)	30 (88.2)	29 (87.9)
Mean (SD)	0.702 (0.229)	0.905 (0.132)	0.526 (0.032)	0.584 (0.056)
CV%	32.6	14.6	6.1	9.6
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.702	0.905	0.515	0.589
Min, Max	0.540, 0.864	0.811, 0.998	0.503, 0.573	0.521, 0.636

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

Timepoint Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
Pre-dose				
n	35	33	35	33
nBLQ (%)	35 (100.0)	33 (100.0)	35 (100.0)	33 (100.0)
Mean (SD)	0.000 (0.000)	0.000 (0.000)	0.000 (0.000)	0.000 (0.000)
CV%	NC	NC	NC	NC
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.000	0.000	0.000	0.000
Min, Max	0.000, 0.000	0.000, 0.000	0.000, 0.000	0.000, 0.000
0.5 hours post-dose				
n	35	33	35	33
nBLQ (%)	0	0	0	0
Mean (SD)	5.92 (3.91)	7.40 (3.74)	2.26 (1.82)	2.81 (1.44)
CV%	66.0	50.6	80.7	51.3
Geometric Mean (CV%)	4.89 (68.2)	6.58 (52.9)	1.74 (83.8)	2.43 (63.3)
Median	4.21	6.64	1.78	2.47
Min, Max	1.58, 14.5	1.87, 17.6	0.398, 8.94	0.479, 6.05

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

Timepoint Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
1 hour post-dose				
n	35	33	35	33
nBLQ (%)	0	0	0	0
Mean (SD)	7.38 (3.21)	8.30 (3.60)	2.90 (1.09)	3.31 (1.35)
CV%	43.5	43.4	37.6	40.7
Geometric Mean (CV%)	6.74 (45.5)	7.67 (40.8)	2.69 (42.4)	3.05 (43.8)
Median	6.99	7.60	2.98	3.13
Min, Max	3.19, 15.0	3.59, 19.9	1.25, 5.40	1.21, 6.50
1.5 hours post-dose				
n	35	32	35	32
nBLQ (%)	0	0	0	0
Mean (SD)	5.78 (2.76)	6.66 (2.52)	2.22 (0.716)	2.66 (1.03)
CV%	47.8	37.8	32.3	38.8
Geometric Mean (CV%)	5.24 (46.9)	6.25 (37.9)	2.10 (34.2)	2.48 (38.4)
Median	5.31	6.04	2.12	2.45
Min, Max	2.20, 15.5	2.17, 14.7	1.10, 3.64	1.35, 4.98

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

Timepoint Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
2 hours post-dose				
n	34	33	34	33
nBLQ (%)	0	0	0	0
Mean (SD)	4.98 (2.46)	5.52 (2.27)	1.94 (0.726)	2.16 (0.852)
CV%	49.3	41.2	37.4	39.4
Geometric Mean (CV%)	4.49 (48.2)	5.12 (40.8)	1.83 (36.1)	2.03 (37.3)
Median	4.27	4.66	1.72	1.94
Min, Max	1.67, 13.4	1.92, 12.9	0.988, 4.41	1.12, 4.53
3 hours post-dose				
n	35	32	35	32
nBLQ (%)	0	0	0	0
Mean (SD)	3.23 (1.73)	3.49 (1.71)	1.22 (0.444)	1.32 (0.587)
CV%	53.6	49.0	36.3	44.3
Geometric Mean (CV%)	2.87 (50.8)	3.15 (47.8)	1.15 (36.6)	1.24 (36.3)
Median	2.94	2.80	1.10	1.20
Min, Max	1.18, 8.83	1.20, 8.41	0.493, 2.73	0.647, 3.94

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

Timepoint Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
4 hours post-dose				
n	35	30	35	30
nBLQ (%)	0	0	0	0
Mean (SD)	2.26 (1.20)	2.55 (1.44)	0.845 (0.304)	0.874 (0.448)
CV%	53.1	56.7	36.0	51.2
Geometric Mean (CV%)	2.02 (50.5)	2.21 (58.7)	0.799 (34.5)	0.800 (41.9)
Median	2.03	2.16	0.747	0.818
Min, Max	0.785, 6.35	0.658, 5.84	0.437, 1.95	0.447, 2.75
6 hours post-dose				
n	35	32	35	32
nBLQ (%)	0	0	0	0
Mean (SD)	1.16 (0.677)	1.37 (0.888)	0.420 (0.181)	0.362 (0.193)
CV%	58.2	64.7	43.0	53.3
Geometric Mean (CV%)	1.02 (56.5)	1.15 (64.4)	0.387 (43.0)	0.324 (48.0)
Median	1.00	1.07	0.402	0.297
Min, Max	0.298, 3.70	0.414, 3.98	0.169, 1.05	0.165, 0.970

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

Timepoint Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
8 hours post-dose				
n	35	33	35	33
nBLQ (%)	0	0	1 (2.9)	3 (9.1)
Mean (SD)	0.774 (0.433)	0.881 (0.570)	0.260 (0.113)	0.218 (0.102)
CV%	56.0	64.6	43.3	46.5
Geometric Mean (CV%)	0.675 (57.8)	0.728 (69.6)	NC	NC
Median	0.668	0.744	0.228	0.186
Min, Max	0.179, 2.30	0.223, 2.24	0.146, 0.749	0.104, 0.547
12 hours post-dose				
n	35	33	35	33
nBLQ (%)	2 (5.7)	2 (6.1)	6 (17.1)	9 (27.3)
Mean (SD)	0.382 (0.263)	0.442 (0.320)	0.160 (0.069)	0.163 (0.059)
CV%	68.8	72.3	43.2	36.4
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.288	0.365	0.133	0.150
Min, Max	0.116, 1.30	0.116, 1.62	0.102, 0.391	0.104, 0.341

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

Timepoint Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
16 hours post-dose				
n	35	32	35	32
nBLQ (%)	6 (17.1)	8 (25.0)	21 (60.0)	25 (78.1)
Mean (SD)	0.265 (0.202)	0.304 (0.270)	0.138 (0.025)	0.156 (0.076)
CV%	76.3	89.0	18.5	48.5
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.196	0.199	0.134	0.112
Min, Max	0.101, 0.887	0.103, 1.37	0.101, 0.203	0.103, 0.298
24 hours post-dose				
n	35	33	35	33
nBLQ (%)	26 (74.3)	25 (75.8)	32 (91.4)	26 (78.8)
Mean (SD)	0.213 (0.115)	0.261 (0.079)	0.115 (0.006)	0.144 (0.022)
CV%	53.9	30.4	5.6	15.0
Geometric Mean (CV%)	NC	NC	NC	NC
Median	0.170	0.237	0.118	0.146
Min, Max	0.105, 0.415	0.188, 0.430	0.108, 0.120	0.120, 0.177

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.1.2
Summary of Plasma CYP Probes (Parent and Metabolite) Concentrations (ng/mL) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

Timepoint Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
48 hours post-dose				
n	35	33	35	33
nBLQ (%)	35 (100.0)	33 (100.0)	35 (100.0)	33 (100.0)
Mean (SD)	NC	NC	NC	NC
CV%	NC	NC	NC	NC
Geometric Mean (CV%)	NC	NC	NC	NC
Median	NC	NC	NC	NC
Min, Max	NC, NC	NC, NC	NC, NC	NC, NC

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; nBLQ = number of observations below the lower limit of quantification (BLQ); SD = standard deviation; LLOQ = lower limit of quantification; NC = not calculated.

Source: Listing 16.2.6.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010102.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.1
Summary of Plasma PK Parameters of CYB003 and Metabolites
PK Population

PK Parameter (Unit) Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
AUC_{0-t} (h*ng/mL)			
n	33	33	33
Mean (SD)	199 (50.3)	1560 (445)	60.4 (19.9)
CV%	25.2	28.4	33.0
Geometric Mean (CV%)	193 (26.9)	1500 (29.1)	57.2 (35.3)
Median	200	1460	58.7
Min, Max	114, 292	879, 2620	29.7, 104
AUC_{0-inf} (h*ng/mL)			
n	33	33	30
Mean (SD)	211 (51.5)	1700 (467)	70.7 (23.4)
CV%	24.4	27.4	33.0
Geometric Mean (CV%)	205 (26.2)	1640 (27.6)	66.8 (36.3)
Median	212	1620	69.4
Min, Max	120, 303	1020, 2940	32.3, 127

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; SD = standard deviation; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2; MR = metabolite ratio.

Source: Listing 16.2.6.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010201.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.1
Summary of Plasma PK Parameters of CYB003 and Metabolites
PK Population

PK Parameter (Unit) Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
C_{last} (ng/mL)			
n	33	33	33
Mean (SD)	1.74 (0.432)	19.0 (6.66)	1.70 (0.479)
CV%	24.8	35.0	28.2
Geometric Mean (CV%)	1.69 (23.7)	18.0 (34.1)	1.64 (27.4)
Median	1.65	18.1	1.54
Min, Max	1.11, 2.85	11.0, 36.9	1.17, 2.81
C_{max} (ng/mL)			
n	33	33	33
Mean (SD)	27.7 (6.80)	173 (47.8)	15.5 (5.20)
CV%	24.5	27.6	33.6
Geometric Mean (CV%)	26.9 (25.4)	167 (28.3)	14.7 (32.4)
Median	26.8	170	14.2
Min, Max	16.7, 43.0	95.6, 280	7.22, 34.0

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; SD = standard deviation; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2; MR = metabolite ratio.

Source: Listing 16.2.6.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010201.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.1
Summary of Plasma PK Parameters of CYB003 and Metabolites
PK Population

PK Parameter (Unit) Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
CL/F (L/h)			
n	33		
Mean (SD)	80.9 (22.0)		
CV%	27.2		
Geometric Mean (CV%)	78.2 (26.2)		
Median	75.5		
Min, Max	52.9, 134		
T_{max} (h)			
n	33	33	33
Mean (SD)	2.83 (0.996)	3.91 (1.11)	1.99 (0.884)
Median	2.93	3.97	2.00
Min, Max	1.05, 5.92	2.00, 5.97	1.00, 3.97

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; SD = standard deviation; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2; MR = metabolite ratio.

Source: Listing 16.2.6.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010201.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.1
Summary of Plasma PK Parameters of CYB003 and Metabolites
PK Population

PK Parameter (Unit) Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
T_{last} (h)			
n	33	33	33
Mean (SD)	20.8 (3.98)	20.8 (3.98)	10.9 (3.35)
Median	23.9	23.9	11.9
Min, Max	15.9, 24.3	15.9, 24.3	5.93, 16.1
K_{el} (1/h)			
n	33	33	31
Mean (SD)	0.156 (0.035)	0.139 (0.025)	0.225 (0.111)
CV%	22.3	18.3	49.3
Geometric Mean (CV%)	0.152 (22.8)	0.137 (18.8)	0.203 (48.1)
Median	0.152	0.141	0.186
Min, Max	0.094, 0.228	0.096, 0.190	0.080, 0.605

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; SD = standard deviation; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2; MR = metabolite ratio.

Source: Listing 16.2.6.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010201.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.1
Summary of Plasma PK Parameters of CYB003 and Metabolites
PK Population

PK Parameter (Unit) Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
$t_{1/2}$ (h)			
n	31	29	17
Mean (SD)	4.61 (0.968)	5.09 (0.919)	3.11 (1.21)
CV%	21.0	18.0	38.8
Geometric Mean (CV%)	4.51 (21.7)	5.01 (18.1)	2.88 (42.9)
Median	4.56	4.91	2.76
Min, Max	3.05, 6.03	3.65, 7.09	1.15, 5.56
V_z/F (L)			
n	33		
Mean (SD)	528 (114)		
CV%	21.6		
Geometric Mean (CV%)	515 (22.6)		
Median	527		
Min, Max	328, 814		

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; SD = standard deviation; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2; MR = metabolite ratio.

Source: Listing 16.2.6.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010201.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.1
Summary of Plasma PK Parameters of CYB003 and Metabolites
PK Population

PK Parameter (Unit) Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
MR_AUC$_{0-inf}$			
n		33	30
Mean (SD)		4.62 (1.42)	0.368 (0.088)
CV%		30.7	23.8
Geometric Mean (CV%)		4.41 (31.5)	0.358 (25.4)
Median		4.49	0.361
Min, Max		2.41, 8.04	0.192, 0.544
MR_AUC$_{0-t}$			
n		33	33
Mean (SD)		4.49 (1.43)	0.338 (0.081)
CV%		31.7	23.8
Geometric Mean (CV%)		4.28 (33.1)	0.329 (24.9)
Median		4.27	0.323
Min, Max		2.14, 8.09	0.178, 0.519

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; SD = standard deviation; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2; MR = metabolite ratio.

Source: Listing 16.2.6.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010201.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.1
Summary of Plasma PK Parameters of CYB003 and Metabolites
PK Population

PK Parameter (Unit) Statistic	CYB003 (N=35)	Psilocin-O-glucuronide (N=35)	4-HIAA (N=35)
MR_C_{max}			
n		33	33
Mean (SD)		3.54 (1.07)	0.642 (0.211)
CV%		30.1	32.8
Geometric Mean (CV%)		3.40 (28.8)	0.607 (35.8)
Median		3.39	0.633
Min, Max		1.99, 7.13	0.254, 1.03

Lower limit of quantification = 1.09 ng/mL for both CYB003 and 4-HIAA-d2, 10.9 ng/mL for Psilocin-O-glucuronide-d10.
Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; SD = standard deviation; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2; MR = metabolite ratio.

Source: Listing 16.2.6.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010201.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
AUC_{0-t} (h*ng/mL)				
n	32	31	22	22
Mean (SD)	17700 (5170)	19800 (6470)	10800 (2230)	11900 (2980)
CV%	29.2	32.7	20.7	25.0
Geometric Mean (CV%)	17000 (30.5)	18700 (37.8)	10500 (21.8)	11500 (27.9)
Median	17200	18600	10700	11500
Min, Max	7960, 27800	6560, 31300	6140, 15900	6150, 17000
AUC_{0-inf} (h*ng/mL)				
n	32	31	21	21
Mean (SD)	18200 (5170)	20500 (6680)	11900 (2430)	12500 (3190)
CV%	28.5	32.6	20.5	25.5
Geometric Mean (CV%)	17400 (29.9)	19300 (38.1)	11600 (21.9)	12100 (28.0)
Median	18200	19800	11700	11700
Min, Max	8020, 28400	6620, 32400	6440, 17500	6370, 18500

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
C_{last} (ng/mL)				
n	32	31	22	22
Mean (SD)	57.8 (36.0)	78.7 (63.5)	83.1 (65.1)	59.5 (51.9)
CV%	62.2	80.6	78.4	87.2
Geometric Mean (CV%)	47.1 (76.7)	57.7 (101.8)	55.3 (135.1)	41.6 (107.5)
Median	57.1	69.4	61.0	36.1
Min, Max	12.3, 153	11.0, 305	10.4, 199	10.7, 175
C_{max} (ng/mL)				
n	32	31	22	22
Mean (SD)	2000 (511)	1910 (469)	578 (99.0)	615 (180)
CV%	25.5	24.5	17.1	29.2
Geometric Mean (CV%)	1940 (24.9)	1860 (22.7)	571 (16.1)	596 (25.0)
Median	1900	1880	548	588
Min, Max	1270, 3240	1060, 3640	449, 849	385, 1280

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
CL/F (L/h)				
n	32	31		
Mean (SD)	5.98 (1.86)	5.57 (2.50)		
CV%	31.2	45.0		
Geometric Mean (CV%)	5.73 (29.9)	5.18 (38.1)		
Median	5.50	5.04		
Min, Max	3.52, 12.5	3.09, 15.1		
T_{max} (h)				
n	32	31	22	22
Mean (SD)	1.35 (0.776)	1.93 (1.47)	7.83 (2.43)	7.07 (2.58)
Median	1.00	1.50	7.97	6.97
Min, Max	0.500, 3.95	0.500, 7.95	2.93, 12.0	2.00, 12.0

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
T_{last} (h)				
n	32	31	22	22
Mean (SD)	33.7 (11.9)	34.0 (12.0)	38.1 (12.1)	42.5 (10.3)
Median	24.0	24.0	47.8	47.8
Min, Max	23.9, 48.1	23.9, 48.1	23.9, 48.3	23.9, 48.1
K_{el} (1/h)				
n	32	31	22	22
Mean (SD)	0.126 (0.036)	0.121 (0.038)	0.089 (0.021)	0.088 (0.029)
CV%	28.3	31.3	23.9	32.7
Geometric Mean (CV%)	0.121 (29.1)	0.116 (31.5)	0.087 (24.6)	0.084 (34.4)
Median	0.124	0.116	0.093	0.086
Min, Max	0.072, 0.215	0.066, 0.210	0.056, 0.144	0.036, 0.161

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
$t_{1/2}$ (h)				
n	32	30	14	18
Mean (SD)	5.96 (1.69)	6.26 (1.93)	8.67 (2.37)	8.39 (2.47)
CV%	28.4	30.9	27.4	29.5
Geometric Mean (CV%)	5.73 (29.1)	5.97 (32.1)	8.36 (28.9)	8.04 (31.2)
Median	5.57	5.97	8.56	8.00
Min, Max	3.22, 9.60	3.31, 10.4	4.81, 12.4	4.31, 13.0
V_z/F (L)				
n	32	31		
Mean (SD)	48.2 (8.66)	45.3 (8.04)		
CV%	18.0	17.8		
Geometric Mean (CV%)	47.4 (18.7)	44.6 (17.2)		
Median	49.2	43.7		
Min, Max	31.5, 66.4	28.1, 73.6		

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
MR_AUC$_{0-inf}$				
n			21	21
Mean (SD)			0.776 (0.093)	0.731 (0.128)
CV%			11.9	17.5
Geometric Mean (CV%)			0.770 (13.3)	0.720 (18.7)
Median			0.777	0.729
Min, Max			0.487, 0.918	0.418, 1.04
MR_AUC$_{0-t}$				
n			22	22
Mean (SD)			0.733 (0.103)	0.704 (0.123)
CV%			14.1	17.5
Geometric Mean (CV%)			0.725 (14.8)	0.694 (18.6)
Median			0.719	0.699
Min, Max			0.474, 0.939	0.404, 1.01

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Caffeine (LLOQ = 10 ng/mL for Caffeine, 10 ng/mL for Paraxanthine)

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)
MR_C_{max}				
n			22	22
Mean (SD)			0.333 (0.074)	0.355 (0.094)
CV%			22.1	26.6
Geometric Mean (CV%)			0.325 (24.1)	0.343 (29.0)
Median			0.332	0.365
Min, Max			0.183, 0.469	0.161, 0.589

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
AUC_{0-t} (h*ng/mL)				
n	30	29	30	29
Mean (SD)	134 (37.4)	157 (40.9)	1910 (884)	1890 (877)
CV%	28.0	26.1	46.4	46.5
Geometric Mean (CV%)	129 (28.6)	151 (28.8)	1750 (42.0)	1740 (40.3)
Median	128	151	1650	1800
Min, Max	70.7, 206	74.9, 238	914, 5090	962, 5430
AUC_{0-inf} (h*ng/mL)				
n	29	28	7	4
Mean (SD)	147 (41.0)	172 (42.3)	2220 (1040)	2250 (613)
CV%	27.8	24.6	46.9	27.3
Geometric Mean (CV%)	142 (28.5)	167 (27.0)	2040 (46.5)	2180 (27.3)
Median	151	166	2070	2150
Min, Max	77.4, 242	88.8, 252	1170, 4120	1630, 3050

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
C_{last} (ng/mL)				
n	30	29	30	29
Mean (SD)	0.833 (0.252)	0.856 (0.266)	19.8 (12.3)	21.2 (12.0)
CV%	30.3	31.1	61.9	56.5
Geometric Mean (CV%)	0.797 (30.9)	0.819 (30.7)	17.4 (53.2)	19.0 (47.1)
Median	0.868	0.778	16.8	18.2
Min, Max	0.506, 1.52	0.510, 1.57	5.34, 73.5	9.93, 71.0
C_{max} (ng/mL)				
n	30	29	30	29
Mean (SD)	25.9 (9.74)	24.5 (6.32)	75.1 (32.1)	72.6 (32.2)
CV%	37.6	25.8	42.7	44.4
Geometric Mean (CV%)	24.2 (39.4)	23.7 (26.9)	69.4 (41.3)	67.4 (38.6)
Median	26.8	23.1	66.9	65.5
Min, Max	9.78, 54.3	12.4, 39.1	34.2, 157	37.4, 196

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
CL/F (L/h)				
n	29	28		
Mean (SD)	238 (67.5)	202 (58.4)		
CV%	28.4	28.9		
Geometric Mean (CV%)	229 (28.5)	195 (27.0)		
Median	216	196		
Min, Max	134, 420	129, 366		
T_{max} (h)				
n	30	29	30	29
Mean (SD)	1.85 (0.809)	2.61 (1.14)	4.00 (1.43)	5.60 (2.11)
Median	1.53	2.03	3.97	5.92
Min, Max	1.00, 4.02	0.983, 5.92	2.00, 7.97	2.93, 12.1

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
T_{last} (h)				
n	30	29	30	29
Mean (SD)	30.4 (10.7)	35.5 (12.2)	47.9 (0.134)	47.0 (4.45)
Median	24.0	24.0	47.8	47.9
Min, Max	23.9, 48.0	23.9, 48.0	47.7, 48.3	23.9, 48.1
K_{el} (1/h)				
n	29	28	29	25
Mean (SD)	0.076 (0.030)	0.072 (0.034)	0.030 (0.010)	0.028 (0.007)
CV%	38.8	47.9	32.2	25.2
Geometric Mean (CV%)	0.070 (45.1)	0.064 (50.8)	0.028 (35.2)	0.027 (25.0)
Median	0.079	0.056	0.029	0.026
Min, Max	0.028, 0.146	0.029, 0.141	0.012, 0.056	0.018, 0.045

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
$t_{1/2}$ (h)				
n	8	15	8	2
Mean (SD)	10.7 (5.10)	9.77 (4.39)	17.5 (2.57)	16.1 (0.917)
CV%	47.8	44.9	14.7	5.7
Geometric Mean (CV%)	9.52 (56.6)	8.85 (48.8)	17.3 (16.2)	16.1 (5.7)
Median	10.8	7.00	18.0	16.1
Min, Max	4.74, 15.9	4.90, 16.4	12.4, 20.1	15.5, 16.8
V_z/F (L)				
n	29	28		
Mean (SD)	3440 (1190)	3290 (1410)		
CV%	34.5	42.8		
Geometric Mean (CV%)	3250 (35.4)	3040 (41.5)		
Median	3170	3160		
Min, Max	1750, 6410	1510, 7390		

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
MR_AUC$_{0-inf}$				
n			7	4
Mean (SD)			19.3 (14.5)	13.9 (3.82)
CV%			75.1	27.6
Geometric Mean (CV%)			16.0 (68.6)	13.4 (29.3)
Median			13.3	13.9
Min, Max			6.95, 49.9	9.47, 18.1
MR_AUC$_{0-t}$				
n			30	29
Mean (SD)			14.3 (7.78)	11.8 (5.43)
CV%			54.5	46.1
Geometric Mean (CV%)			12.7 (49.5)	10.8 (43.3)
Median			11.8	10.5
Min, Max			5.23, 44.5	4.91, 25.9

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Bupropion (LLOQ = 0.5 ng/mL for Bupropion, 1 ng/mL for Hydroxybupropion)

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)
MR_C_{max}				
n			30	29
Mean (SD)			3.16 (2.35)	2.94 (1.34)
CV%			74.4	45.6
Geometric Mean (CV%)			2.69 (58.9)	2.67 (47.0)
Median			3.04	2.72
Min, Max			0.972, 14.2	1.10, 6.76

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
AUC_{0-t} (h*ng/mL)				
n	34	33	32	30
Mean (SD)	142 (56.5)	147 (61.3)	1040 (337)	1140 (457)
CV%	39.7	41.7	32.3	40.1
Geometric Mean (CV%)	132 (41.2)	137 (39.5)	988 (36.0)	1060 (42.0)
Median	137	142	1050	1110
Min, Max	53.2, 312	62.7, 364	408, 1830	502, 2570
AUC_{0-inf} (h*ng/mL)				
n	30	32	31	29
Mean (SD)	155 (59.7)	157 (63.7)	1090 (319)	1180 (455)
CV%	38.6	40.6	29.3	38.5
Geometric Mean (CV%)	145 (38.0)	146 (38.8)	1040 (31.2)	1100 (40.5)
Median	151	153	1110	1140
Min, Max	59.2, 359	66.9, 382	546, 1840	520, 2590

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
C_{last} (ng/mL)				
n	34	33	32	30
Mean (SD)	2.08 (0.956)	1.81 (0.825)	3.31 (2.43)	4.57 (9.68)
CV%	45.9	45.5	73.7	211.6
Geometric Mean (CV%)	1.91 (43.4)	1.67 (40.4)	2.59 (79.3)	2.44 (110.9)
Median	1.78	1.58	1.82	1.76
Min, Max	1.04, 4.70	1.01, 5.06	1.00, 9.32	1.04, 54.2
C_{max} (ng/mL)				
n	34	33	32	30
Mean (SD)	38.8 (22.8)	32.1 (16.1)	110 (42.9)	116 (53.0)
CV%	58.6	50.2	39.1	45.7
Geometric Mean (CV%)	33.9 (55.5)	29.0 (47.2)	101 (46.6)	104 (53.6)
Median	34.3	31.3	105	113
Min, Max	10.8, 123	13.7, 84.9	33.1, 217	30.9, 256

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
CL/F (L/h)				
n	32	32		
Mean (SD)	173 (66.5)	167 (61.8)		
CV%	38.4	36.9		
Geometric Mean (CV%)	162 (38.3)	157 (38.8)		
Median	154	150		
Min, Max	63.8, 387	60.0, 342		
T_{max} (h)				
n	34	33	32	30
Mean (SD)	1.37 (1.29)	2.56 (1.78)	5.46 (1.42)	6.18 (1.95)
Median	1.00	2.07	5.93	5.95
Min, Max	0.500, 5.93	0.500, 7.98	2.93, 8.07	2.95, 12.0

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
T_{last} (h)				
n	34	33	32	30
Mean (SD)	14.9 (7.03)	15.3 (6.93)	37.5 (12.1)	40.4 (11.6)
Median	11.9	12.0	47.8	47.8
Min, Max	7.97, 47.8	7.95, 47.8	23.9, 48.3	16.1, 48.0
K_{el} (1/h)				
n	32	32	32	29
Mean (SD)	0.224 (0.073)	0.234 (0.074)	0.118 (0.040)	0.108 (0.039)
CV%	32.4	31.7	33.9	36.3
Geometric Mean (CV%)	0.213 (32.9)	0.221 (38.5)	0.112 (34.4)	0.103 (32.4)
Median	0.200	0.235	0.098	0.092
Min, Max	0.099, 0.412	0.063, 0.424	0.063, 0.193	0.071, 0.224

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
$t_{1/2}$ (h)				
n	26	28	31	29
Mean (SD)	3.23 (1.06)	3.25 (1.73)	6.48 (2.09)	7.03 (1.92)
CV%	33.0	53.3	32.2	27.3
Geometric Mean (CV%)	3.08 (31.5)	3.00 (37.9)	6.15 (34.4)	6.73 (32.4)
Median	3.16	2.88	7.02	7.51
Min, Max	1.68, 7.03	1.64, 11.0	3.59, 11.1	3.10, 9.80
V_z/F (L)				
n	32	32		
Mean (SD)	837 (378)	762 (281)		
CV%	45.1	36.8		
Geometric Mean (CV%)	761 (47.1)	709 (41.2)		
Median	810	792		
Min, Max	328, 1910	328, 1430		

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
MR_AUC$_{0-inf}$				
n			30	29
Mean (SD)			7.94 (2.69)	8.28 (2.97)
CV%			33.8	35.8
Geometric Mean (CV%)			7.46 (39.0)	7.66 (45.7)
Median			7.97	8.39
Min, Max			3.01, 14.5	2.10, 15.2
MR_AUC$_{0-t}$				
n			32	30
Mean (SD)			8.21 (3.02)	8.51 (3.10)
CV%			36.8	36.4
Geometric Mean (CV%)			7.58 (45.9)	7.87 (45.7)
Median			8.26	8.75
Min, Max			2.20, 15.4	2.11, 16.1

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC$_{0-inf}$, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC$_{0-inf}$ was excluded from summary if %AUC$_{extrap} > 20$%. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Losartan (LLOQ = 1 ng/mL for Losartan, 1 ng/mL for Losartan Carboxylic Acid)

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)
MR_C_{max}				
n			32	30
Mean (SD)			3.45 (1.93)	3.91 (1.76)
CV%			55.9	45.1
Geometric Mean (CV%)			2.97 (62.5)	3.47 (56.6)
Median			3.17	4.17
Min, Max			0.885, 9.41	0.941, 8.13

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
AUC_{0-t} (h*ng/mL)				
n	35	31	24	27
Mean (SD)	645 (718)	586 (537)	369 (81.2)	332 (128)
CV%	111.4	91.7	22.0	38.6
Geometric Mean (CV%)	458 (93.5)	416 (103.3)	361 (20.6)	305 (46.5)
Median	518	423	358	329
Min, Max	109, 4150	77.8, 2590	257, 638	120, 578
AUC_{0-inf} (h*ng/mL)				
n	29	10	23	17
Mean (SD)	746 (764)	979 (630)	376 (83.6)	399 (100)
CV%	102.3	64.3	22.3	25.1
Geometric Mean (CV%)	555 (85.7)	846 (59.7)	368 (20.8)	387 (26.4)
Median	602	885	360	354
Min, Max	153, 4180	358, 2600	259, 647	215, 581

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
C_{last} (ng/mL)				
n	35	31	24	27
Mean (SD)	6.82 (4.34)	9.56 (8.76)	3.53 (2.13)	2.33 (1.42)
CV%	63.6	91.6	60.4	60.8
Geometric Mean (CV%)	5.60 (71.6)	6.84 (96.3)	2.93 (71.9)	2.02 (55.7)
Median	5.38	6.02	3.56	1.93
Min, Max	2.03, 17.7	2.10, 35.1	1.04, 9.56	1.02, 6.33
C_{max} (ng/mL)				
n	35	31	24	27
Mean (SD)	267 (156)	224 (177)	149 (45.5)	108 (70.6)
CV%	58.5	79.0	30.5	65.2
Geometric Mean (CV%)	225 (68.8)	154 (122.1)	143 (32.3)	87.2 (82.6)
Median	259	201	148	85.7
Min, Max	64.9, 755	23.1, 616	79.7, 243	18.1, 323

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
CL/F (L/h)				
n	29	10		
Mean (SD)	46.0 (32.9)	26.9 (14.3)		
CV%	71.6	53.3		
Geometric Mean (CV%)	36.0 (85.7)	23.6 (59.7)		
Median	33.2	22.7		
Min, Max	4.79, 131	7.69, 55.9		
T_{max} (h)				
n	35	31	24	27
Mean (SD)	2.56 (1.11)	4.65 (1.95)	2.61 (1.29)	4.71 (2.05)
Median	2.02	4.00	2.00	4.12
Min, Max	1.50, 5.93	2.00, 8.08	1.50, 5.93	2.00, 8.08

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
T_{last} (h)				
n	35	31	24	27
Mean (SD)	8.24 (3.68)	11.4 (4.35)	10.1 (2.33)	14.3 (3.38)
Median	7.97	11.9	9.92	12.0
Min, Max	3.97, 23.9	5.92, 24.0	7.93, 15.9	11.9, 24.0
K_{el} (1/h)				
n	29	10	23	17
Mean (SD)	0.729 (0.248)	0.470 (0.131)	0.527 (0.102)	0.404 (0.108)
CV%	34.1	27.8	19.3	26.7
Geometric Mean (CV%)	0.685 (39.1)	0.452 (30.7)	0.518 (20.1)	0.390 (28.7)
Median	0.716	0.425	0.534	0.368
Min, Max	0.213, 1.43	0.244, 0.647	0.335, 0.745	0.211, 0.593

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
$t_{1/2}$ (h)				
n	29	10	23	17
Mean (SD)	1.09 (0.519)	1.60 (0.525)	1.36 (0.279)	1.85 (0.559)
CV%	47.5	32.8	20.5	30.2
Geometric Mean (CV%)	1.01 (39.1)	1.53 (30.7)	1.34 (20.1)	1.78 (28.7)
Median	0.968	1.63	1.30	1.89
Min, Max	0.484, 3.26	1.07, 2.84	0.931, 2.07	1.17, 3.29
V_z/F (L)				
n	29	10		
Mean (SD)	58.2 (28.1)	56.6 (24.8)		
CV%	48.2	43.8		
Geometric Mean (CV%)	52.6 (48.2)	52.3 (42.6)		
Median	51.9	46.5		
Min, Max	22.5, 139	31.5, 106		

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
MR_AUC$_{0-inf}$				
n			20	10
Mean (SD)			0.946 (0.528)	0.602 (0.402)
CV%			55.8	66.8
Geometric Mean (CV%)			0.807 (65.8)	0.508 (66.5)
Median			0.812	0.496
Min, Max			0.205, 1.87	0.177, 1.55
MR_AUC$_{0-t}$				
n			24	26
Mean (SD)			1.05 (0.551)	0.936 (0.616)
CV%			52.4	65.9
Geometric Mean (CV%)			0.898 (66.8)	0.726 (96.8)
Median			0.984	0.787
Min, Max			0.202, 1.98	0.065, 2.58

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Omeprazole (LLOQ = 2 ng/mL for Omeprazole, 1 ng/mL for 5-Hydroxy Omeprazole)

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)
MR_C_{max}				
n			24	26
Mean (SD)			0.789 (0.404)	0.702 (0.451)
CV%			51.2	64.1
Geometric Mean (CV%)			0.683 (62.7)	0.549 (95.0)
Median			0.758	0.505
Min, Max			0.170, 1.63	0.045, 1.68

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
AUC_{0-t} (h*ng/mL)				
n	34	33	34	33
Mean (SD)	202 (132)	216 (143)	147 (43.9)	147 (45.7)
CV%	65.5	66.2	29.8	31.1
Geometric Mean (CV%)	164 (77.5)	172 (82.4)	141 (30.8)	140 (36.1)
Median	192	187	138	140
Min, Max	37.2, 645	33.5, 611	65.1, 260	45.3, 270
AUC_{0-inf} (h*ng/mL)				
n	34	32	30	28
Mean (SD)	210 (135)	221 (148)	171 (41.7)	175 (45.0)
CV%	64.4	67.0	24.4	25.7
Geometric Mean (CV%)	172 (75.1)	176 (81.6)	166 (24.5)	170 (25.7)
Median	198	192	160	169
Min, Max	39.4, 656	36.3, 620	103, 266	95.1, 310

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
C_{last} (ng/mL)				
n	34	33	34	33
Mean (SD)	1.24 (0.511)	1.34 (0.780)	1.76 (0.703)	1.79 (0.780)
CV%	41.4	58.4	40.0	43.6
Geometric Mean (CV%)	1.14 (42.3)	1.16 (57.2)	1.58 (54.3)	1.59 (57.0)
Median	1.10	1.15	1.73	1.80
Min, Max	0.540, 2.55	0.508, 3.17	0.503, 3.15	0.521, 3.50
C_{max} (ng/mL)				
n	34	33	34	33
Mean (SD)	30.4 (15.5)	26.5 (13.1)	14.2 (6.61)	12.2 (5.27)
CV%	51.2	49.5	46.6	43.1
Geometric Mean (CV%)	26.5 (59.3)	23.3 (58.1)	12.8 (49.0)	11.1 (50.0)
Median	28.4	25.5	13.3	11.3
Min, Max	6.92, 72.7	6.24, 51.8	4.11, 32.0	2.76, 29.3

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
CL/F (L/h)				
n	34	32		
Mean (SD)	143 (110)	143 (113)		
CV%	76.8	79.3		
Geometric Mean (CV%)	114 (75.1)	111 (81.6)		
Median	98.5	101		
Min, Max	29.7, 495	31.4, 537		
T_{max} (h)				
n	34	33	34	33
Mean (SD)	2.03 (0.954)	3.09 (1.57)	2.39 (1.61)	3.63 (2.38)
Median	1.98	2.93	1.52	2.93
Min, Max	0.983, 4.02	1.50, 7.98	0.967, 7.95	0.500, 7.98

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
T_{last} (h)				
n	34	33	34	33
Mean (SD)	21.5 (8.29)	22.4 (7.93)	26.8 (7.82)	26.8 (7.93)
Median	23.9	23.9	24.0	24.0
Min, Max	7.97, 47.9	11.9, 47.9	23.9, 48.0	23.7, 47.9
K_{el} (1/h)				
n	34	32	34	32
Mean (SD)	0.171 (0.048)	0.171 (0.051)	0.085 (0.017)	0.088 (0.022)
CV%	28.2	29.8	20.4	25.1
Geometric Mean (CV%)	0.164 (29.3)	0.164 (31.6)	0.084 (20.1)	0.086 (23.6)
Median	0.156	0.158	0.081	0.086
Min, Max	0.079, 0.270	0.076, 0.279	0.054, 0.135	0.058, 0.150

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
$t_{1/2}$ (h)				
n	33	31	13	15
Mean (SD)	4.40 (1.35)	4.35 (1.40)	7.95 (1.57)	7.75 (2.30)
CV%	30.7	32.2	19.8	29.7
Geometric Mean (CV%)	4.22 (29.8)	4.16 (30.3)	7.80 (21.3)	7.44 (30.4)
Median	4.52	4.33	7.84	7.75
Min, Max	2.56, 8.72	2.48, 9.09	5.13, 10.3	4.64, 12.0
V_z/F (L)				
n	34	32		
Mean (SD)	762 (372)	768 (429)		
CV%	48.9	55.8		
Geometric Mean (CV%)	691 (45.3)	678 (52.6)		
Median	622	658		
Min, Max	301, 1830	323, 2250		

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
MR_AUC$_{0-inf}$				
n			30	27
Mean (SD)			1.32 (1.13)	1.37 (1.08)
CV%			85.2	78.9
Geometric Mean (CV%)			0.999 (87.4)	1.04 (89.6)
Median			1.04	1.08
Min, Max			0.197, 5.17	0.196, 4.65
MR_AUC$_{0-t}$				
n			34	33
Mean (SD)			1.18 (1.17)	1.12 (1.04)
CV%			99.2	92.5
Geometric Mean (CV%)			0.811 (110.0)	0.764 (119.6)
Median			0.760	0.826
Min, Max			0.095, 5.46	0.078, 4.85

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Metoprolol (LLOQ = 0.5 ng/mL for Metoprolol, 0.5 ng/mL for Alpha-Hydroxymetoprolol)

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)
MR_C_{max}				
n			34	33
Mean (SD)			0.640 (0.604)	0.643 (0.590)
CV%			94.4	91.8
Geometric Mean (CV%)			0.455 (102.6)	0.449 (111.1)
Median			0.423	0.457
Min, Max			0.058, 2.53	0.051, 2.84

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
AUC_{0-t} (h*ng/mL)				
n	35	33	35	33
Mean (SD)	26.4 (13.8)	30.1 (15.2)	9.56 (3.44)	10.4 (4.21)
CV%	52.1	50.5	36.0	40.4
Geometric Mean (CV%)	23.6 (50.4)	27.1 (47.1)	9.01 (35.9)	9.69 (39.7)
Median	23.5	25.3	9.26	9.33
Min, Max	9.82, 69.6	11.5, 81.2	4.96, 18.6	4.78, 23.2
AUC_{0-inf} (h*ng/mL)				
n	35	33	31	22
Mean (SD)	27.7 (14.5)	31.4 (15.8)	10.5 (3.67)	10.5 (4.74)
CV%	52.5	50.2	35.0	45.3
Geometric Mean (CV%)	24.7 (50.2)	28.4 (46.9)	9.94 (33.8)	9.64 (41.6)
Median	25.1	26.5	9.66	8.81
Min, Max	10.3, 73.1	12.0, 84.8	6.19, 20.8	5.04, 23.8

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
C_{last} (ng/mL)				
n	35	33	35	33
Mean (SD)	0.186 (0.079)	0.216 (0.094)	0.136 (0.033)	0.144 (0.031)
CV%	42.2	43.8	24.5	21.6
Geometric Mean (CV%)	0.173 (39.1)	0.199 (41.3)	0.133 (21.0)	0.141 (21.4)
Median	0.170	0.200	0.130	0.146
Min, Max	0.101, 0.415	0.103, 0.495	0.101, 0.267	0.104, 0.224
C_{max} (ng/mL)				
n	35	33	35	33
Mean (SD)	8.11 (3.62)	8.96 (3.56)	3.26 (1.53)	3.59 (1.41)
CV%	44.7	39.7	46.9	39.4
Geometric Mean (CV%)	7.36 (47.4)	8.41 (36.1)	2.96 (47.1)	3.32 (41.8)
Median	7.38	7.87	3.29	3.40
Min, Max	3.70, 15.5	4.60, 19.9	1.25, 8.94	1.55, 6.50

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
CL/F (L/h)				
n	35	33		
Mean (SD)	112 (51.8)	96.5 (40.8)		
CV%	46.1	42.2		
Geometric Mean (CV%)	101 (50.2)	88.2 (46.9)		
Median	99.7	94.4		
Min, Max	34.2, 243	29.5, 208		
T_{max} (h)				
n	35	33	35	33
Mean (SD)	0.962 (0.330)	0.884 (0.304)	0.976 (0.342)	0.915 (0.314)
Median	1.00	1.00	1.00	1.00
Min, Max	0.500, 2.00	0.500, 1.52	0.500, 2.00	0.500, 1.52

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
T_{last} (h)				
n	35	33	35	33
Mean (SD)	17.1 (4.60)	16.6 (4.78)	13.6 (4.24)	13.8 (5.92)
Median	15.9	15.9	11.9	12.0
Min, Max	7.97, 24.0	7.95, 24.0	5.95, 23.9	5.92, 24.3
K_{el} (1/h)				
n	35	33	31	23
Mean (SD)	0.186 (0.069)	0.191 (0.073)	0.224 (0.106)	0.305 (0.140)
CV%	37.0	38.5	47.4	46.0
Geometric Mean (CV%)	0.172 (42.6)	0.177 (41.4)	0.197 (58.8)	0.265 (66.2)
Median	0.193	0.186	0.224	0.312
Min, Max	0.073, 0.374	0.079, 0.376	0.053, 0.581	0.049, 0.546

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
$t_{1/2}$ (h)				
n	26	23	20	18
Mean (SD)	3.69 (1.23)	3.75 (1.34)	2.80 (0.780)	2.16 (0.783)
CV%	33.3	35.7	27.8	36.3
Geometric Mean (CV%)	3.52 (31.5)	3.53 (36.2)	2.70 (29.6)	2.04 (34.2)
Median	3.43	3.58	2.61	1.96
Min, Max	1.85, 6.85	1.84, 7.10	1.19, 4.54	1.27, 4.10
V_z/F (L)				
n	35	33		
Mean (SD)	637 (278)	537 (209)		
CV%	43.6	38.9		
Geometric Mean (CV%)	587 (42.1)	498 (41.7)		
Median	570	488		
Min, Max	298, 1350	235, 930		

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
MR_AUC$_{0-inf}$				
n			31	22
Mean (SD)			0.415 (0.180)	0.374 (0.212)
CV%			43.5	56.7
Geometric Mean (CV%)			0.379 (45.5)	0.326 (56.4)
Median			0.411	0.295
Min, Max			0.146, 0.987	0.153, 0.831
MR_AUC$_{0-t}$				
n			35	33
Mean (SD)			0.402 (0.187)	0.381 (0.188)
CV%			46.5	49.3
Geometric Mean (CV%)			0.364 (48.7)	0.340 (51.2)
Median			0.375	0.340
Min, Max			0.140, 0.927	0.153, 0.819

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.2.2
Summary of Plasma PK Parameters of CYP Probes (Parent and Metabolite) With and Without CYB003
PK Population

CYP Probe: Midazolam (LLOQ = 0.1 ng/mL for Midazolam, 0.1 ng/mL for 1-Hydroxy Midazolam)

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)
MR_C_{max}				
n			35	33
Mean (SD)			0.423 (0.192)	0.412 (0.192)
CV%			45.4	46.6
Geometric Mean (CV%)			0.383 (48.1)	0.377 (44.0)
Median			0.390	0.356
Min, Max			0.157, 0.883	0.174, 1.07

Geometric mean and Geometric CV% were not presented when the minimum concentration was zero at the respective time point.
Values that were BLQ were imputed to zero for all measurements taken prior to dosing and before the first measurable concentration. All other BLQ values were set to missing. K_{el} and dependent parameters (e.g., AUC_{0-inf}, $t_{½}$, CL/F, V_z/F) were excluded from summary if the adjusted $r^2 < 0.8$. AUC_{0-inf} was excluded from summary if $\%AUC_{extrap} > 20\%$. $t_{½}$ was excluded from summary if Span < 2. All estimates excluded in this manner are presented in listings.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; SD = standard deviation; LLOQ = lower limit of quantification; MR = metabolite ratio; NC = not calculated.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t1402010202.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Caffeine

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=33)	CYB003 + Caffeine (Day 8) (N=33)	Comparison of CYB003 + Caffeine (Test) versus Caffeine Alone (Reference)	Paraxanthine Alone (Day 1) (N=33)	CYB003 + Paraxanthine (Day 8) (N=33)	Comparison of CYB003 + Paraxanthine (Test) versus Paraxanthine Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	29	29		17	17	
Geometric LS Mean	17200	19500		11600	12300	
Geometric Mean Ratio [a]			1.13			1.06
Two One-sided 90% CI Limits			(1.05, 1.22)			(0.976, 1.15)
Intra-Participant CV% [b]			16.8			13.5
AUC_{0-t} (h*ng/mL)						
n	29	29		19	19	
Geometric LS Mean	16700	18800		10600	11600	
Geometric Mean Ratio [a]			1.13			1.10
Two One-sided 90% CI Limits			(1.05, 1.22)			(1.02, 1.18)
Intra-Participant CV% [b]			16.9			13.4

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Caffeine

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=33)	CYB003 + Caffeine (Day 8) (N=33)	Comparison of CYB003 + Caffeine (Test) versus Caffeine Alone (Reference)	Paraxanthine Alone (Day 1) (N=33)	CYB003 + Paraxanthine (Day 8) (N=33)	Comparison of CYB003 + Paraxanthine (Test) versus Paraxanthine Alone (Reference)
C_{max} (ng/mL)						
n	29	29		19	19	
Geometric LS Mean	1950	1870		569	604	
Geometric Mean Ratio [a]			0.958			1.06
Two One-sided 90% CI Limits			(0.871, 1.06)			(0.984, 1.15)
Intra-Participant CV% [b]			21.8			13.5

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Bupropion

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=33)	CYB003 + Bupropion (Day 8) (N=33)	Comparison of CYB003 + Bupropion (Test) versus Bupropion Alone (Reference)	Hydroxybupropion Alone (Day 1) (N=33)	CYB003 + Hydroxybupropion (Day 8) (N=33)	Comparison of CYB003 + Hydroxybupropion (Test) versus Hydroxybupropion Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	26	26		3	3	
Geometric LS Mean	145	169		1780	2130	
Geometric Mean Ratio [a]			1.17			1.20
Two One-sided 90% CI Limits			(1.08, 1.27)			(0.907, 1.58)
Intra-Participant CV% [b]			17.0			11.7
AUC_{0-t} (h*ng/mL)						
n	28	28		28	28	
Geometric LS Mean	130	152		1740	1770	
Geometric Mean Ratio [a]			1.17			1.02
Two One-sided 90% CI Limits			(1.08, 1.27)			(0.953, 1.08)
Intra-Participant CV% [b]			18.0			14.3

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Bupropion

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=33)	CYB003 + Bupropion (Day 8) (N=33)	Comparison of CYB003 + Bupropion (Test) versus Bupropion Alone (Reference)	Hydroxybupropion Alone (Day 1) (N=33)	CYB003 + Hydroxybupropion (Day 8) (N=33)	Comparison of CYB003 + Hydroxybupropion (Test) versus Hydroxybupropion Alone (Reference)
C_{max} (ng/mL)						
n	28	28		28	28	
Geometric LS Mean	24.5	23.7		68.4	68.6	
Geometric Mean Ratio [a]			0.966			1.00
Two One-sided 90% CI Limits			(0.852, 1.10)			(0.941, 1.07)
Intra-Participant CV% [b]			28.2			13.9

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Losartan

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=33)	CYB003 + Losartan (Day 8) (N=33)	Comparison of CYB003 + Losartan (Test) versus Losartan Alone (Reference)	Losartan Carboxylic Acid Alone (Day 1) (N=33)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=33)	Comparison of CYB003 + Losartan Carboxylic Acid (Test) versus Losartan Carboxylic Acid Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	28	28		28	28	
Geometric LS Mean	143	147		1040	1130	
Geometric Mean Ratio [a]			1.03			1.09
Two One-sided 90% CI Limits			(0.966, 1.09)			(1.03, 1.15)
Intra-Participant CV% [b]			13.3			12.0
AUC_{0-t} (h*ng/mL)						
n	32	32		30	30	
Geometric LS Mean	130	135		978	1060	
Geometric Mean Ratio [a]			1.04			1.08
Two One-sided 90% CI Limits			(0.970, 1.11)			(1.02, 1.15)
Intra-Participant CV% [b]			15.5			14.0

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Losartan

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=33)	CYB003 + Losartan (Day 8) (N=33)	Comparison of CYB003 + Losartan (Test) versus Losartan Alone (Reference)	Losartan Carboxylic Acid Alone (Day 1) (N=33)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=33)	Comparison of CYB003 + Losartan Carboxylic Acid (Test) versus Losartan Carboxylic Acid Alone (Reference)
C_{max} (ng/mL)						
n	32	32		30	30	
Geometric LS Mean	33.2	28.6		100	104	
Geometric Mean Ratio [a]			0.862			1.04
Two One-sided 90% CI Limits			(0.737, 1.01)			(0.975, 1.10)
Intra-Participant CV% [b]			38.4			14.0

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Omeprazole

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=33)	CYB003 + Omeprazole (Day 8) (N=33)	Comparison of CYB003 + Omeprazole (Test) versus Omeprazole Alone (Reference)	5-Hydroxy Omeprazole Alone (Day 1) (N=33)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=33)	Comparison of CYB003 + 5-Hydroxy Omeprazole (Test) versus 5-Hydroxy Omeprazole Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	9	9		13	13	
Geometric LS Mean	781	931		398	410	
Geometric Mean Ratio [a]			1.19			1.03
Two One-sided 90% CI Limits			(1.03, 1.37)			(0.918, 1.16)
Intra-Participant CV% [b]			16.3			16.8
AUC_{0-t} (h*ng/mL)						
n	31	31		21	21	
Geometric LS Mean	475	416		364	326	
Geometric Mean Ratio [a]			0.876			0.897
Two One-sided 90% CI Limits			(0.723, 1.06)			(0.768, 1.05)
Intra-Participant CV% [b]			46.7			29.6

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Omeprazole

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=33)	CYB003 + Omeprazole (Day 8) (N=33)	Comparison of CYB003 + Omeprazole (Test) versus Omeprazole Alone (Reference)	5-Hydroxy Omeprazole Alone (Day 1) (N=33)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=33)	Comparison of CYB003 + 5-Hydroxy Omeprazole (Test) versus 5-Hydroxy Omeprazole Alone (Reference)
C_{max} (ng/mL)						
n	31	31		21	21	
Geometric LS Mean	234	154		145	99.8	
Geometric Mean Ratio [a]			0.659			0.690
Two One-sided 90% CI Limits			(0.497, 0.873)			(0.529, 0.899)
Intra-Participant CV% [b]			72.9			53.1

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Metoprolol

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=33)	CYB003 + Metoprolol (Day 8) (N=33)	Comparison of CYB003 + Metoprolol (Test) versus Metoprolol Alone (Reference)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=33)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=33)	Comparison of CYB003 + Alpha-Hydroxymetoprolol (Test) versus Alpha-Hydroxymetoprolol Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	31	31		25	25	
Geometric LS Mean	170	180		167	164	
Geometric Mean Ratio [a]			1.06			0.984
Two One-sided 90% CI Limits			(0.992, 1.13)			(0.946, 1.02)
Intra-Participant CV% [b]			15.1			8.1
AUC_{0-t} (h*ng/mL)						
n	32	32		32	32	
Geometric LS Mean	163	176		140	137	
Geometric Mean Ratio [a]			1.08			0.974
Two One-sided 90% CI Limits			(1.01, 1.16)			(0.927, 1.02)
Intra-Participant CV% [b]			16.1			11.7

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Metoprolol

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=33)	CYB003 + Metoprolol (Day 8) (N=33)	Comparison of CYB003 + Metoprolol (Test) versus Metoprolol Alone (Reference)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=33)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=33)	Comparison of CYB003 + Alpha-Hydroxymetoprolol (Test) versus Alpha-Hydroxymetoprolol Alone (Reference)
C_{max} (ng/mL)						
n	32	32		32	32	
Geometric LS Mean	26.7	23.6		12.9	10.7	
Geometric Mean Ratio [a]			0.884			0.836
Two One-sided 90% CI Limits			(0.778, 1.01)			(0.764, 0.914)
Intra-Participant CV% [b]			31.0			21.3

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Midazolam

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=33)	CYB003 + Midazolam (Day 8) (N=33)	Comparison of CYB003 + Midazolam (Test) versus Midazolam Alone (Reference)	1-Hydroxy Midazolam Alone (Day 1) (N=33)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=33)	Comparison of CYB003 + 1-Hydroxy Midazolam (Test) versus 1-Hydroxy Midazolam Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	33	33		20	20	
Geometric LS Mean	24.2	28.4		9.81	10.3	
Geometric Mean Ratio [a]			1.17			1.05
Two One-sided 90% CI Limits			(1.10, 1.25)			(0.965, 1.13)
Intra-Participant CV% [b]			15.6			14.9
AUC_{0-t} (h*ng/mL)						
n	33	33		33	33	
Geometric LS Mean	23.1	27.1		9.04	9.69	
Geometric Mean Ratio [a]			1.17			1.07
Two One-sided 90% CI Limits			(1.10, 1.25)			(0.999, 1.15)
Intra-Participant CV% [b]			15.6			16.8

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.3
Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Midazolam

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=33)	CYB003 + Midazolam (Day 8) (N=33)	Comparison of CYB003 + Midazolam (Test) versus Midazolam Alone (Reference)	1-Hydroxy Midazolam Alone (Day 1) (N=33)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=33)	Comparison of CYB003 + 1-Hydroxy Midazolam (Test) versus 1-Hydroxy Midazolam Alone (Reference)
C_{max} (ng/mL)						
n	33	33		33	33	
Geometric LS Mean	7.21	8.41		2.98	3.32	
Geometric Mean Ratio [a]			1.17			1.12
Two One-sided 90% CI Limits			(1.06, 1.28)			(0.999, 1.25)
Intra-Participant CV% [b]			22.8			27.0

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020103.sas FINAL (2026-08-28,12:53)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Caffeine

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Comparison of CYB003 + Caffeine (Test) versus Caffeine Alone (Reference)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)	Comparison of CYB003 + Paraxanthine (Test) versus Paraxanthine Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	32	31		21	21	
Geometric LS Mean	17300	19500		11500	12100	
Geometric Mean Ratio [a]			1.13			1.05
Two One-sided 90% CI Limits			(1.05, 1.21)			(0.975, 1.13)
Intra-Participant CV% [b]			16.8			13.2
AUC_{0-t} (h*ng/mL)						
n	32	31		22	22	
Geometric LS Mean	16800	18900		10500	11500	
Geometric Mean Ratio [a]			1.12			1.09
Two One-sided 90% CI Limits			(1.04, 1.21)			(1.02, 1.18)
Intra-Participant CV% [b]			16.8			13.1

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Caffeine

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=35)	CYB003 + Caffeine (Day 8) (N=34)	Comparison of CYB003 + Caffeine (Test) versus Caffeine Alone (Reference)	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)	Comparison of CYB003 + Paraxanthine (Test) versus Paraxanthine Alone (Reference)
C_{max} (ng/mL)						
n	32	31		22	22	
Geometric LS Mean	1940	1860		567	597	
Geometric Mean Ratio [a]			0.958			1.05
Two One-sided 90% CI Limits			(0.875, 1.05)			(0.979, 1.13)
Intra-Participant CV% [b]			21.3			13.5

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Bupropion

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Comparison of CYB003 + Bupropion (Test) versus Bupropion Alone (Reference)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)	Comparison of CYB003 + Hydroxybupropion (Test) versus Hydroxybupropion Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	29	28		7	4	
Geometric LS Mean	141	165		2030	2400	
Geometric Mean Ratio [a]			1.17			1.18
Two One-sided 90% CI Limits			(1.08, 1.27)			(0.912, 1.53)
Intra-Participant CV% [b]			17.2			11.6
AUC_{0-t} (h*ng/mL)						
n	30	29		30	29	
Geometric LS Mean	128	151		1730	1760	
Geometric Mean Ratio [a]			1.17			1.01
Two One-sided 90% CI Limits			(1.08, 1.27)			(0.950, 1.08)
Intra-Participant CV% [b]			18.2			14.3

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Bupropion

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=35)	CYB003 + Bupropion (Day 8) (N=34)	Comparison of CYB003 + Bupropion (Test) versus Bupropion Alone (Reference)	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)	Comparison of CYB003 + Hydroxybupropion (Test) versus Hydroxybupropion Alone (Reference)
C_{max} (ng/mL)						
n	30	29		30	29	
Geometric LS Mean	24.2	23.6		68.4	68.3	
Geometric Mean Ratio [a]			0.976			0.998
Two One-sided 90% CI Limits			(0.856, 1.11)			(0.937, 1.06)
Intra-Participant CV% [b]			29.9			13.9

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Losartan

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Comparison of CYB003 + Losartan (Test) versus Losartan Alone (Reference)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)	Comparison of CYB003 + Losartan Carboxylic Acid (Test) versus Losartan Carboxylic Acid Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	30	32		31	29	
Geometric LS Mean	144	148		1020	1110	
Geometric Mean Ratio [a]			1.02			1.09
Two One-sided 90% CI Limits			(0.964, 1.09)			(1.03, 1.15)
Intra-Participant CV% [b]			13.3			12.1
AUC_{0-t} (h*ng/mL)						
n	34	33		32	30	
Geometric LS Mean	133	138		988	1070	
Geometric Mean Ratio [a]			1.04			1.08
Two One-sided 90% CI Limits			(0.971, 1.11)			(1.02, 1.15)
Intra-Participant CV% [b]			15.5			14.0

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Losartan

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=35)	CYB003 + Losartan (Day 8) (N=34)	Comparison of CYB003 + Losartan (Test) versus Losartan Alone (Reference)	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)	Comparison of CYB003 + Losartan Carboxylic Acid (Test) versus Losartan Carboxylic Acid Alone (Reference)
C_{max} (ng/mL)						
n	34	33		32	30	
Geometric LS Mean	34.1	29.2		101	105	
Geometric Mean Ratio [a]			0.858			1.04
Two One-sided 90% CI Limits			(0.735, 1.00)			(0.975, 1.10)
Intra-Participant CV% [b]			38.2			14.0

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Omeprazole

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	Comparison of CYB003 + Omeprazole (Test) versus Omeprazole Alone (Reference)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)	Comparison of CYB003 + 5-Hydroxy Omeprazole (Test) versus 5-Hydroxy Omeprazole Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	29	10		23	17	
Geometric LS Mean	544	656		363	378	
Geometric Mean Ratio [a]			1.21			1.04
Two One-sided 90% CI Limits			(1.05, 1.39)			(0.939, 1.15)
Intra-Participant CV% [b]			16.4			16.6
AUC_{0-t} (h*ng/mL)						
n	35	31		24	27	
Geometric LS Mean	458	405		355	305	
Geometric Mean Ratio [a]			0.886			0.858
Two One-sided 90% CI Limits			(0.733, 1.07)			(0.738, 0.997)
Intra-Participant CV% [b]			46.4			30.9

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Omeprazole

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=35)	CYB003 + Omeprazole (Day 8) (N=34)	Comparison of CYB003 + Omeprazole (Test) versus Omeprazole Alone (Reference)	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)	Comparison of CYB003 + 5-Hydroxy Omeprazole (Test) versus 5-Hydroxy Omeprazole Alone (Reference)
C_{max} (ng/mL)						
n	35	31		24	27	
Geometric LS Mean	225	152		141	87.1	
Geometric Mean Ratio [a]			0.677			0.619
Two One-sided 90% CI Limits			(0.515, 0.890)			(0.475, 0.807)
Intra-Participant CV% [b]			72.0			58.0

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Metoprolol

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Comparison of CYB003 + Metoprolol (Test) versus Metoprolol Alone (Reference)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)	Comparison of CYB003 + Alpha-Hydroxymetoprolol (Test) versus Alpha-Hydroxymetoprolol Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	34	32		30	28	
Geometric LS Mean	168	178		170	168	
Geometric Mean Ratio [a]			1.06			0.988
Two One-sided 90% CI Limits			(0.991, 1.13)			(0.951, 1.03)
Intra-Participant CV% [b]			15.1			8.1
AUC_{0-t} (h*ng/mL)						
n	34	33		34	33	
Geometric LS Mean	160	173		143	140	
Geometric Mean Ratio [a]			1.08			0.976
Two One-sided 90% CI Limits			(1.01, 1.15)			(0.929, 1.03)
Intra-Participant CV% [b]			16.1			11.7

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Metoprolol

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=35)	CYB003 + Metoprolol (Day 8) (N=34)	Comparison of CYB003 + Metoprolol (Test) versus Metoprolol Alone (Reference)	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)	Comparison of CYB003 + Alpha-Hydroxymetoprolol (Test) versus Alpha-Hydroxymetoprolol Alone (Reference)
C_{max} (ng/mL)						
n	34	33		34	33	
Geometric LS Mean	26.3	23.2		13.1	11.0	
Geometric Mean Ratio [a]			0.882			0.842
Two One-sided 90% CI Limits			(0.777, 1.00)			(0.770, 0.921)
Intra-Participant CV% [b]			30.8			21.4

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Midazolam

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	Comparison of CYB003 + Midazolam (Test) versus Midazolam Alone (Reference)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)	Comparison of CYB003 + 1-Hydroxy Midazolam (Test) versus 1-Hydroxy Midazolam Alone (Reference)
AUC_{0-inf} (h*ng/mL)						
n	35	33		31	22	
Geometric LS Mean	24.7	28.8		9.60	9.88	
Geometric Mean Ratio [a]			1.17			1.03
Two One-sided 90% CI Limits			(1.09, 1.25)			(0.950, 1.11)
Intra-Participant CV% [b]			15.6			14.9
AUC_{0-t} (h*ng/mL)						
n	35	33		35	33	
Geometric LS Mean	23.6	27.6		9.01	9.66	
Geometric Mean Ratio [a]			1.17			1.07
Two One-sided 90% CI Limits			(1.10, 1.25)			(1.00, 1.15)
Intra-Participant CV% [b]			15.5			16.7

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.4
Sensitivity Analysis of Comparison of Plasma CYP Probes (Parent and Metabolite) PK Parameters With and Without CYB003
PK Population

CYP Probe: Midazolam

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=35)	CYB003 + Midazolam (Day 8) (N=34)	Comparison of CYB003 + Midazolam (Test) versus Midazolam Alone (Reference)	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)	Comparison of CYB003 + 1-Hydroxy Midazolam (Test) versus 1-Hydroxy Midazolam Alone (Reference)
C_{max} (ng/mL)						
n	35	33		35	33	
Geometric LS Mean	7.36	8.52		2.96	3.31	
Geometric Mean Ratio [a]			1.16			1.12
Two One-sided 90% CI Limits			(1.05, 1.27)			(1.00, 1.25)
Intra-Participant CV% [b]			22.9			27.0

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone, CYB003 + <Metabolite>/<Metabolite> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020104.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Caffeine

PK Parameter (Units) Statistic	Paraxanthine Alone (Day 1) (N=33)	CYB003 + Paraxanthine (Day 8) (N=33)	Comparison of CYB003 + Paraxanthine (Test) versus Paraxanthine Alone (Reference)
MR_AUC_{0-inf}			
n	17	17	
Geometric LS Mean	0.767	0.712	
Geometric Mean Ratio [a]			0.928
Two One-sided 90% CI Limits			(0.885, 0.973)
Intra-Participant CV% [b]			8.0
MR_AUC_{0-t}			
n	19	19	
Geometric LS Mean	0.718	0.685	
Geometric Mean Ratio [a]			0.954
Two One-sided 90% CI Limits			(0.906, 1.00)
Intra-Participant CV% [b]			9.2

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Caffeine

PK Parameter (Units) Statistic	Paraxanthine Alone (Day 1) (N=33)	CYB003 + Paraxanthine (Day 8) (N=33)	Comparison of CYB003 + Paraxanthine (Test) versus Paraxanthine Alone (Reference)
MR_C_{max}			
n	19	19	
Geometric LS Mean	0.324	0.346	
Geometric Mean Ratio [a]			1.07
Two One-sided 90% CI Limits			(0.927, 1.23)
Intra-Participant CV% [b]			25.7

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Bupropion

PK Parameter (Units) Statistic	Hydroxybupropion Alone (Day 1) (N=33)	CYB003 + Hydroxybupropion (Day 8) (N=33)	Comparison of CYB003 + Hydroxybupropion (Test) versus Hydroxybupropion Alone (Reference)
MR_AUC_{0-inf}			
n	3	3	
Geometric LS Mean	13.0	12.7	
Geometric Mean Ratio [a]			0.980
Two One-sided 90% CI Limits			(0.583, 1.65)
Intra-Participant CV% [b]			22.1
MR_AUC_{0-t}			
n	28	28	
Geometric LS Mean	12.5	10.9	
Geometric Mean Ratio [a]			0.867
Two One-sided 90% CI Limits			(0.802, 0.938)
Intra-Participant CV% [b]			17.3

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Bupropion

PK Parameter (Units) Statistic	Hydroxybupropion Alone (Day 1) (N=33)	CYB003 + Hydroxybupropion (Day 8) (N=33)	Comparison of CYB003 + Hydroxybupropion (Test) versus Hydroxybupropion Alone (Reference)
MR_C_{max}			
n	28	28	
Geometric LS Mean	2.62	2.71	
Geometric Mean Ratio [a]			1.04
Two One-sided 90% CI Limits			(0.924, 1.16)
Intra-Participant CV% [b]			25.7

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Losartan

PK Parameter (Units) Statistic	Losartan Carboxylic Acid Alone (Day 1) (N=33)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=33)	Comparison of CYB003 + Losartan Carboxylic Acid (Test) versus Losartan Carboxylic Acid Alone (Reference)
MR_AUC_{0-inf}			
n	28	28	
Geometric LS Mean	7.52	7.89	
Geometric Mean Ratio [a]			1.05
Two One-sided 90% CI Limits			(0.989, 1.11)
Intra-Participant CV% [b]			12.7
MR_AUC_{0-t}			
n	30	30	
Geometric LS Mean	7.64	7.87	
Geometric Mean Ratio [a]			1.03
Two One-sided 90% CI Limits			(0.964, 1.10)
Intra-Participant CV% [b]			15.4

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Losartan

PK Parameter (Units) Statistic	Losartan Carboxylic Acid Alone (Day 1) (N=33)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=33)	Comparison of CYB003 + Losartan Carboxylic Acid (Test) versus Losartan Carboxylic Acid Alone (Reference)
MR_C_{max}			
n	30	30	
Geometric LS Mean	3.02	3.47	
Geometric Mean Ratio [a]			1.15
Two One-sided 90% CI Limits			(0.998, 1.32)
Intra-Participant CV% [b]			32.9

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Omeprazole

PK Parameter (Units) Statistic	5-Hydroxy Omeprazole Alone (Day 1) (N=33)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=33)	Comparison of CYB003 + 5-Hydroxy Omeprazole (Test) versus 5-Hydroxy Omeprazole Alone (Reference)
MR_AUC$_{0-inf}$			
n	6	6	
Geometric LS Mean	0.547	0.482	
Geometric Mean Ratio [a]			0.880
Two One-sided 90% CI Limits			(0.828, 0.936)
Intra-Participant CV% [b]			5.2
MR_AUC$_{0-t}$			
n	20	20	
Geometric LS Mean	0.885	0.821	
Geometric Mean Ratio [a]			0.928
Two One-sided 90% CI Limits			(0.851, 1.01)
Intra-Participant CV% [b]			15.8

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Omeprazole

PK Parameter (Units) Statistic	5-Hydroxy Omeprazole Alone (Day 1) (N=33)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=33)	Comparison of CYB003 + 5-Hydroxy Omeprazole (Test) versus 5-Hydroxy Omeprazole Alone (Reference)
MR_C_{max}			
n	20	20	
Geometric LS Mean	0.674	0.629	
Geometric Mean Ratio [a]			0.934
Two One-sided 90% CI Limits			(0.852, 1.02)
Intra-Participant CV% [b]			16.9

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Metoprolol

PK Parameter (Units) Statistic	Alpha-Hydroxymetoprolol Alone (Day 1) (N=33)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=33)	Comparison of CYB003 + Alpha-Hydroxymetoprolol (Test) versus Alpha-Hydroxymetoprolol Alone (Reference)
MR_AUC$_{0-inf}$			
n	24	24	
Geometric LS Mean	1.07	1.01	
Geometric Mean Ratio [a]			0.942
Two One-sided 90% CI Limits			(0.886, 1.00)
Intra-Participant CV% [b]			12.4
MR_AUC$_{0-t}$			
n	32	32	
Geometric LS Mean	0.811	0.731	
Geometric Mean Ratio [a]			0.902
Two One-sided 90% CI Limits			(0.842, 0.965)
Intra-Participant CV% [b]			16.1

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Metoprolol

PK Parameter (Units) Statistic	Alpha-Hydroxymetoprolol Alone (Day 1) (N=33)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=33)	Comparison of CYB003 + Alpha-Hydroxymetoprolol (Test) versus Alpha-Hydroxymetoprolol Alone (Reference)
MR_C_{max}			
n	32	32	
Geometric LS Mean	0.454	0.429	
Geometric Mean Ratio [a]			0.945
Two One-sided 90% CI Limits			(0.876, 1.02)
Intra-Participant CV% [b]			18.1

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Midazolam

PK Parameter (Units) Statistic	1-Hydroxy Midazolam Alone (Day 1) (N=33)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=33)	Comparison of CYB003 + 1-Hydroxy Midazolam (Test) versus 1-Hydroxy Midazolam Alone (Reference)
MR_AUC_{0-inf}			
n	20	20	
Geometric LS Mean	0.377	0.337	
Geometric Mean Ratio [a]			0.893
Two One-sided 90% CI Limits			(0.816, 0.978)
Intra-Participant CV% [b]			16.7
MR_AUC_{0-t}			
n	33	33	
Geometric LS Mean	0.372	0.340	
Geometric Mean Ratio [a]			0.914
Two One-sided 90% CI Limits			(0.861, 0.971)
Intra-Participant CV% [b]			14.6

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.5
Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
DDI PK Population

CYP Probe: Midazolam

PK Parameter (Units) Statistic	1-Hydroxy Midazolam Alone (Day 1) (N=33)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=33)	Comparison of CYB003 + 1-Hydroxy Midazolam (Test) versus 1-Hydroxy Midazolam Alone (Reference)
MR_C_{max}			
n	33	33	
Geometric LS Mean	0.394	0.377	
Geometric Mean Ratio [a]			0.957
Two One-sided 90% CI Limits			(0.892, 1.03)
Intra-Participant CV% [b]			17.1

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020105.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Caffeine

PK Parameter (Units) Statistic	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)	Comparison of CYB003 + Paraxanthine (Test) versus Paraxanthine Alone (Reference)
MR_AUC_{0-inf}			
n	21	21	
Geometric LS Mean	0.775	0.721	
Geometric Mean Ratio [a]			0.930
Two One-sided 90% CI Limits			(0.889, 0.973)
Intra-Participant CV% [b]			7.8
MR_AUC_{0-t}			
n	22	22	
Geometric LS Mean	0.730	0.697	
Geometric Mean Ratio [a]			0.955
Two One-sided 90% CI Limits			(0.909, 1.00)
Intra-Participant CV% [b]			9.1

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Caffeine

PK Parameter (Units) Statistic	Paraxanthine Alone (Day 1) (N=35)	CYB003 + Paraxanthine (Day 8) (N=34)	Comparison of CYB003 + Paraxanthine (Test) versus Paraxanthine Alone (Reference)
MR_C_{max}			
n	22	22	
Geometric LS Mean	0.324	0.343	
Geometric Mean Ratio [a]			1.06
Two One-sided 90% CI Limits			(0.931, 1.20)
Intra-Participant CV% [b]			24.6

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Bupropion

PK Parameter (Units) Statistic	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)	Comparison of CYB003 + Hydroxybupropion (Test) versus Hydroxybupropion Alone (Reference)
MR_AUC_{0-inf}			
n	7	4	
Geometric LS Mean	16.1	15.3	
Geometric Mean Ratio [a]			0.950
Two One-sided 90% CI Limits			(0.591, 1.53)
Intra-Participant CV% [b]			21.9
MR_AUC_{0-t}			
n	30	29	
Geometric LS Mean	12.7	10.9	
Geometric Mean Ratio [a]			0.864
Two One-sided 90% CI Limits			(0.799, 0.935)
Intra-Participant CV% [b]			17.5

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Bupropion

PK Parameter (Units) Statistic	Hydroxybupropion Alone (Day 1) (N=35)	CYB003 + Hydroxybupropion (Day 8) (N=34)	Comparison of CYB003 + Hydroxybupropion (Test) versus Hydroxybupropion Alone (Reference)
MR_C_{max}			
n	30	29	
Geometric LS Mean	2.66	2.73	
Geometric Mean Ratio [a]			1.03
Two One-sided 90% CI Limits			(0.912, 1.15)
Intra-Participant CV% [b]			26.4

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Losartan

PK Parameter (Units) Statistic	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)	Comparison of CYB003 + Losartan Carboxylic Acid (Test) versus Losartan Carboxylic Acid Alone (Reference)
MR_AUC_{0-inf}			
n	30	29	
Geometric LS Mean	7.28	7.61	
Geometric Mean Ratio [a]			1.05
Two One-sided 90% CI Limits			(0.987, 1.11)
Intra-Participant CV% [b]			12.7
MR_AUC_{0-t}			
n	32	30	
Geometric LS Mean	7.58	7.82	
Geometric Mean Ratio [a]			1.03
Two One-sided 90% CI Limits			(0.965, 1.10)
Intra-Participant CV% [b]			15.3

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Losartan

PK Parameter (Units) Statistic	Losartan Carboxylic Acid Alone (Day 1) (N=35)	CYB003 + Losartan Carboxylic Acid (Day 8) (N=34)	Comparison of CYB003 + Losartan Carboxylic Acid (Test) versus Losartan Carboxylic Acid Alone (Reference)
MR_C_{max}			
n	32	30	
Geometric LS Mean	2.97	3.43	
Geometric Mean Ratio [a]			1.16
Two One-sided 90% CI Limits			(1.01, 1.33)
Intra-Participant CV% [b]			32.7

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Omeprazole

PK Parameter (Units) Statistic	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)	Comparison of CYB003 + 5-Hydroxy Omeprazole (Test) versus 5-Hydroxy Omeprazole Alone (Reference)
MR_AUC_{0-inf}			
n	20	10	
Geometric LS Mean	0.774	0.678	
Geometric Mean Ratio [a]			0.876
Two One-sided 90% CI Limits			(0.825, 0.931)
Intra-Participant CV% [b]			5.2
MR_AUC_{0-t}			
n	24	26	
Geometric LS Mean	0.809	0.745	
Geometric Mean Ratio [a]			0.921
Two One-sided 90% CI Limits			(0.845, 1.00)
Intra-Participant CV% [b]			15.9

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Omeprazole

PK Parameter (Units) Statistic	5-Hydroxy Omeprazole Alone (Day 1) (N=35)	CYB003 + 5-Hydroxy Omeprazole (Day 8) (N=34)	Comparison of CYB003 + 5-Hydroxy Omeprazole (Test) versus 5-Hydroxy Omeprazole Alone (Reference)
MR_C_{max}			
n	24	26	
Geometric LS Mean	0.609	0.564	
Geometric Mean Ratio [a]			0.926
Two One-sided 90% CI Limits			(0.844, 1.01)
Intra-Participant CV% [b]			17.0

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Metoprolol

PK Parameter (Units) Statistic	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)	Comparison of CYB003 + Alpha-Hydroxymetoprolol (Test) versus Alpha-Hydroxymetoprolol Alone (Reference)
MR_AUC_{0-inf}			
n	30	27	
Geometric LS Mean	1.03	0.975	
Geometric Mean Ratio [a]			0.945
Two One-sided 90% CI Limits			(0.889, 1.00)
Intra-Participant CV% [b]			12.4
MR_AUC_{0-t}			
n	34	33	
Geometric LS Mean	0.844	0.762	
Geometric Mean Ratio [a]			0.903
Two One-sided 90% CI Limits			(0.844, 0.966)
Intra-Participant CV% [b]			16.1

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Metoprolol

PK Parameter (Units) Statistic	Alpha-Hydroxymetoprolol Alone (Day 1) (N=35)	CYB003 + Alpha-Hydroxymetoprolol (Day 8) (N=34)	Comparison of CYB003 + Alpha-Hydroxymetoprolol (Test) versus Alpha-Hydroxymetoprolol Alone (Reference)
MR_C_{max}			
n	34	33	
Geometric LS Mean	0.474	0.449	
Geometric Mean Ratio [a]			0.947
Two One-sided 90% CI Limits			(0.878, 1.02)
Intra-Participant CV% [b]			18.1

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Midazolam

PK Parameter (Units) Statistic	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)	Comparison of CYB003 + 1-Hydroxy Midazolam (Test) versus 1-Hydroxy Midazolam Alone (Reference)
MR_AUC_{0-inf}			
n	31	22	
Geometric LS Mean	0.372	0.331	
Geometric Mean Ratio [a]			0.887
Two One-sided 90% CI Limits			(0.812, 0.970)
Intra-Participant CV% [b]			16.6
MR_AUC_{0-t}			
n	35	33	
Geometric LS Mean	0.364	0.333	
Geometric Mean Ratio [a]			0.916
Two One-sided 90% CI Limits			(0.863, 0.974)
Intra-Participant CV% [b]			14.6

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.6
Sensitivity Analysis of Comparison of Plasma CYP Probes Metabolite Ratios With and Without CYB003
PK Population

CYP Probe: Midazolam

PK Parameter (Units) Statistic	1-Hydroxy Midazolam Alone (Day 1) (N=35)	CYB003 + 1-Hydroxy Midazolam (Day 8) (N=34)	Comparison of CYB003 + 1-Hydroxy Midazolam (Test) versus 1-Hydroxy Midazolam Alone (Reference)
MR_C_{max}			
n	35	33	
Geometric LS Mean	0.383	0.368	
Geometric Mean Ratio [a]			0.961
Two One-sided 90% CI Limits			(0.896, 1.03)
Intra-Participant CV% [b]			17.1

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; LS = least squares; CI = confidence interval; MR = metabolite ratio.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020106.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Caffeine

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=33)	CYB003 + Caffeine (Day 8) (N=33)	Comparison of CYB003 + Caffeine (Test) versus Caffeine Alone (Reference)
AUC_{0-inf} (h*ng/mL)			
n	29	29	
Geometric LS Mean	17200	19500	
Geometric Mean Ratio [a]			1.13
Two One-sided 90% CI Limits			(1.05, 1.22)
Intra-Participant CV% [b]			16.8
AUC_{0-t} (h*ng/mL)			
n	29	29	
Geometric LS Mean	16700	18800	
Geometric Mean Ratio [a]			1.13
Two One-sided 90% CI Limits			(1.05, 1.22)
Intra-Participant CV% [b]			16.9

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Caffeine

PK Parameter (Units) Statistic	Caffeine Alone (Day 1) (N=33)	CYB003 + Caffeine (Day 8) (N=33)	Comparison of CYB003 + Caffeine (Test) versus Caffeine Alone (Reference)
C_{max} (ng/mL)			
n	29	29	
Geometric LS Mean	1950	1870	
Geometric Mean Ratio [a]			0.958
Two One-sided 90% CI Limits			(0.871, 1.06)
Intra-Participant CV% [b]			21.8

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Bupropion

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=33)	CYB003 + Bupropion (Day 8) (N=33)	Comparison of CYB003 + Bupropion (Test) versus Bupropion Alone (Reference)
AUC_{0-inf} (h*ng/mL)			
n	26	26	
Geometric LS Mean	145	169	
Geometric Mean Ratio [a]			1.17
Two One-sided 90% CI Limits			(1.08, 1.27)
Intra-Participant CV% [b]			17.0
AUC_{0-t} (h*ng/mL)			
n	28	28	
Geometric LS Mean	130	152	
Geometric Mean Ratio [a]			1.17
Two One-sided 90% CI Limits			(1.08, 1.27)
Intra-Participant CV% [b]			18.0

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Bupropion

PK Parameter (Units) Statistic	Bupropion Alone (Day 1) (N=33)	CYB003 + Bupropion (Day 8) (N=33)	Comparison of CYB003 + Bupropion (Test) versus Bupropion Alone (Reference)
C_{max} (ng/mL)			
n	28	28	
Geometric LS Mean	24.5	23.7	
Geometric Mean Ratio [a]			0.966
Two One-sided 90% CI Limits			(0.852, 1.10)
Intra-Participant CV% [b]			28.2

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Losartan

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=33)	CYB003 + Losartan (Day 8) (N=33)	Comparison of CYB003 + Losartan (Test) versus Losartan Alone (Reference)
AUC_{0-inf} (h*ng/mL)			
n	28	28	
Geometric LS Mean	143	147	
Geometric Mean Ratio [a]			1.03
Two One-sided 90% CI Limits			(0.966, 1.09)
Intra-Participant CV% [b]			13.3
AUC_{0-t} (h*ng/mL)			
n	32	32	
Geometric LS Mean	130	135	
Geometric Mean Ratio [a]			1.04
Two One-sided 90% CI Limits			(0.970, 1.11)
Intra-Participant CV% [b]			15.5

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Losartan

PK Parameter (Units) Statistic	Losartan Alone (Day 1) (N=33)	CYB003 + Losartan (Day 8) (N=33)	Comparison of CYB003 + Losartan (Test) versus Losartan Alone (Reference)
C_{max} (ng/mL)			
n	32	32	
Geometric LS Mean	33.2	28.6	
Geometric Mean Ratio [a]			0.862
Two One-sided 90% CI Limits			(0.737, 1.01)
Intra-Participant CV% [b]			38.4

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Omeprazole

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=33)	CYB003 + Omeprazole (Day 8) (N=33)	Comparison of CYB003 + Omeprazole (Test) versus Omeprazole Alone (Reference)
AUC_{0-inf} (h*ng/mL)			
n	9	9	
Geometric LS Mean	781	931	
Geometric Mean Ratio [a]			1.19
Two One-sided 90% CI Limits			(1.03, 1.37)
Intra-Participant CV% [b]			16.3
AUC_{0-t} (h*ng/mL)			
n	31	31	
Geometric LS Mean	475	416	
Geometric Mean Ratio [a]			0.876
Two One-sided 90% CI Limits			(0.723, 1.06)
Intra-Participant CV% [b]			46.7

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Omeprazole

PK Parameter (Units) Statistic	Omeprazole Alone (Day 1) (N=33)	CYB003 + Omeprazole (Day 8) (N=33)	Comparison of CYB003 + Omeprazole (Test) versus Omeprazole Alone (Reference)
C_{max} (ng/mL)			
n	31	31	
Geometric LS Mean	234	154	
Geometric Mean Ratio [a]			0.659
Two One-sided 90% CI Limits			(0.497, 0.873)
Intra-Participant CV% [b]			72.9

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Metoprolol

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=33)	CYB003 + Metoprolol (Day 8) (N=33)	Comparison of CYB003 + Metoprolol (Test) versus Metoprolol Alone (Reference)
AUC_{0-inf} (h*ng/mL)			
n	31	31	
Geometric LS Mean	170	180	
Geometric Mean Ratio [a]			1.06
Two One-sided 90% CI Limits			(0.992, 1.13)
Intra-Participant CV% [b]			15.1
AUC_{0-t} (h*ng/mL)			
n	32	32	
Geometric LS Mean	163	176	
Geometric Mean Ratio [a]			1.08
Two One-sided 90% CI Limits			(1.01, 1.16)
Intra-Participant CV% [b]			16.1

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Metoprolol

PK Parameter (Units) Statistic	Metoprolol Alone (Day 1) (N=33)	CYB003 + Metoprolol (Day 8) (N=33)	Comparison of CYB003 + Metoprolol (Test) versus Metoprolol Alone (Reference)
C_{max} (ng/mL)			
n	32	32	
Geometric LS Mean	26.7	23.6	
Geometric Mean Ratio [a]			0.884
Two One-sided 90% CI Limits			(0.778, 1.01)
Intra-Participant CV% [b]			31.0

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Midazolam

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=33)	CYB003 + Midazolam (Day 8) (N=33)	Comparison of CYB003 + Midazolam (Test) versus Midazolam Alone (Reference)
AUC_{0-inf} (h*ng/mL)			
n	33	33	
Geometric LS Mean	24.2	28.4	
Geometric Mean Ratio [a]			1.17
Two One-sided 90% CI Limits			(1.10, 1.25)
Intra-Participant CV% [b]			15.6
AUC_{0-t} (h*ng/mL)			
n	33	33	
Geometric LS Mean	23.1	27.1	
Geometric Mean Ratio [a]			1.17
Two One-sided 90% CI Limits			(1.10, 1.25)
Intra-Participant CV% [b]			15.6

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap} > 20\%$, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.7
Comparison of Plasma CYP Probes (Parent Only) PK Parameters With and Without CYB003
DDI PK Population

CYP Probe: Midazolam

PK Parameter (Units) Statistic	Midazolam Alone (Day 1) (N=33)	CYB003 + Midazolam (Day 8) (N=33)	Comparison of CYB003 + Midazolam (Test) versus Midazolam Alone (Reference)
C_{max} (ng/mL)			
n	33	33	
Geometric LS Mean	7.21	8.41	
Geometric Mean Ratio [a]			1.17
Two One-sided 90% CI Limits			(1.06, 1.28)
Intra-Participant CV% [b]			22.8

a. Ratio expressed as percentage of CYB003 + <Parent>/<Parent> Alone.
b. Calculated as 100 * sqrt[exp(MSE) − 1] where MSE is the modelled mean square error.
All values are estimated from a mixed effects regression model on natural log (ln) transformed values of the parameter, including treatment as fixed effect and participant as random effect. Participants must have evaluable parents in both periods to be included in the model. AUC_{0-inf} was excluded from the analysis if $\%AUC_{extrap}$ > 20%, or if the associated adjusted r^2 value was < 0.8.
n is the number of participants with the measurement available.
N is the number of participants in the respective treatment.
CV = coefficient of variation; CI = confidence interval; LS = least squares.

Source: Listing 16.2.6.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020107.sas FINAL (2026-08-28,12:54)

Table 14.2.1.8
Summary of Percent Total Drug-Related Exposures for CYB003 Metabolites
PK Population

PK Parameter (Unit) Statistic	Psilocin-O-glucuronide-d10 (N=35)	4-HIAA-d2 (N=35)
%TRE		
n	33	33
Mean (SD)	75.8 (5.58)	6.01 (1.57)
CV%	7.4	26.0
Geometric Mean (CV%)	75.6 (7.5)	5.81 (27.3)
Median	76.1	5.92
Min, Max	63.8, 86.0	3.36, 9.65

n is the number of participants with the measurement available.
N is the number of participants in the respective analyte.
CV = coefficient of variation; SD = standard deviation; 4-HIAA-d2 = 4-hydroxindole-3-acetic acid-d2; %TRE = percent total drug-related exposure.

Source: Listing 16.2.6.5 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14020108.sas FINAL (2026-08-28,12:54)

Table 14.3.1.1
Treatment-Emergent Adverse Event Overview
Safety Population

Number (%) of participants reporting	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Any TEAE	4 (11.1) [4]	34 (100) [196]	34 (94.4) [200]
Any non-SMRE TEAE	4 (11.1) [4]	14 (41.2) [24]	16 (44.4) [28]
Any SMRE TEAE	0	34 (100) [172]	34 (94.4) [172]
Any TEAE by Greatest Intensity Level [a]	4 (11.1) [4]	34 (100) [196]	34 (94.4) [200]
Severe	0	0	0
Moderate	0	13 (38.2) [36]	13 (36.1) [36]
Mild	4 (11.1) [4]	21 (61.8) [88]	21 (58.3) [90]
Any TEAE by Intensity Level [b]	4 (11.1) [4]	34 (100) [196]	34 (94.4) [200]
Severe	0	0	0
Moderate	0	13 (38.2) [36]	13 (36.1) [36]
Mild	4 (11.1) [4]	34 (100) [160]	34 (94.4) [164]
Any TEAE with an onset date within 48 hours from CYB003 administration on Day 8		34 (100) [195]	34 (94.4) [195]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event category for a participant were counted once in the frequency for the adverse event category.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related TEAEs are defined as events reported as related or possibly related. TEAEs include all SMREs and potential abuse-related events.
a. Maximum intensity for a participant is included along with the number of events with that intensity for the participant.
b. The number of participants and the number of events are counted within each intensity level, regardless of whether that intensity represents the participant's maximum intensity.
AE = adverse event; TEAE = treatment emergent adverse event; SMRE = session monitor-reported event; E = number of events.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030101.sas FINAL (2026-08-28,12:54)

Table 14.3.1.1
Treatment-Emergent Adverse Event Overview
Safety Population

Number (%) of participants reporting	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Any related TEAE	3 (8.3) [3]	34 (100) [194]	34 (94.4) [197]
Any non-SMRE related TEAE	3 (8.3) [3]	12 (35.3) [22]	13 (36.1) [25]
Any SMRE related TEAE	0	34 (100) [172]	34 (94.4) [172]
Any serious TEAE	0	0	0
Death	0	0	0
Any related serious TEAE	0	0	0
Any TEAE leading to study discontinuation	0	0	0
Any potential abuse-related AE	0	34 (100) [93]	34 (94.4) [93]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event category for a participant were counted once in the frequency for the adverse event category.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related TEAEs are defined as events reported as related or possibly related. TEAEs include all SMREs and potential abuse-related events.
a. Maximum intensity for a participant is included along with the number of events with that intensity for the participant.
b. The number of participants and the number of events are counted within each intensity level, regardless of whether that intensity represents the participant's maximum intensity.
AE = adverse event; TEAE = treatment emergent adverse event; SMRE = session monitor-reported event; E = number of events.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030101.sas FINAL (2026-08-28,12:54)

Table 14.3.1.2
Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any TEAE	4 (11.1) [4]	34 (100) [196]	34 (94.4) [200]
Ear and labyrinth disorders	0	2 (5.9) [2]	2 (5.6) [2]
Hyperacusis	0	2 (5.9) [2]	2 (5.6) [2]
Gastrointestinal disorders	0	7 (20.6) [7]	7 (19.4) [7]
Nausea	0	7 (20.6) [7]	7 (19.4) [7]
General disorders and administration site conditions	1 (2.8) [1]	12 (35.3) [15]	13 (36.1) [16]
Feeling of body temperature change	0	3 (8.8) [3]	3 (8.3) [3]
Feeling of relaxation	0	3 (8.8) [3]	3 (8.3) [3]
Feeling cold	0	2 (5.9) [2]	2 (5.6) [2]
Feeling hot	0	2 (5.9) [2]	2 (5.6) [2]
Chest discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Gait disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Malaise	0	1 (2.9) [1]	1 (2.8) [1]
Pain	0	1 (2.9) [1]	1 (2.8) [1]
Vessel puncture site pain	1 (2.8) [1]	0	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse even.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030102.sas FINAL (2026-08-28,12:54)

Table 14.3.1.2
Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Investigations	0	2 (5.9) [2]	2 (5.6) [2]
Blood pressure increased	0	2 (5.9) [2]	2 (5.6) [2]
Metabolism and nutrition disorders	0	1 (2.9) [1]	1 (2.8) [1]
Decreased appetite	0	1 (2.9) [1]	1 (2.8) [1]
Nervous system disorders	0	18 (52.9) [33]	18 (50.0) [33]
Dizziness	0	9 (26.5) [9]	9 (25.0) [9]
Hyperaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Synaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Balance disorder	0	3 (8.8) [3]	3 (8.3) [3]
Coordination abnormal	0	2 (5.9) [2]	2 (5.6) [2]
Dysaesthesia	0	2 (5.9) [2]	2 (5.6) [2]
Headache	0	2 (5.9) [2]	2 (5.6) [2]
Cognitive disorder	0	1 (2.9) [1]	1 (2.8) [1]
Dyskinesia	0	1 (2.9) [1]	1 (2.8) [1]
Parosmia	0	1 (2.9) [1]	1 (2.8) [1]
Psychomotor hyperactivity	0	1 (2.9) [1]	1 (2.8) [1]
Sensory disturbance	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse even.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030102.sas FINAL (2026-08-28,12:54)

Table 14.3.1.2
Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Nervous system disorders (Cont.)			
Tremor	0	1 (2.9) [1]	1 (2.8) [1]
Visual processing disorder	0	1 (2.9) [1]	1 (2.8) [1]
Psychiatric disorders	0	33 (97.1) [126]	33 (91.7) [126]
Hallucination, visual	0	24 (70.6) [24]	24 (66.7) [24]
Affect lability	0	13 (38.2) [15]	13 (36.1) [15]
Euphoric mood	0	10 (29.4) [10]	10 (27.8) [10]
Illusion	0	8 (23.5) [9]	8 (22.2) [9]
Hallucination	0	6 (17.6) [6]	6 (16.7) [6]
Agitation	0	5 (14.7) [6]	5 (13.9) [6]
Paranoia	0	5 (14.7) [5]	5 (13.9) [5]
Time perception altered	0	5 (14.7) [5]	5 (13.9) [5]
Derealisation	0	4 (11.8) [4]	4 (11.1) [4]
Dissociation	0	4 (11.8) [6]	4 (11.1) [6]
Somatic hallucination	0	4 (11.8) [4]	4 (11.1) [4]
Disorientation	0	3 (8.8) [4]	3 (8.3) [4]
Restlessness	0	3 (8.8) [3]	3 (8.3) [3]
Depersonalisation/derealisation disorder	0	2 (5.9) [2]	2 (5.6) [2]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse even.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030102.sas FINAL (2026-08-28,12:54)

Table 14.3.1.2
Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders (Cont.)			
Inappropriate affect	0	2 (5.9) [2]	2 (5.6) [2]
Psychomotor retardation	0	2 (5.9) [2]	2 (5.6) [2]
Acute stress disorder	0	1 (2.9) [1]	1 (2.8) [1]
Anxiety	0	1 (2.9) [1]	1 (2.8) [1]
Confusional state	0	1 (2.9) [1]	1 (2.8) [1]
Delusion	0	1 (2.9) [1]	1 (2.8) [1]
Depressed mood	0	1 (2.9) [1]	1 (2.8) [1]
Disinhibition	0	1 (2.9) [1]	1 (2.8) [1]
Fear	0	1 (2.9) [1]	1 (2.8) [1]
Fear of death	0	1 (2.9) [1]	1 (2.8) [1]
Flight of ideas	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, auditory	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, tactile	0	1 (2.9) [2]	1 (2.8) [2]
Logorrhoea	0	1 (2.9) [1]	1 (2.8) [1]
Loose associations	0	1 (2.9) [1]	1 (2.8) [1]
Nervousness	0	1 (2.9) [1]	1 (2.8) [1]
Perseveration	0	1 (2.9) [1]	1 (2.8) [1]
Sexually inappropriate behaviour	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse even.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030102.sas FINAL (2026-08-28,12:54)

Table 14.3.1.2
Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders (Cont.)			
Somatic delusion	0	1 (2.9) [1]	1 (2.8) [1]
Thinking abnormal	0	1 (2.9) [1]	1 (2.8) [1]
Renal and urinary disorders	0	1 (2.9) [1]	1 (2.8) [1]
Urinary incontinence	0	1 (2.9) [1]	1 (2.8) [1]
Respiratory, thoracic and mediastinal disorders	0	2 (5.9) [2]	2 (5.6) [2]
Dyspnoea	0	1 (2.9) [1]	1 (2.8) [1]
Hyperventilation	0	1 (2.9) [1]	1 (2.8) [1]
Skin and subcutaneous tissue disorders	0	3 (8.8) [3]	3 (8.3) [3]
Hyperhidrosis	0	2 (5.9) [2]	2 (5.6) [2]
Ingrown hair	0	1 (2.9) [1]	1 (2.8) [1]
Surgical and medical procedures	0	1 (2.9) [1]	1 (2.8) [1]
Analgesic therapy	0	1 (2.9) [1]	1 (2.8) [1]
Vascular disorders	3 (8.3) [3]	3 (8.8) [3]	5 (13.9) [6]
Orthostatic hypotension	2 (5.6) [2]	3 (8.8) [3]	4 (11.1) [5]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse even.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030102.sas FINAL (2026-08-28,12:54)

Table 14.3.1.2
Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Vascular disorders (Cont.)			
Hypotension	1 (2.8) [1]	0	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse even.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030102.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any TEAE	4 (11.1) [4]	34 (100) [196]	34 (94.4) [200]
Severe	0	0	0
Moderate	0	13 (38.2) [36]	13 (36.1) [36]
Mild	4 (11.1) [4]	21 (61.8) [88]	21 (58.3) [90]
Ear and labyrinth disorders	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Hyperacusis	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Gastrointestinal disorders	0	7 (20.6) [7]	7 (19.4) [7]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	6 (17.6) [6]	6 (16.7) [6]
Nausea	0	7 (20.6) [7]	7 (19.4) [7]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	6 (17.6) [6]	6 (16.7) [6]
General disorders and administration site conditions	1 (2.8) [1]	12 (35.3) [15]	13 (36.1) [16]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	1 (2.8) [1]	11 (32.4) [14]	12 (33.3) [15]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Feeling of body temperature change	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	0	0
Mild	0	3 (8.8) [3]	3 (8.3) [3]
Feeling of relaxation	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	0	0
Mild	0	3 (8.8) [3]	3 (8.3) [3]
Feeling cold	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Feeling hot	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Chest discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Gait disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Malaise	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Pain	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Vessel puncture site pain	1 (2.8) [1]	0	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	1 (2.8) [1]	0	1 (2.8) [1]
Investigations	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Blood pressure increased	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Metabolism and nutrition disorders	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Decreased appetite	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Nervous system disorders	0	18 (52.9) [33]	18 (50.0) [33]
Severe	0	0	0
Moderate	0	3 (8.8) [4]	3 (8.3) [4]
Mild	0	15 (44.1) [26]	15 (41.7) [26]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Dizziness	0	9 (26.5) [9]	9 (25.0) [9]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	8 (23.5) [8]	8 (22.2) [8]
Hyperaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Severe	0	0	0
Moderate	0	0	0
Mild	0	4 (11.8) [4]	4 (11.1) [4]
Synaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Severe	0	0	0
Moderate	0	0	0
Mild	0	4 (11.8) [4]	4 (11.1) [4]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Balance disorder	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	0	0
Mild	0	3 (8.8) [3]	3 (8.3) [3]
Coordination abnormal	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Dysaesthesia	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Headache	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Cognitive disorder	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Dyskinesia	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Parosmia	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Psychomotor hyperactivity	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Sensory disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Tremor	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Visual processing disorder	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Psychiatric disorders	0	33 (97.1) [126]	33 (91.7) [126]
Severe	0	0	0
Moderate	0	7 (20.6) [26]	7 (19.4) [26]
Mild	0	26 (76.5) [81]	26 (72.2) [81]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Hallucination, visual	0	24 (70.6) [24]	24 (66.7) [24]
Severe	0	0	0
Moderate	0	2 (5.9) [2]	2 (5.6) [2]
Mild	0	22 (64.7) [22]	22 (61.1) [22]
Affect lability	0	13 (38.2) [15]	13 (36.1) [15]
Severe	0	0	0
Moderate	0	2 (5.9) [3]	2 (5.6) [3]
Mild	0	11 (32.4) [12]	11 (30.6) [12]
Euphoric mood	0	10 (29.4) [10]	10 (27.8) [10]
Severe	0	0	0
Moderate	0	0	0
Mild	0	10 (29.4) [10]	10 (27.8) [10]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Illusion	0	8 (23.5) [9]	8 (22.2) [9]
Severe	0	0	0
Moderate	0	2 (5.9) [2]	2 (5.6) [2]
Mild	0	6 (17.6) [7]	6 (16.7) [7]
Hallucination	0	6 (17.6) [6]	6 (16.7) [6]
Severe	0	0	0
Moderate	0	0	0
Mild	0	6 (17.6) [6]	6 (16.7) [6]
Agitation	0	5 (14.7) [6]	5 (13.9) [6]
Severe	0	0	0
Moderate	0	3 (8.8) [4]	3 (8.3) [4]
Mild	0	2 (5.9) [2]	2 (5.6) [2]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Paranoia	0	5 (14.7) [5]	5 (13.9) [5]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	4 (11.8) [4]	4 (11.1) [4]
Time perception altered	0	5 (14.7) [5]	5 (13.9) [5]
Severe	0	0	0
Moderate	0	0	0
Mild	0	5 (14.7) [5]	5 (13.9) [5]
Derealisation	0	4 (11.8) [4]	4 (11.1) [4]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	3 (8.8) [3]	3 (8.3) [3]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Dissociation	0	4 (11.8) [6]	4 (11.1) [6]
Severe	0	0	0
Moderate	0	2 (5.9) [2]	2 (5.6) [2]
Mild	0	2 (5.9) [4]	2 (5.6) [4]
Somatic hallucination	0	4 (11.8) [4]	4 (11.1) [4]
Severe	0	0	0
Moderate	0	0	0
Mild	0	4 (11.8) [4]	4 (11.1) [4]
Disorientation	0	3 (8.8) [4]	3 (8.3) [4]
Severe	0	0	0
Moderate	0	3 (8.8) [4]	3 (8.3) [4]
Mild	0	0	0

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Restlessness	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	0	0
Mild	0	3 (8.8) [3]	3 (8.3) [3]
Depersonalisation/derealisation disorder	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Inappropriate affect	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychomotor retardation	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Acute stress disorder	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Anxiety	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Confusional state	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Delusion	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Depressed mood	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Disinhibition	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Fear	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Fear of death	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Flight of ideas	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Hallucination, auditory	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, tactile	0	1 (2.9) [2]	1 (2.8) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [2]	1 (2.8) [2]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Logorrhoea	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Loose associations	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Nervousness	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Perseveration	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Sexually inappropriate behaviour	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Somatic delusion	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Thinking abnormal	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Renal and urinary disorders	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Urinary incontinence	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Respiratory, thoracic and mediastinal disorders	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Dyspnoea	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Hyperventilation	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Skin and subcutaneous tissue disorders	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Hyperhidrosis	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Ingrown hair	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Surgical and medical procedures	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Analgesic therapy	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Vascular disorders	3 (8.3) [3]	3 (8.8) [3]	5 (13.9) [6]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	3 (8.3) [3]	2 (5.9) [2]	4 (11.1) [5]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.3
Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Orthostatic hypotension	2 (5.6) [2]	3 (8.8) [3]	4 (11.1) [5]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	2 (5.6) [2]	2 (5.9) [2]	3 (8.3) [4]
Hypotension	1 (2.8) [1]	0	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	1 (2.8) [1]	0	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030103.sas FINAL (2026-08-28,12:54)

Table 14.3.1.4

Treatment-Emergent Adverse Events with an Onset Date within 48 Hours from CYB003 Administration on Day 8 by MedDRA System Organ Class and Preferred Term

Safety Population

System Organ Class/ Preferred Term	CYB003 + CYP Probe N=34 n (%) [E]
Participants with any TEAE with an onset date within 48 hours from CYB003 administration on Day 8	34 (100) [195]
Ear and labyrinth disorders	2 (5.9) [2]
Hyperacusis	2 (5.9) [2]
Gastrointestinal disorders	7 (20.6) [7]
Nausea	7 (20.6) [7]
General disorders and administration site conditions	12 (35.3) [15]
Feeling of body temperature change	3 (8.8) [3]
Feeling of relaxation	3 (8.8) [3]
Feeling cold	2 (5.9) [2]
Feeling hot	2 (5.9) [2]
Chest discomfort	1 (2.9) [1]
Discomfort	1 (2.9) [1]
Gait disturbance	1 (2.9) [1]
Malaise	1 (2.9) [1]
Pain	1 (2.9) [1]

CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030104.sas FINAL (2026-08-28,12:54)

Table 14.3.1.4
Treatment-Emergent Adverse Events with an Onset Date within 48 Hours from CYB003 Administration on Day 8 by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYB003 + CYP Probe N=34 n (%) [E]
Investigations	2 (5.9) [2]
Blood pressure increased	2 (5.9) [2]
Metabolism and nutrition disorders	1 (2.9) [1]
Decreased appetite	1 (2.9) [1]
Nervous system disorders	18 (52.9) [33]
Dizziness	9 (26.5) [9]
Hyperaesthesia	4 (11.8) [4]
Synaesthesia	4 (11.8) [4]
Balance disorder	3 (8.8) [3]
Coordination abnormal	2 (5.9) [2]
Dysaesthesia	2 (5.9) [2]
Headache	2 (5.9) [2]
Cognitive disorder	1 (2.9) [1]
Dyskinesia	1 (2.9) [1]
Parosmia	1 (2.9) [1]
Psychomotor hyperactivity	1 (2.9) [1]
Sensory disturbance	1 (2.9) [1]

CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030104.sas FINAL (2026-08-28,12:54)

Table 14.3.1.4
Treatment-Emergent Adverse Events with an Onset Date within 48 Hours from CYB003 Administration on Day 8 by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYB003 + CYP Probe N=34 n (%) [E]
Nervous system disorders (Cont.)	
Tremor	1 (2.9) [1]
Visual processing disorder	1 (2.9) [1]
Psychiatric disorders	33 (97.1) [126]
Hallucination, visual	24 (70.6) [24]
Affect lability	13 (38.2) [15]
Euphoric mood	10 (29.4) [10]
Illusion	8 (23.5) [9]
Hallucination	6 (17.6) [6]
Agitation	5 (14.7) [6]
Paranoia	5 (14.7) [5]
Time perception altered	5 (14.7) [5]
Derealisation	4 (11.8) [4]
Dissociation	4 (11.8) [6]
Somatic hallucination	4 (11.8) [4]
Disorientation	3 (8.8) [4]
Restlessness	3 (8.8) [3]
Depersonalisation/derealisation disorder	2 (5.9) [2]

CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030104.sas FINAL (2026-08-28,12:54)

Table 14.3.1.4
Treatment-Emergent Adverse Events with an Onset Date within 48 Hours from CYB003 Administration on Day 8 by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYB003 + CYP Probe N=34 n (%) [E]
Psychiatric disorders (Cont.)	
Inappropriate affect	2 (5.9) [2]
Psychomotor retardation	2 (5.9) [2]
Acute stress disorder	1 (2.9) [1]
Anxiety	1 (2.9) [1]
Confusional state	1 (2.9) [1]
Delusion	1 (2.9) [1]
Depressed mood	1 (2.9) [1]
Disinhibition	1 (2.9) [1]
Fear	1 (2.9) [1]
Fear of death	1 (2.9) [1]
Flight of ideas	1 (2.9) [1]
Hallucination, auditory	1 (2.9) [1]
Hallucination, tactile	1 (2.9) [2]
Logorrhoea	1 (2.9) [1]
Loose associations	1 (2.9) [1]
Nervousness	1 (2.9) [1]
Perseveration	1 (2.9) [1]
Sexually inappropriate behaviour	1 (2.9) [1]

CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030104.sas FINAL (2026-08-28,12:54)

Table 14.3.1.4
Treatment-Emergent Adverse Events with an Onset Date within 48 Hours from CYB003 Administration on Day 8 by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYB003 + CYP Probe N=34 n (%) [E]
Psychiatric disorders (Cont.)	
Somatic delusion	1 (2.9) [1]
Thinking abnormal	1 (2.9) [1]
Renal and urinary disorders	1 (2.9) [1]
Urinary incontinence	1 (2.9) [1]
Respiratory, thoracic and mediastinal disorders	2 (5.9) [2]
Dyspnoea	1 (2.9) [1]
Hyperventilation	1 (2.9) [1]
Skin and subcutaneous tissue disorders	2 (5.9) [2]
Hyperhidrosis	2 (5.9) [2]
Surgical and medical procedures	1 (2.9) [1]
Analgesic therapy	1 (2.9) [1]
Vascular disorders	3 (8.8) [3]
Orthostatic hypotension	3 (8.8) [3]

CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030104.sas FINAL (2026-08-28,12:54)

Table 14.3.1.5
Related Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any related TEAE	3 (8.3) [3]	34 (100) [194]	34 (94.4) [197]
Ear and labyrinth disorders	0	2 (5.9) [2]	2 (5.6) [2]
Hyperacusis	0	2 (5.9) [2]	2 (5.6) [2]
Gastrointestinal disorders	0	7 (20.6) [7]	7 (19.4) [7]
Nausea	0	7 (20.6) [7]	7 (19.4) [7]
General disorders and administration site conditions	0	12 (35.3) [15]	12 (33.3) [15]
Feeling of body temperature change	0	3 (8.8) [3]	3 (8.3) [3]
Feeling of relaxation	0	3 (8.8) [3]	3 (8.3) [3]
Feeling cold	0	2 (5.9) [2]	2 (5.6) [2]
Feeling hot	0	2 (5.9) [2]	2 (5.6) [2]
Chest discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Gait disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Malaise	0	1 (2.9) [1]	1 (2.8) [1]
Pain	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related AEs are defined as events reported as related or possibly related.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030105.sas FINAL (2026-08-28,12:54)

Table 14.3.1.5
Related Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Investigations	0	2 (5.9) [2]	2 (5.6) [2]
Blood pressure increased	0	2 (5.9) [2]	2 (5.6) [2]
Metabolism and nutrition disorders	0	1 (2.9) [1]	1 (2.8) [1]
Decreased appetite	0	1 (2.9) [1]	1 (2.8) [1]
Nervous system disorders	0	18 (52.9) [32]	18 (50.0) [32]
Dizziness	0	8 (23.5) [8]	8 (22.2) [8]
Hyperaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Synaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Balance disorder	0	3 (8.8) [3]	3 (8.3) [3]
Coordination abnormal	0	2 (5.9) [2]	2 (5.6) [2]
Dysaesthesia	0	2 (5.9) [2]	2 (5.6) [2]
Headache	0	2 (5.9) [2]	2 (5.6) [2]
Cognitive disorder	0	1 (2.9) [1]	1 (2.8) [1]
Dyskinesia	0	1 (2.9) [1]	1 (2.8) [1]
Parosmia	0	1 (2.9) [1]	1 (2.8) [1]
Psychomotor hyperactivity	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related AEs are defined as events reported as related or possibly related.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030105.sas FINAL (2026-08-28,12:54)

Table 14.3.1.5
Related Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Nervous system disorders (Cont.)			
Sensory disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Tremor	0	1 (2.9) [1]	1 (2.8) [1]
Visual processing disorder	0	1 (2.9) [1]	1 (2.8) [1]
Psychiatric disorders	0	33 (97.1) [126]	33 (91.7) [126]
Hallucination, visual	0	24 (70.6) [24]	24 (66.7) [24]
Affect lability	0	13 (38.2) [15]	13 (36.1) [15]
Euphoric mood	0	10 (29.4) [10]	10 (27.8) [10]
Illusion	0	8 (23.5) [9]	8 (22.2) [9]
Hallucination	0	6 (17.6) [6]	6 (16.7) [6]
Agitation	0	5 (14.7) [6]	5 (13.9) [6]
Paranoia	0	5 (14.7) [5]	5 (13.9) [5]
Time perception altered	0	5 (14.7) [5]	5 (13.9) [5]
Derealisation	0	4 (11.8) [4]	4 (11.1) [4]
Dissociation	0	4 (11.8) [6]	4 (11.1) [6]
Somatic hallucination	0	4 (11.8) [4]	4 (11.1) [4]
Disorientation	0	3 (8.8) [4]	3 (8.3) [4]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related AEs are defined as events reported as related or possibly related.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030105.sas FINAL (2026-08-28,12:54)

Table 14.3.1.5
Related Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders (Cont.)			
Restlessness	0	3 (8.8) [3]	3 (8.3) [3]
Depersonalisation/derealisation disorder	0	2 (5.9) [2]	2 (5.6) [2]
Inappropriate affect	0	2 (5.9) [2]	2 (5.6) [2]
Psychomotor retardation	0	2 (5.9) [2]	2 (5.6) [2]
Acute stress disorder	0	1 (2.9) [1]	1 (2.8) [1]
Anxiety	0	1 (2.9) [1]	1 (2.8) [1]
Confusional state	0	1 (2.9) [1]	1 (2.8) [1]
Delusion	0	1 (2.9) [1]	1 (2.8) [1]
Depressed mood	0	1 (2.9) [1]	1 (2.8) [1]
Disinhibition	0	1 (2.9) [1]	1 (2.8) [1]
Fear	0	1 (2.9) [1]	1 (2.8) [1]
Fear of death	0	1 (2.9) [1]	1 (2.8) [1]
Flight of ideas	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, auditory	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, tactile	0	1 (2.9) [2]	1 (2.8) [2]
Logorrhoea	0	1 (2.9) [1]	1 (2.8) [1]
Loose associations	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related AEs are defined as events reported as related or possibly related.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030105.sas FINAL (2026-08-28,12:54)

Table 14.3.1.5
Related Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders (Cont.)			
Nervousness	0	1 (2.9) [1]	1 (2.8) [1]
Perseveration	0	1 (2.9) [1]	1 (2.8) [1]
Sexually inappropriate behaviour	0	1 (2.9) [1]	1 (2.8) [1]
Somatic delusion	0	1 (2.9) [1]	1 (2.8) [1]
Thinking abnormal	0	1 (2.9) [1]	1 (2.8) [1]
Renal and urinary disorders	0	1 (2.9) [1]	1 (2.8) [1]
Urinary incontinence	0	1 (2.9) [1]	1 (2.8) [1]
Respiratory, thoracic and mediastinal disorders	0	2 (5.9) [2]	2 (5.6) [2]
Dyspnoea	0	1 (2.9) [1]	1 (2.8) [1]
Hyperventilation	0	1 (2.9) [1]	1 (2.8) [1]
Skin and subcutaneous tissue disorders	0	2 (5.9) [2]	2 (5.6) [2]
Hyperhidrosis	0	2 (5.9) [2]	2 (5.6) [2]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related AEs are defined as events reported as related or possibly related.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030105.sas FINAL (2026-08-28,12:54)

Table 14.3.1.5
Related Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Surgical and medical procedures	0	1 (2.9) [1]	1 (2.8) [1]
Analgesic therapy	0	1 (2.9) [1]	1 (2.8) [1]
Vascular disorders	3 (8.3) [3]	3 (8.8) [3]	5 (13.9) [6]
Orthostatic hypotension	2 (5.6) [2]	3 (8.8) [3]	4 (11.1) [5]
Hypotension	1 (2.8) [1]	0	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related AEs are defined as events reported as related or possibly related.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030105.sas FINAL (2026-08-28,12:54)

Table 14.3.1.6
Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any SMRE	0	34 (100) [172]	34 (94.4) [172]
Ear and labyrinth disorders	0	2 (5.9) [2]	2 (5.6) [2]
Hyperacusis	0	2 (5.9) [2]	2 (5.6) [2]
Gastrointestinal disorders	0	1 (2.9) [1]	1 (2.8) [1]
Nausea	0	1 (2.9) [1]	1 (2.8) [1]
General disorders and administration site conditions	0	10 (29.4) [11]	10 (27.8) [11]
Feeling of body temperature change	0	3 (8.8) [3]	3 (8.3) [3]
Feeling of relaxation	0	3 (8.8) [3]	3 (8.3) [3]
Feeling cold	0	2 (5.9) [2]	2 (5.6) [2]
Discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Feeling hot	0	1 (2.9) [1]	1 (2.8) [1]
Gait disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Metabolism and nutrition disorders	0	1 (2.9) [1]	1 (2.8) [1]
Decreased appetite	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030106.sas FINAL (2026-08-28,12:54)

Table 14.3.1.6
Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Nervous system disorders	0	16 (47.1) [28]	16 (44.4) [28]
Dizziness	0	5 (14.7) [5]	5 (13.9) [5]
Hyperaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Synaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Balance disorder	0	3 (8.8) [3]	3 (8.3) [3]
Coordination abnormal	0	2 (5.9) [2]	2 (5.6) [2]
Dysaesthesia	0	2 (5.9) [2]	2 (5.6) [2]
Cognitive disorder	0	1 (2.9) [1]	1 (2.8) [1]
Dyskinesia	0	1 (2.9) [1]	1 (2.8) [1]
Headache	0	1 (2.9) [1]	1 (2.8) [1]
Parosmia	0	1 (2.9) [1]	1 (2.8) [1]
Psychomotor hyperactivity	0	1 (2.9) [1]	1 (2.8) [1]
Sensory disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Tremor	0	1 (2.9) [1]	1 (2.8) [1]
Visual processing disorder	0	1 (2.9) [1]	1 (2.8) [1]
Psychiatric disorders	0	33 (97.1) [125]	33 (91.7) [125]
Hallucination, visual	0	24 (70.6) [24]	24 (66.7) [24]
Affect lability	0	13 (38.2) [15]	13 (36.1) [15]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030106.sas FINAL (2026-08-28,12:54)

Table 14.3.1.6
Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders (Cont.)			
Euphoric mood	0	10 (29.4) [10]	10 (27.8) [10]
Illusion	0	8 (23.5) [9]	8 (22.2) [9]
Hallucination	0	6 (17.6) [6]	6 (16.7) [6]
Agitation	0	5 (14.7) [5]	5 (13.9) [5]
Paranoia	0	5 (14.7) [5]	5 (13.9) [5]
Time perception altered	0	5 (14.7) [5]	5 (13.9) [5]
Derealisation	0	4 (11.8) [4]	4 (11.1) [4]
Dissociation	0	4 (11.8) [6]	4 (11.1) [6]
Somatic hallucination	0	4 (11.8) [4]	4 (11.1) [4]
Disorientation	0	3 (8.8) [4]	3 (8.3) [4]
Restlessness	0	3 (8.8) [3]	3 (8.3) [3]
Depersonalisation/derealisation disorder	0	2 (5.9) [2]	2 (5.6) [2]
Inappropriate affect	0	2 (5.9) [2]	2 (5.6) [2]
Psychomotor retardation	0	2 (5.9) [2]	2 (5.6) [2]
Acute stress disorder	0	1 (2.9) [1]	1 (2.8) [1]
Anxiety	0	1 (2.9) [1]	1 (2.8) [1]
Confusional state	0	1 (2.9) [1]	1 (2.8) [1]
Delusion	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030106.sas FINAL (2026-08-28,12:54)

Table 14.3.1.6
Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders (Cont.)			
Depressed mood	0	1 (2.9) [1]	1 (2.8) [1]
Disinhibition	0	1 (2.9) [1]	1 (2.8) [1]
Fear	0	1 (2.9) [1]	1 (2.8) [1]
Fear of death	0	1 (2.9) [1]	1 (2.8) [1]
Flight of ideas	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, auditory	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, tactile	0	1 (2.9) [2]	1 (2.8) [2]
Logorrhoea	0	1 (2.9) [1]	1 (2.8) [1]
Loose associations	0	1 (2.9) [1]	1 (2.8) [1]
Nervousness	0	1 (2.9) [1]	1 (2.8) [1]
Perseveration	0	1 (2.9) [1]	1 (2.8) [1]
Sexually inappropriate behaviour	0	1 (2.9) [1]	1 (2.8) [1]
Somatic delusion	0	1 (2.9) [1]	1 (2.8) [1]
Thinking abnormal	0	1 (2.9) [1]	1 (2.8) [1]
Respiratory, thoracic and mediastinal disorders	0	1 (2.9) [1]	1 (2.8) [1]
Hyperventilation	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030106.sas FINAL (2026-08-28,12:54)

Table 14.3.1.6
Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Skin and subcutaneous tissue disorders	0	2 (5.9) [2]	2 (5.6) [2]
Hyperhidrosis	0	2 (5.9) [2]	2 (5.6) [2]
Surgical and medical procedures	0	1 (2.9) [1]	1 (2.8) [1]
Analgesic therapy	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030106.sas FINAL (2026-08-28,12:54)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any SMRE	0	34 (100) [172]	34 (94.4) [172]
Severe	0	0	0
Moderate	0	7 (20.6) [29]	7 (19.4) [29]
Mild	0	27 (79.4) [116]	27 (75.0) [116]
Ear and labyrinth disorders	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Hyperacusis	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Gastrointestinal disorders	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Nausea	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
General disorders and administration site conditions	0	10 (29.4) [11]	10 (27.8) [11]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	9 (26.5) [10]	9 (25.0) [10]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Feeling of body temperature change	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	0	0
Mild	0	3 (8.8) [3]	3 (8.3) [3]
Feeling of relaxation	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	0	0
Mild	0	3 (8.8) [3]	3 (8.3) [3]
Feeling cold	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Feeling hot	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Gait disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Metabolism and nutrition disorders	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Decreased appetite	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Nervous system disorders	0	16 (47.1) [28]	16 (44.4) [28]
Severe	0	0	0
Moderate	0	2 (5.9) [3]	2 (5.6) [3]
Mild	0	14 (41.2) [23]	14 (38.9) [23]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Dizziness	0	5 (14.7) [5]	5 (13.9) [5]
Severe	0	0	0
Moderate	0	0	0
Mild	0	5 (14.7) [5]	5 (13.9) [5]
Hyperaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Severe	0	0	0
Moderate	0	0	0
Mild	0	4 (11.8) [4]	4 (11.1) [4]
Synaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Severe	0	0	0
Moderate	0	0	0
Mild	0	4 (11.8) [4]	4 (11.1) [4]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Balance disorder	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	0	0
Mild	0	3 (8.8) [3]	3 (8.3) [3]
Coordination abnormal	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Dysaesthesia	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Cognitive disorder	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Dyskinesia	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Headache	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Parosmia	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Psychomotor hyperactivity	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Sensory disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Tremor	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Visual processing disorder	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Psychiatric disorders	0	33 (97.1) [125]	33 (91.7) [125]
Severe	0	0	0
Moderate	0	7 (20.6) [25]	7 (19.4) [25]
Mild	0	26 (76.5) [81]	26 (72.2) [81]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Hallucination, visual	0	24 (70.6) [24]	24 (66.7) [24]
Severe	0	0	0
Moderate	0	2 (5.9) [2]	2 (5.6) [2]
Mild	0	22 (64.7) [22]	22 (61.1) [22]
Affect lability	0	13 (38.2) [15]	13 (36.1) [15]
Severe	0	0	0
Moderate	0	2 (5.9) [3]	2 (5.6) [3]
Mild	0	11 (32.4) [12]	11 (30.6) [12]
Euphoric mood	0	10 (29.4) [10]	10 (27.8) [10]
Severe	0	0	0
Moderate	0	0	0
Mild	0	10 (29.4) [10]	10 (27.8) [10]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Illusion	0	8 (23.5) [9]	8 (22.2) [9]
Severe	0	0	0
Moderate	0	2 (5.9) [2]	2 (5.6) [2]
Mild	0	6 (17.6) [7]	6 (16.7) [7]
Hallucination	0	6 (17.6) [6]	6 (16.7) [6]
Severe	0	0	0
Moderate	0	0	0
Mild	0	6 (17.6) [6]	6 (16.7) [6]
Agitation	0	5 (14.7) [5]	5 (13.9) [5]
Severe	0	0	0
Moderate	0	3 (8.8) [3]	3 (8.3) [3]
Mild	0	2 (5.9) [2]	2 (5.6) [2]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Paranoia	0	5 (14.7) [5]	5 (13.9) [5]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	4 (11.8) [4]	4 (11.1) [4]
Time perception altered	0	5 (14.7) [5]	5 (13.9) [5]
Severe	0	0	0
Moderate	0	0	0
Mild	0	5 (14.7) [5]	5 (13.9) [5]
Derealisation	0	4 (11.8) [4]	4 (11.1) [4]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	3 (8.8) [3]	3 (8.3) [3]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Dissociation	0	4 (11.8) [6]	4 (11.1) [6]
Severe	0	0	0
Moderate	0	2 (5.9) [2]	2 (5.6) [2]
Mild	0	2 (5.9) [4]	2 (5.6) [4]
Somatic hallucination	0	4 (11.8) [4]	4 (11.1) [4]
Severe	0	0	0
Moderate	0	0	0
Mild	0	4 (11.8) [4]	4 (11.1) [4]
Disorientation	0	3 (8.8) [4]	3 (8.3) [4]
Severe	0	0	0
Moderate	0	3 (8.8) [4]	3 (8.3) [4]
Mild	0	0	0

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Restlessness	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	0	0
Mild	0	3 (8.8) [3]	3 (8.3) [3]
Depersonalisation/derealisation disorder	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Inappropriate affect	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychomotor retardation	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Acute stress disorder	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Anxiety	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Confusional state	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Delusion	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Depressed mood	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Disinhibition	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Fear	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Fear of death	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Flight of ideas	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Hallucination, auditory	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, tactile	0	1 (2.9) [2]	1 (2.8) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [2]	1 (2.8) [2]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Logorrhoea	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Loose associations	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Nervousness	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Perseveration	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Sexually inappropriate behaviour	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Somatic delusion	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Thinking abnormal	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Respiratory, thoracic and mediastinal disorders	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Hyperventilation	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Skin and subcutaneous tissue disorders	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Hyperhidrosis	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Surgical and medical procedures	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.7
Session Monitor-Reported Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Analgesic therapy	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030107.sas FINAL (2026-08-28,12:55)

Table 14.3.1.8
Related Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any related SMRE	0	34 (100) [172]	34 (94.4) [172]
Ear and labyrinth disorders	0	2 (5.9) [2]	2 (5.6) [2]
Hyperacusis	0	2 (5.9) [2]	2 (5.6) [2]
Gastrointestinal disorders	0	1 (2.9) [1]	1 (2.8) [1]
Nausea	0	1 (2.9) [1]	1 (2.8) [1]
General disorders and administration site conditions	0	10 (29.4) [11]	10 (27.8) [11]
Feeling of body temperature change	0	3 (8.8) [3]	3 (8.3) [3]
Feeling of relaxation	0	3 (8.8) [3]	3 (8.3) [3]
Feeling cold	0	2 (5.9) [2]	2 (5.6) [2]
Discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Feeling hot	0	1 (2.9) [1]	1 (2.8) [1]
Gait disturbance	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related SMREs are defined as events reported as related or possibly related.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030108.sas FINAL (2026-08-28,12:55)

Table 14.3.1.8
Related Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Metabolism and nutrition disorders	0	1 (2.9) [1]	1 (2.8) [1]
Decreased appetite	0	1 (2.9) [1]	1 (2.8) [1]
Nervous system disorders	0	16 (47.1) [28]	16 (44.4) [28]
Dizziness	0	5 (14.7) [5]	5 (13.9) [5]
Hyperaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Synaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Balance disorder	0	3 (8.8) [3]	3 (8.3) [3]
Coordination abnormal	0	2 (5.9) [2]	2 (5.6) [2]
Dysaesthesia	0	2 (5.9) [2]	2 (5.6) [2]
Cognitive disorder	0	1 (2.9) [1]	1 (2.8) [1]
Dyskinesia	0	1 (2.9) [1]	1 (2.8) [1]
Headache	0	1 (2.9) [1]	1 (2.8) [1]
Parosmia	0	1 (2.9) [1]	1 (2.8) [1]
Psychomotor hyperactivity	0	1 (2.9) [1]	1 (2.8) [1]
Sensory disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Tremor	0	1 (2.9) [1]	1 (2.8) [1]
Visual processing disorder	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related SMREs are defined as events reported as related or possibly related.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030108.sas FINAL (2026-08-28,12:55)

Table 14.3.1.8
Related Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders	0	33 (97.1) [125]	33 (91.7) [125]
Hallucination, visual	0	24 (70.6) [24]	24 (66.7) [24]
Affect lability	0	13 (38.2) [15]	13 (36.1) [15]
Euphoric mood	0	10 (29.4) [10]	10 (27.8) [10]
Illusion	0	8 (23.5) [9]	8 (22.2) [9]
Hallucination	0	6 (17.6) [6]	6 (16.7) [6]
Agitation	0	5 (14.7) [5]	5 (13.9) [5]
Paranoia	0	5 (14.7) [5]	5 (13.9) [5]
Time perception altered	0	5 (14.7) [5]	5 (13.9) [5]
Derealisation	0	4 (11.8) [4]	4 (11.1) [4]
Dissociation	0	4 (11.8) [6]	4 (11.1) [6]
Somatic hallucination	0	4 (11.8) [4]	4 (11.1) [4]
Disorientation	0	3 (8.8) [4]	3 (8.3) [4]
Restlessness	0	3 (8.8) [3]	3 (8.3) [3]
Depersonalisation/derealisation disorder	0	2 (5.9) [2]	2 (5.6) [2]
Inappropriate affect	0	2 (5.9) [2]	2 (5.6) [2]
Psychomotor retardation	0	2 (5.9) [2]	2 (5.6) [2]
Acute stress disorder	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related SMREs are defined as events reported as related or possibly related.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030108.sas FINAL (2026-08-28,12:55)

Table 14.3.1.8
Related Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders (Cont.)			
Anxiety	0	1 (2.9) [1]	1 (2.8) [1]
Confusional state	0	1 (2.9) [1]	1 (2.8) [1]
Delusion	0	1 (2.9) [1]	1 (2.8) [1]
Depressed mood	0	1 (2.9) [1]	1 (2.8) [1]
Disinhibition	0	1 (2.9) [1]	1 (2.8) [1]
Fear	0	1 (2.9) [1]	1 (2.8) [1]
Fear of death	0	1 (2.9) [1]	1 (2.8) [1]
Flight of ideas	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, auditory	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, tactile	0	1 (2.9) [2]	1 (2.8) [2]
Logorrhoea	0	1 (2.9) [1]	1 (2.8) [1]
Loose associations	0	1 (2.9) [1]	1 (2.8) [1]
Nervousness	0	1 (2.9) [1]	1 (2.8) [1]
Perseveration	0	1 (2.9) [1]	1 (2.8) [1]
Sexually inappropriate behaviour	0	1 (2.9) [1]	1 (2.8) [1]
Somatic delusion	0	1 (2.9) [1]	1 (2.8) [1]
Thinking abnormal	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related SMREs are defined as events reported as related or possibly related.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030108.sas FINAL (2026-08-28,12:55)

Table 14.3.1.8
Related Session Monitor-Reported Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Respiratory, thoracic and mediastinal disorders	0	1 (2.9) [1]	1 (2.8) [1]
Hyperventilation	0	1 (2.9) [1]	1 (2.8) [1]
Skin and subcutaneous tissue disorders	0	2 (5.9) [2]	2 (5.6) [2]
Hyperhidrosis	0	2 (5.9) [2]	2 (5.6) [2]
Surgical and medical procedures	0	1 (2.9) [1]	1 (2.8) [1]
Analgesic therapy	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes SMREs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes SMREs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related SMREs are defined as events reported as related or possibly related.
MedDRA version 27.1.
SMRE = Session Monitor-reported event.

Source: Listing 16.2.7.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030108.sas FINAL (2026-08-28,12:55)

Table 14.3.1.9
Potential Abuse-Related Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any potential abuse-related AE	0	34 (100) [93]	34 (94.4) [93]
General disorders and administration site conditions	0	3 (8.8) [3]	3 (8.3) [3]
Feeling of relaxation	0	3 (8.8) [3]	3 (8.3) [3]
Nervous system disorders	0	2 (5.9) [2]	2 (5.6) [2]
Psychomotor hyperactivity	0	1 (2.9) [1]	1 (2.8) [1]
Sensory disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Psychiatric disorders	0	33 (97.1) [88]	33 (91.7) [88]
Hallucination, visual	0	24 (70.6) [24]	24 (66.7) [24]
Affect lability	0	13 (38.2) [15]	13 (36.1) [15]
Euphoric mood	0	10 (29.4) [10]	10 (27.8) [10]
Hallucination	0	6 (17.6) [6]	6 (16.7) [6]
Agitation	0	5 (14.7) [6]	5 (13.9) [6]
Paranoia	0	5 (14.7) [5]	5 (13.9) [5]
Derealisation	0	4 (11.8) [4]	4 (11.1) [4]
Disorientation	0	3 (8.8) [4]	3 (8.3) [4]

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030109.sas FINAL (2026-08-28,12:55)

Table 14.3.1.9
Potential Abuse-Related Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders (Cont.)			
Restlessness	0	3 (8.8) [3]	3 (8.3) [3]
Inappropriate affect	0	2 (5.9) [2]	2 (5.6) [2]
Anxiety	0	1 (2.9) [1]	1 (2.8) [1]
Confusional state	0	1 (2.9) [1]	1 (2.8) [1]
Delusion	0	1 (2.9) [1]	1 (2.8) [1]
Disinhibition	0	1 (2.9) [1]	1 (2.8) [1]
Flight of ideas	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, auditory	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, tactile	0	1 (2.9) [2]	1 (2.8) [2]
Thinking abnormal	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030109.sas FINAL (2026-08-28,12:55)

Table 14.3.1.10
Potential Abuse-Related Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any potential abuse-related AE	0	34 (100) [93]	34 (94.4) [93]
Severe	0	0	0
Moderate	0	7 (20.6) [17]	7 (19.4) [17]
Mild	0	27 (79.4) [62]	27 (75.0) [62]
General disorders and administration site conditions	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	0	0
Mild	0	3 (8.8) [3]	3 (8.3) [3]
Feeling of relaxation	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	0	0
Mild	0	3 (8.8) [3]	3 (8.3) [3]

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030110.sas FINAL (2026-08-28,12:55)

Table 14.3.1.10
Potential Abuse-Related Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Nervous system disorders	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Psychomotor hyperactivity	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Sensory disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030110.sas FINAL (2026-08-28,12:55)

Table 14.3.1.10
Potential Abuse-Related Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders	0	33 (97.1) [88]	33 (91.7) [88]
Severe	0	0	0
Moderate	0	7 (20.6) [17]	7 (19.4) [17]
Mild	0	26 (76.5) [57]	26 (72.2) [57]
Hallucination, visual	0	24 (70.6) [24]	24 (66.7) [24]
Severe	0	0	0
Moderate	0	2 (5.9) [2]	2 (5.6) [2]
Mild	0	22 (64.7) [22]	22 (61.1) [22]
Affect lability	0	13 (38.2) [15]	13 (36.1) [15]
Severe	0	0	0
Moderate	0	2 (5.9) [3]	2 (5.6) [3]
Mild	0	11 (32.4) [12]	11 (30.6) [12]

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030110.sas FINAL (2026-08-28,12:55)

Table 14.3.1.10
Potential Abuse-Related Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Euphoric mood	0	10 (29.4) [10]	10 (27.8) [10]
Severe	0	0	0
Moderate	0	0	0
Mild	0	10 (29.4) [10]	10 (27.8) [10]
Hallucination	0	6 (17.6) [6]	6 (16.7) [6]
Severe	0	0	0
Moderate	0	0	0
Mild	0	6 (17.6) [6]	6 (16.7) [6]
Agitation	0	5 (14.7) [6]	5 (13.9) [6]
Severe	0	0	0
Moderate	0	3 (8.8) [4]	3 (8.3) [4]
Mild	0	2 (5.9) [2]	2 (5.6) [2]

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030110.sas FINAL (2026-08-28,12:55)

Table 14.3.1.10
Potential Abuse-Related Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Paranoia	0	5 (14.7) [5]	5 (13.9) [5]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	4 (11.8) [4]	4 (11.1) [4]
Derealisation	0	4 (11.8) [4]	4 (11.1) [4]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	3 (8.8) [3]	3 (8.3) [3]
Disorientation	0	3 (8.8) [4]	3 (8.3) [4]
Severe	0	0	0
Moderate	0	3 (8.8) [4]	3 (8.3) [4]
Mild	0	0	0

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030110.sas FINAL (2026-08-28,12:55)

Table 14.3.1.10
Potential Abuse-Related Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Restlessness	0	3 (8.8) [3]	3 (8.3) [3]
Severe	0	0	0
Moderate	0	0	0
Mild	0	3 (8.8) [3]	3 (8.3) [3]
Inappropriate affect	0	2 (5.9) [2]	2 (5.6) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	2 (5.9) [2]	2 (5.6) [2]
Anxiety	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030110.sas FINAL (2026-08-28,12:55)

Table 14.3.1.10
Potential Abuse-Related Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Confusional state	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Delusion	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Disinhibition	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030110.sas FINAL (2026-08-28,12:55)

Table 14.3.1.10
Potential Abuse-Related Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Flight of ideas	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	1 (2.9) [1]	1 (2.8) [1]
Mild	0	0	0
Hallucination, auditory	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, tactile	0	1 (2.9) [2]	1 (2.8) [2]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [2]	1 (2.8) [2]

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030110.sas FINAL (2026-08-28,12:55)

Table 14.3.1.10
Potential Abuse-Related Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Thinking abnormal	0	1 (2.9) [1]	1 (2.8) [1]
Severe	0	0	0
Moderate	0	0	0
Mild	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030110.sas FINAL (2026-08-28,12:55)

Table 14.3.1.11
Related Potential Abuse-Related Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any related potential abuse-related AE	0	34 (100) [93]	34 (94.4) [93]
General disorders and administration site conditions	0	3 (8.8) [3]	3 (8.3) [3]
Feeling of relaxation	0	3 (8.8) [3]	3 (8.3) [3]
Nervous system disorders	0	2 (5.9) [2]	2 (5.6) [2]
Psychomotor hyperactivity	0	1 (2.9) [1]	1 (2.8) [1]
Sensory disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Psychiatric disorders	0	33 (97.1) [88]	33 (91.7) [88]
Hallucination, visual	0	24 (70.6) [24]	24 (66.7) [24]
Affect lability	0	13 (38.2) [15]	13 (36.1) [15]
Euphoric mood	0	10 (29.4) [10]	10 (27.8) [10]
Hallucination	0	6 (17.6) [6]	6 (16.7) [6]
Agitation	0	5 (14.7) [6]	5 (13.9) [6]
Paranoia	0	5 (14.7) [5]	5 (13.9) [5]
Derealisation	0	4 (11.8) [4]	4 (11.1) [4]

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related AEs are defined as events reported as related or possibly related.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030111.sas FINAL (2026-08-28,12:55)

Table 14.3.1.11
Related Potential Abuse-Related Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders (Cont.)			
Disorientation	0	3 (8.8) [4]	3 (8.3) [4]
Restlessness	0	3 (8.8) [3]	3 (8.3) [3]
Inappropriate affect	0	2 (5.9) [2]	2 (5.6) [2]
Anxiety	0	1 (2.9) [1]	1 (2.8) [1]
Confusional state	0	1 (2.9) [1]	1 (2.8) [1]
Delusion	0	1 (2.9) [1]	1 (2.8) [1]
Disinhibition	0	1 (2.9) [1]	1 (2.8) [1]
Flight of ideas	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, auditory	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, tactile	0	1 (2.9) [2]	1 (2.8) [2]
Thinking abnormal	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes potential abuse-related AEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes potential abuse-related AEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related AEs are defined as events reported as related or possibly related.
MedDRA version 27.1.
AE = adverse event.

Source: Listing 16.2.7.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030111.sas FINAL (2026-08-28,12:55)

Table 14.3.1.12
Non-SMRE Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any non-SMRE TEAE	4 (11.1) [4]	14 (41.2) [24]	16 (44.4) [28]
Gastrointestinal disorders	0	6 (17.6) [6]	6 (16.7) [6]
Nausea	0	6 (17.6) [6]	6 (16.7) [6]
General disorders and administration site conditions	1 (2.8) [1]	4 (11.8) [4]	5 (13.9) [5]
Chest discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Feeling hot	0	1 (2.9) [1]	1 (2.8) [1]
Malaise	0	1 (2.9) [1]	1 (2.8) [1]
Pain	0	1 (2.9) [1]	1 (2.8) [1]
Vessel puncture site pain	1 (2.8) [1]	0	1 (2.8) [1]
Investigations	0	2 (5.9) [2]	2 (5.6) [2]
Blood pressure increased	0	2 (5.9) [2]	2 (5.6) [2]
Nervous system disorders	0	4 (11.8) [5]	4 (11.1) [5]
Dizziness	0	4 (11.8) [4]	4 (11.1) [4]
Headache	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes non-SMRE TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes non-SMRE TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event; TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030112.sas FINAL (2026-08-28,12:55)

Table 14.3.1.12
Non-SMRE Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders	0	1 (2.9) [1]	1 (2.8) [1]
Agitation	0	1 (2.9) [1]	1 (2.8) [1]
Renal and urinary disorders	0	1 (2.9) [1]	1 (2.8) [1]
Urinary incontinence	0	1 (2.9) [1]	1 (2.8) [1]
Respiratory, thoracic and mediastinal disorders	0	1 (2.9) [1]	1 (2.8) [1]
Dyspnoea	0	1 (2.9) [1]	1 (2.8) [1]
Skin and subcutaneous tissue disorders	0	1 (2.9) [1]	1 (2.8) [1]
Ingrown hair	0	1 (2.9) [1]	1 (2.8) [1]
Vascular disorders	3 (8.3) [3]	3 (8.8) [3]	5 (13.9) [6]
Orthostatic hypotension	2 (5.6) [2]	3 (8.8) [3]	4 (11.1) [5]
Hypotension	1 (2.8) [1]	0	1 (2.8) [1]

CYP Probe refers to Period 1 and includes non-SMRE TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes non-SMRE TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
SMRE = Session Monitor-reported event; TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030112.sas FINAL (2026-08-28,12:55)

Table 14.3.1.13
Non-SMRE Related Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any non-SMRE related TEAE	3 (8.3) [3]	12 (35.3) [22]	13 (36.1) [25]
Gastrointestinal disorders	0	6 (17.6) [6]	6 (16.7) [6]
Nausea	0	6 (17.6) [6]	6 (16.7) [6]
General disorders and administration site conditions	0	4 (11.8) [4]	4 (11.1) [4]
Chest discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Feeling hot	0	1 (2.9) [1]	1 (2.8) [1]
Malaise	0	1 (2.9) [1]	1 (2.8) [1]
Pain	0	1 (2.9) [1]	1 (2.8) [1]
Investigations	0	2 (5.9) [2]	2 (5.6) [2]
Blood pressure increased	0	2 (5.9) [2]	2 (5.6) [2]
Nervous system disorders	0	3 (8.8) [4]	3 (8.3) [4]
Dizziness	0	3 (8.8) [3]	3 (8.3) [3]
Headache	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes non-SMRE related TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes non-SMRE related TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related TEAEs are defined as events reported as related or possibly related.
MedDRA version 27.1.
SMRE = Session Monitor-reported event; TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030113.sas FINAL (2026-08-28,12:55)

Table 14.3.1.13
Non-SMRE Related Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Psychiatric disorders	0	1 (2.9) [1]	1 (2.8) [1]
Agitation	0	1 (2.9) [1]	1 (2.8) [1]
Renal and urinary disorders	0	1 (2.9) [1]	1 (2.8) [1]
Urinary incontinence	0	1 (2.9) [1]	1 (2.8) [1]
Respiratory, thoracic and mediastinal disorders	0	1 (2.9) [1]	1 (2.8) [1]
Dyspnoea	0	1 (2.9) [1]	1 (2.8) [1]
Vascular disorders	3 (8.3) [3]	3 (8.8) [3]	5 (13.9) [6]
Orthostatic hypotension	2 (5.6) [2]	3 (8.8) [3]	4 (11.1) [5]
Hypotension	1 (2.8) [1]	0	1 (2.8) [1]

CYP Probe refers to Period 1 and includes non-SMRE related TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes non-SMRE related TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related TEAEs are defined as events reported as related or possibly related.
MedDRA version 27.1.
SMRE = Session Monitor-reported event; TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030113.sas FINAL (2026-08-28,12:55)

Table 14.3.1.14
Treatment-Emergent Adverse Events by MedDRA Preferred Term
Safety Population

Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any TEAE	4 (11.1) [4]	34 (100) [196]	34 (94.4) [200]
Hallucination, visual	0	24 (70.6) [24]	24 (66.7) [24]
Affect lability	0	13 (38.2) [15]	13 (36.1) [15]
Euphoric mood	0	10 (29.4) [10]	10 (27.8) [10]
Dizziness	0	9 (26.5) [9]	9 (25.0) [9]
Illusion	0	8 (23.5) [9]	8 (22.2) [9]
Nausea	0	7 (20.6) [7]	7 (19.4) [7]
Hallucination	0	6 (17.6) [6]	6 (16.7) [6]
Agitation	0	5 (14.7) [6]	5 (13.9) [6]
Paranoia	0	5 (14.7) [5]	5 (13.9) [5]
Time perception altered	0	5 (14.7) [5]	5 (13.9) [5]
Derealisation	0	4 (11.8) [4]	4 (11.1) [4]
Dissociation	0	4 (11.8) [6]	4 (11.1) [6]
Hyperaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Orthostatic hypotension	2 (5.6) [2]	3 (8.8) [3]	4 (11.1) [5]
Somatic hallucination	0	4 (11.8) [4]	4 (11.1) [4]
Synaesthesia	0	4 (11.8) [4]	4 (11.1) [4]
Balance disorder	0	3 (8.8) [3]	3 (8.3) [3]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT.n is the number of participants with the measurement available.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030114.sas FINAL (2026-08-28,12:55)

Table 14.3.1.14
Treatment-Emergent Adverse Events by MedDRA Preferred Term
Safety Population

Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Disorientation	0	3 (8.8) [4]	3 (8.3) [4]
Feeling of body temperature change	0	3 (8.8) [3]	3 (8.3) [3]
Feeling of relaxation	0	3 (8.8) [3]	3 (8.3) [3]
Restlessness	0	3 (8.8) [3]	3 (8.3) [3]
Blood pressure increased	0	2 (5.9) [2]	2 (5.6) [2]
Coordination abnormal	0	2 (5.9) [2]	2 (5.6) [2]
Depersonalisation/derealisation disorder	0	2 (5.9) [2]	2 (5.6) [2]
Dysaesthesia	0	2 (5.9) [2]	2 (5.6) [2]
Feeling cold	0	2 (5.9) [2]	2 (5.6) [2]
Feeling hot	0	2 (5.9) [2]	2 (5.6) [2]
Headache	0	2 (5.9) [2]	2 (5.6) [2]
Hyperacusis	0	2 (5.9) [2]	2 (5.6) [2]
Hyperhidrosis	0	2 (5.9) [2]	2 (5.6) [2]
Inappropriate affect	0	2 (5.9) [2]	2 (5.6) [2]
Psychomotor retardation	0	2 (5.9) [2]	2 (5.6) [2]
Acute stress disorder	0	1 (2.9) [1]	1 (2.8) [1]
Analgesic therapy	0	1 (2.9) [1]	1 (2.8) [1]
Anxiety	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT.n is the number of participants with the measurement available.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030114.sas FINAL (2026-08-28,12:55)

Table 14.3.1.14
Treatment-Emergent Adverse Events by MedDRA Preferred Term
Safety Population

Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Chest discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Cognitive disorder	0	1 (2.9) [1]	1 (2.8) [1]
Confusional state	0	1 (2.9) [1]	1 (2.8) [1]
Decreased appetite	0	1 (2.9) [1]	1 (2.8) [1]
Delusion	0	1 (2.9) [1]	1 (2.8) [1]
Depressed mood	0	1 (2.9) [1]	1 (2.8) [1]
Discomfort	0	1 (2.9) [1]	1 (2.8) [1]
Disinhibition	0	1 (2.9) [1]	1 (2.8) [1]
Dyskinesia	0	1 (2.9) [1]	1 (2.8) [1]
Dyspnoea	0	1 (2.9) [1]	1 (2.8) [1]
Fear	0	1 (2.9) [1]	1 (2.8) [1]
Fear of death	0	1 (2.9) [1]	1 (2.8) [1]
Flight of ideas	0	1 (2.9) [1]	1 (2.8) [1]
Gait disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, auditory	0	1 (2.9) [1]	1 (2.8) [1]
Hallucination, tactile	0	1 (2.9) [2]	1 (2.8) [2]
Hyperventilation	0	1 (2.9) [1]	1 (2.8) [1]
Hypotension	1 (2.8) [1]	0	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event.
E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT.n is the number of participants with the measurement available.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030114.sas FINAL (2026-08-28,12:55)

Table 14.3.1.14
Treatment-Emergent Adverse Events by MedDRA Preferred Term
Safety Population

Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Ingrown hair	0	1 (2.9) [1]	1 (2.8) [1]
Logorrhoea	0	1 (2.9) [1]	1 (2.8) [1]
Loose associations	0	1 (2.9) [1]	1 (2.8) [1]
Malaise	0	1 (2.9) [1]	1 (2.8) [1]
Nervousness	0	1 (2.9) [1]	1 (2.8) [1]
Pain	0	1 (2.9) [1]	1 (2.8) [1]
Parosmia	0	1 (2.9) [1]	1 (2.8) [1]
Perseveration	0	1 (2.9) [1]	1 (2.8) [1]
Psychomotor hyperactivity	0	1 (2.9) [1]	1 (2.8) [1]
Sensory disturbance	0	1 (2.9) [1]	1 (2.8) [1]
Sexually inappropriate behaviour	0	1 (2.9) [1]	1 (2.8) [1]
Somatic delusion	0	1 (2.9) [1]	1 (2.8) [1]
Thinking abnormal	0	1 (2.9) [1]	1 (2.8) [1]
Tremor	0	1 (2.9) [1]	1 (2.8) [1]
Urinary incontinence	0	1 (2.9) [1]	1 (2.8) [1]
Vessel puncture site pain	1 (2.8) [1]	0	1 (2.8) [1]
Visual processing disorder	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event.
E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT.n is the number of participants with the measurement available.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030114.sas FINAL (2026-08-28,12:55)

Table 14.3.1.15
Treatment-Emergent Adverse Events by MedDRA High Level Term
Safety Population

High Level Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Participants with any TEAE	4 (11.1) [4]	34 (100) [196]	34 (94.4) [200]
Hallucinations (excl sleep-related)	0	28 (82.4) [37]	28 (77.8) [37]
Perception disturbances NEC	0	15 (44.1) [18]	15 (41.7) [18]
Affect alterations NEC	0	14 (41.2) [17]	14 (38.9) [17]
Emotional and mood disturbances NEC	0	10 (29.4) [10]	10 (27.8) [10]
Feelings and sensations NEC	0	10 (29.4) [10]	10 (27.8) [10]
Neurological signs and symptoms NEC	0	9 (26.5) [9]	9 (25.0) [9]
Paraesthesias and dysaesthesias	0	8 (23.5) [10]	8 (22.2) [10]
Behaviour and socialisation disturbances	0	7 (20.6) [7]	7 (19.4) [7]
Nausea and vomiting symptoms	0	7 (20.6) [7]	7 (19.4) [7]
Anxiety symptoms	0	6 (17.6) [8]	6 (16.7) [8]
Dissociative states	0	6 (17.6) [8]	6 (16.7) [8]
Coordination and balance disturbances	0	5 (14.7) [5]	5 (13.9) [5]
Vascular hypotensive disorders	3 (8.3) [3]	3 (8.8) [3]	5 (13.9) [6]
Confusion and disorientation	0	4 (11.8) [5]	4 (11.1) [5]
Increased physical activity levels	0	3 (8.8) [3]	3 (8.3) [3]
Mood alterations with depressive symptoms	0	3 (8.8) [3]	3 (8.3) [3]
Pain and discomfort NEC	0	3 (8.8) [3]	3 (8.3) [3]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (High Level Term [HLT]) for a participant were counted once in the frequency for that adverse event.
E represents the number of adverse event occurrences; participants may contribute multiple events to E within each HLT.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030115.sas FINAL (2026-08-28,12:55)

Table 14.3.1.15
Treatment-Emergent Adverse Events by MedDRA High Level Term
Safety Population

High Level Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Thinking disturbances	0	3 (8.8) [4]	3 (8.3) [4]
Apocrine and eccrine gland disorders	0	2 (5.9) [2]	2 (5.6) [2]
Breathing abnormalities	0	2 (5.9) [2]	2 (5.6) [2]
Delusional symptoms	0	2 (5.9) [2]	2 (5.6) [2]
Dyskinesias and movement disorders NEC	0	2 (5.9) [2]	2 (5.6) [2]
Headaches NEC	0	2 (5.9) [2]	2 (5.6) [2]
Hyperacusia	0	2 (5.9) [2]	2 (5.6) [2]
Vascular tests NEC (incl blood pressure)	0	2 (5.9) [2]	2 (5.6) [2]
Administration site reactions NEC	1 (2.8) [1]	0	1 (2.8) [1]
Analgesia supportive care	0	1 (2.9) [1]	1 (2.8) [1]
Appetite disorders	0	1 (2.9) [1]	1 (2.8) [1]
Asthenic conditions	0	1 (2.9) [1]	1 (2.8) [1]
Bladder and urethral symptoms	0	1 (2.9) [1]	1 (2.8) [1]
Cortical dysfunction NEC	0	1 (2.9) [1]	1 (2.8) [1]
Fear symptoms and phobic disorders (incl social phobia)	0	1 (2.9) [2]	1 (2.8) [2]
Gait disturbances	0	1 (2.9) [1]	1 (2.8) [1]
Mental impairment (excl dementia and memory loss)	0	1 (2.9) [1]	1 (2.8) [1]
Olfactory nerve disorders	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (High Level Term [HLT]) for a participant were counted once in the frequency for that adverse event.
E represents the number of adverse event occurrences; participants may contribute multiple events to E within each HLT.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030115.sas FINAL (2026-08-28,12:55)

Table 14.3.1.15
Treatment-Emergent Adverse Events by MedDRA High Level Term
Safety Population

High Level Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]
Pilar disorders NEC	0	1 (2.9) [1]	1 (2.8) [1]
Sensory abnormalities NEC	0	1 (2.9) [1]	1 (2.8) [1]
Speech and language usage disturbances	0	1 (2.9) [1]	1 (2.8) [1]
Stress disorders	0	1 (2.9) [1]	1 (2.8) [1]
Tremor (excl congenital)	0	1 (2.9) [1]	1 (2.8) [1]

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (High Level Term [HLT]) for a participant were counted once in the frequency for that adverse event.
E represents the number of adverse event occurrences; participants may contribute multiple events to E within each HLT.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030115.sas FINAL (2026-08-28,12:55)

Table 14.3.2.1
Serious Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]

There is no observation in this table.

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030201.sas FINAL (2026-08-28,12:55)

Table 14.3.2.2
Serious Treatment-Emergent Adverse Events by MedDRA System Organ Class, Preferred Term and Intensity
Safety Population

System Organ Class/ Preferred Term/ Intensity	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]

There is no observation in this table.

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC. Each participant will be counted only once within a system organ class or a preferred term by using the event with the highest intensity within each classification for each treatment.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030202.sas FINAL (2026-08-28,12:55)

Table 14.3.2.3
Related Serious Treatment-Emergent Adverse Events by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]

There is no observation in this table.

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
Related TEAEs are defined as events reported as related or possibly related.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030203.sas FINAL (2026-08-28,12:55)

Table 14.3.2.4
Treatment-Emergent Adverse Events Leading to Study Discontinuation by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]

There is no observation in this table.

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030204.sas FINAL (2026-08-28,12:55)

Table 14.3.2.5
Treatment-Emergent Adverse Events Resulting in Death by MedDRA System Organ Class and Preferred Term
Safety Population

System Organ Class/ Preferred Term	CYP Probe N=36 n (%) [E]	CYB003 + CYP Probe N=34 n (%) [E]	Total N=36 n (%) [E]

There is no observation in this table.

CYP Probe refers to Period 1 and includes TEAEs occurring on or after Day 1 CYP Probe and prior to Day 8 dosing.
CYB003 + CYP Probe refers to Period 2 and includes TEAEs occurring with an onset on or after Day 8 CYP Probe + CYB003 dosing.
Multiple occurrences of a specific adverse event (Preferred Term [PT]) for a participant were counted once in the frequency for that adverse event. Likewise, multiple occurrences of adverse events within a specific System Organ Class (SOC) for a participant were counted once in the frequency for that SOC. E represents the number of adverse event occurrences; participants may contribute multiple events to E within each PT and SOC.
n is the number of participants with the measurement available.
The number of participants in the respective trial arms, N, is used as the denominator for percentages.
MedDRA version 27.1.
TEAE = treatment-emergent adverse event.

Source: Listing 16.2.7.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030205.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Alanine Aminotransferase (U/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	23.6 (15.34)	23.9 (13.43)
Median	20.0	20.0
Q1, Q3	15.0, 27.5	17.0, 29.0
Min, Max	9, 96	10, 82
Post-baseline Visit, n	36	34
Mean (SD)	22.7 (18.85)	20.8 (9.82)
Median	18.0	18.5
Q1, Q3	13.0, 24.5	15.0, 24.0
Min, Max	9, 116	9, 61
Change from baseline, n	36	34
Mean (SD)	-0.9 (5.76)	-3.1 (5.14)
Median	-2.5	-1.5
Q1, Q3	-4.0, 0.5	-4.0, 0.0
Min, Max	-8, 20	-21, 2

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Albumin (g/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	44.4 (2.84)	45.2 (2.39)
Median	45.0	45.0
Q1, Q3	42.5, 46.5	43.0, 47.0
Min, Max	38, 49	41, 50
Post-baseline Visit, n	36	34
Mean (SD)	42.8 (2.89)	43.9 (3.14)
Median	43.0	44.0
Q1, Q3	41.0, 45.0	43.0, 46.0
Min, Max	37, 48	36, 51
Change from baseline, n	36	34
Mean (SD)	-1.6 (2.55)	-1.3 (2.76)
Median	-2.0	-1.0
Q1, Q3	-3.5, 0.0	-3.0, 1.0
Min, Max	-6, 5	-9, 3

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Alkaline Phosphatase (U/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	70.3 (21.29)	73.3 (22.92)
Median	66.5	72.0
Q1, Q3	56.0, 79.5	59.0, 81.0
Min, Max	41, 131	40, 135
Post-baseline Visit, n	36	34
Mean (SD)	68.3 (22.41)	69.4 (22.50)
Median	64.5	66.0
Q1, Q3	53.0, 75.5	52.0, 76.0
Min, Max	38, 133	37, 145
Change from baseline, n	36	34
Mean (SD)	-1.9 (5.18)	-3.9 (6.96)
Median	-2.0	-3.0
Q1, Q3	-5.0, 1.0	-8.0, -1.0
Min, Max	-12, 11	-24, 10

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Amylase (U/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	86.4 (33.90)	85.0 (33.22)
Median	84.5	81.0
Q1, Q3	67.5, 100.5	58.0, 102.0
Min, Max	37, 195	37, 180
Post-baseline Visit, n	36	34
Mean (SD)	94.8 (36.48)	84.6 (30.25)
Median	95.5	84.5
Q1, Q3	71.5, 106.0	70.0, 104.0
Min, Max	41, 206	32, 148
Change from baseline, n	36	34
Mean (SD)	8.4 (12.32)	-0.4 (15.25)
Median	6.5	-1.5
Q1, Q3	-1.0, 15.5	-5.0, 6.0
Min, Max	-12, 41	-35, 41

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Aspartate Aminotransferase (U/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	23.1 (9.29)	22.9 (7.69)
Median	20.5	21.5
Q1, Q3	17.0, 26.5	17.0, 25.0
Min, Max	11, 56	13, 49
Post-baseline Visit, n	36	34
Mean (SD)	18.7 (7.26)	17.7 (4.45)
Median	17.5	18.0
Q1, Q3	14.0, 21.5	14.0, 19.0
Min, Max	9, 49	11, 28
Change from baseline, n	36	34
Mean (SD)	-4.4 (5.38)	-5.2 (5.40)
Median	-4.0	-4.5
Q1, Q3	-6.5, -1.0	-8.0, -2.0
Min, Max	-18, 8	-22, 5

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Bilirubin (umol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	10.50 (5.059)	9.76 (5.165)
Median	10.26	10.26
Q1, Q3	7.70, 13.68	6.84, 10.26
Min, Max	0.0, 23.9	0.0, 29.1
Post-baseline Visit, n	36	34
Mean (SD)	9.17 (4.627)	10.56 (4.563)
Median	7.70	8.55
Q1, Q3	5.99, 13.68	8.55, 11.97
Min, Max	0.0, 22.2	3.4, 23.9
Change from baseline, n	36	34
Mean (SD)	-1.33 (3.837)	0.80 (5.125)
Median	-1.71	0.86
Q1, Q3	-3.42, 1.71	-1.71, 3.42
Min, Max	-10.3, 6.8	-12.0, 13.7

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Calcium (mmol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	2.35 (0.059)	2.34 (0.074)
Median	2.35	2.34
Q1, Q3	2.33, 2.39	2.28, 2.38
Min, Max	2.2, 2.5	2.2, 2.5
Post-baseline Visit, n	36	34
Mean (SD)	2.31 (0.077)	2.33 (0.093)
Median	2.30	2.33
Q1, Q3	2.26, 2.35	2.28, 2.38
Min, Max	2.2, 2.6	2.1, 2.6
Change from baseline, n	36	34
Mean (SD)	-0.04 (0.066)	-0.01 (0.102)
Median	-0.03	0.00
Q1, Q3	-0.10, 0.00	-0.10, 0.08
Min, Max	-0.2, 0.1	-0.2, 0.2

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Chloride (mmol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	101.9 (2.45)	101.6 (2.16)
Median	102.0	102.0
Q1, Q3	101.0, 103.0	100.0, 103.0
Min, Max	95, 107	97, 106
Post-baseline Visit, n	36	34
Mean (SD)	102.3 (2.14)	101.6 (1.72)
Median	102.0	102.0
Q1, Q3	101.0, 103.0	100.0, 103.0
Min, Max	98, 108	98, 104
Change from baseline, n	36	34
Mean (SD)	0.4 (2.68)	0.0 (2.16)
Median	0.0	0.0
Q1, Q3	-1.0, 2.0	-2.0, 2.0
Min, Max	-5, 9	-4, 5

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Creatinine (umol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	90.98 (14.403)	88.45 (16.160)
Median	92.38	89.28
Q1, Q3	82.21, 100.33	82.21, 99.89
Min, Max	60.1, 115.8	52.2, 120.2
Post-baseline Visit, n	36	34
Mean (SD)	90.27 (14.028)	92.40 (16.009)
Median	91.05	94.15
Q1, Q3	80.44, 100.33	79.56, 104.31
Min, Max	60.1, 114.0	54.8, 118.5
Change from baseline, n	36	34
Mean (SD)	-0.71 (7.624)	3.95 (8.012)
Median	0.00	4.86
Q1, Q3	-6.19, 4.42	-0.88, 9.72
Min, Max	-23.9, 11.5	-17.7, 16.8

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Gamma Glutamyl Transferase (U/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	22.9 (17.80)	24.4 (19.97)
Median	18.0	17.0
Q1, Q3	12.5, 26.5	12.0, 28.0
Min, Max	6, 101	8, 110
Post-baseline Visit, n	36	34
Mean (SD)	23.4 (22.35)	23.6 (18.56)
Median	17.0	17.5
Q1, Q3	10.5, 26.0	11.0, 29.0
Min, Max	6, 132	6, 103
Change from baseline, n	36	34
Mean (SD)	0.4 (5.89)	-0.7 (3.29)
Median	0.0	-0.5
Q1, Q3	-2.0, 1.0	-2.0, 1.0
Min, Max	-6, 31	-10, 9

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Glucose (mmol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	5.16 (0.596)	5.05 (0.630)
Median	5.16	5.02
Q1, Q3	4.75, 5.55	4.66, 5.33
Min, Max	4.0, 6.9	3.9, 7.3
Post-baseline Visit, n	36	34
Mean (SD)	5.15 (0.467)	5.26 (0.591)
Median	5.08	5.19
Q1, Q3	4.88, 5.38	4.94, 5.44
Min, Max	4.2, 7.0	4.1, 7.7
Change from baseline, n	36	34
Mean (SD)	0.00 (0.462)	0.20 (0.475)
Median	0.06	0.17
Q1, Q3	-0.33, 0.33	-0.17, 0.50
Min, Max	-0.9, 0.9	-0.8, 1.2

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Lactate Dehydrogenase (U/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	173.3 (42.56)	171.5 (36.59)
Median	167.0	159.5
Q1, Q3	144.0, 185.5	147.0, 194.0
Min, Max	101, 300	104, 262
Post-baseline Visit, n	36	34
Mean (SD)	154.4 (36.24)	161.8 (36.62)
Median	147.5	155.5
Q1, Q3	130.5, 175.0	127.0, 189.0
Min, Max	90, 266	106, 237
Change from baseline, n	36	34
Mean (SD)	-18.8 (20.03)	-9.7 (26.82)
Median	-17.0	-11.0
Q1, Q3	-32.5, -3.0	-32.0, -2.0
Min, Max	-73, 15	-43, 72

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Phosphate (mmol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	1.12 (0.157)	1.12 (0.128)
Median	1.11	1.15
Q1, Q3	1.00, 1.23	1.03, 1.20
Min, Max	0.8, 1.5	0.8, 1.4
Post-baseline Visit, n	36	34
Mean (SD)	1.11 (0.170)	1.08 (0.194)
Median	1.11	1.03
Q1, Q3	1.00, 1.26	0.97, 1.23
Min, Max	0.8, 1.5	0.8, 1.6
Change from baseline, n	36	34
Mean (SD)	-0.01 (0.121)	-0.04 (0.180)
Median	0.00	-0.05
Q1, Q3	-0.10, 0.03	-0.13, 0.06
Min, Max	-0.3, 0.3	-0.4, 0.4

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Potassium (mmol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	4.53 (0.360)	4.50 (0.411)
Median	4.50	4.40
Q1, Q3	4.25, 4.75	4.30, 4.60
Min, Max	3.8, 5.7	3.8, 5.8
Post-baseline Visit, n	36	34
Mean (SD)	4.64 (0.331)	4.52 (0.324)
Median	4.60	4.50
Q1, Q3	4.40, 4.80	4.30, 4.70
Min, Max	4.1, 5.9	4.0, 5.3
Change from baseline, n	36	34
Mean (SD)	0.11 (0.313)	0.02 (0.446)
Median	0.10	0.05
Q1, Q3	-0.10, 0.35	-0.10, 0.30
Min, Max	-0.7, 0.7	-1.1, 1.0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Protein (g/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	70.0 (4.99)	69.7 (4.00)
Median	71.0	69.5
Q1, Q3	69.0, 73.0	67.0, 73.0
Min, Max	57, 78	62, 78
Post-baseline Visit, n	36	34
Mean (SD)	67.5 (5.00)	68.0 (5.60)
Median	68.0	68.0
Q1, Q3	65.0, 71.0	65.0, 72.0
Min, Max	55, 76	57, 81
Change from baseline, n	36	34
Mean (SD)	-2.5 (3.31)	-1.7 (5.22)
Median	-3.0	-2.0
Q1, Q3	-5.0, 0.0	-6.0, 2.0
Min, Max	-7, 5	-12, 10

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Sodium (mmol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	139.3 (1.95)	138.8 (1.90)
Median	140.0	139.0
Q1, Q3	138.0, 140.5	138.0, 140.0
Min, Max	135, 143	135, 144
Post-baseline Visit, n	36	34
Mean (SD)	139.7 (1.80)	139.6 (1.46)
Median	140.0	140.0
Q1, Q3	138.5, 141.0	139.0, 141.0
Min, Max	136, 143	136, 142
Change from baseline, n	36	34
Mean (SD)	0.4 (2.27)	0.7 (1.71)
Median	0.0	1.0
Q1, Q3	-1.0, 2.0	-1.0, 2.0
Min, Max	-3, 7	-2, 5

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Urate (umol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	312.27 (64.275)	326.44 (60.901)
Median	312.27	330.11
Q1, Q3	273.61, 339.04	297.40, 368.78
Min, Max	178.4, 452.0	178.4, 463.9
Post-baseline Visit, n	36	34
Mean (SD)	325.98 (60.622)	324.52 (61.294)
Median	318.22	327.14
Q1, Q3	300.37, 353.91	285.50, 356.88
Min, Max	190.3, 463.9	184.4, 463.9
Change from baseline, n	36	34
Mean (SD)	13.71 (37.587)	-1.92 (44.335)
Median	11.90	-5.95
Q1, Q3	-8.92, 41.64	-23.79, 29.74
Min, Max	-71.4, 89.2	-119.0, 101.1

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.1
Chemistry Test Results by Visit
Safety Population

Urea Nitrogen (mmol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	4.67 (1.127)	4.89 (1.293)
Median	4.64	5.18
Q1, Q3	4.11, 5.36	4.28, 5.71
Min, Max	1.8, 6.8	2.1, 6.8
Post-baseline Visit, n	36	34
Mean (SD)	5.48 (1.347)	5.03 (1.458)
Median	5.71	5.00
Q1, Q3	4.82, 6.43	3.93, 6.07
Min, Max	2.5, 7.5	2.5, 8.2
Change from baseline, n	36	34
Mean (SD)	0.81 (1.124)	0.14 (1.307)
Median	0.71	-0.18
Q1, Q3	0.00, 1.43	-1.07, 0.71
Min, Max	-1.4, 3.6	-1.8, 3.6

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030401.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Basophils (10^9/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	0.04 (0.014)	0.03 (0.014)
Median	0.03	0.03
Q1, Q3	0.03, 0.04	0.02, 0.05
Min, Max	0.0, 0.1	0.0, 0.1
Post-baseline Visit, n	36	33
Mean (SD)	0.03 (0.013)	0.03 (0.013)
Median	0.03	0.03
Q1, Q3	0.03, 0.04	0.03, 0.04
Min, Max	0.0, 0.1	0.0, 0.1
Change from baseline, n	36	33
Mean (SD)	0.00 (0.009)	0.00 (0.010)
Median	0.00	0.00
Q1, Q3	-0.01, 0.00	-0.01, 0.01
Min, Max	0.0, 0.0	0.0, 0.0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Basophils/Leukocytes (%)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	0.68 (0.342)	0.63 (0.283)
Median	0.60	0.60
Q1, Q3	0.50, 0.75	0.40, 0.80
Min, Max	0.1, 1.9	0.2, 1.5
Post-baseline Visit, n	36	33
Mean (SD)	0.57 (0.244)	0.55 (0.224)
Median	0.50	0.50
Q1, Q3	0.40, 0.60	0.40, 0.70
Min, Max	0.2, 1.5	0.2, 1.2
Change from baseline, n	36	33
Mean (SD)	-0.11 (0.225)	-0.07 (0.175)
Median	-0.10	-0.10
Q1, Q3	-0.20, 0.00	-0.10, 0.00
Min, Max	-0.8, 0.3	-0.4, 0.3

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Eosinophils (10^9/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	0.13 (0.083)	0.13 (0.084)
Median	0.12	0.11
Q1, Q3	0.08, 0.14	0.07, 0.16
Min, Max	0.0, 0.4	0.0, 0.4
Post-baseline Visit, n	36	33
Mean (SD)	0.15 (0.090)	0.12 (0.082)
Median	0.13	0.09
Q1, Q3	0.10, 0.18	0.06, 0.19
Min, Max	0.0, 0.4	0.0, 0.3
Change from baseline, n	36	33
Mean (SD)	0.03 (0.055)	-0.01 (0.046)
Median	0.02	-0.01
Q1, Q3	0.00, 0.04	-0.02, 0.01
Min, Max	-0.1, 0.2	-0.1, 0.2

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Eosinophils/Leukocytes (%)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	2.36 (1.423)	2.40 (1.521)
Median	2.05	2.10
Q1, Q3	1.35, 3.15	1.20, 3.30
Min, Max	0.3, 6.9	0.7, 6.4
Post-baseline Visit, n	36	33
Mean (SD)	2.66 (1.445)	1.99 (1.427)
Median	2.35	1.60
Q1, Q3	1.60, 3.50	1.00, 2.60
Min, Max	0.4, 6.1	0.3, 5.3
Change from baseline, n	36	33
Mean (SD)	0.30 (0.833)	-0.38 (0.800)
Median	0.20	-0.50
Q1, Q3	-0.20, 0.70	-0.80, -0.10
Min, Max	-1.3, 3.2	-1.6, 2.3

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Ery. Mean Corpuscular HGB Concentration (g/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	331.3 (11.17)	331.3 (11.50)
Median	333.5	333.0
Q1, Q3	325.0, 340.0	324.0, 338.0
Min, Max	306, 347	306, 352
Post-baseline Visit, n	36	33
Mean (SD)	328.9 (10.70)	329.5 (11.11)
Median	329.0	330.0
Q1, Q3	322.5, 333.5	322.0, 336.0
Min, Max	311, 356	308, 355
Change from baseline, n	36	33
Mean (SD)	-2.3 (7.43)	-2.1 (8.79)
Median	-3.0	-4.0
Q1, Q3	-6.5, 3.0	-7.0, 4.0
Min, Max	-19, 12	-20, 18

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Ery. Mean Corpuscular Hemoglobin (pg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	29.76 (1.968)	29.86 (1.943)
Median	29.75	29.80
Q1, Q3	28.40, 31.30	29.10, 31.30
Min, Max	25.2, 33.7	25.0, 33.7
Post-baseline Visit, n	36	33
Mean (SD)	29.69 (1.824)	29.85 (1.937)
Median	29.55	29.90
Q1, Q3	28.10, 31.20	28.70, 31.30
Min, Max	25.6, 33.3	25.3, 33.4
Change from baseline, n	36	33
Mean (SD)	-0.07 (0.481)	-0.02 (0.572)
Median	-0.10	0.00
Q1, Q3	-0.45, 0.30	-0.30, 0.30
Min, Max	-1.0, 1.0	-1.4, 1.2

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Ery. Mean Corpuscular Volume (fL)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	89.79 (4.280)	90.11 (4.000)
Median	89.60	89.55
Q1, Q3	86.35, 91.20	87.50, 92.50
Min, Max	82.1, 99.4	81.6, 99.0
Post-baseline Visit, n	36	33
Mean (SD)	90.23 (4.448)	90.60 (4.413)
Median	89.95	90.10
Q1, Q3	87.65, 92.00	88.00, 92.70
Min, Max	81.6, 100.7	79.7, 101.6
Change from baseline, n	36	33
Mean (SD)	0.44 (1.644)	0.50 (1.607)
Median	0.40	0.40
Q1, Q3	-0.30, 1.35	-0.70, 1.00
Min, Max	-4.7, 3.3	-1.9, 5.1

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Erythrocytes (10^6/uL)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	5.00 (0.451)	4.84 (0.415)
Median	4.91	4.84
Q1, Q3	4.66, 5.30	4.55, 5.11
Min, Max	4.2, 6.0	4.0, 5.6
Post-baseline Visit, n	36	33
Mean (SD)	4.98 (0.399)	4.81 (0.370)
Median	4.97	4.82
Q1, Q3	4.66, 5.28	4.55, 5.13
Min, Max	4.2, 5.8	4.2, 5.5
Change from baseline, n	36	33
Mean (SD)	-0.02 (0.232)	-0.02 (0.236)
Median	-0.09	-0.01
Q1, Q3	-0.17, 0.14	-0.14, 0.14
Min, Max	-0.6, 0.6	-0.7, 0.4

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Hematocrit (L/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	0.45 (0.033)	0.44 (0.031)
Median	0.45	0.43
Q1, Q3	0.43, 0.46	0.41, 0.46
Min, Max	0.4, 0.5	0.4, 0.5
Post-baseline Visit, n	36	33
Mean (SD)	0.45 (0.031)	0.44 (0.031)
Median	0.45	0.43
Q1, Q3	0.42, 0.46	0.42, 0.46
Min, Max	0.4, 0.5	0.4, 0.5
Change from baseline, n	36	33
Mean (SD)	0.00 (0.022)	0.00 (0.025)
Median	0.00	0.00
Q1, Q3	-0.02, 0.02	-0.02, 0.02
Min, Max	0.0, 0.1	-0.1, 0.1

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Hemoglobin (mmol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	9.21 (0.714)	8.94 (0.677)
Median	9.28	8.94
Q1, Q3	8.63, 9.59	8.32, 9.31
Min, Max	7.9, 10.9	7.8, 10.4
Post-baseline Visit, n	36	33
Mean (SD)	9.16 (0.681)	8.90 (0.688)
Median	9.15	8.94
Q1, Q3	8.72, 9.53	8.38, 9.50
Min, Max	7.8, 10.6	7.5, 10.2
Change from baseline, n	36	33
Mean (SD)	-0.06 (0.393)	-0.04 (0.464)
Median	-0.12	-0.06
Q1, Q3	-0.37, 0.12	-0.25, 0.19
Min, Max	-0.9, 1.1	-1.2, 0.9

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Leukocyte Cell Differential (10^9/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	5.62 (1.366)	5.46 (1.436)
Median	5.75	5.26
Q1, Q3	4.47, 6.55	4.35, 6.50
Min, Max	3.3, 8.5	3.2, 8.6
Post-baseline Visit, n	36	33
Mean (SD)	6.12 (2.587)	6.35 (1.481)
Median	5.69	5.92
Q1, Q3	4.76, 6.54	5.42, 7.26
Min, Max	3.3, 18.5	3.8, 9.6
Change from baseline, n	36	33
Mean (SD)	0.50 (2.140)	0.83 (0.966)
Median	0.16	0.99
Q1, Q3	-0.31, 0.69	0.49, 1.34
Min, Max	-1.2, 12.2	-1.5, 3.1

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Lymphocytes (10^9/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	1.77 (0.470)	1.82 (0.457)
Median	1.71	1.78
Q1, Q3	1.45, 1.97	1.57, 2.01
Min, Max	1.0, 2.9	1.0, 3.4
Post-baseline Visit, n	36	33
Mean (SD)	1.89 (0.640)	1.92 (0.542)
Median	1.86	1.78
Q1, Q3	1.47, 2.18	1.60, 2.20
Min, Max	0.7, 4.1	1.1, 4.0
Change from baseline, n	36	33
Mean (SD)	0.11 (0.392)	0.09 (0.273)
Median	0.10	0.08
Q1, Q3	-0.06, 0.27	-0.07, 0.26
Min, Max	-0.7, 1.2	-0.5, 0.7

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Lymphocytes/Leukocytes (%)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	33.01 (10.053)	34.70 (9.014)
Median	32.30	35.85
Q1, Q3	25.95, 38.60	26.40, 42.20
Min, Max	15.8, 59.0	19.1, 51.7
Post-baseline Visit, n	36	33
Mean (SD)	32.99 (9.755)	30.84 (7.380)
Median	33.60	30.50
Q1, Q3	27.20, 37.40	25.70, 34.10
Min, Max	3.9, 57.4	16.4, 47.9
Change from baseline, n	36	33
Mean (SD)	-0.02 (6.083)	-3.63 (5.573)
Median	0.35	-3.80
Q1, Q3	-3.50, 3.95	-6.80, 0.00
Min, Max	-17.2, 15.6	-14.3, 7.8

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Monocytes (10^9/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	0.45 (0.095)	0.47 (0.121)
Median	0.46	0.48
Q1, Q3	0.39, 0.50	0.36, 0.54
Min, Max	0.3, 0.7	0.2, 0.8
Post-baseline Visit, n	36	33
Mean (SD)	0.50 (0.177)	0.54 (0.123)
Median	0.46	0.53
Q1, Q3	0.40, 0.55	0.48, 0.61
Min, Max	0.2, 1.1	0.3, 0.9
Change from baseline, n	36	33
Mean (SD)	0.04 (0.152)	0.06 (0.082)
Median	0.04	0.07
Q1, Q3	-0.03, 0.08	0.00, 0.12
Min, Max	-0.2, 0.6	-0.1, 0.2

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Monocytes/Leukocytes (%)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	8.36 (2.049)	8.80 (2.192)
Median	8.00	8.30
Q1, Q3	7.25, 9.20	7.50, 9.50
Min, Max	4.5, 13.3	4.9, 15.5
Post-baseline Visit, n	36	33
Mean (SD)	8.44 (2.324)	8.61 (1.842)
Median	7.80	8.40
Q1, Q3	6.80, 9.75	7.40, 9.60
Min, Max	5.6, 16.7	5.8, 13.9
Change from baseline, n	36	33
Mean (SD)	0.08 (1.944)	-0.21 (1.348)
Median	0.00	-0.20
Q1, Q3	-1.10, 0.65	-1.00, 0.50
Min, Max	-3.5, 7.7	-2.7, 3.3

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Neutrophils/Leukocytes (%)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	55.34 (11.524)	53.20 (9.917)
Median	56.30	51.70
Q1, Q3	48.80, 63.40	45.10, 61.50
Min, Max	29.8, 73.7	34.1, 70.3
Post-baseline Visit, n	36	33
Mean (SD)	55.08 (10.544)	57.71 (8.397)
Median	55.50	57.20
Q1, Q3	49.80, 59.65	52.50, 63.40
Min, Max	31.2, 89.2	42.8, 76.3
Change from baseline, n	36	33
Mean (SD)	-0.26 (7.454)	4.26 (6.636)
Median	-0.35	4.50
Q1, Q3	-5.10, 3.95	-0.30, 9.10
Min, Max	-19.5, 20.1	-8.0, 16.4

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.2
Hematology Test Results by Visit
Safety Population

Platelets (10^9/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	249.8 (43.63)	252.3 (47.33)
Median	245.5	246.0
Q1, Q3	219.0, 288.0	214.0, 297.0
Min, Max	171, 335	172, 350
Post-baseline Visit, n	36	32
Mean (SD)	239.8 (43.55)	247.5 (42.26)
Median	240.0	239.5
Q1, Q3	205.5, 264.0	212.0, 281.0
Min, Max	161, 329	176, 346
Change from baseline, n	36	32
Mean (SD)	-10.0 (16.73)	-8.9 (24.57)
Median	-10.0	-7.0
Q1, Q3	-19.0, 5.5	-22.5, 7.5
Min, Max	-47, 26	-93, 32

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030402.sas FINAL (2026-08-28,12:56)

Table 14.3.4.3
Urinalysis Test Results by Visit
Safety Population

Ketones (mmol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	0	3
Mean (SD)		0.10 (0.000)
Median		0.10
Q1, Q3		0.10, 0.10
Min, Max		0.1, 0.1

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030403.sas FINAL (2026-08-28,12:56)

Table 14.3.4.3
Urinalysis Test Results by Visit
Safety Population

Specific Gravity

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	1.02 (0.009)	1.02 (0.010)
Median	1.02	1.02
Q1, Q3	1.01, 1.02	1.01, 1.03
Min, Max	1.0, 1.0	1.0, 1.0
Post-baseline Visit, n	36	34
Mean (SD)	1.02 (0.006)	1.02 (0.007)
Median	1.02	1.02
Q1, Q3	1.01, 1.02	1.01, 1.02
Min, Max	1.0, 1.0	1.0, 1.0
Change from baseline, n	36	34
Mean (SD)	0.00 (0.009)	0.00 (0.009)
Median	0.00	0.00
Q1, Q3	-0.01, 0.01	-0.01, 0.01
Min, Max	0.0, 0.0	0.0, 0.0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030403.sas FINAL (2026-08-28,12:56)

Table 14.3.4.3
Urinalysis Test Results by Visit
Safety Population

Urine Protein (g/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	3	3
Mean (SD)	0.01 (0.006)	0.01 (0.006)
Median	0.01	0.01
Q1, Q3	0.01, 0.02	0.01, 0.02
Min, Max	0.0, 0.0	0.0, 0.0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030403.sas FINAL (2026-08-28,12:56)

Table 14.3.4.3
Urinalysis Test Results by Visit
Safety Population

Urobilinogen (umol/L)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	5.32 (4.798)	7.85 (6.496)
Median	3.42	3.42
Q1, Q3	3.42, 3.42	3.42, 17.10
Min, Max	3.4, 17.1	3.4, 17.1
Post-baseline Visit, n	36	34
Mean (SD)	5.70 (5.171)	7.04 (6.126)
Median	3.42	3.42
Q1, Q3	3.42, 3.42	3.42, 17.10
Min, Max	3.4, 17.1	3.4, 17.1
Change from baseline, n	36	34
Mean (SD)	0.38 (6.106)	-0.80 (4.692)
Median	0.00	0.00
Q1, Q3	0.00, 0.00	0.00, 0.00
Min, Max	-13.7, 13.7	-13.7, 13.7

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030403.sas FINAL (2026-08-28,12:56)

Table 14.3.4.3
Urinalysis Test Results by Visit
Safety Population

pH

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline, n	36	34
Mean (SD)	6.57 (0.829)	6.37 (0.810)
Median	6.50	6.00
Q1, Q3	6.00, 7.00	5.50, 7.00
Min, Max	5.5, 9.0	5.5, 9.0
Post-baseline Visit, n	36	34
Mean (SD)	6.22 (0.603)	6.38 (0.628)
Median	6.00	6.50
Q1, Q3	6.00, 6.50	6.00, 6.50
Min, Max	5.5, 8.5	5.5, 8.0
Change from baseline, n	36	34
Mean (SD)	-0.35 (0.674)	0.01 (0.764)
Median	-0.50	0.00
Q1, Q3	-0.75, 0.00	-0.50, 0.00
Min, Max	-1.5, 1.0	-1.0, 2.0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.3 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030403.sas FINAL (2026-08-28,12:56)

Table 14.3.4.4
Thyroid-Stimulating Hormone Test Results by Visit
Safety Population

	Total N=36
Baseline, n	36
Mean (SD)	1.52 (0.710)
Median	1.36
Q1, Q3	1.01, 1.89
Min, Max	0.5, 3.6
Post-baseline Visit, n	30
Mean (SD)	2.27 (0.909)
Median	2.09
Q1, Q3	1.58, 2.76
Min, Max	1.0, 4.5
Change from baseline, n	30
Mean (SD)	0.80 (0.798)
Median	0.62
Q1, Q3	0.32, 1.28
Min, Max	-1.1, 2.8

Baseline refers to Screening and Post-baseline refers to Day 10.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.8.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030404.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Alanine Aminotransferase (U/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	33 (100)	0	0	33 (100)	0
	High	0	0	1 (33.3)	2 (66.7)	0	3 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	34 (94.4)	2 (5.6)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	31 (96.9)	1 (3.1)	0	32 (100)	0
	High	0	0	1 (50.0)	1 (50.0)	0	2 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	32 (94.1)	2 (5.9)	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Albumin (g/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	36 (100)	0	0	36 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	36 (100)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	34 (100)	0	0	34 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	34 (100)	0	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Alkaline Phosphatase (U/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	2 (5.6)	34 (94.4)	0	0	36 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	2 (5.6)	34 (94.4)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	1 (2.9)	33 (97.1)	0	0	34 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	1 (2.9)	33 (97.1)	0	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Amylase (U/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	22 (81.5)	5 (18.5)	0	27 (100)	0
	High	0	0	1 (11.1)	8 (88.9)	0	9 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	23 (63.9)	13 (36.1)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	23 (95.8)	1 (4.2)	0	24 (100)	0
	High	0	0	2 (20.0)	8 (80.0)	0	10 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	25 (73.5)	9 (26.5)	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Aspartate Aminotransferase (U/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	33 (100)	0	0	33 (100)	0
	High	0	0	2 (66.7)	1 (33.3)	0	3 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	35 (97.2)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	33 (100)	0	0	33 (100)	0
	High	0	0	1 (100)	0	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	34 (100)	0	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Bilirubin (umol/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	34 (100)	0	0	34 (100)	0
	High	0	0	1 (50.0)	1 (50.0)	0	2 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	35 (97.2)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	31 (96.9)	1 (3.1)	0	32 (100)	0
	High	0	0	1 (50.0)	1 (50.0)	0	2 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	32 (94.1)	2 (5.9)	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Calcium (mmol/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	36 (100)	0	0	36 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	36 (100)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	1 (2.9)	33 (97.1)	0	0	34 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	1 (2.9)	33 (97.1)	0	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Chloride (mmol/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	1 (100)	0	0	1 (100)	0
	Normal	0	0	35 (100)	0	0	35 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	36 (100)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	34 (100)	0	0	34 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	34 (100)	0	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Creatinine (umol/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	36 (100)	0	0	36 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	36 (100)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	1 (100)	0	0	0	1 (100)	0
	Normal	0	0	32 (100)	0	0	32 (100)	0
	High	0	0	0	1 (100)	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	1 (2.9)	32 (94.1)	1 (2.9)	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Gamma Glutamyl Transferase (U/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	4 (80.0)	1 (20.0)	0	0	5 (100)	0
	Normal	0	3 (10.0)	27 (90.0)	0	0	30 (100)	0
	High	0	0	0	1 (100)	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	7 (19.4)	28 (77.8)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	1 (33.3)	2 (66.7)	0	0	3 (100)	0
	Normal	0	3 (10.0)	27 (90.0)	0	0	30 (100)	0
	High	0	0	0	1 (100)	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	4 (11.8)	29 (85.3)	1 (2.9)	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Glucose (mmol/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	24 (92.3)	2 (7.7)	0	26 (100)	0
	High	0	0	7 (70.0)	3 (30.0)	0	10 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	31 (86.1)	5 (13.9)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	23 (79.3)	6 (20.7)	0	29 (100)	0
	High	0	0	4 (80.0)	1 (20.0)	0	5 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	27 (79.4)	7 (20.6)	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Lactate Dehydrogenase (U/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	6 (100)	0	0	0	6 (100)	0
	Normal	0	8 (30.8)	17 (65.4)	1 (3.8)	0	26 (100)	0
	High	0	0	3 (75.0)	1 (25.0)	0	4 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	14 (38.9)	20 (55.6)	2 (5.6)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	3 (100)	0	0	0	3 (100)	0
	Normal	0	7 (26.9)	18 (69.2)	1 (3.8)	0	26 (100)	0
	High	0	0	4 (80.0)	1 (20.0)	0	5 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	10 (29.4)	22 (64.7)	2 (5.9)	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Phosphate (mmol/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	1 (100)	0	0	0	1 (100)	0
	Normal	0	2 (5.9)	31 (91.2)	1 (2.9)	0	34 (100)	0
	High	0	0	1 (100)	0	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	3 (8.3)	32 (88.9)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	1 (50.0)	1 (50.0)	0	0	2 (100)	0
	Normal	0	2 (6.3)	29 (90.6)	1 (3.1)	0	32 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	3 (8.8)	30 (88.2)	1 (2.9)	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Potassium (mmol/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	35 (100)	0	0	35 (100)	0
	High	0	0	0	1 (100)	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	35 (97.2)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	33 (100)	0	0	33 (100)	0
	High	0	0	1 (100)	0	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	34 (100)	0	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Protein (g/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	1 (50.0)	1 (50.0)	0	0	2 (100)	0
	Normal	0	2 (5.9)	32 (94.1)	0	0	34 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	3 (8.3)	33 (91.7)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	1 (2.9)	33 (97.1)	0	0	34 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	1 (2.9)	33 (97.1)	0	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Sodium (mmol/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	36 (100)	0	0	36 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	36 (100)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	34 (100)	0	0	34 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	34 (100)	0	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Urate (umol/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	2 (100)	0	0	0	2 (100)	0
	Normal	0	0	34 (100)	0	0	34 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	2 (5.6)	34 (94.4)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	1 (100)	0	0	1 (100)	0
	Normal	0	1 (3.0)	32 (97.0)	0	0	33 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	1 (2.9)	33 (97.1)	0	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.5
Chemistry Results: Shift Table from Baseline to Post-Baseline
Safety Population

Urea Nitrogen (mmol/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	1 (100)	0	0	1 (100)	0
	Normal	0	0	34 (97.1)	1 (2.9)	0	35 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	35 (97.2)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	33 (97.1)	1 (2.9)	0	34 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	33 (97.1)	1 (2.9)	0	34 (100)	0
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030405.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Basophils (10^9/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	36 (100)	0	0	36 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	36 (100)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	33 (100)	0	0	33 (100)	1
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	33 (100)	0	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Basophils/Leukocytes (%)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	32 (100)	0	0	32 (100)	0
	High	0	0	2 (50.0)	2 (50.0)	0	4 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	34 (94.4)	2 (5.6)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	31 (100)	0	0	31 (100)	1
	High	0	0	1 (50.0)	1 (50.0)	0	2 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	32 (97.0)	1 (3.0)	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Eosinophils (10^9/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	36 (100)	0	0	36 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	36 (100)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	32 (100)	0	0	32 (100)	1
	High	0	0	1 (100)	0	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	33 (100)	0	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Eosinophils/Leukocytes (%)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	35 (100)	0	0	35 (100)	0
	High	0	0	1 (100)	0	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	36 (100)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	32 (100)	0	0	32 (100)	1
	High	0	0	1 (100)	0	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	33 (100)	0	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Ery. Mean Corpuscular HGB Concentration (g/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	2 (50.0)	2 (50.0)	0	0	4 (100)	0
	Normal	0	2 (6.3)	29 (90.6)	1 (3.1)	0	32 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	4 (11.1)	31 (86.1)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	2 (50.0)	2 (50.0)	0	0	4 (100)	0
	Normal	0	2 (6.9)	26 (89.7)	1 (3.4)	0	29 (100)	1
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	4 (12.1)	28 (84.8)	1 (3.0)	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Ery. Mean Corpuscular Hemoglobin (pg)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	1 (100)	0	0	0	1 (100)	0
	Normal	0	0	34 (100)	0	0	34 (100)	0
	High	0	0	0	1 (100)	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	1 (2.8)	34 (94.4)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	1 (100)	0	0	0	1 (100)	0
	Normal	0	0	31 (100)	0	0	31 (100)	1
	High	0	0	0	1 (100)	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	1 (3.0)	31 (93.9)	1 (3.0)	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Ery. Mean Corpuscular Volume (fL)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	32 (91.4)	3 (8.6)	0	35 (100)	0
	High	0	0	0	1 (100)	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	32 (88.9)	4 (11.1)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	31 (96.9)	1 (3.1)	0	32 (100)	1
	High	0	0	0	1 (100)	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	31 (93.9)	2 (6.1)	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Erythrocytes (10^6/uL)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	34 (100)	0	0	34 (100)	0
	High	0	0	2 (100)	0	0	2 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	36 (100)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	33 (100)	0	0	33 (100)	1
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	33 (100)	0	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Hematocrit (L/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	32 (97.0)	1 (3.0)	0	33 (100)	0
	High	0	0	0	3 (100)	0	3 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	32 (88.9)	4 (11.1)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	32 (100)	0	0	32 (100)	1
	High	0	0	0	1 (100)	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	32 (97.0)	1 (3.0)	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Hemoglobin (mmol/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	34 (100)	0	0	34 (100)	0
	High	0	0	1 (50.0)	1 (50.0)	0	2 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	35 (97.2)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	33 (100)	0	0	33 (100)	1
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	33 (100)	0	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Leukocyte Cell Differential (10^9/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	2 (66.7)	1 (33.3)	0	0	3 (100)	0
	Normal	0	0	32 (97.0)	1 (3.0)	0	33 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	2 (5.6)	33 (91.7)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	3 (100)	0	0	3 (100)	0
	Normal	0	0	30 (100)	0	0	30 (100)	1
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	33 (100)	0	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Lymphocytes (10^9/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	1 (2.8)	34 (94.4)	1 (2.8)	0	36 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	1 (2.8)	34 (94.4)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	32 (100)	0	0	32 (100)	1
	High	0	0	0	1 (100)	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	32 (97.0)	1 (3.0)	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Lymphocytes/Leukocytes (%)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	1 (2.9)	33 (97.1)	0	0	34 (100)	0
	High	0	0	0	2 (100)	0	2 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	1 (2.8)	33 (91.7)	2 (5.6)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	33 (100)	0	0	33 (100)	1
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	33 (100)	0	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Monocytes (10^9/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	34 (94.4)	2 (5.6)	0	36 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	34 (94.4)	2 (5.6)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	33 (100)	0	0	33 (100)	1
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	33 (100)	0	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Monocytes/Leukocytes (%)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	34 (97.1)	1 (2.9)	0	35 (100)	0
	High	0	0	1 (100)	0	0	1 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	35 (97.2)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	31 (100)	0	0	31 (100)	1
	High	0	0	1 (50.0)	1 (50.0)	0	2 (100)	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	32 (97.0)	1 (3.0)	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Neutrophils/Leukocytes (%)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	2 (100)	0	0	0	2 (100)	0
	Normal	0	0	33 (97.1)	1 (2.9)	0	34 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	2 (5.6)	33 (91.7)	1 (2.8)	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	1 (100)	0	0	1 (100)	0
	Normal	0	0	31 (96.9)	1 (3.1)	0	32 (100)	1
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	32 (97.0)	1 (3.0)	0	33 (100)	1
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.6
Hematology Results: Shift Table from Baseline to Post-Baseline
Safety Population

Platelets (10^9/L)

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
CYP Probe N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	36 (100)	0	0	36 (100)	0
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	36 (100)	0	0	36 (100)	0
	Missing	0	0	0	0	0	0	0
CYB003 + CYP Probe N=34	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	32 (100)	0	0	32 (100)	2
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	32 (100)	0	0	32 (100)	2
	Missing	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period; post-baseline refers to Day 2 in Period 1 and Day 9 in Period 2.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030406.sas FINAL (2026-08-28,12:56)

Table 14.3.4.7
Thyroid-Stimulating Hormone Results: Shift Table from Baseline to Post-Baseline
Safety Population

		Post-Baseline						
Group	Baseline	Very Low n (%)	Low n (%)	Normal n (%)	High n (%)	Very High n (%)	Total n (%)	Missing n
Total N=36	Very Low	0	0	0	0	0	0	0
	Low	0	0	0	0	0	0	0
	Normal	0	0	29 (96.7)	1 (3.3)	0	30 (100)	6
	High	0	0	0	0	0	0	0
	Very High	0	0	0	0	0	0	0
	Total	0	0	29 (96.7)	1 (3.3)	0	30 (100)	6
	Missing	0	0	0	0	0	0	0

Baseline refers to Screening and Post-baseline refers to Day 10.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in both baseline and post-baseline visit; n1 is displayed as n in the Total column for the particular row.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).

Source: Listing 16.2.8.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030407.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Systolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	115.6 (13.02)	116.1 (11.02)
Median	114.0	111.5
Q1, Q3	106.5, 120.5	108.0, 125.0
Min, Max	94, 147	100, 138
1 hour post-dose, n	36	33
Mean (SD)	114.6 (10.20)	128.3 (12.96)
Median	117.0	126.0
Q1, Q3	106.5, 121.5	120.0, 132.0
Min, Max	90, 136	111, 165
Change from baseline, n	36	33
Mean (SD)	-1.0 (10.85)	12.0 (9.50)
Median	2.0	13.0
Q1, Q3	-5.0, 6.5	6.0, 19.0
Min, Max	-27, 18	-7, 33

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Systolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
2 hours post-dose, n	36	31
Mean (SD)	111.1 (9.01)	134.0 (16.34)
Median	110.5	133.0
Q1, Q3	105.5, 116.5	125.0, 141.0
Min, Max	91, 127	95, 164
Change from baseline, n	36	31
Mean (SD)	-4.4 (9.76)	17.9 (11.60)
Median	-4.0	17.0
Q1, Q3	-9.5, 1.5	11.0, 27.0
Min, Max	-36, 13	-8, 42
4 hours post-dose, n	36	30
Mean (SD)	110.8 (9.25)	128.7 (15.29)
Median	110.0	125.5
Q1, Q3	105.0, 118.0	120.0, 142.0
Min, Max	91, 128	94, 158
Change from baseline, n	36	30
Mean (SD)	-4.8 (9.95)	12.3 (11.63)
Median	-4.0	14.5
Q1, Q3	-10.0, 2.0	4.0, 17.0
Min, Max	-36, 13	-10, 38

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Systolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
6 hours post-dose, n	36	34
Mean (SD)	107.6 (10.05)	121.4 (13.78)
Median	107.5	119.5
Q1, Q3	100.0, 115.5	112.0, 133.0
Min, Max	81, 136	94, 150
Change from baseline, n	36	34
Mean (SD)	-8.0 (10.93)	5.3 (9.72)
Median	-7.0	5.5
Q1, Q3	-13.5, 0.0	-1.0, 12.0
Min, Max	-42, 9	-18, 25
8 hours post-dose, n	36	34
Mean (SD)	106.8 (9.70)	114.7 (12.60)
Median	107.0	113.5
Q1, Q3	99.5, 115.5	107.0, 120.0
Min, Max	86, 125	92, 150
Change from baseline, n	36	34
Mean (SD)	-8.8 (11.30)	-1.4 (10.48)
Median	-6.5	-2.0
Q1, Q3	-15.0, 0.5	-11.0, 7.0
Min, Max	-34, 10	-22, 17

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Systolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
12 hours post-dose, n	36	34
Mean (SD)	112.4 (12.05)	110.1 (10.29)
Median	110.0	108.0
Q1, Q3	104.5, 122.5	105.0, 113.0
Min, Max	91, 143	94, 140
Change from baseline, n	36	34
Mean (SD)	-3.1 (12.47)	-6.0 (7.78)
Median	-3.5	-3.5
Q1, Q3	-11.0, 6.5	-11.0, 0.0
Min, Max	-37, 19	-26, 10
24 hours post-dose, n	36	34
Mean (SD)	113.2 (9.81)	109.9 (9.84)
Median	112.5	108.5
Q1, Q3	106.0, 120.0	104.0, 116.0
Min, Max	94, 136	89, 134
Change from baseline, n	36	34
Mean (SD)	-2.4 (9.39)	-6.2 (9.62)
Median	-1.0	-4.5
Q1, Q3	-6.5, 3.0	-9.0, 0.0
Min, Max	-23, 13	-31, 14

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Diastolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	71.6 (7.79)	72.0 (7.53)
Median	71.0	71.5
Q1, Q3	65.5, 76.5	67.0, 78.0
Min, Max	57, 95	58, 89
1 hour post-dose, n	36	33
Mean (SD)	73.2 (6.22)	82.2 (8.43)
Median	74.5	80.0
Q1, Q3	68.0, 77.5	78.0, 86.0
Min, Max	60, 84	68, 103
Change from baseline, n	36	33
Mean (SD)	1.7 (6.90)	10.0 (6.56)
Median	2.0	11.0
Q1, Q3	-1.0, 6.5	5.0, 16.0
Min, Max	-18, 14	-4, 20

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Diastolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
2 hours post-dose, n	36	31
Mean (SD)	70.4 (6.59)	82.8 (9.14)
Median	70.5	83.0
Q1, Q3	67.0, 74.5	77.0, 90.0
Min, Max	53, 82	60, 101
Change from baseline, n	36	31
Mean (SD)	-1.2 (7.03)	10.9 (6.43)
Median	-0.5	11.0
Q1, Q3	-4.5, 3.5	6.0, 15.0
Min, Max	-21, 10	-2, 23
4 hours post-dose, n	36	30
Mean (SD)	70.8 (6.78)	79.1 (10.29)
Median	70.0	79.5
Q1, Q3	65.5, 76.0	74.0, 88.0
Min, Max	56, 84	57, 101
Change from baseline, n	36	30
Mean (SD)	-0.8 (6.52)	6.5 (7.99)
Median	-1.5	7.0
Q1, Q3	-4.0, 4.5	1.0, 12.0
Min, Max	-18, 11	-16, 19

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Diastolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
6 hours post-dose, n	36	34
Mean (SD)	64.6 (6.57)	73.6 (8.61)
Median	65.0	75.0
Q1, Q3	60.5, 68.0	67.0, 79.0
Min, Max	48, 82	52, 89
Change from baseline, n	36	34
Mean (SD)	-6.9 (7.12)	1.6 (5.91)
Median	-7.0	3.0
Q1, Q3	-11.0, -2.0	-2.0, 6.0
Min, Max	-28, 8	-14, 13
8 hours post-dose, n	36	34
Mean (SD)	63.1 (6.37)	69.6 (8.19)
Median	62.0	68.0
Q1, Q3	59.0, 67.5	64.0, 76.0
Min, Max	47, 76	52, 88
Change from baseline, n	36	34
Mean (SD)	-8.4 (7.91)	-2.4 (6.74)
Median	-7.5	0.0
Q1, Q3	-15.5, -2.5	-6.0, 3.0
Min, Max	-26, 4	-16, 7

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Diastolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
12 hours post-dose, n	36	34
Mean (SD)	65.6 (8.21)	65.1 (6.96)
Median	62.5	64.0
Q1, Q3	61.0, 70.5	61.0, 70.0
Min, Max	52, 88	52, 84
Change from baseline, n	36	34
Mean (SD)	-5.9 (7.99)	-6.8 (6.42)
Median	-5.5	-5.5
Q1, Q3	-11.0, -1.5	-10.0, -3.0
Min, Max	-22, 19	-23, 6
24 hours post-dose, n	36	34
Mean (SD)	68.9 (7.03)	66.1 (6.78)
Median	69.0	65.5
Q1, Q3	64.0, 74.5	62.0, 71.0
Min, Max	55, 87	52, 77
Change from baseline, n	36	34
Mean (SD)	-2.7 (5.89)	-5.9 (6.39)
Median	-1.0	-5.5
Q1, Q3	-6.0, 0.5	-11.0, -2.0
Min, Max	-20, 7	-23, 7

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Pulse Rate (beats/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	60.5 (9.51)	62.0 (8.21)
Median	58.5	61.5
Q1, Q3	53.5, 64.5	56.0, 67.0
Min, Max	43, 92	47, 76
1 hour post-dose, n	36	33
Mean (SD)	54.7 (7.79)	61.4 (9.44)
Median	52.0	60.0
Q1, Q3	50.0, 59.0	56.0, 67.0
Min, Max	41, 78	38, 83
Change from baseline, n	36	33
Mean (SD)	-5.8 (10.38)	-1.1 (9.22)
Median	-5.0	-1.0
Q1, Q3	-9.0, -1.0	-8.0, 4.0
Min, Max	-33, 20	-16, 22

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Pulse Rate (beats/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
2 hours post-dose, n	36	31
Mean (SD)	53.9 (6.53)	65.2 (8.72)
Median	52.5	65.0
Q1, Q3	50.5, 57.0	58.0, 73.0
Min, Max	41, 73	48, 82
Change from baseline, n	36	31
Mean (SD)	-6.6 (10.76)	2.7 (9.95)
Median	-5.0	2.0
Q1, Q3	-11.0, 1.0	-3.0, 12.0
Min, Max	-37, 9	-18, 21
4 hours post-dose, n	36	30
Mean (SD)	55.3 (6.50)	72.1 (11.39)
Median	54.0	72.0
Q1, Q3	50.0, 57.5	64.0, 80.0
Min, Max	43, 71	50, 98
Change from baseline, n	36	30
Mean (SD)	-5.3 (8.68)	8.8 (10.62)
Median	-3.0	7.5
Q1, Q3	-9.0, 1.0	2.0, 13.0
Min, Max	-31, 7	-16, 30

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Pulse Rate (beats/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
6 hours post-dose, n	36	34
Mean (SD)	63.3 (8.63)	71.8 (8.91)
Median	61.5	73.0
Q1, Q3	56.5, 68.5	66.0, 77.0
Min, Max	51, 96	53, 87
Change from baseline, n	36	34
Mean (SD)	2.8 (8.66)	9.8 (8.63)
Median	4.0	10.0
Q1, Q3	-1.5, 8.0	3.0, 14.0
Min, Max	-25, 18	-5, 35
8 hours post-dose, n	36	34
Mean (SD)	62.6 (9.13)	68.6 (9.10)
Median	64.0	68.0
Q1, Q3	56.0, 67.5	62.0, 74.0
Min, Max	46, 94	47, 93
Change from baseline, n	36	34
Mean (SD)	2.1 (9.54)	6.6 (10.05)
Median	5.0	6.0
Q1, Q3	-3.5, 8.0	1.0, 13.0
Min, Max	-25, 16	-15, 35

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Supine Pulse Rate (beats/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
12 hours post-dose, n	36	34
Mean (SD)	63.9 (9.10)	67.6 (6.35)
Median	63.0	68.5
Q1, Q3	58.0, 72.0	63.0, 72.0
Min, Max	44, 80	53, 78
Change from baseline, n	36	34
Mean (SD)	3.4 (9.58)	5.6 (7.54)
Median	4.0	5.5
Q1, Q3	0.5, 8.0	2.0, 9.0
Min, Max	-32, 22	-11, 26
24 hours post-dose, n	36	34
Mean (SD)	60.8 (6.99)	60.4 (7.60)
Median	60.0	61.0
Q1, Q3	55.0, 66.0	56.0, 64.0
Min, Max	47, 79	44, 83
Change from baseline, n	36	34
Mean (SD)	0.3 (6.69)	-1.6 (7.27)
Median	1.0	-2.0
Q1, Q3	-5.0, 6.0	-6.0, 4.0
Min, Max	-15, 13	-17, 14

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Respiratory Rate (breaths/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	15.1 (2.00)	14.3 (1.64)
Median	16.0	14.0
Q1, Q3	14.0, 16.0	14.0, 16.0
Min, Max	12, 18	12, 18
1 hour post-dose, n	36	33
Mean (SD)	15.1 (1.75)	14.7 (1.98)
Median	16.0	14.0
Q1, Q3	14.0, 16.0	14.0, 16.0
Min, Max	12, 18	12, 18
Change from baseline, n	36	33
Mean (SD)	0.0 (1.43)	0.3 (1.59)
Median	0.0	0.0
Q1, Q3	0.0, 0.0	0.0, 2.0
Min, Max	-6, 2	-2, 4

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Respiratory Rate (breaths/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
2 hours post-dose, n	36	31
Mean (SD)	15.0 (1.82)	15.0 (1.78)
Median	15.0	14.0
Q1, Q3	14.0, 16.0	14.0, 16.0
Min, Max	12, 18	12, 18
Change from baseline, n	36	31
Mean (SD)	-0.1 (1.26)	0.6 (1.48)
Median	0.0	0.0
Q1, Q3	0.0, 0.0	0.0, 2.0
Min, Max	-4, 2	-2, 4
4 hours post-dose, n	36	30
Mean (SD)	14.9 (1.75)	15.1 (2.01)
Median	14.0	16.0
Q1, Q3	14.0, 16.0	14.0, 16.0
Min, Max	12, 18	12, 18
Change from baseline, n	36	30
Mean (SD)	-0.2 (1.15)	0.7 (1.99)
Median	0.0	0.0
Q1, Q3	0.0, 0.0	0.0, 2.0
Min, Max	-4, 2	-4, 4

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Respiratory Rate (breaths/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
6 hours post-dose, n	36	34
Mean (SD)	15.1 (1.75)	14.6 (1.67)
Median	16.0	14.0
Q1, Q3	14.0, 16.0	14.0, 16.0
Min, Max	12, 18	12, 18
Change from baseline, n	36	34
Mean (SD)	0.0 (1.51)	0.3 (1.40)
Median	0.0	0.0
Q1, Q3	0.0, 0.0	0.0, 0.0
Min, Max	-4, 4	-2, 4
8 hours post-dose, n	36	34
Mean (SD)	14.5 (1.93)	14.9 (2.27)
Median	14.0	16.0
Q1, Q3	13.0, 16.0	12.0, 16.0
Min, Max	12, 18	12, 18
Change from baseline, n	36	34
Mean (SD)	-0.6 (2.23)	0.6 (2.34)
Median	0.0	0.0
Q1, Q3	-2.0, 0.0	-2.0, 2.0
Min, Max	-6, 4	-4, 4

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Respiratory Rate (breaths/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
12 hours post-dose, n	36	34
Mean (SD)	14.9 (1.94)	14.8 (1.98)
Median	15.0	14.0
Q1, Q3	14.0, 16.0	14.0, 16.0
Min, Max	10, 18	12, 18
Change from baseline, n	36	34
Mean (SD)	-0.2 (2.68)	0.5 (2.22)
Median	0.0	0.0
Q1, Q3	-2.0, 2.0	0.0, 2.0
Min, Max	-6, 6	-4, 4
24 hours post-dose, n	36	34
Mean (SD)	14.3 (1.69)	14.3 (1.92)
Median	14.0	14.0
Q1, Q3	14.0, 16.0	12.0, 16.0
Min, Max	12, 18	12, 18
Change from baseline, n	36	34
Mean (SD)	-0.8 (1.74)	0.0 (2.09)
Median	0.0	0.0
Q1, Q3	-2.0, 0.0	-2.0, 0.0
Min, Max	-4, 2	-6, 6

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Oral Temperature (C)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	36.67 (0.197)	36.76 (0.180)
Median	36.60	36.70
Q1, Q3	36.55, 36.75	36.60, 36.90
Min, Max	36.3, 37.2	36.5, 37.1
1 hour post-dose, n	36	33
Mean (SD)	36.63 (0.189)	36.70 (0.202)
Median	36.60	36.70
Q1, Q3	36.50, 36.80	36.50, 36.80
Min, Max	36.3, 37.1	36.4, 37.1
Change from baseline, n	36	33
Mean (SD)	-0.04 (0.213)	-0.05 (0.184)
Median	-0.10	-0.10
Q1, Q3	-0.10, 0.05	-0.20, 0.00
Min, Max	-0.5, 0.4	-0.5, 0.3

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Oral Temperature (C)

	CYP Probe N=36	CYB003 + CYP Probe N=34
2 hours post-dose, n	36	31
Mean (SD)	36.65 (0.229)	36.71 (0.224)
Median	36.60	36.70
Q1, Q3	36.50, 36.80	36.60, 36.90
Min, Max	36.2, 37.2	36.1, 37.2
Change from baseline, n	36	31
Mean (SD)	-0.02 (0.228)	-0.05 (0.254)
Median	0.00	0.00
Q1, Q3	-0.10, 0.10	-0.20, 0.10
Min, Max	-0.6, 0.5	-1.0, 0.4
4 hours post-dose, n	36	30
Mean (SD)	36.69 (0.207)	36.93 (0.303)
Median	36.70	36.90
Q1, Q3	36.60, 36.80	36.70, 37.10
Min, Max	36.3, 37.2	36.4, 37.7
Change from baseline, n	36	30
Mean (SD)	0.02 (0.192)	0.19 (0.297)
Median	0.00	0.30
Q1, Q3	-0.10, 0.10	0.00, 0.40
Min, Max	-0.4, 0.5	-0.5, 0.8

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Oral Temperature (C)

	CYP Probe N=36	CYB003 + CYP Probe N=34
6 hours post-dose, n	35	34
Mean (SD)	36.78 (0.239)	37.02 (0.289)
Median	36.70	36.90
Q1, Q3	36.60, 37.00	36.90, 37.30
Min, Max	36.4, 37.4	36.5, 37.6
Change from baseline, n	35	34
Mean (SD)	0.10 (0.235)	0.27 (0.274)
Median	0.10	0.20
Q1, Q3	0.00, 0.20	0.10, 0.50
Min, Max	-0.4, 0.8	-0.3, 0.9
8 hours post-dose, n	36	34
Mean (SD)	36.87 (0.207)	37.01 (0.301)
Median	36.80	37.00
Q1, Q3	36.70, 37.00	36.80, 37.20
Min, Max	36.5, 37.4	36.4, 37.6
Change from baseline, n	36	34
Mean (SD)	0.20 (0.246)	0.25 (0.298)
Median	0.15	0.30
Q1, Q3	0.10, 0.30	0.00, 0.40
Min, Max	-0.3, 0.8	-0.5, 0.8

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Oral Temperature (C)

	CYP Probe N=36	CYB003 + CYP Probe N=34
12 hours post-dose, n	36	34
Mean (SD)	36.83 (0.244)	36.86 (0.210)
Median	36.80	36.90
Q1, Q3	36.70, 37.00	36.70, 37.00
Min, Max	36.5, 37.7	36.5, 37.2
Change from baseline, n	36	34
Mean (SD)	0.16 (0.258)	0.11 (0.270)
Median	0.10	0.10
Q1, Q3	0.00, 0.25	-0.10, 0.30
Min, Max	-0.3, 1.1	-0.6, 0.7
24 hours post-dose, n	36	34
Mean (SD)	36.76 (0.228)	36.70 (0.180)
Median	36.70	36.70
Q1, Q3	36.60, 36.90	36.60, 36.80
Min, Max	36.4, 37.2	36.4, 37.1
Change from baseline, n	36	34
Mean (SD)	0.09 (0.217)	-0.05 (0.222)
Median	0.00	0.00
Q1, Q3	-0.05, 0.20	-0.20, 0.10
Min, Max	-0.3, 0.6	-0.5, 0.3

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Systolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	124.3 (12.72)	123.9 (12.24)
Median	122.0	124.5
Q1, Q3	116.0, 134.5	117.0, 129.0
Min, Max	103, 161	95, 152
1 hour post-dose, n	36	32
Mean (SD)	117.5 (12.26)	136.2 (12.78)
Median	115.5	133.0
Q1, Q3	108.5, 126.5	129.0, 146.0
Min, Max	91, 152	111, 165
Change from baseline, n	36	32
Mean (SD)	-6.8 (12.77)	12.5 (13.35)
Median	-4.0	10.5
Q1, Q3	-11.0, 1.0	5.5, 20.5
Min, Max	-46, 19	-24, 43

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Systolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
2 hours post-dose, n	36	29
Mean (SD)	117.3 (11.12)	138.7 (18.93)
Median	117.0	141.0
Q1, Q3	111.0, 124.5	133.0, 147.0
Min, Max	83, 137	85, 176
Change from baseline, n	36	29
Mean (SD)	-7.1 (14.85)	16.1 (14.85)
Median	-4.5	15.0
Q1, Q3	-11.5, 2.5	12.0, 23.0
Min, Max	-54, 12	-41, 48
4 hours post-dose, n	36	27
Mean (SD)	117.8 (12.60)	132.9 (18.02)
Median	115.5	130.0
Q1, Q3	109.5, 124.5	118.0, 150.0
Min, Max	94, 155	99, 165
Change from baseline, n	36	27
Mean (SD)	-6.6 (14.10)	10.0 (16.89)
Median	-6.5	6.0
Q1, Q3	-12.5, 0.0	-1.0, 18.0
Min, Max	-43, 33	-17, 48

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Systolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
6 hours post-dose, n	36	30
Mean (SD)	112.3 (14.68)	122.9 (19.90)
Median	111.0	119.0
Q1, Q3	105.0, 121.0	112.0, 141.0
Min, Max	67, 139	80, 165
Change from baseline, n	36	30
Mean (SD)	-12.0 (17.38)	-0.3 (17.23)
Median	-9.5	0.5
Q1, Q3	-19.5, -3.0	-11.0, 13.0
Min, Max	-56, 29	-33, 34
8 hours post-dose, n	36	33
Mean (SD)	112.8 (13.82)	119.1 (12.89)
Median	111.5	121.0
Q1, Q3	105.5, 122.5	113.0, 127.0
Min, Max	82, 144	94, 144
Change from baseline, n	36	33
Mean (SD)	-11.6 (15.64)	-3.9 (12.84)
Median	-9.5	-3.0
Q1, Q3	-18.0, -2.0	-12.0, 7.0
Min, Max	-59, 14	-32, 23

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Systolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
12 hours post-dose, n	36	34
Mean (SD)	120.1 (12.06)	112.6 (13.42)
Median	119.0	110.5
Q1, Q3	113.5, 126.0	106.0, 122.0
Min, Max	95, 143	69, 143
Change from baseline, n	36	34
Mean (SD)	-4.2 (14.27)	-11.3 (16.34)
Median	-2.5	-9.0
Q1, Q3	-11.5, 4.5	-17.0, -1.0
Min, Max	-42, 22	-83, 9

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Diastolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	80.7 (7.55)	80.1 (8.64)
Median	82.5	81.0
Q1, Q3	74.0, 86.5	76.0, 85.0
Min, Max	63, 94	61, 99
1 hour post-dose, n	36	32
Mean (SD)	79.9 (8.17)	89.6 (8.91)
Median	81.5	88.5
Q1, Q3	75.0, 86.0	84.5, 94.0
Min, Max	60, 97	73, 112
Change from baseline, n	36	32
Mean (SD)	-0.8 (6.86)	9.4 (8.98)
Median	1.0	8.5
Q1, Q3	-3.0, 3.0	4.0, 16.0
Min, Max	-27, 10	-11, 26

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Diastolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
2 hours post-dose, n	36	29
Mean (SD)	78.0 (7.34)	90.2 (14.23)
Median	78.0	89.0
Q1, Q3	74.5, 83.0	80.0, 98.0
Min, Max	56, 90	66, 124
Change from baseline, n	36	29
Mean (SD)	-2.7 (7.10)	10.9 (10.48)
Median	-2.0	8.0
Q1, Q3	-6.5, 1.5	5.0, 19.0
Min, Max	-31, 9	-9, 41
4 hours post-dose, n	36	27
Mean (SD)	77.6 (7.06)	86.9 (11.78)
Median	78.5	88.0
Q1, Q3	73.0, 83.0	77.0, 98.0
Min, Max	63, 92	66, 107
Change from baseline, n	36	27
Mean (SD)	-3.0 (7.56)	7.4 (10.03)
Median	-2.0	7.0
Q1, Q3	-7.5, 1.5	1.0, 14.0
Min, Max	-24, 13	-10, 29

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Diastolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
6 hours post-dose, n	36	30
Mean (SD)	72.4 (10.04)	76.9 (12.49)
Median	72.5	77.0
Q1, Q3	66.5, 79.5	69.0, 87.0
Min, Max	40, 90	54, 99
Change from baseline, n	36	30
Mean (SD)	-8.3 (10.78)	-2.3 (11.64)
Median	-6.5	-2.0
Q1, Q3	-13.5, -1.5	-8.0, 6.0
Min, Max	-42, 10	-29, 21
8 hours post-dose, n	36	33
Mean (SD)	73.8 (8.16)	76.7 (9.31)
Median	73.5	79.0
Q1, Q3	68.0, 79.0	70.0, 82.0
Min, Max	56, 93	57, 92
Change from baseline, n	36	33
Mean (SD)	-6.8 (7.61)	-2.8 (9.16)
Median	-6.5	-3.0
Q1, Q3	-11.5, -2.0	-6.0, 4.0
Min, Max	-25, 8	-27, 14

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Diastolic Blood Pressure (mmHg)

	CYP Probe N=36	CYB003 + CYP Probe N=34
12 hours post-dose, n	36	34
Mean (SD)	76.2 (7.35)	72.9 (8.78)
Median	75.5	72.5
Q1, Q3	71.0, 82.5	69.0, 77.0
Min, Max	63, 91	47, 90
Change from baseline, n	36	34
Mean (SD)	-4.4 (6.53)	-7.1 (11.08)
Median	-3.0	-5.0
Q1, Q3	-8.5, -1.0	-12.0, 0.0
Min, Max	-20, 10	-52, 10

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Pulse Rate (beats/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	68.5 (11.84)	69.1 (8.93)
Median	67.0	69.5
Q1, Q3	60.5, 77.0	62.0, 76.0
Min, Max	45, 110	53, 87
1 hour post-dose, n	36	32
Mean (SD)	62.0 (9.19)	67.0 (10.17)
Median	60.5	67.5
Q1, Q3	55.0, 69.0	58.5, 74.5
Min, Max	45, 87	46, 86
Change from baseline, n	36	32
Mean (SD)	-6.5 (7.47)	-1.8 (8.40)
Median	-6.5	-2.5
Q1, Q3	-11.0, -1.0	-8.0, 1.5
Min, Max	-23, 8	-14, 23

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Pulse Rate (beats/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
2 hours post-dose, n	36	29
Mean (SD)	59.5 (8.07)	67.4 (11.13)
Median	58.0	67.0
Q1, Q3	54.5, 62.0	60.0, 76.0
Min, Max	43, 80	41, 97
Change from baseline, n	36	29
Mean (SD)	-9.0 (8.66)	-1.0 (9.15)
Median	-9.0	-3.0
Q1, Q3	-14.0, -2.0	-8.0, 6.0
Min, Max	-35, 6	-16, 20
4 hours post-dose, n	36	27
Mean (SD)	62.0 (7.84)	75.4 (11.67)
Median	63.0	78.0
Q1, Q3	56.5, 68.0	66.0, 83.0
Min, Max	41, 76	52, 98
Change from baseline, n	36	27
Mean (SD)	-6.5 (9.11)	6.0 (9.94)
Median	-8.5	5.0
Q1, Q3	-11.5, -0.5	0.0, 13.0
Min, Max	-36, 12	-20, 22

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Pulse Rate (beats/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
6 hours post-dose, n	36	30
Mean (SD)	72.4 (11.04)	77.6 (11.47)
Median	69.5	78.5
Q1, Q3	65.5, 81.0	67.0, 87.0
Min, Max	49, 98	56, 99
Change from baseline, n	36	30
Mean (SD)	3.9 (9.21)	8.2 (6.57)
Median	3.0	6.5
Q1, Q3	-2.5, 10.0	3.0, 14.0
Min, Max	-20, 23	-1, 22
8 hours post-dose, n	36	33
Mean (SD)	71.6 (11.45)	77.8 (11.95)
Median	70.0	78.0
Q1, Q3	65.0, 79.0	69.0, 85.0
Min, Max	47, 112	54, 98
Change from baseline, n	36	33
Mean (SD)	3.1 (12.84)	8.6 (9.03)
Median	3.0	7.0
Q1, Q3	-3.0, 9.0	3.0, 14.0
Min, Max	-45, 33	-4, 30

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.1
Vital Signs by Timepoint
Safety Population

Standing Pulse Rate (beats/min)

	CYP Probe N=36	CYB003 + CYP Probe N=34
12 hours post-dose, n	36	34
Mean (SD)	73.0 (12.05)	79.8 (14.01)
Median	72.0	75.0
Q1, Q3	63.5, 82.0	70.0, 89.0
Min, Max	45, 98	58, 118
Change from baseline, n	36	34
Mean (SD)	4.5 (9.46)	10.6 (12.87)
Median	3.5	7.5
Q1, Q3	-1.0, 10.5	3.0, 16.0
Min, Max	-23, 25	-8, 47

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030501.sas FINAL (2026-08-28,12:56)

Table 14.3.5.2
Potentially Clinically Significant Vital Signs by Timepoint
Safety Population

	CYP Probe N=36	CYB003 + CYP Probe N=34
Pre-dose, N1	36	34
Heart rate < 50 bpm	3 (8.3)	1 (2.9)
Heart rate > 100 bpm	1 (2.8)	0
Supine SBP < 90 mmHg	0	0
Supine SBP > 140 mmHg	2 (5.6)	1 (2.9)
Supine DBP < 40 mmHg	0	0
Supine DBP > 90 mmHg	1 (2.8)	1 (2.9)
Respiratory rate < 12 breaths/minute	0	0
Respiratory rate > 18 breaths/minute	0	0
Oral temperature < 35.5 °C	0	0
Oral temperature > 37.8 °C	0	0
Orthostatic hypotension [a]	17 (47.2)	17 (50.0)
Reduction in SBP of >= 20 mmHg	6 (16.7)	4 (11.8)
Reduction in DBP of >= 10 mmHg	17 (47.2)	17 (50.0)

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
N is the number of participants in the respective trial arms.
Percentages are based on the number of participants in the treatment group with non-missing values for the timepoint, N1.
a. Orthostatic hypotension is calculated as 3 minute standing BP minus supine BP with orthostatic hypotension defined as a reduction of systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg.
SBP = systolic blood pressure; DBP = diastolic blood pressure.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030502.sas FINAL (2026-08-28,12:56)

Table 14.3.5.2
Potentially Clinically Significant Vital Signs by Timepoint
Safety Population

	CYP Probe N=36	CYB003 + CYP Probe N=34
1 hour post-dose, N1	36	33
Heart rate < 50 bpm	8 (22.2)	4 (12.1)
Heart rate > 100 bpm	0	1 (3.0)
Supine SBP < 90 mmHg	0	0
Supine SBP > 140 mmHg	0	6 (18.2)
Supine DBP < 40 mmHg	0	0
Supine DBP > 90 mmHg	0	4 (12.1)
Respiratory rate < 12 breaths/minute	0	0
Respiratory rate > 18 breaths/minute	0	0
Oral temperature < 35.5 °C	0	0
Oral temperature > 37.8 °C	0	0
Orthostatic hypotension [a]	13 (36.1)	15 (45.5)
Reduction in SBP of >= 20 mmHg	2 (5.6)	3 (9.1)
Reduction in DBP of >= 10 mmHg	13 (36.1)	15 (45.5)

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
N is the number of participants in the respective trial arms.
Percentages are based on the number of participants in the treatment group with non-missing values for the timepoint, N1.
a. Orthostatic hypotension is calculated as 3 minute standing BP minus supine BP with orthostatic hypotension defined as a reduction of systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg.
SBP = systolic blood pressure; DBP = diastolic blood pressure.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030502.sas FINAL (2026-08-28,12:56)

Table 14.3.5.2
Potentially Clinically Significant Vital Signs by Timepoint
Safety Population

	CYP Probe N=36	CYB003 + CYP Probe N=34
2 hours post-dose, N1	36	31
Heart rate < 50 bpm	6 (16.7)	2 (6.5)
Heart rate > 100 bpm	0	1 (3.2)
Supine SBP < 90 mmHg	0	0
Supine SBP > 140 mmHg	0	8 (25.8)
Supine DBP < 40 mmHg	0	0
Supine DBP > 90 mmHg	0	7 (22.6)
Respiratory rate < 12 breaths/minute	0	0
Respiratory rate > 18 breaths/minute	0	0
Oral temperature < 35.5 °C	0	0
Oral temperature > 37.8 °C	0	0
Orthostatic hypotension [a]	15 (41.7)	11 (35.5)
Reduction in SBP of >= 20 mmHg	4 (11.1)	1 (3.2)
Reduction in DBP of >= 10 mmHg	15 (41.7)	11 (35.5)

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
N is the number of participants in the respective trial arms.
Percentages are based on the number of participants in the treatment group with non-missing values for the timepoint, N1.
a. Orthostatic hypotension is calculated as 3 minute standing BP minus supine BP with orthostatic hypotension defined as a reduction of systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg.
SBP = systolic blood pressure; DBP = diastolic blood pressure.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030502.sas FINAL (2026-08-28,12:56)

Table 14.3.5.2
Potentially Clinically Significant Vital Signs by Timepoint
Safety Population

	CYP Probe N=36	CYB003 + CYP Probe N=34
4 hours post-dose, N1	36	31
Heart rate < 50 bpm	2 (5.6)	0
Heart rate > 100 bpm	0	0
Supine SBP < 90 mmHg	1 (2.8)	0
Supine SBP > 140 mmHg	0	8 (25.8)
Supine DBP < 40 mmHg	0	0
Supine DBP > 90 mmHg	0	3 (9.7)
Respiratory rate < 12 breaths/minute	0	0
Respiratory rate > 18 breaths/minute	0	0
Oral temperature < 35.5 °C	0	0
Oral temperature > 37.8 °C	0	0
Orthostatic hypotension [a]	12 (33.3)	13 (41.9)
Reduction in SBP of >= 20 mmHg	5 (13.9)	4 (12.9)
Reduction in DBP of >= 10 mmHg	11 (30.6)	13 (41.9)

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
N is the number of participants in the respective trial arms.
Percentages are based on the number of participants in the treatment group with non-missing values for the timepoint, N1.
a. Orthostatic hypotension is calculated as 3 minute standing BP minus supine BP with orthostatic hypotension defined as a reduction of systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg.
SBP = systolic blood pressure; DBP = diastolic blood pressure.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030502.sas FINAL (2026-08-28,12:56)

Table 14.3.5.2
Potentially Clinically Significant Vital Signs by Timepoint
Safety Population

	CYP Probe N=36	CYB003 + CYP Probe N=34
6 hours post-dose, N1	36	34
Heart rate < 50 bpm	1 (2.8)	1 (2.9)
Heart rate > 100 bpm	0	0
Supine SBP < 90 mmHg	1 (2.8)	0
Supine SBP > 140 mmHg	0	3 (8.8)
Supine DBP < 40 mmHg	0	0
Supine DBP > 90 mmHg	0	0
Respiratory rate < 12 breaths/minute	0	0
Respiratory rate > 18 breaths/minute	0	0
Oral temperature < 35.5 °C	0	0
Oral temperature > 37.8 °C	0	0
Orthostatic hypotension [a]	16 (44.4)	10 (29.4)
Reduction in SBP of >= 20 mmHg	4 (11.1)	2 (5.9)
Reduction in DBP of >= 10 mmHg	16 (44.4)	8 (23.5)

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
N is the number of participants in the respective trial arms.
Percentages are based on the number of participants in the treatment group with non-missing values for the timepoint, N1.
a. Orthostatic hypotension is calculated as 3 minute standing BP minus supine BP with orthostatic hypotension defined as a reduction of systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg.
SBP = systolic blood pressure; DBP = diastolic blood pressure.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030502.sas FINAL (2026-08-28,12:56)

Table 14.3.5.2
Potentially Clinically Significant Vital Signs by Timepoint
Safety Population

	CYP Probe N=36	CYB003 + CYP Probe N=34
8 hours post-dose, N1	36	34
Heart rate < 50 bpm	2 (5.6)	1 (2.9)
Heart rate > 100 bpm	1 (2.8)	0
Supine SBP < 90 mmHg	2 (5.6)	0
Supine SBP > 140 mmHg	0	1 (2.9)
Supine DBP < 40 mmHg	0	0
Supine DBP > 90 mmHg	0	0
Respiratory rate < 12 breaths/minute	1 (2.8)	0
Respiratory rate > 18 breaths/minute	0	0
Oral temperature < 35.5 °C	0	0
Oral temperature > 37.8 °C	0	0
Orthostatic hypotension [a]	21 (58.3)	8 (23.5)
Reduction in SBP of >= 20 mmHg	4 (11.1)	1 (2.9)
Reduction in DBP of >= 10 mmHg	21 (58.3)	8 (23.5)

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
N is the number of participants in the respective trial arms.
Percentages are based on the number of participants in the treatment group with non-missing values for the timepoint, N1.
a. Orthostatic hypotension is calculated as 3 minute standing BP minus supine BP with orthostatic hypotension defined as a reduction of systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg.
SBP = systolic blood pressure; DBP = diastolic blood pressure.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030502.sas FINAL (2026-08-28,12:56)

Table 14.3.5.2
Potentially Clinically Significant Vital Signs by Timepoint
Safety Population

	CYP Probe N=36	CYB003 + CYP Probe N=34
12 hours post-dose, N1	36	34
Heart rate < 50 bpm	2 (5.6)	0
Heart rate > 100 bpm	0	3 (8.8)
Supine SBP < 90 mmHg	0	0
Supine SBP > 140 mmHg	1 (2.8)	1 (2.9)
Supine DBP < 40 mmHg	0	0
Supine DBP > 90 mmHg	0	0
Respiratory rate < 12 breaths/minute	1 (2.8)	0
Respiratory rate > 18 breaths/minute	0	0
Oral temperature < 35.5 °C	0	0
Oral temperature > 37.8 °C	0	0
Orthostatic hypotension [a]	24 (66.7)	17 (50.0)
Reduction in SBP of >= 20 mmHg	4 (11.1)	0
Reduction in DBP of >= 10 mmHg	24 (66.7)	17 (50.0)

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
N is the number of participants in the respective trial arms.
Percentages are based on the number of participants in the treatment group with non-missing values for the timepoint, N1.
a. Orthostatic hypotension is calculated as 3 minute standing BP minus supine BP with orthostatic hypotension defined as a reduction of systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg.
SBP = systolic blood pressure; DBP = diastolic blood pressure.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030502.sas FINAL (2026-08-28,12:56)

Table 14.3.5.2
Potentially Clinically Significant Vital Signs by Timepoint
Safety Population

	CYP Probe N=36	CYB003 + CYP Probe N=34
24 hours post-dose, N1	36	34
Heart rate < 50 bpm	1 (2.8)	3 (8.8)
Heart rate > 100 bpm	0	0
Supine SBP < 90 mmHg	0	1 (2.9)
Supine SBP > 140 mmHg	0	0
Supine DBP < 40 mmHg	0	0
Supine DBP > 90 mmHg	0	0
Respiratory rate < 12 breaths/minute	0	0
Respiratory rate > 18 breaths/minute	0	0
Oral temperature < 35.5 °C	0	0
Oral temperature > 37.8 °C	0	0
Orthostatic hypotension [a]	0	0
Reduction in SBP of >= 20 mmHg	0	0
Reduction in DBP of >= 10 mmHg	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
N is the number of participants in the respective trial arms.
Percentages are based on the number of participants in the treatment group with non-missing values for the timepoint, N1.
a. Orthostatic hypotension is calculated as 3 minute standing BP minus supine BP with orthostatic hypotension defined as a reduction of systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg.
SBP = systolic blood pressure; DBP = diastolic blood pressure.

Source: Listing 16.2.9.1 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030502.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

ECG Mean Ventricular Rate (bpm)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	57.4 (7.92)	59.2 (7.80)
Median	56.5	58.0
Q1, Q3	52.0, 63.0	53.0, 64.0
Min, Max	42, 75	47, 83
2 hours post-dose, n	36	32
Mean (SD)	50.8 (5.93)	64.0 (11.11)
Median	51.0	63.5
Q1, Q3	46.5, 54.5	57.0, 70.5
Min, Max	36, 66	44, 93
Change from baseline, n	36	32
Mean (SD)	-6.6 (6.22)	4.3 (11.53)
Median	-5.5	3.0
Q1, Q3	-10.0, -1.5	-2.5, 7.0
Min, Max	-20, 3	-23, 33

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

ECG Mean Ventricular Rate (bpm)

	CYP Probe N=36	CYB003 + CYP Probe N=34
4 hours post-dose, n	36	28
Mean (SD)	53.3 (6.31)	70.2 (10.88)
Median	53.0	69.5
Q1, Q3	50.0, 56.0	63.5, 76.5
Min, Max	40, 68	52, 93
Change from baseline, n	36	28
Mean (SD)	-4.1 (5.67)	10.6 (11.08)
Median	-4.0	10.0
Q1, Q3	-9.0, 0.0	5.0, 16.5
Min, Max	-16, 7	-18, 40
8 hours post-dose, n	36	34
Mean (SD)	60.7 (8.40)	67.3 (8.73)
Median	60.5	67.5
Q1, Q3	54.0, 65.0	62.0, 73.0
Min, Max	44, 86	46, 90
Change from baseline, n	36	34
Mean (SD)	3.3 (7.12)	8.0 (8.91)
Median	3.0	7.5
Q1, Q3	-2.5, 8.5	3.0, 13.0
Min, Max	-10, 20	-17, 28

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

ECG Mean Ventricular Rate (bpm)

	CYP Probe N=36	CYB003 + CYP Probe N=34
24 hours post-dose, n	36	34
Mean (SD)	57.9 (7.39)	59.9 (7.58)
Median	57.0	60.0
Q1, Q3	53.0, 60.5	56.0, 64.0
Min, Max	46, 79	42, 83
Change from baseline, n	36	34
Mean (SD)	0.5 (4.53)	0.7 (6.90)
Median	1.0	0.0
Q1, Q3	-3.5, 4.0	-5.0, 6.0
Min, Max	-8, 8	-11, 14

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

PR Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	167.2 (24.34)	164.9 (24.56)
Median	166.5	161.0
Q1, Q3	153.0, 183.0	152.0, 181.0
Min, Max	105, 215	113, 220
2 hours post-dose, n	36	32
Mean (SD)	169.7 (25.55)	163.4 (38.41)
Median	167.5	166.5
Q1, Q3	156.5, 183.0	152.0, 175.5
Min, Max	106, 222	0, 230
Change from baseline, n	36	32
Mean (SD)	2.5 (9.16)	-2.5 (23.26)
Median	1.0	2.0
Q1, Q3	-3.0, 6.5	-5.0, 10.0
Min, Max	-12, 33	-113, 24

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

PR Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
4 hours post-dose, n	36	28
Mean (SD)	168.1 (25.33)	163.8 (23.97)
Median	165.0	163.0
Q1, Q3	155.5, 180.0	152.0, 176.0
Min, Max	106, 226	111, 230
Change from baseline, n	36	28
Mean (SD)	0.9 (8.67)	-0.3 (10.91)
Median	0.5	-3.0
Q1, Q3	-5.0, 6.0	-8.0, 9.0
Min, Max	-14, 24	-17, 26
8 hours post-dose, n	36	34
Mean (SD)	161.4 (23.29)	171.0 (41.26)
Median	162.0	164.5
Q1, Q3	148.0, 175.0	146.0, 182.0
Min, Max	110, 210	115, 350
Change from baseline, n	36	34
Mean (SD)	-5.8 (11.48)	6.1 (34.80)
Median	-5.5	1.5
Q1, Q3	-11.5, 1.0	-9.0, 9.0
Min, Max	-34, 18	-23, 193

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

PR Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
24 hours post-dose, n	36	34
Mean (SD)	165.3 (23.03)	162.7 (24.13)
Median	164.5	160.0
Q1, Q3	149.0, 181.0	149.0, 175.0
Min, Max	106, 208	108, 208
Change from baseline, n	36	34
Mean (SD)	-1.9 (8.12)	-2.2 (8.34)
Median	-2.5	-2.5
Q1, Q3	-7.5, 4.0	-6.0, 4.0
Min, Max	-18, 19	-22, 16

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

RR Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	1055.8 (145.67)	1022.3 (128.73)
Median	1051.0	1023.0
Q1, Q3	943.5, 1143.5	930.0, 1131.0
Min, Max	800, 1406	715, 1274
2 hours post-dose, n	36	32
Mean (SD)	1184.7 (141.81)	956.5 (163.17)
Median	1174.5	937.5
Q1, Q3	1092.0, 1268.5	842.5, 1045.0
Min, Max	909, 1636	640, 1360
Change from baseline, n	36	32
Mean (SD)	129.0 (112.47)	-56.4 (154.09)
Median	123.5	-56.0
Q1, Q3	37.5, 216.5	-105.0, 34.5
Min, Max	-66, 337	-360, 271

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

RR Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
4 hours post-dose, n	36	28
Mean (SD)	1130.2 (130.98)	868.5 (132.93)
Median	1115.5	860.0
Q1, Q3	1061.5, 1195.5	777.5, 936.5
Min, Max	879, 1492	645, 1134
Change from baseline, n	36	28
Mean (SD)	74.4 (99.88)	-147.6 (138.93)
Median	88.0	-158.0
Q1, Q3	0.5, 144.0	-209.0, -88.5
Min, Max	-116, 228	-502, 195
8 hours post-dose, n	36	34
Mean (SD)	999.5 (136.91)	900.3 (120.67)
Median	986.0	884.5
Q1, Q3	921.5, 1108.5	816.0, 966.0
Min, Max	693, 1363	665, 1280
Change from baseline, n	36	34
Mean (SD)	-56.2 (114.40)	-122.0 (120.47)
Median	-65.0	-114.0
Q1, Q3	-132.0, 47.0	-201.0, -58.0
Min, Max	-272, 129	-347, 186

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

RR Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
24 hours post-dose, n	36	34
Mean (SD)	1040.6 (122.24)	1007.3 (131.10)
Median	1041.5	992.0
Q1, Q3	983.5, 1122.5	924.0, 1069.0
Min, Max	753, 1282	715, 1409
Change from baseline, n	36	34
Mean (SD)	-15.2 (81.80)	-15.1 (122.15)
Median	-21.0	-2.5
Q1, Q3	-73.0, 54.5	-117.0, 80.0
Min, Max	-209, 157	-253, 203

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

QRS Duration (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	97.0 (8.38)	94.0 (7.18)
Median	95.5	92.0
Q1, Q3	91.5, 100.5	89.0, 99.0
Min, Max	82, 114	77, 110
2 hours post-dose, n	36	32
Mean (SD)	97.2 (7.74)	93.2 (7.71)
Median	95.5	94.5
Q1, Q3	91.0, 103.5	87.5, 98.0
Min, Max	86, 113	75, 107
Change from baseline, n	36	32
Mean (SD)	0.2 (4.56)	-1.2 (4.97)
Median	0.0	-1.0
Q1, Q3	-3.5, 3.0	-2.0, 2.0
Min, Max	-8, 12	-12, 7

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

QRS Duration (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
4 hours post-dose, n	36	28
Mean (SD)	96.1 (7.27)	92.8 (7.68)
Median	94.0	93.0
Q1, Q3	92.0, 101.5	88.0, 97.5
Min, Max	81, 112	77, 106
Change from baseline, n	36	28
Mean (SD)	-0.9 (5.34)	-1.1 (5.09)
Median	-2.0	-1.0
Q1, Q3	-3.5, 2.0	-3.5, 1.5
Min, Max	-13, 17	-13, 9
8 hours post-dose, n	36	34
Mean (SD)	96.1 (8.26)	93.2 (7.53)
Median	96.0	93.0
Q1, Q3	89.5, 101.0	88.0, 98.0
Min, Max	81, 114	80, 110
Change from baseline, n	36	34
Mean (SD)	-0.9 (5.48)	-0.8 (5.37)
Median	-1.0	-0.5
Q1, Q3	-5.0, 4.0	-3.0, 2.0
Min, Max	-11, 9	-16, 13

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

QRS Duration (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
24 hours post-dose, n	36	34
Mean (SD)	96.7 (8.07)	96.4 (8.12)
Median	95.5	95.5
Q1, Q3	90.5, 102.0	92.0, 103.0
Min, Max	82, 114	79, 114
Change from baseline, n	36	34
Mean (SD)	-0.3 (4.86)	2.4 (5.59)
Median	0.0	2.0
Q1, Q3	-4.0, 2.5	1.0, 5.0
Min, Max	-8, 14	-10, 15

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

QT Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	399.7 (20.88)	393.8 (20.13)
Median	398.5	392.0
Q1, Q3	387.0, 414.5	377.0, 410.0
Min, Max	355, 443	361, 437
2 hours post-dose, n	36	32
Mean (SD)	411.9 (17.81)	381.8 (26.61)
Median	411.5	382.0
Q1, Q3	401.5, 421.5	364.5, 398.0
Min, Max	367, 449	321, 445
Change from baseline, n	36	32
Mean (SD)	12.1 (11.40)	-10.1 (20.74)
Median	12.5	-11.5
Q1, Q3	6.0, 18.5	-21.0, -1.0
Min, Max	-15, 32	-61, 33

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

QT Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
4 hours post-dose, n	36	28
Mean (SD)	409.4 (20.33)	368.3 (23.49)
Median	408.5	365.5
Q1, Q3	397.5, 424.5	349.5, 382.5
Min, Max	363, 464	333, 430
Change from baseline, n	36	28
Mean (SD)	9.7 (12.16)	-25.4 (18.17)
Median	12.5	-26.0
Q1, Q3	0.0, 19.0	-33.5, -13.0
Min, Max	-17, 28	-56, 21
8 hours post-dose, n	36	34
Mean (SD)	389.5 (19.33)	374.3 (19.89)
Median	388.0	370.0
Q1, Q3	377.0, 402.0	361.0, 393.0
Min, Max	353, 430	338, 408
Change from baseline, n	36	34
Mean (SD)	-10.3 (15.49)	-19.5 (20.29)
Median	-13.0	-18.0
Q1, Q3	-21.5, 3.0	-29.0, -6.0
Min, Max	-43, 18	-93, 12

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

QT Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
24 hours post-dose, n	36	34
Mean (SD)	399.0 (19.83)	405.4 (20.94)
Median	404.5	407.5
Q1, Q3	381.5, 414.5	393.0, 419.0
Min, Max	359, 432	362, 446
Change from baseline, n	36	34
Mean (SD)	-0.7 (12.49)	11.7 (20.33)
Median	0.0	12.5
Q1, Q3	-10.5, 8.0	-3.0, 22.0
Min, Max	-31, 21	-26, 59

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

QTcF Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), n	36	34
Mean (SD)	393.1 (17.26)	391.1 (15.27)
Median	390.5	388.0
Q1, Q3	379.0, 405.0	382.0, 400.0
Min, Max	363, 431	369, 426
2 hours post-dose, n	36	32
Mean (SD)	389.3 (18.23)	388.2 (15.29)
Median	385.5	386.0
Q1, Q3	378.5, 403.0	377.5, 397.0
Min, Max	357, 423	367, 424
Change from baseline, n	36	32
Mean (SD)	-3.8 (10.33)	-2.2 (9.08)
Median	-2.5	-2.0
Q1, Q3	-11.5, 5.0	-9.5, 2.5
Min, Max	-26, 14	-20, 16

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

QTcF Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
4 hours post-dose, n	36	28
Mean (SD)	393.3 (19.68)	386.5 (14.23)
Median	387.5	386.5
Q1, Q3	378.5, 405.0	376.5, 398.0
Min, Max	369, 462	360, 412
Change from baseline, n	36	28
Mean (SD)	0.3 (10.82)	-5.4 (11.39)
Median	1.0	-5.5
Q1, Q3	-5.5, 5.0	-12.0, 2.0
Min, Max	-22, 43	-31, 15
8 hours post-dose, n	36	34
Mean (SD)	390.0 (13.71)	388.7 (17.49)
Median	388.0	386.0
Q1, Q3	379.5, 397.0	374.0, 402.0
Min, Max	366, 418	360, 431
Change from baseline, n	36	34
Mean (SD)	-3.1 (9.90)	-2.4 (14.50)
Median	-3.0	0.0
Q1, Q3	-10.0, 3.5	-14.0, 5.0
Min, Max	-28, 16	-47, 31

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.1
Electrocardiogram Parameter Results by Timepoint
Safety Population

QTcF Interval (msec)

	CYP Probe N=36	CYB003 + CYP Probe N=34
24 hours post-dose, n	36	34
Mean (SD)	393.9 (14.56)	404.9 (19.35)
Median	392.5	400.5
Q1, Q3	384.0, 400.5	388.0, 418.0
Min, Max	364, 430	376, 447
Change from baseline, n	36	34
Mean (SD)	0.9 (9.73)	13.8 (14.57)
Median	0.0	12.0
Q1, Q3	-2.5, 4.0	5.0, 20.0
Min, Max	-30, 25	-12, 56

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants with the measurement available.
N is the number of participants in the respective trial arms.
SD = standard deviation; Q1 = first quartile; Q3 = third quartile; Min = minimum; Max = maximum.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030601.sas FINAL (2026-08-28,12:56)

Table 14.3.6.2
Potentially Clinically Significant QTcF Findings by Timepoint
Safety Population

Timepoint	CYP Probe N=36	CYB003 + CYP Probe N=34
Baseline (Pre-dose), N1	36	34
<=450 msec, n (%)	36 (100)	34 (100)
>450 to <=480 msec, n (%)	0	0
>480 to <=500 msec, n (%)	0	0
>500 msec, n (%)	0	0
2 hours post-dose, N1	36	32
<=450 msec, n (%)	36 (100)	32 (100)
>450 to <=480 msec, n (%)	0	0
>480 to <=500 msec, n (%)	0	0
>500 msec, n (%)	0	0
Change from baseline, N1	36	32
<=30 msec, n (%)	36 (100)	32 (100)
>30 to <=60 msec, n (%)	0	0
>60 to <=90 msec, n (%)	0	0
>90 msec, n (%)	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants within a specific category.
N is the number of participants in the respective trial arms.
Percentages are based on the number of participants in the treatment group with non-missing values for the timepoint, N1.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030602.sas FINAL (2026-08-28,12:56)

Table 14.3.6.2
Potentially Clinically Significant QTcF Findings by Timepoint
Safety Population

Timepoint	CYP Probe N=36	CYB003 + CYP Probe N=34
4 hours post-dose, N1	36	28
<=450 msec, n (%)	35 (97.2)	28 (100)
>450 to <=480 msec, n (%)	1 (2.8)	0
>480 to <=500 msec, n (%)	0	0
>500 msec, n (%)	0	0
Change from baseline, N1	36	28
<=30 msec, n (%)	35 (97.2)	28 (100)
>30 to <=60 msec, n (%)	1 (2.8)	0
>60 to <=90 msec, n (%)	0	0
>90 msec, n (%)	0	0
8 hours post-dose, N1	36	34
<=450 msec, n (%)	36 (100)	34 (100)
>450 to <=480 msec, n (%)	0	0
>480 to <=500 msec, n (%)	0	0
>500 msec, n (%)	0	0
Change from baseline, N1	36	34
<=30 msec, n (%)	36 (100)	33 (97.1)
>30 to <=60 msec, n (%)	0	1 (2.9)
>60 to <=90 msec, n (%)	0	0
>90 msec, n (%)	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants within a specific category.
N is the number of participants in the respective trial arms.
Percentages are based on the number of participants in the treatment group with non-missing values for the timepoint, N1.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030602.sas FINAL (2026-08-28,12:56)

Table 14.3.6.2
Potentially Clinically Significant QTcF Findings by Timepoint
Safety Population

Timepoint	CYP Probe N=36	CYB003 + CYP Probe N=34
24 hours post-dose, N1	36	34
<=450 msec, n (%)	36 (100)	34 (100)
>450 to <=480 msec, n (%)	0	0
>480 to <=500 msec, n (%)	0	0
>500 msec, n (%)	0	0
Change from baseline, N1	36	34
<=30 msec, n (%)	36 (100)	29 (85.3)
>30 to <=60 msec, n (%)	0	5 (14.7)
>60 to <=90 msec, n (%)	0	0
>90 msec, n (%)	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Baseline is the last available non-missing value prior to treatment within each period.
n is the number of participants within a specific category.
N is the number of participants in the respective trial arms.
Percentages are based on the number of participants in the treatment group with non-missing values for the timepoint, N1.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030602.sas FINAL (2026-08-28,12:56)

Table 14.3.6.3
Electrocardiogram Investigator Assessment: Shift Table by Timepoint
Safety Population

		Baseline (Pre-dose)									
		CYP Probe N=36					CYB003 + CYP Probe N=34				
Timepoint		Normal n (%)	Abnormal – NCS n (%)	Abnormal – CS n (%)	Total n (%)	Missing n	Normal n (%)	Abnormal – NCS n (%)	Abnormal – CS n (%)	Total n (%)	Missing n
2 hours post-dose	Normal	1 (10.0)	1 (3.8)	0	2 (5.6)	0	10 (83.3)	2 (10.0)	0	12 (37.5)	0
	Abnormal – NCS	9 (90.0)	25 (96.2)	0	34 (94.4)	0	2 (16.7)	18 (90.0)	0	20 (62.5)	0
	Abnormal – CS	0	0	0	0	0	0	0	0	0	0
	Total	10 (100)	26 (100)	0	36 (100)	0	12 (100)	20 (100)	0	32 (100)	0
	Missing	0	0	0	0	0	0	2	0	2	0
4 hours post-dose	Normal	5 (50.0)	1 (3.8)	0	6 (16.7)	0	8 (72.7)	2 (11.8)	0	10 (35.7)	0
	Abnormal – NCS	5 (50.0)	25 (96.2)	0	30 (83.3)	0	3 (27.3)	15 (88.2)	0	18 (64.3)	0
	Abnormal – CS	0	0	0	0	0	0	0	0	0	0
	Total	10 (100)	26 (100)	0	36 (100)	0	11 (100)	17 (100)	0	28 (100)	0
	Missing	0	0	0	0	0	1	5	0	6	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Clinical significance is determined by the investigator.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in the baseline category with non-missing data at the given post-baseline timepoint; n1 is displayed as n in the Total row for the particular column.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).
CS = Clinically significant, NCS = Not clinically significant.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030603.sas FINAL (2026-08-28,12:56)

Table 14.3.6.3
Electrocardiogram Investigator Assessment: Shift Table by Timepoint
Safety Population

		Baseline (Pre-dose)									
		CYP Probe N=36					CYB003 + CYP Probe N=34				
Timepoint		Normal n (%)	Abnormal – NCS n (%)	Abnormal – CS n (%)	Total n (%)	Missing n	Normal n (%)	Abnormal – NCS n (%)	Abnormal – CS n (%)	Total n (%)	Missing n
8 hours post-dose	Normal	6 (60.0)	3 (11.5)	0	9 (25.0)	0	8 (66.7)	8 (36.4)	0	16 (47.1)	0
	Abnormal – NCS	4 (40.0)	23 (88.5)	0	27 (75.0)	0	4 (33.3)	14 (63.6)	0	18 (52.9)	0
	Abnormal – CS	0	0	0	0	0	0	0	0	0	0
	Total	10 (100)	26 (100)	0	36 (100)	0	12 (100)	22 (100)	0	34 (100)	0
	Missing	0	0	0	0	0	0	0	0	0	0
24 hours post-dose	Normal	9 (90.0)	5 (19.2)	0	14 (38.9)	0	10 (83.3)	2 (9.1)	0	12 (35.3)	0
	Abnormal – NCS	1 (10.0)	21 (80.8)	0	22 (61.1)	0	2 (16.7)	20 (90.9)	0	22 (64.7)	0
	Abnormal – CS	0	0	0	0	0	0	0	0	0	0
	Total	10 (100)	26 (100)	0	36 (100)	0	12 (100)	22 (100)	0	34 (100)	0
	Missing	0	0	0	0	0	0	0	0	0	0

CYP Probe refers to Period 1 and CYB003 + CYP Probe refers to Period 2.
Clinical significance is determined by the investigator.
n is the number of participants with the measurement available.
n1 is the number of participants in the respective trial arm who had data in the baseline category with non-missing data at the given post-baseline timepoint; n1 is displayed as n in the Total row for the particular column.
N is the numbers of participants in the respective trial arms.
Percentages are calculated as 100 x (n/n1).
CS = Clinically significant, NCS = Not clinically significant.

Source: Listing 16.2.9.2 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030603.sas FINAL (2026-08-28,12:56)

Table 14.3.7.1
Columbia Suicide Severity Rating Scale (C-SSRS) by Visit
Safety Population

Timepoint	Total N=36 n (%)
Screening, N1	36
Suicide Ideation	
Wish to be Dead	1 (2.8)
Non-Specific Active Suicidal Thoughts	1 (2.8)
Active Suicidal Ideation with Any Methods (Not Plan) without Intent to Act	1 (2.8)
Active Suicidal Ideation with Some Intent to Act, without Specific Plan	0
Active Suicidal Ideation with Specific Plan and Intent	0
Suicide Behavior	
Actual Attempt	0
Interrupted Attempt	0
Aborted Attempt	0
Preparatory Acts or Behavior	0
Suicidal Behavior [a]	0
Suicide Ideation or Behavior Present	0
Non-Suicidal Self-Injurious Behavior	0

N1 is the number of participants with non-missing responses at current visit and used as the denominator for percentages.
Note: C-SSRS Lifetime administer at Screening. C-SSRS Since Last Visit administered at Days 7, 8, and Follow-up.
a. Suicidal behavior was present during the assessment period.
b. Death by suicide occurred since last assessment.

Source: Listing 16.2.9.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030701.sas FINAL (2026-08-28,12:56)

Table 14.3.7.1
Columbia Suicide Severity Rating Scale (C-SSRS) by Visit
Safety Population

Timepoint	Total N=36 n (%)
Since Last Visit, N1	35
Suicide Ideation	
Wish to be Dead	0
Non-Specific Active Suicidal Thoughts	0
Active Suicidal Ideation with Any Methods (Not Plan) without Intent to Act	0
Active Suicidal Ideation with Some Intent to Act, without Specific Plan	0
Active Suicidal Ideation with Specific Plan and Intent	0
Suicide Behavior	
Actual Attempt	0
Interrupted Attempt	0
Aborted Attempt	0
Preparatory Acts or Behavior	0
Suicidal Behavior [a]	0
Suicide [b]	0
Suicide Ideation or Behavior Present	0
Non-Suicidal Self-Injurious Behavior	0

N1 is the number of participants with non-missing responses at current visit and used as the denominator for percentages.
Note: C-SSRS Lifetime administer at Screening. C-SSRS Since Last Visit administered at Days 7, 8, and Follow-up.
a. Suicidal behavior was present during the assessment period.
b. Death by suicide occurred since last assessment.

Source: Listing 16.2.9.4 EDC Extraction Date: 2026-04-07
Program: /cyb003-101/final/programs/statout/t14030701.sas FINAL (2026-08-28,12:56)