

"When I lost it, they dragged me out": How Care Encounters Empower Marginalized Young Adults' Mental Health Care-Seeking

JIAYING LIU, The University of Texas at Austin, USA

YAN ZHANG, The University of Texas at Austin, USA

CSCW research has long explored ways to enhance the well-being of marginalized populations. This study examines how young adults with diverse marginalized identities navigate their mental health care-seeking journeys through in-depth interviews and visual elicitation methods. Our research with 18 U.S. participants reveals predominantly *passive* behavioral patterns shaped by participants' lived experiences of marginalization. A key finding centers on "*care encounters*" - serendipitous interactions with mental health resources that occur when individuals are not actively seeking support. These encounters emerged as critical turning points, catalyzing shifts from passive to proactive care-seeking behaviors. The transformative impact of these encounters operated through three primary mechanisms: tangible assistance, supportive discourse, and social connection building. Our analysis illuminates the infrastructural conditions that enable such transformative encounters and their effectiveness in connecting marginalized young adults with mental health care. This research advances our understanding of how marginalization influences care-seeking behaviors while providing concrete design implications for socio-technical interventions. Our findings suggest strategies for intentionally engineering care encounters that can help overcome passive behavioral patterns and strengthen marginalized young adults' engagement with mental health resources.

Additional Key Words and Phrases: Help-Seeking, Socio-technical ecosystem, Qualitative, Pathway to Care, Community Health, Social Ecological Theory

ACM Reference Format:

Jiaying Liu and Yan Zhang. 2025. "When I lost it, they dragged me out": How Care Encounters Empower Marginalized Young Adults' Mental Health Care-Seeking. 1, 1 (February 2025), 24 pages. <https://doi.org/10.1145/nnnnnnnn.nnnnnnnn>

1 INTRODUCTION

The mental health crisis continues to escalate both in the United States and globally. According to the National Institute of Mental Health (2023) [54], approximately 57.8 million U.S. adults lived with a mental illness in 2021. Young adults aged 18-25 demonstrate the highest prevalence of mental health concerns (33.7%), yet paradoxically exhibit the lowest rate of seeking outpatient mental health treatment. This disparity becomes even more pronounced among young adults with marginalized identities, particularly racial minorities and LGBTQ+ individuals, who demonstrate significantly lower care-seeking behaviors [30].

A substantial body of HCI and CSCW has addressed mental health support through various digital interventions, such as self-tracking tools [2, 64, 84] and peer support platforms [81, 86, 90], targeting the general population. However, researchers have increasingly recognized that

Authors' addresses: Jiaying Liu, The University of Texas at Austin, USA, jiayingliu@utexas.edu; Yan Zhang, The University of Texas at Austin, Austin, USA, yanz@utexas.edu.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

© 2025 Association for Computing Machinery.

XXXX-XXXX/2025/2-ART \$15.00

<https://doi.org/10.1145/nnnnnnnn.nnnnnnnn>

certain technological solutions may prove inadequate for marginalized populations, who often face limited access to technological resources [29, 51]. Moreover, these populations encounter multifaceted structural barriers in adopting such tools due to their unique cultural, social, and economic circumstances. For instance, Tachtler et al. [78] identified specific micro and macro-level barriers that impede unaccompanied immigrant children's access to and utilization of mental health applications.

Existing research has documented significantly low technology adoption rates among marginalized young adults seeking mental health support [28]. To design ecologically valid socio-technical interventions, it is crucial to understand the lived experiences shaping their care-seeking journeys. While prior work has examined various barriers of marginalized communities to access mental health care [72, 75], there remains a critical gap in understanding how perceived marginalized identity influences care-seeking practices. This research addresses this gap by investigating two research questions:

- How do marginalized young adults navigate their care-seeking journey?
- How does the socio-tech ecosystem of resources facilitate or hinder care-seeking?

We interviewed 18 U.S. young adults with diverse marginalized identities such as LGBTQ+, racial minorities, and homelessness. Our analysis reveals patterns of passiveness in participants' care-seeking behaviors, demonstrating how their marginalized positions and identities constrain access to care and shape their perceptions of mental health resources. These limitations not only restrict their care aspirations but also create isolation from support systems.

Importantly, we found that passive care-seeking practices can transform into proactive patterns through "care encounters" - serendipitous interactions that provide marginalized young adults with tangible assistance, supportive discourse, and social connection. Our analysis examines how institutional and infrastructural designs facilitate these transformative encounters. However, we also identified two critical barriers within the existing socio-technical ecosystem that can impede transitions toward proactive care-seeking: challenges in establishing trust and limitations in empowering individual agency.

- We reveal through in-depth qualitative analysis how marginalization shapes and constrains young adults' mental health care-seeking journeys.
- We identify and examine how "care encounters" transform care-seeking behaviors, illuminating the infrastructural conditions that enable these pivotal interactions.
- We propose design strategies for engineering effective care encounters within the socio-technical ecosystem, addressing multiple levels of mental health support.

2 RELATED WORK

We first review the literature concerning general behavior models around mental health care-seeking and then center on the practices of marginalized populations. We then elaborate on the ecological view of mental health care-seeking which serves as an overarching methodology approach that supports our analysis.

2.1 Seeking Care for Mental Health Concerns

Care-seeking is also referred to as help-seeking, denoting people's behavior of actively obtaining assistance from various sources [68]. The assistance includes suggestions, information, and treatment that serve pragmatic purposes and also social and emotional support and understanding [38].

One of the earliest models focused on formal help-seeking in the medical system. Huxley [31] outlined pathways that a patient may need to navigate through to get care. The pathway consists of five layers. Potential patients entered the pathway from general practitioners, via whom they were referred to primary care, then secondary psychiatric care, and ended with admission to the hospital. The following studies identified other resources that act as referrals within [25] and outside [4] of the medical system, such as native or religious healers, nurses, psychiatric services, and police institutions. The model does not sufficiently consider external factors such as socio-economic status, cultural influences, or the role of informal support networks, which can significantly impact the care-seeking journey.

Rickwood's Stages of help-seeking framework [67] describes patients' subjective experiences of the help-seeking process. The model delineates four key steps: (1) recognizing symptoms and realizing the need for assistance, (2) expressing symptoms and signaling the need for support, (3) identifying available and accessible sources of help, and (4) the final step contingent on the individual's willingness to disclose difficulties to the chosen source. The stages are presented as static categories, potentially overlooking the dynamic nature of help-seeking over time.

Seeking help for mental health is not a quick and simple decision; rather, it is a complex and dynamic process dubbed with stigmatization and solitary feelings [39]. In the process, people constantly negotiate with themselves "if seeking help is necessary" and "when and how to seek help". Biddle et al. [5] delineated this struggle through the Cycle of Avoidance (COA) model, depicting the tensions involved in making sense of, accepting, and avoiding mental distress. They identified several key actions that postpone help-seeking, including normalizing symptoms, proposing alternative explanations, accommodating increasingly severe distress, pushing the threshold between "normal" and "real" distress, and delaying help-seeking. Other studies also pinpointed similar cognitive strategies such as utilizing self-resilience, denying the effects of professional help, and problematizing help-seeking [1, 48].

While these models effectively capture the complexities of mental health care-seeking behaviors and stages, they do not examine how individuals interact with specific health resources and technologies. Moreover, these models are primarily based on general population studies, leaving unclear how marginalized identities shape care-seeking practices and resource engagement.

2.2 Marginalized Populations' Care-Seeking Behaviors

HCI, CSCW, and ICT4D scholars have increasingly focused their attention on marginality and the empowerment of society's most vulnerable populations [21, 59, 60, 89]. Marginalization manifests in multiple dimensions, encompassing economic hardship, gender discrimination, ethnic exclusion, geographical isolation, educational disparities, and political disenfranchisement [57]. These structural inequalities profoundly shape individuals' aspirations and life trajectories, with intersecting identities of race, class, gender, and disability [19] creating complex barriers that diminish both the intention and capacity to seek care [17].

The reality of limited resources presents immediate and tangible barriers for marginalized populations seeking healthcare services. Evidence demonstrates that individuals in remote rural areas face reduced access to essential services, such as helplines [63], and utilize significantly fewer health services [85]. This perception of resource scarcity may also contribute to a deeper pattern of symptom normalization and treatment avoidance [72]. For example, Pendse et al. [61] found that online mental health narratives from people in low-resource areas tend to emphasize somatic expressions of distress through body-focused language and have their experiences stigmatized and invalidated. Such evidence suggests that the impact of marginality extends beyond mere resource limitations, manifesting in more nuanced ways in people's identity perception and throughout the care-seeking process.

HCI research has documented the complex interactions between individuals with marginalized identities and mental health technologies [55], revealing significant cultural and social barriers to care. For instance, Bhattacharjee et al. [3] illuminated how Indian users express profound skepticism toward talk therapy rooted in Western psychological paradigms, highlighting fundamental cultural disconnects in mental health interventions. Studies also demonstrated how perceived discrimination based on marginalized identities actively deters individuals from seeking clinical care [28, 47]. These negative experiences create lasting psychological barriers to healthcare access [45], initiating self-reinforcing cycles that further alienate individuals from mental health resources. The impact of these barriers extends into digital spaces, as Feuston and Piper [23] documented how social media algorithms and platform affordances shape narratives around mental illness. These technological mechanisms, influenced by community norms that equate mental illness with deviance, inadvertently amplify and perpetuate stigma in online spaces, creating additional obstacles to care-seeking behaviors.

While existing research has examined marginalized populations' interactions with specific technologies, there remains a critical gap in understanding how individuals with marginalized identities navigate their broader care-seeking journey and engage with diverse available resources [69]. This comprehensive understanding is crucial for developing culturally sensitive designs that align with ecological systems [7], cultural values [43], and community preferences [19].

2.3 Ecological Approach for Technology-Mediated Care-Seeking

Recent HCI scholarship has expanded beyond studying users' interactions with isolated personal information systems to examine them within complex social ecologies [53, 56, 74]. For example, Wong et al. [87] examined how workplace environments shape and limit engineers' engagement with mental health resources. Theofanopoulou and Slovak [79] studied tangible toys as mediators for parent-child emotional communication in family settings. Bronfenbrenner's social systems framework [6] has been applied in such HCI research to examine the layered context of health resource interactions. Researchers have used its components—microsystem, macrosystem, and chronosystem—to study social influences on technology-mediated mental health management [7, 53, 78].

This ecological perspective has particular relevance for marginalized populations who face distinct socio-economic considerations [33]. For instance, Tachtler et al. [78] demonstrated how macro-level factors, particularly resettlement policies, constrain unaccompanied migrant youths' engagement with mental health mobile applications. Our study extends this ecological approach to examine how marginalized young adults navigate mental health resources within their lived contexts.

We draw on Social Ecological Theory [77] to examine young adults' mental health navigation. The framework outlined by McLeroy et al. [50] identified five levels affecting health behavior: individual characteristics, social relationships, organizational factors, community characteristics, and societal factors including physical, social, and political environments [49]. This framework has been used to study individuals' interactions with diverse health resources and technologies [70, 82]. Among its variants, we use the version by [11] as a pragmatic framework, which is detailed in 3.3 to analyze our participants' resource interactions.

Building on this ecological perspective, our study investigates marginalized young adults' care-seeking practices across digital platforms, institutional services, and informal support networks. By examining how they choose, access, and perceive these resources within their socioeconomic constraints, we aim to identify barriers and enablers in the current care ecosystem that can inform the design of mental health technologies and services for marginalized young adults [10, 86].

3 METHODS

We adopted in-depth interviews to understand the lived experiences of young adults' mental health help-seeking practices.

3.1 Recruitment and participants

We disseminated recruitment messages on the listserv of a southern university in America as previous studies suggested that university students face high and multifaceted pressure [30]. To recruit young adults from diverse backgrounds, we sourced participants through Reddit, aiming to include individuals from marginalized communities who are often underrepresented in university-based studies. Following Reddit's community guidelines, we posted recruitment messages in 12 regional marketplace and job-focused subreddits with large memberships across the United States, including r/SanAntonioJobs and r/NewOrleansMarketplace.

Interested participants ages 18-25 were invited to finish a screening questionnaire where we asked for their demographic information, such as age, gender, education, marginalized identities (e.g., first-generation college students, low-income, LGBTQ+, race minority), mental illness diagnosis, help-seeking practices (e.g., previously used resources and technologies such as family, friends, social media, and apps), and an open-ended question about the most recent/impressive mental health help-seeking experience. We also include the PHQ-4 [36], a validated screening tool for depression and anxiety to get a rough estimate of their current mental health status. We intentionally chose participants to maximize the sample demographic diversity and mental health status. Following the theoretical sampling strategy [12], our recruitment process proceeded in parallel with data analysis to include participants with diverse help-seeking practices, including varied preferred sources and both satisfying and dissatisfying experiences. For instance, we noticed the prominent role of resources actively reaching out to marginalized young adults based on the first twelve interviews. To collect more cases and further understand the care-outreaching, the team carefully selected subsequent participants who reported relevant experiences in the screening survey and invited them to join the interview.

As shown in Table 1, 18 participants were interviewed (7 females, 9 males, and 2 trans/non-binary; 6 White, 5 Black, 4 Asian, and 3 Hispanic). Half of the participants were still in university at the time of the interviews, 6 had graduated from colleges, and 3 had no college education. Thirteen participants had formal diagnoses of mental illness, mainly depression and anxiety. Their anxiety and depression status indicated by PHQ-4 ranged from mild (3-5), moderate (6-8) to severe (9-12).

3.2 Interview Procedure

The interviews averaged a duration of 1 hour and 20 minutes and each interviewee was compensated with a \$30 Amazon gift card. The interviews took place virtually via Zoom from January to September 2023. Before each interview, we introduced the scope of this study and answered any questions concerning the informed consent form. We also assured the participants that the interviews would be anonymous, confidential, and non-judgmental, encouraging them to be as open as they wanted.

We began the interviews by asking participants' current jobs, residential areas, and other background information. Then, the interviewer asked participants to describe their current mental health status. Example questions included, "Could you tell me about your mental health concerns?" "When did you first notice that?" and "How did it evolve over time?"

We utilized **visual elicitation techniques** for a richer exploration of their (non-)help-seeking journeys [13]. When participants started to dive into specific story-telling, we asked them to draw their experiences of when, where, and how they sought help. Participants were encouraged to take

Table 1. Participants Information

| ID | Age | Sex | Education | Race | Diagnosis | PHQ4 | Resources Used in Mental Health Help-seeking |
|-----|-----|-------|--------------------|----------|---------------------------------|------|--------------------------------------------------------------------------------------------------------------------------------------------------|
| P01 | 20 | F | Some college | Black | depression | 5 | professional help, search engine, friend, family |
| P02 | 24 | M | High school | Black | depression | 8 | social media, online communities, friends, professional help |
| P03 | 20 | F | Some college | Asian | no | 4 | social media, search engine, friends, family, mobile apps |
| P04 | 23 | M | Some college | White | Anxiety | 12 | Family, friends, professionals, online communities, services in the local community |
| P05 | 22 | M | 8 through 11 years | Black | anxiety | 12 | family, friend, professional, social media, online communities, telehealth |
| P06 | 22 | F | Some college | Asian | depression | 8 | telehealth, professional help |
| P07 | 21 | F | Some college | Hispanic | Depression, Anxiety | 4 | Friends, professors, people with similar experiences |
| P08 | 23 | F | College graduate | Asian | Depression | 7 | Family, friends, people with similar experiences, Social media, online mental health forums or communities, Search engines, Teletherapy services |
| P09 | 24 | M | College graduate | White | | 5 | Family, friends, Reddit |
| P10 | 24 | M | College graduate | Asian | no | 4 | Hotline, family |
| P11 | 22 | M | Some college | White | Depression | 10 | Family, professionals, social media, search engine, teletherapy |
| P12 | 20 | F | Some college | White | Depression | 12 | Family, professionals, people with similar experiences, social media, online communities, teletherapy, hotlines, search engines |
| P13 | 22 | M | Some college | White | Depression | 9 | Family, friends, search engines, mobile applications |
| P14 | 21 | F | College graduate | Hispanic | Depression | 5 | Professionals, online communities |
| P15 | 25 | Trans | Some college | Hispanic | Depression, PTSD, anxiety, ADHD | 11 | Family, professionals, social media, search engine |
| P16 | 24 | Trans | College graduate | Black | Depression | 8 | friends, professional, stranger |
| P17 | 21 | M | High school | Black | no | 6 | Family, friends, professionals, social media, search engine, teletherapy, services in the local community |
| P18 | 22 | M | College graduate | White | no | 10 | Family, friends, social media, mobile apps, services in local communities |

Notes: PTSD: Post-traumatic stress disorder; ADHD: Attention-deficit/hyperactivity disorder

some time to recall their experiences and present their stories in any visualization format. Our participants drew their help-seeking experiences from different perspectives, shown in Fig 1. The interviewer then asked follow-up questions based on the drawings to elicit details, for example, "What did you do to cope with [a specific event] in the drawing?" "Did you talk with any people (e.g., family, friends) or use any technology (e.g., social media groups, mobile apps, hotlines)?" and "Why did you decide to do that?" In this process, the interviewer paid special attention to how they perceived and used different resources to seek help. The interviewer also encouraged participants to continually add contexts to the drawings as they narrated. Participants were instructed to focus the camera on the drawing to share with the interviewer during this process.

After the interview, we asked participants to reflect on their experiences and evaluate their satisfaction with each mentioned resource using a five-point scale (1- very unsatisfied; 5- very satisfied). We then invited them to share the most satisfying and unsatisfying experiences, the challenges they encountered during help-seeking, and the ideal help they wished to get.

Ethical concerns This study was approved by the University Institutional Review Board (IRB). To protect the safety of our participants, in the informed consent form and at the beginning of interviews, we reminded the participants that they should feel free to take breaks during the interview and that they could exit the interview at any time. During the interview process, we kept sensitive to participants' emotional changes when difficult experiences were disclosed and checked whether they would like to continue when negative emotions were observed [22]. After the interview, we provided a list of mental health resources to the participants for future use, which was considered helpful and appreciated by many participants.

3.3 Data Analysis

All the interviews were conducted by the first author and were audio recorded and transcribed. The interviewer wrote debriefs immediately after each interview. We used iterative open coding and axial coding methods [16] to systematically analyze the transcripts. The analysis was assisted by using NVivo, a qualitative content analysis software. This iterative process continued until




Fig. 1. Examples of Journey Maps Drawn by Participants. P02 and P07 presented their help-seeking timeline, P09 demonstrated his resources network, and P10 quantified his satisfaction with each resource using self-defined mathematic formulas.

the twelfth interview, at which point the codes reached a stable state, forming the foundation of our initial open coding schema. The open coding schema includes codes about resource type (e.g., technologies, family, professionals), support type (e.g., informational support, distraction), and challenges (e.g., high-cost, hard to talk).

We followed the Social Ecological Theory [77] and categorized these resources into four levels: technological, interpersonal, community, and societal levels. The technological level emerged as a new category through our data analysis, while the other three levels were previously discussed in existing literature [71, 77, 80].

Using this framework, we began axial coding to reorganize codes and generate themes and sub-themes related to participants' practices and challenges in seeking help. The two authors convened weekly meetings to review new interview debriefs, reflect on emerging themes in coding, compare codes with existing literature, and document these reflections in notes and memos. Through discussions, we purposely selected the next interviewees to further test and refine the framework. For instance, since societal resources were less frequently mentioned by earlier participants, we deliberately chose participants who utilized societal resources, focusing on how they defined, accessed, and perceived various societal services in comparison to resources at other levels. The initial open code "support type" evolved into the theme "support mechanisms" with three subthemes in axial coding. In addition, we systematically mapped resources used by participants harnessing the Socio-technical Ecosystem Framework, identifying two types of support systems, various pathways, and barriers in help-seeking.

3.4 Positionality

Our research on mental health help-seeking behaviors and resource utilization is part of a broader effort to support young adults from diverse backgrounds through innovative technological solutions and social support. This work is informed by the lead author's lived experience with mental illness and involvement in peer support groups for young adults facing mental health challenges.

3.5 Limitations

Our study only included 18 participants who were young adults based in the U.S. who were willing to talk about their mental health concerns and help-seeking processes. They might also have less self-stigma. Thus, we acknowledge that their help-seeking behaviors could not represent all young adults, particularly those who did not seek help or were unwilling to talk about their experiences. Future research can examine the non-help-seeking behaviors of young adults.

4 FINDINGS

To understand how marginalized young adults navigate their care-seeking journey (RQ1), we mapped participants' interactions with healthcare resources, revealing complex relationships between marginalized identity, mental health understanding, and care-seeking behaviors. In examining how resources facilitate or hinder care-seeking (RQ2), we identified pivotal "care encounters" that can catalyze more proactive health-seeking behaviors. We also analyze the challenges within these encounters that may hinder participants' care-seeking.

4.1 Participants' Resource Use Practices

Our participants represent a diverse range of backgrounds and varying self-identified marginality. Seven participants identified as first-generation college students, eight identified as racial or gender minorities, seven reported coming from low-income backgrounds, and three reported other forms of marginality, such as disabilities or recent immigration. Notably, five participants reported experiencing intersectional marginality.

All of our participants' care-seeking journey spanned more than a year and they shared both positive and negative experiences with different resources. In Figure 2, we map their satisfaction with each resource they have interacted with based on their responses to the interview question: "To what extent are you satisfied with [a specific resource]?"

As mentioned in Section 3.3, we followed the Social Ecological Theory [77] and categorized these resources into four levels. The *technological level* includes search engines, online forums, video-sharing platforms, and mobile apps. These resources were frequently mentioned by participants and most of the use experiences were satisfying, but it should be noted that these resources had relatively small effects on participants. The *interpersonal level* consists of participants' interpersonal ties. Except for P01, all other participants at least tried to seek help from family, friends, relatives, or partners, although six of them got neutral or unsatisfying feedback. The *community level* contains all resources provided by entities that participants are members of, such as pre-college education institutions, universities, companies, and local neighborhoods. 11/18 participants tried community-level resources and six participants had varying degrees of satisfaction. The *societal level* contains non-profit institutions and professional care including professional help from medical systems, the National Lifeline, and governmental institutions. Only 8 participants utilized these resources.

As shown in Figure 2, seven participants only received satisfying support from a single level of resources (i.e. single-level care). Among them, three (P09, P13, and P15) primarily relied on technological solutions, two (P10 and P18) mainly sought emotional support from family members, and one (P08) only used resources from her community, the university mental health services.




Fig. 2. Participants' Interactions with Resources across Mental Health Care-Seeking Journey. Single Level Care denotes that participants have "satisfied" or "very satisfied" experiences with resources at one level; Multi-Level Care means that they have "satisfied" or "very satisfied" experiences with more than one type of resources.

Notably, one participant (P01) did not report any significant experiences with care resources. The remaining eleven participants were able to navigate and obtain care from multiple levels of resources (i.e., multi-level care). We elaborate on participants' navigation of the socio-technical ecosystem in the following sections.

4.2 Passive Behavioral Pattern of Marginalized Young Adults' Care-Seeking

Our analysis reveals participants' approach in dealing with mental health concerns can be characterized as being passive. These individuals, shaped by their unique experiences and backgrounds, frequently engage in behaviors that delay or avoid the active pursuit of mental health support, that seemed to be rooted in the complex interplay of cultural, personal, and societal factors.

4.2.1 Reality: Social Isolation. For many participants, marginalization extends beyond mental health concerns to encompass broader social isolation where participants don't have access to various types of care resources. Some participants face extreme situations, as illustrated by P02:

when I was 17 years old, my parents separated and I was I had no place to live. I was forced to live with my uncle, who was very abusive to me. It was at that point I developed some mental concerns because of the abuse now I'm being detained being spoken many times being denied food. It's just been so tough for me, you know? I just felt very alone. It was the lowest point of my life. It's just terrible. It's just awful. I was just like, The Dark Side of the Moon you know? (P02)

Others struggle with the unavailability of family support. P03 shared:

It's really busy with school and with family. My mom is struggling with back issues. My grandma, she has Psoriasis, so she has to get shots. My dad had heart surgery last summer. And then the only one that is currently able to help is my eldest sister there, who is now a single mom starting to date a new guy and it's all a struggle. So I try to be at least talking to them. (P03)

Social disparities they are experiencing further exacerbate these challenges, as expressed by P18,

I have a lot of friends that are rich. And they come from a rich family. And it goes around the go with yourself. But I can join them because I always feel I don't belong... discrimination and some other things that might. Of the ceiling. Of the So if one's information, get, so, it's kind of one, I think, they'll use it again somewhere. (P18)

4.2.2 Perception: Low aspirations. A prevalent lack of aspiration for care was observed among many participants, including devaluing the importance of mental health and disbelief in professional help. Cultural influences were one force that perpetuates such internalizations, as one participant shared that “*Mental health isn't really a thing in Asian communities. It's not really like shown by parents. I'm sure that there was one time when I was younger that I was sad and then my mom or dad was just deal with it by yourself. And then that's what I do. I deal with it internally. (P01)*”

The low aspiration also frequently stemmed from a deep-seated disbelief in institutional-level care provisions or negative experiences with organizations. P02 expressed skepticism about governmental resources: “*because you don't see these things are gonna happen. This is from the government, so the question is that these resources are very limited and you don't think you probably can get this. (P02)*” The inaccessibility of information further exacerbates this issue, as noted by P15, “*they [institutions] did not make the information accessible in a way that you can really get the support. (P15)*”

4.2.3 Behavior: Waiting and concealing. Marginalized young adults often conceal their mental health concerns and adopt a passive stance toward seeking help. They frequently wait for what they perceive as the "right opportunity" to address their mental health issues. This ambiguous pattern is common, as exemplified by P10 who expressed that he would treat his occasional depressive episodes more seriously "later" but did not have a concrete plan, “*I mean, I might tell my parents when it's the right time.*” Concealing mental health situations might stem from a view that mental health struggles are personal challenges to be overcome independently, as P03 articulated: “*I think it really also is important to have sort of a self-serving with mental health because you're with yourself for the rest of your life, right? Yes, it's important to have a support system externally, but internally is more important. (P03)*”

4.3 The "Care Encounters" as Moments of Change

While participants' care-seeking behaviors may appear passive, they reflect a nuanced approach of receptive anticipation rather than active avoidance of care resources. Our analysis revealed "care encounters" - serendipitous moments when participants discovered or got exposed to previously unimagined resources. We identified three distinct types of encounters that transformed participants' relationships with care: those providing tangible assistance, those offering supportive narratives, and those facilitating social connections. These encounters helped marginalized young adults expand their aspirations for care and overcome their initial resource limitations. We further examined the affordances of the resource ecosystem that facilitated these transformative encounters.

4.3.1 Encountering Tangible Assistance. The most direct and impactful encounters that reshape participants' perceptions and behavior of care-seeking occur through tangible assistance, such as free therapy sessions, performance accommodations, and hospitalization services. We identified two circumstances that facilitate participants' engagement with tangible assistance: accessibility-driven encounters, where designated resource availability motivates engagement, and emergency-driven encounters, which occur during crisis situations.

Designated Accessibility. Beyond mere availability, designated accessibility refers to how care resources thoughtfully woven into the fabric of participants' daily lives became more inviting and attainable. P08's experience exemplifies this transition in perception and behavior. During the

isolating period of the COVID-19 pandemic, a series of email reminders about depression awareness became her unexpected lifeline:

I think it was like halfway through Covid, I received consecutive emails day by day, saying to focus on your mental health. That was when I felt all the stressors were hitting on me all at once and I wondered why didn't I have a break from school? Let me, just, attend one of these Zoom Meetings, and see what they are about. Let me just check these resources out. It was during that time that I realized that I probably need to get some help because these are the symptoms.

Following these initial encounters, P08 utilized the free therapy sessions introduced in the workshops and emails—a pathway also taken by four other participants.

For instance, P06 benefited from a reduced class load; P07 was offered extended tuition payment options; and P08 maintained access to free therapy sessions even after graduation. These accommodations strengthened the incentive for care-seeking, demonstrating that inclusiveness extends beyond vague promises. Reflecting on this experience, P08 expressed surprise at the abundance of available resources: “[...]I wasn't previously aware that there are so many resources out there possible to help you, and was just waiting for, like reach out and stuff, until I was told I was offered it.”

Reactive interventions in emergencies. Reactive interventions which refer to situation-triggered responses to acute mental health needs, play a crucial role in challenging and changing participants' biases or negative attitudes toward care-seeking. These interventions are particularly vital during emergencies such as suicidal ideation or severe mood episodes. P07's experience with the university crisis line exemplifies the comprehensive approach taken by community resources. Upon learning about her self-harming tendencies, the staff mobilized a multi-faceted response: “*I stayed in the hospital for two days...they diagnosed me with a major depressive disorder and generalized anxiety disorder, and then that's when they gave me medications to start to treat both of those.*” This intervention not only provided immediate care but also set P07 on a path of ongoing treatment and support. It demonstrates how active interventions can bridge the gap between the awareness of resources and the actual engagement with mental health services, especially in critical situations.

Besides the community-provided mental health services, P16's story illustrates the proactive role of interpersonal relationships during a crisis. After both of her parents passed away, P16 isolated herself at home for two months, unaware that she might be experiencing depression. Her friends' persistent check-ins culminated in a crucial intervention:

This was too much for me. They came in when I lost it during that time. They took me to the hospital, checked with the hospital, and showed that I really needed this [professional help]. Then they decided that I needed to change the environment and of course, I listened to them. They are my support network and I tell everyone to find your network. This is what truly supports you no matter what bad happens.

4.3.2 Encounter Supportive Discourses. Supportive discourses can open conversations and messaging that normalize mental health discussions and validate care-seeking behaviors. Participants frequently mentioned encountering these conversations in spaces that created supportive environments for mental health narratives, which, often gradually, mitigate their perceived stigmas of mental health and facilitate their further engagement with mental health resources.

Inclusive Atmosphere. As mentioned in Section 4.1, many participants faced stigmas due to their familial, cultural, or religious backgrounds. However, some eventually encountered inclusive environments and spaces that actively welcomed conversations around mental health experiences.

The university campus was frequently mentioned as a microcosm of this awareness-raising ecosystem. For instance, P10 and P13's observations of omnipresent hotline signs in hallways,

offices, and dining halls paint a picture of an environment saturated with care-seeking encouragement. Additionally, participants reported finding such spaces across various channels, from digital platforms like emails (P08, P16) and social media (P11) to physical spaces hosting awareness events (P15, P06). These messages create a constant, gentle nudge toward mental health consciousness, normalizing conversations around mental health and subtly reinforcing the availability of support. These inclusive encounters also facilitated discussions that were previously deemed impossible. P07, unable to grieve her grandfather's death within her family setting, finally found an opportunity to process her grief in a university event that had been held off since middle school.

Intimate Conversations. One major reason behind the waiting and concealing behaviors among marginalized young adults, mentioned by six participants in our study, was the challenge of initiating open conversations about mental health with family and friends. We observed a delicate tension between their desire for support and the difficulty in articulating their needs. P13's reflection captures this struggle: "*I mean just it's hard to talk about it...I'm not good at talking to people about the hard stuff.*" In such situations, participants found breakthrough moments when family members and friends took the initiative to start these deeper conversations. These proactive gestures often opened doors to meaningful support. P14 shared how her mother's willingness to discuss her depression not only aided her recovery journey but also strengthened their relationship: "*She was like, I was really scared you would do something... I feel like if she ever saw me in such a sad place... that she can understand, she knew about my situation, at least the general situations.*"

4.3.3 Encountering Social Connections. Another crucial type of encounter involves people who truly empower care-seeking. These relationships form a vital layer of care, providing comfort and understanding in navigating mental health challenges.

New Friends from In-person Events. For many marginalized young adults, pivotal moments arise when they connect with peers, friends, or even strangers who offer transformative support, both immediate and long-lasting. P12 found solace in a therapist whose care evolved into a nurturing paternal connection: "*I feel connected with him. He talks to me like a daughter because he has a daughter like me. I enjoyed going to play with his family and his children.*" P15's journey was transformed by a sponsor he met at a peer-support group who became like family, extending both emotional and financial support for treatment. Sometimes, the most profound connections emerge from chance encounters, as P16 discovered during a moment of crisis outside a shopping mall:

She's a stranger, total stranger... I was just getting out of the shopping mall. That's when I broke down. She was like, oh, what's wrong? Can you sit down? I cried a lot, cried, cried and talking and talking and talking and she was there she listened... I felt really very free with her because she was she was a She give me, that, atmosphere for listening.

Online groups. The development of supportive relationships extends beyond immediate circles to online communities, where individuals find connections based on shared experiences. The digital landscape further expands these information-sharing possibilities. Online communities provide platforms for reading others' stories, commenting on posts, and sharing personal experiences. For some, like P05 and P18, these "online friends" even surpass "real friends" in terms of support and understanding. P02's engagement with Facebook groups, where she formed meaningful connections with two online friends, exemplifies this phenomenon. These digital spaces become forums for direct exchange of information about symptoms, potential causes, and coping strategies. The value of these online communities is particularly evident in P12's reflection: "*I will say this online group is safer because the online friends have seen their issues but, the majority of my physical friends don't experience what I'm experiencing.*" This sentiment underscores the unique comfort found

in communities of shared experience, where understanding comes from firsthand knowledge of similar struggles.

4.4 Challenges in the care encounters as transformative moments

4.2 illustrates cases of care encounters that successfully lead to attitudinal and behavioral changes towards more proactive care-seeking, however, we also observe challenges in care encounters that fail to transform marginalized young adults' behaviors. We first summarize the pragmatic barriers at each level in Table 2 and demonstrate the fundamental challenges regarding trust forging and agency raising.

Table 2. Barriers of "Care Encounters" at Each Type of Resources

| Resource Category | Barriers | Examples |
|-------------------|---------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Technological | Aversion of technology | <i>"The search engines, these, online things are too general. They are helpful when you search for a road trip, but it's not the mental health thing"</i> (P10). |
| | Emotional frustration | <i>"having to dig for it can be a little bit demoralizing. I just want instant gratification, but when your brain's all cluttered, the last thing you really want is to have to fight for help."</i> (P15) |
| | Lack experience-based personalization | <i>"I get more worried when I look for it because there's just so much information out there... this information on Google might just be like for everybody. I get more anxious because I can't find information that is tailored to my needs."</i> (P08) |
| Interpersonal | Difficulties in disclosure | <i>"My family don't really talk about things [mental health related] like that."</i> (P03) |
| Community | Varying professionalism | <i>"there are so many people talking while they are not qualified to talk"</i> (P14) |
| | Temporality | <i>"It can be hard when you go from university to an external institution because you have to find one that matches your insurance. The first time, I called at least 10 or so people before I got the practitioner. They were like, 'We are not taking patients or we're not taking certain types of patients.' The whole process can be lengthy and challenging".</i> (P01) |
| | Varying service investment | P15's university provides free and unlimited therapy sessions, while P07's university imposes a three-session limit. |
| Societal | Compatibility | <i>"the first one I had was just really weird and would like tell me her personal problems. And then the second one... she wasn't helpful."</i> (P06) |
| | Impersonal | <i>"She kept talking about sadness and stuff. And I tried to tell her, I'm not sad, I'm tired. I stopped seeing her because I felt like she wasn't really listening to me but just reacting."</i> (P01) |

4.4.1 Trust in Resources. Our analysis reveals complex barriers to trust-building between marginalized populations and care resources.

Marginalized populations' existing distrust in care resources can be further reinforced by negative experiences. When participants used web searches to find mental health resources, poorly designed interfaces often undermined their initial attempts to build trust in these services. For instance, P15, seeking free therapeutic services due to financial constraints, encountered a labyrinthine web of poorly designed websites. She noted, *"every time I've looked it up, it seemed inaccessible. Not every website is very user-friendly. You have to go digging and digging to find what you need to find. And when I'm frustrated and feeling like I need help, that's the last thing I want to do."* Her subsequent

analysis revealed a deeper systemic issue: “*Honestly, I’ve studied how Google search comes up. Results from Google are not mainly from your local clinics. It’s people that pay more money for their website to show up first...But they just look like businesses.*”

Beyond accessibility challenges, trust deficits manifest in concerns about efficacy and cultural compatibility. Participants, particularly those with intersectional marginalized identities, expressed strong preferences for providers who share their lived experiences. P01’s experience exemplifies this challenge: “*I tried to find a therapist who is also a black woman, but I did not see any on the website.*” This scarcity of representation in care providers reinforces existing barriers to trust and engagement, highlighting how identity-based needs remain unmet in current care infrastructures.

The online mental health communities, while intended to facilitate support, can inadvertently amplify feelings of marginalization when users encounter content misaligned with their experiences. P05’s poignant account illustrates this disconnect: “*I’m sorry, mine [situation] was going up for two or three weeks. Most of them are like coincidences. So I feel like I was tragic and traumatized because I was being attacked and my friend was beaten to death.*” When confronted with posts about relatively minor stressors while processing severe trauma, participants like P05 reported feeling further alienated rather than supported. This experiential mismatch not only undermined their engagement with online communities but also eroded their trust in available mental health resources more broadly.

4.4.2 Agency Empowerment. Our analysis reveals how different types of mental health resources vary in their ability to facilitate and sustain agency among marginalized young adults in their care-seeking journey. This variation manifests across societal, community, and technological resource levels, each presenting distinct challenges and opportunities for agency development.

Societal-level resources, while offering professional help, often place a substantial burden on individuals’ agency. Only six participants successfully navigated these resources, with many citing overwhelming requirements for access. As P08 explained: “*Because I felt I could not handle it alone anymore and they were the closest and had anonymity.*” This suggests that societal resources often become a last resort, accessed only when individuals feel compelled by severe circumstances rather than through proactive choice, indicating a reactive rather than empowering approach to agency.

Community-level resources present a complex dynamic where agency support varies significantly across institutions. While some community entities actively facilitate care-seeking, others demonstrate limited investment in mental health support. For instance, P13, a sports official, described the absence of workplace mental health resources: “*There’s no training on how to handle the stress... we’re just expected to deal with it.*” Even within similar institutional contexts, approaches to agency empowerment differ markedly. P05’s university demonstrated minimal proactive engagement, while P10’s institution maintained sustained mental health initiatives through frequent mental health awareness events.

Technological resources, despite their widespread use among participants, primarily served an auxiliary role in agency development. While participants frequently initiated their help-seeking journey through technological platforms, using them as extensions of self-coping through information gathering and peer connection, these resources often proved insufficient as standalone support. Notably, four participants who initially relied exclusively on technological resources ultimately transitioned to other forms of support, suggesting technology’s limitations in sustaining long-term agency in care-seeking. This pattern indicates that while technology can facilitate initial steps toward agency, it may not provide the comprehensive support needed for sustained empowerment in mental health care-seeking.

5 DISCUSSION

Our study contributes to CSCW health literature by revealing how mental health care-seeking practices among young adults with diverse marginalized identities are shaped by their identity perceptions and lived realities. While many participants initially showed passive care-seeking patterns, we identified potential pathways toward proactive health engagement through transformative "care encounters" that helped them transcend limited resource endowments. These encounters served as catalysts for change, though we also documented instances where they failed to initiate behavioral changes. Our findings inform future technology design efforts aimed at empowering vulnerable populations by moving beyond access-focused solutions to address the complex interplay between identity and healthcare engagement.

5.1 Seeking care from the margins: From passive to proactive

This study follows CSCW scholars' mission to care for people on the margin [14] and specifically focuses on how marginalized identities influence the way people seek care.

5.1.1 Marginalized Identity and Passive Care-Seeking Patterns. Prior research has established statistical correlations between marginalized identities (e.g., racial and cultural minorities) and low health resource utilization rates [28]. While some studies attribute this reduced resource use primarily to material constraints within these communities, Pendse et al. [61] revealed how resource scarcity can fundamentally reshape individuals' somatic understanding of mental health, suggesting that marginalization's impact extends beyond mere resource constraints.

Our findings advance this understanding by illuminating how individuals' internalized perceptions of marginalization can suppress care-seeking aspirations, creating barriers that persist even when resources become available. For instance, multiple participants remained inert towards university mental health services despite their accessibility. This internalized skepticism was exemplified by P12's dismissive attitude: "*it's something they [the university] have to do... I don't think they will be helpful*". Such responses reveal how marginalized individuals may interpret institutional support as merely performative rather than genuinely beneficial, contributing to their reluctance to engage with available resources.

In addition to previous research examining how cultural background and racial identities influence care-seeking behaviors [72, 75], our analysis reveals additional dimensions of marginalization that critically impact mental health resource utilization. Participants with unconcealed LGBTQ+ identity, experiences of homelessness, or unemployment often faced compound barriers due to their disconnection from traditional support networks and community resources typically tied to stable affiliations. For example, P15, who identified as LGBTQ+, low-income, and disabled, described feeling "helpless" from care resources, noting how their intersecting identities created multiple layers of disconnection from support systems.

Importantly, our analysis distinguishes these passive care-seeking behaviors, rooted in experiences of marginalization, from active resistance stemming from cultural or religious beliefs [3]. While appearing outwardly resistant, often harbor latent hope for potential change and care access. For instance, P06's experience illustrates this underlying receptivity: despite initially expressing reluctance toward seeking care, she scheduled a therapy session upon learning about free university services. This pattern of dormant hope manifesting when opportunities arise was consistent across multiple participants, suggesting that passive waiting often masks an openness to care when barriers are reduced or removed. This insight offers crucial implications for how mental health resources and outreach efforts might be better structured to engage marginalized populations which we will discuss in 5.2.

5.1.2 "Care Encounters" as Catalysts of Aspiration Changes. Our findings reveal how serendipitous "care encounters" serve as pivotal moments in reshaping marginalized young adults' care-seeking trajectories. These encounters manifest through three distinct mechanisms: tangible assistance (e.g., free therapy sessions, academic accommodations, assisted hospitalization), supportive discourse (e.g., normalized conversations about mental health in both community contexts and intimate interpersonal exchanges), and social connection building that fosters meaningful relationships. While previous research has documented attitudinal changes towards care-seeking in digital spaces [66], our study illuminates broader transformation patterns across both online and offline contexts.

Our findings identify that these encounters operate along two temporal dimensions. In acute situations, encounters can catalyze immediate transformation - notably exemplified by P7 and P12's experiences with emergency interventions that, contrary to previous literature's emphasis on negative impacts, served as positive turning points. More commonly, transformation unfolds gradually through accumulated positive interactions that incrementally expand participants' understanding of available resources and possibilities for care, as exemplified by P07's growing comfort in sharing her grief with her grandfather.

Drawing from Park's [57] work on aspiration, our findings demonstrate how care encounters can reshape aspirations for mental healthcare, particularly paramount given marginalized populations typically express limited future aspirations due to structural constraints. For instance, P16's journey from crisis to advocacy illustrates this transformation: "*Now I am a big advocate of mental health and I tell everyone about my story. My friends, they are my support system and I tell people to find your support system.*" Her evolution from experiencing forced hospitalization to actively building a supportive network demonstrates a fundamental shift toward proactive care-seeking. These findings suggest the importance of aspiration-centered design approaches [24, 37, 60] in creating transformative care experiences for marginalized young adults.

5.2 Empower Marginalized Young Adults via the Socio-Technical Care Ecosystem

While many participants described their care encounters as "coincidence" or "luck," our analysis reveals these experiences are systematically enabled by infrastructural designs that structure and facilitate the possibility of such encounters [62]. We discuss the affordances of various resources in the socio-technical care ecosystem and identify critical design implications for improving mental health support systems.

5.2.1 Socio-technical Care Ecosystem Behind "Care Encounters". Following the ecological approach to care-seeking [8, 53, 56, 74], we adopted Social Ecological Theory [77] to examine participants' encounters with four types of resources: societal, community, interpersonal, and technological. Our analysis reveals that the community sources emerge as principal contexts facilitating proactive care-seeking, while societal and technological resources often struggle to achieve similar engagement levels.

The ecological efficacy of community resources proves particularly powerful in empowering young adults with marginalized identities. Our findings highlight how university-based care outreach efforts foster empowerment through two interrelated channels: resource delivery providing accessible mental health support tools and information, while conversation invitations create safe spaces for open dialogue about mental health concerns. This dual approach proves particularly effective in cultivating supportive microenvironments that encourage care-seeking behaviors among marginalized young adults.

Community resources also serve as crucial bridges to societal resources while expanding marginalized young adults' interpersonal networks. Several participants, after positive experiences with university mental health services, decided to continue therapy post-graduation. For instance, P08's

university extended her free therapy sessions while she searched for post-graduation care options. Communities can host events that foster interpersonal relationships, crucial for marginalized young adults who often experience exclusion from their families and lack friends comfortable with mental health discussions. University-managed peer support groups emerged as critical spaces for connection. In one notable case, P17 encountered an alumnus who sponsored all treatment fees, including therapist visits and medication. Such local support groups were perceived by participants (e.g., P07 and P17) as more trustworthy and engaging compared to online peer support groups, enhancing the potential for positive care experiences [27]. On the downside, it is crucial for future study to investigate support for individuals without readily available community resources. The four non-university-affiliated participants in our study—including individuals experiencing homelessness, unemployment, and workplace isolation—underscore the need for support systems beyond university contexts.

Our analysis reveals limitations in technological-level resources' efficacy in initiating active interventions and motivating care-seeking. P16's experience with online portals demonstrates how poorly designed technological interfaces can reinforce feelings of exclusion and diminish care aspirations. The limited engagement with online resources was also observed in previous studies [65], suggests the disconnection of these tools from young adults' daily lives may explain their relatively low ecological validity [52]. These findings emphasize the need for a more integrated approach to mental health support that combines technological solutions with community-based resources while recognizing the crucial role of interpersonal connections in facilitating care encounters.

5.2.2 Empowering through Technologies. Building on research that integrates mental health technologies within young adults' social ecosystems [40, 41, 76], we discuss several key implications that facilitate future researchers and designers to embed and integrate technologies within marginalized young adults' social, cultural, and material environments [27].

Leveraging Low-Technology Solutions for Effective Outreach. While numerous studies demonstrate engagement patterns with mental health self-management applications [2, 84], our research revealed notably limited adoption of these tools among young adults with marginalized identities, with only two out of twelve participants reporting attempted usage. This low adoption rate likely stems from insufficient trust and motivation [9]. In contrast, low-technology interventions, particularly text messages, and notifications [26], have shown promising results with higher acceptability and seamless integration into daily routines [27]. Our findings identified similar low-technology outreach mechanisms as four participants specifically highlighted the effectiveness of university outreach through traditional channels—emails, posters, and social media—in facilitating service utilization. Moreover, three participants discovered the National Crisis Line through prominent search engine results, underscoring the impact of strategic information placement across commonly accessed platforms. These findings suggest that compared to sophisticated mobile applications attempting to address all mental health needs comprehensively, marginalized young adults may respond more favorably to low-tech communication strategies. Such approaches can serve as effective nudges, anchoring critical information and fostering gradual attitudinal shifts toward help-seeking.

Facilitating Technology-Mediated Disclosure in Complicated Interpersonal Relationships. We found five participants struggle with complicated family dynamics that inhibit trust and open discussion of mental health concerns. These individuals often face cultural barriers that create hesitation in disclosing and addressing their mental health needs [15]. Building intimacy and trust through sustained engagement emerges as crucial [18, 73], particularly when traditional family support structures are strained. Technology presents unique opportunities to bridge these interpersonal gaps,

functioning both as an accessibility enabler and a relationship strengthener [58]. The emergence of Large Language Models (LLMs) with enhanced emotion detection capabilities [83] has enabled the development of mediating tools, such as empathetic chatbots [32], which show promise in facilitating difficult conversations. This technological approach resonates particularly well with young adults [35], with mental health professionals confirming its efficacy in raising awareness and managing negative emotions within complex family dynamics [42]. Notably, our study identified alternative platforms, such as gaming environments, as especially effective in creating safe spaces for emotional expression. Four male participants demonstrated this by utilizing gaming platforms to maintain friendships and share personal thoughts away from family tensions [88]. These technology-mediated environments offer alternative pathways for building trust and maintaining connections when traditional face-to-face family interactions prove challenging or emotionally fraught [27].

Build Trust through Identity and Experience-Based Interactions. Our findings reveal that current platforms inadequately address personalization needs, particularly for marginalized users seeking providers with shared lived experiences. For instance, P05's difficulty relating to posts from more resourced individuals highlights the importance of socioeconomic status alignment in care relationships [44]. These insights suggest new directions for recommendation systems and online peer support group design that prioritize identity-based and experience-based connections, as users demonstrate strong engagement with narratives that mirror their personal circumstances [46]. Future research can investigate how to effectively facilitate digital mental health storytelling, given its documented therapeutic benefits [34], especially among young adults [20].

6 CONCLUSION

Through interviews with 18 young adults with diverse marginalized identities, our study reveals how passive care-seeking behaviors are shaped by experiences of marginalization. We identify "care encounters" as critical turning points that can transform passive patterns into proactive engagement through tangible assistance, supportive discourse, and social connections. These findings inform the design of interventions that can intentionally facilitate such transformative encounters, ultimately improving marginalized young adults' access to mental health care.

7 ACKNOWLEDGEMENTS

We extend our sincere gratitude to the participants for their brave and heartful sharing of their experiences and feelings, which has greatly inspired the authors and made this paper possible. We also thank the reviewers for their valuable insights, which helped refine our findings and clarify our contributions.

REFERENCES

- [1] Rebecca Abavi, Allison Branston, Robin Mason, and Janice Du Mont. 2020. An Exploration of Sexual Assault Survivors' Discourse Online on Help-Seeking. *Violence and Victims* 35, 1 (Feb. 2020), 126–140. <https://doi.org/10.1891/0886-6708.VV-D-18-00148>
- [2] Felwah Alqahtani, Andrea Winn, and Rita Orji. 2021. Co-Designing a Mobile App to Improve Mental Health and Well-Being: Focus Group Study. *JMIR Formative Research* 5, 2 (Feb. 2021), e18172. <https://doi.org/10.2196/18172> Company: JMIR Formative Research Distributor: JMIR Formative Research Institution: JMIR Formative Research Label: JMIR Formative Research Publisher: JMIR Publications Inc., Toronto, Canada.
- [3] Ananya Bhattacharjee, Sharifa Sultana, Mohammad Ruhul Amin, Yeshim Iqbal, and Syed Ishtiaque Ahmed. 2023. "What's the Point of Having This Conversation?": From a Telephone Crisis Helpline in Bangladesh to the Decolonization of Mental Health Services. *ACM Journal on Computing and Sustainable Societies* 1, 2 (Dec. 2023), 15:1–15:29. <https://doi.org/10.1145/3616381>
- [4] Kamaldeep Bhui and Dinesh Bhugra. 2002. Mental illness in Black and Asian ethnic minorities: pathways to care and outcomes. *Advances in Psychiatric Treatment* 8, 1 (Jan. 2002), 26–33. <https://doi.org/10.1192/apt.8.1.26>

- [5] Lucy Biddle, Jenny Donovan, Debbie Sharp, and David Gunnell. 2007. Explaining non-help-seeking amongst young adults with mental distress: a dynamic interpretive model of illness behaviour: Illness behaviour among young adults with mental distress. *Sociology of Health & Illness* 29, 7 (Aug. 2007), 983–1002. <https://doi.org/10.1111/j.1467-9566.2007.01030.x>
- [6] Urie Bronfenbrenner. 1979. The ecology of human development: Experiments by nature and design. *Harvard University Press google schola* 2 (1979), 139–163.
- [7] Eleanor R. Burgess, Kathryn E. Ringland, Jennifer Nicholas, Ashley A. Knapp, Jordan Eschler, David C. Mohr, and Madhu C. Reddy. 2019. "I think people are powerful": The Sociality of Individuals Managing Depression. *Proceedings of the ACM on Human-Computer Interaction* 3, CSCW (Nov. 2019), 1–29. <https://doi.org/10.1145/3359143>
- [8] Eleanor R. Burgess, Renwen Zhang, Sindhu Kiranmai Ernala, Jessica L. Feuston, Munmun De Choudhury, Mary Czerwinski, Adrian Aguilera, Stephen M. Schueller, and Madhu C. Reddy. 2021. Technology ecosystems: rethinking resources for mental health. *Interactions* 28, 1 (Jan. 2021), 66–71. <https://doi.org/10.1145/3434564>
- [9] Lauren Burke, Mary John, and Paul Hanna. 2022. A qualitative exploration of how young men in the UK perceive and experience informal help-seeking for mental health difficulties. *Children and Youth Services Review* 137 (June 2022), 106440. <https://doi.org/10.1016/j.childyouth.2022.106440>
- [10] Nuij C, van Ballegooijen W, de Beurs D, de Winter RFP, Gilissen R, O'Connor RC, Smit JH, Kerkhof A, and Riper H. 2022. The feasibility of using smartphone apps as treatment components for depressed suicidal outpatients. *Frontiers in psychiatry* 13 (2022), 971046. <https://doi.org/10.3389/fpsyg.2022.971046>
- [11] Centers for Disease and Control and Prevention. 2022. The Social-Ecological Model: A Framework for Prevention. <https://www.cdc.gov/violenceprevention/about/social-ecologicalmodel.html>
- [12] Kathy Charmaz. 2000. Grounded theory: Objectivist and constructivist methods. *Handbook of qualitative research* 2, 1 (2000), 509–535.
- [13] Annie T. Chen. 2018. Timeline Drawing and the Online Scrapbook: Two Visual Elicitation Techniques for a Richer Exploration of Illness Journeys. *International Journal of Qualitative Methods* 17, 1 (Dec. 2018), 1609406917753207. Publisher: SAGE Publications Inc.
- [14] Ishita Chordia, Leya Breanna Baltaxe-Admony, Ashley Boone, Alyssa Sheehan, Lynn Dombrowski, Christopher A Le Dantec, Kathryn E. Ringland, and Angela D. R. Smith. 2024. Social Justice in HCI: A Systematic Literature Review. In *Proceedings of the CHI Conference on Human Factors in Computing Systems (CHI '24)*. Association for Computing Machinery, New York, NY, USA, 1–33. <https://doi.org/10.1145/3613904.3642704>
- [15] Kyungmi Chung, Hee Young Cho, Young Ran Kim, Kyungun Jhung, Hwa Seon Koo, and Jin Young Park. 2020. Medical Help-Seeking Strategies for Perinatal Women With Obstetric and Mental Health Problems and Changes in Medical Decision Making Based on Online Health Information: Path Analysis. *Journal of Medical Internet Research* 22, 3 (March 2020), e14095. <https://doi.org/10.2196/14095>
- [16] Juliet Corbin and Anselm Strauss. 2014. *Basics of qualitative research: Techniques and procedures for developing grounded theory*. Sage publications.
- [17] Kimberlé Williams Crenshaw. 2013. Mapping the margins: Intersectionality, identity politics, and violence against women of color. In *The public nature of private violence*. Routledge, 93–118. <https://api.taylorfrancis.com/content/chapters/edit/download?identifierName=doi&identifierValue=10.4324/9780203060902-6&type=chapterpdf>
- [18] Thomas Dalsgaard, Mikael B. Skov, Malthe Stougaard, and Bo Thomassen. 2006. Mediated intimacy in families: understanding the relation between children and parents. In *Proceedings of the 2006 conference on Interaction design and children (IDC '06)*. Association for Computing Machinery, New York, NY, USA, 145–152. <https://doi.org/10.1145/1139073.1139110>
- [19] Dipto Das. 2023. Studying Multi-dimensional Marginalization of Identity from Decolonial and Postcolonial Perspectives. In *Companion Publication of the 2023 Conference on Computer Supported Cooperative Work and Social Computing (CSCW '23 Companion)*. Association for Computing Machinery, New York, NY, USA, 437–440. <https://doi.org/10.1145/3584931.3608920>
- [20] Nadia De Vecchi, Amanda Kenny, Virginia Dickson-Swift, and Susan Kidd. 2016. How digital storytelling is used in mental health: A scoping review. *International Journal of Mental Health Nursing* 25, 3 (2016), 183–193.
- [21] Michael A. Devito, Ashley Marie Walker, Jeremy Birnholtz, Kathryn Ringland, Kathryn Macapagal, Ashley Kraus, Sean Munson, Calvin Liang, and Herman Saksono. 2019. Social Technologies for Digital Wellbeing Among Marginalized Communities. In *Companion Publication of the 2019 Conference on Computer Supported Cooperative Work and Social Computing (CSCW '19 Companion)*. Association for Computing Machinery, New York, NY, USA, 449–454. <https://doi.org/10.1145/3311957.3359442>
- [22] Claire Burke Draucker, Donna S. Martsolf, and Candice Poole. 2009. Developing Distress Protocols for Research on Sensitive Topics. *Archives of Psychiatric Nursing* 23, 5 (Oct. 2009), 343–350. <https://doi.org/10.1016/j.apnu.2008.10.008>
- [23] Jessica L. Feuston and Anne Marie Piper. 2019. Everyday Experiences: Small Stories and Mental Illness on Instagram. In *Proceedings of the 2019 CHI Conference on Human Factors in Computing Systems (CHI '19)*. Association for Computing

- Machinery, New York, NY, USA, 1–14. <https://doi.org/10.1145/3290605.3300495>
- [24] Guo Freeman, Jeffrey Bardzell, and Shaowen Bardzell. 2017. Aspirational Design and Messy Democracy: Partisanship, Policy, and Hope in an Asian City. In *Proceedings of the 2017 ACM Conference on Computer Supported Cooperative Work and Social Computing (CSCW '17)*. Association for Computing Machinery, New York, NY, USA, 404–416. <https://doi.org/10.1145/2998181.2998291>
- [25] R. Gater, De B. Almeida E. Sousa, G. Barrientos, J. Caraveo, C. R. Chandrashekhar, M. Dhadphale, D. Goldberg, A. H. Al Kathiri, M. Mubbashar, K. Silhan, D. Thong, F. Torres-Gonzales, and N. Sartorius. 1991. The pathways to psychiatric care: a cross-cultural study. *Psychological Medicine* 21, 3 (Aug. 1991), 761–774. <https://doi.org/10.1017/S003329170002239X>
- [26] Lynn Gitlow, Blaire Janney, Caitlyn Lemery, Julianne McDonald, Bethany McGinley, and Kristin Rabideau. 2019. How college students diagnosed with anxiety disorders and/or depression use everyday technology. *Disability and Rehabilitation: Assistive Technology* 14, 8 (Nov. 2019), 809–816. <https://doi.org/10.1080/17483107.2018.1499143> Publisher: Taylor & Francis _eprint: <https://doi.org/10.1080/17483107.2018.1499143>
- [27] Christine E. Gould, Julia Loup, Eric Kuhn, Sherry A. Beaudreau, Flora Ma, Mary K. Goldstein, Julie Loebach Wetherell, Aimee Marie L. Zapata, Philip Choe, and Ruth O'Hara. 2020. Technology use and preferences for mental health self-management interventions among older veterans. *International Journal of Geriatric Psychiatry* 35, 3 (March 2020), 321–330. <https://doi.org/10.1002/gps.5252>
- [28] Amelia Gulliver, Kathleen M. Griffiths, and Helen Christensen. 2010. Perceived barriers and facilitators to mental health help-seeking in young people: a systematic review. *BMC Psychiatry* 10, 1 (Dec. 2010), 113. <https://doi.org/10.1186/1471-244X-10-113>
- [29] Blake Anthony Hickey, Taryn Chalmers, Phillip Newton, Chin-Teng Lin, David Sibbritt, Craig S. McLachlan, Roderick Clifton-Bligh, John Morley, and Sara Lal. 2021. Smart Devices and Wearable Technologies to Detect and Monitor Mental Health Conditions and Stress: A Systematic Review. *Sensors* 21, 10 (Jan. 2021), 3461. <https://doi.org/10.3390/s21103461> Number: 10 Publisher: Multidisciplinary Digital Publishing Institute.
- [30] Justin Hunt and Daniel Eisenberg. 2010. Mental Health Problems and Help-Seeking Behavior Among College Students. *Journal of Adolescent Health* 46, 1 (Jan. 2010), 3–10. <https://doi.org/10.1016/j.jadohealth.2009.08.008>
- [31] Peter Huxley. 1996. Mental illness in the community: The Goldberg-Huxley model of the pathway to psychiatric care. *Nordic Journal of Psychiatry* 50, sup37 (Jan. 1996), 47–53. <https://doi.org/10.3109/08039489609099730> Publisher: Taylor & Francis _eprint: <https://doi.org/10.3109/08039489609099730>
- [32] Eunkyoung Jo, Daniel A. Epstein, Hyunhoon Jung, and Young-Ho Kim. 2023. Understanding the Benefits and Challenges of Deploying Conversational AI Leveraging Large Language Models for Public Health Intervention. In *Proceedings of the 2023 CHI Conference on Human Factors in Computing Systems*. ACM, Hamburg Germany, 1–16. <https://doi.org/10.1145/3544548.3581503>
- [33] Elizabeth Kazuiunas, Michael S. Klinkman, and Mark S. Ackerman. 2019. Precarious Interventions: Designing for Ecologies of Care. *Proceedings of the ACM on Human-Computer Interaction* 3, CSCW (Nov. 2019), 113:1–113:27. <https://doi.org/10.1145/3359215>
- [34] Deoksoon Kim and Fang Jia. 2020. “Ever wondered what schizophrenia was?”: Students’ digital storytelling about mental disorders. *Journal of Curriculum Studies Research* 2, 2 (2020), 144–169.
- [35] Theodora Koulouri, Robert D. Macredie, and David Olakitan. 2022. Chatbots to Support Young Adults’ Mental Health: An Exploratory Study of Acceptability. *ACM Transactions on Interactive Intelligent Systems* 12, 2 (June 2022), 1–39. <https://doi.org/10.1145/3485874>
- [36] Kurt Kroenke, Robert L. Spitzer, Janet B. W. Williams, and Bernd Löwe. 2009. An Ultra-Brief Screening Scale for Anxiety and Depression: The PHQ-4. *Psychosomatics* 50, 6 (Nov. 2009), 613–621. [https://doi.org/10.1016/S0033-3182\(09\)70864-3](https://doi.org/10.1016/S0033-3182(09)70864-3)
- [37] Neha Kumar, Marisol Wong-Villacres, Naveena Karusala, Aditya Vishwanath, Arkadeep Kumar, and Azra Ismail. 2019. Aspirations-based design. In *Proceedings of the Tenth International Conference on Information and Communication Technologies and Development (ICTD ’19)*. Association for Computing Machinery, New York, NY, USA, 1–11. <https://doi.org/10.1145/3287098.3287117>
- [38] E Megan Lachmar, Andrea K Wittenborn, Katherine W Bogen, and Heather L McCauley. 2017. #MyDepressionLooksLike: Examining Public Discourse About Depression on Twitter. *JMIR MENTAL HEALTH* (2017), 11.
- [39] Daniel G. Lannin, David L. Vogel, Rachel E. Brenner, W. Todd Abraham, and Patrick J. Heath. 2016. Does self-stigma reduce the probability of seeking mental health information? *Journal of Counseling Psychology* 63, 3 (April 2016), 351–358. <https://doi.org/10.1037/cou0000108>
- [40] Emily G. Lattie, Rachel Kornfield, Kathryn E. Ringland, Renwen Zhang, Nathan Winquist, and Madhu Reddy. 2020. Designing Mental Health Technologies that Support the Social Ecosystem of College Students. In *Proceedings of the 2020 CHI Conference on Human Factors in Computing Systems (CHI ’20)*. Association for Computing Machinery, New York, NY, USA, 1–15. <https://doi.org/10.1145/3313831.3376362>

- [41] Trinh Le, Hashini Senaratne, Michael McQuaid, and Gareth W. Tigwell. 2021. Exploring a Multifaceted Framework to Support the Design of Mobile Apps for Self-Regulating Anxiety. In *Extended Abstracts of the 2021 CHI Conference on Human Factors in Computing Systems (CHI EA '21)*. Association for Computing Machinery, New York, NY, USA, 1–7. <https://doi.org/10.1145/3411763.3451645>
- [42] Yi-Chieh Lee, Naomi Yamashita, Yun Huang, and Wai Fu. 2020. "I Hear You, I Feel You": Encouraging Deep Self-disclosure through a Chatbot. In *Proceedings of the 2020 CHI Conference on Human Factors in Computing Systems*. ACM, Honolulu HI USA, 1–12. <https://doi.org/10.1145/3313831.3376175>
- [43] Guo Li, Xiaomu Zhou, Tun Lu, Jiang Yang, and Ning Gu. 2016. SunForum: Understanding Depression in a Chinese Online Community. In *Proceedings of the 19th ACM Conference on Computer-Supported Cooperative Work & Social Computing (CSCW '16)*. Association for Computing Machinery, New York, NY, USA, 515–526. <https://doi.org/10.1145/2818048.2819994>
- [44] Calvin A. Liang, Sean A. Munson, and Julie A. Kientz. 2021. Embracing Four Tensions in Human-Computer Interaction Research with Marginalized People. *ACM Transactions on Computer-Human Interaction* 28, 2 (April 2021), 1–47. <https://doi.org/10.1145/3443686>
- [45] Jiaying Liu and Yan Zhang. 2024. Exploring Young Adults' Mental Health Help-Seeking Journey: Preliminary Findings on Resource Navigation Behavior. *Proceedings of the Association for Information Science and Technology* 61, 1 (2024), 570–575. <https://doi.org/10.1002/pra2.1060> _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/pra2.1060>
- [46] Jiaying Liu, Yan Zhang, and Yeolib Kim. 2023. Consumer Health Information Quality, Credibility, and Trust: An Analysis of Definitions, Measures, and Conceptual Dimensions. In *Proceedings of the 2023 Conference on Human Information Interaction and Retrieval*. ACM, Austin TX USA, 197–210. <https://doi.org/10.1145/3576840.3578331>
- [47] Wenhua Lu, Abigail Todhunter-Reid, Mary Louise Mitsdarffer, Miguel Muñoz-Laboy, Anderson Sungmin Yoon, and Lei Xu. 2021. Barriers and Facilitators for Mental Health Service Use Among Racial/Ethnic Minority Adolescents: A Systematic Review of Literature. *Frontiers in Public Health* 9 (March 2021), 641605. <https://doi.org/10.3389/fpubh.2021.641605>
- [48] Angel Martínez-Hernández, Susan M. DiGiacomo, Natália Carceller-Maicas, Martín Correa-Urquiza, and María Antonia Martorell-Poveda. 2014. Non-professional-help-seeking among young people with depression: a qualitative study. *BMC Psychiatry* 14, 1 (April 2014), 124. <https://doi.org/10.1186/1471-244X-14-124>
- [49] Donna Jo McCloskey, Tabia Henry Akintobi, Ann Bonham, Jennifer Cook, and Tamera Coyne-Beasley. 2011. Principles of Community Engagement (Second Edition). *COMMUNITY ENGAGEMENT* (2011).
- [50] Kenneth R. McLeroy, Daniel Bibreau, Allan Steckler, and Karen Glanz. 1988. An Ecological Perspective on Health Promotion Programs. *Health Education Quarterly* 15, 4 (Dec. 1988), 351–377. <https://doi.org/10.1177/109019818801500401> Publisher: SAGE Publications Inc.
- [51] Noah Miller, Ekaterina R. Stepanova, John Desnoyers-Stewart, Ashu Adhikari, Alexandra Kitson, Patrick Pennefather, Denise Quesnel, Katharina Brauns, Anika Friedl-Werner, Alexander Stahn, and Bernhard E. Riecke. 2023. Awedyssey: Design Tensions in Eliciting Self-transcendent Emotions in Virtual Reality to Support Mental Well-being and Connection. In *Proceedings of the 2023 ACM Designing Interactive Systems Conference (DIS '23)*. Association for Computing Machinery, New York, NY, USA, 189–211. <https://doi.org/10.1145/3563657.3595998>
- [52] David C. Mohr, Ken R. Weingardt, Madhu Reddy, and Stephen M. Schueller. 2017. Three Problems With Current Digital Mental Health Research . . . and Three Things We Can Do About Them. *Psychiatric Services* 68, 5 (May 2017), 427–429. <https://doi.org/10.1176/appi.ps.201600541> Publisher: American Psychiatric Publishing.
- [53] Elizabeth L. Murnane, Tara G. Walker, Beck Tench, Stephen Voida, and Jaime Snyder. 2018. Personal Informatics in Interpersonal Contexts: Towards the Design of Technology that Supports the Social Ecologies of Long-Term Mental Health Management. *Proceedings of the ACM on Human-Computer Interaction* 2, CSCW (Nov. 2018), 127:1–127:27. <https://doi.org/10.1145/3274396>
- [54] National Institute of Mental Health. 2023. Mental Illness. <https://www.nimh.nih.gov/health/statistics/mental-illness>
- [55] Vanessa O Oguamanam, Natalie Hernandez, Rasheeta Chandler, Dominique Guillaume, Kai McKeever, Morgan Allen, Sabreen Mohammed, and Andrea G Parker. 2023. An intersectional look at use of and satisfaction with digital mental health platforms: a survey of perinatal black women. In *Proceedings of the 2023 CHI Conference on Human Factors in Computing Systems*. 1–20.
- [56] Tom Ongwere, Andrew B.L. Berry, Clara Caldeira, Rosa I. Arriaga, Amid Ayobi, Eleanor R. Burgess, Kay Connelly, Patricia Franklin, Andrew D Miller, Aehong Min, and Nervo Verdezoto. 2022. Challenges, Tensions, and Opportunities in Designing Ecosystems to Support the Management of Complex Health Needs. In *Extended Abstracts of the 2022 CHI Conference on Human Factors in Computing Systems (CHI EA '22)*. Association for Computing Machinery, New York, NY, USA, 1–7. <https://doi.org/10.1145/3491101.3503714>
- [57] Joyojeet Pal, Tawfiq Ammari, Ramaswami Mahalingam, Ana Maria Huaita Alfaro, and Meera Lakshmanan. 2013. Marginality, aspiration and accessibility in ICTD. In *Proceedings of the Sixth International Conference on Information and Communication Technologies and Development: Full Papers - Volume 1 (ICTD '13)*. Association for Computing Machinery,

- New York, NY, USA, 68–78. <https://doi.org/10.1145/2516604.2516623>
- [58] Jasper E. Palmier-Claus, Anne Rogers, John Ainsworth, Matt Machin, Christine Barrowclough, Louise Laverty, Emma Barkus, Shitij Kapur, Til Wykes, and Shón W. Lewis. 2013. Integrating mobile-phone based assessment for psychosis into people's everyday lives and clinical care: a qualitative study. *BMC psychiatry* 13 (Jan. 2013), 34. <https://doi.org/10.1186/1471-244X-13-34>
- [59] Angela Pearce. 2020. Socially-Oriented Technologies Attractability Influence on Individuals Academically and Neurologically. In *2020 12th International Conference on Education Technology and Computers*. ACM, London United Kingdom, 165–171. <https://doi.org/10.1145/3436756.3437040>
- [60] Sachin R. Pendse, Naveena Karusala, Divya Siddarth, Pattie Gonsalves, Seema Mehrotra, John A. Naslund, Mamta Sood, Neha Kumar, and Amit Sharma. 2019. Mental health in the global south: challenges and opportunities in HCI for development. In *Proceedings of the 2nd ACM SIGCAS Conference on Computing and Sustainable Societies (COMPASS '19)*. Association for Computing Machinery, New York, NY, USA, 22–36. <https://doi.org/10.1145/3314344.3332483>
- [61] Sachin R. Pendse, Neha Kumar, and Munmun De Choudhury. 2023. Marginalization and the Construction of Mental Illness Narratives Online: Foregrounding Institutions in Technology-Mediated Care. *Proceedings of the ACM on Human-Computer Interaction* 7, CSCW2 (Oct. 2023), 346:1–346:30. <https://doi.org/10.1145/3610195>
- [62] Sachin R. Pendse, Faisal M. Lalani, Mummun De Choudhury, Amit Sharma, and Neha Kumar. 2020. "Like Shock Absorbers": Understanding the Human Infrastructures of Technology-Mediated Mental Health Support. In *Proceedings of the 2020 CHI Conference on Human Factors in Computing Systems (CHI '20)*. Association for Computing Machinery, New York, NY, USA, 1–14. <https://doi.org/10.1145/3313831.3376465>
- [63] Sachin R Pendse, Amit Sharma, Aditya Vashistha, Mummun De Choudhury, and Neha Kumar. 2021. "Can I Not Be Suicidal on a Sunday?": Understanding Technology-Mediated Pathways to Mental Health Support. In *Proceedings of the 2021 CHI Conference on Human Factors in Computing Systems (CHI '21)*. Association for Computing Machinery, New York, NY, USA, 1–16. <https://doi.org/10.1145/3411764.3445410>
- [64] Justin Petelka, Lucy Van Kleunen, Liam Albright, Elizabeth Murnane, Stephen Voids, and Jaime Snyder. 2020. Being (In)Visible: Privacy, Transparency, and Disclosure in the Self-Management of Bipolar Disorder. In *Proceedings of the 2020 CHI Conference on Human Factors in Computing Systems (CHI '20)*. Association for Computing Machinery, New York, NY, USA, 1–14. <https://doi.org/10.1145/3313831.3376573>
- [65] Claudette Pretorius, Darragh McCashin, Naoise Kavanagh, and David Coyle. 2020. Searching for Mental Health: A Mixed-Methods Study of Young People's Online Help-seeking. In *Proceedings of the 2020 CHI Conference on Human Factors in Computing Systems (CHI '20)*. Association for Computing Machinery, New York, NY, USA, 1–13. <https://doi.org/10.1145/3313831.3376328>
- [66] Yada Pruksachatkun, Sachin R. Pendse, and Amit Sharma. 2019. Moments of Change: Analyzing Peer-Based Cognitive Support in Online Mental Health Forums. In *Proceedings of the 2019 CHI Conference on Human Factors in Computing Systems* (Glasgow, Scotland UK) (CHI '19). Association for Computing Machinery, New York, NY, USA, 1–13. <https://doi.org/10.1145/3290605.3300294>
- [67] Debra Rickwood, Frank P. Deane, Coralie J. Wilson, and Joseph Ciarrochi. 2005. Young people's help-seeking for mental health problems. *Australian e-Journal for the Advancement of Mental Health* 4, 3 (Jan. 2005), 218–251. <https://doi.org/10.5172/jamh.4.3.218>
- [68] Debra Rickwood and Thomas. 2012. Conceptual measurement framework for help-seeking for mental health problems. *Psychology Research and Behavior Management* (Dec. 2012), 173. <https://doi.org/10.2147/PRBM.S38707>
- [69] Fiona Robards, Melissa Kang, Tim Usherwood, and Lena Sanci. 2018. How Marginalized Young People Access, Engage With, and Navigate Health-Care Systems in the Digital Age: Systematic Review. *Journal of Adolescent Health* 62, 4 (April 2018), 365–381. <https://doi.org/10.1016/j.jadohealth.2017.10.018>
- [70] James F. Sallis, Neville Owen, and Edwin B. Fisher. 2008. Ecological models of health behavior. In *Health behavior and health education: Theory, research, and practice, 4th ed.* Jossey-Bass, San Francisco, CA, US, 465–485.
- [71] Martin EP Seligman. 1975. On depression, development, and death. *San Francisco: Freeman* (1975).
- [72] Mehwish Shahid, Nicole H. Weiss, Gary Stoner, and Bryan Dewsbury. 2021. Asian Americans' mental health help-seeking attitudes: The relative and unique roles of cultural values and ethnic identity. *Asian American Journal of Psychology* 12 (2021), 138–146. <https://doi.org/10.1037/aap0000230> Place: US Publisher: Educational Publishing Foundation.
- [73] Ji Youn Shin, Minjin Rheu, Jina Huh-Yoo, and Wei Peng. 2021. Designing Technologies to Support Parent-Child Relationships: A Review of Current Findings and Suggestions for Future Directions. *Proceedings of the ACM on Human-Computer Interaction* 5, CSCW2 (Oct. 2021), 441:1–441:31. <https://doi.org/10.1145/3479585>
- [74] Farheen Siddiqui, Delvin Varghese, Pushpendra Singh, Sunita Bapuji Bayyavarupu, Stephen Lindsay, Dharshani Chandrasekara, Pranav Kulkarni, Ling Wu, Taghreed Alshehri, and Patrick Olivier. 2023. Exploring the digital support needs of caregivers of people with serious mental illness. In *Proceedings of the 2023 CHI Conference on Human Factors in Computing Systems (CHI '23)*. Association for Computing Machinery, New York, NY, USA, 1–16.

<https://doi.org/10.1145/3544548.3580674>

- [75] Ewan Soubutts, Pranita Shrestha, Brittany I Davidson, Chengcheng Qu, Charlotte Mindel, Aaron Sefi, Paul Marshall, and Roisin Mcnaney. 2024. Challenges and Opportunities for the Design of Inclusive Digital Mental Health Tools: Understanding Culturally Diverse Young People's Experiences. In *Proceedings of the CHI Conference on Human Factors in Computing Systems (CHI '24)*. Association for Computing Machinery, New York, NY, USA, 1–16. <https://doi.org/10.1145/3613904.3642641>
- [76] Evropi Stefanidi, Johannes Schöning, Yvonne Rogers, and Jasmin Niess. 2023. Children with ADHD and their Care Ecosystem: Designing Beyond Symptoms. In *Proceedings of the 2023 CHI Conference on Human Factors in Computing Systems (CHI '23)*. Association for Computing Machinery, New York, NY, USA, 1–17. <https://doi.org/10.1145/3544548.3581216>
- [77] Daniel Stokols. 1996. Translating Social Ecological Theory into Guidelines for Community Health Promotion. *American Journal of Health Promotion* 10, 4 (March 1996), 282–298. <https://doi.org/10.4278/0890-1171-10.4.282> Publisher: SAGE Publications Inc.
- [78] Franziska Tachtler, Reem Talhouk, Toni Michel, Petr Slovak, and Geraldine Fitzpatrick. 2021. Unaccompanied Migrant Youth and Mental Health Technologies: A Social-Ecological Approach to Understanding and Designing. In *Proceedings of the 2021 CHI Conference on Human Factors in Computing Systems (CHI '21)*. Association for Computing Machinery, New York, NY, USA, 1–19. <https://doi.org/10.1145/3411764.3445470>
- [79] Nikki Theofanopoulou and Petr Slovak. 2022. Exploring Technology-Mediated Parental Socialisation of Emotion: Leveraging an Embodied, In-situ Intervention for Child Emotion Regulation. In *Proceedings of the 2022 CHI Conference on Human Factors in Computing Systems (CHI '22)*. Association for Computing Machinery, New York, NY, USA, 1–16. <https://doi.org/10.1145/3491102.3502130>
- [80] Beti Thompson and Susan Kinne. 1990. Social change theory: Applications to community health. *Health promotion at the community level* 2 (1990), 29–46.
- [81] Tammy Toscos, Maria Carpenter, Michelle Drouin, Amelia Roebuck, Connie Kerrigan, and Michael Mirro. 2018. College Students' Experiences with, and Willingness to Use, Different Types of Telemental Health Resources: Do Gender, Depression/Anxiety, or Stress Levels Matter? *Telemedicine Journal and E-Health: The Official Journal of the American Telemedicine Association* 24, 12 (Dec. 2018), 998–1005. <https://doi.org/10.1089/tmj.2017.0243>
- [82] Ciaran B. Trace, Yan Zhang, Siqi Yi, and Marian Yvette Williams-Brown. 2023. Information practices around genetic testing for ovarian cancer patients. *Journal of the Association for Information Science and Technology* 74, 11 (Nov. 2023), 1265–1281. <https://doi.org/10.1002/asi.24823>
- [83] Yunlong Wang, Shuyuan Shen, and Brian Y Lim. 2023. RePrompt: Automatic Prompt Editing to Refine AI-Generative Art Towards Precise Expressions. In *Proceedings of the 2023 CHI Conference on Human Factors in Computing Systems*. ACM, Hamburg Germany, 1–29. <https://doi.org/10.1145/3544548.3581402>
- [84] David Wiljer, Jenny Shi, Brian Lo, Marcos Sanches, Elisa Hollenberg, Andrew Johnson, Alexxa Abi-Jaoudé, Gloria Chaim, Kristin Cleverley, Joanna Henderson, Wanrudee Isaranuwatthai, Andrea Levinson, Janine Robb, Howard W. Wong, and Aristotle Voineskos. 2020. Effects of a Mobile and Web App (Thought Spot) on Mental Health Help-Seeking Among College and University Students: Randomized Controlled Trial. *Journal of Medical Internet Research* 22, 10 (Oct. 2020), e20790. <https://doi.org/10.2196/20790>
- [85] Lucretia Williams, Gillian Hayes, Gloria Washington, Rebecca W. Black, Candace Williams, Leah Clements, and Michael Allotey. 2021. Analysis Of Distance-Based Mental Health Support For Underrepresented University Students. In *Extended Abstracts of the 2021 CHI Conference on Human Factors in Computing Systems*. ACM, Yokohama Japan, 1–6. <https://doi.org/10.1145/3411763.3451708>
- [86] Howard W. Wong, Brian Lo, Jenny Shi, Elisa Hollenberg, Alexxa Abi-Jaoude, Andrew Johnson, Gloria Chaim, Kristin Cleverley, Joanna Henderson, Andrea Levinson, Janine Robb, Aristotle Voineskos, and David Wiljer. 2021. Postsecondary Student Engagement With a Mental Health App and Online Platform (Thought Spot): Qualitative Study Assessing Factors Related to User Experience. *JMIR Mental Health* 8, 4 (April 2021), e23447. <https://doi.org/10.2196/23447>
- [87] Novia Wong, Victoria Jackson, André Van Der Hoek, Iftekhar Ahmed, Stephen M. Schueller, and Madhu Reddy. 2023. Mental Wellbeing at Work: Perspectives of Software Engineers. In *Proceedings of the 2023 CHI Conference on Human Factors in Computing Systems (CHI '23)*. Association for Computing Machinery, New York, NY, USA, 1–15. <https://doi.org/10.1145/3544548.3581528>
- [88] Marisol Wong-Villacres and Shaowen Bardzell. 2011. Technology-mediated parent-child intimacy: designing for ecuadorian families separated by migration. In *CHI '11 Extended Abstracts on Human Factors in Computing Systems (CHI EA '11)*. Association for Computing Machinery, New York, NY, USA, 2215–2220. <https://doi.org/10.1145/1979742.1979877>
- [89] Susan P. Wyche, Elisa Oreglia, Morgan G. Ames, Christopher Hoadley, Aditya Johri, Phoebe Sengers, and Charles Steinfield. 2012. Learning from marginalized users: reciprocity in HCI4D. In *Proceedings of the ACM 2012 conference on Computer Supported Cooperative Work Companion (CSCW '12)*. Association for Computing Machinery, New York, NY, USA, 1–16. <https://doi.org/10.1145/2374776.2374800>

- USA, 27–28. <https://doi.org/10.1145/2141512.2141527>
- [90] Renwen Zhang, Jordan Eschler, and Madhu Reddy. 2018. Online Support Groups for Depression in China: Culturally Shaped Interactions and Motivations. *Computer Supported Cooperative Work (CSCW)* 27, 3 (Dec. 2018), 327–354. <https://doi.org/10.1007/s10606-018-9322-4>