

QUERY FORM

JOURNAL: The Astrophysical Journal Supplement Series

AUTHOR: Yang et al.

TITLE: Correction of Star Catalog Positional and Proper-motion Systematic Errors for Natural Satellite Astrometry

ARTICLE ID: apjsae8d01

Page 1

Q1
Your article has been processed in line with the journal style. Your changes will be reviewed by the Production Editor, and any amendments that do not comply with journal style or grammatical correctness will not be applied and will not appear in the published article.

Page 1

Q2
IMPORTANT: This PDF reference document has been locked to prevent comments. Please use our online Proofing tool to answer all queries, make corrections and add comments.

Page 1

Q3
The layout of this article has not yet been finalized. This proof may contain columns that are not fully balanced/matched or other typographical imperfections; these issues will be resolved once the final corrections have been incorporated.

Page 1

Q4
Please check that the **names of all authors as displayed in the proof are correct**, and that all **authors are linked to the correct affiliations**. Please also confirm that the correct corresponding author has been indicated. **Note that this is your last opportunity to review and amend this information before your article is published.**

Page 1

Q5
We have been provided with ORCID iDs for the authors as below. Please confirm whether the numbers are correct.
H. Y. Zhang 0000-0003-0956-2440
K. Tang 0000-0001-7747-5707

Page 1

Q6
Please check that the funding information below is correct for inclusion in the article metadata.
MOST | National Natural Science Foundation of China (NSFC): 12473070.

Page 2

Q7
Please define the acronyms [PPM, ACT, AC] if definitions exist.

Page 4

Q8

Please confirm that the edits made to the sentence [The quantities in (2)] retain the intended sense.

Page 10

Q9

If you have an Acknowledgments section, please check that the information there is complete and correct and that all relevant institutions, grant numbers, programs, and collaborators are mentioned as appropriate.

Page 11

Q10

Please check/updated the link in footnote 6 as it directs to the NADC homepage.



Correction of Star Catalog Positional and Proper-motion Systematic Errors for Natural Satellite Astrometry

Z. Yang¹, H. Y. Zhang¹, K. Tang², Y. Yu^{2,3}, S. L. Liao^{2,3}, D. Yan⁴, and R. C. Qiao⁵
¹School of Mathematics, Physics and Statistics, Shanghai University of Engineering Science, Shanghai 201620, People's Republic of China; zhanghy1009@163.com
²Shanghai Astronomical Observatory, Chinese Academy of Sciences, Shanghai 200030, People's Republic of China
³School of Astronomy and Space Science, University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China
⁴School of Physics and Electronic Information, Huanggang Normal University, Huanggang 438000, People's Republic of China
⁵National Time Service Center, Chinese Academy of Sciences, Lintong, Shaanxi 710600, People's Republic of China
 Received 2026 June 14; revised 2026 July 11; accepted 2026 July 14; published 2026 MM DD

Abstract

Precise orbit determination of natural satellites relies on the consistent combination of heterogeneous astrometric observations spanning long time intervals. However, the use of different reference star catalogs introduces spatially correlated systematic biases that degrade data consistency and limit orbit accuracy. In this work, we construct a unified correction framework based on the Gaia DR3 reference frame and derive position and proper-motion corrections for 17 star catalogs commonly used in natural satellite astrometry. The method employs rigorous epoch propagation, HEALPix-based spatial partitioning, and robust median statistics to estimate local systematics, and reconstructs a continuous correction field using radial basis function interpolation. The derived corrections reveal a clear hierarchy among the catalogs: modern catalogs exhibit systematics at the milliarcsecond level, whereas older photographic catalogs show position offsets reaching the 100 mas to arcsecond level. This framework provides a consistent basis for reprocessing historical observations and improves the astrometric accuracy of natural satellite observations, thereby enhancing the precision of orbit determination and dynamical studies of natural satellite systems.

Unified Astronomy Thesaurus concepts: Natural satellites (Solar system) (1089); Gaia (2360); Astrographic catalogs (77); Astrometry (80)

1. Introduction

Precise orbit determination of natural satellites is fundamental to studies of planetary system dynamics and long-term ephemeris prediction. Optical astrometric observations provide the primary positional constraints for improving satellite orbits. However, such observations typically span long time intervals and originate from multiple observatories, instruments, and reduction schemes. The consistency of these heterogeneous datasets directly impacts the accuracy and stability of the resulting orbital solutions and their uncertainty characterization (J. Desmars et al. 2013).

A large number of astrometric positions of natural satellites have been compiled in the Natural Satellite Database (NSDB), maintained by the Laboratoire Temps Espace at Paris Observatory (J.-E. Arlot et al. 2024). Owing to the diversity of data sources and the long temporal coverage, inconsistencies related to the use of different reference star catalogs have become increasingly evident when these observations are combined for orbit determination. Differences in reference frames, regional systematic biases, and proper-motion accuracy among catalogs introduce systematic biases in the reduced positions, which can accumulate over time and degrade the overall consistency of orbital solutions.

Since 1986, we have conducted continuous observations of natural satellites in the Jovian, Saturnian, Uranian, and Neptunian systems, yielding more than 50,000 astrometric

positions, a portion of which has been incorporated into the NSDB. Based on these data, a series of studies has been carried out on the astrometric reduction and orbit determination of Saturnian satellites, the major Uranian satellites, and the irregular satellite Phoebe, as well as on the digitization and reduction of historical natural-satellite plates (R. Qiao et al. 1994; K. X. Shen et al. 2002; R. C. Qiao et al. 2006; D. Yan et al. 2016). These studies demonstrate that reprocessing historical observations within a uniform high-precision reference frame is an effective approach to improving data consistency.

Systematic biases in star catalogs have been extensively investigated (e.g., R. Teixeira et al. 2014). Early analyses of asteroid astrometry showed that optical observations contain nonnegligible systematic biases (M. Carpino et al. 2003). S. R. Chesley et al. (2010) identified regional systematic biases through residual analysis and introduced a spatially resolved debiasing scheme using the Two Micron All Sky Survey (2MASS) as the reference catalog (M. F. Skrutskie et al. 2006). D. J. Tholen et al. (2013) further found that the absence of proper motions in 2MASS produces regional position offsets in high-precision astrometry. This motivated the proper-motion-corrected scheme of D. Farnocchia et al. (2015), which uses PPMXL as the reference catalog (S. Roeser et al. 2010). More recently, S. Eggl et al. (2020) utilized the Gaia DR2 reference frame to derive position and proper-motion corrections for 26 astrometric catalogs used in asteroid astrometry. However, the catalog selection in that work is tailored to asteroid observations and does not cover several catalogs prevalent in historical natural-satellite astrometry, particularly those spanning earlier observational epochs. As



Original content from this work may be used under the terms of the Creative Commons Attribution 4.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

the raw observational data are in many cases no longer available for reduction, applying catalog corrections to the existing reduced positions is the only viable means of recovering their astrometric accuracy. Taken together, these studies establish that catalog debiasing within a common reference frame is essential for improving the consistency of astrometric data.

In the context of natural satellites, similar approaches have only recently begun to be explored. For example, Y. Yuan et al. (2021a) performed a uniform reanalysis of Neptunian satellite observations based on the Gaia DR2 reference frame and demonstrated the applicability of [catalog-debiasing](#) methods to satellite astrometry. In addition, our group has conducted Gaia-based reduction of selected historical observations, confirming the effectiveness of this approach in improving astrometric consistency. Nevertheless, a systematic debiasing framework covering the full range of reference catalogs commonly used in natural-satellite astrometry is still lacking.

In this work, we adopt Gaia DR3 as the common reference frame and derive position and proper-motion corrections for 17 historical star catalogs. These 17 catalogs are identified, based on a systematic survey of the NSDB archive, as the most prevalent reference catalogs in historical natural-satellite astrometry. Building upon the [catalog-debiasing](#) framework of S. Eggl et al. (2020), we introduce adaptations tailored to [natural-satellite](#) astrometry, including refined [crossmatching](#) criteria and additional validation steps. The resulting correction framework provides a consistent basis for reprocessing historical observations and supports improved astrometric accuracy of [natural-satellite](#) observations, thereby enhancing the precision of orbit determination and dynamical studies of [natural-satellite](#) systems.

2. Reference and Astrometric Catalogs

2.1. Gaia Catalog Series

Since 2013, the Gaia mission has provided a dense all-sky optical astrometric reference frame. Gaia DR1 contained astrometric measurements for more than one billion sources, but a full five-parameter solution was available only for the Tycho–Gaia Astrometric Solution subset, while most sources lacked [proper-motion](#) information (Gaia Collaboration et al. 2016; L. Lindegren et al. 2016). Gaia DR2 subsequently provided homogeneous five-parameter astrometric solutions for a much larger number of sources and became an important reference for studies of systematic errors in historical star catalogs and [solar system](#) astrometry (Gaia Collaboration et al. 2018).

In this work, Gaia DR3 is adopted as the reference catalog for deriving position and [proper-motion](#) corrections. The Gaia DR3 astrometric solution is based on Gaia EDR3 and benefits from a longer observational time span and improved [modeling](#) of the spacecraft attitude, chromaticity, and instrumental calibration (L. Lindegren et al. 2021b; Gaia Collaboration et al. 2023). Compared with earlier Gaia releases, Gaia DR3 provides improved astrometric precision and reference frame stability, making it more suitable for quantifying local systematic differences between historical reference catalogs and a common modern astrometric frame (C. Fabricius et al. 2021).

Residual systematic effects still exist in Gaia [EDR3/DR3](#), including parallax biases, proper-motion biases, and small-scale spatial systematics (T. Cantat-Gaudin & T. D. Brandt 2021; L. Lindegren et al. 2021a). However, for the historical catalogs considered in this work, the dominant correction signals arise from catalog-dependent position and [proper-motion](#) biases rather than from the residual systematics of Gaia DR3 itself. Gaia DR3 is therefore used as the practical reference frame for constructing the catalog corrections.

2.2. Historical Astrometric Catalogs

This work targets catalogs that are frequently employed in historical [natural-satellite](#) observations, including older releases that were prevalent before high-precision space-based astrometry became available.

Optical astrometric observations from different epochs rely on a variety of reference star catalogs that differ significantly in epoch, sky coverage, source density, and available astrometric parameters. The catalogs included in this study are those most prevalently used in historical [natural-satellite](#) observations, as identified from a systematic survey of the NSDB archive.

Early photographic catalogs, such as USNO-A2.0 and GSC 1.2, generally lack [proper-motion](#) information and can introduce time-dependent position offsets (D. G. Monet 1998; J. E. Morrison et al. 2001). Intermediate catalogs, including PPM, ACT, and the UCAC series, provide improved positional accuracy and [proper-motion](#) coverage but may still exhibit regional systematic biases (S. Roeser & U. Bastian 1991; S. E. Urban et al. 1998; N. Zacharias et al. 2004, 2013). High-precision catalogs such as Hipparcos and Tycho-2 provide a stable and well-characterized reference frame but are limited in source density, which can restrict the number of usable reference stars in fields near bright planets (European Space Agency 1997; E. Høg et al. 2000). The final sample consists of 17 star catalogs, including Gaia DR1 and Gaia DR2, whose key properties are summarized in Table 1.

3. Basic Framework of the Correction Method

3.1. Overview of the Method

The overall procedure consists of five main steps:

- (1) transform all catalogs to a common reference frame and comparison epoch;
- (2) partition the celestial sphere using a HEALPix tessellation;
- (3) perform [crossmatching](#) between the target catalog and Gaia DR3 within each tile;
- (4) estimate local systematic offsets in position and proper motion;
- (5) construct a debiasing model and generate tabulated corrections.

3.2. Frame and Epoch Transformation

To ensure consistency, all catalogs are transformed to a common reference frame and comparison epoch. J2000.0 is adopted as the comparison epoch since it is the reference epoch for most historical catalogs considered in this work.

For catalogs already in the [International Celestial Reference System \(ICRS\)](#), positions are propagated to epoch J2000.0 when [proper-motion](#) information is available. For catalogs

Table 1
Summary of Astrometric Catalogs Used in This Work

Catalog	Magnitude	Number of Stars	Epoch	Reference Frame	Proper Motion	Coverage	CDS ID	References
AGK1	$V \approx 2.2\text{--}9.1$	5954	mean epoch	FK4	NO	3%	I/310 ^a	Astronomische Gesellschaft (1890)
FK4 catalog	$V \approx 1\text{--}9.5$	1535	B1950	FK4	YES	3%	I/143 ^b	W. Fricke et al. (1963)
SAO	$V = 0\text{--}11$	258,997	J2000	ICRS	YES	98%	I/131A ^c	Smithsonian Astrophysical Observatory Staff (1966)
AGK3	$B, V \approx 6\text{--}11$	183,145	observation	FK4	YES	52%	I/61B ^d	O. Heckmann & W. Dieckvoss (1975)
ACRS	$V \approx 6\text{--}11$	320,211	J2000	ICRS	YES	100%	I/171 ^e	T. E. Corbin et al. (1991)
AC	$V \approx 6\text{--}11.0$	2,026,347	plate epoch	FK4	NO	30%	I/154 ^f	S. Roeser (1990)
PPM	$V \approx 1\text{--}11.5$	378,910	J2000	ICRS	YES	100%	I/146, ^g I/193 ^h	S. Roeser & U. Bastian (1991)
Yale	$V \approx -1\text{--}6.5$	221,338	observation	FK4	YES	76%	I/141 ⁱ	D. Hoffleit & W. H. Warren (1991)
Hipparcos	$V \approx 1\text{--}12.4$	118,218	J1991.25	ICRS	YES	94%	I/239 ^j	European Space Agency (1997)
ACT	$V \approx 6\text{--}11.5$	988,758	J2000	ICRS	YES	100%	I/246 ^k	S. E. Urban et al. (1998)
USNO-A2.0	$B, R = 11\text{--}20$	526,280,881	mean epoch	ICRS	NO	100%	I/252 ^l	D. G. Monet (1998)
Tycho-2	$V \approx 2\text{--}12.5$	2,539,913	J2000	ICRS	YES	100%	I/259 ^m	E. Høg et al. (2000)
GSC 1.2	$V \approx 6\text{--}15$	18,825,216	plate epoch	ICRS	NO	100%	I/254 ⁿ	J. E. Morrison et al. (2001)
UCAC2	$R \approx 8\text{--}16$	48,330,571	J2000	ICRS	YES	88%	I/289 ^o	N. Zacharias et al. (2004)
UCAC4	$R \approx 7\text{--}16$	113,780,093	J2000	ICRS	YES	100%	I/322A ^p	N. Zacharias et al. (2013)
Gaia DR1	$G = 5\text{--}20.7$	1,142,685,696	J2015	ICRS	NO	100%	I/337 ^q	Gaia Collaboration et al. (2016)
Gaia DR2	$G = 3\text{--}21$	1,331,909,727	J2015.5	ICRS	YES	100%	I/345 ^r	Gaia Collaboration et al. (2018)

Note. Catalogs are listed in approximate chronological order of publication. Gaia DR1 and DR2 are included for completeness; their astrometric properties are described in detail in Section 2.1. The “Proper Motion” column indicates whether the catalog provides proper-motion data. “Coverage” gives the approximate fraction of the sky covered by the catalog. Primary literature references correspond to the original catalog publications listed in the CDS entries.

- ^a <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/310>
^b <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/143>
^c <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/131A>
^d <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/61B>
^e <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/171>
^f <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/154>
^g <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/146>
^h <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/193>
ⁱ <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/141>
^j <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/239>
^k <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/246>
^l <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/252>
^m <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/259>
ⁿ <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/254>
^o <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/289>
^p <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/322A>
^q <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/337>
^r <https://cdsarc.cds.unistra.fr/viz-bin/cat/I/345>

lacking proper motions, Gaia positions are propagated to the catalog epoch, and the comparison is performed at that epoch.

For catalogs based on the FK4 reference frame (AC, AGK1, AGK3, FK4 catalog, and Yale), coordinates are first transformed into the ICRS, and matching is carried out at the corresponding comparison epochs (C. A. Murray 1989; C. Ma et al. 1998; A. L. Fey et al. 2015). This procedure ensures that the measured differences primarily reflect intrinsic catalog systematics and avoids introducing additional uncertainties from inaccurate proper motions in historical catalogs.

Gaia DR3 adopts a reference epoch of J2016.0. Source positions are propagated to the desired comparison epoch using the published astrometric parameters (positions, proper motions, and parallaxes). Radial velocities are included when available and assumed to be zero otherwise, which introduces negligible errors at the precision level considered here.

3.3. Sky Partitioning and Crossmatching

After the reference frame and epoch are unified, the celestial sphere is divided using a HEALPix (K. M. Górski et al. 2005) tessellation with $N_{\text{side}} = 64$, corresponding to $N_{\text{pix}} = 49,152$ equal-area tiles, each covering approximately 0.839 deg^2 .

Crossmatching is performed within each tile, and all angular distances are computed on the celestial sphere. For each source in the target catalog, Gaia DR3 candidates are searched within a search radius of $1''$ (D. Farnocchia et al. 2015). The matching criteria are defined as follows:

- (1) If a single candidate is found, the match is accepted.
- (2) If multiple candidates are present, the nearest neighbor is accepted only if $d_1 < 0.2 d_2$, where d_1 and d_2 are the angular distances to the nearest and second-nearest neighbors, respectively.
- (3) If no candidate is found, the source is rejected.

All matches are performed using coordinates expressed in the same reference frame and epoch.

3.4. Bias Estimation

For each matched pair, the differences in position and proper motion are defined as

$$\begin{aligned}\Delta\alpha &= (\alpha_{\text{catalog}} - \alpha_{\text{Gaia}})\cos\delta_{\text{Gaia}}, & \Delta\delta &= \delta_{\text{catalog}} - \delta_{\text{Gaia}}, \\ \Delta\mu_\alpha &= \mu_{\alpha,\text{catalog}} - \mu_{\alpha,\text{Gaia}}, & \Delta\mu_\delta &= \mu_{\delta,\text{catalog}} - \mu_{\delta,\text{Gaia}}.\end{aligned}\quad (1)$$

Here δ_{Gaia} denotes the Gaia DR3 [decl.](#) of the matched source at the comparison epoch. The $\cos\delta_{\text{Gaia}}$ factor converts the coordinate difference to a great-circle arc length.

For catalogs without [proper-motion](#) information, only position differences are used. In this case, Gaia source positions are propagated to the catalog epoch t_{catalog} , and the differences $\Delta\alpha(t_{\text{catalog}})$ and $\Delta\delta(t_{\text{catalog}})$ are computed at that epoch. The tile-level medians are then referred to J2000.0 using the median Gaia proper motion of the tile.

Within each HEALPix tile, the median of these differences is computed and referenced to epoch J2000.0, yielding the systematic correction parameters. Tiles with fewer than [10](#) matched sources are excluded to ensure statistical reliability.

$$(\Delta\alpha_{\text{J2000}}, \Delta\delta_{\text{J2000}}, \Delta\mu_\alpha, \Delta\mu_\delta). \quad (2)$$

[Q8](#) [The](#) quantities in (2) represent local catalog systematics referenced to epoch J2000.0. The use of median statistics provides robustness against outliers without the need for additional rejection criteria.

3.5. Correction Tables and Application

The resulting corrections are stored on a HEALPix grid. Each tile is associated with the four correction parameters $\Delta\alpha_{\text{J2000}}$, $\Delta\delta_{\text{J2000}}$, $\Delta\mu_\alpha$, and $\Delta\mu_\delta$, indexed by its HEALPix pixel number.

For an arbitrary epoch t , the corrections are applied via a linear model:

$$\begin{aligned}\Delta\alpha(t) &= \Delta\alpha_{\text{J2000}} + \Delta\mu_\alpha \Delta t, \\ \Delta\delta(t) &= \Delta\delta_{\text{J2000}} + \Delta\mu_\delta \Delta t, \\ \Delta t &= t - 2000.0.\end{aligned}\quad (3)$$

In practical applications, the corresponding tile is identified from the sky position of an observation, and the associated parameters are used to compute $\Delta\alpha(t)$ and $\Delta\delta(t)$. The corrected positions are obtained by applying these offsets, thereby aligning the observations with the Gaia DR3 reference frame. For the data format and usage of the released correction tables, see [the Data and Code Availability section](#).

4. Results

4.1. Statistical Properties of the Corrections

The global statistics in Table 2 reveal a clear hierarchy in the amplitudes of the derived catalog corrections. Modern high-precision catalogs, such as Gaia DR2, Hipparcos, Tycho-2, UCAC2, and UCAC4, exhibit median position and [proper-motion](#) corrections at the milliarcsecond level, with small dispersions across the sky. In contrast, older photographic catalogs and catalogs without reliable proper motions show significantly larger offsets and broader sky-to-sky scatter, with

typical position systematics ranging from tens to hundreds of [milliarcseconds](#), and reaching the arcsecond level in the most extreme cases (e.g., AGK1).

USNO-A2.0 provides a representative example. The median corrections at [J2000.0](#) are $\Delta\alpha_{\text{J2000}} = 68.0$ mas and $\Delta\delta_{\text{J2000}} = 151.9$ mas, with MAD values of 104.2 and 121.5 mas, respectively. The corresponding [proper-motion](#) corrections are also [nonnegligible](#), reaching 1.22 mas yr^{-1} in $\Delta\mu_\alpha$ and 3.03 mas yr^{-1} in $\Delta\mu_\delta$.

Figure 1 shows the distributions of the correction terms for USNO-A2.0 across all HEALPix tiles. The position components exhibit pronounced global offsets and non-Gaussian distributions, while the [proper-motion](#) corrections display asymmetric behavior between α and δ . These characteristics indicate that the corrections reflect spatially coherent systematic differences with respect to the Gaia DR3 reference frame rather than random noise.

4.2. Spatial Structure of the Systematic Biases

The correction fields exhibit clear spatial organization across the celestial sphere. As shown in Figure 2, the USNO-A2.0 corrections display large-scale gradients together with [blocklike](#) discontinuities. Similar spatial patterns are observed in other historical catalogs, although their morphology and characteristic scales differ.

These structures demonstrate that the correction fields are dominated by spatially correlated systematics. In particular, the presence of coherent large-scale patterns indicates that the derived corrections trace intrinsic catalog biases rather than stochastic matching errors. Consequently, global summary statistics alone are insufficient to fully characterize the spatial behavior of catalog systematics.

4.3. Interpretation of the Spatial Patterns

The morphology of the correction fields reflects the observational and reduction characteristics of the underlying catalogs. Three representative classes can be identified.

Photographic catalogs (e.g., [GSC 1.2](#), [AGK3](#), [SAO](#), and [Yale](#)) show [blocklike](#) or discontinuous structures associated with independent plate-based reductions ([H. K. Eichhorn 1974](#); [I. Platais et al. 1998](#)). This behavior can be quantified through the [intertile discontinuity](#),

$$D_{\text{disk}}(i, j) = |V(i) - V(j)|, \quad (4)$$

where i and j denote adjacent HEALPix tiles (including both edge- and vertex-sharing neighbors) and V represents a correction component. For these catalogs, D_{disk} values frequently exceed 100 mas and can reach the arcsecond level, indicating globally inconsistent plate-to-plate calibrations.

Catalogs based on scanning strategies (e.g., UCAC2; Figure 3) exhibit stripe-like patterns aligned with the scan direction (N. Zacharias et al. 2017), with amplitudes enhanced in crowded regions. This behavior is largely mitigated in UCAC4, consistent with improved reduction procedures.

Catalogs lacking reliable [proper-motion](#) information (e.g., USNO-A2.0) display smooth large-scale gradients in the propagated position corrections. These patterns arise from the accumulation of [proper-motion](#) mismatches over long temporal baselines (E. Vasiliev 2019).

Table 2
Global Statistics of Position and Proper-motion Corrections for Each Star Catalog at Epoch J2000.0

Catalog	$\Delta\alpha_{J2000}$		$\Delta\delta_{J2000}$		$\Delta\mu_{\alpha^*}$		$\Delta\mu_{\delta^*}$	
	(mas)		(mas)		(mas yr ⁻¹)		(mas yr ⁻¹)	
	MED	MAD	MED	MAD	MED	MAD	MED	MAD
AGK1	382.8	785.1	237.6	582.3	0.63	7.76	6.48	7.26
FK4 <i>catalog</i>	-73.2	157.3	-45.3	201.3	-0.88	2.33	-0.26	3.02
SAO	-5.7	270.0	-19.4	266.1	0.61	3.76	-0.11	3.66
AGK3	-127.9	257.3	-45.5	301.5	-1.82	4.67	-0.59	5.38
ACRS	-39.0	136.6	-47.9	127.8	-0.91	2.87	-0.09	2.56
AC	95.1	322.6	498.2	234.1	0.65	2.76	4.97	1.64
PPM	12.7	128.4	-67.1	127.2	-0.23	2.72	-0.46	2.78
Yale	18.9	409.8	-139.4	439.1	0.57	5.70	-0.64	5.73
Hipparcos	0.7	6.7	0.2	5.7	0.08	0.72	0.01	0.62
ACT	-0.3	11.6	0.3	10.8	-0.09	1.03	-0.01	0.97
USNO-A2.0	68.0	104.2	151.9	121.5	1.22	1.71	3.03	1.28
Tycho-2	-0.3	10.0	-1.0	9.7	-0.07	0.81	-0.26	0.81
GSC 1.2	28.3	113.0	-13.4	104.4	0.98	2.81	3.93	1.51
UCAC2	-0.8	10.3	2.9	9.2	-0.30	1.31	-0.58	1.26
UCAC4	1.0	11.0	-3.0	9.6	0.10	0.83	0.36	0.94
Gaia DR1	-23.3	34.1	-55.6	23.0	1.55	2.28	3.70	1.53
Gaia DR2	-0.1	0.3	-0.3	0.4	0.00	0.02	0.01	0.02

Note. MED denotes the median and MAD the median absolute deviation over all valid tiles, defined as $\text{median}(|x_i - \text{median}(x)|)$. Position corrections $\Delta\alpha_{J2000}$ and $\Delta\delta_{J2000}$ are in *milliarcseconds*; *proper-motion* corrections $\Delta\mu_{\alpha^*}$ and $\Delta\mu_{\delta^*}$ are in mas yr^{-1} .

These results show that both the amplitude and morphology of the correction fields are controlled by the intrinsic properties of the target catalogs.

4.4. Smoothing and Practical Implementation

For practical applications, a higher-resolution correction map better represents the spatial structure of catalog systematics. However, directly estimating corrections at finer resolution (e.g., $N_{\text{side}} = 256$) would significantly reduce the number of matched stars per tile, degrading the statistical robustness of the median estimates.

To balance spatial resolution and statistical stability, we adopt a similar strategy to that of S. Eggl et al. (2020): a continuous correction field is reconstructed from the statistically robust $N_{\text{side}} = 64$ tile estimates using radial basis function (RBF) interpolation with a thin-plate spline kernel (G. Wahba 1990; R. Schaback 1995). The interpolation is performed on the unit-sphere coordinates of the HEALPix pixel centers using a local RBF interpolator with eight nearby valid $N_{\text{side}} = 64$ tiles and a thin-plate spline kernel. The final correction product is provided on a HEALPix grid with $N_{\text{side}} = 256$. Since these values are reconstructed from the coarse-grid estimates rather than independently derived from observations at that resolution, they represent an interpolated continuous field rather than independent measurements.

The effect on spatial continuity is illustrated in Figure 4. The reconstructed field eliminates the *blocklike* structures present in the original tile-based map while preserving the underlying large-scale patterns. To quantify this improvement, we evaluate the *intertile* discontinuity D_{disk} defined in Section 4.3. As shown in Figure 5, the *cumulative distribution functions* shift toward smaller D_{disk} after smoothing, and the high-percentile and maximum values of D_{disk} are reduced by factors of roughly 2–4 after smoothing.

At the same time, the large-scale spatial structure of the corrections is preserved, indicating that the smoothing

primarily suppresses sampling noise at small scales rather than altering the underlying systematic patterns.

5. Verification

5.1. Plate Reduction Verification

To independently evaluate the effectiveness of the *catalog-debiasing* model, we compare two reductions of the same Triton CCD frames obtained during 1996–2006. The earlier reduction, based on UCAC2, yielded 943 positions (R. C. Qiao et al. 2007), whereas a later re-reduction of the same frames using Gaia DR2 produced 1007 positions (D. Yan et al. 2020). Strict epoch-based pairing identifies 930 common positions, for which the Gaia DR2 reduction is used as the comparison benchmark.

The position offset Δ is defined as

$$\Delta = \sqrt{(\Delta\alpha)^2 + (\Delta\delta)^2}, \quad (5)$$

where $\Delta\alpha = (\alpha_{\text{UCAC2}} - \alpha_{\text{Gaia DR2}})\cos\delta$ and $\Delta\delta = \delta_{\text{UCAC2}} - \delta_{\text{Gaia DR2}}$ are the position differences between the UCAC2-based reduction and the Gaia DR2 reduction.

Relative to the Gaia DR2 reduction, the UCAC2-based reduction has a mean position offset of 86.2 mas and a median of 82.5 mas. After application of the Gaia *DR3-based* debiasing model, the mean and median offsets decrease to 76.2 and 67.7 mas, respectively. This corresponds to an 11.6% reduction in the mean offset, with 78.0% of the common positions showing smaller Δ values after correction. The remaining difference should not be interpreted solely as uncorrected catalog bias, because the two reductions also differ in centroiding and reduction procedures (D. Yan et al. 2020).

Figure 6(a) shows that the *postdebiasing* cumulative distribution is shifted toward smaller position offsets, indicating that the corrected positions are generally closer to the Gaia DR2 reduction. Figure 6(b) shows that the running mean of Δ




Figure 1. Statistical distributions of the correction terms for the USNO-A2.0 star catalog.

remains lower after debiasing over most of the observing interval, indicating that the improvement is sustained rather than confined to a narrow epoch range.

The debiasing model is built from Gaia DR3, whereas the benchmark reduction uses Gaia DR2 as the reference catalog. Since both releases share a consistent ICRS realization and their differences are far smaller than UCAC2 systematics, the reduction in position offset constitutes a conservative test of the debiasing model.

5.2. Ephemeris-comparison Validation

As a complementary external test, this section compares the *observed* minus *calculated* ($O - C$) residuals before and after applying the *catalog-debiasing* corrections, using the JPL Horizons DE441 planetary ephemeris (R. S. Park et al. 2021) as an independent dynamical reference.

The validation sample consists of 11,731 absolute astrometric observations of the four Galilean satellites from the NSDB archive, spanning 1962–2019 and covering eight reference catalogs across 19 observation files. Because some files contain segments reduced with multiple catalogs,

observations are grouped by (file, catalog) pairs, giving 21 groups.

For each observation, the $O - C$ residuals in the two coordinate directions are computed from both the original and the debiased positions:

$$\begin{aligned}\Delta\alpha &= (\alpha_{\text{obs}} - \alpha_{\text{eph}})\cos\delta \\ \Delta\delta &= \delta_{\text{obs}} - \delta_{\text{eph}}.\end{aligned}\tag{6}$$

Typical single-observation residuals in this dataset are 100–200 mas, while median catalog corrections are only 10–30 mas. At the individual-observation level, the correction signal is therefore dominated by random measurement noise. Observations within a single file, however, share the same reference catalog and reduction pipeline; averaging over the N observations of a file decreases the random component by $1/\sqrt{N}$. We therefore adopt file-level mean residuals $\overline{\Delta\alpha}$ and $\overline{\Delta\delta}$ as diagnostics of systematic offsets, and within-file standard deviations σ_α and σ_δ as diagnostics of random scatter.

Table 3 lists the file-level statistics for all 21 (file, catalog) groups.




Figure 2. All-sky spatial distributions of the correction terms for the USNO-A2.0 star catalog.




Figure 3. Same as Figure 2 but for the UCAC2 catalog.

To combine the *R.A.* and *decl.* components into a single scalar, we define the file-level offset norm

$$|\vec{r}| = \sqrt{\Delta\alpha^2 + \Delta\delta^2}, \quad (7)$$

and the improvement $\Delta|\vec{r}| = |\vec{r}|_{\text{before}} - |\vec{r}|_{\text{after}}$, where positive values indicate a reduction in systematic offset.

Figure 7 shows the improvement for all 21 groups. Files reduced with catalogs carrying large known biases show substantial positive improvements: [jg0032/AGK3](#) (+101 mas), [jg0043/UCAC2](#) (+24 mas), [jg0059/Gaia DR1](#) (+20 mas), and [jg0023/Tycho-2](#) (+21 mas). The two Gaia DR2 files show no significant change, as expected for a catalog whose corrections are negligible. Negative changes are




Figure 4. Example of RBF smoothing applied to the correction field.




Figure 5. Cumulative distributions of *intertile* discontinuities before and after smoothing. The CDF denotes the cumulative distribution function, $P(D_{\text{disc}} \leq x)$.




Figure 6. (a) Empirical cumulative distributions of the position offset Δ before and after debiasing. For a given value of Δ , the vertical coordinate gives the fraction of observations with Δ smaller than this value. (b) The value of Δ as a function of chronological observation index. Light translucent lines show individual measurements, while the dashed black and solid blue lines show the running-mean trends before and after debiasing, respectively.

confined to files with very limited statistics (jg0042 and

Table 3
File-level Ephemeris-comparison Residual Statistics

File	Catalog	N_{obs}	$\overline{\Delta\alpha}_{\text{before}}$ (mas)	$\overline{\Delta\alpha}_{\text{after}}$ (mas)	$\sigma_{\alpha,\text{before}}$ (mas)	$\sigma_{\alpha,\text{after}}$ (mas)	$\overline{\Delta\delta}_{\text{before}}$ (mas)	$\overline{\Delta\delta}_{\text{after}}$ (mas)	$\sigma_{\delta,\text{before}}$ (mas)	$\sigma_{\delta,\text{after}}$ (mas)
jg0032 ^a	AGK3	346	+84.45	+62.05	680.10	609.15	+1353.20	+1253.26	523.12	401.13
jg0011 ^b	Hipparcos	1297	+33.13	+30.39	181.24	178.89	−21.39	−18.65	163.15	167.65
jg0024 ^c	Hipparcos	643	+16.33	+14.52	188.31	188.24	+34.27	+34.63	368.15	369.88
jg0006 ^d	ACT	224	−34.53	−22.73	95.25	95.34	−40.25	−41.48	148.75	146.56
jg0005 ^e	ACT	218	−13.81	−22.00	124.69	123.83	−31.06	−30.81	162.26	160.77
jg0040 ^f	ACT	26	+35.58	+52.92	79.64	80.13	−46.73	−55.63	93.50	94.10
jg0040 ^g	Tycho-2	1821	+7.62	+12.32	134.14	132.31	−33.92	−30.65	139.46	139.08
jg0019 ^g	Tycho-2	727	+2.59	+4.29	102.47	101.95	−24.44	−21.90	112.55	112.22
jg0041 ^h	Tycho-2	330	+146.74	+140.33	128.94	124.03	−17.94	−15.09	127.44	126.54
jg0022 ⁱ	Tycho-2	229	−3.77	+8.95	112.59	109.88	−39.44	−41.21	124.49	124.85
jg0023 ^j	Tycho-2	85	+52.61	+43.63	84.76	83.44	−55.26	−33.26	136.57	135.40
jg0042 ^k	Tycho-2	9	−254.16	−247.89	170.47	166.82	+357.78	+372.75	84.52	86.57
jg0043 ^l	UCAC2	2523	−4.23	−10.79	421.95	422.43	+46.46	+19.80	498.26	498.28
jg0043 ^l	UCAC4	1238	−16.38	−24.49	461.99	461.39	+51.59	+58.40	466.53	467.22
jg0045 ^m	UCAC4	154	+17.15	−0.72	87.51	85.89	−22.14	−17.59	93.21	89.14
jg0044 ⁿ	UCAC4	136	+28.58	+39.63	188.75	188.08	−41.84	−48.67	175.54	172.35
jg0046 ^o	UCAC4	9	−25.12	−47.69	232.65	231.53	+4.67	−27.96	224.87	225.74
jg0054 ^p	Gaia DR1	761	+33.00	+24.97	69.36	69.75	+18.62	+12.62	83.88	83.97
jg0059 ^q	Gaia DR1	451	+25.59	+10.29	77.88	78.06	+22.04	+9.20	71.34	71.34
jg0073 ^r	Gaia DR2	295	+13.99	+14.01	64.82	64.81	−7.70	−7.74	43.54	43.53
jg0069 ^s	Gaia DR2	209	+44.29	+44.31	97.21	97.20	−12.36	−12.39	45.23	45.22

Note. $\overline{\Delta\alpha}$ and $\overline{\Delta\delta}$: file-level mean residual (systematic offset). σ_{α} and σ_{δ} : within-file standard deviation (random scatter). All values are in **milliarcseconds**.

- ^a https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0032.html
^b https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0011.html
^c https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0024.html
^d https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0006.html
^e https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0005.html
^f https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0040.html
^g https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0019.html
^h https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0041.html
ⁱ https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0022.html
^j https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0023.html
^k https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0042.html
^l https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0043.html
^m https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0045.html
ⁿ https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0044.html
^o https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0046.html
^p https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0054.html
^q https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0059.html
^r https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0073.html
^s https://www.sai.msu.ru/neb/nss/html/obspos/OBS_COLL/J/jg0069.html

jg0046, both with $N = 9$) and to a subset of UCAC4 files (jg0043/UCAC4: −9 mas; jg0044: −12 mas).

Figure 8 aggregates the results at the catalog level. In panel (a), **six** of **eight** catalogs fall below the diagonal: AGK3 decreases from 1356 to 1255 mas, UCAC2 from 47 to 23 mas, and Gaia DR1 from 36 to 23 mas. UCAC4 shows a slight increase of 7 mas, and Gaia DR2 is unchanged. In panel (b), all **eight** catalogs cluster tightly along the diagonal, with scatter ratios ($\sigma_{\text{after}}/\sigma_{\text{before}}$) between 0.85 and 1.00.

The systematic offsets are reduced for the majority of catalogs while the random scatter remains unchanged. This confirms that the debiasing correction acts on catalog-dependent systematic biases without altering the stochastic error structure of the observations. Together with the plate reduction test, the ephemeris comparison supports the effectiveness of the Gaia **DR3-based** debiasing model.

6. Conclusions

Using Gaia DR3 as the astrometric reference, we derive systematic position and proper-motion corrections for 17 historical star catalogs. A correction framework is developed based on HEALPix sky partitioning combined with RBF interpolation, enabling the construction of continuous all-sky correction fields while suppressing the discontinuities introduced by tile-based estimation. The resulting model preserves the intrinsic large-scale structure of catalog systematics while suppressing small-scale sampling noise.

The plate reduction test shows that the debiased data exhibit reduced systematic residuals while preserving the random scatter of the observations. The **ephemeris-comparison** test confirms that the file-level systematic offsets are reduced for the majority of catalogs without altering the within-file scatter. Both tests demonstrate that the proposed approach effectively mitigates catalog-dependent systematic offsets without degrading the random component of the residuals, leading to




Figure 7. File-level improvement in systematic offset $\Delta|\bar{r}|$ for each (file, catalog) group, ordered by catalog. Positive values indicate that the debiasing correction reduces the systematic offset with respect to the ephemeris.




Figure 8. Catalog-level comparison of residual statistics before and after debiasing. (a) Systematic offset $|\bar{r}|$: points below the diagonal indicate reduced offsets. (b) Within-catalog scatter σ : clustering along the diagonal confirms that the random component is preserved. Point sizes scale with the number of observations; colors indicate the reference catalog.

improved consistency in orbit determination solutions. The improvement is particularly relevant for historical observations that cannot be re-reduced in the Gaia reference frame.

Acknowledgments

This work has made use of the Natural Satellite Database (NSDB), maintained by the Laboratoire Temps Espace (LTE), which provides a comprehensive compilation of astrometric observations of planetary satellites. We are grateful to those who collect and maintain these datasets for their efforts.

This work also makes use of data from the European Space Agency (ESA) mission Gaia (<https://www.cosmos.esa.int/gaia>), processed by the Gaia Data Processing and Analysis Consortium (DPAC). Funding for the DPAC has been

provided by national institutions, in particular the institutions participating in the Gaia Multilateral Agreement.

We thank Yuan Ye (Purple Mountain Observatory, Chinese Academy of Sciences) for valuable discussions related to catalog systematics and astrometric data processing. We also thank S. Eggl for making the [catalog-debiasing](#) framework and reference implementation publicly available. This work was supported by the National Natural Science Foundation of China under [grant](#) No. 12473070 and the China Manned Space Program with [grant](#) No. CMS-CSST-2025-A16.

Q9

Data and Code Availability

The catalog correction tables and supporting materials are available for review through the National Astronomical Data Center (NADC) at [doi:10.12149/101824](https://doi.org/10.12149/101824).

For each of the 17 catalogs, two HEALPix correction tables are released: a baseline table at $N_{\text{side}} = 64$ derived directly from the cross-match statistics, and an RBF-smoothed full-sky table at $N_{\text{side}} = 256$. Each row corresponds to one HEALPix pixel (NESTED ordering) and contains the position corrections $\Delta\alpha$ and $\Delta\delta$ at epoch J2000.0 together with the [proper-motion](#) corrections $\Delta\mu_\alpha$ and $\Delta\mu_\delta$, all in the sense of (catalog—Gaia DR3). Sky maps and statistical distributions for every catalog are also provided.

To correct an observation, the relevant tile is identified from its sky position, the position correction at epoch J2000.0 is propagated to the observation epoch via the linear model given in Section 3, and the result is subtracted from the observed coordinates. A reference Python tool `query_bias.py` is included in the release, which performs the pixel lookup and epoch propagation automatically. Full usage instructions are given in the accompanying [README](#).⁶

ORCID iDs

H. Y. Zhang  <https://orcid.org/0000-0003-0956-2440>

K. Tang  <https://orcid.org/0000-0001-7747-5707>

References

Arlot, J.-E., Varfolomeev, M. I., & Emelyanov, N. V. 2024, [A&A](#), **691**, A295
 Astronomische Gesellschaft 1890, Catalog der Astronomischen Gesellschaft (Wilhelm Engelmann)
 Cantat-Gaudin, T., & Brandt, T. D. 2021, [A&A](#), **649**, A124
 Carpino, M., Milani, A., & Chesley, S. R. 2003, [Icar](#), **166**, 248
 Chesley, S. R., Baer, J., & Monet, D. G. 2010, [Icar](#), **210**, 158
 Corbin, T. E., Urban, S. E., & Warren, W. H., Jr. 1991, *Astrographic Catalogue Reference Stars* (Corbin and Urban 1991) NSSDC/WDC-A-R&S 91-10, National Space Science Data Center
 Desmars, J., Bancelin, D., Hestroffer, D., & Thuillot, W. 2013, [A&A](#), **554**, A32
 Eggl, S., Farnocchia, D., Chamberlin, A. B., & Chesley, S. R. 2020, [Icar](#), **339**, 113596
 Eichhorn, H. K. 1974, *Astronomy of Star Positions: A Critical Investigation of Star Catalogues, the Methods of Their Construction, and Their Purpose* (Frederick Ungar)
 European Space Agency 1997, *The Hipparcos and Tycho Catalogues*, ESA SP-1200 (ESA Publications Division)

Fabricius, C., Luri, X., Arenou, F., et al. 2021, [A&A](#), **649**, A5
 Farnocchia, D., Chesley, S. R., Chamberlin, A. B., & Tholen, D. J. 2015, [Icar](#), **245**, 94
 Fey, A. L., Gordon, D., Jacobs, C. S., et al. 2015, [AJ](#), **150**, 58
 Fricke, W., Kopff, A., Gliese, W., et al. 1963, [VeARI](#), **10**, 1
 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2016, [A&A](#), **595**, A2
 Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, [A&A](#), **616**, A1
 Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2023, [A&A](#), **674**, A1
 Górski, K. M., Hivon, E., Banday, A. J., et al. 2005, [ApJ](#), **622**, 759
 Heckmann, O., & Dieckvoss, W. 1975, *AGK3. Star Catalogue of Positions and Proper Motions North of -2.5 degrees Declination* (Hamburger Sternwarte)
 Hoffleit, D., & Warren, W. H., Jr. 1991, *The Bright Star Catalog* (5th ed.; Yale Univ. Observatory)
 Høg, E., Fabricius, C., Makarov, V. V., et al. 2000, [A&A](#), **355**, L27
 Lindegren, L., Bastian, U., Biermann, M., et al. 2021a, [A&A](#), **649**, A4
 Lindegren, L., Klioner, S. A., Hernández, J., et al. 2021b, [A&A](#), **649**, A2
 Lindegren, L., Lammers, U., Bastian, U., et al. 2016, [A&A](#), **595**, A4
 Ma, C., Arias, E. F., Eubanks, T. M., et al. 1998, [AJ](#), **116**, 516
 Monet, D. G. 1998, [BAAS](#), **30**, 1427
 Morrison, J. E., Roeser, S., McLean, B., Bucciarelli, B., & Lasker, B. 2001, [AJ](#), **121**, 1752
 Murray, C. A. 1989, [A&A](#), **218**, 325
 Park, R. S., Folkner, W. M., Williams, J. G., & Boggs, D. H. 2021, [AJ](#), **161**, 105
 Platais, I., Girard, T. M., Kozhurina-Platais, V., et al. 1998, [AJ](#), **116**, 2556
 Qiao, R., Shen, K., Liu, J., & Harper, D. 1999, [A&AS](#), **137**, 1
 Qiao, R. C., Tang, Z. H., Shen, K. X., et al. 2006, [A&A](#), **454**, 379
 Qiao, R. C., Yan, Y. R., Shen, K. X., et al. 2007, [MNRAS](#), **376**, 1707
 Roeser, S., & Bastian, U. 1991, *PPM Star Catalogue* (Spektrum Akademischer Verlag)
 Roeser, S., Demleitner, M., & Schilbach, E. 2010, [AJ](#), **139**, 2440
 Roeser, S. 1990, [yCat](#), **1**/154
 Schaback, R. 1995, in *Mathematical Methods for Curves and Surfaces*, ed. M. Dæhlen, T. Lyche, & L. L. Schumaker (Vanderbilt Univ. Press), 477
 Shen, K. X., Qiao, R. C., Harper, D., Hadjifotinou, K. G., & Liu, J. R. 2002, [A&A](#), **391**, 775
 Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, [AJ](#), **131**, 1163
 Smithsonian Astrophysical Observatory Staff 1966, *Star Catalog: Positions and Proper Motions of 258,997 Stars for the Epoch and Equinox of 1950.0* (Smithsonian Institution)
 Teixeira, R., Galli, P. A. B., Le Campion, J.-F., et al. 2014, [RMxAC](#), **43**, 55
 Tholen, D. J., Micheli, M., & Elliott, G. T. 2013, [Icar](#), **223**, 625
 Urban, S. E., Corbin, T. E., & Wycoff, G. L. 1998, [AJ](#), **115**, 2161
 Vasiliev, E. 2019, [MNRAS](#), **484**, 2832
 Wahba, G. 1990, *Spline Models for Observational Data* (SIAM)
 Yan, D., Qiao, R. C., Cheng, X., & Yu, Y. 2020, [Icar](#), **343**, 113662
 Yan, D., Qiao, R. C., Dourneau, G., et al. 2016, [MNRAS](#), **457**, 2900
 Yuan, Y., Li, F., Fu, Y., & Ren, S. 2021a, [A&A](#), **645**, A48
 Zacharias, N., Finch, C., & Frouard, J. 2017, [AJ](#), **153**, 166
 Zacharias, N., Finch, C. T., Girard, T. M., et al. 2013, [AJ](#), **145**, 44
 Zacharias, N., Urban, S. E., Zacharias, M. I., et al. 2004, [AJ](#), **127**, 3043

⁶ nadc.china-vo.org/res/file_upload/download?id=1109148