
OGRDBstats Report

Contents


1	Novel sequence analysis		2
	1.1	End-nucleotide composition	2
	1.2	Per-nucleotide consensus where previous nucleotides match the consensus	2
	1.3	Whole-sequence composition of each assigned read	2
	1.4	Final three nucleotides: frequency of each observed triplet	2
	1.5	CDR3 length distribution, in assignments to novel alleles	3
2	Variation from germline, in assignments to each allele		4
3	Allele usage in potential haplotype anchor genes		9
4	Нар	plotype plots	10
5	Con	nfiguration settings	11

1 Novel sequence analysis


1.1 End-nucleotide composition




1.2 Per-nucleotide consensus where previous nucleotides match the consensus




1.3 Whole-sequence composition of each assigned read




1.4 Final three nucleotides: frequency of each observed triplet





1.5 CDR3 length distribution, in assignments to novel alleles





2 Variation from germline, in assignments to each allele

















3 Allele usage in potential haplotype anchor genes



4 Haplotype plots

5 Configuration settings

```
## Repertoire file: /work/jenkins/PRJEB26509/S42/S42/45/45/91/8fac70050dc9beb579b3392696c7df/45_Finale_
##
## Germline reference file: /work/jenkins/PRJEB26509/S42/S42/45/45/91/8fac70050dc9beb579b3392696c7df/pe
##
## Novel allele file: /work/jenkins/PRJEB26509/S42/S42/45/91/8fac70050dc9beb579b3392696c7df/novel_se
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
```