

Weekly Clean Bulk Adjustment Protocol

Starting Point: 2,700 kcal/day | 155g protein, 375g carbs, 60g fat

Target: 0.25–0.35 kg weight gain per week

Your Stats: 71 kg, 185 cm, ~10-11% BF

What to Track Each Week

Daily Inputs (every morning)

- **Scale weight** (same time, after bathroom, before food/water)
- **Training performance** (note any strength PRs or reps added)

Weekly Inputs (every Friday)

- **W_avg:** Average of last 7 daily weights
 - **Weekly change:** W_avg this week minus W_avg last week
 - **Waist measurement:** Measure at narrowest point (above navel)
 - **Strength check:** Did you add weight or reps on main lifts?
 - **Recovery quality:** Rate 1-5 (sleep quality, soreness, energy)
-

Decision Rules (Apply Every Friday)

 **HOLD CALORIES (Do Nothing)**

When:

- Weekly change = **0.25–0.35 kg** ✓
- Strength is progressing ✓
- Recovery feels good (3-5/5) ✓

Action: Continue eating 2,700 kcal (or current calories) for another week.

ADD +150 KCAL/DAY

When:

- Weekly change = **<0.25 kg** for 2 consecutive weeks
- AND strength is stalling or declining
- AND recovery is okay (no major sleep/stress issues)

Action:

- New total: Current calories + 150 kcal
 - Add as: **+35g carbs** (or +30g carbs + 5g fat)
 - Apply starting next Monday
-

SUBTRACT -100 KCAL/DAY

When:

- Weekly change = **>0.40 kg** for 2 consecutive weeks
- OR waist increased >0.8 cm in one week
- OR weekly change >0.50 kg consistently

Action:

- New total: Current calories - 100 kcal
 - Remove as: **-25g carbs** (from training day meals)
 - Apply starting next Monday
-

**SMALL ADJUSTMENT +100 KCAL****When:**

- Weekly change = **<0.25 kg**
- BUT strength IS improving
- Body feels recovered

Action:

- New total: Current calories + 100 kcal
 - Add as: **+25g carbs**
 - Apply starting next Monday
-

Special Situations**Week 1 Exception**

- Weight will spike 1-3 kg (water + glycogen)
- **DO NOT ADJUST CALORIES**
- Just track and observe

Suspect Water Retention?

Signs:

- Sudden jump >0.7 kg overnight
- After high-sodium meal, poor sleep, or new supplement
- Waist measurement unchanged

Action: Wait one more week before adjusting

Strength Stalling But Weight Rising?

If: Gaining 0.25-0.35 kg/week but no strength gains for 2+ weeks

Action:

- Redistribute carbs: more pre/post workout, less on rest days
- Consider deload week
- Add +50-100 kcal if needed

Activity Level Changed?

If: Daily steps decreased >2,000 for several days

Action: Expect temporary weight retention; wait one week before adjusting

Weekly Template

WEEK 1 (Oct 10-16, 2025)

Calories: 2,700/day

Macros: 155p / 375c / 60f

Action: Track only, NO adjustments (water retention week)

WEEK 2 (Oct 17-23)

Calculate:

- W_{avg} Week 2 vs Week 1 = ?
- Waist change = ?

Action: Track only, NO adjustments (stabilization week)

WEEK 3 (Oct 24-30)

Calculate:

- W_{avg} Week 3 vs Week 2 = ?
- Waist change = ?

Action: Track only, NO adjustments (need 2 weeks of data)

WEEK 4 (Oct 31 - Nov 6)

Calculate:

- Average weekly change from Week 2→3 and Week 3→4
- Waist trend over 2 weeks

Action: FIRST ADJUSTMENT POINT

Apply decision rules:

- If average gain = 0.25-0.35 kg/week → HOLD
- If average gain <0.25 kg/week → ADD +150 kcal
- If average gain >0.40 kg/week → SUBTRACT -100 kcal

WEEK 5+ (Ongoing)

Every Friday:

1. Calculate W_avg and weekly change
 2. Compare to last 2 weeks' trend
 3. Check waist measurement
 4. Review strength progress
 5. Apply decision rules
 6. Adjust calories if needed (starting next Monday)
 7. Wait 2 weeks before next adjustment
-

Quick Reference Card

Weekly Weight Change	Strength Progress	Action
0.25-0.35 kg ✓	Improving ✓	HOLD
<0.25 kg	Stalling	+150 kcal
<0.25 kg	Improving	+100 kcal
>0.40 kg	Any	-100 kcal
0.25-0.35 kg	Stalling 2+ weeks	Redistribute carbs or +50-100 kcal

Waist Override: If waist jumps >0.8 cm in one week → **-100 kcal** regardless of weight change

Monthly Check-In (Every 4 Weeks)

Take full measurements:

- Neck, shoulders, chest, waist, hips, arms, thighs
- Progress photos (same lighting, same poses)
- Body fat estimate (visual or Navy method)

Goal: Chest/shoulders growing faster than waist

Red Flag: Waist growing as fast or faster than upper body

Notes

- All adjustments apply starting the Monday after your Friday decision
- Never adjust more frequently than every 2 weeks
- When in doubt, wait another week and gather more data
- Log everything: weight, calories, training, recovery, sleep quality