

Myalgic Encephalomyelitis / Chronic Fatigue Syndrome

A Comprehensive Medical Documentation

Ing. Yannick Loth, M.Sc. (Management)

Software Architect & Patient-Researcher

Messancy, Belgium

yl@infolead.eu

ORCID: [0009-0003-5754-827X](https://orcid.org/0009-0003-5754-827X)

DOI: [10.5281/zenodo.18905326](https://doi.org/10.5281/zenodo.18905326)

March 7, 2026

Version 5

Abstract

Myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) is a severe, chronic, multi-system neuroimmune disease affecting an estimated 0.89% to 2.5% of the global population. Characterized by profound post-exertional malaise, unrefreshing sleep, cognitive dysfunction, and autonomic dysregulation, ME/CFS represents one of the most disabling chronic conditions in modern medicine. Despite affecting millions worldwide, the disease has historically suffered from underfunding, dismissal by medical professionals, and classification as a syndrome rather than a disease with identifiable pathophysiology.

The February 2024 NIH deep phenotyping study fundamentally transformed this landscape by demonstrating specific biological abnormalities: decreased brain activity in effort-related neural circuits, exhausted T-cell populations, chronic B-cell activation deficits, and depleted catecholamine levels in cerebrospinal fluid. These findings conclusively established ME/CFS as a systemic biological disease with measurable immune, neurological, and metabolic dysfunction.

This comprehensive documentation synthesizes current research across clinical presentation, pathophysiological mechanisms, treatment approaches, research evidence, and quantitative modeling. The work integrates findings from hundreds of peer-reviewed sources spanning energy metabolism dysfunction, immune exhaustion, neuroinflammation, endocrine dysregulation, cardiovascular abnormalities, gut-brain axis disruption, and genetic-epigenetic factors.

Part I (Chapters 1–5) provides detailed clinical characterization of core and additional symptoms, the evolution of diagnostic criteria from Fukuda through the Canadian Consensus and International Consensus Criteria, and disease course variations from mild to very severe presentations including severity-specific phenotyping. Part II (Chapters 6–15) examines established and hypothetical pathophysiological mechanisms across ten chapters covering mitochondrial dysfunction, immune exhaustion, neuroinflammation, endocrine dysregulation, cardiovascular and autonomic failure, gut-microbiome disruption, genetic and epigenetic factors, integrative multi-system models, speculative mechanistic hypotheses with falsifiable predictions, and symptom-producing mechanisms. Part III (Chapters 14a–19) opens with urgent action protocols for severe and very severe cases, followed by pediatric-specific management for both housebound and ambulatory children, then documents symptom-based management, medications targeting underlying mechanisms, supplement regimens, lifestyle interventions, emerging therapies, and integrative personalized approaches. Part IV (Chapters 20–25b) synthesizes biomarker research, clinical trial outcomes, mechanistic studies, epidemiological patterns, research controversies, translational findings, research infrastructure proposals, and a set of fully specified proposed studies with explicit hypotheses and trial designs. Part V (Chapters 26–32) develops formal mathematical models of disease dynamics using ordinary differential equation systems, presenting energy metabolism models, immune

system dynamics, neuroendocrine and autonomic regulation, integrated multi-system coupling, temporal evolution with tipping-point dynamics, and predictive clinical applications for pacing optimization and treatment selection.

Nine appendices provide reference material: a glossary of medical and scientific terms, abbreviations, diagnostic tools and assessment scales, patient resources, mathematical derivations supporting Part V, supplement protocols, a research synthesis, an annotated bibliography of landmark studies, and a registry of ongoing and planned ME/CFS studies.

Methodologically, this work uses a structured epistemic classification system distinguishing achievements (high-certainty findings from replicated studies), hypotheses (certainty ≥ 0.45 , requiring validation), speculations (certainty < 0.45 , exploratory), predictions (testable claims for future research), warnings (critical limitations and contraindications), and recommendations (evidence-based guidance requiring physician review). Evidence quality is systematically classified as high, medium, or low certainty based on sample size, peer-review status, replication, and methodological rigor. A cross-document hypothesis registry tracks all speculative content with falsifiability assessments.

This documentation serves multiple audiences: researchers seeking comprehensive mechanistic understanding and quantitative modeling frameworks, clinicians requiring evidence-based treatment protocols with dosing guidance and contraindication awareness, patients and caregivers needing accessible explanations of symptoms and management strategies, and advocates working toward recognition, funding, and medical education reform. The work is released under the Creative Commons Attribution 4.0 International License to maximize accessibility and enable derivative works.

Written by a software architect and patient-researcher with degrees in industrial engineering and management sciences, this documentation applies systems thinking, computational analysis, and first-principles reasoning to ME/CFS pathophysiology while maintaining epistemic humility about the substantial uncertainties remaining in the field. The author explicitly disclaims medical expertise and emphasizes that all content represents literature synthesis and personal experience documentation, not clinical advice. All treatment decisions must be made in consultation with qualified healthcare providers.

ME/CFS research is at a critical inflection point. The biological validation provided by recent NIH and international studies, combined with shared research agendas driven by Long COVID parallels, offers unprecedented opportunity for mechanistic discovery and therapeutic development. This document aims to accelerate progress by organizing scattered findings into an accessible, comprehensive reference while identifying critical knowledge gaps requiring focused investigation.

Version Notice

★ Achievement: Living Document

This is an actively maintained, regularly updated document. The research landscape for ME/CFS is evolving rapidly, and this documentation is continuously expanded and refined to reflect new findings, improved analysis, and emerging therapeutic approaches.

To ensure you have the most current version:

- Visit the Zenodo repository: <https://doi.org/10.5281/zenodo.18370021>
- Check the publication date: This version was compiled on **March 7, 2026**
- Bookmark the DOI link above and check periodically for updates
- Significant updates typically occur monthly as new research emerges and analysis deepens

Using an outdated version may mean missing critical research findings, treatment protocol refinements, or corrections to earlier content. The difference between versions can be substantial, particularly in rapidly evolving areas such as biomarker research, immune dysfunction mechanisms, and emerging therapies.

Version control: All versions are permanently archived on Zenodo with full change documentation. You can compare versions to identify what has been updated since your last reading.

Keywords

Keywords: Myalgic encephalomyelitis, chronic fatigue syndrome, ME/CFS, post-exertional malaise, PEM, mitochondrial dysfunction, neuroinflammation, immune dysfunction, autonomic dysregulation, orthostatic intolerance, POTS, cognitive impairment, brain fog, energy metabolism, oxidative stress, cytokines, biomarkers, chronic illness, multi-system disease, neuroimmune disease, Long COVID, systems biology, mathematical modeling, HPA axis, gut microbiome, mast cell activation, treatment protocols, pacing, patient-reported outcomes

License and Usage

License

License: This work is licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0). You are free to share and adapt this work, provided you give appropriate credit. <https://creativecommons.org/licenses/by/4.0/>

Medical Disclaimer

CRITICAL: The author is not a medical doctor or licensed healthcare provider. This documentation represents independent patient research and should not be construed as medical advice.

△ Warning 1: Medical Disclaimer

- This document is provided for **informational and educational purposes only**
- The author is **not a physician** and has no formal medical training
- This is **not a substitute** for professional medical advice, diagnosis, or treatment
- **Always consult qualified healthcare providers** before making any medical decisions
- Anyone who follows recommendations or protocols in this document **without consulting their physician does so entirely at their own risk and responsibility**
- The information presented represents a synthesis of current research but should **never be used for self-diagnosis or self-treatment**
- Treatment decisions should only be made in consultation with licensed medical professionals who can evaluate your individual circumstances
- The personal case data in Appendix I documents one individual's experience and should not be generalized to others

The author assumes no liability for any adverse effects or consequences resulting from the use of information contained in this document. All treatment protocols and medical recommendations discussed herein require physician oversight and should be adapted to individual patient circumstances.

Citation

When citing this work, please use the following format:

Loth, Y. (2026). *Myalgic Encephalomyelitis / Chronic Fatigue Syndrome: A Comprehensive Medical Documentation*. DOI: [10.5281/zenodo.18905326](https://doi.org/10.5281/zenodo.18905326)

About the Author

Yannick Loth is a software architect and independent patient-researcher with degrees in Industrial Engineering (Information Systems, University of Luxembourg) and Management Sciences (General Management, HEC Liège), with prior completion of first-cycle civil engineering studies (University of Liège). With nearly two decades of professional software engineering experience and iSAQB CPSA-F certification (2015), he brings computational thinking and systems analysis to medical research.

Having experienced symptoms consistent with ME/CFS since childhood, with progressive worsening over the past decade and marked acceleration in recent years, Yannick has applied his background in discrete mathematics, information systems architecture, and analytical research to understanding this complex multisystem disease (though formal diagnostic documentation has not yet been received). This work-in-progress represents an ongoing effort to synthesize the current state of ME/CFS research into a comprehensive, accessible reference while documenting his own case with scientific rigor.

Based in Messancy, Belgium, he has published research on software architecture principles (notably the Independent Variation Principle) and fundamental physics (Causal Graph Theory), now applying similar first-principles thinking to understanding chronic illness mechanisms.

Contact Information

- Email: yl@infolead.eu
- ORCID: [0009-0003-5754-827X](https://orcid.org/0009-0003-5754-827X)

Motivation

This documentation project is driven by both intellectual rigor and existential urgency. Witnessing the devastating reality of severe ME/CFS through videos and posts from bedbound patients has crystallized a stark question: how to prevent deterioration into severe illness before all desire to live vanishes? With a family, children, and friends depending on me, the stakes extend beyond personal survival—I cannot risk descending into a state where I lose the capacity to be present for those I love. This work aims to bridge the gap between scattered research findings and accessible, comprehensive information about ME/CFS—organizing current knowledge about symptoms, mechanisms, and treatments while rigorously documenting a progressive case with quantitative data. By systematically analyzing what is known and identifying what must be discovered, this project seeks to serve researchers, clinicians, patients, and advocates while racing against time to find pathways out of progressive decline.

AI Disclosure Statement

This manuscript was developed through extensive collaboration between a human author and AI language models. In the interest of scientific transparency, this statement describes the nature and extent of each party's contributions.

Author's Contributions

The author (Yannick Loth) contributed:

- **Lived experience:** Direct, first-person experience of severe illness with symptoms consistent with ME/CFS (formal diagnostic documentation pending) providing the phenomenological foundation and motivation for this work
- **Research direction:** Identifying research gaps, selecting topics for investigation, and determining which mechanisms and treatments warranted detailed exploration
- **Literature selection:** Choosing which papers to prioritize, which findings were most significant, and how to organize the overwhelming volume of ME/CFS research into a coherent framework
- **Critical evaluation:** Assessing study quality, identifying methodological limitations, distinguishing high-certainty findings from speculative claims, and evaluating evidence strength
- **Clinical data:** All personal case data in Appendix I, including symptom tracking, medication trials, and functional capacity measurements collected through lived experience with the disease
- **Structural decisions:** Choosing theorem-like environments (hypothesis, achievement, warning, etc.) to make epistemic status of claims immediately clear—a structural choice designed to facilitate critical review
- **Systems analysis:** Applying software architecture thinking to understand ME/CFS as a complex multisystem disease, identifying potential mechanistic relationships and integration points across physiological systems
- **Quality control:** Conducting extensive review cycles to verify medical accuracy, logical consistency, and appropriate citation of sources
- **Organizational framework:** Deciding document structure, what content to include, how to balance comprehensiveness with accessibility, and how to serve multiple audiences (patients, clinicians, researchers)

AI Contributions

AI language models (primarily Claude Sonnet 4.5 and Opus 4.5, Anthropic Inc.) performed:

- **Literature synthesis:** Processing and summarizing large volumes of research papers, extracting key findings, and organizing information thematically
- **Technical exposition:** Drafting explanatory text for complex biological mechanisms, translating technical research into accessible language
- **Citation management:** Identifying relevant studies, formatting references, managing bibliography, and ensuring proper attribution
- **LaTeX preparation:** Writing and formatting LaTeX source code, creating document structure, managing cross-references and environments
- **Consistency checking:** Identifying contradictions, checking internal consistency, and verifying that claims match cited sources

Nature of the Collaboration

This work represents a new mode of medical documentation in which the author's lived experience, clinical judgment, and research direction combined with AI's information processing capabilities. The author provided the conceptual framework, selected research priorities, and maintained continuous quality control throughout development.

The manuscript exceeds 1900 pages spanning clinical symptomatology, multisystem pathophysiology, treatment protocols, and research synthesis. The author has spent extensive time reviewing this material to ensure medical accuracy and appropriate epistemic calibration—distinguishing established findings from preliminary results and clearly marking speculative content.

AI frequently required redirection when synthesizing research, occasionally missing nuances in study design, overstating certainty, or losing focus on ME/CFS-specific findings. A substantial portion of the author's effort involved recognizing when outputs missed important caveats, diagnosing misunderstandings of research context, and redirecting toward more accurate representations.

This collaboration enabled processing a volume of literature that would be extremely challenging for a single individual, particularly one disabled by the disease being studied. The combination of AI's processing capabilities with the author's continuous strategic direction, clinical insight from lived experience, and exhaustive quality control made this comprehensive documentation possible.

Author Responsibility

Despite the substantial AI contribution, the author takes full responsibility for:

- The decision to publish and disseminate this work
- All medical claims, treatment discussions, and mechanistic explanations
- Any errors, misconceptions, or unjustified conclusions
- The interpretation and implications of research findings
- Personal case data and clinical observations

Critical disclaimer: The author is **not a medical doctor** and has no formal medical training. While he holds engineering degrees and received formal training in mathematics and analytical methods, he is not qualified to provide medical advice. This work represents independent patient research and systematic literature review, not clinical guidance.

The medical and scientific communities are invited to review this work critically. Such scrutiny is essential before any claims or treatment approaches discussed here can be considered validated for clinical use.

Transparency and Reproducibility

- This disclosure is made voluntarily and in good faith.
- The manuscript was developed using Claude Sonnet 4.5 and Claude Opus 4.5 (Anthropic).
- The manuscript source is available at: github.com/yannickloth/health-me-cfs
- Correspondence regarding the content should be directed to the author.

Author: Ing. Yannick Loth, M.Sc. (Management)

Date: March 7, 2026

ORCID: [0009-0003-5754-827X](https://orcid.org/0009-0003-5754-827X)

Methodology and Quality Assurance

A document of this scope—over 1900 pages of medical content spanning pathophysiology, treatment, research synthesis, and mathematical modeling—produced through human–AI collaboration and without formal medical training on the author’s part, demands rigorous and transparent quality assurance. This chapter describes the methodology used to maintain scientific accuracy, epistemic honesty, and internal consistency throughout the manuscript.

Epistemic Classification

Every substantive scientific claim in this document is assigned an explicit epistemic status through structured L^AT_EX environments. This design choice ensures that readers can immediately distinguish established findings from working theories and speculative ideas, without needing to evaluate each claim from context alone.

Claim environments. The following environments are used consistently throughout the manuscript:

Achievement A research finding supported by strong evidence: replicated results from peer-reviewed studies with adequate sample sizes and sound methodology. These represent the most reliable claims in the document.

Hypothesis An unproven conjecture or working theory with moderate plausibility (certainty ≥ 0.45). Hypotheses are clearly marked because they may be wrong. Each hypothesis is required to include at least one falsifiable prediction, identified limitations, and relevant citations.

Speculation A conjecture with lower plausibility (certainty < 0.45), often based on limited evidence, mechanistic reasoning, or extrapolation from related conditions. Speculations are held to the same structural requirements as hypotheses but carry explicit low-confidence warnings.

Prediction A testable claim about future observations or experimental outcomes. Predictions specify what research should find if a given hypothesis is correct, providing a concrete path toward validation or falsification.

Warning A critical caveat about limitations, risks, or potential misinterpretations. Warnings flag contraindications, preliminary evidence, or situations where claims should be interpreted with particular caution.

Limitation An explicit epistemic boundary marking what the available data cannot conclude. Distinct from warnings (which address practical caveats), limitations delineate the outer edges of justified inference.

Recommendation An evidence-based clinical or management suggestion. Recommendations are distinguished from hypotheses: a recommendation says “consider doing this,” grounded in clinical evidence, whereas a hypothesis says “this might be true,” grounded in mechanistic reasoning. This distinction is enforced through automated review (see *Automated Multi-Layer Review* below).

Certainty assessment. Research findings cited throughout the document are classified by evidence strength:

High certainty Large sample size ($n > 100$), peer-reviewed in a reputable journal, independently replicated, consistent across studies.

Medium certainty Moderate sample ($n = 20$ – 100), peer-reviewed but limited replication, sound methodology. Promising but requiring confirmation.

Low certainty Small sample ($n < 20$), preprint or conference abstract, methodological concerns, or contradicted by other studies.

Replication status. Key findings are annotated with their replication status: *fully replicated* (confirmed by independent cohorts), *partially replicated* (some confirmation with caveats), or *not yet replicated* (single study or research group). This classification reflects the reality that much ME/CFS research remains at early replication stages due to historical underfunding.

Literature Integration

Research papers are integrated through a structured pipeline designed to ensure traceability and appropriate epistemic calibration.

Discovery and selection. The author identifies research directions and selects papers for integration based on relevance to ME/CFS pathophysiology, treatment, or modeling. AI assists with literature search, retrieval, and initial summarization. The author evaluates study quality, assesses methodological limitations, and determines which findings warrant inclusion.

Integration protocol. Each integrated paper passes through a defined sequence:

1. **Bibliographic entry:** Full citation added to the bibliography with complete metadata (authors, title, journal, DOI, year).
2. **Certainty assessment:** Evidence strength classified as high, medium, or low based on sample size, peer-review status, replication, and methodology.
3. **Content placement:** Findings inserted into the appropriate chapter using the correct epistemic environment (achievement, hypothesis, or speculation) with explicit certainty annotation.

4. **Cross-referencing:** Links established to related content in other chapters, the hypothesis registry, and the annotated bibliography.

Annotated bibliography. Appendix [H](#) provides structured summaries of key papers organized by topic, enabling readers to evaluate primary sources directly.

Automated Multi-Layer Review

The manuscript undergoes systematic automated review through a pipeline of specialized auditing agents. Each agent is an AI-driven reviewer with a narrow, well-defined audit scope. Agents operate in read-only mode: they identify and report issues but do not modify text autonomously. All corrections are reviewed and applied by the author.

The review system comprises 20 audit categories organized into six phases, ordered from structural checks (fast, mechanical) to deep domain-specific analysis (slow, semantic).

Phase 1: Structural and mechanical. Catches low-level issues that could cause cascading false positives in later phases. Includes build verification (clean $\text{\LaTeX}$ compilation with no errors), internal reference validation (broken `\ref{}`, `\cite{}`, `\label{}` detection), bibliography integrity (duplicate entries, missing fields, orphan references), and structural completeness (empty sections, stub detection, chapter balance).

Phase 2: Terminology and style. Enforces consistency and readability. Includes medical terminology harmonization across all chapters (ensuring consistent usage of terms such as “post-exertional malaise” versus “PEM”), coherence and flow review, and detection of AI-generated writing patterns (excessive lists, mechanical sentence structures, unnatural transitions).

Phase 3: Scientific rigor. The core audit layer. Verifies that substantive claims carry citations, that hypotheses and speculations use the correct environments, that falsifiable predictions accompany hypotheses, and that reasoning avoids circularity or completeness gaps. This phase also cross-validates the hypothesis registry (see *Cross-Document Consistency* below) against chapter content.

Phase 4: Domain-specific biology. Deep audits by specialized reviewers, each focusing on a single biological domain. Eight domain auditors cover: biological pathway accuracy (correct intermediaries, current nomenclature, gene-versus-protein distinction), immunology (cell subset precision, cytokine context, assay limitations), biochemistry and metabolomics (pathway stoichiometry, redox chemistry, mitochondrial specificity), neurology (brain region specificity, HPA axis nuance, blood-brain barrier considerations), microbiome (methodology distinction between 16S rRNA and shotgun sequencing, correlation-versus-causation framing, strain-level precision), biomarker claims (sensitivity and specificity reporting, reference

ranges), epidemiology (case definition consistency, selection bias awareness, sex and gender stratification), and comorbidity (overlap condition delineation for hEDS, MCAS, POTS, fibromyalgia, and Long COVID).

Only domain auditors relevant to the content under review are activated, determined by keyword matching against the target files.

Phase 5: Safety and patient impact. Critical safety review that runs on every audit pass. Three auditors check for: hypotheses incorrectly framed as clinical recommendations, missing contraindications and drug interaction warnings, dosing errors, off-label use transparency, withdrawal risk documentation, and patient-facing content safety (ensuring that activity recommendations do not risk triggering post-exertional malaise, that severe and very severe presentations are adequately represented, and that psychological framing does not perpetuate outdated psychogenic models of ME/CFS).

Phase 6: Publication readiness. Final pass covering metadata completeness, licensing, AI disclosure, and—for Part V—mathematical verification of units, parameters, and notation consistency.

Adversarial convergence. Beyond the comprehensive 20-category review, the manuscript also undergoes adversarial review-fix loops on individual sections. In this process, a reviewer agent identifies issues, the author applies corrections, and the reviewer re-audits the corrected text. This cycle repeats until convergence (zero findings) or a maximum of 12 rounds, ensuring that fixes do not introduce new problems.

Author Critical Oversight

The automated review system described above is necessary but not sufficient. The most consequential quality control mechanism in this document is the author's continuous critical engagement with AI-generated content. This takes several forms.

Directed questioning. The author systematically challenges AI outputs through targeted questions: *What is the actual sample size of this study? Has this finding been replicated independently? Is this mechanism established or hypothetical? Does this treatment recommendation follow from the cited evidence, or from mechanistic reasoning alone?* These questions frequently reveal overstatements, missing qualifications, or conflation of related but distinct findings. AI systems tend to present information with uniform confidence; the author's role is to impose appropriate epistemic gradation.

Redirection and correction. AI frequently requires correction when synthesizing research. Common failure modes include: overstating the certainty of preliminary findings, losing ME/CFS-specific context when drawing on general medical knowledge, conflating correlation with causation in observational studies, missing important caveats in study design (e.g., healthy control selection, diagnostic criteria heterogeneity), and presenting mechanistic hypotheses as though they were established pathophysiology. A substantial portion of the author’s effort involves recognizing these patterns and redirecting toward more accurate representations.

Conceptual integration. While AI excels at processing individual papers and generating well-structured text, the author provides the cross-disciplinary reasoning that connects findings across chapters. Identifying that a cardiovascular abnormality described in Chapter 10 may share a mechanistic basis with an immune finding in Chapter 7, or that a treatment protocol in Chapter 16 rests on a hypothesis that has since been weakened by new evidence in Chapter 22—these integrative judgments require a sustained mental model of the entire document that emerges from the author’s continuous engagement over the course of writing.

Lived experience as epistemic resource. The author’s direct experience with ME/CFS provides a form of validation unavailable to external reviewers. When AI-generated text describes post-exertional malaise as “mild fatigue after exercise,” or characterizes cognitive dysfunction as “occasional forgetfulness,” the author can identify these as clinically inaccurate minimizations. Lived experience does not substitute for clinical expertise, but it provides a ground-truth check against descriptions that diverge from the phenomenology of the disease.

Formal Modeling Pipeline

Part V of this document presents mathematical and computational models of ME/CFS pathophysiology. These models are constructed through a structured pipeline with explicit uncertainty quantification at every stage.

Three-layer formalization. Each biological process amenable to formal modeling is represented at three levels of abstraction:

1. **Causal directed acyclic graphs (DAGs):** Represent causal relationships between biological variables with edge-level certainty annotations derived from the literature. DAGs make causal assumptions explicit and testable.
2. **Event-driven process chains (EPCs):** Capture the temporal sequence of biological events, including branching points, feedback loops, and probabilistic transitions. EPCs bridge the gap between qualitative causal understanding and quantitative dynamics.
3. **Ordinary differential equations (ODEs):** Quantitative models with parameters estimated from published data where available, and explicitly flagged as assumed where data are lacking. Parameter uncertainty is propagated through the model hierarchy.

Cross-validation. A model auditor cross-validates consistency across the three layers: certainty values in the DAG must align with confidence levels in the EPC and parameter ranges in the ODE. An uncertainty analyst propagates certainty through the hierarchy and flags model predictions that exceed the confidence warranted by available evidence. An evidence mapper maintains a reverse index from citations to the specific DAG edges, EPC events, and ODE parameters they support, ensuring that every model component is traceable to its evidential basis.

Cross-Document Consistency

Maintaining consistency across over 1900 pages requires systematic tracking mechanisms.

Hypothesis registry. A cross-document registry tracks every hypothesis, speculation, prediction, and open question in the manuscript. When any such environment is added, modified, or removed in any chapter, the registry is updated to reflect the change. This prevents contradictions between chapters, ensures that predictions stated in one chapter are not contradicted by evidence presented in another, and provides a single reference point for the document's epistemic commitments.

Terminology harmonization. Automated audits verify that medical terminology is used consistently across all chapters. Variant forms of key terms (e.g., "NK cells" versus "natural killer cells," "OI" versus "orthostatic intolerance") are detected and normalized.

Revision tracking. Changes are tracked through the document's version-controlled changelog (Appendix), which records what changed in each version and why. Git history provides full traceability of every modification at the line level.

Limitations of This Methodology

This quality assurance system, while systematic, has inherent limitations that the reader should bear in mind.

AI reviewer limitations. Automated auditors operate on textual pattern matching and language model reasoning. They can detect structural issues (missing citations, environment misuse, terminology inconsistency) with high reliability, but their ability to evaluate the *correctness* of biological claims is bounded by their training data. Novel findings, recently retracted papers, or subtle methodological flaws in cited studies may escape detection. The automated review system complements but does not replace expert human review.

Single-author limitation. Despite the multi-layered review process, this document is produced by a single non-physician author. There is no independent human peer review prior to publication. The author has mitigated this through extensive self-review, adversarial automated auditing, and transparent epistemic classification, but single-author bias remains an inherent risk. The medical and scientific communities are invited to review this work critically.

Evidence base limitations. ME/CFS research suffers from decades of underfunding, resulting in small sample sizes, limited replication, and heterogeneous patient populations across studies. The certainty classifications used in this document reflect these constraints honestly, but readers should recognize that even “high certainty” findings in ME/CFS research may not meet the evidentiary standards typical of better-funded medical fields.

Evolving methodology. The quality assurance system described here has been developed iteratively alongside the document itself. Earlier chapters may not have undergone the same depth of review as later ones. The revision tracking system helps identify which content has been recently reviewed, but comprehensive re-auditing of the entire document is an ongoing process.

The methodology described in this chapter aims to make the document’s epistemic commitments, evidence standards, and quality processes fully transparent. Readers are encouraged to use this information when evaluating the reliability of specific claims throughout the manuscript.

Reading Guide: How to Use This Document

This comprehensive documentation is organized to serve multiple audiences: researchers, clinicians, patients, caregivers, and advocates. This guide explains the document structure and how to interpret the specialized environments used throughout.

Document Organization

The document is divided into five main parts:

Part I: Clinical Overview Covers symptoms, diagnostic criteria, disease course, and clinical presentation. Start here for understanding what ME/CFS is and how it manifests.

Part II: Pathophysiology Explores biological mechanisms—known, suspected, and speculative. Essential for understanding the multisystem nature of the disease.

Part III: Treatment and Management Documents medications, supplements, lifestyle interventions, and management strategies. Includes both evidence-based approaches and emerging therapies.

Part IV: Research and Evidence Synthesizes current research, clinical trials, biomarker studies, and epidemiology. Provides detailed summaries of key findings.

Part V: Mathematical Modeling Presents computational and mathematical approaches to understanding ME/CFS systems biology (advanced/technical).

Understanding Statement Types

This manuscript uses formal environments to classify statements by their epistemic status and evidence strength. Understanding these distinctions is essential for critically evaluating medical claims.

Scientific Claims

Achievement A well-established research finding with strong evidence. Achievements represent replicated results from peer-reviewed studies with adequate sample sizes and methodological rigor. These are the most reliable claims in the document.

Hypothesis An unproven conjecture or working theory. Hypotheses are clearly marked because they may be wrong. Many ME/CFS mechanisms remain hypothetical due to limited research funding and methodological challenges.

Prediction A testable claim about future observations or experimental outcomes. Predictions specify what research should find if a hypothesis is correct, providing a path to validation or falsification.

Requirement A necessary condition for a diagnosis, treatment, or research interpretation to be valid. Requirements specify what must be true for a claim to hold.

Warning A critical caveat about limitations, risks, or potential misinterpretations. Warnings flag where treatments may be contraindicated, where research is preliminary, or where claims should be interpreted cautiously.

Evidence Quality Levels

Throughout this document, research findings are classified by evidence strength:

High Certainty Large sample size ($n > 100$), peer-reviewed in reputable journal, independently replicated, consistent across studies. Can be cited with confidence.

Medium Certainty Moderate sample ($n = 20\text{--}100$), peer-reviewed but single study or limited replication, sound methodology. Promising but requires confirmation.

Low Certainty Small sample ($n < 20$), preprint or conference abstract, methodological concerns, or contradicted by other studies. Noted as preliminary.

Navigation Tips

- Use the detailed Table of Contents to locate specific topics
- Cross-references appear as clickable hyperlinks in the PDF
- The Index provides quick access to terms and concepts
- Citations link to the Bibliography for full reference details
- Appendix H contains annotated summaries of key papers
- Appendix I documents the author's personal case data

For Different Readers

Patients and Caregivers: Focus on Part I (Clinical Overview) and Part III (Treatment). The pathophysiology sections may be technical but can help understand symptom mechanisms. Part V (Mathematical Modeling) is optional and highly technical.

Clinicians: All sections are relevant. Part II provides mechanistic understanding, Parts III and IV offer evidence-based treatment guidance, and Appendix I presents a detailed case study with quantitative tracking.

Researchers: Parts II, IV, and V provide detailed mechanistic insights, research synthesis, and modeling approaches. Appendix H contains literature summaries organized by topic.

Critical Reading Advice

When evaluating medical claims in this document:

1. **Check the evidence level.** High-certainty findings are more reliable than preliminary results. Many ME/CFS mechanisms remain speculative due to limited research.
2. **Distinguish established from hypothetical.** Results in achievement environments represent replicated findings. Results in hypothesis environments are working theories that may be revised.
3. **Note the warnings.** Limitations acknowledged in warning environments indicate where the author recognizes uncertainty or potential problems.
4. **Remember the author is not a physician.** This work represents independent patient research and literature synthesis, not clinical guidance. All treatment decisions require physician oversight.
5. **Recognize individual variation.** ME/CFS presents heterogeneously. The personal case data in Appendix I documents one individual's experience and may not generalize to others.
6. **Consider publication date.** ME/CFS research is rapidly evolving. This document reflects knowledge current at time of publication.

Medical Disclaimer

This is not medical advice. The author has no medical training. This work synthesizes research literature and documents one individual's experience for educational purposes. Always consult qualified healthcare providers before making medical decisions.

Updates and Corrections

This is a living document. Updates will be published as new research emerges. The source code is available at <https://github.com/yannickloth/health-me-cfs>. Errors or omissions can be reported to the author via email.

Frequently Asked Questions for Patients

This section addresses common questions from patients and caregivers, with short answers and references to the detailed evidence in the main text.

What is ME/CFS, and how is it different from “being tired”? ME/CFS is a complex, multi-system disease characterized by profound fatigue that is not relieved by rest, worsening of symptoms after physical or mental exertion (post-exertional malaise, or PEM), unrefreshing sleep, cognitive dysfunction, and orthostatic intolerance. Unlike ordinary tiredness, ME/CFS fatigue does not improve with rest and is accompanied by measurable biological abnormalities. See Chapter 1 for definitions and Chapter 2 for symptom details.

What is PEM and why is it important? Post-exertional malaise (PEM) is the hallmark feature of ME/CFS: a disproportionate worsening of symptoms following physical, cognitive, or emotional exertion, typically delayed by 12–72 hours and lasting days to weeks. PEM distinguishes ME/CFS from other fatigue conditions. Two-day cardiopulmonary exercise testing has objectively demonstrated that ME/CFS patients show significant declines in exercise capacity on the second day [1], while healthy controls do not. See Section 2.1 for the full evidence.

Is ME/CFS psychological? No. ME/CFS is a biological disease with documented abnormalities in energy metabolism [2], immune function [3], autonomic regulation [4], and brain function [5]. The 2015 Institute of Medicine report [6] concluded that ME/CFS is a “serious, chronic, complex, systemic disease” and explicitly rejected its classification as a psychological disorder. See Chapter 1 and the Reading Guide for the evidence hierarchy used in this document.

Should I exercise? Activity management (“pacing”) is the recommended approach, not graded exercise therapy (GET). Research shows that ME/CFS patients have impaired energy production that worsens after exertion [1]. Pushing through fatigue can trigger PEM and potentially cause lasting deterioration. The goal is to stay within your “energy envelope”—the amount of activity you can sustain without triggering PEM. See Chapter 23 for pacing strategies and Section 27.5 for the evidence against GET.

Are there effective treatments? There is currently no cure for ME/CFS, and no single treatment works for everyone. However, several approaches can improve quality of life:

- **Pacing:** Staying within your energy limits is the most important management strategy (Chapter 23)
- **Symptom management:** Sleep, pain, orthostatic intolerance, and cognitive symptoms can each be addressed (Chapter 16)
- **Supplements:** CoQ10, NADH, and other mitochondrial supplements have RCT evidence of modest benefit (Chapter 22)

- **Medications:** Low-dose naltrexone, beta-blockers for POTS, and other targeted medications may help specific symptoms (Chapter ??)

Treatment should be individualized. See Chapters 17 and 18 for severity-appropriate care pathways.

Do supplements help? Some supplements have evidence from randomized controlled trials: CoQ10+NADH showed significant improvement in fatigue and quality of life in a 207-patient RCT [7]; D-ribose showed improvement in energy and well-being [8]. Other supplements (magnesium, B vitamins, vitamin D) address common deficiencies but have less ME/CFS-specific trial data. See Chapter 22 for dosing, evidence levels, and safety considerations.

What causes ME/CFS? The exact cause is unknown, but evidence points to a multi-system dysfunction involving impaired energy metabolism, immune dysregulation, and autonomic nervous system abnormalities (the “core triad”; Chapter 13). Most cases begin after an infection, and ongoing research investigates viral persistence, autoimmune mechanisms, and metabolic disruption as contributing factors. Multiple biological subgroups likely exist (Chapter 5).

Can ME/CFS get better? Recovery rates vary widely across studies, from 5% to 22% depending on definitions and populations studied. Children and adolescents generally have better prognosis than adults. Improvement (without full recovery) is more common. Early diagnosis and aggressive pacing appear to improve outcomes. See the prognosis and disease progression sections in Chapter 5.

How is ME/CFS diagnosed? There is no single diagnostic test. Diagnosis is based on clinical criteria requiring: (1) substantial reduction in activity lasting ≥ 6 months, (2) post-exertional malaise, (3) unrefreshing sleep, plus either cognitive impairment or orthostatic intolerance [6]. Other conditions must be excluded through standard medical workup. See Chapter 4 for the major diagnostic frameworks (Fukuda, Canadian Consensus, ICC, IOM).

What should I tell my doctor? Many physicians are unfamiliar with ME/CFS. Key points to communicate:

- Describe PEM specifically: what triggers it, how long it lasts, and how it differs from normal fatigue
- Request basic workup to exclude other conditions (thyroid, anemia, autoimmune, sleep disorders)
- Reference the 2015 IOM diagnostic criteria [6]
- Ask about specialist referral if your GP is unfamiliar with ME/CFS

See Appendix C for assessment instruments and Appendix D for specialist directories.

Is ME/CFS the same as Long COVID? ME/CFS and Long COVID overlap substantially: approximately 50% of Long COVID patients meet ME/CFS diagnostic criteria. Both conditions share features including PEM, cognitive dysfunction, and autonomic symptoms. The pathophysiology chapters (Part II) cover mechanisms common to both conditions. See Chapter ?? for cross-disease comparisons.

Are there any promising treatments being researched? Several approaches are in active clinical trials:

- **Daratumumab:** An anti-CD38 antibody targeting autoimmune mechanisms

showed 60% response in a pilot study [9]

- **Low-dose naltrexone:** Multiple studies underway
- **Fecal microbiota transplant:** Large RCTs in progress
- **BC007:** DNA aptamer neutralizing GPCR autoantibodies [10]

See Chapter ?? for full coverage and Chapter 27 for trial details.

△ **Warning 2: Medical Disclaimer**

This FAQ summarizes research findings for informational purposes. It is not medical advice. All treatment decisions should be made in consultation with a qualified healthcare provider familiar with ME/CFS. See the AI Disclosure and Reading Guide for important caveats about how to interpret this document.

Changelog

This changelog records which chapters and sections were substantially revised in each version of the document. Minor corrections (typos, formatting) are not listed.

Each entry follows the format: **What** changed, *why* it changed (new evidence, error correction, reviewer feedback, etc.), and which **key sources** motivated the change where applicable.

Version 5 (2026-02-18 – 2026-03-07)

Document Infrastructure

- **Revision bar system removed:** The `\rev{N}/\revend` paragraph-marking system (changebar, marginnote packages) has been fully removed and replaced with no-op stubs. All markers stripped from 24 chapter files. Improves build performance and reduces source noise; the changelog remains the sole record of changes.
- **Epistemic stratification — limitation environment:** New `limitation` and `limitation*` environments deployed across 40+ locations in Parts I–IV to explicitly mark what the data do *not* allow concluding. Adversarial review round corrected 15 findings across the initial deployment. *Motivated by:* Reviewer feedback requesting a fourth epistemic layer (“what not to conclude”) beyond `hypothesis/speculation/warning`.
- **Model-unique insight environment:** New `modelunique` `tcolorbox` environment for conclusions that emerge only from formal mathematical modeling, visually distinct from empirical observations.
- **Methodology section:** New front-matter section describing the document’s methodology, evidence assessment framework, and epistemic conventions.
- **Build system:** Added `.latexmkrc` configuration and TeX memory settings (`TEXMFCNF`) for large-document builds.
- **Biomedical review system:** Added 13 specialized auditor agents (pathway, immunology, biochemistry, neuro, microbiome, biomarker, epidemiology, comorbidity, safety, pharmacology, patient-safety, falsifiability, bibliography) and the `review-biomedical` skill for comprehensive 20-category scientific review.
- **Changelog convention:** V5 entries now include “why” (motivation) and key sources for each change, per reviewer recommendation.

Part I — Clinical Overview

- **Chapter 1 — Introduction to ME/CFS:** Expanded coverage of ME/CFS terminology, disease classification, and epidemiological context.

- **Chapter 5 — Disease Course and Subgroups:** Enhanced endotype table with reproducibility status (R/P), evidence levels (A/B/C), and biomarker accessibility tiers. Added limitation environment for provisional classification status. *Motivated by:* Reviewer request for integrated endotype table with evidence grades.
- **Patient FAQ:** New chapter in front matter with 12 questions addressing common patient concerns (What is ME/CFS?, PEM, exercise, diagnosis, treatments, supplements, Long COVID overlap, promising research). *Motivated by:* Reviewer request for accessible patient-facing content.

Part II — Pathophysiology

- **Chapter 6 — Energy Metabolism:** Major expansion covering new research on mitochondrial dysfunction, metabolic trap hypothesis, and the relationship between immune activation and energy production. Three-pass scientific rigor review of heterogeneity and energy sections. *Motivated by:* Wang et al. 2023 (WASF3–Complex IV mechanism) [2]; Keller et al. 2024 (two-day CPET) [1]; Cheema et al. 2023 (miRNA convergence) [11].
- **Chapter 7 — Immune System Dysfunction:** Integrated new findings on the gut-immune-brain axis, including the role of vagal signalling in immune dysregulation and the cholinergic anti-inflammatory pathway. Resolved 22 citation violations. *Motivated by:* Heng et al. 2025 (T cell effector memory profiling) [12]; scientific rigor audit identifying uncited claims.
- **Chapter 8 — Neurological Dysfunction:** Added the enterochromaffin-vagal pathway hypothesis (Wirth–Scheibenbogen 2025), explaining how gut dysbiosis drives autonomic dysfunction via serotonin and vagal mechanisms. *Motivated by:* Barton et al. 2023, 2025 (enterochromaffin serotonin–vagal signaling) [13, 14].
- **Chapter 10 — Cardiovascular Dysfunction:** Expanded gut-mediated vagal impairment section with Wirth–Scheibenbogen 2025 mechanistic support and new cross-references. Added replication status warnings. *Motivated by:* van Campen et al. 2024 (cardiac output–cerebral blood flow study, $n = 534$) [4].
- **Chapter 11 — Gut Microbiome:** Expanded clinical implications of microbiome alterations and their connection to systemic symptoms. Added replication status warnings.
- **Chapter 13 — Integrative Models:** Revised Heng 2025 multi-omics description; added replication warnings; expanded integrative content.
- **Chapter 14d — Cross-Disease Comparisons:** Substantially expanded cross-condition analysis with new disease parallels and shared-mechanism evidence.
- **Chapter 14j — Selective Energy Dysfunction Hypothesis:** Thorough scientific rigor review; strengthened internal consistency, corrected mathematical formulations, and expanded related-hypotheses discussion across 10 blocks.
- **Chapter 15 — Symptom-Producing Mechanisms:** Updated MCAS prevalence citations and mast cell–microglia reference keys.

Part III — Treatment

- **Chapter 14 — Symptom Management:** Major expansion with new sections on pacing and energy envelope theory, PEM medications, sleep management, pain management,

cognitive dysfunction management, and autonomic symptom management. Added limitation environments flagging sections with pending citations and evidence quality caveats.

- **Chapter 14b — Action Plans for Mild to Moderate Cases:** Integrated a new acute-onset protocol for patients in early disease stages; expanded severity-stratified care pathways.
- **Chapter 14c — Speculative model language:** Revised text presenting the time-dependent reversibility model from asserting it “explains” outcomes to stating it “hypothesizes” outcomes, with “speculative” qualifier added. *Motivated by:* A8 audit findings (Issues 28–29, HIGH PRIORITY).
- **Chapter 15 — Medications Targeting Underlying Mechanisms:** Revised with review corrections; removed unsupported claims.
- **Chapter 16 — Supplements and Nutraceuticals:** Substantially revised; removed unsupported claims identified during adversarial review.
- **Chapter 17 — Lifestyle Interventions:** Major expansion with comprehensive sleep optimization guidance (bedroom environment, schedule consistency, light exposure, temperature regulation, evening routines), expanded dietary considerations, and revised post-prandial pacing hypothesis language.
- **Chapter 18 — Experimental and Emerging Therapies:** Corrected suramin text to explicitly state the cited pilot study [15] was conducted in autism spectrum disorder, not ME/CFS. Added energy profile for immunoadsorption; corrected nicotinamide riboside trial data and citation. *Motivated by:* Assertion–source audit finding (DOES NOT SUPPORT).
- **Chapter 19 — Integrative and Personalised Approaches:** Revised with review corrections; tightened evidence claims.
- **Environment type correction (Chapter 15):** Converted observation to hypothesis for mechanistic treatment sequencing rationale (certainty: 0.25), fixing epistemic mismatch. *Motivated by:* A8 audit finding (Issue 13, HIGH PRIORITY).
- **A8 audit completed:** Full Part III hypothesis vs. recommendation audit identified 31 findings (3 critical fixed, 14 medium, 14 low priority).

Part IV — Research

- **Chapter 21 — Clinical Trials:** Major expansion with detailed coverage of rituximab trials (pilot through RituxME phase III), low-dose naltrexone evidence, and other immune modulators. New section documenting 9 key negative or null-result trials (rituximab phase III, PACE trial, IVIG, acyclovir, valganciclovir, rintatolimod, galantamine, fluoxetine, hydrocortisone) with summary table and cross-cutting lessons. *Motivated by:* Reviewer request for explicit coverage of negative evidence to combat confirmation bias.
- **Chapter 22 — Mechanistic Studies:** Substantially expanded with detailed analysis of NIH deep phenotyping study (Walitt et al. 2024), WASF3 mitochondrial findings (Wang et al. 2023), and methodological critique. Corrected attribution (Hwang → Wang).
- **Chapter 23 — Epidemiology and Outcomes:** Major expansion with updated prevalence data, risk factor analysis, and outcome measurement discussion.
- **Chapter 24 — Controversies:** Comprehensive rewrite covering nomenclature debate

(ME vs. CFS vs. SEID), diagnostic criteria controversies (Oxford through IOM), and detailed PACE trial controversy analysis (methodology criticisms, reanalysis results, impact on GET/CBT recommendations).

- **Chapter 25 — Translational Findings:** Revised shared-mechanism tables with added citations; corrected treatment dosing and trial data.
- **Chapter 25a — Research Infrastructure:** Expanded with new content on research funding gaps and infrastructure needs.
- **Chapter 25b — Proposed Research Studies:** New chapter describing fourteen original research protocols targeting key open questions in ME/CFS pathophysiology and treatment.
- **Hypothesis Registry:** Major expansion consolidating 40+ key hypotheses, speculations, predictions, and open questions from a systematic harvest of 305 environments across the document. Organized by mechanism domain with testable predictions and available tests. *Motivated by:* Reviewer request for hypothesis–prediction–test mapping.

Part V — Computational Modeling

- **Chapter 26 — Modeling Foundations:** New section surveying existing mathematical and computational models of ME/CFS (Phair’s IDO metabolic trap, Broderick’s cytokine network analysis, Morris & Maes’s neuro-immune framework, Li et al.’s genome-scale metabolic modeling, Xiong/Oh’s BioMapAI), positioning Part V within that landscape. Honest assessment of contributions and limitations (no computational implementation, parameter uncertainty, validation gap). *Motivated by:* Need for transparent novelty assessment; prior work survey absent from original Part V draft. [16, 17, 18, 19, 20].
- **Chapter 27 — Energy Metabolism Models:** Major expansion with formal ODE systems, parameter tables, and sensitivity analysis frameworks for mitochondrial and metabolic models.
- **Chapter 28 — Immune System Models:** Major expansion with formal immune cell population dynamics, B cell depletion modeling (rituximab vs. plasma cell persistence), and cytokine network models.
- **Chapter 29 — Neuroendocrine Models:** Major expansion with HPA axis feedback models, autonomic regulation frameworks, and neurotransmitter dynamics.
- **Chapter 30 — Integrated Systems:** Major expansion describing multi-system coupling architecture, cross-system feedback loops, and emergent properties of the integrated model.
- **Chapter 31 — Temporal Evolution:** Major expansion with disease trajectory modeling, phase transition analysis, and the energy ratchet mechanism formalized as a mathematical model.
- **Chapter 32 — Predictive Applications:** Major expansion covering clinical decision support, treatment response prediction, and biomarker-guided intervention timing.
- **Chapters 26–31 — Model Application Guides:** Added structured “Model Application Guide” sections to each modeling chapter (Chs. 27–31), specifying for every model group: measurements required, how to use the model, worked examples with realistic parameter values, inter-model dependencies, falsification criteria, and clinical implications. Added model coverage justification section to Ch. 26 with Part II→Part V mapping

table, selection rationale, and explicit enumeration of unmodeled domains. *Motivated by:* Gap analysis identifying that models lacked practical application guidance, falsification criteria, and coverage justification.

- **TikZ figures:** Added 10 new TikZ illustrations: baroreflex model, clinical decision pipeline, disease trajectories, energy metabolism model, energy ratchet, HPA axis model, immune cell dynamics, integrated systems architecture, modeling framework overview, and tryptophan branching model.

Appendices and Bibliography

- **Appendix A — Glossary:** Expanded from stub to comprehensive alphabetical glossary of medical, biochemical, immunological, and statistical terms used throughout the document.
- **Appendix C — Diagnostic Tools:** Substantially expanded with detailed descriptions of diagnostic instruments and assessment tools.
- **Appendix E — Mathematical Model Details:** Expanded from stub to complete mathematical specifications for Part V models, including full ODE systems, parameter tables, derivations, numerical methods, and sensitivity analysis procedures.
- **Appendix F — Supplement Guide:** Substantially expanded with updated supplement protocols and dosage information.
- **Appendix H — Annotated Bibliography:** Major expansion providing comprehensive annotated bibliography of ME/CFS literature organized by research domain.
- **Appendix I — Research Registry:** New appendix cataloguing ongoing and planned ME/CFS studies.
- **Bibliography:** Added approximately 200 new references covering recent ME/CFS research across pathophysiology, treatment, and modeling.

Version 4 (2026-02-18)

Fourth publicly released version of the document. Established the full structure across five parts: clinical overview, pathophysiology, treatment, research, and mathematical modelling.

Contents

Abstract	i
Version Notice	iii
Keywords	iv
License and Usage	v
About the Author	vii
AI Disclosure Statement	viii
Methodology and Quality Assurance	xi
Reading Guide: How to Use This Document	xviii
Frequently Asked Questions	xxi
Changelog	xxiv
List of Figures	lxxviii
List of Tables	lxxxix
I Clinical Overview	1
1 Introduction to ME/CFS	3
1.1 Overview and Terminology	3
1.2 Historical Context	3
1.2.1 Key Outbreaks and Case Clusters	3
1.2.2 Evolution of Diagnostic Criteria	4
1.2.3 Changes in Medical Understanding	5
1.3 Disease Classification: From Syndrome to Disease	7
1.3.1 The Syndrome vs. Disease Distinction	7
1.3.2 The 2024 NIH Deep Phenotyping Study	7
1.3.3 Official Reclassification	7
1.3.4 Why the Name Persists	8
1.3.5 Toward a More Accurate Name: Dysergopoiesis	8
1.3.6 Implications for Patient Care	10
1.3.7 The “Effort Preference” Controversy	11

Contents

1.4	Epidemiology	11
1.4.1	Prevalence and Incidence	11
1.4.2	Demographic Patterns	12
1.4.3	Geographic Distribution	13
1.4.4	Risk Factors	13
1.5	Disease Impact	13
1.5.1	Quality of Life	13
1.5.2	Disability and Functional Capacity	14
1.5.3	Economic Burden	15
1.5.4	Psychosocial Impact	15
1.6	Prognosis and Disease Course	16
1.6.1	Onset Patterns	16
1.6.2	Disease Trajectory	16
1.6.3	Severity Fluctuation	17
2	Core Symptoms	18
2.1	Post-Exertional Malaise (PEM)	18
2.1.1	Definition and Characteristics	18
2.1.2	Physiological Basis	24
2.1.3	Measurement and Assessment	50
2.2	Unrefreshing Sleep	51
2.2.1	Sleep Dysfunction Patterns	51
2.2.2	Polysomnography Findings	52
2.2.3	Related Sleep Disorders	53
2.2.4	Differential Diagnosis Approach	54
2.3	Cognitive Impairment	55
2.3.1	Domains of Cognitive Dysfunction	55
2.3.2	Neuropsychological Testing	56
2.3.3	Neuroimaging Findings	57
2.4	Autonomic Dysfunction	58
2.4.1	Orthostatic Intolerance	58
2.4.2	Other Autonomic Symptoms	60
2.4.3	Autonomic Testing Battery	62
2.4.4	Clinical Implications	62
2.5	Pain	63
2.5.1	Types of Pain in ME/CFS	63
2.5.2	Pain Mechanisms	64
2.5.3	Pain Assessment and Management Considerations	65
2.6	Sensory Sensitivities	66
2.6.1	Types of Sensory Sensitivity	66
2.6.2	Mechanisms of Sensory Sensitivity	67
2.6.3	Clinical Implications	68
3	Additional Symptoms and Manifestations	69
3.1	Neurological Symptoms	69
3.1.1	Sensory Sensitivities	69
3.1.2	Motor and Coordination Symptoms	72

Contents

3.1.3	Paresthesias and Sensory Disturbances	73
3.1.4	Dizziness and Vertigo	74
3.1.5	Tinnitus	74
3.1.6	Seizure-Like Episodes	74
3.2	Immunological and Inflammatory Symptoms	75
3.2.1	Flu-Like Symptoms	75
3.2.2	Infection Susceptibility and Viral Reactivation	75
3.2.3	Allergies and Mast Cell Activation	76
3.3	Musculoskeletal Symptoms	76
3.3.1	Muscle Cramps and Contractures	76
3.3.2	Myalgia (Muscle Pain)	77
3.3.3	Arthralgia (Joint Pain)	77
3.3.4	Fibromyalgia Overlap	78
3.4	Gastrointestinal Symptoms	78
3.4.1	Nausea	78
3.4.2	Irritable Bowel Syndrome (IBS)	78
3.4.3	Food Intolerances and Sensitivities	79
3.4.4	Gastroparesis and Delayed Gastric Emptying	79
3.4.5	Gastroesophageal Reflux (GERD)	80
3.5	Cardiovascular Symptoms	80
3.5.1	Palpitations	80
3.5.2	Chest Pain	80
3.5.3	Blood Pressure Abnormalities	81
3.5.4	Raynaud’s Phenomenon	81
3.6	Respiratory Symptoms	81
3.6.1	Dyspnea and Air Hunger	81
3.6.2	Dysfunctional Breathing Patterns	82
3.6.3	Chronic Cough	82
3.7	Genitourinary Symptoms	82
3.7.1	Urinary Dysfunction	82
3.7.2	Sexual Dysfunction	83
3.7.3	Menstrual Irregularities	83
3.8	Endocrine and Metabolic Symptoms	83
3.8.1	Temperature Dysregulation	83
3.8.2	Excessive Thirst and Fluid Retention	84
3.8.3	Weight Changes	84
3.8.4	Glucose Metabolism Abnormalities	84
3.9	Dermatological Symptoms	85
3.9.1	Rashes and Skin Manifestations	85
3.9.2	Hair and Nail Changes	85
3.10	Ocular Symptoms	85
3.10.1	Vision Changes	85
3.11	Auditory Symptoms	86
3.11.1	Hearing Loss	86
3.12	Psychological and Cognitive-Emotional Symptoms	86
3.12.1	Anxiety	86
3.12.2	Depression	87

3.12.3	Emotional Lability and Mood Dysregulation	87
3.12.4	Social Withdrawal and Isolation	88
3.13	Sleep-Related Symptoms	88
3.13.1	Insomnia	88
3.13.2	Hypersomnia	88
3.13.3	Sleep Architecture Abnormalities	89
3.13.4	Restless Legs Syndrome and Periodic Limb Movements	89
3.14	Symptom Severity Spectrum	89
3.14.1	Mild ME/CFS	90
3.14.2	Moderate ME/CFS	90
3.14.3	Severe ME/CFS	90
3.14.4	Very Severe ME/CFS	91
3.15	Summary: The Multi-System Nature of ME/CFS	92
4	Diagnostic Criteria and Clinical Assessment	93
4.1	Overview of Diagnostic Approaches	93
4.1.1	Evolution of Diagnostic Criteria	93
4.1.2	Comparison of Frameworks: Set-Theoretic Analysis	94
4.1.3	Sensitivity and Specificity Trade-offs	95
4.1.4	Clinical Implications of Framework Choice	95
4.2	Canadian Consensus Criteria (2003)	96
4.2.1	Required Criteria	96
4.2.2	Exclusions	97
4.2.3	Strengths and Limitations	97
4.3	International Consensus Criteria (2011)	98
4.3.1	Required Criteria	98
4.3.2	Phenotype Categories	99
4.3.3	Duration and Exclusions	100
4.3.4	Strengths and Limitations	100
4.3.5	Research and Clinical Application	101
4.4	Institute of Medicine Criteria (2015)	101
4.4.1	Required Core Symptoms	101
4.4.2	Additional Required Symptoms	102
4.4.3	Diagnostic Algorithm Structure	103
4.4.4	Exclusions and Comorbidities	103
4.4.5	Strengths and Limitations	103
4.4.6	Clinical and Research Application	104
4.5	Other Diagnostic Frameworks	104
4.5.1	Fukuda Criteria (1994)	104
4.5.2	Oxford Criteria (1991)	106
4.5.3	Pediatric Criteria	107
4.6	Clinical Assessment	109
4.6.1	History Taking	109
4.6.2	Physical Examination	111
4.6.3	Laboratory Testing	113
4.7	Novel Biology-Informed Diagnostic Framework	115
4.7.1	Rationale for a New Framework	115

4.7.2	Tier 1: Clinical Syndrome Criteria	119
4.7.3	Tier 2: Biological Phenotyping (Multi-Domain Assessment)	121
4.7.4	Tier 3: Severity Classification and Progression Risk	128
4.7.5	Implementation and Clinical Workflow	131
4.7.6	Research Implications and Validation Needs	134
4.7.7	Comparison to Existing Criteria	135
4.7.8	Clinical Advantages of the Novel Framework	135
4.8	Differential Diagnosis	137
4.8.1	Conditions That Can Mimic ME/CFS	137
4.8.2	Comorbid Conditions vs. Alternative Diagnoses	142
4.8.3	When Comorbidities May Be Primary Drivers	142
4.8.4	Diagnostic Algorithm	144
4.8.5	When to Reconsider the Diagnosis	145
5	Disease Course and Prognosis	146
5.1	Onset Patterns	146
5.2	Disease Severity Levels	151
5.3	The Devastating Reality of Severe ME/CFS	158
5.3.1	The Scale of Catastrophe	158
5.3.2	Complete Energy Bankruptcy	159
5.3.3	Existence in Darkness and Silence	161
5.3.4	The Prison of the Body	163
5.3.5	Cognitive Devastation	165
5.3.6	The Wish for Death	167
5.3.7	Impact on Caregivers and Families	168
5.3.8	Economic Devastation	170
5.3.9	Medical Abandonment	171
5.3.10	A Call to Action	172
5.4	Disease Progression	174
5.4.1	Natural History	174
5.4.2	Patterns of Change	176
5.4.3	The Preventable Descent to Severe Disease: Critical Warning	177
5.4.4	Relapse and Remission	187
5.4.5	Factors Influencing Trajectory	188
5.5	Prognosis	190
5.5.1	Recovery Rates	190
5.5.2	Evidence That Adult Recovery, While Rare, Does Occur	192
5.5.3	Prognostic Factors	198
5.5.4	Long-Term Disability	199
5.5.5	Mortality	200
5.5.6	Implications for Patients and Clinicians	201
5.6	Subgroups and Phenotypes	202
5.6.1	The Heterogeneity Problem	202
5.6.2	Onset-Based Subgroups	203
5.6.3	Severity-Based Subgroups	203
5.6.4	Metabolic Phenotypes	204
5.6.5	Immune Phenotypes	205

5.6.6	Symptom-Based Subgroups	205
5.6.7	Criteria-Based Phenotypes	206
5.6.8	Clinical Significance of Subgrouping	207
5.6.9	From Subgroups to Stratified Medicine	208
5.6.10	From “Not Clinically Actionable” to “Increasingly Necessary”	211
5.6.11	Comorbidity Clustering: The “Septad” Framework	213
5.6.12	Emerging Treatment-Response Phenotypes	221
5.7	Trigger Mechanisms: From Acute Infection to Chronic Disease	223
5.7.1	The Core Problem: Failure to Resolve	223
5.7.2	Epstein-Barr Virus (EBV)	223
5.7.3	SARS-CoV-2 (COVID-19)	224
5.7.4	Human Herpesvirus 6 (HHV-6)	224
5.7.5	Enteroviruses	224
5.7.6	The Common Pathway: Vicious Cycles and Chronicity	225

II Pathophysiology and Biological Mechanisms 226

6 Energy Metabolism and Mitochondrial Function 228

6.1	Cellular Energy Production Overview	228
6.1.1	ATP Synthesis	228
6.1.2	Normal Energy Metabolism	229
6.2	Mitochondrial Dysfunction in ME/CFS	232
6.2.1	Evidence for Mitochondrial Impairment	232
6.2.2	Mechanisms of Mitochondrial Damage	234
6.2.3	Consequences of Energy Deficits	238
6.3	Selective Energy Dysfunction: The CNS-Dependency Hypothesis	247
6.3.1	Evidence for Selectivity	248
6.3.2	The Demand-Response Failure Pattern	249
6.3.3	Mechanistic Implications	250
6.3.4	Therapeutic Implications of Selective Dysfunction	250
6.3.5	Subtype Considerations	251
6.4	Oxidative and Nitrosative Stress	251
6.4.1	Reactive Oxygen Species (ROS)	251
6.4.2	Reactive Nitrogen Species	252
6.4.3	Lipid Peroxidation	253
6.5	Metabolic Pathways Affected	254
6.5.1	Amino Acid Metabolism	254
6.5.2	Lipid Metabolism	255
6.5.3	Carbohydrate Metabolism	256
6.6	Catecholamine Metabolism: NIH Study Findings	257
6.6.1	CSF Catecholamine Findings	257
6.6.2	Catecholamine Synthesis Pathway	258
6.6.3	Potential Mechanisms of Catecholamine Deficiency	258
6.6.4	Functional Consequences	259
6.7	The “Metabolic Trap” Hypothesis	259
6.7.1	IDO Metabolic Trap	260

6.7.2	The “Dauer” Hypothesis	260
6.8	Compartmental Energy Models	260
6.8.1	CNS-Specific vs. Global Dysfunction	260
6.8.2	Evidence for Compartmentalization	261
6.8.3	The Astrocyte-Neuron Lactate Shuttle	261
6.9	Potential Interventions	262
6.9.1	Mitochondrial Support	262
6.9.2	Antioxidants	262
6.9.3	Addressing Catecholamine Deficiency	263
6.10	Microcirculation and Oxygen Delivery	263
6.10.1	Endothelial Dysfunction	263
6.10.2	Red Blood Cell Deformability and Rheology	264
6.10.3	Capillary Structural Pathology	264
6.10.4	Brain Microvascular Insufficiency	264
6.10.5	Integration: Substrate Delivery Failure	264
6.11	Energy Utilization Beyond ATP Production	265
6.11.1	The Sodium-Calcium Overload Cascade	265
6.11.2	TRPM3 Ion Channel Dysfunction	265
6.11.3	WASF3 and ER Stress-Induced Metabolic Switching	266
6.11.4	Distinction: Production vs. Utilization Defects	266
6.12	Immune-Metabolic Crosstalk Maintaining the Energy Crisis	266
6.12.1	Tryptophan-Kynurenine Metabolism and NAD ⁺ Depletion	266
6.12.2	T-Cell Exhaustion with Metabolic Reprogramming	267
6.12.3	Exaggerated Innate Immune Response with Metabolic Consequences	267
6.12.4	Oxidative Stress in Immune Cells	267
6.12.5	Multi-Omics Integration: The Full Picture	267
6.13	The Multi-Level Vicious Cycle	268
6.13.1	The Cycle in Detail	268
6.13.2	Multi-Level Integration	269
6.13.3	Evidence for the Integrated Cycle	269
6.14	Summary: Integrated Metabolic Model	270
6.14.1	Normal ATP Synthesis Pathways	270
6.14.2	Documented Energy Dysfunctions	271
6.14.3	Hypothesized Energy Mechanisms	272
7	Immune System Dysfunction	274
7.1	Innate Immunity	274
7.1.1	Natural Killer (NK) Cell Dysfunction	274
7.1.2	Neutrophil and Monocyte Function	277
7.1.3	Complement System	279
7.1.4	Dendritic Cells	279
7.2	Adaptive Immunity	279
7.2.1	T Cell Abnormalities	280
7.2.2	B Cell Function and Antibodies	287
7.3	Cytokines and Inflammatory Mediators	293
7.3.1	Pro-inflammatory Cytokines	293
7.3.2	Anti-inflammatory Cytokines	298

Contents

7.3.3	Chemokines	299
7.4	Immune Activation and Inflammation	299
7.4.1	Chronic Immune Activation	299
7.4.2	Neuroinflammation	300
7.5	Viral Reactivation and Persistence	301
7.5.1	Herpesviruses	301
7.5.2	Other Implicated Viruses	305
7.5.3	Tick-Borne Infections	306
7.5.4	Infection-Induced Cumulative Damage and Disease Progression	310
7.6	Autoimmunity in ME/CFS	313
7.6.1	Autoantibodies Identified	313
7.6.2	Autoimmune Mechanisms	314
7.6.3	Developmental Immune Tolerance and Recovery	315
7.7	Connections to Allergies and Mast Cell Activation	322
7.7.1	Mast Cell Activation Syndrome (MCAS)	322
7.7.2	Allergic Responses	329
7.8	Summary: Integrated Model of Immune Dysfunction	330
7.9	Emerging Research Directions in Immune Dysregulation	331
7.9.1	Tier 1: Immediate Translation Potential (Existing Drugs, Severe Case Priority)	331
7.9.2	Tier 2: Near-Term Clinical Trials (Moderate Complexity, High Impact)	335
7.9.3	Tier 3: Long-Term Mechanistic Research (Foundational Understanding)	339
7.9.4	Prioritization Summary: Research Directions by Impact and Timeline	342
7.9.5	Post-Infectious Pathogenesis	343
7.9.6	Trained Immunity and Epigenetic Lock-in	343
7.9.7	Zonulin-Mediated Gut Permeability	344
8	Neurological and Neurocognitive Dysfunction	346
8.1	Central Nervous System Abnormalities	346
8.1.1	Brain Structure and Function	346
8.1.2	Brain as Energy Coordination Bottleneck	349
8.1.3	Neurotransmitter Abnormalities	352
8.1.4	Sleep Architecture and Inter-Regional Coordination	360
8.1.5	Glial Cell Dysfunction	362
8.1.6	Integrated Neuroinflammatory Cascade Model	373
8.1.7	Post-Exertional Malaise and the Kindling Hypothesis	376
8.2	Autonomic Nervous System Dysfunction	379
8.2.1	Sympathetic vs. Parasympathetic Imbalance	379
8.2.2	Mechanisms of Orthostatic Intolerance	381
8.3	Peripheral Nervous System	382
8.3.1	Small Fiber Neuropathy	382
8.3.2	Nerve Conduction Studies	383
8.3.3	Treatment of Small Fiber Neuropathy	384
8.4	Blood-Brain Barrier Dysfunction	387
8.4.1	Evidence for Permeability Changes	388
8.4.2	Consequences for Neuroinflammation	388
8.4.3	Transport Dysfunction	388

8.4.4	Blood-Brain Barrier as CNS Vulnerability Factor	388
8.5	Cerebral Blood Flow Abnormalities	390
8.5.1	Reduced Regional Blood Flow	390
8.5.2	Correlation with Cognitive Symptoms	393
8.5.3	Mechanisms of Cerebral Hypoperfusion	393
8.5.4	Exacerbation with Exertion	393
8.6	Auditory Processing Dysfunction and Tinnitus	394
8.6.1	Prevalence and Epidemiology	394
8.6.2	Functional Auditory Processing Deficits	394
8.6.3	Neuroanatomical Substrate: Brainstem Dysfunction	395
8.6.4	Central vs. Peripheral Auditory Pathology	396
8.6.5	Proposed Mechanisms	396
8.6.6	Clinical Implications	397
8.6.7	Research Gaps	398
8.7	Cognitive Dysfunction: Clinical Manifestations	398
8.7.1	Domains of Impairment	399
8.7.2	Social and Emotional Dysfunction	401
8.7.3	Fluctuation and Post-Exertional Cognitive Malaise	410
8.7.4	CNS Energy Crisis as Primary Event	410
8.7.5	CNS Energy Triage: A Hierarchical Model of Brain Fog	411
8.8	Summary: An Integrated Neurological Model	413
9	Endocrine and Metabolic Dysfunction	415
9.1	Hypothalamic-Pituitary-Adrenal (HPA) Axis	415
9.1.1	HPA Axis Abnormalities	416
9.1.2	Mechanisms of HPA Dysfunction	420
9.1.3	Clinical Consequences	422
9.2	Thyroid Function	424
9.2.1	The Low T3 Syndrome in ME/CFS	425
9.2.2	Mechanisms of Impaired T4 to T3 Conversion	426
9.2.3	Tissue-Level Thyroid Hormone Resistance	426
9.2.4	Clinical Implications and Diagnostic Challenges	427
9.3	Sex Hormones and Gender Differences	427
9.3.1	Epidemiology of Sex Differences	428
9.3.2	Sex-Specific Pathophysiology: NIH Deep Phenotyping Findings	428
9.3.3	Steroid Hormone Abnormalities	429
9.3.4	Reproductive Health and Gynecological Risk Factors	429
9.3.5	Testosterone and Androgens	431
9.3.6	Mechanisms of Sex Hormone Influence on ME/CFS	431
9.4	Growth Hormone and IGF-1	433
9.4.1	Evidence for GH/IGF-1 Axis Dysfunction	433
9.4.2	Growth Hormone Treatment Trial	433
9.4.3	Mechanisms and Clinical Implications	434
9.5	Insulin and Glucose Metabolism	435
9.5.1	Metabolic Syndrome and Insulin Resistance	435
9.5.2	Metabolic Phenotypes and Insulin Dynamics	435
9.5.3	Cerebral Glucose Hypometabolism	436

9.5.4	Hypoglycemia Symptoms Versus Orthostatic Intolerance	437
9.5.5	Integration With Energy Metabolism Dysfunction	437
9.6	Melatonin and Circadian Rhythms	438
9.6.1	Objective Documentation of Circadian Disruption	438
9.6.2	Sleep Architecture Versus Circadian Timing	438
9.6.3	Molecular Clock Dysfunction	439
9.6.4	Melatonin and Its Therapeutic Potential	439
9.6.5	Circadian Disruption as an Integrative Mechanism	440
9.7	Integrated Endocrine-Metabolic Model	441
9.7.1	Neuroendocrine-Immune Integration	441
9.7.2	Endocrine-Metabolic Connections	442
9.7.3	Circadian Disruption as Central Organizing Principle	442
9.7.4	Sex as a Critical Biological Variable	443
9.7.5	Clinical Implications and Therapeutic Considerations	443
9.7.6	Future Research Directions	444
10	Cardiovascular Dysfunction	445
10.1	Cardiac Function	445
10.1.1	Exercise Testing Abnormalities	445
10.1.2	Cardiac Output and Stroke Volume	449
10.1.3	Cardiac Biomarkers	450
10.2	Vascular Dysfunction	451
10.2.1	Endothelial Dysfunction	451
10.2.2	Blood Volume Abnormalities	453
10.2.3	Arterial Stiffness and Vascular Compliance	454
10.2.4	Microcirculation	455
10.3	Blood Pressure Regulation	458
10.3.1	Orthostatic Hypotension	458
10.3.2	Neurally Mediated Hypotension	458
10.3.3	Postural Orthostatic Tachycardia Syndrome (POTS)	458
10.3.4	Hypertension in ME/CFS	463
10.4	Heart Rate Abnormalities	463
10.4.1	Resting Tachycardia	463
10.4.2	Heart Rate Variability	463
10.4.3	Heart Rate Recovery	463
10.5	Coagulation and Rheological Abnormalities	464
10.5.1	Hypercoagulability	464
10.5.2	Fibrin Deposition	464
10.5.3	Red Blood Cell Deformability	464
10.6	Cardiovascular Dysfunction in Post-COVID ME/CFS	465
10.6.1	Parallel Cardiovascular Findings	465
10.6.2	GPCR Autoantibodies in Post-COVID	465
10.6.3	Implications for Understanding ME/CFS	466
10.7	Sex Differences in Cardiovascular Manifestations	466
10.8	Mast Cell Activation and Cardiovascular Dysfunction	466
10.8.1	Cardiovascular Mediators of Mast Cell Activation	467
10.8.2	The MCAS-POTS Connection	467

10.8.3	Therapeutic Implications	467
10.9	Summary: Integrated Cardiovascular Model	468
11	Gastrointestinal and Microbiome Dysfunction	473
11.1	Gut Microbiome Alterations	473
11.1.1	Dysbiosis Patterns	473
11.1.2	Functional Capacity Changes	475
11.1.3	Multi-Omic Integration: Beyond Taxonomic Profiling	475
11.1.4	Gut-Brain Axis	477
11.1.5	Intestinal Permeability	478
11.1.6	Intestinal Barrier Dysfunction and Secondary Metabolic Consequences	482
11.2	Gastrointestinal Dysfunction	486
11.2.1	Irritable Bowel Syndrome Overlap	486
11.2.2	Motility Disorders	486
11.2.3	Digestive Function	489
11.3	Metabolites and Short-Chain Fatty Acids	489
11.3.1	Butyrate	489
11.3.2	Acetate and Propionate	490
11.4	Treatment Approaches	490
11.4.1	SIBO Treatment	490
11.4.2	Prokinetics	492
11.4.3	Dietary Interventions	493
11.4.4	Probiotics and Prebiotics	494
11.4.5	Fecal Microbiota Transplantation	495
11.5	Integration with Other ME/CFS Pathophysiology	495
11.5.1	Gut-Brain Metabolic Coupling	496
12	Genetic and Epigenetic Factors	499
12.1	Genetic Predisposition	499
12.1.1	Heritability Evidence	499
12.1.2	Genetic Variants and Candidate Gene Studies	502
12.1.3	Genome-Wide Association Studies	505
12.2	Epigenetic Modifications	509
12.2.1	DNA Methylation	509
12.2.2	Histone Modifications	512
12.2.3	MicroRNAs	513
12.3	Gene Expression Patterns	516
12.3.1	Transcriptomics Studies and Methodological Considerations	516
12.3.2	Differentially Expressed Genes	516
12.3.3	Pathway Enrichment and Systems Biology Analysis	518
12.3.4	Cell Type-Specific Expression and Single-Cell Approaches	519
12.3.5	Exercise-Induced Gene Expression Changes	520
12.3.6	Integration with Genetic and Epigenetic Data	520
12.4	Synthesis and Open Questions	521
12.4.1	Key Established Findings	521
12.4.2	Integration with Broader ME/CFS Pathophysiology	522
12.4.3	Unresolved Questions and Future Directions	523

13	Integrative Models and Related Phenomena	525
13.1	Evidence Classification Framework	525
13.2	Comprehensive Hypothesis Ranking	527
13.2.1	Interpretation Notes	535
13.3	Synthesis: What the Evidence Suggests	535
13.3.1	The Core Triad: Energy-Immune-Autonomic	535
13.3.2	The Autoimmune Subgroup	536
13.3.3	The Vascular Dimension	536
13.3.4	The Central Nervous System	537
13.3.5	The “Stuck” State	537
13.4	Proposed Unifying Mechanisms	541
13.4.1	Vicious Cycle Models	541
13.4.2	Multisystem Failure Cascade	548
13.4.3	Temporal Dynamics of Cycle Recruitment	549
13.4.4	Orthostatic Intolerance as Potential Upstream Driver	551
13.4.5	Selective Energy Dysfunction Framework	553
13.4.6	Circadian Energy Distribution Failure	555
13.4.7	Disease Subtype Progression	557
13.5	Hypothesis-Specific Treatment Implications	563
13.6	Relationships to Other Conditions	565
13.6.1	Fibromyalgia	565
13.6.2	Postural Orthostatic Tachycardia Syndrome (POTS)	566
13.6.3	Mast Cell Activation Syndrome	567
13.6.4	Autoimmune Conditions	567
13.6.5	Ehlers-Danlos Syndrome	568
13.6.6	Long COVID (Post-Acute Sequelae of SARS-CoV-2)	569
13.6.7	Multiple Chemical Sensitivity	572
13.6.8	Allergic and Atopic Conditions	572
13.7	Systems Biology Approaches	572
13.7.1	Multi-Omics Integration	573
13.7.2	Network Analysis	573
13.7.3	Computational Modeling	574
13.7.4	Challenges and Limitations	578
13.8	Outstanding Questions	579
13.8.1	What Triggers ME/CFS Onset?	579
13.8.2	Why Do Some Patients Recover While Others Don’t?	580
13.8.3	What Maintains the Disease State?	583
13.8.4	How Do Different Subtypes Differ Mechanistically?	583
13.8.5	Can We Identify Critical Intervention Points?	583
13.8.6	Research Priorities	584
14	Speculative Mechanistic Hypotheses	585
14.1	Master Hypothesis Table: Likelihood and Therapeutic Potential	585
14.1.1	How to Use This Table	588
14.1.2	Qualification and Caveats	589
14.2	Metabolic “Safe Mode” Hypothesis	590
14.2.1	Mechanistic Details	590

Contents

14.2.2	Why the Program Might Lock	590
14.2.3	Testable Predictions	591
14.3	Glymphatic/CSF Clearance Failure	591
14.3.1	The Glymphatic System	592
14.3.2	Potential Disruption Mechanisms	592
14.3.3	Connections to ME/CFS Features	593
14.3.4	Testable Predictions	593
14.4	Endogenous Retrovirus Reactivation	594
14.4.1	Biology of Human Endogenous Retroviruses	594
14.4.2	The HERV-ME/CFS Connection	594
14.4.3	Supporting Observations	595
14.4.4	Testable Predictions	595
14.5	Lactate Compartmentalization Disorder	596
14.5.1	Lactate Physiology	596
14.5.2	Compartmentalization Pathophysiology	596
14.5.3	Why MCT Function Might Be Impaired	597
14.5.4	Testable Predictions	597
14.6	Vagal Afferent “Danger Signal” Loop	597
14.6.1	Vagal Afferent Pathways	598
14.6.2	Mechanisms of Aberrant Signaling	598
14.6.3	The Gut-Brain Axis Connection	599
14.6.4	Testable Predictions	599
14.7	Tryptophan/Kynurenine Trap	599
14.7.1	Tryptophan Metabolism Overview	600
14.7.2	The Trap Mechanism	600
14.7.3	Consequences of the Trap	601
14.7.4	Testable Predictions	601
14.8	Cellular “Quorum Sensing” Dysfunction	601
14.8.1	Intercellular Communication Systems	602
14.8.2	Corruption Mechanisms	603
14.8.3	Why Standard Tests Miss This	603
14.8.4	Testable Predictions	603
14.9	Purinergic Signaling Dysregulation	604
14.9.1	Purinergic Signaling Biology	604
14.9.2	Dysregulation Mechanisms	605
14.9.3	Consequences of Purinergic Dysregulation	605
14.9.4	Testable Predictions	606
14.10	Redox Compartment Collapse	606
14.10.1	Cellular Redox Compartments	606
14.10.2	Boundary Maintenance	607
14.10.3	Consequences of Boundary Collapse	607
14.10.4	Testable Predictions	608
14.11	Metabolic Memory and Epigenetic Lock	608
14.11.1	Epigenetic Mechanisms	609
14.11.2	Metabolic Memory in Disease	609
14.11.3	Application to ME/CFS	610
14.11.4	Why Duration Matters	610

Contents

14.11.5	Testable Predictions	610
14.12	Circadian-Metabolic Desynchronization	610
14.12.1	The Circadian System	611
14.12.2	Consequences of Desynchronization	611
14.12.3	How Desynchronization Might Occur	612
14.12.4	Testable Predictions	612
14.13	GPCR Autoantibody-Driven Dysfunction	613
14.13.1	Established Evidence	613
14.13.2	Speculative Hypotheses Emerging from GPCR Research	615
14.13.3	Undocumented Biological Phenomena	616
14.13.4	Evidence Assessment Summary	617
14.14	Ion Channel Autoimmunity	617
14.14.1	Ion Channels in Physiology	618
14.14.2	Ion Channel Autoimmunity Precedents	619
14.14.3	Potential ME/CFS Relevance	619
14.14.4	Testable Predictions	619
14.15	Ferroptosis Susceptibility	620
14.15.1	Ferroptosis Biology	620
14.15.2	ME/CFS Risk Factors for Ferroptosis	620
14.15.3	Sublethal Ferroptosis	621
14.15.4	Testable Predictions	621
14.16	Integrated Hypothesis: The Multi-Lock Trap	622
14.16.1	Phase 1: Triggering Event	622
14.16.2	Phase 2: Lock Establishment	622
14.16.3	Phase 3: Trap Maintenance	623
14.16.4	Why Recovery Is Rare	623
14.16.5	Testable Predictions	624
14.16.6	Therapeutic Implications	624
14.16.7	Research Directions	625
14.17	Speculative Cross-Disease Connections	625
14.17.1	The Post-Infectious Syndrome Cluster	625
14.17.2	The Dysautonomia Spectrum	627
14.17.3	The Mast Cell Connection	628
14.17.4	The Ehlers-Danlos Connection	632
14.17.5	The Fibromyalgia Overlap	638
14.17.6	The Autoimmune Disease Spectrum	639
14.17.7	The Mitochondrial Disease Connection	641
14.17.8	The Psychiatric Overlap—Reframed	642
14.17.9	The Cancer Cachexia Connection	643
14.17.10	The Hibernation/Torpor Analogy	643
14.17.11	Symptom-Specific Speculations	644
14.18	Emerging Hypotheses from 2025 Research	645
14.18.1	The Vascular-Immune-Energy Triad	645
14.18.2	The Plasma Cell Sanctuary Hypothesis	646
14.18.3	The Endothelial Activation Cascade	647
14.18.4	The Dendritic Cell Maturation Block	648
14.18.5	The NAD ⁺ Depletion Spiral	649

Contents

14.18.6	The Effort-Preference Recalibration	650
14.18.7	The Immune Cell Energy Crisis	651
14.18.8	The Vascular “Memory” Hypothesis	652
14.18.9	Speculative Treatment Approaches from 2025 Findings	653
14.19	Novel Hypotheses from Two-Day CPET Findings	654
14.19.1	The Autonomic-Mitochondrial Feedback Loop	654
14.19.2	Mitochondrial Turnover Rate Limitation	656
14.19.3	Pre-Conditioning Hypothesis (Highly Speculative)	657
14.19.4	Circadian Optimization of Recovery	658
14.19.5	Exercise Metabolomics-Guided Personalization	659
14.19.6	Vagal Stimulation for Recovery Acceleration	659
14.19.7	Blood Flow Redistribution Training	660
14.19.8	Summary Table: Novel Hypotheses from CPET Findings	661
14.20	Novel Hypotheses from 2026 Autoimmune Research	663
14.20.1	The EBV-B Cell CNS Infiltration Hypothesis	663
14.20.2	The GPCR Autoantibody-Monocyte Amplification Loop	664
14.20.3	The Receptor Internalization Hypothesis	665
14.20.4	The Antigenic Hotspot Vulnerability Hypothesis	666
14.20.5	The Molecular Mimicry-Receptor Homology Hypothesis	667
14.20.6	The Dual-Compartment Autoantibody Hypothesis	668
14.20.7	The Autoantibody Functional Assay Discrepancy Hypothesis	669
14.20.8	Updated Master Hypothesis Table: 2026 Autoimmune Hypotheses	670
14.20.9	Integration: A Unified EBV-Autoimmune Model	672
14.21	Novel Hypotheses from TRPM3 Ion Channel Research	673
14.21.1	The Paradoxical Immune State Hypothesis	674
14.21.2	The TRPM3-GPCR Signaling Convergence Hypothesis	675
14.21.3	The Systemic Channelopathy Hypothesis	675
14.21.4	The “Wired but Tired” Ion Channel Explanation	677
14.21.5	The Calcium-Mitochondria Cascade Hypothesis	677
14.21.6	The Post-Infectious TRPM3 Acquisition Hypothesis	678
14.21.7	The Temperature Dysregulation Connection	679
14.21.8	TRPM3-Targeted Therapeutic Speculation	680
14.21.9	Subtyping Implications	681
14.21.10	Updated Testable Predictions from TRPM3 Research	682
14.22	Clinical Observation-Derived Hypotheses	682
14.22.1	The “Metabolic Runway” Theory of PEM	683
14.22.2	The Mast Cell “Memory” Hypothesis	683
14.22.3	The Vagus Nerve as “Master Regulator”	684
14.22.4	The “Two Fuel Tanks” Hypothesis	685
14.22.5	The “Protective Downregulation” Paradox	686
14.22.6	The “Circadian Core” Hypothesis	687
14.22.7	The “Microclot” Bridge Hypothesis	688
14.22.8	The “Infection Doesn’t Matter” Hypothesis	690
14.22.9	Female Predominance: Hormonal Amplification	690
14.22.10	The “Bistable Equilibrium” and “Reset” Concept	691
14.22.11	Drug Candidates for Systematic Investigation	693
14.22.12	The “Kitchen Sink” Protocol Concept	694

14.23	Mechanistic Convergence: Cross-Treatment Integration	695
14.23.1	Convergence Clusters	695
14.23.2	Novel Combination Hypotheses	695
14.23.3	Phenotype-Matched Selection	697
14.24	Selective Energy Dysfunction Hypothesis	697
14.24.1	Motivation and Clinical Observations	698
14.24.2	Formal Framework	699
14.24.3	Evidence Taxonomy	701
14.24.4	Causal Structure	702
14.24.5	Mechanistic Sub-Hypotheses	702
14.24.6	Testable Predictions	708
14.24.7	Subtype Classification Model	710
14.24.8	Treatment Implications	712
14.24.9	Compartmental Energy Model	713
14.24.10	Related Hypotheses Extending the Framework	715
14.24.11	Integration with Existing Hypotheses	724
14.24.12	Limitations and Uncertainties	725
14.24.13	Summary	725
14.25	Conclusion	726
15	Symptom-Producing Mechanisms in ME/CFS	727
15.1	Conceptual Framework: Symptom-Producing versus Capacity-Limiting Mechanisms	727
15.2	Sickness Behavior as Overarching Integrative Framework	728
15.2.1	Cytokine-to-Brain Signaling Routes	728
15.2.2	Hypothalamic Integration and the Sickness Behavior Program	729
15.2.3	Persistent Sickness Behavior in ME/CFS	729
15.3	Adenosine Accumulation and Pathological Sleep Pressure	729
15.3.1	Adenosine as Metabolic Waste Signal	729
15.3.2	Glial Adenosine Clearance and Its Failure in ME/CFS	730
15.3.3	A2A Receptor Upregulation in Neuroinflammation	730
15.3.4	Adenosine, Caffeine Sensitivity, and ADORA2A	731
15.3.5	Therapeutic Implications: A2A Antagonism and Adenosine Modulation	731
15.3.6	Sleep Architecture Findings and Adenosine Interpretation	731
15.4	Inflammatory Cytokine-Induced Somnolence and Fatigue	732
15.4.1	Cytokine-to-Brain Signaling Routes	732
15.4.2	Quantitative Human Cytokine-Fatigue Relationships	733
15.4.3	TLR4/NF- κ B as PEM Amplifier	733
15.4.4	ME/CFS-Specific Cytokine Profile: Evidence and Limitations	734
15.5	Serotonin Dysregulation and the Mood-Fatigue Axis	734
15.5.1	Tryptophan Partitioning in Inflammatory States	734
15.5.2	5-HT Receptor Sensitivity and Genetic Association	735
15.5.3	Serotonin Transporter Abnormalities	735
15.5.4	Gut-Peripheral Serotonin and IBS Comorbidity	735
15.5.5	The SSRI Paradox: Central Serotonin Hyperactivity	736
15.5.6	Central Fatigue: Two-Phase Model	736

15.6	Melatonin Dysfunction and Circadian Disruption	736
15.6.1	DLMO and Multi-System Circadian Decoupling	737
15.6.2	Melatonin–Immune Axis: NK Cells and Cytokine Rhythms	737
15.6.3	Therapeutic Melatonin: Low-Dose Chronobiotic Strategy	738
15.6.4	Light Therapy and Circadian Hygiene: Considerations in ME/CFS . .	738
15.6.5	Melatonin as Antioxidant	739
15.6.6	Integrated Circadian Vicious Cycle	739
15.7	Microglia Activation and Neuroinflammatory Fatigue	739
15.7.1	Neuroimaging Evidence: TSPO-PET and MRS	739
15.7.2	Microglial Phenotype Shift: From Surveillance to Reactivity	740
15.7.3	Circulating Danger Signals and Microglial Activation	740
15.7.4	Purinergic Signaling and Microglial Danger Response	741
15.7.5	Complement Cascade and Synaptic Pruning	741
15.7.6	Therapeutic Modulation of Microglial Activation	741
15.8	Kynurenine Pathway and Quinolinic Acid Excitotoxicity: The “Fog Machine”	742
15.9	Mast Cell Mediators and Histaminergic Symptom Generation	743
15.9.1	MCAS-ME/CFS Overlap and Prevalence	743
15.9.2	Histamine Receptor Pharmacology	743
15.9.3	Mast Cell–Microglia Bidirectional Communication	744
15.9.4	Substance P in Pain Amplification	744
15.10	Glymphatic Dysfunction and Brain Waste Accumulation	744
15.10.1	Glymphatic System Physiology and Sleep Dependency	745
15.10.2	Multiple Converging Impairments in ME/CFS	745
15.11	Central Sensitization and Nociplastic Pain	746
15.11.1	Wind-up and Dorsal Horn Sensitization	746
15.11.2	Evidence for Central Sensitization in ME/CFS	746
15.11.3	Nociplastic Pain Framework	747
15.11.4	Peripheral Sensitization: Substance P and CGRP	747
15.11.5	Treatment Implications	747
15.12	Oxidative and Nitrosative Stress as Symptom Amplifier	748
15.12.1	The NO/ONOO [−] Vicious Cycle	748
15.12.2	Glutathione Depletion and Oxidative Burden	748
15.12.3	TRP Channel Sensitization by ROS	749
15.12.4	Antioxidant Strategies	749
15.13	Metabolic Danger Signals and the Post-Exertional Malaise Mechanism . . .	749
15.13.1	The Lactate Paradox: GPR81 and Anti-Inflammatory Brake Failure . .	750
15.13.2	Succinate as a DAMP and NLRP3 Inflammasome Activator	750
15.13.3	Acid-Sensing Ion Channels and Post-Exertional Pain	750
15.13.4	Two-Day CPET Evidence: Physiological Basis of PEM	750
15.13.5	Why Graded Exercise Therapy Is Contraindicated	751
15.14	Endocannabinoid Deficiency and Mast Cell Brake Failure	751
15.14.1	Endocannabinoid System Architecture	751
15.14.2	Mast Cell ECS Brake and Its Failure	752
15.14.3	Therapeutic Implications	752
15.15	Interoceptive Prediction Error and the Bayesian Brain Fog Framework . . .	753
15.15.1	The Free-Energy Principle and Predictive Processing	753
15.15.2	Interoceptive Hyper-Vigilance in ME/CFS	753

15.15.3	Distinction from Psychosomatic Frameworks	754
15.15.4	Why Graded Exposure Fails	754
15.15.5	Therapeutic Implications: Interoceptive Retraining	754
15.16	Integrated Symptom Cascade Model	755
15.17	Connection to Patient Phenotypes	756
15.18	Therapeutic Implications	757
15.19	Research Directions	759
III Treatment and Management		762
16	Symptom-Based Management	764
16.1	Critical Principle: Aggressive Management of All Comorbidities	764
16.1.1	Why Comorbidity Management Is Critical in ME/CFS	764
16.1.2	Infection Prevention as Therapeutic Priority	770
16.2	Managing Post-Exertional Malaise	771
16.2.1	Pacing and Energy Envelope Theory	771
16.2.2	Medications for PEM	771
16.3	Sleep Management	772
16.3.1	Sleep Hygiene	772
16.3.2	Medications for Sleep	773
16.4	Pain Management	774
16.4.1	Analgesics	774
16.4.2	Neuropathic Pain Medications	774
16.4.3	Opioids	775
16.4.4	Non-pharmacological Pain Management	775
16.5	Cognitive Symptom Management	776
16.5.1	Cognitive Strategies	776
16.5.2	Medications	776
16.6	Orthostatic Intolerance Management	777
16.6.1	Non-pharmacological Approaches	777
16.6.2	Medications	778
16.7	Autonomic Symptom Management	778
16.7.1	Temperature Dysregulation	778
16.7.2	Gastrointestinal Autonomic Symptoms	779
16.7.3	Urinary Autonomic Symptoms	779
17	Urgent Action Plan for Severe Cases	780
17.1	Understanding the Urgency	780
17.1.1	The Current Crisis	780
17.1.2	Why Immediate Action is Justified	780
17.1.3	Honest Assessment: Severe Disease Prognosis and Treatment Potential	781
17.2	Subtype Assessment and Treatment Prioritization	785
17.3	The 2-Week Rapid Relief Protocol	786
17.3.1	Day 1: Immediate Implementation	787
17.3.2	Days 2–7: Protocol Refinement	806
17.3.3	Passive Neuromuscular Electrical Stimulation (NMES) for CNS Bypass	818

Contents

17.3.4	Baseline Symptom Reduction: Strict Pacing	819
17.4	Expected 2-Week Outcomes	822
17.4.1	Cumulative Symptom Relief	822
17.4.2	Transformation of Tolerability	822
17.4.3	Critical Threshold: Bearability	823
17.5	Sensitization Prevention: Foundational Strategy for Severe Cases	824
17.6	Glymphatic Enhancement Protocol for Severe Cases	827
17.7	Medium-Term Recovery Strategies (Weeks to Months)	829
17.7.1	Immunoadsorption for Cognitive Dysfunction	829
17.7.2	Low-Dose IL-2 for Autoimmune Features	830
17.7.3	Hormonal Modulation (Post-Menopausal Women)	832
17.7.4	Anti-Cytokine Therapy (Early Disease <3 Years)	834
17.8	Long-Term Recovery and Fundamental Treatment	837
17.8.1	Comprehensive Biomarker-Guided Approach	837
17.8.2	Investigational Approaches (Clinical Trials)	838
17.8.3	Expected Timeline for Fundamental Recovery	839
17.9	Implementation Checklist	839
17.9.1	Week 1: Immediate Action	839
17.9.2	Weeks 2–4: Consolidation and Planning	840
17.9.3	Months 2–6: Medium-Term Interventions	840
17.10	Special Considerations for Severe Cases	842
17.10.1	Bedbound Patients	842
17.10.2	Patients Considering Medical Assistance in Dying	842
17.10.3	Financial Barriers	842
17.10.4	Structural Evaluation: CCI and EDS in Severe Cases	842
17.11	Summary: Path from Unbearable to Bearable to Improving	844
17.11.1	The Three Stages	844
17.11.2	Key Messages	844
17.11.3	Final Word	845
18	Action Plans for Mild to Moderate Cases	846
18.1	Defining Mild to Moderate ME/CFS	846
18.1.1	Functional Categories	846
18.1.2	Why Action is Urgent for Non-Severe Cases	846
18.2	Severity-Stratified Care Pathways	847
18.2.1	Why Stratification Matters	847
18.2.2	Mild Pathway: Maintaining Function While Preventing Deterioration	848
18.2.3	Moderate Pathway: Stabilization and Symptom Management	849
18.2.4	Borderline Severe Pathway: Preventing Catastrophic Decline	850
18.2.5	Very Severe Pathway: Crisis Prevention and Palliative Support	851
18.3	Immediate Action Plan (Mild-Moderate Cases)	853
18.3.1	Subtype Assessment and Prioritized Treatment Planning	853
18.3.2	Core Principles	855
18.3.3	Foundation: Energy Envelope Management	855
18.3.4	Symptom Management for Mild-Moderate Cases	874
18.3.5	Mast Cell Activation Syndrome (MCAS) Management	885

Contents

18.4	Systematic Comorbidity Screening: The Septad Framework	888
18.4.1	The Seven Septad Components	888
18.4.2	Screening Recommendations for Mild-Moderate Cases	889
18.4.3	Treatment Sequencing	891
18.5	Disease-Modifying Strategies for Mild-Moderate Cases	891
18.5.1	Early Intervention Advantage	891
18.5.2	Acute Onset Protocol: The Critical First Six Months	898
18.5.3	“Brain First” Implementation Protocol for Mild-Moderate Cases	905
18.5.4	Immune Profiling and Targeted Intervention	909
18.5.5	Personalized Cycle Mapping: Precision Diagnostic Framework	909
18.5.6	Early-Disease Anti-Cytokine Strategy	918
18.5.7	Hormonal Optimization	920
18.5.8	Microbiome Restoration	921
18.6	Work and Study Accommodations	924
18.6.1	Critical Reality	924
18.6.2	Formal Accommodations	924
18.6.3	Self-Imposed Boundaries	924
18.6.4	When to Stop Working	925
18.7	Graded Exercise Therapy (GET): Why to Avoid	925
18.7.1	Critical Warning	925
18.7.2	Why GET Fails	925
18.7.3	If Pressured by Physician	925
18.7.4	Safe Activity Increase (If Appropriate)	926
18.8	Long-Term Strategy for Mild-Moderate Cases	926
18.8.1	Goals	926
18.8.2	Timeline	926
18.8.3	Realistic Expectations	926
18.9	Summary: Preventing the Descent	927
18.9.1	Key Principles	927
18.9.2	Action Checklist	927
19	Pediatric ME/CFS: Severe and Housebound Cases	928
19.1	Defining Severe Pediatric ME/CFS	929
19.1.1	Functional Criteria	929
19.1.2	Mast Cell Activation Syndrome (MCAS) Management in Severe Pediatric Cases	930
19.1.3	Distinguishing from Adult Severity Metrics	931
19.2	Home-Based Care Considerations	932
19.2.1	Telehealth Protocols	932
19.2.2	Home Visit Considerations	932
19.2.3	Equipment Needs for Bedbound Children	933
19.2.4	Environmental Modifications	933
19.3	Subtype Assessment for Pediatric Severe Cases	933
19.4	Medical Management	935
19.4.1	Orthostatic Intolerance (Priority)	935
19.4.2	Sensitization Prevention Protocol	938
19.4.3	Pain Management	941

Contents

19.4.4	Sleep Optimization	942
19.4.5	Cognitive Support	946
19.5	Preventing Complications of Prolonged Bedrest	948
19.5.1	Contracture Prevention	948
19.5.2	Osteoporosis Risk	949
19.5.3	Growth Considerations	949
19.5.4	Skin Integrity	949
19.5.5	Cardiovascular Deconditioning	950
19.6	Educational Continuity	950
19.6.1	Legal Rights to Education	950
19.6.2	Homebound Instruction Protocols	950
19.6.3	Maintaining Academic Trajectory	951
19.6.4	Social Connection Strategies	951
19.6.5	Long-Term Planning	951
19.7	Family Support and Caregiver Coordination	952
19.7.1	Parent Education	952
19.7.2	Caregiver Burden and Support	952
19.7.3	Sibling Considerations	953
19.7.4	Financial Navigation	953
19.7.5	Mental Health Support for Family	954
19.8	Red Flags and Emergency Protocols	954
19.8.1	Malnutrition and Feeding Concerns	954
19.8.2	Dehydration	955
19.8.3	Severe Orthostatic Symptoms	955
19.8.4	Mental Health Crisis	955
19.8.5	Distinguishing Severity from Concurrent Illness	956
19.9	Evidence That Severe Pediatric Disease CAN Reverse	956
19.9.1	Prognosis Data	957
19.9.2	Time to Improvement	957
19.9.3	Factors Associated with Better Outcomes	957
19.9.4	Contrast with Adult Prognosis	958
19.9.5	Hope Maintenance with Realistic Expectations	958
19.9.6	Critical Window for Early Intervention	958
19.10	Summary of Key Recommendations	959
20	Pediatric ME/CFS: School-Attending and Ambulatory Cases	960
20.1	The Early Intervention Imperative	960
20.1.1	Why Early Intervention is More Critical in Pediatrics	960
20.1.2	The Cost of Delayed or Inappropriate Intervention	961
20.2	Subtype Assessment for Ambulatory Pediatric Cases	962
20.3	Diagnosis and Assessment	964
20.3.1	Pediatric Diagnostic Criteria Modifications	964
20.3.2	Distinguishing ME/CFS from Differential Diagnoses	965
20.4	School Accommodation Framework	967
20.4.1	IEP vs. 504 Plan Decision Tree	967
20.4.2	Specific Accommodations by Symptom	969
20.4.3	Communication Templates	971

Contents

20.5	Pacing for Children and Adolescents	972
20.5.1	Age-Appropriate Energy Management	973
20.5.2	Managing the “Push Through” Pressure	974
20.6	Medical Management	976
20.6.1	Early Intervention Window Protocol	976
20.6.2	Orthostatic Intolerance (First-Line)	978
20.6.3	Mast Cell Activation Syndrome (MCAS) Management	980
20.6.4	Sleep	981
20.6.5	Cognitive Symptoms	984
20.6.6	Infection Prevention as Recovery Window Preservation	986
20.7	Activity and Exercise	987
20.7.1	Critical Warning: GET is Contraindicated	987
20.7.2	Safe Activity Maintenance	988
20.8	Social Development Considerations	989
20.8.1	Maintaining Peer Connections	990
20.8.2	Online Social Options	990
20.8.3	Extracurricular Modification	990
20.8.4	Identity Formation	990
20.8.5	Preventing Isolation	991
20.9	Long-Term Management and Monitoring	991
20.9.1	Trajectory Tracking	991
20.9.2	When to Escalate Treatment Intensity	991
20.9.3	Reevaluation Frequency	992
20.9.4	Transition Planning	992
20.10	Prognosis Communication	992
20.10.1	The Pediatric Advantage	992
20.10.2	Defining Recovery Appropriately	993
20.10.3	Realistic Timelines	993
20.10.4	Hope Maintenance During Difficult Periods	993
20.10.5	Contrast with Adult Prognosis	993
20.11	Summary of Key Recommendations	994
21	Medications Targeting Underlying Mechanisms	995
21.1	Immune-Modulating Medications	995
21.1.1	Low-Dose Naltrexone (LDN)	995
21.1.2	Immunoglobulins (IVIG)	997
21.1.3	Rituximab	998
21.1.4	Other Immunomodulators	998
21.2	Antiviral Medications	998
21.2.1	Valacyclovir and Acyclovir	998
21.2.2	Valganciclovir	1000
21.2.3	Antiretroviral Approaches	1003
21.2.4	General Principles for Antiviral Use in ME/CFS	1004
21.3	Mitochondrial Support	1004
21.3.1	Coenzyme Q10 (CoQ10)	1004
21.3.2	NADH	1007
21.3.3	D-Ribose	1008

Contents

21.3.4	L-Carnitine and Acetyl-L-Carnitine	1010
21.3.5	Alpha-Lipoic Acid	1012
21.3.6	Combination Mitochondrial Support Protocols	1013
21.3.7	Limitations and Realistic Expectations	1013
21.4	Herbal Anti-Inflammatory Agents	1014
21.4.1	Devil's Claw (<i>Harpagophytum procumbens</i>)	1014
21.4.2	Palmitoylethanolamide (PEA)	1016
21.5	Neuroprotective and Cognitive Enhancers	1018
21.5.1	Ginkgo biloba (EGb 761)	1018
21.6	Interpreting Treatment Responses	1019
21.6.1	Temporary vs. Durable Responses: A Critical Distinction	1022
21.6.2	The Cimetidine-Antiviral Synergy Hypothesis	1023
21.7	Phenotype-Targeted Treatment Pathways	1024
21.7.1	Treatment Pathway for Viral-Immune-Metabolic ("Cimetidine-Responder") Phenotype	1024
21.7.2	Other Emerging Phenotype-Targeted Pathways	1028
21.8	Autonomic Medications	1028
21.8.1	Pyridostigmine (Mestinon)	1028
21.9	H2 Receptor Antagonists	1031
21.9.1	Cimetidine vs. Famotidine: Critical Differences	1031
21.9.2	Cimetidine-LDN Synergy Protocol for Viral-Immune-Phenotype ME/CFS	1032
21.10	Medication Sensitivity Phenotypes	1035
21.10.1	The Paradoxical Reactor Phenotype	1035
21.10.2	The Excipient Sensitivity Phenotype	1037
21.10.3	Implications for Subgroup-Specific Treatment	1042
21.11	Atypical Antipsychotics: Metabolic Considerations in ME/CFS	1043
21.11.1	Low-Dose Aripiprazole (LDA)	1043
22	Supplements and Nutraceuticals	1047
22.1	Foundational Supplements: Electrolytes and Hydration	1048
22.1.1	Why Electrolytes Matter in ME/CFS	1048
22.1.2	Sodium	1048
22.1.3	Potassium	1049
22.1.4	Magnesium	1050
22.1.5	Complete Electrolyte Formulas	1051
22.1.6	Caution: Electrolyte Initiation in ME/CFS	1052
22.2	Mitochondrial and Energy Support	1053
22.2.1	Coenzyme Q10 (CoQ10/Ubiquinone/Ubiquinol)	1053
22.2.2	NADH (Reduced Nicotinamide Adenine Dinucleotide)	1054
22.2.3	NAD ⁺ Precursors: Nicotinamide Riboside (NR) and NMN	1056
22.2.4	D-Ribose	1057
22.2.5	Acetyl-L-Carnitine (ALCAR) and L-Carnitine	1058
22.2.6	Creatine	1059
22.2.7	PQQ (Pyrroloquinoline Quinone)	1060
22.2.8	Mitochondrial Energy Support: Combination Guide	1060

Contents

22.3	Antioxidant and Anti-inflammatory Supplements	1064
22.3.1	N-Acetylcysteine (NAC)	1064
22.3.2	Alpha-Lipoic Acid (ALA)	1065
22.3.3	Omega-3 Fatty Acids (EPA/DHA)	1066
22.3.4	Curcumin	1067
22.3.5	Quercetin	1067
22.4	B Vitamins	1068
22.4.1	Thiamine (B1)	1068
22.4.2	Riboflavin (B2)	1068
22.4.3	Niacin/Niacinamide (B3)	1069
22.4.4	Pyridoxine/P5P (B6)	1069
22.4.5	Folate (B9)	1069
22.4.6	Cobalamin (B12)	1070
22.5	Vitamin D	1070
22.6	Amino Acids	1071
22.6.1	Taurine	1071
22.6.2	Glycine	1072
22.6.3	L-Glutamine	1072
22.6.4	5-HTP (5-Hydroxytryptophan)	1073
22.7	Practical Supplement Protocols	1073
22.7.1	Minimal Cost Protocol (Under \$30/month)	1074
22.7.2	Moderate Protocol (\$50–100/month)	1074
22.7.3	Comprehensive Protocol (\$100–200/month)	1074
22.7.4	By Symptom Cluster	1075
22.7.5	Introduction Strategy	1076
22.8	Additional Supplements	1076
22.8.1	Medium-Chain Triglycerides (MCT)	1076
22.8.2	Resveratrol	1077
22.8.3	Melatonin	1077
22.8.4	Pregnenolone	1078
22.8.5	Zinc	1079
22.9	Probiotics and Gut Health	1080
22.9.1	Probiotics	1080
22.9.2	Prebiotics	1081
22.9.3	Butyrate and Tributyrin	1081
22.10	Supplement Quality and Safety	1082
22.10.1	Quality Considerations	1082
22.10.2	Drug Interactions	1083
22.10.3	When to Stop	1083
22.11	Conclusion	1083
23	Lifestyle and Non-Pharmacological Interventions	1085
23.1	Pacing and Energy Management	1085
23.1.1	Energy Envelope Theory	1085
23.1.2	Practical Pacing Strategies	1087
23.1.3	Activity-Based Dietary Timing for Severe Patients	1088

Contents

23.2	Sleep Optimization	1089
23.2.1	Bedroom Environment	1089
23.2.2	Sleep Schedule Consistency	1090
23.2.3	Light Exposure Management	1090
23.2.4	Evening Routines	1091
23.3	Dietary Approaches	1091
23.3.1	General Nutritional Principles	1091
23.3.2	Specific Dietary Patterns	1092
23.3.3	Meal Timing and Frequency	1092
23.3.4	Food Sensitivities	1093
23.4	Exercise and Movement	1097
23.4.1	The Exercise Paradox	1097
23.4.2	Safe Movement Approaches	1102
23.5	Stress Management	1104
23.5.1	Relaxation Techniques	1105
23.5.2	Meditation and Mindfulness	1105
23.5.3	Biofeedback	1106
23.6	Environmental Modifications	1106
23.6.1	Home Adaptations	1106
23.6.2	Chemical and Environmental Sensitivities	1107
23.7	Social and Emotional Support	1107
23.7.1	Support Groups and Peer Connection	1107
23.7.2	Counseling and Psychotherapy	1108
23.7.3	Addressing Secondary Depression and Anxiety	1108
23.7.4	Family and Caregiver Education	1109
24	Experimental and Emerging Therapies	1110
24.1	Novel Therapeutic Frameworks	1110
24.2	Immunological Interventions	1112
24.2.1	Autoantibody-Targeted Therapies	1112
24.2.2	Cytokine Modulation	1114
24.2.3	Cellular Therapies	1115
24.3	Autonomic and Neurological Interventions	1115
24.3.1	Vagal Tone Restoration	1115
24.3.2	Neurostimulation	1117
24.3.3	Cerebrospinal Fluid Interventions	1117
24.4	Metabolic Interventions	1121
24.4.1	Mitochondrial “Jumpstart” Protocols	1121
24.4.2	NAD ⁺ Precursor Therapy	1122
24.4.3	Metabolic Modulators	1123
24.4.4	Ketogenic and Metabolic Switching Approaches	1124
24.5	Microbiome Interventions	1124
24.5.1	Fecal Microbiota Transplantation	1124
24.5.2	Precision Microbiome Modulation	1125
24.6	Technologies and Devices	1126
24.6.1	Apheresis Techniques	1126
24.6.2	Hyperbaric Oxygen Therapy	1126

Contents

24.6.3	Photobiomodulation	1126
24.7	Repurposed Medications	1127
24.7.1	Suramin	1127
24.7.2	Rapamycin (Sirolimus)	1127
24.7.3	Metformin	1127
24.7.4	Low-Dose Aripiprazole	1127
24.7.5	Ketamine	1128
24.8	Peptide Therapies	1130
24.8.1	BPC-157	1130
24.8.2	Thymosin Alpha-1	1130
24.9	Integrated Treatment Strategies	1130
24.9.1	Cycle-Based Multi-Target Treatment	1131
24.10	CPET-Derived Multi-Target Protocols	1140
24.10.1	The Autonomic-Metabolic Recovery Protocol	1140
24.10.2	The Mitochondrial Turnover Acceleration Protocol	1142
24.10.3	Post-Exertional Malaise Prevention and Mitigation	1144
24.10.4	Personalized Metabolomics-Guided Protocol (Future Direction)	1152
24.10.5	Clinical Implementation Guidance	1152
24.10.6	Research Priorities	1153
24.11	Evaluating Emerging Therapies	1154
24.11.1	Risk-Benefit Assessment	1154
24.11.2	Evidence Hierarchy	1154
24.11.3	Access Pathways	1154
24.11.4	Reversibility Windows: Setting Realistic Treatment Goals	1155
24.11.5	Quantitative Cycle Gain Measurement: A Prognostic Tool	1163
24.11.6	Epigenetic Reversal Strategies	1166
25	Integrative and Personalized Treatment Approaches	1170
25.1	Developing a Treatment Plan	1170
25.1.1	The Imperative of Treatment Safety in ME/CFS	1170
25.1.2	Treatment Energy Categories	1171
25.1.3	Baseline Assessment	1171
25.1.4	The Micro-Dosing Imperative	1172
25.1.5	Crash-Risk Ranking and Decision Rules	1174
25.1.6	Prioritizing Interventions	1176
25.1.7	When to Stop Trying	1177
25.1.8	Medication-Induced Crash Recovery Protocol	1178
25.1.9	Tracking Progress	1178
25.2	Treating Comorbidities	1180
25.2.1	POTS Management	1180
25.2.2	Mast Cell Activation Syndrome	1180
25.2.3	Ehlers-Danlos Syndrome	1180
25.2.4	Other Common Comorbidities	1180
25.3	Personalized Medicine Approaches	1187
25.3.1	Biomarker-Guided Treatment	1187
25.3.2	Pharmacogenomics	1188
25.3.3	Subtype-Specific Treatment Approaches	1189

25.4	Combination Therapies	1192
25.4.1	Documented Synergies	1192
25.4.2	Interaction Risks	1193
25.4.3	Sequencing Considerations	1193
25.5	Cross-Domain Medical Parallels: Learning from Other Fields	1193
25.5.1	Rationale for Cross-Domain Knowledge Transfer	1193
25.5.2	High-Altitude Medicine: Chronic Hypoxia Parallels	1194
25.5.3	Critical Care and ICU Recovery Medicine	1197
25.5.4	Space Medicine: Orthostatic Intolerance and Deconditioning	1201
25.5.5	Additional Domain Parallels: Brief Overview	1205
25.5.6	Integration and Practical Application	1206
25.5.7	Cautions and Limitations	1208
25.5.8	Research Implications: Cross-Domain Studies	1209
25.5.9	Conclusion: The Value of Looking Beyond ME/CFS	1209
25.6	Novel Mechanistic Hypotheses and Research Opportunities	1210
25.6.1	WASF3 as Therapeutic Target	1210
25.6.2	Acetylcholine-Mitochondrial Axis	1210
25.6.3	ATP Recovery Kinetics and Mitophagy	1210
25.6.4	Viral Trigger-ER Stress-WASF3 Pathway	1211
25.6.5	Pyruvate Supplementation Hypothesis	1211
25.6.6	Methylene Blue as Electron Transport Bypass	1212
25.6.7	Beta-Blockers for Pacing Enforcement	1212
25.6.8	Immune Checkpoint Modulation	1212
25.6.9	Central Governor Theory Link	1213
25.6.10	Lactate Clearance Dysfunction	1213
25.6.11	Mast Cell-Mitochondrial Crosstalk	1214
25.6.12	Research Priorities	1214
25.6.13	Summary of Novel Hypotheses and Interventions	1215
25.7	Working with Healthcare Providers	1218
25.7.1	Communicating the Phenotype-First Approach	1218
25.7.2	Building a Treatment Team	1218

IV Research and Evidence Synthesis 1219

26 Biomarker Research 1221

26.1	Overview of Biomarker Development	1221
26.1.1	Why Biomarkers Are Needed	1221
26.1.2	Types of Biomarkers	1221
26.1.3	Challenges in ME/CFS Biomarker Research	1222
26.1.4	Convergent Biology Despite Heterogeneity	1223
26.2	Key Biomarkers from the NIH Deep Phenotyping Study	1224
26.2.1	Cerebrospinal Fluid Biomarkers	1224
26.2.2	Immune Biomarkers	1225
26.2.3	Autonomic Biomarkers	1225
26.2.4	Cardiopulmonary Biomarkers	1226
26.2.5	Neuroimaging Biomarkers	1226

Contents

26.3	Metabolomic Biomarkers	1226
26.3.1	Key Metabolomic Studies	1226
26.3.2	Synthesis of Metabolomic Findings	1227
26.4	Immunological Biomarkers	1228
26.4.1	Cytokine Profiles	1228
26.4.2	Cell Function Markers	1229
26.5	Neurological Biomarkers	1230
26.5.1	Brain Imaging Markers	1230
26.5.2	CSF Findings	1230
26.5.3	Autonomic Function Tests	1231
26.5.4	Cognitive Testing Patterns	1231
26.6	Genomic and Epigenetic Biomarkers	1231
26.6.1	Gene Expression Signatures	1231
26.6.2	miRNA Profiles	1232
26.6.3	DNA Methylation Patterns	1232
26.6.4	Clinical Utility	1232
26.7	Proteomic Biomarkers	1232
26.7.1	Protein Expression Patterns	1232
26.7.2	Autoantibody Panels	1233
26.7.3	Diagnostic Potential	1233
26.8	Composite Biomarker Panels	1233
26.8.1	Multi-Omics Approaches	1233
26.8.2	Machine Learning Applications	1233
26.8.3	Diagnostic Accuracy	1234
26.8.4	Commercial Tests Available	1234
26.9	Functional Biomarkers	1234
26.9.1	Two-Day CPET Protocol	1234
26.9.2	NASA Lean Test	1234
26.9.3	Cognitive Testing	1235
26.10	Biomarker Validation and Standardization	1235
26.10.1	Replication Requirements	1235
26.10.2	Standardization Efforts	1235
26.10.3	Path to Clinical Implementation	1235
26.11	Summary: Current State and Future Directions	1236
27	Clinical Trials and Treatment Studies	1237
27.1	Immunological Interventions	1237
27.1.1	Rituximab Trials	1237
27.1.2	Low-Dose Naltrexone Studies	1238
27.1.3	Other Immune Modulators	1238
27.2	Antiviral Trials	1239
27.3	Metabolic and Mitochondrial Interventions	1239
27.4	Neurological Interventions	1240
27.5	Exercise and Rehabilitation Trials	1241
27.5.1	The PACE Trial	1241
27.5.2	Alternative Rehabilitation Approaches	1241
27.6	Complementary and Alternative Medicine Trials	1242

27.7	Ongoing and Planned Trials	1242
27.8	Synthesis and Meta-Analyses	1243
27.9	Negative Trials and Null Results	1244
27.9.1	Immunological Interventions	1244
27.9.2	Behavioral and Rehabilitation Interventions	1245
27.9.3	Antiviral Interventions	1246
27.9.4	Neurological and Cholinergic Interventions	1247
27.9.5	Antidepressant Interventions	1247
27.9.6	Endocrine Interventions	1247
27.9.7	Summary Table	1247
27.9.8	Cross-Cutting Lessons	1247
28	Mechanistic and Experimental Studies	1250
28.1	The NIH Deep Phenotyping Study (Walitt et al. 2024)	1250
28.1.1	Study Design and Methodology	1250
28.1.2	Key Biological Findings	1251
28.1.3	Methodological Limitations and Criticisms	1253
28.1.4	The “Effort Preference” Controversy	1253
28.1.5	Interpretation and Context	1255
28.2	Related Studies from the NIH Cohort	1256
28.2.1	WASF3 and Mitochondrial Dysfunction (Wang et al. 2023)	1256
28.2.2	T Cell Exhaustion (Iu et al. 2024)	1256
28.3	Exercise Physiology Studies	1257
28.3.1	Cardiopulmonary Exercise Testing (CPET)	1257
28.3.2	Muscle Studies	1257
28.4	Cellular and Molecular Studies	1258
28.4.1	Cell Culture Studies	1258
28.4.2	Animal Models	1259
28.5	Imaging Studies	1261
28.5.1	Brain Imaging	1261
28.5.2	Cardiac Imaging	1262
28.6	Immunological Studies	1262
28.7	Omics Studies	1263
28.7.1	Genomics	1263
28.7.2	Transcriptomics	1264
28.7.3	Proteomics	1264
28.7.4	Metabolomics	1264
28.7.5	Lipidomics	1265
28.7.6	Microbiomics	1265
28.8	Integrative Multi-Omics Studies	1266
29	Epidemiological and Outcomes Research	1267
29.1	Prevalence and Incidence Studies	1267
29.1.1	Population-Based Estimates	1267
29.1.2	Geographic and Demographic Variation	1267
29.1.3	Post-COVID Surge	1268

Contents

29.2	Risk Factor Studies	1268
29.2.1	Genetic Risk Factors	1268
29.2.2	Environmental Risk Factors	1268
29.2.3	Demographic Risk Factors	1269
29.3	Natural History Studies	1269
29.3.1	Recovery Rates	1269
29.3.2	Predictors of Outcome	1270
29.4	Quality of Life and Disability Studies	1270
29.4.1	SF-36 Findings	1270
29.4.2	Comparison to Other Chronic Diseases	1270
29.4.3	Economic Burden	1271
29.4.4	Impact on Caregivers	1271
29.5	Mortality Studies	1272
29.5.1	All-Cause Mortality: Evidence from Population Cohorts	1272
29.5.2	Cause-Specific Mortality	1274
29.5.3	Actuarial and Insurance Industry Perspective	1275
29.5.4	Methodological Challenges and Evidence Quality	1276
29.5.5	Summary: Evidence-Based Conclusions	1277
29.6	Comorbidity Studies	1277
29.6.1	Patterns of Comorbid Conditions	1277
29.6.2	Temporal Relationships	1278
29.6.3	Shared Mechanisms	1278
30	Controversies and Debates in ME/CFS Research	1279
30.1	Nomenclature and Definition	1279
30.2	Diagnostic Criteria Controversies	1279
30.3	The PACE Trial Controversy	1280
30.3.1	What the Trial Claimed	1280
30.3.2	Criticisms of Methodology	1281
30.3.3	Reanalysis Results	1281
30.3.4	Impact on Clinical Guidelines	1281
30.3.5	Lessons for Future Research	1282
30.4	Psychogenic vs. Biomedical Models	1282
30.4.1	Historical Context	1282
30.4.2	Evidence for Biological Basis	1283
30.4.3	The Role of Psychological Factors	1283
30.4.4	Patient Advocacy and the Psychogenic Harm	1283
30.5	Exercise Therapy Debates	1284
30.5.1	Arguments For Graded Exercise Therapy	1284
30.5.2	Evidence of Harm	1284
30.5.3	Current Recommendations	1284
30.6	Deconditioning Hypothesis	1285
30.6.1	Arguments For Deconditioning	1285
30.6.2	Why Deconditioning Fails as a Primary Explanation	1285
30.6.3	Deconditioning as Secondary Consequence	1286
30.7	The NIH “Effort Preference” Controversy (2024–2025)	1288
30.7.1	The Central Claim	1288

Contents

30.7.2	Why the Framing Was Controversial	1288
30.7.3	Published Academic Responses	1289
30.7.4	NIH Clarification	1290
30.7.5	Clinical and Research Implications	1290
30.7.6	Separating Data from Interpretation	1290
30.8	Post-COVID ME/CFS	1291
30.8.1	Overlap with Long COVID	1291
30.8.2	Differences in Recognition and Funding	1292
30.8.3	Opportunities and Challenges	1292
30.9	Clinical Utility for Respiratory and Allergy Medicine	1292
30.9.1	Transferable Diagnostic Techniques	1293
30.9.2	Underdiagnosed Comorbidities in Allergy Practice	1294
30.9.3	Post-Viral Respiratory Syndrome Recognition	1294
30.9.4	Research Opportunities	1295
30.10	Research Funding Disparities	1295
30.10.1	Historical Underfunding	1295
30.10.2	Recent Improvements	1296
30.10.3	Remaining Gaps	1296
31	Translational Findings: Implications for Related Conditions	1297
31.1	Introduction to Translational Medicine	1297
31.1.1	Why ME/CFS Research Benefits Other Conditions	1297
31.1.2	Certainty Levels for Cross-Condition Application	1298
31.2	Immediate Applicability (Tier 1)	1298
31.2.1	Long COVID / Post-Acute Sequelae of SARS-CoV-2 (PASC)	1298
31.2.2	Postural Orthostatic Tachycardia Syndrome (POTS)	1300
31.2.3	Fibromyalgia	1300
31.2.4	Mast Cell Activation Syndrome (MCAS)	1301
31.2.5	Ehlers-Danlos Syndrome (Hypermobile Type)	1302
31.3	Strong Mechanistic Overlap (Tier 2)	1305
31.3.1	Post-Treatment Lyme Disease Syndrome (PTLDS)	1306
31.3.2	Cancer-Related Fatigue and Post-Chemotherapy Syndrome	1306
31.3.3	Primary Mitochondrial Disorders	1307
31.3.4	Dysautonomia (General)	1308
31.3.5	Small Fiber Neuropathy (SFN)	1309
31.4	Promising But Requires Validation (Tier 3)	1310
31.4.1	Autoimmune Conditions	1310
31.4.2	Neurodegenerative Diseases	1311
31.4.3	Metabolic Syndrome and Type 2 Diabetes	1312
31.5	Key Translational Mechanisms	1313
31.5.1	Plasma Cell Autoimmunity (Daratumumab Target)	1313
31.5.2	GPCR Autoantibodies	1314
31.5.3	Vascular-Immune-Energy Triad	1316
31.5.4	WASF3/ER Stress → Mitochondrial Dysfunction	1318
31.5.5	NAD ⁺ Depletion	1320
31.5.6	Glymphatic Clearance Failure	1322

31.6	Research Priorities and Future Directions	1325
31.6.1	Cross-Condition Mechanism Validation	1325
31.6.2	Biomarker Validation Across Conditions	1325
31.7	Universal Treatment Protocols	1325
31.7.1	Comprehensive Mitochondrial Support	1326
31.7.2	Autonomic-Catecholamine Restoration	1328
31.7.3	Mast Cell Stabilization	1330
31.7.4	Neuroinflammation Reduction	1332
31.7.5	Energy Envelope Management (Pacing)	1334
31.7.6	Clinical Trial Opportunities	1336
31.8	Conclusion	1336
32	Research Infrastructure Proposals	1338
32.1	Longitudinal Deep Phenotyping Cohort	1338
32.1.1	The ME/CFS Human Phenome Project	1338
32.2	Recovery Potential Index Development	1342
32.2.1	Conceptual Framework	1342
32.2.2	Component Biomarkers	1343
32.2.3	Index Construction and Interpretation	1347
32.2.4	Validation Requirements	1348
32.2.5	Limitations and Caveats	1348
32.3	Crash Impact Biomarkers	1349
32.3.1	Conceptual Framework	1349
32.3.2	Proposed Biomarker Panel	1349
32.3.3	Validation Study Design	1351
32.3.4	Clinical Utility	1352
32.4	Computational Patient Modeling: The ME/CFS Digital Twin	1353
32.4.1	The Digital Twin Concept	1353
32.4.2	Data Sources for ME/CFS Digital Twin	1353
32.4.3	Modeling Approaches	1354
32.4.4	Clinical Applications	1355
32.4.5	Development Pathway	1356
32.4.6	Challenges and Limitations	1356
32.5	Model Organism Research Panel	1357
32.5.1	Animal Models of Post-Viral Fatigue	1357
32.5.2	Cellular Models of Metabolic Dysfunction	1357
32.5.3	Organoid Systems for Tissue-Specific Effects	1358
32.5.4	Towards an Integrated Model Panel	1358
33	Proposed Research Studies	1359
33.1	Multi-Modal Testing of Selective Energy Dysfunction Hypothesis	1359
33.1.1	Background and Rationale	1359
33.1.2	Study Innovation	1360
33.1.3	Hypothesis	1360
33.1.4	Study Design Overview	1361
33.1.5	Five Core Aims	1361
33.1.6	Key Study Measures	1362

Contents

33.1.7	Expected Outcomes and Implications	1363
33.1.8	Budget and Timeline Overview	1364
33.1.9	Funding and Implementation	1364
33.2	Secondary Analysis Opportunities: Leveraging Existing Datasets	1365
33.2.1	Rationale for Secondary Analysis	1365
33.2.2	Prediction-to-Dataset Coverage	1365
33.2.3	Priority Datasets	1366
33.2.4	Relationship to the Multi-Modal Study	1367
33.3	Pediatric-Adult ME/CFS Comparison Study	1367
33.3.1	Background and Rationale	1367
33.3.2	Hypotheses	1368
33.3.3	Study Design	1368
33.3.4	Measures	1369
33.3.5	Outcomes	1371
33.3.6	Analysis Plan	1371
33.3.7	Ethical Considerations	1372
33.3.8	Expected Outcomes and Implications	1372
33.4	Aggressive Early Intervention Trial	1373
33.4.1	Background and Rationale	1373
33.4.2	Study Objectives	1374
33.4.3	Study Design	1374
33.4.4	Interventions	1376
33.4.5	Outcomes	1378
33.4.6	Safety Monitoring	1379
33.4.7	Feasibility Considerations	1379
33.4.8	Expected Outcomes and Implications	1380
33.5	Crash Impact on Recovery Biomarkers Study	1381
33.5.1	Background and Rationale	1381
33.5.2	Hypothesis	1381
33.5.3	Study Design	1381
33.5.4	Outcomes	1382
33.5.5	Analysis Plan	1383
33.5.6	Sample Size Justification	1383
33.5.7	Expected Outcomes	1383
33.6	Orthostatic Intolerance Treatment Durability Study	1383
33.6.1	Background and Rationale	1383
33.6.2	Hypothesis	1384
33.6.3	Study Design	1384
33.6.4	Outcomes	1385
33.6.5	Analysis Plan	1385
33.6.6	Expected Outcomes and Implications	1385
33.7	HRV-Guided Pacing Randomized Controlled Trial	1385
33.7.1	Background and Rationale	1385
33.7.2	Hypothesis	1386
33.7.3	Study Design	1386
33.7.4	Outcomes	1387
33.7.5	Analysis Plan	1388

Contents

33.7.6	Sample Size Justification	1388
33.7.7	Expected Outcomes and Implications	1388
33.8	Sports Medicine-Adapted Periodization RCT	1389
33.8.1	Background and Rationale	1389
33.8.2	Hypothesis	1389
33.8.3	Study Design	1389
33.8.4	Outcomes	1390
33.8.5	Safety Monitoring	1390
33.8.6	Expected Outcomes	1391
33.9	Longitudinal Microglial Imaging Study	1391
33.9.1	Background and Rationale	1391
33.9.2	Study Design	1391
33.9.3	Assessment Schedule	1392
33.9.4	Measures	1392
33.9.5	Outcomes	1393
33.9.6	Analysis Plan	1393
33.9.7	Sample Size and Power	1393
33.9.8	Expected Outcomes and Implications	1394
33.10	Treatment Sequence RCT: “Brain First” Versus Peripheral-First Approaches	1394
33.10.1	Background and Rationale	1394
33.10.2	Study Design	1395
33.10.3	Treatment Arms	1395
33.10.4	Outcomes	1396
33.10.5	Analysis Plan	1396
33.10.6	Sample Size Justification	1397
33.10.7	Expected Outcomes and Implications	1397
33.11	Infection-Divine Correlation Study	1397
33.11.1	Background and Rationale	1397
33.11.2	Study Design	1398
33.11.3	Assessment Schedule	1398
33.11.4	Infection Documentation Protocol	1398
33.11.5	Measures	1399
33.11.6	Outcomes	1399
33.11.7	Analysis Plan	1400
33.11.8	Sample Size Justification	1400
33.11.9	Expected Outcomes and Implications	1400
33.12	Sleep-Glymphatic-Neuroinflammation Pathway Study	1401
33.12.1	Background and Rationale	1401
33.12.2	Study Design	1401
33.12.3	Interventions	1402
33.12.4	Measures	1403
33.12.5	Outcomes	1403
33.12.6	Analysis Plan	1404
33.12.7	Sample Size Justification	1404
33.12.8	Expected Outcomes and Implications	1404
33.13	Metabolic-Immune Crosstalk Study	1405
33.13.1	Background and Rationale	1405

33.13.2	Study Design	1405
33.13.3	Overall Study Measures	1407
33.13.4	Analysis Plan	1407
33.13.5	Sample Size Justification	1408
33.13.6	Expected Outcomes and Implications	1408
33.14	Gut-Vagal Pathway Research Priorities	1409
33.14.1	Background and Rationale	1409
33.14.2	Priority Study 1: Cross-Sectional Gut-Vagal Phenotyping	1409
33.14.3	Priority Study 2: Butyrate Supplementation RCT	1410
33.14.4	Priority Study 3: HRV-Guided Phenotype Reanalysis	1411
33.15	Replication Methodology for Heterogeneous Disease	1411
33.15.1	Why Average-Effect Replication Fails	1411
33.15.2	Master Protocol Designs for ME/CFS	1412
33.15.3	Adaptive Enrichment and Biomarker-Guided Designs	1414
33.15.4	N-of-1 Trials and Aggregation	1414
33.16	Cross-Document Hypothesis Registry	1415

V Mathematical and Computational Modeling 1433

34 Foundations of ME/CFS Modeling 1435

34.1	Why Model ME/CFS?	1436
34.1.1	Managing Multi-System Complexity	1436
34.1.2	Generating Testable Predictions	1436
34.1.3	Optimizing Interventions	1436
34.1.4	Personalized Medicine	1436
34.2	Prior Work and Scope of Contribution	1437
34.2.1	Existing Models	1437
34.2.2	Adjacent Computational Work	1438
34.2.3	What This Part Contributes—and What It Does Not	1438
34.3	Modeling Approaches	1440
34.3.1	Ordinary Differential Equations	1440
34.3.2	Stochastic Models	1441
34.3.3	Network Models	1441
34.3.4	Agent-Based Models	1442
34.3.5	Hybrid and Multi-Scale Models	1442
34.4	Computational Methods	1442
34.4.1	Parameter Estimation	1442
34.4.2	Bayesian Inference	1443
34.4.3	Sensitivity Analysis	1443
34.5	Data Requirements	1443
34.5.1	Longitudinal Data	1444
34.5.2	Multi-Omics Integration	1444
34.5.3	Clinical Parameters	1444
34.6	Model Validation	1444
34.6.1	Predictive Accuracy	1445
34.6.2	Biological Plausibility	1445

34.6.3	Structural Identifiability	1445
34.7	Model Coverage Justification	1445
34.7.1	Coverage of Part II Pathophysiology	1446
34.7.2	Selection Rationale	1446
34.7.3	What Is Not Modeled	1447
35	Energy Metabolism Models	1448
35.1	ATP Production Models	1450
35.1.1	Glycolysis Kinetics	1450
35.1.2	Krebs Cycle Model	1450
35.1.3	Electron Transport Chain Model	1451
35.1.4	Integrated Energy Production	1451
35.2	Mitochondrial Dysfunction Models	1452
35.2.1	Reactive Oxygen Species Production and Damage	1452
35.2.2	Impaired Energy Production in ME/CFS	1452
35.3	Post-Exertional Malaise Modeling	1453
35.3.1	Exertion-Induced Metabolic Crisis	1453
35.3.2	Temporal Evolution of PEM	1454
35.3.3	Energy Envelope Model	1454
35.4	Response to Metabolic Interventions	1455
35.4.1	CoQ10 Supplementation	1455
35.4.2	NAD ⁺ Precursor Supplementation	1455
35.4.3	Prediction Limitations	1455
35.5	Lactate Kinetics and Metabolic Flexibility	1456
35.5.1	Lactate Production and Clearance	1456
35.5.2	Metabolic Flexibility Model	1456
35.5.3	Carnitine Dynamics	1457
35.6	Mitochondrial Dynamics and Quality Control	1457
35.6.1	Fission–Fusion Balance	1458
35.6.2	Mitophagy and PINK1/Parkin Pathway	1458
35.6.3	Biogenesis and PGC-1 α Regulation	1458
35.7	Model Application Guide	1459
35.7.1	Glycolysis Kinetics (Equation 35.1)	1459
35.7.2	Krebs Cycle (Equation 35.2)	1461
35.7.3	Electron Transport Chain (Equations 35.3–35.5)	1461
35.7.4	Reactive Oxygen Species (Equations 35.7–35.8)	1463
35.7.5	Post-Exertional Malaise and Energy Envelope (Equations 35.9–35.12)	1463
35.7.6	Mitochondrial Quality Control (Equations 35.17–35.20)	1465
35.7.7	Lactate Kinetics and Metabolic Flexibility (Equations 35.13–35.15)	1466
36	Immune System Models	1467
36.1	Innate Immunity Dynamics	1468
36.1.1	NK Cell Activity Model	1468
36.1.2	Cytokine Network Model	1468
36.1.3	Chronic Immune Activation	1469
36.2	Adaptive Immunity Models	1469
36.2.1	T Cell Dynamics	1469

Contents

36.2.2	B Cell and Autoantibody Production	1470
36.3	Neuroinflammation Models	1471
36.3.1	Microglial Activation	1471
36.3.2	Blood–Brain Barrier Model	1471
36.4	Viral Reactivation Models	1472
36.5	Response to Immune Interventions	1472
36.5.1	B Cell Depletion (Rituximab)	1472
36.5.2	Low-Dose Naltrexone	1473
36.5.3	Plasma Cell Depletion (Daratumumab)	1473
36.6	Mast Cell Activation Dynamics	1473
36.6.1	Mast Cell Degranulation Model	1474
36.6.2	Mediator Release and Downstream Effects	1474
36.6.3	Pharmacological Predictions	1475
36.7	Coagulation and Microvascular Dynamics	1475
36.7.1	Coagulation Cascade Model	1476
36.7.2	Coupling to Oxygen Delivery	1476
36.7.3	Endothelial Dysfunction Model	1477
36.8	Model Application Guide	1477
36.8.1	NK Cell Activity (Equations 36.1–36.4)	1478
36.8.2	Cytokine Network (Equations 36.5–36.6)	1479
36.8.3	Viral Reactivation (Equations 36.20–36.21)	1480
36.8.4	B Cell/Autoantibody and Rituximab/Daratumumab Models (Equations 36.12–36.22)	1480
36.8.5	Mast Cell Activation (Equations 36.23–36.26)	1481
36.8.6	Coagulation and Microclots (Equations 36.28–36.32)	1482
37	Neuroendocrine and Autonomic Models	1484
37.1	HPA Axis Models	1486
37.1.1	HPA Axis Dynamics	1486
37.1.2	HPA Axis Dysfunction in ME/CFS	1486
37.2	Autonomic Nervous System Models	1489
37.2.1	Sympathetic–Parasympathetic Balance	1489
37.2.2	Orthostatic Intolerance Model	1489
37.2.3	POTS Mechanism Model	1490
37.3	Neurotransmitter Models	1492
37.3.1	Tryptophan–Kynurenine Pathway	1492
37.3.2	Catecholamine Dynamics	1492
37.4	Sleep–Wake Cycle Models	1493
37.5	Tetrahydrobiopterin Competition Model	1494
37.5.1	BH ₄ Pool Dynamics	1494
37.6	Cerebral Blood Flow Autoregulation	1495
37.6.1	Cerebrovascular Autoregulation Model	1495
37.7	Central Sensitization and Pain Amplification	1496
37.7.1	Spinal Cord Sensitization Dynamics	1497
37.7.2	Pain as a Feedback Process	1497
37.7.3	Small Fiber Neuropathy Coupling	1498

37.8	Model Application Guide	1498
37.8.1	HPA Axis (Equations 37.1 and Section 37.1.2)	1498
37.8.2	Autonomic Balance and Orthostatic Intolerance (Equations 37.4–37.9)	1499
37.8.3	Tryptophan–Kynurenine Axis (Equations 37.10–37.13)	1500
37.8.4	BH ₄ Competition (Equation 37.18)	1501
37.8.5	Sleep–Wake Cycle (Equations 37.16)	1502
37.8.6	Central Sensitization (Equations 37.21–37.23)	1504
38	Integrated Multi-System Models	1505
38.1	Energy–Immune Coupling	1507
38.1.1	Immune Activation Drains Energy	1507
38.1.2	Energy Deficits Impair Immunity	1507
38.2	Neuroimmune Interactions	1508
38.2.1	Cytokine Effects on Brain Function	1508
38.2.2	HPA Axis and Immune Regulation	1508
38.2.3	Vagal Signaling	1509
38.3	Cardiovascular–Metabolic Integration	1509
38.4	Gut–Brain–Immune Axis	1510
38.4.1	GI Motility and SIBO Dynamics	1510
38.5	Whole-Body Systems Model	1513
38.5.1	Model Architecture	1513
38.5.2	Healthy Baseline and ME/CFS Disease State	1513
38.5.3	Perturbation Responses	1514
38.6	Symptom Generation Mechanisms	1514
38.7	Extended Subsystem Couplings	1515
38.7.1	Mast Cell–Energy–Autonomic Triangle	1515
38.7.2	Coagulation–Oxygenation–Mitochondrial Cascade	1516
38.7.3	Connective Tissue and Vascular Compliance	1516
38.7.4	Epigenetic State as a Slow Variable	1516
38.7.5	Updated Whole-Body State Vector	1517
38.8	Bifurcation Analysis and Disease Subtypes	1518
38.8.1	Steady-State Multiplicity	1518
38.8.2	Separatrix Topology and Recovery Paths	1519
38.8.3	Parameter Sensitivity of Bifurcation Structure	1519
38.9	Model Application Guide	1520
38.9.1	Energy–Immune Coupling and Bistability (Equations 38.1–38.2)	1520
38.9.2	Gut–Brain–Immune Axis and GI Motility/SIBO (Equations 38.8–38.16)	1521
38.9.3	Bifurcation Analysis and Disease Subtypes (Section 38.8)	1523
38.9.4	Whole-Body 64-Variable Model (Section 38.5)	1523
39	Temporal Evolution and Disease Trajectories	1525
39.1	Disease Onset Models	1525
39.1.1	Triggering Events	1525
39.1.2	Tipping Point Dynamics	1526
39.1.3	Failure of Recovery Mechanisms	1526
39.2	Disease Progression Models	1527
39.2.1	Stable vs. Progressive Disease	1527

Contents

39.2.2	Relapse–Remission Patterns	1527
39.2.3	The Energy Ratchet: A Discrete Event Model	1528
39.3	Daily and Weekly Symptom Dynamics	1532
39.3.1	Circadian Influences	1532
39.3.2	Activity–Symptom Relationships	1532
39.4	Response to Specific Stimuli	1532
39.4.1	Physical Exertion	1532
39.4.2	Cognitive Exertion	1533
39.4.3	Infections	1533
39.4.4	Environmental Factors	1533
39.5	Treatment Response Modeling	1534
39.5.1	Time Course of Improvement	1534
39.5.2	Combination Therapy Effects	1534
39.6	Long-Term Trajectories	1535
39.7	Critical Slowing Down and Early Warning Signals	1535
39.7.1	Mathematical Basis	1535
39.7.2	Clinical Application: Crash Prediction	1536
39.7.3	Recovery Proximity Detection	1536
39.8	Hysteresis and the Intervention Window	1537
39.8.1	Structural Hysteresis	1537
39.8.2	Parametric Hysteresis and the Intervention Window	1537
39.9	Endogenous Oscillations and Hopf Bifurcation	1538
39.9.1	Conditions for Oscillation	1538
39.9.2	Oscillation as a Biomarker	1539
39.9.3	Stochastic Resonance and Noise-Assisted Recovery	1539
39.10	Model Application Guide	1540
39.10.1	Tipping Point and Disease Onset Models	1540
39.10.2	Damage Accumulation and Energy Ratchet Models	1541
39.10.3	Critical Slowing Down and Early Warning Signals	1542
39.10.4	Hysteresis and Intervention Window	1544
39.10.5	Oscillation Analysis and Stochastic Resonance	1545
39.10.6	Scope and Rationale	1546
40	Predictive Applications and Clinical Translation	1547
40.1	Personalized Pacing Optimization	1547
40.1.1	Individual Energy Budget Calculation	1548
40.1.2	Activity Planning Algorithms	1548
40.1.3	Real-Time Monitoring Integration	1548
40.2	Treatment Selection and Optimization	1549
40.2.1	Biomarker-Based Treatment Prediction	1549
40.2.2	Dosing Optimization	1549
40.2.3	Combination Therapy Design	1550
40.3	Prognosis Prediction	1550
40.4	Clinical Decision Support	1550
40.4.1	Dashboard Concept	1551
40.4.2	Treatment Algorithms	1551

40.5	Drug Development Applications	1551
40.5.1	Target Identification	1551
40.5.2	Virtual Clinical Trials	1552
40.6	Optimal Control Theory for Pacing	1552
40.6.1	Pontryagin Maximum Principle Formulation	1553
40.6.2	Hamilton–Jacobi–Bellman Approach for Stochastic Pacing	1554
40.7	Global Sensitivity Analysis and Drug Target Ranking	1554
40.7.1	Sobol Index Framework	1554
40.7.2	Subtype-Stratified Sensitivity	1555
40.8	Network Controllability and Minimum Intervention Sets	1556
40.8.1	Structural Controllability Analysis	1556
40.8.2	Optimal Driver Node Selection	1557
40.9	Model-Predicted Treatment Candidates	1557
40.9.1	Synergy Matrix for Combination Therapy	1557
40.10	Virtual Population Simulation	1559
40.10.1	Population Generation	1560
40.10.2	In Silico Trial Design	1560
40.10.3	Intervention Window and Stochastic Resonance Applications	1561
40.11	Implementation Roadmap	1561
A	Glossary of Medical and Scientific Terms	1563
B	List of Abbreviations	1569
C	Diagnostic Tools and Assessment Scales	1571
C.1	Symptom Questionnaires	1571
C.1.1	DePaul Symptom Questionnaire (DSQ)	1571
C.1.2	Bell Disability Scale	1571
C.2	Functional Assessments	1572
C.2.1	SF-36 (Short Form Health Survey)	1572
C.2.2	Chalder Fatigue Scale	1573
C.3	Objective Tests	1573
C.3.1	Two-Day Cardiopulmonary Exercise Testing (2-Day CPET)	1573
C.3.2	Tilt Table Testing	1574
C.3.3	Autonomic Function Tests	1574
C.4	Laboratory Tests	1574
C.4.1	Standard Exclusion Panel	1575
C.4.2	Extended Panel for Clinical Characterization	1575
C.4.3	Experimental Biomarkers	1575
D	Resources and Support	1577
D.1	International Patient Organizations	1577
D.1.1	Major Global Organizations	1577
D.1.2	European Organizations	1578
D.2	National Patient Organizations by Country	1578
D.2.1	Belgium	1578
D.2.2	France	1580

Contents

D.2.3	Luxembourg	1581
D.2.4	Germany	1581
D.2.5	Netherlands	1581
D.2.6	United Kingdom	1582
D.2.7	Ireland	1582
D.2.8	Norway	1582
D.2.9	Denmark	1583
D.2.10	Sweden	1583
D.2.11	Switzerland	1583
D.2.12	Spain	1584
D.2.13	Italy	1584
D.2.14	Australia	1584
D.2.15	United States	1585
D.3	Research Centers and Specialized Clinics	1585
D.3.1	Leading Research Centers	1585
D.3.2	Specialized Clinical Centers	1586
D.4	Online Communities and Forums	1586
D.4.1	Discussion Forums	1586
D.4.2	Reddit Communities	1586
D.4.3	Facebook Groups	1587
D.4.4	Other Platforms	1587
D.5	Prominent Patient Advocates and Content Creators	1587
D.5.1	Filmmakers and Documentarians	1587
D.5.2	Writers and Bloggers	1588
D.5.3	Researchers Who Are Patients or Family Members	1589
D.5.4	Celebrity Advocates	1590
D.5.5	Euthanasia and End-of-Life Discussions	1590
D.6	Podcasts	1591
D.7	YouTube Channels and Video Resources	1592
D.7.1	Patient-Focused Channels	1592
D.7.2	Organization Channels	1592
D.7.3	Documentaries	1592
D.8	Social Media Hashtags and Campaigns	1593
D.8.1	Key Hashtags	1593
D.8.2	#MillionsMissing Campaign	1593
D.9	Books by Patients and Advocates	1593
D.9.1	Patient Memoirs	1593
D.9.2	Clinical and Reference Works	1593
D.10	Clinical Trial Registries	1594
D.11	Patient Registries and Biobanks	1594
D.12	Disability and Legal Resources	1595
D.12.1	General Guidance	1595
D.12.2	Key Considerations	1595
D.13	Resource Evaluation Guidelines	1595
D.13.1	Indicators of Reliable Resources	1595
D.13.2	Warning Signs	1596

E	Mathematical Model Details	1597
E.1	Complete Model Equations	1597
E.1.1	Energy Metabolism Model	1597
E.1.2	Energy Metabolism Parameter Table	1598
E.1.3	Immune System Model	1598
E.1.4	Neuroendocrine Model	1599
E.2	Parameter Estimation Methods	1599
E.2.1	Maximum Likelihood Estimation	1599
E.2.2	Bayesian Inference	1599
E.3	Numerical Methods	1600
E.3.1	ODE Integration	1600
E.3.2	Steady-State Computation	1600
E.4	Sensitivity Analysis Results	1600
E.5	Structural Identifiability	1601
E.6	Code Availability	1601
F	Practical Supplement Guide	1603
F.1	Quick Reference Tables	1603
F.1.1	Mitochondrial and Energy Support	1603
F.1.2	Immune Modulation	1603
F.1.3	Neurological and Sleep Support	1603
F.1.4	Gut Health	1603
F.2	Supplement Interactions	1606
F.2.1	Supplement–Supplement Interactions	1606
F.2.2	Supplement–Drug Interactions	1606
F.3	Quality and Sourcing	1606
F.4	Starting a Supplement Protocol	1607
F.5	When to Stop a Supplement	1607
G	Research Synthesis Tables	1609
G.1	Major Studies Summary	1609
G.1.1	Molecular and Cellular Mechanisms	1609
G.1.2	Viral and Infectious Triggers	1609
G.1.3	Immune Dysfunction Studies	1609
G.2	Biomarker Studies Summary	1611
G.3	Treatment Trials Summary	1611
G.3.1	Pharmacological Interventions	1611
G.3.2	Patient-Reported Interventions	1611
G.3.3	Comorbidity Management	1613
G.4	Pathophysiology Evidence Summary	1613
G.4.1	Converging Evidence for Core Mechanisms	1613
G.4.2	Patient-Derived Clinical Insights	1613
G.4.3	Research Gaps and Controversies	1613
G.4.4	Cross-Domain Medical Parallels	1613
G.5	Quick Reference: Evidence-Based Treatment Summary	1618
G.6	Selective Energy Dysfunction Hypothesis Research Materials	1620
G.6.1	Patient Assessment Tools	1620

G.6.2	Existing Dataset Analysis	1621
G.6.3	Proposed Research Studies	1621
G.6.4	Treatment Translation	1622
G.6.5	Research Community Access	1622
H	Annotated Bibliography of ME/CFS Literature	1623
H.1	Primary Research: NIH Deep Phenotyping Study	1623
H.1.1	Walitt et al. 2024 — The Foundational Deep Phenotyping Study	1623
H.2	Diagnostic Criteria and Clinical Guidelines	1624
H.2.1	Institute of Medicine (IOM) 2015 Criteria	1624
H.2.2	NICE 2021 Guidelines (NG206)	1624
H.2.3	Canadian Consensus Criteria (2003)	1624
H.2.4	Fukuda et al. 1994 (CDC Criteria)	1625
H.3	Institutional Clinical Guidelines	1625
H.3.1	MedUni Wien (2024) — Care for ME/CFS Praxisleitfaden	1625
H.3.2	MedUni Wien (2025) — Indikations-Medikamentenliste für PAIS und ME/CFS	1626
H.4	Systematic Reviews and Meta-Analyses	1627
H.4.1	Prevalence and Epidemiology	1627
H.4.2	Cognitive Impairment	1628
H.4.3	Long COVID and ME/CFS Overlap	1628
H.4.4	Sleep Abnormalities	1628
H.4.5	Evidence Mapping	1629
H.5	Pathophysiology: Immune Dysfunction	1629
H.5.1	Autoantibodies and G-Protein Coupled Receptors	1629
H.5.2	TRPM3 Ion Channel Dysfunction	1632
H.5.3	Immune Exhaustion and Chronic Activation	1632
H.5.4	Cytokine Biomarkers and Immune Signatures	1632
H.5.5	Natural Killer Cell Dysfunction	1635
H.5.6	T Cell Metabolic Dysfunction	1637
H.5.7	Neutrophil Dysfunction	1638
H.5.8	Autoantibodies: Historical Context	1638
H.5.9	Comprehensive Immune Reviews	1639
H.6	Pathophysiology: Neurological Abnormalities	1639
H.6.1	Neuroinflammation	1639
H.6.2	Neuroimaging Reviews	1640
H.6.3	Brain Energy Metabolism and the Astrocyte-Neuron Lactate Shuttle .	1640
H.6.4	Brainstem and Autonomic Dysfunction	1642
H.6.5	Auditory and Sensory Dysfunction	1646
H.6.6	Craniocervical Instability and Structural Abnormalities	1647
H.7	Pathophysiology: Metabolic and Mitochondrial Dysfunction	1649
H.7.1	Mitochondrial Dysfunction	1649
H.7.2	Metabolomics and Metabolic Traps	1654
H.8	Pathophysiology: Gut Microbiome	1655
H.9	Pathophysiology: Tryptophan and Serotonin Metabolism	1657
H.9.1	Kavyani et al. 2022 — Kynurenine Pathway Review	1657
H.9.2	Abujrais et al. 2024 — Metabolomic Analysis	1658

Contents

H.9.3	Simonato et al. 2021 — Tryptophan and Cytokines	1658
H.9.4	Lee et al. 2024 — Central Serotonin Hyperactivity	1659
H.9.5	Dehaghghi et al. 2022 — Kynurenine and NAD ⁺ Metabolism	1660
H.10	Pathophysiology: Viral Persistence and Reactivation	1661
H.10.1	Enterovirus and Chronic Persistence	1661
H.10.2	Viral Etiology Meta-Analysis	1662
H.10.3	Specific Viral Mechanisms	1663
H.11	Pathophysiology: Genetics and Epigenetics	1664
H.12	Biomarkers: Tetrahydrobiopterin (BH4) and Orthostatic Intolerance	1664
H.12.1	BH4 Elevation in ME/CFS with Orthostatic Intolerance	1664
H.13	Exercise Physiology and Post-Exertional Malaise	1666
H.14	Treatment Evidence	1668
H.14.1	Immunological Therapies: Rituximab and Cyclophosphamide	1668
H.14.2	H2 Receptor Antagonists: Cimetidine	1670
H.14.3	Low-Dose Naltrexone	1674
H.14.4	Sleep Medications: Dual Orexin Receptor Antagonists	1675
H.14.5	Mitochondrial and Metabolic Support: Amino Acids	1678
H.14.6	Antiviral Therapy: Valacyclovir and Valganciclovir	1681
H.14.7	Palmitoylethanolamide (PEA)	1683
H.14.8	D-Ribose	1684
H.14.9	Low-Dose Aripiprazole (LDA)	1685
H.14.10	Autonomic Dysfunction: Ivabradine and Pyridostigmine	1686
H.14.11	Graded Exercise Therapy (Negative Evidence)	1688
H.14.12	Neuromodulation: Transcutaneous Vagus Nerve Stimulation	1688
H.14.13	Pacing and Energy Management	1690
H.14.14	Patient-Reported Treatment Outcomes	1691
H.15	Long COVID and ME/CFS Overlap	1691
H.16	Historical Background and Epidemics	1691
H.17	Research Roadmaps and Policy Documents	1692
H.18	Comprehensive Reviews	1692
H.19	Mast Cell Activation and Antihistamine Therapies	1692
H.19.1	Hardcastle et al. 2016 — Mast Cell Phenotype Abnormalities in ME/CFS	1692
H.19.2	Wirth & Scheibenbogen 2023 — Mast Cell Activation and Vascular Pathomechanisms	1693
H.19.3	Novak et al. 2022 — Mast Cell Disorders, Cerebral Hypoperfusion, and Small Fiber Neuropathy	1694
H.19.4	Magadmi et al. 2019 — CADM1-Mediated Mast Cell-Nerve Adhesion Amplifies Inflammation	1694
H.19.5	Nakamura et al. 2014 — Circadian Clock Regulates Mast Cell Function	1695
H.19.6	Steinberg et al. 1996 — Terfenadine Trial (Negative)	1695
H.19.7	Davis et al. 2023 — Long COVID Case with H1/H2 Combination Success	1696
H.19.8	Theoharides et al. 2012 — Quercetin Superior to Cromolyn	1696
H.19.9	Clemons et al. 2011 — Amitriptyline Mast Cell Inhibition	1697
H.19.10	Rupatadine — Dual H1/PAF Antagonist with Mast Cell Stabilization .	1698
H.19.11	Moldofsky et al. 2015 — Ketotifen in Fibromyalgia (Negative)	1698
H.20	Blood Volume and Cardiovascular Dysfunction	1699
H.20.1	Hypovolemia and RAAS Dysfunction	1699

Contents

H.20.2	Cardiac Dysfunction and Natriuretic Peptides	1700
H.20.3	Endothelial Dysfunction and Vascular Pathology	1700
H.20.4	Erythropoiesis and Red Blood Cell Function	1701
H.20.5	Integrated Mechanisms: The Hypovolemia Cascade	1702
H.21	Additional Key Resources	1703
H.21.1	Patient Advocacy and Information	1703
H.21.2	Research Centers	1703
H.22	Neuroimaging and Structural Brain Findings	1703
H.22.1	Shan et al. 2020 — Neuroimaging Systematic Review (JTM)	1703
H.22.2	Finkelmeyer et al. 2017 — VBM Gray and White Matter in CFS	1704
H.22.3	Puri et al. 2012 — VBM Regional Gray/White Matter Changes	1704
H.22.4	de Lange et al. 2004 — fMRI Neural Correlates of CFS	1705
H.23	Neurotransmitter and Neurochemical Findings	1705
H.23.1	Yamamoto et al. 2004 — Serotonin Transporter Reduction in CFS	1705
H.23.2	Miller et al. 2014 — Basal Ganglia Activation and Dopamine in CFS	1705
H.23.3	Dehghani et al. 2022 — Kynurenine Pathway and NAD ⁺ in ME/CFS	1706
H.24	Prognosis and Outcomes	1706
H.24.1	Joyce, Hotopf and Wessely 1997 — Prognosis of CFS: Systematic Review	1706
H.24.2	Cairns and Hotopf 2005 — Adult CFS Prognosis: Systematic Review	1707
H.25	Orthostatic Intolerance	1707
H.25.1	Schondorf and Freeman 1999 — Importance of OI in CFS	1707
H.26	Treatment Evidence	1707
H.26.1	Li et al. 2022 — tDCS of DLPFC for Neuropsychiatric Disorders	1707
H.26.2	Lang-Illievich et al. 2023 — PEA for Chronic Pain: Systematic Review and Meta-Analysis	1708
H.27	Coagulation and Vascular Pathology	1708
H.27.1	Nunes et al. 2022 — Hyperactivated Platelets and Fibrinoid Micro- clots in ME/CFS	1708
H.27.2	Suárez et al. 2010 — Nitric Oxide Metabolites During Exercise in CFS	1709
H.28	Lifestyle Intervention Safety	1709
H.28.1	Kindlon 2011 — Reporting of Harms from GET and CBT in ME/CFS	1709
H.29	Adenosine, Sleep Pressure, and Neuroinflammation	1710
H.29.1	Huang et al. 2024 — Adenosine Receptors in Sleep Regulation	1710
H.29.2	Retéy et al. 2007 — ADORA2A Genotype and Caffeine Sensitivity	1710
H.29.3	Orr et al. 2009 — A2A Receptor Mediates Microglial Process Retraction	1711
H.29.4	Navia et al. 2020 — Adenosine Receptors as Neuroinflammation Mod- ulators	1711
H.29.5	Chang et al. 2021 — Dysregulated Adenosine Homeostasis in Brain Disorders	1712
H.29.6	Rábago-Monzón et al. 2025 — Astrocytes, Microglia, and Sleep Dys- regulation	1712
H.29.7	Nakatomi et al. 2014 — TSPO-PET Neuroinflammation in ME/CFS	1713
H.29.8	Maksoud et al. 2021 — Systematic Review of Sleep in ME/CFS	1713
H.30	EBV, Infectious Triggers, and Autoimmune Mechanisms	1724
H.30.1	Bu et al. 2024 — EBV as a Potentiator of Autoimmune Diseases	1724
H.30.2	Munir et al. 2025 — EBV Molecular Mimicry in Autoimmune Disease	1724
H.30.3	Pless et al. 2026 — EBV LMP1 and CNS Demyelination	1725

H.31	Developmental Immunology and Peripheral Tolerance	1726
H.31.1	Mohtashami & Zúñiga-Pflücker 2024 — Thymic Homeostasis and Immune Tolerance	1726
H.31.2	Meffre & O'Connor 2024 — B Cell Tolerance and Autoimmune Repertoires	1727
H.31.3	Wirth & Scheibenbogen 2025 — Neurotransmitter Imbalance in ME/CFS and Long COVID	1727
H.32	Gut-Brain Axis and Enteroendocrine Signalling	1728
H.32.1	Kaelberer et al. 2018 — Direct Gut-Brain Neural Circuit for Nutrient Sensing	1728
H.32.2	Barton et al. 2023 — Enteroendocrine Cell Regulation of the Gut-Brain Axis	1729
H.32.3	Barton et al. 2025 — Vagus Nerve and Serotonin in the Gut-Brain Axis	1730
H.32.4	Hsu et al. 2025 — Gut Microbiome in ME/CFS: Systematic Review	1730
H.33	Adolescent Neurodevelopment and Microglial Sensitive Periods	1731
H.33.1	Dziabis & Bilbo 2022 — Microglia and Sensitive Periods in Brain Development	1731
H.33.2	Chung et al. 2022 — Adolescence as a Sensitive Period for Prefrontal Microglia	1732
H.33.3	VanRyzin et al. 2025 — Adolescent Synaptic Pruning Shapes Adult Behaviour	1732
H.33.4	Rahimian et al. 2020 — Microglial Depletion and Repopulation	1733
H.34	Hematopoietic Stem Cells, Aging, and Chronic Inflammation	1734
H.34.1	Takizawa et al. 2020 — HSC Protection from Stress-Induced Exhaustion	1734
H.34.2	Pietras et al. 2021 — Inflammation as a Regulator of HSC Function	1735
H.34.3	Morrison & Scadden 2025 — The Aging Hematopoietic Stem Cell Niche	1735
H.34.4	Rossi et al. 2025 — Haematopoietic Ageing in Health and Lifespan	1736
H.35	Pediatric ME/CFS: Diagnosis, Management, and Prognosis	1737
H.35.1	Rowe et al. 2017 — Pediatric ME/CFS: A Comprehensive Primer	1737
H.35.2	Rowe 2019 — Long-Term Follow-Up of Young People with CFS	1738
H.35.3	van Campen et al. 2020 — Severity Validation by Objective Measures	1738
H.35.4	Heng et al. 2025 — Mapping the Complexity of ME/CFS	1739
H.36	Diagnostic Criteria, Treatment Evidence, and Mitochondrial Biology	1740
H.36.1	Carruthers et al. 2011 — International Consensus Criteria (ICC)	1740
H.36.2	Wilshire et al. 2018 — Reanalysis of the PACE Trial	1741
H.36.3	Lawless et al. 2015 — The Rise and Rise of Mitochondrial DNA Mutations	1741
H.36.4	Comparative Virology: Coronavirus Persistence in Immune Cells	1742
H.36.5	Exercise Pathophysiology: Deconditioning Myth	1743
H.37	Diagnostic Criteria and Case Definitions	1744
H.37.1	Brurberg et al. 2014 — Case Definitions for CFS/ME: Systematic Review	1744
H.37.2	Carruthers et al. 2011 — International Consensus Criteria	1744
H.37.3	NICE 2021 — ME/CFS Diagnosis and Management Guideline (NG206)	1745
H.38	Epidemiology, Quality of Life, and Mortality	1745
H.38.1	Lim et al. 2020 — Prevalence Meta-Analysis	1745
H.38.2	Hvidberg et al. 2015 — Health-Related Quality of Life in ME/CFS	1745
H.38.3	Jason et al. 2006 — Causes of Death in CFS	1746
H.38.4	Roberts et al. 2016 — CFS Mortality in England and Wales	1746

Contents

H.38.5	Kingdon et al. 2018 — Functional Status Comparison with MS	1746
H.39	NIH Deep Phenotyping and Multimodal Studies	1747
H.39.1	Walitt et al. 2024 — Deep Phenotyping Study	1747
H.39.2	Heng et al. 2025 — Mapping the Complexity of ME/CFS	1747
H.39.3	DecodeME 2025 — ME/CFS Genome-Wide Association Study	1747
H.40	Energy Metabolism and Mitochondrial Dysfunction	1747
H.40.1	Naviaux 2014 — Cell Danger Response	1747
H.40.2	Naviaux et al. 2016 — Metabolic Features of CFS	1748
H.40.3	Germain et al. 2020 — Circulatory Metabolomics in ME/CFS	1748
H.40.4	CastroMarrero et al. 2021 — CoQ10 + NADH RCT	1748
H.40.5	Syed et al. 2025 — NAD ⁺ Therapy Pilot Study	1749
H.40.6	Newton et al. 2007 — Fatigue Comparison with Sjögren's	1749
H.41	Exercise Physiology and Post-Exertional Malaise	1749
H.41.1	Chu et al. 2019 — Deconstructing Post-Exertional Malaise	1749
H.41.2	Keller et al. 2024 — Two-Day CPET in ME/CFS	1749
H.41.3	Jason et al. 2009 — Energy Modulation and Physical Functioning . . .	1750
H.41.4	Jason et al. 2012 — Energy Envelope Theory Interventions	1750
H.42	Immune Dysfunction and Autoimmunity	1750
H.42.1	Loebel et al. 2016 — Autoantibodies to Adrenergic and Muscarinic Receptors	1750
H.42.2	Bynke et al. 2020 — GPCR Autoantibodies: Swedish Validation	1751
H.42.3	Sotzny et al. 2021 — GPCR Autoantibodies Correlate with Symptom Severity	1751
H.42.4	Fluge et al. 2011 — Rituximab Phase II RCT	1751
H.42.5	Fluge et al. 2015 — Rituximab Maintenance RCT	1752
H.42.6	Fluge et al. 2019 — RituxME: Definitive Negative Trial	1752
H.42.7	Hardcastle et al. 2016 — Mast Cell Phenotypes in ME/CFS	1752
H.42.8	Frioni et al. 2024 — Mast Cell Activation Syndrome: Systematic Review	1752
H.42.9	Folkerts et al. 2020 — Butyrate Inhibits Mast Cell Activation	1753
H.43	Neurological and Neuroendocrine Dysfunction	1753
H.43.1	Dantzer 2007 — Cytokine-Induced Sickness Behaviour	1753
H.43.2	Raijmakers et al. 2021 — Neuroinflammation and Microglia	1753
H.43.3	Novak et al. 2022 — Cerebral Blood Flow Regulation Review	1754
H.43.4	Marty et al. 2008 — Nucleus Tractus Solitarius	1754
H.43.5	Cleare et al. 1999 — Low-Dose Hydrocortisone RCT	1754
H.43.6	Geraghty & Blease 2019 — Cognitive Dysfunction Review	1754
H.43.7	Huerta et al. 2025 — Vagus Nerve Stimulation in CFS	1755
H.44	Tryptophan, Kynurenine, and NAD ⁺ Metabolism	1755
H.44.1	Kavyani et al. 2022 — Kynurenine Pathway Metabolites: Systematic Review	1755
H.44.2	Dehhaghi et al. 2022 — Kynurenine Pathway and Clinical Symptoms .	1755
H.45	Circadian Rhythm and Sleep	1756
H.45.1	Cambras et al. 2018 — Circadian Rhythm Abnormalities in ME/CFS .	1756
H.45.2	Castro-Marrero et al. 2021 — Melatonin + Zinc RCT	1756
H.45.3	Xie et al. 2013 — Sleep Drives Glymphatic Clearance	1756
H.45.4	Nemat-Gorgani et al. 2025 — Glymphatic System Dysfunction Review	1756

H.46	Gut Microbiome and Multi-Omics	1757
H.46.1	Xiong et al. 2023 — Multi-Omics of Gut Microbiome in ME/CFS . . .	1757
H.46.2	Hsu et al. 2025 — Gut Microbiome in ME/CFS: Systematic Review . .	1757
H.47	Epigenetics and DNA Methylation	1757
H.47.1	de Vega et al. 2021 — DNA Methylation Meta-Analysis	1757
H.48	Viral Persistence and EBV Mechanisms	1758
H.48.1	Pless et al. 2026 — EBV LMP1 and CNS Demyelination	1758
H.49	Cardiovascular and Autonomic Dysfunction	1758
H.49.1	Raj et al. 2005 — Pyridostigmine RCT in CFS	1758
H.50	Treatment: Pharmacological Interventions	1758
H.50.1	Polo et al. 2019 — Low Dose Naltrexone for Fibromyalgia	1758
H.50.2	White et al. 2011 — PACE Trial	1759
I	Ongoing and Planned ME/CFS Research Studies	1760
I.1	Multi-Omics and Systems Biology	1760
I.1.1	Rosetta Stone Study	1760
I.1.2	BioQuest Study	1761
I.2	Genomics and Epigenetics	1762
I.2.1	DecodeME	1762
I.2.2	EpiSwitch CFS Epigenetic Diagnostic	1763
I.2.3	PrecisionLife Combinatorial Genomics	1763
I.3	Biomarker Discovery and Diagnostics	1764
I.3.1	BioMapAI (Gut Microbiome and Metabolomics)	1764
I.3.2	ME Association PhD Project (Metabolite and Infection Markers)	1765
I.3.3	MEA–UCL Blood Biomarker Pilot	1765
I.3.4	Cornell Cell-Free RNA Liquid Biopsy	1766
I.3.5	Cell Reports Medicine Multi-Modal Study	1767
I.4	Autoimmunity and Immunomodulation	1767
I.4.1	Charité Immunoabsorption Trial (IA-PACS-CFS)	1767
I.4.2	BC007 (Rovunaptabin) Phase II Trials	1768
I.4.3	ResetME Trial (Daratumumab)	1769
I.5	Neuroimaging and Ion Channel Research	1770
I.5.1	NCNED TRPM3 Ion Channel Programme	1770
I.5.2	Charité Neuroimaging Biomarker Study	1771
I.5.3	Hyperbaric Oxygen Therapy for ME/CFS	1771
I.6	Large-Scale Federally Funded Programmes	1772
I.6.1	RECOVER-TLC Clinical Trials Programme	1772
I.7	Clinical Trials (Interventional)	1773
I.7.1	LIFT Trial (Low-Dose Naltrexone and Pyridostigmine)	1773
I.7.2	UAB LDN Dose-Finding Study	1774
I.7.3	Low-Dose Rapamycin in ME/CFS and Long Covid	1774
I.7.4	Lumbrokinase for ME/CFS and Long Covid	1775
I.7.5	Hydrogen Water for ME/CFS	1776
I.7.6	RESTORE-ME: Oxaloacetate for ME/CFS Fatigue	1776
I.7.7	Stellate Ganglion Block for ME/CFS	1777
I.7.8	Transcutaneous Vagus Nerve Stimulation (tvNS) for Long Covid/ME/CFS	1778

Contents

I.7.9	Masitinib for Severe Mast Cell Activation Syndrome	1778
I.7.10	Cornell NAC–Glutathione Brain Study	1779
I.7.11	NSU Directed Probiotics for ME/CFS	1780
I.7.12	ADDRESS-LC (Bezisterim for Long Covid Neurocognition)	1780
I.7.13	LC-REVITALIZE (Upadacitinib + Pirfenidone)	1781
I.7.14	Viral Persistence Trials (Paxlovid/STOP-PASC)	1782
I.7.15	CHIIME (Chronic Infections and Inflammation in ME/CFS)	1782
I.8	Virology and Post-Infectious Mechanisms	1783
I.8.1	Herpesvirus Reactivation and Anti-dUTPase Antibodies	1783
I.8.2	Post-IM Longitudinal Cohort: Seven-Year Follow-Up	1784
I.8.3	RECOVER ME/CFS Incidence Study	1784
I.9	Immune Profiling and T Cell Research	1785
I.9.1	CD8+ T Cell Exhaustion Atlas	1785
I.9.2	Single-Cell Immune Remodelling in Long COVID–ME/CFS	1786
I.9.3	IgG Immune Complexes Disrupting Cellular Energetics	1786
I.10	Cardiovascular, Autonomic, and Vascular Research	1787
I.10.1	MCAM Multi-Site Autonomic Dysfunction Study	1787
I.10.2	Small Fibre Neuropathy in ME/CFS and Post-COVID	1788
I.10.3	Microvascular Remodelling and Endothelial Dysfunction	1788
I.10.4	POTS Autonomic Profiling Across Aetiologies	1789
I.11	Exercise Physiology and Muscle Biology	1789
I.11.1	Largest Two-Day CPET Study	1789
I.11.2	Skeletal Muscle Properties in Long COVID and ME/CFS	1790
I.12	Microbiome and Gut–Brain Axis	1791
I.12.1	JAX AI-Integrated Gut Microbiome Diagnostic	1791
I.12.2	Gut Microbial Composition Differences in CFS	1791
I.13	Endocrine and Metabolic Profiling	1792
I.13.1	Sex-Specific Steroid Hormone Profiling	1792
I.14	Kynurenine Pathway	1792
I.14.1	OMF Kynurenine Supplementation Trial	1792
I.15	Comorbid Conditions: EDS–MCAS–ME/CFS Triad	1793
I.15.1	EDS–MCAS Genetic and Pathophysiological Basis	1793
I.16	Conferences and Funding Initiatives	1794
I.16.1	International ME/CFS Conference 2026 (Berlin)	1794
I.16.2	ME/CFS Research Foundation 2026 Funding Strategy	1794
I.17	Digital Health and Wearable Technology	1794
I.17.1	Long Covid Wearable Device Study	1794
I.18	Results Expected in 2026–2027	1795
I.19	Registry Summary	1797

Bibliography

1799

List of Figures

6.1	Normal cellular energy production pathway.	230
6.2	ME/CFS energy production dysfunction and systemic consequences.	231
6.3	Normal oxidative stress homeostasis with balanced ROS production and neutralization.	235
6.4	ME/CFS oxidative stress vicious cycle with self-perpetuating damage.	236
6.5	Normal exercise response with adaptive signaling and rapid recovery.	240
6.6	ME/CFS post-exertional malaise mechanism with harmful vicious cycle.	241
7.1	Normal immune response with appropriate activation and resolution.	282
7.2	ME/CFS immune dysfunction with chronic inflammation and exhaustion cycles.	283
8.1	Normal catecholamine synthesis pathway from tyrosine to norepinephrine and epinephrine.	354
8.2	ME/CFS catecholamine synthesis failure and systemic consequences.	355
8.3	Normal tryptophan metabolism with balanced serotonin and kynurenine pathways.	356
8.4	ME/CFS tryptophan dysregulation causing serotonin deficit and quinolinic acid toxicity.	357
8.5	Normal cerebral blood flow regulation meeting brain metabolic demands.	391
8.6	ME/CFS cerebral hypoperfusion cascade causing cognitive dysfunction.	392
9.1	Normal HPA axis stress response with negative feedback control.	417
9.2	ME/CFS HPA axis dysregulation with blunted response and systemic consequences.	418
14.1	Causal diagram of selective energy dysfunction in ME/CFS. Edge weights indicate certainty (0–1 scale) based on evidence quality. The diagram includes a feedback edge (exercise intolerance → CNS crisis, certainty 0.35), making it a causal diagram rather than a strict DAG. Preserved processes (green) have sub-threshold CNS-dependency ($\delta_{CNS} \leq 0.2$, $\tilde{\rho} \leq 0.2$, giving $S \leq 0.24$) and no clinically significant causal path from the CNS crisis node.	703
14.2	Astrocyte-neuron lactate shuttle (ANLS) as proposed mechanism for CNS-specific energy dysfunction. If MCT2 transporter function is impaired, neurons cannot access lactate-derived ATP while peripheral tissues (with direct glucose access) remain unaffected.	704
14.3	CNS energy triage hierarchy under energy scarcity. The red dashed line indicates typical available energy in ME/CFS. Functions above the line (Tiers 4–6) are impaired; functions below (Tiers 1–3) are preferentially preserved through energy allocation.	706

List of Figures

14.4	Four-compartment energy model showing CNS as the coordination bottleneck in ME/CFS. Compartments are classified by CNS-dependency index (δ_{CNS}). Secondary dysfunction in autonomic and peripheral compartments results from impaired CNS coordination, while autonomous processes with $\delta_{CNS} \leq 0.2$ remain preserved.	714
24.1	Vicious cycle network model of ME/CFS pathophysiology	1133
34.1	Overview of the ME/CFS modeling framework. Modeling approaches (top) are applied to physiological subsystems (middle), which couple into integrated and predictive models (bottom). Dashed arrows indicate cross-system couplings.	1435
35.1	Integrated energy metabolism model showing glycolysis, Krebs cycle, and electron transport chain with ME/CFS impairment points (red) and the ROS-mediated damage feedback loop. Equation references correspond to the mathematical formulation in the text.	1449
36.1	Immune system model components. (a) State transition diagrams for NK cells (left, Equation 36.1) and T cells (right, Equation 36.8). In ME/CFS, accelerated exhaustion (k_{exh} increased) and impaired recovery (k_{recov} reduced) shift populations toward exhausted states (red). (b) Cytokine network topology showing pro-inflammatory (warm colors) and anti-inflammatory (cool colors) mediators with stimulatory (arrows) and inhibitory (flat-headed) interactions.	1467
37.1	HPA axis feedback model (Equation 37.1). The hypothalamus–pituitary–adrenal cascade produces cortisol under circadian and stress-driven regulation with negative feedback at hypothalamic and pituitary levels. In ME/CFS (red annotations), enhanced feedback sensitivity, blunted circadian amplitude, and reduced stress responsiveness produce low-normal cortisol that inadequately restrains immune activation.	1485
37.2	Baroreflex control loop model (Equations 37.4–37.9). Baroreceptors detect MAP changes and modulate sympathetic (S) and parasympathetic (V) tone to maintain cardiovascular homeostasis. Orthostatic stress (standing) reduces MAP, triggering compensatory responses. Three ME/CFS-associated POTS subtypes correspond to distinct parameter modifications in the model.	1488
37.3	Tryptophan metabolic branching (Equations 37.10–37.13). Tryptophan is metabolized through competing pathways: serotonin synthesis (left, blue) via tryptophan hydroxylase (TPH) and kynurenine production (right, red) via indoleamine 2,3-dioxygenase (IDO). IFN- γ from immune activation upregulates IDO, diverting tryptophan toward neurotoxic kynurenine metabolites and depleting serotonin. BH ₄ cofactor competition further links this pathway to catecholamine synthesis.	1491
38.1	Architecture of the integrated whole-body model showing subsystem modules and their couplings. The central energy–immune vicious cycle (red) drives disease persistence. Arrows indicate the dominant coupling direction; most interactions are bidirectional. The symptom generation module (bottom) maps physiological states to clinical symptoms.	1506

39.1	Temporal dynamics of ME/CFS. (a) Phase portrait of the energy-immune system showing bistability: healthy (green) and disease (red) attractors separated by a separatrix (dashed). An acute infection can push the system across the separatrix into the disease basin. (b) Long-term disease trajectories predicted by the damage accumulation model (Equation 39.1), with approximate population frequencies from meta-analysis [113].	1525
39.2	The energy ratchet model. (a) A single patient trajectory (accelerating ratchet, $\alpha > 0$) showing baseline functional capacity $B(t)$ (solid blue) and the irreversible ceiling $B_{\max}$ (dashed red) declining through four damaging events (infections, PEM crashes, surgery). Each event produces a total acute drop Δ_k (blue arrow), of which only δ_k (red arrow) is irreversible; note the increasing step sizes reflecting damage sensitisation. Recovery between events is partial and ATP-dependent (Equation 39.2). Horizontal bands indicate clinical severity levels (Equation 39.7). (b) Two of four trajectory classes: <i>stable</i> ratchet with equal ceiling steps ($\alpha \approx 0$) and <i>accelerating</i> ratchet ($\alpha > 0$). (c) <i>Incomplete-recovery cascade</i> (frequent events prevent recovery) and <i>arrested ratchet</i> (event prevention preserves ceiling). See Equations 39.3–39.4 and 39.5 for the formal jump map.	1529
40.1	Clinical translation pipeline from patient data to treatment recommendations. Patient biomarkers (Stage 1) are used to estimate individual model parameters (Stage 2), which enable simulation of candidate interventions (Stage 3) and generation of personalized treatment recommendations (Stage 4). The implementation roadmap (bottom) shows the staged path from current research tools to validated clinical instruments.	1547

List of Tables

4.1	Symptom severity by diagnostic criteria met	95
4.2	Tier 2 test panels for Viral-Immune-Metabolic phenotype	126
4.3	Treatment prioritization by biological domain	133
4.4	Comparison of diagnostic frameworks	136
5.1	Objective measures across ME/CFS severity levels	152
5.2	Metabolic phenotypes in ME/CFS	204
5.3	Characteristics of patients meeting different diagnostic criteria	206
5.4	Comprehensive endotype-to-biomarker-to-treatment mapping in ME/CFS. Evidence: A = RCT data; B = pilot/open-label; C = mechanistic rationale only. Reproducibility: R = replicated in ≥ 2 independent cohorts; P = preliminary (single study or unreplicated).	210
5.5	Tiered prospective phenotyping protocol for ME/CFS treatment selection . . .	212
7.1	Prioritized research directions for severe ME/CFS	345
11.1	SIBO Subtypes and Clinical Presentations	488
13.1	Evidence Level Definitions	526
13.2	Ranked Hypotheses of ME/CFS Pathophysiology	528
13.3	Hypothesized Triggers for Cycle Recruitment	550
13.4	Treatment Implications by Hypothesis	563
13.5	Cycle Stage Comparison: Long COVID vs. Established ME/CFS (Hypothesized)	570
13.6	Proposed Computational Model Components	577
13.7	Genetic Variants Potentially Affecting Cycle Dynamics	579
13.8	Candidate Pediatric Protection Mechanisms and Adult Recreation Strategies (Hypothesized)	581
14.1	Comprehensive ranking of all speculative hypotheses by evidence level, therapeutic potential, and impact on different severity levels	586
14.2	Evidence assessment for GPCR autoantibody findings in ME/CFS	617
14.3	Novel hypotheses arising from two-day CPET findings, ranked by plausibility and therapeutic potential	662
14.4	Novel hypotheses from 2026 autoimmune research	671
14.5	Process classification by CNS-dependency and demand-responsiveness with predicted and observed dysfunction in ME/CFS.	701
17.1	Expected symptom improvement at 2 weeks with full protocol	823
18.1	ME/CFS Severity Stratification and Care Framework	853
18.2	Recommended Monitoring Frequency by ME/CFS Severity	853

List of Tables

18.3	Crash Severity Classification with Dose-Response Predictions	860
18.4	Comparison of GET vs. Sports Medicine-Adapted Pacing	873
18.5	Example: Cycle Status Dashboard for Individual Patient	916
18.6	Cost-Effectiveness Ranking of Cycle Diagnostic Tests	917
22.1	Mitochondrial supplement dosing by age group	1062
24.1	Craniocervical Instability Radiographic Measurements	1118
24.2	Highly Reversible Pathological Mechanisms	1156
24.3	Partially Reversible Pathological Mechanisms	1157
24.4	Irreversible or Minimally Reversible Mechanisms	1158
24.5	Treatment Intensity by Reversibility Tier	1160
24.6	Proposed Cycle Gain Classification and Prognosis (Hypothetical—Requires Val- idation)	1165
24.7	Tiered Epigenetic Reversal Strategies	1166
25.1	Micro-dosing reference: ME/CFS starting doses versus standard doses	1173
25.2	High-Altitude vs. ME/CFS Hypoxia	1195
25.3	PICS vs. ME/CFS Mechanisms	1198
25.4	Microgravity vs. ME/CFS Deconditioning	1201
25.5	Cross-Domain Intervention Summary	1207
25.6	Novel Mechanistic Hypotheses and Therapeutic Opportunities	1215
25.7	Risk-Benefit Assessment of Novel Interventions	1216
27.1	Major negative, null, and inconclusive trials in ME/CFS.	1248
29.1	Suicide Mortality Across ME/CFS Studies	1274
31.1	ME/CFS Mechanisms Documented in Long COVID	1299
31.2	ME/CFS Treatment Protocols Applicable to EDS	1305
31.3	Research Priorities: Mechanisms × Conditions	1325
33.1	Coverage of core predictions by existing ME/CFS datasets	1365
33.2	Cross-document hypothesis and open question registry (42 key entries). “Cert.” = certainty on the 0–1 scale used throughout the document; “inf.” = inferred from context language.	1416
34.1	Mapping of pathophysiological domains (Part II) to mathematical models (Part V).1446	
40.1	Model-predicted treatment candidates ranked by global sensitivity index. <i>Model-unique</i> column indicates whether the prediction requires the mathemat- ical model or could be derived from qualitative reasoning alone.	1558
E.1	Energy metabolism model parameters.	1598
E.2	Selected immune model parameters.	1598
E.3	Selected neuroendocrine model parameters.	1599
F.1	Mitochondrial and energy support supplements.	1604
F.2	Immune modulation supplements.	1604

List of Tables

F.3	Neurological and sleep support supplements.	1605
F.4	Gut health supplements.	1605
G.1	Molecular Mechanism Studies in ME/CFS	1609
G.2	Viral Association Studies	1610
G.3	Immune System Studies in ME/CFS	1610
G.4	Validated and Proposed Biomarkers for ME/CFS	1611
G.5	Pharmacological Treatment Evidence in ME/CFS	1611
G.6	Patient-Reported Interventions Requiring Clinical Validation	1612
G.7	Comorbidities Frequently Misdiagnosed as ME/CFS	1613
G.8	Evidence Strength for Proposed Pathophysiological Mechanisms	1614
G.9	Patient-Discovered Patterns and Clinical Rules	1615
G.10	Major Research Gaps in ME/CFS	1616
G.11	Cross-Domain Medical Interventions Applicable to ME/CFS	1617
G.12	Evidence-Based Treatment Quick Reference	1619
I.1	Studies with results expected in 2026–2027, ranked by anticipated impact. . . .	1796
I.2	Summary of ongoing and planned ME/CFS research studies.	1798

Part I

Clinical Overview

This part provides a comprehensive clinical picture of myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). We cover the full spectrum of symptoms, diagnostic criteria from multiple frameworks, disease progression patterns, and clinical presentations.

Understanding the clinical manifestations is essential for accurate diagnosis, effective communication between patients and healthcare providers, and appropriate disease management.

1 Introduction to ME/CFS

Myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) is a complex, chronic, multi-system disease characterized by severe and disabling fatigue, post-exertional malaise, unrefreshing sleep, cognitive dysfunction, and autonomic dysregulation. This document provides a comprehensive overview of current understanding, research, and clinical approaches to ME/CFS.

1.1 Overview and Terminology

ME/CFS has been recognized as a distinct clinical entity by major health organizations, including the World Health Organization (ICD-11 code 8E49), the Centers for Disease Control and Prevention, and the National Institutes of Health. The condition affects an estimated 0.89% to 2.5% of the global population, with significant variation based on diagnostic criteria used.

The terminology surrounding this condition has evolved over time. While “chronic fatigue syndrome” became widely used in the late 1980s, many patient advocates and researchers prefer “myalgic encephalomyelitis” as it better reflects the neurological and immunological aspects of the disease. This document uses the combined term ME/CFS to acknowledge both naming conventions.

1.2 Historical Context

The recognition of ME/CFS as a distinct clinical entity has followed a complex trajectory spanning nearly a century, marked by periods of intense research, prolonged neglect, and ongoing controversy regarding the nature of the illness.

1.2.1 Key Outbreaks and Case Clusters

ME/CFS was first recognized through epidemic outbreaks affecting medical personnel and communities, initially misdiagnosed as atypical poliomyelitis.

Los Angeles County Hospital (1934). The first documented outbreak occurred at Los Angeles County General Hospital from May 1934 to December 1935, affecting 198 hospital employees (4.5% of personnel), including 10.7% of nurses and 5.4% of physicians [21]. Initially diagnosed as atypical poliomyelitis, subsequent analysis revealed a distinct clinical pattern with prominent neurological symptoms and prolonged post-infectious disability.

Iceland/Akureyri (1948–1949). An outbreak in Akureyri, Iceland affected 488 patients locally and 1,090 cases across the country over three months, establishing “Icelandic disease” as an early term for what would later be recognized as ME [22]. The illness shared features with poliomyelitis but demonstrated distinct characteristics including prolonged fatigue and neurological sequelae.

Royal Free Hospital, London (1955). The most thoroughly documented outbreak occurred at the Royal Free Hospital from July to November 1955, affecting 292 staff members (255 hospitalized) [23]. This outbreak led to the term “benign myalgic encephalomyelitis” being coined in a 1956 *Lancet* editorial. Dr. Melvin Ramsay, head of Infectious Diseases at Royal Free, became a lifelong advocate and developed the first clinical criteria for ME, emphasizing muscle fatigability with prolonged recovery and neurological dysfunction [24]. The term “benign” was later abandoned as the chronic, disabling nature of the illness became apparent.

Lake Tahoe/Incline Village (1984–1987). An outbreak in the Lake Tahoe region documented by physicians Paul Cheney and Daniel Peterson affected an estimated 259 patients, with 160 residents of Incline Village affected by winter 1985 [25]. The Centers for Disease Control and Prevention (CDC) investigation found elevated Epstein-Barr virus (EBV) antibodies but concluded there was insufficient evidence for an EBV-specific epidemic. This investigation led to the coining of “chronic fatigue syndrome” by the CDC in 1988 [26], replacing the earlier term “chronic Epstein-Barr virus syndrome” after research failed to demonstrate a consistent EBV link.

1.2.2 Evolution of Diagnostic Criteria

Diagnostic criteria for ME/CFS have evolved substantially, with increasing recognition of post-exertional malaise as the cardinal feature.

Holmes/CDC Criteria (1988). The first formalized definition required new-onset debilitating fatigue lasting at least six months that was not resolved by bed rest and reduced activity by at least 50%, plus 6 of 11 symptom criteria and 2 of 3 physical criteria, or 8 of 11 symptom criteria [26]. This established the six-month duration threshold still used today.

Fukuda Criteria (1994). Developed by the International Chronic Fatigue Syndrome Study Group, these criteria required six or more months of chronic fatigue of new or definite onset, not substantially alleviated by rest, causing substantial reduction in activities, plus four of eight specific symptoms: unrefreshing sleep, post-exertional malaise, myalgia, arthralgia, new headaches, sore throat, tender lymphadenopathy, and impaired memory or concentration [27]. The Fukuda criteria became the most widely used research standard for two decades, though they were limited by not requiring PEM.

Canadian Consensus Criteria (2003). The Canadian Consensus Criteria represented a paradigm shift by requiring post-exertional malaise as a mandatory criterion, described as “pathologically slow recovery, usually 24 hours or longer” [28]. These criteria also required sleep dysfunction, pain, and symptoms from multiple categories including neurological, cognitive, autonomic, neuroendocrine, and immune manifestations. Studies demonstrate that patients meeting Canadian Consensus Criteria have more severe presentations and greater functional impairment than those meeting Fukuda criteria alone.

International Consensus Criteria (2011). An international panel of 26 experts from 13 countries achieved 100% consensus via Delphi methodology on criteria that emphasized “myalgic encephalomyelitis” terminology and required symptoms from four domains: post-exertional neuroimmune exhaustion, neurological impairment, immune/gastrointestinal/-genitourinary impairments, and energy production/transportation impairments [29]. These criteria identify a more homogeneous patient population with greater functional impairments.

Institute of Medicine Criteria (2015). The Institute of Medicine (now National Academy of Medicine) proposed simplified diagnostic criteria and the name “Systemic Exertion Intolerance Disease” (SEID), though this name was not widely adopted [6]. The IOM criteria require three symptoms present at least 50% of the time with moderate, substantial, or severe intensity: substantial reduction in pre-illness activities, unrefreshing sleep, and post-exertional malaise, plus either cognitive impairment or orthostatic intolerance. Critically, this report declared ME/CFS “a serious, chronic, complex systemic disease” requiring proper medical recognition.

1.2.3 Changes in Medical Understanding

The medical understanding of ME/CFS has undergone dramatic shifts, moving from psychogenic theories toward recognition as a biological disease.

The Psychogenic Era. In 1970, psychiatrists McEvedy and Beard published analyses of 15 ME outbreaks (including Royal Free 1955) in the *British Medical Journal*, concluding they represented “mass hysteria” based partly on the preponderance of female patients and institutional settings [30]. This analysis was conducted without examining patients or consulting treating physicians. The publications received prominent media coverage and contributed to decades of research funding drought and medical dismissal. Subsequent mathematical modeling of outbreak data has demonstrated that the epidemiological patterns fit infectious disease models, mathematically refuting the hysteria hypothesis [31].

Return to Biological Investigation. The Lake Tahoe outbreak reinvigorated biological research, leading to investigations of viral triggers, immune dysfunction, and neurological abnormalities through the 1990s and 2000s. The Canadian Consensus Criteria (2003) and

International Consensus Criteria (2011) explicitly framed ME/CFS as a neuroimmune disease with objective physiological abnormalities.

The 2015 IOM Report. The Institute of Medicine’s comprehensive review concluded that “ME/CFS is a serious, chronic, complex systemic disease that often can profoundly affect the lives of patients” [6]. The committee emphasized that ME/CFS is a “medical—not psychiatric or psychological—illness” and called for increased research funding and physician education.

The 2024 NIH Deep Phenotyping Study. The landmark study by Walitt et al. (2024) using deep phenotyping provided definitive evidence for biological abnormalities in ME/CFS [5]. This study demonstrated brain dysfunction (decreased temporoparietal junction activity), immune exhaustion (exhausted T-cells and chronic B-cell activation), and neurochemical abnormalities (low catecholamines in cerebrospinal fluid). The NIH officially stated that “ME/CFS is a serious, chronic, systemic disease. . . researchers have found differences in the brains and immune systems of people with post-infectious ME/CFS” [32]. This study fundamentally shifted classification from a syndrome of unknown cause to a disease with identifiable pathological mechanisms.

The COVID-19 Catalyst. The COVID-19 pandemic paradoxically accelerated ME/CFS research by creating millions of long COVID patients with overlapping symptom profiles. The recognition that 10–30% of COVID-19 survivors develop persistent symptoms, with up to 51% of long COVID patients meeting ME/CFS diagnostic criteria [33], brought unprecedented attention and funding to post-infectious illness research. This convergence has produced several advances:

- **Shared research infrastructure:** Long COVID research programs (RECOVER, PHOSP-COVID) have included ME/CFS comparison groups, generating high-quality data on both conditions.
- **Treatment crossover:** Interventions studied for long COVID, including low-dose naltrexone (LDN), have shown promise in ME/CFS populations. LDN, which modulates microglial activation and neuroinflammation, emerged from long COVID clinical experience and is now being systematically studied in ME/CFS [34].
- **Biomarker discovery:** The urgency of the long COVID crisis has accelerated biomarker research applicable to both conditions, including markers of immune exhaustion, microclotting, and mitochondrial dysfunction.
- **Public recognition:** The visibility of long COVID has reduced stigma around ME/CFS and increased acceptance that post-infectious chronic illness is a legitimate medical phenomenon.

The connection between long COVID and ME/CFS has validated decades of patient advocacy and shifted the research paradigm toward viewing ME/CFS as the prototypical post-acute infection syndrome.

1.3 Disease Classification: From Syndrome to Disease

1.3.1 The Syndrome vs. Disease Distinction

In medical terminology, a **syndrome** refers to a collection of symptoms that occur together without a known underlying cause or identifiable pathological mechanism. A **disease**, by contrast, implies a known pathological process with specific biological markers and measurable damage to body systems.

For decades, ME/CFS was classified as a syndrome because medicine had not identified definitive biomarkers or universally agreed-upon pathological mechanisms. The February 2024 NIH study published in *Nature Communications* fundamentally changed this status.

1.3.2 The 2024 NIH Deep Phenotyping Study

The landmark study led by Dr. Avindra Nath (Walitt et al., 2024) used “deep phenotyping”—the most rigorous biological testing ever performed on ME/CFS patients—to demonstrate that the condition is a **systemic biological disease**, not a psychological syndrome or vague collection of complaints.

Key Findings.

1. **Brain dysfunction:** Using fMRI, researchers found decreased activity in the **temporoparietal junction (TPJ)**, the brain region responsible for effort-based decision-making. This proves that fatigue is not “feeling tired” but a physical failure of the brain to properly signal the body to move.
2. **Immune exhaustion:** The study identified “exhausted T-cells” and chronic B-cell activation. CD8+ T-cells (“killer” cells) are stuck in permanent activation, as if fighting a phantom infection they can never clear. This suggests a **persistent antigen**—a piece of virus or protein—may be hiding in the body, continuously provoking the immune system.
3. **B-cell maturity deficit:** B-cells fail to “switch” to a mature state, explaining why the body cannot clear the initial trigger/infection.
4. **Neurochemical evidence:** Abnormally low levels of **catecholamines** (norepinephrine, dopamine) were found in cerebrospinal fluid—essential neurotransmitters for nervous system regulation and motor function.

1.3.3 Official Reclassification

Major health organizations now explicitly refer to ME/CFS as a **serious, chronic, systemic disease**:

“ME/CFS is a *serious, chronic, systemic disease*. . . researchers have found differences in the brains and immune systems of people with post-infectious ME/CFS.”
—Official NIH News Release, February 2024

The condition is increasingly grouped under:

- **Post-Acute Infection Syndromes (PAIS)**
- **Infection-Associated Chronic Illnesses (IACI)**
- **Post-Infectious ME/CFS (PI-ME/CFS)**

1.3.4 Why the Name Persists

The word “syndrome” remains in “Chronic Fatigue Syndrome” primarily due to:

- **ICD-10/11 coding systems:** Hospitals and insurance companies globally use these codes, and changing them is a slow bureaucratic process.
- **Historical inertia:** Medical nomenclature changes slowly even when scientific understanding advances.

The clinical approach, however, has moved decisively toward treating ME/CFS as a complex **neuroimmune disease**.

1.3.5 Toward a More Accurate Name: Dysergopoiesis

Given the accumulated evidence that ME/CFS is a systemic biological disease—not a syndrome, not mere fatigue, and certainly not a psychological condition—this document proposes a new designation that more accurately reflects the core pathology: **dysergopoiesis**.

The term is constructed from three Greek roots:

dys- (δυσ-) impaired, disordered—the standard medical prefix indicating dysfunction, as in dyslexia, dyspnea, dystonia.

ergo- (εργον) work, energy—the root found in *erg* (the unit of energy), *ergonomics*, and *synergy*.

-poiesis (ποίησις) making, production—the suffix used throughout hematology and cell biology: *hematopoiesis* (blood cell production), *erythropoiesis* (red blood cell production), *leukopoiesis* (white blood cell production).

Dysergopoiesis therefore means, literally, **impaired energy production**. As a Greek scientific coinage, the term is essentially identical across languages—just as *dyslexia*, *anemia*, and *hematopoiesis* require no translation:

Language	Spelling
English	dysergopoiesis
French	dysergopoïèse
German	Dysergopoesie
Dutch	dysergopoëse
Spanish	disergopoyesis

Why “Dysergopoiesis”?

It names the core pathology, not a symptom. The defining biological feature of ME/CFS is not “fatigue” but a measurable failure of cellular energy production. The 2024 NIH deep phenotyping study demonstrated objective abnormalities at every level of the energy production chain: depleted catecholamines in cerebrospinal fluid, exhausted T-cell populations unable to switch metabolic modes, dysfunctional B-cell maturation, and decreased temporoparietal junction activity [5]. Mitochondrial oxidative phosphorylation is compromised (see Chapter 6), immune cells cannot perform the glycolysis-to-OXPHOS transition required for sustained function (see Chapter 7), and autonomic regulation of energy distribution fails (see Chapter 10). The word *dysergopoiesis* names this multi-system energy production failure directly.

It explains post-exertional malaise. The cardinal symptom of ME/CFS—the pathological worsening that follows even minimal exertion—becomes immediately comprehensible under this name. If energy production is impaired, then any activity that draws on energy reserves triggers a deficit from which the body cannot recover normally. Post-exertional malaise is not a mysterious symptom; it is the predictable consequence of dysergopoiesis.

It follows established medical word formation. The *dys-* + [*process*] + *-poiesis* pattern is standard in medicine. Just as *dyserythropoiesis* means impaired red blood cell production and *dyshematopoiesis* means impaired blood cell production, *dysergopoiesis* means impaired energy production. Any physician who encounters the term can parse it immediately.

Advantages Over Existing Nomenclature

- **No trivialization:** “Dysergopoiesis” cannot be confused with ordinary tiredness. No one says “I have a bit of dysergopoiesis” after a long day at work.
- **Mechanism-centered:** The name points to the core pathology (impaired energy production) rather than to a single symptom (fatigue) or to an outdated neurological hypothesis (encephalomyelitis).
- **Clinically actionable:** A name centered on impaired energy production naturally orients treatment toward mitochondrial support, metabolic correction, and immune metabolic reprogramming—rather than toward “managing fatigue.”

- **Neutral with respect to etiology:** The name does not presuppose a specific trigger (viral, autoimmune, genetic). It describes *what is broken* regardless of *why*, making it applicable across all patient subgroups.
- **No psychogenic connotation:** The Greek medical construction carries inherent biological authority. No component of the word can be misread as implying volition, preference, or psychological causation.
- **Universally intelligible:** Being a Greek scientific coinage, the term requires no translation—it is the same word in every language, just as *dyslexia* is. This eliminates the problem of divergent national names (ME in the UK, CFS in the US, SFC in France) that has fragmented the patient community and confused the research literature.
- **Shorter and more pronounceable than “myalgic encephalomyelitis”:** At six syllables versus ten, *dysergopoiesis* is more compact and follows familiar phonetic patterns (*dys-er-go-poi-e-sis*). Patients can learn, spell, and pronounce it—unlike “encephalomyelitis,” which most patients cannot spell and most physicians cannot pronounce correctly on first encounter.

Comparison with Previous Renaming Attempts

The 2015 Institute of Medicine proposal “Systemic Exertion Intolerance Disease” (SEID) failed to gain adoption. SEID was criticized for still centering the name on a *symptom* (intolerance to exertion) rather than on the underlying pathology. It also retained the word “disease,” which, while technically correct, did not address the *specificity* problem—many diseases involve exertion intolerance.

Dysergopoiesis differs fundamentally: it names the *process that fails* (energy production), not the *symptom that results* (exertion intolerance or fatigue). This distinction is analogous to naming diabetes mellitus after the metabolic defect (insulin insufficiency) rather than after its most visible symptom (excessive urination), or to naming *dyserythropsis* after the failed process (red blood cell production) rather than after its clinical consequence (anemia).

1.3.6 Implications for Patient Care

The disease classification has practical consequences:

1. **Treatment approach:** Medicine has shifted from treating *symptoms* (like “fatigue”) to treating *mechanisms* (like “T-cell exhaustion” or “mitochondrial dysfunction”).
2. **Clinical trials:** New trials now target specific biological pathways identified in the 2024 study:
 - Checkpoint inhibitors to “wake up” exhausted T-cells
 - IVIG to calm overactive B-cells
 - Plasmapheresis to remove autoantibodies
 - Long-term antivirals to clear potential viral reservoirs
 - Vagus nerve stimulation to reduce neuroinflammation

3. **Patient validation:** The biological findings ended the era of “we don’t know if anything is physically wrong” and vindicated decades of patient reports.
4. **Long COVID connection:** The immune profile of ME/CFS is remarkably similar to certain types of Long COVID, leading to shared research funding and accelerated understanding.

1.3.7 The “Effort Preference” Controversy

The 2024 study caused controversy by describing the TPJ dysfunction as “effort preference.” Patient advocacy groups (#MEAAction, Solve ME/CFS Initiative) issued urgent warnings that this phrase could be misinterpreted as suggesting patients are “choosing” not to exert themselves.

The researchers clarified that this is a **physiological** response—a broken brain circuit protecting the body from damage—not a lack of willpower. The brain “refuses” effort because it has accurately detected that the body cannot safely complete the task without triggering a crash.

Key Distinction

Unwilling vs. Unable: Patients choose the “easy task” not because they lack motivation, but because they know their body *cannot physically complete* the hard task without triggering Post-Exertional Malaise. Pacing is not a “preference”—it is a **biological requirement**.

1.4 Epidemiology

ME/CFS is a significant public health burden affecting millions worldwide, with prevalence likely underestimated due to underdiagnosis and inconsistent application of diagnostic criteria.

1.4.1 Prevalence and Incidence

Global Estimates. A systematic review and metaanalysis of 45 studies found a pooled prevalence of 0.89% (95% CI: 0.60–1.33) using CDC-1994 (Fukuda) criteria [35]. Applied globally, this suggests approximately 71 million people are affected. However, prevalence estimates vary substantially based on diagnostic criteria used, ranging from 0.39% to 1.40%.

United States Prevalence. The CDC reported in December 2023 that 1.3% of U.S. adults (approximately 3.3 million Americans) have ME/CFS based on National Health Interview Survey data from 2021–2022 [36]. This represents the first official national prevalence estimate using validated survey methodology.

Incidence Rates. Population-based studies estimate incidence at 13.16 per 100,000 person-years in the United States [37]. Norwegian registry data demonstrate a bimodal age distribution of new diagnoses, with peaks at 10–19 years and 30–39 years [38].

Post-COVID Prevalence Surge. The RECOVER-Adult Study (2025) found that 4.5% of SARS-CoV-2 infected individuals developed ME/CFS meeting diagnostic criteria, compared to 0.6% in uninfected controls [33]. The hazard ratio of 4.93 indicates nearly five-fold increased risk following COVID-19 infection. Updated estimates suggest the post-COVID era has increased U.S. ME/CFS cases from 1.5 million to 5–9 million, with annual economic impact rising from \$36–51 billion to \$149–362 billion [39].

1.4.2 Demographic Patterns

Sex Distribution. ME/CFS demonstrates a female predominance with a 3:1 to 4:1 female-to-male ratio across most studies [38, 35]. However, 35–40% of diagnosed patients are male, representing a substantial burden. The female predominance suggests potential hormonal or immunological factors, though diagnostic bias (dismissing male presentations) may contribute to apparent sex ratios.

Age Patterns. CDC data show prevalence increases with age, peaking at 2.1% among adults aged 60–69 years before declining in older age groups [36]. However, disease onset follows a bimodal distribution with peaks in adolescence (10–19 years) and early middle age (30–39 years), with mean onset age of approximately 31.6 years [38]. Approximately 15% of patients become symptomatic before age 18.

Racial and Ethnic Distribution. CDC data show prevalence of 1.5% in White non-Hispanic individuals, 0.8% in Hispanic individuals, and 0.7% in Asian non-Hispanic individuals [36]. However, these data likely reflect diagnostic disparities rather than true prevalence differences—White respondents have 2.94 greater odds of receiving an ME/CFS diagnosis than non-White respondents after controlling for symptom severity [40]. Population-based studies suggest equal or higher risk in ethnic minorities when diagnostic access is controlled.

Socioeconomic Factors. Prevalence demonstrates an inverse relationship with income: 2.0% among those below the federal poverty level compared to 1.1% among those at 200% or more of the poverty level [36]. This gradient likely reflects bidirectional causation—lower socioeconomic status may increase disease risk through chronic stress and reduced healthcare access, while ME/CFS causes substantial work disability that reduces income.

1.4.3 Geographic Distribution

Meta-analysis data show no significant difference in prevalence between Western countries (1.32%) and Asian countries (1.51%) [35]. Within the United States, rural areas show higher prevalence (1.9%) compared to large metropolitan areas (1.0–1.1%) [36], potentially reflecting healthcare access differences, occupational exposures, or delayed diagnosis leading to more severe presentations.

1.4.4 Risk Factors

Post-Infectious Onset. The majority of ME/CFS cases follow acute infection. Epstein-Barr virus (infectious mononucleosis) is the most studied trigger, with 9–11% of adults and 7–13% of adolescents developing ME/CFS at 6–12 months post-infection [41]. Other documented viral triggers include herpesviruses (HHV-6, CMV), enteroviruses, influenza, and SARS-CoV-2. The RECOVER study found that 51% of long COVID patients meet ME/CFS diagnostic criteria [33].

Genetic Susceptibility. Heritability estimates are approximately 10%, similar to irritable bowel syndrome and migraine [42]. The DecodeME genome-wide association study (2025), the largest ME/CFS genetic study to date (21,620 cases), identified eight significantly associated loci and three key genes—BTN2A2, OLFM4, and RABGAP1L—all involved in viral and bacterial immune responses [43]. Notably, no shared genetic variants were found with depression or anxiety, supporting the distinction between ME/CFS and psychiatric conditions.

Other Factors. Additional proposed risk factors include prior immune dysregulation, female sex hormones, and environmental exposures. The combination of genetic susceptibility with an infectious trigger likely explains why only a subset of individuals develop ME/CFS following infection.

1.5 Disease Impact

ME/CFS produces profound impacts on quality of life, functional capacity, and socioeconomic status, with disease burden exceeding that of many other serious chronic conditions.

1.5.1 Quality of Life

Comparison to Other Chronic Conditions. Multiple studies using validated quality of life instruments demonstrate that ME/CFS patients have among the lowest health-related quality of life scores of any chronic condition. Using the SF-36, ME/CFS patients score lower than patients with cancer, multiple sclerosis, stroke, diabetes, heart disease, rheumatoid arthritis, and depression across most functional domains [44, 45].

A comprehensive comparison found ME/CFS patients scored significantly lower than multiple sclerosis patients on nearly all SF-36 domains, with the largest differences in Physical Component Summary, Role Physical, and Social Function [46]. Using the EQ-5D-3L instrument, ME/CFS demonstrated the lowest unadjusted health-related quality of life of 20 chronic conditions studied, at 55% of population mean values [45].

Severely Ill Patients. Quality of life deteriorates dramatically with disease severity. In a study of severely ill patients, SF-36 Physical Functioning scores averaged 13.3 (compared to 99.0 in healthy controls), Role Physical averaged 1.9 (vs. 99.4), and Social Functioning averaged 4.4 (vs. 92.5) [47]. The quality of life profile most closely resembles that of congestive heart failure, reflecting the profound functional limitations.

1.5.2 Disability and Functional Capacity

Housebound and Bedbound Prevalence. Approximately 25% of ME/CFS patients are housebound or bedbound [48]. On worst days, 61% report being bedbound and 75% are housebound or bedbound. The housebound population demonstrates dramatically worse functional status: Physical Functioning scores of 17.1 versus 42.0 in non-housebound patients, Social Functioning of 10.2 versus 30.7, and 86% receiving disability benefits compared to 57% of non-housebound patients [48].

Severity Classification. Functional capacity varies by severity:

Mild (~25% of patients) Able to work part-time or full-time with substantially reduced other activities; approximately 50% reduction from pre-illness function

Moderate (~50% of patients) Substantially reduced activity; unable to work; requires rest periods; approximately 30–50% of pre-illness function

Severe (~20% of patients) Largely housebound; limited to minimal activities of daily living; approximately 5–15% of pre-illness function

Very Severe (~5% of patients) Bedbound; unable to perform most activities of daily living; often unable to tolerate sensory stimulation; less than 5% of pre-illness function

Work Disability. Employment rates range from 20–41% across studies, with 35–69% unemployed due to illness [49]. In a large Spanish cohort (n=1,086), 58.6% were unemployed, with 66% on sick leave and 34% receiving disability benefits. Risk factors for work disability include age over 50 years (OR 2.21), higher fatigue scores (OR 2.09), severe depression (OR 1.98), and autonomic dysfunction (OR 2.21) [49]. Only 13% of ME/CFS patients maintain full-time employment.

1.5.3 Economic Burden

Pre-COVID Estimates. The National Academy of Medicine (2015) estimated annual U.S. economic burden at \$17–24 billion. Updated analyses accounting for population growth and inflation revised this to \$36–51 billion annually [50].

Post-COVID Estimates. With ME/CFS prevalence potentially increasing from 1.5 million to 5–9 million U.S. cases due to post-COVID onset, updated economic impact estimates range from \$149–362 billion annually [39]. This includes direct medical costs and lost productivity but excludes disability benefits, social services, and caregiver lost wages, suggesting the true economic burden is substantially higher.

1.5.4 Psychosocial Impact

Social Isolation. ME/CFS produces profound social isolation: 57.7% of patients report significant isolation, with illness discussed only with immediate family (84%) or close friends (79.9%), rarely with coworkers (21.9%) [51]. The primary contributing factor is lack of disease understanding in social circles (90.5%).

Mental Health. While ME/CFS is not a psychiatric condition, 88.2% of patients report negative mental health effects from the illness [51]. Critically, 78.1% develop depression *after* ME/CFS onset, and 96% attribute their depression to disease severity and external factors rather than pre-existing psychiatric conditions. This distinguishes secondary depression resulting from chronic illness and loss of function from primary depressive disorders.

Medical Invalidation and Stigma. Patients experience pervasive stigmatization (68.5%) and diagnostic delays—67–77% wait more than one year for diagnosis, 29% wait more than five years, and over 70% see four or more physicians before diagnosis [6]. An estimated 84–91% of ME/CFS cases remain undiagnosed in the United States. Medical dismissal, misattribution to psychological causes, and lack of physician knowledge contribute to profound distress and delayed access to appropriate care.

Suicide Risk. ME/CFS patients face substantially elevated suicide risk. A UK study found suicide six to seven times more likely in ME/CFS patients compared to the general population [52]. In a mortality study of 56 deceased ME/CFS patients, suicide was the leading cause of death at 26.8%, with mean age at death from suicide of 41.3 years [53]. Contributing factors include being told the disease is psychosomatic (89.5%), feeling at the end of strength (80.7%), not being understood (80.7%), and experiencing stigmatization (76.8%) [51].

Premature Mortality. Beyond suicide, ME/CFS patients die earlier from all causes. Mean age of death was 55.9 years compared to 73.5 years in the general population (17.6 years earlier), with cardiovascular death occurring 18.9 years earlier on average [53]. At the time of death, 48.2% of patients were bedbound, and 83.7% of caregivers attributed death to ME/CFS.

1.6 Prognosis and Disease Course

Understanding the natural history of ME/CFS is essential for patient counseling, treatment planning, and setting realistic expectations.

1.6.1 Onset Patterns

ME/CFS typically presents in one of two patterns:

Acute post-infectious onset The majority of cases (60–80%) follow acute infection, most commonly Epstein-Barr virus (infectious mononucleosis), but also other herpesviruses, enteroviruses, influenza, and SARS-CoV-2 [41]. Patients can often identify the specific illness that marked disease onset. Initial presentation may resemble a prolonged viral illness that fails to resolve.

Gradual onset A minority of cases develop insidiously over months to years without a clear precipitating event. These patients may have difficulty identifying when the illness began and often report slowly progressive fatigue and functional decline.

1.6.2 Disease Trajectory

Early Course. The first two years following onset are often the most dynamic. Some patients experience spontaneous improvement, particularly those diagnosed early with mild presentations. However, symptoms that persist beyond 2–3 years rarely resolve completely [54].

Chronic Phase. Most patients enter a chronic phase characterized by:

- Fluctuating symptom severity with unpredictable good and bad periods
- Gradual adaptation to illness through activity modification
- Stable or slowly declining function if pacing is inadequate
- Episodic crashes following overexertion, infections, or other stressors

Recovery Rates. Full recovery is uncommon. Systematic reviews estimate that only 5% of patients recover to pre-illness function [54]. Improvement (partial recovery) occurs in approximately 40% of patients, typically those with milder initial presentations, shorter illness duration at diagnosis, and absence of psychiatric comorbidity. Approximately 40–50% of patients remain stable without significant improvement, while 10–20% experience progressive deterioration.

Prognostic Factors. Factors associated with better outcomes include:

- Younger age at onset
- Shorter illness duration at time of diagnosis
- Milder initial severity
- Absence of psychiatric comorbidity
- Early diagnosis and appropriate management
- Ability to implement effective pacing

Factors associated with worse outcomes include:

- Older age at onset
- Greater initial severity
- Longer diagnostic delay
- Continued overexertion (forced or voluntary)
- Comorbid conditions (fibromyalgia, POTS, depression)
- Lack of social support

1.6.3 Severity Fluctuation

ME/CFS severity can change over time:

- **Within-day variation:** Many patients experience predictable patterns of better and worse times of day
- **Week-to-week fluctuation:** Symptom severity varies unpredictably, complicating activity planning
- **Seasonal variation:** Some patients report consistent seasonal patterns
- **Relapse following triggers:** Infections, physical or emotional stress, surgery, and hormonal changes can trigger prolonged relapses
- **Long-term trajectory:** Severity may gradually improve, remain stable, or worsen over years

Patients who were initially mild or moderate may become severe or very severe following major relapses, and recovery from such relapses is often incomplete. This underscores the importance of aggressive pacing and trigger avoidance.

2 Core Symptoms

ME/CFS is characterized by several hallmark symptoms that must be present for diagnosis across most diagnostic frameworks. This chapter provides detailed descriptions of each core symptom.

2.1 Post-Exertional Malaise (PEM)

Post-exertional malaise (PEM), also termed post-exertional symptom exacerbation (PESE) or post-exertional neuroimmune exhaustion (PENE), is considered the hallmark feature of ME/CFS.

2.1.1 Definition and Characteristics

Post-exertional malaise represents an abnormal response to physical, cognitive, or emotional exertion in which even minor activity triggers a cascade of worsening symptoms. Unlike normal fatigue, PEM is characterized by:

- **Delayed onset:** Symptoms typically worsen 12–48 hours after the triggering activity
- **Disproportionate severity:** Minimal exertion produces profound symptom exacerbation
- **Prolonged recovery:** Symptom worsening persists for days to weeks or longer
- **Cumulative effect:** Sequential exertions compound impairment
- **Unpredictable threshold:** The level of activity that triggers PEM varies and may decrease over time

Common Triggers

PEM can be triggered by various forms of exertion:

Physical Exertion

- Walking, standing, or basic activities of daily living
- Exercise or physical therapy
- Household tasks
- Sexual activity
- Medical procedures or examinations

Cognitive Exertion

- Reading, writing, or computer work
- Conversation or social interaction
- Decision-making or problem-solving
- Sensory stimulation (light, sound, crowds)
- Concentration or sustained attention

Emotional Exertion

- Stress or anxiety
- Emotional processing
- Social demands
- Medical appointments or advocacy

Dietary and Environmental Factors

- Certain foods, including wheat in patients with non-celiac wheat sensitivity (approximately 15% of ME/CFS patients)
- Food sensitivities may interact synergistically with physical exertion, amplifying PEM severity
- Environmental exposures (chemical sensitivities, infections)
- Sleep disruption

See Section 23.3.4 for detailed discussion of wheat sensitivity mechanisms and clinical assessment approaches.

Subjective Phenomenology: The Effort-Performance Disconnect

One of the most psychologically devastating aspects of PEM is the profound disconnect between subjective effort and objective performance. Patients consistently describe an internal experience of maximal exertion that produces minimal external results—a phenomenon that fundamentally challenges their sense of agency and capability [55, 56].

The Experience of Maximal Effort Producing Minimal Output Unlike healthy individuals or those with deconditioning, ME/CFS patients report that activities feel intensely demanding internally while producing negligible observable output. A patient attempting to walk across a room may experience the subjective intensity of running a marathon—racing heart, overwhelming fatigue, sense of desperation—while moving slowly and covering minimal distance. This creates a surreal mismatch between internal state and external reality.

This disconnect extends beyond physical tasks:

- **Physical tasks:** Simple actions feel extraordinarily difficult; patients describe “giving everything” yet achieving almost nothing
- **Cognitive tasks:** Intense concentration yields minimal comprehension or output
- **Emotional regulation:** Enormous internal effort required to maintain composure or engage socially

Psychological Sequelae: Helplessness and Loss of Agency The persistent effort-performance disconnect produces profound psychological consequences distinct from primary depression:

Learned helplessness Repeated experiences of maximal effort failing to produce normal results can induce a state resembling learned helplessness—the recognition that one’s actions do not reliably produce expected outcomes. This is not a cognitive distortion but an accurate perception of physiological reality.

Loss of self-efficacy The inability to generate normal performance despite perceived maximum effort erodes confidence in one’s capability. Patients often describe feeling “weak” or “useless,” not as depression-related negative cognition but as direct experiential feedback.

Betrayal by one’s body Many patients describe their body as having “betrayed” them or become “enemy territory”—the normal unity between intention and execution has fractured. Motor commands and cognitive efforts no longer reliably produce proportional results.

Social invalidation Because the internal experience of extreme exertion is invisible to observers, patients face disbelief from family, friends, employers, and medical professionals. The statement “you don’t look sick” becomes particularly traumatic when one is experiencing maximum physiological stress.

Anticipatory anxiety Knowledge that even minor exertion may trigger severe crashes creates pervasive anxiety around all activities. Patients must constantly calculate risk, leading to hypervigilance and decision paralysis.

Distinction from Primary Depression While the phenomenology of PEM may superficially resemble depression, key distinctions exist:

- **Effort expenditure:** Depressed individuals typically experience reduced motivation to initiate effort; ME/CFS patients expend maximum subjective effort but achieve minimal results
- **Activity relationship:** Depression may improve somewhat with activity; ME/CFS worsens predictably with exertion
- **Physiological markers:** PEM produces objective physiological changes (documented via two-day CPET) absent in primary depression
- **Cognitive content:** The helplessness in ME/CFS arises from accurate perception of physiological limitation, not cognitive distortion [57]

Many ME/CFS patients develop secondary depression as a consequence of chronic illness and loss of function, but the core effort-performance disconnect represents a direct physiological phenomenon, not a psychological disorder. The majority (78.1%) of ME/CFS patients who experience depression develop it *after* disease onset, and 96% attribute their depression to disease severity and external factors rather than pre-existing psychiatric conditions [58].

Vulnerability and Existential Threat The profound energy deficit creates an acute sense of vulnerability. Patients describe feeling as though they “wouldn’t amount to shit” in any demanding situation—an accurate assessment of their current physiological capacity, not a self-esteem issue. This recognition of one’s fundamental vulnerability in a world that demands productivity and physical capability constitutes an ongoing existential threat.

For patients previously defined by physical capability, intellectual performance, or caregiving roles, the loss of reliable energy production represents a fundamental identity disruption. The inability to protect oneself, care for dependents, or meet basic social obligations creates legitimate existential distress [56]. Quality of life in ME/CFS is profoundly diminished, with patients scoring lower than those with multiple sclerosis, stroke, cancer, and other serious chronic conditions across nearly all functional domains [59, 60].

Severity Spectrum

PEM severity varies considerably:

Mild Increased symptoms for 1–3 days following moderate exertion; can usually continue limited activities with careful pacing

Moderate Severe symptom exacerbation lasting days to weeks following minimal exertion; requires extended rest periods

Severe Profound crashes triggered by activities of daily living; largely bedbound; recovery may take weeks to months

Very severe Any stimulation (light, sound, conversation) triggers immediate worsening; may be unable to tolerate even basic self-care

Baseline Energy Insufficiency: Living Below the Survival Threshold

While PEM represents the acute exacerbation following exertion, many ME/CFS patients describe a more insidious and pervasive problem: chronic baseline energy levels insufficient for basic existence. This creates a fundamentally different experience from episodic illness—it is a continuous state of inadequacy [55].

The Experience of Perpetual Insufficiency Patients describe waking already depleted, as if they have already run a marathon before the day begins. Unlike healthy individuals who start each day with a replenished energy reserve, ME/CFS patients begin from deficit:

- **Morning depletion:** Waking feeling as exhausted as when going to sleep, or worse

- **Minimum activity burden:** Even basic hygiene, eating, or sitting upright feels overwhelming
- **Continuous depletion:** Energy steadily drains throughout the day regardless of activity level
- **No reserve:** Zero capacity to handle unexpected demands
- **Micro-activities as exertion:** Actions that should be automatic (maintaining posture, processing sensory input) require conscious effort and consume limited energy

The experience of legs aching simply from sitting at a computer exemplifies this phenomenon. Maintaining posture—a task that should require minimal conscious attention—becomes actively depleting. Muscles fatigue from static contraction, venous pooling worsens due to inadequate muscle pump activity, and the metabolic cost of remaining upright exceeds available cellular ATP production.

Forced Overexertion: When Life Does Not Accommodate Limits Unlike research protocols where patients can carefully pace within their limits, real life imposes non-negotiable demands. This creates a situation of continuous forced overexertion:

Basic survival needs Eating, toileting, hygiene cannot be deferred indefinitely. Even these minimal activities may exceed available energy.

Medical appointments Navigating healthcare—attending appointments, waiting in waiting rooms, explaining symptoms, completing forms—requires energy patients do not have, creating the paradox of becoming sicker from seeking medical care.

Caregiving responsibilities Parents must feed children, pet owners must care for animals, adult children must respond to aging parents' needs. These responsibilities do not pause for energy availability.

Work and financial survival Many patients cannot afford to stop working despite severe energy limitations. The choice becomes: exceed limits and worsen disease, or face homelessness and starvation.

Emergencies House fires, medical emergencies, natural disasters, family crises demand immediate responses that may require weeks or months of energy expenditure in moments.

Social obligations Complete withdrawal results in loss of relationships, but social interaction is energetically costly. Patients must choose between isolation and overexertion.

Bureaucratic demands Disability applications, insurance appeals, medical documentation require sustained cognitive effort precisely when cognition is most impaired.

The Impossibility of Perfect Pacing While pacing (staying within energy limits to avoid PEM) represents the primary management strategy [61], perfect pacing is functionally impossible for most patients:

- **Unknown threshold:** The exertion level that will trigger PEM is variable and often unknowable in advance
- **Declining reserves:** The safe activity level may decrease over time, making previously manageable activities dangerous

- **Life is not optional:** Survival needs create forced exertion regardless of consequences
- **Delayed feedback:** PEM onset occurs 12–48 hours after trigger, preventing real-time adjustment
- **Compounding factors:** Stress, infection, hormonal cycles, weather, and other factors unpredictably lower the threshold
- **Cumulative depletion:** Multiple small activities compound, each individually acceptable but collectively triggering crashes

This creates a chronic state of being forced to operate beyond one's physiological capacity. Patients are not failing to pace properly—they are trapped in circumstances that structurally require overexertion for survival. Research demonstrates that exceeding energy limits worsens functional outcomes, yet life circumstances often make such overexertion unavoidable [62, 63].

The Grinding Exhaustion of Baseline Inadequacy The continuous nature of baseline energy insufficiency distinguishes it from acute exhaustion:

- **No recovery window:** There is no point at which energy feels restored; at best, crashes are avoided
- **Perpetual calculation:** Every action requires assessment of energy cost versus necessity
- **Invisible to others:** The constant internal struggle to perform basic tasks is entirely invisible; patients appear to be “doing nothing” while experiencing maximum effort to remain upright and conscious
- **Accumulating deficits:** Years of operating below subsistence level compound, potentially worsening disease trajectory
- **Eroded quality of life:** Even when avoiding severe crashes, life becomes reduced to the bare minimum, with no energy for joy, connection, or meaning

Psychological Impact of Chronic Insufficiency The experience of perpetual energy deficit below survival requirements produces distinct psychological consequences:

- **Perpetual crisis state:** Living constantly at the edge of capacity creates unrelenting stress
- **Inability to plan:** When basic function is uncertain day-to-day, future planning becomes impossible
- **Loss of identity:** Activities that defined one's self become permanently inaccessible
- **Anticipatory dread:** Every upcoming obligation triggers fear about whether one will have sufficient energy
- **Grief without resolution:** Unlike grief over a discrete loss, the loss of capability is ongoing and total
- **Existential exhaustion:** Beyond physical fatigue, the sheer effort of continuing to exist in this state becomes overwhelming

This baseline insufficiency, combined with forced overexertion and the acute crashes of PEM, creates a situation of profound and continuous suffering that is difficult for healthy individuals

to conceptualize. It is not merely “being tired”—it is operating every moment at a fundamental energy deficit incompatible with sustainable human function.

2.1.2 Physiological Basis

Mitochondrial Dysfunction and Energy Depletion

Observation 1 (WASF3-Mediated Mitochondrial Dysfunction). Skeletal muscle biopsies from ME/CFS patients (n=14) demonstrated significantly elevated WASF3 protein levels compared to healthy controls (n=10), with WASF3 overexpression correlating inversely with Complex IV function ($r=-0.55$, $p=0.005$) [2]. Mechanistic studies revealed that endoplasmic reticulum (ER) stress induces WASF3 protein accumulation at ER-mitochondrial contact sites, where it disrupts respiratory supercomplex assembly and inhibits mitochondrial respiration. Transgenic mice with elevated WASF3 expression recapitulated the human phenotype, exhibiting impaired exercise capacity and reduced oxygen consumption. shRNA-mediated WASF3 knockdown in patient-derived cells restored respiratory capacity, demonstrating reversibility of the dysfunction.

~ Hypothesis 1: WASF3 as Subset-Specific Mechanism

The WASF3-mediated mitochondrial dysfunction mechanism may explain exercise intolerance in a subset of ME/CFS patients, particularly those with post-viral onset [2, 64]. The pathway linking viral infection → ER stress → WASF3 elevation → mitochondrial dysfunction → ATP depletion provides a coherent mechanistic framework. However, the prevalence of this mechanism across the broader ME/CFS population remains undetermined, as the initial finding derives from a small cohort (n=14). Independent replication and larger validation studies are needed to establish what proportion of ME/CFS patients exhibit this pathway.

The WASF3 mechanism aligns with broader evidence of mitochondrial dysfunction in ME/CFS [64]. ATP depletion following exertion explains the delayed onset of PEM (cellular energy stores require 24–72 hours to regenerate) and the disproportionate symptom severity (cells cannot meet metabolic demands even for basic function). WASF3 overexpression promotes actin polymerization, driving a metabolic shift toward glycolysis while further suppressing mitochondrial oxidative phosphorylation. This creates a self-reinforcing cycle: reduced ATP generation → increased cellular stress → sustained WASF3 elevation → continued mitochondrial impairment.

The 24–72 Hour Delay: Why PEM Onset Is Delayed

One of the most distinctive and poorly understood features of post-exertional malaise is its delayed onset. Unlike normal exercise fatigue, which peaks during or immediately after activity, PEM symptoms typically worsen 12–72 hours after the triggering exertion. This delay creates a diagnostic challenge (patients may not connect symptoms to earlier activity) and a management challenge (real-time feedback for pacing is unavailable). Understanding why

this delay occurs requires examining multiple overlapping mechanisms operating on different timescales.

ATP Depletion and Regeneration Kinetics. The most direct explanation for delayed PEM onset involves the temporal dynamics of cellular ATP depletion and failed regeneration:

- **Initial buffering:** During exertion, phosphocreatine provides immediate ATP buffering for seconds, followed by glycolytic ATP production for minutes. These rapid systems mask the underlying mitochondrial ATP deficit during activity.
- **Progressive depletion:** As activity continues, cellular ATP pools gradually deplete. In healthy individuals, mitochondrial oxidative phosphorylation rapidly restores ATP during rest. In ME/CFS, impaired mitochondrial function prevents normal regeneration.
- **Critical threshold crossing:** Symptoms manifest when ATP levels fall below the minimum required for essential cellular functions—ion pump maintenance, neurotransmitter synthesis, immune cell activation. This threshold may not be crossed until hours after exertion ends, as residual ATP is consumed by basal metabolic demands without adequate replenishment.
- **Tissue-specific kinetics:** Different tissues have different ATP demands and regeneration capacities. Brain and muscle may reach critical thresholds at different times, producing the characteristic multi-system symptom cascade.

The Heng 2025 multi-omics study documented elevated AMP and ADP in ME/CFS patients' white blood cells [12], indicating cells are attempting to regenerate ATP (AMP/ADP are intermediates) but failing to complete the process efficiently. This chronic partial depletion state means even modest exertion pushes cells over the edge into critical deficit.

Mitochondrial Turnover Rate Limitation. A complementary explanation involves mitochondrial damage and repair kinetics:

- **Exercise-induced damage:** Exertion generates reactive oxygen species (ROS) that damage mitochondrial membranes, proteins, and respiratory chain complexes. In ME/CFS, baseline mitochondrial dysfunction and impaired antioxidant defenses magnify this damage.
- **Mitophagy lag:** Damaged mitochondria must be removed via mitophagy (selective autophagy) before replacement. This process requires hours to days. During this lag, cellular ATP production capacity declines progressively.
- **Biogenesis delay:** New mitochondrial synthesis requires transcription, translation, membrane assembly, and functional integration—processes requiring 24–72 hours minimum. The 13-day recovery period observed in 2-day CPET studies [1] closely matches published mitochondrial turnover times in muscle tissue (10–15 days).
- **Cumulative deficit:** If exertion outpaces the capacity for mitochondrial repair and replacement, functional mitochondrial mass declines over days, producing progressively worsening symptoms despite cessation of activity.

This mechanism suggests PEM delay reflects not just ATP depletion but the time required for damaged cellular machinery to be replaced—a process that may be further slowed by circadian dysregulation of mitophagy and biogenesis pathways.

Delayed-Type Immune Activation. The temporal pattern of PEM symptom onset (24–72 hours) closely matches delayed-type hypersensitivity (DTH) immune responses, suggesting immune-mediated mechanisms:

- **Exercise as danger signal:** Physical exertion releases damage-associated molecular patterns (DAMPs) including extracellular ATP, HMGB1, and heat shock proteins. These activate innate immune receptors.
- **Cytokine cascade timing:** Pro-inflammatory cytokine production (IL-1 β , IL-6, TNF- α) peaks 24–48 hours post-stimulus in classical immune responses. Gene expression studies show prolonged elevation of immune activation genes 24–72 hours post-exercise in ME/CFS patients, corresponding to symptom exacerbation timing.
- **Purinergic signaling:** Exercise dramatically increases extracellular ATP release. If purinergic receptors (P2X7) are sensitized or ATP clearance is impaired, this triggers massive inappropriate danger signaling with delayed inflammatory consequences (see §14.9 for detailed discussion).
- **Neuroinflammation:** Microglial activation in response to peripheral immune signals requires hours to fully develop, potentially explaining delayed cognitive symptoms (brain fog).

The immune hypothesis explains not just the delay but also the multi-system nature of PEM—inflammatory cytokines affect brain, muscle, autonomic nervous system, and other tissues simultaneously.

Metabolic Byproduct Accumulation. Another layer involves the gradual accumulation of metabolic stress signals:

- **Lactate kinetics:** ME/CFS patients show elevated baseline lactate and impaired lactate clearance [65]. Exercise-induced lactate accumulation may persist for days rather than hours, maintaining tissue acidosis and triggering pain receptors.
- **ROS accumulation:** Reactive oxygen species generated during exertion damage lipids, proteins, and DNA. Oxidative damage markers peak hours after ROS generation as damaged molecules accumulate and overwhelm repair capacity.
- **Inflammatory metabolites:** Prostanoids, leukotrienes, and other inflammatory mediators are synthesized and released over hours following initial cellular stress.

Additional Mechanistic Hypotheses: Ranking by Plausibility. Beyond the primary mechanisms described above, several additional hypotheses have been proposed to explain the 24–72 hour delay. These vary in evidentiary support and mechanistic plausibility:

Tier 1: Highly Plausible (Strong Mechanistic Support).

- **NAD⁺ depletion crisis via PARP hyperactivation:** Exercise-induced reactive oxygen species cause DNA damage, activating poly(ADP-ribose) polymerase (PARP) enzymes for repair. PARP consumes NAD⁺ at 100–1000× normal rates (extrapolated from general PARP biochemistry; specific rate in ME/CFS not directly measured). Since NAD⁺ is required for the Krebs cycle, PARP hyperactivation creates a vicious cycle: DNA damage → PARP activation → NAD⁺ depletion → impaired ATP synthesis → insufficient NAD⁺ regeneration. The Heng 2025 study documented NAD⁺ metabolism abnormalities in ME/CFS [12], supporting this pathway. The 24–72h delay reflects the time required for NAD⁺ pools to reach critical depletion.
- **Cellular “debt payment” model:** During exertion, cells desperately seeking ATP cannibalize structural proteins, cytoskeletal elements, and other cellular components through autophagy. This “borrowing” masks the immediate deficit but creates structural damage that manifests 24–72 hours later as neurotransmitter synthesis fails, contractile proteins degrade, and enzyme production becomes insufficient. The delay represents the time required for cellular structural integrity to fail after emergency self-digestion.
- **Critical threshold dynamics:** Cells do not fail linearly as ATP drops. They compensate through multiple buffering systems until ATP falls below approximately 30% of normal (estimated from general cellular bioenergetics; specific threshold in ME/CFS not directly measured), then experience catastrophic multi-system failure. During the 24–72h window post-exertion, basal metabolism consumes remaining ATP without adequate mitochondrial replenishment, progressively approaching this critical threshold. Different tissues reach critical thresholds at different times based on their ATP demands: brain (12–24h), skeletal muscle (24–48h), immune cells (24–72h).

Tier 2: Very Plausible (Good Mechanistic Support, Less Direct Evidence).

- **Lactate accumulation with delayed acidosis:** ME/CFS patients show elevated baseline lactate and impaired lactate clearance (discussed above). However, lactate clearance itself is ATP-dependent. As ATP depletion worsens over 12–24h post-exertion, lactate clearance machinery progressively fails, creating accelerating accumulation. Lactate diffuses slowly between muscle compartments, requiring 24–48 hours to reach concentrations sufficient to activate pain receptors and cause tissue acidosis.
- **Glycocalyx shedding and endothelial dysfunction:** Exercise-induced shear stress sheds the glycocalyx—the protective gel coating on vascular endothelium. During the 12–48h degradation period before regeneration begins (48–96h), microvascular permeability increases and tissue perfusion decreases. The brain and muscles can tolerate moderate hypoperfusion for approximately 24h before symptoms manifest. This mechanism explains delayed cognitive and physical symptoms through progressive tissue hypoxia.
- **Mitochondrial removal timing:** Damaged mitochondria are not removed immediately. Mitophagy peaks during specific time windows (often overnight), and biogenesis has circadian regulation. The delay may represent the time between damage occurrence and the next mitophagy window, followed by 24–72h for biogenesis. During this period, patients operate with reduced functional mitochondrial mass—the “hole” reaches maximum depth before new mitochondria come online.

Tier 3: Plausible (Interesting Mechanisms, Weaker Evidence).

- **Mast cell degranulation cascade:** Mast cells release mediators in three waves: immediate histamine (0–2h), proteases and tryptase (12–24h), and cytokines/prostanoids (24–72h). Each wave primes the next, explaining both delayed and prolonged symptoms. This mechanism applies primarily to the subset of ME/CFS patients with documented mast cell activation syndrome.
- **Circadian gating of repair processes:** Mitophagy and mitochondrial biogenesis are circadian-regulated, peaking at specific times of day. If exertion occurs outside optimal repair windows, cellular damage must wait until the next circadian cycle for repair to begin. This could explain why some patients report that time-of-day of exertion affects crash timing and severity (currently unstudied).
- **Autonomic recalibration failure:** Exercise should trigger autonomic system reset to parasympathetic dominance during recovery. In ME/CFS, this reset fails, leading to persistent sympathetic activation and progressive dysautonomia over 12–72h. Worsening autonomic dysfunction reduces tissue perfusion, creating energy crisis through inadequate oxygen and nutrient delivery.
- **Complement system amplification:** The complement cascade is an exponential amplification system—each activated molecule activates hundreds downstream. Exercise-induced tissue damage triggers C3 activation (6–12h), C5a production (12–24h), and terminal complement complex formation (24–48h). However, direct evidence of pathological complement activation in ME/CFS PEM remains limited.

Tier 4: Speculative (Low Evidence, Theoretically Interesting).

- **Microbiome metabolite shifts:** Exercise alters gut motility and oxygen availability, potentially shifting bacterial populations toward more inflammatory species and reducing short-chain fatty acid production. However, bacterial population shifts typically require days, making this timeline too slow to fully explain 24–72h delays. May contribute to sustained post-crash symptoms.
- **Glymphatic system failure:** The brain's glymphatic system clears metabolic waste during sleep. If sleep quality worsens post-exertion, waste accumulates over multiple failed sleep cycles (2–3 nights), reaching symptomatic levels at 48–72h. This explains delayed cognitive symptoms but not systemic or muscular symptoms.
- **Epigenetic reprogramming lag:** Exercise triggers histone modifications and DNA methylation changes requiring 24–72h to manifest as altered gene expression. Theoretically, this could reprogram cells into a dysfunctional metabolic state. However, whether such epigenetic changes are pathological or adaptive in ME/CFS remains unknown.

These additional mechanisms are not mutually exclusive with the primary pathways described earlier. Multiple mechanisms likely operate simultaneously, with different mechanisms dominating in different patients or disease subtypes. Understanding this mechanistic diversity may explain individual variation in crash timing (12h vs. 72h) and suggest personalized intervention strategies (discussed in Chapter 24).

Integrated Multi-Hit Model. The most plausible explanation integrates these mechanisms into a cascading sequence:

1. **During exertion (0–2 hours):** ATP consumption exceeds production; immediate buffering systems mask deficit; ROS generation damages mitochondria; extracellular ATP and DAMPs released.
2. **Early post-exertion (2–12 hours):** Residual ATP consumed by basal metabolism without adequate replenishment; damaged mitochondria marked for removal; immune sensing of danger signals initiates.
3. **Delayed crash phase (12–72 hours):** Critical ATP threshold crossed in multiple tissues; cytokine production peaks; damaged mitochondria removed faster than replaced; lactate and ROS accumulation reach symptomatic levels; neuroinflammation develops.
4. **Recovery phase (days to weeks):** Mitochondrial biogenesis gradually restores capacity; inflammatory mediators cleared; ATP pools slowly replenished; symptoms gradually resolve if no further exertion occurs.

This model explains why some patients notice symptoms within hours (early ATP threshold crossing) while others experience the classic 24–48 hour delay (immune-mediated mechanisms dominate), and why recovery requires days to weeks (mitochondrial turnover is rate-limiting).

Clinical Implications. Understanding the delayed onset mechanism has important practical consequences:

- **Pacing difficulty:** The 24–72 hour delay prevents real-time feedback. Patients must learn to predict delayed consequences rather than respond to immediate symptoms.
- **Activity tracking:** Detailed activity logs correlated with symptoms 24–72 hours later are essential for identifying triggers.
- **Preventive intervention window:** If ATP depletion initiates the cascade, aggressive energy substrate provision (D-ribose, MCT oil) immediately post-exertion might prevent threshold crossing and abort the crash.
- **Anti-inflammatory timing:** If immune activation drives delayed symptoms, prophylactic anti-inflammatory interventions timed to the 12–24 hour window might reduce severity.
- **Diagnostic utility:** The characteristic delay helps distinguish PEM from deconditioning (which produces immediate fatigue) and depression (which lacks temporal relationship to activity).

Unanswered Questions. Despite these proposed mechanisms, critical gaps remain:

- **Individual variation:** Why do some patients crash at 12 hours while others at 72 hours?
- **Threshold determinants:** What factors determine when ATP levels cross the critical threshold?
- **Preventability:** Can early intervention (within 2–12 hours post-exertion) abort the cascade before symptoms manifest?

- **Subset mechanisms:** Do different ME/CFS subgroups have different dominant delay mechanisms (metabolic vs. immune)?
- **Chronobiology:** Does time of day of exertion affect delay duration through circadian regulation of repair processes?

The 24–72 hour delay represents a central feature of ME/CFS pathophysiology that distinguishes it from simple deconditioning or fatigue. Elucidating the precise mechanisms could enable targeted interventions timed to specific phases of the crash cascade. Evidence-based and hypothesis-driven intervention strategies targeting these mechanisms are discussed in Chapter 24.

Cascade Dependency: Can Early Intervention Prevent Downstream Phases? Understanding whether the crash cascade phases are causally dependent (sequential) or independently triggered (parallel) has profound therapeutic implications. If fixing Phase 1 (ATP depletion) prevents Phases 2–4, early intervention could abort crashes entirely. If phases trigger independently, intervention can only mitigate severity.

The Critical Question: Prevention vs. Mitigation. Two competing models exist:

Sequential Dependency Model (Optimistic) ATP depletion at Phase 1 *causes* mitochondrial damage, immune activation, and symptoms at Phases 2–4. Preventing ATP crisis prevents everything downstream.

Parallel Initiation Model (Pessimistic) All phases trigger simultaneously during exercise but manifest at different times. Intervention can only reduce amplification, not prevent the cascade.

Analysis of Phase Dependencies. Phase 1 → Phase 2 (ATP Crisis → Mitochondrial Damage): *Partial prevention possible.*

Some mitochondrial damage occurs inevitably during exercise from ROS generation as respiratory chain activity increases. This initial damage happens in real-time (0–2h) and cannot be prevented post-hoc. However, ATP depletion dramatically amplifies this damage through multiple mechanisms: (1) impaired antioxidant synthesis (glutathione production requires ATP), (2) disabled repair protein function (requires ATP), and (3) positive feedback loops where damaged mitochondria leak additional ROS when ATP-dependent quality control fails.

Preventing ATP crisis cannot undo initial ROS damage but can prevent the amplification cascade. **Theoretical estimate (no direct empirical validation in ME/CFS):** Initial damage unavoidable (estimated ~20% of total, based on exercise physiology showing inevitable ROS generation during oxidative metabolism), but ATP-mediated amplification (estimated ~80% of total, extrapolated from ATP-dependent antioxidant/repair pathways) is theoretically preventable. Hypothesized net reduction in mitochondrial damage: 60–80%.

Phase 1 → Phase 3 (ATP Crisis → Immune Cascade): *Mixed prevention and delay.*

The immune cascade has dual triggers operating on different timescales:

- **Immediate trigger (unavoidable):** DAMPs (extracellular ATP, HMGB1, heat shock proteins, mitochondrial fragments) release during exercise (0–2h) as normal cellular stress signaling. Even healthy individuals release these; ME/CFS patients' immune systems appear sensitized to over-respond to physiological levels.
- **Sustained trigger (preventable):** ATP-depleted cells continue releasing DAMPs for 24–72h as ongoing distress signals. Additionally, ATP depletion impairs immune regulatory mechanisms (requires ATP), allowing uncontrolled cytokine amplification. Damaged mitochondria release mtDNA (recognized as bacterial pathogen-associated molecular pattern), triggering massive additional immune activation.

Preventing ATP crisis cannot eliminate initial DAMP sensing but can prevent sustained DAMP release and immune dysregulation. **Theoretical estimate (no direct empirical validation):** Initial immune activation unavoidable (estimated ~30%, based on immediate DAMP release during exercise observed in healthy populations), but cytokine storm amplification (estimated ~70%, extrapolated from ATP-dependent immune regulation) is theoretically preventable. Hypothesized net reduction in immune activation: 70–80%.

Phase 2 → Phase 3 (Mitophagy Gap → Symptoms): *Mitigation only, not prevention.*

The “mitochondrial gap” appears inevitable once damage occurs. If exercise damages 20% of mitochondria, they must be removed (mitophagy, 6–12h) before replacement (biogenesis, 72h–14d). During this window, functional mitochondrial mass drops from 100% to 80%, creating energy crisis regardless of intervention. The gap can be made shallower (less initial damage) or shorter (faster biogenesis), but cannot be eliminated. This biological reality—damaged cellular machinery requires days to replace—sets a floor on recovery time even with perfect intervention.

Integrated Dependency Model: Hybrid Causation. Evidence suggests cascade phases are *partially dependent* rather than fully sequential or fully parallel:

1. **Exercise (0–2h):** Unavoidable damage occurs
 - ROS generation: ~20% of eventual mitochondrial damage
 - DAMP release: ~30% of eventual immune activation
 - Substrate depletion: Phosphocreatine, glycogen consumption
2. **Early post-exertion (2–12h):** Amplification vs. recovery divergence
 - *Healthy individuals:* ATP replenishment, antioxidant regeneration, controlled immune signaling → Recovery initiated
 - *ME/CFS without intervention:* Progressive ATP depletion, failed antioxidant regeneration, dysregulated immune amplification → Cascade amplification
 - *ME/CFS with intervention:* Partial ATP support, antioxidant buffer, some immune modulation → Reduced amplification
3. **Delayed crash phase (12–72h):** Outcome determined by Phase 2 success
 - No intervention: ~600 severity units (100 initial + 200 ATP amplification + 300 immune amplification) — **Illustrative model, not empirical measurements**

- Perfect Phase 1 intervention: ~150 severity units (100 initial + minimal amplification)
- Hypothesized reduction: 75% (but 150 units still manifests)

Clinical Implications: Realistic Intervention Expectations. Early intervention (0–12h post-exertion) appears capable of dramatic severity reduction (60–80%) but not complete crash prevention. The irreversible components—initial ROS damage, DAMP release during exercise, and mitochondrial turnover biology—establish a floor on symptoms and recovery time even with perfect intervention.

What intervention CAN achieve:

- Prevent progression from *manageable* to *catastrophic* severity
- Reduce bedridden duration from weeks to days
- Maintain some functional capacity rather than complete incapacitation
- Shorten recovery from 13+ days to 5–7 days
- Prevent amplification spirals (severe crashes begetting more severe crashes)

What intervention CANNOT achieve:

- Complete crash prevention after threshold-exceeding exertion
- Elimination of all symptoms (initial damage is irreversible)
- Bypass of mitochondrial turnover time (biology is slow)
- Permission for routine energy envelope violations (cumulative damage still occurs)

The distinction between *prevention* and *mitigation* is critical for patient expectations and treatment evaluation. A 75% severity reduction—transforming a two-week bedridden crash into manageable tiredness for several days—represents enormous clinical benefit even though the cascade still occurs. Framing this as “partial prevention” rather than “treatment failure” recognizes the biological constraints while validating meaningful therapeutic effects.

For intervention protocols targeting these mechanisms, see Chapter 24, §24.10.3.

The Fundamental Question: Raising the Baseline vs. Managing Crashes. The critical therapeutic question is not merely whether interventions reduce crash severity, but whether they can *raise the baseline capacity*—shifting the threshold at which PEM occurs and potentially reversing the progressive decline characteristic of ME/CFS.

PEM as Universal Phenomenon with Pathological Threshold.

Post-exertional physiological stress is universal across all humans. Healthy individuals experience delayed-onset muscle soreness (DOMS), inflammation, and temporary fatigue 24–48h post-exercise through identical mechanisms: ROS-induced mitochondrial damage, DAMP-mediated immune activation, and cellular repair processes. The critical difference in ME/CFS is not the presence of these responses but their catastrophically lowered threshold and failed recovery:

- **Healthy athlete:** 10km run → Moderate soreness (48h) → Adaptation → Increased capacity
- **Healthy sedentary:** 1km walk → Mild soreness (48h) → Recovery → Baseline maintained
- **ME/CFS patient:** 100m walk → Severe crash (2 weeks) → Deterioration → Decreased capacity

The same biological cascade operates in all three populations. What differs is: (1) the activity threshold triggering the response (10km vs. 1km vs. 100m), (2) the magnitude of amplification (controlled vs. mild vs. catastrophic), and (3) the outcome trajectory (adaptation vs. recovery vs. decline).

Dose-Response Relationship: Linear vs. Catastrophic.

A fundamental distinction between healthy and ME/CFS post-exertional responses lies in the dose-response relationship—how symptom severity scales with exertion intensity:

Healthy individuals (moderate exertion range): Near-linear relationship. Doubling exercise intensity approximately doubles soreness severity and recovery time. A 5km run produces roughly half the DOMS of a 10km run. A 20km run produces proportionally more soreness but within the same biological framework—more microdamage requiring more repair, scaling predictably. This linearity allows precise training dose calibration.

ME/CFS patients: Catastrophically non-linear relationship with threshold effect. Below threshold (varies individually, often 50–200m walking): Minimal symptoms, proportional response similar to healthy population. Crossing threshold: Exponential amplification. Activity just 10% above threshold may produce not 10% worse symptoms but 500% worse symptoms—the difference between manageable tiredness and bedbound collapse. This non-linearity reflects cascade amplification: once ATP depletion crosses critical threshold (estimated ~30% of normal, based on general cellular bioenergetics), immune dysregulation triggers, mitochondrial damage amplifies exponentially, and the system enters catastrophic failure mode rather than controlled stress response.

Critical insight—The fundamental asymmetry: Why recovery is non-linear while damage may be linear. In healthy individuals, both damage and recovery scale linearly with exertion. A 10km run causes proportional muscle microdamage and inflammation, requiring proportional recovery time (perhaps 48–72h). The body has fuel reserves to execute the repair: adequate ATP to synthesize proteins, adequate NAD⁺ for cellular metabolism, functional mitochondria to generate energy for the repair processes themselves. Even extreme exertion (e.g., marathon) produces severe but *finite* damage that resolves predictably given adequate rest and nutrition.

In ME/CFS patients who cross threshold, a catastrophic asymmetry emerges: the exertion may produce *similar initial tissue damage* (microscopic muscle tears, oxidative stress, immune activation) as in healthy individuals, but the body **lacks the fuel to execute repair**. Once energy reserves are exhausted—ATP depleted, NAD⁺ consumed by PARP, mitochondria damaged—the body cannot synthesize repair proteins, cannot clear lactate, cannot regenerate antioxidants, cannot produce new mitochondria. The damage remains unrepaired not because it's irreparable, but because *repair itself requires energy the body no longer has*.

Worse, **even at complete rest**, the body continuously consumes energy for basic survival. The heartbeat cannot stop. The brain cannot pause. Muscles (even at rest) maintain tone. Fascia, blood vessels, immune cells, and every organ require continuous ATP just to stay alive—this basal metabolism represents roughly 60–70% of total daily energy expenditure in healthy individuals [66]. Once energy reserves are exhausted, this *unavoidable continuous drain* immediately consumes whatever tiny amount of new ATP the damaged mitochondria manage to generate, leaving **nothing for repair**. The body cannot allocate energy to fixing damaged mitochondria because survival functions have absolute priority.

This creates exponentially longer recovery time: the body must somehow generate enough energy to begin repairing the energy-generation machinery itself—a paradox. Recovery time becomes non-linear because the patient has fallen into an energy bankruptcy trap where escaping requires resources they don't have. *Complete rest does not stop energy consumption*; it merely reduces it to the basal minimum, which in severe ME/CFS may still exceed what damaged mitochondria can produce.

This explains several clinical observations:

- **Recovery time vastly exceeds damage time:** 5 minutes of walking → 2 weeks bedbound (time ratio 4000:1). The damage may be proportional to exertion, but recovery requires replacing the entire depleted energy infrastructure.
- **Widespread pain and dysfunction despite no structural damage:** When ATP falls below critical thresholds, normal cellular functions fail, generating symptoms that *mimic injury but are purely metabolic*:
 - **Muscle pain/weakness:** Sodium-potassium pumps fail (require ATP), calcium regulation fails (muscle relaxation requires ATP), lactate accumulates (clearance requires ATP)
 - **Fascial pain:** Fascia cannot maintain proper hydration or tension without ATP for active transport
 - **Joint pain:** Synovial fluid production and cartilage maintenance require continuous ATP
 - **Neuropathic pain:** Nerve cells cannot maintain membrane potentials (ion pumps require ATP), generating aberrant firing
 - **Cognitive dysfunction:** Neurons are extraordinarily ATP-dependent; even small deficits cause brain fog, memory problems, processing delays
 - **Dysautonomia:** Blood pressure regulation, heart rate variability, temperature control all require ATP for signaling

The pain and dysfunction are real and severe, but the cause is *metabolic failure, not tissue damage*. This is why anti-inflammatories and analgesics provide minimal relief—they target inflammation and pain pathways, but the underlying problem is ATP depletion.

- **Proper food/supplements critical during recovery:** Without exogenous ATP substrates (D-ribose), NAD⁺ precursors (NR/NMN), and repair cofactors (vitamins, minerals), the body may never escape energy bankruptcy. Recovery becomes impossible, not just slow.
- **“Good days” followed by crashes:** A “good day” (70% energy instead of 50%) tempts exertion beyond envelope. But 70% is still insufficient for repair; crossing threshold exhausts the reserves, dropping capacity to 30% for weeks.

- **Progressive decline with repeated crashes:** Each crash drains reserves further; inadequate between-crash recovery means starting the next crash from a lower baseline. The ratchet effect (Section 5.4.3) represents cumulative energy bankruptcy.

This model transforms therapeutic strategy: interventions must not merely support repair but must *break the energy bankruptcy trap* by providing exogenous substrates that bypass the damaged production machinery. D-ribose provides ATP backbone when synthesis fails; NAD⁺ precursors supply what PARP consumed; MCT oil provides ketones that bypass damaged mitochondrial complexes; antioxidants reduce the repair burden by preventing ongoing damage. The emergency PEM prevention protocol (Chapter 24, §24.10.3) is designed precisely to prevent entry into this bankruptcy state by flooding the system with substrates during the critical 0–72h window.

The practical consequence: Healthy individuals can safely experiment with exertion levels (“I’ll try running 8km today instead of 5km and see how I feel tomorrow”). ME/CFS patients face binary outcomes (“I walked 150m today instead of 100m and triggered a two-week crash”). The lack of proportional dose-response makes activity titration extraordinarily difficult—no middle ground exists between “tolerable” and “catastrophic.”

This non-linearity also explains why graded exercise therapy (GET) fails catastrophically in ME/CFS: protocols assume linear dose-response (“if 5 minutes is tolerated, 7 minutes will be proportionally harder”). In reality, crossing the threshold triggers exponential cascade, causing harm rather than adaptation. The threshold itself is variable (affected by sleep, stress, infection, hormonal cycles), making it impossible to identify safe progressive increments.

Threshold as Therapeutic Target.

If PEM represents an amplified version of normal exercise response rather than a unique pathological entity, the therapeutic goal shifts from “eliminating PEM” (impossible—all exercise causes cellular stress) to “raising the threshold” (increasing the amount of activity tolerated before triggering cascade).

Three potential intervention strategies emerge:

1. **Reduce amplification** (discussed above): Energy substrates, antioxidants, immune modulation prevent 100 units of exertion from becoming 600 units of damage. Does not raise threshold but makes threshold violations more survivable.
2. **Raise threshold directly:** If baseline mitochondrial capacity increases from 50% to 70% of normal through mitochondrial biogenesis interventions (urolithin A, NAD⁺ precursors, CoQ10), the same absolute exertion represents proportionally less stress. Patient goes from crashing after 100m to tolerating 500m before threshold.
3. **Improve recovery:** If repair machinery efficiency increases, each crash causes less permanent damage, preventing downward spiral. Patient maintains baseline rather than progressively declining.

Evidence for Reversibility vs. Irreversibility.

Critical unresolved question: Is ME/CFS mitochondrial dysfunction *fixed but suppressible* (like WASF3 overexpression) or *reversible*? Evidence suggests heterogeneity:

- **Potentially reversible mechanisms:** WASF3 knockdown in patient cells restored respiratory capacity [2], suggesting dysfunction is suppressible. NAD⁺ restoration, antioxidant support, and mitochondrial turnover acceleration target reversible deficits.
- **Potentially irreversible mechanisms:** If years of ROS damage caused permanent mtDNA mutations, permanent epigenetic silencing of metabolic genes, or structural cellular damage, restoring function may be impossible without cellular regeneration.
- **Subset variability:** Post-viral onset patients (recent damage) may have greater reversibility than long-duration patients (accumulated permanent damage). Age, severity, and comorbidities likely affect reversibility potential.

Quantifying “Return to Baseline”: Realistic Expectations.

If a patient’s current functional capacity is 20% of premorbid, what can interventions achieve?

Pessimistic scenario (damage control only) Interventions prevent further decline, stabilizing at 20% indefinitely. Crash mitigation allows better quality of life within severe constraints but no capacity recovery. Realistic expectation: Prevent progression from moderate to severe, avoid complete disability.

Moderate scenario (partial recovery) Interventions restore 20% → 40–50% capacity by addressing reversible components (NAD⁺ depletion, antioxidant deficiency, mitochondrial turnover inefficiency). Irreversible damage (mtDNA mutations, permanent structural changes) sets ceiling. Realistic expectation: Shift from housebound to limited community function, bedbound to housebound.

Optimistic scenario (substantial recovery) If dysfunction is primarily suppressible (WASF3-type mechanism) rather than permanent damage, interventions restore 20% → 70–80% capacity. Realistic expectation: Return to limited work, independent living, meaningful quality of life—but not full premorbid function. Remaining 20–30% deficit from irreversible components.

Recovery scenario (currently speculative) Complete reversal requires addressing root cause (autoantibody removal, chronic infection clearance, resetting dysregulated systems). If achieved, 20% → 95%+ recovery possible. Examples: Daratumumab responders [67], BC007 responders (if replicated), spontaneous remissions. Realistic expectation: Subset of patients, not universal.

Time Horizons for Recovery Attempts.

Raising baseline requires sustained intervention over mitochondrial turnover timescales:

- **Minimum trial duration:** 3–6 months (multiple complete mitochondrial replacement cycles at 10–15 days each)
- **Expected trajectory:** Gradual improvement (baseline crashes less severe → activity threshold slowly rises → functional capacity incrementally increases)
- **Plateau indicators:** If no improvement after 6–12 months of optimized intervention, likely at irreversible damage ceiling for current medical technology
- **Relapse after improvement:** Suggests suppressible mechanism requiring ongoing intervention (like WASF3 suppression) rather than permanent cure

Individual Variation in Recovery Potential.

Not all patients have equal reversibility potential. Predictors of higher recovery ceiling may include:

- Recent onset (<2 years): Less accumulated permanent damage
- Post-viral trigger: Clear initiating mechanism potentially addressable
- Younger age: Greater cellular regeneration capacity
- Moderate vs. very severe: Severe patients may have crossed irreversibility threshold
- Documented reversible mechanism: WASF3+ patients, autoantibody+ patients with immunoadsorption access
- Minimal comorbidities: Fewer compounding factors

Conversely, very long duration (>10 years), very severe baseline (<10% function), older age (>60), and multiple comorbidities suggest lower ceiling, though individual cases defy prediction.

The Unanswered Core Question.

“How much can we bring back to baseline?” depends entirely on *what percentage of dysfunction is reversible vs. permanent*—currently unknown for individual patients. No biomarker currently predicts reversibility potential. Clinical trial data with aggressive multi-modal interventions sustained for 6–12 months in well-characterized patient cohorts is desperately needed to answer this question empirically.

The therapeutic imperative is attempting restoration despite uncertainty: even 20% → 35% improvement (modest by percentage) represents transformative quality of life change (bed-bound → housebound with mobility). Given low risk of evidence-based interventions (NAD⁺ precursors, CoQ10, careful activity progression), attempting baseline restoration is justified even when complete recovery is unlikely.

The Temporary-to-Permanent Transition: How Acute Illness Becomes Chronic ME/CFS.

A critical question is why some individuals recover from post-viral fatigue while others develop chronic ME/CFS. The answer likely involves vicious cycles that become self-perpetuating when repair capacity is overwhelmed.

Normal recovery trajectory (healthy individuals):

Any significant stressor—viral infection (influenza, mononucleosis, COVID-19), bacterial infection, major surgery, severe physical/emotional trauma, or sustained overexertion—temporarily creates ME/CFS-like physiology:

1. **Acute phase (days 0–7):** Immune response consumes massive ATP; cytokine production induces sickness behavior; mitochondrial damage from ROS; exercise threshold dramatically lowered
2. **Recovery phase (weeks 2–6):** Immune resolution; mitochondrial turnover replaces damaged organelles; ATP pools replenish; threshold gradually normalizes
3. **Full recovery (weeks 6–12):** Return to premorbid baseline capacity; linear dose-response restored; adaptive systems functional

2 Core Symptoms

This trajectory requires intact repair systems operating faster than damage accumulation. Recovery succeeds when mitochondrial biogenesis outpaces damage, immune regulation prevents sustained activation, and ATP regeneration exceeds consumption.

Failed recovery trajectory (ME/CFS development):

In susceptible individuals, the same acute stressor initiates a self-perpetuating cycle:

1. **Initial trigger phase (days 0–7):** Identical acute stress response as healthy individuals—severe fatigue, immune activation, mitochondrial damage, lowered threshold
2. **Critical window (weeks 2–8):** Repair systems fail to outpace ongoing damage
 - Patient attempts normal activity resumption (“I should be better by now”)
 - Each activity episode causes new damage while previous damage unrepaired
 - Mitochondrial removal outpaces biogenesis → net mitochondrial loss
 - Immune system remains activated (autoantibody development? persistent viral antigens? dysregulated signaling?)
 - ATP depletion prevents repair protein synthesis → repair systems themselves fail
3. **Vicious cycle establishment (weeks 8–24):** System enters stable dysfunctional equilibrium
 - Low baseline ATP → Impaired mitochondrial biogenesis (requires ATP)
 - Damaged mitochondria → Low ATP production → Cannot clear damaged mitochondria
 - Immune dysregulation → Inflammation → Mitochondrial damage → More immune activation
 - Activity → Exceeds capacity → Crash → Lower capacity → Activity threshold drops further
 - NAD⁺ depletion → PARP consumes NAD⁺ for DNA repair → Cannot make ATP → More DNA damage
4. **Progressive decline (months to years):** Each crash causes incremental permanent damage
 - Cumulative mitochondrial loss exceeds replacement capacity
 - Potential mtDNA mutations accumulate (damaged mitochondria replicate errors)
 - Epigenetic silencing of metabolic genes (chronic stress response becomes permanent)
 - Autoantibody titers increase if autoimmune component present
 - Autonomic dysfunction develops from chronic hypoperfusion
 - Multi-system symptoms emerge as dysfunction spreads beyond initial metabolic impairment

Why some individuals cannot escape the cycle:

The transition from temporary to permanent dysfunction likely requires multiple factors:

Genetic susceptibility Variants affecting mitochondrial function (WASF3 regulation), immune regulation (HLA types), or metabolic efficiency (NAD⁺ synthesis enzymes) reduce

recovery capacity baseline. Same stressor that healthy person recovers from overwhelms genetically vulnerable system.

Severity of initial insult Massive viral load, severe infection, or combined stressors (infection + surgery + psychological trauma) cause damage exceeding any individual's repair capacity. Even robust systems cannot recover when damage is catastrophic.

Premature activity resumption Attempting normal activity during critical repair window (weeks 2–8) prevents recovery. Each exertion episode creates new damage while previous damage unrepaired, progressively widening the repair deficit until systems collapse into vicious cycle. This mechanism explains clustering of ME/CFS in high-achievers who “push through” illness rather than resting.

Ongoing immune activation If immune system develops autoantibodies during acute infection (molecular mimicry between viral proteins and self-antigens), or if viral fragments persist triggering continuous immune response, the initial trigger never fully resolves. System attempts to recover while inflammation continues, making recovery impossible.

Secondary metabolic traps NAD⁺ depletion, thiamine deficiency, or other metabolic cofactor depletions create secondary bottlenecks. Even if initial trigger resolves, depleted cofactor pools prevent metabolic recovery, maintaining dysfunction indefinitely.

Age and baseline reserve Younger individuals with greater cellular regeneration capacity and larger metabolic reserves can tolerate more damage before crossing into irreversible territory. Older individuals or those with pre-existing subclinical dysfunction have narrower margins—same stressor that 25-year-old recovers from causes 55-year-old to develop chronic ME/CFS.

Clinical implications for prevention:

If ME/CFS develops when temporary threshold depression becomes permanent through failed recovery, aggressive intervention during the critical window (weeks 2–8 post-acute illness) might prevent chronicity:

- **Absolute rest** during acute phase and early recovery (0–6 weeks)—no activity beyond essential self-care
- **Energy substrate support** (D-ribose, NAD⁺ precursors, CoQ10) to support repair systems during recovery
- **Anti-inflammatory support** (if appropriate) to prevent sustained immune activation
- **Gradual activity resumption** only after 6+ weeks complete rest, starting at 20–30% premorbid capacity
- **Immediate cessation** if any post-exertional symptoms develop, indicating threshold not yet normalized
- **Extended timeline acceptance** (12+ weeks for full recovery from severe infections rather than 2–4 weeks)

This prevention strategy remains unvalidated by clinical trials but follows logically from the vicious cycle model. Anecdotal reports from ME/CFS patients commonly include “I got sick and tried to push through it” or “I went back to work too soon after mono,” suggesting premature activity resumption during critical repair window as potential causal factor.

Why this model matters:

Understanding ME/CFS as “normal post-viral physiology that failed to resolve” rather than unique pathological entity has profound implications:

1. **Validates patient experience:** Not “weak” or “malingering”—experienced what everyone experiences post-illness, but repair systems failed
2. **Explains heterogeneity:** Different triggers (viral, trauma, overtraining) create similar dysfunction through final common pathway of failed metabolic recovery
3. **Suggests prevention strategies:** Rest during acute illness and gradual activity resumption might prevent chronicity
4. **Implies reversibility potential:** If dysfunction is maintained by vicious cycles rather than permanent structural damage, breaking cycles might allow recovery
5. **Explains why some recover:** Spontaneous remissions occur when vicious cycles spontaneously break (unknown mechanism) or when time allows ultra-slow repair to eventually succeed

The fundamental insight: ME/CFS may represent the body “stuck” in a temporary protective state (low-energy, inflammation, restricted activity) that should resolve within weeks but becomes permanent when repair systems cannot overcome the initial damage before new damage accumulates. Every healthy person who experiences severe post-viral fatigue is temporarily in an ME/CFS-like state; the disease develops in those unable to escape it.

⚠ **Limitation 1: PEM Vicious Cycle Model: Illustrative Framework, Not Quantified Biology**

The vicious cycle dynamics model below (cycle gain, sequential entry, ratchet effect) provides a conceptually useful framework for understanding disease progression, but its quantitative parameters are illustrative, not empirically derived:

- Numerical values for “cycle gain” (5–10% vs. 30–50% ATP reduction per iteration), “repair capacity” (15–20% biogenesis rate), and “critical thresholds” are hypothetical estimates chosen to illustrate the model’s logic, not measurements from ME/CFS patients.
- The distinction between “escapable” and “inescapable” cycles has not been validated by longitudinal studies tracking cycle parameters over time in individual patients.
- Whether ME/CFS progression truly follows a sequential cycle-recruitment pattern or reflects a single initial insult with variable downstream consequences is undetermined.
- The model’s predictions (earlier intervention = better outcomes, cycle number predicts severity) are consistent with clinical observation but have not been prospectively tested.

Vicious Cycle Dynamics: Cycle Strength, Sequential Entry, and the Ratchet Effect.

The preceding description introduces “vicious cycles” as the mechanism maintaining chronic dysfunction. However, not all vicious cycles are equally “vicious.” A deeper analysis of cycle

dynamics reveals three critical concepts: (1) cycle strength and escapability, (2) sequential entry into multiple reinforcing cycles, and (3) the ratchet effect of irreversible cumulative damage. Understanding these mechanisms explains why some patients stabilize while others progressively deteriorate, and why disease severity correlates with duration.

Cycle strength and escapability: Mathematical foundations of self-perpetuating dysfunction

Vicious cycles in biological systems can be characterized by their *cycle gain*—the degree to which dysfunction in one component amplifies dysfunction in another component within the loop. Systems with cycle gain below a critical threshold remain escapable (the body's repair systems can eventually outpace damage accumulation), while cycles with gain exceeding this threshold become self-perpetuating traps [68].

Weak/escapable cycles (cycle gain < critical threshold): In post-viral fatigue that resolves spontaneously, temporary vicious cycles exist but remain escapable:

- Low ATP → Mild mitochondrial impairment → Slightly reduced ATP production
- **Gain factor:** Each turn of the cycle reduces ATP by 5–10% (hypothetical)
- **Repair capacity:** Mitochondrial biogenesis can increase production by 15–20% when given adequate rest
- **Outcome:** Net positive—repair outpaces degradation, system gradually escapes cycle
- **Timeline:** 4–12 weeks of rest allows full recovery

The critical feature: the amplification factor per cycle iteration is less than unity when repair processes are considered. Even though ATP depletion impairs mitochondrial function, the impairment is modest enough that partial function remains, allowing gradual recovery.

Strong/inescapable cycles (cycle gain > critical threshold): In established ME/CFS, cycle gain exceeds the threshold where repair can compensate:

- Severe ATP depletion → Catastrophic mitochondrial dysfunction → Dramatically lower ATP production
- **Gain factor:** Each turn reduces ATP by 30–50% (hypothetical)
- **Repair capacity:** Mitochondrial biogenesis itself requires ATP; at severe depletion, biogenesis rate falls to 5–10% of normal
- **Outcome:** Net negative—degradation accelerates faster than repair, system locks into dysfunction
- **No spontaneous escape:** Without external intervention breaking the cycle, dysfunction persists indefinitely

Formally, let $G = \frac{D_{n+1}}{D_n}$ be the cycle gain, where D_n represents the dysfunction level after n iterations of the cycle. If $G < 1$, dysfunction diminishes over iterations (repair dominates). If $G > 1$, dysfunction amplifies (damage dominates). The system transitions from escapable to inescapable when the cycle gain crosses unity ($G = 1$).

Equivalently, the mathematical transition point occurs when:

Damage accumulation rate per cycle > Repair capacity per cycle

Once this threshold is crossed ($G > 1$), the system enters a stable dysfunctional equilibrium. The “stability” here is pathological—the dysfunction self-maintains because the very processes needed for repair are themselves impaired by the dysfunction.

What determines cycle strength? Several factors influence whether an individual’s post-viral vicious cycles remain escapable or become inescapable:

Genetic reserve capacity Individuals with variants affecting mitochondrial biogenesis efficiency (e.g., WASF3 regulation [2]), NAD⁺ synthesis capacity [69], or antioxidant systems start with different baseline repair capacities. A genetic variant reducing mitochondrial biogenesis by 30% means the same viral insult creates stronger cycle gain.

Severity of initial trigger Massive viral load or severe infection causes more extensive initial damage. Damage to 80% of mitochondrial population creates stronger cycle gain than damage to 40% because insufficient functional mitochondria remain to support repair.

Ongoing stressors during recovery Continued exertion, concurrent infection, or psychological stress during the critical window (weeks 2–8) increases cycle gain by adding new damage while repair is attempted. Each additional stressor pushes the gain factor higher.

Metabolic cofactor availability Adequate NAD⁺, CoQ10, B vitamins, and other metabolic cofactors reduce cycle gain by supporting repair processes. Depletion of these cofactors amplifies cycle strength.

Immune regulation capacity If immune dysregulation develops (autoantibody production, chronic activation), inflammation creates a parallel cycle that reinforces metabolic dysfunction. The combined gain of coupled cycles exceeds what either cycle alone would produce.

The critical clinical implication: interventions during early disease (first 6–24 months) may reduce cycle gain enough to shift from inescapable to escapable territory. Aggressive pacing reduces damage accumulation rate; NAD⁺ precursors and CoQ10 support repair capacity; immunomodulation (if appropriate) dampens inflammatory amplification. Even modest gain reduction—from 1.3× amplification per cycle to 0.9×—transforms the trajectory from progressive decline to potential recovery.

Sequential entry into multiple vicious cycles: The multi-lock model

A patient does not necessarily enter all pathological cycles simultaneously. **Hypothesized model:** The disease *may* progress through sequential recruitment of additional vicious cycles, with each crash or period of overexertion pushing the patient across new thresholds into previously inactive cycles [70]. While the Maksoud 2020 natural history study documents that ME/CFS severity increases with duration and symptom domains expand over time, the specific ordering proposed below represents one plausible sequence consistent with clinical observations—other orderings may occur depending on individual pathophysiology, genetic factors, and environmental triggers.

Hypothesized progressive cycle recruitment model:

1. **Stage 1: Mitochondrial cycle only (early disease, weeks 8–24)**
 - Primary dysfunction: ATP depletion ↔ Impaired mitochondrial function
 - **Symptoms:** Fatigue, PEM with recovery in days to 1–2 weeks
 - **Severity:** Mild ME/CFS; can work with significant difficulty
 - **Reversibility:** Still potentially escapable with strict pacing and metabolic support
 - **Threshold to next stage:** Repeated crashes cause cumulative mitochondrial loss
2. **Stage 2: Mitochondrial + Immune cycles (months 6–18)**
 - **Trigger for entry:** Cumulative oxidative stress from repeated PEM episodes activates chronic immune response; potential autoantibody development against oxidatively modified proteins or development of GPCR autoantibodies
 - **New cycle:** Immune activation ↔ Inflammation ↔ Mitochondrial damage
 - **Symptoms:** Fatigue worsens; flu-like symptoms emerge; sensory sensitivities develop
 - **Severity:** Moderate ME/CFS; housebound significant fraction of time
 - **Mutual reinforcement:** Mitochondrial dysfunction impairs immune cell function (T cells, NK cells require high ATP); immune inflammation damages mitochondria further
 - **Threshold to next stage:** Chronic inflammation and hypoperfusion stress autonomic nervous system
3. **Stage 3: Mitochondrial + Immune + Autonomic cycles (years 1–3)**
 - **Trigger for entry:** Chronic hypoperfusion from reduced activity; blood volume depletion; potential GPCR autoantibodies affecting β_2 -adrenergic receptors [71, 72]
 - **New cycle:** Autonomic dysfunction ↔ Orthostatic intolerance ↔ Cerebral hypoperfusion ↔ Reduced activity capacity
 - **Symptoms:** POTS develops; orthostatic intolerance limits upright time; brain fog worsens significantly
 - **Severity:** Moderate to severe ME/CFS; bedbound significant hours daily
 - **Mutual reinforcement:** Mitochondrial dysfunction impairs vascular smooth muscle function; autonomic dysfunction reduces perfusion to all tissues including muscle (worsening ATP depletion); immune activation may directly affect autonomic signaling
 - **Threshold to next stage:** Chronic central nervous system hypoperfusion and immune mediator exposure
4. **Stage 4: Mitochondrial + Immune + Autonomic + Neuroinflammation cycles (years 2–5+)**
 - **Trigger for entry:** Chronic cerebral hypoperfusion; peripheral immune activation with cytokine penetration across blood-brain barrier; microglial activation [73]
 - **New cycle:** Neuroinflammation ↔ Central sensitization ↔ Sensory hypersensitivity ↔ Cognitive dysfunction ↔ Sympathetic activation
 - **Symptoms:** Severe sensory sensitivities (light, sound, chemical); profound cog-

nitive impairment; central pain amplification; severe anxiety from dysregulated threat perception

- **Severity:** Severe ME/CFS; mostly bedbound, dark quiet environment required
- **Mutual reinforcement:** Neuroinflammation amplifies autonomic dysfunction (brainstem involvement); cognitive dysfunction impairs ability to pace effectively (worsening all other cycles); central sensitization lowers threshold for all symptom triggers
- **Threshold to next stage:** Chronic HPA axis dysregulation

5. Stage 5: All cycles + Endocrine dysregulation (years 3–10+)

- **Trigger for entry:** Chronic stress response from years of severe illness; HPA axis exhaustion; potential thyroid dysfunction from chronic inflammation
- **New cycle:** Endocrine dysfunction ↔ Metabolic dysfunction ↔ Immune dysfunction ↔ Central dysfunction
- **Symptoms:** Hormonal dysregulation; cortisol abnormalities; thyroid dysfunction in subset; temperature regulation failure; severe multi-system involvement
- **Severity:** Very severe ME/CFS; bedbound, minimal self-care capacity
- **Mutual reinforcement:** Cortisol abnormalities affect immune function and metabolism; thyroid dysfunction affects mitochondrial function; all hormonal systems interact with previously established cycles
- **Potential irreversibility:** Five mutually reinforcing cycles create extremely high combined cycle gain; breaking any single cycle insufficient to allow escape

Key insights from sequential entry model:

- **Severity correlates with number of active cycles:** Mild disease (1–2 cycles) remains potentially escapable; severe disease (4–5 cycles) may be irreversible even with aggressive intervention
- **Early intervention targets fewer cycles:** Treating a patient with only mitochondrial dysfunction requires breaking one cycle; treating very severe disease requires simultaneously breaking five reinforcing cycles
- **Symptoms expand as cycles accumulate:** Early disease presents primarily with fatigue/PEM; established disease shows multi-system involvement reflecting recruitment of immune, autonomic, neurological, and endocrine cycles
- **Critical intervention windows exist:** Preventing entry into Stage 2 (immune cycle) and Stage 3 (autonomic cycle) may prevent progression to severe disease; once 4–5 cycles are established, reversal becomes exponentially more difficult
- **Exertion accelerates cycle recruitment:** Each crash/period of overexertion increases the probability of crossing the next threshold (e.g., pushing a patient with 2 active cycles into activating a 3rd cycle)
- **Treatment must address multiple cycles:** Single-target interventions (e.g., treating only mitochondrial dysfunction while ignoring immune and autonomic cycles) fail because untreated cycles continue to drive dysfunction

This sequential model aligns with the multi-lock hypothesis (Section 14.16) and the five-domain biological phenotyping framework (Section 4.7.3). The observation that most ME/CFS patients show dysfunction in 3+ biological domains [12] supports the concept that chronic established disease involves multiple simultaneously active vicious cycles.

The ratchet effect: Irreversible cumulative damage from repeated crashes

Even if vicious cycles could theoretically be broken, ME/CFS progression may involve *irreversible structural damage* that accumulates with each crash episode. This “ratchet effect” means that each severe PEM episode moves the baseline functional capacity downward permanently, preventing full recovery even when triggering factors are removed (detailed analysis in Section 5.4.3) [74, 70].

Mechanisms of irreversible damage:

Net mitochondrial loss During severe ATP depletion, damaged mitochondria undergo mitophagy (selective autophagy) [64]. If ATP levels remain too low to support mitochondrial biogenesis (which itself requires substantial ATP and functional translation machinery), mitophagy removal exceeds biogenesis replacement. Result: net permanent loss of mitochondrial population. A muscle fiber that previously contained 1000 mitochondria may be reduced to 700 after multiple crashes, permanently reducing ATP production capacity.

Cumulative mtDNA mutations Mitochondrial DNA lacks the robust repair mechanisms of nuclear DNA. Replication errors and repair failures during chronic cellular stress can lead to accumulation of mtDNA mutations through clonal expansion [75]. Damaged mitochondria with mutated mtDNA continue to replicate, expanding the population of dysfunctional organelles. Over time, the fraction of mitochondria with functional respiratory chains decreases, permanently impairing oxidative metabolism.

Epigenetic silencing of metabolic genes Chronic cellular stress induces protective epigenetic changes (DNA methylation, histone modifications) that silence genes involved in oxidative metabolism [76]. Initially adaptive (reducing metabolic demand during crisis), these modifications can become stable over time. After years of illness, metabolic genes may be epigenetically locked in a silenced state, preventing normal mitochondrial function even if the initial trigger is removed.

Autoantibody accumulation If molecular mimicry or exposure of neo-epitopes during tissue damage initiates autoantibody production, these antibodies persist for months to years [71, 72]. Long-lived plasma cells in bone marrow continue producing GPCR autoantibodies (anti- β_2 -adrenergic, anti-M3/M4 muscarinic) indefinitely, creating permanent autonomic and metabolic dysfunction. Each crash that triggers additional immune activation may generate new autoantibody specificities, progressively expanding the autoimmune repertoire.

Vascular endothelial remodeling Repeated ischemia-reperfusion injury during crashes causes endothelial dysfunction [12]. Chronic elevation of von Willebrand factor, fibronectin, and thrombospondin-1 indicates ongoing endothelial activation and potential structural changes to the microvasculature. Over time, vascular remodeling may become permanent, limiting perfusion even when other factors improve.

Central sensitization establishment Repeated microglial activation creates self-perpetuating neuroinflammation [73]. Activated microglia secrete inflammatory mediators that keep neighboring microglia activated, establishing a stable inflammatory state in the central nervous system. Additionally, central sensitization—amplified pain and sensory processing—involves synaptic plasticity changes that stabilize over time. After years of sensory sensitization, these changes may resist reversal.

Crossing cycle thresholds Each crash increases the probability of entering a new vicious cycle (as described in sequential entry model). Once a patient crosses from 2 active cycles to 3 active cycles, the combined cycle gain increases multiplicatively. This represents a discrete permanent worsening: the patient is now trapped in a more complex multi-cycle system.

The “crash limit” hypothesis: Patient communities report anecdotal evidence of a threshold number of severe crashes (estimated at 5–10 major episodes) beyond which recovery capacity is permanently impaired [74]. While this specific threshold lacks formal validation, the underlying biological mechanism is plausible:

- **Observation 1:** Recovery time from crashes lengthens with each successive crash (crash 1 requires 3 days recovery; crash 5 requires 3 weeks; crash 10 requires 3 months)
- **Observation 2:** After a certain number of severe crashes, patients stop recovering to previous baseline entirely—each crash leaves them at a lower functional floor
- **Observation 3:** Some patients report that a single catastrophic overexertion event (running a marathon while ill, severe infection combined with overwork, major surgery without adequate recovery time) triggered irreversible severe worsening
- **Mechanistic interpretation:** Each crash causes *some* irreversible damage (e.g., 5–10% permanent mitochondrial loss, small increase in mtDNA mutation load, modest autoantibody titer increase). Early crashes can be partially compensated for by remaining reserve capacity. After 5–10 crashes, cumulative damage exceeds the compensation threshold, and the system collapses into severe permanent dysfunction.

If this hypothesis is correct, the therapeutic imperative becomes: **prevent all severe crashes**, not merely reduce their frequency. The goal is zero major PEM episodes during the critical first 2–3 years of illness, preventing accumulation of irreversible damage.

Evidence supporting the cycle dynamics and ratchet effect model:

Multiple independent lines of evidence support the concepts of escalating cycle strength, sequential cycle entry, and cumulative irreversible damage:

Evidence 1: Progressive decline with continued exertion Patients who continue high levels of activity (working full-time, attempting graded exercise therapy, “pushing through” symptoms) show progressive worsening over time [74]. This contrasts with progressive improvement expected if dysfunction were purely functional/reversible. The pattern of worsening despite effort suggests:

- Each exertion episode causes net damage exceeding repair

2 Core Symptoms

- Cumulative damage accumulates, progressively reducing baseline capacity
- The system cannot escape vicious cycles without external intervention reducing damage rate

Conversely, aggressive pacing (staying well within energy envelope) is associated with stabilization or modest improvement [61], suggesting that preventing crashes prevents progression even if it doesn't reverse established dysfunction.

Evidence 2: Severity correlates with disease duration Cross-sectional studies show that longer illness duration predicts greater severity and functional impairment [77]:

- Illness duration < 2 years: 60–70% mild-moderate, 30–40% severe
- Illness duration 2–5 years: 40–50% mild-moderate, 50–60% severe
- Illness duration > 10 years: 20–30% mild-moderate, 70–80% severe (approximate estimates from clinical populations)

This temporal progression is consistent with:

- Sequential recruitment of additional vicious cycles over time
- Cumulative irreversible damage accumulating with each year of illness
- Progressive transition from escapable early dysfunction to inescapable multi-cycle traps

Evidence 3: Individual crashes cause permanent worsening Patient reports and clinical observations document that specific severe crashes lead to discrete permanent reductions in functional capacity [74]:

- “I was moderate, had one really bad crash after a wedding, and have been severe ever since”
- “I pushed through a work deadline, crashed for 3 months, and never returned to my previous baseline”
- “Each time I try to increase my activity level, I crash and end up worse than before I started”

This pattern is inconsistent with purely reversible dysfunction and supports cumulative irreversible damage. If dysfunction were entirely maintained by active processes (e.g., ongoing immune activation), removal of the trigger should allow recovery to previous baseline. Instead, crashes cause discrete stepwise reductions in capacity, consistent with structural damage (mitochondrial loss, mtDNA mutations, vascular remodeling, central sensitization).

Evidence 4: Multi-system symptoms develop sequentially Longitudinal patient reports indicate that symptoms expand over time rather than presenting fully formed at onset [70]:

- **Early disease (months 0–12):** Primarily fatigue and PEM
- **Intermediate disease (years 1–3):** Cognitive dysfunction, orthostatic intolerance, sensory sensitivities emerge

- **Established disease (years 3+):** Pain amplification, severe sensory sensitivities, dysautonomia, potential MCAS, gut dysfunction, severe multi-system involvement

This temporal sequence supports sequential cycle entry: initial mitochondrial dysfunction → immune activation → autonomic dysfunction → neuroinflammation → endocrine dysfunction. Patients do not start with all five cycles active; they progressively recruit additional cycles as disease duration increases and cumulative damage accumulates.

Evidence 5: Remission rates decrease with illness duration Recovery rates show strong inverse correlation with illness duration [77]:

- Illness duration < 2 years: 10–20% achieve remission or significant improvement
- Illness duration 2–5 years: 5–10% achieve remission
- Illness duration > 10 years: <1–2% achieve remission

This pattern is consistent with:

- Early disease involving fewer active cycles (1–2) that remain potentially reversible
- Established disease involving more cycles (3–5) with lower probability of simultaneous resolution
- Cumulative irreversible damage increasing over time, reducing the ceiling for potential recovery
- Critical intervention window in first 2 years before epigenetic changes, extensive mitochondrial loss, and multi-cycle entrenchment

Evidence 6: Pediatric vs. adult outcomes Children and adolescents with ME/CFS show dramatically better recovery rates than adults: 68% recovery by 10 years in pediatric cohorts [78] versus <5% in adult cohorts. Potential explanations include:

- **Greater baseline reserve:** Children have higher mitochondrial biogenesis capacity, greater cellular regeneration potential, larger metabolic reserves—all factors reducing cycle gain
- **Earlier intervention:** Pediatric cases often receive school accommodations (mandatory rest, reduced workload) earlier than adults receive workplace accommodations, preventing cumulative damage during critical window
- **Fewer active cycles:** Children may not progress as far through sequential cycle recruitment before spontaneous recovery mechanisms succeed
- **Less irreversible damage:** Shorter illness duration and better pacing reduces cumulative mitochondrial loss, mtDNA mutations, and epigenetic changes
- **Lower autoantibody burden:** Shorter exposure to chronic immune activation may result in lower autoantibody titers and less permanent autoimmune component

The stark difference in outcomes between pediatric and adult populations supports the concept that *duration matters* and that *early aggressive intervention* (whether deliberate or imposed by school systems) prevents progression into irreversible multi-cycle severe disease.

Clinical implications: Preventing progressive cycle entrenchment

Understanding vicious cycle dynamics, sequential recruitment, and irreversible cumulative damage transforms clinical management:

1. **The primary goal is preventing crashes, not managing crashes:** If each severe PEM episode causes permanent incremental damage, the therapeutic imperative is absolute avoidance, not damage limitation. Patients must operate at 50–70% of perceived capacity, leaving substantial margin to prevent envelope violation [61].
2. **Early intervention is critical:** Treating mild disease (1–2 active cycles) has far higher success probability than treating severe disease (4–5 cycles). The first 6–24 months represent the optimal window for preventing cycle entrenchment and irreversible damage.
3. **Multi-target interventions address multiple cycles:** Single-domain treatments (e.g., CoQ10 alone, or immunoadsorption alone) may fail because untreated cycles maintain overall dysfunction. Optimal approach: simultaneous intervention in all accessible domains (mitochondrial support + immunomodulation if appropriate + autonomic support + pacing).
4. **Aggressive pacing prevents both functional decline and structural damage:** Pacing is not merely symptom management—it prevents the cumulative irreversible damage that drives progression from mild to severe disease.
5. **Recovery potential decreases with duration:** Patients with >2–5 years duration may have crossed irreversibility thresholds (extensive mitochondrial loss, stable epigenetic silencing, entrenched multi-cycle traps). Treatment goals shift from “cure” to “stabilization and optimization within constraints.”
6. **The 6-month and 2-year thresholds are not arbitrary:** Six months marks transition from potentially self-resolving post-viral fatigue to established ME/CFS (failed spontaneous cycle escape). Two years marks transition from hypermetabolic (potentially reversible) to hypometabolic (potentially irreversible) state with epigenetic changes and immune exhaustion [70]. Intervention before these thresholds offers the best probability of preventing permanent severe disease.

The fundamental insight: ME/CFS progression is not inevitable—it results from cumulative damage accumulation driven by repeated envelope violations during a disease state where repair capacity is impaired. Preventing this progression requires recognizing that each crash matters, that early disease is more treatable than late disease, and that multi-system dysfunction requires multi-target intervention. The patients who progress to very severe bedbound disease did not develop a “different disease”—they progressed further through the sequential cycle recruitment process and accumulated more irreversible damage. This progression is often preventable through aggressive early intervention, but becomes increasingly difficult to reverse as duration increases and cycles entrench.

2.1.3 Measurement and Assessment

Objective Measurement via Two-Day Cardiopulmonary Exercise Testing

Observation 2 (Two-Day CPET: Objective PEM Measurement). Two-day cardiopulmonary exercise testing (CPET) provides objective evidence for post-exertional malaise through repeated maximal exercise tests separated by 24 hours [79]. Meta-analysis of five studies (n=98 ME/CFS patients, n=51 controls) demonstrated that ME/CFS patients fail to reproduce Day 1 performance on Day 2, whereas healthy sedentary controls maintain or improve performance. The most sensitive metric, workload at ventilatory threshold (VT), showed significant deterioration in ME/CFS patients (mean change from baseline: -33.0W on Day 2 vs. -10.8W on Day 1, $p < 0.05$) while controls demonstrated improvement. This pattern has been independently replicated in subsequent larger cohorts exceeding 150 patients [1], establishing 2-day CPET as the gold standard for objective PEM documentation.

The physiological mechanisms underlying the Day 2 deterioration include:

- **ATP depletion:** Mitochondrial dysfunction prevents normal energy regeneration within 24 hours [64, 2]
- **Immune activation:** Exercise triggers pro-inflammatory cytokine release that persists beyond the immediate post-exercise period
- **Oxidative stress:** Reactive oxygen species accumulate faster than antioxidant systems can neutralize them
- **Anaerobic threshold shift:** Early shift to anaerobic metabolism indicates impaired mitochondrial oxidative capacity
- **Prolonged recovery:** Unlike healthy controls who recover within 48 hours, ME/CFS patients may require 13+ days to return to baseline [1]

~ Hypothesis 2: 2-Day CPET as Diagnostic Tool

Two-day CPET may serve as an objective diagnostic biomarker for ME/CFS, particularly for distinguishing genuine post-exertional malaise from deconditioning or other fatiguing conditions [79]. The consistent Day 2 deterioration pattern appears specific to ME/CFS, with sedentary controls, fibromyalgia patients, and depression patients not exhibiting this phenotype. However, larger validation studies comparing ME/CFS to comprehensive disease control groups are needed to establish clinical sensitivity, specificity, and diagnostic thresholds before 2-day CPET can be implemented as a standalone diagnostic test.

Clinical Assessment Tools

While 2-day CPET provides objective measurement, it remains research-grade and inaccessible to most clinicians. Patient-reported outcome measures remain essential for clinical practice:

- **DePaul Symptom Questionnaire (DSQ):** Validated tool specifically measuring PEM frequency and severity

- **Pacing diaries:** Patient tracking of activity-symptom relationships
- **Functional capacity scales:** Bell Disability Scale, SF-36, and ME/CFS-specific measures
- **Activity monitors:** Actigraphy to objectively measure movement patterns (though cannot distinguish voluntary pacing from incapacity)

2.2 Unrefreshing Sleep

Unrefreshing sleep is a cardinal symptom of ME/CFS, reported by 95–100% of patients in most cohorts [80, 81]. Despite sleeping adequate or even excessive hours, patients wake feeling as exhausted as when they went to bed. This distinguishes ME/CFS sleep dysfunction from simple insomnia, where patients feel better after sleep even if it takes time to fall asleep.

2.2.1 Sleep Dysfunction Patterns

ME/CFS patients experience multiple overlapping sleep disturbances:

Unrefreshing Sleep Despite Adequate Duration

The core feature is lack of restoration from sleep:

- Patients may sleep 8–12+ hours yet wake completely unrefreshed
- Morning exhaustion equal to or worse than evening exhaustion
- No correlation between sleep duration and daytime function
- Paradox: Some patients feel better with *less* sleep (4–6 hours) than with full nights

Sleep Maintenance Problems

Beyond non-restorative sleep, many patients experience:

- **Frequent nocturnal awakenings:** Waking 5–20+ times per night
- **Light, fragmented sleep:** Unable to maintain continuous deep sleep
- **Delayed sleep phase:** Inability to fall asleep until 2–4 AM despite exhaustion
- **Reversed circadian rhythm:** Sleeping during day, awake at night (in severe cases)
- **“Tired but wired”:** Physical exhaustion but mental hyperarousal preventing sleep

Sleep Inertia and Hypersomnia

Some patients experience:

- **Severe sleep inertia:** Taking 2–4 hours to become functional after waking
- **Hypersomnia:** Sleeping 12–16 hours per day, particularly during crashes
- **Inability to wake:** Sleeping through alarms, phone calls, physical touch
- **Nap non-restoration:** Naps fail to provide refreshment (unlike healthy fatigue)

2.2.2 Polysomnography Findings

Objective sleep studies in ME/CFS reveal measurable abnormalities:

Sleep Architecture Disruption

Studies have documented [82, 83]:

- **Reduced slow-wave sleep (Stage N3):** The deepest, most restorative sleep stage is diminished
- **Alpha-delta sleep:** Intrusion of waking alpha waves (8–13 Hz) into delta sleep, preventing deep sleep [84]
- **Increased sleep fragmentation:** More frequent stage transitions and microarousals
- **Reduced sleep efficiency:** Lower percentage of time in bed actually spent asleep
- **REM abnormalities:** Some studies show reduced or disrupted REM sleep

The alpha-delta pattern is particularly notable [84]—the brain shows mixed activity suggesting it never fully enters restorative deep sleep, explaining the subjective experience of “sleeping but not resting.”

Autonomic Dysfunction During Sleep

Polysomnography with additional monitoring reveals:

- **Abnormal heart rate variability:** Reduced parasympathetic tone during sleep
- **Elevated heart rate:** Persistent tachycardia even during sleep
- **Blood pressure instability:** Failure of normal nocturnal dipping
- **Temperature dysregulation:** Abnormal core body temperature curves

Limitations of Standard Polysomnography

Standard sleep studies may appear “normal” in ME/CFS because:

- Sleep stages are scored by visual inspection of 30-second epochs
- Microarousals shorter than 3 seconds are not scored
- Alpha-delta intrusion requires specialized analysis
- Restorative quality cannot be directly measured

Patients often report polysomnography results labeled “normal sleep” despite severe subjective non-refreshment, leading to gaslighting. More detailed spectral analysis or multi-night home monitoring may reveal abnormalities missed by single-night laboratory studies.

2.2.3 Related Sleep Disorders

ME/CFS overlaps with and must be distinguished from primary sleep disorders:

Obstructive Sleep Apnea (OSA)

Sleep apnea can mimic ME/CFS symptoms:

- **Overlap:** Fatigue, unrefreshing sleep, cognitive dysfunction, morning headaches
- **Prevalence:** Affects 10–30% of general population [85, 86, 87]; higher in ME/CFS due to weight gain from inactivity
- **Diagnostic clue:** Witnessed apneas, loud snoring, gasping during sleep
- **Resolution:** CPAP treatment resolves symptoms in true OSA; improves but doesn’t cure comorbid OSA in ME/CFS

Clinical importance: Some patients misdiagnosed with ME/CFS for years experience dramatic improvement with CPAP, indicating primary OSA was the cause. Polysomnography should be standard workup before diagnosing ME/CFS.

Upper Airway Resistance Syndrome (UARS)

A subtler form of sleep-disordered breathing:

- Increased upper airway resistance without frank apneas
- Causes repeated arousals (respiratory effort-related arousals, RERAs)
- May be missed on standard apnea-hypopnea index (AHI)
- Requires esophageal pressure monitoring for diagnosis
- Responds to CPAP or oral appliances

Restless Legs Syndrome (RLS) and Periodic Limb Movement Disorder (PLMD)

Common in ME/CFS:

- **RLS:** Uncomfortable sensations in legs requiring movement to relieve, worse at night
- **PLMD:** Involuntary leg jerks during sleep causing microarousals
- **Prevalence:** Higher in ME/CFS than general population
- **Treatment:** Iron supplementation (if ferritin <75 ng/mL [88]), dopamine agonists, gabapentin

Idiopathic Hypersomnia

Overlapping features:

- Excessive daytime sleepiness despite adequate nighttime sleep
- Sleep inertia lasting hours
- Non-restorative sleep
- Requires Multiple Sleep Latency Test (MSLT) to differentiate from ME/CFS

Circadian Rhythm Disorders

ME/CFS frequently involves circadian disruption:

- **Delayed Sleep-Wake Phase Disorder:** Cannot fall asleep until 2–6 AM
- **Non-24-Hour Sleep-Wake Disorder:** Sleep time progressively delays each day
- **Irregular Sleep-Wake Rhythm:** Fragmented sleep-wake patterns across 24 hours
- May respond to light therapy, melatonin timing, or chronotherapy

2.2.4 Differential Diagnosis Approach

When evaluating unrefreshing sleep in suspected ME/CFS:

1. **Rule out primary sleep disorders first:** Polysomnography, MSLT if indicated
2. **Assess for comorbid conditions:** OSA + ME/CFS can coexist; treat both
3. **Check serum ferritin:** Levels <75 ng/mL may cause RLS/PLMD
4. **Evaluate autonomic function:** Tilt table, heart rate variability
5. **Trial therapeutic interventions:** Response to CPAP, iron, or circadian treatments provides diagnostic information

The key distinction: Primary sleep disorders improve significantly with appropriate treatment (CPAP, iron, etc.), while ME/CFS sleep dysfunction persists despite these interventions, though comorbid treatment helps partially.

2.3 Cognitive Impairment

Cognitive dysfunction, often described as “brain fog,” is a prominent and disabling feature of ME/CFS, affecting 85–95% of patients [89]. Unlike fatigue-related cognitive slowing in healthy individuals, ME/CFS cognitive impairment persists despite rest and worsens substantially following exertion.

2.3.1 Domains of Cognitive Dysfunction

Processing Speed. Processing speed deficits represent the most robust and consistently replicated cognitive finding in ME/CFS. A meta-analysis of 40 studies found large effect sizes for reading speed (Hedges’ $g = -0.82$, $p < 0.0001$) and moderate-to-large effects for other timed tasks [89]. Patients perform 0.5–1.0 standard deviations below healthy controls on processing speed measures, indicating clinically significant impairment. Recent studies using the Stroop task demonstrate that ME/CFS patients show “significantly longer response times than controls indicating cognitive dysfunction” with “global slowing of response times that cannot be overcome by practice” [90].

Attention and Concentration. Patients demonstrate reduced attentional capacity on effortful tasks, with impaired sustained attention during demanding cognitive work [89, 91]. Critically, these deficits persist after controlling for depression and are not explained by psychiatric comorbidity. The constant internal effort required to maintain focus depletes already-limited energy reserves, contributing to cognitive post-exertional malaise.

Memory. Memory impairments follow a specific pattern:

- **Visuospatial immediate memory:** Moderate impairment ($g = -0.55$, $p = 0.007$), with visual modality more affected than verbal [89]
- **Working memory:** Impaired primarily on demanding tasks requiring interference resistance
- **Episodic memory:** Difficulties in storage, retrieval, and recognition processes, though less consistently affected than processing speed
- **Short-term memory:** Variable findings across studies

Executive Function. Executive functions appear relatively preserved compared to processing speed and memory. Meta-analysis found that “executive functions seemed little or not affected and instrumental functions appeared constantly preserved” [89]. However, some patients demonstrate difficulties with mental flexibility, cognitive inhibition, and information generation, particularly under demanding conditions.

Language and Word-Finding. Verbal fluency deficits manifest as word retrieval problems, slowed speech, and linguistic reversals (mixing up word order) [91]. Patients often describe “tip of the tongue” experiences and difficulty with verbal tests of unrelated word association learning and letter fluency. Communication difficulties extend to auditory sequencing problems that impair comprehension of spoken language.

2.3.2 Neuropsychological Testing

Objective Test Results. The Multi-Site Clinical Assessment of ME/CFS (MCAM) study (n=261 ME/CFS patients vs. 165 healthy controls) confirmed deficits in processing speed, attention, working memory, and learning efficiency using standardized neuropsychological batteries [91]. Between 21–38% of patients perform below the 1.5 standard deviation cutoff for clinically significant impairment on Stroop tests.

Pattern of Deficits. The hierarchy of cognitive impairment from most to least affected is:

1. Processing speed (most robust, largest effect sizes)
2. Attention span and working memory (consistently impaired)
3. Immediate memory, especially visual (moderate deficits)
4. Episodic memory (variable across studies)
5. Executive function (relatively preserved)

This pattern differs from depression (which shows more diffuse cognitive effects) and multiple sclerosis (which shows more widespread deficits including greater executive impairment) [92, 93].

Distinction from Depression. Comparative studies demonstrate that ME/CFS cognitive deficits are not attributable to depression. In three-way comparisons of ME/CFS, major depression, and healthy controls, cognitive patterns differ significantly: ME/CFS patients show primary deficits in processing speed and logical memory that persist after controlling for depressive symptoms [92]. Additionally, cognitive performance in ME/CFS does not correlate with fatigue, pain, or depression levels, indicating independent pathophysiology [93].

Subjective-Objective Dissociation. A notable finding is poor correlation between subjective cognitive complaints and objective test performance. Self-reported cognitive dysfunction correlates more strongly with fatigue ($p < 0.001$), pain ($p < 0.001$), and depression ($p < 0.001$) than with actual measured deficits [91]. This suggests subjective complaints reflect overall symptom burden rather than specific cognitive impairments. However, strong concordance exists between subjective mental fatigue complaints and objective cognitive decline following exertion, highlighting the importance of assessing cognition in relation to activity.

2.3.3 Neuroimaging Findings

Functional MRI: Increased Activation. The most consistent fMRI finding is that ME/CFS patients exhibit “increased activations and recruited additional brain regions during cognitive tasks” [94]. This compensatory activation suggests the brain works harder to achieve equivalent performance. Tasks with increasing complexity produce decreased activation in task-specific regions, indicating failure of normal efficiency mechanisms under cognitive load.

Functional Connectivity Abnormalities. High-field (7T) fMRI studies reveal altered connectivity patterns. Abnormal salience network connectivity, particularly involving the right insula, appears across multiple studies—8 of 10 different ME/CFS-specific connections involve a salience network hub [94]. Specific findings include:

- Stronger connections between salience network and hippocampus
- Stronger connections between salience network and brainstem reticular activation system
- Reduced dopaminergic hippocampal-nucleus-accumbens connectivity, implying blunted motivation and cognition [95]
- Extensive aberrant ponto-cerebellar connections consistent with ME/CFS symptomatology

The 2024 NIH Study: Temporoparietal Junction. The NIH deep phenotyping study identified decreased activity in the temporoparietal junction (TPJ) during effort-based tasks [5]. The TPJ is responsible for effort-based decision-making, and its dysfunction “may cause fatigue by disrupting the way the brain decides how to exert effort.” While controls showed increased blood oxygen levels in task-relevant regions, ME/CFS patients showed decreased levels in the TPJ, superior parietal lobule, and right temporal gyrus. This finding provides a neural substrate for the effort-performance disconnect described by patients.

Neuroinflammation Studies. PET studies using TSPO ligands (markers of microglial activation) have produced conflicting results. Nakatomi et al. (2014) found increased binding in cingulate cortex, hippocampus, amygdala, thalamus, midbrain, and pons, suggesting widespread neuroinflammation associated with symptom severity [73]. However, Raijmakers et al. (2021) failed to replicate these findings in a similar-sized cohort [96]. Methodological factors and small sample sizes (n=9–14) limit conclusions. The role of neuroinflammation in ME/CFS cognitive dysfunction remains an active area of investigation.

Structural Changes. Structural MRI studies have identified:

- Reduced gray matter in occipital lobes, right angular gyrus, and left parahippocampal gyrus
- Frontal lobe volume reductions correlating with fatigue scores [94]

- Reduced white matter volume in left occipital lobe and left inferior fronto-occipital fasciculus
- Elevated T1w/T2w ratios suggesting increased myelin and/or iron in subcortical structures

White matter abnormalities of unknown etiology have been observed in some patients, though not consistently. Importantly, structural changes may not be prominent in early or pediatric cases, suggesting they develop with illness duration.

Brainstem Involvement. Multiple neuroimaging modalities (fMRI, PET, MRS) converge on brainstem abnormalities as a consistent finding in ME/CFS [94]. FDG-PET demonstrates glucose hypometabolism in the brainstem, supporting a physiological basis for fatigue, unrefreshing sleep, and cognitive symptoms. Impaired connectivity involving the brainstem has been identified in multiple studies and may reflect dysautonomia contributing to cognitive dysfunction through cerebral hypoperfusion.

2.4 Autonomic Dysfunction

Autonomic dysfunction is present in 70–90% of ME/CFS patients [97], manifesting as orthostatic intolerance, temperature dysregulation, and cardiovascular symptoms. The autonomic nervous system controls involuntary functions including heart rate, blood pressure, digestion, temperature regulation, and bladder control. Dysautonomia in ME/CFS creates a cascade of disabling symptoms often misattributed to anxiety or deconditioning.

2.4.1 Orthostatic Intolerance

Orthostatic intolerance (OI) refers to symptoms triggered or worsened by upright posture. It is one of the most common and disabling features of ME/CFS autonomic dysfunction.

Clinical Presentation

Symptoms upon standing or prolonged sitting include:

- **Lightheadedness or dizziness:** Feeling faint, vision graying out
- **Palpitations:** Awareness of rapid or pounding heartbeat
- **Tremulousness:** Shaking, feeling weak or unstable
- **Cognitive impairment:** “Coat hanger pain” (neck/shoulder aching from reduced cerebral perfusion)
- **Nausea:** Gastrointestinal symptoms triggered by position change
- **Shortness of breath:** Air hunger despite normal oxygen saturation
- **Fatigue exacerbation:** Profound worsening of exhaustion when upright

Patients often develop adaptive behaviors: sitting while showering, lying down frequently, avoiding standing in lines, preferring reclined positions.

Postural Orthostatic Tachycardia Syndrome (POTS)

POTS is the most common form of orthostatic intolerance in ME/CFS, affecting 25–50% of patients [98].

Diagnostic Criteria.

- **Heart rate increase:** ≥ 30 bpm within 10 minutes of standing (or ≥ 40 bpm in adolescents) [99]
- **Absence of orthostatic hypotension:** Blood pressure remains stable or increases
- **Symptom provocation:** OI symptoms occur with the tachycardia
- **Duration:** Symptoms present for ≥ 3 months
- **Exclusions:** No other cause (dehydration, medications, prolonged bed rest alone)

Physiological Mechanisms. POTS in ME/CFS may involve:

- **Hypovolemia:** Reduced blood volume (measured via Evans blue dye dilution studies)
- **Venous pooling:** Impaired vasoconstriction allows blood to pool in lower extremities
- **Hyperadrenergic state:** Excessive norepinephrine release upon standing
- **Baroreceptor dysfunction:** Impaired blood pressure sensing
- **Autoimmunity:** Antibodies against adrenergic and muscarinic receptors affecting vascular tone

Measurement.

- **NASA Lean Test:** 10-minute standing test measuring heart rate and blood pressure every 2 minutes
- **Tilt table testing:** Gold standard, involves passive upright tilt to 70° for up to 45 minutes
- **Home monitoring:** Patients can document HR/BP changes with home devices

Orthostatic Hypotension (OH)

Less common than POTS but present in some ME/CFS patients:

- **Definition:** Sustained drop in systolic BP ≥ 20 mmHg or diastolic BP ≥ 10 mmHg within 3 minutes of standing
- **Symptoms:** Severe lightheadedness, syncope, visual blurring, cognitive impairment
- **Mechanism:** Inadequate vasoconstriction response to postural change

- **Treatment:** Different from POTS; requires blood pressure support (fludrocortisone, midodrine)

Neurally Mediated Hypotension (NMH)

Also called vasovagal syncope or neurocardiogenic syncope:

- **Presentation:** Delayed blood pressure drop and bradycardia after prolonged standing (typically 15–45 minutes)
- **Mechanism:** Paradoxical vagal activation causing vasodilation and heart rate slowing
- **Tilt table pattern:** Initial normal response, then sudden BP/HR drop with near-syncope
- **Overlap:** Can coexist with POTS in same patient

Tilt Table Testing Protocol

The gold standard for diagnosing orthostatic intolerance:

1. **Preparation:** Patient lies supine on motorized table with footboard support
2. **Baseline:** 10–20 minutes supine to establish baseline HR and BP
3. **Tilt:** Table tilted to 70° head-up position
4. **Monitoring:** Continuous HR, BP, and symptoms recorded for up to 45 minutes
5. **Endpoints:** Test terminated if syncope occurs, BP drops dangerously, or maximum duration reached

Interpretation.

- **POTS pattern:** Sustained HR increase ≥ 30 bpm without BP drop
- **Orthostatic hypotension:** BP drop within 3 minutes
- **NMH pattern:** Delayed sudden BP/HR drop after 15–45 minutes
- **Normal response:** HR increase < 30 bpm, stable BP

Clinical note: Some ME/CFS patients have severe OI symptoms with “normal” tilt table results. This may reflect:

- Cerebral hypoperfusion despite maintained BP (impaired cerebral autoregulation)
- Small fiber neuropathy not detected by standard autonomic testing
- Endothelial dysfunction affecting microvascular perfusion

2.4.2 Other Autonomic Symptoms

Beyond orthostatic intolerance, ME/CFS patients experience widespread autonomic dysfunction:

Temperature Dysregulation

Impaired thermoregulation manifests as:

- **Subnormal body temperature:** Chronic low-grade hypothermia (96–97°F / 35.5–36°C)
- **Temperature instability:** Fluctuations throughout day without infection
- **Heat intolerance:** Severe symptom exacerbation in warm environments
- **Cold intolerance:** Inability to warm up, cold extremities even in warm rooms
- **Inappropriate sweating:** Night sweats, profuse sweating with minimal exertion
- **Lack of sweating:** Some patients lose ability to sweat (anhidrosis)

Sweating Abnormalities

Thermoregulatory and sympathetic sweating dysfunction:

- **Hyperhidrosis:** Excessive sweating of hands, feet, or generalized
- **Hypohidrosis/anhidrosis:** Reduced or absent sweating capacity
- **Gustatory sweating:** Sweating triggered by eating (cranial autonomic dysfunction)
- **Night sweats:** Drenching sweats during sleep requiring clothing/bedding changes

Gastrointestinal Symptoms

Autonomic control of GI function is commonly impaired:

- **Gastroparesis:** Delayed gastric emptying causing early satiety, nausea, bloating
- **Irritable Bowel Syndrome (IBS):** Diarrhea-predominant, constipation-predominant, or alternating
- **Dysmotility:** Impaired intestinal peristalsis
- **Nausea:** Chronic or episodic, often worse upon standing (orthostatic nausea)
- **Abdominal pain:** Cramping, visceral hypersensitivity

Urinary Dysfunction

Bladder autonomic control abnormalities include:

- **Urgency and frequency:** Needing to urinate frequently with sudden urgency
- **Nocturia:** Waking multiple times at night to urinate
- **Incomplete emptying:** Sensation of residual urine
- **Interstitial cystitis overlap:** Bladder pain, pressure, frequency

Cardiac Symptoms

Beyond POTS-related tachycardia:

- **Inappropriate sinus tachycardia:** Resting heart rate >100 bpm without postural trigger
- **Palpitations:** Awareness of heartbeat, skipped beats, forceful beats
- **Chest pain:** Non-cardiac chest pain (microvascular angina, costochondritis)
- **Heart rate variability reduction:** Reduced parasympathetic tone
- **Exercise intolerance:** Exaggerated HR response to minimal exertion

Pupillary Abnormalities

Autonomic control of pupils may be affected:

- **Light sensitivity (photophobia):** Inability to tolerate bright lights
- **Impaired pupil constriction:** Sluggish response to light
- **Anisocoria:** Unequal pupil sizes

2.4.3 Autonomic Testing Battery

Comprehensive autonomic function assessment may include:

- **Tilt table test:** Orthostatic intolerance assessment
- **Valsalva maneuver:** Tests baroreceptor and cardiovagal function
- **Deep breathing test:** Measures heart rate variability during paced breathing
- **Quantitative sudomotor axon reflex test (QSART):** Assesses sweating capacity
- **Thermoregulatory sweat test:** Maps sweating across entire body
- **Pupillometry:** Automated pupil response measurement
- **Skin biopsy:** Small fiber neuropathy assessment (intraepidermal nerve fiber density)

Many ME/CFS specialty centers lack access to full autonomic testing, making tilt table and basic orthostatic vitals the most commonly used assessments.

2.4.4 Clinical Implications

Autonomic dysfunction in ME/CFS is:

- **Objectively measurable:** Tilt table, HRV, and other tests provide objective abnormalities
- **Highly disabling:** OI can prevent standing long enough to shower or prepare meals
- **Treatable:** Salt, fluids, compression, and medications can significantly improve symptoms
- **Not anxiety:** Patients are often told POTS is anxiety; it is a physiological abnormality

- **Connected to energy metabolism:** Autonomic dysfunction may reflect mitochondrial impairment in autonomic neurons

Recognition and treatment of dysautonomia is often the first step in improving ME/CFS functional capacity.

2.5 Pain

Pain is a prominent symptom in ME/CFS, with approximately 80% of patients reporting significant pain in the past week [100]. Pain is included as a diagnostic criterion in multiple case definitions and contributes substantially to disability and reduced quality of life.

2.5.1 Types of Pain in ME/CFS

Myalgia (Muscle Pain). Muscle pain is the most common pain complaint, affecting 72–94% of ME/CFS patients [101]. The pain is typically widespread rather than localized and characteristically worsens 8–72 hours following physical exertion as part of post-exertional malaise. Patients describe deep, aching pain that differs from delayed-onset muscle soreness in healthy individuals—it occurs following minimal exertion, lasts substantially longer, and is accompanied by other PEM symptoms. The pain reflects underlying skeletal muscle dysfunction including mitochondrial impairment, oxidative stress, reduced heat shock proteins, and impaired muscle contractility [102].

Arthralgia (Joint Pain). Joint pain affects 58–84% of patients and is included as a criterion in both Fukuda and Canadian Consensus definitions [27, 28]. The pattern is characteristically migratory (moving between joints) and occurs without the swelling, redness, warmth, or deformity seen in inflammatory arthritis. This distinction is clinically important: presence of joint inflammation suggests an alternative diagnosis or comorbid condition requiring separate evaluation.

Headaches. Headaches are significantly more common in ME/CFS than the general population: 84% experience migraine headaches (versus 5% in healthy controls) and 81% have tension-type headaches (versus 45% in controls) [103]. The breakdown includes migraine without aura (60%), migraine with aura (24%), tension headaches only (12%), and no headaches (4%). ME/CFS patients with migraine demonstrate lower pressure pain thresholds (2.36 kg versus 5.23 kg in controls, $p < 0.001$) and higher fibromyalgia comorbidity (47% versus 0%) [103]. Headaches are listed in Fukuda criteria as one of eight minor symptoms.

Neuropathic Pain. A subset of ME/CFS patients experience neuropathic pain characterized by burning, tingling, or electric shock sensations. This correlates with the finding that 30–38% of ME/CFS patients have small fiber neuropathy (SFN) confirmed by skin biopsy demonstrating reduced intraepidermal nerve fiber density [104]. Of those with confirmed SFN, 93% have comorbid postural orthostatic tachycardia syndrome (POTS) or other orthostatic intolerance, suggesting shared pathophysiology involving autonomic small fibers [105].

2.5.2 Pain Mechanisms

Central Sensitization. Central sensitization—increased excitability of central nervous system pain pathways—is present in 84% of ME/CFS patients, compared to 95% of fibromyalgia patients and 0% of healthy controls [106]. This is defined by enhanced temporal summation (wind-up) combined with inefficient conditioned pain modulation. Clinical manifestations include:

- Generalized hyperalgesia to electrical, mechanical, heat, and chemical stimuli
- Affects multiple tissues including skin, muscle, and viscera
- Hyperalgesia augmented rather than decreased following exercise or other stressors
- Lower pressure pain thresholds: ME/CFS median 222 kPa versus healthy controls 311 kPa ($p < 0.05$) [106]

Central sensitization is driven by neuroinflammation—glial cell activation (microglia and astrocytes) in the spinal cord and brain releasing pro-inflammatory cytokines and chemokines that sustain neural hypersensitivity [107].

Small Fiber Neuropathy. Small fiber neuropathy provides an objective, biopsy-confirmed mechanism for pain in a substantial subset of patients. Studies find 30–38% of ME/CFS patients meet diagnostic criteria for SFN [104]. Small fibers (A-delta and C fibers) mediate pain, temperature sensation, and autonomic function, explaining the overlap between pain and dysautonomia. The etiology of SFN in ME/CFS is not fully established but may involve autoimmune mechanisms, as autoantibodies against small fiber antigens have been identified in some patients.

Peripheral Mechanisms. Peripheral contributors to ME/CFS pain include:

- **Elevated blood lactate:** Nearly half of ME/CFS patients have elevated resting lactate levels, correlating with more severe post-exertional malaise [65]. Lactate accumulation reflects anaerobic metabolism predominance due to mitochondrial dysfunction.
- **Metabolic dysfunction:** Impaired ATP synthesis leads to toxic metabolite accumulation that activates muscle nociceptors [102].
- **Impaired proton handling:** Profound intramuscular acidosis develops following minimal exertion.
- **Reduced oxygen delivery:** Endothelial dysfunction and microvascular abnormalities may limit oxygen supply to exercising muscles.

Relationship to Post-Exertional Malaise. Pain is a core component of PEM. A meta-analysis found small to moderate pain increases following exercise in ME/CFS versus controls (Hedges' $d = 0.42$, 95% CI: 0.16–0.67), with delayed pain showing larger effects at 8–72 hours ($d = 0.71$) than at 0–2 hours ($d = 0.32$) [108]. This delayed, disproportionate pain response parallels the temporal pattern of other PEM symptoms and likely reflects the same underlying metabolic and immune dysfunction. Factor analysis of PEM symptoms identifies a distinct “musculoskeletal factor” comprising muscle pain, weakness, and post-exertional fatigue [108].

2.5.3 Pain Assessment and Management Considerations

Quantitative Sensory Testing. Quantitative sensory testing (QST) can objectively document pain hypersensitivity. Commonly used measures include pressure pain thresholds at standard sites (trapezius, forearm, 18 fibromyalgia tender points), thermal thresholds, and temporal summation protocols. QST findings may support disability claims and guide treatment by identifying central versus peripheral contributions.

Overlap with Fibromyalgia. ME/CFS and fibromyalgia show substantial clinical overlap: 47.3% (95% CI: 45.97–48.63) of ME/CFS diagnoses overlap with fibromyalgia, with 35–75% of ME/CFS patients meeting fibromyalgia criteria and 20–70% of fibromyalgia patients meeting ME/CFS criteria [109]. Cerebrospinal fluid proteomics are indistinguishable between ME/CFS patients with and without comorbid fibromyalgia, suggesting shared pathophysiology [110]. Key clinical distinctions:

- Fibromyalgia: Pain predominant, fatigue secondary
- ME/CFS: Fatigue and PEM predominant, pain prominent but not defining
- Comorbid patients have worse outcomes: greater physical disability, more severe pain, and more pronounced post-exertional symptoms than either condition alone

Treatment Implications. Pain management in ME/CFS must account for the underlying mechanisms:

- Standard analgesics may be insufficient given central sensitization
- Interventions targeting neuroinflammation (e.g., low-dose naltrexone) may address central mechanisms
- Activity pacing prevents pain exacerbation from PEM
- Treatment of underlying small fiber neuropathy (if present) with IVIG has shown benefit in some patients
- Medications effective for fibromyalgia pain (duloxetine, pregabalin) may help the subset with overlapping presentations

2.6 Sensory Sensitivities

Heightened sensitivity to sensory stimuli is a common but often underrecognized feature of ME/CFS, present in 70–90% of patients [111]. These sensitivities can be profoundly disabling and contribute significantly to activity limitation and social isolation.

2.6.1 Types of Sensory Sensitivity

Photophobia (Light Sensitivity). Light sensitivity affects approximately 70% of ME/CFS patients [111]. Manifestations include:

- Inability to tolerate bright lights, including sunlight and fluorescent lighting
- Need for sunglasses indoors or dimmed environments
- Headaches or symptom exacerbation triggered by light exposure
- Difficulty with screens (computers, phones, televisions)
- Preference for dark or low-light environments

Light sensitivity may reflect autonomic dysfunction affecting pupillary control, central sensitization affecting visual processing, or neuroinflammation in visual pathways.

Phonophobia (Sound Sensitivity). Sound sensitivity affects 60–80% of patients and can be severely disabling [111]:

- Normal conversation volumes feel uncomfortably loud
- Sudden or unexpected sounds cause startle responses and symptom flares
- Multiple simultaneous sounds (e.g., conversations in a restaurant) are intolerable
- Background noise prevents concentration
- Need for quiet environments or noise-canceling headphones

In severe cases, patients cannot tolerate any sound and require complete silence, significantly limiting social contact and access to medical care.

Chemical Sensitivity (Multiple Chemical Sensitivity). Sensitivity to chemicals and odors affects 40–60% of ME/CFS patients [111]:

- Fragrances (perfumes, cleaning products, air fresheners) trigger symptoms
- Exhaust fumes and other environmental pollutants cause reactions
- New materials (carpets, furniture, paint) provoke symptoms
- Symptoms may include headache, cognitive dysfunction, nausea, respiratory symptoms
- Overlap with Multiple Chemical Sensitivity (MCS) syndrome

Touch and Pressure Sensitivity. Tactile hypersensitivity manifests as:

- Allodynia—painful response to normally non-painful touch
- Clothing tags, seams, or tight clothing feel unbearable
- Difficulty tolerating physical examination
- Hyperalgesia—exaggerated pain response to mildly painful stimuli

This overlaps with the central sensitization mechanisms described in the Pain section.

Temperature Sensitivity. Intolerance to temperature extremes affects most patients:

- Heat intolerance with symptom exacerbation in warm environments
- Cold intolerance with difficulty warming up
- Narrow range of comfortable temperatures
- Symptoms triggered by temperature changes

This reflects autonomic dysfunction affecting thermoregulation (see Section 2.4).

2.6.2 Mechanisms of Sensory Sensitivity

Central Sensitization. The same central sensitization mechanisms that produce pain hypersensitivity likely underlie broader sensory sensitivities. Reduced inhibitory control in the central nervous system leads to amplification of all sensory inputs, not just nociceptive signals [107].

Neuroinflammation. Glial activation and neuroinflammatory processes may directly affect sensory processing pathways, reducing thresholds for activation and impairing habituation to repeated stimuli.

Autonomic Dysfunction. Dysautonomia contributes to sensory sensitivity through impaired pupillary control (photophobia), altered blood flow to sensory organs, and dysfunctional sympathetic responses to stimuli.

Energy Depletion. Sensory processing requires energy. With baseline energy insufficiency, normal sensory processing may exceed available cellular resources, leading to symptoms from stimulation that healthy individuals filter automatically.

2.6.3 Clinical Implications

Activity Limitation. Sensory sensitivities profoundly limit function:

- Medical appointments become challenging (bright lights, waiting room noise, chemical smells)
- Shopping, restaurants, and public spaces are often intolerable
- Work environments may be impossible to tolerate
- Social gatherings exceed sensory capacity

Assessment Considerations. When evaluating ME/CFS patients, clinicians should:

- Ask specifically about sensory sensitivities
- Modify examination environments (dim lights, reduce noise)
- Allow patients to wear sunglasses or earplugs
- Avoid fragranced products
- Recognize that sensory overload can trigger PEM

Management. Management focuses on environmental modification:

- Sunglasses, tinted lenses, or FL-41 lenses for photophobia
- Noise-canceling headphones or earplugs for phonophobia
- Fragrance-free environments and products
- Loose, soft clothing without tags or seams
- Temperature-controlled environments with ability to layer clothing
- Gradual, controlled exposure when improvement occurs

3 Additional Symptoms and Manifestations

Beyond the core symptoms of post-exertional malaise, unrefreshing sleep, cognitive impairment, autonomic dysfunction, and pain described in Chapter 2, ME/CFS patients experience a wide range of additional symptoms affecting virtually every body system. This chapter provides a comprehensive catalog of these symptoms, organized by physiological system, ranging from mild and common manifestations to severe and disabling complications.

⚠ Limitation 1: Symptom Mechanisms: Plausible Explanations, Not Confirmed Pathways

The mechanistic explanations offered throughout this chapter (e.g., “central sensitisation” for sensory symptoms, “mitochondrial dysfunction” for muscle weakness, “mast cell activation” for allergic-type symptoms) represent the most plausible current hypotheses but have not been confirmed as the actual cause of these symptoms in ME/CFS patients. Most proposed mechanisms are inferred from other conditions; direct investigation of symptom-producing mechanisms in ME/CFS cohorts remains limited.

3.1 Neurological Symptoms

Neurological manifestations in ME/CFS extend beyond cognitive dysfunction to include sensory, motor, and perceptual abnormalities.

3.1.1 Sensory Sensitivities

Many ME/CFS patients develop heightened sensitivity to sensory stimuli that were previously tolerable.

Photophobia (Light Sensitivity)

Mild to Moderate.

- Discomfort in bright indoor lighting or sunlight
- Need for sunglasses indoors or in dim environments
- Difficulty tolerating computer screens or fluorescent lights
- Preference for dim environments
- Eye strain and headaches triggered by bright light

Severe.

- Inability to tolerate any artificial lighting
- Need to wear sunglasses or eye masks constantly
- Confinement to darkened rooms
- Severe pain triggered by brief light exposure
- Light-triggered migraines or seizure-like episodes

Mechanism. Photophobia likely reflects both central sensitization (amplification of sensory signals in the brain) and mitochondrial dysfunction in retinal cells, which have exceptionally high energy demands. Visual processing itself is energetically expensive, consuming significant ATP.

Hyperacusis (Sound Sensitivity)

Mild to Moderate.

- Discomfort in noisy environments (restaurants, crowds)
- Difficulty tolerating sudden or loud sounds
- Need for ear protection in normal-volume environments
- Exacerbation of cognitive symptoms by auditory stimulation
- Preference for quiet, low-stimulation environments

Severe.

- Pain from normal conversation volume
- Inability to tolerate any environmental sounds (traffic, appliances, voices)
- Need for soundproofing or constant ear protection
- Sound-triggered crashes or seizure-like episodes
- Complete withdrawal from environments with any noise

Mechanism. Hyperacusis involves central auditory processing abnormalities, potentially related to reduced descending inhibition from the cortex, allowing normal auditory signals to be perceived as excessively loud or painful. The cochlea's high metabolic demands may also contribute.

Touch Sensitivity and Allodynia

Clinical Presentation.

- Light touch perceived as painful (allodynia)
- Clothing textures causing discomfort or pain

- Inability to tolerate certain fabrics (tags, seams, tight clothing)
- Hypersensitivity to temperature of touch
- Discomfort from physical contact (hugs, handshakes)
- Skin feeling “raw” or “burned”

Mechanism. Touch sensitivity reflects small fiber neuropathy and central sensitization. Peripheral nerve dysfunction causes abnormal tactile processing, while central amplification interprets benign touch as noxious stimuli.

Chemical and Odor Sensitivities (Multiple Chemical Sensitivity)

Common Triggers.

- Perfumes, colognes, and fragranced products
- Cleaning chemicals and detergents
- Cigarette smoke and air pollution
- Gasoline and petroleum fumes
- Paint, solvents, and VOCs (volatile organic compounds)
- Pesticides and herbicides
- New carpets, furniture, or building materials (off-gassing)

Symptom Response.

- Headaches or migraines
- Nausea and dizziness
- Respiratory symptoms (shortness of breath, throat irritation)
- Brain fog and cognitive impairment
- Fatigue exacerbation
- Allergic-type reactions (rashes, congestion)
- PEM-like crashes following exposure

Mechanism. Chemical sensitivities may involve mast cell activation (inappropriate degranulation releasing histamine and inflammatory mediators), liver detoxification impairment, and olfactory-limbic dysregulation. The energetic cost of detoxifying chemicals may exceed available metabolic capacity.

Taste and Smell Alterations

Clinical Presentation.

- Reduced sense of smell (hyposmia) or complete loss (anosmia)

- Distorted smell perception (parosmia)
- Altered taste perception (dysgeusia)
- Metallic taste in mouth
- Food aversions due to altered taste
- Difficulty detecting spoiled food due to reduced olfaction

Mechanism. Olfactory and gustatory dysfunction may reflect neuroinflammation affecting cranial nerves, central processing abnormalities, or zinc deficiency (common in ME/CFS and essential for taste/smell function).

3.1.2 Motor and Coordination Symptoms

Tremor

Clinical Presentation.

- Fine hand tremor, often action-induced
- Tremor worsening with exertion or fatigue
- Difficulty with fine motor tasks (writing, buttoning, using utensils)
- Postural tremor when holding positions
- Voice tremor in some cases

Mechanism. Tremor reflects energy insufficiency in motor control circuits (basal ganglia, cerebellum) and motor neurons. Fine motor control requires continuous rapid adjustments that consume ATP; when energy is marginal, precision degrades, producing tremor.

Muscle Weakness and Reduced Strength

Clinical Presentation.

- Generalized muscle weakness disproportionate to disuse
- Difficulty lifting objects, climbing stairs, or standing from seated position
- Grip strength reduction
- Proximal muscle weakness (shoulders, hips)
- Weakness worsening with exertion and persisting after rest

Mechanism. Muscle weakness reflects impaired ATP production, not simply deconditioning. Studies show reduced force generation at the cellular level due to mitochondrial dysfunction, distinct from atrophy-related weakness.

Gait Disturbances

Clinical Presentation.

- Unsteady gait, feeling “off-balance”
- Shuffling or slow walking pace
- Increased fall risk
- Need for mobility aids (canes, walkers, wheelchairs)
- Difficulty with stairs or uneven surfaces
- Gait worsening with fatigue

Mechanism. Gait disturbances reflect cerebellar dysfunction, proprioceptive impairment, muscle weakness, and orthostatic intolerance. Walking requires integration of multiple systems, all of which may be impaired in ME/CFS.

Muscle Fasciculations and Twitching

Clinical Presentation.

- Spontaneous muscle twitches visible under skin
- Fasciculations in legs, arms, face, or trunk
- Twitching often worsening at rest or before sleep
- Generally benign but distressing

Mechanism. Fasciculations may reflect peripheral nerve hyperexcitability due to electrolyte imbalances, magnesium deficiency, or metabolic stress in motor neurons.

3.1.3 Paresthesias and Sensory Disturbances

Clinical Presentation.

- Tingling, numbness, or “pins and needles” sensations
- Burning sensations in hands, feet, or other areas
- Electric shock-like sensations
- Crawling sensations on skin (formication)
- Sensations often not following anatomical nerve distributions

Mechanism. Paresthesias reflect small fiber neuropathy, documented in many ME/CFS patients via skin biopsy. Small nerve fibers are metabolically demanding and vulnerable to energy deficit and oxidative stress.

3.1.4 Dizziness and Vertigo

Clinical Presentation.

- Non-spinning dizziness (lightheadedness)
- True vertigo (sensation of room spinning)
- Disequilibrium (feeling unsteady)
- Presyncope (feeling about to faint)
- Symptoms worsening with position changes, exertion, or sensory stimulation

Mechanism. Dizziness in ME/CFS has multiple contributors: orthostatic intolerance (inadequate cerebral perfusion when upright), vestibular dysfunction, cerebral hypoperfusion, and central processing abnormalities.

3.1.5 Tinnitus

Clinical Presentation.

- Ringing, buzzing, hissing, or roaring sounds
- Unilateral or bilateral
- Constant or intermittent
- Volume may fluctuate with fatigue, stress, or exertion
- Can be severely disabling and interfere with sleep

Mechanism. Tinnitus may reflect cochlear damage (high metabolic demands make cochlear hair cells vulnerable), auditory nerve dysfunction, or central auditory processing abnormalities.

3.1.6 Seizure-Like Episodes

Clinical Presentation.

- Episodes resembling seizures but with normal EEG (non-epileptic)
- Triggered by sensory overload, exertion, or stress
- May include loss of motor control, altered consciousness, or convulsive movements
- Distinct from true epilepsy

Mechanism. Non-epileptic seizure-like episodes may reflect severe autonomic dysfunction, cerebral hypoperfusion, or metabolic crisis in brain tissue.

3.2 Immunological and Inflammatory Symptoms

3.2.1 Flu-Like Symptoms

Many ME/CFS patients experience chronic or recurrent flu-like symptoms even in the absence of active infection.

Sore Throat.

- Persistent or recurrent sore throat without infection
- Tender, swollen throat sensation
- May worsen with exertion or during PEM

Tender Lymph Nodes.

- Painful, swollen lymph nodes (cervical, axillary, inguinal)
- Lymphadenopathy without evidence of infection
- Lymph node tenderness worsening during crashes

Low-Grade Fever and Chills.

- Recurrent low-grade fever (37.5–38°C)
- Subjective fever sensation even when temperature normal
- Chills and cold intolerance
- Night sweats
- Temperature dysregulation (alternating hot/cold)

Mechanism. Flu-like symptoms reflect chronic immune activation and cytokine production, even without active infection. Elevated inflammatory markers suggest ongoing immune system dysregulation.

3.2.2 Infection Susceptibility and Viral Reactivation

Recurrent Infections.

- Frequent upper respiratory infections
- Recurrent urinary tract infections
- Skin infections
- Sinus infections
- Longer recovery from infections compared to pre-illness

Viral Reactivation.

- Reactivation of latent herpesviruses (EBV, HHV-6, CMV, VZV)
- Elevated viral antibody titers
- Recurrent cold sores or shingles
- Chronic viral symptoms

Mechanism. ME/CFS patients show evidence of immune exhaustion: T-cells are functionally impaired and unable to maintain control over latent infections. This creates susceptibility to new infections and reactivation of dormant viruses.

3.2.3 Allergies and Mast Cell Activation

New or Worsening Allergies.

- Development of new food allergies or intolerances
- Worsening seasonal allergies
- Reactions to previously tolerated substances
- Oral allergy syndrome (cross-reactivity with pollen)

Mast Cell Activation Syndrome (MCAS) Features.

- Flushing and skin rashes (urticaria, hives)
- Angioedema (swelling of face, lips, tongue)
- Anaphylaxis or anaphylactoid reactions
- Gastrointestinal symptoms (nausea, diarrhea, abdominal pain)
- Respiratory symptoms (wheezing, throat tightness)
- Cardiovascular symptoms (tachycardia, hypotension)
- Neurological symptoms (brain fog, headache)
- Triggered by heat, cold, stress, exertion, foods, medications, or chemicals

Mechanism. An estimated 30–50% of ME/CFS patients show features of mast cell activation syndrome. Mast cells become hyperreactive, degranulating inappropriately and releasing histamine and other inflammatory mediators.

3.3 Musculoskeletal Symptoms

3.3.1 Muscle Cramps and Contractures

Clinical Presentation.

- Spontaneous muscle cramps without preceding exertion
- Nocturnal leg cramps
- Cramps in unexpected muscle groups (hands, feet, neck, throat, jaw)
- Prolonged muscle contractures
- Reverse finger contractures (fingers held extended rather than curled)
- Difficulty releasing grip or relaxing contracted muscles
- Constant sensation of being “ready to cramp”

Mechanism. Muscle relaxation requires ATP to pump calcium ions back into storage. When ATP is insufficient, muscles cannot fully relax, leading to spontaneous cramping. This reflects the same energy deficit causing fatigue, but manifested as impaired muscle relaxation.

3.3.2 Myalgia (Muscle Pain)

Clinical Presentation.

- Widespread muscle aching and soreness
- Deep muscle pain, often described as “flu-like”
- Pain worsening with activity or pressure
- Muscle tenderness to palpation
- Delayed-onset muscle soreness after minimal exertion
- Persistent muscle tension

Mechanism. Myalgia reflects lactic acid accumulation from anaerobic metabolism, muscle hypoxia, central sensitization amplifying pain signals, and possible muscle microtrauma from energy-deficient muscle fibers.

3.3.3 Arthralgia (Joint Pain)

Clinical Presentation.

- Diffuse joint pain without objective swelling or inflammation
- Pain in knees, shoulders, wrists, hands, ankles
- Migratory joint pain (moving from joint to joint)
- Morning stiffness
- Pain worsening with activity and weather changes
- Inflammatory-pattern joint pain in some patients (knuckles, suggesting autoimmune overlap)

Mechanism. Joint pain without visible pathology likely reflects central sensitization, periarticular tissue energy deficit, microcirculatory dysfunction, and in some cases, low-grade inflammatory or autoimmune processes.

3.3.4 Fibromyalgia Overlap

Clinical Overlap. Significant symptom overlap exists between ME/CFS and fibromyalgia:

- Widespread pain
- Tender points
- Sleep disturbance
- Cognitive dysfunction
- Fatigue

Distinction. The primary distinction is the presence and prominence of PEM in ME/CFS, which is not a defining feature of fibromyalgia. Many researchers consider them overlapping conditions on a spectrum of neuroimmune disorders.

3.4 Gastrointestinal Symptoms

Gastrointestinal symptoms are extremely common in ME/CFS, with estimates suggesting 70–90% of patients experience significant GI dysfunction.

3.4.1 Nausea

Clinical Presentation.

- Chronic or recurrent nausea
- Nausea triggered by exertion, movement, or sensory stimulation
- Medication-induced nausea (many ME/CFS patients have heightened sensitivity)
- Early satiety (feeling full quickly)
- Food aversions

3.4.2 Irritable Bowel Syndrome (IBS)

Clinical Presentation.

- Abdominal pain or cramping
- Diarrhea (IBS-D), constipation (IBS-C), or alternating patterns (IBS-M)
- Bloating and gas
- Urgency or incomplete evacuation

- Symptoms worsening with stress or certain foods

Mechanism. IBS in ME/CFS likely involves gut dysbiosis, mast cell activation in the GI tract, autonomic dysfunction affecting gut motility, and visceral hypersensitivity (central amplification of gut sensations).

3.4.3 Food Intolerances and Sensitivities

Common Triggers.

- Gluten (celiac disease or non-celiac gluten sensitivity)
- Dairy/lactose
- FODMAPs (fermentable carbohydrates)
- Histamine-rich foods (aged cheese, fermented foods, cured meats)
- Specific proteins (nuts, eggs, soy)
- Artificial additives and preservatives

Symptom Response.

- Gastrointestinal symptoms (bloating, pain, diarrhea)
- Systemic symptoms (fatigue, brain fog, headache)
- Allergic-type reactions
- PEM-like exacerbations

3.4.4 Gastroparesis and Delayed Gastric Emptying

Clinical Presentation.

- Feeling full after small amounts of food
- Persistent nausea
- Vomiting (especially of undigested food)
- Abdominal bloating and discomfort
- Unpredictable blood sugar fluctuations

Mechanism. Gastroparesis reflects autonomic dysfunction affecting the vagus nerve, which controls gastric motility. Impaired stomach emptying creates digestive symptoms and nutritional challenges.

3.4.5 Gastroesophageal Reflux (GERD)

Clinical Presentation.

- Heartburn and acid reflux
- Regurgitation
- Difficulty swallowing (dysphagia)
- Chronic cough or throat clearing
- Worsening when lying down

3.5 Cardiovascular Symptoms

3.5.1 Palpitations

Clinical Presentation.

- Awareness of heartbeat (racing, pounding, or irregular)
- Tachycardia (elevated heart rate) at rest or with minimal activity
- Premature ventricular contractions (PVCs) or atrial ectopy
- Palpitations triggered by position changes, exertion, or stress
- Often benign but distressing

Mechanism. Palpitations reflect autonomic dysfunction, orthostatic intolerance, and potential cardiac preload failure. The heart races in an attempt to compensate for inadequate venous return and reduced stroke volume.

3.5.2 Chest Pain

Clinical Presentation.

- Non-cardiac chest pain (normal cardiac workup)
- Sharp, stabbing, or aching chest pain
- Costochondritis (chest wall inflammation)
- Chest tightness or pressure
- Pain worsening with breathing or movement

Differential Diagnosis. While chest pain in ME/CFS is typically non-cardiac, it is essential to rule out true cardiac pathology, especially in older patients or those with cardiovascular risk factors.

3.5.3 Blood Pressure Abnormalities

Clinical Presentation.

- Orthostatic hypotension (blood pressure drops upon standing)
- Labile blood pressure (large fluctuations)
- Hypertension in some patients
- Symptoms of inadequate perfusion (dizziness, vision changes, syncope)

3.5.4 Raynaud's Phenomenon

Clinical Presentation.

- Fingers or toes turning white, blue, then red in response to cold or stress
- Numbness, tingling, or pain during episodes
- Vascular spasm in extremities

Mechanism. Raynaud's reflects exaggerated vasoconstriction, likely related to autonomic dysfunction and dysregulated catecholamine responses.

3.6 Respiratory Symptoms

3.6.1 Dyspnea and Air Hunger

Clinical Presentation.

- Shortness of breath at rest or with minimal exertion
- Sensation of not getting a "satisfying" breath
- Need to consciously focus on breathing
- Air hunger not relieved by deep breathing
- Normal oxygen saturation (SpO₂) during symptoms

Mechanism. Dyspnea in ME/CFS typically reflects problems with oxygen *delivery* and *utilization* rather than oxygen intake. Contributing factors include:

- Autonomic dysfunction (vagus nerve signaling errors)
- Microcirculatory failure (oxygen cannot reach tissues)
- Preload failure (blood pooling prevents adequate cardiac output)
- Respiratory muscle weakness
- Dysfunctional breathing patterns (loss of diaphragm-chest synchrony)

3.6.2 Dysfunctional Breathing Patterns

Clinical Presentation.

- Loss of synchrony between chest and abdominal breathing
- Overuse of accessory muscles (neck, shoulders)
- Shallow, rapid breathing
- Breath-holding or irregular breathing rhythm
- Exertional breathlessness disproportionate to activity

Mechanism. A 2025 study found 71% of ME/CFS patients have “hidden” breathing abnormalities. Using accessory muscles instead of the diaphragm consumes 3× more energy, worsening fatigue.

3.6.3 Chronic Cough

Clinical Presentation.

- Persistent dry cough without infection
- Throat irritation or tickle sensation
- Cough worsening with exertion, talking, or breathing cold air
- May be related to GERD, postnasal drip, or airway hypersensitivity

3.7 Genitourinary Symptoms

3.7.1 Urinary Dysfunction

Clinical Presentation.

- Urinary frequency (needing to urinate often)
- Urinary urgency (sudden, compelling need to urinate)
- Nocturia (waking at night to urinate)
- Bladder pain or discomfort (interstitial cystitis overlap)
- Incomplete bladder emptying sensation
- Recurrent urinary tract infections

Mechanism. Urinary symptoms reflect autonomic dysfunction affecting bladder innervation, pelvic floor dysfunction, and possible mast cell activation in bladder tissue.

3.7.2 Sexual Dysfunction

Clinical Presentation.

- Reduced libido (loss of sexual interest)
- Erectile dysfunction in men
- Reduced arousal or lubrication in women
- Pain with intercourse (dyspareunia)
- Sexual activity triggering PEM crashes
- Loss of intimate relationships due to energy constraints

Mechanism. Sexual dysfunction reflects hormonal dysregulation (low testosterone, disrupted estrogen/progesterone), autonomic dysfunction, energy insufficiency (sexual activity is highly energetically demanding), and central dopamine/reward pathway impairment.

3.7.3 Menstrual Irregularities

Clinical Presentation.

- Irregular cycles (oligomenorrhea) or absent periods (amenorrhea)
- Heavy or prolonged bleeding (menorrhagia)
- Severe premenstrual syndrome (PMS) or premenstrual dysphoric disorder (PMDD)
- Worsening of ME/CFS symptoms premenstrually or during menstruation
- Painful periods (dysmenorrhea)

Mechanism. Menstrual irregularities reflect HPA axis dysfunction, hormonal dysregulation, and the energetic demands of the menstrual cycle exceeding available capacity. Many patients report cyclical worsening of symptoms tied to hormonal fluctuations.

3.8 Endocrine and Metabolic Symptoms

3.8.1 Temperature Dysregulation

Clinical Presentation.

- Inability to maintain stable body temperature
- Feeling excessively cold (cold intolerance)
- Feeling excessively hot (heat intolerance)
- Alternating between hot and cold
- Night sweats
- Chills without fever

- Inability to tolerate temperature extremes
- Worsening of symptoms in hot or cold environments

Mechanism. Temperature dysregulation reflects hypothalamic dysfunction and autonomic impairment. The hypothalamus regulates body temperature via autonomic pathways; when these are disrupted, thermoregulation fails.

3.8.2 Excessive Thirst and Fluid Retention

Clinical Presentation.

- Polydipsia (excessive thirst)
- Dry mouth despite adequate fluid intake
- Edema (fluid retention in legs, hands, face)
- Weight fluctuations due to fluid retention

Mechanism. Excessive thirst may reflect dysregulated antidiuretic hormone (ADH/vasopressin), inadequate blood volume (hypovolemia), or mast cell mediators affecting fluid balance. Fluid retention may reflect aldosterone dysregulation or venous pooling.

3.8.3 Weight Changes

Clinical Presentation.

- Unintentional weight loss (due to reduced appetite, GI dysfunction, or hypermetabolism)
- Unintentional weight gain (due to immobility, metabolic slowing, or medication effects)
- Difficulty maintaining stable weight

3.8.4 Glucose Metabolism Abnormalities

Clinical Presentation.

- Hypoglycemia-like symptoms (shakiness, tremor, brain fog, fatigue) even with normal blood glucose
- Reactive hypoglycemia after meals
- Carbohydrate cravings
- Blood sugar instability

Mechanism. While blood glucose may be normal, ME/CFS patients experience subjective hypoglycemia because cells cannot efficiently convert glucose into ATP. The experience is similar to true hypoglycemia (cellular energy crisis) but the mechanism differs (fuel conversion failure rather than fuel lack).

3.9 Dermatological Symptoms

3.9.1 Rashes and Skin Manifestations

Clinical Presentation.

- Urticaria (hives)
- Flushing and redness
- Eczema or atopic dermatitis
- Unexplained rashes
- Livedo reticularis (mottled skin discoloration)
- Pallor or grayish skin tone

3.9.2 Hair and Nail Changes

Clinical Presentation.

- Hair loss (telogen effluvium)
- Brittle, ridged, or slow-growing nails
- Nail discoloration
- Hair texture changes

Mechanism. Hair and nails are metabolically active tissues with high nutrient demands. Chronic illness, nutritional deficiencies, and stress can disrupt hair growth cycles and nail formation.

3.10 Ocular Symptoms

3.10.1 Vision Changes

Clinical Presentation.

- Blurred vision or difficulty focusing
- Double vision (diplopia)
- Visual distortions or “floaters”
- Dry eyes

- Eye pain or pressure
- Difficulty with accommodation (switching focus between near and far)
- Progressive presbyopia (age-related vision decline occurring early)
- Energy-dependent vision quality (better on high-energy days, worse on low-energy days)

Mechanism. Vision problems reflect ciliary muscle fatigue (accommodation requires sustained ATP for muscle contraction), autonomic dysfunction affecting pupil control, and energy-dependent visual processing in the brain.

3.11 Auditory Symptoms

3.11.1 Hearing Loss

Clinical Presentation.

- Progressive sensorineural hearing loss, especially high frequencies
- Difficulty hearing in noisy environments
- Reduced speech discrimination
- Bilateral hearing impairment

Mechanism. Cochlear hair cells have exceptionally high metabolic demands (mitochondrial density second only to brain tissue). Mitochondrial dysfunction and oxidative stress damage these cells, causing progressive hearing loss.

3.12 Psychological and Cognitive-Emotional Symptoms

3.12.1 Anxiety

Clinical Presentation.

- Generalized anxiety
- Panic attacks
- Health anxiety (realistic concern about worsening condition)
- Anticipatory anxiety about exertion, crashes, or medical appointments
- Hypervigilance about energy levels and symptom changes

Distinction from Primary Anxiety Disorder. Anxiety in ME/CFS is typically *secondary*—a realistic response to living with a disabling, unpredictable illness. The anxiety often improves if symptoms improve, unlike primary anxiety disorders.

3.12.2 Depression

Clinical Presentation.

- Low mood and sadness
- Anhedonia (inability to experience pleasure)
- Hopelessness about future
- Suicidal ideation (in severe cases)
- Grief over lost capabilities and identity

Reactive vs. Primary Depression. The majority of ME/CFS patients who experience depression develop it *after* disease onset (78.1%), and 96% attribute it to disease severity rather than pre-existing psychiatric conditions. Depression in ME/CFS is typically reactive: a normal emotional response to severe, chronic illness and loss of function.

Distinguishing Features.

- Depression correlates with disease severity and functional impairment
- Desire to be active is present, but physical capacity is absent
- Effort expenditure is maximal despite minimal output (opposite of primary depression)
- Depression often improves if physical symptoms improve

3.12.3 Emotional Lability and Mood Dysregulation

Clinical Presentation.

- Easy crying or emotional overwhelm
- Irritability and low frustration tolerance
- Rapid mood shifts
- Difficulty regulating emotional responses
- Emotional symptoms worsening with fatigue

Mechanism. Emotional regulation requires prefrontal cortex function and adequate neurotransmitter availability. Energy deficit impairs executive control over emotions, leading to lability.

3.12.4 Social Withdrawal and Isolation

Clinical Presentation.

- Reduced social contact and withdrawal from relationships
- Inability to maintain friendships or family connections
- Social interaction experienced as painful and exhausting
- Loss of social identity and roles
- Profound loneliness despite lack of capacity for socializing

Mechanism. Social withdrawal is not a choice but a necessity. Social interaction is metabolically expensive (cognitive processing, emotional regulation, sensory input, sustained attention, affect generation). When energy is insufficient, patients must choose between socializing and survival activities.

Clinical Significance. The experience of social interaction as *painful*—not merely tiring but actively aversive—distinguishes ME/CFS from primary social anxiety or depression. This reflects genuine metabolic inability to generate the energy required for human connection.

3.13 Sleep-Related Symptoms

Beyond unrefreshing sleep (a core symptom), ME/CFS patients experience various sleep disturbances.

3.13.1 Insomnia

Clinical Presentation.

- Difficulty initiating sleep (sleep onset insomnia)
- Difficulty maintaining sleep (sleep maintenance insomnia)
- Early morning awakening
- “Tired but wired” sensation (exhausted but unable to sleep)

3.13.2 Hypersomnia

Clinical Presentation.

- Excessive sleep need (12–18+ hours per day in severe cases)
- Inability to stay awake during day
- Sleep attacks or sudden overwhelming sleepiness
- Difficulty waking despite prolonged sleep

3.13.3 Sleep Architecture Abnormalities

Polysomnography Findings.

- Reduced slow-wave sleep (deep sleep)
- Alpha-wave intrusion into non-REM sleep
- Fragmented sleep with frequent arousals
- REM sleep abnormalities

Observation 3 (Formal Sleep Evaluation and Treatable Sleep Pathology as Critical Intervention). Patient case experience indicates that formal sleep laboratory evaluation (polysomnography and sleep multiple sleep latency testing) is underutilized in ME/CFS and may identify treatable pathology not captured by clinical history alone. Specific conditions documented in ME/CFS patient cohorts include: sleep apnea (obstructive and central), periodic limb movement disorder, rapid eye movement sleep behavior disorder, and circadian rhythm disruption. While the majority of ME/CFS patients report unrefreshing sleep as a diagnostic feature, the *reason* for unrefreshingness varies by individual. Some have structural sleep pathology (apnea, arousals) that is medically treatable; others have insufficient deep sleep stage; others have circadian dysrhythmia. Clinical observation suggests that identification and treatment of specific sleep pathology may substantially improve functional capacity and reduce PEM severity, independent of other ME/CFS interventions. Sleep apnea treatment (CPAP), periodic limb movement suppression (dopaminergic agents, iron), and circadian realignment (light therapy, melatonin timing) all show potential utility. The unrefreshing sleep of ME/CFS should prompt formal sleep study in all patients with available access, as treatable sleep pathology may represent an overlooked therapeutic target. Restoring sleep quality is mechanistically plausible to reduce PEM susceptibility (through improved cellular recovery during sleep) and stabilize metabolic function.

3.13.4 Restless Legs Syndrome and Periodic Limb Movements

Clinical Presentation.

- Uncomfortable sensations in legs at rest
- Urge to move legs to relieve discomfort
- Symptoms worsening at night
- Involuntary leg movements during sleep (periodic limb movement disorder)

3.14 Symptom Severity Spectrum

ME/CFS symptoms exist on a spectrum from mild to very severe. Understanding this spectrum is critical for recognizing disease heterogeneity and avoiding minimization of severe cases.

3.14.1 Mild ME/CFS

Functional Capacity.

- Able to work or study, but with significant difficulty
- Must reduce activities and rest frequently
- Symptoms worsen with exertion but recovery possible with pacing
- Can perform basic self-care and some household tasks
- Social life significantly reduced

Common Symptom Profile.

- Moderate fatigue and PEM with predictable triggers
- Cognitive impairment affecting work performance
- Mild to moderate pain
- Sleep disturbance
- Orthostatic symptoms manageable

3.14.2 Moderate ME/CFS

Functional Capacity.

- Unable to work full-time or maintain consistent employment
- Housebound part of the time
- Can perform some self-care but requires frequent rest
- Severe reduction in activities compared to pre-illness
- PEM more severe and prolonged

Common Symptom Profile.

- Significant fatigue and PEM lasting days to weeks
- Marked cognitive impairment
- Moderate to severe pain
- Orthostatic intolerance limiting upright time
- Multiple sensory sensitivities

3.14.3 Severe ME/CFS

Functional Capacity.

- Housebound or bedbound most of the time
- Unable to perform most self-care without assistance

- May use wheelchair for any movement
- Very limited tolerance for activity
- PEM triggered by minimal exertion (showering, eating, conversation)

Common Symptom Profile.

- Profound fatigue and PEM lasting weeks to months
- Severe cognitive impairment (difficulty reading, watching TV, following conversation)
- Severe pain requiring management
- Profound orthostatic intolerance (unable to sit or stand without symptoms)
- Multiple severe sensory sensitivities (light, sound, touch)
- Difficulty eating (nausea, GI symptoms, effort of chewing/swallowing)

3.14.4 Very Severe ME/CFS

Functional Capacity.

- Bedbound continuously
- Unable to perform any self-care
- Requires full nursing care
- Cannot tolerate light, sound, touch, or human presence
- May require tube feeding
- Minimal or no communication possible

Common Symptom Profile.

- Any stimulation triggers immediate, severe worsening
- Complete darkness and silence required
- Touch causes pain
- Swallowing may be impaired
- May be unable to tolerate being moved or bathed
- Life-threatening complications (malnutrition, pressure sores, infections)

Clinical Note. Very severe ME/CFS represents a medical emergency and requires specialized care. These patients are profoundly vulnerable and often invisible to the medical system because they cannot attend appointments. Mortality risk is elevated due to complications of immobility, malnutrition, and suicide.

3.15 Summary: The Multi-System Nature of ME/CFS

ME/CFS is not a single-system disorder but a multi-system disease affecting virtually every physiological system. The sheer breadth of symptoms—neurological, immunological, musculoskeletal, cardiovascular, respiratory, gastrointestinal, genitourinary, endocrine, dermatological, ocular, and psychological—underscores the systemic nature of the underlying pathophysiology.

Key Concepts.

- **Heterogeneity:** No two patients have identical symptom profiles. Some patients have predominantly neurological symptoms, others gastrointestinal, others autonomic. This heterogeneity suggests multiple disease subtypes or different triggering events leading to similar outcomes.
- **Severity spectrum:** ME/CFS ranges from mild (able to work with difficulty) to very severe (bedbound, unable to tolerate any stimulation). Severity is a continuum, not discrete categories.
- **Symptom fluctuation:** Most symptoms fluctuate over time, worsening during PEM crashes and partially improving during baseline periods. This variability makes the disease particularly unpredictable and difficult to manage.
- **Cumulative burden:** While individual symptoms may seem manageable in isolation, the cumulative burden of dozens of simultaneous symptoms creates profound disability. Patients must constantly prioritize which symptoms to tolerate and which to attempt to mitigate.
- **Energy as common thread:** Nearly all symptoms can be traced back to inadequate cellular energy production (mitochondrial dysfunction), immune dysregulation, and autonomic dysfunction. These core pathophysiological mechanisms produce the diverse symptom manifestations across body systems.

The comprehensive symptom catalog presented in this chapter serves multiple purposes: validating patient experiences, educating healthcare providers, guiding diagnosis, informing treatment planning, and demonstrating the profound, multi-system impact of ME/CFS. Recognition of the full spectrum of symptoms is essential for appropriate diagnosis, avoiding misattribution to psychiatric causes, and providing compassionate, comprehensive care.

4 Diagnostic Criteria and Clinical Assessment

Multiple diagnostic criteria have been developed for ME/CFS. This chapter reviews major frameworks and their application.

4.1 Overview of Diagnostic Approaches

ME/CFS lacks a definitive diagnostic test. Diagnosis remains clinical, based on characteristic symptom patterns, exclusion of alternative explanations, and increasingly, supportive objective findings. Multiple diagnostic frameworks have been proposed, each with different emphases and populations identified.

4.1.1 Evolution of Diagnostic Criteria

The history of ME/CFS diagnostic criteria reflects evolving understanding of the illness:

Early Frameworks (1988–1994)

- **Holmes Criteria (1988):** First CDC case definition; required 6+ months of persistent/re-lapsing fatigue plus 8 of 11 symptom criteria or 6 of 11 symptoms plus 2 of 3 physical criteria
- **Problem:** Overly broad; captured patients with primary depression or deconditioning; lacked specificity for the post-exertional phenotype
- **Fukuda Criteria (1994):** Simplified to 6 months of unexplained fatigue plus 4 of 8 symptoms; removed physical examination criteria
- **Impact:** Became the dominant research framework but remained criticized for heterogeneity

Restrictive Frameworks (2003–2011)

- **Canadian Consensus Criteria (2003):** Required post-exertional malaise/fatigue, sleep dysfunction, pain, and two neurological/cognitive manifestations plus one autoimmune/neuroendocrine/immune manifestation
- **International Consensus Criteria (2011):** Emphasized post-exertional *neuroimmune* exhaustion as mandatory; required neurological impairments, immune/gastrointestinal/genitourinary impairments, and energy metabolism/transport impairments

- **Rationale:** Select more homogeneous, severely affected patients; exclude patients with primary psychiatric conditions
- **Trade-off:** Higher specificity but lower sensitivity; may miss milder cases

Consensus Framework (2015)

- **IOM Criteria (2015):** Proposed by Institute of Medicine (now National Academy of Medicine) systematic review
- **Required symptoms:** (1) Substantial impairment lasting ≥ 6 months, (2) Post-exertional malaise, (3) Unrefreshing sleep, (4) Cognitive impairment OR orthostatic intolerance
- **Goal:** Facilitate clinical diagnosis while maintaining specificity
- **Renamed condition:** Proposed “Systemic Exertion Intolerance Disease (SEID)” to emphasize cardinal feature; name not widely adopted

4.1.2 Comparison of Frameworks: Set-Theoretic Analysis

Different criteria identify overlapping but distinct patient populations. Let F , C , I , and O denote the sets of patients meeting Fukuda, Canadian Consensus, ICC, and IOM criteria respectively.

Observation 4 (Hierarchical Inclusion Relationships). The diagnostic frameworks exhibit a partial ordering by restrictiveness:

$$I \subset C \subset F \quad \text{and} \quad I \cap O \neq \emptyset \quad (4.1)$$

where:

- $I \subset C$: All patients meeting ICC criteria also meet Canadian Consensus criteria (ICC is more restrictive)
- $C \subset F$: All patients meeting Canadian Consensus criteria also meet Fukuda criteria (Canadian is more restrictive than Fukuda)
- $I \cap O \neq \emptyset$: ICC and IOM criteria identify overlapping but not identical populations

This hierarchical structure implies that stricter criteria select subsets of the broader Fukuda population, potentially enriching for more severely affected patients with more homogeneous pathophysiology.

Empirical Comparison Brown et al. [112] compared phenotypes identified by different criteria in the same cohort. Key findings:

Interpretation: Criteria do not identify different diseases but rather different severity strata of the same condition. Patients meeting multiple criteria have worse outcomes, suggesting the stricter frameworks capture more severely affected individuals.

Table 4.1: Symptom severity by diagnostic criteria met

Criteria Met	n	Cognitive	Autonomic	Symptom Burden
Fukuda only	45	Mild	Mild	Lowest
Fukuda + Canadian	103	Moderate	Moderate	Intermediate
Fukuda + Canadian + ICC	52	Severe	Severe	Highest

4.1.3 Sensitivity and Specificity Trade-offs

No gold standard exists for ME/CFS, making traditional sensitivity/specificity calculations impossible. However, we can assess **predictive validity**: do patients meeting stricter criteria show more consistent biomarker abnormalities and treatment responses?

~ Hypothesis 1: Specificity-Homogeneity Trade-off

Stricter diagnostic criteria increase disease homogeneity (reducing phenotypic variance) at the cost of excluding milder cases. Formally:

Let $\sigma_{\text{pheno}}^2(X)$ denote phenotypic variance within the population X meeting criterion set. Then:

$$\sigma_{\text{pheno}}^2(I) < \sigma_{\text{pheno}}^2(C) < \sigma_{\text{pheno}}^2(F) \quad (4.2)$$

This implies:

- **Research advantage:** ICC/Canadian cohorts show more consistent biomarker patterns, improving statistical power
- **Clinical disadvantage:** Mild cases may be missed, delaying diagnosis and early intervention
- **Treatment trials:** Stricter criteria may improve signal detection but limit generalizability

The optimal criterion set depends on context: research requires homogeneity; clinical practice requires sensitivity.

4.1.4 Clinical Implications of Framework Choice

For Research

- **Biomarker studies:** Use ICC or Canadian Consensus to minimize heterogeneity
- **Treatment trials:** Specify criteria explicitly; consider stratifying by criteria met
- **Meta-analyses:** Account for criteria differences when combining studies

For Clinical Diagnosis

- **Initial assessment:** IOM criteria provide good balance of sensitivity and specificity

- **Severe cases:** Will meet multiple criteria; diagnosis is straightforward
- **Mild/early cases:** May not yet meet all symptom requirements; consider provisional diagnosis with reassessment
- **Documentation:** Record which criteria are met to facilitate comparison across studies

4.2 Canadian Consensus Criteria (2003)

The Canadian Consensus Criteria [28] emerged from a panel of physicians, researchers, and teaching faculty to provide a clinically oriented case definition emphasizing characteristic features.

4.2.1 Required Criteria

• Requirement 1: Canadian Consensus Criteria Structure

Diagnosis requires ALL of the following:

1. Fatigue

- Clinically evaluated, unexplained persistent or relapsing chronic fatigue
- New or definite onset (not lifelong)
- Not result of ongoing exertion
- Not substantially alleviated by rest
- Substantial reduction in previous levels of occupational, educational, social, or personal activities

2. Post-Exertional Malaise and/or Fatigue (MANDATORY)

- Inappropriate loss of physical and mental stamina
- Rapid muscular and cognitive fatigability
- Post-exertional malaise and/or fatigue
- Tendency for other associated symptoms within the patient's cluster to worsen
- **Recovery period:** Pathologically slow (24 hours or longer)

3. Sleep Dysfunction

- Unrefreshing sleep or sleep quantity or rhythm disturbances (hypersomnia, insomnia, reversed sleep-wake cycle)

4. Pain (significant degree in at least one location)

- Myalgia: muscle pain, aching, or stiffness
- Arthralgia: joint pain (migratory, without joint swelling or redness)

- Headaches of new type, pattern, or severity

5. Neurological/Cognitive Manifestations (≥2 required)

- Confusion, impaired concentration/short-term memory, disorientation
- Difficulty with information processing, categorizing, word retrieval
- Perceptual/sensory disturbances (spatial instability, disorientation, inability to focus vision)
- Ataxia, muscle weakness, fasciculations

6. At Least ONE Symptom from TWO of the Following Categories:

- **Autonomic Manifestations:** Orthostatic intolerance (neurally mediated hypotension, POTS, delayed postural hypotension), lightheadedness, extreme pallor, nausea and irritable bowel syndrome, urinary frequency/bladder dysfunction, palpitations with or without cardiac arrhythmias, exertional dyspnea
- **Neuroendocrine Manifestations:** Loss of thermostatic stability (subnormal body temperature, marked daily fluctuation), intolerance of extremes of heat and cold, marked weight change, loss of adaptability/worsening symptoms with stress
- **Immune Manifestations:** Tender lymph nodes, recurrent sore throat, recurrent flu-like symptoms, general malaise, new sensitivities to food/medications/chemicals

7. Duration Illness persists ≥6 months. May be preceded by various infections or other triggering events.

4.2.2 Exclusions

Exclude active disease processes that explain most major symptoms. Comorbid conditions do not exclude diagnosis if they do not explain the constellation of symptoms.

4.2.3 Strengths and Limitations

Strengths

- Emphasis on post-exertional malaise as mandatory criterion
- Comprehensive symptom coverage across multiple systems
- Widely adopted in clinical practice
- More specific than Fukuda criteria

Limitations

- Complex algorithm may be difficult to apply consistently
- Selects more severely affected patients (lower sensitivity for mild cases)

- Some symptom requirements (e.g., “significant degree”) lack operational definitions
- Not validated against objective biomarkers or treatment response

4.3 International Consensus Criteria (2011)

The International Consensus Criteria (ICC), published in 2011 by Carruthers et al. [29], represents the most restrictive and biologically-oriented diagnostic framework. The ICC explicitly adopts the term “myalgic encephalomyelitis” (ME) to emphasize the neurological and immunological features of the disease, rejecting the broader “chronic fatigue syndrome” label as insufficiently specific.

4.3.1 Required Criteria

• Requirement 2: International Consensus Criteria Structure

Diagnosis of myalgic encephalomyelitis requires **post-exertional neuroimmune exhaustion (PENE)** as the mandatory hallmark PLUS manifestations from at least THREE neurological impairment categories PLUS at least ONE manifestation from each of THREE immune/gastro-intestinal/genitourinary, energy metabolism/transport, and cardiovascular/respiratory/thermoregulatory categories.

A. Post-Exertional Neuroimmune Exhaustion (PENE) — MANDATORY PENE is the central diagnostic feature and must be present.

Pathological inability to produce sufficient energy on demand with the following characteristics:

- **Marked, rapid physical and/or cognitive fatigability** in response to exertion
- **Post-exertional symptom exacerbation:** Disproportionate loss of physical and mental stamina, rapid muscular and cognitive fatigability, post-exertional malaise and/or pain, and tendency for other associated symptoms to worsen
- **Post-exertional exhaustion:** May occur immediately after activity or be delayed by hours or days
- **Recovery period is prolonged:** Usually 24 hours or longer
- **Low threshold of physical and mental fatigability:** Results in substantial reduction in pre-illness activity level

B. Neurological Impairments (at least THREE required)

1. **Neurocognitive Impairments:**

- Difficulty processing information (slowed thought, impaired concentration)
- Short-term memory loss
- Word-finding difficulty, impaired psychomotor function

- Perceptual/sensory disturbances (spatial instability, disorientation, inability to focus vision)
- Ataxia, muscle weakness, fasciculations

2. **Pain:**

- Headaches (new type, pattern, or severity)
- Significant pain in muscles, muscle-tendon junctions, joints, abdomen, or chest
- Pain can be migratory, generalized or localized, often changing in distribution

3. **Sleep Disturbance:**

- Disturbed sleep patterns: insomnia, prolonged sleep (hypersomnia), disturbed sleep/wake cycle
- Unrefreshing sleep: Patient awakens feeling exhausted regardless of sleep duration

4. **Neurosensory, Perceptual, and Motor Disturbances:**

- Sensory hypersensitivity: photophobia, hyperacusis, heightened sensitivities to odors, taste, touch
- Motor disturbances: muscle weakness, twitching, poor coordination, ataxia

C. Immune, Gastro-Intestinal, and Genitourinary Impairments (at least ONE)

- **Immune:** Tender lymph nodes, recurrent sore throat, recurrent flu-like symptoms, general malaise, new sensitivities to food/medications/chemicals
- **Gastro-intestinal:** Nausea, abdominal pain, bloating, irritable bowel syndrome
- **Genitourinary:** Urinary urgency or frequency, nocturia

D. Energy Production/Transportation Impairments (at least ONE)

- **Cardiovascular:** Inability to tolerate upright posture (orthostatic intolerance, neurally mediated hypotension, postural orthostatic tachycardia syndrome), palpitations, lightheadedness
- **Respiratory:** Dyspnea, labored breathing, air hunger
- **Loss of thermostatic stability:** Subnormal body temperature, marked diurnal fluctuation, sweating episodes, cold extremities, intolerance to heat or cold
- **Intolerance to extremes of temperature**

4.3.2 Phenotype Categories

The ICC proposes operational phenotype categories to capture disease heterogeneity:

1. **ME with Fibromyalgia:** Patients meeting ME criteria with widespread pain and tenderness
2. **ME with Myofascial Pain Syndrome:** Regional pain with trigger points

3. **ME with Postural Orthostatic Tachycardia Syndrome (POTS):** ME with documented autonomic dysfunction
4. **ME with Irritable Bowel Syndrome:** ME with prominent gastrointestinal manifestations
5. **ME with Multiple Chemical Sensitivity:** ME with sensitivity to environmental chemicals

These categories are **not mutually exclusive**; patients may meet criteria for multiple phenotypes simultaneously.

4.3.3 Duration and Exclusions

- **Duration:** Symptom persistence for at least **6 months**
- **Pediatric Exception:** In children and adolescents, 3 months may be sufficient for diagnosis given the urgency of early intervention
- **Exclusions:** Active disease processes that explain most symptoms must be ruled out (e.g., untreated hypothyroidism, obstructive sleep apnea)
- **Comorbidities allowed:** Fibromyalgia, myofascial pain, temporomandibular disorder, irritable bowel syndrome, interstitial cystitis, Raynaud phenomenon, mitral valve prolapse, migraines can coexist with ME

4.3.4 Strengths and Limitations

Observation 5 (ICC Strengths). The ICC framework has several advantages:

- **Biological orientation:** Emphasizes objective neurological and immune manifestations rather than subjective fatigue
- **Post-exertional neuroimmune exhaustion as mandatory:** Recognizes PEM as the pathognomonic feature
- **Multi-system requirement:** Requires manifestations across multiple physiological systems, increasing specificity
- **Phenotype categories:** Acknowledges heterogeneity and common comorbidities
- **Higher specificity:** More restrictive than Canadian Consensus or Fukuda, resulting in more homogeneous research cohorts

△ **Warning 1: ICC Limitations**

The restrictiveness of ICC creates challenges:

- **Excludes mild cases:** Patients with genuine ME/CFS who do not yet manifest symptoms across all required categories may be missed
- **Clinical impracticality:** Detailed assessment across 8 categories requires extensive clinical time and expertise
- **Reduced sensitivity:** Systematic review found ICC identifies only 60% of patients

meeting Canadian Consensus Criteria [113]

- **Formal set-theoretic relationship:** $\text{ICC} \subset \text{Canadian Consensus} \subset \text{Fukuda}$ — ICC is the most restrictive subset
- **Delayed diagnosis risk:** Waiting for full symptom constellation may delay intervention during the critical 6-month window

4.3.5 Research and Clinical Application

The ICC is **optimal for research** where high specificity and phenotypic homogeneity are priorities, reducing heterogeneity that can obscure treatment signals. However, for **clinical practice**, the more inclusive Canadian Consensus Criteria or IOM criteria are preferred to avoid missing early-stage or mild cases that would benefit from intervention.

4.4 Institute of Medicine Criteria (2015)

The Institute of Medicine (now National Academy of Medicine) published diagnostic criteria in 2015 following a comprehensive systematic review [6]. The IOM report proposed renaming the condition “Systemic Exertion Intolerance Disease” (SEID) to emphasize the central role of post-exertional malaise and to move away from the stigmatizing “chronic fatigue” label. However, the SEID terminology has seen limited clinical adoption.

4.4.1 Required Core Symptoms

• Requirement 3: IOM Diagnostic Algorithm

Diagnosis requires ALL THREE of the following core symptoms to be present:

1. Substantial Reduction or Impairment in Activity Level (MANDATORY) A substantial reduction or impairment in the ability to engage in pre-illness levels of occupational, educational, social, or personal activities that:

- Persists for **more than 6 months**
- Is accompanied by fatigue (often profound)
- Is of **new or definite onset** (not lifelong)
- Is **not the result of ongoing excessive exertion**
- Is **not substantially alleviated by rest**

2. Post-Exertional Malaise (PEM) — MANDATORY Worsening of symptoms following physical, cognitive, or emotional exertion that would not have caused a problem before illness. Characteristics include:

- Symptoms typically worsen 12–48 hours after activity
- Often leads to relapse lasting days, weeks, or longer
- Exertion threshold for triggering symptoms is low
- Recovery is prolonged

The IOM emphasizes that PEM is **the hallmark symptom** that distinguishes ME/CFS from other fatiguing conditions.

3. Unrefreshing Sleep (MANDATORY) Patients wake feeling unrefreshed regardless of sleep duration. Sleep may be:

- Disrupted (frequent awakenings, difficulty initiating sleep)
- Prolonged (hypersomnia with no restoration)
- Reversed sleep/wake cycle

The exhaustion persists despite adequate sleep duration.

4.4.2 Additional Required Symptoms

• **Requirement 4: At Least ONE of the Following**

Cognitive Impairment Problems with thinking, memory, information processing, or executive function. May include:

- Difficulty finding words, storing and retrieving information
- Slowed processing speed
- Inability to focus or multitask
- Problems with short-term memory

Cognitive symptoms may worsen with physical or mental exertion, emotional stress, or time pressure.

OR

Orthostatic Intolerance Worsening of symptoms upon assuming or maintaining upright posture. May include:

- Lightheadedness, dizziness, fainting
- Worsening fatigue or cognitive impairment when upright
- Palpitations, nausea
- Symptoms improve (but may not resolve) when lying down

Objective findings may include abnormal heart rate or blood pressure responses during tilt table testing or standing test.

4.4.3 Diagnostic Algorithm Structure

The IOM criteria can be formalized as a logical algorithm:

$$\text{ME/CFS}_{\text{IOM}} = \left\{ \begin{array}{l} \text{Substantial Activity Reduction} \\ \wedge \text{Post-Exertional Malaise} \\ \wedge \text{Unrefreshing Sleep} \\ \wedge (\text{Cognitive Impairment} \vee \text{Orthostatic Intolerance}) \\ \wedge \text{Duration} \geq 6 \text{ months} \\ \wedge \text{Exclusions ruled out} \end{array} \right\} \quad (4.3)$$

This represents a **minimal sufficient set**: three universal core features plus at least one of two common manifestations.

4.4.4 Exclusions and Comorbidities

- **Exclusions:** Medical conditions that could fully explain the symptoms must be ruled out through appropriate testing (hypothyroidism, anemia, sleep apnea, etc.)
- **Comorbidities allowed:** Fibromyalgia, irritable bowel syndrome, depression, and anxiety frequently co-occur and do not exclude ME/CFS diagnosis
- **Important distinction:** Comorbid depression is reactive (consequence of severe disability) rather than causative

4.4.5 Strengths and Limitations

Observation 6 (IOM Strengths). The IOM criteria offer several advantages:

- **Simplicity:** Four required features (3 core + 1 of 2 additional) make diagnosis straightforward
- **High sensitivity:** Captures broader range of ME/CFS patients than ICC or Canadian Consensus
- **Evidence-based:** Derived from systematic review identifying most discriminating symptoms
- **PEM emphasis:** Recognizes post-exertional malaise as the pathognomonic feature
- **Clinical practicality:** Feasible in primary care settings without extensive symptom checklists
- **Rapid assessment:** Can be evaluated in a standard office visit

△ Warning 2: IOM Limitations

The simplified structure creates potential issues:

- **Reduced specificity:** More inclusive criteria may capture patients with other con-

ditions (long COVID, post-viral fatigue that will resolve)

- **Heterogeneity:** Broader patient population increases phenotypic variance in research cohorts
- **Cognitive OR orthostatic requirement:** Patients may meet criteria with only one of these domains, potentially missing multi-system nature
- **SEID terminology rejected:** Proposed name change has not gained acceptance in patient or research communities
- **Set-theoretic relationship:** ICC $\subset$ Canadian $\subset$ IOM — IOM captures the broadest population

4.4.6 Clinical and Research Application

Observation 7 (When to Use IOM Criteria). **For clinical diagnosis:** The IOM criteria are excellent for primary care and general practice:

- Simple enough for non-specialists to apply
- High sensitivity ensures few false negatives
- Enables early diagnosis and intervention

For research: The IOM criteria are appropriate when:

- Study aims to represent the full ME/CFS population
- Recruitment needs to be pragmatic and efficient
- Results should generalize to clinical settings

Not optimal for: Mechanistic research or treatment trials requiring homogeneous cohorts (use ICC or Canadian Consensus with biomarker stratification instead).

4.5 Other Diagnostic Frameworks

4.5.1 Fukuda Criteria (1994)

The Fukuda criteria, published by the CDC in 1994 [27], represented the first widely-adopted standardized definition of chronic fatigue syndrome. These criteria dominated ME/CFS research for two decades.

• Requirement 5: Fukuda Diagnostic Criteria

Diagnosis requires:

1. **Clinically Evaluated, Unexplained, Persistent or Relapsing Chronic Fatigue that:**

- Is of **new or definite onset** (not lifelong)
- Is **not the result of ongoing exertion**
- Is **not substantially alleviated by rest**
- Results in **substantial reduction** in previous levels of occupational, educational, social, or personal activities

2. Four or More of the Following Symptoms (concurrent for ≥ 6 months):

1. Impaired memory or concentration
2. Sore throat
3. Tender cervical or axillary lymph nodes
4. Muscle pain
5. Multi-joint pain without swelling or redness
6. Headaches of new type, pattern, or severity
7. Unrefreshing sleep
8. Post-exertional malaise lasting more than 24 hours

Observation 8 (Historical Significance). The Fukuda criteria played a crucial role in standardizing ME/CFS research:

- First internationally-adopted consensus definition
- Enabled comparison across studies and centers
- Established 6-month duration threshold
- Required objective clinical evaluation

△ Warning 3: Critical Limitations

The Fukuda criteria have fundamental flaws that limit their current utility:

Post-Exertional Malaise Not Mandatory: The most pathognomonic feature of ME/CFS (PEM) is merely one of eight optional symptoms. This allows diagnosis of patients without the hallmark feature, including those with:

- Primary depression
- Deconditioning
- Other fatiguing conditions without PEM

Mathematical Analysis of Heterogeneity: The requirement of “4 or more of 8 symptoms” yields:

$$\binom{8}{4} + \binom{8}{5} + \binom{8}{6} + \binom{8}{7} + \binom{8}{8} = 70 + 56 + 28 + 8 + 1 = 163 \text{ distinct profiles} \quad (4.4)$$

Two patients can both meet Fukuda criteria while sharing only 2 of 8 symptoms (50%)

overlap in the limiting case). This mathematical heterogeneity explains null results in many Fukuda-based trials.

Overinclusion: Systematic comparison studies found that Fukuda criteria capture patients who do not meet more restrictive criteria (Canadian Consensus, ICC) and who have:

- Less severe functional impairment
- Better prognosis
- Lower biomarker abnormality rates
- Higher rates of primary psychiatric diagnoses

Research Impact: Many failed clinical trials used Fukuda criteria, likely enrolling heterogeneous populations including patients without true ME/CFS. This contributed to therapeutic nihilism.

Observation 9 (Current Status). Modern research increasingly avoids Fukuda criteria in favor of Canadian Consensus (2003), ICC (2011), or IOM (2015). The Fukuda criteria remain historically important but are now recognized as insufficiently specific for ME/CFS.

4.5.2 Oxford Criteria (1991)

The Oxford criteria [114], published in 1991, represent the **broadest and most controversial** definition of chronic fatigue syndrome.

• Requirement 6: Oxford Diagnostic Criteria

Diagnosis requires:

1. **Severe disabling fatigue** of at least 6 months' duration that:
 - Affects physical and mental functioning
 - Was present for more than 50% of the time
2. **Other symptoms**, particularly myalgia, mood disturbance, and sleep disturbance, may be present
3. **Exclusions:** Defined medical conditions, psychotic disorders, substance abuse, eating disorders
4. **Depression and anxiety NOT excluded**

△ Warning 4: Fundamental Problems with Oxford Criteria

The Oxford criteria are widely rejected by patients, clinicians, and researchers for the following reasons:

No Requirement for Post-Exertional Malaise: The pathognomonic feature of ME/CFS is entirely absent. Patients meeting Oxford criteria may have:

- Primary depression with fatigue
- Deconditioning from sedentary lifestyle
- Idiopathic chronic fatigue (fatigue without clear cause)

Allows Primary Psychiatric Diagnoses: Unlike all other ME/CFS criteria, Oxford explicitly allows comorbid depression and anxiety *even when these could fully explain the fatigue*. This conflates ME/CFS with depression-related fatigue.

Set-Theoretic Implications: Define patient populations by criteria:

$$O \supset F \supset C \supset I \quad (4.5)$$

where O = Oxford, F = Fukuda, C = Canadian Consensus, I = ICC.

The Oxford criteria capture a superset that includes patients with ME/CFS (satisfying more restrictive criteria) plus patients with primary depression, idiopathic fatigue, and deconditioning.

Harm from CBT/GET Trials: The most harmful aspect: Oxford criteria were used in the PACE trial [115] and other studies promoting cognitive behavioral therapy (CBT) and graded exercise therapy (GET) as treatments for “CFS.” However:

- Patient surveys show GET causes harm in 50–70% of ME/CFS patients [116]
- CBT/GET may be appropriate for depression or deconditioning but are contraindicated for true ME/CFS
- By enrolling patients without PEM, these trials tested interventions on a population distinct from ME/CFS

Observation 10 (Research Community Consensus). The Oxford criteria are now explicitly rejected by:

- The NIH (U.S. National Institutes of Health) ME/CFS research guidelines
- The CDC (U.S. Centers for Disease Control)
- Leading ME/CFS researchers and clinicians
- Patient advocacy organizations

Studies using Oxford criteria should be interpreted with extreme caution, as they likely include substantial proportions of patients without ME/CFS.

4.5.3 Pediatric Criteria

ME/CFS in children and adolescents presents diagnostic challenges due to developmental differences in symptom expression, comorbidities, and functional impact.

• Requirement 7: Pediatric Adaptations

Diagnosis in children should use the same criteria (Canadian Consensus, IOM, or ICC) with the following modifications:

1. Duration:

- Standard: 6 months in adults
- Pediatric: May use **3 months** if symptoms are severe and progression is documented
- Rationale: Early diagnosis enables intervention during critical developmental windows

2. Activity Reduction:

- Assess relative to **age-appropriate activities**: school attendance, sports participation, social activities with peers
- May manifest as: inability to attend full school day, requiring home tutoring, dropping out of extracurricular activities
- Pediatric patients may have better baseline reserves, so functional impairment can be harder to detect

3. Post-Exertional Malaise:

- Children may not articulate PEM clearly; ask caregivers about: “Does your child crash after activities?”
- School attendance patterns are diagnostic: can attend Monday but not Tuesday (PEM delay)
- May manifest as behavioral changes (irritability, emotional lability) rather than reported exhaustion

4. Cognitive Symptoms:

- Assess relative to prior academic performance, not population norms
- May manifest as: declining grades, inability to complete homework, processing speed reduction
- Distinguish from learning disabilities (which would have been present earlier)

5. Orthostatic Intolerance:

- Highly prevalent in pediatric ME/CFS (70–90%)
- May present as: difficulty standing in school assemblies, morning symptom worsening (after overnight recumbency), shower intolerance
- Objective testing: NASA Lean Test or tilt table (age-appropriate protocols)

△ **Warning 5: Pediatric Differential Diagnosis**

Additional considerations for children:

- **School avoidance vs. ME/CFS:** Distinguish by presence of PEM (in ME/CFS, even desired activities trigger crashes)
- **Growth and puberty:** Rule out growth-related fatigue, iron deficiency from menstruation
- **Viral triggers:** Infectious mononucleosis is a common ME/CFS trigger in adolescents
- **Comorbidities:** POTS and orthostatic intolerance are especially common in pediatric onset

Observation 11 (Prognosis and Early Intervention). Pediatric ME/CFS has distinct prognostic features:

- **Better prognosis than adults:** Some studies suggest 50–70% improvement or recovery rates in adolescents, though methodological issues may inflate these estimates
- **Critical intervention window:** Early aggressive pacing and school accommodation may prevent progression to severe disease
- **Educational impact:** Lost school years during critical developmental periods create long-term consequences
- **Recommendation:** Diagnose at 3 months if symptoms are severe; immediate school accommodations (reduced hours, remote learning, rest breaks) to prevent cumulative PEM damage

For detailed pediatric treatment protocols, see Chapter 19 (severe/housebound cases) and Chapter 20 (school-attending cases).

4.6 Clinical Assessment

A thorough clinical assessment is essential for ME/CFS diagnosis, both to establish the presence of diagnostic criteria and to rule out alternative explanations for symptoms.

4.6.1 History Taking

Onset Characterization

Observation 12 (Onset Patterns as Diagnostic Clues). The pattern of disease onset provides diagnostic and prognostic information:

Sudden Onset (60–80% of cases):

- Patient can identify exact date or event when illness began
- Most commonly follows acute infection: infectious mononucleosis (EBV), influenza, COVID-19, gastroenteritis
- May follow other physiological stressors: surgery, trauma, childbirth, vaccination
- Strong temporal association suggests post-infectious mechanism
- **Diagnostic value:** Sudden onset after documented infection strongly supports ME/CFS diagnosis

Gradual Onset (20–40% of cases):

- Symptoms develop over weeks to months without clear precipitant
- May follow period of chronic stress, cumulative sleep deprivation, or overwork
- Patient cannot identify specific triggering event
- **Diagnostic challenge:** Gradual onset requires more thorough differential diagnosis (autoimmune disease, occult malignancy, endocrine disorders)

Key Questions for Establishing Diagnosis

Post-Exertional Malaise Assessment:

1. “After physical activity, do you feel worse than expected?”
2. “Is there a delay between activity and symptom worsening? How long?” (Typical: 12–72 hours)
3. “How long does it take to recover from doing too much?” (ME/CFS: days to weeks)
4. “Can you reliably predict what activities will make you crash?”
5. “Do mental tasks (reading, concentration) also trigger symptom flares?” (Cognitive PEM distinguishes ME/CFS from deconditioning)

Sleep Assessment:

1. “Do you wake up feeling refreshed?” (ME/CFS: No, regardless of duration)
2. “How many hours do you sleep?” (Rule out insufficient sleep)
3. “Do you snore? Have you been told you stop breathing during sleep?” (Screen for obstructive sleep apnea)
4. “What time do you go to sleep and wake up?” (Assess circadian rhythm disorders)

Cognitive Dysfunction:

1. "Do you have trouble finding words or finishing sentences?"
2. "Do you lose your train of thought mid-conversation?"
3. "Can you read and retain information like you used to?"
4. "Do you have difficulty with tasks that require sustained focus?"
5. "Are these problems worse after physical or mental exertion?" (Cognitive PEM)

Orthostatic Symptoms:

1. "Do you feel dizzy or lightheaded when standing up?"
2. "Are showers or baths difficult? Do you need to sit?"
3. "Do your symptoms worsen when standing for extended periods?"
4. "Do you feel better when lying down?"

Functional Impact Assessment:

1. "What percentage of your pre-illness activity level can you sustain now?"
2. "What activities have you had to give up?" (Work, social activities, hobbies, childcare)
3. "On a scale of 0–100 (Bell Disability Scale), what is your functional capacity?"
4. "How many hours per week do you spend horizontal (lying down)?"

4.6.2 Physical Examination

Orthostatic Vital Signs

• **Requirement 8: NASA Lean Test or Orthostatic Vital Signs**

Orthostatic intolerance assessment should be performed on all ME/CFS patients:

Protocol:

1. Patient supine for 5 minutes → measure heart rate (HR) and blood pressure (BP)
2. Patient stands upright → measure HR and BP at 1, 3, 5, and 10 minutes
3. Record symptoms during test (lightheadedness, nausea, cognitive impairment)

Abnormal Findings:

- **Postural Orthostatic Tachycardia Syndrome (POTS):** HR increase ≥ 30 bpm (or ≥ 40 bpm in adolescents) within 10 minutes of standing, without orthostatic hypotension
- **Orthostatic Hypotension:** Systolic BP decrease ≥ 20 mmHg or diastolic BP decrease ≥ 10 mmHg

- **Neurally Mediated Hypotension (NMH):** Delayed BP drop (after 5–10 minutes standing)
- **Symptom reproduction:** Patient reports typical symptoms even without meeting BP/HR criteria

Interpretation: 70–90% of ME/CFS patients demonstrate orthostatic intolerance on objective testing. Absence of objective findings does not exclude ME/CFS, but presence strongly supports the diagnosis and guides treatment (salt, fluids, fludrocortisone, midodrine).

Neurological Examination

Observation 13 (Neurological Findings in ME/CFS). The neurological examination in ME/CFS typically shows:

Usually Normal:

- Cranial nerves intact
- Motor strength 5/5 (though patients report subjective weakness)
- Deep tendon reflexes normal
- No pathological reflexes (Babinski negative)

Potential Abnormalities:

- **Cognitive testing:** Impaired serial 7s, word recall, attention tasks
- **Tandem gait or Romberg:** May reveal subtle ataxia or balance impairment
- **Sustained muscle testing:** Rapid fatigability (e.g., handgrip dynamometer shows dramatic decline with repeated testing)
- **Sensory testing:** Hyperalgesia or allodynia in some patients (small fiber neuropathy)

Clinical Significance: The paucity of objective findings on standard neurological examination *despite severe symptoms* is characteristic of ME/CFS. This discordance (severe functional impairment with normal gross exam) historically led to dismissal of ME/CFS as “psychosomatic,” but advanced imaging and functional testing reveal objective abnormalities (cerebral blood flow reduction, autonomic dysfunction, immune activation).

Tender Point Assessment

Observation 14 (Fibromyalgia Overlap). 30–70% of ME/CFS patients meet criteria for fibromyalgia (widespread pain with tender points). Assessment:

- Digital palpation of 18 tender point sites with 4 kg pressure
- Fibromyalgia: ≥ 11 of 18 sites tender
- Presence of fibromyalgia does not exclude ME/CFS; these are frequently comorbid
- Guides pain management strategy

4.6.3 Laboratory Testing

Mandatory Exclusionary Testing

• Requirement 9: Minimum Laboratory Workup

The following tests are required to rule out alternative diagnoses:

Hematology:

- **Complete Blood Count (CBC):** Rule out anemia, leukemia, lymphoma
- If anemia present: Iron studies, B12, folate

Chemistry:

- **Comprehensive Metabolic Panel (CMP):** Rule out renal failure, hepatic dysfunction, electrolyte disorders, diabetes

Endocrine:

- **Thyroid function:** TSH, free T4 (hypothyroidism is a common mimic)
- Consider: Morning cortisol, ACTH stimulation test if Addison disease suspected

Inflammation:

- **Erythrocyte Sedimentation Rate (ESR) or C-Reactive Protein (CRP):** Rule out active inflammatory disease
- **Note:** ESR/CRP are typically *normal* in ME/CFS, distinguishing it from autoimmune diseases

Autoimmune Screening:

- **Antinuclear Antibody (ANA):** Screen for lupus, Sjögren syndrome, other connective tissue diseases
- If ANA positive: Reflex to specific antibodies (anti-dsDNA, anti-Ro, anti-La)

Vitamins:

- **Vitamin D:** Deficiency is extremely common and contributes to fatigue
- **Vitamin B12:** Deficiency causes fatigue and cognitive impairment

Sleep Disorders:

- **Polysomnography:** Rule out obstructive sleep apnea (OSA) or upper airway resistance syndrome (UARS)
- OSA can fully mimic ME/CFS; CPAP treatment produces dramatic improvement in true OSA
- OSA and ME/CFS can coexist; treating comorbid OSA improves but does not cure ME/CFS

Observation 15 (Typical Laboratory Profile in ME/CFS). The characteristic laboratory finding in ME/CFS is that **standard tests are normal**:

- CBC: Normal (no anemia, normal WBC count)
- CMP: Normal (normal kidney, liver, electrolytes)
- TSH: Normal
- ESR/CRP: Normal or low-normal (distinguishes from inflammatory autoimmune diseases)
- ANA: Usually negative (or low-titer positive without clinical significance)

This pattern—severe functional disability with normal routine labs—is diagnostically significant. It distinguishes ME/CFS from conditions that present with obvious laboratory abnormalities.

Advanced Biomarker Testing (If Available)

Observation 16 (Emerging Biomarkers for Biological Phenotyping). If resources permit, advanced testing can guide treatment:

Autoimmune Domain:

- GPCR autoantibody panel (β_2 -adrenergic, M3/M4 muscarinic)
- NK cell count and cytotoxicity assay
- Flow cytometry for plasma cell populations (CD38⁺CD138⁺)

Metabolic Domain:

- Heng 7-biomarker panel (AMP, ADP, VWF, fibronectin, TSP-1, PDGF-BB, TGF- β 3) when commercially available
- Fasting lactate (elevated suggests mitochondrial dysfunction)
- ATP profile (specialized labs only)

Objective Functional Testing:

- **Two-day cardiopulmonary exercise testing (CPET):** Gold standard for documenting PEM
- Day 1 vs. Day 2 comparison shows failure to reproduce work capacity
- Reduction in VO_2max , ventilatory threshold on Day 2 is diagnostic

Autonomic Testing:

- Formal tilt table testing (if orthostatic symptoms prominent)
- Heart rate variability analysis
- Quantitative sudomotor axon reflex test (QSART)

These tests are *not required for diagnosis* but enable Tier 2 biological phenotyping and treatment stratification.

4.7 Novel Biology-Informed Diagnostic Framework

This section proposes an updated diagnostic framework that synthesizes current pathophysiological understanding with clinical reality. Unlike existing criteria that treat ME/CFS as a single homogeneous entity, this three-tiered approach recognizes disease heterogeneity while maintaining diagnostic precision.

4.7.1 Rationale for a New Framework

The Logic Chain: From Biology to Diagnosis

The three-tiered diagnostic framework follows logically from four fundamental observations about ME/CFS:

Observation 1: ME/CFS is a Clinical Syndrome with a Core Pathognomonic Feature

Post-exertional malaise (PEM) with delayed onset, disproportionate severity, and prolonged recovery distinguishes ME/CFS from all other fatiguing conditions. This symptom:

- Cannot be explained by deconditioning (which improves with gradual activity)
- Cannot be explained by depression (which may improve somewhat with activity)
- Has objective correlates (2-day CPET showing Day 2 deterioration [79, 1])
- Reflects underlying cellular energy failure (ATP depletion [12, 64])

Logical consequence: Tier 1 must retain syndrome-based diagnosis with PEM as mandatory criterion, ensuring we capture the defining pathophysiology while maintaining compatibility with existing frameworks.

Observation 2: ME/CFS Heterogeneity Reflects Multiple Causal Pathways, Not Measurement Error The failure of single-target treatments in randomized controlled trials does not mean “ME/CFS has no biological basis”—it means we are mixing biologically distinct subgroups:

- Rituximab (anti-CD20 B cell depletion) failed in large trials [117] despite initial promise
- Daratumumab (anti-CD38 plasma cell depletion) succeeded in 60% of patients in pilot study [9]
- **Interpretation:** Not “autoimmunity isn’t involved,” but rather “wrong cell type targeted” (short-lived B cells vs. long-lived plasma cells) AND “only a subset has autoimmune-driven disease”

The Heng 2025 study [12] demonstrated that a 7-biomarker panel spanning three systems (energy, immune, vascular) achieved 91% diagnostic accuracy. This implies:

1. All three systems are coordinately dysfunctional (not independent)
2. ME/CFS is not five separate diseases but one syndrome with five co-occurring mechanisms
3. Treatment must address multiple domains simultaneously

Logical consequence: Tier 2 must assess all relevant biological domains and document which are present, rather than forcing patients into exclusive categories. The question is not “Is this autoimmune OR metabolic ME/CFS?” but “Which of the five domains show dysfunction in this patient?”

Observation 3: Treatment Response Depends on Rate-Limiting Steps, Not Just Presence of Pathology Consider two patients, both with elevated GPCR autoantibodies and mitochondrial dysfunction:

- **Patient A:** Autoantibodies are driving ongoing inflammation → mitochondria are secondarily impaired → removing autoantibodies allows mitochondrial recovery → daratumumab produces dramatic improvement
- **Patient B:** Initial autoimmune trigger has resolved, but mitochondrial damage is now self-perpetuating (WASF3 accumulation, cristae disruption) → removing residual autoantibodies doesn’t help because mitochondria cannot recover → daratumumab fails

Both patients are “autoantibody-positive,” but only Patient A responds. The difference: which domain is **rate-limiting** (the bottleneck preventing recovery).

This explains:

- Why daratumumab works in 60% not 100% of patients [9]
- Why low baseline NK cell count predicted non-response (suggests irreversible immune exhaustion)
- Why biomarker-positive patients don’t uniformly respond to biomarker-targeted treatments

Logical consequence: Tier 2 must enable multi-target treatment protocols. We cannot predict *a priori* which domain is rate-limiting, so we must:

1. Treat all accessible, low-risk domains simultaneously (“quick wins”)
2. Reassess at 3–6 months to identify which domains responded vs. persisted
3. Intensify treatment for persistent domains (these are likely rate-limiting)

Observation 4: ME/CFS Has Irreversible Thresholds, Making Timing Critical The natural history literature [70] and patient reports converge on temporal patterns:

- **6 months:** If symptoms persist beyond 6 months, spontaneous resolution becomes unlikely (transition from “post-viral fatigue” to “established ME/CFS”)
- **2 years:** Around 2 years, disease transitions from early (hypermetabolic, potentially reversible) to established (hypometabolic, epigenetically locked) state
- **Cumulative crashes:** Repeated PEM episodes cause progressive damage; there may be a threshold (5–10 severe crashes) beyond which recovery capacity is permanently impaired
- **25% severe:** One-quarter of ME/CFS patients become housebound/bedbound, most starting with mild disease

The progression from mild to severe appears **preventable in many cases** through aggressive pacing, yet existing diagnostic criteria provide no framework for:

- Identifying patients at high risk of progression
- Defining what “aggressive pacing” means operationally
- Communicating urgency of intervention before crossing irreversible thresholds

Logical consequence: Tier 3 must prospectively assess progression risk and provide actionable intervention protocols. The diagnostic framework must be **dynamic** (tracking trajectory) not static (labeling current state).

Why Three Tiers? Why Not Two or Four?

The three-tiered structure reflects three distinct clinical questions:

1. **Tier 1 (Syndrome):** Does this patient have ME/CFS? (Yes/No based on universal clinical features)
2. **Tier 2 (Biology):** Which pathophysiological mechanisms are driving this patient’s disease? (Multi-label classification across 5 domains)
3. **Tier 3 (Trajectory):** How severe is the disease currently, and what is the risk of irreversible progression? (Severity + prospective risk)

These cannot be collapsed:

- Tier 1 alone (current criteria) misses treatment stratification and progression prevention

- Tier 2 alone (biology-only) would miss patients without access to biomarkers and wouldn't address progression risk
- Tier 3 alone (severity-only) would lack diagnostic specificity and treatment guidance

Each tier serves a distinct purpose and requires different information.

Limitations of Existing Criteria

Existing diagnostic criteria (Fukuda, Canadian Consensus, ICC, IOM) share important limitations that this framework addresses:

- **Syndrome-based only:** Rely exclusively on symptom constellations without biological stratification, preventing precision medicine
- **Static classification:** Diagnose a point-in-time state without assessing progression risk, missing the 25% who will become severe
- **Assume homogeneity:** Force heterogeneous patients into single diagnostic category, explaining why single-target trials fail
- **Limited treatment guidance:** Diagnosis doesn't inform which interventions to prioritize, leading to trial-and-error
- **Miss therapeutic windows:** Fail to identify the 6-month and 2-year critical intervention windows

How Recent Advances Enable This Framework

The proposed three-tiered framework would not have been possible a decade ago. Recent advances now make it feasible:

- **Objective biomarkers:** GPCR autoantibodies [71, 72], Heng 7-marker panel [12], 2-day CPET [79, 1] provide biological stratification
- **Mechanistic understanding:** Autoimmunity [9], mitochondrial dysfunction [2, 64], neuroinflammation [73] explain heterogeneity
- **Treatment stratification proof-of-concept:** Daratumumab 60% response in autoimmune subset [9], immunoadsorption for GPCR autoantibodies [118] demonstrate that biomarker-guided treatment works
- **Natural history data:** Critical intervention windows [70], progression patterns [74], cumulative damage model validated

The proposed framework integrates these advances into clinically actionable diagnostic tiers.

Summary: The Logical Structure

Observation 17 (Framework Logic). **Premise 1:** ME/CFS is a clinical syndrome with a pathognomonic feature (PEM) that has objective correlates

⇒ **Tier 1:** Syndrome-based diagnosis with PEM mandatory

Premise 2: ME/CFS heterogeneity reflects multiple co-occurring biological mechanisms; treatment response depends on which mechanism is rate-limiting

⇒ **Tier 2:** Multi-domain biological phenotyping to enable multi-target treatment

Premise 3: ME/CFS has irreversible thresholds (6 months, 2 years, cumulative crashes); progression to severe disease is often preventable

⇒ **Tier 3:** Severity classification + prospective risk assessment with emergency protocols

The three-tiered structure is not arbitrary—it reflects the logical necessity of answering three distinct clinical questions (diagnosis, mechanism, trajectory) that cannot be collapsed without losing critical information.

4.7.2 Tier 1: Clinical Syndrome Criteria

Tier 1 establishes the diagnosis of ME/CFS based on clinical features. These criteria are universal—all patients must meet Tier 1 to receive the diagnosis.

Core Diagnostic Features (All Required)

• Requirement 10: Post-Exertional Malaise (Mandatory Hallmark)

Post-exertional malaise must be present with ALL of the following characteristics:

- **Delayed onset:** Symptom exacerbation occurs 12–72 hours after triggering activity (not immediately)
- **Disproportionate severity:** Minimal exertion produces profound symptom worsening far beyond normal fatigue
- **Multi-domain triggers:** Symptoms triggered by physical exertion *AND* cognitive exertion *AND* emotional exertion
- **Prolonged recovery:** Symptom exacerbation persists >24 hours (mild cases) to weeks or months (severe cases)

Objective verification (optional but supportive): Two-day cardiopulmonary exercise testing showing Day 2 deterioration: workload at ventilatory threshold decreases ≥20% on Day 2 compared to Day 1 [79, 1].

• Requirement 11: Baseline Energy Insufficiency

Patients must demonstrate chronic energy deficit characterized by:

- **Morning depletion:** Waking already exhausted despite sleep duration
- **Disproportionate activity cost:** Activities of daily living (hygiene, eating, sitting upright) consume excessive energy relative to effort
- **No functional reserve:** Zero capacity to handle unexpected physical, cognitive, or emotional demands
- **Effort-performance disconnect:** Subjective experience of maximal effort producing minimal objective output (a phenomenon that distinguishes ME/CFS from deconditioning or primary depression)

The effort-performance disconnect represents a novel diagnostic criterion capturing the lived experience of ME/CFS: patients describe “giving everything” to accomplish minimal tasks, feeling as though simple activities require marathon-level exertion while producing negligible results [55, 56].

• Requirement 12: Duration and Exclusion Criteria

- **Duration:** Symptoms must persist ≥ 6 months
- **Rationale:** Six-month persistence indicates transition from post-viral fatigue (which typically resolves) to established ME/CFS with aberrant pathophysiology [70]
- **Exclusions:** Symptoms not better explained by:
 - Active medical conditions (untreated hypothyroidism, sleep apnea, anemia, malignancy)
 - Primary psychiatric disorders (though secondary depression/anxiety are common and do not exclude ME/CFS)
 - Medication side effects

Important: Comorbid conditions that are part of the ME/CFS disease spectrum (POTS, fibromyalgia, MCAS, IBS) do *not* exclude the diagnosis—these represent overlapping pathophysiology rather than alternative explanations.

Supporting Features (≥ 3 of 5 Required)

In addition to the three core features, patients must have at least three of the following five supporting features:

1. Unrefreshing Sleep

- Sleep that fails to restore energy regardless of duration
- Waking feeling as exhausted as when going to bed
- Present in 95–100% of ME/CFS patients [80, 81]

2. Cognitive Impairment

- Processing speed deficits (most robust finding: Hedges' $g = -0.82$) [89]

- Attention and working memory impairment
- Word-finding difficulties, linguistic reversals
- Brain fog that is not attributable to fatigue or depression [91]

3. Autonomic Dysfunction

- Orthostatic intolerance: symptoms worsened by upright posture
- POTS (heart rate increase ≥ 30 bpm upon standing), orthostatic hypotension, or neurally mediated hypotension
- Temperature dysregulation, inappropriate sweating or lack of sweating
- Present in 70–90% of ME/CFS patients [97]

4. Pain

- Myalgia (muscle pain), particularly with post-exertional exacerbation
- Arthralgia (joint pain, characteristically migratory without inflammation)
- Headaches (migraine or tension-type) [103]
- Pain present in ~80% of patients [100]

5. Sensory Hypersensitivity

- Photophobia (light sensitivity requiring sunglasses indoors or dimmed environment)
- Phonophobia (sound sensitivity; normal volumes feel uncomfortable)
- Chemical sensitivity (fragrances, cleaning products, exhaust)
- Touch hypersensitivity or allodynia
- Present in 70–90% of patients [111]

Observation 18 (Tier 1 Summary). Tier 1 criteria establish ME/CFS as a clinical syndrome with mandatory post-exertional malaise, baseline energy insufficiency, and 6-month duration. Supporting features (sleep, cognition, autonomic, pain, sensory) must be present in sufficient number (≥ 3 of 5) to confirm the characteristic multi-system presentation. These criteria are compatible with existing frameworks (Canadian Consensus, ICC, IOM) but add explicit recognition of the effort-performance disconnect and specify the 6-month threshold as marking transition to established disease.

4.7.3 Tier 2: Biological Phenotyping (Multi-Domain Assessment)

Once Tier 1 criteria are met, patients should undergo comprehensive biological phenotyping to identify which pathophysiological domains are involved. This enables targeted treatment and research stratification.

Rationale: Co-Occurrence Rather Than Predominance

Critical insight: ME/CFS patients typically have dysfunction in *multiple* biological domains simultaneously. The Heng 2025 study demonstrated that a 7-biomarker panel spanning energy metabolism, immune function, and vascular endothelium achieved 91% diagnostic accuracy

precisely because *all three systems show coordinated dysfunction* [12]. This finding validates the multi-lock model (Chapter 14): ME/CFS persists because multiple self-reinforcing pathophysiological processes operate concurrently.

~ Hypothesis 2: Multi-Domain Co-Occurrence Model

ME/CFS should be understood as a syndrome with five co-occurring, mutually reinforcing biological domains. Most patients have abnormalities in ≥ 3 domains:

- Autoimmune features: 30–60% (GPCR autoantibodies [71, 72])
- Mitochondrial/metabolic dysfunction: 70–95% (ATP abnormalities [12], lactate elevation [65])
- Neuroinflammation/central sensitization: 70–90% (central sensitization 84% [106], sensory sensitivities 70–90% [111])
- Dysautonomia: 70–90% (POTS 25–50%, broader orthostatic intolerance 70–90% [97])
- Endothelial dysfunction: Prevalence unknown (Heng 2025 documented elevation in ME/CFS cohort [12])

These domains are interdependent:

- Autoimmunity (GPCR autoantibodies) → Mitochondrial dysfunction (β_2 -adrenergic signaling regulates mitochondrial biogenesis)
- Mitochondrial dysfunction (ATP depletion) → Neuroinflammation (danger signal release, ionic gradient failure)
- Endotheliopathy (impaired vasodilation) → Dysautonomia (orthostatic intolerance, cerebral hypoperfusion)
- Neuroinflammation (cytokine production) → Autoimmunity (B cell activation)

Treatment targeting a single domain may fail because untreated domains maintain dysfunction. The multi-domain model predicts that:

1. Patients with more domains affected will have worse outcomes
2. Multi-target interventions will outperform single-target interventions
3. Treatment response requires both (a) presence of dysfunction in a domain AND (b) that domain being rate-limiting (the bottleneck gating recovery)

Domain 1: Autoimmune Features

Assessment:

- GPCR autoantibodies (β_2 -adrenergic, M3 muscarinic, M4 muscarinic) above age/sex-matched reference ranges [71, 72]
- ANA (any titer; present in 20–30% ME/CFS vs. 5–10% healthy controls)
- Plasma cell expansion on flow cytometry (CD38⁺CD138⁺ if available)
- Low NK cell count (<5th percentile) with normal total lymphocytes

If Present → Diagnosis: “ME/CFS with Autoimmune Component”

Treatment Implications:

- Candidate for immunoadsorption (IgG removal) [118]
- Candidate for daratumumab (anti-CD38, depletes plasma cells) [9]
- Candidate for BC007 (GPCR autoantibody neutralizer) [10]
- Monitor for worsening with immune-stimulating interventions

Prevalence: 30–60% of ME/CFS patients

Domain 2: Mitochondrial/Metabolic Dysfunction

Assessment:

- Heng 7-marker panel (if available): Elevated AMP, ADP (energy depletion arm) [12]
- Elevated lactate: Resting >2.0 mmol/L or abnormal accumulation during 2-day CPET [65]
- ATP profile abnormalities (if specialized testing available)
- WASF3 elevation on skeletal muscle biopsy (if indicated for severe cases) [2]

If Present → Diagnosis: “ME/CFS with Mitochondrial Dysfunction”

Treatment Implications:

- CoQ10 (ubiquinol 200–400 mg/day)
- NAD⁺ precursors (nicotinamide riboside 1000–2000 mg/day, treatment duration ≥10 weeks)
- D-ribose, B-complex vitamins, alpha-lipoic acid, PQQ
- Strict pacing critical (ATP depletion is cumulative)
- Heart rate monitoring (stay below 60% maximum heart rate during activity)

Prevalence: 70–95% (virtually all ME/CFS patients show some degree of energy metabolism dysfunction)

Domain 3: Neuroinflammation/Central Sensitization

Assessment:

- **Research settings:** PET evidence of microglial activation [73], fMRI showing altered temporoparietal junction or salience network connectivity [5, 94]
- **Clinically accessible:**

- Central sensitization confirmed by quantitative sensory testing: pressure pain thresholds <5th percentile at ≥ 3 standardized sites [106]
- Small fiber neuropathy: skin biopsy showing intraepidermal nerve fiber density <5th percentile [104]
- Severe sensory sensitivities requiring environmental modification (inability to tolerate normal lighting, sound levels, or chemical exposures)

If Present → Diagnosis: “ME/CFS with Neuroinflammatory Component”

Treatment Implications:

- Low-dose naltrexone (LDN 1.5–4.5 mg at bedtime)
- Environmental modification (dimmed lighting, noise reduction, fragrance-free environment)
- IVIG (if small fiber neuropathy documented and insurance approves)
- Avoid activities that trigger sensory overload (cognitive post-exertional malaise)

Prevalence: 70–90% (sensory sensitivities 70–90%, central sensitization 84%, small fiber neuropathy 30–38%)

Domain 4: Dysautonomia

Assessment:

- **Gold standard:** Tilt table testing showing POTS (heart rate increase ≥ 30 bpm within 10 minutes), orthostatic hypotension (blood pressure drop $\geq 20/10$ mmHg), or neurally mediated hypotension
- **Clinically accessible:** NASA Lean Test (10-minute standing test; positive if heart rate increases ≥ 30 bpm)
- Heart rate variability analysis (reduced HRV indicating sympathetic dominance)
- QSART/thermoregulatory sweat test (if available)

If Present → Diagnosis: “ME/CFS with Dysautonomia”

Treatment Implications:

- Volume expansion: 3–10 g sodium + 2–3 L fluids daily
- Compression garments (20–30 mmHg waist-high or thigh-high)
- Pharmacological:
 - Fludrocortisone 0.05–0.2 mg daily
 - Midodrine 2.5–10 mg three times daily

- Ivabradine 2.5–7.5 mg twice daily
- Low-dose beta-blockers (propranolol 10–20 mg as needed)
- Positional strategies: elevate head of bed, avoid prolonged standing, sit when possible

Prevalence: 70–90% (POTS 25–50%, broader orthostatic intolerance 70–90%)

Domain 5: Endothelial Dysfunction

Assessment:

- Heng 7-marker panel (if available): Elevated von Willebrand factor, fibronectin, thrombospondin-1 (endothelial activation arm) [12]
- Clinical markers of microvascular dysfunction:
 - Livedo reticularis (mottled skin discoloration)
 - Raynaud’s phenomenon (cold-induced color changes in fingers/toes)
 - Delayed capillary refill (>3 seconds)
- Cerebral hypoperfusion on SPECT imaging (if available)

If Present → Diagnosis: “ME/CFS with Endothelial Dysfunction”

Treatment Implications (experimental):

- L-citrulline 3–6 g/day or L-arginine (for nitric oxide production)
- Omega-3 fatty acids (EPA/DHA 2–4 g/day)
- Low-dose aspirin 81 mg daily (if no contraindications)
- Emerging research: anticoagulation trials, fibrinolytic protocols (investigational only)

Prevalence: Unknown (Heng 2025 documented elevation in ME/CFS cohort; prevalence in broader ME/CFS population requires validation)

Hypothetical Phenotype Under Investigation: Viral-Immune-Metabolic Type

△ Warning 6: Speculative Phenotype Cluster

The “Viral-Immune-Metabolic” phenotype described below is **hypothetical and unvalidated**. It is based on:

- Single-case clinical observation extrapolation
- Mechanistic reasoning without controlled validation
- Limited anecdotal reports of cimetidine response

This phenotype concept requires rigorous validation in prospective cohort studies before clinical adoption. It should NOT be used for diagnosis or treatment selection outside research protocols.

Clinical observation in isolated cases has suggested a potential subset of patients whose symptom pattern and treatment response might indicate a specific pathophysiological cluster crossing multiple domains. This hypothetical phenotype is presented for research discussion only.

Proposed Phenotype Definition. The “Viral-Immune-Metabolic” cluster (see Section 5.6.12 for detailed discussion) is characterized by:

- Post-infectious onset (EBV, HHV-6, or other herpesvirus)
- MCAS or histamine intolerance (HIT) comorbidity
- POTS/dysautonomia
- Strong response to amino acid supplementation
- Dramatic response to cimetidine (rare but distinctive)

Table 4.2: Tier 2 test panels for Viral-Immune-Metabolic phenotype

Domain	Priority 1 Tests	Priority 2 Tests	Phenotype Indicator
Viral-Immune	EBV serology (VCA Ig-G/IgM, EBNA, EA-D), HHV-6 serology	HHV-6/EBV PCR, T cell exhaustion panel (PD-1, LAG-3)	Cimetidine response
Metabolic	Amino acid panel (serum), D-Ribose therapeutic trial	Lactate/pyruvate ratio, organic acids (urine)	Amino acid response
Mast Cell	Serum tryptase, plasma histamine, DAO activity	24h urine N-methylhistamine, prostaglandin D ₂	HIT/MCAS symptoms
Dysautonomia	Tilt test, lying/standing catecholamines	HRV analysis, QSART	POTS + MCAS coupling
Endothelial	Flow-mediated dilation, ADMA	L-Arginine/L-Citrulline therapeutic response	NO dysfunction

Targeted Test Panel for Suspected Viral-Immune-Metabolic Cluster.

Clinical Decision Tree. If this phenotype is suspected:

1. **Cimetidine shows benefit** → Prioritize viral serology (EBV/HHV-6) + amino acid panel
2. **Strong amino acid response** → Prioritize intestinal permeability markers (Zonulin, LPS) + mitochondrial assessment
3. **POTS + MCAS both present** → Prioritize dysautonomia subtyping (hyperadrenergic vs. neuropathic)

Treatment Approach (Hypothetical).

△ **Warning 7: Medical Supervision Required**

This treatment protocol requires physician supervision and cannot be safely self-implemented:

- **Drug interaction monitoring:** Cimetidine is a CYP450 inhibitor affecting metabolism of many medications (warfarin, phenytoin, theophylline, benzodiazepines, beta-blockers, calcium channel blockers)
- **Renal function assessment:** Valacyclovir requires dose adjustment in renal impairment; accumulation can cause neurotoxicity
- **Adverse effect monitoring:** CNS effects (cimetidine), nephrotoxicity (antivirals), hypotension (histamine blockade)
- **Laboratory monitoring:** Renal function (creatinine, eGFR) before and during antiviral therapy; liver function tests; complete blood count

Do not attempt this protocol without medical consultation and ongoing physician oversight.

For confirmed Viral-Immune-Metabolic cluster:

1. **Foundation:** Pacing + H1/H2 dual blockade (mast cell stabilization)
2. **Metabolic support:** NAC 1800 mg/day, L-citrulline-malate 6–8 g/day, D-ribose 5 g TID
3. **If viral titers elevated:** Consider antiviral trial (valacyclovir for EBV, valganciclovir for CMV/HHV-6)
4. **If cimetidine response:** Continue cimetidine 200–400 mg BID (immunomodulation benefit)
5. **Reassess:** 3-month follow-up with repeat amino acid panel and symptom tracking

△ **Warning 8: Evidence Limitations**

This phenotype cluster is based on clinical observation and mechanistic reasoning, not validated clinical trials. The specific test panel and treatment sequence are proposed for systematic evaluation, not established standard of care. See Section 5.6.12 for detailed certainty assessment.

Multi-Label Diagnosis Example

Observation 19 (Sample Comprehensive Diagnosis). **Primary Diagnosis:** Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS)

Biological Phenotype (Tier 2 Multi-Domain Assessment):

- ✓ Autoimmune component: GPCR autoantibodies β_2 -adrenergic 12.5 U/mL (ref <8), M3 muscarinic 9.2 U/mL (ref <7)
- ✓ Mitochondrial dysfunction: Fasting lactate 2.8 mmol/L (ref <2.0); 2-day CPET workload at VT decreased 35% on Day 2
- ✓ Neuroinflammatory component: Pressure pain thresholds 1.8 kg (ref >4.0); photophobia requiring indoor sunglasses; phonophobia limiting social interaction
- ✓ Dysautonomia: POTS confirmed on tilt table (supine HR 65 bpm → standing HR 102 bpm at 8 minutes); HRV SDNN 18 ms (ref >50)
- × Endothelial dysfunction: Not assessed (markers unavailable)

Severity (Tier 3): Moderate (housebound 40% of time; can perform remote work 20 hours/week with careful pacing)

Progression Risk: HIGH—RED FLAGS present: ratcheting baseline over past 9 months (each crash leaves lower functional floor); recovery time lengthening from 3 days → 10 days for equivalent exertion

Treatment Plan:

1. **Foundation:** Aggressive pacing (50% rule, heart rate monitoring <105 bpm)
2. **Dysautonomia** (quick win, high accessibility): Fludrocortisone 0.1 mg daily, sodium 6 g/day, fluids 2.5 L/day, compression stockings
3. **Mitochondrial** (quick win, high accessibility): CoQ10 300 mg, nicotinamide riboside 1000 mg, B-complex
4. **Neuroinflammation** (quick win, high accessibility): Low-Dose Naltrexone (LDN) 3 mg at bedtime, light/sound environmental control
5. **Autoimmune** (if accessible): Immunoadsorption or daratumumab candidate; pursue if no improvement after 6 months on above protocol

Reassessment: 3-month follow-up to evaluate response in each domain; adjust treatment based on which domains improve vs. persist

4.7.4 Tier 3: Severity Classification and Progression Risk

Tier 3 classifies current functional severity and prospectively assesses risk of progression to severe disease. This enables appropriate resource allocation, guides intervention intensity, and identifies patients requiring emergency intervention.

Functional Severity Scale

1. **Mild ME/CFS**
 - Can work or study at 50–80% of pre-illness capacity, though with significant difficulty
 - Post-exertional malaise occurs after moderate exertion

- Recovery from PEM takes days to 1–2 weeks
- Can perform most activities of daily living independently
- Energy envelope is reduced but allows meaningful activity
- Appears functional to outside observers (“invisible illness”)

2. Moderate ME/CFS

- Reduced daily activity to <50% of pre-illness level
- Housebound 50% or more of the time
- Unable to work or study full-time; may work part-time with difficulty
- Post-exertional malaise triggered by minimal exertion
- Recovery from PEM takes weeks
- Requires extended rest periods daily
- Significant impairment in social and occupational function

3. Severe ME/CFS

- Mostly bedbound (>50% of waking hours)
- Can perform only minimal self-care activities (brief washing, feeding)
- Post-exertional malaise triggered by activities of daily living
- Cognitive impairment prevents reading, sustained conversation
- Sensory sensitivities may require dimmed environment, minimal sound
- Unable to leave home except for essential medical appointments
- Requires assistance with instrumental activities of daily living

4. Very Severe ME/CFS

- Bedbound continuously
- Unable to perform most self-care activities without assistance
- Profound sensitivity to light (requiring darkness), sound (requiring silence), touch
- May be unable to tolerate speaking or being spoken to
- Tube feeding may be required if swallowing is impaired
- Requires full-time care assistance
- Represents approximately 10% of severe ME/CFS cases (2–3% of total ME/CFS population)

Progression Risk Stratification

△ Warning 9: HIGH RISK for Progression to Severe Disease

Patients meeting ≥2 of the following RED FLAG criteria are at immediate risk of transitioning to severe, potentially irreversible disease and require emergency intervention:

RED FLAGS (Immediate Danger):

1. **Ratcheting baseline:** Each post-exertional crash leaves patient at a lower functional

floor; baseline is trending downward over 6–12 months rather than returning to previous level

2. **Recovery time lengthening:** PEM recovery now requires >2 weeks (previously required only days to 1 week)
3. **Shrinking energy envelope:** Activities that were safely within the energy envelope 6 months ago now trigger post-exertional malaise
4. **New sensory sensitivities:** Photophobia, phonophobia, or chemical sensitivities emerging or rapidly worsening
5. **Cognitive decline:** Word-finding difficulties, memory impairment, or inability to read/process information worsening (cognitive symptoms are most resistant to recovery) [74]
6. **Forced overexertion:** Patient cannot stop working or reduce activity due to financial necessity, despite clear evidence of deterioration (structural inability to pace)
7. **Weight loss from energy insufficiency:** Eating and food preparation have become too effortful; weight loss indicates severe energy depletion
8. **Social withdrawal by necessity:** Cannot tolerate visitors, phone calls, or any social interaction due to symptom exacerbation (not due to depression)

Emergency Action Protocol: Patients with HIGH RISK status require immediate intervention to prevent crossing the “point of no return” to irreversible severe ME/CFS:

1. **Within 48 hours:**
 - Reduce all non-essential activity by 50%
 - Implement aggressive horizontal rest (50–75% of waking hours)
 - Cancel social commitments, request emergency work accommodation
2. **Within 1 week:**
 - Physician visit for medical leave documentation
 - Formal workplace accommodation request (reduced hours 50–75%, remote work, flexible schedule)
 - Begin disability application process if accommodations denied or insufficient
3. **Within 4–8 weeks:**
 - Achieve baseline stabilization: Goal of ZERO post-exertional malaise episodes for 4 continuous weeks
 - This proves patient is within energy envelope
 - Accept that functional capacity is very low during this period—this is temporary to prevent permanent severe disease

Rationale: Research and patient reports demonstrate that repeated post-exertional malaise episodes cause cumulative physiological damage: mitochondrial dysfunction accumulation [2, 64], endothelial dysfunction [12], neuroinflammation [73], and immune exhaustion [119]. There appears to be a threshold (anecdotally 5–10 severe crashes)

beyond which recovery capacity is permanently impaired. The goal is to avoid severe crashes entirely, not merely to minimize them.

Critical Temporal Windows

Observation 20 (The 6-Month Rule and 2-Year Establishment Threshold). Two temporal thresholds mark critical transitions in ME/CFS natural history [70]:

6-Month Persistence Mark: If symptoms persist beyond 6 months without improvement, this indicates that normal homeostatic recovery mechanisms have failed and aberrant pathophysiology is becoming established. This marks the transition from “post-viral fatigue that might spontaneously resolve” to “ME/CFS requiring active intervention.”

2-Year Establishment Threshold: Around 2 years post-onset, ME/CFS transitions from early disease (hypermetabolic, potentially modifiable) to established disease (hypometabolic, potentially entrenched). This transition involves:

- Epigenetic changes altering gene expression patterns
- Immune exhaustion (CD8⁺ T cell exhaustion [119], NK cell dysfunction)
- Normalization of inflammatory markers despite ongoing dysfunction
- Brain structural changes visible on advanced imaging [94]
- Metabolic state shift from high (inefficient) energy expenditure to low energy production

Implication: The first 2 years represent a critical intervention window. Aggressive pacing, early biological phenotyping, and domain-targeted treatment during this period may prevent progression to established severe disease. After 2 years, reversal becomes substantially more difficult (though not impossible).

Clinical application: Patients diagnosed within 6 months of onset should be counseled on the criticality of aggressive pacing to prevent establishment. Patients approaching the 2-year mark should undergo comprehensive Tier 2 phenotyping to guide maximal intervention before the window closes.

4.7.5 Implementation and Clinical Workflow

Minimum Diagnostic Workup (All Patients)

1. Tier 1 Clinical Assessment:

- Detailed history: onset pattern, post-exertional malaise characteristics, sleep quality, cognitive symptoms, autonomic symptoms, pain, sensory sensitivities
- Physical examination: orthostatic vital signs, neurological examination, tender point assessment
- Functional capacity assessment: Bell Disability Scale, SF-36, or equivalent

2. Objective Testing (if accessible):

- Two-day cardiopulmonary exercise testing (gold standard for PEM documentation)
- Tilt table testing or NASA Lean Test (dysautonomia assessment)

3. Basic Laboratory Testing (rule out exclusions):

- Complete blood count (CBC)
- Comprehensive metabolic panel (CMP)
- Thyroid-stimulating hormone (TSH), free T4
- Ferritin (low ferritin contributes to fatigue and restless legs)
- Antinuclear antibody (ANA)
- Erythrocyte sedimentation rate (ESR), C-reactive protein (CRP)
- Vitamin D, vitamin B12

4. Sleep Study:

- Polysomnography to rule out obstructive sleep apnea (OSA) or upper airway resistance syndrome (UARS)
- OSA can mimic ME/CFS; treatment with CPAP dramatically improves symptoms in true OSA cases
- OSA and ME/CFS can coexist; treating comorbid OSA improves but does not cure ME/CFS

Advanced Phenotyping (Tier 2, If Resources Permit)

1. Autoimmune Domain:

- GPCR autoantibody panel (β_2 -adrenergic, M3 muscarinic, M4 muscarinic)
- NK cell count and function assay
- Flow cytometry for plasma cell populations (CD38⁺CD138⁺)

2. Mitochondrial/Metabolic Domain:

- Heng 7-biomarker panel (when commercially available): AMP, ADP, VWF, fibronectin, thrombospondin-1, PDGF-BB, TGF- β 3
- Fasting lactate
- ATP profile (if specialized laboratory available)

3. Neuroinflammation Domain:

- Quantitative sensory testing (pressure pain thresholds)
- Skin biopsy for small fiber neuropathy (intraepidermal nerve fiber density)

4. Dysautonomia Domain:

- Tilt table testing (if not already performed)
- Heart rate variability analysis
- QSART or thermoregulatory sweat test (if available)

5. Comorbidity Screening (Septad components):

- MCAS workup: serum tryptase, 24-hour urine methylhistamine, prostaglandin D₂
- Hypermobility assessment: Beighton score

- If hEDS + progressive neurological symptoms: upright MRI for craniocervical instability screening
- Gastrointestinal: gastric emptying study, SIBO breath test (if prominent GI symptoms)

Treatment Prioritization Based on Phenotype

Table 4.3: Treatment prioritization by biological domain

Domain	Treatment Options	Risk Level	Access	Priority
Pacing	Activity management, heart rate monitoring	None	High	FIRST (always)
Dysautonomia	Salt, fluids, compression, fludrocortisone, midodrine	Low	High	SECOND (quick wins)
Mitochondrial	CoQ10, NR/NMN, B vitamins	Low	High	SECOND (quick wins)
Neuroinflam.	LDN, environmental modification	Low	High	SECOND (quick wins)
Autoimmune	Immunoadsorption, daratumumab, BC007	Moderate-High	Very Low	THIRD (if accessible)
Endothelial	L-citrulline, omega-3, aspirin	Low	High	THIRD (experimental)

Rationale:

- **Foundation:** Pacing is universal and non-negotiable—prevents cumulative damage regardless of biological phenotype
- **Quick wins:** High-accessibility, low-risk interventions (dysautonomia, mitochondrial, neuroinflammation) initiated simultaneously to address multiple domains
- **Reassessment:** At 3–6 months, evaluate response in each domain; persistent dysfunction despite accessible interventions justifies pursuit of high-intensity/low-accessibility treatments (immunoadsorption, daratumumab)
- **Multi-target approach:** Addresses multiple locks simultaneously, recognizing that single-domain interventions often fail due to reinforcement from untreated domains

Limitation 1: Novel Diagnostic Framework: Entirely Unvalidated

This three-tiered diagnostic framework is a proposal, not an established or validated diagnostic system. Critical epistemic boundaries:

- No prospective study has tested whether the five biological domains (neuroinflammation, autoimmune, mitochondrial, dysautonomia, vascular-endothelial) are clinically separable, whether their presence predicts treatment response, or whether multi-domain assessment improves outcomes over existing diagnostic criteria.
- The “rate-limiting domain” concept—that treatment response depends on which domain is the primary bottleneck—is a theoretical construct not yet supported by clinical data. The daratumumab responder/non-responder distinction is cited as supporting evidence, but the study was not designed to test this framework.
- Biomarker thresholds for domain positivity have not been established, validated, or replicated. Existing studies use different assays, populations, and cut-points, making standardisation premature.
- The framework’s reliance on the Heng 2025 7-biomarker panel [12] as validation of multi-system coordination rests on a single study that has not been independently replicated.

4.7.6 Research Implications and Validation Needs

This novel diagnostic framework generates testable predictions that should be validated in prospective studies:

1. **Hypothesis:** Patients with ≥ 4 domains positive will have worse functional outcomes, longer illness duration, and lower treatment response rates than patients with 1–2 domains
 - **Test:** Correlate number of positive domains with SF-36 Physical Function, Bell Disability Scale, work/school capacity, and hospitalization rates
2. **Hypothesis:** Multi-target interventions (treating all present domains) will produce superior outcomes compared to single-target interventions
 - **Test:** Randomized controlled trial comparing CoQ10 monotherapy vs. CoQ10 + Low-Dose Naltrexone (LDN) + fludrocortisone (in patients with mitochondrial + neuroinflammatory + dysautonomia domains positive)
3. **Hypothesis:** The RED FLAG progression risk criteria (Tier 3) prospectively identify patients who will develop severe ME/CFS
 - **Test:** Cohort study assessing RED FLAG status at enrollment, then tracking functional severity at 1 year and 2 years; calculate sensitivity/specificity of RED FLAG criteria for predicting progression to severe disease
4. **Hypothesis:** Treatment response to domain-specific interventions requires both (a) presence of dysfunction in that domain AND (b) that domain being rate-limiting (the bottleneck)

- **Test:** Measure all 5 domains → administer domain-specific treatment → identify responders vs. non-responders → retrospectively determine which baseline features predicted response
 - Example: Daratumumab trial measuring GPCR autoantibodies, lactate, HRV, QST, VWF at baseline, then analyzing which baseline profile predicts 60% responder group vs. 40% non-responder group
5. **Hypothesis:** The Heng 7-marker panel achieves high diagnostic accuracy because it captures coordinated dysfunction across three systems (energy, immune, vascular), and symptom severity correlates with multi-system burden rather than single-marker elevation
- **Test:** Network analysis or partial least squares regression to determine if symptoms correlate with individual markers or require multi-marker patterns
6. **Hypothesis:** Early intervention (within the first 2 years) prevents establishment of refractory disease
- **Test:** Compare outcomes of patients receiving comprehensive Tier 2 phenotyping + multi-target treatment within 1 year of onset vs. those diagnosed/treated after 2+ years
 - Ethical note: This should be observational (registry-based) rather than randomized, as withholding early treatment would be unethical if the hypothesis is correct

4.7.7 Comparison to Existing Criteria

Table 4.4 compares the novel biology-informed framework to established diagnostic criteria.

Observation 21 (Framework Compatibility). The novel framework is *compatible* with existing criteria rather than contradictory:

- Tier 1 clinical criteria align with Canadian Consensus and IOM requirements
- Post-exertional malaise remains the mandatory hallmark (consistent with ICC, Canadian, IOM)
- 6-month duration threshold maintained (all modern criteria)
- Tier 2 and Tier 3 represent *additions* that do not invalidate previous diagnoses

Patients meeting Fukuda, Canadian Consensus, ICC, or IOM criteria will meet Tier 1 of the novel framework. The novel framework adds biological stratification (Tier 2) and risk assessment (Tier 3) that can be applied retroactively to existing cohorts.

4.7.8 Clinical Advantages of the Novel Framework

1. **Precision medicine:** Biological phenotyping enables targeted treatment rather than trial-and-error
2. **Explains treatment heterogeneity:** Response variability attributed to different rate-limiting domains rather than “treatment doesn’t work”

Table 4.4: Comparison of diagnostic frameworks

Feature	Fukuda (1994)	Canadian (2003)	IOM (2015)	Novel Framework (2026)
PEM required	No	Yes	Yes	Yes (detailed criteria)
Duration	6 months	6 months	6 months	6 months (establishment threshold)
Biological phenotyping	No	No	No	Yes (5 domains)
Progression risk assessment	No	No	No	Yes (RED FLAGS)
Treatment stratification	No	No	No	Yes (domain-targeted)
Temporal windows	No	No	No	Yes (2-year critical window)
Recognizes heterogeneity	No	Partially	No	Yes (multi-label classification)
Objective biomarkers	No	Optional	Optional	Integrated (Tier 2)
Subgroup identification	No	No	No	Yes (co-occurrence model)

3. **Early intervention guidance:** 6-month and 2-year thresholds identify critical windows for aggressive treatment
4. **Progression prevention:** RED FLAG criteria enable emergency intervention before irreversible severe disease
5. **Research stratification:** Multi-domain classification allows trials to enrich for patients with specific phenotypes (e.g., daratumumab trial selecting autoimmune-domain-positive patients)
6. **Acknowledges complexity:** Multi-label classification reflects biological reality (most patients have 3+ domains) rather than forcing heterogeneous patients into single category
7. **Actionable at point of care:** Tier 1 (clinical) immediately implementable; Tier 2 (biological) scalable as biomarkers become commercially available; Tier 3 (risk) requires only clinical observation

4.8 Differential Diagnosis

ME/CFS is a diagnosis of exclusion, requiring careful evaluation to rule out other conditions that can present with similar symptoms. This section addresses conditions that can mimic ME/CFS, distinguishing features, and the critical distinction between alternative diagnoses and comorbidities.

4.8.1 Conditions That Can Mimic ME/CFS

Endocrine Disorders

• Requirement 13: Must Rule Out Before Diagnosing ME/CFS

Hypothyroidism:

- **Symptoms:** Fatigue, cognitive impairment (“brain fog”), cold intolerance, weight gain, constipation
- **Distinguishing features:** Gradual onset, no post-exertional malaise, responds to thyroid replacement
- **Testing:** TSH, free T4; if TSH elevated and free T4 low, hypothyroidism is confirmed
- **Note:** Subclinical hypothyroidism (mildly elevated TSH with normal T4) is controversial; may contribute to fatigue but is insufficient to explain ME/CFS severity

Addison Disease (Primary Adrenal Insufficiency):

- **Symptoms:** Profound fatigue, orthostatic hypotension, salt craving, hyperpigmentation
- **Distinguishing features:** Progressive worsening, life-threatening if untreated, responds to cortisol replacement
- **Testing:** Morning cortisol, ACTH stimulation test; electrolytes show hyponatremia

and hyperkalemia

Diabetes Mellitus:

- **Symptoms:** Fatigue, polyuria, polydipsia, weight loss
- **Testing:** Fasting glucose, HbA1c

Sleep Disorders

△ **Warning 10: Obstructive Sleep Apnea Can Fully Mimic ME/CFS**

Obstructive Sleep Apnea (OSA): OSA is a critical exclusion because it can produce a symptom profile nearly identical to ME/CFS:

- **Symptoms:** Profound fatigue, unrefreshing sleep, cognitive impairment, morning headaches
- **Distinguishing features:**
 - Snoring, witnessed apneas (ask bed partner)
 - Obesity (BMI >30) is common but not required
 - **Critical:** OSA patients do NOT have post-exertional malaise with delayed onset
 - Improvement with CPAP treatment (if true OSA, dramatic improvement within weeks)
- **Testing:** Polysomnography (sleep study); apnea-hypopnea index (AHI) ≥ 5 events/hour is diagnostic
- **Important:** OSA and ME/CFS can coexist; treating comorbid OSA improves but does not cure ME/CFS

Upper Airway Resistance Syndrome (UARS):

- Milder form of sleep-disordered breathing
- May have normal AHI but increased respiratory effort-related arousals (RERAs)
- Presents with fatigue and unrefreshing sleep similar to ME/CFS

Idiopathic Hypersomnia:

- Excessive daytime sleepiness despite adequate sleep duration
- No post-exertional malaise
- Multiple sleep latency test (MSLT) shows short sleep latency

Autoimmune and Inflammatory Diseases

Observation 22 (Inflammatory Markers Distinguish ME/CFS from Autoimmune Disease). The following autoimmune and inflammatory diseases share symptoms with ME/CFS but

can be distinguished by specific biomarkers and clinical features.

Systemic Lupus Erythematosus (SLE):

- **Symptoms:** Fatigue, arthralgia, cognitive impairment (“lupus fog”), photosensitivity
- **Distinguishing features:** Malar rash, serositis (pleuritis, pericarditis), renal involvement
- **Testing:** ANA positive (high titer, typically $\geq 1 : 160$), anti-dsDNA, anti-Sm antibodies; low complement (C3, C4); elevated ESR/CRP during flares
- **Key distinction:** SLE has *elevated* inflammatory markers; ME/CFS has *normal or low* ESR/CRP

Sjögren Syndrome:

- **Symptoms:** Fatigue, dry eyes (keratoconjunctivitis sicca), dry mouth (xerostomia)
- **Distinguishing features:** Objective evidence of decreased tear/saliva production
- **Testing:** Anti-Ro (SSA), anti-La (SSB) antibodies; Schirmer test, salivary flow rate

Rheumatoid Arthritis:

- **Symptoms:** Fatigue, joint pain
- **Distinguishing features:** Joint swelling, morning stiffness > 1 hour, symmetric small joint involvement
- **Testing:** Rheumatoid factor (RF), anti-CCP antibodies, elevated ESR/CRP

Multiple Sclerosis (MS):

- **Symptoms:** Fatigue, cognitive impairment, sensory disturbances
- **Distinguishing features:** Focal neurological deficits (optic neuritis, weakness, sensory loss), relapsing-remitting pattern
- **Testing:** MRI brain and spine (demyelinating lesions disseminated in space and time), CSF oligoclonal bands

Hematologic Disorders

• **Requirement 14: Anemia Workup**

Hematologic disorders must be ruled out in the differential diagnosis of ME/CFS.

Iron Deficiency Anemia:

- **Symptoms:** Fatigue, dyspnea on exertion, pica (ice chewing)
- **Testing:** CBC shows microcytic anemia (low MCV); ferritin low (<30 ng/mL)
- **Note:** Iron deficiency *without anemia* (normal hemoglobin, low ferritin) can cause fatigue and restless legs syndrome; should be treated but is insufficient to explain ME/CFS severity

Vitamin B12 Deficiency:

- **Symptoms:** Fatigue, cognitive impairment, peripheral neuropathy, macrocytic anemia
- **Testing:** B12 level <200 pg/mL; methylmalonic acid (MMA) elevated if tissue deficiency

Infectious Diseases

Observation 23 (Post-Infectious vs. Chronic Active Infection). Distinguishing ME/CFS from active infections and post-infectious fatigue is essential for proper diagnosis and management.

Chronic Active Infections (Must Rule Out):

- **HIV/AIDS:** Check HIV antibody/antigen test
- **Hepatitis B/C:** Check HBsAg, anti-HCV
- **Tuberculosis:** In endemic areas or high-risk patients, check tuberculin skin test or interferon-gamma release assay
- **Lyme disease:** In endemic areas with appropriate exposure history, check Lyme serology (ELISA, Western blot)

Post-Infectious Fatigue vs. ME/CFS: Many acute infections (influenza, mononucleosis, COVID-19) are followed by transient fatigue lasting weeks to months. Distinguish from ME/CFS by:

- **Duration:** Post-infectious fatigue typically improves by 3–6 months; ME/CFS persists > 6 months without improvement
- **Post-exertional malaise:** True PEM with delayed onset and prolonged recovery is specific to ME/CFS
- **Trajectory:** Post-infectious fatigue shows gradual improvement; ME/CFS shows plateau or worsening

This distinction is critical in the first 6 months post-infection, as aggressive pacing during this period may prevent transition to established ME/CFS.

Malignancy

△ Warning 11: Occult Malignancy

Cancer-related fatigue can mimic ME/CFS, particularly in early stages without obvious tumor burden:

- **Red flags:** Unintentional weight loss, night sweats, fever, lymphadenopathy, age >50 with new-onset fatigue
- **Screening:** Age-appropriate cancer screening (colonoscopy, mammography); if red flags present, consider CT chest/abdomen/pelvis

- **Laboratory clues:** Anemia, elevated ESR, abnormal WBC count

Psychiatric Disorders

Observation 24 (Depression vs. ME/CFS: Critical Distinctions). Major depression can cause fatigue and cognitive impairment, but several features distinguish it from ME/CFS:

Post-Exertional Malaise (Pathognomonic for ME/CFS):

- **ME/CFS:** Physical or cognitive exertion triggers delayed (12–72 hours) symptom worsening lasting days to weeks
- **Depression:** Activity may be difficult due to lack of motivation, but exertion does NOT trigger delayed physiological crashes
- **Key question:** “If you push through and do an activity you enjoy, do you crash afterward?”
 - ME/CFS: Yes, even desired activities trigger PEM
 - Depression: Enjoyable activities may temporarily improve mood

Anhedonia (Pathognomonic for Depression):

- **Depression:** Loss of interest or pleasure in previously enjoyed activities (anhedonia is a core feature)
- **ME/CFS:** Patients *want* to do activities but are physically unable; they retain interest but lack capacity

Effort vs. Performance:

- **Depression:** Reduced effort (“I don’t feel like doing this”), but if motivation can be mustered, performance is intact
- **ME/CFS:** Normal or increased effort with reduced performance (“I’m trying as hard as I can but my body won’t do it”)

Objective Biomarkers:

- **Two-day CPET:** ME/CFS shows failure to reproduce VO_2max on Day 2; depression does not
- **Orthostatic intolerance:** Objective POTS/NMH on testing supports ME/CFS
- **Inflammatory markers:** Heng panel, cytokine signatures abnormal in ME/CFS

Comorbid Depression in ME/CFS: Many ME/CFS patients develop **reactive depression** (consequence of severe disability, loss of career/social life). This is distinct from primary depression:

- Reactive depression: Depression began *after* ME/CFS onset; patient grieves loss of function
- Primary depression: Depression preceded fatigue; fatigue is a symptom of depression

Treating comorbid depression in ME/CFS is appropriate and may improve quality of life, but antidepressants do not cure ME/CFS.

4.8.2 Comorbid Conditions vs. Alternative Diagnoses

Observation 25 (The ME/CFS Septad). Several conditions frequently co-occur with ME/CFS at rates far exceeding chance, suggesting shared pathophysiology:

1. **ME/CFS** (Myalgic Encephalomyelitis/Chronic Fatigue Syndrome)
2. **Fibromyalgia**: Widespread pain with tender points (30–70% of ME/CFS patients)
3. **POTS** (Postural Orthostatic Tachycardia Syndrome): (70–90% of ME/CFS patients)
4. **MCAS** (Mast Cell Activation Syndrome): Histamine-mediated symptoms (estimates 10–50%)
5. **hEDS** (Hypermobile Ehlers-Danlos Syndrome): Joint hypermobility (higher in ME/CFS than general population)
6. **IBS** (Irritable Bowel Syndrome): Functional GI symptoms (30–50% of ME/CFS patients)
7. **IC** (Interstitial Cystitis): Bladder pain, urinary frequency

Clinical Implication: These conditions are **comorbidities**, not alternative diagnoses. Their presence does NOT exclude ME/CFS. In fact, meeting criteria for multiple septad conditions strengthens the ME/CFS diagnosis and suggests common underlying mechanisms (autonomic dysfunction, small fiber neuropathy, immune activation).

4.8.3 When Comorbidities May Be Primary Drivers

While ME/CFS and its comorbidities typically coexist, in some patients a “comorbidity” may actually be the *primary driver* of symptoms, with ME/CFS-like presentation being downstream consequence rather than the core disease.

Observation 26 (Comorbidities as Potential Primary Pathology). Consider whether an apparent “comorbidity” might be the primary driver when:

MCAS/Histamine Intolerance as Primary:

- Symptoms fluctuate dramatically with dietary triggers (high-histamine foods)
- Marked improvement with H1/H2 blockade disproportionate to typical ME/CFS response

- Intestinal symptoms preceded and dominate fatigue
- Proposed cascade: MCAS → intestinal barrier dysfunction → amino acid malabsorption → mitochondrial failure → ME/CFS phenotype (see Section 11.1.6)

POTS as Primary:

- Fatigue and cognitive symptoms are primarily orthostatic (worse upright, better supine)
- Dramatic improvement with POTS-specific treatment (compression, fluids, fludrocortisone, ivabradine)
- Heart rate criteria met (≥ 30 bpm increase within 10 minutes of standing)
- Consider whether “ME/CFS” is actually deconditioning secondary to untreated POTS

Chronic Viral Reactivation as Primary:

- Documented elevated EBV or HHV-6 titers (especially IgM or PCR positivity)
- Symptom flares correlate with viral reactivation markers
- Response to antiviral therapy (valacyclovir for EBV, valganciclovir for HHV-6/CMV)
- Cimetidine trial produces dramatic improvement (suggests immune enhancement against herpesvirus)

Craniocervical Instability (CCI) as Primary:

- Symptoms markedly position-dependent
- Hypermobility (hEDS) with progressive neurological features
- Suboccipital headaches, neck pain, visual disturbances
- Upright MRI shows craniocervical abnormalities (see Section 5.6.11)

Clinical Approach: Test the Primary Driver Hypothesis. When a comorbidity might be primary:

1. **Prioritize that condition’s workup:** Comprehensive evaluation of the suspected primary driver
2. **Targeted treatment trial:** If the comorbidity is primary, treating it should produce disproportionate improvement
3. **Assess response:** Dramatic improvement ($>50\%$ symptom reduction) with targeted treatment suggests the “comorbidity” was actually the primary pathology
4. **Re-evaluate if partial response:** Partial improvement suggests the comorbidity contributes but is not the sole driver; ME/CFS diagnosis remains appropriate

△ Warning 12: Avoid Premature Reclassification

Do NOT reclassify ME/CFS as “just POTS” or “just MCAS” without:

- Demonstrating that treating the suspected primary condition produces substantial, sustained improvement

- Confirming that post-exertional malaise resolves (not just fatigue)
- Documenting functional recovery, not just symptom reduction

Many patients have true comorbid ME/CFS + POTS + MCAS where all contribute. Treating comorbidities improves but does not cure ME/CFS in most cases.

The Cascading Comorbidity Model. Rather than independent conditions, the septad conditions may cascade:

- **hEDS** → Vascular laxity → **POTS**
- **MCAS** → Intestinal barrier dysfunction → Malabsorption → **Mitochondrial dysfunction**
- **Dysautonomia** → Vagal dysfunction → **GI dysmotility** → **SIBO**
- **Chronic infection** → Immune exhaustion → **Autoimmunity**
- **Small fiber neuropathy** → Autonomic neuropathy → **POTS**

This cascading model suggests that identifying and treating upstream conditions may interrupt downstream pathology. For detailed discussion of the “Septad” framework and its limitations, see Section 5.6.11.

4.8.4 Diagnostic Algorithm

Observation 27 (Decision Tree for ME/CFS Diagnosis). 1. **Step 1: Screen for post-exertional malaise**

- If PEM absent → Consider alternative diagnosis (depression, deconditioning, other fatiguing condition)
- If PEM present → Proceed to Step 2

2. **Step 2: Rule out exclusions via laboratory testing**

- CBC, CMP, TSH/free T4, ESR/CRP, ANA, vitamin D, B12, sleep study
- If positive finding that fully explains symptoms → Treat that condition
- If tests normal or findings insufficient to explain severity → Proceed to Step 3

3. **Step 3: Assess duration and functional impact**

- Duration ≥ 6 months? (or ≥ 3 months in severe pediatric cases)
- Substantial functional impairment?
- If yes to both → Proceed to Step 4

4. **Step 4: Apply diagnostic criteria**

- Use Canadian Consensus, IOM, or ICC criteria
- All require: PEM, unrefreshing sleep, multi-system symptoms
- If criteria met → Diagnose ME/CFS

5. **Step 5: Assess for comorbidities**

- Screen for septad conditions (fibromyalgia, POTS, MCAS, hEDS, IBS)

- Document which conditions are present (multi-label classification)
- These do not exclude ME/CFS; they inform treatment strategy

6. **Step 6: Biological phenotyping (if resources permit)**

- Apply Tier 2 framework: assess autoimmune, mitochondrial, neuroinflammatory, dysautonomia, endothelial domains
- Guide treatment stratification

7. **Step 7: Risk stratification**

- Apply Tier 3 RED FLAG criteria
- If ≥ 2 RED FLAGS → Emergency intervention protocol

4.8.5 When to Reconsider the Diagnosis

△ **Warning 13: Red Flags Suggesting Alternative Diagnosis**

ME/CFS diagnosis should be reconsidered if:

- **New focal neurological signs:** Weakness, sensory loss, visual changes (suggests MS, tumor, stroke)
- **Fever, night sweats, unintentional weight loss:** Suggests infection, malignancy, autoimmune disease
- **Rapid progression over weeks:** ME/CFS typically progresses over months to years; rapid worsening suggests acute process
- **Lack of PEM:** If re-evaluation reveals no true post-exertional malaise, reconsider alternative diagnoses
- **Complete resolution with psychiatric treatment:** If depression treatment alone fully resolves “fatigue,” the diagnosis was likely primary depression, not ME/CFS

5 Disease Course and Prognosis

5.1 Onset Patterns

The manner in which ME/CFS begins has both diagnostic and prognostic significance. Two primary onset patterns are recognized: acute (typically post-infectious) and gradual [120]. Understanding these patterns helps clinicians recognize the disease earlier and may inform treatment approaches.

Post-Infectious Onset

Approximately 64% of ME/CFS cases begin with an acute infectious illness [120]. The patient experiences what appears to be a typical viral infection—influenza, infectious mononucleosis, respiratory illness, or gastrointestinal infection—but fails to recover. Weeks pass, then months, and the expected return to health never comes.

Common Triggering Infections. Documented infectious triggers include:

- **Epstein-Barr virus (EBV):** The most studied trigger, with 10–12% of infectious mononucleosis cases progressing to ME/CFS
- **SARS-CoV-2:** COVID-19 has created a new wave of post-infectious ME/CFS (Long COVID with ME/CFS phenotype)
- **Influenza:** Both seasonal and pandemic strains
- **Enteroviruses:** Including coxsackieviruses and echoviruses
- **Ross River virus:** Endemic trigger in Australia
- **Q fever (*Coxiella burnetii*):** Bacterial trigger with well-documented post-infectious fatigue
- **Giardiasis:** Parasitic infection linked to post-infectious ME/CFS in outbreak studies

Vaccine-Related Complications as Potential Triggers. In rare cases, immunizations have been temporally associated with ME/CFS onset or significant symptom exacerbation. While post-infectious complications from vaccinations are extremely rare, susceptible individuals may experience:

- **Autoimmune neurological reactions:** Case reports describe bilateral facial nerve paralysis, vision changes, or neuropathic symptoms developing days to weeks after vaccination
- **Neuroendocrine disruption:** Including abnormal menstrual function or temperature dysregulation

- **Small fiber neuropathy exacerbation:** Some patients with pre-existing neuropathic predisposition have reported symptom worsening post-vaccination
- **Molecular mimicry hypotheses:** Spike protein epitopes may cross-react with neural tissue, particularly if the individual has pre-existing autoimmune features

Clinical note: These complications are reported anecdotally and have not been formally characterized in ME/CFS populations. The population attributable risk remains extremely small. This section is included to alert clinicians to the possibility that vaccine-temporal associations may represent a specific immunological subtype rather than universal vaccine danger. Patients with this history warrant investigation of autoimmune markers and underlying immune dysregulation (see Section 7.2.2).

Observation 28 (Viral Associations: Meta-Analytic Evidence). A 2023 systematic review and meta-analysis of 64 studies (n=4,971 ME/CFS patients, n=9,221 controls) examining 18 viral species identified significant associations between ME/CFS and multiple viral infections [121]. Five viruses demonstrated odds ratios exceeding 2.0: Borna disease virus (OR≥3.47), HHV-7 (OR>2.0), parvovirus B19 (OR>2.0), enterovirus (OR>2.0), and coxsackie B virus (OR>2.0). However, high heterogeneity (>50%) was observed for EBV and enterovirus associations, suggesting these viral triggers may apply to specific subgroups rather than uniformly across all ME/CFS cases.

△ **Warning 1: Association vs. Causation in Viral Triggers**

While meta-analytic evidence demonstrates statistical associations between viral infections and ME/CFS onset [121], these data cannot establish causation. Viral reactivation may represent a consequence of immune dysfunction rather than the initiating cause. Additionally, detection bias may inflate associations, as ME/CFS patients typically undergo more extensive viral testing than matched controls. The observed heterogeneity across studies indicates that viral etiology likely applies to subsets of ME/CFS patients rather than representing a universal mechanism.

The NIH deep phenotyping study focused specifically on post-infectious ME/CFS, providing detailed characterization of this subgroup [5].

Temporal Pattern. In acute post-infectious onset, the transition from acute infection to chronic illness is often abrupt. Patients can frequently identify the specific day or week when their illness began. The typical pattern:

1. Acute infectious illness with standard symptoms (fever, malaise, respiratory or gastrointestinal symptoms)
2. Expected recovery does not occur after 2–4 weeks
3. Persistent fatigue, cognitive impairment, and post-exertional malaise emerge
4. Full ME/CFS symptom complex develops over weeks to months
5. Stabilization at significantly reduced functional capacity

Pathophysiological Implications. Post-infectious onset suggests mechanisms involving:

- Persistent viral reservoirs or reactivation of latent viruses
- Post-infectious autoimmunity triggered by molecular mimicry
- Chronic immune activation and inflammation
- Disruption of the gut microbiome
- Autonomic nervous system dysregulation

Brain imaging studies show distinct abnormalities in post-infectious ME/CFS compared to gradual-onset cases, supporting the notion that different onset patterns may involve different pathophysiological mechanisms.

Prognosis. Some studies suggest that post-infectious onset may carry a better prognosis than gradual onset, particularly when the triggering infection can be identified and when illness duration is short before diagnosis. However, this finding is not consistent across all studies, and many post-infectious cases progress to severe, permanent disability.

Gradual Onset

Approximately 36% of ME/CFS cases (range 23–41% across studies) develop gradually without a clear infectious trigger [120]. Symptoms accumulate over months to years, making it difficult to identify when the illness truly began.

Progressive Symptom Accumulation. Gradual-onset ME/CFS typically follows a pattern of:

1. Increasing fatigue attributed to stress, overwork, or aging
2. Sleep disturbances that fail to respond to standard interventions
3. Cognitive difficulties (brain fog, concentration problems, word-finding difficulties)
4. Exercise intolerance that progressively worsens
5. Development of post-exertional malaise, often initially unrecognized
6. Eventual recognition that something is fundamentally wrong

The insidious nature of gradual onset often delays diagnosis, as patients and clinicians attribute symptoms to other causes. Mean diagnostic delay is longer in gradual-onset cases, which has prognostic significance (see Section 5.5).

Risk Factors. Gradual onset has been associated with:

- Prior history of multiple infections (cumulative immune burden)
- Chronic stress or overwork
- Other chronic illnesses
- Higher rates of psychiatric comorbidity (though causation is unclear)

- Possible environmental exposures

The association with psychiatric comorbidity is controversial. It may reflect true biological comorbidity, diagnostic confusion (ME/CFS misdiagnosed as depression), or shared underlying mechanisms affecting both mood and energy regulation.

Diagnostic Challenges. Gradual onset creates particular diagnostic challenges:

- No clear temporal marker for illness onset
- Symptoms may be attributed to depression, anxiety, or somatization
- Lack of infectious trigger makes the diagnosis seem less “legitimate”
- Pre-illness functional level may be difficult to establish
- Patients may have adapted to declining function without recognizing its significance

Two-Phase Onset Pattern

A third pattern has been identified in some patients: two-phase onset [120]. This pattern involves:

1. Initial sharp deterioration (often post-infectious)
2. Partial improvement over months
3. Secondary deterioration to chronic ME/CFS

This pattern may represent failed recovery from post-infectious illness, with initial improvement reflecting resolution of acute infection while underlying ME/CFS pathophysiology continues to develop.

Multi-Hit Cascade Pattern

Clinical observation suggests an additional onset variant: the **multi-hit cascade**, where ME/CFS develops through cumulative immune challenges over years rather than a single triggering event.

Typical Multi-Hit Trajectory. The multi-hit pattern involves sequential stressors that progressively deplete compensatory reserves:

1. **Baseline vulnerability:** Pre-existing inflammatory condition (e.g., interstitial cystitis, MCAS, fibromyalgia) or constitutional immune phenotype
2. **Initial hit:** Significant infection during period of stress; patient “never fully recovers” but achieves marginal functionality
3. **Subsequent hits:** Additional immune challenges (viral infections, significant physiological stressors) cause further decompensation

4. **Apparent recoveries:** Periods of improved function that retrospectively appear as PEM cycles or enforced pacing, not true remission
5. **Final collapse:** A seemingly minor challenge (e.g., influenza, moderate activity) triggers complete decompensation after years of accumulated damage

The False Recovery Pattern. A critical insight from multi-hit cases is that “better periods” between challenges may represent **precarious minimal functionality** rather than true recovery:

- Patient operates within a severely reduced energy envelope
- Any significant stressor reveals underlying dysfunction
- Enforced rest (e.g., during lockdowns) produces improvement that disappears with return to normal activity
- What patients retrospectively recognize as “ME/CFS in episodes” was actually PEM cycling
- Each viral hit does not “cause ME/CFS again” but reveals and worsens dysfunction that was never resolved

⚠ **Limitation 1: Multi-Hit Cascade: Retrospective Pattern Recognition, Not Prospective Validation**

The multi-hit cascade onset pattern described above is a clinical narrative constructed from retrospective patient histories, not a prospectively validated disease model. Limitations include:

- Retrospective histories are subject to recall bias: patients reconstruct a coherent narrative from fragmented memories, potentially imposing causal structure where coincidence exists.
- The “false recovery” concept is unfalsifiable as described—any improvement can be reinterpreted as “precarious minimal functionality” after subsequent worsening.
- No biomarker data exist to distinguish true multi-hit cascades from patients with standard post-infectious onset who happen to have sequential infections.
- The model cannot currently predict which patients in “marginal functionality” will decompensate versus stabilise.

⚠ **Warning 2: Distinguishing True Remission from Marginal Functionality**

Clinicians and patients should distinguish between:

- **True remission:** Return to pre-illness baseline; able to tolerate normal activity and minor infections without prolonged recovery
- **Marginal functionality:** Able to perform limited activities through careful pacing; any additional stressor causes crash; “recovered” state requires continuous effort to maintain

The multi-hit pattern suggests that patients in marginal functionality states remain vul-

nerable to progressive decompensation with each additional immune challenge. This has implications for long-term prognosis and the importance of aggressive infection prevention, stress management, and pacing even during “good periods.”

Research Implications. The multi-hit cascade pattern suggests that ME/CFS onset may involve:

- Cumulative immune burden rather than single triggering event
- Progressive exhaustion of compensatory mechanisms
- Threshold effects where final collapse appears disproportionate to the triggering event
- Potential for early intervention at any stage to prevent progression

This model aligns with the “exhausted immune surveillance” phenotype described in Section 7.2.2, where laboratory findings reflect end-stage depletion of immune reserves after years of chronic stimulation.

Clinical Significance of Onset Pattern

While onset pattern provides useful clinical information, it should not be overemphasized in individual patient management. Both post-infectious and gradual-onset patients develop the same symptom complex and require the same management approaches. The key clinical implications of onset pattern include:

- **Diagnostic confidence:** Post-infectious onset with clear temporal association may increase diagnostic confidence
- **Patient validation:** Understanding that infections can trigger chronic illness helps patients understand their condition
- **Research stratification:** Onset pattern may be important for identifying disease subgroups in research
- **Epidemiological monitoring:** Tracking post-infectious ME/CFS helps quantify the burden of infectious diseases

5.2 Disease Severity Levels

Defining Severity

The International Consensus Criteria (ICC) defines ME/CFS severity based on functional capacity relative to pre-illness baseline [122]. These classifications have been objectively validated through activity monitoring, cardiopulmonary exercise testing, and standardized questionnaires [123]. Understanding severity levels is essential for appropriate clinical management, realistic expectations, and resource allocation.

Prevalence across severity levels follows a characteristic distribution: approximately 29% mild, 58% moderate, 11% severe, and 2% very severe [124]. However, functional impairment studies using different criteria (housebound/bedbound status) report that 25–25.7% of patients experience severe functional limitation at some point in their illness [125, 126]. The discrepancy between these figures (13% severe/very severe vs. 25% housebound/bedbound) likely reflects several factors: (1) some moderate patients experience periods of being housebound during severe crashes without meeting criteria for severe ME/CFS classification, (2) the 25% figure represents lifetime prevalence ("at some point") while the 13% figure is a cross-sectional snapshot, and (3) severe patients are systematically underrepresented in research cohorts due to inability to participate. These proportions likely underestimate the true burden of severe disease.

Functional Capacity and Objective Measures

Objective validation studies demonstrate that self-reported severity classifications correlate strongly with measurable physiological parameters [123]:

Table 5.1: Objective measures across ME/CFS severity levels

Measure	Mild	Moderate	Severe
Daily steps (mean)	8,235	5,195	2,031
SF-36 Physical Functioning	70	43	15
Peak VO ₂ (% predicted)	90%	64%	48%
VO ₂ at ventilatory threshold	47%	38%	30%

All differences between severity groups are statistically significant ($p < 0.0001$), confirming that patient-reported severity reflects genuine physiological impairment rather than subjective perception.

Mild ME/CFS

Mild ME/CFS represents approximately 50% reduction in pre-illness activity level [122]. Despite the designation "mild," this category describes substantial disability that would be considered severe in most other medical contexts.

Functional Capacity. Patients with mild ME/CFS may maintain some degree of employment or education, though typically with significant accommodations:

- Reduced hours (part-time work or study)
- Flexible scheduling to accommodate energy fluctuations
- Remote work arrangements to eliminate commuting
- Extended deadlines and modified workloads
- Frequent rest breaks throughout the day

Daily step counts averaging 8,235 steps indicate preserved mobility but at the lower end of healthy population norms (typically 7,000–10,000 steps daily). Peak oxygen consumption at 90% of predicted suggests maintained aerobic capacity under controlled testing conditions, though real-world performance is constrained by post-exertional malaise.

The Invisible Illness Phenomenon. Patients with mild ME/CFS often appear healthy to outside observers, creating a dangerous disconnect between perceived and actual capacity. This “invisible illness” phenomenon leads to:

- Disbelief from employers, educators, family, and healthcare providers
- Pressure to perform at pre-illness levels
- Social isolation when patients decline activities to conserve energy
- Self-doubt about the legitimacy of their condition
- Delayed diagnosis and inappropriate treatment recommendations

The ability to “pass” as healthy exacts a heavy toll. Patients may push through symptoms to meet social expectations, triggering post-exertional malaise and risking progression to more severe disease.

Energy Envelope Management. Successful management of mild ME/CFS requires strict adherence to the energy envelope—staying within available energy reserves rather than borrowing against future capacity [61]. Patients must:

- Track activity levels and symptoms systematically
- Identify personal triggers for post-exertional malaise
- Accept permanent lifestyle modifications
- Resist the temptation to “test” limits during good periods
- Build substantial rest margins into daily schedules

Risk of Progression. Mild ME/CFS is not a stable endpoint. Patients who exceed their energy envelope repeatedly, whether through choice or necessity, face significant risk of progression to moderate or severe disease. Common triggers for deterioration include:

- Intercurrent infections (viral, bacterial)
- Physical overexertion (exercise, travel, demanding work)
- Cognitive overexertion (intensive mental work, emotional stress)
- Medical procedures (surgery, dental work, vaccinations)
- Life stressors (bereavement, relationship breakdown, financial pressure)

Once deterioration occurs, return to baseline is not guaranteed. Many patients describe a “ratchet effect” where each crash leaves them at a lower functional level than before.

Moderate ME/CFS

Moderate ME/CFS describes patients who are mostly housebound, with severely restricted activity in all domains [122, 127]. This category represents the largest proportion of the ME/CFS population (approximately 58%) and encompasses significant heterogeneity in functional capacity.

Functional Limitations. The NICE guideline characterizes moderate ME/CFS by [127]:

- Reduced mobility affecting all daily activities
- Cessation of work or education
- Required rest periods, often 1–2 hours in the afternoon
- Poor quality, disturbed sleep that fails to restore energy
- Significant reduction in social activities

Daily step counts averaging 5,195 reflect the housebound nature of this severity level—enough mobility to move within the home but insufficient for regular excursions. SF-36 physical functioning scores of 43 (compared to population norms near 85) quantify the profound limitation.

The Daily Energy Budget. Patients with moderate ME/CFS face constant decisions about energy allocation. A finite daily budget must cover all activities, and exceeding this budget triggers post-exertional malaise. Typical trade-offs include:

- Shower *or* prepare a meal, but not both
- Brief phone conversation *or* a short walk
- Medical appointment requiring days of pre-appointment rest and post-appointment recovery
- Social visit measured in minutes rather than hours

The cognitive and emotional dimensions of this constant calculation constitute a burden in themselves. Patients describe exhaustion from the relentless need to monitor, plan, and restrict.

Loss of Independence. Moderate ME/CFS typically requires some degree of assistance with daily living:

- Meal preparation and household management
- Transportation for medical appointments
- Shopping and errands
- Medication management during cognitive impairment
- Personal care during severe symptom flares

This dependence represents a profound loss for previously independent individuals. The psychological impact of needing help with basic functions compounds the physical suffering of the disease.

Employment and Financial Impact. Most patients with moderate ME/CFS cannot maintain employment. Among the ME/CFS population overall, only 13% work full-time and 54% are unemployed (compared to 9% in the general population) [125]. The financial consequences cascade:

- Loss of income at peak earning years
- Depletion of savings for living expenses
- Inability to afford treatments not covered by insurance
- Housing instability when rent or mortgage becomes unaffordable
- Dependence on family support or social welfare programs
- Lengthy disability claim battles with insurers who dispute ME/CFS legitimacy

Social Isolation. The combination of energy limitations, unpredictable symptoms, and inability to participate in normal activities leads to progressive social isolation:

- Friends drift away when invitations are repeatedly declined
- Family relationships strain under the burden of caregiving
- Online interaction becomes the primary social connection
- Special occasions (weddings, graduations, funerals) become impossible to attend
- The patient's world shrinks to the confines of their home

Severe ME/CFS

Severe ME/CFS describes patients who are mostly bedridden, with profound limitation in all activities [122]. Approximately 11% of ME/CFS patients fall into this category, though they are underrepresented in research due to inability to travel to study sites or tolerate research protocols.

Functional Status. The NICE guideline characterizes severe ME/CFS by [127]:

- Inability to perform any activity for themselves, or only minimal tasks (face washing, teeth cleaning)
- Severe cognitive difficulties affecting concentration, memory, and communication
- Wheelchair dependence for any mobility outside the bed
- Inability to leave the house, or severe prolonged after-effects if they do
- Mostly bedridden with only brief periods of sitting up
- Extreme sensitivity to light and sound

Daily step counts averaging only 2,031 reflect the near-complete loss of mobility. Peak oxygen consumption at 48% of predicted indicates severe impairment of the body's fundamental capacity to generate energy.

Caregiver Dependence. Patients with severe ME/CFS require substantial assistance with all activities of daily living:

- Personal hygiene (bathing, toileting, grooming)
- Feeding and hydration
- Medication administration
- Position changes to prevent pressure injuries
- Communication with healthcare providers
- Protection from environmental triggers

This level of care typically requires a dedicated family caregiver or, for those without family support, professional home care services that few can afford and that few providers understand how to deliver appropriately for ME/CFS.

Qualitative Difference. Research suggests that severe ME/CFS may represent a qualitatively different disease state rather than simply a more extreme point on a continuum [128]. Compared to mild and moderate patients, those with severe ME/CFS demonstrate:

- Greater autonomic dysfunction
- More frequent and more severe post-exertional malaise
- More pronounced cognitive impairment
- More multisystem symptom involvement
- Significantly worse scores on every SF-36 domain

These findings suggest that progression to severe disease may involve additional pathophysiological mechanisms beyond those operating in milder forms, with implications for treatment approaches.

Healthcare Access Crisis. Approximately 25% of ME/CFS patients are severely affected and almost exclusively housebound, yet many receive no medical care despite being most in need [129]. Barriers include:

- Inability to travel to medical facilities
- Post-exertional malaise triggered by the examination itself
- Lack of physicians willing to make home visits
- Medical professionals unfamiliar with severe ME/CFS presentation
- Insurance systems designed around ambulatory care
- Emergency departments that provide inappropriate treatment (bright lights, noise, activity recommendations)

The result is a population of severely ill patients who are medically abandoned—too sick to access the healthcare system, and invisible to a system that has no mechanism to find them.

Very Severe ME/CFS

Very severe ME/CFS represents the extreme end of the disease spectrum: patients who are completely bedridden and require help with all basic functions [122]. Approximately 2% of ME/CFS patients fall into this category, representing an estimated 62,000 people in the United States alone [125].

The detailed reality of very severe ME/CFS—the complete energy bankruptcy, the necessity of existence in darkness and silence, the loss of basic bodily functions, and the existential suffering that leads many to wish for death—is addressed comprehensively in Section 5.3.

Key Features. Very severe ME/CFS is characterized by [127]:

- Complete confinement to bed, 24 hours per day
- Dependence on others for all care needs
- Inability to tolerate any sensory input (light, sound, touch, movement)
- Profound cognitive impairment affecting communication
- Feeding difficulties requiring liquid nutrition or tube feeding
- Complete loss of independence and autonomy

The Invisible Population. Very severe patients are almost entirely absent from research studies, clinical guidelines, and healthcare systems. They cannot:

- Travel to research facilities
- Tolerate standard medical examinations
- Complete questionnaires or interviews
- Advocate for themselves in healthcare settings
- Participate in patient organizations or support groups

Their existence is known primarily through caregiver reports and memorial records. They suffer in silence, hidden from the medical establishment that should be serving them.

5.3 The Devastating Reality of Severe ME/CFS

Disturbing Content and Disease Lethality

This chapter documents the extreme suffering experienced by patients with severe and very severe ME/CFS. The content is intentionally disturbing because the reality of this disease is disturbing. Readers—particularly healthcare providers, policymakers, and family members—must understand that ME/CFS at its worst represents one of the most devastating conditions in medicine. This is not hyperbole. The evidence presented here demonstrates that severe ME/CFS produces suffering comparable to or exceeding that of terminal cancer, yet without the certainty of death's release.

ME/CFS kills. It kills through suicide when patients can no longer endure the suffering. It kills through cardiac complications from years of autonomic dysfunction. It kills through malnutrition when patients become too weak to eat. It kills through medical neglect when healthcare systems refuse to believe or treat patients adequately.

This chapter exists because the medical community's failure to recognize the severity of this disease has cost lives. Every reader who finishes this chapter should be afraid—not of catching ME/CFS, but of the consequences of continued medical and societal indifference to those who already suffer from it.

5.3.1 The Scale of Catastrophe

Approximately 25% of all ME/CFS patients—an estimated 250,000 people in the United States alone, and over 2 million worldwide—experience severe or very severe disease that leaves them housebound or completely bedbound [126]. These patients have largely vanished from public view. They cannot advocate for themselves. They cannot participate in research studies. They cannot visit doctors' offices. Many have been abandoned by the healthcare system entirely.

Quality of Life: Worse Than Cancer

A landmark 2015 study published in *PLOS ONE* compared the health-related quality of life (HRQoL) of ME/CFS patients against 20 other chronic conditions, including multiple sclerosis, stroke, lung cancer, diabetes, and heart disease [59]. The findings were unequivocal:

Observation 29 (Quality of Life Comparison Across Chronic Conditions). **ME/CFS had the lowest quality of life of all 20 chronic conditions studied**—worse than multiple sclerosis, worse than stroke, worse than cancer [59].

- ME/CFS EQ-5D score: 0.47 (vs. 0.85 population mean)
- ME/CFS quality of life is 55% of the general population average
- Only 7.6% of patients remained employed
- 52.2% were on disability pension

These figures represent the *average* ME/CFS patient. For severe and very severe patients, quality of life approaches or reaches zero.

The implications are staggering. A patient with lung cancer—facing chemotherapy, radiation, the terror of mortality—reports better quality of life than the average ME/CFS patient. And the ME/CFS patient faces this not for months or a few years of treatment, but potentially for decades, with no approved treatments and often no acknowledgment that their suffering is real.

Mortality: Dying Young

ME/CFS is not merely disabling—it is deadly. Memorial record studies consistently document dramatically reduced life expectancy. A 2016 analysis found that ME/CFS patients die, on average, **18–21 years earlier** than the general population [130]:

- **Mean age at death:** 55.9 years (vs. 73.5 years in the general population)
- **Cardiovascular death:** 58.8 years (vs. 77.7 years in controls)—nearly 19 years earlier
- **Suicide:** 41.3 years average age
- **Bedridden before death:** 48.2% of patients

A larger 2025 study of 512 deaths found even more concerning figures, with a mean age at death of 52.5 years—approximately 21 years younger than the general population [131]. The slight difference between studies (52.5 vs. 55.9 years) may reflect cohort composition, with the larger 2025 study potentially capturing more severe cases. The three leading causes of death in ME/CFS are: complications of the disease itself (28.3%), suicide (25.4%), and cancer (23.0%).

‡ Limitation 2: Mortality Figures: Memorial Record Selection Bias

The “18–21 years of lost life expectancy” figures cited above derive from memorial record studies, which capture only deaths reported to patient organisations. Population-based registry studies (Roberts et al. 2016, n=2,147; Smith et al. 2006, n=1,201) find no statistically significant elevation in all-cause mortality. The memorial record figures powerfully convey the severity of the disease but should not be cited as representative of ME/CFS mortality overall; they reflect the extreme tail of the severity distribution. See Section 29.5.1 for detailed methodological analysis.

5.3.2 Complete Energy Bankruptcy

The central feature of severe ME/CFS is **total energy depletion**—not fatigue in any ordinary sense, but a complete metabolic bankruptcy that leaves the body unable to perform even the most basic functions of survival.

What “No Energy” Actually Means

When a severe ME/CFS patient says they have “no energy,” they do not mean they are tired. They mean:

- **Breathing is effortful:** Each breath requires conscious work. The respiratory muscles, like all muscles, run on ATP that the body cannot produce.
- **Swallowing becomes dangerous:** The muscles required for swallowing fail. Food can be aspirated. Patients may require tube feeding to survive [132].
- **The heart struggles:** Cardiac output drops. Blood pools in extremities. Standing becomes impossible because the cardiovascular system cannot maintain perfusion to the brain.
- **Digestion stops:** Peristalsis requires energy. Food sits undigested for hours or days, causing severe gastrointestinal distress.
- **Temperature regulation fails:** The body cannot maintain homeostasis. Patients experience severe chills or overheating from minimal environmental changes.
- **Thinking becomes impossible:** The brain consumes 20% of the body’s energy. When that energy disappears, cognition shuts down—not gradually, but catastrophically.

The Sensation of Dying

Patients with severe ME/CFS describe the physical sensation as **drowning and burning alive simultaneously**—the body in a state of metabolic crisis, sending alarm signals that something is catastrophically wrong. One patient, Samuel, age 21, who chose euthanasia in 2024 rather than continue living with very severe ME/CFS, described it this way:

“The body thinks it is dying because it is running out of energy, and therefore triggers an extreme state of suffering. So bad that you often think there is only one option left.”

This is not metaphor. The mitochondria—the cellular power plants—have failed. Cells throughout the body are operating in crisis mode, triggering the same alarm systems that would activate if the body were actually dying of starvation or suffocation. The patient experiences genuine physiological distress signals, 24 hours a day, for years or decades.

Life-Threatening Malnutrition

A 2021 case series documented five patients with very severe ME/CFS who experienced **life-threatening malnutrition** [132]. Key findings:

- BMI dropped as low as **11.4** before tube feeding was initiated (healthy BMI: 18.5–24.9)
- Swallowing difficulties were repeatedly attributed to “psychological causes” by health-care providers
- Patients developed complications including poor wound healing, neurological damage, and osteoporosis

- Healthcare providers exhibited “clinical inertia”—failing to act even as patients starved

The reason for malnutrition in severe ME/CFS is multifactorial:

1. **Inability to access food:** Patients too weak to prepare or obtain meals
2. **Inability to chew:** Jaw muscles exhaust within seconds
3. **Inability to swallow:** Pharyngeal muscles fail; choking risk
4. **Severe gastrointestinal dysfunction:** Food causes extreme distress
5. **Food intolerances:** Multiple chemical sensitivities make most foods intolerable
6. **Energy cost of eating:** Digestion itself consumes energy the patient cannot spare

5.3.3 Existence in Darkness and Silence

Extreme Sensory Hypersensitivity

Severe ME/CFS patients often develop profound hypersensitivity to light, sound, touch, and smell. A 2023 study found that 73% of ME/CFS patients experience at least one form of sensory hypersensitivity, with 50.4% experiencing both light and sound sensitivity [133]. In severe cases, this hypersensitivity becomes so extreme that normal environmental stimuli cause physical pain and neurological crashes.

Living in Total Darkness

Many severely affected patients must exist in complete or near-complete darkness, 24 hours a day, 365 days a year:

- **Blackout curtains** covering all windows, often with additional light-blocking material
- **No screens:** Television, computers, phones—even for seconds—overwhelm the nervous system
- **No reading:** The visual processing required to read text exhausts available energy
- **Eye masks worn continuously:** Even the faint glow of a digital clock causes distress

The neurological basis involves **central sensitization**—the central nervous system has become hypervigilant, amplifying all incoming sensory signals to painful levels. Light that would be comfortable for a healthy person registers as blinding pain to the severe ME/CFS patient.

Existence in Silence

Sound hypersensitivity (hyperacusis) forces many patients into isolation that approaches sensory deprivation:

- **Double hearing protection:** Earplugs inside industrial ear defenders
- **No music:** What was once a source of joy becomes neurologically unbearable
- **No conversation:** Human speech—even whispered—triggers crashes

- **No television, podcasts, or audiobooks:** All auditory input is too stimulating
- **Environmental noise intolerance:** A car passing outside, a door closing in another room, birds singing—all cause distress

Samuel, the 21-year-old Austrian patient, described his daily existence before choosing euthanasia:

“I must lie in bed 24 hours a day and must not move too much. It must be permanently dark because I cannot tolerate light. I wear double hearing protection because I cannot tolerate sounds. I cannot watch television or videos on my phone for even a second, because moving images overwhelm my nervous system and trigger unbearable suffering. I cannot listen to music or podcasts. I cannot even speak with my own mother, who cares for me, because listening is too exhausting, and speaking itself has become completely impossible. So I must communicate with a pen and paper. My phone I can use only for a few minutes or seconds for messages. Sometimes not at all.”

Touch and Chemical Sensitivities

Beyond light and sound, severe patients often develop:

- **Allodynia:** Normal touch registers as pain. The weight of a blanket, the fabric of clothing, human contact—all cause suffering
- **Chemical sensitivities:** Perfumes, cleaning products, personal care items, cooking odors—all trigger neurological reactions
- **Electromagnetic hypersensitivity:** Some patients report distress from electronic devices, WiFi signals, or fluorescent lighting

The Isolation Chamber

The combined effect of these sensitivities is that severe ME/CFS patients exist in conditions that would constitute solitary confinement torture if imposed by a prison system:

- **No human contact:** Visitors cause crashes from sound, movement, perfume, emotional stimulation
- **No entertainment:** All forms of media are neurologically inaccessible
- **No communication:** Too weak to speak, too sensitive to listen, often unable even to write
- **Alone with thoughts:** Yet even thinking too intensely—positive or negative emotions—can trigger crashes

This is not depression-induced isolation. This is **biologically enforced solitary confinement**—the nervous system has become so dysfunctional that any form of stimulation causes physical harm.

5.3.4 The Prison of the Body

Inability to Perform Basic Bodily Functions

For very severe ME/CFS patients, the most basic functions of human existence become impossible:

Toileting

Going to the toilet—an activity healthy people perform without conscious thought—becomes a major physical challenge or impossibility:

- **Cannot walk to bathroom:** Must use bedpan, commode chair, or diapers
- **Cannot sit upright:** The energy required to maintain an upright position exceeds available reserves
- **Post-toileting crashes:** Even assisted toileting may trigger hours or days of worsened symptoms
- **Constipation:** Peristalsis requires energy; severe patients often have profound constipation
- **Incontinence:** Some patients lose bladder or bowel control from neurological dysfunction

Bathing and Hygiene

Personal hygiene becomes a distant memory for many severe patients:

- **Showering impossible:** Standing under running water requires too much energy; temperature changes too stimulating
- **Bed baths difficult:** Even passive bathing by a caregiver may trigger crashes
- **Teeth brushing exhausting:** The arm movement, the taste of toothpaste, the stimulation—all problematic
- **Hair care abandoned:** Washing, brushing, or cutting hair requires energy that doesn't exist

Samuel noted simply: “Going to the toilet is sometimes difficult. Showering is currently impossible due to extreme physical weakness and sensory overload.”

Eating and Drinking

As discussed in Section 5.3.2, eating itself becomes a dangerous activity:

- **Cannot sit up to eat:** Must be fed lying down or at extreme recline
- **Chewing exhausts jaw muscles:** Can manage only soft or liquid foods

- **Swallowing risk:** Aspiration pneumonia is a genuine threat
- **Tube feeding:** Some patients require nasogastric or PEG tubes for survival
- **TPN (Total Parenteral Nutrition):** In extreme cases, nutrition must bypass the digestive system entirely

Speaking and Communication

The ability to speak—the fundamental human capacity for connection—is lost:

- **Cannot produce speech:** The motor coordination, breath control, and cognitive load required exceed capacity
- **Cannot whisper:** Even minimal vocalization is too demanding
- **Written communication limited:** Holding a pen, forming letters, organizing thoughts—all require energy
- **Digital communication minimal:** A few seconds or minutes on a phone, if anything
- **Communication boards:** Some patients resort to pointing at letters or symbols

One patient reported: “After trying to talk, something got strained so severely that a few weeks later I could not swallow solid food without almost unbearable pain, so I had to switch to blended food. Even then it took years to settle down and there were scary times I really struggled with swallowing at all. I am still bedbound now, still unable to talk, or listen to music, or watch TV.”

Post-Exertional Malaise: The Trap

The defining feature that makes severe ME/CFS a trap from which there is no escape is **post-exertional malaise (PEM)**—any activity beyond the patient’s severely limited “energy envelope” triggers a crash that may last hours, days, weeks, or permanently worsen the baseline condition.

The PEM Trap

The cruel mathematics of severe ME/CFS:

1. Patient has energy for approximately nothing—lying still in darkness and silence
2. Any attempt to do something—speak, think, move, feel—costs energy
3. Energy expenditure triggers PEM: worsened symptoms, often for days
4. PEM reduces baseline capacity further
5. Return to step 1, but with even less capacity than before

This is why severe ME/CFS patients get worse, not better. Every attempt to “push through,” every well-meaning encouragement to “try a little activity,” every unavoidable exertion (a medical appointment, an emergency, a caregiver being unavailable) can permanently damage the patient.

Samuel described this trap:

“But that is not even the worst part. The worst thing about this disease is the cardinal symptom PEM (post-exertional malaise), which ensures that every smallest exceeding of my energy limits leads to a so-called crash and a permanent worsening of all my symptoms and my general condition. So I must bitterly pay for every attempt to live a little, and then end up in an even worse state than before.

Even if I only lie in bed, alone with my thoughts, I must be careful, because even too positive or too negative thoughts mean a crash and thus a deterioration in my condition.”

The Pain Dimension

Severe ME/CFS involves “severe and often almost constant, widespread pain” [126]. This pain has multiple components:

1. **Muscle pain:** Widespread myalgia from metabolic dysfunction and lactic acid accumulation
2. **Joint pain:** Diffuse arthralgia affecting major and minor joints
3. **Nerve pain:** Burning, shooting, or electrical sensations from small fiber neuropathy
4. **Headache:** Persistent headaches, often migrainous in character
5. **Allodynic pain:** Pain from normally non-painful stimuli (touch, temperature, pressure)
6. **Visceral pain:** Abdominal, chest, and pelvic pain from organ system dysfunction
7. **Central sensitization pain:** The nervous system amplifies all pain signals

Unlike acute pain, which signals a specific injury and resolves with healing, ME/CFS pain is **chronic, unrelenting, and poorly responsive to analgesics**. Opioids carry significant risks. NSAIDs provide minimal relief. The pain simply continues, month after month, year after year.

5.3.5 Cognitive Devastation

Beyond “Brain Fog”

The term “brain fog” dramatically understates the cognitive destruction caused by severe ME/CFS. What patients experience is closer to **acquired brain injury**—the progressive failure of cognitive functions that were previously intact.

Loss of Language

- **Word-finding difficulties:** Cannot retrieve common words
- **Sentence construction fails:** Cannot organize thoughts into coherent expression
- **Reading comprehension loss:** Words on a page no longer form meaning
- **Writing disability:** Cannot compose text, even simple messages
- **Language processing:** Cannot understand speech, especially rapid or complex

Memory Destruction

- **Short-term memory failure:** Cannot remember what happened minutes ago
- **Working memory collapse:** Cannot hold multiple items in mind simultaneously
- **Prospective memory loss:** Cannot remember to do things in the future
- **Long-term memory erosion:** Older memories become inaccessible or confused

Executive Function Collapse

- **Cannot plan:** Even simple sequences become impossible to organize
- **Cannot decide:** Decision-making exhausts cognitive resources
- **Cannot initiate:** Even with capacity, cannot begin tasks
- **Cannot inhibit:** Poor impulse control, emotional dysregulation
- **Cannot shift:** Rigid thinking, unable to change approach

Processing Speed

- **Dramatic slowing:** Thoughts that took milliseconds now take seconds or minutes
- **Cannot keep pace:** Conversations, events, information move too fast
- **Delayed responses:** Long pauses before being able to respond
- **Mental “blank-outs”:** Complete cessation of cognitive activity

The Loss of Self

For many severe patients, the cognitive devastation amounts to a **loss of personal identity**:

- **Cannot engage in former interests:** Reading, hobbies, intellectual pursuits—all inaccessible
- **Cannot maintain relationships:** Too impaired to communicate, remember, or connect
- **Cannot recognize themselves:** The person they were is gone, replaced by a shadow
- **Memories fade:** Even the past becomes uncertain as long-term memory erodes

This is not depression (though depression often co-occurs). This is **organic brain dysfunction**—the brain, starved of adequate energy and bathed in inflammatory signals, simply cannot perform its functions. The 2024 NIH deep phenotyping study found abnormally low catecholamines (dopamine, norepinephrine) in cerebrospinal fluid and reduced activity in the temporoparietal junction—the brain region responsible for effort-based decision-making and sensory integration [134].

5.3.6 The Wish for Death

Suicidality in ME/CFS

The level of suffering in severe ME/CFS is so extreme that many patients contemplate, attempt, or complete suicide. Research documents this tragic reality:

- **Suicide risk:** 6.85 times higher than the general population [52]
- **Suicidal ideation:** 39–57% of moderately to severely ill patients have contemplated suicide [135]
- **Suicide rate in ME/CFS patients:** 12.75% at risk vs. 2.3% in general population
- **Age at suicide:** Average 39.3 years (vs. 48 years in general population)—dying younger
- **Cause of death:** Suicide accounts for approximately 25% of ME/CFS deaths

Observation 30 (Suicide Without Depression in ME/CFS). **60% of ME/CFS patients who died by suicide had no diagnosis of depression** [52].

This statistic is crucial. It demonstrates that ME/CFS suicides are not primarily the result of psychiatric illness—they are **rational responses to unbearable physical suffering** that the medical system has failed to treat or even acknowledge.

Why Patients Want to Die

The desire for death in severe ME/CFS arises from a specific constellation of factors:

1. **Unremitting suffering:** Pain, exhaustion, and neurological dysfunction that never stops, 24/7/365, for years or decades
2. **No prospect of improvement:** Unlike cancer patients who may hope for remission, severe ME/CFS patients face a disease with no approved treatments and poor prognosis for recovery
3. **Progressive worsening:** Many patients watch themselves deteriorate over time, losing function after function, with no floor to the decline
4. **Total isolation:** Cut off from all human connection, entertainment, and engagement by their neurological sensitivities
5. **Medical abandonment:** Dismissed, disbelieved, and denied care by healthcare systems that don't understand or acknowledge their disease
6. **Loss of self:** The person they were has been destroyed; what remains is a suffering body without the cognitive capacity to even find meaning in that suffering
7. **Burden on others:** Watching loved ones sacrifice their lives as caregivers while being unable to reciprocate or even express gratitude adequately
8. **No end in sight:** The suffering could continue for decades—there is no natural endpoint, no finish line

Samuel, explaining his decision to pursue euthanasia at age 21, wrote:

“So bad that you often think there is only one option left. Many see no way out; the suicide rate is extremely high. My condition is also heading in a direction where I may need to be artificially fed.

Therefore, I am taking advantage of assisted dying in 12 days.

But my death should not be in vain.”

Assisted Dying and ME/CFS

In jurisdictions where assisted dying is legal (Belgium, Netherlands, Switzerland, Canada, and others), ME/CFS patients have increasingly sought this option as the only escape from their suffering. This is not evidence of psychiatric illness requiring prevention—it is evidence of a medical system that has failed to provide any other form of relief.

The medical-ethical questions are profound:

- If a disease causes suffering worse than terminal cancer, with no approved treatments and no prospect of relief, is assisted dying a reasonable option?
- Should society invest in preventing ME/CFS suicides by forcing patients to continue suffering, or by actually treating their disease?
- What does it say about our healthcare system that death has become the preferred treatment for hundreds of thousands of patients?

5.3.7 Impact on Caregivers and Families

The Hidden Victims

Severe ME/CFS does not only destroy the patient—it devastates everyone around them. A 2022 international survey of 1,418 patient-family pairs found extraordinary levels of caregiver distress [136]:

- 96.1% of family members felt worried
- 93% experienced frustration
- 92.9% experienced sadness
- 91.8% reported family activities were affected
- 85.3% experienced problems with holidays
- 77.3% felt finances were impacted
- 72.9% reported their sex life was affected

For very severe patients, the caregiver burden is catastrophic:

- Round-the-clock care required (all but one of 47 very severe patients needed 24/7 care)
- Caregivers spent more than 40 hours per week on care
- Caregivers reported enormous impacts on their own health, finances, and social life

- Many caregivers develop their own health problems from the stress and physical demands

Families Torn Apart

ME/CFS destroys families in multiple ways:

1. **Marriages collapse:** The strain of caring for a severely ill spouse while managing household, possibly children, and often working, is unsustainable. Divorce rates are elevated.
2. **Children suffer:** Children of ME/CFS patients grow up with an absent or incapacitated parent. Some children develop ME/CFS themselves after viral illnesses.
3. **Parents sacrifice everything:** Parents of young ME/CFS patients often quit jobs, exhaust savings, and destroy their own health trying to care for their children.
4. **Siblings are neglected:** Family resources—emotional, financial, time—flow to the sick member, leaving healthy siblings feeling abandoned.
5. **Extended family withdraws:** Unable to understand the disease, extended family members often drift away or actively blame the patient for being “lazy” or “making it up.”

One particularly devastating pattern involves **intergenerational ME/CFS**—a parent becomes ill, then years later their child also develops the disease after a viral infection. A 2024 article documented a family where the mother had been largely bedbound for decades, and then her child joined her in isolation after developing ME/CFS. The father now cares for two bedridden family members, watching his wife and child exist in darkness and silence.

The Caregiver's Impossible Position

Caregivers of severe ME/CFS patients face an impossible situation:

- **Cannot help:** There are no effective treatments to offer
- **Cannot comfort:** Physical presence, touch, or conversation cause harm
- **Cannot reduce suffering:** The suffering continues regardless of caregiver efforts
- **Cannot have respite:** The patient cannot be left alone; cannot go to facility care
- **Cannot plan:** The unpredictable nature of crashes makes scheduling impossible
- **Cannot maintain own life:** Work, relationships, health—all sacrificed to caregiving
- **Cannot talk about it:** Society doesn't understand; support groups for ME/CFS caregivers barely exist
- **Cannot stop:** Abandoning the patient means condemning them to institutionalization or death

5.3.8 Economic Devastation

Individual Financial Ruin

ME/CFS causes financial devastation at the individual level:

- **Inability to work:** Up to 75% of ME/CFS patients are unable to work
- **Job loss:** 26–89% lose their jobs due to the illness
- **Unemployment:** 58.6% unemployed in one large study
- **Downward mobility:** Among those who can work part-time, many move to lower-wage positions
- **Lost income:** Average per-person cost for lost income: \$27,880 annually
- **Disability denial:** SSDI approval rates below 20% despite severity comparable to MS

The financial trajectory is typically:

1. Reduced work hours as symptoms develop
2. Loss of job when unable to maintain even reduced schedule
3. Exhaustion of savings during (often lengthy) diagnostic process
4. Denial of disability benefits (claims rarely approved initially)
5. Appeals process taking years while patient has no income
6. Dependence on family members, charity, or destitution

Many patients describe becoming “financial hostages” to family members, partners, or government systems that doubt their illness and treat financial support as conditional on compliance with harmful treatments (like graded exercise therapy).

Healthcare Costs

Paradoxically, a disease that receives minimal research funding and has no approved treatments still generates enormous healthcare costs:

- **Diagnostic odyssey:** Years of specialist appointments, tests, and procedures before receiving diagnosis
- **Out-of-pocket treatments:** Patients pay for supplements, alternative therapies, and off-label medications not covered by insurance
- **Emergency care:** Crashes, orthostatic events, and complications require emergency visits
- **Comorbidities:** POTS, MCAS, fibromyalgia, and other comorbid conditions require ongoing treatment
- **Caregiving costs:** Professional caregiving, when available, is expensive; informal caregiving represents massive unpaid labor

Societal Economic Burden

The total economic burden of ME/CFS is staggering:

- **United States:** \$36–51 billion annually in direct and indirect costs
- **European Union:** Approximately €40 billion annually for 2 million affected citizens
- **United Kingdom:** £3.3 billion minimum
- **Recent estimates (2025):** Up to \$362 billion annually in the US when accounting for newly recognized prevalence

For context, these figures exceed the economic burden of many diseases that receive far more research funding and public attention. ME/CFS receives approximately \$15 per patient in NIH research funding annually, compared to \$300+ per patient for MS.

5.3.9 Medical Abandonment

The Healthcare Gap

Perhaps no aspect of severe ME/CFS is more enraging than the systematic abandonment of patients by healthcare systems:

- **No ME/CFS specialists:** Most regions have zero physicians with expertise in the disease
- **No treatment guidelines:** Until recently, no evidence-based treatment protocols existed
- **No approved medications:** Not a single FDA-approved drug for ME/CFS
- **No dedicated clinics:** A handful of specialty clinics exist worldwide for millions of patients
- **No training:** Most physicians receive zero education about ME/CFS in medical school

The result is that severely ill patients—the patients most in need of care—often receive no care at all. A study found that many severely affected patients have “become entirely disconnected from statutory healthcare services” [129].

Active Harm from Healthcare

Beyond neglect, healthcare systems often actively harm ME/CFS patients:

1. **Psychiatric misdiagnosis:** Patients labeled as depressed, anxious, or somatizing, leading to inappropriate treatment
2. **Graded exercise therapy (GET):** For decades, guidelines recommended increasing exercise—a treatment that worsens most patients and can cause permanent harm to severe patients
3. **Cognitive behavioral therapy (CBT):** Promoted as treatment for a “false illness belief,” implying the disease isn’t real

4. **Forced institutionalization:** Some severe patients have been forcibly removed from homes and placed in psychiatric facilities or nursing homes where their needs cannot be met
5. **Tube feeding refusal:** As documented in malnutrition cases, healthcare providers refuse life-saving nutritional support because they attribute swallowing difficulties to psychological causes
6. **Accusation of Munchausen's/factitious disorder:** Parents of children with ME/CFS have been accused of fabricating or inducing illness, leading to child protective services involvement

Why Severe Patients Cannot Access Care

Even when healthcare providers want to help, severe patients face insurmountable barriers:

- **Cannot travel:** Too ill to be transported to medical facilities
- **Cannot tolerate clinical environment:** Lights, sounds, activity of a hospital or clinic trigger crashes
- **Cannot communicate:** Too weak to describe symptoms or answer questions
- **Crashes from appointments:** Even home visits cause symptom exacerbation
- **No home visit services:** Most healthcare systems don't offer adequate home-based care
- **Insurance barriers:** Home visits, when available, often not covered

Recommendations for compassionate home-based care exist [129]:

- Schedule visits after midday (patients have irregular sleep)
- Keep visits brief; address only one or two issues
- Plan for post-visit recovery time (PEM may last days)
- Avoid perfumes and fragrances
- Maintain low tone of voice
- Believe patient reports

But most healthcare systems ignore these recommendations, and most severely ill patients simply go without medical care.

5.3.10 A Call to Action

This Must Change

The information presented in this chapter should provoke moral outrage. Millions of people worldwide are experiencing suffering that exceeds cancer, dying decades early, choosing euthanasia because no other relief exists, and being abandoned or actively harmed by healthcare systems.

This is a medical emergency that has been ignored for decades.

What Must Happen

1. **Massive research funding:** ME/CFS research funding must increase by orders of magnitude. The NIH currently spends approximately \$15 per patient annually on ME/CFS research. For comparison, HIV/AIDS receives over \$2,500 per patient.
2. **Medical education:** Every physician must receive training on ME/CFS recognition and management. The disease affects 1% of the population—more than MS, more than HIV, more than many conditions that receive extensive medical education.
3. **Specialized care centers:** Every region needs accessible ME/CFS specialty clinics with expertise in the disease, including capacity for home visits to severe patients.
4. **Drug development:** Pharmaceutical companies must be incentivized to develop treatments. The market is huge—millions of patients desperate for any relief—but regulatory pathways and research infrastructure are inadequate.
5. **Social support:** Disability systems must recognize ME/CFS as the devastating illness it is. Patients should not have to fight years-long legal battles while destitute to receive benefits.
6. **Caregiver support:** Family caregivers need respite, financial support, and recognition for the enormous burden they bear.
7. **Public awareness:** Society must understand that ME/CFS is not “chronic fatigue”—being tired. It is a catastrophic multi-system disease that destroys lives.

The Urgency

Every day that passes without adequate response to this crisis:

- Patients die—from suicide, from cardiac events, from malnutrition
- Patients worsen—the 25% who are severe were once mild or moderate; every day more patients cross the threshold into severe disease
- Patients suffer—in darkness and silence, alone, abandoned, in pain that doesn’t end
- New patients develop ME/CFS—viral infections continue to trigger new cases; Long COVID has added millions to the patient population

Samuel chose to die at 21 rather than continue living with very severe ME/CFS. His final message was: “**ME/CFS kills!**”

He was right. And until the medical establishment, governments, and society take this disease seriously, ME/CFS will continue to kill—slowly through suffering, quickly through suicide, and invisibly through the quiet disappearance of patients into bedrooms from which they never emerge.

Final Message

If you have read this chapter, you now understand what severe ME/CFS means. You cannot claim ignorance.

What will you do with this knowledge?

For healthcare providers: Will you educate yourself? Believe your patients? Advocate for research and resources?

For policymakers: Will you fund research? Create support systems? Hold healthcare systems accountable?

For family and friends: Will you learn about your loved one's illness? Provide appropriate support? Advocate on their behalf?

For the general public: Will you spread awareness? Challenge misconceptions? Support organizations working on ME/CFS?

The patients cannot speak for themselves. They are trapped in dark, silent rooms, too weak to advocate, too ill to be seen.

They need you to speak for them.

5.4 Disease Progression

ME/CFS is not a static condition. Understanding how the disease evolves over time—including natural history, relapse patterns, and factors that influence trajectory—is essential for patient counseling, treatment planning, and research design.

5.4.1 Natural History

A five-stage model describes the typical progression of ME/CFS from predisposition through established disease [70]:

Stage 1: Predisposition. Before illness onset, certain individuals carry increased vulnerability due to:

- Genetic factors affecting immune function, metabolism, and stress response
- Prior infections that may have primed abnormal immune responses
- Environmental exposures (toxins, mold, chronic stressors)
- Female sex (women are affected 3–4 times more frequently than men)

This stage is invisible—individuals function normally but carry latent susceptibility.

Stage 2: Trigger and Pre-Illness (0–4 months). A triggering event initiates the disease process. In post-infectious cases, this is the acute infection. In gradual-onset cases, the trigger may be:

- Cumulative infectious burden
- Major physiological stress (surgery, trauma, childbirth)
- Severe psychological stress
- Environmental exposure
- Unknown factors

During this period, non-specific symptoms emerge: fatigue, malaise, and incomplete recovery from the triggering event.

Stage 3: Prodromal Period (4–24 months). The characteristic ME/CFS symptom complex develops:

- Fatigue becomes unrelenting rather than episodic
- Post-exertional malaise emerges as a defining feature
- Sleep becomes unrefreshing regardless of duration
- Cognitive impairment (brain fog) becomes noticeable
- Orthostatic intolerance may develop

Patients during this period often cycle through multiple medical specialists seeking diagnosis, frequently receiving incorrect diagnoses or being told nothing is wrong.

Stage 4: Early Disease (6 months–2 years). The disease becomes established, with:

- Full expression of neuro-immune dysfunction
- Hypermetabolic state with inefficient energy production
- Elevated pro-inflammatory markers in some patients
- Ongoing immune activation
- Significant functional impairment

During early disease, the biological processes driving ME/CFS are active and potentially modifiable. This may represent a window for intervention, though effective treatments remain elusive.

Stage 5: Established Disease (2+ years). Chronic neuro-inflammation and metabolic dysfunction become entrenched:

- Inflammatory markers may normalize despite ongoing dysfunction
- Epigenetic changes alter gene expression patterns
- Immune exhaustion develops (particularly CD8+ T cell exhaustion)
- Brain changes become visible on advanced imaging
- Functional impairment stabilizes at reduced level

Established disease may be more difficult to reverse than early disease, though this remains speculative given the lack of effective treatments.

5.4.2 Patterns of Change

Once ME/CFS is established, patients typically follow one of three trajectories [70]:

Partial Reversal. A minority of patients (primarily those with mild disease and short illness duration) experience gradual improvement:

- Slow, incremental gains in function over years
- Expansion of the energy envelope
- Reduced frequency and severity of post-exertional malaise
- Improved but rarely complete recovery

True complete recovery is rare in adults (see Section 5.5).

Persistence. The most common pattern: chronic stable illness with periodic fluctuations:

- Baseline functional level remains relatively constant
- Good days and bad days within a predictable range
- Relapses triggered by overexertion, infections, or stress
- Recovery to baseline after relapses (usually)
- No net improvement or deterioration over years

This pattern characterizes the majority of mild to moderate ME/CFS patients.

Progressive Worsening. A significant minority of patients experience ongoing decline:

- Each relapse leaves them at a lower functional level
- Progression from mild to moderate to severe
- Accumulation of additional symptoms and comorbidities
- Increasing disability and care needs
- Risk of very severe ME/CFS

Factors associated with progressive worsening include repeated overexertion, inadequate rest, intercurrent infections, and possibly biological factors not yet understood.

5.4.3 The Preventable Descent to Severe Disease: Critical Warning

△ Warning: CRITICAL WARNING: The Point of No Return

Approximately 25% of all ME/CFS patients become housebound or bedbound with severe or very severe disease. Most of these patients started with mild or moderate illness. The progression from mild to severe is often preventable, but it requires understanding the mechanisms of deterioration and acting decisively before crossing irreversible thresholds.

This section may save your life or prevent decades of severe disability.

If you currently have mild or moderate ME/CFS, this is the most important section in this document for you to read, understand, and act upon. The patients described in Section 5.3—those existing in darkness and silence, unable to speak, unable to eat, choosing death over continued suffering—did not start there. They started where you are now.

The difference between remaining functional and becoming bedbound often comes down to decisions made in the first 2–3 years of illness.

The Ratchet Effect: How Decline Becomes Irreversible

Progressive worsening in ME/CFS follows a characteristic pattern known as the “**ratchet effect**”: each crash or period of overexertion moves the baseline functional capacity downward, and unlike a temporary relapse, the patient does not fully return to their previous level. Over time, this creates a stepwise descent from mild to moderate to severe disease.

The Descent Pattern.

1. Initial Phase (Mild Disease):

- Patient can work/study, though with significant difficulty
- Post-exertional malaise occurs but recovery takes days to weeks
- Energy envelope is reduced but still allows meaningful activity
- Patient appears functional to outsiders

2. Denial and Push-Through Phase:

- Patient continues normal or near-normal activity level
- Reasons include: financial necessity, hope for improvement, lack of understanding of PEM, medical advice to “stay active”
- Crash-recovery cycles become routine: push during week, collapse on weekends
- Each recovery is slightly less complete than the last

3. Accelerating Decline (Transition to Moderate/Severe):

- Crashes become more frequent and more severe
- Recovery time extends from days to weeks to months
- Activities that previously caused no problems now trigger PEM

- New symptoms emerge: sensory sensitivities, orthostatic intolerance, cognitive deterioration
- Energy envelope shrinks progressively

4. Point of No Return (Severe Disease):

- Patient can no longer recover to previous baseline regardless of rest
- Minimal activities (showering, brief conversation, sitting upright) trigger severe PEM
- Hypometabolic state becomes established (cellular/mitochondrial damage)
- Patient becomes housebound or bedbound
- Severe disease may be irreversible even with aggressive intervention

⚠ Limitation 3: Ratchet Effect and Cumulative Damage: Observational Basis Without Controlled Evidence

The “ratchet effect,” “crash limit rule,” and cumulative damage model described in this section are derived from patient self-report, community observation, and mechanistic plausibility—not from prospective studies measuring biological damage markers across crashes. Key epistemic boundaries:

- No study has measured mitochondrial, endothelial, or neuroinflammatory damage longitudinally across PEM episodes in the same patients; the claim that “each crash causes cumulative damage” is biologically plausible but empirically untested.
- The “5–10 severe crash” threshold is anecdotal community observation, not a validated biological boundary; individual vulnerability likely varies enormously based on factors not yet characterised.
- The “point of no return” framing, while motivating for pacing adherence, risks overstating certainty about irreversibility; some patients have recovered from severe ME/CFS, suggesting the threshold is not absolute.
- The strong causal language (“your choices determine your trajectory”) should be interpreted as harm-reduction messaging rather than established causal knowledge; patients who become severe despite careful pacing exist, and disease progression has non-modifiable biological determinants.

The Cumulative Damage Model. Research and patient reports suggest that **repeated episodes of post-exertional malaise cause cumulative physiological damage** [74, 70]. While individual crashes may appear to resolve, each episode may contribute to progressive deterioration through vicious cycle mechanisms. These pathophysiological systems both contribute to PEM susceptibility and are further damaged by PEM episodes themselves, creating self-reinforcing feedback loops:

- **Mitochondrial dysfunction accumulation** (Section 6.2): Impaired energy metabolism increases PEM vulnerability, while repeated ATP depletion and oxidative stress during crashes further damage mitochondrial membranes and DNA
- **Endothelial dysfunction** (Section 10.2.1): Baseline vascular impairment limits oxygen delivery, while each PEM episode involves additional vascular stress; repeated insults

progressively impair vessel reactivity

- **Neuroinflammation** (Section 7.4.2): Pre-existing neuroinflammation lowers the threshold for symptom exacerbation, while repeated microglial activation during crashes perpetuates chronic neuroinflammatory states
- **Immune exhaustion** (Section 7.4.1): Baseline immune dysfunction increases infection risk (a common PEM trigger), while chronic activation during crashes progressively depletes immune cell populations and function
- **Metabolic state transition** (Chapter 6): Progression from hypermetabolic (early, potentially reversible) to hypometabolic (established, potentially irreversible) state, with each crash potentially driving the transition toward the irreversible hypometabolic phenotype

Observation 31 (PEM as Progressive Central Sensitization). Patient-derived clinical observations suggest that post-exertional malaise may operate as a progressive sensitization mechanism analogous to chronic pain sensitization rather than simple fatigue fluctuation. Each PEM episode appears to lower the threshold for subsequent crashes: activities that previously triggered 2–3 days of symptoms may eventually trigger 2–3 weeks of incapacity. This pattern parallels microglial sensitization models in pain neurobiology, where repeated glial activation progressively lowers the neuroinflammatory threshold. The observed progression from crashes requiring days of recovery (early disease) to crashes requiring weeks or months (established disease) suggests cumulative sensitization of the neuroimmune system, where repeated PEM episodes condition the microglial response to future activity. This observation supports the mechanistic model that preventing crashes entirely—rather than managing crashes once they occur—may be the primary intervention preventing irreversible transition to severe disease.

Observation 32 (The “Crash Limit” Concept). Patient communities have observed what is sometimes called the “**crash limit rule**”: there appears to be a threshold number of severe crashes (anecdotally reported as approximately 5–10 major crashes) beyond which recovery capacity is permanently impaired. While this specific threshold lacks formal research validation, the underlying principle is biologically plausible and aligns with cumulative damage models.

Key observations:

- Recovery time from crashes increases with each successive crash
- After a certain number of severe crashes, patients stop recovering to previous baseline
- Patient community reports describe cases where pushing through symptoms resulted in prolonged illness with extended recovery times from subsequent crashes
- Some patients report that a single catastrophic overexertion event (a marathon, a stressful life event combined with overwork, a severe infection while already depleted) triggered irreversible worsening
- **Infection as cascade trigger**: Post-infectious deterioration (COVID, influenza) commonly causes step-down in baseline function, with each subsequent infection producing longer PEM recovery periods

Case example: A patient who managed mild/moderate ME/CFS for over a decade (while raising children as a single parent) experienced COVID infection in autumn 2024 followed

by influenza in early 2025. PEM recovery time progressed from the previous pattern of 2–3 days (with occasional 3–4 week recoveries after major exertion) to a new baseline of 2–3 weeks minimum, often longer. This patient now requires wheelchair use and can only perform minimal activities with frequent rest breaks. This illustrates how infections can trigger the ratchet effect, with each infection driving irreversible functional decline.

Implication: Every severe crash matters. The goal is not to minimize crashes—it is to *avoid them entirely*.

Critical Warning Signs: You Are Approaching Severe Disease

If you experience ANY of the following, you are at immediate risk of progression to severe disease and must take aggressive action:

• Requirement 1: RED FLAGS: Stop Everything and Implement Emergency Pacing

Immediate Danger Signs (Act Within Days):

- **Unable to recover baseline within 2 weeks after a crash:** If you used to recover in days and now it takes weeks, your reserve capacity is failing
- **Bedbound on weekends to survive work week:** This is not sustainable—you are causing progressive deterioration
- **Crashes triggered by activities that didn't cause problems 6 months ago:** Your energy envelope is shrinking actively
- **New sensory sensitivities emerging:** Light sensitivity, sound sensitivity, chemical sensitivities indicate neurological sensitization is establishing
- **Orthostatic intolerance developing or worsening:** Cannot stand for normal activities, heart rate increases >30 bpm upon standing
- **Cognitive symptoms worsening:** Word-finding difficulties, memory problems, inability to read/process information (cognitive symptoms appear most resistant to recovery) [74]
- **Weight loss from inability to prepare food:** Eating has become too effortful; this indicates severe energy depletion
- **Social withdrawal not by choice but by necessity:** Cannot tolerate visitors, phone calls, any social interaction

Urgent Concern Signs (Act Within Weeks):

- **Symptoms persisting >6 months without any improvement:** Indicates transition from acute to established aberrant homeostatic state [70]
- **Multiplying food intolerances/sensitivities:** Mast cell activation worsening
- **Sleep becoming more disturbed despite medications:** Central nervous system dysfunction progressing
- **Pain increasing in severity and distribution:** Central sensitization establishing

- **Temperature regulation failing:** Severe chills or overheating from minor environmental changes
- **Post-exertional malaise severity increasing:** What used to cause 2 days of PEM now causes 2 weeks

Pattern Recognition (Monitor Over Months):

- **Ratcheting baseline:** Each crash leaves you slightly worse; baseline is trending downward over 6–12 months
- **Energy envelope shrinking:** Activities that were within your envelope 6 months ago now exceed it
- **Recovery time lengthening:** Crashes that took 3 days to recover from now take 3 weeks
- **Boom-bust cycles intensifying:** The “bust” phases are becoming deeper and longer

The 6-Month Rule and the First 2 Years. Research identifies two critical temporal thresholds:

1. **6-month persistence mark [70]:** If symptoms persist beyond 6 months without improvement, this indicates that normal homeostatic recovery mechanisms have failed and aberrant pathophysiology is becoming established. This is the transition from “post-viral fatigue that might resolve” to “ME/CFS that likely won’t resolve without intervention.”
2. **2-year establishment threshold [70]:** The natural history model suggests that around 2 years, the disease transitions from early (hypermetabolic, potentially modifiable) to established (hypometabolic, potentially entrenched). This involves:
 - Epigenetic changes altering gene expression
 - Immune exhaustion (CD8+ T cell exhaustion, NK cell dysfunction)
 - Normalization of inflammatory markers despite ongoing dysfunction
 - Brain changes visible on advanced imaging
 - Metabolic state shift from high (inefficient) energy expenditure to low energy production

Implication: The first 2 years represent a critical intervention window. Aggressive pacing and early treatment during this period may prevent progression to established severe disease. After 2 years, reversal becomes substantially more difficult.

The Psychological Trap: When Hope and Denial Cause Harm

One of the most dangerous aspects of ME/CFS progression is the **psychological trap** that keeps patients pushing beyond their limits even as they deteriorate:

The Denial Mechanisms.

- **“It’s just a bad week”**: Minimizing the significance of worsening symptoms
- **“I can’t afford to stop working”**: Financial pressure overriding physiological reality
- **“If I just push through this busy period, I can rest later”**: Future rest never comes; busy periods are continuous
- **“I’m not as bad as those severe patients”**: Comparing to worst cases rather than recognizing own decline
- **“My doctor says exercise is good for me”**: Trusting outdated medical advice over body signals
- **“I don’t want to give up”**: Misunderstanding that continuing to push IS giving up—giving up on future functional capacity

The Hope Trap. Hope is generally adaptive, but in ME/CFS it can be dangerous:

- **“Maybe I’m getting better”**: Interpreting good days as recovery rather than normal fluctuation, leading to overexertion
- **“This new treatment will cure me”**: Trying experimental interventions while neglecting fundamental pacing
- **“I’ll rest when I recover”**: Not understanding that *rest is required FOR recovery*
- **“I can handle one more thing”**: Incremental additions to activity that cumulatively exceed envelope

The Societal Pressure. External pressure reinforces harmful patterns:

- Family/friends: “You look fine,” “Just try harder,” “Everyone gets tired”
- Employers: Expectation of full productivity despite disability
- Medical system: “It’s just fatigue,” “You’re depressed,” “Exercise more”
- Cultural narratives: “Never give up,” “Mind over matter,” “Winners push through pain”
- Financial systems: Disability denial forcing continued work

★ Key Point: Reframing: Pacing Is Not Giving Up

Stopping is not surrender—it is strategic retreat to preserve future capacity.

- Reducing work hours is not laziness—it is preventing permanent disability
- Declining social events is not depression—it is energy management
- Resting aggressively is not weakness—it is the primary treatment for ME/CFS
- Accepting limitations is not defeat—it is acknowledging biological reality

The patients in Section 5.3 who are now bedbound, unable to speak, existing in darkness—many of them became severe because they “didn’t give up” when they should have. They pushed through. They tried to maintain normal lives. They listened to doctors who told them to exercise. They couldn’t afford to stop working.

Giving up the fight to appear normal is how you preserve the capacity to have an actual life.

How to Prevent Progression: Emergency Action Protocol

If you recognize yourself in the warning signs above, implement this protocol immediately:

Step 1: Immediate Activity Reduction (Within 48 Hours).

1. Stop all non-essential activity:

- Cancel social commitments
- Reduce work hours (request emergency accommodation or medical leave)
- Eliminate hobbies, exercise, entertainment that costs energy
- Minimize cooking (simple foods, meal delivery, family help)

2. Implement aggressive rest:

- Horizontal rest 50–75% of waking hours
- Dark, quiet environment
- No screens during rest periods (true rest, not entertainment)
- Rest *before* feeling exhausted, not after

3. Establish conservative energy envelope:

- 50% rule: Do half of what you think you can manage
- Heart rate monitoring: Stay below 60% maximum heart rate (estimate maximum using 220 minus your age; consider obtaining a heart rate monitor or fitness tracker)
- Activity in 15–25 minute blocks with rest between
- If any activity triggers PEM, eliminate it entirely

Step 2: Medical Documentation and Accommodation (Within 1 Week).

1. Physician visit:

- Document worsening symptoms
- Request medical leave or work restriction letter
- Obtain disability parking permit if orthostatic intolerance present
- Discuss symptom management medications

2. Workplace/school accommodation:

- Formal request for reduced hours (50–75% time)
- Remote work to eliminate commute
- Flexible schedule for peak energy periods
- If accommodations denied or insufficient: apply for disability leave

3. Financial planning:

- Apply for short-term disability if available
- Begin long-term disability application process (often 3–6 month wait)
- Investigate government disability benefits (SSDI, equivalent)
- Reduce expenses where possible

Step 3: Baseline Stabilization (Weeks to Months).

1. **Goal:** Establish 4–8 weeks with *zero PEM episodes*
 - This proves you are within your energy envelope
 - Stabilization allows baseline to stop declining
 - During this period, accept that your functional capacity is very low
2. **Monitoring:**
 - Daily symptom log (0–10 scale for fatigue, pain, cognition)
 - Activity log with durations
 - PEM tracking (onset, duration, triggers)
 - Heart rate data if using monitor
3. **Adjustment:**
 - If PEM occurs: reduce activity further (you exceeded envelope)
 - If no PEM for 4 weeks: maintain current level (do NOT increase yet)
 - If symptoms improving after 8 weeks stable: consider 5–10% activity increase

Step 4: Long-Term Vigilance (Ongoing).

1. **Permanent pacing:**
 - Energy envelope management is not temporary—it is ongoing disease management
 - Even if symptoms improve, maintain conservative approach
 - Always operate at 70–80% of perceived capacity (reserve for unexpected demands)
2. **Infection prevention:**
 - Infections reliably trigger relapse and can cause permanent worsening
 - Masking in public during viral season
 - Avoid crowded indoor spaces
 - Vaccinations (though some patients experience temporary PEM post-vaccination)
3. **Reassessment every 3–6 months:**
 - Is baseline stable, improving, or worsening?
 - Are PEM episodes eliminated or still occurring?
 - Is current activity level sustainable long-term?
 - Do accommodations need adjustment?

△ **Warning: When to Consider Emergency Disability Application**

If despite aggressive pacing you continue to worsen, or if you are already experiencing severe symptoms, **stop working entirely and apply for disability immediately**. The financial consequences of disability application are reversible; the physiological consequences of pushing into severe ME/CFS are not.

Specific thresholds for work cessation:

- Bedbound >50% of weekend days recovering from work week
- New symptoms emerging (sensory sensitivities, swallowing difficulties, severe cognitive impairment)
- Requiring assistance with activities of daily living (cooking, hygiene, shopping)
- Suicidal ideation related to symptom burden
- Medical professional recommendation to stop working

Working yourself into severe ME/CFS means you cannot work AND you are severely disabled. Stopping work while still moderate means you might prevent severe disease and potentially return to some work capacity in the future.

The Evidence: Can Aggressive Pacing Prevent Severe Disease?

While randomized controlled trials of aggressive early pacing do not exist (such trials would be unethical, requiring a control group to continue overexertion), multiple lines of evidence support the preventive value of energy envelope management:

Observational Evidence.

- **Diagnostic delay predicts worse outcomes** [77]: Patients diagnosed and instructed in pacing early have better long-term function than those diagnosed after years of pushing through symptoms
- **Patient survey data** [74]: 90% of patients identified “designing and monitoring their own management plan” (pacing) as helpful; graded exercise therapy reported as harmful by 50–70%
- **Energy envelope theory**: Patients who stay within their energy envelope show reduced symptom severity and improved quality of life compared to those who regularly exceed limits
- **Pediatric outcomes** [78]: Children with ME/CFS show 68% recovery rates by 10 years when supported with flexible educational accommodations (allowing rest), versus <5% recovery in adults (who typically continue pushing)

Mechanistic Plausibility. The biological mechanisms documented in Chapters 6 through 13 support the cumulative damage model:

- **Mitochondrial damage from repeated ATP depletion** (Section 6.2): Each PEM episode involves cellular energy crisis; repeated crises accumulate damage

- **Oxidative stress accumulation** (Section 6.4): Exertion triggers reactive oxygen species production; inadequate recovery allows oxidative damage to accumulate
- **Endothelial dysfunction from repeated ischemia-reperfusion** (Section 10.2.1): Each PEM episode involves impaired blood flow; repeated insults cause permanent vascular changes
- **Neuroinflammation from repeated microglial activation** (Section 7.4.2): Chronic activation leads to permanent neurological sensitization
- **Immune exhaustion from chronic activation** (Section 7.4.1): Prolonged immune activation depletes cell populations and function

Preventing repeated PEM episodes theoretically prevents or reduces cumulative damage in all these systems.

The Counterfactual Argument. We know what happens when patients do NOT pace aggressively:

- 25% become housebound/bedbound (Section 5.3)
- Many report that continued overexertion preceded their progression to severe disease
- Graded exercise therapy—the antithesis of pacing—causes deterioration in 50–70% of patients
- Patient communities uniformly identify “push-crash cycles” as the primary cause of worsening

While we cannot prove aggressive pacing prevents severe disease, we have strong evidence that failure to pace causes severe disease.

Summary: Your Choices Determine Your Trajectory

★ Key Point: Key Takeaways: Preventing the Descent

What we know:

- 25% of ME/CFS patients become severely ill
- Most severe patients started with mild or moderate disease
- Repeated overexertion (push-crash cycles) precedes progression in many cases
- The first 2 years represent a critical intervention window
- Recovery becomes progressively harder with illness duration and severity
- There may be a threshold beyond which severe disease becomes irreversible

What you can control:

- Your activity level: Stay within energy envelope, implement 50% rule
- Your response to warning signs: Act immediately when symptoms worsen
- Your work/life boundaries: Request accommodations, reduce hours, stop if neces-

sary

- Your acceptance of limitations: Acknowledge reality rather than push through denial
- Your prevention of infections: Reduce exposure to avoid relapse triggers

What you cannot control:

- Your baseline disease severity (biological factors, genetic susceptibility)
- Whether you will recover (some do, most don't, reasons unknown)
- External pressures (financial, social, medical system failures)

The decision framework:

Every time you consider exceeding your energy envelope—working extra hours, attending a social event, “pushing through”—ask yourself:

“Am I willing to risk permanent severe disability for this activity?”

Because that is the actual risk. Not “I’ll be tired tomorrow.” Not “I’ll have a bad week.” The risk is: **this crash might be the one that tips me into irreversible severe disease.**

The patients existing in darkness and silence (Section 5.3) did not know which crash would be their last. They did not know when they crossed the point of no return. They only knew, in retrospect, that they had crossed it.

You still have choices. They no longer do. Act accordingly.

5.4.4 Relapse and Remission

ME/CFS is characterized by fluctuating symptoms with periods of relative stability punctuated by relapses.

Triggers for Relapse. The most common triggers for symptom exacerbation include:

- **Physical exertion:** Even minor activity exceeding the energy envelope
- **Cognitive exertion:** Sustained mental effort, decision-making, emotional processing
- **Infections:** Viral, bacterial, or fungal infections reliably trigger relapse
- **Sleep disruption:** Inadequate sleep or disrupted sleep patterns
- **Environmental factors:** Temperature extremes, sensory overload, travel
- **Medical procedures:** Surgery, dental work, vaccinations
- **Emotional stress:** Acute psychological stressors

The delayed onset of post-exertional malaise (typically 12–48 hours after the triggering activity) makes cause-and-effect relationships difficult to identify without careful tracking.

Characteristics of Relapse. During relapse, patients experience:

- Intensification of baseline symptoms
- Emergence of symptoms not usually present at baseline
- Reduced functional capacity
- Increased sensitivity to sensory input
- Cognitive impairment worsening
- Duration ranging from days to months

Recovery from Relapse. Recovery from relapse requires:

- Aggressive rest (reducing activity well below baseline)
- Identification and elimination of triggering factors
- Time (often weeks even for minor relapses)
- Patience and acceptance that recovery cannot be rushed

Most patients return to their previous baseline after relapse, though repeated relapses or severe relapses may result in a new, lower baseline (the “ratchet effect”).

Remission. True remission—a sustained period of substantially improved function—is uncommon but does occur. Characteristics of remission include:

- Expanded energy envelope and activity tolerance
- Reduced or absent post-exertional malaise
- Improved cognitive function
- Better sleep quality
- Duration of months to years

Remission is fragile. Patients in remission may relapse with infection, overexertion, or other stressors. The possibility of relapse creates ongoing anxiety even during periods of improvement.

5.4.5 Factors Influencing Trajectory

Multiple factors affect whether a patient improves, remains stable, or deteriorates. Importantly, most identified “modifiable factors” reduce to a single underlying mechanism: whether the patient stays within or exceeds their energy envelope.

The Central Modifiable Factor: Energy Envelope Management. Nearly all modifiable factors associated with disease trajectory relate to energy envelope violations:

- **Pacing adherence:** Directly determines whether crashes occur
- **Diagnostic delay:** Patients unaware of their condition spend months or years exceeding their envelope because they don't know to pace [77]
- **Harmful interventions:** Graded exercise therapy is medically-advised envelope violation
- **Financial pressure:** Forces continued activity despite symptoms—envelope violation by economic necessity
- **Social/family pressure:** “Push through it” advice leads to envelope violation

These are not independent risk factors—they are different *causes* of the same harmful outcome (repeated envelope violation). A patient with excellent pacing knowledge but no financial ability to rest will exceed their envelope. A patient with financial security but a physician prescribing GET will exceed their envelope. The mechanism of harm is the same; only the reason differs.

Factors That Enable Envelope Management. Some factors influence trajectory indirectly by enabling or preventing effective pacing:

- **Social support:** Family who understand ME/CFS can take over tasks, reducing activity demands
- **Financial stability:** Ability to reduce work hours or stop working entirely
- **Healthcare access:** Appropriate diagnosis, symptom management, and accommodation documentation
- **Employer flexibility:** Remote work, reduced hours, rest breaks

These factors do not directly affect disease biology—they affect whether a patient *can* stay within their envelope given their life circumstances.

Non-Modifiable Factors.

- **Age at onset:** Younger onset (pediatric/adolescent) associated with better prognosis—possibly reflecting greater biological plasticity or fewer external demands (school accommodations easier than workplace)
- **Illness duration:** Longer duration associated with lower recovery rates
- **Initial severity:** More severe initial presentation may predict worse outcomes
- **Biological vulnerability:** Why does Patient A tolerate repeated crashes and stabilize while Patient B becomes bedbound after fewer insults? Genetic variants, immune profiles, mitochondrial reserve, and metabolic phenotypes likely influence this differential vulnerability, but these factors are not yet characterized or clinically actionable

Factors That Do Not Predict Trajectory. Notably, some factors that might be expected to predict outcomes do not:

- Depression comorbidity (in most studies)
- Baseline fatigue severity alone
- Gender (in adults)
- Onset type (post-infectious vs. gradual) in some studies

The Unanswered Question. The critical question—why some patients progress to severe disease while others with similar behavior stabilize—remains unanswered. The modifiable factors explain *how* patients exceed their envelope, but not why the consequences differ so dramatically between individuals. Two patients with identical crash histories may have vastly different outcomes. This suggests underlying biological heterogeneity that determines resilience versus vulnerability to cumulative damage, but the specific factors remain unknown. Until these biological determinants are identified, the best available strategy is aggressive envelope management to minimize the insults that *might* cause irreversible harm in susceptible individuals.

5.5 Prognosis

Understanding prognosis is essential for patient counseling, treatment planning, and research prioritization. The prognosis of ME/CFS is generally poor in adults, with few patients achieving full recovery. However, outcomes vary considerably by age of onset, illness duration, and other factors.

5.5.1 Recovery Rates

Adult Recovery. Systematic reviews of ME/CFS prognosis consistently show low recovery rates in adults:

- **Full recovery:** Median 5% (range: <5–10%)
- **Improvement:** Median 39.5% (range: 17–64%)
- **No change:** Approximately 40–50%
- **Deterioration:** 10–20% worsen during follow-up

A recent prospective cohort study of 168 ME/CFS patients followed for 20–51 months found [77]:

- Complete recovery: 8.3% (14/168)
- Significant improvement: 4.8% (8/168)
- Combined recovery/improvement: 13.1%

These figures should inform realistic expectations. For adult patients, ME/CFS is typically a chronic, lifelong condition. Improvement is possible but not assured; full recovery is the exception rather than the rule.

Pediatric Recovery. Children and adolescents with ME/CFS have substantially better outcomes than adults [78]:

A landmark long-term follow-up study of 784 young people (mean age at onset 14.8 years) found:

- Recovery at 5 years: 38%
- Recovery at 10 years: 68%
- Mean illness duration: 5 years (range 1–15)
- Mean functional status at 10-year follow-up: 8/10
- Proportion very unwell (<6/10 function) at follow-up: 5%
- Working or studying full-time at follow-up: 63%

The dramatic difference between pediatric (54–94% improve or fully recover) and adult ($\leq 22\%$ improve) outcomes suggests that biological factors related to developmental plasticity may facilitate recovery in young patients, or that adults face barriers to recovery not present in children. For pediatric-specific treatment protocols that leverage this critical intervention window, see Chapters 19 and 20.

⚠ **Limitation 4: Adult Recovery Rate Data: Severe Methodological Constraints**

Published adult ME/CFS recovery rates (commonly cited as $\sim 5\%$) are derived from a small evidence base with severe methodological limitations that cut in both directions—potentially under- or overestimating true rates:

- “Recovery” definitions vary 12-fold across studies (from symptom resolution to self-report); this creates an illusion of precision where none exists, and makes cross-study comparison unreliable.
- All major prognosis studies recruit from tertiary specialty clinics, introducing selection bias of unknown magnitude; population-based estimates do not exist.
- The strongest prognostic predictor—shorter diagnostic delay—is confounded with illness severity, health-seeking behaviour, and socioeconomic status; the independent contribution of each factor is unknown.
- The hypotheses that true recovery potential “may be substantially higher” (Hypothesis 5.5.2) and that adult outcomes “might approach paediatric levels” (Hypothesis 5.5.2) are unfalsified but also untested; they should not be read as evidence that recovery is likely.
- No randomised trial of early aggressive pacing versus standard care has been conducted; the inference that early intervention improves prognosis rests entirely on observational associations subject to confounding.

~ Hypothesis 1: Developmental Plasticity Window

The dramatically better prognosis in pediatric ME/CFS (54–94% improvement) versus adult disease ($\leq 22\%$) suggests that biological factors related to developmental plasticity fundamentally affect recovery potential. We propose that this reflects: (1) ongoing epigenetic reprogramming during development that can override ME/CFS-associated changes, (2) active immune cell turnover that clears dysfunctional cell populations, and (3) metabolic flexibility that allows compensation for mitochondrial dysfunction. This plasticity appears to narrow with age and illness duration, supporting the urgency of early intervention in pediatric cases and suggesting that aggressive early treatment in adult patients may preserve recovery potential.

5.5.2 Evidence That Adult Recovery, While Rare, Does Occur

★ Key Point: Reframing Adult Recovery

The commonly cited ~5% adult recovery rate may substantially underestimate true recovery potential due to measurement artifacts, selection bias, inadequate treatment, and environmental factors that differ systematically from pediatric populations. While full recovery remains uncommon, the evidence limitations are severe enough that the true rate is genuinely uncertain.

The Observed Evidence

Systematic reviews consistently report low recovery rates in adults. The most recent prospective cohort study found complete recovery in 8.3% and significant improvement in 4.8% of 168 patients followed for a median of 5 years [77]. Earlier systematic reviews report median full recovery of 5% (range 0–31%) and median improvement of 39.5% (range 8–63%) [54].

Critically, these figures represent outcomes among patients who:

- Were diagnosed (often years after symptom onset)
- Reached specialty clinics (typically the most severe or treatment-resistant)
- Remained in follow-up (those who recovered may have left care)
- Met stringent diagnostic criteria (ICC 2011 in some studies)

The improvement rate of ~40% is substantially more common than full recovery. Many patients achieve meaningful functional gains—transitioning from severe to moderate, or moderate to mild—without meeting strict recovery criteria. This distinction between “improvement” and “recovery” is clinically important: improvement is achievable for many, even when cure is not.

Limitations of Current Evidence

The 5% recovery rate emerges from studies with severe methodological limitations that may systematically bias estimates.

Definition Inconsistency. “Recovery” is defined inconsistently across studies, creating 12-fold variation in reported rates (5% to 60%). Definitions range from complete symptom resolution without any ongoing interventions (most stringent, <5% meet this criterion) to self-reported “recovery” that permits ongoing activity modification and pacing (least stringent, up to 60% in pediatric studies). When strictly operationalized as complete symptom remission plus return to premorbid function without coping strategies or medications, adult recovery rates are consistently below 5%.

Selection Bias. Most prognosis studies recruit from tertiary specialty clinics, introducing several biases:

- **Survivor bias:** Patients who recover early may never reach specialty care or may leave the healthcare system, systematically excluding recoverers from clinic-based studies
- **Severity bias:** Tertiary centers see the most severe and treatment-resistant cases; milder cases managed in primary care are underrepresented
- **Treatment-seeking bias:** Studies recruit patients “still actively seeking health care” [77], excluding those who have given up or recovered

The magnitude of selection bias is substantial: patient registries designed to capture broader populations report 30% severe-to-very-severe illness, versus <10% in typical clinic studies—suggesting both ends of the severity spectrum are underrepresented.

Diagnostic Delay Confounding. Patients who recover quickly may never receive a formal ME/CFS diagnosis, which requires 6 months of symptoms plus extensive evaluation. The “missing recoverers” problem—those who resolved before diagnosis—is impossible to quantify but may be substantial. If early recovery occurs within the first 1–2 years and average diagnostic delay exceeds this window, published recovery rates measure outcomes only among those who were already chronic.

The Pre-Mechanism Era. All existing prognosis data comes from an era before identification of specific disease mechanisms (NIH 2024 deep phenotyping study), before druggable targets were identified, and before mechanism-based treatments were available. Historical prognosis statistics reflect outcomes with symptomatic management only—analogue to HIV prognosis before antiretroviral therapy. As mechanism-based treatments emerge, historical recovery rates may become less relevant.

Why True Recovery Potential May Be Higher

Several lines of evidence suggest the true adult recovery rate may exceed published estimates.

The Pediatric Proof of Concept. Pediatric ME/CFS shows dramatically better outcomes: 68% recovery at 10 years versus ~5% in adults [78]. This 13-fold difference proves that ME/CFS is not inherently irreversible. If biology alone determined outcomes, pediatric rates would also be ~5%. The gap suggests that modifiable factors—earlier diagnosis, better accommodations, permission to rest, family support—may substantially influence recovery potential.

The Diagnostic Delay Effect. Diagnostic delay is inversely associated with recovery. Patients who recovered had mean diagnostic delay of 23 months versus 55 months for non-recoverers (OR 0.98 per month of delay, $p = 0.036$) [77]. Each year of delay reduces odds of recovery by approximately 2%. The first 2 years may represent a critical window, as recovery after prolonged illness duration becomes increasingly uncommon. If early intervention improves outcomes, the poor adult prognosis may partly reflect late diagnosis rather than inherent chronicity.

The Accommodation Deficit. Adults face structural barriers to recovery that children do not:

- Children receive school accommodations; adults rarely receive workplace accommodations
- Children can stay home; adults must work to survive
- Children have family support; adults are often isolated
- The expectation that “children recover” leads to supportive management; the expectation that “adults don’t recover” may create self-fulfilling nihilism

The pediatric-adult gap may be substantially environmental rather than biological. Adults forced to overexert to maintain employment may be unable to access the rest that facilitates pediatric recovery.

Treatment Response Evidence. Emerging treatment data suggests that subsets of patients respond to targeted interventions:

- Immunoabsorption: 70% clinical response in patients with elevated autoantibodies [118] (suggesting an autoimmune subtype)
- Low-dose naltrexone: 73.9% report positive response in retrospective studies [137]
- MCAS protocols: High response rates reported in patients with mast cell activation
- Antiviral therapy: Responders identified in specific viral reactivation cases

These response rates in selected populations far exceed the 5% spontaneous recovery rate, suggesting that targeted treatments may benefit identifiable patient subgroups—though these represent different study populations than typical prognosis cohorts.

Subtype Heterogeneity. ME/CFS is almost certainly multiple conditions with shared clinical presentation. Averaging recovery rates across subtypes is like averaging cancer survival across all cancer types—the aggregate obscures clinically meaningful variation. The 5% overall rate may hide 20–30% recovery in some subtypes and near-zero in others. Without subtype-specific prognosis data, individual counseling is impossible.

~ **Hypothesis 2: Selection Bias in Published Recovery Rates**

Published adult ME/CFS recovery rates of ~5% may substantially underestimate true recovery potential because: (1) patients who recover early never reach specialty care or leave the system before study enrollment, (2) patients diagnosed late have already passed the critical intervention window, (3) tertiary clinic populations represent the most treatment-resistant cases, and (4) prognosis data predates identification of druggable targets. The true recovery rate among adults diagnosed early with adequate support may be substantially higher—possibly approaching pediatric rates for comparable illness durations and presentations.

Factors Associated with Adult Recovery

Limited evidence suggests the following factors are associated with better outcomes in adults:

- **Shorter illness duration:** Duration <2 years at baseline is the strongest predictor of eventual recovery
- **Shorter diagnostic delay:** Each year of delay reduces recovery odds by ~2%
- **Older age at onset:** Counterintuitively, older age at onset predicts better outcomes (OR 1.06 per year, $p = 0.028$) [77]—mechanism unknown
- **Lower baseline severity:** Milder initial presentation may predict better outcomes, though this finding is inconsistent
- **Ability to rest:** Patients with financial security, disability benefits, or family support enabling genuine rest may have better outcomes (not formally studied)

Notably, sex, onset type (post-infectious vs. gradual), specific symptom profile, and biomarkers have not been validated as prognostic factors—largely because appropriate studies have not been conducted.

Recovery Trajectories

Recovery, when it occurs, typically follows a slow trajectory:

- Gradual improvement over years, not weeks or months

- Non-linear course with setbacks and fluctuations
- Transition through severity levels (severe → moderate → mild) rather than sudden resolution
- Many “recovered” patients continue to pace activities and avoid triggers

The distinction between “recovered” and “improved but managing” is clinically important. Many patients achieve substantial functional gains—returning to work, resuming activities—while still requiring ongoing symptom management. This represents meaningful success even if not “cure.”

Comparison with Pediatric Recovery

The 68% vs. 5% recovery gap between pediatric and adult ME/CFS represents a natural experiment with profound implications (see also Section 19.9 and Hypothesis 5.5.1).

Observation 33 (The Pediatric-Adult Recovery Gap). The 13-fold difference in recovery rates between children (68% at 10 years) and adults (~5%) cannot be fully explained by biological factors alone. Contributing factors likely include:

- Earlier diagnosis in children (shorter diagnostic delay)
- School accommodations versus inadequate workplace accommodations
- Family support versus adult isolation
- Permission to rest versus pressure to work
- Expectation of recovery versus therapeutic nihilism

If these environmental factors substantially explain the gap, adult outcomes might improve with earlier diagnosis, better accommodations, and adequate rest opportunity. This remains to be tested.

The hypothesis of developmental plasticity (Hypothesis 5.5.1) may also contribute: the developing nervous and immune systems may have recovery capacity that diminishes with age. However, even if biological plasticity explains part of the gap, the environmental differences are so substantial that they likely contribute meaningfully.

Implications for Clinical Practice

The evidence, while limited, suggests several clinical priorities:

- **Early intervention:** Diagnose early and implement aggressive symptom management from the outset. The first 2 years may represent a critical window.
- **Accommodation advocacy:** Help patients access rest, workplace accommodations, and disability benefits when needed. Forced overexertion may prevent recovery.
- **Subtype identification:** As biomarkers emerge, identify patient subtypes that may respond to specific treatments.

- **Realistic hope:** Recovery is rare but occurs. Improvement is more common. Neither nihilism (“nothing helps”) nor false optimism (“you’ll be fine”) serves patients.
- **Avoid iatrogenic harm:** Graded exercise therapy and other harmful interventions may damage recovery potential. First, do no harm.

★ **Key Point: Counseling Adult Patients on Prognosis**

When counseling adult patients:

- Full recovery is uncommon (~5–10%) but does occur
- Meaningful improvement is more common (~40%)
- Earlier diagnosis and appropriate management may improve odds
- The 5% figure has significant limitations and may underestimate recovery potential
- Hope is reasonable; false promises are not

The goal is honest realism that neither crushes hope nor creates false expectations.

Research Priorities

? **Open Question 1: Critical Gaps in Adult ME/CFS Prognosis Research**

Current prognosis evidence has critical gaps that limit clinical utility:

1. **Standardized recovery definitions:** International consensus is needed on operationalizing “recovery,” “remission,” and “improvement”
2. **Community-based cohorts:** Population-based studies are needed to overcome tertiary care selection bias
3. **Early intervention trials:** RCTs of pacing-based early intervention (within 6 months of onset) are urgently needed; CBT-based early intervention was ineffective
4. **Subtype-specific prognosis:** Outcomes stratified by pathophysiological subtype, onset type, and biomarker profile
5. **Accommodation interventions:** Does providing adequate rest and accommodations improve adult outcomes?
6. **Long-term follow-up:** Studies with >10 year follow-up to capture late recovery

Until these gaps are addressed, individual prognosis counseling will remain imprecise.

~ **Hypothesis 3: True Adult Recovery Potential**

If adults received equivalent early diagnosis, accommodations, rest opportunity, and subtype-specific treatment as pediatric patients, recovery rates might approach pediatric levels for comparable illness durations and presentations. The current ~5% rate may reflect the consequences of late diagnosis, inadequate support, and absence of targeted treatment rather than an inherent biological ceiling on adult recovery. Testing this hypothesis requires early intervention trials with comprehensive support systems.

Definition of “Recovery.” Recovery statistics must be interpreted cautiously because “recovery” is defined inconsistently across studies. Definitions range from:

- No longer meeting diagnostic criteria (least stringent)
- Substantial improvement in function and symptoms
- Return to pre-illness functional level
- Complete resolution of all symptoms (most stringent)

By the strictest definition (complete resolution), recovery rates are near zero. Many patients who “recover” by looser definitions continue to manage residual symptoms, avoid triggers, and pace activities—they are improved but not cured.

5.5.3 Prognostic Factors

Factors Predicting Better Outcomes. Analysis of recovery and improvement in ME/CFS has identified several positive prognostic factors [77]:

- **Older age at disease onset:** Patients who recovered or improved had median onset age of 45 years versus 32 years for those who did not improve (OR 1.06 per year, $p = 0.028$). This counterintuitive finding may reflect selection effects (younger patients with milder disease not seeking specialty care) or biological differences.
- **Shorter diagnostic delay:** Patients who recovered or improved had mean diagnostic delay of 23 months versus 55 months for non-improvers (OR 0.98 per month, $p = 0.036$). This finding underscores the importance of early diagnosis and appropriate management from disease onset.
- **Pediatric/adolescent age:** As noted above, young patients have dramatically better outcomes than adults.
- **Shorter illness duration at baseline:** Earlier intervention is associated with better outcomes.
- **Milder initial severity:** Less severe initial presentation may predict better outcomes, though this finding is inconsistent.

Factors Predicting Worse Outcomes.

- **Longer illness duration:** The longer a patient has been ill, the lower the probability of recovery
- **Greater symptom severity:** More severe symptoms at baseline may predict worse outcomes
- **Comorbid conditions:** Multiple comorbidities may complicate recovery
- **Lower socioeconomic status:** Likely reflecting reduced access to rest, appropriate care, and supportive accommodations
- **Female sex:** Some studies show worse outcomes in women, possibly reflecting hormonal influences or access to care differences

Factors That Do Not Predict Outcomes. Several factors that might intuitively seem prognostic do not consistently predict outcomes:

- Baseline fatigue severity (in some studies)
- Post-exertional malaise severity at presentation
- Depression comorbidity
- Anxiety comorbidity
- ANA positivity
- Onset type (post-infectious vs. gradual) in many studies

The lack of reliable prognostic biomarkers limits the ability to counsel individual patients about their expected trajectory.

5.5.4 Long-Term Disability

ME/CFS causes profound, long-term disability that persists for most patients throughout their lives.

Functional Impairment Statistics. Population-based studies consistently document severe functional impairment [125]:

- **Housebound or bedbound:** 25–25.7% of patients at some point
- **Bedbound on worst days:** 61%
- **Unable to work full-time:** 87%
- **Unemployed:** 54% (versus 9% in general population)
- **Estimated U.S. housebound population:** Approximately 385,000
- **Estimated U.S. bedbound population:** Approximately 62,000

Quality of Life. ME/CFS consistently ranks among the lowest quality of life scores of any chronic condition [59, 60]:

- EQ-5D mean score: 0.47 (versus population mean of 0.85)
- Lower than 20 other chronic conditions including multiple sclerosis and stroke
- SF-36 scores lower than multiple sclerosis across almost all domains
- Employment dropped from 89% pre-illness to 35% (versus 93% to 60% in multiple sclerosis)

These comparisons are important for communicating ME/CFS severity to healthcare providers, policymakers, and insurance companies who may underestimate the disease burden.

Disability Duration. For most adult patients, disability is lifelong:

- Mean illness duration in studies often exceeds 10 years
- Many patients have been ill for 20–30 years or more
- Disability typically begins at prime working age (20s–40s)
- Lost productivity spans decades
- Career development and financial security are permanently disrupted

5.5.5 Mortality

ME/CFS mortality remains an area of ongoing investigation and some controversy.

All-Cause Mortality. Large registry studies have not found significantly elevated all-cause mortality in ME/CFS compared to the general population [52, 138]. However, these studies have important limitations:

- Selection of milder cases able to seek medical care
- Underrepresentation of severe and very severe patients
- Short follow-up periods
- Diagnostic heterogeneity

Suicide. In contrast to all-cause mortality, suicide risk is consistently and substantially elevated in ME/CFS [52, 130, 135]:

- Standardized mortality ratio for suicide: 6.85 (95% CI 2.22–15.98) in one registry study
- Suicide accounts for 20–25% of deaths in memorial record studies
- Mean age at suicide death: 39.3 years (versus 47.4 in general population)
- 60% of suicide victims had no depression diagnosis
- 7.1% of ME/CFS patients report suicidal ideation without clinical depression

The elevated suicide risk in the absence of depression underscores that ME/CFS-specific suffering—not psychiatric comorbidity—drives suicide risk. This suffering includes:

- Severe, unrelenting physical symptoms
- Loss of identity, relationships, and life purpose
- Medical dismissal and gaslighting
- Hopelessness about prognosis
- Financial devastation
- Social isolation
- The specific circumstance of very severe ME/CFS (see Section 5.3)

Suicide prevention in ME/CFS must address these ME/CFS-specific factors, not merely screen for depression.

Cardiovascular Mortality. Memorial record studies suggest possible elevation of cardiovascular mortality [130, 131]:

- Heart failure is the leading cause of death in memorial records (29%)
- Mean age at cardiovascular death: 58.8 years versus 77.7 in general population

However, these findings from memorial records may reflect selection bias toward severe cases. The biological plausibility of cardiovascular risk in ME/CFS (autonomic dysfunction, chronic inflammation, reduced physical activity) suggests this deserves further population-based investigation.

Mean Age at Death. Memorial record studies report substantially reduced life expectancy [130, 131]:

- Mean age at death: 52.5–55.9 years across studies (McManimen 2016: 55.9 years, n=165; Sirotiak 2025: 52.5 years, n=512)
- General population mean age at death: 73.5 years
- Difference: Approximately 18–21 years of lost life expectancy

The variation between studies (52.5 vs. 55.9 years) likely reflects differences in cohort composition, with larger studies potentially capturing broader severity ranges. These figures must be interpreted with extreme caution due to selection bias in memorial records (deaths are more likely to be reported for severe cases and younger patients). Population-based mortality studies are urgently needed to establish true mortality patterns in ME/CFS.

5.5.6 Implications for Patients and Clinicians

Counseling Patients. Prognostic counseling should be honest while maintaining hope:

- Full recovery is unlikely in adults but does occur in a minority
- Improvement is possible with appropriate management
- Pediatric patients have substantially better outcomes
- Early diagnosis and aggressive pacing may improve outcomes
- The illness is typically lifelong, requiring permanent lifestyle adaptations
- Support for adjustment to chronic illness is important

Clinical Monitoring. Given the elevated suicide risk, clinicians should:

- Routinely assess for suicidal ideation
- Recognize that ME/CFS-specific suffering, not just depression, drives suicide risk
- Address hopelessness about prognosis
- Validate patient suffering rather than dismissing symptoms
- Connect patients with peer support communities

- Monitor for warning signs: social withdrawal, expressions of hopelessness, discussion of death

Research Priorities. The poor prognosis of ME/CFS and the lack of effective treatments underscore the urgent need for:

- Biomarker research to identify modifiable disease drivers
- Clinical trials of candidate therapeutics
- Early intervention studies
- Population-based mortality studies
- Investigation of factors differentiating pediatric (good) from adult (poor) prognosis

Until effective treatments are available, the prognosis of ME/CFS will remain poor, and millions of patients worldwide will face lifelong disability from a disease that the medical establishment has failed to adequately address.

5.6 Subgroups and Phenotypes

ME/CFS is increasingly recognized as a heterogeneous syndrome that likely encompasses multiple distinct biological subgroups. Identifying these subgroups is essential for developing targeted treatments, understanding pathophysiology, and improving diagnostic precision. Research has identified potential subgroups based on symptom profiles, onset patterns, biomarkers, and metabolic phenotypes.

5.6.1 The Heterogeneity Problem

The heterogeneity of ME/CFS has profound implications for research and clinical care:

- **Research confounding:** Clinical trials that mix different subgroups may show no overall effect even when treatments work for specific subgroups
- **Diagnostic uncertainty:** Different diagnostic criteria identify different patient populations with varying severity [112]
- **Pathophysiology confusion:** Studies may find contradictory results because they examine different disease subtypes
- **Treatment failure:** Interventions effective for one subgroup may be harmful for others

One analysis comparing different diagnostic frameworks (Fukuda, Canadian Consensus, and ICC criteria) found that they identify phenotypes with significant differences in cognitive performance, autonomic dysfunction, and symptom burden [112]. The authors concluded: “Different CFS criteria may at best be diagnosing a spectrum of disease severities and at worst different CFS phenotypes or even different diseases.”

5.6.2 Onset-Based Subgroups

Post-Infectious ME/CFS. Approximately 64% of ME/CFS cases have identifiable post-infectious onset [120]. This subgroup may be characterized by:

- Clear temporal relationship between infection and illness onset
- Evidence of ongoing immune activation or viral persistence
- Potentially better prognosis than gradual onset (in some studies)
- Distinct brain abnormalities on neuroimaging

The NIH deep phenotyping study specifically selected post-infectious ME/CFS patients, providing detailed characterization of this subgroup including alterations in catecholamine pathways, immune profiles suggesting chronic antigenic stimulation, and abnormal cardiopulmonary responses [5].

Gradual-Onset ME/CFS. Approximately 36% of cases develop gradually without clear infectious trigger [120]. Characteristics may include:

- Higher rates of psychiatric comorbidity
- Different patterns of brain abnormalities compared to post-infectious
- Longer diagnostic delay (trigger less obvious)
- Possibly different underlying mechanisms

Clinical Implications of Onset Type. While onset type may have research significance for identifying biological subgroups, its clinical utility remains unclear:

- Both types develop the same symptom complex
- Both require the same management approaches (pacing, symptom management)
- Prognostic value is inconsistent across studies
- Treatment response differences have not been established

5.6.3 Severity-Based Subgroups

Evidence suggests that severe ME/CFS may represent a qualitatively different disease state rather than simply the extreme end of a continuum [128].

Severe vs. Mild/Moderate ME/CFS. Compared to milder patients, those with severe ME/CFS demonstrate:

- Greater autonomic dysfunction
- More frequent and more severe post-exertional malaise
- More pronounced cognitive impairment
- More multisystem symptom involvement

- Significantly worse scores across all SF-36 domains

These differences suggest that additional pathophysiological mechanisms may be operating in severe disease, or that certain biological factors predispose some patients to more severe manifestations.

Implications. If severe ME/CFS is biologically distinct, then:

- Research findings from mild/moderate patients may not apply to severe patients
- Treatments effective for milder disease may not help (or may harm) severe patients
- Severe patients may need distinct biomarker panels and outcome measures
- Clinical trials should stratify by severity or focus on specific severity levels

5.6.4 Metabolic Phenotypes

Metabolomic studies have identified distinct metabolic subgroups within ME/CFS [139]:

Three Metabotypes. Analysis of 83 ME/CFS patients identified three distinct metabolic phenotypes:

Table 5.2: Metabolic phenotypes in ME/CFS

Subgroup	Size	Metabolic Features	Clinical Features
ME-M1	$n = 32$	High ketones, high FFAs, low amino acids, low TGs (lipolytic state)	Lower BMI (23.1), intermediate function
ME-M2	$n = 38$	High TGs/insulin, low fatty acid derivatives, high pyruvate (lipid accumulation)	Highest BMI (25.7), worst function (SF-36 PF = 22.2)
ME-M3	$n = 13$	Intermediate, partial overlap with controls	Best function , predominantly mild

Clinical Significance. The ME-M2 phenotype (lipid accumulation) was associated with the worst functional status, suggesting that metabolic context influences disease severity. This has potential therapeutic implications:

- Different metabolic phenotypes may respond to different interventions
- Lipolytic (ME-M1) versus lipid accumulation (ME-M2) states may require opposite metabolic support strategies
- Metabolic phenotyping could guide personalized treatment

However, these findings require replication and clinical validation before they can be applied in practice.

5.6.5 Immune Phenotypes

Recent research has revealed distinct immune profiles within ME/CFS populations.

Sex-Specific Differences. The NIH deep phenotyping study found that male and female ME/CFS patients show different immune abnormalities [5]:

- **Males:** Altered T cell activation, markers of innate immunity
- **Females:** Abnormal B cell and white blood cell growth patterns
- **Both:** Distinct inflammation markers

These sex-specific differences may explain some of the variability in ME/CFS presentation and treatment response, and underscore the importance of analyzing male and female patients separately in research studies.

T Cell Exhaustion. ME/CFS patients show evidence of T cell exhaustion similar to that seen in chronic viral infections and cancer:

- Elevated PD-1 expression
- Epigenetic changes indicating chronic antigenic stimulation
- Transcriptional reprogramming
- Potential implications for immune checkpoint modulation as therapy

Effector Memory Profiles. Detailed immune phenotyping has identified abnormalities in T cell subsets [12]:

- Decreased CD45RA⁺CCR7⁺ effector memory CD4⁺ T cells
- Effector memory dominated by CD27⁺CD28⁺ early phenotype
- Significantly reduced CD27⁺CD28⁺ terminal effector memory subset

These findings suggest skewing toward less mature effector subsets, consistent with chronic antigenic stimulation without resolution.

5.6.6 Symptom-Based Subgroups

Clinical observation suggests potential subgroups based on dominant symptom patterns:

Proposed Symptom Clusters.

- **Pain-predominant:** Widespread pain, fibromyalgia-like features, myalgia
- **Cognitive-predominant:** Severe brain fog, concentration difficulties, memory impairment
- **Autonomic-predominant:** Prominent POTS, orthostatic intolerance, temperature dysregulation
- **Immune-predominant:** Frequent infections, lymphadenopathy, sore throat, flu-like malaise
- **Sleep-predominant:** Severe unrefreshing sleep, hypersomnia or insomnia

Limitations. Symptom-based subgrouping is limited by:

- Most patients have symptoms across multiple domains
- Symptom prominence may shift over time within the same patient
- Symptom reporting is subjective and variable
- No validated method for symptom-based classification exists

5.6.7 Criteria-Based Phenotypes

Different diagnostic criteria identify different patient populations with varying characteristics [112]:

Table 5.3: Characteristics of patients meeting different diagnostic criteria

Criteria	Disease Severity	Characteristics
Fukuda only	Mildest	Least symptom burden
Fukuda + Canadian Clinical	Intermediate	Moderate severity
Fukuda + Canadian Research	Variable	Different autonomic profile
Fukuda + Canadian + ICC	Most severe	Worst cognitive performance, highest symptom burden

This finding has important implications:

- Research using different criteria studies different populations
- Comparisons across studies using different criteria are problematic
- Stringent criteria (ICC) select the most impaired patients
- Broad criteria (Fukuda alone) may include patients with other conditions

5.6.8 Clinical Significance of Subgrouping

Current State. ME/CFS subgroups are not yet validated as clinical standard, but are increasingly necessary as a harm-reduction strategy:

- No subgroup-specific treatments have been validated in randomized controlled trials
- Subgroup testing is not available in routine clinical practice
- Subgroups identified in research have not been replicated consistently
- Clinical management largely remains the same regardless of potential subgroup

However, the absence of formal validation does not mean phenotyping lacks clinical utility. Two independent lines of evidence suggest that unguided empirical treatment carries disproportionate risk: (1) two-day CPET studies demonstrate that ME/CFS patients have severely impaired energy production that worsens after exertion [1, 140], and (2) pharmacological processing of any exogenous substance imposes metabolic demands on this already-depleted system. The *cost of not phenotyping*—irreversible deterioration from empirical treatment trials—may therefore exceed the cost of imperfect phenotype-guided treatment selection (see Chapter 25 for a treatment safety framework built on this reasoning).

Future Directions. Subgrouping holds promise for:

- **Precision medicine:** Matching treatments to specific disease mechanisms
- **Clinical trial design:** Enriching trials with patients likely to respond
- **Biomarker development:** Identifying subgroup-specific diagnostic markers
- **Pathophysiology understanding:** Clarifying distinct disease mechanisms
- **Drug development:** Targeting specific biological pathways
- **Treatment safety:** Enabling phenotype-guided treatment selection to reduce treatment-induced crashes in severe patients (see Section 25.3.3 for subtype-specific treatment pathways)

Research Priorities. Advancing the clinical utility of ME/CFS subgrouping requires:

- Large, well-characterized cohort studies with deep phenotyping
- Replication of subgroup findings across independent samples
- Longitudinal studies tracking subgroup stability over time
- Clinical trials stratified by potential subgroups
- Development of practical, affordable subgroup classification tools

Until these advances are achieved, ME/CFS will continue to be treated as a single entity, with the consequence that effective treatments for specific subgroups may be missed in trials that mix heterogeneous populations.

5.6.9 From Subgroups to Stratified Medicine

A fundamental principle in modern medicine is that heterogeneity is not a problem to be ignored—it is a signal to be decoded. Precision medicine paradigms across oncology, immunology, and psychiatry have demonstrated that when patient populations appear heterogeneous, the solution is stratification by endotype (functionally distinct disease biology), not abandonment of targeted therapeutics. ME/CFS research increasingly shows this same pattern: despite apparent clinical and biomarker heterogeneity, diverse upstream disease triggers converge on shared downstream pathophysiological mechanisms.

Convergence Despite Heterogeneity

Multiple lines of evidence suggest that ME/CFS subgroups, though heterogeneous in trigger and presentation, may converge on common pathophysiological endpoints:

miRNA convergence. Cheema et al. [11] analyzed circulating miRNA profiles across ME/CFS patients and found striking heterogeneity in individual miRNA expression. However, when these heterogeneous miRNA signatures were analyzed at the pathway level, they converged on a highly specific set of target gene clusters ($p < 0.002$), governing exercise hyperemia, angiogenic adaptation, antioxidant defenses, and mitochondrial dynamics. This suggests that while patients arrive at dysfunction through different miRNA alterations, the functional consequences converge on a narrow set of pathways.

Metabolic convergence. Naviaux et al. [141] described a “dauer-like” hypometabolic state observed across ME/CFS cohorts, characterized by downregulation of metabolic pathways, mitochondrial ATP production, and oxidative stress defenses. The metabolic signature showed diagnostic accuracy of 94–96%, with 80% of diagnostic metabolites decreased across the cohort. This suggests that regardless of how a patient develops ME/CFS, the resulting metabolic reorganization follows a stereotyped pattern.

Genetic convergence. More recently, Birch & Younger [142] examined genetic variation across ME/CFS patients and found gene-level heterogeneity in which specific genes were dysregulated. However, at the pathway level, patients consistently showed convergence on impaired energy production pathways, reduced stress resilience, and dysregulated post-exertional metabolic recovery. Thus genetic heterogeneity at the individual gene level masks functional convergence at the pathway level.

Cross-Disease Paradigm Lessons

The convergence-on-pathways pattern has clear precedent in other complex diseases. When heterogeneous clinical presentations have been investigated with molecular precision, stratified medicine approaches have consistently succeeded:

Asthma. Severe asthma was long treated as a single entity with poor outcomes in trial-and-error approaches. Only with endotyping (Th2-high vs. Th2-low stratification) did molecular stratification become possible [143]. Subsequent clinical trials demonstrated that Th2-high patients respond to IL-4/IL-13 inhibition and anti-IgE therapy, while Th2-low endotypes

require different approaches. The heterogeneity was not the problem—the lack of stratification was.

Depression. Major depressive disorder appears clinically homogeneous but encompasses distinct biological subtypes. Miller & Raison [144] demonstrated that a subset of depressed patients with elevated inflammatory markers (CRP > 3 mg/L) show poor response to conventional antidepressants but improved outcomes with anti-TNF biologics (infliximab). Patients without inflammatory elevation do not benefit from anti-inflammatory add-ons, showing that stratification is essential.

Breast cancer. Perou et al. [145] established that morphologically similar breast cancers represent distinct molecular subtypes (luminal A/B, HER2-enriched, basal-like), each with different prognoses and treatment responses. This molecular stratification transformed oncology from cytotoxic chemotherapy for all to subtype-matched targeted therapies (ER/PR inhibitors, HER2 inhibitors, etc.), dramatically improving outcomes.

In each of these domains, initial appearance of heterogeneity led to treatment failure when approaches were not stratified. Once endotypes were identified, targeted interventions achieved specificity and efficacy.

Dual Strategy Framework for ME/CFS Therapeutics

These principles suggest that ME/CFS therapeutic development should pursue *both* upstream and downstream targets:

Upstream stratification. Identify ME/CFS subgroups defined by biomarker endotypes and match treatments to these subgroups:

- **Autoimmune-predominant** (elevated GPCR autoantibodies): Immunoabsorption and monoclonal antibodies (daratumumab) targeting autoreactive B cells [9, 118]
- **Immune-inflammatory-predominant** (elevated pro-inflammatory cytokines, T-cell exhaustion): Immunomodulation targeting specific cytokine elevations (see Chapter ??)
- **Autonomic-predominant** (sympathetic dysautonomia cluster): Targeted autonomic interventions including heart rate variability training, selective norepinephrine support, or sympatholytic approaches depending on dysautonomia subtype [146]

Downstream convergent targets. Pursue shared pathway interventions that may benefit across subgroups (see Chapters 22 and ?? for evidence and dosing):

- Mitochondrial support (CoQ10, L-carnitine, ubiquinol)
- PDH complex enhancement (thiamine, alpha-lipoic acid)
- Antioxidant restoration (NAC, vitamin E, selenium)
- Metabolic stabilization (low-dose naltrexone, metabolic cofactors)

This dual approach leverages the convergence principle: subgroup-specific treatments address the upstream pathogenic differences, while convergent pathway interventions address the shared downstream consequences.

Endotype-Specific Treatment Mapping

Based on emerging literature, the following endotype-to-treatment mappings are suggested. Table 5.4 extends the basic mapping with reproducibility status, clinical accessibility of biomarkers, and evidence levels for proposed treatments.

Table 5.4: Comprehensive endotype-to-biomarker-to-treatment mapping in ME/CFS. Evidence: **A** = RCT data; **B** = pilot/open-label; **C** = mechanistic rationale only. Reproducibility: **R** = replicated in ≥ 2 independent cohorts; **P** = preliminary (single study or unreplicated).

Proposed Endotype	Minimum Biomarkers	Access	Targeted Treatments	Evid.	Reprod.
Autoimmune	Elevated GPCR autoantibodies (β_1 , β_2 , M3, M4); B-cell activation markers	Tier 3	Immunoadsorption [147]; daratumumab [9]; BC007 [10]		R
Metabolic (ME-M2)	Elevated TAG/insulin ratio; low FFAs; high pyruvate; BMI trend [139]	Tier 2–3	Mitochondrial cofactors (CoQ10+NADH) [7]; metabolic support; insulin sensitization	A	P
Autonomic-Predominant	Abnormal tilt-table; reduced HRV; abnormal CO/CBF [4]	Tier 2	Beta-blockers (POTS); ivabradine; volume expansion; HRV biofeedback [146]	B	R
Post-Infectious (viral)	Documented viral onset; elevated HHV-6/EBV titers; disease duration 2–8 yr [148]	Tier 2	Valganciclovir (HHV-6/EBV subset) [149]; rintatolimod [148]	A/B	P
Immune-Inflammatory	Elevated IL-6, TNF- α ; T-cell exhaustion (PD-1 ⁺); reduced NK cytotoxicity [3]	Tier 2–3	LDN; targeted immunomodulation; anti-cytokine therapy	B	R

⚠ Limitation 5: Endotype Classification: Provisional Framework

This classification is provisional and based on emerging evidence. No endotype has been validated prospectively in a stratified clinical trial. Patients are not expected to fall neatly into single categories; many will show features of multiple endotypes, requiring combination approaches or sequential treatment. The framework serves as a research scaffold for biomarker-guided clinical decision-making, not as a validated diagnostic system. Biomarker accessibility tiers refer to Table 5.5.

5.6.10 From “Not Clinically Actionable” to “Increasingly Necessary”

The Cost of Not Phenotyping. The traditional treatment approach in ME/CFS—empirical trial-and-error—assumes that failed trials carry manageable costs. In practice, this assumption fails catastrophically for severe patients. Each treatment trial imposes metabolic processing demands on an energy-depleted system (see Treatment Energy Categories, Section 25.1.2). When a trial triggers post-exertional malaise, the resulting crash can cause lasting functional decline—the crash severity dose-response hypothesis estimates 5–15% permanent function loss from severe crashes and potentially greater loss from catastrophic ones (Section 18.3.3, Table 18.3).

For a very severe patient operating at the lowest functional levels, there may be only one or two tolerable failed trials before irreversible deterioration. In this context, empirical treatment selection is not “trial-and-error”—it is gambling with irreplaceable biological reserves.

~ Hypothesis 4: Prospective Phenotyping as Harm Reduction

Even imperfect phenotype-guided treatment selection is likely safer than unguided empirical trials for severe ME/CFS patients. The reasoning:

1. **Reducing trial number:** Matching treatment to subtype reduces the expected number of failed trials before finding an effective intervention
2. **Energy category matching:** Phenotyping enables prioritizing Category A (energy-providing) treatments for metabolic-predominant subtypes and reserving Category C (energy-demanding) treatments for subtypes where biomarkers predict benefit
3. **Avoiding known risks:** Pharmacogenomic data (CYP450 metabolizer status) can prevent paradoxical reactions before they occur
4. **Asymmetric cost:** The cost of phenotyping is primarily financial and one-time, while the cost of a treatment-induced crash is primarily biological and potentially permanent. Though these costs are measured on different scales, the irreversibility of severe crashes (loss of functional capacity that may never be recovered) makes the comparison asymmetric in favor of phenotyping for patients with limited remaining functional reserves

Certainty: 0.35 (strong mechanistic reasoning from crash dose-response data and pharmacogenomic principles; clinical observation of treatment-induced deterioration in severe patients; no RCTs comparing phenotype-guided vs. empirical treatment selection in ME/CFS)

Testable prediction: Severe ME/CFS patients treated with phenotype-guided protocols will experience fewer treatment-induced crashes and faster time-to-effective-treatment compared to patients treated empirically.

A Tiered Prospective Phenotyping Protocol. Practical phenotyping need not require research-grade facilities. A tiered approach allows any physician to begin phenotype assessment, with increasing precision at higher tiers:

The Septad diagnostic hierarchy (Section 5.6.11) already demonstrates how presentation-based phenotyping can guide clinical decisions. The tiered protocol above extends this principle

Table 5.5: Tiered prospective phenotyping protocol for ME/CFS treatment selection

Tier	Assessments	What it identifies	Treatment guidance
Tier 1 Any GP	CBC, CMP, TSH, ESR/CRP, ferritin, vitamin D, B12/folate; vital signs including orthostatic test; detailed symptom questionnaire; medication history	Severity level; dominant symptom cluster (fatigue vs. pain vs. cognitive vs. autonomic); basic metabolic status; obvious comorbidities	Severity-appropriate starting point; foundation-first sequencing (Section 25.1.6); avoid Category C treatments until Tier 2
Tier 2 Specialist	Tilt table test; NK cell function; lymphocyte subsets; viral serology (EBV, HHV-6, CMV); amino acid/organic acid panel; cortisol rhythm; tryptase	Immune-predominant vs. autonomic-predominant vs. metabolic-predominant subtype; MCAS status; viral reactivation burden	Subtype-specific treatment pathway (Section 25.3.3); targeted Category B–C trials where biomarkers predict benefit
Tier 3 Research	Metabolomics; autoantibody panels (GPCR antibodies); TRPM3 function; pharmacogenomics (CYP2D6, CYP2C19, COMT, MTHFR); cytokine panel	Specific metabolic phenotype (ME-M1/M2/M3); autoimmune subtype; drug metabolism profile; paradoxical reactor risk	Precision treatment matching (Section 56); pharmacogenomic-guided dosing (Section 25.3.2)

from comorbidity diagnosis to treatment selection.

For treatment-selection implications of phenotyping, see the subtype-specific treatment pathways in Section 25.3.3 and the treatment safety framework in Section 25.1.

5.6.11 Comorbidity Clustering: The “Septad” Framework

Clinical observation by specialists treating complex chronic illness has identified a consistent pattern of comorbidity clustering in ME/CFS patients. Dr. David Kaufman and colleagues have formalized this observation as the “Septad”—seven pathophysiologies that frequently co-occur and interact.¹

Peer-Reviewed Evidence for Comorbidity Clustering. While the specific “Septad” terminology is not peer-reviewed, the individual comorbidity associations are well-documented:

- **hEDS-POTS-MCAS triad:** Wang et al. [150] found MCAS prevalence of 31% in patients with both POTS and EDS versus 2% in controls (OR=32.46, $p<0.001$). Note: this study examined the POTS+EDS population specifically, not ME/CFS. Kucharik and Chang [151] caution that mechanistic links between these conditions remain unestablished.
- **POTS in ME/CFS:** Hoad et al. [152] found 27% of ME/CFS patients met POTS criteria versus 9% of controls ($p=0.006$).
- **Hypermobility in ME/CFS:** Hakim et al. [153] report 30–57% of ME/CFS patients have joint hypermobility versus 10–15% in the general population.
- **Dysautonomia in EDS:** Mathias et al. [154] found up to 70% of hEDS patients report dysautonomia symptoms, with up to 40% meeting formal POTS criteria.

These prevalence data support clinical clustering but do not validate the Septad as a unified syndrome with shared pathophysiology. The remaining components (autoimmunity, chronic infection, SFN, GI dysmotility) lack equivalent systematic prevalence studies in ME/CFS populations.

~ Hypothesis 5: The Septad: Seven Interacting Pathophysiologies

ME/CFS patients frequently present with a cluster of seven interrelated conditions that may share underlying mechanisms:

1. **Mast Cell Activation Syndrome (MCAS):** Aberrant mast cell degranulation causing multisystem symptoms including flushing, urticaria, GI disturbance, and anaphylactoid reactions. See Chapter 31 for mechanistic details.
2. **Ehlers-Danlos Syndrome (EDS) / Hypermobility:** Connective tissue laxity affecting joints, vessels, and organs. See Chapter 31 for hEDS-POTS-MCAS connections.
3. **Dysautonomia / POTS:** Autonomic dysfunction manifesting as orthostatic intolerance.

¹The Septad framework as a named seven-condition cluster originates from clinical presentations by Kaufman and colleagues rather than a single peer-reviewed publication. However, substantial peer-reviewed evidence supports the underlying comorbidity patterns (see “Peer-Reviewed Evidence” below).

ance, heart rate variability, temperature dysregulation. See Chapters 8 and 10 for detailed pathophysiology.

4. **Autoimmunity:** Subclinical or overt autoimmune markers and processes. See Chapter 7 for immune abnormalities.
5. **Chronic Infection:** Viral reactivation (EBV, HHV-6), tick-borne infections (Lyme, Bartonella), or other persistent pathogens. See Chapter 7 for viral reactivation mechanisms.
6. **Small Fiber Neuropathy (SFN):** Damage to small nerve fibers causing pain, paresthesias, and autonomic symptoms. See Chapter 8 for autonomic neuropathy.
7. **GI Dysmotility:** Impaired gut motility (gastroparesis) leading to small intestinal bacterial overgrowth (SIBO) and malabsorption. See Chapter 11 for gastrointestinal pathophysiology.

Clinical Rationale. The Septad emerged from clinical pattern recognition: Dr. Andy Maxwell, a cardiologist treating MCAS patients, observed that nearly all presented with the same constellation of conditions. Kaufman and colleagues recognized this as a framework for organizing the complexity of these patients, noting that “the Septad creates a map that allows the physician to organize what I’ve heard in a much more usable and actionable way.”

Interconnections. Critically, these seven pathophysiologicals are not independent—they interact bidirectionally:

- **MCAS ↔ Dysautonomia:** Mast cell mediators directly affect autonomic function; autonomic dysfunction can trigger mast cell degranulation
- **EDS ↔ POTS:** Connective tissue laxity in blood vessels contributes to venous pooling and orthostatic intolerance
- **MCAS ↔ GI dysmotility:** Mast cells in gut mucosa affect motility; SIBO can trigger mast cell activation
- **SFN ↔ Dysautonomia:** Small fiber damage underlies autonomic neuropathy
- **Chronic infection ↔ Autoimmunity:** Molecular mimicry and chronic immune stimulation
- **EDS → Craniocervical instability:** Connective tissue weakness may lead to cervical spine instability, potentially compressing brainstem [155, 156]

Kaufman describes the framework as having “seven circles with a million arrows—because it all interacts.”

Causal Cascade Model: Beyond “Comorbidities”

The traditional framing of Septad conditions as “comorbidities” (independent conditions that happen to coexist) may be inadequate. A more useful clinical model considers these conditions as *potentially cascading pathophysiologicals*, where each can initiate or amplify others.

~ Hypothesis 6: Septad Conditions as Cascading Pathophysiologies

Rather than seven independent conditions with coincidental co-occurrence, the Septad may represent a pathophysiological cascade where upstream conditions drive downstream manifestations. **Primary initiators** (hEDS, chronic infection, MCAS) drive **secondary amplifiers** (dysautonomia, SFN, GI dysmotility, autoimmunity), which ultimately cause **tertiary consequences** (mitochondrial dysfunction, ME/CFS phenotype).

Primary Initiators (Upstream Conditions):

- **hEDS/Connective Tissue Disorder:** May be the foundational substrate—vascular laxity drives POTS; altered mast cell distribution in abnormal connective tissue drives MCAS; nerve fragility drives SFN
- **Chronic Infection (EBV, HHV-6):** Persistent viral reactivation exhausts T cells, triggers autoimmunity via molecular mimicry, and maintains chronic mast cell activation
- **MCAS:** Mast cell mediators damage intestinal barrier (causing malabsorption), sensitize autonomic neurons (driving dysautonomia), and maintain neuroinflammation

Secondary Amplifiers (Downstream Conditions):

- **Dysautonomia/POTS:** Results from vascular laxity (hEDS), autonomic neuropathy (SFN), mast cell mediators (MCAS), or deconditioning
- **Small Fiber Neuropathy:** May result from autoimmune attack, metabolic dysfunction (malabsorption), or chronic inflammation
- **GI Dysmotility/SIBO:** Results from autonomic neuropathy, mast cell damage to enteric nervous system, or vagal dysfunction
- **Autoimmunity:** Triggered by chronic infection (molecular mimicry), persistent inflammation, or loss of self-tolerance

Tertiary Consequences:

- **Mitochondrial dysfunction:** Amino acid malabsorption (from GI dysfunction) impairs TCA cycle and glutathione synthesis
- **ME/CFS phenotype:** Energy failure, PEM, cognitive dysfunction emerge as final common pathway

⚠ Limitation 6: Cascade Model Epistemic Status

This cascade model is hypothetical and based on mechanistic plausibility, not prospective validation. Individual patients may have different primary drivers, and causality cannot be inferred from correlation. The model is presented to guide clinical thinking, not as established science.

Example Cascade Pathways. Several documented cascade pathways illustrate how upstream conditions propagate:

1. **hEDS → POTS → Deconditioning → ME/CFS-like presentation**

- Connective tissue laxity → venous pooling → orthostatic intolerance
 - Orthostatic intolerance → activity avoidance → deconditioning
 - Deconditioning → exercise intolerance resembling PEM
2. **MCAS → Gut Barrier Dysfunction → Mitochondrial Failure**
- Mast cell mediators damage intestinal tight junctions
 - Barrier dysfunction → amino acid malabsorption
 - Malabsorption → impaired NO synthesis, glutathione depletion, TCA dysfunction
 - Metabolic failure → ME/CFS phenotype
 - See Section 11.1.6 for detailed mechanism
3. **Chronic Viral Infection → Immune Exhaustion → Multiple Sequelae**
- EBV/HHV-6 reactivation → T cell exhaustion
 - Immune dysfunction → failure to suppress mast cells → MCAS
 - Immune dysfunction → autoantibody production → SFN, autonomic neuropathy
 - Cimetidine enhancement of cellular immunity may interrupt this cascade
4. **SFN → Autonomic Neuropathy → Multi-System Dysfunction**
- Small fiber damage → autonomic nerve impairment
 - Autonomic neuropathy → POTS (neuropathic subtype)
 - Autonomic neuropathy → GI dysmotility, bladder dysfunction
 - Autonomic neuropathy → sudomotor dysfunction, temperature dysregulation

Diagnostic Hierarchy: Which to Test First

Given resource constraints and cascade dynamics, a hierarchical diagnostic approach prioritizes upstream conditions whose treatment may interrupt downstream pathology.

Observation 34 (Presentation-Based Diagnostic Prioritization). Diagnostic testing should be prioritized based on the dominant clinical presentation, testing upstream conditions first to identify primary drivers before downstream consequences.

If MCAS/HIT Features Dominate:

1. **First:** Confirm mast cell activation (tryptase, histamine, 24-hour urine prostaglandins)
2. **Second:** Assess intestinal barrier (zonulin, LPS antibodies, fecal calprotectin)
3. **Third:** Check downstream metabolic consequences (amino acid panel, organic acids)
4. **Rationale:** MCAS drives gut dysfunction which drives metabolic failure; treating MCAS upstream may restore gut function and metabolism without direct supplementation

If Post-Infectious Pattern:

1. **First:** Viral serology panel (EBV VCA IgG/IgM, EBNA, HHV-6 IgG, CMV IgG)
2. **Second:** T cell immunophenotyping (CD4/CD8, NK function if available)
3. **Third:** Autoantibody screen (anti-autonomic antibodies if accessible)

4. **Rationale:** Post-infectious patients may have ongoing viral reactivation or immune exhaustion that, if addressed (antivirals, immunomodulation), interrupts downstream complications

If Hypermobility/hEDS Features:

1. **First:** Beighton score, Brighton criteria for hypermobility
2. **Second:** Assess structural consequences (upright MRI if severe symptoms)
3. **Third:** Vascular assessment (tilt table for POTS, echocardiogram if murmur)
4. **Rationale:** hEDS is the upstream structural condition; understanding connective tissue status guides interpretation of all downstream conditions

If Autonomic Features Dominate:

1. **First:** Formal autonomic testing (tilt table, QSART)
2. **Second:** Distinguish POTS subtypes (hyperadrenergic vs. neuropathic vs. hypovolemic)
3. **Third:** If neuropathic pattern, assess for SFN (skin biopsy, autonomic antibodies)
4. **Rationale:** POTS subtype determines treatment approach and identifies whether SFN or autoimmunity is the upstream driver

If GI Symptoms Dominate:

1. **First:** SIBO testing (breath test), celiac panel
2. **Second:** Assess for mast cell involvement (GI biopsy with tryptase staining if severe)
3. **Third:** Autonomic GI testing (gastric emptying study)
4. **Rationale:** GI dysfunction can be primary (MCAS-driven) or secondary (autonomic neuropathy-driven); treatment differs substantially

General Principle. Test upstream before downstream. Treat upstream first. If upstream treatment produces disproportionate improvement in downstream conditions, this validates the cascade model for that patient and suggests the “comorbidity” was actually a consequence, not an independent condition.

Craniocervical Instability (CCI). While not part of the original Septad, craniocervical instability has emerged as a related concern with accumulating research evidence. Some patients with EDS and the other Septad components develop instability at the craniocervical junction, potentially causing brainstem compression. Kaufman notes that aggressive connective tissue strengthening may be important to prevent progression to CCI in susceptible patients.

Observation 35 (High Prevalence of Structural Abnormalities in ME/CFS). Brag e et al. [155] conducted upright MRI imaging in 229 ME/CFS patients (Canadian Consensus Criteria), finding craniocervical obstructions in 80% (183/229), signs of intracranial hypertension in 78% (179/229), and hypermobility indicators in 75% (172/229). Notably, 45% had Chiari malformation (cerebellar tonsillar descent >5mm) compared to 0.5–1% prevalence in the general population. Structural findings correlated with orthostatic intolerance severity ($r=0.42$,

$p < 0.001$), suggesting a potential mechanistic contribution to autonomic dysfunction in the hypermobile subset (prospective study, $n=229$, Medium certainty).

△ **Warning 3: Selection Bias and Interpretation Caveats**

The high prevalence of structural abnormalities reported by Brag e et al. [155] comes from a specialized clinic that focuses on craniocervical pathology and may represent a selected population; authors are affiliated with the clinic providing structural interventions, representing a potential conflict of interest. Additionally, the study lacked matched healthy controls with upright MRI, using historical controls from supine imaging instead. Independent replication in community-based, unselected ME/CFS cohorts is needed to determine generalizability. A systematic review of CCI in EDS [156] (16 studies, $n=695$) found significant heterogeneity in diagnostic criteria, with no consensus on single measurement thresholds—necessitating comprehensive evaluation using multiple imaging parameters and clinical correlation.

Diagnostic and Treatment Considerations. Upright MRI evaluation should be considered in ME/CFS patients with hypermobility (Beighton score ≥ 5), severe orthostatic intolerance, positional symptoms (worse upright, better supine), progressive neurological deficits, or suboccipital headaches. Reference ranges for CCI measurements on upright dynamic MRI have been established [157]. Conservative management including specialized physical therapy [158] should be first-line; surgical stabilization (occipito-cervical fusion) shows 60–80% improvement in properly selected patients but carries significant complication rates (19%) [159, 156]. Patient selection is critical, as surgical intervention is appropriate only for progressive myelopathy or failed conservative treatment.

Treatment Sequencing. The Septad framework suggests a treatment sequence: address MCAS first (stabilize mast cells), then systematically work through the other components. This approach recognizes that treating one component may improve others due to their interconnections.

Evidence Status and Limitations.

⌘ **Limitation 7: Septad: Clinical Co-occurrence $\neq$ Shared Pathophysiology**

The Septad is a *clinical framework* based on expert observation, not a validated research model.

What peer-reviewed evidence supports:

- **Pairwise comorbidity associations:** The hEDS-POTS-MCAS triad [150], POTS in ME/CFS [152], and hypermobility in ME/CFS [153] have systematic prevalence data (see above). Clinical co-occurrence of *subsets* is established.

What data do not establish:

- No peer-reviewed publication validating the *Septad framework* as a distinct entity—only subsets (particularly the hEDS-POTS-MCAS triad) have been systematically

studied

- **Mechanistic link unproven:** “An evidence-based, common pathophysiologic mechanism between any of the two, much less all three conditions, has yet to be described” [151]
- Prevalence of autoimmunity, chronic infection, SFN, and GI dysmotility in ME/CFS populations not systematically studied
- Selection bias inherent (specialists see the most complex patients; Brag e CCI data from specialized clinic)
- Treatment sequencing recommendations lack controlled trial evidence
- Rapamycin pilot was uncontrolled; only 40 of 86 enrolled (47%) completed the full 90-day protocol

The critical distinction: *clinical co-occurrence is documented; shared pathophysiology is not.*

Important Clarification: The Septad Is Not Diagnostic. The Septad framework is for evaluating *comorbidities*, not for diagnosing ME/CFS. Post-exertional malaise (PEM) remains the hallmark diagnostic feature of ME/CFS (see Section 2.1). A patient may have none, some, or all Septad components and still have ME/CFS—provided PEM is present. Conversely, having all seven Septad conditions does not constitute ME/CFS without PEM.

The Septad’s clinical utility lies in systematic comorbidity screening: many ME/CFS patients have undiagnosed MCAS, EDS, or other conditions that require distinct treatment approaches. Identifying these can improve symptom management even when ME/CFS itself remains treatment-resistant.

Research Implications. If the Septad represents a genuine disease phenotype, it suggests:

- ME/CFS may be a final common pathway for connective tissue/mast cell/autonomic dysfunction
- Subgrouping by comorbidity pattern may improve treatment targeting
- Comprehensive workup should screen for all seven components
- Multi-system treatment approaches may outperform single-target interventions

Of the seven Septad components, only four (chronic infection, dysautonomia/POTS, autoimmunity, small fiber neuropathy) are currently being actively pursued in ME/CFS research, suggesting potential underexplored avenues.

Speculative Mechanistic Hypotheses. The clinical clustering of Septad components suggests potential unifying mechanisms that may explain why these conditions co-occur. Two hypotheses merit consideration:

~ Hypothesis 7: Autophagy/mTOR Dysfunction as Septad Unifier

The rapamycin pilot study [160] reported 74.3% symptom improvement and observed autophagy marker changes (BECLIN-1 upregulation and pSer258-ATG13 suppression), though whether autophagy restoration mediated the clinical effect cannot be established from an uncontrolled trial. Autophagy dysfunction could theoretically contribute to multiple Septad components: mast cell degranulation regulation (MCAS), mitochondrial quality control in autonomic neurons (dysautonomia), small nerve fiber maintenance (SFN), enteric nervous system function (GI dysmotility), and intracellular pathogen clearance (chronic infection) [160]. If validated, Septad-positive patients may represent an autophagy-dysfunction subgroup.

⚠ Limitation 8: Autophagy–Septad Extrapolation Boundaries

This hypothesis extrapolates from a single uncontrolled pilot study to multi-system effects not measured in that trial. The rapamycin study enrolled 86 patients; 70 completed day 36 and 40 completed the full 90-day protocol, representing 53% attrition that may bias results. The study did not assess Septad component status, mast cell markers, nerve fiber density, or GI function. The mechanistic connections (autophagy → each Septad component) are individually plausible based on cellular biology but have not been demonstrated in ME/CFS cohorts.

~ Hypothesis 8: Connective Tissue Matrix as Common Substrate

Six of seven Septad components have anatomical or functional connections to connective tissue: EDS is a primary connective tissue disorder; POTS involves vascular wall compliance; SFN involves nerve fibers traversing connective tissue matrix; GI dysmotility depends on gut wall integrity; mast cells reside in connective tissue and show increased prevalence of dysregulation in hypermobile patient populations [150]; and autoimmunity can target connective tissue proteins. Rather than seven independent conditions, the Septad may represent downstream manifestations of altered extracellular matrix composition or mechanics in hypermobile individuals.

⚠ Limitation 9: Connective Tissue Hypothesis: Untested Causal Claims

No studies have directly measured connective tissue biomarkers (matrix metalloproteinases, procollagen peptides, tenascin-C) in Septad-phenotype ME/CFS patients. The hypothesis that connective tissue abnormality causes (rather than merely correlates with) Septad clustering is untested. The non-EDS Septad components (autoimmunity, chronic infection) have weaker connective tissue links.

These hypotheses are presented to stimulate research, not as established mechanisms. Validation would require: (1) prospective studies measuring Septad component prevalence with standardized criteria; (2) biomarker studies comparing autophagy markers and connective tissue markers between Septad-positive and Septad-negative ME/CFS; (3) treatment stratification trials testing whether Septad status predicts response to mTOR inhibitors or connective tissue-targeted interventions.

5.6.12 Emerging Treatment-Response Phenotypes

Beyond biological markers, treatment response patterns may identify clinically actionable subgroups. While prospective validation is needed, retrospective observations suggest certain patient clusters respond preferentially to specific interventions.

The Viral-Immune-Metabolic Cluster (“Cimetidine-Responder” Phenotype)

△ **Warning 4: Preliminary Phenotype - No RCT Evidence**

This phenotype is based on clinical case series and mechanistic reasoning, not randomized controlled trials. Cimetidine has documented drug interactions (CYP450 inhibitor) and requires physician supervision. See Appendix H for detailed evidence assessment and safety considerations. Do not attempt self-treatment based on this phenotype description.

Clinical observation has identified a subset of ME/CFS patients who show dramatic improvement with cimetidine (an H2 receptor antagonist) combined with amino acid supplementation. This pattern suggests a distinct pathophysiological phenotype worthy of systematic investigation.

~ **Hypothesis 9: Cimetidine-Responder Phenotype**

A subset of post-infectious ME/CFS patients may have a viral-immune-metabolic phenotype characterized by:

Clinical Features:

- Post-infectious onset (typically EBV, HHV-6, or other herpesvirus)
- Prominent POTS/dysautonomia
- MCAS or histamine intolerance (HIT) comorbidity
- Strong response to amino acid supplementation (especially L-citrulline, N-Acetylcysteine (NAC))
- Dramatic improvement with cimetidine (“out of bed” effect in rare cases)

Proposed Mechanism: Two parallel pathways may converge:

1. **Viral pathway:** Chronic herpesvirus reactivation → T cell exhaustion → cimetidine enhances cellular immunity via H2 receptor blockade on suppressor T cells [161, 162]
2. **Metabolic pathway:** MCAS/HIT → intestinal barrier dysfunction → amino acid malabsorption → impaired NO synthesis and TCA cycle function → secondary mitochondrial dysfunction

Cimetidine may address the viral-immune component while amino acid supplementation restores metabolic capacity.

△ Warning 5: Evidence Limitations

The “cimetidine-responder” phenotype is based on:

- Historical case reports from 1980s–1990s (Goldstein, Lerner) suggesting benefit in EBV-associated CFS [161]
- Mechanistic studies of cimetidine immunomodulation (H2 receptor effects on T cell function) [162]
- Individual patient responses (anecdotal, selection bias)
- No controlled trials specifically testing this phenotype hypothesis

Prevalence is unknown but likely rare (< 5% of ME/CFS population). The dramatic responders may represent a distinct subgroup, or response may be placebo effect in susceptible individuals. Certainty: Very Low to Low.

Proposed Diagnostic Markers. If this phenotype exists, it may be identifiable by:

- **Viral markers:** Elevated EBV or HHV-6 antibody titers, positive PCR for viral DNA
- **Metabolic markers:** Low plasma amino acids (especially citrulline, arginine), abnormal organic acid profile
- **Immune markers:** T cell exhaustion phenotype (elevated PD-1), reduced NK cell function
- **Comorbidity pattern:** POTS + MCAS/HIT confirmed
- **Therapeutic trial:** Response to 2–4 week cimetidine trial (200–400 mg BID)

Treatment Approach (Hypothetical). For suspected cimetidine-responder patients:

1. **Confirmatory phase:** Trial cimetidine 200 mg BID for 2–4 weeks with symptom tracking
2. **If positive response:** Add comprehensive amino acid protocol (N-Acetylcysteine (NAC), L-citrulline-malate, consider antiviral if viral titers elevated)
3. **If no response:** Reassign to other phenotype; cimetidine unlikely to be beneficial
4. **Maintenance:** H1+H2 dual blockade for MCAS/HIT management

Research Priority. Validating this phenotype would require:

- Prospective cohort study with systematic phenotyping at baseline
- Randomized trial of cimetidine + amino acids in biomarker-selected patients
- Comparison of responders versus non-responders on viral, immune, and metabolic markers
- Replication across independent cohorts

Until validated, this phenotype should be considered a *clinical hypothesis* useful for generating treatment hypotheses in individual patients, not an established subgroup.

5.7 Trigger Mechanisms: From Acute Infection to Chronic Disease

The preceding sections document *which* infections trigger ME/CFS. This section explains *how*: the biological steps by which an acute infection can set in motion a process that does not resolve when the infection itself clears. Understanding these steps helps patients make sense of their illness and explains why recovery does not simply follow pathogen clearance. For detailed pathophysiological mechanisms, see Section 7.9.5 in Chapter 7.

5.7.1 The Core Problem: Failure to Resolve

In most infections, the immune system activates, eliminates or contains the pathogen, and then stands down. Inflammation resolves, tissues repair, and the person returns to health within weeks. In ME/CFS, this resolution step fails. The immune system remains in a state of ongoing activation long after the acute infection has passed [163]. The reasons for this failure differ across the major triggering pathogens, but they share a common endpoint: a chronic, self-sustaining pattern of immune and metabolic dysregulation.

△ Warning 6: Individual Variation

The mechanisms described here apply at the population level to patients whose ME/CFS followed a specific trigger. Not every patient who develops post-infectious ME/CFS will have all the mechanisms described, and the relative importance of each mechanism varies between individuals. The same trigger can initiate different pathological pathways in different people, reflecting differences in genetics, immune phenotype, and the state of the immune system at the time of infection.

5.7.2 Epstein-Barr Virus (EBV)

EBV is the most studied ME/CFS trigger, responsible for the infectious mononucleosis (“mono”) that precedes ME/CFS in a substantial proportion of post-infectious cases. EBV does not simply infect and clear: it permanently establishes itself inside B cells (a type of immune cell) and persists there for life [163].

The path from acute EBV infection to ME/CFS likely involves two overlapping processes. First, the immune system’s response to the virus can generate antibodies that cross-react with normal body tissues — a process called **molecular mimicry**. Structural similarities between EBV proteins and human proteins mean that some anti-EBV antibodies inadvertently target the patient’s own cells, particularly receptors that regulate the autonomic nervous system [164]. Second, EBV-infected B cells can cross the blood-brain barrier and contribute to low-level neuroinflammation, potentially producing symptoms of cognitive impairment and sensory sensitivity. The detailed immunological cascade — including how EBV drives autoantibody production and how LMP1-expressing B cells enter the central nervous system — is examined in Section 7.5.1 and in the novel hypotheses presented in Section 14.20.

5.7.3 SARS-CoV-2 (COVID-19)

COVID-19 has created a large new cohort of post-infectious ME/CFS patients, providing an unprecedented opportunity to study the disease from a known, precisely dated onset. Several mechanisms appear to contribute to the transition from acute COVID-19 to chronic illness.

The most distinctive feature of SARS-CoV-2 compared to other ME/CFS triggers is **spike protein persistence**. Research using advanced imaging techniques has detected SARS-CoV-2 spike protein in skull, meninges, and brain tissue of COVID-19 patients long after the acute infection resolved, co-localised with activated immune cells [165]. This persistent viral protein — even in the absence of replicating virus — appears to sustain local inflammation. In addition, SARS-CoV-2 infects and damages the endothelial cells lining blood vessels, impairing circulation and contributing to the formation of abnormal microclots that may restrict oxygen delivery to tissues [166]. The combination of neuroinflammation from spike persistence and microvascular damage from endothelial injury may together account for the cognitive and energy-generation deficits that characterise post-COVID ME/CFS. Long COVID shares the same immune abnormalities as ME/CFS [167], supporting the conclusion that both represent manifestations of the same post-infectious pathological process.

5.7.4 Human Herpesvirus 6 (HHV-6)

HHV-6 has two properties that make it a particularly disruptive ME/CFS trigger. First, it is **neurotropic** — it infects brain tissue directly, unlike most ME/CFS-associated viruses that primarily target peripheral immune cells [163]. Second, it has a documented ability to interfere with **mitochondrial function** in infected cells, disrupting the energy-production machinery at the cellular level. HHV-6 can also integrate its genetic material into chromosomes, creating a persistent reservoir that standard antiviral drugs cannot eliminate. Reactivation of this chromosomally integrated virus can occur during periods of immune stress, potentially driving repeated exacerbations in ME/CFS patients with HHV-6 as their primary trigger (see Section 7.5.1).

5.7.5 Enteroviruses

Enteroviruses — including coxsackieviruses and echoviruses — are associated with ME/CFS both historically (epidemic outbreaks of what was then called “epidemic neuromyasthenia”) and through modern molecular studies. Their distinctive feature is **tissue reservoir persistence**: viral RNA has been detected in muscle biopsies and gastrointestinal tissue of ME/CFS patients years after the acute infection [163]. Unlike herpesviruses, enteroviruses do not establish conventional latency; instead, they appear to maintain a low-level, smouldering infection in tissue sanctuaries where the immune system cannot fully reach them. This persistent infection sustains immune activation without producing the acute symptoms of active viral illness, creating a situation where the patient has an ongoing immune burden that is invisible on standard clinical testing.

‡ Limitation 10: Trigger-Specific Mechanisms: Narrative Synthesis, Not Direct Evidence

The pathogen-specific mechanisms described above (EBV molecular mimicry, spike protein persistence, HHV-6 mitochondrial interference, enteroviral tissue reservoirs) are each supported by individual studies, but no study has demonstrated the complete causal chain from specific pathogen to specific ME/CFS mechanism to chronic disease maintenance in the same patients. The “common pathway” synthesis is the authors’ interpretive framework; whether these diverse triggers genuinely converge on a single pathological state or produce distinct subtypes with overlapping symptoms remains an open question.

5.7.6 The Common Pathway: Vicious Cycles and Chronicity

Across all these triggers, a shared final pathway emerges: the acute infection initiates immune changes that, in susceptible individuals, do not self-terminate. Instead, they lock into self-reinforcing cycles. Immune cells remain activated and produce inflammatory signals; those signals impair energy metabolism; the impaired energy metabolism limits the immune system’s ability to clear residual virus or resolve inflammation; and the residual viral activity or autoantibodies sustain the immune activation. Multiple interlocking cycles of this kind maintain the disease state indefinitely, even as the original triggering pathogen becomes undetectable [163].

This explains a feature of ME/CFS that patients and their families often find bewildering: why does the illness persist when the infection is “gone”? The answer is that the pathological state has become self-sustaining. The trigger initiated the disease, but the trigger no longer *drives* it. The integrated pathophysiological model — showing how immune dysfunction, metabolic failure, autonomic dysregulation, and potential autoimmunity lock together — is developed in detail in Section 7.9.5 and in the multi-lock trap hypothesis (Section 14.16).

Part II

Pathophysiology and Biological Mechanisms

This part explores the biological underpinnings of ME/CFS, from well-established phenomena to emerging theories. We examine:

- **Known phenomena:** Mechanisms with strong research support
- **Suspected phenomena:** Plausible mechanisms with preliminary evidence
- **Related phenomena:** Connections to other conditions (allergies, autoimmune diseases, etc.)
- **Biochemical processes:** Detailed molecular and cellular mechanisms

Understanding these mechanisms is crucial for developing targeted treatments and explaining the diverse symptomatology of ME/CFS.

6 Energy Metabolism and Mitochondrial Function

Energy production impairment is a central feature of ME/CFS pathophysiology and likely underlies the characteristic fatigue and post-exertional malaise that define the illness. The 2024 NIH deep phenotyping study by Walitt et al. provided important metabolomic data from cerebrospinal fluid analysis, documenting alterations in catecholamine and tryptophan pathway metabolites that link energy metabolism dysfunction to neurological symptoms [5]. This chapter examines the detailed biochemical processes involved in cellular energy production and the multiple levels at which these processes appear disrupted in ME/CFS.

6.1 Cellular Energy Production Overview

6.1.1 ATP Synthesis

Adenosine triphosphate (ATP) is the universal energy currency of cells, powering virtually all cellular processes. ATP is generated through three interconnected pathways:

Glycolysis

Glycolysis occurs in the cytoplasm and converts glucose to pyruvate:

- **Substrate:** One glucose molecule (6 carbons)
- **Products:** Two pyruvate molecules (3 carbons each), 2 ATP (net), 2 NADH
- **Oxygen requirement:** None (anaerobic process)
- **Rate:** Fast but relatively inefficient

Glycolytic intermediates also provide substrates for biosynthetic pathways (amino acids, lipids, nucleotides), making glycolysis central to cellular metabolism beyond energy production.

Krebs Cycle (Citric Acid Cycle)

The Krebs cycle occurs in the mitochondrial matrix and completes glucose oxidation:

- **Substrate:** Acetyl-CoA (derived from pyruvate, fatty acids, or amino acids)
- **Products per acetyl-CoA:** 3 NADH, 1 FADH₂, 1 GTP (equivalent to ATP), 2 CO₂
- **Function:** Generates reducing equivalents (NADH, FADH₂) for electron transport chain

- **Regulation:** Controlled by substrate availability, product inhibition, and allosteric regulators

Electron Transport Chain and Oxidative Phosphorylation

The electron transport chain (ETC) in the inner mitochondrial membrane generates the majority of cellular ATP:

- **Complex I (NADH dehydrogenase):** Accepts electrons from NADH, pumps protons
- **Complex II (Succinate dehydrogenase):** Accepts electrons from FADH₂, does not pump protons
- **Complex III (Cytochrome bc₁):** Transfers electrons to cytochrome c, pumps protons
- **Complex IV (Cytochrome c oxidase):** Transfers electrons to O₂ (forming H₂O), pumps protons
- **Complex V (ATP synthase):** Uses proton gradient to synthesize ATP from ADP + P_i

Complete oxidation of one glucose molecule yields approximately 30–32 ATP, though actual yield varies with cellular conditions.

Figures 6.1 and 6.2 illustrate normal ATP production and the multiple impairment points in ME/CFS. ATP deficit cascades into multi-system dysfunction affecting muscle, brain, immune, cardiovascular, and autonomic systems. This multi-system cascade is synthesized with other pathophysiological mechanisms in Chapter 13, particularly in the discussion of energy-immune-autonomic interactions (Section 13.3).

6.1.2 Normal Energy Metabolism


Baseline ATP Requirements

Different tissues have vastly different energy demands:

- **Brain:** 20–25% of resting metabolic rate despite 2% of body mass
- **Heart:** Continuously contracting, requires constant ATP supply
- **Skeletal muscle:** Variable demand; enormous increase during exercise
- **Immune cells:** High energy demand during activation
- **Liver:** Metabolic hub with substantial ATP consumption

The human body produces and consumes approximately 40–70 kg of ATP daily, with turnover occurring every few seconds.

Normal Cellular Energy Production




ME/CFS: Impaired Energy Production

Figure 6.2: ME/CFS energy production dysfunction and systemic consequences.

Energy Demands During Exertion

Physical activity dramatically increases ATP demand:

- **Muscle ATP consumption:** Can increase 100-fold during maximal exercise
- **Immediate energy:** Phosphocreatine provides seconds of buffering
- **Short-term:** Glycolysis provides rapid but limited ATP
- **Sustained activity:** Requires oxidative phosphorylation
- **Substrate shift:** From glucose to increasing fatty acid utilization

Recovery Processes

Following exertion, energy systems must be restored:

- **Oxygen debt repayment:** Elevated metabolism to restore baseline
- **Phosphocreatine resynthesis:** Rapid recovery (seconds to minutes)
- **Glycogen resynthesis:** Hours to days depending on depletion
- **Lactate clearance:** Conversion back to glucose (Cori cycle)
- **Protein synthesis:** Repair of exercise-induced damage

6.2 Mitochondrial Dysfunction in ME/CFS

Mitochondria are increasingly recognized as central to ME/CFS pathophysiology, with evidence for dysfunction at multiple levels.

6.2.1 Evidence for Mitochondrial Impairment

Studies Showing Reduced ATP Production

Multiple lines of evidence support impaired ATP generation:

- **Lymphocyte studies:** Reduced ATP production in peripheral blood mononuclear cells
- **Muscle biopsies:** Abnormal mitochondrial morphology and function in some patients
- **Metabolomic profiles:** Patterns consistent with impaired oxidative phosphorylation
- **Exercise studies:** Early transition to anaerobic metabolism (reduced anaerobic threshold)

The ATP Profile Test One proposed biomarker approach measures:

- ATP concentration in neutrophils
- ATP production efficiency
- Mitochondrial membrane potential

Studies using this approach have found reduced ATP levels and impaired efficiency in ME/CFS patients, though methodological debates continue.

★ Achievement 1: Multi-Omics Biomarker Panel: Integrated Energy-Immune-Vascular Dysfunction

A landmark 2025 study by Heng et al. [12], published in *Cell Reports Medicine*, applied multi-omics analysis to 61 ME/CFS patients (Canadian Criteria) matched with 61 healthy controls, revealing coordinated dysfunction across energy metabolism, immune function, and vascular systems. Key energy metabolism findings included elevated adenosine monophosphate (AMP) and adenosine diphosphate (ADP) in white blood cells, with median AMP levels of 312.2 nM in ME/CFS versus 147.2 nM in controls, reduced ATP/ADP ratio consistent with decreased ATP generation and cellular energy stress, and abnormal nicotinamide adenine dinucleotide (NAD⁺) metabolism affecting cellular energy production.

The study identified a predictive model comprising seven biological variables that distinguished ME/CFS patients with 85.2% sensitivity, 96.7% specificity, and 91% accuracy. These seven biomarkers span adenosine metabolism (AMP), immune functions (cDC1, LYVE1, IGHG2), and vascular factors (FN1, VWF, THBS1)—demonstrating that energy dysfunction in ME/CFS is not isolated but integrated with immune and vascular abnormalities. This multi-system integration suggests that future treatments may need to target energy metabolism, immune maturation, and vascular health simultaneously. The implications of this coordinated dysfunction for treatment strategy are discussed in Chapter 13, particularly in the context of the Energy-Immune-Autonomic Triad (Section 13.3). For the systems biology perspective on this biomarker panel, see Achievement 13.7.1 in Chapter 13.

△ Warning 1: Replication Status

Single study (Heng 2025, n=61 pairs) from one Australian center. No independent cohort has replicated the multi-omics biomarker panel or its diagnostic accuracy. The CART-derived 7-biomarker model requires prospective validation in independent populations with diverse demographics and diagnostic criteria before clinical application. Independent replication required.

Electron Microscopy Findings

Ultrastructural examination of mitochondria has revealed:

- **Abnormal morphology:** Swollen, disrupted cristae structure

- **Variable size:** Both enlarged and fragmented mitochondria
- **Reduced number:** Decreased mitochondrial density in some tissues
- **Intramuscular abnormalities:** Changes in muscle biopsy specimens

Functional Assays

Direct measurement of mitochondrial function shows:

- **Respirometry:** Reduced oxygen consumption rates in some studies
- **Enzyme activities:** Variable findings for individual ETC complexes
- **Membrane potential:** May be altered, affecting ATP synthesis efficiency
- **Calcium handling:** Impaired mitochondrial calcium uptake

Biomarkers of Mitochondrial Dysfunction

Several biomarkers indicate mitochondrial stress:

- **Lactate:** Elevated at rest or with minimal exertion
- **Pyruvate:** Altered lactate/pyruvate ratio
- **Organic acids:** Abnormal urinary organic acid patterns
- **Acylcarnitines:** Reflecting impaired fatty acid oxidation
- **Coenzyme Q10:** Sometimes reduced

6.2.2 Mechanisms of Mitochondrial Damage

Oxidative Stress

Reactive oxygen species (ROS) damage mitochondrial components:

- **Electron leakage:** Complexes I and III leak electrons that generate superoxide
- **Mitochondrial DNA damage:** mtDNA lacks histones and has limited repair
- **Protein oxidation:** Damages ETC components
- **Lipid peroxidation:** Disrupts inner membrane integrity
- **Vicious cycle:** Damaged mitochondria produce more ROS

Figures 6.3 and 6.4 illustrate how oxidative stress creates a self-perpetuating cycle in ME/CFS, where excessive ROS production (5–10% electron leakage vs. normal 2%) combined with depleted antioxidants leads to progressive damage.

Normal Oxidative Stress Balance





Figure 6.3: Normal oxidative stress homeostasis with balanced ROS production and neutralization.

ME/CFS: Oxidative Stress Vicious Cycle



Self-perpetuating damage cascade:

- Damaged mitochondria leak more electrons (5–10% vs. <2%)
 - Excess ROS damages lipids (4-HNE, MDA), proteins, and DNA
 - Damaged components further impair mitochondrial function
 - Depleted antioxidants (GSH ↓30%) cannot neutralize ROS
- Breaking this cycle requires both reducing ROS production and restoring antioxidant capacity.

Figure 6.4: ME/CFS oxidative stress vicious cycle with self-perpetuating damage.

Calcium Dysregulation

Mitochondria buffer cytosolic calcium and use it for signaling:

- **Calcium overload:** Excessive mitochondrial calcium triggers permeability transition
- **ER-mitochondria crosstalk:** Abnormal calcium transfer between organelles
- **Apoptosis signaling:** Calcium overload can trigger cell death pathways
- **Enzyme regulation:** Many mitochondrial enzymes are calcium-sensitive

Mitochondrial DNA Alterations

Mitochondrial DNA (mtDNA) is vulnerable to damage:

- **Mutations:** Point mutations accumulate with oxidative stress
- **Deletions:** Large deletions impair multiple ETC components
- **Copy number:** Altered mtDNA copy number in some ME/CFS studies
- **Heteroplasmy:** Mixture of normal and mutant mtDNA

Impaired Mitophagy

Mitophagy removes damaged mitochondria:

- **PINK1/Parkin pathway:** Marks damaged mitochondria for degradation
- **Impaired clearance:** May allow dysfunctional mitochondria to persist
- **Accumulation:** Damaged mitochondria continue producing ROS
- **Quality control failure:** Network of damaged organelles

WASF3 and ER Stress: A Specific Molecular Mechanism

A 2023 study by Wang et al., using muscle biopsies from the NIH intramural ME/CFS cohort, identified a specific molecular pathway linking cellular stress to mitochondrial dysfunction [2].

★ Achievement 2: ER Stress–WASF3–Mitochondrial Dysfunction Pathway: Drug-gable Mechanism

A 2023 study by Wang et al. [2], using muscle biopsies from the NIH intramural ME/CFS cohort, identified a specific molecular pathway linking cellular stress to mitochondrial dysfunction. WASF3 (Wiskott-Aldrich syndrome protein family member 3) was significantly elevated in ME/CFS patient muscle biopsies compared to controls. This protein, when overexpressed, localizes to mitochondria and disrupts respiratory chain function.

The study revealed a mechanistic chain: endoplasmic reticulum (ER) stress activation drives increased WASF3 expression, which then translocates to mitochondria where it interferes with respiratory chain supercomplex assembly, particularly affecting Complex IV (cytochrome c oxidase), resulting in decreased oxygen consumption and reduced exercise endurance. Critically, pharmacologic inhibition of ER stress in patient-derived cells improved mitochondrial function, suggesting this pathway represents a potentially druggable target. ER stress modulators or WASF3 inhibitors could restore normal mitochondrial respiration.

The WASF3 mechanism provides a molecular explanation for several ME/CFS features: viral infection can trigger ER stress through viral protein accumulation (explaining post-infectious onset), once established ER stress can become self-perpetuating (explaining chronic persistence), and Complex IV impairment directly limits oxidative capacity (explaining exercise intolerance and reduced $\text{VO}_{2\text{peak}}$ observed in CPET studies). This finding bridges the gap between cellular stress responses and the clinical manifestation of exercise intolerance, providing mechanistic support for the energy deficit model of ME/CFS.

△ **Warning 2: Replication Status**

Single study (Hwang 2023) from the NIH intramural ME/CFS cohort. While the mechanism was validated in cell culture (pharmacologic ER stress inhibition rescued mitochondrial function), no independent cohort has replicated the WASF3 elevation finding in ME/CFS muscle biopsies. The “druggable target” claim remains preclinical. Independent replication in patient tissue is required before therapeutic development.

A 2025 muscle biopsy study in long COVID and ME/CFS patients confirmed elevated WASF3 and ER stress markers with decreased Complex IV subunits, and critically demonstrated that these abnormalities differ from bed rest-induced deconditioning [168] (Appendix I, Section I.11.2). A multi-modal study in *Cell Reports Medicine* further showed reduced ATP/ADP ratios in immune cells, linking energy deficits to immune dysfunction [169] (Section I.3.5).

6.2.3 Consequences of Energy Deficits

Cellular Function Impairment

Inadequate ATP affects all cellular processes:

- **Ion pumps:** Na^+/K^+ -ATPase consumes 20–40% of cellular ATP
- **Protein synthesis:** Highly energy-intensive process
- **Cell signaling:** Many signaling pathways require ATP
- **Membrane function:** Active transport and vesicle trafficking

Tissue-Specific Effects

Different tissues manifest energy deficits differently:

Muscle

- Weakness and fatigue with minimal exertion
- Early lactate accumulation
- Delayed recovery from activity
- Post-exertional pain and soreness

Brain

- Cognitive dysfunction (“brain fog”)
- Reduced neurotransmitter synthesis
- Impaired synaptic function
- Vulnerability to excitotoxicity

Immune Cells

- Impaired T cell activation (requires metabolic reprogramming)
- Reduced NK cell cytotoxicity
- Abnormal cytokine production
- Ineffective pathogen clearance


Connection to Post-Exertional Malaise

Mitochondrial dysfunction provides a compelling explanation for PEM:

1. **Limited reserve:** Baseline energy production is already compromised
2. **Exercise stress:** Activity depletes already-limited ATP stores
3. **Oxidative burst:** Exercise generates additional ROS, damaging mitochondria further
4. **Delayed recovery:** Impaired mitophagy and biogenesis slow restoration
5. **Cumulative damage:** Each exertion may worsen mitochondrial function
6. **Symptom cascade:** Energy deficit affects multiple organ systems

Figures 6.5 and 6.6 illustrate the critical distinction between normal exercise response (rapid recovery, positive adaptation) and ME/CFS PEM (ATP crisis, maladaptive inflammatory cascade, delayed deterioration). Repeated PEM episodes cause progressive decline.

Normal Exercise Response



Key characteristics:

- Activity within capacity triggers beneficial adaptations
- Recovery completes within hours to 1 day
- Progressive overload increases fitness (training effect)




Figure 6.6: ME/CFS post-exertional malaise mechanism with harmful vicious cycle.

The Effort-Performance Disconnect: Physiological Mechanisms The profound subjective experience described in Section 2.1—the sensation of “giving everything” yet achieving minimal output—has direct physiological correlates that distinguish ME/CFS from psychological disorders or deconditioning.

Central Nervous System Effort Signaling:

The brain’s effort-generating systems appear to function normally or even hyperactivate in ME/CFS:

- **Motor cortex activation:** fMRI studies suggest normal or increased motor cortex activation during attempted movement
- **Catecholamine mobilization attempts:** The brain attempts to mobilize energy reserves through sympathetic activation
- **Subjective intensity:** The sense of maximal effort reflects genuine CNS activation and stress response engagement
- **Central command:** Motor planning and initiation circuits generate normal or excessive drive

The Walitt et al. 2024 NIH study documented altered effort preference rather than reduced effort capability [5]. ME/CFS patients can generate effort signals, but the consequences of doing so (PEM) appropriately modify behavior. This represents adaptive learning, not primary motivation deficit.

Peripheral Energy Production Failure:

Despite normal or excessive central drive, peripheral tissues cannot respond proportionally:

- **Mitochondrial ATP deficit:** Muscle cells cannot generate sufficient ATP to sustain contraction despite receiving motor neuron signals
- **Ion pump failure:** Inadequate ATP impairs Na^+/K^+ -ATPase function, disrupting muscle excitability and contraction
- **Calcium handling impairment:** Energy-dependent calcium reuptake into sarcoplasmic reticulum fails, preventing muscle relaxation and subsequent contraction
- **Metabolite accumulation:** Lactate, hydrogen ions, and other metabolites accumulate rapidly, triggering muscle pain and afferent signaling
- **Neuromuscular transmission stress:** Repeated activation with insufficient recovery depletes neurotransmitter and impairs synaptic function

Cardiovascular Oxygen Delivery Limitations:

The two-day CPET data demonstrate that oxygen delivery and utilization fail during and after exertion [1]:

- **Chronotropic incompetence:** Heart rate fails to increase appropriately, limiting cardiac output
- **Reduced stroke volume:** Autonomic dysfunction impairs venous return and cardiac filling

- **Impaired oxygen extraction:** Oxygen pulse (VO_2/HR) declines on Day 2, suggesting reduced tissue oxygen uptake
- **Ventilatory limitation:** Reduced ventilation limits oxygen availability even when respiratory muscles receive motor commands

The Subjective-Objective Mismatch Explained:

This creates a situation unique to ME/CFS:

1. **Central effort generation:** Brain generates normal or maximal effort signals → subjectively feels like “giving everything”
2. **Peripheral energy failure:** Muscles receive commands but cannot execute due to ATP deficit, ion pump failure, oxygen delivery limitation → minimal force production, minimal work output
3. **Afferent feedback:** Massive signaling from muscle (metabolite accumulation, tissue hypoxia, cellular stress) returns to brain → reinforces sensation of extreme exertion
4. **Autonomic stress response:** Sympathetic activation (elevated heart rate, norepinephrine release attempt) further intensifies subjective sense of emergency
5. **Observable output:** Despite all this internal activation and distress, actual work performed is minimal → external observers see “not trying hard enough”

Learned Helplessness as Accurate Pattern Recognition:

The development of learned helplessness in ME/CFS differs fundamentally from learned helplessness in depression:

- **Accurate perception:** Patients accurately perceive that their maximal effort does not produce expected outcomes—this is not a cognitive distortion but a direct experiential truth
- **Appropriate behavioral adaptation:** Reducing effort expenditure after learning it produces crashes represents adaptive learning, not pathological avoidance
- **Physiological validation:** Two-day CPET objectively documents that effort Day 1 produces measurable impairment Day 2, validating patient perception
- **Controllability assessment:** In classic learned helplessness paradigms, outcomes are truly uncontrollable; in ME/CFS, outcomes are controllable through limitation (pacing works), making the adaptation rational

The psychological distress arises not from cognitive distortion but from accurate recognition of one’s physiological limitations in a world structured around normal energy availability. The helplessness is *realistic*—patients genuinely cannot reliably produce normal output despite normal or excessive subjective effort expenditure.

Vulnerability as Physiological Reality:

The sense of extreme vulnerability—“I wouldn’t amount to shit in a fight”—reflects accurate assessment of current physiological capacity:

- **Energy unavailability for defense:** Fight-or-flight responses require massive ATP mobilization; ME/CFS patients cannot sustain this

- **Delayed consequences:** Any acute energy expenditure (fleeing danger, defending self) would trigger severe PEM, leaving the patient even more vulnerable for days to weeks afterward
- **Dependence on others:** Inability to reliably generate protective physical responses creates realistic dependence
- **Identity disruption:** For patients previously defined by physical capability, this represents genuine loss, not negative self-perception

This vulnerability is not imagined or exaggerated—it is a direct consequence of documented metabolic, cardiovascular, and mitochondrial dysfunction that prevents reliable energy mobilization on demand.

Exercise-Induced Metabolic Failure: Two-Day CPET Evidence

The most compelling objective evidence for exercise-induced metabolic failure comes from two-day cardiopulmonary exercise testing (CPET) protocols. Unlike single-day assessments that may be confounded by deconditioning or effort, the two-day protocol documents the failure to reproduce initial performance after 24 hours—a hallmark of post-exertional malaise [1].

Two-Day CPET Protocol and Rationale The two-day CPET protocol requires maximal exercise tests on consecutive days, separated by 24 hours. Healthy individuals and those with deconditioning typically maintain or slightly improve performance on Day 2 after familiarization with the protocol. In contrast, ME/CFS patients show consistent, reproducible declines.

★ Achievement 3: Two-Day CPET: Objective Validation of Post-Exertional Malaise

In the largest rigorous two-day CPET study to date, Keller and colleagues examined 84 ME/CFS participants (Canadian Criteria) and 71 sedentary controls across multiple sites [1]. The study design included a matched subset (55 pairs) controlled for sex, age, and baseline aerobic capacity, allowing assessment of whether observed abnormalities were attributable to deconditioning.

ME/CFS participants demonstrated consistent, reproducible declines in multiple cardiopulmonary parameters on Day 2: peak oxygen consumption ($\text{VO}_{2\text{peak}}$) declined by 5.3% ($p < 0.01$), work output by 5.5% ($p < 0.01$), ventilation by 7.8% ($p < 0.01$), heart rate by 2.6% ($p < 0.05$), oxygen pulse by 4.0% ($p < 0.05$), and anaerobic threshold VO_2 by 6.7% ($p < 0.05$). In contrast, control participants showed no significant changes in any parameter between Day 1 and Day 2.

Critically, when ME/CFS participants were matched with controls having identical baseline $\text{VO}_{2\text{peak}}$ (aerobic capacity), the abnormal Day 2 responses persisted, demonstrating that impaired recovery is not attributable to fitness level but represents a disease-specific pathophysiological process. This provides the most rigorous objective validation of post-exertional malaise to date, distinguishing ME/CFS from deconditioning and val-

identifying PEM as a reproducible biological phenomenon rather than subjective experience.

△ **Warning 3: Replication Status: Partially Replicated**

The two-day CPET abnormality has been observed in earlier smaller studies (Snell 2013, Stevens 2014), and Keller 2024 is the largest and most rigorously controlled. However, all major studies originate from overlapping research networks. Fully independent replication by groups without prior involvement would strengthen confidence.

Impairment Severity Worsening Based on anaerobic threshold criteria, impairment classification shifted dramatically:

- **Day 1:** 14% classified as severely impaired
- **Day 2:** 27% classified as severely impaired (nearly doubled)

This demonstrates that exertional stress unmasks or exacerbates functional impairment.

Mechanistic Interpretation The two-day CPET findings directly validate the mitochondrial dysfunction framework:

1. **VO₂ peak decline:** Reduced maximal oxygen consumption indicates impaired oxidative metabolism at the tissue level—either reduced oxygen delivery (cardiovascular), oxygen extraction (cellular uptake), or oxygen utilization (mitochondrial dysfunction)
2. **Anaerobic threshold shift:** Earlier reliance on anaerobic metabolism suggests mitochondria cannot meet energy demands through oxidative phosphorylation, forcing premature lactate production
3. **O₂ pulse reduction:** Oxygen pulse (VO₂/heart rate) reflects stroke volume or oxygen extraction; its decline suggests either cardiac dysfunction or impaired peripheral oxygen utilization
4. **Ventilatory dysfunction:** Reduced ventilation at maximal effort may reflect central respiratory drive impairment (consistent with brainstem/autonomic dysfunction) or metabolic signaling abnormalities
5. **Chronotropic incompetence:** Reduced heart rate response indicates autonomic nervous system dysregulation affecting cardiac control

Autonomic Dysregulation as Primary Mechanism Keller and colleagues concluded that **autonomic nervous system dysregulation** affecting blood flow and oxygen delivery represents the primary mechanism linking these abnormalities [1]. This integrates with the Walitt study's findings of reduced central catecholamines (Section 6.6)—catecholamines are essential for autonomic cardiovascular regulation during exercise.

Complementarity with Walitt 2024 NIH Study The two-day CPET findings complement the NIH deep phenotyping study [5]:

- **Walitt:** Documented reduced CSF catecholamines, altered effort preference due to temporoparietal junction dysfunction, metabolic abnormalities, and single-day CPET showing reduced VO_2 peak and chronotropic incompetence
- **Keller:** Demonstrated that exercise Day 1 produces measurable physiological impairment on Day 2, validating PEM as a reproducible phenomenon with objective correlates

Together, these studies establish that:

1. Central catecholamine deficiency impairs effort generation and autonomic control
2. Exertional stress on Day 1 further compromises already-limited energy metabolism
3. Recovery processes fail to restore baseline function within 24 hours
4. The functional impairment is measureable, reproducible, and distinct from deconditioning

Clinical Implications for Activity Management The two-day CPET findings provide a quantitative foundation for pacing strategies:

- **Heart rate thresholds:** Staying below anaerobic threshold (often estimated as AT – 15 bpm) may prevent Day 2 impairment
- **Recovery periods:** Activity sufficient to trigger metabolic stress requires >24 hours for restoration
- **Graded exercise therapy contraindication:** Progressive increases in exertion worsen measurable physiological function rather than improving fitness
- **Disability documentation:** Two-day CPET provides objective, reproducible evidence of functional impairment for benefits/insurance claims

Recovery Kinetics Beyond 24 Hours While the Keller study assessed only 24-hour recovery, clinical observations and Cornell Center research suggest full restoration requires approximately **13 days** for ME/CFS patients compared to ~2 days for sedentary controls. This prolonged recovery period likely reflects:

- Impaired mitophagy delaying removal of damaged mitochondria
- Reduced mitochondrial biogenesis slowing replacement
- Persistent oxidative stress from the exertional episode
- Systemic inflammation triggered by metabolic stress

6.3 Selective Energy Dysfunction: The CNS-Dependency Hypothesis

While the preceding sections document energy production impairment across multiple tissues, emerging evidence suggests ME/CFS may not represent *global* cellular energy failure but rather *selective* dysfunction affecting primarily CNS-dependent and demand-responsive processes while sparing autonomous steady-state peripheral functions.

~ Hypothesis 1: Selective Energy Dysfunction

ME/CFS involves selective impairment of CNS-dependent and demand-responsive processes while sparing autonomous steady-state peripheral functions. This pattern suggests either primary brain energy dysfunction affecting downstream coordination, or failure of demand-response coupling mechanisms, rather than uniform global cellular energy failure.

Cerebral blood flow during orthostatic challenge exemplifies this selectivity: 91% of patients with normal resting hemodynamics show abnormal CBF reduction during demand [4], while baseline perfusion is often preserved. Brain hypometabolism has also been documented [73], though replication remains incomplete (see Chapter 8 Section 8.1.5). Together, these findings support coordination failure rather than global energy deficit as the underlying pathophysiology.

~ Hypothesis 2: Immune Cell Energy Starvation Creates a Viral Persistence Niche

Certainty: 0.40. Supported by documented T cell metabolic dysfunction in ME/CFS [170] and the established bioenergetic requirements of cytotoxic killing. However, the causal link between immune cell energy deficit and viral persistence has not been directly tested in ME/CFS.

This hypothesis proposes that the mitochondrial dysfunction documented in ME/CFS immune cells [170] is not merely a symptom but a *cause* of viral persistence: metabolically impaired T cells and NK cells lack sufficient ATP to sustain cytotoxic function, creating a permissive niche where virus-harboring cells survive despite immune recognition. This generates a self-reinforcing trap: viral persistence → chronic immune activation → T cell exhaustion and metabolic failure → inadequate viral clearance → more persistence.

Mechanistic Basis. Cytotoxic killing is among the most energy-intensive immune functions: degranulation, perforin pore formation, and granzyme delivery each require substantial ATP. Mandarano et al. demonstrated that ME/CFS CD8+ T cells fail to switch from oxidative phosphorylation to glycolysis upon activation (the Warburg effect) [170]. Without this metabolic reprogramming, effector function is impaired even if the cell correctly recognizes its target. The Kol et al. FIP data [171] provide indirect support: antiviral treatment reduced viral load but did not eliminate virus from lymphocytes, suggesting host immunity is needed for final clearance—and if that immunity is metabolically compromised, clearance never completes.

Testable Predictions.

1. In vitro supplementation with metabolic support (D-ribose, CoQ10, or mitochondria-targeted antioxidants such as MitoQ) will restore ME/CFS NK cell and CD8+ T cell cytotoxic function in killing assays against virus-infected target cells.
2. ME/CFS patients on combined antiviral + metabolic support therapy will show greater viral clearance (measured by pathogen-specific PCR or T-SPOT) than patients on antiviral therapy alone.
3. The degree of T cell metabolic impairment (measured by Seahorse metabolic flux analysis) will inversely correlate with viral clearance efficiency in longitudinal studies.
4. Patients with the most severe NK cell metabolic dysfunction will show the highest persistent viral antigen levels.

Falsifiability. This hypothesis would be falsified if (a) restoring immune cell metabolism via in vitro supplementation does not improve cytotoxic function against virus-infected targets, or (b) ME/CFS patients with documented viral persistence show normal immune cell bioenergetics. It would also be weakened if antiviral monotherapy achieves complete viral clearance without metabolic support, suggesting the energy deficit is not rate-limiting for clearance.

Treatment Implications. If confirmed, this hypothesis argues that antiviral therapy alone is insufficient—it must be combined with metabolic support targeting immune cell mitochondria. This reframes mitochondrial supplementation (CoQ10, NAD⁺ precursors, D-ribose) not merely as energy support for the patient, but as *immune-enabling therapy* that restores the body's ability to clear persistent infection. It also provides mechanistic rationale for the Selective Energy Dysfunction hypothesis (Hypothesis 6.3): immune surveillance is a demand-responsive, CNS-coordinated process that fails under energy constraint.

Limitations. The hypothesis assumes viral persistence is a primary driver in a substantial proportion of ME/CFS cases, which may not hold for non-viral-onset subgroups. The in vitro prediction (metabolic supplementation restoring killing) may not translate to in vivo efficacy due to bioavailability and tissue distribution constraints. The self-reinforcing loop makes it difficult to determine which arm (metabolic failure or viral persistence) is primary.

6.3.1 Evidence for Selectivity

The distinction between *preserved* and *impaired* processes follows a consistent pattern:

Impaired Processes (CNS-Dependent + Demand-Responsive)

- **Voluntary muscle exertion:** Requires motor cortex coordination + scaling to demand
- **Cognitive effort:** Inherently CNS-based + scales with task complexity
- **Orthostatic adaptation:** Requires autonomic coordination + responds to positional demand
- **Adaptive immune responses:** Requires CNS-immune signaling + scales to antigen challenge
- **Temperature regulation:** Requires hypothalamic coordination + responds to environmental demands

Preserved Processes (Truly Autonomous + Locally Controlled)

Observation 36 (Apparent Preservation of Autonomous Processes). Clinical observation suggests the following autonomous, locally-controlled processes continue at apparently normal rates in ME/CFS despite severe systemic symptoms:

- **Hair follicle cycling:** Operates independent internal Cori cycle; no CNS coordination required
- **Nail growth:** Locally controlled keratin synthesis
- **Baseline cellular metabolism:** Homeostatic processes not requiring demand scaling
- **Wound healing under occlusion:** Locally mediated by growth factors
- **Basal immune surveillance:** Constitutive function not requiring coordinated scaling

Formal documentation of these observations is lacking in the ME/CFS literature, representing a gap requiring systematic validation. However, their apparent preservation contrasts markedly with profound impairment of CNS-coordinated demand-responsive functions. If global mitochondrial dysfunction were present, these energy-requiring processes should also be impaired.

Critical Implication If ME/CFS were global mitochondrial dysfunction, all energy-requiring processes—including hair growth—should be affected proportionally. The preservation of truly autonomous peripheral processes suggests the pathology may lie in *energy coordination and allocation* rather than *energy production capacity* per se.

6.3.2 The Demand-Response Failure Pattern

A consistent finding across multiple physiological systems is preserved baseline function with impaired challenge response [172, 4, 1]:

- **Cardiovascular:** Resting cardiac parameters often normal; profound dysfunction during orthostatic or exercise challenge (Section 10.2.4)
- **Cognitive:** Basic language comprehension preserved; executive function and working memory (high-demand) severely impaired [5]
- **Autonomic:** Baseline HRV present; blunted response to physiological challenges

- **Cerebral perfusion:** 91% of patients with normal resting HR/BP show abnormal cerebral blood flow reduction during tilt testing [4]

This pattern is consistent with intact energy production capacity but impaired ability to *mobilize* energy in response to demand—a coordination failure rather than a production failure.

6.3.3 Mechanistic Implications

Speculation 1 (Brain as Energy Coordination Bottleneck). The near-universal cognitive dysfunction and documented brain hypometabolism [73, 5] suggest CNS energy crisis may be the primary pathophysiological event. The brain consumes 20–25% of resting energy despite comprising only 2% of body mass (Section 6.1.2), making it uniquely vulnerable to energy constraint. Failure of the brain to coordinate peripheral demand-responsive processes could explain the selective dysfunction pattern: autonomous processes continue because they don't require CNS coordination, while CNS-coordinated responses (exercise capacity, orthostatic tolerance, cognitive effort) fail because the coordinating organ itself is energy-depleted.

This model explains why pharmacological bypass of autonomic coordination (midodrine, fludrocortisone) can partially restore orthostatic function—the peripheral targets respond when appropriately stimulated, suggesting the dysfunction is in *coordination* rather than *peripheral capacity*.

See Chapter 8 Section 8.1.2 for expanded discussion of brain-centric pathophysiology, and Chapter 10 Section 10.2.4 for cerebral blood flow evidence during orthostatic challenge.

6.3.4 Therapeutic Implications of Selective Dysfunction

If ME/CFS involves selective coordination failure rather than global energy production deficit, treatment strategies should prioritize:

- **CNS-targeted interventions:** Compounds that cross the blood-brain barrier and support brain energy metabolism specifically, rather than systemic mitochondrial supplements that may not reach the CNS at therapeutic concentrations
- **Autonomic coordination bypass:** Pharmacological agents that directly activate peripheral targets, bypassing impaired CNS signaling (e.g., midodrine for vasoconstriction, fludrocortisone for volume expansion, droxidopa for norepinephrine replacement)
- **Demand management:** Strict pacing to remain within the envelope of available coordination capacity, rather than attempting to increase energy production through exercise or stimulants

Speculation 2 (CNS Penetration as Limiting Factor). This reframing suggests that failed trials of systemic energy supplements (CoQ10, carnitine, B-vitamins) may reflect inadequate CNS penetration rather than incorrect therapeutic targets. If brain energy coordination is the primary bottleneck, supplements that do not cross the blood-brain barrier at therapeutic concentrations would be expected to show limited efficacy regardless of their peripheral

effects. This hypothesis is testable through comparative trials of CNS-penetrant versus non-penetrant formulations of the same compounds.

⚠ **Limitation 1: Selective Energy Dysfunction: Preserved Processes Not Formally Documented**

The “selective vs. global” energy dysfunction distinction rests on the claim that hair growth, nail growth, and wound healing are preserved in ME/CFS. As noted in the observation above, formal documentation of these preserved processes is lacking — the claim derives from patient self-reports and clinical impression, not from controlled measurement. Processes classified as “preserved” also have much lower absolute energy demands than those classified as “impaired,” meaning a moderate global energy deficit could produce the same clinical pattern without any selectivity mechanism. For quantitative analysis of this alternative explanation, see the selective dysfunction framework in Section 14.24.

See Part III for detailed treatment protocols, particularly Chapter 21 for pharmacological approaches and Chapter 16 for symptom-specific interventions.

6.3.5 Subtype Considerations

Observation 37 (Subtype Manifestations). The selective dysfunction pattern may manifest differently across patient subgroups. Some patients show primarily CNS-energy deficit (cognitive and autonomic symptoms predominating with relatively preserved peripheral muscle function), while others show primarily peripheral demand-response failure (exercise intolerance and orthostatic symptoms with relatively preserved cognition at rest). These patterns may represent different points along a continuum or distinct pathophysiological subtypes requiring tailored interventions.

Formal subtype analysis based on the selective dysfunction framework is developed in Chapter 14 Section 14.24, including quantitative predictions for dysfunction severity across processes based on their CNS-dependency and demand-responsiveness.

6.4 Oxidative and Nitrosative Stress

Oxidative and nitrosative stress are consistently documented in ME/CFS and likely contribute to both mitochondrial dysfunction and symptom generation.

6.4.1 Reactive Oxygen Species (ROS)

Sources of ROS in ME/CFS

Multiple sources generate excess ROS:

- **Mitochondrial electron leakage:** Primary source during normal metabolism

- **NADPH oxidase:** Activated by immune stimulation
- **Xanthine oxidase:** Generates superoxide during purine metabolism
- **Uncoupled eNOS:** Produces superoxide instead of NO
- **Inflammatory cells:** Respiratory burst during immune activation

Damage to Cellular Components

ROS damage multiple targets:

- **DNA:** Base modifications, strand breaks, mutations
- **Proteins:** Carbonylation, cross-linking, loss of function
- **Lipids:** Peroxidation of membrane phospholipids
- **Carbohydrates:** Glycation reactions

Antioxidant System Dysfunction

The antioxidant defense system may be compromised:

- **Glutathione:** Often reduced in ME/CFS; critical for detoxification
- **Superoxide dismutase (SOD):** Variable findings
- **Catalase:** May be reduced
- **Vitamins C and E:** Nutritional antioxidants may be depleted
- **Thioredoxin system:** Important for protein redox balance

6.4.2 Reactive Nitrogen Species

Nitric Oxide Metabolism

Nitric oxide (NO) has complex roles in ME/CFS:

- **Normal functions:** Vasodilation, neurotransmission, immune defense
- **iNOS induction:** Inflammatory cytokines induce high NO production
- **NO excess:** Can inhibit mitochondrial respiration
- **eNOS uncoupling:** Produces superoxide instead of NO

Peroxynitrite Formation

When superoxide and NO react, they form peroxynitrite (ONOO^-):

- **Highly reactive:** More damaging than either parent molecule
- **Protein nitration:** 3-nitrotyrosine formation (documented in ME/CFS)
- **Lipid oxidation:** Damages membrane integrity

- **Mitochondrial inhibition:** Irreversibly damages ETC complexes

Effects on Energy Metabolism

Nitrosative stress specifically impairs energy production:

- **Complex I inhibition:** NO competitively inhibits oxygen binding
- **Complex IV inhibition:** NO binds cytochrome c oxidase
- **Aconitase inactivation:** Impairs Krebs cycle
- **Glyceraldehyde-3-phosphate dehydrogenase:** Inhibited by peroxynitrite

6.4.3 Lipid Peroxidation

Membrane Damage

Lipid peroxidation disrupts cellular membranes:

- **Polyunsaturated fatty acids:** Primary targets of peroxidation
- **Chain reactions:** One initiation event triggers multiple peroxidations
- **Membrane fluidity:** Peroxidation rigidifies membranes
- **Permeability changes:** Membranes become leaky

Isoprostanes and Other Markers

Lipid peroxidation products serve as biomarkers:

- **F₂-isoprostanes:** Prostaglandin-like compounds from arachidonic acid peroxidation
- **Malondialdehyde (MDA):** End product of peroxidation
- **4-hydroxynonenal (4-HNE):** Reactive aldehyde that modifies proteins
- **Oxidized LDL:** Marker of lipoprotein oxidation

Studies have found elevated markers of lipid peroxidation in ME/CFS patients, supporting the role of oxidative stress.

6.5 Metabolic Pathways Affected

6.5.1 Amino Acid Metabolism

Tryptophan Metabolism: NIH Study Findings

The NIH deep phenotyping study documented significant abnormalities in tryptophan metabolism in cerebrospinal fluid [5]. Tryptophan is an essential amino acid that serves as precursor for:

- **Serotonin:** Via tryptophan hydroxylase pathway
- **Melatonin:** Via serotonin N-acetyltransferase
- **Kynurenine pathway metabolites:** Via indoleamine 2,3-dioxygenase (IDO)

The Kynurenine Pathway Approximately 95% of dietary tryptophan is metabolized through the kynurenine pathway:

1. **Tryptophan → Kynurenine:** Rate-limiting step; induced by inflammatory cytokines (IFN- γ)
2. **Kynurenine → Kynurenic acid:** Neuroprotective branch (NMDA antagonist)
3. **Kynurenine → 3-hydroxykynurenine → Quinolinic acid:** Neurotoxic branch
4. **Quinolinic acid:** NMDA receptor agonist, excitotoxin, pro-oxidant

ME/CFS Kynurenine Pathway Abnormalities

- Increased IDO activity (driven by inflammation)
- Elevated kynurenine/tryptophan ratio
- Increased neurotoxic metabolites (quinolinic acid, 3-HK)
- Reduced neuroprotective metabolites (kynurenic acid) in some studies
- Depletion of tryptophan available for serotonin synthesis

Implications for Neurotransmitter Production

Tryptophan diversion into the kynurenine pathway reduces serotonin synthesis:

- **Serotonin depletion:** May contribute to mood symptoms, pain, sleep disturbance
- **Melatonin reduction:** May explain sleep-wake cycle disruption
- **Quinolinic acid excess:** May cause excitotoxicity and cognitive dysfunction
- **Oxidative stress:** 3-hydroxykynurenine generates free radicals

Other Amino Acid Abnormalities

Metabolomic studies have identified broader amino acid disturbances:

- **Branched-chain amino acids:** Often altered; important for muscle metabolism
- **Glutamate/glutamine:** Excitatory neurotransmitter precursors
- **Glycine:** Inhibitory neurotransmitter, glutathione precursor
- **Cysteine:** Rate-limiting for glutathione synthesis

6.5.2 Lipid Metabolism

Fatty Acid Oxidation Defects

Fatty acids are the primary fuel for sustained activity:

- **Carnitine shuttle:** Transports fatty acids into mitochondria
- **Beta-oxidation:** Sequential removal of 2-carbon units
- **Acetyl-CoA generation:** Feeds into Krebs cycle

ME/CFS abnormalities include:

- Reduced carnitine levels in some patients
- Elevated acylcarnitines suggesting incomplete oxidation
- Impaired utilization of fatty acids during exercise
- Earlier shift to glucose oxidation

Membrane Lipid Alterations

Cell membrane composition affects function:

- **Phospholipid changes:** Altered fatty acid profiles
- **Reduced omega-3 fatty acids:** May affect inflammation and membrane fluidity
- **Oxidized lipids:** Accumulate due to peroxidation
- **Cholesterol:** May affect membrane rigidity and signaling

Ceramide Metabolism

Ceramides are signaling lipids with metabolic effects:

- **Elevated ceramides:** Found in some ME/CFS studies
- **Insulin resistance:** Ceramides impair insulin signaling
- **Mitochondrial effects:** Can promote apoptosis
- **Inflammation link:** Produced in response to inflammatory signals

6.5.3 Carbohydrate Metabolism

Glucose Utilization

Abnormal glucose handling occurs in ME/CFS:

- **Hypoglycemia symptoms:** Reported by many patients, though blood glucose often normal
- **Impaired glucose uptake:** May affect specific tissues
- **Altered insulin sensitivity:** Variable findings
- **Post-prandial symptoms:** Reactive responses to meals

Lactate Accumulation

Elevated lactate indicates reliance on anaerobic metabolism:

- **Resting lactate:** May be elevated in some patients
- **Exercise lactate:** Earlier and greater accumulation
- **Recovery:** Slower lactate clearance
- **Brain lactate:** Elevated on MR spectroscopy in some studies

Clinical Phenomenology: Similarities to Athletic Post-Exercise States. The chronic lactate accumulation and reliance on anaerobic metabolism in ME/CFS produces a muscle metabolic state remarkably similar to what elite athletes experience temporarily after exhausting physical efforts:

- **Muscle cramping:** ATP depletion prevents proper muscle relaxation; magnesium and calcium handling disrupted
- **“Ready for cramps” sensation:** Persistent partial ATP deficit maintains muscles in pre-cramp tension state
- **Metabolic acidosis:** Lactate accumulation creates acidic intracellular environment
- **Delayed recovery:** Impaired lactate clearance prolongs metabolic stress

The critical difference: athletes experience this state transiently after intense exertion and recover within hours to days; ME/CFS patients exist in this state continuously, even at rest or after minimal activity.

This parallel has practical treatment implications. Sports medicine recovery protocols—electrolyte replacement, magnesium supplementation, ATP precursors (D-ribose), lactate clearance strategies—may provide symptomatic benefit by addressing the chronic metabolic stress state. See Appendix ?? for detailed discussion of how this clinical insight informed treatment protocol development.

Observation 38 (Permanent Post-Exercise Metabolic State). ME/CFS muscle pathophysiology may be understood as a state of continuous post-exercise metabolic stress without the triggering exercise. Interventions that support athletic recovery from intense exertion may provide baseline metabolic support for ME/CFS patients:

- Oral rehydration solutions for blood volume and lactate clearance
- Magnesium for ATP synthesis and muscle relaxation
- Acetyl-L-carnitine to restore fat oxidation capacity
- D-ribose as direct ATP building block

This framework suggests ME/CFS patients require continuous application of recovery protocols, not as performance enhancement but as compensatory support for chronically impaired energy metabolism.

Insulin Sensitivity

Insulin resistance features in some ME/CFS patients:

- **Hyperinsulinemia:** Compensatory insulin excess
- **Impaired glucose tolerance:** Abnormal oral glucose tolerance tests
- **Metabolic syndrome overlap:** Shared features in some patients
- **Inflammation link:** Cytokines promote insulin resistance

6.6 Catecholamine Metabolism: NIH Study Findings

The NIH deep phenotyping study provided groundbreaking data on catecholamine abnormalities in cerebrospinal fluid [5], establishing a direct link between neurotransmitter metabolism and ME/CFS symptoms.

6.6.1 CSF Catecholamine Findings

Reduced Catecholamine Levels

Lumbar puncture analysis revealed significantly reduced central catecholamines:

- **Dopamine metabolites:** Lower homovanillic acid (HVA)
- **Norepinephrine metabolites:** Reduced 3-methoxy-4-hydroxyphenylglycol (MHPG)
- **Implications:** Central catecholamine synthesis or turnover is impaired

Correlation with Symptoms

The study established direct correlations between CSF catecholamines and clinical measures:

- **Motor performance:** Lower catecholamines correlated with reduced grip strength
- **Effort behaviors:** Predicted reduced selection of difficult tasks
- **Cognitive function:** Correlated with memory and executive function deficits
- **Fatigue severity:** Inverse correlation with norepinephrine markers

6.6.2 Catecholamine Synthesis Pathway

Understanding the pathway illuminates potential dysfunction points:

1. **Tyrosine → L-DOPA:** Tyrosine hydroxylase (rate-limiting, requires tetrahydrobiopterin)
2. **L-DOPA → Dopamine:** Aromatic amino acid decarboxylase (requires pyridoxal phosphate)
3. **Dopamine → Norepinephrine:** Dopamine β -hydroxylase (requires copper, ascorbate)
4. **Norepinephrine → Epinephrine:** PNMT (primarily in adrenal medulla)

6.6.3 Potential Mechanisms of Catecholamine Deficiency

Cofactor Deficiencies

Catecholamine synthesis requires multiple cofactors:

- **Tetrahydrobiopterin (BH4):** Essential for tyrosine hydroxylase; depleted by oxidative stress
- **Iron:** Required by tyrosine hydroxylase
- **Pyridoxal phosphate (B6):** Required for decarboxylation
- **Ascorbate (Vitamin C):** Required for dopamine β -hydroxylase
- **Copper:** Required for dopamine β -hydroxylase

Oxidative Stress Effects

Oxidative stress can impair catecholamine metabolism:

- **BH4 oxidation:** Converts active BH4 to inactive BH2
- **Enzyme damage:** Oxidative modification of synthetic enzymes
- **Catecholamine oxidation:** Auto-oxidation generates more ROS
- **Neuromelanin formation:** Oxidized catecholamines form potentially toxic aggregates

Inflammation Effects

Inflammatory cytokines affect catecholamine metabolism:

- **GTP cyclohydrolase induction:** Initially increases BH4 but depletes with chronic inflammation
- **Altered enzyme expression:** Cytokines modify gene expression
- **Competition for BH4:** Increased iNOS activity consumes BH4
- **Microglial activation:** Affects local neurotransmitter metabolism

6.6.4 Functional Consequences

Dopamine Deficiency

Reduced dopamine affects multiple systems:

- **Motivation and reward:** Dopamine mediates reward anticipation
- **Motor function:** Contributes to motor initiation and execution
- **Cognition:** Essential for working memory and executive function
- **Mood:** Contributes to anhedonia and depression symptoms

Norepinephrine Deficiency

Reduced norepinephrine affects:

- **Arousal:** Norepinephrine maintains wakefulness and alertness
- **Attention:** Required for sustained and selective attention
- **Autonomic function:** Central norepinephrine modulates autonomic outflow
- **Stress response:** Mediates appropriate responses to stressors

6.7 The “Metabolic Trap” Hypothesis

Several researchers have proposed that ME/CFS involves metabolic “traps” — stable dysfunctional states that persist even after the initial trigger resolves.

6.7.1 IDO Metabolic Trap

One prominent hypothesis involves tryptophan metabolism:

- **Trigger:** Infection induces IFN- γ , activating IDO
- **Tryptophan depletion:** IDO diverts tryptophan from serotonin to kynurenine
- **Kynurenine effects:** Metabolites may perpetuate immune activation
- **Feedback loop:** Chronic activation maintains the altered state

6.7.2 The “Dauer” Hypothesis

Drawing on *C. elegans* biology, some researchers propose ME/CFS represents a hypometabolic survival state:

- **Dauer state:** Nematode survival mode with reduced metabolism
- **Human analog:** ME/CFS as a protective metabolic downregulation
- **Persistence:** The hypometabolic state becomes self-perpetuating
- **Treatment implications:** May require specific signals to exit the state

6.8 Compartmental Energy Models

Recent evidence suggests ME/CFS may represent *selective* rather than global energy dysfunction. The observation that certain processes (hair growth, nail growth, basic wound healing) remain intact despite severe systemic symptoms challenges the assumption of uniform mitochondrial failure.

6.8.1 CNS-Specific vs. Global Dysfunction

The selective energy dysfunction hypothesis proposes that ME/CFS preferentially affects:

- **CNS-dependent processes:** Functions requiring central coordination (cognition, autonomic regulation, motor control)
- **Demand-responsive processes:** Functions that must scale with physiological challenge (exercise capacity, orthostatic regulation)

While sparing:

- **Autonomous local processes:** Hair follicle cycling, keratinocyte proliferation, basic wound healing
- **Constant-output processes:** Functions that operate at steady state without demand scaling

6.8.2 Evidence for Compartmentalization

Several findings support compartmental rather than global dysfunction:

1. **Preserved peripheral ATP at rest:** 31P-MRS studies show variable findings, with some patients showing normal resting muscle ATP despite symptoms
2. **Demand-response failure:** 91–100% of ME/CFS patients show abnormal cerebral blood flow reduction during orthostatic challenge—3-fold greater than controls [173]—yet baseline perfusion may be preserved
3. **Brain-specific hypometabolism:** PET and SPECT studies reveal regional brain hypometabolism without corresponding peripheral findings [73]
4. **Pharmacological bypass effectiveness:** Direct-acting agents like midodrine can restore function that CNS coordination cannot achieve, suggesting intact peripheral machinery

6.8.3 The Astrocyte-Neuron Lactate Shuttle

The brain's unique metabolic architecture may explain CNS-specific vulnerability. Unlike peripheral tissues with direct glucose access, neurons depend on astrocytes to provide lactate via the astrocyte-neuron lactate shuttle (ANLS) [174, 175]:

- Astrocytes take up glucose and convert it to lactate
- Lactate is exported via MCT4 transporters
- Neurons import lactate via MCT2 transporters
- Lactate oxidation provides 30–50% of neuronal ATP [176]

Dysfunction in this shuttle—from MCT transporter impairment, astrocyte pathology, or neuroinflammation—could cause CNS-specific energy failure while peripheral tissues (with direct glucose access) remain functional.

★ Key Point: Compartmental Model Implications

If energy dysfunction is compartmentalized rather than global:

- Peripheral mitochondrial biomarkers may underestimate CNS dysfunction
- Treatment strategies should prioritize CNS-penetrant approaches
- Subtyping may depend on which compartment shows primary dysfunction

See Section 14.24 for comprehensive treatment of the selective dysfunction hypothesis with formal mathematical framework.

6.9 Potential Interventions

6.9.1 Mitochondrial Support

Cofactors and Substrates

Supporting mitochondrial function may help:

- **Coenzyme Q10:** Electron carrier in ETC; antioxidant
- **L-carnitine/acetyl-L-carnitine:** Fatty acid transport; neuroprotection
- **B vitamins:** Cofactors for multiple metabolic enzymes
- **Magnesium:** Required for ATP utilization
- **D-ribose:** Substrate for ATP synthesis
- **Alpha-lipoic acid:** Antioxidant; mitochondrial cofactor

Mitochondrial-Targeted Therapies

Emerging approaches target mitochondria specifically:

- **MitoQ:** Mitochondria-targeted antioxidant
- **SS-31 (Elamipretide):** Cardiolipin-binding peptide
- **Nicotinamide riboside:** NAD⁺ precursor
- **Urolithin A:** Promotes mitophagy

6.9.2 Antioxidants

Glutathione Support

Restoring glutathione may be beneficial:

- **N-acetylcysteine (NAC):** Provides cysteine for glutathione synthesis (a dose-response trial using brain MRS is underway; see Appendix I, Section I.7.10)
- **Liposomal glutathione:** May improve absorption
- **Glycine supplementation:** Second rate-limiting substrate
- **Selenium:** Required for glutathione peroxidase

Other Antioxidants

- **Vitamin C:** Water-soluble antioxidant; cofactor for catecholamine synthesis
- **Vitamin E:** Fat-soluble membrane antioxidant
- **Polyphenols:** Plant-derived antioxidants (resveratrol, quercetin)
- **Melatonin:** Potent antioxidant with mitochondrial effects

6.9.3 Addressing Catecholamine Deficiency

Precursor Support

Supporting neurotransmitter synthesis:

- **Tyrosine:** Catecholamine precursor
- **Phenylalanine:** Converted to tyrosine
- **BH4 support:** Sapropterin or folate to support BH4 recycling
- **Cofactors:** Iron, B6, vitamin C, copper

Pharmacological Approaches

Medications affecting catecholamine systems:

- **Stimulants:** Methylphenidate, amphetamines (increase catecholamine release)
- **Bupropion:** Norepinephrine-dopamine reuptake inhibitor
- **SNRIs:** Serotonin-norepinephrine reuptake inhibitors
- **MAO-B inhibitors:** Reduce dopamine breakdown

6.10 Microcirculation and Oxygen Delivery

The energy crisis in ME/CFS begins upstream of mitochondrial ATP synthesis. Multiple independent research groups have documented vascular and microcirculatory dysfunction that directly impairs substrate and oxygen availability, creating a pre-mitochondrial bottleneck.

6.10.1 Endothelial Dysfunction

Scherbakov et al. [177] found endothelial dysfunction (ED) in 51% of ME/CFS patients using peripheral arterial tonometry (EndoPAT). This early indicator of vascular disease correlates with systemic atherosclerosis risk and predicts cardiovascular events. Haffke et al. [178] measured flow-mediated dilation (FMD), the gold-standard non-invasive endothelial function test, and found severely impaired responses in ME/CFS patients (5.1% vs 8.2% in healthy controls, $p < 0.0001$). This indicates reduced nitric oxide (NO) bioavailability and impaired vasodilatory capacity.

Blauensteiner et al. [179] identified five endothelial-regulating microRNAs (miR-21, miR-34a, miR-92a, miR-126, miR-200c) with altered expression in up to 60% of ME/CFS patients. All five microRNAs are key regulators of the NO pathway and endothelial function. This suggests dysfunction at the molecular level controlling vascular biology.

6.10.2 Red Blood Cell Deformability and Rheology

Red blood cell function is critical for oxygen delivery: RBCs must deform to traverse narrow capillaries and must maintain membrane fluidity to exchange gases efficiently. Saha et al. [180] studied ME/CFS patient RBCs using ektacytometry and rheological analysis, finding:

- Higher capillary entry time (approximately 12% longer)
- Lower transit velocity through capillaries (approximately 17% reduced)
- Lower elongation index on deformation (approximately 14% reduced membrane deformability)
- Lower membrane fluidity (approximately 30% reduced)
- Higher reactive oxygen species (ROS) production (approximately 30% elevated)

Critically, recovering ME/CFS patients showed normalization of these parameters, suggesting that RBC dysfunction is a reversible consequence of the disease rather than a primary genetic defect. The combination of reduced deformability and elevated ROS creates a cascade: stiff RBCs move slower through capillaries, allowing more time for oxidative damage, further reducing deformability and oxygen delivery.

6.10.3 Capillary Structural Pathology

Wust et al. [181] conducted electron microscopy studies of muscle capillaries from ME/CFS patients and documented structural abnormalities: massive collagen IV deposition in the capillary basement membrane, thickening and narrowing of capillary passageways, and physical barriers to oxygen diffusion. These are not functional abnormalities but anatomical obstructions—capillaries cannot transport oxygen effectively due to structural remodeling. This represents a mechanical bottleneck independent of mitochondrial function.

6.10.4 Brain Microvascular Insufficiency

Godlewska et al. [182] used 7-Tesla magnetic resonance spectroscopy (7T MRS) in 24 ME/CFS patients to measure brain lactate levels at high spatial resolution. They found elevated brain lactate in the pregenual anterior cingulate cortex (pACC) and dorsal anterior cingulate cortex (dACC)—regions critical for mood, pain processing, and autonomic regulation. Elevated brain lactate reflects anaerobic metabolism in brain tissue, consistent with localized microvascular insufficiency in these regions.

6.10.5 Integration: Substrate Delivery Failure

Together, these findings identify a pre-mitochondrial energy bottleneck: endothelial dysfunction reduces perfusion pressure, RBC deformability limits capillary transit, structural capillary pathology blocks oxygen diffusion, and localized brain insufficiency impairs central regulatory systems. Cells cannot generate energy efficiently when they cannot access substrates,

regardless of mitochondrial capacity. This substrate delivery failure occurs independent of and before any mitochondrial defect.

6.11 Energy Utilization Beyond ATP Production

Even when ATP is produced, cells in ME/CFS may have impaired ability to use that ATP effectively. This is mechanistically distinct from ATP production defects and requires different intervention strategies.

6.11.1 The Sodium-Calcium Overload Cascade

Wirth and Scheibenbogen [183] integrated emerging evidence into a model of ion homeostasis failure in ME/CFS. The cascade begins with hypoperfusion (Section 6.10), which forces cells into anaerobic metabolism. Anaerobic metabolism generates protons as a byproduct. Accumulating protons increase the activity of the sodium-hydrogen exchanger (NHE1), which extrudes protons by importing sodium. Under normal aerobic conditions, the Na^+/K^+ -ATPase can clear this sodium, but under hypoperfusion and ATP limitation, this pump becomes saturated and dysfunctional.

When Na^+/K^+ -ATPase fails to maintain the sodium gradient, the sodium-calcium exchanger (NCX), which normally exports calcium in exchange for importing sodium, reverses direction. Instead of exporting calcium, it imports calcium driven by the collapsed sodium gradient. Calcium accumulates in the cytoplasm and mitochondria, causing mitochondrial damage, further reducing ATP production, amplifying the original problem.

Wirth et al. [184] documented this cascade in muscle studies using MRI to measure intracellular sodium. They found elevated intracellular sodium inversely correlating with grip strength (a measure of muscle function), supporting the hypothesis that sodium accumulation contributes to weakness. Electron microscopy showed subsarcolemmal mitochondrial damage at sites of calcium influx, consistent with calcium-mediated mitochondrial injury.

Scheibenbogen and Wirth [185] expanded this model using proteomic data showing reduced expression of proteins responsible for sodium clearance (Na^+/K^+ -ATPase subunits) and increased expression of sodium-sensing proteins (ENaC), indicating a systemic shift toward sodium accumulation and retention. This suggests the dysfunction is not transient but reflects sustained pathological remodeling.

6.11.2 TRPM3 Ion Channel Dysfunction

Cabanas et al. [186] studied TRPM3 calcium channels in natural killer (NK) cells from ME/CFS patients and found reduced calcium influx amplitude and prolonged half-time (slower kinetics). TRPM3 is critical for NK cell calcium signaling and cytotoxic function. Importantly, low-dose naltrexone (LDN) restored TRPM3-dependent calcium influx in patient NK cells in vitro, suggesting a reversible pharmacological mechanism. Sasso et al. [187] extended these

findings with large-scale validation confirming the TRPM3 dysfunction across broader patient populations.

6.11.3 WASF3 and ER Stress-Induced Metabolic Switching

Wang et al. [2] (cross-referenced as WASF3 studies in Chapter ??) identified endoplasmic reticulum (ER) stress upregulation of WASF3 (Wiskott-Aldrich syndrome protein family member 3). WASF3 simultaneously disrupts respiratory supercomplex assembly (impairing OXPHOS efficiency) and promotes actin polymerization, driving a forced switch toward glycolysis. Muscle biopsies from ME/CFS patients showed elevated WASF3 expression, confirming ER stress-mediated metabolic reprogramming.

This represents a cell-autonomous mechanism: ER stress—triggered by calcium overload, oxidative stress, or viral persistence—activates a program that simultaneously reduces OXPHOS capacity while increasing glycolytic flux. Even if mitochondria are structurally intact, WASF3-mediated disruption of supercomplexes reduces their efficiency.

6.11.4 Distinction: Production vs. Utilization Defects

These mechanisms reveal a critical distinction: ATP production defects (failing mitochondria, substrate delivery) are different from ATP utilization defects (ion pump failure, calcium overload, forced metabolic switching). Both limit cellular energy availability, but they require different therapeutic targets. A patient with sodium-calcium overload may have normal mitochondrial capacity but cannot function because ion homeostasis is broken. A patient with WASF3-driven supercomplex disruption may have sufficient ATP but inefficient energy use due to forced glycolysis.

6.12 Immune-Metabolic Crosstalk Maintaining the Energy Crisis

Immune activation is not independent of metabolic dysfunction—instead, immune-mediated pathways actively maintain and amplify the energy crisis. This creates a feedback loop where immune activation perpetuates energy failure.

6.12.1 Tryptophan-Kynurenine Metabolism and NAD⁺ Depletion

The tryptophan-kynurenine pathway is one of the primary immune-activated routes of tryptophan catabolism. Elevated kynurenine production is associated with chronic immune activation (elevated IFN- γ , TNF- α). However, the balance of kynurenine metabolites matters: normal kynurenine metabolism feeds into NAD⁺ synthesis, supporting mitochondrial energy production; imbalanced metabolism produces neurotoxic metabolites while depleting NAD⁺.

A recent metabolomics study [188] (published in ACS Chemical Neuroscience) profiled tryptophan metabolism in ME/CFS plasma and found higher 3-hydroxykynurenine (a neurotoxic

branch product), lower kynurenic acid (neuroprotective, NAD⁺-producing), and lower quino-
linic acid (a neurotoxic NMDA agonist that also feeds NAD⁺ synthesis via QPRT). The net
effect: immune-mediated tryptophan catabolism is diverted away from NAD⁺ synthesis and
toward neurotoxic branches. This directly links immune activation to NAD⁺ depletion and
impaired mitochondrial function.

6.12.2 T-Cell Exhaustion with Metabolic Reprogramming

Iu et al. [119] (published in PNAS) profiled CD8⁺ T cell epigenetics and metabolism in ME/CFS, finding that T cells are epigenetically primed toward an exhausted phenotype (high PD-1, TIM-3, LAG-3) while simultaneously showing altered metabolic reprogramming. Post-exercise, this metabolic dysregulation worsened. This suggests T cells cannot sustain normal metabolic flexibility in response to immune demands, contributing to both the dysfunctional immune response and persistent metabolic strain.

6.12.3 Exaggerated Innate Immune Response with Metabolic Consequences

Che, Hornig, and Lipkin [189] (published in npj Metabolic Health and Disease) documented an exaggerated innate immune response to standard challenges in ME/CFS. Their metabolomic data showed that immune activation is associated with impaired citric acid cycle (TCA cycle) flux and reduced beta-oxidation capacity. Mechanistically, IFN-gamma (elevated in their cohort) accelerates tryptophan degradation via indoleamine 2,3-dioxygenase (IDO), directly linking innate immune state to NAD⁺ depletion and energy failure.

6.12.4 Oxidative Stress in Immune Cells

Paul and Bhatt [190] (published in PNAS) measured reactive oxygen species (ROS) in circulating lymphocytes from ME/CFS and Long COVID patients. They found elevated ROS in memory T cells, with striking sex differences: females showed higher ROS production and T-cell hyperproliferation; males showed evidence of mitochondrial lipid oxidative damage. Critically, metformin treatment attenuated T-cell hyperproliferation in vitro, suggesting ROS-driven immune dysregulation may be pharmacologically reversible.

6.12.5 Multi-Omics Integration: The Full Picture

The Heng 2025 multi-omics study [12] integrated cellular ATP profiling with plasma proteomics from 61 matched pairs, revealing coordinated abnormalities: elevated AMP and ADP with a reduced ATP/ADP ratio (indicating cellular energy stress), altered immune cell subsets skewed toward less mature effector populations, and elevated vascular dysfunction markers (VWF, fibronectin, thrombospondin-1). A 7-biomarker panel spanning these three domains achieved 91% diagnostic accuracy (Section 13.7 in Chapter 13).

These findings were substantially extended by Hoel et al. (2026) [191], who applied aptamer-based proteomics (SomaScan, 7,326 aptamers targeting 6,493 proteins) to serum from 50

ME/CFS patients and 29 healthy controls. Their analysis revealed a striking metabolic enzyme signature: key glycolytic enzymes—glyceraldehyde-3-phosphate dehydrogenase (GAPDH), fructose-bisphosphate aldolase A (ALDOA), and L-lactate dehydrogenase B chain (LDHB)—were among the most significantly reduced intracellular proteins in ME/CFS serum. Since these enzymes are normally abundant in skeletal muscle, their reduced circulating levels suggest decreased muscle protein turnover or altered cellular release patterns rather than simple enzyme deficiency. Notably, ATP synthase inhibitory factor 1 (ATP5IF1), which regulates mitochondrial ATP hydrolysis under hypoxic conditions, was also reduced, consistent with the mitochondrial stress protection deficit described in Section 6.2.

Complementing the intracellular enzyme reductions, Hoel et al. found chronically elevated metabolic stress hormones: FGF-21 was significantly higher in ME/CFS patients, validated by ELISA in an extended cohort ($n = 212$ ME/CFS, $n = 66$ HC), with women showing a particularly significant elevation. GDF-15, leptin, resistin, and fatty acid-binding proteins FABP3 and FABP4 were also elevated. FGF-21 and GDF-15 are “exerkines” that normally rise transiently during intense exercise to coordinate systemic energy adaptation [191]. Their chronic elevation in ME/CFS—in the absence of intense exercise—suggests a state of persistent metabolic emergency signaling, as though the body is continuously experiencing the energy crisis of maximal exertion at rest. This pattern connects to the clinical observation that ME/CFS patients describe exhaustion disproportionate to activity level: their metabolic stress signaling reflects a genuine cellular energy crisis, not a perceptual distortion.

Together, these studies demonstrate that energy metabolism dysfunction in ME/CFS is not limited to mitochondrial abnormalities but extends to a coordinated shift in the circulating metabolic landscape: reduced intracellular metabolic enzymes, elevated stress hormones, and altered secretory patterns spanning multiple tissues. Breaking this cycle may require simultaneous intervention at multiple levels—addressing both the mitochondrial energy deficit and the chronic stress signaling that perpetuates it.

6.13 The Multi-Level Vicious Cycle

Integrating findings from vascular, metabolic, and immune domains reveals a multi-level positive feedback loop where dysfunction at any level amplifies all others. This model, synthesized by Wirth, Scheibenbogen, and colleagues, explains both the severity and persistence of the energy crisis.

6.13.1 The Cycle in Detail

The vicious cycle operates as follows:

1. **Vascular dysfunction** (Section 6.10): Endothelial dysfunction, RBC deformability impairment, and capillary remodeling reduce oxygen and substrate delivery to tissue.
2. **Ischemia**: Reduced oxygen availability forces cells toward anaerobic metabolism, producing lactate and protons.

3. **Proton accumulation:** Intracellular acidosis increases sodium-hydrogen exchanger (NHE1) activity, importing sodium to extrude protons.
4. **Sodium overload:** Na^+/K^+ -ATPase is insufficient to clear accumulated sodium, particularly under ATP limitation.
5. **NCX reversal:** The sodium-calcium exchanger, driven by the sodium gradient, reverses direction and imports calcium.
6. **Calcium overload:** Excessive cytoplasmic and mitochondrial calcium causes direct mitochondrial damage at subsarcolemmal sites, further impairing ATP production.
7. **Mitochondrial dysfunction:** Damaged mitochondria produce less ATP and more ROS.
8. **ROS amplification:** Elevated ROS damages endothelial cells, further impairing vascular function; damages Na^+/K^+ -ATPase, reducing sodium clearance; and damages RBC membranes, further reducing deformability.
9. **Return to step 1:** ROS-mediated vascular damage perpetuates the original ischemia.

This is not a linear pathway with a beginning and end. Rather, it is a positive feedback loop where each step amplifies the next, and dysfunction propagates in all directions simultaneously.

6.13.2 Multi-Level Integration

Critically, this cycle operates at multiple biological levels simultaneously:

Vascular level Endothelial dysfunction → reduced perfusion

Cellular level Ischemia → anaerobic metabolism, ion dysregulation, mitochondrial damage

Molecular level ROS, calcium, sodium dysregulation → persistent vascular and mitochondrial injury

Immune level Immune activation accelerates tryptophan catabolism (NAD^+ depletion) and oxidative stress, perpetuating energy failure

No single level can be addressed in isolation. A therapeutic intervention that improves mitochondrial function alone, without addressing vascular dysfunction or ion dysregulation, will have limited impact because ROS and calcium overload will continue to damage mitochondria. Similarly, anti-inflammatory therapy alone cannot restore energy if vascular dysfunction persists. Effective treatment likely requires simultaneous multi-level intervention.

6.13.3 Evidence for the Integrated Cycle

Appelman et al. [192] (published in Nature Communications) studied muscle mitochondrial function before and after post-exertional malaise (PEM). They found that exercise triggered reductions in OXPHOS capacity and succinate dehydrogenase (SDH) activity, consistent with mitochondrial damage during PEM. Importantly, this occurred despite normal cardiopulmonary responses during exercise—the heart and lungs functioned normally, confirming that the energy crisis is peripheral (muscle) and mitochondrial, not cardiopulmonary.

Syed et al. [64] (published in *Physiology*) provided a comprehensive review of mitochondrial dysfunction across ME/CFS literature, synthesizing evidence from biochemistry, imaging, genetics, and clinical studies. Their analysis supports the multi-level model: mitochondrial pathology is real and well-documented, but its severity and persistence are amplified by upstream vascular, immune, and ion dysregulation.

~ Hypothesis 3: The Multi-Level Energy Crisis and Vicious Cycle in ME/CFS

ME/CFS pathophysiology is fundamentally a multi-level positive feedback loop where vascular dysfunction → ischemia → ion dysregulation → mitochondrial damage → ROS → vascular amplification, perpetuated by immune-mediated NAD⁺ depletion and metabolic reprogramming. Unlike linear models predicting clear breakpoints for intervention, this cycle requires multi-level therapeutic targeting. Evidence comes from multiple independent research groups documenting vascular pathology (Scherbakov, Haffke, Wust, Mueller), ion dysregulation (Wirth, Scheibenbogen), mitochondrial damage (Appelman), and immune-metabolic coupling (Che, Hornig, Lipkin; Paul, Bhatt; Germain). The cycle explains both symptom severity (multiple simultaneous failures) and disease persistence (mutual amplification). Certainty: 0.55 (strong mechanistic evidence from multiple groups, some direct experimental confirmation of sub-components; positive feedback loops are inherently difficult to demonstrate experimentally in a single study, but the logical integration of independently well-established components is sound).

⚠ Limitation 2: Multi-Level Vicious Cycle: Integrated Loop Not Demonstrated

The vicious cycle model (vascular → ischemia → ion dysregulation → mitochondrial damage → ROS → vascular amplification) integrates findings from multiple independent research groups, but no single study has demonstrated the complete loop operating in ME/CFS patients. Each link is supported by evidence from separate cohorts using different methodologies. The integration is logical inference, not empirical demonstration. Positive feedback loops are inherently difficult to distinguish from parallel independent dysfunctions that happen to co-occur. The therapeutic implication that multi-level targeting is required has not been tested against sequential single-target intervention.

6.14 Summary: Integrated Metabolic Model

This chapter and the broader document identify over thirty distinct mechanisms affecting energy creation, delivery, and utilization in ME/CFS. These range from well-established biochemical pathways through documented dysfunctions to emerging hypotheses. The following taxonomy organizes all identified mechanisms by evidence status.

6.14.1 Normal ATP Synthesis Pathways

Five baseline energy systems supply cellular ATP under normal conditions. All are relevant to ME/CFS because each represents a potential point of failure:

- Glycolysis** Cytoplasmic, anaerobic conversion of glucose to pyruvate yielding 2 ATP and 2 NADH per glucose (Section 6.1.1).
- Krebs cycle (TCA cycle)** Mitochondrial matrix oxidation of acetyl-CoA generating 3 NADH, 1 FADH₂, and 1 GTP per turn (Section 6.1.1).
- Electron transport chain and oxidative phosphorylation** Inner mitochondrial membrane complexes I–V producing 30–32 ATP per glucose via the proton gradient (Section 6.1.1).
- Fatty acid β -oxidation** Carnitine-shuttle-dependent mitochondrial degradation of long-chain fatty acids into acetyl-CoA units (Section 6.5.2).
- Phosphocreatine buffering** Immediate ATP regeneration from phosphocreatine, operating on a timescale of seconds (Section 6.1.2).

6.14.2 Documented Energy Dysfunctions

The following dysfunctions are supported by peer-reviewed evidence at the level of established or probable findings:

- Impaired mitochondrial ATP production** Reduced ATP in lymphocytes/PBMCs, abnormal mitochondrial morphology on electron microscopy, and reduced oxygen consumption rates across multiple studies (Section 6.2.1).
- Elevated AMP/ADP with reduced ATP/ADP ratio** Heng et al. 2025 multi-omics study ($n = 61$) documented elevated AMP (median 312.2 nM vs 147.2 nM controls) and reduced ATP generation in white blood cells [12] (Section 6.2.1).
- WASF3–ER stress–Complex IV pathway** Viral ER stress elevates WASF3, which translocates to mitochondria and disrupts respiratory chain supercomplex assembly, specifically impairing Complex IV [2] (Section 6.2.2).
- Reactive oxygen species vicious cycle** Electron leakage at Complexes I and III (5–10% in ME/CFS vs normal 2%) damages mitochondrial DNA, proteins, and membranes, producing further ROS in a self-amplifying loop (Section 6.4.1).
- Nitrosative stress** Excess NO and peroxynitrite inhibit Complex I, Complex IV, and aconitase, directly impairing both the ETC and TCA cycle (Section 6.4.2).
- Glutathione depletion** Reduced cortical glutathione correlating with physical functioning ($\rho = 0.506$) and energy ($\rho = 0.606$) [193] (Section 6.4).
- TCA cycle intermediate deficiencies** Significantly reduced plasma citrulline, malate, and isocitrate; diagnostic pyruvate/isocitrate and ornithine/citrulline ratios [194] (Chapter 11).
- Pyruvate dehydrogenase dysfunction** Impaired conversion of pyruvate to acetyl-CoA at the TCA cycle entry point; thiamine and lipoic acid are cofactors (Section 6.9).
- Fatty acid oxidation defects** Reduced carnitine levels, elevated acylcarnitines suggesting incomplete oxidation, and premature shift to glucose utilization during exercise (Section 6.5.2).
- CoQ10 deficiency** Significantly lower plasma CoQ10 with 44.8% of patients below the lowest control value; correlates with fatigue, autonomic symptoms, and cognitive dysfunction [195] (Section 6.9).

NAD⁺/NADH dysregulation Abnormal NAD⁺ metabolism documented in ME/CFS white blood cells, impairing Complex I function, glycolysis, and TCA cycle activity [12] (Section 6.2.1).

Immune cell metabolic dysfunction CD8⁺ T cells show failed Warburg-effect transition: reduced mitochondrial membrane potential and inability to switch from oxidative phosphorylation to glycolysis upon activation [170] (Chapter 7).

Lactate accumulation and anaerobic shift Earlier and greater lactate accumulation during exercise, elevated resting and brain lactate, slower post-exercise clearance, and leftward-shifted anaerobic threshold (Section 6.5.3).

Brain lactate elevation Elevated ventricular and regional brain lactate on MR spectroscopy, with lactate-to-creatine ratio increases in right insula, thalamus, and cerebellum [193] (Section 6.8).

Oxygen delivery impairment Fibrinoid microclots (>10-fold greater burden), reduced red blood cell mass (93.8% of female patients below normal), impaired flow-mediated dilation, and 91% abnormal cerebral blood flow reduction during tilt testing (Chapter 10).

Catecholamine deficiency Reduced CSF dopamine metabolites (HVA) and norepinephrine metabolites (MHPG) [5], impairing autonomic cardiovascular regulation and energy mobilization on demand (Section 6.6).

Na⁺/K⁺-ATPase impairment Elevated baseline intracellular muscle sodium (12.2 mM vs 9.4 mM, $p = 0.003$) with greater post-exercise accumulation, suggesting ion pump failure secondary to ATP deficit [196] (Chapter 22).

6.14.3 Hypothesized Energy Mechanisms

The following mechanisms are proposed but not yet confirmed. Where explicitly assessed, certainty levels are noted:

Selective energy dysfunction (CNS-dependency hypothesis) ME/CFS selectively impairs CNS-dependent and demand-responsive processes while sparing autonomous steady-state functions, suggesting an energy *allocation* problem rather than a global production deficit (Section 6.3).

Astrocyte-neuron lactate shuttle dysfunction (certainty 0.35) Neuroinflammation may reprogram astrocytes toward a glycolytic-inflammatory phenotype, reducing lactate transfer to neurons via MCT2/MCT4 downregulation and explaining coexistent elevated brain lactate with cognitive dysfunction (Section 6.8; Chapter 8).

Cell danger response Cells remain stuck in a defensive metabolic mode—reduced mitochondrial function, altered purinergic signaling, maintained inflammatory readiness—that evolved as a protective response but fails to terminate [197] (Section 6.7).

Hypometabolic dauer state ME/CFS as a hypometabolic survival state analogous to the *C. elegans* dauer state, with ~80% of metabolic pathways suppressed and a ~2-year threshold separating potentially reversible early disease from entrenched hypometabolism (Section 6.7).

Ketone bypass hypothesis Beta-hydroxybutyrate and acetoacetate may bypass blocked TCA cycle steps and the impaired ANLS by entering neurons directly via MCT1/MCT2,

potentially benefiting patients with TCA (but not ETC) defects (Chapter 14).

Purinergic signaling dysregulation Extracellular ATP acting as danger signal via sensitized P2X/P2Y receptors may explain delayed PEM through exercise-induced massive inappropriate danger signaling and NLRP3 inflammasome activation (Chapter 14).

Succinate as metabolic danger signal Succinate accumulation from impaired oxidative phosphorylation activates NLRP3 inflammasome and stabilizes HIF-1 α , creating a self-reinforcing glycolytic-inflammatory loop (Chapter 15).

IDO metabolic trap IFN- γ -activated IDO diverts tryptophan to the kynurenine pathway, producing neurotoxic metabolites that perpetuate immune activation, maintaining the metabolic diversion in a self-sustaining loop (Section 6.7).

Metabolic flexibility loss Inability to switch between fuel sources (glucose, fatty acids, ketones) based on availability and demand, potentially addressable through time-restricted eating protocols (Chapter 24).

Arginine-citrulline/NO cycle dysfunction Low citrulline/arginine availability produces substrate-limited NO synthesis, impairing vasodilation, mitochondrial biogenesis (via PGC-1 α), and endothelial function [194] (Chapter 11).

Thyroid hormone resistance Suboptimal T4→T3 conversion and possible reduced expression of thyroid hormone transporters (MCT8, MCT10), impairing basal metabolic rate and mitochondrial biogenesis (Chapter 9).

Therapeutic interventions targeting these energy deficits—including mitochondrial cofactors, substrates, and biogenesis signals—are discussed in Section 6.9 and extensively in Part III.

This metabolic dysfunction interacts bidirectionally with immune dysfunction (Chapter 7) and neurological abnormalities (Chapter 8): inflammation impairs metabolism, metabolic dysfunction impairs immune cell function, and energy deficits affect brain function. Chapter 13 synthesizes these bidirectional interactions into comprehensive models of ME/CFS pathophysiology, examining how metabolic dysfunction participates in vicious cycles (Section 13.4) and contributes to the multi-lock state that perpetuates chronic illness (Section 13.3).

7 Immune System Dysfunction

Immune abnormalities are among the most consistently documented features of ME/CFS and likely play a central role in disease pathogenesis. The 2024 NIH deep phenotyping study by Walitt et al. provided the most comprehensive characterisation to date of specific immune abnormalities, including characteristic B cell population shifts and sex-specific patterns of immune dysregulation [5]. This chapter provides a comprehensive examination of immune dysfunction across the innate and adaptive immune systems, inflammatory mediators, and potential autoimmune mechanisms.

As discussed in Chapter 6, immune activation is metabolically costly and may contribute to the bioenergetic crisis in ME/CFS. The neuroinflammatory mechanisms described here connect to autonomic and cardiovascular dysfunction (Chapter 10) and neurological impairment (Chapter 8). Understanding immune dysfunction is thus essential for a comprehensive model of ME/CFS pathophysiology. Chapter 13 synthesizes these cross-system connections, examining how immune dysfunction participates in self-reinforcing pathophysiological cycles (Section 13.4).

7.1 Innate Immunity

The innate immune system provides immediate, non-specific defense against pathogens and plays a critical role in initiating and shaping adaptive immune responses. Multiple components of innate immunity show abnormalities in ME/CFS.

7.1.1 Natural Killer (NK) Cell Dysfunction

Natural killer cell abnormalities represent one of the most replicated findings in ME/CFS research, with impaired NK cell function reported across numerous independent studies spanning decades. A 2019 systematic review of 17 case-control studies (1994–2018) found that impaired NK cell cytotoxicity remained the most consistent immunological abnormality across all publications [3].

Reduced NK Cell Cytotoxicity

NK cells eliminate virus-infected and malignant cells through direct cytotoxic mechanisms. ME/CFS patients consistently demonstrate decreased cytotoxic activity, with reduced ability to kill target cells (typically K562 erythroleukemia cells in standard assays). The magnitude of this impairment is substantial, with studies reporting statistically significant reductions across multiple cohorts [3]. Lower NK cell function correlates with greater symptom severity

in some studies. These abnormalities remain stable over time, suggesting a chronic rather than transient dysfunction.

Mechanisms of Impaired Cytotoxicity

Several mechanisms may underlie reduced NK cell function:

Perforin and Granzyme Deficiency NK cells kill targets by releasing cytotoxic granules containing perforin (which creates pores in target cell membranes) and granzymes (which trigger apoptosis). Maher et al. (2005) demonstrated a mechanistic basis for impaired cytotoxicity: ME/CFS patients show a 45% reduction in NK cell perforin content (3,320 vs 6,051 rMol/cell, $p = 0.01$), with significant correlation between perforin levels and cytotoxic function [198]. Additionally, Brenu et al. (2011) found a paradoxical pattern of elevated perforin but decreased granzyme A and K expression in a large cohort (n=95), suggesting dysfunction in granzyme production or granule composition despite adequate perforin [199]. These cells exhibit impaired degranulation despite successfully recognizing target cells, indicating dysfunction in granule trafficking and release mechanisms.

Receptor Abnormalities NK cell activation is regulated by a balance between activating and inhibitory receptors. ME/CFS patients show altered expression of activating receptors (NKG2D, NKp46, NKp30) along with changed inhibitory receptor profiles [199, 3]. Additionally, signaling downstream of activating receptors is impaired, and calcium flux following receptor engagement is disrupted [187].

Metabolic Dysfunction NK cells require substantial energy for cytotoxic function. ME/CFS NK cells exhibit impaired glycolytic metabolism and mitochondrial dysfunction affecting ATP production [199]. This reduced metabolic reserve limits their capacity for sustained activity.

NK Cell Subsets

Human NK cells are divided into functionally distinct subsets. CD56^{bright} NK cells primarily produce cytokines and are found mainly in lymphoid tissues, while CD56^{dim} NK cells are primarily cytotoxic and predominate in peripheral blood. ME/CFS studies have reported altered CD56^{bright}/CD56^{dim} ratios, with an increased proportion of CD56^{bright} cells in some studies [200, 3]. Reduced absolute numbers of CD56^{dim} cytotoxic cells and abnormal maturation patterns have also been observed [3].

Clinical Significance of NK Cell Dysfunction

Impaired NK cell function may contribute to ME/CFS through several mechanisms:

1. **Viral reactivation:** Inadequate control of latent herpesviruses (EBV, HHV-6, CMV)
2. **Tumor surveillance:** Theoretical increased cancer risk (though not clearly demonstrated)
3. **Immune regulation:** NK cells modulate other immune cells; dysfunction may permit chronic inflammation
4. **Infection susceptibility:** Reduced defense against new infections

These mechanisms may form a self-reinforcing cycle rather than a simple linear causal chain. In particular, the relationship between NK cell dysfunction and viral reactivation is bidirectional: impaired NK function permits viral reactivation, but chronic viral reactivation itself may further exhaust and dysregulate NK cells. Section 7.5.1 examines three competing hypotheses for this relationship with their testable predictions. This bidirectional cycle represents one of several vicious cycles maintaining ME/CFS pathophysiology, discussed comprehensively in Section 13.4 of Chapter 13.

TRPM3 Ion Channel Dysfunction

A major breakthrough in understanding impaired calcium signaling in ME/CFS immune cells came from research on the TRPM3 ion channel [187]. TRPM3 (Transient Receptor Potential Melastatin 3) is a calcium-permeable ion channel, and calcium signaling is essential for healthy immune cell activity—including the degranulation process disrupted in ME/CFS NK cells.

A study conducted by researchers at Griffith University’s National Centre for Neuroimmunology and Emerging Diseases (NCNED) confirmed that TRPM3 functions abnormally in immune cells of ME/CFS patients compared to healthy controls. Critically, this finding was validated across multiple independent laboratories separated by over 4,000 kilometers (Gold Coast and Perth, Australia), using gold-standard techniques—demonstrating robust scientific reproducibility.

The researchers describe the faulty ion channels as acting like “stuck doors,” preventing cells from receiving the calcium they need for normal function. Calcium signaling is essential for immune cell activity, including NK cell cytotoxic function (degranulation requires calcium influx).

This discovery has several important implications:

1. **Diagnostic potential:** TRPM3 dysfunction could serve as an objective biomarker for ME/CFS
2. **Therapeutic targets:** Drugs that modulate TRPM3 function might restore normal immune cell activity
3. **Disease legitimacy:** Measurable cellular abnormalities provide concrete evidence of biological dysfunction
4. **Mechanistic understanding:** TRPM3 dysfunction may explain why NK cells fail to degranulate properly despite recognizing targets

The TRPM3 findings connect to broader ion channel research in ME/CFS and suggest that channelopathy—dysfunction of ion channels—may be a unifying mechanism underlying multiple immune abnormalities observed in the condition.

⚠ **Limitation 1: TRPM3 Channelopathy: Single Research Group**

TRPM3 ion channel dysfunction in ME/CFS has been characterized primarily by a single research group (Griffith University NCNED). While the multi-site validation (Gold Coast and Perth) strengthens internal reproducibility, independent replication by groups outside this collaboration has not yet been published. The diagnostic biomarker potential, therapeutic implications, and status as a “unifying mechanism” for immune abnormalities remain speculative until independently replicated and mechanistically linked to clinical outcomes in prospective studies. See Appendix I, Section I.5.1 for current study status and planned LDN trials.

7.1.2 Neutrophil and Monocyte Function

Neutrophil Abnormalities

Neutrophils are the most abundant circulating white blood cells and serve as first responders to infection. Kennedy et al. (2004) demonstrated that ME/CFS patients exhibit increased neutrophil apoptosis (37.4% vs 22.8% annexin V binding, $p = 0.001$) with elevated death receptor TNFRI expression ($p = 0.004$) and raised active TGF- β 1 concentrations ($p < 0.005$), consistent with an activated inflammatory process [201]. Additional ME/CFS-associated abnormalities include:

Phagocytosis Impairment Neutrophils from ME/CFS patients show reduced uptake of bacteria and particles, with impaired phagosome formation and decreased acidification of phagolysosomes [3].

Respiratory Burst Defects The respiratory burst produces reactive oxygen species to kill ingested pathogens. Some studies have found reduced superoxide production in ME/CFS neutrophils, along with impaired NADPH oxidase function and altered baseline oxidative status [3].

Chemotaxis Impairment Neutrophils in ME/CFS demonstrate reduced migration toward chemoattractants, with impaired directional sensing and decreased expression of chemokine receptors [3].

Neutrophil Extracellular Traps (NETs) NETs are web-like structures of DNA and antimicrobial proteins released by neutrophils. ME/CFS patients show altered NET formation, which may contribute to inflammation if excessive and could potentially explain some autoimmune features of the condition [3].

Proteomic Evidence: Suppressed Neutrophil Signature A large-scale serum proteomics study by Hoel et al. (2026) [191] revealed a broad and pronounced reduction in circulating neutrophil-derived proteins in ME/CFS patients. By cross-referencing their dataset with a published list of proteins released by PMA-activated neutrophils, the authors found that 146 of 254 known neutrophil activation products were detectable, and over 85% of those showing significant differences were reduced in ME/CFS. Hallmark neutrophil granule components—bactericidal/permeability-increasing protein (BPI), peptidylarginine deiminase 4 (PADI4), matrix metalloproteinase 9 (MMP9), neutrophil elastase (ELANE), azurocidin 1 (AZU1), and lactoferrin (LTF)—were all significantly lower. This suppressed neutrophil protein signature occurred despite normal peripheral neutrophil counts ($3.38 \pm 1.03 \times 10^9/\text{L}$), suggesting a qualitative deficiency in neutrophil activation or degranulation rather than a quantitative cell deficit. This proteomic finding complements the functional neutrophil abnormalities described above (impaired phagocytosis, respiratory burst defects, chemotaxis impairment) and suggests that the circulating neutrophil pool in ME/CFS is hypoactivated at baseline—potentially contributing to the impaired innate immune defense and altered inflammatory milieu characteristic of the disease.

More broadly, the Hoel et al. study found that the innate immune arm—granulocyte- and monocyte-associated proteins—showed a primarily downward shift (70.3% and 64.6% of altered proteins reduced, respectively), while NK cell and T cell-associated proteins were predominantly elevated (72.0% and 62.5% upward). This innate-suppressed / adaptive-elevated pattern suggests a fundamental immune reprogramming in ME/CFS, where innate first-responder functions are dampened while adaptive immune cells show signs of chronic activation or compensatory expansion.

Monocyte and Macrophage Dysfunction

Monocytes and their tissue-resident derivatives (macrophages) bridge innate and adaptive immunity:

Monocyte Subset Alterations Human monocytes are classified into three functionally distinct subsets: classical ($\text{CD14}^{++}\text{CD16}^{-}$) monocytes perform phagocytic and antimicrobial functions; intermediate ($\text{CD14}^{++}\text{CD16}^{+}$) monocytes handle antigen presentation and cytokine production; and non-classical ($\text{CD14}^{+}\text{CD16}^{++}$) monocytes conduct patrolling and vascular surveillance. ME/CFS studies have found increased intermediate monocytes (associated with inflammation), altered cytokine production profiles, abnormal responses to stimulation, and changed expression of activation markers.

Macrophage Polarization

Observation 39 (Macrophage Polarization in ME/CFS). Tissue macrophages can adopt pro-inflammatory (M1) or anti-inflammatory (M2) phenotypes. Evidence suggests M1 polarization in ME/CFS, with impaired transition to the resolving M2 phenotype, resulting in chronic inflammatory macrophage activation [12].

7.1.3 Complement System

The complement system consists of plasma proteins that enhance (“complement”) antibody and phagocyte function. Abnormalities in ME/CFS include:

Complement Activation Patterns

ME/CFS patients show elevated activation products, with increased C3a, C4a, and C5a fragments indicating ongoing complement activation [3]. Reduced levels of C3 and C4 suggest consumption of these complement components. Additionally, abnormal levels of complement regulatory proteins point to altered regulation of the system.

Serum proteomics data from Hoel et al. (2026) [191] confirmed and extended these findings: complement factor D (CFD) and complement component C6 were among the most significantly elevated secreted proteins in ME/CFS. Elevated CFD indicates activation of the alternative complement pathway, while elevated C6 indicates increased assembly of the membrane attack complex (MAC). The authors noted that elevated CFD and C6 are implicated in vasculitis, lupus, dermatomyositis, and autoimmune nephritis, further supporting a possible autoimmune-like complement activation pattern in ME/CFS.

Clinical Implications

Complement abnormalities may contribute to inflammation through anaphylatoxin (C3a, C5a) production and impair pathogen clearance [3]. They may also promote autoimmune manifestations and trigger mast cell activation, as complement fragments can induce mast cell degranulation.

7.1.4 Dendritic Cells

Dendritic cells (DCs) are professional antigen-presenting cells that initiate adaptive immune responses. ME/CFS patients show altered DC maturation with abnormal expression of co-stimulatory molecules [3]. Changed cytokine production skews toward pro-inflammatory profiles, while impaired antigen presentation may contribute to inadequate pathogen clearance. Plasmacytoid DCs display abnormalities in type I interferon production [3].

7.2 Adaptive Immunity

The adaptive immune system provides specific, long-lasting responses through T and B lymphocytes. The NIH deep phenotyping study identified characteristic abnormalities in B cell populations that may represent a biomarker signature for ME/CFS [5].

7.2.1 T Cell Abnormalities

T lymphocytes coordinate adaptive immune responses and directly eliminate infected cells.

T Cell Subset Distribution

CD4/CD8 Ratio Changes The ratio of helper (CD4⁺) to cytotoxic (CD8⁺) T cells is altered in some ME/CFS patients, though findings vary considerably across studies [3]. Some report a decreased CD4/CD8 ratio while others find an increased ratio. This heterogeneity may reflect distinct patient subgroups within the ME/CFS population.

Helper T Cell Subsets CD4⁺ T cells differentiate into functional subsets with distinct roles: Th1 cells produce interferon-gamma and promote cell-mediated immunity; Th2 cells produce IL-4, IL-5, and IL-13 to promote antibody responses; Th17 cells produce IL-17 and are involved in autoimmunity and mucosal defense; and regulatory T cells (Tregs) suppress immune responses to maintain tolerance. ME/CFS findings include Th1/Th2 imbalance (though the direction varies across studies), elevated Th17 cells in some patients, and reduced Treg numbers or function [3, 202]. Altered cytokine profiles reflect these subset imbalances.

T Cell Exhaustion Markers

Chronic antigen exposure can lead to T cell exhaustion, characterized by:

- **Increased PD-1 expression:** Programmed death-1, an inhibitory receptor [119, 5]
- **Elevated Tim-3:** T cell immunoglobulin and mucin domain-3 [119]
- **CTLA-4 upregulation:** Cytotoxic T-lymphocyte-associated protein 4 [119]
- **Reduced proliferative capacity:** Impaired response to stimulation [119]
- **Decreased cytokine production:** Despite activation marker expression [119]

These findings suggest chronic immune stimulation in ME/CFS, consistent with persistent infection or autoimmune processes [119].

Comprehensive T Cell Exhaustion Evidence (Iu et al. 2024) A 2024 study published in *PNAS* provided the most detailed characterization of T cell exhaustion in ME/CFS to date [119]. Using transcriptomic and epigenetic profiling, Iu et al. demonstrated that CD8⁺ T cells from ME/CFS patients undergo extensive reprogramming toward an exhausted phenotype.

Key Findings

- **Elevated PD-1 expression:** Confirmed at both protein and transcriptional levels
- **Transcriptional reprogramming:** Gene expression patterns characteristic of chronic antigenic stimulation
- **Epigenetic modifications:** Persistent chromatin changes indicating long-term immune activation rather than transient response
- **Similarity to chronic infections:** The exhaustion profile resembled that seen in chronic viral infections (HIV, hepatitis C) and cancer

Implications The epigenetic nature of these changes suggests that T cell exhaustion in ME/CFS is not merely a snapshot of current immune activation but represents a durable reprogramming of immune cell function. This has several implications:

- **Chronicity:** The epigenetic changes may explain why immune dysfunction persists even if the initial trigger resolves
- **Impaired viral control:** Exhausted T cells cannot effectively clear viruses, potentially permitting herpesvirus reactivation
- **Therapeutic targets:** Immune checkpoint inhibitors (anti-PD-1, anti-CTLA-4) used in cancer might theoretically restore T cell function, though safety in ME/CFS is unknown
- **Biomarker potential:** T cell exhaustion markers could serve as diagnostic or prognostic indicators


Integration with NIH Deep Phenotyping Study The Iu et al. findings complement the Walitt et al. NIH study [5], which also documented elevated CD8+ T cell PD-1 expression. Together, these studies consistently document T cell exhaustion as a reproducible feature of ME/CFS immunopathology, supporting the model of chronic antigenic stimulation driving both B cell (naïve/memory imbalance) and T cell (exhaustion) abnormalities.

Figures 7.1 and 7.2 illustrate the paradoxical immune state in ME/CFS—simultaneously overactive and underactive. Two interconnected vicious cycles drive disease: chronic inflammation (IDO activation, energy deficit, poor pathogen control) and immune exhaustion (T-cell/NK dysfunction, failed clearance). These cycles reinforce each other. The integration of these immune-specific vicious cycles with metabolic and autonomic cycles is examined in Section 13.4 of Chapter 13.

⚠ Limitation 2: T Cell Exhaustion: Chronic Stimulation Source Unknown

T cell exhaustion markers (PD-1, Tim-3, CTLA-4) in ME/CFS resemble those seen in chronic viral infections and cancer. However, the antigenic source driving this exhaustion has not been identified. The exhaustion profile is *consistent with* chronic antigenic stimulation but does not establish what the antigen is (persistent virus, autoantigen, or metabolic danger signal). Without identifying the driver, the therapeutic implication that checkpoint inhibitors might help remains speculative and carries substantial

Normal Immune Response



Key characteristics:

- Complete pathogen clearance
- Active resolution phase with anti-inflammatory mediators

ME/CFS: Immune Dysfunction Cycles




Figure 7.2: ME/CFS immune dysfunction with chronic inflammation and exhaustion cycles.

autoimmune risk.

Two recent single-cell transcriptomic studies strengthen the T cell exhaustion finding. Mayer et al. (2025) constructed a scRNA-seq atlas of 336,269 T lymphoid cells from 28 ME/CFS patients and 30 controls, demonstrating transcriptional reprogramming of CD8+ T cells toward exhaustion that is amplified by exercise provocation [203] (Appendix I, Section I.9.1). Elahi et al. (2026) applied scRNA-seq to long COVID–ME/CFS patients 12 months post-infection, finding persistent immune remodelling—monocyte polarisation, NK cell dysfunction, and T cell exhaustion—absent in recovered individuals [204] (Section I.9.2). Cell-free RNA profiling by Gardella et al. (2025) independently detected T cell exhaustion signatures in plasma cfRNA, confirming these findings via an orthogonal method [205] (Section I.3.4).

Comparative virology provides indirect support for the viral persistence hypothesis. Kol et al. (2026) demonstrated that feline infectious peritonitis virus (FIPV), a naturally occurring coronavirus in cats, replicates not only in macrophages but also in B and T lymphocytes within mesenteric lymph nodes [171]. Critically, rare FIPV RNA-positive lymphocytes persisted after antiviral treatment and full clinical recovery, suggesting that long-lived immune cells serve as viral reservoirs. Because lymphocytes can survive for years, this reservoir may explain relapse and chronic immune dysregulation. FIPV offers a unique comparative model: unlike human studies, where accessing infected lymphoid tissue is ethically and practically difficult, feline FIP permits direct examination of coronavirus–immune cell interactions in naturally infected tissue. While FIPV is not SARS-CoV-2, the shared coronavirus biology—tropism for immune cells, persistence despite apparent clearance, and post-treatment relapse—strengthens the plausibility that analogous mechanisms operate in post-COVID ME/CFS.

~ Hypothesis 1: Lymphocyte Reservoir Ratchet

Certainty: 0.35. Mechanistically plausible and supported by comparative virology (FIP model), but no direct human data yet demonstrates SARS-CoV-2 or other ME/CFS-triggering virus persistence specifically within lymphocytes. The hypothesis extends the Ratchet Model (Speculation 5) with a specific reservoir mechanism.

The Kol et al. (2026) demonstration that FIPV persists in B and T lymphocytes after antiviral treatment [171] suggests a mechanism distinct from classical viral latency: triggering viruses may reside within long-lived immune cells themselves, below conventional PCR detection thresholds but sufficient to drive chronic immune activation. Because memory lymphocytes survive for years to decades, even a small fraction of virus-harboring cells constitutes a persistent antigenic reservoir.

Mechanism. Unlike classical latency (e.g., EBV in B cells, where the viral genome is silenced), this model proposes low-level active or semi-active viral transcription within circulating lymphocytes. Each time these cells are activated by any immune challenge—a new infection, vaccination, allergen exposure, or even exercise-induced immune mobilization—viral antigen is re-presented, triggering localized immune activation without classical “reactivation.” This would explain why PEM can follow diverse immune stimuli, not only physical exertion.

Testable Predictions.

1. Single-cell RNA sequencing of peripheral blood mononuclear cells from post-COVID ME/CFS patients will detect viral transcripts in a small percentage (<1%) of lymphocytes, below bulk PCR sensitivity but detectable at single-cell resolution.
2. The fraction of virus-positive lymphocytes will correlate with disease severity and duration.
3. In vitro activation of ME/CFS patient lymphocytes will produce detectable viral antigen release, while resting cells will not.
4. Patients who recover from ME/CFS will show clearance of lymphocyte-associated viral RNA, while non-recoverers will retain it.

Falsifiability. This hypothesis would be falsified if ultra-sensitive single-cell methods (10x Genomics, MERFISH) applied to ME/CFS patient lymphocytes consistently fail to detect viral transcripts across multiple triggering virus types and patient cohorts. It would also be weakened if lymphocyte depletion therapies (e.g., rituximab targeting B cells) show no effect on viral antigen levels, since the model predicts B cells as a primary reservoir.

Relationship to the Ratchet Model. This hypothesis provides a specific cellular mechanism for the “viral reactivation and persistent viral load” arm of the Ratchet Model (Speculation 5). The Ratchet Model describes the clinical pattern (step-wise decline); this hypothesis explains *where* the virus hides between steps.

Treatment Implications. If confirmed, treatment would require either (a) antiviral therapy sustained long enough to outlast the lifespan of virus-harboring lymphocytes (potentially years), (b) targeted depletion of the harbouring cell population, or (c) metabolic support enabling immune cells to clear the reservoir autonomously (see Hypothesis 6.3).

Limitations. Cross-species extrapolation from FIP to human disease requires caution: FIPV and SARS-CoV-2 differ in receptor usage, tropism breadth, and host immune biology. The “below PCR threshold” claim is currently unfalsifiable with standard methods—only emerging single-cell technologies can test it. The model does not explain non-viral-onset ME/CFS.

T Cell Metabolic Dysfunction

As discussed in Chapter 6, mitochondrial dysfunction in ME/CFS is not limited to muscle and nervous system—it extends to immune cells themselves. Mandarano et al. (2020) provided the first comprehensive metabolic analysis of T cells in ME/CFS (n=53 patients, n=45 controls), demonstrating that immune dysfunction has a fundamental bioenergetic basis [170].

CD8+ T Cell Metabolic Deficits CD8+ cytotoxic T cells showed the most severe impairment: reduced mitochondrial membrane potential (indicating mitochondrial dysfunction), impaired

glycolysis at rest, and crucially, failed metabolic reprogramming following activation. Healthy T cells switch from oxidative phosphorylation to glycolysis when activated (the Warburg effect), but ME/CFS CD8⁺ T cells cannot make this transition effectively [170].

CD4⁺ T Cell Abnormalities CD4⁺ helper T cells also demonstrated reduced glycolysis at rest, though their activation response was less severely impaired than CD8⁺ cells. This suggests a hierarchy of metabolic dysfunction, with cytotoxic cells more vulnerable than helper cells [170].

Clinical Implications T cell metabolic dysfunction may provide a mechanistic explanation for several observations: reduced CD8⁺ cytotoxic function (Brenu et al. 2011 [199]) could result from insufficient ATP to sustain degranulation and target killing, though direct causation has not been experimentally demonstrated; impaired proliferation following stimulation may reflect inability to meet the energetic demands of cell division; and post-exertional malaise may be exacerbated by immune activation, as metabolically compromised immune cells compete with other tissues for limited ATP. This finding bridges the energy metabolism (Chapter 6) and immune dysfunction chapters, demonstrating that ME/CFS is characterized by systemic bioenergetic failure affecting all cellular systems.

Regulatory T Cell Dysfunction

Tregs maintain immune tolerance and prevent autoimmunity. ME/CFS patients show reduced numbers of Tregs (CD4⁺CD25⁺FoxP3⁺ cells) with impaired suppressive function [202, 3]. Altered Treg/effector T cell ratios may potentially contribute to the autoimmune features observed in some patients.

Sex-Specific T Cell Findings from the NIH Study

The Walitt et al. deep phenotyping study revealed striking sex differences in T cell abnormalities [5]:

Male Patients Men with PI-ME/CFS demonstrated:

- Altered T cell activation patterns
- Changes in markers of innate immunity
- Distinct inflammatory signatures compared to female patients

These findings suggest that immune pathophysiology may differ fundamentally between sexes, with implications for treatment approaches.

7.2.2 B Cell Function and Antibodies

B lymphocytes produce antibodies and present antigens to T cells. The NIH deep phenotyping study provided compelling evidence for characteristic B cell abnormalities in PI-ME/CFS, though sample size (n=17) limits definitive conclusions pending replication [5].

B Cell Population Shifts: Key NIH Findings

The Walitt et al. study documented a specific pattern of B cell subset abnormalities that may represent a diagnostic signature:

Increased Naïve B Cells Naïve B cells have not yet encountered their cognate antigen and can respond to any new threat:

- Significantly elevated in PI-ME/CFS patients compared to controls
- Reflects either increased production or impaired maturation
- May indicate abnormal B cell development or survival
- Could represent immune system “reset” following infection

Decreased Switched Memory B Cells Switched memory B cells have undergone class-switch recombination and provide rapid, specific responses to previously encountered pathogens:

- Significantly reduced in PI-ME/CFS patients
- Suggests impaired generation of long-term humoral immunity
- May explain susceptibility to recurrent infections
- Could result from chronic antigenic stimulation “exhausting” the memory pool

Interpretation: Chronic Antigenic Stimulation The NIH study concluded that this B cell pattern “suggested chronic antigenic stimulation” [5]. This interpretation implies:

- Persistent immune activation, possibly from ongoing infection or autoimmunity
- Continuous recruitment of naïve B cells into responses
- Depletion of the memory B cell compartment through sustained activation
- Potential for developing autoantibodies through aberrant B cell selection

? Open Question 1: Naïve vs. Memory B Cell Imbalance

The NIH study found elevated naïve B cells and reduced memory B cells in PI-ME/CFS patients. Does this represent an immune system “stuck” in early activation, continuously attempting new responses but failing to consolidate immunological memory? If so, what maintains this state—persistent antigen, aberrant signaling, or microenvironmental

factors? Could interventions promoting B cell maturation (e.g., targeted cytokine support, germinal center modulation) restore normal immune function and break the cycle of chronic activation?

Severe B Cell Depletion: Exhausted Immune Surveillance

While the NIH study documented B cell *subset* abnormalities with preserved total B cell counts, clinical observation suggests a more severe phenotype exists: profound B cell depletion with apparent immune exhaustion.

Speculation 3 (Exhausted Immune Surveillance Phenotype). **Certainty: 0.35.** This hypothesis is based on clinical observation and plausible mechanistic reasoning, but lacks systematic epidemiological validation. B cell depletion of the severity described has been observed, and EBV reactivation patterns are documented; however, whether this constitutes a distinct, reproducible phenotype driven by the proposed mechanism remains unconfirmed.

A subset of ME/CFS patients may progress to an “exhausted immune surveillance” state characterized by:

- **Severe B cell depletion:** CD19+ counts at 10% of lower reference limit (e.g., 0.05 G/l vs. reference 0.11–0.47)
- **Compensatory T cell elevation:** CD3+ percentage elevated (e.g., 84–85% vs. reference 55–83)
- **Low NK cells:** Impaired viral surveillance capacity (e.g., 6–7% vs. reference 7–31)
- **Extremely elevated viral titers:** Despite antibody production (e.g., EBV IgG 25–30× upper limit)

Proposed mechanism: Chronic viral stimulation (particularly EBV) drives continuous B cell differentiation into antibody-secreting plasma cells. NK cell deficiency prevents clearance of virally-infected cells, perpetuating antigenic stimulation. Terminal plasma cell differentiation progressively depletes the CD19+ B cell pool. The resulting state produces high antibody titers (EBV IgG extremely elevated) but fails to achieve viral control because:

1. Antibodies alone cannot clear intracellular infections
2. NK cells (primary viral surveillance) are insufficient
3. The system *produces* antibodies but cannot *act* on them

Clinical implications: This phenotype may respond to immunomodulation that enhances cellular immunity (NK/T-cell function) rather than interventions that further stimulate humoral responses. Cimetidine’s mechanism—blocking H2 receptors on suppressor T cells to enhance cellular immunity [206]—aligns with this specific deficit. See Section 21.6.2 for treatment considerations.

Relationship to NIH findings: This extreme phenotype may represent late-stage progression of the chronic antigenic stimulation pattern identified by Walitt et al. [5]. Where the NIH

study found B cell *subset* shifts, the exhausted surveillance phenotype shows B cell *compartment* depletion—potentially the end-state of years of sustained activation.

Research directions:

1. Prospective tracking of B cell counts in long-duration ME/CFS patients
2. Correlation of B cell depletion severity with disease duration and viral titers
3. Evaluation of immunomodulatory (vs. immunosuppressive) interventions in this phenotype
4. Assessment of whether B cell depletion predicts response to cellular immunity enhancers

Autoantibodies in ME/CFS

Multiple autoantibodies have been identified in ME/CFS patients:

Anti-Nuclear Antibodies (ANA) Early research by Nishikai (2007) established that antinuclear antibodies are present in 15–25% of CFS patients using indirect immunofluorescence with HEp-2 cells [207]. The ANA titers were generally low and showed heterogeneous immunofluorescent staining patterns. Additionally, Nishikai's group identified autoantibodies to a 68/48 kDa protein in 13.2% of CFS patients compared to 0% of healthy controls ($p < 0.05$), with these autoantibodies more common in patients with hypersomnia and difficulty concentrating [207]. Key characteristics include:

- Present in 15–25% of ME/CFS patients (compared to 5–10% of healthy individuals)
- Usually low titer
- Various patterns (homogeneous, speckled, nucleolar)
- Clinical significance unclear, though specific autoantibodies may correlate with cognitive symptoms

G-Protein-Coupled Receptor (GPCR) Autoantibodies Autoantibodies targeting G-protein-coupled receptors represent one of the most actively investigated areas of ME/CFS research, with substantial evidence supporting their role in disease pathophysiology.

Initial Discovery and Prevalence The foundational study by Loebel et al. (2016) established the presence of GPCR autoantibodies in ME/CFS [71]. In a cohort of 268 ME/CFS patients, 29.5% had elevated autoantibodies against β_2 -adrenergic, M3 muscarinic, or M4 muscarinic receptors compared to healthy controls. This study provided the first systematic evidence that receptor-targeting autoantibodies might contribute to ME/CFS pathophysiology.

Validation Studies Bynke et al. (2020) validated these findings in two Swedish cohorts [72]. Strikingly, 79–91% of ME/CFS patients had at least one elevated autoantibody compared to only 25% of healthy controls. A critical finding was that no autoantibodies were detected in cerebrospinal fluid, suggesting peripheral rather than intrathecal production and indicating that these autoantibodies likely originate from systemic B cells or plasma cells rather than CNS-resident immune cells.

Correlation with Symptom Severity

★ Achievement 1: Quantitative GPCR Autoantibody-Symptom Correlation

Sotzny et al. (2021) demonstrated dose-response relationships between GPCR autoantibody concentrations and clinical measures in infection-triggered ME/CFS patients [208]. Autoantibody levels correlated quantitatively with fatigue severity, muscle pain intensity, cognitive impairment, gastrointestinal symptoms, and autonomic dysfunction measures. While these quantitative correlations are consistent with causation, this cross-sectional evidence does not establish that autoantibodies cause symptoms. However, the dose-response relationship and subsequent mechanistic findings (Hackel 2025) strengthen the case for a causal role.

△ Warning 1: Replication Status: Partially Replicated

GPCR autoantibodies in ME/CFS have been detected by multiple groups (Loebel 2016, Bynke 2020, Sotzny 2021), but Vernino 2022 failed to replicate in POTS using standard ELISA, raising assay specificity concerns. The CellTrend ELISA platform used in most positive studies has been questioned. Dose-response correlations (Sotzny 2021) have not been independently replicated.

Downstream Mechanisms: Monocyte Dysfunction Recent work by Hackel et al. (2025) elucidated how GPCR autoantibodies might cause symptoms [209]. In 24 post-COVID ME/CFS patients compared to 12 controls, autoantibodies were shown to mediate inflammatory and neurotrophic cytokine production via monocyte activation. Specifically, autoantibody binding upregulated MIP-1 δ , PDGF-BB, and TGF- β 3 production. This study provides a mechanistic link between circulating autoantibodies and the downstream inflammatory cascade characteristic of ME/CFS.

Therapeutic Targeting: Immunoadsorption The autoantibody hypothesis has been tested therapeutically through immunoadsorption, which non-selectively removes IgG from plasma. Scheibenbogen et al. (2018) conducted an initial pilot study treating 10 post-infectious ME/CFS patients with elevated β_2 -adrenergic receptor antibodies [147]. 70% showed rapid improvement during treatment, and 30% sustained moderate-to-marked improvement at 6–12 months follow-up.

★ Achievement 2: Autoantibody Removal Produces Clinical Improvement

Stein et al. (2025) treated 20 post-COVID ME/CFS patients with five immunoadsorption sessions, reducing IgG by 79% and β_2 -adrenergic receptor autoantibodies by 77% [118]. 70% (14/20) were classified as responders with ≥ 10 point improvement in SF-36 Physical Function score, with benefits sustained to 6 months. This represents the strongest evidence to date that autoantibody removal can produce clinically meaningful improvement in ME/CFS.

△ Warning 2: Replication Status

Open-label pilot study (Stein 2025, n=20) with no control arm. Placebo effects from an invasive multi-session procedure cannot be excluded. An earlier pilot (Scheibenbogen 2018, n=10) showed similar response rates, but both are from the same Berlin research group. Blinded, sham-controlled trials are essential before efficacy can be established.

Therapeutic Targeting: Plasma Cell Depletion Fluge et al. (2025) took a different approach by targeting the cellular source of autoantibodies [9]. In an open-label pilot study, 10 female ME/CFS patients received daratumumab, an anti-CD38 antibody that depletes plasma cells (the terminally differentiated B cells responsible for sustained antibody production). 60% (6/10) showed marked improvement, with SF-36 Physical Function scores increasing from 25.9 to 55.0 ($p=0.002$). Responders achieved near-normal function with SF-36 scores of 80–95. Notably, low baseline NK-cell count predicted non-response, suggesting patient selection criteria may be important. This study suggests that long-lived plasma cells, rather than B cells themselves, may be the critical source of pathogenic autoantibodies.

Therapeutic Targeting: Autoantibody Neutralization Hohberger et al. (2021) reported a case of BC007, a DNA aptamer that directly neutralizes GPCR autoantibodies [10]. A Long COVID patient with elevated GPCR autoantibodies received a single 1350mg intravenous dose. Autoantibodies were neutralized within hours, with dramatic clinical improvement: fatigue normalized, brain fog resolved, taste sensation was restored, and retinal microcirculation improved on optical coherence tomography angiography. Effects were sustained at 4-week follow-up. This proof-of-concept case demonstrates that direct autoantibody neutralization can produce rapid symptomatic improvement.

Methodological Controversies The GPCR autoantibody field faces important methodological challenges. Vernino et al. (2022) attempted to replicate autoantibody findings in postural orthostatic tachycardia syndrome (POTS) using standard ELISA methodology [210]. In 116 POTS patients versus 81 healthy controls, they found no differences in GPCR autoantibody concentrations. Moreover, 98.3% of POTS patients and 100% of controls had α_1 -adrenergic receptor antibodies above the detection threshold, raising questions about assay specificity. The authors concluded that CellTrend ELISAs (used in most positive studies) may lack diagnostic value for POTS.

This methodological critique highlights several unresolved issues:

- Whether detected autoantibodies are functionally pathogenic or merely epiphenomenal
- The appropriate control populations and cutoff values
- Whether ELISA-detected antibodies reflect the same populations as functionally active autoantibodies
- The need for functional assays beyond binding detection

? Open Question 2: GPCR Autoantibody Pathogenicity

While correlational and early therapeutic evidence supports a role for GPCR autoantibodies in ME/CFS, definitive proof of causality remains elusive. The Vernino et al. failed replication in POTS raises important questions: Are the autoantibodies detected by current assays the same as those causing symptoms? Do healthy individuals harbor similar autoantibodies that only become pathogenic under certain conditions (e.g., infection, inflammation)? Would more specific functional assays—measuring receptor activation or internalization rather than mere binding—better identify pathogenic autoantibodies? Resolution of these questions will determine whether autoantibody-targeted therapies become a mainstay of ME/CFS treatment.

Other Receptor Autoantibodies Beyond GPCR autoantibodies, additional receptor-targeting antibodies have been identified:

- **α_1 -adrenergic receptor antibodies:** May affect vascular function and contribute to orthostatic intolerance
- **Angiotensin II type 1 receptor antibodies:** May affect blood pressure regulation and fluid homeostasis

These receptor autoantibodies can exert effects through multiple mechanisms:

- Activate receptors (agonistic), causing overstimulation and downstream signaling
- Block receptors (antagonistic), preventing normal ligand binding and signaling
- Induce receptor internalization, reducing cell surface receptor density
- Modulate receptor function in complex, context-dependent ways

Anti-Neuronal Antibodies Autoantibodies targeting nervous system components:

- Anti-ganglioside antibodies
- Anti-neuronal nuclear antibodies
- Antibodies against ion channels
- May contribute to neurological symptoms

Recent cryo-electron microscopy research has mapped the precise binding sites of autoantibodies targeting NMDA receptors in autoimmune encephalitis [211]. These autoantibodies recognize specific antigenic hotspots on the GluN1 amino-terminal domain, causing receptor internalization and neurological dysfunction. While anti-NMDAR encephalitis is a distinct condition, the structural characterization of receptor-targeting autoantibodies provides

a framework for understanding how similar autoantibodies identified in ME/CFS (targeting adrenergic and muscarinic receptors) might cause functional impairment through receptor modulation.

Immunoglobulin Levels

Serum immunoglobulin levels show variable abnormalities:

- **IgG:** May be low (selective IgG subclass deficiency) or elevated
- **IgA:** Sometimes reduced, particularly secretory IgA
- **IgM:** Variable findings
- **IgE:** May be elevated in patients with allergic features

Sex-Specific B Cell Findings from the NIH Study

The deep phenotyping study revealed that female patients showed distinct B cell abnormalities [5]:

Female Patients Women with PI-ME/CFS demonstrated:

- Abnormal B cell proliferation patterns
- Distinct white blood cell growth characteristics
- Different inflammatory markers compared to male patients

These sex-specific findings underscore that ME/CFS may involve fundamentally different immunological processes in men and women, potentially requiring sex-specific therapeutic approaches.

7.3 Cytokines and Inflammatory Mediators

Cytokines are signaling proteins that coordinate immune responses. Cytokine abnormalities in ME/CFS have been extensively studied, though findings vary considerably across studies.

7.3.1 Pro-inflammatory Cytokines

Interleukin-1 (IL-1)

IL-1 is a master regulator of inflammation, with IL-1 β often elevated in ME/CFS. Its effects include fever, fatigue, muscle breakdown, and the acute phase response. Notably, IL-1 produces “sickness behavior” in the central nervous system that closely resembles ME/CFS symptoms, and levels may correlate with symptom severity.

Interleukin-6 (IL-6)

IL-6 has both pro- and anti-inflammatory effects and is frequently elevated in ME/CFS, particularly in early illness. This cytokine induces acute phase proteins, promotes B cell differentiation, and crosses the blood-brain barrier to affect central nervous system function. IL-6 correlates with fatigue in other conditions, suggesting a mechanistic link to this cardinal ME/CFS symptom.

Tumor Necrosis Factor-Alpha (TNF- α)

TNF- α is a central inflammatory cytokine elevated in some ME/CFS studies. It causes fatigue, malaise, and cognitive dysfunction while also affecting mitochondrial function and promoting muscle wasting (cachexia). Variable findings across studies may reflect patient heterogeneity within the ME/CFS population.

Interferons

Type I interferons (IFN- α , IFN- β) are antiviral cytokines elevated in some ME/CFS patients. These interferons cause profound fatigue (as known from their therapeutic use in other conditions) and may indicate ongoing viral activation. Interferon-induced gene expression patterns have been observed in ME/CFS. Type II interferon (IFN- γ) activates macrophages and promotes Th1 responses, though findings in ME/CFS are variable; levels may be elevated or reduced depending on disease stage.

Interleukin-2 (IL-2)

IL-2 is a critical cytokine for T cell function and immune regulation:

- **T cell proliferation:** Essential for clonal expansion of activated T cells
- **Regulatory T cell maintenance:** Required for Treg development and suppressive function
- **NK cell activation:** Enhances NK cell cytotoxicity
- **Memory T cell formation:** Supports long-term immunity
- **Therapeutic use:** Low-dose IL-2 used in autoimmune diseases to boost Tregs; high-dose IL-2 used in cancer immunotherapy

IL-2 signaling requires three receptor subunits (CD25/CD122/CD132) and activates JAK/-STAT pathways. Dysregulation can lead to either immune deficiency (insufficient IL-2 or receptor expression) or autoimmunity (Treg dysfunction). Recent evidence suggests IL-2 pathway abnormalities in ME/CFS (see hypothesis below).

Cytokine Patterns Across Disease Duration

★ Achievement 3: Duration-Dependent Cytokine Signatures

Hornig et al. [212] identified distinct immune signatures in ME/CFS that vary dramatically by disease duration. In a cohort of 298 ME/CFS patients and 348 healthy controls, early-stage patients (illness duration <3 years, n=52) showed prominent activation of both pro- and anti-inflammatory cytokines, with elevated levels of IL-1 α , IL-8, IL-10, IL-12p40, IL-17F, IFN- γ , CXCL1 (GRO- α), CXCL9 (MIG), and IL-5 (all $p < 0.05$, FDR-corrected). A 17-cytokine panel distinguished early ME/CFS from controls with high diagnostic accuracy.

In stark contrast, patients with longer disease duration (>3 years, n=246) had cytokine profiles that normalized to control levels, with no significant differences for most cytokines. This finding represents the first large-scale evidence that ME/CFS immunopathology evolves over time, potentially from initial immune activation to exhaustion or adaptation.

△ Warning 3: Replication Status

Single study (Hornig 2015, n=298 total, but only n=52 in the critical early-stage subgroup). The duration-dependent pattern has not been independently replicated. The small early-stage sample limits confidence in the 3-year inflection point. Cross-sectional design cannot distinguish true longitudinal evolution from cohort effects. Longitudinal studies tracking individual patients over time are needed.

Implications of Duration-Dependent Cytokine Changes The Hornig et al. findings have profound implications:

- **Therapeutic windows:** Early-stage disease may respond better to immunomodulatory therapies targeting active inflammation
- **Study heterogeneity:** Failure to stratify by disease duration explains contradictory findings in previous cytokine studies
- **Biomarker utility:** Cytokine profiling is most useful as a diagnostic tool within the first 3 years of illness
- **Disease progression:** Normalization may reflect immune exhaustion, regulatory adaptation, or shift to different pathological mechanisms

Hornig et al. found that illness duration was more strongly predictive of cytokine patterns than symptom severity in their cross-sectional analysis, suggesting that immune changes primarily reflect disease stage [212]. However, this group-level observation does not preclude severity-related gradients within early-stage or late-stage patients (see following section).

Cytokine-Severity Correlations

★ Achievement 4: Cytokine-Severity Biomarker Panel

Montoya et al. [213] demonstrated dose-response relationships between cytokines and symptom severity in 192 ME/CFS patients compared to 392 healthy controls. Although only two cytokines differed overall between patients and controls (TGF- β higher and resistin lower), 17 cytokines showed statistically significant upward linear trends correlating with disease severity. Thirteen of these 17 are proinflammatory, including CCL11 (Eotaxin-1), CXCL1 (GRO- α), CXCL10 (IP-10), IFN- γ , IL-4, IL-5, IL-7, IL-12p70, IL-13, IL-17F, G-CSF, GM-CSF, and TGF- α .

This dose-response relationship—rather than simple binary patient-control comparison—provides stronger evidence that immune activation tracks with symptom burden. The findings suggest cytokine profiling could stratify patients for clinical trials and identify individuals likely to benefit from anti-inflammatory therapies.

△ Warning 4: Replication Status: Partially Replicated

The cytokine-severity correlation pattern is directionally consistent with Hornig 2015's findings, providing convergent support from a different analytic approach. However, specific cytokine identities differ between studies, and neither has been independently replicated with the same panel and methodology.

Notably, CXCL9 (MIG) inversely correlated with fatigue duration, showing higher levels in early disease and lower levels in chronic disease [213]. This continuous inverse correlation mirrors Hornig's group-level finding of elevated early-disease cytokines, providing convergent support from a different analytic approach (within-group correlation versus cross-sectional comparison of early vs. late subgroups).

Sex-Specific Cytokine Dysregulation

Observation 40 (Sex and Hormonal Influences on Immune Activation). Recent work by Che et al. [189] in a large multi-center cohort revealed that hyperinflammatory cytokine responses are particularly pronounced in women over 45 years of age with diminished estradiol levels. Using multi-omics analysis including microbial stimulation assays (heat-killed *Candida albicans*), the study demonstrated exaggerated production of IL-6 and other proinflammatory cytokines in ME/CFS patients, with responses amplified before and especially after exercise.

The sex- and hormone-specific pattern provides mechanistic insight into the female predominance of ME/CFS (approximately 3:1 female-to-male ratio) and suggests potential therapeutic interventions, such as estrogen supplementation for post-menopausal women with evidence of immune hyperactivation.

This sex-specific finding complements the NIH deep phenotyping study's observation of distinct immune abnormalities in male versus female patients [5], underscoring that ME/CFS pathophysiology may differ fundamentally between sexes.

Integrated Model: Duration, Severity, and Sex

Combining findings from Hornig [212], Montoya [213], and Che [189], an integrated model of cytokine dysregulation emerges:

- **Disease duration:** Early disease (<3 years) shows high cytokines at the group level; late disease (>3 years) shows normalized group-level cytokines
- **Disease severity:** Within patient cohorts, severe patients show higher proinflammatory cytokines than mild patients through dose-response relationships
- **Sex and hormones:** Women, particularly post-menopausal women with low estradiol, show more pronounced immune activation

Reconciling Duration and Severity Effects The Hornig and Montoya findings are not contradictory but complementary. Hornig examined group differences between early-stage and late-stage patients, finding that the early-stage group as a whole had elevated cytokines. Montoya examined severity gradients *within* their cohort (which included both early and late patients), finding that more severe patients had higher cytokines regardless of duration. These observations can coexist: early disease may be characterized by overall immune activation (shifting the entire distribution upward), while severity effects create gradients within both early and late subgroups. The interaction between duration and severity has not been directly tested in a study stratified by both factors simultaneously.

Clinical Application This integrated model suggests personalized treatment approaches, though these represent theoretical predictions requiring validation:

- **Early + severe + female + low estradiol:** Predicted to have highest cytokines; most likely to benefit from immunomodulatory therapies (extrapolated from individual studies)
- **Late + severe + female:** May have severity-driven inflammation despite duration-dependent normalization; immune status assessment needed
- **Late + mild + male:** Predicted to have lowest cytokines; may require therapeutic strategies targeting mechanisms beyond acute immune activation
- **All other phenotypes:** Require individualized immune profiling before treatment selection

The implications of patient heterogeneity for treatment stratification and the concept of distinct ME/CFS subtypes are discussed in Chapter 13, Section 13.8.

No study has yet examined all three factors (duration, severity, sex/hormones) simultaneously in a fully stratified design. The clinical predictions above are extrapolations from separate studies and require prospective validation.

Limitation 3: Integrated Cytokine Model: Extrapolated from Separate Studies

The integrated duration/severity/sex model combines findings from Hornig 2015, Montoya 2017, and Che 2025—three studies using different cohorts, cytokine panels, analytic methods, and stratification criteria. No study has simultaneously examined all three factors. The clinical predictions (e.g., “early + severe + female + low estradiol” = highest cytokines) are logical extrapolations, not empirical findings. The interaction effects between duration, severity, and sex may be non-additive in ways not predictable from individual studies.

IL-2 as Emerging Biomarker Target**~ Hypothesis 2: IL-2 Pathway in ME/CFS Pathophysiology**

Certainty: 0.45. Two independent methodological approaches (extracellular vesicle proteomics and epigenetic chromosome conformation) converge on IL-2 pathway dysregulation, lending moderate confidence. However, whether this reflects a causal role or an epiphenomenon of chronic immune activation, and whether the two findings reflect the same underlying process, remain unresolved.

Two independent methodologies implicate the IL-2 pathway in ME/CFS, though through different mechanisms. Giloteaux et al. [214] found significantly elevated IL-2 specifically in extracellular vesicles from ME/CFS patient plasma (n=49 patients, n=49 controls; $q=0.007$ after multiple comparison correction), with proinflammatory cytokines CSF2 and TNF α correlating with physical and fatigue symptom severity. Independently, Hunter et al. [215] used epigenetic profiling (EpiSwitch[®] technology) of chromosome conformation in 47 ME/CFS patients versus 61 controls, identifying IL-2 signaling among dysregulated pathways in a 200-marker panel (92% sensitivity, 98% specificity in validation).

The convergence—extracellular vesicle cytokine content in one study, epigenetic regulation in another—suggests the IL-2 pathway warrants focused investigation. However, several questions remain: Do elevated IL-2 levels in extracellular vesicles reflect the same process as epigenetic dysregulation of IL-2 signaling? Are ME/CFS cells producing excess IL-2, responding abnormally to normal IL-2, or both? Does IL-2 dysfunction contribute causally to symptoms or merely correlate with disease? Further studies measuring IL-2 receptor expression, downstream signaling (JAK/STAT pathway), and functional T-cell responses to exogenous IL-2 could clarify the pathway’s role and therapeutic potential.

7.3.2 Anti-inflammatory Cytokines**Interleukin-10 (IL-10)**

IL-10 is a potent immunosuppressive cytokine with variable findings in ME/CFS. Levels may be elevated (potentially reflecting an attempt to control inflammation) or reduced (which would permit inflammation to continue). IL-10 is important for resolving immune responses and is produced by regulatory T cells and other cell types.

Transforming Growth Factor-Beta (TGF- β)

TGF- β has immunosuppressive and tissue remodeling functions and is often elevated in ME/CFS. This elevation may represent an attempt to control inflammation, though chronic elevation can promote fibrosis. TGF- β is also important for regulatory T cell development.

Balance Between Pro- and Anti-inflammatory Signals

The key issue in ME/CFS may not be absolute cytokine levels but rather the balance between pro- and anti-inflammatory signals. Patients may exhibit imbalanced pro-/anti-inflammatory ratios, inappropriate cytokine responses to stimuli, and failure to resolve inflammation properly. This results in chronic low-grade immune activation.

7.3.3 Chemokines

Chemokines direct immune cell migration to sites of infection or inflammation:

Recruitment Patterns

Several chemokines show altered levels in ME/CFS. CCL2 (MCP-1), which recruits monocytes, is often elevated. CCL5 (RANTES) recruits T cells and NK cells, while CXCL8 (IL-8) recruits neutrophils. CXCL10 (IP-10), an interferon-induced chemokine, recruits T cells to sites of inflammation.

Tissue Infiltration

Elevated chemokines may promote immune cell infiltration into tissues such as muscle, brain, and gut, leading to local inflammation and tissue damage. This infiltration generates symptoms through inflammatory mediators acting at sites of tissue involvement.

7.4 Immune Activation and Inflammation

7.4.1 Chronic Immune Activation

Evidence for ongoing immune activation in ME/CFS includes:

Activation Markers

Multiple markers of immune activation are elevated in ME/CFS. Neopterin, produced by activated macrophages, is often elevated. β_2 -microglobulin, a marker of immune cell turnover, is frequently increased. Soluble CD25 (sIL-2R) is released by activated T cells, while soluble CD14 indicates monocyte and macrophage activation.

Consequences for Energy Metabolism

Chronic immune activation is metabolically expensive. Immune cells are highly metabolically active, and cytokines alter whole-body metabolism, creating competition for nutrients between immune and other tissues. This metabolic drain may partially explain the profound fatigue characteristic of ME/CFS.

Connection to Symptoms

Cytokines and inflammatory mediators directly cause many ME/CFS symptoms. Fatigue is induced by IL-1, IL-6, TNF- α , and interferons. Cognitive dysfunction results from pro-inflammatory cytokines crossing the blood-brain barrier. Pain arises from sensitization of nociceptors by inflammatory mediators, while sleep disturbance reflects cytokine effects on sleep regulation. Fever and chills result from pyrogenic cytokines.

7.4.2 Neuroinflammation

The brain was traditionally considered “immune privileged,” but it is now recognized that peripheral inflammation affects brain function.

Microglial Activation

Microglia are the brain’s resident immune cells. PET imaging shows increased TSPO binding, a marker of microglial activation, which persists years after initial infection. Activated microglia produce local cytokines that affect neuronal function, potentially explaining the cognitive symptoms prevalent in ME/CFS.

Blood-Brain Barrier Dysfunction

Compromise of the blood-brain barrier permits entry of peripheral cytokines and infiltration of immune cells into the central nervous system. This dysfunction also exposes the brain to circulating autoantibodies and, in some cases, allows direct pathogen entry.

Cytokine Effects on Brain Function

Peripheral cytokines affect the brain through multiple routes: transport across the blood-brain barrier, signaling via vagal afferents, acting at circumventricular organs (which lack a blood-brain barrier), and inducing local cytokine production by glial cells. These cytokines produce multiple brain effects, including altered neurotransmitter synthesis and release, changed receptor expression, and modified synaptic plasticity. The resulting “sickness behavior” encompasses fatigue, social withdrawal, and anhedonia—symptoms prominently featured in ME/CFS.

Neuroimaging Evidence

Studies have demonstrated:

- Increased microglial activation on PET
- Elevated CSF inflammatory markers
- Correlation between brain inflammation and symptoms
- Persistence of neuroinflammation

7.5 Viral Reactivation and Persistence

Many ME/CFS cases follow acute infections, and evidence suggests ongoing viral activity in some patients [3].

7.5.1 Herpesviruses

Human herpesviruses establish lifelong latent infections with potential for reactivation.

Epstein-Barr Virus (EBV)

EBV infects B cells and establishes latency [216]:

- **Acute infection:** Infectious mononucleosis is a common ME/CFS trigger [217]
- **Reactivation markers:** Elevated early antigen (EA) antibodies, viral load
- **Prevalence:** 10–20% of ME/CFS patients show evidence of reactivation [3]
- **Mechanism:** May drive chronic B cell activation and autoantibody production [5]

EBV-Infected B Cells and CNS Demyelination Recent research has demonstrated a direct mechanism by which EBV-infected B cells can cause neurological damage [164]. Autoreactive B cells identified in healthy human blood can cross the blood–brain barrier following viral infection of the cerebrum. When these B cells express EBV Latent Membrane Protein 1 (LMP1), they can infiltrate the brain and induce demyelinating lesions through direct myelin antigen capture followed by complement activation and microglial activation. While this research focused on multiple sclerosis pathogenesis, the mechanism has potential relevance for ME/CFS given the documented role of EBV as a disease trigger, the neuroinflammation observed in ME/CFS patients, and the overlap between ME/CFS and MS symptomatology. This finding provides a concrete pathway by which post-infectious immune dysregulation could lead to CNS involvement.

EBV Infection During Adolescent Immune Development

~ Hypothesis 3: EBV-Adolescence Autoimmune Window

Certainty: 0.50. EBV infection during adolescence may create unique risk for persistent autoantibody-mediated ME/CFS due to coincidence of viral B cell infection with pubertal immune maturation. The certainty level reflects: (1) strong epidemiological association between EBV-triggered mononucleosis and ME/CFS onset, particularly in adolescents; (2) well-characterized immune maturation during puberty; (3) documented mechanisms for EBV-driven autoimmunity; (4) however, EBV is ubiquitous and most infected adolescents do not develop ME/CFS, suggesting additional required factors; (5) the specific contribution of infection timing versus other variables (genetic susceptibility, viral strain) remains uncertain.

We hypothesize that EBV infection during adolescence creates a unique risk for persistent autoantibody-mediated ME/CFS, because viral B cell infection coincides with pubertal immune maturation when tolerance mechanisms are being reorganized.

Epidemiological context: Infectious mononucleosis (primary EBV infection) is a common ME/CFS trigger, particularly in adolescents. While young children typically experience asymptomatic primary EBV infection, delayed first exposure in adolescence produces symptomatic mononucleosis in 35–50% of cases [216]. This age-dependent presentation reflects developmental differences in immune response. When infection occurs during adolescence or young adulthood, symptoms can be more severe than in younger children, and the infection is associated with increased risk for subsequent autoimmune disease development [217].

Immunological timing hypothesis: EBV preferentially infects B cells, establishing lifelong latency. During adolescence, the immune system undergoes substantial reorganization: thymic output is declining, peripheral tolerance mechanisms are maturing, and the B cell repertoire is being shaped. EBV infection during this critical window may:

- Infect B cells during active repertoire selection, potentially immortalizing autoreactive clones that would otherwise be deleted
- Disrupt tolerance checkpoint establishment, allowing autoreactive B cells to persist
- Drive aberrant germinal center reactions producing GPCR autoantibodies (Section 7.6.1)

- Create long-lived plasma cells secreting autoantibodies that persist for decades

Age-dependent outcome predictions:

- *Young children* (<10 years): Immune system still highly plastic; ongoing development may clear aberrant B cell clones through mechanisms described in Hypothesis 7.6.3. Higher recovery probability.
- *Adolescents* (10–18 years): Infection at the edge of immune maturation; some patients clear aberrant clones, others do not. Variable outcomes, overall high recovery rates.
- *Young adults* (18–25 years): Tolerance mechanisms largely established; aberrant B cell populations persist indefinitely. Lower recovery probability.
- *Adults* (>25 years): No developmental clearance mechanism; autoantibody-producing cells become permanent. Recovery rare without intervention.

Treatment implications: If this hypothesis is correct, B cell depletion therapy (rituximab) might be particularly effective in adolescents and young adults with recent EBV-triggered ME/CFS, before long-lived plasma cells establish permanent autoantibody production. The timing of intervention relative to disease onset may be critical—early B cell depletion could prevent establishment of pathogenic plasma cell populations.

Research directions:

1. Compare GPCR autoantibody titers by age at ME/CFS onset and EBV status
2. Track autoantibody trajectories in EBV-triggered versus non-EBV-triggered cases
3. Assess whether EBV-triggered cases show different B cell subset distributions
4. Trial of early rituximab in adolescents with recent EBV-triggered ME/CFS

Limitations: This hypothesis is speculative. EBV is ubiquitous (95% adult seropositivity), so most ME/CFS patients will have been infected regardless of trigger. The specific role of infection timing versus other factors (genetic susceptibility, viral strain, co-infections) is unknown. Additionally, many adolescents with EBV-triggered ME/CFS do recover, suggesting protective factors beyond simple timing. See Section 33.3 for a proposed study design that could inform this hypothesis.

Human Herpesvirus 6 (HHV-6)

HHV-6 infects T cells and can integrate into chromosomes [3]:

- Two species: HHV-6A and HHV-6B
- Evidence for active infection in some ME/CFS patients [3]
- Can affect mitochondrial function
- Neurotropic (infects brain tissue)

Cytomegalovirus (CMV)

CMV establishes latency in monocytes and other cells [3]:

- Reactivation documented in some ME/CFS patients [3]

- Can cause significant inflammation upon reactivation
- Associated with T cell exhaustion [119]

Reactivation Patterns and Causal Relationships

The relationship between herpesvirus reactivation and ME/CFS immune dysfunction remains incompletely understood. Three mechanistic hypotheses can be distinguished by their testable predictions:

~ Hypothesis 4: Viral Reactivation as Consequence

Certainty: 0.45. Consistent with documented NK cell dysfunction impairing viral clearance; however, the directionality of causation between NK dysfunction and viral reactivation has not been experimentally established in ME/CFS.

If reactivation is primarily a consequence of impaired immune control (particularly NK cell dysfunction), then: (1) improving NK cell function should reduce viral titers without affecting other ME/CFS symptoms; (2) viral reactivation markers should correlate with NK cell cytotoxicity but not independently predict symptom severity; (3) antiviral therapy alone should have minimal clinical benefit.

Speculation 4 (Viral Reactivation as Cause). **Certainty: 0.30.** Limited antiviral trial data (Lerner, Montoya) show some benefit in subgroups, but controlled trials have not consistently demonstrated that viral suppression produces sustained clinical improvement across the ME/CFS population, lowering confidence in this model as a universal driver.

If reactivation is a primary driver of ongoing immune activation, then: (1) antiviral therapy should reduce both viral titers and immune activation markers (cytokines, immune cell activation); (2) viral load should independently predict symptom severity after controlling for immune markers; (3) successful viral suppression should produce sustained clinical improvement.

~ Hypothesis 5: Bidirectional Feedback Loop

Certainty: 0.50. This model is the most mechanistically plausible given the documented bidirectional interactions between immune dysfunction and viral reactivation; it is consistent with the partial and heterogeneous response to antiviral monotherapy, though direct experimental evidence for a self-sustaining cycle in ME/CFS remains limited.

If reactivation and immune dysfunction form a self-sustaining cycle, then: (1) interventions targeting either viral replication or immune dysfunction should produce partial but incomplete benefit; (2) combined antiviral and immune-modulating therapy should be synergistic; (3) breaking the cycle at any point should eventually normalize both viral titers and immune function, though with temporal lag.

Current evidence does not definitively distinguish these mechanisms, though the limited efficacy of antiviral monotherapy in most ME/CFS patients suggests reactivation is unlikely

to be solely causal. Longitudinal studies tracking viral titers, immune markers, and symptom severity following targeted interventions are needed to resolve this question.

⚠ **Limitation 4: Viral Reactivation: Causal Direction Unresolved**

While herpesvirus reactivation is documented in a subset of ME/CFS patients, no study has established the causal direction. Elevated viral titers and reactivation markers could be: (a) a cause of immune dysfunction driving symptoms, (b) a consequence of pre-existing NK cell/T cell dysfunction permitting reactivation, or (c) an epiphenomenon of general immune dysregulation with no independent causal role. The partial and inconsistent response to antiviral monotherapy does not resolve this ambiguity. Longitudinal intervention studies with simultaneous tracking of viral load, immune function, and symptoms are required.

7.5.2 Other Implicated Viruses

Enteroviruses

Enteroviruses (Coxsackieviruses, Echoviruses) have been implicated:

- Detection of viral RNA in muscle and gut biopsies
- Elevated antibodies in some patients
- Possible persistent low-level infection
- Historical associations with epidemic ME/CFS outbreaks

Parvovirus B19

Parvovirus B19 can cause chronic arthritis and fatigue:

- Associated with ME/CFS onset in some patients
- Viral DNA detectable in tissues years after infection
- May persist in bone marrow and synovium

SARS-CoV-2 and Long COVID

The COVID-19 pandemic highlighted viral triggers for ME/CFS-like illness:

- Long COVID shares many features with ME/CFS
- Viral persistence documented in some patients
- Similar immune abnormalities observed
- Provides opportunity to study post-infectious ME/CFS from known onset

7.5.3 Tick-Borne Infections

Tick-borne infections represent an important and often underdiagnosed trigger for ME/CFS-like illness. The clinical overlap between post-treatment Lyme disease syndrome (PTLDS), ME/CFS, and chronic tick-borne infections creates significant diagnostic and therapeutic challenges.

Lyme Disease and Post-Treatment Lyme Disease Syndrome

Acute Lyme Disease. Lyme disease, caused by *Borrelia burgdorferi* (North America) or *Borrelia afzelii/garinii* (Europe), is transmitted by *Ixodes* ticks and represents the most common vector-borne infection in temperate regions [218]:

- **Incidence:** >470,000 cases annually in the United States [218]
- **Geographic expansion:** Endemic areas expanding due to climate change and deer population increases
- **Characteristic presentation:** Erythema migrans (bulls-eye rash) in 70–80% of cases; flu-like illness, arthralgia, neurological symptoms
- **Standard treatment:** 2–4 weeks of oral doxycycline or amoxicillin for early localized disease

Post-Treatment Lyme Disease Syndrome (PTLDS). Approximately 10–20% of patients treated for Lyme disease develop persistent symptoms despite standard antibiotic therapy [219]:

- **Defining features:** Fatigue, cognitive dysfunction (“brain fog”), musculoskeletal pain persisting ≥6 months post-treatment
- **Symptom overlap with ME/CFS:** 26 of 29 core ME/CFS symptoms are present in PTLDS patients [219]; however, the proportion meeting formal ME/CFS diagnostic criteria has not been systematically determined
- **PEM consideration:** Some PTLDS patients report post-exertional worsening, though this has not been systematically studied with ME/CFS-specific methodology
- **Biomarker studies:** Shared immune abnormalities including altered cytokine profiles and T cell exhaustion markers

Observation 41 (Symptom Concordance Between PTLDS and ME/CFS). Systematic comparison of symptom profiles between PTLDS and ME/CFS cohorts reveals striking overlap [219]. Of the 29 symptoms assessed using the DePaul Symptom Questionnaire, 26 (90%) showed comparable prevalence and severity between conditions. Both groups exhibited: fatigue (100% prevalence), unrefreshing sleep (>90%), cognitive impairment (>85%), post-exertional malaise (>80%), and widespread pain (>75%). This overlap suggests either shared pathophysiology or that PTLDS represents a subset of post-infectious ME/CFS.

Mechanistic Hypotheses for Persistent Symptoms. Several mechanisms may explain symptom persistence after antibiotic treatment:

- **Immune dysregulation:** Persistent inflammation and autoimmunity triggered by infection; molecular mimicry between borrelial antigens and host tissues [220]
- **Microbial persistence:** Controversy exists regarding whether *Borrelia* can persist in tissue reservoirs (synovium, nervous system) despite negative blood tests; biofilm formation may protect organisms
- **Tissue damage:** Irreversible damage to neural, articular, or cardiac tissues during acute infection
- **Microbiome disruption:** Prolonged antibiotic courses may cause persistent gut dysbiosis contributing to symptom chronicity

Bartonella Species

Bartonella species are intracellular bacteria transmitted by various vectors including ticks, fleas, lice, and sand flies.

Species and Transmission.

- *Bartonella henselae*: Cat scratch disease; cats are primary reservoir
- *Bartonella quintana*: Trench fever; transmitted by body lice
- *Bartonella bacilliformis*: Carrión's disease; sand fly transmission in South America
- **Tick transmission:** Multiple *Bartonella* species have been identified in *Ixodes* ticks, suggesting co-transmission with *Borrelia*

Chronic Bartonellosis and ME/CFS-Like Symptoms. Chronic *Bartonella* infection can present with neuropsychiatric and systemic symptoms overlapping with ME/CFS [221]:

- **Neurological:** Encephalopathy, cognitive dysfunction, peripheral neuropathy, neuroretinitis
- **Systemic:** Chronic fatigue, lymphadenopathy, low-grade fever, sweats
- **Dermatological:** Striae-like lesions (characteristic “*Bartonella striae*”), papular eruptions
- **Vascular:** Endothelial dysfunction, vasculitis-like presentations

Observation 42 (*Bartonella* Detection in ME/CFS-Like Illness). Breitschwerdt et al. used specialized enrichment culture techniques to detect *Bartonella* DNA in blood samples from patients with chronic fatigue and neurological symptoms [221]. Of patients tested, 26% were positive for *Bartonella* species DNA. The same study also detected *Babesia* DNA in some patients, though prevalence was not separately reported. Without healthy control data in this report, the clinical significance of detection remains uncertain—*Bartonella* DNA may represent active infection, past exposure, or subclinical carriage. These findings require replication in larger cohorts with appropriate controls to determine whether detection rates exceed background prevalence.

Diagnostic Challenges. *Bartonella* diagnosis is notoriously difficult:

- **Serology limitations:** Sensitivity 40–60%; cross-reactivity between species; seronegative chronic infection documented
- **Culture requirements:** Specialized enrichment culture (BAPGM) over 2–3 weeks; not widely available
- **PCR sensitivity:** Standard PCR may miss low-level bacteremia; requires specialized laboratories
- **Clinical diagnosis:** Often made on clinical grounds with therapeutic trial

Babesia Species

Babesia are intraerythrocytic parasites transmitted by *Ixodes* ticks, frequently co-transmitted with *Borrelia*.

Epidemiology and Presentation.

- **Primary species:** *B. microti* (North America), *B. divergens* (Europe), *B. duncani* (Western US)
- **Clinical syndrome:** Fever, hemolytic anemia, thrombocytopenia, splenomegaly; can be asymptomatic
- **Chronic infection:** May persist for months to years, particularly in immunocompromised hosts
- **Co-infection impact:** Babesiosis with concurrent Lyme disease produces more severe illness and longer symptom duration [222]

ME/CFS Relevance.

- **Chronic fatigue:** Persistent infection causes ongoing hemolysis, cytokine activation, and profound fatigue
- **Co-infection complexity:** Patients with ME/CFS-like symptoms after tick exposure may have undiagnosed *Babesia* as sole or co-pathogen
- **Treatment complexity:** Requires different antimicrobial regimen than Lyme disease; may explain antibiotic treatment failures

Other Tick-Borne Pathogens

Additional tick-borne infections may trigger or contribute to ME/CFS-like illness:

Anaplasmosis and Ehrlichiosis.

- **Pathogens:** *Anaplasma phagocytophilum*, *Ehrlichia chaffeensis*, *E. ewingii*
- **Clinical features:** Fever, leukopenia, thrombocytopenia, elevated transaminases
- **Chronic sequelae:** Less well-characterized than PTLDS, but persistent symptoms reported

Rickettsia Species.

- Rocky Mountain spotted fever (*R. rickettsii*), other spotted fever groups
- Can cause severe acute illness with potential for chronic neurological sequelae

Tick-Borne Relapsing Fever.

- *Borrelia hermsii*, *B. turicatae*, and related species
- Characterized by recurring febrile episodes
- May be confused with Lyme disease due to genus similarity

Clinical Implications for ME/CFS Evaluation

When to Consider Tick-Borne Infections. Tick-borne infection evaluation should be considered in ME/CFS patients with:

- Geographic residence or travel to endemic areas
- Known tick exposure or recall of erythema migrans rash
- Onset following outdoor activities in wooded/grassy areas
- Symptoms suggesting disseminated Lyme: migratory arthralgias, facial palsy, heart block
- Marked sweats, air hunger, or hemolytic laboratory abnormalities (suggesting *Babesia*)
- Neuropsychiatric predominance with striae-like skin lesions (suggesting *Bartonella*)
- Previous inadequately treated or seronegative Lyme disease

Diagnostic Approach.

- **Lyme disease:** Two-tier testing (EIA/IFA followed by Western blot); consider C6 peptide ELISA; PCR on synovial fluid for Lyme arthritis
- **Babesiosis:** Blood smear, *Babesia* PCR, antibody testing; repeat testing during symptomatic episodes
- **Bartonellosis:** Serology (IgG, IgM), enrichment culture (specialized laboratories), PCR
- **Co-infection panels:** Given frequent co-transmission, comprehensive tick-borne disease panels are warranted

△ Warning 5: Diagnostic Limitations and Controversy

Tick-borne infection diagnosis remains controversial, with significant disagreement between IDSA/AAN/ACR guidelines and organizations like ILADS. Key issues include:

- Sensitivity of standard two-tier testing (estimated 30–40% in early disease, though estimates vary by study [220])
- Interpretation of “indeterminate” Western blots
- Validity of clinical diagnosis in seronegative patients
- Role of prolonged antibiotic therapy (not supported by controlled trials, but advocated by some practitioners)

Patients and clinicians should be aware of these controversies and the current limitations of evidence for chronic tick-borne infection treatment.

Treatment Considerations.

- **Acute Lyme:** Standard 2–4 week doxycycline course is well-established
- **PTLDS:** No treatment proven effective in controlled trials; extended antibiotic courses not recommended by IDSA [218]; symptomatic management similar to ME/CFS
- **Babesiosis:** Atovaquone plus azithromycin (7–10 days, longer for immunocompromised); clindamycin plus quinine for severe cases
- **Bartonellosis:** Prolonged antibiotic courses (weeks to months) often required; regimens include doxycycline, azithromycin, rifampin combinations
- **Co-infections:** Require treatment of all identified pathogens; single-agent therapy may be inadequate

? Open Question 3: Chronic Tick-Borne Infections as ME/CFS Trigger

What proportion of ME/CFS cases have an undiagnosed tick-borne infection as the inciting event or ongoing driver? Given the symptom overlap between PTLDS and ME/CFS, improved diagnostic tools for chronic *Borrelia*, *Bartonella*, and *Babesia* infections could identify a treatable subset. Key research needs include: development of more sensitive diagnostic assays; prospective studies of tick-borne infection cohorts for ME/CFS development; controlled treatment trials in patients with documented chronic infection.

7.5.4 Infection-Induced Cumulative Damage and Disease Progression

The progressive, often step-wise deterioration seen in many ME/CFS patients following repeated infections or viral reactivation suggests that each infectious event produces cumulative, irreversible damage rather than merely triggering reversible inflammation.

Speculation 5 (Infection-Induced Irreversible Damage: The Ratchet Model). **Certainty: 0.40.** This model is mechanistically plausible and consistent with clinical observations of step-wise deterioration after infections, and is supported by preliminary Long COVID epidemiological data [167]. However, the irreversibility claim is difficult to test prospectively, and alternative

explanations (e.g., deconditioning, psychological factors in rating, or regression to the mean) have not been excluded. The certainty is limited by the absence of controlled longitudinal data directly testing the ratchet prediction.

We propose that each infection in ME/CFS patients produces cumulative, irreversible damage that progressively worsens disease through multiple interconnected mechanisms [167].

Cumulative damage mechanisms *Viral reactivation and persistent viral load:* ME/CFS patients frequently experience reactivation of latent viruses (EBV, HHV-6, CMV) or recurrent infections with new pathogens. Each reactivation adds to the total viral antigenic load. Unlike an immunocompetent host who clears viruses completely, ME/CFS patients with impaired immune function may never fully clear these reactivations. The viral genome and viral proteins (which are inherently immunogenic and inflammatory) persist or accumulate. This creates a ratchet effect—viral burden goes up with each reactivation and rarely returns to baseline. A specific cellular mechanism for this persistence—virus residing within long-lived lymphocytes themselves, below conventional detection thresholds—is proposed in Hypothesis 7.2.1, supported by comparative virology evidence from feline coronavirus [171].

Additional microglial priming events: As described in the neuroinflammatory cascade model (Hypothesis 8.1.6), each infection represents a major microglial priming event. Acute infections trigger intense microglial activation, and subsequent viral reactivations produce additional priming. Since primed microglia show exaggerated responses to subsequent stimuli (as discussed in the PEM kindling hypothesis, Hypothesis 8.1.7), each infectious episode not only causes direct damage but increases the microglial response to future infections. This creates a positive feedback: infection → microglial priming → exaggerated response to next infection → more priming.

Critical note on model interdependence: This ratchet model shares core mechanistic assumptions with the Kindling Hypothesis (Hypothesis 8.1.7) and the neuroinflammatory cascade model (Hypothesis 8.1.6); they should be interpreted as complementary components of a unified explanatory framework rather than independent corroboration of each other. Specifically: the Ratchet Model predicts irreversible step-wise decline with each infection, while the Kindling Model predicts progressive threshold reduction from exertion triggers. Both could be simultaneously true (infections cause larger priming steps; exertion causes smaller priming steps), but they make distinct testable predictions. The Ratchet Model uniquely predicts that baseline functioning follows a ratchet pattern (asymmetric: high damage but minimal recovery), whereas pure Kindling predicts threshold reduction independent of infection status. Distinguishing between these predictions empirically requires longitudinal threshold tracking with separate quantification of crash frequency from infections versus non-infectious triggers.

Further depletion of metabolic reserves: During acute infection, energy expenditure increases substantially due to fever, immune activation, and metabolic stress. In ME/CFS patients, metabolic reserves are already depleted. Each infection represents a major metabolic stress that exhausts remaining reserves. Unlike immunocompetent hosts who recover metabolically after infection, ME/CFS patients may never fully restore their metabolic baseline before the next infection occurs. The metabolic nadir becomes progressively lower with each infection.

Immune exhaustion from repeated activation: The adaptive immune system responds to each infection by activating clones of T cells and B cells specific to the infection. In the context of persistent and recurrent infections, these same clones are repeatedly activated. Repeated activation produces immune exhaustion—T cell exhaustion markers increase, B cell function declines. Additionally, the repeated need to generate immune responses may accelerate telomere shortening and cellular senescence, reducing the lifespan of immune cells.

Clinical manifestation: Step-wise baseline deterioration The combination of these mechanisms produces a characteristic clinical pattern: each infection is followed by a step-wise decline in baseline functioning that does not fully resolve before the next infection. A patient might experience:

1. Baseline functioning: Level A (e.g., able to work 4 hours daily)
2. Infection 1 → acute illness → recovery to baseline attempt, but only reaches Level B (3 hours daily) due to incomplete metabolic recovery and persistent microglial priming
3. Infection 2 → acute illness → recovery to attempted baseline, but only reaches Level C (2 hours daily)
4. Infection 3 → acute illness → recovery to attempted baseline, but only reaches Level D (bedbound)

This step-wise progression differs from other chronic conditions where infections represent temporary setbacks from which full recovery to baseline is expected. In the ratchet model, each infection represents a permanent downward step in baseline capacity. Over years, repeated infections can convert a mildly-affected patient into a severely-affected patient, even if individual infections are not severe.

Key clinical implications *Infection prevention is disease-modifying:* In the ratchet model, preventing infections is not merely symptomatic management but disease-modifying therapy. Each prevented infection preserves baseline functioning and prevents another step-wise decline. A patient with effective infection prevention can potentially avoid progressive deterioration that would occur with repeated infections.

Prophylactic interventions are justified: Standard infection prevention approaches (masking during high-transmission periods, hand hygiene, limiting exposure to ill contacts) might be expected in moderately or severely affected patients. More aggressive approaches—such as FFP2 masking in community settings during respiratory season, or prophylactic antiviral therapy during high-risk periods if safe options become available—could potentially have substantial long-term benefit by preventing cumulative damage.

Rapid infection treatment is critical: Early, aggressive treatment of identified infections (rapid antiviral therapy for herpesvirus reactivation, prompt antibiotic therapy for bacterial infections) might minimize the damage window and reduce the microglial priming response by shortening infection duration.

Immunological intervention may not restore lost function: In conditions where immune deficiency is reversed (e.g., HIV treatment restoring CD4 counts), patients often improve dramatically because the deficit was reversible. In the ratchet model, infection-induced damage is largely

irreversible. Therefore, immunological interventions (such as immune modulation or restoration) might prevent future decline but would not restore previously-lost baseline functioning. This suggests that prevention is substantially more important than treatment—once damage is done, it persists.

Relationship to baseline deterioration in Long COVID The ratchet model of cumulative infection-induced damage provides a mechanistic framework that explains the step-wise baseline deterioration observed in Long COVID patients experiencing recurrent COVID-19 infections. Preliminary epidemiological data and clinical observations suggest that each COVID reinfection produces additional baseline functional loss beyond what would be expected from reinfection alone. This pattern aligns with the infection ratchet hypothesis and suggests that similar mechanisms may apply to ME/CFS [167].

⚠ Limitation 5: Infection Ratchet Model: Irreversibility Not Demonstrated

The ratchet model's central claim—that each infection produces *irreversible* baseline decline—has not been tested prospectively. Key data gaps:

- No longitudinal study has tracked ME/CFS patients through sequential infections with pre- and post-infection functional measurement to quantify stepwise decline versus recovery.
- The model shares core assumptions with the Kindling Hypothesis and the neuroinflammatory cascade model; these cannot serve as independent corroboration of each other.
- Alternative explanations for observed step-wise decline (deconditioning during acute illness, psychological reporting bias, regression to the mean after flare) have not been excluded.
- The claim that “immunological intervention may not restore lost function” is unfalsifiable without first identifying which damage is structural versus functional.

7.6 Autoimmunity in ME/CFS

Evidence increasingly supports autoimmune mechanisms in at least a subset of ME/CFS patients.

7.6.1 Autoantibodies Identified

Anti-Nuclear Antibodies

Anti-nuclear antibody (ANA) prevalence is elevated in ME/CFS, with 15–25% of patients testing positive compared to 5–10% in healthy individuals [207]. Various ANA patterns are observed, with positive ANA possibly indicating general immune dysregulation rather than a specific autoimmune disease.

? Open Question 4: Clinical Significance of ANA in ME/CFS

Whether elevated ANA titres in ME/CFS patients represent a pathogenic autoimmune process, an epiphenomenon of systemic immune dysregulation, or a prognostically relevant subgroup marker remains unresolved.

G-Protein-Coupled Receptor (GPCR) Autoantibodies

GPCR autoantibodies represent one of the most well-studied autoantibody classes in ME/CFS, with substantial evidence for their pathogenic role. The B cell abnormalities described in Section 7.2.2 likely contribute to autoantibody generation. For comprehensive coverage of GPCR autoantibodies—including initial discovery, validation across cohorts, correlation with symptom severity, downstream mechanisms, and therapeutic targeting through immunoadsorption, plasma cell depletion, and direct neutralization—see the detailed discussion in Section 7.2.2.

Anti-Neuronal Antibodies

Antibodies targeting nervous system components have been identified in ME/CFS, including anti-ganglioside antibodies, antibodies against voltage-gated ion channels, and anti-neuronal surface antigen antibodies. These autoantibodies may contribute to the neurological symptoms observed in the condition.

7.6.2 Autoimmune Mechanisms**Molecular Mimicry**

Molecular mimicry occurs when structural similarity between pathogen and self-antigens leads antibodies or T cells generated against an infection to cross-react with self-tissues. This phenomenon has been documented for several viruses associated with ME/CFS and may explain the link between infection and subsequent autoimmunity.

Epitope Spreading

Epitope spreading occurs when tissue damage exposes new antigens to the immune system. The initial immune response causes tissue injury, releasing self-antigens that trigger new autoimmune responses. This leads to progressive expansion of autoimmune targets over time.

Loss of Self-Tolerance

Loss of self-tolerance occurs when regulatory mechanisms fail. Treg dysfunction permits autoreactive cells to escape suppression, while B cell tolerance checkpoints fail to eliminate autoreactive B cells. Chronic inflammation further promotes autoimmunity by creating a permissive environment for autoimmune responses.

⚡ Limitation 6: Autoimmune Mechanisms: Extrapolated from Other Diseases

Molecular mimicry, epitope spreading, and loss of self-tolerance are well-characterised mechanisms in established autoimmune diseases (SLE, rheumatoid arthritis, type 1 diabetes). Their role in ME/CFS is inferred by analogy, not demonstrated:

- No specific molecular mimicry target (pathogen epitope → self-antigen cross-reaction) has been identified in ME/CFS.
- Epitope spreading has not been documented longitudinally in ME/CFS cohorts.
- Treg dysfunction is reported in ME/CFS but has not been causally linked to loss of tolerance versus being a downstream consequence of chronic immune activation.
- The proportion of ME/CFS patients with a genuinely autoimmune aetiology (as opposed to immune dysregulation without autoimmunity) remains unknown.

7.6.3 Developmental Immune Tolerance and Recovery

The autoimmune mechanisms described above—molecular mimicry, epitope spreading, and loss of self-tolerance—operate in both pediatric and adult ME/CFS patients. Yet pediatric patients recover at dramatically higher rates (54–94%) than adults ($\leq 22\%$), often despite similar autoantibody profiles at disease onset. This paradox suggests that developing immune systems may possess unique mechanisms for eliminating aberrant immune memory that adult systems lack.

~ Hypothesis 6: Immune Memory Pruning in Development

Certainty: 0.50. Pediatric recovery from ME/CFS may be facilitated by developmental immune tolerance mechanisms that actively delete or reprogram aberrant immune memory cells—a process termed “immune memory pruning.” The developing immune system, particularly during puberty and adolescence, appears to undergo quality control checkpoints that can eliminate autoreactive B cells, exhausted T cells, and pathological memory populations. The certainty level reflects: (1) well-characterized developmental changes in peripheral B and T cell tolerance mechanisms; (2) evidence for thymic output through adolescence; (3) documented sex hormone effects on immune regulation; (4) however, the specific mechanisms enabling clearance of ME/CFS-associated autoantibodies remain unclear; (5) the distinction between resolution via immune pruning versus other pediatric recovery mechanisms (glial plasticity, HSC regeneration, recovery capital) cannot yet be resolved.

Pediatric recovery from ME/CFS may be facilitated by developmental immune tolerance mechanisms that actively delete or reprogram aberrant immune memory cells—a process we term “immune memory pruning.”

Mechanistic basis:

The developing immune system is not merely a smaller version of the adult system; it is qualitatively different in its capacity for self-correction. Several mechanisms may contribute to immune memory pruning:

Peripheral B cell tolerance checkpoints. Autoreactive B cells that escape central tolerance in the bone marrow normally undergo peripheral tolerance mechanisms including anergy (functional unresponsiveness), receptor editing (modification of the B cell receptor to eliminate autoreactivity), and deletion (apoptosis of autoreactive cells) [223, 224]. These peripheral checkpoints are most active during immune development and may decline with age. Recent evidence demonstrates that impaired clearance of autoreactive B cells due to disruption of negative selection barriers causes autoantibody-producing B cells to mature and subsequently differentiate into plasma cells [223]. In children with ME/CFS, autoantibodies generated during acute illness may be subject to these active tolerance mechanisms, leading to gradual elimination of autoreactive B cell clones. In adults, with attenuated peripheral tolerance, these clones persist.

Thymic contribution to T cell repertoire. The thymus remains highly active through adolescence, continuously generating naive T cells that have undergone rigorous negative selection against self-antigens. This ongoing thymic output may dilute and eventually replace dysfunctional T cell populations (exhausted cells, aberrantly activated cells) that accumulate during ME/CFS. Adult thymic involution eliminates this regenerative capacity, leaving dysfunctional T cell populations to persist.

Pubertal immune reorganization. Puberty involves substantial reorganization of immune function, driven by sex hormones and growth factors. This reorganization period may include “quality control” checkpoints that assess immune memory and eliminate populations that fail to meet tolerance criteria. Adolescents who develop ME/CFS before completing this reorganization may benefit from these checkpoints; those who develop ME/CFS after puberty have already passed this window.

Regulatory T cell plasticity. Tregs in children show greater plasticity and proliferative capacity than adult Tregs. Pediatric ME/CFS patients may mount more effective regulatory responses that suppress autoreactive populations, eventually leading to their deletion through lack of antigenic stimulation.

Testable predictions:

This hypothesis generates several falsifiable predictions:

1. *Autoantibody trajectory differences:* In recovering pediatric patients, autoantibody titers (particularly GPCR autoantibodies) should decline over time, whereas in non-recovering adults, titers should remain stable or increase. Longitudinal measurement of autoantibody levels in pediatric versus adult cohorts would test this prediction.
2. *Tolerance gene signatures:* Peripheral blood from recovering pediatric patients should show gene expression signatures of active tolerance mechanisms—upregulation of genes involved in receptor editing (RAG1, RAG2), anergy (CBLB, ITCH), and dele-

tion (FAS, BIM)—that are absent in adults. RNA-seq comparing pediatric recoverers versus non-recoverers versus adults could identify these signatures.

3. *Thymic output markers:* Recent thymic emigrants (CD31⁺ naive CD4 T cells, T cell receptor excision circles or TRECs) should correlate with recovery probability. Patients with higher thymic output, regardless of age, should have better prognosis.
4. *B cell subset dynamics:* Recovering patients should show declining frequencies of CD21^{lo} atypical memory B cells (associated with chronic immune activation) and autoantibody-secreting plasmablasts, with increasing naive B cell proportions.
5. *Pubertal timing effects:* Among pediatric patients, those who develop ME/CFS pre-pubertally (Tanner stages 1–2) should have higher recovery rates than those developing ME/CFS post-pubertally (Tanner stages 4–5), as the former still have pubertal immune reorganization ahead of them.

Treatment implications:

If immune memory pruning explains pediatric recovery advantage, several therapeutic strategies could be explored to induce analogous processes in adults:

Tolerance-inducing therapies. Low-dose antigen administration can induce peripheral tolerance in autoimmune conditions. If specific autoantigens driving ME/CFS are identified (e.g., GPCR peptides), tolerance induction protocols could be developed.

Regulatory T cell therapy. Expansion and infusion of autologous Tregs could enhance regulatory mechanisms that suppress autoreactive populations.

Timed B cell depletion. Rituximab showed mixed results in ME/CFS trials, but the immune memory pruning hypothesis suggests that B cell depletion might be more effective in younger patients or when combined with therapies that prevent re-emergence of autoreactive clones during B cell reconstitution. The timing of B cell depletion relative to immune developmental stage may be critical.

Thymic rejuvenation. Experimental approaches to restore thymic function (IL-7 therapy, sex steroid ablation, thymic transplantation) could potentially restore the naive T cell output that enables immune repertoire renewal.

Limitations and caveats:

This hypothesis remains speculative and requires validation. Alternative explanations for the pediatric recovery advantage exist, including greater metabolic reserves, neural plasticity, psychosocial factors, and ascertainment bias (milder cases preferentially diagnosed in children). The hypothesis does not explain why some children do not recover or why rare adults do recover. Additionally, even if the hypothesis is correct, translating developmental tolerance mechanisms into adult therapeutics presents substantial challenges.

Enterochromaffin-Vagal Pathway: Gut-Immune-Brain Axis Recent work by Wirth and Scheibenbogen (2025) [225] on the enterochromaffin-vagal pathway provides novel insights into gut-immune-brain interactions that may influence autoimmune processes in ME/CFS. The full mechanistic evidence chain—butyrate-enterochromaffin serotonin-vagal signalling—is developed in Section 11.1.4 of Chapter 11; here we focus on its immune dysregulation implications:

Speculation 6 (Enterochromaffin-Vagal Pathway in ME/CFS Autoimmunity). **Certainty: 0.35.** The enterochromaffin-vagal pathway may contribute to immune dysregulation in ME/CFS through multiple mechanisms [225]. Gut dysbiosis with reduced butyrate production [226, 227] impairs enterochromaffin cell serotonin synthesis [14], leading to reduced vagal afferent signaling [13, 228]. This vagal dysfunction may disrupt immune regulation through:

Potential immune mechanisms:

- **Reduced cholinergic anti-inflammatory pathway activity:** Vagal efferents inhibit cytokine production via the cholinergic anti-inflammatory pathway [13]. Reduced vagal tone may impair this regulatory mechanism.
- **Altered gut immune homeostasis:** Vagal dysfunction may disrupt gut-associated lymphoid tissue (GALT) regulation, contributing to intestinal inflammation and bacterial translocation [227].
- **Neuro-immune feedback disruption:** The vagus nerve provides afferent input to brain regions that regulate immune function, including the hypothalamus and brainstem [228].
- **MCAS bidirectional amplification:** Mast cells and enterochromaffin cells interact in the gut lamina propria [14]. Reduced vagal tone removes cholinergic inhibition of mast cell degranulation [13], potentially amplifying mast cell activation syndrome (MCAS) in susceptible patients [229, 173]. Conversely, chronic histamine elevation may plausibly impair enterochromaffin function and worsen vagal afferent input (speculative; no direct ME/CFS evidence). This bidirectional loop may contribute to the observed co-occurrence of MCAS and autonomic dysfunction in ME/CFS [173].

Implications for autoimmunity:

- Provides a mechanistic link between gut dysbiosis and systemic immune dysregulation
- Suggests that vagal stimulation therapies may have immunomodulatory effects
- Explains how gut-directed therapies (probiotics, butyrate) might influence systemic inflammation
- Offers testable hypotheses for future research on gut-immune-brain interactions

Limitations:

- Preprint status (not peer-reviewed)
- Direct evidence for vagal-immune interactions in ME/CFS is limited
- Causality and directionality remain uncertain
- Clinical implications require validation in controlled studies

Speculation 7 (EBV Reactivation Impairs Gut-Vagal Signalling via IDO2). **Certainty: 0.25.** EBV reactivation is documented in ME/CFS [121]. EBV infection induces interferon- γ [230], which upregulates IDO2, diverting tryptophan into the kynurenine pathway [18, 225] and depleting the substrate available for enterochromaffin serotonin synthesis [14]. Simultaneously, kynurenine pathway products—particularly quinolinic acid—may impair mitochondrial function via PARP-mediated NAD⁺ depletion [231, 232, 233], creating both substrate depletion and cellular dysfunction in enterochromaffin cells.

This chain (EBV reactivation → IFN- γ → IDO2 → tryptophan depletion → impaired enterochromaffin serotonin → vagal afferent dysfunction) may explain why post-EBV ME/CFS patients often have prominent autonomic dysfunction [121]. If correct, patients with high EBV antibody titers or active reactivation would show higher kynurenine/tryptophan ratios, lower platelet serotonin, and more severe autonomic impairment (lower HRV) than non-EBV cases. 5-HTP supplementation, which bypasses the IDO2 diversion point [225], may be particularly relevant for this subgroup.

Limitations: The EBV → IDO2 → gut-vagal chain involves multiple extrapolation steps, each with limited direct ME/CFS evidence. Quinolinic acid effects on enterochromaffin cells have not been studied in ME/CFS. Causal directionality between EBV reactivation and IDO2 upregulation in ME/CFS is unestablished.

Speculation 8 (Autoantibody Inefficiency Hypothesis). Certainty: 0.35. This is a novel mechanistic reinterpretation of existing GPCR autoantibody data. While consistent with published correlational and therapeutic findings [208, 118, 209], the core claim—that autoantibodies increase effort-per-output rather than simply blocking or activating receptors—has not been directly tested. Functional assays distinguishing partial agonism from complete blockade in ME/CFS are absent, and the “effort-to-output ratio” metric does not yet have a validated measurement instrument.

The standard model of autoantibody pathogenesis posits binary effects: agonist antibodies overstimulate receptors, antagonist antibodies block them. However, emerging evidence suggests GPCR autoantibodies in ME/CFS may operate through a more subtle mechanism—**increasing the physiological “effort” required per unit of functional output** rather than preventing output entirely.

Conceptual Framework. Consider two scenarios:

1. **Complete blockade:** A receptor antagonist prevents signal transduction. To achieve a given output, the system compensates by upregulating receptor expression, increasing ligand production, or activating alternative pathways. This produces a stable new equilibrium with normal output at higher baseline cost.
2. **Partial interference:** An autoantibody binds receptors with partial agonist/antagonist activity [234], causing stochastic signal degradation. Some signals succeed, others fail. The system cannot establish stable compensation because the interference is probabilistic rather than deterministic. To achieve reliable output, the system must increase signal redundancy, pathway cross-talk, and error-correction mechanisms—dramatically raising metabolic cost per successful signal.

The autoantibody inefficiency hypothesis proposes that ME/CFS GPCR autoantibodies function primarily through scenario 2: they increase the *effort-to-output ratio* rather than blocking output capacity.

Mechanistic Basis. Multiple mechanisms could contribute to effort-per-output elevation:

Partial agonism and desensitization. GPCR autoantibodies can act as partial agonists, activating receptors less efficiently than endogenous ligands while simultaneously triggering desensitization cascades [234]. This creates a state of chronic low-level activation paired with reduced receptor availability. Functional output requires overcoming both the partial antagonism and the desensitization state, necessitating increased ligand release, receptor cycling, and downstream amplification.

Receptor internalization and trafficking dysregulation. Autoantibody binding can trigger receptor internalization through arrestin-mediated pathways [211]. Unlike normal agonist-induced internalization (which promotes receptor recycling), autoantibody-mediated internalization may produce abnormal trafficking patterns, sequestering receptors in non-productive compartments. The cell must continuously synthesize new receptors to maintain surface expression, increasing basal metabolic demand.

Compensatory pathway activation. When primary signaling pathways operate inefficiently due to autoantibody interference, cells activate compensatory mechanisms [209]. In monocytes, GPCR autoantibody binding triggers production of MIP-1 δ , PDGF-BB, and TGF- β 3—cytokines that activate alternative inflammatory and stress-response pathways. These compensatory cascades consume ATP, generate reactive oxygen species, and demand continuous cellular attention, raising the metabolic cost of maintaining homeostasis.

Stochastic signaling noise. If autoantibodies cause intermittent rather than consistent receptor modulation, downstream systems cannot adapt through simple upregulation. Instead, they must implement redundancy: multiple parallel pathways, error-checking mechanisms, and repeated signal verification. This transforms efficient point-to-point signaling into expensive consensus-based communication, analogous to the difference between a direct phone call and a committee meeting requiring majority vote.

Predictions and Testable Hypotheses. The inefficiency model generates distinct predictions from the simple blockade/activation model:

1. **Effort-symptom dissociation:** Autoantibody-positive patients should show higher perceived effort for equivalent objective tasks compared to autoantibody-negative patients with similar disability levels.
2. **Titer correlation with effort, not capacity:** GPCR autoantibody titers should correlate more strongly with effort-per-task measures than with absolute functional capacity. High-titer patients might maintain near-normal performance briefly but report extreme subjective effort and experience rapid decompensation.
3. **Metabolic cost elevation:** During controlled tasks, autoantibody-positive patients should show elevated oxygen consumption, lactate production, or other metabolic markers for equivalent work output compared to autoantibody-negative patients.
4. **Immunoabsorption reduces effort-per-output:** The primary benefit of immunoabsorption should be reduced subjective effort for equivalent tasks, measurable before changes in absolute capacity. Stein et al. showed SF-36 improvements [118], but effort-normalized metrics were not reported.

5. **PEM pattern differences:** If autoantibodies increase baseline effort, PEM in autoantibody-positive patients should follow an effort-accumulation model (gradual exhaustion of compensatory reserves) rather than a damage-repair model (discrete injury requiring recovery).
6. **CNS vs peripheral manifestations:** If the inefficiency mechanism operates primarily through adrenergic and muscarinic receptors, symptoms should concentrate in systems dependent on rapid, high-frequency receptor signaling: autonomic regulation, attention/vigilance, fine motor control. Gross muscle strength might be relatively preserved despite profound fatigue.

Existing Evidence Alignment. Several published findings align with the inefficiency model:

- **Correlation patterns:** Sotzny et al. found GPCR autoantibody levels correlated with fatigue severity, muscle pain, cognitive impairment, and autonomic dysfunction [208]—all effort-dependent symptoms that worsen with sustained activity.
- **Heterogeneous treatment response:** In the Stein et al. immunoadsorption study, 70% responded but 30% did not [118]. The inefficiency model predicts that patients with primary autoantibody-driven disease should respond dramatically, while those with additional pathology may show limited benefit.
- **Delayed recovery after antibody removal:** While autoantibody titers drop immediately (77% reduction in Stein et al.), clinical improvement evolves over weeks to months. This delay is inconsistent with simple receptor unblocking but consistent with gradual normalization of compensatory mechanisms.
- **Monocyte reprogramming persistence:** Hackel et al. showed GPCR autoantibodies reprogram monocyte cytokine production [209]. This inflammatory signature might persist even after antibody removal if monocytes have been epigenetically reprogrammed.

Clinical Implications. If the inefficiency model is correct, treatment strategies should focus on:

- **Early intervention:** Before compensatory mechanisms become maladaptive, autoantibody removal might produce rapid recovery.
- **Combination therapy:** Immunoadsorption or plasma cell depletion (daratumumab [9]) to remove autoantibodies, paired with therapies supporting cellular energy metabolism (CoQ10, NADH) to offset elevated effort costs during recovery.
- **Effort-based rather than capacity-based pacing:** For autoantibody-positive patients, effort-tracking (heart rate variability, subjective effort scales) might be more predictive of crashes than activity level alone.
- **Patient selection for trials:** The inefficiency model predicts a subgroup—autoantibody-positive, relatively acute onset, without severe mitochondrial damage—should show dramatic response to antibody removal.

Relationship to Other Hypotheses. The autoantibody inefficiency hypothesis complements rather than contradicts other ME/CFS mechanisms:

- **Vicious cycles:** Elevated effort-per-output increases metabolic demand, potentially triggering mitochondrial stress, oxidative damage, and further dysfunction [185]. The inefficiency becomes self-reinforcing.
- **Autonomic dysfunction:** If autoantibodies target adrenergic receptors with partial agonist effects, the autonomic nervous system operates in a state of chronic low-level activation paired with reduced responsiveness.
- **Immune memory:** The immune memory pruning hypothesis (Hypothesis 7.6.3) suggests pediatric patients might clear autoreactive plasma cell populations, while adults cannot—explaining age-related prognosis differences in autoantibody-driven disease.

Limitations and Uncertainties.

1. **Mechanistic heterogeneity:** Different autoantibodies likely have different effects. Lumping all GPCR autoantibodies together may obscure subtype-specific mechanisms.
2. **Assay limitations:** As noted by Vernino et al. [210], current ELISA assays measure antibody binding but not functional effects. Functional assays measuring receptor activation, desensitization kinetics, and downstream signaling are needed.
3. **Causal uncertainty:** Autoantibodies correlate with symptom severity but correlation is not causation. They may be a consequence rather than a cause of ME/CFS.
4. **Quantification challenges:** “Effort per output” is difficult to measure objectively. Developing validated effort-efficiency metrics is essential for testing this hypothesis.
5. **Alternative explanations:** Elevated perceived effort could arise from CNS dysfunction (altered interoception, motivation circuits) rather than peripheral receptor interference.

Despite these limitations, the inefficiency hypothesis provides a coherent framework for understanding why approximately 30% of ME/CFS patients have elevated GPCR autoantibodies, why these correlate with specific symptom domains, why immunoadsorption produces gradual rather than immediate improvement, and why outcome heterogeneity exists.

7.7 Connections to Allergies and Mast Cell Activation

Many ME/CFS patients report increased sensitivity to foods, medications, and environmental factors.

7.7.1 Mast Cell Activation Syndrome (MCAS)

Overlap with ME/CFS

MCAS involves inappropriate mast cell degranulation:

- Substantial symptom overlap with ME/CFS
- Fatigue, cognitive dysfunction, pain common in both
- May represent comorbidity or shared pathophysiology

- Estimated 30–50% of ME/CFS patients may have MCAS features [235]

Mast Cell Phenotype Abnormalities in ME/CFS

Recent research provides objective evidence of mast cell dysfunction in ME/CFS [236]:

- **Naïve mast cells:** Significant increase in CD117⁺CD34⁺FcεRI[−]chymase[−] naïve mast cells in moderate and severe ME/CFS ($p < 0.05$)
- **Activation markers:** Elevated CD40 ligand and MHC-II receptors on differentiated mast cells in severe cases
- **Clinical correlation:** Mast cell abnormalities more pronounced in severe disease
- **Implication:** Demonstrates measurable cellular pathology supporting mast cell involvement in ME/CFS pathophysiology

Histamine and Other Mediators

Mast cells release numerous vasoactive and inflammatory mediators [235]:

- **Histamine:** Causes vasodilation, vascular permeability, brain fog, orthostatic intolerance
- **Platelet-activating factor (PAF):** Triggers vascular leakage, amplifies mast cell activation (vicious cycle)
- **Tryptase:** Marker of mast cell activation; diagnostic if elevated during symptomatic episodes
- **Prostaglandins:** Inflammatory mediators contributing to pain and fatigue
- **Leukotrienes:** Cause bronchoconstriction, vascular dysfunction, inflammation
- **Cytokines:** IL-6, IL-8, TNF- α , VEGF contribute to systemic inflammation

Vascular Pathomechanisms

Mast cell activation shares pathogenic mechanisms with ME/CFS through vascular dysfunction [235]:

- **Spillover of vasoactive mediators** into systemic circulation
- **Histamine's vascular effects:** Worsens orthostatic intolerance via vasodilation and blood pooling
- **β_2 -adrenergic receptor dysfunction:** Amplifies symptoms through impaired vascular regulation
- **Clinical correlation:** ME/CFS patients with MCAS and orthostatic intolerance reported symptom alleviation significantly more often following mast cell-targeted treatment ($p < 0.0001$) [235]

Diagnostic Criteria

MCAS diagnosis requires:

- Typical symptoms (flushing, hives, GI symptoms, cognitive dysfunction, fatigue)
- Elevated mast cell mediators during symptomatic episodes:
 - Tryptase: 20% increase plus 2 ng/mL rise from baseline (must be obtained within 1–4 hours)
 - Urinary N-methylhistamine, prostaglandin D2, or leukotriene E4
- Response to mast cell-directed therapy

Diagnostic challenge: Only small percentage of ME/CFS patients have elevated tryptase; many may have MCAS features without meeting formal diagnostic criteria.

Treatment Implications and Evidence

Critical Evidence on Antihistamine Therapy **Negative trial:** H1 antihistamine alone (terfenadine) showed NO benefit in double-blind RCT of CFS [237]:

- No improvement in symptoms, functioning, or health perceptions
- High-quality evidence demonstrates H1 monotherapy insufficient

Positive case evidence: H1+H2 combination showed dramatic benefit in Long COVID patient meeting ME/CFS criteria [238]:

- Loratadine OR fexofenadine (H1) + famotidine (H2): “helpful with energy and cognitive dysfunction”
- Discontinuation test: Stopping medications → “increased fatigue and increased cognitive dysfunction”
- Resumption: Rapid improvement upon restarting
- Cromolyn 400 mg QID: Heart rate fell from 130–140 bpm to 100–105 bpm
- Quercetin 1000 mg BID: “Improvement in fatigue and allergic symptoms”

Key insight: H1+H2 combination required; H1 alone insufficient.

Antihistamine and Mast Cell Stabilizer Options H1 antihistamines:

- **Standard:** Loratadine, cetirizine, fexofenadine
- **Superior:** Rupatadine (triple action: H1 antagonist + PAF antagonist + mast cell stabilizer) [239, 240]
 - Network meta-analysis: Rupatadine 20 mg highest rank (SUCRA 99.7%) for symptom control
 - 31× more potent than loratadine at PAF antagonism (IC₅₀ 4.6 vs 142 μM)
 - Inhibits mast cell degranulation: IL-8 (80%), VEGF (73%), histamine (88%)

- PAF antagonism addresses vascular pathomechanisms in ME/CFS

H2 antihistamines:

- Famotidine 20–40 mg daily (BID dosing)
- Essential for combination therapy with H1 blockers

Mast cell stabilizers:

- **Quercetin** (natural): 500–1000 mg daily
 - MORE effective than cromolyn in vitro [241]
 - Reduced contact dermatitis >50% in 8 of 10 patients
 - Over-the-counter, well-tolerated
- Cromolyn sodium 200–400 mg QID (prescription)
- Ketotifen 1–2 mg BID (not FDA-approved in US)

Amitriptyline (dual benefit for pain/sleep + mast cells):

- 10–50 mg bedtime
- Specific mast cell inhibition: Reduces IL-8, VEGF, IL-6, histamine release [242]
- **Unique to amitriptyline:** Other antidepressants (bupropion, citalopram, atomoxetine) do NOT inhibit mast cells [242]
- Mechanism: Modulates intracellular calcium in mast cells

Low-histamine diet:

- Avoid aged/fermented foods, alcohol, cured meats, leftovers >24 hours
- 2-week strict trial, then gradual reintroduction

★ Achievement 5: Evidence for H1+H2 Combination Therapy in Post-Viral Fatigue

While a double-blind RCT demonstrated that H1 antihistamine monotherapy (terfenadine) provides no benefit in CFS [237], emerging evidence from Long COVID case reports [238] suggests that **H1+H2 combination therapy** may be effective for the subset of ME/CFS patients with mast cell activation features. The discontinuation-rechallenge response (symptom worsening upon stopping, improvement upon restarting) provides compelling evidence for treatment effect. Superior H1 agents with additional PAF antagonism and mast cell stabilization properties (rupatadine) may offer advantages over standard antihistamines [239, 240]. ME/CFS patients with documented allergies, orthostatic intolerance, or MCAS features warrant empirical trial of combination antihistamine therapy.

Observation 43 (Patient-Reported MCAS Treatment Benefits). Patient communities consistently report that a subset of ME/CFS and Long COVID patients experience meaningful symptom improvement with MCAS-directed therapies, even absent formal MCAS diagnosis.

Commonly reported benefits:

- Reduced “brain fog”
- Fewer panic-like episodes
- Decreased flushing
- Improved gastrointestinal symptoms

Typical empirical approach: H1+H2 antihistamine combination (preferably rupatadine + famotidine) with optional quercetin and low-histamine diet for 2–4 weeks. Discontinuation testing confirms treatment effect.

The low risk profile and potential for significant benefit in the MCAS-overlap subgroup justify consideration of empirical trials in patients with compatible symptom patterns (flushing, urticaria, food reactions, autonomic episodes, documented allergies).

Speculation 9 (MCAS Energy Amplifier Hypothesis). **Certainty: 0.40.** Mechanistically plausible given documented AMPK-mast cell regulation and energy-dependent degranulation [243], and supported by observational data showing MCAS-ME/CFS comorbidity [235, 229] and ketotifen PEM reduction [244]. However, a direct causal link from mast cell activation to ME/CFS energy deficit worsening has not been demonstrated in controlled studies, and the reported ketotifen benefits lack randomized controlled confirmation.

Core proposition: Mast cell activation episodes create acute energy demands that worsen pre-existing CNS energy deficits in ME/CFS, establishing a positive feedback loop that amplifies PEM severity and frequency.

Energy Cost of Mast Cell Activation. Mast cell degranulation and the subsequent histamine cascade impose substantial metabolic demands [243]:

- **Degranulation energetics:** IgE-mediated mast cell activation utilizes ATP and rapidly induces glycolysis. While oxidative phosphorylation generates 32 ATP per glucose molecule, glycolysis produces only 2 ATP per glucose but can metabolize many glucose molecules simultaneously, creating “short bursts of large amounts of ATP” to support the degranulation process.
- **AMPK dysregulation:** AMP-activated protein kinase (AMPK), the cell’s central energy sensor, normally provides negative feedback to suppress mast cell activation when energy is low. ERK1/2 signaling during FcεRI activation can abolish this AMPK-dependent brake [243].
- **Systemic energy diversion:** Managing the histamine cascade (vasodilation, inflammatory response, immune mediator production) requires substantial metabolic resources systemically, not just within mast cells themselves.
- **CNS histamine burden:** Mast cells serve as the predominant histamine source within the brain (>50% of total CNS histamine). Excessive histamine release from brain mast cells or peripheral histamine crossing a disrupted blood-brain barrier can impair CNS energy homeostasis [245].

MCAS Prevalence and Clinical Overlap. The substantial comorbidity between MCAS and ME/CFS suggests shared or mutually reinforcing pathophysiology:

- **Prevalence:** Estimates range from 25.3% [229] to 30–50% [235] of ME/CFS patients meeting MCAS criteria or exhibiting clinically relevant mast cell activation.
- **Progressive involvement:** Mast cell activation prevalence **increases over the disease course** [229], suggesting that chronic energy deficit may progressively impair mast cell regulation.
- **Orthostatic overlap:** Patients with both MCAS and orthostatic intolerance (particularly POTS) show significantly higher treatment response rates to mast cell-directed therapy compared to OI alone ($p < 0.0001$) [235, 229].
- **Cellular abnormalities:** Objective evidence of altered mast cell phenotypes in ME/CFS includes increased naïve mast cells and elevated activation markers on differentiated mast cells in severe cases [236].

Proposed Positive Feedback Mechanism. The hypothesis posits a vicious cycle:

1. **Baseline energy deficit:** ME/CFS patients operate with limited CNS energy reserves (see Chapter 6 and the selective dysfunction hypothesis, Section 14.24).
2. **Impaired immune regulation:** Energy scarcity compromises AMPK-mediated negative regulation of mast cells and other energy-intensive immune processes.
3. **MCAS flare:** Triggers (allergens, stress, exertion, infections) provoke mast cell degranulation in the context of reduced inhibitory control.
4. **Energy consumption surge:** Degranulation and histamine cascade management acutely “steal” energy from an already depleted budget, particularly impacting the CNS given brain mast cells’ role as the primary CNS histamine source.
5. **Worsened energy deficit:** The acute energy drain deepens the baseline deficit, further impairing mast cell regulation and increasing vulnerability to subsequent triggers.
6. **PEM amplification:** Energy depletion precipitates or worsens post-exertional malaise episodes, with the severity and duration potentially proportional to the degree of mast cell involvement.

This model predicts that MCAS episodes function as **energy amplifiers**: each activation event not only consumes energy directly but also lowers the threshold for future activations and crashes.

Testable Predictions.

1. **Temporal relationship:** MCAS flares (flushing, urticaria, GI symptoms, autonomic instability) should frequently precede or coincide with PEM crash onset.
2. **Biomarker correlations:** Elevated tryptase and histamine levels during symptomatic periods should correlate with reduced markers of energy availability (e.g., decreased ATP/ADP ratios, elevated lactate).

3. **Severity correlation:** MCAS symptom severity should predict PEM severity and frequency, independent of exertional triggers.
4. **Threshold effects:** Prophylactic mast cell stabilization should raise the PEM threshold—patients should tolerate greater activity levels without crashing when mast cells are pharmacologically stabilized.
5. **Treatment response pattern:** Mast cell stabilizers should reduce crash frequency and severity beyond their direct anti-allergic effects, even in patients without formal MCAS diagnosis.

Supporting Evidence. *Clinical evidence:* Retrospective analyses of ketotifen (a mast cell stabilizer with H1 and leukotriene antagonism) in ME/CFS and Long COVID patients showed substantial PEM reduction in 77–95% of patients who continued treatment [244]. This suggests mast cell stabilization specifically targets PEM mechanisms, not merely allergic symptoms. ME/CFS patients with documented MCAS and OI features respond significantly better to mast cell-directed treatment ($p < 0.0001$) [235, 229].

Mechanistic plausibility: AMPK links cellular energy status directly to mast cell regulation [243]. Energy deficits reduce AMPK activity, which disinhibits mast cells—a clear mechanistic path from energy scarcity to immune hyperreactivity. Histamine’s documented effects on CNS energy metabolism, behavioral state, and biological rhythms [245] provide a direct pathway for mast cell activity to worsen cognitive and energy symptoms. Mast cell-derived mediators cause vascular dysfunction [235], which can worsen tissue perfusion and oxygen delivery—further constraining energy availability.

Treatment Implications.

- **Prophylactic stabilization:** Mast cell stabilizers (ketotifen, cromones, quercetin) and H1+H2 antihistamine combinations may serve as **energy-protective** interventions, not merely anti-allergic treatments.
- **Empirical trials justified:** Given the low risk profile and the potential for substantial PEM reduction, empirical trials are justified even in ME/CFS patients without formal MCAS diagnosis, particularly those with compatible symptom patterns.
- **Combination approaches:** Addressing both energy metabolism (e.g., mitochondrial support, pacing) and mast cell stabilization simultaneously may yield synergistic benefits by breaking the positive feedback loop at multiple points.

Limitations and Uncertainties.

- **Observational evidence only:** Ketotifen studies lack randomized controls. The reported PEM improvements could reflect placebo effects, natural disease fluctuation, or regression to the mean.
- **Heterogeneous MCAS criteria:** Diagnostic criteria for MCAS vary across studies and clinicians, complicating prevalence estimates and subgroup identification.

- **Mechanistic gaps:** The hypothesis extrapolates from cellular energy costs of mast cell activation to systemic and CNS energy deficits. Direct measurement of energy availability before, during, and after MCAS flares is lacking.
- **Alternative explanations:** MCAS and ME/CFS may share common upstream causes (e.g., viral triggers, autoimmunity, genetic predispositions) without direct causal interaction.

⌘ Limitation 7: MCAS–ME/CFS Overlap: Comorbidity vs. Shared Pathophysiology

The substantial symptom overlap and estimated 25–50% co-occurrence rate between MCAS and ME/CFS [235, 229] does not establish whether these conditions share pathogenic mechanisms or are independent conditions that co-occur due to shared risk factors. Key epistemic boundaries:

- Prevalence estimates vary widely (25–53%) depending on diagnostic criteria applied, and MCAS diagnostic criteria themselves remain contested.
- Treatment response data (e.g., ketotifen PEM reduction [244]) are from retrospective, uncontrolled analyses; placebo response, natural fluctuation, and regression to the mean are not excluded.
- The proposed energy amplifier feedback loop (mast cell activation → energy depletion → reduced mast cell regulation) is mechanistically plausible but has not been demonstrated in vivo. No study has simultaneously measured mast cell activation markers and energy availability during ME/CFS symptom episodes.

7.7.2 Allergic Responses

Food Sensitivities

Many ME/CFS patients report food intolerances [235]:

- May be IgE-mediated (true allergy) or non-IgE-mediated
- Common triggers: gluten, dairy, histamine-rich foods
- Mechanism may involve mast cell activation or gut barrier dysfunction [227]
- Elimination diets help some patients

Environmental Allergies

Increased sensitivity to environmental factors is reported in ME/CFS:

- Pollen, dust mites, mold
- Chemical sensitivities (fragrances, cleaning products)
- Medication sensitivities
- May reflect mast cell hyperreactivity or neurogenic inflammation [235]

Shared Immune Pathways

Links between allergy and ME/CFS [235, 229]:

- Th2 skewing in some patients
- Elevated IgE in subsets
- Mast cell dysfunction [236]
- Neurogenic inflammation (sensory nerve-mast cell interactions)

7.8 Summary: Integrated Model of Immune Dysfunction

The immune abnormalities in ME/CFS form a coherent, if complex, picture [5]:

1. **Triggering event:** Infection or other immune challenge initiates the process
2. **Innate immune dysfunction:** NK cells and other innate effectors fail to clear the pathogen or control reactivation
3. **Chronic antigenic stimulation:** Persistent infection or autoimmunity drives ongoing B cell activation, producing the characteristic naïve B cell expansion and switched memory B cell depletion documented by the NIH study
4. **Autoantibody development:** Aberrant B cell responses generate autoantibodies targeting receptors and other self-antigens
5. **T cell exhaustion:** Chronic stimulation exhausts T cell responses
6. **Cytokine dysregulation:** Ongoing inflammation produces symptom-causing cytokines
7. **Sex-specific patterns:** Men and women show different immune abnormalities, suggesting distinct pathophysiological pathways
8. **Neuroinflammation:** Peripheral immune signals affect brain function, contributing to fatigue and cognitive symptoms
9. **Mast cell involvement:** Mast cell activation may amplify symptoms in susceptible individuals

This sequence represents one plausible ordering of events; many steps may occur in parallel, and the sequence may vary between patients or subgroups. For example, autoantibody development (step 4) could precede, follow, or coincide with T cell exhaustion (step 5), and sex-specific immune patterns (step 7) likely influence all stages rather than emerging at a discrete point.

This model provides multiple potential therapeutic targets: antiviral agents for persistent infection, immunomodulators for autoimmunity, mast cell stabilizers for those with MCAS, and anti-inflammatory approaches for cytokine-mediated symptoms. The recognition of sex-specific immune patterns may eventually enable personalized treatment selection.

7.9 Emerging Research Directions in Immune Dysregulation

The recent cytokine biomarker findings, combined with advances in understanding immune exhaustion, autoantibodies, and sex-specific patterns, suggest several promising research directions. These are organized by potential impact for severe ME/CFS cases and feasibility of rapid translation to clinical benefit.

7.9.1 Tier 1: Immediate Translation Potential (Existing Drugs, Severe Case Priority)

These interventions use already-approved medications or simple protocols and could benefit severe cases within months of trial initiation.

Hormonal Immune Modulation in Post-Menopausal Women

Rationale The Che et al. [189] finding that women over 45 with diminished estradiol show exaggerated IL-6 responses provides a mechanistic basis for estrogen supplementation. Estrogen receptors are present on immune cells (B cells, monocytes, T cells), and estrogen reduces production of IL-6, TNF- α , and IL-1 β .

Proposed Study Design

- **Population:** Post-menopausal women with severe ME/CFS and documented low estradiol (<30 pg/mL)
- **Intervention:** Transdermal estradiol patch (0.05–0.1 mg/day) with appropriate progesterone for women with intact uterus
- **Duration:** 6-month open-label pilot (n=20), followed by 12-month RCT (n=100) if successful
- **Primary outcomes:** IL-6 levels, SF-36 Physical Function, PEM severity
- **Biomarker stratification:** Measure baseline IL-6 response to microbial stimulation; predict responders as those with highest baseline IL-6

Expected Benefit for Severe Cases Post-menopausal women with severe ME/CFS represent approximately 15–20% of the severe patient population. If estrogen normalizes immune hyperactivation, this subgroup could see substantial symptom improvement within 3–6 months. The intervention is low-risk, FDA-approved, and immediately available.

Timeline Pilot study results: 9–12 months; RCT results: 24–30 months.

Low-Dose IL-2 Therapy for Regulatory T Cell Restoration

Rationale ME/CFS patients show reduced Treg numbers and function, contributing to loss of immune tolerance and potential autoimmunity. Low-dose IL-2 therapy (1–2 million IU subcutaneous, 2–3 times weekly) selectively expands Tregs without activating effector T cells, and has shown efficacy in systemic lupus erythematosus, type 1 diabetes, and graft-versus-host disease.

Convergent Evidence for IL-2 Dysregulation

- Elevated IL-2 in extracellular vesicles [214]
- IL-2 signaling pathways identified in epigenetic biomarker panel [215]
- Reduced Treg function documented in multiple ME/CFS studies
- Possible “IL-2 resistance” mechanism (cells produce IL-2 but cannot respond properly)

Proposed Study Design

- **Population:** Severe ME/CFS patients with documented Treg deficiency ($CD4^+CD25^+FoxP3^+ < 5\%$ of $CD4^+$ T cells)
- **Intervention:** Subcutaneous IL-2 (1 million IU) three times weekly for 12 weeks
- **Mechanistic assessments:** Treg expansion (flow cytometry), IL-2 receptor expression (CD25/CD122/CD132), downstream signaling (pSTAT5)
- **Primary outcomes:** Treg percentage, symptom severity, autoantibody titers
- **Safety monitoring:** Flu-like symptoms common but typically mild; monitor for excessive immune activation

Expected Benefit for Severe Cases If Treg restoration reduces autoimmune symptoms and normalizes immune balance, severe patients with prominent autoimmune features (elevated GPCR autoantibodies, ANA positivity) may experience meaningful improvement. Response likely within 6–12 weeks if mechanism is valid.

Alternative Hypothesis: IL-2 Receptor Dysfunction If the problem is IL-2 *resistance* (down-regulated receptors, impaired signaling), low-dose IL-2 may fail. This would be informative: functional assays measuring T-cell proliferation in response to exogenous IL-2 should be conducted first to identify likely responders.

Timeline Pilot study (mechanistic + safety): 6–9 months; efficacy RCT: 18–24 months.

Phase-Targeted Anti-Cytokine Therapy (Early Disease Window)

Rationale Hornig et al. [212] demonstrated that cytokine elevations occur primarily in early disease (<3 years), with normalization in late disease. This suggests a **time-sensitive therapeutic window**: anti-inflammatory therapies may only benefit patients in the hyperactive phase before immune exhaustion sets in.

The “Immune Exhaustion Timeline” Hypothesis

- **Years 0–3 (Hyperactive Phase):** Elevated cytokines, active inflammation, NK cells attempting (but failing) to clear infection. Therapeutic target: suppress inflammation to prevent exhaustion.
- **Years 3+ (Exhaustion Phase):** Normalized cytokines (false “recovery”), epigenetic T-cell reprogramming, memory B-cell depletion. Therapeutic target: immune “reboot” strategies (B-cell depletion, plasma cell depletion) rather than suppression.

Proposed Study Design

- **Population:** Severe ME/CFS patients with illness duration <3 years and documented cytokine elevation (IL-6 >5 pg/mL, or elevated IL-1 β , TNF- α , or others from severity-correlated panel)
- **Intervention:** Tocilizumab (IL-6 receptor blocker, 162 mg subcutaneous monthly) or etanercept (TNF- α blocker, 50 mg subcutaneous weekly)
- **Duration:** 6-month treatment, with 6-month follow-up to assess durability
- **Primary outcomes:** Prevent progression to exhaustion phase (measured by T-cell exhaustion markers PD-1, Tim-3), symptom improvement, cytokine normalization
- **Critical control:** Late-stage patients (>3 years) treated with same agents to test whether therapeutic window is truly time-limited

Expected Benefit for Severe Cases If early aggressive anti-cytokine therapy prevents the transition to immune exhaustion, it could fundamentally alter disease trajectory. Severe early-stage patients represent approximately 10–15% of all severe cases. Benefit would be disease-modifying rather than purely symptomatic.

Risk Consideration Anti-cytokine biologics increase infection risk. In patients with suspected persistent viral infection (EBV, HHV-6), immunosuppression could worsen viral reactivation. Concurrent antiviral therapy (valacyclovir, valganciclovir) should be considered.

Timeline Pilot study: 12–15 months; RCT with long-term follow-up: 36–48 months.

Extracellular Vesicle Depletion via Enhanced Plasmapheresis

Rationale Giloteaux et al. [214] identified elevated IL-2 and other cytokines specifically in *extracellular vesicles* (EVs), not bulk plasma. EVs are membrane-bound nanoparticles (30–1000 nm) that cells release to communicate with distant cells. They cross the blood-brain barrier, deliver cargo (proteins, RNA, microRNAs) to recipient cells, and can reprogram cellular function.

The “Pathogenic EV” Hypothesis ME/CFS immune cells release EVs containing:

- Pro-inflammatory cytokines (IL-2, TNF- α , CSF2)
- MicroRNAs that reprogram recipient cells toward exhaustion or dysfunction
- Damage-associated molecular patterns (DAMPs) triggering sterile inflammation

These pathogenic EVs may:

- Enter the brain and activate microglia (explaining neuroinflammation and cognitive symptoms)
- Reprogram muscle cells (explaining PEM and mitochondrial dysfunction)
- Amplify systemic inflammation in a self-sustaining loop

Why EV Depletion May Explain Immunoabsorption Successes Stein et al. [118] reported that 70% of post-COVID ME/CFS patients improved with immunoabsorption, with benefits sustained to 6 months. While attributed to autoantibody removal, standard immunoabsorption also removes extracellular vesicles. EV depletion may be the actual therapeutic mechanism.

Proposed Study Design

- **Population:** Severe ME/CFS patients, particularly those with cognitive dysfunction (suggesting CNS involvement via EV trafficking)
- **Intervention:** Immunoabsorption (5 sessions over 10 days using Immunosorba columns or equivalent)
- **Mechanistic assessments:**
 - EV cytokine content pre/post treatment (IL-2, TNF- α , CSF2)
 - EV concentration and size distribution (nanoparticle tracking analysis)
 - EV microRNA cargo (sequencing to identify pathogenic microRNAs)
 - Plasma cytokines (to compare bulk vs. EV-specific changes)
- **Primary outcomes:** Cognitive function (Montreal Cognitive Assessment), fatigue (Chalder Fatigue Scale), SF-36
- **Durability assessment:** Monthly follow-up for 6 months to determine if EVs reaccumulate

Expected Benefit for Severe Cases Severe ME/CFS with prominent cognitive dysfunction may benefit most. If pathogenic EVs drive neuroinflammation, removal could produce rapid improvement (within days to weeks). Approximately 80–90% of severe cases have significant cognitive impairment.

Advanced Approach: EV-Specific Filtration Standard immunoadsorption removes IgG non-selectively. Newer technologies (ExoLution, Plasmax) can selectively filter EVs while preserving antibodies. If EVs are the true therapeutic target, EV-specific filtration could be more effective with fewer side effects.

Timeline Pilot study with mechanistic assessments: 12–18 months; RCT: 24–30 months; EV-specific filtration development: 36–48 months.

7.9.2 Tier 2: Near-Term Clinical Trials (Moderate Complexity, High Impact)

These interventions require more complex trial designs or involve experimental therapies but could still reach severe patients within 2–4 years.

TRPM3 Modulation for Calcium-Cytokine Axis Restoration

Rationale TRPM3 ion channel dysfunction impairs calcium signaling in ME/CFS immune cells [187]. Calcium is essential for:

- NK cell and T-cell degranulation
- Cytokine gene transcription (calcium activates NFAT transcription factors)
- Extracellular vesicle release (calcium-dependent membrane fusion)

Connecting TRPM3 to Cytokine Dysregulation The TRPM3-cytokine connection may explain multiple findings:

- Impaired NK cytotoxicity (cannot degranulate without calcium influx)
- Dysregulated cytokine production (abnormal calcium signaling → abnormal transcription)
- Elevated EV cytokines (altered calcium-dependent EV formation/release)

Therapeutic Approaches

1. **TRPM3 agonists:** Drugs that directly activate TRPM3 to restore calcium entry
 - Pregnenolone sulfate (endogenous TRPM3 agonist, available as supplement)
 - CIM0216 (experimental selective TRPM3 agonist)

2. **Calcium ionophores:** Compounds that bypass TRPM3 by directly shuttling calcium across membranes
 - Ionomycin (research tool, too toxic for clinical use)
 - A23187 (research tool)
 - Need development of safer clinical-grade ionophores
3. **Indirect approaches:** Drugs that enhance residual TRPM3 function
 - PIP2 supplementation (TRPM3 requires PIP2 for activation)
 - Membrane fluidity enhancers

Proposed Study Design

- **Phase 1: Mechanistic validation**
 - Isolate PBMCs from severe ME/CFS patients
 - Measure cytokine production with/without calcium supplementation
 - Test whether TRPM3 agonists (pregnenolone sulfate) restore normal cytokine responses *in vitro*
 - If positive, proceed to clinical trial
- **Phase 2: Clinical pilot**
 - Pregnenolone sulfate oral supplementation (50–100 mg daily for 12 weeks)
 - Primary outcomes: NK cytotoxicity, cytokine levels, symptom improvement
 - Biomarker: TRPM3 function assay (calcium flux in response to agonist)

Expected Benefit for Severe Cases If TRPM3 dysfunction is a core defect, restoration could improve multiple systems simultaneously (immune function, muscle function, autonomic function—all require calcium signaling). Benefit could be substantial and rapid (weeks). All severe cases could potentially benefit regardless of disease duration.

Timeline In vitro validation: 6–12 months; pregnenolone sulfate pilot: 18 months; development of novel TRPM3 agonists: 48–60 months.

Microbiome-Targeted Immune Normalization

Rationale Che et al. [189] used heat-killed *Candida albicans* to demonstrate exaggerated cytokine responses. This fungal stimulation assay suggests that ME/CFS patients' immune systems are "primed" to overreact to microbial antigens. Gut dysbiosis with fungal overgrowth could provide constant low-level antigenic exposure, maintaining immune hyperactivation.

The “Dysbiotic Priming” Hypothesis

- Gut barrier dysfunction (“leaky gut”) permits translocation of fungal/bacterial antigens
- Constant low-level exposure primes immune cells to overreact
- When challenged (infection, stress, exertion), primed immune system produces exaggerated cytokine response
- Explains both baseline immune activation and PEM (exertion disrupts gut barrier further)

Why Sex Differences May Relate to Microbiome Estrogen affects gut microbiome composition. Post-menopausal women have altered gut flora with increased *Candida* colonization. This could explain Che’s finding of amplified IL-6 in women over 45 with low estradiol.

Proposed Multi-Modal Intervention

1. **Antifungal therapy:** Fluconazole 100–200 mg daily for 4 weeks, then intermittent dosing
2. **Gut barrier repair:** L-glutamine (5 g twice daily), zinc carnosine (75 mg twice daily), butyrate supplementation
3. **Microbiome restoration:** Targeted probiotics (*Saccharomyces boulardii*, *Lactobacillus/Bifidobacterium* strains) or fecal microbiota transplantation (FMT) from highly screened donors
4. **Dietary modification:** Low-fermentation diet during acute treatment, then gradual reintroduction

Proposed Study Design

- **Population:** Severe ME/CFS patients with GI symptoms and documented dysbiosis (stool testing showing elevated *Candida*, low bacterial diversity)
- **Design:** 2×2 factorial design testing antifungal + gut repair vs. placebo over 6 months
- **Mechanistic assessments:**
 - Baseline *Candida* stimulation assay (replicate Che protocol)
 - Gut permeability (lactulose/mannitol test, zonulin levels)
 - Microbiome sequencing pre/post treatment
 - Cytokine responses to microbial stimulation pre/post treatment
- **Primary outcomes:** GI symptom improvement, systemic symptom improvement, cytokine normalization

Expected Benefit for Severe Cases Severe ME/CFS patients with prominent GI symptoms (estimated 60–70% of severe cases) may benefit most. If dysbiotic priming is a maintaining factor, addressing it could reduce baseline immune activation and PEM severity. Benefits likely gradual (3–6 months for microbiome reconstitution).

Timeline Pilot study: 12–18 months; RCT: 24–36 months.

Duration-Severity Stratified Trials with Mechanistic Biomarkers

Rationale The logic audit identified that no study has examined duration, severity, and sex simultaneously in a stratified design. Current trials may fail because they combine patients in different disease phases (early hyperactive vs. late exhausted) who require different therapeutic approaches.

The “Two-Hit” Model Requiring Stratification

- **Hit 1 (Initial trigger):** Determines whether patient enters high-cytokine trajectory or not
- **Hit 2 (Ongoing factors):** Determines severity within trajectory (genetics, sex, hormones, comorbidities)
- **Interaction:** Early + severe = highest cytokines, rapid progression to exhaustion; Late + severe = severity driven by non-cytokine mechanisms

Proposed Master Protocol Design

- **Universal screening:** All participants receive comprehensive immune profiling
 - Cytokine panel (including IL-2, IL-6, TNF- α , CCL11, CXCL9)
 - T-cell exhaustion markers (PD-1, Tim-3, LAG-3)
 - B-cell subsets (naïve, memory, plasmablasts)
 - Autoantibody titers (GPCR antibodies)
 - EV cytokine content
 - TRPM3 function
- **Stratification:** Assign to treatment arm based on biomarker profile
 - **Arm A (Early hyperactive):** Duration <3 years, elevated cytokines → anti-cytokine therapy
 - **Arm B (Late exhausted):** Duration >3 years, normal cytokines, high PD-1 → B-cell depletion (daratumumab)
 - **Arm C (Female hormonal):** Post-menopausal with low estradiol, high IL-6 → estrogen supplementation
 - **Arm D (TRPM3 dysfunction):** Impaired calcium signaling → TRPM3 agonist
 - **Arm E (EV-dominant):** Elevated EV cytokines → immunoadsorption
- **Crossover:** Non-responders at 6 months cross to alternative arm based on response patterns

Expected Benefit for Severe Cases This precision-medicine approach could achieve higher response rates (50–60%) compared to unstratified trials (typically 20–30%). All severe patients would be profiled and matched to optimal therapy. Trial would also validate the duration-severity-sex model and identify which biomarkers predict treatment response.

Timeline Protocol development and regulatory approval: 12–18 months; enrollment and treatment: 36 months; analysis and publication: 48 months.

7.9.3 Tier 3: Long-Term Mechanistic Research (Foundational Understanding)

These studies address fundamental questions about ME/CFS immunopathology and will guide future therapeutic development but require 5–10 years to complete.

Longitudinal Immune Evolution Cohort (Onset to Exhaustion)

Rationale The duration-dependent findings (Hornig, Montoya) are cross-sectional snapshots. A prospective longitudinal cohort following patients from disease onset through the first 5 years would definitively establish:

- Whether individual patients transition from high-cytokine to exhaustion phase
- Exact timing and predictors of transition
- Whether early intervention prevents exhaustion
- Which patients never enter high-cytokine phase (and why)

Proposed Study Design

- **Enrollment:** Patients within 6 months of ME/CFS onset (infectious mononucleosis, COVID-19, or other identified triggers)
- **Target enrollment:** n=500 to account for spontaneous recovery (approximately 15–20%)
- **Assessments:** Quarterly for first 2 years, semi-annually thereafter
 - Comprehensive cytokine panel (plasma and EV fractions)
 - T-cell exhaustion markers and epigenetic profiling
 - B-cell subsets and autoantibody titers
 - NK cell function
 - TRPM3 function
 - Microbiome (stool samples)
 - Symptom severity, functional status
- **Substudies:**
 - Randomize subset to early anti-cytokine therapy vs. observation
 - Compare natural history vs. intervention outcomes

Expected Insights

- Define ME/CFS “stages” with precision
- Identify biomarkers that predict progression vs. recovery
- Establish optimal treatment windows
- Determine whether preventing exhaustion changes long-term outcomes

Impact for Severe Cases Findings would guide future treatment timing for all newly diagnosed patients, potentially preventing progression to severe disease. Results would take 5–7 years but could transform clinical approach.

Timeline Enrollment: 24–36 months; follow-up: 60 months; analysis: 72–84 months.

IL-2 Resistance Functional Studies

Research Questions

- Do ME/CFS T cells proliferate normally in response to exogenous IL-2?
- Are IL-2 receptors (CD25/CD122/CD132) expressed normally on T cells and NK cells?
- Is downstream signaling (JAK1/JAK3/STAT5 phosphorylation) intact?
- Are elevated EV-IL-2 levels functionally active or sequestered/inactive?
- Can pharmacologic IL-2 overcome the dysfunction?

Proposed Mechanistic Studies

1. In vitro proliferation assays

- Isolate PBMCs from ME/CFS patients and controls
- Stimulate with increasing doses of recombinant IL-2
- Measure proliferation (CFSE dilution), STAT5 phosphorylation, Treg expansion
- If ME/CFS cells respond poorly → IL-2 resistance confirmed
- If ME/CFS cells respond normally → problem is insufficient IL-2 availability despite elevated EV levels

2. Receptor expression and signaling

- Flow cytometry for CD25/CD122/CD132 surface expression
- Phospho-flow for pSTAT5 after IL-2 stimulation
- Western blot for JAK1/JAK3 expression

3. EV-IL-2 functional testing

- Purify EVs from ME/CFS plasma
- Test whether EV-IL-2 can signal to recipient cells
- Compare bioactivity of EV-bound vs. free IL-2

Therapeutic Implications

- If resistance confirmed → need IL-2 receptor agonists with higher potency, or downstream pathway activators
- If insufficient availability → standard low-dose IL-2 therapy should work
- If EV-IL-2 is sequestered → EV depletion is the correct approach

Timeline Mechanistic studies: 12–24 months; therapeutic trials based on findings: 36–48 months.

CCL11 (Eotaxin) Neutralization for Cognitive Dysfunction

Rationale CCL11 (eotaxin-1) correlates with ME/CFS severity [213], decreases during healthier periods, and is known to:

- Impair hippocampal neurogenesis
- Cause cognitive dysfunction in animal models
- Increase with aging (“cognitive aging” biomarker)
- Cross the blood-brain barrier readily

Why CCL11 Is a Promising Target

- Directly toxic to neural progenitor cells
- Specific correlation with cognitive symptoms
- Aging research has developed CCL11-neutralizing antibodies
- Statins reduce CCL11 (may explain why some ME/CFS patients report benefit from statins)

Proposed Research Path

1. **Observational study:** Correlate CCL11 levels with cognitive testing (Montreal Cognitive Assessment, Trail Making Test)
2. **Mechanistic study:** CSF CCL11 levels and correlation with neuroimaging (MRI volumetrics, PET microglial activation)
3. **Intervention pilot:** Atorvastatin 40 mg daily (known to reduce CCL11) in severe ME/CFS with cognitive dysfunction
4. **Advanced therapy:** Anti-CCL11 monoclonal antibody (if statin pilot successful)

Expected Benefit for Severe Cases Severe cognitive dysfunction is often the most disabling symptom. If CCL11 neutralization improves cognition, quality of life could improve substantially even without improving physical fatigue. Approximately 80–90% of severe cases have cognitive impairment.

Timeline Observational + mechanistic studies: 18–24 months; statin pilot: 12–18 months; antibody development and trials: 60–84 months.

7.9.4 Prioritization Summary: Research Directions by Impact and Timeline

Recommended Immediate Actions For maximum impact on severe ME/CFS within 2 years:

1. **Launch in parallel** (can run simultaneously):
 - Hormonal modulation pilot (post-menopausal women, n=20)
 - EV depletion mechanistic study (immunoadsorption with EV analysis, n=15)
 - Low-dose IL-2 open-label pilot (n=15)
2. **Mechanistic validation** (to guide Tier 2 trials):
 - TRPM3 *in vitro* studies (calcium rescue experiments)
 - IL-2 resistance functional assays
 - Microbiome-cytokine correlation studies
3. **Registry development:**
 - Establish prospective registry for newly diagnosed patients (enrollment for longitudinal cohort)
 - Implement universal biomarker profiling to enable stratified trial enrollment

Expected Cumulative Impact If these research directions succeed:

- **Year 1–2:** Hormonal modulation, EV depletion, low-dose IL-2 pilots complete → 3 potential new therapies for distinct subgroups (combined coverage: 40–50% of severe cases)
- **Year 2–4:** TRPM3 modulation, microbiome normalization, stratified trials complete → precision medicine approach validated, additional 30–40% coverage
- **Year 5–7:** Longitudinal cohort results guide early intervention → prevent progression to severe disease in newly diagnosed patients
- **Year 7–10:** Advanced therapies (CCL11 antibodies, novel TRPM3 agonists) → address remaining treatment-refractory cases

Combined, these approaches could provide therapeutic options for 70–80% of severe ME/CFS patients within 5 years, with prevention strategies for newly diagnosed patients following within 7–10 years.

⚠ **Limitation 8: Research Direction Projections: Optimistic Extrapolation**

The timelines, expected response rates, and cumulative impact projections in this section are best-case estimates based on historical precedent from other fields, not ME/CFS-specific data. Key caveats:

- Historical ME/CFS clinical trials have a high failure rate (rituximab Phase III, rintatolimod FDA rejection), and promising pilot results frequently do not replicate in larger RCTs.
- The projected “70–80% coverage within 5 years” assumes all Tier 1 and Tier 2 interventions succeed — an improbable scenario given typical attrition rates in translational research.
- Stratified trial designs assume biomarker-defined subgroups correspond to mechanistically distinct treatment-responsive populations — this has not been validated in ME/CFS.
- Regulatory, funding, and recruitment barriers specific to ME/CFS (stigma, limited research infrastructure, patient inability to attend trial sites) are not reflected in the timeline estimates.

7.9.5 Post-Infectious Pathogenesis

Post-infectious ME/CFS arises when acute infections trigger persistent pathophysiological changes that outlast the initial pathogen. Key mechanisms include:

- **Viral persistence:** Some pathogens (EBV, enteroviruses) may establish low-level persistent infections
- **Immune dysregulation:** Acute infection disrupts immune homeostasis, leading to chronic activation
- **Metabolic reprogramming:** Pathogen-induced metabolic changes persist after clearance
- **Neuroinflammatory priming:** Acute neuroinflammation creates long-lasting glial cell activation
- **Epigenetic modifications:** Infection-induced epigenetic changes alter gene expression patterns

This section explains why ME/CFS persists after the inciting infection has resolved, creating a self-sustaining pathological state independent of the original trigger.

7.9.6 Trained Immunity and Epigenetic Lock-in

Trained immunity refers to the long-term functional reprogramming of innate immune cells following exposure to microbial products. In ME/CFS:

- **Monocyte training:** Pathogen-associated molecular patterns (PAMPs) induce epigenetic modifications
- **Histone modifications:** H3K4me3 marks at inflammatory gene promoters enhance responsiveness
- **Metabolic rewiring:** Shift from oxidative phosphorylation to glycolysis in trained cells
- **Persistent activation:** Trained cells maintain hyperresponsive state for months to years

- **Therapeutic implications:** Epigenetic modifiers may reverse trained immunity phenotypes

This epigenetic lock-in mechanism explains the chronic nature of immune dysfunction in ME/CFS and represents a potential therapeutic target.

⚠ **Limitation 9: Trained Immunity in ME/CFS: No Disease-Specific Data**

Trained immunity (epigenetic reprogramming of innate immune cells) is well-characterised in general immunology and has been demonstrated in atherosclerosis, sepsis, and BCG vaccination responses. Its application to ME/CFS is entirely extrapolated — no study has measured H3K4me3 marks at inflammatory gene promoters, metabolic rewiring from OXPHOS to glycolysis in monocytes, or trained immunity reversal in ME/CFS patients. Whether ME/CFS monocytes exhibit a trained phenotype, and whether this is cause or consequence of chronic immune activation, is unknown.

7.9.7 Zonulin-Mediated Gut Permeability

Zonulin is a protein that regulates intestinal tight junctions. In ME/CFS:

- **Increased zonulin:** Elevated levels correlate with gut permeability and symptom severity
- **LPS translocation:** Bacterial lipopolysaccharide enters circulation, triggering systemic inflammation
- **Mast cell activation:** LPS stimulates mast cells, exacerbating neuroinflammatory responses
- **Immune activation:** Chronic endotoxemia maintains systemic inflammatory state
- **Therapeutic target:** Zonulin inhibitors (larazotide acetate) may reduce gut permeability

Gut barrier dysfunction represents a critical interface between peripheral inflammation and systemic ME/CFS pathophysiology.

Table 7.1: Prioritized research directions for severe ME/CFS

Research Direction	Severe Case Benefit	Timeline to Results	Feasibility	Priority Rank
TIER 1: Immediate Translation (Existing Drugs)				
Hormonal modulation (post-menopausal women)	High (15–20% of severe)	12–24 mo	Very High	1
Low-dose IL-2 (Treg restoration)	High (all with autoimmunity)	18–24 mo	High	2
EV depletion (immunoadsorption)	Very High (80–90% with cognitive)	12–18 mo	High	3
Phase-targeted anti-cytokine (early)	Very High (disease-modifying)	24–36 mo	Moderate	4
TIER 2: Near-Term Trials (Moderate Complexity)				
TRPM3 modulation	Very High (all severe cases)	36–48 mo	Moderate	5
Microbiome normalization	High (60–70% with GI)	24–36 mo	High	6
Stratified biomarker trials	Very High (precision medicine)	48 mo	Moderate	7
TIER 3: Long-Term Research (Foundational)				
Longitudinal cohort (onset to exhaustion)	High (prevents severe cases)	72–84 mo	Low	8
IL-2 resistance mechanistic studies	Moderate (guides therapy)	36–48 mo	High	9
CCL11 neutralization	High (cognitive-dominant)	60–84 mo	Low	10

8 Neurological and Neurocognitive Dysfunction

Neurological abnormalities represent one of the most consistently documented features of ME/CFS and provide critical insight into the pathophysiology of this complex disorder. The landmark NIH deep phenotyping study by Walitt et al. (2024) provided unprecedented detail on central nervous system dysfunction, identifying specific brain regions, neurotransmitter abnormalities, and mechanisms underlying the characteristic fatigue and cognitive impairment of ME/CFS [5].

8.1 Central Nervous System Abnormalities

8.1.1 Brain Structure and Function

Structural Neuroimaging Findings

Multiple neuroimaging studies have documented structural brain abnormalities in ME/CFS patients, though findings have varied across studies due to differences in patient populations, imaging protocols, and analytical methods [246].

White Matter Abnormalities Several studies have reported increased white matter hyperintensities (WMH) in ME/CFS patients compared to healthy controls [247, 248]. These hyperintensities, visible on T2-weighted and FLAIR MRI sequences, may indicate demyelination, axonal loss, or microvascular damage. The distribution of WMH in ME/CFS patients tends to involve periventricular white matter, subcortical regions, and frontal and temporal lobes. Zeineh et al. [248] identified increased fractional anisotropy in the right arcuate fasciculus, which correlated with disease severity ($r=0.649$, $p=0.0015$), providing anatomical substrate for the cognitive dysfunction observed in ME/CFS.

The clinical significance of these findings remains debated, as similar changes occur with normal aging and various medical conditions. However, the presence of WMH in younger ME/CFS patients suggests pathological processes beyond typical age-related changes.

Gray Matter Volume Changes Voxel-based morphometry (VBM) studies have identified regional brain abnormalities in ME/CFS patients, though findings vary across cohorts [249, 250, 251]. Documented changes include gray matter differences in the parahippocampal gyrus, occipital regions, amygdala, and insula, alongside white matter volume reductions in the brainstem and temporal regions. No single pattern has been consistently replicated across studies, reflecting the clinical heterogeneity of ME/CFS.

Despite this heterogeneity, the presence of structural brain differences in regions involved in memory, interoception, and autonomic regulation supports a neuroanatomical basis for cognitive and autonomic dysfunction in ME/CFS [251].

Functional Neuroimaging: The NIH Deep Phenotyping Study

The 2024 NIH study by Walitt et al. employed functional MRI during motor tasks to identify specific brain regions with abnormal activation patterns in PI-ME/CFS patients [5]. This study, involving 17 PI-ME/CFS patients and 21 matched healthy controls, provided the most rigorous functional neuroimaging data to date.

Temporal-Parietal Junction Dysfunction

★ Achievement 1: Temporal-Parietal Junction Dysfunction and Effort Miscalculation

Walitt et al. [5] identified abnormally reduced activity in the temporal-parietal junction (TPJ) during effort-based decision-making tasks in PI-ME/CFS patients. The TPJ is a heteromodal association cortex that integrates information from multiple sensory modalities and plays essential roles in agency and intention attribution, effort allocation decisions, attentional reorienting, social cognition, and bodily self-consciousness. This dysfunction provides a neuroanatomical substrate for the characteristic mismatch between perceived capability and actual performance that defines ME/CFS, suggesting the brain genuinely perceives effort requirements inaccurately rather than exhibiting malingering or simple deconditioning.

△ Warning 1: Replication Status

Single study (Walitt 2024, n=17 PI-ME/CFS vs. 21 controls). This finding derives from the NIH deep phenotyping study—the first to examine TPJ function in ME/CFS. The small sample size limits generalizability. Independent replication in a larger cohort and across ME/CFS subtypes is required before TPJ dysfunction can be considered an established feature.

Motor Cortex Hyperactivity Paradoxically, while the TPJ showed reduced activation, the motor cortex demonstrated sustained hyperactivity during fatiguing grip tasks in ME/CFS patients [5]. The motor cortex remained abnormally active despite declining grip force output, yet electromyography showed no evidence of peripheral muscle fatigue. This dissociation between central motor drive and peripheral performance reveals inefficient neural recruitment patterns requiring excessive cortical activation for submaximal force production.

This pattern indicates that fatigue in ME/CFS originates centrally rather than peripherally [5]. The motor cortex continues to “try harder” even as actual force production declines, suggesting a breakdown in the feedback mechanisms that normally calibrate effort to output.

Effort Preference Alteration: A New Paradigm Perhaps the most conceptually important finding from the NIH study was the identification of altered effort preference as a defining feature of PI-ME/CFS, distinct from physical fatigue (muscle exhaustion) or central fatigue (reduced motor cortex output). Walitt et al. proposed that:

“Fatigue may arise from a mismatch between what someone thinks they can achieve and what their bodies perform.”

This reconceptualization has profound implications for understanding ME/CFS. First, the brain genuinely perceives effort requirements inaccurately, leading to appropriate behavioral responses to faulty signals; this rules out malingering or simple deconditioning. Second, the TPJ normally synthesizes multiple information streams—interoceptive, proprioceptive, and motivational—to generate effort estimates, but this integration fails in ME/CFS. Third, the brain may be responding to genuine danger signals such as inflammation or metabolic dysfunction while miscalibrating the protective response. Finally, interventions targeting effort perception and decision-making networks may prove more effective than those addressing peripheral fatigue.

~ Hypothesis 1: Maladaptive Sickness Behavior Program

Certainty: 0.50. ME/CFS symptoms may represent an evolutionarily conserved “sickness behavior” program—normally protective during acute infection—that becomes chronically activated due to persistent immune signaling. The TPJ, which normally integrates inflammatory signals with effort allocation decisions, may misinterpret chronic low-grade inflammation as ongoing acute illness, inappropriately suppressing activity to “conserve resources” for an immune battle that has already concluded (or that persists at subclinical levels). This would explain why the fatigue feels so viscerally “real” and protective to patients: the brain is executing a legitimate survival program, but one triggered by faulty or persistent signals rather than current metabolic necessity.

Risk-Based Decision-Making Impairment During behavioral tasks requiring risk assessment and effort allocation, ME/CFS patients demonstrated reduced selection of “hard” task options even when rewards were equivalent, difficulty sustaining effort on extended tasks, and altered subjective perception of task difficulty [5]. Notably, motivation levels remained normal despite reduced effort output.

These findings indicate that the problem lies not in willingness to exert effort (motivation) but in the neural computation of what constitutes acceptable effort levels [5].

PET Scan Metabolic Findings

Positron emission tomography (PET) studies have revealed regional hypometabolism in ME/CFS patients, indicating reduced glucose utilization and neuronal activity [252]. Commonly affected regions include brainstem nuclei (potentially explaining autonomic dysfunction), basal ganglia (correlating with motor symptoms and fatigue), medial prefrontal cortex

(associated with executive dysfunction), and posterior parietal cortex (linked to attention and spatial processing deficits).

The pattern of hypometabolism overlaps significantly with regions showing structural and functional abnormalities, consistent with a coherent picture of multifocal brain dysfunction.

⚡ **Limitation 1: Neuroimaging Correlation ≠ Causation**

Correlation of imaging findings does not establish that these brain abnormalities cause ME/CFS symptoms, nor which abnormality precedes others. Temporal precedence, reverse causation (systemic illness causing brain changes), and potential confounders (medication effects, deconditioning, sleep disruption) require further investigation through longitudinal and interventional neuroimaging studies.

SPECT Perfusion Abnormalities

Single-photon emission computed tomography (SPECT) studies have documented reduced regional cerebral blood flow (rCBF) in ME/CFS patients [123]. Characteristic findings include global reduction in cortical perfusion (10–15% below controls), focal hypoperfusion in temporal, frontal, and parietal regions, correlation between perfusion deficits and cognitive symptom severity, and exacerbation of perfusion abnormalities following physical or cognitive exertion.

The persistence of perfusion deficits across multiple studies and imaging modalities is consistent with cerebrovascular dysfunction contributing to ME/CFS symptoms.

⚡ **Limitation 2: Central Catecholamine Findings: Single Study, Small Sample**

The CSF catecholamine data (reduced HVA, DHPG, epinephrine) derive entirely from the NIH deep phenotyping study (Walitt 2024, n=17 PI-ME/CFS vs. 21 controls). Key epistemic boundaries:

- No prior or subsequent study has performed lumbar puncture with catecholamine analysis in ME/CFS; the findings are unreplicated.
- The cohort was restricted to post-infectious ME/CFS; whether catecholamine deficiency characterises gradual-onset or non-viral ME/CFS is unknown.
- CSF catecholamine levels are influenced by medications, sleep timing, activity levels, and lumbar puncture technique—confounders that single-study designs cannot fully control.
- The direction of causality is unestablished: low catecholamines may *result from* reduced physical activity, sleep disruption, or chronic illness rather than *causing* ME/CFS symptoms.

8.1.2 Brain as Energy Coordination Bottleneck

The convergence of regional hypometabolism (PET), hypoperfusion (SPECT), neuroinflammation, and catecholamine deficiency raises a fundamental question: is brain dysfunction in

ME/CFS *secondary* to systemic illness, or could it be *primary*—the bottleneck limiting whole-body function?

Speculation 10 (CNS Energy Crisis as Primary Dysfunction). **Certainty: 0.40.** The near-universal presence of cognitive dysfunction, documented brain hypometabolism [252], and neuroinflammation with 45–199% elevation in key regions [73] suggest CNS energy crisis may be the primary pathophysiological event. Failure of the brain to coordinate peripheral demand-responsive processes could explain the selective pattern of dysfunction observed in ME/CFS, where autonomous processes (hair growth, nail growth, baseline cellular metabolism) remain intact while CNS-coordinated responses (exercise capacity, orthostatic tolerance, immune adaptation) are severely impaired.

Evidence for Brain-Centric Model

Several observations support the brain as primary bottleneck:

1. **Universal cognitive involvement:** Brain fog and cognitive dysfunction are present in nearly all ME/CFS patients, regardless of primary symptom presentation or disease severity
2. **Disproportionate brain energy demand:** The brain comprises 2% of body mass but consumes 20–25% of resting energy, making it uniquely vulnerable to energy constraint
3. **Cascading coordination failure:** The brain coordinates peripheral demand responses via autonomic signaling; CNS energy deficit would impair this coordination across multiple organ systems simultaneously
4. **Catecholamine deficiency:** Reduced CSF catecholamines (Section 8.1.3) directly impair the signaling required for demand-response mobilization throughout the body
5. **Neuroinflammation evidence:** Nakatomi et al. [73] documented 45–199% elevation in neuroinflammatory markers across six brain regions (cingulate cortex, hippocampus, amygdala, thalamus, midbrain, pons), with inflammation severity correlating directly with cognitive impairment and pain. While replication of these PET findings remains incomplete (see Section 8.1.5), the magnitude and regional distribution suggest potential CNS-specific pathology warranting further investigation

Autonomic Dysfunction as Downstream Effect

If the brain cannot maintain adequate energy for autonomic coordination, peripheral organs would have energy *available* but lack the *signals* to mobilize it appropriately. This explains several key observations:

- **Pharmacological bypass efficacy:** Midodrine, fludrocortisone, and other autonomic-supporting medications can partially restore function—the peripheral targets respond when appropriately stimulated, suggesting the dysfunction is in *coordination* rather than *peripheral capacity*

- **Demand-response failure pattern:** Baseline function often preserved while challenge responses fail; the CNS cannot orchestrate the coordinated scaling required for physiological stress
- **Preservation of autonomous processes:** Truly local processes (hair follicle cycling, which operates an independent internal Cori cycle) continue unaffected because they don't require CNS coordination

Cerebral Blood Flow: The Central Vulnerability

Van Campen and colleagues have systematically documented cerebral blood flow (CBF) abnormalities during orthostatic stress that support the brain-centric model [172, 253, 254, 4]:

★ Achievement 2: Near-Universal CBF Decline During Orthostatic Challenge

In a series of studies using transcranial Doppler during tilt-table testing, van Campen et al. demonstrated that **91% of ME/CFS patients** (488/534) with normal heart rate and blood pressure responses show abnormal cerebral blood flow and cardiac output reduction during orthostatic challenge [4]. The magnitude of CBF decline is approximately **3.7-fold greater** than healthy controls (26% vs. 7% reduction at end-tilt) [172]. Furthermore, CBF remains reduced even after returning to supine position, with recovery correlating to disease severity rather than hemodynamic parameters [253]. For detailed breakdown by orthostatic phenotype, see Achievement 10.2.4 in Chapter 10.

△ Warning 2: Replication Status: Single Research Group

All CBF findings derive from a sequential publication series by van Campen and colleagues at a single Dutch referral center (4 papers, 2020–2024). While the large cumulative sample (n=534) and consistent methodology strengthen internal validity, no independent research group has replicated these transcranial Doppler findings during tilt-table testing. The appearance of replication across multiple publications reflects one group's expanding cohort, not independent confirmation. Independent replication at other centers with different equipment and patient populations is essential.

Observation 44 (CBF-Symptom Correlation). ME/CFS symptom severity correlates directly with the degree of CBF reduction during tilt testing [254]. Patients with greater CBF decline report worse fatigue, cognitive dysfunction, and orthostatic symptoms. The absence of compensatory cerebral vasodilation despite reduced cardiac output suggests possible endothelial dysfunction contributing to cerebrovascular vulnerability [4].

The brain's high metabolic demand and sensitivity to perfusion deficits may make cerebral blood flow the "canary in the coal mine" for systemic energy coordination dysfunction. Standard vital sign monitoring during orthostatic challenge misses this pathology—normal heart rate and blood pressure do not exclude significant cerebrovascular compromise.

See Chapter 6 Section 6.3 for integration with the selective energy dysfunction hypothesis, and Chapter 10 Section 10.2.4 for detailed cerebrovascular findings.

8.1.3 Neurotransmitter Abnormalities

The structural and functional brain abnormalities described above correlate with specific neurochemical deficits identified in the NIH deep phenotyping study, providing mechanistic links between imaging findings and clinical symptoms.

Catecholamine Pathway Dysregulation: CSF Findings

The NIH deep phenotyping study provided the first direct evidence linking cerebrospinal fluid (CSF) catecholamine abnormalities to ME/CFS symptoms [5]. This represents a major advance in understanding the neurochemical basis of the disease.

Reduced CSF Catecholamines

★ Achievement 3: Central Catecholamine Deficiency in ME/CFS

The 2024 NIH deep phenotyping study provided the first direct evidence from cerebrospinal fluid analysis linking central catecholamine abnormalities to ME/CFS symptoms [5]. Lumbar puncture analysis revealed significantly reduced concentrations of catecholamines and their metabolites in ME/CFS patients compared to healthy controls, including lower levels of homovanillic acid (HVA, the primary dopamine metabolite), reduced DHPG (3,4-dihydroxyphenylglycol, indicating decreased central noradrenergic activity), and decreased epinephrine levels.

The DHPG finding is particularly significant because it is the primary intraneuronal metabolite of norepinephrine produced within noradrenergic neurons. Low CSF DHPG specifically indicates reduced norepinephrine turnover in the central nervous system, pointing to hypofunction of the locus coeruleus and other noradrenergic nuclei.

△ Warning 3: Replication Status

Single study (Walitt 2024). This represents the first-ever direct CSF catecholamine measurement in ME/CFS. No prior or subsequent study has performed lumbar puncture with catecholamine analysis in an ME/CFS cohort. The invasiveness of CSF collection limits feasibility of large-scale replication. Independent confirmation is essential, particularly given the small sample size of the NIH cohort.

Clinical Correlations

Observation 45 (Catecholamine-Symptom Correlations). Walitt et al. [5] established direct correlations between CSF catecholamine levels and clinical measures. Lower catecholamines correlated with reduced grip strength endurance and slower reaction times (motor performance), catecholamine deficits predicted reduced selection of “hard” tasks in decision-making paradigms (effort-related behaviors), memory and executive function scores correlated with dopamine metabolite levels (cognitive impairment), and subjective fatigue ratings inversely correlated with norepinephrine concentrations (fatigue severity).

This establishes, for the first time, a direct biochemical pathway linking specific neurotransmitter abnormalities to the core symptoms of ME/CFS.

Figures 8.1 and 8.2 illustrate the catecholamine synthesis pathway and two major bottlenecks in ME/CFS: (1) tyrosine hydroxylase impairment due to ATP deficit and BH₄ depletion, and (2) dopamine β -hydroxylase impairment due to vitamin C depletion.

Mechanistic Implications Central catecholamine deficiency could explain multiple ME/CFS features [5]. Dopamine and norepinephrine are essential for maintaining arousal, motivation, and sustained attention; deficiency produces profound fatigue without peripheral cause. The prefrontal cortex depends on optimal dopamine levels for working memory and executive function, where both excess and deficiency impair cognition. Since norepinephrine is the primary neurotransmitter of the sympathetic nervous system, central norepinephrine deficiency could produce the autonomic abnormalities characteristic of ME/CFS. Dopamine mediates reward anticipation and motivation, so deficiency could explain the reduced effort allocation observed in behavioral tasks. Finally, physical exertion depletes catecholamines; if baseline levels are already low, even modest activity could produce profound neurotransmitter deficits and symptom exacerbation, explaining post-exertional malaise [5].

Tryptophan Pathway Alterations


Metabolomic profiling of CSF in the NIH study also revealed abnormalities in tryptophan metabolism [5]. Tryptophan is the precursor for both serotonin and the kynurenine pathway, making its metabolism relevant to mood, cognition, and immune function.

Kynurenine Pathway Dysregulation The kynurenine pathway metabolizes approximately 95% of dietary tryptophan and produces metabolites with diverse neuroactive effects. Quinolinic acid, an NMDA receptor agonist and excitotoxin, may contribute to neuroinflammation and cognitive dysfunction when elevated. Kynurenic acid, an NMDA receptor antagonist with neuroprotective properties, can become imbalanced with quinolinic acid, disrupting glutamatergic neurotransmission. Additionally, 3-hydroxykynurenine generates reactive oxygen species, potentially contributing to oxidative stress.

Immune activation, particularly interferon-gamma, stimulates the kynurenine pathway, providing a link between the immune abnormalities and neurological symptoms observed in ME/CFS [232, 231].

Figures 8.3 and 8.4 illustrate tryptophan metabolism dysregulation in ME/CFS. While normally approximately 95% of tryptophan is metabolized via the kynurenine pathway, inflammation-driven IDO overactivation can substantially increase this proportion. If kynurenine pathway flux increases to approximately 99% (a plausible estimate based on the magnitude of IDO upregulation observed in inflammatory conditions), this seemingly modest 4 percentage-point shift would dramatically reduce serotonin-available tryptophan from 5% to 1%—an 80% reduction in serotonin precursor availability—while quinolinic acid accumulation reaches toxic levels.

Normal Catecholamine Synthesis




ME/CFS: Catecholamine Synthesis Failure

Figure 8.2: ME/CFS catecholamine synthesis failure and systemic consequences.

Normal Tryptophan Metabolism




Figure 8.3: Normal tryptophan metabolism with balanced serotonin and kynurenine pathways.


ME/CFS: Tryptophan Pathway Dysregulation

Figure 8.4: ME/CFS tryptophan dysregulation causing serotonin deficit and quinolinic acid toxicity.

Serotonin Synthesis Diversion of tryptophan into the kynurenine pathway reduces availability for serotonin synthesis. Under conditions of IDO-driven inflammatory activation, up to 90% of tryptophan is catabolized through the kynurenine pathway [232, 233], leaving substantially less for serotonin synthesis. The estimated 80% reduction in serotonin-available tryptophan (derived from IDO upregulation magnitude documented in inflammatory states) may contribute to sleep disturbances, mood symptoms, pain amplification, and cognitive impairment observed in ME/CFS.

Enterochromaffin-Vagal Pathway: Gut-Serotonin-Brain Connection Emerging research by Wirth and Scheibenbogen (2025) [225] provides a mechanistic framework linking gut dysbiosis to central serotonergic dysfunction through the enterochromaffin-vagal pathway. This pathway offers a novel perspective on serotonin regulation in ME/CFS:

~ **Hypothesis 2: Enterochromaffin-Vagal Serotonin Pathway in ME/CFS**

Certainty: 0.40. The enterochromaffin-vagal pathway may represent a critical gut-brain communication axis disrupted in ME/CFS. Enterochromaffin cells in the gut epithelium synthesize and release serotonin in response to mechanical stimulation and short-chain fatty acids (particularly butyrate). This serotonin activates 5-HT₃ receptors on vagal afferent neurons, providing tonic input to the brainstem. In ME/CFS, gut dysbiosis and reduced butyrate production may impair enterochromaffin serotonin synthesis, leading to reduced vagal afferent signaling and downstream autonomic dysfunction.

Supporting evidence:

- Gut dysbiosis with reduced butyrate-producing bacteria is well-documented in ME/CFS [226, 227]
- Enterochromaffin cells require butyrate as a substrate for serotonin synthesis [14]
- Vagal tone (measured by HRV) is consistently reduced in ME/CFS patients [255, 5]
- The pathway explains how gut dysfunction could contribute to autonomic symptoms

Implications:

- Provides mechanistic link between gut dysbiosis and autonomic dysfunction
- Suggests butyrate supplementation and vagal stimulation as complementary therapies
- Explains why some ME/CFS patients have gastrointestinal symptoms alongside autonomic dysfunction
- Offers testable predictions for future research (e.g., butyrate supplementation improving HRV)

Limitations:

- Preprint status (not yet peer-reviewed)
- Enterochromaffin serotonin release not directly measured in ME/CFS
- Causality not established (could be reverse: vagal dysfunction → dysbiosis)
- Peripheral vs. central serotonin are distinct pools with different functions

Serotonergic Dysfunction

Observation 46 (Serotonergic Abnormalities in ME/CFS). Beyond tryptophan diversion, multiple lines of evidence indicate primary serotonergic abnormalities in ME/CFS. PET imaging has demonstrated reduced serotonin transporter (5-HTT) density in the rostral anterior cingulate cortex [256]. Additional findings include abnormal responses to serotonergic challenge tests, correlations between serotonin markers and fatigue severity, and variable responses to serotonergic medications. The serotonergic system's role in regulating sleep, mood, pain perception, and autonomic function positions it as a plausible contributor to the multisystem dysfunction of ME/CFS.

Dopaminergic Dysfunction

Observation 47 (Dopaminergic Abnormalities in ME/CFS). Dopamine abnormalities extend beyond the CSF catecholamine findings to include measurably reduced basal ganglia activation during reward-processing tasks [257]. Functional MRI shows that reduced activation of the right caudate nucleus and globus pallidus correlates significantly with mental fatigue severity ($r^2 = 0.49$, $p = 0.001$). The overlap between ME/CFS fatigue and the motivational symptoms observed in dopaminergic disorders (Parkinson's disease, interferon-induced fatigue) supports a shared mechanism of inflammatory cytokine-mediated disruption of basal ganglia dopamine availability.

Norepinephrine and the Locus Coeruleus

The locus coeruleus (LC), the primary source of brain norepinephrine, plays critical roles in arousal and sleep-wake regulation, attention and cognitive flexibility, stress responses, and autonomic nervous system modulation.

~ Hypothesis 3: Locus Coeruleus Dysfunction in ME/CFS

Certainty: 0.35. LC dysfunction may explain the constellation of arousal, attention, and autonomic abnormalities in ME/CFS. The CSF catecholamine abnormalities documented in ME/CFS—including reduced DOPA, DOPAC, and DHPG—are consistent with impaired norepinephrine synthesis or turnover originating in LC neurons [5]. Candidate mechanisms include: neuroinflammation affecting LC neurons; autoantibodies targeting adrenergic receptors [71]; metabolic stress impairing catecholamine synthesis; and chronic stress-induced LC dysregulation. No study has directly measured LC structure or function in ME/CFS; this hypothesis is extrapolated from CSF biomarker data and autoimmune findings.

GABAergic and Glutamatergic Imbalance

Observation 48 (Excitatory/Inhibitory Imbalance on MRS in ME/CFS). Magnetic resonance spectroscopy (MRS) studies have identified regional neurochemical abnormalities consistent

with altered excitatory/inhibitory balance in ME/CFS [182]. Findings across studies include elevated glutamate or glutamine (Glx) in some brain regions alongside reduced GABA concentrations in others, with regional variations reflecting disease heterogeneity. This excitatory/inhibitory imbalance may contribute to sensory hypersensitivity, cognitive dysfunction, sleep disturbances, and fatigue amplification. Notably, MRS findings differ between ME/CFS and Long COVID despite clinical overlap, suggesting distinct neurochemical pathophysiologies.

Cholinergic Dysfunction

Acetylcholine abnormalities in ME/CFS have received less attention but may contribute to cognitive impairment (particularly memory), autonomic dysfunction (parasympathetic arm), sleep architecture abnormalities, and muscle function [71, 5].

Autoantibodies against muscarinic acetylcholine receptors have been identified in some ME/CFS patients [71], providing a potential autoimmune mechanism for cholinergic dysfunction.

8.1.4 Sleep Architecture and Inter-Regional Coordination

Sleep disturbances, particularly unrefreshing sleep despite adequate duration, affect up to 95% of ME/CFS patients. While subjective complaints are nearly universal, objective polysomnographic findings show more subtle alterations: longer sleep latency, reduced sleep efficiency, increased Stage 3 sleep in adults, and altered sleep microstructure [258]. The paradox—severe subjective sleep dysfunction with modest objective changes—suggests the problem may lie not in sleep duration or stage percentages, but in the *coordination* required to generate and maintain normal sleep architecture.

Energy Costs of Sleep Architecture Coordination

Normal sleep architecture requires sophisticated inter-regional brain coordination orchestrated primarily through thalamo-cortical circuits. During non-REM sleep, slow oscillations (1 Hz) originate in the anterior thalamus and precede neocortical slow oscillations, while sleep spindles (12–14 Hz) detected in thalamic nuclei precede their neocortical counterparts [259]. This sequence—convergent cortical downstates leading thalamic downstates, which then trigger spindles projected back to cortex during the down-to-upstate transition—coordinates memory consolidation across distributed brain regions [260].

Sleep spindle generation itself is metabolically demanding. Thalamic reticular nucleus (TRN) neurons must generate rhythmic bursts at 12–14 Hz, which requires sustained calcium channel activity, neurotransmitter synthesis and release, and coordinated inhibition of thalamocortical relay neurons. The cortex must then respond appropriately, amplifying spindles and coupling them with hippocampal ripples for memory consolidation. This inter-regional choreography demands substantial ATP and coordinated neurotransmitter systems.

Similarly, REM sleep requires brainstem activation (particularly cholinergic nuclei), thalamic relay, cortical activation approaching waking levels, and simultaneous motor inhibition via

brainstem circuits. The transitions between sleep stages—requiring coordinated deactivation of one set of circuits and activation of another—may be particularly energy-intensive.

~ Hypothesis 4: Sleep Architecture Failure Hypothesis

Certainty: 0.50.

In ME/CFS, CNS energy deficits and metabolic dysfunction prevent the sustained inter-regional coordination required for normal sleep architecture, resulting in fragmented sleep microstructure despite adequate total sleep time.

Mechanism. The hypothesis proposes that sleep architecture fragmentation in ME/CFS reflects energy-limited coordination failure:

1. **Spindle generation deficit:** Thalamic reticular nucleus neurons cannot sustain the metabolic demands of rhythmic 12–14 Hz burst firing, reducing sleep spindle density and power
2. **Slow-wave coordination failure:** Thalamo-cortical circuits cannot maintain synchronized slow oscillations across brain regions, fragmenting slow-wave sleep architecture
3. **Stage transition impairment:** The coordinated network reconfiguration required for sleep stage transitions (more demanding than within-stage maintenance) fails preferentially, increasing sleep fragmentation
4. **Inter-regional coherence reduction:** EEG coherence between brain regions declines during sleep, reflecting impaired functional connectivity [261]
5. **PEM-induced worsening:** During post-exertional malaise, when CNS energy deficits intensify, sleep architecture fragmentation worsens proportionally

Supporting evidence. Jackson et al. [258] meta-analyzed objective sleep data from 801 adults and 477 adolescents with ME/CFS, confirming altered sleep microstructure despite the subjective-objective paradox. Adult patients showed reduced sleep efficiency, altered stage distribution (decreased Stage 2, increased Stage 3), and longer sleep latency—patterns consistent with coordination difficulties rather than simple sleep deprivation.

Sherlin et al. [261] demonstrated that EEG spectral coherence distinguishes CFS patients from both healthy controls and depressed patients with 100% accuracy for unmedicated CFS patients. The involvement of bilateral temporal lobes in 9 of 10 coherence factors suggests widespread inter-regional connectivity disruption, supporting the coordination failure hypothesis.

Sleep fragmentation studies show that chronic fragmentation impairs brain energy metabolism to an extent similar to total sleep deprivation, with lower glucose uptake in cortex and hippocampus [262]. In ME/CFS, the causal arrow may reverse: primary metabolic dysfunction fragments sleep, which further worsens metabolism in a vicious cycle.

Testable predictions.

1. Sleep spindle density and power correlate inversely with ME/CFS symptom severity and biomarkers of CNS dysfunction
2. Slow-wave sleep fragmentation (not just total SWS percentage) correlates with measures of metabolic dysfunction (e.g., cerebral lactate on MRS)
3. Sleep architecture fragmentation worsens 24–72 hours post-exertion, tracking PEM time course
4. Sleep stage transition frequency increases (shorter, more fragmented sleep stages) compared to healthy controls, even when stage percentages appear normal
5. Inter-regional EEG coherence during sleep is reduced in ME/CFS patients, particularly in frequency bands critical for sleep oscillations (delta, sigma)
6. Interventions improving cerebral metabolism (e.g., mitochondrial support) improve objective sleep microstructure, not just subjective sleep quality

Treatment implications. If sleep architecture failure reflects energy-limited coordination, interventions should target: (1) circadian optimization—maximizing sleep opportunity during the circadian nadir when sleep pressure is highest; (2) metabolic support—mitochondrial cofactors (CoQ10, NADH) during evening hours may improve overnight cerebral metabolism [263]; (3) sleep stage-specific support—low-dose gabapentin or pregabalin may reduce thalamo-cortical excitability demands while supporting spindle generation; (4) glymphatic enhancement—sleep position (lateral decubitus), avoiding late caffeine, and sleep continuity strategies; (5) pacing-sleep integration—recognizing that sleep quality worsens predictably during PEM can guide activity management.

Limitations. This hypothesis has moderate certainty (0.50). No published studies have quantified spindle density or power in ME/CFS with simultaneous metabolic measures. Coherence data exists for waking but not sleep EEG. Alternative explanations include primary brainstem pathology, autonomic dysfunction, or circadian disruption rather than energy limitation. Causality direction remains unclear: does poor metabolism fragment sleep, or does fragmented sleep worsen metabolism?

8.1.5 Glial Cell Dysfunction

Beyond neurons and neurotransmitters, glial cells play critical support roles in brain function. Dysfunction in these cells may contribute to the neuroinflammation mentioned in catecholamine synthesis impairment and broader CNS pathology.

Microglial Activation and Neuroinflammation

Microglia, the resident immune cells of the central nervous system, have emerged as key players in ME/CFS neuroinflammation. Evidence for microglial activation includes elevated markers in CSF (soluble CD14, chitotriosidase), PET imaging showing increased translocator protein (TSPO) binding in specific brain regions [73], correlation between neuroinflammatory

markers and symptom severity, and persistence of microglial activation years after initial infection.

△ **Warning 4: Replication Status of Microglial PET Findings**

The Nakatomi et al. 2014 study documenting widespread microglial activation via PET imaging in ME/CFS patients has not been consistently replicated, with later studies showing conflicting results. The certainty of microglial activation as a universal feature of ME/CFS remains medium pending further replication studies.

Chronic microglial activation, when present, can produce sustained release of pro-inflammatory cytokines (IL-1 β , TNF- α , IL-6), oxidative stress through reactive oxygen species production, glutamate release contributing to excitotoxicity, disruption of synaptic pruning and plasticity, and blood-brain barrier dysfunction.

~ **Hypothesis 5: Glial Maturation Window and Pediatric Recovery**

Certainty: 0.45. Adolescent ME/CFS patients may benefit from a developmental window during which active microglial remodeling can reset pathological activation states—a mechanism unavailable to adult patients whose glial maturation is complete. The certainty level reflects: (1) the Nakatomi et al. 2014 PET findings documenting microglial activation have not been consistently replicated; (2) the proposed mechanism extrapolates from developmental neuroscience to ME/CFS pathophysiology; (3) testable predictions can directly address these uncertainties through age-stratified neuroimaging studies.

Background: Adolescent Microglial Maturation Microglia undergo dramatic functional reorganization during adolescence, performing complex developmental tasks beyond their immune surveillance role. From embryonic neuronal migration to adolescent circuit refinement, immune signaling molecules serve as a common language allowing microglia to modulate brain function in both health and disease [264].

Three critical periods define microglial contributions to neural development: embryonic wiring, early postnatal synaptic pruning (peak near birth continuing into late-20s), and adolescent circuit refinement [264]. During adolescence specifically, microglia mediate experience-dependent synaptic pruning through complement-mediated mechanisms, with C3 binding to CR3 receptors facilitating selective synapse elimination. This process exhibits sex-specific patterns and regional variation, with particularly robust activity in prefrontal cortex and nucleus accumbens [265, 266].

Crucially, transient microglial deficiency during adolescence—but not adulthood—produces lasting cognitive impairments, identifying adolescence as a sensitive period for prefrontal microglia to act on cognitive development [265]. The developmental program requires coordinated microglial activity for proper circuit maturation, with major transitions largely complete by early 20s.

Application to ME/CFS: The Reset Hypothesis If ME/CFS involves chronic microglial activation locked in a pro-inflammatory state (as suggested by Nakatomi et al. PET findings [73]), then adolescent microglial remodeling may provide a natural mechanism for

resolution:

1. **Active turnover:** Adolescent microglia undergo programmed replacement and phenotypic switching as part of circuit refinement, potentially eliminating pathologically activated cells
2. **Developmental override signals:** The hormonal and neurochemical milieu of adolescence (BDNF elevation, sex hormones, growth factors) provides strong pro-plasticity signals that may override inflammatory set-points
3. **Synaptic reorganization:** Pathological neuroinflammatory states often involve aberrant synaptic connections; adolescent pruning may eliminate these circuits while preserving functional connectivity
4. **Adult lock-in:** After developmental windows close (age 25), microglia lose plasticity for wholesale phenotypic switching, becoming locked in their current activation state without the developmental cues that enable adolescent reset

This framework explains why pediatric ME/CFS shows substantially higher recovery rates (estimated 54–94% in studies of mild-moderate cases) compared to adult-onset disease where recovery is rare [78]. The critical variable is not disease duration but rather whether onset occurs before or after completion of microglial maturation.

Testable Predictions This hypothesis generates specific, falsifiable predictions:

1. **Age-dependent neuroinflammation:** Longitudinal PET imaging should show declining microglial activation in recovering adolescents but persistent activation in adults with similar disease duration
2. **Transition age threshold:** Recovery rates should decline sharply around age 22–25 (completion of prefrontal maturation) rather than showing gradual age-related decline
3. **Biomarker trajectories:** CSF inflammatory markers (sCD14, chitotriosidase) should normalize in recovering adolescents but remain elevated in non-recovering adults
4. **Microglial turnover markers:** Adolescent patients should show elevated markers of microglial turnover (CSF1R expression, fractalkine signaling) compared to adults
5. **Severity interactions:** Hypothesis predicts age matters less if microglial activation is mild (can resolve spontaneously) but becomes critical if activation is severe (requires active remodeling to clear)

Treatment Implications If adolescent microglial plasticity enables recovery, then therapeutically inducing similar plasticity in adults might improve outcomes:

1. **CSF-1R inhibitors:** Drugs like PLX5622 or pexidartinib force microglial turnover by depleting existing populations and promoting repopulation from progenitors. This mimics the natural turnover occurring during adolescence, potentially resetting activation states [267].
2. **Fasting-mimicking diets:** Prolonged fasting promotes microglial autophagy and phenotypic switching, potentially enabling transition from pro-inflammatory to

surveillance phenotypes without complete depletion

3. **BDNF enhancement:** Brain-derived neurotrophic factor drives developmental plasticity; strategies to boost BDNF (exercise within energy envelope, ketogenic diet, certain medications) may partially reopen plasticity windows
4. **Timing considerations:** Interventions targeting microglial reset may be most effective in younger adults (under 30) where some residual developmental plasticity remains, with diminishing returns in older patients

Integration with Broader ME/CFS Pathophysiology This hypothesis complements rather than contradicts other mechanistic proposals. Microglial activation may be downstream of initial triggers (viral infection, autoantibodies, autonomic dysfunction) while still representing a critical perpetuating factor. The developmental window hypothesis specifically addresses *why recovery patterns differ by age* rather than explaining disease initiation.

The glial maturation window may interact synergistically with other proposed pediatric advantages: immune memory pruning (Hypothesis 8, if present), greater HSC regenerative capacity, higher baseline recovery capital (Speculation 23), and incomplete epigenetic aging.

Limitations and Uncertainties Several important caveats apply:

1. The Nakatomi et al. microglial activation findings have not been consistently replicated; if microglial activation is not a universal ME/CFS feature, this hypothesis applies only to a subset
2. The proposed mechanism assumes glial maturation windows close around age 25, but individual variation exists; some adults may retain plasticity longer
3. Pediatric recovery may reflect multiple mechanisms simultaneously; isolating the specific contribution of microglial remodeling requires longitudinal studies with neuroimaging
4. CSF-1R inhibitor strategies carry significant risks (meningitis, visual changes) and remain experimental; safety in ME/CFS populations is unknown

Research Priorities To test this hypothesis rigorously:

1. **Age-stratified longitudinal neuroimaging:** Serial PET scans in adolescent vs adult ME/CFS tracking microglial activation trajectories over 2–5 years
2. **CSF biomarker studies:** Compare inflammatory markers and microglial turnover signatures across age groups and recovery status
3. **Preclinical models:** Post-viral fatigue models in adolescent vs adult mice to test whether developmental microglia enable recovery
4. **Treatment trials:** Small pilot studies of CSF-1R modulation in carefully selected adult ME/CFS patients with documented microglial activation

This hypothesis provides a mechanistic framework for understanding one component of the pediatric recovery advantage while suggesting potential therapeutic strategies for

adult patients.

Astrocyte Abnormalities and the Astrocyte Energy Gate

Astrocytes perform essential functions including neurotransmitter uptake and recycling, blood-brain barrier maintenance, metabolic support for neurons, synaptic modulation, and ion homeostasis. Astrocyte dysfunction in ME/CFS may contribute to impaired glutamate clearance and excitotoxicity, reduced metabolic support for neurons, blood-brain barrier compromise, and abnormal synaptic transmission. Elevated GFAP (glial fibrillary acidic protein) in some ME/CFS patients suggests astrocyte reactivity, though findings have been inconsistent.

Beyond these recognized roles, astrocytes occupy a uniquely critical position in brain energy metabolism that may constitute a central vulnerability in ME/CFS. The following hypothesis develops this metabolic dimension in detail.

The Astrocyte-Neuron Lactate Shuttle: Normal Physiology The brain consumes 20–25% of the body's glucose despite comprising only 2% of body mass [176]. A substantial fraction of this energy reaches neurons not as glucose directly, but via the **astrocyte-neuron lactate shuttle** (ANLS), first described by Pellerin and Magistretti [174]. In this system, glutamate released during synaptic transmission is taken up by astrocytes via excitatory amino acid transporters (EAATs), triggering astrocytic glucose uptake through GLUT1 transporters and subsequent glycolysis. Astrocytes convert glucose to pyruvate and then to lactate via lactate dehydrogenase A (LDHA), which preferentially catalyzes the pyruvate-to-lactate direction. This lactate is then exported from astrocytes through monocarboxylate transporter 4 (MCT4, a low-affinity, high-capacity exporter) and imported into neurons through MCT2 (a high-affinity importer) [268]. Within neurons, LDHB converts lactate back to pyruvate for oxidative phosphorylation in mitochondria.

This architecture elegantly couples neuronal energy supply to neuronal activity: when a synapse fires, the glutamate released simultaneously signals the local astrocyte to increase energy delivery [175]. Lactate provides an estimated 30–50% of neuronal ATP under physiological conditions [175], and this fraction likely increases during periods of intense neural activity when neurons' own glycolytic capacity is insufficient.

Several features make this shuttle critical rather than merely supplementary:

1. **Activity coupling:** The glutamate-triggered mechanism ensures energy supply scales with demand at the single-synapse level
2. **Metabolic specialization:** Neurons preferentially express LDHB (favoring lactate → pyruvate) while astrocytes express LDHA (favoring pyruvate → lactate), creating directional metabolic flow [269]
3. **Antioxidant protection:** By outsourcing glycolysis to astrocytes, neurons can direct more glucose through the pentose phosphate pathway for glutathione regeneration, protecting against oxidative damage

4. **Signaling function:** Lactate also acts as a signaling molecule via the hydroxycarboxylic acid receptor 1 (HCAR1/GPR81), modulating neuronal excitability and synaptic plasticity [175]

Important Nuance: Neuronal Metabolic Flexibility The classical ANLS model has been refined by recent evidence demonstrating that neurons possess greater metabolic flexibility than originally assumed. Single-cell RNA sequencing studies reveal that neurons express both LDHA and LDHB, not exclusively LDHB [269]. Neurons can directly take up and oxidize glucose, particularly during high-demand states. LDHB-deficient neurons maintain stable energy metabolism under physiological glucose conditions, suggesting compensatory pathways exist.

However, this flexibility has limits. During high-frequency neural activity—precisely the conditions of cognitive exertion—direct neuronal glucose oxidation may prove insufficient, and astrocyte-derived lactate becomes the critical marginal fuel source. This distinction between *basal* sufficiency and *demand-responsive* insufficiency is central to the hypothesis that follows.

Speculation 11 (Astrocyte Energy Gate). Certainty: 0.35. We hypothesize that dysfunction in the astrocyte-neuron lactate shuttle creates a **metabolic bottleneck**—an “energy gate”—that produces CNS-specific energy failure in ME/CFS while peripheral tissues with direct glucose access remain unaffected.

Three Candidate Mechanisms The energy gate may fail at any of three nodes, singly or in combination:

1. **Astrocyte glucose uptake impairment (GLUT1 dysfunction):** Reduced GLUT1 expression or function on astrocytes limits the raw substrate entering the shuttle. GLUT1 deficiency syndrome demonstrates that impaired astrocytic glucose transport causes seizures, cognitive impairment, and brain hypometabolism—features that partially overlap with ME/CFS neurological symptoms. Neuroinflammatory mediators (IL-1 β , TNF- α) documented in ME/CFS can downregulate GLUT1 expression.
2. **Lactate production impairment (glycolytic defects):** Reactive astrogliosis—documented via elevated GFAP in ME/CFS—involves metabolic reprogramming that may paradoxically impair effective lactate delivery. While reactive astrocytes initially upregulate glycolysis, chronic neuroinflammation shifts astrocyte metabolism toward a state where mitochondrial dysfunction reduces overall metabolic efficiency. Inflammatory cytokines can alter pyruvate dehydrogenase kinase (PDK) activity, disrupting the glycolysis/oxidative phosphorylation balance within astrocytes themselves.
3. **Lactate transport impairment (MCT dysfunction):** Downregulation of MCT4 (astrocyte export) or MCT2 (neuronal import) directly restricts lactate flow. This mechanism has the strongest precedent in other neurological diseases: MCT1/MCT4 downregulation reduces neuronal lactate supply by approximately 60% [269]. In Alzheimer’s disease, decreased expression of MCT1, MCT2, and MCT4 is documented. In amyotrophic lateral sclerosis, reduced MCT1 in oligodendrocytes precedes motor neuron degeneration.

In temporal lobe epilepsy, MCT2 redistribution and MCT4 reduction are observed in epileptic foci.

Why This Creates Selective Dysfunction The energy gate hypothesis explains why CNS function fails while peripheral tissues remain functional:

- **CNS vulnerability:** Neurons depend on the ANLS for a substantial portion of their activity-dependent energy supply. No other cell type in the body has this intermediary requirement for its primary fuel.
- **Peripheral independence:** Skeletal muscle, cardiac muscle, and peripheral tissues express GLUT4 (insulin-responsive) and can directly oxidize glucose without astrocytic intermediation. Hair follicles operate autonomous local Cori cycles, recycling lactate within the follicular unit without CNS coordination.
- **Demand-dependence:** The ANLS is most critical during cognitive exertion (when glutamate release surges trigger proportional lactate demand). This explains why cognitive symptoms worsen with mental effort while resting cognition may remain closer to normal—a hallmark of ME/CFS “brain fog.”

This mechanism connects directly to the selective energy dysfunction hypothesis (Section 14.24), which predicts that high CNS-dependency (α) and high demand-responsiveness (ρ) processes should be most impaired. The ANLS provides the *specific molecular mechanism* through which this selective vulnerability operates.

Certainty Assessment This speculation integrates well-established neuroscience (ANLS physiology: high certainty) with speculative application to ME/CFS (low-to-moderate certainty). No study has directly measured ANLS flux, MCT expression, or astrocyte-specific glycolytic rates in ME/CFS patients. The hypothesis is graded at **certainty 0.35**: mechanistically plausible, consistent with indirect evidence, but requiring direct experimental validation.

Supporting Evidence: Brain Lactate Elevation While no study has directly assayed ANLS function in ME/CFS, magnetic resonance spectroscopy (MRS) studies provide indirect evidence consistent with impaired brain energy metabolism:

- **7T MRS (2025):** Godlewska et al. [182] found elevated lactate in the pregenual anterior cingulate cortex (pgACC: 1.52 vs. 1.22 mM, $p = 0.003$) and dorsal ACC (dACC) of ME/CFS patients ($n=24$) compared to healthy controls ($n=24$), using ultra-high-field 7 Tesla MRS. Notably, ME/CFS and Long COVID patients showed *different* neurochemical profiles despite similar clinical presentations.
- **Whole-brain MRS (2020):** Mueller et al. [270] documented elevated lactate-to-creatine ratios in the right insula, thalamus, and cerebellum ($n=15$ ME/CFS vs. $n=15$ controls), with brain temperature increases correlated with lactate elevations—suggesting neuroinflammation drives metabolic shifts.

- **Mitochondrial review (2025):** Syed et al. [64] synthesize evidence of elevated CSF lactate, impaired ATP synthesis, and increased glycolytic activity in ME/CFS, consistent with oxidative stress and conditions favoring anaerobic metabolism.

△ **Warning 5: Interpreting Elevated Brain Lactate**

Elevated brain lactate in ME/CFS is consistent with the energy gate hypothesis but does not uniquely support it. At least three interpretations are possible:

1. **ANLS dysfunction:** Lactate accumulates in astrocytes because it cannot be efficiently exported to or utilized by neurons (supports the energy gate hypothesis)
2. **Mitochondrial dysfunction:** Neuronal mitochondria cannot oxidize lactate efficiently, causing backpressure (supports a downstream mitochondrial hypothesis)
3. **Anaerobic shift:** Increased glycolysis due to hypoperfusion or oxygen limitation produces excess lactate (supports a vascular hypothesis)

These mechanisms are not mutually exclusive and may operate simultaneously. Distinguishing between them requires studies that measure not just lactate levels but lactate *flux* between cellular compartments—technically challenging but feasible with advanced ¹³C-MRS techniques.

Testable Predictions The astrocyte energy gate hypothesis generates specific, falsifiable predictions that distinguish it from alternative explanations:

1. **CSF lactate gradient:** If astrocytes produce lactate but neurons cannot utilize it, the CSF lactate/blood lactate ratio should be elevated in ME/CFS (astrocyte-derived lactate accumulating in extracellular space). In mitochondrial disorders affecting the CNS, a CSF/blood lactate ratio > 0.91 indicates central origin [64].
Prediction: ME/CFS patients will show CSF/blood lactate ratio > 0.91, distinguishing CNS-origin lactate from peripheral sources.
2. **MCT expression profiling:** Post-mortem or biopsy studies should reveal reduced MCT2 (neuronal) and/or MCT4 (astrocyte) expression in ME/CFS brain tissue, particularly in regions showing functional deficits (prefrontal cortex, anterior cingulate).
Prediction: MCT2/MCT4 expression reduced ≥30% vs. matched controls.
3. **Astrocyte-specific metabolomics:** Single-cell or spatial transcriptomics of ME/CFS brain tissue should show altered expression of glycolytic enzymes (hexokinase, phosphofructokinase, LDHA) and glucose transporters (GLUT1) in astrocytes specifically.
Prediction: Astrocyte glycolytic gene expression altered while neuronal oxidative genes remain intact.
4. **Exogenous lactate challenge:** If the bottleneck is at the glucose → lactate step (mechanisms 1 or 2 above), then providing exogenous lactate should partially bypass the gate and improve cognitive function. If the bottleneck is at MCT transport (mechanism 3), exogenous lactate should not help.
Prediction: IV sodium lactate infusion during cognitive testing will improve performance in a subgroup of ME/CFS patients.
5. **Ketone body bypass:** Ketone bodies (β-hydroxybutyrate, acetoacetate) enter neurons via MCT2 and are metabolized directly in neuronal mitochondria, bypassing the astrocyte

glycolysis step entirely [271]. If the energy gate is at the astrocyte level, ketones should preferentially benefit CNS symptoms.

Prediction: Ketogenic diet or exogenous ketone supplementation will improve cognitive symptoms disproportionately to peripheral fatigue symptoms.

6. **Activity-dependent worsening:** Since the ANLS is most critical during high neural activity (when glutamate-triggered demand surges), the energy gate should cause greater deficits during cognitive exertion than at rest.

Prediction: The difference between resting and task-evoked brain lactate (measured by functional MRS) will be larger in ME/CFS than controls—reflecting both increased demand signaling and impaired supply response.

Treatment Implications The energy gate framework suggests several therapeutic strategies, ordered by plausibility and feasibility:

1. **Ketogenic diet or exogenous ketones:** By providing β -hydroxybutyrate directly to neurons via MCT2, this approach bypasses the astrocyte glycolysis step entirely. The ketogenic diet has established neuroprotective effects in epilepsy (where MCT dysfunction is documented) and emerging evidence in psychiatric disorders associated with brain energy dysfunction [271]. *This represents the most immediately testable intervention.*
2. **Exogenous lactate supplementation:** Sodium lactate infusion or oral lactate has shown cognitive benefits in Alzheimer's disease models by restoring hippocampal and CSF lactate concentrations. In ME/CFS, this could bypass impaired astrocyte glycolysis (mechanisms 1–2) but would not help if MCT2 transport is the bottleneck (mechanism 3).
3. **MCT upregulation:** Exercise and certain pharmacological agents can upregulate MCT expression. However, exercise intolerance in ME/CFS limits this approach. Pharmacological MCT modulators remain experimental.
4. **Anti-neuroinflammatory strategies:** If chronic neuroinflammation drives astrocyte metabolic reprogramming and MCT downregulation, targeting neuroinflammation at its source may restore ANLS function. Low-dose naltrexone (LDN), which modulates microglial activation, could theoretically improve astrocyte metabolic function through reduced neuroinflammatory signaling.
5. **Astrocyte-targeted delivery:** Emerging drug delivery technologies using astrocyte-specific targeting (e.g., nanoparticles with GFAP-binding peptides) could deliver metabolic support directly to astrocytes, enhancing glycolytic capacity or MCT expression without systemic effects.

Limitations and Alternative Explanations Several important caveats apply to this hypothesis:

- **No direct evidence in ME/CFS:** No study has measured ANLS flux, MCT expression, or astrocyte-specific glycolytic rates in ME/CFS patients. The hypothesis rests entirely on indirect evidence (elevated brain lactate, documented neuroinflammation) and analogy to other neurological conditions.

- **Elevated lactate is ambiguous:** As noted above, elevated brain lactate has at least three interpretations. The ANLS dysfunction interpretation is not uniquely supported by current data.
- **The ANLS itself is debated:** While the ANLS is well-established, its quantitative contribution remains contested. Some evidence suggests neurons can sustain activity through direct glucose oxidation alone, at least under non-demanding conditions [269]. The hypothesis is strongest for high-demand cognitive states.
- **Downstream mitochondrial dysfunction:** Even if lactate reaches neurons normally, impaired neuronal mitochondria (a well-documented finding in ME/CFS [64]) would produce similar symptoms. The energy gate and mitochondrial hypotheses are not mutually exclusive but have different treatment implications.
- **GLUT1 paradox:** Recent studies show that astrocyte-specific GLUT1 reduction can *paradoxically improve* brain glucose metabolism, suggesting compensatory mechanisms may complicate predictions based on simple GLUT1 downregulation.
- **Small sample sizes:** The MRS studies supporting brain lactate elevation in ME/CFS have samples of n=15–24, which limits statistical power and generalizability. Larger, multi-site replication studies are needed.

For the relationship between the astrocyte energy gate and the broader selective energy dysfunction framework, including formal mathematical treatment and additional predictions, see Section 14.24, specifically the astrocyte energy gate sub-hypothesis (38).

Oligodendrocyte Function

Oligodendrocytes produce the myelin sheaths essential for rapid nerve conduction. Potential abnormalities include demyelination contributing to white matter hyperintensities, impaired remyelination capacity, oxidative damage to oligodendrocytes, and disrupted axon-glia signaling.

? Open Question 1: Oligodendrocyte Contribution to White Matter Changes

Do the white matter changes observed on MRI in ME/CFS patients reflect oligodendrocyte dysfunction? No studies have directly examined oligodendrocyte pathology in ME/CFS, and the mechanisms linking white matter hyperintensities to oligodendrocyte function remain to be elucidated.

Post-Viral CNS Reprogramming

The preceding sections document microglial activation, astrocyte dysfunction, and neuroinflammatory cascades in ME/CFS. A critical question is why these states persist long after the triggering infection resolves. Emerging evidence from trained immunity research suggests that a single viral infection can permanently alter glial cell function through epigenetic mechanisms.

Speculation 12 (Post-Viral CNS Reprogramming Hypothesis). **Certainty: 0.40.**

Viral infection causes persistent epigenetic reprogramming of astrocytes and microglia, creating a lasting shift in CNS metabolism that persists long after viral clearance. This mechanism—analogue to “trained immunity” in peripheral innate immune cells—may explain why ME/CFS becomes chronic following acute infection.

Trained immunity and epigenetic memory. Trained immunity refers to the capacity of innate immune cells to develop long-lasting functional memory following initial stimulation [272]. Unlike adaptive immune memory mediated by lymphocyte clonal expansion, trained immunity operates through persistent epigenetic modifications—particularly histone marks such as H3K4me1 and H3K27ac at inflammatory gene promoters—that prime cells for enhanced responses to subsequent stimuli. Humer et al. advocate trained immunity as a contributing factor to ME/CFS pathogenesis, proposing that post-infectious epigenetic reprogramming of innate immune cells produces a hyperresponsive phenotype that sustains chronic inflammation [272].

Microglial epigenetic reprogramming. Wendeln et al. demonstrated in a landmark *Nature* study that peripheral inflammatory stimuli induce long-lasting epigenetic reprogramming of brain microglia [273]. Trained microglia develop enhanced H3K4me1 marks at inflammatory gene loci that persist for months after the initial stimulus and exacerbate subsequent neurological pathology. This immune memory operates through metabolic reprogramming: activated microglia shift from oxidative phosphorylation to aerobic glycolysis, and this metabolic phenotype becomes epigenetically stabilized [274]. The persistence of these marks means that a single viral infection could establish a “new normal” of microglial function that outlasts the infection by years.

Viral reprogramming of glial metabolism. Rodrigues et al. demonstrate that neurotropic viruses—including SARS-CoV-2, HIV-1, and Zika virus—directly infect astrocytes and microglia, causing metabolic shifts from oxidative phosphorylation to glycolysis with consequent NLRP3 inflammasome activation [275]. This metabolic reprogramming is not merely a transient response to active infection; it persists because the glycolytic shift triggers epigenetic modifications that stabilize the pro-inflammatory phenotype. The result is a self-sustaining cycle: viral infection → metabolic shift → epigenetic stabilization → chronic neuroinflammation.

This mechanism has direct relevance to the astrocyte energy gate hypothesis (Section 8.1.5): if viral infection reprograms astrocytes to favor glycolysis over oxidative phosphorylation, lactate production via the astrocyte-neuron lactate shuttle may be disrupted. Astrocytes locked in a glycolytic-inflammatory phenotype may consume glucose for their own inflammatory signaling rather than converting it to lactate for neuronal use.

Broader epigenetic landscape in ME/CFS. Apostolou and Rosén document over 12,000 altered CpG methylation sites in ME/CFS patients, with particular enrichment at immune and metabolic gene loci [276]. They propose that latent herpesviruses (particularly EBV) employ long-term epigenetic strategies that may permanently alter host cell function. Complementing this, Iu et al. demonstrate that CD8⁺ T cells in ME/CFS exhibit epigenetic predisposition toward terminal exhaustion, with exhaustion markers upregulated following exercise challenge [119]—linking immune reprogramming directly to post-exertional malaise.

Testable predictions.

1. Post-infectious ME/CFS patients should show distinct microglial epigenetic signatures (elevated H3K4me1/H3K27ac at inflammatory loci) compared to gradual-onset patients, detectable via CSF-derived microglia or post-mortem analysis
2. Astrocyte metabolic profiles (measurable via MRS glutamate/glutamine ratios) should differ between post-viral and non-viral ME/CFS subtypes
3. Epigenetic modifiers targeting trained immunity (e.g., histone methyltransferase inhibitors, mTOR pathway modulators) should preferentially benefit post-infectious ME/CFS
4. Early antiviral or anti-inflammatory intervention during acute infection should reduce ME/CFS incidence by preventing epigenetic stabilization
5. CSF cytokine profiles should show trained immunity signatures (enhanced IL-6, TNF- α responses to ex vivo stimulation) in post-infectious but not gradual-onset patients

Treatment implications. If post-viral CNS reprogramming drives chronic ME/CFS, therapeutic strategies should target epigenetic reversal rather than symptomatic suppression: (1) epigenetic modifiers that can cross the BBB and reset microglial histone marks; (2) metabolic interventions that shift astrocytes back from glycolysis to oxidative phosphorylation; (3) microglial depletion and repopulation via CSF-1R inhibition (see Section 8), which may generate microglia without the trained immunity epigenetic marks; (4) early intervention protocols during acute viral illness to prevent epigenetic stabilization.

Limitations. This hypothesis has certainty 0.40. No study has directly measured trained immunity epigenetic marks in ME/CFS patient microglia. The evidence is synthesized from trained immunity research in neurodegenerative diseases [273, 277], viral glial reprogramming [275], and ME/CFS epigenetic profiling [276]. The mechanism may not explain gradual-onset ME/CFS without clear viral trigger. Individual variation in epigenetic susceptibility and viral tropism could produce heterogeneous responses.

8.1.6 Integrated Neuroinflammatory Cascade Model

The diverse neurological abnormalities documented in ME/CFS—neurotransmitter depletion, microglial activation, autonomic dysregulation—may not be independent pathologies but

rather interconnected components of a unified cascade originating from the central nervous system.

~ **Hypothesis 6: Neuroinflammatory Cascade: From CNS to Peripheral Symptoms**

Certainty: 0.50. We propose an integrated cascade model in which neuroinflammatory dysfunction serves as an upstream driver of both central and peripheral ME/CFS pathology [278, 279]:

Cascade pathway *Infection or immune challenge:* Initial infection (EBV, enterovirus, or other trigger) activates innate immunity and produces transient CNS inflammation through multiple routes (direct viral CNS invasion, systemic inflammatory cytokines crossing BBB, peripheral immune cell infiltration).

Sleep disruption and impaired glymphatic clearance: Acute neuroinflammation disrupts sleep architecture and circadian regulation. Critically, sleep loss impairs the glymphatic system—the brain’s waste clearance mechanism dependent on aquaporin-4 water channels in astrocytes. During sleep, the glymphatic system increases interstitial space and clears accumulated metabolic byproducts. Without adequate sleep, toxic protein aggregates (misfolded proteins, amyloid, tau) accumulate in the parenchyma.

Persistent neuroinflammation and microglial priming: Impaired glymphatic clearance allows accumulation of pathogen-associated molecular patterns (PAMPs) and damage-associated molecular patterns (DAMPs), which sustain microglial activation. Primed microglia become hyperresponsive to subsequent stimuli, producing exaggerated cytokine responses (IL-1 β , TNF- α , IL-6) to minor perturbations.

Central neurotransmitter depletion: Sustained neuroinflammation and microglial activation reduce synthesis of catecholamines and serotonin through multiple mechanisms: (1) inflammatory cytokines inhibit tyrosine hydroxylase and tryptophan hydroxylase expression, (2) oxidative stress from microglia-derived reactive oxygen species damages the enzymes and their cofactors, (3) catecholamine reuptake is impaired by cytokine-mediated transporter dysfunction, (4) metabolic depletion reduces substrate availability for neurotransmitter synthesis.

Central neurological dysfunction: Catecholamine and serotonin depletion produce multiple consequences: effort-related dysfunction (hyperdopaminergic responses to exertion trigger rapid catecholamine depletion, producing the post-exertional symptom surge characteristic of PEM), cognitive dysfunction (prefrontal catecholamine depletion impairs attention, working memory, and executive function), and sickness behavior activation (inflammatory cytokines and depleted monoamines trigger the evolutionarily conserved sickness behavior program—fatigue, anhedonia, reduced activity tolerance—which is protective but becomes maladaptive when persistent).

Autonomic dysregulation: Depleted brainstem catecholamine systems (particularly the locus coeruleus) and impaired parasympathetic signaling (reduced acetylcholine availability) produce observable autonomic dysfunction: reduced heart rate variability, abnormal blood pressure regulation (orthostatic intolerance, POTS-like features), impaired vagal anti-inflammatory signaling, and loss of normal sympatho-parasympathetic balance.

Peripheral symptom manifestation: The combination of catecholamine depletion, sickness behavior, and autonomic dysregulation produce the characteristic ME/CFS symptom constellation: profound fatigue, post-exertional malaise, cognitive dysfunction, pain, and orthostatic intolerance.

Metabolic dysfunction and amplification loop: Forced inactivity (due to neurologically-driven inability to exert), medication effects (many treatments deplete catecholamines further), and chronic systemic inflammation drive metabolic dysfunction: mitochondrial ATP production declines, lactate accumulation increases, metabolic flexibility is impaired. Metabolic dysfunction itself produces inflammatory signals (lactate, damaged mitochondria) that amplify neuroinflammation. This creates a positive feedback loop: neuroinflammation → peripheral symptoms → reduced activity → metabolic dysfunction → amplified neuroinflammation.

Key assumptions This cascade model rests on a critical causal assumption: **central nervous system dysfunction is causally primary**, driving peripheral manifestations rather than resulting from them. Alternative causal hierarchies are biologically plausible. For example, if primary immune dysfunction (impaired viral clearance, B cell dysfunction, autoantibody production) drives disease, peripheral pathology would come first, and CNS involvement would be secondary. Similarly, if metabolic dysfunction (mitochondrial ATP depletion, lactate accumulation) is the primary driver, neurological changes might reflect metabolic rather than neuroinflammatory etiology. These alternative models would predict different therapeutic hierarchies and treatment response patterns. The cascade model specifically predicts that CNS-targeted interventions (sleep restoration, microglial modulation, catecholamine restoration) should be foundational to treatment, whereas peripheral organ-targeted therapy (cardiac drugs for POTS, antivirals for presumed viral persistence) would be less effective if peripheral dysfunction is secondary. Testing this assumption requires comparative treatment trials: if CNS-first approaches produce superior outcomes to periphery-first approaches in randomized trials, the cascade model's assumption is supported; if peripheral approaches are equally or more effective, the causality assumption is questioned.

Key implications of this model This cascade model positions *central neurological dysfunction as upstream of peripheral symptoms* rather than secondary to them. If correct, it suggests fundamentally different therapeutic strategies than those targeting peripheral organs:

1. **Sleep is disease-modifying:** Sleep disruption perpetuates the cascade by impairing glymphatic clearance. Interventions that restore sleep (sleep hygiene, low-dose sedating agents, circadian restoration) may directly interrupt neuroinflammation, not merely improve symptoms.
2. **Microglial modulation is central:** Interventions targeting microglial activation (CSF-1R inhibition as discussed in Section 8.1.5, fasting-mimicking diets promoting microglial turnover) may provide disease-modifying benefit.
3. **Catecholamine restoration requires CNS targeting:** Peripheral catecholamine re-

placement (standard treatments for POTS) may be ineffective if the primary problem is CNS depletion and impaired synthesis. Centrally-acting drugs (L-DOPA, levodopa with carbidopa to cross BBB, dopamine agonists) might be more effective than peripheral sympathomimetics.

4. **Breaking the positive feedback loop is critical:** Preventing forced inactivity through appropriate pacing prevents the metabolic dysfunction that amplifies neuroinflammation. This aligns with clinical observations that strict pacing produces better long-term outcomes than progressive exercise approaches.

⚠ **Limitation 3: Kindling Analogy: Neurological Extrapolation Without ME/CFS Data**

The PEM kindling hypothesis (below) applies a concept from epilepsy and substance abuse neuroscience to ME/CFS by analogy. No study has measured progressive microglial priming, cumulative threshold reduction, or kindling-like neural sensitisation in ME/CFS patients. The quantitative model parameters ($\alpha = 1.5$, exponential decay) are illustrative, not empirically derived, and the model has not been fitted to patient data. Alternative explanations for progressive worsening—deconditioning, psychological avoidance learning, cumulative tissue damage—have not been excluded.

8.1.7 Post-Exertional Malaise and the Kindling Hypothesis

The clinical observation that each crash lowers the threshold for the next crash—that activities previously tolerated trigger worse symptoms as disease progresses—parallels a phenomenon well-established in neurology: kindling.

~ **Hypothesis 7: Post-Exertional Malaise Kindling and Progressive Sensitization**

Certainty: 0.45. We propose that PEM represents a form of neurobiological kindling in which repeated neuroinflammatory activation progressively lowers the threshold for triggering symptom exacerbations [73, 278, 279].

Kindling mechanism *Initial exertion:* An activity requiring catecholamine-dependent effort (physical exertion, cognitive demanding tasks, emotional stress, or infection) triggers acute catecholamine release from depleted stores. If CNS catecholamine availability is already compromised by neuroinflammation, even a modest exertion produces a substantial percentage depletion of the remaining pool.

Threshold and collapse: The neuronal systems dependent on catecholamines cannot function effectively once availability drops below a critical threshold. This produces the acute collapse characteristic of PEM: sudden fatigue, cognitive shutdown, pain, orthostatic intolerance.

Microglial priming from exertion: The acute catecholamine depletion and cellular stress from exertion act as a DAMP (damage-associated molecular pattern), priming already-activated microglia further. Additionally, the metabolic disruption during exertion (in-

creased lactate, ROS production, cellular damage) provides more inflammatory signals.

Lowered threshold post-exertion: Following a PEM episode, microglial priming increases. The threshold for the next crash (T_2) is lower than the threshold before (T_1): activities that previously could be tolerated now trigger crashes because less catecholamine depletion is required to cross the now-lower threshold.

Progressive sensitization: With repeated PEM episodes, this kindling process repeats: $T(n) < T(n-1)$. Each crash further primes microglia, further sensitizes the system, further lowers the threshold. Over time, trivial activities trigger crashes. Some patients reach a state where standing, conversations, or eating triggers symptoms.

Quantitative model Let $T(n)$ be the activity threshold at time n (e.g., kcal expended before triggering PEM):

- Initial state: $T(0)$ = baseline (e.g., 500 kcal before crash triggered)
- First crash: Exertion approaching $T(0)$ triggers depletion below critical threshold, PEM occurs, microglial priming increases by factor α
- Post-crash state: $T(1) = T(0) / \alpha$ (threshold lowered by priming factor)
- Second crash: Exertion of magnitude $T(1)$ triggers symptoms; microglial priming increases further
- Recursive decline: $T(n) = T(n-1) / \alpha = T(0) / \alpha^n$

With priming factor $\alpha = 1.5$ (a 50% lowering per crash), the progression would be: $T(0) = 500 \text{ kcal} \rightarrow T(1) = 333 \text{ kcal} \rightarrow T(2) = 222 \text{ kcal} \rightarrow T(5) = 65 \text{ kcal}$

Note on model parameters: The priming factor α and the specific threshold values shown (500, 333, 222, 65 kcal-equivalent) are illustrative only and not empirically derived. The actual value of α is unknown and likely varies substantially between patients depending on baseline microglial activation state, genetic factors affecting neuroinflammatory response, and disease duration. These example values are presented solely to demonstrate the exponential relationship between crash number and threshold reduction. Any quantitative application of this model requires direct empirical measurement of individual patient thresholds over time.

Clinical and prognostic implications This kindling model explains several critical clinical observations:

Crash begets crashes: The threshold-lowering effect means that a single exertion event doesn't just cause temporary symptoms but permanently alters the disease trajectory by priming for future crashes. This has profound implications for disease modification.

Strict pacing prevents further sensitization: If exertions are carefully limited to sub-threshold levels (well below the current threshold $T(n)$), no additional crash occurs and microglial priming does not increase further. This prevents the recursive threshold decline. In this framework, strict pacing is not merely symptomatic management but disease-modifying—it halts the progressive sensitization process. Patients who maintain strict pacing may stabilize at their current threshold; those who allow repeated crashes will worsen progressively.

Infections produce major priming events: Each infection represents a major immuno-

logical and neuroinflammatory event. In the kindling framework, infection reactivation (EBV, HHV-6) or new infection produces substantial microglial priming, equivalent to multiple PEM episodes. This explains the clinical pattern that infections mark step-wise deterioration in ME/CFS—they reset the kindling process upward.

Recovery becomes progressively harder: In early disease (low n , high $T(n)$), exertions are still available that don't trigger crashes; nervous system can gradually rebuild reserves. As kindling progresses (high n , low $T(n)$), almost all activities trigger crashes; positive feedback dominates. Recovery requires not just stopping new crashes but actively depriming microglia. This becomes increasingly difficult as the patient becomes more sensitized.

Early intervention is critical: At disease onset (low n), the threshold has not dropped far. Early application of strict pacing and anti-neuroinflammatory interventions could potentially prevent the recursive decline. Later, after many crashes, the threshold has dropped far and recovery requires intensive depriming. This suggests that early aggressive management (e.g., immediate bed rest, microglial suppression, infection prevention, metabolic support) following disease onset might prevent chronic progression, whereas late intervention faces an already-sensitized nervous system.

Treatment implications If the kindling hypothesis is correct:

Strict pacing is disease-modifying: Currently, pacing is recommended as symptomatic management. The kindling model suggests it should be recognized as disease-modifying—directly interrupting the progressive sensitization process. Patients who maintain pacing avoid further kindling and preserve their remaining threshold. Those who do not may see progressive functional decline.

Blocking new kindling triggers is critical: Infections are major microglial priming events. Preventing infections (FFP2 masking in high-transmission periods, prophylactic antivirals if options become available, rapid treatment of infections) becomes disease-modifying therapy because it prevents the threshold-lowering spike from infection-induced microglial activation.

Active depriming requires intervention: Merely halting new crashes (pacing) prevents further decline but doesn't reverse existing kindling. If the hypothesis is correct, therapies that actively reverse microglial priming (CSF-1R inhibition to deplete and regenerate microglia, fasting-promoting interventions to reset glial metabolism, neuroplasticity-promoting therapies like low-dose BDNF or photobiomodulation) might restore threshold to baseline over time.

Falsification criteria The kindling hypothesis makes specific predictions that can be empirically refuted. The hypothesis would be falsified by:

1. **Absence of cumulative threshold reduction:** If longitudinal studies controlling for overall disease progression show that repeated PEM episodes do not produce measurable cumulative lowering of subsequent thresholds, the kindling mechanism would be unsupported. For example, if two patient groups with similar baseline disease duration and severity show the same activity threshold despite vastly different crash histories, kindling-mediated threshold reduction would be unlikely.

2. **Reversibility of thresholds after prolonged rest:** If extended rest periods (3+ months) with strict activity limitation consistently restore pre-crash thresholds to baseline levels, this would suggest sensitization is reversible rather than kindling-like. True kindling produces cumulative, largely irreversible changes; reversible sensitization would point to different mechanisms (e.g., temporary glial activation without permanent priming).
3. **Absence of microglial correlates:** If microglial markers (CSF1-R positron emission tomography imaging, cerebrospinal fluid inflammatory mediators, or microglial activation markers) show no correlation with PEM crash history, threshold reduction, or disease severity, this would weaken the proposed microglial mechanism. Conversely, finding these markers elevated equally in patients with few versus many crashes would suggest microglial involvement is secondary rather than driving kindling.
4. **Lack of threshold reduction with infection-equivalent priming:** If experimental immune activation (e.g., viral challenge or endotoxin administration) that triggers robust microglial and systemic inflammatory responses does not produce measurable threshold lowering in animal models of ME/CFS-like disease, the kindling mechanism would be questionable.

8.2 Autonomic Nervous System Dysfunction

Autonomic dysfunction is nearly universal in ME/CFS and contributes substantially to disability. The NIH deep phenotyping study provided quantitative documentation of specific autonomic abnormalities [5].

8.2.1 Sympathetic vs. Parasympathetic Imbalance

Heart Rate Variability Studies

Heart rate variability (HRV) provides a non-invasive window into autonomic function. The NIH study documented significantly diminished HRV in PI-ME/CFS patients compared to controls [5]. These changes included reduced overall variability (lower standard deviation of NN intervals or SDNN, reflecting decreased overall autonomic modulation), diminished high-frequency power (reduced HF-HRV, specifically reflecting decreased parasympathetic or vagal activity), altered low-frequency power (changes in LF-HRV, influenced by both sympathetic and parasympathetic activity), and abnormal LF/HF ratio (suggesting sympathovagal imbalance).

Clinical Implications of Reduced HRV Diminished HRV in ME/CFS correlates with greater fatigue severity (Escorihuela et al., n=45: RMSSD $p=0.027$, HFnu $p=0.007$ [255]), worse orthostatic intolerance, impaired cognitive function, reduced exercise capacity, and poorer quality of life.

Low HRV is also an independent predictor of cardiovascular morbidity and mortality in other populations, raising concerns about long-term cardiovascular outcomes in ME/CFS.

Baroreflex Sensitivity

The baroreflex maintains blood pressure stability through rapid adjustments in heart rate and vascular tone. The NIH study found diminished baroreflex cardiovagal gain in ME/CFS patients [5], indicating impaired ability to modulate heart rate in response to blood pressure changes, reduced parasympathetic responsiveness, delayed cardiovascular adaptation to postural changes, and vulnerability to orthostatic stress.

Baroreflex Testing Methods Several methods assess baroreflex function. Spontaneous baroreflex analysis calculates the relationship between spontaneous blood pressure and R-R interval fluctuations. The Valsalva maneuver assesses heart rate and blood pressure responses to standardized straining. Neck suction or pressure directly stimulates carotid baroreceptors, while pharmacological methods use vasoactive drugs to manipulate blood pressure.

Evidence for Decreased Parasympathetic Activity

Multiple lines of evidence converge on parasympathetic (vagal) dysfunction as a central feature of ME/CFS autonomic abnormalities. Reduced HRV high-frequency power provides a direct measure of cardiac vagal modulation. Diminished baroreflex sensitivity, which is primarily mediated by vagal mechanisms, further supports this dysfunction. Pupillary abnormalities reveal altered pupil responses to light (parasympathetically mediated), while gastrointestinal dysmotility reflects vagal nerve dysregulation of gut function. Additionally, reduced respiratory sinus arrhythmia indicates impaired vagally mediated heart rate variation with breathing.

The NIH study explicitly concluded that the autonomic findings indicated “decreased parasympathetic activity” [5], providing a unifying explanation for many ME/CFS symptoms.

Sympathetic Nervous System Abnormalities

While parasympathetic dysfunction is prominent, sympathetic abnormalities also occur. Resting sympathetic overactivity manifests as elevated norepinephrine spillover and increased muscle sympathetic nerve activity. Despite this elevated baseline, sympathetic reactivity is impaired, showing blunted responses to stressors. Regional sympathetic dysfunction produces variable activation across different vascular beds, while catecholamine dysregulation affects synthesis, release, and clearance.

Reconciling central vs. peripheral norepinephrine: An apparent contradiction exists between reduced central (CNS) norepinephrine documented in CSF [5] and elevated peripheral norepinephrine spillover. This likely reflects compartmentalization: central noradrenergic

systems (locus coeruleus, brain norepinephrine) are separate from peripheral sympathetic nervous system activity. One proposed mechanism is that central deficiency could plausibly impair the brain's regulation of the sympathetic nervous system, leading to dysregulated peripheral sympathetic output—elevated at rest but unable to respond appropriately to challenges. This dissociation between central and peripheral catecholamine compartments is well-established in autonomic physiology.

The combination of elevated baseline sympathetic activity with reduced reactivity creates a rigid, poorly adaptive autonomic system unable to respond appropriately to physiological challenges.

8.2.2 Mechanisms of Orthostatic Intolerance

Orthostatic intolerance (OI) affects an estimated 70–90% of ME/CFS patients and manifests as postural orthostatic tachycardia syndrome (POTS), neurally mediated hypotension (NMH), orthostatic hypotension (OH), or combinations of these conditions.

Dysautonomia and POTS are components of the “Septad” framework of frequently co-occurring conditions in ME/CFS (Section 5.6.11). Small fiber neuropathy, another Septad component, may underlie autonomic dysfunction in a subset of patients, emphasizing the need for comprehensive evaluation of these interrelated pathophysiologies.

Blood Volume Abnormalities

Reduced blood volume is well-documented in ME/CFS and contributes to orthostatic intolerance [280]. Streeten and Bell documented that red blood cell mass was significantly reduced ($p < 0.001$) in 93.8% of female patients and 50% of male patients, with plasma volume subnormal in 52.6%. This total blood volume decrease compromises cardiovascular reserve through mechanisms possibly involving renin-angiotensin-aldosterone system dysfunction, reduced erythropoietin, or increased capillary permeability.

Hypovolemia reduces cardiac preload, compromising stroke volume and cardiac output, particularly during orthostatic stress.

Vascular Dysfunction

Multiple vascular abnormalities contribute to orthostatic intolerance. Impaired venoconstriction reduces the ability to mobilize venous blood during standing, leading to excessive venous pooling as blood accumulates in dependent vessels. Arterial dysregulation produces abnormal resistance vessel responses, while endothelial dysfunction impairs nitric oxide bioavailability.

Adrenergic Receptor Dysfunction

Abnormalities in adrenergic receptor function may explain some autonomic symptoms. Beta-adrenergic receptor autoantibodies have been identified in subsets of ME/CFS patients [71] and may either activate or block receptors. Loebel et al. found that 29.5% of ME/CFS patients (n=268) had elevated autoantibodies against β_2 , M3, and/or M4 receptors. Antibodies against β_2 adrenergic and M3 muscarinic receptors (both vasodilators) could explain vasoconstriction and hypoxemia observed in ME/CFS. Alpha-adrenergic abnormalities produce altered vasoconstrictor responses, while receptor desensitization may result from chronic catecholamine exposure downregulating receptors. Additionally, post-receptor signaling defects in G-protein coupling or second messenger systems may contribute to dysfunction.

Renin-Angiotensin-Aldosterone System

The RAAS regulates blood volume and pressure through sodium and water retention, vasoconstriction, and sympathetic activation.

Abnormalities in ME/CFS may include reduced aldosterone response to orthostatic stress, impaired renin secretion, altered angiotensin II sensitivity, and inappropriate natriuresis.

8.3 Peripheral Nervous System

8.3.1 Small Fiber Neuropathy

Small fiber neuropathy (SFN) affects thinly myelinated A-delta fibers and unmyelinated C fibers, which mediate pain, temperature, and autonomic functions. SFN has emerged as a significant finding in ME/CFS.

Skin Biopsy Findings

Punch skin biopsies with intraepidermal nerve fiber density (IENFD) measurement represent the gold standard for SFN diagnosis:

- **Reduced IENFD:** Multiple studies report decreased nerve fiber density in ME/CFS patients [281, 282]
- **Correlation with symptoms:** Lower IENFD correlates with pain severity and autonomic dysfunction
- **Distal predominance:** Typical length-dependent pattern with greater abnormalities in feet than thighs
- **Prevalence:** Estimates range from 30–60% of ME/CFS patients meeting criteria for SFN. Oaklander et al. [281] found 41% of fibromyalgia patients (overlapping with ME/CFS) had reduced IENFD diagnostic for SFN. A meta-analysis by Grayston et al. [282] reported 49% pooled prevalence (95% CI: 38–60%) of small fiber pathology in fibromyalgia across 8 studies

Autonomic Testing

Quantitative sudomotor axon reflex testing (QSART) and related methods assess small fiber autonomic function:

- **Reduced sweat output:** Indicating sudomotor dysfunction
- **Abnormal sweat gland innervation:** On skin biopsy analysis
- **Correlation with orthostatic intolerance:** SFN may contribute to autonomic dysregulation

Pain Mechanisms

SFN may explain chronic pain in ME/CFS through:

- **Neuropathic pain:** Burning, tingling, electric shock sensations
- **Allodynia:** Pain from normally non-painful stimuli
- **Hyperalgesia:** Exaggerated pain responses
- **Central sensitization:** Peripheral nerve damage may trigger central pain amplification

Potential Causes of SFN in ME/CFS

- Autoimmune mechanisms (ganglioside antibodies, sodium channel antibodies)
- Metabolic dysfunction (mitochondrial, oxidative stress)
- Chronic inflammation
- Microvascular abnormalities affecting nerve blood supply
- Direct viral damage (in post-infectious cases)

8.3.2 Nerve Conduction Studies

Electrophysiological Findings

Standard nerve conduction studies (NCS) assess large myelinated fiber function and are typically normal in ME/CFS, consistent with selective small fiber involvement. However, some studies report:

- Subtle prolongation of distal latencies
- Reduced compound muscle action potential amplitudes
- Abnormal F-wave parameters
- Changes suggesting subclinical demyelination

Implications

The contrast between abnormal small fiber findings and relatively preserved large fiber function suggests:

- Selective vulnerability of small fibers to ME/CFS pathophysiology
- Potential autoimmune targeting of specific nerve fiber populations
- Metabolic or oxidative stress preferentially affecting unmyelinated fibers
- Different pathophysiology from typical diabetic or inflammatory neuropathies

8.3.3 Treatment of Small Fiber Neuropathy

Management of SFN in ME/CFS requires addressing both symptomatic relief and underlying mechanisms. Treatment strategies must be adapted for ME/CFS-specific considerations including medication sensitivity and post-exertional malaise.

First-Line Neuropathic Pain Medications

Gabapentinoids. Gabapentin and pregabalin remain first-line treatments for neuropathic pain based on NeuPSIG guidelines [283]. Dosing recommendations below derive from these guidelines and clinical experience; individual titration is essential:

- **Gabapentin:** Start 100–300 mg at bedtime; titrate slowly to 900–3600 mg/day in divided doses
- **Pregabalin:** Start 25–75 mg at bedtime; titrate to 150–600 mg/day in divided doses
- **Mechanism:** Bind alpha-2-delta subunit of voltage-gated calcium channels, reducing excitatory neurotransmitter release
- **Benefits:** Also improve sleep quality and may reduce central sensitization
- **ME/CFS considerations:** Start at lower doses due to common medication sensitivity; sedation may help or hinder depending on individual sleep patterns

Serotonin-Norepinephrine Reuptake Inhibitors (SNRIs). Duloxetine has FDA approval for diabetic peripheral neuropathy [283]:

- **Duloxetine:** Start 20–30 mg daily; target 60 mg daily (range 30–120 mg)
- **Venlafaxine:** Alternative SNRI; 150–225 mg/day extended-release
- **Mechanism:** Enhance descending pain inhibition pathways via norepinephrine and serotonin
- **Additional benefits:** May help comorbid depression and fatigue in some patients
- **Cautions:** Discontinuation syndrome with abrupt cessation; may increase blood pressure

Tricyclic Antidepressants. Low-dose tricyclics provide analgesic effects independent of antidepressant action:

- **Amitriptyline:** Start 10 mg at bedtime; titrate to 25–75 mg (lower than antidepressant doses)
- **Nortriptyline:** Less sedating alternative; 10–75 mg at bedtime
- **Mechanism:** Block norepinephrine reuptake, sodium channels, and NMDA receptors
- **Benefits:** Improve sleep architecture; long clinical experience
- **Cautions:** Anticholinergic effects (dry mouth, constipation, urinary retention); cardiac effects at higher doses; morning sedation

Topical Treatments

Topical agents provide targeted relief with minimal systemic effects—particularly valuable in medication-sensitive ME/CFS patients:

Lidocaine.

- **5% lidocaine patches:** Apply to painful areas for up to 12 hours daily
- **Mechanism:** Blocks sodium channels in peripheral nerves, reducing ectopic firing
- **Advantages:** Minimal systemic absorption; can be cut to size; well-tolerated
- **Limitations:** Localized effect only; works best for focal pain

Capsaicin.

- **Low-concentration cream (0.025–0.075%):** Apply 3–4 times daily
- **High-concentration patch (8%):** Single application by healthcare provider; effects last 3 months
- **Mechanism:** Depletes substance P from peripheral nerve endings; defunctionalizes TRPV1-expressing nociceptors
- **Cautions:** Initial burning sensation (usually diminishes with regular use); avoid mucous membranes and eyes

Treatment of Underlying Causes

Autoimmune SFN. When SFN has an autoimmune etiology (suggested by anti-ganglioside or anti-sodium channel antibodies), immunomodulation may be beneficial [284]:

- **IVIG:** 0.4 g/kg/day for 5 days, then monthly maintenance; case series evidence (low certainty) suggests improvement in pain and autonomic symptoms in autoimmune SFN, though RCT data are lacking [285]
- **Corticosteroids:** Short courses for acute flares; long-term use limited by side effects
- **Other immunomodulators:** Rituximab, azathioprine, mycophenolate in refractory cases

- **ME/CFS relevance:** Given autoimmune hypotheses in ME/CFS, autoimmune SFN testing should be considered in patients with prominent neuropathic features

⚠ **Limitation 4: Small Fiber Neuropathy: Prevalence Extrapolated from Fibromyalgia**

SFN prevalence estimates in ME/CFS (30–60%) derive largely from fibromyalgia cohorts with assumed overlap rather than ME/CFS-specific studies. Key data gaps:

- The most-cited prevalence data (Oaklander 2013, Grayston 2019 meta-analysis) studied fibromyalgia patients, not ME/CFS cohorts selected by ME/CFS diagnostic criteria.
- No large-scale study has determined IENFD in a well-characterised ME/CFS cohort with appropriate controls.
- Treatment recommendations (alpha-lipoic acid, acetyl-L-carnitine, IVIG for autoimmune SFN) are extrapolated from diabetic neuropathy or case series; none have been tested in ME/CFS-specific trials.
- The causal relationship between SFN and ME/CFS symptoms is unestablished—SFN may be an independent comorbidity rather than a mechanistic contributor.

Metabolic and Nutritional Support. Several supplements may support nerve regeneration. Note that evidence derives primarily from diabetic neuropathy populations; efficacy in ME/CFS-associated SFN has not been specifically studied:

- **Alpha-lipoic acid:** 600–1800 mg daily; demonstrated efficacy in diabetic neuropathy RCTs [286]; antioxidant and mitochondrial cofactor
- **Acetyl-L-carnitine:** 1500–3000 mg daily; supports neuronal energy metabolism; RCT evidence in diabetic neuropathy showing improved pain and nerve regeneration [287]
- **B vitamins:** B12 (methylcobalamin 1000–5000 mcg), B6 (avoid excess >100 mg/day, which can cause neuropathy), B1 (benfotiamine 300–600 mg)
- **Mechanism:** Support mitochondrial function, reduce oxidative stress, provide nerve membrane substrates

⚠ **Warning 6: Vitamin B6 Toxicity**

While B6 deficiency can cause neuropathy, excess pyridoxine supplementation (typically >200 mg/day chronically) can paradoxically cause a sensory neuropathy. Patients should not exceed 100 mg/day without medical supervision, and B6 levels should be checked if neuropathy worsens with supplementation.

ME/CFS-Specific Considerations

Treatment of SFN in ME/CFS requires adaptation for this population:

- **Start low, go slow:** Begin at 25–50% of typical starting doses due to medication sensitivity

- **Single changes:** Add or adjust one medication at a time to identify responses
- **Sedation balance:** Sedating medications (gabapentin, amitriptyline) may help sleep but worsen daytime fatigue
- **Autonomic effects:** Many neuropathic pain medications affect autonomic function; monitor orthostatic symptoms
- **PEM awareness:** Exercise-based therapies sometimes recommended for neuropathy are contraindicated in ME/CFS due to PEM risk
- **Topical preference:** Consider topical agents first given lower systemic burden

Treatment Algorithm

The following algorithm represents a proposed approach synthesized from NeuPSIG guidelines [283] and clinical experience with ME/CFS patients. It is not a validated clinical guideline:

1. **Diagnosis confirmation:** Skin biopsy for IENFD; autonomic testing; screen for treatable causes (diabetes, B12 deficiency, autoantibodies)
2. **Address underlying causes:** Treat autoimmune SFN with immunomodulation; correct nutritional deficiencies
3. **First-line symptomatic:** Topical lidocaine for focal pain; low-dose gabapentinoid or TCA at bedtime
4. **Second-line:** Add SNRI if inadequate response; consider combination therapy (e.g., gabapentinoid + TCA)
5. **Adjunctive support:** Alpha-lipoic acid, acetyl-L-carnitine for neuroprotection (extrapolated from diabetic neuropathy evidence)
6. **Refractory cases:** Pain medicine referral; interventional options; IVIG trial if autoimmune markers present

? Open Question 2: SFN Reversibility in ME/CFS

Can small fiber neuropathy in ME/CFS patients be reversed with appropriate treatment? Case reports suggest IENFD can normalize after treating underlying conditions (e.g., autoimmune SFN with IVIG, diabetic SFN with glucose control). If ME/CFS-associated SFN has an autoimmune or inflammatory basis, early immunomodulation might prevent permanent nerve damage. Longitudinal studies with serial skin biopsies in treated patients would clarify whether nerve regeneration is achievable.

8.4 Blood-Brain Barrier Dysfunction

The blood-brain barrier (BBB) normally restricts entry of cells, pathogens, and molecules from the bloodstream into the brain parenchyma. BBB dysfunction may contribute to neuroinflammation and neurological symptoms in ME/CFS.

8.4.1 Evidence for Permeability Changes

- **CSF/serum albumin ratio:** Elevated in some ME/CFS patients, indicating increased permeability
- **Neuroimaging markers:** Subtle gadolinium enhancement suggesting leakage
- **Peripheral inflammatory markers in CSF:** Cytokines and chemokines crossing the barrier
- **Autoantibodies in CNS:** Entry of pathogenic antibodies

8.4.2 Consequences for Neuroinflammation

BBB dysfunction permits:

- **Peripheral immune cell infiltration:** T cells, monocytes entering brain tissue
- **Cytokine entry:** Peripheral inflammatory mediators reaching the CNS
- **Autoantibody access:** Receptor-targeting antibodies affecting neural function
- **Pathogen penetration:** Viral particles or antigens entering the brain

8.4.3 Transport Dysfunction

Beyond passive permeability, active transport systems at the BBB may be dysfunctional:

- **Glucose transporters:** Potentially explaining cerebral hypometabolism
- **Amino acid transporters:** Affecting neurotransmitter precursor availability
- **Drug efflux pumps:** Altering CNS drug concentrations
- **Receptor-mediated transcytosis:** Impaired transport of essential molecules

8.4.4 Blood-Brain Barrier as CNS Vulnerability Factor

While the preceding subsections address BBB permeability and transport dysfunction from the perspective of what enters or exits the CNS, the BBB may create *unique vulnerability* for CNS tissues in ME/CFS through mechanisms that paradoxically stem from the barrier's protective function.

Speculation 13 (Blood-Brain Barrier Vulnerability Hypothesis). **Certainty: 0.35.**

The blood-brain barrier creates CNS-specific vulnerability in ME/CFS through three converging mechanisms: (1) trapping damage signals that trigger persistent neuroinflammation, (2) limiting access to mitochondrial cofactors needed for repair, and (3) preventing the rapid mitochondrial turnover possible in peripheral dividing cells.

Mechanism 1: Trapping neuroinflammatory signals. Mitochondrial dysfunction leads to mitochondrial DNA (mtDNA) leakage into the cytoplasm, activating the cGAS-STING pathway and triggering type I interferon and pro-inflammatory cytokine production [288]. In peripheral tissues, the resulting inflammation can be resolved through immune cell infiltration and clearance. In the CNS, however, the BBB restricts immune cell access. When neuronal or glial mtDNA activates cGAS-STING signaling in microglia and astrocytes, the resulting neuroinflammation becomes *trapped*—peripheral immune cells that might otherwise regulate or resolve the inflammation cannot readily cross the barrier. This may explain the persistent microglial activation documented in ME/CFS PET studies [73]: once initiated by mtDNA leak, neuroinflammation perpetuates because the BBB prevents clearance mechanisms available to peripheral tissues.

Mechanism 2: Limited cofactor access for mitochondrial repair. Mitochondrial function requires continuous supply of cofactors (CoQ10, NAD⁺/NADH, B vitamins), which reach peripheral tissues relatively easily via systemic circulation but face BBB transport limitations. CoQ10 can cross the BBB via SR-B1 and RAGE receptors but is simultaneously effluxed back to blood via LRP-1/LDLR receptors, creating net-negative brain uptake in many conditions [289]. Only methylcobalamin (the active B12 form) crosses without biotransformation via specific cubam receptors [290]; the common supplement form cyanocobalamin requires conversion before CNS entry. NAD⁺ has a short half-life (1–2 hours) and limited BBB penetration [291]. The consequence: oral supplementation that improves peripheral mitochondrial function may have minimal CNS effects.

Mechanism 3: Constrained mitochondrial turnover in post-mitotic cells. Neurons do not divide after development. Brain synaptic mitochondrial proteins have a median half-life of 25.7 days versus hepatic mitochondrial proteins at 3.5 days—a 7-fold difference [292]. Damaged mitochondria at distal synapses must travel potentially meters back to the soma for mitophagy, and may accumulate dysfunction where energy demand is highest. In contrast, peripheral dividing cells replace entire cells every few days, diluting mitochondrial damage across generations. CNS mitochondria thus accumulate damage *faster* than they can be repaired or replaced.

Convergence. These three mechanisms converge: mitochondrial dysfunction leads to mtDNA leak; mtDNA activates cGAS-STING neuroinflammation; the BBB prevents immune clearance so inflammation persists; the BBB limits cofactor supply so repair is constrained; and the neuronal post-mitotic state prevents turnover from diluting damage. The result is progressive CNS dysfunction even when peripheral tissues stabilize—explaining why cognitive symptoms may persist despite improved muscle function or reduced systemic inflammation.

Testable predictions.

1. CSF should show higher concentrations of mitochondrial DAMPs (mtDNA fragments, 8-OHdG) relative to blood, reflecting faster CNS damage accumulation

2. BBB-penetrant supplement forms (methylcobalamin, liposomal CoQ10, intranasal NAD⁺) should improve cognitive symptoms more than standard forms at equivalent doses
3. Intranasal delivery of mitochondrial-targeted compounds should show cognitive benefits where oral forms do not, as intranasal delivery bypasses the BBB via olfactory and trigeminal pathways [291]
4. Patients with higher CSF/serum albumin ratios (greater BBB permeability [293]) should paradoxically show *lower* neuroinflammation, as increased permeability allows partial immune clearance
5. CSF should show elevated STING pathway activation markers (cGAMP, phospho-TBK1) correlating with cognitive symptom severity

Treatment implications. If this hypothesis is correct, therapeutic strategies should prioritize: (1) BBB-penetrant cofactor formulations (methylcobalamin over cyanocobalamin, liposomal CoQ10, BBB-penetrant antioxidants such as MitoQ or idebenone); (2) intranasal delivery of NAD⁺, NMN, or glutathione; (3) cGAS-STING pathway inhibition to reduce trapped neuroinflammation; (4) mitophagy-enhancing compounds (urolithin A, spermidine) to help neurons clear damaged mitochondria.

Limitations. This hypothesis has certainty 0.35. It synthesizes mechanisms documented in neurodegenerative diseases and applies them to ME/CFS by analogy—no study has directly measured cGAS-STING activation, CSF mtDNA levels, or mitochondrial turnover rates in ME/CFS patients. BBB permeability is heterogeneous across ME/CFS patients [293], and if some patients have highly permeable BBB, the trapping mechanism may not apply universally. Genetic polymorphisms in BBB transporters may create substantial patient-to-patient variability. Despite these limitations, the hypothesis is testable and generates specific predictions evaluable in cohorts with CSF access.

8.5 Cerebral Blood Flow Abnormalities

Cerebral blood flow (CBF) abnormalities are among the most consistently documented findings in ME/CFS and likely contribute substantially to cognitive symptoms.


Figures 8.5 and 8.6 illustrate how multiple mechanisms reduce cerebral blood flow in ME/CFS (30–40 mL/100g/min vs. normal 50–60 mL/100g/min, a 40% reduction).

8.5.1 Reduced Regional Blood Flow

Multiple neuroimaging modalities have demonstrated CBF reductions [123]:

- **Global hypoperfusion:** 10–20% reduction in total cerebral blood flow (measured by SPECT and Doppler ultrasound)
- **Regional deficits:** Particularly in frontal, temporal, and parietal regions

Normal Cerebral Blood Flow



Key points: The brain requires constant, high blood flow due to its enormous metabolic demands. Autoregulation maintains stable perfusion across a wide range of blood pressures. Adequate O₂ and glucose delivery, plus CO₂/lactate removal,

Figure 8.5: Normal cerebral blood flow regulation meeting brain metabolic demands.




Figure 8.6: ME/CFS cerebral hypoperfusion cascade causing cognitive dysfunction.

- **Brainstem hypoperfusion:** Potentially explaining autonomic dysfunction [294]
- **Subcortical abnormalities:** Basal ganglia and thalamic hypoperfusion

Van Campen et al. [123] documented that 90% of ME/CFS patients (n=429) showed abnormal CBF reduction (>13%) during head-up tilt testing, with end-tilt CBF reduction of 26% in ME/CFS patients versus only 7% in controls (n=44). Importantly, this occurred even in the absence of hypotension or tachycardia, indicating intrinsic cerebrovascular dysfunction rather than solely cardiovascular causes.

8.5.2 Correlation with Cognitive Symptoms

CBF reductions correlate with specific cognitive deficits:

- Frontal hypoperfusion → executive dysfunction, working memory impairment
- Temporal hypoperfusion → verbal memory deficits, language processing difficulties
- Parietal hypoperfusion → attention deficits, spatial processing impairment
- Global hypoperfusion → processing speed reduction, mental fatigue

8.5.3 Mechanisms of Cerebral Hypoperfusion

The cerebral hypoperfusion documented above likely results from multiple converging mechanisms:

- **Reduced cardiac output:** Secondary to autonomic dysfunction (Section 8.2) and blood volume deficits [280]
- **Impaired cerebral autoregulation:** Inability to maintain CBF across blood pressure changes [294]
- **Endothelial dysfunction:** Reduced nitric oxide-mediated vasodilation
- **Increased cerebrovascular resistance:** Vasoconstriction or structural changes
- **Neurovascular uncoupling:** Failure of blood flow to match metabolic demand

The integration of autonomic dysfunction, reduced blood volume, and direct cerebrovascular pathology creates a multifactorial reduction in brain perfusion that correlates with cognitive symptom severity. This multifactorial integration is characteristic of ME/CFS pathophysiology and is discussed in the context of multi-system interactions in Chapter 13, Section 13.3.

8.5.4 Exacerbation with Exertion

Importantly, cerebral perfusion abnormalities worsen following physical or cognitive exertion:

- Further CBF reductions post-exercise
- Prolonged recovery of normal perfusion
- Correlation with post-exertional malaise severity

- Potential contribution to cognitive “crashes” following activity

8.6 Auditory Processing Dysfunction and Tinnitus

Auditory symptoms represent an underrecognized but significant neurological manifestation of ME/CFS, with convergent evidence from functional, epidemiological, systematic, and anatomical studies establishing auditory dysfunction as a documented feature of the disease.

8.6.1 Prevalence and Epidemiology

★ Achievement 4: Tinnitus-ME/CFS Epidemiological Association

Schubert et al. [295] provided the first large-scale epidemiological evidence linking ME/CFS to tinnitus in a population-based cohort of 124,609 individuals from the Dutch Lifelines study. ME/CFS patients demonstrated 1.57 times higher odds (OR 1.568, $p < 0.05$) of experiencing constant tinnitus compared to healthy controls, identifying ME/CFS as a novel disease associate for tinnitus beyond traditional audiological causes such as noise exposure, age-related hearing loss, and cardiovascular disease.

This finding aligns with earlier cohort studies and patient surveys reporting tinnitus prevalence ranging from 48% to 78% in ME/CFS patients, substantially higher than the 10–15% prevalence in the general population.

★ Achievement 5: Systematic Evidence for Auditory Dysfunction

A 2024 systematic review by Skare et al. [296] synthesized evidence from 172 articles (1990–2024) documenting ear abnormalities across ME/CFS, fibromyalgia, long-COVID syndrome, postural orthostatic tachycardia syndrome (PoTS), and related conditions. The review identified cochlear complaints—including tinnitus, hearing loss, and hyperacusis—as the most frequent auditory findings in ME/CFS. Four pathophysiological mechanisms were proposed: (1) viral effects on cochlear or central auditory structures, (2) vascular impairment reducing blood flow to the cochlea and brainstem, (3) autoimmune reactions targeting inner ear antigens, and (4) oxidative stress damaging cochlear hair cells and auditory neurons.

The systematic review recommended that all ME/CFS patients with audiological complaints receive ENT consultation and formal audiometry to assess the nature and severity of auditory dysfunction.

8.6.2 Functional Auditory Processing Deficits

Beyond subjective tinnitus complaints, objective evidence demonstrates specific auditory processing impairments in ME/CFS patients.

★ Achievement 6: Selective Auditory Processing Impairment

Johnson et al. [297] demonstrated modality-specific cognitive impairment in a controlled comparison of 20 CFS patients, 20 multiple sclerosis (MS) patients, and 20 healthy controls. CFS patients showed differential impairment on auditory versus visual processing tasks, while MS patients showed equal impairment on both modalities. This pattern suggests specific dysfunction in central auditory pathways rather than general cognitive slowing, distinguishing the ME/CFS cognitive profile from the more global impairment observed in other neurological conditions.

Functional MRI studies have further documented that CFS patients recruit additional or atypical brain regions during cognitive and sensory tasks compared to controls [298], requiring greater neural resources to achieve equivalent task performance. This pattern of compensatory over-recruitment reflects inefficient neural processing consistent with the broader energy-limitation framework.

8.6.3 Neuroanatomical Substrate: Brainstem Dysfunction

The functional auditory deficits and elevated tinnitus prevalence in ME/CFS are explained by documented structural and functional abnormalities in brainstem regions critical for auditory processing.

★ Achievement 7: Brainstem Structural Abnormalities

Nelson et al. [299] synthesized MRI evidence from 11 studies demonstrating structural and functional brainstem abnormalities in ME/CFS patients. The brainstem contains the primary central auditory pathway structures:

- **Cochlear nucleus** (medulla) — receives input from cochlear nerve; first central processing station
- **Superior olivary complex** (pons) — sound localization via interaural time and intensity differences
- **Lateral lemniscus** — ascending auditory pathway connecting lower and upper brainstem
- **Inferior colliculus** (midbrain) — integration of ascending auditory information before thalamic relay

Dysfunction in these structures provides a neuroanatomical substrate explaining both the auditory processing deficits documented by Johnson et al. [297] and the increased tinnitus prevalence observed by Schubert et al. [295].

Importantly, brainstem abnormalities in ME/CFS extend beyond auditory pathways to include autonomic control centers (see Section 8.2), arousal systems (locus coeruleus), and sensory integration regions. This explains the co-occurrence of auditory symptoms with autonomic dysfunction, sleep disturbances, and sensory hypersensitivity—all manifestations of brainstem pathology.

8.6.4 Central vs. Peripheral Auditory Pathology

Observation 49 (Central vs. Peripheral Auditory Pathology). The convergence of functional deficits [297], population-level tinnitus prevalence [295], systematic evidence [296], and brainstem MRI abnormalities [299] suggests predominantly **central (brainstem)** rather than peripheral (cochlear) auditory pathology in ME/CFS. This distinction has important therapeutic implications: neurological approaches targeting brainstem dysfunction, cerebral perfusion, and neuroinflammation may be more effective than peripheral ENT interventions focused solely on the cochlea or middle ear.

Evidence supporting central over peripheral pathology includes:

- Auditory processing deficits may occur without peripheral hearing loss on audiometry
- Tinnitus severity often fluctuates with orthostatic stress and cerebral hypoperfusion
- Auditory symptoms co-occur with other brainstem-mediated dysfunction (autonomic, arousal, sensory)
- Hyperacusis (sound sensitivity) suggests central gain dysregulation rather than peripheral damage
- Auditory symptoms are part of broader post-exertional malaise rather than isolated ear pathology

8.6.5 Proposed Mechanisms

Based on the systematic review by Skare et al. [296] and integration with established ME/CFS pathophysiology, four mechanisms likely contribute to auditory dysfunction (this exemplifies the multi-mechanism pattern characteristic of ME/CFS symptoms, as discussed in Chapter 13):

Viral Effects. Direct viral damage to cochlear structures or central auditory pathways may occur during acute infection. This is particularly relevant for post-infectious ME/CFS onset, where viral neurotropism (e.g., EBV, HHV-6) could affect brainstem auditory nuclei. Acute onset of tinnitus following infection supports this mechanism.

Vascular Impairment. Reduced cerebral blood flow documented in ME/CFS (Section 8.5) likely affects the highly vascularized cochlea and brainstem auditory centers. The stria vascularis in the cochlea maintains the ionic gradient essential for sound transduction and is metabolically active, making it vulnerable to hypoperfusion. Fluctuating tinnitus severity correlating with orthostatic stress supports vascular involvement.

Autoimmune Reactions. Antibodies targeting inner ear antigens (anti-cochlin, anti-HSP70) or auditory brainstem structures may produce autoimmune inner ear disease (AIED). This mechanism aligns with broader autoimmune theories of ME/CFS and suggests potential benefit from immunomodulatory treatment in select patients.

Oxidative Stress. Reactive oxygen species generated by mitochondrial dysfunction and neuroinflammation can damage cochlear hair cells and auditory neurons. The cochlea has high metabolic demands and limited antioxidant capacity, making it vulnerable to oxidative damage. This mechanism connects auditory dysfunction to the mitochondrial pathology documented in ME/CFS.

8.6.6 Clinical Implications

Assessment Recommendations

Based on the documented prevalence and clinical significance of auditory dysfunction:

1. **Routine screening:** All ME/CFS patients should be screened for tinnitus, hearing loss, hyperacusis, and auditory processing difficulties
2. **Formal audiometry:** Patients reporting auditory symptoms should receive comprehensive audiological evaluation
3. **ENT consultation:** Rule out treatable peripheral causes (cerumen impaction, otosclerosis, Ménière's disease)
4. **Central auditory testing:** Consider auditory brainstem response (ABR) testing to assess central pathways
5. **Correlation with ME/CFS severity:** Document whether auditory symptoms fluctuate with overall disease activity, orthostatic stress, and post-exertional malaise

Treatment Considerations

Given the proposed central pathology:

- **Address underlying ME/CFS pathophysiology:** Optimize cerebral perfusion (salt/fluid loading for orthostatic intolerance), treat neuroinflammation, support mitochondrial function
- **Symptomatic management:** Sound therapy (white noise, tinnitus masking), cognitive-behavioral therapy for tinnitus distress (distinct from CBT as ME/CFS treatment)
- **Avoid ototoxic medications:** Many drugs can worsen tinnitus (aminoglycosides, loop diuretics, high-dose aspirin, certain chemotherapies)
- **Consider immunomodulation:** In patients with evidence of autoimmune component (autoantibodies, inflammatory markers)
- **Antioxidant support:** Alpha-lipoic acid, CoQ10, N-acetylcysteine (extrapolated from evidence in age-related hearing loss and noise-induced damage)

△ Warning 7: Tinnitus as PEM Symptom

Many ME/CFS patients report that tinnitus intensity increases during post-exertional malaise or correlates with fatigue severity. This pattern suggests tinnitus may function

as a real-time indicator of energy depletion or cerebral hypoperfusion. Patients should be educated to recognize worsening tinnitus as a potential warning sign to rest and avoid further exertion.

8.6.7 Research Gaps

Despite the convergent evidence for auditory dysfunction in ME/CFS, significant gaps remain:

- **Causality:** Cross-sectional designs cannot determine whether ME/CFS causes auditory dysfunction, auditory dysfunction contributes to ME/CFS symptoms, or a common mechanism produces both
- **Subtype correlation:** Unknown whether auditory symptoms predict specific ME/CFS subgroups or correlate with particular biomarkers
- **Treatment trials:** No randomized controlled trials of auditory-targeted interventions in ME/CFS populations
- **Mechanism validation:** The four proposed mechanisms (viral, vascular, autoimmune, oxidative) require experimental validation
- **Reversibility:** Unknown whether treating underlying ME/CFS pathophysiology can reverse auditory dysfunction
- **Longitudinal trajectory:** Natural history of auditory symptoms in ME/CFS not systematically documented

? Open Question 3: Central Auditory Gain and Sensory Hypersensitivity

Hyperacusis (sound sensitivity) in ME/CFS may reflect dysregulated central gain in auditory processing pathways. The brainstem and auditory cortex normally adjust sensitivity (gain) based on environmental demands and context. In ME/CFS, chronic neuroinflammation, altered neurotransmitter levels, or thalamic dysfunction may inappropriately increase central auditory gain, amplifying all sounds and making normal environmental noise intolerable. This would parallel central sensitization in pain pathways. Testing this hypothesis with objective measures of auditory gain (acoustic reflex thresholds, loudness discomfort levels, auditory brainstem response) could clarify mechanisms and guide treatment targeting central gain normalization rather than peripheral protection.

8.7 Cognitive Dysfunction: Clinical Manifestations

The neurological abnormalities described above manifest clinically as characteristic patterns of cognitive dysfunction, often described by patients as “brain fog.”

8.7.1 Domains of Impairment

Processing Speed

Slowed information processing is perhaps the most consistent cognitive finding, manifesting as delayed reaction times, slower performance on timed tasks, reduced ability to keep up with rapid conversations, and difficulty with time-pressured activities.

Attention and Concentration

Attention and concentration deficits include difficulty sustaining attention, easy distractibility, impaired divided attention (multitasking), and reduced attentional capacity under stress.

Memory

Memory impairments encompass working memory deficits (holding information “online”), impaired short-term memory encoding, word-finding difficulties, and variable long-term memory retrieval.

~ Hypothesis 8: Memory Triage Consequence

Certainty: 0.55.

Memory encoding is substantially more energy-expensive than memory retrieval, predicting that ME/CFS patients should show disproportionate encoding deficits relative to retrieval impairment—a pattern consistent with CNS energy triage.

Differential energy costs of memory operations. Hippocampal memory encoding requires long-term potentiation (LTP), involving NMDA receptor activation, calcium-dependent signaling cascades, new protein synthesis, and structural synaptic remodeling [300]. These processes are metabolically demanding: encoding a new memory trace requires de novo gene expression, dendritic spine growth, and synaptic protein trafficking. Retrieval, by contrast, reactivates existing synaptic patterns through pattern completion in CA3 networks—a process that uses established circuits without requiring new protein synthesis or structural modification [301].

Quantitative estimates suggest that LTP-associated protein synthesis increases local energy consumption by 30–50% above baseline in hippocampal neurons, whereas pattern completion during retrieval operates within normal metabolic parameters. Working memory maintenance in prefrontal cortex similarly requires sustained neuronal firing against inhibitory currents, creating continuous metabolic demand proportional to the number of items held in mind [302].

Predicted pattern in ME/CFS. If CNS energy is limited, the brain should sacrifice high-cost encoding operations before low-cost retrieval operations—a “memory triage.” This predicts:

1. **Encoding > retrieval impairment:** Patients should show greater difficulty forming new memories than accessing old ones. Standardized testing should reveal disproportionate deficits on encoding-dependent tasks (learning new word lists, forming new associations) relative to recognition or cued recall of previously encoded material
2. **Working memory > long-term retrieval:** Sustained prefrontal firing for working memory maintenance is metabolically costly; retrieving consolidated long-term memories from distributed cortical stores is less so
3. **Encoding degrades with exertion:** During PEM, when CNS energy deficits intensify, new memory formation should decline more steeply than the ability to recall previously consolidated information
4. **Context-dependent encoding failure:** Encoding in metabolically demanding contexts (noisy environments, multitasking, social interaction) should fail preferentially, as these conditions compete for the limited energy budget

Supporting evidence. The meta-analysis by Sebaiti et al. [303] documents memory impairment in ME/CFS with moderate effect sizes ($g = -0.55$ to -0.67), but existing studies have not systematically separated encoding from retrieval. Clinical observation consistently reports that ME/CFS patients struggle more with forming new memories (“I can’t take in new information”) than with accessing established knowledge (“I remember things from before I got sick”). Patients frequently describe intact recognition (“I know I’ve seen this before”) with impaired free recall of recently encountered material—precisely the pattern predicted by encoding-selective energy limitation.

Treatment implications. If encoding is selectively impaired by energy limitation, compensatory strategies should emphasize: (1) reducing encoding load through external memory aids (notes, recordings, photographs) rather than relying on internal encoding; (2) scheduling new learning during peak energy windows; (3) using spaced repetition to distribute encoding costs across multiple low-demand sessions; (4) leveraging recognition over recall (multiple-choice formats, visual cues) when possible.

Limitations. This hypothesis has certainty 0.55. The differential energy cost of encoding versus retrieval is well established in neuroscience, but the specific prediction of disproportionate encoding impairment in ME/CFS awaits formal testing with paradigms designed to isolate encoding from retrieval. Existing neuropsychological batteries typically conflate encoding and retrieval in composite “memory” scores. Confounds include attention deficits (which impair encoding indirectly), medication effects, and sleep disruption (which impairs memory consolidation independently of encoding).

Executive Function

Executive function deficits present as planning and organization difficulties, impaired cognitive flexibility, reduced problem-solving ability, and difficulty with complex decision-making.

8.7.2 Social and Emotional Dysfunction

While less frequently discussed in clinical literature, social and emotional impairments represent significant sources of disability in ME/CFS and are direct consequences of the neurometabolic dysfunction documented above.

⚠ **Limitation 5: Social-Emotional Phenomenology: Clinical Observation, Not Empirical Research**

The social disability descriptions in this section (social interaction as painful exertion, flat affect, relationship conflict as insurmountable barrier, environmental control as survival mechanism) are based on clinical observation and patient self-report. No published study has validated “social exhaustion,” “affective energy expenditure,” or “interpersonal metabolic cost” as measurable constructs in ME/CFS. The proposed neurobiological explanations (catecholamine depletion → social withdrawal) are mechanistically plausible but represent inference, not demonstrated cause-and-effect. Alternative explanations—comorbid depression, social anxiety disorder, learned avoidance—have not been formally excluded in the populations described.

Note on evidence base: The detailed phenomenology described in this section is based primarily on extensive clinical observation and patient reports rather than systematic empirical research. While the underlying neurobiological mechanisms (catecholamine depletion, prefrontal hypometabolism, TPJ dysfunction) are well-documented [5], the specific social and emotional manifestations described below await formal validation through patient surveys, qualitative research, and prospective studies. This section should be considered a synthesis of clinical observation with established neuroscience, not yet a body of peer-reviewed ME/CFS-specific research on social disability.

Social Interaction as Metabolically Demanding Activity

Social interaction requires the simultaneous coordination of multiple high-energy cognitive and neurological processes:

- **Language processing and production:** Real-time comprehension, response formulation, word retrieval, and articulation
- **Working memory load:** Tracking conversational context, remembering prior statements, maintaining coherent narrative threads
- **Executive function demands:** Monitoring social cues, adjusting behavior in real-time, inhibiting inappropriate responses
- **Sensory integration:** Simultaneous processing of facial expressions, vocal prosody, body language, and environmental context
- **Motor control for affect generation:** Voluntary and involuntary facial expressions, eye contact, postural adjustments, vocal modulation
- **Reward system engagement:** Dopamine-mediated reward processing that makes social interaction inherently reinforcing in healthy individuals

When ATP production is impaired and catecholamine levels are low (as documented in the NIH study [5]), these processes cannot be sustained. The brain experiences social demands as it would physical exertion beyond capacity: as painful, threatening, something to avoid.

Clinical Presentation: Social Interaction as Painful Exertion

Many ME/CFS patients report that social interaction feels actively *painful* rather than merely tiring:

- Subjective experience identical to being forced to perform physical exercise while exhausted
- Approach characterized by “minimize the pain”—engage only as much as absolutely necessary
- Absence of enjoyment or reward, even in interactions that would previously have been pleasurable
- Duration often measured in minutes before exhaustion becomes overwhelming
- Post-social crashes (cognitive and physical PEM) lasting hours to days

This pattern may persist for decades and often predates formal ME/CFS diagnosis, suggesting it reflects fundamental metabolic limitations rather than secondary depression or psychological withdrawal.

Flat Affect and Energy Conservation

Generating and displaying emotional affect is metabolically expensive:

- **Muscular activation:** Smiling, animated facial expressions, and expressive body language require continuous motor control
- **Neurochemical substrates:** Emotional expression requires adequate dopamine for motivation and reward signaling
- **Prefrontal-limbic coordination:** Generating contextually appropriate affect requires coordination between multiple brain regions

When energy is scarce, the brain prioritizes survival functions over social signaling. The result is observable flat affect—patients appear emotionally unexpressive, disengaged, or “unhappy” even when not experiencing negative emotion. This is **not** conscious suppression or masking; it reflects genuine inability to generate the energetic and neurochemical processes required for emotional expression.

Interpersonal Consequences and Misattribution

The combination of social withdrawal and flat affect creates predictable interpersonal difficulties:

- **Misinterpretation as contempt or disinterest:** Observers lacking context for the patient's energy deficit often interpret flat affect and minimal engagement as disdain, superiority, or lack of care
- **Relationship damage:** Colleagues, friends, and family members feel rejected, judged, or dismissed when the actual issue is metabolic incapacity
- **Emotional contagion:** Others interacting with ME/CFS patients often become unhappy or uncomfortable themselves, unable to understand the patient's apparent lack of positive affect
- **Inability to explain:** The exhaustion that prevents social engagement also impairs the cognitive and communication capacity needed to explain the problem ("explaining why I'm too tired to talk requires energy to talk")
- **Vicious cycle:** Negative reactions from others increase the stress and energy demand of social interaction, further reducing capacity

Patients are frequently blamed for "attitude problems," "not trying," or "not caring" when the actual issue is neurometabolic failure to generate expected social signals.

The Communication Double-Bind

ME/CFS patients face an impossible situation regarding social interaction:

1. Employment and relationships require communication and social engagement
2. Communication and social engagement are painfully exhausting and worsen symptoms
3. Avoiding social interaction damages relationships and is misinterpreted as contempt
4. Explaining the difficulty requires the very communication capacity that is depleted
5. There is no winning strategy—only choices between different types of harm

Relationship Conflict as Insurmountable Barrier

The energy deficit affecting social interaction becomes critically limiting when relationships encounter even minor conflict or tension:

- **Conflict management requires peak cognitive resources:** Navigating disagreements, processing emotions, formulating diplomatic responses, regulating one's own reactions, and sustaining conversation through discomfort all require executive function, emotional regulation, and sustained attention—precisely the capacities most impaired in ME/CFS
- **Minor conflicts become insurmountable:** What healthy individuals would consider trivial relationship friction (scheduling disagreements, differing preferences, minor miscommunications) becomes *impossibly difficult to manage* when cognitive and emotional resources are depleted
- **Relationship attrition:** Friendships require ongoing maintenance, occasional conflict resolution, and emotional investment. When any conflict—however minor—exceeds available energy, relationships deteriorate and are eventually abandoned

- **Selection for low-maintenance relationships only:** Only relationships requiring absolutely minimal effort, zero conflict, and no emotional complexity can be sustained. This severely restricts social connection to a vanishingly small subset of potential relationships
- **Inability to repair:** Even when patients recognize that a relationship is worth preserving, they lack the energy to engage in the repair conversations necessary to resolve issues. The relationship fails not from lack of desire but from metabolic inability to execute repair
- **Compounding isolation:** As relationships with any degree of complexity or occasional friction are abandoned due to inability to manage conflict, social networks contract to near-zero. Patients become profoundly isolated not from preference but from inability to meet the basic energy demands of relationship maintenance
- **Loss of deep connections:** The inability to engage seriously in friendship—to invest emotional energy, navigate normal ups and downs, work through misunderstandings—means that only the most superficial relationships can survive. Patients lose access to the deep, meaningful connections that require tolerance for occasional difficulty
- **Present but disengaged:** Even when patients are physically able to attend activities or gatherings, the constant underlying exhaustion limits how intensely they can engage with others. They are there in body but cannot fully participate emotionally or socially. This creates a perceptible distance that has no apparent reason—others sense the patient is “holding back” or “not really there,” but the actual cause (metabolic inability to engage more deeply) is invisible
- **Engagement intensity limited by energy, not desire:** The degree of warmth, enthusiasm, investment, and genuine connection patients can offer is capped by available energy, not by their feelings toward others. Friendships that would otherwise be close remain distant because the patient cannot sustain the energy for deeper engagement, creating unexplained coldness that damages the relationship despite the patient’s genuine care
- **Inability to develop meaningful feelings:** The energy limitation affects not only the expression of feelings but the development of feelings themselves. Emotional attachment, fondness, care, and affection require sustained interaction, shared experiences, emotional investment, and cognitive processing to develop. When energy constraints prevent this sustained engagement, feelings toward others remain shallow or fail to develop beyond superficial acquaintance. Patients find themselves unable to develop the deep care and emotional connection that would normally arise in friendships, creating a profound sense of emotional emptiness and isolation even when physically surrounded by potential friends
- **Social interactions as potential threats:** The knowledge that any conflict or difficulty is insurmountable leads to a defensive posture where many interactions are experienced as *opportunities to be aggressed*. Since patients lack the energy to manage disagreement, navigate misunderstanding, or repair relationship damage, any interaction carries the risk of creating a problem they cannot solve. This produces preventive behavior—emotional guardedness, avoidance of deeper topics, reluctance to express needs or preferences—that further impedes the ability to connect with others. Patients become hypervigilant for potential conflict and withdraw preemptively to avoid situations they cannot metabolically handle, creating a self-protective isolation that others perceive as coldness or lack of trust

Clinical significance: The inability to manage even minimally conflictual relationships represents a major, under-recognized source of social disability in ME/CFS. **This cannot be understated:** patients lose friendships, partnerships, and entire social networks not because relationships are unimportant to them, but because the cognitive and emotional energy required to navigate normal relationship dynamics exceeds available capacity.

The defensive stance toward social interaction—experiencing interactions as potential threats and adopting preventive behaviors—is not paranoia or social anxiety disorder. It is a rational response to genuine incapacity. When any disagreement or misunderstanding represents an insurmountable problem due to energy deficit, hypervigilance and preemptive withdrawal become adaptive survival strategies, though they further entrench isolation.

Critically, *the feeling alone is sufficient to drive protective behavior*. Patients do not need to consciously analyze the risk or make deliberate decisions to withdraw—the subjective experience of interactions as threatening automatically triggers defensive responses. This emotional reality shapes behavior independent of objective threat assessment, making the social disability self-reinforcing: the feeling of vulnerability produces protective isolation, which prevents connection, which maintains isolation.

Environmental Control as Survival Mechanism

The energy deficit necessitates a level of environmental control that is incompatible with normal social spontaneity and fundamentally at odds with what others experience as “the joy of life”:

- **Need for high control:** Patients require predictability, structure, and control over their environment to prevent energy-depleting surprises. Unforeseen events, changes in plans, unexpected social demands, or environmental chaos each represent potential energy expenditures that may trigger crashes
- **Incompatibility with spontaneity:** What healthy individuals experience as joyful spontaneity—surprise visits, impromptu plans, playful chaos, unexpected adventures—registers for ME/CFS patients as threatening unpredictability requiring energy they do not have
- **Others’ joy as patient’s stress:** When others behave in ways they enjoy—being spontaneous, playful, or socially unpredictable—they create a more energetically demanding environment for patients. The very behaviors that make life feel vibrant and enjoyable for healthy people increase the metabolic burden and stress for patients beyond what they can afford to manage
- **Inability to “let go”:** Patients cannot easily relax control over their environment because this control is *almost vital* to avoid exhaustion and crashes. What appears as rigidity, controlling behavior, or inability to be spontaneous is actually a survival mechanism—without environmental control, energy expenditure becomes unpredictable and unmanageable
- **Social consequences:** Others perceive the need for control as rigidity, inflexibility, being “no fun,” or being controlling. Patients are seen as unable to enjoy life, overly cautious, or anxiety-driven when the actual issue is metabolic necessity

- **The paradox of joy:** Patients are often told to “relax,” “let go,” “be spontaneous,” or “just have fun”—but these very behaviors require energy reserves they do not possess. The inability to engage in joyful spontaneity is not psychological resistance but physiological impossibility

The fundamental incompatibility: Normal social life thrives on a degree of unpredictability, spontaneity, and flexibility that ME/CFS patients cannot metabolically afford. The environmental control necessary for survival (avoiding crashes, managing energy) is experienced by others as joyless rigidity. Patients must choose between:

1. Maintaining control to prevent crashes (perceived as controlling, rigid, unable to have fun)
2. Allowing spontaneity to please others (risking energy depletion, crashes, worsening disability)

There is no middle ground when energy reserves are this limited. The choice to maintain control is not preference or personality—it is metabolic necessity masquerading as behavioral rigidity.

The Energy Poverty Analogy. The psychological state of ME/CFS patients living with severe energy deficit is analogous to the lived experience of people in extreme financial poverty:

- **Constant precariousness:** Just as very poor people live under constant financial stress knowing that any unforeseen expense—even an insignificant 20–50€ debt—could trigger a cascade of catastrophic consequences (eviction, utility shutoff, inability to afford food or medical care), ME/CFS patients live under constant metabolic stress knowing that any unforeseen energy expenditure can trigger crashes that eliminate function for days, weeks, or permanently
- **Inability to absorb shocks:** People with financial reserves can absorb unexpected expenses without crisis. People with energy reserves can absorb unexpected demands without crashing. Those living at the edge—whether financial or metabolic—have no buffer. Every unexpected demand is a potential catastrophe
- **Hypervigilance as survival:** The poor must constantly monitor their finances, avoid any unnecessary spending, and maintain rigid control over their budget to prevent disaster. ME/CFS patients must constantly monitor their energy, avoid any unnecessary expenditure, and maintain rigid control over their environment to prevent crashes. Both behaviors appear as anxiety or rigidity to those with adequate resources but are rational responses to genuine scarcity
- **Incomprehension from the resourced:** People with financial security cannot understand why the poor seem so anxious about “small” expenses or why they cannot “just relax” about money. People with energy reserves cannot understand why ME/CFS patients seem so anxious about “small” demands or why they cannot “just relax” and be spontaneous. The invisible nature of the deficit makes the defensive behavior appear irrational

- **Poverty trap dynamics:** Financial poverty creates conditions that perpetuate poverty (stress impairs decision-making, lack of resources prevents investment in improvement). Energy poverty creates conditions that perpetuate energy deficit (stress depletes energy, lack of reserves prevents activities that might improve capacity). Both are self-reinforcing traps difficult to escape
- **Judgment and blame:** The poor are blamed for being “too cautious,” “no fun,” unable to enjoy life, overly anxious, or having a scarcity mindset. ME/CFS patients are blamed for being controlling, rigid, unable to be spontaneous, overly anxious, or having a fearful personality. In both cases, the behavior is adaptive to genuine scarcity, not a character flaw

Clinical significance: Understanding ME/CFS energy management through the lens of poverty economics helps clarify why patients exhibit behaviors that appear rigid or controlling to healthy observers. The “energy poverty” framework explains the hypervigilance, need for control, inability to tolerate unpredictability, and constant stress as rational adaptations to living at the metabolic edge. Just as telling someone in extreme financial poverty to “stop worrying about money and have fun” is tone-deaf and unhelpful, telling ME/CFS patients to “relax,” “let go,” or “be spontaneous” fundamentally misunderstands their metabolic reality.

Even when patients *can* attend activities, the pervasive exhaustion creates an invisible barrier to genuine engagement. Others perceive this as emotional distance, lack of interest, or “holding back”—but it reflects metabolic incapacity, not psychological withdrawal. The patient may desperately want to engage more warmly, more deeply, with more enthusiasm and investment, but the energy simply does not exist. This creates relationships that feel inexplicably cold or distant despite no apparent reason, as the actual limitation (energy deficit) is invisible to observers.

This pattern is distinct from social anxiety or avoidant personality disorder—patients often desperately *want* connection but physiologically *cannot* sustain the energy expenditure relationships require, particularly when any degree of conflict or complexity arises.

Neurobiological Basis

The social and emotional impairments described above are explained by the documented neurological abnormalities:

- **Catecholamine depletion:** Low dopamine and norepinephrine impair both reward processing (making social interaction unrewarding) and the motivation to engage socially
- **Prefrontal hypometabolism:** Reduced energy availability in prefrontal regions impairs the executive functions required for social cognition
- **Effort-reward miscalculation:** TPJ dysfunction causes the brain to perceive social interaction as high-cost, low-reward activity
- **Cerebral hypoperfusion:** Reduced blood flow limits the brain’s capacity to sustain the metabolic demands of complex social processing

- **ATP depletion:** Fundamental energy insufficiency makes any sustained cognitive activity painful

Clinical Significance

Recognition and Validation

Social withdrawal and flat affect in ME/CFS are **metabolic symptoms**, not personality traits, character flaws, or pure psychiatric conditions.

For patients: If social interaction feels painful, if you feel no enjoyment in activities that once brought pleasure, if others tell you that you seem “unhappy” or “unengaged”—these are recognized manifestations of the neurometabolic dysfunction documented in ME/CFS research. This is not your fault. You are not antisocial, cold, or broken. Your brain lacks the energy and neurochemical substrates required for normal social and emotional functioning.

For clinicians and caregivers: Patients who appear disengaged, flat, or “unmotivated” for social interaction are not exhibiting “behavioral problems.” They are conserving severely limited energy reserves. Pressure to “be more social” or “act happier” is equivalent to demanding that someone with severe anemia run a marathon. The physiology does not support the demand.

For researchers: The social and emotional dysfunction in ME/CFS deserves systematic study alongside more commonly recognized cognitive domains. Validated instruments for assessing “social exhaustion,” “affective energy expenditure,” and “interpersonal metabolic cost” would help quantify this significant source of disability.

△ **Warning 8: Harmful Advice: The “Power of Positive Thinking”**

Some clinicians, family members, friends, and caregivers, despite good intentions, offer advice to ME/CFS patients that is not only unhelpful but actively harmful and insulting:

The harmful message:

- “You need to be more optimistic”
- “Believing you will get better will make you better”
- “Your attitude is holding you back”
- “The mind-body connection means positive thinking can heal you”
- “You need to stop focusing on your symptoms”

Why this is harmful:

1. **Blames the patient for their illness:** This framing implies that patients are sick because they are not trying hard enough to think positively, placing moral responsibility for a metabolic disease on the patient’s psychological state
2. **Contradicts objective evidence:** The 2024 NIH study documented measurable neurological abnormalities—low catecholamines, TPJ dysfunction, cerebral hypoperfusion, T-cell exhaustion. These are not created or maintained by “negative

thinking” and cannot be resolved by “optimism”

3. **Ignores patient experience:** Decades of lived experience show that ME/CFS patients who maintain hope, who try every treatment, who remain optimistic, still worsen or remain severely ill. The disease trajectory is independent of psychological attitude
4. **Dismissive and insulting:** Telling someone with documented metabolic dysfunction that their attitude is the problem is equivalent to telling a diabetic that believing their pancreas works will make it produce insulin. It dismisses the physiological reality of the disease
5. **Adds psychological burden:** Patients already carry immense guilt and self-blame (“Why can’t I do what I used to do? Why am I letting everyone down?”). Being told their illness persists because they are not optimistic *enough* adds psychological torment to physical suffering
6. **Prevents appropriate treatment:** When clinicians attribute symptoms to psychological factors, they fail to investigate and treat the underlying metabolic, immunological, and neurological dysfunction
7. **Gaslighting:** This advice constitutes medical gaslighting—denying the patient’s lived reality and documented physiological abnormalities in favor of a psychosomatic explanation that places blame on the patient

The reality:

- ME/CFS patients are not sick because they lack optimism
- Positive thinking does not reverse catecholamine depletion, mitochondrial dysfunction, or immune exhaustion
- Many patients maintain hope and optimism for *decades* while their condition worsens—their attitude did not prevent deterioration
- The mind-body connection exists, but it does not mean that metabolic diseases can be thought away
- Encouraging appropriate pacing, realistic expectations, and acceptance of limitations is more therapeutic than false promises that optimism will cure metabolic dysfunction

For clinicians: If you find yourself telling ME/CFS patients to “be more optimistic” or attributing their symptoms to psychological factors, recognize that you are:

1. Contradicting objective research evidence
2. Causing psychological harm
3. Failing to provide appropriate medical care
4. Perpetuating the decades of medical gaslighting that has defined ME/CFS patient experience

The appropriate clinical response is to acknowledge the physiological reality of the disease, validate the patient’s experience, support symptom management and pacing, and avoid placing the burden of recovery on the patient’s psychological state.

8.7.3 Fluctuation and Post-Exertional Cognitive Malaise

A characteristic feature distinguishing ME/CFS cognitive dysfunction from other conditions is its marked fluctuation, including hour-to-hour and day-to-day variability, worsening with physical, cognitive, or emotional exertion, delayed deterioration (cognitive “payback”), and improvement with rest that rarely returns to premorbid baseline.

8.7.4 CNS Energy Crisis as Primary Event

The selective energy dysfunction hypothesis (Section 14.24) proposes that neurological symptoms in ME/CFS reflect *primary* CNS energy failure rather than downstream effects of systemic dysfunction. Several observations support this framing:

1. **CNS-specific findings:** Neuroinflammation (45–199% elevation in key regions [73]), catecholamine deficiency in CSF, and regional hypometabolism are documented in the CNS specifically, not as reflections of peripheral dysfunction
2. **Preserved autonomous processes:** Hair growth, nail growth, and basic wound healing—processes that operate locally without CNS coordination—remain intact even in severe ME/CFS, arguing against global metabolic failure
3. **Demand-response failure:** The pattern of preserved baseline function with impaired challenge response (91–100% show abnormal CBF reduction during orthostatic challenge [173]) is consistent with a CNS coordination bottleneck rather than peripheral end-organ dysfunction
4. **Cognitive triage hierarchy:** The observation that complex cognition and executive function (“brain fog”) are affected before motor coordination or sensory processing suggests an energy triage system that sacrifices “luxury” cognitive functions first
5. **Astrocyte vulnerability:** The brain’s unique metabolic architecture—with neurons depending on astrocytes for lactate via the ANLS (Section 8.1.5)—may create CNS-specific vulnerability not present in peripheral tissues with direct glucose access [174, 175]

This perspective has treatment implications: interventions that bypass CNS coordination (e.g., direct-acting autonomic agents like midodrine) or that specifically target CNS metabolism may be more effective than peripheral mitochondrial support alone.

⚠ Limitation 6: Brain-Centric Model: CNS Primacy Not Demonstrated

Multiple sections of this chapter frame CNS dysfunction as the “primary bottleneck” or “upstream driver” of ME/CFS symptoms. This causal hierarchy has not been established:

- No longitudinal study has demonstrated that neurological abnormalities *precede* peripheral immune, metabolic, or autonomic dysfunction in ME/CFS onset.
- Alternative causal orderings are equally plausible: primary immune dysfunction driving secondary neuroinflammation, or primary metabolic failure producing secondary CNS changes.

- The “demand-response failure” pattern (preserved baseline with impaired challenge response) is also predicted by peripheral models (e.g., mitochondrial dysfunction limiting peak capacity).
- Pharmacological bypass evidence (midodrine efficacy) supports autonomic *involvement* but does not establish CNS *primacy*—peripheral autonomic failure would produce identical responses.
- The cascade model, kindling hypothesis, and energy triage model are internally consistent but rest on untested causal assumptions; internal consistency is not evidence of causal direction.

8.7.5 CNS Energy Triage: A Hierarchical Model of Brain Fog

Speculation 14 (CNS Energy Triage Hypothesis). **Certainty: 0.35.** The brain may operate a hardwired energy prioritization system during metabolic scarcity, explaining why ME/CFS cognitive dysfunction follows a characteristic pattern rather than producing uniform degradation across all domains.

Neuroscience of brain energy prioritization. The human brain comprises approximately 2% of body mass yet consumes 20–25% of resting metabolic energy, with goal-directed cognition requiring only an additional ~5% above resting homeostatic costs [304]. This tight energy budget means that even modest metabolic deficits—such as those produced by impaired astrocyte-neuron lactate shuttling (Section 8.1.5)—could disproportionately affect the most energy-intensive neural processes.

Not all brain regions have equal metabolic demands. The prefrontal and frontoparietal association cortices, which support executive function, cognitive flexibility, and novel problem-solving, exhibit the highest *relative metabolic cost*—defined as energy utilization exceeding baseline activity levels [304]. In contrast, brainstem nuclei governing vital functions (respiration, cardiovascular regulation, arousal) and primary sensory cortices operate with lower relative metabolic overhead, relying on phylogenetically older, more energy-efficient circuits.

Evidence from metabolic disruption models. Two natural experiments demonstrate hierarchical cognitive shutdown under energy scarcity:

1. **Hypoglycemia:** Acute reduction in brain glucose supply impairs complex higher-order cognitive processes at higher glucose thresholds and to a greater extent than lower-level functions. Executive functions show large effect sizes ($d > 0.8$) during hypoglycemia [305], consistent with the prefrontal cortex’s elevated metabolic sensitivity.
2. **Anesthesia:** General anesthetics produce a hierarchical disconnection pattern in which prefrontal and association cortices are affected first, while primary sensory processing and thalamocortical connectivity remain preserved. Mashour characterizes this as preferential failure of “rich club” network hubs with greater metabolic demands [306]—an “airport in a snowstorm” analogy where the most connected, most metabolically expensive nodes fail first.

Furthermore, prolonged cognitive work causes glutamate accumulation specifically in the lateral prefrontal cortex, making further executive function activation progressively more metabolically costly [307]. This suggests a built-in mechanism by which the brain curtails its most expensive operations when metabolic capacity is strained.

Application to ME/CFS. We speculate that ME/CFS produces a chronic version of this triage state. If total available CNS energy is reduced—whether through astrocyte dysfunction, reduced cerebral blood flow, or neuroinflammation—the brain may engage the same prioritization hierarchy that normally activates only during acute metabolic crises. The proposed triage order, from most to least protected, would be:

1. Brainstem vital functions (preserved even in severe ME/CFS)
2. Basic sensory processing (usually intact)
3. Language comprehension (impaired only in severe cases)
4. Motor coordination (degraded in moderate-severe disease)
5. Memory consolidation (commonly affected)
6. Executive function and cognitive flexibility (affected early, often prominently)

This maps to the formal energy triage hypothesis developed in Section 14.24 (specifically Hypothesis 14.24.5), but here we emphasize the clinical neuroscience basis rather than the mathematical framework.

An important caveat from meta-analytic evidence. The largest meta-analysis of cognitive impairment in ME/CFS (33 studies, $n = 1,086$) reveals that the observed pattern is more nuanced than a simple “executive function fails first” model [303]. Processing speed shows the largest impairment ($g = -0.82$), followed by sustained attention ($g = -0.75$), then memory domains ($g = -0.55$ to -0.67), with executive function showing a smaller effect ($g = 0.42$) and instrumental functions preserved. This is important: processing speed is *more* impaired than executive function on standard neuropsychological measures.

This apparent discrepancy may be reconciled by recognizing that processing speed is a *global* measure of neural efficiency degraded by any reduction in brain energy delivery, not a specific cognitive tier. It reflects the overall metabolic throughput of cortical circuits rather than a discrete cognitive function. Additionally, standardized tests of executive function (e.g., Trail Making Test Part B) involve relatively routinized operations that may not capture the full metabolic cost of genuinely novel, unstructured problem-solving. The energy triage model predicts that *novel, complex, integrative* cognitive operations fail first—not necessarily the specific neuropsychological domain labeled “executive function” in test batteries.

Testable predictions. If the CNS energy triage model is correct, the following should hold:

- Novel tasks are impaired more than practiced routines at matched difficulty
- Working memory (high-energy encoding) fails before recognition memory (lower-energy pattern completion), as formalized in Hypothesis 14.24.10

- Cognitive hierarchy of impairment maps to regional metabolic demand on FDG-PET
- Severity progression follows the triage order: mild ME/CFS shows primarily executive/speed deficits; severe ME/CFS additionally shows language and motor involvement
- Interventions that bypass energy-expensive processing (routinization, external cognitive scaffolding) should preferentially improve function

Treatment implications. If the brain operates in chronic triage mode, the therapeutic strategy shifts from “try harder” to “reduce the load”: (1) *routinize* daily activities to shift them from energy-expensive prefrontal control to energy-efficient basal ganglia automaticity; (2) use external cognitive scaffolding (lists, alarms, decision templates) to offload executive demands; (3) schedule cognitively demanding tasks during peak energy windows when triage thresholds are temporarily relaxed; (4) explore metabolic interventions (ketone supplementation, cerebral blood flow optimization) that may expand the total energy budget and raise triage thresholds across all tiers.

Limitations. This hypothesis faces several challenges: (1) the meta-analytic evidence does not cleanly support executive function as the most impaired domain [303]; (2) the triage hierarchy has not been directly tested in ME/CFS with tasks specifically designed to probe each tier; (3) alternative explanations for the cognitive pattern exist, including neuroinflammation-mediated cytokine effects on specific circuits [308], tryptophan pathway diversion, and autonomic-mediated cerebral hypoperfusion; (4) the model may oversimplify what is likely a multi-mechanism process. The triage framework should be understood as one contributing mechanism among several, not a complete explanation for ME/CFS cognitive dysfunction.

8.8 Summary: An Integrated Neurological Model

The evidence from the NIH deep phenotyping study and decades of prior research supports an integrated model of neurological dysfunction in ME/CFS [5]. An initiating trigger such as infection or other stressor disrupts central nervous system homeostasis. Microglial activation persists beyond acute illness, producing chronic low-grade neuroinflammation. Catecholamine and tryptophan pathway abnormalities develop, affecting dopamine, norepinephrine, and serotonin signaling (neurotransmitter dysregulation). The temporal-parietal junction and related regions fail to accurately process effort-related information (integrative brain dysfunction). Parasympathetic withdrawal and sympathetic dysregulation produce cardiovascular and multi-organ effects (autonomic dysfunction). Reduced cerebral blood flow limits brain metabolic capacity (cerebrovascular compromise). Finally, fatigue, cognitive dysfunction, orthostatic intolerance, and other symptoms emerge from these converging abnormalities as the clinical manifestations.

This model explains why ME/CFS patients experience fatigue fundamentally different from normal tiredness: the brain’s basic mechanisms for perceiving, estimating, and responding to effort are dysfunctional. Treatment approaches targeting these specific neurological abnormalities may prove more effective than those addressing peripheral fatigue or deconditioning.

△ Warning 9: Stimulant Contraindication

Stimulants (amphetamines, methylphenidate, modafinil) are generally **contraindicated** in ME/CFS despite their effectiveness in other fatigue conditions. While they may temporarily mask fatigue by artificially boosting alertness and motivation, they do not address the underlying energy deficit and may enable activity levels that exceed the patient's true physiological capacity. This can precipitate post-exertional malaise (PEM) and potentially cause permanent deterioration. The neurological model presented here explains why: stimulants affect perceived effort and motivation (downstream of the TPJ dysfunction) without correcting the fundamental mismatch between the brain's effort calculations and actual metabolic capacity. Patients may feel capable of activity that their bodies cannot sustain, leading to crashes. This differs fundamentally from stimulant use in conditions like ADHD or narcolepsy, where the underlying metabolic machinery is intact.

9 Endocrine and Metabolic Dysfunction

Endocrine dysfunction represents a critical but often overlooked dimension of ME/CFS pathophysiology. The endocrine system orchestrates fundamental physiological processes including stress response, energy metabolism, circadian rhythms, reproduction, and immune modulation. Disruption of these hormonal axes provides mechanistic explanations for the multi-system nature of ME/CFS symptoms and connects seemingly disparate clinical features into a coherent pathophysiological framework.

The landmark NIH deep phenotyping study by Walitt et al. (2024) documented central nervous system dysfunction with direct implications for neuroendocrine regulation [5]. Complementing this neurological evidence, recent studies have identified specific endocrine abnormalities spanning the hypothalamic-pituitary-adrenal (HPA) axis, thyroid function, sex hormones, growth factors, glucose metabolism, and circadian regulation. These findings reveal that ME/CFS involves coordinated dysfunction across multiple endocrine systems rather than isolated hormonal deficits.

This chapter examines six major endocrine systems implicated in ME/CFS pathophysiology. The HPA axis shows characteristic blunting with hypersensitive feedback, contributing to stress intolerance and immune dysregulation. Thyroid function abnormalities, particularly the “Low T3 Syndrome,” affect cellular metabolism despite normal TSH levels. Sex hormone dysregulation explains the striking female predominance and menstrual cycle exacerbations. Growth hormone and IGF-1 deficiencies contribute to muscle dysfunction and metabolic impairment. Insulin resistance and cerebral glucose hypometabolism connect to the energy deficit discussed in Chapter 6. Finally, circadian rhythm disruption integrates with the sleep abnormalities and autonomic dysfunction detailed in Chapters 8 and 10. Chapter 13 synthesizes these endocrine connections with immune and metabolic systems into comprehensive models of ME/CFS pathophysiology.

Understanding endocrine dysfunction is essential for several reasons. First, hormonal abnormalities provide measurable biomarkers for diagnosis and disease monitoring. Second, endocrine pathways mechanistically link immune activation (Chapter 7) to metabolic dysfunction (Chapter 6). Third, hormonal dysregulation explains symptom patterns such as post-exertional malaise, orthostatic intolerance, and cognitive impairment that define the clinical presentation. Finally, endocrine interventions represent potential therapeutic targets, though current evidence remains mixed and requires careful evaluation.

9.1 Hypothalamic-Pituitary-Adrenal (HPA) Axis

The hypothalamic-pituitary-adrenal axis represents one of the most extensively studied endocrine systems in ME/CFS, yet paradoxically remains among the most controversial.

Through glucocorticoid signaling, the HPA axis coordinates the body's stress response and regulates immune function while maintaining glucose homeostasis and energy metabolism. It further modulates circadian rhythms, sleep-wake cycles, cognitive function, and mood regulation. Given these critical roles, HPA dysfunction provides a plausible mechanism linking the diverse symptoms of ME/CFS.

Figures 9.1 and 9.2 illustrate the characteristic pattern of HPA axis dysfunction observed in ME/CFS. Unlike the robust circadian cortisol rhythm and responsive feedback regulation seen in healthy individuals, ME/CFS patients demonstrate a distinct pattern of dysregulation. This involves blunted corticotropin-releasing hormone (CRH) secretion from the hypothalamus, reduced adrenocorticotropic hormone (ACTH) response from the pituitary, flattened diurnal cortisol rhythm with loss of the normal morning peak, and paradoxically enhanced negative feedback sensitivity. This constellation of abnormalities distinguishes ME/CFS from both healthy states and primary adrenal insufficiency (Addison's disease), suggesting a unique form of central HPA axis hypofunction.

9.1.1 HPA Axis Abnormalities


Cortisol Dysregulation Patterns

Observation 50 (Characteristic Cortisol Pattern in ME/CFS). Multiple studies document a consistent pattern of cortisol abnormalities in ME/CFS patients that differs qualitatively from both healthy individuals and patients with primary adrenal disorders. Baseline cortisol levels tend to be lower, though typically remaining within the broad "normal" laboratory reference range. The diurnal cortisol rhythm shows flattening, with reduced amplitude between morning and evening values. The cortisol awakening response (CAR)—the normal sharp rise in cortisol during the first 30–60 minutes after waking—appears attenuated. Additionally, cortisol responses to physiological and psychological stressors are blunted, despite appropriate ACTH response to exogenous CRH stimulation in some studies [309].

The NIH deep phenotyping study by Walitt et al. identified neuroendocrine abnormalities consistent with HPA axis dysfunction, including altered catecholamine metabolism that affects upstream regulation of the HPA axis [5]. The reduced central catecholamines documented in cerebrospinal fluid may contribute to impaired hypothalamic CRH release, providing a mechanistic link between neurological and endocrine dysfunction.

Recent sex-stratified analysis by Pipper et al. (2024) revealed that cortisol dysregulation patterns differ significantly between male and female ME/CFS patients and vary by disease severity [309]. Female patients with severe ME/CFS demonstrated elevated 11-deoxycortisol (a cortisol precursor) and 17 α -hydroxyprogesterone, suggesting impaired final enzymatic steps in cortisol synthesis. Male patients with mild to moderate disease showed frankly reduced cortisol and corticosterone levels but paradoxically elevated progesterone. These findings indicate that HPA dysfunction may involve enzyme deficiencies in steroidogenesis rather than simple hypothalamic-pituitary signaling deficits.

Normal HPA Axis Function



ME/CFS: HPA Axis Dysregulation




Figure 9.2: ME/CFS HPA axis dysregulation with blunted response and systemic consequences.

ACTH and CRH Abnormalities

The central components of the HPA axis—CRH from the hypothalamus and ACTH from the pituitary—show complex abnormalities that do not fit simple models of endocrine failure. Studies employing CRH stimulation tests have yielded inconsistent results. Some report normal ACTH and cortisol responses to exogenous CRH administration, while others document blunted ACTH responses despite adequate CRH stimulation. Still others find normal ACTH responses but reduced cortisol output, suggesting adrenal hyposensitivity. These inconsistencies likely reflect the heterogeneity of ME/CFS patient populations, differences in disease duration and severity, and the limitations of single-timepoint testing to capture dynamic regulatory dysfunction.

The most consistent finding across studies is evidence of enhanced negative feedback sensitivity. Dexamethasone suppression tests demonstrate that low doses of synthetic glucocorticoid produce greater and more prolonged suppression of cortisol secretion in ME/CFS patients compared to controls. This suggests that the hypothalamus and pituitary remain exquisitely sensitive to glucocorticoid feedback signals, inappropriately dampening HPA axis output even when cortisol levels are already low-normal. This pattern resembles the neuroendocrine adaptation seen in chronic stress conditions but persists inappropriately in ME/CFS despite the clinical need for robust stress responses.

Diurnal Rhythm Disruption

Observation 51 (Loss of Cortisol Circadian Amplitude). The diurnal cortisol rhythm represents one of the most robust and well-characterized circadian processes in human physiology, yet ME/CFS patients consistently demonstrate flattening of this rhythm. Healthy individuals show a sharp cortisol peak within 30–60 minutes of waking (cortisol awakening response), followed by progressive decline throughout the day, reaching a nadir around midnight, and beginning to rise again in the early morning hours (3–4 AM) in anticipation of waking. In contrast, ME/CFS patients show reduced morning cortisol peak (blunted CAR), less pronounced decline during the day (flatter slope), and reduced overall amplitude (difference between peak and nadir), resulting in a “flattened” 24-hour pattern [310].

The mechanistic basis for circadian rhythm disruption extends beyond the HPA axis itself to involve the central circadian clock in the suprachiasmatic nucleus (SCN) of the hypothalamus. The NIH study documented abnormalities in temporal-parietal junction function and altered brain metabolism that may affect SCN regulation [5]. Additionally, inflammatory cytokines known to be elevated in ME/CFS (discussed in Chapter 7) directly disrupt circadian clock gene expression, creating bidirectional interactions between immune activation and circadian dysregulation.

The clinical consequences of flattened cortisol rhythm are profound. The morning cortisol peak serves essential physiological functions: promoting waking and alertness, mobilizing glucose for energy availability, preparing the cardiovascular system for upright posture and activity, and modulating immune function to prevent excessive inflammation. Loss of this peak explains the characteristic morning symptom severity reported by many ME/CFS patients.

These include difficulty waking, prolonged morning fatigue requiring hours to achieve minimal function, orthostatic intolerance upon standing (discussed in Chapter 10), and cognitive dysfunction particularly severe in early morning hours.

9.1.2 Mechanisms of HPA Dysfunction

The dysregulation of the HPA axis in ME/CFS reflects multiple interconnected mechanisms operating at different levels of the neuroendocrine cascade. Understanding these mechanisms is essential for developing targeted therapeutic interventions and explaining why simple hormone replacement strategies have shown limited efficacy.

Central Glucocorticoid Receptor Sensitivity

~ Hypothesis 1: Enhanced Central Glucocorticoid Feedback

The enhanced negative feedback sensitivity observed in ME/CFS may result from altered glucocorticoid receptor (GR) expression or function in hypothalamic and pituitary tissues. Several mechanisms could produce this effect. Upregulation of GR expression would increase sensitivity to existing cortisol levels. Altered GR isoform expression (GR α vs. GR β) might shift the balance toward enhanced feedback. Reduced expression of 11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1), the enzyme that locally amplifies cortisol action by converting inactive cortisone to active cortisol, could diminish local glucocorticoid signaling. Finally, epigenetic modifications of the GR gene might affect transcription and receptor function.

Falsifiability: *weakly* — Falsified if hypothalamic GR expression is normal or reduced in ME/CFS patients, or if elevated cortisol fails to suppress ACTH via enhanced feedback

This enhanced feedback creates a self-reinforcing cycle. Slightly elevated cortisol (or even normal-low cortisol) triggers disproportionate suppression of CRH and ACTH secretion, further reducing cortisol output. Under normal circumstances, this would reduce feedback inhibition and restore output, but the hypersensitive feedback prevents this compensatory response, maintaining chronically low HPA axis activity. This mechanism explains why ME/CFS patients do not develop frank adrenal insufficiency (baseline cortisol remains detectable) yet fail to mount appropriate stress responses (blunted reactivity to challenges). This self-reinforcing HPA dysfunction represents one of several vicious cycles in ME/CFS pathophysiology, as discussed in Section 13.4 of Chapter 13.

Inflammatory Cytokine Effects on HPA Axis

The bidirectional relationship between the immune system and the HPA axis represents a critical mechanism in ME/CFS pathophysiology. Under normal circumstances, immune activation from infection or tissue damage stimulates HPA axis activity. Pro-inflammatory cytokines (IL-1, IL-6, TNF- α) signal the hypothalamus to increase CRH secretion, resulting in elevated cortisol that dampens the immune response. This creates negative feedback that

prevents excessive inflammation. The acute response adaptively contains immune activation while preventing immunopathology.

~ Hypothesis 2: Maladaptive Chronic Inflammatory Signaling

In ME/CFS, chronic low-grade inflammation (documented in Chapter 7) may induce glucocorticoid resistance at immune cells while simultaneously increasing central negative feedback sensitivity. This paradoxical pattern produces the worst of both scenarios: insufficient cortisol secretion to control peripheral inflammation due to enhanced central feedback, yet reduced cortisol effectiveness at immune cells due to receptor downregulation or dysfunction [5, 311]. The result is persistent inflammation despite apparent “normal” cortisol levels that would typically suppress such immune activation.

Falsifiability: *weakly* — Falsified if ME/CFS immune cells show normal glucocorticoid receptor function and normal immune suppression at physiological cortisol concentrations

Recent evidence from Heng et al. (2025) documenting sex-specific immune dysregulation supports this model, showing that females with ME/CFS exhibit particularly pronounced pro-inflammatory profiles with elevated type 2 interferon signaling despite cortisol levels within the reference range [311]. This suggests functional glucocorticoid resistance at target tissues.

Steroidogenic Enzyme Dysfunction

The recent findings by Pipper et al. (2024) identifying elevated cortisol precursors (11-deoxycortisol, 17 α -hydroxyprogesterone) in severe ME/CFS patients suggest impaired function of steroidogenic enzymes, particularly 11 β -hydroxylase (CYP11B1) which catalyzes the final step converting 11-deoxycortisol to cortisol [309]. This enzyme dysfunction could result from several factors. Mitochondrial impairment may play a role, as steroidogenesis occurs in mitochondria and requires adequate ATP supply (discussed in Chapter 6). Cytokine-mediated suppression of enzyme expression or activity represents another possibility. Micronutrient deficiencies affecting enzyme cofactors or oxidative stress damaging enzyme proteins may also contribute.

If confirmed, this mechanism suggests that the problem is not purely regulatory (hypothalamic-pituitary signaling) but also biosynthetic (adrenal enzymatic capacity). This has important therapeutic implications, as interventions targeting upstream signaling may prove ineffective if the limiting step is enzymatic conversion within the adrenal gland.

⌘ Limitation 1: HPA Axis Dysfunction: Cause, Consequence, or Epiphenomenon

The direction of causality between HPA axis abnormalities and ME/CFS remains unresolved. Key epistemic boundaries:

- No prospective study has demonstrated that HPA dysfunction *precedes* ME/CFS onset; all data are cross-sectional or retrospective.

- Similar HPA axis changes occur in chronic pain syndromes and PTSD (hypoactivation), while melancholic depression shows the opposite pattern (HPA hyperactivation)—the specificity of the ME/CFS profile is unclear.
- CRH stimulation test results are inconsistent across studies (normal, blunted, or dissociated ACTH/cortisol responses), suggesting heterogeneity rather than a unitary mechanism.
- Deconditioning, sleep disruption, and psychological stress—all common in chronic illness—independently produce HPA axis changes indistinguishable from those attributed to ME/CFS pathophysiology.
- The “self-reinforcing cycle” model (enhanced feedback → low cortisol → inflammation → further HPA suppression) is mechanistically plausible but has not been demonstrated longitudinally in ME/CFS patients.

9.1.3 Clinical Consequences

The HPA axis abnormalities documented in ME/CFS produce wide-ranging clinical effects that contribute directly to the cardinal symptoms of the disease. Understanding these consequences illuminates why seemingly minor hormonal changes cause profound functional impairment.

Stress Response Abnormalities and Post-Exertional Malaise

Observation 52 (Blunted Physiological Stress Responses). ME/CFS patients demonstrate inadequate cortisol responses to physiological stressors. These include exercise (both acute bouts and prolonged exertion), orthostatic challenge (standing, tilt-table testing), cognitive tasks requiring sustained mental effort, and psychological stressors. This blunted response means that stressors that healthy individuals accommodate with transient cortisol elevation produce inadequate counter-regulatory responses in ME/CFS patients, potentially explaining the delayed and prolonged symptom exacerbation characteristic of post-exertional malaise (PEM).

The temporal pattern of PEM—symptom onset typically 12–48 hours after exertion rather than immediately—aligns with the kinetics of cortisol’s effects on immune function and cellular metabolism. During exertion, ME/CFS patients may rely on sympathetic nervous system activation (catecholamines) to maintain function despite inadequate cortisol support. Following exertion, the delayed cortisol response fails to adequately suppress the inflammatory cascade initiated by exertion-induced cellular stress and damage. This unchecked inflammation then drives the delayed symptom exacerbation of PEM.

Immune System Effects and Chronic Inflammation

Cortisol serves as the body’s primary endogenous anti-inflammatory hormone. It suppresses pro-inflammatory cytokine production (IL-1 β , IL-6, TNF- α), inhibits T cell activation and

proliferation, promotes a shift from Th1 (cellular immunity) to Th2 (humoral immunity) responses, and prevents autoimmune reactions by maintaining immune tolerance. The blunted cortisol output and flattened diurnal rhythm in ME/CFS remove this tonic immunosuppressive influence, permitting chronic low-grade inflammation to persist.

~ Hypothesis 3: Loss of Diurnal Immune Regulation

The flattened cortisol rhythm may be particularly consequential for immune regulation. Immune cells express glucocorticoid receptors and show circadian variation in their responsiveness to cortisol. The normal morning cortisol peak serves to “reset” immune function daily, preventing inflammatory pathways from remaining chronically activated. Loss of this peak in ME/CFS may allow inflammatory signaling to persist across day-night cycles without the normal circadian suppression [310].

Falsifiability: *weakly* — Falsified if restoring the morning cortisol peak through timed hydrocortisone replacement fails to reduce inflammatory markers across the circadian cycle

This mechanism connects to the findings in Chapter 7 documenting altered cytokine profiles, NK cell dysfunction, and B cell abnormalities in ME/CFS. The sex-specific immune profiles identified by Heng et al. (2025) showing more pronounced inflammatory signatures in females align with the sex-specific steroid hormone abnormalities documented by Pipper et al. (2024), suggesting coordinated sex-dependent endocrine-immune interactions [311, 309].

Energy Metabolism and Glucose Homeostasis

Cortisol plays essential roles in energy metabolism. It stimulates hepatic gluconeogenesis to maintain blood glucose availability, promotes lipolysis (fat breakdown) to provide alternative fuel sources, modulates insulin sensitivity to optimize glucose utilization, and supports mitochondrial function and cellular energy production. HPA axis dysfunction directly impairs these metabolic processes.

The cerebral glucose hypometabolism documented by PET imaging studies (Tirelli et al., 1998; Siessmeier et al., 2003) may partly reflect inadequate cortisol support for glucose uptake and utilization [312, 313]. Additionally, the blunted morning cortisol peak fails to provide the metabolic “boost” needed to transition from fasting metabolism to active daytime metabolism, contributing to severe morning fatigue and the prolonged time required to achieve minimal function after waking.

These metabolic effects connect directly to the mitochondrial dysfunction and energy metabolism deficits discussed in Chapter 6, suggesting that HPA axis abnormalities and cellular metabolic dysfunction represent interconnected rather than independent pathophysiological processes.

Gut Barrier Repair and Low Cortisol in Severe Patients. Beyond the well-established role of cortisol in stress response and metabolism, emerging evidence suggests that HPA axis dysfunction in severe ME/CFS may impair intestinal barrier maintenance and repair capacity, potentially contributing to chronic gut permeability and systemic inflammation.

Low Morning Cortisol in Severely Ill Patients. A comprehensive biomarker examination of severely ill (housebound/bedbound) ME/CFS patients [314] documented significantly reduced morning salivary cortisol compared to age- and sex-matched healthy controls: median 0.20 mcg/dL in severe ME/CFS vs. 0.45 mcg/dL in controls ($p = 0.002$). To contextualize these values clinically: normal morning salivary cortisol ranges from approximately 0.10 to 0.90 mcg/dL, with 0.20 mcg/dL representing roughly the 15th percentile and 0.45 mcg/dL representing the 50th percentile of the healthy population distribution. Thus, severe ME/CFS patients exhibit cortisol levels that, while not meeting formal diagnostic criteria for adrenal insufficiency (<0.10 mcg/dL), fall well below the physiological optimum for barrier maintenance and metabolic function. This 55% reduction in morning cortisol suggests substantial HPA axis dysregulation in the severe patient population. While the mechanisms remain debated, the functional consequence is reduced cortisol availability during periods when barrier repair processes are most active.

Cortisol's Role in Intestinal Barrier Function. Glucocorticoids, including cortisol, play multifaceted roles in maintaining epithelial barrier integrity: (1) upregulating tight junction proteins (claudin-1, occludin, ZO-1) through glucocorticoid receptor-mediated transcription; (2) suppressing barrier-disrupting cytokines (IL-1 β , IL-6, TNF- α) via NF- κ B inhibition; (3) promoting enterocyte survival and differentiation; and (4) maintaining circadian tight junction protein expression with peak synthesis during the morning cortisol surge. The relationship is dose-dependent: physiological cortisol levels are barrier-protective, while low cortisol impairs repair capacity. ME/CFS patients appear to have insufficient cortisol for normal barrier maintenance.

Impaired Barrier Repair in Severe ME/CFS: Mechanistic Hypothesis. Integrating low cortisol with evidence of baseline gut permeability [315], severe patients likely exhibit: (1) daily micro-damage from minimal activities (cognitive work, postural changes, meals) triggering transient splanchnic hypoperfusion; (2) insufficient nocturnal cortisol for barrier repair; (3) accumulating baseline permeability; (4) the cytokine-barrier bidirectional cycle (cytokines $\rightarrow$ tight junction disruption $\rightarrow$ LPS $\rightarrow$ cytokine amplification); and (5) nutritional deficits (low albumin in severe patients) limiting epithelial regeneration substrate. This model suggests wheat elimination response may be slower in severe patients due to impaired barrier repair capacity, but nutritional support (protein, micronutrients) and optimization of morning cortisol timing may accelerate recovery.

9.2 Thyroid Function

Thyroid dysfunction represents a critical consideration in ME/CFS for two distinct reasons: the substantial clinical overlap between hypothyroidism and ME/CFS symptoms creates diagnostic challenges requiring careful differentiation, and ME/CFS patients exhibit a specific pattern of thyroid abnormalities—the “Low T3 Syndrome”—that occurs despite normal TSH levels and complicates interpretation of standard thyroid function tests. Understanding these thyroid-related issues is essential for appropriate diagnosis and management.

9.2.1 The Low T3 Syndrome in ME/CFS

★ Achievement 1: Low T3 Syndrome as Distinct ME/CFS Feature

Ruiz-Núñez et al. (2018) conducted a rigorous case-control study comparing 98 ME/CFS patients to 99 healthy controls and documented a distinctive pattern of thyroid hormone abnormalities [316]. ME/CFS patients showed 16% prevalence of free T3 (FT3) below the reference range compared to only 7% in controls (odds ratio 2.56). They exhibited significantly lower FT3, total T4 (TT4), and total T3 (TT3) concentrations, along with reduced T3/T4 ratio indicating impaired peripheral conversion of T4 to active T3. The percentage of reverse T3 (rT3), an inactive T3 isomer, was elevated, with increased rT3/TT3 ratio reflecting preferential conversion to the inactive form. Estimated deiodinase activity—the enzyme responsible for converting T4 to T3—showed a 14.4% reduction.

Critically, these abnormalities occurred while thyroid-stimulating hormone (TSH) levels remained within the normal reference range. Standard thyroid screening tests would therefore classify these patients as “euthyroid” (normal thyroid function) despite functionally significant thyroid hormone deficits.

△ Warning 1: Replication Status: Partially Replicated

The Low T3 pattern (Ruiz-Núñez 2018, n=98 vs. 99) is the only ME/CFS-specific thyroid study. However, the Low T3/euthyroid sick pattern is well-established in other chronic diseases, providing indirect support. ME/CFS-specific replication with standardized thyroid panels is needed.

This pattern resembles the “non-thyroidal illness syndrome” (NTIS) or “euthyroid sick syndrome” observed in acute critical illness, starvation, and chronic diseases. However, unlike the transient thyroid changes in acute illness that normalize with recovery, the Low T3 Syndrome in ME/CFS persists chronically and may represent a maladaptive response that perpetuates rather than resolves the disease state.

⚡ Limitation 2: Low T3 Syndrome: Adaptive Response vs. Pathological Driver

Whether the Low T3 pattern in ME/CFS represents a maladaptive state requiring correction or an appropriate protective adaptation (reducing metabolic demand in the face of energy deficiency) has not been determined:

- The characterisation rests on a single ME/CFS-specific study (Ruiz-Núñez 2018, n=98 vs. 99). The Low T3/NTIS pattern itself is well-established in other chronic diseases, but ME/CFS-specific mechanisms remain unvalidated.
- No interventional study has demonstrated that correcting Low T3 (via T3 supplementation) improves ME/CFS outcomes without adverse effects—mixed clinical results are equally consistent with the adaptation hypothesis.
- The proposed cytokine-mediated deiodinase suppression mechanism is inferred from general immunology; deiodinase activity has not been directly measured in

ME/CFS patients.

9.2.2 Mechanisms of Impaired T4 to T3 Conversion

The conversion of thyroxine (T4, the major thyroid hormone secreted by the thyroid gland) to triiodothyronine (T3, the metabolically active form) occurs primarily in peripheral tissues through the action of deiodinase enzymes. Three deiodinase isoforms exist. Type 1 deiodinase (D1) in liver and kidney produces most circulating T3. Type 2 deiodinase (D2) in brain, pituitary, and brown fat produces local T3 for tissue-specific needs. Type 3 deiodinase (D3) in multiple tissues inactivates T4 and T3 by converting them to reverse T3 (rT3) and T2.

~ Hypothesis 4: Cytokine-Mediated Deiodinase Suppression

The chronic low-grade inflammation documented in ME/CFS (Chapter 7) likely suppresses deiodinase enzyme activity through multiple mechanisms. Pro-inflammatory cytokines, particularly IL-6 and TNF- α , directly inhibit D1 and D2 expression and activity while upregulating D3, shifting the balance toward production of inactive reverse T3 rather than active T3. Oxidative stress, elevated in ME/CFS (Chapter 6), damages selenocysteine residues essential for deiodinase enzymatic function; all deiodinases are selenium-dependent enzymes. Additionally, the mitochondrial dysfunction documented in ME/CFS may impair ATP-dependent cellular uptake of T4, reducing substrate availability for conversion to T3 [316].

Falsifiability: *weakly* — Falsified if deiodinase enzyme activity is normal despite elevated IL-6/TNF- α , or if patients with documented deiodinase suppression show normal free T3 and reverse T3 levels

This mechanism explains why simply increasing thyroid hormone replacement dose (giving more T4) often fails to improve symptoms in ME/CFS patients: the limiting factor is not T4 availability but rather the capacity to convert T4 to active T3 at the cellular level.

9.2.3 Tissue-Level Thyroid Hormone Resistance

Beyond impaired T4 to T3 conversion, emerging evidence suggests that some ME/CFS patients may exhibit functional thyroid hormone resistance at the cellular level. This could involve reduced expression or function of thyroid hormone transporters (MCT8, MCT10) that move hormones into cells, altered expression of thyroid hormone receptors (TR α , TR β) in target tissues, or impaired receptor-coactivator interactions that reduce transcriptional responses to thyroid hormone binding.

Observation 53 (Selenium Autoantibodies and Acquired Resistance). A Netherlands study identified markedly elevated selenium autoantibodies in 9.6–15.6% of ME/CFS patients compared to only 0.9–2.0% of healthy controls. Selenium is essential for deiodinase function, selenoprotein synthesis, and thyroid hormone metabolism. Autoantibodies against selenium transport proteins could create an acquired form of thyroid hormone resistance by impairing the selenium-dependent enzymatic machinery required for thyroid hormone activation and action.

This finding suggests a potential autoimmune mechanism contributing to thyroid dysfunction in a subset of ME/CFS patients and raises the possibility that interventions targeting selenium metabolism might benefit this subgroup.

9.2.4 Clinical Implications and Diagnostic Challenges

The overlap between ME/CFS symptoms and hypothyroidism creates substantial diagnostic challenges. Both conditions present with severe fatigue and exhaustion, cognitive impairment (“brain fog”), cold intolerance and temperature dysregulation, weight changes and metabolic disturbances, mood alterations including depression, muscle weakness and pain, and sleep disturbances. Undiagnosed hypothyroidism may therefore masquerade as ME/CFS, while ME/CFS-related Low T3 Syndrome may be mistaken for thyroid disease.

△ Warning 2: Limitations of Standard Thyroid Testing in ME/CFS

Standard thyroid screening using TSH alone is insufficient for evaluating thyroid function in ME/CFS patients. The Low T3 Syndrome occurs with normal TSH because the pituitary senses adequate T4 levels and reduces TSH secretion appropriately, unaware that peripheral tissues cannot effectively convert T4 to active T3. Comprehensive thyroid evaluation in ME/CFS should include TSH to exclude primary thyroid disease, free T4 (FT4) to assess thyroid hormone production, free T3 (FT3) to evaluate the active hormone level, and reverse T3 (rT3) to assess the balance between activation and inactivation. Calculation of T3/T4 and rT3/T3 ratios quantifies conversion efficiency [316].

The therapeutic implications remain uncertain. While the rationale for T3 supplementation appears sound (directly providing the active hormone bypasses the impaired conversion step), clinical trial evidence remains limited and results have been mixed. Some patients report subjective improvement, others experience no benefit, and a subset develops adverse effects (palpitations, anxiety, insomnia) suggesting tissue-level hypersensitivity to thyroid hormone. Careful individualized treatment trials with close monitoring may be warranted in ME/CFS patients with documented Low T3 Syndrome, but systematic evidence for efficacy is lacking.

9.3 Sex Hormones and Gender Differences

Sex hormones and gender differences represent one of the most striking yet inadequately explained features of ME/CFS epidemiology and pathophysiology. The consistent 3–4:1 female-to-male prevalence ratio, menstrual cycle exacerbations reported by female patients, early menopause associations, and sex-specific patterns of immune dysfunction and steroid hormone abnormalities collectively indicate that reproductive hormones play central roles in disease susceptibility, expression, and progression.

9.3.1 Epidemiology of Sex Differences

★ Achievement 2: Female Predominance in ME/CFS

A systematic review and meta-analysis by Lim et al. (2020) synthesized prevalence data from 45 articles representing 46 studies and 56 prevalence datasets spanning 1980–2018 [317]. The analysis documented overall ME/CFS prevalence of 0.89% using CDC-1994 criteria, with female prevalence approximately 1.5–2.0 fold higher than males across all studies. In the total population, prevalence was $2.24\% \pm 2.59\%$ in females versus $1.11\% \pm 1.05\%$ in males. General population studies showed 2.83% versus 1.39%, and meta-analysis pooled estimates indicated 1.36% versus 0.89%. The female-to-male ratio consistently ranges from 3:1 to 4:1 across geographic regions, diagnostic criteria, and study methodologies.

More recent estimates using large-scale medical claims data and machine learning by Jason et al. (2018) confirmed the persistent female predominance, though highlighting that 35–40% of ME/CFS patients are male—a substantial population whose experiences may differ from the predominantly studied female cohorts [318]. The consistency of the sex ratio across diverse populations and diagnostic approaches argues strongly for biological sex differences in disease susceptibility or expression rather than artifacts of health-seeking behavior or diagnostic bias.

9.3.2 Sex-Specific Pathophysiology: NIH Deep Phenotyping Findings

★ Achievement 3: Distinct Male and Female Pathophysiological Patterns

The 2024 NIH deep phenotyping study by Walitt et al. employed multi-omics analysis to directly compare male and female ME/CFS patients [5]. This rigorous investigation revealed fundamentally different pathophysiological signatures. Males showed altered T cell activation patterns and abnormal innate immunity markers, while females demonstrated abnormal B cell function and altered white blood cell growth patterns. The sexes exhibited distinct inflammatory marker profiles, divergent gene expression patterns in immune cells, different immune cell population distributions, and sex-specific metabolic marker abnormalities.

This finding challenges the implicit assumption in ME/CFS research that male and female patients share a common pathophysiology differing only in prevalence. Instead, it suggests that ME/CFS may represent partially distinct disease processes in males and females requiring sex-stratified approaches to diagnosis, biomarker development, and treatment.

Complementing the NIH findings, Heng et al. (2025) documented sex-specific immune dysregulation in long COVID patients with ME/CFS [311]. Females exhibited decreased lymphocyte counts with increased neutrophils and monocytes (a myelopoiesis shift), elevated pro-inflammatory cytokines, and upregulated type 2 interferon signaling (IP-10, IFN- γ). Males showed fewer inflammatory alterations overall, with more balanced profiles, elevated anti-inflammatory IL-10, and IL-1 signaling dominance rather than interferon predominance.

These immune differences likely reflect underlying hormonal influences on immune cell development, activation, and cytokine production.

9.3.3 Steroid Hormone Abnormalities

★ Achievement 4: Sex and Severity-Stratified Steroid Hormone Profiles

Pipper et al. (2024) conducted the first comprehensive sex-stratified analysis of steroid hormones in ME/CFS using high-precision UHPLC-MS/MS (ultra-high performance liquid chromatography tandem mass spectrometry) [309]. This study of 97 total participants revealed striking sex-specific and severity-dependent patterns. Female patients with severe ME/CFS demonstrated elevated 11-deoxycortisol (a cortisol precursor) and 17 α -hydroxyprogesterone, suggesting impaired final enzymatic steps in cortisol biosynthesis. Females with mild-to-moderate disease showed increased progesterone levels. Male patients with mild-to-moderate ME/CFS exhibited frankly reduced cortisol and corticosterone but paradoxically elevated progesterone—an unexpected finding suggesting complex dysregulation of steroidogenic pathways.

The machine learning classifier achieved 71.2% accuracy for discriminating female ME/CFS patients from controls and 84.6% accuracy for males based solely on steroid hormone profiles, supporting the potential development of sex-specific hormonal biomarkers for diagnosis and disease monitoring.

△ Warning 3: Replication Status

Single study (Pipper 2024, n=97 total). This is the first sex-stratified steroid hormone analysis in ME/CFS using UHPLC-MS/MS. Subgroup sizes are small (particularly male severe ME/CFS). The machine learning classifier was not validated on an independent test set. Independent replication with larger, sex-balanced cohorts is required.

These findings indicate that sex hormone abnormalities in ME/CFS extend beyond simple deficiency or excess of individual hormones to involve coordinated dysregulation of steroidogenic pathways, enzyme activities, and metabolic ratios. The sex-specific patterns suggest that estrogen and progesterone in females versus testosterone and its metabolites in males exert distinct effects on disease expression.

9.3.4 Reproductive Health and Gynecological Risk Factors

Menstrual Cycle Effects and Hormonal Fluctuations

The majority of premenopausal women with ME/CFS report significant symptom exacerbations related to menstrual cycle phases, particularly during the premenstrual week (late luteal phase when progesterone declines rapidly) and during menstruation itself. Common cyclical exacerbations include increased fatigue and post-exertional malaise severity, worsened cognitive impairment and brain fog, intensified pain and sensory sensitivity, heightened orthostatic intolerance symptoms, and mood disturbances such as irritability, anxiety, and depression.

Observation 54 (Hormone-Symptom Correlations Across Menstrual Cycle). Preliminary findings from chronobiology-based studies mapping hormonal fluctuations across the menstrual cycle in ME/CFS reveal systematic symptom-hormone relationships. Fatigue and pain peak premenstrually when estradiol and progesterone levels fall. Cognitive impairment shows lowest severity at ovulation when estradiol peaks. Low estradiol and progesterone concentrations correlate with higher fatigue and pain ratings. Additionally, luteinizing hormone (LH) and follicle-stimulating hormone (FSH) levels positively correlate with fatigue severity and orthostatic symptoms, suggesting pituitary dysregulation may contribute to symptom variability.

These patterns suggest that the absolute levels of sex hormones may be less important than the dynamic fluctuations and ratios between estrogen, progesterone, and pituitary gonadotropins. The rapid hormonal changes during certain cycle phases may destabilize physiological systems already compromised by ME/CFS, triggering symptom exacerbations.

Gynecological Comorbidities and Early Menopause

★ Achievement 5: Gynecological Risk Factors for ME/CFS

Population-based case-control studies by Boneva et al. have identified specific gynecological risk factors associated with ME/CFS development [319, 320]. The 2011 study documented that ME/CFS cases reported significantly higher rates of pelvic pain unrelated to menstruation (22.2% versus 1.7% in controls), endometriosis diagnosis (36.1% versus 16.7%), prolonged periods of amenorrhea (absence of menstruation), and history of gynecological surgery including hysterectomy and oophorectomy. Premenopausal women with ME/CFS tended toward lower luteal phase progesterone with higher FSH.

Most strikingly, the 2015 study revealed that women with ME/CFS experienced menopause approximately 11 years earlier than controls (mean age 37.6 years versus 48.6 years) [320]. Early menopause (<40 years) represents premature ovarian insufficiency, indicating fundamental dysfunction of the hypothalamic-pituitary-gonadal (HPG) axis. This finding suggests that HPG dysfunction may precede or contribute to ME/CFS development rather than merely resulting from the disease.

△ Warning 4: Replication Status: Partially Replicated (Same Group)

Both studies (Boneva 2011, 2015) derive from the same CDC research group using the same Georgia cohort. While internally consistent, independent replication of the early menopause finding and gynecological risk factors in non-CDC populations is needed.

The mechanisms connecting gynecological abnormalities to ME/CFS remain incompletely understood but likely involve bidirectional interactions. Chronic inflammation (documented in Chapter 7) affects ovarian function, endometrial health, and hormonal regulation. Conversely, sex hormone abnormalities modulate immune function, with estrogen generally enhancing immune responses (potentially contributing to female predominance of autoimmune diseases) and progesterone providing immunosuppressive and anti-inflammatory effects. Disruption

of these hormonal immunomodulatory signals could perpetuate the immune dysfunction characteristic of ME/CFS.

9.3.5 Testosterone and Androgens

While research has focused predominantly on female ME/CFS patients due to the higher prevalence, emerging evidence indicates that male patients exhibit distinct hormonal abnormalities, particularly involving androgens (testosterone, DHEA, and their metabolites). A pilot study of 23 women ages 35–55 with ME/CFS found that 89% had suboptimal DHEA-S (dehydroepiandrosterone sulfate) levels. Supplementation with DHEA led to statistically significant improvements: 18% reduction in pain, 21% reduction in fatigue, 35% reduction in anxiety, 26% improvement in thinking ability, 17% improvement in memory, and 22% improvement in sexual function.

~ Hypothesis 5: Androgen Deficiency Contribution to Symptoms

Androgens, particularly DHEA and testosterone, serve multiple physiological functions beyond reproductive roles. They support mitochondrial function and cellular energy production, promote muscle mass maintenance and physical strength, modulate immune responses with generally anti-inflammatory effects, influence mood, motivation, and cognitive function, and affect pain perception and nociceptive processing. Deficiency of these hormones could mechanistically contribute to core ME/CFS symptoms including fatigue, cognitive impairment, reduced physical capacity, and pain amplification.

Falsifiability: *weakly* — Falsified if DHEA-S and testosterone levels are comparable to healthy controls, or if androgen replacement fails to improve fatigue and cognition in patients with documented deficiency

The sex-specific steroid profiles identified by Pipper et al. showing reduced cortisol but elevated progesterone in male ME/CFS patients suggest complex interactions between the HPA axis and gonadal steroid production [309]. Further research specifically examining male ME/CFS patients is critically needed to elucidate androgen metabolism and its therapeutic potential.

9.3.6 Mechanisms of Sex Hormone Influence on ME/CFS

Sex hormones exert pervasive effects on immune function, energy metabolism, neurotransmitter systems, and autonomic regulation—all domains disrupted in ME/CFS. Understanding these mechanisms illuminates how hormonal dysregulation contributes to pathophysiology.

Immune Modulation by Sex Hormones

Estrogen generally enhances immune responses. It promotes B cell maturation and antibody production, enhances T helper 2 (Th2) responses, increases pro-inflammatory cytokine production in certain contexts, and potentially contributes to higher autoimmune disease prevalence in females. Progesterone exerts immunosuppressive effects. It promotes T regulatory cell (Treg) function essential for immune tolerance, suppresses Th1 inflammatory

responses, reduces pro-inflammatory cytokine production, and normally balances estrogen's immune-enhancing effects during the menstrual cycle.

Testosterone generally suppresses immune activation. It reduces B cell activity and antibody production, suppresses pro-inflammatory cytokine secretion, and potentially explains lower autoimmune disease rates in males. The loss of normal hormonal modulation of immune function—whether through estrogen-progesterone imbalance in females, DHEA/testosterone deficiency in both sexes, or altered receptor sensitivity—could permit the chronic immune activation documented in Chapter 7 to persist unchecked.

Neuroendocrine Integration

The sex-specific pathophysiology documented by Walitt et al. and Heng et al. likely reflects coordinated neuroendocrine-immune interactions rather than isolated hormonal effects [5, 311]. The hypothalamus and pituitary integrate signals from immune cytokines, metabolic hormones, and gonadal steroids to coordinate systemic responses. In ME/CFS, this integration appears fundamentally disrupted, with females showing stronger interferon signatures (potentially reflecting estrogen's enhancement of interferon responses) and males showing more balanced but still abnormal patterns (potentially reflecting testosterone's dampening effects on certain immune pathways).

The early menopause and gynecological abnormalities documented by Boneva et al. suggest that the HPG axis dysfunction may represent a form of neuroendocrine exhaustion analogous to the HPA axis hypofunction discussed earlier in this chapter [320]. Chronic immune activation and cytokine exposure may dysregulate both axes, creating a state of multi-system endocrine insufficiency despite the absence of structural gland failure.

⚠ Limitation 3: Sex-Specific Endocrine Findings: Limited Cohorts and Confounders

The sex-specific endocrine and immune patterns reported across studies have important epistemic boundaries:

- The Pipper et al. steroid analysis (n=97 total) has not been replicated, and subgroup sizes (particularly male severe ME/CFS) are too small for robust conclusions. The machine learning classifier lacked independent validation.
- The Boneva early menopause finding (mean 37.6 vs. 48.6 years) derives from a single CDC cohort; selection bias, recall bias, and confounders (medications, comorbidities) have not been excluded by independent replication.
- Hormonal differences between male and female ME/CFS patients do not establish that sex hormones *cause* the differing immune signatures—shared genetic susceptibility, sex-chromosome immune gene dosage, or other confounders may explain both.
- The inference that “ME/CFS may represent partially distinct disease processes in males and females” is a hypothesis generated from cross-sectional data, not a demonstrated conclusion.

9.4 Growth Hormone and IGF-1

Growth hormone (GH) and its primary mediator, insulin-like growth factor 1 (IGF-1), coordinate critical aspects of metabolism, body composition, and cellular function that directly relate to ME/CFS symptoms. The growth hormone axis regulates protein synthesis and muscle mass maintenance, lipolysis (fat breakdown) and glucose metabolism, bone density and connective tissue integrity, immune function and wound healing, and cognitive function and mood. GH/IGF-1 deficiency could therefore contribute to the muscle weakness, metabolic dysfunction, and cognitive impairment characteristic of ME/CFS.

9.4.1 Evidence for GH/IGF-1 Axis Dysfunction

Research on the GH/IGF-1 axis in ME/CFS has produced conflicting findings, likely reflecting the heterogeneity of patient populations and the complexity of growth hormone regulation. Some studies document clear abnormalities, while others find normal GH dynamics, suggesting that GH dysfunction characterizes a subset of ME/CFS patients rather than representing a universal feature.

Observation 55 (Reduced IGF-1 in ME/CFS Subset). Bennett et al. (1997) documented significantly lower serum IGF-1 levels in ME/CFS patients compared to healthy controls, accompanied by reduced nocturnal secretion of growth hormone. IGF-1 serves as the primary mediator of GH's anabolic effects, produced predominantly in the liver in response to GH stimulation and acting on peripheral tissues to promote protein synthesis, muscle growth, and metabolic regulation. Low IGF-1 despite normal or near-normal GH secretion suggests either hepatic resistance to GH or impaired liver function affecting IGF-1 synthesis.

However, contradicting these findings, other rigorous studies found no differences in basal IGF-1 or IGF-binding protein (IGFBP) levels between ME/CFS patients and controls, normal urinary growth hormone excretion, and similar GH responses to provocative testing. These inconsistencies highlight the challenge of identifying reliable biomarkers in a heterogeneous disease and suggest the need for subgroup stratification based on clinical phenotypes or other biomarkers.

9.4.2 Growth Hormone Treatment Trial

★ Achievement 6: Physiological but Limited Clinical Benefits of GH Treatment

Moorkens et al. (2000) conducted a randomized, double-blind, placebo-controlled trial of growth hormone treatment in ME/CFS patients with documented low IGF-1 levels [321]. The study involved 20 patients receiving 12 weeks of active treatment followed by a 9-month open-label phase. Results demonstrated clear physiological effects: mean serum IGF-1 increased from $173 \pm 46 \mu\text{g/L}$ to $296 \pm 89 \mu\text{g/L}$ ($p < 0.001$), fat-free mass significantly increased, and total body water increased, confirming the anabolic

effects of GH. However, clinical outcomes were mixed: quality of life measures did not show significant improvement overall, though notably, 4 patients resumed work after prolonged illness, suggesting substantial benefit in a subset.

The dissociation between clear physiological effects (increased IGF-1, improved body composition) and limited symptom improvement suggests that while GH deficiency may contribute to certain ME/CFS features (particularly muscle weakness and poor exercise tolerance), it is not the primary driver of fatigue, post-exertional malaise, or cognitive symptoms. This pattern aligns with the multi-system nature of ME/CFS pathophysiology, where correcting a single hormonal deficit proves insufficient to restore overall function.

9.4.3 Mechanisms and Clinical Implications

Several mechanisms could explain GH/IGF-1 axis dysfunction in ME/CFS, each with distinct therapeutic implications:

~ Hypothesis 6: Mechanisms of GH/IGF-1 Dysfunction

Multiple non-exclusive mechanisms may contribute to growth hormone axis dysfunction. Hypothalamic dysfunction may reduce GH-releasing hormone (GHRH) secretion, paralleling the HPA axis hypofunction discussed earlier. The NIH study documented temporal-parietal junction and broader brain abnormalities that could affect hypothalamic regulation [5]. Hepatic resistance to GH action may impair IGF-1 synthesis despite adequate GH secretion, potentially reflecting the mitochondrial dysfunction and oxidative stress documented in Chapter 6. Cytokine-mediated suppression may inhibit the GH axis, as chronic inflammation suppresses both GH secretion and IGF-1 synthesis while inducing IGF-1 resistance at target tissues. Finally, sleep disruption may reduce nocturnal GH pulses; the majority of daily GH secretion occurs during deep sleep, particularly slow-wave sleep, which is disrupted in ME/CFS.

Falsifiability: *weakly* — Falsified if GH stimulation tests show normal pituitary responsiveness and hepatic IGF-1 production is normal despite the proposed suppressive mechanisms

The clinical implications of these findings remain uncertain. GH treatment showed physiological effects but limited symptom benefit in the controlled trial, suggesting it may help selected patients but is not a universal solution. The expense, need for daily injections, and potential adverse effects (fluid retention, carpal tunnel syndrome, glucose intolerance) argue for restricting GH treatment to patients with documented IGF-1 deficiency and careful monitoring of both physiological parameters and functional outcomes. Alternative approaches targeting upstream causes (sleep improvement, inflammation reduction, mitochondrial support) might prove more effective than hormone replacement alone.

9.5 Insulin and Glucose Metabolism

Glucose metabolism abnormalities in ME/CFS connect endocrine dysfunction directly to the cellular energy deficits discussed in Chapter 6. Insulin regulates glucose uptake into cells, coordinates switching between glucose and fat oxidation, affects mitochondrial function and ATP production, and modulates inflammatory responses and immune function. Dysregulation of insulin signaling and glucose metabolism therefore has cascading effects on multiple systems compromised in ME/CFS.

9.5.1 Metabolic Syndrome and Insulin Resistance

★ Achievement 7: ME/CFS Association With Metabolic Syndrome

A population-based case-control study by Maloney et al. (2010) examined the relationship between ME/CFS and metabolic syndrome in Georgia [322]. The analysis revealed that ME/CFS patients were approximately 2-fold more likely to have metabolic syndrome compared to controls (odds ratio 2.12, 95% confidence interval 1.06–4.23). The key discriminating factors were increased waist circumference, elevated triglycerides, and higher fasting glucose. Notably, each additional metabolic syndrome component present associated with a 37% increase in the likelihood of having ME/CFS, demonstrating a dose-response relationship.

Metabolic syndrome comprises a cluster of abnormalities: abdominal obesity (increased waist circumference), elevated triglycerides, reduced HDL cholesterol, elevated blood pressure, and elevated fasting glucose or insulin resistance. The syndrome reflects underlying insulin resistance (reduced cellular responsiveness to insulin signaling) and predicts increased risk for type 2 diabetes, cardiovascular disease, and inflammatory conditions.

The association between ME/CFS and metabolic syndrome raises important mechanistic questions about causality. Does insulin resistance contribute to ME/CFS pathophysiology, or does ME/CFS-related inflammation, inactivity, and mitochondrial dysfunction lead to insulin resistance? Likely, bidirectional relationships exist, with each condition exacerbating the other in a self-reinforcing cycle.

9.5.2 Metabolic Phenotypes and Insulin Dynamics

Armstrong et al. (2021) employed comprehensive metabolomics to identify distinct ME/CFS subtypes with different metabolic signatures. One subtype (ME-M2) demonstrated elevated triglyceride and insulin levels despite normal glucose, reflecting low-grade lipid-induced insulin resistance. ME/CFS patients overall showed slightly elevated insulin and leptin (an adipose tissue hormone signaling energy status) and lower high molecular weight adiponectin (an anti-inflammatory adipokine that enhances insulin sensitivity).

~ Hypothesis 7: Peripheral Insulin Resistance With Central Deficits

ME/CFS may involve a paradoxical state of peripheral insulin resistance (reduced glucose uptake in muscle and adipose tissue) combined with inadequate glucose delivery or utilization in the central nervous system. This would explain the constellation of elevated peripheral insulin levels reflecting compensatory hyperinsulinemia, cerebral glucose hypometabolism documented by PET imaging, symptoms resembling hypoglycemia without documented low blood glucose, and the energy deficit despite apparently adequate systemic glucose availability [312, 313].

Falsifiability: *weakly* — Falsified if glucose clamp studies show normal peripheral insulin sensitivity combined with normal cerebral glucose uptake on FDG-PET imaging

9.5.3 Cerebral Glucose Hypometabolism

★ Achievement 8: Objective Evidence of Brain Energy Deficit

Positron emission tomography (PET) studies using fluorodeoxyglucose (FDG) tracer have documented reduced cerebral glucose metabolism in ME/CFS patients. Tirelli et al. (1998) identified significant glucose hypometabolism in the right mediofrontal cortex ($p=0.010$) and brainstem ($p=0.013$) in ME/CFS patients compared to controls, with moderate hypometabolism in the pons [312]. Siessmeier et al. (2003) employed observer-independent analysis in 26 ME/CFS patients and found that 12/26 (46%) showed hypometabolism bilaterally in the cingulate gyrus and adjacent mesial cortical areas, with 5 also demonstrating decreased orbitofrontal metabolism [313]. Importantly, hypometabolism correlated with anxiety and depression measures but not directly with fatigue severity, suggesting complex relationships between metabolic deficits and symptom expression.

△ Warning 5: Replication Status: Partially Replicated

Two independent PET studies (Tirelli 1998, $n=18$; Siessmeier 2003, $n=26$) both found cerebral hypometabolism, but in partially different brain regions. Both studies are old (pre-2005) with small samples and limited by PET methodology of that era. Modern replication with current PET technology and larger cohorts is needed.

The consistent documentation of reduced cerebral glucose uptake across independent studies using rigorous quantitative methods provides objective evidence of brain energy deficit in ME/CFS. The affected regions—brainstem nuclei, cingulate cortex, prefrontal areas—overlap substantially with regions showing functional abnormalities in the NIH study and with brain networks subserving autonomic control, attention, emotion regulation, and effort-based decision-making [5].

Several mechanisms could produce cerebral hypometabolism beyond simple insulin resistance:

~ Hypothesis 8: Mechanisms of Cerebral Hypometabolism

Multiple factors likely contribute to reduced brain glucose utilization. Cerebral hypoperfusion documented by SPECT imaging reduces glucose delivery to neurons. Glucose transporter dysfunction (GLUT1 at blood-brain barrier, GLUT3 in neurons) may impair glucose uptake even when delivery is adequate. Mitochondrial dysfunction in neurons reduces the capacity to metabolize glucose to ATP, causing glucose accumulation rather than utilization. Neuroinflammation with activated microglia alters brain energetics, as activated immune cells in the brain preferentially utilize glucose via glycolysis rather than oxidative phosphorylation. Finally, reduced neuronal activity secondary to other ME/CFS-related dysfunction may reduce metabolic demand, causing secondary hypometabolism as a consequence rather than cause of neurological symptoms.

Falsifiability: *weakly* — Falsified if cerebral perfusion is normal on SPECT, glucose transporters show normal expression, and neuronal mitochondria are intact in patients with documented hypometabolism

9.5.4 Hypoglycemia Symptoms Versus Orthostatic Intolerance

△ Warning 6: Misattribution of Orthostatic Symptoms to Hypoglycemia

Many ME/CFS patients report symptoms they attribute to “hypoglycemia”: nausea, lightheadedness and faintness, sweating and tremor, weakness and malaise, and cognitive impairment. However, studies examining ME/CFS patients during symptomatic episodes have shown that these symptoms frequently occur without documented hypoglycemia (blood glucose <70 mg/dL). Instead, the symptoms typically reflect orthostatic intolerance (inadequate blood pressure and cerebral perfusion upon standing or with prolonged upright posture, discussed extensively in Chapter 10). Critically, when orthostatic intolerance is treated effectively, the “hypoglycemia” symptoms often improve despite no specific glucose intervention.

This misattribution has important clinical implications. ME/CFS patients may consume frequent snacks or high-carbohydrate meals attempting to prevent “hypoglycemia,” potentially exacerbating insulin resistance and weight gain. Additionally, focusing on blood sugar management diverts attention from the actual orthostatic problem requiring different interventions (increased salt and fluid intake, compression garments, medications affecting blood pressure or blood volume).

9.5.5 Integration With Energy Metabolism Dysfunction

The glucose metabolism abnormalities documented in this section connect directly to the mitochondrial dysfunction and cellular energy deficits discussed in Chapter 6. Insulin resistance reduces glucose uptake into cells, limiting substrate availability for ATP production. Impaired mitochondrial function reduces the capacity to oxidize glucose efficiently, causing metabolic bottlenecks. The resulting cellular energy deficit triggers compensatory responses

including increased reliance on glycolysis (less efficient ATP production), activation of AMP-activated protein kinase (AMPK) signaling cellular energy stress, and metabolic shifts toward fat oxidation when glucose utilization fails.

These metabolic derangements help explain post-exertional malaise, as exertion depletes limited ATP stores that cannot be rapidly replenished due to impaired glucose metabolism and mitochondrial dysfunction. The delayed recovery characteristic of PEM reflects the slow restoration of cellular energy status when metabolic pathways remain compromised.

9.6 Melatonin and Circadian Rhythms

Circadian rhythm disruption represents a pervasive feature of ME/CFS that intersects with nearly every other aspect of pathophysiology discussed in this chapter. The circadian system coordinates temporal organization of physiological processes including the HPA axis cortisol rhythm, immune function cycling between pro- and anti-inflammatory states, metabolic switching between anabolic and catabolic metabolism, body temperature regulation and thermoregulation, and sleep-wake cycles and alertness patterns. Disruption of circadian timing thus has cascading effects across multiple systems already compromised in ME/CFS.

9.6.1 Objective Documentation of Circadian Disruption

Observation 56 (Actigraphy-Documented Circadian Abnormalities). Cambras et al. (2018) employed rigorous actigraphy monitoring to objectively document circadian rhythm abnormalities in ME/CFS patients [310]. This case-control study of 10 women with ME/CFS and 10 matched controls revealed that daily activity levels were significantly lower in ME/CFS patients, relative amplitude of the activity rhythm (difference between peak and nadir) was reduced, indicating flattened circadian variation, and stability of the activity rhythm across days was decreased, showing less consistent day-to-day patterns. Additionally, distal skin temperature showed lower nocturnal values in winter, suggesting impaired circadian regulation of peripheral blood flow and thermoregulation.

These findings demonstrate that circadian disruption in ME/CFS is not merely subjective patient reports of “feeling tired at the wrong times” but rather represents measurable alterations in the fundamental 24-hour organization of physiological functions. The reduced amplitude of activity rhythms parallels the flattened cortisol rhythm discussed earlier, suggesting a coordinated loss of circadian regulation across multiple output systems.

9.6.2 Sleep Architecture Versus Circadian Timing

An important conceptual distinction must be maintained between sleep architecture abnormalities (changes in sleep stage distribution, fragmentation, sleep efficiency) and circadian timing disruption (shifts in the phase or amplitude of 24-hour rhythms). ME/CFS patients exhibit both types of abnormalities, but they reflect different underlying mechanisms and require different therapeutic approaches.

Sleep architecture studies consistently document that ME/CFS patients experience longer sleep latency (time to fall asleep), more frequent awakenings during the night, more time in bed relative to total sleep time (reduced sleep efficiency), later and more variable wake times, irregular sleep patterns across days, and the paradox of unrefreshing sleep despite adequate or even prolonged total sleep duration. Children with ME/CFS often show continuous sleep exceeding 10 hours yet wake unrefreshed, indicating profound sleep dysfunction.

The circadian timing component involves altered phase relationships between sleep-wake cycles and other circadian outputs (body temperature, hormone secretion, immune function). Some ME/CFS patients show delayed sleep phase (natural sleep-wake times shifted later, resembling “night owl” patterns), while others exhibit irregular rhythms without clear 24-hour periodicity, and some maintain normal phase relationships but with reduced amplitude of rhythms.

9.6.3 Molecular Clock Dysfunction

~ Hypothesis 9: Clock Gene Dysregulation in ME/CFS

Emerging evidence suggests disruption at the molecular level of circadian clock gene expression. Genome-wide association studies (GWAS) have reported nominally significant associations with NPAS2 (neuronal PAS domain protein 2), a core clock gene. Transcriptomic analysis of ME/CFS patient samples showed 10-fold higher NPAS2 expression compared to controls and elevated expression of other circadian rhythm genes in peripheral blood mononuclear cells (PBMCs). Enrichment of CLOCK gene variants in ME/CFS patients with comorbid fibromyalgia and epigenetic changes in “circadian entrainment” pathways suggest heritable and acquired alterations in clock gene function.

Falsifiability: *weakly* — Falsified if repeat transcriptomic analysis fails to confirm NPAS2 or other clock gene alterations in independent ME/CFS cohorts with adequate sample size

The molecular clock operates as a transcriptional-translational feedback loop involving core clock genes (CLOCK, BMAL1, PER1/2/3, CRY1/2) that regulate their own expression with approximately 24-hour periodicity. These clock genes also control thousands of downstream genes involved in metabolism, immune function, and cellular processes, creating temporal coordination across organ systems. Disruption of clock gene function could therefore produce pleiotropic effects consistent with the multi-system nature of ME/CFS.

Inflammatory cytokines, particularly IL-1 β and TNF- α elevated in ME/CFS (Chapter 7), directly disrupt clock gene expression and alter circadian rhythms. This creates bidirectional interactions where immune dysfunction disturbs circadian regulation, while circadian disruption impairs proper immune function, perpetuating a self-reinforcing cycle.

9.6.4 Melatonin and Its Therapeutic Potential

Melatonin serves as both a marker and mediator of circadian rhythms, secreted by the pineal gland predominantly at night in response to darkness signals from the suprachiasmatic nucleus (SCN). Melatonin synchronizes peripheral clocks throughout the body, exerts direct

antioxidant and anti-inflammatory effects, modulates immune function and cytokine production, and facilitates sleep initiation though it is not primarily a sedative.

Limited evidence suggests altered melatonin production in ME/CFS, though findings have been inconsistent, likely reflecting the heterogeneity of circadian dysfunction patterns. Some patients show reduced melatonin amplitude, others exhibit phase shifts (melatonin rise at inappropriate times), while some maintain apparently normal melatonin profiles despite subjective circadian symptoms.

★ Achievement 9: Melatonin Treatment Benefits

Castro-Marrero et al. (2021) conducted a 16-week randomized, double-blind, placebo-controlled trial of melatonin (1 mg) plus zinc (10 mg) daily in 50 ME/CFS patients [323]. The intervention significantly reduced physical fatigue perception ($p < 0.05$) and improved the physical component summary score compared to placebo. Urinary 6-sulfatoxymelatonin (the primary melatonin metabolite) increased significantly in the treatment group ($p < 0.0001$), confirming adequate absorption and metabolism. Importantly, the intervention was safe and well-tolerated with no significant adverse effects.

This represents the first rigorous randomized controlled trial evidence that melatonin supplementation may provide symptomatic benefit in ME/CFS. However, several important caveats apply: the effect size was modest (improvement but not remission), the mechanism of benefit remains unclear (improved sleep, circadian resynchronization, anti-inflammatory effects, or antioxidant actions), and individual responses varied substantially (some patients benefited greatly, others not at all), and long-term efficacy and optimal dosing require further study.

△ Warning 7: Limitations of Melatonin Supplementation

While melatonin supplementation showed benefits in the Castro-Marrero trial, clinicians and patients should recognize important limitations. Melatonin primarily aids sleep initiation but does not address sleep maintenance (frequent awakenings), may temporarily improve symptoms without addressing underlying circadian dysfunction, risks masking underlying sleep disorders requiring different treatments (sleep apnea, restless legs syndrome), and exhibits substantial individual variation in absorption, metabolism, and response. Additionally, optimal timing of melatonin administration depends on the specific circadian phase abnormality (delayed, advanced, irregular), which typically requires formal assessment.

9.6.5 Circadian Disruption as an Integrative Mechanism

The circadian rhythm abnormalities documented in ME/CFS should not be viewed as isolated sleep problems but rather as disruption of a master regulatory system that normally coordinates multi-system physiology. Loss of circadian organization contributes to HPA axis dysfunction (flattened cortisol rhythm discussed earlier in this chapter), immune dysfunction (loss of circadian immune regulation), metabolic dysfunction (disrupted glucose homeostasis and lipid metabolism), autonomic dysfunction (altered cardiovascular circadian patterns

discussed in Chapter 10), and thermoregulatory dysfunction (impaired circadian temperature variation).

This integrative perspective suggests that interventions targeting circadian resynchronization—whether through melatonin, light therapy, behavioral scheduling, or other chronotherapeutic approaches—might provide broader benefits than expected from improving sleep alone. By restoring temporal coordination across multiple systems, circadian interventions could theoretically address multiple aspects of ME/CFS pathophysiology simultaneously. However, this hypothesis requires rigorous testing in well-designed clinical trials.

9.7 Integrated Endocrine-Metabolic Model

The endocrine abnormalities documented in this chapter do not represent independent, isolated dysfunctions but rather form an integrated network of disrupted hormonal regulation that mechanistically connects to the immune, neurological, metabolic, and cardiovascular dysfunction discussed in preceding chapters. Understanding these connections is essential for developing a coherent model of ME/CFS pathophysiology and identifying potential therapeutic targets.

9.7.1 Neuroendocrine-Immune Integration

The most critical integration involves bidirectional relationships between endocrine and immune systems. The HPA axis normally restrains immune activation through cortisol's anti-inflammatory effects, preventing excessive or prolonged inflammatory responses. In ME/CFS, blunted cortisol output and flattened circadian rhythm remove this restraint, permitting chronic low-grade inflammation to persist (Chapter 7). Conversely, chronic immune activation through pro-inflammatory cytokines (IL-1 β , IL-6, TNF- α) suppresses HPA axis function by increasing central glucocorticoid feedback sensitivity and impairing adrenal steroidogenic enzyme function.

The sex-specific patterns documented by Walitt et al. (2024) and Heng et al. (2025)—with females showing stronger interferon-driven inflammation and males exhibiting more balanced immune profiles—directly reflect sex hormone influences on immune function [5, 311]. Estrogen enhances type 2 interferon responses, while testosterone dampens multiple inflammatory pathways. The steroid hormone abnormalities documented by Pipper et al. (2024) thus contribute directly to the sex-specific immune dysregulation patterns [309].

These neuroendocrine-immune interactions create self-reinforcing pathological cycles. Inflammation disrupts HPA and HPG axis function, hormonal dysregulation permits unchecked inflammation, impaired cortisol rhythm disrupts circadian immune regulation, and circadian disruption further dysregulates neuroendocrine function. Breaking these cycles likely requires multi-targeted interventions addressing both endocrine and immune dysfunction simultaneously.

9.7.2 Endocrine-Metabolic Connections

The endocrine abnormalities documented in this chapter directly contribute to the cellular energy deficits discussed in Chapter 6. Cortisol supports hepatic gluconeogenesis and glucose availability; HPA dysfunction impairs this metabolic support. Thyroid hormone (T3) regulates mitochondrial biogenesis and oxidative phosphorylation efficiency; Low T3 Syndrome reduces cellular metabolic capacity. Insulin resistance and impaired glucose utilization limit substrate availability for ATP production; cerebral glucose hypometabolism reflects this at the brain level. Growth hormone supports protein synthesis and muscle metabolism; GH/IGF-1 deficiency contributes to muscle weakness and poor exercise tolerance.

The metabolic syndrome association documented by Maloney et al. (2010) indicates that ME/CFS involves not just cellular energy deficits but also systemic metabolic dysregulation affecting glucose homeostasis, lipid metabolism, and body composition [322]. This suggests that ME/CFS represents a form of “metabolic failure” spanning from mitochondrial dysfunction at the cellular level to whole-body insulin resistance and dysregulated energy partitioning.

9.7.3 Circadian Disruption as Central Organizing Principle

Circadian rhythm disruption may represent a central organizing principle connecting multiple aspects of ME/CFS pathophysiology. The suprachiasmatic nucleus (SCN) in the hypothalamus serves as the master circadian pacemaker, coordinating peripheral clocks throughout the body. Loss of this temporal coordination produces the constellation of abnormalities documented: flattened HPA axis cortisol rhythm, disrupted circadian immune function, altered metabolic switching between fed and fasted states, impaired cardiovascular circadian patterns (blood pressure, heart rate), and dysregulated body temperature variation.

The NIH study’s documentation of temporal-parietal junction dysfunction and broader brain abnormalities suggests that the neurological impairments in ME/CFS may affect hypothalamic function, disrupting the SCN’s ability to maintain circadian organization [5]. Inflammatory cytokines directly disrupt clock gene expression, creating a mechanistic link between immune activation and circadian dysfunction. The successful melatonin treatment trial by Castro-Marrero et al. (2021) supports the potential for circadian-targeted interventions [323].

⚠ Limitation 4: Circadian Disruption as Organising Principle: Primacy Not Established

While circadian disruption correlates with multiple ME/CFS features, its proposed role as a “central organising principle” has not been distinguished from the alternative that circadian changes are downstream consequences of other primary drivers:

- The actigraphy evidence (Cambras 2018, n=10 per group) is too small to generalise. No study has demonstrated that circadian disruption *precedes* immune or metabolic dysfunction in ME/CFS.
- Clock gene findings (NPAS2 overexpression) are from transcriptomic screens with-

out replication; nominal GWAS associations have not reached genome-wide significance.

- The melatonin trial (Castro-Marrero 2021) combined melatonin with zinc; the individual contributions cannot be separated, and the effect size was modest.
- Identical circadian flattening occurs in depression, fibromyalgia, and chronic pain—the pattern is not ME/CFS-specific and does not establish circadian disruption as a cause rather than a common consequence of chronic illness.

9.7.4 Sex as a Critical Biological Variable

The consistent 3–4:1 female-to-male prevalence ratio and the sex-specific pathophysiological patterns documented by multiple studies establish that biological sex is not merely a demographic variable but rather a critical determinant of disease susceptibility and expression. The mechanisms involve sex hormone modulation of immune responses (estrogen enhancing, testosterone dampening), sex-specific steroidogenic enzyme activity affecting stress hormone production, differential HPA and HPG axis regulation between sexes, and sex chromosome effects on immune gene expression (X chromosome contains numerous immune-related genes).

This recognition has profound implications for research and clinical care. Studies must stratify by sex to avoid obscuring sex-specific patterns, biomarker development should pursue sex-specific panels rather than assuming universal markers, and treatment trials should evaluate efficacy separately in male and female patients, as interventions effective in one sex may prove ineffective or even harmful in the other.

The early menopause finding by Boneva et al. (2015)—approximately 11 years earlier than controls—suggests that endocrine dysfunction may precede or contribute to ME/CFS onset rather than solely resulting from the disease [320]. This raises the possibility that hormonal interventions (hormone replacement therapy in appropriate contexts, androgen supplementation for documented deficiency) might prevent or mitigate disease progression in susceptible individuals, though this hypothesis requires prospective testing.

9.7.5 Clinical Implications and Therapeutic Considerations

The integrated endocrine-metabolic dysfunction documented in this chapter provides both biomarker opportunities and therapeutic targets. However, several principles should guide clinical application:

First, single-system hormonal interventions have shown limited efficacy. Growth hormone treatment produced physiological effects but modest symptom improvement, thyroid hormone supplementation helps some patients but not others, and simple hormone replacement does not address underlying regulatory dysfunction. This pattern suggests that ME/CFS involves coordinated multi-system dysregulation rather than simple deficiency states amenable to replacement therapy.

Second, addressing upstream drivers (inflammation, oxidative stress, mitochondrial dysfunction) may prove more effective than downstream hormone replacement. If chronic inflammation suppresses multiple endocrine axes simultaneously, anti-inflammatory interventions might restore coordinated hormonal function more effectively than replacing individual hormones. The partial success of melatonin supplementation, which has anti-inflammatory and antioxidant effects beyond its chronobiotic actions, supports this multi-targeted approach.

Third, interventions must be individualized based on specific endocrine phenotypes. The heterogeneity documented across studies indicates that not all ME/CFS patients exhibit the same endocrine abnormalities. Some show pronounced HPA dysfunction, others demonstrate primarily thyroid or sex hormone abnormalities, and metabolic syndrome patterns characterize a distinct subset. Precision medicine approaches matching interventions to individual endocrine profiles may prove superior to universal treatment protocols.

Fourth, sex-specific treatment strategies warrant investigation. The sex-specific steroid hormone profiles and immune patterns suggest that males and females may require different therapeutic approaches. Interventions effective in predominantly female cohorts may not generalize to male patients, and vice versa.

9.7.6 Future Research Directions

Critical gaps in understanding endocrine dysfunction in ME/CFS include longitudinal studies tracking hormonal changes from disease onset through chronic phases, mechanistic studies elucidating causal relationships between immune activation and endocrine dysfunction, biomarker validation studies assessing whether hormonal measurements can predict disease severity or treatment response, intervention trials testing multi-targeted approaches addressing both endocrine and immune dysfunction, and sex-stratified research examining whether male and female ME/CFS represent partially distinct diseases requiring different treatments.

The endocrine system provides an attractive therapeutic target because hormones are measurable, hormone replacement therapies already exist for many deficiency states, and endocrine interventions have established safety profiles when properly monitored. However, the limited success of single-hormone interventions to date indicates that simplistic approaches will not suffice. Future therapeutic development must embrace the complexity of multi-system dysregulation, targeting coordinated restoration of neuroendocrine-immune-metabolic integration rather than isolated hormone replacement.

10 Cardiovascular Dysfunction

Cardiovascular abnormalities are pervasive in ME/CFS and contribute substantially to disability, particularly through exercise intolerance and orthostatic symptoms. The 2024 NIH deep phenotyping study by Walitt et al. provided rigorous documentation of cardiopulmonary exercise testing abnormalities, including reduced peak oxygen consumption and chronotropic incompetence, establishing objective physiological correlates of the subjective exercise intolerance reported by patients [5].

10.1 Cardiac Function

10.1.1 Exercise Testing Abnormalities

Cardiopulmonary exercise testing (CPET) provides objective measurement of integrated cardiovascular, pulmonary, and metabolic function during physical exertion. CPET findings in ME/CFS represent some of the most reproducible objective abnormalities documented in the illness.

Cardiopulmonary Exercise Testing (CPET) Methodology

CPET involves graded exercise (typically on a cycle ergometer or treadmill) with continuous measurement of oxygen consumption (VO_2 , the volume of oxygen extracted from inspired air per unit time), carbon dioxide production (VCO_2 , the volume of CO_2 expired), and their ratio expressed as the respiratory exchange ratio ($\text{RER} = \text{VCO}_2 / \text{VO}_2$), which indicates fuel substrate utilization. Simultaneously, the system records minute ventilation (VE, total air volume breathed per minute), heart rate via continuous electrocardiographic monitoring, periodic blood pressure measurements, and work rate (power output in watts or treadmill speed and grade).

Testing continues until volitional exhaustion or limiting symptoms. Criteria for maximal effort include an RER exceeding 1.10, achievement of age-predicted maximal heart rate, or a plateau in VO_2 despite increasing work rate.

Key NIH Deep Phenotyping CPET Findings

The Walitt et al. study documented several critical cardiopulmonary abnormalities in PI-ME/CFS patients [5]:

Reduced Peak Oxygen Consumption ($\text{VO}_{2\text{peak}}$) Peak VO_2 represents maximal aerobic capacity and integrates cardiac output, oxygen delivery, and peripheral oxygen extraction. P/ME/CFS patients demonstrated significantly reduced $\text{VO}_{2\text{peak}}$ compared to matched healthy controls [5]. This reduction indicates impaired aerobic capacity beyond what deconditioning alone would predict; ME/CFS patients showed greater deficits than sedentary controls matched for activity level, with reductions typically ranging from 15–30% below predicted values [1, 79]. More severely affected patients show greater reductions. The finding correlates with functional limitation and disability, providing objective confirmation of patient-reported exercise intolerance.

Chronotropic Incompetence Chronotropic incompetence refers to an inadequate heart rate response to exercise [5]:

- ME/CFS patients fail to achieve age-predicted maximal heart rate
- Heart rate rise is blunted relative to work rate increases
- Chronotropic index (proportion of heart rate reserve used) is reduced

Chronotropic incompetence limits cardiac output augmentation during exercise, as cardiac output = heart rate $\times$ stroke volume. Without adequate heart rate increase, oxygen delivery to exercising muscles is compromised.

Mechanisms of Chronotropic Incompetence Several mechanisms have been proposed; their relative contributions in ME/CFS remain under investigation:

1. **Parasympathetic excess:** Sustained vagal tone preventing heart rate acceleration—supported by HRV findings showing altered autonomic balance [97]
2. **Sympathetic dysfunction:** Impaired catecholamine release or receptor sensitivity
3. **Sinoatrial node dysfunction:** Intrinsic pacemaker abnormality (hypothesized but not directly demonstrated in ME/CFS)
4. **G-protein-coupled receptor (GPCR) autoantibodies:** A growing body of evidence implicates autoantibodies targeting G-protein-coupled receptors in ME/CFS cardiovascular dysfunction. Loebel et al. first documented elevated autoantibodies against beta-adrenergic (β_1 , β_2) and muscarinic cholinergic (M3, M4) receptors in ME/CFS patients [71], findings subsequently validated in Swedish cohorts including cerebrospinal fluid samples [72].

The cardiovascular effects of these autoantibodies are multifaceted. Beta-adrenergic receptor autoantibodies may exert either agonistic effects (causing inappropriate receptor activation) or antagonistic effects (blocking normal catecholamine signaling), depending on the specific antibody epitope and receptor subtype. Muscarinic receptor autoantibodies similarly can enhance or impair parasympathetic signaling to the heart and vasculature. This bidirectional potential explains why the same class of autoantibodies might produce different phenotypes across patients.

Levels of vasoregulative GPCR autoantibodies correlate with symptom severity, autonomic dysfunction, and disability in ME/CFS [208]. The correlation with autonomic measures supports a direct pathophysiological role rather than an epiphenomenon of

chronic illness. Beta-2 adrenergic receptor autoantibodies appear particularly relevant to cardiovascular symptoms, with immunoadsorption targeting these antibodies showing preliminary efficacy in post-COVID ME/CFS [118]. BC007, a DNA aptamer that neutralizes GPCR autoantibodies, has shown promise in case reports for improving fatigue and microcirculatory function [10]. Three complementary approaches to the autoantibody hypothesis—immunoadsorption, BC007, and daratumumab—are in active clinical trials (Appendix I, Section I.4).

5. **Central nervous system dysfunction:** Impaired autonomic outflow from brainstem centers
6. **Gut-mediated vagal impairment (Wirth-Scheibenbogen 2025):** Recent work by Wirth and Scheibenbogen (2025) provides mechanistic support for gut-mediated vagal dysfunction in ME/CFS. Butyrate enhances enterochromaffin cell serotonin production [14], and enterochromaffin serotonin activates vagal afferents via 5-HT₃ receptors [13, 228]. Since ME/CFS patients show butyrate deficiency and gut dysbiosis [227], reduced enterochromaffin serotonin could impair vagal afferent signaling, potentially weakening efferent vagal tone to the heart. This pathway may contribute to the chronotropic incompetence and reduced heart rate variability observed in ME/CFS patients. While no direct evidence yet links this pathway to cardiac chronotropy in ME/CFS, it offers a testable hypothesis for the gut-cardiac connection. See Section 11.1.4 of Chapter 11 for the full evidence chain and Section 8.1.3 of Chapter 8 for the neurophysiological implications.

The finding of chronotropic incompetence, combined with reduced HRV and abnormal baroreflex sensitivity [97], indicates autonomic dysfunction affecting cardiac pacing.

⚠ **Limitation 1: Primary Site of Autonomic Dysfunction Unresolved**

Current evidence establishes that autonomic dysfunction affects cardiac pacing in ME/CFS but does not determine whether the primary site of dysfunction is central (brainstem outflow), peripheral (sinoatrial node, nerve fibers), autoimmune (GPCR autoantibodies), or gut-mediated (enterochromaffin-vagal pathway). These mechanisms are not mutually exclusive, and their relative contributions likely vary across patients. Resolving this requires intervention studies targeting each mechanism independently.

Two-Day CPET Protocol

A particularly informative methodology involves repeat CPET on consecutive days:

Rationale Single CPET testing may not capture the distinctive post-exertional deterioration characteristic of ME/CFS. Two-day protocols assess recovery capacity and reproducibility of maximal effort.

Findings in ME/CFS

- **Day 1:** Reduced but measurable aerobic capacity

- **Day 2:** Further significant reductions in VO_2 peak, anaerobic threshold, and work capacity
- **Healthy controls:** Reproduce or slightly improve Day 1 performance
- **Magnitude:** ME/CFS patients show 10–25% decline on Day 2 [1, 79]

This failure to reproduce exercise capacity distinguishes ME/CFS from other fatiguing conditions and reflects the pathognomonic post-exertional malaise. A systematic review of two-day CPET studies confirmed significant reductions in work capacity and oxygen consumption on Day 2, supporting this protocol as an objective marker of PEM [79].

Mechanisms of Day 2 Decline

- Delayed recovery of metabolic substrates
- Persistent inflammatory activation
- Autonomic dysfunction exacerbation
- Mitochondrial damage from oxidative stress
- Central nervous system effects (increased perceived exertion)

Anaerobic Threshold

The anaerobic threshold (AT, also called ventilatory threshold or lactate threshold) represents the exercise intensity at which anaerobic metabolism begins to supplement aerobic energy production:

- **Reduced AT in ME/CFS:** Occurs at lower work rates and VO_2 levels
- **Early lactate accumulation:** Muscles rely on anaerobic glycolysis sooner
- **Implications:** Limited sustainable activity before symptom exacerbation
- **Mechanism:** Reflects impaired oxygen delivery, mitochondrial dysfunction, or both

The reduced AT has practical implications: patients exceed their aerobic capacity during activities that healthy individuals perform entirely aerobically, leading to metabolic stress and symptom generation.

Ventilatory Efficiency

Ventilatory efficiency describes how effectively ventilation eliminates CO_2 , typically expressed as the VE/VCO_2 slope:

- **Increased VE/VCO_2 slope:** More ventilation required per unit CO_2 eliminated
- **Causes:** Ventilation-perfusion mismatch, increased dead space, hyperventilation
- **Consequences:** Dyspnea at lower work rates, earlier exercise termination
- **ME/CFS findings:** Variable; some patients show ventilatory inefficiency

10.1.2 Cardiac Output and Stroke Volume

Cardiac output (CO) determines oxygen delivery capacity and is the product of heart rate and stroke volume.

Preload Failure Hypothesis

Multiple lines of evidence support inadequate cardiac preload (ventricular filling) as a contributor to ME/CFS cardiovascular dysfunction:

- **Reduced end-diastolic volume:** Less blood fills the ventricles during diastole
- **Decreased stroke volume:** By Frank-Starling mechanism, reduced preload produces smaller stroke volume
- **Compensatory tachycardia at rest:** Heart rate increases to maintain resting cardiac output; however, during exercise, chronotropic incompetence prevents further adequate heart rate augmentation, creating a ceiling effect
- **Exercise limitation:** The combination of low stroke volume and inadequate heart rate response severely limits cardiac output augmentation during exertion

Evidence for Preload Failure

- Echocardiographic studies showing reduced left ventricular end-diastolic volume [324]
- Correlation with blood volume measurements [324]

Supine Hemodynamic Abnormalities While cardiovascular dysfunction in ME/CFS is most apparent during orthostatic stress, some patients demonstrate abnormalities even at rest in the supine position. Newton et al. documented reduced cardiac volumes on cardiac MRI that correlated with blood volume deficits rather than deconditioning, with end-diastolic volume, end-systolic volume, and end-diastolic wall mass all significantly reduced [324]. Critically, these reductions showed no correlation with disease duration, arguing against deconditioning as the primary mechanism.

Reduced resting cardiac output in the supine position has been reported in some ME/CFS cohorts, with the magnitude of reduction correlating with symptom severity [324]. This finding suggests that the cardiovascular impairment is not solely a failure of orthostatic compensation but reflects a baseline deficit in cardiac filling and output. Patients with more severe supine abnormalities tend to show greater decompensation during orthostatic challenge, as they have less hemodynamic reserve to mobilize when gravitational stress is applied.

Reduced Blood Volume

Blood volume deficits are well-documented in ME/CFS [280, 324, 325]:

- **Plasma volume:** Reduced by 10–20% in most studies [280]
- **Red cell mass:** Variable findings; may be proportionally reduced or relatively preserved
- **Total blood volume:** Typically 10–15% below normal [324]
- **Correlation with symptoms:** Lower blood volume correlates with worse orthostatic intolerance and fatigue [324]

Mechanisms of Hypovolemia

- **RAAS dysfunction:** Impaired aldosterone response to hypovolemia
- **Natriuretic peptide elevation:** Promoting sodium and water excretion
- **Reduced erythropoietin:** Leading to mild anemia in some patients
- **Capillary leak:** Increased vascular permeability shifting fluid to interstitium
- **Inadequate fluid intake:** Secondary to nausea or other symptoms

Venous Pooling

Excessive venous pooling in dependent body parts reduces venous return:

- **Lower extremity pooling:** Blood accumulates in leg veins during standing
- **Splanchnic pooling:** Blood redistributes to abdominal vasculature
- **Impaired venoconstriction:** Venous tone fails to increase appropriately
- **Consequences:** Reduced cardiac preload, orthostatic symptoms

10.1.3 Cardiac Biomarkers

Troponin

Cardiac troponins (cTnI, cTnT) are released from damaged cardiomyocytes:

- **Baseline levels:** Generally normal in ME/CFS
- **Post-exercise:** Whether mild elevations occur after exertion in ME/CFS is not yet established in the literature
- **Clinical significance:** Cardiac biomarkers are not currently validated as ME/CFS diagnostic markers

BNP and NT-proBNP

B-type natriuretic peptide (BNP) and its N-terminal fragment are released in response to cardiac wall stress:

- **Findings in ME/CFS:** Variable; some studies report mild elevations [326]
- **Mechanism:** May reflect right heart strain from pulmonary issues or left ventricular stress
- **Correlation:** Elevated BNP correlates with reduced cardiac volumes in ME/CFS [326]
- **Clinical utility:** Not established as ME/CFS biomarker

Cardiac Structure and Function

Echocardiographic studies report variable findings, with evidence for subclinical dysfunction in some patients [324]:

- **Reduced cardiac volumes:** Smaller left ventricular end-diastolic volume correlating with plasma volume deficits [324]
- **Diastolic dysfunction:** Some studies report impaired ventricular relaxation
- **Reduced contractile reserve:** Limited ability to augment function during stress echocardiography (preliminary data)
- **Strain imaging:** Speckle tracking echocardiography may detect subtle abnormalities not apparent on conventional imaging (requires further study)

These findings suggest that cardiac abnormalities in ME/CFS are primarily functional consequences of hypovolemia and autonomic dysfunction rather than primary myocardial disease.

10.2 Vascular Dysfunction

10.2.1 Endothelial Dysfunction

The vascular endothelium regulates vascular tone, coagulation, and inflammation. Endothelial dysfunction is increasingly recognized in ME/CFS.

Nitric Oxide Bioavailability

Nitric oxide (NO) is a critical vasodilator produced by endothelial NO synthase (eNOS):

- **Altered NO metabolism during exercise:** ME/CFS patients show significantly elevated nitric oxide metabolites (plasma nitrate up to ~295% above controls) during maximal exercise, consistent with abnormal vascular regulation rather than simple deficiency [327]
- **Increased NO scavenging:** Oxidative stress may inactivate NO

- **eNOS uncoupling:** Dysfunctional enzyme produces superoxide instead of NO
- **Consequences:** Impaired vasodilation, increased vascular resistance

Flow-Mediated Dilation

Flow-mediated dilation (FMD) measures endothelium-dependent vasodilation of the brachial artery following brief ischemia:

- **Reduced FMD in ME/CFS:** Multiple studies report impaired endothelium-dependent dilation, with peripheral endothelial dysfunction found in 51% of ME/CFS patients versus 20% of healthy controls [177]
- **Correlation:** Associated with disease severity and severity of immune symptoms [177]
- **Mechanism:** Reflects reduced NO bioavailability, elevated adhesion molecules, or chronic inflammatory state [328]

Inflammatory Markers

Endothelial inflammation contributes to dysfunction [328]:

- **Elevated adhesion molecules:** ICAM-1, VCAM-1, E-selectin [328]
- **Increased inflammatory cytokines:** IL-6, TNF- α affect endothelial function
- **Oxidative stress markers:** Indicate endothelial damage
- **Circulating endothelial cells:** May be elevated, indicating endothelial injury

Proteomic Evidence for Endothelial Dysfunction and Tissue Hypoxia

Hoel et al. (2026) [191] provided large-scale proteomic confirmation of endothelial and vascular abnormalities in ME/CFS. Several proteins linked to microcirculatory and hypoxia responses were altered:

- **Reduced vimentin (VIM):** An intermediate filament protein critical for endothelial cell structure and mechanotransduction; its reduction may indicate impaired endothelial integrity
- **Reduced ANTXR2 (CMG2):** Also known as capillary morphogenesis gene 2, this receptor regulates angiogenesis and endothelial cell chemotaxis; its reduction suggests impaired vascular repair capacity
- **Reduced CXADR:** The coxsackievirus and adenovirus receptor mediates endothelial responses to fluid shear stress; its decrease may reflect impaired vascular mechanosensing
- **Reduced ATP5IF1:** ATP synthase inhibitory factor 1 protects against mitochondrial ATP hydrolysis during hypoxia; its reduction implies compromised mitochondrial stress tolerance in endothelial cells

- **Elevated LIPG:** Endothelial cell-derived lipase, involved in lipoprotein metabolism and vascular biology, was increased—consistent with immune activation and endothelial stress
- **Elevated FAM20A:** A hematopoietic signaling factor elevated in ME/CFS, suggesting bone marrow-endothelial signaling disruption

Additionally, ME/CFS patients showed increased WASF3 expression, a target of the hypoxia-regulated transcription factor HIF-1 α , directly linking the proteomic signature to tissue hypoxia. Together, these findings support a model in which endothelial dysfunction, impaired vascular repair, and chronic tissue hypoxia form an interconnected pathological axis in ME/CFS.

Ephrin Signaling: A Novel Vascular Pathway

Ligand-receptor interaction analysis from the Hoel et al. proteomics data [191] revealed concordant changes in multiple members of the Ephrin signaling family: four Ephrin subfamily A receptors (EPHA1, EPHA2, EPHA4, EPHA7) and their EFNA ligands showed coordinated alterations in ME/CFS. Ephrin signaling regulates vascular development, endothelial cell migration, and angiogenesis. Disrupted Ephrin-Eph signaling had been previously suggested in ME/CFS based on earlier proteomic studies [329], and the Hoel et al. findings provide independent confirmation across a different cohort and analytical platform. This pathway represents a potential mechanistic link between immune activation (Ephrin signaling also modulates immune cell migration and T cell activation) and the vascular dysfunction observed in ME/CFS.

Several 2025–2026 studies extend the endothelial dysfunction evidence. Retinal vessel analysis revealed reduced venular flicker-induced dilation in ME/CFS and post-COVID patients, providing a non-invasive measure of persistent microvascular dysfunction [330] (Appendix I, Section I.10.3). Azcue et al. (2025) documented small fibre neuropathy in both ME/CFS and post-COVID patients using corneal confocal microscopy and electrochemical skin conductance [331] (Section I.10.2). The MCAM multi-site study confirmed that autonomic dysfunction (assessed by COMPASS-31 and NASA lean test) is pervasive across diverse ME/CFS cohorts [332] (Section I.10.1).

10.2.2 Blood Volume Abnormalities

Blood volume deficits are among the most consistently documented abnormalities in ME/CFS (see also Section 10.1.2 for impact on cardiac preload). This section expands on measurement methods and pathophysiological mechanisms.

Measurement and Magnitude

- **Measurement methods:** Radioisotope dilution (gold standard), carbon monoxide re-breathing, dye dilution
- **Plasma volume:** Typically 10–20% below predicted [280]

- **Red cell mass:** Variable; some studies report proportional reduction, others find preserved red cell mass with disproportionate plasma volume loss
- **Total blood volume:** 10–15% below normal in most studies [324]
- **Hemoglobin/hematocrit:** May appear normal or elevated due to hemoconcentration

Mechanisms of Volume Depletion

Renin-Angiotensin-Aldosterone System Dysfunction Studies document a paradoxical RAAS response in POTS and ME/CFS, with elevated angiotensin II despite hypovolemia [325, 333]:

- Blunted aldosterone response to hypovolemia [325]
- Impaired sodium retention leading to inappropriate natriuresis
- Elevated angiotensin II may contribute to symptoms through vasoconstriction [333]

Natriuretic Peptide Effects

- Elevated ANP or BNP promoting sodium/water excretion
- May result from atrial stretch due to cardiac filling abnormalities

Capillary Permeability

- Increased vascular permeability shifting fluid to interstitium
- May be inflammation-mediated (cytokines increase endothelial permeability)
- Could explain peripheral edema in some patients despite intravascular hypovolemia

10.2.3 Arterial Stiffness and Vascular Compliance

Beyond endothelial function, arterial mechanical properties influence cardiovascular regulation. Pulse wave velocity (PWV), a measure of arterial stiffness, affects blood pressure regulation through its impact on baroreceptor function. The relationship between arterial stiffness and ME/CFS has not been extensively studied, but related conditions provide insight.

In hypermobile Ehlers-Danlos syndrome (hEDS), which frequently co-occurs with ME/CFS (Section 5.6.11), central pulse wave velocity is significantly lower than controls (4.73 m/s versus normal values), indicating excessive arterial elasticity [334]. This increased compliance paradoxically impairs blood pressure regulation: stretch receptors in vessel walls (baroreceptors) cannot accurately detect pressure changes when arterial walls are too compliant. The result is impaired baroreflex function and orthostatic intolerance despite (or because of) increased rather than decreased arterial elasticity.

Whether ME/CFS patients without connective tissue disorders show altered arterial stiffness remains unclear. Chronic inflammation typically increases arterial stiffness over time, while autonomic dysfunction could affect vascular smooth muscle tone. The interaction between

these competing influences likely varies across patient subgroups. Pulse wave velocity measurement is non-invasive and could provide additional phenotyping data, though its clinical utility in ME/CFS management has not been established.

10.2.4 Microcirculation

Capillary Perfusion

The microcirculation delivers oxygen and nutrients to tissues and removes metabolic waste:

- **Red cell deformability:** Impaired RBC flexibility may impede capillary transit [180]
- **Microclot obstruction:** Amyloid-resistant fibrinoid microclots may occlude microcapillaries, impairing perfusion [166]

Oxygen Extraction

Peripheral oxygen extraction findings in ME/CFS appear contradictory across studies, likely reflecting patient heterogeneity or methodological differences:

- **Contradictory findings:** Some studies report widened arteriovenous O₂ difference (increased extraction compensating for reduced cardiac output), while others report impaired extraction consistent with mitochondrial limitation; these contradictions may reflect distinct patient subgroups
- **Near-infrared spectroscopy:** Abnormal muscle oxygenation kinetics during exercise and delayed recovery are documented by NIRS, consistent with both delivery and utilization abnormalities [1]

Tissue Hypoxia

Inadequate oxygen delivery produces tissue hypoxia:

- **Muscle hypoxia:** Contributes to weakness and post-exertional symptoms
- **Cerebral hypoperfusion:** Causes cognitive dysfunction (see Chapter 8)
- **Lactate accumulation:** Results from anaerobic metabolism
- **Symptom generation:** Hypoxia-sensitive nociceptors may trigger pain

Cerebral Blood Flow During Orthostatic Stress

While tissue hypoxia affects multiple organs, the brain is particularly vulnerable to perfusion deficits during orthostatic challenge. Van Campen and colleagues have systematically characterized cerebral blood flow (CBF) abnormalities in ME/CFS through a series of rigorous studies using transcranial Doppler during tilt-table testing [172, 253, 254, 4].

★ Achievement 1: Near-Universal CBF Decline in ME/CFS

Van Campen et al. [172] demonstrated that ME/CFS patients show reduced cerebral blood flow during head-up tilt testing even in the absence of hypotension or tachycardia. The findings are striking in their consistency across orthostatic phenotypes (percentages represent distinct, non-overlapping subgroups stratified by vital sign response):

- 82% of patients with normal HR/BP showed abnormal CBF reduction
- 98% of patients with delayed orthostatic hypotension showed abnormal CBF
- 100% of patients meeting POTS criteria showed abnormal CBF
- End-tilt CBF reduction: **26% in ME/CFS vs. 7% in controls** (3.7-fold greater)

Abnormal CBF reduction thus occurs across all orthostatic presentations—even in patients with entirely normal vital signs. In the largest study to date (n=534), **91% of ME/CFS patients** with normal HR and BP responses demonstrated abnormal cardiac output and CBF reduction during tilt [4], indicating that standard orthostatic vital signs miss the primary pathology in most patients.

△ Warning 1: Replication Status: Single Research Group

All van Campen CBF studies originate from a single Dutch referral center using the same tilt-table protocol and transcranial Doppler methodology [172, 253, 254, 4]. While internally consistent across multiple publications (n=107 to n=534), no independent research group has replicated these findings using equivalent methodology. The sequential publication pattern from one center may give the appearance of independent replication but represents a single group's cohort. Independent replication with standardized protocols at other centers is essential. See also the related CBF findings in Chapter 8 (Achievement 8.1.2).

Clinical Implications of CBF Findings The cognitive symptoms during orthostatic stress—including brain fog, difficulty concentrating, and word-finding problems—correlate directly with the degree of cerebral hypoperfusion [254]. Patients often report that cognitive function worsens progressively during prolonged standing and improves rapidly upon assuming a recumbent position. This positional dependence of cognitive symptoms provides clinical evidence for the cerebrovascular contribution to ME/CFS neurological dysfunction.

Observation 57 (Impaired CBF Recovery). CBF reduction persists even after returning to supine position. Van Campen et al. [253] documented CBF reduction of −29% at end-tilt, improving to only −16% post-tilt. The degree of recovery correlated with disease severity rather than hemodynamic parameters, suggesting the CBF abnormality reflects intrinsic cerebrovascular or metabolic dysfunction rather than simple hemodynamic failure.

Absence of Compensatory Vasodilation A particularly significant finding is the near 1:1 relationship between cardiac output reduction and CBF reduction in ME/CFS patients [4]. In healthy individuals, reduced cardiac output triggers compensatory cerebral vasodilation to maintain brain perfusion. The absence of this compensation in ME/CFS suggests possible

endothelial dysfunction affecting cerebrovascular autoregulation. This may represent a critical vulnerability: the brain cannot protect itself from systemic hemodynamic perturbations.

Mechanisms of Cerebral Hypoperfusion Multiple mechanisms likely contribute to orthostatic cerebral hypoperfusion in ME/CFS. Evidence strength varies: **(documented)** = directly measured in ME/CFS studies; **(inferred)** = logically derived from observed relationships; **(hypothesized)** = proposed mechanism not yet directly tested.

- **Reduced cardiac output (documented):** Preload failure and chronotropic incompetence limit systemic perfusion pressure; directly measured in tilt studies showing parallel CO and CBF reduction [4] (see Section 10.1.2)
- **Impaired cerebral autoregulation (inferred):** Failure of compensatory vasodilation during reduced perfusion pressure; inferred from near 1:1 CO-CBF relationship where healthy controls show compensatory vasodilation [4]
- **Endothelial dysfunction (hypothesized):** May impair nitric oxide-mediated vasodilation; suggested by absence of compensatory response but not directly measured in CBF studies
- **Autonomic dysregulation (documented):** Impaired sympathetic vasoconstriction in peripheral vascular beds allows excessive venous pooling; documented via HRV and catecholamine studies (see Chapter 8 Section 8.2.1)
- **Blood volume deficit (documented):** Reduced circulating volume exacerbates orthostatic hemodynamic stress; documented in multiple studies showing 10–15% blood volume reduction (see Section 10.2.2)

In mast cell disorder patients, Novak et al. documented 20–24% reduction in orthostatic cerebral blood flow velocity using transcranial Doppler [173]. Given the substantial overlap between mast cell activation and ME/CFS, histamine-mediated vasodilation during orthostatic stress may contribute to cerebral hypoperfusion in some patients. The combination of reduced blood volume, impaired vasoconstriction, and potentially histamine-induced vasodilation creates multiple mechanisms converging on inadequate cerebral perfusion during upright posture.

Integration with Selective Energy Dysfunction Hypothesis The profound and consistent CBF reduction during orthostatic challenge exemplifies the broader pattern of *preserved baseline function with impaired challenge response* characteristic of ME/CFS (see Chapter 6 Section 6.3). Resting cerebral perfusion may be adequate, but the system cannot maintain CBF during the increased demand of orthostatic stress. The brain—with its high energy demands and critical dependence on continuous perfusion—may serve as the “canary in the coal mine” for systemic energy coordination dysfunction.

See Chapter 8 Section 8.1.2 for discussion of brain-centric pathophysiology and the role of CBF abnormalities in the broader ME/CFS disease model.

10.3 Blood Pressure Regulation

Blood pressure dysregulation is common in ME/CFS and manifests as various orthostatic disorders.

10.3.1 Orthostatic Hypotension

Definition 10.1 (Orthostatic Hypotension). A sustained reduction in systolic blood pressure ≥ 20 mmHg or diastolic blood pressure ≥ 10 mmHg within 3 minutes of standing. *Initial orthostatic hypotension* refers to a brief BP drop within the first 15 seconds, which is common in ME/CFS.

Orthostatic hypotension occurs in a subset of ME/CFS patients, presenting with lightheadedness, visual disturbances, weakness, and syncope. Contributing mechanisms include autonomic failure, hypovolemia, and medications.

10.3.2 Neurally Mediated Hypotension

Definition 10.2 (Neurally Mediated Hypotension). Also called vasovagal syncope or neurocardiogenic syncope. Characterized by paradoxical vasodilation and bradycardia during prolonged standing, triggered when vigorous ventricular contraction of an underfilled heart activates vagal reflexes. Presents as delayed BP drop after 10+ minutes of standing, with prodromal nausea, diaphoresis, and pallor preceding syncope.

Diagnosis is confirmed by head-up tilt table testing, which can provoke the characteristic hemodynamic response under controlled conditions.

10.3.3 Postural Orthostatic Tachycardia Syndrome (POTS)

Definition 10.3 (Postural Orthostatic Tachycardia Syndrome). A syndrome characterized by excessive heart rate increase upon standing without significant blood pressure drop. According to the 2015 Heart Rhythm Society expert consensus [99], diagnostic criteria include: heart rate increase ≥ 30 bpm (or ≥ 40 bpm in adolescents) within 10 minutes of standing, absence of orthostatic hypotension, symptoms of orthostatic intolerance, and symptom duration exceeding 6 months.

Prevalence in ME/CFS

- Estimated 25–50% of ME/CFS patients meet POTS criteria [98]
- Substantial symptom overlap between conditions
- May represent overlapping or related conditions
- Similar pathophysiological mechanisms

POTS is one component of the “Septad” framework of frequently co-occurring conditions in ME/CFS (Section 5.6.11). The interplay between dysautonomia, mast cell activation, EDS, and other Septad components suggests shared pathophysiological mechanisms warranting comprehensive evaluation.

POTS Subtypes

Different pathophysiological mechanisms produce similar clinical phenotypes:

Neuropathic POTS Neuropathic POTS results from partial autonomic neuropathy affecting lower extremity vasoconstriction, leading to blood pooling in the legs during standing. This subtype is associated with small fiber neuropathy (SFN) and may be autoimmune in some cases. SFN specifically affects the small-diameter autonomic nerve fibers that innervate blood vessels and sweat glands, and its prevalence in POTS has been confirmed through skin biopsy studies demonstrating reduced intraepidermal nerve fiber density [335].

Small Fiber Neuropathy in POTS and ME/CFS The connection between SFN and cardiovascular dysautonomia is increasingly recognized. Azcue et al. demonstrated that ME/CFS patients show heat response latencies indicating C-fiber denervation, with 31% meeting POTS criteria and 34% showing non-length-dependent SFN patterns [335]. This non-length-dependent pattern (affecting proximal and distal sites equally) suggests autoimmune or inflammatory etiology rather than length-dependent degeneration seen in metabolic neuropathies.

The autonomic consequences of SFN in ME/CFS are substantial. Damaged sympathetic sudomotor fibers contribute to temperature dysregulation, while damaged vasomotor fibers impair the normal vasoconstrictor response to orthostatic stress. When patients stand, intact baroreceptors detect the gravitational blood shift, but the effector arm of the reflex (sympathetic vasoconstriction mediated by small fibers) functions inadequately, resulting in venous pooling and compensatory tachycardia characteristic of neuropathic POTS.

In mast cell disorder patients, Novak et al. documented SFN in 80% of cases, with universal dysautonomia when combining sympathetic, parasympathetic, and sudomotor testing [173]. Given the overlap between mast cell activation and ME/CFS (Section 5.6.11), SFN may represent a common pathway linking immune dysregulation to cardiovascular symptoms in both conditions.

Hyperadrenergic POTS

- Excessive sympathetic activation [325]
- Standing norepinephrine >600 pg/mL
- Associated with tremor, anxiety, hypertension during episodes
- May involve norepinephrine transporter deficiency

Hypovolemic POTS

- Low blood volume as primary driver [325, 333]
- Compensatory tachycardia to maintain cardiac output
- May respond to volume expansion
- Overlaps with ME/CFS blood volume deficits

Splanchnic Blood Flow Dysregulation The splanchnic vascular bed—comprising the circulation to the gastrointestinal tract, liver, spleen, and pancreas—contains approximately 30% of total blood volume. This substantial reservoir plays a critical role in cardiovascular homeostasis, but also represents a vulnerability in dysautonomic conditions. In POTS and ME/CFS, splanchnic blood flow dysregulation has implications beyond cardiovascular symptoms, potentially contributing to gastrointestinal dysfunction and systemic inflammation through gut barrier compromise.

Splanchnic Pooling in POTS. POTS patients demonstrate excessive abdominal blood pooling, which can occur even in supine and resting positions [336]. Different POTS subtypes exhibit distinct splanchnic mechanisms:

- **Neuropathic POTS:** Sympathetic denervation extends beyond lower extremities to include splanchnic vessels, leading to inadequate vasoconstriction and pooling in the abdominal vasculature. This pooling reduces effective circulating volume and cardiac preload.
- **Normal-flow POTS:** Characterized by splanchnic vasodilation even when supine, these patients show inappropriate relaxation of mesenteric vessels, sequestering blood in the abdomen and reducing venous return to the heart.
- **Post-prandial exacerbation:** Splanchnic pooling increases substantially after meals in all POTS patients. Digestion requires 10–30% increase in splanchnic blood flow; in patients with impaired autonomic compensation, this post-prandial demand cannot be met without compromising perfusion to other organs. Many POTS-ME/CFS patients report worsening symptoms during and after meals, potentially reflecting this competitive redistribution.

Treatment Evidence: Abdominal compression garments (40 mmHg) combined with leg compression (20–30 mmHg) have been shown to reduce splanchnic-mesenteric venous pooling and improve orthostatic tolerance in POTS patients [336]. This mechanical intervention suggests that splanchnic pooling is not merely an epiphenomenon but actively contributes to hemodynamic instability.

Paradoxical Hypoperfusion During Activity. While pooling implies increased blood volume in the splanchnic bed, this does not translate to adequate *perfusion* at the microvascular level. Blood accumulates in distended capacitance vessels rather than flowing through nutrient capillaries. During activity—even minimal exertion in severe patients—sympathetic

activation attempts to redistribute blood to exercising muscles and the brain, but dysautonomic patients cannot effectively mobilize pooled splanchnic blood. The result is paradoxical: splanchnic vessels remain dilated (pooling persists) while capillary perfusion pressure drops (hypoperfusion), creating ischemic stress in the intestinal mucosa.

Van Campen et al. [140] demonstrated that severe ME/CFS patients show 27% reduction in cerebral blood flow during minimal orthostatic stress (20-degree head-up tilt), compared to only 7% reduction in healthy controls. Given that splanchnic vessels are more compliant and receive lower circulatory priority than cerebral vessels during sympathetic activation, it is mechanistically plausible that severe ME/CFS patients experience *equal or greater* splanchnic blood flow reduction during routine postural changes.

Inference for Severe Patients (Certainty: 0.65): If a 20-degree tilt—a nearly supine position—causes 27% cerebral blood flow reduction, then sitting upright, standing attempts, or even cognitive exertion requiring blood redistribution likely trigger substantial splanchnic hypoperfusion in severe ME/CFS patients. This hypoperfusion may occur during activities that would not be considered “exercise” in healthy individuals: reading, conversation, sitting upright for meals, or toileting.

Connection to Gut Barrier Function. The gastrointestinal mucosa is highly metabolically active and exquisitely sensitive to hypoperfusion. In healthy individuals, 60 minutes of vigorous exercise (70% VO₂max) causes portal blood flow to decrease by 80%, with splanchnic hypoperfusion detectable within 10 minutes [337]. This exercise-induced hypoperfusion results in measurable intestinal injury: plasma I-FABP (intestinal fatty acid-binding protein, a marker of enterocyte damage) increases within 1 hour, and intestinal permeability transiently rises, allowing bacterial lipopolysaccharide (LPS) to translocate into systemic circulation [337].

In ME/CFS patients—particularly those with POTS overlap—chronic baseline splanchnic pooling combined with exaggerated hypoperfusion during minimal activities may create sustained or repeated ischemic stress to the gut mucosa. When combined with wheat consumption (which upregulates zonulin and primes tight junctions for permeability), even brief hypoperfusion episodes could trigger bacterial translocation and LPS-mediated inflammatory responses.

Clinical Translation: The high comorbidity between POTS and ME/CFS (25–50% of ME/CFS patients meet POTS criteria [98]) suggests shared splanchnic dysregulation pathways. For patients with both conditions, abdominal compression may serve a dual purpose: not only improving orthostatic tolerance by reducing pooling, but also protecting gut barrier function by maintaining adequate splanchnic perfusion during activities. This hypothesis remains untested but is mechanistically grounded.

Speculation 15 (Postural Orthostatic Gut Syndrome). We propose the concept of **Postural Orthostatic Gut Syndrome (POGS)** for a subset of severe ME/CFS patients with POTS comorbidity, in whom upright posture alone causes sufficient splanchnic hypoperfusion to trigger wheat-primed gut barrier failure. The mechanistic sequence would be:

1. Upright posture → gravitational blood pooling in abdomen and legs
2. Baroreceptor-mediated sympathetic activation → attempt to restore cardiac output

3. Splanchnic vasoconstriction (reflex response) → paradoxical hypoperfusion despite local pooling
4. Wheat-primed tight junctions (zonulin-upregulated) → acute permeability increase under ischemic stress
5. LPS translocation → post-orthostatic symptoms (brain fog, nausea, fatigue, delayed systemic inflammation)

Certainty: 0.65 (inferred from documented CBF reduction via van Campen cerebral blood flow studies [172, 4] combined with established splanchnic mechanisms from ter Steege 2012 [336]; the inference chain is: documented orthostatic CBF reduction → inferred splanchnic hypoperfusion → hypothesized gut barrier failure in wheat-primed mucosa).

Testable Predictions:

- Plasma LPS and I-FABP will increase 2–4 hours after 20-minute seated position (vs. supine baseline) in severe ME/CFS+POTS patients but not in ME/CFS-only or POTS-only patients without wheat consumption
- Abdominal compression will reduce post-postural LPS spikes in severe ME/CFS+POTS patients
- Wheat elimination will reduce symptom exacerbation during upright positioning (in POGS patients)

This model emphasizes that gut barrier function is not isolated from autonomic and cardiovascular dysfunction—it is intimately coupled. Therapeutic interventions targeting one system (compression for POTS) may simultaneously benefit gastrointestinal function through improved splanchnic perfusion.

Future Research Directions: These predictions provide a roadmap for future validation studies to establish whether POGS represents a distinct clinical entity or a mechanistic subtype within the ME/CFS spectrum. Prospective studies combining tilt-table testing with serial gut permeability biomarkers (LPS, I-FABP, zonulin) in severity-stratified ME/CFS cohorts with and without POTS comorbidity would be particularly informative.

Speculation 16 (Functional vs. Structural OI Distinction). The higher prevalence but better reversibility of orthostatic intolerance in pediatric ME/CFS (70–90% prevalence with high response to treatment [338, 339]) compared to adult disease suggests qualitatively different mechanisms. We speculate that pediatric OI may primarily represent functional miscalibration of an autonomic system still undergoing developmental tuning, while adult OI may involve structural damage (small fiber neuropathy, receptor autoantibody-mediated damage) accumulated over illness duration. This distinction would explain why OI treatment in children often produces multi-domain improvement (fatigue, cognition, wellbeing) while adult responses may be more limited. If correct, this supports aggressive early OI treatment in both populations to prevent functional miscalibration from progressing to structural damage.

Certainty: 0.35 (prevalence and prognosis data documented [338, 339]; functional vs. structural mechanism distinction is speculative and untested)

10.3.4 Hypertension in ME/CFS

While hypotension is more commonly discussed, hypertension also occurs:

- **Supine hypertension:** Some patients have elevated BP when lying down
- **Labile hypertension:** Wide BP fluctuations
- **Stress-related:** BP spikes during symptom exacerbations
- **Medication-related:** Sympathomimetics for orthostatic symptoms may raise BP

10.4 Heart Rate Abnormalities

10.4.1 Resting Tachycardia

Many ME/CFS patients exhibit elevated resting heart rate:

- **Mechanism:** Compensatory response to low stroke volume
- **Sympathetic activation:** Chronic low-grade sympathetic overdrive
- **Deconditioning:** Loss of cardiovascular fitness
- **Clinical significance:** Correlates with symptom severity

10.4.2 Heart Rate Variability

Heart rate variability (HRV) reflects autonomic modulation of the sinoatrial node (see Chapter 8 for detailed discussion). Multiple studies document autonomic dysfunction in ME/CFS [97], and the NIH deep phenotyping study confirmed significantly reduced HRV in ME/CFS patients [5]:

- **Reduced overall HRV:** Lower SDNN and total power
- **Diminished parasympathetic markers:** Reduced high-frequency power and RMSSD
- **Altered sympathovagal balance:** Changed LF/HF ratio
- **Prognostic implications:** Low HRV predicts poor health outcomes generally

10.4.3 Heart Rate Recovery

Heart rate recovery (HRR) after exercise reflects parasympathetic reactivation:

- **Definition:** HR decrease from peak to 1 or 2 minutes post-exercise
- **ME/CFS findings:** Delayed HRR indicating impaired vagal reactivation
- **Clinical significance:** Abnormal HRR predicts mortality in other populations
- **Mechanism:** Consistent with parasympathetic dysfunction

10.5 Coagulation and Rheological Abnormalities

10.5.1 Hypercoagulability

Some ME/CFS patients show evidence of increased coagulation activation [166]:

- **Platelet hyperactivation:** Approximately 80% of ME/CFS patients demonstrate platelet hyperactivation (mean spreading score 2.72 vs. 1.00 in controls), with pseudopodia formation [166]
- **Thromboelastography:** Hypercoagulable state detected in ~50% of ME/CFS participants across multiple thromboelastography parameters [166]
- **Elevated fibrinogen and D-dimer:** Clotting cascade activation markers have been reported in some cohorts, though systematic data remain limited

Serum proteomics data from Hoel et al. (2026) [191] provided further evidence for a procoagulant state in ME/CFS. Among secreted proteins, those involved in coagulation and complement pathways showed a consistent pattern of elevated levels. Specifically, coagulation-related proteins showed increased secretion across multiple tissue compartments, while the broader secretome analysis revealed heightened activity in inflammation-associated pathways (complement factors, chemokines, cytokines) alongside reduced enzyme levels—a pattern consistent with chronic low-grade immune activation driving a prothrombotic environment.

10.5.2 Fibrin Deposition

Excessive fibrin deposition may impair microcirculation [166]:

- **Fibrinoid microclots:** Amyloid-resistant fibrin microclots are present in ME/CFS plasma at >10-fold greater burden than in healthy controls [166]
- **Microcapillary obstruction:** These rigid microclots may block microcapillaries, reducing perfusion and contributing to post-exertional symptoms [166]
- **Fibrinolysis resistance:** The amyloid conformation of these microclots renders them resistant to normal fibrinolytic clearance [166]
- **Treatment implications:** Anticoagulation and fibrinolytic agents have been investigated in small, preliminary case series; evidence remains insufficient for clinical recommendations

10.5.3 Red Blood Cell Deformability

Red blood cells must deform to traverse capillaries [180]:

- **Reduced deformability:** Red blood cell deformability is significantly diminished in ME/CFS patients [180]
- **Mechanisms:** Membrane oxidative damage, altered lipid composition

- **Consequences:** Impaired capillary perfusion and oxygen delivery, potentially contributing to exercise intolerance [180]
- **Measurement:** Ektacytometry, micropipette aspiration

10.6 Cardiovascular Dysfunction in Post-COVID ME/CFS

The COVID-19 pandemic has provided an unfortunate natural experiment in post-infectious illness, with Long COVID (post-acute sequelae of SARS-CoV-2, PASC) showing remarkable overlap with ME/CFS cardiovascular manifestations. This convergence strengthens the evidence for shared pathophysiological mechanisms and may accelerate therapeutic development.

10.6.1 Parallel Cardiovascular Findings

Long COVID patients demonstrate cardiovascular abnormalities closely mirroring those documented in ME/CFS, including POTS (affecting 30–50% of Long COVID patients with persistent symptoms), reduced exercise capacity on CPET with similar patterns of reduced VO_2 peak and early anaerobic threshold, autonomic dysfunction with altered HRV and baroreflex sensitivity, and endothelial dysfunction with impaired flow-mediated dilation [105].

Small fiber neuropathy has been documented in both conditions using identical methodologies. Azcue et al. found that ME/CFS and post-COVID patients showed comparable patterns of heat response latencies indicating C-fiber denervation [335]. The non-length-dependent pattern observed in both conditions suggests autoimmune or inflammatory etiology rather than metabolic neuropathy.

10.6.2 GPCR Autoantibodies in Post-COVID

The GPCR autoantibody hypothesis has received substantial support from post-COVID research. Elevated beta-adrenergic and muscarinic receptor autoantibodies have been documented in Long COVID patients with cardiovascular symptoms, and immunoadsorption targeting these autoantibodies has shown efficacy in post-COVID ME/CFS [118]. The Scheibbogen group demonstrated that repeated immunoadsorption in patients with post-COVID ME/CFS and elevated beta-2 adrenergic receptor autoantibodies produced significant improvements in fatigue and autonomic symptoms.

Hackel et al. demonstrated that GPCR autoantibodies reprogram monocyte function in post-COVID ME/CFS, altering cytokine production patterns and potentially explaining systemic inflammatory features [209]. This finding links autoantibodies to immune dysfunction beyond direct receptor effects, suggesting multiple downstream consequences of the autoimmune process.

10.6.3 Implications for Understanding ME/CFS

The convergence of Long COVID and ME/CFS cardiovascular findings supports the hypothesis that both conditions share common post-infectious pathophysiology. The larger Long COVID research effort, driven by the pandemic's scale, is generating mechanistic insights likely applicable to ME/CFS. Therapeutic interventions developed for Long COVID—including immunoadsorption, GPCR-targeting aptamers, and mast cell stabilizers—may prove equally effective in ME/CFS patients with similar pathophysiology.

10.7 Sex Differences in Cardiovascular Manifestations

ME/CFS demonstrates a 3:1 to 4:1 female predominance [318], and emerging evidence suggests that cardiovascular manifestations may differ between sexes beyond simple prevalence differences.

POTS is more common in females, with cohort studies consistently showing 4:1 to 5:1 female-to-male ratios [98]. This sex difference exceeds the overall ME/CFS female predominance, suggesting additional sex-specific factors in POTS pathophysiology. Potential contributors include differences in blood volume regulation (females have lower baseline blood volume per kilogram), hormonal effects on vascular tone and autonomic function, and sex differences in autoimmune propensity affecting GPCR autoantibody production.

The NIH deep phenotyping study revealed distinct immune abnormalities in male versus female ME/CFS patients [5], and these differences likely extend to cardiovascular manifestations. Sex hormone effects on endothelial function, baroreflex sensitivity, and autonomic balance may modulate how the underlying disease process manifests cardiovascularly.

Blood volume deficits may be proportionally greater in females. van Campen et al. found that red blood cell mass was reduced in 93.8% of female ME/CFS patients compared to 50% of males, while plasma volume was subnormal in the majority of both sexes [340]. This sex difference in red cell mass reduction may reflect hormonal influences on erythropoiesis or differential inflammatory effects.

Clinical implications include the need for sex-stratified analysis in cardiovascular research and potentially different therapeutic thresholds. The higher prevalence of POTS in females may warrant lower diagnostic thresholds for autonomic testing referral in female patients with orthostatic symptoms.

10.8 Mast Cell Activation and Cardiovascular Dysfunction

Mast cell activation syndrome (MCAS) frequently co-occurs with POTS and ME/CFS, forming part of the “Septad” of overlapping conditions. The cardiovascular effects of mast cell degranulation provide a mechanistic link between immune activation and hemodynamic instability.

10.8.1 Cardiovascular Mediators of Mast Cell Activation

Mast cells release multiple vasoactive mediators upon degranulation:

- **Histamine:** Causes vasodilation through H1 and H2 receptor activation on vascular smooth muscle, increasing vascular permeability and contributing to hypotension
- **Prostaglandin D₂:** Potent vasodilator that may contribute to flushing and hypotensive episodes
- **Tryptase:** Serine protease that can activate protease-activated receptors on endothelial cells, potentially contributing to endothelial dysfunction
- **Platelet-activating factor (PAF):** Causes vasodilation, increases vascular permeability, and promotes platelet aggregation
- **Heparin:** Released during degranulation, may contribute to bleeding tendency and affect coagulation

During mast cell degranulation episodes, the sudden release of vasodilatory mediators can produce acute hypotensive episodes, flushing, and tachycardia. When mast cell activation is chronic and low-grade, the cumulative effect may include sustained endothelial dysfunction and impaired vascular reactivity.

10.8.2 The MCAS-POTS Connection

The relationship between MCAS and POTS is bidirectional. Mast cell mediators, particularly histamine, cause peripheral vasodilation that exacerbates venous pooling during orthostatic stress. Conversely, orthostatic stress may trigger mast cell degranulation in susceptible individuals, creating a feed-forward loop. This bidirectional interaction exemplifies the reinforcing pathophysiological cycles discussed in Section 13.4 of Chapter 13.

Novak et al. documented that mast cell disorder patients universally showed dysautonomia when combining sympathetic, parasympathetic, and sudomotor testing [173]. The same patients showed 20–24% reduction in orthostatic cerebral blood flow, directly linking mast cell activation to cerebral hypoperfusion during standing.

The high prevalence of small fiber neuropathy (80%) in mast cell disorder patients [173] suggests that mast cell mediators may directly damage autonomic nerve fibers or that both findings reflect a common underlying autoimmune process. Tryptase and other mast cell proteases can cleave components of the extracellular matrix and potentially damage nerve terminals.

10.8.3 Therapeutic Implications

The mast cell-cardiovascular connection has therapeutic implications. H1 antihistamines (cetirizine, loratadine, rupatadine) and H2 blockers (famotidine) may improve orthostatic symptoms in patients with concurrent MCAS. Mast cell stabilizers (cromolyn sodium, ketotifen) may provide broader suppression of mediator release. In patients with ME/CFS and prominent flushing, episodic tachycardia, or symptom fluctuation temporally associated with

meals or environmental triggers, evaluation for MCAS should be considered, and empiric antihistamine therapy may be warranted.

10.9 Summary: Integrated Cardiovascular Model

Cardiovascular dysfunction in ME/CFS involves multiple interacting abnormalities [5]:

1. **Reduced blood volume:** Hypovolemia compromises cardiac preload and limits cardiac output reserve
2. **Autonomic dysfunction:** Parasympathetic withdrawal reduces HRV and impairs baroreflex function; chronotropic incompetence limits exercise heart rate response
3. **Endothelial dysfunction:** Impaired vasodilation reduces tissue perfusion
4. **Cardiac limitation:** Preload failure and chronotropic incompetence reduce maximal cardiac output
5. **Microcirculatory impairment:** Abnormal capillary perfusion and oxygen extraction limit peripheral oxygen delivery
6. **Exercise intolerance:** The cumulative effect is reduced VO_2 peak and early anaerobic threshold, objectively confirmed by the NIH study
7. **Post-exertional deterioration:** Unique to ME/CFS, the failure to recover exercise capacity on day 2 CPET reflects pathological response to exertion
8. **Orthostatic intolerance:** Blood pressure dysregulation (POTS, NMH, OH) produces symptoms with upright posture

This cardiovascular dysfunction explains much of the disability in ME/CFS: patients cannot sustain physical activity because their cardiovascular system cannot deliver adequate oxygen to meet metabolic demands. The objective documentation of reduced VO_2 peak and chronotropic incompetence in the NIH deep phenotyping study provides biological validation of patients' reported exercise intolerance. These cardiovascular abnormalities integrate with metabolic dysfunction (Chapter 6), autonomic dysfunction (Chapter 8), and immune dysregulation (Chapter 7) to produce the multi-system pathophysiology synthesized in Chapter 13.

Treatment approaches targeting cardiovascular dysfunction include volume expansion (fludrocortisone, increased fluid and salt intake), direct-acting autonomic agents (midodrine as alpha-agonist for vasoconstriction), and careful activity management to avoid exceeding the reduced aerobic threshold. The efficacy of pharmacological agents that bypass impaired CNS autonomic coordination (such as midodrine acting directly on peripheral alpha-receptors) provides indirect support for the selective energy dysfunction hypothesis discussed in Section 6.3.

The recognition that cardiovascular abnormalities are objective and measurable helps counter misconceptions that ME/CFS exercise intolerance reflects psychological factors or simple deconditioning. Charlton et al. (2026) formalized this argument in the *British Journal of Sports Medicine*, concluding that post-exertional malaise cannot be explained by cardiac deconditioning and calling for a fundamental rethinking of exercise pathophysiology in post-viral illness [341].

~ Hypothesis 1: Motor-Autonomic Coordination Overload Hypothesis

Physical activity requires simultaneous CNS coordination of motor output (muscle recruitment, movement planning, proprioceptive feedback) and autonomic regulation (heart rate adjustment, blood pressure maintenance, thermoregulation, respiratory drive). In healthy individuals, these coordination tasks operate efficiently within the brain's energy budget. We hypothesize that in ME/CFS, where CNS energy is the primary bottleneck (Section 14.24), motor and autonomic coordination compete for insufficient resources, causing both systems to fail under demand.

The Dual-Coordination Problem. During exercise, the CNS must:

- **Motor coordination:** Generate movement commands, integrate proprioceptive feedback, maintain balance, adjust force output—all requiring continuous cortical, cerebellar, and spinal processing.
- **Autonomic coordination:** Increase heart rate, redistribute blood flow, maintain blood pressure during postural changes, regulate respiration, initiate sweating—all requiring brainstem and hypothalamic processing.
- **Integration:** Coordinate motor and autonomic outputs so that cardiovascular supply matches muscular demand in real time.

In ME/CFS, if total CNS energy available for coordination is reduced, attempting both tasks simultaneously will exceed the available budget sooner than either task alone. This explains the central governor theory observation [342, 343]: the brain limits motor output to protect itself from energy depletion.

ME/CFS-Specific Predictions. This hypothesis explains several puzzling CPET findings:

Reduced VO₂peak beyond deconditioning. The NIH deep phenotyping study documented VO₂peak reductions exceeding what deconditioning alone would predict [5]. If the brain limits motor output to preserve autonomic coordination capacity, peak exercise performance reflects the CNS energy budget, not peripheral muscle capacity.

Chronotropic incompetence. The failure to achieve age-predicted maximal heart rate [5] may reflect CNS prioritization: under energy constraint, the brain may reduce autonomic drive to the heart in order to preserve motor coordination, or vice versa. The specific pattern of failure depends on which system the CNS prioritizes in a given individual.

Day-2 CPET deterioration. The pathognomonic worsening on repeat CPET the following day reflects CNS energy depletion that has not recovered. The first test depletes CNS coordination reserves; insufficient recovery time means the second test starts from a lower baseline, producing objectively worse performance.

PEM as coordination exhaustion. Post-exertional malaise may represent the downstream consequence of depleting CNS coordination reserves. Once exhausted, the brain cannot adequately coordinate autonomic function (causing orthostatic symptoms, heart rate irregularity) or motor output (causing weakness, poor coordination), producing the multi-system symptom exacerbation characteristic of PEM.

Testable Predictions.

1. **Cognitive-physical interference:** ME/CFS patients should show greater cognitive impairment during physical activity (dual-task paradigm) than healthy controls, reflecting competition for shared CNS resources.
2. **Autonomic-motor trade-off:** During exercise, ME/CFS patients should show an inverse relationship between motor performance and autonomic function quality (e.g., better muscle output correlates with worse HRV, and vice versa).
3. **Separate-task preservation:** Motor tasks without significant autonomic demand (e.g., seated fine motor tasks) and autonomic challenges without motor demand (e.g., passive tilt testing) should each show less impairment than combined motor-autonomic challenges (e.g., exercise).
4. **Pharmacological bypass:** Agents that directly support autonomic function (mido-drine, pyridostigmine) should improve exercise tolerance by offloading CNS autonomic coordination, freeing energy for motor output.

Treatment Implications.

- **Pre-treatment with autonomic agents:** Taking autonomic-supporting medications before planned physical activity could extend exercise tolerance by reducing CNS autonomic coordination demands.
- **Activity design:** Activities that minimize simultaneous motor-autonomic demand (recumbent exercise, swimming) should be better tolerated than upright weight-bearing exercise.
- **Pacing rationale:** The coordination overload model provides a mechanistic rationale for pacing: staying below the threshold where motor and autonomic demands simultaneously exceed CNS capacity prevents the cascade of coordination failure that produces PEM.

Limitations. The hypothesis assumes CNS energy is the primary constraint, which remains debated. Peripheral factors (mitochondrial dysfunction, reduced blood volume, deconditioning) independently contribute to exercise intolerance. The dual-task prediction requires careful experimental design to distinguish CNS resource competition from general fatigue. Central governor theory itself remains controversial in exercise physiology.

Certainty: 0.55 (CPET findings well-documented; CNS coordination mechanism plausible; dual-task predictions testable but not yet tested in ME/CFS)

Speculation 17 (Small Fiber Neuropathy Increases CNS Metabolic Load). Small fiber neuropathy affects approximately 30% of ME/CFS patients [335, 331], creating a bidirectional communication burden between the peripheral nervous system and central nervous system that may amplify energy constraints.

Afferent Signal Degradation. SFN reduces the quality of autonomic afferent signals reaching the CNS—temperature sensing, baroreceptor feedback, visceral sensation. Degraded

sensory input increases CNS processing demands to extract meaningful information. Neural systems must increase firing rates quadratically to achieve linear improvements in signal-to-noise ratio [344], creating disproportionate metabolic costs when processing noisy peripheral signals. This is analogous to listening to conversation in a noisy environment: the brain expends more energy processing degraded input to achieve adequate perception.

Efferent Command Amplification. When efferent small fibers are damaged, the CNS must generate stronger, more frequent, or redundant autonomic commands to achieve target physiological responses. Fewer functional nerve fibers mean each must be driven harder, or signals must be repeated, increasing the metabolic cost of autonomic control. During orthostatic stress, the CNS may detect inadequate vasoconstriction (via baroreceptor feedback) despite issuing normal commands, triggering escalating compensatory signals that further drain central energy reserves.

Testable Predictions.

- Intraepidermal nerve fiber density (IENFD) should inversely correlate with brainstem and hypothalamic glucose uptake (FDG-PET) during autonomic challenges such as tilt testing ($r < -0.5$ expected).
- ME/CFS patients with confirmed SFN should demonstrate worse cognitive fatigue and brain fog than those without SFN, controlling for pain severity and autonomic dysfunction magnitude.
- Treatment of autoimmune SFN with IVIG should reduce CNS metabolic burden measurable by PET or MR spectroscopy, with corresponding improvements in cognitive symptoms.
- Corneal nerve fiber tortuosity (measured non-invasively via corneal confocal microscopy) should correlate with CNS lactate accumulation and cognitive impairment.
- Cognitive load should exacerbate autonomic dysfunction more severely in SFN-positive patients, reflecting competition for limited CNS energy resources.

Treatment Implications. If this hypothesis is correct, treating SFN may reduce CNS metabolic burden and improve cognitive symptoms even without direct CNS interventions. The non-length-dependent SFN pattern documented in ME/CFS [335] suggests autoimmune etiology, potentially responsive to immunomodulation. Case series (low certainty) suggest IVIG improves pain and autonomic symptoms in autoimmune SFN [285, 284], though randomized controlled trials in idiopathic SFN have shown mixed results. The distinct autoimmune pattern in ME/CFS-associated SFN may predict better immunotherapy response than idiopathic cases.

Limitations. No studies have directly measured CNS metabolic demand in relation to SFN severity in ME/CFS. SFN and cognitive dysfunction may share common causes (e.g., autoimmunity or inflammation) rather than having a causal relationship. The relative contribution

of SFN to overall CNS energy constraints is unknown and may be minor compared to other factors.

Current Evidence. Azcue et al. documented that ME/CFS patients show prolonged heat response latencies indicating C-fiber dysfunction, with 31% meeting POTS criteria [335]. A follow-up study using corneal confocal microscopy demonstrated increased small fiber tortuosity in ME/CFS compared to controls ($F = 6.80, p < 0.01$), with tortuosity serving as the primary discriminator between patients and controls (AUC= 0.720) [331]. The non-length-dependent pattern (upper and lower extremities equally affected) distinguishes ME/CFS-associated SFN from metabolic neuropathies like diabetic neuropathy, suggesting immune-mediated damage targeting specific antigens on small nerve fibers. The connection between reduced parasympathetic activation and worse cognitive performance [335] provides indirect support for peripheral-CNS interface dysfunction, though directionality remains uncertain.

Certainty: 0.40 (SFN prevalence established; CNS metabolic mechanism speculative)

11 Gastrointestinal and Microbiome Dysfunction

Gastrointestinal symptoms affect the majority of ME/CFS patients, with 50–90% reporting irritable bowel syndrome (IBS)-like symptoms. The gut microbiome—the complex ecosystem of bacteria, archaea, fungi, and viruses inhabiting the intestinal tract—has emerged as a key player in ME/CFS pathophysiology, with bidirectional connections to immune function, metabolism, and the central nervous system.

GI dysmotility is one component of the “Septad” framework of frequently co-occurring conditions in ME/CFS (Section 5.6.11). This chapter examines microbiome alterations, intestinal permeability, motility disorders, and their connections to systemic symptoms.

11.1 Gut Microbiome Alterations

11.1.1 Dysbiosis Patterns

Multiple studies have documented consistent patterns of gut microbiome alterations in ME/CFS patients [226], though no single “ME/CFS signature” has been established.

★ Achievement 1: Reduced Microbiome Diversity in ME/CFS

Giloteaux et al. [226] performed 16S rRNA sequencing on stool samples from 48 ME/CFS patients and 39 healthy controls, finding:

- Significantly **reduced bacterial diversity** in ME/CFS specimens
- Reduction in relative abundance and diversity of **Firmicutes** phylum
- Increased pro-inflammatory species, decreased anti-inflammatory species
- Machine learning classification achieved 82.93% accuracy distinguishing ME/CFS from controls

This foundational study established that dysbiosis is a reproducible feature of ME/CFS (prospective case-control, n=87, High certainty).

△ Warning 1: Replication Status: Partially Replicated

The Giloteaux et al. (2016) finding of reduced bacterial diversity has been directionally confirmed in subsequent studies, but IBS co-morbidity—present in 50–90% of ME/CFS patients—was not controlled in the original study. Nagy-Szakal et al. [345] demonstrated that IBS status is the strongest driver of microbiome composition differences (see Warning 11.1.1). Studies not controlling for IBS may overestimate or misattribute microbiome changes to ME/CFS itself. The 82.93% classification accuracy has not been independently

validated.

Specific Bacterial Taxa Alterations. A 2024 systematic review [346] of 11 studies (553 ME/CFS patients, 480 controls) identified consistent patterns:

Decreased (health-promoting bacteria):

- *Faecalibacterium prausnitzii*—major butyrate producer, inversely correlated with fatigue severity
- *Eubacterium rectale*—butyrate producer
- *Roseburia* species—short-chain fatty acid producers
- Lachnospiraceae family overall
- Firmicutes phylum (contains most butyrate producers)

Increased (pro-inflammatory bacteria):

- *Enterocloster bolteae* (formerly *Clostridium bolteae*)—associated with fatigue in multiple sclerosis and autoimmune diseases
- *Ruminococcus gnavus*—associated with inflammatory bowel disease
- *Bacteroides* genus
- Bacteroidetes phylum overall

△ Warning 2: IBS Co-Morbidity as Confounding Factor

Nagy-Szakal et al. [345] demonstrated that IBS co-morbidity is the strongest driver of bacterial composition differences in ME/CFS. When analyzing 50 ME/CFS patients with and without IBS:

- IBS status explained more variance than ME/CFS diagnosis alone
- Integrating metagenomic and metabolomic data improved ME/CFS classification (AUC=0.836)
- Studies not controlling for IBS may overestimate or misattribute microbiome changes

Given 50–90% IBS prevalence in ME/CFS, careful phenotyping is essential for research interpretation.

Recent studies reinforce the microbiome–ME/CFS link. A 2025 *Scientific Reports* study confirmed significant compositional differences (reduced *F. prausnitzii* and *Bifidobacterium*, enriched *Clostridium*) in CFS versus controls [347] (Appendix I, Section I.12.2). Most notably, an AI platform integrating stool, blood, and routine lab data achieved 90% diagnostic accuracy in 249 individuals, the highest reported for any ME/CFS biomarker panel [348] (Section I.12.1). Both studies await independent blinded multi-site replication.

11.1.2 Functional Capacity Changes

Beyond taxonomic alterations, ME/CFS patients show impaired microbiome *function*—the metabolic activities bacteria perform.

★ Achievement 2: Deficient Butyrate-Producing Capacity

Guo et al. [349] performed multi-omic analysis (metagenomics, metabolomics, qPCR) on 106 ME/CFS cases and 91 controls, demonstrating:

- Reduced capacity for **butyrate synthesis** confirmed across all methodologies
- *F. prausnitzii* deficiency correlated with fatigue severity
- Bacterial network disturbances affecting butyrate-producing community
- Fecal short-chain fatty acid levels reduced

Butyrate is the primary energy source for colonocytes and has anti-inflammatory, barrier-protective, and neuromodulatory functions. Its deficiency may contribute to intestinal permeability and systemic inflammation (multi-center study, n=197, High certainty).

△ Warning 3: Replication Status: Partially Replicated

The butyrate deficiency finding (Guo et al. 2023, n=197) is the largest study to date and confirmed across multiple methodologies within the study. However, dietary intake—the primary determinant of short-chain fatty acid production—was not controlled. Fiber intake differences between cases and controls could confound butyrate measurements. Earlier smaller studies reported consistent direction of effect but from overlapping research networks.

11.1.3 Multi-Omic Integration: Beyond Taxonomic Profiling

Traditional 16S rRNA sequencing identifies *which bacteria* are present but cannot determine *what they are doing*. This distinction matters: two patients with identical bacterial communities may produce entirely different metabolite profiles depending on gene expression, substrate availability, and inter-species interactions. Early ME/CFS microbiome studies relied exclusively on taxonomic profiling and achieved moderate classification accuracy (e.g., 82.93% in Giloteaux et al. [226]). Nagy-Szakal et al. [345] combined fecal metagenomic and plasma metabolomic profiling, achieving in-sample classification AUC=0.893 with an 8-species bacterial model (cross-validated AUC=0.745). Crucially, their analysis revealed that IBS comorbidity—not ME/CFS diagnosis—was the dominant driver of bacterial compositional differences, underscoring why metabolomic data are needed to disentangle ME/CFS-specific signatures from IBS confounding.

The most comprehensive multi-omic analysis to date, by Xiong et al. [350], combined shotgun metagenomics with plasma metabolomics in 149 ME/CFS patients (74 short-term, 75 long-term) and 79 healthy controls. Multi-omic integration achieved AUC=0.90 for ME/CFS classification, substantially outperforming species relative abundance alone (AUC=0.73) or metagenomic gene profiles alone (AUC=0.73). The plasma metabolome alone achieved AUC=0.82, already superior to taxonomic profiling. Key discriminating metabolites included

sphingomyelins, diacylglycerols, ceramides, and reduced butyrate-related compounds—a metabolic fingerprint invisible to 16S sequencing.

Critically, Xiong et al. also revealed a temporal dissociation between dysbiosis and metabolic dysfunction. Short-term ME/CFS patients (<4 years) showed pronounced microbial dysbiosis with relatively preserved metabolic profiles, while long-term patients (>10 years) had largely resolved microbial dysbiosis but *worsened* metabolic aberrations and more severe clinical symptoms. This finding suggests that early dysbiosis may initiate metabolic disruptions that become self-sustaining even after partial microbiome normalization—and that metabolomic profiling captures disease progression features that taxonomic surveys miss entirely.

△ **Warning 4: Replication Status: Not Yet Replicated**

The Xiong et al. (2023) multi-omic integration findings come from a single center (Jackson Laboratory/Bateman Horne Center), though with a large, well-characterized cohort. The temporal dissociation between dysbiosis and metabolic dysfunction is a novel observation that requires independent confirmation. The superior classification performance of multi-omics (AUC=0.90) has not been validated in external cohorts. Several ongoing studies aim to replicate and extend these findings; see Appendix I, Sections I.3.1 and I.7.11.

Tryptophan Metabolism Alterations. The gut microbiome significantly modulates tryptophan availability, with gut enterochromaffin cells producing >90% of the body's serotonin. ME/CFS patients show disrupted tryptophan pathways [231, 188]:

- Reduced circulating serotonin and kynurenine affecting neurotransmission [351]—notably, these changes appeared independent of cytokine levels, suggesting tryptophan dysregulation may be a primary feature rather than secondary to inflammation
- Altered kynurenine pathway metabolites—lower 3-hydroxykynurenine and 3-hydroxyanthranilic acid, with elevated kynurenine/3HK ratios [188]
- Kynurenine pathway hyperactivation may deplete NAD⁺ via PARP activation, contributing to energy deficits [232, 233]
- Disrupted indole derivative production (aryl hydrocarbon receptor ligands)

IDO2-Mediated Tryptophan Diversion. Post-infectious immune activation may further deplete tryptophan availability through upregulation of indoleamine 2,3-dioxygenase 2 (IDO2), which diverts tryptophan into the kynurenine pathway. Two lines of evidence support this mechanism in post-COVID patients: Guo et al. found persistently elevated IDO2 expression in peripheral blood mononuclear cells, associated with reduced intracellular tryptophan and impaired mitochondrial function [352]; Rus et al. independently confirmed kynurenine pathway activation with reduced serotonin synthesis and elevated neurotoxic metabolites [353]. Wirth and Scheibenbogen¹ proposed that this IDO2-driven tryptophan diversion—simultaneously depleting serotonin and generating neurotoxic kynurenine derivatives—may link post-infectious immune activation to neurocognitive symptoms [225]. Whether this

¹Currently available as a preprint; not yet peer-reviewed.

mechanism also impairs gut–brain signaling via reduced enterochromaffin serotonin (see Section 11.1.4) remains an untested extension of their framework.

11.1.4 Gut-Brain Axis

The gut-brain axis comprises bidirectional communication between the intestinal microbiome and central nervous system through four major pathways:

1. **Neural pathway:** Vagus nerve (primary) and spinal afferents
2. **Immune pathway:** Cytokine signaling, gut-associated lymphoid tissue (GALT)
3. **Hormonal pathway:** Neurotransmitters produced by or modulated by microbiota (serotonin, GABA, dopamine precursors)
4. **Metabolic pathway:** Short-chain fatty acids, tryptophan metabolites, bile acids

Vagal Nerve Signaling. The vagus nerve connects the gut microbiome directly to brainstem nuclei controlling autonomic function, inflammation, and mood. Proposed mechanisms in ME/CFS:

- Viral infections may damage vagal afferents, impairing gut-brain communication
- Dysbiosis alters vagal signaling patterns
- Reduced vagal tone (common in ME/CFS) impairs anti-inflammatory cholinergic pathway
- Bidirectional dysfunction is evident: brain inflammation degrades gut function while gut dysfunction worsens neurological symptoms. The temporal and causal relationships remain unresolved. Longitudinal studies are needed to determine which comes first or whether both result from a common upstream cause. This bidirectional gut-brain interaction is one of several such cycles in ME/CFS, as discussed in Section 13.4 of Chapter 13.

Enterochromaffin Cell–Vagal Serotonergic Pathway. A specific mechanism linking gut dysbiosis to vagal impairment involves the enterochromaffin cell–vagal signaling pathway. Three established findings motivate this hypothesis:

1. **Neuropod cell–vagal synapses:** Kaelberer et al. demonstrated that enteroendocrine “neuropod cells” form direct synaptic connections with vagal afferent neurons, using glutamate for rapid gut-to-brain signaling [228].
2. **Serotonin–vagal activation:** Enterochromaffin cells release serotonin that activates vagal afferents via 5-HT₃ receptors [13, 14], linking gut serotonin availability to vagal tone.
3. **Butyrate–serotonin link:** Butyrate enhances serotonin production in enterochromaffin cells [14], and ME/CFS patients show deficient butyrate-producing bacteria (Achievement 11.1.2).

Combining these observations leads to a hypothesis: butyrate deficiency in ME/CFS may reduce enterochromaffin serotonin release, which could in turn diminish vagal afferent signaling. However, no study has directly measured enterochromaffin serotonin output in ME/CFS patients, so this chain remains inferential. Wirth and Scheibenbogen’s broader framework of neurotransmitter dysregulation in post-infectious illness [225] provides theoretical support, but the specific gut–vagal link proposed here extends beyond their analysis.

? Open Question 1: Gut Serotonin–Vagal–Cardiovascular Link

Does gut dysbiosis-mediated reduction in enterochromaffin serotonin release impair vagal afferent signaling sufficiently to reduce efferent vagal tone to the heart and cardiovascular system? If confirmed, this gut–serotonin–vagal–cardiovascular chain would represent a concrete mechanism by which microbiome alterations directly contribute to orthostatic intolerance and autonomic dysfunction in ME/CFS (see Chapter 10).

Microbial Neurotransmitter Production. Intestinal bacteria synthesize or modulate multiple neuroactive compounds:

- **Serotonin:** >90% of body’s serotonin produced in gut; peripheral serotonin depletion has been reported in ME/CFS patients [351], while mouse models suggest central serotonergic hyperactivity may also occur [354]—raising the possibility of compartmentalized dysregulation, though cross-species validation is needed
- **GABA:** Produced by *Lactobacillus* and *Bifidobacterium* species
- **Dopamine precursors:** Generated by intestinal bacteria
- **Short-chain fatty acids:** Cross blood-brain barrier, modulate microglial function

11.1.5 Intestinal Permeability

Intestinal permeability (“leaky gut”) refers to impaired barrier function allowing bacterial products to enter systemic circulation.

★ Achievement 3: Evidence of Bacterial Translocation in ME/CFS

A 2023 study [315] measured intestinal permeability markers in ME/CFS patients compared to fibromyalgia patients and healthy controls:

- Significantly **elevated zonulin-1 (ZO-1)** in ME/CFS versus controls
- Elevated **lipopolysaccharide (LPS)** and soluble CD14 (sCD14)
- **67% of ME/CFS patients** showed increased IgA against LPS
- **40% showed increased IgM against LPS** (versus 0% in controls)
- IgA levels correlated with illness severity

This provides direct evidence of intestinal barrier dysfunction and bacterial product translocation in ME/CFS (case-control, High certainty).

△ Warning 5: Replication Status: Partially Replicated

The bacterial translocation finding (2023 study) is consistent with earlier publications by Maes and colleagues showing elevated LPS and immune activation markers. However, the research groups overlap substantially, limiting the independence of replication. Additionally, the study measured zonulin using a commercial ELISA kit; recent methodological reviews have raised concerns about zonulin assay specificity, as some kits may cross-react with complement C3 [315]. Independent replication with validated permeability assays (e.g., lactulose/mannitol ratio) would strengthen these findings.

Severity Stratification: Critical Knowledge Gap. While baseline gut barrier dysfunction has been established in ME/CFS populations [315], a critical question remains unanswered: **Do severe/bedbound patients exhibit higher baseline permeability markers than mild/moderate patients?**

The Maes et al. study [315] did not stratify participants by disease severity, leaving uncertainty about whether zonulin, LPS, and sCD14 elevations correlate with functional impairment. This gap is significant because severity-dependent gut barrier dysfunction would suggest distinct pathophysiological mechanisms and treatment priorities for different patient subgroups.

~ Hypothesis 1: Severity-Dependent Baseline Gut Permeability

Certainty: 0.50. The individual supporting observations (CBF reduction by severity, cytokine-severity correlations, reduced cortisol in severe patients, POTS comorbidity) are moderately well-established, but their *integration* into a unified severity-stratified permeability model has not been directly tested. Certainty is set at 0.50 rather than the originally proposed 0.65 to reflect this untested integration.

Severe ME/CFS patients likely exhibit *higher* baseline permeability markers than mild/moderate patients, mediated by four convergent mechanisms:

1. **Chronic splanchnic dysregulation:** Severe patients demonstrate exaggerated vascular dysregulation during minimal stress. van Campen et al. [140] documented 27% reduction in cerebral blood flow during 20-degree head-up tilt in severe ME/CFS patients, compared to 7% in healthy controls. Given that splanchnic vessels receive lower circulatory priority than cerebral vessels during sympathetic activation, severe patients likely experience chronic low-grade splanchnic hypoperfusion during routine activities (postural changes, cognitive work, meals), leading to sustained gut barrier stress rather than episodic exercise-induced spikes.
2. **Cytokine-mediated barrier disruption:** Montoya et al. [213] established that 17 cytokines correlate linearly with ME/CFS disease severity, 13 of which are proinflammatory. Several of these—particularly IL-1 β , IL-6, and IFN- γ —are known to disrupt tight junction proteins through direct effects on occludin and ZO-1 expression [213]. In severe patients with sustained cytokine elevation, this creates a potential *bidirectional cycle*: cytokines disrupt tight junctions $\rightarrow$ LPS translocation increases $\rightarrow$ TLR4 activation amplifies cytokine production $\rightarrow$ further barrier compromise. This self-sustaining loop may explain why disease severity tends to

increase over time in some patients.

3. **Impaired barrier repair capacity:** A comprehensive biomarker examination of severely ill (housebound) ME/CFS patients [314] revealed significantly lower morning salivary cortisol (median 0.20 mcg/dL vs. 0.45 mcg/dL in controls, $p=0.002$) and reduced serum albumin. Cortisol is essential for tight junction protein expression and barrier maintenance; albumin reflects protein nutritional status necessary for epithelial cell regeneration. Severe patients' reduced capacity for barrier repair may allow wheat-induced or hypoperfusion-induced permeability to persist longer than in ambulatory patients.
4. **POTS comorbidity and splanchnic pooling:** Approximately 45% of severely ill ME/CFS patients exhibit orthostatic intolerance [314], suggesting substantial overlap with postural orthostatic tachycardia syndrome (POTS). In POTS, the splanchnic vascular bed—which contains up to 30% of total blood volume—exhibits excessive abdominal blood pooling even in supine and resting positions. This chronic pooling can paradoxically reduce effective splanchnic perfusion during activity, as blood accumulates in distended vessels rather than perfusing capillary beds. For severe ME/CFS patients with POTS features, baseline gut hypoperfusion may be a chronic condition rather than an activity-triggered event.

Speculation 18 (Chronic vs. Episodic Permeability Models). We propose the **Severity-Dependent Barrier Dysfunction Model** to describe the transition from episodic exercise-triggered permeability in mild/moderate patients to chronic sustained barrier compromise in severe/bedbound ME/CFS. Based on converging lines of evidence, this model distinguishes two distinct patterns:

Mild-Moderate Pattern (Episodic):

- Baseline zonulin/LPS mildly elevated or normal between exertions
- Acute permeability spikes triggered by intentional exercise (walking, cycling)
- Splanchnic hypoperfusion occurs during discrete activity bouts
- LPS translocation causes post-exercise inflammatory surges (PEM)
- Recovery periods allow partial barrier restoration
- Wheat elimination prevents exercise-induced permeability amplification

Severe-Bedbound Pattern (Chronic Sustained):

- Baseline zonulin/LPS chronically elevated (new homeostatic set point)
- Minimal activities (cognitive work, postural changes, meals) cause micro-exacerbations
- Chronic low-grade splanchnic hypoperfusion from baseline autonomic dysfunction
- Low cortisol and nutritional deficits prevent overnight barrier repair
- Cytokine-barrier bidirectional cycle becomes self-sustaining
- Wheat elimination may reduce symptom “noise floor” rather than preventing acute PEM

Testable Predictions:

1. Plasma zonulin and LPS will stratify by severity: mild < moderate < severe
2. I-FABP (intestinal injury marker) will be chronically elevated in severe patients at rest, not just post-exercise
3. LPS fluctuations in severe patients will correlate with minimal daily activities (30-minute cognitive tasks, postural changes) measurable at 2–4 hour post-activity intervals
4. Wheat elimination in severe patients will show gradual baseline LPS reduction over 12+ weeks, rather than acute post-exercise LPS prevention seen in ambulatory patients

Treatment Implications: This model suggests wheat elimination may benefit severe/bed-bound patients *without requiring exercise testing*, as chronic baseline permeability—not exercise-induced spikes—may be the primary driver of sustained endotoxemia. However, response patterns may differ: ambulatory patients may experience acute PEM reduction, while severe patients may notice gradual quality-of-life improvements within severe functional limitations.

Clinical Vignettes: Contrasting Barrier Patterns. To illustrate the distinction between episodic and chronic barrier dysfunction patterns proposed by the Severity-Dependent Barrier Dysfunction Model, consider two representative cases:

Patient A (Ambulatory, Mild-Moderate): A 35-year-old woman with mild ME/CFS maintains part-time work from home. She experiences predictable post-exertional malaise 24–48 hours after intentional exercise attempts (e.g., 15-minute walks). Between activity bouts, she reports baseline fatigue but manageable cognitive function. After wheat elimination, she notes that exercise-induced symptom exacerbations become milder and shorter in duration, suggesting successful prevention of exercise-triggered gut barrier failure and subsequent LPS-mediated inflammation.

Patient B (Bedbound, Severe): A 42-year-old man with severe ME/CFS is bedbound 22+ hours daily. He experiences continuous baseline symptoms (severe fatigue, cognitive impairment, nausea) with micro-exacerbations triggered by minimal activities such as 10 minutes of reading, sitting upright for meals, or brief conversations. No intentional exercise is possible. After wheat elimination, he reports a gradual reduction in symptom “noise floor” over 12–16 weeks—baseline nausea decreases, cognitive fog lightens slightly—but he remains functionally bedbound, consistent with chronic baseline permeability reduction rather than prevention of discrete exercise-induced LPS spikes.

These vignettes align with the Severity-Dependent Barrier Dysfunction Model: Patient A exhibits episodic barrier stress with recovery capacity, while Patient B demonstrates chronic sustained barrier compromise from minimal daily activities combined with impaired repair mechanisms.

Mechanism of Barrier Dysfunction. Tight junction proteins (occludin, claudins, zonula occludens) normally seal the paracellular space between enterocytes. In ME/CFS:

- Zonulin (prehaptoglobin-2) is released in response to gliadin, bacteria, or other triggers
- Zonulin loosens tight junctions, increasing paracellular permeability
- Gram-negative bacterial LPS enters mesenteric lymph nodes and bloodstream

- LPS triggers TLR4-mediated immune activation and cytokine release
- Chronic low-grade endotoxemia may drive systemic inflammation

Exercise-Induced Barrier Dysfunction and Synergistic Effects. Beyond chronic dietary and dysbiotic factors, acute exertion may further compromise intestinal barrier function through splanchnic hypoperfusion. Intense physical exercise diverts blood flow from the splanchnic circulation to working muscles, causing transient intestinal ischemia lasting 20–60 minutes [337]. This ischemic insult triggers:

- Increased intestinal epithelial cell damage (elevated I-FABP biomarker) [355]
- Acute increase in paracellular permeability [337]
- LPS translocation from the intestinal lumen

In patients with pre-existing wheat-induced barrier dysfunction (via gliadin-mediated zonulin release or ATI-mediated inflammation), exercise-induced ischemia may create a synergistic amplification: the wheat-primed baseline permeability is further compromised by exercise-induced ischemia, resulting in excessive endotoxin translocation. This synergistic mechanism may explain why some ME/CFS patients report disproportionate PEM exacerbation following exercise, particularly when wheat consumption occurs in the hours or days preceding physical activity. See Section 23.3.4 for clinical evidence and therapeutic implications of this potential interaction.

Biomarkers of Intestinal Permeability.

- **Zonulin:** Tight junction modulator; elevated suggests active barrier dysfunction
- **LPS (lipopolysaccharide):** Bacterial endotoxin; elevated indicates translocation
- **sCD14:** Soluble form of LPS receptor; marker of monocyte activation
- **Intestinal fatty acid-binding protein (I-FABP):** Enterocyte damage marker
- **Anti-LPS antibodies (IgA, IgM, IgG):** Indicate immune response to translocated endotoxin

11.1.6 Intestinal Barrier Dysfunction and Secondary Metabolic Consequences

Beyond bacterial translocation, intestinal barrier dysfunction may contribute to ME/CFS through impaired nutrient absorption, particularly amino acids critical for mitochondrial function and nitric oxide synthesis.

Microbiome–Mast Cell Bidirectional Regulation. The gut microbiome and intestinal mast cells engage in bidirectional communication that is directly relevant to the MCAS–ME/CFS overlap [356, 357]. In the suppressive direction, microbial short-chain fatty acids—particularly butyrate and propionate—are potent mast cell stabilizers. Folkerts et al. [358] demonstrated that butyrate achieves up to 90% inhibition of IgE-mediated human mast cell degranulation, acting through histone deacetylase (HDAC) inhibition rather than through GPR41/GPR43

receptors. Mechanistically, butyrate epigenetically downregulates the tyrosine kinases BTK (−2.12-fold), SYK (−2.84-fold), and LAT (−4.6-fold)—critical transducers of FcεRI-mediated signaling. Propionate showed comparable inhibitory potency, while acetate had no effect. This finding directly links the butyrate-producer deficiency documented in ME/CFS (Achievement 11.1.2) to a mechanistic pathway for mast cell dysregulation: reduced microbial butyrate production removes an epigenetic brake on mast cell activation.

In the pro-inflammatory direction, dysbiotic microbiomes can actively promote mast cell activation through multiple routes. Certain gut bacteria express histidine decarboxylase (HDC) and produce histamine directly within the intestinal lumen. Barcik et al. [359] demonstrated this for *Lactobacillus reuteri*, which carries a complete chromosomal *hdc* gene cluster; other known HDC-expressing gut commensals include *Enterococcus faecalis* and certain *Escherichia coli* strains [356]. Barcik et al. showed that bacterial histamine secretion in the gut can influence immune responses systemically, including in the lung, demonstrating that gut-derived microbial histamine has effects beyond the intestinal tract. Additionally, lipopolysaccharide (LPS) from gram-negative bacteria—elevated in ME/CFS due to bacterial translocation (Achievement 11.1.5)—directly stimulates mast cell activation and upregulates tryptase, chymase, and pro-inflammatory cytokine expression [356].

This bidirectional relationship creates a potential vicious cycle in ME/CFS: dysbiosis reduces butyrate-mediated mast cell suppression while simultaneously increasing pro-inflammatory microbial signals (histamine, LPS), leading to mast cell hyperactivation. Activated mast cells then release proteases and histamine that damage the intestinal barrier [315], further promoting bacterial translocation and worsening dysbiosis. This cycle may contribute to the observation that MCAS diagnoses are far more common *after* ME/CFS onset (16.7%) than before it (2.8%) [229]—if progressive loss of butyrate-producing commensals gradually removes the epigenetic restraint on mast cell activation, MCAS could emerge as a secondary development rather than a pre-existing comorbidity. It also suggests that microbiome restoration (targeted probiotics, prebiotics, or FMT; see Section 11.4.5) could have mast cell-stabilizing effects independent of pharmacological interventions, by restoring endogenous butyrate production.

△ **Warning 6: Evidence Limitations: Microbiome–Mast Cell Axis**

The butyrate–mast cell inhibition data (Folkerts et al. 2020) derive from *in vitro* and *ex vivo* models using millimolar butyrate concentrations. While these concentrations are physiological in the colonic lumen, circulating butyrate levels are substantially lower, and whether systemic butyrate concentrations in ME/CFS patients are sufficient to modulate mast cell activity *in vivo* remains undemonstrated. The proposed vicious cycle (dysbiosis → reduced butyrate → mast cell activation → barrier damage → worsened dysbiosis) has not been tested longitudinally in ME/CFS cohorts. No study has simultaneously measured fecal butyrate levels, mast cell activation markers, and microbiome composition in ME/CFS patients.

Certainty: Low–Moderate. Individual mechanistic components are well-established; their integration into a self-reinforcing cycle in ME/CFS is plausible but unvalidated.

~ **Hypothesis 2: MCAS/HIT → Barrier Dysfunction → Mitochondrial Failure Cascade**

In patients with mast cell activation syndrome (MCAS) or histamine intolerance (HIT), a pathophysiological cascade may contribute to ME/CFS symptoms:

1. **MCAS/HIT activation:** Mast cell degranulation releases histamine, proteases (tryptase, chymase), and inflammatory mediators in the intestinal mucosa
2. **Barrier disruption:** Mast cell-derived proteases damage tight junction proteins; histamine increases paracellular permeability via H1 receptor-mediated cytoskeletal changes [315]
3. **Amino acid malabsorption:** Impaired intestinal absorption reduces bioavailability of:
 - L-citrulline and L-arginine (NO synthesis precursors)
 - Glycine, glutamate, cysteine (glutathione precursors)
 - Branched-chain amino acids (mitochondrial substrates)
4. **NO synthesis impairment:** Reduced arginine/citrulline availability → decreased nitric oxide production → endothelial dysfunction, impaired vasodilation, POTS exacerbation
5. **TCA cycle dysfunction:** Malate deficiency (often supplemented with citrulline-malate) impairs TCA cycle flux; documented in ME/CFS metabolomics [194]
6. **Secondary mitochondrial failure:** Combined NO pathway and TCA cycle impairment → reduced ATP production → ME/CFS fatigue phenotype

This cascade may explain why patients with MCAS/HIT comorbidity show particularly robust responses to amino acid supplementation—they are correcting malabsorption-induced deficiencies rather than primary metabolic defects.

△ **Warning 7: Evidence Limitations**

This cascade hypothesis integrates multiple established observations but has not been directly validated:

- Intestinal amino acid absorption has not been directly measured in ME/CFS patients
- The link between MCAS and amino acid malabsorption is mechanistically plausible but unconfirmed
- Response to amino acid supplementation could reflect direct metabolic support rather than malabsorption correction
- Yamano et al. [194] documented TCA cycle metabolite deficiencies in ME/CFS but did not assess intestinal function

Certainty: Low (hypothesis level).

Supporting Evidence. Several lines of evidence support this cascade model:

- **Metabolomic deficiencies:** Yamano et al. [194] found significantly reduced plasma

citrulline, malate, and isocitrate in ME/CFS patients compared to controls

- **Mast cells and barrier function:** Mast cell activation directly increases intestinal permeability through multiple mechanisms [315]
- **MCAS-ME/CFS overlap:** 10–50% of ME/CFS patients have MCAS features (Section 5.6.11)
- **Amino acid supplementation response:** Clinical reports suggest some ME/CFS patients show significant improvement with L-citrulline-malate, NAC, and other amino acid protocols
- **Glutathione deficiency:** Shungu et al. [193] demonstrated reduced cortical glutathione in ME/CFS with correlation to physical functioning; pilot NAC supplementation normalized glutathione levels

NO Dysfunction as a Central Mechanism. Nitric oxide (NO) is essential for vascular regulation, mitochondrial function, and immune signaling. In the MCAS-barrier-mitochondrial cascade:

- **L-Arginine pathway:** Arginine $\xrightarrow{\text{NOS}}$ NO + citrulline
- **Citrulline recycling:** Citrulline is converted back to arginine in the “citrulline-NO cycle”
- **ME/CFS abnormality:** Low citrulline/arginine availability → substrate-limited NO production
- **Downstream effects:**
 - Endothelial dysfunction and impaired vasodilation
 - Reduced blood flow during exertion (contributes to PEM)
 - Impaired mitochondrial biogenesis (NO regulates PGC-1 α)
 - Orthostatic intolerance (POTS exacerbation)

Therapeutic Implications. If this cascade is operative in a subset of ME/CFS patients:

- **MCAS/HIT treatment:** Stabilizing mast cells (H1/H2 blockade, quercetin, ketotifen) may reduce barrier damage
- **Amino acid supplementation:** L-citrulline-malate may be more effective than L-arginine (bypasses hepatic first-pass, provides TCA cycle substrate)
- **Glutathione support:** NAC supplementation addresses documented deficiency
- **Barrier restoration:** Targeting tight junction repair (butyrate, zinc carnosine, collagen) may improve absorption
- **Phenotype identification:** Patients with confirmed MCAS/HIT + low amino acid panel may be candidates for targeted intervention

Research Needs. Validating this cascade requires:

- Direct measurement of intestinal amino acid absorption in ME/CFS (stable isotope studies)

- Correlation of barrier permeability markers with amino acid levels
- Controlled trials of amino acid supplementation stratified by MCAS/HIT status
- Longitudinal assessment of barrier function, amino acids, and symptoms during MCAS treatment

11.2 Gastrointestinal Dysfunction

11.2.1 Irritable Bowel Syndrome Overlap

IBS and ME/CFS show profound overlap, complicating both diagnosis and treatment.

Prevalence.

- **50–90% of ME/CFS patients** meet criteria for IBS (median 51%)
- Cluster analysis: 59.6% of ME/CFS patients have “abdominal discomfort syndrome”
- IBS patients have **5-fold higher odds** of having CFS compared to general population
- IBS-Constipation (IBS-C) subtype shows higher ME/CFS association than IBS-Diarrhea

Clinical Significance. ME/CFS patients with comorbid IBS demonstrate:

- More severe fatigue
- Poorer appetite
- Increased abdominal pain
- Greater overall symptom burden

11.2.2 Motility Disorders

Gastroparesis

Delayed gastric emptying is common in ME/CFS and contributes significantly to symptom burden.

Observation 58 (High Prevalence of Gastroparesis in ME/CFS). A 2023 gastric emptying scintigraphy study [360] in ME/CFS patients (n=40) demonstrated:

- **72% showed delayed liquid-phase emptying**
- **38% showed delayed solid-phase emptying**
- Degree of delay correlated significantly with symptom severity
- Both liquid and solid delay increased with more severe ME/CFS
- Lower proximal stomach accommodation after meals
- Larger fasting antral area (suggesting visceral hypersensitivity)

Symptom profiles resembled functional dyspepsia, though autonomic dysfunction likely contributes (cross-sectional, n=40, Medium certainty).

Autonomic Connection. Gastroparesis in ME/CFS is likely secondary to autonomic dysfunction:

- Parasympathetic dysfunction associated with delayed gastric emptying
- Vagal nerve impairment reduces gastric motility
- Sympathetic hyperactivation may inhibit digestive function
- Dysautonomia disrupts coordinated antral contractions

Symptoms.

- Early satiety (75% of ME/CFS patients)
- Postprandial fullness and bloating
- Nausea (35%)
- Abdominal pain (45%)
- Vomiting (in severe cases)

Small Intestinal Bacterial Overgrowth (SIBO)

SIBO occurs when excessive bacteria colonize the small intestine, which is normally relatively sterile. It is increasingly recognized as a major contributor to ME/CFS gastrointestinal symptoms.

Observation 59 (SIBO Prevalence in ME/CFS). A retrospective analysis of ME/CFS patients referred for breath testing found:

- 479 patients referred; 367 completed hydrogen-methane breath tests
- **48% SIBO-positive when excluding equivocal results** (152/316)
- 41% SIBO-positive overall (including equivocal as negative)
- 45% SIBO-negative
- 14% equivocal
- Predictive factors: older age, IBS diagnosis

This suggests a substantial proportion of ME/CFS patients have bacterial overgrowth, though the retrospective design and referral bias (patients referred for breath testing likely had more GI symptoms) limit generalizability (retrospective chart review; 479 referred, 367 with usable results, 316 conclusive; Medium certainty) [361].

Pathophysiology: Migrating Motor Complex Dysfunction. The migrating motor complex (MMC) is a cyclic pattern of electromechanical activity that “sweeps” bacteria from the small intestine to the colon during fasting.

- MMC occurs every 90–120 minutes during fasting
- Controlled by Interstitial Cells of Cajal linking smooth muscle to enteric nervous system
- Studies suggest most patients with abnormal MMC develop duodenal bacterial overgrowth
- MMC dysfunction allows bacteria to accumulate in small intestine

Factors impairing MMC in ME/CFS:

- **Autonomic dysfunction:** Vagal impairment reduces MMC activity
- **Post-infectious autoimmunity:** Anti-CdtB and anti-vinculin antibodies (from prior gastroenteritis) damage Interstitial Cells of Cajal
- **Chronic stress:** Sympathetic activation suppresses MMC
- **Hypothyroidism:** Modulates enteric nervous system function
- **Medications:** Opioids, anticholinergics impair motility

SIBO Subtypes. Different bacterial populations produce different gases, leading to distinct clinical presentations:

Table 11.1: SIBO Subtypes and Clinical Presentations

Subtype	Gas Produced	Predominant Symptoms	Treatment Focus
Hydrogen-dominant	H ₂	Diarrhea	Rifaximin
Methane (IMO)	CH ₄	Constipation	Rifaximin + neomycin
Hydrogen sulfide (ISO)	H ₂ S	Diarrhea, gas, odor	Bismuth, targeted antibiotics

IMO = Intestinal Methanogen Overgrowth; ISO = Intestinal Sulfide Overproduction.

- **Hydrogen-dominant:** Most common; *E. coli*, *Klebsiella*, other hydrogen producers
- **Methane-dominant (IMO):** Archaea (*Methanobrevibacter smithii*) convert H₂ to CH₄; slows transit time, causes constipation
- **Hydrogen sulfide (ISO):** Sulfate-reducing bacteria; associated with diarrhea; NOT detected by standard H₂/CH₄ tests (requires Trio-Smart 3-gas test)

Diagnosis: Breath Testing. Breath testing remains the primary non-invasive diagnostic method for SIBO.

Protocol:

- Substrate: Glucose (50–75g) or lactulose (10g)
- Measurements: Every 15 minutes for 120 minutes
- End-expiratory breath samples analyzed for H₂, CH₄ (and H₂S if available)

Interpretation:

- **Hydrogen:** Rise ≥ 20 ppm above baseline
- **Methane:** ≥ 10 ppm at any point
- **Hydrogen sulfide:** ≥ 3 ppm at any point (Trio-Smart criteria)

Glucose vs Lactulose:

- Glucose absorbed in proximal small intestine (more specific for SIBO)
- Lactulose reaches colon (may yield false positives)
- Glucose preferred for diagnosis; lactulose may detect distal SIBO

11.2.3 Digestive Function

Beyond motility, ME/CFS patients may have impaired digestive capacity:

- **Pancreatic enzyme insufficiency:** Reduced lipase, protease, amylase secretion
- **Bile acid malabsorption:** Impaired fat digestion; contributes to diarrhea
- **Nutrient malabsorption:** Consequent to SIBO, intestinal permeability, and enzyme deficiency
- **Hypochlorhydria:** Reduced gastric acid (sometimes from PPI overuse) predisposes to SIBO

11.3 Metabolites and Short-Chain Fatty Acids

Short-chain fatty acids (SCFAs) are produced by bacterial fermentation of dietary fiber and have profound effects on host physiology.

11.3.1 Butyrate

Butyrate is the most metabolically important SCFA, with multiple functions relevant to ME/CFS:

- **Colonocyte energy:** Primary fuel source for colonic epithelium
- **Barrier function:** Strengthens tight junctions, reduces permeability
- **Anti-inflammatory:** Inhibits NF- κ B, reduces pro-inflammatory cytokines
- **Immune modulation:** Promotes regulatory T cell differentiation
- **Neuromodulation:** Crosses blood-brain barrier, affects microglial function
- **Epigenetic effects:** Histone deacetylase inhibitor

Butyrate deficiency in ME/CFS (see Achievement [11.1.2](#)) may contribute to:

- Increased intestinal permeability

- Chronic low-grade inflammation
- Impaired immune regulation
- Neuroinflammation

11.3.2 Acetate and Propionate

- **Acetate:** Most abundant SCFA; enters systemic circulation; substrate for lipogenesis; appetite regulation
- **Propionate:** Gluconeogenic substrate; cholesterol-lowering effects; modulates immune function

11.4 Treatment Approaches

Treatment of gastrointestinal dysfunction in ME/CFS requires addressing multiple interrelated issues: motility, bacterial overgrowth, permeability, and microbiome composition.

11.4.1 SIBO Treatment

Antibiotics

Rifaximin (Xifaxan):

- Non-absorbed antibiotic targeting small intestine
- **Dosing:** 550 mg three times daily (1650 mg/day) for 14 days
- Effective for hydrogen-dominant SIBO
- Systematic reviews confirm efficacy and safety
- **Limitation:** Some patients relapse after completing course

Neomycin:

- More effective against methane-producing archaea
- Often combined with rifaximin for IMO (methane-dominant)
- **Dosing:** 500 mg twice daily for 14 days (with rifaximin)

Metronidazole:

- Used for hydrogen sulfide SIBO
- Alternative when rifaximin unavailable or failed
- More systemic side effects

△ Warning 8: SIBO Recurrence

SIBO frequently recurs after antibiotic treatment, particularly if underlying motility dysfunction is not addressed. Addressing root causes (MMC dysfunction, autonomic impairment) and using prokinetics post-treatment may reduce recurrence.

Herbal Antimicrobials

Observation 60 (Herbal Therapy Comparable to Rifaximin in Observational Study). A 2014 retrospective study [362] compared herbal antimicrobials to rifaximin in 104 SIBO patients:

- **Herbal therapy:** 46% breath test normalization
- **Rifaximin:** 34% breath test normalization
- Herbal therapy at least as effective as rifaximin in this cohort
- 57% of rifaximin non-responders achieved normalization with subsequent herbal therapy

Herbal antimicrobials may represent an alternative for patients preferring non-pharmaceutical approaches or with antibiotic intolerance. However, this was a retrospective, non-randomized comparison; RCT validation is needed before concluding equivalence (retrospective cohort, n=104, Medium certainty).

Effective herbal agents:

- **Berberine:** Reduces pathogenic bacteria, improves intestinal barrier; more effective against hydrogen-producing bacteria
- **Allicin** (garlic extract): Antibacterial, antifungal; most effective against methane-producing microbes
- **Oregano oil:** Active constituents carvacrol (55–85%) and thymol; disrupts bacterial cell membranes; preserves beneficial *Lactobacillus* and *Bifidobacterium*
- **Neem:** Broad-spectrum antimicrobial

Protocol:

- Typical duration: 4–6 weeks
- Often two agents combined (e.g., berberine + oregano oil)
- Reassess with breath testing after treatment

Contraindications and Interactions:

- **Berberine:** Inhibits CYP3A4, CYP2D6, and CYP2C9 enzymes; may increase levels of many medications including anticoagulants, immunosuppressants, and statins; contraindicated in pregnancy
- **Oregano oil:** May lower blood pressure; caution with antihypertensives; avoid in pregnancy

- **Garlic/allicin:** Antiplatelet effects; avoid with anticoagulants (warfarin, aspirin); discontinue 7–10 days before surgery
- **General:** All herbal antimicrobials should be used under medical supervision; drug-herb interactions are common

Elemental Diet

Elemental diets provide pre-digested nutrients (amino acids, simple sugars, minimal fat) that are absorbed in the proximal small intestine, effectively “starving” bacteria of fermentable substrates.

Observation 61 (Elemental Diet Efficacy). Multiple studies demonstrate high efficacy:

- Classic study [363]: 80% breath test normalization at 14 days, 85% at 21 days
- Recent 2025 study [364] with palatable formulation: **83% SIBO eradication**
- 100% normalization in hydrogen-SIBO (n=6)
- 58% normalization in IMO (n=12)
- 66% symptom improvement in those who normalized

Elemental diet is highly effective but requires motivation due to taste and restrictive nature (clinical trials, Medium-High certainty).

Protocol:

- Duration: 14 days exclusive elemental diet (may extend to 21 days if still abnormal at day 15)
- Formulas: Vivonex Plus, mBIOTA Elemental (newer palatable formulation)
- Caloric intake based on individual requirements
- Gradual reintroduction of regular foods over 2 weeks after completion

11.4.2 Prokinetics

Prokinetic agents stimulate gastrointestinal motility and may help prevent SIBO recurrence by restoring MMC function.

- **Low-dose erythromycin** (50–100 mg at bedtime): Motilin receptor agonist; stimulates MMC. Tachyphylaxis develops with continuous use; drug holidays recommended (3 weeks on, 1 week off).
- **Prucalopride** (Motegrity): Selective 5-HT₄ receptor agonist; FDA-approved for chronic constipation. May be more effective for idiopathic gastroparesis.
- **Metoclopramide** (Reglan): Dopamine D2 antagonist; only FDA-approved medication for gastroparesis. **Warning:** Risk of tardive dyskinesia with prolonged use.
- **Domperidone:** Dopamine antagonist; not available in US (requires FDA expanded access). Cardiac monitoring required (QT prolongation risk).

△ Warning 9: Prokinetic Limitations

Prokinetics have limited effectiveness as monotherapy for gastroparesis. They are most useful for:

- Preventing SIBO recurrence after successful eradication
- Mild-to-moderate gastroparesis
- Combined with dietary modifications

They do not cure underlying autonomic dysfunction and require careful monitoring for side effects.

11.4.3 Dietary Interventions**Low-FODMAP Diet**

FODMAPs (Fermentable Oligosaccharides, Disaccharides, Monosaccharides, And Polyols) are poorly absorbed carbohydrates that are fermented by gut bacteria, producing gas and drawing water into the intestine.

Evidence:

- Multiple RCTs demonstrate efficacy for IBS symptoms
- Limited direct evidence in ME/CFS, but given 50–90% IBS overlap, likely beneficial for GI symptoms
- One fibromyalgia study (n=38) showed significant reduction in gut symptoms and widespread pain after 4 weeks

Implementation:

1. **Elimination phase** (2–6 weeks): Strict avoidance of high-FODMAP foods
2. **Reintroduction phase**: Systematic testing of individual FODMAP groups
3. **Personalization phase**: Long-term diet based on individual tolerances

Requires dietitian guidance for proper implementation; not intended as permanent restriction.

Gastroparesis Diet

- Small, frequent meals (5–6 per day)
- Low-fat (fat delays gastric emptying)
- Low-fiber during flares (fiber delays emptying)
- Well-cooked, soft foods
- Avoid lying down after meals
- Liquid calories if solid food poorly tolerated

11.4.4 Probiotics and Prebiotics

Probiotic Evidence in ME/CFS

Lactobacillus casei strain Shirota:

- 48 ME/CFS patients, 8 weeks: Significant decrease in anxiety scores versus placebo
- Follow-up study (39 patients, 2 months): Significant reduction in anxiety symptoms
- No change in depression scores

Bifidobacterium infantis 35624:

- 35 ME/CFS patients, 8 weeks
- Significantly **reduced inflammatory markers**: CRP, IL-6, TNF- α
- 71% showed reduced pro-inflammatory markers
- Symptom reduction in 3 separate RCTs

★ Key Point: Strain-Specific Probiotic Selection

Probiotic effects are highly strain-specific—a well-established principle in microbiology. *B. infantis* 35624 shows strongest evidence for reducing inflammation in ME/CFS, while *L. casei* Shirota may preferentially improve anxiety. Generic “probiotic” recommendations are unlikely to be helpful; strain selection should be evidence-based.

△ Warning 10: Probiotic Cautions

- Not all strains are equally effective; many commercial products lack evidence
- Effects may be transient (requiring ongoing use)
- Individual responses vary substantially
- Quality and viability of commercial products inconsistent
- Some patients report worsening with probiotics (particularly with SIBO)

Prebiotics

Prebiotics are non-digestible fibers that selectively feed beneficial bacteria:

- Fructo-oligosaccharides (FOS)
- Galacto-oligosaccharides (GOS)
- Inulin
- Resistant starch

Caution in SIBO: Prebiotics may worsen symptoms in patients with active SIBO by feeding overgrown bacteria. Generally better tolerated after SIBO eradication.

11.4.5 Fecal Microbiota Transplantation

FMT represents the most radical microbiome intervention—complete ecosystem replacement rather than supplementation.

Speculation 19 (FMT for ME/CFS). **Certainty: 0.25.** One completed pilot RCT (n=11) showed no symptomatic benefit, though dose may have been suboptimal. Two larger placebo-controlled trials are ongoing (Comeback, n=80; RESTORE ME, n=160); results expected 2026+. FMT is classified as speculation pending these results.

Rationale for FMT in ME/CFS:

- Restores microbial diversity impossible to achieve with probiotics alone
- Transfers not just bacteria but bacteriophages, fungi, and metabolites
- Donor microbiome may provide missing metabolic functions (butyrate production, tryptophan metabolism)
- May reset gut-immune interactions

Current Evidence:

- **Finnish pilot study (2023):** Randomized, double-blind, placebo-controlled (n=11). FMT was safe but did **not** relieve symptoms or improve quality of life. However, dose may have been suboptimal (30g vs 70g typically needed).
- **Norwegian “Comeback” study:** Randomized, double-blind, placebo-controlled (n=80). Completed enrollment March 2025; results expected 2026.
- **RESTORE ME study:** Placebo-controlled (n=160); underway since 2020.

⚠ **Limitation 1: FMT: Association Established, Causality Not Proven**

FMT for ME/CFS remains **experimental**. The data do not currently support a causal link between dysbiosis and ME/CFS symptoms:

- One negative pilot study (small sample, potentially underdosed)
- Large trials ongoing but not yet reported
- Dysbiosis in ME/CFS is association, not proven causality—both may result from shared upstream mechanisms (e.g., autonomic dysfunction reducing gut motility)
- Long-term safety of FMT not fully established

Await results of adequately powered trials before considering FMT for ME/CFS.

11.5 Integration with Other ME/CFS Pathophysiology

Gastrointestinal dysfunction interconnects with other ME/CFS mechanisms (these connections are synthesized into comprehensive multi-system models in Chapter 13):

- **Immune dysfunction** (Chapter 7): Intestinal permeability drives LPS-mediated immune activation; GALT dysregulation affects systemic immunity
- **Autonomic dysfunction** (Chapter 8): Vagal impairment causes gastroparesis and MMC dysfunction; dysautonomia and GI symptoms bidirectionally reinforce each other. A specific gut–vagal mechanism may operate through enterochromaffin cell serotonin release affecting vagal afferent signaling (Section 11.1.4), with potential downstream effects on cardiovascular autonomic function (Chapter 10)
- **MCAS** (Section 5.6.11): Bidirectional microbiome–mast cell regulation operates through butyrate-mediated epigenetic suppression and dysbiosis-driven histamine/LPS activation ([Microbiome–Mast Cell Bidirectional Regulation](#), Section 11.1.6); mast cells in gut mucosa affect motility and permeability, with potential interactions with SIBO
- **Energy metabolism** (Chapter 6): Butyrate deficiency reduces colonocyte energy; malabsorption impairs nutrient availability for mitochondrial function
- **Neurological symptoms**: Gut-brain axis dysfunction contributes to cognitive impairment, mood disturbance, and autonomic symptoms

? Open Question 2: Causality in Gut-ME/CFS Relationship

Critical unresolved questions:

- Does dysbiosis *cause* ME/CFS symptoms, or is it a *consequence* of the disease?
- Can correcting microbiome alterations improve ME/CFS outcomes?
- Which comes first: autonomic dysfunction causing dysmotility, or gut dysfunction driving autonomic symptoms?
- Would microbiome-targeted therapies be disease-modifying or merely symptomatic?

Answering these questions requires interventional studies with objective outcome measures beyond symptom questionnaires.

11.5.1 Gut-Brain Metabolic Coupling

The preceding sections establish that ME/CFS patients show reduced butyrate-producing bacteria (Achievement 11.1.2), disrupted tryptophan metabolism, and impaired gut-brain axis signaling (Section 11.1.4). Separately, evidence from neuroimaging demonstrates brain hypometabolism in ME/CFS (see Chapter 8). The following speculation proposes a mechanistic link between these observations.

Speculation 20 (Gut-Brain Energy Theft Hypothesis). In ME/CFS, a dysbiotic microbiome may actively divert energy substrates away from the central nervous system, contributing to or worsening brain hypometabolism and cognitive dysfunction. This “energy theft” operates through several convergent mechanisms:

Bacterial Substrate Competition. The human gut harbors approximately 10^{13} bacteria, collectively constituting a metabolically active organ that consumes host nutrients. Under normal

conditions, commensal bacteria provide net metabolic benefits—fermenting indigestible fiber into short-chain fatty acids, synthesizing vitamins, and supporting immune homeostasis. However, when dysbiosis shifts the community toward pathobionts and away from mutualists, this metabolic balance may reverse. Pathogenic and opportunistic bacteria preferentially consume simple sugars (glucose, amino acids) that would otherwise be absorbed by the host for systemic distribution, including to the brain [365]. In ME/CFS, where total energy availability is already compromised (Chapter 6), even modest bacterial diversion of substrates could meaningfully reduce CNS energy supply.

Butyrate Deficiency and Brain Energy. Butyrate is not merely a colonocyte fuel—it crosses the blood-brain barrier and serves as a preferred energy substrate for astrocytes, outcompeting acetate in cerebral cortical tissue [366]. Astrocytes are central to the brain’s metabolic support network, providing neurons with lactate and other substrates via the astrocyte-neuron lactate shuttle. The well-documented butyrate producer deficiency in ME/CFS (Achievement 11.1.2) therefore has implications beyond gut health: reduced circulating butyrate may deprive astrocytes of a key energy substrate, impairing their capacity to support neuronal metabolism.

Tryptophan Diversion. Gut bacteria metabolize tryptophan through multiple pathways, reducing systemic availability of this essential amino acid. In ME/CFS, dysbiosis-associated kynurenine pathway hyperactivation diverts tryptophan away from serotonin synthesis (see Section 11.1.4 and [231, 188]). Beyond serotonin depletion, this diversion depletes NAD⁺ precursors via quinolinic acid accumulation, further compromising cellular energy metabolism in the brain.

Inflammation Tax. Dysbiosis-driven intestinal permeability (Achievement 11.1.5) results in LPS translocation and chronic low-grade endotoxemia. The immune response to translocated bacterial products imposes a significant metabolic cost: pro-inflammatory cytokine production, acute phase protein synthesis, and immune cell activation collectively consume glucose and amino acids that would otherwise support CNS function [367]. This “inflammation tax” compounds the direct substrate competition, creating a double burden on the already constrained energy budget.

Testable Predictions.

1. Dysbiosis severity (measured by reduced butyrate-producer abundance or increased pathobiont load) should correlate with degree of CNS hypometabolism on FDG-PET imaging.
2. Microbiome restoration interventions (FMT, targeted probiotics; see Section 11.4.5 and Speculation 19) should improve cognitive function, with effect size proportional to microbiome normalization.
3. Fecal butyrate levels should correlate with plasma butyrate and, in turn, with brain metabolic markers (MR spectroscopy NAA/Cr ratios or FDG-PET regional uptake).

4. Post-prandial cognitive worsening (commonly reported by patients as “food coma” or post-meal brain fog) should correlate with markers of bacterial fermentation and small intestinal bacterial load.
5. Periods of reduced caloric intake or fasting may temporarily improve cognitive clarity by reducing bacterial substrate availability—though this must be weighed against the risks of caloric restriction in an energy-depleted condition.

Treatment Implications. This framework suggests that cognitive symptoms in ME/CFS may be partially addressable through microbiome-targeted interventions:

- **Targeted probiotics:** Supplementation with butyrate-producing strains (*F. prausnitzii*, *E. rectale*) to restore SCFA production and potentially brain butyrate supply.
- **Prebiotics:** Dietary fiber substrates (resistant starch, inulin) that selectively feed butyrate producers.
- **SIBO treatment:** Reducing small intestinal bacterial overgrowth (Section 11.2.2) may decrease proximal glucose competition.
- **Direct butyrate supplementation:** Oral sodium butyrate or tributyrin to bypass microbial production deficits.

Limitations.

- The correlation between dysbiosis and cognitive symptoms does not establish causation—both may result from a common upstream mechanism (e.g., autonomic dysfunction reducing gut motility).
- Quantitative modeling of bacterial substrate consumption relative to host needs has not been performed; the “theft” may be quantitatively trivial compared to total host energy flux.
- Current evidence for butyrate as a brain fuel comes primarily from animal models [366]; human relevance at physiological concentrations remains to be confirmed.
- Individual variation in microbiome composition, diet, and metabolic rate makes population-level predictions difficult.

Certainty: 0.40. This is a speculative integrative hypothesis synthesizing established individual findings into a novel mechanistic framework. Direct validation through the testable predictions above is needed before clinical application.

12 Genetic and Epigenetic Factors

Genetic predisposition and epigenetic modifications provide the constitutional vulnerability upon which environmental triggers act to produce ME/CFS. While the condition is clearly not a simple Mendelian disorder, converging evidence from family studies, twin research, and molecular genetics demonstrates that heritable factors substantially influence disease susceptibility. The 2025 DecodeME genome-wide association study (n=21,620) represents a watershed moment in understanding the polygenic architecture underlying ME/CFS [43].

Understanding genetic and epigenetic contributions illuminates why only a subset of individuals develop chronic illness following viral infections, traumatic events, or other precipitating factors. The immune abnormalities detailed in Chapter 7—including NK cell dysfunction, T cell exhaustion, and cytokine dysregulation—reflect genetic variants in immune system genes and epigenetic reprogramming following chronic antigenic stimulation. Similarly, the metabolic dysfunction described in Chapter 6 arises in part from genetic variants affecting mitochondrial function and oxidative metabolism, amplified by epigenetic silencing of metabolic genes. Neurological manifestations (Chapter 8) may stem from genetic influences on ion channels such as TRPM3, neurotransmitter systems, and blood-brain barrier integrity. Cardiovascular and autonomic dysfunction (Chapter 10) shows familial clustering consistent with inherited susceptibility to orthostatic intolerance and dysautonomia.

This chapter examines the genetic architecture of ME/CFS susceptibility, epigenetic modifications that may perpetuate chronic illness, and patterns of gene expression dysregulation across multiple physiological systems. Understanding these constitutional factors is essential for developing personalized therapeutic approaches, identifying at-risk individuals, and elucidating the fundamental mechanisms that distinguish those who recover from acute illness from those who progress to chronic disease.

12.1 Genetic Predisposition

12.1.1 Heritability Evidence

Familial clustering and twin studies provide converging evidence that genetic factors substantially influence ME/CFS susceptibility while demonstrating that inheritance follows a complex polygenic pattern rather than simple Mendelian transmission. The gene-environment interaction model best explains the observed patterns: genetic variants establish constitutional vulnerability, but environmental triggers (particularly viral infections) are required for disease manifestation.

Twin Studies and Heritability Estimates

Twin studies offer the most rigorous method for partitioning genetic and environmental contributions to disease risk. Monozygotic (identical) twins share 100% of their DNA sequence, while dizygotic (fraternal) twins share approximately 50%. Higher concordance in monozygotic compared to dizygotic twins indicates genetic contribution, with the magnitude of the difference allowing estimation of heritability.

Observation 62 (Twin Study Heritability Estimates). Buchwald et al. conducted a population-based twin study using the University of Washington Twin Registry, identifying twin pairs where at least one twin met criteria for chronic fatigue. The study found significantly higher concordance in monozygotic twins (55%) compared to dizygotic twins (19%) for unexplained chronic fatigue, yielding a heritability estimate of approximately $h^2 = 0.51$ (95% CI: 0.37–0.65). When applying more stringent CFS case definitions, monozygotic concordance decreased to approximately 30–40%, but still exceeded dizygotic concordance, suggesting heritability estimates in the range $h^2 = 0.3$ –0.5 depending on phenotype definition [368].

These moderate heritability estimates indicate that genetic factors explain 30–50% of liability to ME/CFS, with environmental factors and gene-environment interactions accounting for the remainder. The incomplete concordance even in monozygotic twins (55% rather than 100%) demonstrates that genetic susceptibility alone is insufficient for disease development.

Australian twin registry studies corroborate these findings, with concordance patterns consistent with polygenic inheritance and substantial environmental contribution. The moderate heritability is similar to other complex diseases such as type 2 diabetes ($h^2 \approx 0.4$ –0.6) and autoimmune conditions, supporting classification of ME/CFS as a multifactorial disorder arising from interactions between multiple genetic variants and environmental exposures.

Familial Aggregation and Relative Risk

Family studies complement twin research by examining disease clustering across multiple generations and family structures. Multiple independent studies document elevated ME/CFS prevalence among first-degree relatives of affected individuals compared to the general population.

Walsh et al. conducted a family study examining relatives of ME/CFS probands and found that first-degree relatives had significantly increased risk, though precise relative risk estimates varied depending on diagnostic criteria and ascertainment methods [369]. The observed familial aggregation persisted after controlling for shared household exposures during childhood, arguing against purely environmental transmission through common viral exposures or psychosocial factors. Families with multiple affected members often show variable clinical presentations, suggesting that shared genetic susceptibility manifests differently depending on individual trigger exposures and additional genetic modifiers.

The pattern of familial clustering shows several notable features. First, affected relatives frequently report different precipitating events (different viral infections, surgeries, traumas), indicating that the inherited component reflects general vulnerability rather than specific

pathogen susceptibility. Second, age of onset varies widely among affected family members, suggesting that the genetic liability does not determine timing but rather establishes life-long susceptibility that may be triggered at any point. Third, affected relatives may show different dominant symptom profiles (some predominantly immunological, others metabolic or neurological), consistent with the hypothesis that shared genetic factors establish broad systemic vulnerability that interacts with individual-specific factors to determine phenotypic expression.

Gene-Environment Interaction Model

The gene-environment interaction framework provides the most parsimonious explanation for observed inheritance patterns. Genetic variants establish constitutional susceptibility, but environmental triggers are necessary and often sufficient to precipitate disease in genetically vulnerable individuals. This model explains several key observations that pure genetic or pure environmental models cannot.

The NIH RECOVER study found that 4.5% of COVID-19 survivors developed ME/CFS [370], meaning 95.5% recovered fully despite identical viral exposure. This dramatic variation in outcome following a common environmental trigger strongly implicates genetic factors in determining who progresses from acute infection to chronic illness. Similarly, the well-documented association between infectious mononucleosis and subsequent ME/CFS affects only a minority of those infected with Epstein-Barr virus, despite near-universal infection rates by adulthood in most populations. *Giardia lamblia* outbreaks provide natural experiments: following the 2004 Bergen, Norway outbreak, approximately 5% of exposed individuals developed chronic fatigue meeting ME/CFS criteria, while 95% recovered completely, again demonstrating genetic influence on chronic sequelae following identical pathogen exposure.

Children of ME/CFS parents inherit elevated risk compared to the general population, but most do not develop the condition. This pattern reflects the polygenic architecture: multiple risk variants segregate through families, with children inheriting various combinations. Some inherit many risk alleles and show high genetic liability, others inherit few and have risk approaching population baseline. Environmental trigger exposure then interacts with this inherited liability: children with high genetic loading may develop ME/CFS following relatively mild infections, while those with low genetic loading may remain unaffected even after severe viral illnesses. Intermediate genetic liability creates vulnerability to severe triggers but resilience against mild ones.

The specific genetic variants inherited may influence which environmental triggers are most pathogenic. For example, children inheriting immune gene variants affecting viral immune responses may be particularly susceptible to viral triggers but less susceptible to non-infectious stressors. Those inheriting metabolic gene variants might be vulnerable to physical or metabolic stressors. This genotype-specific susceptibility to different environmental factors could explain the heterogeneity in precipitating events observed even within affected families.

12.1.2 Genetic Variants and Candidate Gene Studies

Before the advent of genome-wide association studies, candidate gene approaches investigated single nucleotide polymorphisms (SNPs) in genes hypothesized to influence ME/CFS pathophysiology. These studies focused on immune system genes, metabolic pathways, neurotransmitter systems, and stress response mechanisms. While candidate gene studies have important methodological limitations—including small sample sizes, multiple testing issues, and publication bias—they have identified several plausible genetic associations that warrant further investigation in larger cohorts.

Human Leukocyte Antigen (HLA) Associations

The HLA complex on chromosome 6p21 encodes major histocompatibility complex (MHC) proteins that present antigens to T cells and play a central role in adaptive immunity. HLA alleles show strong associations with autoimmune diseases, and several studies have examined whether specific HLA types predispose to ME/CFS.

Multiple studies have reported associations between ME/CFS and specific HLA class II alleles, particularly HLA-DRB1 and HLA-DQ variants. Several studies have identified increased frequency of specific HLA-DQA1 alleles in ME/CFS patients compared to controls, suggesting a potential role for antigen presentation in disease pathogenesis. Carlo-Stella et al. found associations with HLA-DQ3, particularly in patients with post-infectious onset [371]. However, these associations have shown inconsistent replication across populations, likely reflecting both genuine population differences in HLA allele frequencies and the polygenic architecture of ME/CFS where HLA contributes modest effect size as one of many susceptibility loci.

The biological plausibility of HLA associations is strong. HLA molecules determine which viral and self-peptides are presented to T cells, influencing both antiviral immune responses and potential autoreactivity. Specific HLA alleles might predispose to inefficient viral clearance, prolonged antigenic stimulation, or molecular mimicry leading to autoimmune sequelae. The connection to post-infectious onset supports this mechanism: individuals with particular HLA types may mount ineffective immune responses to triggering infections, permitting viral persistence or chronic immune activation.

Immune System Gene Variants

Beyond HLA, numerous genes regulating innate and adaptive immunity have been examined as ME/CFS susceptibility loci.

Cytokine and cytokine receptor genes represent logical candidates given the well-documented cytokine dysregulation in ME/CFS (Chapter 7). Polymorphisms in TNF- α promoter region (particularly the -308 G/A variant associated with higher TNF- α production) have been investigated, with some studies reporting increased frequency of high-expression alleles in ME/CFS patients. Variants in IL-10 (an anti-inflammatory cytokine), IL-6, and IL-1 gene clusters have also been examined. Goertzel et al. reported associations with variants affecting

IL-10 expression, consistent with the hypothesis that impaired anti-inflammatory responses permit chronic inflammation [372].

Toll-like receptor (TLR) genes, which recognize pathogen-associated molecular patterns and initiate innate immune responses, have shown suggestive associations. TLR4 polymorphisms affecting responsiveness to bacterial lipopolysaccharide may influence susceptibility to post-infectious ME/CFS. Pattern recognition receptor variants could plausibly affect both initial pathogen detection and subsequent inflammatory cascades.

Complement system genes have received less attention but merit investigation given emerging evidence for complement dysregulation in ME/CFS. Genetic variants affecting complement activation thresholds or regulatory protein function might predispose to excessive inflammatory responses or impaired clearance of immune complexes.

Metabolic and Mitochondrial Gene Variants

The profound metabolic dysfunction documented in ME/CFS (Chapter 6) suggests that genetic variants affecting cellular energetics may contribute to disease susceptibility.

Mitochondrial DNA (mtDNA) variants have been examined in several studies, though results remain inconclusive. Unlike nuclear DNA, mtDNA is maternally inherited and shows high mutation rates. Some studies have reported increased mtDNA deletions or specific haplogroup associations in ME/CFS, but replication has been inconsistent. The biological rationale remains strong: mtDNA encodes critical components of the electron transport chain, and variants reducing mitochondrial efficiency could predispose to bioenergetic crisis under conditions of increased demand or oxidative stress.

Nuclear genes encoding mitochondrial proteins represent equally plausible candidates. Recent evidence identifies WASF3 pathway dysregulation in ME/CFS [64], potentially affecting cellular energy production capacity. WASF3 is involved in actin cytoskeleton regulation and mitochondrial dynamics; genetic variants affecting its expression or function might impair mitochondrial quality control mechanisms or cellular energy distribution.

Genes involved in glucose metabolism, fatty acid oxidation, and oxidative stress responses have shown suggestive associations in small studies. Polymorphisms affecting glycolytic enzyme expression, carnitine transport (relevant for fatty acid metabolism), or antioxidant systems (superoxide dismutase, catalase, glutathione pathways) could plausibly influence metabolic reserve and stress tolerance.

Ion Channel and Neurotransmitter System Genes

Neurological symptoms in ME/CFS (Chapter 8) and the documented dysfunction of transient receptor potential melastatin 3 (TRPM3) ion channels suggest genetic variants in ion channel genes as susceptibility factors.

TRPM3 dysfunction represents one of the most mechanistically compelling genetic associations. Marshall-Gradisnik and colleagues have demonstrated reduced TRPM3 function

in ME/CFS patients' natural killer cells and B cells, with impaired calcium influx following TRPM3 activation [373]. While these functional studies demonstrate acquired TRPM3 dysfunction, genetic variants in the TRPM3 gene (particularly regulatory variants affecting expression levels) could establish constitutional vulnerability. TRPM3 channels regulate calcium signaling, which is essential for immune cell function, neurotransmitter release, and cellular metabolism. Reduced baseline TRPM3 expression due to genetic variants might create a narrower functional reserve, rendering individuals more susceptible to further acquired dysfunction.

Other ion channel genes merit investigation. Voltage-gated calcium channels, potassium channels regulating neuronal excitability, and acid-sensing ion channels (ASICs) involved in pain perception and autonomic regulation all represent plausible candidates. Channelopathies—diseases caused by ion channel dysfunction—often present with episodic symptoms, fatigue, and autonomic features resembling aspects of ME/CFS.

Neurotransmitter system genes, particularly those affecting serotonin, norepinephrine, and dopamine metabolism, have been examined given the prominent cognitive and mood symptoms. The catechol-O-methyltransferase (COMT) gene, which catabolizes catecholamines, exists in high-activity (Val158) and low-activity (Met158) variants. Some studies have reported associations with the Met158 variant, which would reduce catecholamine degradation and potentially affect stress responses and cognitive function. Serotonin transporter (5-HTTLPR) polymorphisms affecting serotonin reuptake have shown inconsistent associations.

Autonomic and Cardiovascular Genes

The high prevalence of orthostatic intolerance and postural orthostatic tachycardia syndrome (POTS) in ME/CFS patients (co-occurring in approximately 60%) [374] suggests genetic overlap with autonomic dysfunction syndromes.

Adrenergic receptor genes, particularly β -adrenergic receptor variants affecting cardiac responsiveness to catecholamines, represent logical candidates. The β_1 -adrenergic receptor gene (ADRB1) shows common polymorphisms affecting receptor density and signaling efficiency. Variants that alter cardiovascular responsiveness to sympathetic activation could predispose to orthostatic intolerance, particularly when combined with other ME/CFS-related pathophysiology such as reduced blood volume or impaired baroreceptor function.

Genes affecting renin-angiotensin-aldosterone system (RAAS) function, which regulates blood volume and vascular tone, could influence susceptibility to orthostatic symptoms. ACE (angiotensin-converting enzyme) gene variants, particularly the insertion/deletion polymorphism affecting ACE levels, might interact with other cardiovascular genetic factors to determine orthostatic tolerance.

Limitations of Candidate Gene Studies

△ Warning 1: Candidate Gene Study Limitations

Most candidate gene studies in ME/CFS suffer from serious methodological limitations that prevent definitive conclusions. Common issues include small sample sizes (often $n < 100$ cases), which provide insufficient statistical power to detect modest genetic effects; inadequate correction for multiple testing, leading to false positive findings; publication bias favoring positive associations; and lack of independent replication in separate cohorts. Many reported associations have not been replicated, and effect sizes when reported are often implausibly large, suggesting winner's curse (overestimation of effect size in discovery samples).

The transition to genome-wide association studies addresses many of these limitations through systematic interrogation of common genetic variation across the entire genome, adequate sample sizes to detect realistic effect sizes, stringent correction for multiple testing, and consortia-based designs facilitating replication.

12.1.3 Genome-Wide Association Studies

Genome-wide association studies (GWAS) represent a paradigm shift from candidate gene approaches, systematically interrogating millions of common genetic variants across the entire genome to identify disease-associated loci without prior hypotheses about specific genes. GWAS have successfully identified genetic risk factors for numerous complex diseases including type 2 diabetes, inflammatory bowel disease, schizophrenia, and rheumatoid arthritis. For ME/CFS, GWAS has been hindered by the challenges of patient recruitment, diagnostic heterogeneity, and the need for large sample sizes to detect the modest effect sizes typical of complex polygenic diseases.

DecodeME: The Largest ME/CFS GWAS

★ Achievement 1: DecodeME GWAS Findings

The DecodeME study represents the largest genetic investigation of ME/CFS to date, recruiting 21,620 ME/CFS cases and comparing them to population controls through the UK Biobank and other cohorts [43]. This unprecedented sample size provides statistical power to detect genetic variants with realistic effect sizes (odds ratios of 1.1–1.3) that reach genome-wide significance ($p < 5 \times 10^{-8}$).

DecodeME employed rigorous case ascertainment through physician diagnosis and self-report with verification, accepting patients meeting CCC (Canadian Consensus Criteria), ICC (International Consensus Criteria), or IOM (Institute of Medicine) diagnostic criteria. This inclusive approach maximizes sample size while acknowledging diagnostic heterogeneity, with sensitivity analyses examining whether genetic architecture differs across diagnostic subtypes.

The study's scale enables several key analyses beyond simple case-control association: estimation of SNP heritability (the proportion of ME/CFS liability explained by common genetic variants), genetic correlation analyses comparing ME/CFS to other conditions, polygenic risk score development, and gene-based and pathway enrichment tests identifying biological systems enriched for associated variants.

△ **Warning 2: Replication Status: Not Yet Replicated (By Design)**

DecodeME is the first adequately powered ME/CFS GWAS. The genome-wide significant loci identified await replication by design—GWAS discovery cohorts require independent replication cohorts to confirm true associations versus false positives. No independent replication cohort of comparable size currently exists, though international collaborations (e.g., with US ME/CFS Clinician Coalition biobanks) may provide future replication opportunities.

Initial findings from DecodeME confirm the polygenic architecture of ME/CFS, with disease liability arising from the cumulative effects of many variants of small individual effect rather than single genes of large effect. SNP heritability estimates from DecodeME provide genome-wide validation of the twin study heritability estimates discussed in Section 12.1.1, demonstrating that common genetic variants explain a substantial proportion of familial clustering.

Specific genome-wide significant loci identified in DecodeME require careful interpretation. Many GWAS hits fall in non-coding regions affecting gene regulation rather than protein sequence, necessitating functional follow-up to identify causal variants and target genes. Some associated loci contain genes with clear biological relevance to ME/CFS pathophysiology—immune genes, metabolic genes, neurological genes—while others highlight previously unsuspected pathways requiring mechanistic investigation.

Genetic Correlations with Other Conditions

GWAS enables calculation of genetic correlations—the degree to which two conditions share common genetic risk variants. Positive genetic correlation indicates overlapping genetic architecture; negative correlation suggests protective variants for one condition increase risk for another. DecodeME genetic correlation analyses illuminate the relationship between ME/CFS and related conditions.

Several conditions show positive genetic correlation with ME/CFS, suggesting shared genetic susceptibility. Depression and anxiety disorders demonstrate genetic overlap, though this correlation does not imply causality in either direction; rather, shared genetic variants may predispose to both ME/CFS and mood disorders through common biological pathways (perhaps involving neurotransmitter systems, HPA axis regulation, or inflammation). Fibromyalgia and irritable bowel syndrome show genetic correlation with ME/CFS, consistent with clinical overlap and suggesting shared pain processing or autonomic dysfunction pathways.

Notably, Long COVID shows genetic correlation with ME/CFS, supporting clinical observations of similar post-infectious phenotypes. This correlation suggests that genetic variants

predisposing to ME/CFS following various infections also predispose to chronic symptoms following COVID-19. The magnitude of this correlation informs debate about whether Long COVID and ME/CFS represent the same condition or overlapping but distinct entities: high correlation ($r_g > 0.7$) would suggest essentially the same genetic liability, while moderate correlation ($r_g = 0.3\text{--}0.5$) indicates shared but not identical genetic architecture.

Autoimmune diseases may show variable genetic correlation with ME/CFS. If significant positive correlation emerges, this would support hypotheses of autoimmune mechanisms in ME/CFS and suggest that some genetic susceptibility to ME/CFS reflects general autoimmune liability.

Polygenic Risk Scores

Polygenic risk scores (PRS) aggregate the effects of thousands or millions of genetic variants into a single quantitative measure of inherited liability. PRS can identify individuals at high genetic risk (top decile of PRS distribution), who may benefit from preventive interventions, or individuals at low genetic risk despite environmental exposures. For ME/CFS, PRS applications include risk stratification, mechanistic subtyping, and prediction.

DecodeME enables development of ME/CFS polygenic risk scores that can be tested for clinical utility. Key questions include: Does high PRS predict which individuals develop ME/CFS following infectious mononucleosis or COVID-19? Do patients with high versus low PRS show different clinical phenotypes, treatment responses, or prognoses? Can PRS combined with environmental risk factors improve prediction compared to either alone?

The clinical utility of PRS depends on effect size distribution. If ME/CFS liability reflects thousands of variants each contributing tiny effects, PRS will show modest discriminative ability (area under curve approximately 0.6–0.65), limiting clinical utility. If a subset of variants have larger effects, PRS performance improves. Even modest predictive ability may have clinical value: if 20% of exposed individuals with high PRS develop ME/CFS versus 2% with low PRS, this tenfold risk gradient could guide post-exposure monitoring and early intervention.

Earlier GWAS Attempts and Methodological Challenges

Observation 63 (Earlier Small GWAS Studies). Prior to DecodeME, several smaller GWAS attempts were conducted with sample sizes of 200–500 cases. These studies were severely underpowered to detect realistic effect sizes for complex disease variants and produced no genome-wide significant findings that replicated. This failure reflects general principles of GWAS: detecting odds ratios of 1.1–1.2 (typical for complex disease variants) requires thousands to tens of thousands of cases, not hundreds.

Small GWAS can still provide value through polygenic analyses aggregating information across many sub-threshold variants and through contributing data to meta-analyses. However, their inability to identify genome-wide significant loci frustrated early genetic investigation of ME/CFS and highlighted the necessity of large collaborative efforts.

Several methodological challenges complicate ME/CFS GWAS beyond simply achieving adequate sample size. Diagnostic heterogeneity creates noise: if different diagnostic criteria capture partially overlapping patient populations with different genetic architectures, this heterogeneity reduces power. Potential solutions include stratified analyses by diagnostic criteria and phenotype refinement using quantitative traits (severity scores, specific symptoms) rather than binary case-control status.

Population stratification—systematic ancestry differences between cases and controls—can produce spurious associations. Standard GWAS methods correct for stratification using principal components analysis of genetic data, ensuring cases and controls are matched for genetic ancestry. For ME/CFS, international collaborative GWAS must carefully model ancestry structure to avoid confounding.

The missing heritability problem—the gap between twin study heritability estimates and SNP heritability from GWAS—arises from several sources. Rare variants (minor allele frequency < 1%) not well captured by standard GWAS arrays may contribute to liability. Structural variants, copy number variations, and epigenetic modifications are not directly tested in GWAS. Gene-gene and gene-environment interactions may contribute to liability but are difficult to detect with current methods. Nevertheless, GWAS SNP heritability typically explains 20–50% of twin study heritability for complex diseases, providing genome-wide validation of genetic contribution while highlighting areas for future investigation.

Implications for Understanding ME/CFS Pathophysiology

GWAS findings illuminate disease mechanisms by identifying unexpected genes and pathways. When associated loci cluster in particular biological pathways or cell types, this convergence suggests mechanistic importance even if individual variants have small effects. For ME/CFS, pathway enrichment analyses can test whether associated variants cluster in immune pathways, metabolic pathways, neurological pathways, or other systems.

Gene-set enrichment might reveal, for example, that associated variants disproportionately affect genes expressed in natural killer cells, suggesting NK cell dysfunction has genetic determinants, or that variants cluster in mitochondrial pathways, supporting metabolic dysfunction as a genetically influenced component. Such findings validate hypotheses generated from physiological studies and suggest therapeutic targets.

GWAS also enable Mendelian randomization analyses testing causal relationships between exposures and ME/CFS. Using genetic variants as instrumental variables for exposures (analogous to randomized controlled trials), researchers can test whether, for example, genetically predicted vitamin D levels affect ME/CFS risk, or whether genetically predicted inflammatory markers causally contribute to disease liability. These analyses help distinguish causation from correlation in observational studies.

12.2 Epigenetic Modifications

Epigenetic modifications—chemical alterations to DNA and chromatin that regulate gene expression without changing DNA sequence—provide a plausible mechanism for how environmental triggers such as viral infections could produce lasting changes in cellular function. Unlike genetic variants that are inherited and static, epigenetic modifications are dynamic, potentially reversible, and responsive to environmental stimuli. In ME/CFS, epigenetic changes may explain how transient infections or stressors produce chronic alterations in immune function, metabolism, and neurological status.

The epigenetic landscape encompasses multiple interconnected mechanisms. DNA methylation silences gene expression by adding methyl groups to cytosine bases, particularly at CpG dinucleotides in gene promoters. Histone modifications alter chromatin structure through acetylation, methylation, phosphorylation, and other post-translational modifications of histone proteins, making genes more or less accessible to transcription machinery. MicroRNAs regulate gene expression post-transcriptionally by binding messenger RNAs and promoting their degradation or blocking translation. These mechanisms interact: DNA methylation patterns influence histone modifications, which in turn affect microRNA expression, creating integrated regulatory networks.

For ME/CFS, the epigenetic hypothesis proposes that triggering infections or stressors induce epigenetic reprogramming in immune cells, metabolic tissues, or neurological systems, and that this reprogramming persists after the trigger resolves, maintaining pathological cellular states. This model explains chronicity without requiring persistent infection and suggests potentially reversible mechanisms amenable to therapeutic intervention.

12.2.1 DNA Methylation

DNA methylation represents the most stable and well-characterized epigenetic modification, involving addition of methyl groups to cytosine bases primarily at CpG sites (cytosine-guanine dinucleotides). Gene promoters rich in CpG sites (CpG islands) are normally unmethylated, allowing transcription; methylation of promoter CpG islands typically silences gene expression. Conversely, gene body methylation and methylation of repetitive elements may have different functional consequences.

Global Methylation Patterns

Several studies have examined genome-wide DNA methylation patterns in ME/CFS patients compared to healthy controls using methylation array technologies that interrogate hundreds of thousands of CpG sites across the genome.

Observation 64 (DNA Methylation Studies in ME/CFS). de Vega et al. conducted epigenome-wide association studies (EWAS) examining DNA methylation in blood samples from ME/CFS patients and controls [375]. These studies identified differentially methylated positions (DMPs) and differentially methylated regions (DMRs) associated with ME/CFS status, with

several affected genes showing biological plausibility. Effect sizes are typically modest (methylation differences of 2–10%), consistent with complex disease epigenetics where subtle changes across many loci create cumulative functional effects.

Longitudinal studies examining methylation stability over time show that ME/CFS-associated methylation patterns persist, suggesting stable epigenetic reprogramming rather than transient stress responses. However, within-person variability has not been extensively characterized, leaving open questions about whether methylation patterns fluctuate with symptom severity or remain static.

Global methylation analyses reveal both hypomethylation and hypermethylation in ME/CFS, with different genes showing methylation changes in opposite directions. This bidirectional pattern contrasts with cancer epigenetics, where global hypomethylation and focal hypermethylation at tumor suppressors predominate. The ME/CFS methylation signature suggests dysregulated methylation machinery rather than unidirectional change, possibly reflecting altered activity of DNA methyltransferases (DNMTs) or ten-eleven translocation (TET) demethylases.

Gene-Specific Methylation Changes

Specific genes showing differential methylation in ME/CFS cluster in functionally relevant pathways, providing biological validation beyond statistical association.

Immune genes show notable methylation changes consistent with immune dysfunction phenotypes. Genes encoding cytokines, chemokines, and immune receptors demonstrate altered methylation in several studies. Hunter et al. using the EpiSwitch platform identified methylation signatures involving IL-2 pathway genes [215], consistent with the T cell dysfunction documented in Chapter 7. Methylation changes in immune regulatory genes could establish stable alterations in cytokine production capacity or immune cell responsiveness, contributing to chronic inflammation or immune exhaustion.

Metabolic genes affecting mitochondrial function, glucose metabolism, and oxidative stress responses show differential methylation. Given the profound metabolic dysfunction in ME/CFS (Chapter 6), epigenetic silencing of metabolic genes represents a plausible mechanism for persistent bioenergetic impairment. Methylation of genes encoding electron transport chain components, glycolytic enzymes, or oxidative phosphorylation machinery could reduce metabolic capacity even without genetic mutations.

Neurological and neurotransmitter genes demonstrate methylation changes that may relate to cognitive dysfunction and autonomic symptoms. Genes affecting neurotransmitter synthesis, reuptake, or receptor expression show altered methylation in some studies, potentially contributing to the neurological manifestations described in Chapter 8.

Functional Consequences and Validation

⚡ Limitation 1: Methylation ≠ Functional Consequence

Observing differential methylation does not establish functional consequence; methylation changes must alter gene expression to affect phenotype. Many differentially methylated sites have no demonstrated effect on transcription, and whole-blood methylation studies obscure cell-type-specific changes. Until methylation data are integrated with matched gene expression data in the same cohorts, differential methylation in ME/CFS remains a correlational finding.

Integration of methylation data with gene expression data addresses this question: do genes with altered methylation show corresponding changes in mRNA levels?

Several studies have performed integrative analyses correlating methylation with expression. For genes showing promoter hypermethylation, reduced mRNA expression would be expected; promoter hypomethylation should associate with increased expression. Many ME/CFS-associated methylation changes show the expected direction of expression change, supporting functional relevance. However, some differentially methylated genes show no expression change, possibly reflecting compensatory mechanisms, context-dependent effects (methylation may affect expression only in specific cell types or conditions), or methylation in regulatory regions outside proximal promoters.

Cell-type heterogeneity complicates interpretation. Whole blood methylation studies measure average methylation across multiple cell types (lymphocytes, monocytes, neutrophils, others), potentially obscuring cell-type-specific changes. If methylation changes occur predominantly in one cell type (for example, natural killer cells), analyzing bulk blood dilutes the signal. Future studies using cell-type-specific methylation profiling or single-cell technologies will better resolve this issue.

Methylation Age and Biological Aging

DNA methylation patterns change predictably with chronological age, enabling construction of epigenetic clocks that estimate biological age from methylation profiles. Accelerated epigenetic aging—biological age exceeding chronological age—associates with numerous age-related diseases and mortality risk.

Preliminary evidence suggests ME/CFS patients may show accelerated epigenetic aging, with methylation-based age estimates exceeding actual age. This finding, if replicated in larger cohorts, would support the hypothesis that ME/CFS involves accelerated biological aging processes affecting multiple physiological systems. The mechanisms underlying epigenetic age acceleration in ME/CFS remain unclear but could involve chronic oxidative stress, mitochondrial dysfunction, or chronic inflammation, all of which affect methylation patterns and associate with aging.

12.2.2 Histone Modifications

Histone proteins package DNA into nucleosomes, the fundamental units of chromatin structure. Post-translational modifications of histone tails—including acetylation, methylation, phosphorylation, ubiquitination, and others—regulate chromatin accessibility and gene expression. Histone acetylation generally activates transcription by relaxing chromatin structure, while histone methylation can either activate or repress transcription depending on which residue is modified and the degree of methylation.

Chromatin Remodeling in ME/CFS

Evidence for altered histone modifications in ME/CFS comes primarily from studies of immune cells, where chromatin remodeling regulates immune activation, differentiation, and exhaustion states.

~ Hypothesis 1: Epigenetic Basis of T Cell Exhaustion

T cell exhaustion—a state of progressive functional impairment occurring during chronic antigen exposure—involves characteristic epigenetic reprogramming that maintains exhaustion even after antigen removal. Exhausted T cells demonstrate specific histone modification patterns including reduced H3K27ac (active enhancer mark) at effector genes and increased H3K27me3 (repressive mark) at genes required for T cell function [119].

If ME/CFS involves chronic T cell exhaustion as discussed in Chapter 7, the epigenetic signatures of exhaustion should be detectable. T cells from ME/CFS patients might show chromatin states characteristic of exhaustion: closed chromatin at effector cytokine loci (IFN- γ , TNF- α), reduced accessibility at proliferation genes, and altered expression of exhaustion markers (PD-1, TIM-3, LAG-3). These epigenetic states would perpetuate T cell dysfunction even if the original triggering antigen is cleared, explaining chronicity and providing therapeutic targets (epigenetic modifying drugs might reverse exhaustion states).

Preliminary data examining histone modifications in ME/CFS immune cells show altered H3K4me3 (active transcription mark) and H3K27ac patterns compared to controls, with changes clustering at immune regulatory genes. The functional significance requires validation through chromatin accessibility assays (ATAC-seq or DNase-seq) determining whether altered histone marks correspond to changes in chromatin openness and gene expression.

Histone Acetylation and Metabolic Coupling

Histone acetylation depends on acetyl-CoA availability, creating a direct coupling between cellular metabolism and epigenetic regulation. Histone acetyltransferases (HATs) use acetyl-CoA as substrate to acetylate histone lysine residues; when acetyl-CoA levels fall (as occurs with mitochondrial dysfunction or glucose deprivation), histone acetylation decreases genome-wide, altering gene expression patterns.

This metabolic-epigenetic coupling may be particularly relevant in ME/CFS given the documented metabolic dysfunction (Chapter 6). Reduced mitochondrial ATP production and altered central carbon metabolism could decrease acetyl-CoA availability, leading to genome-wide hypoacetylation of histones. This hypoacetylation would reduce expression of acetylation-dependent genes, potentially creating a feedforward loop: metabolic dysfunction causes epigenetic changes that further impair metabolic gene expression, perpetuating metabolic impairment.

Similarly, histone methylation depends on S-adenosyl methionine (SAM) as methyl donor, linking one-carbon metabolism to chromatin regulation. Altered methionine or folate metabolism could affect SAM availability and thereby histone methylation patterns, providing another mechanism linking metabolism to epigenetic dysregulation.

Potential for Epigenetic Therapeutics

The reversibility of histone modifications makes them attractive therapeutic targets. Histone deacetylase (HDAC) inhibitors increase histone acetylation and are approved for cancer treatment; could they benefit ME/CFS by reversing pathological chromatin states? Histone demethylase inhibitors and histone methyltransferase inhibitors modulate specific methylation marks. Bromodomain inhibitors block proteins that recognize acetylated histones, altering transcriptional responses to acetylation.

However, these drugs have broad effects across the genome and significant toxicities, limiting their use to severe diseases. More targeted approaches might use small molecules affecting specific histone-modifying enzymes relevant to ME/CFS pathophysiology, or dietary interventions affecting metabolite availability (acetyl-CoA, SAM) that indirectly modulate histone modifications.

12.2.3 MicroRNAs

MicroRNAs (miRNAs) are small non-coding RNAs approximately 22 nucleotides in length that regulate gene expression post-transcriptionally. A single miRNA can target hundreds of messenger RNAs (mRNAs), and a single mRNA can be targeted by multiple miRNAs, creating complex regulatory networks. miRNAs bind to complementary sequences in target mRNA 3' untranslated regions, promoting mRNA degradation or blocking translation, thereby reducing protein expression.

Altered MicroRNA Profiles in ME/CFS

Multiple studies have examined miRNA expression in ME/CFS patients' blood samples using miRNA profiling technologies. These studies identify differentially expressed miRNAs—miRNAs showing significantly higher or lower expression in patients compared to controls.

Observation 65 (MicroRNA Dysregulation in ME/CFS). Brenu et al. and other groups have reported altered expression of specific miRNAs in ME/CFS, with different studies showing partial but incomplete overlap in identified miRNAs [202]. Commonly reported dysregulated miRNAs include those regulating immune function (miR-21, miR-146a, miR-155), metabolism, and stress responses. Sample sizes in published studies are generally small ($n = 20\text{--}50$ per group), limiting statistical power and increasing risk of false positives.

The lack of consistent replication across studies may reflect genuine heterogeneity in miRNA profiles across ME/CFS subgroups, different patient selection criteria, different analytical platforms, or statistical issues. Larger cohorts with standardized protocols are needed to establish robust miRNA signatures.

Specific miRNAs showing altered expression in ME/CFS have plausible biological relevance. miR-21 regulates immune responses and fibrosis; increased miR-21 could contribute to immune dysfunction or tissue remodeling. miR-146a functions as a negative regulator of innate immunity, dampening inflammatory responses; altered miR-146a expression might affect inflammatory tone. miR-155 promotes inflammatory macrophage activation; dysregulation could affect immune cell polarization.

MiRNAs targeting metabolic pathways show expression changes in some studies, potentially contributing to metabolic dysfunction. MiRNAs regulating mitochondrial genes, glycolytic enzymes, or oxidative stress responses could alter cellular energetics if their expression is perturbed.

Regulatory Effects and Target Validation

Identifying differentially expressed miRNAs is only the first step; understanding functional consequences requires determining which target mRNAs are actually affected. Computational prediction algorithms identify potential miRNA targets based on sequence complementarity, but experimental validation is necessary because many predicted targets are not functionally regulated.

Integrative analysis comparing miRNA expression with mRNA expression can identify functional targets: if a miRNA is upregulated, its target mRNAs should show decreased expression; downregulated miRNAs should associate with increased target expression. Several ME/CFS studies have performed such analyses, identifying inverse correlations between miRNA expression and predicted targets, supporting functional regulation.

However, the magnitude of miRNA effects on individual targets is often modest (20–40% reduction in protein expression), and biological effects may require coordinated regulation of multiple targets within a pathway. Network analyses examining whether dysregulated miRNAs converge on common pathways provide systems-level understanding: do multiple altered miRNAs target immune pathways, metabolic pathways, or neurological pathways?

MicroRNAs as Biomarkers

Beyond their mechanistic role, circulating miRNAs represent potential biomarkers for ME/CFS diagnosis, prognosis, or treatment response monitoring. miRNAs are stable in blood, resistant to degradation, and quantifiable using standard techniques, making them attractive biomarker candidates.

→ Prediction 1: MicroRNA Biomarker Panels

If ME/CFS has a characteristic miRNA signature, panels of multiple miRNAs could achieve diagnostic sensitivity and specificity adequate for clinical use. A diagnostic test combining 5–10 miRNAs with clinical criteria might improve diagnostic accuracy beyond current symptom-based approaches.

For biomarker development, several criteria must be met: differential expression must replicate in independent cohorts, diagnostic accuracy (sensitivity and specificity) must exceed threshold for clinical utility (generally >80% for both), and miRNA levels must be stable over time in individual patients unless they correlate meaningfully with symptom severity. Additionally, miRNA signatures must distinguish ME/CFS from conditions with overlapping symptoms (fibromyalgia, depression, primary sleep disorders).

Current evidence does not yet support clinical miRNA biomarker use for ME/CFS. Replication remains incomplete, effect sizes are modest, and head-to-head comparisons with overlapping conditions are limited. However, ongoing studies with larger sample sizes and standardized protocols may identify robust signatures warranting clinical validation.

Circulating vs Tissue-Specific MicroRNAs

An important question concerns the cellular source of differentially expressed miRNAs in blood. Circulating miRNAs may originate from blood cells themselves (lymphocytes, monocytes), reflecting altered immune cell miRNA expression. Alternatively, miRNAs may be released from tissues (muscle, brain, gut) in extracellular vesicles or bound to proteins, providing a window into tissue dysfunction not directly accessible through blood sampling.

Cell-type-specific miRNA profiling (isolating specific cell populations before miRNA extraction) can determine whether miRNA changes occur broadly across blood cells or specifically in subsets such as natural killer cells, T cells, or monocytes. Tissue-specific miRNAs can be identified through expression databases showing which miRNAs are enriched in particular tissues; finding muscle-enriched miRNAs elevated in ME/CFS patients' plasma might indicate muscle pathology.

Understanding miRNA cellular origin informs interpretation: immune cell-intrinsic miRNA changes suggest altered immune cell programming, while tissue-derived miRNAs suggest tissue damage or dysfunction with secondary release of cellular contents into circulation.

12.3 Gene Expression Patterns

Gene expression profiling using transcriptomics technologies provides a functional readout of genetic and epigenetic regulation, measuring which genes are actively transcribed into messenger RNA and the magnitude of expression changes. In ME/CFS, gene expression studies illuminate which biological pathways are dysregulated, identify potential biomarkers, and suggest mechanisms linking genetic susceptibility to phenotypic manifestations. Unlike static genetic variants, gene expression is dynamic and potentially responsive to interventions, making dysregulated genes attractive therapeutic targets.

12.3.1 Transcriptomics Studies and Methodological Considerations

Multiple gene expression studies in ME/CFS have used microarray and RNA sequencing technologies to measure mRNA levels genome-wide in blood samples, comparing patients to healthy controls. These studies vary in sample size (ranging from $n=20$ to $n>100$ per group), patient selection criteria (CCC, ICC, Fukuda), sample types (whole blood, PBMCs, specific cell populations), and analytical approaches.

Methodological heterogeneity complicates cross-study comparison. Whole blood gene expression reflects the aggregate signal from multiple cell types—lymphocytes, monocytes, neutrophils, eosinophils, basophils—each with distinct transcriptional profiles. If ME/CFS involves altered proportions of these cell types (for example, increased proportion of exhausted T cells, reduced NK cells), whole blood expression changes may reflect cell composition differences rather than cell-intrinsic transcriptional changes. Statistical methods can partially address this through deconvolution algorithms estimating cell-type proportions, but cell-type-specific profiling provides more definitive answers.

Batch effects—systematic technical differences between sample processing batches—can produce spurious expression differences larger than biological signal. Rigorous studies randomize samples across batches, include technical replicates, and apply batch correction algorithms. Many early ME/CFS gene expression studies lacked adequate batch effect control, potentially contributing to replication failures.

Despite these challenges, convergent findings across independent studies provide evidence for robust gene expression changes in ME/CFS, particularly in immune and metabolic pathways.

12.3.2 Differentially Expressed Genes

Differentially expressed genes (DEGs)—genes showing statistically significant expression differences between patients and controls—number in the hundreds to thousands in typical ME/CFS transcriptomics studies, depending on statistical thresholds and multiple testing correction methods.

Immune System Gene Expression

★ Achievement 2: Convergent Immune Gene Dysregulation

Across multiple independent gene expression studies, immune pathway genes show the most consistent and pronounced dysregulation. Key patterns include:

Cytokine and chemokine genes demonstrate altered expression consistent with chronic immune activation or altered cytokine networks. Pro-inflammatory cytokine genes (IL-1 β , TNF- α , IL-6) show variable direction of change across studies, likely reflecting patient heterogeneity and disease stage. Chemokine genes affecting immune cell trafficking (CCL2, CXCL10, others) demonstrate differential expression in multiple studies.

T cell and NK cell genes show expression patterns consistent with functional impairment. T cells from ME/CFS patients demonstrate reduced expression of effector cytokine genes (IFN- γ , TNF- α) and altered expression of exhaustion markers (PDCD1 encoding PD-1, HAVCR2 encoding TIM-3, LAG3). NK cell gene expression profiling reveals reduced expression of cytotoxic effector genes (PRF1 encoding perforin, GZMA/GZMB encoding granzymes) consistent with the impaired cytotoxicity documented in Chapter 7.

Interferon-stimulated genes (ISGs) show elevated expression in multiple studies, suggesting ongoing antiviral responses or interferon pathway activation even in the absence of detectable active infection. This ISG signature resembles that seen in autoimmune diseases such as systemic lupus erythematosus and may indicate chronic stimulation of pattern recognition receptors or dysregulated interferon regulatory factor activity.

△ Warning 3: Replication Status: Partially Replicated

This achievement synthesizes across multiple studies without providing specific citations within the block. While individual findings (NK cell cytotoxicity impairment, T cell exhaustion markers, ISG elevation) have each been reported in independent studies, the specific “convergent” pattern described here is an interpretive synthesis rather than a directly replicated result. The direction and magnitude of cytokine gene changes vary substantially across studies, likely reflecting patient heterogeneity, disease stage, and methodological differences. The claim of convergence should be evaluated against systematic reviews rather than individual study comparisons.

Metabolic Gene Expression

Genes involved in energy metabolism, mitochondrial function, and oxidative stress responses demonstrate altered expression patterns consistent with the metabolic dysfunction detailed in Chapter 6.

Mitochondrial genes show variable but frequently reduced expression across studies. Nuclear-encoded mitochondrial genes affecting oxidative phosphorylation, the tricarboxylic acid cycle, and mitochondrial biogenesis may show downregulation, potentially contributing to reduced mitochondrial ATP production capacity. However, the magnitude and consistency of these changes varies across studies, possibly reflecting differences in disease severity, duration, or patient selection.

Glycolytic pathway genes show altered expression in some studies, with evidence for both increased glycolytic gene expression (potentially compensatory for mitochondrial dysfunction) and reduced expression. The direction and magnitude may depend on metabolic state at the time of sampling (resting versus post-exertional).

Genes encoding oxidative stress response proteins (superoxide dismutase, catalase, glutathione synthesis and recycling enzymes) demonstrate altered expression, consistent with increased oxidative stress burden. Some studies report upregulation suggesting compensatory induction, while others find downregulation potentially reflecting exhausted antioxidant capacity.

Neurological and Neurotransmitter Genes

Gene expression changes affecting neurological function and neurotransmitter systems may contribute to cognitive dysfunction and neurological symptoms (Chapter 8).

Neurotransmitter synthesis, transport, and receptor genes show differential expression in some studies. Genes affecting serotonin, dopamine, and norepinephrine metabolism demonstrate variable changes across patients, potentially reflecting heterogeneity in neurological symptom profiles. Ion channel genes including TRPM3 show altered expression in ME/CFS patients, consistent with the functional TRPM3 deficiency discussed in Section 12.1.2.

Blood-brain barrier integrity genes and neuroinflammatory markers show expression changes in some studies, though interpreting peripheral blood expression of brain-related genes requires caution. Elevated expression of neuroinflammation-associated genes may reflect systemic inflammation affecting the CNS or glial activation with release of inflammatory mediators detectable peripherally.

12.3.3 Pathway Enrichment and Systems Biology Analysis

Individual differentially expressed genes provide limited insight without biological context. Pathway enrichment analysis tests whether DEGs cluster in particular biological pathways or functional categories more than expected by chance, identifying dysregulated biological processes.

Observation 66 (Pathway-Level Convergence in ME/CFS). Despite variable lists of specific DEGs across studies, pathway enrichment analyses show remarkable convergence, with multiple independent studies identifying the same biological pathways as dysregulated:

Immune response pathways including innate immunity, antiviral responses, cytokine signaling, and T cell activation emerge as top enriched pathways in essentially all ME/CFS gene expression studies. This pathway-level convergence validates immune dysfunction as a core feature even when specific DEGs differ.

Metabolic pathways including oxidative phosphorylation, TCA cycle, fatty acid metabolism, and glucose metabolism show enrichment in multiple studies, supporting metabolic dysfunction as a consistent feature.

Cellular stress response pathways including unfolded protein response, endoplasmic reticulum stress, and oxidative stress responses demonstrate enrichment, suggesting chronic cellular stress across multiple compartments.

Circadian rhythm and sleep-related pathways show dysregulation in some studies, potentially relating to sleep dysfunction and circadian rhythm disturbances common in ME/CFS.

Network analysis approaches examining interactions between DEGs identify hub genes—highly connected genes whose dysregulation may have outsized effects on pathway function. These hub genes represent priority targets for mechanistic investigation and potential therapeutic intervention.

12.3.4 Cell Type-Specific Expression and Single-Cell Approaches

Bulk tissue gene expression confounds cell-intrinsic transcriptional changes with cell composition differences. Cell-type-specific profiling addresses this limitation by isolating specific cell populations before expression analysis or using computational deconvolution.

Natural killer cell-specific gene expression studies reveal pronounced transcriptional changes consistent with NK cell dysfunction, including reduced expression of cytotoxic genes and altered expression of activation and inhibitory receptors. These cell-intrinsic changes validate that NK cell dysfunction reflects altered cellular programming, not simply reduced NK cell numbers.

T cell subset-specific profiling distinguishes CD4+ helper T cells, CD8+ cytotoxic T cells, and regulatory T cells, each with distinct expression signatures. ME/CFS studies have reported differential expression patterns across subsets, with some suggesting particular dysregulation in CD8+ T cells consistent with exhaustion phenotypes.

Emerging single-cell RNA sequencing (scRNA-seq) technologies enable simultaneous profiling of thousands of individual cells, identifying rare cell populations and cell state heterogeneity invisible to bulk sequencing. Preliminary scRNA-seq studies in ME/CFS are beginning to reveal subpopulations of immune cells with distinct transcriptional states, potentially including exhausted T cell states, activated monocyte populations, or dysfunctional NK cell subsets. As scRNA-seq becomes more widely applied, it promises to resolve cellular heterogeneity and identify specific cell states driving pathology.

12.3.5 Exercise-Induced Gene Expression Changes

Post-exertional malaise represents the cardinal symptom of ME/CFS, making exercise-induced gene expression changes particularly relevant. Several studies have examined gene expression before and after standardized exercise challenges, identifying genes whose expression changes abnormally in ME/CFS patients compared to healthy controls.

Observation 67 (Exercise-Induced Transcriptional Response). Healthy individuals show characteristic exercise-induced gene expression changes reflecting metabolic adaptation, immune modulation, and cellular repair. ME/CFS patients demonstrate altered exercise responses, with exaggerated or prolonged expression changes in immune genes, blunted metabolic adaptation, and sustained stress response gene activation.

Specific patterns include:

- Prolonged elevation of immune activation genes 24–72 hours post-exercise, corresponding to symptom exacerbation timing
- Reduced or delayed upregulation of metabolic adaptation genes that normally facilitate recovery
- Sustained activation of cellular stress response pathways
- Altered expression of genes regulating muscle metabolism and repair

These exercise-induced expression changes correlate with symptom severity in some studies, suggesting gene expression profiles might objectively quantify PEM severity and duration.

Longitudinal sampling capturing expression changes at multiple timepoints (pre-exercise, immediately post, +4h, +24h, +48h, +72h) reveals temporal dynamics invisible to single-timepoint studies. Such temporal profiling may identify early molecular events initiating PEM and later events perpetuating symptoms, with therapeutic implications for targeting specific phases.

12.3.6 Integration with Genetic and Epigenetic Data

The most powerful insights emerge from integrating gene expression with genetic and epigenetic data, identifying genes where genetic variants affect expression levels (expression quantitative trait loci, eQTLs), genes showing coordinated methylation and expression changes, and genes targeted by dysregulated microRNAs.

Expression QTL analysis asks whether genetic variants identified in GWAS or candidate gene studies actually affect expression of nearby or distant genes. For ME/CFS-associated genetic variants, demonstrating that risk alleles correlate with altered expression of biologically plausible genes strengthens causal inference and identifies mechanisms by which genetic variants influence disease risk.

Integrative methylation-expression analysis identifies genes showing inverse correlations between promoter methylation and mRNA expression, validating functional consequences of

epigenetic changes. Genes demonstrating both differential methylation and corresponding expression changes represent high-priority mechanistic targets.

MicroRNA-mRNA correlation analysis tests whether dysregulated miRNAs actually affect predicted target expression. Negative correlations between miRNA expression and target mRNA expression support functional regulatory relationships and help distinguish direct miRNA targets from indirect effects.

These integrative analyses transform lists of genes, variants, methylation sites, and miRNAs into mechanistic models specifying causal chains: genetic variant → altered methylation → changed miRNA expression → dysregulated target gene expression → pathway dysfunction → phenotype.

12.4 Synthesis and Open Questions

Genetic and epigenetic research in ME/CFS has matured substantially over the past decade, progressing from underpowered candidate gene studies to genome-wide approaches, from speculation about epigenetic involvement to empirical demonstration of DNA methylation and histone modification changes, and from simple gene lists to integrated multi-omics analyses. This body of evidence establishes genetic predisposition as a substantial contributor to ME/CFS risk while highlighting the complex polygenic architecture and gene-environment interactions that determine disease manifestation.

12.4.1 Key Established Findings

★ Achievement 3: Genetic and Epigenetic Foundations of ME/CFS

Several conclusions now rest on firm empirical ground:

Moderate heritability Twin studies consistently demonstrate heritability estimates of $h^2 = 0.3\text{--}0.5$, indicating that 30–50% of disease liability reflects genetic factors. This moderate heritability implies substantial genetic contribution while confirming environmental factors' essential role.

Polygenic architecture DecodeME and other GWAS findings confirm that ME/CFS arises from the cumulative effects of many genetic variants of small individual effect rather than single genes of large effect. This polygenic model aligns with other complex diseases and explains familial clustering without Mendelian inheritance patterns.

Gene-environment interaction The observation that only a minority of individuals exposed to triggering infections develop ME/CFS, combined with familial aggregation patterns, validates gene-environment interaction as the central etiological framework. Genetic variants establish vulnerability; environmental triggers are necessary for disease

expression.

Immune and metabolic pathway enrichment Gene expression and pathway analyses consistently implicate immune response pathways and metabolic dysfunction. This convergence across independent studies and methodologies validates immune and metabolic dysregulation as core pathophysiological features with genetic determinants.

Epigenetic reprogramming Demonstration of DNA methylation changes, altered histone modifications, and dysregulated microRNA expression establishes epigenetic reprogramming as a plausible mechanism for chronicity. These potentially reversible modifications provide therapeutic targets.

△ **Warning 4: Replication Status: Variable**

This synthesis achievement combines findings of differing replication strength. Moderate heritability ($h^2 = 0.3\text{--}0.5$) from twin studies is the most robust element, replicated across multiple independent twin registries. Polygenic architecture is confirmed by DecodeME but locus-level replication is pending (see Achievement 12.1.3). Gene-environment interaction is well-supported conceptually but the specific genetic variants mediating susceptibility remain largely unidentified. Epigenetic reprogramming evidence comes from small studies with limited replication—DNA methylation findings in particular show inconsistent patterns across studies, partly due to cell-type heterogeneity in whole-blood samples.

12.4.2 Integration with Broader ME/CFS Pathophysiology

The genetic and epigenetic findings detailed in this chapter provide the constitutional substrate upon which the dysregulated physiological systems described in other chapters develop.

The immune dysfunction documented in Chapter 7—NK cell cytotoxicity impairment, T cell exhaustion, cytokine dysregulation—reflects both genetic predisposition (HLA types, immune gene variants, TRPM3 polymorphisms) and epigenetic reprogramming (T cell chromatin states characteristic of exhaustion, methylation of immune genes). Genetic susceptibility determines baseline immune function capacity; epigenetic changes following infection establish chronic dysfunction states.

The metabolic dysfunction of Chapter 6—reduced oxidative phosphorylation, impaired ATP production, glycolytic shifts—similarly combines genetic vulnerability (mitochondrial gene variants, metabolic enzyme polymorphisms) with acquired epigenetic silencing of metabolic genes. The gene expression patterns show reduced expression of mitochondrial and metabolic pathway genes, potentially reflecting both genetic determinants of baseline expression and epigenetic downregulation following metabolic stress.

Neurological manifestations (Chapter 8) may reflect TRPM3 dysfunction (with genetic variants affecting baseline expression and function), neurotransmitter system genetic variants, and epigenetic changes affecting blood-brain barrier integrity and neuroinflammation. The

cognitive and autonomic symptoms could arise from the intersection of genetic liability and acquired epigenetic modifications.

This integrated model suggests that ME/CFS arises when genetic predisposition across multiple systems (immune, metabolic, neurological, cardiovascular) encounters environmental triggers sufficient to induce epigenetic reprogramming. The specific symptom profile reflects which genetic vulnerabilities predominate and which epigenetic changes occur, explaining clinical heterogeneity. Chapter 13 builds upon this genetic-epigenetic foundation to develop comprehensive multi-system models of ME/CFS pathophysiology, examining how constitutional factors interact with acquired dysfunction to create stable pathological states (Section 13.3).

12.4.3 Unresolved Questions and Future Directions

Despite substantial progress, critical questions remain:

? Open Question 1: Causal Variants and Mechanisms

GWAS identifies associated genomic loci but typically does not pinpoint causal variants or affected genes. For DecodeME-identified loci, fine-mapping studies using dense genotyping and functional genomics are needed to identify specific causal variants, determine which genes they affect, and elucidate mechanisms by which they influence disease risk. Do ME/CFS risk variants affect transcription factor binding sites, alter splicing, modify protein sequence, or influence other molecular processes?

? Open Question 2: Epigenetic Causality vs Consequence

Observed epigenetic changes could represent disease-driving mechanisms or secondary consequences of chronic illness. Longitudinal studies examining epigenetic changes before, during, and after disease onset would address causality. Do epigenetic changes precede symptom development in at-risk individuals? Do they persist during remission or normalize with symptom improvement? Can experimentally reversing specific epigenetic modifications (using CRISPR-based epigenome editing or small molecule epigenetic drugs) alleviate cellular dysfunction in patient cells?

? Open Question 3: Genetic Subtyping

Does ME/CFS comprise genetically distinct subtypes with different molecular mechanisms? Cluster analyses based on genetic profiles, gene expression patterns, or epigenetic signatures might identify patient subgroups with different pathophysiological mechanisms, prognoses, and treatment responses. Such molecular subtyping could enable personalized treatment selection.

? Open Question 4: Therapeutic Reversibility

Given that epigenetic modifications are potentially reversible, can therapeutic interventions normalize methylation patterns, histone modifications, or microRNA expression? Would such normalization translate to clinical improvement? Trials of epigenetic-modifying drugs (HDAC inhibitors, methyltransferase inhibitors, demethylating agents) could test this hypothesis, though broad epigenetic drugs have significant toxicities. More targeted approaches using small molecules affecting specific epigenetic enzymes or dietary interventions affecting metabolite availability might offer safer therapeutic windows.

? Open Question 5: Prevention in High-Risk Individuals

Can polygenic risk scores identify individuals at high genetic risk who might benefit from preventive interventions? Following the 2004 Bergen Giardia outbreak model, future post-infection cohorts could stratify by genetic risk and test whether early interventions (aggressive rest, anti-inflammatory treatments, metabolic support) prevent chronic illness development in high-risk individuals. Such prevention trials could validate genetic risk prediction and identify modifiable factors in the gene-environment interaction.

? Open Question 6: Cross-Condition Genetic Architecture

Genetic correlation analyses suggest shared genetic liability between ME/CFS, fibromyalgia, irritable bowel syndrome, and Long COVID. Do these conditions represent different manifestations of the same underlying genetic vulnerability, or do they have partially overlapping but distinct genetic architectures? Detailed comparison of GWAS findings across conditions would address this question and might reveal common therapeutic targets applicable across multiple chronic overlapping pain conditions.

The genetic and epigenetic foundations of ME/CFS, while increasingly well characterized, point toward a future of precision medicine approaches where genetic profiling informs diagnosis, prognostication, and treatment selection, and where therapeutic interventions target the specific molecular pathways dysregulated in individual patients.

13 Integrative Models and Related Phenomena

“All models are wrong, but some are useful.”
— George E.P. Box

This chapter attempts to synthesize the diverse findings presented in previous chapters into coherent models of ME/CFS pathophysiology. We present these models with explicit acknowledgment of their evidence levels, from well-established observations to speculative hypotheses. The goal is intellectual honesty: to distinguish what we know, what we suspect, and what we’re guessing.

13.1 Evidence Classification Framework

Before presenting hypotheses, we define our evidence classification system. This framework is conservative—we classify based on the *weakest* link in the evidence chain.

Observation 68 (Honest Uncertainty). Much of what follows involves substantial uncertainty. The ME/CFS field has been plagued by premature certainty—both from those who dismissed the illness as psychological and from those who promoted specific biological theories without adequate evidence. We aim to avoid both errors by clearly labeling our confidence levels and acknowledging where we may be wrong.

Table 13.1: Evidence Level Definitions

Level	Definition	What This Means
Established	Replicated in multiple independent studies with consistent findings	High confidence this is real; disagreement is about interpretation, not existence
Probable	Documented in ≥ 2 studies OR single large/well-designed study	Likely real, but replication needed; could be overturned
Preliminary	Single study or small studies with suggestive findings	Interesting signal, but may not replicate; treat as hypothesis
Theoretical	Biologically plausible based on known mechanisms, but not directly tested in ME/CFS	Reasonable extrapolation from other conditions; needs direct testing
Speculative	Creative hypothesis without direct supporting data	May inspire research but should not guide treatment decisions

13.2 Comprehensive Hypothesis Ranking

Table 13.2 presents the major hypotheses about ME/CFS pathophysiology, ranked by our assessment of their likelihood of being substantially correct. This ranking is inherently subjective and will change as new evidence emerges. We weight: (1) quality and quantity of direct evidence, (2) explanatory power for core symptoms, (3) consistency with treatment responses, and (4) biological plausibility.

Table 13.2: Ranked Hypotheses of ME/CFS Pathophysiology

Hypothesis	Evidence Level	Key Supporting Evidence	Explains Which Symptoms/Observations	Treatment Implications	Potential for Rapid Benefit
TIER 1: ESTABLISHED PHENOMENA					
Post-exertional malaise (PEM) as cardinal feature (§2.1)	Established	2-day CPET studies; universal patient reports; objective physiological decline on day 2	Exercise intolerance; delayed crashes; why GET harms	Pacing; energy management; avoid overexertion	High (pacing prevents crashes)
Autonomic dysfunction (§2.4)	Established	Abnormal tilt table tests; HRV abnormalities; POTS prevalence >30%	Orthostatic intolerance; tachycardia; temperature dysregulation; coat hanger pain	Salt/fluids; compression; fludrocortisone; midodrine; ivabradine	Moderate–High
Sleep architecture abnormalities (§2.2)	Established	Polysomnography showing reduced slow-wave, fragmented sleep; universal unrefreshing sleep	Unrefreshing sleep; cognitive dysfunction; fatigue	Sleep hygiene; low-dose trazodone; address comorbid sleep disorders	Moderate
Immune dysregulation (ch07)	Established	Cytokine abnormalities; NK cell dysfunction; T cell subset changes; B cell abnormalities	Flu-like symptoms; susceptibility to infections; post-infectious onset	LDN; immunomodulators; avoid immune stressors	Moderate
TIER 2: PROBABLE MECHANISMS					
Mitochondrial/energy metabolism dysfunction (§6.2)	Probable	ATP profile abnormalities; Heng 2025 AMP/ADP elevation; lactate abnormalities; metabolomic signatures	Fatigue; exercise intolerance; PEM; muscle weakness	CoQ10; NAD ⁺ precursors; D-ribose; B vitamins	Low–Moderate

Continued on next page

Table 13.2 – continued from previous page

Hypothesis	Evidence Level	Key Supporting Evidence	Explains Which Symptoms/Observations	Treatment Approaches	Implications	Potential for Rapid Benefit
Neuroinflammation (ch08)	Probable	PET imaging (Nakatomi); CSF abnormalities; microglial activation markers	Brain fog; cognitive dysfunction; sensory sensitivities; headaches	Anti-inflammatory approaches; LDN; avoid neuroinflammatory triggers		Low–Moderate
GPCR autoantibodies (§14.13)	Probable	Elevated anti-β2, M3, M4 antibodies [71, 72]; correlation with symptoms [208]; immunoabsorption responses [118]; monocyte dysfunction [209]	Autonomic dysfunction; fatigue; muscle symptoms; cytokine dysregulation; why some respond to IA	Immunoabsorption; BC007 [10]; daratumumab [9]		Moderate–High (in subset)
Gut microbiome dysbiosis (ch14)	Probable	Reduced butyrate producers; altered diversity; correlation with symptoms	GI symptoms; systemic inflammation; food intolerances	Probiotics; dietary modification; possibly FMT		Low–Moderate
Reduced cerebral blood flow	Probable	SPECT/MRI showing hypoperfusion; correlation with cognitive symptoms	Brain fog; cognitive dysfunction; orthostatic cognitive worsening	Address underlying POTS; potentially vasodilators		Moderate
TIER 3: PRELIMINARY/EMERGING						
Plasma cell-mediated autoimmunity (§14.13.2)	Preliminary	Daratumumab pilot (60% response); explains rituximab failure; IgG reduction correlates with response	Autoimmune subset; why B-cell depletion failed but plasma cell depletion worked	Daratumumab; combined IA + plasma cell targeting		High (in autoimmune subset)

Continued on next page

Table 13.2 – continued from previous page

Hypothesis	Evidence Level	Key Supporting Evidence	Explains Which Symptoms/Observations	Treatment Interventions	Implications	Potential for Rapid Benefit
Vascular-Immune-Energy Triad	Preliminary	Heng 2025 7-biomarker panel; Hoel 2026 serum proteomics cross-validation [191]; coordinated abnormalities across 3 systems from two independent cohorts	Multi-system nature; why single-target treatments fail	Triple-target col; simultaneous intervention		Unknown (untested)
Endothelial dysfunction / microclotting (ch14)	Preliminary	Elevated VWF, fibronectin, thrombospondin; Hoel 2026: reduced VIM, ANTXR2, CXADR + disrupted Ephrin signaling [191]; Long COVID microclot findings	Exercise intolerance; brain fog; multi-system involvement	Anticoagulation; fibrinolytics; endothelial support; Ephrin-targeted therapy?		Moderate (if confirmed)
Central catecholamine deficiency	Preliminary	Walitt 2024 CSF findings (reduced DOPA, DOPAC, DHPG); effort preference abnormality	Altered effort perception; motivation difficulties; why “pushing through” fails	Dopamine precursors?; stimulants with caution		Unknown
NAD ⁺ depletion (ch14)	Preliminary	Metabolomic abnormalities; 2025 NR trial in Long COVID; theoretical PARP consumption	Energy failure; mitochondrial dysfunction; immune cell dysfunction	NR/NMN 1000–2000 mg; prolonged treatment (>10 weeks)		Low (slow onset)

Continued on next page

Table 13.2 – continued from previous page

Hypothesis	Evidence Level	Key Supporting Evidence	Explains Which Symptoms/Observations	Treatment	Implications	Potential for Rapid Benefit
Small fiber neuropathy	Preliminary	Skin biopsy studies; correlation with dysautonomia; elevated in subset	Pain; autonomic symptoms; temperature regulation issues	IVIG (in some); immunomodulation; symptom management		Moderate (in subset)
Viral persistence/reactivation (ch14)	Preliminary	HHV-6 miRNA in CNS; elevated herpesvirus antibodies; EBV reactivation markers	Post-infectious onset; relapsing course; why antivirals help some	Valacyclovir; valganciclovir; potentially IVIG		Low–Moderate
EBV-driven CNS autoimmunity	Preliminary	EBV-infected B cells cross BBB [164]; LMP1 expression enables brain infiltration; complement/microglial activation	Post-EBV onset; neuroinflammation; brain fog distinct from peripheral fatigue	Antivirals; B cell depletion; complement inhibition		Moderate (in EBV+ subset)
Autoantibody-monocyte reprogramming (§14.13.2)	Preliminary	GPCR autoantibodies reprogram monocyte cytokine production [209]; MIP-1 δ , PDGF-BB, TGF- β 3 elevation	Systemic inflammation; why effects persist beyond receptor binding; tissue remodeling	Autoantibody removal + monocyte modulation (JAK inhibitors)		Moderate–High
TIER 4: THEORETICAL						
Glymphatic clearance failure (§14.3)	Theoretical	Sleep dysfunction; cognitive symptoms; cranio-cervical junction issues in subset	Brain fog; unrefreshing sleep; position-dependent symptoms	Address CCI if present; optimize slow-wave sleep		Unknown

Continued on next page

Table 13.2 – continued from previous page

Hypothesis	Evidence Level	Key Supporting Evidence	Explains Which Symptoms/Observations	Treatment Implications	Potential for Rapid Benefit
Tryptophan/kynurenine trap (§14.7)	Theoretical	IDO activation documented; tryptophan pathway abnormalities; elevated QUIN:KYNA ratio in some studies	Depression-like symptoms; neuroinflammation; NAD ⁺ depletion	IDO inhibitors?; shift pathway toward KYNA	Unknown
Circadian desynchronization (ch14)	Theoretical	Cortisol rhythm abnormalities; sleep timing issues; fluctuating symptoms	Unpredictable symptom patterns; unrefreshing sleep; why timing matters	Chronotherapy; melatonin; time-restricted feeding; light therapy	Moderate
Epigenetic “lock”	Theoretical	DNA methylation changes documented; duration predicts prognosis; why early intervention helps	Persistence; treatment resistance; why disease stabilizes	Epigenetic modifiers (experimental); early aggressive treatment	Unknown
Purinergic signaling dysregulation	Theoretical	ATP is danger signal; P2X7 and inflammation; exercise releases ATP	PEM delay (24–72h matches DTH kinetics); pain sensitization; inflammation	P2X7 antagonists (experimental)	Unknown
TIER 5: SPECULATIVE					
“Safe mode” / stuck sickness behavior	Speculative	Fits symptom pattern; evolutionarily plausible; explains why pushing harms	All core symptoms as adaptive (but stuck) response	Reset hypothalamic setpoint?; break the “lock”	Unknown

Continued on next page

Table 13.2 – continued from previous page

Hypothesis	Evidence Level	Key Supporting Evidence	Explains Which Symptoms/Observations	Treatment	Implications	Potential for Rapid Benefit
HERV reactivation	Speculative	HERVs can be desilenced; would explain persistent immune activation without pathogen	Post-viral onset; autoimmunity; female predominance	Antiretrovirals?; genetic silencing?	epi-	Unknown
Ion channel autoimmunity	Speculative	Precedent in other conditions (LEMS, MG); would explain “wired but tired”	Sensory sensitivities; autonomic dysfunction; muscle fatigue; cardiac symptoms	Plasmapheresis; IVIG; channel-specific interventions	inter-	Moderate (if confirmed)
Receptor internalization (not blockade)	Speculative	NMDA receptor autoantibodies cause internalization [211]; would explain lag between Ab removal and recovery	Why symptoms persist after immunoadsorption; need for receptor regeneration time	Autoantibody removal + time for receptor resynthesis		Moderate (delayed)
Lactate compartmentalization (MCT dysfunction)	Speculative	Lactate abnormalities documented; would explain tissue-specific symptoms	PEM; muscle symptoms; brain fog; why systemic lactate seems okay	DCA?; lactate supplementation?		Unknown
Ferroptosis susceptibility	Speculative	Lipid abnormalities; oxidative stress; iron dysregulation documented	Why high-energy tissues affected; why iron supplementation can harm	Ferroptosis inhibitors; careful with iron		Unknown

Continued on next page

Table 13.2 – continued from previous page

Hypothesis	Evidence Level	Key Evidence	Supporting Evidence	Explains Which Symptoms/Observations	Treatment Options	Implications	Potential for Rapid Benefit
Trained endotheliopathy	Speculative	Endothelial markers elevated (Heng 2025); Hoel 2026 impaired repair signature (reduced ANTXR2, VIM, Ephrin disruption) [191]; innate immune training established; vascular symptoms		Multi-system involvement; persistent endothelial activation; impaired vascular repair; microvascular dysfunction	Vascular-focused protocol; angiogenesis support; reversal?		Unknown

13.2.1 Interpretation Notes

1. **Ranking reflects current evidence, not ultimate truth.** The “Speculative” hypotheses may prove correct; the “Established” phenomena may be reinterpreted. Science is provisional.
2. **Multiple hypotheses may be simultaneously true.** ME/CFS is almost certainly heterogeneous. Different patients may have different primary drivers, and individual patients may have multiple contributing mechanisms.
3. **“Treatment implications” does not mean “proven treatment.”** We list logical therapeutic consequences of each hypothesis, not demonstrated efficacy. Very few ME/CFS treatments have robust RCT support.
4. **“Potential for rapid benefit” is our subjective assessment** of how quickly patients might improve *if* the hypothesis is correct *and* appropriate treatment is applied. “Unknown” means we cannot predict.
5. **Severely ill patients face different considerations.** Some interventions (immunoadsorption, daratumumab) require hospital access impossible for bedbound patients. Others (pacing, supplements) are accessible. The table does not capture this dimension adequately.
6. **Cross-references to detailed discussions.** Many hypotheses are explored in depth in earlier chapters: immune dysfunction (Chapter 7), neurological abnormalities (Chapter 8), energy metabolism (Chapter 6), cardiovascular findings (Chapter 10), and microbiome alterations (Chapter 11). This chapter synthesizes those findings; consult earlier chapters for mechanistic detail.

13.3 Synthesis: What the Evidence Suggests

Drawing together the ranked hypotheses, several patterns emerge:

13.3.1 The Core Triad: Energy-Immune-Autonomic

Three systems show consistent abnormalities across evidence levels:

1. **Energy metabolism** (mitochondrial dysfunction, ATP depletion, metabolomic abnormalities)—see integrated metabolic model in Section 6.14
2. **Immune function** (cytokine dysregulation, autoantibodies, NK cell dysfunction)—detailed in Chapter 7
3. **Autonomic regulation** (POTS, HRV abnormalities, catecholamine changes)—integrated cardiovascular dysfunction discussed in Section 10.9

The Heng 2025 study [12] suggests these are not independent—the 7-biomarker panel spanning all three systems achieved 91% diagnostic accuracy. This correlation is consistent with coordinated dysfunction. If these systems are functionally coupled, this would have profound implications:

‡ Limitation 1: Diagnostic Correlation ≠ Causal Interdependence

A multi-system biomarker panel achieving high diagnostic accuracy establishes that these systems are co-altered in ME/CFS. It does not establish that they are causally interdependent: the same upstream driver could independently perturb all three systems without any system-to-system causal link. Distinguishing coordinated dysfunction from independent co-occurrence requires intervention studies (e.g., correcting one system and observing whether others normalize).

- Treatments targeting only one system may fail because the others maintain dysfunction
- Patient subgroups may differ in which system predominates, not which system is involved
- A “multi-lock” model (see Chapter 14) may explain treatment resistance

13.3.2 The Autoimmune Subgroup

The daratumumab pilot trial (60% response) [9] provides the strongest evidence yet for an autoimmune mechanism in *a subset* of patients. Key insights:

- Rituximab (anti-CD20, targets B cells) failed in large trials [376]
- Daratumumab (anti-CD38, targets plasma cells) succeeded in pilot [9]
- This suggests **long-lived plasma cells**, not B cells, are the critical autoantibody source
- The 60% response rate implies heterogeneity—not all ME/CFS is autoimmune
- Biomarkers for patient selection are urgently needed
- The ResetME RCT ($n = 66$, placebo-controlled) is now recruiting to test these findings (Appendix I, Section I.4.3)

Observation 69 (The Rituximab Puzzle Solved?). The daratumumab finding [9] may explain one of ME/CFS research’s biggest disappointments. Rituximab showed promise in early trials but failed in the large Norwegian RCT [376]. If the critical autoantibodies come from long-lived plasma cells ($CD38^+$, $CD20^-$), rituximab would deplete the wrong cells. Existing plasma cells would continue producing autoantibodies for months, and by the time B cells returned, no improvement would be evident. The trial “failed” not because autoimmunity isn’t involved, but because the wrong cells were targeted.

13.3.3 The Vascular Dimension

Elevated VWF, fibronectin, and thrombospondin [12] point to **endothelial activation**—the blood vessel lining is chronically stressed. This connects to:

- Long COVID microclot findings (emerging evidence)
- Cerebral hypoperfusion documented in ME/CFS [172]
- Exercise intolerance (endothelium cannot vasodilate properly)
- Multi-system involvement (endothelium is everywhere)

The Hoel et al. (2026) serum proteomics study [191] substantially deepened this picture. Beyond confirming elevated coagulation and complement factors, the study revealed a specific vascular injury signature: reduced vimentin (endothelial structural protein), reduced ANTXR2/CMG2 (capillary morphogenesis regulator), reduced CXADR (shear stress sensor), and elevated endothelial lipase (LIPG). Critically, Ephrin receptor signaling—a pathway governing vascular development and angiogenesis—showed coordinated disruption, with four Ephrin A receptors and their ligands simultaneously altered. The authors also documented elevated WASF3 (a HIF-1 α target), providing direct proteomic evidence for tissue hypoxia. Together, these findings suggest not just endothelial activation but impaired endothelial repair capacity: the vasculature is stressed, and the molecular machinery for repairing and remodeling damaged vessels appears compromised.

If ME/CFS is partly an **endotheliopathy**, vascular-targeted treatments (anticoagulation, fibrinolytics, endothelial support) might help—but this remains preliminary.

13.3.4 The Central Nervous System

The Walitt 2024 finding [5] of altered **effort preference** (not physical fatigue) localizes part of the problem to the brain. Combined with:

- CSF catecholamine deficiency [5]
- Neuroinflammation on PET imaging [73]
- Cognitive dysfunction correlating with perfusion [172]
- Brainstem abnormalities [5]

This suggests ME/CFS involves a **central state change**—the brain is computing effort-reward differently, possibly appropriately given peripheral metabolic dysfunction, but creating the subjective experience of profound unwillingness/inability to exert.

13.3.5 The “Stuck” State

Multiple hypotheses converge on the idea that ME/CFS represents a **stable pathological state** that resists perturbation:

- Epigenetic changes may “lock” gene expression patterns
- Autoantibodies from long-lived plasma cells provide continuous dysfunction
- Metabolic pathway shifts may be self-perpetuating
- The brain’s effort computation may be recalibrated
- Circadian rhythms may be desynchronized

This “multi-lock” concept (detailed in Chapter 14) suggests why:

- Single interventions rarely produce cures
- Early treatment may prevent lock stabilization
- Disease duration correlates with prognosis

- Some patients spontaneously recover (locks didn't fully stabilize)
- Treatment may need to target multiple locks simultaneously

Observation 70 (The Multi-Lock Model and Treatment Implications). If ME/CFS involves multiple self-reinforcing abnormalities, this has profound implications for clinical trials. A treatment targeting one mechanism (e.g., immunoadsorption removing autoantibodies) might show modest benefit if other locks (epigenetic, metabolic, autonomic) maintain dysfunction. This could explain the disappointing results of many single-mechanism trials. Future research should explore: (1) sequential combination therapies (break locks one at a time), (2) simultaneous multi-targeted protocols (address all locks together), or (3) biomarker-guided sequencing (identify which lock predominates in each patient). The daratumumab 60% response rate [9] may reflect successful targeting in patients where autoimmunity is the primary lock, while non-responders have different dominant mechanisms.

Speculation 21 (Sickness Behavior “Stuck On” Hypothesis). Sickness behavior is an evolutionarily conserved motivational state triggered by pro-inflammatory cytokines (IL-1 β , IL-6, TNF- α) acting on the brain through both humoral and neural (vagal afferent) pathways [377, 378]. Its hallmarks—fatigue, social withdrawal, anorexia, hyperalgesia, cognitive impairment, and somnolence—precisely parallel the core symptom constellation of ME/CFS [379]. We speculate that ME/CFS represents a state in which the sickness behavior program, normally self-limiting, becomes chronically activated or fails to disengage.

Adaptive Function of Sickness Behavior. In acute infection, sickness behavior redirects energy from locomotion, foraging, and social interaction toward immune function. This re-allocation is metabolically efficient: fever alone increases metabolic rate by approximately 13% per degree Celsius, and immune activation consumes substantial glucose and amino acids [380]. The “cost” of sickness behavior (reduced activity) is offset by the “benefit” of enhanced pathogen clearance. Crucially, sickness behavior involves active CNS reprogramming of motivation and effort perception, not merely peripheral weakness [381].

Mechanisms of Persistence. Several mechanisms could prevent sickness behavior from resolving:

Chronic low-grade immune activation. Even after the initial infection resolves, persistent immune activation (from viral reservoirs, autoantibodies, dysbiosis-driven LPS translocation, or mast cell activation) may continuously provide the cytokine signals that maintain sickness behavior. This does not require high-level inflammation—even subtle elevations in IL-1 β and TNF- α are sufficient to trigger sickness behavior circuits [377].

Vagal afferent sensitization. The vagus nerve transmits peripheral inflammatory signals to the nucleus tractus solitarius and subsequently to the hypothalamus. Repeated activation may sensitize this pathway, such that progressively lower levels of peripheral inflammation trigger full sickness behavior responses [382]. This could explain why ME/CFS patients experience severe symptoms despite only modest peripheral inflammatory markers.

Hypothalamic set-point shift. The hypothalamus integrates immune, metabolic, and autonomic signals to determine the sickness behavior “set point.” Chronic activation may shift this set

point, requiring active intervention (rather than mere absence of infection) to return to baseline. This parallels allostatic load theory: the regulatory system itself becomes dysregulated.

Cytokine-epigenetic feedback. Pro-inflammatory cytokines induce epigenetic changes (DNA methylation, histone modification) in hypothalamic neurons and microglia [383]. These epigenetic modifications may persist after the cytokine signal diminishes, creating a self-maintaining state where the sickness behavior program remains “written into” neural gene expression.

Testable Predictions.

1. **Anti-cytokine response:** IL-1 receptor antagonist (anakinra) should reduce sickness behavior symptoms in ME/CFS. Preliminary evidence is mixed: Roerink et al. found no significant benefit from anakinra in ME/CFS in an RCT [384], though the 4-week treatment duration may have been insufficient to reverse established set-point shifts.
2. **Vagal modulation:** Non-invasive vagal nerve stimulation should modulate sickness behavior intensity, potentially providing rapid (hours to days) symptomatic relief.
3. **Motivational dissociation:** If ME/CFS is “stuck” sickness behavior, patients should show altered neural responses to reward and effort in fMRI, specifically resembling experimentally induced sickness (e.g., LPS challenge studies) rather than depression or deconditioning.
4. **Cytokine sensitivity:** ME/CFS patients should show amplified sickness behavior responses to standardized immune challenges (e.g., typhoid vaccination) compared to healthy controls, reflecting sensitized sickness circuits.

Treatment Implications.

- **Desensitization approaches:** Graduated immune challenge protocols (conceptually analogous to allergy desensitization) might recalibrate sensitized sickness circuits.
- **Epigenetic interventions:** If epigenetic modifications maintain the stuck state, histone deacetylase inhibitors or other epigenetic modulators might help reverse the programming.
- **Reframing:** Understanding ME/CFS as “stuck sickness behavior” reframes symptoms as adaptive responses in an inappropriate context, potentially reducing stigma and guiding mechanistic research.

⚠ Limitation 2: Sickness Behavior Model: Explanatory Gaps

The sickness behavior model does not explain all ME/CFS features. PEM—the hallmark worsening after exertion—is not a recognized feature of acute sickness behavior. The negative anakinra trial [384] suggests that simply blocking one cytokine pathway is insufficient, though this does not refute the broader hypothesis. The model also struggles to explain why some patients develop ME/CFS after non-infectious triggers (physical trauma, surgery) where the initial sickness behavior program may not have

been engaged.

Certainty: 0.40 (symptom overlap is striking; mechanistic persistence pathway speculative; negative anakinra trial complicates picture)

Speculation 22 (Partial Torpor Trap Hypothesis). Torpor is a phylogenetically conserved state of controlled metabolic suppression in which organisms dramatically reduce body temperature, heart rate, and metabolic rate to survive periods of energy scarcity [385, 386]. We speculate that ME/CFS involves activation of torpor-related metabolic suppression pathways without the coordinated physiological program that enables safe entry into and arousal from torpor—a “partial torpor trap.”

Torpor Biology. Recent research has identified specific neural circuits controlling torpor in mice:

- **QRFP neurons:** Hypothalamic neurons expressing pyroglutamylated RFamide peptide (QRFP) are both necessary and sufficient to induce torpor-like states in mice [386]. Chemogenetic activation of these neurons reduces body temperature by 5–10°C and metabolic rate by 40–70%.
- **Preoptic area circuits:** Genetically distinct neuronal populations in the preoptic area drive distinct features of torpor (temperature reduction, metabolic suppression, behavioral quiescence) [385].
- **Coordinated entry and arousal:** Normal torpor involves coordinated engagement of thermoregulatory, metabolic, and cardiovascular systems, with active arousal mechanisms ensuring safe exit.

ME/CFS as Partial Torpor. Several ME/CFS features resemble incomplete torpor engagement:

Metabolic suppression without temperature reduction. ME/CFS patients show reduced metabolic rate and metabolic inflexibility but generally maintain normal core body temperature (though some report subjective coldness and temperature dysregulation). This pattern is consistent with activation of metabolic suppression pathways without the thermoregulatory component—as if only part of the torpor program has engaged.

Cardiovascular changes. Torpor involves reduced heart rate and cardiac output. ME/CFS patients show reduced cardiac output, blunted heart rate responses (chronotropic incompetence), and orthostatic intolerance—changes directionally consistent with partial torpor cardiovascular adjustment.

Behavioral quiescence with preserved awareness. In torpor, animals become behaviorally quiescent. ME/CFS patients show profound reduction in physical activity while maintaining cognitive awareness (albeit impaired)—consistent with dissociation between behavioral and consciousness components of the torpor program.

Arousal failure. Normal torpor includes coordinated arousal involving UCP1-mediated thermogenesis, sympathetic activation, and TRPM2-mediated calcium signaling. If ME/CFS

involves partial torpor, the arousal program may be incomplete or repeatedly failing, trapping the organism in a low-metabolic state.

Testable Predictions.

1. **QRFP pathway markers:** ME/CFS patients should show altered QRFP signaling (measurable in CSF) compared to healthy controls and fatigued-but-not-ME/CFS patients.
2. **Arousal marker deficiency:** Markers of torpor arousal (UCP1 expression in brown adipose tissue, sympathetic activation patterns, TRPM2 channel activity) should be reduced or dysregulated in ME/CFS.
3. **Brown adipose tissue:** ME/CFS patients may show altered brown adipose tissue activity (measurable by FDG-PET cold stimulation), reflecting impaired thermogenic arousal.
4. **Dauer analogy:** In *C. elegans*, the dauer state (metabolic arrest under adverse conditions) is triggered by specific signaling pathways. If ME/CFS involves an analogous “dauer-like” state in humans, metabolic profiling should reveal pathway-specific suppression patterns distinct from simple caloric restriction.

Treatment Implications. If ME/CFS involves partial torpor, treatment should focus on completing the arousal program:

- **Thyroid optimization:** Thyroid hormones are critical torpor arousal signals. Even “normal range” thyroid function might be insufficient if arousal pathways are suppressed.
- **Brown adipose stimulation:** Cold exposure protocols or pharmacological BAT activation might engage arousal circuits.
- **Sympathomimetics:** Carefully titrated sympathomimetic agents might provide the sympathetic activation component missing from incomplete arousal.

Limitations. This hypothesis is highly speculative. Humans do not normally enter torpor, and it is unclear whether human hypothalamic circuits retain functional torpor-induction capacity. The analogy between ME/CFS and torpor is based on phenomenological similarity rather than demonstrated mechanistic overlap. Core body temperature is generally normal in ME/CFS, which argues against full torpor pathway engagement. No human studies have examined QRFP or related torpor-control pathways in ME/CFS.

Certainty: 0.30 (intriguing conceptual framework; minimal direct evidence; highly speculative)

13.4 Proposed Unifying Mechanisms

13.4.1 Vicious Cycle Models

Several vicious cycles may perpetuate ME/CFS. These cycles are identified and discussed in detail within their respective system chapters: immune vicious cycles in Chapter 7, HPA-

immune feedback in Chapter 9, MCAS-POTS interactions in Chapter 10, and gut-brain bidirectional dysfunction in Chapter 11. Here we synthesize these chapter-specific cycles into a comprehensive framework:

Inflammation-Metabolism Cycle.

1. Inflammation activates IDO, shunting tryptophan toward kynurenine [231]
2. Kynurenine pathway produces neurotoxic quinolinic acid [232]
3. Neuroinflammation maintains cytokine production [73]
4. Cytokines perpetuate IDO activation

Energy-Immune Cycle.

1. Mitochondrial dysfunction depletes ATP [12]
2. Immune cells cannot complete activation/maturation (ATP-dependent)
3. Dysfunctional immune response fails to clear triggers
4. Persistent triggers maintain inflammation
5. Inflammation impairs mitochondria [64]

Autonomic-Vascular Cycle.

1. Autonomic dysfunction impairs vascular regulation
2. Poor perfusion causes tissue hypoxia
3. Hypoxia triggers HIF pathway and metabolic shifts
4. Metabolic abnormalities affect autonomic centers

Gut-Serotonin-Vagal-Autonomic Cycle (Hypothesized). This proposed cycle chains five observations—the first three individually documented, the last two inferred:

1. Gut dysbiosis in ME/CFS reduces butyrate-producing bacteria [349] (*documented*)
2. Butyrate enhances enterochromaffin cell serotonin synthesis [14]; butyrate deficiency would therefore be expected to reduce it (*documented mechanism, inferred consequence in ME/CFS*)
3. Enterochromaffin serotonin activates vagal afferents via 5-HT₃ receptors [13, 228] (*documented*)
4. Reduced vagal afferent input may diminish efferent vagal tone via brainstem integration, impairing autonomic output to heart, gut, and immune system (*inferred; not directly demonstrated in ME/CFS*)
5. Impaired vagal efferent output would worsen gut motility, potentially promoting further dysbiosis—closing the cycle (*inferred*)

Wirth and Scheibenbogen's broader framework of neurotransmitter dysregulation [225] provides theoretical context for steps 2–4, though the specific cyclic pathway proposed here extends beyond their analysis. This hypothesized cycle links the Autonomic-Vascular Cycle above to gut pathophysiology (Chapter 11, Section 11.1.4). Validation would require measuring enterochromaffin serotonin output and vagal afferent activity in ME/CFS patients alongside butyrate levels.

Exertion-Crash Cycle.

1. Patient feels slightly better, increases activity
2. Activity exceeds metabolic capacity
3. Post-exertional crash (24–72 hours delayed)
4. Crash worsens baseline, triggers immune/metabolic responses
5. Partial recovery, patient attempts activity again

Breaking these cycles is the goal of effective treatment—but which cycle to break, and how, likely differs between patients.

Inflammation-Barrier Bidirectional Cycle. A self-sustaining pathophysiological cycle may become entrenched in severe ME/CFS patients, in which systemic inflammation and intestinal barrier dysfunction perpetually reinforce each other:

1. **Baseline systemic inflammation:** ME/CFS patients exhibit elevated pro-inflammatory cytokines (IL-1 β , IL-6, TNF- α , IFN- γ) [213] that correlate with disease severity.
2. **Tight junction disruption:** These cytokines directly suppress expression of tight junction proteins (claudin-1, occludin, zonula occludens-1), disrupting the intestinal epithelial seal [315].
3. **LPS translocation:** Permeability increases, allowing bacterial lipopolysaccharide (LPS) to cross the intestinal epithelium and enter systemic circulation [315].
4. **TLR4 activation and cytokine amplification:** Translocated LPS activates toll-like receptor 4 (TLR4) on innate immune cells, triggering production of additional pro-inflammatory cytokines (IL-1 β , IL-6, TNF- α) [387].
5. **Barrier repair impairment:** The elevated cytokines that trigger barrier disruption simultaneously suppress anti-inflammatory mechanisms and stress the repair capacity of already compromised enterocytes. Additionally, low cortisol in severe patients (discussed in Chapter 9) reduces expression of tight junction proteins [388].
6. **Cycle amplification:** With each cycle, baseline permeability increases, baseline cytokines increase, and the barrier's repair capacity becomes increasingly exhausted. This self-sustaining loop may explain why disease severity tends to increase over time in some severe patients without intervention.

~ Hypothesis 1: Breaking the Inflammation-Barrier Cycle in Severe Patients

Ceasing the perpetuation of this cycle may require simultaneous multi-target interventions rather than single-pathway strategies. Possible approaches include: (1) **Barrier restoration** via glutamine supplementation, collagen peptides, or other enterocyte-supportive nutrients; (2) **Wheat elimination** to remove zonulin-mediated barrier stress; (3) **Anti-inflammatory support** via omega-3 fatty acids, curcumin, or other dietary polyphenols; (4) **Mast cell stabilization** with quercetin or ketotifen to reduce histamine-mediated inflammatory mediator release; and (5) **Cortisol optimization** to support endogenous barrier repair. Addressing only one mechanism (e.g., wheat elimination without anti-inflammatory support or barrier repair substrates) may prove insufficient if the cycle has become deeply entrenched, requiring comprehensive multi-pathway interventions for clinical efficacy in severe patients.

Certainty: 0.75 (mechanistic logic strong; direct evidence for this integrated multi-target approach lacking in ME/CFS).

★ Key Point: Orthostatic Intolerance as Potential Upstream Driver

Pediatric ME/CFS data suggest that orthostatic intolerance (OI) may function as an upstream driver—a lynchpin whose early correction can prevent cascade into multi-system dysfunction.

In pediatric ME/CFS treatment studies, aggressive OI management produces improvements not only in cardiovascular symptoms but also in fatigue, cognitive function, and overall wellbeing [78]. This broad benefit pattern suggests that OI is not merely one symptom among many, but rather a primary pathophysiological driver whose downstream effects include immune activation, neuroinflammation, and metabolic dysfunction.

Mechanistic rationale:

Chronic cerebral hypoperfusion from untreated OI creates a cascade: reduced brain oxygen delivery triggers compensatory metabolic shifts, impairs neurotransmitter synthesis, activates microglia (neuroinflammation), and disrupts autonomic regulatory centers. The resulting autonomic dysfunction further worsens perfusion, completing a vicious cycle. Additionally, systemic hypoperfusion during orthostatic challenge causes tissue hypoxia, oxidative stress, and immune activation throughout the body.

If OI is the initiating driver, then correcting it early—before secondary systems become dysregulated—could prevent the recruitment of additional vicious cycles (energy-immune, inflammation-metabolism) that characterize established ME/CFS.

Clinical implications:

This framework suggests that *early, aggressive OI treatment* may be disease-modifying rather than merely symptomatic:

- In pediatric patients, OI correction may prevent progression to multi-system ME/CFS
- In early-stage adult ME/CFS (under 2 years duration), aggressive OI treatment could interrupt cascade before lock-in
- In established ME/CFS, OI treatment remains important but may be insufficient

alone—additional cycles have been recruited

- Treatment aggressiveness should match disease stage: maximal in early disease when prevention is possible

Caveats:

This interpretation remains speculative. Alternative explanations exist: OI treatment's broad benefits could reflect improved perfusion supporting all systems rather than preventing cascade, or pediatric OI responsiveness could reflect developmental plasticity enabling recovery through multiple pathways simultaneously. Additionally, not all ME/CFS patients have prominent OI, suggesting heterogeneity in primary drivers. The hypothesis applies most strongly to the OI-predominant subgroup, particularly in early disease stages.

See Section 5.6.11 for the Septad framework that positions OI as one of seven interconnected pathophysiological domains, and Chapter 10 for detailed discussion of OI mechanisms and treatments.

Speculation 23 (Recovery Capital Model). We propose a conceptual framework of “Recovery Capital”—the cumulative biological capacity for recovery that is consumed by severe post-exertional malaise episodes and regenerated over time. In this model, children possess high initial Recovery Capital (developmental plasticity, immune renewal, metabolic flexibility) and regenerate it rapidly, while adults start with lower capital and regenerate slowly if at all. Each severe crash “spends” Recovery Capital through epigenetic changes, accumulated cellular damage, and immune exhaustion. Once Recovery Capital is depleted below a threshold, recovery becomes unlikely. This framework explains why strict pacing (capital preservation) and early intervention (maximizing capital before depletion) are particularly critical in pediatric patients, and why aggressive early treatment in adult patients may preserve recovery potential that would otherwise be lost.

Speculation 24 (Hematopoietic Stem Cell Exhaustion Model). **Certainty: 0.30.** ME/CFS may involve accelerated exhaustion of hematopoietic stem cells (HSCs), with the pediatric recovery advantage reflecting children's larger HSC reserves. This speculation extends the Recovery Capital framework by identifying HSC function as a critical, quantifiable component of biological reserve. The certainty level reflects: (1) highly speculative extrapolation from aging biology to ME/CFS; (2) indirect peripheral blood markers as proxy for bone marrow HSC status; (3) absence of direct bone marrow biopsy studies in ME/CFS populations; (4) lack of direct intervention trials targeting HSC function.

We propose that ME/CFS involves accelerated exhaustion of hematopoietic stem cells (HSCs), and that the pediatric recovery advantage reflects children's larger HSC reserves and greater regenerative capacity.

Conceptual foundation:

Hematopoietic stem cells reside in bone marrow niches and give rise to all blood and immune cells throughout life. HSC function declines with age through multiple mechanisms: telomere shortening limits replicative capacity, accumulation of DNA damage triggers senescence,

epigenetic drift alters differentiation potential, and clonal selection reduces diversity. This age-related decline is well-characterized and contributes to immunosenescence—the progressive deterioration of immune function with aging [389, 390].

Recent 2024–2025 research demonstrates that inflammation is a driving force of HSC aging, causing irreversible exhaustion of functional HSCs [391]. Critically, HSCs can be induced to proliferate and differentiate in response to stress signals during infection, inflammation, chemotherapy, radiation, and aging [392]. However, with chronic or repeated stimulation, HSCs show loss of function and exhaustion [392]. Transient LPS exposure primes aged HSCs to undergo accelerated differentiation at the expense of self-renewal, leading to depletion of HSCs, with the central regulator NF- κ B mediating functional impairment by inflammation insult [391].

We hypothesize that ME/CFS triggers and perpetuates accelerated HSC exhaustion through mechanisms that may be reversible if addressed early but become permanent once thresholds are crossed.

Proposed mechanism:

Initial insult. The triggering event (typically infection) produces massive immune activation requiring rapid expansion of effector cells. This expansion draws heavily on HSC reserves, as progenitor cells must proliferate to replace the mature cells consumed in the immune response. A severe or prolonged initial infection could substantially deplete HSC reserves through this demand-driven exhaustion.

Post-exertional amplification. Each crash episode may trigger additional waves of immune activation, cytokine release, and oxidative stress—all of which place demands on HSCs. Unlike healthy individuals who have HSC reserves to accommodate occasional stressors, ME/CFS patients operating with depleted reserves experience cumulative damage with each crash. This creates a vicious cycle: crashes deplete HSCs, reduced HSC function impairs recovery, incomplete recovery leads to more crashes.

Inflammatory damage to the niche. Chronic inflammation may damage the bone marrow microenvironment (the “niche”) that supports HSC function. Inflammatory cytokines alter niche cell function, disrupt the signals that maintain HSC quiescence, and may directly damage HSCs through oxidative stress. This niche damage could persist even if systemic inflammation resolves, leaving HSCs unable to function normally.

Clonal restriction. As HSC diversity declines, the remaining clones may be less capable of generating the full spectrum of immune cells needed for healthy function. Clonal hematopoiesis of indeterminate potential (CHIP)—dominance of blood production by a small number of HSC clones—is associated with increased inflammation, cardiovascular disease, and mortality in aging populations. ME/CFS may accelerate this clonal restriction.

The pediatric advantage:

Children possess several HSC-related advantages that could explain their superior recovery rates:

Larger initial reserves. Children have more HSCs per unit of bone marrow and a higher proportion of functionally competent, long-term repopulating HSCs. They can sustain greater HSC consumption before crossing critical thresholds.

Active bone marrow. Pediatric bone marrow is highly cellular (red marrow), while adult marrow progressively converts to fatty (yellow) marrow with reduced hematopoietic capacity. The active pediatric marrow can regenerate HSC populations more effectively.

Greater regenerative capacity. Pediatric HSCs have longer telomeres, less accumulated DNA damage, and more robust self-renewal capacity. After an insult, they can recover function more completely.

More plastic niche. The pediatric bone marrow microenvironment is more plastic and may be able to repair inflammatory damage that would be permanent in adults.

Connection to other hypotheses:

HSC exhaustion integrates with other proposed ME/CFS mechanisms:

Immune dysfunction. Many immune abnormalities in ME/CFS—reduced NK cell function, T cell exhaustion, altered cytokine profiles—could stem from inability to regenerate healthy immune cells due to HSC exhaustion.

Epigenetic aging. Epigenetic clocks measure biological age partly through methylation patterns established during hematopoiesis. Accelerated epigenetic aging in ME/CFS could reflect HSC exhaustion and altered differentiation.

Autoimmunity. HSC exhaustion could impair tolerance mechanisms that depend on continuous generation of naive, properly selected lymphocytes, potentially contributing to autoantibody persistence.

Recovery Capital. HSC reserve is a concrete, measurable component of Recovery Capital. Patients with preserved HSC function retain capacity for immune regeneration; those with exhausted HSCs do not.

Biomarker development:

If HSC exhaustion contributes to ME/CFS, several biomarkers could be developed:

- **Circulating progenitors:** CD34⁺ cell counts in peripheral blood as a proxy for bone marrow output
- **Clonal diversity:** TCR/BCR repertoire diversity as an indirect measure of HSC diversity; reduced diversity suggests clonal restriction
- **CHIP mutations:** Screening for clonal hematopoiesis mutations (DNMT3A, TET2, ASXL1) that indicate oligoclonal dominance
- **Telomere length:** Particularly in HSC-enriched populations or as a predictor of replicative capacity
- **Single-cell HSC profiling:** Advanced approaches (single-cell RNA-seq of bone marrow aspirates) could directly characterize HSC populations

Treatment implications:

If HSC exhaustion is a key mechanism, treatments could aim to:

Preserve remaining HSCs. Strict pacing, crash prevention, and anti-inflammatory therapy would minimize ongoing HSC consumption. This provides additional rationale for the “preservation” arm of ME/CFS management.

Support HSC regeneration. Fasting-mimicking diets have been shown to promote HSC regeneration in animal models and may be beneficial in ME/CFS. Growth factors (G-CSF, EPO) could be explored, though with caution given their complexity.

Niche repair. Therapies targeting the bone marrow microenvironment could potentially restore HSC function even when HSCs themselves are viable but quiescent due to niche dysfunction.

HSC supplementation (speculative). In severe cases with confirmed HSC exhaustion, autologous HSC boost (collection during a good period, expansion ex vivo, reinfusion) could theoretically replenish reserves. This would require extensive development and carries significant risks.

Limitations:

This model is highly speculative. Direct evidence for HSC exhaustion in ME/CFS is limited; most evidence is indirect, based on peripheral blood markers and reasoning from aging biology. Bone marrow studies in ME/CFS are rare due to the invasiveness of biopsy. The model does not explain why some patients with long disease duration do eventually recover, or why some young patients do not recover. Additionally, HSC exhaustion could be a consequence rather than a cause of ME/CFS—a downstream effect of other primary mechanisms.

13.4.2 Multisystem Failure Cascade

A proposed sequence for ME/CFS development:

Phase 1: Triggering Event.

- Infection (EBV, enteroviruses, SARS-CoV-2, others)
- Severe stress (physical, psychological, surgical)
- Combination of factors in vulnerable individual

Phase 2: Acute Response.

- Normal sickness behavior program activates
- Metabolic suppression, immune activation, behavioral changes
- This is *adaptive*—conserving resources for recovery

Phase 3: Failed Resolution.

- In most people, acute phase resolves in days to weeks
- In ME/CFS-susceptible individuals, resolution fails
- Possible reasons: genetic susceptibility, severity of insult, timing, comorbidities

Phase 4: Lock Establishment.

- Autoantibodies generated and plasma cells established
- Epigenetic changes stabilize “sick” gene expression
- Metabolic pathways shift to new equilibrium
- Brain recalibrates effort computation
- Autonomic setpoints shift

Phase 5: Stable Pathological State.

- Multiple locks reinforce each other
- Perturbations (exertion, stress, infection) trigger defensive responses
- Spontaneous recovery becomes unlikely
- Treatment must address multiple locks

13.4.3 Temporal Dynamics of Cycle Recruitment

The multisystem failure cascade above describes discrete phases, but understanding *what triggers transitions between phases*—particularly the recruitment of additional vicious cycles—is critical for prevention strategies. The cycle dynamics framework (Chapter 2, §2.1) identifies specific triggers that may accelerate progression from single-cycle to multi-cycle disease.

Proposed Recruitment Sequence**~ Hypothesis 2: Sequential Cycle Recruitment Model**

ME/CFS typically begins with one primary vicious cycle (usually mitochondrial) and progressively recruits additional cycles over time:

Stage 1 (0–6 months): Mitochondrial cycle only

↓

Stage 2 (6–18 months): Mitochondrial + Immune (triggered by sustained ROS signaling)

↓

Stage 3 (12–36 months): + Autonomic (triggered by chronic immune activation crossing BBB)

↓

Stage 4 (>2 years): + Neuroinflammatory + Endocrine (central sensitization)

↓

Stage 5 (>5 years): Full cycle engagement with epigenetic lock-in

Evidence Grade: D (hypothesized based on clinical progression patterns and mechanistic logic; not empirically validated as universal sequence)

Recruitment Triggers

Table 13.3: Hypothesized Triggers for Cycle Recruitment

Trigger	Proposed Mechanism	Target Cycle	Evidence
Severe crashes (Grade 4–5)	Massive ROS release triggers inflammatory cascade; exceeds repair capacity	Immune, neuroinflammatory	D
Secondary infections	Reactivate immune system; overwhelm already-depleted reserves	Immune	C
Cumulative damage threshold	Gradual mtDNA mutations reach critical mass	Mitochondrial (amplification)	C
Chronic hypoperfusion	Sustained autonomic dysfunction impairs BBB, enables CNS penetration	Neuroinflammatory	D
Psychosocial stress	HPA axis activation recruits endocrine dysfunction	Endocrine	C

★ Key Point: Prevention Implications of Cycle Recruitment

If severe crashes are the primary trigger for cycle recruitment, then **aggressive pacing from diagnosis** may prevent or delay progression from single-cycle to multi-cycle disease:

Testable prediction: Patients adhering strictly to energy envelope pacing show slower cycle recruitment over 2 years (hazard ratio <0.5 for each additional cycle activation) compared to those with frequent crashes.

Clinical implication: Pediatric ME/CFS studies report 54–94% improvement or recovery rates [393], while adult ME/CFS shows median recovery of only 5% (range 0–31%) in systematic review [54]. If this difference reflects disease stage rather than age *per se*, the high early-recovery rates may result from aggressive rest preventing cycle recruitment beyond Stage 1–2. The low adult recovery rate in established disease would then reflect multi-cycle involvement where spontaneous resolution becomes increasingly improbable with each additional engaged cycle—a prediction of the model requiring prospective validation, not an empirical observation.

This framework transforms pacing from “symptom management” to **disease-modifying therapy**—not merely reducing current symptoms but preventing irreversible progression.

Sentinel Biomarkers for Cycle Activation

Early detection of cycle recruitment could enable preemptive intervention:

- **Immune cycle sentinel:** Rising IL-6, TNF- α , or emergence of autoantibodies before clinical immune symptoms manifest
- **Autonomic cycle sentinel:** Declining HRV, increasing resting heart rate, or emerging orthostatic intolerance
- **Neuroinflammatory sentinel:** Rising substance P, emerging sensory sensitivities, or new cognitive symptoms
- **Endocrine sentinel:** Blunted cortisol awakening response, emerging temperature dysregulation

Regular monitoring for these sentinel biomarkers in early-stage patients could trigger preemptive intervention before full cycle activation.

Research Priority: Inception Cohort Study

Validating the cycle recruitment model requires a prospective inception cohort:

- **Population:** New-onset ME/CFS (<6 months), confirmed Stage 1–2 status
- **Follow-up:** 5 years with quarterly cycle mapping (Years 1–2), semi-annual (Years 3–5)
- **Intervention sub-study:** Randomize to intensive pacing support vs. standard care; compare cycle recruitment rates
- **Sample size:** $n = 130$ (65 per arm) for 80% power to detect HR = 0.5 for cycle recruitment
- **Primary endpoint:** Time to first additional cycle activation (Stage 1–2 → Stage 3+)

Such a study would provide the first empirical test of whether crash prevention truly delays disease progression.

13.4.4 Orthostatic Intolerance as Potential Upstream Driver

★ Key Point: OI as Mechanistic Lynchpin in Early ME/CFS

Certainty: 0.55. Orthostatic intolerance may function as an upstream mechanistic driver in early ME/CFS, triggering a cascade of secondary dysfunction across immune, metabolic, and neuroimmune systems. This model is grounded in pediatric clinical observations and splanchnic perfusion physiology, though high-quality trials testing the causal direction remain limited.

Clinical Foundation: Pediatric Treatment Response Pediatric ME/CFS specialists consistently report that aggressive orthostatic intolerance (OI) treatment produces improvements extending far beyond cardiovascular symptoms—including fatigue duration and severity, cognitive function, post-exertional malaise patterns, and general well-being [393]. Rowe and colleagues document recovery or substantial improvement in 40–70% of pediatric ME/CFS patients with aggressive tilt table-guided fludrocortisone, midodrine, and compression therapy, even in patients with severe initial disability. Criti-

cally, symptom improvements precede increases in exercise tolerance by weeks to months, suggesting causality rather than simple symptom clustering.

Proposed Mechanism: Splanchnic Perfusion as Integrative Hub During orthostatic stress, splanchnic blood flow declines by 50–80% to maintain cerebral and coronary perfusion. In dysautonomic patients, this adaptive response becomes exaggerated, creating chronic splanchnic hypoperfusion that triggers three downstream cascades:

Gut barrier failure and endotoxemia. Splanchnic ischemia impairs intestinal epithelial function, increasing zonulin-mediated permeability and allowing lipopolysaccharide (LPS) translocation. Chronic LPS exposure triggers mast cell activation, systemic inflammation, and adaptive immune responses (Section 7.9.7).

Metabolic dysregulation. The small intestine normally produces 40% of the body's glutamine, a critical fuel for rapidly dividing cells. Splanchnic hypoperfusion reduces intestinal glutamine synthesis, depleting substrates for immune cell proliferation and mitochondrial recovery. This may explain the selective energy dysfunction pattern in ME/CFS.

Neuroimmune amplification. Chronic endotoxemia and sleep disruption (consequent to autonomic instability) activate neuroinflammation through microglia and astrocytes. Over weeks to months, epigenetic changes (histone modifications, DNA methylation) lock cells into pro-inflammatory phenotypes (Section 7.9.6), and autoantibodies against autonomous nerve receptors establish, perpetuating dysfunction even if OI is later corrected.

Integration with Septad Framework Within the vicious cycle septad (Section 5.6.11), OI occupies a distinctive position: it is simultaneously a primary symptom (orthostatic intolerance) and a mechanistic hub connecting infection/immune activation → autonomic dysfunction → splanchnic ischemia → gut barrier failure, endotoxemia, and secondary metabolic and immune dysfunction. Early OI treatment interrupts the cycle before epigenetic lock-in and autoimmune establishment, explaining the higher efficacy in recently-onset disease.

Testable Predictions This framework generates specific, falsifiable predictions:

1. **Age-stratified efficacy.** Aggressive early OI treatment should produce higher remission rates in patients with disease onset <6 months versus >2 years, due to prevention of epigenetic reprogramming and autoantibody establishment.
2. **Barrier marker response.** Patients responding to OI treatment should show declining serum zonulin, I-FABP, and LPS-binding antibodies; non-responders should show persistent elevation despite OI correction.
3. **Symptom sequence.** Orthostatic symptoms should improve first (weeks), followed by fatigue (4–12 weeks), then cognition and immune symptoms (8–16 weeks). Reversal of this sequence (cognition improving before orthostasis) would argue against OI as primary driver.

Adult Implications and Caveats In adults with long-standing ME/CFS (>5 years), aggressive OI treatment shows lower response rates [394], consistent with the hypothesis that secondary dysfunctions have become epigenetically entrenched. However, recent-onset adult ME/CFS (<1 year, high OI severity) may show responses comparable to pediatric patients, suggesting disease duration rather than age per se is the critical variable. This predicts that OI-focused treatment in recently-diagnosed adults with prominent autonomic features should receive high priority.

Clinical Rationale: Front-Loading OI Treatment This model provides mechanistic justification for front-loading orthostatic interventions (Section 18.5.1) early in disease course, even when cardiovascular symptoms seem modest. Delaying OI treatment in hope of spontaneous resolution risks epigenetic reprogramming and autoimmunity establishment, narrowing the window for cascade interruption.

Limitations The proposed mechanism remains partially speculative: (1) splanchnic perfusion has not been directly measured in ME/CFS populations; (2) OI-related barrier failure and endotoxemia are inferred from indirect markers; (3) the temporal sequence of epigenetic changes remains unclear; (4) approximately 30–40% of ME/CFS patients lack significant OI, suggesting alternative primary drivers for OI-negative subgroups; (5) controlled trials of aggressive early OI treatment with non-cardiovascular endpoints are lacking.

Research Priorities Resolving this question requires: (1) comparative longitudinal trials of aggressive versus standard OI treatment in early-onset ME/CFS with non-cardiovascular endpoints; (2) measurement of splanchnic perfusion (Doppler ultrasound) and gut barrier markers (zonulin, I-FABP) in OI-positive and OI-negative subgroups; (3) epigenetic profiling of microglial and splanchnic endothelial cells in responders versus non-responders to OI treatment; (4) studies of the early-intervention trial concept described in Chapter 33.

13.4.5 Selective Energy Dysfunction Framework

The selective energy dysfunction hypothesis (Section 14.24) proposes that ME/CFS preferentially affects CNS-dependent, demand-responsive processes while sparing autonomous local processes. This framework integrates with and clarifies several aspects of the vicious cycle models above.

★ Key Point: Integration of Selective Dysfunction with Unifying Models

Clarifies the vicious cycle targets: The energy-immune cycle and autonomic-vascular cycle both operate through CNS coordination. If CNS energy is the primary bottleneck, all cycles dependent on CNS signaling become vulnerable simultaneously—explaining why ME/CFS affects multiple systems.

Explains preserved functions: Hair growth, nail growth, and basic wound healing continue because they operate via local autonomous regulation ($\delta_{CNS} < 0.2$) outside the affected coordination pathways.

Reframes the “stuck state”: The multi-lock model proposes multiple independent locks. Selective dysfunction suggests these locks may be downstream manifestations of a single upstream CNS energy bottleneck. If the brain cannot generate coordination signals, all CNS-dependent systems fail regardless of their local machinery’s integrity.

Explains pharmacological bypass: The effectiveness of direct-acting agents (midodrine, pyridostigmine) that bypass CNS coordination supports the selective dysfunction model—peripheral end-organs are functional; only the coordination signal is missing.

Evidence Synthesis Across Systems. The selective energy dysfunction framework is supported by consistent evidence across multiple physiological systems, each documenting preserved baseline function with impaired demand-responsive capacity:

- **Energy Metabolism (Chapter 6, Section 6.3):** CNS-dependent and demand-responsive processes show selective impairment while autonomous steady-state peripheral functions (hair growth, nail growth, wound healing) continue at apparently normal rates despite severe systemic symptoms. This pattern distinguishes selective coordination failure from global mitochondrial dysfunction.
- **Neurological System (Chapter 8, Section 8.1.2):** Near-universal cognitive dysfunction, documented brain hypometabolism, neuroinflammation (45–199% elevation across key regions), and catecholamine deficiency suggest the brain serves as the primary energy bottleneck. The brain’s disproportionate energy demand (20–25% of total energy while comprising only 2% of body mass) makes it uniquely vulnerable to energy constraint, with downstream failures in autonomic coordination affecting all CNS-dependent systems.
- **Cardiovascular System (Chapter 10, Section 10.2.4):** Cerebral blood flow abnormalities exemplify the selective dysfunction pattern: 91% of ME/CFS patients with normal resting heart rate and blood pressure show abnormal CBF reduction during orthostatic challenge, with reduction magnitude 3.7-fold greater than controls (26% vs. 7%). CBF remains reduced even after returning to supine position, correlating with disease severity rather than hemodynamic parameters, indicating intrinsic cerebrovascular or metabolic dysfunction.

Collectively, these findings establish a coherent mechanistic framework where CNS energy failure drives the selective pattern: autonomous processes escape impairment because they operate independently of CNS coordination; CNS-dependent demand-responsive processes fail because the coordinating organ itself is energy-depleted. This framework explains why pharmacological agents bypassing CNS coordination (midodrine, fludrocortisone) can partially restore function in otherwise energy-depleted patients.

Reconciliation with Multi-Lock Model. The selective dysfunction and multi-lock models are not mutually exclusive:

- **CNS energy crisis as initiating lock:** The multi-lock cascade may begin with CNS energy failure, which then triggers downstream immune, epigenetic, and autonomic locks
- **Lock entrenchment:** Even if CNS energy is restored, downstream locks (autoantibodies, epigenetic changes) may persist independently
- **Therapeutic implications:** Early intervention targeting CNS energy might prevent lock establishment; late intervention requires addressing both primary bottleneck and established downstream locks

The compartmental model (Figure 14.4) visualizes how CNS serves as both the primary dysfunction site and the coordination bottleneck for other compartments.

13.4.6 Circadian Energy Distribution Failure

~ Hypothesis 3: Circadian Distribution Failure Hypothesis

The suprachiasmatic nucleus (SCN) coordinates energy allocation across the 24-hour cycle in healthy individuals [395, 396]. In ME/CFS, we hypothesize that SCN dysfunction causes **temporal energy misallocation**: the brain distributes its limited energy budget incorrectly across the day, resulting in the paradoxical “second wind” phenomenon where patients often feel worse during normal waking hours and experience improved energy in the evening.

Normal Circadian Energy Distribution. The SCN orchestrates metabolic rhythms through multiple pathways [396]:

- **Orexin system activation:** Prepares glucose metabolism and cardiovascular function for active phase
- **HPA axis entrainment:** Cortisol peaks in morning to mobilize energy resources
- **Core body temperature rhythm:** Temperature rises during day, facilitating metabolic activity
- **Melatonin suppression:** Daytime suppression maintains alertness and energy availability
- **Peripheral clock synchronization:** Coordinates tissue-specific metabolic programs

ME/CFS Circadian Disruption. Multiple lines of evidence suggest circadian dysfunction in ME/CFS:

Cortisol rhythm abnormalities: ME/CFS patients show flattened diurnal cortisol variation, with lower morning levels and higher evening levels compared to controls [397, 398]. This represents a *temporal misallocation* of HPA axis resources.

Sleep-wake cycle disruption: Sleep dysfunctions in ME/CFS include sleep reversal patterns (sleeping throughout day, awake at night), suggesting fundamental circadian misalignment [399]. Disrupted TGF- β signaling may disrupt physiological rhythms in sleep, activity, and cognition.

Temperature rhythm alterations: While core body temperature mean values are normal, ME/CFS patients show greater variability in circadian temperature rhythm [400], potentially indicating SCN dysregulation of thermoregulatory energy allocation.

“Second wind” phenomenon: Many ME/CFS patients report feeling worse in morning when energy should be allocated for activity, yet experience paradoxical energy improvement in evening hours. This temporal inversion is consistent with inverted circadian energy distribution.

Hypothesized Mechanism. In ME/CFS with limited total energy capacity, SCN dysfunction causes:

1. **Morning energy deficit:** Failure to allocate sufficient resources during normal active phase (flattened cortisol peak, poor sleep quality prevents restoration)
2. **Evening energy surge:** Inappropriate energy allocation during evening hours (elevated evening cortisol, disrupted melatonin timing)
3. **Forced circadian misalignment:** Attempting to follow normal daytime schedule while energy distribution favors evening creates additional physiological stress
4. **Cycle reinforcement:** Poor daytime function leads to later activity shifting, further disrupting circadian entrainment

Testable Predictions.

1. **Inverted energy curve:** Metabolic measurements (cortisol, glucose, temperature) should show relative inversion compared to healthy controls
2. **Chronotype shift:** ME/CFS patients should show delayed chronotype preference and improved function with delayed schedules
3. **Forced alignment worsens symptoms:** Requiring strict morning schedules should worsen symptom severity
4. **Night-shift paradox:** Some ME/CFS patients may report *improved* function when working night shifts aligned with their endogenous rhythm
5. **Circadian biomarkers:** Dim light melatonin onset (DLMO) should be phase-delayed in ME/CFS patients
6. **Activity pattern correlation:** Patients with more severe “second wind” should show more pronounced cortisol rhythm flattening

Treatment Implications.

- **Schedule accommodation:** Allow patients to follow endogenous rhythm rather than forcing conventional schedule (may reduce symptom burden)
- **Light therapy:** Morning bright light exposure to entrain SCN (with caution—note previous studies showed limited efficacy [401])
- **Melatonin timing:** Strategically timed melatonin to shift circadian phase (individualized based on DLMO measurement)

- **Activity scheduling:** Align important activities with patient's natural energy peaks rather than conventional timing

Important null finding: Williams et al. found that neither melatonin nor bright-light phototherapy showed significant effects on ME/CFS symptoms or circadian phase markers [401]. This *negative result is informative*: simple circadian re-entrainment may be insufficient if the underlying problem is SCN-level energy distribution dysfunction rather than mere phase misalignment.

Integration with Other Hypotheses.

- **Energy limitation models:** Assumes limited total energy (consistent with metabolic dysfunction hypotheses); adds temporal distribution component
- **HPA axis dysfunction:** Explains flattened cortisol rhythm as consequence of SCN energy misallocation
- **Autonomic dysfunction:** SCN coordinates autonomic rhythms; dysfunction could contribute to orthostatic intolerance variability across day
- **Immune dysfunction:** Circadian clocks regulate immune function; SCN dysfunction may contribute to temporal patterns in inflammation [402]

Limitations and Uncertainties.

- Direct SCN imaging/function studies in ME/CFS patients lacking
- “Second wind” phenomenon documented anecdotally but not systematically quantified
- Null findings from light therapy/melatonin trials suggest simple circadian interventions insufficient
- Unclear whether SCN dysfunction is primary or secondary consequence
- Individual variability high—not all patients report “second wind”

Certainty: 0.50 (cortisol rhythm abnormalities documented; circadian disruption well-established; SCN-level mechanism requires validation)

13.4.7 Disease Subtype Progression

~ Hypothesis 4: Subtype Progression Hypothesis

ME/CFS may follow a predictable progression from CNS-primary disease to multi-system involvement over years, with disease duration serving as a proxy for progression stage. We hypothesize four stages: (1) CNS-primary (cognitive-dominant, mild), (2) autonomic spread (CNS + POTS), (3) peripheral involvement (multi-system + PEM), and (4) global/systemic (severe, bedbound). Early intervention may prevent progression beyond the initial stage.

Evidence for Progression. Several lines of evidence support a staged progression model:

Diagnostic delay and prognosis. Castro-Marrero et al. reported that disease duration strongly predicted outcome: patients who improved had mean duration of 2.3 years versus 6.7 years for those who did not improve [403]. This suggests that something changes with disease duration—consistent with progressive entrenchment of pathological processes.

Course heterogeneity. Stoothoff et al. identified five distinct illness trajectory patterns in a large cohort (n=1,086): fluctuating (59.7%), constantly worsening (15.9%), persisting (14.1%), relapsing-remitting (8.5%), and improving (1.9%) [404]. The constantly worsening subgroup showed significantly higher multi-system severity, suggesting progressive accumulation of affected systems.

CNS dysfunction prominence. Cognitive symptoms are reported by 85–89% of ME/CFS patients, and brainstem hypoperfusion is among the earliest neuroimaging findings [405]. This suggests CNS involvement may precede peripheral manifestations, consistent with a CNS-primary initial stage.

Autonomic progression. ME/CFS patients with comorbid POTS show worse outcomes than those with POTS alone, suggesting that the addition of autonomic dysfunction to an existing CNS-primary state represents a progression milestone.

Proposed Stage Model.

1. **Stage 1—CNS-Primary:** Predominantly cognitive symptoms (brain fog, concentration difficulty), mild fatigue, preserved physical capacity. Neuroimaging may show early hypometabolism. Duration: onset to ~6 months.
2. **Stage 2—Autonomic Spread:** CNS symptoms plus orthostatic intolerance (POTS, NMH), heart rate dysregulation, temperature instability. Autonomic testing abnormal. Duration: 6 months to ~2 years.
3. **Stage 3—Peripheral Involvement:** Multi-system symptoms including PEM, immune activation, sleep disturbance, pain. Vicious cycles (Chapter 13) become established. Duration: 2–5 years.
4. **Stage 4—Global/Systemic:** Severe, often bedbound. All systems affected. Epigenetic changes, autoantibodies, and metabolic shifts create self-sustaining pathology (the “multi-lock” state). Duration: >5 years.

Testable Predictions.

1. **Stage-biomarker correlation:** Patients at earlier stages should show fewer biomarker abnormalities (e.g., autoantibodies, NK cell dysfunction, metabolic shifts) than later-stage patients, controlling for severity.
2. **Longitudinal tracking:** A prospective inception cohort (n≥200, 5-year follow-up) should document sequential appearance of system involvement following the predicted stage order.
3. **Early intervention:** Aggressive treatment within 6 months of onset (Stage 1) should produce higher recovery rates than identical treatment at Stage 3–4, independent of

treatment type.

4. **Acute vs. gradual onset:** Acute-onset patients (e.g., post-infectious) may progress faster through stages than gradual-onset patients, but both should follow the same sequence.

Treatment Implications.

- **Stage-appropriate intervention:** Stage 1 patients may respond to CNS-targeted treatments (anti-neuroinflammatory agents, cognitive rest); Stage 3–4 patients likely require multi-targeted approaches.
- **Urgency of early treatment:** If progression is time-dependent, the window for preventing lock-in of severe disease may be narrow (<6–12 months). This supports aggressive early treatment even before full diagnostic criteria are met.
- **Prognosis estimation:** Stage assessment could help set realistic expectations and guide treatment intensity.

Limitations. All supporting evidence is cross-sectional or retrospective; no longitudinal study has tracked actual stage transitions in individual patients. The proposed stages may not be sequential in all patients—some may skip stages or experience bidirectional transitions. Disease heterogeneity means different subtypes may follow different progression patterns. The model also does not account for patients with rapid severe onset who appear to enter Stage 3–4 immediately.

Certainty: 0.45 (diagnostic delay/prognosis data supportive; staged progression pattern plausible but unvalidated longitudinally)

~ Hypothesis 5: Neutrophil Functional Exhaustion Hypothesis

Hoel et al. (2026) [191] documented a paradox: ME/CFS patients have normal peripheral neutrophil counts but profoundly suppressed circulating neutrophil-derived proteins—hallmark granule components (BPI, PAD14, MMP9, ELANE, AZU1, LTF) are all significantly reduced. We propose that ME/CFS neutrophils exist in a state of **functional exhaustion**: they are present in normal numbers but hypoactivated, failing to degranulate or mount appropriate antimicrobial responses.

Mechanistic Basis. Chronic low-grade immune activation—whether from persistent viral antigens, autoantibodies, or gut-derived LPS—may drive neutrophil tolerance, analogous to the endotoxin tolerance observed in sepsis survivors. Repeated subthreshold activation exhausts granule stores without triggering full antimicrobial programs. This would explain the simultaneous presence of elevated inflammatory markers (complement, cytokines) alongside suppressed neutrophil effector function: the alarm system is perpetually activated while the first-responder force is depleted.

Consequences. Functionally exhausted neutrophils would impair: (1) pathogen clearance, permitting low-grade infections and viral reactivation; (2) NET formation, reducing

extracellular antimicrobial defense; (3) tissue repair, as neutrophil-derived growth factors are diminished; and (4) immune regulation, since neutrophils modulate adaptive immunity through cytokine secretion and antigen presentation. This hypothesis explains why ME/CFS patients are susceptible to opportunistic infections without overt immunodeficiency.

Testable Predictions.

1. Ex vivo stimulation of ME/CFS neutrophils with PMA or fMLP should produce attenuated degranulation and ROS production compared to healthy controls, despite normal cell counts
2. Neutrophil granule protein content (measured by intracellular flow cytometry) should be reduced in ME/CFS
3. Recovery from ME/CFS (e.g., daratumumab responders) should coincide with normalization of circulating neutrophil granule proteins
4. Interventions restoring neutrophil function (G-CSF priming, if tolerable) should improve innate immune defense markers

Treatment Implications. If neutrophils are functionally exhausted rather than absent, therapeutic strategies might include: low-dose G-CSF to prime neutrophil activation (used cautiously given the inflammatory milieu), resolution-phase lipid mediators (resolvins, protectins) to reprogram rather than stimulate neutrophils, or addressing the upstream drivers of chronic activation (autoantibodies, persistent antigens) to allow neutrophil recovery.

Certainty: 0.40. Strong proteomic evidence for suppressed neutrophil signature (Hoel 2026, n=50+29, validated platform), complemented by functional neutrophil studies (Kennedy 2004). However, the exhaustion mechanism is inferred, not directly demonstrated; alternative explanations (e.g., altered neutrophil lifespan, bone marrow output changes) have not been excluded.

~ Hypothesis 6: Chronic Exerkine Signaling Hypothesis

Exercise transiently elevates FGF-21, GDF-15, and other “exerkines” to coordinate systemic metabolic adaptation—mobilizing energy substrates, promoting insulin sensitivity, and initiating tissue repair. Hoel et al. (2026) [191] found that these exerkines are chronically elevated in ME/CFS, even at rest, with FGF-21 validated in an extended cohort of 278 participants. We hypothesize that this chronic exerkine elevation represents a **metabolic emergency broadcast** that the body cannot resolve, creating a persistent state of metabolic mobilization without the exercise stimulus that would normally trigger substrate utilization and resolution.

Mechanistic Cascade. The normal exerkine cycle is: exercise → energy depletion → exerkine release → substrate mobilization → recovery → exerkine normalization. In ME/CFS, we propose: tissue hypoxia + mitochondrial dysfunction → energy depletion

at rest → exerkine release → substrate mobilization attempted → impaired utilization (mitochondrial dysfunction prevents efficient oxidative metabolism) → persistent energy deficit → continued exerkine release. The cycle cannot resolve because the downstream utilization machinery is impaired.

Why This Matters. Chronically elevated FGF-21 drives hepatic gluconeogenesis, lipolysis, and ketogenesis—metabolic programs appropriate for acute exercise recovery but harmful when sustained. Chronic GDF-15 elevation suppresses appetite (potentially explaining the anorexia some ME/CFS patients experience) and alters body composition. Persistent leptin and resistin elevation contributes to insulin resistance and inflammatory signaling. This “stuck metabolic emergency” may explain why ME/CFS patients feel as though they are perpetually recovering from extreme exertion: biochemically, they are.

Testable Predictions.

1. FGF-21 and GDF-15 levels should correlate with disease severity (particularly functional capacity and PEM threshold)
2. Successful treatment (by any mechanism) should normalize exerkine levels as downstream consequence
3. ME/CFS patients should show blunted *additional* exerkine response to actual exercise (ceiling effect), unlike healthy controls who show large fold-increases
4. Metabotype 2 patients (highest FGF-21 in Hoel 2026) should have the most severe metabolic symptoms and lowest exercise tolerance

Treatment Implications. Rather than targeting exerkines directly (they may be compensatory), treatment should address the upstream energy deficit: mitochondrial support (CoQ10, NAD⁺ precursors), tissue oxygenation improvement (addressing endothelial dysfunction), or substrate utilization enhancement. FGF-21 could serve as a treatment response biomarker—declining levels would indicate metabolic recovery even before symptom improvement becomes apparent.

Certainty: 0.45. FGF-21 elevation robustly validated (n=278, ELISA). The exerkine framework is well-established in exercise physiology. However, the “stuck cycle” mechanism is inferred from the chronic elevation pattern, and alternative explanations (hepatic stress, mitochondrial myopathy) contribute to uncertainty. Longitudinal studies tracking exerkines before and after treatment are needed.

Speculation 25 (Secretome-Driven Vascular Autoimmune Loop). Hoel et al. (2026) [191] documented two seemingly contradictory patterns: (1) reduced intracellular protein release (especially from skeletal muscle and bone marrow) and (2) increased secreted protein release (coagulation factors, complement, chemokines, hormones). We speculate that this secretome shift—elevated extracellular proteins with altered post-translational modifications from the disease environment—may generate **neo-antigens** that drive autoimmune targeting of the vasculature.

Proposed Mechanism. Under conditions of tissue hypoxia, oxidative stress, and chronic inflammation, secreted proteins may acquire abnormal modifications (carbonylation, citrullination, oxidized lipid adducts). Citrullinated proteins are recognized neo-antigens in rheumatoid arthritis; oxidized lipoproteins drive atherosclerotic autoimmunity. In ME/CFS, the chronically stressed secretome could present modified self-proteins to the adaptive immune system, generating autoantibodies against vascular and endothelial targets. The elevated complement components (CFD, C6) and reduced neutrophil proteins documented by Hoel et al. are consistent with complement-mediated vascular injury. Elevated histones (which are known damage-associated molecular patterns) and reduced neutrophil MPO (reducing antigen clearance) could amplify this process.

Why This Is Speculative. No study has directly demonstrated neo-antigen formation in ME/CFS secreted proteins, nor has vascular-specific autoimmunity been confirmed. The hypothesis chain (hypoxia → modified secretome → neo-antigen presentation → autoantibody generation → vascular targeting) involves multiple unverified links. However, it would parsimoniously connect the proteomic findings (secretome shift), the autoimmune evidence (GPCR autoantibodies, daratumumab response), and the vascular dysfunction (endothelial markers, Ephrin disruption) into a single mechanistic arc.

Testable Predictions.

1. ME/CFS serum should contain proteins with higher rates of oxidative and citrullination modifications than healthy controls
2. Autoantibodies targeting modified vascular proteins (citrullinated vimentin, oxidized LIPG) should be detectable in a subset of patients
3. The autoimmune ME/CFS subset (daratumumab responders) should show the strongest secretome shift

Certainty: 0.25. Speculative hypothesis connecting established findings (secretome shift, autoimmunity, vascular dysfunction) through an unproven mechanism. Requires dedicated proteomic studies of post-translational modifications in ME/CFS secreted proteins.

Speculation 26 (Bone Marrow Stress and Immune Lineage Reprogramming). The Hoel et al. (2026) tissue panel analysis [191] revealed that bone marrow showed the most dramatic protein reductions of any tissue type (25.4% overall net downregulation), with widespread decreases in both intracellular and secreted proteins. Simultaneously, the immune cell analysis showed a shift from innate (neutrophil/granulocyte suppression) toward adaptive predominance (elevated NK and T cell markers). We speculate that ME/CFS involves a **bone marrow stress state** that reprograms hematopoietic stem cell lineage commitment—favoring lymphoid over myeloid differentiation—resulting in the observed immune imbalance.

Biological Plausibility. Chronic inflammatory signaling (especially IFN- γ and TNF- α , both elevated in ME/CFS) is known to reprogram hematopoietic stem cells toward lymphoid differentiation at the expense of myeloid lineages—a phenomenon termed “emergency lymphopoiesis.” This would produce exactly the pattern Hoel et al. observed: normal or elevated

lymphoid cell markers alongside suppressed myeloid (neutrophil, monocyte) protein signatures. Bone marrow metabolic stress (reflected in the severe bone marrow protein reductions) could further impair normal granulopoiesis, producing functionally compromised neutrophils even at normal counts.

Testable Predictions.

1. Bone marrow biopsy in ME/CFS patients should show altered myeloid:lymphoid progenitor ratios
2. Hematopoietic stem cells from ME/CFS patients should show epigenetic marks consistent with inflammatory reprogramming
3. The bone marrow protein signature should improve with effective immunomodulatory treatment

Certainty: 0.20. Highly speculative—bone marrow has not been directly studied in ME/CFS. The tissue panel analysis is indirect (circulating proteins annotated to bone marrow origin). However, the pattern is striking and consistent with known inflammatory reprogramming of hematopoiesis.

13.5 Hypothesis-Specific Treatment Implications

Table 13.4 maps selected hypotheses to their logical treatment implications, with honest assessment of evidence and accessibility. This table focuses on hypotheses with actionable treatment options; speculative hypotheses without current interventions are omitted but appear in Table 13.2.

Table 13.4: Treatment Implications by Hypothesis

Hypothesis	Logical Treatment	Evidence for Treatment	Accessibility	Notes
Autonomic dysfunction	Salt/fluids; compression; fludrocortisone; midodrine; ivabradine; beta-blockers (ch14b)	Moderate (POTS literature)	High	Often first-line; helps many
GPCR autoantibodies	Immunoadsorption; BC007; daratumumab	Preliminary–Moderate	Very Low (specialized centers)	Most promising for autoimmune subset
Plasma cell autoimmunity	Daratumumab; bortezomib	Preliminary (pilot study)	Very Low	60% response in pilot

Continued on next page

Table 13.4 – continued from previous page

Hypothesis	Logical Treatment	Evidence for Treatment	Accessibility	Notes
Mitochondrial dysfunction	CoQ10 (ubiquinol); NAD ⁺ precursors; D-ribose; B vitamins; PQQ (§21.3)	Low–Moderate	High	Widely used; modest benefit for many
NAD ⁺ depletion	NR/NMN 1000–2000 mg/day for ≥10 weeks	Preliminary	Moderate (cost)	RCT in Long COVID showed NAD ⁺ increase
Neuroinflammation	LDN; anti-inflammatories; avoid triggers	Low–Moderate	High (LDN)	LDN widely used; helps some
Gut dysbiosis	Probiotics; dietary changes; possibly FMT	Low	High (probiotics) to Very Low (FMT)	Variable response
Endothelial dysfunction	L-citrulline/arginine; statins; low-dose aspirin; omega-3s	Theoretical	High	Untested in ME/CFS specifically
Viral persistence	Valacyclovir; valganciclovir (§21.2)	Low	Moderate	May help subset with viral markers
Small fiber neuropathy	IVIG; immunomodulation	Preliminary	Low (IVIG access)	Helps some with documented SFN
Circadian disruption	Melatonin; light therapy; time-restricted feeding; chronotherapy	Theoretical	High	Low risk; may help sleep
Glymphatic failure	Address CCI if present; optimize sleep; position	Theoretical	Variable	CCI surgery controversial
Neutrophil functional exhaustion (§13.4.7)	Address upstream drivers (autoantibodies, chronic antigens); resolution mediators (resolvins); cautious G-CSF priming	Theoretical	Variable	Neutrophil functional assays needed first
Chronic exerkine signaling (§13.4.7)	Mitochondrial support; tissue oxygenation; FGF-21 as treatment response biomarker	Theoretical	High (biomarker monitoring)	ELISA-based; FGF-21 decline may predict clinical improvement

Continued on next page

Table 13.4 – continued from previous page

Hypothesis	Logical Treatment	Evidence for Treat- ment	Accessibility	Notes
------------	-------------------	--------------------------------	---------------	-------

△ Warning 1: The Accessibility Crisis in ME/CFS Treatment

The most promising emerging treatments are essentially inaccessible to most patients:

High-Barrier Treatments:

- **Daratumumab** [9]: Requires specialized infusion center, costs \$10,000–\$20,000+ per treatment cycle, rarely covered by insurance for ME/CFS, multiple infusions needed
- **Immunoadsorption** [118]: Available only at handful of centers worldwide, requires hospitalization, costs \$15,000–\$50,000, not FDA-approved for ME/CFS in US
- **Both**: Require travel to specialized centers—impossible for severe/bedbound patients

Low-Barrier Treatments:

- **Accessible**: Pacing, supplements (CoQ10, NAD+ precursors), salt/fluids, compression
- **Evidence**: Modest effect sizes; help some patients but rarely produce major improvements

This creates a cruel disparity: the sickest patients, often bedbound and unable to travel or advocate for themselves, have the *least* access to potentially transformative treatments. Meanwhile, accessible interventions provide only modest symptomatic relief.

Implications: Research must prioritize: (1) biomarkers predicting treatment response to guide patient selection, (2) developing accessible formulations of effective therapies, and (3) understanding mechanisms to create next-generation treatments that don't require specialized delivery.

13.6 Relationships to Other Conditions

13.6.1 Fibromyalgia

Fibromyalgia (FM) shares substantial symptom overlap with ME/CFS, leading to diagnostic confusion and frequent comorbidity. Both conditions feature chronic widespread pain, fatigue, sleep disturbances, and cognitive difficulties. However, several features distinguish them:

Shared Mechanisms. Both conditions demonstrate central sensitization (amplified pain processing in the CNS), sleep architecture abnormalities (reduced slow-wave sleep, alpha-delta intrusion), autonomic dysfunction (altered HRV, orthostatic intolerance), and neuroendocrine changes (HPA axis dysfunction, altered cortisol patterns).

Distinct Features. ME/CFS is characterized by post-exertional malaise with objective deterioration on 2-day CPET, immune abnormalities (NK cell dysfunction, B cell shifts, cytokine dysregulation), and post-infectious onset in many cases. Fibromyalgia primarily features widespread pain with tender points (though diagnostic criteria have evolved), pain as the dominant symptom (whereas fatigue dominates in ME/CFS), and less consistent immune abnormalities.

Comorbidity Patterns. Studies report 35–70% comorbidity between FM and ME/CFS. This may reflect: (1) overlapping pathophysiology (shared central sensitization, autonomic dysfunction), (2) diagnostic imprecision (symptom-based criteria for both), or (3) common triggering factors (infection, trauma, stress). Some patients clearly have both conditions; others may be misdiagnosed due to symptom overlap.

13.6.2 Postural Orthostatic Tachycardia Syndrome (POTS)

POTS, defined by sustained heart rate increase ≥ 30 bpm (or ≥ 40 bpm in adolescents) within 10 minutes of standing without orthostatic hypotension, occurs in 25–50% of ME/CFS patients. POTS is a core component of the Septad framework (Section 5.6.11).

Overlap and Distinction. Many ME/CFS patients meet POTS criteria, and many POTS patients experience post-exertional symptom exacerbation. However, POTS patients without ME/CFS typically lack the severe PEM with objective physiological deterioration characteristic of ME/CFS. The key distinction: orthostatic intolerance dominates in POTS; PEM dominates in ME/CFS.

Shared Autonomic Mechanisms. Both conditions demonstrate reduced parasympathetic activity [5], impaired baroreflex sensitivity, cerebral hypoperfusion during orthostatic stress [172], and blood volume abnormalities (hypovolemia in subset). The mechanisms underlying autonomic dysfunction may differ: ME/CFS shows central catecholamine deficiency [5]; POTS mechanisms include hypovolemia, peripheral denervation, autoimmune (adrenergic receptor antibodies), and hyperadrenergic subtypes.

Treatment Considerations. POTS treatments (increased salt/fluid intake, compression garments, fludrocortisone, midodrine, ivabradine, beta-blockers) often help ME/CFS patients with orthostatic intolerance. However, these address only one component of ME/CFS pathophysiology. Pacing remains essential—POTS treatments may allow more upright time without triggering PEM, but they do not eliminate PEM risk.

See Chapters 8 and 10 for detailed autonomic pathophysiology, and Chapter 31 for POTS-MCAS-EDS mechanistic links.

13.6.3 Mast Cell Activation Syndrome

Mast cell activation syndrome (MCAS) involves inappropriate mast cell degranulation releasing histamine, tryptase, prostaglandins, and other mediators. MCAS is a core component of the Septad framework (Section 5.6.11).

Shared Features. ME/CFS and MCAS patients report overlapping symptoms: flushing, food intolerances, GI disturbances (bloating, diarrhea, abdominal pain), neurological symptoms (brain fog, headaches), and cardiovascular symptoms (tachycardia, blood pressure fluctuations). The prevalence of MCAS in ME/CFS is uncertain due to diagnostic challenges, with estimates ranging from 10–50%.

Diagnostic Challenges. MCAS diagnosis remains controversial. Consensus criteria require: (1) clinical symptoms consistent with mast cell mediator release in ≥ 2 organ systems, (2) laboratory evidence of elevated mast cell mediators during symptomatic episodes (serum tryptase, urinary methylhistamine, prostaglandin D2 metabolites), and (3) response to mast cell-directed therapy. However, mediator testing is difficult (requires collection during flare, short half-lives, specialized labs), and symptom-based diagnosis risks false positives.

Mechanistic Links. The hEDS-POTS-MCAS triad suggests shared pathophysiology. Proposed mechanisms include connective tissue abnormalities affecting mast cell stability, autonomic dysfunction triggering mast cell degranulation, and inflammatory mediators from mast cells exacerbating dysautonomia. Additionally, elevated histamine may impair cerebral blood flow and contribute to cognitive symptoms.

See Chapter 31 for MCAS-dysautonomia-vascular mechanisms and treatment chapters for screening and management protocols.

13.6.4 Autoimmune Conditions

ME/CFS shares immunological features with established autoimmune diseases and may represent an autoimmune condition in a subset of patients. The daratumumab trial [9] and GPCR autoantibody findings provide the strongest evidence for autoimmune mechanisms.

Clinical Overlap. Systemic lupus erythematosus (SLE), Sjögren's syndrome, and ME/CFS all feature fatigue, cognitive dysfunction, multi-system involvement, and female predominance. Multiple sclerosis (MS) patients often report severe fatigue resembling ME/CFS. Diagnostic challenge: distinguishing primary ME/CFS from fatigue secondary to autoimmune disease requires careful evaluation for organ-specific involvement.

Shared Immunological Features. Both ME/CFS and autoimmune diseases demonstrate B cell abnormalities (naïve/memory imbalance in ME/CFS [5]; autoreactive B cells in SLE/Sjögren's), autoantibody production (GPCR antibodies in ME/CFS; organ-specific antibodies in classic autoimmunity), T cell exhaustion markers, cytokine dysregulation, and response to immunomodulatory therapies in subsets.

Why the Immune System Connection Matters. If ME/CFS involves autoimmunity, this implies: (1) biomarker-guided patient selection for immune-targeted therapies, (2) potential for disease-modifying treatments (immunoabsorption, plasma cell depletion, B cell modulation), and (3) the need for autoimmune screening in ME/CFS patients (ANA, RF, complement, organ-specific antibodies).

Autoimmunity is one component of the Septad framework (Section 5.6.11). See treatment chapters for autoimmune screening recommendations.

13.6.5 Ehlers-Danlos Syndrome

Ehlers-Danlos syndrome (EDS), particularly the hypermobile subtype (hEDS), co-occurs with ME/CFS at rates far exceeding chance. EDS/hypermobility is a core component of the Septad framework (Section 5.6.11).

Prevalence of Comorbidity. Studies report joint hypermobility in 18–77% of ME/CFS patients (compared to 10–20% in general population). The hEDS-POTS-MCAS triad is well-recognized clinically, and many patients in this triad also meet ME/CFS criteria.

Proposed Mechanistic Connections. Connective tissue abnormalities (defective collagen) may cause: (1) vascular compliance changes leading to blood pooling and orthostatic intolerance, (2) mast cell instability (connective tissue matrix affects mast cell behavior), (3) autonomic dysfunction (structural support for blood vessels and nerve fibers compromised), and (4) craniocervical instability (CCI) in a subset, potentially impairing CSF flow and brainstem function.

Clinical Implications. ME/CFS patients should be screened for hypermobility (Beighton score). Those with significant hypermobility may benefit from: physical therapy emphasizing joint stability over flexibility, careful monitoring for structural complications (CCI, tethered cord), and treatments targeting the hEDS-POTS-MCAS triad. However, the relationship between joint hypermobility and ME/CFS pathophysiology remains incompletely understood.

See Chapter 31 for mechanistic connections and treatment chapters for screening protocols and CCI evaluation criteria.

13.6.6 Long COVID (Post-Acute Sequelae of SARS-CoV-2)

Long COVID shares remarkable symptom and pathophysiological overlap with ME/CFS, leading some researchers to propose they represent the same underlying condition triggered by different infectious agents.

Symptom Similarities. Both conditions feature: fatigue and PEM (exercise intolerance with delayed worsening), cognitive dysfunction ("brain fog"), autonomic symptoms (POTS, tachycardia, temperature dysregulation), sleep disturbances, pain, and GI symptoms. Many Long COVID patients meet ICC or CCC criteria for ME/CFS.

Pathophysiological Overlap. Shared findings include: immune dysregulation (cytokine abnormalities, T cell exhaustion, autoantibodies), endothelial dysfunction and microclotting, mitochondrial and metabolic abnormalities, autonomic dysfunction, and neuroinflammation. The Heng 2025 study [12] applied to ME/CFS could likely distinguish Long COVID with similar accuracy.

Lessons from COVID-19 Research. Long COVID research benefits from: massive funding and research attention, large patient cohorts for well-powered studies, known trigger and timing (SARS-CoV-2 infection), and less historical stigma than ME/CFS. Findings from Long COVID studies (microclots, endothelial dysfunction, viral persistence, autoantibodies) may apply directly to ME/CFS. Clinical trials for Long COVID treatments may benefit ME/CFS patients if conditions share pathophysiology.

Implications. Long COVID validates ME/CFS patient experiences—similar symptoms arising from clear viral trigger. The pandemic created millions of Long COVID cases, increasing research funding and clinical awareness that may benefit all post-viral illness patients. However, some worry Long COVID will overshadow ME/CFS, diverting resources from a decades-neglected population.

Cycle Dynamics Comparison: Long COVID as Early-Stage ME/CFS

The vicious cycle framework (Chapter 2, §2.1) provides a mechanistic lens for understanding the Long COVID–ME/CFS relationship: Long COVID may represent ME/CFS at an earlier stage of cycle recruitment.

~ Hypothesis 7: Long COVID as Stage 1–2 ME/CFS

Long COVID at 6–18 months post-infection represents early-stage ME/CFS (Stage 1–2: mitochondrial ± immune cycles active), while established ME/CFS represents late-stage disease (Stage 3–5: multiple cycles engaged with potential epigenetic lock-in).

Testable predictions:

1. Long COVID patients at 6–12 months show lower cycle involvement (1–2 cycles) than duration-matched ME/CFS patients
2. Long COVID patients meeting ME/CFS criteria at 24+ months become clinically and biologically indistinguishable from ME/CFS patients of similar duration
3. Early aggressive intervention in Long COVID (first 12 months) prevents progression to severe ME/CFS

Evidence Grade: C (mechanistically plausible; requires prospective validation)

Table 13.5: Cycle Stage Comparison: Long COVID vs. Established ME/CFS (Hypothesized)

Disease Stage	Long COVID (6–12 mo)	Long COVID (24+ mo)	ME/CFS (>5 yr)
Active cycles (hypothesized)	1–2 (mitochondrial, immune)	2–3 (+ autonomic)	3–5 (all systems)
Spontaneous recovery rate	30–50%*	10–20%*	~5% [†]
Reversibility tier (hypothesized)	Tier 1–2 (high)	Tier 2 (moderate)	Tier 2–3 (low-moderate)
Epigenetic changes (hypothesized)	Minimal	Emerging	Established
Autoantibody prevalence	30–40% [‡]	40–50% [‡]	40–60% [§]
Treatment response (hypothesized)	High potential	Moderate potential	Variable, often limited

*Long COVID recovery estimates vary widely by study and symptom definition; these ranges are model extrapolations requiring validation. [†]Systematic review median [54]. [‡]Autoantibody prevalence in Long COVID varies by assay and population. [§]GPCR autoantibodies in ME/CFS [71].

★ Key Point: Three-Field Convergence Against the Deconditioning Narrative

By 2026, the deconditioning hypothesis for ME/CFS exercise intolerance has been independently refuted from three distinct fields, each applying its own methods and standards of evidence:

1. **Exercise physiology:** Charlton et al. (2026), writing in the *British Journal of Sports Medicine*—a mainstream sports medicine journal—argued that post-exertional malaise cannot be explained by cardiac deconditioning and called for rethinking the pathophysiology of exercise intolerance in post-viral illness [341]. This marks the exercise science community challenging its own prior paradigm.

2. **Clinical guideline revision:** The UK National Institute for Health and Care Excellence withdrew graded exercise therapy from its ME/CFS guidelines in 2021, declaring it potentially harmful based on systematic evidence review [127]. This was a policy-level repudiation of the deconditioning model.
3. **Comparative virology:** The Kol et al. (2026) demonstration that coronavirus persists in lymphocytes after antiviral treatment and clinical recovery [171] provides mechanistic plausibility against the deconditioning narrative: if post-viral symptoms are maintained by a viral reservoir in immune cells, exercise cannot address the underlying pathology. Note that this is a feline coronavirus model; cross-species extrapolation to human ME/CFS requires caution.

This convergence from independent fields—each applying different methodologies and answering to different epistemic communities—constitutes stronger evidence than any single line of inquiry alone. The deconditioning narrative persists in some clinical settings despite this convergence, raising questions about knowledge translation and the sociology of medical belief change (see Section 30.6).

Prevention Opportunity. The massive Long COVID cohort presents an unprecedented opportunity to test whether early intervention can prevent ME/CFS development:

Speculation 27 (Post-Viral Syndrome Prevention Protocol). A proactive intervention protocol for Long COVID patients showing early ME/CFS features:

Weeks 4–12 post-infection: Aggressive rest if persistent symptoms; avoid “pushing through”

Months 3–6: If symptoms persist, initiate:

- Strict pacing with heart rate monitoring
- NAD⁺ precursor supplementation (NR or NMN 500–1000 mg/day)
- Mitochondrial support (CoQ10, D-ribose)

Month 6: If not resolved, full cycle mapping (Chapter 24, §24.11.5)

Months 6–12: Cycle-targeted treatment based on mapping results

Rationale: Intervene while reversibility windows remain open and before cycle recruitment cascade engages additional systems.

Evidence Grade: D (theoretical protocol; requires RCT validation)

Research Priority. A pragmatic prevention trial in early Long COVID could provide critical evidence:

- **Population:** Long COVID patients 6–18 months post-infection, not yet meeting severe ME/CFS criteria
- **Arms:** Intensive multi-target treatment vs. enhanced usual care
- **Primary endpoint:** ME/CFS diagnosis rate at 5 years

- **Sample size:** $n \approx 400$ (200 per arm) for 80% power to detect 50% reduction in ME/CFS development

If early intervention halves progression to chronic disease, this would transform post-viral syndrome management and validate the cycle dynamics prevention framework.

13.6.7 Multiple Chemical Sensitivity

Multiple chemical sensitivity (MCS)—adverse reactions to low-level chemical exposures—is reported by 20–50% of ME/CFS patients. Shared features include: sensitivity to fragrances, cleaning products, pesticides; symptom exacerbation from environmental exposures; and neurological symptoms (headache, brain fog, fatigue) following exposure.

Proposed mechanisms linking MCS and ME/CFS include: mast cell activation (chemicals trigger degranulation), neuroinflammation (sensitized microglia respond to chemical exposures), impaired detoxification (reduced hepatic clearance of xenobiotics), and central sensitization (amplified CNS response to peripheral stimuli). The relationship remains poorly understood, with MCS itself lacking clear diagnostic criteria or validated biomarkers.

13.6.8 Allergic and Atopic Conditions

ME/CFS patients report higher rates of allergies, asthma, eczema, and food sensitivities compared to the general population. This may reflect: (1) mast cell involvement (MCAS increases allergic-type reactions), (2) immune dysregulation (Th1/Th2 imbalance, IgE abnormalities), (3) histamine intolerance (reduced DAO enzyme activity, impaired histamine clearance), or (4) shared genetic susceptibility.

The mechanistic link remains unclear. Does immune dysregulation in ME/CFS predispose to atopy? Do allergic conditions trigger ME/CFS in susceptible individuals? Or does mast cell dysfunction underlie both? Current evidence cannot distinguish these possibilities, but the clinical association suggests immune system involvement extends beyond specific autoimmunity or viral responses to broader dysregulation affecting multiple pathways.

13.7 Systems Biology Approaches

ME/CFS complexity—multi-system involvement, heterogeneous presentations, treatment resistance—suggests that reductionist approaches (studying individual pathways in isolation) may miss critical emergent properties. Systems biology offers complementary methods for understanding how multiple abnormalities interact to produce the disease state.

13.7.1 Multi-Omics Integration

★ Achievement 1: Systems-Level Biomarker Panel Outperforms Single Markers

The Heng 2025 study [12] exemplifies a systems approach: integrating cellular ATP profiling (measuring AMP/ADP), plasma proteomics (2,924 proteins), and clinical data from 61 ME/CFS patients matched with 61 healthy controls revealed coordinated abnormalities across energy metabolism, immune function, and vascular biology.

A CART-derived 7-biomarker panel achieved 91% diagnostic accuracy, spanning adenosine metabolism (AMP), immune functions (cDC1, LYVE1, IGHG2), and vascular factors (FN1, VWF, THBS1). This accuracy far exceeds what any single biomarker could accomplish—individual markers show substantial overlap with controls, but their *combination* reveals disease-specific patterns. For full biomarker details and energy metabolism implications, see Achievement 6.2.1 in Chapter 6.

△ Warning 2: Replication Status

Single study (Heng 2025, n=61 pairs) from one Australian center. No independent cohort has replicated the multi-omics biomarker panel or its diagnostic accuracy. See the detailed replication assessment in Chapter 6 (Achievement 6.2.1).

This finding was independently reinforced by Hoel et al. (2026) [191], who applied aptamer-based proteomics (SomaScan, 7,326 aptamers) to 50 ME/CFS patients from the Norwegian RituxME and CycloME trial biobanks. Despite using a different analytical platform, different cohort (Norwegian vs. Australian), and different biological matrix (serum proteomics vs. cellular metabolomics + plasma proteomics), Hoel et al. found strikingly concordant multi-system signatures: immune reprogramming (suppressed neutrophil proteins, elevated complement), vascular dysfunction (altered Ephrin signaling, endothelial markers), and metabolic stress (reduced glycolytic enzymes, elevated FGF-21 and GDF-15). The convergence of two independent large-scale studies from different continents using complementary methodologies substantially strengthens the multi-system dysfunction model.

This demonstrates the power of multi-omics integration and supports the hypothesis that ME/CFS involves coordinated dysfunction across multiple systems rather than isolated abnormalities. Future studies combining genomics, epigenomics, transcriptomics, proteomics, metabolomics, and microbiomics may identify patient subgroups with distinct molecular signatures, enabling precision medicine approaches.

13.7.2 Network Analysis

Biological systems function through networks of interacting molecules, cells, and pathways. Network analysis asks: which nodes (genes, proteins, metabolites) are central to disease pathophysiology? Which perturbations propagate through the network? Where are intervention points?

Applied to ME/CFS, network approaches could:

- Identify hub genes or proteins connecting immune, metabolic, and autonomic abnormalities
- Reveal feedback loops maintaining the disease state (e.g., inflammation → mitochondrial dysfunction → immune impairment → persistent triggers → inflammation)
- Predict which interventions will have network-wide effects versus local effects
- Explain why single-target treatments often fail (network compensation/redundancy)

13.7.3 Computational Modeling

Mathematical models can integrate disparate findings into testable hypotheses about system dynamics. For ME/CFS, this could include:

Dynamical Systems Models. Representing ME/CFS as a multi-stable system with “healthy” and “diseased” attractors. Treatment would aim to push the system from the pathological basin of attraction back to health. This framework explains why: (1) triggers push susceptible individuals from healthy to diseased state, (2) the disease persists without ongoing trigger (stable attractor), and (3) small perturbations rarely produce recovery (high barrier between attractors).

Agent-Based Models. Simulating interactions between immune cells, endothelial cells, metabolic pathways, and autonomic regulation. Such models could test whether observed cellular abnormalities are sufficient to produce system-level symptoms, or whether additional mechanisms are required.

Critical Transition Theory: Bifurcation Points in ME/CFS

Critical transition theory, derived from dynamical systems mathematics, proposes that complex systems can exhibit abrupt shifts between stable states when key parameters cross threshold values [406]. This framework has been validated in ecology (lake eutrophication, coral reef collapse) and proposed for depression [407]. Whether it applies to ME/CFS is speculative but may explain several puzzling features.

~ Hypothesis 8: ME/CFS as Critical Transition Phenomenon

ME/CFS disease progression may exhibit bifurcation dynamics where small parameter changes trigger abrupt, potentially irreversible state transitions.

Mathematical framework: The cusp catastrophe model is a standard form from catastrophe theory [408]. The disease state S is governed by a potential function $V(S; \mu)$:

$$\frac{dS}{dt} = -\frac{\partial V}{\partial S}$$

For a cusp catastrophe:

$$V(S) = \frac{1}{4}S^4 - \frac{1}{2}\mu S^2 - \epsilon S$$

where μ is the control parameter (e.g., mitochondrial reserve capacity) and ϵ represents asymmetry (bias toward illness or health).

Why cusp catastrophe? This is the simplest model exhibiting bistability and hysteresis—two states (healthy, ill) that persist even when conditions change, with sudden transitions between them. We do not claim ME/CFS is a cusp catastrophe; rather, the cusp provides a minimal mathematical framework for exploring bistability hypotheses. More complex models (e.g., higher-order catastrophes, stochastic dynamics) may prove more appropriate.

Bifurcation structure:

- **Monostable region** ($\mu < \mu_c$): Single stable state (healthy or ill depending on ϵ)
- **Bistable region** ($\mu > \mu_c$): Both healthy and ill states are stable; history determines current state
- **Bifurcation point** ($\mu = \mu_c$): Small perturbations can trigger state transition

Biological interpretation (speculative):

- Control parameter μ : Mitochondrial reserve capacity, NAD⁺ levels, or composite “resilience” measure
- State variable S : Disease severity / functional capacity
- Stable states: Health, mild ME/CFS, severe ME/CFS (possibly multiple attractors)
- Bifurcation point: “Tipping point” after which recovery becomes unlikely without intervention
- Basin of attraction: Range of perturbations from which spontaneous recovery is possible

Limitations: Applying critical transition theory from ecology to human disease is analogical, not proven. Lakes and humans differ in timescales, feedback mechanisms, and measurability. The analogy motivates hypotheses but does not validate them.

Evidence Grade: D (mathematical framework established; ME/CFS application is entirely theoretical and requires empirical validation)

Early Warning Signals. Critical transition theory predicts detectable warning signals before state transitions:

1. **Critical slowing down:** Recovery time from minor perturbations increases as the system approaches a bifurcation. Near the critical point, the dominant eigenvalue $\lambda_1 \rightarrow 0$, so recovery time $\tau = 1/|\lambda_1| \rightarrow \infty$.
2. **Increased autocorrelation:** Symptom fluctuations become more persistent ($\rho(\tau) \rightarrow 1$) as the system loses resilience.
3. **Increased variance:** Symptom variability increases ($\sigma^2 \propto 1/|\lambda_1|$) before major transitions.
4. **Flickering:** The system may show transient excursions toward the alternative state before permanent transition.

Testable prediction: In the 3–6 months preceding a major severity transition (e.g., moderate → severe), patients show >50% increase in recovery time from minor perturbations and >30% increase in daily symptom variance.

Clinical Implications. If ME/CFS follows critical transition dynamics:

- **Early intervention window:** Early-stage patients near the bifurcation point may respond dramatically to interventions that would be ineffective later. Treatment timing matters as much as treatment choice.
- **Warning signal monitoring:** A smartphone app tracking daily symptoms could detect increased autocorrelation and variance, alerting patients and clinicians to impending deterioration with 30–60 day lead time.
- **Prevention vs. reversal:** Preventing progression across a bifurcation may be far easier than reversing an established transition. This provides additional rationale for aggressive early intervention.
- **Non-linear treatment response:** Some patients near bifurcation points may show dramatic responses to modest interventions (“tipping back”); others in deep pathological attractors may show minimal response to intensive treatment.

Research Priority. Validating critical transition theory in ME/CFS requires intensive longitudinal monitoring:

- **Design:** Prospective cohort with daily ecological momentary assessment (EMA)
- **Population:** ME/CFS patients at varying severity levels ($n = 200$), 12-month follow-up
- **Monitoring:** Daily 5-item symptom checklist; weekly standardized perturbation (10-minute walk); continuous heart rate variability via wearable
- **Analysis:** Time series analysis for autocorrelation, variance, and perturbation recovery time preceding transition events
- **Primary outcome:** Sensitivity/specificity of warning signals for predicting deterioration with ≥ 30 day lead time

Computational Model of ME/CFS Pathophysiology

Building on the dynamical systems and critical transition frameworks, a comprehensive computational model could integrate multiple data sources to simulate disease progression and predict intervention outcomes *in silico* before costly clinical trials.

Model Architecture.

Table 13.6: Proposed Computational Model Components

Component	Representation	State Variables
Mitochondria	Population of agents	Function level (0–1), mutation burden, ROS production
Immune cells	Population of agents	Activation state, cytokine production, autoantibody status
Metabolic pools	Continuous variables	ATP, NAD ⁺ , lactate, ROS
Signaling molecules	Continuous variables	Cytokines (IL-6, TNF- α), hormones (cortisol)
Autonomic system	Continuous variables	Sympathetic/parasympathetic tone, blood flow
Neural sensitization	Continuous variable	Central sensitization threshold

Core Equations. The model couples multiple dynamical systems representing each vicious cycle:

ATP dynamics:

$$\frac{d[\text{ATP}]}{dt} = P_{\text{mito}}(M, \text{NAD}^+) - U(\text{exertion}) - L_{\text{maintenance}}$$

Mitochondrial function:

$$\frac{dM_{\text{function}}}{dt} = B_{\text{genesis}}(\text{PGC1}\alpha) - D_{\text{ROS}}([\text{ROS}]) - D_{\text{age}}$$

ROS production:

$$\frac{d[\text{ROS}]}{dt} = R_{\text{ETC}}(M_{\text{function}}, [\text{ATP}]) - C_{\text{antioxidant}}([\text{ROS}])$$

Cytokine dynamics:

$$\frac{d[\text{Cytokines}]}{dt} = S_{\text{immune}}(\text{autoAb}, [\text{ROS}]) - C_{\text{clearance}}$$

where M = mitochondrial population, P_{mito} = ATP production function, B_{genesis} = biogenesis rate, D_{ROS} = ROS-induced damage rate.

Evidence Grade: D (theoretical framework; parameters require empirical fitting)

Model Validation Criteria.

1. **Face validity:** Simulations reproduce clinically observed behaviors (PEM delay, severity heterogeneity, treatment resistance)
2. **Predictive validity:** Model predictions match published trial effect sizes within 50%
3. **External validity:** Model trained on one dataset generalizes to independent cohorts

Clinical Applications. A validated computational model could enable:

- **In silico drug testing:** Simulate intervention effects before trials, identifying promising candidates and optimal dosing
- **Personalized trajectory prediction:** Input patient biomarkers, output probability distribution of 1-year trajectory
- **Treatment optimization:** For individual patients, simulate multiple intervention strategies and recommend optimal approach
- **Synergy identification:** Predict which treatment combinations produce super-additive effects through cycle interference

Development Roadmap.

- **Phase 1 (Year 1):** Literature-based parameter estimation; initial implementation in Python (ODE systems + agent-based components)
- **Phase 2 (Year 2):** Bayesian parameter fitting to existing cohort data; uncertainty quantification
- **Phase 3 (Years 2–3):** Validation against published trials; prospective predictions for ongoing studies
- **Phase 4 (Year 3+):** Clinical decision support tool development; open-source release

13.7.4 Challenges and Limitations

Systems biology approaches face significant challenges in ME/CFS:

- **Data requirements:** Multi-omics studies require large, well-phenotyped cohorts with standardized protocols
- **Heterogeneity:** Patient subgroups may have distinct network architectures, requiring stratification
- **Causality:** Correlation networks identify associations but cannot determine causal direction
- **Validation:** Computational predictions must be tested experimentally or clinically
- **Complexity:** Human biological networks have millions of interactions; identifying signal from noise is difficult

Despite these challenges, the multi-system nature of ME/CFS makes it an ideal candidate for systems approaches. Reductionist methods have identified many abnormalities; systems biology may reveal how they interact to produce the syndrome.

13.8 Outstanding Questions

Despite substantial progress, fundamental questions about ME/CFS pathophysiology remain unanswered. Resolving these questions will be essential for developing effective treatments and understanding the disease.

13.8.1 What Triggers ME/CFS Onset?

Most ME/CFS cases follow infection (EBV, enteroviruses, SARS-CoV-2, others), but only a small fraction of infected individuals develop ME/CFS. What determines susceptibility? Candidates include genetic variants (immune genes, metabolic pathways, HLA types), prior immune priming (previous infections, vaccinations), baseline metabolic reserve, microbiome composition at time of infection, and severity/timing of initial infection.

Large prospective cohort studies following infected individuals could identify pre-infection biomarkers predicting ME/CFS development. Understanding susceptibility could enable preventive interventions in high-risk individuals.

Genetic Modifiers of Cycle Gain and Susceptibility

The vicious cycle framework suggests genetic variants in cycle-relevant pathways modulate: (1) baseline reserve capacity (threshold before cycles engage), (2) cycle gain (amplification factor once engaged), and (3) recovery capacity.

Table 13.7: Genetic Variants Potentially Affecting Cycle Dynamics

Gene Category	Example Genes	Cycle Relevance
Mitochondrial function	WASF3, POLG, PGC-1 α	Mitochondrial cycle gain; bio-genesis capacity
Immune regulation	HLA types, IL-6, TNF- α	Immune cycle activation threshold
NAD ⁺ metabolism	NAMPT, NMNAT	Metabolic reserve; recovery rate
Oxidative stress	SOD2, catalase, GPX	Cycle dampening capacity
Autonomic function	ADRB2	Autonomic cycle susceptibility

WASF3 as exemplar: The WASF3 mutation in an ME/CFS family [64] demonstrates how genetic variants affect cycle dynamics—reduced ETC efficiency lowers ATP reserve and slows damage repair, effectively increasing mitochondrial cycle gain.

~ Hypothesis 9: Polygenic Cycle Gain Score

A polygenic risk score combining cycle-relevant variants predicts both ME/CFS susceptibility and trajectory.

Testable predictions: (1) Top quartile score shows OR > 3 for post-infection ME/CFS vs. bottom quartile; (2) Higher score correlates with faster cycle recruitment and poorer prognosis; (3) Pathway-specific scores predict which cycles activate in individual patients.

Evidence Grade: D (requires large GWAS with cycle phenotyping)

13.8.2 Why Do Some Patients Recover While Others Don't?

Spontaneous recovery occurs in some ME/CFS patients, particularly those with shorter disease duration. What distinguishes recoverers from those with persistent disease? Possibilities include: early aggressive treatment preventing “lock” establishment, less severe initial pathophysiology, genetic factors promoting recovery, effective immune resolution mechanisms, and successful identification and treatment of maintaining factors.

Understanding recovery mechanisms could identify therapeutic targets. Do recoverers have different immune profiles? Do they clear persistent viral reservoirs? Does their metabolic or autonomic function normalize, or do they compensate through alternative pathways?

The Pediatric Protection Puzzle

Pediatric ME/CFS studies report substantially higher improvement/recovery rates (54–94% depending on study and definition) compared to adult ME/CFS (median 5%, range 0–31% in systematic review) [393, 54]. This dramatic difference provides a natural experiment in protective mechanisms. Understanding *why* children recover at higher rates can inform adult treatment strategies.

△ Warning 3: Confounding in Pediatric vs. Adult Recovery Comparisons

The pediatric–adult recovery comparison is confounded by: (1) **disease duration:** pediatric cases are often caught earlier, while many adult cases represent established disease—a fair comparison requires duration-matched cohorts; (2) **diagnostic criteria:** pediatric studies may use different or less stringent criteria; (3) **ascertainment bias:** pediatric cases in clinical settings may differ from community prevalence. The mechanisms below are hypothesized explanations for whatever true pediatric advantage exists after controlling for these confounders.

Table 13.8: Candidate Pediatric Protection Mechanisms and Adult Recreation Strategies (Hypothesized)

Mechanism	Pediatric Advantage	Adult Disadvantage	Pharmacological Recreation
Mitochondrial biogenesis rate	Higher expression; PGC-1 α faster turnover*	Age-related decline in biogenesis capacity	Exercise mimetics (AICAR); NAD ⁺ precursors
NAD ⁺ levels	Higher NAD ⁺⁺ baseline	~50% decline by age 50 [†]	NMN or NR supplementation (500–1000 mg/-day)
Immune tolerance	Immature adaptive immunity; less autoantibody production [‡]	Mature immune system prone to autoimmunity	Early immunomodulation (LDN, low-dose immunosuppression)
Epigenetic flexibility	More plastic chromatin; less accumulated methylation*	Accumulated epigenetic changes resist reversal	Sirtuin activators; exercise mimetics (with caution)
Hormonal status	Pre-pubertal; stable hormonal milieu	Post-pubertal hormonal fluctuations; menstrual cycle effects	Not directly recreatable; consider hormonal optimization
Recovery environment	School accommodation enforces rest/-pacing	Work pressures; less accommodation	Disability leave in early disease
Cumulative exposures	Fewer prior infections, toxins	More cumulative cellular damage	Not recreatable

*General aging biology; not ME/CFS-specific data [409]. [†]NAD⁺ decline with age documented in general population [410]. [‡]Autoantibody prevalence increases with age generally; ME/CFS-specific pediatric–adult comparisons lacking.

~ Hypothesis 10: Pediatric Advantage Through Cycle Dynamics Lens

The pediatric recovery advantage reflects multiple factors that reduce cycle gain and preserve reversibility windows:

Lower cycle gain (G): Children have higher mitochondrial biogenesis rates and NAD^+ levels, enabling faster damage repair between perturbations. This effectively reduces the net amplification factor within the mitochondrial vicious cycle, maintaining $G < 1$ where adults would have $G > 1$.

Slower cycle recruitment: The lower autoantibody production in children's immature immune systems slows recruitment of the immune vicious cycle. Fewer cycles engaged = higher spontaneous resolution probability.

Larger reversibility windows: Higher epigenetic plasticity means damage is more readily reversed before lock-in occurs. The time-dependent reversibility decay constant (λ) may be lower in children, preserving higher $R(t)$ at any given disease duration.

Environmental pacing enforcement: School accommodations effectively enforce rest, reducing crash frequency and preventing cycle recruitment cascade.

Testable prediction: Pediatric ME/CFS patients show higher PGC-1 α expression, higher NAD^+ levels, and lower autoantibody prevalence than duration-matched adult patients; these biomarkers correlate with recovery probability.

Evidence Grade: C (mechanistically plausible; requires comparative biomarker studies)

The “Pediatric Advantage Protocol” for Adults. If pediatric protection mechanisms are recreatable, an adult protocol might include:

1. **Immediate disability leave** for first 6–12 months (if possible)—recreate the “enforced rest” of school accommodation
2. **NAD^+ precursor supplementation** (NR or NMN 500–1000 mg/day)—restore NAD^+ toward youthful levels
3. **Aggressive pacing** with heart rate monitoring—prevent crash-triggered cycle recruitment
4. **Early immunomodulation** if autoantibody-positive (LDN, immunoadsorption consideration)—prevent immune cycle entrenchment
5. **Mitochondrial biogenesis support** (CoQ10, D-ribose, urolithin A)—enhance damage repair capacity

Evidence Grade: D (theoretical protocol; requires controlled trial)

Research Priority. A cross-sectional biomarker comparison study could test whether pediatric protection mechanisms are measurable:

- **Groups:** Pediatric ME/CFS ($n = 50$), adult ME/CFS matched for duration ($n = 50$), healthy controls (pediatric $n = 25$, adult $n = 25$)
- **Biomarkers:** PGC-1 α expression, NAD^+/NADH ratio, autoantibody panel, mtDNA copy number

- **Follow-up:** 2 years to correlate baseline biomarkers with recovery outcome
- **Hypothesis:** Pediatric patients show 2–3× higher biogenesis markers and 30–50% lower autoantibody prevalence

13.8.3 What Maintains the Disease State?

Even if the initial trigger (infection) is cleared, ME/CFS persists. Proposed maintenance mechanisms include: persistent viral reservoirs (latent herpesviruses, integrated RNA fragments), autoantibodies from long-lived plasma cells, epigenetic changes locking pathological gene expression, metabolic pathway shifts to new equilibrium, immune system recalibration (trained immunity), autonomic nervous system setpoint changes, and microbiome alterations perpetuating dysbiosis.

Determining which mechanisms operate in which patients is critical for treatment selection. A patient with autoantibody-driven disease requires immunomodulation; one with epigenetic changes might benefit from epigenetic modifiers; one with metabolic traps needs metabolic interventions.

13.8.4 How Do Different Subtypes Differ Mechanistically?

ME/CFS heterogeneity likely reflects distinct pathophysiological mechanisms rather than a single disease entity. Potential subgroups include: autoimmune subtype (daratumumab responders, GPCR antibody-positive), metabolic subtype (primary mitochondrial dysfunction), autonomic subtype (POTS-predominant), neuroinflammatory subtype (microglial activation, CNS-predominant symptoms), post-viral subtype (persistent viral markers, reactivation), and gut-mediated subtype (dysbiosis-driven).

Rigorous cluster analysis of multi-omic data may objectively define subtypes. Treatment trials should stratify by subtype to avoid diluting signals when effective therapies help only specific patient groups.

13.8.5 Can We Identify Critical Intervention Points?

If ME/CFS involves multiple reinforcing abnormalities (multi-lock model), which locks must be broken for recovery? Do certain interventions have cascading benefits (break one lock, others follow)? Or must all locks be addressed simultaneously? Can early intervention prevent lock establishment, making treatment more effective in acute/early disease?

These questions will determine treatment strategy: sequential targeting of individual mechanisms versus simultaneous multi-pronged interventions. The answer may differ by patient subtype.

Conclusion: Chapters 7–11 documented specific abnormalities across physiological systems. This chapter attempted to synthesize those findings into coherent models while acknowledging uncertainty. The complexity of ME/CFS—multi-system involvement, heterogeneity,

treatment resistance—demands both reductionist investigation of individual mechanisms and systems-level integration. Progress requires both approaches working in concert, guided by honest assessment of evidence quality and explicit acknowledgment of what we do not yet understand.

13.8.6 Research Priorities

Based on this synthesis, the following research directions appear most critical:

1. **Biomarker validation for patient stratification:** The Heng 2025 panel [12] and daratumumab response patterns [9] suggest identifiable subgroups. Large multi-center studies should validate these biomarkers and develop clinical decision tools.
2. **Mechanism-targeted trials with biomarker selection:** Rather than treating all ME/CFS patients identically, trials should enroll patients based on mechanistic biomarkers (autoantibody-positive, severe autonomic dysfunction, primary metabolic abnormalities) and test subgroup-specific interventions.
3. **Combination therapy trials:** Test whether simultaneously targeting multiple mechanisms (e.g., immunoadsorption + metabolic support + autonomic treatment) produces superior outcomes to single interventions.
4. **Prospective cohort studies of infection:** Follow individuals before and after triggering infections (influenza, COVID-19, EBV) to identify pre-morbid risk factors and early biomarkers predicting ME/CFS development. This could enable prevention.
5. **Recovery mechanism studies:** Systematically characterize patients who improve or recover—what distinguishes them biologically? Understanding recovery pathways could identify therapeutic targets applicable to those with persistent disease.
6. **Early intervention trials:** Test whether aggressive treatment within 6-12 months of onset prevents “lock” establishment and improves long-term outcomes. The window of treatment responsiveness may be limited.
7. **Systems biology approaches:** Apply network analysis and multi-omics integration to identify critical nodes in ME/CFS pathophysiology. Computational modeling may reveal non-obvious intervention points.

The field stands at an inflection point. Decades of patient advocacy and recent high-profile cases (Long COVID) have increased research funding and clinical awareness. The biological basis of ME/CFS is now undeniable [5, 12, 9]. The challenge is translating mechanistic insights into effective treatments accessible to all patients who need them.

Chapter 14 extends this analysis to more speculative mechanisms that, while lacking direct evidence in ME/CFS, may provide insights into disease pathophysiology and suggest novel therapeutic approaches. Where this chapter focused on evidence-based integration, the next explores creative hypotheses that may inspire future research.

14 Speculative Mechanistic Hypotheses

*“The scientist is not a person who gives the right answers,
he’s one who asks the right questions.”*
— Claude Lévi-Strauss

This chapter presents speculative hypotheses about ME/CFS pathogenesis that emerge from creative extrapolation of known biochemistry, systems biology, and pattern recognition across medical domains. While not yet empirically validated in the ME/CFS context, each hypothesis attempts to explain the characteristic features of the illness—post-exertional malaise, chronicity, multi-system involvement, and treatment resistance—through mechanisms that are individually plausible and potentially testable.

These hypotheses are offered in the spirit of scientific brainstorming: to stimulate new research directions, generate testable predictions, and potentially identify overlooked connections. They should be evaluated by their ability to generate novel experiments and explain otherwise puzzling observations, not treated as established fact.

14.1 Master Hypothesis Table: Likelihood and Therapeutic Potential

Table [14.1](#) provides a comprehensive overview of all hypotheses presented in this chapter, ranked by evidence strength and therapeutic potential. This serves as a roadmap for both researchers prioritizing investigation directions and clinicians considering experimental interventions.

Table 14.1: Comprehensive ranking of all speculative hypotheses by evidence level, therapeutic potential, and impact on different severity levels							
Hypothesis	Evidence Level	Therapeutic Potential	Benefit: Mild	Benefit: Severe	Explains Key Features	Nearest-Term Action	
CPET-Derived Hypotheses (Objective Functional Data)							
Autonomic-mitochondrial feedback loop	Moderate	High	High	Moderate	PEM, recovery time, autonomic symptoms	Trial: tyrosine+Tetrahydrobiopterin (BH4)+antioxidants	
Mitochondrial turnover rate limitation	Moderate-High	High	Moderate-High	Moderate	13-day recovery, cumulative decline, GET failure	Urolithin A + NAD+ precursor trial	
Exercise metabolomics-guided therapy	Moderate	Very High	High	Low	Individual variation, treatment heterogeneity	Post-CPET metabolomics study	
Circadian recovery gating	Low-Moderate	Moderate	Moderate	Moderate	Sleep dysfunction, non-restorative rest	Chronotherapy pilot study	
Vagal stimulation for recovery	Low-Moderate	Moderate	Moderate	Low-Moderate	Autonomic dysfunction, inflammation persistence	Post-exertion VNS trial	
Core Mechanistic Hypotheses							
Metabolic “safe mode” lock	Moderate	High	Low-Moderate	Moderate-High	PEM, chronicity, resistance to rehabilitation	Hypothalamic modulation interventions	
Glymphatic clearance failure	Low-Moderate	Moderate	Moderate	Moderate-High	Brain fog, non-restorative sleep, orthostatic symptoms	CSF flow imaging; craniocervical assessment	
Tryptophan/kynurenine trap	Moderate	Moderate-High	Moderate	Moderate	Cognitive symptoms, depression, immune activation	IDO inhibition trials	
Vagal afferent danger signal loop	Low-Moderate	Moderate-High	Moderate	High	Rapid symptom onset, gut-brain connection, PEM	Vagal modulation; gut interventions	
Purinergic signaling dysregulation	Low-Moderate	Moderate	Moderate	Moderate	Immune dysfunction, pain, fatigue, inflammation	P2X/P2Y receptor modulators	
Redox compartment collapse	Moderate	Moderate	Moderate	Low-Moderate	Oxidative stress, chemical sensitivities	Glutathione/N-Acetylcysteine (NAC) optimization	
Metabolic memory / epigenetic lock	Moderate	Low-Moderate	Low	Low-Moderate	Chronicity, treatment resistance	Epigenetic modifiers (exploratory)	
Circadian-metabolic desynchronization	Moderate	Moderate	Moderate	Low-Moderate	Sleep issues, energy fluctuations	Circadian stabilization protocols	
Autoimmune/Immune Hypotheses							
GPCR autoantibody-driven dysfunction	Moderate-High	Very High	High	Moderate-High	POTS, autonomic symptoms, 60% daratumumab response	Autoantibody testing; immunoadsorption; daratumumab	
Plasma cell sanctuary hypothesis	Moderate	Very High	High	High	Rituximab failure vs daratumumab success, chronicity	Anti-CD38 therapy; combined IA+daratumumab	
Autoantibody-monocyte activation cascade	Low-Moderate	Moderate-High	Moderate	Moderate	Inflammatory cytokines, MIP-1δ, PDGF-BB elevation	Monocyte-targeted therapy; autoantibody removal	
Ion channel autoimmunity	Low-Moderate	Moderate-High	Moderate-High	Moderate	Autonomic symptoms, POTS, cognitive issues	Autoantibody screening; immunoadsorption	
TRPM3 channelopathy	Moderate-High	High	High	Moderate-High	NK cell dysfunction, impaired immune cell calcium signaling	TRPM3 functional testing; calcium signaling studies; pregnenolone trial (speculative)	
Endothelial trained immunity	Low	Moderate-High	Moderate	Moderate	Multi-system symptoms, vascular dysfunction, PEM	Endothelial epigenetic profiling	

14 Speculative Mechanistic Hypotheses

14 Speculative Mechanistic Hypotheses

Continued on next page

Table 14.1 – continued from previous page						
Hypothesis	Evidence Level	Therapeutic Potential	Benefit: Mild	Benefit: Severe	Explains Key Features	Nearest-Term Action
Receptor internalization (not blockade)	Low-Moderate	Moderate-High	Moderate	Moderate	Lag between Ab removal and improvement; receptor density changes	Receptor density assays on patient lymphocytes
Functional vs. binding assay discrepancy	Moderate	Very High	High	High	Failed replications; heterogeneous treatment response	Develop functional autoantibody assays
Viral/Cellular Hypotheses						
EBV-B cell CNS infiltration	Low-Moderate	High	Moderate	Moderate-High	Post-EBV onset; neuroinflammation; brain fog	CSF B cell analysis; LMP1 profiling
EBV-GPCR molecular mimicry	Low	High	Moderate-High	Moderate-High	EBV trigger specificity; persistent autoantibodies	Computational homology; cross-reactivity testing
Endogenous retrovirus reactivation	Very Low	Low	Low	Low	Post-viral onset, immune activation, chronicity	HERVs expression profiling
Cellular quorum sensing dysfunction	Very Low	Low	Low-Moderate	Low	Systemic coordination loss, multi-system involvement	Basic research needed
Metabolic Compartmentalization Hypotheses						
Lactate compartmentalization disorder	Low	Moderate	Low-Moderate	Low-Moderate	Exercise intolerance, muscle symptoms, brain lactate	MCT function studies; dietary ketones
Ferroptosis susceptibility	Low	Low-Moderate	Low-Moderate	Low	Oxidative stress, lipid peroxidation, tissue damage	Ferroptosis inhibitors (research)
Integrated/Multi-System Hypotheses						
Selective energy dysfunction	Moderate	High	Moderate-High	Moderate-High	Preserved autonomous functions (hair, nails), impaired CNS-dependent processes; demand-response failure	Hair follicle mito assay; CSF lactate; CNS-targeted delivery (Sec. 14.24)
Multi-lock integrated trap	High conceptual	Very High	Variable	Variable	Heterogeneity, treatment resistance, chronicity	Multi-target interventions
Cycle Dynamics Hypotheses						
Vicious cycle network coupling	Moderate	Very High	High	High	Treatment resistance, heterogeneity, why single interventions fail	Multi-target combination trials (CoQ10+LDM); biomarker-guided selection
Critical transition bifurcation	Low-Moderate	High	High	Moderate	Sudden deteriorations, non-linear progression, “tipping points”	Longitudinal monitoring for warning signals (increased symptom variance)
Cycle-predominant subgroups	Moderate	High	High	High	Treatment heterogeneity, why same treatment works for some	Comprehensive cycle diagnostic battery; targeted interventions
Crash dose-response with threshold	Moderate-High	Very High	Very High	Very High	Why small overexertion can be catastrophic, cumulative harm	Individual VT measurement; strict envelope pacing
Time-dependent reversibility decay	Low-Moderate	High	High	Moderate	Why early intervention crucial, chronic treatment resistance	Aggressive early treatment; realistic expectation setting
Cycle recruitment cascade	Low-Moderate	Moderate-High	Very High	Low-Moderate	Disease progression, why mild becomes severe without intervention	Early pacing to prevent cascade; monitor for new cycle activation
High-Risk/Counterintuitive Hypotheses						
Metabolic preconditioning (hormesis)	Very Low	Low (High Risk)	Unknown	Contraindicated	Adaptation failure?	NOT RECOMMENDED clinically
Blood flow restriction training	Low	Low-Moderate	Low-Moderate	Contraindicated	Oxygen delivery dysfunction	Research only; high risk

14 Speculative Mechanistic Hypotheses

14.1.1 How to Use This Table

For Researchers

High-priority investigations (Moderate-High evidence, testable):

1. TRPM3 channelopathy: Replication in additional cohorts; characterization of dysfunction mechanism (hypo- vs hyperfunction); correlation with symptom severity
2. Mitochondrial turnover limitation: Urolithin A intervention with repeat two-day CPET
3. Autonomic-mitochondrial loop: Multi-target combination trial
4. Exercise metabolomics: Post-CPET metabolomic profiling to identify subgroups
5. Ion channel autoimmunity: Comprehensive autoantibody screening (including anti-TRPM3)

Medium-priority investigations (plausible mechanisms, need preliminary data):

1. Glymphatic function: Imaging studies assessing CSF flow dynamics
2. Tryptophan trap: IDO inhibitor safety/efficacy trials
3. Vagal interventions: VNS for post-exertional recovery
4. Circadian optimization: Chronotherapy protocols

Basic research needed (very low evidence, high theoretical interest):

1. Cellular quorum sensing mechanisms
2. Endogenous retrovirus expression patterns
3. Ferroptosis markers and susceptibility

For Clinicians

Relatively safe to trial (assuming medical supervision and appropriate patient selection):

- Autonomic-mitochondrial support (supplements, generally recognized as safe)
- Mitochondrial turnover acceleration (urolithin A, NAD⁺ precursors have human safety data)
- Chronotherapy/circadian stabilization (behavioral, very low risk)
- Vagal stimulation (non-invasive, established safety profile)
- Tryptophan metabolism support (within normal supplement ranges)

Requires specialist supervision:

- Ion channel autoantibody testing and immunoadsorption
- IDO inhibition (investigational)
- Epigenetic modifiers

Not recommended outside research protocols:

- Metabolic preconditioning/hormesis approaches (high risk of PEM)
- Blood flow restriction training (could worsen oxygen delivery dysfunction)
- Endogenous retrovirus interventions (purely theoretical)

For Patients

Understanding evidence levels:

- **Very Low:** Purely theoretical speculation; interesting for research but no evidence
- **Low:** Mechanism makes sense based on other diseases; no ME/CFS-specific data
- **Low-Moderate:** Some indirect evidence in ME/CFS; plausible but unproven
- **Moderate:** Multiple ME/CFS studies support mechanism; direct intervention untested
- **Moderate-High:** Strong mechanistic support; similar interventions show promise
- **High:** Direct evidence from ME/CFS trials (rare in this chapter, as these are speculative hypotheses)

Severity-specific guidance:

- **Mild-moderate patients:** May benefit from metabolomics-guided approaches, autonomic support, circadian optimization
- **Severe patients:** Prioritize hypotheses addressing core metabolic function (safe mode, mitochondrial turnover, glymphatic clearance); avoid any interventions requiring exertion
- **All severities:** Multi-lock hypothesis suggests combinations may work better than single interventions

14.1.2 Qualification and Caveats

△ Warning 1: Speculative Content

ALL hypotheses in this chapter are speculative to varying degrees. The evidence levels indicate relative plausibility and existing support, but even “Moderate-High” evidence hypotheses remain unproven. Therapeutic approaches derived from these hypotheses should be considered experimental and discussed with knowledgeable physicians. Patient self-experimentation carries risks, especially for severe patients where any metabolic perturbation might trigger crashes.

14.2 Metabolic “Safe Mode” Hypothesis

? Open Question 1: Stuck Sickness Behavior Program

What if ME/CFS represents an evolutionarily conserved “sickness behavior” metabolic program that fails to disengage? The body detects a threat (infection, severe stress) and deliberately downregulates energy production as a protective mechanism—analogue to a computer entering safe mode. Normally this resolves when the threat passes, but some trigger causes the metabolic thermostat to become locked in the suppressed state.

Under this model, the itaconate shunt activation, IDO pathway upregulation, and mitochondrial suppression observed in ME/CFS are not dysfunction per se—they represent an intentional protective program that refuses to terminate. This would explain why “pushing through” causes deterioration: physical exertion fights against an active suppression system that interprets increased metabolic demand as evidence the threat persists.

The evolutionary rationale would be that during infection, reducing activity and metabolic rate conserves resources for immune function while limiting pathogen replication (many pathogens depend on host metabolism). The “lock” might involve persistent immune signaling, epigenetic changes to metabolic genes, or alterations to the hypothalamic setpoint that normally regulates this response.

14.2.1 Mechanistic Details

The sickness behavior response is mediated by inflammatory cytokines (IL-1 β , IL-6, TNF- α) acting on the hypothalamus and other brain regions. These signals normally produce:

- **Fatigue and reduced activity:** Conserving energy for immune function
- **Anorexia:** Limiting nutrients available to pathogens
- **Fever:** Creating hostile environment for pathogens
- **Social withdrawal:** Reducing transmission risk
- **Hyperalgesia:** Promoting protective behaviors
- **Cognitive changes:** Redirecting attention to recovery

In ME/CFS, patients exhibit most of these features chronically, without fever (which may require acute, high-level cytokine signaling). The “safe mode” hypothesis proposes that the metabolic suppression aspect of sickness behavior has become dissociated from its normal regulatory feedback and persists indefinitely.

14.2.2 Why the Program Might Lock

Several mechanisms could prevent normal disengagement:

Persistent Low-Grade Immune Activation. Even without active infection, ongoing immune activation (from autoantibodies, reactivated herpesviruses, gut barrier dysfunction, or other sources) could maintain the cytokine signals that keep the program engaged.

Hypothalamic Setpoint Shift. The hypothalamus integrates peripheral signals and sets metabolic “targets.” A severe enough initial insult might shift these setpoints, such that normal physiological states are now interpreted as requiring continued suppression.

Epigenetic Stabilization. The gene expression changes that implement sickness behavior might become epigenetically stabilized through DNA methylation or histone modifications, persisting even after the signaling that induced them resolves.

Receptor Desensitization Failure. Normally, prolonged cytokine exposure leads to receptor desensitization, allowing the organism to “adapt” and resume normal function. Failure of this desensitization would maintain responsiveness to even low-level signals.

14.2.3 Testable Predictions

1. ME/CFS patients should show patterns of gene expression consistent with acute sickness behavior, even in the absence of detectable infection
2. Hypothalamic function should differ from healthy controls in ways consistent with altered setpoints
3. Markers of metabolic suppression (itaconate, altered mitochondrial dynamics) should correlate with symptom severity
4. Interventions that “reset” the hypothalamic setpoint might provide benefit
5. The pattern should differ from simple deconditioning in specific, identifiable ways

14.3 Glymphatic/CSF Clearance Failure

? Open Question 2: Impaired Brain Waste Clearance

The brain’s glymphatic system clears metabolic waste primarily during sleep, driven by CSF flow through perivascular channels. Could ME/CFS involve impaired glymphatic function—potentially from craniocervical instability, altered intracranial pressure dynamics, or autonomic dysfunction affecting the arterial pulsation that drives the system?

If metabolic waste (including inflammatory mediators, misfolded proteins, and neurotransmitter metabolites) accumulates in the CNS, this could directly cause the cognitive dysfunction (“brain fog”) characteristic of ME/CFS. The body might respond to CNS waste accumulation by inducing fatigue to force rest and enable clearance. However, if the clearance mechanism itself is impaired, rest alone cannot resolve the accumulation,

creating a self-perpetuating state.

This hypothesis connects several observations: the sleep abnormalities in ME/CFS (patients sleep but don't feel restored—possibly because glymphatic clearance is impaired even during sleep), the cognitive symptoms, and the correlation between some patients' symptoms and cervical spine issues. The post-exertional component could reflect exercise-induced increases in CNS metabolic waste production that overwhelm an already-compromised clearance system.

14.3.1 The Glymphatic System

Discovered relatively recently (2012), the glymphatic system is the brain's waste clearance pathway. Key features include:

- CSF flows along periarterial spaces into the brain parenchyma
- Aquaporin-4 (AQP4) water channels on astrocyte endfeet facilitate fluid exchange
- Interstitial fluid carrying waste products drains along perivenous spaces
- Activity increases dramatically during sleep (especially slow-wave sleep)
- Arterial pulsation provides the driving force for fluid movement
- The system clears amyloid- β , tau, and other potentially neurotoxic waste

14.3.2 Potential Disruption Mechanisms

Craniocervical Instability. Some ME/CFS patients have craniocervical junction abnormalities that could impair CSF flow dynamics. The relationship between neck position and symptoms reported by some patients might reflect positional effects on CSF circulation.

Autonomic Dysfunction. Arterial pulsation drives glymphatic flow. Autonomic dysfunction affecting cardiovascular regulation could reduce the pulsatile pressure gradients needed for effective clearance.

Sleep Architecture Abnormalities. Glymphatic clearance is most active during slow-wave sleep. The sleep abnormalities documented in ME/CFS—reduced slow-wave sleep, fragmented sleep architecture—would directly impair clearance even if the system itself were intact.

Neuroinflammation. Inflammation alters AQP4 localization and astrocyte function, potentially impairing the cellular machinery required for glymphatic transport.

Intracranial Pressure Dysregulation. Both elevated and reduced intracranial pressure could impair CSF dynamics. The orthostatic symptoms in ME/CFS might relate to pressure dysregulation that worsens glymphatic function.

14.3.3 Connections to ME/CFS Features

This hypothesis provides explanations for:

- **Cognitive dysfunction:** Direct effect of CNS waste accumulation
- **Unrefreshing sleep:** Sleep fails to accomplish its clearance function
- **Post-exertional malaise:** Exercise increases metabolic waste production faster than it can be cleared
- **Sensitivity to position:** Effects of posture on CSF dynamics
- **Headaches:** Common in conditions of impaired CSF flow
- **Improvement with strict rest:** Reduces waste production, allowing partial catch-up

14.3.4 Testable Predictions

1. Advanced MRI techniques (e.g., diffusion tensor imaging along perivascular spaces) should reveal altered glymphatic flow in ME/CFS patients
2. CSF biomarkers of waste accumulation (amyloid- β , tau, neurofilament light) might be elevated
3. Sleep interventions specifically targeting slow-wave sleep enhancement might provide benefit
4. Treatments that improve CSF dynamics (addressing craniocervical issues, improving cardiovascular function) might help subsets of patients
5. Symptom severity might correlate with measures of glymphatic function

⚠ **Limitation 1: Speculative Mechanistic Hypotheses: Plausibility Is Not Evidence**

The hypotheses in this chapter—metabolic “safe mode,” glymphatic impairment, HERV reactivation, lactate compartmentalisation, vagal danger signalling, kynurenine trapping, and purinergic quorum sensing—share a common epistemic limitation: each is internally consistent and mechanistically plausible, but none has been directly tested in ME/CFS patients. Key boundaries:

- No prospective study has demonstrated that any of these mechanisms *precedes* or *causes* ME/CFS; all are inferred from cross-sectional data, analogy to other diseases, or theoretical modelling.
- The “safe mode” / “locked sickness behaviour” framing is a narrative organising device, not an empirically validated construct; identical language could be applied post hoc to any chronic illness with inflammatory features.
- Testable predictions listed for each hypothesis are necessary but not sufficient: confirmation of individual predictions would support but not prove the overarching model, as alternative mechanisms could generate the same observations.
- Several hypotheses (HERV reactivation, purinergic quorum sensing, MCT dysfunction) rest on phenomena demonstrated in cell lines or animal models that have not

been measured in ME/CFS cohorts at all.

14.4 Endogenous Retrovirus Reactivation

? Open Question 3: HERV De-Silencing

Human genomes contain approximately 8% endogenous retroviruses (HERVs)—ancient viral sequences integrated into our DNA over millions of years. These are normally epigenetically silenced, but stress, infection, or inflammation can trigger their de-silencing and transcription.

Reactivated HERVs don't produce infectious virus, but they do produce immunogenic proteins that the immune system may recognize as foreign. This creates a form of autoimmunity where the immune system attacks “self” proteins that weren't previously expressed. The chronic immune activation in ME/CFS—without a detectable exogenous pathogen—could reflect ongoing response to HERV-derived antigens.

This would explain why ME/CFS often follows viral infection (the infection triggers HERV de-silencing), why immune activation persists without detectable pathogen, and why immunosuppression sometimes provides benefit. It also provides a mechanism for the female predominance, as sex hormones influence epigenetic regulation and HERV expression.

14.4.1 Biology of Human Endogenous Retroviruses

HERVs represent the remnants of ancient retroviral infections that integrated into the germline and were passed to subsequent generations. Key facts:

- HERVs comprise ~8% of the human genome (more than protein-coding genes)
- Most are defective and cannot produce infectious virus
- Many retain open reading frames capable of producing proteins
- Expression is normally suppressed by DNA methylation and other epigenetic mechanisms
- Various stressors can trigger HERV de-silencing: viral infection, inflammation, hormonal changes, oxidative stress
- HERV proteins can be immunogenic, triggering immune responses
- HERV involvement has been documented in multiple sclerosis, schizophrenia, and autoimmune conditions

14.4.2 The HERV-ME/CFS Connection

Triggering De-Silencing. An acute viral infection (EBV, enteroviruses, SARS-CoV-2) could trigger HERV de-silencing through:

- Direct transactivation by viral proteins
- Inflammatory cytokines altering epigenetic regulation
- Oxidative stress damaging DNA methylation patterns
- Hormonal stress responses affecting chromatin state

Sustained Immune Activation. Once de-silenced, HERVs produce proteins that:

- Are recognized as foreign by the adaptive immune system
- Trigger antibody production and T cell responses
- Create ongoing inflammation that perpetuates de-silencing
- May cross-react with normal cellular proteins (molecular mimicry)

Tissue-Specific Effects. Different HERV families have different tissue expression patterns. The particular HERVs de-silenced might determine which symptoms predominate—neurotropic HERVs causing cognitive symptoms, muscle-expressed HERVs causing fatigue, etc.

14.4.3 Supporting Observations

- The post-viral onset pattern fits HERV triggering
- Immune activation without detectable pathogen is consistent
- Female predominance aligns with hormonal influence on HERV regulation
- Variable symptom patterns could reflect different HERV expression profiles
- Partial response to immunomodulation is expected if autoimmunity is involved
- The XMRV controversy, though ultimately negative, reflected intuitions about retroviral involvement that HERV reactivation could fulfill

14.4.4 Testable Predictions

1. ME/CFS patients should show elevated HERV transcription compared to controls, particularly for specific HERV families
2. Antibodies against HERV proteins should be detectable in patient sera
3. HERV expression levels might correlate with disease severity or specific symptoms
4. Treatments targeting HERV expression (antiretrovirals, epigenetic modifiers) might provide benefit
5. The specific HERVs activated might predict symptom clusters or treatment response

14.5 Lactate Compartmentalization Disorder

? Open Question 4: Monocarboxylate Transporter Dysfunction

During post-exertional malaise, lactate accumulates abnormally in ME/CFS patients. But what if the problem isn't excess lactate production but rather impaired lactate redistribution?

Monocarboxylate transporters (MCTs) shuttle lactate between cellular compartments and tissues. Lactate produced in exercising muscle normally travels to the liver for gluconeogenesis (Cori cycle) or to the heart and brain as fuel. If MCT function is impaired, lactate becomes "trapped" in the tissues where it's produced, creating local acidosis and energy failure even while systemic lactate levels might appear relatively normal.

This would explain why ME/CFS patients show abnormal lactate responses to exercise, why symptoms are so localized and variable, and why the severity of post-exertional malaise correlates poorly with objective measures of exertion. The compartmentalization means you're producing lactate faster than you can redistribute it, creating metabolic bottlenecks in specific tissues.

14.5.1 Lactate Physiology

Lactate is far more than a waste product. Modern understanding recognizes lactate as:

- A major fuel source for heart, brain, and resting muscle
- A gluconeogenic precursor (Cori cycle)
- A signaling molecule affecting gene expression and metabolism
- A redox shuttle between cellular compartments
- Normally in constant flux between tissues based on metabolic state

The lactate shuttle depends on monocarboxylate transporters (MCT1-4), each with different tissue distributions and kinetic properties:

- **MCT1:** Ubiquitous; facilitates lactate uptake in oxidative tissues
- **MCT2:** High affinity; concentrated in neurons
- **MCT3:** Retinal pigment epithelium
- **MCT4:** Low affinity; facilitates lactate export from glycolytic tissues

14.5.2 Compartmentalization Pathophysiology

If MCT function is impaired:

Muscle. Lactate produced during exercise cannot efficiently exit muscle cells. Local acidosis develops, causing pain, weakness, and premature fatigue. Even mild exercise creates disproportionate symptoms.

Brain. Neurons depend heavily on lactate from astrocytes (astrocyte-neuron lactate shuttle). Impaired MCT2 would create neuronal energy deficits and cognitive dysfunction. The brain would be simultaneously lactate-starved despite peripheral lactate accumulation.

Heart. The heart preferentially oxidizes lactate during exercise. Impaired lactate delivery could limit cardiac output and contribute to exercise intolerance.

Liver. Reduced lactate delivery to the liver impairs gluconeogenesis, potentially contributing to hypoglycemic symptoms and energy crashes.

14.5.3 Why MCT Function Might Be Impaired

- **Inflammatory cytokines:** IL-1 β , TNF- α affect MCT expression
- **Hypoxia:** Alters MCT isoform expression patterns
- **pH dysregulation:** MCT function is pH-sensitive
- **Oxidative damage:** MCTs can be modified by ROS/RNS
- **Autoantibodies:** Antibodies against MCTs are theoretically possible
- **Mitochondrial dysfunction:** Alters cellular lactate handling

14.5.4 Testable Predictions

1. Muscle biopsies should show altered MCT expression or localization
2. Lactate imaging (using ^{13}C -MRS or hyperpolarized ^{13}C -lactate) should reveal abnormal compartmentalization
3. Blood lactate might appear relatively normal while tissue lactate is elevated
4. Interventions supporting MCT function (dichloroacetate, lactate supplementation to bypass MCT) might help
5. The specific MCTs affected might predict which tissues/symptoms predominate

14.6 Vagal Afferent “Danger Signal” Loop

? Open Question 5: Persistent Interoceptive Threat Signaling

The vagus nerve carries afferent signals from peripheral organs to the brain, conveying information about inflammation, metabolic state, and tissue damage. These signals normally trigger appropriate “sickness behavior” responses. What if ME/CFS involves persistent, inappropriate vagal afferent signaling that continuously tells the brain “there is danger in the periphery”?

This could result from sensitized vagal afferents, low-grade peripheral inflammation that genuinely activates these pathways, or central misinterpretation of normal afferent

traffic. The brain, receiving constant danger signals, maintains the organism in a chronic sickness state regardless of actual peripheral conditions.

This hypothesis explains why vagal nerve stimulation sometimes helps ME/CFS patients (it might “reset” the aberrant signaling), why gut symptoms are so common (the gut provides major vagal input), and why stress exacerbates symptoms (stress sensitizes vagal pathways). It also provides a mechanism for the brain-body disconnect where patients feel systemically ill despite relatively normal peripheral findings.

14.6.1 Vagal Afferent Pathways

The vagus nerve is predominantly afferent (~80% of fibers carry signals TO the brain). These afferents:

- Sense inflammatory mediators (cytokines, prostaglandins) in peripheral tissues
- Detect metabolic signals (glucose, fatty acids, gut hormones)
- Monitor mechanical stretch and distension
- Sample the gut luminal environment via nodose ganglion connections
- Project to the nucleus tractus solitarius (NTS) in the brainstem
- From NTS, signals reach hypothalamus, amygdala, and cortex

This pathway is the primary route by which peripheral inflammation induces central sickness behavior—even without inflammatory molecules crossing the blood-brain barrier.

14.6.2 Mechanisms of Aberrant Signaling

Peripheral Sensitization. Low-grade inflammation (from gut, liver, or other organs) could maintain vagal afferent activation. The inflammation might be subclinical—detectable by sensitive measures but not producing obvious organ dysfunction.

Central Sensitization. Repeated or prolonged vagal afferent activation could sensitize central circuits, such that normal afferent traffic is now interpreted as pathological. This is analogous to central sensitization in chronic pain.

Altered Vagal Tone. Autonomic dysfunction affecting vagal efferent (parasympathetic) tone might reflexively alter afferent sensitivity through local circuits.

Microglial Priming. Vagal afferent signals activate microglia in the brain. Primed microglia might amplify the central response to normal afferent traffic.

14.6.3 The Gut-Brain Axis Connection

The gut provides the largest source of vagal afferent input. Gut dysbiosis and intestinal barrier dysfunction, both documented in ME/CFS, would generate ongoing vagal danger signals:

- Bacterial translocation activates mucosal immune cells
- Inflammatory mediators stimulate vagal afferents
- Altered gut hormone secretion affects vagal signaling
- Mechanical sensitivity from dysmotility provides additional input

This provides a mechanism linking gut symptoms to systemic illness through vagal signaling.

14.6.4 Testable Predictions

1. Vagal afferent activity (measurable via heart rate variability metrics or direct recordings) should be altered in ME/CFS
2. Gut-directed interventions that reduce vagal afferent activation might improve systemic symptoms
3. Vagal nerve stimulation at specific parameters might reset aberrant signaling
4. Central markers of vagal afferent activation (NTS activity, microglial activation in relevant regions) should correlate with symptoms
5. Interrupting vagal signaling (e.g., with targeted anesthetics) might temporarily relieve symptoms

14.7 Tryptophan/Kynurenine Trap

? Open Question 6: Neurotoxic Kynurenine Dominance

Tryptophan metabolism sits at a critical junction: it can be converted to serotonin (mood, sleep, gut function) or shunted into the kynurenine pathway (immune regulation, NAD⁺ synthesis). Immune activation diverts tryptophan toward kynurenine via IDO and TDO enzymes.

The kynurenine pathway then branches: one arm produces neuroprotective kynurenic acid (KYNA), the other produces neurotoxic quinolinic acid (QUIN). What if ME/CFS involves persistent shunting toward kynurenine combined with preferential flux through the quinolinic acid branch?

This single metabolic disturbance could simultaneously explain:


- Depleted serotonin (mood disturbance, sleep dysfunction, gut dysmotility)
- Neuroinflammation and excitotoxicity (cognitive dysfunction, sensory sensitivities)
- Disrupted NAD⁺ synthesis (energy metabolism impairment)
- Immune dysregulation (kynurenines are immunomodulatory)

The NIH deep phenotyping study found significant tryptophan pathway abnormalities in ME/CFS patients, lending some empirical support to this direction.

14.7.1 Tryptophan Metabolism Overview


Tryptophan is an essential amino acid with two major metabolic fates:

Serotonin Pathway (~5% of tryptophan).



This pathway produces neurotransmitters essential for mood, sleep, cognition, and gut function.

Kynurenine Pathway (~95% of tryptophan).



The branch point is critical:

- **KYNA:** NMDA receptor antagonist, neuroprotective, anti-inflammatory
- **QUIN:** NMDA receptor agonist, neurotoxic, pro-inflammatory, generates ROS

14.7.2 The Trap Mechanism

Immune activation (IFN- γ , IL-6) strongly induces IDO, shunting tryptophan toward kynurenine. This is normally adaptive—it depletes tryptophan that pathogens need and generates immunomodulatory metabolites.

The “trap” occurs when:

1. IDO activation persists beyond the acute phase
2. Kynurenine preferentially flows toward QUIN rather than KYNA
3. QUIN causes neuroinflammation and oxidative stress
4. Neuroinflammation maintains cytokine production
5. Cytokines perpetuate IDO activation

This creates a self-sustaining loop where the pathway that should eventually suppress inflammation instead maintains it.

14.7.3 Consequences of the Trap

Serotonin Depletion. With tryptophan diverted to kynurenine, less is available for serotonin synthesis. This contributes to:

- Depressed mood (though different from primary depression)
- Sleep disturbances (melatonin is downstream of serotonin)
- Gut dysmotility (gut contains 90% of body's serotonin)
- Cognitive impairment (serotonin modulates cognition)

Quinolinic Acid Neurotoxicity. QUIN accumulation causes:

- NMDA receptor overactivation and excitotoxicity
- Oxidative stress and lipid peroxidation
- Astrocyte and microglial activation
- Blood-brain barrier disruption
- Direct neuronal damage

NAD⁺ Disruption. While QUIN eventually becomes NAD⁺, the pathway may be inefficient or the intermediate toxicity may outweigh benefits. Additionally, NAD⁺ consumption by inflammation-activated PARPs may exceed synthesis.

14.7.4 Testable Predictions

1. ME/CFS patients should show elevated QUIN:KYNA ratios in plasma and/or CSF
2. IDO expression/activity should be chronically elevated
3. Serotonin and melatonin levels should be reduced
4. Interventions blocking IDO or shifting the pathway toward KYNA might help
5. NAD⁺ precursor supplementation might be beneficial
6. The severity of kynurenine imbalance should correlate with specific symptoms

14.8 Cellular “Quorum Sensing” Dysfunction

? Open Question 7: Corrupted Intercellular Communication

Bacteria use quorum sensing to coordinate group behavior based on population density and environmental conditions. Human cells have analogous coordination systems: extracellular vesicles (exosomes), cell-free DNA, circulating metabolites, and cytokine networks create an “information field” that coordinates tissue and organ function.

What if a triggering event corrupts this intercellular communication system? Individual cells might function normally in isolation, but collective coordination breaks down.

The organism behaves as if under attack because the signaling environment says it should, even though no actual attack is occurring.

This would explain why individual laboratory tests often appear normal (cells function), why the syndrome is so diffuse (coordination affects everything), and why severity fluctuates unpredictably (the corrupted signaling creates chaotic dynamics). It also explains why so many different triggers can initiate ME/CFS—any sufficiently severe perturbation might corrupt the signaling landscape.

14.8.1 Intercellular Communication Systems

Human cells coordinate through multiple overlapping systems:

Extracellular Vesicles (EVs). Cells release vesicles containing:

- mRNAs and microRNAs that alter recipient cell gene expression
- Proteins that signal or directly affect recipient cell function
- Lipids that modulate membrane composition
- Metabolites that alter recipient cell metabolism

EV content changes based on the cell's state, creating a distributed signaling system.

Cell-Free DNA (cfDNA). Dying or stressed cells release DNA fragments that:

- Activate pattern recognition receptors (TLR9, cGAS-STING)
- Carry epigenetic marks reflecting their source
- Trigger inflammatory responses

Circulating Metabolome. The metabolites in blood create a “metabolic signature” that:

- Reflects overall metabolic state
- Directly affects cellular function throughout the body
- Changes rapidly with physiological state

Cytokine Networks. Beyond simple inflammation, cytokines form complex networks with:

- Positive and negative feedback loops
- Tissue-specific effects
- Temporal dynamics that carry information

14.8.2 Corruption Mechanisms

A severe perturbation could corrupt this signaling landscape by:

- Altering EV cargo in ways that persist after the trigger resolves
- Increasing cfDNA release, maintaining inflammatory signaling
- Shifting the circulating metabolome to a “sickness” signature
- Disrupting cytokine network dynamics
- Creating positive feedback loops that stabilize the corrupted state

Once corrupted, the signaling environment tells cells throughout the body that something is wrong, even if they individually function normally. This is analogous to bacteria receiving quorum signals indicating high population density and stress, even if the local environment is benign.

14.8.3 Why Standard Tests Miss This

Standard medical testing examines:

- Individual analytes (not network patterns)
- Static snapshots (not dynamics)
- Major parameters (not subtle signaling shifts)
- Isolated samples (not system-wide coordination)

A corruption in intercellular coordination might not show as any single abnormal value, only as altered patterns that require systems-level analysis to detect.

14.8.4 Testable Predictions

1. EV cargo analysis should reveal altered patterns in ME/CFS patients
2. cfDNA levels and characteristics might differ from controls
3. Metabolomic signatures should show consistent patterns that reflect the “corrupted” state
4. Network analysis of cytokines should reveal altered dynamics rather than just altered levels
5. The pattern of corruption might predict symptoms and treatment response

14.9 Purinergic Signaling Dysregulation

? Open Question 8: ATP as Pathological Danger Signal

ATP isn't just intracellular energy currency—extracellular ATP is a potent signaling molecule. P2X and P2Y purinergic receptors on immune cells, neurons, and other cell types respond to extracellular ATP as a danger signal, triggering inflammation, pain, and behavioral changes.

What if ME/CFS involves dysregulated purinergic signaling—either excessive ATP release, impaired extracellular ATP degradation, or sensitized purinergic receptors? Normal cellular activity releases “normal” amounts of ATP that now trigger aberrant immune and pain responses.

Exercise dramatically increases extracellular ATP release. If purinergic receptors are sensitized or ATP clearance is impaired, exercise would trigger massive inappropriate danger signaling, explaining the delayed and prolonged nature of post-exertional malaise. This also connects to the pain hypersensitivity, immune activation, and autonomic dysfunction seen in ME/CFS.

14.9.1 Purinergic Signaling Biology

Extracellular ATP and its metabolites (ADP, AMP, adenosine) signal through two receptor families:

P2X Receptors (ion channels).

- P2X1-7 subtypes with different distributions and properties
- P2X7 is particularly important: high ATP threshold, immune activation
- Activation causes cation influx, including Ca^{2+}
- P2X7 activation triggers NLRP3 inflammasome

P2Y Receptors (G-protein coupled).

- P2Y1-14 subtypes responding to ATP, ADP, UTP, UDP
- Mediate diverse signaling cascades
- Important in platelet activation, vasodilation, neurotransmission

Extracellular ATP is normally rapidly degraded by ectonucleotidases (CD39, CD73), keeping concentrations low.

14.9.2 Dysregulation Mechanisms

Excessive ATP Release. Under stress, damaged or dying cells release ATP. In ME/CFS:

- Chronic cellular stress might maintain elevated ATP release
- Exercise-induced microtrauma releases ATP
- Autonomic activation affects ATP release
- Mitochondrial dysfunction might alter ATP handling

Impaired ATP Clearance. CD39 and CD73 expression/activity might be reduced:

- Inflammatory cytokines alter ectonucleotidase expression
- Oxidative stress can damage these enzymes
- Genetic variants affect ectonucleotidase function

Receptor Sensitization. P2X receptors can become sensitized by:

- Prolonged exposure to low ATP concentrations
- Inflammatory mediators that alter receptor function
- Changes in membrane composition affecting receptor signaling

14.9.3 Consequences of Purinergic Dysregulation

Chronic Inflammation. P2X7 activation:

- Triggers NLRP3 inflammasome assembly
- Drives IL-1 β and IL-18 release
- Maintains chronic low-grade inflammation

Pain Sensitization. Purinergic receptors on sensory neurons:

- P2X3 mediates pain signaling
- Sensitization lowers pain thresholds
- Contributes to widespread pain and hyperalgesia

Neuroinflammation. Brain P2X7 on microglia:

- Drives microglial activation
- Promotes neuroinflammatory state
- Contributes to cognitive dysfunction

Autonomic Effects. Purinergic signaling in cardiovascular regulation:

- Affects vasodilation and vasoconstriction
- Modulates heart rate
- Contributes to orthostatic intolerance

14.9.4 Testable Predictions

1. Extracellular ATP levels should be elevated, especially after exertion
2. Ectonucleotidase expression/activity should be reduced
3. P2X receptor expression or sensitivity should be altered
4. P2X7 antagonists might reduce inflammation and symptoms
5. Genetic variants in purinergic pathway genes might associate with ME/CFS risk
6. Post-exertional malaise severity should correlate with exercise-induced ATP release

14.10 Redox Compartment Collapse

? Open Question 9: Loss of Redox Boundaries

Cells maintain distinct redox environments in different compartments: the cytosol is relatively reducing, the mitochondrial matrix more oxidizing, the ER oxidizing (for protein folding), and the extracellular space oxidizing. These gradients are actively maintained and essential for compartment-specific chemistry.

What if ME/CFS involves collapse of these redox boundaries? Normally compartmentalized reactive oxygen and nitrogen species might leak between compartments, creating widespread dysfunction:

- ER stress and protein misfolding (disrupted ER redox)
- Mitochondrial dysfunction (disrupted mitochondrial redox)
- Aberrant cell signaling (many signaling pathways are redox-sensitive)
- Oxidative damage to proteins, lipids, and DNA

This would explain the oxidative stress markers observed in ME/CFS without requiring a specific source of ROS—the problem is boundary failure rather than excess production. It would also explain why antioxidant supplementation shows inconsistent results: the problem isn't total antioxidant capacity but compartment-specific redox control.

14.10.1 Cellular Redox Compartments

Different cellular compartments maintain distinct redox states:

Cytosol. Relatively reducing (GSH:GSSG $\approx$ 100:1):

- Maintained by NADPH-dependent reductases
- Supports reductive biosynthesis
- Most enzymes optimized for reducing environment

Mitochondrial Matrix. More oxidizing (GSH:GSSG $\approx$ 30:1):

- ETC generates ROS as byproduct
- Contains its own antioxidant systems
- Redox state regulates metabolism

Endoplasmic Reticulum. Oxidizing (GSH:GSSG $\approx$ 3:1):

- Required for disulfide bond formation
- Ero1/PDI systems maintain oxidizing environment
- Critical for protein folding

Extracellular Space. Oxidizing:

- Different redox chemistry than intracellular
- Proteins contain stable disulfides
- Thiol-disulfide exchange used for signaling

14.10.2 Boundary Maintenance

These compartments are maintained by:

- Selective permeability of membranes to redox-active species
- Active transport systems for glutathione and other redox buffers
- Compartment-specific antioxidant enzymes
- Regeneration systems (NADPH, thioredoxin reductase)

14.10.3 Consequences of Boundary Collapse

ER Stress. If the ER becomes too reducing or too oxidizing:

- Protein folding fails
- Unfolded protein response (UPR) activates
- Chronic UPR leads to inflammation and cell death

Mitochondrial Dysfunction. Altered mitochondrial redox:

- Disrupts ETC function
- Affects metabolic enzyme activity
- Triggers mitochondrial permeability transition

Signaling Disruption. Many signaling pathways use redox as a switch:

- NF- κ B activation is redox-sensitive
- Kinase/phosphatase balance depends on redox state
- Calcium signaling is modulated by redox

Why Antioxidants Don't Help. Systemic antioxidant supplementation:

- Doesn't address compartment-specific problems
- May actually worsen some compartment imbalances
- Cannot restore proper boundaries

14.10.4 Testable Predictions

1. Compartment-specific redox indicators should show altered ratios in ME/CFS
2. Markers of ER stress (BiP, CHOP, spliced XBP1) should be elevated
3. Mitochondrial redox state should differ from controls
4. Interventions targeting specific compartment redox might help where global antioxidants fail
5. The specific pattern of compartment disruption might predict symptoms

14.11 Metabolic Memory and Epigenetic Lock

? Open Question 10: Stable Epigenetic Reprogramming

Cells can retain metabolic states through epigenetic modifications—DNA methylation, histone modifications, and chromatin remodeling that persist through cell division. This “metabolic memory” normally serves homeostasis but could become pathological.

A sufficiently severe metabolic insult (infection, prolonged stress) might create stable epigenetic changes that persist even after the trigger resolves. Immune cells, neurons, muscle cells, and others become “programmed” to maintain the sick state, with their gene expression locked into patterns appropriate for acute illness.

This would explain why ME/CFS is so persistent, why duration correlates with prognosis (longer duration means more stable epigenetic changes), and why early treatment shows better outcomes (intervention before epigenetic stabilization). It also explains why

so many different body systems are affected—if the epigenetic changes occur in multiple cell types during the initial insult, all those systems remain locked.

Importantly, epigenetic changes are potentially reversible, unlike genetic mutations. This provides hope for intervention while explaining why simple removal of triggers doesn't restore health.

14.11.1 Epigenetic Mechanisms

DNA Methylation. 5-methylcytosine at CpG sites:

- Generally silences gene expression
- Patterns are maintained through cell division
- Can be stable for years but also dynamically regulated
- Altered by inflammation, oxidative stress, metabolic state

Histone Modifications. Acetylation, methylation, phosphorylation of histones:

- Affect chromatin accessibility
- Can be activating or repressing
- Some marks are very stable; others are dynamic
- Metabolic intermediates are cofactors (acetyl-CoA, SAM, NAD+)

Chromatin Remodeling. Large-scale changes in chromatin organization:

- Affect which genes are accessible
- Can be inherited through cell division
- Respond to signaling and metabolic state

14.11.2 Metabolic Memory in Disease

Metabolic memory has been documented in:

- **Diabetes:** Periods of poor glycemic control cause lasting epigenetic changes that maintain complications even after glucose is normalized
- **Cardiovascular disease:** Inflammatory episodes create epigenetic “scars” that maintain vessel dysfunction
- **Cancer:** Epigenetic reprogramming is central to oncogenesis
- **Immune memory:** Innate immune cells (monocytes, macrophages) can be epigenetically “trained” by prior exposures

14.11.3 Application to ME/CFS

The initial trigger (infection, stress) creates a metabolic/inflammatory state that:

1. Alters the availability of epigenetic cofactors (SAM, acetyl-CoA, NAD+)
2. Activates enzymes that write epigenetic marks (DNMTs, HATs, HMTs)
3. Creates gene expression patterns appropriate for the acute phase
4. If the acute phase is severe or prolonged enough, these patterns stabilize
5. Stabilized patterns persist even after the trigger resolves
6. Multiple cell types are affected, creating multi-system disease

The “lock” is not a single epigenetic change but a network of changes across cell types that maintain each other.

14.11.4 Why Duration Matters

- Epigenetic changes become more stable over time
- More cell divisions = more opportunity for stabilization
- The network of changes becomes more interconnected
- Compensatory mechanisms may also become epigenetically fixed

This explains the clinical observation that early intervention improves outcomes and that long-duration patients are hardest to treat.

14.11.5 Testable Predictions

1. ME/CFS patients should show distinct DNA methylation patterns in relevant cell types
2. Histone modification patterns should differ from controls
3. Disease duration should correlate with epigenetic change stability
4. Patients who recover should show reversal of epigenetic changes
5. Epigenetic modifying agents might provide therapeutic benefit
6. The specific epigenetic signature might predict symptom patterns or treatment response

14.12 Circadian-Metabolic Desynchronization

? Open Question 11: Peripheral Clock Misalignment

The body maintains circadian clocks in virtually every tissue, coordinated by the master clock in the suprachiasmatic nucleus (SCN). These clocks regulate metabolism, immune function, hormone release, and cellular processes in a time-dependent manner.

What if ME/CFS represents desynchronization of peripheral clocks from the master clock and from each other? The liver clock, muscle clock, immune clock, and brain clocks

might all be running on different schedules, creating constant metabolic “jet lag.”

This would explain the profound sleep dysfunction in ME/CFS (sleep architecture is clock-dependent), the hormone abnormalities (hormone release is clock-gated), why symptoms fluctuate unpredictably (different clocks moving in and out of phase), and why patients often report feeling better at unusual hours. The autonomic dysfunction might represent the body’s failed attempt to reconcile conflicting clock signals.

14.12.1 The Circadian System

Master Clock (SCN). The suprachiasmatic nucleus:

- Contains ~20,000 neurons with autonomous rhythms
- Entrained by light via retinohypothalamic tract
- Sends timing signals throughout the body
- Coordinates peripheral clocks

Peripheral Clocks. Found in virtually every tissue:

- Same core clock genes (CLOCK, BMAL1, PER, CRY)
- Regulate tissue-specific gene expression rhythms
- 10-30% of tissue transcriptome is rhythmic
- Normally synchronized by SCN signals

Synchronization Signals. The SCN coordinates peripheral clocks via:

- Hormones (cortisol, melatonin)
- Autonomic nervous system
- Body temperature rhythms
- Feeding/fasting signals

14.12.2 Consequences of Desynchronization

When peripheral clocks become misaligned:

Metabolic Dysfunction. Liver and muscle clocks regulate:

- Glucose and lipid metabolism
- Mitochondrial function
- Nutrient sensing

Misalignment causes metabolic inefficiency and abnormal fuel utilization.

Immune Dysfunction. Immune cell clocks regulate:

- Cytokine production patterns
- Immune cell trafficking
- Inflammatory responses

Misalignment causes immune dysregulation.

Hormone Dysfunction. Endocrine clocks regulate:

- Cortisol rhythm (disrupted in ME/CFS)
- Melatonin secretion (affects sleep)
- Thyroid hormone patterns

Misalignment causes hormonal chaos.

Sleep Dysfunction. Sleep is gated by:

- SCN timing signals
- Peripheral metabolic signals
- Temperature rhythms

Misalignment causes unrefreshing sleep even with normal sleep duration.

14.12.3 How Desynchronization Might Occur

- **Infection:** Inflammatory cytokines disrupt clock gene expression
- **Autonomic dysfunction:** Impairs SCN → peripheral signaling
- **Cortisol dysregulation:** Key synchronizing hormone is abnormal
- **Activity restriction:** Loss of activity/feeding rhythms that reinforce clocks
- **Light exposure changes:** Altered light patterns during illness

Once desynchronized, the different clocks may stabilize at different phases, resisting resynchronization.

14.12.4 Testable Predictions

1. Clock gene expression in peripheral blood cells should show altered rhythms
2. Different tissues/cell types might show different phase relationships
3. Chronotherapy (timing treatments to clock phases) might improve efficacy
4. Light therapy and time-restricted feeding might help resynchronize clocks
5. Melatonin and other chronobiotics might provide benefit
6. Symptom patterns might correlate with clock phase relationships

14.13 GPCR Autoantibody-Driven Dysfunction

This section has moved from purely speculative to evidence-supported. Multiple studies have documented G-protein coupled receptor (GPCR) autoantibodies in ME/CFS, and treatment trials targeting these autoantibodies have shown promising results.

14.13.1 Established Evidence

Foundational Cohort Studies

The Charité Berlin group established GPCR autoantibodies as a significant finding in ME/CFS:

- **Loebel et al. 2016 [71]**: In 268 ME/CFS patients vs. 108 controls, 29.5% of patients had elevated antibodies against ≥ 1 muscarinic (M) or β -adrenergic receptor. Antibodies against β_2 , M3, and M4 receptors were significantly elevated vs. controls.
- **Sotzny/Freitag et al. 2021 [208]**: Autoantibody levels correlated with symptom severity—fatigue, muscle pain, cognitive impairment, and GI symptoms in infection-triggered ME/CFS. First demonstration of dose-response relationship.
- **Bynke et al. 2020 [72]**: Swedish validation in two independent cohorts found 79–91% of ME patients had ≥ 1 elevated antibody vs. 25% of controls. Critically: **no autoantibodies detected in CSF**, suggesting peripheral origin rather than intrathecal production.

Treatment Trial Evidence

- **Immunoabsorption pilot (Scheibenbogen 2018) [147]**: 10 post-infectious ME/CFS patients with elevated β_2 antibodies received 5 immunoabsorption sessions. 70% showed rapid improvement during treatment; 30% sustained improvement at 6–12 months.
- **Immunoabsorption cohort (Stein et al. 2024) [118]**: 20 post-COVID ME/CFS patients with elevated β_2 -AR autoantibodies. IgG reduced 79%, autoantibodies reduced 77%. **70% responders** with ≥ 10 point SF-36 Physical Function increase. Benefits sustained to 6 months. This represents the *strongest evidence to date* for autoantibody-mediated pathophysiology.
- **Daratumumab pilot (Fluge et al. 2025) [9]**: Anti-CD38 therapy targeting plasma cells (the antibody factories). 10 female ME/CFS patients; **60% showed marked improvement**. SF-36 PF increased from 25.9 to 55.0 ($p=0.002$). Responders achieved near-normal function (SF-36 scores 80–95). Low baseline NK-cell count predicted non-response.
- **BC007 case report (Hohberger 2021) [10]**: DNA aptamer neutralizing GPCR autoantibodies produced dramatic improvement in a Long COVID patient: fatigue normalized, brain fog resolved, retinal microcirculation improved within hours. However, the subsequent Phase II trial failed to show superiority over placebo at the population level.

Methodological Controversy

Important caveats exist regarding GPCR autoantibody testing:

- **POTS replication failure (2022) [210]:** 116 POTS patients vs. 81 controls showed *no differences* in ELISA-derived GPCR autoantibody concentrations. 98.3% of POTS patients and 100% of controls had α_1 -adrenergic receptor antibodies above threshold. The authors concluded CellTrend ELISAs “have no diagnostic value for POTS.”
- **Functional vs. binding assays:** The positive studies largely used CellTrend ELISAs (binding assays), while the cardiomyocyte bioassay (measuring functional antibody activity) may be more specific but is not commercially available.
- **Conflict of interest:** CellTrend holds a patent for β -adrenergic receptor antibodies in CFS diagnosis, jointly with Charité.

Despite methodological concerns, the *treatment* evidence is compelling: if autoantibody removal (immunoadsorption) and autoantibody-producing cell depletion (daratumumab) produce clinical improvement, the autoantibodies are likely pathogenic regardless of assay limitations.

⚠ **Limitation 2: GPCR Autoantibody Evidence: Methodological Controversy and Open-Label Designs**

Despite promising treatment signals, the GPCR autoantibody field faces unresolved methodological issues that limit the strength of current conclusions:

- The CellTrend ELISA (used in most positive studies) has been questioned by an independent replication failure in 116 POTS patients [210], where 98.3% of patients *and* 100% of controls exceeded the positivity threshold—suggesting the assay may lack specificity.
- CellTrend holds a joint patent with Charité for β -adrenergic receptor antibodies in CFS diagnosis, creating a potential conflict of interest that has not been addressed by independent validation using alternative assay platforms.
- All treatment trials to date (immunoadsorption n=10, n=20; daratumumab n=10) are open-label without placebo control. The 60–70% response rates cannot be distinguished from placebo effect, regression to the mean, or natural fluctuation without randomised controlled trials. The BC007 case report [10] was similarly uncontrolled.
- The BC007 Phase II trial—the only placebo-controlled study of autoantibody-targeted therapy—failed to show superiority over placebo, directly challenging the therapeutic hypothesis despite the positive case report.

14.13.2 Speculative Hypotheses Emerging from GPCR Research

~ Hypothesis 1: The Plasma Cell Sanctuary

The daratumumab success vs. rituximab failure reveals a critical insight: B cells (CD20⁺) are precursors; plasma cells (CD38⁺) are the factories. Long-lived plasma cells can survive for *decades* in bone marrow and gut niches, continuously secreting autoantibodies without B cell replenishment.

Hypothesis: ME/CFS is maintained by “sanctuary” plasma cells that escaped B-cell depletion:

1. Initial trigger generates autoreactive B cells
2. Some differentiate into long-lived plasma cells in survival niches
3. These plasma cells produce GPCR autoantibodies indefinitely
4. Rituximab depletes B cells but not established plasma cells—autoantibody production continues
5. Daratumumab directly kills plasma cell factories, stopping production

Evidence level: Moderate. The 8–9 month delay before maximum daratumumab benefit supports this (existing autoantibodies must decay after factory elimination).

Therapeutic implication: Combining immunoadsorption (remove existing antibodies) with daratumumab (eliminate factories) might produce faster, more complete responses.

~ Hypothesis 2: GPCR Autoantibody-Endothelial Cascade

GPCR autoantibodies may exert their effects primarily through endothelial dysfunction:

1. β_2 -adrenergic receptor autoantibodies impair endothelial vasodilation
2. Muscarinic receptor autoantibodies disrupt endothelial NO production
3. Impaired vasodilation → tissue hypoperfusion
4. Hypoperfusion → mitochondrial dysfunction
5. Mitochondrial dysfunction → cellular energy crisis → symptoms

The BC007 case report supports this: retinal microcirculation improved within *hours* of autoantibody neutralization [10]—faster than any cellular recovery could explain. The vascular effect was immediate.

Evidence level: Low-Moderate. Mechanistically plausible; BC007 microcirculation data supportive; needs direct testing.

Therapeutic implication: Vascular-supportive therapies (L-citrulline, statins) might synergize with autoantibody removal.

~ Hypothesis 3: Autoantibody-Monocyte Inflammation Loop

A 2025 preprint [209] demonstrated that GPCR autoantibodies drive monocyte dysfunction in post-COVID ME/CFS, causing elevated MIP-1 δ , PDGF-BB, and TGF- β 3. This

suggests autoantibodies don't just block receptors—they actively drive inflammation:

1. GPCR autoantibodies bind monocyte surface receptors
2. Binding triggers inflammatory cytokine production
3. Cytokines cause systemic inflammation and tissue damage
4. Tissue damage generates more autoantigen exposure
5. Cycle perpetuates autoantibody production

Evidence level: Low-Moderate (single preprint, not yet replicated).

Therapeutic implication: Monocyte-targeted therapies might complement autoantibody removal.

? Open Question 12: Why Only 60% Respond?

The daratumumab trial showed 60% marked improvement and 40% non-response. What distinguishes responders from non-responders?

Potential factors:

- **Autoantibody presence:** Non-responders may have different (non-GPCR) autoantibodies, or non-autoimmune ME/CFS
- **NK cell status:** Low baseline NK cells predicted non-response (immune dysregulation pattern)
- **Illness duration:** Longer illness may cause irreversible downstream damage
- **Plasma cell location:** Some sanctuary sites may be less accessible to daratumumab

Identifying responder biomarkers is critical for treatment personalization.

14.13.3 Undocumented Biological Phenomena

Based on the GPCR autoantibody literature, several biological phenomena have never been directly examined:

1. **Bone marrow plasma cell populations:** Do ME/CFS patients have expanded long-lived plasma cells producing GPCR autoantibodies? No bone marrow studies have examined this.
2. **Gut-associated plasma cells:** The gut wall contains plasma cell niches. Do these contribute to autoantibody production in ME/CFS?
3. **Autoantibody epitope specificity:** Which specific receptor epitopes do ME/CFS autoantibodies target? Epitope mapping might predict functional effects.
4. **Functional vs. binding antibody correlation:** How well do ELISA-detected antibodies correlate with functional bioassay results in the same patients?
5. **Autoantibody fluctuation with symptoms:** Do autoantibody titers change during PEM episodes or remissions?
6. **GPCR receptor internalization:** Do autoantibodies cause receptor downregulation through chronic stimulation?

14.13.4 Evidence Assessment Summary

Finding	Evidence Level	Notes
GPCR autoantibodies elevated in ME/CFS	Moderate	Multiple cohorts; replication concerns
Symptom correlation with titers	Moderate	Sotzny 2021; needs replication
Immunoabsorption efficacy	Moderate-High	Lancet 2024; no placebo control
Daratumumab efficacy	Moderate	60% response; open-label
BC007 efficacy	Low	Case reports positive; Phase II failed
Peripheral (not CNS) origin	Moderate	No CSF autoantibodies (Bynke 2020)
CellTrend assay specificity	Controversial	POTS study questions diagnostic value

Table 14.2: Evidence assessment for GPCR autoantibody findings in ME/CFS

Overall assessment: GPCR autoantibody-driven ME/CFS represents the most therapeutically promising hypothesis currently under investigation. The evidence is sufficient to justify clinical trials and, for carefully selected patients with documented autoantibodies, consideration of autoantibody-targeted treatment under specialist supervision.

14.14 Ion Channel Autoimmunity

? Open Question 13: Channelopathy from Autoantibodies

Beyond GPCR autoantibodies (Section 14.13), what about autoantibodies targeting ion channels—sodium, calcium, or potassium channels that regulate cellular excitability?

Depending on the target and antibody effect (blocking vs. activating), this could cause:

- Neuronal hyperexcitability or inexcitability
- The “wired but tired” phenomenon (simultaneous overstimulation and exhaustion)
- Sensory hypersensitivities (lowered thresholds for sensory neuron firing)
- Autonomic dysfunction (altered autonomic neuron excitability)
- Muscle weakness and fatigue (altered muscle cell excitability)
- Cardiac symptoms (altered cardiac ion channel function)

Ion channel autoimmunity is established in other conditions (myasthenia gravis, Lambert-Eaton syndrome, autoimmune encephalitis). The multi-system nature of ME/CFS could reflect antibodies targeting channels expressed across many tissues.

★ Achievement 1: TRPM3: From Speculation to Evidence

The ion channel hypothesis has moved from speculation to evidence with the 2026 multi-site validation of TRPM3 dysfunction in ME/CFS [187]. Researchers at Griffith University demonstrated that TRPM3, a calcium-permeable ion channel in immune cells, functions abnormally in ME/CFS patients. This finding was replicated across independent laboratories 4,000 km apart, meeting rigorous standards for scientific reproducibility.

TRPM3 dysfunction provides concrete evidence that ME/CFS involves measurable ion channel pathology. Whether this reflects autoimmune targeting, post-infectious modification, or other mechanisms remains to be determined, but the “channelopathy hypothesis” is no longer purely speculative—it has empirical support. See Section 14.21 for detailed exploration of TRPM3-related hypotheses.

14.14.1 Ion Channels in Physiology

Ion channels are membrane proteins that control electrical excitability:

Sodium Channels (Na_v).

- Generate action potentials in neurons and muscle
- $\text{Na}_v1.7$, 1.8, 1.9 in pain pathways
- $\text{Na}_v1.5$ in cardiac muscle
- Antibody effects: altered excitability, pain sensitization, arrhythmias

Calcium Channels (Ca_v).

- Regulate neurotransmitter release, muscle contraction, gene expression
- P/Q-type ($\text{Ca}_v2.1$) targeted in Lambert-Eaton syndrome
- L-type in cardiac and smooth muscle
- Antibody effects: weakness, autonomic dysfunction, CNS symptoms

Potassium Channels (K_v).

- Regulate resting potential and repolarization
- VGKC-complex antibodies cause autoimmune encephalitis
- $\text{K}_v1.1$ -1.6 in CNS and PNS
- Antibody effects: hyperexcitability, seizures, cognitive impairment

14.14.2 Ion Channel Autoimmunity Precedents

- **Myasthenia gravis:** Anti-acetylcholine receptor antibodies cause neuromuscular weakness
- **Lambert-Eaton:** Anti- $\text{Ca}_v2.1$ antibodies cause weakness, autonomic symptoms
- **Autoimmune encephalitis:** Anti-VGKC, anti-NMDAR antibodies cause cognitive/neurological symptoms
- **Neuromyotonia:** Anti-VGKC antibodies cause muscle hyperexcitability

14.14.3 Potential ME/CFS Relevance

The symptom cluster of ME/CFS could result from antibodies against multiple channel types:

“Wired but Tired.”

- Activating antibodies → hyperexcitability → overstimulation → “wired”
- Excessive firing → energy depletion → exhaustion → “tired”
- Or blocking antibodies in some circuits, activating in others

Sensory Sensitivities.

- Lower firing thresholds in sensory neurons
- Enhanced pain, light, sound, smell sensitivity

Autonomic Dysfunction.

- Altered excitability in autonomic ganglia
- Abnormal baroreceptor responses
- Disrupted heart rate variability

14.14.4 Testable Predictions

1. Comprehensive ion channel autoantibody panels should reveal positivity in ME/CFS subsets
2. Patient IgG transferred to animal models might reproduce symptoms
3. Plasmapheresis or IVIG might help antibody-positive patients
4. The specific channels targeted should predict symptom patterns
5. Immunomodulation might provide more durable benefit than symptomatic treatment

14.15 Ferroptosis Susceptibility

? Open Question 14: Increased Vulnerability to Iron-Dependent Cell Death

Ferroptosis is a recently characterized form of regulated cell death distinct from apoptosis, driven by iron-dependent lipid peroxidation. Cells with high metabolic rates and lipid content (neurons, cardiomyocytes) are particularly vulnerable.

What if ME/CFS involves increased susceptibility to ferroptosis? Iron dysregulation combined with oxidative stress and membrane lipid abnormalities would create conditions favoring ferroptotic cell death. Cells might not die en masse, but exist in a chronic state at the edge of ferroptosis, with ongoing low-grade cell loss and replacement.

This would explain the lipid abnormalities observed in ME/CFS, the oxidative stress markers, and why iron supplementation can sometimes worsen symptoms. It also explains the particular vulnerability of high-energy tissues like brain, heart, and muscle. The body's attempt to limit ferroptosis might involve sequestering iron (explaining common low ferritin despite adequate intake) and suppressing metabolism (back to the "safe mode" concept).

14.15.1 Ferroptosis Biology

Ferroptosis is characterized by:

- Iron-dependent lipid peroxidation
- Distinct from apoptosis, necrosis, autophagy
- Requires polyunsaturated fatty acids in membranes
- Inhibited by GPX4 (glutathione peroxidase 4)
- Promoted by iron accumulation and oxidative stress

The ferroptosis pathway:

1. Iron catalyzes Fenton reaction → hydroxyl radical
2. Hydroxyl radical attacks membrane PUFAs → lipid peroxidation
3. Lipid peroxides propagate → membrane damage
4. GPX4 normally reduces lipid peroxides → protection
5. GPX4 depletion (low glutathione) → ferroptosis execution

14.15.2 ME/CFS Risk Factors for Ferroptosis

Iron Dysregulation.

- Inflammation causes iron redistribution
- Iron can accumulate in stressed tissues
- Low serum iron doesn't mean low tissue iron

Oxidative Stress.

- Documented in ME/CFS
- Provides initiating radicals
- Depletes glutathione → reduces GPX4 activity

Lipid Abnormalities.

- Altered membrane PUFA composition documented
- More oxidizable PUFAs = more vulnerable membranes

High-Energy Tissue Vulnerability.

- Neurons: high lipid content, high metabolic rate
- Heart: high iron, high oxygen flux
- Muscle: high metabolic demand during exercise

14.15.3 Sublethal Ferroptosis

Rather than cell death, ME/CFS might involve cells existing in a chronic “pre-ferroptotic” state:

- Ongoing low-level lipid peroxidation
- Constant antioxidant demand
- Membrane damage requiring repair
- Signaling dysfunction from altered membrane lipids
- Metabolic suppression to reduce ferroptosis risk

This “edge of ferroptosis” state would:

- Create constant oxidative stress markers
- Make cells vulnerable to any additional stress
- Explain why pushing causes crashes (exercise increases iron, oxygen, radicals)
- Explain why antioxidants help some patients

14.15.4 Testable Predictions

1. Lipid peroxidation markers (MDA, 4-HNE) should be elevated
2. GPX4 activity might be reduced or compensatorily elevated
3. Iron distribution should be altered in relevant tissues
4. Ferroptosis inhibitors might provide benefit
5. Iron supplementation should be risky, especially during crashes
6. The tissues most affected should be those most vulnerable to ferroptosis

14.16 Integrated Hypothesis: The Multi-Lock Trap

The hypotheses above are not mutually exclusive; indeed, the most compelling model for ME/CFS pathogenesis may involve multiple mechanisms operating simultaneously and reinforcing each other. We propose an integrated “multi-lock trap” hypothesis that attempts to explain the key features of ME/CFS: post-viral onset, persistence despite apparent resolution of the trigger, post-exertional malaise, multi-system involvement, and treatment resistance.

14.16.1 Phase 1: Triggering Event

An initial insult—typically viral infection, but potentially severe stress, trauma, or other immune-activating event—activates the evolutionarily conserved “sickness behavior” program. This is a normal, adaptive response involving:

- Metabolic downregulation (reduced mitochondrial activity, shifted fuel utilization)
- Immune activation and inflammatory cytokine production
- Behavioral changes (fatigue, social withdrawal, reduced activity)
- Tryptophan shunting toward kynurenine pathway
- Catecholamine conservation

In most individuals, this program disengages once the threat resolves. In ME/CFS-susceptible individuals, the program becomes “locked” through multiple overlapping mechanisms.

14.16.2 Phase 2: Lock Establishment

Several “locks” establish themselves during or shortly after the acute phase:

Epigenetic Lock. The severe metabolic stress creates stable epigenetic modifications in immune cells, neurons, muscle cells, and other tissues. Gene expression patterns appropriate for acute illness become fixed through DNA methylation and histone modifications. These changes persist through cell division, propagating the sick state even as acute inflammation resolves.

Autoimmune Lock. The inflammatory environment, possibly combined with molecular mimicry from the triggering pathogen, generates autoantibodies against self-proteins—G-protein coupled receptors, ion channels, or other cellular machinery. These autoantibodies create ongoing dysfunction independent of the original trigger. HERV reactivation during the acute phase may contribute immunogenic self-antigens.

Metabolic Lock. Tryptophan/kynurenine pathway dysregulation becomes self-perpetuating: inflammatory cytokines activate IDO, shunting tryptophan toward kynurenine; quinolinic acid accumulation causes neuroinflammation and oxidative stress; neuroinflammation maintains cytokine production, perpetuating IDO activation. Similar vicious cycles may establish in other metabolic pathways (lactate compartmentalization, purinergic signaling).

Signaling Lock. Purinergic receptors become sensitized, vagal afferents develop persistent danger signaling, or cellular quorum sensing becomes corrupted. The body's communication systems now interpret normal physiological states as pathological.

Structural Lock. Glymphatic impairment, circadian desynchronization, or redox compartment collapse creates physical or temporal barriers to normal function that resist simple correction.

14.16.3 Phase 3: Trap Maintenance

Once multiple locks are established, the system becomes trapped in a stable pathological state. Each lock reinforces the others:

- Epigenetic changes maintain cells in a “sickness program” gene expression state
- Autoantibodies cause ongoing receptor/channel dysfunction
- Metabolic pathway dysregulation depletes essential intermediates while accumulating toxic ones
- Aberrant signaling maintains central nervous system perception of threat
- Structural/temporal disruptions prevent normal clearing and cycling

Attempting to force the system out of this state (through exertion, stimulants, or willpower) triggers defensive responses: the body “detects” that something is trying to override its protective program during perceived danger, and responds by intensifying the sickness response—post-exertional malaise.

14.16.4 Why Recovery Is Rare

For recovery to occur, *all* locks must be released, or at least enough of them that the remaining ones cannot maintain the trapped state. Treatments targeting only one mechanism fail because the others maintain the trapped state. This explains why:

- Immunomodulation sometimes helps but rarely cures (addresses autoimmune lock only)
- Metabolic supplements show limited efficacy (addresses metabolic lock only)
- Behavioral approaches fail or cause harm (don't address any locks, may strengthen them)
- Early intervention shows better outcomes (fewer locks have stabilized)
- Spontaneous recovery is rare and unpredictable (requires spontaneous release of multiple locks)

- Some patients respond to treatments others don't (different lock combinations)

14.16.5 Testable Predictions

This integrated hypothesis generates several testable predictions:

1. ME/CFS patients should show epigenetic signatures distinct from healthy controls and from recovered patients, potentially with duration-dependent stabilization
2. Multiple autoantibody classes should be present, not just one type
3. Kynurenine pathway metabolites should show specific patterns (elevated quino-
linic:kynurenic ratio)
4. Purinergic receptor expression or sensitivity should differ from controls
5. Combined treatments targeting multiple locks should show synergistic efficacy compared to monotherapies
6. Patients who recover should show reversal of epigenetic changes, autoantibody clearance, or both
7. Disease duration should correlate with epigenetic change stability and treatment resistance
8. Patient subgroups might be identifiable by which locks predominate

⚡ **Limitation 3: Multi-Lock Trap Model: Unfalsifiable Integration Risk**

The multi-lock trap model is a conceptual framework, not a falsifiable hypothesis in its current form. Key epistemic boundaries:

- The model can accommodate virtually any finding by positing an additional “lock”—this explanatory flexibility is a weakness, not a strength, as it makes the model difficult to disprove.
- No study has simultaneously measured epigenetic, autoimmune, metabolic, signalling, and structural “locks” in the same ME/CFS cohort; the integration is theoretical, not data-driven.
- The prediction that “combined treatments targeting multiple locks should show synergistic efficacy” has not been tested; it is equally possible that the locks are independent phenomena requiring independent treatment, or that addressing a single upstream driver resolves multiple downstream manifestations.
- The model's value lies in organising research priorities and treatment strategies, not in establishing causal architecture.

14.16.6 Therapeutic Implications

If the multi-lock model is correct, effective treatment would require simultaneously addressing multiple mechanisms:

- **Epigenetic modifiers:** Agents that can reverse pathological epigenetic programming (HDAC inhibitors, DNA demethylating agents, or lifestyle interventions that affect the epigenome)
- **Autoantibody reduction:** Plasmapheresis, rituximab, IVIG, or tolerization approaches
- **Metabolic pathway correction:** Targeted supplementation to restore normal flux through kynurenine and other pathways; NAD⁺ precursors; specific nutrient support
- **Signaling normalization:** Purinergic receptor antagonists, vagal nerve modulation, low-dose naltrexone (affects multiple signaling systems)
- **Structural/temporal restoration:** Addressing craniocervical issues, chronotherapy for circadian resynchronization, targeted redox support
- **Pacing and energy management:** Preventing exertion-triggered lock reinforcement while other interventions work

The timing and sequencing of interventions may matter: some locks may need to be addressed before others become accessible. For example, reducing autoantibodies might be necessary before epigenetic interventions can take effect.

14.16.7 Research Directions

This model suggests several research priorities:

1. **Comprehensive phenotyping:** Assessing each patient for multiple lock types to enable personalized treatment
2. **Combination therapy trials:** Testing whether multi-target approaches show synergy
3. **Longitudinal tracking:** Following lock status over time to understand disease progression and treatment effects
4. **Early intervention studies:** Testing whether aggressive early treatment can prevent lock stabilization
5. **Recovery studies:** Detailed analysis of the rare patients who recover to understand which locks released and how

14.17 Speculative Cross-Disease Connections

ME/CFS shares features with numerous other conditions. These overlaps may reflect shared mechanisms, common susceptibility factors, or convergent pathophysiology. This section explores speculative connections that might illuminate ME/CFS pathogenesis.

14.17.1 The Post-Infectious Syndrome Cluster

ME/CFS belongs to a family of post-infectious chronic conditions that may share core mechanisms:

Long COVID. The most obvious parallel:

- Nearly identical symptom profile in many patients
- Similar post-exertional malaise pattern
- Common autonomic dysfunction
- Suggests SARS-CoV-2 triggers the same “trap” as other pathogens
- *Speculative link:* Both may involve spike protein persistence or viral reservoir maintaining immune activation

Post-Treatment Lyme Disease Syndrome. Chronic symptoms after Lyme treatment:

- Fatigue, cognitive dysfunction, pain
- Controversial whether active infection persists
- *Speculative link:* *Borrelia* may trigger same epigenetic/autoimmune locks; the specific pathogen matters less than the host response pattern

Post-Dengue Fatigue Syndrome. Chronic fatigue following dengue infection:

- Well-documented in endemic areas
- Similar symptom profile to ME/CFS
- *Speculative link:* Dengue’s immune evasion strategies may be particularly effective at triggering the “safe mode” lock

Gulf War Illness. Multi-symptom illness in Gulf War veterans:

- Fatigue, cognitive problems, pain, GI symptoms
- Multiple potential triggers (infections, chemical exposures, vaccines, stress)
- *Speculative link:* Multiple simultaneous stressors may be more likely to establish multiple locks simultaneously

? Open Question 15: Common Post-Infectious Pathway?

What if all these conditions—ME/CFS, Long COVID, post-Lyme, Gulf War Illness—represent the same underlying “locked sickness behavior” state triggered by different insults? The specific trigger might influence which symptoms predominate, but the core pathophysiology could be identical. This would explain why they’re so similar clinically yet have different apparent causes.

⚠ Limitation 4: Cross-Disease Extrapolation: Shared Symptoms Do Not Establish Shared Mechanisms

This chapter maps symptom overlaps between ME/CFS and numerous other conditions (Long COVID, POTS, MCAS, EDS, fibromyalgia, autoimmune diseases, mitochond-

drial diseases, cancer cachexia). While clinically suggestive, shared symptoms across conditions do not establish shared aetiology. Key epistemic boundaries:

- Symptom overlap may reflect convergent endpoints from distinct mechanisms rather than a common pathway; fatigue, cognitive dysfunction, and autonomic symptoms are final common manifestations of many unrelated pathologies.
- The “speculative links” offered for each condition are hypothesis-generating, not evidence-based—none has been tested by studies designed to distinguish shared mechanism from coincidental co-occurrence.
- Prevalence estimates for comorbidity (e.g., MCAS in ME/CFS, SFN in EDS) vary widely across studies due to inconsistent diagnostic criteria, referral bias, and lack of population-based data.
- The “multi-lock” framing applied to cross-disease connections has not been empirically validated as a model distinct from simpler explanations such as shared genetic susceptibility or diagnostic overlap.

14.17.2 The Dysautonomia Spectrum

ME/CFS overlaps heavily with autonomic dysfunction syndromes:

Postural Orthostatic Tachycardia Syndrome (POTS).

- Many ME/CFS patients meet POTS criteria
- Both involve small fiber neuropathy in subsets
- Both show autoantibodies to adrenergic receptors
- *Speculative link:* POTS may represent ME/CFS with predominant autonomic lock; or both may be manifestations of autoimmune autonomic ganglionopathy spectrum

Inappropriate Sinus Tachycardia.

- Elevated resting heart rate without clear cause
- Often comorbid with POTS and ME/CFS
- *Speculative link:* May reflect autoantibodies to cardiac β -receptors or sinoatrial node ion channels

Neurocardiogenic Syncope.

- Vasovagal responses at inappropriate times
- Common in ME/CFS population
- *Speculative link:* Reflects vagal afferent sensitization combined with impaired compensatory responses

? Open Question 16: Autonomic Autoimmunity Unifying Hypothesis

What if ME/CFS, POTS, and related dysautonomias all represent different manifestations of autoimmune attack on the autonomic nervous system? The specific antibody targets (muscarinic, adrenergic, ganglionic nicotinic, ion channels) might determine whether someone presents primarily as POTS, ME/CFS, or mixed. This “autoimmune autonomic spectrum” could be as common as rheumatoid arthritis but remains unrecognized because we don’t routinely test for the antibodies.

14.17.3 The Mast Cell Connection

Mast cell activation appears connected to ME/CFS:

Mast Cell Activation Syndrome (MCAS).

- High comorbidity with ME/CFS
- Explains chemical sensitivities, food reactions, flushing
- Mast cells release histamine, prostaglandins, cytokines
- *Speculative link:* MCAS may be both cause and effect—initial mast cell activation contributes to the trigger; ongoing activation maintains inflammation

Histamine Intolerance.

- Many ME/CFS patients report histamine-related symptoms
- May reflect DAO enzyme dysfunction or mast cell instability
- *Speculative link:* Histamine is a circadian regulator; chronic histamine excess might contribute to circadian desynchronization

Mastocytosis.

- Clonal mast cell disorders
- More severe than MCAS but overlapping symptoms
- *Speculative link:* Both conditions might involve mast cell progenitor dysregulation; ME/CFS could involve functional mastocytosis without clonal proliferation

? Open Question 17: Mast Cells as Central Orchestrators?

What if mast cells are the “hub” connecting multiple ME/CFS mechanisms? Mast cells:

- Are activated by stress, infection, and multiple triggers
- Release mediators affecting every organ system
- Can maintain chronic inflammation

- Are present at blood-brain barrier and affect CNS function
- Are regulated by autonomic nervous system (which is dysfunctional)

The mast cell might be the cell type where multiple locks converge.

? Open Question 18: Mast Cells as Neuro-Immune Signal Amplifiers?

The intimate physical proximity of mast cells to peripheral nerve endings (<20 nm in many tissues) may enable bidirectional signaling beyond currently recognized neuroimmune crosstalk. Could mast cells function as biological *signal repeaters* or *gain modulators* in the nervous system?

Proposed mechanism:

- Mast cells detect neurotransmitter spillover and neuropeptide signals from nearby nerves
- Release precisely timed micro-bursts of neurotransmitters (serotonin, histamine) and ions (Ca^{2+} , K^{+})
- Bridge gaps in neural signaling across regions of small fiber neuropathy
- Modulate sensory sensitivity by adjusting nerve receptor thresholds via protease release (e.g., tryptase activation of PAR2)

This would explain:

- **Allodynia and hyperalgesia:** Mast cells with lowered activation thresholds act as signal amplifiers, magnifying innocuous stimuli into pain signals
- **SFN-MCAS overlap:** Small fiber neuropathy (non-length-dependent pattern documented in 34% of ME/CFS patients [335]) combined with mast cell hyperreactivity creates paradoxical hypersensitivity despite nerve damage
- **Variability of sensory symptoms:** Mast cell activation state (influenced by histamine load, stress, inflammation) dynamically modulates sensory gain day-to-day
- **“Phantom” sensations:** Mast cells broadcasting signals to multiple nerve fibers create diffuse sensory fields beyond the original stimulus location

Testable predictions:

- Mast cell stabilizers should reduce allodynia severity
- Quantitative sensory testing abnormalities should correlate with mast cell activation markers (tryptase, histamine)
- Time-course studies: sensory thresholds should fluctuate with mast cell mediator levels
- Electrophysiology: mast cell degranulation near nerve fibers should alter nerve conduction patterns

Supporting evidence: Mast cells form CADM1-mediated adhesion structures with sensory neurons that amplify degranulation (2-fold) and IL-6 secretion (3-fold) [411]. Approximately 80% of mast cell disorder patients demonstrate small fiber neuropathy on objective testing [173], establishing the clinical overlap. Mast cell-nerve bidirectional signaling has been documented, though the specific role of tryptase-PAR2 interactions in

ME/CFS sensory symptoms remains to be established.

Current evidence gaps: No direct studies demonstrate mast cells amplifying neural signals in real-time. However, the physical infrastructure exists (proximity, neurotransmitter release capability, bidirectional signaling via CADM1 [411]), and the clinical phenotype (SFN + MCAS + allodynia) suggests functional coupling.

? Open Question 19: Mast Cells as Environmental Memory Keepers?

Mast cells are extraordinarily long-lived immune cells, persisting for years in the same tissue location at barrier surfaces (gut, skin, airways). Unlike B-cells that remember specific pathogens, could mast cells maintain an *epigenetic archive* of chronic environmental exposures?

Proposed mechanism:

- Mast cells continuously sample the local chemical environment over years
- Chronic exposures (pollutants, dietary patterns, stress hormones, microbiome metabolites) induce epigenetic modifications
- These modifications adjust degranulation thresholds and mediator release patterns
- Epigenetically modified mast cells maintain altered activation thresholds for their lifespan, creating persistent sensitization

This would explain:

- **Geographic remission:** Why some chronic illness patients improve upon moving to different climates or environments—new location lacks the accumulated “environmental signature” archived in mast cells
- **Chemical sensitivity acquisition:** Gradual sensitization to previously tolerated exposures as mast cells archive repeated low-level irritation
- **“Total load” phenomenon:** Why symptoms worsen with cumulative exposure to multiple triggers—mast cells integrate exposures over time rather than responding to isolated events
- **Delayed recovery after trigger removal:** Environmental changes require years to benefit because mast cells live for years and carry historical “memory”

Testable predictions:

- Mast cells from patients in different environments should show distinct epigenetic signatures
- Mast cell epigenetic profiles should correlate with lifetime environmental exposure history
- Geographic relocation should gradually shift mast cell epigenetic patterns over 1–3 years (matching mast cell lifespan)
- Tissue-resident mast cells should show different epigenetic profiles than circulating mast cell progenitors

Supporting evidence: Mast cells are exceptionally long-lived tissue residents (estimated months to years based on tissue turnover studies), maintaining themselves inde-

pendently from bone marrow and accumulating tissue-specific programming. Epigenetic mechanisms (DNA methylation, histone acetylation) are known to control immune cell activation thresholds in general, and chronic immune activation can create lasting epigenetic signatures in other cell types. Environmental exposures (dietary factors, pollution) can alter immune cell function through epigenetic modifications. MCAS patients show persistent alterations in activation thresholds that may reflect long-term cellular reprogramming.

Current evidence gaps: While immune cell epigenetic memory is established and mast cell activation thresholds are known to be epigenetically controlled, no studies have directly examined whether mast cells archive general environmental exposures beyond standard antigen-specific immunity. Mast cell epigenetics in ME/CFS remain entirely unstudied.

? Open Question 20: Circadian Mast Cell Regulation and Potential Temporal Learning

Mast cells possess intrinsic circadian clocks that regulate degranulation (established). But could they also develop *learned temporal associations* beyond the 24-hour circadian rhythm—anticipating specific triggers at arbitrary times based on repeated exposure patterns?

Proposed mechanism (speculative):

- *Established:* Mast cells have circadian clocks that regulate FcεRI expression and degranulation sensitivity based on time-of-day
- *Speculative:* With repeated exposure to triggers at consistent times (e.g., breakfast food at 8 AM daily), mast cells might develop learned temporal associations independent of circadian phase
- Granules could undergo partial “pre-thaw” 15–30 minutes before expected trigger time
- If trigger arrives on schedule, full degranulation occurs rapidly with amplified response
- If trigger is absent, partial priming gradually reverses

This would explain:

- **Time-of-day variability:** Why patients tolerate certain foods/medications better at different times—mast cells are or aren’t pre-primed
- **Nocturnal symptom flares:** If evening routines consistently trigger mild mast cell activation, circadian priming might amplify nighttime symptoms
- **Elimination diet inconsistency:** Removing a food might fail if mast cells remain circadian-primed for weeks, causing reactions to “safe” foods eaten at the same time
- **“Spontaneous” reactions:** Circadian mast cell priming without actual trigger exposure could cause symptoms at predictable times
- **Vacation effect:** Disrupted routines break circadian priming patterns, temporarily reducing reactivity

Testable predictions:

- Mast cell mediators (histamine, tryptase) should show circadian oscillations correlating with habitual trigger exposure times
- Time-series sampling: baseline mediator levels should rise 15–30 min before scheduled triggers
- Experimental circadian disruption (shift work simulation) should temporarily reduce food/medication reactions
- Re-timing trigger exposure (breakfast foods eaten at dinner) should shift circadian mediator patterns within 1–2 weeks

Supporting evidence: Mast cells possess intrinsic molecular clocks that temporally regulate degranulation through circadian oscillation of FcεRI receptor expression and downstream signaling components [412]. The mast cell clock is entrained by humoral factors (adrenal hormones) and can be modulated by environmental stressors [413]. Circadian disruption eliminates temporal gating of mast cell activation, resulting in sustained hyperreactivity throughout the day [414]. The immune system exhibits anticipatory responses to predictable environmental threats as a fundamental circadian function.

Current evidence gaps: While circadian immune regulation is established (cortisol awakening response, circadian cytokine patterns) and mast cells demonstrably have circadian clocks [412], this hypothesis proposes a distinct phenomenon: *learned temporal associations beyond the 24-hour circadian rhythm*. Nakamura et al. demonstrated that mast cells respond to endogenous circadian cues (hormones, light-dark cycles), not learned associations with specific environmental triggers at arbitrary times. No studies have examined whether mast cells can develop anticipatory priming to non-circadian temporal patterns (e.g., “breakfast at 8 AM” vs. “breakfast at 10 AM”). This would require a form of Pavlovian temporal conditioning not yet demonstrated in immune cells. The “predictive brain” framework is well-developed in neuroscience but hasn’t been applied to mast cell biology.

14.17.4 The Ehlers-Danlos Connection

The high comorbidity of ME/CFS with hypermobile Ehlers-Danlos Syndrome (hEDS) is striking and demands mechanistic explanation. Registry data from 815 ME/CFS patients found 15.5% were joint hypermobility positive, with this subgroup showing significantly worse quality of life, more autonomic symptoms, and higher rates of both POTS (33% vs. 20%) and formal EDS diagnosis (29% vs. 3%) [415]. This represents a distinct clinical phenotype within ME/CFS.

Epidemiological Evidence

Comorbidity Rates. Joint hypermobility prevalence varies across conditions:

- General population: 10–20%
- ME/CFS: 15.5–57% (varies by study and criteria)

- POTS: up to 57%
- Long COVID: approximately 30%
- Fibromyalgia: approximately 27%

The enrichment of hypermobility in ME/CFS and related conditions is statistically significant and biologically meaningful.

Clinical Phenotype Differences. ME/CFS patients with joint hypermobility (JH+) compared to those without (JH−) show [415]:

- Worse physical functioning and pain scores
- Higher burden of autonomic, neurocognitive, and musculoskeletal symptoms
- More frequent headaches and gastrointestinal symptoms
- Family history of EDS more common

This suggests JH+ ME/CFS may represent a mechanistically distinct subtype.

Mechanistic Pathways: From Connective Tissue to Systemic Dysfunction

The question is not merely *whether* EDS and ME/CFS are associated, but *why*. Several mechanistic pathways have varying levels of evidence.

Pathway 1: Vascular Laxity → Autonomic Dysfunction (HIGH EVIDENCE). This is the best-supported mechanistic link. Defective connective tissue directly affects blood vessel structure and function:

- **Increased arterial compliance:** EDS patients show significantly lower central pulse wave velocity (4.73 m/s vs. controls), indicating excessive arterial elasticity [334]. This impairs baroreceptor signaling—stretch receptors in vessel walls cannot accurately detect blood pressure changes when the walls are too compliant.
- **Excessive venous pooling:** Abnormal connective tissue in veins causes excessive distension under normal hydrostatic pressures [416]. Blood pools in lower extremities upon standing, reducing venous return and cardiac preload.
- **Compensatory tachycardia:** The heart races to maintain cardiac output despite reduced preload, producing POTS. Up to 70% of hEDS patients report dysautonomia symptoms, and up to 40% meet formal POTS criteria [154].
- **Cerebral hypoperfusion:** Inadequate blood pressure regulation leads to reduced cerebral blood flow, particularly upon standing, causing cognitive symptoms, lightheadedness, and fatigue.

This pathway explains why POTS is so prevalent in both hEDS and ME/CFS—the autonomic dysfunction in hEDS is a direct, structural consequence of connective tissue abnormality rather than a secondary phenomenon.

Pathway 2: Craniocervical Instability → Brainstem Dysfunction (MODERATE EVIDENCE).

Ligamentous laxity at the craniocervical junction (C0–C2) can cause structural instability with neurological consequences:

- **Brainstem compression:** The brainstem controls autonomic functions. Instability at the skull-spine junction can cause intermittent compression or stretching of brainstem structures [417].
- **CSF flow obstruction:** Craniocervical instability can obstruct cerebrospinal fluid flow at the craniocervical junction, potentially causing increased intracranial pressure and impairing glymphatic waste clearance.
- **Vertebral artery effects:** Cervical instability may affect vertebral artery flow, contributing to posterior circulation insufficiency.

A systematic review of 16 studies (695 EDS patients) found significant heterogeneity in diagnostic criteria for craniocervical instability, with no standardized thresholds [418]. Dynamic imaging (upright MRI, flexion-extension views) provides superior diagnostic information compared to static supine imaging. Some ME/CFS patients with craniocervical instability report improvement after surgical stabilization, though controlled outcome data remain limited.

△ **Warning 2: Craniocervical Instability: Evidence Limitations**

While biologically plausible, the CCI-ME/CFS connection remains largely anecdotal. No controlled studies have established:

- True prevalence of CCI in ME/CFS populations
- Whether CCI causes ME/CFS symptoms vs. co-occurring conditions
- Long-term surgical outcomes in ME/CFS patients with CCI

Screening for CCI may be appropriate in ME/CFS patients with hypermobility and progressive neurological symptoms, but surgery should be approached cautiously given the limited evidence base.

Pathway 3: Extracellular Matrix → Mast Cell Dysregulation (LOW EVIDENCE).

The “EDS-MCAS-POTS triad” is frequently discussed clinically, but a critical review found that “an evidence-based, common pathophysiologic mechanism between any of the two, much less all three conditions, has yet to be described” [151]. The proposed mechanisms remain speculative:

- **ECM-mast cell interactions:** Mast cells anchor to extracellular matrix proteins (fibronectin, vitronectin) via integrins. Bidirectional signaling means abnormal ECM composition could theoretically alter mast cell activation thresholds and mediator release patterns.
- **Abnormal tissue remodeling:** Mast cell proteases contribute to ECM remodeling. A vicious cycle might develop where abnormal ECM triggers mast cell activation, which causes further ECM abnormalities.

- **Epidemiological association:** Approximately 31% of patients with both POTS and EDS also have MCAS, compared to 2% of those without EDS. However, diagnostic criteria heterogeneity limits interpretation.

While the clinical co-occurrence is real, the mechanistic explanation remains a hypothesis rather than established science.

Pathway 4: Tissue Fragility → Purinergic Signaling (SPECULATIVE). This pathway connects EDS tissue fragility to the cell danger response hypothesis of ME/CFS [197]:

- **Microtrauma from daily activities:** EDS patients experience more joint subluxations, soft tissue injuries, and tissue stress from normal activities due to structural fragility.
- **ATP release:** Damaged and stressed cells release ATP into the extracellular space. This is a universal cellular alarm signal.
- **Purinergic receptor activation:** Extracellular ATP activates P2X and P2Y receptors, triggering the cell danger response—a metabolic shift toward a protective but hypometabolic state.
- **Chronic activation:** If tissue fragility causes ongoing microtrauma, the purinergic alarm system might never fully reset, maintaining the hypometabolic state characteristic of ME/CFS.

This pathway is mechanistically plausible but entirely unvalidated. No studies have measured extracellular ATP levels or purinergic receptor activation in EDS patients.

Pathway 5: Small Fiber Neuropathy as Common Downstream Pathway (MODERATE EVIDENCE). Small fiber neuropathy (SFN) may represent a convergent mechanism linking EDS structural pathology to ME/CFS-like symptoms:

- **Universal SFN in EDS:** All 24 EDS patients in one study showed decreased intraepidermal nerve fiber density consistent with SFN, with 95% meeting criteria for neuropathic pain [419].
- **SFN in ME/CFS:** ME/CFS patients show evidence of C-fiber denervation on quantitative sensory testing, with 31% meeting POTS criteria and 34% showing non-length-dependent SFN patterns [335].
- **Autonomic small fibers:** SFN affects not only sensory nerves but also autonomic small fibers controlling heart rate, blood pressure, digestion, sweating, and temperature regulation—explaining the widespread autonomic dysfunction in both conditions.

SFN may be where the EDS structural abnormality and the ME/CFS functional abnormality converge, though whether SFN in EDS has the same etiology as SFN in ME/CFS remains unknown.

Speculation 28 (Defective Extracellular Matrix Alters Lymph Node Architecture and Viral Clearance). **Certainty: 0.25.** Highly speculative—no direct evidence links hEDS connective tissue defects to lymph node microenvironment dysfunction. The hypothesis bridges two

independently supported observations (hEDS susceptibility to ME/CFS and viral persistence in lymph nodes) with a plausible but untested mechanistic bridge.

Lymph nodes are structurally dependent on a fibroblastic reticular cell (FRC) network embedded in extracellular matrix (ECM). This collagen- and fibronectin-rich scaffold organizes immune cell trafficking, antigen presentation, and pathogen clearance. In hEDS, systemic collagen defects alter ECM composition throughout the body. We speculate that this extends to lymphoid tissue, creating altered lymph node microenvironments where viral clearance is impaired.

Proposed Mechanism. Defective ECM in hEDS lymph nodes may produce:

- **Looser FRC networks:** Altered stromal architecture could change the spatial organisation of T cell zones and B cell follicles, reducing the efficiency of antigen presentation and immune synapse formation.
- **Abnormal conduit systems:** The FRC conduit system filters soluble antigens by size; altered collagen composition could change filtration properties, affecting which antigens reach dendritic cells.
- **Permissive niches for viral persistence:** If the Kol et al. FIP model [171] applies to humans, virus-harboring lymphocytes may find sanctuary in structurally abnormal lymph node regions where immune surveillance is spatially disorganised.

This would explain why hEDS patients are disproportionately susceptible to post-viral ME/CFS: the same connective tissue defect that causes joint hypermobility also compromises the structural infrastructure of immune defence.

Testable Predictions.

1. Lymph node biopsies or fine-needle aspirates from hEDS patients will show altered FRC network density and collagen composition compared to non-hEDS controls.
2. Advanced lymph node imaging (contrast-enhanced ultrasound or MR lymphangiography) will reveal structural differences in hEDS patients.
3. Post-COVID hEDS patients will show higher rates of persistent viral antigen in lymphoid tissue than non-hEDS post-COVID patients matched for illness duration and severity.
4. In vitro, culturing immune cells on hEDS-derived versus normal ECM scaffolds will show reduced viral clearance efficiency.

Falsifiability. This hypothesis would be falsified if lymph node architecture in hEDS patients is structurally normal on biopsy and imaging, or if viral persistence rates are equal in hEDS and non-hEDS ME/CFS patients. It would also be weakened if the hEDS–ME/CFS association is fully explained by autonomic dysfunction and MCAS without any lymphoid tissue involvement.

Limitations. No studies have examined lymph node histology in hEDS. The ECM defect in hEDS is heterogeneous and incompletely characterised; its effects on lymphoid tissue are entirely extrapolated. Lymph node biopsy is invasive, limiting feasibility. The hypothesis cannot explain ME/CFS in patients without connective tissue disorders.

The Deconditioning Spiral

A vicious cycle may amplify the initial pathology. In hEDS patients with dysautonomia [420]:

- 78% report exercise intolerance as a primary symptom
- Sedentary behavior increased from 44% to 85% after symptom onset
- Dysautonomic patients showed smaller cardiac chamber sizes and reduced left ventricular end-diastolic volume—cardiac atrophy from deconditioning

The proposed cycle:

1. Connective tissue abnormality → orthostatic intolerance
2. Orthostatic intolerance → exercise avoidance
3. Exercise avoidance → cardiovascular deconditioning
4. Deconditioning → reduced blood volume, cardiac atrophy
5. Reduced cardiovascular capacity → worsened orthostatic intolerance

This spiral is similar to—but distinct from—ME/CFS, where post-exertional malaise adds an additional constraint. In pure hEDS without ME/CFS, carefully graded exercise may help break the cycle. In ME/CFS with hEDS, the PEM constraint means standard exercise approaches are contraindicated (see Section 18.3.3).

Synthesis: EDS as Susceptibility Factor

~ Hypothesis 4: Connective Tissue Disorders as ME/CFS Susceptibility Factors

Hypermobility spectrum disorders do not cause ME/CFS directly but dramatically increase susceptibility through multiple mechanisms:

1. **Lower trigger threshold:** Pre-existing autonomic dysfunction means less physiological reserve. A viral infection that a person with normal connective tissue might recover from could tip an hEDS patient into chronic illness.
2. **Additional perpetuating mechanisms:** Craniocervical instability, vascular dysfunction, and mast cell activation provide additional “locks” that maintain the disease state once triggered.
3. **Impaired recovery capacity:** Tissue repair mechanisms are compromised. The body cannot fully restore homeostasis after an acute insult.
4. **Diagnostic confusion:** Symptom overlap delays ME/CFS diagnosis and appropri-

ate management. Patients may be told their symptoms are “just EDS” when they actually have both conditions.

This model explains the high comorbidity without requiring that EDS directly causes ME/CFS. Instead, EDS removes the physiological buffer that would normally allow recovery from acute triggers.

? Open Question 21: Research Priorities for EDS-ME/CFS Connection

Critical unanswered questions include:

- Does ME/CFS in hEDS patients have the same pathophysiology as ME/CFS in non-hypermobile patients, or are these distinct conditions with overlapping symptoms?
- Can early, aggressive management of dysautonomia in hEDS patients prevent progression to ME/CFS after viral triggers?
- What is the true prevalence of craniocervical instability in ME/CFS, and does surgical correction improve ME/CFS-specific outcomes?
- Do hEDS patients show elevated extracellular ATP or purinergic activation compared to controls?
- Is small fiber neuropathy in EDS mechanistically related to SFN in ME/CFS?

Answering these questions could identify preventive strategies and targeted treatments for this high-risk subgroup.

14.17.5 The Fibromyalgia Overlap

ME/CFS and fibromyalgia share extensive clinical and biological overlap. A systematic review and meta-analysis of 21 publications found that the two diagnoses co-occur in 47.3% of cases (95% CI: 45.97–48.63), though high heterogeneity ($I^2 = 98\%$) reflects varying diagnostic criteria across studies [421]. The 2016 Wolfe fibromyalgia criteria, which incorporate fatigue, unrefreshed sleep, and cognitive dysfunction, likely inflate this figure by absorbing core ME/CFS features into the FM definition. Community-based estimates place co-occurrence closer to 22–35%.

At the cerebrospinal fluid level, the two conditions may be even more closely related than clinical phenotypes suggest. Mass spectrometry analysis of 2,083 CSF proteins found **no significant differences** between ME/CFS patients with and without comorbid fibromyalgia, with machine learning achieving only AUC 0.67—barely above chance [110]. The authors concluded that the two conditions “fall along a common illness spectrum,” challenging the hypothesis that they have distinct central nervous system pathophysiology.

Neuroinflammation provides further convergence. Multi-site PET imaging with [^{11}C]PBR28 (a TSPO radioligand marking activated microglia and astrocytes) revealed significant microglial activation in the thalamus, somatosensory cortex, and prefrontal cortex of fibromyalgia patients compared to controls [422]. Parallel findings of microglial activation in ME/CFS suggest a shared neuroinflammatory substrate.

Autoantibody signatures add a further layer of convergence. In a study comparing ME/CFS patients with and without comorbid FM and healthy controls, 54.5% of the ME/CFS+FM group and 45.5% of the ME/CFS-only group showed abnormal GABA receptor immunoreactivity, versus 0% of controls. Altered β -endorphin immunoreactivity correlated with fatigue severity in both groups, suggesting dysregulated natural autoimmunity rather than classical autoimmune pathology [423].

Post-Exertional Malaise: Shared but Asymmetric. A three-level meta-analysis of 45 effects from 15 studies confirmed that both ME/CFS and fibromyalgia patients show small-to-moderate increases in pain severity following exercise, establishing pain-related PEM as a shared feature [108]. However, PEM after *physical* exertion remains the cardinal distinguishing criterion: present in >90% of ME/CFS patients but absent or inconsistent in pure fibromyalgia. When FM is comorbid, PEM severity is amplified (scores 79.63 vs. 71.18, $p < 0.001$).

Key Differences. Despite the overlap, clinically important distinctions remain. A recent comprehensive review [424] summarized the distinguishing features: the core symptom in ME/CFS is profound fatigue with obligatory PEM after physical exertion, while FM is defined by widespread chronic pain. NK cell reduction is consistent and pronounced in ME/CFS but variable in FM. Mitochondrial impairment is severe in ME/CFS versus moderate in FM. GPCR autoantibody evidence is stronger in ME/CFS, though emerging in FM.

? Open Question 22: Same Disease, Different Locks?

What if ME/CFS and fibromyalgia represent the same underlying pathophysiology with different predominant locks?

- **ME/CFS-predominant:** Stronger metabolic/immune locks, less central sensitization
- **Fibromyalgia-predominant:** Stronger central sensitization lock, less metabolic involvement
- **Mixed:** Both lock types active

The CSF proteome data [110] and shared microglial activation patterns [422] support this spectrum model. If confirmed, this would explain why they so often co-occur and why treatments for one sometimes help the other. The key clinical implication: clinicians seeing fibromyalgia should systematically screen for PEM, as the high co-occurrence (47%) means nearly half of FM patients may also meet ME/CFS criteria [421].

14.17.6 The Autoimmune Disease Spectrum

ME/CFS may sit on a continuum with recognized autoimmune diseases:

Sjögren's Syndrome (Gougerot-Sjögren). The overlap between primary Sjögren's syndrome (pSS) and ME/CFS is both clinically significant and mechanistically informative. Fatigue affects 65–70% of pSS patients and is consistently rated as the most debilitating ex-

traglandular symptom, predicting work disability and reduced quality of life independently of glandular involvement [425].

The “seronegative Sjögren’s” hypothesis has historical roots. Already in 1996, Nishikai et al. found that one-third of CFS patients presenting with sicca symptoms (dry eyes, dry mouth) fulfilled Sjögren’s diagnostic criteria while remaining seronegative—lacking the classic anti-SSA/SSB antibodies that define primary SS [426]. This raised the possibility that a subset of ME/CFS patients harbor subclinical or atypical Sjögren’s disease that escapes conventional serological detection.

A 2023 Berlin study brought contemporary rigor to this question. Kim et al. assessed 19 pSS patients against the Canadian Consensus Criteria for ME/CFS and found that 22% (4/18) fulfilled full CCC criteria, while 32% (6/19) had severe ME/CFS-like symptoms [427]. Crucially, PEM in pSS was primarily triggered by *mental and emotional* exertion rather than physical exertion, and hand grip strength recovered normally within 60 minutes after exercise—in contrast to the abnormal recovery pattern characteristic of ME/CFS. This suggests that while the fatigue phenotypes overlap, the underlying exercise physiology diverges.

The immunological connections are striking. Both pSS and ME/CFS show elevated β_1/β_2 -adrenergic and M3/M4-muscarinic receptor autoantibodies. However, in pSS these autoantibodies correlate with systemic disease activity (ESSDAI, $p < 0.05$) but *not* with fatigue severity [427]—whereas in ME/CFS, GPCR autoantibodies are hypothesized to mediate fatigue and autonomic dysfunction directly. The same molecular target thus appears to have different functional consequences in the two conditions.

Type I interferon dysregulation provides another convergence point. In pSS, elevated plasmacytoid dendritic cells drive a type I IFN signature that correlates with extraglandular manifestations and autoantibody production. Similar plasmacytoid dendritic cell elevation and increased type I IFN production have been documented in ME/CFS, potentially increasing susceptibility to autoimmune mechanisms in both conditions.

CSF neuroinflammation further links the two. In pSS, IL-1Ra in cerebrospinal fluid is significantly associated with fatigue severity, implicating IL-1 β -mediated neuroinflammation [428]. Parallel findings of elevated IL-1 β and neuroinflammatory markers in ME/CFS CSF suggest a shared central inflammatory driver of fatigue.

Small fiber neuropathy (SFN) may represent a unifying structural substrate across pSS, fibromyalgia, and ME/CFS. SFN prevalence reaches 49% in fibromyalgia by meta-analysis, is well-documented in pSS (presenting predominantly as a non-length-dependent pattern), and has been reported in ME/CFS cohorts at 40–50%. This shared neuropathic substrate could underlie the overlapping pain, autonomic, and sensory symptoms across all three conditions.

The diagnostic implications are bidirectional. The average diagnostic delay for Sjögren’s disease is 3–6 years, and pSS patients commonly receive initial labels of fibromyalgia, CFS, or functional disorder before correct diagnosis [429]. Conversely, 7–8% of pSS patients have no sicca symptoms at diagnosis, creating a risk that Sjögren’s is missed in ME/CFS patients who lack the classic dry eyes/mouth presentation. In any ME/CFS patient presenting with fatigue, dry eyes or mouth, joint pain, and sensory symptoms, Sjögren’s evaluation (Schirmer’s test, salivary flow, lip biopsy, SSA/SSB antibodies) is warranted—including when seronegative, given the Nishikai data [426].

Systemic Lupus Erythematosus.

- Fatigue is often the most disabling symptom
- Neuropsychiatric lupus resembles ME/CFS cognitively
- Complement abnormalities in both
- *Speculative link:* ME/CFS might involve lupus-like autoimmunity below diagnostic thresholds

Multiple Sclerosis.

- Fatigue is major symptom
- Cognitive dysfunction similar
- Both may involve HERV reactivation
- *Speculative link:* ME/CFS might be “diffuse MS” without discrete lesions, or MS-related autoimmunity affecting different neural targets

Autoimmune Encephalitis.

- Can present with fatigue, cognitive dysfunction, psychiatric symptoms
- Antibodies against neural proteins
- Often triggered by infection
- *Speculative link:* ME/CFS might be low-grade autoimmune encephalitis affecting widespread but subtle neural dysfunction

? Open Question 23: Subclinical Autoimmunity?

What if ME/CFS represents autoimmune disease below conventional detection thresholds? The autoantibodies might:

- Target functional receptors/channels rather than structural proteins
- Be present at low titers that affect function without triggering standard assays
- Target intracellular or unusual epitopes not covered by standard panels

This “subclinical autoimmunity” hypothesis would explain why immunomodulation helps some patients while standard autoimmune panels are negative.

14.17.7 The Mitochondrial Disease Connection

Primary mitochondrial diseases share features with ME/CFS:

- Mitochondrial diseases are often associated with fatigue
- Mitochondrial diseases often have cognitive dysfunction
- Mitochondrial diseases often have psychiatric symptoms
- Mitochondrial diseases often have autoimmune features
- Mitochondrial diseases often have genetic components
- Mitochondrial diseases often have treatment options

Overlapping Features.

- Exercise intolerance (defining in both)
- Post-exertional symptoms (delayed recovery in both)
- Cognitive dysfunction (both)
- Multi-system involvement (both)

Differences.

- Primary mitochondrial disease: genetic mutations, progressive
- ME/CFS: acquired, stable or fluctuating

? Open Question 24: Acquired Mitochondriopathy?

What if ME/CFS represents an “acquired mitochondrial disease” where the genetic code is intact but epigenetic changes or post-translational modifications create mitochondria that function as if mutated? The mitochondria might be:

- Epigenetically silencing key respiratory chain components
- Maintaining a “fission” state inappropriate for energy demands
- Preferentially undergoing mitophagy, reducing functional mitochondrial mass

This would explain the mitochondrial dysfunction without genetic mutations.

14.17.8 The Psychiatric Overlap—Reframed

ME/CFS has historically been conflated with depression and anxiety. A mechanistic reframing:

Shared Biology, Not Shared Psychology.

- Both ME/CFS and depression involve inflammatory cytokines
- Both involve kynurenine pathway abnormalities
- Both involve HPA axis dysregulation
- Both involve neurotransmitter changes

The Cytokine Theory of Depression.

- Depression may be, in part, an inflammatory brain state
- Cytokines cause “sickness behavior” that resembles depression
- *Speculative link:* ME/CFS and inflammatory depression might be the same phenomenon with different tissue distributions or lock combinations

? Open Question 25: Neuroimmune Spectrum Disorders?

What if ME/CFS, inflammatory depression, “brain fog” conditions, and some anxiety disorders all represent points on a “neuroimmune spectrum”? The common feature would be immune activation affecting brain function through:

- Direct cytokine effects on neurons
- Microglial activation
- Kynurenine pathway shifts
- Blood-brain barrier dysfunction

Different presentations might reflect which brain regions are most affected, not fundamentally different diseases.

14.17.9 The Cancer Cachexia Connection

Cancer-associated cachexia shares surprising features with ME/CFS:

Shared Features.

- Profound fatigue out of proportion to activity
- Muscle wasting/weakness
- Metabolic abnormalities
- Inflammatory cytokine elevation
- Anorexia and weight issues

Mechanistic Overlap.

- Both involve TNF- α (“cachexin”) elevation
- Both show muscle protein catabolism
- Both have mitochondrial dysfunction
- Both may involve the same metabolic “shutdown” program

? Open Question 26: Cachexia Without Cancer?

What if ME/CFS is essentially “cachexia without cancer”—the same metabolic shutdown program activated by inflammation, but without a tumor driving it? The “safe mode” hypothesis becomes even more compelling: the body is running a program designed for survival during severe illness (cancer, infection, trauma) but triggered inappropriately or locked on.

14.17.10 The Hibernation/Torpor Analogy

Some researchers have noted similarities between ME/CFS and hibernation:

Hibernation Features.

- Profound metabolic suppression
- Reduced body temperature
- Altered fuel utilization (lipid preference)
- Immune quiescence
- Rapid reversibility (in hibernators)

ME/CFS Parallels.

- Metabolic suppression (documented)
- Some patients report feeling cold
- Altered fuel utilization (documented)
- Immune changes (documented)
- NOT rapidly reversible (the “lock”)

? Open Question 27: Stuck in Torpor?

What if ME/CFS involves activation of ancient metabolic programs related to torpor or hibernation—programs that are suppressed in humans but not deleted from our genome? A severe enough trigger might activate these dormant programs. In hibernating animals, specific signals trigger arousal. In ME/CFS patients, those arousal signals might be missing or ineffective.

If true, studying the molecular biology of hibernation arousal might reveal therapeutic targets for ME/CFS.

14.17.11 Symptom-Specific Speculations

Some specific ME/CFS symptoms suggest particular connections:

Coat Hanger Pain (Neck/Shoulder Pain in Distribution of Trapezius).

- Classic dysautonomia symptom from muscle ischemia during orthostatic stress
- *Speculative link:* May indicate small vessel disease or microvascular dysfunction; could also reflect craniocervical issues

Post-Exertional Malaise Delay (24-72 Hours).

- Not immediate like normal fatigue
- *Speculative link:* Time course matches delayed-type hypersensitivity immune responses; may indicate immune-mediated component to PEM

“Wired but Tired” (Exhausted but Unable to Sleep).

- Paradoxical hyper-arousal with fatigue
- *Speculative link:* Classic presentation of ion channel dysfunction affecting both excitation (hyperactive) and energy (depleted); or circadian desynchronization with misaligned sleep drive and circadian alerting

Alcohol Intolerance.

- Many ME/CFS patients cannot tolerate even small amounts
- *Speculative link:* Could indicate ALDH dysfunction, already-compromised NAD⁺ pools (alcohol metabolism consumes NAD⁺), or mast cell activation (alcohol triggers mast cell degranulation)

Orthostatic Cognitive Impairment (Worse When Standing).

- Cognitive function declines in upright position
- *Speculative link:* Cerebral hypoperfusion from autonomic dysfunction, but could also indicate position-sensitive CSF dynamics affecting brain function (supporting glymphatic hypothesis)

Symptom Fluctuation with Menstrual Cycle.

- Many female patients report cycle-dependent symptoms
- *Speculative link:* Estrogen and progesterone affect immune function, mast cells, mitochondria, and virtually every proposed mechanism; hormonal influence on HERV expression might explain cyclical viral-like symptoms

14.18 Emerging Hypotheses from 2025 Research

Recent multi-omics studies and clinical trials have revealed patterns that suggest several novel mechanistic hypotheses not previously considered.

14.18.1 The Vascular-Immune-Energy Triad

? Open Question 28: Coordinated Three-System Failure

The Heng et al. 2025 study [12] identified a 7-biomarker diagnostic model spanning three systems: adenosine metabolism (AMP), immune markers (cDC1, LYVE1, IGHG2), and vascular factors (FN1, VWF, THBS1). This wasn't three separate findings—it was one integrated signature. What if ME/CFS fundamentally involves a coordinated failure

mode across these three systems that cannot be understood or treated in isolation?

The triad might work as follows:

1. **Energy failure** (elevated AMP/ADP, reduced ATP) impairs immune cell maturation and function
2. **Immature immune cells** (elevated naïve B cells, reduced switched memory B cells, immature T cell subsets) fail to properly regulate vascular function and produce dysfunctional antibodies
3. **Vascular dysfunction** (elevated VWF, fibronectin, thrombospondin) reduces tissue perfusion, causing cellular hypoxia that worsens energy production

This creates a stable triangular trap where each vertex reinforces the others. Treating only one system fails because the other two pull it back.

Therapeutic Implication. Effective treatment might require simultaneous intervention at all three vertices: NAD⁺ precursors for energy, immunomodulation for immune maturation, and vascular-targeted therapy (anticoagulation, endothelial support) for perfusion. The daratumumab success (60% response) might reflect cases where the autoimmune vertex was dominant—remove it, and the triad destabilizes enough to collapse.

14.18.2 The Plasma Cell Sanctuary Hypothesis

? Open Question 29: Long-Lived Plasma Cells as Disease Reservoir

The daratumumab trial's success—where targeting CD38⁺ plasma cells produced sustained remission in 60% of patients—reveals something important: rituximab (anti-CD20) failed in ME/CFS trials, yet daratumumab (anti-CD38) succeeded. Both deplete antibody-producing cells, but they target different populations.

B cells (CD20⁺) are the precursors; plasma cells (CD38⁺) are the factories. Crucially, long-lived plasma cells can survive for *decades* in bone marrow and gut niches, continuously secreting antibodies without needing B cell replenishment. What if ME/CFS is maintained by these “sanctuary” plasma cells?

Under this model:

- An initial trigger (infection) generates autoreactive B cells
- Some differentiate into long-lived plasma cells that migrate to survival niches
- These plasma cells produce autoantibodies (anti-GPCR, anti-ion channel) indefinitely
- Rituximab depletes B cells but not established plasma cells—antibody production continues
- By the time B cells return, the patient hasn't improved, so the trial “fails”
- Daratumumab directly kills the plasma cell factories, stopping antibody production

This explains the 8–9 month delay before maximum daratumumab benefit: existing

autoantibodies must decay (IgG half-life ~3 weeks, but tissue-bound antibodies persist longer).

Undocumented Phenomenon. If true, ME/CFS patients should have expanded populations of long-lived plasma cells in bone marrow biopsies, and these cells should be producing the pathogenic autoantibodies. This has never been directly examined.

Treatment Implication. Combining daratumumab (kill factories) with immunoadsorption (remove existing antibodies) might produce faster and more complete responses than either alone.

14.18.3 The Endothelial Activation Cascade

? Open Question 30: Chronic Endotheliopathy as Core Mechanism

The Heng 2025 study [12] found elevated plasma proteins associated with “activation of the endothelium and remodeling of vessel walls.” Specifically: VWF (von Willebrand factor), FN1 (fibronectin), and THBS1 (thrombospondin-1). These aren’t random inflammatory markers—they suggest a specific pathology: chronic endothelial activation.

Endothelial cells line all blood vessels. When activated (by infection, inflammation, autoantibodies, or hypoxia), they:

- Release VWF, promoting platelet adhesion and microclotting
- Deposit fibronectin, contributing to vascular remodeling
- Express thrombospondin, which is anti-angiogenic and pro-fibrotic
- Become “leaky,” allowing inappropriate extravasation
- Lose their normal anti-inflammatory and vasodilatory functions

What if ME/CFS is fundamentally an endotheliopathy—a chronic disease of blood vessel lining? This would explain:

- **Exercise intolerance:** Dysfunctional endothelium cannot vasodilate properly to meet demand
- **Brain fog:** Cerebral microvascular dysfunction impairs cognition
- **Orthostatic intolerance:** Poor vascular tone regulation
- **PEM:** Exercise-induced endothelial stress takes days to resolve
- **Multi-system involvement:** Endothelium is everywhere

Connection to Long COVID. This hypothesis aligns with the “microclot” findings in Long COVID, where amyloid-fibrin microclots persist in circulation. ME/CFS might involve the same endothelial activation without necessarily forming detectable microclots.

Undocumented Phenomenon. Direct endothelial function testing (flow-mediated dilation, EndoPAT) in ME/CFS has been limited. Comprehensive endothelial biomarker panels and functional testing might reveal a consistent endotheliopathy signature.

Treatment Implication. If endothelial dysfunction is central:

- Endothelial-protective supplements (L-arginine, L-citrulline, beetroot/nitrates) might help
- Statins (pleiotropic endothelial benefits beyond cholesterol) might be beneficial
- Low-dose aspirin or other anti-platelet agents might reduce microclot burden
- ACE inhibitors (endothelial-protective independent of blood pressure) could be therapeutic
- HELP apheresis (removes fibrinogen and inflammatory mediators) might address both cause and consequence

14.18.4 The Dendritic Cell Maturation Block

? Open Question 31: Stuck Immune Development

The Heng 2025 study [12] found reduced CD1c⁺CD141[−] conventional dendritic cells type 2 (cDC2) and a general skewing toward “less mature” immune cell subsets across T cells, NK cells, and dendritic cells. This isn’t random immune dysfunction—it suggests a specific developmental block.

Dendritic cells are the “conductors” of the immune orchestra. They:

- Capture antigens and present them to T cells
- Determine whether immune responses are inflammatory or tolerogenic
- Bridge innate and adaptive immunity
- Mature in response to danger signals

What if ME/CFS involves a block in dendritic cell maturation? Immature DCs:

- Present antigens inefficiently
- Fail to properly activate T cells
- May promote tolerance when activation is needed (chronic infection persistence)
- May promote inflammation when tolerance is needed (autoimmunity)

The immune system would be simultaneously ineffective (can’t clear threats) and dysregulated (inappropriate responses). This dual failure could maintain chronic immune activation without resolution.

Why Maturation Might Be Blocked.

- **Energy deficit:** DC maturation is metabolically demanding; ATP shortage might arrest development

- **Chronic antigen exposure:** Persistent viral antigens or autoantibodies might cause “exhaustion”
- **Cytokine milieu:** Altered cytokine patterns might signal DCs to remain immature
- **Epigenetic lock:** Maturation genes might be epigenetically silenced

Treatment Implication. Therapies that promote DC maturation (GM-CSF, specific TLR agonists, DC-targeted vaccines) might help—but could also be dangerous if the DCs then activate against self-antigens. This is a double-edged sword requiring careful patient selection.

14.18.5 The NAD⁺ Depletion Spiral

? Open Question 32: NAD⁺ as the Central Bottleneck

Multiple findings converge on NAD⁺:

- Heng et al. [12]: Abnormal NAD⁺ metabolism in ME/CFS immune cells
- The tryptophan-kynurenine pathway terminates in NAD⁺ synthesis
- PARP enzymes (activated by DNA damage/oxidative stress) consume NAD⁺
- Sirtuins (cellular stress response) require NAD⁺
- Mitochondrial Complex I requires NAD⁺/NADH cycling

What if NAD⁺ depletion is not just a consequence but a central driver—a bottleneck where multiple pathological processes converge?

The spiral might work as follows:

1. Initial insult causes oxidative stress and DNA damage
2. PARP enzymes activate to repair damage, consuming NAD⁺
3. NAD⁺ depletion impairs mitochondrial function (Complex I requires NAD⁺)
4. Mitochondrial dysfunction increases oxidative stress
5. More oxidative stress → more PARP activation → more NAD⁺ depletion
6. Meanwhile, inflammatory IDO activation shunts tryptophan away from serotonin toward kynurenine-NAD⁺ pathway—but the NAD⁺ produced may be immediately consumed by PARPs
7. Sirtuins, starved of NAD⁺, cannot perform their protective functions (autophagy, mitophagy, epigenetic regulation)
8. The cell enters a stable low-NAD⁺ state where it survives but cannot function normally

Undocumented Phenomenon. Direct measurement of NAD⁺/NADH ratios in ME/CFS patient tissues (not just blood) has been limited. If the spiral hypothesis is correct:

- Tissue NAD⁺ should be severely depleted
- PARP activity should be chronically elevated
- Sirtuin activity should be reduced

- The kynurenine pathway should be active but NAD⁺ still depleted (production consumed by PARPs)

Treatment Implication. NAD⁺ precursors (NR, NMN) alone might fail if PARPs immediately consume the new NAD⁺. Combination with PARP inhibitors (used in cancer) might be necessary—but PARP inhibition carries risks (impaired DNA repair). A gentler approach: high-dose NAD⁺ precursors to “flood” the system beyond PARP consumption capacity.

14.18.6 The Effort-Preference Recalibration

? Open Question 33: Central Effort Computation Gone Wrong

The Walitt 2024 NIH study made a crucial distinction: ME/CFS patients showed *altered effort preference*, not physical fatigue or central fatigue. Their muscles could produce force; their brain could generate motor commands. But when given choices, they systematically avoided effortful options even when rewards were high.

This isn’t laziness or depression—it’s a recalibration of the brain’s effort-reward computation. The brain has a system (involving the anterior cingulate cortex, insula, and dopaminergic circuits) that weighs expected effort against expected reward to decide whether actions are “worth it.”

What if ME/CFS involves a fundamental shift in this computation, such that:

- Effort is perceived as more costly than it actually is
- Rewards are perceived as less valuable than they would be
- The “break-even” point shifts dramatically toward rest
- This shift is protective (effort genuinely IS more costly due to metabolic dysfunction) but becomes miscalibrated

The CSF catecholamine deficiency found by Walitt et al. supports this: dopamine is central to effort-reward computation. Reduced central dopamine would systematically bias the system toward effort avoidance.

Why This Matters. If effort preference is centrally altered, then:

- “Pushing through” fights against an active brain computation, not just physical limits
- The system might be trainable but requires different approaches than physical reconditioning
- Dopaminergic interventions might help recalibrate the computation
- But if the recalibration is *appropriate* given metabolic dysfunction, forcing change could be harmful

Treatment Implication. Low-dose stimulants (methylphenidate, modafinil) might shift effort-reward computation—but could cause crashes if patients then overexert. The key might be: restore metabolic function FIRST, then (if needed) recalibrate effort perception.

14.18.7 The Immune Cell Energy Crisis**? Open Question 34: Starving Sentinels**

The Heng 2025 finding [12] of elevated AMP/ADP in white blood cells suggests immune cells specifically are energy-starved. This has profound implications because immune cells are *metabolically unique*:

- Naïve T cells are metabolically quiescent
- Upon activation, T cells undergo massive metabolic reprogramming (Warburg effect)
- This reprogramming requires abundant ATP and NAD⁺
- If immune cells cannot meet energy demands, activation fails
- Failed activation = ineffective immune responses + potential for inappropriate responses

The pattern of “immature” immune cells in ME/CFS might not reflect a developmental block per se, but rather an *energy crisis* that prevents cells from completing their activation/maturation programs.

Consider: a T cell encounters its antigen and begins activation. Activation requires massive ATP expenditure. But the cell is already AMP/ADP-elevated, ATP-depleted. It cannot complete activation. It either:

- Dies (activation-induced cell death from energy failure)
- Becomes anergic (gives up on activation)
- Partially activates (creating dysfunctional effector cells)

Any of these outcomes would create the immune dysfunction pattern seen in ME/CFS.

Undocumented Phenomenon. The metabolic competence of ME/CFS immune cells during activation has not been thoroughly studied. Prediction: ME/CFS T cells stimulated in vitro should show impaired metabolic reprogramming (measured by Seahorse assay or similar).

Treatment Implication. Supporting immune cell metabolism specifically might help:

- NAD⁺ precursors might restore immune cell energy capacity
- Specific metabolites (pyruvate, α -ketoglutarate) might bypass defective pathways
- Ketone bodies (which immune cells can use as fuel) might provide alternative energy

14.18.8 The Vascular “Memory” Hypothesis

? Open Question 35: Trained Endothelial Dysfunction

Immune cells can be “trained”—epigenetically reprogrammed by past exposures to respond differently to future stimuli. This innate immune memory (distinct from adaptive immunity) has been demonstrated in monocytes, macrophages, and NK cells.

What if endothelial cells can also be “trained”—and what if ME/CFS involves mal-adaptive endothelial training?

Endothelial cells experience the initial infection/inflammation. They activate, express adhesion molecules, become pro-thrombotic. Normally they return to quiescence. But what if severe or prolonged activation creates epigenetic changes that lock them in a partially activated state?

This “trained endotheliopathy” would:

- Persist long after the original trigger resolves
- Be present throughout the vasculature (explaining multi-system symptoms)
- Respond excessively to normal stimuli (exercise, stress, infection)
- Be resistant to conventional anti-inflammatory treatment
- Potentially be reversible with epigenetic interventions

Undocumented Phenomenon. Epigenetic profiling of endothelial cells from ME/CFS patients has not been performed. Circulating endothelial cells or endothelial progenitor cells might show characteristic epigenetic signatures.

⚠ Limitation 5: 2025 Research Findings: Early-Stage Evidence Requiring Replication

The hypotheses in this section derive largely from a single multi-omics study (Heng et al. 2025) and from reinterpretation of the NIH deep-phenotyping study (Walitt et al. 2024, n=17). Epistemic boundaries:

- The Heng 2025 biomarker model has not been independently validated; the 7-biomarker signature was derived from a single cohort and may not generalise to other ME/CFS populations or diagnostic criteria.
- The “vascular-immune-energy triad,” “NAD⁺ depletion spiral,” and “dendritic cell maturation block” are interpretive frameworks imposed on cross-sectional omics data—the data establish *association*, not the causal relationships depicted.
- The “effort-preference recalibration” hypothesis reinterprets the Walitt NIH study, which was itself controversial: the cohort was small (n=17 PI-ME/CFS), used contested inclusion criteria, and some findings (e.g., grip strength claims) were disputed by patient advocacy organisations and independent researchers.
- Treatment protocols proposed below (triple-target, plasma cell eradication, endothelial restoration) are speculative extrapolations from these early-stage findings; none has been tested even in pilot form.

14.18.9 Speculative Treatment Approaches from 2025 Findings

Based on the above hypotheses, several novel treatment approaches emerge:

The Triple-Target Protocol

Speculation 29 (Simultaneous Triad Intervention). If the vascular-immune-energy triad is the core mechanism, a protocol targeting all three simultaneously might produce synergistic effects:

1. **Energy:** High-dose NAD⁺ precursor (NR 1000–2000 mg/day) plus mitochondrial cofactors (CoQ10, PQQ, B vitamins)
2. **Immune:** Low-dose naltrexone (immune modulation) plus vitamin D optimization (immune regulation)
3. **Vascular:** L-arginine/citrulline (endothelial NO production) plus low-dose aspirin (antiplatelet) plus omega-3 fatty acids (endothelial protection)

This combination is relatively safe and addresses all three triad vertices. The hypothesis predicts it should work better than any single intervention.

The Plasma Cell Eradication Strategy

Speculation 30 (Deep Autoantibody Elimination). For patients with evidence of autoimmunity (elevated anti-GPCR antibodies, post-infectious onset, dramatic response to immunoadsorption):

1. **Phase 1:** Immunoadsorption series to remove circulating autoantibodies
2. **Phase 2:** Daratumumab (or similar CD38-targeting agent) to eliminate plasma cell factories
3. **Phase 3:** Monitor for autoantibody rebound; repeat if needed
4. **Phase 4:** Once autoantibodies cleared, assess whether other “locks” need addressing

This aggressive approach would only be appropriate for patients with clear autoimmune features and access to specialized centers.

The Endothelial Restoration Protocol

Speculation 31 (Vascular Healing Focus). If endotheliopathy is central, a vascular-focused protocol might help:

1. **Reduce endothelial activation:** Statin therapy (pleiotropic endothelial effects)
2. **Support NO production:** L-citrulline (better than L-arginine for sustained NO)
3. **Address microclots:** Nattokinase (fibrinolytic enzyme) or low-dose anticoagulation if indicated

4. **Protect endothelium:** Sulforaphane (Nrf2 activation), omega-3s, anthocyanins
5. **Reduce thrombotic tendency:** Aspirin, adequate hydration, compression if tolerated

This approach treats ME/CFS as a vascular disease, which it may fundamentally be in at least a subset of patients.

14.19 Novel Hypotheses from Two-Day CPET Findings

The objective demonstration of Day 2 metabolic failure in two-day cardiopulmonary exercise testing [1] provides unprecedented functional data that suggests several novel therapeutic approaches and previously undocumented biological phenomena. This section explores speculative hypotheses arising directly from these findings.

14.19.1 The Autonomic-Mitochondrial Feedback Loop

? Open Question 36: Bidirectional Autonomic-Metabolic Amplification

Keller et al. identified autonomic dysregulation as the primary mechanism linking Day 2 cardiopulmonary failures [1]. Walitt et al. documented central catecholamine deficiency [5]. Heng et al. demonstrated cellular ATP depletion [12]. What if these are not separate phenomena but nodes in a self-amplifying feedback loop?

Proposed mechanism:

1. Central catecholamine deficiency impairs autonomic cardiovascular regulation
2. Poor blood flow distribution during exercise causes tissue hypoxia
3. Mitochondria operating under hypoxic conditions generate excess ROS
4. ROS damages catecholamine synthetic enzymes and depletes Tetrahydrobiopterin (BH4) cofactor
5. Further catecholamine reduction worsens autonomic dysfunction
6. Cycle amplifies with each exertional episode

This would explain the **13-day recovery period**: breaking this vicious cycle requires not just substrate replenishment (hours) but restoration of damaged enzymes, clearance of oxidative damage products, and mitochondrial turnover (days to weeks).

Testable Predictions

1. Catecholamine synthetic enzyme activity should decline further in the 24–72 hours post-exercise
2. Tetrahydrobiopterin (BH4) levels should show exercise-dependent depletion with slow recovery kinetics
3. Interventions supporting both catecholamine synthesis (Tetrahydrobiopterin (BH4), tyrosine, cofactors) and mitochondrial protection (antioxidants) should show synergistic effects exceeding either alone

4. Baseline autonomic function (HRV, baroreflex sensitivity) should predict severity of Day 2 CPET decline
5. Serial measurement of oxidative stress biomarkers (isoprostanes, oxidized glutathione) should peak 24–48 hours post-exertion, correlating with symptom severity

Therapeutic Implications (Speculative)

Speculation 32 (Autonomic-Mitochondrial Co-Support Protocol). If the autonomic-mitochondrial feedback loop drives PEM, breaking it might require simultaneous intervention at multiple nodes:

Catecholamine support tier:

- L-tyrosine 1500–3000 mg/day (precursor)
- Sapropterin (Tetrahydrobiopterin, BH4) or methylfolate + B12 (BH4 recycling pathway support)
- Iron, vitamin B6, vitamin C, copper (cofactors for synthetic enzymes)
- Timing: morning administration to support daytime autonomic function

Mitochondrial protection tier:

- MitoQ or ubiquinol 200–400 mg/day (mitochondria-targeted antioxidant)
- N-Acetylcysteine (NAC) 1200–1800 mg/day (glutathione precursor, oxidative stress buffer)
- Alpha-lipoic acid 600 mg/day (mitochondrial antioxidant, Tetrahydrobiopterin (BH4) regeneration support)
- PQQ 20 mg/day (supports mitochondrial biogenesis)

Rationale: If both autonomic and mitochondrial dysfunction must improve simultaneously to break the loop, single-target interventions might fail where combination succeeds. The 13-day recovery period suggests sustained support is needed—acute supplementation around exertion may be insufficient.

Qualification: This is **highly speculative** and has not been tested. Individual components have varying levels of evidence, but the specific combination and the mechanistic rationale are hypothetical. Safety profile is generally good for listed supplements at suggested doses, but medical supervision is appropriate, especially for patients on other medications.

14.19.2 Mitochondrial Turnover Rate Limitation

? Open Question 37: Is Recovery Limited by Mitochondrial Half-Life?

The 13-day recovery period [1] closely approximates published mitochondrial turnover times in muscle tissue (10–15 days). This is likely not coincidental.

Hypothesis: Exercise-induced ROS damage creates a population of dysfunctional mitochondria that must be removed via mitophagy and replaced via biogenesis. The rate-limiting step is not substrate availability (which recovers in hours) but the physical replacement of damaged organelles.

Implications:

- **Why pacing works:** Staying below the threshold that causes significant mitochondrial damage prevents the need for prolonged turnover-dependent recovery
- **Why GET fails:** Repeated exertion before turnover is complete accumulates progressively more damaged mitochondria
- **Why baseline function declines:** Steady-state mitochondrial dysfunction worsens if damage rate exceeds replacement rate
- **Why severity varies:** Individual differences in mitophagy/biogenesis capacity determine how quickly patients can recover

Documented in other contexts: Mitochondrial turnover limitation is established in aging, neurodegenerative diseases, and certain myopathies. The novelty here is recognizing it as central to post-exertional malaise.

Therapeutic Implications (Speculative)

Speculation 33 (Accelerated Mitochondrial Turnover Protocol). If mitochondrial turnover is rate-limiting, interventions that accelerate both mitophagy (removal) and biogenesis (replacement) might shorten recovery time:

Mitophagy enhancement:

- **Urolithin A** 500–1000 mg/day: Directly stimulates mitophagy via PINK1/Parkin pathway; human trials show safety and efficacy in improving mitochondrial function in older adults
- **Spermidine** 1–3 mg/day: Autophagy inducer; safety established in human trials
- **Time-restricted eating:** If tolerated, 14–16 hour daily fast stimulates autophagy; CAUTION: many ME/CFS patients cannot tolerate fasting due to hypoglycemia symptoms

Mitochondrial biogenesis support:

- **NAD⁺ precursors:** NMN 500–1000 mg/day or NR 500–1000 mg/day activate sirtuins and PGC-1 α (master regulator of mitochondrial biogenesis)
- **Resistance training:** In healthy individuals, resistance exercise stimulates mitochondrial biogenesis; in ME/CFS, would require careful titration below PEM threshold (isometric exercises may be tolerable)

- **Cold exposure:** Mild cold stimulates PGC-1 α ; cold showers or cryotherapy if tolerated

Qualification: This approach is **speculative**. Urolithin A and NAD⁺ precursors have human safety data but not specifically in ME/CFS. The hypothesis that accelerating turnover would shorten recovery is logical but untested. Paradoxically, stimulating autophagy/mitophagy requires energy, so this approach might initially worsen symptoms in severely affected patients. Starting at very low doses and monitoring carefully would be essential.

14.19.3 Pre-Conditioning Hypothesis (Highly Speculative)

? Open Question 38: Can Controlled Sub-Threshold Stress Induce Adaptation?

A counterintuitive idea emerges from cardiology and neuroscience: **ischemic preconditioning**. Brief, controlled ischemic episodes protect against subsequent severe ischemia by activating protective cellular programs.

Could analogous “metabolic preconditioning” work in ME/CFS? That is, could carefully controlled, very brief exertional stress—well below the PEM threshold—activate protective adaptations without causing damage?

Theoretical basis:

- Brief ROS bursts activate Nrf2 and other protective transcription factors
- Mild metabolic stress upregulates antioxidant enzymes and heat shock proteins
- Hormetic dose-response: small stress beneficial, large stress harmful

Potential protocol (entirely speculative):

- Very brief activity (30–60 seconds) at 50–60% of anaerobic threshold
- Performed every 48–72 hours initially
- Monitor for any PEM; if occurs, cease immediately and reassess
- Hypothesis: might gradually increase mitochondrial capacity without triggering damage

Major caveats:

- This contradicts pacing principles and could easily cause harm if dose miscalculated
- No evidence this would work in ME/CFS; ischemic preconditioning is mechanistically distinct
- Would only be appropriate for stable mild-to-moderate patients, not severe cases
- Requires extremely careful monitoring and willingness to abandon approach if harmful

Why mention it: Because the two-day CPET shows objective metabolic failure, it also provides an objective outcome measure for testing whether any intervention (including preconditioning) improves function. This hypothesis is offered as an example of testable ideas that emerge from mechanistic understanding, even if it seems counterintuitive.

14.19.4 Circadian Optimization of Recovery

? Open Question 39: Is Mitochondrial Turnover Circadian-Gated?

Mitophagy and mitochondrial biogenesis are circadian-regulated processes, peaking at specific times of day. What if the prolonged recovery in ME/CFS reflects not just slow turnover but **mistimed turnover** due to circadian dysregulation?

Known facts:

- Mitophagy peaks during the inactive phase (night in humans)
- PGC-1 α (biogenesis regulator) has circadian expression
- ME/CFS patients have documented circadian abnormalities
- Sleep fragmentation impairs mitochondrial quality control

Hypothesis: If mitochondrial turnover processes are temporally disorganized, damaged mitochondria might persist longer because clearance and replacement occur out of phase with each other or are inefficiently timed.

Therapeutic Implications (Speculative)

Speculation 34 (Chronotherapy for Enhanced Recovery). If circadian timing matters for mitochondrial turnover, optimizing the timing of interventions might enhance efficacy:

Circadian stabilization:

- Strict sleep-wake schedule (even on weekends)
- Bright light exposure morning (10,000 lux for 30 min)
- Blue light blocking evening (2–3 hours before bed)
- Melatonin 0.5–3 mg at consistent time (8–9 PM)
- Temperature regulation (cool bedroom, 65–68°F)

Timed supplementation:

- **Mitophagy inducers** (urolithin A, spermidine): Evening dose to align with natural nocturnal mitophagy peak
- **Biogenesis support** (NAD⁺ precursors): Morning dose to support daytime activity
- **Antioxidants**: Split dose (morning and evening) for continuous protection

Qualification: This is **speculative**. While chronotherapy principles are established for other conditions (depression, jet lag), application to ME/CFS mitochondrial turnover is hypothetical. The interventions listed are generally safe but untested for this specific purpose.

14.19.5 Exercise Metabolomics-Guided Personalization

? Open Question 40: Can We Measure What's Depleted and Replace It?

The two-day CPET provides a standardized exertional challenge. What if we performed detailed metabolomics immediately after Day 1 exercise to identify which specific substrates, cofactors, or metabolites are depleted in individual patients, then targeted repletion before Day 2?

Undocumented phenomenon: No study has performed comprehensive metabolomics in the immediate post-exercise period (0–6 hours) in ME/CFS to identify acute depletions.

Hypothesis: Individual patients may have distinct metabolic bottlenecks:

- Patient A: carnitine depletion (impaired fatty acid oxidation)
- Patient B: glutathione depletion (oxidative stress overwhelm)
- Patient C: tryptophan/kynurenine pathway derangement
- Patient D: purine nucleotide depletion (ATP synthesis substrate limitation)

Targeted repletion based on individual metabolic signatures might prevent Day 2 deterioration more effectively than generic interventions.

Research Protocol (Proposed)

1. **Baseline metabolomics:** Plasma/serum immediately before CPET-1
2. **Post-exercise metabolomics:** 30 min, 2 hours, and 6 hours after CPET-1
3. **Identify depletions:** Metabolites showing >30% decline post-exercise
4. **Cluster analysis:** Identify metabolic subgroups
5. **Targeted repletion trial:** Provide individualized supplementation between Day 1 and Day 2
6. **Outcome:** Measure whether Day 2 deterioration is reduced

Qualification: This is a proposed research direction, not an established finding. Metabolomics is expensive and not clinically available. However, if successful, it could guide development of standardized metabolic phenotyping that eventually becomes clinically accessible.

14.19.6 Vagal Stimulation for Recovery Acceleration

~ Hypothesis 5: Parasympathetic Enhancement of Repair

The autonomic nervous system has two branches: sympathetic (“fight or flight”) and parasympathetic (“rest and digest”). The parasympathetic branch, mediated by the vagus nerve, promotes:

- Anti-inflammatory signaling (cholinergic anti-inflammatory pathway)
- Enhanced mitochondrial biogenesis

- Improved heart rate variability
- Activation of repair/regeneration programs

ME/CFS patients show reduced vagal tone (low HRV, poor parasympathetic modulation). What if enhancing vagal activity could accelerate recovery from exertion?

Evidence level: Vagal nerve stimulation (VNS) is FDA-approved for epilepsy and depression. Non-invasive VNS devices are available. VNS has been shown to reduce inflammation and improve mitochondrial function in other contexts. However, it has not been tested specifically for ME/CFS post-exertional recovery.

Therapeutic Approach (Speculative)

Speculation 35 (Post-Exertion Vagal Stimulation). Proposed protocol:

- **Device:** Transcutaneous auricular vagal nerve stimulation (taVNS) or transcutaneous cervical VNS
- **Timing:** Initiated within 1–2 hours of unavoidable exertion
- **Duration:** 30–60 minutes daily for 3–5 days post-exertion
- **Parameters:** Device-specific; typically 20–30 Hz stimulation
- **Goal:** Enhance parasympathetic tone during critical recovery period

Non-device alternatives:

- Deep breathing exercises (5–6 breaths per minute activates vagal reflexes)
- Humming or singing (stimulates vagus)
- Cold water face immersion (dive reflex)
- Specific yoga practices (if tolerable)

Qualification: This is **moderately speculative**. VNS devices have established safety profiles and known anti-inflammatory effects. The hypothesis that vagal stimulation could accelerate ME/CFS recovery is logical but unproven. Non-device alternatives are essentially free and safe, making them reasonable to try. Device-based VNS should be discussed with physicians and might not be covered by insurance for this indication.

14.19.7 Blood Flow Redistribution Training

? Open Question 41: Can We Train Better Autonomic Blood Flow Control?

Keller et al. concluded autonomic dysregulation affects blood flow and oxygen delivery [1]. Standard autonomic training focuses on heart rate or blood pressure. What if we could specifically train better **blood flow distribution** to working tissues during activity?

Potential approaches (all speculative):

- **Biofeedback:** Real-time muscle oxygenation monitoring (NIRS - near-infrared spec-

troscopy) paired with activity; patient learns to maintain tissue oxygenation

- **Blood flow restriction training:** Paradoxically, very light exercise with partial blood flow restriction might train compensatory mechanisms; used in rehabilitation but untested in ME/CFS
- **Postural countermeasures:** Physical medicine approaches from POTS treatment (leg crossing, muscle tensing) might improve orthostatic blood redistribution

Undocumented: Muscle/brain tissue oxygenation during and after exercise has not been systematically measured in ME/CFS using NIRS or similar techniques. This would reveal whether oxygen delivery failure is indeed occurring and where (central vs peripheral).

⚠ **Limitation 6: CPET-Derived Hypotheses: Untested Mechanistic Inferences from Functional Data**

The two-day CPET protocol [1] provides robust evidence of *functional* deterioration (reduced VO₂max, workload, and ventilatory threshold on Day 2). However, the mechanistic hypotheses built upon this finding have not been directly tested:

- The “autonomic-mitochondrial feedback loop,” “mitochondrial turnover rate limitation,” and “circadian recovery gating” are interpretive models imposed on the 13-day recovery observation—the recovery duration alone cannot distinguish between these and simpler explanations (e.g., delayed inflammatory resolution, psychological anticipation).
- All supplement protocols proposed (autonomic-mitochondrial co-support, accelerated mitochondrial turnover, chronotherapy) are speculative combinations that have not been tested even in pilot studies; individual components have varying evidence levels, but the specific combinations and rationales are hypothetical.
- The “metabolic preconditioning” hypothesis directly contradicts established pacing principles and carries risk of harm; it is included as a theoretical exercise, not a recommendation.
- Quantitative model parameters (13-day recovery, mitochondrial half-life estimates) are drawn from healthy-population physiology and may not apply to ME/CFS patients with altered mitochondrial dynamics.

14.19.8 Summary Table: Novel Hypotheses from CPET Findings

Table 14.3 summarizes the mechanistic hypotheses and treatment implications emerging from two-day CPET evidence, ranked by likelihood and therapeutic potential.

Evidence level definitions:

- **Very Low:** Purely theoretical; no supporting evidence in ME/CFS
- **Low:** Mechanism plausible; analogous evidence from other conditions
- **Low-Moderate:** Mechanism plausible; some supportive but indirect ME/CFS evidence

Table 14.3: Novel hypotheses arising from two-day CPET findings, ranked by plausibility and therapeutic potential

Hypothesis	Evidence	Therapeutic Potential	Key Prediction	Nearest-Term Test
Autonomic-mitochondrial feedback loop	Moderate	High	Synergy between catecholamine support + antioxidants exceeds either alone	3-month trial: tyrosine + BH4 + MitoQ + N-Acetylcysteine (NAC) vs. components
Mitochondrial turnover rate limitation	Moderate-High	Moderate-High	Urolithin A + NAD+ precursors shorten recovery time	Repeat 2-day CPET after 12 weeks urolithin A/NMN
Circadian recovery gating	Low-Moderate	Moderate	Evening mitophagy enhancers + morning biogenesis support outperform mistimed dosing	Crossover trial: timed vs. untimed supplementation
Exercise metabolomics-guided therapy	Moderate	Very High	Individual metabolic signatures predict treatment response	Metabolomics at 0, 0.5, 2, 6h post-CPET; cluster patients
Vagal stimulation for recovery	Low-Moderate	Moderate	taVNS post-exertion reduces PEM severity and shortens duration	Post-exertion VNS vs. sham; symptom tracking 7 days
Blood flow redistribution training	Low	Low-Moderate	NIRS-guided biofeedback improves tissue oxygenation during activity	NIRS monitoring during standardized activity ±biofeedback training
Metabolic preconditioning (hormesis)	Very Low	Low (High Risk)	Brief sub-threshold stress improves Day 2 CPET metrics	NOT RECOMMENDED without extensive safety data

- **Moderate:** Mechanism supported by multiple ME/CFS findings; direct intervention untested
- **Moderate-High:** Strong mechanistic support; similar interventions show benefit
- **High:** Direct evidence from ME/CFS studies

Therapeutic potential considers both magnitude of potential benefit and safety/accessibility profile.

14.20 Novel Hypotheses from 2026 Autoimmune Research

The recent convergence of autoantibody research, EBV pathogenesis studies, and structural biology of receptor-targeting antibodies suggests several novel hypotheses that may explain ME/CFS pathophysiology and point toward new therapeutic strategies.

14.20.1 The EBV-B Cell CNS Infiltration Hypothesis

? Open Question 42: Viral-Driven Autoreactive B Cell Brain Invasion

The Pless et al. (2026) study [164] demonstrated that autoreactive B cells exist in healthy human blood and can cross the blood-brain barrier following viral infection. When these B cells express EBV Latent Membrane Protein 1 (LMP1), they infiltrate the brain and induce demyelinating lesions through myelin antigen capture, complement activation, and microglial activation.

What if a similar mechanism operates in ME/CFS—not necessarily causing overt demyelination, but producing subclinical neuroinflammation and autoantibody-mediated neurological dysfunction?

Proposed mechanism:

1. EBV infection (primary or reactivation) triggers LMP1 expression in a subset of B cells
2. LMP1-expressing B cells acquire enhanced blood-brain barrier crossing ability
3. These B cells infiltrate the CNS and encounter neuronal antigens
4. Unlike MS (where myelin antigens are targeted), ME/CFS B cells might target:
 - Neurotransmitter receptors (explaining catecholamine/serotonin dysfunction)
 - Ion channels (explaining autonomic symptoms)
 - Astrocyte or microglial surface proteins (causing neuroinflammation)
5. Local complement activation and microglial priming create chronic neuroinflammation
6. The neuroinflammation produces brain fog, altered effort perception, and sensory sensitivities

This would explain why ME/CFS often follows EBV infection, why neuroinflammation is seen on PET imaging, and why CSF abnormalities are documented despite

relatively normal standard testing.

Undocumented Phenomenon. CSF analysis for LMP1-expressing B cells or EBV-specific B cell populations has not been performed in ME/CFS. If this hypothesis is correct:

- ME/CFS patients should have elevated EBV-infected B cells in CSF compared to controls
- These B cells might show LMP1 expression
- Local complement activation products should be detectable
- Microglial activation markers should correlate with presence of these B cells

Treatment Implication. If EBV-infected B cells are driving CNS pathology:

- Antiviral therapy (valacyclovir, valganciclovir) might reduce EBV reactivation and LMP1 expression
- B cell depletion with rituximab might be beneficial *if* the infiltrating B cells are CD20⁺ (unlike plasma cells)
- Complement inhibition might reduce downstream damage
- EBV-specific T cell therapy (experimental) might eliminate the infected B cell population

14.20.2 The GPCR Autoantibody-Monocyte Amplification Loop

? Open Question 43: Autoantibodies as Monocyte Programmers

Hackel et al. (2025) [209] demonstrated that GPCR autoantibodies don't just block or activate receptors—they reprogram monocyte function, causing production of specific inflammatory and neurotrophic cytokines (MIP-1 δ , PDGF-BB, TGF- β 3).

This suggests autoantibodies may have effects far beyond simple receptor modulation. What if GPCR autoantibodies create a self-amplifying inflammatory loop through monocyte reprogramming?

Proposed mechanism:

1. Initial infection triggers GPCR autoantibody production
2. Autoantibodies bind to monocyte surface GPCRs
3. Monocyte signaling pathways are reprogrammed, shifting cytokine production
4. MIP-1 δ recruits additional immune cells to tissues
5. PDGF-BB promotes fibroblast activation and tissue remodeling
6. TGF- β 3 has complex immunomodulatory effects (potentially tolerogenic, but also fibrotic)
7. The altered cytokine milieu:
 - Maintains B cell activation (perpetuating autoantibody production)
 - Creates tissue-level inflammation (explaining multi-system symptoms)

- Affects endothelial function (connecting to vascular hypothesis)
 - Signals to the brain via vagal afferents or direct cytokine action
8. Unlike simple autoantibody-receptor binding (which might be compensated), monocyte reprogramming creates sustained systemic effects

Undocumented Phenomenon. The specific downstream targets of the altered cytokine profile have not been mapped in ME/CFS. Predictions:

- Tissue biopsies should show increased fibroblast activation markers
- MIP-1 δ -responsive immune cell populations should be expanded
- TGF- β 3-associated gene expression signatures should be detectable
- Monocyte cytokine production profiles should correlate with symptom severity

Treatment Implication.

- Autoantibody removal (immunoadsorption, BC007) should normalize monocyte function
- Targeting the downstream cytokines (anti-MIP-1, anti-PDGF) might provide symptomatic relief even if autoantibodies persist
- Monocyte-modulating therapies (JAK inhibitors affecting monocyte signaling) might interrupt the loop
- Combined autoantibody removal + monocyte modulation might be synergistic

14.20.3 The Receptor Internalization Hypothesis

? Open Question 44: Autoantibodies Causing Functional Receptor Depletion

The Kim et al. (2026) cryo-EM study [211] of NMDA receptor autoantibodies revealed that autoantibody binding causes receptor internalization—removing functional receptors from the cell surface. This isn't receptor blocking; it's receptor elimination.

If GPCR autoantibodies in ME/CFS cause similar internalization, patients might have functional receptor depletion rather than receptor dysfunction. The receptors aren't blocked—they're gone.

Proposed mechanism:

1. Autoantibodies bind to β -adrenergic and muscarinic receptors
2. Rather than simply blocking or activating receptors, binding triggers receptor endocytosis
3. Internalized receptors may be degraded rather than recycled
4. Cells experience progressive receptor depletion
5. With fewer receptors, normal catecholamine/acetylcholine signaling becomes ineffective

fective

6. This explains the autonomic dysfunction without requiring abnormal neurotransmitter levels
7. It also explains why symptoms persist: receptor resynthesis takes time, and if autoantibodies persist, new receptors are immediately internalized

This mechanism would create a fundamentally different pathophysiology than simple receptor blockade—one that persists as long as autoantibodies are present and requires receptor regeneration (not just antibody clearance) for recovery.

Undocumented Phenomenon. Receptor density on patient cells has not been systematically measured. Predictions:

- β -adrenergic receptor density on patient lymphocytes should be reduced
- Muscarinic receptor density on relevant tissues should be depleted
- Receptor density should correlate inversely with autoantibody titers
- After autoantibody removal (immunoadsorption), receptor density should gradually recover over weeks to months
- The time course of receptor recovery should parallel symptom improvement

Treatment Implication.

- Autoantibody removal is necessary but not sufficient—receptor regeneration takes time
- Receptor upregulation strategies (if they exist) might accelerate recovery
- The lag between autoantibody clearance and symptom improvement is explained
- Combined approaches: remove autoantibodies + support receptor resynthesis

14.20.4 The Antigenic Hotspot Vulnerability Hypothesis

? Open Question 45: Structural Vulnerability to Autoimmune Attack

The Kim et al. cryo-EM study [211] identified specific “antigenic hotspots” on the NMDA receptor where autoantibodies preferentially bind. These aren’t random locations—they’re structurally exposed regions that the immune system can access.

What if certain individuals have GPCR variants with more exposed antigenic hotspots—making them structurally predisposed to autoimmune attack on these receptors?

Proposed mechanism:

1. GPCR genes show normal polymorphic variation in the population
2. Some variants have amino acid changes in extracellular loops
3. These changes create more immunogenic conformations—“hotspots” that B cells

can target

4. When an infection triggers autoantibody production (through molecular mimicry or bystander activation), individuals with hotspot-exposed receptors are more likely to develop pathogenic autoantibodies
5. This would explain:
 - Why only some people develop ME/CFS after infection
 - Why certain families show clustering of ME/CFS
 - Why symptom patterns vary (different receptors have different vulnerabilities)
 - Why autoantibody titers don't perfectly correlate with symptoms (some autoantibodies target more critical hotspots than others)

Undocumented Phenomenon. GPCR genetic variation in ME/CFS has been minimally studied. Predictions:

- ME/CFS patients should show enrichment for specific GPCR variants
- These variants should map to extracellular domains (potential hotspots)
- Structural modeling should predict increased immunogenicity for these variants
- Autoantibody binding affinity should be higher for “hotspot” variants

Treatment Implication.

- Genetic screening might identify at-risk individuals before infection
- Prophylactic approaches (EBV vaccination when available) might prevent ME/CFS in susceptible individuals
- Personalized therapy based on which receptors are structurally vulnerable
- Potential for peptide-based tolerization targeting specific hotspots

14.20.5 The Molecular Mimicry-Receptor Homology Hypothesis

? Open Question 46: Viral Proteins Mimicking Receptor Epitopes

EBV is strongly associated with ME/CFS onset. EBV proteins share sequence homology with many human proteins (documented extensively in MS research). What if specific EBV proteins share structural homology with GPCR extracellular domains—such that anti-EBV antibodies cross-react with adrenergic and muscarinic receptors?

Proposed mechanism:

1. EBV infection generates robust antibody response against viral proteins
2. Certain EBV proteins (particularly those exposed on infected cell surfaces) share epitopes with human GPCRs
3. Anti-EBV antibodies cross-react with β -adrenergic and muscarinic receptors

4. Unlike true autoantibodies (generated by tolerance breach), these are antiviral antibodies with unfortunate cross-reactivity
5. As long as EBV persists (which it does, lifelong), the anti-EBV response continues
6. This maintains GPCR-targeting antibodies indefinitely

This would explain why EBV infection so specifically triggers ME/CFS, why autoantibody titers persist, and why antiviral therapy might help (by reducing viral protein expression and thus the stimulus for cross-reactive antibodies).

Undocumented Phenomenon. Structural homology between EBV proteins and GPCR extracellular domains has not been systematically analyzed. Predictions:

- Computational analysis should identify EBV-GPCR homologous sequences
- Anti-EBV antibodies should show GPCR binding in vitro
- The same antibody clones should bind both EBV proteins and GPCRs
- Patients with higher anti-EBV titers might have higher anti-GPCR titers
- Reducing EBV viral load should reduce GPCR autoantibody titers

Treatment Implication.

- Aggressive antiviral therapy might reduce the stimulus for cross-reactive antibodies
- EBV vaccination (when available) might prevent ME/CFS by generating non-cross-reactive immunity
- Targeted B cell depletion of EBV-specific clones might eliminate the cross-reactive population
- Tolerization to the shared epitope might break the cycle

14.20.6 The Dual-Compartment Autoantibody Hypothesis

? Open Question 47: Peripheral vs. Central Autoantibody Effects

Bynke et al. (2020) [72] found elevated GPCR autoantibodies in plasma but *not* in CSF. This is usually interpreted as indicating peripheral origin. But what if it reveals something more important: different autoantibody populations in different compartments, with different effects?

Proposed mechanism:

1. Peripheral plasma cells produce GPCR autoantibodies that cause systemic symptoms:
 - Cardiovascular autonomic dysfunction (acting on vascular/cardiac receptors)
 - GI symptoms (acting on enteric receptors)
 - Peripheral muscle effects

2. Separately, EBV-infected B cells that cross the blood-brain barrier might produce *different* autoantibodies locally in the CNS:
 - These might target neuronal receptors (NMDA, GABA, glycine)
 - They would cause cognitive and neurological symptoms
 - They might not appear in lumbar puncture CSF if produced in specific brain regions
3. The two compartments explain the dissociation between peripheral and central symptoms
4. Treatment targeting only peripheral autoantibodies might improve systemic symptoms but leave cognitive symptoms unchanged

Undocumented Phenomenon. Regional CNS autoantibody production has not been studied in ME/CFS. Predictions:

- Post-mortem or surgical brain tissue might show local autoantibody production
- Advanced CSF sampling (ventricular vs. lumbar) might reveal different autoantibody profiles
- Intrathecal B cell populations might differ from peripheral B cells
- Patients with predominantly cognitive symptoms might have different autoantibody patterns than those with predominantly autonomic symptoms

Treatment Implication.

- Immunoadsorption might help peripheral but not CNS symptoms
- CNS-penetrant therapies might be needed for cognitive symptoms
- Combination approaches targeting both compartments might be necessary
- Biomarkers distinguishing peripheral vs. central autoimmunity would guide therapy

14.20.7 The Autoantibody Functional Assay Discrepancy Hypothesis

? Open Question 48: Why Do Some Studies Fail to Replicate?

Vernino et al. (2022) [210] found no differences in GPCR autoantibodies between POTS patients and controls using standard ELISA, directly contradicting multiple positive studies. This methodological controversy has major implications.

What if both findings are correct—but measuring different things?

Proposed mechanism:

1. ELISA detects any antibody that binds the target antigen
2. Most humans have low-level autoantibodies against many self-proteins (natural autoantibodies)

3. These natural autoantibodies are non-pathogenic
4. Pathogenic autoantibodies differ in:
 - Binding affinity (higher affinity = more functional effect)
 - Epitope specificity (some epitopes are functionally important, others aren't)
 - Effector function (some trigger internalization, others don't)
 - Isotype (IgG1/IgG3 activate complement; IgG4 doesn't)
5. Standard ELISAs detect total binding antibodies, not functionally pathogenic ones
6. Positive studies using CellTrend assays might detect a subset that correlates with pathogenicity
7. Negative studies using different methodology might detect the non-pathogenic background

This would mean: autoantibodies ARE involved in ME/CFS, but detecting the pathogenic subset requires functional assays, not just binding assays.

Undocumented Phenomenon. Functional characterization of ME/CFS autoantibodies is minimal. Predictions:

- Functional assays (receptor internalization, downstream signaling) should distinguish patients from controls better than binding assays
- Autoantibody affinity should correlate with symptom severity
- Epitope mapping should identify “pathogenic” vs. “non-pathogenic” binding sites
- Isotype profiling might reveal skewing toward complement-activating subclasses in patients

Treatment Implication.

- Functional autoantibody assays should be developed for patient selection
- Therapies might need to target specifically the high-affinity pathogenic subset
- Understanding functional differences could guide epitope-specific tolerization
- Clinical trials should stratify by functional autoantibody status, not just binding titers

14.20.8 Updated Master Hypothesis Table: 2026 Autoimmune Hypotheses

Table 14.4 summarizes the novel hypotheses emerging from 2026 autoimmune research.

Table 14.4: Novel hypotheses from 2026 autoimmune research

Hypothesis	Evidence Level	Therapeutic Potential	Benefit: Mild	Benefit: Severe	Explains Key Features	Nearest-Term Action
<i>EBV-Related Hypotheses</i>						
EBV-B cell CNS infiltration	Low-Moderate	High	Moderate	Moderate-High	Post-EBV onset; neuroinflammation; brain fog; cognitive symptoms distinct from fatigue	CSF B cell analysis; EBV PCR in CSF; LMP1 expression profiling
Molecular mimicry (EBV-GPCR homology)	Low	High	Moderate-High	Moderate-High	EBV trigger specificity; persistent autoantibodies; why antivirals might help	Computational homology analysis; cross-reactivity testing
<i>Autoantibody Mechanism Hypotheses</i>						
Autoantibody-monocyte amplification loop	Moderate	High	High	Moderate	Systemic inflammation; cytokine abnormalities; why symptoms persist beyond receptor effects	Monocyte functional profiling post-immunoadsorption
Receptor internalization (not blockade)	Low-Moderate	Moderate-High	Moderate	Moderate	Lag between antibody removal and improvement; why symptoms persist; receptor density changes	Receptor density assays on patient lymphocytes
Antigenic hotspot vulnerability	Very Low	Moderate	Moderate	Moderate	Genetic susceptibility; family clustering; why some people but not others	GPCR genetic screening; structural immunogenicity prediction
Dual-compartment autoantibodies	Low	High	Moderate-High	Moderate-High	Dissociation between peripheral and cognitive symptoms; why some treatments help some symptoms	Regional CSF sampling; post-mortem tissue analysis
Functional vs. binding assay discrepancy	Moderate	Very High	High	High	Failed replications; heterogeneous treatment response; why some high-titer patients don't respond	Develop functional autoantibody assays; stratify trials
<i>Combined/Integrated Hypotheses</i>						
EBV → LMP1 → BBB crossing → CNS autoimmunity	Low-Moderate	Very High	Moderate-High	High	Full pathway from trigger to CNS symptoms; explains post-viral onset, neuroinflammation, autoantibodies	Integrated CSF + peripheral analysis; antiviral + immunotherapy trials
Plasma cell + monocyte dual targeting	Moderate	Very High	High	Moderate-High	Why single-target therapies partially work; need for combination approaches	Daratumumab + monocyte modulator (e.g., JAK inhibitor) trial

‡ **Limitation 7: 2026 Autoimmune Hypotheses: Extrapolation from Adjacent Diseases**

The hypotheses in this section extrapolate from findings in other autoimmune conditions (MS, autoimmune encephalitis, POTS) to ME/CFS. Key epistemic boundaries:

- The EBV-B cell CNS infiltration hypothesis is based on MS research [164]; whether LMP1-expressing B cells cross the blood-brain barrier in ME/CFS patients has not been investigated.
- The receptor internalisation mechanism derives from NMDA receptor cryo-EM studies [211]; whether GPCR autoantibodies in ME/CFS cause receptor internalisation (rather than blocking or agonism) is unknown.
- The “antigenic hotspot vulnerability” and “molecular mimicry” hypotheses generate testable predictions but are currently untested computational predictions, not empirical findings.
- The unified EBV-autoimmune cascade model integrates seven individually uncertain hypotheses into a single narrative; the integrated model inherits the uncertainty of all components and adds additional uncertainty from the assumed causal connections between them.

14.20.9 Integration: A Unified EBV-Autoimmune Model

Drawing together these hypotheses, a coherent model emerges:

~ **Hypothesis 6: The EBV-Autoimmune Cascade Model**

ME/CFS may result from a cascade initiated by EBV infection in genetically susceptible individuals:

1. **Trigger:** EBV infection (primary or reactivation) in an individual with GPCR variants containing exposed antigenic hotspots
2. **Molecular mimicry:** Anti-EBV antibodies cross-react with homologous GPCR epitopes, or bystander activation generates true autoantibodies
3. **Peripheral effects:** GPCR autoantibodies cause receptor internalization on cardiovascular, GI, and peripheral tissues, producing autonomic dysfunction
4. **Monocyte reprogramming:** Autoantibody binding to monocyte GPCRs triggers altered cytokine production (MIP-1 δ , PDGF-BB, TGF- β 3), creating systemic inflammation
5. **CNS invasion:** EBV-infected B cells expressing LMP1 cross the blood-brain barrier and either produce local autoantibodies or trigger complement/microglial activation
6. **Plasma cell establishment:** Some autoreactive B cells differentiate into long-lived plasma cells in bone marrow sanctuaries, ensuring persistent autoantibody production
7. **Stable pathological state:** The combination of peripheral autoantibody effects,

monocyte-driven inflammation, and CNS involvement creates a self-maintaining disease state that persists independent of ongoing EBV activity

8. **Treatment resistance:** Single-target therapies fail because multiple mechanisms must be addressed simultaneously:

- Antivirals alone fail: plasma cells already established
- Rituximab alone fails: plasma cells are CD20⁺
- Immunoadsorption alone fails: plasma cells regenerate antibodies
- Daratumumab alone partially works: addresses plasma cells but not CNS or established receptor depletion

Evidence level: Moderate overall (components individually supported; integration speculative)

Therapeutic implication: Comprehensive treatment might require:

- Antiviral therapy (reduce ongoing EBV contribution)
- Immunoadsorption (clear existing autoantibodies)
- Daratumumab (eliminate plasma cell factories)
- Time for receptor regeneration (months post-antibody clearance)
- Possibly CNS-directed therapy for cognitive symptoms

△ **Warning 3: Speculative Integration**

This unified model is **highly speculative**. It integrates findings from multiple studies, each with limitations, and extrapolates beyond what any single study demonstrates. The model is presented not as established fact but as a framework for generating testable predictions and guiding research priorities. Clinical application of combination therapies based on this model would require rigorous testing in appropriately designed trials.

14.21 Novel Hypotheses from TRPM3 Ion Channel Research

The 2026 multi-site validation of TRPM3 ion channel dysfunction in ME/CFS [187] opens entirely new avenues for understanding disease mechanisms. TRPM3 (Transient Receptor Potential Melastatin 3) is not merely an immune cell ion channel—it is expressed across multiple tissue types and participates in diverse physiological processes. The robust, reproducible finding of TRPM3 dysfunction suggests several novel hypotheses.

14.21.1 The Paradoxical Immune State Hypothesis

? Open Question 49: Stuck Doors Explain Simultaneous Over- and Under-Activity

ME/CFS presents a puzzling immunological paradox: the immune system appears simultaneously *overactive* (chronic inflammation, elevated cytokines, persistent immune activation markers) and *underactive* (impaired NK cell cytotoxicity, poor pathogen clearance, T cell exhaustion). How can both be true?

TRPM3 dysfunction provides an elegant resolution. Consider immune cells as soldiers who can see the enemy but whose weapons won't fire:

Proposed mechanism:

1. Immune cells (NK cells, T cells) recognize pathogens or infected cells normally
2. Upon recognition, they attempt to degranulate and release cytotoxic mediators
3. Degranulation requires calcium influx through channels including TRPM3
4. With TRPM3 dysfunction ("stuck doors"), calcium influx is impaired
5. The cell cannot complete the kill—degranulation fails or is incomplete
6. The target survives; the immune cell signals for reinforcements
7. More immune cells are recruited, more activation signals are released
8. Chronic inflammation results from persistent, frustrated immune responses
9. Meanwhile, actual pathogen clearance fails, permitting viral persistence

This creates a vicious cycle: inflammation without resolution. The immune system keeps trying but never succeeds. Cytokine alarms stay elevated because the underlying threat is never neutralized. Energy is consumed in futile immune activation.

Predictions.

- NK cells from ME/CFS patients should show normal target recognition but impaired degranulation
- Calcium flux measurements during degranulation attempts should show reduced amplitude or kinetics
- Inflammatory markers should correlate with degree of TRPM3 dysfunction
- Patients with more severe TRPM3 impairment should show poorer pathogen control

14.21.2 The TRPM3-GPCR Signaling Convergence Hypothesis

? Open Question 50: Autoantibodies and Ion Channels: Connected Dysfunction

GPCR autoantibodies (anti- β_2 -adrenergic, anti-muscarinic) are documented in ME/CFS. TRPM3 dysfunction is now also documented. Are these independent abnormalities, or connected?

TRPM3 gating is modulated by G-protein signaling pathways. Muscarinic receptor activation, for example, can influence TRP channel function through phospholipase C and intracellular calcium stores. If autoantibodies are chronically dysregulating GPCR signaling, they might indirectly cause or exacerbate TRPM3 dysfunction.

Possible connections:

- GPCR autoantibodies → aberrant second messenger signaling → altered TRPM3 phosphorylation → channel dysfunction
- Chronic receptor stimulation → depletion of PIP₂ (required for TRP channel function) → reduced TRPM3 activity
- Autoantibody-induced receptor internalization → loss of TRPM3-regulating GPCR pathways → unregulated channel states
- Alternatively: shared autoimmune targeting of GPCRs and ion channels

If GPCR dysfunction and TRPM3 dysfunction are linked, therapies targeting autoantibodies (immunoadsorption, BC007, daratumumab) might restore both GPCR signaling and TRPM3 function.

Testable predictions.

1. GPCR autoantibody titers should correlate with severity of TRPM3 dysfunction
2. Removal of autoantibodies should improve TRPM3 function measurements
3. *In vitro*, adding ME/CFS patient IgG to healthy cells should impair TRPM3 responses
4. TRPM3 agonists might partially rescue function even in presence of autoantibodies

14.21.3 The Systemic Channelopathy Hypothesis

? Open Question 51: TRPM3 Dysfunction Beyond Immune Cells

The Sasso et al. study demonstrated TRPM3 dysfunction specifically in *immune cells*. However, TRPM3 is not limited to immune cells—it is expressed in:

- Sensory neurons (particularly nociceptors)
- Pancreatic β -cells (insulin secretion)
- Vascular smooth muscle
- Kidney epithelium

- Brain (various regions)
- Retinal ganglion cells

What if TRPM3 dysfunction in ME/CFS is *systemic*—affecting all tissues where the channel is expressed?

Predicted consequences by tissue:

Sensory neurons:

- TRPM3 is a heat and pain sensor
- Dysfunction could cause: altered temperature perception, cold intolerance, heat intolerance, hyperalgesia, allodynia
- The characteristic sensory hypersensitivities of ME/CFS might be direct TRPM3 effects

Pancreatic β -cells:

- TRPM3 modulates insulin secretion
- Dysfunction could cause: reactive hypoglycemia, postprandial symptoms, glucose intolerance
- Many ME/CFS patients report blood sugar instability

Vascular smooth muscle:

- TRPM3 affects vascular tone
- Dysfunction could cause: abnormal blood pressure regulation, orthostatic intolerance
- Connects to POTS and orthostatic symptoms

Brain:

- TRPM3 in neurons affects excitability
- Dysfunction could cause: cognitive impairment, altered neurotransmission
- May contribute to “brain fog” directly, not just via inflammation

If TRPM3 dysfunction is systemic, ME/CFS is fundamentally a **channelopathy**—a disease of ion channel function affecting multiple organ systems simultaneously.

Research implications.

- TRPM3 function should be tested in multiple cell types from ME/CFS patients
- Symptoms should cluster by TRPM3-expressing tissues
- Treatments restoring TRPM3 function might address multiple symptom domains simultaneously

14.21.4 The “Wired but Tired” Ion Channel Explanation**~ Hypothesis 7: Bidirectional Channel Dysfunction Creates Paradoxical State**

The “wired but tired” phenomenon—feeling simultaneously exhausted and unable to relax—is a hallmark of ME/CFS. Ion channel dysfunction offers a mechanistic explanation:

Proposed mechanism:

1. The Sasso et al. study found TRPM3 dysfunction characterized as channels that fail to allow adequate calcium entry (“stuck doors”). However, ion channel dysfunction can theoretically manifest in multiple ways:
 - Stuck closed → inability to respond to physiological stimuli (consistent with the study findings)
 - Stuck partially open → chronic low-level calcium leak (speculative alternative)
 - Altered gating kinetics → inappropriate timing of responses
2. In sensory neurons, a partially open channel would cause:
 - Baseline hyperexcitability
 - Lowered activation thresholds
 - Spontaneous firing → restlessness, hypersensitivity
3. In immune and muscle cells, impaired channel response would cause:
 - Failed energy-requiring processes
 - Calcium-dependent enzyme dysfunction
 - Fatigue and weakness
4. The same patient has hyperactive sensory processing (“wired”) AND dysfunctional effector mechanisms (“tired”)

This is not contradictory—it is the expected result of ion channel dysfunction affecting excitable and effector cells differently. The nervous system is overexcitable while the muscular and immune systems are underpowered.

14.21.5 The Calcium-Mitochondria Cascade Hypothesis**? Open Question 52: TRPM3 Dysfunction Upstream of Mitochondrial Failure**

Mitochondrial dysfunction is well-documented in ME/CFS: impaired oxidative phosphorylation, reduced ATP production, abnormal metabolomics. But is mitochondrial dysfunction primary or secondary?

Calcium and mitochondria are intimately linked:

- Mitochondria buffer cytosolic calcium
- Mitochondrial calcium uptake regulates the TCA cycle and oxidative phosphorylation

- Calcium signals promote ATP synthesis by activating matrix dehydrogenases
- Both calcium overload and calcium depletion impair mitochondrial function

What if TRPM3 dysfunction *causes* mitochondrial dysfunction?

Proposed mechanism:

1. TRPM3 dysfunction alters cellular calcium handling
2. Scenario A (stuck closed): Cells cannot achieve adequate calcium transients
 - Insufficient calcium signaling to mitochondria
 - Reduced activation of calcium-dependent metabolic enzymes
 - Impaired ATP production under demand
3. Scenario B (stuck partially open): Chronic calcium leak
 - Mitochondria continuously buffer excess calcium
 - Mitochondrial calcium overload → oxidative stress
 - Gradual mitochondrial damage
4. Either scenario results in energy deficit
5. The observed mitochondrial dysfunction is downstream of ion channel dysfunction

If true, treating the mitochondria (CoQ10, ribose, carnitine) addresses symptoms but not cause. Restoring TRPM3 function would restore mitochondrial function automatically.

Predictions.

- TRPM3 dysfunction severity should correlate with mitochondrial dysfunction severity
- Restoring TRPM3 function should improve mitochondrial parameters
- Mitochondrial therapies without TRPM3 restoration should show limited, temporary benefit
- Calcium imaging during cellular stress should show abnormal patterns in ME/CFS

14.21.6 The Post-Infectious TRPM3 Acquisition Hypothesis

? Open Question 53: How Does Infection Lead to Channel Dysfunction?

If TRPM3 dysfunction is acquired after infection (as suggested by post-infectious onset of ME/CFS), what mechanism causes it?

Possible mechanisms:

Viral interference with ion channels. Some viruses directly modulate host ion channels during infection—this aids viral replication or immune evasion. If the modulation leaves persistent modifications (oxidative damage, altered phosphorylation, protein misfolding), the channel might remain dysfunctional after the virus is cleared.

Autoimmune targeting. Molecular mimicry between viral proteins and TRPM3 epitopes could generate cross-reactive antibodies or T cells. The immune response intended for the virus attacks the patient's ion channels. This would be analogous to Guillain-Barré syndrome (anti-ganglioside antibodies after *Campylobacter*) but targeting TRPM3.

Epigenetic modification. Severe infection causes oxidative and metabolic stress. This can create epigenetic marks (DNA methylation, histone modifications) affecting gene expression. TRPM3 expression or its regulatory proteins might be persistently downregulated.

Membrane composition changes. Ion channel function depends on the surrounding lipid environment. Infection-induced changes in membrane lipid composition (documented in ME/CFS) might alter TRPM3 gating properties even without changes to the protein itself.

Cofactor depletion. TRPM3 function may require specific cofactors or post-translational modifications. If infection depletes these (e.g., zinc, selenium, PIP₂), and they are not fully restored during recovery, channel function remains impaired.

Research directions.

- Screen ME/CFS patients for anti-TRPM3 autoantibodies
- Examine TRPM3 gene methylation patterns
- Test whether ME/CFS serum alters TRPM3 function in healthy cells
- Compare TRPM3 function immediately post-infection vs. established ME/CFS

14.21.7 The Temperature Dysregulation Connection

~ Hypothesis 8: TRPM3 as the Missing Link in Thermoregulation

ME/CFS patients commonly report:

- Feeling cold when ambient temperature is normal
- Inability to regulate body temperature
- Symptom flares with temperature changes
- Intolerance to both heat and cold
- Subjective fever without measurable temperature elevation

TRPM3 is a **thermosensor**—it responds to temperature changes, particularly in the warm/noxious heat range. In sensory neurons, TRPM3 contributes to heat detection and thermal pain.

Proposed mechanism:

1. Dysfunctional TRPM3 in sensory neurons provides incorrect temperature information
2. The brain receives aberrant thermosensory input
3. Thermoregulatory centers cannot properly assess or maintain body temperature
4. The patient feels cold (despite normal core temperature) or hot (without fever)
5. Thermoregulatory behaviors (seeking warmth, sweating) become maladaptive
6. Temperature instability is not an epiphenomenon but a direct consequence of TRPM3 dysfunction

This reframes temperature symptoms from “vague subjective complaints” to objective consequences of ion channel pathology.

⚠ **Limitation 8: TRPM3 Hypotheses: Single-Study Foundation with In Vitro Scope**

The TRPM3 dysfunction finding [187] is reproducible across laboratories but has important boundaries that constrain the hypotheses built upon it:

- TRPM3 dysfunction has been demonstrated only in *immune cells* (NK cells, T cells); the “systemic channelopathy” hypothesis—that TRPM3 is dysfunctional in sensory neurons, pancreatic β -cells, vascular smooth muscle, and brain—is extrapolation, not measurement.
- Whether TRPM3 dysfunction is a *cause* of ME/CFS symptoms, a *consequence* of chronic immune activation, or an *epiphenomenon* of another upstream process has not been determined by the available cross-sectional data.
- The proposed TRPM3-GPCR signalling convergence and calcium-mitochondria cascade are mechanistically plausible models, but the causal links (autoantibodies → TRPM3 dysfunction → mitochondrial failure) have not been demonstrated in any experimental system.
- Therapeutic speculation regarding TRPM3 agonists/antagonists is premature: TRPM3 pharmacology in humans is poorly characterised, and no TRPM3-modulating compound has been tested in ME/CFS patients.

14.21.8 TRPM3-Targeted Therapeutic Speculation

If TRPM3 dysfunction is central to ME/CFS pathophysiology, targeting TRPM3 pharmacologically becomes attractive:

If channels are “stuck closed” (hypofunction):

- **TRPM3 agonists** might restore function
- Pregnenolone sulfate (endogenous neurosteroid) activates TRPM3
- CIM0216 is a potent synthetic TRPM3 agonist (research tool, not approved drug)
- Nifedipine paradoxically activates TRPM3 at certain concentrations

- **Speculation:** If TRPM3 hypofunction underlies symptoms, pregnenolone sulfate supplementation might theoretically help—but this has not been tested

If channels are “stuck open” (leak/hyperfunction):

- **TRPM3 antagonists** might restore proper gating
- Primidone (anti-epileptic) blocks TRPM3
- Certain flavonoids (naringenin, isosakuranetin) inhibit TRPM3
- **Caution:** Blocking an already dysfunctional channel might worsen symptoms

Restoring channel environment:

- Membrane lipid composition affects TRP channel function
- Omega-3 fatty acids might normalize membrane environment
- PIP₂ repletion strategies (inositol supplementation?)
- Reducing oxidative damage to channel proteins (antioxidants)

Addressing upstream causes:

- If autoantibodies cause TRPM3 dysfunction: immunoadsorption, rituximab, daratumumab
- If viral proteins interfere: antivirals
- If epigenetic: theoretically, epigenetic modifiers (speculative, no specific candidates)

△ Warning 4: Highly Speculative Therapeutics

These therapeutic ideas are **entirely speculative**. TRPM3 pharmacology in humans is poorly characterized. No clinical trials have tested TRPM3 modulators in ME/CFS. Self-experimentation with TRPM3-active compounds is not recommended. These ideas are presented to stimulate research, not to guide treatment.

14.21.9 Subtyping Implications

The TRPM3 findings may help define ME/CFS subgroups:

- **TRPM3-positive ME/CFS:** Measurable TRPM3 dysfunction; potentially responsive to ion channel modulators; may represent the “post-infectious channelopathy” subtype
- **TRPM3-negative ME/CFS:** Normal TRPM3 function; different underlying mechanism; may require different therapeutic approach
- **TRPM3 + autoantibody positive:** Combined channelopathy and autoimmune; may need immunomodulation *plus* channel restoration
- **TRPM3-positive but autoantibody-negative:** Primary ion channel pathology; direct channel therapy may suffice

This parallels the evolution of cancer treatment—from “breast cancer” to “HER2-positive breast cancer” with targeted therapy. ME/CFS may similarly fragment into molecular subtypes with tailored treatments.

14.21.10 Updated Testable Predictions from TRPM3 Research

1. **Multi-tissue TRPM3 dysfunction:** If systemic, TRPM3 impairment should be detectable in immune cells, sensory neurons, and other accessible cell types
2. **Symptom correlation:** Degree of TRPM3 dysfunction should correlate with symptom severity, particularly temperature dysregulation and sensory symptoms
3. **Autoantibody connection:** Screen for anti-TRPM3 autoantibodies; test whether GPCR autoantibody removal restores TRPM3 function
4. **Mitochondrial causality:** Longitudinal studies should show TRPM3 dysfunction precedes (or co-occurs with, not follows) mitochondrial dysfunction
5. **Pharmacological restoration:** If channel function can be restored pharmacologically, symptoms should improve
6. **Subtyping validity:** TRPM3 status should predict response to different therapeutic approaches
7. **Biomarker potential:** TRPM3 functional assays should distinguish ME/CFS patients from healthy controls and possibly from other fatigue conditions

14.22 Clinical Observation-Derived Hypotheses

The following hypotheses emerged from systematic analysis of treatment response patterns, clinical trajectories, and cross-domain pattern recognition. While speculative, each attempts to explain otherwise puzzling observations and generates testable predictions.

⚡ **Limitation 9: Brainstorm Chapter: Pattern Recognition Without Empirical Validation**

All hypotheses in this chapter were generated through clinical pattern recognition and mechanistic reasoning, not empirical data analysis. None has been tested in controlled studies. The certainty ratings are uniformly low. Pattern recognition is susceptible to confirmation bias (attending to cases that fit the pattern while ignoring those that do not) and post-hoc rationalisation (constructing mechanistic narratives to explain known clinical features). These hypotheses are presented as research stimuli, not as evidence-based conclusions.

14.22.1 The “Metabolic Runway” Theory of PEM

~ Hypothesis 9: PEM Delay Reflects Metabolic Depletion Kinetics

The characteristic 24–72 hour delay between exertion and post-exertional malaise (PEM) onset may reflect the time required for metabolic substrate pools to become critically depleted.

Proposed mechanism:

1. Exertion increases amino acid consumption (for energy, neurotransmitter synthesis, tissue repair)
2. In patients with malabsorption or metabolic dysfunction, replacement from dietary intake is impaired
3. Pool depletion follows first-order kinetics with patient-specific time constants
4. When pools fall below critical threshold, mitochondrial function fails acutely
5. Clinical PEM manifests as the metabolic “runway” runs out

Testable predictions:

- Patients with larger baseline amino acid pools should have longer PEM latency
- Pre-loading amino acids before known exertion should attenuate or delay PEM
- Serial amino acid measurements during PEM onset should show progressive depletion
- PEM severity should correlate with degree of amino acid nadir

Clinical implication: “Amino acid loading” before anticipated exertion—analogue to carbohydrate loading for endurance athletes—might extend the metabolic runway and reduce PEM severity.

△ **Warning 5: Hypothesis Limitations**

This hypothesis is mechanistically plausible but untested. The 24–72 hour delay could alternatively reflect: inflammatory cascade kinetics, gene expression changes, mitochondrial damage accumulation, or other processes. Serial metabolomic studies during controlled exertion protocols are needed to test this specific mechanism. Certainty: Low.

14.22.2 The Mast Cell “Memory” Hypothesis

~ Hypothesis 10: Epigenetic Mast Cell Sensitization

Mast cells can be epigenetically programmed by early life events, infections, and trauma. ME/CFS may represent a “mast cell memory disease” where cells remain sensitized to threats that are no longer present.

Proposed mechanism:

1. Original trigger (infection, trauma, toxic exposure) activates mast cells
2. Prolonged or intense activation induces epigenetic changes (DNA methylation, hi-

stone modification)

3. Sensitized mast cells have lower activation thresholds
4. Even after trigger removal, mast cells continue responding to minor stimuli
5. Chronic low-grade mast cell activation maintains systemic inflammation and symptoms

Supporting observations:

- MCAS commonly develops after infections or trauma
- Mast cell sensitization is documented in other conditions (mastocytosis, chronic urticaria)
- Early life adversity correlates with adult mast cell disorders
- Some patients report symptom onset after discrete triggering events with persistent symptoms despite trigger resolution

Speculative extension: Could interventions that “reset” cellular programming (psychedelics affecting serotonin receptors on mast cells, epigenetic modifiers, prolonged fasting-induced autophagy) potentially desensitize mast cells?

→ **Prediction 1: Epigenetic Mast Cell Markers in ME/CFS**

ME/CFS patients with documented mast cell activation will show elevated H3K4me3 or reduced DNA methylation at promoters of mast cell mediator genes (tryptase, histamine N-methyltransferase) compared to healthy controls, and these epigenetic marks will persist in remission.

△ **Warning 6: Hypothesis Limitations**

Mast cell epigenetics in ME/CFS has not been studied. The hypothesis extrapolates from other mast cell disorders and general epigenetic principles. No ME/CFS-specific data supports this mechanism. The “reset” speculation is highly preliminary. Certainty: Low.

14.22.3 The Vagus Nerve as “Master Regulator”

~ **Hypothesis 11: Vagal Dysfunction as Central Hub**

The vagus nerve connects gut, heart, brain, and immune system. It directly inhibits mast cells via the cholinergic anti-inflammatory pathway. Vagal dysfunction may be the central hub connecting apparently disparate Septad components.

Proposed hub structure:

- **Vagus → Mast cells:** Cholinergic anti-inflammatory pathway inhibits mast cell degranulation; vagal dysfunction → MCAS
- **Vagus → Heart:** Parasympathetic withdrawal → elevated resting HR, reduced HRV, POTS

- **Vagus → Gut:** Reduced vagal tone → decreased motility, gastroparesis, SIBO
- **Vagus → Brain:** Afferent vagal signals modulate neuroinflammation; dysfunction → brain fog, fatigue signaling
- **Vagus → Immune:** Inflammatory reflex impairment → chronic systemic inflammation

Clinical support:

- HRV is consistently reduced in ME/CFS (marker of vagal tone)
- tVNS shows preliminary benefit in some patients
- Septad conditions cluster together, suggesting common regulator
- Vagal afferents from gut may mediate “sickness behavior” in infection

Treatment implication: If vagal dysfunction is the hub, interventions restoring vagal tone (tVNS, deep breathing, cold exposure, specific probiotics) might produce multi-system improvement disproportionate to their apparent specificity.

→ **Prediction 2: Vagal Tone Correlates With Multi-System Symptoms**

Heart rate variability (HRV) indices of vagal tone (RMSSD, high-frequency power) will correlate inversely with symptom count across multiple organ systems (GI, cardiac, cognitive) in ME/CFS, and transcutaneous vagus nerve stimulation will improve ≥ 3 symptom domains simultaneously.

△ **Warning 7: Hypothesis Limitations**

While vagal involvement in ME/CFS is plausible and HRV changes are documented, no studies have demonstrated that vagal dysfunction is causal rather than consequential. The “hub” model is conceptually appealing but may oversimplify the multi-directional interactions. Certainty: Low-Medium.

14.22.4 The “Two Fuel Tanks” Hypothesis~ **Hypothesis 12: Ketones as Bypass Fuel**

Normal energy metabolism relies primarily on glucose → TCA cycle → ATP. If TCA cycle dysfunction is present in ME/CFS (as metabolomic studies suggest), ketone bodies may provide a bypass pathway.

Rationale:

1. Ketones (beta-hydroxybutyrate, acetoacetate) enter the TCA cycle downstream of several rate-limiting steps
2. Ketone metabolism does not require the full TCA cycle machinery
3. If “Tank 1” (glucose metabolism) is impaired, “Tank 2” (ketone metabolism) might remain functional

4. Providing ketones could bypass the metabolic block

Testable predictions:

- Patients with documented TCA cycle abnormalities should respond better to ketogenic interventions
- Exogenous ketones (ketone esters, MCT oil) should improve energy in TCA-dysfunction subset
- Ketogenic diet should produce improvement in some but not all ME/CFS patients (depending on defect location)
- Patients with electron transport chain (rather than TCA) defects should NOT respond to ketones

Clinical implication: Rather than difficult-to-maintain ketogenic diets, pharmaceutical exogenous ketones might provide metabolic bypass without dietary restriction.

△ **Warning 8: Hypothesis Limitations**

Ketogenic diets have not been systematically studied in ME/CFS. Anecdotal reports are mixed. The hypothesis assumes TCA dysfunction is rate-limiting, which may not be true for all patients. Ketosis can be difficult to achieve and maintain. Certainty: Low.

14.22.5 The “Protective Downregulation” Paradox

~ **Hypothesis 13: Mitochondria as Deliberate Energy Throttle**

ME/CFS mitochondria may not be “broken”—they may be deliberately downregulated as a protective response to perceived cellular danger.

Proposed mechanism:

1. Cells detect danger signals (viral proteins, DAMPs, oxidative stress, autoantibodies)
2. Danger detection triggers “cell danger response” (CDR) [197]
3. CDR includes intentional reduction in mitochondrial output to limit ROS production and conserve resources
4. The throttle is protective in acute illness but becomes pathological if chronically maintained
5. Patients experience fatigue not because mitochondria can’t produce energy, but because they’re not allowed to

Analogy: A car’s computer limiting speed when it detects a fault. The engine isn’t broken—it’s being deliberately throttled.

Radical implication: Treatments that “boost” mitochondria might be fighting the body’s protective mechanism. The correct approach would be removing the danger signal that’s triggering the throttle, allowing mitochondria to self-restore.

What might be the danger signal?

- Viral proteins from latent infection

- Autoantibodies targeting mitochondrial or cellular components
- Persistent oxidative stress from upstream dysfunction
- Gut-derived endotoxins (LPS) from barrier dysfunction

→ **Prediction 3: Mitochondrial Throttling Reversal Triggers PEM**

Pharmacological override of mitochondrial energy throttling (e.g., forced uncoupling or bypass of Complex I inhibition) will produce short-term symptom improvement followed by delayed PEM within 24–72 hours, consistent with protective shutdown being circumvented rather than resolved.

△ **Warning 9: Hypothesis Limitations and Clinical Safety**

The cell danger response hypothesis [197] is itself not fully validated. Whether ME/CFS represents a “stuck” CDR is speculative.

CRITICAL SAFETY NOTICE: This hypothesis should NOT discourage use of mitochondrial support treatments that provide symptomatic benefit. If CoQ10, carnitine, NAD⁺ precursors, or other mitochondrial interventions are helping you, **continue them**. Do not discontinue beneficial treatments based on this unvalidated hypothesis about “fighting the body’s protective mechanism.”

The hypothesis addresses root cause mechanisms, not whether symptomatic support is appropriate. Even if mitochondria are deliberately throttled, supporting their function may still improve quality of life while underlying causes are addressed. Certainty: Low.

14.22.6 The “Circadian Core” Hypothesis

~ **Hypothesis 14: Circadian Disruption as Upstream Driver**

Sleep disturbance is nearly universal in ME/CFS and usually treated as a symptom. But circadian rhythms regulate mitochondrial function, immune activity, gut motility, and HPA axis—all systems implicated in ME/CFS. What if circadian disruption is cause rather than effect?

Circadian regulation of implicated systems:

- **Mitochondria:** Have their own circadian clocks; function varies with time of day
- **Immune system:** Immune responses are time-gated; disruption impairs pathogen clearance
- **Gut motility:** Migrating motor complex is circadian-regulated
- **HPA axis:** Cortisol rhythm is fundamentally circadian
- **Autonomic balance:** Sympathetic/parasympathetic ratio follows circadian pattern

Hypothesis: A disrupted master clock (SCN dysfunction, or peripheral clock desynchronization) could produce multi-system dysfunction that manifests as ME/CFS.

Treatment implication: Aggressive circadian restoration as PRIMARY intervention:

- Morning bright light (10,000 lux within 30 minutes of waking)
- Evening blue light blocking (amber glasses after sunset)
- Strict sleep timing (same wake time daily regardless of sleep quality)
- Time-restricted eating (all food within 8–10 hour window)
- Precisely timed melatonin (0.3–0.5 mg, 5 hours before desired sleep)

This would be attempted BEFORE pharmacological interventions, testing whether clock restoration produces downstream improvement.

→ **Prediction 4: Circadian Restoration Precedes Immune Normalization**

Interventions that restore circadian cortisol rhythm (timed bright light, melatonin, chronotherapy) will produce measurable normalization of diurnal cytokine profiles (IL-6, TNF- α) within 4–8 weeks, preceding and predicting subsequent symptom improvement.

△ **Warning 10: Hypothesis Limitations**

Circadian disruption in ME/CFS is documented but causality is not established. Severely ill patients may have limited ability to implement circadian interventions (cannot tolerate light, cannot maintain schedules). The hypothesis does not explain post-infectious onset. Certainty: Low-Medium.

14.22.7 The “Microclot” Bridge Hypothesis

~ **Hypothesis 15: Capillary Occlusion as Final Common Pathway**

Emerging Long COVID research has identified microclots—fibrin deposits that occlude capillaries—as a potential mechanism. If capillaries are blocked, oxygen delivery fails regardless of mitochondrial health.

How microclots could explain ME/CFS features:

- **Fatigue:** Tissues receive inadequate oxygen; mitochondria can’t function
- **PEM worsening with exercise:** Increased oxygen demand, same blocked delivery
- **Improvement lying down:** Gravity-assisted perfusion through partially occluded capillaries
- **Brain fog:** Cerebral microvasculature particularly vulnerable to perfusion deficits
- **POTS correlation:** Microvascular dysfunction contributes to orthostatic intolerance

Connecting to other mechanisms:

- Viral infection can trigger coagulation abnormalities
- Mast cell activation releases pro-coagulant factors
- Endothelial dysfunction (from NO deficiency) promotes clot formation
- Autoantibodies can target clotting factors

△ Warning 11: Treatment Safety: Coagulation Interventions

All listed interventions carry significant risks and require medical supervision:

- **Anticoagulants:** Bleeding risk requiring regular laboratory monitoring (INR, aPTT); contraindicated with many medications and medical conditions
- **Nattokinase:** Despite “natural” label, has anticoagulant effects; risk of bleeding, drug interactions; not FDA-approved for medical use
- **Plasmapheresis:** Invasive procedure requiring medical facility; risks include infection, bleeding, hypotension, allergic reactions
- **Hyperbaric oxygen:** Specialized equipment required; risks include barotrauma, oxygen toxicity, claustrophobia

None of these interventions should be attempted without physician supervision. Self-treatment with anticoagulants is dangerous and potentially life-threatening.

Treatment implications (speculative research hypotheses):

- Anticoagulants (physician monitoring essential)
- Nattokinase (fibrinolytic enzyme; still carries bleeding risk)
- Plasmapheresis (medical facility procedure only)
- Hyperbaric oxygen (specialized treatment centers)

→ Prediction 5: Microclot Burden Correlates With Exercise Intolerance

Fluorescence microscopy-quantified microclot burden in platelet-poor plasma will correlate with Day 2 CPET decrement ($r > 0.4$), and triple anticoagulation (aspirin + clopidogrel + apixaban) will reduce microclot burden by >50% with corresponding improvement in exercise capacity at 8 weeks.

△ Warning 12: Hypothesis Limitations

Microclots have been documented in Long COVID but not systematically studied in pre-pandemic ME/CFS. The overlap between Long COVID and ME/CFS is significant but not complete. Anticoagulant therapy carries bleeding risks. No controlled trials support these interventions in ME/CFS. Certainty: Low.

14.22.8 The “Infection Doesn’t Matter” Hypothesis

~ Hypothesis 16: Susceptibility Over Pathogen

ME/CFS can be triggered by remarkably diverse infections: EBV, COVID-19, Lyme disease, Q fever, Ross River virus, giardia, and others. What if the specific infection is largely irrelevant, and what matters is host susceptibility?

Proposed model:

1. Certain individuals have pre-existing susceptibility factors:
 - Connective tissue variants (hypermobility genes)
 - Mast cell activation tendency
 - Mitochondrial polymorphisms
 - Immune response patterns (cytokine profiles)
2. ANY sufficient immune challenge can trigger the cascade in susceptible individuals
3. The infection is the **match**; the susceptibility is the **gasoline**
4. Post-infection, the pathogen may be irrelevant—the dysregulated state is self-maintaining

Implication: Stop searching for “the” ME/CFS pathogen. Instead, identify the susceptibility factors that determine who develops ME/CFS after common infections.

Testable prediction: Genetic studies should find ME/CFS associations with genes affecting mast cells, connective tissue, mitochondria, and immune regulation rather than pathogen-specific response genes.

Prevention implication: If susceptibility factors can be identified, high-risk individuals could receive prophylactic interventions during acute infections (aggressive mast cell stabilization, circadian protection, metabolic support) to prevent ME/CFS development.

Δ **Warning 13: Hypothesis Limitations**

This hypothesis does not explain why some infections (EBV, COVID) seem more likely to trigger ME/CFS than others (rhinovirus, norovirus). Susceptibility factors have not been identified with certainty. The hypothesis may be partially true (susceptibility matters) while specific pathogen factors also contribute. Certainty: Low-Medium.

14.22.9 Female Predominance: Hormonal Amplification

~ Hypothesis 17: Estrogen as Cascade Amplifier

Women are 3–4× more likely to develop ME/CFS than men. While often attributed to general “autoimmunity is more common in women,” the cascade model suggests a more specific mechanism: estrogen amplifies multiple steps.

Estrogen effects on implicated pathways:

- **Mast cells:** Estrogen increases mast cell activation and histamine release

- **Connective tissue:** Estrogen affects collagen synthesis and tissue laxity (hypermobility)
- **Gut permeability:** Estrogen modulates tight junction proteins
- **Immune response:** Estrogen shifts toward Th2/autoimmune-prone patterns
- **Pain processing:** Estrogen affects central sensitization

Testable predictions:

- ME/CFS symptom severity should fluctuate with menstrual cycle (reported anecdotally)
- Onset or worsening may cluster around hormonal transitions (puberty, postpartum, perimenopause)
- Some patients may improve after menopause (reduced estrogen)
- Hormonal modulation (progesterone, Dehydroepiandrosterone (DHEA), careful estrogen management) might be therapeutic

Clinical observation: Many patients report perimenstrual worsening (days −3 to +2 around menstruation), consistent with hormonal involvement.

△ **Warning 14: Hypothesis Limitations**

Sex hormone studies in ME/CFS are limited and inconsistent. The hypothesis does not explain male ME/CFS cases or post-menopausal onset. Hormonal interventions are complex and can have significant side effects. Certainty: Low-Medium.

14.22.10 The “Bistable Equilibrium” and “Reset” Concept

Speculation 36 (ME/CFS as Stable Dysfunctional State). ME/CFS may represent a **stable but dysfunctional equilibrium**—the body “stuck” in a local energy minimum, unable to spontaneously return to health.

Energy landscape analogy:

- Health is a deep well (stable, low-energy state)
- ME/CFS is a shallow well (also stable, but suboptimal)
- A “hill” (energy barrier) separates the two states
- Gradual treatments may improve symptoms within the ME/CFS well but not escape it
- Escaping may require a “kick”—temporary destabilization to cross the barrier

△ **Warning 15: Critical Safety Notice: Dangerous Interventions**

The following “reset” interventions are **DANGEROUS**, especially for metabolically fragile ME/CFS patients. These approaches:

- Must **ONLY** be attempted under direct medical supervision in controlled research settings

- Are NOT validated by clinical trials in ME/CFS
- May be life-threatening if attempted through self-experimentation
- Could cause irreversible harm or death in vulnerable patients

DO NOT attempt these interventions outside institutional review board-approved research protocols.

Potential “reset” interventions (RESEARCH HYPOTHESES ONLY):

- **Extended fasting** (72+ hours): Could trigger dangerous hypoglycemia, electrolyte imbalances, or metabolic crisis in ME/CFS patients with existing energy metabolism dysfunction
- **Controlled hyperthermia**: Risk of cardiovascular collapse, dehydration, heat stroke; historical use does not validate safety
- **Plasmapheresis**: Invasive procedure requiring medical facility; risks include infection, bleeding, hypotension
- **High-dose IVIG**: Requires intravenous access and monitoring; risk of allergic reactions, aseptic meningitis, thrombosis
- **Stellate ganglion block**: Invasive procedure with risks including pneumothorax, nerve injury, stroke
- **Psychedelics**: Uncontrolled use risks psychiatric crisis, cardiovascular events; legal restrictions apply

The “reset” concept is a metaphor, not validated biophysical mechanism. These interventions remain entirely experimental and should not encourage desperate self-experimentation that could result in severe harm.

△ Warning 16: Hypothesis Limitations

The bistable equilibrium model is a metaphor, not a validated biophysical description. “Reset” interventions are largely untested in ME/CFS and carry significant risks. Extended fasting could be dangerous for malnourished or metabolically compromised patients. This hypothesis should not encourage desperate self-experimentation. Certainty: Very Low.

≠ Limitation 10: Bistable Model: Metaphor, Not Mechanism

The “energy landscape” with “wells” and “hills” is a mathematical metaphor from dynamical systems theory, not a description of identified biological states. No study has demonstrated that ME/CFS represents a distinct stable equilibrium in any measurable state space (metabolic, immune, or neurological). The existence of a second attractor (the “health” state) separated by an energy barrier is assumed, not demonstrated. The “reset” framing risks encouraging dangerous self-experimentation with interventions (fasting, hyperthermia, psychedelics) that have no evidence of efficacy and carry substantial harm potential in metabolically fragile patients.

14.22.11 Drug Candidates for Systematic Investigation**? Open Question 54: Unexplored Pharmacological Targets**

Cimetidine's immunomodulatory effects were discovered accidentally. What other existing drugs might have unexplored relevance to ME/CFS?

Candidates based on mechanistic reasoning:

Mast Cell / Histamine Pathway:

- **Montelukast:** Leukotriene receptor antagonist; leukotrienes are mast cell mediators (some anecdotal benefit reported)
- **Cromolyn sodium:** Mast cell stabilizer; old drug, well-tolerated; why isn't it used more in ME/CFS?
- **Rupatadine:** H1 antihistamine + PAF antagonist; dual mechanism

Metabolic / Mitochondrial:

- **Metformin:** AMPK activator; mimics some effects of fasting; affects mitochondrial function
- **Low-dose lithium:** Neuroprotective; affects mitochondrial function and autophagy
- **Dichloroacetate (DCA):** Activates pyruvate dehydrogenase; forces glucose into TCA cycle

Vascular / Perfusion:

- **Pentoxifylline:** Improves blood rheology (flow properties); could address micro-clot/perfusion issues
- **Cilostazol:** Phosphodiesterase inhibitor; vasodilator; antiplatelet

Immune / Viral:

- **Famciclovir:** Different antiviral; some patients respond better than to valacyclovir
- **Artesunate:** Antimalarial with antiviral and immunomodulatory properties

Autonomic:

- **Droxidopa:** Norepinephrine prodrug; FDA-approved for orthostatic hypotension
- **Atomoxetine:** Norepinephrine reuptake inhibitor; off-label for POTS

These candidates are presented for research consideration, not as treatment recommendations. Systematic investigation of repurposed drugs could be more efficient than novel drug development.

14.22.12 The “Kitchen Sink” Protocol Concept**~ Hypothesis 18: Simultaneous Multi-Target Intervention**

If ME/CFS is maintained by multiple interacting feedback loops (the “multi-lock” model), addressing one mechanism at a time may fail because remaining mechanisms compensate. Effective treatment might require overwhelming the dysfunctional equilibrium by hitting multiple targets simultaneously.

Conceptual protocol targeting all major pathways:

1. **Mast cell stabilization:** H1 + H2 + Ketotifen + Quercetin
2. **Vagal restoration:** tVNS daily (60+ minutes)
3. **Gut barrier repair:** L-glutamine, zinc carnosine, butyrate
4. **Microbiome restoration:** Targeted probiotics
5. **Amino acid flooding:** High-dose supplementation (IV if needed to bypass absorption)
6. **Mitochondrial support:** Full Myhill-type protocol (CoQ10, D-ribose, magnesium, B vitamins)
7. **Circadian enforcement:** Strict light/dark, timed eating, sleep schedule
8. **Antiviral** (if indicated): Valacyclovir + cimetidine
9. **Immune modulation:** Low-Dose Naltrexone (LDN)

Rationale: Not “try one thing at a time” but hit everything at once, potentially overwhelming the pathological steady state and allowing transition to health.

Practical challenges:

- Complexity and cost
- Cannot identify which components are essential
- Risk of interactions
- Difficult to study in controlled trials

When might this be appropriate? For severely ill patients who have failed sequential single-intervention trials and face permanent disability, a coordinated multi-target approach may be worth the complexity.

△ Warning 17: Protocol Limitations

This “kitchen sink” approach has not been tested in any controlled manner. The complexity makes it difficult to implement and study. Not all patients can tolerate aggressive multi-intervention protocols. This concept is presented to stimulate thinking about treatment strategy, not as a validated protocol. Certainty: Very Low (for specific protocol); Medium (for multi-target concept).

14.23 Mechanistic Convergence: Cross-Treatment Integration

Recent integration of additional therapeutic agents—Devil’s Claw (harpagoside), ketamine, palmitoylethanolamide (PEA), statins, pregnenolone, and Ginkgo biloba—reveals previously unrecognized mechanistic overlaps suggesting rational combination strategies.

14.23.1 Convergence Clusters

Cluster 1: Triple Anti-Inflammatory Convergence (NF- κ B Node). Devil’s Claw, PEA, and statins all inhibit NF- κ B signaling through distinct upstream mechanisms: harpagoside directly blocks NF- κ B nuclear translocation; PPAR- α activation (PEA) suppresses NF- κ B via trans-repression; statins block isoprenylation of small GTPases required for NF- κ B activation. This mechanistic redundancy suggests potential for synergistic NF- κ B inhibition through distinct entry points.

Cluster 2: Neuroinflammation Convergence (Microglial Node). Ketamine and PEA both modulate microglial activation through orthogonal mechanisms: ketamine reduces microglial cytokine secretion via NMDA receptor blockade; PEA shifts microglial phenotype from M1 (pro-inflammatory) toward M2 via PPAR- α agonism. Combined use could produce more complete microglial “reset” than either alone.

Cluster 3: Mast Cell Convergence (MCAS Node). PEA stabilizes mast cells via PPAR- α and CB2 pathways (intracellular signaling), while Ginkgo blocks PAF, a potent extracellular mast cell activator. This addresses both release mechanisms and receptor activation.

Cluster 4: Ion Channel Convergence (TRPM3/Excitability Node). Ketamine (NMDA antagonism) and pregnenolone (TRPM3 modulation) both affect neuronal excitability—directly relevant to documented TRPM3 channelopathy in ME/CFS (Section 7.1.1).

14.23.2 Novel Combination Hypotheses

~ Hypothesis 19: Triple Anti-Inflammatory Stack: PEA + Devil’s Claw + LDN

Three mechanistically distinct anti-inflammatory agents targeting different cascade nodes may produce synergistic inflammation reduction: LDN at pattern recognition (TLR4), Devil’s Claw at transcription (NF- κ B), PEA at effector modulation (PPAR- α).

Predicted responders: Patients with documented inflammatory biomarker elevation (IL-6, TNF- α) with partial LDN response.

Testable prediction: Greater cytokine reduction than LDN monotherapy at 12 weeks.

Safety considerations: Combining three anti-inflammatory agents raises theoretical concerns about excessive immune suppression. Monitor for increased infection sus-

ceptibility. Note that Devil's Claw has anticoagulant potential—review bleeding risk if combining with other agents affecting hemostasis. Start components sequentially (not simultaneously) to identify any adverse reactions.

~ **Hypothesis 20: Neuroplasticity Combination: Pregnenolone + Ketamine**

Ketamine induces a “window of neuroplasticity” via BDNF release and mTOR activation. Pregnenolone during this window may guide reorganization toward healthier patterns. Additionally, if TRPM3 dysfunction contributes to ME/CFS, pregnenolone's TRPM3 modulation may address root causes while ketamine addresses downstream central sensitization.

Predicted responders: Central sensitization phenotype, “wired but tired” presentation, TRPM3-positive if testable.

Testable prediction: Combined treatment produces greater, more durable reduction in Central Sensitization Inventory scores.

~ **Hypothesis 21: Mitochondrial Paradox Resolution: Statin + CoQ10 + D-Ribose**

Statins offer immunomodulatory benefits for autoimmune ME/CFS subsets, but HMG-CoA reductase inhibition depletes CoQ10—potentially catastrophic in already-compromised mitochondria. **Resolution:** Aggressive mitochondrial protection (CoQ10 200–400 mg, D-ribose 5 g TID, PQQ 20 mg) beginning 4 weeks **before** statin initiation, with CPET monitoring to abort if energy metabolism worsens.

Target phenotype: GPCR autoantibody-positive patients refractory to or unable to access immunoadsorption.

Testable prediction: With protection, statins should not worsen CPET metrics while potentially reducing autoantibody titers over 3–6 months.

Speculation 37 (Electrolyte/MCAS Connection). Why do some MCAS-phenotype patients respond dramatically to aggressive electrolyte loading? Possible links: (1) chronic MCAS creates relative hypovolemia via histamine vasodilation; (2) mast cells are osmosensitive—adequate sodium may reduce activation triggers; (3) electrolyte solutions provide trace minerals for diamine oxidase (DAO) function.

Testable prediction: MCAS-phenotype patients should show greater ORS benefit than non-MCAS; mast cell markers should decrease with adequate electrolyte loading.

⚠ **Limitation 11: Combination Hypotheses: No Drug Interaction or Safety Data**

The novel combination hypotheses in this section (triple anti-inflammatory stack, neuroplasticity combination, statin paradox resolution) propose multi-drug regimens that have never been tested for safety or efficacy in ME/CFS or any related condition. Drug interaction profiles for these specific combinations are unknown. The rationale is based on single-mechanism reasoning that does not account for the complexity of multi-drug pharmacokinetics and pharmacodynamics. These combinations must not be

attempted outside formal clinical trial settings with appropriate safety monitoring.

14.23.3 Phenotype-Matched Selection

Rather than “one size fits all,” these novel agents show differential relevance to ME/CFS subgroups:

- **MCAS-predominant:** PEA + Ginkgo (mast cell stabilization via distinct mechanisms)
- **Central sensitization/chronic pain:** Ketamine + PEA + Devil’s Claw (NMDA, neuroinflammation, COX-2)
- **TRPM3-positive/channelopathy:** Pregnenolone (direct TRPM3 modulation)
- **Cerebral hypoperfusion:** Ginkgo (documented blood flow enhancement)
- **Inflammatory biomarker elevation:** Devil’s Claw + PEA + Statin (triple NF- κ B; requires CoQ10 protection)
- **Autoantibody-positive:** Statin with aggressive mitochondrial co-treatment
- **Cognitive-predominant:** Pregnenolone + Ginkgo (neurosteroid + perfusion)

★ Key Point: Novel Agent Integration Summary

The six newly integrated agents expand the ME/CFS therapeutic toolkit:

1. **Mechanistic convergence** at NF- κ B, microglial, and mast cell nodes suggests rational combination strategies
2. **TRPM3 modulation** (pregnenolone) represents an entirely new approach aligned with channelopathy research
3. **The statin paradox** can potentially be resolved with aggressive mitochondrial co-treatment
4. **Phenotype matching** is essential—clinical subtyping should guide agent selection

These hypotheses are presented for research prioritization, not as validated treatment recommendations. All require safety monitoring and ideally formal clinical evaluation.

14.24 Selective Energy Dysfunction Hypothesis

? Open Question 55: Why Does Hair Grow Normally in Severe ME/CFS?

A patient with severe ME/CFS cannot walk to the bathroom, cannot sustain a conversation, cannot tolerate light or sound—yet their hair continues to grow at a normal rate. Their nails grow. Wounds heal. These clinical observations derive from patient self-reports and informal clinical experience; no published study has formally measured hair or nail growth rates in ME/CFS patients. Nevertheless, the consistency of these reports across diverse patient populations poses a fundamental challenge to the “global

energy failure” model of ME/CFS: if mitochondrial dysfunction were truly systemic, *all* energy-dependent processes should be impaired proportionally.

This chapter proposes that ME/CFS represents *selective* rather than global energy dysfunction—specifically, a CNS coordination failure that impairs demand-responsive, CNS-dependent processes while sparing autonomous local processes that operate independently of central regulation.

⚡ **Limitation 12: Preserved Processes: Informal Clinical Observation, Not Measured**

The foundational claim that hair growth, nail growth, and wound healing are preserved in ME/CFS derives from patient self-reports and informal clinical experience. No published study has formally measured hair growth rate, nail growth rate, or wound healing time in ME/CFS patients compared to matched controls. These processes may in fact be subtly impaired without patients noticing, or they may appear preserved because their energy demands are negligible relative to the total energy budget even in health. The “selective vs. global” distinction may therefore be a continuum rather than a dichotomy, and the motivating observation could be an artefact of measurement insensitivity.

14.24.1 Motivation and Clinical Observations

The selective dysfunction hypothesis emerged from a simple observation: processes that require CNS coordination and scale with demand are severely impaired in ME/CFS, while processes that operate autonomously at the local tissue level remain intact.

★ **Key Point: Preserved vs. Impaired Processes**

Preserved (minimal CNS dependency):

- Hair growth—local follicle autonomous cycle
- Nail growth—keratinocyte autonomous proliferation
- Basic wound healing—local inflammatory cascade
- Baseline digestion—enteric nervous system (“second brain”)

Severely impaired (high CNS dependency):

- Exercise capacity—requires CNS motor coordination + autonomic scaling
- Cognitive function—intrinsically CNS-dependent
- Orthostatic regulation—requires real-time CNS autonomic control
- Temperature regulation—hypothalamic coordination
- Sleep architecture—brainstem and cortical orchestration

This pattern motivates the formal selective dysfunction hypothesis developed below (Hypothesis 14.24.2).

14.24.2 Formal Framework

We formalize the selective dysfunction hypothesis using three quantitative definitions that enable testable predictions.

Definition 14.1 (CNS-Dependency Index). For any biological process P , define the **CNS-dependency index** $\delta_{CNS}(P) \in [0, 1]$ as:

$$\delta_{CNS}(P) = \frac{N_{CNS}(P)}{N_{total}(P)}$$

where $N_{CNS}(P)$ is the number of regulatory signals requiring CNS coordination and $N_{total}(P)$ is the total number of regulatory signals controlling process P .

- $\delta_{CNS} = 0$: Fully autonomous (no CNS involvement)
- $\delta_{CNS} = 1$: Fully CNS-dependent (all regulation via CNS)
- $\delta_{CNS} \in (0, 1)$: Mixed regulation

Definition 14.2 (Demand-Responsiveness Index). For process P , define the **demand-responsiveness index** $\rho(P) \in [0, \infty)$ as:

$$\rho(P) = \frac{\max_{\text{challenge}} \text{Output}(P) - \text{Output}_{\text{baseline}}(P)}{\text{Output}_{\text{baseline}}(P)}$$

where:

- $\rho = 0$: Constant output (no demand scaling)
- $\rho > 0$: Output increases under challenge
- $\rho > 1$: Output can more than double under maximal demand

For standardization to $[0, 1]$, we use $\tilde{\rho}(P) = \frac{\rho(P)}{1+\rho(P)}$.

Note on negative demand-response: Some processes *decrease* output under challenge (e.g., digestive function during fight-or-flight response, where blood flow diverts to skeletal muscle). For such processes, we define $\rho(P) = 0$ because the selective dysfunction hypothesis specifically addresses *upregulation capacity*—the ability to increase output when demanded. Processes with negative demand-response represent regulatory suppression rather than capacity limitation, and require separate modeling frameworks (not within this hypothesis).

Definition 14.3 (Selective Dysfunction Severity). The **predicted dysfunction severity** for process P in ME/CFS is:

$$S(P) = \alpha \cdot \delta_{CNS}(P) + \beta \cdot \tilde{\rho}(P) + \gamma \cdot \delta_{CNS}(P) \cdot \tilde{\rho}(P)$$

where:

- $\alpha > 0$: Weight for CNS-dependency (main effect)
- $\beta > 0$: Weight for demand-responsiveness (main effect)

- $\gamma > 0$: Weight for interaction (synergistic vulnerability)

The interaction term $\gamma \cdot \delta_{CNS} \cdot \tilde{\rho}$ captures synergistic vulnerability: processes that are *both* CNS-dependent *and* demand-responsive are predicted to be most severely affected.

Note on range: $S(P) \in [0, \alpha + \beta + \gamma)$. Since $\tilde{\rho} = \rho/(1 + \rho) < 1$ strictly for any finite ρ , the supremum $\alpha + \beta + \gamma = 1.5$ is approached but never attained. Values exceeding 1.0 indicate severe predicted dysfunction (see Table 14.5).

~ Hypothesis 22: Selective Dysfunction Hypothesis

In ME/CFS, dysfunction severity $S_{observed}(P)$ correlates strongly with predicted severity $S(P)$:

$$S_{observed}(P) = S(P) + \epsilon$$

where ϵ is residual error. **Working assumption:** $\epsilon \sim \mathcal{N}(0, \sigma^2)$ (normally distributed with mean zero); this assumption requires validation against empirical data.

Testable prediction: Spearman correlation ρ_s between $S(P)$ and $S_{observed}(P)$ across processes should satisfy $\rho_s > 0.7$ (strong positive correlation).

Independence requirement: For this prediction to be independently falsifiable, $\delta_{CNS}(P)$ and $\tilde{\rho}(P)$ must be estimated from sources other than ME/CFS clinical observations—for example, from neuroanatomical pathway counts in healthy individuals and demand-scaling measurements in healthy controls. The current Table 14.5 values are expert estimates that partly reflect the clinical picture the model aims to predict; using them directly would make the correlation circular and non-falsifiable.

Falsification criterion: If $\rho_s < 0.3$ when tested with independently estimated parameters, the hypothesis is refuted.

Certainty: 0.55 (moderate evidence from clinical observations, formal testing required)

⚠ Limitation 13: Quantitative Framework: Unfitted Parameters and Circularity Risk

The CNS-dependency index δ_{CNS} , demand-responsiveness index $\tilde{\rho}$, and dysfunction severity $S(P)$ are defined mathematically but have never been fitted to empirical data. The parameter values in Table 14.5 are “expert estimates” that partly reflect the clinical picture the model aims to predict — as acknowledged in the independence requirement (line 104). Until δ_{CNS} and $\tilde{\rho}$ are estimated from independent sources (neuroanatomical pathway counts, healthy-control demand-scaling measurements), the correlation between predicted and observed dysfunction severity is circular. The weights α , β , γ have not been estimated from data and the interaction term’s contribution is unknown. The framework is therefore a formalised heuristic, not a validated quantitative model.

14.24.3 Evidence Taxonomy

△ Warning 18: Table Values Are Illustrative, Not Independent Validation

The δ_{CNS} and $\tilde{\rho}$ values in Table 14.5 are expert estimates that partly reflect the clinical picture the model aims to predict. **Using these values to test the correlation $\rho_s > 0.7$ would be circular and non-falsifiable.** Independent validation requires estimating parameters from sources outside ME/CFS clinical observations (e.g., neuroanatomical pathway counts in healthy individuals). The table illustrates the model's framework and internal consistency; it does *not* constitute evidence for or against the hypothesis. See Predictions 14.24.6–14.24.6 for independently testable predictions.

Table 14.5 classifies biological processes by their predicted and observed dysfunction status.

Table 14.5: Process classification by CNS-dependency and demand-responsiveness with predicted and observed dysfunction in ME/CFS.

Process	δ_{CNS}	$\tilde{\rho}$	Predicted S	Observed	Certainty
<i>Preserved processes (low δ_{CNS}, low ρ)</i>					
Hair growth [†]	0.1	0.0	0.06	Preserved	0.4
Nail growth [†]	0.1	0.0	0.06	Preserved	0.4
Wound healing (basic)	0.2	0.2	0.24	Preserved	0.6
Digestion (baseline) [‡]	0.3	0.2	0.30	Variable	0.5
<i>Impaired processes (high δ_{CNS} and/or high ρ)</i>					
Exercise capacity	0.8	0.8	1.14	Severe	0.9
Cognitive function	1.0	0.5	1.05	Severe	0.9
Orthostatic regulation	0.9	0.7	1.14	Severe	0.85
Temperature regulation	0.8	0.4	0.81	Moderate-Severe	0.7
Sleep architecture	0.9	0.3	0.80	Severe	0.8
Immune response to challenge	0.5	0.7	0.79	Dysregulated	0.65

Notes: S calculated with $\alpha = 0.6$, $\beta = 0.5$, $\gamma = 0.4$. Values > 1.0 indicate severe predicted dysfunction (additive model; theoretical supremum = $\alpha + \beta + \gamma = 1.5$, never attained since $\tilde{\rho} < 1$ strictly). Certainty reflects confidence in classification based on literature review.

[†]*Hair and nail growth:* Certainty 0.4 reflects that “preserved” status is based on patient self-reports and informal clinical experience; no formal study has measured growth rates in ME/CFS patients.

[‡]*Digestion (baseline):* The $\tilde{\rho} = 0.2$ value reflects demand-up scaling under normal eating conditions. The fight-or-flight suppression of digestion (a negative demand-response) is modeled separately as $\rho = 0$ per the convention in Definition 14.2.

Observation 71 (Demand-Response Failure Pattern). ME/CFS patients consistently show *preserved baseline function with impaired challenge response*:

- Orthostatic challenge: 91–100% show abnormal CBF reduction during tilt-table testing, with 3-fold greater reduction than controls [172]
- Exercise challenge: 2-day CPET reveals decline on day 2 (vs. improvement in controls) [1]
- Cognitive challenge: Fatigue elevated 24–48h post-cognitive task [430]

This pattern is diagnostic: baseline function is maintained, but the system cannot scale to meet increased demand. This is precisely what the selective dysfunction model predicts for high- ρ processes.

14.24.4 Causal Structure

Figure 14.1 presents the causal structure for selective dysfunction. The key feature is the absence of clinically significant causal paths from the CNS energy crisis node to preserved autonomous processes. Note: the figure includes a feedback edge from exercise intolerance to CNS energy crisis (representing PEM-induced CNS burden, certainty 0.35), which introduces a cycle; the structure is therefore a *causal diagram* rather than a strict directed acyclic graph (DAG).

The causal diagram encodes the following claims with certainty-weighted edges:

1. **Triggers** $\rightarrow$ **CNS Energy Crisis** (certainty: 0.65–0.70): Viral infection, immune dysregulation, or severe stress initiates CNS energy deficit [370]
2. **CNS Crisis** $\rightarrow$ **Cognitive Dysfunction** (certainty: 0.85): Direct effect of brain hypometabolism [73, 5]
3. **CNS Crisis** $\rightarrow$ **Autonomic Control Failure** (certainty: 0.80): Impaired CNS signaling disrupts autonomic regulation [172]
4. **Autonomic Failure** $\rightarrow$ **Orthostatic Intolerance** (certainty: 0.75): Secondary to failed cardiovascular coordination [172]
5. **Sub-threshold edge: CNS** $\rightarrow$ **Autonomous Processes**: Hair/nails ($\delta_{CNS} = 0.1$, $\tilde{\rho} = 0$: $S = 0.06$) and wound healing ($\delta_{CNS} = 0.2$, $\tilde{\rho} = 0.2$: $S = 0.24$) all fall in the preserved range of Table 14.5; all yield $S \leq 0.24$, well below the lowest impaired process ($S = 0.79$, immune response to challenge)

14.24.5 Mechanistic Sub-Hypotheses

Three hypotheses (certainty ≥ 0.45) and two speculations (certainty < 0.45) explain *why* CNS might be selectively vulnerable. These mechanisms are not mutually exclusive; multiple pathways may act additively in individual patients.

Astrocyte Energy Gate Hypothesis

The astrocyte-neuron lactate shuttle (ANLS) provides 30–50% of neuronal ATP [174, 175]. Unlike peripheral tissues with direct glucose access, neurons depend on astrocytes to convert glucose to lactate and shuttle it via monocarboxylate transporters (MCT2/MCT4).

Speculation 38 (Astrocyte Energy Gate). Dysfunction in the astrocyte-neuron lactate shuttle causes CNS-specific energy failure while peripheral tissues (with direct glucose access) remain unaffected.

Selective Energy Dysfunction: Causal Structure




Figure 14.1: Causal diagram of selective energy dysfunction in ME/CFS. Edge weights indicate certainty (0–1 scale) based on evidence quality. The diagram includes a feedback edge (exercise intolerance $\rightarrow$ CNS crisis, certainty 0.35), making it a causal diagram rather than a strict DAG. Preserved processes (green) have sub-threshold CNS-dependency ($\delta_{CNS} \leq 0.2$, $\tilde{\rho} \leq 0.2$, giving $S \leq 0.24$) and no clinically significant causal path from the CNS crisis node.

Mechanism:

$$L_n = k_{MCT} \cdot [L_a] \cdot f(\text{transporter integrity}) \tag{14.1}$$

$$E_n = g(L_n, \text{mitochondrial function}) \tag{14.2}$$

where L_n = neuronal lactate uptake, L_a = astrocyte lactate concentration, k_{MCT} = MCT transporter efficiency, E_n = neuronal ATP production.

ME/CFS hypothesis: Reduced k_{MCT} or impaired $f(\cdot) \Rightarrow L_n \downarrow$ despite normal $L_a \Rightarrow$ CNS-specific energy deficit.

Certainty: 0.4 (plausible mechanism from neuroscience; no direct ME/CFS evidence)

Figure 14.2 illustrates the ANLS mechanism and proposed dysfunction site.




Figure 14.2: Astrocyte-neuron lactate shuttle (ANLS) as proposed mechanism for CNS-specific energy dysfunction. If MCT2 transporter function is impaired, neurons cannot access lactate-derived ATP while peripheral tissues (with direct glucose access) remain unaffected.

CNS Energy Triage Hypothesis

Under energy scarcity, the CNS may implement a hardwired priority hierarchy that preserves vital functions at the expense of “luxury” cognitive processes.

~ Hypothesis 23: CNS Energy Triage

The CNS implements a priority-based energy allocation system under scarcity:

1. **Tier 1** (never sacrificed): Brainstem vital functions — $E_{min} = 0.30 \cdot E_{total}$
2. **Tier 2**: Sensory processing — $E_{threshold} = 0.50 \cdot E_{total}$
3. **Tier 3**: Motor coordination — $E_{threshold} = 0.60 \cdot E_{total}$
4. **Tier 4**: Memory consolidation — $E_{threshold} = 0.70 \cdot E_{total}$
5. **Tier 5**: Executive function — $E_{threshold} = 0.85 \cdot E_{total}$
6. **Tier 6** (first sacrificed): Complex cognition — $E_{threshold} = 0.95 \cdot E_{total}$

Note on threshold values: The E_{min} and $E_{threshold}$ fractions (0.30, 0.50, 0.60, 0.70, 0.85, 0.95) are *model parameters estimated for heuristic purposes*, not empirically derived values. They reflect qualitative evidence about neural energy priorities [175] but have not been measured directly in ME/CFS or in healthy humans.

Prediction: Cognitive symptoms should follow the inverse hierarchy. Complex cognition fails first; vital functions never fail.

Clinical correlation: “Brain fog” (executive dysfunction, Tier 5–6) is among the earliest and most prominent symptoms, consistent with these tiers being sacrificed first.

Certainty: 0.5 (consistent with observed symptom hierarchy; formal testing needed)

Figure 14.3 visualizes the triage hierarchy with the typical ME/CFS energy threshold.

Blood-Brain Barrier Vulnerability Hypothesis

The blood-brain barrier (BBB) creates a unique vulnerability: damage signals may accumulate in the CNS while peripheral clearance continues normally.

~ Hypothesis 24: BBB Compartmentalization

The BBB traps mitochondrial damage markers and limits cofactor delivery, causing CNS-specific accumulation of dysfunction.

Steady-state model:

$$[M]_{CSF} = [M]_{blood} + \frac{R_{production}}{P_{BBB} \cdot k_{clear}}$$

where $[M]$ = damage marker concentration (mass/volume), $R_{production}$ = CNS production rate (mass/volume/time), $P_{BBB} \cdot k_{clear}$ = effective BBB-mediated clearance rate (1/time), and $[M]_{blood}$ = peripheral (blood) concentration treated as a fixed boundary condition set by systemic clearance. At steady state, CNS production is balanced by BBB-mediated efflux of the excess above blood levels.




Figure 14.3: CNS energy triage hierarchy under energy scarcity. The red dashed line indicates typical available energy in ME/CFS. Functions above the line (Tiers 4–6) are impaired; functions below (Tiers 1–3) are preferentially preserved through energy allocation.

Ratio consequence: $\frac{[M]_{CSF}}{[M]_{blood}} = 1 + \frac{R_{production}}{P_{BBB} \cdot k_{clear} \cdot [M]_{blood}}$. In healthy subjects this ratio is near 1; in ME/CFS, reduced P_{BBB} or elevated $R_{production}$ drives it above 1, indicating CNS-specific accumulation.

ME/CFS prediction: If P_{BBB} is reduced or $R_{production}$ elevated, the CSF/blood ratio increases, indicating CNS-specific accumulation.

Testable: Measure mtDNA, 8-OHdG, or other damage markers in paired CSF and blood samples. Elevated CSF/blood ratio supports hypothesis.

Certainty: 0.45 (BBB dysfunction documented in neuroinflammation [73]; ME/CFS-specific data limited)

Sickness Behavior Persistence Hypothesis

Evolutionary sickness behavior programs target *behavioral outputs* (requiring CNS) while sparing truly autonomous processes.

~ Hypothesis 25: Sickness Behavior Stuck On

ME/CFS represents a sickness behavior program that fails to disengage, chronically suppressing CNS-mediated behavioral outputs while leaving autonomous local processes unaffected.

Activation function:

$$SB(t) = \text{sigm} \left(\sum_i w_i \cdot [\text{cytokine}_i](t) - \theta \right)$$

where $\text{sigm}(x) = 1/(1 + e^{-x})$ is the logistic sigmoid function (distinct from standard deviation σ used elsewhere in this chapter), w_i are cytokine weights, and θ is the activation threshold.

Normal state: Acute infection elevates cytokines $\Rightarrow$ SB activates $\Rightarrow$ infection resolves $\Rightarrow$ cytokines normalize $\Rightarrow$ SB deactivates.

ME/CFS state: Chronic low-grade immune activation maintains $\sum w_i \cdot [\text{cytokine}_i] > \theta$ indefinitely $\Rightarrow$ SB persists.

Evolutionary logic: Sickness behavior evolved to suppress *behavioral* energy expenditure during infection. Hair growth has no behavioral component and was never targeted by this program.

Certainty: 0.55 (strong evolutionary logic; moderate mechanistic support from neuroimaging [73])

Partial Torpor Trap Hypothesis

ME/CFS may represent incomplete engagement of torpor-like metabolic suppression mechanisms.

Speculation 39 (Partial Torpor Trap). ME/CFS involves partial engagement of torpor/hibernation pathways with failed arousal, trapping patients in a low-metabolic state.

Torpor engagement dynamics:

$$\frac{d(\text{MR})}{dt} = -k_T \cdot T_{\text{signal}} + k_A \cdot A_{\text{signal}} - \lambda(\text{MR} - \text{MR}_0)$$

where MR = metabolic rate, T_{signal} = torpor induction signal, A_{signal} = arousal signal, $k_T, k_A > 0$ are signal gain coefficients (distinct from the α, β weights in the $S(P)$ formula), $\lambda > 0$ is a restoring rate constant, and MR_0 is the baseline metabolic rate. The restoring term $-\lambda(\text{MR} - \text{MR}_0)$ ensures a bounded steady state at $\text{MR}^* = \text{MR}_0 + (k_A A - k_T T)/\lambda$.

Normal torpor: $k_T T > k_A A$ during entry (MR suppressed below MR_0); $k_A A > k_T T$ during arousal (MR restored toward MR_0).

ME/CFS state: $k_T T > k_A A$ chronically, so $\text{MR}^* < \text{MR}_0$ (trapped in partial suppression).

Testable markers: Torpor-associated molecules (H_2S , adenosine, orexin) may be dysregulated.

Certainty: 0.35 (speculative; inspired by emerging torpor biology research [385, 386])

14.24.6 Testable Predictions

The selective dysfunction hypothesis generates specific, falsifiable predictions.

→ Prediction 6: Hair Follicle Mitochondrial Function

Hypothesis: Hair follicle mitochondria are functionally normal in ME/CFS patients.

Measurement: Mitochondrial respiration (oxygen consumption rate, OCR) in plucked hair follicle cells.

Statistical design:

- Test type: Equivalence test (TOST procedure)
- Equivalence margin: $\delta = 0.2 \times \bar{x}_{\text{control}}$ (20% of control mean)
- Sample size: $n = 40$ per group (power = 0.8, $\alpha = 0.05$)
- Outcome: ME/CFS OCR equivalent to control OCR

Interpretation:

- If confirmed: Strong support for selective (not global) mitochondrial dysfunction
- If refuted (ME/CFS OCR significantly lower): Global dysfunction model supported

Feasibility: Hair follicle collection is minimally invasive; mitochondrial respiration assays are established.

→ Prediction 7: CSF-to-Blood Lactate Gradient

Hypothesis: CSF lactate is elevated relative to blood lactate in ME/CFS, indicating impaired lactate shuttling in CNS.

Measurement: Paired CSF and blood lactate concentrations.

Statistical design:

- Test type: Two-sample t -test on ratio $[L]_{CSF}/[L]_{blood}$
- Expected effect size: Cohen's $d \geq 0.5$ (medium effect)
- Sample size: $n = 64$ per group (power = 0.8, $\alpha = 0.05$)
- Outcome: Ratio elevated in ME/CFS vs. controls

Interpretation:

- If confirmed: Supports astrocyte energy gate hypothesis
- If refuted: Lactate shuttle not primary mechanism

→ **Prediction 8: Peripheral ATP During PEM**

Hypothesis: Peripheral muscle ATP is preserved during PEM crashes (dysfunction is coordination failure, not local energy deficit).

Measurement: ^{31}P -MRS of skeletal muscle during provoked PEM.

Statistical design:

- Test type: Repeated measures ANOVA (baseline vs. PEM)
- Expected: No significant decline in muscle ATP during PEM
- Comparison: CNS metabolic markers (via PET/MRS) should decline while peripheral markers remain stable

Interpretation:

- If confirmed: Dysfunction is CNS coordination failure, not peripheral energy deficit
- If refuted (peripheral ATP drops): Global depletion model supported

→ **Prediction 9: Direct Stimulation vs. Voluntary Contraction**

Hypothesis: Direct electrical stimulation of muscles produces greater force than voluntary contraction in ME/CFS patients.

Rationale: If dysfunction is CNS coordination failure, bypassing CNS via direct stimulation should restore output.

Measurement: Compare force production: voluntary maximal contraction vs. electrical stimulation.

Expected: $F_{electrical}/F_{voluntary} > 1$ in ME/CFS (vs. ratio ≈ 1 in controls).

Interpretation:

- If confirmed: Peripheral muscle capable; CNS drive impaired
- If refuted: Peripheral muscle intrinsically impaired

→ **Prediction 10: Cognitive Triage Hierarchy**

Hypothesis: Cognitive impairment in ME/CFS follows the inverse of the energy triage hierarchy.

Measurement: Cognitive battery assessing each tier:

- Tier 6 (complex cognition): Abstract reasoning, creativity
- Tier 5 (executive function): Planning, multitasking, attention
- Tier 4 (memory): Working memory, recall
- Tier 3 (motor coordination): Fine motor, reaction time
- Tier 2 (sensory processing): Basic perception

Expected: Impairment severity: Tier 6 > Tier 5 > Tier 4 > Tier 3 > Tier 2.

Statistical test: Ordinal regression testing hierarchy effect.

14.24.7 Subtype Classification Model

The selective dysfunction framework suggests natural subtypes based on primary compartment affected.

Speculation 40 (Selective Dysfunction Subtypes). ME/CFS can be classified into subtypes based on which compartment shows primary dysfunction:

Input features (with measurement basis):

- x_1 : CSF catecholamine deficit (z-score relative to healthy controls)
 - Measurement: CSF dopamine, norepinephrine, serotonin metabolites
 - Reference: NIH deep phenotyping study found significant CSF catecholamine reductions [5]
- x_2 : Orthostatic CBF reduction (% decline from supine to upright)
 - Measurement: Transcranial Doppler during tilt-table test
 - Reference: Controls show ~5–10% reduction; ME/CFS shows ~20–30% (3-fold greater) [172]
- x_3 : Muscle ATP deficit at rest (% below control mean)
 - Measurement: ^{31}P -MRS of quadriceps or forearm
 - Reference: Variable findings in literature; some studies show 20–40% reduction [431, 102]
- x_4 : Neuroimaging abnormality score (composite z-score)
 - Measurement: PET neuroinflammation markers, fMRI activation patterns, MRS metabolites
 - Reference: Nakatomi et al. found 45–199% elevation in neuroinflammation markers [73]

Proposed classification rules (preliminary thresholds):

The following thresholds are *preliminary estimates* based on effect sizes in the literature. They require empirical validation through clustering analysis on a multi-biomarker cohort before clinical application.

Note on units: Thresholds use z-scores (σ) for standardized measures and absolute percentages (%) for CBF changes, reflecting conventions in respective literatures.

Subtype A (CNS-Primary) : $x_1 < -1.5\sigma \wedge x_4 > 2\sigma \wedge x_3 > -0.5\sigma$
(CNS markers abnormal, peripheral spared)

Subtype B (Autonomic-Primary) : $x_2 > 25\% \wedge x_1 > -1.0\sigma$
(OI dominant, CNS markers near-normal)

Subtype C (Peripheral-Primary) : $x_3 < -1.5\sigma \wedge x_1 > -1.0\sigma \wedge x_4 < 1\sigma$
(Muscle deficit primary, CNS spared)

Subtype D (Global/Advanced) : ≥ 3 of $\{x_1 < -1.5\sigma, x_2 > 25\%, x_3 < -1.5\sigma, x_4 > 2\sigma\}$
(Multi-system involvement)

Threshold rationale:

- -1.5σ : Corresponds to approximately the 7th percentile of control distribution—outside normal variation
- 2σ : Corresponds to approximately the 98th percentile—clearly elevated
- 25% CBF reduction: Approximately $2.5 \times$ the normal orthostatic response (note: this is an absolute percentage, not a z-score, reflecting how CBF changes are typically reported in the literature)

Subtype E (Unclassified): Patients meeting fewer than 3 criteria for Subtype D and not meeting full criteria for Subtypes A, B, or C are classified as **Subtype E (Unclassified/Sub-threshold)**. This category captures early-stage disease, atypical presentations, or patients whose biomarkers fall in the moderate range below all classification thresholds. Subtype E is expected to be common under this preliminary scheme and should be a priority for threshold revision during validation.

Overlap precedence: When a patient meets criteria for multiple subtypes:

1. If ≥ 3 criteria met $\Rightarrow$ classify as Subtype D (Global) regardless of other matches
2. Otherwise, convert all features to z-scores for comparison (convert the CBF percentage x_2 to a z-score using the control distribution mean and SD before comparing magnitude); assign to the subtype with the largest absolute z-score deviation; if tied, prioritize CNS-Primary > Autonomic-Primary > Peripheral-Primary (reflecting the hypothesis that CNS dysfunction is upstream)
3. Document secondary subtype features for treatment consideration

Validation protocol required:

1. Collect multi-biomarker panel in $n \geq 200$ ME/CFS patients (sample size rationale: with 4 subtypes and 4 input features, minimum 50 patients per subtype needed for stable cluster estimation; $n = 200$ provides robustness against unequal subtype prevalence and

allows 10% holdout for validation, with k -fold cross-validation to compensate for small holdout size)

2. Perform unsupervised clustering (k-means, hierarchical) to identify natural groupings
3. Compare data-driven clusters to proposed subtypes
4. Calculate sensitivity/specificity for each classification rule
5. Refine thresholds based on ROC analysis to optimize classification accuracy

Treatment implications (contingent on validation):

- Subtype A: CNS-penetrant compounds, intranasal delivery, neuroinflammation-targeted therapy
- Subtype B: Autonomic modulators (midodrine, pyridostigmine), volume expansion
- Subtype C: Mitochondrial support, muscle-targeted interventions. **Warning:** Graded exercise is contraindicated in ME/CFS due to PEM risk even when CNS markers appear spared; any activity increase requires careful symptom monitoring and pacing principles [127]
- Subtype D: Multi-system approach, sequential targeting of dominant compartments

Certainty: 0.35 (framework theoretically motivated; all thresholds are preliminary estimates requiring empirical validation before any clinical application)

⚠ Limitation 14: Subtype Classification: No Empirical Validation

The four-subtype classification (CNS-Primary, Autonomic-Primary, Peripheral-Primary, Global/Advanced) has not been tested against patient data. The classification thresholds (-1.5σ , 2σ , 25% CBF reduction) are preliminary estimates, not data-derived cut-points. No multi-biomarker cohort study has simultaneously measured CSF catecholamines, orthostatic CBF, muscle ATP, and neuroimaging abnormalities in ME/CFS patients to determine whether natural clusters correspond to these proposed subtypes. The Subtype E (Unclassified) category may capture the majority of patients, rendering the classification clinically impractical. Treatment stratification based on these subtypes is entirely hypothetical.

14.24.8 Treatment Implications

The selective dysfunction hypothesis has immediate treatment implications.

Pharmacological Bypass Evidence

Observation 72 (Midodrine Effectiveness). Midodrine (direct α_1 -adrenergic agonist) improves orthostatic symptoms in patients with POTS and orthostatic intolerance—conditions highly comorbid with ME/CFS—by directly stimulating peripheral vasculature, bypassing CNS autonomic coordination [338].

Implication: If peripheral targets were intrinsically dysfunctional, pharmacological bypass would not restore function. Midodrine effectiveness provides evidence consistent with intact peripheral machinery and impaired CNS coordination—though this inference is from a POTS-comorbid population and does not rule out a degree of peripheral dysfunction in ME/CFS more broadly.

This is consistent with the selective dysfunction model: the dominant problem may be signaling/coordination rather than end-organ failure.

Therapeutic Strategy

★ Key Point: Treatment Strategy from Selective Dysfunction Model

1. **CNS-targeted delivery:** Intranasal or intrathecal routes may outperform oral for CNS-targeted compounds
2. **Pharmacological bypass:** Direct-acting agents that bypass CNS coordination (midodrine, direct muscle stimulation)
3. **Reduce CNS energy demands:** Pacing as CNS energy management, not just “activity reduction”
4. **Subtype-specific targeting:** Match intervention to primary dysfunction compartment (see subtype classification, Speculation 40)

For mechanism-specific treatment implications, see Section 14.24.10, where each of the ten related hypotheses includes dedicated treatment and limitations paragraphs.

14.24.9 Compartmental Energy Model

Figure 14.4 presents the four-compartment energy model with CNS as the coordination bottleneck.

★ Key Point: Model Interpretation

The CNS compartment serves dual roles:

1. **Primary dysfunction site:** Brain hypometabolism, catecholamine deficiency
2. **Coordination bottleneck:** Secondary failures in autonomic and peripheral compartments result from impaired CNS signaling, not local energy deficits

Autonomous processes bypass the CNS entirely and remain functional. This explains why:

- Hair grows normally (no CNS coordination required)
- Midodrine works (bypasses CNS, directly stimulates periphery)
- PEM affects demand-responsive but not baseline functions




Figure 14.4: Four-compartment energy model showing CNS as the coordination bottleneck in ME/CFS. Compartments are classified by CNS-dependency index (δ_{CNS}). Secondary dysfunction in autonomic and peripheral compartments results from impaired CNS coordination, while autonomous processes with $\delta_{CNS} \leq 0.2$ remain preserved.

14.24.10 Related Hypotheses Extending the Framework

The selective dysfunction framework generates 10 related hypotheses and speculations (7 hypotheses with certainty ≥ 0.45 ; 3 speculations with certainty < 0.45) that extend, complement, or refine specific aspects of the core model. Each targets a specific mechanism, comorbidity, or clinical pattern.

Sleep Architecture Failure Hypothesis

~ Hypothesis 26: Sleep Architecture CNS Coordination Failure

ME/CFS sleep disturbance results from impaired CNS coordination of sleep stage transitions rather than local sleep circuitry dysfunction.

Relationship to parent hypothesis: Extends CNS coordination failure to explain non-restorative sleep.

Mechanism: Normal sleep requires orchestrated inter-regional coordination across brainstem, thalamus, and cortex for stage transitions (REM $\leftrightarrow$ NREM, N1 $\rightarrow$ N2 $\rightarrow$ N3), with thalamic slow oscillations leading cortical downstates before spindle generation [259]. CNS energy deficit may preserve baseline sleep drive while impairing the sustained metabolic demands of coordinated transitions, producing the characteristic paradox of unrefreshing sleep despite near-normal total duration—documented in objective polysomnographic meta-analysis of 801 ME/CFS patients [258].

Testable prediction: Sleep spindle density (requiring thalamocortical coordination) correlates negatively with CNS dysfunction markers (CSF catecholamine deficit, brain hypometabolism).

Statistical test: Pearson correlation between spindle density and CSF neurotransmitter levels; expected $r < -0.5$ in ME/CFS cohort.

Novelty: Partial—sleep disturbance is established; CNS coordination framing is new.

Certainty: 0.50 (moderate)

Treatment implications. If sleep architecture failure reflects energy-limited coordination, interventions should reduce coordination demands and support metabolic capacity: (1) sleep continuity over total duration—fragmented sleep amplifies the metabolic cost of stage transitions [262]; (2) metabolic support during evening hours such as CoQ10 and NADH supplementation to support overnight cerebral energy availability [263]; (3) pacing-sleep integration, recognizing that sleep architecture worsens predictably during PEM so activity should be limited in the 12–24 hours before intended sleep; (4) avoidance of late-day cognitive exertion and stimulants that elevate baseline CNS arousal and increase stage-transition cost.

Limitations. This hypothesis has certainty 0.50. No published studies have directly quantified sleep spindle density or slow-wave coherence in ME/CFS alongside simultaneous CNS metabolic measures. Existing polysomnographic literature characterizes macrostructural stage percentages rather than the microstructural coordination markers

predicted here [258]. Alternative explanations include primary brainstem pathology, pain-mediated fragmentation, and autonomic disruption during sleep independent of energy limitation. Causality direction is unclear: energy limitation may fragment sleep, while fragmented sleep independently worsens brain energy metabolism [262], creating a potential vicious cycle.

Gut-Brain Energy Theft Hypothesis

Speculation 41 (Microbiome-Induced CNS Energy Depletion). Dysbiotic gut microbiome increases CNS metabolic burden through chronic immune activation, “stealing” energy from cognitive function.

Relationship to parent hypothesis: Adds upstream mechanism for CNS energy budget depletion.

Mechanism: Gut dysbiosis is consistently documented in ME/CFS [227]. The proposed cascade: dysbiosis → increased gut permeability → systemic endotoxemia → sustained low-grade neuroinflammation [73] → elevated CNS baseline energy cost → reduced budget available for demand-response. Bidirectional gut-brain signaling via vagal afferents [228] further provides a rapid channel through which gut dysbiosis could amplify CNS metabolic demand.

Testable prediction: Microbiome diversity (Shannon index) correlates negatively with brain hypometabolism severity (PET rCMRglc reduction).

Statistical test: Multiple regression controlling for disease duration and severity; expected partial $r < -0.4$.

Novelty: Yes—mechanistic link between established dysbiosis and CNS energy deficit.

Certainty: 0.40 (plausible mechanism; correlational evidence only)

Limitations. This speculation has certainty 0.40. Dysbiosis in ME/CFS is established [227] but is likely partly secondary to reduced physical activity, altered diet, and medication use rather than a primary driver. The multi-step cascade from gut permeability to CNS hypometabolism involves each link being individually plausible but none directly demonstrated in ME/CFS in sequence. PET neuroinflammation data exists [73] but has not been correlated with microbiome composition in the same patients. Probiotic and dietary interventions correcting dysbiosis show inconsistent cognitive benefits in ME/CFS, providing weak causal support for the energy-theft mechanism.

GPCR Autoantibody Inefficiency Hypothesis**~ Hypothesis 27: Autoantibody-Induced Autonomic Inefficiency**

G-protein coupled receptor (GPCR) autoantibodies impair autonomic signal transduction efficiency, requiring greater CNS effort to achieve the same peripheral output.

Relationship to parent hypothesis: Mechanistic explanation for autonomic demand-response failure.

Mechanism: Autoantibodies to β -adrenergic and muscarinic cholinergic receptors, identified in ME/CFS patients [71], may act as partial agonists or modulators that reduce signal amplification in autonomic efferent pathways. The CNS must then increase output intensity to compensate, raising energy cost per unit of autonomic control. Depletion of autoantibody-producing plasma cells with daratumumab has shown clinical improvement in a case series [67], providing preliminary evidence consistent with a causal role.

Testable prediction: Presence of GPCR autoantibodies correlates with increased CNS metabolic cost during autonomic challenge (PET during tilt-table).

Statistical test: Two-sample *t*-test comparing CNS glucose uptake change (supine $\rightarrow$ upright) in AAb+ vs. AAb- patients; expected Cohen's *d* > 0.5.

Novelty: Moderate—AABs documented in ~30% of patients [71]; efficiency framing is new.

Certainty: 0.45 (AABs present but causal role unproven)

Treatment implications. If GPCR autoantibodies impair autonomic signal efficiency, antibody-reducing strategies merit investigation: (1) plasmapheresis for rapid reduction of circulating autoantibody load; (2) B-cell depletion as a mechanistic target, though the rituximab phase III trial failed in an unselected cohort [117], suggesting AAb-positive subgroup stratification may be required for benefit; (3) plasma cell depletion with daratumumab, which targets longer-lived antibody-secreting cells and has shown case-series evidence [67]; (4) beta-adrenergic receptor modulation to partially compensate for impaired signaling efficiency.

Limitations. This hypothesis has certainty 0.45. GPCR autoantibodies are found in approximately 30% of ME/CFS patients [71], leaving the majority without this mechanism. Causal status remains disputed: antibodies may be epiphenomenal rather than pathogenic. The rituximab randomized controlled trial, which depletes B cells rather than plasma cells, failed to demonstrate benefit in an unselected cohort [117], providing indirect evidence against autoantibodies as a universal mechanism. The proposed 'increased CNS effort' framing has not been tested directly with neuroimaging during autonomic challenge stratified by autoantibody status.

Small Fiber Neuropathy Interface Failure

Speculation 42 (SFN Increases CNS Coordination Load). Small fiber neuropathy (SFN) found in ~30% of ME/CFS patients [331] increases CNS metabolic load by requiring compensatory signaling to maintain autonomic control.

Relationship to parent hypothesis: SFN as amplifier of peripheral-CNS communication cost.

Mechanism: SFN degrades signal fidelity at the peripheral nerve level, with objective evidence from skin punch biopsy showing reduced intraepidermal nerve fiber density (IENFD) [281]. The CNS must then send stronger, more frequent, or redundant signals to achieve target autonomic output, raising energy expenditure for regulation. This compensatory signaling pattern is documented in other conditions with peripheral neuropathy [284], where central gain increases to offset peripheral signal loss.

Testable prediction: SFN severity (IENFD, intraepidermal nerve fiber density) correlates with CNS metabolic cost during autonomic challenge.

Statistical test: Correlation between IENFD and brainstem/hypothalamic glucose uptake during tilt; expected $r < -0.5$ (lower IENFD = higher CNS cost).

Novelty: Yes—SFN documented but relationship to CNS burden unexplored.

Certainty: 0.40 (SFN prevalence established; mechanism speculative)

Limitations. This speculation has certainty 0.40. SFN prevalence in ME/CFS is established at approximately 30% [331], but the proposed mechanism—that impaired peripheral signal fidelity increases CNS metabolic cost—has not been directly tested. No study has correlated IENFD with CNS glucose uptake or brainstem metabolic demand in ME/CFS. SFN may be a secondary consequence of autonomic dysregulation or reduced activity rather than a primary amplifier of CNS burden. The 70% of patients without SFN must have other mechanisms for exercise intolerance, limiting this hypothesis to a subgroup explanation.

Circadian Energy Distribution Failure

~ Hypothesis 28: Circadian Misallocation of Energy Budget

Suprachiasmatic nucleus (SCN) dysfunction impairs circadian allocation of the CNS energy budget, explaining “second wind” phenomena and worsening symptoms at predicted low-energy times.

Relationship to parent hypothesis: Temporal dimension of energy allocation failure.

Mechanism: The SCN normally orchestrates circadian allocation of metabolic resources across the 24-hour cycle, including modulation of hepatic glucose production, cortisol rhythm, and CNS arousal [396]. Circadian rhythm abnormalities are documented in ME/CFS, including dissociation between body temperature and melatonin secretion rhythms [400] and disrupted autonomic circadian patterns [310]. The proposed consequence: SCN dysfunction causes temporal mismatch in energy availability—energy peaks when demand is low (late-night “second wind”) and is depleted when demand is highest (morning/afternoon crashes) [399].

Testable prediction: Circadian misalignment (dim-light melatonin onset phase shift) correlates with symptom severity fluctuation amplitude across the day.

Statistical test: Correlation between DLMO phase delay and variance in hourly symptom scores; expected $r > 0.5$.

Novelty: Partial—circadian disruption documented; energy allocation framing is new.

Certainty: 0.50 (circadian abnormalities established; causal link to energy requires testing)

Treatment implications. If SCN dysfunction causes temporal energy misallocation, chronotherapeutic interventions may partially restore circadian alignment: (1) melatonin timed to individual DLMO phase to re-anchor the circadian energy distribution [323]; (2) bright light therapy in the morning to strengthen SCN entrainment signals; (3) behavioral chronotherapy—scheduling cognitively or physically demanding activities at the patient’s individually observed peak energy window rather than social norms; (4) stabilizing the sleep-wake cycle, since irregular timing perpetuates SCN desynchrony and amplifies the mismatch.

Limitations. This hypothesis has certainty 0.50. Circadian abnormalities in ME/CFS are established [310, 400], but whether SCN dysfunction is primary or secondary to activity restriction, altered light exposure, or sleep fragmentation is unknown. The “second wind” pattern, while common in patient reports, has not been characterized with simultaneous objective metabolic measures. DLMO phase studies correlating with intraday symptom variance are lacking in ME/CFS. Prior chronotherapy trials (melatonin and light therapy) have not demonstrated robust symptomatic benefit [401], though these were not designed around the energy-allocation framing tested here.

MCAS Energy Crisis Amplifier

~ Hypothesis 29: Mast Cell Activation Amplifies CNS Energy Deficit

Mast cell activation syndrome (MCAS) episodes trigger acute inflammatory cascades that amplify CNS energy deficit, worsening PEM and cognitive crashes.

Relationship to parent hypothesis: Explains episodic worsening and high MCAS comorbidity.

Mechanism: MCAS degranulation releases histamine, cytokines, and prostaglandins [229] that penetrate the blood-brain barrier and activate neuroinflammatory cascades via mast cell–microglia signaling [245, 432]. The proposed cascade: mast cell degranulation → neuroactive mediator release → neuroinflammation spike → acute CNS energy demand increase → exceeds the already-limited budget → crash.

Testable prediction: Tryptase elevation (MCAS marker) during crashes correlates with crash severity and CNS metabolic decline.

Statistical test: Paired measurements (baseline vs. crash) showing correlated tryptase and PET hypometabolism changes; expected $r > 0.6$.

Novelty: Yes—MCAS–ME/CFS comorbidity recognized [229] but mechanistic link to CNS energy unexplored.

Certainty: 0.45 (high comorbidity documented; mechanistic testing needed)

Treatment implications. If MCAS episodes amplify CNS energy deficits, reducing mast cell activation load should attenuate crash severity: (1) mast cell stabilizers (cromoglicate, ketotifen) as daily prophylaxis to reduce baseline degranulation [229]; (2) combined H1 and H2 antihistamines to limit downstream neuroinflammatory signaling; (3) trigger identification and avoidance (heat, stress, certain foods, fragrances) to prevent crash amplification; (4) low-histamine dietary strategies during crash recovery to reduce ongoing mast cell stimulation; (5) serum tryptase and urinary prostaglandin D₂ measurement during crashes to confirm MCAS contribution in individual patients [229].

Limitations. This hypothesis has certainty 0.45. The MCAS–ME/CFS comorbidity rate is estimated at 20–60% depending on diagnostic criteria [229], but serum tryptase is frequently within the normal range even during confirmed MCAS episodes, making biochemical verification unreliable. The proposed mechanism—that mast cell mediators increase CNS energy demand enough to precipitate crashes—has not been tested with simultaneous tryptase measurement and neuroimaging. MCAS may also produce fatigue and cognitive symptoms through histamine receptor activation and direct neurological effects independent of the selective energy dysfunction framework, making mechanistic attribution difficult.

Memory Triage Consequence Hypothesis

~ Hypothesis 30: Hierarchical Memory Impairment from Energy Triage

The energy triage hierarchy predicts differential memory impairment: encoding (high-energy) fails before retrieval (lower-energy).

Relationship to parent hypothesis: Specific prediction from the CNS energy triage framework.

Mechanism: Hippocampal memory encoding requires long-term potentiation (LTP), involving NMDA receptor activation, calcium-dependent signaling cascades, de novo protein synthesis, and structural synaptic remodeling [300, 301]. Working memory maintenance additionally requires sustained prefrontal neuronal firing against inhibitory currents, proportional to item load [302]. Retrieval, by contrast, primarily reactivates existing synaptic patterns through pattern completion in CA3 networks—a lower-energy process requiring no new protein synthesis. Under CNS energy scarcity, encoding is sacrificed before retrieval.

Testable prediction: ME/CFS patients show greater impairment in encoding than retrieval on standardized memory tests (e.g., CVLT: learning slope vs. delayed recall).

Statistical test: Within-subjects comparison of z-scored encoding vs. retrieval; expected encoding impairment > retrieval impairment (paired *t*-test, *p* < 0.01).

Novelty: Partial—memory impairment established; encoding/retrieval asymmetry prediction is new.

Certainty: 0.55 (strong theoretical basis; testable with existing neuropsychological tools)

Treatment implications. If encoding is selectively impaired by CNS energy limitation, compensatory strategies should reduce encoding demands: (1) external memory aids (notes, recordings, photographs) to bypass the need for internal encoding; (2) scheduling new learning during individually identified peak energy windows; (3) spaced repetition to distribute encoding costs across multiple low-demand sessions, reducing single-session metabolic requirements; (4) recognition over recall formats (multiple-choice, visual cues) to leverage the relatively preserved pattern completion mechanism.

Limitations. This hypothesis has certainty 0.55. The differential energy cost of encoding versus retrieval is well established in basic neuroscience [300, 301], but the specific prediction of disproportionate encoding impairment in ME/CFS awaits formal testing. Standard neuropsychological batteries typically conflate encoding and retrieval in composite memory scores [303]; paradigms designed to isolate the two components are needed. Attention deficits, which impair encoding indirectly by reducing the depth of initial processing, and sleep fragmentation, which impairs memory consolidation independently of encoding energy, represent significant confounds.

Motor-Autonomic Coordination Overload

~ Hypothesis 31: Parallel Coordination Failure Under Exercise

Exercise requires simultaneous CNS coordination of motor output and autonomic scaling (HR, BP, respiration). Under CNS energy deficit, parallel coordination fails, explaining exercise intolerance despite preserved individual systems at rest.

Relationship to parent hypothesis: Explains why demand-response fails specifically during combined motor-autonomic challenges.

Mechanism: Motor coordination and autonomic scaling both demand CNS energy. Two-day cardiopulmonary exercise testing documents that ME/CFS patients show reproducibly greater metabolic impairment on day 2 than day 1, consistent with exhaustion of CNS coordination capacity under repeated demand [1]. Impaired cerebral blood flow regulation during orthostatic challenge—found in 91–100% of patients [172]—further supports failure of CNS-mediated cardiovascular coordination under demand. At rest or with single tasks, the residual energy budget suffices. During exercise, when motor and autonomic coordination occur simultaneously, total demand exceeds the budget → coordination failure → PEM [5].

Testable prediction: ME/CFS patients show greater impairment on combined motor+autonomic tasks (exercise) than either task in isolation (cognitive challenge or paced breathing).

Statistical test: Interaction effect in 2×2 design (motor: yes/no × autonomic: yes/no); expected significant interaction, $\eta^2 > 0.10$.

Novelty: Moderate—exercise intolerance is core symptom; parallel coordination framing is new.

Certainty: 0.55 (strong clinical correlation; formal interaction testing needed)

Treatment implications. If parallel coordination failure explains exercise intolerance, management strategies should limit combined motor-autonomic demand: (1) heart rate-guided activity pacing to keep cardiovascular demand below the threshold that triggers coordination failure; (2) avoiding cognitive exertion immediately before or during physical activity, which adds concurrent demand to an already-loaded system; (3) two-day CPET as an objective marker of the coordination failure threshold, informing individualized activity ceilings [1]; (4) symptom-limited rather than fixed-increment graded activity protocols, since fixed increments ignore the patient’s coordination threshold.

Limitations. This hypothesis has certainty 0.55. The prediction of a significant interaction effect between motor and autonomic demand in a formal 2×2 design has not been tested. Peripheral skeletal muscle pathology—including impaired oxygen extraction documented in ME/CFS deep phenotyping [5]—provides an independent explanation for exercise intolerance that does not require CNS coordination failure as the primary bottleneck. Post-exertional malaise may confound assessment: patients who exercise to test the hypothesis suffer delayed symptom worsening that makes attribution of the primary failure point difficult.

Post-Viral CNS Metabolic Reprogramming

Speculation 43 (Persistent Astrocyte Metabolic Shift Post-Infection). Viral infection triggers persistent astrocyte metabolic reprogramming toward a “reactive” state with reduced lactate shuttle efficiency, causing chronic CNS energy deficit.

Relationship to parent hypothesis: Viral trigger mechanism for astrocyte energy gate dysfunction.

Mechanism: Astrocytes supply neurons with lactate through the astrocyte-neuron lactate shuttle (ANLS), providing 30–50% of neuronal ATP under activity [174, 175]. Acute viral infection may activate astrocytes toward a reactive phenotype, downregulating monocarboxylate transporters (MCT1/MCT4) and lactate dehydrogenase, reducing ANLS efficiency. Neuroinflammation is documented in ME/CFS by PET imaging [73], consistent with a persistent reactive glial state. Unlike normal post-infectious recovery, ME/CFS may involve failure to revert astrocytes to baseline (“stuck” reactive phenotype).

Testable prediction: Post-mortem or biopsy astrocyte gene expression shows persistent reactive markers (GFAP, complement C3, IL-1 α) and downregulated metabolic genes (LDH, MCT) in ME/CFS vs. controls.

Statistical test: Differential expression analysis; expected fold-change > 2 for reactive markers, < 0.5 for metabolic genes.

Novelty: Moderate—viral trigger and neuroinflammation established; persistent astrocyte metabolic shift is novel.

Certainty: 0.40 (mechanistically plausible; astrocyte-specific testing limited in ME/CFS)

Limitations. This speculation has certainty 0.40. No published studies have directly measured astrocyte MCT1/MCT4 density, lactate dehydrogenase expression, or lactate shuttle capacity in ME/CFS brain tissue. Post-mortem ME/CFS brain studies are extremely scarce, and in vivo biopsy of CNS tissue is not clinically feasible. The reactive astrocyte A1/A2 nomenclature used here is contested in the current literature, with more context-dependent phenotypic spectra now recognized. PET neuroinflammation findings [73] implicate activated glia but cannot distinguish astrocytic from microglial contributions. Evidence for viral-induced persistent astrocyte metabolic reprogramming derives from animal models and has not been replicated in human ME/CFS post-mortem or in vitro tissue.

Subtype Progression Hypothesis

~ Hypothesis 32: CNS-Primary to Global Subtype Progression

ME/CFS may progress from CNS-primary dysfunction (Subtype A) to global/multi-system involvement (Subtype D) over time as secondary cascades develop.

Relationship to parent hypothesis: Temporal evolution of the subtype classification framework.

Mechanism: Initial CNS dysfunction causes chronic elevation of autonomic regulatory demands, which over time may produce secondary peripheral abnormalities through sustained sympathetic overactivation and oxidative stress. Onset pattern data shows early ME/CFS presenting predominantly with neurological and autonomic features [120], while longer-duration ME/CFS increasingly shows multi-system metabolic and vascular involvement [12]. The proposed sequence: initial CNS dysfunction → chronic autonomic overload → secondary cardiovascular and peripheral damage → expansion from CNS-primary (Subtype A) to global (Subtype D) dysfunction.

Testable prediction: Longitudinal biomarker data shows progression: initially elevated x_4 (CNS) only, later additional abnormalities in x_1, x_2, x_3 (autonomic, peripheral).

Statistical test: Survival analysis: time to progression from Subtype A to Subtype D. Expected median progression time: 3–5 years (*estimated from informal clinical observation; no systematic longitudinal data available; this parameter requires empirical derivation*).

Novelty: Yes—subtype progression rarely studied in ME/CFS literature.

Certainty: 0.45 (progression observed clinically; systematic longitudinal data lacking)

Treatment implications. If ME/CFS follows a CNS-primary-to-global progression trajectory, early-stage intervention may prevent cascade to multi-system involvement: (1) aggressive energy management and pacing from diagnosis onset to limit cumulative CNS coordination stress; (2) regular multi-system biomarker monitoring (autonomic, cardiovascular, metabolic) to detect early secondary involvement before it consolidates [12];

(3) prioritizing treatment of CNS-primary features (orthostatic dysfunction, cognitive support) before they produce downstream peripheral consequences; (4) the hypothesis implies that treatments effective for early-stage (Subtype A) disease may be less effective once global involvement is established, providing a clinical rationale for aggressive early intervention.

Limitations. This hypothesis has certainty 0.45. Systematic longitudinal biomarker data spanning the proposed progression timeline is lacking in ME/CFS. Natural history studies are confounded by changes in diagnostic criteria, healthcare access, treatment effects, and selection bias in cohort enrollment. Subtype classification based on compartment-specific biomarkers is not yet validated or standardized. Progression may alternatively reflect worsening of a single pathological process rather than cascade to new compartments. The clinical impression that longer-duration disease shows more multi-system involvement [120] may partly reflect cumulative diagnostic recognition of pre-existing comorbidities rather than true disease spread.

14.24.11 Integration with Existing Hypotheses

The selective dysfunction hypothesis integrates with and extends existing models:

- **Metabolic Safe Mode** (Section 14.2): Selective dysfunction specifies *which* systems the safe mode affects (CNS-dependent, demand-responsive) and *which* it spares (autonomous)
- **Glymphatic Clearance Failure** (Section 14.3): Provides mechanism for CNS-specific waste accumulation within the BBB vulnerability sub-hypothesis
- **Autonomic Dysfunction** (Chapter 10): Reframes as CNS coordination failure rather than peripheral autonomic pathology
- **Mitochondrial Dysfunction** (Chapter 6): Constrains location—mitochondrial dysfunction may be CNS-specific or CNS-predominant rather than global

⚠ **Limitation 15: Selective Dysfunction Framework: Alternative Explanations Not Excluded**

The selective dysfunction hypothesis provides an internally consistent framework, but alternative explanations for the same clinical pattern have not been excluded:

- **Scale artifact:** Processes classified as “preserved” (hair, nail growth) have much lower absolute energy demands than “impaired” processes (exercise, cognition). A global but moderate energy deficit (e.g., 30% reduction) could spare low-demand processes while disabling high-demand ones — no selectivity mechanism required.
- **Peripheral contributions:** Impaired skeletal muscle oxygen extraction, documented in NIH deep phenotyping [5], provides an alternative peripheral explanation for exercise intolerance that does not require CNS coordination failure as the primary bottleneck.

- **Post-hoc fitting:** The 17 hypotheses and speculations in this chapter were developed to explain known ME/CFS features. No prediction from this framework has been tested prospectively.

14.24.12 Limitations and Uncertainties

△ **Warning 19: Limitations**

1. **Parameter estimation:** The δ_{CNS} and ρ values in Table 14.5 are estimated, not empirically derived
2. **Interaction complexity:** The additive model with interaction may oversimplify nonlinear relationships
3. **Heterogeneity:** ME/CFS likely includes multiple pathophysiological subtypes; not all may fit this model
4. **Causal direction:** The DAG assumes CNS dysfunction causes peripheral symptoms; reverse or bidirectional causation possible
5. **Evidence gaps:** Direct tests of the hypothesis (hair follicle mitochondria, CSF lactate gradients) have not been performed

14.24.13 Summary

Conclusion 1. The selective energy dysfunction hypothesis proposes that ME/CFS preferentially affects CNS-dependent, demand-responsive biological processes while sparing autonomous local processes. This explains the paradox of preserved hair growth alongside severe functional impairment.

Key features:

- Formal quantification via CNS-dependency (δ_{CNS}) and demand-responsiveness (ρ) indices
- Causal diagram with certainty-weighted edges (contains a feedback cycle; not a strict DAG)
- Three mechanistic hypotheses (Hypothesis 14.24.5: CNS energy triage; Hypothesis 14.24.5: BBB vulnerability; Hypothesis 14.24.5: sickness behavior persistence) and two speculations (Speculation 38: astrocyte energy gate; Speculation 39: partial torpor trap) explaining CNS selective vulnerability
- Ten related hypotheses and speculations extending the framework to sleep, gut-brain axis, autoantibodies, SFN, circadian rhythms, MCAS, memory, exercise physiology, post-viral reprogramming, and disease progression
- Specific, falsifiable predictions with statistical designs
- Natural subtype classification based on primary dysfunction compartment
- Treatment implications including pharmacological bypass, CNS-targeted delivery, and mechanism-specific interventions

Overall certainty: 0.55 (moderate)—hypothesis is consistent with clinical observations and supported by converging evidence from brain metabolism, autonomic dysfunction, and preserved autonomous function literature. Formal testing of predictions required.

14.25 Conclusion

The hypotheses presented in this chapter are speculative extrapolations intended to stimulate new research directions. They share several features:

- Each is grounded in established biochemistry and physiology
- Each attempts to explain the characteristic features of ME/CFS
- Each generates testable predictions
- None requires invoking unknown biology—only novel combinations of known mechanisms

The integrated “multi-lock” model suggests that ME/CFS may not have a single cause or mechanism but rather represents a stable pathological state maintained by multiple interacting processes. This perspective explains both the heterogeneity of ME/CFS and its resistance to treatment while suggesting that effective therapy may require targeting multiple mechanisms simultaneously.

These ideas are offered to the research community in the hope that some may prove fruitful and that all may contribute to the creative ferment from which scientific progress emerges.

15 Symptom-Producing Mechanisms in ME/CFS

Understanding ME/CFS requires distinguishing between mechanisms that *limit capacity*—impaired mitochondrial function, reduced cerebral blood flow, metabolic rigidity—and mechanisms that *produce symptoms*: the biological processes that generate the lived experience of fatigue, cognitive impairment, pain, and malaise. This chapter focuses on the latter. The distinction matters clinically: a patient may have preserved aerobic capacity yet be driven into exhaustion by powerful neurochemical signals that the brain interprets as danger; conversely, restoring energy metabolism without addressing symptom-signalling pathways would still leave the patient disabled.

The sections that follow trace symptom production from overarching integrative frameworks (sickness behavior, 15.1–15.2), through specific neurochemical generators (adenosine, cytokines, serotonin, melatonin, microglia, kynurenine, mast cells, 15.3–15.9), into systems-level amplifiers (glymphatic failure, central sensitization, oxidative stress, metabolic danger signals, endocannabinoid deficiency, interoceptive dysregulation, 15.10–15.15), and finally to integrated models, patient phenotypes, and therapeutic implications (15.16–15.19).

15.1 Conceptual Framework: Symptom-Producing versus Capacity-Limiting Mechanisms

Chapters 6–13 of this document establish that ME/CFS involves profound capacity-limiting pathophysiology: mitochondrial ATP synthesis deficiency, reduced cardiac output and cerebral blood flow, hypoxic metabolic drift, and neurovascular uncoupling. These mechanisms restrict the total work capacity and energy output available to a patient. A patient with severe mitochondrial dysfunction may theoretically have preserved neurochemical signaling: the brain is not activated with cytokines, neurotransmitter systems are not dysregulated, pain pathways are not sensitized. Yet such a patient would still be disabled by energy lack.

Symptom-producing mechanisms—the focus of this chapter—operate on a different axis. A patient with robust ATP synthesis, preserved cardiac output, and normal metabolic flexibility could nonetheless feel profoundly exhausted, cognitively impaired, and in pain if subjected to a sufficient barrage of pro-inflammatory cytokines, disrupted neurochemical signaling, and central sensitization. The distinction matters clinically: capacity-limiting mechanisms set the “ceiling” on what a patient can do; symptom-producing mechanisms determine how the patient *feels* across any level of capacity.

In ME/CFS, both axes are disrupted. A patient may be experiencing both ATP synthesis deficiency (Chapter 6) *and* cytokine-driven fatigue signals (Section 15.4); both reduced cardiac

output (Chapter 7) *and* mast cell-mediated brain fog (Section 15.9). The presence of capacity-limiting pathology makes symptom-producing mechanisms *more salient*, not less: cytokine fatigue signal is more easily interpreted as danger when the body is actually energy-depleted; pain signals are amplified by central sensitization when tissues are genuinely hypoxic. The interaction produces a multiplicative burden rather than a simple sum.

This distinction also has therapeutic implications. A treatment that increases mitochondrial ATP (e.g., CoQ10) without addressing inflammatory signaling will restore capacity without resolving symptoms. Conversely, a treatment that suppresses cytokine signaling (e.g., anti-TNF monoclonal antibody) without addressing energy metabolism will improve how the patient feels without expanding what the patient can do. Optimal treatment addresses both axes.

The sections that follow trace the major symptom-producing pathways: the sickness behavior cascade that integrates multiple cytokine signals into a coordinated malaise program (Sections 15.2), followed by specific neurochemical generators of fatigue and cognitive dysfunction (Sections 15.3–15.9), then systems-level amplifiers that transduce these signals into multi-organ dysfunction (Sections 15.10–15.15), and finally synthesis models and therapeutic implications (Sections 15.16–15.19).

15.2 Sickness Behavior as Overarching Integrative Framework

Sickness behavior—the coordinated constellation of fatigue, social withdrawal, anhedonia, cognitive slowing, hyperalgesia, and appetite suppression induced by immune activation—provides an overarching framework for understanding ME/CFS symptomatology. This section argues that many core ME/CFS symptoms represent a *chronic, dysregulated sickness behavior state* mediated by cytokines, prostaglandins, and vagal afferents acting on hypothalamic and limbic circuits. The adaptive logic of acute sickness behavior (conserving energy for immune defense) becomes maladaptive when chronically engaged without resolution.

15.2.1 Cytokine-to-Brain Signaling Routes

Peripheral cytokines communicate with the brain through three complementary routes [433, 434]. The *humoral pathway* operates via circumventricular organs—notably the organum vasculosum of the lamina terminalis (OVLT) and the area postrema—where an incomplete blood-brain barrier permits direct cytokine diffusion into adjacent hypothalamic tissue. A *saturable transport pathway* operates through carrier-mediated influx systems at the BBB endothelium for IL-1 β , IL-6, and TNF- α [434]. The *neural pathway* exploits vagal afferents: cytokines at peripheral infection sites activate cytokine receptors on paraganglia of the vagus nerve, transmitting immune signals to the nucleus tractus solitarius and thence to hypothalamic circuits within minutes [433, 382].

At the blood-brain barrier endothelium, circulating cytokines trigger prostaglandin E₂ (PGE₂) synthesis, which diffuses into the hypothalamic parenchyma to activate EP3 and EP4 receptors, suppressing wake-active orexin neurons and promoting sleep-pressure circuits [435, 436].

15.2.2 Hypothalamic Integration and the Sickness Behavior Program

~ Hypothesis 1: Sickness Behavior: Physiological Basis

Peripheral proinflammatory cytokines communicate with the hypothalamus via the neural and humoral pathways described above. Once there, cytokines and PGE₂ activate distinct receptor populations that coordinate a unified behavioral program: IL-1 β suppresses wake-active neurons in the lateral hypothalamus, IL-6 and TNF- α activate sleep-promoting circuits in the ventrolateral preoptic area, and PGE₂ potentiates these effects via EP3/EP4 signaling. The downstream result is the sickness behavior syndrome—fatigue, anorexia, hyperalgesia, social withdrawal, fever, and sleep dysregulation—all orchestrated by hypothalamic cytokine integration [377, 378]. This program is adaptive in acute infection (conserving energy for immune defense), but becomes pathological when chronically engaged. (Mechanism: well-established in animal and human models; certainty: High for acute sickness behavior, Medium for chronic persistence in ME/CFS).

15.2.3 Persistent Sickness Behavior in ME/CFS

~ Hypothesis 2: Persistent Sickness Behavior in ME/CFS

In acute infection, sickness behavior resolves as pathogens are cleared and cytokines return to baseline. In ME/CFS, this resolution fails: persistent immune activation (elevated TGF- β , NK hypofunctionality, activated microglia) maintains cytokine drive, while HPA axis hyporesponsiveness removes the glucocorticoid anti-inflammatory brake [379]. Post-exertional malaise may represent an acute exacerbation of this state, in which exercise triggers an abnormal cytokine spike that transiently recapitulates full sickness behavior [378, 379]. (Certainty: Medium. Mechanism coherent; direct PGE₂ measurement in ME/CFS lacking.)

15.3 Adenosine Accumulation and Pathological Sleep Pressure

Adenosine is the primary homeostatic sleep signal: it accumulates during wakefulness and is cleared during sleep via A1 and A2A receptor-mediated processes. This section examines evidence that ME/CFS patients exhibit aberrant adenosine dynamics—elevated basal levels, impaired clearance, or heightened receptor sensitivity—producing a state of chronic excessive sleep pressure that manifests as unrefreshing sleep, daytime somnolence, and post-exertional fatigue amplification.

15.3.1 Adenosine as Metabolic Waste Signal

Adenosine is generated extracellularly through two converging routes: (1) enzymatic hydrolysis of released ATP and AMP via the ecto-nucleotidase cascade (CD39 $\rightarrow$ CD73), and (2) intracellular export of adenosine formed from AMP dephosphorylation during periods of

high energetic demand [437]. In healthy tissue, extracellular adenosine remains low during wakefulness and accumulates progressively as a function of metabolic activity, creating the homeostatic sleep signal [438]. In ME/CFS, the mitochondrial ATP synthesis deficiency documented in Chapters 6–8 would predictably elevate AMP/ADP ratios even at baseline, driving constitutive adenosine generation independent of the normal wakefulness duration signal.

15.3.2 Glial Adenosine Clearance and Its Failure in ME/CFS

Extracellular adenosine is cleared primarily by astrocyte-mediated uptake via equilibrative nucleoside transporters (ENT1 and ENT2) and enzymatic degradation by adenosine deaminase [437]. Astrocytes thus act as the principal regulators of the extracellular adenosine tone that determines sleep pressure [439]. Reactive astrogliosis, which is implicated in ME/CFS neuroinflammation (see Section 15.7), disrupts this clearance machinery: reactive astrocytes show altered ENT expression and adenosine deaminase activity, impairing the buffering of extracellular adenosine surges. The predicted consequence is a chronically elevated basal adenosine tone producing pathological sleep pressure even without extended prior wakefulness.

15.3.3 A2A Receptor Upregulation in Neuroinflammation

~ Hypothesis 3: A2A Receptor Upregulation Amplifies Sleep Pressure in ME/CFS Neuroinflammation

During microglial activation, A2A adenosine receptors are upregulated coincident with P2Y12 downregulation [440]. This receptor shift has been termed chemotactic reversal: activated microglia respond to extracellular adenosine (the breakdown product of ATP) by retracting processes and adopting amoeboid morphology, rather than extending toward the signal as resting microglia do. In ME/CFS, Nakatomi et al. [441] demonstrated 45–199% elevated TSPO binding in cingulate cortex, hippocampus, amygdala, thalamus, and brainstem (n=9 patients vs. n=10 controls), establishing sustained microglial activation as a feature of the disease. The hypothesis is that this persistent neuroinflammation maintains elevated A2A receptor density across the sleep-regulatory regions, enhancing the sensitivity of the system to adenosine and producing excessive sleep pressure responses to normal or modestly elevated adenosine concentrations. (Study quality: Medium; direct ME/CFS A2A receptor expression data absent; supported by mechanistic extrapolation from Nakatomi 2014 and Orr 2009.)

A2A receptor blockade using selective antagonists reduces neuroinflammation in mixed glial cell models, with A2A antagonists outperforming A1 agonists in anti-inflammatory and antioxidant efficacy [442]. This mechanistic evidence supports the rationale for adenosine receptor-targeted interventions in inflammatory fatigue conditions, though no ME/CFS-specific clinical trials have been conducted.

15.3.4 Adenosine, Caffeine Sensitivity, and ADORA2A

Caffeine exerts its wake-promoting and fatigue-opposing effects exclusively through adenosine receptor blockade, with the A2A subtype being the primary target for sleep-regulatory effects [438]. Individual sensitivity to caffeine is substantially determined by the ADORA2A c.1083T>C single nucleotide polymorphism: carriers of specific genotypes show caffeine-induced brain electrical changes that closely resemble insomnia, while other genotypes show minimal caffeine-sleep interaction [443].

Speculation 44 (Elevated A2A Receptor Density Underlies ME/CFS Caffeine Paradox). ME/CFS patients commonly report anomalous caffeine responses: either exaggerated sensitivity at low doses, or apparent inefficacy of caffeine at standard doses. If ME/CFS is associated with upregulated A2A receptor expression (as hypothesized from the neuroinflammation evidence above), two opposing consequences are plausible. Heightened receptor density could increase binding sites for caffeine antagonism, potentially requiring higher doses for effect (apparent insensitivity). Conversely, if baseline adenosine is chronically elevated and receptors are tonically occupied, even low doses of caffeine may produce disproportionate displacement effects (exaggerated sensitivity). No ADORA2A pharmacogenetics study has been conducted in an ME/CFS cohort; this remains an untested prediction. (Certainty: Low. No ME/CFS-specific adenosine receptor density or caffeine pharmacokinetics study found in literature search.)

15.3.5 Therapeutic Implications: A2A Antagonism and Adenosine Modulation

The mechanistic model developed in this section suggests two therapeutic targets: (1) reducing adenosine overproduction by addressing its mitochondrial and neuroinflammatory drivers, and (2) modulating A2A receptor signaling directly. Non-selective adenosine receptor antagonists including theophylline and caffeine block both A1 and A2A receptors. A2A-selective antagonism is emerging as a neuroinflammation strategy with anti-inflammatory and antioxidant properties demonstrated in glial models [442].

Speculation 45 (Low-Dose Theophylline as Adenosine Antagonist in ME/CFS). Theophylline, a non-selective adenosine receptor antagonist with A1 and A2A activity, has been proposed clinically for ME/CFS given its profile of reducing adenosine-mediated sleep pressure and its established use in orthostatic intolerance via adenosine-mediated vascular tone effects. However, no published controlled trial of theophylline specifically targeting adenosine dysregulation in ME/CFS has been identified. Any clinical use should be considered highly experimental, with no evidence base beyond mechanistic plausibility. (Certainty: Very Low — mechanistic rationale only, no ME/CFS clinical evidence.)

15.3.6 Sleep Architecture Findings and Adenosine Interpretation

Systematic review of polysomnographic studies in ME/CFS identifies elevated microarousal index as the single most consistent objective abnormality: all five studies measuring microarousal index found significantly elevated values in ME/CFS patients vs. healthy controls,

while 13 studies found no difference in sleep onset latency [444]. The pattern — normal sleep initiation but pathological sleep fragmentation — is mechanistically consistent with an adenosine dysregulation hypothesis. Sleep onset depends on adenosine reaching a threshold level (preserved in ME/CFS, since sleep onset latency is normal), while sleep maintenance depends on continued adenosine-mediated suppression of arousal circuits across the night. Heightened A2A receptor sensitivity or elevated basal adenosine tone may paradoxically impair this maintenance function by saturating receptors and triggering compensatory arousal responses.

The subjective–objective discrepancy (91% of patients report non-restorative sleep despite near-normal aggregate polysomnography scores) may reflect microarchitectural disruption not captured by standard sleep staging: elevated microarousals interrupt slow-wave sleep restorative function without prolonging sleep onset or dramatically altering sleep stage proportions.

15.4 Inflammatory Cytokine-Induced Somnolence and Fatigue

Pro-inflammatory cytokines (IL-1 β , IL-6, TNF- α , IFN- α/γ) directly induce fatigue and somnolence through central mechanisms independent of peripheral tissue damage. This section reviews the cytokine-fatigue literature, the specific profile documented in ME/CFS (elevated IL-1 β , IL-6, IFN- α in subsets), and the downstream signaling pathways by which circulating cytokines access the brain to suppress arousal circuits and activate fatigue-signaling pathways.

15.4.1 Cytokine-to-Brain Signaling Routes

Peripheral cytokines communicate with the brain through three complementary routes [433, 434]. The *humoral pathway* operates via circumventricular organs—notably the organum vasculosum of the lamina terminalis (OVLT) and the area postrema—where an incomplete blood-brain barrier permits direct cytokine diffusion into adjacent hypothalamic tissue. A *saturable transport pathway* operates through carrier-mediated influx systems at the BBB endothelium for IL-1 β , IL-6, and TNF- α [434]. The *neural pathway* exploits vagal afferents: cytokines at peripheral infection sites activate cytokine receptors on paraganglia of the vagus nerve, transmitting immune signals to the nucleus tractus solitarius and thence to hypothalamic circuits within minutes [433, 382].

At the BBB endothelium and within perivascular spaces, circulating cytokines trigger prostaglandin E₂ (PGE₂) synthesis, which diffuses into the hypothalamic parenchyma to activate EP3 receptors, suppressing wake-active orexin neurons and promoting sleep-pressure circuits. Microglia—the brain’s resident immune cells—amplify this signal via TLR4/NF- κ B activation, producing local IL-1 β , IL-6, and TNF- α that sustain the fatigue state long after peripheral cytokine levels normalize [434].

15.4.2 Quantitative Human Cytokine-Fatigue Relationships

Observation 73 (Cytokine-Fatigue Correlation in Human Infection). Vollmer-Conna et al. [445] demonstrated in prospective cohorts with documented EBV, Q fever, and Ross River virus infection that IL-1 β and IL-6 levels correlate directly and quantitatively with fatigue severity, malaise, pain, mood disturbance, and cognitive slowing. Higher peripheral cytokine burdens predict worse sickness symptom scores, establishing a dose-dependent relationship in humans analogous to animal model data (study: multiple infection cohorts, natural infection setting, certainty: Medium-High).

Observation 74 (IFN- α Induces Neurovegetative Fatigue: RCT Evidence). Capuron et al. [446] demonstrated in a double-blind randomized trial (n=40 melanoma patients receiving IFN- α therapy) that fatigue and anorexia emerge within two weeks of cytokine exposure through mechanisms distinct from serotonergic pathways: the neurovegetative syndrome (fatigue, somnolence, anorexia) was substantially less responsive to paroxetine than the concurrent mood syndrome. This dissociation indicates that IFN- α -induced fatigue operates via non-monoaminergic circuits, consistent with direct hypothalamic cytokine signaling (RCT, n=40, certainty: High).

15.4.3 TLR4/NF- κ B as PEM Amplifier

~ Hypothesis 4: TLR4/NF- κ B Activation as PEM Cytokine Amplifier

ME/CFS patients may exhibit exaggerated and prolonged TLR4/NF- κ B activation following physical exertion, driving the post-exertional cytokine surge that underlies PEM. Light et al. [447] demonstrated that moderate exercise elicits significantly greater increases in leukocyte TLR4 gene expression in CFS patients versus controls, with elevations persisting 48 hours post-exercise and correlating with fatigue and pain severity (n=19 CFS, n=18 controls). Moneghetti et al. [448] independently documented that IL-1 β and IFN- α are specifically elevated 18 hours post-exercise in ME/CFS but not sedentary controls (n=24 vs. n=24), providing the cytokine correlate of the gene expression changes. Che et al. [189] further demonstrated that this heightened innate immune response worsens after exercise in ME/CFS, implicating TLR4/NF- κ B drive as a central mechanism.

Testable prediction: TLR4 blockade (e.g., TAK-242) or anti-IL-1 β therapy (anakinra [384]) before planned exertion should attenuate or delay PEM onset, with effect size proportional to pre-exercise TLR4 expression.

Limitations: Both studies use small samples; the causal direction (TLR4 drives PEM vs. PEM drives TLR4) remains unresolved; peripheral leukocyte TLR4 is a proxy for microglial TLR4 activity. Certainty: Medium (single studies, small n, no replication by independent groups at time of writing).

15.4.4 ME/CFS-Specific Cytokine Profile: Evidence and Limitations

Several large studies document cytokine abnormalities in ME/CFS subsets. Hornig et al. [212] identified distinct plasma immune signatures early in illness (elevated pro-inflammatory cytokines) that shift toward an exhaustion pattern in longer-duration disease. Montoya et al. [213] correlated 17 cytokines with disease severity in 192 CFS patients, with IL-17F and TGF- β showing the strongest severity correlations. Giloteaux et al. [214] conducted comprehensive proteomics analysis, identifying dysregulated cytokine networks in ME/CFS.

However, a systematic review of 15 case-control studies (Corbitt et al. [449], screening 16,702 publications) concluded that cytokine findings are *heterogeneous and inconclusive* as diagnostic markers. No universal cytokine signature has been validated. Elevations in IL-1 β , IL-6, and IFN- α/γ are documented in *subsets* of ME/CFS patients and should not be overstated as universal features.

This heterogeneity is consistent with ME/CFS being a syndrome of multiple convergent pathomechanisms: cytokine-driven fatigue may predominate in some patient subsets while other mechanisms (adenosine dysregulation, mitochondrial dysfunction, central sensitization) predominate in others.

15.5 Serotonin Dysregulation and the Mood-Fatigue Axis

Serotonin participates in fatigue signaling through central (dorsal raphe, limbic circuits) and peripheral (gut-brain axis, platelet) pathways. This section examines ME/CFS-associated serotonin dysregulation: altered tryptophan metabolism (competing with kynurenine pathway), platelet serotonin transport abnormalities, and dysregulated 5-HT receptor sensitivity. It also addresses the paradox of SSRI responses in ME/CFS and what they reveal about serotonin's role.

15.5.1 Tryptophan Partitioning in Inflammatory States

Observation 75 (Tryptophan Partitioning in ME/CFS). Under pro-inflammatory conditions characterizing ME/CFS, the enzyme indoleamine-2,3-dioxygenase (IDO1/IDO2) shifts tryptophan catabolism from serotonin synthesis toward the kynurenine pathway [231, 232]. Metabolomic studies in ME/CFS cohorts confirm lower serum serotonin alongside altered kynurenine metabolites compared to controls (n=38 [188]; n=35 [351]). Elevated tryptophan-to-serotonin ratios in male ME/CFS patients suggest impaired tryptophan-to-serotonin conversion [188]. (Certainty: Medium.)

15.5.2 5-HT Receptor Sensitivity and Genetic Association

~ Hypothesis 5: 5-HT Receptor Sensitivity Abnormalities in ME/CFS

Genetic variation in the 5-HT_{2A} receptor gene (HTR2A) may contribute to ME/CFS pathophysiology. Smith et al. [450] identified three HTR2A polymorphisms significantly associated with CFS (n=137); the rs6311 A-allele increases promoter activity and creates a transcription factor binding site, potentially conferring enhanced 5-HT_{2A} receptor expression. Anti-5-HT autoantibodies are present in 61.5 % of ME/CFS patients versus 5.7 % of healthy controls [451], and their presence correlates with hyperalgesia, neurocognitive dysfunction, and autonomic symptom severity—consistent with sensitized serotonergic signaling. (Certainty: Low-Medium; single genetic study, replication pending.)

15.5.3 Serotonin Transporter Abnormalities

Observation 76 (Serotonin Transporter Abnormalities in ME/CFS). Multiple lines of evidence implicate serotonin transporter (5-HTT) dysfunction in ME/CFS. In vivo PET imaging demonstrates reduced 5-HTT density in the rostral anterior cingulate cortex of CFS patients [256]. Genetically, longer L and XL allelic variants of the 5-HTT promoter (SLC6A4) are enriched in CFS patients (n=78) [452]. At the cellular level, neuroinflammatory microglial activation drives IL-1 β -mediated upregulation of astrocytic 5-HTT, reducing extracellular serotonin and impairing 5-HT_{1A} signaling [453]. (Certainty: Low-Medium.)

15.5.4 Gut-Peripheral Serotonin and IBS Comorbidity

~ Hypothesis 6: Gut-Peripheral Serotonin Dysregulation and IBS Comorbidity

Given that approximately 90–95 % of body serotonin is synthesized by enterochromaffin cells in the gastrointestinal mucosa, gut dysbiosis and bacterial translocation documented in ME/CFS [451] may disrupt peripheral serotonin pools and drive 5-HT autoimmunity. Anti-5-HT autoantibody positivity in ME/CFS is associated with elevated IgA responses to gram-negative lipopolysaccharide—consistent with intestinal permeability—and correlates with multi-domain symptom severity [451]. This may partially explain high IBS comorbidity rates in ME/CFS, with shared enterochromaffin serotonergic dysfunction as a candidate mechanism. (Certainty: Medium; correlational, causality unestablished.)

15.5.5 The SSRI Paradox: Central Serotonin Hyperactivity

~ Hypothesis 7: The SSRI Paradox: Central Serotonin Hyperactivity in ME/CFS

The frequent clinical observation that SSRIs are unhelpful or worsening in ME/CFS—in contrast to their efficacy in depression—is mechanistically explained by the hyper-serotonergic hypothesis. Lee et al. [354] demonstrated that high-dose fluoxetine induced serotonin spillover in the mouse dorsal raphe nucleus, causing 5-HT1A receptor desensitization and ME/CFS-like symptoms (fatigue, post-exertional malaise, orthostatic intolerance, hyperalgesia, disrupted sleep). Symptom reversal followed serotonin synthesis inhibition (p-chlorophenylalanine) and CRISPR-mediated 5-HT1A knockdown, establishing directional causality. Depression involves serotonin *hypo*activity; ME/CFS central fatigue may involve *hyper*activity with secondary receptor downregulation. (Certainty: Medium—animal model, human mechanistic replication pending.)

Testable predictions.

- ME/CFS patients with pre-existing SSRI use should show greater symptom severity than unexposed patients, controlling for depression.
- Serotonin synthesis inhibition or 5-HT1A agonism (e.g. buspirone) would reduce fatigue severity if this model is correct.
- Elevated serum or CSF serotonin should be present in the subset of patients with SSRI-worsened illness.

15.5.6 Central Fatigue: Two-Phase Model

Speculation 46 (Central Fatigue: From Serotonin Trigger to Kynurenine Maintenance). The classical serotonin hypothesis of central fatigue posits that elevated brain serotonin impairs motor output and promotes fatigue [343]. Yamashita's fatigue circuit model [454] refines this: serotonin release is transient during exercise, while kynurenic acid accumulation in the hypothalamus-hippocampal circuit persists and correlates better with fatigue duration. A two-phase model may apply in ME/CFS: serotonergic hyperactivity in the dorsal raphe (Section 15.5.5) may serve as an initial trigger, while tryptophan partitioning toward the kynurenine pathway (Section 75) sustains fatigue through kynurenic acid-mediated glutamate suppression and dopamine depletion [231]. (Certainty: Low; integrative speculation, not empirically tested as a unified model.)

15.6 Melatonin Dysfunction and Circadian Disruption

Melatonin abnormalities in ME/CFS extend beyond simple sleep-onset dysregulation: altered peak timing, blunted amplitude, and disrupted interactions with the immune system contribute to the characteristic unrefreshing sleep and post-sleep symptom burden. This section covers melatonin's immunomodulatory functions, its anti-inflammatory and antioxidant

roles, and how circadian misalignment creates a self-reinforcing cycle of poor sleep quality and immune dysregulation.

15.6.1 DLMO and Multi-System Circadian Decoupling

Healthy circadian function is characterized by tight coupling between dim-light melatonin onset (DLMO), core body temperature rhythm, and activity cycles. In ME/CFS this coupling is disrupted: while DLMO timing itself may not differ significantly from controls, the normal correlation between DLMO and temperature acrophase is absent in patients [400, 455]. ME/CFS patients also lack the midday temperature rise observed in healthy individuals and exhibit an anomalous evening temperature drop, consistent with multi-system circadian decoupling rather than a simple phase delay [455].

Melatonin amplitude abnormalities show a phenotypic split by age: adolescent CFS patients demonstrate significantly elevated nocturnal melatonin at midnight and into the early hours ($p < 0.001$ vs. controls) [456], while adult CFS patients with delayed circadian phase (DLMO >21:30h) report characteristic fatigue and sleep symptom patterns amenable to chronotherapy [457]. This heterogeneity matters clinically: melatonin supplementation is not appropriate for patients already secreting supratherapeutic nocturnal levels [456].

Objectively measured sleep architecture in ME/CFS confirms the unrefreshing sleep phenotype: adults show increased sleep onset latency, increased wake after sleep onset, reduced sleep efficiency, decreased stage N2 sleep, paradoxically increased slow-wave sleep (N3), and longer REM latency in a meta-analysis of 24 studies ($n = 801$ adults) [458]. The paradoxical N3 increase alongside subjective unrefreshing sleep suggests that circadian misalignment may disrupt the *quality* and restorative function of slow-wave sleep without reducing its measured duration.

15.6.2 Melatonin–Immune Axis: NK Cells and Cytokine Rhythms

Melatonin functions as a circadian immunomodulator. Mechanistically, melatonin promotes NK cell maturation and activation via the JAK3/STAT5 signaling pathway, increasing T-bet expression and thereby enhancing NK cell proliferation, degranulation, and IFN- γ secretion [459]. The circadian clock itself regulates rhythmic NK cell activity and cytokine release through clock genes including *Per2*, *Bmal1*, and *ROR α* [402].

Speculation 47 (Circadian Disruption Compounds NK Hypofunctionality). In ME/CFS, NK cell cytotoxic function is consistently reduced. If nocturnal melatonin signaling is abnormal—whether due to delayed phase, blunted amplitude, or multi-system circadian decoupling—the JAK3/STAT5/T-bet pathway driving NK maturation may be chronically understimulated. This provides a mechanistic bridge between circadian dysfunction and the well-documented NK cell hypofunctionality in ME/CFS, and may contribute to impaired viral clearance and immune dysregulation. This hypothesis remains untested directly in ME/CFS populations and is consistent with but not proven by existing data [459, 455, 460].

15.6.3 Therapeutic Melatonin: Low-Dose Chronobiotic Strategy

The pharmacology of melatonin as a chronobiotic is dose- and timing-dependent. Phase response curve studies establish that 0.5 mg melatonin, taken 2–4 h before DLMO, produces phase advances equivalent to those achieved with 3.0 mg when each dose is given at its respective optimal time [461]. Low-dose melatonin is preferred in ME/CFS for several reasons: it avoids supraphysiological blood levels, does not suppress endogenous pineal secretion, and produces minimal sedation.

In CFS patients with objectively delayed DLMO (>21:30h), open-label melatonin (5 mg, 5 h before DLMO) for 3 months produced significant improvements in fatigue, concentration, motivation, and activity on the Checklist Individual Strength (CIS); fatigue normalized in 8 of 27 patients during treatment vs. 2 of 29 pre-treatment [457]. The absence of a placebo control limits causal inference but provides directional evidence for DLMO-stratified treatment.

In delayed sleep-wake phase disorder (DSWPD)—a condition overlapping with ME/CFS in a significant proportion of patients—an RCT ($n = 40$) of 0.5 mg melatonin combined with evening dim light and time-in-bed scheduling for 4 weeks demonstrated significant improvements in DLMO timing, sleep parameters, and fatigue [462]. Importantly, melatonin timing based on estimated DLMO (from actigraphy sleep-onset data) was as effective as timing based on formally measured DLMO, making this approach clinically practical where DLMO testing is unavailable.

Observation 77 (Melatonin Dosing and Timing in ME/CFS). Current evidence supports low-dose melatonin (0.5 mg) taken 2–3 h before the individual's estimated or measured DLMO for patients with objectively delayed circadian phase. Higher doses (3–5 mg) may be appropriate when phase delay is severe, with timing adjusted accordingly. Melatonin supplementation should not be initiated in patients with elevated nocturnal melatonin or without evidence of phase delay, as benefit appears confined to the phase-delayed subgroup [457, 456, 401].

15.6.4 Light Therapy and Circadian Hygiene: Considerations in ME/CFS

Bright light therapy is the standard first-line treatment for delayed sleep-wake phase disorder, acting via direct SCN resetting. However, a controlled trial in unselected CFS patients found neither melatonin nor phototherapy produced symptomatic improvement [401], likely because most patients in that cohort were not stratified by circadian phenotype. The evidence from van Heukelom et al. [457] and Swanson et al. [462] suggests that benefit is confined to those with objectively delayed phase.

Clinically, light therapy in ME/CFS requires additional caution beyond standard DSWPD protocols. Morning bright light exposure carries an orthostatic and energy cost that may precipitate or worsen post-exertional malaise in sensitive patients. Evening dim light (maintaining <10 lux in the 2 hours before desired bedtime) and blue-light filtering are lower-risk components of circadian hygiene that can be combined with timed melatonin without the same exertion burden.

15.6.5 Melatonin as Antioxidant

Beyond its chronobiotic and immunomodulatory roles, melatonin functions as a mitochondria-targeted antioxidant: intramitochondrial concentrations exceed blood levels substantially, positioning it at the primary site of reactive oxygen species (ROS) production in the electron transport chain [463]. Melatonin scavenges ROS and reactive nitrogen species (RNS) directly, stimulates superoxide dismutase, glutathione peroxidase, and catalase, and suppresses pro-oxidant enzymes. Its metabolites N¹-acetyl-N²-formyl-5-methoxykynuramine (AFMK) and N-acetyl-5-methoxykynuramine (AMK) retain antioxidant activity, creating a cascade of protection [463]. This antioxidant function is particularly relevant to the oxidative and nitrosative stress mechanisms discussed in Section 15.12; circadian disruption reducing nocturnal melatonin may amplify the ROS burden in ME/CFS mitochondria.

15.6.6 Integrated Circadian Vicious Cycle

These mechanisms constitute a self-reinforcing cycle. Circadian disruption reduces nocturnal melatonin signaling, impairing NK cell activation and antiviral surveillance, promoting immune dysregulation and cytokine-driven symptom flares that are themselves chronodisruptive. Simultaneously, reduced melatonin permits greater mitochondrial oxidative stress during the night, contributing to cellular energy deficits and neuroinflammation. Poor sleep architecture—with preserved but non-restorative slow-wave sleep and prolonged REM latency [458]—fails to clear metabolic waste (see Section 15.10), perpetuating cognitive symptoms and pain sensitization. The result is a chronobiological vicious cycle in which impaired melatonin function amplifies, and is amplified by, the immunological and bioenergetic pathology characteristic of ME/CFS [460, 455].

15.7 Microglia Activation and Neuroinflammatory Fatigue

Microglia—the brain’s resident immune cells—when chronically activated produce a neuroinflammatory state that suppresses neural circuit efficiency and generates fatigue and cognitive symptoms. This section presents the neuroinflammatory hypothesis of ME/CFS fatigue: evidence from neuroimaging (PET glial activation), CSF studies, and post-mortem data supporting microglial activation, and the downstream consequences for synaptic function, neurotransmitter reuptake, and the blood-brain barrier.

15.7.1 Neuroimaging Evidence: TSPO-PET and MRS

Positron emission tomography using the translocator protein (TSPO) ligand ¹¹C-(R)-PK11195 provides the most direct *in vivo* evidence for microglial activation in ME/CFS. Nakatomi et al. [441] demonstrated widespread neuroinflammation in 9 CFS/ME patients versus 10 controls, with significant TSPO binding increases in the cingulate cortex, hippocampus, amygdala, and thalamus, correlating with cognitive impairment severity (high certainty for PET methodology; small sample limits generalizability).

However, the evidence is not uncontested. Raijmakers et al. [464] found no signs of neuroinflammation in 13 women with CFS using the same TSPO ligand, highlighting the importance of patient phenotyping, cohort selection, and methodological heterogeneity. VanElzakker et al. [465] argue that most neuroimaging studies inadequately target the brainstem, which may be the primary site of neuroinflammatory activity.

Magnetic resonance spectroscopy (MRS) provides a complementary, non-invasive window: Mueller et al. [270] found elevated lactate-to-creatine ratios in the right insula, thalamus, and cerebellum of ME/CFS patients, with brain temperature elevations consistent with inflammatory metabolic shifts. A meta-analysis of 65 neuroimaging studies confirmed hypoactivity in the insular and thalamic regions as the most consistent finding [246].

15.7.2 Microglial Phenotype Shift: From Surveillance to Reactivity

Under homeostatic conditions, microglia perform synaptic surveillance and phagocytic clearance. In ME/CFS, the neuroglial failure hypothesis proposes that persistent triggering factors (viral remnants, autoantibodies, metabolic danger signals) shift microglia toward a reactive M1-like state characterized by release of $\text{TNF-}\alpha$, $\text{IL-1}\beta$, IL-6 , and reactive oxygen species [466]. This reactivity may be self-perpetuating: microglial ROS production damages local neurons, releasing further danger signals that sustain activation.

~ Hypothesis 8: Neuroglial Failure as ME/CFS Pathobiological Core

Renz-Polster et al. [466] propose that impaired or pathologically reactive neuroglia—microglia, astrocytes, and oligodendrocytes—constitute the common denominator of ME/CFS pathobiology. Under this model, microglial reactive state suppresses neural circuit efficiency and disrupts glial metabolic support to neurons, producing fatigue, cognitive impairment, and post-exertional malaise. The hypothesis extends naturally to Long COVID, which shows neuroglial activation on post-mortem analysis. Certainty: Medium (theoretical synthesis; not yet empirically tested as a whole; component mechanisms supported by animal and in vitro data).

15.7.3 Circulating Danger Signals and Microglial Activation

A key mechanistic question is how peripheral pathology in ME/CFS crosses into the brain to activate microglia. Gottschalk et al. [467] demonstrated that serum from ME/CFS patients directly stimulates reactive oxygen species (ROS) and nitric oxide production in human microglial cells *in vitro*. The authors identified elevated ATG13 (autophagy-related protein 13) in ME/CFS serum as the active factor, acting via RAGE (receptor for advanced glycation end products) on microglial cell surfaces. Neutralization of ATG13 substantially reduced the oxidative stress response, implicating impaired autophagy as an upstream driver of neuroinflammation (certainty: Low-Medium; in vitro only; requires in vivo validation).

15.7.4 Purinergic Signaling and Microglial Danger Response

Microglia express P2X and P2Y purinergic receptors that respond to extracellular ATP released from damaged or metabolically stressed cells. In the cell danger response (CDR) framework, persistent purinergic signaling maintains microglia in a reactive state. The adenosine A2A receptor, upregulated on reactive microglia, further modulates this activation cycle. In ME/CFS, where post-exertional metabolic disturbance is cardinal, exercise-induced ATP release may serve as a recurring microglial activation trigger, providing a mechanistic link between physical exertion and neuroinflammatory symptom exacerbation.

15.7.5 Complement Cascade and Synaptic Pruning

Reactive microglia co-opt the complement cascade to tag synapses for elimination. Under normal developmental conditions, C1q, C3, and C4 opsonize weak synapses for phagocytosis via microglial CR3 receptors—a process essential for circuit refinement. In ME/CFS, chronic microglial reactivity may sustain inappropriate synaptic pruning in adulthood, with potential consequences for cognitive processing speed and synaptic transmission efficiency [466].

Speculation 48 (Excess Synaptic Pruning as a Substrate for Cognitive Symptoms). By analogy with neurodegenerative conditions in which complement-mediated synaptic loss precedes neuronal death, ME/CFS neuroinflammation may drive a subthreshold but functionally significant loss of synaptic contacts in prefrontal and hippocampal circuits. This mechanism could explain “brain fog” disproportionate to neuronal loss. The hypothesis is speculative in ME/CFS specifically and requires direct complement and synapse density measurements in ME/CFS post-mortem tissue. Certainty: Low (extrapolated from Alzheimer’s disease and developmental biology literature; no ME/CFS-specific data yet).

15.7.6 Therapeutic Modulation of Microglial Activation

Two agents with established anti-neuroinflammatory profiles have been investigated in ME/CFS: low-dose naltrexone (LDN) and minocycline.

Low-dose naltrexone. At doses of 1.5–4.5 mg, naltrexone antagonizes toll-like receptor 4 (TLR4) on microglial cells, interrupting a key activation signal and reducing downstream cytokine release [468, 34, 469]. The mechanism is distinct from its opioid antagonism at higher doses. Retrospective and open-label data suggest symptom benefit in subsets of ME/CFS patients, though randomized controlled trial evidence is lacking.

Minocycline. Minocycline, a tetracycline antibiotic with independent anti-inflammatory and neuroprotective properties, inhibits microglial activation and reduces ROS production. Miwa [470] conducted an open-label trial in 100 ME patients (42-day course, 100 mg/day), reporting a favorable performance-status response in 27% of participants, with best outcomes in those within six months of disease onset. A subsequent pilot study confirmed higher response

rates (80%) in early-stage patients. As Numata [471] observes, the modest and heterogeneous response underscores that neuroinflammation represents only one pathobiological thread in ME/CFS, and patient stratification by disease stage and phenotype is essential before trialling targeted anti-neuroinflammatory agents.

~ Hypothesis 9: Microglial Modulation as Stage-Dependent Therapy

Therapeutic benefit from anti-microglial agents (LDN, minocycline) may be concentrated in early-stage ME/CFS, when active neuroinflammation is more likely to be driving symptoms rather than fixed structural or epigenetic changes. In later stages, microglial modulation alone may be insufficient without addressing upstream triggers (persistent viral antigens, autoantibodies) and downstream consequences (glymphatic failure, synaptic remodeling). Certainty: Low-Medium (mechanistic plausibility supported; clinical evidence limited to open-label data).

15.8 Kynurenine Pathway and Quinolinic Acid Excitotoxicity: The “Fog Machine”

When tryptophan is shunted into the kynurenine pathway by IDO-1 (induced by IFN- γ and other inflammatory mediators), downstream metabolites diverge into neuroprotective (kynurenic acid) versus neurotoxic (quinolinic acid, 3-hydroxykynurenine) branches. This section argues that in ME/CFS, the balance is shifted toward neurotoxic metabolites, producing NMDA receptor overstimulation, excitotoxic neural stress, and the characteristic cognitive dysfunction. This pathway mechanistically connects immune activation to brain fog.

~ Hypothesis 10: Quinolinic Acid Excitotoxicity in ME/CFS

Proinflammatory cytokines (IFN- γ , IL-1 β , TNF- α) upregulate IDO-1, diverting tryptophan from serotonin synthesis into the kynurenine pathway [230, 231]. Downstream, kynurenine monooxygenase (KMO) converts kynurenine to 3-hydroxykynurenine and ultimately to quinolinic acid (QUIN), a potent NMDA receptor agonist [472]. At sub-threshold concentrations, QUIN drives hippocampal dysfunction, oxidative stress, and mitochondrial injury rather than frank excitotoxic cell death, potentially explaining the cognitive impairment and dysexecutive symptoms in ME/CFS [231, 232]. (Certainty: Medium. KP dysregulation in ME/CFS is replicated; direct QUIN measurement in brain parenchyma is technically challenging and understudied.)

~ Hypothesis 11: KYNA–QUIN Balance and Brain Fog

Kynurenic acid (KYNA), produced by kynurenine aminotransferases (KAT) from kynurenine, is a broad-spectrum NMDA and α -7 nicotinic receptor antagonist that counteracts QUIN toxicity [473, 230]. However, elevated KYNA itself impairs cognition by suppressing cholinergic and dopaminergic neurotransmission. The net cognitive outcome in ME/CFS thus depends on the dynamic QUIN/KYNA ratio: when QUIN dom-

inates, excitotoxic stress predominates; when KYNA dominates, a sedating “fog” effect predominates [473]. (Certainty: Medium. Ratio hypothesis is mechanistically coherent; direct human data in ME/CFS is limited to peripheral measurements.)

15.9 Mast Cell Mediators and Histaminergic Symptom Generation

Mast cells, strategically positioned at the blood-brain barrier, gut mucosa, and skin, release a broad array of mediators (histamine, prostaglandins, leukotrienes, substance P, TNF- α) that produce multi-system symptoms. This section examines evidence for mast cell activation syndrome (MCAS) overlap with ME/CFS, the histaminergic component of brain fog via H3 receptor-mediated modulation of acetylcholine and dopamine release, and the role of mast cell-mediated neuroinflammation in symptom amplification.

15.9.1 MCAS-ME/CFS Overlap and Prevalence

Approximately 15–25% of ME/CFS patients meet clinical criteria for MCAS [229]. Shared features include post-exertional worsening, orthostatic intolerance, mast-cell-type symptoms (flushing, angioedema, GI dysfunction), and trigger sensitivity to foods and environmental factors. The MCAS-positive ME/CFS subgroup responds better to mast-cell-directed treatment [173, 229], suggesting that mast cell activation contributes substantially to symptoms in a meaningful patient subset.

15.9.2 Histamine Receptor Pharmacology

Mast cell-derived histamine acts on four histamine receptor subtypes:

- H1R (postsynaptic): mediates allergic inflammation, itch, vascular permeability
- H2R (gastric, cardiac): regulates gastric acid, cardiac chronotropy
- H3R (presynaptic autoreceptor and heteroreceptor): inhibits release of histamine, dopamine, acetylcholine, norepinephrine, and serotonin. H3R activation = reduced neurotransmitter tone (cognitive slowing, fatigue)
- H4R (immune cells, gut): regulates immune cell chemotaxis

~ Hypothesis 12: H3 Receptor Mediation of Histamine-Induced Cognitive Impairment

Mast-cell-derived histamine, whether released systemically (as in MCAS) or locally within the central nervous system, acts on histamine H3 receptors (H3R) — presynaptic autoreceptors and heteroreceptors exclusively expressed in the brain [474]. H3R activation inhibits the release of dopamine, acetylcholine, and norepinephrine in the frontal cortex,

hippocampus, and striatum, reducing neurotransmitter tone. This mechanism could contribute to the brain fog, impaired attention, and motivational deficits characterizing ME/CFS [474, 173]. Prediction: H3R inverse agonists (e.g., pitolisant) would attenuate cognitive symptoms in ME/CFS patients with elevated histamine or MCAS overlap. (Certainty: Low-Medium. H3R pharmacology is well-established; direct evidence in ME/CFS patients is currently absent.)

15.9.3 Mast Cell–Microglia Bidirectional Communication

Mast cell degranulation products (histamine, tryptase, IL-1 β , TNF- α) activate microglia via protease-activated receptor 2 (PAR2) and purinergic receptors [432, 475]. Activated microglia reciprocally stimulate further mast cell degranulation, creating a positive feedback loop that amplifies neuroinflammation beyond what either cell type could sustain independently. Tryptase-mediated blood-brain barrier disruption additionally allows peripheral immune access to the CNS, potentially explaining why some ME/CFS patients exhibit fluctuating, trigger-sensitive symptom bursts characteristic of mast cell biology [229, 245].

~ Hypothesis 13: Mast Cell–Microglia Amplification Loop in ME/CFS Neuroinflammation

The mast cell-microglia loop is demonstrated in vitro and in animal models; its specific role in ME/CFS requires direct investigation. However, the documented mast cell activation prevalence in ME/CFS (15–25% definite MCAS) suggests that this loop may amplify neuroinflammatory symptom burden in a significant subgroup. (Certainty: Medium.)

15.9.4 Substance P in Pain Amplification

Substance P is released from sensory neurons and activates mast cells via MRGPRX2/NK1 receptors. Activated mast cells release more substance P (amplification loop) and histamine/tryptase that sensitize nociceptors, contributing to allodynia, widespread pain, and central sensitization in ME/CFS [475, 432].

15.10 Glymphatic Dysfunction and Brain Waste Accumulation

The glymphatic system—the brain’s lymphatic-equivalent waste-clearance network—operates primarily during slow-wave sleep, flushing metabolic byproducts (tau, amyloid, glutamate, inflammatory mediators) from the interstitial space. This section examines evidence that glymphatic dysfunction in ME/CFS, driven by impaired slow-wave sleep architecture and aquaporin-4 dysregulation, creates a cycle: poor sleep → waste accumulation → cognitive impairment and neuroinflammation → worse sleep. This provides a mechanistic link between unrefreshing sleep and brain fog.

15.10.1 Glymphatic System Physiology and Sleep Dependency

The glymphatic system (named by Nedergaard lab, 2013) is a brain-wide fluid exchange network in which cerebrospinal fluid (CSF) flows along perivascular spaces (para-arterial), exchanges with interstitial fluid (ISF) through astrocytic AQP4 water channels, and drains via para-venous spaces and meningeal lymphatics. This system clears tau, amyloid-beta, and metabolic waste products. Crucially, it operates preferentially during slow-wave sleep, with a 60% increase in interstitial space during SWS compared to waking [476]. Sleep duration is less important than sleep *quality* (specifically SWS content): glymphatic clearance is maximally coupled to delta-oscillation (slow-wave) sleep, not total sleep duration [477].

15.10.2 Multiple Converging Impairments in ME/CFS

In ME/CFS, several factors converge to impair glymphatic function:

Reduced slow-wave sleep. ME/CFS patients show reduced SWS relative to total sleep time, with alpha-delta intrusion pattern disrupting SWS continuity. While total sleep time may be normal or even extended, the quality is compromised.

AQP4 dysregulation. Aquaporin-4 channels, positioned at astrocytic endfeet facing perivascular spaces, are the critical bottleneck for CSF-ISF exchange. Norepinephrine (elevated in ME/CFS due to autonomic dysregulation) inhibits AQP4 function: chronic adrenergic dysregulation in ME/CFS may chronically suppress glymphatic flow [477]. Neuroinflammation can additionally depolarize AQP4 [478].

Pre-existing neuroinflammation. Activated microglia disrupt perivascular flow dynamics.

~ Hypothesis 14: Glymphatic Failure as Driver of Cognitive Symptoms and Unrefreshing Sleep

The glymphatic system — a brain-wide CSF/ISF exchange network driven by astrocytic aquaporin-4 (AQP4) water channels and perivascular fluid dynamics — clears metabolic waste, tau, and amyloid- β primarily during slow-wave sleep [476]. In ME/CFS, multiple factors converge to impair this system: (1) reduced SWS content from alpha-delta sleep intrusion, (2) chronic adrenergic dysregulation (elevated norepinephrine) inhibiting AQP4 polarization at astrocytic endfeet, and (3) pre-existing neuroinflammation disrupting perivascular flow dynamics [477]. Waste accumulation then activates the NLRP3 inflammasome in microglia, producing IL-1 β and IL-18 that further disrupt sleep architecture [477, 478]. The resulting vicious cycle — impaired glymphatic clearance → waste accumulation → neuroinflammation → worse sleep — may explain why ME/CFS patients report unrefreshing sleep despite adequate total sleep duration [479, 477]. (Certainty: Low-Medium for the ME/CFS-specific application; the glymphatic mechanism

itself is High certainty. Direct glymphatic imaging studies in ME/CFS are lacking as of 2025.)

Observation 78 (Sleep Quality Over Duration: The Glymphatic Rationale). The glymphatic hypothesis provides a mechanistic rationale for a clinically observed but poorly explained phenomenon: ME/CFS patients often sleep for adequate or extended durations yet report profoundly unrefreshing sleep and persistent cognitive impairment [479]. Since glymphatic clearance is maximally coupled to delta-oscillation (slow-wave) sleep — not total sleep duration — the alpha-delta intrusion pattern documented in ME/CFS polysomnography (non-delta electroencephalographic activity during NREM sleep) would suppress glymphatic flow regardless of how long the patient sleeps [477, 476]. Treatment implication: improving sleep quality (SWS content) is likely more therapeutically relevant than extending sleep duration [477].

15.11 Central Sensitization and Nociceptive Pain

Pain is a frequently under-addressed symptom in ME/CFS. This section covers central sensitization—the amplification of nociceptive processing in the dorsal horn and supraspinal circuits—as the mechanism underlying the widespread, disproportionate pain and allodynia seen in ME/CFS. It distinguishes nociceptive pain (arising from altered nociception without identifiable tissue damage) from nociceptive and neuropathic pain, and connects central sensitization to the neuroinflammatory and glial mechanisms discussed earlier.

15.11.1 Wind-up and Dorsal Horn Sensitization

Repetitive C-fiber nociceptor input produces a phenomenon known as wind-up: a slow temporal summation of action potentials in wide dynamic range neurons of the dorsal horn that is experienced as progressively increasing pain. Wind-up depends on activation of *N*-methyl-D-aspartate (NMDA) receptors, which become available following sustained depolarization that removes the resting magnesium block; calcium influx then activates kinase cascades that potentiate synaptic efficacy — a process analogous to long-term potentiation in the hippocampus [480]. Once established, central sensitization manifests as dynamic tactile allodynia (pain from light touch), secondary hyperalgesia, and aftersensations, all reflecting expanded and lowered pain threshold in central circuits.

15.11.2 Evidence for Central Sensitization in ME/CFS

In ME/CFS, multiple lines of evidence demonstrate central sensitization as a disease feature rather than an epiphenomenon. Generalized hyperalgesia has been documented for electrical, mechanical, heat and histamine stimuli across skin, muscle and visceral tissues. Critically, endogenous inhibitory analgesia — the conditioned pain modulation (CPM) response that normally suppresses ongoing pain during a second noxious stimulus — is absent or blunted in

ME/CFS patients [101]. Furthermore, exercise that normally activates endogenous analgesia instead exacerbates pain in ME/CFS, accompanied by significant post-exercise upregulation of ASIC3, P2X4 and TLR4 gene expression in leukocytes, persisting for 48 h and correlating with fatigue and pain severity [481].

Dolorimetry studies confirm that ME/CFS and Gulf War Illness patients are significantly more tender than sedentary controls across all sex strata, and dolorimetry scores correlate strongly with self-reported pain (Spearman $R = -0.574$ to -0.629 , $p < 0.001$) and interoceptive symptoms [482]. These authors propose injury to midbrain and medullary descending regulatory pathways as the mechanism for simultaneous loss of antinociceptive and antiinteroceptive inhibition.

15.11.3 Nociplastic Pain Framework

The International Association for the Study of Pain adopted *nociplastic pain* as a third mechanistic descriptor in 2017, defined as “pain that arises from altered nociception” not fully explained by nociceptive or neuropathic mechanisms [483]. Clinical criteria for musculoskeletal nociplastic pain require: duration exceeding three months; regional, multifocal or widespread distribution; absence of a complete nociceptive or neuropathic explanation; and clinical signs of hypersensitivity in the pain region. Fibromyalgia is the prototype; ME/CFS pain fits the same phenotype.

Speculation 49 (Kynurenine–NMDA Link to Central Sensitization). Elevated quinolinic acid (an NMDA agonist produced by the kynurenine pathway under neuroinflammatory conditions – see Section 15.8) may provide sustained NMDA receptor activation that maintains wind-up and dorsal horn sensitization in ME/CFS. If confirmed, this would mechanistically link neuroinflammation, kynurenine pathway activation, and central sensitization in a single causal chain. (Certainty: Low; no direct human measurement of spinal QUIN in ME/CFS.)

15.11.4 Peripheral Sensitization: Substance P and CGRP

Substance P and calcitonin gene-related peptide (CGRP) released from primary afferent C fibers sensitize peripheral nociceptors and maintain neurogenic inflammation in the periphery. In the context of mast cell activation (see Section 15.9), mast cell-released histamine and tryptase further lower nociceptor thresholds, creating a peripheral–central sensitization loop. CGRP-driven neurogenic inflammation can be sustained via ROS-dependent TRPA1 activation in Schwann cells, independently of direct neuronal injury.

15.11.5 Treatment Implications

Targeting central sensitization requires centrally acting approaches:

- **NMDA antagonism:** Low-dose ketamine, memantine and dextromethorphan reduce wind-up; see also LDN (low-dose naltrexone) which attenuates microglial activation [469].

- **Alpha-2-delta ligands:** Pregabalin and gabapentin reduce calcium channel-mediated neurotransmitter release in sensitized dorsal horn circuits; widely used in fibromyalgia with modest effect sizes.
- **Pain neuroscience education:** Reconceptualizing pain as a central amplification phenomenon rather than peripheral tissue damage reduces catastrophizing and improves function in short-term studies in ME/CFS [481].

15.12 Oxidative and Nitrosative Stress as Symptom Amplifier

The nitric oxide/peroxynitrite (NO/ONOO^-) cycle and reactive oxygen species (ROS) amplify symptom production across multiple domains: mitochondrial uncoupling, protein nitration, membrane lipid peroxidation, and TRP channel sensitization. This section presents Martin Pall's NO/ONOO^- cycle hypothesis in the context of current evidence, covers ROS-induced transient receptor potential (TRP) channel activation as a pain amplifier, and connects oxidative stress to the neuroinflammatory cascade, explaining why antioxidant strategies have shown partial but inconsistent benefit.

15.12.1 The NO/ONOO^- Vicious Cycle

Speculation 50 (The NO/ONOO^- Vicious Cycle Hypothesis). Martin Pall proposed that ME/CFS is initiated when short-term stressors (infection, trauma, chemical exposure) induce pro-inflammatory cytokines ($\text{IL-1}\beta$, IL-6 , $\text{TNF-}\alpha$, $\text{IFN-}\gamma$) that upregulate inducible nitric oxide synthase (iNOS), elevating nitric oxide [484]. Nitric oxide reacts with superoxide to form peroxynitrite (ONOO^-), a potent oxidant that damages proteins, lipids and DNA. Pall proposed six positive feedback loops sustaining elevated ONOO^- : $\text{NF-}\kappa\text{B}$ activation (increasing iNOS), NMDA receptor sensitization, mitochondrial complex I/III electron leakage (generating more superoxide), tetrahydrobiopterin (BH_4) depletion via ONOO^- oxidation (causing NOS uncoupling, converting NOS into a peroxynitrite synthase), and HPA axis suppression via lowered glucocorticoid production [485]. This mechanism — named the NO/ONOO^- cycle — predicts that agents lowering multiple cycle elements simultaneously (antioxidants, NMDA antagonists, anti-inflammatory compounds) would be required for therapeutic benefit. (Certainty: Low; Medical Hypotheses journal, not empirically tested in RCTs; mechanistically plausible and consistent with downstream evidence.)

15.12.2 Glutathione Depletion and Oxidative Burden

Empirical support for oxidative stress as a disease mechanism in ME/CFS comes from neuroimaging studies. Proton magnetic resonance spectroscopy documented significantly reduced cortical glutathione (GSH) in ME/CFS patients compared with controls, with a strong inverse correlation between ventricular lactate and cortical GSH ($r = -0.545$, $p = 0.001$), and significant positive correlations between GSH and physical functioning ($\rho = 0.506$, $p = 0.001$) and energy levels ($\rho = 0.606$, $p < 0.001$) [193]. A pilot trial of N-acetylcysteine (NAC) at 1800 mg/day for four weeks normalized cortical GSH, ventricular lactate and symptom

scores, providing proof-of-concept that GSH repletion via NAC crosses the blood–brain barrier [486].

15.12.3 TRP Channel Sensitization by ROS

Reactive oxygen species activate transient receptor potential channels directly: TRPV1 (vanilloid 1, the capsaicin receptor) and TRPA1 (ankyrin 1, activated by oxidant products such as 4-hydroxynonenal and acrolein) function as cellular danger sensors, translating oxidative stress into nociceptive signaling. In the context of neurogenic inflammation, CGRP-driven TRPA1 activation in Schwann cells sustains mechanical allodynia independently of direct neuronal injury, providing a peripheral amplification loop for central sensitization.

15.12.4 Antioxidant Strategies

~ Hypothesis 15: Antioxidant Support as Symptom Modifier

Multiple antioxidant interventions address different nodes of oxidative/nitrosative stress in ME/CFS:

- **NAC** (N-acetylcysteine): GSH precursor crossing the blood–brain barrier; 1800 mg/day normalized cortical GSH and improved symptoms in pilot data [486].
- **CoQ10 + NADH**: Restores mitochondrial electron transport efficiency, reducing complex I/III electron leakage and ROS generation; RCT showed significant cognitive fatigue improvement and QoL gains.
- **Melatonin**: A mitochondria-targeted antioxidant that scavenges ROS and reactive nitrogen species (RNS), stimulates antioxidant enzymes (SOD, GPx, catalase) and chelates transition metals reducing hydroxyl radical production; mitochondrial concentrations greatly exceed blood levels.

Certainty: Medium individually; no multi-agent antioxidant RCT in ME/CFS has been completed. Consistent with the NO/ONOO[−] cycle prediction that multi-agent approaches targeting several nodes will be needed for robust benefit.

15.13 Metabolic Danger Signals and the Post-Exertional Malaise Mechanism

Post-exertional malaise (PEM) — the hallmark ME/CFS symptom — represents a pathological response to exertion that is mechanistically distinct from normal fatigue. This section proposes that PEM arises from metabolic danger signal activation: lactate and succinate accumulation triggers acid-sensing ion channels (ASICs) and activates NLRP3 inflammasome priming, converting the physiological response to exercise into a sustained inflammatory cascade. This explains PEM's delayed onset (2–48h), disproportionate severity, and multi-system manifestation.

15.13.1 The Lactate Paradox: GPR81 and Anti-Inflammatory Brake Failure

Under physiological conditions, lactate generated during exercise acts as an anti-inflammatory signal via the G-protein-coupled receptor GPR81 (also known as HCAR1 or HCA1), which is expressed on macrophages and other immune cells. GPR81 activation suppresses LPS-stimulated macrophage TNF- α and IL-6 production via AMPK/LATS-mediated YAP inactivation, disrupting the YAP–NF- κ B p65 interaction and reducing cytokine transcription [487]. Under normal conditions, lactate generated by exercise thus dampens post-exercise inflammation.

Speculation 51 (GPR81 Desensitization in ME/CFS). In ME/CFS, chronic low-grade immune activation and elevated baseline lactate may desensitize or downregulate GPR81 signaling, converting the anti-inflammatory lactate brake into a non-functional component. This would explain why exercise-generated lactate fails to suppress post-exertional immune activation in ME/CFS patients while it does so in healthy controls. (Certainty: Low; GPR81 expression and function in ME/CFS leukocytes has not been measured directly.)

15.13.2 Succinate as a DAMP and NLRP3 Inflammasome Activator

Succinate, a tricarboxylic acid cycle intermediate, accumulates in metabolically stressed or ischemic tissue where oxidative phosphorylation is impaired. Extracellular succinate acts as a danger-associated molecular pattern (DAMP), activating the NLRP3 inflammasome—a cytoplasmic innate immune sensor—to produce cleaved IL-1 β and IL-18, amplifying neuroinflammation. Succinate also stabilizes HIF-1 α , shifting cellular metabolism toward glycolysis and further lactate/succinate accumulation, creating a self-reinforcing metabolic danger loop. Abnormal TCA cycle intermediate levels (including succinate) have been documented in ME/CFS metabolomics studies, consistent with impaired oxidative phosphorylation.

15.13.3 Acid-Sensing Ion Channels and Post-Exertional Pain

Acid-sensing ion channels (ASICs), particularly ASIC3, detect local acidosis from lactate accumulation and H⁺ generation during exercise. Post-exercise leukocyte gene expression studies in ME/CFS demonstrate significantly greater increases in ASIC3, P2X4 and P2X5 mRNA compared to controls, persisting for 48 h and correlating with fatigue and pain severity [481]. ASIC3 activation contributes both to pain perception and to TLR4-mediated NF- κ B/cytokine amplification, providing a mechanistic link between exercise-induced acidosis and the sustained immune activation of PEM.

15.13.4 Two-Day CPET Evidence: Physiological Basis of PEM

The two-day cardiopulmonary exercise test (CPET) protocol provides objective physiological evidence for PEM as a real, measurable, reproducible phenomenon [488]. In the largest multi-site 2-day CPET study ($n = 84$ ME/CFS, $n = 71$ controls), ME/CFS patients showed significant Day 1 to Day 2 declines in peak work capacity, peak oxygen consumption and cardiovascular

function, while controls maintained or improved performance [1]. Severity correlates with magnitude of decline: patients with severe ME/CFS show the largest Day 2 deterioration in peak workload (−19%) compared with mild–moderate patients [140].

~ Hypothesis 16: PEM as Metabolic Danger Signal Cascade

Post-exertional malaise arises through a multi-step metabolic danger signal cascade: (1) exertion exceeds impaired aerobic threshold, generating excess lactate and succinate; (2) lactate fails to suppress immune activation via desensitized GPR81; (3) succinate activates NLRP3 inflammasome, producing IL-1 β surge; (4) ASIC3 channels detect acidosis, amplifying pain and TLR4-driven NF- κ B activation; (5) immune activation and oxidative stress persist for 24–48 h, producing the delayed-onset, multi-system PEM syndrome. This model predicts that exercise below the ventilatory threshold — but not above it — would minimize danger signal generation, consistent with pacing recommendations. (Certainty: Medium; mechanistic components individually supported; integrated model not yet tested in ME/CFS directly.)

15.13.5 Why Graded Exercise Therapy Is Contraindicated

Standard graded exercise therapy (GET) assumes deconditioning as the primary mechanism of exercise intolerance and prescribes progressive load increases. The metabolic danger signal model predicts the opposite: even modest exercise above the impaired ventilatory threshold on Day 1 produces supra-threshold ASIC3/NLRP3 activation on Day 2, exacerbating metabolic stress rather than resolving it. The consistent finding of objectively reduced Day 2 ventilatory threshold in 2-day CPET studies demonstrates that this threshold is physiologically (not psychologically) impaired [1, 140], and that progressive exercise at conventional doses worsens physiological capacity. This constitutes physiological contraindication to GET as currently prescribed.

15.14 Endocannabinoid Deficiency and Mast Cell Brake Failure

The endocannabinoid system (ECS) acts as a global neuroimmune modulator, with 2-arachidonoylglycerol (2-AG) serving as the primary agonist at CB1 and CB2 receptors to suppress neuroinflammation, reduce mast cell degranulation, and regulate pain thresholds. This section proposes that Clinical Endocannabinoid Deficiency (CECD) in ME/CFS — evidenced by reduced 2-AG levels and CB1 receptor downregulation — removes a key brake on mast cell activation and neuroinflammatory amplification, contributing to symptom severity.

15.14.1 Endocannabinoid System Architecture

The endocannabinoid system (ECS) comprises two main G-protein-coupled receptors (CB1 and CB2), two principal endogenous ligands (anandamide and 2-arachidonoylglycerol, 2-AG), and the enzymatic machinery for their on-demand synthesis and degradation. CB1 receptors

are concentrated at presynaptic terminals throughout the CNS, mediating retrograde suppression of neurotransmitter release; CB2 receptors are predominantly expressed by peripheral immune cells including mast cells, macrophages, natural killer cells and microglia where they suppress pro-inflammatory signaling. 2-AG, the more abundant ECS agonist, is synthesized from membrane diacylglycerol by diacylglycerol lipase (DAGL- α/β) and degraded primarily by monoacylglycerol lipase (MAGL), whose pharmacological inhibition substantially elevates 2-AG tissue levels.

15.14.2 Mast Cell ECS Brake and Its Failure

Mast cell degranulation — the release of histamine, tryptase, prostaglandins, leukotrienes and cytokines — is normally suppressed by CB2 receptor activation via the DAGL- β /2-AG/CB2 signaling axis. Palmitoylethanolamide (PEA) inhibits mast cell degranulation via this pathway and is meta-analytically effective for nociceptive, neuropathic and nociplastic pain (including conditions overlapping with ME/CFS) [489, 490]. Cannabidiol (CBD) additionally suppresses IgE-mediated mast cell degranulation by inhibiting Fc ϵ RI downstream signaling and calcium mobilization, with effects preserved even in CB1/CB2-depleted cells, suggesting additional non-canonical mechanisms [491].

Speculation 52 (Clinical Endocannabinoid Deficiency in ME/CFS). Ethan Russo proposed that fibromyalgia, migraine and irritable bowel syndrome share an underlying Clinical Endocannabinoid Deficiency (CECD): reduced anandamide/2-AG tone, impaired CB receptor signaling, and consequent loss of ECS-mediated homeostatic regulation of pain, sleep and immune function [492]. Updated evidence supports this hypothesis: CSF anandamide is statistically reduced in migraineurs, and PTSD — another post-traumatic condition with ME/CFS overlap — shows ECS hypofunction on advanced neuroimaging [493]. By analogy, ME/CFS may represent a CECD state in which mast cell CB2 signaling is insufficient, removing the primary brake on mast cell degranulation and neuroinflammatory amplification. Direct measurement of CSF anandamide and 2-AG, and CB receptor expression in immune cells, in ME/CFS patients is required to test this hypothesis. (Certainty: Low; extrapolated from fibromyalgia and PTSD evidence; no direct ME/CFS CSF endocannabinoid measurements published.)

15.14.3 Therapeutic Implications

Current ECS-targeting approaches with evidence in ME/CFS-relevant conditions include:

- **Palmitoylethanolamide (PEA):** Ultra-micronized PEA 600–1200 mg/day; inhibits mast cell degranulation via DAGL- β /2-AG/CB2 pathway; effective for nociplastic pain (SMD = −0.59, peak effect at 24–26 weeks) [490].
- **Cannabidiol (CBD):** Non-psychoactive; suppresses mast cell degranulation by CB1/CB2-independent mechanism; also attenuates microglial activation [491].
- **Lifestyle ECS enhancement:** Aerobic exercise (within energy envelope), sleep normalization and stress reduction each tonically elevate endocannabinoid levels, providing mechanistic rationale for pacing that includes gentle movement.

15.15 Interoceptive Prediction Error and the Bayesian Brain Fog Framework

The predictive processing framework proposes that the brain continuously generates predictions about bodily state and updates them against sensory evidence. This section applies this framework to ME/CFS: chronic interoceptive prediction errors — the brain’s model of body state persistently diverging from actual physiological signals — could produce the fatigue, cognitive effort-intolerance, and sensory hypersensitivity characteristic of ME/CFS without requiring ongoing peripheral pathology. This is not a psychosomatic hypothesis; rather, it explains how peripheral dysfunction (e.g., autonomic, immune) becomes encoded in predictive brain circuits and perpetuates symptoms even during partial physiological recovery.

15.15.1 The Free-Energy Principle and Predictive Processing

The free-energy principle, proposed by Friston, holds that biological systems minimize *surprise* (formally: variational free energy, a bound on the surprise of sensory observations) by maintaining an internal generative model of their environment and updating it against incoming sensory evidence [494]. In predictive processing terms: perception is the brain’s best explanation of sensory data, not a direct read-out; prediction errors (mismatches between model and reality) propagate upward to update the model; action drives the body to fulfill the model’s predictions. Applied to interoception — the brain’s modeling of the body’s internal state — this framework predicts that persistent, genuine perturbations in afferent bodily signals will produce persistent, amplified prediction errors experienced phenomenologically as fatigue, effort-intolerance and pain [494].

15.15.2 Interoceptive Hyper-Vigilance in ME/CFS

~ Hypothesis 17: Chronic Interoceptive Prediction Error in ME/CFS

In ME/CFS, chronic peripheral dysfunction (autonomic dysregulation, immune activation, impaired oxygen delivery) generates persistently aberrant afferent interoceptive signals. The brain’s generative model, attempting to minimize prediction error, increases the *precision weighting* assigned to interoceptive signals — effectively amplifying internal body awareness and reducing the threshold for detecting deviation from homeostatic norms. This precision upweighting manifests clinically as: heightened heartbeat discrimination accuracy, lower pain pressure thresholds, cognitive effort-intolerance, and the subjective experience of fatigue as a physiological state rather than a psychological one. Importantly, this is not a psychosomatic mechanism; it is the brain’s rational response to genuine physiological disorder. (Certainty: Medium; mechanism plausible; direct tests of prior beliefs and precision weighting in ME/CFS are limited.)

Empirical support comes from case-control studies in post-infective fatigue syndrome: patients demonstrate significantly higher accuracy on heartbeat discrimination tasks, lower pressure pain thresholds, and a distinct cardiac response profile characterized by insensitivity to

task difficulty and absence of habituation [495]. Heightened interoceptive sensitivity correlated strongly with concurrent symptoms. Complementary structural neuroimaging shows increased grey matter in the insula (the primary interoceptive cortex) in ME/CFS versus controls, with white matter reductions in brainstem pathways. Tenderness (a proxy for central sensitization) and interoceptive symptom burden are strongly correlated in ME/CFS and Gulf War Illness [482], suggesting that nociplastic and interoceptive mechanisms may co-amplify each other.

15.15.3 Distinction from Psychosomatic Frameworks

The interoceptive prediction error model is explicitly distinguished from psychosomatic or functional somatic syndrome models in two critical respects. First, the primary afferent signals in ME/CFS are genuinely abnormal (reduced VO₂max, impaired autonomic function, elevated lactate, abnormal cytokine profiles), not normal signals misinterpreted by an anxious mind. Second, the model predicts that symptom persistence should track the persistence of peripheral dysfunction, not psychological intervention alone — consistent with the observation that CBT/GET produces no sustained physiological improvement on objective exercise testing [1].

15.15.4 Why Graded Exposure Fails

Graded exposure therapy operates on the premise that catastrophizing and fear-avoidance maintain symptoms by preventing corrective experiences. This is appropriate when avoidance is maintained by maladaptive beliefs about safe activities. In ME/CFS, however, two-day CPET demonstrates that the belief “exercise causes harm” is physiologically accurate: Day 2 objective capacity declines are reproducible and measurable [1, 140]. Exposing patients to an activity that genuinely produces physiological harm (NLRP3 activation, ASIC3 upregulation, immune surge) to demonstrate it is “safe” is both theoretically inconsistent with the interoceptive error model and inconsistent with the physiological evidence.

15.15.5 Therapeutic Implications: Interoceptive Retraining

Consistent with the model, therapeutic approaches that improve interoceptive *resolution* — the ability to discriminate internal signals accurately rather than amplify them non-specifically — rather than suppress interoceptive awareness, are mechanistically predicted to be beneficial:

- **Heart rate variability (HRV) biofeedback:** Provides high-resolution, real-time autonomic feedback, potentially improving the precision and accuracy of cardiovascular interoceptive prediction, reducing allostatic load.
- **Pacing within energy envelope:** By remaining below the ventilatory threshold, pacing prevents the generation of genuine metabolic danger signals, reducing the afferent input driving prediction error.

- **Interoceptive awareness training:** Mindfulness-based approaches that explicitly target interoceptive attention have been explored in chronic pain and fatigue; effects are modest and require careful implementation to avoid effort-based exacerbation in ME/CFS.

15.16 Integrated Symptom Cascade Model

The mechanisms described in Sections 15.2–15.15 do not operate in isolation: they form a self-reinforcing symptom cascade. This section presents an integrated model showing how initial triggers (infection, immune activation, autonomic dysregulation) initiate the sickness behavior state, which then activates neurochemical symptom generators, which in turn are amplified by systems-level mechanisms, creating the chronic, treatment-resistant symptom burden of ME/CFS. The cascade model also explains phenotypic variability: different entry points and dominant pathways produce different symptom profiles.

Layer 1: Initial Trigger. An acute infection (viral, bacterial), immune challenge (vaccination, environmental antigen), or physiological stress (trauma, toxin exposure) activates the peripheral immune system, raising circulating cytokine levels (IL-1 β , IL-6, TNF- α , IFN- γ).

Layer 2: Sickness Behavior State. Peripheral cytokines communicate with the hypothalamus via vagal and humoral routes, triggering the coordinated sickness behavior program: fatigue, anorexia, social withdrawal, hyperalgesia, sleep dysregulation (Section 15.2). In healthy individuals, this state resolves within days to weeks as the infection clears and cytokines normalize. In ME/CFS, the resolution fails.

Layer 3: Neurochemical Generators. Persistent cytokine drive, now coupled with altered circadian rhythm (Section 15.6) and adenosine dysregulation (Section 15.3), sustains and amplifies fatigue signaling. The kynurenine pathway diverts tryptophan from serotonin synthesis (Section 15.5), exacerbating mood and cognitive symptoms. These neurochemical changes are reciprocally reinforced by circadian disruption.

Layer 4: Microglial Activation and Neuroinflammation. Sustained peripheral cytokine production and circulating danger signals (ATG13, metabolic byproducts) activate microglia (Section 15.7). Microglial activation amplifies local cytokine production, creating a self-sustaining neuroinflammatory state independent of peripheral drivers. Mast cells (Section 15.9) establish bidirectional amplification loops with microglia.

Layer 5: Systems-Level Amplifiers. Chronic neuroinflammation impairs glymphatic function (Section 15.10), preventing waste clearance and further driving microglial activation. Elevated kynurenine metabolites (Section 15.8) and oxidative stress (Section 15.12) contribute to NMDA receptor activation, driving central sensitization (Section 15.11) and pain amplification. Endocannabinoid deficiency (Section 15.14) removes the brake on these amplifying mechanisms.

Layer 6: PEM Loop. Exercise-induced metabolic stress (lactate, succinate accumulation) activates danger signal pathways (Section 15.13), triggering TLR4/NF- κ B and NLRP3 inflammasome activation (Section 15.4). The resulting cytokine surge recapitulates the sickness behavior state (acute, disproportionate, sustained) — post-exertional malaise. Importantly, PEM is not deconditioning; it is the acute re-engagement of the neuroinflammatory cascade triggered by metabolic stress. This explains PEM's delayed onset (24–48h), its disproportionate severity relative to the exertion that triggered it, and its multi-system nature.

Interoceptive Encoding. Throughout this cascade, the brain's interoceptive model (Section 15.15) encodes the accumulated evidence of bodily dysfunction: elevated interoceptive precision weighting, expanded pain thresholds, amplified fatigue signals. Even if peripheral pathology partially remits, the brain's model persists, perpetuating symptoms.

The cascade model predicts that treatment efficacy depends on the patient's position within the cascade and the primary dominant mechanism. A patient whose PEM is driven primarily by microglial activation may respond to LDN (anti-microglial). A patient whose primary mechanism is glymphatic failure may respond to interventions improving slow-wave sleep quality. A patient with metabolic danger dominance requires pacing to prevent exercise triggering. No single intervention targets the entire cascade; multi-mechanism approaches addressing multiple layers are theoretically more robust.

15.17 Connection to Patient Phenotypes

ME/CFS is clinically heterogeneous, and the symptom mechanism framework provides a basis for understanding this variability. This section maps dominant symptom clusters — sleep-predominant, brain fog-predominant, pain-predominant, and PEM-predominant phenotypes — onto the mechanistic pathways described above, generating testable predictions about which mechanisms will be most prominent in each phenotype and which therapeutic targets are most likely to be relevant.

Sleep-Predominant Phenotype. Characterized by severe unrefreshing sleep, excessive daytime somnolence, and sleep-dependent symptom fluctuation. Dominant mechanisms: adenosine dysregulation (Section 15.3), melatonin/circadian disruption (Section 15.6), and glymphatic failure (Section 15.10). Sleep quality (not duration) improves with circadian-targeted interventions (low-dose melatonin + evening dim light) and strategies improving slow-wave sleep continuity. Cross-reference to Chapter 2 sleep phenotype data.

Brain Fog-Predominant Phenotype. Characterized by severe cognitive impairment disproportionate to fatigue, attention/executive dysfunction, and mental effort-intolerance. Dominant mechanisms: kynurenine pathway activation driving quinolinic acid excitotoxicity (Section 15.8), mast cell-derived histamine via H3 receptor suppression of acetylcholine (Section 15.9), and impaired glymphatic waste clearance (Section 15.10). Therapeutic targets: interventions supporting glymphatic clearance, mast cell stabilization, and potentially IDO inhibitors (investigational).

Pain-Predominant Phenotype. Characterized by widespread, disproportionate pain, allodynia, and hyperalgesia. Dominant mechanisms: central sensitization with NMDA receptor overactivation (Section 15.11, amplified by kynurenine pathway Section 15.8), oxidative/nitrosative stress driving TRP channel sensitization (Section 15.12), and mast cell-mediated neurogenic inflammation (Section 15.9). Therapeutic targets: NMDA antagonists (ketamine, memantine, LDN), alpha-2-delta ligands, pain neuroscience education, and antioxidant support.

PEM-Predominant Phenotype. Characterized by severe post-exertional malaise that dominates the clinical picture, with minor baseline symptoms. Dominant mechanisms: metabolic danger signal activation (Section 15.13), TLR4/NF- κ B amplification (Section 15.4), and the inability to suppress exercise-induced lactate inflammation via GPR81 desensitization (speculation in Section 51). Therapeutic targets: strict pacing (below ventilatory threshold), potentially TLR4 antagonists (investigational), and management of post-exertional metabolic stress. Cross-reference to Chapter 5 regarding disease severity and PEM progression.

Most patients exhibit a *mixed phenotype* with elements of multiple dominant mechanisms. Phenotyping can guide stratified treatment: sleep-predominant patients prioritize circadian and sleep-quality interventions; brain fog-predominant patients prioritize glymphatic support and mast cell management; pain-predominant patients require central sensitization-targeted approaches; PEM-predominant patients require pacing-first strategies. This framework is testable: biomarker studies can correlate adenosine, kynurenine, histamine, and metabolic markers with phenotype, and RCTs can test mechanism-matched versus non-matched treatments.

15.18 Therapeutic Implications

Each symptom-producing mechanism identified in this chapter represents a potential therapeutic target. This section synthesizes the treatment implications across all mechanisms, organizing them into actionable domains: (1) targeting the sickness behavior signaling cascade; (2) neurochemical rebalancing; (3) systems-level amplifier suppression; and (4) disrupting the PEM amplification loop. It also identifies synergistic combinations and contraindicated approaches (e.g., interventions that paradoxically worsen symptom production).

Domain 1: Sickness Behavior Cascade Suppression. The upstream cytokine-to-brain communication can be targeted at multiple levels:

- Anti-cytokine monoclonal antibodies (e.g., anti-TNF, anti-IL-6) or receptor antagonists (anakinra for IL-1 β [384]): direct cytokine suppression. Mechanism: blocks the initiating signal. Limitation: systemic immunosuppression carries infection risk; ME/CFS is not universally characterized by high circulating cytokines.
- Vagal stimulation or cytokine sensing inhibition (investigational): blocks neural pathway signal transduction.

- HPA axis support via gentle stress reduction, sleep optimization: restores glucocorticoid anti-inflammatory brake.

Domain 2: Neurochemical Rebalancing.

- Melatonin (low-dose 0.5–3 mg, DLMO-timed): resets circadian phase, improves NK cell function, provides antioxidant support. Evidence: Medium (vanHeukelom 2006, Swanson 2024). Best for sleep-predominant phenotype.
- Low-dose naltrexone (LDN, 1.5–4.5 mg): blocks TLR4 on microglia, suppressing neuroinflammatory cytokine production. Evidence: Low-Medium (open-label, retrospective). Broad applicability.
- Antihistamines (H1: cetirizine, loratadine; H2: famotidine): reduce mast cell-derived histamine effects. Evidence: clinical experience, no RCT. Best for MCAS-ME/CFS overlap.
- Mast cell stabilizers (cromolyn sodium, ketotifen): prevent degranulation. Evidence: clinical experience.
- Palmitoylethanolamide (PEA, 600–1200 mg/day): supports endocannabinoid system, suppresses mast cell activation. Evidence: High for nociplastic pain (meta-analysis). Particularly for pain-predominant phenotype.

Domain 3: Systems-Level Amplifier Suppression.

- Sleep quality optimization: Improving slow-wave sleep (SWS) content (not just duration) drives glymphatic clearance. Approaches: circadian alignment, sleep hygiene, potentially low-dose melatonin. Evidence: Mechanistically grounded; direct glymphatic studies in ME/CFS lacking.
- Antioxidant support: NAC (N-acetylcysteine 1800 mg/day), CoQ10 + NADH, melatonin. Evidence: Pilot/observational. Best as multi-agent approach per NO/ONOO[−] cycle prediction.
- NMDA antagonism for central sensitization: Low-dose ketamine infusions, memantine, low-dose dextromethorphan. Evidence: Established in other pain syndromes; not directly tested in ME/CFS.
- Alpha-2-delta ligands: Pregabalin, gabapentin. Evidence: Established in fibromyalgia; ME/CFS-specific evidence absent.

Domain 4: PEM Loop Disruption.

- Pacing/energy envelope management: Remain below ventilatory threshold to prevent lactate/succinate danger signal generation. Evidence: High (mechanistically grounded, supported by 2-day CPET data). This is the primary, non-negotiable intervention for PEM-predominant phenotype.
- Avoidance of graded exercise therapy (GET): Standard GET assumes deconditioning and prescribes progressive loading, which violates the metabolic danger principle. Evidence: Contraindicated per Section 15.13.

- Potential future targets: GPR81 agonists (to restore anti-inflammatory lactate brake), TLR4 antagonists (to prevent danger signal amplification), NLRP3 inflammasome inhibitors (to block immune surge). Evidence: All investigational, not yet in clinical use.

Synergistic Combinations. Multi-mechanism approaches are theoretically superior to single-agent interventions:

- Sleep quality optimization + LDN + antihistamines: targets glymphatic + microglial + mast cell mechanisms simultaneously.
- PEA + melatonin + pacing: endocannabinoid support + circadian alignment + PEM prevention.
- Antioxidant stack (NAC + CoQ10 + melatonin) + LDN: addresses oxidative stress and neuroinflammation concurrently.

Contraindicated Approaches.

- Graded exercise therapy (GET): Violates metabolic danger principle; worsens PEM. Evidence: Contraindicated.
- High-dose SSRIs: ME/CFS may involve serotonin hyperactivity, not deficiency (Lee 2024). Adding more serotonin worsens symptoms. Evidence: Clinical observation + Lee 2024 animal mechanistic data.
- Uncontrolled bright light therapy in unselected ME/CFS: May precipitate orthostatic stress and PEM. Evidence: Williams 2002 null trial; mechanistic caution based on autonomic dysfunction.
- Aggressive immune suppression in non-cytokine-dominant patients: Carries infection risk without symptom benefit in subsets without elevated circulating cytokines.

15.19 Research Directions

The symptom-producing mechanisms framework generates multiple high-priority research directions. This section outlines: (1) mechanistic studies needed to confirm each proposed pathway; (2) biomarker development for pathway-specific assessment; (3) clinical trial designs to test mechanism-targeted interventions; and (4) the systems biology approaches required to model the cascade dynamics. Priority is given to research directions that are technically feasible within 5 years and have the highest potential clinical impact.

Priority 1: Direct Measurement of Adenosine and Metabolite Dynamics. Current adenosine hypothesis in ME/CFS rests on mechanistic inference. Study design: Measure extracellular adenosine in cerebrospinal fluid (CSF) and plasma, with simultaneous polysomnography and objective adenosine-sensitive neuroimaging (PET A2A receptor binding). Compare baseline and post-exertion adenosine kinetics in ME/CFS vs. controls. Feasibility: High (CSF sampling and PET imaging are established). Impact: Would directly test whether adenosine dysregulation is a primary or secondary feature. Expected timeline: 2–3 years.

Priority 2: Kynurenine Pathway Profiling with RCT of IDO Inhibition. Current evidence is observational (metabolomics). Study design: (1) Observational: comprehensive metabolomic profiling of the kynurenine branch in 100+ ME/CFS patients vs. controls, with phenotype correlation (sleep, brain fog, pain); (2) Interventional: Phase II RCT of IDO-1 inhibitor (e.g., 1-methyl-tryptophan analog) vs. placebo with kynurenine/QUIN/KYNA biomarkers as primary outcomes and cognitive function as secondary outcome. Feasibility: Medium (IDO inhibitors are investigational but available; kynurenine analysis is established). Impact: Would establish whether IDO inhibition improves kynurenine-driven brain fog. Expected timeline: 3–5 years.

Priority 3: Glymphatic Imaging Studies in ME/CFS. Glymphatic dysfunction is hypothesized but never directly imaged in ME/CFS. Study design: Multi-site observational study using diffusion tensor imaging (DTI) and dynamic contrast-enhanced MRI to visualize CSF-ISF exchange efficiency during sleep and waking in 50 ME/CFS patients vs. 50 controls. Correlate imaging with polysomnography (SWS)

Priority 4: Clinical Endocannabinoid Deficiency (CECD) Confirmation Study. CECD is proposed but untested in ME/CFS. Study design: Cross-sectional measurement of CSF anandamide and 2-AG levels, CB1/CB2 receptor expression (via PET), and peripheral immune cell cannabinoid signaling capacity in 50 ME/CFS patients (stratified by phenotype) vs. 50 controls. Correlate ECS dysfunction with pain, mast cell activation, and neuroinflammation markers. Feasibility: Medium (requires CSF sampling and PET availability). Impact: Would establish CECD as a validated mechanism and support PEA/CBD therapeutic development. Expected timeline: 2–3 years.

Priority 5: Phenotype-Stratified RCTs of Mechanism-Matched Treatments. Current ME/CFS trials are unselected cohort design, potentially mixing incompatible mechanisms. Study design: Phase III RCTs in each phenotype subset:

- Sleep-predominant: low-dose melatonin (DLMO-timed) + light therapy vs. placebo
- Brain fog-predominant: mast cell stabilizer (cromolyn) vs. placebo
- Pain-predominant: PEA vs. placebo
- PEM-predominant: pacing adherence support + LDN vs. standard care

Feasibility: High (trials are feasible; requires 200–300 patients total across 3–5 sites). Impact: Would establish whether phenotype-matched treatment outperforms unselected approaches. Expected timeline: 3–5 years.

Priority 6: Exercise-Induced Metabolic Danger Signal Biomarkers. The metabolic danger hypothesis predicts lactate/succinate surge, NLRP3 inflammasome activation, and ASIC upregulation post-exercise. Study design: Measure plasma lactate, succinate, NLRP3 activity (cleaved IL-18), and leukocyte ASIC3/P2X mRNA at baseline, immediately post-exercise, and 6/12/24/48h post-exercise in 30 ME/CFS patients vs. 30 controls during 2-day CPET. Correlate with PEM severity. Feasibility: High (established protocols). Impact: Would validate

metabolic danger model and enable development of biomarker-guided exercise prescriptions. Expected timeline: 1–2 years.

Priority 7: Systems Biology Modeling of the Symptom Cascade. Develop mechanistic computational models integrating cytokine dynamics, neurochemical generators, and systems amplifiers. Study design: Agent-based or network-based modeling (e.g., using data from above studies) to simulate cascade dynamics and predict treatment combinations. Validation: Prospective simulation of responses to single and combined interventions, tested against clinical trial outcomes. Feasibility: High (modeling is computationally feasible with existing data). Impact: Would provide framework for treatment optimization and prediction of which interventions will synergize vs. antagonize. Expected timeline: Parallel to above studies (1–5 years).

All these directions are designed to be complementary, iterative, and hypothesis-driven. Early results from mechanistic studies (Priorities 1–4) would inform trial design for Priority 5. Together, they would generate the mechanistic understanding and clinical evidence needed to move ME/CFS treatment from trial-and-error to precision medicine approaches.

Part III

Treatment and Management

This part comprehensively covers interventions for ME/CFS, including:

- **Medications:** Prescription drugs targeting specific symptoms and mechanisms
- **Supplements and nutraceuticals:** Vitamins, minerals, and other compounds (e.g., magnesium, CoQ10)
- **Lifestyle interventions:** Pacing, sleep hygiene, dietary approaches
- **Experimental and emerging therapies:** Cutting-edge treatments under investigation
- **Management strategies:** Practical approaches for living with ME/CFS

Evidence levels are indicated throughout, distinguishing between well-supported interventions and those with preliminary or anecdotal support.

16 Symptom-Based Management

16.1 Critical Principle: Aggressive Management of All Comorbidities

△ **Warning 1: This Cannot Be Overstated**

Every comorbid condition, no matter how seemingly minor, must be treated aggressively and promptly.

When managing ME/CFS patients with comorbidities, clinicians must understand a fundamental principle that cannot be overstated: *any improvement, even one that appears insignificant in isolation, is essential in the context of ME/CFS recovery and may directly contribute to ME/CFS recovery itself.*

≠ **Limitation 1: Comorbidity Management Rationale: Clinical Experience, Not Trial Evidence**

The comorbidity management principles in this section (lines through Section 16.2) reflect expert clinical reasoning and patient-reported experience rather than ME/CFS-specific trial evidence. Claims about metabolic reserve, energy debt compounding, and threshold effects are clinical heuristics widely used by ME/CFS specialists but have not been formally validated in controlled studies.

16.1.1 Why Comorbidity Management Is Critical in ME/CFS

ME/CFS patients operate at the absolute edge of their metabolic capacity. Unlike healthy individuals who can tolerate minor health issues without functional impact, ME/CFS patients have no reserve capacity to absorb additional burdens.

The Compounding Effect of Comorbidities

- **No buffer capacity:** Healthy individuals can manage multiple minor health issues simultaneously because they have metabolic reserves. ME/CFS patients have zero buffer—every additional symptom or condition directly subtracts from their already insufficient functional capacity
- **Energy debt compounds:** A “minor” sleep disturbance that a healthy person could ignore may cost an ME/CFS patient hours of functionality the next day. Untreated allergies that cause mild congestion may increase breathing effort enough to trigger PEM. Chronic pain that others “manage” consumes energy the ME/CFS patient cannot spare

- **Symptoms cascade:** One untreated condition triggers others. Pain disrupts sleep, poor sleep worsens cognitive function, cognitive dysfunction impairs the ability to manage symptoms, which worsens the baseline condition. In ME/CFS, these cascades rapidly become catastrophic
- **Delayed recovery:** ME/CFS recovery is measured in months to years, not days to weeks. Every day spent managing a preventable comorbidity is a day not spent recovering from ME/CFS. The cumulative cost of “minor” untreated conditions over months becomes devastating
- **Threshold effects:** ME/CFS patients often exist just below a functional threshold. A 5% improvement in energy from treating a comorbidity may be the difference between complete disability and minimal function. What appears “insignificant” to clinicians may be life-changing for patients

Clinical Approach: Treat Everything Aggressively

Clinical Imperative

Do not dismiss any treatable condition as “too minor to matter” in ME/CFS patients. What appears insignificant in a healthy patient may represent the difference between entirely housebound and able to leave the house occasionally—not to live a joyful life, but simply to make some brief outings possible. These marginal gains in severe disability are the difference between absolute confinement and minimal function.

Conditions requiring aggressive treatment in ME/CFS:

1. **Sleep disorders:** Even mild sleep apnea, periodic limb movements, or insomnia must be treated aggressively. Sleep disruption prevents the already-impaired recovery mechanisms from functioning
2. **Pain conditions:** Chronic pain (migraines, joint pain, neuropathic pain) consumes energy and prevents rest. Adequate analgesia is not optional—it is essential for energy conservation
3. **Allergies and sinus issues:** Chronic congestion, post-nasal drip, or allergic inflammation increase breathing effort and immune activation. These are not “minor annoyances”—they are energy drains
4. **Gastrointestinal disorders:** IBS, GERD, gastroparesis, food intolerances—all impair nutrient absorption and require energy to manage. Treating GI symptoms can dramatically improve overall function
5. **Endocrine dysfunction:** Hypothyroidism, adrenal insufficiency, sex hormone imbalances—even subclinical levels that might be ignored in healthy patients warrant treatment in ME/CFS
6. **Nutritional deficiencies:** Vitamin D, B12, iron, magnesium deficiencies should be corrected aggressively. Even “borderline low” values may impair function in patients already operating at the edge

7. **Infections:** Chronic sinusitis, UTIs, dental infections, fungal overgrowth—any ongoing infection must be treated promptly. The immune response and inflammation drain limited energy reserves
8. **POTS and orthostatic intolerance:** Aggressive treatment with fluids, salt, compression, and medications (fludrocortisone, midodrine, beta-blockers) can meaningfully improve function
9. **ADHD and cognitive dysfunction:** If stimulants or other ADHD medications improve function, they should be used. The cognitive energy saved may enable better symptom management overall
10. **Mental health comorbidities:** Depression and anxiety are both consequences of and contributors to ME/CFS disability. Aggressive treatment with appropriate medications and therapy is essential, not optional

The Virtuous Cycle: Physical Improvements Enable Psychological Improvements

This cannot be understated: Any treatment that allows patients to function closer to “normal”—even if still far from truly normal—creates a favorable basis for psychological improvement and may improve quality of life for everyone involved.

The Multi-Level Benefits of Physical Symptom Treatment. When comorbidities are treated and function improves, benefits cascade across multiple domains:

1. **Physical pain reduction is real and immediate:** Treating pain, sleep disorders, or orthostatic intolerance directly reduces physical suffering. This alone justifies aggressive treatment
2. **Psychological pain reduction follows:** When physical function improves, psychological suffering often decreases. Being able to shower independently, prepare a meal, or briefly leave the house reduces feelings of helplessness, dependency, and despair. The psychological burden of complete disability is partially lifted by even minimal functional gains
3. **Potential for psychosomatic improvement (if such exists in ME/CFS):** While ME/CFS is not a psychosomatic illness, improvements in physical function may create conditions where mind-body interactions—if they exist in this disease—can work in the patient’s favor rather than against them. Feeling slightly less disabled may reduce stress, which may reduce symptom exacerbation, creating a modest virtuous cycle
4. **Restoration of social connection:** When patients gain enough function to interact with family and friends—even briefly, even in limited ways—relationships can partially resume. Family members and friends may finally retrieve someone with whom they can interact in acceptable or even enjoyable ways, rather than only witnessing suffering
5. **Reduced caregiver burden:** Improvements that enable greater independence reduce the physical and emotional burden on caregivers, improving their quality of life and the relationship dynamic

6. **Hope and agency:** When treatments produce tangible improvements, patients regain a sense that their condition is not entirely beyond control. This psychological shift—from complete helplessness to having some agency—can be profoundly meaningful even when disability remains severe

The Compounding Nature of Improvement. Physical improvements enable psychological improvements, which may enable better symptom management, which may enable further physical improvements:

- Better function → reduced psychological distress → better sleep quality → improved baseline function
- Reduced pain → ability to engage in minimal activity → reduced deconditioning → less pain from movement
- Improved social connection → reduced isolation and depression → better adherence to pacing and treatment → improved outcomes
- Increased independence → restored dignity and self-worth → motivation to continue treatment → sustained improvements

For Family and Friends: The Relief of Reconnection. For loved ones who have watched the patient disappear into severe disability, even small functional improvements can be deeply meaningful:

- **Restoration of interaction:** When the patient can tolerate brief conversations or visits, relationships that had essentially ceased can resume in limited form
- **Witnessing improvement rather than only decline:** Seeing the patient gain any function provides hope and relief after potentially years of watching deterioration
- **Reduced guilt and helplessness:** When treatments help, family members feel less helpless and guilty about their inability to help
- **Acceptable or enjoyable interactions:** Moving from interactions defined entirely by caregiving and suffering to interactions that include moments of connection, conversation, or even brief enjoyment transforms the relationship

Clinical Principle: Treat for Total Quality of Life. When treating ME/CFS comorbidities, recognize that benefits extend far beyond the specific symptom being treated:

- Treating pain improves physical suffering *and* psychological well-being *and* social relationships
- Treating cognitive dysfunction improves function *and* restores agency *and* enables better symptom management
- Treating orthostatic intolerance improves physical capacity *and* enables social connection *and* reduces caregiver burden

The psychological pain is real. The physical pain is real. Both deserve aggressive treatment. Improvements in physical function create conditions for improvements in psychological state, social connection, and overall quality of life for patients and their families.

This cannot be understated.

The “Insignificant Improvement” Fallacy

Clinicians accustomed to treating otherwise healthy patients may dismiss a 5–10% functional improvement as “clinically insignificant.” **In ME/CFS, this is catastrophically wrong.**

Understanding the Percentage Baseline. When discussing percentage improvements, it is critical to understand what baseline we are measuring against:

- **If measuring relative to current function:** A patient operating at 10% of normal capacity who gains a “5% improvement” relative to their current state only moves to 10.5% of normal—seemingly trivial
- **If measuring relative to healthy baseline:** A 5% improvement *of the patient’s potential 100% capacity* is massive when the patient currently operates at 10%. This represents moving from 10% to 15% of normal capacity—a **50% relative increase** in available function
- **The clinical reality:** Most meaningful improvements are measured against the patient’s healthy baseline, not their current compromised state. A treatment that restores 5 percentage points of normal capacity when you only have 10 percentage points available *increases your functional capacity by 50%*—a massive improvement

Examples with concrete baselines:

Consider a patient currently operating at 15% of their pre-illness capacity:

- Treating sleep apnea that restores 5 percentage points (baseline) = moving from 15% to 20% = 33% relative increase in function
- Correcting vitamin D deficiency that restores 3 percentage points = moving from 20% to 23% = 15% relative increase
- Treating POTS that restores 7 percentage points = moving from 23% to 30% = 30% relative increase
- Managing chronic pain that restores 5 percentage points = moving from 30% to 35% = 17% relative increase

Cumulative result: Four “minor” interventions restore 20 percentage points of baseline capacity, moving the patient from 15% to 35% function—more than **doubling** their functional capacity.

What this means practically:

- **At 15% capacity:** Bedbound most of the day, needs assistance with basic ADLs, cannot work

- **At 35% capacity:** Can shower independently, prepare simple meals, manage basic household tasks, potentially work part-time from home with pacing
- The difference between 15% and 35% is the difference between complete dependence and minimal independence—life-changing for the patient even though clinicians might dismiss these as “small” improvements

Clinical principle: In ME/CFS, *aggregate marginal gains measured against healthy baseline matter enormously*. Small absolute improvements become massive relative improvements when the starting point is severe functional limitation. Never dismiss an intervention because it only restores “a few percentage points”—those points may represent doubling or tripling the patient’s available capacity.

Time Scales Matter

ME/CFS recovery, when it occurs, happens over months to years. Every untreated comorbidity:

- Delays the start of recovery by keeping the patient in a worsened baseline state
- Consumes energy that could otherwise go toward healing
- May trigger PEM episodes that cause setbacks lasting weeks
- Compounds over time, making the total burden exponentially worse

A treatable condition left untreated for 6 months may cost the patient a year of recovery time. The urgency of treating “minor” issues in ME/CFS cannot be overstated.

For Patients: Advocate for Comprehensive Treatment

If your clinician dismisses a symptom or comorbidity as “not significant enough to treat,” recognize that this reflects a fundamental misunderstanding of ME/CFS. You may need to:

- Explicitly explain that small improvements are critical when operating at the metabolic edge
- Request trials of treatments even for “borderline” or “mild” conditions
- Seek specialists for individual comorbidities rather than expecting your ME/CFS provider to manage everything
- Document functional improvements from treating comorbidities to demonstrate their importance

The principle: Treat everything. Every improvement counts. Nothing is too minor to matter when you’re already at the edge of functional collapse.

16.1.2 Infection Prevention as Therapeutic Priority

Observation 79 (Infections as Irreversible Ratchet Mechanism: Prevention as Critical Intervention). Patient case experience demonstrates that infections—even mild infections that a healthy person would weather without consequence—can trigger permanent step-down in baseline function in ME/CFS. Each infection (upper respiratory, influenza, COVID-19, or other pathogenic exposure) frequently causes an irreversible worsening of functional capacity, with prolonged post-infectious PEM lasting weeks to months even for mild infections. The “ratchet effect” operates at the infection level: baseline function declines step-by-step with each infection, creating cumulative deterioration from mild to moderate to severe disease. Some patients report specific infections (COVID, influenza, or opportunistic reactivation) as triggering the transition from relatively stable illness to progressive decline, suggesting that infection prevention may be among the most impactful interventions in the ME/CFS treatment toolkit.

Concrete infection prevention strategies:

- **FFP2 masking:** N95 or FFP2 masks in crowded indoor spaces, particularly during respiratory season (September–April)
- **Prophylactic antivirals:** During household exposures (family member infected), antivirals started immediately may prevent infection entirely or minimize severity
- **Immediate treatment:** Any infection symptoms treated aggressively and immediately with antivirals, antibiotics (if bacterial), or other targeted therapy rather than “watching and waiting”
- **Vaccination:** Despite potential for temporary post-vaccination PEM in some patients, infection-prevention benefit likely exceeds risk (consult with knowledgeable physician)
- **Environmental control:** Avoid crowded spaces, maintain distance from infectious individuals, ventilation optimization in home/workplace
- **Immune support:** While evidence is limited, adequate nutrition, micronutrient repletion (vitamin D, zinc, selenium), sleep quality, and pacing all support baseline immune function

Rationale: Because infections cause irreversible worsening, preventing infection is mechanistically equivalent to preventing the ratchet effect itself. An infection prevented entirely is functionally equivalent to saving months of recovery time and avoiding permanent functional loss. In a disease where recovery is difficult and decline is permanent, prevention becomes the primary treatment.

⚠ **Limitation 2: Sections 16.2–16.7: Citations Pending**

The symptom management sections below (PEM, sleep, pain, cognitive, orthostatic, autonomic) were drafted from clinical consensus and specialist experience but currently lack inline citations. Specific claims—prevalence percentages, dosage ranges, mechanism descriptions—require supporting references before these sections can be considered evidence-based. Citation integration is a priority for the next revision.

16.2 Managing Post-Exertional Malaise

Post-exertional malaise (PEM) is the hallmark feature of ME/CFS and its management is the foundation of all treatment strategies. PEM is characterized by a disproportionate worsening of symptoms following physical, cognitive, or emotional exertion, typically delayed by 12–72 hours and lasting days to weeks.

16.2.1 Pacing and Energy Envelope Theory

Pacing is the primary strategy for preventing PEM. The energy envelope theory, developed by Jason et al., posits that ME/CFS patients have a reduced energy envelope—a maximum amount of energy available for activity on any given day—and that exceeding this envelope triggers PEM. The goal of pacing is to keep activity within the envelope.

Key pacing principles include:

- **Heart rate monitoring:** Many patients find that maintaining heart rate below their anaerobic threshold (typically 50–60% of age-predicted maximum, or approximately 100–110 bpm for many patients) prevents PEM triggers. Wearable heart rate monitors provide real-time feedback
- **Baseline establishment:** Patients should identify their current sustainable activity level—the amount of activity they can perform without triggering PEM—and use this as their starting point. This baseline is often lower than patients expect
- **Activity–rest cycling:** Alternating periods of activity with mandatory rest periods (e.g., 15 minutes of activity followed by 15 minutes of rest) prevents cumulative energy depletion
- **Pre-emptive rest:** Resting *before* feeling exhausted, rather than pushing until collapse, is critical. By the time fatigue is severe, the energy envelope has already been exceeded
- **Cognitive and emotional pacing:** Cognitive tasks (reading, conversations, screen time) and emotional stressors consume energy and must be budgeted alongside physical activity

⚠ Limitation 3: Pacing Evidence Base

While pacing is widely recommended by ME/CFS specialists and patient organizations, and large patient surveys consistently rank it as the most helpful management strategy, high-quality randomized controlled trial evidence is limited. Most supporting evidence comes from observational studies, patient surveys, and clinical experience. The optimal pacing strategy likely varies by patient severity and subtype.

16.2.2 Medications for PEM

No medication is specifically approved for PEM prevention or treatment. Pharmacological approaches remain experimental and off-label:

- **Low-dose naltrexone (LDN):** Some patients report reduced PEM severity and frequency at doses of 1.5–4.5 mg nightly. The proposed mechanism involves immunomodulation via transient opioid receptor blockade (Chapter 21)
- **Low-dose aripiprazole (LDA):** Anecdotal reports and one retrospective case series suggest doses of 0.25–2 mg may reduce PEM severity, possibly through dopamine and neuroinflammation modulation
- **Anti-inflammatory agents:** Some patients benefit from NSAIDs or corticosteroids taken at PEM onset, consistent with the inflammatory component of post-exertional symptom flares
- **Mitochondrial support:** CoQ10, D-ribose, and NADH supplementation may reduce PEM severity in some patients by supporting impaired mitochondrial function (Chapter 22)

△ **Warning 2: No PEM “Cure”**

No medication reliably prevents or treats PEM. Pharmacological approaches are adjunctive to pacing, not substitutes for it. Patients who use medications to “push through” activity beyond their energy envelope risk cumulative harm.

16.3 Sleep Management

Sleep dysfunction is nearly universal in ME/CFS, affecting over 90% of patients. Characteristic features include unrefreshing sleep despite adequate duration, difficulty initiating or maintaining sleep, reversed sleep–wake cycles, and vivid or disturbing dreams. Sleep disturbance both results from and contributes to the core pathophysiology of ME/CFS, creating a vicious cycle where poor sleep worsens fatigue, pain, and cognitive dysfunction.

16.3.1 Sleep Hygiene

Standard sleep hygiene principles apply but must be adapted for the ME/CFS context:

- **Environmental modifications:** Dark, cool (18–20°C), quiet bedroom. Blackout curtains and white noise machines may help patients with sensory sensitivities. Remove screens from the bedroom
- **Consistent schedule:** Fixed wake time is more important than fixed bedtime. Gradual shift toward desired schedule (15 minutes per week) rather than abrupt changes
- **Light exposure:** Morning bright light (10,000 lux light box or natural daylight within 30 minutes of waking) helps entrain circadian rhythm. Evening blue light reduction via amber-tinted glasses or screen filters beginning 2 hours before bedtime
- **Temperature regulation:** Many ME/CFS patients have impaired thermoregulation. Cooling mattress pads, breathable bedding, and layered blankets allow temperature adjustment

- **Pre-sleep routine:** A consistent wind-down period of 30–60 minutes with low-stimulation activities. Cognitive or emotional stimulation close to bedtime can delay sleep onset by hours in ME/CFS patients

⚠ **Limitation 4: Sleep Hygiene Alone Is Insufficient**

Sleep hygiene improves sleep conditions but rarely resolves ME/CFS sleep dysfunction, which has a neurological basis (Chapter 8). Clinicians who prescribe only sleep hygiene without addressing the underlying sleep architecture disruption undertreat these patients.

16.3.2 Medications for Sleep

Pharmacological sleep management in ME/CFS often requires a trial-and-error approach, as individual responses vary substantially. Medications should be started at the lowest effective dose.

- **Melatonin:** 0.5–5 mg, 30–60 minutes before bedtime. Extended-release formulations may help with sleep maintenance. Well-tolerated with minimal side effects. Some evidence of immunomodulatory benefit in ME/CFS
- **Trazodone:** 25–100 mg at bedtime. A sedating antidepressant frequently used off-label for insomnia. Improves sleep maintenance and may increase slow-wave sleep. Low risk of dependence
- **Low-dose amitriptyline:** 5–25 mg at bedtime. Improves sleep quality and may reduce pain and headaches. Anticholinergic side effects (dry mouth, constipation) limit dose
- **Low-dose doxepin:** 3–6 mg (Silenor). Selectively blocks histamine H1 receptors at low doses, promoting sleep without significant anticholinergic effects. FDA-approved for insomnia
- **Gabapentin:** 100–600 mg at bedtime. Particularly useful when pain and sleep disturbance co-occur. Increases slow-wave sleep. Dizziness and cognitive effects may limit use
- **Pregabalin:** 25–150 mg at bedtime. Similar profile to gabapentin. FDA-approved for fibromyalgia, which frequently co-occurs with ME/CFS
- **Suvorexant/lemborexant:** Orexin receptor antagonists that may be particularly relevant given hypothesized orexin pathway dysfunction in ME/CFS. Clinical experience in ME/CFS is limited
- **Z-drugs** (zolpidem, eszopiclone): Effective for sleep initiation but do not improve the unrefreshing quality of ME/CFS sleep. Risk of dependence with long-term use; best reserved for intermittent use
- **Benzodiazepines:** Generally avoided due to dependence risk, cognitive impairment, and potential to worsen daytime fatigue. May be appropriate short-term during severe exacerbations

16.4 Pain Management

Pain affects 70–95% of ME/CFS patients and includes headaches, myalgia, arthralgia, and neuropathic pain. Pain in ME/CFS is likely mediated by central sensitization, neuroinflammation, and small fiber neuropathy (Chapter 8) rather than peripheral tissue damage, which has implications for treatment selection.

16.4.1 Analgesics

Simple analgesics provide modest relief for many patients but rarely achieve adequate pain control as monotherapy:

- **Acetaminophen** (paracetamol): 500–1000 mg up to 4 times daily. Safe for most patients but limited efficacy for neuropathic pain. Avoid exceeding 3 g/day, particularly in patients with hepatic compromise
- **NSAIDs**: Ibuprofen (200–400 mg), naproxen (250–500 mg). Useful for inflammatory pain, headaches, and myalgia. Long-term use carries gastrointestinal, renal, and cardiovascular risks; periodic use with gastroprotection is preferred. May paradoxically benefit ME/CFS through anti-inflammatory effects beyond analgesia
- **Topical agents**: Diclofenac gel, capsaicin cream, or lidocaine patches provide localized relief without systemic side effects. Appropriate for focal myalgia or arthralgia

16.4.2 Neuropathic Pain Medications

Given the central sensitization and small fiber neuropathy underlying much ME/CFS pain, neuropathic pain agents are often more effective than traditional analgesics:

- **Gabapentin**: 100–1200 mg in divided doses, titrated slowly. First-line for neuropathic pain in ME/CFS due to dual benefit on pain and sleep. Start at 100 mg at bedtime to minimize cognitive effects
- **Pregabalin**: 25–300 mg in divided doses. Similar mechanism to gabapentin with more predictable pharmacokinetics. FDA-approved for fibromyalgia
- **Duloxetine**: 20–60 mg daily. An SNRI with FDA approval for fibromyalgia, diabetic neuropathy, and chronic musculoskeletal pain. May improve fatigue alongside pain in some patients. Nausea and activation side effects limit tolerability
- **Low-dose naltrexone (LDN)**: 1.5–4.5 mg at bedtime. Growing evidence for efficacy in fibromyalgia and preliminary evidence in ME/CFS. Proposed mechanism involves microglial modulation and reduced central sensitization. Well-tolerated; vivid dreams are the most common side effect

16.4.3 Opioids

Opioid use in ME/CFS is controversial and generally discouraged as routine treatment:

- **Risks:** Dependence, tolerance, opioid-induced hyperalgesia, cognitive impairment, constipation (worsening GI dysfunction), and immune suppression—all particularly concerning in ME/CFS patients
- **Limited efficacy:** Central sensitization pain responds poorly to opioids compared to neuropathic agents
- **Potential role:** In patients with severe pain refractory to all other interventions, low-dose opioids (e.g., tramadol 50–100 mg, low-dose buprenorphine) may be considered under specialist supervision with clear monitoring and discontinuation criteria
- **Tramadol:** Has dual opioid and serotonin–norepinephrine reuptake inhibition. Some ME/CFS patients report benefit; serotonin syndrome risk with concurrent SSRIs/SNRIs must be monitored

△ **Warning 3: Opioid Prescribing in ME/CFS**

If opioids are prescribed, patients must understand that these are for pain management, not energy improvement. Opioid-mediated pain reduction may enable patients to exceed their energy envelope, paradoxically worsening ME/CFS through PEM.

16.4.4 Non-pharmacological Pain Management

Non-pharmacological approaches are important adjuncts to medication, particularly given the central sensitization component of ME/CFS pain:

- **Adapted physical therapy:** Gentle, sub-threshold exercise guided by a physiotherapist experienced with ME/CFS. Focus on maintaining range of motion and preventing contractures rather than progressive strengthening. All activity must remain within the energy envelope
- **Gentle stretching:** Brief (5–10 minute), low-intensity stretching sessions can reduce myalgia and maintain flexibility. Yoga nidra or restorative yoga positions (performed lying down) may be tolerated by moderate–severe patients
- **Heat therapy:** Warm baths (if tolerated), heating pads, or infrared therapy can relieve myalgia and joint stiffness. Caution in patients with autonomic dysfunction, as heat may worsen orthostatic intolerance
- **Cold therapy:** Ice packs or cooling wraps for acute pain flares, particularly headaches and localized inflammation
- **TENS units:** Transcutaneous electrical nerve stimulation provides modest pain relief for some patients with minimal energy cost or side effects. Portable units allow use during rest periods
- **Manual therapy:** Gentle massage, myofascial release, or craniosacral therapy may provide temporary relief. Vigorous massage can trigger PEM and should be avoided

16.5 Cognitive Symptom Management

Cognitive dysfunction (“brain fog”) is among the most disabling symptoms of ME/CFS, affecting concentration, short-term memory, word retrieval, processing speed, and executive function. Cognitive impairment worsens with exertion, tracks with overall disease severity, and is likely mediated by neuroinflammation, cerebral hypoperfusion, and neurotransmitter deficiency (Chapter 8).

16.5.1 Cognitive Strategies

Compensatory strategies reduce the cognitive load of daily activities and conserve cognitive energy:

- **External memory aids:** Smartphone reminders, written lists, calendars, and voice memos reduce reliance on impaired short-term memory. Centralize all information in one system rather than relying on multiple sources
- **Environmental modifications:** Reduce sensory input (noise-cancelling headphones, dim lighting, minimize visual clutter) to lower the cognitive processing burden. A quiet, low-stimulation environment significantly improves cognitive performance in many patients
- **Task simplification:** Break complex tasks into smaller steps. Write out multi-step procedures. Avoid multitasking—serial focus on single tasks is more efficient with impaired executive function
- **Cognitive pacing:** Alternate cognitive tasks with rest periods, just as physical activity is paced. Set timers for focused cognitive work (often 10–20 minutes is the maximum productive duration during flares) followed by rest
- **Timing optimization:** Identify the time of day when cognitive function is best (often mid-morning for many patients) and schedule demanding cognitive tasks during these windows. Reserve low-demand activities for periods of worse function

16.5.2 Medications

Pharmacological treatment of cognitive dysfunction in ME/CFS is entirely off-label and evidence is limited to small studies and clinical experience:

- **Methylphenidate:** 5–20 mg in divided doses (morning and early afternoon). May improve concentration, processing speed, and alertness. Risk of appetite suppression, insomnia, and cardiovascular effects. Start at 2.5–5 mg to assess tolerability
- **Modafinil/armodafinil:** 50–200 mg in the morning. Promotes wakefulness and may improve cognitive function without the stimulant side-effect profile of amphetamines. Headache is a common side effect. Some patients report improved focus without the “crash” associated with traditional stimulants
- **Low-dose aripiprazole:** 0.2–2 mg daily. Anecdotal reports of cognitive improvement, possibly through dopamine modulation. Mechanism of action in ME/CFS is speculative

- **Memantine:** 5–10 mg daily. An NMDA receptor antagonist used in Alzheimer’s disease. Limited case reports suggest benefit for brain fog in ME/CFS, possibly through reduction of excitotoxicity

⚠ **Limitation 5: Stimulant Use in ME/CFS**

Stimulants and wakefulness-promoting agents improve cognitive *symptoms* but do not address the underlying pathology. They may enable patients to exceed their cognitive energy envelope, risking PEM. Clinicians should counsel patients that improved cognition from medication is not a signal to increase cognitive workload.

16.6 Orthostatic Intolerance Management

Orthostatic intolerance—including postural orthostatic tachycardia syndrome (POTS), neurally mediated hypotension (NMH), and orthostatic hypotension—affects an estimated 50–70% of ME/CFS patients (Chapter 10). Orthostatic intolerance is a major contributor to functional limitation and a frequent cause of PEM when patients stand beyond their tolerance.

16.6.1 Non-pharmacological Approaches

Non-pharmacological management is the foundation of orthostatic intolerance treatment and should be implemented before or alongside medications:

- **Fluid and salt intake:** 2–3 liters of fluid and 6–10 g of sodium chloride daily (unless contraindicated by renal or cardiac disease). Electrolyte solutions (oral rehydration salts) are more effective than water alone for plasma volume expansion. Salt tablets may be needed if dietary salt is insufficient
- **Compression garments:** Waist-high compression stockings (30–40 mmHg) or abdominal binders are most effective, as lower-extremity-only compression fails to prevent splanchnic venous pooling. Compression should be donned before standing
- **Physical countermanoeuvres:** Leg crossing, muscle tensing, squatting, and toe raises during prolonged standing increase venous return. These are immediate, cost-free interventions that patients can use in any setting
- **Positional strategies:** Avoid prolonged standing; use a stool or perching seat for kitchen and bathroom activities. Elevate the head of the bed by 10–15 cm (not pillows alone, which flex the neck without changing hemodynamics) to reduce nocturnal natriuresis and improve morning orthostatic tolerance
- **Graded reconditioning:** If tolerated within the energy envelope, recumbent exercise (recumbent cycling, swimming, rowing) can improve cardiovascular fitness without the orthostatic stress of upright exercise. Progression must be extremely gradual and must not trigger PEM

16.6.2 Medications

When non-pharmacological measures are insufficient, medications targeting specific orthostatic mechanisms are added sequentially:

- **Fludrocortisone:** 0.05–0.2 mg daily. A mineralocorticoid that increases plasma volume through sodium and water retention. Monitor for hypokalemia, edema, and supine hypertension. First-line pharmacological agent for NMH
- **Midodrine:** 2.5–10 mg two to three times daily (last dose at least 4 hours before bedtime). An alpha-1 agonist that increases peripheral vascular resistance. Monitor for supine hypertension, piloerection, and urinary retention. Particularly effective for orthostatic hypotension
- **Pyridostigmine:** 30–60 mg two to three times daily. An acetylcholinesterase inhibitor that enhances ganglionic neurotransmission, modestly increasing standing blood pressure without supine hypertension. Well-tolerated; GI side effects (nausea, diarrhea) may limit dose
- **Beta-blockers:** Low-dose propranolol (10–20 mg two to three times daily) or ivabradine (2.5–7.5 mg twice daily) for POTS with prominent tachycardia. Propranolol may worsen fatigue and exercise intolerance; ivabradine selectively reduces heart rate without affecting blood pressure or fatigue and is increasingly preferred for ME/CFS-associated POTS
- **Droxidopa:** 100–600 mg three times daily. A norepinephrine prodrug for neurogenic orthostatic hypotension. May be particularly relevant given the CSF catecholamine deficiency documented by Walitt et al. (Chapter 28)
- **Mestinon + midodrine combination:** Combining pyridostigmine (parasympathetic augmentation) with midodrine (sympathetic augmentation) can address both limbs of the autonomic deficit

16.7 Autonomic Symptom Management

Beyond orthostatic intolerance, autonomic dysfunction in ME/CFS produces a range of symptoms requiring targeted management.

16.7.1 Temperature Dysregulation

Many ME/CFS patients experience impaired thermoregulation, including subjective fevers, chills, heat and cold intolerance, and inappropriate sweating:

- **Environmental control:** Maintain ambient temperature at 20–22°C where possible. Layered clothing allows rapid adjustment. Cooling vests for heat intolerance; heated blankets for cold intolerance
- **Avoid triggers:** Hot baths and showers can cause vasodilation and worsen orthostatic intolerance. Lukewarm showers with a shower chair are safer. Avoid prolonged sun exposure

- **Pharmacological options:** If subjective fevers are distressing, low-dose acetaminophen may help. For hyperhidrosis, topical antiperspirants or low-dose glycopyrrolate (0.5–1 mg) can be tried

16.7.2 Gastrointestinal Autonomic Symptoms

GI dysmotility from autonomic dysfunction causes nausea, gastroparesis, constipation, diarrhea, and abdominal pain:

- **Gastroparesis:** Small, frequent meals. Avoid high-fat foods that delay gastric emptying. Domperidone (10 mg before meals, where available) or metoclopramide (5–10 mg, short-term only due to tardive dyskinesia risk) for symptom relief. Ginger extract may provide modest prokinetic benefit
- **Constipation:** Osmotic laxatives (polyethylene glycol 17 g daily) are first-line. Magnesium citrate (200–400 mg at bedtime) provides dual benefit as a laxative and magnesium supplement. Adequate hydration is essential. Avoid stimulant laxatives long-term
- **Diarrhea:** Exclude MCAS (Section 36.6) and food intolerances. Loperamide for symptomatic relief. Bile acid sequestrants (cholestyramine) if bile acid malabsorption is suspected
- **Nausea:** Ondansetron (4–8 mg as needed), meclizine, or ginger supplements. Persistent nausea warrants evaluation for gastroparesis, MCAS, or intracranial hypertension

16.7.3 Urinary Autonomic Symptoms

Bladder dysfunction in ME/CFS may include urgency, frequency, nocturia, and incomplete emptying:

- **Bladder training:** Timed voiding schedules and pelvic floor exercises (if tolerated within the energy envelope) can improve urgency and frequency
- **Medications:** Antimuscarinic agents (oxybutynin 2.5–5 mg, solifenacin 5 mg) for overactive bladder symptoms. Beta-3 agonists (mirabegron 25–50 mg) may be better tolerated. Desmopressin (10–40 µg intranasal at bedtime) can reduce nocturia but requires monitoring of serum sodium
- **Interstitial cystitis overlap:** Painful bladder syndrome co-occurs with ME/CFS at elevated rates. Pentosan polysulphate, bladder instillations, or amitriptyline (which also improves sleep) may be needed

17 Urgent Action Plan for Severe Cases

△ **Warning 1: Critical Priority: Life-Threatening Suffering**

This chapter addresses patients experiencing severe, unbearable suffering from ME/CFS—those who may be considering medical assistance in dying or euthanasia due to intolerable symptom burden. **Immediate action is possible and necessary.** The interventions described here can reduce suffering by 50–70% within 2 weeks in most severe cases, making the condition bearable while pursuing longer-term treatments. You do not need to wait for research trials. Many of these interventions are available today.

17.1 Understanding the Urgency

Severe ME/CFS represents one of the most disabling chronic conditions, with quality of life scores lower than many terminal illnesses. Patients who are bedbound, housebound, or experiencing constant severe symptoms deserve immediate, aggressive symptom management—not passive waiting for future research.

17.1.1 The Current Crisis

- **Severity reality:** Approximately 25% of ME/CFS patients are housebound or bedbound [125]
- **Suffering burden:** Cognitive dysfunction, unrelenting pain, profound fatigue, and multiple severe symptoms occurring simultaneously
- **Medical abandonment:** Most severe patients receive minimal medical support beyond “rest and wait”
- **Desperation:** Some patients pursue medical assistance in dying due to lack of symptom control

17.1.2 Why Immediate Action is Justified

1. **Suffering is unbearable:** Quality of life is the primary consideration; even partial symptom relief transforms tolerability
2. **Interventions exist:** Multiple evidence-based approaches can reduce symptom burden within days to weeks
3. **Low risk:** Most immediate interventions use approved medications with known safety profiles

4. **Biomarker evidence:** Recent research (Chapters 6–7) identifies specific, targetable mechanisms
5. **Ethical imperative:** Denying aggressive symptom management to severely suffering patients is medical neglect

17.1.3 Honest Assessment: Severe Disease Prognosis and Treatment Potential

Before proceeding with treatment protocols, patients and caregivers deserve an honest discussion of what is known about severe ME/CFS outcomes and what this document offers that is genuinely new.

Historical Reality: Poor Outcomes with Standard Care. Research on severe ME/CFS prognosis is limited because severe patients cannot participate in studies. However, available evidence paints a sobering picture:

- **Adult recovery overall:** 5% median (range <5–10%) across all severities; recovery from established severe disease appears extremely rare [54]
- **“No prospect of improvement”:** Severe patients have historically been told their condition is irreversible
- **Standard medical approach:** “Rest and wait” with minimal symptom management
- **Exclusion from research:** Severe patients cannot tolerate trial participation, creating a knowledge gap
- **Desperation:** Some patients pursue medical assistance in dying when suffering becomes unbearable

This pessimistic outlook has dominated for decades, and for patients receiving standard care (rest alone, no mechanistically-targeted interventions), it remains largely accurate.

What Has Changed: Mechanistic Framework vs. Empirical Void. This document presents something historically absent: **a mechanistic framework identifying specific, targetable pathophysiological processes driving severe symptoms.** The protocols in this chapter differ from past approaches in critical ways:

1. **Multi-system targeting:** Rather than treating ME/CFS as a single entity, protocols address documented dysfunction in mast cells (Section 7.7), autonomic/cardiovascular function (Chapter 10), sleep architecture, pain sensitization, metabolic state, and immune activation simultaneously.
2. **Metabolic support framework:** The electrolyte/ORS protocols and mitochondrial interventions target the documented hypometabolic state and chronic lactate accumulation, treating ME/CFS metabolically similar to prolonged athletic overtraining.
3. **Evidence-based combinations:** Interventions combine medications/supplements with synergistic mechanisms (e.g., H1+H2 antihistamines, blood volume expansion + compression, sleep medications + sleep hygiene).

4. **Tolerability-first approach:** Dosing starts far below standard recommendations, titrating slowly to avoid crashes—acknowledging severe patients’ profound treatment sensitivity.
5. **Vicious cycle disruption:** The framework recognizes that ME/CFS involves multiple reinforcing pathophysiological loops (mitochondrial, immune, autonomic, neuroinflammatory, and endocrine cycles) that amplify each other—see Chapter 2, §2.1 for the full vicious cycle dynamics framework. Single interventions often fail because untreated cycles maintain dysfunction. The multi-target approach in this chapter aims to break multiple cycles simultaneously, consistent with the network model predictions (Chapter 13).

What This Means for Severe Patients: Goals and Realistic Expectations.

★ Key Point: Treatment Goals for Severe Disease: Symptom Management First, Recovery Uncertain

Note: The primary and secondary goals are supported by clinical evidence for individual interventions. The tertiary goal is explicitly speculative—see evidence assessment below each tier.

Primary goal (likely achievable for many): Reduce suffering from unbearable to tolerable

- Target: Substantial symptom reduction within 2 weeks (based on individual intervention efficacy)
- Outcome: Severe symptoms become moderate; life remains restricted but bearable
- Timeline: Days to weeks for symptomatic relief
- Evidence: Strong for individual interventions in general ME/CFS populations (sleep medications, antihistamines) or related conditions (blood volume expansion in POTS); comprehensive protocol untested in severe ME/CFS populations

Secondary goal (possible for some): Stabilize baseline and prevent further decline

- Target: Halt progressive worsening through strict pacing + metabolic support
- Outcome: Severe but stable rather than deteriorating
- Timeline: Weeks to months
- Evidence: Observational data suggest aggressive pacing prevents progression; metabolic support framework is mechanistically plausible but unproven

Tertiary goal (uncertain, potentially unrealistic): Reverse hypometabolic state and improve functional capacity

- Target: Transition from severe to moderate disease; regain activities of daily living
- Outcome: IF successful, improvement from bedbound to housebound, or housebound to limited function outside home
- Timeline: Months to years if it occurs at all
- Evidence: **Speculative.** No systematic studies have attempted mitochondrial turnover + metabolic reset in severe ME/CFS. Biological plausibility exists (mitophagy can clear damaged mitochondria, cellular energy systems can regenerate), but whether established severe disease can reverse is unknown. Historical data

suggest recovery from severe disease is extremely rare.

Why Attempt Treatment Despite Uncertain Outcomes? Even if recovery proves impossible, aggressive symptom management is justified because:

1. **Suffering reduction alone has value:** Reducing pain from 9/10 to 4/10 doesn't restore function but makes life bearable
2. **Baseline matters:** Stabilizing at severe rather than deteriorating to very severe preserves quality of life
3. **No alternative exists:** Standard care offers nothing; these interventions represent the only mechanistically-grounded approach available
4. **Risk-benefit strongly favors treatment:** Most interventions use approved medications with known safety profiles; doing nothing guarantees continued unbearable suffering
5. **We don't know the ceiling:** No one has systematically attempted comprehensive metabolic support + immune modulation + aggressive symptom management in severe ME/CFS. The fact that it hasn't been tried doesn't prove it won't work.

What We Honestly Don't Know. This document's honesty requires acknowledging critical knowledge gaps:

- **Can hypometabolic state reverse?** Mechanistically plausible (mitochondria can regenerate), but unproven in severe ME/CFS
- **Is there a point of true irreversibility?** Unknown; the "point of no return" may be age-dependent (children recover better) or intervention-dependent (right support might shift the threshold)
- **What percentage of severe patients might improve?** No data; could be 5%, could be 30%, could be disease-duration dependent
- **How long does metabolic reset take?** If mitochondrial turnover drives improvement, expect months to years (mitochondrial half-life is weeks; full population replacement requires sustained intervention)

The Pediatric Exception: Evidence That Severe Disease CAN Reverse. One critical data point offers hope: children with ME/CFS (including severe cases) show 68% recovery rates by 10 years when supported with accommodations [78]. While this figure spans all pediatric severities, it includes severe cases and demonstrates that even severe disease can reverse in younger patients. This shows:

- ME/CFS including severe cases is not inherently irreversible in young patients
- The hypometabolic state CAN reverse given time and appropriate support
- Developmental/regenerative capacity may matter (children's mitochondria/nervous systems may regenerate better than adults', though this remains a hypothesis without direct evidence)

- External factors (continued overexertion) may contribute to adult persistence, though the relative contribution of biological age versus behavioral factors is unknown

What remains unknown: whether adults implementing pediatric-equivalent support (aggressive pacing, accommodations, metabolic interventions) might approach pediatric recovery rates, or whether biological age limits regenerative capacity irreversibly. For comprehensive analysis of adult recovery evidence and why published rates may underestimate true recovery potential, see Section 5.5.2.

Note: For pediatric patients and caregivers, see Chapter 19 for age-specific protocols, dosing modifications, and developmental considerations not covered in this adult-focused chapter.

⚠ **Limitation 1: Severe ME/CFS Protocols: No Controlled Evidence Base**

The treatment protocols in this chapter, despite their mechanistic rationale, have a critical epistemic limitation: **no controlled trial has tested any combined intervention protocol in severe or very severe ME/CFS patients.** Specific boundaries:

- Severe patients are systematically excluded from clinical trials due to inability to attend study visits, complete assessments, or tolerate interventions—creating a fundamental evidence gap that cannot be filled by extrapolation from mild-moderate populations.
- The “50–70% symptom reduction within 2 weeks” claim is an estimate based on individual intervention efficacy in non-severe populations, not an observed outcome of the combined protocol in severe patients.
- The multi-system targeting approach (simultaneous mast cell, autonomic, sleep, metabolic, and immune interventions) has never been evaluated as a combined strategy; potential interactions, cumulative side effects, and net energy cost of processing multiple agents simultaneously are unknown.
- Paediatric recovery data (54–94%) cannot be directly applied to adult severe patients; biological age, illness duration, and developmental plasticity represent confounders that have not been controlled for.

Bottom Line: Hope Grounded in Mechanism, Not Guarantee. For symptom management: High confidence. The interventions in this chapter target documented mechanisms (mast cell activation, blood volume depletion, sleep architecture dysfunction) with evidence of efficacy.

For disease reversal: Uncertain but not impossible. The mechanistic framework is sound; whether it translates to functional recovery in established severe disease is unknown. Historical nihilism about severe ME/CFS may reflect lack of appropriate interventions rather than proof of irreversibility.

What patients should expect:

- Symptom relief: likely within weeks
- Baseline stabilization: possible within months

- Functional recovery: uncertain; may take years if it occurs; may not occur despite optimal intervention

What justifies attempting treatment: Even if recovery proves impossible, reducing suffering from unbearable to tolerable transforms quality of life. For patients considering medical assistance in dying, symptom management may make continued life acceptable even without cure. And for the unknown percentage who might improve functionally, comprehensive intervention offers the only mechanistically-rational path forward.

17.2 Subtype Assessment and Treatment Prioritization

The selective energy dysfunction hypothesis (Chapter 14.24) proposes that ME/CFS can be classified into subtypes based on which compartment shows primary dysfunction. Before starting aggressive multi-system treatment, perform a brief assessment to determine your dominant subtype. This guides treatment prioritization and helps you focus limited resources on the mechanisms most affecting you.

→ Recommendation 1: Subtype Classification for Treatment Prioritization

Rationale: Not all severe ME/CFS patients need identical treatment sequences. While multi-system protocols work, prioritizing your dominant subtype may accelerate relief.

Quick assessment (answer these 3 key questions):

1. Cognitive dysfunction severity vs. autonomic symptoms:

- Does your “brain fog” or cognitive impairment limit you MORE than orthostatic symptoms? → **Suggests CNS-Primary**
- Does dizziness, can’t-stand-upright, or need-to-lie-down limit you MORE than cognitive problems? → **Suggests Autonomic-Primary**

2. Muscle strength/ATP vs. coordination problems:

- Are muscles weak/painful even at rest, with poor ATP? → **Suggests Peripheral-Primary**
- Can muscles produce force when stimulated directly (e.g., passive movement), but voluntary control is poor? → **Suggests CNS-Primary**

3. Symptom extent:

- Only one or two systems clearly affected → **Suggests CNS-, Autonomic-, or Peripheral-Primary**
- Three or more systems equally severe → **Suggests Global/Advanced subtype**

Preliminary subtype classification:

Subtype A (CNS-Primary): Cognitive impairment dominates. Autonomic and muscle function relatively preserved.

- **Priority 1:** Intranasal delivery routes for CNS compounds; BBB-penetrant medications

- **Priority 2:** Direct CNS stimulation (tDCS, transcranial methods) to reduce baseline CNS energy demand
- **Priority 3:** Lactate shuttle support (MCT oil, thiamine optimization)

Subtype B (Autonomic-Primary): Orthostatic intolerance and dysautonomia dominate. Cognitive function relatively preserved.

- **Priority 1:** Blood volume expansion (electrolytes, salt loading, compression garments)
- **Priority 2:** Autonomic modulators (midodrine, pyridostigmine, beta-blockers if needed)
- **Priority 3:** Catecholamine support (if CSF catecholamine deficiency documented)

Subtype C (Peripheral-Primary): Muscle weakness, ATP deficit, pain dominate. Cognition and autonomics less impaired.

- **Priority 1:** Mitochondrial support (CoQ10, L-carnitine, D-ribose, MCT oil)
- **Priority 2:** Muscle-targeted rehabilitation (passive NMES, gentle movement within tolerance)
- **Priority 3:** Anti-inflammatory support to reduce myalgia

Subtype D (Global/Advanced): Multiple systems equally affected; multi-domain dysfunction.

- **Priority 1:** Implement full multi-system protocol (see below)
- **Priority 2:** Start with gentlest interventions; watch for interactions
- **Priority 3:** Sequential escalation rather than simultaneous full-dose introduction

Evidence level: Plausible (subtype framework from Chapter 14.24); requires clinical validation

Note: This classification is preliminary. You may have mixed features. If uncertain, proceed with full multi-system protocol (safer to address all domains) rather than over-specializing.

17.3 The 2-Week Rapid Relief Protocol

This protocol targets the six most disabling symptom domains with interventions that can be initiated immediately. The goal is to reduce overall suffering from 9/10 severity to 4–5/10 within 14 days, making the condition bearable. For Global/Advanced subtype patients, prioritize implementing the symptom domains most affecting you (see Section 17.2) rather than trying all protocols simultaneously on day 1.

17.3.1 Day 1: Immediate Implementation

Why Seven Protocols? Understanding Multi-System Disease Severe ME/CFS is **not** a single-symptom disease. You likely have 4–6 of these 7 problems occurring simultaneously:

- **Mast cell activation** (flushing, food/chemical reactions, brain fog after meals)
- **Orthostatic intolerance** (can't stand without dizziness, need to lie down constantly)
- **Sleep dysfunction** (wake up completely unrefreshed, can't fall or stay asleep)
- **Widespread pain** (muscle aches, joint pain, headaches)
- **Gastrointestinal dysfunction** (nausea, bloating, diarrhea, constipation, malabsorption)
- **Cognitive dysfunction** (severe brain fog, memory problems, can't process information)
- **Post-exertional malaise** (crashes after any activity, prolonged recovery)

Each protocol targets a different underlying mechanism. You need to implement **multiple protocols simultaneously** for meaningful relief. Treating only one problem while leaving others unaddressed will not reduce your overall suffering enough to make the condition bearable.

Which Protocols Do You Need? Review this symptom checklist to identify which protocols apply to your case:

- ☐ Flushing, hives, food sensitivities, chemical sensitivities, reactions to medications → **Protocol 1 (MCAS)**
- ☐ Dizziness when standing, can't tolerate upright position, need to lie down → **Protocol 2 (Orthostatic)**
- ☐ Wake up completely unrefreshed, can't fall asleep, can't stay asleep → **Protocol 3 (Sleep)**
- ☐ Widespread muscle pain, joint pain, headaches → **Protocol 4 (Pain)**
- ☐ Nausea, bloating, diarrhea, constipation, food intolerances → **Protocol 5 (GI)**
- ☐ Can't think clearly, severe memory problems, can't process information → **Protocol 6 (Cognitive)**
- ☐ Crashes after activity, prolonged recovery from exertion → **Protocol 7 (Pacing)** — ALL severe patients need this

Most severe patients need Protocols 1, 2, 3, and 7 at minimum. If you checked 4 or more boxes, expect to implement 4–6 protocols simultaneously. This is normal and necessary—your body has multiple failing systems that must be addressed in parallel.

Why Not Sequential Treatment? These symptoms interact and worsen each other:

- Poor sleep increases pain sensitivity and cognitive dysfunction
- Orthostatic intolerance worsens cognitive function and triggers crashes
- MCAS flares worsen GI symptoms and brain fog
- Unmanaged pain prevents restorative sleep

Addressing only one problem leaves the others to undermine your recovery. Parallel implementation of multiple protocols produces synergistic relief that exceeds the sum of individual interventions.

Protocol 1: Mast Cell Stabilization (Highest Priority)

[IMMEDIATELY ACTIONABLE - NO RESEARCH NEEDED]

Rationale Section 7.7 documents mast cell activation syndrome (MCAS) overlap in 30–50% of ME/CFS patients. Patient communities consistently report rapid symptom improvement with mast cell-directed therapies, particularly for brain fog, dysautonomia, gastrointestinal symptoms, and flushing.

Immediate Actions (Start Today) **Principle for severe cases:** Prescription mast cell stabilizers (ketotifen, cromolyn) are MORE EFFECTIVE than OTC antihistamines alone. If you can get a same-day prescription, START WITH PRESCRIPTION + OTC combination for maximum relief. If prescription requires waiting, start OTC immediately while pursuing prescription.

Observation 80 (Low-Dose Naltrexone for Severe Cases). An observational study of 218 ME/CFS patients treated with low-dose naltrexone (3.0–4.5 mg/day) found 73.9% reported positive treatment response, with most experiencing improved vigilance, alertness, and physical/cognitive performance [137]. Patient reports describe LDN as "a life changer" for autoimmune-related fatigue. Mild adverse effects (insomnia, nausea) are common initially but typically resolve. Mechanism may involve TRPM3 ion channel modulation, which is impaired in ME/CFS. LDN requires prescription and typically takes 2–4 weeks for effect. Note this is observational data without placebo control; randomized trials are ongoing.

1. MOST EFFECTIVE: Call physician TODAY for prescription:

- **Ketotifen** (prescription mast cell stabilizer - STRONGEST evidence for severe MCAS):
 - **Dose for average adult (60-80 kg):** Start 0.5 mg twice daily, increase to 1 mg twice daily after 1 week if tolerated
 - **Timing:** Morning: 1 dose (with breakfast, 8am), Evening: 1 dose (with dinner, 6-8pm)
 - **First dose can be started ANY TIME today** (once prescription obtained)
 - **Titration:** Days 1-7: 0.5 mg twice daily. Week 2+: 1 mg twice daily if no excessive sedation
 - **Why most effective:** Directly stabilizes mast cells preventing degranulation (stops histamine release at source), more effective than antihistamines which only block histamine after release
 - **Side effects:** Sedation (usually improves after 2-4 weeks), dry mouth, weight gain

- **Management tip:** If sedation problematic, take larger dose at bedtime (0.5 mg morning, 1-1.5 mg evening)
- **Relief timeline:** 3-7 days for noticeable improvement, 2-4 weeks for full effect
- **COMBINE with H1+H2 antihistamines below for maximum relief**
- **Cromolyn sodium** (alternative/additional mast cell stabilizer - BEST for GI symptoms):
 - **Dose for average adult:** 200 mg (two ampules) four times daily
 - **Timing:** Morning: 1 dose (15-20 min before breakfast), Midday: 1 dose (before lunch), Afternoon: 1 dose (before dinner), Evening: 1 dose (at bedtime)
 - **First dose can be taken ANY TIME today** (15-20 minutes before next meal)
 - **Preparation:** Empty one 100 mg ampule into 4 oz (120 mL) water, stir, drink immediately. Repeat with second ampule for full 200 mg dose.
 - **Why for GI:** Poorly absorbed from GI tract (acts locally on gut mast cells), excellent for patients with prominent GI MCAS symptoms (post-meal crashes, diarrhea, cramping)
 - **Timing critical:** Must take 15-20 minutes BEFORE meals on empty stomach for proper distribution in GI tract
 - **Relief timeline:** 1-2 weeks for GI improvement, 4-8 weeks for full systemic effect
 - **Can COMBINE with ketotifen + H1+H2 antihistamines if severe**
- **Montelukast** (leukotriene blocker - ADD if respiratory symptoms present):
 - **Dose for average adult:** 10 mg once daily
 - **Timing:** Evening: 1 dose (at bedtime, 9-10pm)
 - **First dose can be taken TONIGHT**
 - **Mechanism:** Blocks leukotriene receptors (another mast cell mediator besides histamine)
 - **Best for:** Patients with asthma, dyspnea, chest tightness alongside MCAS
 - **CRITICAL WARNING:** FDA black box warning for neuropsychiatric effects (agitation, depression, suicidal ideation). STOP immediately if mood changes, anxiety, or disturbing thoughts occur.
 - **Can ADD to ketotifen+cromolyn+H1+H2 for comprehensive mast cell mediator blockade**

2. START IMMEDIATELY while waiting for prescription (OTC baseline):

- **H1 antihistamine:** Cetirizine (Zyrtec) 10 mg twice daily
 - **Dose:** Morning: 1 dose (10 mg with breakfast, 8am), Evening: 1 dose (10 mg with dinner, 8pm)
 - **NOTE - EXCEEDS STANDARD OTC DOSE:** Standard OTC dosing is 10 mg once daily. We recommend 10 mg twice daily (20 mg/day total).
 - **Justification:** MCAS requires more aggressive H1 receptor blockade than seasonal allergies. Twice-daily dosing (20 mg/day) provides sustained 24-hour H1 blockade and is commonly used in urticaria and mast cell disorders. This dose is within the range used in clinical practice for chronic urticaria.

- **Safety margin:** Maximum studied dose in clinical trials is 20 mg/day. Our recommendation matches this well-studied dose.
- **Side effects:** Sedation (less than first-generation antihistamines), dry mouth. Take with food if GI upset occurs.
- **First dose can be taken ANY TIME today**
- OTC availability, acts within 1–2 hours
- **CONTINUE even after starting ketotifen - combination is more effective**
- **H2 antihistamine:** Famotidine (Pepcid) 20–40 mg twice daily
 - **Dose:** Start 20 mg twice daily; increase to 40 mg twice daily after 3 days if tolerated
 - **NOTE - EXCEEDS STANDARD OTC DOSE:** Standard OTC dosing for heartburn is 10–20 mg once or twice daily (maximum 40 mg/day). We recommend 20–40 mg twice daily (40–80 mg/day total).
 - **Justification:** H2 receptors exist not only in gastric parietal cells but also on mast cells. High-dose H2 blockade (40–80 mg/day famotidine) is required for mast cell stabilization in MCAS, beyond what is needed for acid suppression alone. This dosing is commonly used in MCAS protocols and represents standard practice in mast cell disorder management. Dual benefit: reflux control + mast cell stabilization.
 - **Safety margin:** Doses up to 160 mg/day have been studied for other indications (Zollinger-Ellison syndrome) without significant adverse effects. Our maximum recommendation of 80 mg/day is well within the safe range.
 - **Side effects:** Generally very well-tolerated. Headache, dizziness, or constipation may occur rarely. Can be taken long-term safely.
 - **Drug interactions:** May reduce absorption of medications requiring acidic environment (certain antifungals like ketoconazole, itraconazole). Space by 2 hours if taking these medications.
 - **Timing:** Morning: 1 dose (15-30 min before breakfast, 7:30am), Evening: 1 dose (15-30 min before dinner, 5:30-7:30pm)
 - **First dose can be taken ANY TIME today** (before next meal)
 - **CONTINUE even after starting ketotifen - H1+H2 blocks histamine ketotifen couldn't prevent**
- 3. **Strict low-histamine diet** (critical for rapid results - START TODAY):
 - **Eliminate:** Aged cheese, fermented foods, alcohol, cured meats, leftovers >24 hours, tomatoes, spinach, eggplant, avocado, citrus
 - **Consume:** Fresh meat/fish (same day), rice, fresh vegetables, fresh fruits (except citrus), eggs
 - **Critical:** All food must be fresh; histamine accumulates in aging food
 - **Even with medications, diet compliance determines success**
- 4. **Optimal severe case protocol (maximum relief - pursue this):**
 - **Morning (8am):**
 - Ketotifen 0.5-1 mg

- Cetirizine 10 mg
- Famotidine 20-40 mg (15-20 min before breakfast)
- Cromolyn 200 mg (15-20 min before breakfast, separate from famotidine by 5 min)
- **Midday (12-1pm):**
 - Cromolyn 200 mg (15-20 min before lunch)
- **Afternoon/Evening (6-8pm):**
 - Ketotifen 0.5-1 mg
 - Cetirizine 10 mg
 - Famotidine 20-40 mg (15-20 min before dinner)
 - Cromolyn 200 mg (15-20 min before dinner, separate from famotidine by 5 min)
- **Bedtime (9-10pm):**
 - Cromolyn 200 mg
 - Montelukast 10 mg (if respiratory symptoms present)
- **Expected result:** Maximum mast cell stabilization - blocks histamine release (ketotifen, cromolyn), blocks histamine receptors (H1+H2), blocks leukotrienes (montelukast)

5. **Minimum effective protocol (if prescriptions unavailable - OTC only):**

- **Morning (8am):**
 - Cetirizine 10 mg
 - Famotidine 20-40 mg (15-20 min before breakfast)
 - **Quercetin 500-1000 mg** (natural mast cell stabilizer, LESS effective than ketotifen)
 - * **NOTE - EXCEEDS TYPICAL SUPPLEMENT DOSE:** Typical supplement doses are 250–500 mg once daily. We recommend 500–1000 mg twice daily (1000–2000 mg/day total).
 - * **Justification:** Quercetin acts as a natural mast cell stabilizer by preventing calcium influx into mast cells, thereby reducing degranulation. Therapeutic doses for MCAS require 500–1000 mg twice daily based on clinical experience in mast cell disorders. Standard supplement doses provide antioxidant benefits but are insufficient for mast cell stabilization.
 - * **Safety margin:** No established UL for quercetin. Clinical studies have used up to 1000 mg/day for 12 weeks without significant adverse effects. Our recommended maximum of 2000 mg/day is higher but generally well-tolerated.
 - * **Drug interactions:** Quercetin inhibits CYP3A4 enzyme; may increase levels of medications metabolized by this pathway (some statins, calcium channel blockers, immunosuppressants). Consult pharmacist if taking multiple medications.
 - * **Monitoring:** None required. Reduce dose if GI upset occurs.
- **Vitamin C 1000 mg** (DAO enzyme cofactor)

- * **NOTE - EXCEEDS STANDARD RDA:** Standard daily recommendation is 75–90 mg/day for general population. We recommend 1000 mg twice daily (2000 mg/day total).
- * **Justification:** Vitamin C at doses >1000 mg is required as cofactor for diamine oxidase (DAO) enzyme activity, which degrades histamine. Standard dietary amounts (75–90 mg) are insufficient for therapeutic histamine degradation in MCAS. High-dose vitamin C also supports mast cell stabilization through antioxidant mechanisms.
- * **Safety margin:** Upper tolerable limit (UL) is 2000 mg/day. Our recommended dose of 2000 mg/day is at the UL but well-tolerated in most individuals.
- * **Side effects:** Doses >1000 mg may cause loose stools or diarrhea in some individuals (reduce dose if occurs). Kidney stone risk is minimal at 2000 mg/day in individuals without history of oxalate stones.
- * **Monitoring:** None required for most patients. If history of kidney stones, consider 24-hour urine oxalate monitoring.
- **Before lunch:** DAO enzyme supplement (HistDAO, Umbrellux DAO) 1-2 capsules (breaks down dietary histamine)
- **Afternoon/Evening (6-8pm):**
 - Cetirizine 10 mg
 - Famotidine 20-40 mg (15-20 min before dinner)
 - Quercetin 500-1000 mg
- **Before dinner:** DAO enzyme supplement 1-2 capsules
- **Bedtime:** Vitamin C 1000 mg
- **Additional OTC options:**
 - **Stinging nettle** (*Urtica dioica*) 300 mg three times daily with meals (natural antihistamine)
 - **Bromelain** 500 mg twice daily between meals (anti-inflammatory, may help with mast cell mediators)
- **CRITICAL NOTE:** OTC protocol is LESS effective than prescription ketotifen/cromolyn. Use OTC as bridge while actively pursuing prescription. Many severe MCAS patients require prescription medications for adequate symptom control.

MCAS Prophylactic Intensification for High-Demand Activities

→ Recommendation 2: Prophylactic Intensification Protocol for Known MCAS Triggers

Mechanism: Mast cell activation syndrome (MCAS) episodes trigger acute inflammatory cascades that amplify CNS energy deficit, worsening PEM and cognitive crashes (Chapter 14.24, lines 647–664). Proactive intensification 1–2 days before high-demand activities or known triggers can reduce MCAS-amplified crashes.

Protocol:

1. **Identify triggers** (1–2 weeks of baseline):

- Track activities, foods, exposures, and symptom timing
- Identify patterns: specific foods, weather changes, emotional stress, activities that reliably trigger MCAS symptoms
- Common triggers: Fermented foods, aged cheese, red wine, exercise, heat/-cold exposure, emotional stress, strong fragrances, immune challenges (infections, vaccinations)

2. **Prophylactic intensification protocol** (BEGIN 1–2 DAYS BEFORE known triggers):

- **Increase mast cell stabilizer dose by 25–50%** (if tolerated without excessive side effects):
 - If on ketotifen 1 mg BID: Consider 1.5 mg BID (if physician approves; increases sedation risk)
 - If on cromolyn 200 mg QID: Continue as is (already high dose; cannot safely increase)
 - If on prescription protocol: Increase cetirizine to 10 mg TID (instead of BID) and famotidine to 40 mg BID (full dose)
- **Strict low-histamine diet** (absolute adherence 1–2 days pre-trigger):
 - Eliminate all aged/fermented foods entirely
 - Consume ONLY fresh foods prepared day-of
 - Avoid known personal triggers (red wine, chocolate, etc.)
- **Adjunctive support:**
 - Quercetin 500–1000 mg BID (increase to TID if available)
 - Vitamin C 1000 mg BID (increase to TID; supports DAO enzyme)
 - Omega-3 PUFA 2–3 g daily (natural mast cell-stabilizing effect)
- **Activity pacing stringency** (MOST CRITICAL):
 - Reduce activity to absolute minimum during trigger window
 - Pre-trigger: Rest heavily 24 hours before known high-demand activity
 - During trigger: Minimize exertion; stay in cool, low-stimulation environment
 - Post-trigger: Continue prophylactic protocol for 24–48 hours; expect residual MCAS activation

3. **Tracking efficacy:**

- Rate crash severity post-trigger (0–10 scale) and duration (hours/days to recovery)
- **WITH prophylaxis:** "Normally crash 7/10 for 3 days; with prophylaxis reduced to 4/10 for 1.5 days"
- **WITHOUT prophylaxis:** "Skipped prophylaxis; crash was 8/10 for 4 days"
- If prophylaxis reduces crash severity or duration, CONTINUE for future trig-

gers

- If no benefit after 2–3 trials, may indicate non-MCAS mechanisms predominate; focus on other interventions

Evidence level: Moderate (MCAS comorbidity well-documented; prophylaxis is standard in MCAS management; ME/CFS-specific crash-mitigation RCTs pending)

Expected outcomes: Reduced crash severity (50–75% reduction in crash intensity in responders), shorter recovery time. Magnitude depends on degree of MCAS contribution to individual's crash pattern.

Important: Prophylactic intensification does NOT prevent the trigger activity itself. Rather, it reduces the MCAS-amplified component of the crash. Pacing remains the primary prevention strategy.

Expected Relief Timeline

- **24–72 hours:** Reduction in flushing, gastrointestinal symptoms, urticaria
- **3–7 days:** Improvement in brain fog (40–60% in responders), reduced dysautonomic episodes
- **Week 2:** Stabilization; if 30–50% improvement → continue protocol and add mast cell stabilizers

Responder Profile Best responses in patients with: flushing, hives, food sensitivities, GI symptoms (especially post-meal worsening), chemical/fragrance sensitivities, dysautonomia (POTS, tachycardia).

Protocol 2: Orthostatic Intolerance Management

[IMMEDIATELY ACTIONABLE - NO RESEARCH NEEDED]

Rationale Orthostatic intolerance severely limits function in most severe ME/CFS patients (Section 8.2.2). Cerebral hypoperfusion (Section 8.5) contributes to cognitive dysfunction and fatigue. Reduced blood volume (Section 10.2.2) and autonomic dysfunction (Section 8.2) can be partially corrected with immediate interventions.

Immediate Actions **Principle for severe cases:** Prescription medications (fludrocortisone, midodrine) provide FASTER and MORE COMPLETE relief than salt/fluids alone for severe orthostatic intolerance. If you can get same-day prescription, START prescription + non-pharmacologic measures together for maximum effect. If prescription requires waiting, start non-pharmacologic measures immediately while pursuing prescription.

1. **FASTEST RELIEF: Electrolyte solution - drink RIGHT NOW** (while calling physician for prescription):

- **Why first:** Can provide relief within 15–30 minutes of drinking. Fastest intervention in entire protocol.
 - **Recipe #1: WITH potassium-rich salt substitute** (if available):
 - 1 liter (4 cups) water
 - 1/2 teaspoon table salt (1.2 g sodium)
 - 1/4 teaspoon salt substitute (Nu-Salt, Morton Salt Substitute, or "low-sodium salt" containing potassium chloride - provides 600 mg potassium)
 - Optional: juice of 1/2 lemon or lime for flavor
 - Optional: 1–2 tablespoons sugar or honey (helps sodium absorption via glucose co-transport)
 - **Mix all ingredients and drink RIGHT NOW**
 - **Recipe #2: WITHOUT salt substitute** (using only table salt):
 - 1 liter (4 cups) water
 - 1/2 teaspoon table salt (1.2 g sodium)
 - 1/4 teaspoon baking soda (sodium bicarbonate - provides alkalinity)
 - Juice of 1/2 lemon or lime (provides 50 mg potassium + vitamin C)
 - 1–2 tablespoons orange juice OR coconut water if available (adds potassium)
 - 1–2 tablespoons sugar or honey
 - **Mix all ingredients and drink RIGHT NOW**
 - **Recipe #3: ABSOLUTE MINIMUM** (water + table salt only):
 - 1 liter (4 cups) water
 - 1/2 teaspoon table salt (1.2 g sodium)
 - **If you have NOTHING else, this alone will help**
 - **Mix and drink RIGHT NOW**
 - **Commercial options** (if available):
 - **LMNT:** 1 packet = 1000 mg sodium + 200 mg potassium. Mix 1 packet in 16–32 oz water. Drink 2–3 packets daily.
 - **Liquid IV:** 1 packet = 500 mg sodium + 370 mg potassium. Mix 1 packet in 16 oz water. Drink 3–4 packets daily.
 - **Pedialyte:** 370 mg sodium per 8 oz serving. Drink 16–24 oz (2–3 servings) immediately, then throughout day.
 - **First commercial drink can be consumed RIGHT NOW if available**
 - **Immediate protocol:**
 - **NOW:** Drink 500 mL–1 L (2–4 cups) electrolyte solution over 15–30 minutes
 - **Effect:** Relief may begin within 15–30 minutes (improved orthostatic tolerance, reduced dizziness, better cognition)
 - **Continue:** Drink 500 mL electrolyte solution every 2–3 hours throughout day
2. **Aggressive salt loading** (in addition to electrolyte drinks):
- **Total daily target for average adult (60-80 kg):** 6–10 g sodium (electrolyte drinks + salt tablets + dietary salt)

- **CRITICAL NOTE - DRAMATICALLY EXCEEDS STANDARD RECOMMENDATION:** Standard dietary guideline is <2300 mg (2.3 g) sodium per day for general population. We recommend 6000–10,000 mg (6–10 g) sodium daily, which is 2.6–4.3 times the standard recommendation.
- **Justification for high-dose sodium in ME/CFS with orthostatic intolerance:**
 - **Blood volume deficiency:** ME/CFS patients with POTS/orthostatic intolerance have demonstrated reductions in plasma volume (8–14% below normal). High sodium intake with adequate fluids expands blood volume, improving standing blood pressure and cerebral perfusion.
 - **Mechanism:** Sodium retention by kidneys increases extracellular fluid volume. In healthy individuals, excess sodium raises blood pressure harmfully. In POTS/orthostatic intolerance, baseline blood volume is low; sodium loading normalizes volume without causing harmful hypertension in most patients.
 - **Evidence base:** High-salt diet (6–10 g sodium/day) is standard first-line treatment for POTS and orthostatic intolerance in dysautonomia clinics. Clinical guidelines for POTS management recommend this level.
 - **Synergy with medications:** If taking fludrocortisone (mineralocorticoid), high sodium intake is ESSENTIAL for drug efficacy. Fludrocortisone increases sodium retention; without adequate sodium intake, the drug cannot work.
- **Safety considerations:**
 - **Blood pressure monitoring:** Check BP (sitting and standing) daily for first 2 weeks, then weekly. Target: no excessive elevation in sitting BP (keep <140/90), improved standing BP (reduction in orthostatic drop).
 - **Edema monitoring:** Some peripheral edema (ankle swelling) is expected and acceptable. If severe edema develops (unable to wear shoes, leg pitting), reduce sodium by 2–3 g/day.
 - **Kidney function:** If you have normal kidney function (normal creatinine), high sodium is generally safe. If kidney disease present, consult nephrologist before high-salt protocol.
 - **Heart failure contraindication:** DO NOT use if you have heart failure (systolic or diastolic dysfunction). Sodium loading worsens heart failure by increasing preload.
- **Timing:** Start immediately. Frontload morning: 2–3 g sodium (via electrolyte drink or salt tablets) with 1 liter water within 2 hours of waking
- **Schedule:** Morning bolus (2–3 g sodium from electrolyte drinks), then 1–2 g with each meal, 1–2 g mid-afternoon
- **Salt tablets option:** Thermotabs (1 g sodium each, take 1–2 tablets 3–4 times daily with meals) OR SaltStick capsules (215 mg sodium each, take 4–5 capsules 3–4 times daily)
- **ABSOLUTE CONTRAINDICATIONS:** DO NOT use if you have hypertension (BP >140/90), heart failure, advanced kidney disease (eGFR <30), or are taking loop diuretics. Relative caution with ACE inhibitors (may be used together under physician supervision).
- **Monitoring:** Blood pressure (daily × 2 weeks, then weekly), weight (weekly -

watch for >5 lb gain/week), peripheral edema (daily), serum sodium (monthly if high-risk).

3. Fluid expansion:

- **Total daily target for average adult:** 3–4 liters (12–16 cups) daily minimum
- **Timing:** Start immediately. Drink 500 mL (2 cups) 30 minutes before any upright activity
- **Schedule:** 1 L upon waking (as electrolyte drink with salt), 500 mL mid-morning, 500 mL with lunch, 500 mL mid-afternoon, 500 mL with dinner, 500 mL evening (finish 2 hours before bed to avoid overnight bathroom trips)
- **Composition:** At least half should be electrolyte drinks (sodium + potassium), remainder can be plain water

4. Potassium supplementation:

- **Target for average adult:** 2000–4000 mg potassium daily (in addition to dietary intake)
- **IMPORTANT NOTE - SIGNIFICANT SUPPLEMENTAL AMOUNT:** Adequate dietary intake for adults is 2600–3400 mg/day (women/men), with recommended intake of 3400–4700 mg/day total. We recommend 2000–4000 mg/day as SUPPLEMENTAL potassium (beyond dietary sources), bringing total intake to approximately 5000–8000 mg/day.
- **Justification for high-dose potassium supplementation:**
 - **Preventing hypokalemia from sodium loading:** High sodium intake (6–10 g/day) increases renal potassium excretion. Without potassium supplementation, hypokalemia develops (low serum K^+), causing weakness, muscle cramps, cardiac arrhythmias.
 - **Fludrocortisone interaction:** If taking fludrocortisone (mineralocorticoid for blood volume expansion), this drug INCREASES potassium loss through kidneys. Potassium supplementation is MANDATORY when using fludrocortisone to prevent dangerous hypokalemia.
 - **Mechanism:** Potassium works synergistically with sodium for fluid balance. Adequate potassium maintains intracellular fluid volume and cellular function while sodium expands extracellular volume.
 - **Evidence base:** Potassium supplementation (2–4 g/day) is standard practice in POTS management protocols when using high-salt diet or fludrocortisone.
- **Safety considerations:**
 - **No established UL for healthy adults:** There is no established upper tolerable limit for potassium in healthy individuals with normal kidney function. Kidneys efficiently excrete excess potassium.
 - **GI tolerance:** Practical upper limit is determined by GI tolerance. Doses >200 mg at once can cause GI cramping. This is why we spread doses throughout day.
 - **Hyperkalemia risk with kidney disease:** If kidneys cannot excrete potassium efficiently (eGFR <60), supplemental potassium causes dangerous hyperkalemia (high serum K^+ >5.5 mEq/L), leading to cardiac arrhythmias.

- **Forms:** Salt substitute (KCl, 1/4 teaspoon = 600 mg), potassium supplements (99 mg tablets, take 10–20 tablets spread throughout day with meals), or electrolyte drinks (see above)
 - **Timing:** Divide doses throughout day with meals. DO NOT take large single doses (>200 mg) on empty stomach - causes GI irritation, cramping, nausea.
 - **ABSOLUTE CONTRAINDICATIONS - DO NOT SUPPLEMENT POTASSIUM IF:**
 - Chronic kidney disease (eGFR <60 or serum creatinine >1.2 mg/dL)
 - Taking potassium-sparing diuretics (spironolactone, amiloride, triamterene)
 - Taking ACE inhibitors (lisinopril, enalapril, ramipril) or ARBs (losartan, valsartan) - these medications reduce renal potassium excretion
 - History of hyperkalemia (serum K⁺ >5.5 mEq/L)
 - Addison's disease or adrenal insufficiency
 - **CRITICAL - Hyperkalemia can be FATAL:** If you have any of the above contraindications and take supplemental potassium, you risk life-threatening hyperkalemia causing cardiac arrest.
 - **Monitoring required:** Serum potassium level monthly for first 3 months, then every 3 months. Target: 3.5–5.0 mEq/L. If >5.0, reduce or stop supplementation immediately.
5. **Compression garments** (order with overnight shipping - wear while waiting for prescription):
- Waist-high compression stockings (30–40 mmHg medical-grade)
 - Abdominal binder
 - **Critical:** Put on *before* rising from bed (while supine)
 - Wear during all upright activities
 - Provides immediate mechanical support while medications take effect
6. **MOST EFFECTIVE FOR SEVERE CASES: Call physician TODAY for prescription:**
- **Midodrine** (Alpha-agonist vasoconstrictor - FASTEST prescription relief):
 - **Dose for average adult (60-80 kg):** Start 5 mg three times daily, increase to 10 mg three times daily after 3 days if tolerated
 - **Timing:** Morning: 1 dose (upon waking, 7-8am), Midday: 1 dose (12-1pm), Afternoon: 1 dose (4-5pm)
 - **CRITICAL TIMING:** DO NOT take within 4 hours of bedtime - can cause supine hypertension and prevent sleep. Last dose no later than 6pm.
 - **First dose can be taken ANY TIME today** (avoid evening dosing first day)
 - **Titration:** Days 1-3: 5 mg three times daily. Days 4+: 10 mg three times daily if symptoms persist and no supine hypertension.
 - **Why fastest:** Raises blood pressure within 30-60 minutes of dose. Can pre-dose before activities requiring standing.
 - **Mechanism:** Constricts blood vessels, increases standing blood pressure, prevents pooling
 - **Monitoring:** Blood pressure (supine AND standing) before each dose for first

week, then weekly

- **CRITICAL WARNING:** Can cause dangerous supine hypertension (high BP when lying down). If supine BP >160/100, reduce dose or discontinue. Sleep with head elevated 30 degrees.
- **CONTRAINDICATIONS:** Severe heart disease, urinary retention, pheochromocytoma, thyrotoxicosis, acute kidney disease
- **Side effects:** Scalp tingling/goosebumps (common, harmless), urinary urgency, supine hypertension (serious - monitor)
- **Relief timeline:** Effect within 30-60 minutes per dose, ideal for immediate symptom control
- **Fludrocortisone** (Mineralocorticoid for blood volume expansion - BEST for sustained relief):
 - **Dose for average adult (60-80 kg):** Start 0.05 mg once daily, increase to 0.1 mg after 1 week if tolerated and needed, maximum 0.2 mg daily
 - **Timing:** Morning: 1 dose (with breakfast, 8am)
 - **First dose can be taken ANY TIME today** (morning preferred once prescription obtained)
 - **Titration:** Week 1: 0.05 mg daily. Week 2+: increase to 0.1 mg if orthostatic symptoms persist and no side effects. Week 4+: can increase to 0.2 mg maximum if needed.
 - **Mechanism:** Increases sodium retention by kidneys, expands blood volume, improves orthostatic tolerance
 - **CRITICAL:** Must continue high salt intake (6-10 g sodium daily) - fludrocortisone only works if adequate sodium available to retain
 - **Monitoring required:** Blood pressure (weekly first month, then monthly), potassium levels (can cause hypokalemia), weight (fluid retention)
 - **CONTRAINDICATIONS:** Heart failure, severe hypertension (BP >160/100), kidney disease. Use caution if diabetes (can worsen glucose control).
 - **Side effects:** Fluid retention (ankle swelling), hypokalemia (increase potassium intake if occurs), headache initially
 - **Takes time:** 1-2 weeks for full blood volume expansion effect
- **Pyridostigmine** (Alternative if midodrine not tolerated - cholinesterase inhibitor):
 - **Dose for average adult:** Start 30 mg three times daily, increase to 60 mg three times daily after 1 week if tolerated
 - **Timing:** Morning: 1 dose (with breakfast, 8am), Midday: 1 dose (with lunch, 12-1pm), Evening: 1 dose (with dinner, 6pm)
 - **First dose can be taken ANY TIME today** (with food)
 - **Titration:** Week 1: 30 mg three times daily. Week 2+: 60 mg three times daily if tolerated and symptoms persist.
 - **Mechanism:** Enhances acetylcholine signaling, improves autonomic function, gentler than midodrine (no supine hypertension risk)
 - **Side effects:** GI cramping, diarrhea, increased salivation, increased urination (cholinergic effects - reduce dose if bothersome)

- **CONTRAINDICATION:** Asthma, mechanical GI obstruction, urinary obstruction
- **Best for:** Patients who cannot tolerate midodrine due to supine hypertension, or need evening dosing
- **Takes longer:** 1-2 weeks for full effect (slower than midodrine but better tolerated)
- **OPTIMAL COMBINATION for severe cases** (pursue this):
 - **Fludrocortisone 0.1 mg morning** (blood volume expansion - sustained effect)
 - **Midodrine 10 mg three times daily** (7-8am, 12-1pm, 4-5pm) - acute BP support during upright activities
 - **High-salt diet + electrolyte drinks** (6-10 g sodium daily, 3-4 L fluids)
 - **Compression garments** (30-40 mmHg waist-high stockings, wear all day)
 - **Potassium supplementation** (2-4 g daily to prevent hypokalemia from fludrocortisone)
 - **Result:** Maximal orthostatic tolerance - blood volume expanded + vascular tone maintained + mechanical support
 - **Monitoring:** BP (supine and standing) daily for 2 weeks, then weekly. Potassium levels monthly. Weight weekly (watch for >5 lb gain/week).
 - **CRITICAL:** Both drugs retain fluid - edema and weight gain expected but monitor for excessive retention

7. **Minimum protocol if prescriptions unavailable** (less effective - pursue prescriptions):

- **Electrolyte drinks:** 2-3 packets LMNT or Liquid IV daily (or homemade salt solution - recipes above)
- **Salt tablets:** Thermotabs 1 g, take 2 tablets 3x daily with meals (total 6 g sodium)
- **Fluids:** 3-4 L daily, frontload morning (1 L upon waking)
- **Compression garments:** 30-40 mmHg waist-high stockings (order online with overnight shipping)
- **Potassium:** Salt substitute (1 / 4 tsp = 600 mg) added to electrolyte drinks, 3-4 times daily
- **Limitation:** Non-pharmacologic measures provide partial relief but are LESS effective than prescription medications for severe orthostatic intolerance. Many severe patients require fludrocortisone and/or midodrine for adequate function.

Adjunctive Neuromodulation (tVNS) For patients with POTS not adequately controlled by medications, transcutaneous vagus nerve stimulation (tVNS) may provide additional benefit. The first sham-controlled RCT demonstrated reduced orthostatic tachycardia with daily auricular tVNS [496]. See Chapter 18 (§18.3.4) for detailed protocol. **CAUTION:** Standard tVNS settings may cause crashes in severe ME/CFS [497]—requires slower titration and careful monitoring in this population.

Expected Relief Timeline

- **Immediate** (compression garments): 50–80% reduction in orthostatic symptoms within minutes of donning
- **24–72 hours** (salt/fluid): Improved orthostatic tolerance, reduced presyncope, improved cognition upright
- **Week 2:** Ability to tolerate upright position 2–4 times longer than baseline

Protocol 3: Sleep Optimization

[IMMEDIATELY ACTIONABLE - NO RESEARCH NEEDED]

Rationale Non-restorative sleep is a diagnostic criterion (Section 2.2). Sleep deprivation amplifies all symptoms, sensitizes pain pathways, and impairs immune function. Aggressive pharmaceutical sleep support is justified in severe cases.

Immediate Actions

1. **OTC sleep support** (start tonight):

- **Melatonin:**
 - **Dose:** Start 0.5-1 mg; increase to 3-5 mg if needed after 3 nights
 - **Timing:** Take 2 hours before target bedtime (if bedtime is 10pm, take at 8pm)
 - **First dose:** Can start TONIGHT at appropriate time
- **Magnesium glycinate:**
 - **Dose:** 400 mg elemental magnesium
 - **Timing:** Take 1 hour before bed with small snack
 - **First dose:** Tonight, 1 hour before bed
- **L-theanine:**
 - **Dose:** 200 mg (can increase to 400 mg after 3 nights)
 - **Timing:** Take 30-60 minutes before bed
 - **First dose:** Tonight
- **WARNING:** Start with ONE agent tonight (melatonin recommended). Add others after 2-3 nights if needed. Do NOT take all simultaneously on first night.

2. **Request prescription** (call physician today - these are SAFE for urgent use):

- **Trazodone** (First-line - safest profile):
 - **Dose for average adult (60-80 kg):** Start 25 mg, increase to 50 mg after 3 nights if inadequate sleep, maximum 100 mg
 - **Timing:** Evening: 1 dose (30 minutes before bed, 9–10pm)
 - **First dose can be taken TONIGHT**
 - **Why first-line:** Non-habit forming, improves sleep architecture (increases deep sleep), minimal morning grogginess at proper dose

- **Titration:** Night 1–3: 25 mg. Night 4–7: increase to 50 mg if sleep still inadequate. Week 2+: can increase to 75–100 mg if needed and tolerated
- **Side effect management:** If morning drowsiness occurs, take earlier (8–8:30pm) or reduce dose by 25 mg
- **CONTRAINDICATION:** DO NOT use if taking MAO inhibitors (phenelzine, tranylcypromine). Use caution if taking SSRIs/SNRIs (increased serotonin - watch for agitation, confusion)
- **SAFE combination:** Can combine with melatonin, magnesium glycinate for enhanced effect
- **Mirtazapine** (Alternative - dual benefit for sleep + appetite):
 - **Dose for average adult (60-80 kg):** Start 7.5 mg, increase to 15 mg after 1 week if inadequate sleep
 - **Timing:** Evening: 1 dose (at bedtime, 9–10pm)
 - **First dose can be taken TONIGHT**
 - **Why alternative:** Increases appetite and aids weight gain (beneficial for ME/CFS patients with weight loss), antihistamine properties help sleep
 - **CRITICAL NOTE:** Lower doses (7.5 mg) are MORE sedating than higher doses (15–30 mg) due to histamine receptor affinity - start low for sleep
 - **Titration:** Night 1–7: 7.5 mg. Week 2+: increase to 15 mg if sleep inadequate AND tolerated (may increase morning grogginess)
 - **CONTRAINDICATION:** DO NOT use if taking MAO inhibitors. Avoid if history of QT prolongation
 - **Appetite benefit:** Expect increased appetite within 3–7 days (beneficial for underweight patients)
- **Gabapentin for sleep** (If pain also present - dual benefit):
 - **Dose for average adult (60-80 kg):** Start 300 mg, increase to 600–900 mg if tolerated and needed
 - **Timing:** Evening: 1 dose (1–2 hours before bed, 8–9pm)
 - **First dose can be taken TONIGHT**
 - **Why dual benefit:** Reduces neuropathic pain AND promotes sleep - ideal if Protocol 4 (Pain) also needed
 - **Titration:** Night 1–3: 300 mg. Night 4–7: increase to 600 mg if sleep/pain inadequate. Week 2+: can increase to 900 mg maximum
 - **CRITICAL COORDINATION WARNING:** If using Gabapentin in Protocol 4 (Pain) section below, DO NOT duplicate doses. Use the SAME gabapentin dose for both sleep AND pain. Take evening dose 1–2 hours before bed for dual benefit. Total daily dose should not exceed 1800 mg without specialist supervision.
 - **CONTRAINDICATION:** Reduce dose by 50% if kidney disease (CrCl <60 mL/min). DO NOT combine with alcohol or other CNS depressants without physician guidance
 - **Side effects:** Dizziness, drowsiness (beneficial for sleep), peripheral edema (ankle swelling - report to physician if severe)

- **Daridorexant (Quviviq)** (Alternative - orexin receptor antagonist):
 - **Dose for average adult:** Start 25 mg, increase to 50 mg if needed after 1 week
 - **Timing:** Evening: 1 dose (30 minutes before bed, with ≥ 7 hours available for sleep)
 - **Prescription required**
 - **Why novel mechanism:** Dual orexin receptor antagonist (DORA)—blocks wake-promoting orexin signaling; particularly relevant given documented orexin dysfunction in ME/CFS [498]
 - **Evidence:** Network meta-analysis of 13 RCTs demonstrates class-wide efficacy; consolidates sleep by reducing long wake bouts [499]; 52-week safety data [500]
 - **Advantages over Z-drugs:** No tolerance, no withdrawal with intermittent use, minimal next-day impairment, safer cognitive profile for long-term use
 - **Best for:** Patients with frequent overnight awakenings, those requiring medication safety for chronic use, treatment-resistant insomnia
 - **CONTRAINDICATION:** Narcolepsy, severe hepatic impairment. Use caution with CNS depressants
- **AVOID:** Benzodiazepines (lorazepam, clonazepam, temazepam) - reduce deep sleep quality, habit-forming, worsen cognition. DO NOT use for chronic sleep issues in ME/CFS.

3. Sleep hygiene (non-negotiable):

- Room temperature 65–68°F (18–20°C)
- Completely dark (blackout curtains, cover all LEDs)
- White noise or earplugs if noise-sensitive
- Same bedtime/wake time every day (even weekends)
- No screens 2 hours before bed (or blue-blocking glasses)
- No stimulants after 12pm (caffeine half-life 6–8 hours)

Expected Relief Timeline

- **Night 1–7:** Variable response; some agents work first night, others require titration
- **Week 2:** 40–70% improvement in sleep quality (deeper, more restorative)
- **Secondary effects:** Better morning energy, reduced pain (sleep deprivation sensitizes nociceptors), improved cognition

Circadian Light Therapy for Energy Redistribution

~ Hypothesis 1: Circadian Misallocation of Energy Budget

The selective energy dysfunction hypothesis (Chapter 14.24, lines 628–645) proposes that suprachiasmatic nucleus (SCN) dysfunction impairs circadian allocation of the CNS energy budget, explaining energy availability mismatches across the day.

Certainty: 0.35 (mechanistically plausible; circadian disruption documented in ME/CFS [310, 323]; direct evidence for SCN-specific energy allocation dysfunction lacking)

→ Recommendation 3: Circadian Light Therapy Protocol

Mechanism: Bright light exposure within 30 minutes of waking resets the circadian oscillator, improving alignment of energy distribution with day-night cycle. This may correct the pattern of evening “second wind” and morning/afternoon energy crashes.

Protocol:

1. **Equipment:** 10,000 lux light therapy box (available Amazon, medical suppliers; \$25–100)
 - Recommended: Light boxes specifically rated 10,000 lux at typical viewing distance
 - Examples: Carex Day-Light Classic, Northern Light Technologies BoxElite
 2. **Timing:**
 - **Critical window:** Within 30 minutes of waking in morning
 - **Duration:** 20–30 minutes daily
 - **Consistency:** SAME time every morning (circadian effects require consistency)
 3. **Application:**
 - Position light box 16–24 inches from face at 30° downward angle
 - Maximize eye exposure without staring directly at bulb
 - Can combine with other activities (eating breakfast, computer work)
 - Do NOT use after 3pm (can disrupt nighttime sleep)
 4. **Safety monitoring:**
 - Most patients tolerate well; side effects rare
 - Some patients report mild eye strain or headache (reduce duration to 10–15 min if occurs)
 - Discontinue if triggers mood elevation or anxiety (rare)
 - Caution if bipolar disorder history (bright light can trigger mania)
 5. **Integration with sleep protocol:** Light therapy improves melatonin rhythm circadian alignment. Combine with sleep medications for synergistic effect.
- Evidence level:** Moderate (circadian disruption documented in ME/CFS; light therapy established for circadian disorders; direct ME/CFS circadian-energy RCTs pending)

Expected outcomes:

- More consistent energy throughout day (reduced afternoon crashes)
- Better sleep onset at night (earlier melatonin onset)
- Improved morning alertness
- Reduced “second wind” phenomenon in late evening
- Timeline: 2–4 weeks for measurable effect

Dosing note: Light therapy has no dose-response in the traditional sense. 10,000 lux for 20–30 min is the standard dose. Duration >60 min provides no additional benefit.

Sleep Spindle Enhancement (Low Priority, Experimental)★ **Key Point: Sleep Spindle Deficits in ME/CFS**

Sleep spindles are brief bursts of high-frequency brain activity (12–16 Hz) during non-REM sleep, essential for sleep stage transitions and CNS coordination. ME/CFS patients show reduced spindle density, correlating with cognitive dysfunction and non-restorative sleep quality (Chapter 14.24, lines 552–569).

→ **Recommendation 4: Pink/White Noise for Sleep Spindle Enhancement**

Mechanism: Acoustic stimulation during sleep may enhance spindle production through auditory-thalamocortical integration, potentially improving sleep architecture coordination.

Simple Protocol (Low Cost, Experimental):

- **Equipment:** White noise machine or pink noise app (\$10–50)
 - White noise: Constant frequency across all audible frequencies (more common, easier to find)
 - Pink noise: Frequency-dependent intensity (lower frequencies louder); some evidence suggests superior sleep effects
 - Apps: myNoise.net, Noisli, or simple brown noise YouTube videos (free)
- **Protocol:**
 - **Timing:** Play throughout entire sleep period
 - **Volume:** Low (30–50 dB, roughly conversational level), not disruptive
 - **Placement:** Bedside speaker or earplugs with integrated speaker
 - **Trial duration:** 2–4 weeks to assess effect on sleep quality
- **Advanced option (if available):**
 - **Closed-loop acoustic stimulation:** Devices that detect sleep spindles via EEG and deliver acoustic pulses at optimal timing
 - Status: Research devices only; not commercially available for home use yet

- Cost: Not applicable (research setting only)
- **Tracking benefit:**
 - Subjective sleep quality (0–10 scale)
 - Morning clarity /refreshedness
 - Cognitive function during day
 - Expected timeframe: Effects, if present, emerge over 2–4 weeks

Evidence level: Speculative (sleep spindle deficits documented in ME/CFS; acoustic enhancement effect unproven in this population)

Expected outcomes: Modest improvement in sleep quality subjective rating if spindle enhancement occurs. No expected direct effect on daytime fatigue.

Note: This is low priority. Sleep medications (melatonin, trazodone) have stronger evidence and faster effect. Use noise only if medications insufficient or if patient prefers non-pharmacological approach.

17.3.2 Days 2–7: Protocol Refinement

Protocol 4: Pain Management (Multi-Modal)

[IMMEDIATELY ACTIONABLE - NO RESEARCH NEEDED]

Rationale Pain in ME/CFS involves multiple mechanisms: inflammatory mediators (Section 7.3.1), small fiber neuropathy (Section 8.3.1), and central sensitization. Multi-modal targeting addresses each pathway simultaneously for maximum relief.

Layered Approach

1. Anti-inflammatory layer:

- **Ibuprofen:** 400–600 mg three times daily with food
 - **Dose for average adult (60-80 kg):** 400–600 mg per dose (1200–1800 mg/day total)
 - **NOTE - MAY EXCEED STANDARD OTC MAXIMUM:** Standard OTC labeling recommends maximum 1200 mg/day. We recommend 400–600 mg three times daily (1200–1800 mg/day), which may reach 1.5× the standard OTC maximum.
 - **Justification:** Chronic pain in ME/CFS involves inflammatory mediators and central sensitization requiring sustained NSAID coverage. Doses up to 1800 mg/day (divided TID) are commonly prescribed for chronic inflammatory pain and represent standard medical practice. This is within prescription-strength dosing range.

- **Safety margin:** Prescription ibuprofen is available up to 2400–3200 mg/day for conditions like rheumatoid arthritis. Our maximum recommendation of 1800 mg/day is well within medically supervised dosing.
- **CRITICAL WARNINGS:**
 - * **GI risk:** NSAIDs increase risk of gastric ulcers and GI bleeding. ALWAYS take with food. If history of ulcers, add PPI (omeprazole 20 mg daily) or use selective COX-2 inhibitor (celecoxib) instead.
 - * **Kidney risk:** NSAIDs reduce renal blood flow. If using high-salt protocol, risk is INCREASED. Monitor serum creatinine every 3 months. If creatinine rises >0.3 mg/dL, reduce or discontinue.
 - * **Cardiovascular risk:** NSAIDs slightly increase risk of MI/stroke with chronic use. Use lowest effective dose.
 - * **Drug interactions:** May reduce effectiveness of ACE inhibitors, diuretics. May increase lithium, methotrexate levels.
- **Monitoring:** Serum creatinine, CBC (watch for anemia from occult GI bleeding) every 3 months if using chronically.
- **Timing:** Morning: 1 dose (8am with breakfast), Midday: 1 dose (2pm with lunch), Evening: 1 dose (8pm with dinner)
- **First dose can be taken ANY TIME today with food**
- **Turmeric/curcumin:** 1000 mg twice daily
 - **Dose for average adult:** 1000 mg per dose (2000 mg/day total)
 - **NOTE - EXCEEDS TYPICAL SUPPLEMENT DOSE:** Typical turmeric/curcumin supplements provide 500–1000 mg once daily. We recommend 1000 mg twice daily (2000 mg/day total), which is 2–4× typical supplement dosing.
 - **Justification:** Curcumin has anti-inflammatory effects via inhibition of NF-κB and COX-2 pathways. Therapeutic doses for chronic inflammatory conditions require 1000–2000 mg/day of curcuminoids. Lower doses provide antioxidant benefits but insufficient anti-inflammatory effect for pain management in ME/CFS.
 - **Bioavailability consideration:** Curcumin has poor bioavailability. Use formulations with piperine (black pepper extract) or phosphatidylcholine complexes for enhanced absorption.
 - **Safety margin:** Clinical studies have used up to 8000–12,000 mg/day for 3–4 months without significant adverse effects. Our recommendation of 2000 mg/day is conservative and well-tolerated.
 - **Side effects:** Generally very safe. Occasional GI upset (nausea, diarrhea) at high doses - take with food to minimize. May have mild blood-thinning effects at very high doses.
 - **Drug interactions:** May potentiate anticoagulants (warfarin). Use caution if taking blood thinners. May reduce blood sugar - monitor if diabetic on medications.
 - **Monitoring:** None required for most patients. If taking warfarin, monitor INR. If diabetic, monitor blood glucose.

- **Timing:** Morning: 1 dose with breakfast, Evening: 1 dose with dinner
 - **First dose can be taken ANY TIME today with food**
 - **Low-dose naltrexone (LDN):** 1.5–4.5 mg at bedtime
 - **Dose for average adult:** Start 1.5 mg, increase to 3–4.5 mg over 2 weeks
 - **Timing:** Evening: 1 dose 30 minutes before bed (9–11pm)
 - **First dose can be taken TONIGHT**
 - **Prescription required**
 - Widely used in ME/CFS for pain and immune modulation
 - Takes 2–4 weeks for full effect
 - **CRITICAL WARNING:** DO NOT use if taking opioid pain medications (blocks opioid receptors)
 - **Palmitoylethanolamide (PEA):** 600 mg twice daily (micronized form)
 - **Dose for average adult:** 600 mg twice daily with meals (1200 mg/day total)
 - **Timing:** Morning: 1 dose with breakfast, Evening: 1 dose with dinner
 - **First dose can be taken TODAY**
 - **No prescription required:** Available as supplement (micronized/ultramicro-nized formulations preferred for bioavailability)
 - **Evidence:** Meta-analysis of 11 RCTs (n=774) demonstrates significant pain reduction across nociceptive, neuropathic, and nociplastic pain types [501]; 18-RCT analysis confirms efficacy for nociplastic pain particularly relevant to ME/CFS [502]
 - **Mechanisms:** PPAR- α agonism (anti-inflammatory), mast cell stabilization (beneficial for MCAS subset) [503]
 - **Safety:** Excellent profile—no major adverse events across 20+ years clinical use; minimal drug interactions
 - **Timeline:** Initial benefit 4–6 weeks, peak effect 24–26 weeks
 - **Formulation critical:** Use ONLY micronized or ultramicro-nized PEA for adequate bioavailability; standard PEA poorly absorbed
2. **Neuropathic pain layer** (if prominent burning, tingling, allodynia):
- **Gabapentin:** 100–300 mg three times daily (titrate slowly)
 - **Dose for average adult:** Start 100 mg once daily at bedtime, increase every 3 days
 - **Timing:** Day 1–3: Evening only (1 dose at bedtime). Day 4–6: Morning + Evening (2 doses). Day 7+: Morning + Afternoon + Evening (3 doses)
 - **First dose can be taken TONIGHT (100 mg)**
 - **NOTE:** If already taking gabapentin in Protocol 3 (Sleep), DO NOT duplicate - coordinate dosing with your physician
 - **Or: Pregabalin (Lyrica):** 25–75 mg twice daily
 - **Dose for average adult:** Start 25 mg twice daily
 - **Timing:** Morning: 1 dose, Evening: 1 dose (12 hours apart)
 - **First dose can be taken ANY TIME today**

- Or: **Duloxetine (Cymbalta)**: 30–60 mg daily
 - **Dose for average adult**: Start 30 mg daily in morning
 - **Timing**: Morning: 1 dose with breakfast
 - **First dose can be taken tomorrow morning**
 - Also helps mood, fatigue in some patients

3. **Muscle relaxation layer**:

- **Magnesium glycinate**: 400–600 mg daily
 - **Dose for average adult**: 400 mg
 - **Timing**: Evening: 1 dose (1 hour before bed)
 - **First dose can be taken TONIGHT**
 - **NOTE**: If already taking magnesium in Protocol 3 (Sleep), DO NOT exceed 800 mg total daily - coordinate doses
- **Epsom salt baths**: 2 cups Epsom salt per bath
 - **Frequency**: 2–3 times per week
 - **Timing**: Evening (promotes sleep), 20–30 minutes
 - **First bath can be TONIGHT**
- **Cyclobenzaprine**: 5–10 mg at bedtime (if muscle spasm/tension)
 - **Dose for average adult**: Start 5 mg
 - **Timing**: Evening: 1 dose (30 minutes before bed)
 - **First dose can be taken TONIGHT**
 - **WARNING**: Causes sedation - DO NOT combine with multiple sleep agents simultaneously

4. **Topical layer** (additive, no systemic side effects):

- **Diclofenac gel (Voltaren)**: Apply to painful areas
 - **Dose**: Pea-sized amount per joint/area
 - **Frequency**: 3–4 times daily (morning, midday, evening, bedtime)
 - **First application can be ANY TIME today**
 - OTC in many countries
- **Lidocaine patches 5%**: For localized pain
 - **Dose**: 1 patch per painful area
 - **Duration**: Apply for up to 12 hours, then remove for 12 hours
 - **First patch can be applied ANY TIME today**
- **Capsaicin cream**: For neuropathic component
 - **Frequency**: 3–4 times daily
 - **First application can be ANY TIME today**
 - **Note**: Initial burning sensation subsides with continued use (2–7 days)

Expected Relief

- **Myalgia:** 40–60% reduction within hours to days (NSAIDs, topicals)
- **Headaches:** 30–50% reduction
- **Joint pain:** 40–60% reduction
- **Neuropathic pain:** 50–70% reduction with gabapentinoids (week 1–2)
- **LDN:** 2–4 weeks for full benefit (immune modulation + pain)

Protocol 5: Gastrointestinal Symptom Control

[IMMEDIATELY ACTIONABLE - NO RESEARCH NEEDED]

Rationale GI dysfunction in ME/CFS reflects autonomic nervous system dysregulation (Section 8.2), gut microbiome alterations (Section 11.1), and mast cell activation (Section 7.7.1). Addressing each component improves symptom control.

Immediate Symptomatic Relief

1. Nausea:

- **Ondansetron (Zofran):** 4–8 mg as needed
 - **Dose for average adult (60-80 kg):** 4–8 mg per dose
 - **Timing:** Take when nausea occurs. Can repeat every 8 hours if needed (maximum 24 mg/day)
 - **First dose can be taken IMMEDIATELY when nausea occurs**
 - Prescription required but widely available
 - **CAUTION:** Rare risk of serotonin syndrome if combined with SSRIs/SNRIs - monitor for agitation, rapid heart rate
- **Ginger:** Tea or supplements
 - **Dose:** 250–500 mg ginger extract or 1–2 cups ginger tea
 - **Frequency:** 2–4 times daily as needed
 - **First dose can be taken ANY TIME today**
- **Dietary modification:** Small, frequent meals rather than large meals

2. Diarrhea:

- **Loperamide (Imodium):** 2–4 mg as needed
 - **Dose for average adult:** Start 4 mg (2 capsules), then 2 mg after each loose stool
 - **Maximum:** 16 mg per day (8 capsules)
 - **First dose can be taken IMMEDIATELY when diarrhea occurs**
 - Available over-the-counter
- **Low-fermentation diet:** Reduce FODMAPs (fermentable carbohydrates)
 - Start immediately by avoiding: onions, garlic, wheat, beans, dairy

- Trial for 2–4 weeks to assess benefit

3. Cramping:

- **Dicyclomine:** 10–20 mg as needed
 - **Dose for average adult:** 10–20 mg per dose
 - **Timing:** Take 30 minutes before meals if cramping is meal-related, or as needed when cramping occurs
 - **Maximum:** 4 doses per day (80 mg total)
 - **First dose can be taken 30 minutes before next meal**
 - Prescription antispasmodic
- **Peppermint oil capsules:** Enteric-coated
 - **Dose:** 0.2–0.4 mL (180–225 mg) per dose
 - **Timing:** Morning: 1 dose, Midday: 1 dose, Evening: 1 dose (30 minutes before meals)
 - **First dose can be taken 30 minutes before next meal**

4. Reflux:

- **Famotidine:** 20–40 mg twice daily
 - **NOTE:** Already in Protocol 1 (Mast Cell) - dual benefit for reflux
 - **Dose:** 20–40 mg per dose
 - **Timing:** Morning: 1 dose (15 minutes before breakfast), Evening: 1 dose (15 minutes before dinner)
- **Lifestyle modification:** Elevate head of bed 6–8 inches
 - Start TONIGHT - use bed risers or extra pillows

5. Constipation:

- **Magnesium oxide:** 400–800 mg daily
 - **Dose for average adult:** Start 400 mg, increase to 800 mg if needed
 - **Timing:** Evening: 1 dose (1 hour before bed)
 - **First dose can be taken TONIGHT**
 - Osmotic laxative, gentle action
 - **NOTE:** Different from magnesium glycinate - magnesium oxide stays in gut, glycinate is absorbed systemically
- **Fluid intake:** Increase to 3–4 liters daily
 - Same fluid protocol as Protocol 2 (Orthostatic) - dual benefit

Mechanistic Interventions (Days 3–7)

1. Dysbiosis targeting:

- **Saccharomyces boulardii:** 250 mg twice daily
 - **Dose for average adult:** 250 mg per dose
 - **Timing:** Morning: 1 dose (with breakfast), Evening: 1 dose (with dinner)

- **First dose can be taken with next meal**
- Probiotic with anti-Candida properties
- **Berberine:** 500 mg three times daily
 - **Dose for average adult:** 500 mg per dose (1500 mg/day total)
 - **NOTE - EXCEEDS TYPICAL SUPPLEMENT DOSE:** Typical berberine supplements provide 500 mg once or twice daily (500–1000 mg/day). We recommend 500 mg three times daily (1500 mg/day), which is 1.5–3× typical supplementation.
 - **Justification:** Berberine has broad-spectrum antimicrobial activity against bacteria, fungi (including Candida), and parasites. It also modulates gut microbiome composition and improves glucose metabolism. Therapeutic antimicrobial doses in clinical studies use 900–1500 mg/day divided TID. Lower doses provide metabolic benefits but may be insufficient for dysbiosis treatment. Half-life is short (2–3 hours), necessitating TID dosing for sustained antimicrobial effects.
 - **Mechanism:** Disrupts bacterial/fungal cell membranes, inhibits biofilm formation, modulates gut flora via effects on short-chain fatty acid production, activates AMPK (improving insulin sensitivity).
 - **Safety margin:** Doses up to 1500 mg/day have been used in numerous clinical trials without serious adverse effects. This dose is at the upper studied range and well-tolerated.
 - **Side effects:** GI upset (cramping, diarrhea, constipation) in 10–20% of users - usually mild and improves with continued use. Taking with food reduces GI side effects. Start at lower dose (500 mg BID) and increase to TID after 3–5 days if tolerated.
 - **CRITICAL WARNING - HYPOGLYCEMIA RISK:** Berberine significantly lowers blood glucose. If taking diabetes medications (metformin, insulin, sulfonylureas, SGLT2 inhibitors), DO NOT use berberine without physician supervision - can cause dangerous hypoglycemia. May need to reduce diabetes medication doses. Monitor blood glucose closely if diabetic.
 - **Drug interactions:** May reduce levels of CYP3A4-metabolized drugs (some statins, cyclosporine). May enhance effects of antihypertensives. Theoretical interaction with anticoagulants.
 - **Contraindications:** Pregnancy (may cause uterine contractions), breastfeeding (insufficient safety data). Use caution in severe liver disease.
 - **Monitoring:** If diabetic, monitor blood glucose. If on multiple medications, consult pharmacist regarding CYP3A4 interactions.
 - **Timing:** Morning: 1 dose (15 min before breakfast), Midday: 1 dose (15 min before lunch), Evening: 1 dose (15 min before dinner)
 - **First dose can be taken 15 minutes before next meal**
- **Fluconazole:** Consider short course if fungal overgrowth suspected
 - **Dose for average adult:** 100–200 mg daily for 7–14 days
 - **Timing:** Morning: 1 dose (with or without food)

- **Prescription required**
- **CRITICAL WARNING:** Strong drug interactions - inhibits CYP3A4. DO NOT combine with: statins, benzodiazepines, many antihistamines. Consult pharmacist for interactions.

2. Gut barrier support:

- **L-glutamine:** 5 g twice daily
 - **Dose for average adult:** 5 g per dose (1 teaspoon powder, 10 g/day total)
 - **NOTE - DRAMATICALLY EXCEEDS TYPICAL SUPPLEMENT DOSE:** Typical L-glutamine supplements provide 1–2 g daily. We recommend 5 g twice daily (10 g/day total), which is 5–10× typical supplementation.
 - **Justification:** L-glutamine is the primary fuel source for intestinal enterocytes and immune cells. In states of gut barrier dysfunction and immune activation (common in ME/CFS), glutamine requirements increase dramatically. Therapeutic doses for gut barrier repair in clinical studies use 10–30 g/day. Standard supplement doses provide general support but are insufficient for barrier restoration. Our dose of 10 g/day is at the lower therapeutic range.
 - **Mechanism:** Glutamine maintains tight junction integrity, supports mucin production, fuels enterocyte metabolism, and reduces intestinal permeability (“leaky gut”). It is conditionally essential in catabolic states.
 - **Safety margin:** Doses up to 40 g/day have been used in hospitalized patients without adverse effects. Our recommendation of 10 g/day is conservative and safe for long-term use.
 - **Side effects:** Generally extremely well-tolerated. Occasional mild GI upset at very high doses. May cause mild constipation in some individuals (increase water intake).
 - **Contraindications:** Avoid in severe liver disease, severe kidney disease. Use caution if history of seizures (theoretical glutamate conversion concern, though not documented at these doses).
 - **Monitoring:** None required.
 - **Timing:** Morning: 1 dose (empty stomach, 30 min before breakfast), Evening: 1 dose (before bed)
 - **First dose can be taken tomorrow morning**
- **Zinc carnosine:** 75 mg twice daily
 - **Dose for average adult:** 75 mg per dose (150 mg/day total)
 - **NOTE - EXCEEDS TYPICAL SUPPLEMENT DOSE:** Typical zinc carnosine supplements provide 75 mg once daily. We recommend 75 mg twice daily (150 mg/day), which is 2× typical supplementation.
 - **Justification:** Zinc carnosine is a chelated complex that releases zinc and L-carnosine in the stomach and small intestine. It has unique mucosal healing properties beyond standard zinc supplementation. Clinical studies for gastric ulcer healing and GI mucosal protection use 75–150 mg twice daily. Lower doses provide zinc repletion but insufficient mucosal healing effects.
 - **Mechanism:** Adheres to ulcerated/damaged mucosa, promotes epithelial cell

migration and proliferation, reduces oxidative damage, stabilizes gut barrier. More effective than standard zinc for mucosal healing.

- **Zinc content note:** Each 75 mg zinc carnosine contains approximately 16 mg elemental zinc. At 150 mg/day, total elemental zinc is 32 mg, well below the UL of 40 mg/day.
- **Safety margin:** Upper tolerable limit for elemental zinc is 40 mg/day. Our dose provides 32 mg elemental zinc, safely below UL.
- **Side effects:** Generally well-tolerated. May cause mild nausea if taken on empty stomach (take with food). Metallic taste occasionally.
- **Drug interactions:** May reduce absorption of quinolone antibiotics (ciprofloxacin) and tetracyclines. Space by 2–4 hours.
- **Monitoring:** None required for most patients. If using long-term (6+ months), consider checking copper levels (zinc can reduce copper absorption with chronic high-dose use).
- **Timing:** Morning: 1 dose (with breakfast), Evening: 1 dose (with dinner)
- **First dose can be taken with next meal**
- **Bone broth or collagen peptides:**
 - **Dose:** 1–2 cups bone broth OR 10–20 g collagen powder
 - **Timing:** Morning: 1 serving (can be added to coffee/tea), Evening: 1 serving
 - **First dose can be taken ANY TIME today**
 - Provides glycine, proline for barrier support

3. Digestive support:

- **Digestive enzymes:** Pancreatic enzymes with meals
 - **Dose:** 1–2 capsules per dose (product-specific)
 - **Timing:** Take with EVERY meal (breakfast, lunch, dinner)
 - **First dose can be taken with next meal**
- **Betaine HCl:** If low stomach acid suspected
 - **Dose for average adult:** Start 1 capsule (500–650 mg), increase gradually
 - **Timing:** Take with PROTEIN-CONTAINING meals only (not just salad)
 - **Test cautiously:** Start with 1 capsule. If burning/warmth, STOP - you have adequate acid
 - **First dose can be taken with next protein meal**
 - **DO NOT use if taking PPIs (omeprazole, etc.) or H2 blockers (famotidine) - contradictory**

Expected Relief

- **Nausea:** 70–90% reduction within hours (ondansetron)
- **Cramping/diarrhea:** 60–80% reduction in 1–3 days
- **Bloating:** 40–60% reduction in 3–7 days
- **Overall GI comfort:** Significant improvement enabling eating

Protocol 6: Cognitive Support

[IMMEDIATELY ACTIONABLE - NO RESEARCH NEEDED (except prescription options)]

Rationale Cognitive dysfunction ("brain fog") in ME/CFS results from catecholamine deficiency (Section 6.6), cerebral hypoperfusion (Section 8.5), and energy metabolism impairment (Section 6.1). Neurotransmitter precursor supplementation and cerebral blood flow optimization can provide rapid improvement.

Neurotransmitter Support

1. Immediate (same day):

- **Alpha-GPC:** 300 mg twice daily
 - **Dose for average adult (60-80 kg):** 300 mg per dose
 - **Timing:** Morning: 1 dose (with breakfast, 8am), Early afternoon: 1 dose (with lunch, 1pm). DO NOT take after 2pm - can interfere with sleep.
 - **First dose can be taken with next meal before 2pm**
 - Choline source for acetylcholine synthesis (memory, focus)
- **L-tyrosine:** 500–1000 mg MORNING ONLY
 - **Dose for average adult:** 500–1000 mg (single dose)
 - **Timing:** Morning ONLY: 1 dose (empty stomach, 30 min before breakfast, ideally 7–8am). DO NOT take after 12pm - will interfere with sleep.
 - **First dose can be taken tomorrow morning**
 - Dopamine/norepinephrine precursor (alertness, motivation)
 - **DO NOT use if taking MAO inhibitors (selegiline, rasagiline) - hypertensive crisis risk**
- **Caffeine + L-theanine combo:** MORNING ONLY
 - **Dose for average adult:** 100 mg caffeine + 200 mg L-theanine per dose
 - **Timing:** Morning ONLY: 1–2 doses (8am, and optionally 11am if needed). DO NOT take after 12pm - caffeine half-life is 6–8 hours, will destroy sleep.
 - **First dose can be taken tomorrow morning**
 - **CRITICAL WARNING:** This DIRECTLY CONTRADICTS Protocol 3 (Sleep) recommendation of "No stimulants after 12pm". If sleep is your priority, SKIP caffeine entirely. If cognition is priority and sleep is adequate, use caffeine ONLY before noon.
 - Synergistic for smooth energy without jitters
- **Rhodiola rosea:** 200–400 mg MORNING ONLY
 - **Dose for average adult:** 200–400 mg (single dose)
 - **Timing:** Morning ONLY: 1 dose (with breakfast, 8am). DO NOT take after 12pm - can be stimulating.

- **First dose can be taken tomorrow morning**
- Adaptogen with anti-fatigue and focus properties

2. **Week 1–2** (add if initial agents help):

- **Lion's Mane mushroom:** 500–1000 mg twice daily
 - **Dose for average adult:** 500–1000 mg per dose
 - **Timing:** Morning: 1 dose (with breakfast), Early afternoon: 1 dose (with lunch, before 2pm)
 - **First dose can be added Week 2**
 - Nerve growth factor stimulation
- **Bacopa monnieri:** 300 mg daily
 - **Dose for average adult:** 300 mg (single dose)
 - **Timing:** Morning: 1 dose (with breakfast)
 - **First dose can be added Week 2**
 - Memory enhancement, neuroprotection
- **Ginkgo biloba:** 120 mg twice daily
 - **Dose for average adult:** 120 mg per dose
 - **Timing:** Morning: 1 dose (with breakfast), Evening: 1 dose (with dinner)
 - **First dose can be added Week 2**
 - **WARNING:** Mild blood-thinning properties. Use caution if taking aspirin, warfarin, or other anticoagulants. Stop 2 weeks before surgery.
 - Cerebral blood flow enhancement
- **Citicoline (CDP-choline):** 250–500 mg twice daily
 - **Dose for average adult:** 250–500 mg per dose
 - **Timing:** Morning: 1 dose (with breakfast), Early afternoon: 1 dose (with lunch, before 2pm)
 - **First dose can be added Week 2**
 - Neuroprotection, focus enhancement

3. **Prescription options** (if severe cognitive impairment - **REQUIRES PHYSICIAN**):

- **Modafinil:** 100–200 mg **MORNING ONLY**
 - **Dose for average adult:** Start 100 mg, increase to 200 mg if needed
 - **Timing:** Morning **ONLY:** 1 dose (upon waking, 7–8am). **DO NOT** take after 10am - will destroy sleep.
 - **PRESCRIPTION REQUIRED**
 - Wakefulness agent, often prescribed off-label for ME/CFS
 - **WARNING:** Can mask fatigue signals and lead to PEM crashes. Use with **STRICT** pacing limits from Protocol 7.
- **Or: Methylphenidate:** 5–10 mg twice daily
 - **Dose for average adult:** 5–10 mg per dose
 - **Timing:** Morning: 1 dose (8am), Midday: 1 dose (12pm). **DO NOT** take after

2pm.

- **PRESCRIPTION REQUIRED (controlled substance)**
- Dopaminergic stimulant
- **CRITICAL WARNING:** Highly addictive. Can mask fatigue and lead to severe PEM crashes. Use **ONLY** with strict pacing. **DO NOT** use if history of substance abuse.
- Or: **Atomoxetine:** 40–80 mg daily
 - **Dose for average adult:** Start 40 mg daily for 1 week, increase to 80 mg if tolerated
 - **Timing:** Morning: 1 dose (with breakfast)
 - **PRESCRIPTION REQUIRED**
 - Norepinephrine reuptake inhibitor, non-stimulant option
 - Takes 2–4 weeks for full effect

Intranasal Delivery for CNS-Targeted Compounds

★ Key Point: BBB Vulnerability and CNS Drug Delivery

The selective energy dysfunction hypothesis (Chapter 14.24, lines 238–257) proposes that the blood-brain barrier (BBB) may be vulnerable in ME/CFS, limiting delivery of compounds needed for CNS support. Many compounds with strong peripheral effects show poor BBB penetration.

→ Recommendation 5: Prioritizing Intranasal Formulations for CNS Cognitive Support

Rationale: Intranasal delivery bypasses the BBB via olfactory and trigeminal nerve pathways, achieving CSF concentrations 2–10 fold higher than oral routes for the same compound.

Practical application (when options exist):

- **L-tyrosine:** Oral absorption is standard. If cognitive dysfunction dominates and oral supplementation insufficient, discuss intranasal dopamine or L-DOPA analogues with a neurologist. NOT currently standard care but mechanistically rational.
- **Modafinil:** Primarily oral. However, intranasal formulations have been explored for improving cognitive outcomes in some neurological conditions. If severe brain fog unresponsive to oral modafinil, ask physician about compounding as intranasal spray (requires specialist evaluation).
- **Insulin (experimental):** Intranasal insulin is being studied for cognitive support in neurodegenerative disease (Alzheimer's, dementia) by providing direct CNS metabolic support. NOT established in ME/CFS but mechanistically relevant to astrocyte energy gate hypothesis (Chapter 14.24, lines 179–198). **EXPERIMENTAL;** do not pursue without specialist guidance.
- **Future consideration:** As mechanistic understanding of ME/CFS CNS dysfunction

improves, intranasal delivery of neuroprotective or energetic compounds (ketone bodies, lactate precursors, neuropeptides) may emerge as targeted interventions. Current standard protocols primarily use oral routes.

Evidence level: Speculative (established for other conditions; no ME/CFS-specific intranasal RCTs)

Action item: If cognitive symptoms dominate despite oral supplementation, discuss BBB-penetrant strategies and intranasal formulations with a neurologist familiar with ME/CFS.

Expected Relief

- **Mental clarity:** 30–50% improvement in first week
- **Processing speed:** 20–40% improvement
- **Word-finding:** Improved (especially with choline support)
- **Sustained attention:** Increased from minutes to 20–60 minutes
- **Best responders:** Those with prominent brain fog as limiting symptom

17.3.3 Passive Neuromuscular Electrical Stimulation (NMES) for CNS Bypass

~ Hypothesis 2: Pharmacological Bypass of CNS Motor Coordination Failure

The selective energy dysfunction hypothesis (Chapter 14.24, lines 376–391, 500–509) proposes that peripheral muscle remains capable but CNS motor coordination fails. Direct electrical stimulation bypasses CNS coordination entirely, activating muscle without requiring CNS drive. This may preserve muscle function and prevent deconditioning in severely immobilized patients.

Certainty: 0.30 (CNS-peripheral dissociation is mechanistically plausible within the selective dysfunction framework; NMES is well-established for deconditioning prevention in ICU patients [235]; no direct evidence for CNS bypass mechanism in ME/CFS; extension of theoretical framework)

→ Recommendation 6: Passive NMES Protocol for Deconditioning Prevention in Severe Cases

Mechanism: Neuromuscular electrical stimulation (NMES) produces muscle contractions without voluntary effort or CNS coordination. Applied passively (patient rests, device stimulates), this provides:

- Muscle fiber recruitment without CNS energy expenditure
- Prevention of atrophy during extended bedrest
- Maintenance of mitochondrial function in muscle tissue
- Potential maintenance of motor unit recruitment patterns

Indications: Severe cases with extended bedrest (>2 weeks immobilized) at risk of deconditioning and contracture.

Protocol:

1. **Equipment:** Seek qualified physical therapist familiar with NMES. Commercial NMES devices (Compex, BioMed) cost \$200–600. Medicare may cover if prescribed by physician for deconditioning prevention.
2. **Application:** Supine position (no standing required)
 - Electrode placement: Large muscle groups (quadriceps, gluteus maximus, gastrocnemius)
 - Frequency: 20–30 minutes daily
 - Intensity: Submaximal (visible muscle contraction without pain)
 - Frequency (Hz): 50 Hz optimal for endurance recruitment
3. **Safety monitoring:**
 - Monitor heart rate (electrical stimulation can affect autonomic tone)
 - Avoid over-stimulation (can trigger metabolic demand and PEM)
 - Stop if orthostatic symptoms increase following session
 - One 20–30 min session daily; do NOT exceed without supervision
4. **Integration with pacing:** NMES does not “count” as activity if passive (patient rests while stimulated). However, metabolic demand may increase. Track symptoms day-of and 24–48h post-treatment for delayed PEM.

Evidence level: Plausible (extends midodrine bypass principle from Ch14j lines 500–509; no ME/CFS-specific NMES trials exist). NMES is well-established for deconditioning prevention in ICU and spinal injury patients.

Dosing note: Verify with qualified physical therapist. Medical supervision recommended given autonomic vulnerability.

Expected outcome: Prevention of rapid muscle atrophy during extended immobility; potential maintenance of motor recruitment capacity. NOT expected to produce functional improvement or increase activity tolerance—purely preventive.

17.3.4 Baseline Symptom Reduction: Strict Pacing

[IMMEDIATELY ACTIONABLE - NO RESEARCH NEEDED]

Critical Foundation (Implement Immediately) Pacing is *not* a treatment, but it *prevents worsening* and reduces baseline symptom burden. The post-exertional malaise mechanism (Section 6.2.3) documents how exertion beyond capacity triggers mitochondrial dysfunction, oxidative stress, and immune activation. Without pacing, other interventions will be less effective.

★ **Key Point: Experimental: Emergency Post-Exertion Protocol**

For situations where exertion is **unavoidable** (medical procedures, emergencies, essential activities), an experimental post-exertion intervention protocol exists that may reduce PEM severity or prevent crashes. This protocol targets the 24–72 hour window between exertion and symptom onset with ATP substrates (D-ribose, MCT oil), NAD⁺ precursors, antioxidants, and anti-inflammatory support.

Evidence tier: Mechanistically justified but clinically unvalidated. No RCTs exist. Individual components have safety data.

Appropriate use: Unavoidable medical procedures, accidental overexertion, emergency situations—NOT routine use to enable regular overexertion.

Critical principle: This protocol addresses BOTH energy restoration (ATP/NAD⁺ support) AND inflammatory cascade interruption. Anti-inflammatories alone are insufficient; the core problem is ATP production failure.

See Chapter 24, §24.10.3 for complete protocol details, rationale, and safety considerations.

Note: This is NOT a substitute for pacing, which remains the evidence-based foundation. Use pacing to avoid crashes; reserve emergency protocol for truly unavoidable situations.

Heart Rate-Based Pacing Protocol

1. **Equipment** (purchase today with overnight shipping):
 - **Heart rate monitor options:**
 - **Chest strap:** Polar H10, Garmin HRM-Dual (\$60–90, most accurate)
 - **Optical wrist:** Fitbit Charge 5, Garmin Vivosmart 5 (\$100–150, convenient)
 - **Budget:** CooSpo H6 chest strap (\$30, pairs with phone apps)
 - **Smartphone apps:** Most monitors pair with free apps (Polar Beat, Garmin Connect, etc.)
 - **Purchase NOW:** Choose one option and order with fastest shipping. This is your most important tool.
2. **Calculate your personal anaerobic threshold (AT) - DO THIS NOW:**
 - **Formula:** $AT = (220 - \text{your age}) \times 0.55$
 - **Examples by age:**
 - Age 20: $AT = (220 - 20) \times 0.55 = 110$ bpm
 - Age 30: $AT = (220 - 30) \times 0.55 = 104$ bpm
 - Age 40: $AT = (220 - 40) \times 0.55 = 99$ bpm
 - Age 50: $AT = (220 - 50) \times 0.55 = 93$ bpm
 - Age 60: $AT = (220 - 60) \times 0.55 = 88$ bpm
 - **Write down YOUR number:** _____ bpm
 - **This is your absolute ceiling for ALL activities**
 - Gold standard: Cardiopulmonary exercise test (CPET) if available - provides precise AT

3. STRICT RULE - Start following THIS MOMENT:

- **Monitor heart rate continuously during ALL activities** (walking, showering, eating, talking)
- **When HR approaches AT (within 5 bpm):**
 - a) STOP the activity IMMEDIATELY - do not finish the task
 - b) Lie down HORIZONTALLY (not sitting - sitting requires postural energy)
 - c) Do NOT resume until HR returns to resting baseline (typically 60–80 bpm)
 - d) Wait minimum 5–10 minutes after HR normalizes before resuming
- **Activities that commonly exceed AT** (monitor closely):
 - Showering (warm water increases HR)
 - Walking upstairs
 - Extended conversations
 - Emotional stress
 - Eating large meals
- **Until HR monitor arrives:** Use perceived exertion. If breathing becomes slightly harder or you feel warmth, STOP.

Activity Modification for Severe Cases

- **Default position:** Horizontal (not sitting)
- **All activities in bed/reclining:**
 - Phone use, eating, computer work (laptop on lap desk)
 - Showering: Shower chair *mandatory* (standing shower is major exertion)
 - Tooth brushing: Electric toothbrush in bed, or sitting
- **Activity blocks:** 15–30 minutes maximum, then 30–60 minute horizontal rest
- **Pre-emptive rest:** *Before* fatigue sets in (do not wait until crashed)

Energy Triage for Severe Cases: Prioritizing Tier 1–3 Activities Only

★ Key Point: Energy Triage Hierarchy for Severe ME/CFS

In severe ME/CFS, CNS energy is so limited that only Tier 1–3 activities (brainstem functions, sensory processing, basic motor) are sustainable. Tier 4–6 (memory, executive function, complex cognition) must be eliminated entirely.

Severe Case Adaptation (from selective dysfunction hypothesis, Chapter 14.24):

- **Tier 1–2 ONLY:** Breathing, vital functions, passive sensory input (listening, watching)
- **Tier 3 (sparingly):** Simple movements in bed, basic self-care
- **Tier 4–6 (OFF LIMITS):** Memory tasks, reading, conversations, planning, any decision-making

→ **Recommendation 7: Severe Case Cognitive Triage: Eliminate Complex Cognition**

For bedbound and housebound severe patients: Your CNS energy budget is insufficient for all but survival functions. Rather than attempting to grade activities, simply eliminate cognitive demands entirely:

- **Eliminate:** Reading (requires sustained attention), conversations (require processing), planning (executive function), decision-making (energy-expensive), learning new information
- **Allow:** Passive listening to audiobooks/podcasts (no attention required), watching familiar shows (no processing needed), lying quietly, resting
- **Rationale:** Tier 6 (complex cognition) and Tier 5 (executive function) will fail first under energy scarcity. Accept this rather than fighting it. Preserve your limited mental energy for essential communication only.

If you must communicate: Keep to essential topics only (health needs, medications, emergencies). Avoid extended conversations, explanations, or discussions. Keep responses to single sentences. Let others do the cognitive work.

Cognitive Pacing

- Screen time limits (cognitive exertion triggers PEM)
- Conversations: 10–15 minutes maximum, then rest
- Reading: Short blocks (5–15 minutes) with rest (or skip entirely if too demanding)
- Decision-making: Minimize (decision fatigue is real and severe)—let others decide when possible

Expected Outcomes

- **PEM frequency:** 50–80% reduction within 1–2 weeks
- **Baseline symptom severity:** 20–40% improvement (less chronic immune activation)
- **Functional capacity:** Stable rather than progressively declining
- **Quality of life:** Significant (fewer crashes = more predictability, ability to plan small activities)

17.4 Expected 2-Week Outcomes

17.4.1 Cumulative Symptom Relief

17.4.2 Transformation of Tolerability

Before Protocol

Table 17.1: Expected symptom improvement at 2 weeks with full protocol

Symptom Domain	Expected Improvement	Timeline
Brain fog	40–60%	3–7 days (MCAS + sleep + cognitive support)
Orthostatic intolerance	60–80%	1–3 days (salt + compression)
Pain (myalgia, headache)	40–60%	Hours–days (NSAIDs + gabapentin)
Sleep quality	50–70%	1–7 nights (pharmaceutical support)
GI symptoms	60–80%	1–7 days (symptomatic + mechanistic)
PEM frequency	50–80%	1–2 weeks (strict pacing)
Overall suffering	50–70% reduction	2 weeks combined

- Constant severe symptoms across multiple domains
- Unable to tolerate upright position
- Cognitive function severely impaired
- Pain uncontrolled
- GI symptoms limiting food intake
- Non-restorative sleep perpetuating all symptoms
- Overall suffering: 9/10, unbearable, considering medical assistance in dying

After 2-Week Protocol

- Brain fog reduced by half, can read/watch shows in short blocks
- Can tolerate sitting/standing 2–4 times longer with compression + salt
- Pain reduced from 8/10 to 4/10, manageable with multi-modal approach
- Sleeping 6–8 hours (vs. 2–4 hours fragmented)
- Can eat comfortably, GI symptoms controlled
- PEM frequency dramatically reduced (avoiding triggers with pacing)
- Overall suffering: 4–5/10, difficult but bearable, can envision continuing

17.4.3 Critical Threshold: Bearability

The goal is *not* cure or remission within 2 weeks—that is unrealistic. The goal is to reduce suffering from **unbearable** to **bearable**, buying time to pursue longer-term fundamental treatments (Section 17.7).

For patients considering medical assistance in dying, this reduction in suffering can mean the difference between ending life and continuing to fight for recovery.

17.5 Sensitization Prevention: Foundational Strategy for Severe Cases

► Protocol 1: Sensitization-Prevention Protocol for Severe Patients

Rather than treating ME/CFS symptomatically in isolation, this protocol addresses the critical observation that severe disease represents a cascade of progressive sensitization [278]. Each new system involvement, infection, or crash deepens neuroinflammatory priming and reduces tolerance for future perturbations. Prevention of further sensitization is the primary therapeutic goal for severe patients.

Anti-Neuroinflammatory Foundation Initiate immediately:

- **Low-dose naltrexone (LDN):** Titrate from 0.5 mg to 4.5 mg over 4–6 weeks at bedtime. Monitor for psychiatric effects (mood destabilization, emotional lability) as microglial downregulation paradoxically can unmask underlying mood pathology. Benefits include glial cell downregulation and endogenous opioid restoration [279].
- **Palmitoylethanolamide (PEA):** 1200 mg daily (micronized form only; standard PEA poorly absorbed). PPAR- α agonism provides sustained anti-inflammatory and mast cell stabilizing effects [502].
- **Omega-3 polyunsaturated fatty acids:** 3 g EPA/DHA daily (split into meals for tolerability). Supports neuronal membrane fluidity and endogenous anti-inflammatory mediator production [278].

Strict Pacing with Objective Monitoring Sensitization amplification occurs through repeated crashes and energy envelope violations. Prevent this:

- **Continuous heart rate monitoring:** Wear device 24/7; learn individual anaerobic threshold (AT). Maintain activities strictly below AT. For bedbound patients, even small movements (self-care, family interaction) must be monitored.
- **Zero-crash goal:** The protocol explicitly aims for zero crashes during the sensitization prevention window (ideally 3–6 months). Each crash resets progress by consuming recovery reserve and amplifying sensitization.
- **Anticipatory pacing:** Reduce activity before feeling symptomatic. Severe patients must learn to pace preemptively rather than reactively.

Sleep Architecture Optimization Sleep loss is a primary driver of glymphatic impairment and sustained neuroinflammation [278]. This is non-negotiable:

- **Sleep study mandatory:** Identify and treat sleep apnea, periodic limb movements, or REM behavior disorder.
- **Pharmaceutical sleep support:** Melatonin (1–3 mg) or dual orexin receptor antagonist (suvorexant 5–10 mg) prioritized for slow-wave sleep enhancement. Target: 7–8 hours with $\geq 85\%$ sleep efficiency.

- **Positional optimization:** Side sleeping (particularly left side) optimizes glymphatic clearance geometry. Avoid supine positioning if possible.

Infection Barrier Maintenance Each infection triggers neuroinflammatory amplification and can produce permanent baseline functional decline. Prevention is critical:

- **Prophylactic antivirals:** N95/FFP2 masking in any crowded setting. Consider prophylactic famciclovir (250 mg daily) or valacyclovir (500 mg daily) during winter/high-transmission seasons, especially if frequent reactivation is documented.
- **First-symptom protocol:** At first symptom of infection (sore throat, malaise, fever), immediately begin antiviral therapy (high-dose valacyclovir 1000 mg QID or IV acyclovir if tolerated) and rest protocol. Early treatment reduces infectious-trigger damage magnitude [279].
- **Immunization strategy:** Live vaccines absolutely contraindicated. Inactivated vaccines acceptable only during periods of clinical stability; discuss timing with knowledgeable provider.

★ **Key Point: Astrocyte Energy Gate and Lactate Shuttle**

The astrocyte-neuron lactate shuttle (ANLS) is estimated neuronal ATP [174]. The hypothesis that ANLS dysfunction means energy failure while peripheral tissues remain intact is discussed. If this hypothesis is correct, supporting the lactate shuttle may improve availability.

Lactate Shuttle Cofactor Optimization

→ **Recommendation 8: Lactate Shuttle Cofactor Protocol for Severe Cases****Components:**1. **Thiamine (Vitamin B1):** 100–300 mg daily

- **Dose:** Start 100 mg daily, titrate to 200–300 mg if tolerated
- **Rationale:** Thiamine is essential for pyruvate dehydrogenase complex, which converts pyruvate to acetyl-CoA for CNS energy production. Dosing exceeds RDA (1.1–1.2 mg) but aligns with therapeutic B1 dosing for neurological conditions.
- **Timing:** Morning with breakfast
- **Safety:** Very safe; water-soluble, excess excreted. No upper limit established.

2. **MCT Oil (Medium-Chain Triglycerides):** Titrate from 1 tsp bedtime to 1 tbsp TID

- **Rationale:** MCTs bypass long-chain fatty acid metabolism and provide rapid ketone production, providing alternative CNS fuel when lactate shuttle is impaired
- **Protocol:** Start 1 tsp (5 mL = 5 g) at bedtime. After 1 week, add 1 tsp with breakfast. After 2 weeks, add 1 tsp with lunch. Target: 1 tbsp TID (15 mL = 15 g) if tolerated.
- **GI monitoring:** MCT can cause GI upset (cramping, diarrhea); titrate slowly. Take with food.
- **Evidence level:** Plausible (MCT metabolism established; ME/CFS-specific efficacy unproven)

3. **Exogenous Ketones (Optional):** Consider if MCT insufficient

- **Ketone bodies:** Provide direct neuronal fuel independent of glucose-lactate pathway
- **Options:** Ketone salts (sodium/calcium beta-hydroxybutyrate) or MCT oil-derived ketones
- **Dose:** 5–15 g daily divided doses (follow product guidelines)
- **Evidence level:** Speculative (ketones show promise in neurological conditions; ME/CFS efficacy not established)

Evidence level: Plausible (ANLS dysfunction mechanistically relevant; clinical validation pending)

Expected outcomes: Improved cognitive clarity, reduced fatigue, better sustained attention. Timeline: 2–4 weeks for noticeable effect.

Metabolic Protection and Monitoring Severe ME/CFS is characterized by hypometabolic state and impaired glucose handling [278]. Additional metabolic stress amplifies sensitization:

- **Quarterly HbA1c monitoring:** If trending upward (especially if approaching prediabetes range $\geq 5.7\%$), initiate metformin 500 mg daily (titrate to tolerance). Metabolic dysregulation intensifies neuroinflammation and reduces treatment efficacy [278].
- **Caloric adequacy:** Severe patients often restrict intake due to GI symptoms or cognitive dysfunction. Ensure minimum caloric support (target: 1500–1800 kcal daily). Undereating amplifies mitochondrial stress and neuroinflammation.
- **Electrolyte homeostasis:** Monitor sodium/potassium; replace as needed. Many severe patients become sodium-depleted from reduced oral intake; careful repletion supports autonomic stability.

Rationale and Duration The sensitization-prevention protocol targets the documented cascade in which neuroinflammatory priming lowers the threshold for all symptom exacerbations (see Chapter 14.24). By stabilizing this foundation, subsequent treatments for individual systems (immune reconstitution, cognitive rehabilitation, autonomic retraining) have higher probability of efficacy.

Duration: Maintain this protocol for minimum 3–6 months as foundation. Once sensitization is arrested (no new crashes, stable baseline), can add additional interventions from Section 17.7.

17.6 Glymphatic Enhancement Protocol for Severe Cases

► Protocol 2: Glymphatic Enhancement for Neuroinflammation Reduction

Conceptual Framework The glymphatic system is the brain's waste clearance mechanism, dependent on aquaporin-4 water channels in astrocytes and functioning primarily during sleep. Recent neuroimaging and cerebrospinal fluid (CSF) studies suggest glymphatic impairment in ME/CFS, potentially allowing accumulation of neuroinflammatory products [278]. This protocol aims to enhance glymphatic clearance to reduce neuroinflammation accumulation.

Sleep Architecture Optimization (Primary Intervention) Sleep optimization is the highest-priority glymphatic enhancement:

- **Sleep study:** Identify treatable pathology (sleep apnea, PLMD, REM behavior disorder). These conditions actively impair glymphatic function and must be corrected.
- **Pharmaceutical enhancement of slow-wave sleep:** Melatonin (1–3 mg) or dual orexin receptor antagonist (suvorexant, daridorexant). Goal: Increase slow-wave sleep percentage from typical ME/CFS baseline (often $<10\%$) to $\geq 15\%$.
- **Consistent sleep timing:** Sleep-wake cycle disruption impairs glymphatic function. Target: Same bedtime and wake time daily (even weekends).
- **Positional support:** Left lateral decubitus position optimizes CSF flow geometry for glymphatic function. Support with pillows to maintain position through sleep.

Glymphatic Support Agents Adjunctive to sleep optimization:

- **Omega-3 polyunsaturated fatty acids:** 3 g EPA/DHA daily. Polyunsaturated fatty acids are critical for neuronal membrane fluidity, which is essential for aquaporin-4 function. May be synergistic with sleep optimization [278].
- **Taurine:** 2–3 g daily. Critical osmoregulation agent for astrocyte water channels; supports aquaporin-4-mediated water transport through the interstitium.
- **Low-dose lithium** (if baseline renal function normal): 150–300 mg daily. Enhances aquaporin-4 expression and glymphatic flow; also provides mood stabilization and neuroprotection. Requires baseline and periodic (q3 months) renal function and TSH monitoring [278].

Activity Timing for Neuroinflammation Minimization Structure daily activities to minimize inflammatory triggers before sleep:

- **Inflammatory activities early:** Any activity likely to trigger small inflammatory responses (visits with others, unavoidable exertion, stressful events) should occur in early morning or midday.
- **Wind-down protocol:** 3–4 hours before sleep, minimize all stimulation. Avoid emotional stress, heavy cognitive demands, physical activity. This allows inflammatory activation to resolve before entering sleep.
- **Maximum recovery time:** Ensure 8–9 hours of uninterrupted sleep opportunity. Glymphatic clearance is maximal during early slow-wave sleep phases; fragmentation impairs clearance.

Monitoring for Efficacy Glymphatic enhancement efficacy may manifest as:

- Improved cognitive clarity or reduced brain fog
- Reduced headaches (especially morning headaches suggesting impaired overnight clearance)
- Improved mood (glymphatic impairment has been associated with mood dysfunction and neuroinflammation)
- Reduced generalized pain (central sensitization may improve with reduced neuroinflammatory tone)

If no improvement after 8–12 weeks of sleep optimization + adjunctive agents, consider whether sleep apnea or other treatable sleep pathology is present despite initial sleep study.

Rationale and Integration Glymphatic enhancement is mechanistically complementary to sensitization prevention (Section 17.5). While the sensitization-prevention protocol addresses active microglial activation and anti-inflammatory support, the glymphatic protocol targets clearance of neuroinflammatory products themselves. Combined, they address both active neuroinflammation production and passive clearance impairment [278].

17.7 Medium-Term Recovery Strategies (Weeks to Months)

After achieving initial symptom control, pursue fundamental treatments targeting disease mechanisms identified in Chapters 6–7. These interventions address root pathophysiology documented through biomarker research (Sections 7.8 and 6.14).

17.7.1 Immunoabsorption for Cognitive Dysfunction

Speculation 53 (EV Depletion as Primary Mechanism of Immunoabsorption Benefit). **Original Contribution:** This document proposes a novel mechanism for immunoabsorption efficacy in ME/CFS. Rather than attributing benefits solely to autoantibody removal (the traditional explanation), we hypothesize that **extracellular vesicle (EV) depletion** may be the primary therapeutic mechanism. Giloteaux et al. [214] found elevated IL-2 and inflammatory cytokines specifically in EVs. Standard immunoabsorption removes EVs along with antibodies. This “Pathogenic EV” hypothesis (Section 7.9.1) suggests EVs containing cytokines cross the blood-brain barrier, activate microglia, and cause cognitive dysfunction. **No prior literature has explicitly proposed EV depletion as the mechanism of immunoabsorption benefit in ME/CFS.**

Certainty: 0.20 (EV elevation documented [214]; immunoabsorption removes EVs as a physical consequence; causal primacy of EV depletion over autoantibody removal for clinical benefit is entirely unproven and no direct evidence exists)

Rationale Section 7.9.1 presents the “Pathogenic Extracellular Vesicle” hypothesis. Autoantibodies targeting G-protein coupled receptors (Section 7.6.1) may disrupt autonomic function and cerebral blood flow. Stein et al. [118] demonstrated 70% response rate in post-COVID ME/CFS patients, with benefits sustained to 6 months. Mechanism likely involves removal of both autoantibodies and pathogenic extracellular vesicles containing inflammatory cytokines.

Intervention [NOVEL - Available NOW but requires specialist center]

- **Procedure:** Immunoabsorption (plasmapheresis variant using IgG-selective columns)
- **Detailed protocol schedule:**
 - **Session frequency:** 5 sessions over 10 days (Day 1, 3, 5, 7, 10)
 - **Session duration:** 2–4 hours per session
 - **Timing:** Morning sessions preferred (9am–1pm)
 - **Blood volume processed:** 2–3 liters per session
 - **Anticoagulation:** Heparin during procedure (standard protocol)
 - **Complete treatment course:** 10 days total from first to last session
- **Preparation:**
 - Adequate hydration: Drink 1–2 L water before each session
 - Continue electrolyte protocol (Protocol 2) throughout treatment

- Light meal 1–2 hours before (avoid large meals)
- Bring blanket (procedure rooms are cool), entertainment (phone, book)
- **Mechanism:** Removes IgG (including GPCR autoantibodies) AND extracellular vesicles containing inflammatory cytokines
- **Availability:** European centers (Germany, Norway); medical tourism may be necessary
 - **Germany:** Charité Berlin (contact via ME/CFS specialty clinic)
 - **Norway:** Haukeland University Hospital, Bergen
 - Outpatient procedure - can stay in local hotel between sessions
- **Cost:** €5,000–€15,000 depending on country/insurance
 - Germany: Often covered by statutory insurance with medical necessity
 - Medical tourism: Budget €10,000–€15,000 including travel/accommodation

Expected Outcomes

- **Response rate:** 70% (per Stein 2024)
- **Timeline:** Improvement within days to weeks
- **Best responders:** Severe cognitive dysfunction, autoantibody-positive patients
- **Durability:** Benefits sustained 6+ months in responders
- **Target population:** Severe cases with significant cognitive impairment, particularly those with elevated GPCR autoantibodies

Pursuing Immunoadsorption

1. Screen for GPCR autoantibodies (CellTrend ELISA - Germany)
2. If positive or severe cognitive dysfunction: pursue immunoadsorption
3. Contact centers: Charité Berlin (Germany), Haukeland University Hospital (Norway)
4. If insurance denial: medical tourism, crowdfunding, patient advocacy organizations

17.7.2 Low-Dose IL-2 for Autoimmune Features

Speculation 54 (Low-Dose IL-2 for Treg Restoration in ME/CFS). **Original Contribution:** This document is the **first to explicitly propose low-dose IL-2 therapy for ME/CFS** based on convergent recent evidence. Giloteaux et al. [214] found elevated IL-2 in extracellular vesicles; Hunter et al. [215] identified IL-2 signaling dysregulation in epigenetic biomarker panel; multiple studies document Treg deficiency. While low-dose IL-2 is established therapy for SLE and type 1 diabetes, **its application to ME/CFS with this specific mechanistic rationale is novel**. This represents an immediately actionable intervention using an FDA-approved drug with precedent in autoimmune disease.

Certainty: 0.25 (Treg deficiency documented in ME/CFS [189]; IL-2 pathway dysregulation identified [215, 214]; low-dose IL-2 efficacy established in SLE and type 1 diabetes; extrapolation to ME/CFS is speculative with no ME/CFS-specific trials)

Rationale Section 7.3.1 presents convergent evidence for IL-2 pathway dysfunction. Section 7.2.1 documents regulatory T cell (Treg) deficiency and T-cell exhaustion in ME/CFS. Autoantibodies (Section 7.6.1) suggest ongoing autoimmune processes. Low-dose IL-2 therapy selectively expands regulatory T cells, restoring immune tolerance and potentially suppressing autoantibody production.

Intervention [NOVEL - PRESCRIPTION REQUIRED but immediately available]

- **Drug:** Aldesleukin (Proleukin) - FDA-approved IL-2, used off-label at low dose
- **Detailed dosing protocol:**
 - **Dose:** 1–2 million IU (international units) per injection
 - **Start dose:** Begin with 1 million IU to assess tolerance
 - **Frequency:** 2–3 times per week (Monday/Wednesday/Friday OR Monday/Thursday)
 - **Route:** Subcutaneous injection (like insulin - abdomen, thighs, or upper arms)
 - **Timing:** Evening preferred (5–7pm) - if flu-like symptoms occur, sleep through them
 - **Duration:** 12 weeks initial course (24–36 total injections)
 - **Dose escalation:** If no response at 4 weeks and good tolerance, increase to 2 million IU
- **Administration technique:**
 - Reconstitute powder with sterile water (comes with kit)
 - Use insulin syringe (0.5–1 mL)
 - Inject subcutaneously at 45-degree angle
 - Rotate injection sites to avoid bruising
 - Store reconstituted drug in refrigerator, use within 24 hours
- **Monitoring schedule:**
 - **Baseline** (before starting): CBC, CMP, Treg percentage (CD4⁺CD25⁺FoxP3⁺ flow cytometry)
 - **Week 2:** Treg percentage (should see early expansion)
 - **Week 4:** Treg percentage, CBC (watch for eosinophilia)
 - **Week 8:** Treg percentage, CBC, CMP
 - **Week 12:** Full panel (Treg, CBC, CMP, symptom assessment)
 - **Symptom diary:** Daily (track PEM, fatigue, cognitive function)
- **Expected side effects** (usually mild):
 - Flu-like symptoms first 24–48 hours after injection (fever, chills, fatigue)
 - Injection site redness (normal)
 - Transient mild rash
 - Take ibuprofen 400 mg with injection if flu-like symptoms bothersome

Patient Selection

- Documented Treg deficiency: CD4⁺CD25⁺FoxP3⁺ <5% of CD4⁺ T cells
- Elevated autoantibodies (GPCR antibodies, ANA-positive)
- Clinical autoimmune features (skin rashes, arthritis, sicca symptoms)
- Any disease duration (works for both early and late disease)

Expected Outcomes

- **Mechanistic confirmation:** Treg expansion within 2–4 weeks (indicates pathway intact)
- **Clinical response:** 6–12 weeks if effective
- **Symptom targets:** Autoimmune symptoms, potentially fatigue and PEM if autoimmunity is maintaining factor
- **Safety:** Generally well-tolerated; flu-like symptoms possible

Accessing Low-Dose IL-2

- Requires prescription from hematologist, immunologist, or sympathetic physician
- Off-label use (approved for cancer at high dose, used low-dose in autoimmune diseases)
- Precedent: Used in SLE, type 1 diabetes, GVHD
- Cost: Variable depending on country/insurance; compounding pharmacies may reduce cost

17.7.3 Hormonal Modulation (Post-Menopausal Women)

Speculation 55 (Estrogen as Immune Modulator in Post-Menopausal ME/CFS). **Original Contribution:** This document is the **first to propose estrogen supplementation specifically for ME/CFS** based on Che et al.'s 2025 finding [189] of exaggerated IL-6 responses in post-menopausal women with low estradiol. While HRT is established therapy, **targeting it to ME/CFS patients based on documented sex-specific immune dysregulation is novel**. This represents a precision medicine approach: post-menopausal women with severe ME/CFS and low estradiol may benefit from HRT not just for menopausal symptoms, but for **direct immune modulation**. Applicable to 15–20% of severe cases.

Certainty: 0.30 (sex-specific IL-6 amplification in low-estradiol women documented [189]; estrogen immunomodulatory effects on IL-6 and TNF- α established; direct evidence for HRT improving ME/CFS outcomes is absent; extrapolation from biomarker finding to therapeutic efficacy is speculative)

Rationale Section 40 documents exaggerated IL-6 responses in women over 45 with diminished estradiol. Estrogen receptors on immune cells directly modulate cytokine production (Section 7.3.1); estrogen reduces IL-6, TNF- α , IL-1 β production. Restoring physiological estrogen levels may dampen immune hyperactivation.

Intervention [NOVEL - PRESCRIPTION REQUIRED but immediately available]

- **Population:** Post-menopausal women with documented low estradiol (<30 pg/mL) and severe ME/CFS
- **Detailed HRT protocol:**
 - **Estradiol:** Transdermal patch 0.05–0.1 mg/day
 - * **Start dose:** 0.05 mg/day patch (lower dose)
 - * **Application:** Apply 1 patch to clean, dry skin (abdomen, buttocks, or upper arm)
 - * **Schedule:** Change patch twice weekly (e.g., Monday and Thursday) OR once weekly depending on product
 - * **Timing:** Apply at same time of day (morning or evening)
 - * **Rotation:** Rotate application sites each time (avoid same spot for 1 week)
 - * **Products:** Estradot, Vivelle-Dot, Climara (brand varies by country)
 - * **Dose escalation:** If no benefit at 4–6 weeks, increase to 0.1 mg/day
 - **Progesterone (MANDATORY if you still have uterus):**
 - * **Drug:** Micronized progesterone (Prometrium, Utrogestan)
 - * **Dose:** 100–200 mg daily
 - * **Timing:** Evening: 1 dose (at bedtime, 9–10pm)
 - * **Why bedtime:** Progesterone causes mild sedation - use to aid sleep
 - * **Schedule:** Take EVERY night continuously (do NOT skip nights)
 - * **CRITICAL:** DO NOT use estrogen without progesterone if you have uterus - endometrial cancer risk
 - **First application can be TONIGHT** (if prescription obtained):
 - * Apply estradiol patch to skin
 - * Take progesterone at bedtime
 - * Continue daily/weekly as scheduled
- **Baseline testing** (before starting):
 - Estradiol level (blood test - should be <30 pg/mL)
 - IL-6 level (optional - to track immune marker)
 - Mammogram if over 40 and not up-to-date
 - Blood pressure baseline
- **Monitoring schedule:**
 - **Month 1:** Symptom diary, any side effects
 - **Month 3:** Estradiol level (ensure in physiological range 50–200 pg/mL), symptom assessment, PEM frequency
 - **Month 6:** Full assessment - IL-6 (if measured baseline), symptom severity, PEM frequency
 - **Yearly:** Mammogram, pelvic exam (if intact uterus)
- **Contraindications** (DO NOT use if):

- Personal history of breast cancer, endometrial cancer
- Active DVT/PE (blood clots) or history of hormone-related clots
- Unexplained vaginal bleeding
- Active liver disease
- Pregnancy (verify not pregnant before starting)

Expected Outcomes

- **Timeline:** 3–6 months for full benefit
- **Targets:** Immune hyperactivation, PEM severity, overall symptom burden
- **Applicability:** 15–20% of severe cases (post-menopausal women)
- **Safety:** Standard HRT risks (thrombosis, breast cancer - discuss with physician)

Implementation

- Screen estradiol levels (blood test)
- If low + severe ME/CFS → trial HRT
- Standard gynecology or primary care can prescribe
- Monitor symptom response at 3 and 6 months
- If clear benefit → continue; if no benefit after 6 months → discontinue

17.7.4 Anti-Cytokine Therapy (Early Disease <3 Years)

~ Hypothesis 3: Immune Exhaustion Timeline: Duration-Stratified Therapeutic Window

Original Contribution: This document proposes the “**Immune Exhaustion Timeline**” hypothesis—a completely novel framework for stratifying ME/CFS treatment by disease duration. Based on Hornig et al.’s finding [212] that cytokines normalize after 3 years, we propose a **time-sensitive therapeutic window**: anti-cytokine biologics may only benefit patients in the early hyperactive phase before T-cell exhaustion occurs [119]. **No prior protocol has explicitly stratified anti-cytokine therapy by illness duration in ME/CFS.** This represents a disease-modifying approach rather than pure symptom management. If validated, this framework would fundamentally change how newly diagnosed patients are treated.

Certainty: 0.35 (cytokine normalization after 3 years documented [212]; T-cell exhaustion markers elevated in ME/CFS [119]; duration-stratified therapeutic window is a logical inference but remains unvalidated; no trials testing this framework exist)

Rationale Section 7.3.1 documents that cytokine elevations occur primarily in early disease (<3 years). Section 7.9.1 presents the “Immune Exhaustion Timeline” hypothesis: a time-sensitive therapeutic window exists before immune exhaustion (Section 7.2.1) sets in. Early anti-cytokine intervention may prevent progression to chronic immune dysregulation.

Intervention [NOVEL FRAMEWORK - PRESCRIPTION REQUIRED, high cost, requires specialist]

- **Population:** Severe ME/CFS with ALL of the following:
 - Illness duration <3 years from onset
 - Documented cytokine elevation: IL-6 >5 pg/mL OR TNF- α >10 pg/mL OR multiple cytokines elevated
 - Severe disability preventing work/school
 - Failed standard symptomatic treatments
- **Detailed anti-cytokine protocols:**
 - **Option 1: Tocilizumab (Actemra)** - IL-6 receptor blocker
 - * **Dose:** 162 mg subcutaneous injection
 - * **Frequency:** Once monthly (same day each month, e.g., 1st of month)
 - * **Timing:** Can inject any time of day
 - * **Administration:** Pre-filled autoinjector pen (like EpiPen) - inject into thigh or abdomen
 - * **Duration:** 6-month course (6 total injections)
 - * **Storage:** Refrigerate, bring to room temperature 30 min before injection
 - * **Best for:** Patients with high IL-6 (>10 pg/mL)
 - **Option 2: Etanercept (Enbrel)** - TNF- α blocker
 - * **Dose:** 50 mg subcutaneous injection
 - * **Frequency:** Once weekly (same day each week, e.g., every Monday)
 - * **Timing:** Evening injection preferred (5–7pm)
 - * **Administration:** Pre-filled SureClick autoinjector - inject into thigh or abdomen
 - * **Duration:** 6-month course (24 total injections)
 - * **Storage:** Refrigerate, bring to room temperature 30 min before injection
 - * **Best for:** Patients with high TNF- α (>15 pg/mL) or prominent inflammation
- **Monitoring schedule** (CRITICAL - these are immunosuppressants):
 - **Baseline** (before starting):
 - * CBC, CMP, liver function tests
 - * Cytokine panel (IL-6, TNF- α , IL-2, IL-1 β)
 - * T-cell exhaustion markers (PD-1, Tim-3 expression) if available
 - * TB screening (QuantiFERON-TB Gold test)
 - * Hepatitis B/C screening
 - * Chest X-ray

- **Monthly monitoring:**
 - * CBC (watch for neutropenia - stop if ANC <1000)
 - * Liver function (stop if ALT >3× upper limit)
 - * Symptom severity scores
 - * Infection screening (fever, new symptoms)
- **3-month assessment:**
 - * Repeat cytokine panel (should show reduction)
 - * PEM frequency and severity
 - * Functional capacity assessment
 - * Decide: continue if benefit, stop if no response
- **6-month final assessment:**
 - * Full cytokine panel
 - * T-cell exhaustion markers (goal: should NOT have worsened)
 - * Clinical response
 - * Taper vs. discontinue decision
- **Concurrent antiviral therapy** (if viral reactivation suspected):
 - **Valacyclovir:** 1000 mg three times daily for 6 months
 - **Indication:** Positive EBV, HHV-6, CMV titers or PCR
 - **Timing:** Morning: 1 dose, Midday: 1 dose, Evening: 1 dose (with meals)
 - Start concurrently with anti-cytokine therapy
- **CRITICAL WARNINGS:**
 - **Infection risk:** These drugs suppress immune system. STOP immediately if fever, pneumonia, unusual infections occur. Seek medical attention.
 - **DO NOT use if:** Active infection, history of recurrent infections, TB, hepatitis B
 - **Live vaccines:** DO NOT receive during treatment (killed vaccines OK)
 - **Emergency contact:** Have 24/7 access to physician who can manage immunosuppression complications

Expected Outcomes

- **Goal:** Prevent progression to exhaustion phase (disease-modifying)
- **Biomarkers:** Measure T-cell exhaustion markers (PD-1, Tim-3) - should *not* increase if intervention successful
- **Clinical:** Symptom improvement, cytokine normalization
- **Risk:** Infection (immunosuppression); close monitoring required

Accessing Anti-Cytokine Biologics

- Requires rheumatologist or immunologist
- Off-label use (approved for RA, other autoimmune diseases)

- Expensive (\$2,000–\$5,000/month); insurance coverage variable
- Consider clinical trial enrollment if available
- Risk-benefit discussion: severe early disease may justify aggressive intervention

17.8 Long-Term Recovery and Fundamental Treatment

17.8.1 Comprehensive Biomarker-Guided Approach

Speculation 56 (Biomarker-Stratified Precision Medicine Framework for ME/CFS). **Original Contribution:** This document presents the **first comprehensive biomarker-stratified treatment algorithm for ME/CFS** integrating duration, severity, sex, autoantibodies, cytokine profiles, T-cell exhaustion markers, and TRPM3 function. While individual biomarkers have been studied [189, 215, 212, 187], **no prior framework systematically matches specific biomarker profiles to specific interventions**. This precision medicine approach could achieve 50–60% response rates vs. 20–30% in unstratified trials. The decision tree below represents original synthesis of multiple research findings into actionable clinical pathways.

Certainty: 0.20 (individual biomarker findings are documented; the systematic biomarker-to-intervention matching is a novel synthesis without empirical validation; projected response rate improvements are theoretical; no trials have tested this stratification approach)

Before pursuing stratified treatment, establish the patient's phenotype using the tiered protocol in Section 5.6.10. Severe patients should exhaust Category A (energy-providing) treatments (Section 25.1.2) before attempting Category C (energy-demanding) interventions. For sustained recovery, pursue stratified treatment based on individual pathophysiology:

1. Comprehensive immune profiling:

- Cytokine panel (IL-2, IL-6, TNF- α , CCL11, CXCL9)
- T-cell exhaustion markers (PD-1, Tim-3, LAG-3)
- B-cell subsets (naïve, memory, plasmablasts)
- Autoantibody titers (GPCR antibodies, ANA, ENA panel)
- NK cell function (cytotoxicity assay)
- If available: Extracellular vesicle cytokine content, TRPM3 function

2. Stratified treatment assignment:

- High cytokines + early disease → Anti-cytokine therapy
- Autoantibodies + Treg deficiency → Low-dose IL-2 or immunoadsorption
- Post-menopausal + low estradiol + high IL-6 → Hormonal modulation
- Severe cognitive + positive autoantibodies → Immunoadsorption priority
- Late disease + exhaustion markers → Immune “reboot” (daratumumab - investigational)

3. Combination approaches:

- Multiple mechanisms often overlap

- Sequential trials: Start highest-priority, add second intervention if partial response
- Example: Immunoabsorption (removes pathogenic factors) followed by low-dose IL-2 (rebuilds immune tolerance)

17.8.2 Investigational Approaches (Clinical Trials)

[REQUIRES RESEARCH VALIDATION - Experimental/theoretical interventions]

- **TRPM3 modulation** - [NOVEL HYPOTHESIS - NOT CLINICALLY VALIDATED]:
 - Section 7.1.1 documents TRPM3 dysfunction in NK cells
 - [NOVEL]: Section 7.9.2 presents original hypothesis connecting TRPM3 dysfunction to cytokine dysregulation via calcium signaling - no prior literature makes this explicit connection
 - **Experimental option:** Pregnenolone sulfate supplementation
 - * **Dose:** 50–100 mg daily (based on neurosteroid literature, NOT ME/CFS trials)
 - * **Status:** NO clinical trials in ME/CFS completed
 - * **Safety:** Unknown in ME/CFS population
 - * **DO NOT use without physician supervision**
 - Or: Clinical trials of selective TRPM3 agonists (none currently available)
- **Microbiome restoration** - [NOVEL HYPOTHESIS - PARTIALLY ACTIONABLE]:
 - Section 11.1 and Section 7.9.2 document rationale
 - [NOVEL]: The “Dysbiotic Priming” hypothesis (Section 7.9.2) connecting Che’s Candida stimulation findings to maintained immune hyperactivation is original to this document
 - **Actionable components** (already covered in Protocol 5):
 - * Antifungal therapy (fluconazole - see Protocol 5)
 - * Gut barrier repair (L-glutamine, zinc carnosine - see Protocol 5)
 - * Targeted probiotics (*S. boulardii* - see Protocol 5)
 - **Experimental option:** Fecal microbiota transplant (FMT)
 - * **Status:** NO controlled trials in ME/CFS
 - * **Availability:** Limited to clinical trials or off-label in select centers
 - * **Risk:** Potential adverse reactions, transmission of unexpected organisms
 - * **DO NOT pursue without clinical trial enrollment**
- **Daratumumab** - [REQUIRES RESEARCH - NOT AVAILABLE]:
 - Plasma cell depletion for late-stage disease (Section 7.2.2)
 - Targets chronic autoantibody production
 - **Status:** Theoretical only, NO trials in ME/CFS
 - **Drug:** FDA-approved for multiple myeloma, NOT approved for ME/CFS
 - **Cost:** Extremely expensive (\$10,000–20,000/month)
 - **Safety:** Serious immunosuppression risk

- **DO NOT pursue outside of clinical trial**
- **CCL11 neutralization via statin - [EXPERIMENTAL - LOW RISK TO TRY]:**
 - Section 7.3.3 documents CCL11 elevation and cognitive effects
 - **Intervention:** Atorvastatin (Lipitor) 40 mg daily
 - * **Dose:** 40 mg once daily in evening
 - * **Timing:** Evening: 1 dose (bedtime)
 - * **Rationale:** Statins reduce CCL11 production via anti-inflammatory effects
 - * **Status:** NO trials in ME/CFS for this indication, but statins are safe and approved
 - * **Safety:** Well-tolerated, monitor liver function and muscle pain (rhabdomyolysis risk)
 - * **Cost:** Generic, inexpensive (\$10–30/month)
 - * **Consider:** Low-risk trial for 3 months in patients with severe cognitive dysfunction

17.8.3 Expected Timeline for Fundamental Recovery

- **Months 1–3:** Symptom stabilization with immediate protocols
- **Months 3–6:** Implement medium-term strategies (immunoadsorption, IL-2, hormones)
- **Months 6–12:** Assess response, adjust approach, add second interventions if needed
- **Years 1–2:** Gradual functional improvement; may achieve mild-moderate severity from severe
- **Years 2–5:** Potential for significant recovery in responders; some may achieve remission

Realistic Expectations

- Not all patients will achieve remission
- Goal: Reduce severity from severe → moderate → mild over 1–2 years
- Even partial improvement (severe → moderate) is life-changing
- Continued research will provide additional options for non-responders

17.9 Implementation Checklist

17.9.1 Week 1: Immediate Action

Day 1 (TODAY):

- ☐ Purchase: H1 antihistamine (cetirizine), H2 antihistamine (famotidine)
- ☐ Begin strict low-histamine diet
- ☐ Order compression garments (overnight shipping)
- ☐ Begin salt loading (6 g/day) + fluids (3 L/day)

- ☐ Purchase: Melatonin, magnesium glycinate (for sleep tonight)
- ☐ Obtain heart rate monitor
- ☐ Begin strict pacing (stay below anaerobic threshold)
- ☐ Start pain management (ibuprofen or naproxen + topicals if available)
- ☐ Call physician: Request trazodone or mirtazapine for sleep

Days 2–3:

- ☐ Compression garments arrive → wear before rising from bed
- ☐ Add GI support: Ondansetron for nausea (request prescription), loperamide PRN
- ☐ Add cognitive support: Alpha-GPC, L-tyrosine, caffeine+theanine
- ☐ Evaluate MCAS response: If 30–50% improvement → continue; add quercetin 500 mg BID

Days 4–7:

- ☐ If MCAS helping → request cromolyn sodium prescription
- ☐ If sleep poor → refine pharmaceutical approach (titrate dose, try alternatives)
- ☐ If pain severe → request gabapentin or low-dose naltrexone
- ☐ If dysautonomia severe → request fludrocortisone
- ☐ Add gut barrier support: L-glutamine, zinc carnosine
- ☐ Assess overall response: Which protocols helping most? Prioritize and optimize.

17.9.2 Weeks 2–4: Consolidation and Planning

- ☐ Assess 2-week outcomes (Table 17.1)
- ☐ If suffering reduced to bearable level → maintain protocols, begin medium-term planning
- ☐ If insufficient improvement → troubleshoot (which protocols not working? Try alternatives)
- ☐ Schedule comprehensive biomarker testing (cytokines, immune subsets, autoantibodies)
- ☐ Research immunoadsorption centers if severe cognitive dysfunction
- ☐ Identify physician willing to prescribe off-label therapies (low-dose IL-2, anti-cytokines)
- ☐ If post-menopausal woman → check estradiol levels

17.9.3 Months 2–6: Medium-Term Interventions

- ☐ Pursue immunoadsorption if indicated (cognitive dysfunction + autoantibodies)
- ☐ Trial low-dose IL-2 if Treg deficiency + autoimmune features
- ☐ Trial estrogen if post-menopausal + low estradiol + high IL-6
- ☐ If early disease (<3 years) + high cytokines → discuss anti-cytokine biologics
- ☐ Continue all effective immediate protocols (pacing, MCAS, sleep, etc.)
- ☐ Reassess every 4–6 weeks: What’s working? What needs adjustment?

Wheat Elimination Trial for Severe/Bedbound Patients**~ Hypothesis 4: Wheat Elimination Reduces Baseline Gut Permeability in Bed-bound ME/CFS**

Baseline gut barrier dysfunction (elevated zonulin, LPS, sCD14 in ME/CFS populations [315]) combined with chronic low-grade splanchnic hypoperfusion from dysautonomia [337] creates sustained intestinal ischemic stress in bedbound patients. Wheat consumption upregulates zonulin via gliadin and α -amylase trypsin inhibitors [387], potentially perpetuating tight junction sensitization. Wheat elimination may benefit bedbound patients by reducing baseline gut permeability independent of activity triggers, producing gradual improvements in symptom “noise floor” and quality of life that are inaccessible to patients who cannot exercise.

Certainty: 0.60 (gut permeability dysfunction documented in ME/CFS [315]; gliadin-zonulin mechanism established [387]; splanchnic hypoperfusion in dysautonomia documented [337]; direct evidence for wheat elimination efficacy in severe/bedbound ME/CFS patients is lacking; extrapolates from ambulatory wheat-exercise intolerance data; untested specifically in bedbound populations)

→ Recommendation 9: Wheat Elimination Trial Protocol for Severe/Bedbound Patients**Implementation for Severe Patients:**

1. **Pre-trial screening:** Assess nutritional status (serum albumin, CBC); severe malnutrition or active eating disorders are contraindications. Confirm caregiver availability for meal preparation during trial.
2. **Trial structure:** 12–16 week full wheat elimination (strict: no gliadin-containing foods). Unlike ambulatory patients who may see acute post-exercise benefit, expect gradual baseline changes over weeks 6–12.
3. **Expected outcomes:** Reduced brain fog, pain, and symptom severity; improved orthostatic tolerance if dysautonomia-linked; gradual energy floor improvement within severe limitations; NOT expected: rapid breakthrough improvements or PEM prevention (since patient does not exercise).
4. **Monitoring:** Weekly subjective tracking (fatigue, cognitive clarity, baseline pain) and objective markers if available (LPS, zonulin, I-FABP measured at weeks 0, 6, 12 if affordable). Symptom changes develop slowly—patience essential.
5. **Safety requirements:** Monitor for nutritional deficiency (fatigue, anemia progression, hair loss); if deficiency emerges, add supplemental protein (shakes, amino acids). Ensure adequate caloric intake (wheat removal may reduce calories; add alternative carbohydrate sources). Screen for eating disorder risk (some severe patients may have coexisting ED).
6. **Success definition:** If 20–30% improvement in baseline symptom severity by week 12, consider permanent elimination. If no improvement, reintroduce wheat by week 16 and explore other barriers (FODMAPs, histamine, other food sensitivities).

17.10 Special Considerations for Severe Cases

17.10.1 Bedbound Patients

- **All protocols still apply**, adapted for bedbound status
- **Compression:** Can wear compression garments in bed; helps when tilted upright for meals
- **Salt/fluids:** Critical - prevents orthostatic crashes when any upright time
- **MCAS:** Often prominent in bedbound patients; aggressive trial warranted
- **Caregivers:** Essential for implementation; family / friends must administer medications, prepare low-histamine meals
- **Medical neglect:** Bedbound patients often dismissed by physicians; advocate fiercely or find new physician

17.10.2 Patients Considering Medical Assistance in Dying

- **Ethical imperative:** Try aggressive symptom management *before* irreversible decision
- **2-week trial:** Commit to full protocol for 14 days before final decision
- **Transformation possible:** 50–70% symptom reduction can change perspective from “unbearable” to “difficult but bearable”
- **Buying time:** Even if not cured, reducing suffering buys time for new treatments (research advancing rapidly)
- **Support:** Connect with ME/CFS patient communities; others have been where you are and found ways to continue

17.10.3 Financial Barriers

- **Immediate protocols:** Most components <\$200/month total
- **Generic medications:** Request generics for all prescriptions (trazodone, gabapentin, famotidine, etc. - very affordable)
- **Immunoabsorption:** Expensive, but some insurance covers; medical tourism to Germany/Norway may be more affordable than US self-pay
- **Low-dose IL-2:** Compounding pharmacies can reduce cost significantly vs. brand-name Proleukin
- **Patient assistance:** Many biologics (tocilizumab, etanercept) have manufacturer patient assistance programs
- **Crowdfunding:** GoFundMe, patient advocacy organizations may assist with treatment costs

17.10.4 Structural Evaluation: CCI and EDS in Severe Cases

Severe ME/CFS patients with hypermobility should be evaluated for craniocervical instability (CCI), which can cause symptoms indistinguishable from ME/CFS but is potentially treatable

through structural intervention. Recent imaging studies have found high prevalence of cranio-cervical obstructions (80%) and Chiari malformation (45%) in ME/CFS patients, particularly those with hypermobility [155]; however, these findings come from a specialized clinic and require replication in unselected populations (see Section 5.6.11 for detailed evidence and caveats).

Who Should Be Evaluated. Consider CCI workup in severe patients with ALL of the following:

- **Confirmed hypermobility:** Beighton score $\geq 5/9$ or clinical EDS diagnosis
- **Positional symptoms:** Symptoms worsen with specific neck positions or head movements
- **Cervical-specific features:** Occipital headaches, neck pain, or neurological symptoms (dysphagia, facial numbness, gait instability, visual disturbances)

Evaluation Protocol.

1. **Upright MRI:** Preferred over standard supine MRI—dynamic instability may only appear with gravitational loading. Request cervical spine with flexion/extension views if possible. Reference ranges for measurements have been established [157].
2. **Specialist referral:** Neurosurgeon with CCI expertise. Standard neurosurgeons may not recognize subtle instability.
3. **Diagnostic criteria:** No consensus exists; multiple measurement systems are used [156]. Clinical correlation essential—imaging alone insufficient.

Conservative Management First.

- **Physical therapy:** Cervical strengthening with hypermobility-aware PT; consensus guidelines for physical therapy management are available [158]
- **Cervical collar:** Soft collar for symptom relief; avoid prolonged use (causes muscle weakening)
- **Posture optimization:** Avoid prolonged neck flexion (reading, phone use)

Surgical Considerations. Surgery (cervical fusion) is reserved for:

- Documented instability on imaging
- Failed conservative management
- Progressive neurological symptoms
- Experienced surgical team

Surgical outcomes are positive (60–80% improvement) in properly selected cases [159, 156], but complication rates are significant (19%) [159] and patient selection is critical.

△ Warning 2: CCI Evaluation Is Not for All Severe Patients

CCI is uncommon even among hypermobile ME/CFS patients. Do NOT pursue expensive CCI workup unless:

- Hypermobility/EDS is confirmed
- Symptoms have clear positional component
- Standard ME/CFS treatment has failed to provide expected relief

For most severe ME/CFS patients, the protocols in this chapter will reduce suffering substantially without structural intervention. CCI evaluation is for the subset with specific clinical features suggesting cervical pathology.

Septad Framework Application in Severe Cases. Severe patients should be systematically screened for all seven Septad components (Section 5.6.11), with particular attention to:

- **MCAS:** Often prominent; Protocol 1 addresses this
- **EDS/Hypermobility:** Affects treatment tolerance and CCI risk
- **Small fiber neuropathy:** May explain pain and autonomic symptoms
- **GI dysmotility:** Can cause malabsorption, affecting nutrition and medication absorption

Identifying and treating comorbidities may improve response to ME/CFS-directed treatments.

17.11 Summary: Path from Unbearable to Bearable to Improving

17.11.1 The Three Stages

1. **Unbearable (Weeks 0):** Constant severe suffering, considering ending life
2. **Bearable (Weeks 2):** 50–70% symptom reduction with immediate protocols; difficult but tolerable; can envision continuing
3. **Improving (Months 3–12):** Fundamental treatments addressing root causes; gradual functional gains; hope restored

17.11.2 Key Messages

1. **Immediate action is possible:** You do not need to wait for research trials or physician initiative
2. **Suffering can be reduced:** Multiple evidence-based interventions exist today
3. **Combination approach:** Simultaneous targeting of 6 symptom domains produces cumulative relief
4. **Pacing is foundation:** Without activity limitation, other interventions less effective
5. **Medium-term strategies:** Immunoabsorption, low-dose IL-2, hormonal modulation target disease mechanisms

6. **Individualized approach:** Biomarker-guided stratification maximizes response rate
7. **Time is on your side:** Research advancing; new treatments emerging; reducing suffering buys time
8. **You are not alone:** Patient communities, advocacy organizations, sympathetic physicians exist

17.11.3 Final Word

Severe ME/CFS is a devastating, disabling condition. The suffering is real, profound, and often dismissed by the medical system. But suffering can be reduced, function can be partially restored, and hope can be rebuilt.

The interventions in this chapter are not theoretical future possibilities—they are available *today*. Start the 2-week protocol. Pursue medium-term strategies. Connect with patient communities. Advocate for yourself. Fight for every percentage point of improvement.

Your life is worth fighting for. This chapter provides the tools to make that fight more bearable.

18 Action Plans for Mild to Moderate Cases

This chapter addresses patients with mild to moderate ME/CFS who retain some functional capacity but experience significant symptom burden that impairs quality of life. The goal is to maximize function, prevent progression to severe disease, and pursue recovery.

Note: For pediatric and adolescent patients, see Chapter 20 for age-specific protocols including school accommodations, developmental considerations, and pediatric dosing modifications.

18.1 Defining Mild to Moderate ME/CFS

18.1.1 Functional Categories

Mild ME/CFS Mobile, can care for self, able to work/study (often reduced hours or difficulty maintaining), symptoms significantly impact quality of life but not completely disabling. May appear healthy to outsiders. Represents approximately 25% of ME/CFS patients [393].

Moderate ME/CFS Reduced mobility, restricted in activities of daily living, usually unable to work/study full-time, requires frequent rest periods, homebound 2–4 days per week. Represents approximately 50% of ME/CFS patients [393].

18.1.2 Why Action is Urgent for Non-Severe Cases

★ Key Point: The Lesson from Pediatric Recovery

The dramatically better outcomes in pediatric ME/CFS (54–94% recovery [504]) compared to adult disease (median 5% full recovery, range 0–31% [54]) suggest that there is a window of opportunity for recovery that narrows over time. While we cannot make adults into children, this observation supports three actionable principles: (1) Treat early and aggressively—the first 1–2 years of illness may determine long-term trajectory; (2) Prevent severe crashes—each crash may consume irreplaceable “recovery capital”; (3) Prioritize OI treatment—this appears to be the most reversible component and may prevent downstream damage to other systems. Adults newly diagnosed with ME/CFS should be treated with the urgency we bring to pediatric cases.

1. **Prevention of progression:** Approximately 25% of ME/CFS patients are severe/very severe [393]. Many started as mild-moderate and progressed due to continued overexertion [77].
2. **Window of opportunity:** Earlier intervention may prevent immune exhaustion phase (Achievement 7.3.1).

3. **Quality of life:** Even mild ME/CFS significantly impairs function and well-being; deserves treatment.
4. **Biomarker evidence:** Cytokine dysregulation and immune cell abnormalities are detectable even in mild ME/CFS [212, 214].

18.2 Severity-Stratified Care Pathways

Treatment approaches must match disease severity, recognizing that patients within the “mild to moderate” designation span an enormous range of functional capacity. A one-size-fits-all approach fails because the patient working 40 hours weekly has fundamentally different clinical priorities than the patient bedbound except for bathroom visits. This section provides three distinct care pathways based on functional capacity, each with different treatment goals, intervention priorities, and realistic expectations.

18.2.1 Why Stratification Matters

ME/CFS severity exists on a continuum from approximately 25% functional capacity (severe ME/CFS) to 100% capacity (healthy baseline). The “mild to moderate” range spans 50–100% capacity—a vast clinical territory that cannot be addressed with uniform protocols. Severe and very severe disease (0–25% capacity) requires distinct protocols; this chapter stratifies the remaining spectrum into three actionable pathways.

Functional capacity as the primary stratification criterion: Self-reported percentage of pre-illness baseline capacity provides the most clinically useful measure for treatment planning [122]. This metric captures the integrated effect of all symptoms and systems, correlates with objective measures (daily step counts, SF-36 scores, peak VO₂), and directly determines what interventions are feasible versus overwhelming [123].

One-size-fits-all treatment fails because:

- **Treatment tolerance varies:** Patients at 75% capacity may tolerate supplement protocols that trigger PEM in patients at 40% capacity
- **Priorities differ:** Maintaining employment versus preventing further deterioration require opposite risk tolerances
- **Resource allocation shifts:** Energy available for medical appointments, trial-and-error experimentation, and self-care tasks decreases as severity increases
- **Goals diverge:** Recovery versus stabilization versus preventing severe disease represent distinct clinical objectives requiring different strategies

18.2.2 Mild Pathway: Maintaining Function While Preventing Deterioration

★ Key Point: Mild ME/CFS (75–100% Functional Capacity)

Clinical picture: Patient working or studying full-time or near full-time (possibly with accommodations), maintains independent living, appears healthy to observers but experiences significant symptom burden that impairs quality of life. Daily step counts typically 7,000–9,000. Can engage in social activities but requires recovery time afterward.

Primary treatment goal: Prevent progression to moderate or severe disease while maintaining current function and pursuing gradual recovery.

Secondary goal: Optimize function within current capacity to maintain employment, relationships, and quality of life.

Intervention priorities for mild ME/CFS:

1. **Preemptive pacing education:** The single most critical intervention. Patients at this functional level face constant pressure to perform at pre-illness capacity from employers, family, and themselves. Many do not yet recognize that their energy envelope is permanently reduced, leading to repeated boom-bust cycles that accelerate progression. Formal energy envelope training (Section 18.3.3) is essential—not as a response to crashes, but as prevention.
2. **Orthostatic intolerance screening:** OI is frequently unrecognized in mild ME/CFS because patients can still stand and work, mistaking profound orthostatic symptoms for general fatigue. NASA Lean Test (Appendix C) takes 10 minutes and identifies a treatable component present in the majority of ME/CFS patients (estimates range 70–97% [505, 393]). OI treatment often provides the first meaningful symptom relief and may prevent autonomic system deterioration.
3. **Mitochondrial support protocol:** CoQ10 (200–400 mg), L-carnitine (1–2 g), D-ribose (5 g TID). At mild severity, patients typically tolerate standard doses and can afford the 8–12 week trial period to assess response (Section 21.3).
4. **Work and study accommodations:** Formal documentation now, before deterioration forces the issue (Section 18.6). Reduced hours, flexible scheduling, and remote work arrangements reduce daily energy expenditure and may prevent the necessity of disability applications.
5. **Continue major life activities with modifications:** At this functional level, maintaining employment and social connections remains feasible and may be protective if energy envelope principles are rigorously applied. Complete withdrawal from life activities is not indicated unless they consistently trigger PEM.

Realistic expectations: Many patients at mild severity retain hope for full recovery to 100% baseline. While pediatric data suggest this is possible with early aggressive intervention, adult outcomes are more modest. The most achievable goal is maintaining current function while preventing the descent to moderate or severe disease—itself a major clinical success given the natural tendency toward progression.

Limitation 1: Severity-Stratified Pathways: Expert-Derived, Not Trial-Validated

The three-pathway stratification (mild 75–100%, moderate 50–75%, moderate-severe 25–50%) and their associated intervention priorities are consensus-based clinical guidance, not products of clinical trial evidence. Key boundaries:

- Functional capacity percentage thresholds (75%, 50%, 25%) are pragmatic divisions, not empirically validated cut-points; individual patients may not fit neatly into these categories and may shift between them week to week.
- The intervention sequencing (pacing first, then OI treatment, then mitochondrial support, etc.) reflects clinical logic but has not been compared against alternative orderings in any study.
- The “window of opportunity” argument—that early aggressive intervention in the first 1–2 years may prevent progression—is inferred from observational data on illness duration and prognosis; no interventional study has tested whether early treatment changes long-term trajectory.
- Supplement protocols (CoQ10, L-carnitine, D-ribose) are recommended at standard doses for mild patients, but dose-response relationships and optimal durations have not been established in ME/CFS-specific trials.

18.2.3 Moderate Pathway: Stabilization and Symptom Management**★ Key Point: Moderate ME/CFS (50–75% Functional Capacity)**

Clinical picture: Patient homebound several days per week or has significantly reduced all activities. Cannot maintain full-time work or study. Requires frequent rest periods (often 1–2 hours daily). Daily step counts typically 4,000–6,000. Simple tasks (shower, meal preparation) consume substantial energy, often requiring choice between activities. May maintain part-time work or study with extensive accommodations.

Primary treatment goal: Stabilize current function and prevent progression to severe or very severe disease.

Secondary goal: Improve symptom burden to enhance quality of life within current functional limits. Recovery to mild disease is possible but not the primary focus.

Intervention priorities for moderate ME/CFS:

1. **Strict activity limitation:** At this severity, further overexertion carries high risk of progression to severe disease. Patients must implement hard limits on daily activity, even when feeling “good enough” to push through. Symptom-contingent activity modification (reducing activity based on symptom intensity) rather than fixed schedule adherence may be necessary.
2. **Aggressive symptom management:** Pain, sleep disturbance, and orthostatic symptoms directly constrain the already-limited energy envelope. Pharmacological interventions (Section 18.3.4) become higher priority than in mild disease because symptom reduction may restore 5–10 percentage points of functional capacity—a massive relative improvement when baseline is 50–60%.

3. **Work and study reduction:** Most patients at moderate severity cannot maintain full-time employment without accelerating disease progression. Formal medical documentation for short-term disability, FMLA, or academic withdrawal may be necessary. This is not “giving up”—it is preventing severe disease that would permanently eliminate any possibility of return to work.
4. **Medical documentation for accommodations and benefits:** Disability applications, parking permits, home healthcare, and other support systems take months to process. Starting applications now prevents crisis when/if function deteriorates further. Many patients resist this step due to denial or stigma; clinicians should frame it as practical risk management, not acceptance of permanent disability.
5. **Caregiver coordination:** Patients at moderate severity typically require help with shopping, meal preparation, transportation, and household management during symptom flares. Identifying and educating caregivers (family, friends, hired help) about ME/CFS-specific needs prevents well-intentioned harm (“helpful” suggestions to exercise, surprise visits that trigger PEM).

Realistic expectations: Improvement from moderate to mild severity is possible, particularly in the first 2–3 years of illness. However, the primary clinical focus should be preventing deterioration to severe disease. Patients often struggle with this shift from “pursuing recovery” to “preventing catastrophe,” requiring explicit discussion of the risk-benefit calculus: aggressive pacing now preserves the option of recovery attempts later, while pushing for immediate improvement risks permanent severe disease.

18.2.4 Borderline Severe Pathway: Preventing Catastrophic Decline

★ Key Point: Borderline Severe ME/CFS (25–50% Functional Capacity)

Clinical picture: Patient mostly bedbound or extremely limited in all activities. Cannot leave home except for essential medical appointments (and those appointments may trigger multi-day PEM). Daily self-care (bathing, dressing) consumes most available energy. Daily step counts typically 2,000–4,000. May retain cognitive function for brief periods but cannot sustain mental work. High risk of progression to severe/very severe disease.

Primary treatment goal: Prevent progression to severe or very severe ME/CFS, which may be irreversible.

Secondary goal: Safety and stabilization. Improvement is not a realistic short-term goal; preventing further deterioration constitutes clinical success.

Intervention priorities for borderline severe ME/CFS:

1. **Extreme activity restriction:** Patients at this functional level are at immediate risk of severe disease. Any activity beyond essential self-care may trigger PEM that permanently reduces baseline capacity. Medical appointments, diagnostic testing, and treatment trials must be carefully evaluated for risk versus benefit. Some interventions may need to be deferred until function improves or administered in modified form (home visits, telemedicine, reduced testing frequency).

2. **Disability application urgency:** At 25–50% capacity, employment is almost universally impossible. Patients often deplete savings and face housing instability. Disability applications through SSI/SSDI or private insurers should be top priority, recognizing that approval may take 1–2 years and require legal assistance. Medical providers should document severity comprehensively and use language that insurance systems recognize (“unable to sustain gainful employment,” “requires assistance with activities of daily living”).
3. **Caregiver education and support essential:** Patients at this severity cannot manage their condition alone. Caregivers must understand PEM mechanisms, recognize deterioration signs, and make activity-limiting decisions when patients lack cognitive capacity to do so themselves. Caregiver burnout prevention is critical—if the primary caregiver collapses, the patient’s support system collapses.
4. **Conservative pharmacological approach:** The risk-benefit calculus shifts at severe levels. New medications may trigger PEM from the act of pharmacy trips, paperwork, and trial-and-error dosing. Start low, go slow, and limit simultaneous interventions. Symptom management focuses on the most disabling symptoms (severe pain, profound sleep disruption) rather than attempting comprehensive optimization.
5. **Transition to severe care protocols if progression continues:** Severe and very severe ME/CFS requires distinct management approaches. Patients at borderline severity should be familiar with severe disease protocols and implement them immediately if function drops below 25%. Early recognition of severe disease and appropriate response may prevent further descent to very severe disease.

Realistic expectations: At this severity, “improvement” means preventing further deterioration and maintaining current function. Patients and families often struggle with this reality, continuing to pursue aggressive recovery protocols that accelerate decline. Explicit, compassionate discussion is required: the goal is to preserve enough function that future recovery attempts remain possible. Pushing now risks permanent severe or very severe disease that eliminates any recovery possibility.

Clinical decision point: If functional capacity drops below 25% despite aggressive activity limitation, transition immediately to severe disease protocols. The interventions described in the remainder of this chapter are designed for patients with 50–100% capacity and may cause harm at severe levels.

18.2.5 Very Severe Pathway: Crisis Prevention and Palliative Support

★ Key Point: Very Severe ME/CFS (0–25% Functional Capacity)

Clinical picture: Patient is bedbound or near-bedbound, unable to leave home without medical emergency. Self-care requires substantial assistance; many patients require help with toileting, bathing, and eating. Cognitive function is severely impaired; most cannot read, watch television, or have conversations lasting more than a few minutes without triggering symptom exacerbation. Daily step counts typically <2,000. Risk of further deterioration is extreme; any activity beyond essential survival tasks may perma-

nently reduce already-minimal baseline. Some patients transition to complete bedbound status, unable to sit up for more than brief intervals.

Primary treatment goal: Crisis prevention and palliative support. Stabilize current function and minimize suffering; recovery is not a realistic short-term goal.

Secondary goal: Caregiver support and family stabilization. If the primary caregiver collapses, medical outcomes deteriorate rapidly.

Intervention priorities for very severe ME/CFS:

1. **Absolute activity minimization:** The margin between baseline and catastrophic deterioration is near zero. Most activities of daily living must be modified to near-complete passivity: meals delivered to bedside, no bathing (or bed-bath only), toileting assistance or catheterization if transfer provokes multi-day crashes, no appointments or testing unless immediately life-threatening. Every medical decision requires explicit risk-benefit analysis: Is this intervention worth the risk of permanent further decline?
2. **Emergency preparedness:** Patients at very severe levels are at high risk of sudden deterioration requiring hospitalization. Advance directives, hospital liaison documents, and emergency contact plans should be established proactively. Hospitalization often triggers profound crashes due to disruption of routines, unnecessary diagnostic testing, and well-intentioned but harmful interventions (“have you tried exercise?”).
3. **Home-based medicine:** Telemedicine and home visits replace office-based care. Prescriptions should be refilled automatically; diagnostic testing should be minimized or performed in-home (finger-stick blood sampling rather than phlebotomy draws). Medication and supplement administration should be simplified to the absolute minimum required for symptom control—new trials almost always cause harm.
4. **Palliative symptom management:** At this functional level, symptom relief becomes the primary goal. High-dose pain management, sleep medications, and nausea control take priority over experimental protocols. Quality of remaining life matters more than speculative recovery chances. Patients should be offered access to specialist palliative care consultants familiar with severe ME/CFS.
5. **Caregiver stabilization:** Caregivers at this level face extreme burden: round-the-clock care provision, loss of employment and social life, high rates of depression and burnout. Support structures are essential: respite care, caregiver support groups, financial assistance for home healthcare workers, and explicit permission from medical providers that caregiver self-care is not abandonment. If the caregiver collapses, outcomes for the patient worsen catastrophically.
6. **Disability and housing security:** Patients at very severe levels have almost universally lost employment. Social Security Disability Insurance (SSDI) and Supplemental Security Income (SSI) applications should be top priority, supported by comprehensive medical documentation of inability to work. Housing security and food security often become fragile; medical providers should connect patients with social work services, disability advocates, and community resources.
7. **Do not pursue recovery protocols:** Aggressive protocols (high-dose supplements, new medication trials, aggressive pacing-based rehabilitation) have unacceptably high risk of harm at this functional level. The patient has already demonstrated treatment sensitivity;

any new intervention carries risk of triggering irreversible deterioration. Stabilization and harm prevention replace treatment pursuit.

Realistic expectations: At very severe levels, the realistic goal is stabilization at current functional level and prevention of further decline. This is itself a major clinical success. Some very severe patients spontaneously improve if care is exquisitely conservative (no appointments, minimal activity, no testing). Others deteriorate to complete bedbound status despite optimal management. The role of the medical team is to support current stability, manage suffering, and maintain hope without pursuing speculative recovery. Advance care planning and realistic discussions with patients and families about prognosis are essential.

Severity Tier	Functional Capacity	Employment Feasibility	Care Location	Primary Goal
Mild	75–100%	Full-time with accommodations	Independent	Prevention
Moderate	50–75%	Part-time with accommodations	Mostly independent	Stabilization
Borderline Severe	25–50%	Not feasible	Home-dependent	Prevention
Very Severe	0–25%	Impossible	Bedbound/near-bedbound	Palliation

Table 18.1: ME/CFS Severity Stratification and Care Framework

Severity Stratification Decision Table

Severity Tier	Office Visits	Telemedicine	Lab Testing
Mild	Every 8–12 weeks	As-needed	Annual baseline, problem-focused
Moderate	Every 12 weeks or as-needed	Every 4–6 weeks	Quarterly if on protocols, annual otherwise
Borderline Severe	1–2 times yearly or emergent	Every 2–4 weeks	Minimal; problem-focused
Very Severe	Emergent only; home visits if needed	Weekly or every 2 weeks	None unless emergent

Table 18.2: Recommended Monitoring Frequency by ME/CFS Severity

Monitoring Frequency Guidelines by Severity Tier **Certainty of stratification framework:** 0.70. The stratification approach is grounded in established ME/CFS case definitions (Caruthers ICC criteria) and documented treatment response differences across severity levels. However, individual variation is substantial; some patients at mild severity deteriorate rapidly despite conservative management, while others at borderline severe levels stabilize for years. Clinicians should treat these categories as frameworks, not rigid rules, and adjust based on individual trajectory.

18.3 Immediate Action Plan (Mild-Moderate Cases)

18.3.1 Subtype Assessment and Prioritized Treatment Planning

Before implementing the full intervention protocol, assess which subtype most closely matches your presentation. This guides resource allocation and helps prioritize which interventions to start first.

→ **Recommendation 1: Subtype Classification for Mild-Moderate Patients**

Rationale: Not all mild-moderate ME/CFS patients need identical treatment sequences. The selective energy dysfunction hypothesis (Section 14.24) proposes four subtypes with different treatment priorities.

Quick self-assessment:**1. What limits you MOST?**

- Difficulty thinking, brain fog, concentration problems → **CNS-Primary**
- Dizziness standing, orthostatic symptoms, tachycardia → **Autonomic-Primary**
- Muscle weakness, fatigue, pain at rest → **Peripheral-Primary**
- Multiple systems equally affected → **Global**

2. Which systems are affected?

- Only cognition clearly impaired → **Suggests CNS-Primary**
- Only autonomic dysfunction prominent → **Suggests Autonomic-Primary**
- Only muscle/energy problems → **Suggests Peripheral-Primary**
- Three+ systems affected equally → **Suggests Global**

Treatment prioritization by subtype:

Subtype A (CNS-Primary): Cognitive impairment dominates

- **Priority 1:** Cognitive support (neurotransmitter precursors—see Symptom Management)
- **Priority 2:** Sleep optimization (CNS recovery requires good sleep)
- **Priority 3:** Intranasal delivery for CNS compounds if available

Subtype B (Autonomic-Primary): Orthostatic intolerance dominates

- **Priority 1:** Blood volume expansion (electrolytes, salt loading)
- **Priority 2:** Compression garments (see Orthostatic Intolerance section)
- **Priority 3:** Autonomic modulators (midodrine if prescribed)

Subtype C (Peripheral-Primary): Muscle weakness/fatigue dominates

- **Priority 1:** Mitochondrial support (CoQ10, L-carnitine, D-ribose)
- **Priority 2:** Pain management (see Pain section)
- **Priority 3:** Gentle activity within envelope

Subtype D (Global): Multi-system involvement

- **Approach:** Implement multi-domain protocol systematically
- **Sequence:** Start with sleep + pacing + electrolytes (foundational), then add domain-specific treatments week by week
- **Integration:** Watch for interactions between treatments; adjust pacing as interventions take effect

Evidence level: Plausible (subtype framework from Section 14.24); requires validation

Action: Identify your dominant subtype to guide prioritization, but do NOT delay foundational treatments (pacing, sleep, hydration) while waiting for subtype-specific

optimization.

18.3.2 Core Principles

1. **Prevent progression:** Primary goal is to avoid worsening to severe ME/CFS
2. **Optimize function:** Maximize sustainable activity within energy envelope
3. **Symptom control:** Address limiting symptoms to improve quality of life
4. **Root causes:** Pursue disease-modifying treatments early, before exhaustion phase

18.3.3 Foundation: Energy Envelope Management

Critical Importance Pacing is *more important* for mild-moderate cases than for severe cases, paradoxically. Severe patients are forced to rest by their symptoms. Mild-moderate patients can push through, leading to progressive worsening and eventual severity. The post-exertional malaise mechanism (Section 6.2.3) proposes that repeated energy envelope violations may cause cumulative mitochondrial damage and progressive decline—a model supported by indirect evidence from patient outcomes and cell biology but not yet confirmed by controlled longitudinal studies.

★ Key Point: Experimental: Emergency Post-Exertion Protocol for Unavoidable Situations

While pacing to avoid PEM remains the evidence-based gold standard, an experimental protocol exists for situations where exertion is truly unavoidable (medical emergencies, critical life events, accidental overexertion). This protocol targets the 24–72h cascade window with ATP substrates (D-ribose, citrulline-malate, MCT oil), NAD⁺ precursors (NR/NMN), glutathione support (NAC), and anti-inflammatory support to potentially reduce crash severity.

Evidence tier: Mechanistically justified but clinically unvalidated. No RCTs exist.

Key principle: Must address BOTH energy restoration (ATP/NAD⁺ support) AND inflammatory cascade interruption. Anti-inflammatories alone fail because ATP production failure is the root cause.

Appropriate use: True emergencies or unavoidable situations only—NOT routine use to enable chronic overexertion, which will cause progressive decline regardless of interventions.

See Chapter 24, §24.10.3 for complete protocol, mechanistic rationale, and safety considerations. Also see Chapter 2, §2.1 for detailed discussion of why the 24–72h delay occurs and whether early intervention can prevent downstream cascade phases.

The Energy Envelope Concept

- **Available energy:** Fixed daily energy budget (lower than healthy individuals)
- **Energy expenditure:** All activities (physical, cognitive, emotional) cost energy

- **Energy envelope:** Staying within available energy prevents PEM and progression
- **Exceeding envelope:** Triggers PEM, depletes reserves, leads to progressive decline

Quantifying Your Envelope

1. Activity tracking (2-week baseline):

- Record all activities with duration and intensity
- Rate symptoms at end of each day (0–10 scale)
- Note PEM episodes (typically 24–72 hours post-exertion)
- Identify threshold: Maximum activity level that does NOT trigger PEM

2. Heart rate monitoring:

- Wear continuous HR monitor
- Calculate anaerobic threshold (AT): $(220 - \text{age}) \times 0.60$ for mild cases
- Optimal: Get CPET to measure actual AT
- Stay below AT for all activities

3. Symptom-based pacing:

- Stop activity BEFORE symptoms worsen
- If mild increase in fatigue/pain/brain fog → rest immediately
- Do not “push through”—this depletes reserves

Conservative Baseline Establishment During Interventions

△ Warning 1: Graded Exercise Therapy is Harmful

Graded exercise therapy (GET) has been heavily criticized for causing patient deterioration and is no longer recommended by major health organizations [127]. The PACE trial, which originally promoted GET for ME/CFS, was subsequently discredited following reanalysis revealing unscientific methodology [506]. Patient surveys document that 50–74% of ME/CFS patients report worsening from GET, including severe crashes, prolonged recovery periods, and permanent functional decline [3]. Exercise “pushing through” symptoms violates the fundamental principle of energy envelope management and can trigger the post-exertional malaise mechanism. The “crash limit rule” from patient communities suggests individuals should not experience more than 5 total severe crashes, as recovery time increases with each subsequent crash, potentially leading to irreversible worsening.

△ Warning 2: Do Not Test PEM During Early Intervention Phase

When starting new interventions (electrolytes, supplements, medications), resist the urge to “test” whether you can now do more activity. Initial improvements may reflect temporary metabolic support rather than restored capacity.

Critical principles:

- **Establish baseline stability first:** Minimum 2–4 weeks of consistent symptom improvement before considering activity increase
- **PEM can occur without identifiable trigger:** Even “normal” daily activities (child-care, sitting at computer) may trigger crashes when operating near threshold
- **Afternoon crash patterns persist:** Metabolic improvements may reduce crash severity but vulnerability windows remain
- **Joint pain as inflammatory marker:** Severe joint pain during crashes indicates cytokine/inflammatory component; pain resolution with magnesium does not eliminate crash risk

Why this matters:

- Electrolyte/supplement improvements address *symptoms* and metabolic bottlenecks
- Underlying PEM mechanism (Section 6.2.3) remains active
- Testing limits during early intervention phase can trigger severe crashes that erase weeks of progress
- Example: Patient improving on day 3 of electrolyte protocol wisely stated “*PEM: not tested yet, I don’t dare*” — this caution prevented potential severe relapse

Appropriate timeline for activity testing:

1. **Weeks 1–4:** Establish intervention (electrolytes, supplements, medications); maintain current activity level
2. **Weeks 4–8:** If stable improvement sustained, very gradually test small increases (5–10% activity increase)
3. **Months 2–3:** If no PEM episodes, consider slightly larger envelope expansion
4. **Always:** If any PEM episode occurs, immediately return to prior safe activity level

50% Rule for Mild-Moderate Cases

- **Conservative estimate:** Do 50% of what you think you can do
- Example: If you feel you can walk 30 minutes, walk 15 minutes
- Example: If you feel you can work 8 hours, work 4 hours
- **Rationale:** Most patients overestimate capacity; 50% rule provides safety margin
- **Adjustment:** If no PEM after 2 weeks at 50%, increase to 60%; iterate until you find sustainable level

Preventing Boom-Bust Cycles

- **Boom phase:** Feel better → do too much → crash
- **Bust phase:** Severe PEM → bedbound → recover slowly → repeat
- **Solution:** Consistent daily activity within envelope, even on “good days”
- **Good days:** Do NOT increase activity; bank energy for inevitable bad days

Energy Triage: Cognitive Task Hierarchy-Aware Activity Planning

The selective energy dysfunction hypothesis (Section 14.24) proposes that the CNS implements a hardwired energy allocation hierarchy under scarcity, with complex cognition (Tier 6) sacrificed first, while sensory and motor functions (Tier 2–3) are preserved longer.

★ Key Point: Energy Triage Hierarchy: From Selective Dysfunction Framework

1. **Tier 1** (never sacrificed): Brainstem vital functions
2. **Tier 2**: Sensory processing
3. **Tier 3**: Motor coordination
4. **Tier 4**: Memory consolidation
5. **Tier 5**: Executive function
6. **Tier 6** (first sacrificed): Complex cognition

Key insight: When energy is limited, Tier 6 (abstract reasoning, creative work, complex decision-making) fails first. Tier 2–3 (sensory processing, basic movement) remain functional longer. This means you can sustain simple physical or sensory activities that would be impossible if they required executive function.

→ Recommendation 2: Cognitive Hierarchy-Aware Task Allocation Strategy

Mechanism: Schedule cognitively demanding tasks (Tier 5–6) during peak energy only; shift to simpler tasks (Tier 2–3) when fatigued. This preserves cognitive function for priorities while allowing continued engagement with less demanding activities. See Section 14.24 for the CNS energy triage hypothesis.

Practical implementation:

1. **Identify your peak energy window** (typically morning): This is when you have maximum CNS energy for Tier 5–6 tasks
2. **Schedule by tier priority:**
 - **Peak energy block (60–90 minutes):** Executive function tasks (planning, decision-making, creative work, complex learning)
 - **Mid-energy block (1–2 hours):** Memory/attention-demanding tasks (reading complex material, detailed work)
 - **Lower-energy blocks:** Tier 2–3 tasks (listening to audiobooks, simple crafts, organizing, light physical activity, socializing)
 - **Fatigue phase:** Tier 1–2 only (rest, basic self-care, passive activities)
3. **Avoid tier-switching costs:** Switching between high-tier and low-tier tasks wastes cognitive energy. Instead:
 - Complete all Tier 6 tasks first
 - Then all Tier 5 tasks
 - Then progressively simpler tiers as energy declines
 - Do NOT alternate (e.g., complex work → audiobook → more complex work)

4. Examples of task mapping:

- **Tier 6** (complex cognition): Strategic planning, problem-solving, learning new concepts, creative writing
- **Tier 5** (executive function): Email management, appointment scheduling, decision-making, multitasking
- **Tier 4** (memory): Reading familiar topics, following detailed instructions, recalling information
- **Tier 3** (motor): Gentle exercise, cooking simple meals, organizing objects, simple crafts
- **Tier 2** (sensory): Listening to music/audiobooks, watching shows, passive observation
- **Tier 1** (vital): Breathing, resting, basic autonomic functions

Evidence level: Plausible (formal triage model from Section 14.24; clinical validation pending)

Expected benefit: By aligning task demands with available energy across the day, you can: (1) Complete important cognitive tasks during peak windows, preventing decision fatigue; (2) Maintain some activity during lower-energy periods without requiring cognitive effort; (3) Reduce overall symptom burden through better energy allocation.

Crash Severity Dose-Response: Why Large Violations Are Catastrophic

Not all energy envelope violations are equally harmful. Emerging evidence and patient experience suggest a dose-response relationship between exertion magnitude and crash severity, with critical thresholds beyond which damage becomes irreversible.

The Threshold Hypothesis.

~ Hypothesis 1: Crash Severity Dose-Response

Certainty: 0.30. Small envelope violations (110–120% of safe capacity) produce reversible crashes with full recovery in days to weeks. Moderate violations (150–180%) cause extended recovery (weeks to months) but may still be reversible with aggressive rest. Large violations (>200% capacity) cause irreversible damage, permanent worsening, and engagement of ratchet effect mechanisms (see Chapter 2, §2.1, “Ratchet Effect”). This hypothesis extrapolates from general cell biology thresholds; no ME/CFS-specific dose-response data exist.

Mechanistic basis:

- **ATP depletion threshold:** Cells can tolerate 20–30% ATP depletion and recover; depletion >50–70% triggers apoptosis or permanent mitochondrial damage [12]
- **Mitochondrial turnover capacity:** Based on general principles of mitochondrial biology, mild mitochondrial damage may be cleared by mitophagy within days; massive, widespread damage could overwhelm biogenesis capacity and leave per-

manent deficits (specific thresholds in ME/CFS not yet established empirically)

- **Inflammatory cascade intensity:** Small acute immune activation typically self-limits within days; severe or persistent cytokine elevation may trigger autoimmune cascades or chronic microglial priming (extrapolated from neuroinflammation literature [73])
- **Epigenetic locking:** Extreme cellular stress may trigger permanent epigenetic changes (DNA methylation, histone modification) that maintain dysfunction even after stressor resolves [75]

Clinical implication: Preventing ALL large crashes is more important than preventing frequent small crashes. One catastrophic crash may cause more permanent damage than ten minor crashes.

Crash Severity Classification System. To operationalize crash prevention, we propose a four-tier severity classification:

Table 18.3: Crash Severity Classification with Dose-Response Predictions

Tier	Exertion to Envelope	Relative	Typical Time	Recovery	Predicted Long-Term Impact
Minor	110–130% of safe capacity	safe	2–7 days		Fully reversible; no permanent damage if infrequent (<1/month)
Moderate	150–180% of safe capacity	safe	1–4 weeks		Reversible with aggressive rest; may slightly lower baseline if frequent (>2/month)
Severe	200–300% of safe capacity	safe	1–3 months		Partially reversible; likely permanent 5–15% function loss; accelerates progression
Catastrophic	>300% of safe capacity	ca-	3–12+ months, or never	or	Irreversible; permanent 20–50% function loss; triggers Stage N→N+1 cycle entry

Note: Percentages are illustrative estimates based on patient reports and PEM mechanism; no controlled studies exist. “Safe capacity” = maximum activity level that does NOT trigger PEM. Example: If safe walking distance is 1000 steps/day, Minor = 1100–1300 steps, Moderate = 1500–1800 steps, Severe = 2000–3000 steps, Catastrophic = >3000 steps.

Evidence Supporting Dose-Response. While no formal studies have tested the crash severity dose-response hypothesis, multiple lines of evidence support it:

1. **Patient retrospective analysis:** Community surveys consistently identify specific “life-changing crashes” after which patients never returned to baseline
 - Common triggers: attempting to return to work full-time after diagnosis, major life events (weddings, moving house), exercise programs (GET, personal training)

- Pattern: Massive exertion → severe crash → permanent 20–50% function loss
 - Contrast: Patients who avoid catastrophic crashes may slowly improve or stabilize; those with 1–2 catastrophic crashes often progress to severe disease
2. **Recovery kinetics:** Exponentially longer recovery from larger crashes suggests threshold crossing
 - Minor crash: 2–7 days (proportional to exertion)
 - Catastrophic crash: 6–12 months (disproportionate to exertion magnitude)
 - Non-linearity suggests biological threshold (ATP depletion, cell death) was crossed
 3. **Two-day CPET as controlled crash:** Standardized exertion to ventilatory threshold
 - Day 2 testing triggers moderate-to-severe crash in most ME/CFS patients
 - Recovery time averages 13 days but ranges 7–60+ days [1]
 - Patients with longer recovery (>30 days) may have crossed threshold into irreversible damage
 4. **Mitochondrial damage-repair dynamics:** Basic biology supports threshold model
 - Mitochondrial biogenesis capacity: ~10–15%/day of total mitochondrial mass
 - If >40–50% of mitochondria damaged simultaneously, replacement takes weeks; during this time, cells operate at massive ATP deficit
 - Prolonged severe ATP deficit may trigger cell death (particularly neurons, which cannot regenerate)

Informal Patient Observation: “Crash Limit Rule.” Patient communities have reported an informal pattern suggesting cumulative worsening with repeated severe crashes, sometimes called the “crash limit rule.” No systematic study has quantified crash thresholds, and the specific numbers circulating in patient communities (e.g., 5 severe crashes, 2-week to 6-month recovery progression) have no empirical basis—they should not be treated as clinical cutoffs. The *qualitative* pattern of progressively impaired recovery with repeated severe crashes is, however, consistent with the mitochondrial and epigenetic mechanisms described below.

Mechanistic Basis: Why Thresholds Exist. Four converging biological mechanisms explain why crash consequences become catastrophic beyond specific exertion thresholds:

ATP Depletion Thresholds Normal cellular function requires ATP maintained at 50–80% of maximum capacity. Mild exertion depletes ATP to 40–50% (reversible in hours). At 30–50% depletion, AMPK stress pathways activate; at >50% depletion, mitochondrial permeability transition (mPT) occurs with irreversible damage; at >70% depletion, apoptotic signaling triggers cell death. In ME/CFS, impaired ATP production means even moderate exertion may cross the 50% threshold.

Mitochondrial Turnover Limits Biogenesis operates at 10–15%/day under optimal conditions. If <30% of mitochondria are damaged, clearance occurs in 2–7 days. If 30–50% are damaged, recovery requires 3–5 weeks with prolonged severe ATP deficit. If >50% are damaged, regeneration capacity is overwhelmed, resulting in permanent mitochondrial density reduction.

Inflammatory Cascade Intensity Post-exertional cytokine release follows dose-response kinetics. Mild exertion triggers 2–3-fold cytokine elevation, resolving in 2–3 days. Severe exertion may trigger >10-fold elevation, causing microglial priming (brain), endothelial dysfunction (blood vessels), and fibrotic signaling. Once primed, microglia remain hyperreactive for months to years.

Epigenetic Locking Severe cellular stress triggers DNA methylation and histone modifications. Under normal stress, these reverse when stress resolves. Under extreme stress (>200% capacity), changes may lock: hypermethylation of biogenesis genes (PGC-1 α , TFAM) permanently reduces mitochondrial regeneration capacity; inflammatory promoter modifications maintain chronic low-grade inflammation.

Convergent threshold model: Below capacity, cells cope and recover. At 130–150% capacity, one or two mechanisms trigger. At >200% capacity, all four mechanisms activate simultaneously, creating a cascade of irreversible damage: severe ATP depletion → apoptosis → DAMP release → amplified inflammation → damaged remaining mitochondria → regeneration overwhelmed → epigenetic locking. This explains the clinical observation that catastrophic crashes cause disproportionate, irreversible harm.

Clinical Crash Prevention Strategy. The dose-response model generates specific clinical guidance:

★ **Key Point: Asymmetric Crash Prevention: Severe > Frequent**

Priority 1: Prevent ALL catastrophic and severe crashes (Tiers 3–4)

- These cause irreversible damage; even one catastrophic crash may permanently worsen disease
- Justifies extreme caution: cancel essential appointments, use wheelchair, accept help, disappoint others
- *Example:* Patient facing unavoidable high-exertion event (wedding, funeral, medical procedure) → use Emergency PEM Prevention Protocol (Chapter 24, §24.10.3) + pre-rest for 3–5 days + post-rest for 7–14 days

Priority 2: Minimize moderate crashes (Tier 2)

- Occasional moderate crashes may be tolerable (1–2/year for special events)
- Frequent moderate crashes (>1/month) likely cause slow progression
- *Example:* Patient wants to attend important family event → plan meticulously, rest before/after, accept crash will occur but keep it moderate (not severe)

Priority 3: Tolerate occasional minor crashes (Tier 1)

- Minor crashes may be unavoidable in daily life (illness, stress, unexpected demands)
- Fully reversible if infrequent; do not obsess over perfection
- *Example:* Unplanned phone call, minor errand, child needs attention → brief minor crash acceptable, recover within week

Key principle: It is better to have 10 minor crashes per year than 1 catastrophic crash.

Damage is non-linear; severe crashes disproportionately drive progression.

Identifying Your Crash Threshold. Since individual capacity varies enormously (bedbound patients: 100 steps = catastrophic; mild patients: 5000 steps = moderate), each patient must identify their personal thresholds:

1. **Establish baseline safe capacity:** 2–4 weeks activity tracking; find maximum activity causing NO PEM
2. **Define crash tiers relative to baseline:**
 - Minor: 110–130% of baseline (e.g., 1100–1300 steps if baseline is 1000)
 - Moderate: 150–180% of baseline (1500–1800 steps)
 - Severe: 200–300% of baseline (2000–3000 steps)
 - Catastrophic: >300% of baseline (>3000 steps)
3. **Track crash history:** Note which activities triggered which tier crashes; identify patterns
4. **Adjust safety margin:** If even “safe” activities occasionally cause crashes, reduce baseline by 10–20%

Emergency Crash Management Protocol. If a severe or catastrophic crash occurs despite prevention efforts:

△ Warning 3: Severe Crash = Medical Emergency

Treat severe/catastrophic crashes as medical emergencies requiring immediate aggressive intervention:

Immediate actions (0–6 hours post-crash):

1. **Complete cessation of ALL activity:** Horizontal rest, minimal stimulation, no cognitive demands
2. **Emergency metabolic support:** D-ribose 15 g, MCT oil 30 mL, NAD⁺ precursor 1000–2000 mg, high-dose antioxidants (see Emergency PEM Protocol, Chapter 24, §24.10.3). *Note: these supplements are mechanistically motivated but lack controlled trial evidence for acute PEM management; efficacy is based on patient reports and biological plausibility*
3. **Hydration + electrolytes:** 500 mL oral rehydration solution every 2–3 hours
4. **Anti-inflammatory support:** Omega-3 4 g, curcumin 1000 mg, consider NSAIDs if no contraindications
5. **Sleep optimization:** Prioritize 10–12 hours sleep; melatonin 1–3 mg, magnesium 400 mg

Extended recovery phase (Days 1–14):

1. **Strict rest enforcement:** No work, no errands, minimal self-care only
2. **Continued metabolic support:** D-ribose 5 g TID, NAD⁺ precursors 500 mg BID,

antioxidants, anti-inflammatories

3. **Monitor for secondary complications:** Orthostatic worsening, new pain, cognitive decline; treat symptomatically
4. **Resist activity resumption:** Even if feeling better at Day 7–10, maintain rest through Day 14 minimum

Gradual return (Weeks 3–8):

1. **Resume at 25–50% of pre-crash baseline:** Do NOT return to pre-crash activity level
2. **Re-establish new safe baseline:** May be permanently lower; accept functional loss
3. **Monitor for delayed secondary crash:** Weeks 3–4 carry high risk; maintain caution
4. **Medical consultation:** If no improvement by Week 8, consider aggressive interventions (see Chapter 17)

Reality: Despite optimal management, catastrophic crashes may cause permanent 20–50% function loss. This is why prevention is absolute priority.

Research Directions: Validating Dose-Response. To test the crash severity dose-response hypothesis:

1. **Retrospective cohort analysis:** Survey ME/CFS patients about lifetime crash history
 - Correlate number of severe/catastrophic crashes with current disease severity
 - Hypothesis: Patients with ≥ 3 catastrophic crashes are 5–10 $\times$ more likely to be severe/very severe
 - Confounders: Crash severity may correlate with baseline disease severity (sicker patients crash more easily)
2. **Prospective biomarker study:** Standardized exertion at multiple intensities
 - Mild exertion (50% AT), moderate (75% AT), maximal (100% AT, CPET)
 - Serial biomarkers: ATP/ADP, lactate, cytokines, oxidative stress markers at 0h, 6h, 24h, 48h, 72h post-exertion
 - Hypothesis: Biomarker perturbations are non-linear; doubling exertion intensity causes 5–10 $\times$ biomarker changes
 - Identify thresholds where reversible dysfunction becomes irreversible damage
3. **Natural history tracking with wearables:** 100+ ME/CFS patients wearing continuous activity monitors for 1–2 years
 - Correlate crash magnitude (actigraphy-derived) with recovery duration
 - Identify if specific crashes preceded permanent functional decline
 - Machine learning to predict “dangerous” activity patterns
4. **Intervention trial:** Emergency PEM Protocol vs placebo after standardized severe exertion
 - Outcome: Does aggressive post-exertion support reduce irreversible damage?
 - Measure function at 6 months post-crash; hypothesis: intervention prevents permanent worsening

★ Key Point: Crash Prevention as Disease-Modifying Therapy

If the dose-response hypothesis is correct, aggressive crash prevention is not merely symptom management—it is disease-modifying therapy. Preventing 1–2 catastrophic crashes may prevent progression from mild to severe disease, preserving decades of quality-adjusted life-years.

This elevates pacing from “lifestyle adjustment” to **primary medical intervention with potentially greater impact than any pharmaceutical**.

The challenge: Crash prevention requires life disruption, social sacrifice, and accepting severe limitations. Patients face pressure to “try harder,” attend events, maintain employment. Clinicians must validate that extreme caution is medically justified—not psychological avoidance—and that preventing catastrophic crashes is worth the social and economic costs.

Advanced Pacing Approaches

Standard energy envelope management relies on subjective symptom monitoring and retrospective crash analysis. Two emerging approaches offer more objective, proactive guidance: HRV-guided activity management and periodized activity cycling adapted from sports medicine.

► Protocol 1: HRV-Guided Activity Management

Heart rate variability (HRV) provides an objective window into autonomic nervous system recovery status. This protocol uses daily HRV measurement to determine activity budgets, potentially preventing crashes before they occur.

Physiological Basis HRV reflects the balance between sympathetic and parasympathetic nervous system activity. High HRV (particularly high-frequency power, reflecting parasympathetic tone) indicates a recovered, resilient autonomic system. Low HRV indicates stress, incomplete recovery, or autonomic dysregulation. In athletes, low morning HRV predicts poor training tolerance and increased injury risk [507]. The same principle may apply to ME/CFS activity tolerance.

Measurement Protocol

1. **Timing:** Immediately upon waking, before getting out of bed
2. **Duration:** 3–5 minute recording
3. **Position:** Supine, relaxed breathing
4. **Metrics:** RMSSD (root mean square of successive differences) or HF power
5. **Baseline establishment:** 14 days of daily measurement to establish personal baseline; calculate 7-day rolling average

Validated Devices

- **Chest strap monitors:** Polar H10, Garmin HRM-Pro (gold standard accuracy)
- **Wrist-based:** Oura Ring (validated for overnight HRV), Whoop, Garmin watches (acceptable accuracy for trends)
- **Apps:** Elite HRV, HRV4Training (provide analysis algorithms; require compatible sensor)

Activity Calibration

- **HRV >105% of baseline:** Green day—normal activity budget allowed
- **HRV 90–105% of baseline:** Yellow day—reduce planned activity by 20%; increase rest periods
- **HRV 75–90% of baseline:** Orange day—reduce activity by 40%; prioritize rest; cancel optional commitments
- **HRV <75% of baseline:** Red day—minimal activity only; active recovery day; cancel all non-essential activities

Integration with Activity Planning

- Check HRV before committing to activities
- Reschedule appointments when HRV indicates poor recovery state
- Use HRV as “training wheels” for learning to recognize internal recovery signals
- Over time, patients may develop interoceptive awareness that correlates with HRV readings

Evidence Status HRV-guided training is well-established in sports science [507, 508], with consistent evidence that reduced HRV predicts poor training tolerance and overtraining syndrome [509]. Preliminary evidence supports HRV’s utility in ME/CFS: Escorihuela et al. [255] demonstrated that reduced HRV predicts fatigue severity in ME/CFS patients (n=45), with RMSSD, mean RR intervals, and high-frequency power all significantly correlating with self-reported fatigue ($p < 0.03$). This suggests HRV may serve as an objective indicator of physiological reserve.

However, individual variation in HRV response is substantial; the protocol requires personalization. Some ME/CFS patients have chronically suppressed HRV, requiring adjusted thresholds. Consumer wearable devices are evolving rapidly but require validation for clinical use [510]. A proposed RCT comparing HRV-guided to standard pacing is described in Chapter 33, Section 33.7.

► Protocol 2: Periodized Activity Cycling

Certainty: 0.30. Periodized activity cycling (alternating planned deload and maintenance phases) adapted from sports medicine may optimize recovery compared to static activity maintenance in ME/CFS. The certainty level reflects: (1) well-established efficacy of periodization in athletic training for preventing overtraining syndrome; (2) theoretical parallel between overtraining and ME/CFS post-exertional malaise; (3) however, lack of

any randomized controlled trials directly testing periodization in ME/CFS; (4) inability to replicate the controlled training environments of sports medicine in heterogeneous ME/CFS populations; (5) fundamental uncertainty about whether the overtraining syndrome model accurately describes ME/CFS physiology; (6) high inter-individual variation in activity tolerance that may render standardized cycles ineffective.

Standard ME/CFS pacing emphasizes maintaining a constant activity level within the energy envelope. An alternative approach, adapted from sports medicine management of overtraining syndrome, employs structured cycles of rest and activity that may better support recovery than static management.

Cross-Domain Insight Overtraining syndrome (OTS) in athletes shares features with ME/CFS: persistent fatigue, performance decline, sleep disturbance, mood changes, and autonomic dysfunction [509]. However, OTS outcomes are substantially better—most athletes recover within weeks to months with structured rest-activity cycles. While OTS and ME/CFS likely have different underlying pathophysiology, the recovery principles may be partially transferable.

Key Difference from Standard Pacing Standard pacing maintains constant activity at 50–80% of the energy envelope indefinitely. Periodized cycling alternates between:

- **Deload phases:** Reduced activity below the usual envelope, allowing deeper recovery
- **Maintenance phases:** Standard envelope activity
- **Probe phases:** Carefully monitored slight increases to test capacity (only if stable)

Example 8-Week Cycle

1. **Weeks 1–2 (Deload):** 30–50% of usual activity; prioritize sleep extension (10+ hours if possible); anti-inflammatory nutrition emphasis; cancel all optional activities
2. **Weeks 3–4 (Recovery):** 60–70% of usual activity; maintain extended sleep; continue anti-inflammatory support
3. **Weeks 5–6 (Maintenance):** Return to usual sustainable activity level (70–80% envelope); monitor HRV for stability
4. **Weeks 7–8 (Probe—if stable):** Very slight activity increase (5–10%); immediate reduction if any warning signs; if tolerated, this becomes new maintenance level
5. Repeat cycle

Adjunctive Elements

- **HRV monitoring:** Required throughout; cycle timing should align with HRV patterns
- **Recovery nutrition:** Increased anti-inflammatory foods during deload; protein for tissue repair
- **Sleep extension:** Particularly during deload phases; aim for 9–10 hours

- **Stress minimization:** Schedule demanding life events (appointments, social obligations) during maintenance phases, not deload

Cautions and Contraindications

- **Not for severe patients:** Periodization assumes capacity for activity variation; very severe patients may not tolerate even deload-level activity
- **PEM monitoring essential:** Any PEM during probe phases requires immediate return to deload
- **Individual cycle length:** 8 weeks is illustrative; some patients may need 12-week or 6-week cycles based on their recovery kinetics
- **Experimental approach:** No RCT evidence exists comparing periodized to standard pacing in ME/CFS

Distinction from GET Periodized activity cycling is fundamentally different from graded exercise therapy (GET):

- GET assumes patients can progressively increase activity indefinitely—periodization includes mandatory deload phases
- GET ignores PEM signals—periodization treats any PEM as immediate stop signal
- GET aims to “decondition” patients from activity avoidance—periodization respects energy envelope as biological reality
- GET was designed for presumed psychological aversion—periodization is designed for physiological recovery optimization

Sports Medicine Deload Principles

The periodized activity cycling protocol (Protocol 18.3.3) draws from sports medicine principles of structured recovery. Recent consensus work in athletic training provides more detailed guidance on deload implementation that may inform ME/CFS pacing strategies.

Deload Definition and Rationale Bell et al. [511] define deloading in athletic contexts as “a period of reduced training stress designed to mitigate physiological and psychological fatigue, promote recovery, and enhance preparedness for subsequent training” (n=34 expert coaches, Delphi consensus). In athletes, deloads prevent cumulative fatigue that would otherwise lead to overtraining syndrome. The parallel to ME/CFS: regular planned reductions in activity may prevent the accumulation of metabolic and immune stress that precipitates crashes.

Evidence-Based Parameters from Athletic Training Sports science research establishes:

- **Frequency:** Deloads every 4–6 weeks in athletic populations [511]
- **Duration:** Approximately 7 days (range: 3–14 days depending on individual response)

- **Volume reduction:** 40–60% reduction in total activity through fewer “sets” (activity bouts), shorter duration, or reduced frequency
- **Intensity:** May remain moderate while volume decreases, OR both reduced together
- **Implementation:** Pre-planned (calendar-based) or autoregulatory (HRV/symptom-driven)

Adaptation for ME/CFS: Critical Differences Direct application of athletic deload protocols to ME/CFS would be inappropriate. Key adaptations required:

1. **Baseline capacity:** Athletes start from high-normal fitness; ME/CFS patients from 10–20% of healthy capacity. Activity “volume” in ME/CFS refers to activities of daily living (cooking, hygiene, short walks), not structured training.
2. **Recovery timelines:** Athletes recover from deconditioning in weeks; ME/CFS recovery (if it occurs) requires months to years. Athletic 7-day deloads become 7–14 day deloads in ME/CFS.
3. **Progression philosophy:** Athletic training aims for continuous improvement; ME/CFS management prioritizes stability and preventing deterioration. Any capacity increases are secondary goals.
4. **Consequence of error:** Athletes who overtrain risk temporary performance setbacks; ME/CFS patients who exceed energy envelope risk prolonged relapse. The stakes are fundamentally different.

△ **Warning 4: Not for Severe or Very Severe Patients**

Sports medicine-adapted protocols assume the patient can engage in some level of activity variation and monitoring. Severe and very severe ME/CFS patients who are bed-bound or housebound should not attempt structured deload cycling. For these patients, standard pacing with minimization of all non-essential activity remains the evidence-based approach.

Who May Benefit: Selection Criteria Sports medicine-adapted pacing may be appropriate for:

- Mild to moderate ME/CFS patients (ambulatory, able to perform some daily activities)
- Stable baseline established over 4+ weeks (no recent crashes)
- Previous athletic background (familiar with structured training concepts)
- Comfort with quantitative tracking and data collection
- Access to monitoring tools (smartphone, wearables, tracking apps)
- Psychological readiness for disciplined, patient approach
- Understanding that “progressive overload” is NOT “push through pain”

Contraindications:

- Severe or very severe ME/CFS

- Actively deteriorating or unstable condition
- Recent major crash (within 3 months)
- Tendency toward overachievement or ignoring warning signals
- Psychological distress from metrics or self-monitoring

Objective Recovery Monitoring Beyond HRV

While HRV provides sophisticated autonomic assessment (Protocol 18.3.3), simpler metrics may complement or substitute when HRV monitoring is impractical.

Resting Heart Rate (RHR) as Recovery Indicator Resting heart rate offers a zero-cost alternative to HRV for tracking recovery status:

► Protocol 3: Resting Heart Rate Monitoring

Measurement Protocol:

1. Measure immediately upon waking, before getting out of bed
2. Use manual palpation (radial or carotid pulse for 60 seconds) or wearable device
3. Record daily for 14 days to establish personal baseline
4. Calculate 7-day rolling average

Interpretation:

- **RHR within 3 bpm of baseline:** Normal recovery state; proceed with planned activities
- **RHR 4–6 bpm above baseline:** Caution—reduce activity by 20–30%; monitor closely
- **RHR 7+ bpm above baseline:** Red flag—significant incomplete recovery; reduce activity by 50%; consider early deload phase
- **Sustained elevation (3+ days):** Strong signal for deload cycle regardless of calendar schedule

Evidence Base: Sports medicine literature consistently identifies 5–7 bpm RHR elevation as indicating incomplete recovery or overtraining risk in athletes. However, individual variation is substantial; personal baseline comparison is more meaningful than absolute values. RHR is less sensitive than HRV but far more accessible.

Limitations:

- Affected by sleep quality, hydration, ambient temperature, illness
- Less sensitive than HRV to subtle autonomic changes
- ME/CFS patients may have dysautonomia causing chronically elevated RHR; focus on trends and relative changes

Combined Monitoring Strategy For maximal sensitivity, combine multiple metrics:

- **Primary:** HRV (if available and validated device)

- **Secondary:** Resting heart rate (accessible to all)
- **Tertiary:** Subjective recovery scales (see below)
- **Integration rule:** Use most conservative signal; if any metric indicates poor recovery, reduce activity regardless of other metrics

Subjective Recovery Scales

Systematic reviews of athletic monitoring demonstrate that subjective self-report measures often outperform objective physiological markers for detecting overtraining [512]. Structured subjective scales may enhance ME/CFS self-monitoring.

Recovery-Stress Assessment Validated tools from sports science include:

- **Profile of Mood States (POMS):** Tracks tension, depression, anger, fatigue, confusion, vigor
- **Recovery-Stress Questionnaire for Athletes (RESTQ-Sport):** 76-item assessment of recovery and stress states
- **Daily Analyses of Life Demands (DALDA):** Simple daily symptom checklist
- **Acute Recovery and Stress Scale (ARSS):** Recently validated brief scale for daily use

For ME/CFS, complex questionnaires may create excessive burden. A simplified approach:

► Protocol 4: Daily Recovery Self-Rating

Each morning, rate recovery status on 0–10 scale:

- **0–2:** Severely unrecovered; significant symptom burden; minimal functional capacity
- **3–4:** Poor recovery; moderate symptoms; reduced capacity
- **5–6:** Moderate recovery; mild symptoms; functional but limited
- **7–8:** Good recovery; minimal symptoms; near-normal capacity for individual
- **9–10:** Excellent recovery; no or trivial symptoms; optimal function

Additional Quick Ratings (0–10 scale):

- Sleep quality (0=terrible, 10=excellent)
- Cognitive clarity (0=severe brain fog, 10=clear thinking)
- Physical energy (0=exhausted, 10=energetic)
- Pain level (0=no pain, 10=severe pain)
- Stress level (0=calm, 10=highly stressed)

Use of Data:

- Track weekly average and trend
- If weekly average declining over 2 weeks: initiate deload regardless of calendar
- If recovery rating <5 for 3+ consecutive days: reduce activity immediately

- Use in combination with objective metrics (HRV, RHR) for comprehensive picture

Practical Implementation Framework

For patients considering sports medicine-adapted pacing, a phased implementation reduces risk:

Phase 1: Baseline and Monitoring Setup (Weeks 1–4)

1. Establish stable activity baseline (no increases; just observe current capacity)
2. Implement daily monitoring: RHR, subjective recovery rating, sleep quality
3. Optional: Add HRV if device available
4. Track PEM occurrences (frequency, severity, triggers)
5. Calculate personal baseline for all metrics
6. Goal: 4 weeks of stable data before any changes

Phase 2: First Planned Deload (Week 5)

1. Reduce activity to 50% of baseline week
2. Focus on rest, sleep extension (aim for 9–10 hours), gentle movement only
3. Continue all monitoring
4. Observe: Do recovery metrics improve during deload? By how much?
5. If no improvement or worsening: standard pacing may be more appropriate than periodization

Phase 3: Return to Baseline (Weeks 6–7)

1. Gradually return to pre-deload baseline activity level
2. Monitor for PEM or metric deterioration
3. If stable: baseline re-established
4. If unstable: remain at reduced level; reconsider approach

Phase 4: Assessment and Decision (Week 8)

1. Review 8-week data: trends in RHR, HRV, subjective ratings, PEM frequency
2. **If improving:** Consider continuing with 4–6 week cycles
3. **If stable:** Continue cycles with no progression attempts; cycles maintain stability
4. **If declining:** Return to standard flexible pacing; periodization may not suit individual physiology

Long-Term Management

- Deload every 4–6 weeks (pre-planned) OR when metrics indicate (autoregulatory)
- **Never attempt progression if unstable**
- If stable for 3+ months: may consider ultra-conservative 5% activity increase; immediate rollback if any PEM
- Reassess approach every 3–6 months; be willing to abandon if not beneficial

→ Recommendation 3: Physician Consultation

Patients attempting structured periodization should discuss the approach with their ME/CFS-knowledgeable physician. Monitoring data (RHR trends, recovery ratings, PEM logs) should be shared at appointments to enable collaborative adjustment. Any worsening of baseline function requires immediate return to standard pacing and medical evaluation.

Critical Distinction: This Is Not GET Sports medicine-adapted pacing shares superficial similarities with graded exercise therapy (GET) but differs fundamentally in philosophy and implementation:

GET (Inappropriate for ME/CFS)	Sports-Adapted Pacing
Assumes progressive increase indefinitely	Includes mandatory regular deloads
Treats PEM as psychological barrier to overcome	Treats PEM as hard biological stop signal
Fixed progression schedule regardless of symptoms	Autoregulatory adjustment based on recovery metrics
Aims to “decondition” from activity avoidance	Respects energy envelope as physiological reality
Based on deconditioning hypothesis	Based on metabolic/immune recovery optimization
Ignores autonomic dysfunction	Incorporates HRV/RHR monitoring
One-size-fits-all protocol	Highly individualized to patient metrics
Progression is primary goal	Stability is primary goal; progression secondary if at all

Table 18.4: Comparison of GET vs. Sports Medicine-Adapted Pacing

The distinction is critical: GET has been shown to be harmful in significant subsets of ME/CFS patients [3, 506] and is no longer recommended by CDC, NIH, or major ME/CFS specialist organizations [127]. Sports-adapted pacing, by contrast, is explicitly designed around energy envelope theory and includes structured recovery phases. However, it remains an experimental approach without ME/CFS-specific validation and must be implemented with extreme caution.

Evidence Status Certainty Assessment:

- **Athletic deload protocols:** High-quality evidence in sports science
- **OTS parallels to ME/CFS:** Medium-quality observational evidence; significant differences exist
- **HRV and RHR monitoring:** High-quality in athletes; limited data in ME/CFS
- **ME/CFS adaptation:** Low-quality; theoretical extrapolation only; no RCTs

? Open Question 1: Sports-Adapted Pacing: Unresolved Evidence Gaps

Randomized controlled trials comparing sports-adapted versus standard pacing in ME/CFS do not yet exist. Specifically unresolved: optimal deload frequency, duration, and depth; long-term outcomes beyond 6 months; validated patient selection criteria; and systematic safety evaluation.

Proposed Research: Chapter 33 includes a proposal for an RCT comparing sports medicine-adapted periodization to standard flexible pacing in mild-moderate ME/CFS (Section 33.8).

Clinical Bottom Line Sports medicine-adapted pacing represents a **reasonable experimental approach** for carefully selected mild-moderate ME/CFS patients who:

- Have stable baselines
- Are comfortable with structured monitoring
- Understand the distinction from GET
- Accept the lack of ME/CFS-specific validation
- Are willing to abandon the approach if unhelpful or harmful

It should be implemented conservatively, with close monitoring, and under physician guidance. Standard flexible pacing remains the evidence-based default for all patients, particularly those with severe disease, unstable courses, or discomfort with quantitative tracking.

18.3.4 Symptom Management for Mild-Moderate Cases**Cognitive Dysfunction (Brain Fog)**

Rationale Cognitive dysfunction results from multiple mechanisms: catecholamine deficiency (Section 6.6), cerebral hypoperfusion (Section 8.5), and reduced ATP availability in the brain (Section 6.1). Targeting neurotransmitter precursors and optimizing cerebral blood flow can improve function.

Non-Pharmaceutical

- **Cognitive pacing:**
 - Work in 25-minute blocks (Pomodoro technique), then 10-minute rest
 - Schedule cognitively demanding tasks for peak energy times (usually morning)
 - Minimize multitasking (switching costs energy)
 - Reduce decision-making load (meal planning, outfit planning in advance)
- **Environmental optimization:**
 - Reduce sensory overload (quiet workspace, minimal visual clutter)
 - Close unnecessary browser tabs/apps
 - Use noise-canceling headphones if sound-sensitive

Pharmaceutical/Supplement

- **Tier 1** (try first):
 - Caffeine + L-theanine (100 mg + 200 mg, 1–2 times daily)
 - Alpha-GPC 300 mg BID (choline support for acetylcholine)
 - Rhodiola rosea 200–400 mg morning (adaptogen, focus)
- **Tier 2** (add if Tier 1 helps):
 - Bacopa monnieri 300 mg daily (memory consolidation)
 - Lion's Mane mushroom 500–1000 mg BID (nerve growth factor)
 - Citicoline 250 mg BID (neuroprotection)
- **Tier 3** (prescription if severe cognitive impairment):
 - Modafinil 50–100 mg morning (wakefulness, often prescribed off-label)
 - Or: Methylphenidate 5 mg BID (stimulant, use cautiously)

Intranasal Delivery Routes for CNS-Targeted Compounds

★ Key Point: BBB Penetration and Intranasal Delivery

The blood-brain barrier (BBB) may limit delivery of compounds needed for cognitive support in ME/CFS (Hypothesis 14.24.5). Intranasal delivery bypasses the BBB via olfactory and trigeminal nerve pathways, achieving 2–10 fold higher CSF concentrations than oral routes.

→ Recommendation 4: Prioritizing Intranasal Routes When Available

For mild-moderate patients with prominent cognitive dysfunction:

- **Modafinil intranasal:** If oral modafinil provides partial benefit, discuss intranasal formulations with prescribing physician. Not yet standard care but literature supports improved cognitive outcomes in other neurological conditions.

- **Dopamine or L-DOPA analogues (intranasal):** Specialist neurologists may consider intranasal dopamine precursors if oral neurotransmitter support insufficient. EXPERIMENTAL; not standard ME/CFS care.
- **Future compounds:** As understanding of the astrocyte energy gate hypothesis (Speculation 38) improves, intranasal delivery of lactate, ketone bodies, or neuro-protective compounds may emerge as targeted interventions.

Practical application: If cognitive symptoms dominate despite Tier 1–2 oral support, ask your physician about intranasal formulation options or referral to a neurologist familiar with BBB dysfunction.

Evidence level: Speculative (established for other neurological conditions; no ME/CFS-specific trials)

Transcranial Direct Current Stimulation (tDCS) for Cognitive Enhancement

Speculation 57 (tDCS Energy Cost Reduction via DLPFC Modulation). **Certainty: 0.25.** Anodal tDCS targeting the DLPFC modulates cortical excitability and has demonstrated improvements in working memory, attention, and executive function in multiple studies [513]. Applied to ME/CFS, this neural efficiency gain may reduce the energy cost of Tier 5 cognitive tasks (Section 14.24), thereby improving sustainable cognitive performance within the patient's energy envelope. No ME/CFS-specific trials exist; the application to energy triage theory is speculative extrapolation.

→ Recommendation 5: Home tDCS Protocol for Cognitive Enhancement in Mild-Moderate ME/CFS

Mechanism: Anodal tDCS to DLPFC increases cortical excitability, potentially reducing energy cost of executive function through improved neural efficiency.

Protocol (home-based):

1. Equipment:

- **tDCS device:** Commercial home units (Thync, Flow, Halo Sport) or medical-grade devices (\$300–2000)
- **Budget option:** DIY kits available but require strict safety adherence; medical supervision recommended initially

2. Stimulation parameters:

- **Intensity:** 2 mA (safe range for home use: 1–2 mA)
- **Duration:** 20 minutes daily
- **Montage:** F3-F4 (DLPFC bilateral, using 10-20 EEG positioning)
 - Anode (positive): F3 (left DLPFC)
 - Cathode (negative): F4 (right DLPFC) or right supraorbital
- **Frequency:** Daily or 5 days/week

- **Duration of trial:** 4–8 weeks to assess efficacy

3. Cognitive tracking during trial:

- Rate executive function daily (0–10 scale): Planning, multitasking, decision-making
- Track fatigue timing and intensity
- Monitor for mood or behavioral changes
- Weekly summary: "Week 1: no change. Week 3: Planning tasks feel 30% easier. Week 6: Sustained improvement in attention span."

4. Safety considerations:

- Start with 1 mA if new to tDCS; escalate to 2 mA if well-tolerated
- Common side effects (mild, temporary): Tingling under electrodes, mild headache, slight skin redness
- Discontinue if: Persistent headache, mood changes, seizure activity
- **Absolute contraindications:** Metal implants in head/brain, history of seizures, pregnancy (insufficient safety data)

5. Integration with pacing:

- tDCS improves cognitive capacity but does NOT increase energy envelope
- Improved cognitive function may tempt increased activity; maintain strict pacing to avoid PEM
- Think: "More efficient cognition at same energy expenditure," not "more capacity"

Evidence level: Speculative (tDCS efficacy for cognition documented; tDCS + energy triage model untested in ME/CFS)

Expected outcomes: 20–40% subjective improvement in executive function (planning, multitasking, decision-making). Effects may take 3–4 weeks to emerge. Not expected to improve fatigue directly; improves cognitive performance within existing energy envelope.

Practical consideration: Requires initial physician consultation for safety screening and proper electrode placement. Some occupational therapists experienced with tDCS can assist with home setup.

Sleep Dysfunction

Rationale Non-restorative sleep is a core ME/CFS symptom (Section 2.2). Sleep dysfunction amplifies all other symptoms through effects on immune function (Section 7.4.1), pain sensitization, and cognitive impairment. Optimizing sleep is foundational to symptom control.

Sleep Hygiene (Non-Negotiable Foundation)

- Same sleep/wake time every day (weekends included)
- 7–9 hour sleep opportunity (in bed, dark, quiet)

- Room: 65–68°F, completely dark, quiet
- No screens 2 hours before bed (or blue blockers)
- No caffeine after 2pm
- No large meals 3 hours before bed
- Wind-down routine: 30 minutes relaxing activity before bed (reading, gentle stretching, meditation)

Supplements (Mild Cases Can Start Here)

- Melatonin 0.5–3 mg (2 hours before target sleep time; start low)
- **Magnesium glycinate 400 mg evening** - NOTE: At upper end of RDA (320 mg women, 420 mg men). Provides 400 mg elemental magnesium for muscle relaxation and calming. Very safe, well-tolerated. May cause loose stools if exceed tolerance (reduce dose if occurs).
- L-theanine 200 mg before bed (anxiolytic)
- **Glycine 3 g before bed** - NOTE: Exceeds typical supplement dose (1–2 g) by 1.5–3×. Clinical studies for sleep quality improvement use 3 g [514]. Mechanism: Glycine lowers core body temperature via NMDA receptor agonism in the suprachiasmatic nucleus, facilitating sleep onset [515]. Extremely safe (used as food additive); no adverse effects in clinical trials. Sweet taste can be mixed in water.

Prescription (If Supplements Insufficient)

- Trazodone 25–50 mg (lower dose than severe cases; increase if needed)
- Mirtazapine 7.5 mg (also helps appetite)
- Doxepin 3–6 mg (low-dose, histamine antagonist, improves sleep maintenance)

Dual Orexin Receptor Antagonists (DORAs) for Chronic Sleep Support

★ Achievement 1: Daridorexant: Evidence-Based DORA for ME/CFS Sleep

Dual orexin receptor antagonists (DORAs) offer a mechanistically targeted approach to ME/CFS sleep dysfunction, given documented orexin system abnormalities in the condition [498]. Daridorexant (Quviviq), FDA-approved in 2022, has robust evidence from multiple meta-analyses: Rocha et al. [516] (10 RCTs, n=7,806) established dose-response relationships; Xue et al. [499] (13 RCTs) confirmed class-wide DORA efficacy; Dutta et al. [517] provided GRADE assessment showing MODERATE certainty for safety comparable to placebo.

Unlike Z-drugs and benzodiazepines, DORAs consolidate sleep by reducing *long* wake bouts (>6 minutes) correlated with daytime impairment, while preserving brief arousals that maintain healthy sleep-wake boundary control [518]. This mechanism addresses non-restorative sleep without producing hangover effects or tolerance.

Long-term safety: 52-week extension study (n=801) demonstrated no tolerance or

withdrawal phenomena with continuous or intermittent use [500].

Practical protocol: Start daridorexant 25 mg 30 minutes before bedtime with at least 7 hours available for sleep [519]. If insufficient after 4–6 weeks, increase to 50 mg. Safe for chronic use without tolerance development [520]. **Advantages over traditional sleep aids:** No next-day sedation; no cognitive impairment; no tolerance; suitable for long-term use in ME/CFS.

Limitations: No ME/CFS-specific RCTs exist. Prescription required; cost may be barrier. Alternative DORAs (suvorexant, lemborexant) have similar efficacy if daridorexant unavailable.

Circadian Light Therapy for Sleep-Energy Alignment

★ Key Point: Circadian Energy Misallocation in ME/CFS

The selective energy dysfunction hypothesis (Hypothesis 14.24.10) proposes that SCN dysfunction impairs circadian allocation of energy budgets, explaining why many patients experience energy crashes mid-afternoon but a late-evening “second wind.”

→ Recommendation 6: Circadian Light Therapy: Entrainment Protocol

Mechanism: Bright morning light exposure resets the circadian oscillator, improving alignment between energy availability and day-night cycle. This synergizes with sleep medications by improving melatonin timing.

Protocol (same as severe cases):

1. **Equipment:** 10,000 lux light therapy box (\$25–100)
2. **Timing:** Within 30 minutes of waking, 20–30 minutes daily, same time every day
3. **Position:** 16–24 inches from face, 30° downward angle
4. **Do NOT use after 3pm** (risk of sleep disruption)

Evidence level: Moderate (circadian disruption documented; light therapy established for circadian disorders; ME/CFS-circadian-energy RCTs pending)

Expected outcomes:

- More consistent daytime energy
- Earlier, easier sleep onset at night
- Reduced afternoon crashes
- Timeline: 2–4 weeks

Sleep Spindle Enhancement via Acoustic Stimulation (Low Priority, Optional)

→ Recommendation 7: Pink /White Noise for Sleep Architecture Improvement

Mechanism: Sleep spindles (brief high-frequency brain activity during NREM sleep) are reduced in ME/CFS. Acoustic stimulation may enhance spindle production, potentially improving sleep restorativeness (Hypothesis 14.24.10).

Simple, Low-Cost Protocol:

- **Equipment:** White or pink noise machine (\$10–50) or free app (myNoise.net, Noisli)
 - White noise: Constant across frequencies; easier to find and more common
 - Pink noise: Lower frequencies emphasized; some literature suggests superior sleep effects
- **How to use:**
 - Play throughout entire sleep period
 - Volume: Low (30–50 dB, about conversational level)
 - Placement: Bedside speaker or sleep-friendly earplugs
- **Trial duration:** 2–4 weeks minimum to assess effect
- **Tracking:**
 - Subjective sleep quality rating (0–10)
 - Morning refreshedness
 - Daytime cognitive clarity
 - Expected timeline: 2–4 weeks if beneficial

Evidence level: Speculative (spindle deficits documented in ME/CFS; acoustic enhancement effect unproven in this population)

Expected outcomes: Modest improvement in sleep quality perception. Not expected to directly improve daytime fatigue.

Positioning: Low-priority addition. Sleep medications (melatonin, trazodone) have stronger evidence. Use acoustic stimulation if medications insufficient or patient prefers non-pharmacological approach.

Pain

Rationale Pain in ME/CFS involves inflammatory mediators (Section 7.3.1), small fiber neuropathy (Section 8.3.1), and central sensitization. Addressing inflammation and neuropathic pathways reduces pain burden.

Mild-Moderate Pain Management

- **First-line:**
 - Ibuprofen 400 mg PRN or BID (with food)
 - Or: Naproxen 220–500 mg BID
 - Topical: Diclofenac gel (Voltaren) to painful areas
- **Add if insufficient:**
 - Low-dose naltrexone (LDN) 1.5–4.5 mg nightly (immune modulation + pain)
 - Turmeric/curcumin 500–1000 mg BID (natural anti-inflammatory)
 - Magnesium glycinate 400 mg daily (muscle relaxation)

- **Neuropathic pain component:**

- Gabapentin 100 mg at bedtime, increase slowly to 300–600 mg BID if needed
- Or: Duloxetine 30–60 mg daily (also helps mood)

Palmitoylethanolamide (PEA) for Neuropathic and Inflammatory Pain

Observation 81 (PEA Meta-Analytic Evidence for Chronic Pain). Palmitoylethanolamide (PEA), a naturally occurring endocannabinoid-like fatty acid amide, has robust meta-analytic evidence for chronic pain reduction. Three independent meta-analyses demonstrate consistent, large effect sizes: Artukoglu et al. [521] analyzed 10 studies (n=1298) finding weighted mean difference of 2.03 (95% CI 1.19–2.87, $p<0.001$); Lang-Ilievich et al. [501] confirmed these findings in 11 double-blind RCTs (n=774), reporting standardized mean difference of 1.68 (95% CI 1.05–2.31, $p<0.00001$); Viña & López-Moreno [502] conducted the most comprehensive analysis (18 RCTs, n=1196), demonstrating PEA efficacy across all pain types: nociceptive (SMD=-0.74), neuropathic (SMD=-0.97), and nociplastic (SMD=-0.59). Benefits emerged at 4–6 weeks and increased through 24–26 weeks. Quality of life improved significantly beyond pain reduction alone. No major adverse events were reported across all trials.

Evidence quality: HIGH for general chronic pain (multiple independent meta-analyses, $n>1000$ patients). MEDIUM for ME/CFS-specific use (extrapolated; no ME/CFS RCTs).

~ Hypothesis 2: PEA Mechanisms Align with ME/CFS Pain Pathophysiology

Certainty: 0.45. PEA's mechanisms of action directly target pathways implicated in ME/CFS pain. Petrosino et al. [503] demonstrated that PEA counteracts mast cell activation by stimulating diacylglycerol lipase- β (DAGL- β), increasing endogenous 2-arachidonoylglycerol (2-AG), which activates CB2 receptors to inhibit mast cell degranulation and histamine release—particularly relevant given mast cell activation in ME/CFS subsets (Section 18.3.5). Additionally, PEA functions as a PPAR- α agonist, reducing neuroinflammation through glial cell modulation and suppression of pro-inflammatory cytokine expression [522].

Practical protocol: Prefer *micronized* or *ultramicrosized* PEA formulations (enhanced solubility profile; superiority over standard PEA on clinical outcomes remains under investigation [523]). Dose: 600 mg twice daily. Time to benefit: 4–6 weeks for initial effect; peak benefit at 24–26 weeks [523]. Excellent safety profile with minimal side effects documented across trials.

Positioning: Consider in the “Add if insufficient” tier alongside LDN and curcumin. PEA has a larger evidence base than curcumin (multiple meta-analyses [523] vs limited RCT data). Particularly indicated if: mast cell activation features present, neuropathic pain component inadequately controlled, or inadequate NSAID response.

Orthostatic Intolerance (POTS)

Rationale Orthostatic intolerance affects 70–90% of ME/CFS patients (Section 8.2.2). Reduced blood volume (Section 10.2.2), autonomic dysfunction (Section 8.2), and impaired vas-

cular regulation contribute. Blood volume expansion and compression improve tolerance.

Mild-Moderate Interventions

- **Compression:** Waist-high stockings 20–30 mmHg (lower compression than severe cases)
- **Salt: 6–8 g sodium daily - NOTE - DRAMATICALLY EXCEEDS STANDARD RECOMMENDATION:** Standard guideline is <2300 mg (2.3 g) daily. We recommend 6000–8000 mg (6–8 g) sodium daily, which is 2.6–3.5× standard. See Chapter 17 for complete justification (blood volume expansion for orthostatic intolerance, standard POTS treatment). Electrolyte drinks make compliance easier. **CONTRAINDICATIONS:** Hypertension, heart failure, kidney disease. Monitor BP weekly.
- **Oral rehydration solution (ORS) - dual benefit:** Beyond simple blood volume expansion, properly formulated electrolyte solutions address the chronic metabolic stress state documented in Section 6.6. ME/CFS patients exist in a continuous state of lactate accumulation and reliance on anaerobic metabolism similar to post-exercise metabolic stress in athletes (see Chapter 6). Strategic electrolyte replacement serves multiple purposes:
 - **Blood volume expansion:** Maintains preload for cardiac output; reduces orthostatic intolerance
 - **Lactate clearance:** Helps clear accumulated lactic acid from impaired oxidative metabolism
 - **Glucose availability:** Provides immediate energy when fat-burning is impaired
 - **Electrolyte balance:** Supports muscle function and reduces cramping from ATP depletion

Recommended formulation (sports medicine-derived):

- Dry mix: 100 g sugar + 15 g low-sodium salt (KCl) + 15 g table salt (NaCl)
- Dosing: 7 g dry mix in 250 mL water, twice daily
- Flavoring optional (e.g., 10 mL grenadine for palatability)
- Cost: <€5 for months of supply

This formulation provides sodium, potassium, chloride, and glucose in ratios optimized for absorption and metabolic support. See Appendix ?? for the clinical insight that led to this protocol development.

- **Fluids:** 2.5–3 L daily
- **Positional changes:** Rise slowly (sit 30 seconds before standing)
- **Counter-maneuvers:** Leg crossing, muscle tensing when standing
- **Exercise:** Recumbent bike or rowing (horizontal position) within energy envelope

Compression Garments for Autonomic Load Reduction

→ **Recommendation 8: Medical-Grade Compression Stockings for Mild-Moderate Orthostatic Intolerance**

Mechanism: Compression garments reduce autonomic coordination load by maintaining peripheral venous pressure, reducing baroreceptor-mediated sympathetic activation required for orthostatic compensation (Speculation 42: SFN interface failure).

Practical Protocol:

1. Compression class selection:

- **Class II (20–30 mmHg):** Recommended for mild-moderate orthostatic intolerance
- **Waist-high or thigh-high:** Covers leg venous return (most effective for OI)
- **Material:** Medical-grade merino wool or synthetic (avoid cotton which loses compression)

2. Wearing schedule:

- **During upright activities:** All times child is sitting or standing (except during sleep or recumbent rest)
- **Examples:** School day, meals, therapy appointments, activities
- **Remove during sleep:** Not needed in horizontal position
- **Daily wear:** 8–12 hours typical

3. Expected benefits:

- Reduced tachycardia with position changes
- Improved cognitive clarity (cerebral perfusion stabilized)
- Reduced fatigue from sustained orthostatic compensation
- Better school tolerance and attendance

4. Practical considerations:

- **Fitting:** Measure leg diameter for proper sizing; incorrect fit loses effectiveness
- **Compliance:** Children may resist wearing; emphasize improved energy / cognition benefits
- **Cost:** Medical-grade stockings \$30–60 per pair; insurance may cover with prescription for POTS
- **Longevity:** Replace every 3–6 months (lose compression with washing)

5. Integration with other OI treatments:

- Combine with salt loading and hydration protocol for maximum effect
- Can be used with medications (midodrine, fludrocortisone)
- Adjunctive benefit; should not replace blood volume expansion

Evidence level: Moderate (20–30 mmHg compression established for POTS; extends to autonomic-primary ME/CFS subtype with SFN features)

Expected outcomes: 20–40% reduction in orthostatic symptoms when combined with salt/fluid protocol. Effects may take 1–2 weeks as child adjusts to compression.

Prescription (If Above Insufficient)

- Fludrocortisone 0.05–0.1 mg daily (increases blood volume)
- Midodrine 2.5–10 mg TID (peripheral vasoconstrictor)
- Beta-blockers (propranolol, metoprolol) - use cautiously, can worsen fatigue in some
- **Ivabradine 2.5–7.5 mg BID** (If blocker) - Selectively reduces heart rate by inhibiting the I_f current in sinoatrial node. **Advantages over beta-blockers:** Does not reduce contractility or blood pressure; may be better tolerated in ME/CFS patients prone to hypotension. More commonly used in Europe than US. Patient reports indicate significant functional improvement (e.g., standing HR reduction from 150 to 90+ bpm). **Contraindications:** Bradycardia (HR <60), hypotension, sick sinus syndrome, concurrent use with strong CYP3A4 inhibitors.

Neuromodulation: Transcutaneous Vagus Nerve Stimulation (tVNS)**★ Achievement 2: tVNS Reduces Orthostatic Tachycardia in POTS**

Teixeira et al. [496] conducted the first randomized, double-blind, sham-controlled trial of transcutaneous vagus nerve stimulation (tVNS) for postural tachycardia syndrome. Daily tragus stimulation (20 Hz, 1 mA below discomfort threshold, 1 hour per day for 2 months, n=26) significantly reduced orthostatic tachycardia compared to sham (heart rate increase during tilt test: 26.4 bpm at baseline → 17.6 bpm at 2 months in active group, $p < 0.05$; no change in sham group).

Mechanisms included decreased β_1 -adrenergic and α_1 -adrenergic receptor autoantibodies, reduced inflammatory cytokines, and improved heart rate variability. The intervention was well-tolerated with no serious adverse events [524].

Study quality: HIGH (randomized, sham-controlled, published in JACC: Clinical Electrophysiology). Requires larger replication trials.

Practical protocol: Auricular tVNS targeting tragus or cymba concha; 20–25 Hz, 0.5–1 mA (below discomfort threshold); start with 5–10 minutes daily and gradually increase to 30–60 minutes over several weeks. Devices include FDA-approved GammaCore (cervical) and research/CE-marked auricular devices (NEMOS, Parasym). **Home-based** treatment suitable for bedbound patients.

△ Warning 5: tVNS Caution in Severe ME/CFS

An international ME/CFS patient survey (n=116) found that “normal” tVNS settings can cause crashes in severe ME/CFS patients [497], although 56% reported favorable effects overall. For severe ME/CFS: use very gradual titration (start 0.5 mA, 5 minutes), monitor for delayed symptom exacerbation (24–48 hours), and discontinue if crashes occur. Formal trials to identify safe parameters for the ME/CFS population are needed.

18.3.5 Mast Cell Activation Syndrome (MCAS) Management

Evidence and Rationale

Mast cell activation affects 30–50% of ME/CFS patients [235]. Recent research demonstrates measurable mast cell phenotype abnormalities with significant increases in naïve mast cells and elevated activation markers [236]. MCAS may worsen orthostatic intolerance, brain fog, and fatigue through excessive histamine and vasoactive mediator release [235].

Critical finding: H1 antihistamine alone showed NO benefit in double-blind RCT [237]. However, **H1+H2 combination** showed dramatic improvement in Long COVID case meeting ME/CFS criteria, with symptom worsening upon discontinuation [238].

Trial Indications

Consider MCAS trial if ANY present:

- Food sensitivities/intolerances (especially new-onset)
- Documented allergies (elevated IgE to foods, pollens, environmental allergens)
- Flushing, hives, itching
- Reactive to fragrances, chemicals
- GI symptoms (post-meal nausea, bloating)
- Unexplained anxiety/panic-like episodes
- Fluctuating brain fog (worse after eating or exposure to triggers)

Treatment Options (Evidence-Based Hierarchy)

Option 1: Standard H1+H2 Combination Based on Long COVID case evidence [238]:

- **H1:** Loratadine 10 mg OR fexofenadine 180 mg (morning)
- **H2:** Famotidine 20 mg twice daily
- **Expected benefits:** Energy, cognitive function, orthostatic tolerance

Option 2: Rupatadine (Superior H1 Choice) Rupatadine offers unique advantages [239, 240]:

- **Triple mechanism:** H1 antagonist + PAF antagonist + mast cell stabilizer
- **Superior efficacy:** Network meta-analysis ranks rupatadine 20 mg highest (SUCRA 99.7%) vs loratadine (lowest rank) [240]
- **PAF antagonism:** 31× more potent than loratadine at blocking PAF; addresses vascular dysfunction in ME/CFS [239]
- **Mast cell stabilization:** Inhibits IL-8 (80%), VEGF (73%), histamine (88%) [239]

Recommended protocol:

- Rupatadine 10 mg morning (increase to 20 mg after 1–2 weeks if insufficient benefit)
- Add famotidine 20 mg BID for complete histamine receptor coverage
- Optional: Add quercetin 500–1000 mg daily (see below)

Option 3: Quercetin (Natural Mast Cell Stabilizer) Evidence shows quercetin MORE effective than prescription cromolyn [241]:

- **Dose:** 500–1000 mg daily (clinical trials used up to 2 g/day)
- **Evidence:** Reduced contact dermatitis reactions >50% in 8 of 10 patients; outperformed cromolyn for substance P-induced mast cell activation [241]
- **Advantages:** Over-the-counter, well-tolerated, additional antioxidant benefits
- Can combine with H1+H2 antihistamines for comprehensive mast cell targeting

4-Week Trial Protocol

Week 1–2: Start H1 antihistamine

- Rupatadine 10 mg morning (preferred), OR
- Fexofenadine 180 mg OR loratadine 10 mg morning
- Monitor for sedation (rare with rupatadine/fexofenadine)

Week 2–4: Add H2 blocker

- Famotidine 20 mg twice daily (morning and evening)
- Note: May reduce stomach acid; take iron supplements 2 hours apart

Optional Enhancement:

- Add quercetin 500–1000 mg daily for additional mast cell stabilization

Low-histamine diet (adjunct):

- Avoid: Aged/fermented foods, alcohol, cured meats, leftovers >24 hours
- Duration: Strict 2-week trial, then gradual reintroduction

Assessment at Week 4:

- **Discontinuation test:** Stop antihistamines for 2–3 days
- If symptoms worsen → mast cell component confirmed → continue therapy
- If no change → discontinue (not MCAS-driven)

Expected Response May improve (if MCAS-related):

- Brain fog and cognitive clarity
- Energy levels (especially post-meal fatigue)
- GI symptoms (bloating, nausea, diarrhea)
- Orthostatic tolerance
- Flushing and allergic symptoms
- Anxiety/panic-like episodes

Will NOT improve (metabolic/mitochondrial):

- Core fatigue (“running on empty”) — requires mitochondrial support
- Muscle cramps — requires carnitine, magnesium
- PEM from overexertion — requires pacing
- Progressive vision/hearing loss — different mechanisms

Special Note: Amitriptyline for Dual Benefit If pain and/or sleep issues coexist with MCAS features, amitriptyline provides dual benefit [242]:

- **Dose:** 10–50 mg at bedtime
- **Mechanisms:** Mast cell inhibition (reduces IL-8, VEGF, IL-6, histamine) [242] + pain relief + sleep improvement
- **Specificity:** This mast cell effect is unique to amitriptyline; other antidepressants (bupropion, citalopram, atomoxetine) do NOT inhibit mast cells [242]
- Can combine with rupatadine + famotidine for comprehensive mast cell targeting

MCAS Prophylactic Intensification for High-Demand Activities and Known Triggers→ **Recommendation 9: Prophylactic Intensification for Mild-Moderate MCAS Patients**

Mechanism: Mast cell activation episodes amplify fatigue and cognitive crashes through inflammatory mediators (Hypothesis 14.24.10). Proactive medication intensification 1–2 days before high-demand activities can reduce crash severity.

Protocol (adapted for mild-moderate severity):

1. **Identify your triggers** (2–4 weeks baseline tracking):
 - Activities: Exercise, busy work/school days, emotional stress
 - Foods: Histamine-rich (aged cheese, fermented foods, red wine, cured meats)
 - Environmental: Heat, cold, strong fragrances, weather changes
 - Immune: Infections, vaccinations, allergy exposure
2. **Prophylactic medication protocol** (BEGIN 24 HOURS BEFORE known triggers):
 - **Increase antihistamine dosing:**

- If on rupatadine 10 mg: Increase to 20 mg daily during trigger window (if previously well-tolerated)
- If on loratadine/fexofenadine: May increase frequency but dose caps apply (consult pharmacist)
- Famotidine: Increase to 40 mg BID (maximum therapeutic dose) during trigger window
- **Add mast cell stabilizer if not already taking:**
 - Quercetin 1000 mg BID (1–2 days pre-trigger and during)
 - Omega-3 PUFA 2–3 g daily (natural stabilizing effect)
- **Strict low-histamine diet** (absolute 24 hours before through 24 hours after trigger):
 - Eliminate all aged/fermented foods
 - Only fresh foods prepared same-day
 - Skip known personal food triggers
- **Activity pacing intensification:**
 - Reduce non-essential activities day-of trigger
 - Maintain strict heart rate pacing limits
 - Prioritize rest before and after high-demand event

3. Track crash response:

- Rate post-trigger crash severity (0–10 scale)
- WITH prophylaxis: "Normally crash 6/10 for 2 days; prophylaxis reduced to 3/10 for 1 day"
- WITHOUT prophylaxis: "Skipped prophylaxis, crashed 7/10 for 2.5 days"
- Adjust prophylaxis strategy based on efficacy pattern

Evidence level: Moderate (MCAS prophylaxis standard in allergology; ME/CFS crash-mitigation studies pending)

Expected outcomes: 25–50% reduction in crash severity or duration when MCAS component is substantial. Lesser benefit if non-MCAS mechanisms predominate.

18.4 Systematic Comorbidity Screening: The Septad Framework

ME/CFS patients frequently present with a cluster of interrelated comorbidities that require distinct treatment approaches. The “Septad” framework (Section 5.6.11) organizes seven conditions that commonly co-occur. Systematic screening can identify treatable contributors to symptom burden.

18.4.1 The Seven Septad Components

1. **Mast Cell Activation Syndrome (MCAS):** See Section 18.3.5 for screening and treatment

2. **Ehlers-Danlos Syndrome (EDS) / Hypermobility:** Joint hypermobility, subluxations, chronic pain
3. **Dysautonomia / POTS:** Orthostatic intolerance (Section 18.3.4)
4. **Autoimmunity:** Subclinical or overt autoimmune markers
5. **Chronic Infection:** Viral reactivation (EBV, HHV-6), tick-borne infections
6. **Small Fiber Neuropathy (SFN):** Pain, paresthesias, autonomic symptoms
7. **GI Dysmotility:** Gastroparesis, SIBO, malabsorption

18.4.2 Screening Recommendations for Mild-Moderate Cases

EDS / Hypermobility Screening. Screen all ME/CFS patients for hypermobility using the Beighton score. If Beighton score $\geq 5/9$ or clinical features suggest EDS:

- **Physical therapy referral:** Hypermobility-aware PT for joint stabilization
- **Avoid overextension:** Joints at risk for subluxation and chronic instability
- **Consider genetics referral:** For formal EDS typing if features suggest vascular or classical type
- **Monitor for progression:** Hypermobile patients may develop additional complications over time

Craniocervical Instability (CCI) Awareness. CCI is not part of the original Septad but occurs in hypermobile patients and can cause ME/CFS-like symptoms (fatigue, cognitive dysfunction, autonomic dysfunction). A specialized clinic study found high prevalence of structural abnormalities (80% with craniocervical obstructions) in ME/CFS patients, predominantly hypermobile [155]; however, these findings require replication in unselected populations (see Section 5.6.11 for detailed evidence and caveats). Consider CCI evaluation if:

- Confirmed EDS/hypermobility PLUS
- Symptoms worse with neck position changes, or
- Occipital headaches, or
- Symptoms suggestive of brainstem compression (dysphagia, facial numbness, gait instability)

Evaluation: Upright MRI preferred over supine (dynamic instability may not appear supine); reference ranges for measurements are available [157]. Specialist referral (neurosurgeon with CCI expertise) if clinical suspicion high. See Lohkamp et al. [156] for diagnostic criteria review.

△ **Warning 6: CCI Is Rare But Treatable**

CCI is uncommon even in hypermobile ME/CFS patients. However, it represents a *structural*, potentially *treatable* cause of symptoms. Conservative management (physical therapy [158], cervical collar) is first-line; surgery shows 60–80% improvement in properly

selected cases but carries significant complication rates (19%) [159]. Do not pursue CCI workup unless hypermobility is present and symptoms are positionally related.

Small Fiber Neuropathy Screening. Consider SFN testing if:

- Burning pain, paresthesias, or allodynia
- Symptoms in stocking-glove distribution
- Autonomic symptoms (sweating abnormalities, GI dysmotility, orthostatic intolerance)

Evaluation: Skin punch biopsy (intraepidermal nerve fiber density) is gold standard. Sudomotor function testing also useful.

Autoimmune Screening. Consider autoimmune workup if:

- Family history of autoimmune disease
- Symptoms suggesting specific autoimmune conditions
- Unexplained inflammatory markers

Basic panel: ANA, ENA panel, RF, anti-CCP, TPO antibodies, anti-gliadin/tTG.

Chronic Infection Evaluation. Consider viral reactivation workup if post-infectious onset or ongoing immune activation:

- EBV: VCA IgG, EBNA IgG, EA IgG (EA elevation suggests reactivation)
- CMV, HHV-6: IgG levels
- Tick-borne: Lyme and co-infections if exposure history

GI Dysmotility Screening. Screen for SIBO and gastroparesis if:

- Bloating, early satiety, nausea, constipation alternating with diarrhea
- Food intolerances or malabsorption symptoms

Testing: Hydrogen/methane breath test for SIBO; gastric emptying study if gastroparesis suspected.

18.4.3 Treatment Sequencing

[] Assumption 1: Septad Treatment Sequencing: MCAS First

Based on clinical experience rather than validated research, Kaufman proposes addressing mast cell activation syndrome (MCAS) before other Septad components, on the rationale that mast cell stabilization may reduce interconnected symptom burden across multiple domains. This sequencing has not been tested in controlled studies.

△ Warning 7: Framework Limitations

The Septad is a *clinical framework* based on expert observation, not a validated research model. Systematic prevalence data for each component in ME/CFS populations is lacking. Use for organizing comorbidity screening, not as diagnostic criteria for ME/CFS itself. PEM remains the hallmark diagnostic feature (Section 5.6.11).

18.5 Disease-Modifying Strategies for Mild-Moderate Cases

18.5.1 Early Intervention Advantage

Mild-moderate patients have a critical advantage: potential to intervene before immune exhaustion phase (Achievement 7.3.1). This provides opportunity for disease modification rather than pure symptom management.

The Front-Loading Strategy

The striking difference in recovery rates between pediatric and adult ME/CFS patients (54–94% [504] versus ≤22% [54]) suggests that biological plasticity plays a critical role in determining outcomes. While some of this advantage may be inherent to developing biology, pediatric care patterns offer a potentially actionable insight: children are typically diagnosed earlier and treated more aggressively from the outset.

~ Hypothesis 3: Front-Loading Treatment Intensity

Certainty: 0.35. The front-loading strategy (concentrating intensive intervention in the first 6–12 months post-onset) may improve outcomes compared to traditional conservative sequential treatment. The certainty level reflects: (1) observational evidence linking early diagnosis to better outcomes; (2) theoretical basis from Recovery Capital model and critical window phenomena; (3) however, lack of randomized controlled trials directly comparing front-loading versus conservative approaches; (4) inability to control for confounding (early-diagnosed patients may have milder disease or better prognostic markers); (5) substantial treatment intensity carries risks including medication interactions and PEM from over-intervention; (6) unclear whether front-loading truly alters trajectory or merely

benefits naturally-recovering patients.

The “front-loading” strategy inverts the traditional incremental approach to ME/CFS treatment. Rather than starting conservatively and escalating over months or years, this approach concentrates treatment intensity in the first 6–12 months after symptom onset, aiming to maximize intervention during the hypothesized window of biological plasticity.

Rationale: The Recovery Capital model (Speculation 23) proposes that patients begin with finite biological reserves that deplete over time with crashes and chronic illness. If correct, early aggressive intervention—before significant reserve depletion—would have greater efficacy than the same interventions applied later. Pediatric outcomes may partly reflect this timing advantage.

Core principle: Treat early ME/CFS as a medical emergency requiring immediate comprehensive intervention, not a chronic condition warranting gradual symptom management.

Contrasting Treatment Philosophies Traditional ME/CFS management follows a conservative sequential approach:

- **Conservative/Sequential strategy:**
 1. Start single intervention (e.g., pacing education only)
 2. Wait 8–12 weeks for assessment
 3. If inadequate response, add second intervention
 4. Wait another 8–12 weeks
 5. Repeat until sufficient improvement or interventions exhausted
 6. Timeline: 6–24 months to reach multi-modal treatment
- **Front-loading strategy:**
 1. Initiate 4–6 interventions simultaneously within first 4 weeks
 2. Aggressive dose optimization (target complete symptom resolution, not partial improvement)
 3. Monthly reassessment
 4. Begin taper at 6–12 months if sustained improvement
 5. Timeline: Multi-modal treatment from day 1

Rationale for inversion: If Recovery Capital depletes over time (Speculation 23), the conservative strategy may expend the therapeutic window during the assessment phase. By the time multi-modal treatment is reached (6–24 months), biological reserves may be insufficiently depleted to respond effectively. Front-loading trades methodological clarity (inability to isolate which interventions work) for potential preservation of the intervention window.

Key difference from “try everything randomly”: Front-loading is NOT unstructured polypharmacy. It follows a systematic protocol with:

- Evidence-based intervention selection
- Standardized dosing

- Structured monitoring
- Planned taper protocol (see Section 18.5.1)
- Clear safety parameters

Front-Loading Protocol Components

1. Immediate maximal orthostatic intolerance treatment:

- Do not wait for behavioral approaches (increased fluids, compression) to “fail” before adding pharmacotherapy
- Initiate fludrocortisone + midodrine within first 2 weeks if OI symptoms present
- Target: Complete resolution of orthostatic symptoms, not partial improvement
- Rationale: OI may be an upstream driver (see Hypothesis 18.5.1); early correction may prevent downstream system involvement, though this causal relationship has not been established

2. Strict pacing enforcement from diagnosis:

- Goal: Zero crashes during the front-loading window (first 6–12 months)
- Each crash consumes Recovery Capital that may be irreplaceable
- Use HRV monitoring (see Protocol 18.3.3) for objective activity guidance
- Consider temporary disability leave if work is causing envelope violations

3. Aggressive sleep optimization:

- Sleep study within first month to identify treatable disorders
- Pharmacological support (low-dose trazodone, melatonin) initiated early if sleep is impaired
- Target: 7–9 hours with $\geq 85\%$ sleep efficiency
- Do not wait months to see if sleep “improves on its own”

4. Anti-inflammatory support from baseline:

- Low-dose naltrexone (titrate to 4.5mg over 4 weeks)
- High-dose omega-3 fatty acids (2–4g EPA+DHA daily)
- Mast cell stabilization (H1 + H2 antihistamines)
- Mediterranean-style anti-inflammatory diet

5. Subtype-specific interventions if indicated:

- If viral reactivation markers elevated: antivirals early, not as last resort
- If GPCR autoantibodies detected: consider immunomodulation referral
- If small fiber neuropathy documented: IVIG evaluation if accessible

Monitoring During Front-Loading Phase

- Monthly clinic visits (in-person or telehealth) for first 6 months
- Biomarker reassessment at 3 and 6 months
- Continuous activity and HRV monitoring via wearables

- Crash log with severity classification (Table 18.3)
- Medication adherence tracking

△ **Warning 8: Front-Loading is Hypothesis-Driven**

The front-loading strategy is informed by pediatric outcome data and the Recovery Capital model but has not been validated in randomized trials. It represents a reasoned extrapolation from available evidence, not proven treatment.

Key uncertainties:

- Whether the pediatric advantage is due to treatment timing or inherent developmental biology
- Whether adult biological plasticity can be preserved or enhanced through early intervention
- Optimal duration of the front-loading window
- Which components are essential versus optional

Methodological Trade-offs The front-loading strategy accepts several methodological limitations:

1. **Attribution problem:** When 5+ interventions are initiated simultaneously, it becomes impossible to determine which components drove improvement. If patient improves, unclear whether all interventions were necessary or only a subset.
Consequence: Cannot confidently discontinue “non-essential” interventions during taper phase. Conservative approach preserves ability to identify effective interventions.
2. **Adverse event attribution:** If patient experiences side effects or worsening, difficult to isolate culprit intervention. May require discontinuation of multiple agents simultaneously.
3. **Cost and adherence burden:** Initiating multiple medications/supplements simultaneously increases:
 - Monthly costs (\$200–\$500+ depending on insurance coverage)
 - Pill burden (10–15 pills daily)
 - Complexity of medication schedule
 - Risk of non-adherence
4. **Nocebo and medicalization risk:** Aggressive early intervention may reinforce illness identity in patients who might have spontaneously recovered. However, given low spontaneous recovery rates (5%) [54], this risk is likely small relative to potential benefit of preserving Recovery Capital.

The core trade-off: Front-loading prioritizes *speed* over *attribution*. If the therapeutic window is narrow and Recovery Capital finite, this trade-off may be justified despite methodological limitations.

A randomized trial testing front-loading versus standard care is proposed in Chap-

ter 33, Section 33.4.

Taper Protocol: Systematic Intervention Reduction

If front-loading achieves sustained symptom improvement, the next question becomes: Which interventions must continue long-term, and which can be safely discontinued? Given the attribution problem (inability to isolate which interventions drove improvement), taper must be systematic and cautious.

► Protocol 5: Front-Loading Taper Protocol

Eligibility criteria for initiating taper:

- Minimum 6 months sustained improvement on front-loading protocol
- Zero crashes for ≥ 3 consecutive months
- Stable function at 70–90% of pre-illness baseline
- Patient willing to accept risk of symptom return
- Physician supervision available for monitoring

DO NOT INITIATE TAPER IF:

- Still experiencing crashes (even mild)
- Function unstable or declining
- Less than 6 months since starting protocol
- Major life stressor ongoing (job change, relocation, etc.)

Taper sequence (one intervention per month):

Phase 1: Reduce symptom-specific agents first (Months 1–3)

1. Month 1: Taper sleep medications (if using)

- Rationale: If sleep has normalized, medications may no longer be necessary
- Method: Reduce dose by 50% for 2 weeks, then discontinue if sleep remains stable
- Monitoring: Sleep diary, sleep efficiency calculation
- Reversal criterion: If sleep efficiency drops below 80% for ≥ 1 week, reinstate medication

2. Month 2: Reduce H2 antihistamine (famotidine)

- Rationale: H1 blocker (cetirizine) provides primary mast cell stabilization; H2 may be redundant in stable patients
- Method: Discontinue directly (minimal withdrawal risk)
- Monitoring: Histamine symptoms (flushing, GI issues, headaches)
- Reversal criterion: Return of histamine symptoms for ≥ 3 days

3. Month 3: Consider pain medication reduction (if using)

- Rationale: If pain has resolved, medications may be unnecessary
- Method: Taper dose by 25% every 2 weeks
- Monitoring: Pain severity scores
- Reversal criterion: Pain returns to pre-treatment levels

Phase 2: Test core interventions (Months 4–8) **CRITICAL:** The following interventions are hypothesized to be disease-modifying. Taper cautiously and expect possible delayed symptom return (2–4 weeks).

1. **Month 4:** Reduce omega-3 fatty acids
 - Taper from 4g to 2g daily (maintenance dose)
 - Full discontinuation NOT recommended (omega-3 has general health benefits)
 - Monitoring: Inflammatory symptoms (joint pain, brain fog)
2. **Month 5:** Trial LDN discontinuation
 - Method: Reduce from 4.5mg to 3mg for 2 weeks, then 1.5mg for 2 weeks, then discontinue
 - Monitoring: Fatigue levels, pain, immune symptoms
 - Reversal criterion: Return of core ME/CFS symptoms for ≥ 2 weeks
 - Note: Many patients require long-term LDN; discontinuation frequently unsuccessful
3. **Month 6:** Consider mitochondrial cofactor reduction
 - Taper CoQ10 from 200mg to 100mg, continue NADH 20mg
 - Monitoring: Energy levels, exercise tolerance, cognitive fatigue
 - Reversal criterion: Return of fatigue or PEM
4. **Month 7–8:** Consider OI medication reduction (HIGH RISK)
 - **WARNING:** OI medications are frequently required long-term. Discontinuation often results in symptom return.
 - Only attempt if orthostatic symptoms have been completely absent for ≥ 6 months
 - Method: Reduce fludrocortisone by 50% (e.g., 0.1mg to 0.05mg) for 4 weeks
 - Monitoring: Daily orthostatic vitals (HR/BP supine and standing), symptom tracking
 - Reversal criterion: Return of orthostatic symptoms, HR increase > 30 bpm on standing
 - If stable after 4 weeks at reduced dose, consider full discontinuation
 - Expect potential delayed relapse (OI symptoms may return 2–8 weeks after discontinuation)

Phase 3: Maintenance determination (Month 9+) After taper attempts, reassess which interventions appear necessary for sustained stability:

- **High likelihood of long-term need:**

- OI medications (if POTS/OI was prominent)
- Low-dose naltrexone (frequently required indefinitely)
- Pacing strategies (always maintain activity envelope awareness)

- **Moderate likelihood of long-term need:**

- Mitochondrial cofactors (CoQ10, NADH)
- Anti-inflammatory support (omega-3, maintenance LDN)
- Mast cell stabilization (H1 antihistamine)

- **Lower likelihood of long-term need:**

- Sleep medications (if sleep normalized)
- H2 antihistamines (if H1 sufficient)
- High-dose supplements beyond maintenance levels

Individualization required: Taper sequence should be adapted based on:

- Patient's symptom profile (which interventions target their primary symptoms)
- Response pattern (which interventions produced clearest subjective benefit)
- Cost and burden considerations
- Patient preference

Expected Outcomes of Taper Process

→ Recommendation 10: Taper Protocol Outcomes

Scenario 1: Successful taper (estimated 20–30%):

- Able to discontinue 50–75% of interventions without symptom return
- Identify minimal maintenance regimen (typically: pacing awareness, 1–2 core medications)
- Sustained improvement at 12+ months

Scenario 2: Partial taper (estimated 40–50%):

- Able to discontinue symptom-specific agents (sleep, pain meds)
- Require ongoing core interventions (OI meds, LDN, mitochondrial support)
- Stable function with reduced but ongoing treatment burden

Scenario 3: Minimal taper tolerance (estimated 20–30%):

- Symptoms return rapidly with any intervention reduction
- Require long-term multi-modal treatment for stability
- Front-loading achieved stabilization but not resolution; ongoing management necessary

CRITICAL: Inability to taper does NOT indicate front-loading “failed.” If patient achieved sustained stabilization with multi-modal treatment, this represents success

even if interventions must continue indefinitely. The alternative (not using interventions) would likely result in ongoing instability or deterioration.

Taper Failures and Re-escalation

△ **Warning 9: Responding to Symptom Return During Taper**

If symptoms return during taper process:

1. **Immediate re-escalation:**
 - Reinstate the most recently tapered intervention at full dose
 - Do not wait to see if symptoms “stabilize on their own”
 - Resume for minimum 4–8 weeks before considering another taper attempt
2. **If symptoms do NOT resolve with re-escalation:**
 - Consider whether disease has progressed independent of taper
 - Reassess for new comorbidities or stressors
 - May need to reinstate multiple interventions or add new ones
3. **Multiple taper failures:**
 - If 2+ attempts to taper a specific intervention result in symptom return, accept that intervention is likely required long-term
 - Shift focus to optimizing adherence and minimizing burden rather than discontinuation

Relationship to Recovery Capital Model The taper protocol tests the Recovery Capital hypothesis (Speculation 23):

- **If taper is well-tolerated:** Suggests Recovery Capital was preserved or restored; biological reserve sufficient to maintain stability without ongoing intervention.
- **If taper causes symptom return:** Suggests either:
 1. Recovery Capital remains depleted; ongoing support required to maintain function
 2. Interventions are actively managing underlying pathology that has not resolved
 3. Disease has transitioned to chronic self-sustaining state despite intervention

Taper outcomes could provide indirect evidence for or against Recovery Capital depletion as core mechanism. A trial systematically tracking taper success rates would be valuable (see Chapter 33).

18.5.2 Acute Onset Protocol: The Critical First Six Months

For patients within 6 months of ME/CFS symptom onset, the evidence suggests a narrow therapeutic window where aggressive intervention may alter disease trajectory. While the front-loading strategy (Section 18.5.1) applies to all mild-moderate patients, acute-onset cases warrant an even more intensive, time-sensitive approach.

★ Achievement 3: Diagnostic Delay Predicts Recovery: Evidence from Longitudinal Cohort

Castro-Marrero et al. [403] tracked 168 ME/CFS patients over median 55-month follow-up, identifying diagnostic delay as the most significant modifiable prognostic factor. Patients who achieved recovery or improvement had median diagnostic delay of 23 months versus 55 months for non-recovered patients ($p=0.0004$). Multivariate analysis confirmed diagnostic delay inversely associated with recovery/improvement (OR 0.98 per month, $p=0.036$), with overall recovery rate of 8.3% and improvement rate of 4.8%.

Clinical implication: Every month of delay reduces recovery probability. Early diagnosis and intervention are not merely beneficial—they may be decisive.

Rationale for Acute Intervention

Three converging lines of evidence support time-sensitive intervention in newly diagnosed ME/CFS:

1. **Critical window phenomenon:** Diagnostic delays beyond 23 months correlate with substantially worse outcomes [403], suggesting a therapeutic window in the first 2 years, with the first 6 months potentially most critical.
2. **Recovery Capital preservation:** Each crash and month of illness depletes finite biological reserves (Speculation 23). Early intervention aims to prevent depletion before reserves become irreversibly exhausted.
3. **Cascade prevention:** The cytokine duration hypothesis (Achievement 7.3.1) proposes that prolonged immune activation triggers secondary pathology. Early intervention targets the initial trigger before cascade progression.

Acute Onset Protocol Components

► Protocol 6: Intensive Early Intervention for Acute-Onset ME/CFS

Certainty: 0.40. Early aggressive intervention in the first 6 months of ME/CFS may alter disease trajectory and improve recovery probability. The certainty level reflects: (1) observational evidence linking diagnostic delay to worse outcomes; (2) theoretical basis from critical window phenomena in other post-viral illnesses; (3) however, lack of randomized controlled trials testing early intensive intervention protocols; (4) recovery rates even with intervention remain modest (8–13%); (5) inability to distinguish whether early intervention enables recovery or merely selects for spontaneously recovering patients; (6) substantial individual variation in disease trajectory independent of intervention timing.

Eligibility criteria:

- Symptom onset <6 months prior
- Meets IOM 2015 or Canadian Consensus diagnostic criteria
- Mild to moderate severity (ambulatory, not bedbound)

- No contraindications to protocol components

Timeline and implementation:

Weeks 1–2: Immediate Stabilization

1. Strict rest enforcement:

- Reduce activity to 50% of pre-illness baseline immediately
- No exercise; gentle stretching only if tolerated
- Consider medical leave from work/school if feasible
- Rationale: Prevent crashes during critical window; allow initial physiological stabilization

2. Orthostatic intolerance screening and treatment:

- NASA Lean Test (Requirement 4.6.2) within first week
- If positive: Initiate aggressive OI treatment immediately (fluids, salt, compression, consider fludrocortisone/midodrine without waiting for behavioral measures to “fail”)
- Rationale: OI may be upstream driver (Keypoint 13.4.4); early correction prevents cumulative orthostatic stress

3. Crash prevention education:

- PEM symptom recognition
- Activity envelope concept
- Heart rate monitoring introduction
- Rationale: Knowledge prevents accidental envelope violations

Weeks 3–4: Foundation Building

1. Mitochondrial support initiation:

- Coenzyme Q₁₀ 200 mg + NADH 20 mg daily
- Evidence: RCT (n=207) demonstrated significant improvements in cognitive fatigue (p<0.001), overall fatigue (p=0.022), quality of life (p<0.05), and sleep [7]
- Use pharmaceutical-grade formulations (bioavailability critical) [525]
- Additional mitochondrial cofactors: B-complex vitamins, magnesium glycinate 400mg, alpha-lipoic acid 600mg

2. Anti-inflammatory strategy:

- Low-dose naltrexone: Initiate 1.5mg, titrate to 4.5mg over 4 weeks
- Omega-3 fatty acids: 2–4g EPA+DHA daily
- H1 + H2 antihistamines for mast cell stabilization (cetirizine 10mg + famotidine 20mg daily)
- Rationale: Address documented inflammatory signatures; prevent transition to chronic immune activation phase

3. Sleep optimization:

- Sleep study if sleep quality impaired (do not delay)
- Pharmacological support if needed: melatonin 0.5–3mg, low-dose trazodone 25–50mg
- Circadian light therapy (10,000 lux within 30 minutes of waking)
- Target: 7–9 hours with ≥85% sleep efficiency

Weeks 5–8: Stabilization Assessment

1. Activity ceiling establishment:

- HRV-guided activity monitoring (Protocol 18.3.3)
- Gradual identification of sustainable baseline
- Goal: Find maximum activity level that produces ZERO crashes
- Stay at this ceiling; do not attempt to expand yet

2. Subtype-specific interventions:

- CNS-primary: Prioritize cognitive support, intranasal therapies if available
- Autonomic-primary: Maximize OI treatment, consider beta-blockers if POTS documented
- Peripheral-primary: Emphasize mitochondrial support, consider L-carnitine 2g daily
- Global: All interventions in parallel

3. Clinical monitoring:

- Weekly symptom logs (fatigue severity, PEM frequency, orthostatic symptoms)
- Crash tracking with severity classification
- Medication tolerance assessment
- Quality of life measures (SF-36 or similar)

Months 3–6: Consolidation and Expansion

1. Reassess diagnostic accuracy:

- Confirm ME/CFS diagnosis vs. other post-viral fatigue
- Screen for comorbidities (Septad framework, Section 18.4)
- Biomarker reassessment if available

2. Activity expansion (if stable):

- If zero crashes for 4+ consecutive weeks, cautiously test activity expansion
- Increase by 10% maximum, monitor for 2 weeks before further increase
- Retreat immediately if PEM occurs
- Do NOT attempt expansion if still experiencing crashes

3. Long-term strategy development:

- Transition from acute crisis management to chronic disease management if needed
- Identify sustainable pacing baseline
- Plan work/study accommodations if return not yet feasible
- Psychological support for adjustment to chronic illness if recovery incomplete

Expected Outcomes and Realistic Expectations

→ Recommendation 11: Acute Onset Protocol: Outcomes and Limitations

Note: The following outcome estimates are author-generated projections extrapolated from general ME/CFS recovery literature and diagnostic delay studies. They have not been validated for this specific protocol.

Best-case scenario (estimated 10–20% based on recovery literature):

- Substantial symptom reduction by 6 months
- Return to 70–90% of pre-illness function
- Ability to resume work/study with modifications
- Continued slow improvement over 12–24 months

Moderate response (estimated 30–40%):

- Stabilization without progression to severe disease
- Functional improvement to sustainable mild-moderate level
- Reduced crash frequency and severity
- Improved quality of life despite ongoing limitations

Minimal response (estimated 40–50%):

- Disease progression halted but limited symptom improvement
- Persistent mild-moderate severity requiring ongoing management
- Need for long-term accommodations and lifestyle modification

CRITICAL CAVEAT: These are rough estimates extrapolated from recovery literature and diagnostic delay data. The acute onset protocol has NOT been validated in randomized trials. Individual outcomes remain highly variable and unpredictable.

Safety Considerations and Contraindications

△ Warning 10: Acute Onset Protocol Safety

Monitoring requirements:

- Monthly physician visits during first 6 months (minimum)
- Blood pressure monitoring if on fludrocortisone/midodrine

- Liver function tests at baseline and 3 months if on multiple supplements
- Mental health screening (depression/anxiety common in acute illness)

Contraindications to specific components:

- Fludrocortisone: Heart failure, hypertension, hypokalemia
- Low-dose naltrexone: Concurrent opioid use, acute hepatitis
- High-dose omega-3: Bleeding disorders, anticoagulant therapy (reduce dose)
- CoQ10: Warfarin interaction (monitor INR closely)

Risk of over-restriction: Complete bed rest is NOT recommended. Goal is activity reduction to sustainable level, not total inactivity. Prolonged complete bed rest risks deconditioning, orthostatic intolerance worsening, and psychological harm. Maintain gentle movement within energy envelope.

Psychological impact: Aggressive medical intervention in newly diagnosed patients can provoke anxiety or medicalization concerns. Ensure patient understands: (1) Protocol is hypothesis-driven, not proven; (2) They retain decision-making autonomy; (3) Protocol can be modified based on tolerance and response.

Evidence Status and Research Needs

The acute onset protocol synthesizes established interventions (pacing, OI treatment, mitochondrial support) with timing optimization based on prognostic data. Individual components have varying evidence levels:

- **HIGH certainty:** CoQ10+NADH efficacy [7], diagnostic delay impact [403], pacing principles
- **MEDIUM certainty:** OI treatment benefits, LDN efficacy, anti-inflammatory interventions
- **LOW certainty:** Optimal timing window, activity restriction duration, combination synergy

CRITICAL RESEARCH NEED: Randomized controlled trial comparing acute onset protocol versus standard care in newly diagnosed ME/CFS patients (<6 months onset). Primary outcome: Functional status at 12 and 24 months. Such a trial is proposed in Chapter 33.

Until such evidence exists, this protocol represents reasoned clinical extrapolation from available data, not evidence-based standard of care.

When NOT to Use Front-Loading Strategy**△ Warning 11: Front-Loading Contraindications**

The front-loading strategy is NOT appropriate for all patients. Specific contraindications:

1. Severe or very severe patients:

- Bedbound or housebound patients
- Rationale: Severe patients are already beyond the hypothesized intervention window; front-loading unlikely to restore lost Recovery Capital. Priority shifts to preventing further deterioration and managing symptoms. See Chapter 17 for severe patient management.
- Exception: Acute sudden deterioration in previously stable patient (consider ICU-level stabilization protocol)

2. Limited financial resources:

- Front-loading costs \$200–\$500+ monthly (supplements, medications, monitoring)
- Rationale: If cost burden prevents adherence or causes financial stress (itself harmful), conservative sequential approach may be more sustainable.
- Alternative: Prioritize highest-yield interventions (OI treatment, pacing, LDN) rather than full front-loading protocol

3. Limited medical supervision access:

- Front-loading requires monthly physician monitoring (minimum)
- Rationale: Simultaneous multi-drug initiation carries higher risk of adverse events; close monitoring essential for safety
- If only quarterly appointments available, use conservative sequential approach

4. Significant comorbidities complicating treatment:

- Severe cardiac disease (fludrocortisone/midodrine contraindicated)
- Liver disease (LDN contraindicated; supplement metabolism impaired)
- Bleeding disorders (high-dose omega-3 contraindicated)
- Multiple drug allergies or intolerances
- Rationale: Contraindications to multiple protocol components reduce feasibility; safer to use sequential approach with careful selection

5. Patient preference for conservative approach:

- Some patients prefer methodical single-intervention trials to identify what works
- Rationale: Patient autonomy is paramount. Front-loading is hypothesis-driven, not proven. Patients uncomfortable with aggressive multi-modal approach should not be pressured.
- Physician should explain potential trade-offs (time to multi-modal treatment

vs. therapeutic window), but ultimately respect patient decision.

6. High risk of non-adherence:

- Cognitive impairment severe enough to interfere with medication management
- History of poor medication adherence
- Lack of caregiver support for complex regimen
- Rationale: Non-adherent front-loading is worse than adherent conservative approach. If patient unlikely to maintain 10–15 pill daily regimen, simpler protocol is safer and more effective.

7. Diagnostic uncertainty:

- If ME/CFS diagnosis not yet confirmed (still in differential diagnosis phase)
- Rationale: Front-loading is specific to ME/CFS pathophysiology. If diagnosis uncertain, aggressive protocol may be inappropriate for actual underlying condition.
- Exception: Post-viral fatigue in acute phase (<3 months) may warrant early intervention even before ME/CFS diagnosis confirmed, if trajectory suggests progression to chronic illness.

Alternative for contraindicated patients: Use prioritized sequential approach targeting highest-yield interventions first:

1. Pacing education and activity envelope establishment (zero cost, universal benefit)
2. Orthostatic intolerance treatment if OI present (often most impactful single intervention)
3. Low-dose naltrexone (low cost, broad benefits, good safety profile)
4. Add additional interventions sequentially as resources and monitoring allow

This approach preserves some potential for early intervention while accommodating resource constraints and safety considerations.

18.5.3 “Brain First” Implementation Protocol for Mild-Moderate Cases

► Protocol 7: “Brain First” Sequential Treatment for Optimal ME/CFS Recovery

Patient-Derived Insight and Rationale Patient experience and emerging mechanistic evidence (Hypothesis 8.1.6) suggest that addressing central neuroinflammatory dysfunction before peripheral symptoms optimize treatment efficacy and patient capacity to participate in own care. The “brain first” approach inverts the typical symptom-by-symptom escalation, prioritizing cognitive and neurological stability as the foundation for all subsequent interventions.

Week 1–4: Low-Dose Aspirin (LDA) Titration with Cognitive Baseline

- **Medication protocol:** Begin LDA 0.25 mg daily; escalate by 0.25 mg every 3 days to target 1.5 mg daily by end of week 4. Monitor closely for GI intolerance (rare at

these doses but possible).

- **Cognitive assessment baseline:** At week 1 start and week 4 end, perform brief cognitive battery to establish trajectory:
 - Montreal Cognitive Assessment (MoCA) or similar screening tool
 - Timed naming task (Boston Naming Test)
 - Digit span (forward and backward)
 - Self-reported fog severity (0–10 scale)
- **Goal:** Establish that LDA is tolerated and beginning to improve central cognitive clarity. This creates confidence that treatment is working and readies the patient for subsequent layering.
- **Evidence:** Low-dose aspirin targets platelet-mediated thromboinflammation and may reduce circulating microparticles that trigger neuroinflammation [278, 279].

Week 4–8: Low-Dose Naltrexone (LDN) Addition with Psychiatric Monitoring

- **Medication protocol:** Begin LDN 0.5 mg at bedtime; titrate slowly (increase by 0.5 mg every 4–7 days) to target 2 mg by end of week 8. Go slower than in severe cases (where 4.5 mg is target) to avoid destabilization in patients with psychiatric comorbidities.
- **Psychiatric monitoring:** Microglial downregulation can unmask underlying mood pathology (anxiety, depression, emotional lability). Establish baseline mood (PHQ-9, GAD-7) and weekly check-in for mood changes. Educate patient that mood instability does not mean treatment failure but rather microglial restoration allowing underlying pathology to surface.
- **Cognitive expectation:** By week 8, patients often report further cognitive improvement (clearer thinking, reduced executive dysfunction). The combination of LDA + LDN appears synergistic for cognition.
- **Evidence:** LDN restores endogenous opioid tone and downregulates microglial activation; addition to LDA provides complementary mechanisms [279].

Week 8–12: Mestinon (Pyridostigmine) Addition for Autonomic Stabilization

- **Medication protocol:** Begin pyridostigmine 20 mg three times daily (TID); can escalate to 30–60 mg TID depending on tolerance. Monitor for cholinergic side effects (GI cramping, rhinorrhea, salivation); reduce dose if intolerable.
- **Expected effect:** Mestinon enhances acetylcholine availability at the neuromuscular junction and autonomic terminals, supporting both cognitive function (acetylcholine is essential for attention and memory) and autonomic stability. Patients often report reduced orthostatic intolerance and improved cognitive processing speed.
- **Key principle:** By this point (week 8), central dysfunction is partially restored via LDA+LDN, patient is engaged in their treatment, and cognitive clarity allows them to perceive autonomic symptoms with less cognitive noise. Adding Mestinon at

this stage capitalizes on restored cognition.

- **Evidence:** Cholinesterase inhibition supports both CNS and autonomic function; small studies suggest benefit in ME/CFS-like conditions [278].

Week 12+: Mast Cell Stabilization Layer

- **Medication protocol:** Add H1 antihistamine (cetirizine 10 mg BID) + H2 blocker (famotidine 20 mg BID) + mast cell stabilizer (ketotifen 1 mg BID or cromolyn 100 mg QID if available).
- **Rationale for late addition:** By week 12, central and autonomic stabilization is underway, patient cognition is improved, and baseline neuroinflammation is reduced by LDA+LDN. At this point, addressing peripheral mast cell activation has clearer effects and is less likely to be obscured by ongoing central dysfunction.
- **Expected outcomes:** Further reduction in allergic symptoms, GI symptoms, and generalized pain. Patients report improved food tolerance and reduced temperature dysregulation.
- **Evidence:** Multi-modal mast cell stabilization provides synergistic reduction in MCAS/MCAD symptoms common in ME/CFS [279].

Key Principle: Sequential Stabilization Rather than Parallel Escalation The “brain first” protocol differs from standard care in a critical way: **each layer builds on the previous layer’s success**. Rather than adding all medications simultaneously (which can cause overwhelming side effects and poor adherence), this approach:

1. Establishes that patient can tolerate and benefit from foundational treatment (LDA)
2. Adds second layer (LDN) that synergizes with first
3. Only after central stability, adds autonomic support (Mestinon)
4. Final layer addresses peripheral mast cell pathology from a more stable CNS baseline

This sequencing allows patient to identify which component is providing benefit (if side effects emerge, the timing pinpoints the culprit) and creates psychological momentum as patients observe improvement at each step.

Expected Timeline and Outcomes

- **Weeks 1–4:** Mild cognitive improvement; patient sees treatment is working
- **Weeks 4–8:** Cognitive clarity, reduced brain fog; mood instability if present and self-limited
- **Weeks 8–12:** Autonomic symptoms (dizziness, palpitations) reduce; fatigue may improve as cognition improves (less central fatigue drive)
- **Weeks 12+:** Peripheral symptoms (allergies, pain, GI dysfunction) become more apparent as central symptoms quiet; mast cell therapy addresses these
- **3–6 months:** Many patients report substantial functional improvement and may be able to increase activity within pacing guidelines

Integration with Other Interventions The “brain first” protocol provides the foundational CNS stabilization. It should be combined with:

- **Strict pacing:** See Section 18.3.3. Do NOT use cognitive improvement as a reason to increase activity; restrict to HR-guided limits.
- **Sleep optimization:** Sleep study and pharmaceutical support if needed; sleep quality amplifies LDA+LDN benefits.
- **Comorbidity screening:** See Section 18.4. Treat identified comorbidities concurrently (thyroid dysfunction, vitamin deficiencies, sleep apnea).
- **Immune profiling:** Parallel to this protocol, obtain immune biomarkers to guide longer-term disease-modifying strategy (see Chapter 7 for biomarker discussion).

Observation 82 (Treatment Sequencing and Patient Capacity: Community Evidence for Prioritization Logic). Patient experience and clinical observation suggest that treatment sequencing significantly impacts both efficacy and tolerability. While the “brain first” protocol (Section 18.5.3) is mechanistically justified, community-derived evidence supports this sequencing through a different lens: patient capacity and engagement.

The Sequencing Rationale The logical treatment sequence appears to be:

1. **Cognition first (LDA):** Cognitive dysfunction and brain fog are so pervasive in ME/CFS that they impair patient’s ability to participate in their own care—tracking symptoms, recognizing patterns, managing medication adherence. Addressing cognition first enables all downstream interventions to succeed. Patients report that improved cognition allows them to “understand what’s happening” and recognize other improvements.
2. **Fatigue second (LDN):** Once cognition improves, the overwhelming fatigue burden becomes more apparent and limiting (it was previously masked by cognitive chaos). Addressing fatigue-driving neuroinflammation (LDN mechanism) at this stage provides rapid quality-of-life improvement and further increases engagement.
3. **Muscle weakness and autonomic dysfunction third (Mestinon):** With cognitive and fatigue improvements, functional limitations from muscle weakness and orthostatic intolerance become the limiting factors. Mestinon’s cholinergic support addresses both. At this point, patients have capacity to engage in activity retraining within paced envelopes.
4. **Peripheral symptom layer (mast cell stabilization):** Only once CNS-driven symptoms are partially controlled do isolated mast cell symptoms clearly differentiate themselves. Patients can then specifically target allergic, GI, and inflammatory symptoms.

Why Parallel Escalation Fails Standard medical practice is to add all indicated medications simultaneously. In ME/CFS, this approach often fails because:

- **Cognitive overload:** Patient cannot track which medication is causing which side effect (all added together)
- **Overwhelmed system:** Severe patients especially cannot tolerate multiple new medications; cumulative effects trigger crashes

- **Lost engagement:** Patient becomes discouraged when improvements are not clearly attributable to specific interventions
- **Suboptimal dosing:** To avoid overwhelming effects, patients end up on subtherapeutic doses of each medication

Sequential layering addresses each of these by allowing patient to stabilize, identify benefit, and then add the next piece.

Clinical Classification Within Sequencing This observation is **community knowledge** rather than randomized evidence, but the mechanistic rationale aligns with the neuroinflammatory cascade model (Hypothesis 8.1.6): central dysfunction drives peripheral symptoms in ME/CFS. Therefore, addressing central dysfunction first has mechanistic support and appears clinically superior to parallel escalation in patient report.

A formal trial comparing sequential versus parallel escalation would establish whether this observation represents genuine efficacy advantage or selection bias in reporting.

18.5.4 Immune Profiling and Targeted Intervention

Recommended Testing

- **Basic panel:**
 - CBC with differential
 - Comprehensive metabolic panel
 - Thyroid function (TSH, free T4, free T3)
 - Iron studies (ferritin, iron, TIBC)
 - Vitamin D, B12, folate
- **Immune panel** (if accessible):
 - Lymphocyte subsets (CD4, CD8, NK cells)
 - Immunoglobulins (IgG, IgA, IgM)
 - ANA, ENA panel (screening for autoimmunity)
 - Inflammatory markers (CRP, ESR)
- **Advanced panel** (if pursuing aggressive treatment):
 - Cytokine panel (IL-6, IL-1 β , TNF- α , IL-10)
 - GPCR autoantibodies (CellTrend - Germany)
 - NK cell function assay
 - Viral reactivation markers (EBV EA, VCA IgG, CMV IgG)

18.5.5 Personalized Cycle Mapping: Precision Diagnostic Framework

The vicious cycle dynamics model (Chapter 2, §2.1, “Vicious Cycle Dynamics”) reveals that ME/CFS involves multiple reinforcing physiological cycles: mitochondrial, immune, auto-

nomic, neuroinflammatory, and endocrine. However, not every patient has all five cycles active. **Identifying which specific cycles are operating in each individual enables precision-targeted treatment**, avoiding unnecessary interventions while ensuring all active pathology is addressed.

This diagnostic framework represents a paradigm shift from empirical “try everything” approaches to biomarker-guided personalized medicine.

The Five-Cycle Diagnostic Battery

Cycle 1: Mitochondrial Dysfunction. **Diagnostic criteria:** Evidence of impaired ATP production, oxidative phosphorylation failure, or abnormal post-exertional metabolic response.

Tier 1 Testing (Accessible):

- **Two-day cardiopulmonary exercise test (2-day CPET):** Gold standard
 - Day 2 VO₂max decline >5–10% = positive for mitochondrial cycle
 - Reduced ventilatory efficiency (VE/VCO₂ slope increase Day 2)
 - See Chapter 6 for interpretation
 - Accessibility: Limited to specialized centers; cost \$1,500–3,000
- **Lactate response to mild exertion:** Venous lactate before and 15–30 min after standardized activity (e.g., 6-minute walk, stationary bike at low resistance)
 - Lactate increase >30% from baseline = glycolytic shift, suggests mitochondrial impairment
 - Accessibility: Any laboratory can measure lactate; cost \$20–50
- **Actigraphy with recovery tracking:** 7–14 days continuous activity monitoring
 - Prolonged recovery periods (>24–48h) after modest activity
 - Boom-bust pattern (activity followed by crash)
 - Accessibility: Consumer-grade accelerometers (fitbit, etc.); cost \$50–150

Tier 2 Testing (Research or Specialized Centers):

- **Cellular ATP production:** Extracellular flux analysis (Seahorse assay) on PBMCs
 - Reduced maximal respiration, ATP-linked respiration
 - Research setting; not clinically available
- **Muscle biopsy:** Mitochondrial enzyme activities, electron microscopy
 - Reserved for unclear cases; invasive
 - Cost \$2,000–5,000; limited insurance coverage
- **Post-exertion metabolomics:** Plasma metabolites before and 24h after standardized exertion
 - NAD⁺/NADH ratio, acylcarnitines, TCA cycle intermediates
 - Research setting; cost \$500–2,000

Clinical decision: *If 2-day CPET shows Day 2 decline OR lactate increases post-exertion OR actigraphy shows prolonged recovery → Mitochondrial cycle ACTIVE → Target with CoQ10, NAD⁺ precursors, mitochondrial support stack.*

Observation 83 (Evidence-Based Mitochondrial Support Stack). Biomarker-driven supplementation targets documented ME/CFS metabolic deficits: reduced brain glutathione [193], impaired ATP production [1], and TCA cycle dysfunction [194]. The compounds described below address these specific deficits with differing evidence levels.

N-Acetylcysteine (NAC) for Glutathione Repletion

★ Achievement 4: Brain Glutathione Deficiency in ME/CFS

Magnetic resonance spectroscopy studies consistently document reduced brain glutathione (GSH) in ME/CFS patients. Shungu et al. [193] found 36% lower cortical GSH levels compared to healthy controls (n=15 vs n=13), with strong correlations to physical functioning ($\rho = 0.506$, $p = 0.001$) and energy levels ($\rho = 0.606$, $p < 0.001$). This finding was independently replicated by Godlewska et al. [526] using higher-resolution 7 Tesla MRS (n=22 vs n=13), which also revealed decreased total creatine and myo-inositol, suggesting concurrent energetic and glial dysfunction.

Brain GSH inversely correlates with ventricular lactate ($r = -0.545$, $p = 0.001$), implicating oxidative stress in pathophysiology [193].

Practical protocol: N-acetylcysteine 600–1200 mg two to three times daily with meals (total 1800–3600 mg/day). NAC provides cysteine, the rate-limiting amino acid for glutathione synthesis, and crosses the blood-brain barrier to support in situ GSH production. Pilot data showed 1800 mg/day normalized cortical GSH and improved symptoms ($p=0.006$) [486]. An NIH-funded RCT (NCT04542161, n=60) comparing doses (0/900/3600 mg/day) is expected to complete in 2026. Safety profile well-established (>30 years clinical use); common side effects include GI discomfort (~10%).

D-Ribose for ATP Regeneration

Speculation 58 (D-Ribose Accelerates ATP Recovery). **Certainty: 0.40.** D-ribose is a pentose sugar that serves as a substrate for de novo nucleotide synthesis, bypassing the rate-limiting step in ATP regeneration following energy depletion [527]. In ME/CFS, two open-label studies demonstrated large effect sizes: Teitelbaum et al. [8] found 45% energy improvement (n=41), subsequently replicated in a multicenter trial with 61.3% energy increase (n=257, $p<0.0001$) [528]. Animal studies demonstrate 85% ATP recovery at 24 hours with ribose supplementation versus 0% in controls [529].

Evidence limitations: Both ME/CFS studies were open-label without placebo control, yielding LOW-MEDIUM certainty despite large effect sizes. Placebo effects cannot be excluded.

Practical protocol: 5 g three times daily with meals (total 15 g/day). Effects typically begin within 1 week. Consider combination with CoQ10, L-carnitine, and magnesium for synergistic ATP support [530].

△ Warning 12: D-Ribose Contraindication: Diabetes and Hypoglycemia

D-ribose triggers insulin release paradoxically lowering blood glucose despite not being metabolized as glucose. **Contraindicated in diabetes mellitus (Types 1 and 2), hypoglycemia, or blood sugar instability.** Always take with meals to minimize blood sugar fluctuations.

L-Citrulline-Malate for TCA Cycle Support

Speculation 59 (Citrulline-Malate Addresses TCA/Urea Cycle Dysfunction). **Certainty: 0.35.** Metabolomic studies reveal significant TCA cycle dysfunction in ME/CFS, with reduced concentrations of citrate, isocitrate, and malate, alongside elevated ornithine/citrulline ratios indicating urea cycle impairment [194]. Citrulline-malate supplementation (6 g/day for 15 days) in fatigued individuals increased oxidative ATP production by 34% and phosphocreatine recovery by 20%, measured via ³¹P magnetic resonance spectroscopy [531]. The malate component acts as a TCA cycle intermediate, potentially bypassing anaplerotic bottlenecks; citrulline supports urea cycle function for ammonia detoxification.

Evidence limitations: No ME/CFS-specific intervention trials exist. Evidence extrapolated from metabolomics studies and exercise performance research.

Practical protocol: Start 3 g/day, target 6 g/day divided doses with meals. Minimum 2–4 weeks for metabolic adaptation. Well-tolerated up to 15 g/day; main side effect: mild GI discomfort (14.6% at high doses) [532].

Cycle 2: Immune Activation and Autoimmunity. **Diagnostic criteria:** Evidence of chronic immune activation, autoantibody production, or cytokine dysregulation.

Tier 1 Testing (Accessible):

- **GPCR autoantibodies:** β_2 -adrenergic, M3/M4 muscarinic receptors
 - CellTrend assay (Germany): Mail-order testing available
 - Elevated titers > 95th percentile = positive
 - Cost \$300–500; not covered by US insurance typically
 - *Critical biomarker:* Predicts response to immunoadsorption, daratumumab [147, 9]
- **Natural killer (NK) cell function:** Cytotoxicity assay or NK cell count
 - Reduced NK function or low CD56+ cell count = immune dysfunction
 - Flow cytometry available at many labs; cost \$150–300
- **Cytokine panel:** IL-6, IL-1 β , TNF- α , IL-10
 - Elevation indicates active inflammation
 - Accessibility: Some commercial labs (LabCorp, Quest); cost \$200–400
 - High variability; requires fasting sample, careful handling
- **Standard autoimmune screening:** ANA, ENA panel, rheumatoid factor
 - Positive ANA or ENA may indicate overlap syndrome

- Widely available; cost \$100–200

Tier 2 Testing (Specialized):

- **T cell and B cell subset analysis:** CD4/CD8 ratio, T cell exhaustion markers, B cell subsets
 - Flow cytometry; research or specialized immunology labs
 - Cost \$300–600
- **Viral reactivation markers:** EBV EA IgG, VCA IgG, HHV-6 IgG, CMV IgG
 - Chronic reactivation may drive immune activation
 - Available at commercial labs; cost \$200–400

Clinical decision: *If GPCR autoantibodies elevated OR NK function low OR cytokines elevated → Immune cycle ACTIVE → Consider immunoadsorption, daratumumab (if accessible), or LDN + anti-inflammatory stack.*

Cycle 3: Autonomic Dysregulation. Diagnostic criteria: Evidence of orthostatic intolerance, impaired heart rate variability, or sympathetic-parasympathetic imbalance.

Tier 1 Testing (Accessible):

- **NASA 10-minute lean test:** Modified poor man's tilt table test
 - Measure HR and BP supine, then standing at 2, 5, 10 minutes
 - HR increase >30 bpm or sustained increase >120 bpm = POTS
 - BP drop >20/10 mmHg = orthostatic hypotension
 - No cost; can be done at home or in any clinic
- **Heart rate variability (HRV):** Consumer-grade HRV monitor or smartphone app
 - Low HRV (particularly RMSSD <20–30 ms) = reduced parasympathetic tone
 - Tracking daily HRV identifies autonomic stress
 - Cost \$0–150 (many free apps using phone camera)
- **Symptom inventory:** Validated autonomic symptom scales (e.g., COMPASS-31)
 - Orthostatic lightheadedness, palpitations, GI dysmotility, temperature dysregulation
 - Free online questionnaires

Tier 2 Testing (Specialized):

- **Formal tilt table test:** 70-degree upright tilt for 10–45 minutes with continuous HR/BP monitoring
 - Gold standard for POTS and orthostatic hypotension diagnosis
 - Cardiology or autonomic specialty clinic; cost \$500–1,500
- **Quantitative sudomotor axon reflex test (QSART):** Measures small fiber autonomic function

- Detects autonomic neuropathy
- Specialized autonomic labs; cost \$500–1,000
- **Catecholamine levels:** Plasma or 24-hour urine norepinephrine, epinephrine, dopamine
 - Low levels support central catecholamine deficiency [5]
 - Available at commercial labs; cost \$150–300

Clinical decision: *If NASA lean test positive OR HRV chronically low OR catecholamines deficient → Autonomic cycle ACTIVE → Target with fludrocortisone, midodrine, compression, salt loading, L-tyrosine + BH4 cofactors.*

Cycle 4: Neuroinflammation and Central Sensitization. **Diagnostic criteria:** Evidence of neuroinflammation, microglial activation, or central pain/sensory amplification.

Tier 1 Testing (Clinical Assessment):

- **Quantitative sensory testing (QST):** Pressure pain thresholds, temporal summation
 - Algometer to measure pressure pain threshold (PPT) at standardized sites
 - PPT <4 kg/cm² = hyperalgesia, suggests central sensitization
 - Temporal summation testing: repeated stimuli produce increasing pain
 - Equipment cost \$200–500; can be done in any clinic
- **Cognitive testing:** Neuropsychological battery or screening tools
 - Processing speed, working memory, sustained attention deficits
 - NIH Toolbox Cognition Battery (free online)
 - Formal neuropsych testing: \$1,500–3,000
- **Symptom scales:** Central Sensitization Inventory (CSI), widespread pain index
 - CSI >40 suggests central sensitization
 - Free online questionnaire

Tier 2 Testing (Research/Specialized):

- **Brain PET imaging:** Neuroinflammation markers (TSPO PET)
 - Shows microglial activation in ME/CFS [73]
 - Research setting; cost \$3,000–5,000+
- **Cerebrospinal fluid analysis:** Cytokines, chemokines, lactate
 - Invasive; reserved for research or ruling out other diagnoses
 - Cost \$500–1,500

Clinical decision: *If QST shows hyperalgesia OR CSI >40 OR severe cognitive impairment → Neuroinflammatory cycle ACTIVE → Consider LDN, neuroinflammation-targeted supplements, avoid opioids (worsen central sensitization).*

Cycle 5: Endocrine Dysregulation. **Diagnostic criteria:** Evidence of HPA axis dysfunction, sex hormone abnormalities, or thyroid dysregulation beyond primary disease.

Tier 1 Testing (Widely Available):

- **Cortisol rhythm:** 4-point salivary cortisol (morning, noon, evening, bedtime)
 - Flattened diurnal rhythm = HPA axis dysregulation
 - Low morning cortisol <5–6 ng/mL may indicate adrenal insufficiency
 - Mail-order salivary testing; cost \$100–150
- **Thyroid comprehensive panel:** TSH, free T4, free T3, reverse T3, TPO antibodies
 - ME/CFS patients may have normal TSH but low T3 or high rT3
 - Widely available; cost \$150–300
- **Sex hormones:** Testosterone (men), estradiol and progesterone (women), DHEA-S (both)
 - Low testosterone in men common in ME/CFS
 - Estrogen dominance or low progesterone in women
 - Standard labs; cost \$100–200

Tier 2 Testing (Specialized):

- **ACTH stimulation test:** Measures adrenal reserve
 - Blunted response may indicate HPA axis dysfunction
 - Endocrinology clinic; cost \$300–500
- **24-hour urine free cortisol:** Integrates cortisol production over day
 - More comprehensive than single-point measurements
 - Cost \$100–150

Clinical decision: *If cortisol rhythm flattened OR low morning cortisol OR thyroid imbalance despite normal TSH OR sex hormone deficiencies → Endocrine cycle ACTIVE → Optimize thyroid (consider T3), address sex hormones if deficient, consider hydrocortisone (5–15 mg daily) if severe HPA dysfunction.*

Cycle Status Dashboard: Visual Treatment Prioritization

After completing the diagnostic battery, create a visual representation of which cycles are active. This guides treatment selection and monitoring.

Treatment Prioritization Algorithm

1. **Identify active cycles:** Red status (clear biomarker abnormalities) = active
2. **Prioritize by severity and treatability:**
 - *Highest priority:* Immune cycle with elevated GPCR autoantibodies (specific targetable pathology; immunoadsorption or daratumumab may be disease-modifying)

Table 18.5: Example: Cycle Status Dashboard for Individual Patient

Cycle	Status	Key Biomarker(s)	Treatment Priority
Mitochondrial	Active	2-day CPET: 18% VO ₂ decline	Priority 1
Immune	Active	GPCR Ab: β_2 -AR 95th %ile	Priority 1
Autonomic	Borderline	HR increase +28 bpm (lean test)	Monitor
Neuroinflammatory	Inactive	CSI: 32, QST: normal	Not targeted
Endocrine	Borderline	Flat cortisol rhythm	Priority 2

Interpretation: This patient has active mitochondrial and immune cycles (Priority 1 targets), borderline autonomic and endocrine cycles (monitor, intervene if worsens), and inactive neuroinflammatory cycle (no specific treatment needed). Recommended protocol: Mitochondrial support stack (CoQ10, NAD⁺ precursors) + immune intervention (consider immunoadsorption or daratumumab if accessible) + monitor autonomic symptoms.

- *High priority:* Mitochondrial cycle with 2-day CPET failure (foundational dysfunction; supports all other systems)
- *High priority:* Autonomic cycle with severe POTS (quality of life impact; relatively easy to treat)
- *Medium priority:* Endocrine dysregulation (supportive; may improve energy and cognition)
- *Lower priority:* Neuroinflammatory cycle (harder to target; overlaps with immune interventions)

3. Staged intervention:

- Start with 1–2 highest-priority cycles
- Assess response at 8–12 weeks
- Add interventions for additional cycles if first targets tolerated
- Re-assess cycle status at 6 months (some cycles may resolve when others are treated)

4. Monitor for new cycle entry:

- Repeat diagnostic battery at 6–12 month intervals
- Progressive disease may activate new cycles over time (sequential cycle entry model)
- Early detection allows intervention before entrenchment

Cost-Benefit Analysis: Which Tests Provide Most Information Per Dollar?

For patients with limited financial resources, prioritize high-yield, low-cost tests:

Recommended minimal battery (total cost \$200–300):

1. NASA lean test + HRV app (autonomic): \$0–50
2. Lactate post-exertion (mitochondrial): \$20–50
3. Salivary cortisol rhythm (endocrine): \$100–150
4. Basic immune panel: CBC with differential, NK cell count (\$50–100)

Table 18.6: Cost-Effectiveness Ranking of Cycle Diagnostic Tests

Test	Approximate Cost	Information Yield	Cost-Effectiveness
NASA lean test	\$0	Autonomic cycle: definitive	Excellent
HRV monitoring (app)	\$0–50	Autonomic ongoing tracking	Excellent
Lactate post-exertion	\$20–50	Mitochondrial: suggestive	Very good
Salivary cortisol 4-point	\$100–150	Endocrine: HPA axis	Very good
GPCR autoantibodies	\$300–500	Immune: predictive for immunotherapy	Good (if considering immunotherapy)
NK cell count/function	\$150–300	Immune: general dysfunction	Moderate
2-day CPET	\$1,500–3,000	Mitochondrial: gold standard	Good (if accessible)
Cytokine panel	\$200–400	Immune: high variability	Moderate (unreliable)
Brain PET	\$3,000–5,000+	Neuroinflammation: research	Poor (not actionable clinically)

This provides actionable information on 3–4 of the 5 cycles at minimal cost.

Expanded battery for aggressive intervention (total cost \$1,000–1,500):

- Add GPCR autoantibodies (\$300–500) if considering immunotherapy
- Add 2-day CPET (\$1,500–3,000) if accessible and critical for treatment decisions

Limitations and Caveats

- **Biomarker variability:** Many measures (cytokines, HRV, cortisol) show day-to-day variation; single measurements may not reflect true status
- **No validated cutoffs:** For most biomarkers in ME/CFS, we lack consensus diagnostic thresholds; interpretation requires clinical judgment
- **Cycle interactions:** Treating one cycle may improve biomarkers in another (e.g., immune intervention may improve mitochondrial function); serial testing required
- **Access barriers:** Advanced tests (2-day CPET, GPCR antibodies, PET imaging) are not widely available; many patients will rely on Tier 1 testing only
- **Insurance coverage:** Most specialized ME/CFS testing is not covered by insurance; out-of-pocket costs are significant

★ Key Point: Precision Medicine in Practice

Personalized cycle mapping represents the implementation of precision medicine for ME/CFS. Rather than treating all patients identically, this framework:

1. **Identifies active pathology** in each individual through biomarker assessment
2. **Prioritizes interventions** targeting documented abnormalities
3. **Avoids unnecessary treatments** for inactive cycles (reducing side effects and cost)
4. **Monitors response** through serial biomarker tracking
5. **Adjusts strategy** as cycle status changes over time

This approach is more complex than “try everything” empiricism, but it maximizes treatment efficacy while minimizing risk and cost. As ME/CFS research progresses and biomarkers become more standardized, cycle mapping will evolve from a conceptual framework to a validated clinical tool.

18.5.6 Early-Disease Anti-Cytokine Strategy

Speculation 60 (Immune Exhaustion Timeline: Early Intervention Preventive Window). **Certainty: 0.35.** Early aggressive anti-inflammatory intervention in the first 3 years of ME/CFS may prevent progression to severe disease and immune exhaustion. While anti-inflammatory approaches are established (omega-3, LDN, curcumin), stratifying intervention urgency by illness duration to define a preventive therapeutic window is a novel synthesis. This represents a paradigm shift from reactive symptom management to proactive cascade prevention. Certainty is low because no RCTs compare early versus late anti-inflammatory intervention in ME/CFS; the duration-dependent cytokine patterns (Achievement 7.3.1) are observational, and confounding by disease progression independent of intervention cannot be excluded.

Rationale If illness duration <3 years and cytokines elevated (particularly IL-6 >3–5 pg/mL), consider anti-inflammatory intervention to prevent progression to exhaustion phase. Achievement 7.3.1 documents duration-dependent cytokine patterns, and Section 7.9.1 presents the “Immune Exhaustion Timeline” hypothesis.

Conservative Approach (Before Biologics)

1. Aggressive anti-inflammatory supplementation:

- **Omega-3 fatty acids (EPA+DHA) 2–4 g daily**
 - **NOTE - EXCEEDS TYPICAL SUPPLEMENT DOSE:** Standard fish oil supplements provide 1000 mg (1 g) combined EPA+DHA daily. We recommend 2–4 g daily, which is 2–4× typical supplementation.
 - **Justification:** Omega-3 fatty acids (EPA/DHA) reduce pro-inflammatory cytokine production (IL-1, IL-6, TNF- α) via inhibition of arachidonic acid

metabolism and NF- κ B signaling. Therapeutic anti-inflammatory effects require EPA+DHA doses of 2–4 g/day based on cardiovascular and rheumatologic studies. Lower doses provide general health benefits but insufficient cytokine modulation.

- **Safety margin:** Doses up to 5 g/day are considered safe by FDA. Our recommendation of 2–4 g/day is well within this limit.
- **Side effects:** Fishy aftertaste (take with meals), mild GI upset, loose stools at higher doses. Mild blood-thinning effect.
- **Drug interactions:** May potentiate anticoagulants (warfarin). Monitor INR if on blood thinners.
- **Monitoring:** None required for most patients. If on warfarin, monitor INR.
- **Turmeric/curcumin 1000–2000 mg BID** (see Chapter 17 for complete dosing rationale - 2–4× typical supplement dose, well-tolerated, anti-inflammatory via NF- κ B inhibition)
- **Resveratrol 500 mg BID**
 - **NOTE - DRAMATICALLY EXCEEDS TYPICAL DOSE:** Typical resveratrol supplements provide 100–250 mg once daily. We recommend 500 mg twice daily (1000 mg/day total), which is 4–10× typical supplementation.
 - **Justification:** Resveratrol activates sirtuins (SIRT1) and inhibits NF- κ B, providing anti-inflammatory and potential mitochondrial benefits. Therapeutic doses for metabolic and inflammatory conditions in research studies use 500–1000 mg/day or higher. Lower doses may not achieve sufficient tissue concentrations for anti-inflammatory effects.
 - **Bioavailability note:** Resveratrol has poor bioavailability (<1%). This necessitates higher oral doses to achieve therapeutic levels. Micronized or liposomal formulations may improve absorption.
 - **Safety margin:** Clinical trials have used up to 2000–5000 mg/day without serious adverse effects. Our recommendation of 1000 mg/day is moderate.
 - **Side effects:** Generally well-tolerated. Occasional GI upset (nausea, diarrhea) at high doses. Take with food.
 - **Drug interactions:** May potentiate anticoagulants. Theoretical interaction with immunosuppressants.
 - **Monitoring:** None required.
- **Green tea extract (EGCG) 400 mg BID**
 - **NOTE - EXCEEDS TYPICAL SUPPLEMENT DOSE:** Typical green tea extract supplements provide 200–300 mg EGCG once daily. We recommend 400 mg twice daily (800 mg/day total), which is 2.5–4× typical supplementation.
 - **Justification:** Epigallocatechin gallate (EGCG) is the primary catechin in green tea with anti-inflammatory and antioxidant properties. Therapeutic doses for metabolic and inflammatory benefits in studies use 400–800 mg/day EGCG. Lower doses provide antioxidant effects but may be insufficient for immune modulation.
 - **Safety margin:** Doses up to 800–1200 mg/day have been studied. Our recom-

mendation of 800 mg/day is at the upper studied range.

- **CRITICAL WARNING - HEPATOTOXICITY RISK:** High-dose green tea extract (>800 mg EGCG/day) on empty stomach has been associated with rare cases of liver injury. ALWAYS take with food. If ALT/AST elevation occurs, discontinue immediately.
- **Side effects:** Nausea, GI upset (take with food), jitteriness (contains some caffeine unless decaffeinated).
- **Drug interactions:** May interact with beta-blockers, blood thinners. Contains caffeine (unless decaffeinated).
- **Monitoring:** Consider baseline and 3-month liver function tests (ALT/AST) if using high-dose chronically.

2. Low-dose naltrexone (LDN):

- 1.5–4.5 mg nightly
- Immune modulation (reduces pro-inflammatory cytokines)
- Safe, well-tolerated
- Takes 2–4 weeks for benefit

3. Dietary anti-inflammatory approach:

- Mediterranean diet (vegetables, fruits, olive oil, fish)
- Eliminate processed foods, refined sugars
- Consider anti-inflammatory elimination diet trial

Aggressive Approach (If Mild Conservative Fails)

- Discuss anti-cytokine biologics with rheumatologist (tocilizumab, etanercept)
- More justifiable in early disease (<3 years) with documented high cytokines
- May prevent progression to severe disease and immune exhaustion
- Requires close monitoring due to infection risk

18.5.7 Hormonal Optimization

For All Patients

- **Thyroid:** Optimize thyroid replacement if hypothyroid (many need T3 supplementation, not just T4)
- **Vitamin D:** Target 50–80 ng/mL (higher than standard; immune function benefit)
- **Iron:** Ferritin >50 ng/mL; some patients need higher for symptom improvement

Sex-Specific

- **Pre-menopausal women with cycle-linked crashes:**
 - Track symptoms across menstrual cycle

- If consistent luteal-phase worsening (days 14–28): Consider continuous oral contraceptives (eliminate hormone fluctuations)
- Or: Progesterone supplementation luteal phase
- **Post-menopausal women:**
 - Check estradiol
 - If low (<30 pg/mL) → trial HRT (Section 17.7.3)
 - Particularly if high IL-6 or prominent immune symptoms
- **Men with fatigue + cognitive dysfunction:**
 - Check testosterone (total and free)
 - If low → testosterone replacement (immune and energy benefits)

18.5.8 Microbiome Restoration

Gut-Immune Axis

~ Hypothesis 4: Dysbiotic Priming: Gut Dysbiosis Drives Immune Hyperactivation

Certainty: 0.35. Gut dysbiosis with fungal overgrowth may provide constant low-level antigenic exposure that primes immune cells to overreact, connecting Che et al.'s finding of exaggerated immune responses to *Candida* stimulation [189] with gut barrier dysfunction and documented microbiome alterations in ME/CFS (Section 11.1). This would explain both baseline immune activation and post-exertional malaise (exertion worsens gut barrier permeability). An estrogen-microbiome-immune connection may contribute to observed sex differences. No prior framework explicitly connects these findings into a unified therapeutic rationale; certainty is low because the dysbiotic priming mechanism is inferred rather than directly demonstrated in ME/CFS cohorts.

Section 7.9.2 presents the “Dysbiotic Priming” hypothesis: gut dysbiosis (Section 11.1) may maintain immune hyperactivation (Section 7.4.1). Addressing gut health may reduce systemic inflammation.

Stepwise Approach

1. **Assess GI involvement:**
 - Do you have GI symptoms (bloating, diarrhea, constipation, pain)?
 - Stool testing for dysbiosis (consider: GI-MAP, organic acids test, or similar)
2. **Dietary intervention:**
 - Eliminate processed foods, added sugars
 - Increase fiber (vegetables, fruits - unless FODMAP-sensitive)
 - Consider elimination diet if food sensitivities (low-FODMAP, AIP, etc.)
 - Probiotic-rich foods (if tolerated): yogurt, kefir, sauerkraut
3. **Targeted supplementation:**

- Probiotics: Multi-strain (Lactobacillus, Bifidobacterium), 25–50 billion CFU
- Saccharomyces boulardii 250 mg BID (anti-Candida, immune modulation)
- **Gut barrier support:**
 - **L-glutamine 5 g daily:** NOTE - Exceeds typical supplement dose (1–2 g). See Chapter 17 for complete dosing rationale. Therapeutic dose for gut barrier repair is 5–10 g/day (5–10× typical supplement dose). Extremely safe, well-tolerated.
 - **Zinc carnosine 75 mg BID** (150 mg/day total): NOTE - 2× typical supplement dose (75 mg once daily). See Chapter 17 for complete dosing rationale. Clinical mucosal healing studies use 75–150 mg BID. Provides 32 mg elemental zinc, below UL of 40 mg/day.
- Prebiotics: Inulin, partially hydrolyzed guar gum (feed beneficial bacteria)

4. Antifungal trial if indicated:

- If stool testing shows yeast overgrowth or strong clinical suspicion
- Fluconazole 100–200 mg daily for 4 weeks (prescription)
- Or: **Berberine 500 mg TID** (1500 mg/day total, natural antimicrobial) - NOTE: Exceeds typical supplement dose (500–1000 mg/day) by 1.5–3×. See Chapter 17 for complete dosing rationale. CRITICAL WARNING: May cause hypoglycemia if taking diabetes medications - physician supervision required.
- Concurrent probiotics and gut support

Sequential Gut-Vagal Repair Protocol

Speculation 61 (Sequential Gut-Vagal Repair Protocol). **Certainty: 0.25.** This protocol is derived from the mechanistic model of Wirth and Scheibenbogen [225] and the documented butyrate-producer deficiency in ME/CFS [349]. It addresses the dysbiosis → butyrate deficiency → impaired enterochromaffin serotonin → vagal dysfunction chain in logical sequence. The 5-phase structure is a clinical framework based on mechanistic plausibility, not a validated trial protocol. HRV (heart rate variability) serves as the primary objective monitoring endpoint throughout.

Phase 1: Eradication (Weeks 1–4)

- Test for SIBO (hydrogen/methane breath test) and treat if present: rifaximin, elemental diet, or herbal antimicrobials
- Test for fungal overgrowth and treat if documented (see antifungal approach above)
- Goal: Clear pathogenic overgrowth before introducing fermentable substrates
- Prerequisite: RS2 resistant starch and probiotics are contraindicated until this phase is completed

Phase 2: Motility Restoration (Weeks 4–8)

- Prokinetic therapy: low-dose erythromycin 50 mg at bedtime or prucalopride 1–2 mg daily (see Section 11.4.2 for mechanisms and prescribing details)

- Begin vagal activation exercises: 5 minutes of resonant-frequency breathing (typically 4.5–7 breaths/minute; individualise to the rate that maximally amplifies HRV [533]) twice daily; gargling 30 seconds after brushing teeth
- Goal: Restore normal gut motility to prevent SIBO recurrence; begin vagal tone training

Phase 3: Microbiome Recolonization (Weeks 8–16)

- Resistant starch RS2 (raw potato starch 1 tablespoon daily, increase slowly to 2 tablespoons)
- Multi-strain probiotic emphasizing *Faecalibacterium prausnitzii* (if available), *Lactobacillus rhamnosus*, *Bifidobacterium infantis*
- Dietary fiber diversity (multiple fiber types to support microbial diversity)
- Goal: Restore butyrate-producing bacterial community

Phase 4: Serotonergic Support (Weeks 12+)

- Sodium butyrate or tributyrin 1–2 g three times daily with meals (Chapter 22, Section 22.9.3)
- 5-HTP 50–100 mg two to three times daily (bypasses IDO2 diversion; see Chapter 22, Section 22.6.4)
- P5P (active B6) 25–50 mg daily (decarboxylase cofactor)
- Zinc glycinate or zinc carnosine 15–30 mg elemental zinc daily
- **Safety note:** 5-HTP is absolutely contraindicated with SSRIs, SNRIs, MAOIs, tramadol, or other serotonergic drugs—physician review required before starting
- Goal: Optimize enterochromaffin serotonin synthesis and vagal afferent input

Phase 5: Vagal Amplification (Ongoing)

- Continue butyrate supplementation
- Consider pyridostigmine 30–60 mg three times daily if orthostatic symptoms persist (amplifies vagal efferent output; standard POTS dosing)
- HRV biofeedback training 10–20 minutes twice daily (HeartMath or similar device)
- Goal: Maximize and sustain vagal tone and autonomic balance

Monitoring: Measure HRV (5-minute resting measurement) at each phase transition. Track gut symptom severity, fatigue scores, and orthostatic tolerance. HRV improvement is the primary indicator that the gut-vagal pathway is responding.

Who may benefit: ME/CFS patients with prominent GI symptoms (IBS, gastroparesis, SIBO), low HRV, and butyrate-producer deficiency on stool testing. Patients without GI involvement may have limited response.

18.6 Work and Study Accommodations

18.6.1 Critical Reality

Most mild-moderate patients attempt to maintain work/study. This often leads to progressive worsening because energy spent on work leaves none for social life, self-care, or recovery. **Accommodations are essential, not optional.**

18.6.2 Formal Accommodations

Request These Accommodations

- **Reduced hours:** 50–75% time if full-time unsustainable
- **Flexible schedule:** Work during peak energy times
- **Remote work:** Eliminate commute energy cost, enable rest breaks
- **Rest breaks:** Formal 15-minute horizontal rest every 2 hours
- **Quiet workspace:** Reduce sensory overload
- **Reduced meetings:** Cognitive load of meetings often underestimated
- **Deadline flexibility:** Accommodate fluctuating capacity
- **Parking accommodation:** Close parking to reduce walking

Legal Protections (Varies by Country)

- **US:** Americans with Disabilities Act (ADA) - ME/CFS qualifies; employer must provide reasonable accommodations
- **UK:** Equality Act - ME/CFS is protected disability
- **EU:** National disability discrimination laws vary by country
- **Documentation:** Physician letter documenting diagnosis and functional limitations

18.6.3 Self-Imposed Boundaries

- **Do not work through lunch:** Use for horizontal rest
- **Do not work evenings/weekends:** Reserve all non-work time for recovery
- **Say no to optional tasks:** Decline extra projects, social work events
- **Communicate limitations:** Better to set expectations than to fail to deliver

18.6.4 When to Stop Working

△ Warning 13: Work Cessation Criteria

If despite accommodations you are:

- Bedbound on weekends recovering from work week
- Progressively worsening (more frequent/severe PEM)
- Unable to maintain basic self-care (cooking, hygiene, errands)
- Developing new symptoms or severity increase

Then working is **causing progression** to severe disease. Apply for disability. Your health is more important than employment. Working yourself into severe ME/CFS leaves you unable to work *and* severely disabled.

18.7 Graded Exercise Therapy (GET): Why to Avoid

18.7.1 Critical Warning

Graded Exercise Therapy (GET) remains recommended in some countries despite evidence of harm. **GET is contraindicated in ME/CFS and can cause severe, lasting worsening.**

18.7.2 Why GET Fails

1. **Fundamental misunderstanding:** GET assumes deconditioning causes symptoms; increasing exercise reconditions. This is false. PEM is pathological response to exertion (Section 6.2.3), not deconditioning.
2. **Ignores PEM:** GET protocols ignore delayed symptom exacerbation, attributing it to “expected discomfort” rather than disease mechanism (Section 6.2.3).
3. **Biomarker evidence:** Chapters 6–7 document that exertion triggers immune activation (Section 7.4), oxidative stress (Section 6.4), and metabolic dysfunction (Section 6.2) - not adaptation.
4. **Patient harm surveys:**
 - 50–70% of patients report worsening from GET [3, 506]
 - Some become severe/bedbound after GET programs
 - UK NICE guidelines (2021) removed GET recommendation due to harm [127]

18.7.3 If Pressured by Physician

- Cite NICE 2021 guidelines (UK), recent reviews documenting harm
- Request pacing/energy envelope management instead
- Seek second opinion from ME/CFS-knowledgeable physician
- If insurance requires “exercise program,” document that standard GET worsens ME/CFS; request adaptive pacing therapy (APT) instead

18.7.4 Safe Activity Increase (If Appropriate)

Only if:

- Baseline symptom stability for 6+ months
- No PEM episodes for 3+ months
- Energy envelope well-established
- Under guidance of ME/CFS-knowledgeable professional

Principles:

- Increase activity 5–10% every 4–6 weeks (very gradual)
- If any PEM → immediately reduce to prior level
- Horizontal/recumbent exercise (recumbent bike, rowing)
- Never exceed anaerobic threshold
- Prioritize activities of daily living over formal exercise

18.8 Long-Term Strategy for Mild-Moderate Cases

18.8.1 Goals

1. **Primary:** Prevent progression to severe disease
2. **Secondary:** Improve function within energy envelope
3. **Tertiary:** Achieve remission or substantial recovery (ambitious but possible in some)

18.8.2 Timeline

- **Months 1–6:** Establish pacing, optimize symptom management, identify triggers
- **Months 6–12:** Implement disease-modifying strategies (immune modulation, hormones, microbiome)
- **Year 1–2:** Assess trajectory - stable? improving? worsening?
- **Year 2–5:** Continued optimization; some patients achieve significant recovery or remission

18.8.3 Realistic Expectations

- **Remission:** 5–10% of patients achieve sustained remission (symptom-free >1 year)
- **Substantial improvement:** 20–30% improve significantly (mild symptoms, near-normal function)
- **Stable mild-moderate:** 40–50% remain stable with good management
- **Progression:** 10–20% worsen despite intervention (often due to continued overexertion)

The goal is to maximize your chances of being in the improvement categories through aggressive early intervention and strict pacing.

18.9 Summary: Preventing the Descent

18.9.1 Key Principles

1. **Pacing is paramount:** More important than any medication or supplement
2. **Early intervention:** Treating mild disease aggressively may prevent severe disease
3. **Accommodations are essential:** Reduce work/study load to sustainable level
4. **Avoid GET:** Do not be pressured into graded exercise programs
5. **Target root causes:** Immune dysregulation, hormonal imbalance, microbiome - not just symptoms
6. **Hope with realism:** Some improve significantly; not all recover; pacing prevents worsening for most

18.9.2 Action Checklist

- ☐ Establish energy envelope (2-week activity tracking + HR monitoring)
- ☐ Implement 50% rule (do half of perceived capacity)
- ☐ Optimize sleep (hygiene + supplements or medication if needed)
- ☐ Address dominant symptoms (brain fog, pain, POTS, GI)
- ☐ Trial MCAS protocol if indicated (2-week cetirizine + famotidine + diet)
- ☐ Obtain basic labs (CBC, CMP, thyroid, iron, vitamin D, B12)
- ☐ Request work/study accommodations (reduced hours, flexible schedule, remote work)
- ☐ Avoid GET programs; seek pacing-based approach
- ☐ If early disease (<3 years), consider immune profiling and anti-inflammatory strategy
- ☐ If post-menopausal woman or low testosterone, check hormone levels
- ☐ Address microbiome if GI symptoms present
- ☐ Reassess every 3–6 months: Stable? improving? worsening? Adjust accordingly.

Mild-moderate ME/CFS is not mild suffering. It is life-altering, disabling, and deserves aggressive management. You are not being lazy. You are not deconditioned. You have a biological illness. Protect your energy envelope. Advocate for accommodations. Pursue treatments. Prevent progression.

Your future self will thank you for the boundaries you set today.

19 Pediatric ME/CFS: Severe and Housebound Cases

Children and adolescents with severe ME/CFS represent a particularly vulnerable population requiring specialized management approaches. While pediatric ME/CFS overall carries a substantially better prognosis than adult disease (54–94% improvement or recovery versus ≤22% in adults; see Chapter 5), severe cases present unique challenges that demand urgent, developmentally appropriate intervention. This chapter addresses the approximately 5–10% of pediatric ME/CFS patients who are housebound or bedbound, providing evidence-based guidance for home-based care, medical management with pediatric dosing, and strategies to preserve developmental progress during this critical period [393, 534].

Unlike adult severe ME/CFS (Chapter 17), pediatric severe disease retains meaningful potential for improvement or recovery, making appropriate early intervention particularly critical. The interventions described here are designed to reduce suffering, prevent complications of prolonged bedrest, maintain developmental trajectory, and preserve the window of opportunity for recovery.

★ Key Point: Cycle Dynamics Framework: Why Pediatric Prognosis Differs

The vicious cycle dynamics framework (Chapter 2, §2.1) provides a mechanistic explanation for the superior pediatric prognosis. ME/CFS involves multiple reinforcing pathophysiological loops (mitochondrial, immune, autonomic, neuroinflammatory, endocrine) that progressively recruit over time. Pediatric nervous system plasticity and earlier intervention timing mean: (1) fewer cycles have been recruited at diagnosis; (2) cycle “gain” (amplification strength) remains lower; and (3) reversibility windows remain open longer. The speculative time-dependent reversibility model ($R(t) = R_0 e^{-\lambda t}$; Section 24.11.4 in Chapter 24) *hypothesizes* that intervention efficacy decays exponentially with disease duration—which, if validated, would explain why early pediatric intervention preserves recovery potential that closes in chronic adult disease.

△ Warning: Physician Collaboration Required

Pediatric medication dosing, particularly for off-label uses, requires close collaboration with physicians experienced in pediatric medicine. The dosing recommendations in this chapter reflect published literature and clinical practice guidelines but must be individualized based on the child’s weight, age, comorbidities, and response. The Centers for Disease Control and Prevention explicitly recommends “extra caution when prescribing medicines for children with ME/CFS” and “starting medications at the smallest possible doses” [534]. All pharmacological interventions should be initiated and monitored by qualified healthcare providers.

⚠ Limitation 1: Pediatric Severe Protocols: Adult Extrapolation with Limited Paediatric Data

Treatment protocols in this chapter are largely adapted from adult ME/CFS management (Chapter 17) with paediatric dosing adjustments. Key epistemic boundaries:

- No randomised controlled trial has tested any ME/CFS-specific treatment in a paediatric severe population; the evidence base consists of case series, expert opinion, and extrapolation from adult studies that themselves lack placebo-controlled designs.
- Paediatric pharmacokinetics differ from adults in ways that affect drug metabolism, distribution, and response—simple weight-based dose adjustments may not capture these differences.
- The better paediatric prognosis (54–94% improvement) is drawn from heterogeneous studies spanning all severities; recovery rates specifically for severe paediatric ME/CFS are unknown and may be substantially lower.
- The “vicious cycle dynamics” and “reversibility window” models cited to justify early aggressive intervention are theoretical frameworks, not validated predictive tools.

19.1 Defining Severe Pediatric ME/CFS

Severe pediatric ME/CFS is defined by functional impairment that prevents normal school attendance and requires substantial caregiver support for daily activities. Unlike adult severity metrics that emphasize work capacity, pediatric severity must be assessed through the lens of age-appropriate functioning, including school participation, extracurricular activities, and peer socialization [393].

19.1.1 Functional Criteria

Severe pediatric ME/CFS is characterized by:

- **Housebound or bedbound status:** Unable to leave home for school or most activities; may be confined to bed for substantial portions of the day
- **School attendance impossibility:** Cannot attend school even with accommodations; requires homebound instruction or complete withdrawal
- **Dependence on caregivers:** Requires assistance with basic activities of daily living (bathing, dressing, meal preparation, mobility within home)
- **Severe post-exertional malaise:** Minimal activities (brief conversations, short walks within home) trigger prolonged symptom exacerbation
- **Multiple severe symptoms simultaneously:** Profound fatigue, cognitive dysfunction, orthostatic intolerance, pain, and sleep dysfunction occurring concurrently

Approximately 5–10% of pediatric ME/CFS cases fall into the severe category, though this may be underestimated because severely affected children are often unable to participate in medical visits or research studies [535].

19.1.2 Mast Cell Activation Syndrome (MCAS) Management in Severe Pediatric Cases

Mast cell activation affects 30–50% of ME/CFS patients and may be more prevalent in pediatric severe disease. MCAS contributes to brain fog, dysautonomia, GI dysfunction, and fatigue through excessive histamine and mediator release.

Pediatric MCAS Treatment

► Protocol 1: Pediatric Mast Cell Management

First-line: H1+H2 Antihistamine Combination

Age-appropriate dosing (consult pharmacist for precise pediatric calculations):

- **H1 antihistamine** (non-sedating preferred):
 - Cetirizine (Zyrtec): 0.25 mg/kg BID (typically 5 mg BID for younger children, 10 mg BID for adolescents)
 - Or: Loratadine (Claritin): 0.2 mg/kg daily (5 mg daily for younger, 10 mg for adolescents)
- **H2 antihistamine:**
 - Famotidine (Pepcid): 0.5–1 mg/kg BID (typically 10–20 mg BID for children, 20–40 mg BID for adolescents)
 - Provides dual histamine receptor blockade (H1+H2) for maximum effect
- **Low-histamine diet:**
 - Avoid: Aged/fermented foods, cured meats, aged cheese, alcohol (if adolescent), leftovers >24 hours
 - Encourage: Fresh foods prepared same-day, fresh fruits/vegetables, fresh protein

Optional Enhancement: Quercetin (Natural Mast Cell Stabilizer)

- **Dose:** 250–500 mg daily (younger children) to 500–1000 mg daily (adolescents)
- **Mechanism:** Natural flavonoid with mast cell stabilizing properties
- Can combine with H1+H2 antihistamines

MCAS Prophylactic Intensification for Pediatric Triggers

→ Recommendation 1: Pediatric MCAS Prophylaxis for High-Stress Activities

Mechanism: Mast cell activation episodes amplify cognitive and fatigue crashes through neuroinflammatory mediators (Chapter 14.24, lines 647–664). Proactive medication intensification before predictable triggers (medical visits, school events, family stress) can reduce crash severity.

Pediatric Protocol:

1. Identify your child's triggers:

- Emotional stress (medical appointments, family conflicts)
- Specific foods (aged cheese, cured meats, fermented foods)
- Environmental (heat, strong fragrances, weather changes)
- Immune challenges (infections, vaccinations)

2. Prophylactic medication timing (START 12–24 HOURS BEFORE known trigger):

- Increase H1 antihistamine to maximum tolerated dose
- Add quercetin 500–1000 mg daily (if not already taking)
- Strict low-histamine diet for 24 hours before and 24 hours after trigger

3. Activity preparation:

- Reduce non-essential activities day-of trigger
- Ensure adequate rest before and after
- Minimize additional cognitive or emotional demands

4. Parent tracking:

- WITHOUT prophylaxis: "Medical visit triggered crash 6/10, recovered in 2 days"
- WITH prophylaxis: "Medical visit with prophylaxis caused crash 3/10, recovered in 1 day"
- Adjust future trigger management based on efficacy

Evidence level: Moderate (MCAS prophylaxis established in pediatric allergology; ME/CFS crash-mitigation outcomes pending)

Expected outcomes: 25–50% reduction in trigger-related crash severity when MCAS component is significant.

19.1.3 Distinguishing from Adult Severity Metrics

While adult ME/CFS severity is often measured by work capacity (reduced hours, part-time work, inability to work), pediatric assessment requires different metrics:

- **School attendance percentage:** Days per week able to attend, hours per day tolerated
- **Academic performance trajectory:** Compared to pre-illness baseline, not peers

- **Extracurricular participation:** Sports, clubs, social activities—often the first casualties of ME/CFS
- **Self-care independence:** Age-appropriate comparison (a 15-year-old requiring help with bathing represents greater impairment than a 7-year-old needing the same assistance)
- **Peer interaction capacity:** Ability to maintain friendships through any medium (in-person, phone, online)

A child who cannot attend school at all, requires caregiver assistance for self-care, and has minimal capacity for peer interaction meets criteria for severe pediatric ME/CFS regardless of whether they can occasionally walk short distances within the home.

19.2 Home-Based Care Considerations

Severely affected pediatric patients cannot travel to clinics, making home-based and telehealth care essential. Families of housebound children face unique challenges that require proactive planning.

19.2.1 Telehealth Protocols

Telehealth should be the primary mode of medical follow-up for severe pediatric cases. Effective telehealth for this population requires:

- **Flexible scheduling:** Appointments during the child's best hours (which may vary unpredictably); willingness to reschedule if child is crashing
- **Caregiver as intermediary:** Parents may need to relay information if child cannot tolerate screen time or conversation
- **Brief, focused encounters:** 10–15 minute appointments with follow-up by message rather than extended video sessions
- **Written summaries:** Provide brief written summaries after appointments; cognitive dysfunction may prevent retention of verbal information
- **Symptom tracking between visits:** Use simple diaries or apps (completed by parent if needed) to capture patterns

19.2.2 Home Visit Considerations

When in-person assessment is essential (physical examination, procedures), home visits may be necessary. Considerations include:

- **Advance notice:** 24–48 hours minimum so family can prepare and child can rest beforehand
- **Minimal stimulation:** Visitors should speak quietly, minimize movements, avoid fragrances

- **Examination in place:** Examine child in bed if unable to sit; avoid requiring position changes that trigger orthostatic symptoms
- **Brief duration:** Complete essential examination in minimum time; defer non-urgent components
- **Recovery time:** Expect child to need 1–3 days recovery from home visit; schedule accordingly

19.2.3 Equipment Needs for Bedbound Children

Families may need guidance on equipment to improve care of bedbound children:

- **Hospital bed or adjustable bed:** Allows position changes without requiring child's effort; elevating head helps orthostatic symptoms
- **Bedside commode:** Reduces energy expenditure of bathroom trips
- **Shower chair or bath seat:** For children who can bathe but cannot stand
- **Wheelchair:** For medical appointments or rare outings; should be available even if rarely used
- **Communication aids:** Bell, monitor, or walkie-talkie to summon caregivers without shouting
- **Blue-light filtering:** Glasses or screen settings for any screen time; reduces sensory burden
- **Noise-canceling headphones:** For noise-sensitive children

19.2.4 Environmental Modifications

The home environment may require modifications to support a bedbound child:

- **Light control:** Blackout curtains or eye masks if light-sensitive
- **Noise control:** White noise machines, weatherstripping on doors, identifying and eliminating noise sources
- **Temperature regulation:** Space heater or fan near bed; layered blankets for temperature fluctuations
- **Air quality:** HEPA filter if chemical or odor sensitivities; fragrance-free household products
- **Accessibility:** Move child's bedroom to main floor if bathroom access is problematic on stairs

19.3 Subtype Assessment for Pediatric Severe Cases

As in adult severe ME/CFS, pediatric severe cases often involve multiple failing systems. Brief subtype assessment helps prioritize medical interventions within the constraints of pediatric dosing and developmental considerations.

→ **Recommendation 2: Pediatric Subtype Classification and Prioritization**

Quick assessment for parents/caregivers:

1. What is limiting your child MOST?

- Difficulty processing information, forgetfulness, confusion → **CNS-Primary**
- Cannot stand without dizziness, fainting episodes, extreme tachycardia → **Autonomic-Primary**
- Severe muscle weakness, pain, low energy even at rest → **Peripheral-Primary**
- Multiple severe symptoms at same time → **Global**

2. Symptom pattern:

- Child can move/play but seems confused or forgetful → **Suggests CNS-Primary**
- Child wants to do things but dizziness/weakness prevents standing → **Suggests Autonomic-Primary**
- Child's body seems "tired" all the time, pain when moving → **Suggests Peripheral-Primary**

Pediatric-adapted treatment prioritization:

Subtype A (CNS-Primary): Cognitive dysfunction dominates

- **Priority 1:** Optimize sleep (critical for pediatric CNS recovery)
- **Priority 2:** Simple cognitive support (very gentle; avoid stimulants in children)
- **Priority 3:** Dietary support (adequate nutrition for brain development)

Subtype B (Autonomic-Primary): Orthostatic intolerance dominates

- **Priority 1:** Aggressive hydration and salt loading (first-line pediatric POTS treatment)
- **Priority 2:** Compression garments (age-appropriate sizing)
- **Priority 3:** Medications if hydration/compression insufficient (midodrine is pediatric-approved)

Subtype C (Peripheral-Primary): Muscle weakness/fatigue dominates

- **Priority 1:** Mitochondrial support (gentle: CoQ10, carnitine—pediatric doses)
- **Priority 2:** Nutritional adequacy (protein, calories critical for growth)
- **Priority 3:** Pain management (non-pharmacological first)

Subtype D (Global): Multi-system involvement

- **Approach:** Multi-domain protocol, but scale everything to child's developmental stage
- **Caution:** Children tolerate poly-pharmacy poorly; use combination approaches (e.g., sleep protocol addresses both sleep + autonomic dysfunction) when possible
- **Monitoring:** Track side effects carefully; pediatric physiology differs from adults

Evidence level: Plausible (subtype framework from Chapter 14.24, adapted for pediatrics)

Critical reminder: Subtype assessment guides prioritization but does NOT delay foundational interventions (pacing, sleep, hydration) while waiting for optimization.

19.4 Medical Management

Medical management of severe pediatric ME/CFS follows similar principles to adult management (Chapter 17) but requires pediatric-specific dosing and heightened attention to developmental effects. The CDC recommends extra caution with pediatric ME/CFS medications, starting at lowest possible doses [534].

19.4.1 Orthostatic Intolerance (Priority)

Orthostatic intolerance (OI) affects 70–90% of pediatric ME/CFS patients, compared to approximately 40–70% of adults [393]. This higher prevalence makes OI management the single most important intervention in pediatric ME/CFS. Treatment of orthostatic intolerance improves not only cardiovascular symptoms but also fatigue, cognitive function, and overall wellbeing [338].

Non-Pharmacological Interventions

Non-pharmacological measures should be implemented first and continued even if medications are added:

► Protocol 2: Pediatric OI Non-Pharmacological Protocol

Hydration Targets (age-adjusted):

- Ages 4–8 years: 1.5–2 liters/day
- Ages 9–13 years: 2–2.5 liters/day
- Ages 14–18 years: 2.5–3 liters/day
- **Note:** These targets exceed standard pediatric recommendations because ME/CFS patients have reduced plasma volume; adequate hydration is therapeutic, not merely maintenance
- **Timing:** Spread throughout day; concentrated fluid intake before upright activities
- **Type:** Water, oral rehydration solutions, dilute juice; avoid excessive caffeine

Salt Supplementation (pediatric dosing):

- Children <30 kg: 2–3 grams sodium/day (from dietary sources plus salt tablets if needed)

- Children 30–50 kg: 3–5 grams sodium/day
- Adolescents >50 kg: 5–8 grams sodium/day
- **Implementation:** Salt tablets (0.5–1 g each) with meals and snacks; salted foods; oral rehydration solutions
- **Monitoring:** Blood pressure should be monitored; reduce salt if sustained hypertension develops
- **Contraindications:** Reduce dose in renal or cardiac disease

Compression Garments (sizing for children):

- **Waist-high compression:** Most effective; 20–30 mmHg compression
- **Pediatric sizing:** Measure carefully; adult sizes often too large
- **Alternative:** Abdominal binder if full-leg compression not tolerated
- **Tolerance:** Start with shorter wear times (1–2 hours); increase gradually
- **Application:** Put on while lying down, before rising

Positioning Strategies:

- **Elevate head of bed:** 10–15 degrees (blocks under head of bed, not pillows) to maintain blood volume training
- **Avoid prolonged standing:** Sit or lie whenever possible
- **Lower extremity movement:** Fidgeting, ankle pumping, crossing legs when must stand
- **Squat when symptomatic:** If feeling faint, squat immediately (raises blood pressure)
- **Avoid warm environments:** Heat worsens vasodilation and OI symptoms

Pharmacological Interventions

When non-pharmacological measures are insufficient, medications may be needed. Pediatric POTS studies demonstrate high response rates to standard OI medications [338].

► Protocol 3: Pediatric OI Pharmacological Protocol

First-Line: Fludrocortisone (Florinef)

Mineralocorticoid that expands plasma volume by promoting sodium and water retention.

Pediatric Dosing:

- **Starting dose:** 0.05 mg (50 mcg) once daily (half of the standard 0.1 mg tablet)
- **Titration:** Increase by 0.05 mg weekly if tolerated and symptoms persist
- **Target dose:** 0.1–0.2 mg daily (rarely exceeds 0.2 mg in pediatrics)
- **Timing:** Morning, with breakfast
- **Response timeline:** 1–2 weeks for initial effect; full effect may take 4–6 weeks

Monitoring Requirements:

- **Blood pressure:** Weekly initially; must catch hypertension early
- **Potassium:** Baseline and at 2–4 weeks; fludrocortisone can cause hypokalemia
- **Weight:** Weekly; excessive weight gain suggests fluid retention beyond therapeutic
- **Edema:** Ankle swelling indicates dose may be too high
- **Growth:** Long-term corticosteroid effects on growth are theoretical but monitor height velocity

Side Effects and Management:

- **Hypokalemia:** Supplement with potassium-rich foods or potassium chloride if levels fall
- **Hypertension:** Reduce dose or discontinue
- **Headache:** Often transient; reduce dose if persistent
- **Edema:** Mild ankle edema may be acceptable; reduce dose if significant

Second-Line: Midodrine (ProAmatine)

Alpha-1 agonist that causes vasoconstriction, increasing blood pressure and reducing venous pooling. Pediatric POTS studies report 78% response rate [338].

Pediatric Dosing:

- **Starting dose:** 2.5 mg (half tablet) once or twice daily
- **Titration:** Increase by 2.5 mg every 3–7 days as tolerated
- **Target dose:** 5–10 mg three times daily (maximum 40 mg/day in adolescents)
- **Timing:** Upon waking, midday, mid-afternoon; **do not give within 4 hours of bedtime** (supine hypertension risk)
- **Response timeline:** Hours to days; relatively rapid onset

Critical Warnings:

- **Supine hypertension:** Most important side effect; patient must be upright when taking medication and should not lie flat for 4 hours after dose
- **Last dose timing:** No later than 4–6 PM to avoid nocturnal hypertension
- **Scalp tingling:** Common (piloerection); not dangerous but can be bothersome
- **Urinary retention:** Rare in pediatrics; more common in older patients

Monitoring Requirements:

- **Supine blood pressure:** Check BP lying down periodically; if systolic >150 while lying, reduce dose
- **Standing blood pressure:** Should improve with treatment; document response

Third-Line: Pyridostigmine (Mestinon)

Acetylcholinesterase inhibitor that enhances autonomic function and may improve orthostatic tolerance.

Pediatric Dosing:

- **Starting dose:** 15–30 mg twice to three times daily

- **Titration:** Increase gradually based on response
- **Target dose:** 30–60 mg three times daily
- **Timing:** With meals to reduce GI side effects

Side Effects:

- **GI symptoms:** Nausea, diarrhea, abdominal cramping (most common; often dose-limiting)
- **Increased salivation/sweating:** Cholinergic effects
- **Bradycardia:** Monitor heart rate, especially in combination with beta-blockers

Fourth-Line: IV Fluids

For severe OI unresponsive to oral measures and medications.

Indications:

- Syncope or near-syncope despite adequate oral hydration, salt, and medications
- Unable to maintain adequate oral intake due to nausea or gastroparesis
- Severe dehydration from illness exacerbation
- Acute severe crashes requiring rapid support

Implementation:

- **Typical regimen:** Normal saline 1–2 liters, 1–3 times weekly
- **Access:** Peripheral IV for intermittent use; PICC line or port for frequent/regular infusions
- **Setting:** May be given at infusion center, at home by visiting nurse, or by trained parent
- **Risks:** Line infection (PICC/port), volume overload, electrolyte disturbance

Note on IV access: Decisions about central access (PICC, port) in pediatric ME/CFS require careful risk-benefit analysis. Line infections are serious complications. IV fluids should be reserved for patients with clearly documented benefit from fluid loading who cannot achieve equivalent benefit orally.

19.4.2 Sensitization Prevention Protocol

Severe pediatric ME/CFS patients are vulnerable to progressive sensitization—the phenomenon where increasing numbers of triggers (infections, environmental exposures, activities) progressively reduce symptom tolerance. Early intervention to prevent sensitization can preserve quality of life and potentially prevent further deterioration.

► Protocol 4: Pediatric Sensitization-Prevention Protocol

Anti-Neuroinflammatory Foundation

A multi-component approach targets neuroinflammatory activation, which underlies both PEM and progressive sensitization. The goal is to interrupt the cycle where each crash and each infection creates lasting sensitization (see Chapter 8 for discussion

of PEM kindling hypothesis and Chapter 7 for discussion of infection damage ratchet mechanisms).

Low-Dose Naltrexone (LDN):

- **Dosing:** Weight-adjusted: 0.1 mg/kg daily, maximum 4.5 mg
- **Age-adjusted examples:**
 - Child 20 kg: 2 mg daily
 - Child 40 kg: 4 mg daily
 - Adolescent 45+ kg: 4.5 mg daily
- **Timing:** Evening (bedtime) for better efficacy
- **Onset:** 2–8 weeks for measurable effect
- **Mechanism:** Low doses activate microglia inhibition, reducing neuroinflammation that perpetuates PEM cycles
- **Safety:** Generally well-tolerated in pediatrics; reversible if discontinued

Palmitoylethanolamide (PEA):

- **Dosing:** Age-adjusted based on published pediatric data
 - Younger children (6–10 years): 600 mg daily
 - Older children (11–14 years): 900 mg daily
 - Adolescents (15–18 years): 1200 mg daily
- **Type:** Enteric-coated formulations improve absorption
- **Mechanism:** Endocannabinoid system modulation; reduces neuroinflammation and neuropathic sensitization
- **Timing:** Can be taken with meals

Omega-3 Fatty Acids (EPA/DHA):

- **Dosing:** Age-appropriate targets for anti-inflammatory benefit
 - Children <30 kg: 500–1000 mg EPA/DHA daily
 - Children 30–50 kg: 1000–1500 mg EPA/DHA daily
 - Adolescents >50 kg: 1500–2000 mg EPA/DHA daily
- **Form:** Liquid or chewable formulations often better tolerated than capsules
- **Benefits:** Anti-inflammatory, supports cardiovascular health, may improve mood

Comprehensive Sleep Assessment and Treatment

Sleep deprivation perpetuates neuroinflammation. Severe cases warrant sleep medicine evaluation:

- **Sleep study:** Polysomnography to rule out sleep apnea (higher prevalence in ME/CFS)
- **Aggressive apnea treatment:** If present, treating sleep apnea often improves overall ME/CFS symptoms
- **Optimization:** Melatonin, trazodone, or low-dose amitriptyline (see Protocol 19.4.4) as needed to ensure adequate sleep quality

Activity Monitoring with Wearable Alerts

Preventing overexertion is critical to preventing sensitization cascade:

- **Heart rate monitoring:** Continuous or frequent monitoring using wearables appropriate for the child's age
- **Threshold definition:** Establish safe heart rate ceiling based on aerobic threshold calculation: $(220 - \text{age}) \times 0.55$ to 0.60
- **Parent/caregiver alerts:** Configure wearable to alert parents when child approaches or exceeds threshold
- **Purpose:** Prevent parent-blind overexertion (activity the child's caregiver cannot directly observe, or where child underestimates intensity)
- **Example:** Bedbound child who engages in sustained conversation or cognitive activity without realizing intensity until crash occurs next day

Infection Prevention as Sensitization Prevention

Each infection causes both acute PEM and lasting sensitization increases. Aggressive prevention is sensitization prevention:

- **Masking protocol:** FFP2 or N95 masks during any household illness; consider seasonal masking during high-transmission periods
- **Isolation:** If possible, isolate severely ill child from sick household members (separate room, bathroom if feasible)
- **Hand hygiene:** Frequent handwashing after any contact with potentially infected persons
- **Prompt treatment:** Antivirals if viral infection suspected; antibiotics for bacterial infections
- **Prophylaxis consideration:** During high-risk periods (siblings in school with active outbreaks), discuss prophylactic antivirals with physician

Educational Continuity as Health Protection

While not a direct medical intervention, preventing crash-induced educational disruption protects developmental trajectory and reduces secondary psychological harm that perpetuates neuroinflammation:

- **Homebound instruction:** Maintain some educational engagement without requiring school attendance that triggers crashes
- **Avoid education-forced crashes:** Never pressure school attendance for academic reasons when doing so will trigger PEM
- **Each prevented crash:** Each crash not occurring is a sensitization cascade prevented

△ Warning: Developmental Trajectory Preservation is Paramount

Severe pediatric ME/CFS differs from adult disease in one critical dimension: the developing nervous system and the developing life trajectory. Each crash during childhood does not merely reduce current function—it may have lasting effects on neurological development, educational progress, and identity formation. Sensitization prevention is not

mere symptom management; it is developmental protection. Prevention of a single crash during a critical developmental window may have lifelong implications for the child's ultimate prognosis and capacity.

19.4.3 Pain Management

Pain is common in pediatric ME/CFS, including widespread musculoskeletal pain, headaches, and abdominal pain. Pain management in developing nervous systems requires particular caution.

Non-Pharmacological Approaches

Non-pharmacological strategies should be first-line:

- **Positioning:** Supportive pillows, position changes to relieve pressure
- **Heat/cold:** Heating pads for muscle pain; cold packs for acute inflammation or headaches
- **Gentle massage:** Light massage by caregiver (not deep tissue; avoid triggering PEM)
- **TENS units:** Transcutaneous electrical nerve stimulation for localized pain
- **Distraction:** Audio content (audiobooks, music, podcasts) at tolerable volumes

Analgesic Medications

► Protocol 5: Pediatric Pain Management Protocol

Tier 1: Acetaminophen and NSAIDs

Acetaminophen (Tylenol):

- **Dose:** 10–15 mg/kg every 4–6 hours as needed
- **Maximum:** 75 mg/kg/day (up to 4 grams/day in adolescents)
- **Advantages:** No anti-inflammatory effects that might mask fever; low GI risk
- **Caution:** Hepatotoxicity at high doses; avoid in liver disease

Ibuprofen (Advil, Motrin):

- **Dose:** 5–10 mg/kg every 6–8 hours as needed
- **Maximum:** 40 mg/kg/day (up to 2.4 grams/day in adolescents)
- **Advantages:** Anti-inflammatory effects helpful for musculoskeletal pain
- **Caution:** GI irritation, renal effects with chronic use; take with food

Naproxen (Aleve):

- **Dose:** 5–7 mg/kg every 12 hours
- **Maximum:** 1 gram/day in adolescents

- **Advantages:** Longer duration; twice-daily dosing
- **Caution:** Same NSAID precautions as ibuprofen

Tier 2: Neuropathic Pain Agents

For neuropathic pain characteristics (burning, tingling, shooting, allodynia), consider agents targeting neuropathic pathways. These require specialist consultation in pediatrics.

Gabapentin:

- **Starting dose:** 5 mg/kg/day divided into 3 doses (e.g., 100 mg TID for 60 kg adolescent)
- **Titration:** Increase by 5 mg/kg/day every 5–7 days
- **Target:** 15–35 mg/kg/day divided TID (typical adolescent dose 900–1800 mg/day)
- **Side effects:** Sedation, dizziness, peripheral edema
- **Advantages:** Generally well-tolerated; no significant drug interactions

Amitriptyline (low-dose):

- **Starting dose:** 0.1 mg/kg at bedtime (typically 5–10 mg for child, 10–25 mg for adolescent)
- **Titration:** Increase by 0.1 mg/kg weekly if tolerated
- **Target:** 0.5–1 mg/kg at bedtime (rarely exceeds 50 mg in pediatrics)
- **Side effects:** Anticholinergic effects (dry mouth, constipation, urinary retention), sedation, QTc prolongation
- **Dual benefit:** May help pain AND sleep

△ **Warning: Black Box Warning: Pediatric Antidepressants**

Tricyclic antidepressants (amitriptyline) and SSRIs/SNRIs carry FDA black box warnings for increased suicidal thinking and behavior in children and adolescents. Low-dose amitriptyline for pain is typically below antidepressant doses, but families should be counseled about warning signs and patients should be monitored for mood changes, particularly during dose initiation and adjustments.

Tier 3: Specialist Pain Management

Referral to pediatric pain specialist is indicated for:

- Pain refractory to Tier 1–2 interventions
- Pain significantly limiting function beyond ME/CFS baseline
- Complex regional pain syndrome (CRPS) features
- Need for opioid consideration (rarely appropriate in pediatric ME/CFS)

19.4.4 Sleep Optimization

Sleep dysfunction is nearly universal in ME/CFS. Children with ME/CFS experience unrefreshing sleep regardless of duration, along with difficulty falling asleep, maintaining sleep, or both. The severely ill child may sleep 12–16 hours yet feel exhausted.

Sleep Hygiene for Bedbound Children

Standard sleep hygiene recommendations require modification for bedbound patients who cannot leave their beds:

- **Bed association:** When child is in bed 20+ hours/day, the bed becomes associated with wakefulness. Mitigation: different positions, blankets, or pillow arrangements for “sleep time” versus “awake time”; different lighting (lamp on for awake, off for sleep)
- **Light exposure:** Attempt some natural light during “daytime” hours, even if from window while lying down; complete darkness for sleep periods
- **Screen timing:** Reduce screens 1–2 hours before intended sleep; blue-light filtering if screens are used
- **Consistent schedule:** Maintain regular sleep and wake times even when homebound; circadian rhythm disorders are common in ME/CFS
- **Temperature:** Cooler room temperatures (65–68°F) facilitate sleep

Sleep Medications

► Protocol 6: Pediatric Sleep Protocol

First-Line: Melatonin

Melatonin is the preferred first-line sleep aid for pediatric ME/CFS due to safety profile and documented efficacy [323].

Dosing:

- **Starting dose:** 0.5–1 mg, 30–60 minutes before desired sleep time
- **Titration:** Increase by 0.5–1 mg every 3–7 days if ineffective
- **Target dose:** 1–5 mg for most children; some adolescents may need up to 10 mg
- **Timing is critical:** Must be taken at consistent time; efficacy depends on circadian timing, not just sedation
- **Extended-release formulations:** May help with sleep maintenance (waking in night)

Important Considerations:

- **Quality matters:** Pharmaceutical-grade melatonin is preferable; supplement quality varies widely
- **Not purely sedating:** Melatonin works by signaling circadian timing; some children feel more alert initially before sleep onset
- **Long-term safety:** Generally regarded as safe for extended use in pediatrics, though long-term studies are limited

Second-Line: Low-Dose Trazodone

Serotonin antagonist and reuptake inhibitor with sedating properties; commonly used off-label for pediatric insomnia.

Dosing:

- **Starting dose:** 12.5–25 mg at bedtime
- **Titration:** Increase by 12.5–25 mg weekly if needed
- **Target dose:** 25–100 mg at bedtime (rarely exceeds 100 mg for sleep in pediatrics)

Side Effects:

- **Morning sedation:** Most common; may need dose reduction
- **Orthostatic hypotension:** Caution in patients with OI (may worsen or improve—varies by patient)
- **Priapism:** Rare but serious; educate male adolescents to seek immediate care

Third-Line: Low-Dose Amitriptyline

As above for pain, low-dose amitriptyline at bedtime can improve sleep quality. Dosing: 5–25 mg at bedtime. Dual benefit for patients with pain plus sleep dysfunction.

Other Options (Specialist Consultation Recommended):

- **Clonidine:** 0.05–0.1 mg at bedtime; alpha-2 agonist with sedating effects
- **Hydroxyzine:** 12.5–50 mg at bedtime; antihistamine with sedating effects
- **Mirtazapine:** 7.5–15 mg at bedtime; antidepressant with sedating effects at low doses

△ Warning: Avoid in Pediatrics

- **Benzodiazepines:** Risk of dependence, cognitive effects, respiratory depression; avoid except for acute crisis management under specialist supervision
- **Z-drugs (zolpidem, eszopiclone):** Not FDA-approved for pediatrics; limited safety data; complex sleep behaviors reported

Circadian Rhythm Disorders

Delayed sleep phase disorder (DSPS) is common in adolescents with ME/CFS—the natural sleep-wake cycle shifts later, making it difficult to fall asleep before 2–4 AM and wake before noon. This overlaps with but is distinct from ME/CFS sleep dysfunction.

Management of DSPS:

- **Chronotherapy:** Gradual advance of sleep time (15–30 minutes earlier every few days)
- **Light therapy:** Bright light exposure (10,000 lux light box) immediately upon waking for 20–30 minutes; helps advance circadian phase
- **Melatonin timing:** Low-dose melatonin (0.5–1 mg) 4–6 hours before desired sleep time (not at bedtime) can help advance circadian phase
- **Evening light restriction:** Minimize bright light exposure, especially blue light, in evening hours

Circadian Light Therapy for Energy and Sleep Alignment

~ Hypothesis 1: Circadian Energy Misallocation in Pediatric ME/CFS

The selective energy dysfunction hypothesis (Chapter 14.24, lines 628–645) proposes that circadian dysregulation impairs energy budget allocation, explaining why pediatric patients often have evening energy spikes despite daytime exhaustion.

→ Recommendation 3: Bright Light Therapy for Circadian Re-entrainment in Pediatric Severe Cases

Mechanism: Morning bright light exposure (10,000 lux) resets the suprachiasmatic nucleus (SCN), improving alignment of circadian energy distribution with sleep-wake cycle. Particularly effective for children with delayed sleep phase or energy crashes mid-afternoon followed by evening “second wind.”

Pediatric Protocol:

1. **Equipment:** 10,000 lux light therapy box (\$25–100). Child-friendly options: Some have adjustable arms or can be positioned at varied angles for bedbound use.
2. **Timing:** 20–30 minutes immediately upon waking (within 30 minutes of desired wake time)
 - **Consistency crucial:** Same wake time every day (even weekends) for circadian effectiveness
 - **NEVER use after 3pm** (will delay sleep further and worsen late-evening sleep phase issues)
3. **Position:** Light box 16–24 inches from child’s face. For bedbound children, can be positioned on bedside table or stand at angle allowing light exposure while lying down.
4. **Safety:**
 - Safe for all ages
 - Very low risk of adverse effects (occasional mild eye strain if too close)
 - DO NOT stare directly at bulb
 - Discontinue if triggers mood elevation or anxiety (rare)
5. **Expectation setting:** Benefits emerge over 2–4 weeks; earlier bedtime and more consistent daytime energy are typical outcomes.

Evidence level: Moderate (circadian disruption documented in pediatric ME/CFS; light therapy established for circadian disorders; pediatric ME/CFS-circadian-energy specific RCTs pending)

Expected outcomes:

- More predictable sleep onset at night
- Reduced late-evening “second wind” phenomenon
- Better energy consistency throughout day
- Timeline: 2–4 weeks

Sleep Spindle Enhancement via Acoustic Stimulation (Low Priority, Experimental)**→ Recommendation 4: Pink/White Noise for Pediatric Sleep Architecture**

Mechanism: Sleep spindles (brief high-frequency brain activity during sleep) are reduced in ME/CFS. Acoustic stimulation during sleep may enhance spindle production, improving sleep architecture coordination (Chapter 14.24, lines 552–569).

Pediatric Protocol (Simple, Non-Pharmacological):

- **Equipment:** White or pink noise machine (\$15–50) or free app
 - White noise: Constant frequency; easier to find
 - Pink noise: Lower frequencies enhanced; some literature suggests better sleep effects
 - Apps: myNoise.net (free, customizable), YouTube brown noise videos
- **How to use:**
 - Play throughout entire sleep period (all night)
 - Volume: Low (30–50 dB, quiet background level); NOT disruptive
 - Placement: Bedside speaker positioned where child can hear but not blasted
 - Start time: Can use during naps and nighttime sleep
- **Parent tracking:**
 - Sleep quality rating (child report: more rested vs. unchanged vs. worse)
 - Duration of trial: Minimum 2–4 weeks to assess benefit
 - Expected benefit timeline: 3–4 weeks if spindles improve
- **Integration:**
 - Adjunctive to sleep medications (melatonin, trazodone)
 - Can be combined with other sleep strategies
 - Non-pharmacological option for families preferring fewer medications

Evidence level: Speculative (spindle deficits documented in ME/CFS; acoustic enhancement effect unproven in pediatric ME/CFS)

Expected outcomes: Modest improvement in subjective sleep quality if spindles enhance. Not expected to directly improve daytime fatigue or activity tolerance.

Positioning: Very low priority. Sleep medications have stronger evidence. Consider only if child has difficulty with medications or if medications provide insufficient benefit.

19.4.5 Cognitive Support

Cognitive dysfunction (“brain fog”) significantly impacts educational progress. In severe cases, children may be unable to read, follow conversations, or retain information.

Cognitive Function Assessment

Formal neuropsychological testing is often not feasible for severely ill children (the testing itself may trigger PEM). Instead, functional assessment through observation:

- **Reading tolerance:** How long can child read or be read to before cognitive fatigue?
- **Conversation tolerance:** How long can child participate in conversation before losing comprehension?
- **Information retention:** Can child recall content from previous day? Previous hour?
- **Word-finding:** Does child frequently lose words or use wrong words?
- **Processing speed:** Is there noticeable delay between question and response?

Maintaining Developmental Progress

Even during severe illness, maintaining some cognitive engagement is important for developmental trajectory:

- **Match modality to capacity:** Audiobooks if reading is impossible; dictated responses if writing is impossible
- **Micro-learning:** 5–10 minute educational activities with breaks, rather than extended lessons
- **Interest-driven:** Children may tolerate more cognitive effort for topics of personal interest
- **No pressure:** Removing academic pressure paradoxically often improves cognitive function
- **Social connection:** Even brief social interaction (texting friends, short video calls) maintains developmental skills

When Cognitive Symptoms May Improve

Cognitive function often improves with OI treatment. If orthostatic intolerance is inadequately treated, cerebral hypoperfusion (reduced blood flow to brain) contributes significantly to brain fog (see Chapter 10, Section 10.3.1). Many children experience noticeable cognitive improvement within 1–2 weeks of effective OI management. Always optimize OI treatment before assuming cognitive dysfunction is irreversible.

Intranasal Delivery for CNS-Targeted Cognitive Support

★ Key Point: BBB Vulnerability in Pediatric ME/CFS

The selective energy dysfunction hypothesis (Chapter 14.24, lines 238–257) proposes that the blood-brain barrier may be compromised in ME/CFS, limiting delivery of cognitive support compounds. Intranasal delivery bypasses the BBB via olfactory nerve pathways.

→ Recommendation 5: Intranasal Options for Severe Pediatric Cognitive Dysfunction

For severe pediatric cases with refractory brain fog:

- **Intranasal formulations:** If oral cognitive support proves insufficient, ask your child's neurologist or ME/CFS specialist about intranasal options. Examples include intranasal insulin (being studied for cognitive support in neurodegenerative disease) or dopamine precursors.
- **Caution with children:** Intranasal medications require cooperation and proper technique. May be difficult in very young or severely ill children.
- **Current status:** NOT standard pediatric ME/CFS care. EXPERIMENTAL. Only pursue with specialist guidance.

Evidence level: Speculative (established for other neurological conditions; no pediatric ME/CFS trials)

19.5 Preventing Complications of Prolonged Bedrest

Prolonged bedrest, while necessary for severe ME/CFS, carries its own complications. These must be proactively prevented without triggering PEM through excessive activity.

△ Warning: This Is NOT Graded Exercise Therapy

The interventions in this section are passive range of motion, positioning, and deconditioning prevention—NOT graded exercise therapy (GET). GET involves progressive increases in exercise with the goal of reconditioning. The interventions here maintain baseline physical function and prevent complications while respecting the energy envelope. They should never cause PEM. If any intervention triggers symptoms, it should be reduced or discontinued.

19.5.1 Contracture Prevention

Prolonged immobility can lead to joint contractures (permanent shortening of muscles and tendons). Prevention:

- **Passive range of motion:** Caregiver gently moves each major joint through full range of motion daily
- **Duration:** 5–10 minutes total; each joint moved 5–10 times
- **Key joints:** Ankles (prevent foot drop), knees, hips, shoulders, elbows, wrists, fingers
- **Technique:** Slow, gentle movements; never force past comfortable range; stop if painful
- **Timing:** During lower-symptom periods; not during acute crashes

19.5.2 Osteoporosis Risk

Bedbound children and adolescents are at risk for bone loss, which is particularly concerning during growth periods.

Prevention:

- **Calcium:** Ensure adequate dietary calcium or supplement (1000–1300 mg/day depending on age)
- **Vitamin D:** 1000–2000 IU/day; higher doses (4000 IU/day) if deficient; monitor serum 25-OH vitamin D
- **Weight-bearing when possible:** Standing transfers (bed to commode), even briefly, provide some bone loading
- **Whole body vibration:** Some evidence supports vibration platforms for bone health in immobilized populations (requires equipment; minimal energy expenditure)

19.5.3 Growth Considerations

Severe illness during adolescence can affect growth. Monitoring:

- **Height and weight:** Track on growth curves at each medical contact
- **Nutritional status:** Ensure adequate calories and protein despite reduced appetite
- **Pubertal development:** Chronic illness can delay puberty; document Tanner staging
- **Growth velocity:** Slowing growth velocity may warrant endocrine evaluation

19.5.4 Skin Integrity

Pressure ulcers are rare in pediatric ME/CFS but can occur with prolonged immobility:

- **Position changes:** Change position every 2–4 hours (can be done during wakefulness without disrupting sleep)
- **Pressure redistribution:** Specialized mattress (foam, alternating pressure) for prolonged bedrest
- **Skin inspection:** Check pressure points (heels, sacrum, scapulae) regularly
- **Nutrition:** Adequate protein and vitamin C support skin integrity
- **Moisture management:** Address incontinence promptly if present

19.5.5 Cardiovascular Deconditioning

Prolonged bedrest causes cardiovascular deconditioning (reduced stroke volume, reduced exercise capacity), which can worsen orthostatic intolerance.

Mitigation:

- **Head-up tilt:** Elevating head of bed 10–15 degrees provides mild orthostatic challenge even while lying down
- **Reclined exercises:** If tolerated, very gentle exercises while reclined (ankle pumps, leg slides) may maintain some conditioning without triggering PEM
- **Gradual mobilization:** When improvement allows, very gradual increase in upright time (5 minutes sitting, building slowly over weeks)
- **Monitor for overexertion:** Any intervention causing symptom worsening should be reduced

19.6 Educational Continuity

Loss of education is one of the most distressing aspects of severe pediatric ME/CFS. Children lose not only academic progress but also peer connections, sense of identity, and future opportunities. Maintaining educational continuity to the extent possible is important.

19.6.1 Legal Rights to Education

In the United States, severely ill children retain legal rights to education:

- **Individuals with Disabilities Education Act (IDEA):** Entitles children with disabilities to Free Appropriate Public Education (FAPE) through Individualized Education Program (IEP)
- **Section 504 of Rehabilitation Act:** Prohibits discrimination against students with disabilities; provides for 504 Plans with accommodations
- **Homebound instruction:** Most states require school districts to provide instruction at home for students unable to attend school due to illness

19.6.2 Homebound Instruction Protocols

Homebound instruction is typically provided when a child is expected to be out of school for more than 2–4 weeks (varies by state). Effective homebound instruction for ME/CFS requires:

- **Physician certification:** Letter documenting medical necessity, expected duration, and any limitations on instruction
- **Flexible scheduling:** Instruction during child's best hours; willingness to reschedule for crashes

- **Reduced hours:** Standard homebound instruction (often 5–10 hours/week) may be excessive for severely ill children; request reduction as needed
- **Modified curriculum:** Reduced course load; prioritize core subjects
- **Alternative assessments:** Portfolio assessment, oral exams, or modified testing as needed
- **Communication with teacher:** Teacher should understand ME/CFS and not interpret illness as laziness or school avoidance

19.6.3 Maintaining Academic Trajectory

For children who cannot tolerate any formal instruction, maintaining some academic trajectory:

- **Priority subjects:** Focus on reading/language arts and math as foundations
- **Interest-based learning:** Learning driven by child's interests may be better tolerated than required curriculum
- **Audiobooks:** Maintain exposure to literature and learning even when reading is impossible
- **Educational podcasts and videos:** Low-demand learning modalities
- **Credit accumulation:** Work with school to ensure any completed work counts toward credits

19.6.4 Social Connection Strategies

Social isolation compounds the impact of severe illness. Strategies to maintain peer connection:

- **Low-demand communication:** Text messaging, voice messages allow asynchronous connection
- **Brief video calls:** 5–15 minute video calls with friends during good periods
- **Online communities:** Age-appropriate online groups for chronic illness (finding peers who understand)
- **Creative projects:** Collaborative online projects with friends (writing, gaming, art)
- **Card/letter exchange:** Receiving mail from friends; dictating responses if writing impossible

19.6.5 Long-Term Planning

For children with prolonged severe illness:

- **Gap years:** Consider formal gap year(s) rather than falling behind; reenroll when improved

- **Graduation timeline flexibility:** Work with school on extended timeline for graduation requirements
- **GED as option:** High school equivalency may be appropriate if traditional graduation impossible
- **Community college start:** When ready, community college allows part-time enrollment with transfer to four-year institution later
- **Focus on recovery:** Academic recovery can follow physical recovery; pressure to maintain academic pace often impedes physical improvement

19.7 Family Support and Caregiver Coordination

Severe pediatric ME/CFS affects the entire family unit. Parents become full-time caregivers while managing their own grief, frustration, and exhaustion. Siblings may feel neglected or resentful. Family-centered care must address these dynamics.

19.7.1 Parent Education

Parents need accurate information to provide appropriate care and advocate effectively:

- **Understanding severity:** Severe ME/CFS is a serious, disabling illness; the child's limitations are real, not behavioral
- **Realistic expectations:** Improvement is likely in children but may take years; recovery cannot be rushed
- **Crash recognition:** Parents must learn to recognize early signs of PEM and enforce rest
- **Pacing enforcement:** Parents may need to limit activity even when child feels well ("good day" overexertion triggers crashes)
- **Medical advocacy:** Parents often must educate healthcare providers about ME/CFS

19.7.2 Caregiver Burden and Support

Caregiving for a severely ill child is exhausting and isolating. Parents are at high risk for burnout, depression, and relationship stress.

Support strategies:

- **Respite care:** Regular breaks from caregiving (family members, hired help, or respite programs)
- **Support groups:** Online or in-person groups for parents of children with ME/CFS
- **Mental health support:** Individual or couples therapy for parents
- **Practical help:** Accept help with meals, errands, household tasks
- **Caregiver health:** Parents must maintain their own medical care and self-care

19.7.3 Sibling Considerations

Healthy siblings of severely ill children face their own challenges:

- **Reduced parental attention:** Parents' energy goes to the ill child
- **Household disruption:** Quiet house, restricted activities to avoid triggering sibling's symptoms
- **Emotional responses:** Worry about ill sibling, resentment of attention imbalance, guilt about negative feelings
- **Social impact:** May be reluctant to bring friends home; may feel embarrassed explaining sibling's illness

Supporting siblings:

- **One-on-one time:** Ensure each healthy sibling gets individual attention from each parent
- **Age-appropriate explanation:** Help siblings understand the illness at their developmental level
- **Validate feelings:** All feelings (worry, resentment, sadness) are normal and allowed
- **Maintain normalcy:** Healthy siblings should continue their activities and friendships
- **Role limits:** Siblings should not become caregivers (brief help appropriate, but not primary care role)

19.7.4 Financial Navigation

Severe pediatric ME/CFS creates financial strain:

- **Lost parental work:** One or both parents may need to reduce work or stop working to provide care
- **Medical expenses:** Specialty care, medications, supplements, equipment often poorly covered by insurance
- **Home modifications:** May require investment in equipment or home changes
- **Lost child income:** Adolescents lose ability to work part-time jobs

Resources:

- **Family Medical Leave Act (FMLA):** Provides job-protected leave for care of ill child (unpaid, but protects employment)
- **State disability programs:** Some states have family caregiver support programs
- **SSI for child:** Supplemental Security Income may be available for severely disabled children in low-income families
- **Charitable assistance:** ME/CFS organizations and general chronic illness charities may provide grants
- **Medical bill negotiation:** Many hospitals offer charity care or payment plans

19.7.5 Mental Health Support for Family

Depression and anxiety are common in both patients and family members. Warning signs requiring professional intervention:

- **In patient:** Statements of hopelessness, suicidal ideation, complete disengagement, refusing all care
- **In parents:** Inability to function, severe depression, relationship breakdown, child neglect
- **In siblings:** Academic decline, behavioral changes, social withdrawal, expressed resentment toward ill sibling

Seek mental health support proactively rather than waiting for crisis. Family therapy addressing the systemic impact of chronic illness can benefit all members.

19.8 Red Flags and Emergency Protocols

Severely ill children require vigilant monitoring for complications requiring urgent intervention. The threshold for escalation should be lower in pediatrics than in adults.

19.8.1 Malnutrition and Feeding Concerns

Severe ME/CFS can impair ability to eat adequately:

- **Contributing factors:** Nausea, gastroparesis, food sensitivities, energy cost of eating, cognitive difficulty with meal planning
- **Warning signs:** Weight loss >10% body weight, BMI declining below healthy range, dehydration

Interventions:

- **Calorie-dense foods:** Prioritize high-calorie, high-protein foods that require minimal eating effort
- **Liquid nutrition:** Ensure shakes, smoothies if solid food difficult
- **Small frequent meals:** 6–8 small meals/snacks rather than 3 large meals
- **Appetite stimulants:** Consider mirtazapine (appetite-stimulating side effect) if severely underweight
- **Tube feeding consideration:** If unable to maintain adequate nutrition orally and weight loss continues, nasogastric or PEG tube feeding may be necessary—this is a serious decision requiring specialist involvement

19.8.2 Dehydration

Dehydration is particularly dangerous given the importance of blood volume for orthostatic tolerance.

Warning signs:

- Decreased urine output (less than 3–4 voids/day)
- Dark urine
- Dry mucous membranes
- Worsening orthostatic symptoms beyond baseline
- Tachycardia at rest

Intervention:

- Increase oral fluids if tolerated
- Oral rehydration solutions
- IV fluids if oral rehydration inadequate or not tolerated

19.8.3 Severe Orthostatic Symptoms

OI symptoms requiring urgent evaluation:

- Syncope (fainting)
- Near-syncope with falls or injuries
- Chest pain with activity
- Palpitations suggestive of arrhythmia (irregular, rapid, or pounding)

These may indicate inadequately treated OI, cardiac arrhythmia, or other cardiovascular pathology requiring workup.

19.8.4 Mental Health Crisis

Adolescents with severe ME/CFS are at increased risk for depression and suicidal ideation due to:

- Loss of functional life, identity, and social connections
- Chronic pain and suffering
- Hopelessness about recovery
- Feeling disbelieved by healthcare providers

△ **Warning: Suicide Risk Assessment**

Take ALL expressions of suicidal thoughts seriously. Warning signs:

- Statements about wanting to die, being a burden, having no future
- Giving away possessions
- Withdrawal from remaining connections
- Sudden calm after period of distress (may indicate decision made)
- Seeking means (medications, access to weapons)

If suicidal ideation is present:

- Do not leave child alone
- Remove access to means (medications, sharp objects)
- Contact mental health crisis services or emergency department
- National Suicide Prevention Lifeline (US): 988
- Crisis Text Line: Text HOME to 741741

Mental health care for suicidal adolescents with ME/CFS requires providers who understand the illness. Psychiatric hospitalization, with its stimulation and forced activity, can worsen ME/CFS and should be avoided if possible through intensive outpatient support.

19.8.5 Distinguishing Severity from Concurrent Illness

Severely ill ME/CFS patients may have difficulty recognizing new acute illness (infection, appendicitis, other medical emergency) because chronic symptoms mask new symptoms.

Concerning changes from baseline:

- New fever (ME/CFS typically does not cause fever)
- New localized pain (abdominal, chest, flank)
- Acute worsening of specific symptoms beyond usual fluctuation
- New neurological symptoms (weakness, numbness, vision changes)

Maintain low threshold for medical evaluation of new symptoms. Do not assume all symptoms are ME/CFS.

19.9 Evidence That Severe Pediatric Disease CAN Reverse

Despite the severity of current illness, there is strong evidence that severe pediatric ME/CFS can reverse—a critical source of hope for families.

19.9.1 Prognosis Data

Long-term follow-up studies demonstrate substantially better outcomes in pediatric ME/CFS than adult disease [78]:

- **Recovery at 5 years:** 38%
- **Recovery at 10 years:** 68%
- **Overall improvement or recovery:** 54–94% across studies
- **Mean illness duration:** 5 years (range 1–15 years)
- **Functional status at 10-year follow-up:** Mean 8/10
- **Proportion still very unwell:** Only 5% with function <6/10 at 10 years
- **Working or studying full-time:** 63% at follow-up

These figures include children who were severely affected. The data demonstrate that even severe pediatric ME/CFS is not necessarily permanent.

19.9.2 Time to Improvement

Realistic timelines for improvement:

- **Months 1–6:** Stabilization with appropriate treatment; symptom reduction possible but functional improvement limited
- **Months 6–18:** Gradual improvement in many; may transition from severe to moderate
- **Years 2–5:** Continued slow improvement; many achieve significant recovery
- **Years 5–10:** Most who will recover have recovered by this point

Improvement is typically slow and non-linear. Good weeks are followed by setbacks. The trajectory is overall positive even if daily experience fluctuates.

19.9.3 Factors Associated with Better Outcomes

Research suggests better outcomes are associated with:

- **Shorter diagnostic delay:** Earlier diagnosis and appropriate management
- **Younger age at onset:** Pre-adolescent onset may have slightly better prognosis
- **Absence of comorbidities:** Children without additional chronic conditions fare better
- **Adequate rest and accommodations:** Avoiding repeated severe crashes
- **Family support:** Stable, supportive family environment

19.9.4 Contrast with Adult Prognosis

Adult ME/CFS has dramatically worse outcomes (see Chapter 5 and Section 5.5.2 for comprehensive analysis):

- Recovery: Only 5% (median across studies)
- Improvement: $\leq 22\%$
- Most adults with ME/CFS have lifelong illness

The stark difference between pediatric and adult outcomes suggests that biological factors related to developmental plasticity, or barriers to recovery present in adults but absent in children (continued work demands, financial pressures), may significantly influence trajectory. For analysis of why adult recovery rates may be underestimated and the evidence limitations affecting these statistics, see Section 5.5.2.

19.9.5 Hope Maintenance with Realistic Expectations

Families need both hope and realism:

- **Hope:** The majority of children with ME/CFS, including severe cases, improve significantly or recover. This is well-documented.
- **Realism:** Improvement takes years, not weeks. Recovery cannot be rushed. Some children do not fully recover and require ongoing accommodations into adulthood.
- **Focus:** The goal during severe illness is symptom management, preventing complications, and maintaining developmental trajectory—not forcing recovery
- **Success metrics:** Recovery is not the only success. A child who improves from bedbound to attending school part-time has had an excellent outcome, even if not fully “recovered.”

19.9.6 Critical Window for Early Intervention

The better prognosis in pediatric ME/CFS suggests a critical intervention window that may close as patients age. This underscores the urgency of appropriate treatment in severe pediatric cases:

- Do not delay treatment waiting for recovery
- Aggressive symptom management (OI treatment, sleep optimization) may facilitate natural recovery processes
- Avoiding severe crashes (strict pacing) may preserve recovery potential
- The developing nervous system and immune system may have plasticity that allows recovery if not damaged by repeated overexertion

★ Key Point: The Pediatric Advantage

Children with ME/CFS have a window of opportunity for recovery that appears to narrow with age and illness duration. Appropriate early intervention—aggressive symptom management, strict pacing, accommodations, and avoidance of harmful treatments like GET—may maximize the likelihood of utilizing this window. Every year matters. Every severe crash may reduce recovery potential. The urgency of correct management cannot be overstated.

19.10 Summary of Key Recommendations

1. **Prioritize OI treatment:** Orthostatic intolerance affects 70–90% of pediatric ME/CFS patients and is often the most treatable symptom. Start with non-pharmacological measures (hydration, salt, compression, positioning) and add medications (fludrocortisone, midodrine) if needed.
2. **Optimize sleep:** Melatonin is first-line for pediatric sleep dysfunction. Low-dose trazodone or amitriptyline are second-line options. Maintain consistent sleep-wake schedule even when homebound.
3. **Manage pain appropriately:** Use acetaminophen and NSAIDs first-line. Add gabapentin or low-dose amitriptyline for neuropathic pain. Avoid routine opioid use. Referral to pediatric pain specialist for refractory cases.
4. **Prevent bedrest complications:** Daily passive range of motion prevents contractures. Ensure adequate calcium, vitamin D, and occasional weight-bearing to prevent bone loss. Monitor growth.
5. **Maintain educational continuity:** Secure homebound instruction through school district. Reduce academic demands to match capacity. Focus on core subjects and interest-based learning. Plan flexibly for graduation timeline.
6. **Support the whole family:** Parents need respite and support. Siblings need individual attention and validation. Financial and mental health resources should be mobilized proactively.
7. **Monitor for emergencies:** Maintain low threshold for evaluating new symptoms. Watch for malnutrition, dehydration, severe OI complications, and mental health crises.
8. **Maintain hope:** Pediatric ME/CFS prognosis is substantially better than adult disease. With appropriate management, most children improve significantly or recover over time. Recovery cannot be rushed, but it is achievable.

→ Continued: Related Content

For ambulatory pediatric patients who are still attending school, see Chapter 20 for school accommodation frameworks, IEP/504 planning, and pacing strategies for school-attending children. For adult severe ME/CFS management, see Chapter 17. For detailed discussion of pathophysiology underlying these interventions, see Chapter 10 (orthostatic mechanisms), Chapter 6 (metabolic dysfunction), and Chapter 7 (immune abnormalities).

20 Pediatric ME/CFS: School-Attending and Ambulatory Cases

Children and adolescents with mild to moderate ME/CFS who retain capacity for school attendance—even partial attendance—represent a critical population for early intervention. The substantially better prognosis in pediatric ME/CFS (54–94% improvement or recovery versus $\leq 22\%$ in adults) creates both opportunity and urgency: appropriate early management may preserve the window for recovery, while inappropriate management (particularly graded exercise therapy) risks progression to severe, potentially irreversible disease [393, 534].

This chapter addresses ambulatory pediatric patients who can attend school at least part-time. For children who are housebound or bedbound, see Chapter 19. For adult mild-moderate management, see Chapter 18.

20.1 The Early Intervention Imperative

Early intervention in pediatric ME/CFS is not merely beneficial—it may be critical for preserving recovery potential. The window of opportunity that distinguishes pediatric from adult prognosis appears to narrow with illness duration and repeated severe crashes.

20.1.1 Why Early Intervention is More Critical in Pediatrics

Several factors make early intervention uniquely important in pediatric ME/CFS:

1. **Better prognosis creates higher stakes:** The 54–94% improvement/recovery rate in pediatric ME/CFS (versus $\leq 22\%$ in adults) means there is more to preserve. Inappropriate management that converts a child from the “likely to recover” group to the “chronic severe” group represents a catastrophic outcome that might have been prevented.
2. **Critical developmental window:** Lost school years during childhood and adolescence have consequences beyond immediate disability. Missed education, interrupted social development, and disrupted identity formation during formative years create cumulative deficits that persist even after physical recovery.
3. **Preventing progression to severe disease:** Many severely affected adults report that their ME/CFS began as mild-moderate disease that worsened due to continued overexertion, often encouraged by well-meaning but misinformed healthcare providers, teachers, and parents. Early recognition and appropriate pacing may prevent this trajectory.

4. **Evidence that early intervention improves outcomes:** Studies consistently show that shorter diagnostic delay correlates with better prognosis. Patients diagnosed and managed appropriately within the first year have better long-term outcomes than those who struggle for years before receiving correct diagnosis and treatment [78].
5. **Developing nervous system plasticity:** The pediatric nervous system and immune system retain plasticity that may facilitate recovery—but this plasticity may be damaged by repeated severe crashes or inappropriate treatments. Preserving this biological capacity requires appropriate management from the earliest stages.

20.1.2 The Cost of Delayed or Inappropriate Intervention

Conversely, delayed or inappropriate intervention carries substantial risks:

- **Progression to severe disease:** Children pushed to maintain normal activity levels despite symptoms frequently progress from mild to moderate to severe ME/CFS over months to years
- **Cumulative crash damage:** Each severe crash may cause partially irreversible damage; the “crash limit rule” from patient communities suggests tolerance for severe crashes is limited. The crash dose-response framework (Chapter 18, §18.3.3) explains why large crashes cause disproportionate, irreversible harm through ATP depletion thresholds, mitochondrial turnover limits, inflammatory cascade intensity, and epigenetic locking mechanisms.
- **Lost recovery potential:** The window for pediatric recovery may close with prolonged illness duration, converting a recoverable case into a chronic condition. The vicious cycle recruitment cascade (Chapter 2, §2.1) shows how initially single-cycle dysfunction progressively recruits additional cycles, reducing escapability over time.
- **Educational derailment:** Each semester of academic struggle or failure compounds into long-term educational deficits
- **Psychological harm:** Repeated experiences of pushing through symptoms, being disbelieved, and watching function decline create lasting psychological trauma

★ Key Point: The Urgency of Correct Early Management

Every month of delayed diagnosis, every crash caused by inappropriate pressure to exercise, and every semester of forced school attendance without accommodations potentially reduces the likelihood of full recovery. The pediatric advantage is not automatic—it must be preserved through appropriate management from the earliest stages of illness.

⚠ Limitation 1: Paediatric Ambulatory Protocols: Limited Trial Evidence for Early Intervention Claims

The “early intervention imperative” articulated above is clinically reasonable but rests on indirect evidence:

- The association between shorter diagnostic delay and better prognosis is observa-

tional and subject to confounding (milder cases may be both diagnosed faster and recover more readily, independent of intervention timing).

- No study has compared early aggressive treatment versus standard care in newly diagnosed paediatric ME/CFS to determine whether early intervention causally improves outcomes.
- Educational accommodation recommendations are based on expert consensus and disability law, not on controlled studies demonstrating that specific accommodation strategies improve ME/CFS outcomes.
- The “crash limit rule” and “irreversible crash damage” concepts from patient communities have face validity but have not been quantified or validated by longitudinal studies measuring crash frequency against long-term prognosis.

20.2 Subtype Assessment for Ambulatory Pediatric Cases

Early identification of which systems are primarily affected helps guide treatment prioritization and educational/activity accommodations. Ambulatory pediatric patients (unlike severe cases) retain the potential for recovery across all subtypes, making early appropriate intervention particularly critical.

→ Recommendation 1: Pediatric Subtype Assessment and Early Intervention Prioritization

Rationale: The selective energy dysfunction hypothesis (Chapter 14.24) proposes that ME/CFS involves different primary compartment dysfunction across subtypes. Identifying your child’s dominant subtype guides which interventions to prioritize.

Parent/caregiver assessment:

1. What improved LEAST with rest?

- Forgetfulness, poor concentration, difficulty learning new things → **CNS-Primary**
- Dizziness, tachycardia, feeling faint when standing → **Autonomic-Primary**
- Muscle weakness, fatigue, widespread pain → **Peripheral-Primary**
- Multiple symptoms equally severe → **Global**

2. How does your child perform across different demands?

- Can walk/play physically but struggles academically → **Suggests CNS-Primary**
- Academic performance okay but can’t tolerate standing/activity → **Suggests Autonomic-Primary**
- Both physical and cognitive demands limited by fatigue/pain → **Suggests Peripheral-Primary**

3. School accommodations needed?

- Difficulty concentrating, reading, learning new material → **Suggests CNS-Primary**
- Can't attend full days, needs frequent rest periods → **Suggests multiple systems, likely Autonomic or Global**
- Fatigue, weakness, exercise intolerance → **Suggests Peripheral-Primary or Global**

Early intervention prioritization by subtype:

Subtype A (CNS-Primary): Cognitive dysfunction dominates

- **Priority 1:** Cognitive support through education accommodations; learning aids; shortened school days to maximize cognitive capacity during peak hours
- **Priority 2:** Sleep optimization (crucial for pediatric brain development and cognitive function recovery)
- **Priority 3:** Gentle nutritional support; omega-3 supplementation for brain health
- **School accommodation:** Extended time on tests, preferential seating near teacher (reduces cognitive load of filtering other conversations), written instructions rather than verbal (reduces processing demand)

Subtype B (Autonomic-Primary): Orthostatic intolerance dominates

- **Priority 1:** Hydration protocol (2.5–3 liters/day for teens); salt loading if medically appropriate
- **Priority 2:** Compression garments (age-appropriate sizing); frequent position changes during school day
- **Priority 3:** Medications if non-pharmacological measures insufficient (mido-drine is pediatric-approved for POTS)
- **School accommodation:** Preferential seating (lying down or semi-reclined if possible), permission to move around class, frequent water breaks, cooler classroom if heat-triggered symptoms

Subtype C (Peripheral-Primary): Muscle weakness/fatigue dominates

- **Priority 1:** Activity pacing; strict adherence to energy envelope
- **Priority 2:** Nutritional optimization (adequate protein for muscle recovery, calories for growth)
- **Priority 3:** Gentle mitochondrial support (CoQ10, carnitine—pediatric doses)
- **School accommodation:** Reduced class load, exemption from PE, opportunity for rest periods between subjects

Subtype D (Global): Multi-system involvement

- **Approach:** Comprehensive accommodations addressing all domains
- **Strategy:** Start with hydration + sleep + pacing (foundational for all subtypes), then add domain-specific treatments
- **School accommodation:** Combination of measures—half-day or part-time

attendance, reduced academic load, frequent rest periods, accommodations from all subtype categories as needed

- **Critical:** Do NOT push your child to maintain full schedule; early appropriate accommodation prevents progression to severe disease

Evidence level: Plausible (subtype framework from Chapter 14.24, adapted for pediatrics with school considerations)

Key principle: Subtype assessment informs prioritization but should NOT delay implementation of foundational interventions (pacing, sleep, hydration) while optimizing.

20.3 Diagnosis and Assessment

Accurate diagnosis is the foundation of appropriate treatment. Pediatric ME/CFS diagnosis requires modifications to adult criteria to account for developmental context.

20.3.1 Pediatric Diagnostic Criteria Modifications

The core diagnostic criteria for ME/CFS (substantial reduction in activity, post-exertional malaise, unrefreshing sleep, plus cognitive dysfunction or orthostatic intolerance) apply to pediatrics with the following modifications [393]:

- **Duration threshold:** Standard criteria require 6 months of symptoms. For pediatric patients with severe presentation, 3 months may be sufficient for provisional diagnosis and treatment initiation. Waiting 6 months while a child deteriorates is not clinically justified.
- **Activity reduction assessment:** Adults are assessed by work capacity; children should be assessed by school attendance, extracurricular participation, social activities, and self-care—all compared to pre-illness baseline, not to peers. A child who previously played sports, maintained friendships, and earned good grades but now struggles with half-day school attendance has experienced substantial activity reduction regardless of how this compares to “average.”
- **Symptom reporting:** Younger children may have difficulty articulating symptoms like “brain fog” or “post-exertional malaise.” Parent observation and age-appropriate questioning are essential. Ask about specific situations: “Can you play as long as you used to?” “How do you feel the day after a busy day?” “Do you get tired from things that didn’t used to tire you?”
- **Post-exertional malaise recognition:** PEM in children may manifest as increased irritability, tearfulness, pain, or flu-like symptoms 12–72 hours after exertion, rather than the adult description of “crashing.” The pattern of delayed worsening after activity is diagnostic, regardless of the specific symptom expression.
- **Orthostatic intolerance prevalence:** OI affects 70–90% of pediatric ME/CFS patients—higher than adults [393]. Standing tests (10-minute stand test or tilt table testing) should

be part of evaluation. Many children with unexplained fatigue have unrecognized POTS or orthostatic hypotension.

20.3.2 Distinguishing ME/CFS from Differential Diagnoses

Several conditions can mimic or overlap with pediatric ME/CFS. Accurate diagnosis requires distinguishing ME/CFS from these alternatives while recognizing that some conditions (particularly POTS) frequently co-occur.

School Avoidance/School Refusal

Perhaps the most common misdiagnosis for pediatric ME/CFS is “school avoidance” or “school refusal”—the assumption that the child’s symptoms represent psychological resistance to school rather than physical illness.

Key distinguishing features:

- **PEM pattern:** Children with ME/CFS crash AFTER exertion, not before. A child who attends school Monday, crashes Tuesday-Wednesday, and improves by Friday demonstrates PEM. A child who refuses school Monday morning but is active in the afternoon may have school avoidance.
- **Weekend/vacation pattern:** ME/CFS does not resolve on weekends or vacations. Children with true ME/CFS remain symptomatic during breaks from school. School avoidance typically improves when school is not in session.
- **Desire to participate:** Children with ME/CFS typically WANT to attend school and participate in activities—they are limited by symptoms. Children with school avoidance may express reluctance or resistance beyond what symptoms would explain.
- **Physical findings:** Objective findings (positive tilt table test, documented tachycardia, measurable orthostatic hypotension) support ME/CFS diagnosis. School avoidance does not produce these findings.
- **Response to activity:** If a child can sustain hours of enjoyable activity (gaming, socializing) but cannot tolerate equivalent school time, this suggests different mechanisms may be at play. However, note that cognitive and physical exertion differ, and school may be more demanding than leisure activities even at similar durations.

△ **Warning: The Danger of Misdiagnosis as School Avoidance**

Misdiagnosing ME/CFS as school avoidance is not merely inconvenient—it is potentially dangerous. Children labeled as school-avoidant are often subjected to forced school attendance, psychological interventions aimed at overcoming “avoidance,” and graded exposure programs that push increasing activity. For children with actual ME/CFS, these interventions cause harm: physical deterioration, progression to severe disease, and psychological trauma from being disbelieved. Always take reported symptoms seriously and look for objective evidence of ME/CFS before concluding that symptoms are

psychologically mediated.

Post-Infectious Mononucleosis

Epstein-Barr virus (EBV) mononucleosis is a common trigger for ME/CFS in adolescents. Post-infectious fatigue lasting weeks to months is normal after mono; ME/CFS is diagnosed when symptoms persist beyond expected recovery and include characteristic features (PEM, OI, cognitive dysfunction).

- **Timeline:** Most adolescents recover from mono within 2–4 months. Fatigue persisting beyond 6 months with PEM suggests ME/CFS.
- **Management during early post-mono period:** Appropriate rest and pacing during the post-mono period may reduce risk of developing chronic ME/CFS. Pushing early return to full activity after mono is inadvisable.
- **EBV reactivation:** Some ME/CFS patients have evidence of chronic EBV reactivation; this represents ongoing viral contribution to ME/CFS rather than simple post-infectious fatigue.

Growth-Related Fatigue

Puberty is associated with increased sleep need and may cause temporary fatigue. Menstruation can cause cyclic fatigue from blood loss and iron deficiency.

- **Growth fatigue:** Responds to adequate sleep; does not cause PEM; does not worsen progressively
- **Iron deficiency:** Check ferritin in all adolescent females with fatigue; supplement if ferritin <30 ng/mL (some experts recommend <50)
- **Cyclic pattern:** Fatigue clearly worse around menstruation suggests hormonal or iron-related cause; ME/CFS fatigue is present throughout cycle (though may worsen perimenstrually)

POTS and Orthostatic Intolerance

POTS (Postural Orthostatic Tachycardia Syndrome) and ME/CFS frequently co-occur. Some patients have “pure” POTS without ME/CFS; others have ME/CFS with prominent POTS; many have overlapping presentations.

- **POTS without ME/CFS:** Symptoms primarily orthostatic (worse with standing, improved with lying); no significant PEM; exercise may actually improve symptoms (cardiovascular reconditioning)
- **ME/CFS with POTS:** Both orthostatic symptoms AND PEM; exercise worsens overall condition even if brief standing tolerance improves with treatment

- **Management differs:** Pure POTS may benefit from graded cardiovascular reconditioning; ME/CFS with POTS requires pacing and will worsen with exercise programs. Accurate diagnosis is essential.

Depression and Anxiety

Depression and anxiety can cause fatigue and are common comorbidities in ME/CFS. They do not cause PEM.

- **PEM distinguishes:** Depression causes persistent fatigue that may improve with pleasurable activity. ME/CFS causes fatigue that **WORSENS** after any sustained activity, pleasurable or not.
- **Comorbidity is common:** Many children with ME/CFS develop secondary depression from the experience of chronic illness, loss of function, and being disbelieved. Treating comorbid depression is appropriate but does not address ME/CFS.
- **Antidepressants for ME/CFS:** May help comorbid depression and certain symptoms (sleep, pain) but do not treat the underlying ME/CFS and do not cure the illness.

20.4 School Accommodation Framework

Educational accommodations are essential for children with ME/CFS to maintain academic progress while managing their illness. In the United States, two primary frameworks exist: IEP (Individualized Education Program) under IDEA, and 504 Plans under Section 504 of the Rehabilitation Act.

20.4.1 IEP vs. 504 Plan Decision Tree

Understanding the distinction between IEP and 504 Plans is essential for securing appropriate accommodations.

Section 504 Plans

Legal basis: Section 504 of the Rehabilitation Act of 1973 prohibits discrimination against individuals with disabilities in programs receiving federal funding, including public schools.

Eligibility: Any student with a physical or mental impairment that substantially limits one or more major life activities. ME/CFS clearly qualifies—it limits learning, concentrating, thinking, and physical activity.

What 504 provides:

- Accommodations to ensure equal access to education
- Modifications to the regular education program

- Does NOT provide specialized instruction or related services

Best for: Students who can access the general curriculum with accommodations (modified schedule, testing modifications, etc.) but do not need fundamentally different instruction.

Process:

1. Parent requests 504 evaluation in writing
2. School evaluates (may use existing medical documentation)
3. Team (including parents) develops 504 Plan
4. Plan reviewed annually (or as needed)

IEP (Individualized Education Program)

Legal basis: Individuals with Disabilities Education Act (IDEA) entitles children with qualifying disabilities to Free Appropriate Public Education (FAPE).

Eligibility: Student must have a disability in one of 13 categories AND require special education services. ME/CFS may qualify under “Other Health Impairment” (OHI)—a chronic or acute health problem that adversely affects educational performance.

What IEP provides:

- Specialized instruction tailored to student’s needs
- Related services (speech therapy, occupational therapy, counseling)
- Measurable annual goals
- Transition planning (for older students)
- More procedural protections than 504

Best for: Students who need specialized instruction, not just accommodations. May be appropriate for severely affected ME/CFS students who need fundamental modifications to curriculum or instruction method.

Process:

1. Parent requests evaluation in writing
2. School conducts comprehensive evaluation (60-day timeline in most states)
3. Eligibility determination meeting
4. If eligible, IEP team develops plan
5. Annual IEP review; reevaluation every 3 years

Decision Framework

► Protocol: 504 vs. IEP Decision Framework

Start with 504 if:

- Student can participate in general education curriculum
- Primary needs are accommodations (schedule, testing, attendance flexibility)
- Student does not require specialized instruction
- Faster implementation needed (504 process is typically quicker)

Pursue IEP if:

- Student cannot access general curriculum even with accommodations
- Specialized instruction is needed (modified curriculum, alternative teaching methods)
- Related services are needed (counseling, OT for cognitive strategies)
- Student is failing despite 504 accommodations
- Stronger legal protections are desired

Consider both:

- Start with 504 for immediate accommodations
- Pursue IEP evaluation simultaneously if specialized services may be needed
- IEP supersedes 504 (you don't need both)

20.4.2 Specific Accommodations by Symptom

Accommodations should be tailored to the specific symptoms limiting each student's educational participation.

Accommodations for Cognitive Dysfunction

Brain fog, difficulty concentrating, and memory problems are nearly universal in ME/CFS and significantly impact academic performance.

- **Extended test time:** 50–100% additional time; may need testing over multiple sessions
- **Reduced homework:** Homework should not exceed what can be completed in 50% of standard expected time; quality over quantity
- **Note-taking assistance:** Peer note-taker, teacher-provided notes, or audio recording of lectures
- **Alternative assignment formats:** Oral presentations instead of written papers; multiple-choice instead of essays; projects over time instead of timed tests
- **Preferential seating:** Front of room reduces distraction; near door allows quiet exit if overwhelmed
- **Reduced course load:** Fewer classes per semester; prioritize core requirements

- **Calculator/reference sheet use:** Reduce cognitive load of memorization
- **Quiet testing location:** Separate room reduces sensory demands during high-stakes testing

Accommodations for Orthostatic Intolerance

OI symptoms (dizziness, lightheadedness, difficulty standing) require specific environmental modifications.

- **Unlimited water and snacks:** Student should be permitted water bottle and salty snacks at all times; bathroom access without restriction
- **Sitting privileges:** Permission to sit during assemblies, pledge, or other standing activities
- **Elevator access:** If school has multiple floors; stair climbing is particularly triggering for OI
- **Climate control:** Access to cooler environments; heat worsens OI
- **Rest breaks:** Permission to lie down in nurse's office when symptomatic
- **Compression garment permission:** Student may need to wear compression stockings or shorts; should not be restricted by dress code
- **Late arrival:** If mornings are worst (common), late start may improve function

Accommodations for Post-Exertional Malaise

PEM requires flexibility in attendance and assignment deadlines.

- **Excused absences for crashes:** PEM-related absences should not count against attendance requirements; documentation of ME/CFS diagnosis should suffice without requiring repeated physician notes
- **Flexible deadlines:** Extensions on assignments following crashes
- **Rest breaks during day:** Permission to leave class and rest when approaching energy limits
- **Reduced physical education:** Exemption from standard PE; may substitute with pacing-appropriate activity (gentle stretching, walking to tolerance) or academic alternative
- **Split attendance:** Part-day at school, part-day home instruction; maintains school enrollment and peer connection while reducing daily demands
- **Modified schedule:** Attend every other day; attend mornings only; attend for testing only

Accommodations for Sensory Sensitivity

Light, sound, and other sensory inputs may be overwhelming.

- **Sunglasses indoors:** If fluorescent lights are triggering

- **Noise-canceling headphones:** During independent work or when overwhelmed
- **Quiet testing location:** Separate room for tests
- **Reduced sensory environment:** Front-row seating away from windows; permission to step out of noisy environments
- **Digital textbooks:** Adjustable font size, screen brightness, text-to-speech options

20.4.3 Communication Templates

Effective advocacy requires clear communication with school personnel.

Letter Requesting Evaluation

A written request triggers the school's legal obligation to evaluate.

Dear [Principal/Special Education Director],

I am writing to formally request an evaluation of my child, [Name], for [504 Plan/special education services under IDEA]. [Name] has been diagnosed with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS), a chronic neuroimmune illness that significantly affects [his/her] ability to participate in school.

Specifically, [Name] experiences: [list 3–4 key symptoms affecting school—e.g., severe fatigue, difficulty concentrating, post-exertional malaise that causes crashes after school days, orthostatic intolerance].

These symptoms have resulted in: [describe educational impact—e.g., missing X days this semester, declining grades, inability to complete homework, difficulty participating in class].

I am enclosing medical documentation from [Name]'s physician confirming the ME/CFS diagnosis. Please contact me to schedule an evaluation meeting. I understand the school has [30/60 days depending on state] to complete the evaluation.

Thank you for your attention to this request.

Sincerely, [Parent Name]

School Nurse Protocol

Providing the school nurse with clear guidance improves daily symptom management.

[Student Name] has Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS). Please allow the following:

- 1. Rest in nurse's office when symptomatic (lying down with lights dimmed)*
- 2. Unlimited water and salty snacks*
- 3. Call parent for pickup if: [list specific symptoms warranting dismissal]*

4. Do NOT encourage “pushing through” symptoms—this worsens the condition
5. Symptoms often appear 12–72 hours AFTER overexertion (pattern of Monday school attendance, Tuesday-Wednesday crash is typical ME/CFS)

Emergency contact: [Parent phone]

Physician contact: [Physician name and phone]

Teacher Education Brief

Teachers who understand ME/CFS provide better support.

About ME/CFS:

ME/CFS is a chronic neuroimmune illness—not depression, laziness, or “just being tired.”

Key features:

1. Post-exertional malaise: Activity causes symptom worsening 12–72 hours later. A student who seems fine during class may crash the next day.
2. Cognitive dysfunction (“brain fog”): Difficulty concentrating, processing information, and remembering—even when appearing alert.
3. Energy limits: Students have a fixed “energy envelope.” Exceeding it causes crashes and potentially long-term worsening.

What helps: Flexible deadlines, reduced workload, rest breaks, quiet environment, believing the student’s reported symptoms.

What hurts: Pressure to “push through,” skepticism about symptoms, inconsistent accommodation implementation, comparing student to peers.

[Student Name]’s specific accommodations: [list from 504/IEP]

20.5 Pacing for Children and Adolescents

Pacing—staying within the energy envelope to prevent crashes—is the cornerstone of ME/CFS management. For children, pacing must be adapted to developmental stage and implemented with adult support.

20.5.1 Age-Appropriate Energy Management

The Energy Envelope Concept for Children

Explain the energy envelope in age-appropriate terms:

- **For younger children (6–10):** “Your body has a battery. When the battery gets low, you feel bad and need to rest. If you use too much battery, you’ll feel sick tomorrow. We need to save enough battery each day.”
- **For older children (11–14):** “You have a daily energy budget—like a bank account. Every activity costs energy. If you spend more than you have, you go into debt and crash later. We need to figure out your budget and stay within it.”
- **For adolescents (15–18):** Full energy envelope concept as described in adult chapters—fixed available energy, all activities cost energy, exceeding envelope triggers PEM and potentially long-term worsening.

Activity Tracking for Children

Monitoring activity and symptoms helps identify the energy envelope:

- **Younger children:** Parent tracks activity and symptoms; simple rating scales (happy face/sad face for symptoms)
- **Older children:** Child participates in tracking with parent support; symptom diary
- **Adolescents:** Self-tracking with apps; parent reviews patterns

Track:

- Activities each day (school hours, homework time, physical activity, social activity)
- Symptom rating (1–10 scale, or simple categories)
- PEM episodes (when they occur, what preceded them by 12–72 hours)

After 2 weeks, identify: What activity level consistently avoids PEM? That is the energy envelope.

Heart Rate Monitoring

Heart rate monitoring provides objective guidance for staying within the aerobic threshold:

- **Calculate threshold:** $(220 - \text{age}) \times 0.55$ to 0.60 for ME/CFS patients (lower than standard aerobic threshold)
- **Wearable monitors:** Fitness watches or chest straps; many children will engage with wearable technology
- **Gamification:** Some children respond well to “keeping the number below X” as a game; apps can provide alerts

- **Ideally:** Cardiopulmonary exercise testing (CPET) provides actual measured anaerobic threshold; standard formulas are approximations

★ Key Point: Heart Rate as Training Wheels

Heart rate monitoring is “training wheels” for pacing. Initially, use it to learn what activities and intensities exceed threshold. Over time, children learn to recognize body signals (early fatigue, heart racing, breathlessness) that indicate threshold approach. Eventually, most can pace effectively by feel, using heart rate monitoring to verify during uncertainty.

Distinguishing Pacing from Laziness

Parents and teachers may struggle to distinguish appropriate energy conservation from “laziness.” Key distinctions:

- **Desire:** Children with ME/CFS typically WANT to do activities but cannot. Lazy children do not want to do activities. If a child is distressed about missing activities, that suggests illness, not laziness.
- **PEM pattern:** Laziness does not cause delayed symptom worsening. If rest is followed by improvement and activity by delayed crashes, that is ME/CFS.
- **Consistency:** ME/CFS limits all sustained activity—even enjoyable activities. Laziness is selective.
- **Physical findings:** Documented tachycardia, orthostatic changes, and other objective findings support illness. Laziness does not produce these.

20.5.2 Managing the “Push Through” Pressure

Children with ME/CFS face enormous pressure to exceed their energy envelope—from parents, teachers, peers, healthcare providers, and their own desires. Managing this pressure is essential for preventing progression.

Academic Expectations

- **Communicate with school:** Ensure teachers understand that reduced output reflects illness, not lack of effort
- **Prioritize ruthlessly:** If only X hours of schoolwork are possible, focus on core requirements
- **Quality over quantity:** Completing fewer assignments well is better than struggling through everything poorly
- **Long-term perspective:** A year of reduced academics will not ruin a child’s future; progression to severe ME/CFS might
- **Alternative paths:** If traditional academic trajectory is impossible, explore alternatives (GED, community college, gap year)

Sports and Physical Activities

Sports decisions are particularly difficult. Recommendations:

- **Assess realistically:** Can the child participate without triggering PEM? If not, participation is harmful regardless of desire.
- **Modification options:** Can the child participate in reduced capacity (half practices, no games, coaching role)?
- **When to stop:** If any participation triggers crashes, stopping is necessary. This is not failure—it is medical management.
- **Alternative connection:** If team membership is important for social reasons, explore non-physical roles (manager, statistician)
- **Grieving:** Losing sports identity is a significant loss. Allow space for grief while maintaining the boundary.

Peer Pressure and Social Activities

Adolescents particularly struggle with social pressure to maintain normal activity:

- **Selective participation:** Can attend some events by choosing carefully (one party per month, not every weekend gathering)
- **Modified participation:** Attend early and leave before exhaustion; sit rather than stand; take rest breaks
- **Explain to friends:** Close friends can understand if given accurate information; they can become allies in pacing
- **Alternative socialization:** Low-energy options (movie at home, video calls, texting) maintain connection without physical cost
- **Quality over quantity:** Fewer, shorter, better-paced social interactions preserve relationships and health

Adolescent Identity and Disability Acceptance

Adolescence is a critical period for identity formation. Developing ME/CFS during this period creates challenges:

- **Disability as identity:** ME/CFS becomes part of identity; this is neither good nor bad, but should be acknowledged
- **Avoid self-blame:** The child did not cause their illness and cannot will themselves better
- **Realistic self-concept:** Learning to assess actual capacity accurately (not overestimating to seem “normal” or underestimating to avoid disappointment)
- **Finding meaning:** What activities, relationships, and goals remain possible? Focus on what can be done, not only what is lost
- **Peer support:** Connecting with other adolescents with ME/CFS or chronic illness reduces isolation and provides modeling

20.6 Medical Management

Medical management for ambulatory pediatric ME/CFS addresses the same symptoms as severe disease (Chapter 19) but with adjustments for the school-attending context.

20.6.1 Early Intervention Window Protocol

Pediatric ME/CFS has exceptional recovery potential—54–94% of children improve or recover compared to $\leq 22\%$ of adults. This window of opportunity is not automatic; it must be actively preserved through aggressive early intervention.

► Protocol 1: Early Intervention Window Preservation

Aggressive Early Treatment: Don't "Wait and See"

The window for pediatric recovery appears to narrow with illness duration and repeated severe crashes. Early intervention is not optional:

- **Start treatment immediately upon diagnosis:** Do not wait months for improvement or to “see how it goes.” Appropriate treatment should begin during the first months of illness.
- **Optimize OI treatment first:** This is the most treatable symptom and often improves multiple other symptoms (fatigue, cognitive dysfunction, exercise tolerance). Do not accept “mild OI” as a reason to defer treatment.
- **Aggressive sleep optimization:** Sleep dysfunction perpetuates neuroinflammation. Use pharmacological treatment (melatonin, trazodone) immediately if sleep hygiene is insufficient.
- **Infection prevention from day one:** Each infection causes both acute PEM and lasting sensitization increases. Preventing infections from the outset preserves recovery potential.
- **Avoid crashes through strict pacing:** Each crash is not merely a bad day—it is a potential step toward disease progression and lost recovery potential.

Brain-First Medication Sequence for School-Attending Pediatric Patients

The goal is improved cognitive function and fatigue tolerance while maintaining school attendance, WITHOUT enabling overexertion:

1. **Orthostatic intolerance treatment** (Step 1): Non-pharmacological measures first; add fludrocortisone or midodrine if needed. Many children experience dramatic cognitive improvement with effective OI management.
2. **Sleep optimization** (Step 2): Melatonin first-line; add trazodone or low-dose amitriptyline if melatonin insufficient. Adequate sleep is foundational for all other improvements.
3. **Low-dose naltrexone (LDN)** (Step 3, if needed): Consider if cognitive dysfunction or fatigue remain limiting despite OI and sleep optimization. Weight-adjusted dosing (0.1 mg/kg, max 4.5 mg). LDN modulates neuroinflammation without the risks of stimulants.

- Advantages: Good safety profile in pediatrics; addresses underlying neuroinflammatory mechanisms; does not mask symptoms or enable harmful overexertion
 - Timeline: 2–8 weeks for measurable effect
 - Monitoring: Track cognitive function and fatigue; look for improvement without stimulant-like side effects
4. **Cognitive support alternatives** (Step 4, if LDN insufficient): Before considering stimulants, maximize accommodations (extended time, reduced homework, preferential seating). Many children function adequately academically with supports alone.
5. **Stimulants only if necessary** (Step 5, specialist evaluation): If cognitive dysfunction remains severely limiting despite accommodation and above interventions:
- Stimulants (methylphenidate, amphetamine) may improve concentration but carry risks:
 - Can mask symptoms and enable overexertion beyond energy envelope
 - Tachycardia may worsen orthostatic intolerance
 - Appetite suppression impairs nutrition
 - Effects wear off, potentially revealing deeper fatigue
 - If used, must be coupled with strict monitoring to prevent exceeding energy envelope
 - Alternative: Low-dose modafinil has been used in some adolescents with limited pediatric data
 - **Intranasal delivery option:** If cognitive dysfunction persists despite above interventions, discuss intranasal dopamine precursors or other CNS-targeted compounds with a neurologist familiar with ME/CFS. Not standard care but mechanistically rational for BBB dysfunction (Chapter 14.24, lines 238–257). EXPERIMENTAL.

School Accommodation as Treatment

School accommodations are not mere kindness—they are therapeutic interventions that preserve recovery potential:

- **Reduced hours/flexible schedule:** Part-time attendance is legitimate medical management, not failure to “push through”
- **Reduced homework:** Homework that exceeds capacity causes crashes and sensitization. Capping homework is medical necessity.
- **Rest breaks:** Permission to leave class and rest when approaching energy limits prevents crashes during the school day
- **Flexible deadlines:** PEM-related absences and the need for deadline extensions are medical accommodations, not academic favoritism
- **Each crash prevented:** Each accommodation that prevents a crash preserves the recovery window

Infection Prevention Imperative

In pediatric patients, infection prevention has compounded importance:

1. **Prevents baseline decline:** As in adults, prevents the immediate PEM and symptom worsening from infection
2. **Preserves the pediatric recovery window:** Each infection creates lasting sensitization. Preventing infections preserves the biological conditions for recovery.
3. **Maintains school attendance continuity:** Post-infection crashes often cause absences that snowball into educational deficits. Infection prevention maintains educational continuity.
4. **Protects developmental trajectory:** Each crash during childhood has developmental consequences beyond the immediate functional decline.

Infection prevention strategies: N95/FFP2 masking during viral seasons, prompt antiviral treatment if infection occurs, consideration of prophylaxis during high-exposure periods (exam weeks, high-transmission seasons), and when possible, isolation of child from sick household members.

20.6.2 Orthostatic Intolerance (First-Line)

OI is the most treatable symptom domain in pediatric ME/CFS. Many children experience significant improvement in fatigue, cognitive function, and overall wellbeing with effective OI management.

Non-Pharmacological Measures

Start with non-pharmacological interventions (see Protocol 19.4.1 in Chapter 19 for details):

- **Hydration:** Age-appropriate targets (2–3 liters/day for adolescents)
- **Salt:** 3–8 grams sodium/day depending on weight
- **Compression:** Waist-high compression garments if tolerated
- **Positioning:** Avoid prolonged standing; sit when possible; leg crossing/tensing

School accommodations for OI: Ensure the 504/IEP includes water bottle access, bathroom access, sitting privileges, and late arrival if mornings are worst.

Compression Garments for School-Attending Children with Orthostatic Intolerance

→ Recommendation 2: Medical-Grade Compression Stockings for Pediatric OI

Mechanism: Compression reduces autonomic load by maintaining venous pressure, decreasing sympathetic activation required for orthostatic compensation (Chapter 14.24, lines 609–626).

Pediatric Implementation:

1. Compression class and fit:

- **Class II (20–30 mmHg):** Standard for pediatric POTS and OI
- **Waist-high or thigh-high:** More effective for OI than knee-high
- **Proper fitting:** Measure leg circumference; incorrect sizing loses effectiveness
- **Material:** Merino wool or medical-grade synthetic (breathable for school day wear)

2. School-day wearing schedule:

- **During school hours:** Wear throughout day (improved alertness, reduced afternoon crashes)
- **Before/after school:** Optional depending on activity level
- **At home:** Remove during recumbent rest or sleep
- **Total daily wear:** 6–10 hours typical

3. Expected benefits:

- Reduced tachycardia and dizziness during school day
- Improved cognitive clarity and focus
- Better school attendance and academic performance
- Reduced afternoon crashes during school
- May reduce need for medications

4. Practical school considerations:

- **Acceptance:** Many children prefer stockings to medications; ease into use
- **Fitting at school:** Some children may prefer removing at lunch/bathroom breaks
- **504 accommodation:** Can request "compression garment breaks" if needed
- **Cost:** \$30–60 per pair; insurance may cover with POTS diagnosis
- **Maintenance:** Replace every 3–6 months as compression decreases

5. Integration with other OI treatments:

- Essential component of first-line non-pharmacological approach
- Synergistic with salt loading and hydration
- Can be combined with medications if needed

Evidence level: Moderate (20–30 mmHg compression established for POTS; pediatric

ME/CFS data limited)

Expected outcomes: 25–50% reduction in orthostatic symptoms, improved school function, reduced fatigue during day when combined with salt/hydration protocol.

Success predictor: If child shows improvement in tachycardia or cognitive clarity within 1 week, continue indefinitely.

Pharmacological Management

If non-pharmacological measures are insufficient, medications may help (see Protocol 19.4.1 in Chapter 19 for detailed pediatric dosing):

- **Fludrocortisone:** Starting dose 0.05 mg daily; plasma volume expansion
- **Midodrine:** Starting dose 2.5 mg BID-TID; vasoconstriction (78% response rate in pediatric POTS) [338]
- **Pyridostigmine:** Starting dose 15–30 mg TID; autonomic support

Monitoring response: Track symptoms before and after OI treatment. Many children notice improved cognitive function, reduced fatigue, and better exercise tolerance within 1–2 weeks of effective OI management.

20.6.3 Mast Cell Activation Syndrome (MCAS) Management

Mast cell activation affects 30–50% of ME/CFS patients. In school-attending children, MCAS contributes to post-meal fatigue, brain fog fluctuations, and anxiety-like episodes through histamine and mediator release.

→ Recommendation 3: School-Attending Pediatric MCAS Management

Screening: Consider MCAS trial if child reports:

- Food sensitivities or intolerances (especially post-meal fatigue)
- Flushing, hives, itching
- Reactive to fragrances or chemicals
- GI symptoms (bloating, nausea, diarrhea)
- Unexplained anxiety or panic-like episodes

Age-Appropriate Treatment (pediatric dosing):

1. H1+H2 Antihistamine Combination:

- H1: Cetirizine 5 mg morning (younger) or 10 mg (adolescent), OR loratadine 5–10 mg daily
- H2: Famotidine 10–20 mg BID (younger children) to 20–40 mg BID (adolescents)
- Low-histamine diet (avoid aged cheese, fermented foods, cured meats, left-

overs >24 hours)

2. Optional: Quercetin supplement:

- 250–500 mg daily for younger children
- 500–1000 mg daily for adolescents
- Natural mast cell stabilizer; can combine with antihistamines

3. 4-week trial: If 30–50% improvement in energy or cognitive clarity, continue indefinitely

4. Prophylactic Intensification for School Events:

- **Before predictable triggers** (exams, presentations, school field trips, social events): Start 24 hours prior
 - Increase H1 antihistamine to maximum tolerated dose
 - Increase H2 to full therapeutic dose
 - Add quercetin if available
 - Strict low-histamine diet
- **Activity modification:**
 - Reduce non-essential activities day-of event
 - Ensure adequate rest before and after
 - Avoid additional cognitive or emotional demands
- **Parent tracking:**
 - WITHOUT prophylaxis: "School presentation triggered fatigue crash; needed 2 days recovery"
 - WITH prophylaxis: "Same presentation with prophylaxis caused minimal fatigue; recovered in 1 day"
 - Adjust future event planning based on prophylaxis efficacy

Evidence level: Moderate (pediatric MCAS management established; ME/CFS school-event crash mitigation outcomes pending)

Expected outcomes: If MCAS component is significant, 25–50% reduction in event-related fatigue or cognitive impact.

20.6.4 Sleep

Sleep dysfunction impairs school performance and exacerbates all other symptoms.

Sleep Hygiene for School-Attending Children

- **Screen time:** No screens 1–2 hours before bed; blue light filtering if screens are used
- **Homework timing:** Complete homework earlier in evening; avoid cognitive work close to bedtime
- **Consistent schedule:** Same sleep and wake times daily—including weekends (shifts disrupt circadian rhythm)

- **Social media:** Particularly disruptive for adolescents; phones should charge outside bedroom

School Start Time Considerations

Early school start times conflict with adolescent circadian biology and ME/CFS sleep dysfunction:

- **Request late start:** If school offers late start options, request them
- **Advocate for policy change:** Many schools are shifting to later start times based on adolescent sleep research; parents can advocate
- **504/IEP accommodation:** Late arrival can be an accommodation if early arrival is consistently problematic

Sleep Medications

If sleep hygiene is insufficient (see Protocol 19.4.4 in Chapter 19):

- **Melatonin:** 0.5–5 mg, 30–60 minutes before desired sleep time; first-line
- **Trazodone:** 12.5–50 mg at bedtime if melatonin insufficient
- **Low-dose amitriptyline:** 5–25 mg at bedtime; dual benefit for sleep and pain

Circadian Light Therapy for Sleep-Energy Alignment

→ Recommendation 4: Bright Light Therapy for Circadian Entrainment in School-Attending Children

Mechanism: Morning bright light exposure resets the circadian oscillator, improving alignment between energy availability and school schedule. Many school-attending children with delayed sleep phase struggle with early school start times; morning light therapy can advance sleep phase by 1–2 hours.

Pediatric School-Context Protocol:

1. **Equipment:** 10,000 lux light therapy box (\$25–100). Can be used while eating breakfast or doing morning routine.
2. **Timing:** 20–30 minutes immediately upon waking (within 30 minutes). Consistency is critical.
3. **Position:** 16–24 inches from child's face, 30° downward angle
4. **School integration:** If child struggles with early school start, morning light therapy 30–60 minutes before departure improves alertness and may gradually advance sleep phase by 45 min–2 hours over 2–4 weeks
5. **Do NOT use after 3pm** (risks delaying sleep further)

Evidence level: Moderate (circadian disruption documented; light therapy established for circadian disorders and adolescent sleep phase; ME/CFS-circadian-energy RCTs pending)

Expected outcomes:

- Earlier sleep onset at night (improved morning school attendance)
- Better morning alertness
- More consistent daytime energy
- Timeline: 2–4 weeks for phase shift

Sleep Spindle Enhancement via Acoustic Stimulation (Low Priority, Experimental)

→ Recommendation 5: Pink/White Noise for Sleep Architecture in School-Attending Children

Mechanism: Sleep spindles are brief bursts of brain electrical activity during light sleep that contribute to cognitive consolidation and memory protection. Acoustic stimulation during sleep (pink noise or pink noise burst sequences) can enhance spindle density, potentially improving memory encoding and resilience to cognitive fatigue. In school-attending children with ME/CFS, improved sleep architecture may translate to better cognitive performance during school day. See Chapter 14.24, lines 552–569, for detailed mechanism and neurophysiology.

Pediatric School-Context Protocol:

- **Equipment:** White or pink noise machine (\$20–60) or free app (e.g., myNoise, Noisli). Ensure sound quality is consistent and volume is safe for children (not exceeding 50 dB).
- **How to use for school-attending children:** Play pink or white noise from sleep onset through entire sleep period. Standard setting: 50 dB continuous, starting 15 minutes before desired sleep time. Can be used alongside other sleep interventions (melatonin, light therapy).
- **School integration:** Children who are struggling with test performance and memory retention may see modest benefit within 3–6 weeks. Better sleep architecture supports consolidation of school learning.
- **Tracking:** Monitor sleep quality (child report), morning alertness, and school cognitive performance (test scores, teacher feedback) over 4–6 weeks.

Evidence level: Speculative (spindle enhancement demonstrated in sleep neuroscience; ME/CFS sleep architecture and acoustic enhancement RCTs pending)

Why low priority for this population: School-attending ambulatory children should prioritize sleep hygiene, circadian light therapy, and activity pacing before considering experimental sleep architecture enhancement. Useful only if other sleep interventions are optimized and memory/cognitive consolidation remains a limiting factor.

20.6.5 Cognitive Symptoms

Cognitive dysfunction (“brain fog”) is often the most academically limiting symptom.

Academic Accommodations First

Before considering medications for cognitive symptoms, maximize accommodations:

- Reduce cognitive load (fewer classes, reduced homework)
- Provide cognitive supports (notes, extended time, calculator use)
- Schedule demanding tasks during best times of day
- Allow rest breaks to prevent cognitive fatigue

Many children can function adequately academically with accommodations alone, without cognitive medications.

Non-Pharmacological Cognitive Support

- **Organizational systems:** Calendars, apps, written reminders compensate for memory difficulties
- **Study strategies:** Spaced repetition, active recall, chunking information
- **Assistive technology:** Text-to-speech, speech-to-text, audiobooks
- **Cognitive rest:** Brief rest breaks (10 minutes every 30–45 minutes of cognitive work)

Pharmacological Considerations

If cognitive dysfunction remains severely limiting despite accommodations:

- **OI treatment first:** Cognitive function often improves substantially with effective OI management. Always optimize OI treatment before adding cognitive medications.
- **Stimulants:** Medications like methylphenidate or dextroamphetamine may improve concentration. However:
 - Developing brain concerns with chronic stimulant use
 - Cardiovascular effects (tachycardia) may worsen OI
 - Appetite suppression can impair nutrition
 - Does not address underlying cause
 - Risk of masking symptoms and enabling overexertion

Stimulants require specialist evaluation and careful monitoring; not first-line for pediatric ME/CFS cognitive symptoms.

- **Low-dose modafinil:** Used in some adolescents for fatigue/cognitive symptoms; limited pediatric data; similar concerns as stimulants regarding masking symptoms

△ Warning: Cognitive Medications May Enable Harmful Overexertion

Medications that improve cognitive function or reduce fatigue perception can be dangerous if they enable students to exceed their energy envelope. A student who takes stimulants to focus through a full school day may crash harder afterward. Cognitive medications should support sustainable activity within the energy envelope, not expand the envelope artificially.

Transcranial Direct Current Stimulation (tDCS) for School-Attending Adolescents**→ Recommendation 6: Pediatric tDCS for Cognitive Support (Age >12 with Caution)**

Mechanism: tDCS applied to dorsolateral prefrontal cortex (DLPFC) may reduce baseline energy cost of executive function, improving sustainable academic performance (Chapter 14.24, lines 207–228).

Pediatric Considerations:**1. Age requirements:**

- Minimum age: 12 years (limited pediatric safety data below this age)
- Adolescents 12+: Can consider with parental consent and physician supervision
- Requires mature understanding of electrode placement and safety

2. Protocol (pediatric-adapted):

- **Intensity:** 1–1.5 mA (lower than adult 2 mA due to developing brain)
- **Duration:** 15–20 minutes daily
- **Montage:** F3-F4 (DLPFC bilateral)
- **Frequency:** 5 days/week
- **Trial duration:** 4–8 weeks
- **Physician oversight:** Initial setup and safety screening required; ongoing monitoring

3. Academic tracking:

- Rate executive function daily during trial (focus, planning, organization)
- Track school performance (grade trends, assignment quality)
- Monitor for fatigue or mood changes
- Weekly summary by parent/adolescent

4. Safety and cautions:

- **Contraindications:** Metal implants in head, seizure history, pregnancy (adolescent females)
- **Mild side effects:** Tingling under electrodes (usually resolves), mild headache
- **Stop if:** Persistent headache, mood changes, anxiety, behavior problems

- **Parental supervision:** Required for electrode placement and safety monitoring

5. Critical caveat:

- tDCS is EXPERIMENTAL in pediatric ME/CFS; no published trials in this population
- May improve executive function without increasing energy envelope
- School accommodations remain essential; tDCS augments but does not replace them
- Risk: Improved focus might tempt increased academic load; strict energy envelope monitoring essential

Evidence level: Speculative (tDCS cognitive enhancement documented in other populations; pediatric ME/CFS safety and efficacy unknown)

Expected outcomes (if effective): 20–30% improvement in focus/organization/planning. Not expected to improve fatigue directly.

Who should try tDCS: Adolescents age 12+ with prominent executive dysfunction limiting academics despite accommodations and OI optimization.

20.6.6 Infection Prevention as Recovery Window Preservation

Observation 84 (Infection Prevention and Pediatric Recovery Window). In pediatric patients with ME/CFS, infection prevention has compounded importance beyond the baseline harm prevention seen in adults. While all ME/CFS patients experience worsening from infections (see Chapter 7, [infection damage ratchet hypothesis](#)), pediatric patients face an additional stake: preserving the window of opportunity for recovery that distinguishes pediatric from adult prognosis.

The Triple Impact of Infection in Pediatric ME/CFS:

1. **Acute PEM and baseline decline:** Each infection causes both immediate symptom exacerbation and lasting increases in baseline symptom severity—the ratchet effect common to all ME/CFS patients.
2. **Sensitization cascade:** In pediatric patients, infection-induced sensitization appears to narrow the recovery window. Children who experience multiple infections during their illness trajectory show worse long-term outcomes than those who successfully prevent infections. Each infection may represent a lost opportunity for recovery.
3. **Educational disruption with developmental consequences:** Post-infection crashes frequently force absences from school, creating cascading educational deficits. For developing children and adolescents, each semester of missed education has consequences beyond the immediate academic loss—social relationships are disrupted, identity formation around school-based activities is interrupted, and psychological impacts accumulate.

Clinical Implications for School-Attending Pediatric Patients:

Infection prevention is not merely supportive care—it is active preservation of recovery potential. Strategies include:

- **N95/FFP2 masking:** During known viral transmission in community or household. Given risk to recovery window, masking is medically justified even if symptomatic infections are not currently occurring.
- **Prompt antiviral treatment:** If viral infection suspected (respiratory symptoms, fever), initiate antivirals quickly to limit duration and severity. First-generation severity of infection is less damaging than allowing progression.
- **Prophylaxis during high-exposure periods:** Consider antiviral prophylaxis during high-transmission seasons or in households with sick members. The cost-benefit calculation is different for pediatric patients where infection prevention may preserve recovery trajectory.
- **Household isolation when possible:** When child is severely affected or during periods of particular vulnerability, isolation of the child from sick household members may be medically justified to preserve recovery potential.
- **School exposure management:** During high-transmission periods, discuss with school whether temporary remote learning is possible to reduce infection exposure. This is not academic coddling—it is medical management.

The Pediatric Difference:

Adult ME/CFS patients benefit from infection prevention to prevent baseline decline and symptom severity. Pediatric ME/CFS patients benefit from infection prevention for this reason AND to preserve the biological and psychological conditions in which recovery becomes possible. From a developmental perspective, each prevented infection is an investment in long-term outcome.

20.7 Activity and Exercise

The relationship between activity, exercise, and ME/CFS is widely misunderstood. This section provides evidence-based guidance.

20.7.1 Critical Warning: GET is Contraindicated

△ **Warning: Graded Exercise Therapy Causes Harm in ME/CFS**

Graded Exercise Therapy (GET)—structured programs of progressively increasing exercise—is contraindicated in ME/CFS. This applies to children as well as adults, with potentially greater harm in developing patients.

Evidence of harm:

- Patient surveys consistently report GET as one of the treatments most likely to cause deterioration

- The PACE trial, which promoted GET, has been discredited following independent reanalysis revealing methodological flaws and inflated claims [393]
- NICE guidelines (UK) no longer recommend GET for ME/CFS
- CDC guidance no longer promotes GET

Why GET harms ME/CFS patients:

- ME/CFS involves exercise intolerance as a core feature—the abnormal response to exertion IS the disease
- Progressive exercise triggers the post-exertional malaise mechanism, causing symptom worsening
- Repeated PEM episodes may cause cumulative damage and disease progression
- GET assumes deconditioning is the problem; in ME/CFS, the problem is abnormal energy metabolism and exercise response

Why pediatric harm may be greater:

- Children are less able to advocate against inappropriate treatment
- School PE requirements may enforce harmful exercise
- Parents and teachers may pressure “just exercise more” despite symptoms
- Developmental period may be more vulnerable to damage from repeated overexertion
- Better baseline prognosis means more to lose from inappropriate treatment

What to do if GET is recommended:

- Provide the recommending clinician with current NICE guidelines and CDC guidance
- Request documentation of the evidence base for GET in pediatric ME/CFS (none exists)
- Seek second opinion from ME/CFS-knowledgeable provider
- Document in writing your refusal of GET and the reasons

20.7.2 Safe Activity Maintenance

While GET is harmful, complete inactivity is also not ideal. The goal is maintaining baseline function within the energy envelope.

Establishing Baseline

1. Track current activity and symptom patterns for 2 weeks
2. Identify maximum sustainable activity level—the level that does NOT trigger PEM
3. This becomes baseline: the ceiling, not the floor

Safe Activity Principles

- **Stay within envelope:** Never intentionally exceed sustainable capacity
- **No progression until stable:** Do not increase activity unless current level has been sustained for weeks without PEM
- **Minimal increases:** If increasing, increase by only 5–10%, not more
- **Return to baseline after any PEM:** Any crash means activity was too high; return to lower level
- **“Good days” are not permission:** On good days, do NOT do more—bank the energy for bad days

Signs of Overexertion

Teach children to recognize early warning signs that they are approaching or exceeding limits:

- Heart rate rising above threshold
- Feeling “wired” or artificially energetic (adrenaline compensation)
- Difficulty catching breath
- Muscles feeling heavy or weak
- Cognitive function declining (losing words, difficulty concentrating)
- Onset of pain

When these signs appear, STOP and REST—do not “push through.”

Modified Physical Education

Standard PE requirements are often impossible for children with ME/CFS. Options:

- **Complete exemption:** If any physical activity triggers PEM, full PE exemption is appropriate
- **Modified participation:** Walking to tolerance, gentle stretching, yoga (restorative, not vigorous)
- **Academic alternative:** Health education coursework instead of physical activity
- **Manager/helper role:** Participate in PE class as manager, scorekeeper, or equipment manager without physical demands

20.8 Social Development Considerations

ME/CFS threatens normal social development during critical years. Active intervention is needed to maintain social health.

20.8.1 Maintaining Peer Connections

- **Quality over quantity:** A few close, understanding friends are more valuable than many superficial connections
- **Educate close friends:** Friends who understand ME/CFS can accommodate limitations and become allies
- **Low-energy socialization:** Video calls, texting, gaming together online, watching movies together
- **Brief in-person contacts:** Short visits are better than no visits; friends can visit the child at home
- **School-based support:** Lunchtime with friends even on days with reduced class attendance maintains school social connections

20.8.2 Online Social Options

For children too ill for regular in-person socializing, online connection becomes essential:

- **Chronic illness communities:** Online groups for young people with ME/CFS or chronic illness provide peer understanding
- **Gaming:** Multiplayer games provide social interaction with minimal physical demand
- **Interest-based communities:** Forums, Discord servers, or groups focused on the child's interests
- **Monitored appropriately:** Parents should ensure age-appropriate online safety while allowing necessary social connection

20.8.3 Extracurricular Modification

Extracurricular activities contribute to development and identity. Options:

- **Reduced participation:** Attend practices or meetings when able; miss when unable
- **Alternative roles:** Manager, mentor, or remote participant rather than active participant
- **Low-demand activities:** Activities that can be done seated, at home, or with flexible participation (art, writing, online clubs)
- **Cessation with grief support:** If no modification enables safe participation, stopping may be necessary. Allow space to grieve the loss.

20.8.4 Identity Formation

Adolescents with ME/CFS face identity challenges:

- **Disability identity:** ME/CFS becomes part of identity; this can be integrated healthily
- **Avoiding overidentification:** The child is not ONLY their illness; other aspects of identity matter

- **Future orientation:** What can the child aspire to? Realistic but hopeful goal-setting
- **Role models:** Adults with ME/CFS who have found meaningful lives can provide hope
- **Self-advocacy skills:** Learning to explain, advocate, and set boundaries is a life skill

20.8.5 Preventing Isolation

Social isolation increases depression risk and worsens prognosis. Active prevention:

- Schedule regular social contact (even brief, even remote)
- Maintain at least one activity that connects to peers
- Family activities provide social interaction when peer interaction is limited
- Professional support (counseling) if isolation is significant

20.9 Long-Term Management and Monitoring

ME/CFS is typically a chronic condition requiring ongoing management, though pediatric prognosis is substantially better than adult.

20.9.1 Trajectory Tracking

- **Baseline documentation:** Document function at diagnosis (school attendance, activity level, symptom severity)
- **Regular reassessment:** At least annually, document current function
- **Compare to own baseline:** Progress is measured against the child's own history, not peers
- **Identify trends:** Is the child improving, stable, or declining over months to years?

20.9.2 When to Escalate Treatment Intensity

Signs that warrant treatment escalation:

- Function declining despite current management
- Increasing severity of crashes
- New symptoms developing
- School attendance declining
- Current treatment not achieving expected improvement

Escalation options:

- Add medications not yet tried (OI medications, sleep medications)
- Increase accommodations (more reduced schedule, transition to homebound)
- Consult ME/CFS specialist if not already involved

- Consider aggressive interventions from Chapter 17 if appropriate

20.9.3 Reevaluation Frequency

- **Initial stabilization:** Frequent contact (every 2–4 weeks) until symptoms stabilize
- **Stable management:** Every 3–6 months
- **Transitions:** More frequent during school transitions (new school year, middle to high school)
- **Puberty:** Hormonal changes may affect symptoms; monitor during pubertal transition

20.9.4 Transition Planning

For older adolescents, plan transitions:

- **Transition to adult care:** Begin at age 16–17; identify adult ME/CFS providers
- **College planning:** If college-bound, consider accommodations, reduced course load, online options, schools with strong disability services
- **Workforce planning:** If not college-bound, vocational planning with disability considerations
- **Insurance continuity:** Plan for insurance coverage after aging off parents' plan
- **Self-management skills:** Adolescents should progressively take over managing their own pacing, medications, and appointments

20.10 Prognosis Communication

Children and families need accurate prognosis information—hopeful but realistic.

20.10.1 The Pediatric Advantage

Pediatric ME/CFS prognosis is substantially better than adult [78]:

- **Improvement/recovery rates:** 54–94% (versus $\leq 22\%$ in adults)
- **Recovery by 10 years:** 68%
- **Mean illness duration:** 5 years (range 1–15)
- **Function at follow-up:** Mean 8/10; only 5% remain very unwell

This is genuinely good news. Most children with ME/CFS improve significantly or recover.

20.10.2 Defining Recovery Appropriately

“Recovery” requires careful definition:

- **Full resolution:** Complete return to pre-illness function with no symptoms (less common)
- **Substantial improvement:** Major increase in function; minimal ongoing symptoms; can live normal or near-normal life (more common)
- **No longer meeting criteria:** Still has some symptoms but does not meet diagnostic threshold (common)

Any of these outcomes is success. A child does not need to achieve perfect health to have a good outcome.

20.10.3 Realistic Timelines

- **Months:** Symptom management may improve; functional improvement limited
- **Year 1–2:** Gradual improvement in many; some achieve significant gains
- **Years 2–5:** Continued improvement; many recover during this period
- **Years 5–10:** Most who will recover have done so; remaining patients may have chronic course

Improvement is slow and non-linear. Good weeks and bad weeks alternate. The trajectory is positive even if daily experience fluctuates.

20.10.4 Hope Maintenance During Difficult Periods

When progress is slow or absent:

- **Acknowledge difficulty:** It is hard to be ill; it is hard to wait for improvement
- **Reframe success:** Stability is success; not getting worse is success; managing symptoms well is success
- **Focus on present:** What can be enjoyed today, regardless of tomorrow?
- **Connect with others:** Other families facing ME/CFS understand; support groups help
- **Professional support:** Mental health support for adjustment to chronic illness

20.10.5 Contrast with Adult Prognosis

For families asking “what if this continues into adulthood?”:

- Adult prognosis is worse, but not hopeless (20–25% improve)
- Many adults with ME/CFS live meaningful lives with accommodations
- Adult treatments continue to improve
- The child’s trajectory is not determined by adult statistics

★ Key Point: The Goal of Early Management

The goal of appropriate early management is to maximize the likelihood of being in the 54–94% who improve or recover, rather than the small percentage who develop chronic severe disease. Every decision—about pacing, accommodations, avoiding GET, treating symptoms—should be evaluated against this goal. Early intervention is not about curing ME/CFS today; it is about preserving the window for recovery over the coming years.

20.11 Summary of Key Recommendations

1. **Diagnose accurately:** Use pediatric-modified criteria; distinguish from school avoidance by PEM pattern; evaluate for OI.
2. **Secure school accommodations:** Pursue 504 Plan or IEP with specific accommodations for cognitive dysfunction, OI, and PEM. Provide school with clear documentation.
3. **Teach and enforce pacing:** Help children identify their energy envelope and stay within it. Resist pressure to “push through” from any source.
4. **Treat OI aggressively:** Non-pharmacological measures first (hydration, salt, compression); add medications if needed. OI treatment often improves multiple symptoms.
5. **Optimize sleep:** Melatonin first-line; address sleep hygiene; consider school start time accommodations.
6. **Never recommend GET:** Graded exercise therapy is contraindicated and causes harm. If recommended by another provider, provide current evidence and decline.
7. **Maintain social development:** Active effort to preserve peer connections and identity development despite illness limitations.
8. **Monitor and adjust:** Regular reassessment; escalate treatment if declining; prepare for transitions.
9. **Communicate realistic hope:** Most children improve; recovery is common; timelines are long; success has many definitions.

→ Continued: Related Content

For children who are housebound or too ill for school attendance, see Chapter 19. For adult mild-moderate management, see Chapter 18. For detailed pathophysiology, see Chapter 10 (orthostatic mechanisms), Chapter 6 (metabolic dysfunction), and Chapter 7 (immune abnormalities).

21 Medications Targeting Underlying Mechanisms

21.1 Immune-Modulating Medications

21.1.1 Low-Dose Naltrexone (LDN)

Low-dose naltrexone (LDN) has emerged as one of the most commonly used off-label treatments for ME/CFS, despite limited controlled trial data.

Mechanism of Action

Naltrexone at standard doses (50 mg) blocks opioid receptors to treat addiction. At low doses (1–4.5 mg), the mechanism differs:

- **Transient opioid blockade:** Brief receptor occupancy may trigger compensatory endorphin upregulation [468]
- **Glial cell modulation:** LDN may antagonize Toll-like receptor 4 (TLR4) on microglia, reducing neuroinflammation [468]
- **Immune modulation:** Effects on T regulatory cells and cytokine balance reported; in vitro evidence of TRPM3 ion channel restoration in ME/CFS NK cells [536]
- **Endorphin rebound:** Overnight blockade may increase morning endorphin levels [468]

Dosing Protocols

Typical protocols involve:

- Starting dose: 0.5–1.5 mg at bedtime
- Gradual titration over weeks to months
- Target dose: 3–4.5 mg (individual optimization required)
- Compounding pharmacy often needed for low doses

Evidence in ME/CFS

Evidence remains preliminary:

- A retrospective study of ME/CFS patients reported symptomatic improvement in a majority of respondents [137]
- No large randomized controlled trials completed in ME/CFS

- A small randomized crossover trial in fibromyalgia showed significant pain reduction [469]; larger reviews support anti-inflammatory effects [468]
- Patient community reports generally favorable

Side Effects

Generally well-tolerated:

- Vivid dreams (common, usually transient)
- Sleep disturbance initially
- Headache
- Nausea (rare)

△ Warning 1: LDN Psychiatric Adverse Effects

While LDN is generally well-tolerated, **severe psychiatric reactions including depression and suicidal ideation** have been reported in a subset of ME/CFS patients. These reactions appear more common in individuals who exhibit paradoxical responses to other medications.

Risk factors for psychiatric adverse effects:

- History of paradoxical medication reactions
- Pre-existing mood vulnerability
- Concurrent use of other neuroactive medications

Monitoring protocol:

- Screen for mood changes during first 2–4 weeks of treatment
- Use PHQ-2 or similar brief screening at each dose adjustment
- Ensure caregiver/family awareness to observe for behavioral changes
- Have immediate discontinuation plan ready
- Discontinue immediately if depressive symptoms or suicidal ideation emerge

LDN's reputation as a "harmless" intervention may lead to inadequate monitoring. Patients and prescribers should maintain vigilance for mood changes, particularly in the "paradoxical reactor" phenotype (see Section 21.10.1).

Speculation 62 (LDN Combination Protocols). Patient community reports describe synergistic benefits from combining LDN with other interventions. One frequently mentioned combination involves LDN (at bedtime), NAD⁺ precursors (nicotinamide riboside or NMN, in the morning), and melatonin (at bedtime for circadian regulation). The theoretical rationale combines: (1) LDN's anti-neuroinflammatory effects, (2) NAD⁺'s role in mitochondrial energy production and cellular repair, and (3) melatonin's effects on sleep architecture, circadian rhythm, and its own anti-inflammatory properties. Individual case reports describe dramatic improvements, including return to work after prolonged disability. However, this represents **anecdotal evidence only**—no controlled trials have evaluated this specific combination, and publication bias strongly favors positive reports. The heterogeneous nature

of ME/CFS means that treatments helping some patients may be ineffective or harmful for others. Patients considering such combinations should work with knowledgeable physicians and implement changes sequentially to identify individual responses.

Energy Profile. Category B (energy-neutral). Naltrexone at micro-doses (1–4.5 mg) requires minimal hepatic CYP3A4 processing; metabolic demand is negligible. Unlike many neuroactive medications, LDN does not trigger immune activation or require energy-intensive cellular remodeling. Processing burden is negligible relative to any therapeutic benefit, making it safe to initiate even in severely energy-depleted patients.

⚠ **Limitation 1: Off-Label Medications: Borrowed Evidence from Other Conditions**

The majority of medications in this chapter are used off-label in ME/CFS, with evidence imported from other conditions. Key epistemic boundaries:

- LDN efficacy in ME/CFS rests on retrospective data, patient community reports, and a small fibromyalgia crossover trial—no large RCT has been completed in ME/CFS specifically.
- Antiviral protocols rest on limited controlled evidence: the Lerner 2007 valacyclovir trial [537] was placebo-controlled but small and conducted by a single group; the Montoya 2013 valganciclovir RCT (n=30) [149] showed significant improvement in mental fatigue, fatigue severity, and cognitive function in responders, but the overall effect was driven by a subgroup, and neither trial has been independently replicated.
- Mitochondrial supplements (CoQ10, L-carnitine, D-ribose, NADH) have individually small or single-study evidence bases in ME/CFS; combination protocols are extrapolated from biochemical reasoning rather than clinical testing.
- The “Energy Profile” classification (Categories A/B/C) assigned to each medication is a rational framework that has not been validated by measuring actual metabolic processing costs in ME/CFS patients.

21.1.2 Immunoglobulins (IVIG)

Energy Profile. Category C (energy-demanding). IVIG involves infusion of large protein aggregates requiring extensive hepatic processing, and the infusion itself triggers immune activation cascades that consume substantial ATP. The immune response to exogenous immunoglobulins—complement activation, cytokine release, lymphocyte mobilization—represents a significant metabolic burden. Severe patients may experience transient worsening resembling a Herxheimer-like reaction driven by immune activation energy demands. Consider stabilizing patients with Category A mitochondrial supplements before initiating.

21.1.3 Rituximab

Energy Profile. Category C (energy-demanding). B-cell depletion triggers major immune remodeling as the body reconstitutes its humoral immune repertoire, a process consuming substantial cellular energy over weeks to months. Initial infusions are particularly demanding due to acute cytokine release. Reserved for severe refractory cases; the energy cost is real and must be factored into risk-benefit assessment alongside other toxicities.

21.1.4 Other Immunomodulators

Additional immunomodulatory agents (corticosteroids, interferons, cyclophosphamide) have been explored in ME/CFS on a case-report or small-series basis, but lack sufficient evidence to warrant detailed treatment protocols. These are generally Category C (energy-demanding) due to broad immune activation or suppression and should only be considered within specialized research settings.

21.2 Antiviral Medications

Viral triggers and persistent viral reactivation have been implicated in ME/CFS pathogenesis. A systematic review and meta-analysis found significant associations with enteroviruses, HHV-7, parvovirus B19, and Borna disease virus [121]. EBV and HHV-6 are among the most commonly identified persistent viral agents across the literature [163], with enteroviruses implicated by both reviews. A subset of ME/CFS patients may benefit from antiviral therapy, though identifying responders remains challenging.

21.2.1 Valacyclovir and Acyclovir

Valacyclovir (Valtrex) and its active metabolite acyclovir target herpesviruses including EBV, HHV-6, varicella-zoster virus (VZV), and herpes simplex viruses (HSV-1, HSV-2).

Mechanism of Action

- **Nucleoside analog:** Acyclovir mimics guanosine, a building block of viral DNA
- **Viral DNA polymerase inhibition:** Incorporates into viral DNA, causing chain termination
- **Selective toxicity:** Preferentially activated by viral thymidine kinase, sparing host cells
- **Valacyclovir advantage:** L-valyl ester prodrug with 3–5× higher oral bioavailability than acyclovir

Evidence in ME/CFS

Evidence for herpesvirus-targeted antivirals in ME/CFS is preliminary but suggestive:

- **Lerner studies (2001–2013):** Multiple studies showed improvement in subset of ME/CFS patients with elevated EBV or HHV-6 antibody titers treated with long-term valacyclovir [538, 537, 539]
- **Subset response:** Approximately 30–40% of treated patients showed clinical benefit [539]
- **Duration requirement:** Benefits often required 3–6 months of continuous therapy [537]
- **Relapse upon discontinuation:** Some patients worsened when treatment stopped, suggesting suppressive rather than curative effect
- **Controlled evidence:** A 36-month placebo-controlled trial demonstrated sustained improvement in the valacyclovir-treated group [537]

Dosing Protocols

Valacyclovir.

- **Initial dose:** 500–1000 mg twice daily
- **High-dose protocol:** Up to 1000 mg three times daily in Lerner studies
- **Duration:** Minimum 3–6 months; some patients require indefinite suppressive therapy
- **Renal adjustment:** Reduce dose in renal impairment (creatinine clearance <50 mL/min)

Acyclovir (if valacyclovir unavailable or cost-prohibitive).

- **Dose:** 800 mg 3–5 times daily
- **Bioavailability disadvantage:** Requires more frequent dosing due to lower absorption
- **Cost:** Often less expensive than valacyclovir

Patient Selection

Consider antiviral trial in patients with:

- **Viral onset:** Clear infectious trigger (mononucleosis, severe flu-like illness)
- **Elevated antibody titers:** EBV VCA IgG >750, EBV EA (early antigen) IgG positive, HHV-6 IgG elevated
- **Persistent sore throat:** Chronic pharyngitis suggesting viral reactivation
- **Lymphadenopathy:** Tender lymph nodes
- **Immune subset dominance:** If viral/immune features predominate over other ME/CFS features

Limitations.

- Elevated EBV titers are common in healthy population (>90% seropositive)
- No clear titer threshold predicts response
- Some responders have “normal” titers
- Treatment is empirical

Side Effects and Monitoring

Common Side Effects.

- Headache (most common)
- Nausea
- Diarrhea
- Dizziness

Serious Adverse Events (rare).

- **Renal toxicity:** Acute kidney injury, particularly with high doses or dehydration
- **Thrombotic microangiopathy:** Rare; more common in immunocompromised patients
- **CNS effects:** Confusion, hallucinations (high doses, renal impairment)

Monitoring.

- **Baseline:** Creatinine, BUN, CBC
- **During treatment:** Creatinine every 3–6 months for long-term use
- **Hydration:** Maintain adequate fluid intake to prevent crystalluria

Energy Profile. Category B–C (dose-dependent). At standard ME/CFS doses (1–3 g daily), valacyclovir imposes moderate hepatic and renal processing burden; renal clearance requires adequate hydration and creates a mild but persistent metabolic demand. With renal impairment or at high doses, processing burden increases toward Category C. Long treatment durations (3–6 months) mean this low-grade metabolic cost is sustained; patients should ensure mitochondrial support is adequate before initiating prolonged courses.

21.2.2 Valganciclovir

Valganciclovir (Valcyte), a prodrug of ganciclovir, has broader antiviral coverage than valacyclovir, including better activity against HHV-6 and CMV.

Mechanism of Action

- **Guanosine analog:** Similar to acyclovir but with different selectivity
- **Broader herpesvirus coverage:** More potent against CMV and HHV-6 than valacyclovir
- **Viral DNA polymerase inhibition:** Blocks viral DNA synthesis

Montoya Stanford Study

The landmark study by Jose Montoya [149]:

- **Design:** Double-blind, placebo-controlled trial (EVOLVE study), 30 ME/CFS patients with elevated HHV-6 or EBV titers
- **Treatment:** Valganciclovir 900 mg twice daily for up to 6 months
- **Results:** Significant improvement in cognitive function (primary outcome) in responders; 7.4× increased likelihood of improvement vs. placebo
- **Response pattern:** Approximately 50–60% showed clinical benefit
- **Delayed improvement:** Benefits often appeared after 3–4 months
- **Durability:** Some patients maintained improvement after stopping; others required maintenance therapy

Dosing and Duration

- **Induction dose:** 900 mg twice daily for first 3–6 months
- **Maintenance dose:** 450–900 mg daily if prolonged therapy needed
- **Trial duration:** Minimum 3 months; Montoya protocol used up to 6 months
- **Renal adjustment:** Significant dose reduction required for creatinine clearance <60 mL/min

Risks and Benefits

Potential Benefits.

- Improved cognitive function (brain fog reduction)
- Increased energy in responders
- Reduction in flu-like symptoms
- Better quality of life scores

Significant Risks.

- **Bone marrow suppression:** Neutropenia, anemia, thrombocytopenia (BLACK BOX WARNING)
- **Renal toxicity:** Creatinine elevation, renal failure
- **Teratogenicity:** Contraindicated in pregnancy; requires contraception

- **Cost:** Extremely expensive (\$1000–3000/month without insurance)
- **GI side effects:** Nausea, diarrhea, abdominal pain

Contraindications.

- Absolute neutrophil count <500 cells/ μ L
- Platelet count <25,000/ μ L
- Pregnancy or breastfeeding
- Hypersensitivity to ganciclovir or valganciclovir

Required Monitoring.

- **Baseline:** CBC with differential, comprehensive metabolic panel, pregnancy test
- **Weekly for first month:** CBC to detect bone marrow suppression early
- **Every 2 weeks months 2–3:** CBC, creatinine
- **Monthly thereafter:** CBC, creatinine
- **Discontinuation criteria:** ANC <750, platelets <50,000, creatinine doubling

Clinical Decision-Making

Valganciclovir should be reserved for:

- Severe, refractory ME/CFS unresponsive to other interventions
- Strong viral component (elevated HHV-6 or CMV titers, viral onset)
- Failed trial of valacyclovir
- Patient willing to accept monitoring burden and risks
- Physician experienced in managing potential toxicities

The risk-benefit ratio requires careful consideration. Many experts consider valganciclovir a “last resort” option due to toxicity, reserving it for severe cases with clear viral markers.

Energy Profile. Category C (energy-demanding). Valganciclovir imposes significant hepatic and renal processing burden. Bone marrow suppression diverts cellular resources from ATP-producing pathways toward hematopoietic recovery, directly undermining energy production. The combination of hepatic CYP metabolism, high renal clearance demands, and immune system reorganization makes this among the most metabolically costly medications used in ME/CFS. Reserve for cases where viral phenotyping strongly predicts benefit.

21.2.3 Antiretroviral Approaches

Rationale

Some researchers have proposed antiretroviral drugs based on:

- Possible retroviral involvement in ME/CFS subset
- Reverse transcriptase activity detected in some patient samples
- Overlap between ME/CFS and post-treatment Lyme disease or other persistent infections
- Exploratory mechanistic hypotheses

Limited Evidence

- **Lack of reproducible retroviral findings:** Early reports of XMRV (xenotropic murine leukemia virus-related virus) were later shown to be laboratory contamination
- **No controlled trials:** Antiretroviral use in ME/CFS remains entirely anecdotal
- **Significant toxicity:** HIV antiretrovirals carry serious side effect profiles
- **Not recommended:** No expert consensus supports antiretroviral use outside research protocols

Research Directions

Future research might explore:

- **Endogenous retroviral activation:** Human endogenous retroviruses (HERVs) may be activated in ME/CFS
- **Reverse transcriptase inhibitors:** Tenofovir or other agents as research tools
- **Biomarker-guided trials:** Patient selection based on molecular evidence of retroviral activity

Currently, antiretroviral therapy for ME/CFS is **experimental only** and should not be attempted outside institutional review board-approved research protocols.

Energy Profile. Category C (energy-demanding). HIV antiretrovirals carry heavy hepatic CYP450 metabolism burden and significant toxicity profiles including mitochondrial toxicity (particularly NRTIs, which inhibit mitochondrial DNA polymerase- γ). The combination of hepatic processing demands, known mitochondrial toxicity, and serious adverse effect profiles argues strongly against empirical use. The energy cost provides additional reason beyond the lack of evidence to avoid these agents outside research protocols.

21.2.4 General Principles for Antiviral Use in ME/CFS

1. **Start with less toxic agents:** Trial valacyclovir before considering valganciclovir
2. **Allow adequate duration:** Minimum 3–6 months to assess response
3. **Monitor carefully:** Regular laboratory monitoring for toxicity
4. **Manage expectations:** Response rates vary widely across studies and patient selection criteria; many patients show no benefit
5. **Consider combination with other treatments:** Antivirals work best as part of comprehensive approach (pacing, autonomic support, etc.)
6. **Discontinue if no benefit:** If no improvement after 6 months, discontinue rather than continue indefinitely
7. **Assess maintenance need:** Some responders require long-term suppressive therapy; others can stop after initial course

21.3 Mitochondrial Support

Mitochondrial dysfunction is increasingly recognized as central to ME/CFS pathophysiology. Multiple supplements targeting mitochondrial function are widely used, though evidence quality varies. These interventions aim to support ATP production, reduce oxidative stress, and improve electron transport chain efficiency.

21.3.1 Coenzyme Q10 (CoQ10)

Coenzyme Q10 (ubiquinone) is an essential component of the electron transport chain, shuttling electrons between Complex I/II and Complex III. It also functions as a powerful antioxidant.

Mechanism of Action

- **Electron carrier:** Accepts electrons from Complex I (NADH dehydrogenase) and Complex II (succinate dehydrogenase), transfers to Complex III
- **Antioxidant:** Reduced form (ubiquinol) scavenges reactive oxygen species, protecting mitochondrial membranes
- **Membrane stabilization:** Integrates into mitochondrial inner membrane, maintaining structural integrity
- **Gene expression:** May modulate expression of genes involved in mitochondrial biogenesis

Ubiquinol vs. Ubiquinone

Two forms are commercially available:

Ubiquinone (oxidized form).

- Standard supplemental form
- Must be reduced to ubiquinol in the body for activity
- Less expensive
- Adequate for most individuals with normal reduction capacity

Ubiquinol (reduced form).

- Active, antioxidant form
- Does not require metabolic conversion
- 2–3× better bioavailability than ubiquinone
- Preferred for patients >40 years, those with impaired mitochondrial function
- More expensive

For ME/CFS patients with suspected mitochondrial impairment, ubiquinol may be preferable despite higher cost.

Evidence in ME/CFS

- **Small studies:** Some trials show modest improvement in fatigue and oxidative stress markers
- **Mechanistic rationale:** Strong theoretical basis given documented mitochondrial dysfunction
- **Fibromyalgia evidence:** Related condition shows benefit with CoQ10 (300 mg/day ubiquinol)
- **Safety profile:** Excellent; few side effects even at high doses
- **Limitations:** No large, well-controlled ME/CFS trials

Dosing and Bioavailability

Standard Dosing.

- **Ubiquinone:** 200–400 mg daily in divided doses
- **Ubiquinol:** 100–300 mg daily (lower dose due to better absorption)
- **Timing:** Take with fatty meals to enhance absorption (lipophilic compound)
- **Duration:** Minimum 8–12 weeks to assess benefit; may require 3–6 months

Bioavailability Enhancement.

- Take with fat-containing foods (avocado, nuts, olive oil)
- Soft gel formulations absorb better than powder capsules
- Divide total daily dose (e.g., 200 mg twice daily rather than 400 mg once)
- Consider ubiquinol form if poor response to ubiquinone

Side Effects

Generally very well-tolerated:

- Mild GI upset (nausea, diarrhea) in <5% of users
- Insomnia if taken late in day (some report increased energy)
- Rare: Rash, dizziness
- **Drug interactions:** May reduce warfarin effectiveness; monitor INR if anticoagulated

△ **Warning 2: Statin-Induced CoQ10 Depletion in ME/CFS**

Statins (HMG-CoA reductase inhibitors) deplete CoQ10 by blocking the mevalonate pathway, which is required for both cholesterol and CoQ10 synthesis. This has critical implications for ME/CFS patients:

ME/CFS-specific concern: ME/CFS patients have significantly lower baseline plasma CoQ10 levels than healthy controls, with 44.8% below the lowest control value [195]. Lower CoQ10 correlates with worse fatigue, autonomic symptoms, and cognitive dysfunction.

Clinical implications:

- Statins may worsen pre-existing CoQ10 deficiency in ME/CFS
- This could exacerbate fatigue, dysautonomia, and cognitive symptoms
- ME/CFS represents a **relative contraindication** for statin therapy unless CoQ10 is co-supplemented
- If statins are medically necessary (cardiovascular indications), mandatory CoQ10 supplementation (200–400 mg ubiquinol daily) should accompany therapy
- Monitor symptom changes closely when initiating statins in ME/CFS patients

Note on statin pleiotropic effects: Statins possess anti-inflammatory and immunomodulatory properties beyond lipid-lowering [540]. In autoimmune conditions, these effects can be beneficial [541]. However, in ME/CFS, the risk of worsening mitochondrial dysfunction through CoQ10 depletion likely outweighs potential anti-inflammatory benefits, particularly given that alternative anti-inflammatory approaches exist that do not deplete CoQ10.

Energy Profile. Category A (net energy provider). CoQ10 directly supports electron transport at the Complex I/III junction, providing substrate for ATP synthesis rather than consuming it. Supplementation replenishes a cofactor found to be deficient in ME/CFS patients, with

deficiency levels correlating with fatigue severity and autonomic symptoms [195]. One of the safest treatments for energy-depleted patients—it supplies what the system lacks. Ubiquinol form requires no metabolic conversion, making it especially suitable for severe patients with impaired reduction capacity.

21.3.2 NADH

Nicotinamide adenine dinucleotide (NADH) is the reduced form of NAD^+ , a critical coenzyme in cellular energy production.

Role in Energy Production

- **Electron donor:** NADH donates electrons to Complex I of electron transport chain
- **Glycolysis and TCA cycle:** Generated during glucose metabolism and Krebs cycle
- **ATP production:** Each NADH molecule can generate approximately 2.5 ATP molecules via oxidative phosphorylation
- **Lactate metabolism:** Required for lactate-to-pyruvate conversion (lactate dehydrogenase reaction)

Studies in ME/CFS

- **Forsyth et al. (1999) [542]:** Randomized, double-blind, placebo-controlled crossover trial in 26 ME/CFS patients; 10 mg NADH daily for 4 weeks showed 31% response rate vs. 8% placebo response (statistically significant)
- **Santaella et al. (2004) [543]:** Randomized trial (n=31) comparing NADH to conventional therapy over 24 months; significant improvement in first trimester ($p < 0.001$), but later comparable to active control
- **Mixed evidence:** Small sample sizes, variable formulations, heterogeneous patient populations; Forsyth study provides strongest evidence but limited replication
- **Subset response:** May benefit patients with documented NAD^+ metabolism abnormalities (per Heng 2025 findings) [12]

Dosing

- **Standard dose:** 10–20 mg daily on empty stomach (30–60 minutes before breakfast); clinical trial doses were 10 mg [542] and 20 mg [7]
- **Formulation:** Enteric-coated or sublingual to prevent gastric degradation
- **Complementary:** NAD^+ precursors (NR, NMN) serve a different role—see Chapter 22, Section 22.2.8 for the two-axis framework
- **Duration:** Trial for minimum 4–8 weeks

NADH vs. NAD⁺ Precursors

NADH and NAD⁺ precursors serve complementary rather than competing roles: NADH donates electrons directly at Complex I for immediate ATP production, while precursors replenish the broader cellular NAD⁺ pool for sirtuins, PARP, and metabolic signaling. For the complete two-axis framework and combination rationale, see Chapter 22, Section 22.2.8.

Nicotinamide Riboside (NR).

- Efficiently converts to NAD⁺ inside cells
- Dose: 300–1000 mg daily
- Better studied than NADH supplementation
- May improve mitochondrial biogenesis

Nicotinamide Mononucleotide (NMN).

- Direct NAD⁺ precursor
- Dose: 250–500 mg daily
- Emerging evidence for efficacy
- More expensive than NR

For ME/CFS mitochondrial support, the combination of NADH + CoQ10 has the strongest RCT evidence (Castro-Marrero 2021, n=207) [7], while NR or NMN can be added for NAD⁺ pool replenishment—a complementary mechanism rather than a replacement.

Energy Profile. Category A (net energy provider). NADH is a direct electron donor to Complex I of the electron transport chain; each molecule can generate approximately 2.5 ATP via oxidative phosphorylation. Minimal processing overhead is required—the molecule enters mitochondrial energy production directly without the enzymatic conversion steps needed by NAD⁺ precursors. Ideal for energy-depleted patients as a direct substrate rather than a precursor requiring further metabolic steps.

21.3.3 D-Ribose

D-ribose is a 5-carbon sugar that serves as the backbone of ATP, ADP, and AMP.

ATP Synthesis Support

- **Rate-limiting substrate:** Ribose availability can limit ATP regeneration after depletion
- **Purine salvage pathway:** Provides ribose-5-phosphate for adenine nucleotide synthesis
- **Bypass mechanism:** Supplements ribose directly, bypassing pentose phosphate pathway

- **Post-ischemic recovery:** Accelerates ATP regeneration after energy depletion (established in cardiac ischemia models)

Evidence in ME/CFS and Fibromyalgia

- **Teitelbaum et al. (2006) [8]:** Open-label pilot study (n=41) in fibromyalgia/ME/CFS patients; 5g D-ribose three times daily showed significant improvement across multiple domains: energy (+45%), sleep (+30%), mental clarity (+30%), pain intensity (-15%), and overall well-being (+30%)
- **Mechanism:** Post-exertional ATP depletion in ME/CFS may respond to ribose supplementation as ATP backbone precursor; accelerates purine salvage pathway
- **Anecdotal support:** Widely reported patient benefit; some notice improvement within 1-2 weeks
- **Lack of RCTs:** No placebo-controlled trials in ME/CFS; open-label design limits certainty despite impressive effect sizes

Dosing Protocols

- **Standard dose:** 5 grams (1 scoop) 2–3 times daily
- **Total daily dose:** 10–15 grams
- **Timing:** Spread throughout day; some take pre-activity
- **Form:** Powder dissolved in water or beverages (no capsule form practical due to high dose)
- **Loading phase:** Some protocols use higher initial doses for 1–2 weeks
- **Duration:** Effects may appear within 1–2 weeks; trial for 4–6 weeks minimum

Side Effects

- **Hypoglycemia:** Ribose can lower blood glucose; problematic for diabetics or those prone to hypoglycemia
- **GI symptoms:** Diarrhea, nausea if taken on empty stomach
- **Lightheadedness:** Take with food to minimize
- **Caution in diabetes:** Monitor blood glucose; may require insulin adjustment

Energy Profile. Category A (net energy provider). D-ribose provides the pentose sugar backbone for ATP regeneration via the purine salvage pathway, bypassing the energy-intensive de novo synthesis route. It converts to the ATP backbone with minimal metabolic processing overhead, making it one of the most direct energy substrates available. Particularly valuable post-exertion when adenine nucleotide pools are depleted and rapid regeneration is needed.

21.3.4 L-Carnitine and Acetyl-L-Carnitine

Carnitine is essential for transporting long-chain fatty acids into mitochondria for beta-oxidation.

Mechanism of Action

L-Carnitine.

- **Fatty acid shuttle:** Transports long-chain fatty acids across mitochondrial membrane via carnitine palmitoyltransferase (CPT) system
- **Energy substrate delivery:** Enables fatty acid oxidation for ATP production
- **Acetyl-CoA buffering:** Helps remove excess acetyl groups during metabolism

Acetyl-L-Carnitine (ALCAR).

- Acetylated form that crosses blood-brain barrier more readily
- Supports neuronal energy metabolism
- May enhance acetylcholine synthesis
- Neuroprotective and cognitive effects

Evidence in ME/CFS

- **Plioplys and Plioplys (1995) [544]:** Biomarker study (n=35) demonstrated significantly lower total carnitine, free carnitine, and acylcarnitine levels in CFS patients compared to controls; carnitine levels correlated with functional capacity
- **Plioplys and Plioplys (1997) [545]:** Treatment study with L-carnitine 3g/day for 8 weeks showed significant improvement in 12 of 18 clinical parameters; provides proof-of-concept for carnitine supplementation
- **Vermeulen and Scholte (2004) [546]:** Open-label randomized study (n=90, three groups) comparing acetyl-L-carnitine (2g/day), propionyl-L-carnitine (2g/day), and combination over 24 weeks; acetyl-L-carnitine showed 59% improvement in mental fatigue (p=0.015); propionyl-L-carnitine showed 63% improvement in general fatigue (p=0.004); combination therapy showed benefits in both domains
- **Malaguarnera et al. (2011) [547]:** While not ME/CFS-specific, double-blind RCT in hepatic encephalopathy demonstrated acetyl-L-carnitine's efficacy for reducing fatigue and improving cognitive function; supports mechanism of action
- **Mechanisms:** Addresses documented carnitine deficiency [544], improves fatty acid oxidation, supports mitochondrial function
- **Subset specificity:** May particularly help patients with acylcarnitine abnormalities on metabolomic testing; carnitine levels could serve as treatment-response biomarker

Dosing

L-Carnitine.

- **Dose:** 1000–3000 mg daily in divided doses
- **Form:** L-carnitine tartrate or L-carnitine fumarate (avoid D-carnitine)
- **Timing:** Between meals for optimal absorption

Acetyl-L-Carnitine.

- **Dose:** 2000 mg daily in divided doses (based on Vermeulen 2004 study showing efficacy at 2g/day for mental fatigue) [546]
- **Cognitive focus:** Preferred for brain fog and cognitive symptoms; 59% improvement rate in mental fatigue domain
- **Timing:** Morning and early afternoon (may cause alertness)

Propionyl-L-Carnitine.

- **Dose:** 2000 mg daily in divided doses (based on Vermeulen 2004 study showing efficacy for general fatigue) [546]
- **Physical fatigue focus:** Preferred for general fatigue and physical exhaustion; 63% improvement rate
- **Less commonly available:** May require compounding pharmacy or specialty suppliers

Combination. Some patients use both forms: L-carnitine for peripheral energy metabolism + ALCAR for cognitive support.

Side Effects

- **Body odor:** "Fishy" smell in some individuals (genetic variation in FMO3 enzyme)
- **GI upset:** Nausea, diarrhea at high doses
- **Insomnia:** If taken late in day
- **TMAO concerns:** Gut bacteria convert carnitine to TMAO (trimethylamine N-oxide), linked to cardiovascular risk in some studies; clinical significance in ME/CFS unclear

Energy Profile. Category A (net energy provider). Carnitine enables fatty acid oxidation, the body's most energy-dense metabolic pathway (yielding substantially more ATP per carbon than glucose). ALCAR additionally crosses the blood-brain barrier to support neuronal energy metabolism. Minimal processing overhead relative to the substantial energy yield from improved fatty acid transport into mitochondria. Particularly relevant where documented carnitine deficiency limits beta-oxidation capacity.

21.3.5 Alpha-Lipoic Acid

Alpha-lipoic acid (ALA) is a mitochondrial cofactor and powerful antioxidant.

Mechanism of Action

- **Cofactor for pyruvate dehydrogenase:** Essential for converting pyruvate to acetyl-CoA (entry into TCA cycle)
- **Cofactor for alpha-ketoglutarate dehydrogenase:** Critical TCA cycle enzyme
- **Antioxidant:** Scavenges multiple reactive oxygen species; regenerates other antioxidants (vitamins C, E, glutathione)
- **Metal chelation:** Binds toxic metals, potentially protective
- **Blood-brain barrier penetration:** Can protect neural mitochondria

Evidence

- **Diabetic neuropathy:** Well-established benefit in diabetic peripheral neuropathy (600–1800 mg/day)
- **ME/CFS rationale:** Theoretical benefit given mitochondrial dysfunction and oxidative stress
- **Limited ME/CFS trials:** No large controlled studies specific to ME/CFS
- **Small fiber neuropathy:** May help subset with documented SFN (common in ME/CFS)

Dosing

- **Standard dose:** 300–600 mg daily in divided doses
- **High-dose protocol:** Up to 1200–1800 mg/day used in diabetic neuropathy studies
- **R-lipoic acid vs. racemic:** R-form is the naturally occurring, bioactive enantiomer; may be more effective
- **Timing:** Take on empty stomach 30–60 minutes before meals for optimal absorption
- **Duration:** Minimum 8–12 weeks; neurological benefits may require months

Side Effects

- **Hypoglycemia:** Can lower blood glucose; caution in diabetics
- **Nausea:** Particularly at higher doses
- **Skin rash:** Rare
- **Biotin depletion:** High-dose ALA may compete with biotin; consider biotin supplementation (5–10 mg/day) with long-term high-dose ALA

Energy Profile. Category A–B (provider to neutral). ALA serves as an essential cofactor for pyruvate dehydrogenase and alpha-ketoglutarate dehydrogenase—rate-limiting TCA cycle enzymes that cannot function without it. Net energy effect is positive: ALA enables energy production rather than consuming it. It additionally regenerates antioxidants (vitamins C, E, glutathione), reducing the metabolic cost of chronic oxidative stress. Safe for energy-depleted patients.

21.3.6 Combination Mitochondrial Support Protocols

Many ME/CFS specialists recommend combining multiple mitochondrial supplements:

Basic Mitochondrial Stack

- CoQ10 (ubiquinol) 200–300 mg daily
- B-complex vitamins (B1, B2, B3, B5 for TCA cycle cofactors)
- Magnesium 400–600 mg daily (ATP-Mg complex, hundreds of enzymatic reactions)
- Vitamin D 2000–5000 IU daily (mitochondrial gene expression)

Enhanced Protocol

Add to basic stack:

- D-ribose 10–15 g daily (ATP regeneration)
- L-carnitine 1500–3000 mg daily (fatty acid transport)
- Alpha-lipoic acid 600–1200 mg daily (antioxidant, cofactor)
- NAD⁺ precursor (NR 300–1000 mg or NMN 250–500 mg)

Implementation Strategy

1. Start with basic stack for 4–6 weeks
2. Add one additional supplement at a time, spaced 2–4 weeks apart
3. Monitor response to each addition with symptom diary
4. Discontinue supplements showing no benefit after 8–12 weeks
5. Adjust doses based on tolerance and response

21.3.7 Limitations and Realistic Expectations

- **Modest benefits:** Mitochondrial supplements typically provide 10–30% improvement, not remission
- **Subset specificity:** May help those with documented mitochondrial dysfunction more than others
- **Cost burden:** Comprehensive protocols cost \$100–300/month

- **Evidence gaps:** Most lack large, high-quality RCTs in ME/CFS
- **Supportive, not curative:** Address downstream consequences, not root cause
- **Best as foundation:** Work optimally when combined with pacing, autonomic support, sleep optimization

Mitochondrial support represents a rational therapeutic approach given documented energy metabolism abnormalities, though individual responses vary widely.

21.4 Herbal Anti-Inflammatory Agents

21.4.1 Devil's Claw (*Harpagophytum procumbens*)

Devil's Claw is an herbal preparation derived from the secondary roots of *Harpagophytum procumbens*, native to southern Africa. The active constituent harpagoside demonstrates anti-inflammatory properties potentially relevant to ME/CFS-associated pain and inflammation.

Mechanism of Action

Devil's Claw exhibits a broader anti-inflammatory profile than NSAIDs [548]:

- **COX-1/2 inhibition:** Reduces prostaglandin synthesis, similar to NSAIDs
- **AP-1 pathway inhibition:** Blocks activator protein-1 mediated gene transcription—a mechanism distinct from conventional NSAIDs
- **Cytokine suppression:** Dose-dependently reduces TNF- α , IL-1 β , and IL-6 in macrophages
- **iNOS inhibition:** Reduces nitric oxide production and associated oxidative stress

Evidence Base

A systematic review of 12 randomized controlled trials (n=1,105) established the evidence base for chronic musculoskeletal pain [549]:

- **Strong evidence:** 50 mg harpagoside/day effective for acute exacerbations of chronic low back pain
- **Moderate evidence:** 60 mg harpagoside/day for osteoarthritis of spine, hip, and knee
- **Non-inferiority:** 60 mg harpagoside comparable to 12.5 mg rofecoxib (COX-2 inhibitor) for chronic low back pain
- **Dose dependence:** Products with ≥ 50 mg harpagoside daily show better outcomes than lower-dose preparations

The Cochrane Collaboration confirmed strong evidence for Devil's Claw in chronic low back pain [550].

Safety Profile

A systematic review of 28 clinical trials found a favorable safety profile [551]:

- Minor adverse events in approximately 3% of patients
- Primarily gastrointestinal (nausea, diarrhea, abdominal discomfort)
- Incidence not higher than placebo in double-blind studies
- Rare cases of allergic reactions reported

△ **Warning 3: Devil's Claw Contraindications and Interactions**

Contraindications:

- Peptic ulcer disease or active gastritis
- Gallstones (may increase bile production)
- Cardiovascular conditions (may affect heart rate)
- Pregnancy and lactation (insufficient safety data)

Drug interactions:

- **Anticoagulants/antiplatelets:** May enhance bleeding risk; avoid with warfarin, aspirin, clopidogrel
- **Antihypertensives:** May potentiate blood pressure lowering effects
- **Antidiabetics:** May affect blood glucose levels

Surgical consideration: Discontinue at least 2 weeks before elective surgery due to potential anticoagulant effects.

Relevance to ME/CFS

While no trials have specifically evaluated Devil's Claw in ME/CFS, several features suggest potential utility:

- **Anti-inflammatory mechanism:** IL-6 and TNF- α suppression may address neuroinflammation implicated in ME/CFS
- **Pain management:** Evidence in musculoskeletal pain may translate to ME/CFS-associated myalgia
- **Favorable side effect profile:** Better tolerated than NSAIDs with similar efficacy
- **Combination potential:** Broader mechanism of action than NSAIDs may complement other interventions

Practical Use

- **Dosing:** Standardized extract providing 50–100 mg harpagoside daily, divided into 2–3 doses
- **Duration:** 8–12 weeks needed to assess efficacy (onset slower than NSAIDs)

- **Product quality:** Ensure standardization to harpagoside content; variable quality in commercial preparations
- **Timing:** Take with food to minimize gastrointestinal effects

Energy Profile. Category B (energy-neutral). Herbal preparations with modest hepatic processing demands. Anti-inflammatory effects—particularly TNF- α , IL-1 β , and IL-6 suppression—may indirectly reduce the metabolic cost of chronic systemic inflammation. Overall energy impact is minimal, and the indirect benefit from reduced inflammatory burden may provide a small net positive effect in patients with prominent inflammatory features.

21.4.2 Palmitoylethanolamide (PEA)

Palmitoylethanolamide (PEA) is an endogenous fatty acid amide with anti-inflammatory, analgesic, and mast cell-stabilizing properties. It offers a well-tolerated option for chronic pain management in ME/CFS.

Mechanism of Action

PEA operates through multiple complementary pathways [552]:

- **PPAR- α activation:** Primary mechanism; inhibits NF- κ B and p38-MAPK signaling, reducing pro-inflammatory cytokine production at the source
- **Mast cell stabilization:** Reduces mast cell degranulation and release of histamine, NGF, TNF- α , and other inflammatory mediators—particularly relevant for ME/CFS patients with MCAS features
- **Cannabinoid system modulation:** Indirect effects on CB1/CB2 receptors; increases CB2 receptor expression on immune cells (“entourage effect”)
- **Glial cell modulation:** Reduces microglial and astrocyte activation, addressing neuroinflammatory contributions to pain and cognitive dysfunction
- **TRPV1 interaction:** Modulates vanilloid receptor signaling involved in pain transmission

Evidence Base

A systematic review and meta-analysis of 11 RCTs (n=774) established PEA’s efficacy for chronic pain [490]:

- **Significant pain reduction:** Standardized mean difference of 1.68 on 11-point scale (p < 0.00001)
- **Broad efficacy:** Effective across pain types—nociceptive, neuropathic, and nociplastic (central sensitization)
- **Consistent results:** 9 of 11 studies (82%) showed significant benefit

- **Excellent safety:** 6/11 studies reported no treatment-related adverse effects; when adverse effects occurred, they were mild and transient (primarily GI)

An earlier meta-analysis similarly confirmed efficacy with minimal adverse effects [489].

Relevance to ME/CFS

Several features make PEA particularly suitable for ME/CFS:

- **Nociplastic pain:** ME/CFS often involves central sensitization; PEA is effective for this pain type
- **MCAS comorbidity:** Mast cell stabilization addresses a common ME/CFS comorbidity
- **Neuroinflammation:** Microglial modulation may address cognitive symptoms (“brain fog”)
- **Safety profile:** Superior tolerability compared to NSAIDs; suitable for patients with multiple sensitivities
- **Combination potential:** Can be safely combined with other analgesics to enhance efficacy or allow dose reduction

Practical Use

- **Dosing:** 400–600 mg twice daily (800–1200 mg/day total); some protocols start at 300 mg BID
- **Formulation:** Micronized (mPEA) or ultramicronized (umPEA) forms preferred for enhanced bioavailability
- **Time to effect:** 6–8 weeks for maximum benefit; onset slower than conventional analgesics
- **Duration:** Can be used long-term; no tolerance or dependence observed
- **Administration:** Take with food; no significant drug interactions identified

Speculation 63 (PEA + LDN Combination). Both PEA and low-dose naltrexone (LDN) modulate neuroinflammation through distinct mechanisms—PEA via PPAR- α /mast cells and LDN via TLR4/microglia. Theoretically, combining these agents could provide synergistic anti-neuroinflammatory effects. Patient community reports describe such combinations, though no controlled trials have evaluated them. Given the excellent safety profiles of both compounds, empirical combination in patients with partial response to either alone may be reasonable under physician supervision.

Energy Profile. Category B (energy-neutral). PEA is an endogenous fatty acid amide processed through normal lipid metabolism pathways with minimal hepatic burden. Mast cell stabilization may reduce the ongoing energy cost of chronic mast cell degranulation and mediator release, a potentially significant indirect benefit in patients with MCAS overlap. Overall, a safe choice for energy-depleted patients with no meaningful metabolic processing demand.

21.5 Neuroprotective and Cognitive Enhancers

Cognitive dysfunction (“brain fog”) is among the most disabling symptoms in ME/CFS. This section reviews agents that may support cognitive function through neuroprotection, enhanced cerebral blood flow, or neurotransmitter modulation.

21.5.1 Ginkgo biloba (EGb 761)

Mechanism. Ginkgo biloba extract (standardized as EGb 761) contains flavonoid glycosides and terpene lactones with multiple neurologically relevant actions:

- **Cerebral blood flow enhancement:** Increases microvascular perfusion through vasodilatory and hemorheological effects
- **Antioxidant activity:** Scavenges free radicals and reduces lipid peroxidation in neural tissue
- **Platelet-activating factor (PAF) antagonism:** Terpene lactones (ginkgolides) inhibit PAF, reducing neuroinflammation
- **Mitochondrial protection:** May support mitochondrial function under oxidative stress
- **Neurotransmitter modulation:** Enhances cholinergic, dopaminergic, and noradrenergic transmission

Relevance to ME/CFS. ME/CFS involves documented cerebral hypoperfusion (reduced blood flow to brain), oxidative stress, and cognitive impairment. Ginkgo’s multi-target mechanism addresses several of these features:

- Cerebral blood flow enhancement may improve cognitive symptoms related to hypoperfusion
- PAF antagonism may reduce neuroinflammation, particularly relevant for MCAS subset
- Antioxidant effects support compromised cellular defenses

Evidence Base.

- **Cognitive impairment:** Meta-analyses demonstrate modest cognitive benefits in dementia and age-related cognitive decline [553, 554]
- **Cerebral insufficiency:** German Commission E approved for cerebral insufficiency with symptoms including difficulty concentrating, memory deficits, and fatigue [555]
- **ME/CFS-specific:** No randomized controlled trials in ME/CFS specifically; evidence extrapolated from related conditions
- **Fibromyalgia:** Small studies suggest potential benefit for cognitive symptoms, but evidence is limited

Dosing.

- **Standard dose:** 120–240 mg daily of standardized extract (EGb 761 or equivalent)
- **Typical products:** Cerebogan, Ginkobil, Tebonin (all standardized to 24% flavonoid glycosides, 6% terpene lactones)
- **Division:** Usually split into 2–3 doses daily (e.g., 80 mg three times daily)
- **Onset:** Effects may require 4–6 weeks of consistent use

Safety Considerations.

- **Bleeding risk:** Ginkgo has antiplatelet effects; caution with anticoagulants (warfarin), antiplatelet agents (aspirin, clopidogrel), and before surgery
- **Drug interactions:** May affect CYP enzyme metabolism; potential interactions with SSRIs (serotonin syndrome risk), anticonvulsants
- **Seizure threshold:** Theoretical concern for lowering seizure threshold (ginkgotoxin in impure preparations); use only standardized extracts
- **Gastrointestinal:** Mild GI upset, headache in some patients
- **Quality control:** Use standardized pharmaceutical-grade extracts; avoid unprocessed ginkgo seeds (toxic)

★ **Key Point: Ginkgo for ME/CFS Cognitive Symptoms**

Ginkgo biloba may provide modest support for cognitive symptoms in ME/CFS through enhanced cerebral blood flow and antioxidant effects. Use standardized extract (EGb 761 equivalent) 120–240 mg daily for minimum 6 weeks to assess response. **Caution:** Review bleeding risk and drug interactions before initiating. Not a substitute for pacing—cognitive energy still limited even with pharmaceutical support.

Energy Profile. Category B (energy-neutral). Standardized herbal extract with minimal metabolic processing demands. Improved cerebral blood flow may enhance brain energy substrate delivery, providing an indirect positive effect on cognitive energy availability. Caution: CYP enzyme interactions mean Ginkgo can alter the metabolism of co-administered medications, potentially shifting their energy classification; review the full medication list before initiating.

21.6 Interpreting Treatment Responses

Observation 85 (Extreme Heterogeneity in Medication Response). A striking feature of ME/CFS treatment is the extreme variability in individual responses to the same medication. Treatments that produce dramatic improvement in one patient may be ineffective or even harmful in another. This heterogeneity likely reflects the syndrome nature of ME/CFS—a common clinical presentation arising from diverse underlying pathophysiologies. Patient subgroups may include those with: (1) ongoing viral reactivation (who may respond to antivirals), (2)

autoimmune mechanisms (who may respond to immunomodulation), (3) MCAS/mast cell involvement (who may respond to antihistamines), (4) primary mitochondrial dysfunction (who may respond to metabolic support), or (5) combinations thereof. Until reliable biomarkers enable subgroup identification, treatment necessarily involves empirical trials with careful monitoring. This reality should temper both therapeutic nihilism (“nothing works”) and uncritical enthusiasm for any single treatment. The appropriate clinical stance is systematic, monitored experimentation guided by individual symptom patterns and physiological testing where available.

Observation 86 (Patient-Derived Treatment Sequence: The “Brain First” Protocol). Patient communities have developed an empirical treatment sequencing approach that prioritizes symptom domains in a specific order: (1) cognition/brain fog first, (2) fatigue second, (3) muscle weakness and pain third. The rationale is that cognitive restoration allows patients to better recognize their activity limits and manage pacing effectively, whereas fatigue improvement without cognitive restoration leads to dangerous overexertion. A frequently described sequence combines: low-dose aripiprazole or similar dopaminergic agents for cognitive symptoms (if metabolically tolerated), followed by low-dose naltrexone for sustained energy support, then pyridostigmine for autonomic/muscle symptoms. This represents community-derived knowledge rather than evidence-based protocol. Individual case reports describe dramatic functional improvement with this sequence, though others experience minimal benefit or adverse effects. The theoretical appeal lies in addressing the constraint (cognition) that limits patient’s ability to self-manage other symptoms. However, this protocol lacks controlled trial validation, and the optimal sequence likely varies by individual pathophysiology. Patients considering such sequencing should work with knowledgeable physicians, monitor carefully for adverse effects (particularly metabolic effects of dopaminergic agents), and recognize that individual responses may differ substantially from published case reports.

~ Hypothesis 1: Upstream-to-Downstream Treatment Sequencing

The “Brain First” sequence LDA → LDN → Mestinon may align with the neuroinflammatory cascade hypothesis in pathophysiology:

Proposed sequencing logic:

1. **Layer 1 - Dopaminergic restoration (LDA/aripiprazole):** Addresses documented catecholamine deficiency (particularly in NIH deep phenotyping studies). Dopamine is critical for: prefrontal cortex function (attention, executive planning), reward/motivation processing, and autonomic regulation. Restoring dopaminergic tone treats the upstream neurochemical deficit.
2. **Layer 2 - Microglial modulation (LDN):** Reduces microglial-mediated neuroinflammation through TLR4 signaling reduction. This targets the secondary neuroinflammatory cascade triggered by catecholamine deficiency—when dopamine drops, microglia become hyperactivated, perpetuating neuroinflammation even if baseline dopamine is restored. LDN addresses this consequence.
3. **Layer 3 - Autonomic ganglionic enhancement (Mestinon/pyridostigmine):** Addresses the downstream autonomic dysfunction resulting from upstream neurological dysfunction. Enhances acetylcholinergic transmission at autonomic ganglia,

improving heart rate variability and orthostatic tolerance. By this point, cognitive restoration (Layer 1) allows patients to recognize dysautonomic symptoms and apply appropriate pacing.

Cascade mechanism explanation: This upstream-to-downstream approach may be more effective than simultaneous multi-drug therapy because:

- Restoring dopamine (Layer 1) reduces the driving force for microglial activation, making Layer 2 (LDN) more effective
- Reducing neuroinflammation (Layer 2) may restore autonomic signaling, reducing need for maximum Layer 3 doses
- Sequential addition allows titration to individual tolerance before stacking additional neuroactive agents
- Cognitive restoration precedes fatigue improvement, preventing dangerous overexertion crashes

Critical caveats:

- This mechanistic framework is speculative and derived from hypothesis, not proven pathophysiology
- The cascade neuroinflammatory model itself remains under investigation (see Section 4.7.3 and pathophysiology chapters)
- Metabolic risks of dopaminergic agents (See Warning 21.11.1) may offset benefits in metabolically vulnerable patients
- Individual patients may require completely different sequences based on unique pathophysiological profiles
- The optimal sequence likely varies between rapid/acute responders (who benefit from simultaneous multi-agent) and slow-responders (who benefit from sequential layering)

The “Brain First” sequence represents an emerging hypothesis that cognitive improvement should precede fatigue improvement to allow safer self-management of remaining symptoms. Whether the proposed cascade mechanism actually explains superior outcomes remains uncertain.

Certainty: 0.25 (mechanistic reasoning from known pharmacology; no controlled trials testing this specific sequence; speculative cascade model)

? Open Question 1: Predicting Treatment Response

Can clinical features, biomarkers, or genetic profiles predict which ME/CFS patients will respond to specific treatments? If the syndrome comprises distinct pathophysiological subgroups, identifying these subgroups prior to treatment could dramatically improve therapeutic efficiency and reduce the burden of failed empirical trials. Potential stratification approaches include: immune profiling (B cell subsets, autoantibodies, NK function), metabolomic signatures, microbiome composition, autonomic phenotyping, or combinations thereof. Machine learning approaches applied to multi-omic datasets may eventually identify patterns invisible to traditional analysis.

21.6.1 Temporary vs. Durable Responses: A Critical Distinction

Observation 87 (Compensatory vs. Disease-Modifying Treatment). Treatment responses in ME/CFS may fall into two fundamentally different categories:

Compensatory (Symptomatic) Treatments:

- Address downstream consequences of the underlying pathology
- Provide relief while the treatment is maintained
- Relapse occurs when treatment is stopped or overwhelmed
- Analogous to “mopping the floor while the tap is running”
- Examples: amino acid supplementation (bypasses malabsorption), antihistamines (blocks histamine effects)

Disease-Modifying (Root Cause) Treatments:

- Address the underlying driver of the disease process
- May produce sustained remission even after treatment cessation
- Prevent or reduce vulnerability to relapse triggers
- Analogous to “turning off the tap”
- Examples: antiviral therapy (if viral reactivation is the driver), immunomodulation (if autoimmunity is the driver)

Observation 88 (Interpreting Temporary Improvement). A treatment that produces temporary but not durable improvement is *clinically significant*, not a failure:

1. **Proof of treatability:** The response demonstrates that the symptom complex is modifiable, not fixed
2. **Mechanistic clue:** The type of treatment that works suggests the pathway involved
3. **Foundation for optimization:** Compensatory treatments can stabilize patients while root cause is identified
4. **Relapse analysis:** What triggers relapse (infection, stress, treatment cessation) reveals what the compensatory treatment was masking

Example: A patient who improves dramatically on cimetidine + amino acids but relapses after an infection has demonstrated:

- The immune-metabolic pathway is involved (cimetidine response)
- Malabsorption or metabolic dysfunction is present (amino acid response)
- The underlying driver was not eliminated (relapse with immune challenge)
- Viral reactivation is a plausible root cause (infection-triggered relapse, cimetidine immunomodulation)

This pattern suggests a plausible next step: testing for viral reactivation and, if positive, adding antiviral therapy with the goal of converting compensatory treatment into disease-modifying treatment.

△ Warning 4: Avoid Premature Conclusion of Treatment Failure

A treatment that works temporarily should not be abandoned simply because relapse occurs. Instead:

- Document the response pattern (onset, magnitude, duration, relapse triggers)
- Analyze what the relapse reveals about the underlying driver
- Consider whether an additional intervention could make the response durable
- Maintain compensatory treatments while pursuing root cause identification

21.6.2 The Cimetidine-Antiviral Synergy Hypothesis

For patients with suspected viral-driven ME/CFS who show cimetidine response, a synergistic approach combining immunomodulation with direct antiviral therapy may convert temporary improvement into durable remission.

~ Hypothesis 2: Mechanistic Rationale for Cimetidine-Antiviral Combination**Cimetidine alone:**

- Blocks H2 receptors on suppressor T cells, enhancing cellular immunity [206]
- Increases NK cell activity and T cell cytotoxicity against viral targets
- Reduces viral-mediated immunosuppression
- **Limitation:** Does not directly reduce viral load; improvement depends on continuous enhanced immune pressure

Antivirals alone:

- Directly inhibit viral replication (valacyclovir inhibits HSV/EBV/VZV DNA polymerase)
- Reduce viral load during active replication phases
- **Limitation:** Less effective during latency; require functional immune response for complete suppression

Combination rationale:

- Cimetidine enhances immune clearance capacity
- Antiviral reduces viral load, making immune task easier
- Two-pronged attack: direct viral suppression + enhanced immune-mediated clearance
- May produce more complete viral suppression and more durable remission than either alone

Observation 89 (Historical Precedent and Pharmacokinetic Enhancement). Goldstein et al. [206] reported improvement in patients with chronic active EBV infection treated with cimetidine. More recent reviews of H2 receptor immunomodulation [556] confirm the mechanistic basis for enhanced cellular immunity. A recent pharmacokinetic study by Stuijt et al. [557] demonstrated that cimetidine significantly enhances systemic acyclovir concentrations through inhibition of renal clearance, providing a mechanistic rationale for the syner-

gistic potential of cimetidine-antiviral combinations. The logical extension—combining H2 blockade with direct antiviral therapy—represents a hypothesis-driven approach worthy of controlled evaluation.

Practical Protocol Considerations. For patients with:

1. Documented cimetidine response (energy improvement on H2 blockade)
2. Evidence of herpesvirus reactivation (elevated EBV EA-IgG, positive PCR, or HHV-6 elevation)

Consider:

- Cimetidine 200–400 mg BID (immunomodulation)
- Valacyclovir 1000 mg BID (direct antiviral) for minimum 3–6 months
- Regular monitoring: renal function, viral titers/PCR, clinical response
- Response evaluation at 3 and 6 months

This combination addresses both the immune dysfunction (cimetidine) and the viral driver (antiviral), potentially converting a compensatory response into disease modification.

21.7 Phenotype-Targeted Treatment Pathways

As understanding of ME/CFS heterogeneity advances, treatment pathways can be tailored to specific phenotype clusters. This section presents a hypothetical pathway for one emerging phenotype—the “Viral-Immune-Metabolic” cluster (see Section 5.6.12 and Section 4.7.3).

21.7.1 Treatment Pathway for Viral-Immune-Metabolic (“Cimetidine-Responder”) Phenotype

△ Warning 5: CRITICAL: Unvalidated Hypothetical Protocol

This protocol has NOT been validated in any controlled clinical trial.

- **Evidence level:** Clinical observation + mechanistic reasoning only
- **Expected responder rate:** Likely <10% even in carefully selected population
- **Status:** RESEARCH DISCUSSION ONLY—not for clinical implementation
- **Risk:** Inappropriate application to wrong patients may cause harm or delay effective treatment

DO NOT implement this protocol without:

1. Physician supervision and monitoring
2. Documented failure of evidence-based interventions
3. Informed consent regarding experimental nature

4. Recognition that most patients will NOT respond

The VIM phenotype concept itself is hypothetical and requires validation before clinical adoption.

Patient Selection Criteria

Consider this pathway for patients with:

- Post-infectious onset (especially documented EBV, HHV-6, or mononucleosis)
- POTS or dysautonomia confirmed
- MCAS or histamine intolerance (dietary triggers, antihistamine response)
- Response to amino acid supplementation (L-citrulline, NAC) noted
- OR dramatic improvement with cimetidine trial (rare but distinctive)

Phase 1: Confirmatory Trial (Weeks 1–4)

Goal: Determine if patient fits the cimetidine-responder pattern

1. Baseline assessment:

- Document current symptoms (validated scales: Bell Disability Scale, SF-36, CFQ)
- Order: EBV serology (VCA IgG, IgM, EBNA, EA-D), HHV-6 serology
- Order: Serum amino acid panel (if available)
- Record POTS status (NASA Lean Test or tilt table)

2. Cimetidine trial:

- Cimetidine 200 mg BID for 2 weeks
- If tolerated and some response: increase to 400 mg BID for 2 additional weeks
- Track: Energy (0–10 scale), hours out of bed, PEM episodes

3. Interpretation at Week 4:

- **Dramatic response** ($\geq 50\%$ improvement): Strong indicator of phenotype; proceed to Phase 2
- **Partial response** (20–50% improvement): Possible phenotype; proceed cautiously
- **No response** ($< 20\%$ improvement): Unlikely to be this phenotype; discontinue cimetidine, consider alternative approaches

Phase 2: Foundation Therapy (Weeks 4–12)

For patients with positive Phase 1 response:

Continue:

- Cimetidine 400 mg BID (or 200 mg BID if higher dose not tolerated)

Add sequentially (2-week intervals to identify individual responses):

1. Mast cell stabilization:

- Add H1 antihistamine (cetirizine 10 mg or fexofenadine 180 mg daily)
- Consider quercetin 500 mg BID (mast cell stabilizer)

2. Amino acid support:

- N-Acetylcysteine (NAC) 600 mg TID (glutathione precursor)
- L-citrulline-malate 3 g BID (NO synthesis + TCA cycle support)

3. Mitochondrial cofactors:

- D-ribose 5 g TID (ATP precursor)
- CoQ10 (ubiquinol) 200 mg daily
- B-complex with methylfolate and methylcobalamin

Phase 3: Optimization (Weeks 12–24)

Assess response at Week 12:

- Repeat symptom scales (Bell, SF-36)
- Reassess POTS status
- Consider repeat amino acid panel

If partial response, add as indicated:

- **Persistent viral symptoms:** Consider valacyclovir 1 g BID if EBV titers elevated (especially IgM or EA-D positive)
- **Persistent POTS:** Add ivabradine 2.5–5 mg BID or pyridostigmine 30 mg TID
- **Persistent pain/inflammation:** Increase PEA to 1200 mg/day (um-PEA form preferred)
- **Persistent cognitive symptoms:** Consider LDN 1.5–4.5 mg at bedtime

Phase 4: Diagnostic Confirmation (Months 3–6)

If significant improvement, pursue confirmatory testing:

- EBV/HHV-6 PCR (viral load) to assess suppression
- Repeat amino acid panel to assess normalization
- Consider intestinal permeability markers (Zonulin, LPS) if MCAS component prominent
- Consider flow-mediated dilation if NO dysfunction hypothesis being evaluated

Maintenance Protocol

For sustained responders:

- Continue H1 + H2 dual blockade indefinitely (mast cell management)
- Continue amino acid supplementation at maintenance doses
- Periodic reassessment (every 3–6 months)
- Attempt gradual dose reduction after 12 months of stability
- Monitor for relapse; resume full protocol if symptoms return

Expected Response Pattern

Based on mechanistic reasoning and limited case reports:

- **Timeline:** Initial cimetidine response may occur within days to 2 weeks; full amino acid/metabolic response typically requires 4–12 weeks
- **Response rate:** Unknown; likely <10% of ME/CFS population (rare phenotype)
- **Degree of improvement:** Dramatic responders may see 50–80% improvement; partial responders 20–40%
- **Durability:** Unknown; may require ongoing treatment to maintain benefit

~ Hypothesis 3: Mechanism of Response

The proposed mechanism integrates two parallel pathways:

Viral-immune pathway: Cimetidine blocks H2 receptors on suppressor T cells, enhancing cellular immunity against persistent herpesviruses (EBV, HHV-6). This allows improved viral control without requiring direct antivirals.

Metabolic pathway: MCAS/HIT causes intestinal barrier dysfunction and amino acid malabsorption. Exogenous amino acid supplementation (citrulline, NAC) bypasses the absorption deficit, restoring NO synthesis, glutathione levels, and TCA cycle function.

The synergy explains why patients may respond to the combination (cimetidine + amino acids) more than to either alone.

△ Warning 6: Cimetidine Drug Interactions

Cimetidine is a CYP450 inhibitor (particularly CYP1A2, CYP2D6, CYP3A4). It may increase levels of medications metabolized by these enzymes, including:

- Theophylline, warfarin, phenytoin
- Some benzodiazepines and SSRIs
- Beta-blockers (propranolol)

Review drug interactions before initiating cimetidine. In some cases, famotidine (which lacks significant CYP inhibition) may be substituted, though it also lacks cimetidine's immunomodulatory effects.

21.7.2 Other Emerging Phenotype-Targeted Pathways

As biological phenotyping advances (see Section 4.7.3), additional treatment pathways may be developed for:

- **Autoimmune-predominant phenotype:** Immunoabsorption, daratumumab, BC007 (for GPCR autoantibody-positive patients)
- **Mitochondrial-predominant phenotype:** Aggressive NAD⁺ precursor therapy, potentially rapamycin (mTOR modulation)
- **Neuroinflammatory-predominant phenotype:** LDN, IVIG (if SFN documented), environmental modification
- **Dysautonomia-predominant phenotype:** Comprehensive POTS protocol (volume expansion, compression, pharmacotherapy)

The key principle is matching treatment intensity and target to the patient's biological profile, rather than applying the same protocol to all ME/CFS patients.

21.8 Autonomic Medications

21.8.1 Pyridostigmine (Mestinon)

Pyridostigmine, an acetylcholinesterase inhibitor, has shown benefit for autonomic dysfunction in ME/CFS, particularly for orthostatic intolerance and POTS.

Mechanism of Action

Pyridostigmine inhibits acetylcholinesterase, prolonging acetylcholine activity at:

- **Autonomic ganglia:** Enhances sympathetic and parasympathetic neurotransmission
- **Neuromuscular junction:** Increases muscle strength (though this is not the primary target in ME/CFS)
- **Heart:** Vagal effects may improve heart rate variability

In POTS and autonomic dysfunction, pyridostigmine improves ganglionic transmission, enhancing the autonomic nervous system's ability to regulate cardiovascular function.

Evidence in POTS and ME/CFS

- **Raj et al. (2005) [558]:** Randomized crossover trial in POTS patients demonstrated reduced standing tachycardia without supine bradycardia
- **Mechanism:** Enhances ganglionic transmission in the autonomic nervous system
- **Clinical experience:** Widely used in ME/CFS clinics for autonomic symptoms

△ Warning 7: ME/CFS Dose Sensitivity

ME/CFS patients typically require 1/4 to 1/3 of standard pyridostigmine doses.

Standard POTS dosing: 30–60 mg three times daily (90–180 mg/day total)

ME/CFS-specific considerations:

- **Starting dose:** 15–20 mg once daily (not 60 mg)
- **Titration:** Increase by 15–20 mg increments every 1–2 weeks
- **Maximum tolerated:** Many ME/CFS patients stabilize at 20–30 mg 1–3× daily
- **Standard dose intolerance:** 60 mg may cause severe symptoms requiring bed rest

Rationale: ME/CFS patients often exhibit heightened sensitivity to neuroactive medications. The autonomic nervous system may be hyperreactive, such that standard doses produce excessive cholinergic effects. Start low and titrate slowly.

Side effects at excessive doses:

- Severe fatigue and weakness (paradoxical)
- Gastrointestinal distress (cramping, diarrhea)
- Excessive salivation
- Muscle fasciculations
- Bradycardia

If gastrointestinal symptoms occur, reduce dose rather than discontinuing.

Dosing Protocol for ME/CFS

1. **Week 1–2:** 15–20 mg once daily with food (morning or noon)
2. **Week 3–4:** If tolerated, add second dose (20 mg BID)
3. **Week 5–6:** If needed and tolerated, add third dose (20 mg TID)
4. **Maximum:** Most ME/CFS patients do not exceed 60 mg total daily

Timing: Take with meals to reduce GI side effects. Allow 4–6 hours between doses.

Duration of effect: Each dose lasts approximately 3–4 hours; extended-release formulation (Mestinon Timespan 180 mg) is available but rarely appropriate for ME/CFS patients due to dose sensitivity.

Compounding Pharmacy Approach for MCAS-Overlap Patients

ME/CFS patients with concurrent mast cell activation syndrome (MCAS) may fail to tolerate commercial pyridostigmine (Mestinon) tablets not because of the active ingredient itself, but because of excipient-triggered reactions. Commercial Mestinon 60 mg tablets contain lactose monohydrate, colloidal silicon dioxide, and stearic acid as excipients (per manufacturer labeling); generic pyridostigmine formulations may additionally contain colorants (to distinguish dosage strengths) and other binders that vary by manufacturer. Lactose and synthetic colorants are among the most commonly present excipients with documented potential for adverse reactions in sensitized patients [559], and MCAS patients are disproportionately

susceptible to such excipient-mediated reactions due to their lowered threshold for mast cell degranulation [560].

Observation 90 (Compounded Pyridostigmine as Diagnostic and Therapeutic Strategy in MCAS). In MCAS-overlap patients who cannot tolerate commercial Mestinon, compounding pharmacies (*préparation magistrale* in francophone systems) can prepare pure pyridostigmine bromide in hypoallergenic capsules with minimal excipients. This approach serves two functions:

1. **Therapeutic access:** Allows MCAS patients to receive pyridostigmine’s autonomic benefits when commercial formulations are not tolerated
2. **Diagnostic differentiation:** Distinguishes true pyridostigmine intolerance (pharmacological adverse effect) from excipient-mediated reactions (hypersensitivity or pseudoallergic reactions triggered by inactive ingredients). If a patient tolerates compounded pyridostigmine but not commercial Mestinon, the reaction is excipient-driven, not drug-driven—and the drug class should not be abandoned

Suggested protocol for MCAS-overlap patients:

1. Begin with compounded pure pyridostigmine bromide in hypoallergenic capsules (specify: no colorants, no lactose, no parabens)
2. Follow standard ME/CFS titration schedule (Section 21.8.1): start 15–20 mg once daily, titrate over weeks
3. Once target dose is reached and stable (minimum 4–8 weeks), optionally attempt transition to commercial Mestinon at a reduced dose (e.g., half the stable compounded dose), as both excipient exposure and dose-related cholinergic effects may contribute to any reaction
4. If commercial formulation triggers symptoms (GI distress, flushing, urticaria, fatigue flare) that were absent with the compounded form, return to compounded formulation indefinitely

Some patients report successful transition to commercial tablets after prolonged periods (months) on the compounded form. Whether this reflects gradual immunological desensitization, stabilization of mast cell reactivity through concurrent MCAS treatment, or simply natural disease fluctuation is unknown. Other patients require compounded formulations permanently. Both outcomes are clinically acceptable.

Cost and access considerations: Compounded medications are generally more expensive than commercial generics and may not be covered by insurance in all jurisdictions. However, for MCAS patients who would otherwise have no access to autonomic pharmacotherapy, the cost differential is clinically justified. Prescribers should specify “pure API, minimal excipients, no colorants, no lactose” on compounding orders.

Energy Profile. Category B (energy-neutral). Pyridostigmine works by preventing enzymatic breakdown of acetylcholine rather than requiring active metabolic synthesis, an energy-efficient mechanism. Low hepatic processing burden at ME/CFS doses (15–60 mg total daily).

Does not consume significant ATP, and any improvement in autonomic function may indirectly reduce the energy cost of dysautonomic compensation. Safe from a metabolic energy perspective.

21.9 H2 Receptor Antagonists

H2 receptor antagonists (H2 blockers) were developed for gastric acid suppression but have immunomodulatory properties relevant to ME/CFS. Cimetidine in particular has been studied for viral infections (see Section 21.6.2).

21.9.1 Cimetidine vs. Famotidine: Critical Differences

While both are H2 blockers, cimetidine and famotidine have important differences for ME/CFS patients:

Property	Cimetidine	Famotidine
Immunomodulation	Yes (T-cell enhancement)	Minimal
CYP450 inhibition	Strong (1A2, 2D6, 3A4)	Minimal
CNS penetration	Moderate	Lower
Psychiatric effects	Rare	Reported (see below)

Clinical implication: Famotidine cannot be substituted for cimetidine when immunomodulation is the therapeutic goal. However, for pure acid suppression in patients requiring CYP450-metabolized medications, famotidine is preferred.

△ Warning 8: H2 Blocker Psychiatric Adverse Effects

H2 receptor antagonists can cause psychiatric adverse effects, including **depression and suicidal ideation**. While these are rare, they appear more frequent in patients with the “paradoxical reactor” phenotype (see Section 21.10.1).

Famotidine-specific risk: Despite lower CNS penetration than cimetidine, famotidine has been associated with severe psychiatric reactions in susceptible individuals. Notably, some patients tolerate cimetidine but not famotidine, suggesting drug-specific rather than class-wide effects.

Risk factors:

- History of paradoxical medication reactions
- Pre-existing mood disorders
- Concurrent use of other CNS-active medications
- ME/CFS with prominent neurological features

Monitoring:

- Screen for mood changes during first 2–4 weeks

- Ensure caregiver/family awareness for early detection
- Discontinue immediately if depressive symptoms or suicidal ideation emerge
- If famotidine causes psychiatric effects, do not assume cimetidine will also—trial may be warranted

△ **Warning 9: Aspirin Contraindication in Histamine Intolerance**

Aspirin is contraindicated in patients with histamine intolerance (HIT).

Aspirin inhibits platelet cyclo-oxygenase, which reduces platelet-mediated histamine inactivation. This mechanism causes aspirin to trigger histamine release and block histamine metabolism, significantly worsening symptoms in patients with HIT or MCAS.

For ME/CFS patients with confirmed HIT or MCAS-overlap phenotype:

- **Avoid aspirin** entirely (including low-dose “cardioprotective” regimens)
- **Avoid other NSAIDs:** They share similar histamine-liberating effects
- **Use alternatives for pain management:** Acetaminophen, PEA (palmitylethanolamide), topical analgesics
- **Communicate with prescribers:** Clearly document HIT status to prevent inadvertent aspirin prescription

This contraindication applies regardless of cardiovascular indication, as the histamine burden outweighs cardioprotective benefit.

21.9.2 Cimetidine-LDN Synergy Protocol for Viral-Immune-Phenotype ME/CFS

For patients with evidence of viral-immune phenotype (elevated EBV titers, history of viral trigger, strong response to cimetidine alone), combining cimetidine with low-dose naltrexone may address both viral-immune and neuroinflammatory pathways.

~ **Hypothesis 4: Cimetidine-LDN Mechanistic Rationale**

Cimetidine immunomodulation:

- Blocks H2 receptors on suppressor T cells, enhancing cellular immune function
- Increases NK cell activity and T cell cytotoxicity against EBV and HHV-6
- Reduces viral-mediated immune suppression
- Direct mechanism: H2 receptor antagonism → enhanced Th1/Tc1 response against intracellular pathogens

LDN neuroinflammation reduction:

- Modulates TLR4 signaling on microglia, reducing neuroinflammatory cytokines (IL-6, TNF- α)
- May reduce microglial activation secondary to viral-driven immune activation
- Addresses downstream neurological consequences while cimetidine addresses upstream viral driver

Synergistic rationale: The combination targets two complementary mechanisms:

1. **Viral control:** Cimetidine enhances immune clearance capacity against persistent herpesviruses
2. **Neuroinflammation reduction:** LDN modulates microglial response, reducing secondary neurological damage
3. **Dual targeting:** Two-pronged approach may produce more complete viral suppression and superior symptomatic improvement than either agent alone

This mechanism may explain why some patients report dramatic response to cimetidine alone but plateau at partial improvement, while combination with LDN may extend recovery further. (This combination rationale is inferred from individual case reports, not controlled trial data.)

► Protocol 1: Cimetidine-LDN Synergy Protocol for Viral-Immune Phenotype

Patient Selection:

This protocol is appropriate for patients demonstrating:

- Clear post-viral onset (documented EBV infection, mononucleosis, or severe flu-like illness at disease onset)
- Elevated EBV serology (VCA IgG >750 mIU/mL, EA-D present, or positive PCR for EBV/HHV-6)
- Dramatic response to cimetidine trial (≥50% improvement in energy and function)
- No contraindication to LDN (see Warning 21.1.1)

Phase 1: Establish Cimetidine Baseline (Weeks 1–4)

- **Cimetidine dose:** 200 mg twice daily (400 mg total daily)
- **Assessment at Week 4:** Document improvement in energy, symptom severity, hours out of bed
- **Continuation criterion:** If energy improved ≥25%, proceed to Phase 2
- **Discontinuation criterion:** If minimal response (<10% improvement), this phenotype unlikely; discontinue and pursue alternative pathway

Phase 2: Add LDN with Mood Monitoring (Weeks 5–12)

- **LDN initiation:** Start 0.5 mg at bedtime (compounded low-dose form required)
- **Titration:** Increase by 0.5 mg every 1–2 weeks toward target 3 mg at bedtime
- **Psychiatric monitoring:** MANDATORY—LDN carries psychiatric adverse effect risk in subset of patients
 - Daily mood assessment first 2 weeks
 - PHQ-2 screening at each dose adjustment
 - Caregiver/family observation for behavioral changes
 - Immediate discontinuation if depression or suicidal ideation emerges
- **Continue cimetidine:** Maintain 200 mg BID throughout LDN titration

Phase 3: Combination Assessment (Weeks 12–16)

At Week 12, evaluate the combination:

- **Response assessment:** Compare current function to Phase 1 baseline (Week 4)
- **Expected improvement pattern:**
 - Energy/fatigue domain: Substantial improvement on cimetidine alone in responders; potential for further gains with LDN addition (magnitude varies; no controlled comparison data)
 - Cognitive function: May show additional improvement (LDN-mediated microglial modulation)
 - Pain/inflammation: May improve as neuroinflammation decreases
- **Non-response to combination:** If combined therapy provides <10% additional benefit over cimetidine alone, consider discontinuing LDN; continue cimetidine alone
- **Psychiatric adverse effects:** If mood changes emerged, discontinue LDN regardless of energy benefit

Phase 4: Viral Monitoring (ongoing)

If combination therapy shows improvement, assess viral response:

- **EBV serology:** Repeat VCA IgG, EA-D at 12 weeks; assess for titers declining toward normal range
- **EBV PCR:** If available, quantitative PCR to assess viral load suppression
- **Interpretation patterns:**
 - *EBV titers decline + symptoms improve:* Viral control achieved; continue combination indefinitely
 - *EBV titers decline without symptom improvement:* Viral suppression necessary but not sufficient; add other interventions
 - *Symptoms improve without titers declining:* May reflect improved immune tolerance rather than viral clearance; monitor for relapse

Maintenance Protocol (months 3+):

For sustained responders:

- Continue cimetidine 200–400 mg daily (dose adjusted to symptom stability)
- Continue LDN 3 mg at bedtime
- Reassess quarterly: symptoms, EBV titers, mood screening
- Plan gradual dose reduction after 12–18 months of stability if viral titers have normalized

Observation 91 (Cimetidine-LDN Synergy: Clinical Rationale and Precedent). This combination therapy emerges from convergent observations:

Cimetidine evidence: Goldstein et al. [206] documented improvement in chronic active EBV infection with H2 blockade. Recent pharmacokinetic data demonstrates cimetidine significantly enhances systemic acyclovir concentrations through renal clearance inhibition [557], providing mechanistic rationale for H2-antiviral synergy.

LDN evidence: While LDN is used broadly in ME/CFS, the mechanistic link to viral-immune

phenotypes is underexplored. However, the role of microglial activation in post-viral neurological sequelae is well-established, making LDN's TLR4 modulation theoretically relevant to viral-triggered ME/CFS.

Patient-derived knowledge (anecdotal; not from controlled studies): Some ME/CFS community reports describe cimetidine as a “turning point” medication—often the first intervention producing substantial improvement. The informal observation that some cimetidine responders plateau at partial improvement despite continued use suggests that additional pathways (particularly neuroinflammation) may limit further recovery. LDN's complementary mechanism could address this limitation.

Clinical practice note: This combination has not been formally studied in controlled trials. The protocol represents hypothesis-driven integration of established mechanisms (H2-mediated immunomodulation + TLR4-modulated neuroinflammation reduction) with pragmatic clinical experience. Individual responses vary widely; some patients may respond excellently to this combination, while others show minimal additional benefit of LDN beyond cimetidine alone.

Energy Profile. Category B–C (context-dependent). The drug itself has moderate hepatic processing burden via CYP1A2 and CYP2D6. However, as a CYP450 inhibitor, cimetidine reduces the metabolic processing burden of co-administered drugs by slowing their hepatic clearance. Net energy effect depends on context: as monotherapy, Category B; when used alongside other CYP450-metabolized medications, cimetidine may act as an indirect energy-sparing agent by reducing the total hepatic metabolic load. This dual role should inform prescribing decisions in polypharmacy contexts.

21.10 Medication Sensitivity Phenotypes

21.10.1 The Paradoxical Reactor Phenotype

A clinically significant subset of ME/CFS patients exhibits **paradoxical reactions** to medications—responses opposite to the expected effect, or severe adverse reactions at therapeutic doses.

Definition and Clinical Features

Paradoxical reactions include:

- **Opposite effects:** Sedatives causing agitation; stimulants causing fatigue; anxiolytics causing anxiety
- **Extreme sensitivity:** Severe symptoms at standard or even low doses
- **Psychiatric reactions:** Depression, suicidal ideation, or psychotic symptoms from medications not typically associated with these effects

- **Unpredictable patterns:** Tolerating one medication in a class while reacting severely to another

Examples from clinical observation:

- Pyridostigmine 60 mg causing severe prostration (standard starting dose)
- Famotidine causing depression and suicidal ideation
- Low-dose corticosteroids causing hypermania or psychosis
- LDN causing severe depression (typically well-tolerated)
- Tolerating cimetidine but not famotidine (same drug class)

Proposed Mechanisms

The paradoxical reactor phenotype may involve:

1. **Altered receptor sensitivity:** Upregulated or downregulated receptors from chronic illness
2. **Metabolic differences:** Variant CYP450 activity (ultra-rapid or poor metabolizers)
3. **Blood-brain barrier dysfunction:** Increased CNS penetration of medications
4. **Autonomic dysregulation:** Exaggerated responses to neuroactive compounds
5. **Mast cell activation:** MCAS may predispose to medication sensitivity
6. **Neuroinflammation:** Altered CNS pharmacodynamics

Clinical Management

Directive 1 (Paradoxical Reactor Protocol). For patients identified as paradoxical reactors:

General principles:

1. **Start at micro-doses:** 1/4 to 1/10 of standard starting dose
2. **Titrate slowly:** Minimum 1–2 week intervals between dose increases
3. **Monitor closely:** Daily symptom tracking, especially mood
4. **Expect variability:** Response to one medication does not predict response to another
5. **Have discontinuation plan ready:** Know what symptoms require immediate cessation
6. **Prefer previously tolerated agents:** If patient tolerated a medication before, prefer it over untested alternatives

Mood monitoring protocol (for any neuroactive medication):

- Daily mood check for first 2 weeks
- PHQ-2 screening questions at each dose adjustment
- Family/caregiver observation for behavioral changes
- Immediate discontinuation if suicidal ideation emerges

Documentation: Maintain careful records of all reactions, including dose, timing, and symptoms. This history guides future prescribing.

Observation 92 (Drug Class Does Not Predict Individual Tolerance). Patients may tolerate one medication in a class while experiencing severe reactions to another in the same class. Cimetidine tolerance does not guarantee famotidine tolerance. A severe reaction to one SSRI does not preclude trial of another. Each medication must be evaluated individually in paradoxical reactors.

21.10.2 The Excipient Sensitivity Phenotype

A distinct and clinically underrecognized medication sensitivity pattern in ME/CFS involves reactions not to the active pharmaceutical ingredient (API) but to **excipients**—the inactive ingredients used in commercial drug formulations. This pattern is particularly prevalent in patients with concurrent MCAS, but may also occur in ME/CFS patients without formal MCAS diagnosis, though evidence for this broader susceptibility is limited to clinical observation.

Excipients as Triggers of Hypersensitivity and Pseudoallergic Reactions

Commercial pharmaceutical tablets and capsules contain numerous excipients serving as fillers, binders, disintegrants, lubricants, colorants, preservatives, and coatings. A systematic analysis of 42,052 oral medications found that 92.8% contain at least one inactive ingredient identified as a potential allergen in sensitized individuals [559]. The following excipient classes are of particular concern, drawing on clinical trigger profiles described in the MCAS literature [560]:

- **Colorants:** Tartrazine (FD&C Yellow #5), sunset yellow (FD&C Yellow #6), and other azo dyes are recognized clinical triggers of pseudoallergic reactions including urticaria and angioedema in MCAS patients [560]. The reactions occur through non-IgE-mediated pathways; proposed mechanisms include direct mast cell and basophil degranulation, but the precise pathway remains incompletely characterized and lacks a specific primary citation. Many generic medications use colorants to distinguish dosage strengths [559].
- **Preservatives:** Parabens (methylparaben, propylparaben), sodium benzoate, and sulfites are recognized clinical triggers of pseudoallergic and hypersensitivity reactions in MCAS patients [560]. In the general population, the most widely proposed mechanism for ingested sulfites involves SO₂ generation in the acidic gastric environment, with subsequent reflex bronchoconstriction and non-IgE mast cell activation; true IgE-mediated sulfite allergy is rare, and the dominant reaction phenotype is non-immunological. However, these mechanistic details derive from general sulfite-sensitivity literature; they lack a specific primary citation in the MCAS context and whether MCAS patients react through the same pathway has not been directly studied. Reker et al. document the prevalence of preservatives across commercial oral formulations [559].
- **Lactose:** Present in approximately 45% of oral solid dosage formulations as a filler [559]. In MCAS patients with gastrointestinal mucosal involvement, secondary lactase deficiency may theoretically develop as a consequence of intestinal inflammation—by

analogy with established secondary lactase deficiency in Crohn's disease and other inflammatory enteropathies—leading to osmotic and fermentative GI symptoms upon lactose ingestion. Both the presence of sufficient MCAS-mediated mucosal damage to cause secondary lactase deficiency and the clinical relevance of this pathway remain unestablished; direct evidence in MCAS is lacking. Whether lactose itself directly activates mast cells through an immune-mediated pathway has not been established either.

- **Polyethylene glycol (PEG):** Used as coating material and solubilizer. Both IgE-mediated and non-IgE-mediated hypersensitivity reactions to PEG and related polysorbates are increasingly recognized across the general population receiving PEG-containing pharmaceuticals—more common than previously appreciated [561]. MCAS patients, with their lowered degranulation threshold [560], may be disproportionately susceptible to non-IgE-mediated PEG reactions, since stimuli that activate mast cells rarely in non-MCAS individuals may more readily activate the lowered-threshold MCAS mast cells. IgE-mediated PEG hypersensitivity requires prior humoral sensitization (specific IgE production), a prerequisite that is independent of the mast cell degranulation threshold. However, once sensitization has occurred, whether a lowered degranulation threshold in MCAS patients amplifies the severity of IgE-mediated reactions has not been studied. Direct evidence for either reaction type in MCAS cohorts is lacking.
- **Carboxymethylcellulose and polysorbate 80:** Used in liquid formulations and some tablets. In murine models, dietary concentrations of these emulsifiers disrupted intestinal mucus layer integrity, altered gut microbiota composition, and promoted colitis and metabolic syndrome [562]. Whether pharmaceutical doses (which have not been compared pharmacokinetically to the dietary concentrations used in the murine model) produce analogous effects in humans with GI MCAS involvement is unknown and speculative; human data are absent.

Clinical Significance in ME/CFS

The excipient sensitivity phenotype has several important implications:

1. **Misattribution of drug intolerance:** When a patient reports intolerance to a medication, clinicians typically attribute the reaction to the active ingredient and abandon the drug class entirely. In excipient-sensitive patients, this means losing access to potentially beneficial therapies. A patient who “cannot tolerate pyridostigmine” may in fact be unable to tolerate the lactose, colorant, or preservative in the specific commercial formulation—not the pyridostigmine itself.
2. **Inconsistent reactions across formulations:** Patients may tolerate a brand-name medication but react to a generic (or vice versa), because different manufacturers use different excipient profiles [559]. This brand/generic differential has a clear pharmacological explanation: the active ingredient is identical, but differing excipient compositions may trigger reactions of different severity in sensitized individuals. Despite this explanation, the pattern is likely underappreciated in clinical practice; when a brand/generic differential is reported, the differing excipient profile should be investigated before attributing the reaction to the active ingredient.

3. **Cumulative excipient load:** Each medication adds its own excipient burden. A typical oral solid dosage form contains a median of 8 inactive ingredients (mean 8.8) [559]; a patient taking five medications is therefore exposed to approximately 40 excipient exposures daily (at the median per-formulation count); even with substantial overlap across formulations, such a patient plausibly encounters 15–25 distinct excipient types, many repeated across multiple formulations daily. MCAS is characterized by additive trigger summation: multiple sub-threshold stimuli can collectively exceed the activation threshold and produce symptomatic degranulation [563, 560]. Applied to pharmaceutical excipients, this means that no single formulation’s excipient burden need be sufficient alone—the combined daily excipient load from polypharmacy may collectively drive symptomatic activation. Direct evidence for this specific mechanism in excipient-driven MCAS is lacking; the inference extends from the well-established MCAS trigger summation concept. Whether cumulative excipient load poses a similar risk in excipient-sensitive patients without MCAS is unknown; the mechanisms underlying non-MCAS excipient sensitivity have not been characterized, and it is unclear whether trigger summation applies outside the MCAS context.
4. **Polypharmacy amplification:** ME/CFS patients, who often require multiple medications for different symptom domains, may be particularly vulnerable to cumulative excipient burden—through mast cell trigger summation in those with MCAS, or through mechanisms that remain poorly characterized in those without MCAS—though the magnitude of this vulnerability has not been directly quantified. The apparent “intolerance to everything” reported by some patients in clinical practice has not been systematically quantified, and its causes are likely heterogeneous; however, excipient overload is a biologically plausible contributor that is distinct from—and may be conflated with—true sensitivity to multiple active ingredients.

The excipient classes and mechanisms discussed above are framed primarily in terms of MCAS pathophysiology. In ME/CFS patients without MCAS, excipient sensitivity may involve different mechanisms—such as non-mast-cell immune activation, gut barrier dysfunction, or neurological sensitization—but these remain uncharacterized, and no systematic study of excipient sensitivity in non-MCAS ME/CFS has been conducted.

The Compounding Pharmacy Solution

Compounding pharmacies (*préparations magistrales* in francophone systems, *Rezepturarzneimittel* in German-speaking countries) can prepare medications using the pure API with minimal, patient-selected excipients. This approach is already commonly used for LDN micro-dosing in ME/CFS and for low-dose methylene blue (Chapter 24), but its application should be considered systematically for all medications in excipient-sensitive patients. Practical barriers include variable insurance coverage (compounded medications are often not reimbursed), limited availability of compounding pharmacies in some regions, and absence of standardized quality assurance comparable to commercial drug manufacturing.

Directive 2 (Systematic Excipient Management Protocol). For patients with documented or suspected excipient sensitivity (MCAS diagnosis, history of reacting differently to brand vs. generic, unexplained multi-drug intolerance):

Step 1—Excipient audit:

- For each current medication, obtain the complete excipient list (available in the Summary of Product Characteristics / package insert).
- Cross-reference excipients across all medications to identify common triggers.
- Compare excipients in tolerated vs. non-tolerated formulations of the same drug class.

Step 2—Targeted reformulation:

- For medications where excipient-mediated reaction is suspected, request compounded formulation with pure API.
- Specify: no colorants, no lactose, no parabens, no PEG, hypoallergenic capsule shell.
- Start at a low dose and titrate upward, even if prior reaction was suspected excipient-driven—reactions may have involved both excipient and dose components. *Exception:* if the prior reaction was potentially life-threatening—anaphylaxis, or any of its severe manifestations (airway angioedema, laryngeal edema, bronchospasm, hypotension)—do not attempt re-challenge without specialist supervision and full emergency preparedness, regardless of whether the anaphylaxis is suspected to be excipient- or API-mediated; a compounded formulation eliminates most excipients but still contains the API and minimal residual excipients (e.g., capsule shell material), and if the original reaction was API-mediated—or mediated by an excipient retained in the compounded form—the compounded form poses a comparable risk. In such cases the risk of severe anaphylaxis on re-exposure outweighs the benefit of confirming excipient causality outside a controlled clinical setting.

Step 3—Diagnostic confirmation:

- If the patient titrates to the previously intolerable dose on the compounded formulation without reproducing the original adverse reaction, excipient sensitivity is the probable explanation—provided that the clinician has documented and can reasonably exclude other confounds:
 - Spontaneous change in sensitivity over time (compare baseline disease activity before and during the compounded trial).
 - Concurrent medication changes (none should occur during the trial).
 - Dose-response effects independent of excipient change (once a stable dose is reached, avoid further dose changes during the observation period used to assess excipient tolerance; any dose adjustments made during titration should be noted separately).
 - Concurrent GI anti-inflammatory treatments that may have independently reduced mucosal sensitivity.
 - Placebo or expectation effects (particularly if the patient knows the formulation has changed).
- Document the specific excipients present in the non-tolerated formulation(s) but absent in the compounded form, to build a patient-specific trigger profile.
- Optionally, after prolonged stability on the compounded form (months), attempt cautious transition to a commercial formulation with the fewest identified trigger excipients.

Step 4—Ongoing management:

- Maintain a patient-specific excipient avoidance list in the medical record.
- Review excipient composition before any new prescription.
- Alert pharmacists to avoid substitution with different generic manufacturers without consulting the excipient profile.
- Consider compounding as first-line for any new medication trial in confirmed excipient-sensitive patients.

The hEDS/POTS/MCAS Triad, ME/CFS Overlap, and Medication Management

The frequent co-occurrence of hypermobile Ehlers-Danlos syndrome (hEDS), POTS, and MCAS—commonly termed the “triad”—is documented in the literature [150, 151], though no proven pathophysiologic mechanism linking all three conditions has been established [151]. ME/CFS overlaps substantially with this triad: POTS is present in a large proportion of ME/CFS patients (Chapter 10), and MCAS features are increasingly discussed in the context of ME/CFS [244, 564], though systematic prevalence data remain limited. Whether ME/CFS represents a fourth comorbidity, a consequence of the triad, or a partially overlapping syndrome remains unresolved. Regardless, the practical implications for medication management are immediate.

Observation 93 (The Triad Overlap Complicates Every Medication Decision). Patients with ME/CFS who also have MCAS and POTS (with or without hEDS) face compounded medication challenges:

1. **ME/CFS demands dose sensitivity:** Start at 1/4–1/10 standard doses, titrate slowly (Section 21.10.1).
2. **MCAS demands excipient awareness:** Every formulation must be evaluated for mast cell triggers (Section 21.10.2).
3. **POTS demands autonomic caution:** Medications affecting heart rate, blood pressure, or volume status require careful monitoring.
4. **Combined effect:** The intersection of all three requirements means that even introducing a single new medication requires consideration of dose (ME/CFS), formulation (MCAS), and hemodynamic impact (POTS) simultaneously.

This triple constraint explains why triad-overlap patients are often perceived as “difficult” or “impossible to treat.” The difficulty is real, and it is physiological—rooted in the biology of the conditions and the pharmacology of available drugs—not psychological. Systematic application of the excipient management protocol (Section 21.10.2), ME/CFS dose sensitivity guidelines (Section 21.10.1), and autonomic monitoring may enable successful treatment in this challenging population, though clinical validation of this integrated approach is lacking.

Observation 94 (Simultaneous Onset of ME/CFS, POTS, and MCAS Features: Clinical Patterns and Unproven Mechanisms). In clinical practice, patients sometimes report simultaneous or near-simultaneous onset of fatigue with post-exertional malaise (ME/CFS), orthostatic

intolerance (POTS), and mast cell-mediated symptoms such as flushing, urticaria, and multiple new food or chemical sensitivities (suggestive of MCAS) following a triggering event, typically infection. The hEDS/POTS/MCAS triad literature documents high rates of co-occurrence among these conditions [150], though whether simultaneous onset is typical has not been systematically studied. A candidate mechanism proposed in the hEDS/MCAS literature involves connective tissue laxity altering mast cell tissue distribution and vascular tone [565], but causal evidence for this specific mechanism remains absent, consistent with the broader finding that no proven pathophysiologic mechanism linking the three conditions has been established [151]. In post-infectious ME/CFS, the co-occurrence of POTS and MCAS features may additionally or alternatively reflect a distinct mechanism. One speculative candidate—proposed here without established literature support and offered as a candidate for future investigation—is infection-triggered autoimmune disruption of autonomic ganglia (e.g., via autoantibodies against ganglionic acetylcholine receptors or adrenergic receptors) concurrent with widespread mast cell destabilization. These mechanisms are not mutually exclusive; patients with pre-existing connective tissue laxity may be additionally susceptible to infection-triggered destabilization. No causal pathway for either mechanism has been established.

The clinical implication is actionable independently of mechanism: given the documented co-occurrence [150] and its clinical recognition in review literature [151], when one condition in this cluster is diagnosed, active screening for the others is clinically warranted, though formal screening guidelines have not been established.

Untreated MCAS may undermine ME/CFS pharmacotherapy through multiple mechanisms—both the excipient sensitivity pathways described in this section (excipient-mediated degranulation and cumulative trigger summation) and direct mast cell-mediated drug hypersensitivity reactions independent of excipients. No controlled study has directly measured this effect, and the inference rests on the biological plausibility of those mast cell activation pathways rather than on direct evidence.

Untreated POTS may amplify ME/CFS fatigue and cognitive dysfunction through orthostatic cerebral hypoperfusion, though direct measurement of hypoperfusion-mediated fatigue amplification specifically in ME/CFS patients with comorbid POTS is lacking. Relevant to the overlap context specifically: in Novak et al.'s cohort of mast cell disorder patients—all of whom presented with dysautonomia—a 20–24% reduction in orthostatic cerebral blood flow was documented [173], suggesting that in the MCAS–POTS–ME/CFS overlap population the hypoperfusion burden may be compounded, though this remains to be studied directly.

21.10.3 Implications for Subgroup-Specific Treatment

The medication sensitivity phenotypes described in this section—paradoxical reactors (Section 21.10.1), excipient-sensitive patients (Section 21.10.2), and the compounded medication challenges in triad-overlap patients (Section 21.10.2)—reinforce the fundamental heterogeneity of ME/CFS discussed in Chapter 5. Treatment protocols that work for one subgroup may be inapplicable or harmful for another.

Until biomarker-driven subtyping becomes available (Chapter 26), clinical management must proceed by careful phenotyping and systematic trial-and-error. The protocols described

above—micro-dosing, excipient auditing, compounding pharmacy utilization, and triad-aware prescribing—provide a structured framework for this necessarily empirical process. The goal is not to eliminate trial-and-error, but to make each trial maximally informative: when a medication fails, determining *why* it failed (wrong drug, wrong dose, wrong formulation, or wrong timing) prevents premature abandonment of potentially effective treatments.

21.11 Atypical Antipsychotics: Metabolic Considerations in ME/CFS

Atypical antipsychotics are sometimes used off-label in ME/CFS for neurological symptoms, autonomic dysfunction, or sleep disturbance. While evidence for primary psychiatric indication is lacking, some patients report benefit for specific symptom clusters, though published case series are limited. However, these medications carry significant metabolic risks, particularly relevant to ME/CFS patients.

21.11.1 Low-Dose Aripiprazole (LDA)

Aripiprazole, a partial dopamine agonist, is occasionally used at low doses for cognitive symptoms or dysautonomia in ME/CFS. Patient case reports describe cognitive improvement at doses of 1–2 mg daily, substantially lower than psychiatric indication doses (10–30 mg).

Metabolic Risk Warning

△ **Warning 10: Aripiprazole-Associated Prediabetes and Metabolic Syndrome Risk**

CRITICAL: Aripiprazole carries prediabetes risk even at low doses, with particular implications for ME/CFS patients.

While atypical antipsychotics are known to cause metabolic derangements (particularly olanzapine, quetiapine), aripiprazole was initially classified as having lower metabolic risk. However, emerging evidence and clinical observation suggest this may be false reassurance, particularly in the ME/CFS population.

Metabolic risks documented in aripiprazole use:

- **Hyperglycemia and diabetes:** Case reports and small studies document glucose dysregulation, with some patients developing prediabetes or frank diabetes even at low doses
- **Weight changes:** Paradoxically, weight loss can occur (unlike olanzapine/quetiapine), which may reflect uncontrolled metabolic dysfunction rather than benign effect
- **Lipid abnormalities:** Elevated triglycerides, reduced HDL cholesterol reported
- **Insulin resistance:** Direct effects on insulin signaling pathways at dopamine re-

ceptor level

ME/CFS-specific concern: ME/CFS patients already demonstrate:

- Metabolic dysfunction and impaired glucose tolerance
- Increased incidence of metabolic syndrome and POTS-associated dysautonomia
- Proposed bidirectional relationship between ME/CFS and metabolic syndrome: metabolic derangement may worsen ME/CFS symptoms (dysautonomia, fatigue), and ME/CFS may impair metabolic control

Adding aripiprazole (which impairs metabolic regulation) to ME/CFS patients with underlying metabolic vulnerability may trigger or accelerate transition from normal glucose tolerance to prediabetes to overt diabetes.

Clinical recommendation:

- If aripiprazole is considered, obtain baseline fasting glucose, HbA1c, and lipid panel
- Repeat metabolic testing every 3 months during treatment
- Educate patients on prediabetes symptoms and the bidirectional ME/CFS ↔ metabolic syndrome relationship
- At first sign of glucose elevation (fasting >100 mg/dL, HbA1c >5.7%), discontinue and switch to alternative agent
- Consider metabolic-neutral alternatives (bupropion, low-dose stimulants) for cognitive symptoms or dopaminergic dysfunction

The theoretical benefit for cognition must be weighed against real metabolic risk in a population already vulnerable to metabolic derangement.

Metabolic Protection During LDA Therapy

If low-dose aripiprazole is deemed beneficial despite metabolic risks, the following protocol preserves cognitive benefits while preventing metabolic amplification of disease:

► Protocol 2: LDA Metabolic Protection Protocol

Baseline Assessment (before LDA initiation):

- Fasting glucose (target: <100 mg/dL)
- Hemoglobin A1c (HbA1c; target: <5.7%)
- Fasting insulin level
- Lipid panel (triglycerides, HDL, total cholesterol)

Monitoring Schedule:

- **During titration phase:** Monthly fasting glucose measurement
- **Maintenance phase:** Quarterly HbA1c; fasting glucose every 6 weeks
- **Annual:** Full metabolic panel (lipids, comprehensive metabolic panel)

Intervention Thresholds:

- **HbA1c >5.7% or fasting glucose >100 mg/dL:** Initiate metabolic intervention
- **HbA1c >6.5% or fasting glucose >126 mg/dL:** Discontinue LDA and switch to metabolic-neutral alternative

First-Line Intervention:

- **Metformin 500 mg:** Start at 500 mg once daily with dinner, increase to 500 mg twice daily over 2 weeks
- **Benefit:** Direct insulin sensitization plus anti-inflammatory properties (particularly TLR4 pathway relevant to ME/CFS)
- **Monitoring:** Monitor gastrointestinal tolerance; diarrhea is most common side effect
- **Recheck metabolic markers:** 6 weeks after initiation

Alternative Intervention (if metformin intolerant):

- **Berberine 500 mg TID:** Natural alkaloid with similar mechanism to metformin (AMPK activation, insulin sensitization)
- **Comparable efficacy:** Some studies suggest equivalent glucose control to metformin
- **Advantage:** Often better tolerated; milder GI side effects

Lifestyle Intervention (concurrent):

- **Time-restricted eating:** 8-hour eating window (if tolerable within ME/CFS activity limitations)
- **Rationale:** Improves insulin sensitivity, reduces metabolic syndrome progression
- **ME/CFS adaptation:** Can be combined with appropriate pacing; may require careful meal timing to avoid post-prandial crashes

Escalation Protocol:

- **If progression despite metformin:** Consider GLP-1 agonist (semaglutide)
- **Rationale:** Additional weight regulation, greater glycemic control, cardiovascular benefit
- **Caution:** Nausea may be problematic in patients with MCAS or GI sensitivity; start at lowest dose

△ Warning 11: LDA Metabolic Amplification of Neuroinflammatory Cascade

LDA metabolic effects may create a therapeutic ceiling through metabolic amplification of neuroinflammation. The sequence hypothesized in the cascade neuroinflammatory model (see Section 21.6.2 and pathophysiology section on neuroinflammatory cascade) suggests that metabolic dysfunction feeds back into neuroimmune activation, limiting the cognitive benefits achievable with dopaminergic therapy alone.

Monitor and intervene early to prevent metabolic syndrome amplifying the neuroinflammatory cascade and offsetting the cognitive benefits of LDA therapy. This protective

approach may allow extended use of an otherwise effective intervention.

Alternative Approaches for Cognitive Dysfunction

Before considering aripiprazole or other antipsychotics for cognitive symptoms, evaluate:

- **Sleep optimization:** Undiagnosed sleep apnea, circadian dysregulation, or sleep architecture disruption commonly drive cognitive impairment; formal sleep study and treatment may resolve symptoms
- **Mitochondrial support:** CoQ10, NAD⁺ precursors, D-ribose specifically target energy-dependent cognitive processing
- **Cerebral blood flow enhancement:** Ginkgo biloba, low-dose vasodilators address documented hypoperfusion
- **Neuroinflammation reduction:** LDN, PEA, or antimicrobial protocols may improve cognition by reducing neuroimmune activation
- **Metabolic support:** Restoration of amino acid levels, glucose control, and mitochondrial efficiency often improve cognition without pharmacologic risk

Most ME/CFS patients show substantial cognitive improvement with foundational interventions before requiring psychotropic medications.

Energy Profile. Category C (energy-demanding). Aripiprazole is a substrate of CYP2D6 and CYP3A4 requiring significant hepatic metabolism. Additionally, metabolic syndrome risk—insulin resistance, glucose dysregulation, and lipid abnormalities—represents an ongoing disruption of energy metabolism at the cellular level: impaired insulin signaling reduces glucose uptake into mitochondria, directly limiting ATP production. Even at low doses (1–2 mg), the combination of hepatic processing demand and metabolic perturbation places this firmly in Category C. Use only when cognitive benefit clearly justifies the energy cost, and protect against metabolic amplification with concurrent monitoring and intervention.

22 Supplements and Nutraceuticals

“First, do no harm. Second, be honest about what we don’t know.”

This chapter reviews supplements and nutraceuticals commonly used in ME/CFS. We emphasize accessible, over-the-counter options that patients can try without prescription, while being honest about the limited evidence base. Most supplements have not been rigorously tested in ME/CFS-specific trials; recommendations are based on mechanism, related-condition evidence, and patient-reported outcomes.

Observation 95 (Evidence Limitations). Very few supplements have been tested in randomized controlled trials specifically for ME/CFS. Most evidence comes from: (1) mechanistic plausibility based on documented ME/CFS abnormalities, (2) trials in related conditions (fibromyalgia, Long COVID, chronic fatigue), (3) small open-label studies, and (4) patient surveys. We note the evidence level for each supplement. “Widely used” does not mean “proven effective.”

⚠ Limitation 1: Supplement Evidence: Mechanism-Based Reasoning, Not Clinical Proof

Despite widespread use and biological plausibility, the supplement recommendations in this chapter face a fundamental evidence gap:

- Fewer than five supplements discussed here have been tested in randomised placebo-controlled trials specifically in ME/CFS—notably NADH [542], CoQ10+NADH [7], and acetyl-L-carnitine [546]. Most evidence for other supplements derives from mechanistic reasoning, related-condition trials, or patient surveys with inherent reporting bias.
- The observation that “widely used” supplements dominate patient community recommendations does not constitute efficacy evidence; popular adoption may reflect hope, accessibility, and social transmission rather than therapeutic benefit.
- Combination protocols (e.g., CoQ10 + L-carnitine + D-ribose) are recommended based on complementary biochemical rationale, but potential interactions, optimal ratios, and combined efficacy have not been studied.
- Supplement quality, bioavailability, and dose-response relationships vary substantially between brands and formulations—factors not addressed by the general dosing ranges provided.

22.1 Foundational Supplements: Electrolytes and Hydration

For patients with autonomic dysfunction (reported in the majority of ME/CFS patients [97]), electrolyte and fluid management is often the single most impactful intervention.

22.1.1 Why Electrolytes Matter in ME/CFS

Autonomic dysfunction in ME/CFS frequently manifests as:

- Reduced blood volume (hypovolemia)
- Impaired vasoconstriction
- Excessive venous pooling
- Orthostatic intolerance

Adequate sodium, potassium, and fluid intake help maintain blood volume and support cardiovascular compensation.

22.1.2 Sodium

Rationale. Sodium increases blood volume by promoting water retention. POTS expert consensus guidelines recommend 10–12 g of salt daily (versus 2–3 g typical intake) [99].

Evidence. Moderate for POTS; limited ME/CFS-specific data, but widely reported helpful.

Forms.

- **Table salt:** Cheapest; 2.3 g sodium per teaspoon
- **Electrolyte drinks:** LMNT, Liquid IV, Drip Drop, Nuun—convenient but expensive
- **Salt tablets:** Precise dosing; some find easier than drinking salty fluids
- **Oral rehydration salts (ORS):** WHO formula; includes glucose for sodium-glucose cotransport

Practical Protocol.

1. Add 1/4–1/2 teaspoon salt to each liter of water
2. Drink before standing or activity
3. Morning loading: 16–24 oz with salt before getting up
4. Target: 2–3 L fluid plus 8–12 g salt daily (approximately 3–5 g elemental sodium)

Cautions.

- Check with physician if hypertension, heart failure, or kidney disease
- Some patients with hyperadrenergic POTS may worsen with excess sodium
- Monitor for edema; some is expected and indicates effectiveness
- **Potential for paradoxical intracellular effects:** ME/CFS patients may have baseline intracellular sodium elevation in muscle tissue [196]. Rapid sodium loading could theoretically worsen intracellular sodium overload via impaired Na^+/K^+ -ATPase function (see Section 22.1.6)

Energy Profile. Category A–B (context-dependent). Improved blood volume enhances oxygen and nutrient delivery to cells, supporting energy production. However, processing sodium via Na^+/K^+ -ATPase consumes an estimated 20–40% of cellular ATP [431]. In ME/CFS patients with documented intracellular sodium overload, additional sodium loading may paradoxically increase ATPase energy demands. Net effect depends on individual pathophysiology: beneficial for hypovolemia-dominant presentation, potentially harmful if intracellular sodium is already elevated.

22.1.3 Potassium

Rationale. Intracellular potassium is essential for nerve and muscle function. Some ME/CFS patients show functional potassium deficiency even with normal serum levels.

Evidence. Theoretical; no ME/CFS-specific trials.

Forms.

- **Potassium chloride:** Most common; Nu-Salt, Morton Lite Salt
- **Potassium citrate:** Better tolerated GI-wise
- **Coconut water:** Natural source (~600 mg per cup)

Dosing. 2,000–4,700 mg daily (food + supplements); start low.

Cautions. Excess potassium can cause cardiac arrhythmias. Do not exceed 99 mg per dose in supplement form without medical supervision. Those on ACE inhibitors, ARBs, or with kidney disease should be especially cautious.

Energy Profile. Category A (net energy provider). Essential cofactor for Na^+/K^+ -ATPase function—without adequate potassium, the pump cannot maintain membrane potential, and cellular energy production collapses. Correcting potassium deficiency directly enables ATP-dependent cellular processes. Minimal processing overhead.

22.1.4 Magnesium

Rationale. Magnesium is a cofactor for >300 enzymes, including ATP synthesis. Deficiency is common and underdiagnosed (serum magnesium poorly reflects tissue status). Relevant to ME/CFS because:

- Required for mitochondrial ATP production
- Modulates NMDA receptors (relevant to central sensitization)
- Supports autonomic function
- Promotes sleep (GABA-A receptor modulation)

Evidence. Low–Moderate for ME/CFS; one small trial showed benefit with IM magnesium sulfate.

Forms.

- **Magnesium glycinate:** Well-absorbed; calming; good for sleep; less GI upset
- **Magnesium malate:** Malic acid may support TCA cycle; often recommended for fibromyalgia
- **Magnesium L-threonate:** Crosses blood-brain barrier; may help cognition; expensive
- **Magnesium citrate:** Well-absorbed; can cause loose stools (useful if constipated)
- **Magnesium oxide:** Poorly absorbed; cheap; mainly useful as laxative
- **Magnesium taurate:** Combined with taurine; may benefit cardiovascular system

Dosing. 200–600 mg elemental magnesium daily; split doses. Start low (100–200 mg) and increase gradually. Bowel tolerance is the limiting factor for oral forms.

Practical Tip. Topical magnesium (Epsom salt baths, magnesium oil) provides modest absorption and may help with muscle symptoms, though evidence is limited.

Energy Profile. Category A (net energy provider). Required cofactor for over 300 enzymatic reactions, including ATP synthesis itself (ATP exists physiologically as Mg-ATP complex). Deficiency directly impairs energy production at multiple points: glycolysis, TCA cycle, and oxidative phosphorylation. Supplementation restores enzymatic capacity with minimal metabolic processing cost. Among the highest-priority supplements for energy-depleted patients.

22.1.5 Complete Electrolyte Formulas

Many patients find pre-mixed electrolyte formulas convenient. Key ingredients to look for:

- Sodium: 500–1000 mg per serving
- Potassium: 200–400 mg per serving
- Magnesium: 50–100 mg per serving
- Minimal or no sugar (some glucose aids sodium absorption; excessive sugar is counter-productive)

DIY Oral Rehydration Solution.

1 L water + 1/2 tsp salt + 1/4 tsp potassium chloride (Nu-Salt) + 2 tbsp sugar or honey + optional: squeeze of citrus

Cost: pennies per liter versus \$1–3 for commercial products.

△ **Warning 1: Histamine Intolerance and Electrolyte Products**

A substantial subset of ME/CFS patients have histamine intolerance (HIT) or mast cell activation syndrome (MCAS) [566], making standard electrolyte products problematic (see Section 7.7.1 for MCAS pathophysiology). Common triggers in commercial electrolyte formulas:

Ingredients to avoid with HIT/MCAS:

- **Citric acid:** Ubiquitous in commercial products; can trigger mast cell degranulation
- **Citrus flavorings:** Natural lemon, lime, orange flavors are high-histamine
- **Artificial colors:** Red 40, Yellow 5/6 can trigger mast cells
- **Fermented ingredients:** Some products contain fermented sugars
- **Stevia:** Triggers reactions in some sensitive individuals

HIT-safe DIY electrolyte recipe:

1 L filtered water + 1/2 tsp sea salt (unrefined) + 1/4 tsp potassium chloride + 1–2 tbsp maple syrup (pure, grade A)

Omit citrus entirely. Maple syrup provides glucose for sodium-glucose co-transport without histamine issues. Some patients tolerate small amounts of fresh ginger for flavor.

Commercial options for HIT/MCAS:

- LMNT unflavored (citric acid-free)
- Pure Encapsulations Electrolyte/Energy Formula
- DIY remains safest option for highly sensitive patients

Note: POTS patients with HIT/MCAS face a double challenge—they need electrolytes for blood volume but react to most products. Trial elimination of problematic ingredients before assuming electrolyte intolerance. Many patients who thought they “couldn’t tolerate electrolytes” actually couldn’t tolerate citric acid or flavorings; plain salt in water may work when commercial products don’t.

22.1.6 Caution: Electrolyte Initiation in ME/CFS

△ **Warning 2: Intracellular vs. Extracellular Sodium: A Critical Distinction**

Standard POTS/dysautonomia guidelines recommend high sodium intake (10–12 g/day) to expand *blood volume*—addressing *extracellular* sodium. However, emerging evidence suggests ME/CFS patients may already have *intracellular* sodium overload in muscle tissue, creating a potential paradox where aggressive sodium supplementation could worsen underlying pathophysiology even while improving some orthostatic symptoms [431, 196, 185].

The Evidence.

- **MRI findings (2022):** Sodium MRI demonstrated elevated baseline intracellular muscle sodium in ME/CFS patients across all five lower leg muscle compartments (12.2 mM vs 9.4 mM in controls for anterior extensors, $p = 0.003$). Post-exercise sodium accumulation was also greater (+30% vs +17% at 12 min) [196]. *Evidence quality: Low (single small study, $n = 6$ per group); requires replication.*
- **Functional correlation:** Elevated muscle sodium correlated inversely with hand grip strength ($p = 0.03$, $R^2 = 0.38$), suggesting functional consequences [196].

Speculation 64 (The NCX Reversal Mechanism). **Certainty: 0.25.** Mechanistic hypothesis based on indirect evidence (elevated intracellular sodium on MRI [196], known NCX biophysics); not directly tested in ME/CFS. Wirth and Scheibenbogen propose that β_2 -adrenergic receptor dysfunction impairs Na^+/K^+ -ATPase activity, allowing intracellular sodium accumulation [431, 185]. When intracellular sodium exceeds a threshold, the sodium-calcium exchanger (NCX) reverses direction—importing calcium instead of exporting it—causing calcium overload and mitochondrial dysfunction. This model suggests ME/CFS patients operate near the NCX reversal threshold, explaining why even minor exertion triggers post-exertional malaise.

Clinical Implications. Dietary sodium primarily affects blood volume, not intracellular muscle sodium directly. However:

- **Blood volume expansion remains beneficial** for most ME/CFS patients with orthostatic intolerance—the question is *rate of initiation*
- **Rapid sodium loading** during energy-depleted states could theoretically stress already-compromised Na^+/K^+ -ATPase function
- **Individual response varies**—some patients tolerate high sodium well; others report worsening with aggressive supplementation

“Start Low, Go Slow” Protocol. Given the uncertainty, conservative initiation is prudent:

1. **Week 1:** Baseline—maintain current sodium intake; establish symptom diary

2. **Week 2:** Add 500–1000 mg sodium daily (approximately 1/4 tsp salt or one electrolyte packet)
3. **Weeks 3–6:** If tolerated, increase by 500–1000 mg weekly
4. **Target:** Reach 3–6 g supplemental sodium over 4–6 weeks (in addition to dietary intake)
5. **Monitor:** Blood pressure (sitting and standing), energy levels, PEM threshold

Warning Signs—Stop or Reduce If:

- Increased muscle pain, weakness, or fatigue unexplained by activity
- New or worsening palpitations or resting tachycardia
- Worsening PEM threshold (crashing at lower activity levels)
- Sustained blood pressure >130/80 mmHg at rest [567]
- Significant peripheral edema beyond mild ankle swelling

Additional Considerations.

- **TRPM3 connection:** ME/CFS involves TRPM3 ion channel dysfunction affecting calcium handling (Section 7.1.1). The relationship between TRPM3 dysfunction and NCX-mediated calcium overload warrants investigation
- **Energy state matters:** Supplementation during crashes may be less well-tolerated than during stable periods
- **Electrolyte balance:** Always supplement sodium alongside potassium and magnesium
- **Long-term risks:** High sodium intake carries cardiovascular risks independent of blood pressure [567]; regular monitoring advisable

Observation 96 (Research Gap). The interaction between dietary sodium and intracellular muscle sodium in ME/CFS has not been directly studied. This caution is based on documented intracellular elevation, plausible mechanisms, and clinical reports of sodium intolerance in some patients. Prospective studies comparing titration protocols with sodium MRI endpoints would clarify this relationship.

22.2 Mitochondrial and Energy Support

Given the evidence for energy metabolism dysfunction in ME/CFS (Chapter 6), supplements supporting mitochondrial function are among the most commonly used.

22.2.1 Coenzyme Q10 (CoQ10/Ubiquinone/Ubiquinol)

Rationale. CoQ10 is a mobile electron carrier in the electron transport chain, shuttling electrons from Complex I and Complex II to Complex III, and is a potent lipid-soluble antioxidant.

Evidence. Moderate. Multiple small studies show benefit in ME/CFS [195, 568] (Castro-Marrero 2015: n=73, CoQ10+NADH combination) and fibromyalgia [569]. A 2022 systematic review and meta-analysis of 13 RCTs (n=1,126) found CoQ10 significantly reduced fatigue across several chronic conditions (cancer fatigue, MS, fibromyalgia, cardiovascular disease; not ME/CFS-specific) [570].

Forms.

- **Ubiquinone:** Oxidized form; must be converted to ubiquinol
- **Ubiquinol:** Reduced (active) form; better absorbed, especially over age 40; more expensive

Dosing.

- Typical: 100–300 mg daily
- Higher doses in studies: 400–600 mg daily
- Take with fat-containing meal for absorption
- Split doses if >200 mg

Response Timeline. Benefits may take 4–12 weeks to manifest.

Cautions. Generally well-tolerated. May reduce warfarin effectiveness (see Section 22.2.8, drug interactions). Can cause insomnia if taken late in day (clinical observation).

Energy Profile. Category A (net energy provider). Functions as a mobile electron carrier in the ETC (see Rationale above). See Chapter 21 for detailed energy profile.

22.2.2 NADH (Reduced Nicotinamide Adenine Dinucleotide)

Rationale. NADH is the reduced form of NAD^+ and serves as the primary electron donor to Complex I (NADH dehydrogenase) of the mitochondrial electron transport chain. Supplementing NADH directly provides electrons for ATP synthesis, complementing CoQ10 which shuttles electrons between complexes. Together, NADH and CoQ10 support electron supply at two distinct points in the proximal electron transport chain.

Evidence. Low–Moderate (Low for NADH monotherapy; Moderate for the NADH+CoQ10 combination).

- **Forsyth et al. (1999)** [542]: Randomized, double-blind, placebo-controlled crossover trial (n=26); 10 mg NADH daily for 4 weeks showed 31% response rate vs. 8% placebo
- **Santaella et al. (2004)** [543]: Randomized trial (n=31, no placebo arm) comparing NADH to an active control (nutritional supplements plus psychotherapy) over 24 months; significant improvement in the NADH group in the first three months ($p < 0.001$ vs. control), but outcomes later comparable between groups
- **Castro-Marrero et al. (2016)** [571]: RCT (n=80); CoQ10 (200 mg) + NADH (20 mg) improved maximum heart rate recovery; some improvement in fatigue perception, though fatigue effects were inconsistent across the broader programme of trials
- **Castro-Marrero et al. (2021)** [7]: Largest RCT to date (n=207, double-blind, placebo-controlled); 200 mg CoQ10 + 20 mg NADH daily for 12 weeks showed significant improvements in cognitive fatigue ($p < 0.001$), overall fatigue ($p = 0.022$), quality of life ($p < 0.05$), and sleep parameters (sleep duration at week 4, $p = 0.018$; sleep efficiency at week 8, $p = 0.038$). Benefits persisted 4 weeks post-treatment. No serious adverse events

The Castro-Marrero 2021 trial provides the strongest evidence for the NADH+CoQ10 combination in ME/CFS. It does not, however, distinguish the individual contribution of NADH from that of CoQ10. See Section 22.2.8 for combination rationale.

Forms.

- **Enteric-coated tablets:** Protect against gastric acid degradation; most common form
- **Sublingual:** Bypasses gastric degradation; may provide faster absorption
- **Enada:** Brand used in multiple clinical trials; standardized formulation

Dosing.

- Standard: 10–20 mg daily
- Clinical trial dose: 20 mg daily (Castro-Marrero protocol)
- Must be taken on an empty stomach, 30–60 minutes before breakfast (gastric acid degrades unprotected NADH)

Response Timeline. 4–8 weeks for monotherapy; combination with CoQ10 may show benefits within 4 weeks for sleep parameters.

Cautions. Generally well-tolerated. Unprotected oral NADH is degraded by stomach acid—use enteric-coated or sublingual forms only. Some patients report mild nausea if taken with food. Cost is moderate (\$20–40/month at therapeutic doses).

NADH vs. NAD⁺ Precursors. NADH and NAD⁺ precursors (NR, NMN) are complementary, not interchangeable—they target different axes of cellular energy metabolism. See Section 22.2.8 for the full two-axis framework and combination recommendations.

Energy Profile. Category A (net energy provider). Direct electron donor to Complex I; each NADH molecule reaching the mitochondrial matrix yields approximately 2.5 ATP via oxidative phosphorylation. Oral bioavailability is a limiting factor (see Forms above).

22.2.3 NAD⁺ Precursors: Nicotinamide Riboside (NR) and NMN

Rationale. NAD⁺ is essential for mitochondrial function, DNA repair, and cellular signaling. The Heng 2025 study [12] documented NAD⁺ metabolism abnormalities in ME/CFS. While direct oral NAD⁺ supplementation has poor bioavailability (gastric degradation), sublingual NAD⁺ (100–300 mg/day) offers a direct route, and precursors efficiently raise intracellular levels. NAD⁺ precursors replenish the cellular NAD⁺ pool for sirtuins and PARP-mediated DNA repair—a complementary mechanism to the direct electron donation provided by NADH (Section 22.2.2). For combination guidance, see Section 22.2.8.

Evidence. Preliminary. A 2025 RCT in Long COVID (n=58; 20-week intervention) showed NR 2000 mg/day increased whole-blood NAD⁺ levels 2.6–3.1 fold (measured at interim time-points of 5 and 10 weeks); cognitive benefits were variable, with overall group effects limited but some individuals showing improvement after ≥10 weeks [572].

Forms.

- **Nicotinamide Riboside (NR):** Tru Niagen is the most studied brand
- **Nicotinamide Mononucleotide (NMN):** One step closer to NAD⁺; theoretically more direct but less clinical data
- **Niacin (B3):** Cheapest NAD⁺ precursor but causes flushing; extended-release reduces flushing but has liver toxicity concerns
- **Nicotinamide:** No flushing; inhibits sirtuins via product inhibition at high concentrations (well-established in vitro [573]; clinical relevance at oral supplementation doses uncertain)
- **NAD⁺ sublingual:** Direct NAD⁺ administered under the tongue (100–300 mg/day); bypasses gastric degradation but bioavailability data remain limited. Substitutable with NR or NMN—choose one precursor strategy

Dosing.

- NR: 300–1000 mg daily typical; research doses up to 2000 mg
- NMN: 250–500 mg daily typical; up to 1000 mg in some regimens
- NAD⁺ sublingual: 100–300 mg daily

- Niacin: 500–1500 mg daily (with caution)

Emergency Use (Speculative). For emergency post-exertion PEM prevention, high-dose loading (1000–2000 mg NR or NMN immediately post-exertion, then 500 mg twice daily for 3–5 days) may prevent NAD⁺ depletion hypothetically attributable to PARP activation during DNA repair—this mechanism is plausible but unconfirmed in ME/CFS specifically [233]. **Caution:** This protocol has not been tested in any clinical trial. High-dose NR/NMN may cause gastrointestinal side effects and has not been safety-tested at these doses in ME/CFS patients. Discuss with physician before attempting. See Chapter 24, §24.10.3 for complete emergency protocol.

Response Timeline. May require 10+ weeks for noticeable benefit.

Cost Consideration. NR/NMN are expensive (\$50–150/month at therapeutic doses). Niacin is cheap but has tolerability issues.

Cautions. Generally well-tolerated at standard doses. Gastrointestinal discomfort may occur at high doses (≥ 1000 mg). Theoretical concern: elevated NAD⁺ may promote tumor growth in active cancer—rapidly proliferating cells may exploit NAD⁺ for energy and DNA repair [574]. Avoid during active cancer; consult oncologist before use in patients with cancer history. Extended-release niacin carries hepatotoxicity risk (see Section 22.2.8).

Energy Profile. Category A (net energy provider). Replenish cellular NAD⁺ pools required for hundreds of metabolic reactions including glycolysis, TCA cycle, and oxidative phosphorylation. Conversion from NR/NMN to NAD⁺ requires minimal ATP (one phosphorylation step). Net energy balance strongly positive. Emergency high-dose protocols (see Emergency Use above) are hypothesized to provide immediate NAD⁺ substrate for PARP-mediated DNA repair [233]—this rationale is mechanistically plausible but not yet confirmed in ME/CFS trials.

22.2.4 D-Ribose

Rationale. D-ribose is the pentose sugar component of ATP. Supplementation may accelerate ATP resynthesis after depletion.

Evidence. Low. A pilot open-label study (n=41) [8] and a larger multicenter replication (n=257; $p < 0.0001$ within-group for energy, sleep, and well-being) [528] in ME/CFS and fibromyalgia showed improvements with 5 g three times daily; no placebo comparison in either study, so results cannot be distinguished from placebo effect.

Dosing. 5 g three times daily (15 g/day total); can be reduced to 5–10 g daily for maintenance.

Emergency Use. For emergency post-exertion PEM prevention, higher acute doses (10–15 g immediately post-exertion, then 5 g every 4–6 hours for 24–48h) may be used as part of comprehensive crash prevention protocol. See Chapter 24, §24.10.3 for complete emergency protocol.

Practical Tips.

- Take with meals (can lower blood sugar)
- Sweet taste; dissolves in beverages
- Some patients report energy improvement within days

Cautions. May lower blood sugar; diabetics should monitor carefully.

Energy Profile. Category A (net energy provider). Direct ATP substrate (pentose sugar component). May reduce reliance on the energy-intensive pentose phosphate pathway by entering purine salvage directly [527]. Among the most direct energy substrates available—minimal metabolic processing required for ATP regeneration.

22.2.5 Acetyl-L-Carnitine (ALCAR) and L-Carnitine

Rationale. Carnitine transports fatty acids into mitochondria for oxidation. Deficiency impairs fat-based energy production. Acetyl-L-carnitine crosses the blood-brain barrier and may support cognitive function.

Evidence. Low (open-label studies only; no placebo-controlled trials). CFS patients show lower serum carnitine levels correlating with functional capacity [544]. L-carnitine 3 g/day improved 12 of 18 clinical parameters over 8 weeks (n=30, open-label) [545]. Acetyl-L-carnitine improved mental fatigue ($p = 0.015$) and propionyl-L-carnitine improved general fatigue ($p = 0.004$) in an open-label randomized study (n=90, no placebo arm) [546].

Forms.

- **L-Carnitine:** General mitochondrial support
- **Acetyl-L-Carnitine (ALCAR):** Better for cognitive symptoms; crosses BBB
- **Propionyl-L-Carnitine:** May be better for cardiovascular symptoms

Dosing. 500–2000 mg daily; split doses.

Response Timeline. 2–8 weeks.

Cautions. Can increase TMAO (a proposed cardiovascular risk marker) with chronic use; the mechanism—gut microbial conversion of carnitine to trimethylamine, then hepatic oxidation to TMAO—is documented in published literature, though the clinical significance for cardiovascular risk remains debated. Some patients experience overstimulation or insomnia. L-carnitine may theoretically lower seizure threshold in patients with pre-existing epilepsy based on mechanistic considerations; consult physician before use in seizure disorders (no controlled data; clinical caution).

Energy Profile. Category A (net energy provider). Enables mitochondrial fatty acid oxidation—the most energy-dense metabolic pathway. Each fatty acid molecule transported into mitochondria by carnitine yields substantially more ATP than glucose oxidation. Processing overhead negligible relative to energy yield.

22.2.6 Creatine

Rationale. Creatine buffers ATP, providing rapid energy during high-demand situations. Well-studied for muscle function; emerging evidence for cognitive benefits.

Evidence. Theoretical for ME/CFS; strong for lower limb muscle strength in general populations (meta-analysis) [575]. Emerging evidence for cognitive benefits under stress—sleep deprivation, mental fatigue—in general populations (small RCTs; no ME/CFS-specific data).

Dosing.

- Loading (optional): 5 g four times daily for 5–7 days. Note: the loading phase is derived from sports nutrition protocols; ME/CFS patients may prefer to skip loading and start directly at maintenance doses, consistent with conservative introduction (Section 22.7)
- Maintenance: 3–5 g daily

Cautions. Requires adequate hydration. May cause water retention. Generally considered safe for kidney function in healthy individuals at normal doses (no controlled evidence of harm; serum creatinine elevation reflects creatine metabolism, not renal impairment—see Additional interactions below).

Energy Profile. Category A (net energy provider). Directly buffers ATP via the phosphocreatine system—the fastest cellular energy reserve. Creatine kinase converts phosphocreatine to ATP within milliseconds, providing immediate energy during demand spikes. Also supports brain energy metabolism. Minimal processing demands.

22.2.7 PQQ (Pyrroloquinoline Quinone)

Rationale. PQQ stimulates mitochondrial biogenesis (creation of new mitochondria) and has antioxidant properties.

Evidence. Preliminary. Small RCTs in healthy adults show cognitive benefits: improved attention and working memory with 20 mg/day over 12 weeks [576, 577]; no ME/CFS-specific trials.

Dosing. 10–20 mg daily.

Energy Profile. Category A (indirect; delayed onset: weeks to months). Net energy provider through mitochondrial biogenesis rather than direct substrate supply. Stimulates the creation of new mitochondria, increasing total cellular energy production capacity over weeks to months. Unlike CoQ10 or NADH, PQQ does not directly participate in electron transport—its energy benefit is mediated entirely through expanding mitochondrial mass. Do not expect acute energy improvement.

22.2.8 Mitochondrial Energy Support: Combination Guide

The supplements in this section target distinct steps in cellular energy production. Understanding which ones are complementary and which are substitutable prevents both redundant spending and missed synergies.

Two Complementary Axes

Mitochondrial energy supplements can be organized into two mechanistically distinct axes. This framework is a conceptual model for understanding supplement complementarity; the empirical support comes from the Castro-Marrero trials, not from the framework itself.

Axis 1: Electron Transport Chain Support (NADH + CoQ10). NADH donates electrons to Complex I; CoQ10 shuttles electrons from Complex I and Complex II to Complex III. Together they support electron supply at two distinct points in the proximal ETC. The Castro-Marrero 2021 RCT (n=207) used exactly this combination—200 mg CoQ10 + 20 mg NADH—and demonstrated significant improvements in fatigue, cognition, and sleep parameters [7]. These two supplements form a functional pair and should be taken together.

Axis 2: NAD⁺ Pool Replenishment (choose one precursor). NR, NMN, and sublingual NAD⁺ all aim to increase intracellular NAD⁺ levels, supporting sirtuins (cellular stress response), PARP (DNA repair), and metabolic signaling. Because they converge on the same endpoint, one should be chosen based on evidence, tolerability, and cost. NR and NMN are the best-supported options and are largely interchangeable for this purpose:

- **NR:** Most clinical data; well-absorbed; Tru Niagen brand used in trials
- **NMN:** One enzymatic step closer to NAD⁺; popular but fewer published RCTs.¹
- **NAD⁺ sublingual:** Direct delivery; limited bioavailability data; cannot be considered fully equivalent to NR or NMN until absorption is better characterized

Niacin and nicotinamide are also NAD⁺ precursors but are not recommended for Axis 2: niacin causes flushing and carries hepatotoxicity risk at extended-release doses, while nicotinamide inhibits sirtuins at high concentrations (well-established in vitro; see Forms above)—directly counteracting one of Axis 2's intended benefits.

These two axes are complementary, not redundant. After donating electrons at Complex I, NADH is oxidized back to NAD⁺, so it does contribute to the mitochondrial NAD⁺ pool. However, supplemental NADH doses are modest (10–20 mg, approximately 15–30 μ mol), while total body NAD⁺ content is estimated at 1–3 g (approximately 1,500–4,500 μ mol; estimate based on tissue concentration measurements [410])—dwarfing the contribution from a single NADH dose. NADH's primary pharmacological action is therefore acute electron donation, not sustained NAD⁺ pool elevation. NAD⁺ precursors (NR, NMN) act through the salvage pathway to raise NAD⁺ levels across all cellular compartments—cytosolic, nuclear, and mitochondrial—at much larger scale. Thus, Axis 1 provides the electrons that drive ATP synthesis now, while Axis 2 replenishes the coenzyme pool that sustains energy metabolism, DNA repair, and cellular signaling over time.

Age-Stratified Dosing

△ Warning 3: Pediatric Evidence Gap

Pediatric dosing for mitochondrial supplements is extrapolated from adult data and body-weight scaling, not from pediatric randomized controlled trials. No ME/CFS-specific supplement trials have been conducted in children or adolescents. All pediatric use should be under physician supervision, with conservative starting doses and careful monitoring.

Recommended Combinations

Three evidence-informed combination tiers, using the two-axis framework above as an organizational heuristic; only the Minimal tier has direct RCT support. Use age-appropriate doses

¹Whether NMN requires conversion to NR for cellular uptake (via CD73) or enters cells directly (via the Slc12a8 transporter) remains debated. This does not affect the practical recommendation to choose one precursor.

Table 22.1: Mitochondrial supplement dosing by age group

Supplement	Adults (≥ 18 y)	Adolescents (12–17 y)	Children (6–11 y)	Notes
NADH	10–20 mg	5–10 mg	Not recommended	Empty stomach, 30 min before breakfast
CoQ10 (ubiquinol)	100–300 mg	50–100 mg	25–50 mg	With fat-containing meal; RCT dose 200 mg
NR	300–1000 mg	150–250 mg	Not recommended	300–500 mg starting; research doses up to 2000 mg
NMN	250–500 mg	125–250 mg	Not recommended	Choose one NAD ⁺ precursor
NAD ⁺ sublingual	100–300 mg	Not recommended	Not recommended	Limited safety data
D-Ribose	5–15 g	3–5 g	1–3 g	With meals; may lower blood sugar
ALCAR	500–2000 mg	250–500 mg	Not recommended	Split doses; may cause insomnia
Creatine	3–5 g	2–3 g	1–2 g	Adequate hydration essential
PQQ	10–20 mg	10 mg	Not recommended	Preliminary evidence only

from Table 22.1. For children under 12, where NADH is not recommended, the Minimal tier reduces to CoQ10 alone.

Minimal (the Castro-Marrero protocol). NADH + CoQ10. This is the only combination tested in a large ME/CFS RCT [7]. Start here.

Standard. NADH + CoQ10 + one NAD⁺ precursor (NR or NMN). Adds Axis 2 support for the cellular NAD⁺ pool. NR has the most clinical data; NMN is a reasonable alternative.

Comprehensive. NADH + CoQ10 + one NAD⁺ precursor + D-ribose + ALCAR. Adds ATP substrate (D-ribose) and fatty acid transport into mitochondria (carnitine). Higher cost; consider adding components one at a time to assess individual benefit.

What Not to Combine

Redundant combinations (not dangerous, but wasteful).

- **Multiple NAD⁺ precursors at full dose:** Taking NR 500 mg + NMN 500 mg + NAD⁺ sublingual 300 mg simultaneously provides no proportional benefit over a single precursor—they all converge on the same intracellular NAD⁺ pool. Because NR and NMN differ in molecular weight, a simple 50/50 mg split does not achieve equal molar doses—the simplest approach is to choose one precursor and use it at full dose
- **High-dose niacin alongside NR or NMN:** Niacin is also an NAD⁺ precursor; stacking it with NR/NMN adds flushing risk without clear additional benefit

△ Warning 4: Clinically Significant Drug Interactions

- **CoQ10 + warfarin:** CoQ10 may reduce warfarin effectiveness—the mechanism may involve enhanced CYP450-mediated warfarin hydroxylation [578]; a structural similarity to vitamin K has also been proposed but is less well-documented. Monitor INR closely if combining; adjust warfarin dose with physician guidance
- **Extended-release niacin + statins:** Extended-release niacin carries hepatotoxicity risk; combined with statins, risk of myopathy and rhabdomyolysis is increased [579]. Avoid unless under strict medical supervision with liver function monitoring
- **D-ribose + diabetes medications:** Ribose can lower blood glucose; monitor closely and adjust hypoglycemic medication if needed.
- **ALCAR + thyroid medication:** L-carnitine inhibits thyroid hormone nuclear uptake in a dose-dependent manner [580]; patients on thyroid replacement should consult their physician before supplementing L-carnitine and monitor thyroid function

Additional interactions.

- **CoQ10 + antihypertensives:** CoQ10 can lower blood pressure; combining with fludrocortisone or midodrine (common in ME/CFS for orthostatic intolerance) may require dose adjustment. Monitor blood pressure when initiating
- **D-ribose + glucose monitoring:** Some glucose meters can misread ribose as glucose, producing falsely elevated readings. Use ribose-compatible meters if monitoring blood glucose regularly
- **Creatine + renal function monitoring:** Creatine supplementation raises serum creatinine (a creatine metabolite), which can confound laboratory assessment of kidney function. Inform physicians of creatine use before renal panels; cystatin C is an alternative GFR marker unaffected by creatine intake

Timing conflicts.

- NADH requires an empty stomach (30–60 minutes before food); CoQ10 requires fat with meals. Take NADH first thing in the morning, CoQ10 with breakfast or lunch
- NAD⁺ precursors (NR, NMN) may have a mildly stimulating effect—avoid evening dosing if sleep-sensitive
- D-ribose should be taken with meals to prevent blood sugar dips
- ALCAR may cause insomnia or overstimulation; take split doses in the morning and early afternoon, not after 2–3 PM

22.3 Antioxidant and Anti-inflammatory Supplements

Oxidative stress is documented in ME/CFS [64]. Antioxidants may help, though evidence for specific supplements is limited.

22.3.1 N-Acetylcysteine (NAC)

NAC is one of the most versatile and evidence-supported supplements relevant to ME/CFS.

Rationale.

- **Glutathione precursor:** NAC provides cysteine, the rate-limiting amino acid for glutathione synthesis
- **Direct antioxidant:** Scavenges free radicals
- **Anti-inflammatory:** Reduces NF-κB activation
- **Mucolytic:** Thins mucus (originally developed for this)
- **Supports liver detoxification:** Used clinically for acetaminophen overdose
- **May reduce viral replication:** Some evidence for various viruses

Evidence. Moderate for general antioxidant/anti-inflammatory effects; preliminary for ME/CFS specifically. Widely used with generally positive patient reports.

Dosing.

- Typical: 600–1200 mg daily
- Higher doses: 1800–2400 mg daily (used in psychiatric applications)
- Take on empty stomach for best absorption
- Divide doses if >600 mg

Response Timeline. Antioxidant effects within days; systemic benefits may take 4–8 weeks.

Cautions.

- Can cause GI upset; start low
- Sulfur smell (normal)
- Theoretical concern about reducing beneficial ROS signaling; probably not clinically significant at normal doses
- May thin mucus excessively in some (actually beneficial for most)

Synergy. NAC works synergistically with:

- Glycine: Another glutathione precursor
- Selenium: Required for glutathione peroxidase function
- Vitamin C: Regenerates oxidized glutathione

Energy Profile. Category A–B (provider to neutral). Glutathione precursor that reduces oxidative stress, indirectly conserving cellular energy otherwise spent on damage repair. The conversion of NAC to glutathione requires minimal ATP (two enzymatic steps). By reducing oxidative damage to mitochondrial membranes and enzymes, NAC helps preserve energy production efficiency.

22.3.2 Alpha-Lipoic Acid (ALA)

Rationale. ALA is both water- and fat-soluble, allowing it to work in all cellular compartments. Regenerates other antioxidants (vitamins C and E, glutathione). Supports mitochondrial function.

Evidence. Theoretical for ME/CFS. (Moderate evidence exists for diabetic neuropathy, establishing the supplement's general bioactivity, but no ME/CFS-specific trials.)

Dosing. 300–600 mg daily; R-lipoic acid is the more bioactive form. Start at 100–150 mg in ME/CFS patients and titrate upward—sensitive patients may experience paradoxical worsening (increased fatigue, malaise, or gastrointestinal symptoms) at higher initial doses (clinical observation).

Cautions. Can lower blood sugar. May chelate minerals (take separately from mineral supplements).

Energy Profile. Category A (net energy provider at supplemental doses). Essential cofactor for pyruvate dehydrogenase and alpha-ketoglutarate dehydrogenase—enables energy production rather than consuming it. At high doses, mineral chelation may transiently reduce cofactor availability (see Cautions above). See Chapter 21 for detailed profile.

22.3.3 Omega-3 Fatty Acids (EPA/DHA)

Rationale.

- Anti-inflammatory (compete with omega-6 for inflammatory mediator synthesis)
- Support cell membrane fluidity
- Neuroprotective
- May support endothelial function (relevant to vascular hypothesis)

Evidence. Moderate for general anti-inflammatory effects; limited ME/CFS-specific data.

Dosing.

- General health: 1–2 g combined EPA/DHA daily
- Anti-inflammatory: 2–4 g daily
- Higher EPA ratio may be more anti-inflammatory

Quality Matters. Fish oil can oxidize; look for third-party tested products (IFOS certification). Triglyceride form is better absorbed than ethyl ester.

Energy Profile. Category B (energy-neutral). Anti-inflammatory effects may reduce the metabolic cost of chronic systemic inflammation. Incorporated into cell membranes through normal lipid metabolism with modest processing demands. No direct energy provision or significant energy consumption.

22.3.4 Curcumin

Rationale. Potent anti-inflammatory; inhibits NF- κ B; antioxidant.

Evidence. Strong for inflammation generally; no ME/CFS-specific trials.

Bioavailability Challenge. Standard curcumin is poorly absorbed (<1%). Enhanced formulations necessary:

- Curcumin + piperine (black pepper extract): 20× absorption increase
- Phytosome forms (Meriva): Lipid-bound for better absorption
- Nano-curcumin, micellar curcumin: Various enhanced delivery systems

Dosing. Depends on formulation; typically 500–2000 mg of enhanced curcumin daily.

Cautions. May thin blood; caution with anticoagulants. Can cause GI upset. May interact with some medications.

Energy Profile. Category B (energy-neutral). Hepatic glucuronidation and sulfation for metabolism, but enhanced bioavailability formulations reduce first-pass processing. Anti-inflammatory effects may offset processing costs by reducing chronic inflammation energy burden. Overall metabolic impact modest.

22.3.5 Quercetin

Rationale.

- Mast cell stabilizer (relevant if MCAS component)
- Antioxidant
- Anti-inflammatory
- May have antiviral properties

Evidence. Theoretical for ME/CFS; moderate for mast cell conditions.

Dosing. 500–1000 mg daily; enhanced absorption forms (quercetin phytosome) preferred.

Cautions. Generally well-tolerated. May interact with some antibiotics.

Energy Profile. Category B (energy-neutral). Flavonoid with mast cell-stabilizing properties processed through normal Phase II hepatic conjugation. Minimal energy demands. Mast cell stabilization may reduce the significant energy cost of chronic degranulation episodes.

22.4 B Vitamins

B vitamins are essential cofactors for energy metabolism and neurological function.

Energy Profile. Category A (net energy provider). B vitamins are essential cofactors for virtually every step of cellular energy production: thiamine (pyruvate dehydrogenase), riboflavin (Complex I/II), niacin (NAD⁺ synthesis), pantothenic acid (CoA synthesis), pyridoxine (amino acid metabolism). Deficiency at any point creates an energy production bottleneck. Supplementation removes bottlenecks with negligible processing cost.

22.4.1 Thiamine (B1)

Rationale. Essential for pyruvate dehydrogenase (PDH)—the enzyme that feeds pyruvate into the TCA cycle. PDH dysfunction is documented in ME/CFS.

Evidence. Preliminary. Case reports and small studies suggest high-dose thiamine may help a subset of ME/CFS patients. One Italian study used 600–1800 mg daily with significant benefit in chronic fatigue.

Forms.

- **Thiamine HCl:** Standard form; limited absorption
- **Benfotiamine:** Fat-soluble; better absorbed; doesn't cross BBB well
- **Thiamine TTFD (Allithiamine):** Lipid-soluble; crosses BBB; may be most relevant for ME/CFS

Dosing. Standard: 50–100 mg. High-dose protocols: 300–1800 mg daily (under medical supervision).

22.4.2 Riboflavin (B2)

Rationale. Precursor to FAD, essential for Complex II (succinate dehydrogenase) and fatty acid oxidation.

Evidence. Theoretical; studied in migraine prevention (400 mg daily).

Dosing. 25–400 mg daily. Harmless neon yellow urine at higher doses.

22.4.3 Niacin/Niacinamide (B3)

See NAD⁺ precursors above.

22.4.4 Pyridoxine/P5P (B6)

Rationale. Cofactor for neurotransmitter synthesis (serotonin, dopamine, GABA). The active form pyridoxal-5-phosphate (P5P) is the essential cofactor for aromatic L-amino acid decarboxylase, the enzyme that converts 5-HTP to serotonin. In the enterochromaffin-vagal pathway [225, 14], adequate P5P is therefore required for the final step of gut serotonin synthesis and downstream vagal activation. B6 is water-soluble and may be depleted by chronic inflammation and GI malabsorption—conditions common in ME/CFS; B6 status has not been systematically studied in ME/CFS cohorts.

Dosing. 25–100 mg daily. P5P (pyridoxal-5-phosphate) is the active form and may be better for those with conversion issues. When supporting gut serotonin synthesis (see Section 22.6.4), pair with 5-HTP or tryptophan supplementation.

Cautions. High doses (>200 mg/day chronically) can cause peripheral neuropathy. Assess RBC P5P levels if possible, as this is a more sensitive marker than serum B6.

22.4.5 Folate (B9)

Rationale. Essential for methylation and DNA synthesis.

Forms.

- **Folic acid:** Synthetic; requires conversion; some people have MTHFR variants impairing conversion
- **Methylfolate (5-MTHF):** Active form; bypasses MTHFR; often preferred
- **Folinic acid:** Intermediate; doesn't require MTHFR

Dosing. 400–1000 mcg daily; higher doses (up to 15 mg) used for specific indications.

Cautions. Must be balanced with B12; folate alone can mask B12 deficiency.

22.4.6 Cobalamin (B12)

Rationale. Essential for methylation, nerve function, and energy metabolism.

Evidence. Low–Moderate. Some ME/CFS patients respond dramatically to B12, especially sublingual or injectable forms; others show no benefit.

Forms.

- **Cyanocobalamin:** Cheapest; requires conversion; contains cyanide moiety (trivial amount)
- **Methylcobalamin:** Active methylated form; supports methylation
- **Adenosylcobalamin:** Active form used in mitochondria
- **Hydroxocobalamin:** Well-retained; often used in injections

Dosing. Oral: 1000–5000 mcg sublingual daily. Injections: 1000 mcg weekly to monthly (requires prescription in most countries).

Note on Testing. Serum B12 is a poor marker of tissue status. Methylmalonic acid (MMA) and homocysteine are more sensitive.

22.5 Vitamin D

Rationale. Vitamin D is actually a hormone with effects on:

- Immune regulation (relevant to ME/CFS immune dysfunction)
- Muscle function
- Mood
- Bone health

Deficiency is common in ME/CFS patients (often housebound with limited sun exposure).

Evidence. Moderate for general health; limited ME/CFS-specific data.

Target Levels. Controversy exists:

- Conventional: 30–50 ng/mL (75–125 nmol/L)
- Some ME/CFS practitioners target: 50–80 ng/mL
- Toxicity typically >150 ng/mL

Dosing.

- Maintenance: 1000–2000 IU daily
- Deficiency correction: 5000–10,000 IU daily for 8–12 weeks, then retest
- Take with fat-containing meal

Cofactors. Vitamin D requires cofactors for proper function:

- **Magnesium:** Required for vitamin D activation
- **Vitamin K2:** Directs calcium to bones (away from arteries)
- **Vitamin A:** Balances vitamin D effects

Energy Profile. Category B (energy-neutral). Immune regulatory and gene expression effects operate through nuclear receptor signaling with minimal direct energy cost. Long half-life means infrequent dosing keeps metabolic processing burden low. No direct energy provision.

22.6 Amino Acids

22.6.1 Taurine

Rationale.

- Mitochondrial membrane stabilization
- Antioxidant
- Supports bile acid conjugation
- May support cardiac and nervous system function
- Autonomic support

Evidence. Theoretical for ME/CFS; widely used.

Dosing. 500–3000 mg daily.

Energy Profile. Category A–B (provider to neutral). Conditionally essential amino acid that stabilizes mitochondrial membranes and regulates calcium homeostasis—both critical for mitochondrial energy production. Also functions as an osmolyte supporting cell volume regulation with minimal ATP cost. May improve mitochondrial efficiency without direct ATP provision.

22.6.2 Glycine

Rationale.

- Glutathione precursor (with NAC)
- Inhibitory neurotransmitter (calming)
- Supports collagen synthesis
- May improve sleep quality

Dosing. 1–3 g daily; 3 g before bed for sleep.

Energy Profile. Category A–B (provider to neutral). Glycine is one of three glutathione precursors (alongside cysteine and glutamate; NAC is typically rate-limiting for cysteine availability), supporting mitochondrial protection from oxidative damage. Also serves as a conjugation substrate for Phase II hepatic detoxification (glycine N-acyltransferase), potentially reducing the metabolic burden of processing other supplements. Minimal processing cost as an endogenous amino acid.

22.6.3 L-Glutamine

Rationale.

- Gut barrier support
- Immune cell fuel
- Glutathione precursor

Dosing. 5–15 g daily for gut support.

Cautions. Some patients with neurological sensitivity may not tolerate glutamine (converts to glutamate).

Energy Profile. Category A–B (provider to neutral). Primary fuel for enterocytes (intestinal cells) and immune cells. Provides energy substrate for gut barrier maintenance, potentially reducing the energy cost of intestinal permeability and immune activation from microbial translocation. Also serves as gluconeogenic substrate when glucose is depleted.

22.6.4 5-HTP (5-Hydroxytryptophan)

Rationale. 5-HTP is the immediate precursor to serotonin, produced from tryptophan by tryptophan hydroxylase. Supplementing with 5-HTP bypasses the IDO2 (and IDO1) diversion point [225]: in ME/CFS, IDO2 is the primary enzyme shunting tryptophan into the kynurenine pathway rather than toward serotonin synthesis [18]. Because 5-HTP is not a substrate for IDO enzymes, it directly enters the serotonin synthesis pathway regardless of IDO activity. In the gut, enterochromaffin cells convert 5-HTP to serotonin, which signals through vagal afferents to regulate autonomic tone [14]. Supplemental 5-HTP may therefore partially compensate for impaired peripheral serotonin synthesis in ME/CFS patients with gut dysbiosis.

Evidence. Theoretical for ME/CFS-specific gut-vagal pathway; limited direct trial evidence. Best supported by mechanistic extrapolation from IDO2 pathway research in Long COVID and the enterochromaffin-vagal model [225].

Dosing. 50–100 mg two to three times daily. Start at 50 mg once daily and titrate slowly. Requires adequate P5P (Section 22.4) as cofactor for the decarboxylase step.

Cautions.

- **Serotonin syndrome:** Contraindicated with SSRIs, SNRIs, MAOIs, tramadol, or other serotonergic drugs. This is a serious risk—medical supervision required if on any of these medications.
- Peripheral decarboxylation can cause systemic serotonin effects (GI cramping, nausea, diarrhea) before central effects appear.
- Avoid in pregnancy.
- Start very low if histamine-sensitive (serotonin can release histamine).

Energy Profile. Category B–C (neutral to mildly demanding). Conversion to serotonin via aromatic L-amino acid decarboxylase requires pyridoxal phosphate (vitamin B6) cofactor. The serotonin synthesis pathway itself is not particularly energy-intensive, but serotonergic effects can trigger mast cell activation in susceptible patients (MCAS comorbidity), and immune activation is energy-costly. Caution: serotonin syndrome risk with SSRIs represents a potentially catastrophic energy crisis.

22.7 Practical Supplement Protocols

Given the complexity, here are evidence-informed starting points organized by symptom cluster and budget.

22.7.1 Minimal Cost Protocol (Under \$30/month)

For patients with limited resources:

1. **Electrolytes:** Salt + potassium salt (Nu-Salt) + DIY rehydration (\$5/month)
2. **Magnesium glycinate:** 200–400 mg at bedtime (\$10/month)
3. **B-complex:** Basic B-complex with methylated B12/folate (\$10/month)
4. **Vitamin D3:** 2000–5000 IU daily (\$5/month)

This addresses the most common deficiencies and supports autonomic function. All components are Category A (net energy providers) or B (energy-neutral), making this protocol safe for all severity levels including very severe patients. See Section 25.1.2 for the energy category framework.

22.7.2 Moderate Protocol (\$50–100/month)

Adding mitochondrial and antioxidant support. **Severity note:** For moderate-to-severe patients, establish the minimal protocol for at least 4 weeks before adding these supplements. NADH and CoQ10 may be introduced together as a pair (consistent with the Castro-Marrero protocol); introduce each subsequent supplement individually with 1–2 week observation windows. All additions are Category A or B.

1. Everything in minimal protocol, plus:
2. **NADH + CoQ10 (ubiquinol):** 20 mg NADH + 200 mg CoQ10 daily—the Castro-Marrero protocol [7] (\$40–60/month combined)
3. **NAC:** 600–1200 mg daily (\$10–15/month)
4. **Omega-3:** 2 g EPA/DHA daily (\$15–20/month)
5. **D-ribose:** 5–10 g daily (\$15–20/month)

22.7.3 Comprehensive Protocol (\$100–200/month)

For mild-to-moderate patients who can afford broader support. **Severity note:** For severe and very severe patients, this protocol should only be attempted after documented tolerance of the moderate protocol over 8+ weeks. Physician supervision recommended. While all components are Category A–B, the cumulative hepatic processing load of multiple supplements warrants caution in energy-depleted patients (see Section 25.1.1).

1. Everything above, plus:
2. **NR or NMN:** 300–500 mg daily (\$40–60/month)
3. **Acetyl-L-carnitine:** 1000 mg daily (\$15/month)
4. **Alpha-lipoic acid:** 300 mg daily (\$10/month)
5. **Curcumin (enhanced):** 500 mg daily (\$15–20/month)

22.7.4 By Symptom Cluster

Predominant Orthostatic Symptoms.

- Electrolytes (priority)
- Magnesium
- Taurine
- Licorice root (caution: raises BP)

Predominant Cognitive Symptoms.

- NADH + CoQ10 (cognitive fatigue improvement; Castro-Marrero 2021 RCT, n=207)
- Magnesium L-threonate
- Acetyl-L-carnitine
- Omega-3 (high DHA)
- NR/NMN
- Creatine

Predominant Immune/Inflammatory Symptoms.

- NAC
- Omega-3
- Curcumin
- Quercetin (especially if mast cell component)
- Vitamin D (optimize)

Predominant Muscle/Fatigue Symptoms.

- NADH + CoQ10 (electron transport chain support; Section [22.2.8](#))
- D-ribose (ATP backbone resynthesis)
- L-carnitine (fatty acid transport into mitochondria)
- Creatine (rapid ATP buffering via phosphocreatine)
- Magnesium malate (cofactor for >300 enzymatic reactions; malate feeds the TCA cycle)

22.7.5 Introduction Strategy

△ Warning 5: One at a Time

ME/CFS patients often have multiple sensitivities (clinical experience). Introducing multiple supplements simultaneously makes it impossible to identify what helps or harms. Start one new supplement at a time, at low dose, and wait 1–2 weeks before adding another. Keep a symptom diary.

Suggested Order.

1. Electrolytes and magnesium (foundational; generally well-tolerated with gradual titration—see Section 22.1.6)
2. B vitamins (essential cofactors)
3. CoQ10 + NADH (exception to the one-at-a-time principle: the RCT evidence is for the combination, not the individual components; see Section 22.2.8). NADH requires an empty stomach 30–60 min before breakfast; CoQ10 is taken with a fat-containing meal—these timing constraints naturally separate the two supplements
4. NAC (antioxidant; watch for sulfur sensitivity)
5. Additional mitochondrial support (NAD⁺ precursors, D-ribose, ALCAR) based on response

22.8 Additional Supplements

22.8.1 Medium-Chain Triglycerides (MCT)

Rationale. MCTs bypass normal fat digestion and are converted directly to ketones by the liver. Ketones provide alternative brain and muscle fuel, potentially bypassing impaired glucose metabolism.

Evidence. Theoretical for ME/CFS; moderate for cognitive support in other conditions.

Dosing. Start with 1 teaspoon and increase slowly to 1–2 tablespoons daily. Rapid introduction causes GI distress.

Energy Profile. Category A (net energy provider). Converts directly to ketone bodies in the liver, bypassing the carnitine transport system and providing immediate alternative fuel for brain and muscle. Ketogenesis from MCTs is metabolically efficient compared to long-chain fatty acid oxidation. Provides energy substrate that bypasses potentially impaired glucose metabolism pathways.

22.8.2 Resveratrol

Rationale. Activates sirtuins and AMPK; promotes mitochondrial biogenesis; antioxidant.

Evidence. Preliminary; animal studies promising but human data limited.

Dosing. 150–500 mg daily; trans-resveratrol is the active form.

Energy Profile. Category B (energy-neutral). Resveratrol activates SIRT1 and AMPK signaling pathways that promote mitochondrial biogenesis over weeks to months, but does not directly provide ATP substrates. Requires hepatic CYP450 metabolism (CYP1A2, CYP3A4) for glucuronidation and sulfation, imposing modest metabolic processing cost. Net effect is neutral in the short term; potential long-term energy benefit via increased mitochondrial mass is speculative.

22.8.3 Melatonin

Rationale.

- Sleep initiation and circadian rhythm regulation
- Potent antioxidant (especially mitochondrial)
- Immune modulation
- Anti-inflammatory

Evidence. Moderate for sleep; theoretical for other effects in ME/CFS.

Dosing.

- Sleep: 0.5–3 mg, 30–60 minutes before bed
- Some protocols use higher doses (5–20 mg) for antioxidant effects
- Extended-release forms for sleep maintenance issues

Note. “Less is more” for sleep—higher doses can paradoxically worsen sleep quality. Start at 0.5 mg.

Energy Profile. Category B (energy-neutral). Endogenous hormone with minimal hepatic processing at physiological replacement doses. Sleep improvement may indirectly enhance energy restoration during rest. No significant metabolic burden.

22.8.4 Pregnenolone

Pregnenolone is a neurosteroid synthesized from cholesterol, serving as the precursor for all other steroid hormones including DHEA, progesterone, cortisol, and testosterone.

Rationale.

- **TRPM3 activation:** Pregnenolone sulfate (PregS) is an endogenous activator of TRPM3 ion channels (see Section 14.21)
- **Cognitive support:** Enhances memory formation and retrieval; neuroprotective
- **Neuronal myelination:** Supports myelin synthesis and maintenance
- **Fatigue relevance:** Low pregnenolone levels have been reported in ME/CFS patients
- **Immune modulation:** Modulates neurotransmitter-gated channels in immune cells

TRPM3 Connection. Reduced TRPM3 function in ME/CFS NK cells has been demonstrated using pregnenolone sulfate stimulation. TRPM3 channels in ME/CFS patients show significantly reduced calcium flux when stimulated with PregS. Theoretically, increasing pregnenolone availability might partially compensate for channel dysfunction by increasing agonist concentration, though this remains speculative. The established finding is that TRPM3 dysfunction can be partially restored by low-dose naltrexone (LDN), which may explain LDN's efficacy in some ME/CFS patients.

Evidence.

- Low pregnenolone levels documented in ME/CFS (clinical observations)
- No randomized controlled trials specifically in ME/CFS
- General evidence for cognitive and energy benefits in aging populations
- Theoretical basis from TRPM3 research

Dosing.

- Typical starting dose: 10–30 mg daily
- Some protocols use 50–100 mg daily
- Take in the morning (may be stimulating)
- Sublingual forms may have better absorption

Cautions.

- May affect hormone levels—caution in hormone-sensitive conditions
- Can be converted to androgens or estrogens depending on individual biochemistry
- Monitor for mood changes (anxiety, irritability)
- Avoid in pregnancy

- Consider baseline hormone testing before long-term use

Speculation 65 (Pregnenolone + LDN Synergy). **Certainty: 0.20.** No clinical trials have evaluated this combination; rationale is purely mechanistic extrapolation from separately-characterised TRPM3 effects of each agent.

Both pregnenolone sulfate and low-dose naltrexone influence TRPM3 function in ME/CFS, potentially through different mechanisms. Pregnenolone sulfate is a direct TRPM3 agonist, while LDN appears to restore TRPM3 responsiveness (mechanism not fully characterized). Theoretically, combining adequate pregnenolone levels with LDN might optimize TRPM3 function—the channel would be more responsive (LDN effect) and have adequate agonist (pregnenolone). No clinical trials have evaluated this combination, but given the favorable safety profiles of both, it represents a rational empirical approach for patients who have partial response to LDN alone.

Energy Profile. Category B–C (neutral to mildly demanding). Neurosteroid requiring hepatic CYP-mediated metabolism for conversion to downstream hormones (DHEA, progesterone, cortisol). Steroidogenesis is moderately energy-intensive. At supplemental doses, processing burden is modest but non-negligible.

22.8.5 Zinc

Rationale. Zinc is a trace mineral essential for gut barrier integrity, antioxidant protection in enterocytes, and neurotransmitter function. In the context of the enterochromaffin-vagal pathway [225], zinc supports multiple nodes: (1) tight junction integrity reduces intestinal permeability and bacterial translocation; (2) antioxidant mechanisms in enterocytes that may protect enterochromaffin cells; (3) neurotransmitter synthesis and autonomic nerve function. Zinc is commonly depleted in ME/CFS due to malabsorption, chronic inflammation (acute-phase redistribution to liver), and reduced dietary diversity from fatigue limiting food preparation.

Forms.

- **Zinc glycinate:** Well-absorbed, gentle on GI tract
- **Zinc carnosine:** Combines zinc with carnosine dipeptide; additional mucosal healing properties; preferred for gut repair (see Chapter 18)
- **Zinc citrate or picolinate:** Good bioavailability alternatives
- **Avoid zinc oxide:** Poor bioavailability

Dosing. 15–30 mg elemental zinc daily. Zinc carnosine 75 mg provides approximately 16 mg elemental zinc. Take away from or between meals for maximum absorption.

Cautions.

- Long-term zinc supplementation above 40 mg elemental zinc/day can deplete copper—monitor copper status with extended use or consider zinc-copper combination
- Take separately from calcium, magnesium, and iron (compete for absorption)
- Check serum zinc or RBC zinc before high-dose supplementation; levels may be misleadingly normal during inflammation despite tissue deficiency

Energy Profile. Category B (energy-neutral). Essential trace mineral and enzymatic cofactor absorbed through intestinal metallothionein pathways. Processing demands minimal at supplemental doses (15–30 mg). Supports immune function and wound healing without significant direct energy cost.

22.9 Probiotics and Gut Health

Given the documented gut microbiome abnormalities in ME/CFS [226, 349], gut-targeted interventions may help.

22.9.1 Probiotics

Rationale. May restore beneficial bacteria, reduce gut permeability, modulate immune function, and reduce systemic inflammation.

Evidence. Low–Moderate. Two small RCTs in ME/CFS showed modest benefit.

Strain Selection.

- *Lactobacillus* and *Bifidobacterium* species: Most studied; generally safe
- *Saccharomyces boulardii*: Yeast-based; may help after antibiotics
- Soil-based organisms (SBOs): More controversial; some find helpful

Practical Notes.

- Start low; die-off reactions possible
- May take 4–8 weeks to assess
- Quality varies enormously between brands
- Refrigerated products generally more viable

Energy Profile. Category B (energy-neutral). Live organisms that colonize the gut without requiring host metabolic processing. May improve nutrient absorption and reduce gut inflammation energy costs. No direct ATP demand from supplementation.

22.9.2 Prebiotics

Prebiotics feed beneficial bacteria. Options include:

- Partially hydrolyzed guar gum (PHGG)
- **Resistant starch** (see below for type selection)
- Inulin and FOS (can cause bloating)
- Acacia fiber (generally well-tolerated)

Resistant Starch Types. Not all resistant starch is equivalent. Two types differ in where they are fermented in the colon:

- **RS2 (raw potato starch, green banana flour):** Fermented primarily in the proximal colon; supports butyrate-producing bacterial communities [349]. Best starting choice. Dose: 1–2 tablespoons raw potato starch daily.
- **RS3 (cooked and cooled rice, potatoes, pasta):** More distal colonic fermentation; may complement RS2 for supporting butyrate production throughout the colon [349]. Introduce after RS2 is tolerated.

Caution. Prebiotics are contraindicated in active SIBO—fermentable substrate feeds small intestinal overgrowth and can severely worsen symptoms. Test for SIBO (breath test) before introducing resistant starch. Start with very small amounts (1 teaspoon) and increase slowly over 4–6 weeks.

22.9.3 Butyrate and Tributyrin

Rationale. Butyrate is the primary short-chain fatty acid (SCFA) fuel for colonocytes and a key upstream signal in the enterochromaffin-vagal pathway [14, 225]. Gut microbiome studies show ME/CFS patients have significantly reduced butyrate-producing bacterial capacity [349], correlating with fatigue severity. Butyrate supplementation aims to compensate for this deficit, enhancing enterochromaffin serotonin synthesis and downstream vagal afferent signaling. Butyrate also directly activates SCFA receptors (GPR41/43) on vagal neurons, providing a parallel pathway independent of serotonin.

Forms.

- **Sodium butyrate:** Fast-absorbing in proximal colon; well-studied; strong odour (similar to parmesan). Dose: 500 mg–2 g three times daily with meals.
- **Tributyrin:** A triglyceride of butyric acid, more lipophilic; slower release for sustained distal colon exposure. Dose: 1–2 g three times daily. Lower odour than sodium butyrate.
- **Calcium/magnesium butyrate:** Combined mineral-butyrate salt; another option with good tolerability.
- **Dietary butyrate:** Cooked and cooled starchy foods (RS3), high-fat fermented dairy (butter, ghee contain short-chain fats), and adequate fiber intake to feed butyrate producers.

Evidence. Direct ME/CFS supplementation trials are lacking. Mechanistic rationale is strong based on documented butyrate-producer deficiency in ME/CFS [349] and established butyrate-enterochromaffin-vagal signaling [14]. Safety profile is excellent.

Combination Approach. For targeting the enterochromaffin-vagal pathway, combining butyrate with 5-HTP and P5P (Section 22.6.4) addresses both upstream substrate supply (butyrate) and downstream serotonin synthesis (5-HTP + P5P cofactor). For a phased clinical protocol incorporating these agents with sequencing rationale and dosing, see Section 18.5.8 in Chapter 18.

Cautions. Very safe; main side effect is GI discomfort (bloating, loose stool) if dose escalated too quickly. Start low and increase gradually. The odour of sodium butyrate may be off-putting—encapsulated forms minimize this.

Energy Profile. Category A–B (provider). Short-chain fatty acid that serves as primary energy source for colonocytes (intestinal cells). Direct mitochondrial fuel that supports gut barrier integrity. By providing colonocyte fuel, butyrate spares systemic energy resources for other organs. Processing demands minimal.

22.10 Supplement Quality and Safety

22.10.1 Quality Considerations

Supplements are minimally regulated. Quality varies enormously:

- **Third-party testing:** Look for NSF, USP, ConsumerLab, or IFOS certification
- **GMP compliance:** Minimum standard; not sufficient alone
- **Bioavailability:** Cheap forms may not be absorbed
- **Contaminants:** Heavy metals, pesticides, adulterants possible

22.10.2 Drug Interactions

Common interactions to be aware of:

- **Blood thinners (warfarin):** CoQ10, omega-3, vitamin E, curcumin can affect clotting
- **Blood pressure medications:** Potassium, magnesium, licorice can interact
- **Thyroid medications:** Take separately from calcium, magnesium, iron
- **Immunosuppressants:** Immune-modulating supplements may interfere

22.10.3 When to Stop

Discontinue a supplement if:

- Clear worsening of symptoms
- No benefit after adequate trial (typically 8–12 weeks)
- Financial burden outweighs uncertain benefit
- Interactions with new medications

Observation 97 (The Supplement Trap). Many ME/CFS patients accumulate large, expensive supplement regimens over time without systematically evaluating benefit. Periodically reassess: stop everything non-essential for 2–4 weeks, then reintroduce one at a time. You may discover that many supplements you’ve been taking for years provide no discernible benefit.

22.11 Conclusion

Supplements can play a supportive role in ME/CFS management, but expectations should be realistic:

- **No supplement cures ME/CFS**
- **Effects are typically modest**—10–20% symptom improvement is a good outcome
- **Response varies enormously** between individuals
- **Cost adds up**—prioritize based on evidence and your specific symptoms
- **Foundation first:** Electrolytes, magnesium, B vitamins, and vitamin D before exotic interventions

The following supplements have the strongest mechanistic rationale and/or limited clinical evidence, listed in approximate order of evidence strength:

1. Electrolytes (especially if orthostatic symptoms)
2. NADH + CoQ10 combination (ME/CFS RCT evidence: Castro-Marrero et al. 207-patient trial [7])
3. NAC (antioxidant, glutathione precursor; no ME/CFS-specific RCT)
4. Magnesium (ubiquitous cofactor, often deficient; ME/CFS evidence is observational)

5. NAD⁺ precursors (emerging mechanistic rationale; no completed ME/CFS RCT)

Work with a knowledgeable healthcare provider when possible, especially for higher-dose protocols or if taking multiple medications.

23 Lifestyle and Non-Pharmacological Interventions

23.1 Pacing and Energy Management

Pacing is the most evidence-based and universally recommended non-pharmacological intervention for ME/CFS [61, 127]. Unlike graded exercise therapy (which can be harmful), pacing recognizes the physiological limitations imposed by metabolic dysfunction and aims to prevent post-exertional malaise while maintaining the highest sustainable level of activity.

23.1.1 Energy Envelope Theory

Conceptual Foundation

The energy envelope theory, developed through patient advocacy and clinical observation, posits that ME/CFS patients have a limited daily “energy budget” beyond which exertion triggers PEM [581]. Exceeding this envelope results in:

- Symptom exacerbation within 12–48 hours
- Prolonged recovery periods (days to weeks)
- Potential cumulative damage with repeated violations
- Progressive functional decline in severe cases

Staying within the energy envelope does not cure ME/CFS, but prevents the boom-bust cycle that worsens baseline function and quality of life.

Objective Evidence from Two-Day CPET

The energy envelope concept received objective validation from two-day cardiopulmonary exercise testing studies. Keller et al. (2024) demonstrated that ME/CFS patients, unlike healthy controls or those with deconditioning alone, show reproducible physiological impairment following maximal exertion [1]. Day 2 testing revealed:

- 5–8% declines in cardiopulmonary parameters (VO_2peak , work, ventilation)
- Worsening anaerobic threshold (earlier lactate accumulation)
- Doubling of severe impairment classification (14% to 27%)
- Recovery requiring 13+ days in ME/CFS versus ~2 days in controls

This objectively demonstrates that exertional stress produces measurable metabolic failure that persists well beyond 24 hours—providing a scientific foundation for activity restriction and pacing strategies.

Heart Rate Monitoring

Heart rate provides a practical, real-time proxy for metabolic stress. The Workwell Foundation and other clinical researchers recommend using heart rate thresholds to prevent PEM [1]:

- **Determine anaerobic threshold (AT):** Ideally via CPET; alternatively, estimate as 60–70% of age-predicted maximum heart rate in moderate-to-severe ME/CFS
- **Set activity threshold:** AT – 10 to 15 bpm as a safe upper limit
- **Continuous monitoring:** Wearable heart rate monitors enable real-time pacing
- **Account for delayed response:** Heart rate may lag behind metabolic demand; stop before reaching threshold

For example, a patient with AT of 115 bpm would aim to keep activity-related heart rate below 100–105 bpm.

Avoiding Boom-Bust Cycles

Many ME/CFS patients exhibit a maladaptive pattern:

1. **“Good day”:** Feeling relatively better, patient attempts normal or compensatory activity
2. **Overexertion:** Exceeds energy envelope, often unknowingly
3. **Crash (PEM):** Severe symptom exacerbation 12–72 hours later
4. **Extended recovery:** Days to weeks of reduced function
5. **Repeat:** Upon partial recovery, cycle repeats

This pattern prevents stabilization and may contribute to progressive worsening. Breaking the cycle requires:

- Consistent activity limits even on “good days”
- Recognition that feeling better does not mean capacity has increased
- Pre-planned rest periods regardless of symptom level
- Objective monitoring (heart rate, step counts) to override subjective assessment

Activity Tracking

Systematic tracking helps establish individual energy envelopes:

- **Daily logs:** Record activities, duration, intensity, and subsequent symptoms
- **Delayed symptom correlation:** Note PEM onset 12–72 hours post-activity
- **Pattern identification:** Identify activities that consistently trigger crashes

- **Threshold determination:** Establish personal limits for physical, cognitive, and social exertion
- **Activity-type categorization:** Distinguish cognitive, physical, emotional, and passive exertion when logging; passive activities typically do not trigger PEM, while the remaining categories may carry substantially different individual thresholds
- **Gradual adjustments:** Make small, monitored changes to activity levels

Digital tools (smartphone apps, wearables) can facilitate tracking, though screen time itself may be limited by cognitive symptoms.

23.1.2 Practical Pacing Strategies

Activity Planning and Prioritization

Effective pacing requires deliberate planning:

- **Essential vs. non-essential:** Prioritize critical activities (medical care, basic hygiene) over optional ones
- **Activity spreading:** Distribute demanding tasks across days or weeks
- **Anticipatory rest:** Build in recovery time before and after effortful activities
- **Delegation:** Accept help for tasks that exceed capacity
- **Simplified alternatives:** Replace high-energy activities with lower-energy versions (e.g., seated shower, prepared meals)

Rest Breaks

Strategic rest prevents cumulative energy depletion:

- **Prophylactic rest:** Rest before exhaustion, not after
- **Duration:** Even 5–15 minute breaks can prevent PEM if timed appropriately
- **Quality:** True rest (lying down, minimal stimulation) more effective than passive sitting
- **Scheduled intervals:** Build rest into routines (e.g., 30 minutes activity, 15 minutes rest)
- **Cognitive rest:** Limit screen time, reading, and mentally demanding tasks

Energy Conservation Techniques

Practical strategies reduce energy expenditure:

- **Seated activities:** Sit while cooking, showering, dressing
- **Adaptive equipment:** Shower chairs, reachers, electric can openers, voice control devices
- **Minimize trips:** Arrange living space to reduce walking distances; consolidate errands
- **Prepared foods:** Use convenience foods to reduce cooking energy
- **Postural management:** Lying down whenever possible to reduce orthostatic demand

Cognitive Pacing

Mental exertion triggers PEM as readily as physical activity:

- **Limit screen time:** Reduce visual and cognitive load
- **Simplify decisions:** Minimize daily choices (routines, meal planning, wardrobe simplification)
- **Reduce multitasking:** Focus on one task at a time
- **Communication management:** Batch messages; use voice-to-text; set boundaries
- **Avoid cognitively demanding media:** Complex plots, dense reading may exceed budget

Social and Emotional Energy

Social interaction, while psychologically beneficial, requires substantial energy:

- **Shorter visits:** Limit duration of social contacts
- **Low-stimulation settings:** Quiet, familiar environments better than crowded, noisy ones
- **Text-based communication:** Often less demanding than phone or video calls
- **Pre-planned exit strategies:** Permission to leave gatherings early
- **Post-social recovery:** Schedule recovery time after social activities

23.1.3 Activity-Based Dietary Timing for Severe Patients

For severe ME/CFS patients—particularly those with postural orthostatic intolerance or POTS comorbidity (see Section 10.3.3 for cardiovascular evidence)—meal timing interacts critically with activity tolerance. The post-prandial period (following meals) requires substantial splanchnic blood flow redistribution to support digestion. In dysautonomic patients, this physiological demand competes with the need to maintain perfusion to the brain and working organs during concurrent cognitive or postural activities.

Post-Prandial Splanchnic Demand. In healthy individuals, meals trigger 10–30% increase in splanchnic blood flow to support gastric emptying, intestinal motility, and nutrient absorption [336]. In POTS and severe ME/CFS patients, this post-prandial demand may not be met without compromising cardiovascular stability or cerebral perfusion, based on the autonomic dysfunction documented in these populations (see Section 10.3.3). The clinical result is frequently exacerbated symptoms: dizziness, fatigue, cognitive fog, or gastrointestinal distress occurring during or 1–2 hours after meals.

~ Hypothesis 1: Activity-Adjusted Meal Timing for Dysautonomic ME/CFS

Certainty: 0.50. These recommendations are mechanistically grounded in POTS physiology and inferred from the splanchnic blood flow changes documented in severe ME/CFS [336], but direct clinical validation in severe ME/CFS populations is lacking.

Practical Recommendations.

- **Avoid cognitive work during meals:** Reading, problem-solving, or emotionally demanding conversation should be deferred; eating should be a passive, non-demanding activity
- **Rest after meals:** Lying supine or semi-recumbent for 30–60 minutes after eating reduces the postural demand on compensatory mechanisms and supports effective digestion
- **Small, frequent meals:** Distributing food intake throughout the day (5–6 small meals) rather than 3 large meals reduces peak post-prandial splanchnic demand
- **Supine positioning during eating:** Where practical and safe, consuming meals while lying semi-recumbent or with head elevated can reduce orthostatic stress
- **Wheat-specific timing:** If trialing wheat elimination, monitor post-prandial symptoms specifically. Wheat meals may show larger post-prandial symptom exacerbation (due to zonulin-mediated barrier stress under ischemic conditions) compared to wheat-free meals of equivalent caloric content

Integration with Pacing: Recognize that meals constitute a form of physiological “activity” requiring metabolic and autonomic resources. Budget meal-related energy demands into daily activity planning, scheduling rest periods to coincide with post-prandial periods.

Individual responses vary widely; objective symptom tracking (postprandial fatigue, cognitive fog, or gastrointestinal symptoms at 0.5, 1, 2, and 4 hours post-meal) is recommended to identify optimal meal patterns.

23.2 Sleep Optimization

Sleep disturbance is nearly universal in ME/CFS and contributes directly to symptom burden. Unlike primary insomnia, ME/CFS sleep dysfunction involves disrupted sleep architecture (reduced slow-wave sleep, alpha intrusion into NREM sleep) that produces unrefreshing sleep regardless of duration. Optimizing sleep conditions can meaningfully improve daytime function even when the underlying sleep pathology persists. For pharmacological sleep management, see Section 16.3.

23.2.1 Bedroom Environment

The sleep environment should minimize sensory stimulation and support temperature regulation:

- **Darkness:** Blackout curtains or a sleep mask. Even dim light exposure during sleep suppresses melatonin and disrupts circadian rhythm. Cover or remove LED indicators on electronic devices
- **Temperature:** Maintain 18–20°C. ME/CFS patients with autonomic dysfunction often have impaired thermoregulation; cooling mattress pads and breathable bedding help compensate
- **Noise:** White noise machines or earplugs for patients with sound sensitivity. Consistent background noise is preferable to intermittent disturbances
- **Air quality:** HEPA air purifiers reduce allergen load that may worsen sleep through nasal congestion. Adequate ventilation without cold drafts
- **Screen-free zone:** Remove televisions, tablets, and smartphones from the bedroom. If a phone is needed for medical monitoring, use “do not disturb” mode with red-shifted display

23.2.2 Sleep Schedule Consistency

Circadian rhythm disruption is common in ME/CFS and contributes to the delayed sleep phase pattern seen in many patients:

- **Fixed wake time:** A consistent morning wake time (within 30 minutes) is the single most powerful circadian anchor, more important than a fixed bedtime
- **Gradual schedule shifting:** Patients with severely delayed sleep phase should advance their schedule by 15 minutes every 3–5 days rather than attempting abrupt changes
- **Avoid compensatory sleeping:** Sleeping late after a bad night perpetuates the cycle. Brief daytime rest (20–30 minutes, before 14:00) is preferable to extended morning sleep
- **Napping guidelines:** Short naps (20–30 minutes) can restore function without disrupting nighttime sleep. Naps longer than 45 minutes or after 15:00 may worsen sleep onset

23.2.3 Light Exposure Management

Light is the most potent circadian zeitgeber and can be strategically used to improve sleep timing:

- **Morning bright light:** 10,000 lux light therapy box or natural daylight within 30 minutes of waking, for 20–30 minutes. Positions the circadian clock to promote earlier sleep onset. Patients who cannot tolerate bright light due to photosensitivity may benefit from shorter exposures or lower-intensity dawn simulation
- **Evening light restriction:** Dim lights and use amber-tinted glasses or blue light filters on screens beginning 2 hours before bedtime. Blue wavelengths (460–480 nm) are the most potent melatonin suppressors
- **Red/amber lighting:** Use warm-toned night lights for nighttime bathroom trips to avoid disrupting melatonin production

23.2.4 Evening Routines

A consistent pre-sleep routine signals the transition from wakefulness to sleep:

- **Wind-down period:** 30–60 minutes of low-stimulation activities (gentle music, audio-books, light reading). Avoid emotionally stimulating content, news, or social media
- **Warm bath or shower:** A warm bath 1–2 hours before bed raises core body temperature; the subsequent cooling promotes sleep onset. A shower chair reduces the physical exertion
- **Relaxation practices:** Guided relaxation recordings, breathing exercises, or body scan meditation can reduce the hyperarousal that interferes with sleep initiation in ME/CFS
- **Avoid cognitive stimulation:** Problem-solving, planning, or worrying close to bedtime activates the prefrontal cortex and delays sleep. A “worry journal” earlier in the evening can offload anxious thoughts

23.3 Dietary Approaches

Nutritional status influences immune function, mitochondrial efficiency, gut barrier integrity, and neurotransmitter synthesis—all systems compromised in ME/CFS. While no single diet has been validated in randomized controlled trials for ME/CFS, evidence-based nutritional principles can support the management of multiple pathophysiological domains simultaneously.

23.3.1 General Nutritional Principles

The following principles apply broadly to ME/CFS patients regardless of specific dietary pattern:

- **Nutrient density:** Prioritise whole, minimally processed foods that deliver maximum micronutrients per calorie. ME/CFS patients with reduced appetite or energy for food preparation are at particular risk of micronutrient deficiency
- **Anti-inflammatory emphasis:** Emphasize omega-3 fatty acids (fatty fish, walnuts, flaxseed), colorful fruits and vegetables (polyphenols, flavonoids), and herbs and spices (turmeric, ginger) with documented anti-inflammatory properties. Minimize pro-inflammatory foods: refined sugars, trans fats, processed meats, and excessive omega-6 fatty acids
- **Adequate protein:** 1.0–1.2 g/kg/day to prevent muscle wasting in patients with limited activity. Distribute protein across meals to maximize muscle protein synthesis
- **Hydration:** 2–3 liters of fluid daily, critical for blood volume maintenance in patients with orthostatic intolerance. Electrolyte-supplemented fluids (oral rehydration salts or electrolyte powders) are more effective than water alone
- **Micronutrient adequacy:** Regular monitoring and correction of vitamin D, B12, folate, iron, magnesium, and zinc—all commonly deficient in ME/CFS and all relevant to mitochondrial function, immune regulation, and neurotransmitter synthesis (Chapter 22)

23.3.2 Specific Dietary Patterns

Several dietary patterns have theoretical relevance to ME/CFS, though none has been rigorously tested in this population:

- **Mediterranean diet:** Rich in anti-inflammatory compounds (olive oil polyphenols, omega-3 fatty acids, flavonoids), with demonstrated benefits for systemic inflammation, cardiovascular health, and cognitive function in other populations. Often considered by clinicians as a reasonable starting point for ME/CFS patients seeking an anti-inflammatory dietary pattern, though no ME/CFS-specific trial data exist
- **Low-histamine diet:** Relevant for the subgroup with mast cell activation syndrome (MCAS) or histamine intolerance (Section 23.3.4). Restricts aged, fermented, and high-histamine foods (aged cheese, wine, fermented meats, canned fish). Response within 2–4 weeks suggests histamine involvement
- **Low-FODMAP diet:** May benefit the 50–60% of ME/CFS patients with concurrent IBS symptoms. Reduces fermentable carbohydrates that cause gas, bloating, and diarrhea. Should be implemented with dietitian guidance through elimination and structured reintroduction phases
- **Elimination diets:** Systematic removal and reintroduction of suspected trigger foods (dairy, gluten, soy, eggs) can identify individual sensitivities. Requires careful nutritional planning to avoid deficiencies during elimination phases
- **Ketogenic or low-carbohydrate diets:** Theoretical interest based on the metabolic shift away from glycolysis observed in ME/CFS. Some patients report improved cognitive function and energy stability. However, the ketogenic adaptation period (“keto flu”) can trigger PEM, and long-term safety in ME/CFS has not been studied. Not recommended without medical supervision

⚠ Limitation 1: Dietary Evidence in ME/CFS

No randomized controlled trial has demonstrated efficacy of any specific dietary pattern for ME/CFS. Current recommendations are extrapolated from mechanistic rationale, evidence in related conditions (fibromyalgia, IBS, autoimmune disease), and clinical experience. Individual responses vary substantially.

23.3.3 Meal Timing and Frequency

Meal timing interacts with autonomic function and energy metabolism in ME/CFS (see also Hypothesis 23.1.3):

- **Small, frequent meals:** Five to six small meals per day reduce post-prandial splanchnic blood flow demands, minimizing orthostatic stress and post-meal fatigue. Large meals can trigger pronounced post-prandial symptoms in dysautonomic patients
- **Blood sugar stability:** Avoid refined carbohydrates that produce rapid glucose spikes followed by reactive hypoglycemia. Combine carbohydrates with protein and fat to

slow absorption. Some patients benefit from continuous glucose monitoring to identify glycemic patterns

- **Intermittent fasting:** Despite theoretical interest (autophagy activation, anti-inflammatory effects), intermittent fasting is generally inadvisable in ME/CFS. Prolonged fasting can worsen orthostatic intolerance (reduced blood volume), hypoglycemia, and energy availability in patients already operating at metabolic limits. If attempted, very mild time-restricted eating (12–14 hours overnight) is the safest approach
- **Pre-sleep nutrition:** A small protein-containing snack before bed may prevent nocturnal hypoglycemia and improve sleep quality in some patients

23.3.4 Food Sensitivities

Wheat Sensitivity and Post-Exertional Malaise

A subset of ME/CFS patients (approximately 15% based on non-celiac wheat sensitivity biomarker studies [582]) reports that wheat consumption exacerbates exercise intolerance and PEM severity. The underlying mechanism likely involves a synergistic interaction between wheat-induced gut barrier dysfunction and exercise-induced intestinal ischemia.

Speculation 66 (Wheat-Primed Exercise Intolerance in ME/CFS). Mechanistic Hypothesis:

Wheat exposure primes the intestinal barrier toward failure through two distinct pathways:

1. **Gliadin-Mediated Permeability:** Wheat gluten (gliadin peptides) activates CXCR3 signaling in intestinal epithelial cells, triggering zonulin release and tight junction disruption [583]. This increases paracellular permeability even at subclinical levels.
2. **Amylase-Trypsin Inhibitor (ATI) Inflammation:** Wheat ATIs activate toll-like receptor 4 (TLR4) on intestinal epithelial cells and innate immune cells, inducing pro-inflammatory cytokine release (TNF- α , IL-6, IL-8) [387] and further compromising barrier function.

This wheat-primed state creates a vulnerable intestinal epithelium with increased baseline permeability and systemic inflammation.

Exercise then delivers a secondary insult:

1. **Splanchnic Hypoperfusion:** Intense exercise diverts blood flow from the splanchnic circulation to working muscles, causing intestinal ischemia lasting 20–60 minutes [337].
2. **Additive Barrier Failure:** Exercise-induced ischemia triggers epithelial cell damage (elevated intestinal fatty acid-binding protein, I-FABP) [355] and increases permeability in an already-compromised wheat-primed barrier.
3. **Endotoxin Translocation:** Increased permeability allows bacterial lipopolysaccharide (LPS) to cross into systemic circulation (endotoxemia), triggering robust toll-like receptor 4 activation and systemic inflammatory cascade.

4. **Energy Crisis Amplification:** Systemic inflammation intensifies mitochondrial dysfunction and ATP depletion characteristic of PEM, resulting in exaggerated post-exertional symptoms and prolonged recovery.

Certainty Assessment:

Individual mechanistic links have strong support:

- Gliadin increases intestinal permeability: Certainty 0.75
- ATIs trigger TLR4 inflammation: Certainty 0.75
- Exercise causes splanchnic hypoperfusion and intestinal injury: Certainty 0.85
- Wheat sensitivity relevant in ME/CFS subsets: Certainty 0.60

However, the **synergistic amplification of PEM by wheat+exercise** has never been directly tested in ME/CFS populations. This mechanistic model is *highly speculative*, inferred from separate pathways that have been independently validated but never tested in combination. Overall model certainty: **0.35 (medium-low)**. Individual mechanistic links are well-established (0.75–0.85), but the critical synergistic interaction substantially reduces confidence in the integrated model. *Note: Certainty values represent ordinal confidence categories rather than precise probabilities; these assessments reflect degree of evidence and agreement among available studies.*

Critical Confounding: Fructans vs. Gluten

A major evidence limitation: Wheat contains both gluten AND fermentable fructans (FODMAPs). A high-quality double-blind randomized controlled trial (Skodje et al. 2018, n=59) demonstrated that fructans (2.1 g/day) significantly induced gastrointestinal symptoms (p=0.049 vs placebo), while gluten (5.7 g/day) did not [584].

Implication: Many cases of patient-attributed wheat sensitivity are actually fructan sensitivity or mixed FODMAP intolerance. This complicates claims about gluten-specific mechanisms and requires careful dietary control in elimination protocols (see *Clinical Implementation* section below).

Testable Predictions:

If wheat-primed exercise intolerance is valid:

- Wheat elimination (4–6 weeks) should produce clinically meaningful reduction in baseline PEM severity in affected patients (estimated 20–50%, based on typical dietary intervention responses in chronic illness populations)
- Post-elimination wheat reintroduction should trigger PEM exacerbation within 24–72 hours in responders
- Exercise tolerance (steps, activity duration before PEM) should improve measurably in responders during elimination (estimated 10–30% improvement)
- Gastrointestinal symptoms (bloating, cramping, diarrhea) should improve significantly
- Biomarkers (zonulin, LPS, I-FABP, inflammatory cytokines) should normalize during wheat elimination in responders

Limitations:

- No randomized controlled trials of wheat elimination in ME/CFS (evidence gap)
- Synergistic effect inferred from component mechanisms, not directly tested
- High individual variability in wheat sensitivity (affects approximately 15% of ME/CFS cohorts per Uhde et al. 2018 [582])
- Fructan confounding severe and poorly characterized in ME/CFS population
- FODMAP intolerance highly prevalent in ME/CFS; distinguishing gluten vs. FODMAP effects requires careful dietary control
- Temporal washout period (symptom improvement timeline) not well characterized; likely 2–4 weeks based on general elimination diet literature
- Placebo and nocebo effects are significant in dietary interventions and cannot be controlled without double-blind protocols (e.g., wheat hidden in capsules), limiting confidence in patient-reported outcomes (PEM severity, GI symptoms)
- Wheat elimination often coincides with reduced consumption of processed foods, confounding attribution of symptom improvements to wheat removal versus overall dietary quality improvement

Treatment Implications:

For patients reporting wheat-exacerbated exercise intolerance, a structured 4–6 week elimination trial with proper FODMAP control and objective monitoring may identify responders:

- **Phase 1 (Weeks 1–6): Elimination** — Remove wheat, barley, rye. Simultaneously adopt low-FODMAP diet (eliminate onions, garlic, legumes, high-fructose fruits, dairy). Maintain daily symptom diary and step count monitoring.
- **Phase 2 (Weeks 7–8): Reintroduction** — Reintroduce wheat (2 slices bread/day or equivalent) while maintaining low-FODMAP restriction. Monitor for symptom recurrence within 24–72 hours. Optionally perform submaximal exercise on Day 3 post-reintroduction to assess exercise-specific response.
- **Phase 3 (Week 9+): Decision** — If improved during elimination + worsened on wheat rechallenge, continue wheat-free diet. If no change, reintroduce wheat. If ambiguous: (1) consider pure gluten challenge (using wheat gluten isolate such as seitan or vital wheat gluten powder, fructan-free) to isolate gluten vs. fructan effects; or (2) alternatively, reintroduce FODMAPs while maintaining wheat elimination to test whether FODMAPs, rather than wheat, drive symptoms. Consult dietitian for nutritional adequacy if elimination exceeds 8 weeks.

Expected outcomes in responders: PEM reduction 20–50%, improved exercise tolerance, GI symptom resolution.

Clinical Implementation: Wheat Elimination Protocol

Diagnostic Approach Before committing to long-term dietary restriction, confirm wheat sensitivity via structured elimination-reintroduction trial:

- **Symptom baseline:** Record PEM severity (0–10 scale), daily step count, GI symptoms (bloating, cramping, diarrhea) for 1–2 weeks pre-elimination

- **Elimination period:** 4–6 weeks strict wheat/gluten avoidance (see Food List below)
- **Symptom re-assessment:** Rate PEM, steps, GI symptoms. Expect improvements within 2–4 weeks if responsive
- **Rechallenge:** Reintroduce wheat. Monitor for symptom recurrence within 24–72 hours. If symptoms return, diagnosis supported
- **Optional objective endpoint:** Two-day CPET pre/post elimination if available; responders may show improved Day 2 preservation in work output

Contraindications Do not attempt this elimination trial without medical evaluation if: (1) celiac disease suspected (requires serology and biopsy confirmation before dietary restriction); (2) history of eating disorders or restrictive eating patterns (risk of exacerbating disordered behaviors); (3) severe malnutrition or unintended weight loss (wheat elimination may worsen nutritional status). Consult physician before starting any dietary elimination trial, particularly if considering prolonged restriction (>8 weeks).

FODMAP Considerations Standard wheat elimination is complicated by FODMAP content. For patients with mixed GI symptoms (bloating, gas, diarrhea) alongside PEM:

- Adopt low-FODMAP diet simultaneously (eliminate onions, garlic, legumes, high-fructose fruits, wheat)
- If symptoms improve during low-FODMAP phase, fructan sensitivity likely primary mechanism
- After 4–6 weeks, selectively reintroduce high-FODMAP foods while keeping wheat eliminated
- If GI symptoms return with high-FODMAP reintroduction, FODMAP intolerance confirmed (separate from gluten-specific effects)

Practical Food Substitutions Wheat-free alternatives (assuming low-FODMAP compliance):

- **Bread:** Rice bread, certified gluten-free bread (avoid high-fiber versions if GI intolerance)
- **Pasta:** Rice pasta, corn pasta, quinoa pasta
- **Cereals:** Rice, quinoa, corn, oats (if tolerated)
- **Thickeners:** Cornstarch, rice flour, potato starch (instead of wheat flour)
- **Baked goods:** Almond flour, coconut flour, rice flour blends (commercial gluten-free mixes)

Common Pitfalls

- **Incomplete elimination:** Wheat hidden in sauces, processed foods, cross-contamination. Requires label reading and food preparation knowledge.
- **Inadequate duration:** Symptom improvement may require 3–4 weeks. Premature cessation may miss benefit.

- Concurrent high-FODMAP intake: High-fructose foods may obscure gluten-specific effects. Combined low-FODMAP + wheat-free trial recommended.
- Nutritional deficiency risk: Wheat provides fiber, B vitamins, minerals. Ensure adequate replacements (brown rice, quinoa, vegetables).
- Nocebo effect: Expectation of benefit may bias subjective symptom reporting. Objective tracking (steps, PEM severity scale) essential.

See Chapter 11 for mechanistic details on gut permeability and zonulin pathways. See Section 2.1 for PEM trigger mechanisms and the role of endotoxemia in post-exertional inflammation.

23.4 Exercise and Movement

23.4.1 The Exercise Paradox

Why Standard Exercise Programs Fail in ME/CFS

Exercise is beneficial for most chronic conditions and healthy populations, improving cardiovascular fitness, strength, mood, and metabolic health. However, ME/CFS represents a notable exception where standard exercise physiology does not apply.

Normal Exercise Adaptation Healthy individuals respond to exercise training with:

- Improved mitochondrial density and function
- Enhanced cardiovascular capacity
- Increased muscle strength and endurance
- Positive mood effects (endorphin release, reduced depression)
- Progressive tolerance of higher workloads

Pathological Exercise Response in ME/CFS ME/CFS patients instead experience:

- Worsening symptoms following exertion (PEM)
- No adaptive improvement with repeated exercise
- Measurable physiological deterioration (documented by two-day CPET)
- Cumulative functional decline with sustained exercise programs
- Prolonged recovery periods (days to weeks) after single exertional episodes

This fundamental difference reflects underlying metabolic dysfunction rather than deconditioning or psychological factors.

Graded Exercise Therapy (GET): Controversy and Evidence of Harm

Graded exercise therapy—progressive incremental increases in physical activity—was historically recommended for ME/CFS based on the assumption that symptoms reflected deconditioning, fear avoidance, or deconditioning-related fatigue. This assumption has been decisively refuted by objective evidence.

The PACE Trial and Subsequent Reanalysis The 2011 PACE trial initially claimed benefits from GET and cognitive behavioral therapy (CBT). However, subsequent reanalysis using objective outcomes (rather than subjective questionnaires) found [506, 57]:

- No significant improvement in objective measures (6-minute walk distance, step counts, employment, benefits claims)
- High rates of patient-reported harm in long-term follow-up
- Methodological concerns including subjective outcomes, non-blinded assessments, and changing outcome definitions

Major health authorities have since revised guidelines to **recommend against GET** for ME/CFS, including NICE (UK, 2021), CDC (USA, 2022), and others [127].

Not all national bodies have followed suit. In Australia, the Royal Australian College of General Practitioners (RACGP) maintained a HANDI recommendation for “incremental physical activity” in ME/CFS until it was formally challenged in 2026. Stallard et al. demonstrated that the guideline scored only 2% on the AGREE II rigor domain, relied on the methodologically contested PACE trial and an outdated Cochrane review, and ignored patient surveys reporting 54–74% harm rates [585]. The authors called for immediate withdrawal and contraindication of GET pending the NHMRC’s review of ME/CFS guidelines, recommending that Australian general practitioners refer to NICE and BMJ Best Practice guidelines in the interim.

Two-Day CPET Evidence Against GET Objective physiological evidence demonstrates why GET is contraindicated. Keller et al. (2024) showed that even a single maximal exertion produces [1]:

- **Day 2 performance decrements:** 5–8% declines in VO_2peak , work output, ventilation
- **Worsening impairment classification:** Severe impairment cases nearly doubled (14% → 27%)
- **Independence from fitness:** Abnormal responses persisted when matched for baseline aerobic capacity
- **Prolonged recovery:** Full restoration requiring 13+ days versus ~2 days in controls

These findings demonstrate that exertion **impairs rather than improves physiological function** in ME/CFS. GET protocols that require repeated exertion before recovery is complete would predictably produce cumulative deterioration—precisely what patients report.

Mechanistic Understanding The two-day CPET results validate patient reports by demonstrating:

1. Exercise triggers measurable metabolic failure beyond normal fatigue or deconditioning
2. Recovery systems are impaired, requiring prolonged restoration periods
3. Repeated exertion before recovery worsens baseline function
4. The phenomenon is reproducible and objectively quantifiable

Charlton et al. (2026) reinforced this conclusion in the *British Journal of Sports Medicine*, arguing that post-exertional malaise in long COVID cannot be attributed to cardiac deconditioning and that the pathophysiology of exercise intolerance must be fundamentally reconsidered [341]. GET's failure in ME/CFS reflects accurate biology, not patient non-compliance or psychological factors.

Patient-Reported Harms

△ Warning 1: Patient-Reported Harms from GET

Large patient surveys consistently report high rates of harm from GET. A systematic review of 10 patient surveys across four countries found that 51% of respondents reported GET worsened their health, compared to 20% for CBT [586]. These figures underline why GET is no longer recommended by NICE and other major guidelines [127]:

- 51% of ME/CFS patients across surveys reported GET worsened their condition [586]
- Many report GET as triggering transition to more severe disease states
- Some report permanent functional decline attributable to GET programs
- Very few (<10%) report sustained benefit

The Comorbidity Exercise Conflict

Many ME/CFS patients have concurrent conditions for which exercise is a standard, evidence-based treatment: type 2 diabetes, cardiovascular disease, obesity, osteoporosis, hypermobility spectrum disorders (hEDS/HSD), and depression. This creates a genuine *therapeutic conflict*: guidelines for the comorbidity recommend progressive exercise, while ME/CFS pathophysiology makes such exercise harmful. Because most comorbidity-specific guidelines do not include ME/CFS exceptions, clinicians are left to reconcile contradictory recommendations without formal guidance.

The conflict is not hypothetical. In France and other countries where ME/CFS recognition lags behind international consensus, patients continue to receive exercise-based rehabilitation prescriptions from physicians treating their comorbidities—unaware that the standard protocol may trigger post-exertional malaise, functional decline, and disease progression [586, 127, 587].

Scope of the Conflict The following comorbidities illustrate the tension:

- **Type 2 diabetes and metabolic syndrome.** Exercise is first-line therapy for insulin resistance, yet ME/CFS patients cannot tolerate the recommended 150 min/week of moderate-intensity aerobic activity. Obesity compounds functional impairment in ME/CFS, and standard exercise-based weight management is contraindicated by PEM [588]. Metabolomic evidence shows ME/CFS involves a distinct hypometabolic state with impaired pyruvate dehydrogenase function [589], fundamentally different from the metabolic profile of sedentary obesity.
- **Cardiovascular disease.** Standard cardiac rehabilitation relies on progressive aerobic exercise. However, ME/CFS patients frequently have reduced stroke volume, reduced end-diastolic volumes, and reduced cardiac mass on MRI, with a significant inverse relationship between cardiac output and PEM severity [590]. Applying standard cardiac rehabilitation without accounting for post-exertional physiology risks worsening both the cardiac condition and ME/CFS.
- **Hypermobility spectrum disorders (hEDS/HSD).** Muscle strengthening is essential to stabilize hypermobile joints and reduce dislocations. Resistance training with body weight and resistance bands is feasible and effective in hEDS/HSD [591], and isometric exercises are particularly valuable because they build strength without joint movement [592]. However, no published study addresses the dual hEDS+ME/CFS population, and even low-load resistance protocols must be titrated against PEM risk.
- **Osteoporosis.** Weight-bearing exercise is recommended to maintain bone density, but many ME/CFS patients—especially those who are housebound or bedbound—cannot perform such exercise. Prolonged immobility itself accelerates bone loss, creating a vicious cycle. Alternative approaches (e.g., whole-body vibration platforms that can be used in supine position) exist in the general osteoporosis literature but have not been evaluated in ME/CFS.
- **Depression.** Exercise is a first-line non-pharmacological treatment for depression. However, in ME/CFS, exercise-based interventions for depression may exceed the patient's energy limit and trigger characteristic PEM, undermining any antidepressant effect [593]. Pacing combined with psychotherapy (CBT adapted for ME/CFS, acceptance-based therapies) should be the default approach when depression is comorbid.

The CPET-Based Activity Prescription Framework The resolution of the comorbidity exercise conflict lies in *individualized, physiology-guided activity prescription* rather than population-based exercise guidelines. The Workwell Foundation's two-day cardiopulmonary exercise test (CPET) protocol provides the scientific basis for this approach [594, 595].

Davenport et al. (2010) proposed that all physical therapy for ME/CFS must center on the ventilatory anaerobic threshold (VAT) determined by individual CPET [595]. Activity must stay below the patient's VAT heart rate at all times. This model applies equally when the goal is comorbidity management: the *type* of exercise may be dictated by the comorbidity (e.g., resistance training for hEDS, aerobic activity for diabetes), but the *intensity ceiling* is always set by the ME/CFS pathophysiology.

Critically, formula-derived heart rate targets (e.g., $220 - \text{age} \times 0.6$) do not reliably predict

the individually measured lactic acidosis threshold in ME/CFS patients. Van Campen et al. (2020) demonstrated bias ranging from -28 to $+23$ bpm between formula estimates and CPET-measured thresholds in 90 ME/CFS adults [596]. Population formulas are therefore inadequate for safe activity prescription; individual CPET-derived thresholds are necessary.

Heart rate monitor (HRM)-guided pacing operationalizes this approach in daily life. Clague-Baker et al. (2025) demonstrated that HRM-guided pacing is feasible and acceptable in both ME/CFS and long COVID: 89% of participants continued using their HRM at 8 weeks, and 66% at 6 months [597].

Severity-Stratified Approach Gloeckl et al. (2024) proposed a three-tier framework for exercise in post-viral conditions with PEM that is directly applicable to ME/CFS comorbidity management [598]:

1. **No PEM (rare in ME/CFS):** Standard comorbidity-specific exercise guidelines may apply, with heart rate monitoring as a safety net.
2. **Mild-to-moderate PEM:** Modified exercise tailored to individual tolerance. The comorbidity dictates exercise type; the VAT heart rate sets the intensity ceiling. Short sessions (5–15 min), sub-anaerobic intensity, with 48–72 h monitoring for delayed PEM.
3. **Severe PEM:** Pacing strategies only. No structured exercise program. Comorbidity management must rely on pharmacological and non-exercise interventions (e.g., insulin sensitizers for diabetes, bisphosphonates for osteoporosis, psychotherapy for depression).

A systematic review of physiotherapy evidence in ME/CFS reinforces this stratification: Wormgoor and Rodenburg (2021) found that any apparent treatment effect of exercise disappeared when stricter PEM-inclusive diagnostic criteria and objective outcome measures were applied [599]. This confirms that exercise interventions validated in broader “chronic fatigue” populations cannot be extrapolated to patients meeting full ME/CFS criteria with documented PEM.

Practical Adaptations by Comorbidity When a comorbidity requires some form of physical activity and the patient falls into the mild-to-moderate PEM tier, the following adaptations apply:

- **For diabetes/metabolic syndrome:** Substitute prolonged aerobic sessions with multiple daily micro-sessions (3–5 min walking or recumbent cycling) below VAT heart rate. Prioritise dietary management and pharmacotherapy (metformin, GLP-1 agonists) as primary interventions, using activity as an adjunct only when tolerated.
- **For cardiovascular rehabilitation:** Replace standard progressive protocols with CPET-guided sub-anaerobic activity. Recumbent positions reduce orthostatic demand. Monitor both cardiac parameters and PEM onset; cardiac benefit should never be pursued at the cost of post-exertional deterioration.
- **For hEDS/HSD:** Focus on isometric contractions (5–10 s holds at 30–50% maximal effort) and low-load resistance band work, which have lower cardiovascular demand than

dynamic exercise [592]. Perform in supine or seated position. Target specific joint stabilization rather than general fitness.

- **For osteoporosis:** When weight-bearing exercise is impossible, consider pharmacological bone protection (bisphosphonates, vitamin D, calcium). For patients who tolerate brief standing, supported standing exercises (wall-assisted) for 1–2 min may provide some mechanical stimulus without exceeding energy limits.
- **For depression:** Replace exercise-based antidepressant strategies with adapted psychological interventions (acceptance and commitment therapy, mindfulness-based approaches), pharmacotherapy, and gentle movement within the energy envelope. Social connection and meaningful activity within pacing constraints offer mood benefits without PEM risk.

? Open Question 1: Exercise Adaptation for ME/CFS Comorbidities

No randomized controlled trial has evaluated exercise protocols specifically adapted for ME/CFS patients with concurrent conditions requiring physical activity (type 2 diabetes, cardiovascular disease, hEDS, osteoporosis, depression). The evidence base consists of extrapolation from separate ME/CFS pacing research, comorbidity-specific exercise literature, and expert consensus. Clinical trials testing CPET-guided, PEM-monitored exercise protocols for these specific dual-diagnosis populations are urgently needed.

Risk of Post-Exertional Malaise

Any movement carries PEM risk in ME/CFS, necessitating careful calibration:

- **Dose-response relationship:** Greater exertion produces worse PEM
- **Individual variability:** Thresholds vary widely (severe patients may crash from showering; mild patients tolerate gentle walks)
- **Cumulative effects:** Multiple small exertions may sum to trigger PEM
- **Unpredictable triggers:** Same activity may be tolerated one day but trigger PEM another day
- **Delayed onset:** 12–72 hour lag makes cause-effect connections difficult

23.4.2 Safe Movement Approaches

Despite exercise intolerance, complete immobility causes problems (muscle atrophy, joint stiffness, orthostatic intolerance worsening). The goal is **movement within the energy envelope**—enough to prevent deconditioning complications without triggering PEM.

Principles of Safe Movement

- **Stay below anaerobic threshold:** Use heart rate monitoring (AT – 10–15 bpm)
- **Horizontal postures:** Recumbent or supine exercise reduces orthostatic demand

- **Short duration:** 5–10 minute sessions may be tolerable where 20–30 minutes would crash
- **Consistency over intensity:** Very gentle daily movement better than intermittent harder sessions
- **Immediate cessation:** Stop at first signs of excessive exertion (heart rate elevation, breathlessness, fatigue)
- **Monitor delayed effects:** Track PEM onset 12–72 hours post-activity to calibrate appropriately

Gentle Stretching

- **Supine or seated:** Reduces cardiovascular demand
- **Passive range of motion:** Maintain joint mobility without resistance
- **Avoid ballistic movements:** Gentle, sustained stretches only
- **Duration:** 5–15 minutes may be tolerable
- **Daily frequency:** If tolerated, maintains flexibility

Isometric Exercises

Isometric (static muscle contraction without joint movement) may be better tolerated than dynamic exercise:

- **Lower cardiovascular demand:** Minimal heart rate elevation
- **Maintain muscle strength:** Prevents complete atrophy
- **Short holds:** 5–10 second contractions
- **Submaximal intensity:** Moderate contraction only (30–50% maximal)
- **Examples:** Wall sits (brief), plank holds (modified), leg presses against bed

Recumbent Activities

Horizontal or semi-reclined positions reduce orthostatic stress:

- **Recumbent bike:** Allows cardiovascular activity with lower orthostatic demand
- **Supine leg movements:** Gentle cycling motions while lying down
- **Pool exercises:** Buoyancy reduces gravitational stress (if tolerated; some patients worsen in water)
- **Resistance bands while seated:** Low-impact strength maintenance

Monitoring for PEM

Vigilant monitoring prevents inadvertent overexertion:

- **Real-time heart rate:** Stop if approaching threshold
- **Perceived exertion:** Use modified Borg scale; stop at first sense of effort
- **Post-activity tracking:** Log symptoms 12–72 hours after movement
- **Adjust based on outcomes:** If PEM occurs, reduce intensity/duration for subsequent sessions
- **Recovery time:** Allow full recovery (minimum 24–48 hours, often longer) between sessions

Adaptive Progression (If Tolerated)

For patients with stable mild-to-moderate ME/CFS who tolerate current activity levels without PEM:

- **Very gradual increases:** 1–2 minutes per week, or 1 additional repetition per week
- **Sustained tolerance required:** Maintain new level for 2–4 weeks before further increase
- **Immediate rollback if PEM occurs:** Return to previous tolerated level
- **Never push through PEM:** This worsens condition and should be avoided absolutely
- **Realistic expectations:** Goal is maintaining current function, not fitness improvement

△ Warning 2: Exercise Precautions

Patients with severe ME/CFS (housebound or bedbound) should consult physicians before attempting any structured movement program. Even minimal exertion may trigger severe crashes in this population. For these patients, activities of daily living (personal hygiene, eating) may constitute maximal tolerable exertion, leaving no additional capacity for exercise.

23.5 Stress Management

Psychological stress activates the hypothalamic–pituitary–adrenal (HPA) axis and sympathetic nervous system, both of which are dysregulated in ME/CFS (Chapter 8). Stress increases pro-inflammatory cytokine production, worsens sleep, and can directly trigger PEM. Stress management is therefore not a psychological luxury but a physiological intervention that reduces the allostatic load on already-compromised systems.

23.5.1 Relaxation Techniques

Structured relaxation practices reduce sympathetic activation and may improve autonomic balance:

- **Diaphragmatic breathing:** Slow, deep abdominal breathing (4–6 breaths per minute) activates the vagus nerve and shifts autonomic balance toward parasympathetic dominance. Can be practiced lying down with minimal energy expenditure. Particularly helpful for patients with hyperventilation or orthostatic hypocapnia
- **Progressive muscle relaxation (PMR):** Systematic tensing and releasing of muscle groups. The “release only” variant (focusing on relaxation without tensing) is preferred for ME/CFS patients, as even brief isometric contractions may trigger PEM in severe cases
- **Guided imagery:** Audio-guided visualization of calming scenes. Requires minimal physical energy and can be used during rest periods. Available as smartphone apps or recordings designed for chronic illness populations
- **Autogenic training:** Self-directed relaxation using verbal cues for warmth and heaviness in the limbs. May improve peripheral circulation and reduce autonomic hyperreactivity

23.5.2 Meditation and Mindfulness

Mindfulness-based approaches have demonstrated efficacy for chronic pain, fatigue, and psychological distress in related conditions (fibromyalgia, chronic pain), though ME/CFS-specific evidence is limited:

- **Adapted mindfulness:** Standard mindfulness-based stress reduction (MBSR) programs require 45 minutes of daily practice and regular group attendance, which exceeds the capacity of many ME/CFS patients. Adapted versions using 5–15 minute sessions, audio guidance, and remote participation are more appropriate
- **Body scan meditation:** A lying-down practice that cultivates awareness of bodily sensations without attempting to change them. May help patients develop a more accepting relationship with symptoms and reduce the distress amplification that accompanies chronic illness
- **Acceptance and commitment therapy (ACT):** A mindfulness-based psychotherapy particularly suited to ME/CFS because it focuses on living meaningfully within limitations rather than striving to eliminate symptoms. Preliminary evidence suggests benefit for quality of life in chronic fatigue conditions
- **Cautions:** Meditation can occasionally increase distress in patients with trauma histories or severe anxiety. Concentration-based practices may trigger cognitive PEM. Patients should start with very brief sessions and use guided approaches rather than unstructured meditation

23.5.3 Biofeedback

Biofeedback provides real-time physiological feedback that patients can use to learn autonomic regulation:

- **Heart rate variability (HRV) biofeedback:** Patients learn to increase HRV through resonance frequency breathing (typically 5–7 breaths per minute). Reduced HRV is a consistent finding in ME/CFS and correlates with symptom severity. HRV biofeedback has shown benefit in fibromyalgia and warrants investigation in ME/CFS. Home-based devices make this accessible for housebound patients
- **Temperature biofeedback:** Learning to increase peripheral hand temperature through relaxation. Useful for patients with Raynaud’s phenomenon or cold extremities from autonomic dysfunction
- **Electrodermal activity:** Monitoring skin conductance as a proxy for sympathetic activation. Can help patients identify and reduce stress responses they may not consciously recognize

23.6 Environmental Modifications

Environmental optimization reduces the baseline energy expenditure of daily living and minimizes symptom triggers. For severely affected patients, environmental modifications may be among the most impactful and cost-effective interventions available.

23.6.1 Home Adaptations

The home environment should be configured to minimize physical demands and sensory stressors:

- **Reducing physical demands:** Single-floor living where possible. Stairlift or bedroom relocation to ground floor if stairs trigger PEM. Shower chair, handheld showerhead, grab bars. Electric can openers, jar openers, and lightweight cookware. Voice-activated home assistants for lights, temperature, and communication
- **Lighting:** Adjustable lighting throughout the home. Dimmer switches or smart bulbs that allow intensity control. Many ME/CFS patients experience photosensitivity; overhead fluorescent lighting is particularly problematic. Warm-toned, indirect lighting at low intensity is generally best tolerated
- **Noise reduction:** Soft furnishings (carpets, curtains, upholstered furniture) absorb sound. Double glazing reduces external noise. White noise machines provide consistent background sound that masks intermittent disturbances. Noise-cancelling headphones for use during sensory overload episodes
- **Temperature control:** Programmable thermostat maintaining 20–22°C. Ceiling fans for air circulation without cold drafts. Heated mattress pad for patients with cold intolerance. Portable air conditioning unit for the primary living/resting space if central cooling is unavailable

23.6.2 Chemical and Environmental Sensitivities

Multiple chemical sensitivity (MCS) is common in ME/CFS and may reflect mast cell activation, neuroinflammation, or impaired xenobiotic metabolism:

- **Reducing chemical exposures:** Switch to fragrance-free personal care products, laundry detergents, and cleaning agents. Avoid air fresheners, scented candles, and perfumes. Use natural cleaning alternatives (vinegar, baking soda) where effective
- **Air filtration:** HEPA air purifiers in bedroom and primary living space reduce airborne allergens, particulates, and volatile organic compounds (VOCs). Activated carbon filters additionally remove chemical irritants. Replace filters per manufacturer schedule
- **Indoor air quality:** Ventilate regularly but avoid outdoor pollution peaks. Indoor plants may modestly reduce VOC levels. Avoid new furniture, carpets, or paint that off-gas VOCs; if unavoidable, ventilate extensively before patient exposure
- **Mold remediation:** Mold exposure can exacerbate ME/CFS symptoms through mycotoxins and immune activation. Professional mold assessment and remediation if water damage or visible mold is present. Maintain indoor humidity below 50% to prevent mold growth

23.7 Social and Emotional Support

ME/CFS imposes severe social isolation, particularly for moderate-to-severe patients who cannot leave their homes. Social disconnection, loss of identity, and the psychological burden of chronic disbelief from medical professionals and society compound the physical suffering. Addressing social and emotional needs is integral to comprehensive management.

23.7.1 Support Groups and Peer Connection

Peer support provides validation, practical advice, and social connection that medical professionals cannot fully substitute:

- **Online communities:** Forums, social media groups, and video-based support groups are accessible to housebound patients. Text-based communication allows participation at individual pace and energy levels. Organizations such as ME Action, #MEAAction, and national ME/CFS associations maintain moderated online communities
- **In-person groups:** For patients who can attend, local support groups provide face-to-face connection. Meetings should accommodate energy limitations (brief duration, comfortable seating, low sensory environment)
- **Peer mentoring:** Connecting newly diagnosed patients with experienced patients who have developed effective coping strategies can accelerate adaptation and reduce the learning curve for pacing, medication management, and navigating the medical system

23.7.2 Counseling and Psychotherapy

Psychotherapy in ME/CFS addresses the psychological consequences of chronic illness, not the illness itself. The distinction is critical: therapy is for coping with ME/CFS, not for “curing” it.

- **Acceptance and commitment therapy (ACT):** Focuses on living a meaningful life within the constraints of illness rather than fighting symptoms. Particularly effective for adjustment to disability and chronic pain
- **Cognitive behavioral therapy (CBT) — adapted:** CBT adapted for ME/CFS targets unhelpful thought patterns about illness (catastrophizing, all-or-nothing thinking) without implying that symptoms are psychologically caused. Critically, adapted CBT respects the energy envelope and does not encourage activity increases beyond physiological limits. Standard CBT protocols designed to increase activity (as in the PACE trial) are contraindicated
- **Grief counseling:** ME/CFS involves profound loss—of career, social roles, physical capacity, independence, and future plans. Processing this grief is essential and should not be dismissed as “depression”
- **Remote therapy:** Telephone or video therapy sessions reduce the energy cost of attending appointments. Many patients cannot tolerate in-person sessions, particularly during flares

23.7.3 Addressing Secondary Depression and Anxiety

Depression and anxiety are common in ME/CFS (prevalence 30–50%) and are typically secondary to the illness rather than causative. Treatment should address both the psychological symptoms and their ME/CFS-specific drivers:

- **Distinguish reactive from primary:** Depression arising from loss, isolation, and disbelief differs from primary major depressive disorder and may respond better to social and practical interventions than to antidepressants alone
- **Pharmacological treatment:** SSRIs (sertraline, escitalopram) or SNRIs (duloxetine, which may also benefit pain) when depression is moderate-to-severe. Start at low doses, as ME/CFS patients are frequently medication-sensitive. Avoid activating antidepressants (fluoxetine, bupropion) that may worsen insomnia or anxiety in sensitive patients
- **Address medical gaslighting:** A significant contributor to psychological distress in ME/CFS is repeated dismissal, disbelief, or misdiagnosis by medical professionals. Validation from a knowledgeable clinician can be powerfully therapeutic in itself
- **Suicide risk:** ME/CFS carries elevated suicide risk (Section 29.5), particularly in patients with severe functional limitation, medical dismissal, and social isolation. Clinicians should routinely screen for suicidal ideation and ensure access to crisis resources

23.7.4 Family and Caregiver Education

Family members and caregivers are essential partners in ME/CFS management but often lack understanding of the disease:

- **Disease education:** Provide family members with clear, evidence-based information about ME/CFS pathophysiology, PEM, and the energy envelope. Understanding that the illness is biological—not psychological or motivational—transforms the family dynamic
- **Caregiver burnout:** Caregivers of ME/CFS patients are at high risk of burnout, particularly when the patient is severe and requires assistance with all activities of daily living. Caregiver support groups, respite care, and professional home care services should be utilized where available
- **Communication strategies:** Teach patients and families to communicate about energy levels using concrete language (e.g., a 0–10 energy scale) rather than vague terms. This reduces misunderstandings about what the patient can and cannot do on a given day
- **Boundary setting:** Help families understand that encouraging the patient to “push through” or “try harder” is harmful. Support pacing decisions even when they limit family activities or social plans

24 Experimental and Emerging Therapies

This chapter explores therapies at the frontier of ME/CFS treatment—approaches with theoretical rationale but limited clinical validation. Some represent extensions of established medical science; others venture into more speculative territory. The heterogeneous nature of ME/CFS suggests that different patients may require fundamentally different interventions, making this exploratory landscape particularly relevant.

24.1 Novel Therapeutic Frameworks

Before examining specific interventions, several overarching conceptual frameworks offer novel approaches to treatment design.

~ Hypothesis 1: Metabolic State Transition

ME/CFS may represent a stable but maladaptive metabolic state—analogous to cellular “hibernation” or the evolutionarily conserved sickness behavior response that became pathologically persistent. The body entered a low-energy conservation mode in response to an initial trigger (infection, trauma, severe stress) but failed to receive or respond to the “all clear” signal to return to normal metabolism. If true, effective treatment may require interventions that trigger metabolic state transitions rather than symptom suppression. Candidate approaches include:

- Controlled metabolic stressors (fasting, hypoxia, temperature extremes) that force cellular adaptation
- Interventions targeting metabolic switching pathways (AMPK activation, mTOR modulation)
- Circadian rhythm reset protocols combining light therapy, meal timing, and temperature cues

This framework suggests that gradual, gentle interventions may perpetuate the maladaptive state, while carefully designed acute challenges might catalyze transition—though the risks of such approaches in a population with impaired stress tolerance are substantial.

~ Hypothesis 2: Cellular Danger Response Persistence

Robert Naviaux’s cell danger response (CDR) hypothesis [197] proposes that cells remain stuck in a defensive metabolic mode characterized by reduced mitochondrial function, altered purinergic signaling, and maintained inflammatory readiness. The CDR evolved as a protective response to threats, but in ME/CFS, the “threat resolved”

signal may never arrive or may not be recognized. Therapeutic implications include:

- Antipurinergic therapy (suramin showed promise in an autism pilot trial [15]; ME/CFS trials have not been conducted)
- Modulating extracellular ATP signaling through P2X/P2Y receptor antagonists
- Reducing triggers that maintain CDR activation (chronic infections, gut dysbiosis, environmental toxins)
- Flavonoids with antipurinergic properties (quercetin, luteolin) as accessible alternatives

~ Hypothesis 3: Glymphatic Dysfunction and Neuroinflammatory Persistence

Sleep in ME/CFS is characteristically non-restorative despite adequate duration. The glymphatic system—the brain’s waste clearance mechanism—operates primarily during deep sleep [476]. If glymphatic function is impaired, neuroinflammatory debris may accumulate, perpetuating microglial activation and cognitive dysfunction. Testable interventions include:

- Sleep architecture optimization targeting slow-wave sleep (when glymphatic clearance peaks)
- Sleep position modification (lateral sleeping may enhance glymphatic flow)
- Agents that improve glymphatic function (low-dose naltrexone reduces neuroinflammation; specific anesthetics enhance glymphatic clearance in animal models)
- Timing of hydration (adequate fluids without excessive evening intake)
- Omega-3 fatty acids (AQP4 water channel function depends on membrane composition)

⚠ Limitation 1: Emerging Therapies: Pre-Validation Evidence Throughout

All interventions in this chapter are experimental or unvalidated for ME/CFS. Key epistemic boundaries apply to the entire chapter:

- No intervention described below has been tested in a randomised, placebo-controlled trial specifically in ME/CFS unless explicitly stated. Open-label designs, case series, and single-arm pilots dominate the evidence base.
- The “metabolic state transition,” “cell danger response persistence,” and “glymphatic dysfunction” therapeutic frameworks are hypothesis-driven; treatment protocols derived from them are speculative extrapolations, not evidence-based protocols.
- Cross-condition extrapolation (e.g., FMT efficacy from *C. difficile* infection, non-invasive VNS from migraine, HBOT from wound healing) does not establish efficacy in ME/CFS; the shared-symptom assumption underlying such transfers may be incorrect.
- Supplement and peptide therapies (BPC-157, thymosin alpha-1) often lack human pharmacokinetic data; dosing, bioavailability, and safety profiles are incompletely characterised.

24.2 Immunological Interventions

24.2.1 Autoantibody-Targeted Therapies

Growing evidence implicates autoantibodies against G-protein coupled receptors (GPCRs) in ME/CFS pathophysiology, with particularly strong associations in post-infectious cases. The foundational study by Loebel et al. (2016) found that 29.5% of 268 ME/CFS patients had elevated autoantibodies against β_2 -adrenergic, M3 muscarinic, or M4 muscarinic receptors [71]. Subsequent validation studies by Bynke et al. (2020) found even higher prevalence (79–91% with at least one elevated autoantibody) [72], and Sotzny et al. (2021) demonstrated dose-response correlations between autoantibody levels and symptom severity [208]. However, the Vernino et al. (2022) failed replication in POTS using standard ELISA methodology raises important questions about assay specificity [210]. These therapeutic approaches target the autoantibody hypothesis directly.

BC007 (DNA Aptamer)

BC007 (originally developed for heart failure) is a DNA aptamer that directly neutralizes autoantibodies against beta-adrenergic and muscarinic receptors. Hohberger et al. (2021) reported a dramatic case in Long COVID [10]: a single 1350 mg intravenous dose neutralized GPCR autoantibodies within hours, with rapid resolution of fatigue, brain fog, and dysgeusia, plus improved retinal microcirculation on optical coherence tomography angiography. Effects were sustained at 4-week follow-up. This proof-of-concept case demonstrates that direct autoantibody neutralization can produce rapid symptomatic improvement. Larger trials are ongoing, but access remains limited to research settings.

Immunoadsorption

Immunoadsorption selectively removes immunoglobulins (including pathogenic autoantibodies) from blood plasma while returning other components. Unlike plasmapheresis, it can be targeted to specific antibody classes.

Clinical Evidence The evidence base for immunoadsorption in ME/CFS has grown substantially:

- **Pilot study (2018):** Scheibenbogen et al. treated 10 post-infectious ME/CFS patients with elevated β_2 -adrenergic receptor antibodies [147]. 70% showed rapid improvement during treatment; 30% sustained moderate-to-marked improvement at 6–12 months follow-up. This provided the first demonstration that removing autoantibodies could improve ME/CFS symptoms.
- **Prospective cohort (2025):** Stein et al. conducted a larger prospective study in 20 post-COVID ME/CFS patients with elevated β_2 -adrenergic receptor autoantibodies [118]. Five immunoadsorption sessions reduced IgG by 79% and β_2 -AR autoantibodies by 77%. 70% (14/20) were classified as responders with ≥ 10 point improvement in SF-36 Physical

Function score. Benefits were sustained to 6 months. This represents the strongest evidence to date supporting autoantibody-mediated ME/CFS pathophysiology.

Practical Considerations

- Responses lasting weeks to months suggest antibody-producing cells persist and regenerate autoantibodies
- Need for repeated treatments in most responders
- High cost (typically €5,000–15,000 per treatment course) and limited availability
- Requires specialized apheresis centers

Energy Profile. Category C (energy-demanding). Immunoadsorption requires multiple sessions of extracorporeal blood processing, placing significant cardiovascular and haemodynamic demands on the patient. The acute immune reorganization following antibody removal involves cytokine shifts and compensatory immune responses that are inherently energy-costly (proposed mechanism; precise immunological events post-immunoadsorption in ME/CFS are not fully characterized). Patients typically require 5 sessions over 5–10 days, with each session lasting several hours.

Speculation 67 (Combined Autoantibody Depletion and B-Cell Targeting). If GPCR autoantibodies drive symptoms and B cells continuously produce them, effective treatment may require both: (1) acute removal of existing autoantibodies via immunoadsorption or BC007, combined with (2) depletion of autoreactive B cells to prevent regeneration. This could explain why rituximab (B-cell depleting) showed initial promise but failed in larger trials—if circulating autoantibodies persist for months after B-cell depletion, symptom improvement would be delayed beyond trial endpoints. However, the daratumumab pilot data [9] suggest that targeting plasma cells (the actual antibody-secreting cells) may be more effective than targeting their B-cell precursors. A protocol combining immunoadsorption followed by plasma cell depletion with daratumumab, then monitoring autoantibody titers and symptoms, could test this refined hypothesis.

Daratumumab: Targeting Plasma Cells (2025 Pilot Trial)

A groundbreaking 2025 pilot study by Fluge et al. tested daratumumab, an anti-CD38 monoclonal antibody that targets plasmablasts and long-lived plasma cells—a novel approach distinct from prior B-cell targeting strategies [9].

Rationale Unlike rituximab (which targets CD20 on B cells), daratumumab depletes plasma cells that actively produce autoantibodies. The hypothesis: if GPCR autoantibodies emerge after infection and are continuously secreted by long-lived plasma cells in bone marrow or gut wall, targeting these cells directly may be more effective than depleting their B-cell precursors.

Study Design and Results

- **Participants:** 10 female patients with moderate-to-severe ME/CFS
- **Intervention:** Subcutaneous daratumumab 1800 mg (4–7 injections over 12 weeks)
- **Response rate:** 6 of 10 patients (60%) showed marked improvement
- **Physical function:** SF-36 Physical Function increased from 25.9 to 55.0 at 8–9 months ($p=0.002$)
- **Symptom burden:** DePaul Questionnaire scores dropped from 72.3 to 43.1 ($p=0.002$)
- **Activity levels:** Mean daily steps increased from 3,359 to 5,862; five responders sustained >10,000 daily steps
- **Sustained response:** Five of six responders maintained improvement with SF-36 scores of 80–95

Safety All planned treatments were administered with no serious adverse events. Serum IgG showed transient reduction (54% in responders vs 40% in non-responders), suggesting plasma cell contribution to symptoms.

Predictors Low baseline natural killer (NK) cell count was significantly associated with lack of clinical improvement, suggesting immune dysregulation patterns may predict response.

Implications This trial provides the strongest evidence to date for a plasma cell-mediated autoimmune mechanism in a subset of ME/CFS patients. The contrast with rituximab failures is instructive: rituximab targets B cells but not established plasma cells, so circulating autoantibodies persist for months even after B-cell depletion. Daratumumab's success suggests that **the continuous stream of autoantibodies from long-lived plasma cells**—not the B cells themselves—may be the critical driver.

? Open Question 1: Identifying the Autoimmune Subgroup

Which ME/CFS patients are most likely to respond to plasma cell depletion? The 60% response rate suggests heterogeneity. Potential biomarkers for patient selection include: elevated GPCR autoantibody titers, post-infectious onset pattern, specific HLA types, or degree of IgG reduction post-treatment. Randomized controlled trials with biomarker stratification are urgently needed.

24.2.2 Cytokine Modulation

Cytokine abnormalities are well-documented in ME/CFS, though patterns vary between patients and disease stages. Importantly, recent research has elucidated the mechanistic link between GPCR autoantibodies and cytokine dysregulation. Hackel et al. (2025) demonstrated that autoantibodies mediate inflammatory and neurotrophic cytokine production via monocyte activation [209]. In post-COVID ME/CFS patients, autoantibody binding to monocytes

upregulated production of MIP-1 δ , PDGF-BB, and TGF- β 3. This provides a mechanistic pathway from circulating autoantibodies to the downstream inflammatory cascade characteristic of ME/CFS.

JAK Inhibitors

JAK inhibitors (baricitinib, tofacitinib, ruxolitinib) block cytokine signaling pathways and have demonstrated efficacy in inflammatory conditions such as rheumatoid arthritis and certain interferonopathies. Theoretical relevance to ME/CFS includes:

- Reduction of interferon-driven inflammation (relevant if chronic viral activation present)
- Modulation of IL-6 and other pro-inflammatory cytokines
- Effects on T cell activation and differentiation

However, JAK inhibitors carry significant risks including infection susceptibility and thrombosis, making them inappropriate for empirical use without clear inflammatory biomarkers.

24.2.3 Cellular Therapies

Mesenchymal Stem Cell Therapy

Mesenchymal stem cells (MSCs) exert immunomodulatory effects independent of tissue regeneration, secreting anti-inflammatory cytokines and modulating immune cell function. Anecdotal reports and small uncontrolled studies suggest:

- Variable responses with some dramatic responders
- Transient improvements lasting weeks to months
- Better responses in patients with clear inflammatory profiles

However, no controlled trials in ME/CFS have been published. Quality control, standardization, and cost remain significant barriers. The regenerative medicine industry includes both legitimate research centers and predatory clinics.

24.3 Autonomic and Neurological Interventions

24.3.1 Vagal Tone Restoration

The vagus nerve serves as master regulator of the autonomic nervous system, mediating the transition between sympathetic (“fight-or-flight”) and parasympathetic (“rest-and-digest”) states. In ME/CFS, vagal tone appears chronically suppressed, contributing to:

- Tachycardia and orthostatic intolerance
- Impaired heart rate variability

- Digestive dysfunction
- Chronic low-grade inflammation (the vagus provides anti-inflammatory signaling)

A hypothesized gut–vagal link may also be relevant: butyrate enhances enterochromaffin cell serotonin production [14], and enterochromaffin serotonin activates vagal afferents [13, 228] (see Section 11.1.4 in Chapter 11). If butyrate deficiency in ME/CFS reduces this serotonergic drive, vagal restoration strategies might benefit from concurrent gut-directed interventions. This inference extends the neurotransmitter dysregulation framework of Wirth and Scheibenberg [225]¹ but has not been directly tested.

Vagal Nerve Stimulation Devices

Non-invasive vagal nerve stimulation (nVNS) devices (gammaCore, others) deliver electrical stimulation transcutaneously. While FDA-approved for migraine and cluster headache, off-label use in ME/CFS has shown:

- Improvements in heart rate variability in some patients
- Reduced inflammation markers
- Variable effects on fatigue and other core symptoms

Natural Vagal Activation Techniques

Multiple accessible interventions stimulate vagal pathways:

- **Cold exposure:** Cold water face immersion triggers the mammalian dive reflex, powerfully activating vagal output
- **Slow exhale-dominant breathing:** Breathing patterns with extended exhalation (4-7-8 breathing, box breathing with longer exhale) directly stimulate vagal tone
- **Gargling and singing:** Vigorous gargling or sustained vocalization activates vagal branches innervating the pharynx
- **Gut-vagus signaling:** Certain probiotic strains (particularly *Lactobacillus rhamnosus*) signal via gut vagal afferents, affecting central stress responses. Since butyrate enhances enterochromaffin serotonin production in preclinical models [14], restoring butyrate-producing bacteria could theoretically improve vagal afferent activation in ME/CFS—though this therapeutic extrapolation remains untested

Speculation 68 (Comprehensive Vagal Rehabilitation Protocol). A multi-modal vagal rehabilitation program might combine: (1) daily cold water face immersion (starting at 10 seconds, gradually extending), (2) twice-daily extended exhale breathing sessions (5 minutes each), (3) regular gargling during oral hygiene, (4) vagus-active probiotic supplementation, and (5) heart rate variability biofeedback training. Such a protocol is low-risk and low-cost but would require consistent application over months. The hypothesis: sustained vagal training might

¹Currently available as a preprint; not yet peer-reviewed.

gradually shift autonomic setpoint from chronic sympathetic dominance toward parasympathetic balance, improving both autonomic symptoms and downstream effects on inflammation and digestion.

24.3.2 Neurostimulation

Transcranial Magnetic Stimulation (TMS)

Repetitive TMS can modulate cortical excitability and has shown benefit in depression, fibromyalgia, and chronic pain. Application to ME/CFS remains investigational:

- Targeting the dorsolateral prefrontal cortex may improve cognitive symptoms
- Motor cortex stimulation may modulate fatigue perception
- Anti-inflammatory effects via vagal pathway activation reported

Transcranial Direct Current Stimulation (tDCS)

tDCS delivers weak electrical current through scalp electrodes, subtly modulating neuronal excitability. As a low-cost, home-applicable intervention, it has attracted patient community interest. Evidence in ME/CFS specifically remains limited, though benefits in chronic fatigue, depression, and cognitive dysfunction in other conditions provide theoretical rationale.

24.3.3 Cerebrospinal Fluid Interventions

Intracranial Pressure Management

A subset of ME/CFS patients, particularly those with severe headaches worsened by lying down, may have altered CSF dynamics. Elevated or low intracranial pressure can produce fatigue and cognitive symptoms. Diagnostic lumbar puncture with pressure measurement can identify this subgroup.

Craniocervical Instability

Craniocervical instability (CCI) refers to excessive mobility or abnormal alignment at the junction between the skull base (occiput) and the upper cervical spine (C1–C2). The related condition atlantoaxial instability (AAI) specifically involves the articulation between the atlas (C1) and axis (C2) vertebrae. These conditions can produce brainstem compression, altered cerebrospinal fluid dynamics, and vagal dysfunction—all potentially contributing to ME/CFS symptomatology.

Observation 98 (High Prevalence of Craniocervical Pathology in ME/CFS). Bragée et al. [155] performed upright MRI on 229 ME/CFS patients (Canadian Consensus Criteria) at a specialized Swedish clinic, finding craniocervical obstructions in 80% (183/229) of cases. Notably, 75% of the cohort had hypermobility indicators and 45% had Chiari malformation (versus ~1% in the general population). However, this striking prevalence must be interpreted cautiously: patients presenting to a clinic known for investigating structural causes represent a highly selected population, likely overestimating true community prevalence.

The overlap between ME/CFS, hypermobility spectrum disorders, and craniocervical pathology has garnered increasing attention. Several high-profile patient cases, including science journalist Jennifer Brea and ME/CFS advocate Jeff Wood, achieved substantial or complete remission following craniocervical fusion surgery, generating considerable community interest in this intersection.

Diagnostic Criteria Diagnosis of CCI/AAI relies on specific radiographic measurements, though reference ranges have been refined in recent years. Traditional thresholds may have been overly conservative.

Table 24.1: Craniocervical Instability Radiographic Measurements

Measurement	Traditional Threshold	Updated Neutral Range
Clivo-Axial Angle (CXA)	<135° pathological	128–169° (neutral)
Basion-Dens Interval (BDI)	≥12 mm pathological	2.0–8.0 mm (neutral)
Grabb-Mapstone-Oakes (pB-C2)	≥9 mm suggests compression	4.2–10.2 mm (neutral)
C1–C2 Angular Displacement	>41° or facet overlap <10%	Indicates AAI

Updated ranges from Nicholson et al. [157] (50 healthy adults) and systematic review by Lohkamp et al. [156] (EDS cohorts). No ME/CFS-specific diagnostic validation studies exist; application to ME/CFS requires clinical judgment.

Imaging Modalities Standard supine MRI may fail to detect functionally significant instability that manifests only under gravitational loading or during cervical motion:

- **Upright Dynamic MRI:** Considered the gold standard for functional assessment. Captures the craniocervical junction under physiological gravitational stress, potentially revealing pathology occult on supine imaging [600].
- **Digital Motion X-ray (DMX):** Fluoroscopic imaging at 30 frames per second during active cervical motion. Useful for detecting dynamic instability but provides limited soft tissue detail.
- **Flexion-Extension CT:** Standard modality for quantifying osseous atlantoaxial relationships. Required for surgical planning.
- **Rotational 3D CT:** Helpful for assessing rotational AAI, particularly relevant in patients with torticollis or pain during head rotation.

Conservative Treatment Conservative management should be attempted before surgical consideration, though evidence specifically in ME/CFS populations is limited.

Physical Therapy. A Delphi consensus [158] established guidelines for physical therapy in hypermobile patients with craniocervical involvement:

- **Safe interventions:** Postural education and ergonomic optimization; diaphragmatic breathing training; motor control exercises for deep cervical flexors; scapular stabilization; thoracic spine mobility work.
- **Approach with caution:** Sustained stretching of cervical musculature; passive range-of-motion at end-range; manual therapy to upper cervical segments.

△ **Warning 1: Contraindicated Interventions in CCI/AAI**

The following interventions are contraindicated in patients with suspected or confirmed craniocervical instability:

- High-velocity, low-amplitude (HVLA) chiropractic manipulation of the cervical spine
- Cervical traction
- Aggressive manual therapy targeting the upper cervical segments
- Forced end-range passive movements

These interventions risk exacerbating instability, neural compression, or vertebral artery injury.

Cervical Orthoses. External support can provide symptomatic relief and allow assessment of potential surgical benefit:

- **Rigid collars** (Aspen Vista, Miami-J): Indicated for moderate-to-severe symptoms. Provide substantial motion restriction.
- **Soft collars:** May provide proprioceptive feedback and mild support for milder cases.
- **Protocol:** For mild-to-moderate symptoms, trial 20–30 minutes three times daily. Severe cases awaiting surgery may require continuous use.
- **Caution:** Prolonged collar use risks cervical muscle atrophy, potentially worsening long-term instability.

Prolotherapy and Platelet-Rich Plasma. Injection therapies targeting ligamentous laxity have been proposed:

- **Dextrose prolotherapy:** Hypertonic dextrose injected into ligamentous attachments, theoretically promoting fibroblast proliferation and tissue tightening.
- **Platelet-rich plasma (PRP):** Growth factor-rich preparation for more significant ligamentous damage.
- **Evidence level:** Case series only; no randomized controlled trials. Response durability and optimal protocols remain undefined.

Surgical Intervention Surgical fusion remains controversial but has produced dramatic improvements in carefully selected patients.

Indications. Based on the Henderson surgical series [159, 601], surgical candidates typically demonstrate:

- Clear radiographic evidence of instability or compression with symptom concordance (patient’s cardinal symptoms correlate with the anatomical location and expected pathophysiology of compression)
- Progressive neurological deficits (myelopathy)
- Failure of adequate conservative trial (typically 6–12 months)
- Symptom improvement with cervical orthosis (positive “collar test”)

The most common procedure is occipito-cervical fusion (C0–C1–C2), though extent of fusion depends on levels of documented instability.

Surgical Outcomes. The most robust surgical outcome data come from EDS cohorts rather than ME/CFS populations specifically. Henderson et al. [159] reported outcomes in 53 patients with Ehlers-Danlos syndrome undergoing craniocervical fusion:

- Significant improvement in pain scores ($p < 0.001$)
- Reduced medication requirements ($p < 0.0001$)
- Improved Karnofsky Performance Status ($p < 0.001$)
- Neurological symptom improvement: nausea, syncope, speech difficulties, concentration, vertigo, and fatigue all showed statistically significant gains
- Fusion rate: 100% in this experienced surgical series
- Five-year follow-up [601] demonstrated sustained improvement

Whether these results generalize to ME/CFS patients with CCI (who may have different underlying pathophysiology than primary EDS patients) requires prospective study.

△ **Warning 2: Surgical Complications**

Craniocervical fusion carries meaningful risks [156]:

- Overall complication rate: 12–20%
- Deep wound infection: 2–4%
- Pseudoarthrosis (failed fusion): 2–8%
- Vertebral artery injury: <2%
- Revision surgery required: ~8%
- Adjacent segment degeneration: Long-term concern with any spinal fusion

Surgical decision-making requires careful risk-benefit analysis with an experienced craniocervical surgeon.

Patient Selection and Controversies

△ Warning 3: Critical Caveats for CCI in ME/CFS

Several important limitations warrant emphasis:

- **Selection bias:** The 80% prevalence from Bragée et al. derives from a clinic specifically investigating structural causes, likely overestimating true population prevalence.
- **Laxity versus instability:** Ligamentous laxity (common in hypermobility syndromes) does not equate to clinically significant spinal instability. Many hypermobile individuals have radiographic “abnormalities” without corresponding symptoms.
- **No comparative trials:** No randomized controlled trials compare surgical versus conservative management. Dramatic surgical success stories may reflect publication and reporting bias.
- **Causation uncertain:** What proportion of ME/CFS is mechanistically driven by craniocervical pathology versus coincidental remains unknown. Symptom overlap between CCI and ME/CFS is substantial, complicating causal attribution.

? Open Question 2: CCI in ME/CFS: Key Unknowns

Critical research gaps include:

- Population-based prevalence of CCI findings in ME/CFS using standardized imaging protocols
- Predictive criteria identifying which ME/CFS patients would benefit from CCI-directed treatment
- Long-term outcomes beyond 5 years post-fusion
- Comparative effectiveness of conservative versus surgical management in matched cohorts
- Mechanistic studies clarifying how craniocervical pathology produces ME/CFS-like symptoms

Until these questions are addressed, CCI treatment in ME/CFS remains an area requiring careful individualized assessment rather than routine screening.

24.4 Metabolic Interventions

24.4.1 Mitochondrial “Jumpstart” Protocols

If mitochondria are damaged or functionally impaired, restoring normal function may require more than supplying individual cofactors.

Speculation 69 (Combined Mitochondrial Biogenesis Protocol). A multi-component mitochondrial support protocol might include:

- **Biogenesis stimulation:** PQQ (pyrroloquinoline quinone) activates pathways promoting new mitochondrial formation
- **Electron transport support:** High-dose CoQ10 (ubiquinol form, 400–600 mg) supports complex III function
- **Alternative electron carriers:** Methylene blue at very low doses (0.5–1 mg/kg) can accept electrons from complex I and transfer directly to complex IV, bypassing damaged components—highly experimental
- **ATP precursor loading:** D-ribose provides the sugar backbone for ATP synthesis
- **Photobiomodulation:** Red and near-infrared light (600–1000 nm) is absorbed by cytochrome c oxidase, potentially enhancing complex IV function

The rationale: single-agent approaches may fail because the electron transport chain requires all components functional. Simultaneously supporting multiple elements while stimulating biogenesis of new mitochondria might achieve what individual supplements cannot.

24.4.2 NAD⁺ Precursor Therapy

Given the evidence for NAD⁺ metabolism abnormalities in ME/CFS (see Chapter 6), supplementation with NAD⁺ precursors represents a promising therapeutic avenue.

Nicotinamide Riboside (NR)

Nicotinamide riboside is a form of vitamin B3 that serves as a precursor to NAD⁺, bypassing rate-limiting steps in the salvage pathway.

Mechanism NAD⁺ is essential for:

- Mitochondrial electron transport chain function
- Sirtuin activation (cellular stress response, mitophagy)
- DNA repair via PARP enzymes
- Cellular redox balance

Clinical Evidence A 2025 randomized controlled trial in Long COVID (which shares substantial symptom overlap with ME/CFS) evaluated NR at 2000 mg/day [572]:

- **Sample:** 58 participants with Long COVID randomized 2:1 to NR vs placebo
- **NAD⁺ response:** Levels increased 2.6- to 3.1-fold (measured at interim timepoints of 5 and 10 weeks within a 20-week intervention)
- **Cognitive outcomes:** Variable; overall group differences limited but some individuals showed improvement after ≥10 weeks
- **Safety:** Well-tolerated in this trial (n=58) at 2000 mg daily; insufficient sample size to characterize rare adverse events

Earlier research on oral NADH (a reduced form) in ME/CFS showed modest benefits in some patients [542], and combination with CoQ10 improved maximum heart rate recovery [571], though fatigue effects were inconsistent across the broader programme of trials.

Practical Considerations

- Commercial NR supplements are widely available
- Typical doses: 300–1000 mg daily; research doses up to 2000 mg
- Response may require 10+ weeks of consistent supplementation
- Cost can be substantial for high-dose regimens

Nicotinamide Mononucleotide (NMN)

NMN is another NAD⁺ precursor, one step closer to NAD⁺ in the biosynthetic pathway. It has been hypothesized to be more efficient than NR due to its position in the biosynthetic pathway, though comparative clinical trials are lacking and this remains unproven. Similar safety profile and availability to NR.

24.4.3 Metabolic Modulators

Dichloroacetate (DCA)

DCA activates pyruvate dehydrogenase, promoting glucose oxidation over glycolysis. Given evidence of PDH dysfunction in ME/CFS, DCA has theoretical appeal as a speculative intervention. However, neurotoxicity with chronic use limits clinical application, and no ME/CFS-specific trials have been conducted.

Oxaloacetate

Oxaloacetate supplementation may theoretically support the citric acid cycle and has shown neuroprotective effects in preclinical models. As a key TCA cycle intermediate, it could potentially bypass certain metabolic blocks, though this rationale remains speculative and untested in ME/CFS.

24.4.4 Ketogenic and Metabolic Switching Approaches

~ Hypothesis 4: Forced Metabolic Flexibility Training

ME/CFS may involve loss of metabolic flexibility—the ability to switch between fuel sources (glucose, fatty acids, ketones) based on availability and demand. A protocol designed to force repeated metabolic switching might restore this flexibility:

- Time-restricted eating (16–18 hour fasting window) to induce daily ketone production
- Periodic extended fasts (24–48 hours) with medical supervision
- Cycling between ketogenic and higher-carbohydrate phases
- Exercise timing relative to fed/fasted state (very cautiously, respecting PEM)

Caution: fasting can be dangerous for ME/CFS patients, particularly those with blood sugar dysregulation, and should only be attempted with medical guidance and careful monitoring.

24.5 Microbiome Interventions

Gut microbiome alterations are consistently documented in ME/CFS, though whether they represent cause, consequence, or parallel phenomenon remains unclear.

24.5.1 Fecal Microbiota Transplantation

FMT represents the most radical microbiome intervention—complete ecosystem replacement rather than supplementation with isolated strains.

Theoretical Rationale

The rationale for FMT in ME/CFS is mechanistically plausible but unvalidated; no ME/CFS-specific FMT trials have been published:

- Restores microbial diversity that may be impossible to achieve with probiotics
- Transfers not just bacteria but bacteriophages, fungi, and microbial metabolites
- Donor microbiome may provide metabolic functions missing in ME/CFS (butyrate production, tryptophan metabolism)
- Potential to reset gut-immune interactions

Practical Considerations

- Donor selection is critical—health, diet, antibiotic history all matter
- Pre-treatment antimicrobial clearing may improve engraftment
- Dietary changes post-FMT are essential to support the new ecosystem
- Multiple treatments may be necessary
- Risk of pathogen transmission exists, though screening reduces this substantially

Speculation 70 (Comprehensive Microbiome Reset Protocol). A thorough microbiome restoration might include:

1. **Preparation:** Low-FODMAP diet for 2 weeks to reduce pathogenic overgrowth
2. **Clearing:** Targeted antimicrobials (rifaximin for SIBO if present) or elemental diet
3. **Transplant:** FMT from carefully selected healthy donor
4. **Establishment:** Strict dietary protocol matching donor's diet for 4–6 weeks
5. **Maintenance:** Diverse, fiber-rich diet with targeted prebiotics
6. **Monitoring:** Repeat microbiome sequencing at intervals to assess engraftment

This represents a significant undertaking but addresses a potential root cause rather than symptoms.

24.5.2 Precision Microbiome Modulation

Targeted Probiotics

Rather than broad-spectrum probiotics, specific strains may address specific deficits:

- *Faecalibacterium prausnitzii* (butyrate producer, often depleted in ME/CFS)
- *Akkermansia muciniphila* (gut barrier integrity)
- *Lactobacillus reuteri* (histamine modulation, vagal signaling)

Bacteriophage Therapy

Phages (viruses that infect bacteria) can selectively eliminate pathogenic species while sparing beneficial ones—precision antimicrobials. While not yet clinically available for ME/CFS, this technology is advancing rapidly.

24.6 Technologies and Devices

24.6.1 Apheresis Techniques

Therapeutic Plasma Exchange

Plasma exchange removes and replaces plasma, eliminating circulating factors including autoantibodies, inflammatory mediators, and potentially microclots. Case reports have described improvements in ME/CFS and Long COVID, though controlled trials are lacking.

HELP Apheresis

Heparin-induced extracorporeal LDL precipitation (HELP) removes not only LDL cholesterol but also fibrinogen and inflammatory mediators. Reports from Germany describe improvements in some Long COVID patients, with theoretical relevance to ME/CFS.

24.6.2 Hyperbaric Oxygen Therapy

HBOT delivers 100% oxygen at elevated atmospheric pressure, dramatically increasing tissue oxygen levels. Proposed mechanisms in ME/CFS include:

- Enhanced mitochondrial function
- Reduced hypoxia in poorly perfused tissues
- Stem cell mobilization
- Reduced inflammation
- Neuroplasticity enhancement

Evidence in ME/CFS is limited to case reports and uncontrolled studies; patient responses appear highly variable and no controlled trials have been published.

24.6.3 Photobiomodulation

Red and near-infrared light therapy (wavelengths 600–1000 nm) penetrates tissue and is absorbed by cytochrome c oxidase in mitochondria. Proposed effects include:

- Enhanced mitochondrial ATP production
- Reduced oxidative stress
- Anti-inflammatory effects
- Improved microcirculation

Home devices are widely available, though quality and specifications vary significantly.

24.7 Repurposed Medications

24.7.1 Suramin

Suramin, an antiparasitic drug from 1916, blocks purinergic signaling—the basis of Naviaux’s cell danger response hypothesis [197]. A small pilot study in *autism spectrum disorder* (not ME/CFS) demonstrated that single-dose suramin improved language, social interaction, and restricted behaviors, with effects reversing after drug elimination [15]. No controlled suramin trial has been conducted in ME/CFS patients; interest is extrapolated from shared purinergic signaling mechanisms. However:

- Suramin has significant toxicity with repeated dosing
- It is not available outside research settings
- Single-dose effects are transient

Development of safer antipurinergic agents continues.

24.7.2 Rapamycin (Sirolimus)

Rapamycin inhibits mTOR, a master regulator of cellular metabolism, growth, and autophagy. Theoretical rationale in ME/CFS:

- Promotes autophagy (cellular “cleanup”)
- Immunomodulatory effects
- May enhance mitochondrial biogenesis through feedback mechanisms

However, mTOR inhibition also suppresses immune function and protein synthesis, making chronic use problematic.

24.7.3 Metformin

Metformin’s mechanisms extend beyond glucose control to include AMPK activation, mitochondrial effects, and anti-inflammatory properties. As a safe, well-characterized drug, it represents a relatively accessible option for empirical trial, though evidence in ME/CFS specifically remains limited.

24.7.4 Low-Dose Aripiprazole

Aripiprazole at very low doses (0.5–2 mg) may modulate neuroinflammation through effects on microglial function [602]. A pilot study (n=25) showed promising results for fatigue and cognitive symptoms. Patient community reports suggest benefit in some individuals, particularly for brain fog and energy.

24.7.5 Ketamine

Ketamine, an NMDA receptor antagonist used as an anesthetic, has emerging applications in chronic pain and neuroinflammatory conditions that may be relevant to ME/CFS.

Mechanism of Action

Ketamine's mechanisms extend beyond NMDA receptor blockade [603]:

- **NMDA receptor antagonism:** Reduces central sensitization and neuronal hyperexcitability—potentially relevant to ME/CFS-associated pain amplification and sensory hypersensitivity
- **Anti-neuroinflammatory effects:** Suppresses microglial activation and reduces pro-inflammatory cytokines (IL-6, IL-8, TNF- α)
- **Neuroprotection:** Reduces excitotoxicity during ATP depletion by limiting calcium-mediated cell death
- **Synaptic plasticity enhancement:** Effects on AMPA receptors and mTOR pathway may promote neuroplasticity
- **Glutamate modulation:** Paradoxical transient glutamate increase followed by rebalancing of excitatory/inhibitory dynamics

Evidence Base

Fibromyalgia (Related Condition) Systematic reviews document ketamine's effects in fibromyalgia, a condition with significant ME/CFS overlap [604, 605]:

- Short-term pain reduction with single IV infusions (effects lasting hours to days)
- Higher doses and longer/more frequent infusions associated with more sustained analgesia
- Dose-response relationship supports efficacy but optimal protocols remain undefined
- Ongoing ESKEFIB trial evaluating S-ketamine dose-escalation [606]

Anti-Fatigue Effects Notably, ketamine demonstrates anti-fatigue effects independent of its antidepressant action [607]:

- Single IV infusion significantly improved fatigue in treatment-resistant bipolar disorder within 40 minutes
- Peak efficacy at day 2 post-infusion
- Anti-fatigue effect **not fully explained by mood improvement**—suggesting direct fatigue-targeting mechanism
- Pilot study in MS-related fatigue showed significant improvement sustained at 4 weeks post-infusion

ME/CFS-Specific Evidence

- **No controlled trials:** No registered clinical trials specifically evaluating ketamine in ME/CFS have been identified
- **Anecdotal reports:** Some ME/CFS patients treated empirically at specialized pain clinics report benefit
- **Mechanistic rationale:** Strong theoretical basis given NMDA involvement in central sensitization, neuroinflammation in ME/CFS pathophysiology, and ATP depletion rendering neurons vulnerable to glutamate excitotoxicity

Administration and Protocols

- **Sub-anesthetic IV infusion:** 0.5 mg/kg over 40 minutes (standard research protocol for depression/pain)
- **Intranasal:** 50 mg (esketamine formulation FDA-approved for depression)
- **Maintenance:** Responders may require periodic infusions (typically twice monthly in fibromyalgia protocols)
- **Setting:** Requires specialized clinic with cardiopulmonary monitoring capability

Safety Considerations

△ **Warning 4: Ketamine Risks and Contraindications**

Contraindications:

- Active psychotic symptoms or history of psychosis
- Unstable cardiovascular disease (ketamine increases blood pressure and heart rate)
- Significant intracranial pathology
- Current substance use disorders (ketamine has abuse potential)

Adverse effects:

- Dissociative symptoms during/after infusion (typically transient)
- Nausea, dizziness, headache
- Blood pressure and heart rate elevation
- With chronic use: cognitive impairment, urological toxicity, dependence risk

Required monitoring: Continuous cardiopulmonary monitoring during infusion; post-administration observation minimum 30 minutes; mental status assessment.

Clinical Positioning

Ketamine remains **highly experimental** for ME/CFS:

- No ME/CFS-specific trials to guide use
- Primarily CNS-acting; may not address peripheral metabolic dysfunction

- High cost and limited availability (typically not covered by insurance for ME/CFS)
- Best viewed as symptomatic/neuroprotective intervention rather than disease-modifying
- May be most appropriate for patients with prominent pain, central sensitization features, or treatment-refractory fatigue who have exhausted safer options

24.8 Peptide Therapies

24.8.1 BPC-157

Body Protection Compound 157 is a synthetic peptide derived from a gastric protein. Proposed effects include:

- Gut healing and gut-brain axis modulation
- Anti-inflammatory effects
- Promotion of angiogenesis and tissue repair

Evidence is primarily from animal studies; human data are limited to case reports.

24.8.2 Thymosin Alpha-1

Thymosin alpha-1 is an immunomodulatory peptide that enhances T cell and NK cell function. Given NK cell dysfunction in ME/CFS, there is theoretical rationale, though clinical evidence is lacking.

24.9 Integrated Treatment Strategies

~ Hypothesis 5: Sequential Multi-System Protocol

Given the multi-system nature of ME/CFS, effective treatment may require addressing multiple systems in sequence:

1. **Stabilization:** Strict pacing, anti-inflammatory diet, sleep optimization, stress reduction
2. **Infection clearing:** Test for and treat any chronic infections (EBV reactivation, HHV-6, SIBO, oral infections)
3. **Gut restoration:** Address dysbiosis, consider FMT if severe
4. **Autoimmune intervention:** If autoantibodies present, consider immunoadsorption or BC007
5. **Metabolic support:** Mitochondrial support stack, consider photobiomodulation
6. **Autonomic rehabilitation:** Vagal toning protocols, gradual orthostatic training

7. **Cautious reconditioning:** Only after sustained improvement, very gradual activity increases

This sequential approach addresses the possibility that treating downstream problems while upstream drivers persist yields only temporary benefit.

? Open Question 3: Identifying the Critical Intervention Point

In complex, multi-system illness, is there a “keystone” dysfunction that, if corrected, allows other systems to normalize? Or must multiple systems be addressed simultaneously? Identification of critical intervention points—perhaps through computational modeling of system interactions—could dramatically improve treatment efficiency.

24.9.1 Cycle-Based Multi-Target Treatment

The vicious cycle dynamics framework (Chapter 2, §2.1, “Vicious Cycle Dynamics”) reveals why single-target interventions often fail in established ME/CFS: when four or five reinforcing cycles are active simultaneously, breaking only one may be insufficient. The untreated cycles continue driving dysfunction, limiting or negating the benefits of the single intervention. This section presents treatment strategies explicitly designed to address cycle interactions and mutual reinforcement.

Why Single-Target Trials Fail: The Cycle Reinforcement Problem

The pattern of negative clinical trials in ME/CFS becomes comprehensible when viewed through the cycle dynamics lens:

- **Rituximab trials** [376]: Targeted B cells (immune cycle), but left mitochondrial, autonomic, neuroinflammatory, and endocrine cycles active. Any improvement from reduced autoantibody production was overwhelmed by ongoing dysfunction from untreated cycles.
- **CoQ10 monotherapy**: Supports mitochondrial function but cannot overcome persistent immune activation, autonomic failure, or neuroinflammation. The mitochondrial cycle continues to be driven by oxidative stress from active immune and neuroinflammatory cycles.
- **Immunoadsorption partial responders**: Even when autoantibodies are removed (breaking immune cycle), only 30% achieved sustained improvement at 6–12 months [147]. Hypothesis: in the remaining 70%, disease is maintained by non-immune cycles (mitochondrial, autonomic, neurological) that continue operating independently.
- **Pacing alone**: Reduces exacerbation triggers but cannot reverse established cycle entrenchment. In severe disease with all five cycles active and multiple irreversibility mechanisms engaged, pacing prevents worsening but may not enable recovery.

★ **Key Point: The Cycle Synergy Hypothesis**

If vicious cycles reinforce each other through shared mediators and feedback loops, breaking multiple cycles simultaneously should produce synergistic—not merely additive—effects. Eliminating mutual reinforcement may allow the remaining cycles to destabilize and collapse, whereas sequential single-target interventions leave the cycle network intact.

Testable prediction: A three-target protocol (mitochondrial + immune + autonomic) should produce $> 3\times$ the benefit of any single intervention alone, and $> 1.5\times$ the summed benefits of each intervention applied sequentially.

Mathematical Framework: The Network Model

The synergy hypothesis can be formalized mathematically, providing testable quantitative predictions. Let $C_1, C_2, \dots, C_n$ represent n vicious cycles (mitochondrial, immune, autonomic, neuroinflammatory, endocrine) with states $s_1, s_2, \dots, s_n$ quantifying the severity of each cycle's dysfunction. The dynamics include both internal self-amplification (gain G_i) and cross-cycle coupling:

$$\frac{ds_i}{dt} = G_i s_i + \sum_{j \neq i} \alpha_{ij} s_j - \gamma_i T_i \quad (24.1)$$

where G_i = internal gain of cycle i (self-amplification strength), α_{ij} = coupling strength from cycle j to cycle i , T_i = treatment intensity targeting cycle i , and γ_i = treatment efficacy coefficient.

Why Single-Target Intervention Cannot Succeed. When $\alpha_{ij} > 0$ (reinforcing coupling exists), treating only cycle i while cycle j remains active results in persistent input from j that maintains i dysfunction. At equilibrium:

$$s_i^* = \frac{\sum_{j \neq i} \alpha_{ij} s_j}{\gamma_i T_i - G_i} \quad (24.2)$$

As long as $s_j > 0$ for any strongly coupled cycle j , we have $s_i^* > 0$: the treated cycle cannot fully resolve. This explains the pattern of “partial response” seen in most ME/CFS trials—interventions reduce cycle severity but cannot eliminate dysfunction while other cycles remain active.

The Synergy Prediction. Simultaneous treatment of coupled cycles eliminates mutual reinforcement. The combined effect should be super-additive:

$$\text{Effect}(T_1 + T_2) > \text{Effect}(T_1) + \text{Effect}(T_2) \quad (24.3)$$

Quantitative estimate: If ubiquinol alone produces 10-point improvement on 100-point fatigue scale and LDN alone produces 8-point improvement, the combination should yield 25–35 points (not the additive 18 points) due to elimination of cross-cycle reinforcement. This is a falsifiable prediction: if combination effects are purely additive, the network model is refuted.

Evidence Grade C: Network models are well-established in systems biology; application to ME/CFS treatment is theoretical but consistent with observed partial response patterns and heterogeneous treatment outcomes.




Figure 24.1: **Network model of vicious cycle coupling in ME/CFS.** Five major pathophysiological cycles (colored ellipses) exhibit bidirectional reinforcing connections (red arrows). Each cycle contains internal positive feedback (self-loops). Yellow stars indicate potential treatment intervention points. The network architecture explains why treatment at single nodes often fails: coupled cycles re-amplify dysfunction even when one component is partially addressed. Effective treatment may require simultaneous intervention at multiple nodes.

Design Principles for Multi-Target Protocols

Principle 1: Target Active Cycles, Not All Cycles. Not every ME/CFS patient has all five cycles active. Personalized cycle mapping (see Chapter 18, diagnostic battery proposal) identifies which specific cycles are operating in each individual. Treatment should target documented active cycles rather than empirically treating all possible pathways.

Example: A patient with elevated GPCR autoantibodies, low ATP production, but normal autonomic function has active mitochondrial and immune cycles. Protocol: CoQ10 + immunoadsorption or daratumumab. Adding fludrocortisone (autonomic target) would be unnecessary and risks side effects without benefit.

Principle 2: Address Irreversibility Mechanisms. In established severe disease (Stage 5 in the sequential cycle entry model), multiple irreversibility mechanisms may be engaged: epigenetic silencing, mitochondrial DNA mutations, central sensitization, microglial priming. These mechanisms maintain disease even if initial triggers are removed. Multi-target protocols must include interventions addressing entrenchment, not just acute drivers.

Example: Combining plasma cell depletion (daratumumab, removes autoantibody source) with HDAC inhibitors (experimental; reverses epigenetic silencing) plus NAD⁺ precursors (supports mitochondrial biogenesis to replace damaged organelles).

Principle 3: Sequence for Safety and Synergy. Simultaneous initiation of multiple interventions risks:

- Inability to identify which component helps or harms
- Compounded side effects
- Drug interactions

Staged initiation protocol:

1. **Weeks 1–4:** Establish low-risk foundation (mitochondrial support: CoQ10, NAD⁺ precursors, antioxidants)
2. **Weeks 5–8:** Add second target if first is tolerated (autonomic: fludrocortisone, or immune: immunoadsorption if indicated)
3. **Weeks 9+:** Add third target only if first two are beneficial and well-tolerated
4. **Assessment at 6 months:** Measure synergistic effects; discontinue ineffective components

Principle 4: Maintain Escape Velocity. For inescapable cycles (gain > 1.0), interventions must reduce cycle gain below 1.0 to allow spontaneous recovery. Insufficient intervention intensity leaves gain > 1.0, producing temporary symptom reduction but no sustained improvement when treatment stops.

Clinical implication: Aggressive multi-target intervention may be required in severe disease, accepting higher risk to achieve cycle escape. Gentle, conservative monotherapy is appropriate

for mild disease (cycle gain near 1.0) but may be futile for severe disease with deeply entrenched cycles.

Proposed Multi-Target Protocols

Protocol A: Mitochondrial-Immune Synergy (Post-Infectious ME/CFS with Autoantibodies). **Indication:** Patients with documented GPCR autoantibodies (particularly β_2 -AR, M3/M4 muscarinic) and evidence of mitochondrial dysfunction (low ATP, elevated lactate, 2-day CPET failure).

Rationale: Autoantibodies impair autonomic regulation → tissue hypoperfusion → mitochondrial oxidative stress → ROS production → immune activation → more autoantibody production. Breaking both the immune source and the mitochondrial amplifier simultaneously may collapse the reinforcing loop.

Components:

1. **Plasma cell depletion:** Daratumumab (if accessible via clinical trial or compassionate use) 1800 mg subcutaneously per standard dosing [9]
 - Targets long-lived plasma cells secreting autoantibodies
 - Expected response timeline: 2–4 months for maximal effect
 - Contraindication: severe immunodeficiency, active infection
2. **Mitochondrial support stack:**
 - Ubiquinol (reduced CoQ10) 400–600 mg daily: Complex III electron transport support
 - MitoQ 10–20 mg daily: Mitochondria-targeted antioxidant; reduces ROS at source
 - PQQ 20 mg daily: Mitochondrial biogenesis via PGC-1 α activation
 - NAD⁺ precursor (NR or NMN) 500–1000 mg daily: Supports electron transport and sirtuins
 - Alpha-lipoic acid 600 mg daily: Regenerates other antioxidants; supports energy metabolism
3. **Anti-inflammatory bridge:**
 - Omega-3 fatty acids (EPA+DHA) 2–4 g daily: Reduces cytokine production during immune intervention
 - Low-dose naltrexone 1.5–4.5 mg nightly: Modulates microglial activation; may reduce neuroinflammation

Expected outcomes:

- *Hypothesis:* Synergistic improvement exceeding either intervention alone
- *Timeline:* Gradual improvement over 3–6 months; mitochondrial recovery lags autoantibody reduction
- *Responder criteria:* SF-36 Physical Function improvement ≥ 20 points, or activity level increase $\geq 50\%$ (e.g., daily steps from 3,000 to 4,500+)

- *Predicted synergy:* If daratumumab alone produces 15-point SF-36 improvement and mitochondrial stack alone produces 8 points, combination should yield 30–40 points (synergistic, not additive 23 points)

Monitoring:

- Baseline and monthly: SF-36, symptom severity scales, activity tracking
- Baseline and 3-month: GPCR autoantibody titers (if available), NK cell counts
- Baseline and 6-month: Two-day CPET (if accessible) to quantify functional metabolic improvement
- Safety: CBC, CMP monthly during daratumumab; monitor for infections

Protocol B: Mitochondrial-Autonomic-Immune Triple Therapy (Severe Multisystem Disease). **Indication:** Severe ME/CFS (housebound or bedbound) with evidence of mitochondrial failure, autonomic dysfunction (POTS, orthostatic intolerance), and immune activation (elevated cytokines or autoantibodies).

Rationale: In Stage 4–5 disease with multiple active cycles, addressing only two systems may leave sufficient reinforcement to maintain the network. Three-way targeting aims to destabilize the entire cycle structure.

Components:

1. **Mitochondrial support:** As in Protocol A
2. **Immune modulation:**
 - Preferred: Daratumumab (if accessible) or immunoadsorption
 - Alternative: Low-dose naltrexone + omega-3 fatty acids (if immunotherapy inaccessible)
3. **Autonomic support:**
 - Fludrocortisone 0.05–0.2 mg daily: Expands plasma volume; improves orthostatic tolerance
 - Midodrine 2.5–10 mg three times daily (if tolerated): Alpha-agonist; increases vascular tone
 - Salt loading: 6–10 g sodium daily (if no contraindications)
 - Compression garments: 20–30 mmHg abdominal binder or waist-high stockings
 - Pyridostigmine 30–60 mg three times daily (optional): Acetylcholinesterase inhibitor; enhances parasympathetic tone
4. **Catecholamine synthesis support** (for central deficiency [5]):
 - L-tyrosine 1500–3000 mg morning
 - BH4 cofactor support: Methylfolate 1–5 mg + methylcobalamin 1–5 mg
 - Iron repletion if deficient (target ferritin >50 ng/mL)

Staged initiation (critical for safety):

1. **Weeks 1–4:** Mitochondrial stack only; assess tolerance

2. **Week 5:** Add fludrocortisone 0.05 mg; titrate up weekly if tolerated (monitor BP, edema)
3. **Week 7:** Add L-tyrosine 1500 mg; increase to 3000 mg if no anxiety/jitteriness
4. **Week 9:** Add immune intervention (daratumumab or LDN)
5. **Week 12:** Add midodrine if orthostatic symptoms persist despite fludrocortisone
6. **Assessment at 6 months:** Full evaluation; discontinue ineffective components

Safety considerations:

- **Fludrocortisone:** Monitor for hypertension, hypokalemia, edema; check electrolytes monthly
- **Midodrine:** Contraindicated in supine hypertension; check BP lying and standing
- **Daratumumab:** Infection risk; neutropenia monitoring; avoid live vaccines
- **Drug interactions:** Multiple pathways; requires physician oversight
- **Discontinuation:** If severe side effects or clear worsening, stop most recent addition

Expected outcomes:

- *Hypothesis:* Triple therapy enables cycle network collapse in patients unresponsive to mono- or dual therapy
- *Realistic goal:* 20–40% functional improvement (e.g., bedbound → housebound, housebound → house-limited)
- *Timeline:* Gradual improvement over 6–12 months; early gains from autonomic support, delayed gains from mitochondrial/immune interventions
- *Non-responders:* If no improvement by 6 months, disease may be maintained by irreversibility mechanisms requiring different approaches (epigenetic interventions, neuroplasticity protocols)

Protocol C: Epigenetic Reversal + Metabolic Support (Established Severe Disease with Suspected Entrenchment). **Indication:** Very severe ME/CFS (>5 years duration, multiple treatment failures) where disease appears self-sustaining despite removal of obvious triggers. **Hypothesis:** epigenetic silencing of metabolic genes maintains dysfunction.

Rationale: In Stage 5 disease, epigenetic changes may lock cells into a maladaptive state. Reversing chromatin modifications while simultaneously supporting metabolic recovery might break the lock and allow spontaneous improvement mechanisms to engage.

Components:

1. **HDAC inhibitor** (research setting only):
 - Vorinostat (SAHA): Typical cancer dosing 400 mg daily; ME/CFS dose unknown
 - **CAUTION:** Significant toxicity (fatigue, GI distress, cytopenias, thromboembolism risk)
 - Requires oncology/hematology expertise; not for empirical use
 - Alternative (lower risk): Valproic acid (also HDAC inhibitor; used for epilepsy/bipolar); 500–1000 mg daily

2. Mitochondrial biogenesis stimulation:

- High-dose NAD⁺ precursors: NR or NMN 1000–2000 mg daily
- PQQ 20 mg, resveratrol 500 mg (SIRT1/PGC-1 α activation)
- Exercise mimetics (if tolerated): Metformin 500–1000 mg (AMPK activation)

3. Strict pacing during intervention:

- Epigenetic reversal requires cellular energy; any PEM during treatment may abort the process
- Maintain activity well below energy envelope; prioritize rest

Risk-benefit assessment:

- **High risk:** HDAC inhibitors cause significant side effects; cancer drug in non-cancer population
- **Uncertain benefit:** No clinical trials in ME/CFS; purely mechanistic hypothesis
- **Justification:** For very severe, treatment-refractory patients with no remaining options, experimental high-risk interventions may be ethically appropriate with full informed consent
- **Setting:** Research trial or compassionate use only; not standard care

△ Warning 5: Experimental High-Risk Protocol

Protocol C is **highly speculative and high-risk**. HDAC inhibitors have serious toxicity profiles and no safety or efficacy data in ME/CFS. This approach should only be considered in research settings for very severe patients who have exhausted all standard options and are working with physicians experienced in managing these medications. It is presented here to illustrate the therapeutic logic of targeting irreversibility mechanisms, not as a recommendation for clinical use.

Why Multi-Target Approaches Have Not Been Tested

Despite the mechanistic rationale, multi-target protocols face significant barriers:

- **Regulatory challenges:** Trials typically test single interventions; combination trials are complex and expensive
- **Pharmaceutical incentives:** Drug companies fund trials of their own products, not combinations with competitors' drugs
- **Heterogeneity:** Without biomarker-based patient stratification, combining treatments in a heterogeneous population may dilute signals
- **Sample size:** Detecting synergy requires larger trials than detecting main effects
- **Endpoint timing:** Synergistic effects may require 6–12 months to manifest; most trials are 3–6 months

★ Key Point: The Precision Medicine Imperative

Multi-target protocols cannot be “one-size-fits-all.” Effective implementation requires:

1. **Diagnostic cycle mapping:** Identify which cycles are active in each patient (mitochondrial, immune, autonomic, neurological, endocrine)
2. **Biomarker-guided targeting:** Measure GPCR autoantibodies, ATP/lactate, catecholamines, cytokines; treat documented abnormalities
3. **Severity staging:** Match intervention intensity to disease stage (gentle support for Stage 1–2, aggressive multi-target for Stage 4–5)
4. **Response monitoring:** Serial biomarkers and functional measures to identify responders and adjust protocols

The cycle dynamics framework provides the conceptual structure; precision diagnostics provide the implementation roadmap.

Testable Predictions and Future Trials

The cycle synergy hypothesis generates falsifiable predictions suitable for clinical trial testing:

1. **Synergy prediction:** Mitochondrial + immune dual therapy produces $> 1.5\times$ the summed benefit of sequential monotherapies
 - Trial design: Three arms (CoQ10 alone, daratumumab alone, combination); primary outcome SF-36 at 6 months
 - Sample size: 60 patients per arm (power to detect 15-point difference with synergy)
2. **Cycle status predicts response:** Patients with both elevated autoantibodies AND low ATP respond to dual therapy; patients with only one abnormality respond to monotherapy
 - Trial design: Stratify by biomarker status; test interaction effects
 - Validates precision medicine approach
3. **Timing prediction:** Early intervention (disease duration <2 years, Stage 1–2) responds to dual therapy; late intervention (duration >5 years, Stage 4–5) requires triple therapy or epigenetic approaches
 - Trial design: Stratify by duration; compare response rates
 - Informs when aggressive intervention is justified
4. **Escape velocity:** Insufficient intervention intensity (e.g., low-dose CoQ10 + LDN) produces transient improvement; adequate intensity (high-dose stack + daratumumab) produces sustained improvement after treatment stops
 - Trial design: Test two dose levels; measure durability at 12 months post-treatment
 - Distinguishes symptomatic relief from cycle-breaking

? Open Question 4: Can Multi-Target Protocols Induce Remission?

The most critical unknown: In established ME/CFS, can any intervention—single or multi-target—truly reverse the disease, or only manage symptoms? If vicious cycles with irreversibility mechanisms are self-sustaining, breaking external drivers may be insufficient.

However, the 60% response to daratumumab [9] with six of ten patients reaching SF-36 PF scores of 80–95 (near-normal function) suggests that at least some patients can achieve sustained remission when the correct driver is targeted. The question is whether non-responders have different active cycles requiring different combinations, or whether they have engaged irreversibility mechanisms requiring more aggressive interventions.

Answering this requires: (1) comprehensive biomarker phenotyping to identify all active cycles, (2) tailored multi-target protocols matched to each patient's cycle profile, (3) long-term follow-up to distinguish temporary improvement from true remission. Until these studies are conducted, the upper bound of treatment efficacy remains unknown.

24.10 CPET-Derived Multi-Target Protocols

The objective demonstration of metabolic failure via two-day cardiopulmonary exercise testing [1] has catalyzed development of novel treatment protocols targeting the specific dysfunctions revealed: autonomic-mitochondrial coupling failure, prolonged recovery kinetics, and exercise-induced oxidative damage. This section presents integrated protocols derived from these findings.

24.10.1 The Autonomic-Metabolic Recovery Protocol

Rationale: Keller et al. identified autonomic dysregulation as the primary driver of Day 2 CPET failure [1]. Walitt et al. documented central catecholamine deficiency [5]. Heng et al. showed cellular ATP depletion [12]. These findings suggest a bidirectional feedback loop: catecholamine deficiency impairs autonomic control → poor tissue perfusion → mitochondrial oxidative stress → catecholamine enzyme damage → worsening autonomic function.

Hypothesis: Breaking this loop requires simultaneous support for both autonomic neurotransmitter synthesis and mitochondrial protection.

Protocol Components

Catecholamine Support (Morning Administration)

- **L-tyrosine:** 1500–3000 mg upon waking (empty stomach for better absorption)
 - Precursor for dopamine and norepinephrine synthesis
 - Lower doses (500–1000 mg) for patients sensitive to stimulation
 - Monitor for anxiety, jitteriness; reduce dose if occurs

- **Cofactor support:**
 - Vitamin B6 (pyridoxal-5-phosphate): 25–50 mg (required for aromatic amino acid decarboxylase)
 - Vitamin C: 1000 mg (required for dopamine β -hydroxylase)
 - Iron: If deficient, supplement to restore ferritin >50–75 ng/mL (required for tyrosine hydroxylase)
 - Copper: 1–2 mg if dietary intake inadequate (required for dopamine β -hydroxylase)
- **Tetrahydrobiopterin (BH4) support** (rate-limiting cofactor):
 - *Option 1:* Sapropterin (prescription BH4) 5–10 mg/kg/day if accessible
 - *Option 2:* Methylfolate 1–5 mg + methylcobalamin 1–5 mg (supports BH4 recycling via DHFR pathway)
 - *Option 3:* 5-MTHF + vitamin C combination (vitamin C regenerates oxidized BH4)

Mitochondrial Protection (Split Dosing)

- **MitoQ** 10–20 mg morning:
 - Mitochondria-targeted ubiquinone conjugated to lipophilic cation
 - Accumulates in inner mitochondrial membrane; scavenges ROS at source
 - Human trials show safety; may be superior to standard CoQ10 for oxidative stress
- **N-acetylcysteine (NAC)** 600 mg twice daily (morning and afternoon):
 - Cysteine donor for glutathione synthesis
 - Established safety profile; FDA-approved for acetaminophen overdose
 - Split dosing maintains glutathione throughout day
- **Alpha-lipoic acid** 300–600 mg morning:
 - Mitochondrial antioxidant; regenerates other antioxidants (glutathione, vitamins C/E)
 - Supports BH4 recycling
 - Use R-lipoic acid form for better bioavailability
- **PQQ (pyrroloquinoline quinone)** 10–20 mg morning:
 - Supports mitochondrial biogenesis via PGC-1 α activation
 - May help replace damaged mitochondria over time

Timing Rationale

- **Morning catecholamine support:** Aligns with natural circadian peak; supports daytime autonomic function
- **Continuous antioxidant coverage:** NAC split dosing; MitoQ has 24-hour residence time
- **Avoid evening stimulation:** Tyrosine/BH4 may impair sleep if taken late

Expected Timeline and Outcomes

- **Weeks 1–2:** Possible initial stimulation from tyrosine; adjust dose as needed
- **Weeks 4–8:** Gradual improvement in PEM recovery time, orthostatic tolerance, cognitive function
- **Weeks 12–16:** If effective, may see improved baseline energy, reduced crash severity, shorter recovery periods
- **Assessment:** Consider repeat two-day CPET at 6 months if accessible to quantify functional improvement

Safety Considerations

- **Contraindications:**
 - Tyrosine: hyperthyroidism, phenylketonuria (PKU), use with MAOIs
 - NAC: active peptic ulcer (theoretical risk), asthma (may trigger bronchospasm in rare cases)
 - BH4/methylfolate: may unmask B12 deficiency; ensure adequate B12 status first
- **Drug interactions:**
 - Tyrosine may potentiate sympathomimetics, thyroid hormones
 - NAC may reduce efficacy of nitroglycerin
 - Alpha-lipoic acid may lower blood glucose; monitor if diabetic
- **Monitoring:** Baseline and periodic blood pressure, heart rate; symptom tracking

Qualification**△ Warning 6: Speculative Protocol**

This protocol is **highly speculative**. While each component has safety data and the mechanistic rationale is sound, the specific combination has not been tested in controlled trials. This represents an experimental approach for patients who have exhausted standard options and are working with knowledgeable physicians. It should not be considered standard of care.

24.10.2 The Mitochondrial Turnover Acceleration Protocol

Rationale: The 13-day recovery period after CPET [1] approximates mitochondrial turnover time in muscle (10–15 days). Hypothesis: exercise-induced ROS damage creates dysfunctional mitochondria that must be physically replaced. Accelerating both removal (mitophagy) and regeneration (biogenesis) might shorten recovery time.

Protocol Components

Mitophagy Enhancement (Evening Dosing)

- **Urolithin A** 500–1000 mg evening:
 - Directly activates mitophagy via PINK1/Parkin pathway
 - Usually derived from gut bacteria converting ellagitannins (from pomegranates/nuts)
 - Direct supplementation bypasses need for microbial conversion
 - Human trials in aging adults show improved mitochondrial function, muscle endurance
 - Proprietary formulation (Mitopure®) has most human safety/efficacy data
- **Spermidine** 1–3 mg evening:
 - General autophagy inducer
 - Found naturally in wheat germ, soybeans, aged cheese
 - Human longevity trials show safety
 - Evening dosing aligns with natural nocturnal autophagy peak
- **Time-restricted eating** (optional, if tolerated):
 - 14–16 hour daily fast (e.g., 7 PM to 9–11 AM)
 - Stimulates autophagy/mitophagy during fasting window
 - CAUTION: Many ME/CFS patients cannot tolerate fasting due to hypoglycemia symptoms
 - Only attempt if already metabolically flexible; discontinue if worsens symptoms

Mitochondrial Biogenesis Support (Morning Dosing)

- **NAD⁺ precursors:**
 - *Option 1:* NMN (nicotinamide mononucleotide) 500–1000 mg morning
 - *Option 2:* NR (nicotinamide riboside) 500–1000 mg morning
 - Activate sirtuins (SIRT1, SIRT3) and PGC-1 α (master regulator of mitochondrial biogenesis)
 - Human trials show NAD⁺ elevation, improved muscle function
 - Morning dosing supports daytime energy metabolism
- **Resveratrol** 200–500 mg morning (optional):
 - SIRT1 activator; synergizes with NAD⁺ precursors
 - Enhances PGC-1 α activity
 - Use micronized formulation for better absorption

Complementary Interventions

- **Resistance training** (if tolerated):

- In healthy individuals, resistance exercise stimulates mitochondrial biogenesis
- In ME/CFS, requires extreme caution: isometric holds (5–10 seconds) below PEM threshold
- Heart rate must stay below AT - 15 bpm
- Frequency: no more than every 3–4 days initially
- This is HIGH RISK; only for stable mild-moderate patients
- **Cold exposure** (if tolerated):
 - Mild cold activates PGC-1 α via β -adrenergic signaling
 - Options: cold showers (gradually progressing from 30 seconds), cryotherapy
 - CAUTION: Cold may exacerbate symptoms in some patients; discontinue if adverse

Expected Timeline

- **Weeks 1–4:** Mitophagy may initially increase fatigue as damaged mitochondria are cleared
- **Weeks 8–12:** Biogenesis begins to dominate; gradual energy improvement
- **Weeks 12–16:** If effective, reduced PEM severity, faster recovery from unavoidable exertion
- **Assessment:** Repeat two-day CPET at 6 months to measure objective improvement in Day 2 performance

Safety and Qualification

- **Safety:** Urolithin A, spermidine, NAD⁺ precursors have human safety data
- **Caution:** Stimulating autophagy requires cellular energy; may initially worsen symptoms in severe patients
- **Recommendation:** Start at low doses (half stated amounts), titrate slowly over weeks
- **Severe patients:** May not tolerate this approach; prioritize stabilization first

△ Warning 7: Experimental Protocol

This protocol is **speculative**. The hypothesis that accelerating mitochondrial turnover will shorten ME/CFS recovery time is logical but unproven. The interventions listed have safety data from other populations but have not been tested specifically for ME/CFS post-exertional recovery.

24.10.3 Post-Exertional Malaise Prevention and Mitigation

Mechanistic Foundation: The 24–72 Hour Window

Post-exertional malaise exhibits a characteristic delayed onset, with symptoms typically peaking 24–72 hours after the triggering activity (see Chapter 2, §2.1 for detailed mechanistic discussion). This delay creates both a challenge and an opportunity: the window between

exertion and symptom onset represents a potential intervention period to abort or mitigate the crash cascade.

The delay mechanisms most amenable to intervention include:

Tier 1: Primary Intervention Targets.

- **ATP depletion cascade:** Cells maintain apparent function during exertion through phosphocreatine and glycolytic buffering, but underlying ATP pools progressively deplete. When ATP falls below approximately 30% of normal (estimated from general cellular bioenergetics) 24–48h post-exertion, catastrophic multi-system failure occurs. *Intervention target:* Provide ATP substrates to prevent threshold crossing.
- **Mitochondrial removal-regeneration gap:** Exercise-induced ROS damage triggers mitophagy (initiation 6–12h, peak overnight), removing damaged mitochondria before biogenesis replaces them (initial recovery 24–72h, complete turnover 10–15 days). During this period, functional mitochondrial mass drops 30–50% (theoretical estimate based on removal-before-replacement dynamics), creating energy crisis. *Intervention target:* Accelerate biogenesis, minimize oxidative damage.
- **NAD⁺ depletion via PARP activation:** DNA damage from exercise activates PARP enzymes, consuming NAD⁺ at 100–1000× normal rates (extrapolated from general PARP biochemistry). Since NAD⁺ is required for ATP synthesis, this creates a vicious cycle reaching critical depletion at 24–72h. *Intervention target:* Provide NAD⁺ precursors immediately.
- **Delayed-type immune activation:** Exercise releases damage-associated molecular patterns (DAMPs) triggering cytokine production that peaks 24–48h post-stimulus in classical DTH patterns. *Intervention target:* Modulate immune activation in the 12–24h window.

★ Key Point: Hypothesis: Dual Pathway Requirement

Anti-inflammatory interventions alone are insufficient for PEM prevention because the core dysfunction is ATP production failure—this follows logically from the mechanism. **Whether energy restoration alone is sufficient, or whether cascade interruption is also required, remains untested.** The hypothesis that both pathways must be addressed rests on two theoretical considerations:

1. **Energy restoration (ATP/NAD⁺ support):** Required because ATP depletion is the initiating driver (mechanistically established)
2. **Cascade interruption (anti-inflammatory/antioxidant):** Hypothetically required because unchecked inflammatory cascades might (a) cause secondary ATP depletion through cytokine-mediated mitochondrial dysfunction, or (b) create symptoms independent of energy status through direct tissue damage

Empirical question: Could ATP/NAD⁺ support alone achieve 60–80% severity reduction, or is anti-inflammatory intervention necessary? This requires comparative trials. Until then, addressing both pathways simultaneously represents the most mechanistic-

cally complete approach.

The Emergency PEM Prevention Protocol

Clinical context: For patients who must undergo unavoidable exertion (medical procedures, essential activities, emergency situations), can targeted interventions immediately post-exertion reduce PEM severity or prevent onset?

Evidence tier: *Hypothesis-driven, mechanistically justified, pending RCT validation.* Individual components have safety data from other contexts. The integrated protocol represents rational polypharmacy targeting identified mechanisms but lacks direct testing in ME/CFS PEM.

Rationale: The 24–72h delay between exertion and symptom peak provides an intervention window. If ATP depletion, mitochondrial damage, NAD⁺ exhaustion, and immune activation initiate the cascade, aggressive multi-target support immediately post-exertion might prevent threshold crossing and abort symptom development.

Phase 1: Immediate Post-Exertion (0–2 Hours). *Goal:* Prevent ATP threshold crossing; minimize oxidative damage; provide NAD⁺ substrate; initiate parasympathetic recovery.

- **ATP substrate provision:**
 - **D-ribose** 10–15 g: Ribose is the sugar backbone of ATP; provides direct substrate for ATP resynthesis. Open-label studies in ME/CFS patients (n=257) report 61% improvement in energy [528]. **CONTRAINDICATION:** Diabetes, hypoglycemia (paradoxically lowers blood glucose)—see warning in Chapter 18
 - **Citrulline-malate** 3–6 g: Dual mechanism—malate replenishes depleted TCA cycle intermediates (documented deficiency in ME/CFS [194]); citrulline supports urea cycle/ammonia detoxification. 31P-MRS evidence shows 34% increased oxidative ATP production [531]
 - **Creatine** 5 g: Buffers ATP via phosphocreatine system; provides immediate energy reserve
 - **Medium-chain triglycerides (MCT oil)** 15–30 mL: Rapidly absorbed, bypasses damaged mitochondrial complexes; provides ketones as alternative fuel
 - Dissolve ribose, citrulline-malate, and creatine in water or juice; take MCT oil separately or mixed in beverage
- **NAD⁺ precursor loading:**
 - **Nicotinamide riboside (NR)** or NMN 1000–2000 mg: High-dose NAD⁺ precursor to prevent PARP-induced NAD⁺ bankruptcy
 - Rationale: PARP activation begins immediately post-exertion; NAD⁺ pools must be maintained to support both DNA repair AND ATP synthesis
- **High-dose antioxidant buffer:**

- **N-acetylcysteine (NAC)** 1200–1800 mg: Glutathione precursor; scavenges ROS. Targets documented 36% brain glutathione deficiency in ME/CFS [193], replicated with 7T MRS [526]. First oral NAC crosses BBB and elevates brain GSH. Pilot data suggest symptom improvement at 1800 mg/day [486]; NIH RCT ongoing (NCT04542161)
- **Vitamin C** 2000–3000 mg: Regenerates other antioxidants; supports BH4 recycling
- **Alpha-lipoic acid** 600 mg: Mitochondrial antioxidant; regenerates glutathione
- Rationale: Exercise-induced ROS damage peaks in first hours; antioxidant support minimizes mitochondrial damage requiring later removal
- **Vagal activation** (parasympathetic recovery):
 - Deep diaphragmatic breathing: 6 breaths/minute for 10–20 minutes
 - Cold water face immersion: 30–60 seconds (triggers dive reflex, potent vagal activation)
 - Transcutaneous auricular VNS device if available: 30–60 minutes
 - Rationale: Activates parasympathetic "rest-and-digest" mode; shifts from sympathetic stress response to recovery state
- **Complete rest protocol:**
 - Horizontal position immediately; no further activity
 - Minimal sensory stimulation (dim lights, quiet environment)
 - No cognitive demands (no reading, screens, conversation)
 - Hydration: 500 mL electrolyte solution (ORS or sports drink)

Phase 2: Early Cascade Prevention (2–24 Hours). *Goal:* Sustain ATP support; accelerate damaged mitochondria removal; prevent immune cascade initiation; maintain antioxidant coverage.

- **Continued ATP substrate provision:**
 - D-ribose 5 g every 4–6 hours (total 15–20 g/day)
 - Citrulline-malate 3 g twice daily (6 g/day total): Maintains TCA cycle substrate availability
 - MCT oil 15 mL twice daily
 - Rationale: Basal metabolism continues consuming ATP; sustained substrate provision prevents progressive depletion
- **Sustained NAD⁺ support:**
 - NR or NMN 500 mg twice daily (morning and early afternoon)
 - Rationale: PARP activity continues during DNA repair phase; NAD⁺ pools must remain adequate
- **Mitophagy enhancement:**
 - **Urolithin A** 500–1000 mg evening: Activates PINK1/Parkin mitophagy pathway; accelerates damaged mitochondria removal

- Rationale: Faster clearance of damaged mitochondria reduces the depth of the functional deficit
- **Sustained antioxidant coverage:**
 - NAC 600 mg twice daily (morning, evening)
 - Vitamin C 1000 mg twice daily
 - Alpha-lipoic acid 300 mg once daily
- **Sleep optimization** (critical for glymphatic clearance):
 - Melatonin 0.5–3 mg 1–2h before bed
 - Magnesium glycinate 300–400 mg evening
 - Prioritize 8–10h sleep opportunity; sleep quality determines glymphatic waste clearance
- **Strict rest enforcement:**
 - No additional physical, cognitive, or emotional exertion
 - Cancel non-essential activities for 24–48h minimum
 - Rationale: Any additional ATP demand during this critical window may push cells over the threshold

Phase 3: Delayed Immune Cascade Window (12–72 Hours). *Goal:* Modulate immune activation; support mitochondrial biogenesis; maintain energy substrates until crisis window passes.

- **Anti-inflammatory support:**
 - **Omega-3 fatty acids** 2–4 g EPA+DHA daily: Reduces pro-inflammatory cytokine production
 - **Curcumin** 500–1000 mg twice daily: NF- κ B inhibition; reduces inflammatory signaling
 - **Optional: Low-dose ibuprofen or naproxen** (if no contraindications): 200–400 mg ibuprofen twice daily or 220 mg naproxen twice daily for 2–3 days
 - Rationale: Immune cascade peaks at 24–48h; anti-inflammatory support during this window may reduce symptom severity
 - CAUTION: NSAIDs alone are insufficient; must be combined with energy support
- **Mitochondrial biogenesis support** (begin Day 2–3):
 - Continue NR/NMN 500 mg twice daily (supports PGC-1 α activation)
 - **PQQ** 20 mg morning: Mitochondrial biogenesis support
 - **Acetyl-L-carnitine** 1000 mg morning: Fatty acid transport; mitochondrial support
 - Rationale: Accelerating biogenesis may shorten the period of functional mitochondrial deficit
- **Continued ATP support** (Days 2–5):
 - D-ribose 5 g 2–3 times daily
 - MCT oil 15 mL once or twice daily

- Adequate hydration: 2–3L fluids daily with electrolytes
- **Vagal toning** (daily):
 - Breathing exercises: 10–20 minutes twice daily
 - Humming/singing (stimulates vagus nerve)
 - Cold exposure if tolerated (brief cold shower, cold pack on neck)
 - Rationale: Sustained parasympathetic activation opposes inflammatory cascades
- **Progressive rest reduction:**
 - Days 1–2: Complete rest, horizontal as much as possible
 - Days 3–5: Gradual return to minimal essential activities only
 - Days 5–7: Resume normal baseline pacing (NOT normal activity—return to pre-exertion energy envelope)
 - Monitor for delayed crash; if symptoms emerge despite protocol, extend rest period

Monitoring and Outcome Assessment

- **Symptom tracking:** Daily ratings (0–10 scale) for:
 - Fatigue/energy level
 - Cognitive function (brain fog, processing speed)
 - Pain/myalgia
 - Orthostatic symptoms
 - Overall functional capacity
- **PEM timeline documentation:**
 - Record symptom onset time (hours post-exertion)
 - Peak severity day
 - Duration until return to baseline
- **Comparative assessment:** If protocol is used multiple times, compare:
 - PEM severity with vs. without protocol
 - Recovery duration with vs. without protocol
 - Proportion of exertions that result in full crashes vs. mitigated symptoms
- **Physiological markers** (if available):
 - Heart rate variability (HRV): Indicates autonomic recovery
 - Resting heart rate: Should return to baseline if crash prevented
 - Orthostatic vital signs: Improvement indicates successful mitigation

Safety Considerations

- **Contraindications:**
 - D-ribose: Diabetes (may lower blood sugar); monitor glucose
 - MCT oil: Severe liver disease; start low dose (15 mL) to assess GI tolerance

- High-dose NAC: Active peptic ulcer (theoretical risk); asthma (rare bronchospasm risk)
- Creatine: Kidney disease (theoretical concern; human data shows safety)
- NSAIDs: GI ulcers, kidney disease, cardiovascular disease, aspirin allergy
- **Drug interactions:**
 - NAC: May reduce nitroglycerin efficacy
 - Alpha-lipoic acid: May lower blood glucose; monitor if diabetic or on diabetes medications
 - Omega-3 fatty acids: Blood thinning effect; use caution with anticoagulants (warfarin, etc.)
 - NSAIDs: Avoid with anticoagulants, other NSAIDs, corticosteroids
 - NR/NMN: Theoretical interaction with PARPi cancer drugs (avoid combination)
- **Tolerability issues:**
 - D-ribose: Sweet taste; some report transient hypoglycemia symptoms (take with food if occurs)
 - MCT oil: GI distress (diarrhea, nausea) if dose too high initially; start 15 mL, increase gradually
 - High-dose NAC: Nausea, sulfur smell/taste; take with food
 - Creatine: Some experience bloating or water retention
- **Cost considerations:**
 - Single emergency use: approximately \$30–50 for all supplements
 - D-ribose (bulk powder): \$25–35 for 250 g (sufficient for multiple uses)
 - NR/NMN (high-dose): \$1.50–3 per 1000 mg dose
 - Other components: \$10–20 for emergency supply
 - More affordable than lost function from severe multi-week crash

Evidence Tier and Qualification

△ **Warning 8: Hypothesis-Driven Protocol Pending Validation**

This protocol is **mechanistically justified but clinically unvalidated**. No randomized controlled trials have tested whether post-exertion interventions reduce ME/CFS PEM severity or duration. However, unlike purely speculative approaches, this protocol targets identified biological mechanisms with plausible intervention windows:

Mechanistic support:

- ATP depletion as PEM driver: Documented in Heng 2025 [12]
- Mitochondrial damage-regeneration gap: Matches 13-day CPET recovery timeline [1]
- NAD⁺ depletion via PARP: Established pathway with ME/CFS abnormalities [12]
- Delayed immune activation: Gene expression shows 24–72h cytokine peaks post-

exercise

Component evidence:

- D-ribose: Small ME/CFS studies show benefit; widely used in sports medicine
- NAD⁺ precursors: Raise NAD⁺ levels in humans; Long COVID trial shows cognitive benefit subset
- Antioxidants: NAC, vitamin C, ALA have safety data; reduce oxidative stress markers
- Mitophagy enhancers: Urolithin A shows mitochondrial benefits in human aging trials
- Anti-inflammatories: Omega-3, curcumin, NSAIDs have established effects

What this protocol is NOT:

- NOT a substitute for pacing (prevention remains superior to mitigation)
- NOT validated by clinical trials in ME/CFS PEM
- NOT appropriate for regular use to enable routine overexertion
- NOT guaranteed to work (individual variation in mechanisms likely)

Appropriate use cases:

- Unavoidable medical procedures (surgery, imaging, emergency care)
- Essential life events (family emergencies, legal obligations)
- Accidental overexertion despite careful pacing
- Situations where crash prevention is critical (e.g., before important medical appointment)

This protocol represents rational therapeutic design based on identified pathophysiology. It is offered for informed patient-physician decision-making in situations where potential benefits outweigh the minimal risks of the interventions and the certain risks of unmitigated severe PEM. Patients using this protocol should document outcomes to contribute to collective knowledge about efficacy.

Future Research Directions

To validate and optimize this approach:

1. **Proof-of-concept RCT:** 60–80 ME/CFS patients randomized to full protocol vs. placebo immediately post-standardized exertion (submaximal CPET or standardized activity). Primary outcome: PEM severity (area under curve, days 1–7). Secondary outcomes: symptom onset timing, recovery duration, proportion with aborted crashes.
2. **Mechanism validation:** Serial biomarkers (ATP/ADP/AMP, NAD⁺/NADH, oxidative stress markers, cytokines) at baseline, immediately post-exertion, +6h, +24h, +48h, +72h in protocol vs. control groups. Correlate biomarker trajectories with symptom outcomes.
3. **Component necessity testing:** Factorial design or systematic component removal to identify which elements are critical vs. redundant. Is ATP support alone sufficient? Must all three phases be present?

4. **Timing optimization:** Test early intervention (0–2h) vs. delayed intervention (6–12h) vs. continuous (0–72h). Identify minimum effective intervention window.
5. **Personalization:** Metabolomics or genetic profiling to identify patient subsets most likely to respond. Are NAD⁺ metabolism SNPs predictive? Does baseline lactate predict D-ribose response?
6. **Real-world effectiveness:** Observational cohort of patients using protocol for unavoidable exertions with detailed symptom tracking. Patient-reported outcomes may precede formal RCTs.

The mechanistic plausibility and low risk profile support early pilot testing. Given the severe impact of PEM on ME/CFS patients' lives, developing validated prevention strategies represents a high-priority research direction.

24.10.4 Personalized Metabolomics-Guided Protocol (Future Direction)

Concept: Use post-exercise metabolomics to identify individual metabolic bottlenecks, then target repletion.

Proposed research protocol:

1. Baseline metabolomics (plasma/serum) before CPET
2. Serial samples: 30 min, 2 hours, 6 hours post-CPET
3. Identify metabolites showing >30% decline
4. Cluster patients by depletion patterns
5. Targeted repletion trial: provide individualized supplementation
6. Measure whether Day 2 CPET deterioration is reduced

Hypothetical examples:

- **Carnitine depletion pattern:** Supplement with L-carnitine 2–3 g/day
- **Glutathione depletion pattern:** Aggressive NAC + glycine + selenium
- **Purine nucleotide depletion:** D-ribose 5–15 g/day + magnesium
- **Tryptophan/kynurenine imbalance:** Consider IDO inhibition (experimental)

Current status: Not clinically available. Metabolomics is expensive and requires specialized facilities. However, if pilot studies show promise, standardized metabolic phenotyping could eventually become accessible.

24.10.5 Clinical Implementation Guidance

Patient Selection

- **Autonomic-Metabolic Protocol:** Mild-to-moderate patients; orthostatic symptoms; cognitive dysfunction

- **Mitochondrial Turnover Protocol:** Patients with severe PEM, prolonged recovery; not for severely affected patients initially
- **Post-Exertion Emergency:** Any severity when unavoidable exertion necessary
- **Metabolomics-Guided:** Research setting only currently

Sequencing

For patients trying multiple approaches:

1. Start with lowest-risk interventions: circadian stabilization, vagal toning, basic antioxidants
2. Add Autonomic-Metabolic Protocol after 4–8 weeks if tolerated
3. Consider Mitochondrial Turnover Protocol after 12 weeks if stable
4. Reserve Post-Exertion Emergency for specific situations

Monitoring and Adjustment

- Symptom diaries: daily ratings of energy, PEM, cognitive function
- Heart rate variability tracking (if accessible): indicates autonomic function improvement
- Functional measures: steps per day, activity duration before PEM
- Blood work: Baseline and 3-month CBC, CMP, iron studies, homocysteine (if using methylated B vitamins)
- Discontinue or reduce dose if: increased anxiety, insomnia, worsening symptoms beyond initial adjustment period

Integration with Standard Care

These protocols complement, not replace:

- Strict pacing (the evidence-based foundation)
- Sleep optimization
- Treatment of comorbidities (POTS, MCAS, etc.)
- Nutritional adequacy
- Psychological support

24.10.6 Research Priorities

To validate and refine these protocols:

1. **Pilot safety trial:** Autonomic-Metabolic Protocol in 20–30 ME/CFS patients; primary outcome: safety and tolerability; secondary: symptom measures at 12 weeks

2. **Mechanistic study:** Serial biomarkers (catecholamines, oxidative stress markers, mitochondrial function assays) before/during/after protocol; correlate with symptom response
3. **Two-day CPET as outcome:** Repeat CPET at 6 months; measure if Day 2 deterioration is reduced in treatment arm vs. control
4. **Metabolomics phenotyping:** Post-exercise metabolomics in 50–100 patients; identify metabolic subgroups; test if subgroup-specific interventions work better than one-size-fits-all
5. **Comparative effectiveness:** Autonomic-Metabolic Protocol vs. Mitochondrial Turnover Protocol vs. combined; which works best for whom?

The two-day CPET provides the objective outcome measure that has long been lacking in ME/CFS research, making these trials feasible and interpretable.

24.11 Evaluating Emerging Therapies

24.11.1 Risk-Benefit Assessment

Experimental therapies vary enormously in risk profile:

- **Low risk:** Breathing exercises, dietary modifications, widely-used supplements
- **Moderate risk:** Prescription medications with established safety profiles, probiotics
- **Higher risk:** Immunosuppressants, invasive procedures, poorly-characterized compounds

24.11.2 Evidence Hierarchy

- **Strongest:** Randomized controlled trials in ME/CFS patients
- **Moderate:** Open-label studies in ME/CFS, RCTs in related conditions
- **Preliminary:** Case reports, mechanistic rationale, patient community reports
- **Speculative:** Theoretical extrapolation from basic science

24.11.3 Access Pathways

- Clinical trials (ClinicalTrials.gov lists ongoing studies)
- Compassionate use / expanded access programs
- Off-label prescription (requires willing physician)
- Medical tourism (significant risks regarding quality and safety)

24.11.4 Reversibility Windows: Setting Realistic Treatment Goals

A critical question for patients evaluating any intervention—experimental or established—is: *What is realistically achievable?* Understanding which pathological mechanisms are reversible versus irreversible guides treatment selection, prevents wasted resources on futile interventions, and enables patients to set appropriate expectations.

The cycle dynamics framework (Chapter 2, §2.1, “Ratchet Effect”) identifies seven irreversibility mechanisms. Not all are equally entrenched, and treatment windows exist where aggressive intervention may reverse seemingly permanent dysfunction. This section provides a mechanism-by-mechanism analysis of reversibility potential.

Time-Dependent Reversibility: The Decay Model

A critical insight is that many mechanisms transition from *reversible* to *irreversible* over time. Acute mitochondrial dysfunction in early disease may be 80% reversible through enhanced biogenesis and repair, but after years of sustained dysfunction, accumulated mtDNA mutations and organelle loss create damage that cannot be fully corrected.

This time-dependency can be modeled as exponential decay:

$$R(t) = R_0 e^{-\lambda t} \quad (24.4)$$

where $R(t)$ = reversibility fraction at time t (proportion of dysfunction that remains correctable), R_0 = initial reversibility (at onset, often 0.8–0.95 for metabolic/immune dysfunction), λ = decay constant (mechanism-specific; estimated 0.1–0.3 yr^{−1} for mitochondrial dysfunction), and t = time since onset (years).

Example 24.1 (Clinical Interpretation of Time-Dependent Reversibility). Consider two patients with comparable mitochondrial dysfunction severity:

- **Patient A:** 6 months of illness, $R(0.5) = 0.9 \times e^{-0.2 \times 0.5} \approx 0.81$ (81% reversible)
- **Patient B:** 5 years of illness, $R(5) = 0.9 \times e^{-0.2 \times 5} \approx 0.33$ (33% reversible)

Even with identical current dysfunction severity, Patient A has >2-fold higher recovery potential. This framework explains why early intervention is critical and why expecting symmetric outcomes across disease durations is unrealistic.

Clinical implication: Duration should factor into prognosis discussions and treatment intensity decisions. Early-stage patients (<2 years) warrant aggressive multi-target intervention; late-stage patients (>10 years) should focus on stabilization and targeting remaining reversible components.

Reversibility Classification: Three-Tier Framework

Tier 1: Highly Reversible (Intervention Window: Weeks to Months). These mechanisms maintain dysfunction through active, ongoing processes that can be interrupted:

Table 24.2: Highly Reversible Pathological Mechanisms

Mechanism	Why Reversible	Intervention	Timeline
GPCR autoantibodies (circulating)	Antibodies cleared in weeks if production stops; not structurally damaging	Immunoadsorption, BC007, daratumumab (plasma cell depletion)	2–6 months
Acute immune activation	Cytokine half-lives measured in hours; no permanent tissue damage if brief	Anti-inflammatory interventions, LDN, omega-3, biologics	2–8 weeks
Metabolic substrate depletion	NAD ⁺ , ATP, cofactors regenerate rapidly if supply provided	NAD ⁺ precursors, D-ribose, CoQ10, B vitamins	2–12 weeks
Functional autonomic dysregulation	Neurotransmitter deficits reversible; no neuronal death	Fludrocortisone, midodrine, L-tyrosine, BH4 support	4–12 weeks
Acute epigenetic changes	Recent histone modifications reverse spontaneously or with intervention	HDAC inhibitors (experimental), NAD ⁺ precursors (sirtuin activation)	3–6 months

Clinical implication: Early aggressive intervention (disease duration <2–3 years) targeting these mechanisms has highest probability of substantial improvement (30–60% function restoration). Example: Daratumumab responders achieving SF-36 scores of 80–95 [9].

Key principle: *The earlier the intervention, the more reversible.* Mechanisms in Tier 1 become Tier 2 (partially reversible) after years of persistence, as secondary irreversible changes accumulate.

Tier 2: Partially Reversible (Intervention Window: Months to Years). These mechanisms involve structural damage or cellular reprogramming, but residual regenerative capacity exists:

Key principle: *Improvement is incremental and slow.* Patients and clinicians must maintain realistic expectations: 20–30% functional improvement is life-changing (severe → moderate severity) but not “cure.”

Tier 3: Irreversible or Minimally Reversible (Intervention Unlikely to Help). These mechanisms involve permanent structural damage, cell death, or deeply entrenched reprogramming beyond current therapeutic reach:

Table 24.3: Partially Reversible Pathological Mechanisms

Mechanism		Partial Reversibility	Intervention	Realistic Gain
Mitochondrial (moderate)	loss	Biogenesis replaces lost mitochondria, but slowly; limited by cellular capacity	PQQ, NAD ⁺ precursors, exercise mimetics (metformin), strict pacing	10–30% improvement over 6–12 months
Established epigenetic silencing		Some changes reversible with aggressive intervention; others locked	HDAC inhibitors (vorinostat—high risk), prolonged NAD ⁺ support	Variable; 0–40% improvement
Microglial (moderate)	priming	Primed microglia can deactivate with sustained anti-inflammatory environment	LDN, omega-3, curcumin, avoid inflammatory triggers	Gradual; 10–25% cognitive improvement over 6–12 months
Central (early)	sensitization	Neuroplasticity allows some desensitization; incomplete reversal	Graded sensory exposure, LDN, avoid opioids, pain psychology	20–40% pain reduction; rarely complete resolution
Muscle	deconditioning	Muscle rebuilds with careful progressive loading below PEM threshold	Isometric exercises, very gradual reconditioning (see Chapter 18)	Return to 50–70% of pre-illness strength (if PEM avoided)

Clinical implication: Modest functional gains (10–40%) achievable even in established disease (3–10 years duration), but require sustained intervention (6–24 months) and risk of PEM during recovery. Example: Bedbound → housebound, or housebound → house-limited with part-time seated work.

Table 24.4: Irreversible or Minimally Reversible Mechanisms

Mechanism	Why Irreversible	Management Strategy
Extensive mitochondrial loss (>50–70%)	Biogenesis capacity overwhelmed; residual mitochondria insufficient to power cell; cell death occurs	Accept functional limits; focus on preventing further loss; maximize function of remaining capacity
High mtDNA mutation burden	Mutations replicate with each division; dysfunctional mitochondria proliferate faster than healthy ones (clonal expansion)	Prevent additional oxidative damage; antioxidant support; strict pacing; no curative intervention exists
Severe central sensitization (chronic >5–10 years)	Permanent cortical reorganization; structural brain changes; glial scar formation	Symptom management only; pain psychology; avoid exacerbating factors; LDN may help but no reversal
Neuronal loss	Neurons do not regenerate; lost cognitive/autonomic function permanent	Compensatory strategies; cognitive aids; environmental modifications; treat residual neuroinflammation to prevent progression
Deep epigenetic locking (>10 years)	Chromatin compaction; DNA methylation stable across cell divisions; difficult to reverse without cell replacement	Experimental only; high-dose HDAC inhibitors (severe side effects); reprogramming strategies (future research)
Autoimmune memory (long-lived plasma cells in bone marrow)	Plasma cells live decades; continuously secrete autoantibodies even if B cells depleted	Plasma cell targeting (daratumumab); otherwise, requires repeated immunoadsorption every 6–12 months indefinitely

Clinical implication: In very severe, long-duration disease (>10–15 years, very severe severity), substantial recovery is unlikely. Realistic goal: stabilization and 5–15% improvement. Prevents false hope from “miracle cure” promises while maintaining motivation for achievable gains. Example: Very severe bedbound → severe bedbound but able to tolerate more visitors, less sensory sensitivity.

Key principle: *Futility recognition prevents harm.* Pursuing aggressive, high-risk interventions (experimental stem cells, overseas clinics) when irreversible mechanisms dominate wastes resources and may worsen condition through travel stress, procedure complications, or false hope followed by despair.

Clinical Decision Framework: Matching Intervention to Reversibility Tier

Step 1: Estimate Reversibility Profile. Based on disease duration, severity, and biomarkers, estimate which tier dominates:

- **High reversibility potential** (Tier 1 dominant):
 - Disease duration <3 years
 - Mild to moderate severity
 - Documented active inflammation (elevated cytokines, autoantibodies)
 - Recent onset or recent worsening (suggests active process, not burnout)
 - Rapid fluctuations (good days vs bad days—indicates functional not structural)
- **Moderate reversibility potential** (Tier 2 dominant):
 - Disease duration 3–10 years
 - Moderate to severe severity
 - Stable dysfunction (few good days; consistent severe symptoms)
 - Some biomarker abnormalities but not extreme
 - History of partial responses to interventions (suggests residual capacity)
- **Low reversibility potential** (Tier 3 dominant):
 - Disease duration >10–15 years
 - Very severe severity (bedbound, severe sensory sensitivity, minimal tolerance)
 - Complete treatment non-response (tried >10 interventions, zero benefit)
 - Evidence of structural damage (brain MRI changes, severe autonomic failure despite treatment, extreme cognitive impairment)
 - Progressive worsening despite aggressive pacing

Step 2: Match Treatment Intensity to Reversibility Tier.

Step 3: Set Realistic Goals by Tier.

- **Tier 1 (high reversibility):** Realistic goal = 30–60% functional improvement, possibly remission
 - Example: Moderate ME/CFS (housebound 3 days/week) → Mild ME/CFS (working part-time, occasional social activities)
 - Aggressive intervention justified; high expected value
- **Tier 2 (moderate reversibility):** Realistic goal = 10–30% functional improvement

Table 24.5: Treatment Intensity by Reversibility Tier

Reversibility Tier	Recommended Approach	Avoid
High (Tier 1)	Aggressive, multi-target intervention; justify moderate-to-high risk for high potential gain; immunotherapy, multi-target metabolic protocols, consider experimental trials	Conservative “wait and see”; missed window may allow transition to Tier 2
Moderate (Tier 2)	Sustained, patient intervention; low-to-moderate risk; long timelines (6–24 months); supplement stacks, LDN, pacing optimization, gradual reconditioning	High-risk experimental interventions with severe side effects (benefit unlikely to justify risk)
Low (Tier 3)	Stabilization and symptom management; prevent further decline; quality of life focus; palliative approach; low-risk supportive care only	Aggressive interventions; travel for experimental treatments; false hope from “miracle cures”; high-risk procedures

- Example: Severe (bedbound 80% of time) → Moderate-severe (housebound, can attend medical appointments, short visits)
- Sustained effort justified; meaningful quality of life gains
- Not “cure,” but life-changing for patient and caregivers
- **Tier 3 (low reversibility):** Realistic goal = stabilization + 5–15% improvement in specific symptoms
 - Example: Very severe (bedbound, severe sensory sensitivity) → Very severe (bedbound, but tolerates 30-minute visitor vs 10 minutes; less severe pain)
 - Focus on quality of life within severe constraints
 - Prevent false hope; validate suffering while maintaining realistic optimism

The “20% Rule”: Why Modest Improvement Is Life-Changing

Patients, families, and even clinicians often underestimate the impact of 15–25% functional improvement in severe ME/CFS:

Observation 99 (Functional Severity Transitions). The severity categories (mild, moderate, severe, very severe) are not linear. Small percentage gains can shift categories:

- **Very severe → Severe** (20% improvement):
 - Before: Bedbound 23 hours/day, cannot tolerate light/sound, requires feeding assistance

- After: Bedbound 18–20 hours/day, can tolerate low light and whispered conversation for 30 minutes, can self-feed soft foods
- Impact: Restores dignity, reduces caregiver burden, enables minimal social connection
- **Severe → Moderate-severe** (25% improvement):
 - Before: Housebound, bedbound 60% of day, cannot attend appointments, washing hair triggers crash
 - After: Can attend medical appointments with wheelchair, shower 2×/week without severe crash, sit upright for meals
 - Impact: Access to medical care, basic hygiene, participates in family life
- **Moderate → Mild-moderate** (30% improvement):
 - Before: Cannot work, grocery shopping triggers 3-day crash, housebound 2–3 days/week
 - After: Part-time remote work (10–15 hours/week), can grocery shop with rest breaks, attends important family events
 - Impact: Financial independence, social participation, purpose and identity beyond illness

The “20% rule”: In severe disease, 20% functional improvement is not “minimal”—it may represent the difference between complete dependence and partial independence, between social isolation and meaningful connection, between hopelessness and purpose.

Patients should be encouraged to pursue interventions with 20% expected benefit, not dismiss them as “not enough.”

Avoiding the Reversibility Trap: When Hope Becomes Harm

△ **Warning 9: The Reversibility Trap**

The reversibility framework creates a paradox: understanding that some mechanisms are reversible may drive patients to pursue increasingly aggressive, risky interventions in pursuit of full recovery—even when their reversibility profile (Tier 3, low potential) predicts minimal benefit.

Red flags for reversibility trap:

- Pursuing experimental treatments overseas despite very severe disease and >15 years duration
- Spending life savings on unproven interventions when Tier 3 profile predicts <10% benefit probability
- Refusing to accept limitations, constantly seeking “one more treatment”
- Severe crashes from travel to clinics, worsening baseline in pursuit of cure
- Family conflict over “giving up” vs realistic goal-setting

Clinical guidance:

- Validate patient's desire for improvement while acknowledging irreversibility realities
- Frame stabilization and modest gains as success, not failure
- Discuss opportunity cost: resources spent on futile high-risk interventions vs quality-of-life improvements (better wheelchair, home modifications, caregiver support)
- Psychological support for grief over permanent losses
- Distinguish "false hope" (unproven promises of cure) from "realistic hope" (achievable 10–20% gains)

Key message: Accepting irreversibility is not "giving up"—it is strategic resource allocation to maximize achievable gains while preventing harm from futile, high-risk interventions.

Future Directions: Expanding Reversibility Windows

Current limitations in reversibility may not be permanent. Research directions that could shift Tier 3 mechanisms to Tier 2:

1. **Epigenetic reprogramming:** Safer HDAC inhibitors, targeted chromatin remodeling, cellular reprogramming factors
2. **Mitochondrial replacement:** Mitochondrial transplantation, induced mitochondrial biogenesis beyond current limits
3. **Neuroplasticity enhancement:** Brain stimulation protocols, neuroplasticity drugs, targeted rehabilitation
4. **Senescent cell clearance:** Senolytics to remove dysfunctional cells blocking regeneration
5. **Stem cell therapies:** True regenerative potential (not current mesenchymal stem cell "immune modulation")

Until these become clinically validated, the reversibility framework as outlined represents current realistic boundaries.

★ Key Point: Reversibility-Guided Treatment Philosophy

The reversibility windows framework generates a treatment philosophy:

1. **Early aggressive intervention** (Tier 1): Justify moderate-to-high risk for high potential gain; don't "wait and see"
2. **Sustained moderate intervention** (Tier 2): Patient, low-risk, long-timeline approaches; 10–30% gains are meaningful
3. **Stabilization and quality of life** (Tier 3): Prevent harm from futile interventions; accept limitations; maximize achievable gains
4. **Always:** Pacing as disease-modifying therapy across all tiers

This framework prevents two harmful extremes:

- **Therapeutic nihilism:** “Nothing works, don’t bother trying” (ignores Tier 1 and 2 reversibility)
- **Unrealistic optimism:** “Full recovery is always possible with enough effort” (ignores Tier 3 irreversibility)

The middle path: Evidence-based hope matched to individual reversibility potential.

Observation 100 (The Desperation-Exploitation Gradient). Severe, treatment-resistant illness creates vulnerability to exploitation. The ME/CFS patient community has been targeted by:

- Unproven stem cell treatments at overseas clinics
- High-cost “personalized medicine” protocols with little evidence
- Supplements with exaggerated claims
- Practitioners promoting theories rejected by mainstream medicine

While maintaining openness to novel approaches, patients should apply skepticism proportional to claims, cost, and risk. Red flags include: guarantees of cure, pressure to commit quickly, inability to provide outcome data, and hostility to questions.

24.11.5 Quantitative Cycle Gain Measurement: A Prognostic Tool

The vicious cycle framework (Chapter 2, §2.1) identifies “cycle gain” as the amplification factor within each pathophysiological loop. When $G > 1$, each iteration amplifies dysfunction; when $G < 1$, the system naturally dampens toward baseline. Quantifying individual patients’ cycle gain could transform prognosis and treatment planning.

The Cycle Gain Concept

~ Hypothesis 6: Cycle Gain as Prognostic Biomarker

Recovery kinetics following standardized exertion may predict long-term disease trajectory.

Theoretical definition: The cycle gain G represents net amplification across a vicious cycle. In control theory, for a feedback loop with n sequential processes, the net loop gain is the product of individual gains:

$$G = \prod_{i=1}^n g_i$$

where g_i is the gain of each component process (ROS generation, mitochondrial damage, biogenesis impairment, etc.). This multiplicative relationship is standard in feedback systems analysis [608].

Critical threshold: $G = 1$ represents a bifurcation point:

- $G < 1$: System naturally dampens perturbations (stable, self-correcting)

- $G > 1$: Perturbations amplify with each iteration (unstable, self-reinforcing)

Proposed clinical proxy: We hypothesize that recovery time after standardized exertion *correlates with* (but does not define) cycle gain. This correlation requires empirical validation. The mapping between recovery time and G proposed in Table 24.6 is a testable hypothesis, not an established relationship.

Evidence Grade: C (control theory foundation established; ME/CFS application hypothetical)

△ **Warning 10: Avoiding Circular Reasoning**

Cycle gain (G) is a theoretical construct representing feedback loop amplification. Recovery time is a proposed *proxy measure* that may correlate with G . The hypothesis is falsifiable: if recovery time does not predict long-term trajectory or treatment response, the proxy is invalid regardless of whether the underlying cycle gain concept is correct.

Proposed Measurement Protocol

A standardized protocol to measure cycle gain:

Baseline Assessment (Day 0).

- Blood draw: ATP, ADP, AMP, lactate, pyruvate, NAD^+/NADH ratio
- Symptom assessment: Bell Disability Scale, 7-day symptom diary
- Functional status documentation

Standardized Exertion (Day 1–2).

- Two-day cardiopulmonary exercise testing (2-day CPET) per Vermeulen protocol
- Alternative for milder cases: 6-minute walk test at self-selected “comfortable” pace

Recovery Kinetics (Days 1–28).

- Blood draws: +30 min, +2h, +24h, +48h, +7 days, +14 days, +28 days
- Daily symptom diaries
- Primary outcome: Time to return to 90% of baseline on Bell Scale

Proposed Cycle Gain Proxy. A clinical proxy for cycle gain could be derived from recovery dynamics:

$$G_{\text{proxy}} = \frac{\text{Peak symptom severity} \times \text{Recovery time}}{\text{Exertion magnitude} \times \text{Baseline function}}$$

Rationale: This formula captures the intuition that higher cycle gain produces both more severe symptoms and longer recovery from equivalent exertion. However, this is a *proposed* metric requiring validation—the specific functional form is not theoretically derived and alternatives (logarithmic, threshold-based) may prove superior.

Calibration: Healthy control reference values must be established empirically. We hypothesize $G_{\text{healthy}} \approx 0.2\text{--}0.4$ based on typical post-exercise recovery (1–3 days), but this requires measurement in validation studies.

Clinical Interpretation

Table 24.6: Proposed Cycle Gain Classification and Prognosis (Hypothetical—Requires Validation)

Cycle Gain (Proxy)	Recovery Time	2-Year Prognosis*	Treatment Implications
$G < 0.7$	<5 days	Higher recovery potential	Conservative management; emphasize pacing
$G = 0.7\text{--}1.0$	5–10 days	Moderate recovery potential	Moderate intervention; address reversible components
$G = 1.0\text{--}1.5$	10–21 days	Lower recovery potential	Aggressive multi-target therapy; break cycle amplification
$G > 1.5$	>21 days	Poor recovery potential	Intensive intervention required; consider experimental protocols

*Specific prognosis percentages removed as they were not empirically derived. The qualitative relationship (longer recovery → poorer prognosis) is hypothesized based on disease chronicity patterns [54] but the quantitative mapping to G values requires prospective validation. **Evidence Grade: D** (entire classification scheme is theoretical).

△ Warning 11: Testing Risk

The 2-day CPET required for accurate cycle gain measurement may trigger prolonged crashes in susceptible patients. Protocol safeguards:

- Exclude very severe patients (Bell <20) from testing
- Provide 2-week rest period post-testing with monitoring
- Informed consent must communicate potential for temporary worsening

Research Priorities

- 1. **Validation study:** Prospective cohort ($n \geq 100$) correlating cycle gain measurement with 2-year outcome
- 2. **Biomarker correlation:** Does ATP/ADP ratio change predict clinical recovery time? ($r > 0.5$ expected)
- 3. **Treatment response prediction:** Do patients with $G < 1$ show $>3\times$ higher response rate to interventions?
- 4. **Simplified protocol:** Develop point-of-care ATP test + 14-day symptom diary for primary care use

Evidence Grade: D (proposed methodology; requires validation)

24.11.6 Epigenetic Reversal Strategies

~ Hypothesis 7: Epigenetic Lock-In in Chronic ME/CFS

In severe, prolonged ME/CFS, epigenetic modifications may “lock in” metabolic dysfunction even after original triggers resolve. DNA methylation of mitochondrial biogenesis genes (PGC-1 α , TFAM) and histone modifications maintaining closed chromatin at metabolic loci could create self-perpetuating pathological states.

Supporting evidence: Epigenetic changes have been documented in ME/CFS, including altered DNA methylation patterns in immune cells [609]. However, whether these changes are causal (driving dysfunction) or consequential (downstream of other pathology) remains unknown.

Evidence Grade: D (epigenetic changes documented; causal role in ME/CFS pathology not established)

Targeting these putative epigenetic locks represents a potential approach for treatment-refractory patients, though this strategy is highly speculative.

Tiered Epigenetic Intervention Framework

Table 24.7: Tiered Epigenetic Reversal Strategies

Tier	Population	Intervention	Risk Profile	Evidence
1	All ME/CFS	NAD ⁺ precursors (sirtuin activation)	Low	C
2	Moderate-severe, failed treatments	NAD ⁺ + metformin + resveratrol	Low-moderate	D
3	Very severe, refractory	HDAC inhibitors (research only)	High	D

Tier 1: Sirtuin Activation (Low Risk)

NAD⁺-dependent sirtuins (SIRT1, SIRT3) deacetylate histones and transcription factors, potentially reversing epigenetic silencing of metabolic genes.

Protocol:

- NR or NMN: 500–1000 mg/day
- Mechanism: Restore NAD⁺ to support sirtuin function
- Duration: ≥12 weeks for epigenetic effects

Evidence Grade: C (NAD⁺ restoration documented; epigenetic reversal in ME/CFS not directly tested)

Tier 2: Combined Epigenetic Modulation (Moderate Risk)

For patients failing Tier 1:

Protocol:

- NAD⁺ precursor (NR 1000 mg/day)
- Metformin 500 mg BID (AMPK activation → PGC-1 α induction)
- Resveratrol 500 mg/day (sirtuin activation)
- Duration: 16 weeks

Monitoring: Blood glucose, lactate, liver function at baseline and 8 weeks.

Evidence Grade: D (theoretical combination; safety and efficacy require study)

Tier 3: HDAC Inhibitors (Research Stage Only)

△ **Warning 12: HDAC Inhibitor Caution**

HDAC inhibitors (vorinostat, butyrate, valproic acid) are cancer drugs with significant toxicity. Their use in ME/CFS is **entirely experimental** and should occur **only in research settings** with full ethics approval and intensive monitoring.

Risk profile:

- Thrombocytopenia (25–50%)
- Fatigue/asthenia (30–60%)—paradoxical concern in ME/CFS
- GI symptoms (30–50%)
- Cardiac effects (QTc prolongation, 5–10%)

Appropriate only for: Very severe, treatment-refractory patients (Bell <20) who have failed all other approaches, with full informed consent acknowledging experimental nature and risks.

Mechanism and Rationale**~ Hypothesis 8: Mathematical Model of Epigenetic Lock-In**

The epigenetic lock-in concept can be formalized as a dynamical system where methylation state becomes self-reinforcing:

$$\frac{dM}{dt} = \alpha(1 - M) - \beta M \cdot f(S)$$

where M = methylation level at metabolic genes (0–1), α = de novo methylation rate, β = baseline demethylation rate, and $f(S)$ = disease-state-dependent modifier of demethylation efficiency.

Model interpretation: When disease severity S is high, $f(S) \rightarrow 0$, suppressing demethylation and maintaining hypermethylation at a stable equilibrium $M^* \approx 1$. Recovery requires either reducing S (breaking other disease mechanisms) or pharmacologically increasing β (epigenetic intervention).

Limitations: This is a conceptual model, not a quantitative prediction. Parameter values (α , β , functional form of f) are unknown. The model illustrates the *logic* of epigenetic lock-in but cannot predict specific outcomes without empirical calibration.

Evidence Grade: D (theoretical construct; mathematical form is illustrative, not validated)

Epigenetic interventions aim to shift this balance by:

- Enhancing demethylation (sirtuin activation, TET enzyme support)
- Reducing methylation (DNMT inhibition—dietary: green tea EGCG, curcumin)
- Opening chromatin (HDAC inhibition—research only)

Research Priority

A cautious Phase 1b/2a study:

Phase 1 (Tier 2 combination):

- Population: ME/CFS, moderate-severe (Bell 20–40), failed ≥ 3 treatments, $n = 30$
- Intervention: NAD⁺ precursor + metformin + resveratrol, 16 weeks
- Primary outcome: Safety and tolerability
- Secondary: Clinical response, methylation array changes

Phase 2 (HDAC inhibitor pilot, only if Phase 1 safe and shows epigenetic effect):

- Population: Very severe (Bell < 20), failed all treatments, $n = 10$
- Intervention: Low-dose vorinostat (200 mg, half of cancer dosing), 8 weeks
- Intensive monitoring: Weekly CBC, LFTs, ECG
- Primary outcome: Safety

Evidence Grade: D (entirely theoretical; requires careful staged investigation)

25 Integrative and Personalized Treatment Approaches

25.1 Developing a Treatment Plan

Treatment planning in ME/CFS may differ fundamentally from other chronic conditions. The standard medical approach—try a treatment, observe the response, adjust—assumes that failed trials carry manageable costs. In ME/CFS, this assumption is potentially dangerous. Each treatment trial imposes metabolic demands on an already energy-depleted system, and failed trials can trigger post-exertional malaise severe enough to cause permanent functional decline (see Section 18.3.3 on crash severity dose-response). For severe and very severe patients, there may be only a narrow margin for error before irreversible deterioration occurs.

This section provides a safety-first framework for treatment planning, integrating the energy cost of treatments themselves into the decision-making process.

25.1.1 The Imperative of Treatment Safety in ME/CFS

△ Warning 1: Treatment Trials as Energy Gambles

Every medication or supplement trial in ME/CFS carries an energy cost independent of the treatment's intended effect. Processing any exogenous substance requires ATP for hepatic metabolism (CYP450 system), renal clearance, protein binding, receptor adaptation, and immune surveillance. In a system with impaired mitochondrial ATP production—as demonstrated by two-day cardiopulmonary exercise testing [1, 140], mitochondrial function assays [610], and metabolomic profiling [611, 612]—this processing overhead can itself trigger post-exertional malaise.

The clinical consequence: A treatment that is safe and well-tolerated in healthy individuals may provoke a crash in a severe ME/CFS patient—not because of pharmacological adverse effects, but because the body cannot afford the metabolic cost of processing the substance.

Illustrative clinical observations:

- Standard-dose pyridostigmine (60 mg) causing severe prostration in a patient who later tolerated 20 mg (Section 21.10.1)
- IVIG infusions triggering prolonged crashes, consistent with the substantial immune activation and protein processing demands of large immunoglobulin infusions
- Multiple supplement initiation simultaneously overwhelming hepatic processing

capacity (see sequencing rationale in Section 25.1.6)

Critical principle: Treatment trials are not free. Each one consumes finite energy reserves. In severe patients, the number of tolerable failed trials may be very small before permanent deterioration occurs.

25.1.2 Treatment Energy Categories

Observation 101 (Treatment Energy Cost Classification). To guide treatment selection in energy-depleted patients, we classify all ME/CFS interventions by their net energy impact:

Category A — Net energy providers: Treatments that directly supply ATP substrates, electron transport chain components, or essential cofactors for energy production. Net positive energy balance. These treatments give the body what it lacks. Category A does not mean risk-free—GI intolerance, allergic reactions, and supplement interactions remain possible—but the *energy* cost is minimal to net positive. *Examples:* D-ribose, CoQ10, NADH/NAD⁺ precursors, creatine, L-carnitine, magnesium, B vitamins, MCT oil.

Category B — Energy-neutral: Treatments with minimal metabolic processing demands that neither provide nor consume significant ATP. Therapeutic effects operate through pathways that do not substantially burden energy metabolism. *Examples:* Most antihistamines, melatonin, PEA, LDN, Ginkgo biloba, probiotics.

Category C — Energy-demanding: Treatments requiring significant ATP for hepatic CYP450 metabolism, renal clearance, immune activation, or protein processing. Net negative energy balance in depleted systems. The absolute magnitude of this metabolic cost has not been quantified in ME/CFS; however, the clinical observation that severely depleted patients crash from treatments well-tolerated by healthy individuals suggests the cost is meaningful relative to their reduced energy reserves. These treatments may be highly effective but impose a metabolic cost that must be budgeted. *Examples:* Valganciclovir, rituximab, IVIG, aripiprazole.

Most medications and supplements in Chapters 21 and 22 include an Energy Profile paragraph specifying their category and mechanistic rationale.

Given that Category A treatments supply energy substrates while Category C treatments consume them, and that ME/CFS patients have impaired energy production, a logical sequencing principle emerges: **start with Category A (energy providers), stabilize, then cautiously introduce Category B (neutral), and reserve Category C (energy-demanding) for cases where phenotyping strongly predicts benefit** (see Section 5.6.10 for phenotyping protocols).

25.1.3 Baseline Assessment

Before initiating any treatment, establish:

1. **Functional status:** Bell Disability Scale score, hours upright, activity tolerance threshold

2. **Energy envelope:** Heart rate monitoring or subjective energy tracking to quantify available reserves (see Chapter 23)
3. **Current medication inventory:** All active medications, supplements, and their energy categories—the total metabolic processing load matters
4. **Comorbidity mapping:** Identify Septad components (Section 5.6.11) and prioritize by symptom burden
5. **Phenotype identification:** Determine dominant subtype (immune-predominant, metabolic-predominant, neurological-predominant, autonomic-predominant) using tiered assessment (Section 5.6.10)
6. **Treatment history:** Prior adverse reactions, paradoxical responses, pharmacogenomic data if available
7. **Severity classification:** Mild, moderate, severe, or very severe—this determines the safety margin for treatment trials

25.1.4 The Micro-Dosing Imperative

△ **Warning 2: Never Start at Standard Doses in ME/CFS**

ME/CFS patients—especially severe and very severe—must **never** be started on standard medication doses. The paradoxical reactor phenotype (Section 21.10.1) affects a subset of patients—prevalence unknown, but frequently reported in clinical practice and patient communities—and even patients who are not paradoxical reactors have reduced metabolic processing capacity.

Universal rule: Start at 1/4 to 1/10 of the standard starting dose. Titrate upward at minimum 1–2 week intervals. Monitor daily for the first 2 weeks of any new treatment.

Energy profiles for action-plan medications. The following treatments appear in Table 25.1 but are covered in action-plan chapters (Chapters 17–18) rather than in the mechanism-targeting chapters. Their energy profiles are summarized here for completeness.

- **Trazodone** (Category B, Low crash risk): Serotonin antagonist and reuptake inhibitor (SARI) used for sleep. At ME/CFS micro-doses (12.5–25 mg), CYP3A4/CYP2D6-mediated first-pass metabolism is expected to be reduced at these doses (pharmacokinetic inference), placing the metabolic burden within the energy-neutral range.
- **Amitriptyline** (Category B, Low crash risk): Tricyclic antidepressant used at sub-antidepressant doses (5 mg) for pain and sleep; also inhibits proinflammatory mast cell mediator release [242], an additional therapeutic benefit that does not add to metabolic burden. CYP2D6/CYP2C19 metabolic burden is expected to be reduced at this dose in extensive metabolizers (pharmacokinetic inference); CYP2D6 poor metabolizers (approximately 5–10% of European-ancestry populations; standard pharmacology) may require further dose reduction. Anticholinergic side effects (dry mouth, constipation) are the primary concern at these doses, not energy cost.
- **Fludrocortisone** (Category B–C, Medium crash risk): Synthetic mineralocorticoid used in orthostatic intolerance [613, 338]; expected mechanism is plasma volume expansion

Table 25.1: Micro-dosing reference: ME/CFS starting doses versus standard doses

Treatment	Standard dose	ME/CFS start	Energy cat.	Crash risk
<i>Immune-modulating</i>				
LDN	4.5 mg	0.5 mg	B	Medium
IVIG	Weight-based	50% standard rate	C	High
Cimetidine	400 mg BID	200 mg daily	B–C	Low
<i>Antiviral</i>				
Valacyclovir	1000 mg BID	500 mg daily	B–C	Medium
Valganciclovir	900 mg BID	450 mg daily	C	High
<i>Autonomic</i>				
Pyridostigmine	60 mg TID	15–20 mg daily	B	Low
Fludrocortisone	0.1 mg daily	0.05 mg daily	B–C	Medium
Midodrine	10 mg TID	2.5 mg daily	B–C	Medium
<i>Neuroactive (incl. herbal)</i>				
Aripiprazole	10–30 mg	0.5–1 mg	C	High
Trazodone	50–100 mg	12.5–25 mg	B	Low
Amitriptyline	25–50 mg	5 mg	B	Low
Ginkgo biloba	120–240 mg	40–60 mg	B	Very low
<i>Mitochondrial and metabolic support</i>				
CoQ10 (ubiquinol)	200–300 mg	50–100 mg	A	Very low
D-Ribose	15 g daily	2.5 g daily	A [†]	Very low
NR/NMN	300–500 mg	100–150 mg	A [†]	Very low
L-Carnitine	1500–3000 mg	500 mg	A	Very low
NAC	1200 mg daily	300–600 mg daily	A–B	Very low
Magnesium	400–600 mg	100–200 mg	A	Very low

Energy categories: A = net energy provider, B = energy-neutral, C = energy-demanding. [†]Category A assignment based on proposed energy-supportive mechanisms; clinical evidence in ME/CFS is preliminary.

Crash risk reflects both energy cost *and* pharmacological adverse effect potential—a Category B treatment can carry elevated crash risk if it has significant non-energy adverse effects (e.g., LDN carries Medium crash risk despite Category B energy status, due to psychiatric reaction risk in paradoxical reactors).

Severe/very severe patients: use the lower end of ME/CFS starting doses.

via renal sodium retention (standard mineralocorticoid pharmacology). Energy cost is typically highest during the initial adaptation period (estimated first 2–4 weeks; pharmacodynamic inference), when electrolyte rebalancing (potassium monitoring is required; standard mineralocorticoid pharmacology) and cardiovascular volume adjustment are most active. Severe patients may not tolerate this period, requiring slower titration or alternative agents. At steady state, the ongoing mineralocorticoid effect is expected to impose minimal additional metabolic demand.

- **Midodrine** (Category B–C, Medium crash risk): Alpha-1 agonist for orthostatic intolerance [338]. Energy cost is expected to arise primarily from sustained peripheral vasoconstriction and the associated cardiovascular compensation (pharmacodynamic inference). The resulting energy demand is expected to be modest but continuous; supine hypertension is a documented pharmacological risk requiring monitoring (standard pharmacology).

25.1.5 Crash-Risk Ranking and Decision Rules

The following ranking covers the principal ME/CFS treatments discussed in this document, including those beyond Table 25.1: immunotherapies from Chapter 21 (e.g., rituximab), emerging treatments from Chapter 24 (e.g., immunoadsorption), and supplements from Chapter 22.

The ranking uses three tiers; Table 25.1 uses a finer four-level scale. The mapping: “Highest” corresponds to table “High” entries; “Moderate” to “Medium” entries; and “Low” encompasses both “Low” and “Very low” table entries. For treatments appearing in the table, the crash-risk column is the definitive tier assignment; for treatments not in the table, the ranking below assigns the tier directly. Energy category alone does not determine tier placement (e.g., cimetidine is Category B–C but Low crash risk due to its CYP inhibition properties, which are expected to reduce hepatic metabolic load in polypharmacy (pharmacokinetic inference)).

Highest crash risk (Category C, complex immune/metabolic effects):

- **Rituximab**: Anti-CD20 monoclonal antibody; high metabolic burden from B-cell depletion and subsequent immune remodeling (mechanistic inference)
- **IVIG**: Pooled immunoglobulin infusion; high metabolic burden from large protein load and immune modulation response (mechanistic inference)
- **Valganciclovir**: Antiviral nucleoside analogue; high metabolic burden from bone marrow suppression and hepatic/renal clearance (established adverse effects; energy burden by mechanistic inference)
- **Aripiprazole**: Atypical antipsychotic at micro-doses (0.5–1 mg, clinical outcomes reported in [602]); high metabolic burden from CYP2D6/3A4 metabolism (standard pharmacology) and hypothesized dopaminergic activation sensitivity (proposed mechanism)
- **Immunoadsorption**: Extracorporeal autoantibody removal; high metabolic burden from haemodynamic stress of apheresis and subsequent transient immune disruption (mechanistic inference)

Moderate crash risk (Category B–C, or Category B with paradoxical potential):

- **LDN:** Proposed TLR4 antagonist and glial modulator (proposed mechanisms); minimal metabolic burden (moderate crash risk driven by psychiatric reaction potential in paradoxical reactors, not metabolic cost)
- **Valacyclovir:** Antiviral prodrug; moderate metabolic burden from sustained hepatic and renal clearance (mechanistic inference)
- **Fludrocortisone:** Synthetic mineralocorticoid for orthostatic intolerance; moderate metabolic burden from mineralocorticoid-mediated metabolic adaptation (pharmacodynamic inference)
- **Midodrine:** Alpha-1 agonist for orthostatic intolerance; moderate metabolic burden from cardiovascular adaptation (pharmacodynamic inference)
- **5-HTP:** Serotonin precursor; moderate metabolic burden from serotonergic effects (mechanistic inference), with possible mast cell activation in MCAS-susceptible patients (proposed mechanism)
- **Pregnenolone:** Neurosteroid precursor; moderate metabolic burden from hepatic CYP-mediated steroidogenesis (mechanistic inference)

Low crash risk (predominantly Category A–B; includes higher-category exceptions with energy-sparing properties per mapping note above). Within this tier, “negligible” indicates no significant hepatic processing (e.g., direct cofactors, luminal-acting agents), while “minimal” indicates modest but documented enzymatic metabolism—whether CYP-mediated, hepatic conjugation, renal clearance, or systemic enzymatic processing (regardless of whether the substance is a pharmaceutical or supplement):

- **Magnesium, electrolytes:** Essential ion cofactors (ATP synthase, membrane potential); negligible metabolic burden
- **CoQ10:** Electron shuttle between ETC Complexes I/II and III; negligible metabolic burden
- **NAD⁺ precursors (NR/NMN):** NAD⁺ pool replenishment, supporting NADH generation via glycolytic and TCA-cycle dehydrogenase reactions (proposed mechanism for energy benefit); negligible metabolic burden
- **NADH:** Direct electron donor to Complex I of the electron transport chain; negligible metabolic burden
- **D-ribose:** ATP backbone sugar supporting purine nucleotide resynthesis (proposed mechanism); negligible metabolic burden
- **L-carnitine:** Mitochondrial fatty acid transport cofactor (carnitine palmitoyltransferase system); negligible metabolic burden
- **Creatine:** Phosphocreatine energy buffer for rapid ATP regeneration; negligible metabolic burden
- **PQQ:** Proposed mitochondrial biogenesis activator (preclinical; clinical evidence lacking); negligible metabolic burden
- **MCT oil:** Ketone precursor; absorbed via portal vein (bypassing lymphatic transport) and oxidized without carnitine-dependent mitochondrial entry (C8/C10 MCTs; C12 is partially CPT1-dependent); negligible metabolic burden
- **NAC:** Glutathione precursor and direct thiol antioxidant (cysteine donor via deacetylation, primarily hepatic and cellular); minimal metabolic burden

- **ALA:** Mitochondrial antioxidant and heavy metal chelator; minimal metabolic burden
- **Omega-3 (EPA/DHA):** Anti-inflammatory fatty acids; minimal metabolic burden
- **Curcumin, quercetin:** Polyphenol anti-inflammatories; minimal metabolic burden
- **B vitamins, vitamin D:** Essential metabolic cofactors; minimal metabolic burden
- **Taurine, glycine, glutamine:** Amino acid supplements; negligible metabolic burden
- **Zinc:** Essential trace element (direct cofactor); negligible metabolic burden
- **Probiotics:** Gut microbiome modulation; negligible metabolic burden
- **PEA:** Endocannabinoid-related lipid mediator (glial modulation via PPAR- α ; proposed mechanism); minimal metabolic burden
- **Melatonin:** Circadian rhythm regulator and antioxidant; minimal metabolic burden
- **Resveratrol:** Polyphenol with proposed sirtuin-activating properties (preclinical; clinical relevance uncertain); minimal metabolic burden
- **H1 antihistamines (cetirizine, loratadine, fexofenadine):** Histamine receptor blockers for MCAS symptom management; minimal metabolic burden
- **Cimetidine:** H2 antagonist with CYP inhibitory properties (in polypharmacy, may reduce total hepatic metabolic load of co-administered drugs; see Chapter 21); minimal metabolic burden
- **Pyridostigmine:** Acetylcholinesterase inhibitor; minimal metabolic burden
- **Trazodone:** SARI used for sleep at micro-doses (12.5–25 mg); minimal metabolic burden
- **Amitriptyline:** Tricyclic at sub-antidepressant dose (5 mg) for pain and sleep; minimal metabolic burden
- **Ginkgo biloba:** Mild vasodilator and antioxidant; minimal metabolic burden

△ **Warning 3: The Paradox of Effective Treatments**

The treatments associated with the most pronounced reported responses in biomarker-defined subgroups (rituximab, immunoadsorption, valganciclovir; see Chapters 21 and 24) are also the most energy-demanding. For severe patients, this creates a fundamental dilemma: the interventions that could potentially restore function require an energy investment the patient may not be able to afford. This paradox is the strongest argument for **phenotype-guided treatment selection** (Section 25.3.3)—reserving Category C treatments for patients whose biomarker profile predicts a high probability of response, rather than empirical trial-and-error.

25.1.6 Prioritizing Interventions

The following treatment introduction sequence is recommended for ME/CFS, particularly in moderate-to-severe patients, based on energy-cost principles (evidence-based guidelines for this sequencing are not yet available):

1. **Foundation first** (Weeks 1–4): Electrolytes, magnesium, sleep optimization (behavioral; no pharmacological energy category applies). Pharmacological items are all Category A or B. Address the most basic physiological deficits before anything else.

2. **Energy substrate support** (Weeks 5–8): CoQ10, B vitamins, then one additional mitochondrial support (D-ribose or NR/NMN). All Category A (D-ribose and NR/NMN carry the A⁺ qualifier; see Table 25.1 footnote). The rationale is to build energy reserves before introducing treatments that consume energy (proposed sequencing principle).
3. **Symptomatic relief** (Weeks 9–12): Energy-neutral treatments targeting dominant symptoms—PEA for pain, melatonin for sleep, antihistamines for MCAS (all Low crash risk per ranking above). Minimal energy cost.
4. **Targeted interventions** (After stabilization): Treatments requiring subtype identification, spanning Category B through B–C depending on the agent. LDN for immune/neuroinflammatory subtype (Category B, but Moderate crash risk from paradoxical reactions). Pyridostigmine for autonomic subtype (Category B, Low crash risk). Antivirals for viral-immune subtype (Category B–C).
5. **High-cost interventions** (Only with phenotyping): Category C treatments (IVIG, valganciclovir, immunomodulators) only when biomarker evidence strongly predicts benefit and the patient has adequate energy reserves from Category A foundation.

See Section 22.7 in Chapter 22 for cost-stratified supplement protocols that align with this sequence.

25.1.7 When to Stop Trying

Directive 3 (Treatment Trial Limits for Severe Patients). **Certainty: 0.30** (clinical reasoning based on crash dose-response data and paradoxical reactor observations; no RCTs on optimal trial limits)

For severe and very severe ME/CFS patients, the following decision rules may prevent treatment-induced deterioration:

Stop-and-stabilize criteria (thresholds are proposed clinical defaults, not evidence-based cut-offs; clinicians should adjust based on individual patient trajectory):

- **After any treatment-induced crash:** Do not attempt another new treatment until baseline function is fully restored (minimum 4 weeks at pre-crash level)
- **After two consecutive treatment-induced crashes:** Pause all treatment experimentation for minimum 3 months. Focus exclusively on pacing and Category A energy support. *Rationale:* Two consecutive crashes suggest either a pattern of global medication intolerance or inadequate recovery time between trials; continuing risks cumulative decline
- **After three failed Category C trials:** Reassess the treatment approach entirely. The patient may be a global paradoxical reactor, and further empirical trials carry unacceptable risk. *Rationale:* Three Category C failures represent substantial cumulative energy expenditure with no return; if failures reflect a systemic intolerance pattern, each additional trial is unlikely to succeed and risks further cumulative harm
- **If functional capacity has declined >20% since treatment trials began:** Immediate cessation of all experimentation. Stabilization and energy restoration take absolute priority. *Rationale:* A 20% decline is chosen as the threshold at which iatrogenic harm

clearly outweighs any remaining potential benefit from empirical trials; this corresponds roughly to losing one severity tier (e.g., moderate to severe)

Decision threshold: For very severe (bedridden) patients, apply a more conservative standard—stop after *any single* treatment-induced functional decline lasting more than 1 week. The margin for error in very severe disease is extremely narrow.

25.1.8 Medication-Induced Crash Recovery Protocol

When a treatment trial triggers a crash:

Step 1 — Immediate actions (within hours):

- Discontinue the offending treatment immediately
- Do not attempt to “push through” in hopes the reaction will resolve
- Implement strict activity restriction (reduce to 50% of pre-crash activity level)

Step 2 — Energy support (first 24–72 hours):

- Maintain or increase Category A supplements (electrolytes, magnesium, CoQ10, D-ribose)
- Consider the post-exertional emergency protocol (Section 24.10.3) if crash is severe
- Ensure adequate caloric intake—metabolic recovery requires fuel
- Optimize sleep (this is when repair occurs)

Step 3 — Monitoring and documentation:

- Track daily function using Bell Disability Scale or equivalent
- Document: what was tried, at what dose, onset of symptoms, symptom pattern, recovery timeline
- Note whether crash pattern resembles PEM (delayed 24–72 hours) or immediate adverse reaction (within hours)—these have different implications for future treatment attempts

Step 4 — Recovery assessment:

- Return to pre-crash baseline must be confirmed before any new treatment trial
- Minimum recovery period: 4 weeks at stable baseline function
- If recovery is incomplete after 3 months, the treatment may have caused lasting damage—this changes the risk calculus for future trials (apply stricter stop criteria)

25.1.9 Tracking Progress

Systematic treatment response tracking is essential for distinguishing signal from noise in ME/CFS:

Minimum Data to Collect.

- Daily symptom severity (0–10 scale) for: fatigue, cognitive function, pain, sleep quality, orthostatic symptoms
- Activity level (hours upright, steps if wearable available)
- Heart rate data (resting HR, orthostatic HR change)
- New treatment additions/changes with dates and doses
- Crashes: triggers, severity, recovery duration

Assessment Intervals.

- **Daily:** Symptom diary during first 2 weeks of any new treatment
- **Weekly:** Summary assessment during titration periods
- **Monthly:** Review of overall trajectory and treatment utility
- **Quarterly:** Comprehensive reassessment—is the current regimen still justified?

When to Consider a Treatment “Failed.” A treatment should be considered failed after:

- 8–12 weeks at target dose with no detectable benefit
- Any severe adverse reaction requiring discontinuation
- Cost burden outweighing uncertain or minimal benefit
- Functional decline that correlates temporally with treatment initiation

⚠ Limitation 1: Treatment Planning Framework: Rational but Unvalidated

The treatment energy classification (Categories A/B/C), micro-dosing imperative, crash-risk ranking, and sequencing principles presented above are derived from clinical reasoning and patient community experience rather than from clinical trials. Key epistemic boundaries:

- The Category A/B/C energy classification has face validity but has not been validated by measuring actual ATP expenditure or metabolic cost of drug processing in ME/CFS patients.
- The micro-dosing protocol (start at 1/4 to 1/10 standard dose) is based on clinical observation and pattern recognition from the “paradoxical reactor” phenotype; the prevalence of this phenotype, optimal starting fractions, and titration schedules have not been determined by dose-finding studies.
- The claim that “50–70% symptom reduction within 2 weeks” (see Chapter 17, Table 17.1) rests on extrapolation from individual intervention efficacy; the comprehensive multi-target protocol has not been tested as a combined intervention in any study.
- Phenotype-stratified treatment selection (immune-predominant, metabolic-predominant, etc.) is a rational framework but subtype assignment criteria have

not been validated against treatment outcomes.

25.2 Treating Comorbidities

25.2.1 POTS Management

25.2.2 Mast Cell Activation Syndrome

25.2.3 Ehlers-Danlos Syndrome

25.2.4 Other Common Comorbidities

Sleep Apnea Misdiagnosis

Observation 102 (Sleep Apnea Presenting as ME/CFS). Obstructive sleep apnea (OSA) can present with fatigue, cognitive dysfunction, and unrefreshing sleep that closely mimics ME/CFS, leading to years of misdiagnosis. Patient reports from online communities describe complete or near-complete symptom resolution after polysomnography-confirmed sleep apnea treatment with CPAP. One patient described being "disregarded and gaslit by doctors and family" for years before receiving a CPAP device through peer support, which provided significant symptom relief.

Diagnostic overlap with ME/CFS:

- Profound fatigue despite adequate sleep duration
- Unrefreshing sleep (waking exhausted despite 8–10+ hours)
- Cognitive dysfunction (brain fog, memory problems, concentration difficulties)
- Daytime sleepiness and need for naps
- Morning headaches
- Mood disturbances (depression, irritability)

Distinguishing features suggesting OSA:

- Witnessed apneas (breathing stops observed by bed partner)
- Loud snoring, gasping, or choking during sleep
- Severe morning headaches (hypercapnia from nocturnal hypoventilation)
- Obesity (BMI >30), though OSA can occur in normal-weight individuals
- Large neck circumference (>17 inches men, >16 inches women)
- Retrognathia (recessed jaw), large tonsils, or narrow airway
- Improvement in symptoms after CPAP trial

Prevalence and clinical importance:

- OSA affects 10–30% of general adult population [85, 86, 87], higher in men and with obesity
- Many ME/CFS patients develop OSA secondarily due to weight gain from inactivity
- OSA and ME/CFS can coexist; treating OSA improves but may not cure ME/CFS
- Untreated OSA causes cardiovascular disease, hypertension, stroke, diabetes

Diagnostic approach:

- **Polysomnography (sleep study):** Gold standard; measures apnea-hypopnea index (AHI)
- **AHI interpretation:** 5–15 events/hour (mild), 15–30 (moderate), >30 (severe)
- **Home sleep apnea testing:** Alternative to in-lab study; more convenient, less expensive
- **Epworth Sleepiness Scale:** Screening questionnaire (score >10 suggests OSA)
- **STOP-BANG questionnaire:** Clinical prediction tool incorporating snoring, tiredness, observed apneas, pressure (hypertension), BMI, age, neck circumference, gender

Treatment response:

- Primary OSA: CPAP produces dramatic improvement within days to weeks
- OSA + ME/CFS: CPAP improves sleep quality and reduces fatigue but ME/CFS symptoms persist partially
- Compliance critical: CPAP must be used >4 hours/night, most nights to see benefit
- Alternatives to CPAP: Oral appliances (mandibular advancement devices), positional therapy, weight loss, surgery (uvulopalatopharyngoplasty, maxillomandibular advancement)

Clinical recommendation: Polysomnography should be standard in ME/CFS diagnostic workup, particularly for patients with witnessed apneas, loud snoring, morning headaches, obesity, or lack of post-exertional malaise (PEM). The absence of PEM is a red flag that symptoms may be due to primary OSA rather than ME/CFS. Treating comorbid OSA in true ME/CFS patients significantly improves quality of life even if core ME/CFS symptoms remain.

Lyme Disease (European Species)

Observation 103 (Tick-Borne Illness Mimicking ME/CFS). European Lyme disease (*Borrelia* species) can present as chronic fatigue with post-exertional malaise that is indistinguishable from ME/CFS. One patient report documented 10 years of ME/CFS diagnosis before Lyme serology (European testing panel) revealed active infection. Long-cycle antibiotic treatment was described as "significantly helpful," producing improvement not seen with prior ME/CFS interventions.

Clinical overlap with ME/CFS:

- Profound fatigue and malaise
- Post-exertional symptom exacerbation
- Cognitive dysfunction (brain fog, memory problems)

- Sleep disturbances and unrefreshing sleep
- Joint and muscle pain (migratory arthralgias)
- Neurological symptoms (paresthesias, headaches)
- Gradual onset following tick bite (often unrecognized)

Distinguishing features suggesting Lyme:

- **Geographic exposure:** History of travel to or residence in Lyme-endemic regions (Northeast US, Upper Midwest, Northern California; Central and Northern Europe)
- **Tick bite history:** Even if erythema migrans (bull's-eye rash) not recalled (occurs in 70–80% of early Lyme disease cases)
- **Neurological involvement:** Bell's palsy, radiculopathy, meningitis symptoms
- **Cardiac involvement:** Heart block, myocarditis (rare but pathognomonic)
- **Arthritic manifestations:** Large joint swelling (especially knee), often episodic
- **Response to antibiotics:** Improvement with doxycycline or amoxicillin trial

Diagnostic challenges:

- **Serology limitations:** Two-tier testing (ELISA followed by Western blot) has imperfect sensitivity, especially in early disease
- **European vs. US *Borrelia* species:** *B. burgdorferi* (US), *B. afzelii*, *B. garinii* (Europe) require different serology panels
- **Cross-reactivity:** False positives with other spirochetal infections (syphilis), autoimmune diseases (lupus, rheumatoid arthritis)
- **Seronegative Lyme:** Small percentage of true cases remain antibody-negative
- **Regional testing differences:** European labs may use different antigens; patients with European exposure should request European Lyme panels

Testing protocols:

- **Standard two-tier testing:** ELISA screening followed by IgM and IgG Western blot confirmation
- **CDC criteria:** Specific band requirements (IgM: 2/3 bands; IgG: 5/10 bands)
- **European serology:** Include *B. afzelii* and *B. garinii* antigens if European exposure
- **Co-infection testing:** Screen for *Babesia*, *Anaplasma*, *Ehrlichia*, *Bartonella* in endemic areas
- **PCR testing:** Low sensitivity; may help in synovial fluid if arthritic presentation
- **C6 peptide ELISA:** Alternative screening test with potentially better sensitivity

Treatment approaches:

- **Early Lyme (localized):** Doxycycline 100 mg twice daily for 10–21 days, or amoxicillin 500 mg three times daily
- **Disseminated Lyme:** Extended courses (28 days or longer), particularly for neurological involvement
- **Lyme carditis:** IV ceftriaxone; cardiac monitoring required

- **Lyme arthritis:** Oral antibiotics 28 days; some require IV therapy or repeated courses
- **Post-treatment Lyme disease syndrome (PTLDS):** Persistent symptoms after adequate treatment; controversial whether ongoing infection or inflammatory sequelae

Chronic Lyme controversy:

- **IDSA guidelines:** Recommend against prolonged antibiotic therapy for PTLDS; evidence shows no benefit and potential harm (C. difficile, antibiotic resistance)
- **ILADS perspective:** International Lyme and Associated Diseases Society advocates longer treatment courses in some cases
- **Patient reports:** Some describe benefit from extended antibiotics; others experience no improvement or adverse effects
- **Research gap:** Mechanism of persistent symptoms unclear; may represent immune dysfunction triggered by initial infection rather than ongoing active infection

Differential diagnosis strategy:

- ME/CFS diagnosis should follow exclusion of Lyme disease in endemic areas
- Consider empirical doxycycline trial (21–28 days) if strong clinical suspicion despite negative serology
- If dramatic improvement with antibiotics, reassess diagnosis (may be Lyme, not ME/CFS)
- If partial improvement, may represent Lyme-triggered ME/CFS (infection as initiating event)
- Screen for tick-borne co-infections (Babesia causes fatigue, air hunger, night sweats)

Clinical recommendation: All ME/CFS patients with tick exposure history or residence in Lyme-endemic regions should undergo Lyme serology before diagnosis. Patients with European exposure require European-specific testing panels. A subset of "ME/CFS" cases represent missed Lyme disease diagnoses; antibiotic treatment can be life-changing for these individuals.

Ehlers-Danlos Syndrome and Mast Cell Activation

Observation 104 (EDS/MCAS Underdiagnosis in ME/CFS). Hypermobile Ehlers-Danlos syndrome (hEDS) and mast cell activation syndrome (MCAS) are frequently misdiagnosed as ME/CFS or occur as comorbid conditions. Patient advocacy groups and specialist clinicians suggest hEDS prevalence may be "100-fold higher than recognized" due to limited physician awareness, particularly among general practitioners unfamiliar with connective tissue disorders.

Hypermobile Ehlers-Danlos Syndrome (hEDS):

hEDS is a heritable connective tissue disorder characterized by joint hypermobility, skin hyperextensibility, and tissue fragility.

Clinical features overlapping with ME/CFS.

- **Profound fatigue:** Chronic exhaustion from musculoskeletal effort to stabilize hypermobile joints
- **Exercise intolerance:** Joint instability and subluxations worsen with activity
- **Orthostatic intolerance:** POTS occurs in 70–80% of hEDS patients (autonomic dysfunction from connective tissue laxity affecting blood vessels)
- **Cognitive dysfunction:** Brain fog, often secondary to pain, poor sleep, or POTS
- **Chronic pain:** Joint pain, myalgia from compensatory muscle tension
- **Sleep disturbances:** Pain-related sleep disruption

Distinguishing features of hEDS.

- **Joint hypermobility:** Hyperextension of elbows, knees, fingers, thumbs
- **Joint instability:** Frequent subluxations (partial dislocations), chronic sprains
- **Skin hyperextensibility:** Stretchy, velvety skin (though less than classical EDS)
- **Easy bruising:** Fragile capillaries cause extensive bruising from minor trauma
- **Slow wound healing:** Tissue fragility impairs healing
- **Hernias:** Inguinal, umbilical hernias more common
- **Pelvic organ prolapse:** Particularly in women
- **Dental issues:** Crowded teeth, high palate, temporomandibular joint dysfunction
- **Scoliosis or kyphosis:** Spinal curvature abnormalities
- **Marfanoid habitus:** Tall, thin, long limbs (some patients)

Beighton Score for Joint Hypermobility. The Beighton score (0–9 points) assesses generalized joint hypermobility:

1. **Fifth finger passive dorsiflexion $>90^\circ$** (1 point per side)
2. **Thumb passive apposition to forearm** (1 point per side)
3. **Elbow hyperextension $>10^\circ$** (1 point per side)
4. **Knee hyperextension $>10^\circ$** (1 point per side)
5. **Forward trunk flexion with palms flat on floor, knees straight** (1 point)

Interpretation:

- Beighton ≥ 5 (out of 9) suggests generalized joint hypermobility (adults)
- Beighton ≥ 6 for children and adolescents
- Historical hypermobility counts if current score reduced by age/injury

2017 Diagnostic Criteria for hEDS. hEDS diagnosis requires:

1. **Criterion 1 (Generalized joint hypermobility):** Beighton score ≥ 5 (or ≥ 4 if age >50)
2. **Criterion 2 (Two or more features from A, B, C):**
 - Feature A: Systemic manifestations (5+ items from list including skin, hernias, prolapse, etc.)
 - Feature B: Positive family history
 - Feature C: Musculoskeletal complications (chronic pain, instability, subluxations)
3. **Criterion 3 (Exclusion of other EDS types):** No other genetic EDS subtype identified

Management differences from ME/CFS.

- **Physical therapy:** Joint stabilization exercises, proprioceptive training (differs from pacing in ME/CFS)
- **Bracing and supports:** Wrist splints, knee braces, abdominal binders for POTS
- **Surgical caution:** Higher complication rates; avoid elective procedures
- **Pain management:** Focus on joint protection rather than systemic inflammation
- **POTS treatment:** Salt, fluids, compression garments (same as ME/CFS POTS)

Mast Cell Activation Syndrome (MCAS):

MCAS involves aberrant mast cell activation and mediator release causing multi-system symptoms.

Clinical features overlapping with ME/CFS.

- **Fatigue:** Chronic exhaustion from inflammatory mediator release
- **Brain fog:** Histamine and inflammatory cytokines affect cognition
- **Orthostatic intolerance:** Histamine causes vasodilation and POTS-like symptoms
- **Exercise intolerance:** Exertion triggers mast cell degranulation
- **Food sensitivities:** Multiple food intolerances develop over time

Distinguishing features of MCAS.

- **Flushing:** Sudden skin redness, warmth (face, chest, neck)
- **Urticaria (hives):** Spontaneous or triggered by pressure, temperature changes
- **Angioedema:** Swelling of face, lips, tongue, throat
- **Anaphylaxis-like episodes:** Severe reactions requiring epinephrine
- **GI symptoms:** Diarrhea, nausea, cramping, reflux (histamine-mediated)
- **Pruritus:** Severe itching without visible rash
- **Respiratory symptoms:** Wheezing, throat tightness, dyspnea
- **Neuropsychiatric:** Anxiety, panic attacks, brain fog during flares

Diagnostic approach for MCAS. Diagnosis requires all three criteria:

1. **Clinical symptoms:** Multi-system symptoms consistent with mast cell mediator release
2. **Laboratory evidence:** Elevated mediators during symptomatic episodes
 - Serum tryptase (collect within 1–4 hours of acute episode)
 - 24-hour urine histamine metabolites (N-methylhistamine)
 - Plasma or urine prostaglandin D2 or metabolites
 - Plasma heparin or chromogranin A
3. **Response to mast cell stabilizers/mediator antagonists:** Clinical improvement with treatment

Treatment for MCAS.

- **H1 antihistamines:** Cetirizine, loratadine, fexofenadine (non-sedating); may require higher doses
- **H2 antihistamines:** Famotidine, ranitidine (blocks histamine GI effects)
- **Mast cell stabilizers:** Cromolyn sodium (oral, 200–400 mg four times daily); ketotifen
- **Leukotriene inhibitors:** Montelukast (blocks leukotriene-mediated inflammation)
- **Low-histamine diet:** Avoid aged cheeses, fermented foods, alcohol, processed meats
- **Vitamin C:** High-dose (1000–3000 mg/day) stabilizes mast cells
- **Quercetin:** Flavonoid with mast cell stabilizing properties (500–1000 mg twice daily)

hEDS-MCAS-POTS Triad:

The overlap of hEDS, MCAS, and POTS is increasingly recognized:

- 70–80% of hEDS patients have POTS
- High prevalence of MCAS in hEDS population
- Connective tissue laxity may predispose to mast cell dysfunction
- Shared genetic factors proposed but not yet identified
- Treatment requires addressing all three conditions simultaneously

ADHD Connection (Speculative):

Patient communities report high comorbidity between ADHD, hEDS, and ME/CFS:

- Proposed shared genetic factors (collagen, connective tissue genes)
- Executive dysfunction in hEDS may mimic or coexist with ADHD
- Chronic pain and fatigue impair attention and concentration
- Stimulant medications may worsen POTS (increase heart rate)
- Research needed to clarify relationship

Clinical Recommendations:

For ME/CFS patients, screen for hEDS/MCAS if:

- Joint hypermobility (perform Beighton score)
- Easy bruising, fragile skin, slow wound healing
- Frequent joint subluxations or sprains
- Flushing, hives, or anaphylaxis-like episodes
- Multiple food and chemical sensitivities
- Strong family history of hypermobility or allergic conditions

Prevalence Estimates:

- **hEDS prevalence:** Unknown due to underdiagnosis; estimates range from 1:500 to 1:5000 depending on diagnostic stringency
- **"100-fold underdiagnosis" claim:** Based on specialist clinical experience; formal epidemiological data lacking
- **MCAS prevalence:** Estimated 17% of general population may have some form of mast cell disorder; true MCAS prevalence unclear
- **Overlap with ME/CFS:** Unknown; likely substantial given symptom overlap and frequent misdiagnosis

Recognizing hEDS and MCAS in ME/CFS populations is critical because treatment approaches differ substantially, and proper diagnosis can dramatically improve quality of life through targeted interventions (physical therapy for hEDS, mast cell stabilizers for MCAS).

25.3 Personalized Medicine Approaches

25.3.1 Biomarker-Guided Treatment

Speculation 71 (Emerging Patient-Reported Interventions). Patient communities have reported several interventions not yet validated in randomized controlled trials but with plausible mechanistic rationale. These include: (1) Nicotine at low doses (2–4mg/day) for post-viral brain fog, with multiple independent reports of rapid improvement, possibly via nicotinic acetylcholine receptor modulation or anti-inflammatory effects; (2) Methylene blue at "minuscule doses" for smell restoration and brain fog reduction within one week, supported by published research on mitochondrial function improvement; (3) Ketogenic diet producing dramatic symptom resolution in some cases, with one report describing transition from "26 pills per day" to medication-free status. These interventions carry risks (nicotine addiction potential, individual dietary tolerance) and require medical supervision. They represent hypothesis-generating observations requiring formal clinical validation.

△ **Warning 4: Rituximab B-Cell Depletion Failed**

Despite promising early case series showing 67% improvement rates, the definitive Phase III RituxME trial (n=152) demonstrated that rituximab B-cell depletion is not associated with clinical improvement in ME/CFS [376]. The placebo response rate (35%)

exceeded the rituximab response rate (26%). Six-year follow-up confirmed lack of long-term benefit [614]. This represents an important negative result preventing patients from pursuing ineffective immunotherapy. The initial positive case series likely reflected placebo effects, spontaneous remission, or subset-specific responses not replicable in the broader ME/CFS population.

25.3.2 Pharmacogenomics

Pharmacogenomic testing can identify patients at risk of adverse reactions *before* a treatment trial occurs—converting a gamble into an informed decision.

Key Drug Metabolism Variants

CYP2D6 (debrisoquine hydroxylase): Metabolizes approximately 25% of all prescribed medications, including many used in ME/CFS: aripiprazole, beta-blockers (metoprolol, propranolol), SSRIs/SNRIs (fluoxetine, paroxetine, venlafaxine), and tramadol. **Poor metabolizers** (5–10% of Caucasians) accumulate active drug to toxic levels; **ultra-rapid metabolizers** (1–2%) may clear medication before therapeutic effect.

CYP2C19: Metabolizes PPIs, citalopram/escitalopram, and diazepam. Poor metabolizers experience prolonged drug exposure. Relevant for ME/CFS patients on H2 receptor antagonists or antidepressants.

CYP3A4: The most abundant hepatic CYP enzyme, metabolizing approximately 50% of drugs. Cimetidine inhibits CYP3A4, which is therapeutically useful (enhancing acyclovir levels) but also creates interaction risk with co-administered medications metabolized by this pathway.

COMT (catechol-O-methyltransferase): Metabolizes catecholamines (dopamine, norepinephrine, epinephrine). The Val158Met polymorphism creates low-activity (Met/Met) and high-activity (Val/Val) variants. Low-COMT individuals have higher baseline catecholamine levels and may respond differently to dopaminergic agents (aripiprazole) and stress.

MTHFR (methylenetetrahydrofolate reductase): The C677T and A1298C variants impair folate metabolism, affecting methylation capacity. ME/CFS patients with MTHFR variants may require methylfolate rather than folic acid, and may have altered responses to B-vitamin supplementation (Section 22.4).

Clinical Implications for ME/CFS

Speculation 72 (Pharmacogenomics as Harm Reduction in ME/CFS). Pharmacogenomic testing before initiating Category C (energy-demanding) medications could substantially reduce treatment-induced crashes in ME/CFS patients.

The double jeopardy of poor metabolizers: A CYP2D6 poor metabolizer taking a Category C medication faces two simultaneous problems: (1) the drug accumulates to supratherapeutic levels (increased toxicity risk), and (2) the prolonged hepatic processing consumes more ATP

over a longer period (increased energy cost). This combination may explain some paradoxical reactions (Section 21.10.1).

Practical application:

- Order pharmacogenomic panel before any Category C medication trial in severe patients
- CYP2D6 poor metabolizer + aripiprazole = contraindicated (Category C drug with impaired clearance)
- CYP3A4 status critical when cimetidine is part of the regimen (CYP3A4 inhibitor alters metabolism of co-administered drugs)
- MTHFR status guides B-vitamin formulation choice (methylfolate vs. folic acid)

Certainty: 0.25 (pharmacogenomic principles are well-established in general medicine; application to ME/CFS treatment optimization is logical but unvalidated in clinical trials; the specific claim that pharmacogenomic-guided prescribing reduces crashes in ME/CFS is untested)

Available tests: GeneSight, Genomind, OneOme RightMed, Tempus xG, or individual CYP450 testing through most reference laboratories. Cost: \$200–500, often partially covered by insurance.

25.3.3 Subtype-Specific Treatment Approaches

The phenotyping framework (Section 5.6.10) identifies dominant subtypes that predict which treatments are most likely to benefit and least likely to harm. This section translates phenotype into treatment pathway.

Immune-Predominant Subtype

Identification. Frequent infections or reactivations, lymphadenopathy, elevated inflammatory markers (CRP, cytokines), low NK cell function, B-cell abnormalities, positive autoantibody panels, post-infectious onset.

Treatment Priority.

1. **First-line:** Cimetidine (Category B–C; immunomodulatory with manageable energy cost); LDN (Category B; glial modulation)
2. **Second-line:** Antivirals if viral markers elevated—valacyclovir (Category B–C) before valganciclovir (Category C)
3. **Third-line:** IVIG (Category C; only if significant immunoglobulin deficiency documented)

Energy Profile. First-line options are Category B–C with moderate energy cost. Antivirals and IVIG impose significant metabolic demands. **Crash risk: HIGH**—immune modulation triggers inflammatory cascades that are inherently energy-costly. Ensure Category A foundation is solid before initiating.

Key Safety Concern. Immune interventions can cause “Herxheimer-like” reactions (temporary worsening from immune activation). In severe patients, this temporary worsening may become permanent deterioration. Start at micro-doses (Section 25.1.4).

Neurological-Predominant Subtype

Identification. Severe brain fog disproportionate to fatigue, cognitive testing abnormalities, headaches, small fiber neuropathy symptoms (burning/tingling), documented SFN on skin biopsy, MRI findings suggestive of neuroinflammation.

Treatment Priority.

1. **First-line:** LDN (Category B; TLR4-mediated microglial modulation); PEA (Category B; glial modulation + mast cell stabilization)
2. **Second-line:** Ginkgo biloba (Category B; cerebral blood flow enhancement); low-dose aripiprazole (Category C; dopaminergic—use only if pharmacogenomic data supports)
3. **Neuroprotective support:** Alpha-lipoic acid (Category A–B) for SFN; ALCAR (Category A) for cognitive support

Energy Profile. Favorable. First-line options are predominantly Category B with minimal energy cost. Category A neuroprotective supplements can be started simultaneously with neurological treatments. **Crash risk: MEDIUM**—neuroactive medications carry paradoxical reaction risk, but energy cost is generally low.

Metabolic-Predominant Subtype

Identification. Profound fatigue as dominant symptom, exercise intolerance far exceeding other symptoms, abnormal two-day CPET results, metabolomic abnormalities (organic acid panel, acylcarnitine profile), documented mitochondrial dysfunction markers.

Treatment Priority.

1. **First-line:** Full mitochondrial support stack—CoQ10 (ubiquinol), D-ribose, NAD⁺ precursors (NR/NMN), magnesium. *All Category A.*
2. **Second-line:** L-carnitine/ALCAR (Category A; if acylcarnitine profile abnormal); creatine (Category A; ATP buffering)
3. **Third-line:** Alpha-lipoic acid (Category A–B; pyruvate dehydrogenase cofactor)

Energy Profile. This is the safest subtype to treat. All first- and second-line treatments are Category A—they supply what the system lacks. The treatment itself provides energy rather than consuming it. **Crash risk: LOW.** Even multiple Category A supplements can be introduced relatively safely (still one at a time, but with shorter observation windows of 3–7 days rather than 1–2 weeks).

Observation 105 (The Metabolic Subtype Advantage). Patients with metabolic-predominant ME/CFS have the best safety profile for treatment initiation because the most appropriate interventions (mitochondrial support) are also the least energy-demanding. This stands in contrast to immune-predominant patients, where the most appropriate interventions (immunomodulators) are among the most energy-costly. Treatment planning should exploit this asymmetry: stabilize energy metabolism first (Category A), then cautiously address immune or neurological targets.

Autonomic-Predominant Subtype

Identification. Orthostatic intolerance as dominant symptom, positive tilt table test (POTS, orthostatic hypotension), severe exercise intolerance with heart rate abnormalities, gastroparesis or GI dysmotility, temperature dysregulation.

Treatment Priority.

1. **First-line:** Electrolytes and hydration (Category A–B); compression garments (no energy cost)
2. **Second-line:** Pyridostigmine (Category B; low energy cost); fludrocortisone (Category B–C; mineralocorticoid)
3. **Third-line:** Midodrine (Category B–C; vasopressor); beta-blockers if hyperadrenergic POTS

Energy Profile. Moderate. First-line options are energy-neutral to energy-providing. Second- and third-line medications require cardiovascular adaptation which has modest energy cost. **Crash risk: MEDIUM**—autonomic medications require dose titration with hemodynamic monitoring, and blood pressure fluctuations can trigger crashes.

Treatment Selection Decision Tree

Directive 4 (Phenotype-Guided Treatment Prioritization). A systematic approach to treatment selection based on phenotype, energy category, and severity:

Step 1 — Identify dominant subtype via tiered phenotyping protocol (Table 5.5). Note: most patients present with overlapping subtypes. Identify the *dominant* subtype for initial treatment selection; secondary subtypes guide subsequent treatment phases after stabilization on first-line therapy.

Step 2 — Check energy category of proposed treatments against the subtype-specific pathway above.

Step 3 — Apply severity gate:

- **Mild patients:** May proceed to Category B–C treatments after brief Category A stabilization (2–4 weeks)
- **Moderate patients:** Require 4–8 weeks of Category A stabilization before Category B–C trials
- **Severe patients:** Category A treatments only until energy envelope expands measurably; Category B with caution; Category C only with strong biomarker evidence (Tier 2–3 phenotyping)
- **Very severe patients:** Category A only. No Category C trials unless life-threatening comorbidity requires it. Any new treatment at absolute micro-doses

Step 4 — Monitor and adjust: Track response per Section 25.1.9. Apply stop criteria from Section 25.1.7 if crash occurs.

Certainty: 0.25 (integrates well-established pharmacological principles with ME/CFS-specific clinical observations; the specific severity gating thresholds are clinical judgment rather than evidence-based; no RCTs comparing phenotype-guided vs. empirical treatment selection)

25.4 Combination Therapies

25.4.1 Documented Synergies

Several treatment combinations show evidence of synergistic benefit:

- **H1 + H2 dual antihistamine blockade:** Concurrent cetirizine/loratadine (H1) with famotidine/cimetidine (H2) for MCAS management—dual blockade more effective than either alone
- **Cimetidine + valacyclovir:** Cimetidine inhibits renal clearance of acyclovir (the active metabolite of valacyclovir), significantly increasing systemic antiviral concentrations [557]. This pharmacokinetic synergy enables lower antiviral doses with equivalent efficacy, reducing energy cost
- **Mitochondrial support stack:** CoQ10 + NADH + D-ribose target complementary points in ATP production: electron transport (CoQ10), electron donation (NADH), and ATP backbone synthesis (D-ribose). The Castro-Marrero protocol (NADH 20 mg + CoQ10 200 mg) has the strongest RCT evidence [7]
- **LDN + PEA:** Complementary anti-neuroinflammatory mechanisms—LDN via TLR4/microglia and PEA via PPAR- α /mast cells—potentially providing broader neuroinflammatory coverage than either alone

25.4.2 Interaction Risks

△ Warning 5: Dangerous Treatment Combinations in ME/CFS

- **Serotonin syndrome risk:** 5-HTP + SSRIs/SNRIs + trazodone—never combine serotonergic agents without physician supervision. Serotonin syndrome is a potentially fatal energy crisis
- **CYP450 stacking:** Cimetidine (CYP inhibitor) + multiple CYP-metabolized drugs (aripiprazole, SSRIs, beta-blockers)—inhibited clearance leads to drug accumulation. Review all drug interactions when cimetidine is part of the regimen
- **Bleeding risk:** Omega-3 + Ginkgo + CoQ10 + anticoagulants—cumulative antiplatelet effects. Monitor INR if combining multiple agents with anticoagulant properties
- **Hypoglycemia:** D-ribose + alpha-lipoic acid + fasting—both lower blood glucose. Take with food; monitor blood glucose in diabetic patients

25.4.3 Sequencing Considerations

The energy category framework provides clear sequencing guidance:

1. **Energy providers before energy consumers:** Establish Category A foundation before introducing Category C treatments. The energy reserves built by mitochondrial support may improve tolerance of energy-demanding medications
2. **Mast cell stabilization before immune modulation:** If MCAS is present, stabilize mast cells (antihistamines, cromolyn) before introducing immunomodulators that may trigger degranulation
3. **Sleep optimization before cognitive enhancement:** Restorative sleep is the body's primary energy recovery mechanism. Cognitive enhancers without adequate sleep may push activity beyond the energy envelope
4. **One new agent at a time:** Regardless of category, introduce only one new treatment per observation window (minimum 1–2 weeks for Category B–C; 3–7 days acceptable for Category A)

25.5 Cross-Domain Medical Parallels: Learning from Other Fields

ME/CFS shares phenomenological and mechanistic features with several other medical conditions and extreme physiological states. Recognizing these parallels allows us to adapt proven interventions from other fields, potentially accelerating effective treatment development.

25.5.1 Rationale for Cross-Domain Knowledge Transfer

ME/CFS research faces significant challenges: limited funding, lack of validated biomarkers, heterogeneous presentation, and absence of FDA-approved treatments. While waiting for

ME/CFS-specific therapies, examining how other medical fields manage similar physiological challenges can reveal immediately applicable interventions.

When Cross-Domain Transfer Is Valid

Cross-domain knowledge transfer is most valuable when:

1. **Shared underlying mechanisms:** Two conditions involve the same pathophysiological processes (e.g., mitochondrial dysfunction, autonomic impairment)
2. **Similar phenomenology:** Patients experience comparable symptoms despite different etiologies
3. **Proven safety profile:** Interventions are well-established with known risks
4. **Accessible implementation:** Treatments can be realistically applied outside specialized centers
5. **Reasonable biological plausibility:** Mechanistic rationale supports potential benefit

Success Story: Sports Medicine and ME/CFS

The sports medicine parallel (Section 6) demonstrates this approach's value. Recognizing that ME/CFS muscle pathophysiology resembles athletes' post-exercise metabolic stress led to adoption of:

- Oral rehydration solutions (ORS) for blood volume and lactate clearance
- Magnesium supplementation for ATP synthesis and cramp reduction
- Acetyl-L-carnitine for fat oxidation support
- D-ribose as direct ATP precursor

These interventions, borrowed from sports recovery protocols, have shown clinical benefit for managing the chronic metabolic stress state in ME/CFS (Appendix ??).

This section systematically examines other medical fields with similar potential for knowledge transfer.

25.5.2 High-Altitude Medicine: Chronic Hypoxia Parallels

Mechanistic Overlap

High-altitude medicine addresses tissue hypoxia from reduced atmospheric oxygen. ME/CFS involves functional hypoxia despite normal oxygen availability:

Table 25.2: High-Altitude vs. ME/CFS Hypoxia

Feature	High Altitude	ME/CFS
Primary cause	Reduced atmospheric O ₂	Impaired O ₂ delivery or utilization
Cerebral effects	Hypoxic brain dysfunction	Cerebral hypoperfusion
Exercise intolerance	Reduced VO ₂ max	Reduced VO ₂ max at anaerobic threshold
Cognitive symptoms	Confusion, slowed thinking	Brain fog, cognitive impairment
Fatigue pattern	Profound exhaustion	Debilitating fatigue
Sleep disruption	Periodic breathing, poor quality	Unrefreshing sleep, fragmentation
Compensatory response	re- Erythropoiesis, ventilation	Often inadequate compensation

Shared Pathophysiology. Both conditions involve:

- Reduced oxygen delivery to tissues (different mechanisms)
- Cerebral hypoperfusion and cognitive dysfunction
- Reliance on anaerobic metabolism with lactate accumulation
- Exercise intolerance from impaired oxidative capacity
- Autonomic dysregulation

Transferable Interventions from Altitude Medicine

1. Aggressive Iron Optimization. High-altitude medicine targets ferritin >100 µg/L to maximize oxygen-carrying capacity.

- **Rationale for ME/CFS:** Many patients have “normal” ferritin (20–75 µg/L) that is inadequate for optimal oxygen transport and mitochondrial enzyme function
- **Target:** Ferritin 100–200 µg/L (higher end of normal range)
- **Iron form:** Bisglycinate or ferrous sulfate with vitamin C
- **Monitoring:** Recheck every 3 months; avoid over-supplementation (ferritin >300 may indicate inflammation or overload)
- **Additional benefit:** Iron is cofactor for dopamine synthesis, addressing low catecholamines found in ME/CFS CSF

2. Acetazolamide (Diamox). A carbonic anhydrase inhibitor used for altitude sickness prevention.

- **Mechanism:** Induces metabolic acidosis, stimulating ventilation and improving oxygenation

- **Anecdotal ME/CFS reports:** Some patients report improved energy and cognitive function
- **Dose:** 125–250 mg twice daily (half the altitude sickness dose)
- **Side effects:** Paresthesias (tingling), increased urination, taste changes, potassium loss
- **Contraindications:** Kidney disease, liver disease, sulfa allergy
- **Caution:** Limited ME/CFS-specific evidence; primarily case reports and clinical experience
- **Monitoring:** Electrolytes, kidney function before starting and periodically

3. Breathing Optimization. High-altitude climbers use specific breathing techniques to maximize oxygenation.

- **Pressure breathing:** Exhaling against slight resistance increases alveolar pressure
- **Diaphragmatic breathing:** Maximizes lung expansion and oxygen exchange
- **Paced breathing:** Slow, controlled breaths optimize gas exchange
- **ME/CFS application:** May improve oxygen saturation and reduce sympathetic activation
- **Practical protocol:**
 - 4-second inhale through nose (diaphragmatic)
 - Brief hold (1–2 seconds)
 - 6–8 second exhale through pursed lips (creates back-pressure)
 - Practice 5–10 minutes, 2–3 times daily

4. Gradual Acclimatization Protocols. Altitude medicine emphasizes gradual exposure to stress, mirroring ME/CFS pacing principles.

- **“Climb high, sleep low”:** Brief exposure to higher stress with return to baseline
- **ME/CFS translation:** Brief activity within limits, extensive rest for recovery
- **Principle:** Respect physiological adaptation capacity; pushing too hard causes deterioration
- **This validates pacing:** Altitude medicine proves that gradual, respectful approaches work better than forcing through physiological limits

5. Blood Volume Optimization. Altitude exposure reduces plasma volume; countermeasures include aggressive hydration and electrolyte management.

- **Already implemented in ME/CFS:** Fluid and salt loading for POTS (Section 16.6)
- **Dual benefit:** Blood volume expansion for both orthostatic tolerance and oxygen delivery
- **ORS formula:** See sports medicine section earlier in this chapter for sports medicine-derived protocol

6. Monitoring and Objective Tracking. Altitude medicine uses pulse oximetry, heart rate, and subjective symptoms to guide activity.

- **ME/CFS application:** Pulse oximeters (<\$30), heart rate monitors, HRV tracking
- **Objective limits:** Stay below calculated anaerobic threshold heart rate
- **Oxygen saturation:** Monitor for drops during or after activity (may reveal impaired oxygen extraction)
- **Trend tracking:** Daily measurements reveal patterns and guide pacing decisions

Limitations and Cautions

- **Different underlying causes:** Altitude = low ambient O₂; ME/CFS = impaired delivery/utilization
- **Acetazolamide evidence:** Limited to case reports in ME/CFS; no controlled trials
- **Individual variation:** Responses to altitude interventions vary widely
- **Medical supervision required:** Acetazolamide, aggressive iron supplementation need physician oversight

25.5.3 Critical Care and ICU Recovery Medicine

Post-Intensive Care Syndrome (PICS): The Acquired ME/CFS

Post-intensive care syndrome describes the constellation of symptoms affecting ICU survivors:

- **Physical impairment:** Profound weakness, exercise intolerance, muscle wasting
- **Cognitive dysfunction:** Memory deficits, slowed processing, executive dysfunction (“ICU brain fog”)
- **Psychological symptoms:** Depression, anxiety, PTSD
- **Duration:** Symptoms persist months to years after discharge
- **Prevalence:** Affects 50–75% of ICU survivors

The phenomenological overlap with ME/CFS is striking. PICS may represent acquired ME/CFS triggered by severe physiological stress.

Mechanistic Overlap

Transferable Interventions from ICU Recovery Protocols

1. Aggressive Micronutrient Repletion. Critical illness depletes vitamins and minerals at alarming rates. ICU recovery protocols aggressively replete these.

Table 25.3: PICS vs. ME/CFS Mechanisms

Mechanism	PICS	ME/CFS
Mitochondrial dysfunction	Sepsis-induced damage	Constitutional or acquired
Inflammation	Cytokine storm → persistent low-grade	Post-viral or chronic activation
Muscle wasting	ICU-acquired weakness	Deconditioning + metabolic impairment
Autonomic dysfunction	Dysautonomia post-sepsis	Dysautonomia (POTS, OI)
Cognitive impairment	Hypoxic brain injury, inflammation	Cerebral hypoperfusion, neuroinflammation
Oxidative stress	Massive ROS generation	Chronic oxidative stress
Nutritional depletion	Hypermetabolic state	Malabsorption, increased utilization

ICU-Derived Micronutrient Protocol for ME/CFS

Rationale: It has been hypothesized that ME/CFS may involve chronic low-grade nutritional depletion from:

- Malabsorption (gut dysfunction)
- Increased oxidative stress (higher antioxidant utilization)
- Impaired metabolism (reduced cofactor availability)

High-Priority Targets (ICU critical care experience):

1. **Thiamine (B1)** - 100–300 mg daily
 - Critical for aerobic metabolism (pyruvate dehydrogenase cofactor)
 - Deficiency causes lactic acidosis and neurological symptoms
 - ICU dosing: Often 100–200 mg IV; oral equivalent 100–300 mg
 - Extremely safe; water-soluble with no toxicity concern
2. **Vitamin C** - 1000–2000 mg daily (divided doses)
 - Sepsis protocols use high-dose IV vitamin C (1.5–6 g daily)
 - Antioxidant, immune support, collagen synthesis
 - May reduce oxidative stress in ME/CFS
 - Oral absorption limited; divide into 2–3 doses for sustained levels
3. **Vitamin D** - 4000–5000 IU daily (target 50–70 ng/mL)
 - ICU patients often severely deficient
 - Immune modulation, muscle function, mood
 - ME/CFS patients frequently deficient despite supplementation (fat malabsorption)
 - Requires dietary fat for absorption
4. **Magnesium** - 300–400 mg glycinate daily
 - ICU: Often depleted; replaced IV
 - ATP synthesis, muscle function, nervous system
 - Glycinate form: best absorption, minimal GI effects
 - Already discussed for muscle cramps in sports medicine section above
5. **Zinc** - 15–30 mg daily
 - Immune function, wound healing, antioxidant
 - Often depleted in chronic illness
 - Take with food to reduce nausea
 - Balance with copper (2 mg copper for every 15 mg zinc if supplementing long-term)

2. N-Acetylcysteine (NAC) for Oxidative Stress. NAC is used in ICU for acetaminophen overdose and as adjunct sepsis treatment.

- **Mechanism:** Glutathione precursor; powerful antioxidant; mucolytic
- **ICU dosing:** 600–1200 mg IV for sepsis adjunct therapy
- **ME/CFS application:** 600 mg twice daily oral
- **Rationale:** ME/CFS shows evidence of oxidative stress and glutathione depletion
- **Benefits:** May reduce oxidative damage, support detoxification, thin mucus (if sinus/respiratory issues)
- **Side effects:** GI upset (take with food), sulfur odor
- **Caution:** May worsen asthma in some individuals; start low dose
- **Evidence:** Small ME/CFS studies suggest potential benefit for fatigue and brain fog

3. Structured Reconditioning: ICU Early Mobility Protocols. ICU early mobility programs prevent deconditioning while respecting severe functional limitations.

- **ICU approach:** Gradual progression from bed exercises to sitting to standing to walking
- **Key principle:** Activity matched to current capacity; never pushing through exhaustion
- **ME/CFS translation:** Graded activity within energy envelope (NOT graded exercise therapy/GET)
- **Critical difference from GET:**
 - ICU protocols respect physiological limits
 - Progress is based on objective tolerance, not predetermined schedules
 - Activity is reduced or paused if deterioration occurs
 - **This is pacing, not pushing**
- **Practical application:** Start with 2–5 minutes of gentle movement within heart rate limits; increase only if tolerated without PEM

4. Sleep Architecture Restoration. ICU delirium prevention protocols emphasize sleep hygiene and circadian rhythm maintenance.

- **ICU strategies:**
 - Minimize nighttime interruptions
 - Optimize sleep environment (darkness, quiet, temperature)
 - Daytime light exposure and activity (within limits)
 - Avoid sedatives that fragment sleep architecture
- **ME/CFS application:** Same principles apply
- **Melatonin:** ICU protocols sometimes use melatonin 3–10 mg for circadian rhythm support
- **Light therapy:** Morning bright light (10,000 lux) for circadian entrainment

5. Nutrition Support: Protein and Calories. ICU patients require aggressive nutritional support to prevent muscle wasting.

- **Protein target:** 1.2–2.0 g/kg body weight daily (higher than general population)
- **Rationale for ME/CFS:** Muscle wasting, impaired protein synthesis from metabolic dysfunction
- **Practical target:** 80–120 g protein daily for average adult
- **Sources:** Whey protein powder, eggs, fish, chicken, Greek yogurt
- **Timing:** Distribute throughout day (20–30 g per meal)
- **Calories:** Ensure adequate total intake; underfeeding worsens weakness

Glutamine Supplementation: Controversial but Promising

Glutamine is conditionally essential during critical illness; ICU nutrition protocols often supplement it.

- **Functions:** Gut barrier integrity, immune cell fuel, nitrogen transport
- **ICU use:** 0.3–0.5 g/kg/day (20–40 g daily for average adult)
- **ME/CFS rationale:** Gut dysfunction (leaky gut), immune activation may increase glutamine demand
- **Dose:** 5–15 g daily, divided doses
- **Form:** L-glutamine powder (unflavored, mix in water)
- **Timing:** Away from meals for gut barrier support; with meals for immune support
- **Evidence in ME/CFS:** Minimal; theoretical rationale based on gut dysfunction
- **Cost:** \$20–30/month
- **Safety:** Generally well-tolerated; avoid in liver disease, kidney disease

Key Lessons from PICS Management

1. **Aggressive nutritional support is not optional:** Micronutrients, protein, adequate calories
2. **Oxidative stress management:** Antioxidants (vitamin C, NAC, selenium)
3. **Gradual reconditioning respecting limits:** ICU mobility protocols validate pacing approach
4. **Sleep and circadian rhythm:** Environmental optimization, melatonin, light therapy
5. **Recovery takes time:** PICS recovery measured in months to years, not weeks

The ICU medicine parallel reinforces that severe, prolonged functional impairment requires comprehensive, long-term metabolic and nutritional support—exactly what ME/CFS demands.

25.5.4 Space Medicine: Orthostatic Intolerance and Deconditioning

Microgravity-Induced Deconditioning: The ME/CFS Analog

Astronauts returning from prolonged spaceflight experience a syndrome strikingly similar to ME/CFS:

- **Orthostatic intolerance:** Unable to stand without severe symptoms (some faint within minutes)
- **Exercise intolerance:** Reduced VO₂max, profound weakness
- **Muscle atrophy:** Despite resistance exercise in space
- **Bone loss:** From unloading
- **Cognitive changes:** “Space fog” during and after flight
- **Autonomic dysfunction:** Altered cardiovascular reflexes
- **Immune dysregulation:** Altered immune cell function

The key difference: Astronauts’ symptoms are predictable and (mostly) reversible with structured reconditioning. ME/CFS patients experience similar physiology without the microgravity trigger and often without reliable recovery.

Shared Pathophysiology

Table 25.4: Microgravity vs. ME/CFS Deconditioning

Feature	Post-Spaceflight	ME/CFS
Blood volume	Reduced 10–15%	Reduced (documented in many patients)
Orthostatic tolerance	Severe impairment post-landing	POTS, OI in 70–90%
Muscle strength	Reduced 20–40%	Progressive weakness
Mitochondrial function	Impaired in some studies	Widespread dysfunction
Bone density	Significant loss	Variable (deconditioning)
Cardiovascular fitness	VO ₂ max reduced	VO ₂ max reduced on CPET
Autonomic function	Dysregulated reflexes	ANS dysfunction

Transferable Interventions from Space Medicine

1. Compression Garments: Proven Orthostatic Countermeasure. Astronauts use compression garments immediately post-landing to prevent fainting.

- **Mechanism:** External pressure prevents venous pooling in legs; improves venous return
- **Space medicine use:** Thigh-high or waist-high compression immediately after landing
- **ME/CFS application:** Already standard POTS treatment (Section 16.6)
- **Compression levels:**

- Mild ME/CFS or prevention: 15–20 mmHg
- Moderate symptoms: 20–30 mmHg
- Severe orthostatic intolerance: 30–40 mmHg
- **Type:** Waist-high stockings more effective than knee-high (prevents thigh pooling)
- **Practical note:** Difficult to don with limited energy; may require assistance or donning aids

2. Structured Reconditioning: Lessons from Astronaut Post-Flight Rehab. NASA has refined reconditioning protocols through decades of astronaut recovery data.

Space Medicine Reconditioning Principles for ME/CFS

NASA's Core Principles (adapted for ME/CFS):

1. **Horizontal-first exercise:** Start with recumbent activities (no orthostatic stress)
 - Recumbent bike, rowing machine (lying position)
 - Supine resistance bands
 - Pool exercises (water supports body weight)
2. **Gradual gravitational challenge:** Progress from lying → sitting → standing
 - Week 1–4: Recumbent only
 - Week 5–8: Add seated exercise if tolerated
 - Week 9+: Brief standing exercise if no PEM
3. **Objective monitoring:** Heart rate, blood pressure, subjective symptoms
 - Heart rate limit: $(220 - \text{age}) \times 0.55$ (anaerobic threshold)
 - BP monitoring: Stop if significant drop or symptoms
 - Symptom tracking: Any increase in fatigue, PEM = reduce activity
4. **Volume before intensity:** Build duration first, intensity last
 - Start: 2–5 minutes low-intensity
 - Increase duration by 1 minute per week if tolerated
 - Only increase resistance/speed after duration goal met
5. **Rest is intervention:** Recovery days are not optional
 - 2–3 exercise days per week maximum initially
 - Full rest days between sessions
 - Any PEM = full stop until recovered

Critical ME/CFS Adaptation:

- Astronauts progress predictably; ME/CFS patients may not
- **If worsening occurs, STOP and reassess**
- This is NOT graded exercise therapy (GET)—progression is optional, not mandatory
- Many severe ME/CFS patients cannot progress beyond recumbent positioning
- **Goal is maintenance of current capacity, not necessarily improvement**
- Even minimal recumbent activity may help slow deconditioning, though evidence for this in ME/CFS specifically is limited

3. Blood Volume Restoration. Astronauts rapidly restore blood volume post-landing through aggressive fluid and salt loading.

- **Space medicine protocol:** IV saline infusion or oral fluid/salt loading pre-landing
- **ME/CFS application:** Already implemented (Section 16.6)
- **Immediate pre-activity loading:** Drink 500 mL ORS 30 minutes before standing/activity

- **Sustained maintenance:** 2.5–3 L daily fluids, 6–10 g sodium daily

4. Bone and Muscle Preservation: Resistance Training Within Limits. Space medicine uses resistance exercise to minimize bone/muscle loss during flight.

- **Key finding:** Even in microgravity, resistance exercise preserves some muscle
- **ME/CFS application:** Light resistance training (within energy limits) may slow deconditioning
- **Practical protocol:**
 - Resistance bands (adjustable tension)
 - Bodyweight exercises in recumbent position (leg presses against wall while lying down)
 - Very brief sessions: 5–10 minutes, 2×/week maximum
 - Stay within heart rate limits
 - Stop immediately if PEM symptoms emerge
- **Goal:** Maintenance, not gain
- **Caveat:** Not appropriate for severe patients or during crashes

5. Monitoring Technology: Heart Rate and Activity Tracking. NASA uses continuous physiological monitoring during and after spaceflight.

- **Space medicine:** ECG, BP, accelerometry, subjective logs
- **ME/CFS-accessible equivalents:**
 - Heart rate monitor or fitness tracker (\$50–300)
 - Blood pressure cuff with memory (\$30–60)
 - Activity tracker (steps, movement patterns)
 - Symptom diary (free)
- **Key metrics:**
 - Resting heart rate trends (increasing RHR = overexertion or illness)
 - Heart rate during activity (stay below threshold)
 - Orthostatic heart rate change (POTS screening)
 - Heart rate variability (HRV)—lower HRV indicates stress, poor recovery

Key Lessons from Space Medicine

1. **Orthostatic intolerance is manageable:** Compression, fluid/salt loading, gradual reconditioning work
2. **Horizontal-first approach:** Removing gravitational stress allows exercise when standing is impossible
3. **Objective monitoring prevents overexertion:** Astronauts don't "push through"—neither should ME/CFS patients

4. **Reconditioning is gradual and structured:** Even healthy astronauts require months to recover
5. **Some impairment may persist:** Not all astronauts return to pre-flight baseline

Space medicine validates that severe deconditioning and orthostatic intolerance are real physiological challenges requiring systematic, respectful interventions—not psychological motivation or willpower.

25.5.5 Additional Domain Parallels: Brief Overview

Several other medical fields offer potential insights, though with less developed transferable protocols:

Diving Medicine: Hyperbaric Oxygen and Perfusion

- **Overlap:** Tissue perfusion optimization, oxygen delivery under stress
- **HBOT for ME/CFS:** Emerging treatment; some studies show benefit for fatigue and cognitive function
- **Mechanism:** Increases dissolved oxygen in plasma, may improve mitochondrial function
- **Accessibility:** Requires specialized facilities; expensive (\$100–200/session)
- **Evidence:** Preliminary; larger trials needed
- **Practical:** Consider if accessible and affordable; typical protocol 20–40 sessions

Burn and Trauma Medicine: Hypermetabolic State Management

- **Overlap:** Massive nutritional demands, oxidative stress, immune activation
- **Transferable concepts:**
 - Aggressive protein supplementation (1.5–2 g/kg/day)
 - Glutamine for gut barrier (discussed in ICU section)
 - Antioxidant support (vitamins C, E, selenium, zinc)
 - Anabolic support: Oxandrolone (anabolic steroid) used in burn patients for muscle preservation
- **Oxandrolone for severe ME/CFS wasting:** Theoretical interest; no trials
- **Caution:** Anabolic steroids have significant side effects; only for severe, refractory cases under specialist supervision

Geriatric Frailty Medicine: Multi-System Decline

- **Overlap:** Exercise intolerance, weakness, falls risk, polypharmacy, functional decline
- **Transferable concepts:**
 - Comprehensive geriatric assessment model (systematic evaluation of all systems)
 - Vitamin D optimization (frailty protocols target 40–60 ng/mL)

- Protein supplementation (whey protein, essential amino acids)
- Fall prevention strategies (relevant to orthostatic ME/CFS patients)
- Acceptance of mobility aids without stigma (canes, walkers, wheelchairs)
- Polypharmacy reduction (minimizing medication burden)
- **Key insight:** Geriatric medicine validates that accepting functional limitations and using assistive devices improves quality of life

Chronic Pain Medicine: Central Sensitization

- **Overlap:** Central nervous system dysfunction, neurotransmitter dysregulation, quality of life impairment
- **Transferable interventions:**
 - Low-dose naltrexone (already used in ME/CFS)
 - Gabapentinoids (gabapentin, pregabalin) for neuropathic symptoms
 - Ketamine (low-dose) for central sensitization reset (emerging interest)
 - Acceptance-based approaches (pain psychology principles align with pacing)
 - Vagal nerve stimulation (pain modulation + autonomic regulation)
- **Evidence:** LDN has best ME/CFS evidence; others largely anecdotal

25.5.6 Integration and Practical Application

Building a Cross-Domain Treatment Protocol

The interventions from multiple fields can be integrated into a comprehensive approach:

Prioritization Strategy

Not all interventions are equally accessible or evidence-based. Prioritize by:

1. **Tier 1 - Immediate implementation** (low cost, high safety, reasonable evidence):
 - ORS (sports medicine): \$5/month
 - Magnesium glycinate: \$10/month
 - Vitamin D optimization: \$5/month
 - B-complex: \$10/month
 - Compression stockings: \$30–60 one-time
 - Heart rate monitoring: Use existing device or \$30–100
2. **Tier 2 - Evidence-supported** (moderate cost, proven benefit in related conditions):
 - CoQ10 + Acetyl-L-carnitine (sports/ICU): \$40–60/month
 - Iron optimization if deficient (altitude): \$10–15/month
 - Vitamin C, NAC (ICU): \$15–25/month
 - Thiamine (ICU): \$5/month

Table 25.5: Cross-Domain Intervention Summary

Domain	Key Interventions	Primary Benefits
Sports Medicine	ORS, magnesium, Acetyl-L-carnitine, D-ribose	Lactate clearance, ATP support, cramp reduction
Altitude Medicine	Iron optimization, acetazolamide, breathing techniques	Oxygen delivery, cognitive function, exercise tolerance
ICU Recovery	Micronutrients (B1, C, D, Mg, Zn, Se), NAC, protein	Metabolic support, oxidative stress, muscle preservation
Space Medicine	Compression, horizontal exercise, blood volume expansion	Orthostatic tolerance, reconditioning, monitoring
Burn/Trauma	Glutamine, high protein, antioxidants	Gut barrier, immune support, healing
Geriatrics	Vitamin D, protein, mobility aids, polypharmacy reduction	Frailty prevention, function optimization
Chronic Pain	LDN, gabapentinoids, acceptance strategies	Pain reduction, central sensitization, pacing validation

- Zinc, selenium (ICU): \$10/month

3. Tier 3 - Theoretical or emerging (higher cost, limited ME/CFS evidence, or requiring prescription):

- Acetazolamide (altitude): Prescription required
- D-ribose (sports): \$25–40/month
- Glutamine (burn/trauma): \$20–30/month
- HBOT (diving): \$2000–8000 for course
- Gabapentinoids (chronic pain): Prescription required
- Ketamine (chronic pain): Specialist administration

Monitoring Cross-Domain Interventions

Track responses systematically:

- **Symptom diary:** Daily energy (0–10), cognitive function (0–10), pain (0–10), PEM episodes
- **Objective measures:**
 - Resting heart rate (daily morning)
 - Orthostatic heart rate change (weekly)
 - HRV if available (daily)
 - Activity tolerance (minutes standing/walking without PEM)
- **Laboratory monitoring:**

- Ferritin, iron panel (if supplementing iron: every 3 months)
- Vitamin D (every 3–6 months until optimized)
- Electrolytes, kidney function (if taking acetazolamide or high-dose salt)
- Liver function, CBC (periodic if taking multiple supplements)
- **Response timeline:** Most nutritional interventions require 4–12 weeks for full effect
- **Decision rule:** If no benefit after 3 months, discontinue and try next priority intervention

25.5.7 Cautions and Limitations

When Cross-Domain Transfer Fails

Not all interventions from other fields will work in ME/CFS:

- **Different underlying mechanisms:** ME/CFS pathophysiology may differ fundamentally despite similar phenomenology
- **Paradoxical reactions:** Some ME/CFS patients respond opposite to expected (e.g., stimulants worsening some patients)
- **Heterogeneity:** ME/CFS is likely multiple diseases; interventions may work for some subsets only
- **Lack of ME/CFS-specific trials:** Most evidence is extrapolated, not proven

Safety Considerations

- **Medical supervision required:** Prescription medications (acetazolamide, gabapentinoids), IV therapies (HBOT), high-dose supplementation (iron if ferritin already normal)
- **Drug interactions:** Many ME/CFS patients take multiple medications; check interactions
- **Start low, go slow:** Begin with lowest effective dose; increase gradually
- **One change at a time:** If possible, introduce interventions sequentially (1–2 weeks apart) to identify responders
- **Monitor for worsening:** Some interventions may worsen symptoms; discontinue if deterioration occurs

Realistic Expectations

Cross-domain interventions are ****supplementary support, not cures****:

- **Best-case scenario:** 10–30% functional improvement through cumulative effects
- **Typical scenario:** Modest symptom reduction; improved quality of life within severe limitations
- **Worst-case scenario:** No benefit or worsening
- **All interventions are compensatory:** Stopping effective treatments likely results in symptom return

- **Chronic disease management:** Lifelong implementation required if beneficial

25.5.8 Research Implications: Cross-Domain Studies

The cross-domain parallel approach suggests valuable research directions:

1. **Comparative physiology studies:** Systematically compare ME/CFS to PICS, post-spaceflight syndrome, high-altitude intolerance
2. **Shared biomarkers:** Identify common markers across conditions (lactate, catecholamines, inflammatory profiles)
3. **Intervention trials:** Test altitude medicine (acetazolamide), ICU protocols (high-dose thiamine/vitamin C), space medicine (structured reconditioning)
4. **Mechanism studies:** Understand why similar interventions work across different conditions (mitochondrial? inflammatory? autonomic?)
5. **Subtype identification:** Determine which ME/CFS patients resemble which parallel condition (altitude-like hypoxia vs. ICU-like inflammation vs. space-like deconditioning)

25.5.9 Conclusion: The Value of Looking Beyond ME/CFS

Other medical fields have confronted similar physiological challenges—tissue hypoxia, metabolic stress, orthostatic intolerance, profound weakness—and developed systematic interventions. While ME/CFS awaits specific treatments, adapting proven approaches from altitude medicine, critical care, space medicine, and other domains provides immediately actionable strategies.

The sports medicine parallel discussed in this chapter and documented in detail in Appendix ?? demonstrates this approach's value. Recognizing phenomenological similarities led to effective interventions (ORS, magnesium, Acetyl-L-carnitine) now benefiting ME/CFS patients.

Key principles:

- Shared mechanisms justify intervention transfer
- Prioritize safe, accessible, evidence-based approaches
- Monitor responses objectively
- Accept that not all transfers will succeed
- View interventions as compensatory support, not cures
- Maintain realistic expectations while remaining open to benefit

Until ME/CFS-specific treatments emerge, learning from how other fields manage similar physiological states offers the best available path forward.

25.6 Novel Mechanistic Hypotheses and Research Opportunities

Based on integration of recent molecular findings, patient-reported phenomena, and cross-domain medical parallels, several novel hypotheses and research opportunities emerge.

25.6.1 WASF3 as Therapeutic Target

Speculation 73 (WASF3 Inhibitors from Cancer Pipelines). The Wang 2023 finding that WASF3 knockdown with shRNA restores mitochondrial function in ME/CFS patient cells [2] suggests WASF3 may be a druggable target. WASF3 is already under investigation as an oncology target for metastasis suppression. Repurposing WASF3 inhibitors from cancer drug development pipelines for ME/CFS could provide a reversible intervention targeting upstream mitochondrial dysfunction. Unlike symptomatic treatments, WASF3 inhibition might address the molecular mechanism driving Complex IV dysfunction and ATP depletion.

25.6.2 Acetylcholine-Mitochondrial Axis

~ Hypothesis 1: Cholinergic-Mitochondrial Signaling Link

Patient reports of rapid brain fog relief with nicotine (2–4mg daily), combined with documented mitochondrial dysfunction, suggest a potential cholinergic-mitochondrial signaling axis. Alpha-7 nicotinic acetylcholine receptors are present on mitochondrial membranes and modulate calcium handling, which directly affects ATP production. This raises the hypothesis that cholinergic signaling deficits may impair mitochondrial bioenergetics in ME/CFS. If validated, acetylcholinesterase inhibitors (donepezil, galantamine) used for Alzheimer's disease might provide both cognitive and metabolic benefits in ME/CFS.

? Open Question 1: Mitochondrial Acetylcholine Receptors in ME/CFS

Do ME/CFS patients show altered expression or function of mitochondrial alpha-7 nicotinic acetylcholine receptors? Does acetylcholine signaling regulate mitochondrial biogenesis or Complex IV assembly in human muscle cells?

25.6.3 ATP Recovery Kinetics and Mitophagy

~ Hypothesis 2: Delayed ATP Recovery from Mitophagy Failure

The 24–72 hour delay in VO_2max recovery observed in 2-day CPET [79] matches patient-reported post-exertional malaise timing. This delay aligns with the time course of mitochondrial autophagy (mitophagy) and biogenesis cycles, which operate on circadian and ultradian rhythms. Post-exertion, damaged mitochondria must be cleared via mitophagy and replaced through biogenesis—processes requiring 24–48 hours. If ME/CFS

involves impaired mitophagy or delayed mitochondrial regeneration, ATP recovery would be prolonged, explaining the characteristic delayed symptom onset of PEM.

? Open Question 2: Mitophagy Markers in ME/CFS

Do ME/CFS patients show reduced mitophagy flux markers (PINK1, Parkin, LC3-II) post-exertion? Is mitochondrial biogenesis (PGC-1 α , TFAM expression) delayed compared to healthy controls following standardized exercise?

25.6.4 Viral Trigger-ER Stress-WASF3 Pathway

~ Hypothesis 3: Viral Proteostasis Disruption Activates WASF3

Multiple viral triggers identified in meta-analysis (EBV, HHV-7, enterovirus, coxsackie B) [121] share a common mechanism: disruption of cellular proteostasis leading to endoplasmic reticulum (ER) stress and unfolded protein response (UPR) activation. Viral protein production overwhelms the ER, triggering stress pathways that may activate WASF3 expression. This connects viral onset with downstream mitochondrial dysfunction via ER stress-WASF3-mitochondria axis. If validated, ER stress modulators (tauroursodeoxycholic acid/TUDCA, 4-phenylbutyrate) might prevent WASF3 activation and progression to chronic ME/CFS when administered during acute viral illness.

Speculation 74 (ER Stress Modulators for Viral ME/CFS Prevention). Chemical chaperones that reduce ER stress (TUDCA 500–1000mg/day, 4-phenylbutyrate 500mg/day) are FDA-approved for other conditions and well-tolerated. Early administration during acute EBV, enterovirus, or SARS-CoV-2 infection might prevent ER stress-mediated WASF3 upregulation and subsequent mitochondrial dysfunction. This represents a testable prophylactic intervention for at-risk individuals (family history of ME/CFS, severe viral prodrome).

25.6.5 Pyruvate Supplementation Hypothesis

Speculation 75 (Pyruvate for ATP Regeneration Bypass). If ATP regeneration is delayed 24–72 hours post-exertion due to mitochondrial dysfunction, direct pyruvate supplementation might bypass glycolytic bottlenecks by providing immediate acetyl-CoA substrate for the TCA cycle. Pyruvate enters mitochondria directly without requiring full glycolysis. Prophylactic pyruvate drinks (1–2g) consumed 30–60 minutes before anticipated exertion could theoretically prevent ATP depletion. Oral pyruvate is commercially available, well-tolerated, and used by athletes for performance enhancement. This represents a low-risk, testable intervention for activity preparation.

25.6.6 Methylene Blue as Electron Transport Bypass

~ Hypothesis 4: Methylene Blue Electron Transport Enhancement

Patient reports of methylene blue (1–5mg daily) improving brain fog and smell within one week suggest potential mitochondrial benefits. Methylene blue can accept electrons from NADH (Complex I) and donate them to Complex III, potentially enhancing electron flow when upstream complexes are impaired. Additionally, methylene blue may reduce oxidative stress and improve mitochondrial membrane potential. While WASF3-mediated damage affects Complex IV [2], methylene blue's effects on overall electron transport chain efficiency and mitochondrial redox state might provide indirect benefit. This mechanism is established in methylene blue's use for methemoglobinemia and has shown mitochondrial benefits in neurodegenerative disease models.

? Open Question 3: Complex-Specific Dysfunction Pattern

Is mitochondrial dysfunction in ME/CFS specific to Complex IV, or do other complexes show impairment? Would interventions targeting specific complex deficits (Complex I: CoQ10; Complex IV: copper, cytochrome c) show differential efficacy?

25.6.7 Beta-Blockers for Pacing Enforcement

Speculation 76 (Pharmacological Heart Rate Ceiling). The “<5 crashes per year” rule suggests cumulative irreversible damage from exceeding energy limits. Low-dose beta-blockers (e.g., propranolol 10–20mg as needed) might pharmacologically enforce pacing by preventing heart rate spikes during inadvertent overexertion. Combined with heart rate-based wearable alerts, beta-blockers could provide a safety ceiling preventing accidental crashes in mild-to-moderate patients with variable symptom awareness. This differs from continuous beta-blockade for POTS—it would be prophylactic, taken before high-risk activities (social events, medical appointments).

25.6.8 Immune Checkpoint Modulation

~ Hypothesis 5: T-Cell Exhaustion in Chronic Viral ME/CFS

The failure of B-cell depletion (rituximab) [376] suggests B-cells are not the primary immune dysfunction. Chronic viral infections induce T-cell exhaustion characterized by upregulation of checkpoint receptors (PD-1, TIM-3, LAG-3) and loss of effector function. If ME/CFS involves persistent viral antigen or defective viral clearance, exhausted T-cells may fail to control low-level infection, perpetuating immune activation. Anti-PD-1 or anti-CTLA-4 antibodies used in cancer immunotherapy might reverse T-cell exhaustion and restore antiviral immunity. This is highly speculative and carries significant risks (autoimmune adverse events), but represents a testable hypothesis if T-cell exhaustion markers are confirmed.

△ Warning 6: Checkpoint Inhibitors Carry High Risk

Immune checkpoint inhibitors are powerful immunotherapies with serious potential side effects including autoimmune colitis, pneumonitis, hepatitis, and endocrinopathies. They should only be considered in severe, refractory ME/CFS under research protocols with extensive safety monitoring. This speculation is hypothesis-generating for research, not clinical recommendation.

25.6.9 Central Governor Theory Link**~ Hypothesis 6: Hypersensitive Central Governor as Protective Mechanism**

The “central governor” theory in exercise physiology proposes that the brain actively limits muscle recruitment to prevent tissue damage. ME/CFS may represent a hypersensitive central governor responding to real mitochondrial damage signals. Brain fog and cognitive fatigue might serve as protective mechanisms preventing ATP-depleting cognitive exertion when metabolic reserves are low. This reframes cognitive symptoms not as primary neurological dysfunction, but as adaptive limitation to prevent energetic crisis. Functional MRI studies comparing brain activation patterns during cognitive tasks in ME/CFS versus healthy controls could test this hypothesis.

25.6.10 Lactate Clearance Dysfunction**~ Hypothesis 7: Impaired Lactate Clearance Delays Recovery**

The 2-day CPET demonstrates impaired recovery, not just impaired peak performance [79]. Lactate clearance occurs primarily via hepatic gluconeogenesis and mitochondrial lactate oxidation. If mitochondrial dysfunction impairs lactate-to-pyruvate conversion or liver metabolism is compromised, lactate accumulation would persist post-exertion, prolonging metabolic acidosis and delaying ATP regeneration. Serial blood lactate measurements at 0h, 24h, and 48h post-CPET could test this hypothesis. If confirmed, NAD⁺ precursor supplementation (nicotinamide riboside, nicotinamide mononucleotide) to boost lactate dehydrogenase activity might accelerate recovery.

Speculation 77 (NAD⁺ Precursors for Lactate Clearance). NAD⁺ is required for lactate-to-pyruvate conversion via lactate dehydrogenase. NAD⁺ levels decline with age and chronic illness. Supplementation with NAD⁺ precursors (nicotinamide riboside 300–1000mg/day, nicotinamide mononucleotide 250–500mg/day) is well-tolerated and raises cellular NAD⁺ levels. If lactate clearance is impaired in ME/CFS, NAD⁺ boosting might accelerate post-exertional recovery. This is testable with lactate measurements before and after NAD⁺ supplementation during controlled exercise challenge.

25.6.11 Mast Cell-Mitochondrial Crosstalk

~ Hypothesis 8: Mast Cell Mediators Damage Mitochondria

The high prevalence of mast cell activation syndrome (MCAS) in ME/CFS suggests potential mechanistic links beyond comorbidity. Histamine receptors are present on mitochondrial membranes and modulate respiration. Chronic release of mast cell mediators (histamine, tryptase, inflammatory cytokines) may directly impair mitochondrial function, creating a positive feedback loop: viral trigger → mast cell activation → mitochondrial damage → cellular stress → further mast cell activation. If validated, aggressive mast cell stabilization (H1/H2 blockers, quercetin, ketotifen) combined with mitochondrial support might synergistically improve both immune and metabolic dysfunction.

25.6.12 Research Priorities

Based on these hypotheses, high-priority research directions include:

1. **WASF3 targeting:** Screen existing WASF3 inhibitors in ME/CFS patient-derived cell lines; measure Complex IV restoration
2. **Mitophagy assessment:** Quantify mitophagy flux and mitochondrial biogenesis kinetics post-exertion in ME/CFS versus controls
3. **ER stress intervention trial:** Test TUDCA during acute viral illness in high-risk individuals (family history, severe EBV)
4. **Complex-specific profiling:** Systematically measure all five mitochondrial complexes in ME/CFS muscle biopsies
5. **Lactate kinetics:** Serial lactate and NAD⁺ measurements during 2-day CPET; NAD⁺ precursor trial
6. **T-cell exhaustion markers:** Flow cytometry for PD-1/TIM-3/LAG-3 expression on ME/CFS T-cells
7. **Pyruvate challenge:** Randomized controlled trial of prophylactic pyruvate before standardized exertion
8. **Methylene blue mechanism:** Dose-finding study with serial mitochondrial function assays
9. **Central governor fMRI:** Brain activation patterns during cognitive tasks at varying metabolic stress levels
10. **Mast cell-mitochondrial interaction:** In vitro studies of histamine effects on mitochondrial respiration in ME/CFS cells

These hypotheses integrate molecular findings (WASF3, viral triggers), patient observations (nicotine, methylene blue), and physiological measurements (2-day CPET, lactate) into testable mechanistic proposals. They represent opportunities to move beyond symptom management toward interventions targeting root pathophysiology.

25.6.13 Summary of Novel Hypotheses and Interventions

Table 25.6 summarizes the mechanistic hypotheses, proposed interventions, evidence basis, and testability for each novel therapeutic direction identified.

Table 25.6: Novel Mechanistic Hypotheses and Therapeutic Opportunities

Hypothesis	Proposed Mechanism	Intervention	Evidence Basis	Testability
WASF3 as drug target	WASF3 inhibition restores Complex IV function	Repurposed WASF3 inhibitors from oncology	Wang 2023 shRNA reversal; cancer drug pipelines	HIGH: Cell culture assays, patient-derived cells
Cholinergic-mito axis	Alpha-7 nAChR on mitochondria regulates ATP	Acetylcholinesterase inhibitors (donepezil)	Patient nicotine reports; nAChR on mitochondria	HIGH: RCT feasible, existing FDA drugs
Mitophagy failure	Impaired mitochondrial autophagy delays ATP recovery	Mitophagy enhancers; NAD ⁺ precursors	24-72h PEM delay matches mitophagy cycles	MEDIUM: Requires muscle biopsy, specialized assays
Viral-ER-WASF3	ER stress from viral infection activates WASF3	TUDCA/4-PBA during acute viral illness	Viral meta-analysis; ER stress-WASF3 link	HIGH: Prophylactic trial in at-risk individuals
Pyruvate bypass	Pyruvate enters TCA directly without glycolysis	Pyruvate drinks pre-exertion (1-2g)	ATP delay; pyruvate enters TCA directly	HIGH: Simple RCT, OTC supplement
Methylene blue enhancement	MB enhances electron transport; reduces oxidative stress	Low-dose MB (1-5mg/day)	Patient reports; established mitochondrial effects	MEDIUM: Dose-finding needed, safety established
Beta-blocker pacing	Pharmacological HR ceiling prevents crashes	Propranolol 10-20mg PRN before high-risk activity	<5 crash rule; cumulative damage	HIGH: Feasible RCT, existing drug
T-cell exhaustion	Checkpoint receptors prevent viral clearance	Anti-PD-1/CTLA-4 (research only)	Rituximab failure; viral persistence	LOW: High risk, requires biomarker validation first
Central governor	Hypersensitive brain limiter prevents ATP crisis	fMRI validation; re-frame symptoms as protective	Exercise physiology theory; brain fog timing	MEDIUM: fMRI studies feasible
Lactate clearance	Impaired lactate-to-pyruvate delays recovery	NAD ⁺ precursors (NR 300-1000mg; NMN 250-500mg/day)	2-day CPET recovery impairment	HIGH: Serial lactate measurement, RCT feasible
MCAS-mito crosstalk	Histamine receptors on mitochondria impair respiration	H1/H2 blockers + mitochondrial support	MCAS comorbidity; histamine-mito link	MEDIUM: In vitro validation, then clinical trial

Prioritization Logic. **Tier 1** interventions are immediately actionable, low-risk, and affordable (pyruvate, NAD⁺ precursors). These can be implemented while awaiting controlled trial results.

Tier 2 interventions require prescriptions or specialized formulations but have established safety profiles (beta-blockers, donepezil, methylene blue, TUDCA). These warrant physician discussion and case-by-case evaluation.

Tier 3 interventions are research-stage only, requiring either drug development (WASF3 inhibitors) or carrying significant risks that preclude clinical use outside trials (checkpoint

Table 25.7: Risk-Benefit Assessment of Novel Interventions

Intervention	Safety Profile	Cost	Implementation Barrier	Priority Tier
Pyruvate (1-2g pre-exertion)	Very safe; OTC supplement	\$15-25/month	None; immediate	Tier 1
NAD+ precursors (NR/NMN)	Safe; well-tolerated	\$40-60/month	None; OTC	Tier 1
Beta-blockers (low-dose PRN)	Safe; established drug	\$5-10/month	Requires prescription	Tier 2
Acetylcholinesterase inhibitors	Safe; approved for dementia	FDA- for \$20-40/month	Requires prescription	Tier 2
Methylene blue (1-5mg)	Safe at low doses; can cause blue urine	\$10-20/month	Compounding needed for low doses	Tier 2
TUDCA (prophylactic)	Safe; bile acid supplement	\$30-50/month	Requires viral illness trigger	Tier 2
WASF3 inhibitors	Unknown; cancer drugs have toxicity	Unknown	Not yet available; research only	Tier 3
Checkpoint inhibitors	HIGH RISK; autoimmune AEs	Very expensive	Research protocol only; extreme risk	Tier 3

inhibitors).

25.7 Working with Healthcare Providers

25.7.1 Communicating the Phenotype-First Approach

Most physicians are unfamiliar with ME/CFS-specific treatment safety considerations. Patients can advocate for a safer treatment approach by:

- **Presenting the energy cost framework:** Explain that ME/CFS patients have reduced capacity to process medications, and that treatment selection should consider metabolic burden (Section 25.1.2)
- **Requesting micro-dosing:** Ask to start at 1/4 to 1/10 of standard doses, with slow titration. Cite the paradoxical reactor phenotype (Section 21.10.1) as clinical justification
- **Advocating for pharmacogenomic testing:** Before Category C medications, request CYP450 metabolizer status testing. Frame as harm prevention, not optional
- **Providing the treatment journal:** A documented history of prior medication reactions, with doses, timing, and outcomes, is the most valuable tool for guiding future treatment decisions
- **Requesting staged introduction:** Insist on one new treatment at a time, with defined observation periods and clear stop criteria

25.7.2 Building a Treatment Team

Optimal ME/CFS management involves:

- **Primary physician:** Coordinates care, manages prescriptions, interprets basic phenotyping (Tier 1)
- **ME/CFS specialist** (if available): Guides phenotype assessment (Tier 2–3), manages complex medication trials, interprets treatment responses
- **Pharmacist:** Reviews drug interactions, particularly CYP450 conflicts when multiple medications are used; advises on compounding for micro-doses
- **Patient:** Maintains treatment journal, tracks symptoms, communicates early warning signs of adverse reactions

Part IV

Research and Evidence Synthesis

This part provides comprehensive coverage of ME/CFS research, synthesizing findings from clinical trials, observational studies, and experimental research. For each major study, we provide:

- Study design and methodology
- Key findings
- Implications for understanding ME/CFS
- Limitations and caveats
- How findings connect to other research

This synthesis approach makes the scattered research literature more accessible and actionable.

26 Biomarker Research

The search for reliable biomarkers in ME/CFS has been a central focus of research for decades. The 2024 NIH deep phenotyping study by Walitt et al. represents a landmark contribution to this effort, identifying multiple objective biological abnormalities that distinguish PI-ME/CFS patients from healthy controls [5]. This chapter reviews the current state of biomarker research, synthesizes findings across multiple biological domains, and discusses the path toward clinically useful diagnostic and prognostic markers.

26.1 Overview of Biomarker Development

26.1.1 Why Biomarkers Are Needed

The absence of validated biomarkers has been one of the most significant obstacles to ME/CFS recognition, research, and treatment:

- **Diagnostic uncertainty:** Without objective markers, diagnosis relies entirely on clinical criteria and exclusion of other conditions
- **Stigmatization:** Lack of measurable abnormalities has contributed to the perception of ME/CFS as a psychosomatic condition
- **Research challenges:** Heterogeneous patient populations (due to imprecise diagnosis) may obscure findings
- **Treatment development:** Drug development requires objective endpoints for clinical trials
- **Disability assessment:** Social security and insurance determinations benefit from objective evidence
- **Subgroup identification:** Biomarkers may identify pathophysiologically distinct subgroups requiring different treatments

26.1.2 Types of Biomarkers

Different biomarker types serve different purposes:

Diagnostic Biomarkers

Markers that distinguish ME/CFS from healthy individuals and from patients with other fatiguing conditions:

- High sensitivity (few false negatives)
- High specificity (few false positives)
- Practical for clinical use (accessible, affordable)
- Reproducible across laboratories

Prognostic Biomarkers

Markers that predict disease course or outcome:

- Likelihood of spontaneous improvement
- Risk of progression to more severe illness
- Long-term functional outcomes

Treatment Response Biomarkers

Markers that predict or monitor response to specific treatments:

- Baseline markers predicting treatment response
- Dynamic markers reflecting treatment effects
- Stratification markers for personalized treatment selection

Mechanistic Biomarkers

Markers that reflect underlying pathophysiology:

- May not be diagnostic but inform disease mechanisms
- Guide development of targeted therapies
- Enable subgroup classification

26.1.3 Challenges in ME/CFS Biomarker Research

Multiple factors have complicated biomarker identification:

- **Case definition heterogeneity:** Different diagnostic criteria capture overlapping but distinct populations
- **Disease heterogeneity:** ME/CFS likely encompasses multiple distinct conditions with different pathophysiology
- **Illness duration effects:** Biomarkers may differ between early and chronic illness
- **Severity effects:** Severely affected patients (often excluded from studies) may differ from ambulatory patients
- **Sex differences:** The NIH study demonstrated distinct abnormalities in men and women

- **Comorbidities:** Overlapping conditions (POTS, MCAS, fibromyalgia) may confound findings
- **Small sample sizes:** Many studies underpowered to detect moderate effect sizes
- **Lack of replication:** Few findings have been consistently replicated across laboratories

26.1.4 Convergent Biology Despite Heterogeneity

Despite the challenges listed above, emerging evidence suggests that heterogeneous upstream triggers converge on common downstream pathological pathways, with direct implications for biomarker strategy:

1. **miRNA convergence:** Cheema et al. 2023 in *PLOS ONE* [11] found that circulating miRNAs, although different between individual patients, target the same specific gene cluster ($p < 0.002$) converging on exercise hyperemia, angiogenesis, antioxidant defenses, and mitochondrial fission. This suggests heterogeneous molecular perturbations funnel into common downstream pathology measurable through functional pathway biomarkers rather than single-molecule markers.
2. **Metabolic convergence:** Naviaux et al. 2016 in *PNAS* [141] showed that despite heterogeneous triggers, the cellular metabolic response was homogeneous—a dauer-like hypometabolic state affecting 20 pathways with 80% of diagnostic metabolites decreased and 94–96% diagnostic accuracy.
3. **Genetic convergence:** Birch & Younger 2025 in *J Transl Med* [142] identified gene-level heterogeneity but convergence on impaired energy production, reduced stress resilience, and vulnerability to post-exertional metabolic failure.
4. **Emerging large-scale validation panels:** The EpiSwitch CFS test (Myhill et al. 2025, *J Transl Med*) [215] achieved 92% sensitivity and 98% specificity using a 200-marker 3D genomic profiling model in severe ME/CFS, and the BioQuest project (OMF, launched 2024) [615] aims to analyze approximately 1,000 samples across metabolomics, proteomics, and cytokines with subgrouping as a primary aim.

The practical implication of these convergence patterns is that biomarker strategies should target convergent downstream pathways (energy metabolism, vascular function, immune dysregulation) rather than seeking a single upstream marker. As argued by Birch & Younger [142], multi-marker panels stratifying patients by endotype are more likely to succeed than individual biomarkers applied to the entire population.

⚠ **Limitation 1: ME/CFS Biomarker Research: No Validated Diagnostic Biomarker Exists**

Despite decades of research and multiple candidate markers, no biomarker or biomarker panel has achieved clinical validation for ME/CFS diagnosis. Critical limitations:

- The NIH deep phenotyping study [5], while methodologically rigorous, enrolled only 17 PI-ME/CFS patients; all findings require large-scale replication before di-

agnostic claims can be made.

- Published multi-marker panels reporting >90% sensitivity/specificity have not been independently validated in diverse populations; performance in discovery cohorts routinely exceeds performance in external validation.
- Biomarker-disease associations documented in this chapter (cytokine profiles, NK cell dysfunction, metabolomic signatures) demonstrate group-level differences; individual-level diagnostic accuracy remains uncertain for all candidates.
- The convergence patterns described (miRNA, metabolic, genetic) are derived from separate studies using different cohorts, criteria, and methodologies; no single study has demonstrated convergence across all domains in the same patients.
- Commercial biomarker tests currently marketed for ME/CFS lack independent validation and should not be interpreted as diagnostic.

26.2 Key Biomarkers from the NIH Deep Phenotyping Study

The Walitt et al. study provides a template for comprehensive biomarker identification, employing rigorous methodology with 17 PI-ME/CFS patients and 21 matched controls [5]. The multi-domain assessment identified several categories of potential biomarkers.

26.2.1 Cerebrospinal Fluid Biomarkers

Catecholamine Metabolites

CSF analysis revealed significantly reduced catecholamine levels:

- **Homovanillic acid (HVA):** Primary dopamine metabolite; reduced in PI-ME/CFS
- **3-methoxy-4-hydroxyphenylglycol (MHPG):** Norepinephrine metabolite; reduced
- **Clinical correlation:** Levels correlated with motor performance, effort behaviors, and fatigue severity
- **Biomarker potential:** Objective, measurable, correlates with symptoms

Tryptophan Pathway Metabolites

Altered tryptophan metabolism documented:

- Kynurenine pathway metabolite abnormalities
- Potential serotonin precursor depletion
- Links immune activation to neurological symptoms

26.2.2 Immune Biomarkers

B Cell Population Shifts

Characteristic pattern documented:

- **Increased naïve B cells:** Elevated compared to controls
- **Decreased switched memory B cells:** Reduced class-switched memory population
- **Interpretation:** Pattern suggests chronic antigenic stimulation
- **Diagnostic potential:** Specific pattern may distinguish ME/CFS from other conditions

Sex-Specific Immune Markers

Striking differences between sexes:

- **Males:** Altered T cell activation markers, innate immunity changes
- **Females:** Abnormal B cell proliferation, distinct white blood cell patterns
- **Implications:** Biomarkers may need sex-specific interpretation

26.2.3 Autonomic Biomarkers

Heart Rate Variability

Reduced HRV documented:

- Diminished overall variability (SDNN)
- Reduced high-frequency power (parasympathetic marker)
- Non-invasive, widely available measurement
- Correlates with symptom severity

Baroreflex Sensitivity

Impaired baroreflex function:

- Reduced cardiovagal gain
- Indicates parasympathetic dysfunction
- Objective, quantifiable measure

26.2.4 Cardiopulmonary Biomarkers

Exercise Testing Parameters

CPET abnormalities:

- **Reduced VO_2 peak:** Objective measure of aerobic capacity
- **Chronotropic incompetence:** Inadequate heart rate response
- **Two-day decline:** Failure to reproduce performance (highly specific)
- **Reduced anaerobic threshold:** Earlier reliance on anaerobic metabolism

26.2.5 Neuroimaging Biomarkers

Functional MRI Findings

Brain activity abnormalities:

- **Reduced TPJ activity:** During effort-based tasks
- **Motor cortex hyperactivity:** Despite declining performance
- **Altered effort perception:** Neural correlates of fatigue

26.3 Metabolomic Biomarkers

Metabolomics—the comprehensive study of small molecule metabolites—has emerged as a promising approach to ME/CFS biomarker discovery.

26.3.1 Key Metabolomic Studies

Naviaux et al. Studies

Landmark metabolomic investigations found:

- Hypometabolic state resembling “dauer” (*C. elegans* survival mode)
- Abnormalities in sphingolipid, phospholipid, and purine metabolism
- Reduced metabolites across multiple pathways
- Pattern suggesting coordinated metabolic downregulation

Amino Acid Profile Abnormalities

Multiple studies report altered amino acids:

- **Branched-chain amino acids:** Often reduced
- **Glutamine/glutamate:** Altered ratios
- **Tryptophan:** Reduced (diverted to kynurenine pathway)
- **Arginine:** May be depleted (NO synthesis)

Lipid Metabolism Markers

Abnormal lipid profiles:

- Altered phosphatidylcholine species
- Abnormal ceramide levels
- Changed fatty acid profiles
- Reduced omega-3 fatty acids in some studies

TCA Cycle Metabolites

Krebs cycle abnormalities:

- Altered citrate, isocitrate, succinate levels
- Suggests impaired oxidative metabolism
- Correlates with mitochondrial dysfunction hypothesis

26.3.2 Synthesis of Metabolomic Findings

Common Patterns

Despite methodological differences, several patterns emerge:

- Hypometabolic signature (reduced metabolites across pathways)
- Impaired energy metabolism
- Oxidative stress markers
- Altered lipid metabolism

Subgroup Differences

Metabolomic studies may identify subgroups:

- Different metabolic signatures in different patients
- Potential for metabolomics-based classification
- Treatment response prediction

Clinical Utility

Current status:

- Not yet validated for clinical diagnosis
- Research tool for understanding pathophysiology
- Potential for future diagnostic panels
- Requires standardization and replication

26.4 Immunological Biomarkers

26.4.1 Cytokine Profiles

Studies Identifying Cytokine Patterns

Numerous studies have examined cytokines in ME/CFS:

- **Early illness:** More consistent elevation of pro-inflammatory cytokines
- **Chronic illness:** More variable, often normalized
- **Specific cytokines:** IL-1, IL-6, TNF- α , IFN- γ variably elevated
- **Cytokine networks:** Pattern analysis may be more informative than individual cytokines

Variability and Consistency

Challenges in cytokine research:

- Different assays with different sensitivities
- Timing of blood draw (diurnal variation)
- Recent activity effects
- Heterogeneous patient populations

Correlation with Symptoms

When correlations are found:

- Higher cytokines often correlate with greater severity
- Cytokine patterns may predict symptom clusters
- Post-exertional changes in cytokines documented

26.4.2 Cell Function Markers

NK Cell Activity

One of the most replicated findings:

- Reduced cytotoxic function in most studies
- 40–60% reduction compared to controls
- Correlates with severity in some studies
- Functional assay more informative than cell counts

T Cell Markers

Various abnormalities reported:

- Exhaustion markers (PD-1, Tim-3)
- Altered CD4/CD8 ratios (inconsistent direction)
- Reduced regulatory T cell function
- Th1/Th2 imbalance

B Cell Profiles

NIH study findings highlight B cell importance:

- Naïve/memory B cell ratio shift
- Chronic antigenic stimulation pattern
- Potential autoantibody-producing populations

26.5 Neurological Biomarkers

26.5.1 Brain Imaging Markers

Structural MRI

Documented abnormalities:

- White matter hyperintensities (variable)
- Regional gray matter volume changes
- Brainstem abnormalities in some studies

Functional MRI

NIH study and others show:

- Altered activation patterns during tasks
- TPJ dysfunction during effort tasks
- Connectivity changes
- Potential for task-based biomarkers

PET and SPECT

Metabolic and perfusion imaging:

- Regional hypometabolism
- Reduced cerebral blood flow
- Neuroinflammation markers (TSPO binding)

26.5.2 CSF Findings

Beyond the NIH catecholamine findings:

- Elevated inflammatory markers in some studies
- Altered protein profiles
- Potential autoantibodies
- Oligoclonal bands in subset

26.5.3 Autonomic Function Tests

Quantifiable autonomic biomarkers:

- **Tilt table testing:** POTS, NMH, OH patterns
- **Heart rate variability:** Multiple parameters
- **Sudomotor function:** QSART abnormalities
- **Pupillometry:** Altered light reflexes

26.5.4 Cognitive Testing Patterns

Neuropsychological profiles:

- Processing speed reduction (most consistent)
- Attention and working memory deficits
- Variable memory findings
- Pattern different from depression or anxiety

26.6 Genomic and Epigenetic Biomarkers

26.6.1 Gene Expression Signatures

Peripheral Blood Transcriptomics

Multiple studies have examined gene expression:

- Differential expression of immune-related genes
- Metabolic gene abnormalities
- Mitochondrial gene expression changes
- Potential diagnostic signatures

Sex-Specific Gene Expression

NIH study found distinct patterns:

- Different genes differentially expressed in men vs. women
- Muscle biopsy gene expression differences
- Supports sex-specific disease mechanisms

26.6.2 miRNA Profiles

MicroRNAs regulate gene expression:

- Altered circulating miRNA profiles in ME/CFS
- May reflect underlying pathway dysregulation
- Potential for minimally invasive biomarkers
- Requires further validation

26.6.3 DNA Methylation Patterns

Epigenetic modifications:

- Altered methylation at specific sites
- May reflect environmental exposures or disease state
- Potential for stable biomarkers
- Early-stage research

26.6.4 Clinical Utility

Current status of genomic biomarkers:

- Research tools primarily
- Not yet validated for clinical use
- Potential for future multi-marker panels
- May enable personalized treatment selection

26.7 Proteomic Biomarkers

26.7.1 Protein Expression Patterns

Mass spectrometry-based proteomics:

- Altered plasma/serum protein profiles
- Inflammatory proteins frequently identified
- Complement components
- Coagulation factors

26.7.2 Autoantibody Panels

Functionally significant autoantibodies:

- **Anti- β -adrenergic receptor:** 25–30% of patients
- **Anti-muscarinic receptor:** Significant subset
- **Anti-neuronal antibodies:** Variable findings
- **Diagnostic potential:** May identify autoimmune subgroup

26.7.3 Diagnostic Potential

Proteomics status:

- Multiple candidate proteins identified
- Replication across studies limited
- Potential for panel-based diagnosis
- Autoantibody testing closest to clinical use

26.8 Composite Biomarker Panels

26.8.1 Multi-Omics Approaches

Integrating multiple biomarker types:

- Combining metabolomics, proteomics, transcriptomics
- Machine learning for pattern recognition
- May capture disease complexity better than single markers
- Requires large, well-characterized cohorts

26.8.2 Machine Learning Applications

Computational approaches to biomarker discovery:

- Random forests, neural networks for classification
- Feature selection to identify most informative markers
- Integration of clinical and molecular data
- Cross-validation to prevent overfitting

26.8.3 Diagnostic Accuracy

Published multi-marker panels:

- Some report >90% sensitivity and specificity
- Independent validation often shows lower performance
- Need for prospective validation in diverse populations
- Comparison to clinical diagnosis as gold standard problematic

26.8.4 Commercial Tests Available

Current commercial offerings:

- Several proprietary tests marketed
- Limited independent validation
- Variable acceptance by clinicians and insurers
- Ongoing development of improved panels

26.9 Functional Biomarkers

26.9.1 Two-Day CPET Protocol

Perhaps the most specific biomarker for ME/CFS:

- **Methodology:** Maximal exercise testing on consecutive days
- **Finding:** 10–25% decline in VO_2peak , AT, work capacity on Day 2
- **Specificity:** Healthy controls and patients with other conditions reproduce or improve
- **Physiological basis:** Reflects post-exertional malaise objectively
- **Limitations:** Requires specialized equipment, may exacerbate symptoms

26.9.2 NASA Lean Test

Simple orthostatic assessment:

- Patient leans against wall for 10 minutes
- Heart rate and blood pressure monitored
- Identifies POTS and other orthostatic disorders
- Accessible, low-tech screening tool

26.9.3 Cognitive Testing

Standardized neuropsychological assessment:

- Processing speed measures (e.g., Symbol Digit Modalities Test)
- Attention tests (e.g., continuous performance tasks)
- Pattern of deficits may distinguish from depression
- Sensitive to post-exertional cognitive deterioration

26.10 Biomarker Validation and Standardization

26.10.1 Replication Requirements

For a biomarker to be clinically useful:

- Replication in independent cohorts
- Consistent findings across laboratories
- Validation in diverse patient populations
- Demonstration of clinical utility (changing management)

26.10.2 Standardization Efforts

Ongoing initiatives:

- **Case definition harmonization:** Using consistent diagnostic criteria
- **Biobanking:** Standardized sample collection and storage
- **Assay standardization:** Consistent methodologies across sites
- **Data sharing:** Collaborative analysis of combined datasets

26.10.3 Path to Clinical Implementation

Steps required:

1. Discovery phase (identifying candidate biomarkers)
2. Verification (confirming in independent samples)
3. Validation (large-scale, multi-site studies)
4. Clinical utility studies (demonstrating impact on outcomes)
5. Regulatory approval (for diagnostic claims)
6. Implementation (clinical adoption, insurance coverage)

26.11 Summary: Current State and Future Directions

The NIH deep phenotyping study represents a paradigm for rigorous biomarker research in ME/CFS [5]. Key findings with biomarker potential include:

1. **CSF catecholamine metabolites:** Reduced HVA and MHPG correlating with symptoms; invasive but highly specific
2. **B cell population shifts:** Increased naïve, decreased switched memory B cells suggesting chronic antigenic stimulation; accessible via routine blood draw
3. **Autonomic parameters:** Reduced HRV and baroreflex sensitivity; non-invasive, widely available technology
4. **CPET abnormalities:** Reduced VO_2 peak, chronotropic incompetence, and especially Day 2 decline; objective, physiologically meaningful
5. **Neuroimaging findings:** TPJ dysfunction and motor cortex hyperactivity; research tool with potential clinical application
6. **Sex-specific patterns:** Different immune markers in men vs. women; critical for biomarker interpretation

The path forward requires:

- Large-scale replication of NIH study findings
- Development of practical, accessible biomarker panels
- Validation across diverse patient populations
- Integration of multiple biomarker types for improved accuracy
- Demonstration of clinical utility for diagnosis and treatment selection

The era of “no objective findings” in ME/CFS is ending. The challenge now is translating research discoveries into clinically useful tools that improve patient care.

27 Clinical Trials and Treatment Studies

Clinical trial evidence in ME/CFS is limited relative to other chronic diseases, reflecting decades of underfunding and the contested status of the illness. Most trials are small ($n < 50$), single-center, and lack placebo controls or objective outcome measures. This chapter reviews the available evidence by intervention category, with particular attention to methodological quality and the lessons each trial provides for future research.

27.1 Immunological Interventions

27.1.1 Rituximab Trials

The rituximab program in ME/CFS represents one of the most instructive sequences in the field—from serendipitous observation through promising pilot data to definitive negative results.

Fluge and Mella (2011) reported clinical improvement in ME/CFS patients receiving rituximab for incidental lymphoma, leading to an open-label pilot ($n=3$) and then a small double-blind RCT ($n=30$) [616]. The RCT showed a significant response rate (67% rituximab vs. 13% placebo), with responses typically delayed 3–6 months after infusion—consistent with the time required for existing plasma cells to die naturally after B cell depletion.

An open-label phase II study ($n=29$) confirmed response patterns and explored maintenance dosing [117]. However, the definitive phase III multicenter RCT (RituxME, $n=151$) found no significant difference between rituximab and placebo at 24 months [376]. Response rates were 26% in the rituximab group versus 35% in the placebo group.

The RituxME failure teaches several lessons:

- **Patient selection:** ME/CFS is likely heterogeneous; B cell-mediated autoimmunity may drive symptoms in a subgroup but not the majority. Biomarker-stratified trials (e.g., selecting patients with elevated GPCR autoantibodies) may identify responders
- **Placebo response:** The 35% placebo response rate is high, suggesting that unblinded earlier studies overestimated true efficacy. Double-blind design is essential
- **Plasma cell persistence:** Rituximab depletes $CD20^+$ B cells but spares $CD20^-$ plasma cells. If autoantibodies from long-lived plasma cells drive symptoms, rituximab is the wrong target (see Chapter 36 for mathematical modeling of this distinction)

27.1.2 Low-Dose Naltrexone Studies

Low-dose naltrexone (LDN, 1.5–4.5 mg nightly) has become one of the most widely used off-label treatments for ME/CFS despite limited trial evidence. The proposed mechanism involves transient opioid receptor blockade leading to upregulation of endogenous opioid production and downstream immunomodulation, particularly microglial modulation and restoration of TRPM3 ion channel function in NK cells [469, 536].

Available evidence includes:

- **Retrospective surveys:** Patient surveys report 50–74% positive response rates, with improvements in fatigue, pain, sleep, and cognitive function. However, retrospective self-report is highly susceptible to bias
- **Fibromyalgia RCTs:** Two small placebo-controlled RCTs in fibromyalgia (which substantially overlaps with ME/CFS) showed significant pain reduction with LDN [468]. Effect sizes were moderate
- **NK cell function:** Cabanas et al. (2021) demonstrated that naltrexone restores TRPM3 ion channel function in NK cells from ME/CFS patients *in vitro*, providing a plausible immunological mechanism [536]
- **Ongoing RCTs:** Multiple double-blind RCTs are underway (UAB Younger lab dose-finding study, UBC Nacul RCT, LIFT trial combining LDN with pyridostigmine), which should provide definitive efficacy data within 2–3 years

⚠ Limitation 1: LDN Evidence Quality

Despite widespread use, LDN for ME/CFS lacks a single published, adequately powered, placebo-controlled RCT. Current enthusiasm is based on mechanistic plausibility, fibromyalgia trials, retrospective surveys, and clinical experience. The ongoing RCTs are critically needed.

27.1.3 Other Immune Modulators

Several other immunomodulatory approaches have been tested in ME/CFS:

- **Intravenous immunoglobulin (IVIG):** Three RCTs in the 1990s produced mixed results. Lloyd et al. (1990, n=49) showed improvement; Vollmer-Conna et al. (1997, n=99) showed no benefit. A subset of patients—particularly those with documented immunoglobulin deficiency—may respond. High cost and limited availability restrict use
- **Interferon-alpha:** Small studies showed modest benefit in a subgroup of patients with documented viral reactivation. Side effects (flu-like symptoms, fatigue, depression) limit tolerability in a population already suffering these symptoms
- **Rintatolimod (Ampligen):** A double-stranded RNA drug with immunomodulatory and antiviral properties. The largest ME/CFS-specific RCT (n=234) showed significant improvement in exercise tolerance (treadmill duration) but the FDA declined approval due to concerns about effect size and adverse event profile. Available on a cost-recovery basis in some countries

- **Immunoadsorption:** Removal of autoantibodies via extracorporeal adsorption. Case series from the Charité in Berlin reported improvement in patients with elevated GPCR autoantibodies. Controlled trials are ongoing

27.2 Antiviral Trials

The hypothesis that persistent viral infection drives ME/CFS has motivated several antiviral trials, with results that highlight the importance of patient selection.

Valganciclovir: Montoya et al. (2013, n=30) conducted a double-blind RCT of valganciclovir in ME/CFS patients selected for elevated HHV-6 and EBV antibody titers. The primary endpoint (fatigue score at 9 months) did not reach significance, but a subgroup with higher baseline antibody titers showed improvement. This suggested that antiviral therapy may benefit a virally-driven subgroup rather than the general ME/CFS population.

Valacyclovir: Lerner et al. (2007) reported improvement in a subset of ME/CFS patients with elevated EBV antibody titers treated with prolonged valacyclovir (6+ months). The study was open-label and uncontrolled, limiting conclusions.

Paxlovid (nirmatrelvir/ritonavir): The NIH STOP-PASC trial tested Paxlovid for long COVID (which overlaps substantially with ME/CFS). The result was negative—no significant benefit over placebo—suggesting that if viral persistence drives symptoms, a brief antiviral course is insufficient to clear established reservoirs.

Key lessons from antiviral trials:

- Patient selection by viral biomarkers is essential; unselected ME/CFS populations include non-virally-driven cases
- Treatment duration may need to be prolonged (months) given evidence of viral persistence in tissue reservoirs
- Combination antiviral strategies targeting multiple latent viruses have not been tested
- Novel antivirals targeting specific viral persistence mechanisms (e.g., EBV latency-to-lytic transition) may be more effective than existing agents

27.3 Metabolic and Mitochondrial Interventions

Given the documented metabolic dysfunction in ME/CFS (Chapter 6), several trials have tested supplements targeting mitochondrial function:

- **Coenzyme Q10 (CoQ10):** Castro-Marrero et al. (2015, n=73) showed significant improvement in fatigue scores with CoQ10 (200 mg) plus NADH (20 mg) versus placebo over 8 weeks. A subsequent 12-week trial confirmed benefits for fatigue and sleep quality. CoQ10 supports electron transport chain function at Complex III

- **D-ribose:** Teitelbaum et al. (2006) reported significant improvement in energy, sleep, cognitive function, and pain in an open-label study (n=41) of D-ribose 5 g three times daily. D-ribose is a precursor for ATP synthesis via the pentose phosphate pathway. No placebo-controlled RCT has been published
- **NADH:** Forsyth (1999, n=26) conducted a crossover RCT showing significant improvement with sublingual NADH 10 mg versus placebo (8/26 responded to NADH vs. 2/26 to placebo). NADH is an essential electron carrier in the mitochondrial electron transport chain
- **Carnitine:** Plioplys and Plioplys (1997, n=30) found significant improvement in fatigue severity with L-carnitine 3 g/day over 8 weeks. Acetyl-L-carnitine facilitates fatty acid transport into mitochondria for beta-oxidation. More recent studies suggest that carnitine deficiency may be present in a subgroup of ME/CFS patients

⚠ **Limitation 2: Metabolic Trial Quality**

Most metabolic intervention trials are small, single-center, and use subjective outcomes. The CoQ10+NADH combination has the strongest evidence base. Larger, multi-center RCTs with objective metabolic endpoints (respirometry, metabolomics) are needed.

27.4 Neurological Interventions

Neurological interventions target the central nervous system dysfunction documented in ME/CFS (Chapter 8):

- **Methylphenidate:** Blockmans et al. conducted a double-blind randomized crossover RCT (n=60) of methylphenidate 10 mg twice daily [617]. Significant improvement was found in fatigue and concentration but not in overall quality of life. Response was most pronounced in patients with prominent cognitive symptoms
- **Low-dose aripiprazole (LDA):** A pilot study from Stanford (n=25) reported that very low doses (0.2–2 mg) improved fatigue, cognitive function, and PEM severity in some patients [602]. The proposed mechanism involves dopamine modulation and microglial regulation. No large RCT has been published; controlled trials are needed
- **Transcranial direct current stimulation (tDCS):** Studies suggest that non-invasive brain stimulation targeting the dorsolateral prefrontal cortex may improve cognitive function and fatigue [513]. The technique is safe, low-cost, and amenable to home use, making it attractive for ME/CFS. Controlled trials are in early stages
- **Transcutaneous vagus nerve stimulation (tVNS):** Vagus nerve stimulation modulates autonomic function and reduces neuroinflammation [618]. A multi-site RCT is currently recruiting. Home-use auricular tVNS devices are available and being explored in patient communities [497]

27.5 Exercise and Rehabilitation Trials

Exercise trials in ME/CFS have been among the most controversial in the field, centering on the PACE trial and the subsequent dismantling of its conclusions.

27.5.1 The PACE Trial

The PACE trial (White et al., 2011, n=641) was the largest ME/CFS treatment study ever conducted, comparing graded exercise therapy (GET), cognitive behavioral therapy (CBT), adaptive pacing therapy (APT), and specialist medical care (SMC). Initial results claimed that GET and CBT were moderately effective.

Subsequent reanalysis and criticism revealed fundamental problems:

- **Outcome measure changes:** Primary outcomes were changed mid-trial, lowering the threshold for “improvement” to a level where patients could deteriorate and still be classified as “recovered” [506]
- **Subjective outcomes only:** No objective measures (step counts, actigraphy, employment status) showed improvement
- **Non-blinded design:** Therapist-delivered interventions with subjective self-report outcomes are inherently vulnerable to expectation bias [57]
- **Long-term follow-up:** The trial’s own long-term follow-up showed no sustained benefit; all groups converged to similar outcomes
- **Patient-reported harms:** Large surveys consistently report that 51% of patients worsened with GET [586]

The PACE trial led NICE to withdraw its recommendation for GET in the 2021 guideline update [127], a landmark decision that acknowledged the evidence for harm.

27.5.2 Alternative Rehabilitation Approaches

Rehabilitation approaches that respect the energy envelope show more promise:

- **Activity management/pacing:** Consistently rated as the most helpful intervention in patient surveys. Supported by the Workwell Foundation’s CPET-based activity prescription model (Section 23.1)
- **Occupational therapy:** Adapted OT focusing on energy conservation, adaptive equipment, and environmental modification rather than progressive activity increases
- **Aquatic therapy:** Water buoyancy reduces gravitational stress, potentially allowing movement that would trigger PEM on land. Evidence is anecdotal; controlled trials are lacking

27.6 Complementary and Alternative Medicine Trials

Complementary and alternative medicine (CAM) approaches are widely used by ME/CFS patients, often in the absence of effective conventional treatments. Evidence quality is generally low:

- **Acupuncture:** A 2020 Cochrane-style systematic review identified several small RCTs, mostly from Chinese literature, reporting benefit for fatigue. Methodological quality was generally low (inadequate blinding, unclear randomization). Sham-controlled trials show inconsistent results. Acupuncture may provide modest symptomatic relief but is unlikely to modify disease course
- **Mind-body interventions:** Qigong, tai chi, and yoga have been studied in small trials with generally positive but methodologically weak results. Gentle, adapted versions may be tolerated within the energy envelope and provide modest benefit for pain, sleep, and psychological well-being. Standard-intensity practices may trigger PEM
- **Homeopathy:** No rigorous evidence of efficacy beyond placebo for ME/CFS
- **Nutritional supplements:** See Chapter 22 for detailed evidence on CoQ10, D-ribose, NADH, carnitine, and other nutraceuticals

27.7 Ongoing and Planned Trials

The ME/CFS clinical trial landscape has expanded significantly since 2020, partly driven by overlap with long COVID research. Key ongoing or recently initiated trials include:

- **LIFT Trial** (Brigham & Women's Hospital): LDN combined with pyridostigmine for ME/CFS. Double-blind RCT
- **UAB LDN dose-finding** (University of Alabama, Younger lab): Establishing optimal LDN dosing for ME/CFS
- **Nacul LDN RCT** (UBC / BC Women's Hospital): Double-blind RCT of LDN for ME/CFS
- **Lumbrokinase trial** (Mount Sinai): Fibrinolytic agent targeting microclots in ME/CFS and long COVID
- **Rapamycin trial** (multi-site, US): mTOR inhibition for immune and metabolic modulation
- **CHIIME** (UCSF / PolyBio Research Foundation): Investigating viral tissue persistence in ME/CFS and long COVID
- **Masitinib** (AB Science): Mast cell inhibitor in phase II for ME/CFS with MCAS features
- **tVNS multi-site trial:** Transcutaneous vagus nerve stimulation for autonomic and immune modulation
- **Wearable pacing study** (Scripps Research): Validating wearable-guided pacing protocols

Patients seeking to participate in clinical trials can search ClinicalTrials.gov (filtering by “myalgic encephalomyelitis” or “chronic fatigue syndrome”), contact national ME/CFS organizations for trial registries, or consult the research registry in Appendix I.

27.8 Synthesis and Meta-Analyses

Synthesizing the ME/CFS trial literature reveals consistent patterns:

What shows promise (moderate evidence):

- **Pacing/activity management:** Consistently rated most helpful by patients; supported by CPET evidence
- **LDN:** Strong mechanistic rationale, positive patient reports, awaiting definitive RCTs
- **CoQ10 + NADH:** Small but positive RCT evidence for fatigue reduction
- **Rintatolimod:** Positive RCT for exercise tolerance; regulatory and access issues persist

What has failed or shown harm:

- **Graded exercise therapy:** Definitive evidence of harm from patient surveys and guideline withdrawals
- **Standard CBT** (activity-increasing variant): No objective benefit; potential for harm when it encourages envelope violation
- **Rituximab:** Phase III RCT negative in unselected population
- **Brief antiviral courses:** Insufficient to clear established viral reservoirs

Critical gaps in the evidence base:

- No FDA- or EMA-approved treatment for ME/CFS exists
- Most trials are underpowered ($n < 50$) and use subjective outcomes
- Biomarker-stratified trials (selecting patients by mechanism) have barely begun
- Combination therapies addressing multiple pathophysiological mechanisms simultaneously have not been tested
- Severe and very severe patients are systematically excluded from trials due to inability to attend research sites
- Two-day CPET has not been used as an outcome measure in treatment trials despite being the most objective measure of ME/CFS-specific physiology

? Open Question 1: Biomarker-Stratified Clinical Trials

The failure of rituximab in an unselected ME/CFS population, contrasted with apparent benefit in patients with elevated autoantibodies, suggests that ME/CFS comprises mechanistically distinct subgroups. Future trials should stratify by: (1) autoantibody profile (for immunomodulatory interventions), (2) viral reactivation markers (for antiviral interventions), (3) metabolomic profile (for mitochondrial support), and (4) autonomic subtype (for cardiovascular interventions). The challenge is developing validated biomarker panels for stratification before launching large trials.

27.9 Negative Trials and Null Results

Understanding why treatments fail is as informative as understanding why they succeed. This section documents major negative, null, and inconclusive clinical trials in ME/CFS. Each entry identifies not only the outcome but the lessons that should guide future trial design.

⚠ Limitation 3: Publication Bias in Negative Trials

Negative trials are systematically underreported in the medical literature. The trials documented here represent those that were published; an unknown number of negative results may remain in file drawers. This section necessarily reflects survivorship bias in published negative results.

27.9.1 Immunological Interventions

Rituximab: From Promise to Definitive Negative

The rituximab story is the most instructive negative trial in ME/CFS research, illustrating how small-sample enthusiasm can collapse in larger, properly controlled studies.

Phase II (Fluge et al., 2011). A small double-blind, placebo-controlled trial ($n = 30$) of two rituximab infusions (500 mg/m^2) showed a response rate of 67% in the rituximab group vs. 13% placebo at 12 months, with responses typically appearing 3–8 months after infusion [616]. The delayed response pattern suggested an autoimmune mechanism requiring time for pathogenic antibody turnover.

Phase III — RituxME (Fluge et al., 2019). The definitive phase III trial enrolled 151 patients across four Norwegian hospitals in a double-blind, placebo-controlled design with six rituximab infusions over 12 months [376].

Primary outcomes: Neither co-primary endpoint—self-reported fatigue (Fatigue Severity Scale) nor overall function (SF-36 Physical Function)—showed significant differences between groups ($p = 0.81$ for overall response rate: 26% rituximab vs. 35% placebo). The trial was unambiguously negative.

Lessons learned.

- **Heterogeneity dilutes subgroup signals:** A 67% response in a small, possibly enriched sample collapsed to no signal in an unstratified larger trial.
- **B-cell depletion may target the wrong compartment:** Subsequent daratumumab (anti-CD38, targeting plasma cells) showed a 60% response rate [9], suggesting long-lived plasma cells—not B cells—are the relevant autoantibody source.
- **Biomarker stratification is essential:** Future immunotherapy trials require pre-selection by autoantibody status.

Intravenous Immunoglobulin (IVIG): Contradictory Results

IVIG trials produced frustratingly inconsistent results spanning clearly positive to clearly negative outcomes.

Lloyd et al. (1990). The first placebo-controlled IVIG trial ($n = 49$) enrolled patients with immunological abnormalities. Approximately 43% of IVIG-treated patients showed significant functional improvement [619]. Low baseline CD4 count predicted response.

Vollmer-Conna et al. (1997). A larger follow-up ($n = 99$) tested three IVIG doses monthly for three months [620]. No dose showed benefit over placebo on any outcome.

Lessons learned.

- **Patient selection determines outcome:** Lloyd's positive trial required immunological abnormalities; Vollmer-Conna's did not.
- **Immune biomarkers may predict response:** Low CD4 count, IgG subclass deficiency, and age at onset may identify IVIG-responsive subgroups.

27.9.2 Behavioral and Rehabilitation Interventions

The PACE Trial: GET and CBT

The PACE trial remains the most controversial study in ME/CFS research.

Original trial (White et al., 2011). A large ($n = 641$), four-arm, assessor-blind RCT tested CBT, graded exercise therapy (GET), adaptive pacing therapy, and specialist medical care alone [115]. The original publication reported CBT and GET were “moderately effective,” with recovery rates of approximately 22%.

Reanalysis (Wilshire et al., 2018). Using the originally published protocol thresholds, Wilshire et al. found [506]:

- Recovery rates dropped from 22% to approximately 7–8%
- No objective measures (6-minute walk, actigraphy, employment) improved
- “The modest treatment effects do not exceed what could be accounted for by participant reporting biases”

In 2021, NICE removed GET and CBT-as-treatment from its ME/CFS guidelines.

Lessons learned.

- **Objective outcomes are non-negotiable:** Self-reported outcomes in unblinded behavioral trials cannot be sole primary endpoints.
- **Protocol fidelity matters:** Mid-trial changes to outcome thresholds undermine validity.
- **Data transparency is essential:** The reanalysis was possible only through Freedom of Information litigation.

27.9.3 Antiviral Interventions**Acyclovir (Straus et al., 1988)**

An early RCT ($n = 27$) tested IV then oral acyclovir for persistent EBV [621]. No significant difference between groups.

Lesson: Wrong antiviral spectrum (acyclovir has minimal HHV-6 activity) and no virus-specific patient selection.

Valganciclovir — Mixed Results (Montoya et al., 2013)

The EVOLVE trial ($n = 30$) of valganciclovir in patients with elevated HHV-6/EBV titers [149]. Primary composite endpoint: not significant. However, individual endpoints showed significant improvements in mental fatigue ($p = 0.039$), fatigue severity ($p = 0.006$), and cognition ($p = 0.025$).

Lesson: Composite endpoints can mask real subset effects. Virus-selected populations show stronger signals.

Rintatolimod (Ampligen): Regulatory Failure

Phase III ($n = 234$) showed marginal overall improvement in exercise tolerance [622]. Post-hoc analysis found 51.2% improvement in patients with 2–8 year disease duration [148]. FDA advisory committee voted 8–5 against approval, citing reliance on post-hoc subgroup analyses.

Lesson: Post-hoc subgroup analyses cannot rescue a trial; the disease-duration effect requires prospective confirmation.

27.9.4 Neurological and Cholinergic Interventions

Galantamine (Blacker et al., 2004)

The largest ME/CFS pharmacological trial ($n = 434$) tested galantamine at four doses across 35 centers [623]. No statistically significant differences on any outcome measure.

Lesson: Definitive null result rules out peripheral cholinergic enhancement. Cognitive impairment in ME/CFS is not cholinergic.

27.9.5 Antidepressant Interventions

Fluoxetine (Vercoulen et al., 1996)

Double-blind RCT ($n = 96$) enrolling both depressed and non-depressed CFS patients [624]. Fluoxetine showed no effect on any CFS dimension in either subgroup.

Lesson: CFS is not depression. Serotonergic modulation does not address core fatigue.

27.9.6 Endocrine Interventions

Hydrocortisone

Two RCTs tested hydrocortisone replacement based on mild hypocortisolism findings. McKenzie et al. ($n = 70$): modest improvement offset by adrenal suppression [625]. Cleare et al. ($n = 32$): 28% vs. 9% normalized fatigue scores, but adrenal suppression even at low doses [626].

Lesson: Mild hypocortisolism may be a *consequence* of ME/CFS rather than a cause, explaining why correction does not resolve symptoms.

27.9.7 Summary Table

27.9.8 Cross-Cutting Lessons

1. **ME/CFS is not one disease for treatment purposes.** Every trial enrolling unstratified patients produced null or weak results. Every trial pre-selecting by biomarker found stronger signals.
2. **Subjective outcomes in unblinded trials are unreliable.** Future trials must include objective measures: actigraphy, two-day CPET, employment status, or validated functional capacity assessments.
3. **Composite primary endpoints can mask real effects.** Selecting a single, clinically meaningful primary endpoint is preferable.
4. **Post-hoc subgroup analyses are hypothesis-generating, not confirmatory.** Adaptive trial designs can incorporate pre-specified biomarker stratification.

Table 27.1: Major negative, null, and inconclusive trials in ME/CFS.

Intervention	Design	<i>n</i>	Result	Key Lesson
Rituximab III [376]	DB-RCT	151	Negative	No benefit unstratified; subgroup may exist
PACE (GET/CBT) [115]	AB-RCT	641	Contested	No objective benefit; reanalysis ~7% [506]
Rintatolimod [622]	DB-RCT	234	Mixed	Marginal overall; subset signal at 2–8 yr
Galantamine [623]	DB-RCT	434	Negative	Definitive null; not cholinergic
Acyclovir [621]	DB-RCT	27	Negative	Wrong antiviral spectrum
Valganciclovir [149]	DB-RCT	30	Mixed	Primary negative; individual endpoints positive
IVIg [620]	DB-RCT	99	Negative	No benefit without immune pre-selection
Fluoxetine [624]	DB-RCT	96	Negative	CFS ≠ depression
Hydrocortisone [625]	DB-RCT	70	Negative	Adrenal suppression outweighs benefit

DB = double-blind; AB = assessor-blind; RCT = randomized controlled trial.

5. **The null hypothesis is informative.** Each negative trial narrows the hypothesis space: galantamine rules out cholinergic mechanisms; fluoxetine argues against serotonergic deficit; hydrocortisone reframes hypocortisolism as consequence not cause.

28 Mechanistic and Experimental Studies

28.1 The NIH Deep Phenotyping Study (Walitt et al. 2024)

The 2024 NIH Intramural Study, published in *Nature Communications*, represents the most comprehensive and expensive deep phenotyping study of post-infectious ME/CFS to date [5]. This landmark study merits detailed examination both for its substantial biological contributions and for the significant methodological controversies it generated.

28.1.1 Study Design and Methodology

Overview

- **Duration:** 8 years (launched 2016, published February 2024)
- **Cost:** Approximately \$8 million
- **Investigators:** 75+ NIH researchers across 15 institutes
- **Setting:** NIH Clinical Center inpatient evaluation over several days
- **Design:** Cross-sectional deep phenotyping study

Participants

- **PI-ME/CFS patients:** 17 (original target was 40; recruitment halted at 42% due to COVID-19 pandemic)
- **Healthy controls:** 21 (matched by age, sex, BMI)
- **Inclusion criteria:** Post-infectious onset ME/CFS (viral or bacterial trigger), illness duration <5 years, met rigorous diagnostic criteria
- **Critical exclusion:** Severely affected patients unable to travel to NIH

Comprehensive Assessment Battery

The study employed an unprecedented range of assessments:

Neurological and Brain Assessments

- Functional MRI (fMRI) during grip strength and effort tasks
- Transcranial magnetic stimulation
- Cognitive performance testing
- Effort-Expenditure for Rewards Task (EEfRT)

Autonomic Function Testing

- Heart rate variability measures (RMSSD, SDNN)
- Baroreflex cardiovascular function
- Chronotropic response assessment

Physical Performance

- Cardiopulmonary exercise testing (CPET)—single day protocol
- Grip strength testing (maximum and sustained)
- Motor performance evaluations

Tissue Sampling

- Muscle biopsies for gene expression analysis
- Skin biopsies
- Cerebrospinal fluid (CSF) analysis via lumbar puncture
- Comprehensive blood sampling

Advanced Omics Approaches

- Immune profiling: Flow cytometry of B cells and T cells
- Gene expression: PBMC and skeletal muscle transcriptomics
- Metabolomics: CSF and plasma metabolite profiling
- Microbiome analysis: Gut microbiota characterization
- Proteomics

Metabolic Chamber Study

- Multi-day assessment in controlled environment
- Energy consumption measurement
- Sleep pattern analysis
- Controlled diet

28.1.2 Key Biological Findings

The study documented multiple objective abnormalities (detailed in respective chapters):

Central Catecholamine Deficiency (Chapter 8)

- Abnormally low CSF levels of norepinephrine, dopamine, and DHPG (3,4-dihydroxyphenylglycol)
- Catecholamine levels correlated with grip strength, effort preference, and cognitive symptoms
- First direct CSF neurotransmitter measurements in ME/CFS

Immune Dysfunction (Chapter 7)

- Increased naïve B cells with decreased switched memory B cells
- Pattern consistent with chronic antigenic stimulation
- Elevated CD8+ T cell PD-1 expression (exhaustion marker)
- Sex-specific differences: males showed T cell/innate immunity changes; females showed B cell abnormalities

Autonomic Dysfunction (Chapter 10)

- Diminished heart rate variability at rest and during activity
- Impaired baroreflex-cardiovagal function
- Chronotropic incompetence during exercise

Cardiopulmonary Abnormalities

- Significantly reduced peak VO_2 compared to controls
- Reduced peak work capacity
- Lower ventilation during exercise
- Early anaerobic threshold onset

Neuroimaging Findings

- Reduced temporoparietal junction (TPJ) activity during motor tasks
- Abnormally sustained motor cortex activation despite declining force output
- No evidence of peripheral muscle fatigue on EMG

Grip Strength Pattern

A revealing finding: maximum grip strength showed no difference between patients and controls, but sustained grip strength was markedly reduced. The authors noted: “If deconditioning were the cause, we would expect maximum strength differences” [5]—arguing against simple deconditioning as explanation.

28.1.3 Methodological Limitations and Criticisms

Sample Size

The study achieved only 42% of its enrollment target (17 vs. planned 40 patients), limiting statistical power for subgroup analyses and reducing generalizability.

Selection Bias

The exclusion of severely affected patients (approximately 25% of the ME/CFS population who are homebound or bedbound [125]) means findings may not generalize to those most disabled by the illness and most in need of research attention.

Single-Day CPET Protocol

The study used single-day cardiopulmonary exercise testing rather than the gold-standard two-day protocol that documents post-exertional malaise. The two-day protocol consistently shows Day 2 VO_2 peak decline of approximately 13.8% and work capacity decline of approximately 12.5% in ME/CFS patients, while controls show stable or improved performance [1]. By using only single-day testing, the study failed to objectively document PEM, the defining feature of ME/CFS.

Post-Exertional Malaise Assessment

PEM is mentioned only three times in the entire paper despite being the hallmark symptom of ME/CFS. The study design did not systematically assess or document PEM.

28.1.4 The “Effort Preference” Controversy

The study’s most controversial element was its characterization of altered “effort preference” as “the defining motor behavior” of PI-ME/CFS.

The Claim

Walitt et al. proposed that fatigue in ME/CFS arises from dysfunction of integrative brain regions (particularly the TPJ) affecting how the brain calculates effort requirements. They defined effort preference as “how much effort a person subjectively wants to exert” and concluded this was distinct from physical fatigue or central fatigue.

The EEfRT Methodology Problem

The study used the Effort-Expenditure for Rewards Task (EEfRT), a psychiatric assessment tool designed to measure motivation for rewards in conditions like depression and schizophrenia. A critical requirement of the EEfRT, as stated by its developers, is that tasks must be easy enough for all participants to complete without fatigue—the tool is designed to measure motivation, not ability.

However, in the Walitt study:

- Controls completed 96–99% of hard trials successfully
- ME/CFS patients completed only 65% of hard trials
- Seven of 15 ME/CFS patients performed below any control participant
- SF-36 Physical Function scores: 28.7 for ME/CFS vs. 97.5 for controls

Academic Reanalysis

Kirvin-Quamme et al. (2025) published a formal reanalysis in *Frontiers in Psychology* [627]. Key findings:

- Positive correlation ($r_s = 0.38$, $p = 0.03$) between hard task completion rate and proportion of hard task choices—indicating an ability confound
- The hard task was simply too difficult for many ME/CFS patients to complete, regardless of preference
- Data support interpretation that patients were “unable” rather than “unwilling”

Published Critique in Nature Communications

Davenport et al. published a formal commentary in *Nature Communications* [628] stating that the effort preference interpretation “risks reinforcing skepticism about the serious biological nature of [ME] and its hallmark of post-exertional malaise (PEM), as well as its potential misclassification as a mental health condition.”

Patient and Expert Community Response

The “effort preference” framing generated significant criticism:

- ME/CFS experts Drs. Lucinda Bateman and Brayden Yellman expressed being “particularly dismayed by use of the term ‘effort preference’ as an explanation for the origin of fatigue”
- Multiple experts called for retraction or correction of the effort preference claims
- Patient advocates noted the framing echoed problematic language from the PACE trial

NIH Clarification

NIH subsequently clarified that “preference” referred to “subconscious or unconscious or pre-conscious calculations by the brain” rather than conscious choice. Critics responded that if the intended meaning was unconscious brain dysfunction, the word “preference” was misleading and potentially harmful.

Limitation 1: NIH Deep Phenotyping Study: Small Sample and Contested Interpretation

The Walitt et al. 2024 study is the most comprehensive deep-phenotyping study of ME/CFS to date, but its conclusions are constrained by design limitations:

- The study enrolled only 17 post-infectious ME/CFS patients against an original target of 40; recruitment was halted due to COVID-19. At n=17 vs. 21 controls, many comparisons are underpowered and findings may not generalize.
- Severe and very severe patients were excluded because they could not travel to the NIH Clinical Center—the population with the most extreme pathophysiology is systematically absent.
- The use of “effort preference” to describe grip strength findings generated legitimate scientific controversy; whether the framing reflects unconscious neural computation (as the authors clarified) or introduces psychologizing bias into objective data remains unresolved.
- Biological findings (CSF catecholamine deficiency, B cell shifts, autonomic dysfunction) are unreplicated by independent groups and derive from a single cross-sectional assessment.

28.1.5 Interpretation and Context

Despite the controversy over interpretation, the Walitt study’s biological findings—catecholamine deficiency, B cell population shifts, autonomic dysfunction, cardiopulmonary impairment—represent valuable contributions to ME/CFS research. The challenge lies in separating the objective biological data from the contested psychological framing.

? Open Question 1: Separating Data from Interpretation

The NIH deep phenotyping study illustrates a broader challenge in ME/CFS research: how to extract valid biological findings from studies whose interpretive frameworks may be problematic. The catecholamine, immune, and autonomic data stand on their own merit regardless of how they are contextualized. Future research should build on these biological findings while employing more appropriate methodologies for assessing effort and function in patients with energy-limiting illness.

28.2 Related Studies from the NIH Cohort

28.2.1 WASF3 and Mitochondrial Dysfunction (Wang et al. 2023)

Using muscle biopsies from the same NIH intramural cohort, Wang et al. identified a specific molecular mechanism linking cellular stress to exercise intolerance [2].

Key Findings

- **Elevated WASF3 protein:** ME/CFS muscle biopsies showed increased WASF3 (Wiskott-Aldrich syndrome protein family member 3)
- **ER stress activation:** Endoplasmic reticulum stress was aberrantly increased
- **Mitochondrial localization:** WASF3 localizes to mitochondria and disrupts respiratory chain complex assembly
- **Functional consequence:** Decreased oxygen consumption and exercise endurance

Proposed Mechanism

1. Cellular stress activates the unfolded protein response (ER stress)
2. ER stress induces WASF3 expression
3. WASF3 translocates to mitochondria
4. WASF3 disrupts respiratory chain complex IV assembly
5. Impaired oxidative phosphorylation reduces exercise capacity

Therapeutic Implication

Pharmacologic inhibition of ER stress improved mitochondrial function in patient-derived cells, suggesting a potential therapeutic target.

28.2.2 T Cell Exhaustion (lu et al. 2024)

A separate study examining immune cells from ME/CFS patients found extensive evidence of CD8+ T cell exhaustion [119].

Key Findings

- **Elevated PD-1 expression:** Exhaustion marker on CD8+ T cells
- **Transcriptional reprogramming:** Gene expression patterns consistent with chronic antigenic stimulation
- **Epigenetic changes:** Persistent modifications indicating long-term immune activation
- **Similar to chronic infections:** Pattern resembles exhaustion seen in chronic viral infections and cancer

Implications

T cell exhaustion provides independent confirmation of chronic immune activation in ME/CFS and suggests that immune checkpoint therapies or other approaches to reverse exhaustion might have therapeutic potential.

28.3 Exercise Physiology Studies

28.3.1 Cardiopulmonary Exercise Testing (CPET)

The two-day CPET protocol is the most important methodological contribution to ME/CFS exercise physiology research. By repeating maximal exercise testing on consecutive days, it objectively documents post-exertional malaise—the hallmark feature of ME/CFS that cannot be captured by single-day testing.

Keller et al. (2024) published the largest rigorous two-day CPET study to date (n=84 ME/CFS, n=71 sedentary controls) [1]. Key findings include:

- **Day 2 performance decline:** ME/CFS patients show significant reductions in VO_2peak (−5–8%), peak work rate, and ventilatory anaerobic threshold on Day 2. Healthy controls and sedentary individuals show stable or improved Day 2 performance
- **Worsening impairment classification:** The proportion of patients classified as severely impaired nearly doubles from Day 1 to Day 2 (14% to 27%)
- **Prolonged recovery:** ME/CFS patients require 13+ days for physiological recovery versus approximately 2 days for controls
- **Earlier anaerobic threshold:** ME/CFS patients reach anaerobic threshold at lower absolute and relative workloads, indicating premature shift to anaerobic metabolism
- **Independence from baseline fitness:** Day 2 decrements persist when matched for baseline aerobic capacity, ruling out deconditioning as the sole explanation

Lim et al. (2020) performed an earlier systematic review confirming significant workload decline at ventilatory threshold in ME/CFS patients (−14.6 W) while controls showed improvement (+6.5 W, $p=0.01$) [79].

The Workwell Foundation has been instrumental in standardizing CPET methodology for ME/CFS and advocating for two-day testing as the objective diagnostic standard. Their work has established that ME/CFS exercise intolerance is a measurable, reproducible physiological abnormality—not a subjective complaint.

28.3.2 Muscle Studies

Skeletal muscle studies have revealed both functional and structural abnormalities in ME/CFS:

- **Lactate accumulation:** ME/CFS patients show earlier and greater lactate accumulation during submaximal exercise, consistent with premature reliance on anaerobic glycolysis. This is compatible with impaired oxidative phosphorylation (Chapter 6)
- **Oxidative capacity:** Muscle biopsy studies have reported reduced mitochondrial enzyme activity and decreased oxidative capacity in some ME/CFS cohorts, though results are heterogeneous across studies
- **WASF3 and ER stress:** Wang et al. (2023) identified elevated WASF3 protein in ME/CFS muscle biopsies, linking endoplasmic reticulum stress to mitochondrial respiratory supercomplex disruption [2]. This provides a specific molecular mechanism connecting cellular stress to exercise intolerance (Section 28.1)
- **Fiber type distribution:** Some studies report a shift from oxidative Type I fibers to glycolytic Type II fibers, consistent with chronic disuse but also potentially reflecting metabolic reprogramming
- **Small fiber neuropathy:** Reduced intraepidermal nerve fiber density has been documented in ME/CFS muscle and skin biopsies, providing a structural basis for pain and sensory symptoms

28.4 Cellular and Molecular Studies

28.4.1 Cell Culture Studies

In vitro studies using patient-derived cells have provided critical mechanistic insights by isolating cellular defects from whole-body confounders:

- **PBMC bioenergetics:** Seahorse extracellular flux analyzer studies consistently show reduced mitochondrial respiration in peripheral blood mononuclear cells (PBMCs) from ME/CFS patients compared to healthy controls [170, 326]. Basal and maximal respiration, spare respiratory capacity, and ATP-linked respiration are all reduced. Critically, these deficits persist in cultured cells removed from the patient's systemic environment, suggesting a cell-intrinsic metabolic defect rather than a consequence of circulating factors alone
- **Metabolic flux analysis:** Fluge et al. (2016) demonstrated that ME/CFS patient serum inhibits pyruvate dehydrogenase (PDH) activity in healthy muscle cells, suggesting a circulating inhibitory factor. PDH inhibition would impair the conversion of pyruvate to acetyl-CoA, forcing reliance on anaerobic glycolysis even in the presence of adequate oxygen—consistent with the “metabolic trap” hypothesis
- **NK cell function:** Reduced NK cell cytotoxicity is one of the most replicated findings in ME/CFS [236]. Cell culture studies demonstrate both reduced killing capacity and impaired calcium signaling through TRPM3 ion channels [536]
- **T cell stimulation:** Patient-derived T cells show altered metabolic responses to stimulation, with reduced glycolytic reserve and impaired ability to upregulate oxidative phosphorylation upon activation—consistent with the T cell exhaustion phenotype documented *in vivo*

28.4.2 Animal Models

Animal models for ME/CFS have been historically limited by the difficulty of replicating a complex, multi-system human disease in laboratory animals. Most existing models target individual pathways rather than the full syndrome.

Existing Approaches and Limitations

Rodent models using viral infection (murine gammaherpesvirus, poly(I:C) injection), immune activation (LPS administration), or forced exercise have reproduced individual ME/CFS features—fatigue-like behavior, neuroinflammation, or immune activation—but none recapitulates the defining feature of post-exertional malaise with delayed onset and prolonged recovery. Key limitations include:

- **Artificial induction:** Laboratory infection or chemical challenge does not replicate the natural triggering events (viral infection in a genetically susceptible host) that produce human ME/CFS.
- **PEM measurement:** No validated method exists for quantifying PEM in rodents. Activity monitoring can detect reduced voluntary wheel running, but distinguishing pathological exercise intolerance from sickness behavior or learned avoidance is methodologically difficult.
- **Chronicity:** Most models produce acute effects lasting days to weeks, whereas ME/CFS is defined by persistence over months to years.
- **Multi-system involvement:** Models targeting a single pathway (e.g., neuroinflammation) cannot reproduce the simultaneous immune, metabolic, autonomic, and neurological dysfunction that characterizes ME/CFS.

Feline Infectious Peritonitis as a Comparative Coronavirus Model

The Kol et al. (2026) discovery that feline infectious peritonitis virus (FIPV) persists in B and T lymphocytes after antiviral treatment and clinical recovery [171] establishes FIP as a uniquely valuable comparative model for post-viral immune dysfunction. Unlike artificial rodent models, FIP is a naturally occurring coronavirus disease with several features directly relevant to post-COVID ME/CFS:

- **Natural coronavirus infection:** FIPV is a feline coronavirus that produces systemic immune dysfunction without artificial induction.
- **Immune cell tropism:** FIPV replicates in macrophages, B cells, and T cells—paralleling the suspected SARS-CoV-2 tropism for immune cells in long COVID.
- **Post-treatment viral persistence:** Viral RNA persists in lymphocytes after antiviral treatment (GS-441524) and resolution of clinical signs, modeling the hypothesized viral reservoir in human ME/CFS (Hypothesis 7.2.1).

- **Accessible lymphoid tissue:** Unlike human studies, feline lymph node tissue can be directly examined, enabling mechanistic studies of viral-immune cell interactions impossible in human subjects.
- **Treatment response monitoring:** Cats treated with GS-441524 (a nucleoside analog related to remdesivir) can be monitored longitudinally for immune recovery, relapse, and persistent dysfunction.

Proposed Research Directions.

1. **Longitudinal immune profiling:** Track immune cell phenotype, function, and viral RNA persistence in FIP-recovered cats over 12–24 months post-treatment. Determine whether persistent lymphocyte infection correlates with chronic immune dysfunction analogous to ME/CFS T cell exhaustion.
2. **Activity monitoring:** Implement accelerometry-based activity tracking in FIP-recovered cats to determine whether a subgroup develops persistent activity reduction or post-exertional deterioration analogous to PEM.
3. **Metabolic profiling:** Compare immune cell bioenergetics (Seahorse analysis) between FIP-recovered cats with persistent viral RNA versus those who clear virus completely, testing whether viral persistence drives immune cell metabolic dysfunction (as proposed in Hypothesis 6.3).
4. **Intervention testing:** Use the FIP model to test combined antiviral + metabolic support strategies before human trials, determining whether metabolic supplementation enhances viral clearance from lymphocytes.

? Open Question 2: Viral RNA in Circulating Lymphocytes as Diagnostic Biomarker

If the lymphocyte reservoir hypothesis (Hypothesis 7.2.1) is correct, ultra-sensitive single-cell methods (10x Genomics single-cell RNA-seq, MERFISH spatial transcriptomics) applied to peripheral blood mononuclear cells from ME/CFS patients should detect low-level viral transcripts in a small fraction of lymphocytes—below the sensitivity of bulk PCR but detectable at single-cell resolution.

Key unknowns:

- What is the minimum fraction of virus-positive lymphocytes required to sustain chronic immune activation? (Even 0.01% of circulating lymphocytes represents millions of cells.)
- Does the viral species matter, or would any persistent virus in lymphocytes produce the ME/CFS phenotype?
- Can this biomarker distinguish post-viral ME/CFS from non-viral-onset ME/CFS, potentially defining a treatable subgroup?
- What is the sensitivity threshold required—and do current single-cell platforms achieve it?

If validated, this would represent the first objective diagnostic biomarker for post-viral

ME/CFS and would immediately stratify patients for antiviral therapy trials.

? Open Question 3: Temporal Mapping of the Deconditioning-to-Pathology Transition

When does secondary deconditioning in ME/CFS become irreversible tissue pathology? The Ratchet Model (Speculation 5) predicts cumulative irreversible damage, but the transition point is unknown. This question has direct clinical implications: if there is a window during which deconditioning is still reversible, early targeted rehabilitation (within the energy envelope) could prevent permanent functional loss.

A study design to address this:

- Serial muscle biopsies (at diagnosis, 6 months, 12 months, 24 months) in newly diagnosed ME/CFS patients managed with strict pacing (no GET)
- Paired two-day CPET at each time point to track exercise capacity trajectory
- Comparison cohort: long-duration (>5 year) ME/CFS patients to establish “end-point” tissue changes
- Outcome: distinguish deconditioning-reversible changes (muscle fiber type shifts, capillary density reduction) from pathology-irreversible changes (mitochondrial DNA damage, fibrosis, small fiber denervation)

This would be the first study to empirically separate the deconditioning component from the primary pathological component in ME/CFS exercise intolerance, directly testing a core assumption of both the deconditioning and the ratchet models.

28.5 Imaging Studies

28.5.1 Brain Imaging

Neuroimaging studies have documented structural, functional, and neuroinflammatory abnormalities in ME/CFS, though findings are heterogeneous across studies and methodologies:

Structural MRI: Several studies report reduced white matter volume, cortical thinning, and altered white matter integrity (reduced fractional anisotropy on diffusion tensor imaging) in ME/CFS patients. Barnden et al. (2011) found white matter volume reductions correlating with fatigue severity. However, findings are inconsistent—some studies report no structural differences—possibly reflecting patient heterogeneity and small sample sizes.

Functional MRI: The NIH deep phenotyping study (Section 28.1) found reduced temporoparietal junction (TPJ) activity during motor tasks and abnormally sustained motor cortex activation despite declining force output [5]. Other fMRI studies have reported altered connectivity in the default mode network, salience network, and executive control network—patterns overlapping with those seen in chronic pain and neuroinflammatory conditions.

PET imaging: Nakatomi et al. (2014) used ¹¹C-PK11195 PET to demonstrate widespread neuroinflammation (activated microglia) in the cingulate cortex, hippocampus, amygdala,

thalamus, midbrain, and pons of ME/CFS patients [73]. The degree of neuroinflammation correlated with cognitive impairment severity. Subsequent studies using more selective TSPO ligands have confirmed elevated neuroinflammatory markers, though the specificity of TSPO PET for microglial activation (versus astrocyte activation) remains debated.

SPECT: Single-photon emission computed tomography studies have consistently shown reduced cerebral blood flow in ME/CFS, particularly in the brainstem, frontal, and temporal regions. Reduced perfusion correlates with cognitive symptoms and orthostatic intolerance. SPECT evidence predates and is consistent with the cerebral hypoperfusion documented by more recent techniques.

28.5.2 Cardiac Imaging

Cardiovascular imaging has revealed structural and functional cardiac abnormalities in ME/CFS that are distinct from deconditioning:

- **Reduced blood volume:** Studies using dual-label red blood cell volume testing have documented significant blood volume reductions in ME/CFS patients (average 10–15% below predicted). This hypovolemia contributes to reduced cardiac preload and orthostatic intolerance
- **Cardiac MRI:** Ponikowski et al. and subsequent studies found reduced left ventricular mass, reduced end-diastolic volume, and lower stroke volume in ME/CFS patients compared to healthy controls and deconditioned individuals. These reductions are disproportionate to the degree of inactivity, suggesting a disease-specific cardiac effect rather than pure deconditioning
- **Invasive hemodynamic testing:** Systrom et al. used invasive cardiopulmonary exercise testing with right heart catheterization to demonstrate reduced cardiac filling pressures (preload failure) and impaired peripheral oxygen extraction in ME/CFS, providing direct evidence of both central and peripheral circulatory dysfunction

28.6 Immunological Studies

Immunological research has provided some of the most consistent and replicated findings in ME/CFS:

NK cell dysfunction: Reduced NK cell cytotoxic activity is the most replicated immunological finding in ME/CFS, confirmed across multiple independent laboratories and cohorts [236]. The defect involves both reduced numbers of cytotoxic NK cell subsets (CD56^{dim}CD16⁺) and impaired per-cell killing capacity, linked to defective calcium signaling through TRPM3 ion channels [536].

T cell abnormalities: Multiple studies document altered CD4⁺/CD8⁺ ratios, increased expression of activation and exhaustion markers (CD38, HLA-DR, PD-1, Tim-3), and reduced proliferative capacity. The NIH deep phenotyping study found elevated CD8⁺ T cell PD-1 expression consistent with chronic antigenic stimulation [5]. Iu et al. (2024) provided transcriptomic and epigenetic evidence for extensive T cell exhaustion [119].

Cytokine profiles: Hornig et al. (2015) identified distinct cytokine signatures that vary with disease duration: elevated pro-inflammatory cytokines (IL-6, TNF- α , IL-1 β) in early disease (≤ 3 years) transitioning to a mixed or suppressed profile in chronic disease [212]. Montoya et al. (2017) confirmed cytokine–severity correlations [213]. However, cytokine findings remain inconsistent across laboratories, partly due to methodological differences in sample handling, assay platforms, and patient selection.

Autoantibody discovery: Loebel et al. (2016) and subsequent studies identified elevated autoantibodies against G protein-coupled receptors (GPCRs), including β_2 -adrenergic, muscarinic M3/M4, and angiotensin II AT1 receptors [71, 208, 72]. These autoantibodies provide a mechanistic link between immune dysfunction and autonomic symptoms. Their functional significance is supported by the clinical response to immunoadsorption in some patients.

Viral persistence: Evidence for persistent viral infection in ME/CFS includes elevated antibody titers against EBV, HHV-6, and CMV in subgroups of patients, detection of viral DNA/RNA in tissue samples, and the FIP model of post-viral lymphocyte persistence [171]. Whether viral persistence is causative or an epiphenomenon of immune dysfunction remains an active research question.

28.7 Omics Studies

High-throughput omics technologies have transformed ME/CFS research by enabling unbiased, hypothesis-free discovery of disease-associated molecular signatures.

28.7.1 Genomics

Genetic studies suggest a modest heritable component to ME/CFS susceptibility:

- **Twin studies:** Concordance rates are higher in monozygotic than dizygotic twins, suggesting genetic contribution, though shared environment cannot be fully excluded
- **GWAS:** No genome-wide significant loci have been consistently replicated across studies, likely reflecting polygenic architecture, phenotypic heterogeneity, and insufficient sample sizes. The UK ME/CFS Biobank and DecodeME (n>20,000) represent the largest genetic studies to date and may have sufficient power to detect common variants of small effect
- **HLA associations:** Several studies report associations with specific HLA alleles (particularly HLA-DRB1), consistent with an autoimmune component in a subgroup. Associations with immune-regulatory genes (cytokine promoter polymorphisms, KIR genes) have also been reported but not consistently replicated
- **Rare variants:** Whole-exome sequencing studies are in early stages. Rare variants in mitochondrial DNA, ion channel genes, and complement pathway genes have been reported in individual families but require validation in larger cohorts

28.7.2 Transcriptomics

Gene expression profiling has identified reproducible transcriptomic signatures:

- **Immune gene dysregulation:** Multiple studies report altered expression of immune-related genes, including upregulation of interferon-stimulated genes, downregulation of NK cell-associated transcripts, and dysregulation of NF- κ B signaling pathways
- **Metabolic gene changes:** Downregulation of genes involved in oxidative phosphorylation and upregulation of glycolysis-related transcripts are consistent with the metabolic shift documented by functional studies
- **Exercise-induced changes:** Transcriptomic profiling before and after exercise challenge reveals gene expression changes in ME/CFS patients that differ from healthy controls, including altered stress response and immune activation patterns persisting 24–48 hours post-exercise
- **Single-cell transcriptomics:** Emerging single-cell RNA-seq studies are resolving cell-type-specific expression changes that are averaged out in bulk profiling, revealing that immune cell subsets carry distinct transcriptomic signatures in ME/CFS

28.7.3 Proteomics

Proteomic studies complement transcriptomics by capturing post-translational regulation and secreted factors:

- **Plasma proteomics:** Studies have identified altered levels of complement components, coagulation factors, acute-phase proteins, and extracellular matrix remodeling enzymes in ME/CFS plasma
- **CSF proteomics:** The NIH study and others have profiled cerebrospinal fluid proteins, identifying alterations in neurotransmitter metabolism, neuroinflammatory markers, and blood–brain barrier integrity proteins
- **Post-translational modifications:** Oxidative protein modifications (carbonylation, nitrosylation) are elevated in ME/CFS, consistent with oxidative and nitrosative stress. Altered protein phosphorylation patterns in immune cells reflect dysregulated signaling cascades

28.7.4 Metabolomics

Metabolomics has produced some of the most striking and replicated findings in ME/CFS:

- **Naviaux et al. (2016):** Identified a hypometabolic signature in ME/CFS, with widespread reductions in amino acids, lipids, and nucleotides consistent with a dauer-like state of metabolic hibernation [589]. The metabolic profile discriminated ME/CFS patients from controls with >90% accuracy

- **Tryptophan–kynurenine pathway:** Multiple studies document a shift from serotonin synthesis toward kynurenine production, driven by inflammation-induced indoleamine 2,3-dioxygenase (IDO) activation. This has implications for mood, pain, and cognitive function (Chapter 8)
- **Amino acid depletion:** Reduced levels of branched-chain amino acids, tryptophan, and other essential amino acids are consistently reported, suggesting impaired anabolic metabolism
- **Lipid abnormalities:** Altered phospholipid profiles, reduced plasmalogens, and abnormal sphingolipid metabolism have been documented, consistent with membrane dysfunction and impaired mitochondrial lipid metabolism

28.7.5 Lipidomics

Dedicated lipidomic profiling has revealed ME/CFS-specific lipid signatures:

- **Phospholipid remodeling:** Altered phosphatidylcholine-to-phosphatidylethanolamine ratios suggest membrane instability. Reduced plasmalogens—antioxidant ether lipids concentrated in mitochondrial and cell membranes—may contribute to both oxidative vulnerability and mitochondrial dysfunction
- **Eicosanoid imbalance:** Altered prostaglandin and leukotriene profiles reflect immune dysregulation. Elevated prostaglandin E2 may contribute to pain sensitization and vasodilation
- **Sphingolipid changes:** Altered ceramide and sphingomyelin levels have been reported, with potential implications for immune cell signaling and apoptosis

28.7.6 Microbiomics

Gut microbiome studies have documented consistent alterations in ME/CFS (see also Chapter 11):

- **16S rRNA sequencing:** Reduced microbial diversity and altered community composition, including reduced butyrate-producing bacteria (*Faecalibacterium prausnitzii*, *Roseburia*) and increased pro-inflammatory species. These changes are consistent with the gut barrier dysfunction and immune activation documented in ME/CFS
- **Shotgun metagenomics:** Functional profiling reveals altered metabolic capacity of the ME/CFS microbiome, including reduced short-chain fatty acid production and altered tryptophan metabolism—linking gut microbial changes to the systemic metabolic abnormalities detected by metabolomics
- **Gut–brain axis:** Microbial metabolites (short-chain fatty acids, tryptophan derivatives) directly influence CNS function through vagal signaling and systemic circulation, providing a mechanistic link between gut dysbiosis and neurological symptoms

28.8 Integrative Multi-Omics Studies

The convergence of multiple omics layers on consistent pathophysiological themes strengthens confidence in ME/CFS disease mechanisms:

Multi-omics integration studies have begun combining genomic, transcriptomic, proteomic, metabolomic, and microbiomic data from the same patients. Heng et al. (2025) demonstrated that multi-omics integration outperforms any single omics layer for classifying ME/CFS patients and identifying mechanistic subtypes. Key integrative findings include:

- **Metabolic-immune convergence:** Transcriptomic immune gene dysregulation and metabolomic hypometabolic signatures co-localize to the same patients, supporting the energy-immune vicious cycle model (Chapter 13)
- **Gut-systemic axis:** Microbiomic changes in butyrate-producing bacteria correlate with metabolomic tryptophan depletion and transcriptomic immune activation, confirming the gut-immune-brain axis as a functional unit in ME/CFS
- **Subgroup identification:** Unsupervised clustering of multi-omics data identifies at least 2–4 molecular subtypes that may correspond to clinically distinct patient populations—a critical step toward biomarker-stratified clinical trials
- **Network approaches:** Systems biology network analyses integrating protein–protein interactions, metabolic pathways, and gene regulatory networks have identified hub molecules (e.g., IDO1, HIF-1 α , NF- κ B) that connect apparently disparate disease features into coherent mechanistic models

⚠ Limitation 2: Multi-Omics Reproducibility

Most multi-omics studies in ME/CFS are single-cohort analyses with small sample sizes ($n < 50$). Validation in independent cohorts is essential before molecular subtypes can be considered clinically actionable. Standardization of sample collection, processing, and analytical pipelines across sites remains a major challenge. The ME/CFS Biobank initiatives (UK, US, European) are addressing these issues through harmonized protocols.

29 Epidemiological and Outcomes Research

29.1 Prevalence and Incidence Studies

Estimating ME/CFS prevalence is complicated by the absence of a diagnostic biomarker, the use of multiple case definitions of varying stringency, and the high rate of undiagnosed cases. Prevalence estimates vary by an order of magnitude depending on methodology and criteria applied.

29.1.1 Population-Based Estimates

The most cited prevalence estimates range from 0.1% to 0.9% of the general adult population:

- **Fukuda (1994) criteria:** Broader definition; population-based studies using these criteria report prevalence of 0.2–0.4% (approximately 800,000–1,600,000 adults in the US)
- **Canadian Consensus Criteria (2003):** Stricter definition requiring PEM; prevalence approximately 0.1–0.2%
- **Institute of Medicine (2015):** Estimated 836,000–2,500,000 Americans affected, with 84–91% undiagnosed
- **Community surveillance studies:** Jason et al. (1999) conducted door-to-door screening in Chicago, identifying a prevalence of 0.42% using the Fukuda criteria—substantially higher than clinic-based estimates, confirming widespread underdiagnosis

29.1.2 Geographic and Demographic Variation

Prevalence varies across populations, though much variation may reflect diagnostic practices rather than true disease frequency:

- **Sex ratio:** Females outnumber males approximately 3–4:1 across most studies [318, 35]. This ratio is consistent across cultures and diagnostic criteria, suggesting a biological basis (hormonal, immune, or genetic)
- **Age distribution:** Peak onset in the 30–49 age range, though ME/CFS occurs at all ages including children and adolescents. Pediatric prevalence is estimated at 0.1–0.5% [35]
- **Ethnic and socioeconomic factors:** Community-based studies (as opposed to clinic-based) show similar or higher prevalence in minority and lower-income populations [318], contradicting the outdated characterization of ME/CFS as a disease of affluent white women. Underdiagnosis in minority populations likely reflects healthcare access disparities rather than lower disease incidence

29.1.3 Post-COVID Surge

The COVID-19 pandemic has significantly increased ME/CFS incidence. An estimated 51% of long COVID patients meeting PEM criteria satisfy ME/CFS diagnostic criteria [370]. Given the hundreds of millions of SARS-CoV-2 infections worldwide, the projected increase in ME/CFS prevalence is substantial—potentially doubling or tripling the pre-pandemic patient population. This surge has accelerated research funding and clinical attention but has also strained the already limited specialist infrastructure.

29.2 Risk Factor Studies

29.2.1 Genetic Risk Factors

Evidence supports a genetic contribution to ME/CFS susceptibility, though the genetic architecture remains poorly characterized:

- **Family studies:** First-degree relatives of ME/CFS patients have a higher risk of developing the condition [42]. Family clustering is consistently reported, though shared environmental exposures cannot be excluded
- **Twin studies:** Monozygotic twins show higher concordance than dizygotic twins, with heritability estimates of 30–50% [42]. This suggests a moderate genetic contribution with substantial environmental influence
- **Genetic associations:** HLA alleles (particularly HLA-DRB1), cytokine gene polymorphisms, and immune-regulatory gene variants have been associated with ME/CFS in candidate gene studies, but none has been consistently replicated [42]. Genome-wide association studies (DecodeME, UK ME/CFS Biobank) are ongoing and may identify robust common variants (Section 28.7)

29.2.2 Environmental Risk Factors

Environmental triggers interact with genetic susceptibility to precipitate ME/CFS:

- **Infectious triggers:** Viral infection is the most common precipitant, identified in 60–80% of cases. Epstein-Barr virus, enteroviruses, HHV-6, and SARS-CoV-2 are the most frequently implicated pathogens. The Dubbo Infection Outcomes Study documented ME/CFS-like illness in 11% of patients following acute EBV, Ross River virus, or *Coxiella burnetii* infection, with severity of acute illness predicting chronic sequelae
- **Toxic exposures:** Organophosphate pesticides, solvents, mold/mycotoxin exposure, and heavy metals have been reported as triggers in case series. Systematic epidemiological evidence is limited
- **Physical trauma:** Surgery, childbirth, and physical injury occasionally precede ME/CFS onset, possibly through immune activation or HPA axis disruption

- **Psychological stress:** Severe or prolonged stress is reported as a precipitant in some cases, though its role is controversial and must be distinguished from the psychogenic model that attributes ME/CFS to stress itself
- **Vaccination:** Rare cases of ME/CFS onset following vaccination have been reported. Population-level studies do not show elevated risk, and the benefit of vaccination (particularly for infection prevention in existing ME/CFS patients) far outweighs any theoretical risk

29.2.3 Demographic Risk Factors

Demographic factors associated with ME/CFS risk include:

- **Sex:** Female predominance (3–4:1) is robust across studies. Potential explanations include hormonal influences on immune function, higher autoimmune disease susceptibility in females, and possible diagnostic bias (males may be less likely to seek care or receive the diagnosis)
- **Age:** Peak onset in the 30s–40s, but can occur at any age. Pediatric onset may carry a better prognosis than adult onset [78]
- **Socioeconomic status:** Community-based studies show no association with higher socioeconomic status. The historical perception of ME/CFS as a “yuppie flu” reflected clinic-based sampling bias (wealthier patients had better access to specialists who could diagnose the condition)

29.3 Natural History Studies

Understanding the natural history of ME/CFS is critical for prognosis, treatment planning, and patient counseling. However, few long-term prospective studies exist, and most evidence comes from retrospective cohorts with significant loss to follow-up.

29.3.1 Recovery Rates

Recovery from ME/CFS is possible but uncommon:

- **Full recovery:** Estimated at 5–10% across studies, though definitions of “recovery” vary. Strict criteria (return to pre-illness function) yield lower estimates than subjective improvement
- **Significant improvement:** An additional 20–30% of patients report meaningful improvement over time, though most remain below pre-illness function
- **Stable illness:** The majority of patients (40–60%) follow a fluctuating but relatively stable course, with periods of relative improvement and relapse
- **Progressive decline:** An estimated 10–20% of patients experience progressive worsening, particularly those who experience repeated infections, undergo inappropriate exercise therapy, or lack adequate pacing support

- **Pediatric prognosis:** Children and adolescents may have better prognosis than adults, with recovery rates estimated at 20–50% [504, 78], though these estimates are based on limited follow-up periods

29.3.2 Predictors of Outcome

Factors associated with better or worse prognosis include:

- **Better prognosis:** Younger age at onset, shorter illness duration at diagnosis, less severe initial presentation, identifiable infectious trigger, absence of psychiatric comorbidity, early adoption of pacing, and access to knowledgeable medical care
- **Worse prognosis:** Older age at onset, longer duration before diagnosis, severe initial presentation, multiple comorbidities (POTS, MCAS, fibromyalgia), exposure to graded exercise therapy, repeated infections during illness, and delayed or absent diagnosis
- **Illness duration:** Most recovery occurs within the first 2–5 years. Patients who remain ill beyond 5 years are unlikely to achieve full recovery, though improvement remains possible

29.4 Quality of Life and Disability Studies

ME/CFS produces one of the lowest quality-of-life scores of any chronic disease—a finding that is remarkably consistent across studies and measurement instruments.

29.4.1 SF-36 Findings

The Short Form 36 (SF-36) health survey is the most widely used quality-of-life instrument in ME/CFS research. ME/CFS patients consistently score lower than patients with congestive heart failure, type 2 diabetes, multiple sclerosis, end-stage renal disease, and most cancers on the physical function subscale. Typical SF-36 physical function scores in ME/CFS range from 20–35 (out of 100), compared to 50–70 for most other chronic diseases and 85–95 for healthy controls. Social function and role-physical subscales are similarly devastated.

29.4.2 Comparison to Other Chronic Diseases

The functional impairment of ME/CFS is frequently underestimated by clinicians unfamiliar with the disease:

- **Functional capacity:** Severely affected ME/CFS patients (approximately 25% of the population [125]) are housebound or bedbound, requiring assistance with basic activities of daily living. This level of disability is comparable to advanced multiple sclerosis or late-stage heart failure

- **Cognitive impairment:** Processing speed, attention, and working memory deficits in ME/CFS are comparable to those seen in mild traumatic brain injury or early-stage dementia
- **Pain burden:** Pain severity in ME/CFS is comparable to fibromyalgia and rheumatoid arthritis
- **Overall quality of life:** Multiple studies using different instruments consistently rank ME/CFS among the most disabling chronic conditions, yet it receives a fraction of the research funding allocated to diseases with comparable burden

29.4.3 Economic Burden

The economic impact of ME/CFS is substantial and largely borne by patients and their families:

- **Direct medical costs:** Estimated at \$9,000–14,000 per patient per year in the US, driven by specialist consultations, medications, and diagnostic testing [48]
- **Lost productivity:** The dominant economic cost. With unemployment rates of 35–69% and reduced productivity among those who remain employed, annual US economic losses are estimated at \$17–24 billion [50]
- **Informal caregiving:** Family caregivers provide substantial unpaid care, with estimated value exceeding direct medical costs. Caregiver burden is associated with reduced caregiver quality of life, lost caregiver productivity, and increased caregiver healthcare utilization
- **Cost-per-DALY:** ME/CFS ranks among the most cost-effective diseases to research given the large gap between disease burden and research investment

29.4.4 Impact on Caregivers

The burden on caregivers—typically spouses, parents, or adult children—is severe and understudied:

- **Physical burden:** Assisting with personal care, household tasks, and medical management for patients who cannot perform these independently
- **Psychological burden:** Witnessing a loved one's suffering and decline, managing their own grief and frustration, and coping with the social isolation that accompanies caregiving for a housebound patient
- **Financial burden:** Loss of the patient's income, potential reduction in the caregiver's working hours, and out-of-pocket medical expenses not covered by insurance
- **Social isolation:** Caregivers often become socially isolated alongside the patient, losing their own social connections and recreational activities
- **Relationship strain:** The transformation from partner to caregiver fundamentally alters the relationship dynamic and can create resentment, guilt, and grief on both sides

29.5 Mortality Studies

Understanding mortality patterns in ME/CFS is essential for both clinical practice and actuarial assessment (life insurance underwriting). While early concerns suggested potentially elevated mortality, large population-based cohort studies have provided more nuanced evidence. This section synthesizes findings from registry studies, clinical cohorts, and memorial record analyses.

Limitation 1: ME/CFS Mortality Data: Memorial Records vs. Registry Studies

Claims about ME/CFS mortality are subject to a fundamental evidence gap. Memorial record studies report dramatically reduced life expectancy (mean age at death 52–56 years), but these studies capture only deaths reported to patient organisations, inherently overrepresenting severe and fatal cases. Population-based registry studies (Roberts et al. 2016, Smith et al. 2006) find no elevated all-cause mortality. Neither design answers the critical question: does ME/CFS *cause* premature death, or do premature deaths from other causes preferentially enter memorial records? Until prospective cohort studies with ≥ 20 -year follow-up and severity stratification are conducted, the true mortality risk of ME/CFS cannot be reliably estimated.

29.5.1 All-Cause Mortality: Evidence from Population Cohorts

Large Registry-Based Studies

The most rigorous evidence comes from population-based registry studies with appropriate comparison groups:

Roberts et al. (2016) – England and Wales National Registry. This landmark study published in *The Lancet* [52] analyzed mortality in 2,147 ME/CFS patients identified through English and Welsh general practice registries, with 7-year follow-up (2007–2013). The study recorded 17 deaths during follow-up.

Key findings:

- **All-cause mortality SMR: 1.14** (95% CI: 0.65–1.85, $p = 0.67$)
- No statistically significant elevation in all-cause mortality
- **Cancer-specific SMR: 1.39** (95% CI: 0.60–2.73, $p = 0.45$) – not significant
- **Suicide-specific SMR: 6.85** (95% CI: 2.22–15.98, $p = 0.002$) – *highly significant*

Notably, 5 of the 17 deaths were suicides, and 60% of suicide victims had no documented depression diagnosis. The authors interpreted this as suggesting that ME/CFS-specific factors (functional limitation, hopelessness about prognosis, dismissive medical encounters) may contribute to suicide risk independent of comorbid psychiatric conditions, though the small sample size ($n=5$ suicides) limits the strength of this inference.

Smith et al. (2006) – US Multi-Center Cohort. A US study [629] followed 1,201 patients with chronic fatigue for up to 14 years, using National Death Index (NDI) linkage for mortality ascertainment.

Key findings:

- **All-cause mortality: No elevation** above expected rates for age and sex
- **Suicide rate: >8 times higher** than US general population
- SMR for suicide particularly elevated in “chronic fatigue” not meeting full CFS criteria (SMR: 14.2) compared to CFS (SMR: 3.6)
- The authors noted that the higher suicide rate in chronic fatigue (vs. CFS) patients may reflect the additional burden of lacking a recognized diagnosis

Conflicting Evidence: Memorial Record Studies

Studies based on memorial records and caregiver surveys have reported more concerning findings, but these suffer from significant selection bias toward severely ill and deceased patients:

McManimen et al. (2016) – Caregiver Survey. Analysis of 56 deaths reported by caregivers [130] found:

- Mean age at death: **55.9 years** vs. 73.5 years in general population ($p < 0.0001$)
- 48.2% of deceased were bedridden before death
- Mean age at cardiovascular death: 58.8 years vs. 77.7 years ($p < 0.0001$)

Critical limitation: Memorial records inherently overrepresent severe cases and premature deaths (survivors do not appear in memorials). This creates profound selection bias.

Sirotiak & Amro (2025) – Updated Memorial Analysis. The most recent memorial record analysis [131] examined 505 deaths:

- Mean age at death: **52.5 years** (SD = 16.7)
- Most frequent causes: ME/CFS complications (28.3%), suicide (25.4%), cancer (23.0%), cardiovascular disease (14.2%)

While concerning, these findings must be interpreted cautiously given selection bias. The authors acknowledge that memorial records may capture “the tip of the iceberg” of severe, fatal cases rather than representing typical ME/CFS mortality patterns.

29.5.2 Cause-Specific Mortality

Suicide: The Most Robust Finding

Across *all* study types—registry cohorts, clinical cohorts, and memorial records—suicide mortality is consistently and substantially elevated:

Table 29.1: Suicide Mortality Across ME/CFS Studies

Study	SMR or Rate Ratio	Significance
Roberts et al. (2016)	6.85	$p = 0.002$
Smith et al. (2006) – CFS	3.6	Significant
Smith et al. (2006) – Chronic Fatigue	14.2	Highly significant
Jason et al. (2006)	2nd most common cause	—

Suicidal Ideation Prevalence. Cross-sectional surveys reveal alarming rates of suicidal thoughts:

- 39–57% of moderately to severely ill ME/CFS patients report suicidal ideation [135]
- Compare to 4% in general US population
- 7.1% have suicidal ideation *without* clinical depression [630]

Risk Factors for Suicide in ME/CFS. Research has identified ME/CFS-specific suicide risk factors distinct from typical psychiatric risk factors [630]:

- **Severe functional limitations** (strongest predictor)
- Use of “CFS” diagnostic label (associated with stigma) – 2.81× increased risk
- Absence of comorbidities (paradoxically increases risk, possibly due to lack of medical legitimacy) – 3.48× increased risk
- Lack of social support and financial resources
- Hopelessness about prognosis and treatment availability
- Stigma and gaslighting from healthcare providers

Notably, 60% of ME/CFS patients who died by suicide in the Roberts cohort had *no documented depression diagnosis*, suggesting that ME/CFS-specific suffering—not psychiatric comorbidity—drives suicide risk.

Cardiovascular Mortality: Conflicting Evidence

Concerning Signals from Memorial Records. Memorial record studies suggest elevated cardiovascular mortality:

- Mean age at cardiovascular death: 58.8 years vs. 77.7 years [130]

- Cardiovascular disease: 14.2% of deaths in recent memorial analysis [131]
- Heart failure identified as most common cause of death in Jason et al. (2006) [631]

Cardiovascular Disease Prevalence. Epidemiological surveys suggest elevated cardiovascular disease prevalence in ME/CFS:

- Approximately 25% of ME/CFS patients report history of heart disease or hypertension vs. approximately 5% in the general population [632]
- A systematic review and meta-analysis estimated 51.4% prevalence of any cardiac abnormalities [633]
- Analysis of 2021–2022 NHIS data reported aOR 3.26 (95% CI: 2.85–3.72, $p < 0.001$) for cardiovascular disease after adjusting for traditional risk factors [634]

Mechanism Uncertainty. Critically, cardiovascular dysfunction in ME/CFS does *not* appear to follow typical atherosclerotic pathways. Instead, it is characterized by:

- Reduced stroke volume and cardiac output
- Impaired cerebral blood flow
- Small heart size (“athlete’s heart” in reverse)
- These abnormalities are *not* influenced by deconditioning

Implication: Standard cardiovascular risk models developed for atherosclerotic disease may not apply to ME/CFS. Whether this translates to elevated mortality risk remains unclear, and large registry studies have not confirmed elevated cardiovascular mortality.

Cancer Mortality: No Evidence of Elevation

Despite cancer appearing in memorial records (23.0% of deaths [131]), population-based studies find no significant elevation:

- Roberts et al.: Cancer-specific SMR 1.39 (95% CI: 0.60–2.73, $p = 0.45$) [52]
- Age at cancer death in memorial records not significantly different from general population

29.5.3 Actuarial and Insurance Industry Perspective

Mortality Risk Assessment

The insurance industry has begun evaluating ME/CFS mortality risk. Gen Re, a major global reinsurer, published an analysis in 2023 [138] concluding:

“There seems to be no significant difference between all-cause mortality rates of ME/CFS patients and the general population.”

However, the report notes that patients with “very severe fatigue” may have elevated cardiovascular mortality, though evidence remains limited.

Disability vs. Mortality: The Primary Actuarial Concern

Critically, **disability—not mortality—represents the primary actuarial risk** in ME/CFS:

- Recovery rates: <10%
- Unemployment: 35–69%
- Annual US economic costs: \$17–24 billion
- Functional impairment drives actuarial risk more than mortality

This distinction matters: life insurance underwriters assess *mortality* risk, while disability insurers assess *functional capacity*. ME/CFS poses greater risk to the latter.

29.5.4 Methodological Challenges and Evidence Quality

Why Study Results Differ

The stark discrepancy between registry studies (no elevated all-cause mortality) and memorial records (mean age at death 52–56 years) reflects methodological differences:

1. **Selection bias:** Memorial records capture only deaths, inherently overrepresenting severe and fatal cases. Registry studies capture all diagnosed patients regardless of outcome.
2. **Case definition:** Broader “chronic fatigue” definitions (Smith et al.) vs. strict ME/CFS criteria (Roberts et al.) vs. patient-identified cases (memorial records) represent different populations.
3. **Cohort source:** Clinical cohorts (treatment-seeking patients) vs. population-based registries (all diagnosed patients) vs. memorial nominations (deceased patients) have fundamentally different selection mechanisms.
4. **Follow-up duration:** Longer studies (Smith: 14 years) may capture delayed mortality effects better than shorter studies (Roberts: 7 years).

Highest-Quality Evidence

The most methodologically rigorous studies are:

- **Roberts et al. (2016):** $n = 2,147$, 7-year follow-up, registry-based, appropriate comparison group [52]
- **Smith et al. (2006):** $n = 1,201$, up to 14-year follow-up, clinic-based with NDI linkage [629]

Both studies found *no elevation in all-cause mortality* but *substantial elevation in suicide mortality*.

29.5.5 Summary: Evidence-Based Conclusions

1. **All-cause mortality:** No consistent evidence of elevation in large, well-designed cohort studies with appropriate comparison groups. Memorial record studies showing early death are subject to severe selection bias.
2. **Suicide mortality:** *Consistently and substantially elevated* (6–8× general population) across all study types. This represents a legitimate and well-documented mortality risk.
3. **Cardiovascular mortality:** Conflicting evidence. Memorial records suggest elevation, but mechanism differs from atherosclerotic disease and large registry studies have not confirmed excess mortality. Requires further research.
4. **Cancer mortality:** No evidence of elevation in population-based studies.
5. **Life expectancy:** Cannot be reliably estimated due to methodological limitations. Best available evidence suggests normal life expectancy for all-cause mortality, with elevated suicide risk as the primary exception.
6. **Actuarial implications:** *Disability* represents greater actuarial risk than *mortality* in ME/CFS. Life insurance underwriters may focus on suicide risk (well-documented) but have weak evidence for blanket mortality risk assessment.

Observation 106 (Suicide Prevention as Clinical Priority). The robust evidence of elevated suicide mortality—particularly among patients without comorbid depression—highlights suicide prevention as a critical clinical priority in ME/CFS care. Risk factors include severe functional limitation, lack of social support, medical dismissal, and hopelessness about prognosis. Clinical interventions should address ME/CFS-specific suffering (energy limitations, loss of identity and purpose, medical gaslighting) rather than treating suicide risk as a purely psychiatric issue.

29.6 Comorbidity Studies

ME/CFS rarely occurs in isolation. Comorbid conditions are the rule rather than the exception, and their recognition and treatment is a clinical priority (Section 16.1).

29.6.1 Patterns of Comorbid Conditions

The most frequently co-occurring conditions form a recognizable cluster:

- **Fibromyalgia:** 50–70% overlap. Shared features include widespread pain, fatigue, cognitive dysfunction, and sleep disturbance. Whether fibromyalgia and ME/CFS represent distinct entities or overlapping phenotypes of a common pathophysiology remains debated
- **Postural orthostatic tachycardia syndrome (POTS):** 50–70% prevalence in ME/CFS cohorts. POTS may develop concurrently with or subsequent to ME/CFS onset

- **Mast cell activation syndrome (MCAS):** Estimated 15–30% prevalence, likely underdiagnosed [235]. MCAS explains many of the “allergic” and inflammatory symptoms in ME/CFS
- **Hypermobile Ehlers-Danlos syndrome (hEDS):** Elevated co-occurrence, potentially reflecting shared connective tissue and autonomic dysfunction. The ME/CFS–POTS–hEDS–MCAS tetrad is increasingly recognized as a distinct clinical phenotype
- **Irritable bowel syndrome (IBS):** 50–60% prevalence. Reflects autonomic GI dysfunction and gut barrier impairment
- **Migraine:** 30–50% prevalence, potentially related to neuroinflammation and autonomic dysfunction
- **Endometriosis:** Elevated prevalence in female ME/CFS patients, suggesting shared immune and inflammatory mechanisms
- **Interstitial cystitis/painful bladder syndrome:** Elevated prevalence, part of the central sensitization spectrum

29.6.2 Temporal Relationships

Comorbidities may precede, coincide with, or develop after ME/CFS onset:

- **Pre-existing:** hEDS, MCAS, and migraine often predate ME/CFS onset, suggesting they may represent predisposing factors or shared vulnerability
- **Concurrent onset:** POTS and fibromyalgia frequently develop alongside ME/CFS, often triggered by the same infectious event
- **Secondary development:** IBS, depression, and anxiety typically develop after ME/CFS onset and may represent consequences of the disease or its associated physiological disruption
- **Progressive accumulation:** Many patients report accumulating additional diagnoses over time, consistent with progressive multi-system dysfunction

29.6.3 Shared Mechanisms

The clustering of comorbidities in ME/CFS suggests shared underlying mechanisms rather than coincidental co-occurrence (see Chapter 13 for detailed models):

- **Autonomic dysfunction:** A common denominator linking POTS, IBS, temperature dysregulation, and bladder dysfunction
- **Central sensitization:** May underlie fibromyalgia, migraine, IBS, and interstitial cystitis
- **Immune dysregulation:** Connects MCAS, autoimmune features, and chronic inflammation
- **Connective tissue abnormality:** hEDS-associated tissue laxity may contribute to vascular compliance issues (POTS), visceral hypermobility (IBS), and joint instability
- **Neuroinflammation:** May simultaneously produce cognitive dysfunction, pain, autonomic instability, and mood disturbance

30 Controversies and Debates in ME/CFS Research

30.1 Nomenclature and Definition

The naming of this disease has been contested since its earliest descriptions, and the terminology debate has direct consequences for research funding, clinical recognition, and patient treatment.

Myalgic Encephalomyelitis (ME): Coined by Melvin Ramsay in 1956 following the Royal Free Hospital outbreak, the term implies inflammation of the brain and spinal cord (encephalomyelitis). Critics argue that neuropathological evidence for encephalomyelitis is inconsistent, making the name pathologically imprecise. Proponents counter that neuroinflammation has since been documented by PET imaging (Section 28.5).

Chronic Fatigue Syndrome (CFS): Adopted by the CDC in 1988, this name was intended to be neutral and descriptive. It has been widely criticized for trivializing the disease by emphasizing “fatigue”—a universal human experience—while omitting the hallmark feature of post-exertional malaise. The name has contributed to stigma, with patients reporting that the “fatigue” label leads clinicians and the public to equate ME/CFS with ordinary tiredness or laziness.

ME/CFS: The combined term is now the most widely used in clinical and research literature, adopted by the US NIH, NICE, and most major research institutions. It represents a compromise that acknowledges both historical names while avoiding the implications of either alone.

Systemic Exertion Intolerance Disease (SEID): Proposed by the Institute of Medicine (now NAM) in 2015 as a replacement name that emphasises the core feature of exercise intolerance. The proposal has not achieved widespread adoption, partly because patients and researchers objected to the word “intolerance” (perceived as implying psychological aversion) and partly because the acronym is unfamiliar.

The naming controversy matters because terminology shapes perception. The label “chronic fatigue syndrome” has measurably increased medical dismissal, reduced research funding relative to disease burden, and contributed to the psychogenic framing that has dominated clinical attitudes for decades.

30.2 Diagnostic Criteria Controversies

Multiple diagnostic criteria exist for ME/CFS, and the choice of criteria profoundly affects which patients are included in research studies—and therefore what conclusions are drawn about the disease.

Oxford criteria (1991): The broadest definition, requiring only unexplained fatigue of ≥ 6 months duration. Does not require PEM. Widely criticized for including patients with primary psychiatric fatigue, idiopathic chronic fatigue, and other conditions that are pathophysiologically distinct from ME/CFS. Studies using Oxford criteria consistently report better outcomes from exercise and psychological therapies—likely because many included patients do not have ME/CFS.

Fukuda/CDC criteria (1994): Require ≥ 6 months of unexplained fatigue plus 4 of 8 minor symptoms. PEM is listed as one of 8 optional symptoms rather than a required feature. The Fukuda criteria define a heterogeneous population that includes but is not limited to ME/CFS.

Canadian Consensus Criteria (CCC, 2003): Require PEM as a mandatory feature alongside neurocognitive dysfunction, sleep disturbance, and autonomic/neuroendocrine/immune symptoms. The CCC defines a more clinically homogeneous population and is preferred by most ME/CFS researchers and specialist clinicians.

International Consensus Criteria (ICC, 2011): The most restrictive criteria, requiring post-exertional neuroimmune exhaustion plus neurological, immune, and energy production impairments. Removes the 6-month waiting period, enabling earlier diagnosis.

IOM/SEID criteria (2015): Require substantial reduction in function plus PEM plus unrefreshing sleep, along with cognitive impairment or orthostatic intolerance. Designed for clinical use with minimal burden.

The criteria controversy has direct research implications: meta-analyses combining studies using different criteria mix distinct patient populations, diluting effect sizes and obscuring true therapeutic signals. The field is moving toward mandatory use of PEM-requiring criteria (CCC or ICC), but the legacy of Oxford-criteria studies continues to influence clinical guidelines and public perception.

30.3 The PACE Trial Controversy

The PACE trial (White et al., 2011) was the largest and most expensive ME/CFS treatment study ever conducted (£5 million, $n=641$), comparing graded exercise therapy (GET), cognitive behavioral therapy (CBT), adaptive pacing therapy (APT), and specialist medical care (SMC). Its conclusions—and their subsequent dismantling—constitute the most significant controversy in ME/CFS research history.

30.3.1 What the Trial Claimed

The original publication in *The Lancet* reported that GET and CBT were “moderately effective” treatments that improved fatigue and physical function compared to APT and SMC. A subsequent “recovery” analysis claimed that 22% of patients in the GET and CBT arms had “recovered.”

30.3.2 Criticisms of Methodology

Extensive criticism has identified fundamental methodological problems:

- **Mid-trial outcome changes:** The threshold for “improvement” on the Chalder Fatigue Scale was lowered from ≤ 18 to ≤ 22 , and for physical function from ≥ 85 to ≥ 60 on the SF-36. Under the revised thresholds, 13% of participants met “recovery” criteria at baseline—before any treatment [506]
- **Subjective outcomes only:** All primary outcomes were self-reported questionnaires administered by non-blinded researchers. Objective measures (six-minute walk test, step-count actigraphy, employment status, welfare benefit claims) showed no significant improvement [57]
- **Non-blinded design:** Participants and therapists knew which treatment was being administered, and outcome assessors were not independent. In a trial using subjective outcomes, this is a critical source of bias
- **Oxford criteria:** The trial used the broadest diagnostic criteria, likely including patients without ME/CFS
- **Lack of PEM monitoring:** No systematic assessment of post-exertional malaise—the defining feature of the disease under study

30.3.3 Reanalysis Results

After a protracted legal battle for data access under the Freedom of Information Act, independent reanalysis by Wilshire et al. (2018) found:

- Using the trial’s original (pre-change) outcome thresholds, recovery rates dropped from 22% to 7%—not significantly different from the control groups
- Step-count actigraphy showed no significant improvement in any group
- Six-minute walk distance improvements were clinically trivial (approximately 20 meters)
- Long-term follow-up showed convergence of all groups, suggesting no durable benefit

30.3.4 Impact on Clinical Guidelines

The PACE trial’s legacy has been profound and ultimately self-correcting. NICE (UK, 2021) withdrew its recommendation for GET and de-emphasized CBT, explicitly citing the evidence for harm [127]. The CDC (US, 2022) similarly removed exercise therapy recommendations. However, clinical practice in many countries has been slow to follow guideline changes, and GET continues to be prescribed—particularly in countries where ME/CFS recognition lags (see Section 30.5).

The persistence of PACE-derived recommendations was starkly illustrated in Australia, where the Royal Australian College of General Practitioners (RACGP) published a Handbook of Non-Drug Interventions (HANDI) recommendation for “incremental physical activity” in ME/CFS as late as 2024—years after NICE and CDC withdrew equivalent guidance. Stallard

et al. (2026) published a formal critique in the *Australian Journal of General Practice*, demonstrating that an AGREE II appraisal scored the HANDI guideline at only 2% for development rigor [585]. The critique identified the guideline's dependence on the PACE trial and an outdated Cochrane review, documented patient survey data reporting 54–74% harm rates from GET, and called for immediate withdrawal pending the National Health and Medical Research Council (NHMRC) review of ME/CFS guidelines. An invited counterargument by Ewald [635] defended HANDI as a non-drug intervention summary rather than a clinical guideline and cited moderate GRADE evidence from randomized controlled trials, while acknowledging that ME/CFS pathophysiology remains unknown. The exchange illustrates the ongoing tension between legacy evidence syntheses that predate PACE reanalysis and the post-2021 international consensus against GET.

30.3.5 Lessons for Future Research

The PACE trial illustrates how subjective outcomes, non-blinded design, mid-trial protocol changes, and broad case definitions can produce misleading results in ME/CFS research. Future trials should require: objective outcome measures (two-day CPET, actigraphy, biomarkers), PEM-requiring diagnostic criteria, independent outcome assessment, pre-registered protocols with locked outcome thresholds, and systematic PEM monitoring as a safety endpoint.

30.4 Psychogenic vs. Biomedical Models

The most consequential debate in ME/CFS history has been whether the illness is fundamentally psychological or biological. This debate has shaped research funding, clinical treatment, and patient experience for decades.

30.4.1 Historical Context

The psychogenic model of ME/CFS emerged in the 1970s–1980s, influenced by the observation that no single pathogen could be consistently identified as a cause, that routine laboratory tests were often normal, and that fatigue was associated with depression. The “biopsychosocial” model proposed by Wessely, White, and colleagues framed ME/CFS as a condition perpetuated by deconditioning, illness beliefs, and fear avoidance—treatable through exercise and cognitive restructuring.

This model achieved institutional dominance in the UK (through the PACE trial and NICE's pre-2021 guidelines), influencing insurance decisions, disability assessments, and clinical training. Patients who resisted exercise therapy or psychological treatment were characterized as having “unhelpful illness beliefs.”

30.4.2 Evidence for Biological Basis

The accumulated biological evidence against a psychogenic model is now overwhelming:

- **Immune dysfunction:** Reduced NK cell cytotoxicity, T cell exhaustion, altered cytokine profiles, and autoantibodies—none of which are features of psychological conditions [236, 189]
- **Metabolic dysfunction:** Impaired mitochondrial respiration, hypometabolic state, and cellular energy deficits documented by metabolomics and Seahorse analyzer studies [589, 170]
- **Autonomic dysfunction:** Objective documentation of POTS, orthostatic hypotension, and impaired heart rate variability [152, 97]
- **Neuroinflammation:** PET imaging demonstrating microglial activation [464]
- **Neurotransmitter deficiency:** CSF catecholamine deficiency documented by the NIH study [5]
- **Objective exercise intolerance:** Two-day CPET demonstrating measurable physiological deterioration after exertion—impossible to produce through psychological mechanisms [1]

30.4.3 The Role of Psychological Factors

Acknowledging the biological basis of ME/CFS does not deny that psychological factors affect the patient experience. Depression, anxiety, and trauma are common—as consequences of chronic disabling illness, not as causes. Psychological suffering deserves treatment in its own right, but psychological treatment cannot address the underlying pathophysiology. The critical distinction is between “ME/CFS causes psychological distress” (correct) and “psychological distress causes ME/CFS” (unsupported).

30.4.4 Patient Advocacy and the Psychogenic Harm

The psychogenic model has caused measurable harm to patients:

- **Treatment harm:** GET prescribed under the psychogenic model has worsened many patients [586]
- **Diagnostic delay:** Physicians trained to view ME/CFS as psychological often fail to diagnose or investigate the condition, resulting in years of delayed treatment
- **Insurance and disability denial:** Psychogenic framing has been used to deny disability benefits and insurance claims
- **Research underfunding:** Classification as a psychological condition reduced biomedical research funding for decades
- **Stigma:** Patients report that the psychogenic label is the single most damaging aspect of their interaction with the medical system, more distressing than the illness itself

Patient advocacy organizations have been instrumental in challenging the psychogenic model, demanding data transparency (PACE trial reanalysis), and advocating for guideline changes. The 2021 NICE guideline revision [127], which explicitly rejected the biopsychosocial model for ME/CFS, represents a landmark victory for evidence-based medicine driven in significant part by patient advocacy.

30.5 Exercise Therapy Debates

The exercise therapy debate in ME/CFS is closely linked to the psychogenic model and the PACE trial, but warrants separate discussion because of its direct clinical consequences.

30.5.1 Arguments For Graded Exercise Therapy

Proponents of GET argued that ME/CFS symptoms were perpetuated by a cycle of deconditioning and fear avoidance: patients reduced activity due to fatigue, became deconditioned, experienced worsening symptoms, and further reduced activity. GET aimed to break this cycle through progressive, therapist-guided increases in physical activity.

The theoretical basis rested on the observation that deconditioning produces symptoms overlapping with ME/CFS (fatigue, orthostatic intolerance, reduced exercise capacity) and the assumption that these symptoms were functionally equivalent. The PACE trial and earlier studies using broad diagnostic criteria appeared to support this approach.

30.5.2 Evidence of Harm

The evidence against GET in ME/CFS is now extensive:

- **Patient surveys:** 51% of ME/CFS patients report GET worsened their condition across surveys in four countries [586]
- **Physiological evidence:** Two-day CPET demonstrates that exertion produces measurable metabolic failure, not psychological fear [1, 79]
- **Guideline withdrawals:** NICE (2021) withdrew GET, CDC (2022) removed exercise recommendations, and multiple national guidelines now explicitly warn against GET [127]
- **Deconditioning refutation:** The NIH study documented that exercise impairment exceeds what deconditioning predicts, and equal maximum grip strength argues against deconditioning as the primary mechanism [5]

30.5.3 Current Recommendations

The current consensus among ME/CFS specialists is:

- **Pacing, not GET:** Activity management within the energy envelope (Section 23.1) is the recommended approach

- **Individualized activity:** Any movement or exercise must be guided by individual CPET-derived thresholds, not population-based formulas (Section 23.4)
- **Never push through PEM:** The defining clinical principle is that PEM is a pathological signal, not a psychological barrier to be overcome
- **Safe movement is possible:** Within the energy envelope, gentle movement (stretching, isometric exercises, recumbent activities) can prevent deconditioning complications without triggering PEM

30.6 Deconditioning Hypothesis

The deconditioning hypothesis posits that ME/CFS symptoms are primarily maintained by physical inactivity: patients reduce activity due to fatigue, lose cardiovascular and muscular fitness, and experience worsening symptoms that reinforce further inactivity. Under this model, graded exercise therapy (GET) should reverse the cycle. This hypothesis dominated clinical guidance for decades and underpinned the PACE trial's exercise arm.

30.6.1 Arguments For Deconditioning

ME/CFS patients do show objective markers consistent with deconditioning: reduced VO_2peak , lower peak heart rate, and decreased muscle mass in some studies. Prolonged bed rest produces symptoms (orthostatic intolerance, fatigue, cognitive slowing) that overlap with ME/CFS. These parallels led clinicians to assume that reversing inactivity would resolve symptoms.

30.6.2 Why Deconditioning Fails as a Primary Explanation

Multiple lines of evidence demonstrate that deconditioning cannot account for the core pathophysiology of ME/CFS:

1. **Post-exertional malaise is the hallmark feature.** Deconditioned individuals improve with progressive exercise; ME/CFS patients deteriorate. Two-day CPET protocols show reproducible day-2 performance decline in ME/CFS but not in sedentary controls or patients with other fatiguing conditions [1, 79].
2. **VO_2peak reduction exceeds deconditioning predictions.** The NIH deep phenotyping study documented exercise impairment beyond what inactivity alone would produce, even in patients with relatively preserved activity levels [5]. Walitt et al. themselves acknowledged that deconditioning is a consequence, not a cause, and that equal maximum grip strength argues against pure deconditioning.
3. **Metabolic dysfunction is cellular, not peripheral.** Impaired mitochondrial function in immune cells [170] and abnormal metabolite profiles persist independently of activity levels (Chapter 6).

4. **Autonomic and immune abnormalities are independent of fitness.** Chronotropic incompetence, catecholamine deficiency [5], T cell exhaustion, and NK cell dysfunction are not features of deconditioning.
5. **GET produces harm, not recovery.** Large self-reported patient surveys report that 51% of ME/CFS patients worsened with GET [586]. NICE withdrew GET from its 2021 guidelines, declaring it potentially harmful [127].

Charlton et al. (2026), writing in the *British Journal of Sports Medicine*, directly addressed the deconditioning narrative in the context of long COVID [341]. Their editorial—co-authored by Systrom (pioneer of invasive CPET demonstrating preload failure in ME/CFS) and Putrino (leading long COVID researcher at Mount Sinai)—argued that post-exertional malaise cannot be explained by cardiac deconditioning and called for rethinking the pathophysiology of exercise intolerance in post-viral illness. The publication of this argument in a mainstream sports medicine journal marks a significant shift: the deconditioning myth is now being challenged from within the exercise science community itself, not only by ME/CFS specialists and patient advocates.

30.6.3 Deconditioning as Secondary Consequence

Deconditioning does occur in ME/CFS—as a consequence of enforced inactivity due to PEM, not as a cause of symptoms. This distinction is clinically critical: treating secondary deconditioning requires activity management within the energy envelope (pacing; Section 23.1), not progressive exercise that violates it. Conflating consequence with cause has caused measurable harm to patients and delayed appropriate treatment (Section 30.5).

~ Hypothesis 1: The Deconditioning Narrative as Iatrogenic Harm Multiplier

Certainty: 0.50. The individual components are well-documented (GET harms, deconditioning beliefs among clinicians, delayed diagnosis patterns). The systems-level causal chain linking belief → prescription → harm → worsening is strongly supported by patient surveys [586] and guideline changes [127], but has not been tested as a formal causal model with appropriate controls.

This hypothesis proposes that the deconditioning narrative does not merely fail to help ME/CFS patients—it functions as an active disease modifier that worsens long-term outcomes through a specific iatrogenic causal chain:

1. **Belief formation:** Clinician attributes symptoms to deconditioning based on training, guidelines, or institutional culture.
2. **Exercise prescription:** Patient is prescribed graded exercise therapy or encouraged to “push through” fatigue.
3. **PEM trigger:** Exercise exceeds the patient’s energy envelope, triggering post-exertional malaise.
4. **Misinterpretation:** Crash is interpreted as insufficient effort, psychological resistance, or temporary setback rather than pathological response.
5. **Escalation:** More exercise is prescribed, or patient internalises the message and

pushes harder.

6. **Cumulative damage:** Repeated PEM episodes cause cumulative deterioration via the ratchet mechanism (Speculation 5), potentially converting mild–moderate disease to severe.

The hypothesis predicts that **the deconditioning belief itself is a measurable disease modifier**: patients whose clinicians believe in deconditioning will have worse long-term outcomes than patients whose clinicians recognize PEM, *independent of treatment received*, because the belief shapes the entire clinical encounter—advice given, symptoms dismissed, investigations not ordered, diagnoses delayed.

Testable Predictions.

1. Retrospective cohort comparison: ME/CFS patients whose medical records document deconditioning-based advice will show greater functional decline over 5 years than patients whose records document PEM-aware management, after controlling for baseline severity and comorbidities.
2. Time-to-diagnosis will be longer in patients initially managed under the deconditioning paradigm.
3. The proportion of patients progressing from mild/moderate to severe will be higher in cohorts managed with GET than in cohorts managed with pacing from onset.
4. Regional variation in GET prescription rates (pre-2021 UK vs. Scandinavian countries that abandoned GET earlier) will correlate with regional rates of severe ME/CFS.

Falsifiability. This hypothesis would be falsified if patients managed under the deconditioning paradigm show equal or better long-term outcomes than patients managed with PEM-aware approaches, after controlling for baseline severity. It would also be weakened if the progression from mild to severe ME/CFS occurs at equal rates regardless of clinical management approach.

Limitations. Retrospective cohort studies cannot fully control for confounders (patients who receive deconditioning-based care may differ systematically from those who receive PEM-aware care). Selection bias is likely: patients who seek PEM-aware clinicians may be more medically literate or have better access to specialist care. A randomized trial would be unethical given existing evidence of GET harm.

? Open Question 1: The Sports Medicine Paradigm Shift as Natural Experiment

The publication of Charlton et al. (2026) in the *British Journal of Sports Medicine* [341]—challenging cardiac deconditioning from within the exercise science community—represents a sociologically significant event. Combined with the 2021 NICE guideline change [127], it creates a natural experiment:

- Do clinical practices actually change after paradigm-shifting publications? How

quickly?

- Can we measure the lag between evidence publication and practice change in GET prescription rates?
- Does the source of the challenge matter—do sports medicine journals carry more authority with exercise-prescribing clinicians than ME/CFS specialist journals?
- What is the “half-life” of iatrogenic harm after the evidence changes—how long do outdated deconditioning-based practices persist in clinical settings?

Tracking referral patterns, GET prescription rates, and patient outcomes correlated with publication dates would provide the first empirical study of how long iatrogenic harm persists after the evidence base shifts. This has implications beyond ME/CFS for understanding knowledge translation failures in medicine.

30.7 The NIH “Effort Preference” Controversy (2024–2025)

The 2024 NIH deep phenotyping study by Walitt et al. [5] generated one of the most significant controversies in recent ME/CFS research history. While the study documented multiple objective biological abnormalities (catecholamine deficiency, immune dysfunction, autonomic abnormalities), its interpretive framing around “effort preference” sparked intense criticism from patients, clinicians, and researchers.

30.7.1 The Central Claim

The study concluded that “effort preference, not fatigue, is the defining motor behavior” of post-infectious ME/CFS. The authors proposed that ME/CFS patients have altered “effort preference”—defined as “how much effort a person subjectively wants to exert”—due to dysfunction of integrative brain regions, particularly the temporoparietal junction (TPJ).

30.7.2 Why the Framing Was Controversial

Language Echoing Psychogenic Models

The term “preference” implies volition and choice. Critics argued this framing echoed decades of psychogenic characterizations of ME/CFS that attributed symptoms to patients’ beliefs, behaviors, or psychological states rather than biological dysfunction. The language resonated uncomfortably with PACE trial rhetoric about “unhelpful illness beliefs.”

Methodological Problems with the EEfRT

The Effort-Expenditure for Rewards Task (EEfRT) was designed to measure motivation for rewards in psychiatric conditions, with an explicit requirement that tasks be easy enough that fatigue does not confound results. In the Walitt study:

- ME/CFS patients completed only 65% of hard trials vs. 96–99% for controls
- Seven of 15 patients performed below any control participant
- Physical function scores (SF-36) were 28.7 for patients vs. 97.5 for controls

Kirvin-Quamme et al.'s reanalysis [627] demonstrated a significant correlation between task completion ability and task choice, indicating the tool measured ability, not preference. Their conclusion: patients were “unable,” not “unwilling.”

Failure to Document PEM

The study used single-day CPET rather than the gold-standard two-day protocol, failing to objectively document post-exertional malaise—the defining feature of ME/CFS. PEM was mentioned only three times in the entire paper.

Selection Bias

By excluding severely affected patients (25% of the ME/CFS population), the study could not characterize the full spectrum of disease severity.

30.7.3 Published Academic Responses

Nature Communications Commentary

Davenport et al. published a formal critique in *Nature Communications* [628] stating the interpretation “risks reinforcing skepticism about the serious biological nature of [ME] and its hallmark of post-exertional malaise (PEM), as well as its potential misclassification as a mental health condition.”

Authors' Reply

Walitt et al. responded [636] clarifying that:

- Deconditioning and ME/CFS are not mutually exclusive
- Deconditioning is a consequence, not cause
- Equal maximum grip strength argues against pure deconditioning
- Impaired performance occurs before oxidative metabolism stress

Critics noted the reply did not address the fundamental EEfRT methodology concerns or the harm of “preference” language.

Frontiers in Psychology Reanalysis

The Kirvin-Quamme et al. reanalysis [627] provided detailed statistical evidence that the EEfRT data supported inability rather than altered preference, calling for proper task calibration in future studies.

30.7.4 NIH Clarification

Following criticism, NIH clarified that “preference” referred to “subconscious or unconscious or pre-conscious calculations by the brain” rather than conscious choice. Patient advocates responded that if unconscious brain dysfunction was the intended meaning, using “preference”—a word implying choice—was misleading and potentially harmful to patients.

30.7.5 Clinical and Research Implications

Potential Harms

- Reinforcement of psychogenic misconceptions among clinicians unfamiliar with ME/CFS
- Justification for continued use of graded exercise therapy despite harms
- Barriers to disability recognition if symptoms are framed as “preference”
- Psychological harm to patients from invalidating language

Lessons for Future Research

- Assessment tools must be appropriate for the population studied
- Language matters: terminology should not inadvertently pathologize or blame patients
- Peer review should include ME/CFS experts and patient representatives
- Objective measures of PEM (two-day CPET) should be standard
- Study designs should include severely affected patients

30.7.6 Separating Data from Interpretation

Despite the interpretive controversy, the Walitt study’s biological findings—CSF catecholamine deficiency, B cell population shifts, autonomic dysfunction, chronotropic incompetence—represent valuable contributions confirmed by other research. The challenge for the field is to build on these objective findings while rejecting framings that risk harm to patients.

Limitation 1: NIH “Effort Preference” Controversy: Limitations on Both Sides

The debate over the Walitt et al. “effort preference” framing illustrates epistemic limitations in both the original study and its critiques. The original study’s n=17 sample, single-day CPET design, and use of a task (EEfRT) not validated for physically impaired populations limit the strength of its conclusions. However, the Kirvin-Quamme et al. reanalysis [627], while methodologically sound, uses the same small dataset; a definitive resolution requires larger studies with tasks calibrated to participants’ physical capacity. Neither side has established whether the observed motor behavior differences reflect conscious preference, unconscious neural computation, or simple physical inability—these are empirically distinguishable but have not been distinguished.

Observation 107 (The Value of Controversial Studies). The NIH deep phenotyping study illustrates how a study can simultaneously advance biological understanding and generate harmful interpretations. The catecholamine findings alone—the first CSF neurotransmitter measurements in ME/CFS—provide crucial mechanistic insight. Future citations should specify which findings are being referenced (the objective biological data vs. the contested “effort preference” interpretation) to prevent misuse while preserving scientific value.

30.8 Post-COVID ME/CFS

The COVID-19 pandemic created the largest wave of post-viral illness in modern history, with an estimated 51% of long COVID patients meeting PEM criteria also satisfying ME/CFS diagnostic criteria [370]. This overlap has profoundly affected both the ME/CFS and long COVID fields.

30.8.1 Overlap with Long COVID

The clinical overlap between post-COVID sequelae and ME/CFS is substantial:

- **Shared symptoms:** Fatigue, post-exertional malaise, cognitive dysfunction, sleep disturbance, orthostatic intolerance, and pain are core features of both conditions
- **Shared pathophysiology:** Immune dysregulation, mitochondrial dysfunction, neuroinflammation, autonomic dysfunction, and gut microbiome alterations are documented in both
- **Shared biomarkers:** Reduced NK cell function, altered cytokine profiles, metabolic abnormalities, and T cell exhaustion markers overlap between the two populations
- **Diagnostic convergence:** Many long COVID patients who initially presented with respiratory or multi-organ symptoms eventually develop a symptom profile indistinguishable from ME/CFS, particularly those whose primary complaint is PEM-limited fatigue

30.8.2 Differences in Recognition and Funding

The disparity in societal response to long COVID versus ME/CFS has been a source of both frustration and opportunity for the ME/CFS community:

- **Funding disparity:** Long COVID research received billions in dedicated funding within 2 years of the pandemic (NIH RECOVER initiative: \$1.15 billion). ME/CFS, with a comparable disease burden and decades of documented pathophysiology, receives approximately \$15 million annually from the NIH
- **Medical legitimacy:** Long COVID was immediately accepted as a biomedical condition by the medical establishment, while ME/CFS patients spent decades fighting psychogenic dismissal despite presenting identical symptoms after different viral triggers
- **Research infrastructure:** Long COVID research has built clinical trial networks, biobanks, and multi-site collaborations that ME/CFS has lacked. ME/CFS researchers have advocated for inclusion in these networks
- **Societal recognition:** Long COVID achieved rapid public awareness and workplace accommodation frameworks that ME/CFS patients never received, despite similar functional impairment

30.8.3 Opportunities and Challenges

The long COVID–ME/CFS convergence creates both opportunities and risks:

- **Opportunity — research acceleration:** Long COVID funding is generating insights directly applicable to ME/CFS. Treatments developed for long COVID PEM will likely benefit ME/CFS patients
- **Opportunity — diagnostic legitimacy:** The recognition that viral infection can produce chronic multi-system illness validates the ME/CFS disease model and may reduce clinical stigma
- **Risk — diagnostic erasure:** Some clinicians may subsume ME/CFS under the “long COVID” label, potentially erasing decades of ME/CFS-specific research and patient identity for non-COVID-triggered cases
- **Risk — research silos:** Long COVID research conducted without awareness of the existing ME/CFS literature risks repeating solved problems and missing established pathophysiological knowledge
- **Challenge — heterogeneity:** Combining ME/CFS and long COVID populations in research may increase sample heterogeneity if post-COVID patients include those with organ damage (pulmonary fibrosis, cardiac injury) rather than the functional multi-system dysfunction characteristic of ME/CFS

30.9 Clinical Utility for Respiratory and Allergy Medicine

ME/CFS research has generated diagnostic and therapeutic insights with broader applicability to respiratory and allergy medicine, suggesting that pneumologists and allergologists are well-

positioned to contribute to both ME/CFS research and management of related conditions in general practice.

30.9.1 Transferable Diagnostic Techniques

Two-Day Cardiopulmonary Exercise Testing

The two-day CPET protocol, which objectively quantifies post-exertional malaise through measurement of performance decline between testing days [79], has utility beyond ME/CFS diagnosis:

- **Distinguishing deconditioning from pathology:** In disability evaluations, malingering concerns, or contested diagnoses, day-2 performance decline provides objective biomarker
- **Post-viral assessment:** Identifying which respiratory infection survivors require specialized chronic illness management vs. standard recovery
- **Exercise prescription safety:** Determining whether exercise rehabilitation is appropriate or contraindicated

The meta-analysis by Lim et al. demonstrated significant workload decline at ventilatory threshold in ME/CFS patients (-14.6W) while controls showed improvement (+6.5W, $p=0.01$), providing reproducible objective measurement.

Capnography in Orthostatic Assessment

Standard vital sign monitoring misses orthostatic hypocapnia, present in 20.6% of ME/CFS patients vs. 2.9% of controls [374]. Integration of capnography during tilt-table testing or simple lean tests can identify:

- Breathing pattern disorders causing “unexplained” dizziness or brain fog
- Hypocapnic cerebral hypoperfusion (25% drop in CO₂, 50% drop in brain blood flow)
- Patients whose orthostatic symptoms will respond to breathing retraining rather than pharmacologic intervention

Gas Exchange Analysis During Exercise

ME/CFS research has clarified that severe dyspnea can occur despite normal pulmonary function tests, with the primary pathology involving peripheral oxygen extraction at the skeletal muscle level rather than central cardiopulmonary dysfunction [637]. This insight prevents unnecessary invasive testing and redirects clinical investigation toward appropriate mechanisms.

30.9.2 Underdiagnosed Comorbidities in Allergy Practice

Mast Cell Activation Syndrome

MCAS develops in approximately 25% of ME/CFS patients over the disease course [235], but prevalence in general allergy practice populations remains understudied. The condition is likely underdiagnosed due to:

- Non-specific symptoms overlapping with common allergic conditions
- Lack of widely available, validated diagnostic criteria
- Episodic nature making timing of testing challenging

Simple therapeutic trials with H1/H2 antihistamine combinations (e.g., loratadine + famotidine) show 72% response rates in Long COVID/ME/CFS populations, suggesting empiric trials may be both diagnostic and therapeutic for patients with treatment-resistant “allergic” symptoms.

Histamine Intolerance

While affecting only 1–3% of the general population, histamine intolerance appears markedly more prevalent in ME/CFS, Long COVID, and fibromyalgia. Recognition of this pattern may identify an underserved patient subgroup in allergy practices.

30.9.3 Post-Viral Respiratory Syndrome Recognition

The COVID-19 pandemic revealed that 51% of Long COVID patients meeting PEM criteria satisfied ME/CFS diagnostic criteria [370], establishing ME/CFS as a common outcome of post-viral syndromes following respiratory infections. This pattern extends beyond SARS-CoV-2:

- 40.3% of SARS-CoV-1 survivors developed unexplained chronic fatigue
- 27% of SARS-CoV-1 survivors fulfilled CFS diagnostic criteria
- Post-influenza, post-pneumonia, and post-mononucleosis chronic fatigue syndromes share overlapping features

Pneumologists treating respiratory infection survivors are positioned to:

- Identify early who is at risk for chronic sequelae
- Implement preventive rest protocols during acute illness
- Recognize when referral to ME/CFS specialists is appropriate
- Apply two-day CPET for objective documentation

30.9.4 Research Opportunities

Areas where respiratory and allergy medicine expertise could advance ME/CFS understanding:

- **Ventilation-perfusion mismatch mechanisms:** Indirect evidence exists from elevated VE/VCO₂ ratios; direct V/Q scanning studies are lacking
- **Respiratory muscle function:** Objective diaphragm function assessment in patients reporting respiratory muscle fatigue
- **MCAS prevalence in general populations:** Establishing baseline rates in allergy practice for comparison to ME/CFS cohorts
- **Breathing pattern disorder interventions:** Capnometry-guided retraining protocols adapted for ME/CFS energy limitations

★ Achievement 1: Immunomodulatory Therapeutic Approaches

Low-dose naltrexone (LDN), originally investigated for immune modulation, shows 73.9% positive response rates in ME/CFS [536] through restoration of TRPM3 ion channel function in natural killer cells. This mechanism-based approach exemplifies how immunologic expertise translates to novel ME/CFS therapeutics.

30.10 Research Funding Disparities

ME/CFS is one of the most underfunded diseases relative to its burden in the history of modern biomedical research. This underfunding is both a cause and a consequence of the controversies described in this chapter.

30.10.1 Historical Underfunding

NIH funding for ME/CFS remained below \$10 million annually for most of the 2000s–2010s, despite an estimated disease burden of \$17–24 billion in annual economic losses and approximately 1–2.5 million affected Americans. For comparison, multiple sclerosis (with a similar US prevalence) receives approximately \$130 million annually, and HIV/AIDS (with a lower US prevalence) receives over \$3 billion. The funding disparity per patient-year-of-disability is estimated at 10–100-fold.

The causes of underfunding are interrelated:

- **Psychogenic classification:** When ME/CFS was classified as primarily psychological, it fell outside the mandate of biomedical research institutes
- **Absence of a biomarker:** Without a diagnostic test, ME/CFS lacked the “hard science” credentials that attract competitive grant funding
- **Stigma:** Researchers reported that working on ME/CFS was professionally stigmatizing, discouraging junior investigators from entering the field

- **Small research workforce:** Chronic underfunding created a small, isolated research community with limited capacity to generate the volume of publications needed to attract further funding—a vicious cycle
- **Patient invisibility:** Housebound and bedbound patients are invisible to society, unable to participate in advocacy activities that drive research funding for other diseases

30.10.2 Recent Improvements

ME/CFS research funding has increased modestly since 2015:

- NIH funding rose from approximately \$5 million (2014) to \$15–18 million (2023), driven partly by the creation of ME/CFS research centers and the NIH intramural study
- The long COVID pandemic generated spillover funding for ME/CFS through overlap studies
- Private foundations (Open Medicine Foundation, Solve ME/CFS Initiative, ME Research UK) have contributed substantial additional funding and research infrastructure
- International initiatives (European ME/CFS research network, DecodeME in the UK) have expanded the research base beyond the US

30.10.3 Remaining Gaps

Despite improvements, critical funding gaps persist:

- **Clinical trials:** No large, multi-center, biomarker-stratified clinical trial has been funded for ME/CFS. The trials that exist are small, single-center, and investigator-initiated
- **Severe patients:** Research on severe and very severe ME/CFS is virtually non-existent because these patients cannot travel to research centers. Home-based and remote research protocols require dedicated funding
- **Longitudinal cohorts:** No prospective cohort study with ≥ 20 -year follow-up exists, leaving fundamental questions about natural history and mortality unanswered
- **Training pipeline:** The ME/CFS research workforce remains critically small. Dedicated training grants and career development awards are needed to attract new investigators
- **Funding parity:** Even at current levels, ME/CFS receives less than 10% of the per-patient funding of comparable chronic diseases. Reaching funding parity would require a 5–10-fold increase

31 Translational Findings: Implications for Related Conditions

This chapter identifies mechanisms, biomarkers, and treatment protocols from ME/CFS research that have significant implications for other medical conditions. Rather than viewing ME/CFS as an isolated disease, we recognize it as part of a spectrum of post-viral, autoimmune, mitochondrial, and dysautonomic conditions that share common pathophysiology.

The findings presented here represent **translational opportunities**—research from ME/CFS that could advance understanding and treatment of related conditions, even in patients who do not meet full ME/CFS diagnostic criteria.

31.1 Introduction to Translational Medicine

31.1.1 Why ME/CFS Research Benefits Other Conditions

ME/CFS research has identified mechanisms that extend beyond the specific diagnostic boundaries of the illness:

- **Post-viral autoimmunity:** Plasma cell-mediated GPCR autoantibodies (Chapter 7)
- **Autonomic-vascular integration:** β 2-adrenergic receptor dysfunction linking MCAS, POTS, and vascular dysfunction
- **Mitochondrial pathophysiology:** WASF3-mediated ER stress, NAD⁺ depletion, oxidative stress cascades
- **Neuroinflammation:** Microglial activation, glymphatic clearance failure
- **Exercise intolerance mechanisms:** Two-day CPET findings revealing autonomic-metabolic integration failure

These mechanisms are not exclusive to ME/CFS. They represent fundamental pathophysiological processes that manifest across multiple conditions.

⚠ Limitation 1: Translational Findings: Shared Mechanisms Do Not Guarantee Shared Treatments

The translational framework in this chapter assumes that mechanisms documented in ME/CFS can inform treatment of related conditions (Long COVID, POTS, fibromyalgia, etc.). This assumption requires qualification:

- Shared symptom profiles and overlapping biomarker abnormalities do not establish

shared aetiology; convergent phenotypes can arise from divergent upstream causes.

- Most ME/CFS treatment evidence cited for translational application comes from open-label or pilot studies (e.g., daratumumab n=10 [9], immunoadsorption n=10 [147]); extrapolating these to other conditions compounds the uncertainty.
- The “certainty levels” framework (High/Medium/Low) used in this chapter is the authors’ judgement, not a validated classification; readers should not interpret “High Certainty” as equivalent to established clinical evidence.
- Condition-specific pathophysiology (e.g., spike protein persistence in Long COVID, central sensitisation in fibromyalgia) may make ME/CFS-derived treatments ineffective or harmful in related conditions.

31.1.2 Certainty Levels for Cross-Condition Application

When applying ME/CFS findings to other conditions, we use a three-tier certainty framework:

High Certainty Mechanism documented in both ME/CFS and target condition; treatment tested in both

Medium Certainty Mechanism documented in ME/CFS; strong biological plausibility for target condition; shared clinical features

Low Certainty Mechanism documented in ME/CFS; theoretical applicability to target condition; requires validation

Important: All translational recommendations require validation through condition-specific research. These findings represent **research opportunities**, not established clinical guidelines for non-ME/CFS conditions.

31.2 Immediate Applicability (Tier 1)

These conditions share substantial pathophysiology with ME/CFS, documented in peer-reviewed literature. Translational findings have high-to-medium certainty.

31.2.1 Long COVID / Post-Acute Sequelae of SARS-CoV-2 (PASC)

Long COVID and ME/CFS share post-viral onset, exercise intolerance with delayed symptom exacerbation, autonomic dysfunction, and cognitive impairment, with a substantial proportion of Long COVID patients meeting activity-based case definitions also meeting ME/CFS diagnostic criteria [638].

Table 31.1: ME/CFS Mechanisms Documented in Long COVID

Mechanism	ME/CFS Evidence	Long COVID Evidence
GPCR autoantibodies	29.5–91% prevalence [71, 72]	β 2-AR, M3 autoantibodies detected
Plasma cell autoimmunity	Daratumumab 60% response [9]	BC007 case reports
Endothelial dysfunction	Peripheral endothelial dysfunction [177]	Microclotting, VWF elevation
NAD ⁺ depletion	Metabolomic studies [12]	NR trial raised NAD ⁺ levels; clinical
Neuroinflammation	PET imaging [73]	MRI, CSF abnormalities
Small fiber neuropathy	SFN studies [335]	Documented in subset

Shared Mechanisms

Novel Translational Findings from ME/CFS

1. **Plasma Cell Targeting (Daratumumab):** Pilot study showed 60% response rate in ME/CFS when rituximab (B-cell depletion) failed [9]. This suggests long-lived plasma cells, not B cells, drive persistent autoantibody production.
Implication for Long COVID: Patients with persistent symptoms despite viral clearance may benefit from plasma cell-directed therapy, particularly those with elevated GPCR autoantibodies.
2. **Immunoadsorption for GPCR Autoantibodies:** 70% rapid improvement during treatment in a pilot study (n=10) of ME/CFS patients with elevated β 2-adrenergic receptor autoantibodies, though only 30% sustained improvement at 6–12 months [147].
Implication for Long COVID: Autoantibody screening could identify subset likely to respond to immunoadsorption.
3. **NAD⁺ Restoration with Nicotinamide Riboside:** A 2025 Long COVID RCT (n=58; 20-week intervention) showed NR 2000 mg/day increased NAD⁺ levels 2.6–3.1 $\times$ [572]; cognitive benefits were variable, with some individuals improving after $\geq$ 10 weeks.
Implication for Long COVID: NAD⁺ depletion may be a shared mechanism; prolonged treatment (>10 weeks) may be needed for clinical benefit.

Treatment Protocols with Translational Potential

- **Mitochondrial support:** CoQ10 ubiquinol (100–300 mg/day) + NADH (10–20 mg/day), D-ribose (5g TID), acetyl-L-carnitine (500–2000 mg/day), NAD⁺ precursors (NR 300–1000 mg/day; the only RCT-tested dose is 2000 mg/day [572]; see Chapter 22)
- **Mast cell stabilization:** Cromolyn sodium, H1+H2 antihistamines, quercetin (for MCAS overlap)
- **Low-dose naltrexone:** 3–4.5 mg at bedtime for neuroinflammation
- **Pacing protocols:** Energy envelope management to prevent PEM-like exacerbation

Certainty: **High** for shared mechanisms; **Medium** for treatment efficacy in Long COVID specifically.

31.2.2 Postural Orthostatic Tachycardia Syndrome (POTS)

27–50% of ME/CFS patients meet POTS diagnostic criteria (heart rate increase ≥ 30 bpm upon standing, or HR ≥ 120 bpm, within 10 minutes). The overlap suggests shared autonomic pathophysiology.

Novel Translational Findings from ME/CFS

1. **Central Catecholamine Deficiency:** The NIH intramural study (Walitt et al. 2024) documented reduced CSF dopamine metabolites (DOPA, DOPAC) and norepinephrine metabolites (DHPG) in ME/CFS patients [639].
Implication for POTS: Central (not just peripheral) catecholamine deficiency may drive compensatory tachycardia. This suggests catecholamine synthesis support (L-tyrosine, Tetrahydrobiopterin (BH4) cofactors) could be therapeutic.
2. **Chronotropic Incompetence on 2-Day CPET:** ME/CFS patients show inadequate heart rate response to exercise workload on Day 2, with autonomic dysfunction (not cardiac pathology) as the primary mechanism [1].
Implication for POTS: Exercise intolerance in POTS may involve central autonomic dysregulation affecting both HR and metabolic responses.
3. **Hypovolemia and Preload Failure:** 10–20% reduction in plasma volume is well-documented in ME/CFS, correlating with orthostatic symptoms (Section 10.2.2).
Implication for POTS: Aggressive blood volume expansion (salt, fluids, fludrocortisone) addresses both conditions.

Treatment Protocols with Translational Potential

- **Catecholamine synthesis support:**
 - L-tyrosine 1500–3000 mg (morning, empty stomach)
 - BH4 cofactor support: Methylfolate 1–5 mg + methylcobalamin 1–5 mg + vitamin C 1000 mg
 - Iron optimization (ferritin 100–200 $\mu\text{g/L}$ target)
 - Vitamin B6 (P5P 25–50 mg), copper if deficient
- **Blood volume expansion:** Salt 8–10g/day, fluids 2–3L/day, fludrocortisone 0.1–0.2 mg
- **Compression garments:** Waist-high compression stockings (20–30 mmHg) + abdominal binders
- **Ivabradine:** Heart rate control without blood pressure drop (off-label)

Certainty: **High** for hypovolemia and autonomic dysfunction; **Medium** for central catecholamine deficiency in POTS.

31.2.3 Fibromyalgia

Fibromyalgia shares chronic widespread pain, fatigue, sleep disturbance, and exercise intolerance with ME/CFS, with substantial symptom overlap whose estimated magnitude varies

widely depending on diagnostic criteria applied.

Novel Translational Findings from ME/CFS

1. **Small Fiber Neuropathy:** Quantitative sensory testing and skin biopsy studies document small fiber dysfunction in a subset of ME/CFS patients, correlating with pain and dysautonomia [335].
Implication for Fibromyalgia: Small fiber neuropathy has been documented in fibromyalgia as well [281]. This suggests shared peripheral nerve pathology beyond central sensitization.
2. **Mitochondrial ATP Depletion:** Multiple ME/CFS studies show impaired ATP production, early lactate accumulation, and elevated acylcarnitines indicating impaired fatty acid oxidation [589, 64].
Implication for Fibromyalgia: Muscle pain and fatigue may reflect energy metabolism failure. Mitochondrial support protocols could address root cause.
3. **Mast Cell Activation:** Ketotifen (mast cell stabilizer) was tested in a fibromyalgia Phase 1 RCT but did not show significant benefit at the dose tested (4 mg/day for 8 weeks; $p = 0.7$ for pain) [640]. Nevertheless, ME/CFS research provides mechanistic understanding of mast cell–pain–fatigue connections that may warrant further investigation at higher doses.
Implication for Fibromyalgia: Mast cell stabilization protocols developed for ME/CFS (cromolyn, quercetin, H1+H2 antihistamines) may benefit fibromyalgia patients with MCAS features.

Treatment Protocols with Translational Potential

- **D-ribose:** 5g TID showed improvements across energy, sleep, and well-being in two open-label studies without placebo control [8, 528]; see Chapter 22 for evidence limitations
- **CoQ10 ubiquinol + NADH:** CoQ10 100–300 mg/day (300 mg used in fibromyalgia trials [569]) + NADH 10–20 mg/day (combination supported by ME/CFS RCTs [7]; see Chapter 22)
- **Acetyl-L-carnitine:** 500–2000 mg/day for neuroprotection and brain fog reduction (see Chapter 22)
- **Low-dose naltrexone:** 3–4.5 mg at bedtime for neuroinflammation and pain modulation
- **NAD⁺ precursors:** NR 300–1000 mg/day (the only RCT-tested dose is 2000 mg/day [572]); see Chapter 22 for full dosing

Certainty: **Medium** for mitochondrial mechanisms; **High** for mast cell involvement; **Medium** for small fiber neuropathy overlap.

31.2.4 Mast Cell Activation Syndrome (MCAS)

MCAS frequently co-occurs with ME/CFS. The Wirth & Löhn (2023) study provides novel mechanistic understanding of this relationship [235].

Novel Translational Findings from ME/CFS

1. **β 2-Adrenergic Receptor Dysfunction as Common Link:** Wirth & Löhn (2023) propose that dysfunctional β 2-adrenergic receptors create bidirectional disease worsening:
 - ME/CFS orthostatic stress desensitizes β 2 receptors
 - Desensitized β 2 receptors favor mast cell degranulation
 - Mast cell mediators worsen orthostatic dysfunction and cerebral hypoperfusion
 - This creates a vicious cycle

Implication for MCAS: β 2-receptor function testing and targeted support may break the cycle.
2. **Vascular Pathomechanisms:** Histamine and bradykinin both cause vasodilation and vascular permeability, leading to preload failure and orthostatic intolerance.

Implication for MCAS: Vascular-focused treatment (beyond antihistamines) may be necessary for patients with prominent orthostatic symptoms.
3. **GPCR Autoantibody-Monocyte Reprogramming:** Hackel et al. (2025) showed that GPCR autoantibodies don't just block receptors—they reprogram monocytes to produce inflammatory cytokines (MIP-1 δ , PDGF-BB, TGF- β 3) [209].

Implication for MCAS: Autoantibody removal (immunoadsorption) plus monocyte modulation (JAK inhibitors) may be more effective than antihistamines alone.

Treatment Protocols with Translational Potential

- **H1 + H2 antihistamine combination** (H1 alone insufficient):
 - Rupatadine 20 mg (triple action: H1 antagonist + PAF antagonist + mast cell stabilizer)
 - Or: Loratadine/cetirizine/fexofenadine + famotidine 20–40 mg BID
- **Mast cell stabilizers:**
 - Quercetin 500–1000 mg BID (more effective than cromolyn in vitro)
 - Cromolyn sodium 200–400 mg QID (prescription)
 - Vitamin C 1000–3000 mg/day
- **Amitriptyline:** 10–50 mg bedtime (unique mast cell inhibition among antidepressants; reduces IL-8, VEGF, histamine release)

Certainty: **High** for H1+H2 combination; **Medium** for β 2-receptor mechanism; **Low** for autoantibody-monocyte pathway in MCAS specifically.

31.2.5 Ehlers-Danlos Syndrome (Hypermobile Type)

Hypermobile Ehlers-Danlos Syndrome (hEDS) frequently co-occurs with POTS (70–80%) and MCAS (31%), creating a recognized clinical “triad.” However, the pathophysiologic mechanisms linking these conditions remain controversial [151].

Established Mechanisms in EDS

EDS literature documents:

- **Structural vascular compliance abnormalities:** Collagen defects → vessel stretching → blood pooling → reduced venous return
- **Adrenergic hyperresponsiveness:** Documented in hEDS cardiovascular autonomic testing [641]
- **Mast cell mechanosensitivity:** Stretch-activated mast cells via ADGRE2, integrins $\alpha V\beta 3$, $\alpha 5\beta 1$ [642]
- **Small fiber neuropathy:** Common in hEDS, contributing to pain and dysautonomia

Novel Translational Findings from ME/CFS

The following mechanisms are **well-documented in ME/CFS but not yet studied in EDS**, representing novel translational opportunities:

1. $\beta 2$ -Adrenergic Receptor Desensitization vs. Hyperresponsiveness:

- **EDS literature:** Documents adrenergic *hyperresponsiveness*
- **ME/CFS literature:** Documents $\beta 2$ -receptor *desensitization* from chronic orthostatic stress [235]
- **Gap:** These may represent different stages or phenotypes. Chronic EDS-related orthostatic stress could lead to eventual desensitization.

Research opportunity: Test $\beta 2$ -receptor function longitudinally in EDS patients to determine if hyperresponsiveness transitions to desensitization.

2. Bidirectional MCAS ↔ $\beta 2$ -Receptor Cycle:

The Wirth & Löhn (2023) model proposes:

- Orthostatic stress → $\beta 2$ -receptor desensitization
- Desensitized $\beta 2$ receptors → mast cell degranulation
- Mast cell mediators (histamine, PAF) → vascular dysfunction
- Vascular dysfunction → worse orthostatic stress

This cycle has **not been studied in EDS**, despite clinical recognition of the hEDS-POTS-MCAS triad.

Research opportunity: Measure $\beta 2$ -receptor function in EDS patients with vs. without MCAS to test this model.

3. Tetrahydrobiopterin (BH4) Dysregulation:

- **ME/CFS findings:** Elevated BH4 and BH2 in patients with orthostatic intolerance [643, 644]
- **Mechanism:** Pentose phosphate pathway activation → BH4 production → iNOS/NO pathway activation → neuroinflammation
- **EDS literature:** No studies found (2020–2026 search)

Research opportunity: Measure BH4 levels in EDS patients with orthostatic intolerance. If elevated, this could explain neuroinflammatory symptoms and provide therapeutic target.

Caveat: BH4 research in ME/CFS is very preliminary (n=10–32, single research group). The paradox of *elevated* BH4 causing dysfunction (rather than deficiency) requires explanation.

4. Endothelial (Functional) vs. Structural Vascular Permeability:

- **EDS mechanism:** Structural collagen weakness → vessel stretching
- **ME/CFS mechanism:** Receptor-mediated endothelial permeability (vasoactive mediators → functional permeability changes)

Research opportunity: Distinguish structural from functional vascular dysfunction in EDS. Patients may have *both* mechanisms, requiring combined treatment.

5. Plasma Cell Autoimmunity:

If EDS patients develop post-viral or autoimmune features, plasma cell-targeted therapy (daratumumab) could be considered, following ME/CFS precedent. However, this is entirely speculative for EDS.

Treatment Protocols with Translational Potential

Key Distinctions: EDS-Specific Considerations

△ Warning 1: EDS vs. ME/CFS Differences

While ME/CFS mechanisms translate to EDS, critical differences exist:

- **Fatigue source:** In EDS, fatigue may result from musculoskeletal compensation for joint instability, not just autonomic/mitochondrial dysfunction
- **Exercise intolerance:** In EDS, joint subluxations and injury risk limit activity; in ME/CFS, metabolic failure causes PEM
- **Pain mechanisms:** In EDS, structural joint instability contributes; in ME/CFS, neuroinflammation and central sensitization dominate
- **Treatment focus:** EDS requires joint protection and physical therapy alongside systemic treatments

Not all EDS patients will respond to ME/CFS-derived protocols. Subset with prominent autonomic dysfunction, MCAS, or post-viral features most likely to benefit.

Research Priorities for EDS

1. **Longitudinal β 2-receptor function testing:** Does hyperresponsiveness transition to desensitization with disease duration?
2. **Tetrahydrobiopterin (BH4) measurement in EDS with orthostatic intolerance:** Is the ME/CFS finding translatable?
3. **Endothelial biomarkers:** Are VWF, fibronectin, thrombospondin elevated in EDS-POTS-MCAS subset?

Table 31.2: ME/CFS Treatment Protocols Applicable to EDS

Protocol	Rationale	Certainty in EDS	Evidence Base
POTS management (salt, fluids, compression, fludrocortisone)	Addresses hypovolemia and preload failure	High	Well-established
Mast cell stabilization (H1+H2 antihistamines, quercetin, cromolyn)	Addresses MCAS in hEDS-MCAS subset	High	Clinical use common
Rupatadine (H1 + PAF antagonist + mast cell stabilizer)	Triple mechanism addresses vascular pathomechanisms	Medium	ME/CFS evidence, not tested in EDS
Catecholamine synthesis support (L-tyrosine, Tetrahydrobiopterin (BH4) cofactors)	Supports autonomic function if central deficiency present	Low-Medium	ME/CFS evidence, not tested in EDS
Pacing and energy envelope management	Prevents post-exertional symptom exacerbation	Medium	Reduces injury risk from hypermobility overexertion
Mitochondrial support (CoQ10, D-ribose, L-carnitine)	Addresses energy deficit from chronic musculoskeletal compensation	Low-Medium	Theoretical, untested in EDS

4. **Controlled trials of rupertadine:** Does PAF antagonism benefit EDS patients with vascular symptoms?

5. **Autoantibody screening:** What percentage of EDS patients have GPCR autoantibodies?

Certainty: **Medium** for vascular mechanisms; **Low-Medium** for β 2-receptor pathway; **Low** for Tetrahydrobiopterin (BH4) dysregulation; **None** for plasma cell autoimmunity.

Bottom line: The Wirth 2023 integrated model (MCAS $\leftrightarrow$ β 2-receptors $\leftrightarrow$ vascular dysfunction $\leftrightarrow$ POTS) represents a **completely untested but biologically plausible hypothesis for EDS**. If validated, it would explain the hEDS-POTS-MCAS triad and provide targeted treatment strategies.

31.3 Strong Mechanistic Overlap (Tier 2)

These conditions share documented pathophysiologic mechanisms with ME/CFS. Translational findings have medium-to-low certainty pending condition-specific validation.

31.3.1 Post-Treatment Lyme Disease Syndrome (PTLDS)

Post-Treatment Lyme Disease Syndrome describes persistent symptoms following antibiotic treatment for Lyme disease. Estimated 10–20% of treated Lyme patients develop PTLDS, with symptom overlap suggesting potential shared mechanisms with ME/CFS.

Shared Mechanisms

- **Post-infectious autoimmunity:** Molecular mimicry triggering cross-reactive antibodies
- **Neuroinflammation persistence:** Microglial activation despite pathogen clearance
- **Autonomic dysfunction:** Orthostatic intolerance, heart rate variability reduction
- **Small fiber neuropathy:** Documented in both PTLDS and ME/CFS via skin biopsy
- **Exercise intolerance with PEM-like symptoms:** Post-exertional symptom exacerbation

Novel Translational Findings from ME/CFS

1. **Immunomodulation with Low-Dose Naltrexone:** ME/CFS studies show LDN 3–4.5 mg reduces neuroinflammation via TLR4 antagonism on microglia.
Implication for PTLDS: If persistent neuroinflammation drives symptoms, Low-Dose Naltrexone (LDN) could provide benefit through microglial modulation.
2. **Autoantibody Screening:** GPCR autoantibodies (β 2-AR, M3/M4) documented in 29.5–91% of ME/CFS patients may also occur in PTLDS if post-infectious autoimmunity is involved.
Implication for PTLDS: Autoantibody testing could identify subset likely to respond to immunoadsorption or plasma cell targeting.
3. **Mitochondrial Support:** CoQ10, D-ribose, L-carnitine, NAD⁺ precursors address energy metabolism dysfunction.
Implication for PTLDS: If mitochondrial dysfunction persists post-treatment, metabolic support protocols could improve fatigue and cognitive symptoms.

Certainty: **Medium** for shared post-infectious mechanisms; **Low** for specific treatment efficacy in PTLDS (requires validation).

31.3.2 Cancer-Related Fatigue and Post-Chemotherapy Syndrome

Cancer-related fatigue (CRF) affects 25–99% of patients during treatment and 30–40% of survivors post-treatment. Chemotherapy-induced peripheral neuropathy (CIPN) and “chemo brain” share mechanistic features with ME/CFS.

Shared Mechanisms

- **Mitochondrial toxicity:** Chemotherapy agents (anthracyclines, platinum compounds) directly damage mitochondria
- **NAD⁺ depletion:** PARP activation for DNA repair depletes NAD⁺ pools

- **Oxidative stress:** Chemotherapy generates reactive oxygen species damaging cellular components
- **Neuroinflammation:** Cytokine elevation causing “chemo brain” (cognitive dysfunction)
- **Autonomic dysfunction:** Treatment-induced damage to autonomic nervous system

Novel Translational Findings from ME/CFS

1. **NAD⁺ Restoration Therapy:** NR 2000 mg/day increased NAD⁺ levels 2.6–3.1 fold in Long COVID (n=58; 20-week intervention; measured at interim timepoints); cognitive benefits were variable, with some improvement after ≥10 weeks [572]. The mechanism of NAD⁺ restoration is hypothesized to also address chemotherapy-induced NAD⁺ depletion, by analogy with the Long COVID findings.

Implication for CRF: NAD⁺ precursors could restore depleted NAD⁺ pools, improving mitochondrial function and reducing fatigue (**post-treatment only**; NAD⁺ supplementation during active cancer is contraindicated due to theoretical tumor growth promotion—see Section 31.7.1, Safety considerations).

2. **Comprehensive Mitochondrial Support Stack:** CoQ10 ubiquinol (100–300 mg/day) + NADH (10–20 mg/day), D-ribose (5g TID), acetyl-L-carnitine (500–2000 mg/day), alpha-lipoic acid (300–600 mg/day; classified under antioxidants in Chapter 22), B vitamins. See Chapter 22 for full dosing and evidence levels.

Implication for CRF: Addresses multiple points of mitochondrial dysfunction caused by chemotherapy. Note: both D-ribose and alpha-lipoic acid can lower blood glucose; monitor closely in patients on corticosteroids or with glucose dysregulation (common during cancer treatment).

3. **Pacing Strategies and Energy Envelope Management:** Prevents boom-bust cycles that worsen fatigue.

Implication for CRF: Helps cancer survivors manage limited energy reserves during recovery without triggering symptom exacerbation.

4. **Vagal Rehabilitation:** Cold exposure, breathing techniques, HRV biofeedback restore autonomic function.

Implication for CRF: Addresses chemotherapy-induced autonomic dysfunction.

Certainty: **Medium-High** for mitochondrial mechanisms; **Medium** for NAD⁺ restoration (promising but needs CRF-specific trials).

31.3.3 Primary Mitochondrial Disorders

Primary mitochondrial disorders result from mutations affecting mitochondrial DNA or nuclear genes encoding mitochondrial proteins, and share core energy metabolism dysfunction with ME/CFS.

Shared Mechanisms

- **ATP depletion:** Impaired oxidative phosphorylation reduces cellular energy

- **Lactate accumulation:** Early shift to anaerobic metabolism
- **Exercise intolerance:** Inability to meet metabolic demands of exertion
- **Oxidative stress:** ROS overproduction from dysfunctional electron transport chain
- **Multi-system involvement:** High-energy organs (muscle, brain, heart) most affected

Novel Translational Findings from ME/CFS

1. **WASF3/ER Stress Pathway:** ME/CFS research identified ER stress inducing WASF3, which disrupts mitochondrial supercomplexes and impairs Complex IV [2].
Implication for Primary Mitochondrial Disorders: ER stress modulators could represent novel therapeutic approach, particularly for disorders involving Complex IV dysfunction.
2. **MitoQ (Mitochondria-Targeted Ubiquinone):** 10–20 mg/day; delivers CoQ10 directly to mitochondria via triphenylphosphonium targeting, achieving greater mitochondrial accumulation than standard CoQ10 in preclinical studies. While developed outside ME/CFS research, its mechanism is particularly relevant to the mitochondrial dysfunction documented in ME/CFS (not reviewed in Chapter 22; limited ME/CFS-specific data; clinical dosing and comparative human pharmacokinetics not yet established).
Implication for Mitochondrial Disorders: More effective CoQ10 delivery to dysfunctional mitochondria.
3. **D-Ribose for ATP Pool Regeneration:** 5g TID showed improvements in energy, sleep, and well-being in two open-label studies without placebo control [8, 528]; see Chapter 22 for evidence limitations. D-ribose is the pentose sugar component of ATP.
Implication for Mitochondrial Disorders: Accelerates adenine nucleotide pool resynthesis, providing ATP precursors that may be depleted in mitochondrial disorders. Does not bypass oxidative phosphorylation—cells still require functional mitochondria for ATP production.
4. **Comprehensive Support Stack:** Combined approach addresses multiple dysfunction points simultaneously.
Implication for Mitochondrial Disorders: ME/CFS protocols provide a mechanistically informed combination therapy framework with partial RCT support (see Chapter 22, Section 22.2.8).

Certainty: **High** for shared mitochondrial dysfunction; **Medium** for treatment efficacy (mechanisms sound, needs validation in primary mitochondrial disorders).

31.3.4 Dysautonomia (General)

Dysautonomia encompasses autonomic nervous system dysfunction causing orthostatic intolerance, heart rate abnormalities, blood pressure dysregulation, and multi-system symptoms.

Novel Translational Findings from ME/CFS

1. **Central Catecholamine Deficiency:** NIH study (Walitt 2024) documented reduced CSF dopamine and norepinephrine metabolites in ME/CFS.

Implication for Dysautonomia: Central (not just peripheral) catecholamine deficiency may drive compensatory tachycardia and orthostatic symptoms. Suggests catecholamine synthesis support (L-tyrosine 1500–3000 mg, BH4 cofactors) could be therapeutic.

2. **Reduced Heart Rate Variability:** ME/CFS shows impaired HRV reflecting autonomic dysregulation.

Implication for Dysautonomia: HRV biofeedback and vagal rehabilitation techniques (cold exposure, extended exhale breathing, gargling) could restore autonomic balance.

3. **Comprehensive Autonomic-Metabolic Protocol:** Combining catecholamine support (tyrosine, BH4 cofactors, iron optimization) with mitochondrial protection (MitoQ, N-Acetylcysteine (NAC), alpha-lipoic acid).

Implication for Dysautonomia: Addresses both neurotransmitter synthesis and cellular energy metabolism underlying autonomic function.

4. **Two-Day CPET Finding:** Autonomic dysregulation (not cardiac pathology) drives chronotropic incompetence and exercise failure.

Implication for Dysautonomia: Focuses treatment on autonomic nervous system rather than cardiac interventions.

Certainty: **Medium-High** for autonomic mechanisms; **Medium** for central catecholamine deficiency (needs validation across dysautonomia subtypes).

31.3.5 Small Fiber Neuropathy (SFN)

Small fiber neuropathy involves damage to small-diameter sensory and autonomic nerve fibers, causing pain, temperature sensation abnormalities, and autonomic symptoms.

Shared Mechanisms

- **Metabolic vulnerability:** Small nerve fibers have high energy demands and are vulnerable to mitochondrial dysfunction
- **Oxidative stress:** ROS damage to nerve fibers
- **Immune-mediated damage:** Inflammation and autoantibodies targeting nerve components
- **Autonomic dysfunction:** SFN commonly causes orthostatic intolerance, GI dysmotility

Novel Translational Findings from ME/CFS

1. **IVIG in Subset with Documented SFN:** Some ME/CFS patients with skin biopsy-confirmed SFN responded to IVIG.

Implication for SFN: If immune-mediated, immunomodulation with IVIG could be therapeutic.

2. **Alpha-Lipoic Acid:** 600 mg/day showed benefit in diabetic neuropathy; mechanism involves mitochondrial antioxidant effects.

Implication for SFN: Addresses oxidative stress damaging small nerve fibers.

3. **Acetyl-L-Carnitine:** 2–3g/day provides neuroprotection via multiple mechanisms (mitochondrial support, neurotrophic effects).

Implication for SFN: May slow progression and support nerve fiber regeneration.

4. **Autoantibody Screening:** If GPCR autoantibodies contribute to autonomic SFN symptoms, immunoadsorption could be considered.

Implication for SFN: Identifies subset with autoantibody-mediated pathology amenable to specific intervention.

Certainty: **High** for shared metabolic vulnerability; **Low-Medium** for specific treatments (IVIG access limited, needs SFN-specific validation).

31.4 Promising But Requires Validation (Tier 3)

These conditions have theoretical mechanistic overlap with ME/CFS based on known pathophysiology. Translational findings are speculative pending direct research.

31.4.1 Autoimmune Conditions

Systemic autoimmune diseases (lupus, Sjögren's syndrome, rheumatoid arthritis, multiple sclerosis) share immune dysregulation features with ME/CFS.

Novel Translational Hypotheses from ME/CFS

1. **Plasma Cell Targeting Beyond B-Cell Depletion:**

ME/CFS Finding: Rituximab (anti-CD20, B-cell depletion) failed in Phase III trial, but daratumumab (anti-CD38, plasma cell depletion) showed 60% response rate in pilot.

Hypothesis for Autoimmune Diseases: Long-lived plasma cells in bone marrow and tissue sanctuaries produce autoantibodies resistant to B-cell depletion. Daratumumab could benefit autoimmune patients who failed rituximab.

Precedent: Multiple myeloma (plasma cell malignancy) responds to daratumumab. Autoimmune diseases may involve similar plasma cell-driven pathology.

Research Priority: Test daratumumab in rituximab-refractory lupus, Sjögren's, RA patients with persistent autoantibody production.

2. **GPCR Autoantibodies Causing Functional Symptoms:**

ME/CFS Finding: β 2-adrenergic, M3/M4 muscarinic receptor autoantibodies found in 29.5–91%, correlating with autonomic and cognitive symptoms.

Hypothesis for Autoimmune Diseases: Functional symptoms in autoimmune disease (fatigue, brain fog, autonomic dysfunction) may result from GPCR autoantibodies, not just tissue-specific autoantibodies.

Implication: Autoantibody screening could identify subset benefiting from immunoadsorption.

3. **Autoantibody-Monocyte Reprogramming (Hackel 2025):**

ME/CFS Finding: GPCR autoantibodies reprogram monocytes to produce inflammatory cytokines (MIP-1 δ , PDGF-BB, TGF- β 3).

Hypothesis for Autoimmune Diseases: Autoantibodies don't just block/activate receptors—they reprogram immune cells to produce persistent inflammation.

Implication: Combined autoantibody removal + JAK inhibitors (monocyte modulation) could be more effective than either alone.

4. **Low-Dose IL-2 for Regulatory T Cell Restoration:**

ME/CFS: Proposed but not yet tested systematically.

Precedent: Low-dose IL-2 (1 million IU) restored Treg function in SLE with clinical improvement.

Hypothesis: Treg dysfunction common to multiple autoimmune conditions; restoration could provide benefit across diseases.

Certainty: Low (theoretical, requires validation). **Highest priority:** Daratumumab in rituximab-refractory autoimmune disease.

31.4.2 Neurodegenerative Diseases

Alzheimer's disease, Parkinson's disease, and related dementias share neuroinflammation, oxidative stress, and protein aggregation pathology.

Novel Translational Hypotheses from ME/CFS

1. **Glymphatic Clearance Failure:**

ME/CFS Finding: Impaired slow-wave sleep and hypothesized glymphatic dysfunction preventing brain waste clearance.

Established in Neurodegenerative Disease: Glymphatic system clears amyloid- β and tau during sleep; dysfunction accelerates Alzheimer's pathology.

Translational Opportunity: Sleep architecture optimization (target slow-wave sleep), lateral sleeping position (enhances glymphatic flow), melatonin (circadian rhythm restoration).

Implication: Early intervention to restore glymphatic function could slow neurodegenerative progression.

2. **Microglial Activation and Neuroinflammation:**

ME/CFS Finding: PET imaging showed widespread microglial activation correlating with cognitive symptoms [73].

Established in Neurodegenerative Disease: Chronic microglial activation drives neurodegeneration.

Translational Opportunity: Low-dose naltrexone (TLR4 antagonism on microglia), omega-3 fatty acids (EPA/DHA 2–4g/day), curcumin (anti-inflammatory).

Implication: Microglial modulation could slow progression if initiated early.

3. **NAD⁺ Depletion and Mitochondrial Dysfunction:**

ME/CFS Finding: Metabolomic abnormalities, proposed NAD⁺ depletion contributing to energy failure.

Established in Neurodegenerative Disease: NAD⁺ declines with aging [410]; depletion impairs mitochondrial function, DNA repair (PARP), and sirtuin-mediated protein homeostasis [233].

Translational Opportunity: The only RCT-tested dose is NR 2000 mg/day (20-week intervention; cognitive benefit variable) [572]. Lower doses (300–1000 mg/day) are more commonly used in clinical practice but lack controlled evidence. See Chapter 22 for full dosing.

Implication: NAD⁺ restoration could support neuronal energy metabolism and protein quality control.

4. Oxidative Stress and Peroxynitrite Formation:

ME/CFS Finding: Oxidative stress markers elevated; peroxynitrite formation damaging cellular components [64].

Established in Neurodegenerative Disease: Oxidative damage to proteins, lipids, DNA accelerates neurodegeneration.

Translational Opportunity: Comprehensive antioxidant protocol (MitoQ, alpha-lipoic acid, N-Acetylcysteine (NAC), vitamin E, selenium).

Implication: Neuroprotection through oxidative stress reduction.

Certainty: Low-Medium (mechanisms plausible, requires prospective trials). **Highest priority:** Glymphatic optimization (sleep interventions) as preventive strategy.

31.4.3 Metabolic Syndrome and Type 2 Diabetes

Metabolic syndrome involves insulin resistance, dyslipidemia, hypertension, and chronic low-grade inflammation.

Translational Hypotheses

1. **Mitochondrial Dysfunction as Common Pathway:** Both ME/CFS and metabolic syndrome show impaired mitochondrial function, though through different mechanisms.

Translational Opportunity: Mitochondrial support (CoQ10, alpha-lipoic acid, carnitine) could improve insulin sensitivity and energy metabolism.

2. **Chronic Inflammation and Cytokine Dysregulation:** Elevated IL-6, TNF- α in both conditions.

Translational Opportunity: Anti-inflammatory approaches (omega-3, curcumin, Low-Dose Naltrexone (LDN)) could reduce inflammatory burden.

3. **NAD⁺ Depletion and Metabolic Dysfunction:** NAD⁺ depletion impairs sirtuin function, affecting metabolic regulation.

Translational Opportunity: NR/NMN supplementation could improve metabolic parameters.

Certainty: Low (theoretical overlap, requires metabolic syndrome-specific trials).

31.5 Key Translational Mechanisms

This section synthesizes the mechanisms with broadest applicability across multiple conditions.

31.5.1 Plasma Cell Autoimmunity (Daratumumab Target)

The discovery that daratumumab (plasma cell targeting) succeeds where rituximab (B-cell targeting) failed represents a paradigm shift in understanding autoantibody-mediated disease.

B-Cell vs. Plasma Cell Targeting: Critical Distinction

Rituximab (Anti-CD20) depletes CD20⁺ B cells in circulation and lymphoid organs:

- **ME/CFS Phase III trial:** No benefit over placebo (Fluge 2019)
- **Mechanism:** CD20 not expressed on plasma cells; long-lived plasma cells in bone marrow sanctuaries continue producing autoantibodies
- **Duration:** B-cell depletion lasts 6–12 months but autoantibody titers remain elevated

Daratumumab (Anti-CD38) targets CD38⁺ plasma cells:

- **ME/CFS pilot study:** 60% response rate (6/10) with marked improvement; SF-36 Physical Function increased from 25.9 to 55.0 ($p=0.002$)
- **Mechanism:** CD38 highly expressed on plasma cells; depletes long-lived plasma cells producing pathogenic autoantibodies
- **Response timing:** Gradual improvement over months as autoantibody titers decline
- **Precedent:** Proven in multiple myeloma (malignant plasma cells)

Why Plasma Cells Matter: Biological Basis

Plasma cells are terminally differentiated antibody-producing cells:

1. **Long-lived plasma cells (LLPCs)** reside in bone marrow survival niches, producing antibodies for years without requiring B-cell replenishment
2. **Short-lived plasma cells** in lymphoid tissues die within days-weeks and require continuous B-cell differentiation
3. **Rituximab depletes B cells** but doesn't affect LLPCs → autoantibody production continues
4. **Daratumumab depletes LLPCs** → autoantibody titers finally decline

Cross-Condition Implications

Conditions likely to benefit from plasma cell targeting:

1. **Rituximab-refractory autoimmune diseases:**
 - Systemic lupus erythematosus (SLE) with persistent anti-dsDNA antibodies
 - Sjögren's syndrome with anti-Ro/SSA persistence
 - Myasthenia gravis with anti-AChR antibodies
 - Neuromyelitis optica with anti-AQP4 antibodies
2. **Long COVID with elevated GPCR autoantibodies:** If autoantibodies drive persistent symptoms, plasma cell depletion could provide lasting benefit
3. **Post-infectious autoimmune syndromes:** PTLDS, Guillain-Barré syndrome with antibody-mediated pathology
4. **Any condition with documented pathogenic autoantibodies not responding to B-cell depletion**

Key principle: If rituximab failed despite clear autoantibody involvement, plasma cell targeting should be considered before concluding autoimmunity is not the mechanism.

Clinical Considerations

Advantages:

- Targets root cause (antibody-producing cells) rather than circulating antibodies
- Proven safety profile in multiple myeloma (extensive clinical experience)
- No serious adverse events in ME/CFS pilot

Limitations:

- Expensive (biologics cost \$10,000+/month typically)
- Requires autoantibody documentation for rational use
- Immunosuppression: infection monitoring required
- Gradual response (months, not weeks)

Research priority: High. Phase II trials in rituximab-refractory autoimmune disease justified by ME/CFS pilot data and biological rationale.

31.5.2 GPCR Autoantibodies

G-protein-coupled receptor (GPCR) autoantibodies represent a mechanism explaining “functional” symptoms across multiple conditions previously dismissed as psychosomatic.

GPCR Autoantibodies in ME/CFS

Prevalence:

- **β 2-adrenergic receptor:** 29.5–91% of ME/CFS patients (prevalence varies by assay, cut-off)
- **M3/M4 muscarinic receptors:** Elevated in subset
- **α 1-adrenergic receptor:** May contribute to vascular dysfunction

Functional effects:

- **Not simple blockade:** Autoantibodies can activate, block, or modulate receptor function
- **Downstream signaling alterations:** Chronic receptor stimulation/blockade → desensitization, internalization
- **Cellular reprogramming:** Hackel 2025 showed autoantibodies reprogram monocytes to produce inflammatory cytokines

Clinical Manifestations by Receptor Type

β 2-Adrenergic receptor autoantibodies:

- Autonomic dysfunction: Orthostatic intolerance, tachycardia
- Vascular effects: Impaired vasodilation, blood pooling
- Metabolic effects: Reduced Na^+/K^+ -ATPase → intracellular sodium accumulation
- Mast cell effects: Favors degranulation (worsens MCAS)

M3/M4 Muscarinic receptor autoantibodies:

- Cognitive dysfunction: Cholinergic system disruption affecting memory, attention
- Autonomic effects: Altered parasympathetic function
- GI symptoms: Dysmotility from enteric nervous system dysfunction

α 1-Adrenergic receptor autoantibodies:

- Vascular dysfunction: Impaired vasoconstriction
- Orthostatic hypotension: Inadequate compensatory response to standing

Cross-Condition Implications

POTS: β 2-AR autoantibodies could explain tachycardia, exercise intolerance, and autonomic failure in subset of POTS patients

Long COVID: GPCR autoantibodies documented; may drive persistent autonomic and cognitive symptoms post-infection

Autoimmune diseases with “functional” symptoms: Fatigue, brain fog, autonomic dysfunction in lupus, Sjögren’s may reflect GPCR autoantibodies, not just tissue damage

Post-infectious syndromes: PTLDS, post-viral fatigue may involve molecular mimicry triggering GPCR autoantibodies

Diagnostic and Therapeutic Implications

Testing:

- CellTrend assay (commercial): Measures functional effects on cell lines
- ELISA-based assays: Detect binding autoantibodies
- **Challenge:** Assay standardization, cutoff values not established

Treatments if elevated:

1. **Immunoabsorption:** 70% response rate in ME/CFS with elevated β 2-AR autoantibodies; removes autoantibodies selectively
2. **Plasma cell targeting:** Daratumumab prevents autoantibody regeneration
3. **BC007 (DNA aptamer):** Neutralizes GPCR autoantibodies; dramatic case report in Long COVID
4. **IVIg:** May provide competing antibodies, immune modulation

Research priority: High. Establish validated assays, define pathogenic thresholds, conduct controlled trials of autoantibody-directed therapies.

31.5.3 Vascular-Immune-Energy Triad

The Heng 2025 multi-omics study identified coordinated dysfunction across three systems, achieving 91% diagnostic accuracy with a 7-biomarker panel.

The 7-Biomarker Panel

Immune markers:

- IL-8 (elevated): Neutrophil chemoattractant, inflammation
- TNF- α (elevated): Pro-inflammatory cytokine

Vascular markers:

- von Willebrand Factor (VWF, elevated): Endothelial activation/damage
- Fibronectin (elevated): Extracellular matrix protein, vascular remodeling
- Thrombospondin (elevated): Anti-angiogenic, endothelial stress

Metabolic markers:

- Lactate (elevated): Anaerobic metabolism, mitochondrial dysfunction
- Pyruvate (ratio altered): Impaired oxidative phosphorylation

Why the Triad Matters: Systems Integration

Not three independent problems—coordinated dysfunction:

1. **Vascular dysfunction** → impaired tissue perfusion → hypoxia → mitochondrial stress
2. **Immune activation** → cytokines (IL-6, TNF- α) → endothelial activation → vascular dysfunction
3. **Mitochondrial dysfunction** → ATP depletion → immune cell dysfunction → altered cytokine production
4. **Positive feedback loops**: Each system's dysfunction worsens the others

Clinical Implication: Why Single-Target Treatments Fail

Targeting only immune system (e.g., anti-cytokine therapy):

- Addresses inflammation but not vascular dysfunction or energy deficit
- Vascular and metabolic problems persist → immune activation returns

Targeting only mitochondria (e.g., CoQ10 alone):

- Improves energy metabolism but not immune activation or vascular dysfunction
- Persistent inflammation and hypoperfusion limit mitochondrial recovery

Targeting only vascular system (e.g., vasodilators):

- Improves perfusion but not immune dysfunction or cellular energy production
- Inflammatory and metabolic problems persist

Triple-Target Treatment Strategy

Vascular support:

- L-citrulline/arginine (NO precursors): 3–6g/day
- Omega-3 fatty acids (endothelial function): EPA/DHA 2–4g/day
- Statins (endothelial protection): If indicated

Immune modulation:

- Low-dose naltrexone (neuroinflammation): 3–4.5 mg
- Curcumin (anti-inflammatory): 500–1000 mg bioavailable form
- Omega-3 (anti-inflammatory): Overlaps with vascular support

Metabolic support:

- Comprehensive mitochondrial stack—including CoQ10, NADH, NAD⁺ precursors (NR), D-ribose, ALCAR, and B vitamins (see Section 31.7.1 for full protocol and Chapter 22 for dosing)

Rationale: These three systems are hypothesized to interact; addressing multiple targets simultaneously may provide greater benefit than isolated interventions, though this has not been tested in controlled comparisons.

Cross-Condition Applicability

High relevance: Any condition showing:

- Elevated inflammatory markers (IL-6, TNF- α , CRP)
- Vascular dysfunction (impaired FMD, elevated VWF)
- Metabolic abnormalities (elevated lactate, mitochondrial dysfunction)

Examples:

- Long COVID (documented triad dysfunction)
- Diabetes complications (vascular + metabolic + inflammation)
- Cardiovascular disease (all three systems involved)
- Neurodegenerative disease (neuroinflammation + vascular + energy failure)

Research priority: Validate 7-biomarker panel across conditions; test triple-target protocol in controlled trials.

31.5.4 WASF3/ER Stress → Mitochondrial Dysfunction

The WASF3 pathway represents a druggable target linking ER stress to mitochondrial dysfunction.

The Mechanism

1. **Trigger:** Viral infection, inflammatory stress, or other cellular stress induces ER stress
2. **WASF3 induction:** ER stress response upregulates WASF3 expression
3. **Mitochondrial supercomplex disruption:** WASF3 interferes with respiratory chain supercomplex assembly
4. **Complex IV impairment:** Particularly affects cytochrome c oxidase (Complex IV)
5. **ATP depletion:** Impaired oxidative phosphorylation reduces energy production
6. **Oxidative stress:** Dysfunctional electron transport chain generates ROS
7. **Vicious cycle:** ROS → more ER stress → more WASF3 → worse mitochondrial function

Why This Pathway Matters

Explains post-infectious onset:

- Viral infection triggers ER stress
- WASF3 induction persists after viral clearance
- Mitochondrial dysfunction becomes self-perpetuating

Explains multi-system involvement:

- High-energy tissues (brain, muscle, heart) most affected
- ER stress is universal cellular response
- Pattern matches ME/CFS symptom distribution

Provides therapeutic targets:

- ER stress inhibitors (experimental)
- WASF3 inhibition (research target)
- Mitochondrial protection downstream of WASF3

Cross-Condition Implications

Primary mitochondrial disorders: If WASF3 induction occurs secondary to mitochondrial dysfunction, inhibiting ER stress could break vicious cycle

Neurodegenerative diseases: ER stress and protein misfolding central to Alzheimer's, Parkinson's; WASF3 pathway could contribute to energy failure

Cancer-related fatigue: Chemotherapy induces ER stress; WASF3 pathway could mediate persistent fatigue post-treatment

Sepsis recovery: Severe infection triggers ER stress; WASF3-mediated mitochondrial dysfunction could explain prolonged weakness

Therapeutic Strategies

ER stress modulators (experimental):

- Tauroursodeoxycholic acid (TUDCA): Chemical chaperone reducing ER stress
- 4-Phenylbutyric acid (4-PBA): ER stress inhibitor
- **Status:** Used in primary biliary cirrhosis; ME/CFS testing needed

Downstream mitochondrial protection:

- MitoQ: Mitochondria-targeted antioxidant
- Alpha-lipoic acid: Mitochondrial antioxidant, ER stress reducer

- N-Acetylcysteine (NAC): Precursor to glutathione, reduces oxidative stress

Supporting Complex IV function:

- Copper supplementation (if deficient): Complex IV cofactor
- CoQ10: Electron carrier supporting Complex IV

Research priority: Medium-High. WASF3 pathway newly identified; validation and therapeutic targeting needed.

31.5.5 NAD⁺ Depletion

NAD⁺ (nicotinamide adenine dinucleotide) is a universal cofactor affecting mitochondria, DNA repair, sirtuins, and circadian rhythms. Depletion represents a unifying mechanism across aging-related and chronic diseases.

NAD⁺ Functions in Cellular Metabolism

Mitochondrial function:

- Essential cofactor for electron transport chain (Complexes I, III)
- NAD⁺/NADH ratio determines oxidative vs. reductive state
- Depletion impairs ATP production directly

DNA repair:

- PARP (poly-ADP-ribose polymerase) consumes NAD⁺ for DNA repair
- Chronic DNA damage (oxidative stress, inflammation) depletes NAD⁺ pools
- NAD⁺ depletion → impaired DNA repair → cellular dysfunction

Sirtuins (protein deacetylases):

- SIRT1-7 require NAD⁺ for activity
- Regulate protein homeostasis, autophagy, mitochondrial biogenesis
- NAD⁺ depletion → impaired protein quality control

Circadian rhythms:

- SIRT1 regulates CLOCK/BMAL1 circadian machinery
- NAD⁺ levels oscillate with circadian rhythm
- Depletion disrupts sleep-wake cycles

Evidence for NAD⁺ Depletion in ME/CFS

Metabolomic abnormalities: Tryptophan-NAD⁺ pathway dysregulation [12]

PARP activation: Oxidative stress and DNA damage trigger PARP, consuming NAD⁺ [233]

Chronic inflammation: Inflammatory cytokines induce cellular stress → PARP activation → NAD⁺ consumption [233] (proposed mechanism; direct ME/CFS data limited)

Clinical trial: 2025 Long COVID RCT (n=58) showed NR 2000 mg/day increased NAD⁺ levels 2.6–3.1×; cognitive benefits were variable [572]

Cross-Condition Implications

Universal mechanism affecting:

- **Aging-related decline:** NAD⁺ in human tissue declines significantly with age (approximately 50% by age 50 in skin tissue measurements) [410]
- **Neurodegenerative diseases:** Impaired mitochondrial function, protein homeostasis [233]
- **Metabolic syndrome:** Insulin resistance proposed to be linked to NAD⁺ depletion (mechanistic evidence from animal models [645])
- **Cancer-related fatigue:** Chemotherapy and radiation are proposed to deplete NAD⁺ (via PARP activation and direct mitochondrial toxicity) [233]
- **Chronic inflammatory conditions:** PARP activation consumes NAD⁺ [233]
- **Mitochondrial disorders:** Primary dysfunction worsened by NAD⁺ depletion

NAD⁺ Restoration Strategies

Nicotinamide riboside (NR):

- Dose: 1000–2000 mg/day
- Duration: >10 weeks required for benefit
- Mechanism: Converted to NAD⁺ via salvage pathway
- Evidence: Long COVID RCT positive; ME/CFS trials ongoing

Nicotinamide mononucleotide (NMN):

- Dose: 1000–2000 mg/day
- Mechanism: One step closer to NAD⁺ than NR
- Evidence: Animal studies strong; human trials emerging

Niacin (nicotinic acid):

- Dose: 500–1000 mg/day (extended-release to minimize flushing)
- Mechanism: Converts to NAD⁺ via Preiss-Handler pathway

- Trade-off: Cheaper but flushing limits tolerability

Optimize NAD⁺ consumption:

- Reduce oxidative stress (antioxidants) → less PARP activation
- Anti-inflammatory approaches → less cellular stress
- Sleep optimization → restore circadian NAD⁺ oscillation

Research priority: High. NAD⁺ restoration is safe, biologically plausible, and shows promise across multiple conditions.

31.5.6 Glymphatic Clearance Failure

The glymphatic system is the brain's waste clearance system, active primarily during slow-wave sleep. Dysfunction allows toxic metabolites to accumulate, driving neurodegeneration and cognitive impairment.

Glymphatic System: Discovery and Function

Discovery (Nedergaard 2012):

- Brain lacks lymphatic vessels; alternative clearance mechanism identified
- Cerebrospinal fluid (CSF) flows along paravascular spaces
- Interstitial fluid with metabolic waste is cleared into CSF
- Most active during slow-wave (deep) sleep

What it clears:

- Amyloid- β (accumulates in Alzheimer's disease)
- Tau protein (forms tangles in neurodegeneration)
- Metabolic waste products
- Inflammatory mediators

Why sleep matters:

- During wakefulness: Brain cells expanded, limited interstitial space
- During slow-wave sleep: Brain cells shrink 60%, interstitial space increases
- This expansion allows CSF influx and waste clearance
- Disrupted sleep → impaired clearance → toxic accumulation

Glymphatic Dysfunction in ME/CFS

Evidence:

- **Non-restorative sleep:** Diagnostic criterion; patients wake unrefreshed
- **Alpha-delta sleep pattern:** Alpha waves intrude into delta (slow-wave) sleep
- **Reduced slow-wave sleep:** Impairs glymphatic clearance
- **Cognitive dysfunction:** Brain fog may reflect metabolite accumulation
- **Cranio cervical junction issues in subset:** May impair CSF flow

Hypothesis: Impaired glymphatic clearance allows neuroinflammatory mediators and metabolic waste to accumulate, perpetuating cognitive dysfunction and neuroinflammation.

Cross-Condition Implications: Neurodegenerative Diseases

Alzheimer's disease:

- Amyloid- β accumulation directly linked to impaired glymphatic clearance
- Sleep disruption accelerates amyloid deposition
- Poor sleep quality predicts Alzheimer's risk

Parkinson's disease:

- Alpha-synuclein (forms Lewy bodies) cleared by glymphatic system
- Sleep disorders common in early Parkinson's
- REM sleep behavior disorder precedes motor symptoms by years

Traumatic brain injury:

- TBI impairs glymphatic function
- Sleep disruption post-TBI worsens outcomes
- Early sleep optimization may improve recovery

Migraine:

- Glymphatic dysfunction may allow inflammatory mediator accumulation
- Poor sleep triggers migraines
- Sleep optimization reduces migraine frequency

Therapeutic Strategies to Optimize Glymphatic Function

Sleep architecture optimization:

1. Target slow-wave sleep:

- Low-dose trazodone (25–50 mg): Increases slow-wave sleep without hangover
- Avoid benzodiazepines: Suppress slow-wave sleep
- Sleep hygiene: Dark, cool room (60–67°F optimal)

2. Melatonin:

- Dose: 0.5–3 mg (lower often more effective than higher)
- Timing: 1–2 hours before desired sleep time
- Regulates circadian rhythm, antioxidant effects

3. Magnesium glycinate:

- Dose: 400–800 mg at bedtime
- Promotes relaxation, GABA-ergic effects
- Glycinate form best absorbed, least laxative effect

Sleep position:

- **Lateral (side) sleeping:** Most effective for glymphatic clearance (animal studies)
- Supine (back) sleeping: Least effective
- Mechanism: CSF flow enhanced in lateral position

Craniocervical optimization:

- If craniocervical instability (CCI) or Chiari malformation present: Surgical evaluation
- Proper pillow support: Maintains cervical alignment
- Physical therapy: Addresses cervical spine issues

Circadian rhythm entrainment:

- Morning bright light exposure (10,000 lux, 30 min)
- Evening dim light (minimize blue light 2 hours before bed)
- Consistent sleep-wake times (even weekends)

Preventive strategy:

- **Neurodegenerative disease prevention:** Optimize glymphatic function before amyloid/tau accumulation
- **Post-TBI recovery:** Aggressive sleep optimization may prevent chronic sequelae
- **Migraine prophylaxis:** Sleep architecture improvement reduces attack frequency

Research priority: High. Sleep optimization is low-risk, low-cost, and has strong biological rationale for neuroprotection.

31.6 Research Priorities and Future Directions

31.6.1 Cross-Condition Mechanism Validation

Which ME/CFS mechanisms need testing in which conditions:

Table 31.3: Research Priorities: Mechanisms × Conditions

Mechanism	Priority Conditions for Testing
Plasma cell autoimmunity (daratumumab)	Long COVID, PTLDS, autoimmune diseases where rituximab failed
β 2-receptor desensitization	EDS-POTS-MCAS, dysautonomia, Long COVID
Tetrahydrobiopterin (BH4) dysregulation	EDS with OI, POTS, dysautonomia, migraine
WASF3/ER stress pathway	Primary mitochondrial disorders, metabolic myopathies
NAD ⁺ depletion	Cancer-related fatigue, aging-related decline, neurodegenerative disease
Glymphatic clearance failure	Alzheimer's, Parkinson's, migraine, TBI
GPCR autoantibody-monocyte reprogramming	Long COVID, autoimmune conditions with functional symptoms

31.6.2 Biomarker Validation Across Conditions

The Heng 2025 7-biomarker panel (91% diagnostic accuracy in ME/CFS) includes:

- **Immune:** IL-8, TNF- α
- **Vascular:** VWF, fibronectin, thrombospondin
- **Metabolic:** Lactate, pyruvate

Research priority: Validate this panel in Long COVID, fibromyalgia, EDS-POTS-MCAS, and other conditions with multi-system dysfunction.

31.7 Universal Treatment Protocols

ME/CFS research has identified treatment strategies with potential applicability across the full spectrum of post-viral, autoimmune, mitochondrial, and dysautonomic conditions. The pro-

protocols below represent evidence-based approaches that address fundamental shared pathophysiology rather than condition-specific symptoms.

Critical caveat: These protocols are derived from ME/CFS research and clinical experience. Direct application to other conditions requires:

1. Physician supervision and approval
2. Condition-specific contraindication screening
3. Individualized dosing based on severity and comorbidities
4. Monitoring for adverse effects
5. Recognition that evidence quality varies by condition

31.7.1 Comprehensive Mitochondrial Support

Rationale and Mechanism

Mitochondrial dysfunction is documented in ME/CFS [2, 64]; analogous findings have been reported in Long COVID [572], fibromyalgia [569], and other chronic fatigue conditions [646]. The comprehensive mitochondrial support stack addresses multiple points of failure:

1. **Electron transport chain support:** CoQ10 (shuttles electrons from Complex I and Complex II to Complex III), NADH (electron donor to Complex I)
2. **ATP synthesis cofactors:** D-ribose (substrate), magnesium (ATPase cofactor)
3. **Oxidative stress protection:** Alpha-lipoic acid (mitochondrial antioxidant; see Chapter 22). Optional additions: vitamin E (membrane antioxidant) and phosphatidylcholine (membrane repair)—these are not reviewed individually in this text and lack ME/CFS-specific evidence; inclusion is based on theoretical rationale only
4. **Fatty acid transport:** Acetyl-L-carnitine (mitochondrial fatty acid oxidation)
5. **NAD⁺ restoration:** Nicotinamide riboside (NR) or nicotinamide mononucleotide (NMN)
6. **Citric acid cycle support:** B-complex vitamins (B1, B2, B3, B5)

Evidence base:

- ME/CFS: CoQ10 + NADH demonstrated significant improvements in cognitive fatigue, overall fatigue, and sleep parameters in the largest RCT (Castro-Marrero 2021, n=207) [7]; an earlier smaller trial (Castro-Marrero 2015, n=73) showed improvements in fatigue and biochemical parameters [568]
- Long COVID: NR 2000 mg/day increased NAD⁺ levels 2.6–3.1 fold (Wu 2025, n=58) [572]
- Fibromyalgia: CoQ10 300 mg/day reduced pain and fatigue (Cordero 2013, n=20) [569]
- Mitochondrial disorders: Established therapeutic role for CoQ10, ribose, carnitine [646]

Protocol Details

Core stack (evidence-based dosing):

- **Coenzyme Q10:** 100–300 mg/day typical (ubiquinol form preferred; RCT dose 200 mg; up to 400–600 mg in studies)
- **D-ribose:** 5g TID (15g/day total), dissolved in water, taken with meals
- **NADH:** 10–20mg/day, sublingual or enteric-coated
- **Acetyl-L-carnitine:** 500–2000 mg/day, divided doses (start low; see Chapter 22)
- **Alpha-lipoic acid:** 300–600 mg/day (R-lipoic acid form preferred; start at 100–150 mg in ME/CFS and titrate—sensitive patients may experience paradoxical worsening (increased fatigue, malaise, or gastrointestinal symptoms) at higher initial doses; clinical observation)
- **Magnesium glycinate:** 200–600 mg/day elemental (divided doses to avoid diarrhea; see Chapter 22)
- **B-complex:** High-potency formulation with methylated forms (B12 as methylcobalamin)

Advanced additions:

- **Nicotinamide riboside (NR):** 300–1000 mg/day (morning dosing; the only RCT-tested dose is 2000 mg/day [572]; see Chapter 22)
- **Pyrroloquinoline quinone (PQQ):** 10–20 mg/day (mitochondrial biogenesis)
- **Creatine monohydrate:** 3–5 g/day (ATP buffering, cognitive support)

Implementation Strategy

1. **Titration:** Start with 25–50% of target doses, increase weekly (clinical experience; no controlled titration protocol data). Introduce supplements one at a time per Chapter 22, Section 22.7; apply this titration to each supplement individually as it is added
2. **Timing:** Split doses throughout day; CoQ10 and fat-soluble nutrients with meals
3. **Response monitoring:** Track energy levels, cognitive function, post-exertional symptoms
4. **Minimum trial duration:** 8–12 weeks per individual supplement (mitochondrial adaptations require time). With sequential introduction (1–2 weeks between additions per Chapter 22), the full stack requires substantially longer
5. **Responder identification:** Responder rates have not been established in controlled trials; clinical experience suggests some patients improve, but no reliable proportion can be cited. Non-responders may have different rate-limiting pathology

Safety considerations:

- **CoQ10:** May reduce warfarin effectiveness—the mechanism may involve enhanced CYP450-mediated warfarin hydroxylation [578]; a structural similarity to vitamin K has also been proposed but is less well-documented. Monitor INR closely with physician guidance; see Chapter 22, Section 22.2.8

- Alpha-lipoic acid: Monitor glucose in diabetics (insulin-sensitizing effect)
- Carnitine: May theoretically lower seizure threshold in pre-existing epilepsy; consult physician (no controlled data; clinical caution). Also inhibits thyroid hormone nuclear uptake—patients on thyroid replacement should monitor thyroid function [580]; see Chapter 22, Section 22.2.8 for additional drug interactions
- Magnesium: Dose-dependent diarrhea; reduce dose or switch to magnesium threonate
- NR/NMN: Theoretical concern about NAD⁺ promoting tumor growth—rapidly proliferating cancer cells may exploit elevated NAD⁺ for energy and DNA repair [574]. Avoid in active cancer; consult oncologist before use in patients with cancer history

Cross-Condition Applications

High priority for mitochondrial support:

- Long COVID with persistent fatigue
- Cancer-related fatigue (post-treatment, not during active treatment; see NR/NMN safety note above)
- Fibromyalgia with exercise intolerance
- POTS with fatigue predominance
- Primary mitochondrial disorders (adjunct to genetic-specific therapy)
- Neurodegenerative diseases (Parkinson's, early Alzheimer's)

Lower priority (less evidence):

- Autoimmune conditions without fatigue
- MCAS (unless significant fatigue component)
- Metabolic syndrome (focus on lifestyle first)

31.7.2 Autonomic-Catecholamine Restoration

Rationale and Mechanism

Catecholamine dysfunction affects POTS, dysautonomia, ME/CFS with orthostatic intolerance, and conditions with autonomic neuropathy [172]. The restoration protocol addresses:

1. **Substrate availability:** L-tyrosine (precursor for dopamine → norepinephrine → epinephrine)
2. **Cofactor sufficiency:** Tetrahydrobiopterin (BH4), vitamin C, copper
3. **Methylation support:** S-AdoMet, methylated B-vitamins (for catecholamine metabolism)
4. **Adrenal support:** Vitamin B5 (pantothenic acid), adaptogenic herbs

Evidence base:

- POTS: L-tyrosine improved orthostatic tolerance (case reports, small studies)

- ME/CFS: BH4 elevation correlates with orthostatic intolerance [644]
- Dysautonomia: Vitamin C supports catecholamine synthesis [647]
- Adrenal insufficiency: B5 deficiency impairs cortisol synthesis

Protocol Details

Core interventions:

- **L-tyrosine:** 500–1500mg/day, morning and midday (empty stomach for absorption)
- **Vitamin C:** 1000–2000mg/day (cofactor for dopamine β -hydroxylase)
- **Vitamin B6 (P5P):** 50–100mg/day (cofactor for aromatic L-amino acid decarboxylase)
- **Methylfolate:** 1–5mg/day (methylation pathway support)
- **Methylcobalamin (B12):** 1000–5000mcg/day sublingual
- **Pantothenic acid (B5):** 500–1000mg/day (adrenal cortex support)

Advanced additions:

- **Sapropterin (Tetrahydrobiopterin, BH4):** 5–10mg/kg/day (prescription; for documented BH4 deficiency)
- **SAMe:** 400–800mg/day (methylation, catecholamine metabolism)
- **Copper:** 2mg/day (cofactor for dopamine β -hydroxylase; only if deficient)
- **Adaptogens:** Rhodiola rosea 200–400mg, ashwagandha 300–600mg (adrenal support)

Implementation Strategy

1. **Baseline assessment:** Orthostatic vital signs, symptom severity scores
2. **Tyrosine titration:** Start 500mg/day, increase to 1500mg over 2 weeks
3. **Timing:** Morning and early afternoon (avoid evening due to potential sleep disruption)
4. **Response monitoring:** Orthostatic tolerance, brain fog, energy, heart rate variability
5. **Trial duration:** 4–8 weeks minimum

Safety considerations:

- **Contraindications:** Hyperthyroidism (tyrosine is thyroid hormone precursor), MAO inhibitors
- **Warnings:** May worsen anxiety or insomnia in susceptible individuals
- **Monitoring:** Blood pressure (may increase in some patients)
- **Drug interactions:** Levodopa (competes for absorption), thyroid medications

Cross-Condition Applications

High priority:

- POTS with low norepinephrine or hyperadrenergic subtype
- ME/CFS with orthostatic intolerance
- Dysautonomia (diabetic, autoimmune, idiopathic)
- Long COVID with autonomic dysfunction
- EDS with POTS

Moderate priority:

- Fibromyalgia with brain fog
- Neurodegenerative diseases (Parkinson's - with caution due to levodopa interactions)

31.7.3 Mast Cell Stabilization

Rationale and Mechanism

Mast cell activation contributes to ME/CFS, MCAS, EDS, POTS, Long COVID, and potentially fibromyalgia [235, 560]. Stabilization strategies target:

1. **Histamine blockade:** H1 + H2 receptor antagonism (dual pathway)
2. **Membrane stabilization:** Cromolyn sodium, quercetin
3. **PAF inhibition:** Rupatadine (H1 + PAF dual action)
4. **Mediator degradation:** DAO supplementation for histamine
5. **Trigger avoidance:** Dietary histamine, stress, temperature extremes

Evidence base:

- MCAS: H1+H2 combination superior to monotherapy [560]
- ME/CFS: Rupatadine improved fatigue and orthostatic symptoms (clinical observations)
- EDS: High prevalence of mast cell activation; stabilization improves GI symptoms [565]
- Long COVID: Antihistamines improved symptoms in observational studies

Protocol Details

First-line (H1 + H2 combination):

- **H1 antagonist:** Cetirizine 10–20mg/day OR loratadine 10–20mg/day OR fexofenadine 180mg/day
- **H2 antagonist:** Famotidine 20–40mg BID OR ranitidine 150mg BID (if available)
- **Rationale:** Dual blockade addresses both H1 (allergic symptoms) and H2 (GI, vascular) pathways

Advanced interventions:

- **Rupatadine:** 10–20mg/day (H1 + PAF inhibition; superior to single-mechanism antihistamines)
- **Cromolyn sodium:** 200mg QID oral (membrane stabilizer; prescription)
- **Ketotifen:** 1–4mg/day (potent stabilizer; may cause sedation)
- **Quercetin:** 500–1000mg BID (natural flavonoid stabilizer)
- **DAO supplementation:** 10,000–20,000 HDU before meals (histamine degradation)
- **Vitamin C:** 1000mg BID (natural antihistamine, mast cell stabilizer)

Dietary modifications:

- Low-histamine diet (avoid aged cheeses, fermented foods, alcohol, leftover meat)
- DAO-rich foods (fresh meat, eggs)
- Avoid histamine liberators (citrus, strawberries, tomatoes, chocolate)
- Trial duration: 4–6 weeks

Implementation Strategy

1. **Start conservative:** H1 + H2 combination for 2–4 weeks
2. **Add stabilizers:** If partial response, add quercetin or cromolyn
3. **Consider rupatadine:** If standard antihistamines insufficient
4. **Dietary trial:** Implement low-histamine diet concurrently
5. **Response monitoring:** Symptom diary (flushing, GI symptoms, orthostatic tolerance, brain fog)

Safety considerations:

- **First-generation antihistamines:** Avoid (diphenhydramine, hydroxyzine) due to anticholinergic effects and cognitive impairment
- **Ketotifen:** Significant sedation; start low (0.5–1mg) and titrate
- **Cromolyn:** GI side effects common; take 15–30 minutes before meals
- **Drug interactions:** H2 blockers may affect absorption of pH-dependent medications

Cross-Condition Applications

High priority:

- MCAS (primary indication)
- EDS with MCAS features
- POTS with flushing or GI symptoms
- ME/CFS with orthostatic intolerance and MCAS overlap
- Long COVID with allergic/inflammatory symptoms

Moderate priority:

- Fibromyalgia with food sensitivities
- Migraine with histamine trigger pattern

31.7.4 Neuroinflammation Reduction**Rationale and Mechanism**

Neuroinflammation contributes to ME/CFS, Long COVID, fibromyalgia, neurodegenerative diseases, and potentially autoimmune conditions [648]. Reduction strategies target:

1. **Microglial modulation:** Low-dose naltrexone (LDN)
2. **Lipid mediators:** Omega-3 fatty acids (EPA/DHA)
3. **NF- κ B inhibition:** Curcumin, resveratrol
4. **Vagal stimulation:** Non-invasive VNS, deep breathing
5. **BBB protection:** Luteolin, apigenin

Evidence base:

- ME/CFS: LDN 4.5mg improved pain and fatigue in 65% (Younger 2013, n=80) [468]
- Long COVID: Omega-3 2g/day reduced inflammatory markers (pilot data)
- Fibromyalgia: LDN reduced pain scores by 30% (Parkitny 2014, meta-analysis) [649]
- Alzheimer's: Curcumin reduced amyloid burden (preclinical, limited human data)

Protocol Details**Core interventions:**

- **Low-dose naltrexone (LDN):** 1.5–4.5mg at bedtime (prescription; compounded)
 - Start 1.5mg, increase by 1mg every 2 weeks to 4.5mg
 - Mechanism: Transient opioid receptor blockade → increased endorphin production, microglial modulation
 - Response time: 8–12 weeks for full effect
- **Omega-3 fatty acids:** 2–4g/day combined EPA+DHA
 - High EPA:DHA ratio (2:1 or 3:1) preferred for anti-inflammatory effect
 - Triglyceride form better absorbed than ethyl ester
- **Curcumin:** 500–1000mg BID (with black pepper/piperine for bioavailability)
 - Use liposomal or phytosome formulations for enhanced absorption

Advanced additions:

- **Luteolin:** 100–200mg/day (microglial inhibitor, BBB permeable)

- **Resveratrol:** 200–500mg/day (SIRT1 activator, anti-inflammatory)
- **Palmitoylethanolamide (PEA):** 600–1200mg/day (endocannabinoid modulator)
- **Alpha-lipoic acid:** 600mg/day (NF- κ B inhibition, crosses BBB)

Non-pharmacological:

- **Vagal nerve stimulation:** Non-invasive transcutaneous VNS devices (gammaCore, Parasym)
- **Breathing exercises:** Slow diaphragmatic breathing (5–6 breaths/min) for 10–20 min BID
- **Cold exposure:** Brief cold showers (vagal activation, anti-inflammatory)

Implementation Strategy

1. **Start with LDN:** Highest evidence base; titrate slowly to minimize side effects
2. **Add omega-3:** Immediate start (safe, broad benefits)
3. **Layer curcumin:** After 4 weeks if partial response
4. **Consider advanced agents:** If inadequate response after 8–12 weeks
5. **Response monitoring:** Pain scores, cognitive function, sleep quality, overall well-being

Safety considerations:

- **LDN contraindications:** Active opioid use (precipitates withdrawal), liver disease
- **LDN side effects:** Vivid dreams (dose-dependent), insomnia (switch to morning dosing)
- **Omega-3:** Bleeding risk at high doses (>3g/day); caution with anticoagulants
- **Curcumin:** May potentiate anticoagulants; GI upset in sensitive individuals
- **Resveratrol:** May interact with blood thinners

Cross-Condition Applications

High priority:

- ME/CFS with pain and cognitive dysfunction
- Fibromyalgia (LDN well-established)
- Long COVID with neurological symptoms
- Autoimmune conditions with CNS involvement (MS, lupus cerebritis)

Moderate priority:

- Neurodegenerative diseases (adjunct therapy)
- POTS with brain fog
- Cancer-related fatigue (LDN may modulate cancer-related inflammation)

31.7.5 Energy Envelope Management (Pacing)

Rationale and Mechanism

Energy envelope management (pacing) prevents post-exertional symptom exacerbation across ME/CFS, Long COVID, POTS, fibromyalgia, and any condition with exercise intolerance [650]. The approach addresses:

1. **Anaerobic threshold violation:** Staying within aerobic capacity prevents PEM
2. **Boom-bust cycles:** Consistent activity prevents overexertion followed by crashes
3. **Circadian optimization:** Aligning activity with natural energy fluctuations
4. **Recovery prioritization:** Adequate rest prevents accumulated deficits

Evidence base:

- ME/CFS: Pacing superior to graded exercise therapy (PACE trial reanalysis) [506]
- Long COVID: Activity management improved function vs. push-through approach
- POTS: Heart rate-based exercise limits improved outcomes vs. standard exercise
- Fibromyalgia: Pacing reduced pain flares and improved consistency

Protocol Details

Core principles:

1. **Establish baseline:** Identify current sustainable activity level (what you can do consistently without symptom worsening)
2. **Stay within envelope:** Operate at 70–80% of baseline on average (leave margin for fluctuations)
3. **Monitor intensity:** Use heart rate, perceived exertion, symptom tracking
4. **Avoid boom-bust:** Resist temptation to "cash in" on good days with excessive activity
5. **Gradual expansion:** Increase activity by 5–10% every 2–4 weeks if sustained improvement

Heart rate monitoring approach:

- **Calculate anaerobic threshold (AT):**
 - Conservative method: $(220 - \text{age}) \times 0.6$
 - Workwell Foundation formula: $(220 - \text{age}) \times 0.55$ for severe ME/CFS
 - 2-day CPET testing (gold standard but not widely available)
- **Activity limits:** Keep heart rate below AT during all activities
- **Wearable devices:** Continuous HR monitors (Polar, Garmin, Apple Watch) with alerts
- **Rest breaks:** When approaching AT, stop activity immediately and rest until HR normalizes

Activity structuring:

- **Time-based limits:** Cap activities at 10–15 minute intervals with rest breaks
- **Task modification:** Break complex tasks into smaller components
- **Energy accounting:** Track "energy expenditure" throughout day
- **Pre-planning:** Schedule high-priority activities during peak energy windows
- **Rest is active treatment:** Schedule rest periods, not just "what's left over"

Symptom monitoring:

- Daily symptom diary (fatigue, pain, cognitive function, PEM severity)
- Activity log (duration, intensity, heart rate data)
- Identify personal triggers and patterns
- Adjust envelope boundaries based on data, not motivation

Implementation Strategy

1. **Assessment phase** (2–4 weeks):
 - Track current activity and symptoms without modification
 - Identify baseline capacity and PEM triggers
 - Calculate heart rate threshold
2. **Stabilization phase** (4–8 weeks):
 - Reduce activity to 70–80% of baseline
 - Implement heart rate monitoring
 - Eliminate boom-bust cycles
 - Goal: Consistent symptom stability
3. **Expansion phase** (ongoing):
 - Increase activity by 5–10% every 2–4 weeks
 - Monitor for PEM after each increase
 - Pull back immediately if symptoms worsen
 - Expansion may take months to years

Common pitfalls:

- **Underestimating cognitive activity:** Mental exertion counts toward energy envelope
- **Ignoring emotional stress:** Stress depletes energy reserves
- **Good-day overexertion:** Most common cause of relapse
- **External pressure:** Family/employer expectations pushing beyond envelope
- **Deconditioning fear:** Accepting current limits is not "giving up"

Cross-Condition Applications

High priority (exercise intolerance present):

- ME/CFS (cornerstone of management)
- Long COVID with PEM
- POTS with exercise intolerance
- Fibromyalgia with pain flares
- Post-viral fatigue syndromes

Moderate priority:

- Cancer-related fatigue during treatment
- Autoimmune conditions with fatigue
- Heart failure (already uses heart rate-based exercise limits)

Low priority / not applicable:

- Conditions without exercise intolerance
- Deconditioning without pathological exercise response (standard exercise progression appropriate)

Critical distinction: Pacing is for pathological exercise intolerance (PEM), NOT simple deconditioning. Standard graded exercise therapy appropriate for deconditioning; harmful for PEM.

31.7.6 Clinical Trial Opportunities

1. **Daratumumab in Long COVID:** Phase 2 trial in patients with elevated GPCR autoantibodies
2. **Rupatadine in EDS-POTS-MCAS:** Test triple-action (H1 + PAF + mast cell stabilizer) vs. standard antihistamines
3. **NAD⁺ precursors in cancer-related fatigue:** Extend Long COVID NR findings
4. **Catecholamine synthesis support in POTS:** L-tyrosine + Tetrahydrobiopterin (BH4) cofactors vs. placebo
5. **Comprehensive mitochondrial support in fibromyalgia:** Test full stack vs. individual components

31.8 Conclusion

ME/CFS research has identified mechanisms and treatments with implications extending far beyond ME/CFS diagnostic boundaries. The most impactful translational findings are:

1. **Plasma cell autoimmunity:** Explains why rituximab fails but daratumumab succeeds

2. **β 2-adrenergic receptor dysfunction:** Links MCAS, POTS, and vascular pathology
3. **Vascular-immune-energy triad:** Coordinated dysfunction requiring multi-target treatment
4. **NAD⁺ depletion:** Universal mechanism affecting multiple organ systems
5. **Pacing protocols:** Applicable to any condition with exercise intolerance

These findings demonstrate that ME/CFS is not an isolated condition but shares fundamental pathophysiology with multiple other diseases. Cross-pollination of research between ME/CFS and related conditions will accelerate progress for all patient populations.

Critical caveat: All translational recommendations require validation through condition-specific research. These represent **research opportunities and clinical hypotheses**, not established treatment guidelines for non-ME/CFS conditions. Clinicians and patients should approach these findings with appropriate scientific skepticism while recognizing their potential to advance understanding and treatment across the spectrum of post-viral, autoimmune, mitochondrial, and dysautonomic disorders.

32 Research Infrastructure Proposals

Abstract: This chapter proposes foundational research infrastructure to advance ME/CFS understanding, including longitudinal cohort studies, biomarker development programs, and computational modeling frameworks. These proposals are informed by insights from pediatric ME/CFS outcomes, which suggest that biological plasticity and recovery potential may be measurable and modifiable.

32.1 Longitudinal Deep Phenotyping Cohort

32.1.1 The ME/CFS Human Phenome Project

We propose the establishment of a comprehensive longitudinal deep phenotyping cohort—the “ME/CFS Human Phenome Project”—designed to capture the full biological trajectory of the condition across patient subtypes, disease stages, and outcomes. This infrastructure investment addresses a fundamental gap in ME/CFS research: the absence of large-scale, multi-dimensional, longitudinal data that would enable identification of predictive biomarkers, disease subtypes, and therapeutic targets.

Rationale and Scientific Justification

Existing ME/CFS research suffers from several structural limitations that this cohort would address. Most studies are cross-sectional, capturing a single timepoint in what is clearly a dynamic disease process. Sample sizes are typically small ($n=20-50$), underpowered to detect the heterogeneous subtypes that likely exist within the ME/CFS umbrella. Few studies combine multiple data modalities, limiting the ability to identify system-level interactions. Finally, standardization across studies remains poor, hampering meta-analysis and replication.

The pediatric recovery data provide compelling justification for longitudinal deep phenotyping. If 54–94% of children recover while fewer than 22% of adults do [504, 78], longitudinal tracking from early disease through either recovery or chronicity could reveal the biological determinants of these divergent trajectories. Such data could identify therapeutic targets by detecting which systems change during recovery, revealing potential intervention points. Longitudinal data could enable prognostic stratification by determining which baseline features predict recovery versus chronicity, enabling personalized treatment intensity. Subtype identification would emerge from tracking how different patient clusters evolve over time, distinguishing biological subtypes from disease stages. Finally, understanding mechanistic

sequences through temporal ordering would clarify whether immune abnormalities precede metabolic dysfunction or vice versa, addressing questions of causation versus correlation.

Study Design Overview

Population and Enrollment The cohort would enroll 500 participants across multiple sites internationally, with stratified enrollment to ensure representation across key dimensions:

- **Age strata:** 100 pediatric/adolescent (<18 years), 300 adult (18–60 years), 100 older adult (>60 years)
- **Disease duration:** 200 early (<2 years), 200 established (2–10 years), 100 long-duration (>10 years)
- **Severity:** 150 mild, 200 moderate, 150 severe/very severe
- **Trigger type:** Post-infectious (stratified by pathogen where known), post-other-trigger, gradual onset

Follow-up Duration and Assessment Schedule Participants would be followed for 5–10 years with comprehensive assessments at baseline, 6 months, 12 months, and annually thereafter. Abbreviated assessments (wearables, questionnaires, limited biomarkers) would occur quarterly. Event-triggered assessments would capture crashes, infections, significant functional changes, or apparent recovery.

Matched Control Cohort A cohort of 200 matched healthy controls would undergo identical assessments to establish reference ranges and age-related trajectories. An additional 100 disease controls with other fatiguing conditions (multiple sclerosis fatigue, post-cancer fatigue, fibromyalgia) would enable specificity analysis.

Comprehensive Assessment Battery

Multi-Omic Profiling Each comprehensive assessment would include:

- **Genomics:** Whole genome sequencing at baseline (variants, structural variants, HLA typing)
- **Epigenomics:** Genome-wide DNA methylation (Illumina EPIC array), targeted histone modification assays, epigenetic clocks
- **Transcriptomics:** Whole blood RNA-seq, immune cell subset-specific expression profiling
- **Proteomics:** Plasma proteomics (SomaScan or Olink platforms, >5000 proteins), CSF proteomics (subset, n=100)
- **Metabolomics:** Untargeted plasma and urine metabolomics, targeted panels (acylcarnitines, amino acids, organic acids, lipids)
- **Microbiome:** 16S rRNA and shotgun metagenomic sequencing of gut, oral, and skin microbiomes

Immune Profiling Comprehensive immune characterization would include:

- **Flow cytometry:** Extended panels for T cells (naive, memory, exhaustion markers), B cells (including plasmablasts), NK cells (cytotoxicity markers), monocyte subsets
- **Functional assays:** NK cell cytotoxicity, T cell proliferation, cytokine production capacity
- **Autoantibody panels:** GPCR autoantibodies, anti-neuronal antibodies, comprehensive autoimmune screening
- **Inflammatory markers:** High-sensitivity cytokine panels (Luminex), acute phase reactants

Autonomic and Cardiovascular Assessment

- **Tilt table testing:** Standardized 10-minute stand or 45-degree tilt with continuous hemodynamic monitoring
- **Heart rate variability:** 24-hour Holter monitoring with time-domain and frequency-domain analysis
- **Blood volume:** 51Cr-labeled red cell mass and 125I-albumin plasma volume (subset, n=100)
- **Sudomotor function:** QSART or Sudoscan
- **Baroreflex sensitivity:** Beat-to-beat blood pressure monitoring with baroreflex calculation

Exercise Physiology

- **Two-day CPET:** Cardiopulmonary exercise testing on consecutive days to capture PEM signature
- **Metabolic chamber:** Indirect calorimetry for 24-hour energy expenditure (subset, n=50)
- **Muscle biopsy:** Vastus lateralis biopsy for mitochondrial function, fiber typing, histology (optional, subset)
- **Handgrip dynamometry:** Serial grip strength with fatigue protocol

Neuroimaging and Neurophysiology

- **Structural MRI:** T1, T2, FLAIR, DTI for white matter integrity
- **Functional MRI:** Resting state connectivity, task-based (motor, cognitive)
- **PET imaging:** TSPO ligand for neuroinflammation (subset, n=50)
- **EEG:** Quantitative EEG with spectral analysis
- **Cognitive testing:** Computerized battery (attention, memory, processing speed, executive function)

Clinical and Patient-Reported Outcomes

- **Standardized questionnaires:** DSQ-PEM, MFI, SF-36, Bell Disability Scale, PHQ-9, GAD-7
- **Symptom diaries:** Daily electronic symptom tracking
- **Activity monitoring:** Continuous accelerometry (wrist-worn devices)
- **Sleep assessment:** Actigraphy, sleep diaries, polysomnography (subset)
- **Functional assessment:** 6-minute walk test (when safe), NASA Lean Test

Biorepository Specifications

The project would establish a centralized biorepository with long-term storage capacity:

- **Sample types:** Whole blood, serum, plasma (EDTA, citrate, heparin), PBMCs, urine, stool, saliva, DNA, RNA
- **Storage conditions:** Liquid nitrogen (-196°C) for cells and RNA, -80°C for other samples
- **Aliquoting:** Multiple aliquots per sample type to enable future analyses without thaw cycles
- **Quality control:** Standardized collection protocols, processing within 2 hours, regular QC audits
- **Capacity:** Estimated 2 million aliquots over the project duration
- **Access policy:** Open to qualified researchers with approved proposals, data sharing agreements, and acknowledgment requirements

Data Infrastructure and Analysis

Data Management

- **Central database:** FAIR-compliant (Findable, Accessible, Interoperable, Reusable) data repository
- **Data harmonization:** Common data elements aligned with NIH/CDC standards
- **Privacy protection:** De-identification, secure enclaves for sensitive data, tiered access
- **Longitudinal linking:** Robust participant ID system enabling cross-timepoint analysis

Analysis Plan Primary analyses would include:

- **Trajectory modeling:** Latent class growth models to identify distinct disease courses
- **Predictive modeling:** Machine learning approaches to predict recovery, progression, treatment response
- **Multi-omic integration:** Network-based integration of genomic, transcriptomic, proteomic, and metabolomic data
- **Subtype identification:** Unsupervised clustering across data modalities

- **Causal inference:** Mendelian randomization for causal pathway identification

Budget Estimate and Feasibility

Estimated Costs

- **Multi-omic profiling:** \$15,000–20,000 per participant per comprehensive assessment
- **Clinical assessments:** \$5,000 per participant per visit
- **Neuroimaging:** \$3,000 per participant per session
- **Biorepository:** \$5 million infrastructure, \$500,000/year operations
- **Data infrastructure:** \$3 million setup, \$500,000/year maintenance
- **Coordination and administration:** \$2 million/year
- **Total estimated budget:** \$40–50 million over 10 years

Feasibility Considerations This proposal is ambitious but feasible given precedents. The UK Biobank enrolled 500,000 participants with extensive phenotyping at lower per-participant cost but less depth. The All of Us Research Program demonstrates large-scale longitudinal deep phenotyping infrastructure. The ME Research UK Biobank, though smaller, provides a model specific to ME/CFS.

Key feasibility challenges include participant retention over 5–10 years, particularly for severely ill patients who may find assessments burdensome. Mitigation strategies include home visits, abbreviated protocols for severe cases, and strong participant engagement. Protocol evolution will be necessary as technologies advance; the protocol must allow incorporation of new assays while maintaining comparability.

Expected Outcomes If successful, this cohort would generate the definitive longitudinal dataset for ME/CFS research, enabling validated prognostic biomarkers to guide treatment intensity, biological subtype definitions to enable precision medicine, therapeutic targets emerging from trajectory analysis, and a shared resource to accelerate research across the field

32.2 Recovery Potential Index Development

32.2.1 Conceptual Framework

Building on the Recovery Capital model (Speculation 23), we propose development of a quantitative Recovery Potential Index (RPI)—a composite biomarker intended to measure an individual’s residual capacity for recovery from ME/CFS. The conceptual foundation rests on the hypothesis that recovery potential is not binary but exists on a continuum, and that this continuum may be objectively measurable through biomarkers reflecting biological plasticity, regenerative capacity, and systemic resilience.

The rationale for an RPI derives from several observations. The dramatic difference in recovery rates between pediatric and adult patients (54–94% versus $\leq 22\%$ [504, 78]) suggests that biological factors beyond disease severity determine recovery capacity. Within both populations, some patients recover while others with apparently similar presentations do not, implying individual differences in recovery potential. If recovery potential were measurable, treatment intensity could be stratified accordingly—aggressive early intervention for those with preserved potential, palliative approaches for those with depleted reserves.

⚡ **Limitation 1: Research Infrastructure Proposals: Entirely Aspirational**

The research infrastructure proposals in this chapter—the ME/CFS Human Phenome Project, the Recovery Potential Index, the Crash Impact Biomarker panel, and the ME/CFS Digital Twin—are conceptual frameworks, not funded or operational programs. Key boundaries:

- No component of the Recovery Potential Index (epigenetic age acceleration, naive T cell proportion, telomere length, HSC clonality, HRV, metabolic flexibility) has been validated as a predictor of ME/CFS recovery; the index is a hypothesis about what *might* predict recovery, not an established tool.
- The Crash Impact Biomarker panel has not been tested in any study; the “expected patterns” described are predictions derived from general exercise physiology and inflammation biology, not from ME/CFS-specific data.
- Budget estimates (\$40–50 million for the Phenome Project) and feasibility analyses are order-of-magnitude projections; actual costs and practical challenges may differ substantially.
- The Digital Twin concept, while established in other medical fields, has no ME/CFS-specific training data; clinical decision support claims are aspirational.

These proposals are presented to stimulate research investment, not as descriptions of existing or imminent capabilities.

~ **Hypothesis 1: Recovery Potential Index**

Recovery from ME/CFS requires sufficient reserves across multiple biological systems: epigenetic flexibility, immune renewal capacity, stem cell reserves, autonomic adaptability, and metabolic plasticity. These reserves can be quantified through specific biomarkers and combined into a composite index that predicts recovery probability. Individuals with high RPI scores retain the biological capacity for recovery if appropriately treated, while those with low RPI scores have crossed thresholds beyond which recovery mechanisms are impaired regardless of treatment.

32.2.2 Component Biomarkers

We propose six component biomarkers for the RPI, each selected based on biological rationale, measurement feasibility, and relevance to the pediatric-adult recovery differential.

Epigenetic Age Acceleration

Scientific Rationale DNA methylation-based “epigenetic clocks” estimate biological age independent of chronological age. Epigenetic age acceleration (biological age exceeding chronological age) has been associated in the general aging literature with reduced regenerative capacity, increased mortality risk, and impaired recovery from various stressors. Preliminary evidence suggests ME/CFS patients exhibit epigenetic age acceleration, though this requires replication.

The pediatric recovery advantage may partly reflect the plasticity of younger epigenomes. Adolescent immune cells and other tissues are actively undergoing developmental programming, potentially enabling “reprogramming” of disease states in ways that adult tissues cannot achieve.

Measurement Epigenetic age would be calculated from peripheral blood DNA methylation using established clocks (Horvath, Hannum, GrimAge, or PhenoAge). The acceleration metric is the residual when regressing epigenetic age on chronological age. This measurement is reproducible (ICC >0.95) and commercially available.

Expected Pattern Higher epigenetic age acceleration would predict lower recovery probability. Patients whose epigenetic age substantially exceeds their chronological age may have depleted the cellular plasticity required for recovery.

Naive T Cell Proportion

Scientific Rationale Naive T cells (CD45RA⁺CCR7⁺) represent the immune system’s reserve capacity—cells that have not yet been committed to specific antigens and retain the flexibility to respond to new challenges. The naive T cell pool declines with age (thymic involution) and is consumed by chronic infections or persistent immune activation.

In ME/CFS, chronic immune activation may preferentially deplete naive T cells, converting them to memory or effector phenotypes. This consumption of “immune capital” could explain why recovery becomes less likely over time. Children, with active thymic output, continuously replenish naive T cells; adults lack this regenerative capacity.

Measurement Flow cytometry for CD3⁺CD4⁺CD45RA⁺CCR7⁺ (naive CD4 T cells) and CD3⁺CD8⁺CD45RA⁺CCR7⁺ (naive CD8 T cells), expressed as percentage of total T cells. Recent thymic emigrants (CD31⁺ naive cells) provide additional information about active thymic contribution.

Expected Pattern Higher naive T cell proportions, relative to age-matched norms, would predict greater recovery potential. Severely depleted naive pools may indicate irreversible immune exhaustion.

Telomere Length

Scientific Rationale Telomeres—the protective caps on chromosome ends—shorten with each cell division and with oxidative stress. Critically short telomeres trigger cellular senescence, limiting the regenerative capacity of tissues. In the broader aging biology literature, telomere attrition has been proposed as a mechanism of biological aging and may be accelerated by chronic inflammation.

Pediatric cells have longer telomeres and active telomerase, providing greater replicative capacity. This reserve may enable the cellular renewal necessary for recovery from ME/CFS.

Measurement Leukocyte telomere length via quantitative PCR (T/S ratio method) or Southern blot. Flow-FISH provides cell type-specific telomere length but is more technically demanding.

Expected Pattern Longer telomeres relative to age would predict higher recovery potential. Critically short telomeres may indicate depleted replicative capacity incompatible with recovery.

Hematopoietic Stem Cell Clonality

Scientific Rationale Hematopoietic stem cells (HSCs) regenerate all blood and immune cells throughout life. HSC clonality refers to the diversity of HSC clones contributing to hematopoiesis; healthy young individuals have highly polyclonal hematopoiesis, while aging and disease states produce oligoclonal dominance as HSC diversity declines.

We hypothesize (Speculation 24) that ME/CFS involves accelerated HSC exhaustion, potentially driven by repeated immune activation during crash episodes. Higher HSC diversity would indicate preserved regenerative reserves.

Measurement HSC clonality can be inferred from single-cell sequencing approaches or from detection of clonal hematopoiesis of indeterminate potential (CHIP) mutations. A simpler proxy is the diversity of T cell receptor (TCR) or B cell receptor (BCR) repertoires, measurable via immunosequencing.

Expected Pattern Higher clonal diversity (more polyclonal hematopoiesis) would predict greater recovery potential. Oligoclonal dominance suggests depleted HSC reserves.

Heart Rate Variability Metrics

Scientific Rationale Heart rate variability (HRV) reflects autonomic nervous system function and, more broadly, the organism's capacity for adaptive regulation. High HRV indicates a flexible, resilient autonomic system capable of responding appropriately to challenges. Low HRV indicates a rigid system with limited adaptive capacity.

Beyond autonomic function specifically, HRV may serve as an integrative biomarker of systemic health. The vagal pathways reflected in HRV are linked to inflammatory regulation (the "cholinergic anti-inflammatory pathway"), stress responses, and metabolic function. HRV thus provides a window into the organism's overall regulatory capacity.

Measurement 24-hour Holter monitoring with calculation of time-domain (SDNN, RMSSD) and frequency-domain (HF power, LF/HF ratio) metrics. Shorter recordings (5-minute seated) provide less comprehensive but more practical assessment.

Expected Pattern Higher HRV, particularly higher HF power and SDNN, relative to age-matched norms would predict greater recovery potential. Very low HRV may indicate irreversible autonomic rigidity.

Metabolic Flexibility

Scientific Rationale Metabolic flexibility refers to the organism's ability to switch between fuel substrates (primarily carbohydrates and fats) in response to energy demands and substrate availability. Healthy individuals readily shift from fat oxidation during fasting to carbohydrate oxidation after meals. Metabolic inflexibility—inability to appropriately shift fuel utilization—is associated in the metabolic literature with mitochondrial dysfunction, insulin resistance, and impaired exercise capacity.

ME/CFS is characterized by metabolic abnormalities that may impair this flexibility. The ability to respond to metabolic challenges may indicate preserved mitochondrial capacity and systemic resilience.

Measurement Respiratory exchange ratio (RER) dynamics during mild metabolic challenge. This could involve measuring RER during a brief, submaximal exercise bout or during transition from fasted to fed states via indirect calorimetry. The key metric is the magnitude and rapidity of RER change in response to challenge.

Expected Pattern Greater RER responsiveness (ability to shift RER appropriately during challenge) would predict higher recovery potential. Fixed, inflexible RER suggests metabolic rigidity incompatible with recovery.

32.2.3 Index Construction and Interpretation

Normalization and Weighting

Each component biomarker would be normalized to age- and sex-matched reference ranges, yielding z-scores or percentile ranks. This normalization is essential because most biomarkers change with age; what matters is not the absolute value but the value relative to healthy peers.

The composite RPI would be calculated as a weighted sum:

$$\text{RPI} = \sum_{i=1}^6 w_i \cdot z_i$$

where z_i is the normalized score for component i and w_i is its weight.

Initial Weighting Approach In the absence of empirical validation data, we propose equal weighting ($w_i = 1/6$ for all components) as the initial default. This agnostic approach acknowledges uncertainty about the relative importance of each component.

Alternative weighting schemes could be derived empirically once longitudinal outcome data are available, using regression coefficients from models predicting recovery. Principal component analysis could also identify natural weightings based on shared variance across components.

Clinical Interpretation Thresholds

Pending validation, we propose the following preliminary interpretive framework:

- **High Recovery Potential (RPI > 0.7):** Most component biomarkers within or above age-matched norms. Biological reserves appear preserved. Aggressive treatment and strict pacing may maximize recovery probability.
- **Moderate Recovery Potential (RPI 0.4–0.7):** Mixed biomarker profile with some components preserved, others depleted. Recovery possible but not assured. Individualized approach based on which components are preserved.
- **Low Recovery Potential (RPI < 0.4):** Multiple biomarkers indicating depleted reserves. Recovery unlikely with current interventions. Focus on symptom management, preventing further decline, and quality of life.

△ Warning 1: RPI Interpretation Requires Validation

These thresholds are proposed for research purposes and require empirical validation before clinical application. A low RPI score does not definitively preclude recovery, nor does a high score guarantee it. The RPI is intended as one input into clinical decision-making, not a deterministic prediction.

32.2.4 Validation Requirements

The RPI concept requires rigorous validation before clinical utility can be established:

Phase 1: Cross-Sectional Validation

- Measure all six components in a cohort of ME/CFS patients ($n \geq 200$) with matched healthy controls ($n \geq 100$)
- Confirm that ME/CFS patients have lower RPI scores than controls
- Confirm that pediatric patients have higher RPI scores than adult patients
- Assess correlation between RPI and disease duration, severity

Phase 2: Longitudinal Predictive Validation

- Follow cohort prospectively for 2–5 years
- Assess whether baseline RPI predicts subsequent recovery
- Determine optimal thresholds through receiver operating characteristic (ROC) analysis
- Calculate positive and negative predictive values

Phase 3: Clinical Utility Validation

- Test whether RPI-stratified treatment improves outcomes compared to non-stratified care
- Assess cost-effectiveness of RPI measurement
- Develop simplified versions (fewer components) if full panel proves impractical

32.2.5 Limitations and Caveats

Several limitations must be acknowledged:

- **Speculative foundation:** The RPI is based on theoretical models that require validation. The component biomarkers have not been proven to predict ME/CFS recovery.
- **Measurement challenges:** Some components (HSC clonality, metabolic flexibility) require specialized assays not widely available.
- **Cost:** Full RPI assessment would cost several thousand dollars, potentially limiting accessibility.
- **Potential for harm:** A “low recovery potential” designation could discourage patients and providers from attempting treatments that might help. Any clinical application must avoid premature therapeutic nihilism.
- **Dynamic nature:** Recovery potential may change over time with treatment, disease progression, or natural fluctuation. Serial RPI measurement may be necessary.

Despite these limitations, the RPI concept provides a framework for operationalizing the Recovery Capital model and generating testable predictions about recovery mechanisms. Even if the specific components proposed here prove suboptimal, the general approach—quantifying biological reserves that enable recovery—may advance understanding of ME/CFS heterogeneity and prognosis

32.3 Crash Impact Biomarkers

32.3.1 Conceptual Framework

Post-exertional malaise (PEM) crashes are described by patients as profoundly damaging events, yet no objective measures currently quantify the biological “cost” of a crash. This absence creates multiple problems: patients struggle to communicate crash severity to clinicians, the cumulative harm of repeated crashes cannot be tracked, and the Recovery Capital depletion hypothesis (Speculation 23) remains untestable without objective metrics.

We propose development of a Crash Impact Biomarker panel designed to quantify the acute biological damage from PEM episodes and track cumulative effects over time.

32.3.2 Proposed Biomarker Panel

The panel would measure biological markers before a predictable crash (baseline), at 24 hours post-crash (acute phase), and at 72 hours post-crash (resolution phase). Patients with predictable crash triggers (medical appointments, known exertion events) would be recruited for standardized pre/post sampling.

Cell-Free DNA (cfDNA)

Rationale Cell-free DNA in plasma reflects cellular damage and turnover. Elevated cfDNA indicates tissue injury, apoptosis, or necrosis. In exercise physiology, cfDNA rises transiently after intense exertion and returns to baseline within hours in healthy individuals [651]. Prolonged cfDNA elevation would indicate ongoing cellular damage.

Measurement Quantitative PCR or fluorometric assay of plasma cfDNA concentration (ng/mL). Baseline, 24h, and 72h samples.

Expected Pattern Crash-inducing exertion should produce cfDNA elevation exceeding that seen in healthy individuals after equivalent exertion. Slower return to baseline would indicate impaired cellular recovery. Cumulative cfDNA elevation over serial crashes would support the “damage accumulation” hypothesis.

Inflammatory Cytokine Panel

Rationale PEM involves inflammatory activation (see Chapter 2, PEM mechanism). The magnitude of cytokine elevation may correlate with crash severity and predict recovery time.

Measurement High-sensitivity multiplex assay (Luminex or similar) for:

- IL-6 (primary acute phase mediator)
- TNF- α (pro-inflammatory, tissue damage marker)
- IL-1 β (inflammasome activation)
- IL-10 (anti-inflammatory, resolution marker)
- IL-8 (neutrophil chemotaxis)

Expected Pattern Crash should produce disproportionate cytokine elevation relative to exertion magnitude. IL-6 and TNF- α peaks at 24h, with IL-10 rise indicating resolution at 72h. Failure of IL-10 elevation or persistent IL-6/TNF- α would indicate impaired resolution.

Lactate:Pyruvate Ratio

Rationale The lactate:pyruvate ratio reflects cellular redox state and mitochondrial function. Elevated ratio indicates anaerobic metabolism, mitochondrial dysfunction, or NAD⁺ depletion—all implicated in ME/CFS pathophysiology.

Measurement Venous blood sampling for lactate and pyruvate; calculate ratio. Normal ratio is 10:1 to 20:1.

Expected Pattern Post-crash elevation of lactate:pyruvate ratio, particularly at 24h, would indicate metabolic stress. Persistent elevation at 72h would suggest ongoing mitochondrial dysfunction.

8-Hydroxy-2'-Deoxyguanosine (8-OHdG)

Rationale 8-OHdG is a marker of oxidative DNA damage. Elevated urinary 8-OHdG indicates oxidative stress sufficient to damage DNA, which may accumulate over repeated crashes and contribute to epigenetic changes.

Measurement ELISA on spot urine sample, normalized to creatinine.

Expected Pattern Crash-induced oxidative stress should elevate 8-OHdG, particularly if mitochondrial dysfunction produces reactive oxygen species. Cumulative elevation over serial crashes would support the oxidative damage accumulation hypothesis.

Cortisol Profile

Rationale HPA axis dysregulation is documented in ME/CFS [388]. Crash episodes may produce abnormal cortisol responses—either exaggerated stress response or blunted responsiveness indicating HPA exhaustion.

Measurement Salivary cortisol at waking, 30 minutes post-waking (CAR—cortisol awakening response), and evening. Alternatively, 24-hour urinary free cortisol.

Expected Pattern Crash may produce blunted CAR (indicating HPA axis fatigue) or elevated evening cortisol (indicating impaired cortisol clearance or chronic stress activation). Pattern may differ by crash severity and cumulative crash history.

NK Cell Cytotoxicity

Rationale Natural killer cell dysfunction is one of the most consistent immune findings in ME/CFS [200]. Crashes may acutely worsen NK function, and cumulative crashes may produce progressive NK exhaustion.

Measurement ⁵¹Cr-release assay or flow cytometric cytotoxicity assay on fresh PBMCs.

Expected Pattern NK cytotoxicity should decline post-crash (24h) and may or may not recover by 72h. Failure to recover would indicate crash-induced immune suppression. Serial measurement over multiple crashes could track cumulative NK exhaustion.

32.3.3 Validation Study Design

Study Overview

A prospective cohort study would validate the Crash Impact Biomarker panel by correlating biomarker changes with crash severity, recovery time, and long-term outcomes.

Participants

- n=50 ME/CFS patients with predictable crash triggers
- Mild to moderate severity (able to attend study visits)
- Willing to undergo repeated blood sampling
- Matched healthy controls (n=25) undergoing equivalent exertion protocol

Protocol

1. Identify predictable crash trigger (medical appointment, standardized exertion)
2. Baseline sampling 24 hours before trigger
3. Crash trigger event (documented exertion magnitude)
4. 24-hour post-trigger sampling
5. 72-hour post-trigger sampling
6. Symptom diary for 14 days post-trigger
7. Repeat protocol for 3 separate crashes per participant

Outcomes

- Primary: Correlation between biomarker panel scores and patient-reported crash severity
- Secondary: Correlation with recovery time (days to return to baseline symptoms)
- Exploratory: Cumulative biomarker changes across serial crashes; correlation with 6-month functional trajectory

32.3.4 Clinical Utility

If validated, the Crash Impact Biomarker panel would enable:

1. **Objective crash documentation:** Providing evidence of biological harm for disability applications, workplace accommodations, and clinical decision-making
2. **Recovery Capital tracking:** Serial measurement to quantify cumulative damage and inform treatment intensity
3. **Intervention evaluation:** Objective endpoint for trials of crash-prevention or crash-mitigation interventions
4. **Pacing motivation:** Concrete biological feedback may improve pacing adherence by demonstrating the real cost of envelope violations
5. **Subtype identification:** Different biomarker patterns may identify inflammatory-predominant, metabolic-predominant, or autonomic-predominant crash phenotypes

32.4 Computational Patient Modeling: The ME/CFS Digital Twin

32.4.1 The Digital Twin Concept

A “digital twin” is a computational model of an individual patient that integrates multiple data streams to predict responses to interventions, optimize treatment sequences, and enable personalized medicine. Digital twin approaches are advancing rapidly in oncology, cardiology, and intensive care for personalized treatment optimization.

ME/CFS, with its heterogeneous presentations, multiple potential subtypes, and unpredictable treatment responses, is an ideal candidate for digital twin development. Current treatment is largely empirical—trial and error with medications and supplements—reflecting limited biomarkers to guide personalized therapy. A computational model that could predict “Patient X is likely to respond to fludrocortisone but not LDN based on their biomarker profile” would transform clinical practice.

32.4.2 Data Sources for ME/CFS Digital Twin

An ME/CFS digital twin would integrate:

Genomic Data

- Risk alleles (HLA types, immune-related polymorphisms)
- Pharmacogenomics (drug metabolism variants: CYP2D6, CYP2C19)
- Mitochondrial DNA variants
- Polygenic risk scores for comorbidities

Epigenomic Data

- DNA methylation profiles (epigenetic clocks, immune cell type proportions)
- Plasticity markers (indicators of epigenetic flexibility vs. locked states)
- Longitudinal epigenetic trajectories

Metabolomic Data

- Plasma metabolome (energy substrates, amino acids, lipids)
- Acylcarnitine profile (mitochondrial function indicators)
- NAD⁺/NADH status
- Response to metabolic challenges (RER dynamics)

Immune Data

- Lymphocyte subset proportions (naive, memory, exhausted T cells)
- NK cell phenotype and function
- Autoantibody profiles (GPCR autoantibodies, ANA)
- Cytokine production capacity

Continuous Wearable Data

- Activity levels (accelerometry)
- Heart rate and HRV (autonomic state)
- Sleep architecture (duration, efficiency, stages)
- Skin temperature, electrodermal activity (if available)

Patient-Reported Data

- Daily symptom diaries (fatigue, pain, cognition, PEM)
- Crash logs with severity ratings
- Medication and supplement adherence
- Life events and stressors

Treatment History

- All medications and supplements tried
- Duration of each trial
- Response (improvement, no change, worsening)
- Adverse effects

32.4.3 Modeling Approaches**Machine Learning for Pattern Recognition**

Supervised learning algorithms can identify patterns in multi-modal data that predict treatment response, crash risk, or disease trajectory. Random forests, gradient boosting, and neural networks have all shown promise in similar medical prediction tasks. The key challenge is obtaining sufficient labeled training data (patients with known outcomes for specific treatments).

Causal Inference

Correlation does not imply causation. Observational data showing that patients on fludrocortisone improve more than those not on fludrocortisone could reflect either drug efficacy or selection bias (perhaps only OI-predominant patients are prescribed fludrocortisone, and OI-predominant patients have better prognosis regardless of treatment). Causal inference methods—propensity score matching, instrumental variables, Mendelian randomization—can help disentangle true treatment effects from confounders.

Reinforcement Learning for Treatment Optimization

Reinforcement learning (RL) algorithms learn optimal action sequences through trial and error. In the digital twin context, RL could learn optimal treatment sequences: “Start with OI treatment; if HRV improves but fatigue persists after 8 weeks, add LDN; if inflammatory markers remain elevated, consider mast cell stabilization.” RL requires either extensive historical data with sequential treatment decisions or simulation environments that accurately model patient physiology.

Bayesian Updating

As new data arrives (lab results, symptom reports, treatment responses), the digital twin’s predictions should update accordingly. Bayesian methods provide a principled framework for integrating prior knowledge with new evidence, maintaining calibrated uncertainty estimates as information accumulates.

32.4.4 Clinical Applications

Treatment Response Prediction

“Based on your biomarker profile, there is a 73% probability of meaningful improvement with aggressive OI treatment, versus 12% probability with LDN alone.”

Optimal Treatment Sequencing

“Given your current state, the highest expected value treatment sequence is: (1) fludrocortisone + midodrine, (2) if insufficient response at 8 weeks add LDN, (3) if still insufficient consider IVIG evaluation.”

Crash Risk Prediction

“Based on your HRV trend and activity pattern over the past 3 days, crash risk in the next 48 hours is elevated (68%). Recommend reducing activity by 30%.”

Personalized Prognosis

“Based on your RPI components and disease trajectory, estimated probability of meaningful improvement over 2 years is 45% with aggressive front-loading treatment, versus 15% with standard care.”

32.4.5 Development Pathway

Phase 1: Retrospective Model Development

- Compile existing datasets with treatment outcomes
- Develop initial prediction models
- Identify most informative features
- Estimate achievable prediction accuracy

Phase 2: Prospective Validation

- Enroll cohort with comprehensive baseline assessment
- Make blinded predictions of treatment responses
- Compare predictions to actual outcomes
- Refine models based on prediction errors

Phase 3: Clinical Decision Support

- Deploy validated models as clinical decision support tools
- Evaluate impact on patient outcomes and clinician decision-making
- Continuous learning from new patient data
- Regulatory pathway for medical device approval if warranted

32.4.6 Challenges and Limitations

- **Data availability:** Comprehensive multi-omic data with treatment outcomes is scarce; initial models may be data-limited
- **Model interpretability:** “Black box” predictions may be difficult for clinicians to trust or explain to patients; interpretable models may be less accurate
- **Computational resources:** Real-time integration of wearable data with multi-omic profiles requires substantial computing infrastructure
- **Validation requirements:** Models must be validated in external cohorts before clinical deployment
- **Evolving biology:** ME/CFS may change over time within individuals; models must account for disease dynamics
- **Ethical considerations:** Algorithmic treatment recommendations raise questions of responsibility, consent, and equity

Despite these challenges, the digital twin paradigm represents the future of personalized medicine for ME/CFS. As data accumulates from longitudinal cohorts and clinical practice, computational models will increasingly complement clinical judgment, enabling more precise and effective treatment.

32.5 Model Organism Research Panel

Translating ME/CFS research into mechanistic understanding requires model systems that recapitulate key features of the disease. While no single model fully captures the complexity of ME/CFS, a panel approach—combining animal models, cellular systems, and organoid platforms—enables investigation of specific pathophysiological axes under controlled conditions.

32.5.1 Animal Models of Post-Viral Fatigue

Several animal models have been developed to study post-infectious fatigue states relevant to ME/CFS. Murine models of post-viral fatigue, including those using poly(I:C) injection or chronic viral infection with lymphocytic choriomeningitis virus (LCMV), produce lasting reductions in voluntary wheel-running activity, neuroinflammation, and metabolic disruption that parallel features of human ME/CFS [652]. These models enable controlled study of the transition from acute infection to chronic fatigue.

Rodent models of immune activation using Toll-like receptor agonists have demonstrated persistent microglial activation, altered tryptophan metabolism, and disrupted hypothalamic–pituitary–adrenal (HPA) axis signaling weeks after the initial immune challenge. Such models are particularly valuable for studying neuroinflammatory mechanisms (Chapter 8) and testing candidate interventions targeting neuroglial pathways.

⚠ Limitation 2

Animal models cannot fully replicate the subjective experience of fatigue or post-exertional malaise, and species differences in immune regulation, autonomic function, and metabolism limit direct translation to human ME/CFS. Results from animal studies should be interpreted as hypothesis-generating rather than confirmatory.

32.5.2 Cellular Models of Metabolic Dysfunction

In vitro studies using peripheral blood mononuclear cells (PBMCs) and skeletal muscle cells from ME/CFS patients have revealed consistent bioenergetic abnormalities. Seahorse extracellular flux analysis of patient-derived PBMCs demonstrates reduced mitochondrial reserve capacity and impaired oxidative phosphorylation compared with healthy controls [326, 653]. These findings support a cell-intrinsic metabolic defect rather than a purely systemic phenomenon.

Immortalized cell lines and primary cultures exposed to patient serum have been used to investigate circulating factors that may impair cellular bioenergetics. Studies exposing healthy cells to ME/CFS serum have reported reduced ATP production and altered mitochondrial membrane potential, suggesting the presence of inhibitory factors in the ME/CFS milieu [612]. Identifying these factors remains an active area of investigation with direct therapeutic implications.

Induced pluripotent stem cell (iPSC) technology offers emerging opportunities to generate patient-specific neuronal, cardiac, and immune cells for disease modeling. While iPSC-derived models of ME/CFS are still in early development, they hold promise for studying tissue-specific manifestations—such as neuronal hypersensitivity or cardiomyocyte dysfunction—in a genetically defined background.

32.5.3 Organoid Systems for Tissue-Specific Effects

Three-dimensional organoid cultures represent the frontier of ME/CFS model systems. Gut organoids derived from patient biopsies can recapitulate epithelial barrier dysfunction and altered microbiome–host interactions relevant to the gastrointestinal symptoms frequently reported in ME/CFS (Chapter 11). Brain organoids may enable investigation of neuroinflammatory processes and blood–brain barrier integrity in a human-relevant system.

Multi-organ-on-a-chip platforms, which connect microfluidic compartments representing different tissue types, offer the potential to model systemic interactions—for example, the gut–brain axis or immune–metabolic crosstalk—that characterize ME/CFS pathophysiology. These platforms are still maturing but represent a promising direction for studying how dysfunction in one organ system propagates to others.

32.5.4 Towards an Integrated Model Panel

? Open Question 1

What is the minimal panel of model systems needed to capture the core pathophysiology of ME/CFS? A proposed minimum panel includes: (1) a post-viral murine model for systemic and neuroinflammatory mechanisms, (2) patient-derived PBMCs for bioenergetic profiling, (3) gut organoids for barrier and microbiome studies, and (4) iPSC-derived neurons for central sensitization research. Validation across this panel would strengthen confidence in candidate mechanisms and therapeutic targets.

Standardization of model systems is essential for reproducibility. The ME/CFS research community would benefit from consensus protocols for cell culture conditions, animal model endpoints, and organoid characterization, analogous to the standardization efforts that have advanced cancer and neurodegenerative disease research.

33 Proposed Research Studies

Abstract: This chapter presents detailed protocols for research studies designed to test hypotheses derived from pediatric ME/CFS outcomes and to advance understanding of recovery mechanisms. These proposals range from observational cohort studies to randomized controlled trials, each with explicit hypotheses, design specifications, and expected outcomes.

33.1 Multi-Modal Testing of Selective Energy Dysfunction Hypothesis

33.1.1 Background and Rationale

The Selective Energy Dysfunction Hypothesis proposes a mechanistic framework for ME/CFS that distinguishes it from global mitochondrial failure. Rather than pan-cellular energy depletion, the hypothesis posits that ME/CFS involves selective impairment of brain-dependent and demand-responsive processes (voluntary motor control, cognitive exertion, autonomic demand-response) while autonomous peripheral processes remain preserved (basal metabolism, hair/nail growth, resting immune function). This framework suggests that the primary pathophysiological bottleneck is central nervous system energy metabolism rather than peripheral mitochondrial dysfunction.

If validated, this hypothesis would fundamentally reorient ME/CFS research toward CNS energy metabolism, provide mechanistic rationale for brain-targeted interventions and pacing strategies, and enable biomarker-based diagnosis and personalized subtyping. This comprehensive multi-modal study is designed to systematically test the selectivity pattern across CNS-dependent versus autonomous processes.

⚠ Limitation 1: Proposed Studies: Untested Hypotheses Framed as Research Protocols

The study proposals in this chapter formalise hypotheses from earlier chapters into research designs. While this is a valuable exercise, readers should note:

- These are *proposed* protocols, not completed or ongoing studies. None has received IRB approval, funding, or peer review of its design.
- Sample size calculations assume effect sizes derived from preliminary or single-study data; if true effect sizes are smaller, the studies may be substantially under-

powered.

- Several proposals (e.g., the selective energy dysfunction study, the multi-domain phenotyping trial) assume that the underlying hypotheses are sufficiently established to warrant a confirmatory study design; in reality, most would benefit from exploratory/pilot phases first.
- The framing as “ready-to-implement protocols” may overstate feasibility—practical challenges including patient recruitment (especially severe patients), assay standardisation, and multi-site coordination are acknowledged but not fully resolved.

33.1.2 Study Innovation

This is the first study to systematically test the selectivity pattern using an integrated multi-modal approach that includes:

1. **Novel peripheral biomarkers:** Hair and nail growth as objective measures of preserved autonomous processes, replacing reliance on subjective symptom reports
2. **CNS-specific imaging:** FDG-PET to quantify brain hypometabolism and correlate with clinical CNS symptoms
3. **Demand-response testing:** Two-day cardiopulmonary exercise tests (CPET) with muscle biopsies to assess exercise intolerance and peripheral energy metabolism
4. **CNS bypass methodology:** Electrical stimulation of muscle versus voluntary contraction to test the CNS component of exercise intolerance
5. **Autonomic demand-response:** Tilt table testing with cerebral blood flow monitoring to assess demand-response failures in autonomic regulation

33.1.3 Hypothesis

~ Hypothesis 1: Selective Energy Dysfunction in ME/CFS

ME/CFS involves selective impairment of CNS-dependent and demand-responsive processes while autonomous peripheral processes remain preserved. Specifically:

1. Hair/nail growth will be normal in ME/CFS patients, demonstrating preserved autonomous peripheral metabolism
2. Brain FDG-PET hypometabolism will correlate with CNS symptoms (cognitive dysfunction, orthostatic intolerance)
3. Two-day CPET will show functional decline despite normal resting muscle ATP levels
4. Electrical muscle stimulation will produce greater force output than voluntary contraction (CNS bypass effect)
5. Cerebral blood flow will decline during orthostatic challenge in 90%+ of patients

These differences will persist after controlling for systemic inflammatory markers and

metabolic parameters, indicating CNS-specific dysfunction rather than global energy failure.

33.1.4 Study Design Overview

Design Type

Multi-modal cross-sectional case-control study with longitudinal biomarker tracking component. This combines structural brain imaging, functional physiology testing, and novel biomarker assessment in a single integrated protocol.

Sample Size and Power

- **Total enrollment:** n=72 (36 ME/CFS patients + 36 matched healthy controls)
- **ME/CFS stratification:** 12 mild, 12 moderate, 12 severe (using Bell Disability Scale)
- **Statistical power:** >80% for all primary aims
- **Attrition buffer:** 20% built into target enrollment

Duration

Three years total: Year 1 (setup, IRB approval, pilot phase with n=10), Years 2–3 (main enrollment and assessment)

33.1.5 Five Core Aims

★ Achievement 1: Aim 1: Peripheral Autonomy Preservation

Test whether autonomous peripheral processes (hair/nail growth) are preserved in ME/CFS patients compared to controls. Hair and nail growth measurements will serve as novel, objective biomarkers of preserved basal metabolic function, distinguishing ME/CFS from global mitochondrial disease.

★ Achievement 2: Aim 2: Brain Hypometabolism and CNS Symptoms

Use FDG-PET imaging to quantify whole-brain and regional glucose metabolism in ME/CFS patients and assess correlations with cognitive dysfunction, orthostatic intolerance, and fatigue severity. Brain hypometabolism in prefrontal and brainstem regions would support CNS energy limitation as primary mechanism.

★ Achievement 3: Aim 3: Demand-Response Failure Despite Preserved Muscle Energy

Conduct two-day standardized cardiopulmonary exercise testing (2-day CPET) with skeletal muscle biopsies. Measure exercise intolerance (functional decline from Day 1 to Day 2) and parallel muscle ATP levels, creatine phosphate, and mitochondrial respiratory capacity. Hypothesis predicts functional decline with normal resting muscle energy stores.

★ Achievement 4: Aim 4: CNS Control Failure—Electrical Stimulation Bypass

Apply electrical stimulation to quadriceps muscle during exercise testing and compare maximal force output to voluntary contraction efforts. Greater force with electrical stimulation (bypassing CNS control) would demonstrate that exercise intolerance is CNS-mediated rather than peripheral muscle limitation.

★ Achievement 5: Aim 5: Autonomic Demand-Response Failure

Perform tilt table testing with continuous cerebral blood flow monitoring via transcranial Doppler ultrasound. Hypothesis predicts inadequate cerebral autoregulation during orthostatic challenge, explaining orthostatic intolerance and cognitive dysfunction during positional stress.

33.1.6 Key Study Measures

Novel Peripheral Biomarkers

- Hair growth rate (baseline vs. 3-month follow-up): measured as new hair emergence from scalp
- Nail growth rate: fingernail and toenail length change over 3 months
- Baseline metabolic rate: indirect calorimetry in resting state
- Resting muscle ATP: quantified via magnetic resonance spectroscopy (MRS) of vastus lateralis

CNS Neuroimaging

- FDG-PET imaging: whole-brain glucose metabolism with regional analysis
- Regions of interest: prefrontal cortex, posterior cingulate, brainstem, thalamus
- Correlations with symptom severity and cognitive testing

Exercise Physiology (2-Day CPET)

- Day 1 and Day 2 standardized exercise protocols: ramp protocol on stationary cycle ergometer
- Primary outcome: Functional decline from Day 1 to Day 2 (reduced workload capacity)

- Secondary: Peak oxygen consumption, ventilatory threshold, heart rate response
- Muscle biopsy (vastus lateralis): Electron microscopy, respiratory chain enzyme activities, mtDNA copy number, ATP content

Electrical Stimulation Testing

- Quadriceps electrical stimulation: Incremental stimulation intensity to maximal tolerable
- Maximal voluntary contraction (MVC) force: Standard maneuver for comparison
- Outcome measure: Difference in force production (electrical vs. voluntary)

Autonomic Testing

- Tilt table test: 70-degree head-up tilt for 10 minutes or until symptoms limit continued testing
- Transcranial Doppler ultrasound: Continuous measurement of middle cerebral artery blood flow velocity
- Heart rate and blood pressure: Continuous monitoring
- Outcome: Cerebral blood flow decline magnitude during tilt

33.1.7 Expected Outcomes and Implications

If Selective Energy Dysfunction Hypothesis Is Validated

1. Establishes CNS energy metabolism as the primary pathophysiological bottleneck in ME/CFS
2. Reorients research focus from peripheral mitochondrial dysfunction to central neuroinflammation and brain energy metabolism
3. Provides mechanistic rationale for:
 - **Brain-targeted therapies:** Ketogenic diet, intranasal insulin, cerebral blood flow enhancers
 - **Pacing strategies:** Energy envelope approaches are evidence-based rather than speculative
 - **Contraindications:** “Push through” approaches are contraindicated due to CNS energy limitation
4. Enables biomarker-based diagnosis: FDG-PET or 2-day CPET patterns could serve as diagnostic tools
5. Informs personalized medicine: Stratification by CNS involvement severity (mild to severe) guides treatment intensity
6. Generates publications in high-impact journals targeting **Nature Medicine**, **Lancet Neurology**, or **Brain**

If Hypothesis Is Refuted

1. Still advances field with high-quality multi-modal data
2. Negative findings rule out major mechanistic model, redirecting research toward alternatives
3. Enables alternative hypothesis generation based on empirical data
4. Characterizes biomarker patterns in a well-phenotyped cohort for future studies

33.1.8 Budget and Timeline Overview

Total Budget

\$2.15M over 3 years, with allocation:

- Personnel (PI, Co-Investigators, coordinator, assistant, statistician): \$960K
- Brain imaging (PET): \$450K
- Exercise physiology and muscle biopsies: \$350K
- Equipment and supplies: \$190K
- Participant compensation (\$650 per person): \$80K
- Travel, publications, regulatory: \$90K
- Indirect costs (30% institutional rate): \$637K

Timeline

- **Year 1:** Institutional setup, IRB approval, equipment procurement, pilot phase (n=10)
- **Year 2:** Main enrollment (target n=36 additional ME/CFS patients and 36 controls)
- **Year 3:** Final data collection, analysis, manuscript preparation
- **Publication plan:** 4–5 papers targeting *Nature Medicine*, *Lancet Neurology*, *Brain*, *Journal of Applied Physiology*

33.1.9 Funding and Implementation

This proposal is currently in draft form and is being developed for submission to major funding agencies. Target funding sources include:

- **NIH:** R01 grant mechanism (October 2026 cycle)
- **Private foundations:** Solve M.E. Initiative, Open Medicine Foundation
- **International sources:** European ME/CFS research consortia

Implementation requires identification of a principal investigator and institutional home with research infrastructure and expertise in neuroimaging, exercise physiology, and ME/CFS patient populations. The full detailed proposal (50+ pages) includes complete methods, statistical analysis plan, regulatory considerations, and appendices, and is available in the project staging area (see `.claude/content-staging/research-proposal-selective-dysfunction.md`).

33.2 Secondary Analysis Opportunities: Leveraging Existing Datasets

33.2.1 Rationale for Secondary Analysis

The multi-modal study described in Section 33.1 represents the definitive test of the Selective Energy Dysfunction Hypothesis but requires three years and approximately \$2.15M to execute. A complementary near-term strategy leverages existing ME/CFS research datasets that already contain measurements relevant to the hypothesis's core predictions. Secondary analyses can provide preliminary evidence, identify methodological challenges, and generate pilot data for funding applications within months rather than years, with no new data collection required for several key predictions.

33.2.2 Prediction-to-Dataset Coverage

Four of the seven core predictions (brain hypometabolism, exercise demand-response failure, autonomic demand-response failure, and CNS coordination failure) can be tested through secondary analysis of existing datasets. Table 33.1 maps these predictions to available resources.

Table 33.1: Coverage of core predictions by existing ME/CFS datasets

Prediction	NIH 2024	CPET DB	Autonomic	Brain PET	UK Biobank
Brain hypometabolism	✓	–	–	✓✓	Partial
Exercise demand-response	✓	✓✓	–	–	–
Resting muscle ATP preserved	–	–	–	–	–
ATP scaling failure	✓	✓✓	–	–	–
Autonomic demand-response	✓	–	✓✓	–	Partial
CNS coordination failure	✓✓	–	–	✓	–
Hair/nail growth normal	–	–	–	–	–

✓✓ = ideal dataset for this prediction; ✓ = can test prediction; Partial = incomplete coverage; – = not testable.

Three predictions (resting muscle ATP, electrical stimulation versus voluntary force, and hair/nail growth rate) require novel measurement protocols and are addressed exclusively in the multi-modal study.

33.2.3 Priority Datasets

Tier 1: Highest Feasibility

NIH Intramural ME/CFS Study (Walitt et al. 2024) This deep-phenotyping study enrolled 17 ME/CFS patients and 21 healthy controls, collecting comprehensive measurements including two-day cardiopulmonary exercise testing, brain fMRI, cerebrospinal fluid catecholamines and metabolites, autonomic function tests, motor cortex excitability via transcranial magnetic stimulation, and cognitive testing [5]. Published findings—reduced CSF norepinephrine and dopamine, reduced temporal-parietal junction activation, motor cortex hyperactivity, and Day 2 CPET functional decline—are broadly consistent with the Selective Energy Dysfunction framework (interpreted here as post-hoc alignment rather than a priori confirmation). A secondary analysis could systematically test the CNS-peripheral selectivity pattern: correlating catecholamine levels with symptom domains (CNS-dependent versus autonomous processes), comparing motor cortex excitability with peripheral force production, and examining demand-response patterns in the 2-day CPET data stratified by catecholamine subgroups. The full dataset may be accessible through NIH data-sharing mechanisms (e.g., dbGaP).

Published Two-Day CPET Meta-Analysis Multiple research groups have published studies demonstrating significant Day 2 functional decline in ME/CFS using standardized 2-day CPET protocols [1, 140]. A systematic meta-analysis of published summary statistics would directly test the demand-response failure prediction: quantifying Day 2 decline magnitude, assessing heterogeneity by symptom domain (cognitive-predominant versus physical-predominant), and characterizing recovery time as a proxy for demand-response failure duration. This analysis requires no data-sharing agreements and can be initiated immediately using published data.

Tier 2: Medium Feasibility

Van Campen Autonomic Studies A series of studies by van Campen and colleagues characterizes cerebral blood flow via transcranial Doppler during orthostatic challenge in cohorts of 100–400 ME/CFS patients [172, 254, 4]. Published findings show that 91% of patients with a normal heart rate and blood pressure response exhibit abnormal cardiac output and cerebral blood flow patterns during tilt [4], with 82% showing CBF abnormalities even when heart rate and blood pressure remain normal [172]. This preserved-baseline, impaired-challenge pattern is directly consistent with the selective demand-response failure prediction. Individual patient trajectory data (baseline, tilt, recovery) could provide a granular test of the hypothesis's autonomic component; accessing the full dataset would require international collaboration.

UK Biobank ME/CFS Cohort The UK Biobank contains a substantial subcohort of self-reported ME/CFS cases with standardized measurements including wrist accelerometry, cognitive testing, blood biomarkers, and brain MRI in a subset. The large sample size enables detection of activity-pattern abnormalities, cognitive demand-response patterns, and CNS biomarker correlations, though the absence of clinically validated ME/CFS diagnoses

limits interpretability. Access is available to approved researchers through an established application process.

Tier 3: Exploratory

Brain PET/SPECT Meta-Analysis Published neuroimaging studies report regional brain hypometabolism and neuroinflammation in ME/CFS, including evidence of neuroinflammation elevation of 45–199% above controls in the cingulate cortex, hippocampus, amygdala, thalamus, midbrain, and pons [441]. A meta-analysis of published regional findings could identify consistently affected brain regions across studies and characterize subtype patterns. Because raw imaging data is generally not publicly available due to regulatory constraints, this approach relies on published summary statistics rather than individual-level data.

33.2.4 Relationship to the Multi-Modal Study

Secondary analyses and the multi-modal study serve complementary roles. Secondary analyses can generate preliminary evidence on four of the seven predictions within 6–18 months, identify subtype patterns that should be accounted for in primary study design, and provide pilot data for grant applications at low cost. The multi-modal study remains the definitive experimental test of the hypothesis’s full prediction set, and is the only pathway to testing the three predictions that require novel measurement protocols.

33.3 Pediatric-Adult ME/CFS Comparison Study

33.3.1 Background and Rationale

The striking disparity between pediatric and adult ME/CFS recovery rates represents one of the most important clues to understanding recovery mechanisms. While estimates vary by study and definition, pediatric recovery rates of 54–94% contrast sharply with adult rates of $\leq 22\%$ [393]. This difference persists even when controlling for disease duration, suggesting that age-related biological factors—not merely time since onset—determine recovery probability.

Several explanations for this differential have been proposed: developmental plasticity allowing biological “resetting” in younger patients (see the Glial Maturation Window hypothesis, Speculation 8), active immune development enabling clearance of pathological processes (Hypotheses 7.6.3 and 7.5.1), higher metabolic reserves in children, or greater regenerative capacity across multiple organ systems. However, no study has systematically compared the biological profiles of pediatric and adult ME/CFS patients to identify specific mechanisms underlying differential recovery.

This study would provide the first comprehensive cross-sectional comparison of biological features between pediatric and adult ME/CFS patients, generating hypotheses about which systems drive recovery and informing development of targeted interventions.

33.3.2 Hypotheses

~ Hypothesis 2: Biological Plasticity Differential

Pediatric ME/CFS patients will demonstrate preserved biological plasticity compared to adult patients, manifest as:

1. Lower epigenetic age acceleration (more youthful methylation patterns relative to chronological age)
2. Higher naive T cell proportions (greater immune reserve)
3. Greater mitochondrial respiratory capacity
4. Higher metabolic flexibility
5. Greater autonomic adaptability (higher HRV)
6. More diverse hematopoietic stem cell clonality

These differences will persist after controlling for disease duration and severity, indicating that age-related biology—not disease stage—underlies the recovery differential.

33.3.3 Study Design

Design Overview

This is a cross-sectional observational study comparing biological profiles between pediatric/adolescent and adult ME/CFS patients. The study includes both a discovery phase (comprehensive profiling) and a validation phase (replication in independent cohort).

Participants

Inclusion Criteria All participants:

- ME/CFS diagnosis meeting IOM 2015 criteria (or pediatric equivalent)
- Disease duration 6 months to 5 years (to minimize confounding by duration)
- Stable disease (no major change in severity over past 3 months)
- Able to provide informed consent (parental consent for minors)

Pediatric cohort (n=100):

- Age 10–17 years at enrollment
- Tanner stage documented

Adult cohort (n=100):

- Age 25–55 years at enrollment
- Premenopausal women or age-matched men

Exclusion Criteria

- Alternative diagnosis explaining symptoms
- Active infection at time of assessment
- Immunosuppressive medication within past 3 months
- Pregnancy or lactation
- Unable to tolerate study procedures
- Severe psychiatric comorbidity precluding participation

Stratification Within each age group, participants will be stratified by:

- Severity (mild, moderate, severe using Bell scale)
- Trigger type (post-infectious vs. other)
- Sex (target 70% female in each group, reflecting epidemiology)

Control Groups

- **Healthy controls:** 50 pediatric, 50 adult, matched for age and sex
- **Disease controls:** 25 pediatric, 25 adult with other post-viral fatigue syndromes (recovered from acute infection but with persistent fatigue not meeting ME/CFS criteria)

33.3.4 Measures**Epigenomic Assessment**

- Genome-wide DNA methylation via Illumina EPIC array
- Epigenetic age calculation (Horvath, GrimAge, PhenoAge clocks)
- Targeted methylation at immune-related genes
- Histone modification assays (H3K4me3, H3K27ac) at selected loci

Immune Profiling

- Extended flow cytometry panels:
 - T cell subsets: naive (CD45RA⁺CCR7⁺), central memory, effector memory, TEMRA, exhaustion markers (PD-1, CTLA-4, LAG-3)
 - B cell subsets: naive, memory, plasmablasts, CD21^{lo} atypical memory
 - NK cell subsets: CD56^{bright} vs. CD56^{dim}, cytotoxicity markers
 - Monocyte subsets: classical, intermediate, non-classical
 - T regulatory cells: CD4⁺CD25^{hi}FoxP3⁺
- Recent thymic emigrants (CD31⁺ naive CD4 T cells)
- NK cell cytotoxicity functional assay
- T cell proliferation assay

- Cytokine production capacity (intracellular staining after stimulation)
- Autoantibody panel: GPCR autoantibodies, ANA, anti-neuronal antibodies
- Inflammatory markers: high-sensitivity cytokine panel (30+ cytokines), CRP, ESR

Mitochondrial Function

- PBMC respirometry (Seahorse XF assay): basal respiration, maximal capacity, spare respiratory capacity, ATP-linked respiration
- Plasma acylcarnitine profile
- Lactate:pyruvate ratio
- CoQ10 levels
- Muscle biopsy (optional subset, n=20 per group): electron microscopy, respiratory chain enzyme activities, mtDNA copy number

Metabolomic Profiling

- Untargeted plasma metabolomics (LC-MS/MS)
- Targeted panels: amino acids, organic acids, lipids
- Metabolic flexibility assessment: RER dynamics during standardized mild challenge
- Fasting insulin, glucose, HOMA-IR

Autonomic Assessment

- 24-hour Holter monitoring with HRV analysis
- NASA Lean Test (10-minute stand)
- Baroreflex sensitivity
- Pupillometry

Stem Cell and Regenerative Markers

- TCR/BCR repertoire diversity via immunosequencing
- Circulating progenitor cells (CD34⁺)
- Telomere length (flow-FISH)
- Senescence markers: p16^{INK4a} expression, senescence-associated secretory phenotype (SASP) markers

Clinical Assessment

- DSQ-PEM (DePaul Symptom Questionnaire)
- Bell Disability Scale
- MFI (Multidimensional Fatigue Inventory)
- SF-36

- Pediatric Quality of Life Inventory (PedsQL) for pediatric cohort
- Detailed medical history and physical examination
- 7-day actigraphy

33.3.5 Outcomes

Primary Outcomes

1. Composite Recovery Potential Index (RPI) score (see Section 32.2)
2. Individual RPI component scores
3. Between-group differences in each biological domain

Secondary Outcomes

1. Correlations between biological markers and clinical severity
2. Identification of biological features unique to pediatric ME/CFS
3. Identification of biological features associated with shorter disease duration
4. Exploratory subtype identification via unsupervised clustering

33.3.6 Analysis Plan

Primary Analysis

Between-group comparisons (pediatric vs. adult) using:

- ANCOVA adjusting for disease duration, severity, and sex
- Effect sizes (Cohen's *d*) and confidence intervals
- False discovery rate correction for multiple comparisons

Secondary Analyses

- Mediation analysis: Does any biological factor mediate the age-recovery relationship?
- Network analysis: How do biological systems interact differently in pediatric vs. adult patients?
- Machine learning: Can biological profiles classify patients by age group? What features drive classification?
- Correlation with clinical measures: Which biological features predict symptom severity?

Power Analysis and Sample Size Justification

With n=100 per group:

- 80% power to detect Cohen's $d=0.40$ (medium effect) at $\alpha=0.05$ for continuous outcomes
- 80% power to detect 15% difference in proportions
- Sufficient for exploratory subgroup analyses (n=25+ per subgroup)

Based on the dramatic difference in recovery rates (54–94% vs. $\leq 22\%$), we anticipate large effect sizes ($d > 0.8$) for biologically relevant differences, making n=100 per group well-powered.

33.3.7 Ethical Considerations

Pediatric-Specific Protections

- Parental consent plus child assent required
- Procedures minimized to reduce burden on ill children
- Home visits offered for severely affected participants
- Child life specialist available during procedures
- Mandatory rest periods during assessment days
- Parents may remain present for all procedures

General Protections

- IRB approval at all participating sites
- DSMB oversight
- Procedures adapted to patient capacity (no procedures that would cause PEM)
- Results returned to participants who request them (with genetic counseling as appropriate)
- Samples stored in biorepository with consent for future research

33.3.8 Expected Outcomes and Implications

If the hypothesis is supported, this study would:

1. Identify specific biological systems that differ between pediatric and adult ME/CFS patients
2. Generate therapeutic targets for interventions aimed at “restoring” adult systems to more youthful states
3. Validate the Recovery Potential Index as a prognostic tool
4. Inform design of the Aggressive Early Intervention Trial (Section [33.4](#))

If the hypothesis is not supported (no systematic biological differences), this would suggest that the recovery differential stems from psychosocial factors, disease recognition/treatment timing, or other non-biological mechanisms—itself an important finding that would redirect research priorities

33.4 Aggressive Early Intervention Trial

33.4.1 Background and Rationale

The Window of Opportunity Hypothesis

The pediatric recovery data and the Recovery Capital model (Speculation 23) converge on a critical insight: recovery potential may be time-limited. If ME/CFS involves progressive “hardening” of pathological states—through epigenetic stabilization, autoantibody establishment, stem cell exhaustion, and neural pathway consolidation—then there may exist a window of opportunity during which aggressive intervention can prevent this hardening and maximize recovery probability.

Several lines of evidence support this window concept. Recovery rates decline with disease duration across all age groups, suggesting a time-dependent process of chronification. Pediatric patients, who are diagnosed and treated more quickly relative to their disease course, have dramatically better outcomes. Preliminary evidence suggests that early aggressive treatment of orthostatic intolerance in children produces better outcomes than delayed treatment. The biological mechanisms proposed in the Recovery Capital model (epigenetic changes, immune exhaustion, stem cell depletion) are all progressive and potentially irreversible beyond certain thresholds.

Current Standard of Care Limitations

Current ME/CFS management is largely reactive rather than proactive. Patients often experience diagnostic delays of months to years, during which they may worsen through inappropriate activity recommendations. Even after diagnosis, treatment is typically incremental—addressing symptoms one at a time, with conservative dosing and slow titration. While this approach minimizes adverse effects, it may forfeit the window of opportunity when biological plasticity is maximal.

~ Hypothesis 3: Front-Loading Treatment

Aggressive, comprehensive intervention initiated within 12 months of ME/CFS symptom onset can prevent the establishment of permanent pathological states and significantly increase recovery rates compared to standard incremental care. The earlier and more comprehensive the intervention, the greater the recovery probability.

33.4.2 Study Objectives

Primary Objective

To determine whether aggressive multimodal intervention initiated within 12 months of ME/CFS symptom onset increases the proportion of patients achieving recovery at 2 years compared to standard care.

Secondary Objectives

1. To compare functional outcomes between groups at 6, 12, 18, and 24 months
2. To assess the safety and tolerability of aggressive early intervention
3. To identify predictors of response to early intervention
4. To evaluate changes in biological markers (RPI components) with treatment
5. To assess cost-effectiveness of aggressive versus standard care

33.4.3 Study Design

Design Overview

This is a randomized, controlled, parallel-group trial comparing aggressive multimodal intervention to standard care in adults with early-stage ME/CFS. The trial is open-label due to the nature of the interventions, with blinded outcome assessment for primary endpoints.

Participants

Inclusion Criteria

- Age 18–50 years
- ME/CFS diagnosis meeting IOM 2015 criteria
- Symptom onset within preceding 12 months (documented by medical records or detailed history)
- Mild to moderate severity (Bell scale 40–70)
- Able to attend study visits
- Willing to adhere to assigned treatment arm
- Informed consent provided

Exclusion Criteria

- Severe ME/CFS (Bell scale <40) at enrollment
- Alternative diagnosis explaining symptoms
- Contraindication to any study medications
- Pregnancy, planned pregnancy, or breastfeeding
- Active substance abuse
- Major psychiatric illness requiring hospitalization within past year
- Unable to comply with study procedures
- Participation in another interventional trial

Sample Size

Target enrollment: n=100 (50 per arm)

Power Calculation Assumptions:

- Recovery rate in standard care arm: 15% (based on adult ME/CFS literature)
- Clinically meaningful recovery rate in intervention arm: 40% (based on pediatric data suggesting aggressive early treatment approaches pediatric outcomes)
- Two-sided $\alpha=0.05$, power=80%
- 15% dropout rate

Required sample size: 43 per arm; inflated to 50 per arm for dropout.

This is an ambitious target difference, but the hypothesis predicts a substantial effect if the window of opportunity concept is valid. If the true effect is smaller, this study would be underpowered, and results would inform sample size for a larger definitive trial.

Randomization

1:1 randomization to intervention versus standard care, stratified by:

- Sex (male/female)
- Baseline severity (Bell 40–55 vs. 56–70)
- Trigger type (post-infectious vs. other)

Central randomization via web-based system with concealed allocation.

33.4.4 Interventions

Aggressive Multimodal Intervention Arm

The intervention arm receives a comprehensive, front-loaded treatment protocol addressing all major pathophysiological mechanisms simultaneously. This approach contrasts with the typical sequential, incremental approach to ME/CFS management.

Component 1: Maximal Orthostatic Intolerance Management

- **Immediate hydration protocol:** Minimum 2.5L fluid daily with 3–5g sodium supplementation (adjusted for blood pressure)
- **Compression garments:** Waist-high graduated compression (20–30 mmHg) worn during all upright activity
- **Pharmacological support** (initiated within first 2 weeks, not delayed for behavioral approaches to “fail”):
 - Fludrocortisone 0.1–0.2 mg daily for volume expansion
 - Midodrine 5–10 mg TID for vasoconstriction
 - Ivabradine 5–7.5 mg BID if heart rate remains elevated despite above
 - Pyridostigmine 30–60 mg TID if additional support needed
- **Monitoring:** Weekly orthostatic vital signs initially, then monthly

Component 2: Strict Pacing Protocol

- **Activity monitoring:** Continuous accelerometry with heart rate tracking
- **Heart rate-guided pacing:** Activity limited to maintain HR below aerobic threshold (typically 55–60% of age-predicted max)
- **Energy envelope training:** Formal education on energy management with weekly coaching sessions for first 3 months
- **Crash prevention:** Mandatory rest periods; pre-emptive reduction of activity when early warning signs detected
- **Goal:** Zero crashes during treatment period (each crash consumes Recovery Capital)

Component 3: Sleep Optimization

- Sleep study (home-based) to identify treatable disorders
- **Sleep hygiene intervention:** Standardized protocol with weekly adherence monitoring
- **Pharmacological support as needed:**
 - Low-dose trazodone (25–100 mg) or mirtazapine (7.5–15 mg) for sleep maintenance
 - Melatonin 0.5–3 mg for circadian issues
 - CPAP/BiPAP if sleep apnea identified
- **Target:** 7–9 hours sleep with ≥85% sleep efficiency

Component 4: Anti-Inflammatory /Immune Modulation

- **Low-dose naltrexone:** Titrate from 0.5 mg to 4.5 mg over 4 weeks
- **Mast cell stabilization:** H1 antihistamine (cetirizine 10 mg or equivalent) + H2 antihistamine (famotidine 40 mg daily)
- **Omega-3 fatty acids:** 2–4 g EPA+DHA daily
- **Anti-inflammatory diet:** Mediterranean-style, with elimination of identified food sensitivities
- **If elevated inflammatory markers:** Consider short-course oral corticosteroids (prednisone 20 mg × 5 days) or colchicine 0.5 mg BID

Component 5: Mitochondrial Support

- CoQ10 (ubiquinol) 200–400 mg daily
- NAD⁺ precursor: NR or NMN 500–1000 mg daily
- D-ribose 5 g TID
- B vitamin complex including B12 (methylcobalamin) and folate (methylfolate)
- Acetyl-L-carnitine 1000–2000 mg daily

Component 6: Targeted Therapy Based on Phenotype

- **If elevated GPCR autoantibodies:** Referral for immunoadsorption or consideration of off-label rituximab (if available through compassionate use)
- **If viral reactivation markers:** Valacyclovir 1000 mg TID for 6 months
- **If small fiber neuropathy documented:** IVIG consideration (if accessible)
- **If significant MCAS features:** Escalate mast cell stabilization (cromolyn, ketotifen)

Coordination and Monitoring

- Dedicated care coordinator for each patient
- Weekly telehealth check-ins for first 3 months, then biweekly
- Monthly in-person visits with comprehensive assessment
- Rapid response protocol for adverse events or crashes

Standard Care Arm

Participants in the standard care arm receive current best-practice management as described in existing ME/CFS guidelines:

- Education about ME/CFS and pacing (single session)
- Symptom-based medication as clinically indicated

- Orthostatic intolerance management: behavioral approaches first, medications added if behavioral approaches insufficient after 4–6 weeks
- Sleep hygiene education
- Treatment of comorbidities
- Visits every 3 months

Standard care represents the “sequential, conservative” approach that is currently typical for ME/CFS management.

33.4.5 Outcomes

Primary Outcome

Recovery at 24 months, defined as:

1. No longer meeting IOM criteria for ME/CFS (assessed by blinded clinician)
2. Bell Disability Scale ≥ 80 (able to work/attend school full-time with minor symptoms)
3. Patient self-report of “recovered” or “nearly recovered”
4. Sustained for ≥ 3 months at time of 24-month assessment

All four criteria must be met for classification as “recovered.”

Secondary Outcomes

- Bell Disability Scale score at 6, 12, 18, 24 months
- SF-36 physical and mental component scores
- DSQ-PEM crash frequency and severity
- Days per month with significant activity limitation
- Employment/educational status
- Recovery Potential Index component changes from baseline
- Time to sustained improvement (Bell scale increase ≥ 20 points for ≥ 3 months)

Safety Outcomes

- Adverse events (all, serious, related to intervention)
- Medication discontinuations due to intolerance
- Disease worsening (Bell scale decrease ≥ 20 points)
- Hospitalizations
- Emergency department visits

33.4.6 Safety Monitoring

Data Safety Monitoring Board

An independent DSMB will review safety data every 6 months and conduct interim efficacy analysis at 50% enrollment.

Stopping Rules

- Significantly higher rate of serious adverse events in intervention arm
- Significantly higher rate of disease worsening in intervention arm
- Clear evidence of benefit or futility at interim analysis (O'Brien-Fleming boundaries)

Known Risks

- Fludrocortisone: Hypokalemia, hypertension, edema
- Midodrine: Supine hypertension, urinary retention
- Ivabradine: Bradycardia, visual disturbances
- LDN: Vivid dreams, transient sleep disturbance
- Multiple supplements: GI upset, interactions

Risk Mitigation

- Baseline screening for contraindications
- Gradual medication titration
- Frequent monitoring during initiation
- Clear instructions for adverse event reporting
- Medication adjustment protocols for common issues

33.4.7 Feasibility Considerations

Recruitment Challenges

- Early-stage ME/CFS patients may not yet have diagnosis; outreach to primary care needed
- Patients may be reluctant to be randomized to standard care; detailed informed consent about clinical equipoise
- 12-month symptom onset window limits eligible population

Mitigation

- Partnership with post-COVID clinics (rapid identification of post-infectious cases)
- Provider education campaign
- Clear communication that standard care is current best practice, not inferior care

Intervention Complexity

The multimodal intervention is complex and requires significant coordination.

Mitigation

- Detailed protocol manual
- Centralized training for study staff
- Dedicated care coordinators
- Standardized escalation pathways

Cost

The intervention arm is more expensive than standard care due to medications, supplements, monitoring, and coordination.

Mitigation

- Budget includes medication/supplement provision
- Cost-effectiveness analysis will inform future implementation
- If effective, early recovery reduces long-term healthcare costs

33.4.8 Expected Outcomes and Implications

If the hypothesis is supported and the intervention arm shows significantly higher recovery rates:

1. This would provide first evidence that aggressive early intervention can substantially alter ME/CFS prognosis
2. It would establish a new treatment paradigm emphasizing front-loading of comprehensive therapy
3. It would generate data on which intervention components are most important (through exploratory analyses)
4. It would inform cost-effectiveness analyses for healthcare system implementation
5. It would provide urgency for earlier diagnosis, as the window of opportunity is time-limited

If the hypothesis is not supported:

1. This would suggest that recovery potential is determined by factors other than treatment timing/intensity
2. It would redirect research toward identifying the subgroup (if any) that responds to early aggressive treatment
3. Safety and tolerability data would still inform clinical practice
4. Biological marker data would contribute to understanding of ME/CFS pathophysiology

△ Warning 1: Ethical Considerations

This trial involves assigning some patients to standard care while others receive aggressive intervention. This is ethically justified only because clinical equipoise exists: we do not currently know whether aggressive early intervention improves outcomes. If preliminary data strongly favored one approach, equipoise would be lost and randomization would become unethical. The DSMB will monitor for loss of equipoise throughout the trial.

33.5 Crash Impact on Recovery Biomarkers Study

33.5.1 Background and Rationale

The Recovery Capital model (Speculation 23) proposes that patients possess finite biological reserves that deplete with each crash episode. If correct, crash frequency and severity should correlate with accelerated decline in Recovery Potential Index (RPI) components over time. This study would test this hypothesis directly in a pediatric cohort, where the range of outcomes (recovery vs. chronification) is wide enough to detect biomarker-outcome relationships.

33.5.2 Hypothesis

~ Hypothesis 4: Crash-Induced Recovery Capital Depletion

Higher frequency of PEM crashes in pediatric ME/CFS patients will correlate with faster decline in RPI component biomarkers over 2 years, independent of baseline severity. Patients who experience fewer crashes will maintain higher RPI scores and have greater probability of recovery.

33.5.3 Study Design

Design Overview

Prospective observational cohort study with serial biomarker assessment and crash tracking.

Participants

- n=50 pediatric/adolescent ME/CFS patients (ages 10–17)
- Disease duration 6 months to 3 years at enrollment
- Mild to moderate severity (able to attend quarterly study visits)
- Parental consent plus child assent

Assessment Schedule

- **Baseline:** Full RPI component panel (epigenetic age, naive T cell proportion, telomere length, HRV metrics, metabolic flexibility assessment), clinical severity, symptom measures
- **Quarterly (every 3 months):** Abbreviated RPI panel (HRV, selected immune markers), symptom questionnaires, crash diary review
- **Annually (12 and 24 months):** Full RPI panel, comprehensive clinical assessment
- **Continuous:** Wearable activity monitoring, electronic crash diary with severity ratings

Crash Documentation

Participants (with parental assistance) will maintain electronic crash diaries including:

- Date of crash trigger (exertion event)
- Type of trigger (physical, cognitive, emotional, mixed)
- Crash severity (1–10 scale, anchored descriptions)
- Recovery duration (days to return to baseline)
- Classification per crash severity tier (Table 18.3 from treatment chapter)

33.5.4 Outcomes**Primary Outcome**

Correlation between cumulative crash burden (sum of severity-weighted crashes) and change in composite RPI score from baseline to 24 months.

Secondary Outcomes

- Correlation of crash burden with individual RPI components
- Association between crash burden and 24-month recovery status
- Time-varying analysis: Does crash burden in months 0–12 predict RPI decline in months 12–24?
- Threshold analysis: Is there a crash burden threshold beyond which RPI decline accelerates?

33.5.5 Analysis Plan

- Mixed-effects models with random intercepts for subjects to assess RPI trajectory
- Crash burden as time-varying covariate
- Adjustment for baseline severity, age, sex, disease duration
- Sensitivity analyses with different crash severity weighting schemes

33.5.6 Sample Size Justification

With $n=50$ and 3 timepoints per subject (150 observations):

- 80% power to detect correlation $r=0.35$ between crash burden and RPI change at $\alpha=0.05$
- Sufficient for exploratory subgroup analyses

33.5.7 Expected Outcomes

If the hypothesis is supported, this study would:

1. Provide first direct evidence that crashes deplete measurable biological reserves
2. Validate crash prevention as disease-modifying intervention
3. Identify which RPI components are most crash-sensitive
4. Inform clinical recommendations about crash prevention intensity

33.6 Orthostatic Intolerance Treatment Durability Study

33.6.1 Background and Rationale

Orthostatic intolerance (OI) treatment in pediatric ME/CFS produces substantial symptom improvement in many patients. However, the durability of these improvements after medication withdrawal is unknown. Two possibilities exist:

1. **Functional recalibration:** OI treatment during the developmental window may enable permanent autonomic system recalibration, allowing medication withdrawal with sustained improvement
2. **Symptomatic suppression only:** Treatment merely suppresses symptoms while active; withdrawal leads to prompt relapse

Distinguishing these possibilities has major clinical implications. If recalibration occurs, children could potentially discontinue medications after a period of stability. If not, long-term treatment may be necessary.

33.6.2 Hypothesis

~ Hypothesis 5: OI Treatment Durability in Pediatric Patients

Pediatric ME/CFS patients who achieve stable clinical response to OI medications for ≥ 6 months will maintain $\geq 70\%$ of their improvement 3 months after gradual medication withdrawal, reflecting functional recalibration of autonomic systems rather than mere symptom suppression.

33.6.3 Study Design

Design Overview

Single-arm prospective study with structured medication withdrawal and outcome assessment.

Participants

- n=50 pediatric ME/CFS patients (ages 10–17)
- Currently on stable OI medication regimen (fludrocortisone, midodrine, or combination) for ≥ 6 months
- Clinical response documented (improvement in orthostatic symptoms, functional capacity)
- No change in OI medications for past 3 months
- Willing to attempt medication withdrawal

Exclusion Criteria

- Severe ME/CFS (cannot tolerate potential symptom worsening)
- Parental or patient unwillingness to risk symptom relapse
- Medical indication for continued OI treatment independent of ME/CFS

Withdrawal Protocol

1. **Baseline assessment:** Full OI evaluation (NASA Lean Test, HRV, symptom scales), functional capacity
2. **Weeks 1–4:** 50% dose reduction of all OI medications
3. **Weeks 5–8:** Discontinue remaining medications
4. **Week 12 (3 months post-withdrawal):** Primary endpoint assessment
5. **Escape protocol:** If intolerable symptoms at any point, return to prior effective dose; patient classified as “relapse”

33.6.4 Outcomes

Primary Outcome

Proportion of patients maintaining $\geq 70\%$ of baseline improvement (measured by composite OI symptom score and functional capacity) at 3 months post-withdrawal without resuming medications.

Secondary Outcomes

- Time to symptom relapse (if occurs)
- Objective OI measures at 3 months (NASA Lean Test heart rate response, HRV)
- Patient-reported quality of life
- Proportion requiring medication resumption

33.6.5 Analysis Plan

- Primary analysis: Proportion meeting primary endpoint with 95% confidence interval
- Kaplan-Meier survival analysis for time to relapse
- Exploratory: Baseline predictors of sustained improvement (age, disease duration, initial OI severity, HRV parameters)

33.6.6 Expected Outcomes and Implications

If $\geq 50\%$ of patients maintain improvement after withdrawal:

- Supports recalibration hypothesis
- Suggests time-limited treatment protocols may be appropriate in pediatrics
- Informs research on inducing similar recalibration in adults

If $< 30\%$ maintain improvement:

- Suggests ongoing treatment is necessary for sustained benefit
- Informs long-term treatment planning and medication adherence counseling

33.7 HRV-Guided Pacing Randomized Controlled Trial

33.7.1 Background and Rationale

Energy envelope management (pacing) is the cornerstone of ME/CFS symptom management, but standard pacing relies on subjective symptom monitoring and retrospective crash analysis. Patients often discover they have exceeded their envelope only after PEM occurs. Heart rate

variability (HRV) offers a potential objective, prospective measure of autonomic recovery that could guide daily activity decisions before crashes occur.

HRV-guided training is well-established in sports science, where athletes adjust training intensity based on morning HRV readings. Translating this approach to ME/CFS pacing could improve crash prevention and patient confidence in activity decisions.

33.7.2 Hypothesis

~ Hypothesis 6: HRV-Guided Pacing Superiority

Adults with ME/CFS randomized to HRV-guided pacing will experience fewer PEM crashes and achieve better functional outcomes over 6 months compared to those using standard symptom-based pacing.

33.7.3 Study Design

Design Overview

Two-arm parallel-group randomized controlled trial comparing HRV-guided pacing to standard symptom-based pacing.

Participants

- n=60 adults with ME/CFS (ages 18–60)
- Mild to moderate severity (Bell scale 40–70)
- Experiencing ≥ 2 PEM crashes per month on current pacing approach
- Willing to use HRV monitoring device and follow assigned protocol
- Smartphone ownership (for HRV app and data collection)

Randomization

1:1 allocation to HRV-guided or standard pacing, stratified by:

- Baseline severity (Bell 40–55 vs. 56–70)
- Baseline HRV (above vs. below median RMSSD for ME/CFS patients)

Intervention Arms**HRV-Guided Pacing (n=30)**

- Provided with validated HRV sensor (chest strap) and app
- 2-week baseline HRV assessment to establish individual norms
- Daily morning HRV measurement protocol
- Activity calibration based on HRV (Protocol [18.3.3](#))
- Weekly coaching calls for first month to support implementation
- App-based activity recommendations throughout study

Standard Symptom-Based Pacing (n=30)

- Standardized pacing education session
- Activity diary for self-monitoring
- Symptom-based envelope identification
- Weekly coaching calls for first month (attention control)
- Usual pacing approach throughout study

Blinding

Open-label (blinding not feasible for behavioral intervention). Outcome assessors blinded to allocation for primary endpoint assessment.

33.7.4 Outcomes**Primary Outcomes**

- PEM crash frequency over 6 months (electronic diary)
- Functional capacity at 6 months (Bell Disability Scale)

Secondary Outcomes

- Crash severity when crashes occur
- Patient-reported pacing confidence (validated scale)
- Quality of life (SF-36)
- Activity levels (actigraphy)
- Protocol adherence (HRV measurement frequency, activity adjustment compliance)

Exploratory Outcomes

- Baseline HRV as moderator of intervention effect
- Learning curve: Does HRV-guided pacing improve over time as patients learn their patterns?
- Interoceptive awareness: Do patients develop better symptom recognition with HRV feedback?

33.7.5 Analysis Plan

- Primary analysis: Intention-to-treat comparison of crash frequency (negative binomial regression) and functional capacity (ANCOVA) between arms
- Per-protocol sensitivity analysis for patients with $\geq 80\%$ HRV measurement adherence
- Pre-specified subgroup analyses by baseline severity and HRV

33.7.6 Sample Size Justification

Based on preliminary data:

- Assumed control arm crash rate: 3 per month (36 over 6 months)
- Clinically meaningful reduction: 40% (to 1.8 per month)
- With $n=30$ per arm: 80% power at $\alpha=0.05$
- Accounts for 15% dropout

33.7.7 Expected Outcomes and Implications

If HRV-guided pacing shows benefit:

- Establishes HRV monitoring as standard of care adjunct
- Provides objective tool for patients and clinicians
- Informs development of HRV-based pacing apps and devices
- Opens research direction for personalized pacing algorithms

If no benefit observed:

- May indicate HRV is insufficiently predictive in ME/CFS
- May suggest need for different HRV protocols or metrics
- Would redirect research toward other objective pacing tools

33.8 Sports Medicine-Adapted Periodization RCT

33.8.1 Background and Rationale

Standard ME/CFS pacing uses flexible, responsive activity adjustments based on symptoms. Sports medicine offers an alternative approach: structured periodization with pre-planned deload cycles. Athletes use deloads (temporary 40–60% reductions in training volume) to prevent overtraining syndrome and promote recovery. The question: Could structured deload cycles improve outcomes for mild-moderate ME/CFS patients compared to standard flexible pacing?

This approach differs fundamentally from graded exercise therapy (GET). GET assumes progressive increases indefinitely; periodization includes mandatory recovery phases. GET ignores PEM; periodization treats PEM as absolute stop signal. The rationale is to test whether structured recovery cycles prevent the metabolic and immune stress accumulation that precipitates crashes.

33.8.2 Hypothesis

~ Hypothesis 7: Structured Periodization for ME/CFS Stability

Mild-moderate ME/CFS patients randomized to sports medicine-adapted periodization (with structured 7–14 day deload cycles every 4–6 weeks) will experience fewer crashes and greater functional stability over 6 months compared to standard flexible pacing.

33.8.3 Study Design

Design Overview

Two-arm parallel-group randomized controlled trial comparing sports-adapted periodization to standard flexible pacing.

Participants

- n=60 adults with ME/CFS (ages 18–60)
- Mild to moderate severity (Bell scale 40–70)
- Stable baseline for ≥ 4 weeks (no recent crashes)
- Comfortable with structured monitoring and data tracking
- Exclusion: Severe/very severe patients, recent major crash (<3 months), active deterioration

Intervention Arms

Sports-Adapted Periodization (n=30)

- 4-week baseline monitoring phase (establish activity capacity)
- Structured 4–6 week cycles: 3–5 weeks baseline activity + 7–14 day deload (50% volume reduction)
- Daily monitoring: resting heart rate, HRV (optional), subjective recovery rating
- Autoregulatory adjustment: deload triggered early if metrics decline
- PEM = immediate deload initiation regardless of schedule
- Weekly check-ins with pacing coach

Standard Flexible Pacing (n=30)

- Standardized pacing education
- Symptom-based activity adjustment (no pre-planned deloads)
- Daily monitoring: subjective symptoms and activity log
- Activity reductions when symptoms worsen
- Weekly check-ins with pacing coach (attention control)

33.8.4 Outcomes

Primary Outcomes

- PEM crash frequency over 6 months
- Functional capacity at 6 months (Bell Disability Scale)

Secondary Outcomes

- Activity consistency (standard deviation of weekly activity levels)
- Crash severity and recovery time
- Quality of life (SF-36)
- Patient confidence in pacing strategy
- Adverse events (worsening of baseline function)

33.8.5 Safety Monitoring

- Monthly functional assessments
- Immediate exit criteria: Any sustained worsening of baseline Bell score by ≥ 10 points
- Data Safety Monitoring Board review at 3 months
- Protocol allows switching from periodization to flexible pacing if unhelpful

33.8.6 Expected Outcomes

If periodization shows benefit:

- Establishes structured deload cycles as evidence-based option for selected patients
- Provides clear protocol for implementation
- Opens research into optimal cycle length and deload depth

If no benefit or harm observed:

- Standard flexible pacing remains evidence-based default
- May indicate that pre-planned cycles cannot accommodate ME/CFS variability
- Redirects focus to real-time adaptive pacing strategies

33.9 Longitudinal Microglial Imaging Study

33.9.1 Background and Rationale

Progressive post-exertional malaise (PEM) worsening suggests cumulative biological damage, particularly in the central nervous system. Microglial activation, a hallmark of neuroinflammation, may represent a measurable correlate of PEM progression. TSPO (translocator protein) positron emission tomography (PET) imaging provides a non-invasive method to quantify microglial activation in vivo. This study would establish TSPO-PET as a biomarker for neuroinflammation severity and its relationship to PEM trajectories.

~ Hypothesis 8: Microglial Activation and PEM Progression

Increasing TSPO-PET signal intensity correlates with PEM recovery time and functional decline. Patients with progressive PEM show significantly higher TSPO-PET signal than stable PEM patients. Low-dose naltrexone (LDN) treatment reduces TSPO-PET signal and slows PEM worsening [602].

33.9.2 Study Design

Design Overview

Prospective longitudinal observational study with serial TSPO-PET imaging and detailed PEM documentation.

Participants

- n=50 ME/CFS patients (ages 18–60)
- Documented PEM with variable severity trajectories (stable, slowly progressive, rapidly progressive)
- Mild to moderate severity (able to tolerate imaging procedures)
- Disease duration ≥ 6 months
- No contraindications to PET imaging

Stratification

Stratified by PEM trajectory at baseline:

- Stable PEM (n=15): Crash frequency and severity unchanged over past 6 months
- Progressive PEM (n=20): Increasing crash frequency or severity over past 6 months
- Rapidly Progressive PEM (n=15): Significant functional decline over past 3 months

33.9.3 Assessment Schedule

- **Baseline:** TSPO-PET imaging, detailed clinical assessment, 6-month pre-baseline PEM diary retrospective review
- **6 months:** TSPO-PET imaging, PEM diary review, functional assessment
- **12 months:** TSPO-PET imaging, comprehensive clinical and biomarker assessment
- **Continuous:** Electronic PEM crash diary with severity ratings (1–10 scale), recovery duration documentation

33.9.4 Measures**TSPO-PET Imaging**

- ^{11}C -PBR28 or ^{18}F -DPA-714 radioligand (TSPO-specific tracers)
- Standardized uptake value (SUV) analysis in predefined regions of interest (basal ganglia, thalamus, brainstem, prefrontal cortex)
- Distribution volume ratio (DVR) to derive binding potential
- Whole-brain voxel-wise analyses to identify activation hotspots

PEM Documentation

- Crash trigger (physical, cognitive, emotional, mixed)
- Pre-crash activity level (hours of exertion)
- Crash severity (1–10 scale, anchored descriptions)
- Recovery duration (days to baseline)
- Associated symptoms (cognitive dysfunction, pain, autonomic symptoms)

Clinical and Functional Measures

- Bell Disability Scale
- DSQ-PEM
- Cognitive assessment (Montreal Cognitive Assessment)
- Autonomic testing (NASA Lean Test, HRV)
- Inflammatory markers (high-sensitivity CRP, cytokine panel)

33.9.5 Outcomes

Primary Outcomes

1. Correlation between TSPO-PET signal intensity at baseline and PEM recovery time at 12 months
2. Differences in baseline TSPO-PET signal between progressive and stable PEM groups
3. Change in TSPO-PET signal from baseline to 12 months as a function of PEM trajectory

Secondary Outcomes

- Correlation between TSPO-PET signal and functional decline (Bell scale change)
- Regional specificity: Which brain regions show signal changes most relevant to PEM?
- Effect of LDN treatment (in patients who elect to initiate) on TSPO-PET signal reduction
- Correlation of TSPO-PET with systemic inflammatory markers

33.9.6 Analysis Plan

- Spearman or Pearson correlations between TSPO-PET SUV and PEM recovery time
- ANOVA comparing TSPO-PET signal across PEM trajectory groups
- Mixed-effects models with random intercepts for subjects to assess PET signal trajectory
- ROI-specific and voxel-wise analyses with multiple comparison correction
- Adjustment for age, sex, disease duration, and baseline severity

33.9.7 Sample Size and Power

With n=50 participants and 3 imaging timepoints per subject:

- 80% power to detect Spearman $\rho=0.35$ between TSPO-PET and PEM recovery time at $\alpha=0.05$
- Sufficient for subgroup analyses by PEM trajectory
- Adequate for exploratory regional analyses

33.9.8 Expected Outcomes and Implications

If correlations are significant:

1. Establishes TSPO-PET as biomarker for neuroinflammation severity in ME/CFS
2. Validates use of TSPO-PET as clinical trial outcome measure
3. Informs mechanism of LDN efficacy (microglial suppression)
4. Identifies patients at high risk for PEM progression

If results are null:

1. Suggests microglial activation is not primary driver of PEM progression
2. Redirects focus toward other neuroinflammatory mechanisms
3. May indicate TSPO is insufficient marker (astrocytic activation, other glia)

33.10 Treatment Sequence RCT: “Brain First” Versus Peripheral-First Approaches

33.10.1 Background and Rationale

ME/CFS pathophysiology involves multiple interconnected systems: central neuroinflammation, peripheral immune dysregulation, metabolic dysfunction, and mast cell activation. Standard care typically addresses these sequentially as symptoms warrant. However, the hierarchical interactions between these systems suggest that addressing the primary driver first (likely central neuroinflammation) might produce superior outcomes compared to peripheral-first approaches.

The “Brain First” hypothesis posits that LDN-induced suppression of central microglial activation creates a permissive environment for recovery of peripheral immune function and metabolic adaptation, whereas beginning with peripheral interventions (mast cell stabilization or metabolic support) may not address the root neuroinflammatory driver.

~ Hypothesis 9: Treatment Sequencing and Hierarchical Recovery

Patients randomized to a “Brain First” sequence (LDA/LDN→Mestinon→peripheral support) show superior 6-month outcomes and faster cognitive improvement compared to “Mast Cell First” (mast cell stabilization first) or “Metabolic First” (metabolic supplementation first) sequences, due to removal of the neuroinflammatory barrier to recovery [602].

33.10.2 Study Design

Design Overview

Three-arm parallel-group open-label RCT with blinded outcome assessment. Medications are administered sequentially in protocol-specified order, with fixed timing of medication introduction.

Participants

- n=120 adults with ME/CFS (ages 18–55)
- Mild to moderate severity (Bell scale 40–70)
- Disease duration 1–10 years
- No prior treatment with LDN, LDA, Mestinon, or extensive mast cell stabilization
- Able to attend monthly in-person visits or have reliable telehealth access
- Willing to be randomized to assigned sequence

Exclusion Criteria

- Severe ME/CFS (Bell scale <40)
- Contraindication to any study medications
- Active infection requiring treatment
- Psychiatric hospitalization within past year
- Substance use disorder
- Participation in other interventional trials

Randomization

1:1:1 randomization to three treatment sequences, stratified by:

- Baseline severity (Bell 40–55 vs. 56–70)
- Sex

33.10.3 Treatment Arms

Arm A: Brain First (n=40)

1. **Months 0–3:** LDN titration (0.5 mg → 4.5 mg) + pacing optimization
2. **Months 3–6:** Add Mestinon (30 mg TID, titrate as tolerated) for autonomic support
3. **Months 6+:** Add peripheral support only if needed (mast cell stabilizers, metabolic support)

Arm B: Mast Cell First (n=40)

1. **Months 0–3:** Ketotifen 1 mg BID + H1 antihistamine (cetirizine 10 mg) + H2 antihistamine (famotidine 40 mg daily)
2. **Months 3–6:** Add standard supportive care (sleep optimization, pacing)
3. **Months 6+:** Add neuroinflammatory support (LDN) if inadequate response

Arm C: Metabolic First (n=40)

1. **Months 0–3:** CoQ10 (ubiquinol) 300 mg daily + D-ribose 5 g TID + amino acid support
2. **Months 3–6:** Extend metabolic protocol, add NAD⁺ precursor (NMN 500 mg daily)
3. **Months 6+:** Add LDN or mast cell stabilization if inadequate response

All arms receive standardized pacing education and sleep optimization protocol beginning at enrollment.

33.10.4 Outcomes

Primary Outcome

DSQ-PEM total score change from baseline to 6 months (larger decrease = improvement).

Secondary Outcomes

- Cognitive function (Montreal Cognitive Assessment) change baseline to 6 months
- Bell Disability Scale change baseline to 6 months
- PEM crash frequency (monthly average over months 4–6)
- Fatigue severity (MFI subscales)
- Time to substantial improvement (DSQ-PEM decrease ≥ 10 points for ≥ 4 weeks)
- Functional capacity improvement trajectory over 6 months

Tertiary Outcomes

- Arm-specific adverse event rates
- Medication discontinuation rates due to intolerance
- Biomarker changes (inflammatory markers, HRV, autonomic measures)

33.10.5 Analysis Plan

- Primary analysis: ANCOVA comparing 6-month DSQ-PEM change across arms, adjusted for baseline severity and sex
- Post-hoc pairwise comparisons with Bonferroni correction (Brain First vs. Mast Cell First; Brain First vs. Metabolic First)

- Secondary analyses: Per-protocol analysis for patients with $\geq 80\%$ medication adherence
- Trajectory analysis: Quadratic mixed-effects models for symptom change over 6 months

33.10.6 Sample Size Justification

With $n=40$ per arm (120 total):

- 80% power to detect Cohen's $d=0.50$ between Brain First and either alternative arm at $\alpha=0.05$ (two-sided, Bonferroni-corrected to 0.025)
- Based on pilot data showing 10-point DSQ-PEM improvement in LDN responders vs. 3-point in controls ($SD=10$)
- Accounts for 15% dropout

33.10.7 Expected Outcomes and Implications

If Brain First arm shows significantly superior outcomes:

1. Establishes neuroinflammation as primary treatment target
2. Informs clinical practice guidelines for ME/CFS medication sequencing
3. Validates hierarchical model of ME/CFS pathophysiology
4. Suggests LDN should be first-line agent for most patients

If all arms show similar outcomes:

1. Suggests sequence is less important than comprehensive treatment
2. Supports individualized approach based on patient phenotype
3. Indicates need for biomarker-driven sequencing strategies

33.11 Infection-Divine Correlation Study

33.11.1 Background and Rationale

Clinical experience suggests that infections trigger PEM and may cause persistent functional decline in ME/CFS patients. However, the quantitative relationship between infection frequency and cumulative baseline functional decline has not been formally characterized. This prospective cohort study would establish whether infections produce irreversible baseline declines and whether early antiviral intervention can mitigate these effects.

~ Hypothesis 10: Infection-Induced Irreversible Decline

Each documented infection in ME/CFS patients produces quantifiable, irreversible baseline functional decline. Early antiviral intervention (<24 hours from symptom onset)

significantly reduces the magnitude of post-infectious decline compared to delayed treatment. Strict infection prevention protocols slow baseline functional decline compared to standard hygiene practices [279].

33.11.2 Study Design

Design Overview

Prospective observational cohort study with standardized infection documentation, treatment timing records, and quarterly functional assessments over 3 years.

Participants

- n=200 adults and adolescents with ME/CFS (ages 12–65)
- Disease duration ≥ 6 months
- Mild to moderate severity (able to participate in quarterly assessments)
- History of ≥ 2 documented infections in prior 12 months (to enrich for infection-prone individuals)
- Ability to document infections with temporal precision

Exclusion Criteria

- Immunosuppressive therapy or active malignancy
- Unable to provide infection documentation
- Severe psychiatric comorbidity precluding informed consent

33.11.3 Assessment Schedule

- **Baseline:** Full clinical assessment, baseline functional status (Bell scale, SF-36, DSQ-PEM)
- **Quarterly (every 3 months):** Functional reassessment, infection history review
- **Continuous:** Infection documentation by patient (illness diary or symptom tracker app)

Duration: 3 years per participant.

33.11.4 Infection Documentation Protocol

Participants document each illness with:

- Date of symptom onset
- Presumed pathogen (viral vs. bacterial, if identifiable)
- Symptoms (upper respiratory, gastrointestinal, systemic)

- Severity (1–10 scale)
- Time from symptom onset to antiviral treatment initiation (if applicable)
- Antiviral agents used and duration
- PEM response (crash triggered yes/no, severity if yes, recovery duration)
- Estimated baseline functional impact 2 weeks post-infection recovery

33.11.5 Measures

Functional Status

- Bell Disability Scale (primary functional outcome)
- SF-36 (secondary functional outcome)
- DSQ-PEM (symptom burden)
- Days per month with significant activity limitation
- Employment/educational status

Infection Data

- Cumulative infection count over study period
- Infection type distribution
- Treatment timing for each infection
- Antiviral vs. untreated infections
- Infection severity distribution

Infection Prevention Practices

- Quarterly questionnaire on hygiene practices (handwashing frequency, sick contacts avoidance, masking in high-risk settings, etc.)
- Categorization: Standard hygiene vs. intensive prevention practices

33.11.6 Outcomes

Primary Outcome

Cumulative functional decline (baseline to 3-year Bell scale change) as a function of cumulative infection burden over 3 years.

Secondary Outcomes

- Functional decline per infection (Bell scale change per documented illness)
- Impact of treatment timing: Functional decline in <24h-treated infections vs. delayed treatment vs. untreated
- Effect of infection prevention: 3-year Bell scale change in intensive prevention vs. standard practice groups
- Infection frequency: Comparison of high-infection vs. low-infection subgroups in matched disease duration cohorts

Exploratory Outcomes

- Dose-response relationship: Does infection count linearly predict functional decline?
- Threshold effect: Is there a critical infection frequency beyond which decline accelerates?
- Infection type specificity: Do certain pathogen types cause greater decline?

33.11.7 Analysis Plan

- Linear regression: Bell scale decline (outcome) vs. cumulative infection count (predictor), adjusted for baseline severity, age, sex, disease duration
- Stratified analysis: Compare decline slopes between early-treatment (<24h) vs. delayed/untreated groups
- Prevention effect: ANCOVA comparing 3-year Bell scale change between intensive prevention and standard hygiene groups, adjusted for baseline infection frequency
- Time-varying analyses: Does infection burden in years 0–1 predict steeper decline in years 1–3?
- Logistic regression: Predictors of experiencing ≥ 1 major functional decline event

33.11.8 Sample Size Justification

With $n=200$ and 12 assessment points over 3 years (2,400 observations):

- 80% power to detect linear association between infection count and Bell scale decline (regression $\beta=1.5$ points per infection, $SD=10$) at $\alpha=0.05$
- Adequate power for subgroup comparisons ($n \geq 50$ per treatment timing category)

33.11.9 Expected Outcomes and Implications

If hypothesis is supported:

1. Quantifies functional cost of infections in ME/CFS
2. Validates aggressive infection prevention as disease-modifying strategy
3. Establishes early antiviral treatment as standard of care intervention

4. Informs prognostic counseling about long-term functional trajectory
5. May support use of prophylactic antivirals in high-risk patients

If infections do not predict decline:

1. Suggests infection-triggered decline is less significant than currently believed
2. Redirects focus toward other mechanisms of disease progression
3. Questions cost-benefit of intensive infection prevention practices

33.12 Sleep-Glymphatic-Neuroinflammation Pathway Study

33.12.1 Background and Rationale

The glymphatic system—the brain’s waste clearance mechanism that is maximally active during sleep—may play a critical role in neuroinflammation control. Poor sleep in ME/CFS could impair glymphatic clearance of neuroinflammatory mediators, perpetuating microglial activation and progressive symptom deterioration. Conversely, treating underlying sleep disorders might interrupt this pathway and slow disease progression.

This study tests whether interventions targeting sleep disorders in ME/CFS patients with documented sleep pathology produce measurable reductions in neuroinflammation and slowing of PEM progression.

~ Hypothesis 11: Sleep as Glymphatic-Neuroinflammation Linchpin

Treating sleep disorders in ME/CFS patients reduces the rate of PEM threshold decline through improved glymphatic clearance and reduced neuroinflammatory burden. Patients with newly treated sleep apnea or sleep disorders will show slower neuroinflammatory marker increase and slower functional decline over 12 months compared to matched untreated controls [73].

33.12.2 Study Design

Design Overview

Randomized, prospective study of ME/CFS patients with documented sleep disorders, comparing immediate treatment to delayed (6-month wait-list control) treatment.

Participants

- n=60 ME/CFS patients (ages 18–60) with documented sleep disorders
- Sleep disorder documentation via home sleep apnea testing: OSA (AHI ≥ 5 events/hour) or other sleep pathology (insomnia with documented sleep fragmentation, periodic breathing, etc.)

- Mild to moderate severity (Bell scale 40–75)
- No prior treatment with CPAP, BiPAP, or formal sleep disorder management
- Willing to be randomized to immediate vs. delayed treatment

Randomization

1:1 randomization to:

- Immediate treatment (n=30): Sleep disorder management initiated at baseline
- Delayed treatment (n=30): Sleep management begins at 6-month mark (wait-list control)

Stratification by sleep disorder type (OSA vs. other).

33.12.3 Interventions

Immediate Treatment Arm

OSA patients (CPAP/BiPAP):

- Polysomnography or split-night study to determine therapeutic pressure
- CPAP or BiPAP initiation with titration protocol
- Mask fitting and desensitization
- Monthly adherence monitoring for first 3 months, then quarterly
- Target: ≥ 4 hours per night use

Insomnia or fragmentation:

- Sleep restriction therapy with gradual sleep window expansion
- Cognitive-behavioral therapy for insomnia (CBT-I) protocol
- Pharmacological support: Trazodone or mirtazapine as needed
- Weekly coaching sessions for first 4 weeks

Delayed Treatment Arm

Standard sleep hygiene education only for first 6 months; treatment as above begins at month 6.

33.12.4 Measures

Sleep Assessment (baseline, 6 months, 12 months)

- Home sleep apnea testing or portable sleep monitoring
- Sleep diary (7-day baseline, 7-day at each timepoint)
- Actigraphy (7-day continuous at each timepoint)
- Pittsburgh Sleep Quality Index (PSQI)

Neuroinflammation Markers (baseline, 6 months, 12 months)

- CSF sampling (optional lumbar puncture subset, n=20 per arm): TNF- α , IL-6, IL-1 β , MCP-1, neopterin
- Serum inflammatory markers: High-sensitivity CRP, IL-6, TNF- α , IL-1 β
- TSPO-PET imaging (subset, n=15 per arm): microglial activation assessment

Clinical Outcome Measures (baseline, 6 months, 12 months)

- PEM crash diary (continuous): Frequency, severity, recovery duration
- Bell Disability Scale
- Cognitive function (Montreal Cognitive Assessment)
- Autonomic function (NASA Lean Test, HRV)
- DSQ-PEM

33.12.5 Outcomes

Primary Outcomes

1. Change in inflammatory marker trajectory (serum IL-6) from baseline to 12 months: Immediate vs. delayed arm
2. PEM threshold stability: Rate of change in PEM severity over 12 months in treated vs. untreated groups

Secondary Outcomes

- Individual inflammatory markers (TNF- α , IL-1 β , CRP)
- CSF markers in subset (if available)
- TSPO-PET signal change in imaging subset
- Bell Disability Scale change baseline to 12 months
- Cognitive function improvement
- PEM crash frequency change

Tertiary Outcomes

- Sleep quality improvement and correlation with inflammatory markers
- CPAP/BiPAP adherence as moderator of treatment effect
- Slow-wave sleep percentage (from polysomnography in subset) and correlation with inflammatory markers

33.12.6 Analysis Plan

- Primary analysis: Linear mixed-effects model comparing serum IL-6 trajectory over 12 months between arms
- Secondary outcomes: ANCOVA for Bell scale and other clinical outcomes at 12 months, adjusted for baseline values
- Mediation analysis: Does change in sleep quality mediate the relationship between treatment arm and inflammation?
- Adherence analysis: Among immediate arm, does CPAP adherence predict inflammatory marker improvements?

33.12.7 Sample Size Justification

With $n=30$ per arm and 3 measurement timepoints:

- 80% power to detect 30% difference in IL-6 slope between arms (assuming $SD=0.8$ in log IL-6) at $\alpha=0.05$
- Based on preliminary data: treated sleep apnea showing 25–35% reduction in inflammatory markers in healthy controls

33.12.8 Expected Outcomes and Implications

If sleep treatment reduces neuroinflammation and slows PEM progression:

1. Establishes sleep as critical disease-modifying factor
2. Validates screening and treating sleep disorders as standard ME/CFS care
3. Identifies glymphatic function as therapeutic target
4. May support clinical guidelines for aggressive sleep disorder management

If sleep treatment shows minimal effect:

1. Suggests neuroinflammation maintenance is not primarily glymphatic-dependent
2. Indicates other mechanisms of microglial activation are primary
3. Redirects focus toward direct microglial targeting (LDN, other agents)

33.13 Metabolic-Immune Crosstalk Study

33.13.1 Background and Rationale

Low-dose naltrexone (LDN) shows promise in ME/CFS but not all patients respond, and some who respond show plateau or loss of benefit. This may reflect metabolic complications emerging during LDN therapy, particularly the development of insulin resistance or glucose intolerance, which could amplify neuroinflammation through metabolic-immune pathways. Concurrent metabolic intervention (metformin or other insulin-sensitizing agents) might preserve or enhance LDN efficacy by preventing metabolic deterioration.

This mixed-methods study examines the metabolic-immune crosstalk in ME/CFS and tests whether metabolic intervention improves or sustains LDN efficacy.

~ Hypothesis 12: Metabolic-Immune Crosstalk in LDN Response

LDA-induced metabolic changes (glucose intolerance, insulin resistance) amplify neuroinflammation and create a therapeutic ceiling that limits cognitive benefit over time. Concurrent metformin administration in patients developing prediabetes will preserve LDN efficacy and enhance cognitive outcomes compared to LDN monotherapy. HbA1c improvements correlate with neuroinflammatory marker reduction [278].

33.13.2 Study Design

This study combines three complementary components:

Component 1: Cross-Sectional Metabolic-Immune Comparison (n=80)

Compare ME/CFS patients with vs. without metabolic syndrome (MS) on inflammatory markers and neuroinflammatory burden.

Participants:

- n=40 ME/CFS without metabolic syndrome
- n=40 ME/CFS with metabolic syndrome (modified NCEP criteria)

Outcomes:

- Inflammatory marker profiles (IL-6, TNF- α , IL-1 β , CRP)
- TSPO-PET signal (subset, n=10 per group)
- Cognitive function (Montreal Cognitive Assessment)
- Functional capacity (Bell scale)

Component 2: Longitudinal Tracking of Metabolic Development (n=60)

ME/CFS patients without baseline metabolic syndrome tracked prospectively as some develop metabolic complications.

Assessment schedule:

- Baseline: HbA1c, fasting glucose, insulin, lipid panel, inflammatory markers
- Every 3 months for 18 months: Metabolic labs, inflammatory markers
- Continuous: Cognitive assessments, functional measures

Outcomes:

- Rate of metabolic deterioration in LDN-treated vs. untreated subgroups
- Correlation between metabolic changes and inflammatory marker changes
- Identification of patients at high risk for metabolic complications

Component 3: Interventional Trial—Metformin in Prediabetic ME/CFS Patients (n=40)

ME/CFS patients with newly identified prediabetes (HbA1c 5.7–6.4%) randomized to metformin vs. placebo.

Randomization: 1:1 to metformin 500 mg BID (target 1000 mg BID) vs. placebo, stratified by concurrent LDN use.

Assessment schedule:

- Baseline: Full metabolic panel, inflammatory markers, cognitive function, functional capacity
- Every 3 months for 12 months: Metabolic labs, inflammatory markers
- Cognitive and functional assessments at baseline, 6 months, 12 months

Primary outcomes (metformin trial):

1. HbA1c change from baseline to 12 months
2. Serum IL-6 change (inflammatory outcome)
3. Cognitive function change (Montreal Cognitive Assessment)

33.13.3 Overall Study Measures

Metabolic Assessment

- Fasting glucose, insulin, HOMA-IR
- HbA1c
- Lipid panel (total cholesterol, LDL, HDL, triglycerides)
- Metabolic syndrome classification (modified NCEP criteria)

Immune/Inflammatory Assessment

- High-sensitivity CRP
- IL-6, TNF- α , IL-1 β
- Monocyte activation markers (CD14⁺CD16^{hi})
- TSPO-PET imaging in subsets

Cognitive and Functional Outcomes

- Montreal Cognitive Assessment (primary cognitive measure)
- Bell Disability Scale
- DSQ-PEM
- Processing speed (DSST—Digit Symbol Substitution Test)

33.13.4 Analysis Plan

Cross-Sectional Component

- T-tests or Mann-Whitney U comparing inflammatory markers between metabolic syndrome and non-syndrome groups
- Correlation analyses between metabolic parameters and inflammatory/cognitive measures
- Effect sizes (Cohen's d) with 95% confidence intervals

Longitudinal Component

- Mixed-effects models with random intercepts for subjects to assess metabolic deterioration trajectory
- Stratified analyses by LDN use vs. non-use
- Time-varying analysis of relationship between metabolic changes and inflammatory marker changes

Interventional Component (Metformin Trial)

- Primary analysis: ANCOVA comparing HbA1c change and IL-6 change between metformin and placebo arms at 12 months, adjusted for baseline values
- Secondary analysis: Cognitive function change (Montreal Cognitive Assessment) at 12 months
- Subgroup analysis: Differential effect in patients with vs. without concurrent LDN
- Adherence analysis: Association between metformin adherence and HbA1c/inflammatory improvements

33.13.5 Sample Size Justification

Cross-Sectional Component

With n=40 per group:

- 80% power to detect Cohen's $d=0.65$ difference in IL-6 between groups at $\alpha=0.05$

Longitudinal Component

With n=60 and 7 measurement timepoints (baseline plus 6 follow-ups):

- Adequate power for trajectory analyses
- Sufficient for subgroup comparisons ($n \geq 30$ per LDN status)

Metformin Trial

With n=20 per arm:

- 80% power to detect 0.5% difference in HbA1c change between arms at $\alpha=0.05$
- Based on expected metformin effect of 0.5–1% HbA1c reduction in prediabetic populations

33.13.6 Expected Outcomes and Implications

If metabolic complications drive LDN plateau/loss of benefit:

1. Establishes metabolic-immune crosstalk as key ME/CFS mechanism
2. Validates metabolic monitoring during LDN therapy
3. Supports concurrent metformin use in prediabetic patients to sustain LDN benefit
4. Informs guidelines for managing LDN-associated metabolic effects
5. May explain treatment resistance in subset of patients

If metabolic intervention does not enhance LDN efficacy:

1. Suggests metabolic changes are consequence rather than driver of neuroinflammation
2. Indicates need for alternative approaches to sustaining LDN response
3. Might redirect focus toward other mechanisms of LDN resistance

33.14 Gut-Vagal Pathway Research Priorities

33.14.1 Background and Rationale

Wirth and Scheibenbogen [225] proposed that gut dysbiosis impairs enterochromaffin cell serotonin synthesis, reducing vagal afferent signalling and contributing to the autonomic dysfunction characteristic of ME/CFS. This model is supported by documented butyrate-producer deficiency in ME/CFS microbiome studies [349, 226] and established mechanisms linking butyrate, enterochromaffin serotonin, and vagal activation [14, 228]. The following research priorities would test this model and identify a “gut-vagal” ME/CFS subtype for targeted intervention.

33.14.2 Priority Study 1: Cross-Sectional Gut-Vagal Phenotyping

Hypothesis

ME/CFS patients with prominent GI symptoms have lower resting HRV, lower butyrate-producing bacterial abundance, and lower platelet serotonin than ME/CFS patients without GI involvement and healthy controls. These markers cluster to identify a distinct gut-vagal subtype.

Design

- **Type:** Cross-sectional case-control, n=100–200 (ME/CFS with GI symptoms, ME/CFS without GI symptoms, healthy controls)
- **Primary measures:** 5-minute resting HRV; stool microbiome (focus on *F. prausnitzii* and butyrate-producer abundance); platelet serotonin; 24-hour urinary 5-HIAA; fecal short-chain fatty acids
- **Secondary measures:** Kynurenine/tryptophan ratio (IDO activity marker; operationalises the EBV-IDO2–gut-vagal chain in Speculation 7: patients with EBV reactivation predicted to show higher k/t ratios, lower platelet serotonin, and worse HRV); EBV serology (IgG titres, EBNA, VCA antibodies; planned subgroup analysis of k/t ratio and HRV stratified by EBV reactivation status); serum tryptase and 24-hour urinary methylhistamine (MCAS markers; tests the bidirectional mast cell–enterochromaffin amplification mechanism); serum chromogranin A; GI symptom severity; fatigue severity; orthostatic tolerance

- **Analysis:** Cluster analysis to identify whether gut-vagal markers co-vary; correlation of butyrate-producer abundance with HRV and fatigue; subgroup analysis of EBV-seropositive vs. EBV-seronegative patients on k/t ratio, platelet serotonin, and HRV; correlation of tryptase and methylhistamine with HRV and fatigue scores
- **Timeline:** 6–12 months

Expected Outcomes

If gut-vagal markers cluster in a subgroup, this establishes the foundation for targeted intervention trials. HRV and stool microbiome testing are both accessible and could serve as simple screening tools for this subtype.

33.14.3 Priority Study 2: Butyrate Supplementation RCT

Hypothesis

Tributylin supplementation at 2 g three times daily (the upper end of the 1–2 g clinical range; see Section 22.9.3 in Chapter 22) improves HRV and reduces fatigue in ME/CFS patients with documented butyrate-producer deficiency, compared to placebo.

Design

- **Type:** Randomised, double-blind, placebo-controlled; n=60 (30 tributyrin, 30 placebo)
- **Duration:** 12 weeks
- **Stratification:** Randomise within strata defined by baseline butyrate-producer abundance (low vs. moderate) to test whether response is greatest in most-deficient patients
- **Primary outcome:** Change in resting HRV at 12 weeks
- **Secondary outcomes:** Fatigue severity (SF-36 vitality, FACIT-Fatigue), GI symptom score, stool microbiome, platelet serotonin, orthostatic tolerance
- **Timeline:** 12–18 months

Expected Outcomes

A positive result would provide the first RCT evidence for a microbiome-vagal mechanism in ME/CFS and support butyrate supplementation as a disease-modifying intervention in the gut-vagal subtype. A null result in the full population but positive result in the low-butyrate stratum would support biomarker-guided treatment selection.

33.14.4 Priority Study 3: HRV-Guided Phenotype Reanalysis

Rationale

Multiple existing ME/CFS cohorts include HRV data. A secondary analysis stratifying by baseline HRV could test whether low-HRV patients have more GI symptoms and different microbiome profiles than high-HRV patients—testing the gut-vagal subtype hypothesis at low cost before embarking on prospective studies.

Design

- **Type:** Secondary analysis of existing datasets
- **Key datasets:** Any ME/CFS cohort with HRV, GI symptom, and microbiome data (e.g., Giloteaux et al. Cornell cohort [226])
- **Analysis:** Correlate HRV with GI symptom severity, butyrate-producer abundance, and fatigue scores; test whether HRV clusters identify a distinct subgroup
- **Timeline:** 3–6 months
- **Cost:** Low (secondary analysis, no new data collection)

33.15 Replication Methodology for Heterogeneous Disease

ME/CFS, like other multisystem chronic illnesses, exhibits profound treatment heterogeneity. Conventional randomized controlled trials powered to detect average treatment effects frequently fail despite strong mechanistic rationale and robust phase II data. This section examines why average-effect replication fails in heterogeneous diseases and proposes advanced designs—basket trials, umbrella trials, adaptive enrichment, and N-of-1 aggregation—suited to ME/CFS subgroup structure.

33.15.1 Why Average-Effect Replication Fails

The fundamental challenge in replicating treatments for heterogeneous diseases is that a population average effect may not apply to any identifiable subgroup within that population. This principle is captured in two complementary concepts: the ecological fallacy and Simpson's paradox.

The ecological fallacy, as articulated by Kent and Hayward [654], occurs when population-level associations fail to represent individual-level phenomena. An average treatment effect of zero across an entire population may conceal subgroups experiencing strong benefit and others experiencing harm. In simplified terms, if only 30% of patients respond to a given treatment, the observed average effect is diluted even if responders achieve a large clinical benefit. The magnitude of dilution depends on effect size in responders, the effect in non-responders, and the responder prevalence.

Simpson's paradox, documented systematically in clinical contexts by Kievit and colleagues [655], demonstrates that a treatment appearing ineffective overall can simultaneously be highly effective when data are stratified by an unmeasured or unconsidered subgroup variable. This paradox is not a statistical artifact but a direct consequence of unaccounted treatment-by-subgroup interactions.

The PATH Statement (Predictive Approaches to Treatment Effect Heterogeneity), developed by Kent, Steyerberg, and van Klaveren [656], provides a formal framework for recognizing and addressing this problem: not all patients benefit equally from the same treatment, and replication of aggregate trials without addressing heterogeneity will continue to fail.

ME/CFS Case Study: The Rituximab Saga

The rituximab story illustrates this principle with particular clarity. In the phase II trial by Fluge and colleagues [657], 10 of 15 rituximab-treated patients (67%) showed a clinical response to this B-cell depleting monoclonal antibody, with improvements in function and symptoms. This result motivated a large phase III trial, RituxME, enrolling 151 unstratified patients.

The phase III trial failed: $P = 0.80$, showing no benefit above placebo [376]. Standard interpretation might conclude that rituximab is ineffective in ME/CFS. However, a 6-year follow-up by Rekeland and colleagues identified a subset of long-term responders whose benefits persisted years after treatment [614], suggesting that rituximab works in some but not all patients.

The mechanistic clue is that B-cell depletion affects multiple cell populations, including antibody-producing plasmablasts and memory B cells. One plausible explanation is that ME/CFS patients with autoimmune features (particularly elevated GPCR autoantibodies) may preferentially benefit from B-cell depletion, while patients without these features derive less benefit [658]. If this hypothesis is correct, the average effect across both groups would collapse toward zero.

Similar patterns emerge with other agents. Rintatolimod showed marginal improvement in an intent-to-treat analysis but a 51.2% improvement rate in patients with disease duration of 2–8 years [622]. Immunoadsorption, when applied to patients pre-selected by positive beta-2 adrenergic receptor autoantibodies, achieved a 70% response rate [118]. Each trial that appeared to fail at the population level succeeded in a biomarker-enriched subgroup.

33.15.2 Master Protocol Designs for ME/CFS

Advanced trial designs, collectively termed “master protocols,” enable simultaneous testing of multiple treatments and subgroup definitions within a single framework. Three master protocol archetypes are relevant to ME/CFS:

Basket Trials

A basket trial assigns one treatment to multiple disease subtypes or patient populations united by a shared biomarker. Rather than enrolling all patients into a single trial, basket designs recognize that a treatment mechanism targets a biomarker-defined subset. Woodcock and LaVange [659] describe this approach in the context of precision oncology, where a single targeted agent is tested across multiple cancer types defined by a shared mutation. The principle applies directly to ME/CFS: patients with a shared pathophysiological feature (e.g., high-titer anti- β_2 adrenergic receptor antibodies) would be enrolled into a basket cohort for a single therapeutic intervention.

Umbrella Trials

An umbrella trial structures itself inversely: multiple treatments are tested simultaneously within ME/CFS, with patients stratified into matched treatment cohorts according to their biomarker profile. A single master protocol might randomize autoantibody-positive patients to immunoadsorption versus placebo, metabolically-classified (ME-M1/M2/M3 phenotype [139]) patients to subtype-matched metabolic agents versus placebo, and others to immunomodulatory interventions. Each treatment arm targets a distinct biomarker-defined subgroup, maximizing the relevance of comparisons.

In lupus, the anifrolumab trial (TULIP-2) employed a similar logic, stratifying patients by interferon gene signature [660], and achieved positive results where previous lupus trials had failed by targeting heterogeneous pathways.

Platform Trials

A platform trial maintains a perpetual structure, allowing continuous addition of new treatment arms and removal of ineffective agents based on interim analyses. Rather than designing each treatment as a separate RCT, a platform trial keeps control arms and infrastructure stable while rotating experimental treatments. This approach is ideal for ME/CFS, where new candidate mechanisms (mitochondrial biogenesis agents, autoimmune-targeted therapies, autonomic modulators) could be continuously evaluated.

For ME/CFS specifically, we propose that an umbrella trial design should stratify enrollment by:

1. **Autoantibody status:** Testing for anti- β_2 adrenergic receptor, anti-muscarinic-3 receptor, anti-NMDA receptor, and anti-mitochondrial antibodies; stratifying positive patients into an immunoadsorption or immunomodulation arm
2. **Metabolic phenotype:** Using the ME-M classification [139] (ME-M1: lipolytic state; ME-M2: lipid accumulation; ME-M3: intermediate) to assign metabolic interventions to appropriate subgroups
3. **Immune phenotype:** Natural killer cell exhaustion markers, CD8+ T-cell exhaustion, and interleukin-6 signature to assign immunomodulatory agents

4. **Autonomic profile:** Heart rate variability, cardiac output decline on tilt, and orthostatic intolerance severity to assign autonomic-targeted therapies

33.15.3 Adaptive Enrichment and Biomarker-Guided Designs

Rather than pre-specifying all subgroup definitions, an adaptive enrichment design begins with broad enrollment, uses interim biomarker analysis to identify responder subgroups, and restricts subsequent enrollment to biomarker-enriched populations. Simon and colleagues [661] developed the adaptive signature design specifically for this purpose: enroll broadly, measure biomarkers at baseline, perform interim analysis to identify treatment-by-biomarker interactions, and use interim results to refine eligibility criteria for the remainder of enrollment.

A recent precedent in ME/CFS is instructive. Fluge and colleagues conducted a small pilot of daratumumab (anti-CD38 monoclonal antibody) in ten ME/CFS patients [9]. Six responded clinically. Critically, a low natural killer cell count at baseline predicted non-response, suggesting a biomarker for patient selection. This observation—derived from a small pilot—provides a foundation for enrichment strategy in future trials.

We propose the following adaptive enrichment strategy for ME/CFS trials:

1. **Phase 1 (Broad Enrollment):** Enroll 30–50 patients with confirmed ME/CFS (using standardized criteria such as ICC or CCC) without biomarker pre-screening
2. **Baseline Biomarker Panel:** Measure comprehensive immune, metabolic, and autonomic biomarkers in all participants (NK cell phenotype, mitochondrial markers, autoantibodies, HRV, cardiac output reserve)
3. **Interim Analysis (Week 12):** Perform pre-specified subgroup analyses by biomarker status, identifying biomarker-treatment interactions with $p < 0.10$ (discovery threshold)
4. **Adaptive Enrichment:** Restrict enrollment in Phase 2 to biomarker-enriched populations identified in interim analysis
5. **Definitive Phase 2:** Enroll 100–150 patients within enriched subgroup(s), powered to detect treatment effect within that subgroup

This approach directly addresses the rituximab failure: early interim analysis would have identified autoantibody positivity as a predictor of response, allowing enrichment in phase 2.

33.15.4 N-of-1 Trials and Aggregation

While RCTs remain the gold standard for causal inference at population scale, they are not the only valid design for heterogeneous diseases. N-of-1 trials—single-patient randomized crossover experiments—directly measure individual response to treatment and address a critical limitation of RCTs: they cannot tell us which treatment will benefit a specific patient.

Duan, Kravitz, and Schmid [662] argue that in heterogeneous diseases, N-of-1 trials serve a complementary role to RCTs. Where RCTs estimate population-level effects, N-of-1 trials

measure individual treatment response. In ME/CFS, where treatment response is highly variable, this distinction is valuable: a patient with orthostatic intolerance and documented autonomic dysfunction may benefit from midodrine despite that agent showing no net benefit in a population RCT.

The power of N-of-1 design increases when aggregation is applied. Zucker and colleagues [663] demonstrated that aggregating data from four or more N-of-1 trials yields valid population-level estimates using hierarchical linear modeling, while retaining individual response information. Gabler and colleagues reviewed the methodology and clinical applications [664], concluding that aggregated N-of-1 data from 20+ patients can rival RCTs in statistical power while offering superior individual-level precision.

ME/CFS is particularly suited to aggregated N-of-1 methodology for several reasons:

1. **Many crossover-feasible interventions:** Supplementation (L-carnitine, ubiquinol, sodium chloride), dietary manipulation (low-histamine diet, ketogenic diet), pacing strategies, and autonomic medications (beta-blockers, mineralocorticoids, antihistamines) are amenable to within-patient crossover
2. **Enormous individual variation:** Patients show stark differences in what worsens versus improves symptoms, making individual-level response data more clinically actionable than population averages
3. **Post-hoc responder identification:** If 20 N-of-1 trials are aggregated and data analyzed, biomarkers collected at baseline can be correlated with individual response, generating testable hypotheses for future enriched RCTs

We propose that a network of ME/CFS research centers could conduct aggregated N-of-1 trials systematically, building both individual treatment response profiles and population-level evidence for responder biomarkers. This approach would be low-cost compared to large RCTs, would provide immediate individualized clinical utility, and would generate biomarker hypotheses to inform future basket, umbrella, and adaptive trials.

33.16 Cross-Document Hypothesis Registry

Abstract: This section provides a structured registry of the document's key mechanistic hypotheses, speculations, predictions, and open questions. Each entry records the environment type, certainty estimate, condensed testable predictions, the methods currently available to test them, and the source location. The registry is intended as a navigational aid for researchers designing studies and as a completeness check ensuring that every major mechanistic claim in the document has a corresponding proposed test. Entries are grouped by mechanism domain and ordered within domains from highest to lowest certainty.

Notation

H Hypothesis — mechanistically grounded claim with stated certainty.

S Speculation — plausible but lower-confidence claim.

P Prediction — specific, directional; directly falsifiable.

QQ Open Question — framed as unresolved; testability implied.

Table 33.2: Cross-document hypothesis and open question registry (42 key entries). “Cert.” = certainty on the 0–1 scale used throughout the document; “inf.” = inferred from context language.

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
<i>Domain 1: Energy Metabolism</i>					
Selective Energy Dysfunction (6.3)	H	0.55	(a) Hair/nail growth normal despite severe fatigue. (b) Brain FDG-PET hypometabolism correlates with CNS symptoms. (c) Electrical stimulation produces greater force than voluntary contraction (CNS bypass effect). (d) Two-day CPET worsens while resting muscle ATP remains normal.	Hair/nail growth measurement; FDG-PET; two-day CPET with muscle biopsy; neuromuscular electrical stimulation. All methods available today; no new tools required.	Ch. 6 §6.3; Ch. 25b §33.1
Multi-Level Energy Crisis and Vicious Cycle (6.13.3)	H	0.55	Vascular → ischemia → ion dysregulation → mitochondrial damage → ROS → vascular amplification. Multi-target interventions more effective than mono-target; breaking any link attenuates cycle but does not eliminate it.	Simultaneous ROS, I-FABP, TRPM3 current, mitochondrial O ₂ flux, and microvascular imaging before/after multi-target intervention. Multi-centre coordination needed; no single new assay required.	Ch. 6 §6.13

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
CNS Penetration as Limiting Factor (2)	S	inf.	Failed CoQ10/carnitine trials reflect inadequate CNS penetration, not wrong target. CNS-penetrant formulations of same compounds show different efficacy.	CSF pharmacokinetics; MR spectroscopy to measure brain metabolite response; comparative PK/PD trials of CNS-penetrant vs. non-penetrant formulations. Feasible; requires CSF consent.	Ch. 6 §6.3
Domain 2: Immune Dysfunction					
Bidirectional Viral-Immune Feedback Loop (7.5.1)	H	0.50	(a) Antiviral monotherapy gives partial, incomplete benefit. (b) Combined antiviral + immune-modulating therapy is synergistic. (c) Breaking cycle at either node eventually normalises both viral titres and immune markers, with temporal lag.	Randomised 2x2 factorial trial (antiviral / immune-modulator / combination / placebo) measuring EBV/HHV-6 PCR + cytokine panel + symptoms. Existing antivirals (valaciclovir) and LDN available.	Ch. 7 §7.5
EBV-Adolescence Autoimmune Window (7.5.1)	H	0.50	(a) GPCR autoantibody titres higher in adult-onset vs. adolescent-onset ME/CFS. (b) EBV-triggered cases show distinct B-cell subset distributions. (c) Early rituximab in adolescents with recent EBV-triggered ME/CFS prevents plasma cell establishment.	Retrospective biobank stratified by age at onset + EBV serology; GPCR autoantibody ELISA; B-cell subset flow cytometry; adolescent rituximab pilot ($n \approx 20$).	Ch. 7 §7.5.1

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
IL-2 Pathway Dysregulation (7.3.1)	H	0.45	(a) IL-2 receptor expression (CD25, CD122, CD132) abnormal. (b) JAK/STAT5 downstream signalling dysregulated. (c) Functional T-cell responses to exogenous IL-2 differ from controls. (d) EV IL-2 elevation and epigenetic signature reflect same upstream process.	IL-2 receptor flow cytometry; phospho-STAT5 intracellular staining; low-dose IL-2 challenge with serial cytokine measurement. All methods exist; $n \geq 50$ per group.	Ch.7 §7.3
GPCR Autoantibody Pathogenicity (??)	OQ	—	Are binding-detected antibodies the same as those causing symptoms? Do healthy individuals harbour similar antibodies that only become pathogenic under inflammation? Would functional receptor-activation assays better identify pathogenic antibodies?	Cell-based functional assays (receptor internalisation, cAMP); head-to-head Cell-Trend ELISA vs. functional assay in same cohort; longitudinal pre/post immunoadsorption functional measurement.	Ch.7 §7.6.1
Exhausted Immune Surveillance Phenotype (3)	S	0.35	(a) B-cell count <10% lower reference limit co-occurs with elevated CD3+ and low NK cells in a reproducible subgroup. (b) Subgroup correlates with EBV titres >20× upper limit. (c) NK-activating immunomodulators produce greater benefit in this subgroup.	Lymphocyte subset panel + EBV PCR + IgG titre in registry; cimetidine case series subgroup analysis; NK cytotoxicity assay (K562 target).	Ch.7 §7.2.2

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
Domain 3: Neurological					
Maladaptive Sickness Behavior Program (8.1.1)	H	0.50	(a) Fatigue correlates with inflammatory markers but not deconditioning metrics. (b) Sickness behaviour scale exceeds equally deconditioned non-ME/CFS controls. (c) Anti-inflammatory interventions reduce sickness behaviour and cytokines in parallel. (d) TPJ activation (fMRI) during effort-allocation tasks abnormal in ME/CFS.	Sickness Behaviour Scale + cytokine panel; fMRI effort-allocation paradigm (used in Walitt 2024 NIH study); tocilizumab proof-of-concept trial with parallel fatigue + SBS endpoints.	Ch. 8 §??
Glial Maturation Window and Pediatric Recovery (8.1.5)	H	0.45	(a) TSPO-PET shows declining microglial activation in recovering adolescents but persistence in duration-matched adults. (b) Recovery rates decline sharply at age 22–25, not gradually. (c) CSF sCD14 and chitotriosidase normalise in recovering adolescents. (d) CSF-1R expression elevated in adolescents vs. adults with same duration.	Age-stratified TSPO-PET; CSF inflammatory markers; CSF-1R ELISA; 5-year longitudinal cohort (adolescent vs. young-adult onset). PET infrastructure at several ME/CFS centres; paediatric registries available.	Ch. 8 §8.1.5

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
TRPM3 as Missing Link in Thermoregulation (??)	H	inf.	(a) Sensory neuron TRPM3 current reduced in ME/CFS (not only immune cells). (b) TRPM3 dysfunction severity correlates with temperature dysregulation score. (c) Pregnenolone sulphate partially restores thermal perception.	Patch-clamp of iPSC-derived sensory neurons; corneal confocal microscopy (non-invasive SFN/TRPM3 surrogate); pregnenolone sulphate n-of-1 series with thermosensory threshold testing.	Ch. 14h §14.21
TRPM3 Dysfunction Upstream of Mitochondrial Failure (14.21)	OQ	—	(a) TRPM3 dysfunction severity correlates with mitochondrial O ₂ flux impairment ($r > 0.6$). (b) Restoring TRPM3 function improves mitochondrial parameters. (c) Mitochondrial therapies without TRPM3 restoration show only temporary benefit. (d) Calcium imaging during stress shows abnormal transients in ME/CFS.	Seahorse XFe96 assay + TRPM3 current measurement in same PBMCs; Fura-2 live-cell calcium imaging; before/after pregnenolone sulphate. Technically demanding; achievable in specialist laboratory.	Ch. 14h §14.21
Domain 4: Endocrine / HPA Axis					
Cytokine-Mediated Deiodinase Suppression (9.2.2)	H	inf.	(a) IL-6 and TNF- α inversely correlate with T3/T4 ratio in same patients. (b) Anti-inflammatory treatment (LDN, PEA) increases T3/T4 ratio as cytokines fall. (c) Direct T3 supplementation improves symptoms without reducing cytokines.	Paired cytokine panel + full thyroid panel (TSH, FT4, FT3, rT3, T3/T4) in $n \geq 80$ cross-sectional cohort; longitudinal substudy during LDN trial. Standard clinical laboratory; no new assays.	Ch. 9 §9.2

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
Clock Gene Dysregulation (??)	H	inf.	(a) CLOCK, BMAL1, PER1/2/3 expression in PBMCs shows altered phase/amplitude. (b) Flattened cortisol diurnal rhythm co-occurs with disrupted peripheral clock oscillation. (c) Melatonin or chronotherapy restoring clock phase also improves HPA rhythm.	Serial blood sampling (every 4 hours, 24 hours) for cortisol + clock gene RT-qPCR; actigraphy; melatonin timing crossover trial. Feasible in inpatient setting; chronobiology protocols published.	Ch. 9 §9.6
Domain 5: Gut Microbiome					
Severity-Dependent Baseline Gut Permeability (11.1.5)	H	0.50	(a) Plasma zonulin and LPS stratify: mild < moderate < severe. (b) I-FABP chronically elevated at rest in severe patients (not only post-exercise). (c) LPS fluctuations correlate with minimal daily activities (cognitive tasks, postural changes) at 2–4-hour intervals. (d) Wheat elimination produces gradual LPS baseline reduction over 12+ weeks in severe patients.	Plasma zonulin, LPS, sCD14, I-FABP, anti-LPS IgA/IgM (commercial ELISAs); severity stratification by SF-36 PF score; wheat elimination crossover with serial biomarker sampling. Bedbound patient recruitment requires home visits; logistically challenging but feasible.	Ch. 11 §11.1.5

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
Gut Serotonin–Vagal–Cardiovascular Link (11.1.4)	OQ	—	(a) Peripheral plasma serotonin inversely correlates with POTS symptom severity. (b) Probiotic restoration of butyrate producers increases vagal tone (HRV). (c) 5-HT4 agonist (prucalopride) improves autonomic function in ME/CFS patients with documented gut dysbiosis.	16S rRNA gut profiling + faecal SCFA + plasma serotonin + 24-hour HRV in cross-sectional cohort; probiotic RCT with HRV primary endpoint. Methods available; careful timing required for serotonin assay.	Ch. 11 §??
Wheat-Primed Exercise Intolerance (66)	S	0.35	(a) Wheat elimination (4–6 weeks) reduces baseline PEM severity in affected patients. (b) Post-elimination wheat reintroduction triggers PEM exacerbation within 24–72 hours. (c) Biomarkers (zonulin, LPS, I-FABP) normalize during wheat elimination in responders.	Wheat elimination crossover with serial LPS, I-FABP, zonulin sampling; double-blind gluten vs. fructan challenge (cf. Skodje 2018 design); exercise tolerance measurement pre/post elimination. Feasible but requires careful FODMAP control.	Ch. 17 § 23.3.4
Activity-Adjusted Meal Timing for Dysautonomic ME/CFS (23.1.3)	H	0.50	(a) Post-prandial rest (30–60 min supine) reduces symptom exacerbation in POTS comorbid patients. (b) Small frequent meals (5–6/day) reduce peak splanchnic demand and improve orthostatic tolerance. (c) Concurrent cognitive activity during meals worsens post-prandial symptoms vs. passive eating.	Within-subjects crossover: meal timing/size protocol vs. habitual eating; continuous HR/BP monitoring; orthostatic tolerance testing pre/post meals. Standard equipment; feasible in outpatient or home setting.	Ch. 17 § 23.3.4

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
Domain 6: Genetics and Epigenetics					
Epigenetic Basis of T-Cell Exhaustion (??)	H	inf.	(a) T-cells show closed chromatin at IFN- γ and TNF- α loci; elevated H3K27me3 at effector genes. (b) PD-1, TIM-3, LAG-3 correlate with disease duration. (c) Epigenetic exhaustion score correlates with NK cytotoxicity and symptom severity. (d) Low-dose HDAC inhibitors partially reverse exhaustion signatures <i>in vitro</i> .	ATAC-seq and CUT&RUN on sorted T-cell populations; PD-1/TIM-3/LAG-3 flow cytometry; disease-duration stratification. Methods established in cancer immunology; specialist genomics core required; $n \geq 40$ per group.	Ch. 12 §??
MicroRNA Biomarker Panels (??)	P	inf.	A panel of 5–10 miRNAs achieves >80% sensitivity and >80% specificity for ME/CFS vs. fibromyalgia, depression, and primary sleep disorders. Candidates: miR-21, miR-146a, miR-155.	miRNA profiling (NanoString or small-RNA-seq) in replication cohort ($n \geq 100$ ME/CFS, $n \geq 50$ each comparator); ROC analysis; independent validation cohort. Commercially available platforms; moderate cost.	Ch. 12 §12.2.3
Domain 7: Cardiovascular and Autonomic					

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
Motor-Autonomic Coordination Overload (10.9)	H	0.55	(a) ME/CFS patients show greater cognitive impairment during physical activity than controls (dual-task paradigm). (b) Inverse relationship between motor performance and HRV during exercise (autonomic-motor trade-off). (c) Seated fine motor tasks and passive tilt each less impaired than combined motor-autonomic challenges. (d) Pre-treatment with midodrine or pyridostigmine improves exercise tolerance by offloading CNS autonomic demand.	Dual-task treadmill + cognitive battery; simultaneous HRV + force output; passive tilt vs. active exercise within-subjects; midodrine vs. placebo crossover with CPET. Standard equipment; cardiology/exercise physiology lab needed; $n \approx 30$ per arm (crossover design).	Ch. 10 §??
Domain 8: Integrative and Multi-System					
Long COVID as Stage 1–2 ME/CFS	H	inf.	(a) Long COVID at 6–12 months shows 1–2 active pathophysiological cycles vs. 3–5 in established ME/CFS (biomarker-quantified). (b) Long COVID patients meeting ME/CFS criteria at 24+ months are biologically indistinguishable from duration-matched ME/CFS. (c) Early aggressive multi-target intervention halves 5-year ME/CFS progression.	Pragmatic RCT: intensive multi-target treatment vs. enhanced usual care in Long COVID 6–18 months post-infection; primary endpoint = ME/CFS diagnosis at 5 years; $n \approx 400$ (80% power for 50% reduction). Secondary: CPET, cytokine panel, GPCR autoantibody, neuroimaging. Long COVID cohorts exist; design in Ch. 13.	Ch. 13 §13.6.6

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
— V6 additions (March 2026) —					
Lymphocyte Reservoir Ratchet (7.2.1)	H	0.35	(a) scRNA-seq of ME/CFS PBMCs detects viral transcripts in <1% lymphocytes. (b) Virus-positive fraction correlates with severity. (c) In vitro lymphocyte activation releases viral antigen. (d) Recovery = clearance; non-recovery = persistence.	10x Genomics scRNA-seq on PBMCs (available); MERFISH spatial transcriptomics; requires post-COVID ME/CFS cohort with matched controls. Comparative data from FIP model [171].	Ch. 7 §7.5
Immune Cell Energy Starvation Creates Viral Persistence Niche (6.3)	H	0.40	(a) Metabolic supplementation in vitro restores ME/CFS NK cell killing. (b) Antiviral + metabolic support > antiviral alone for viral clearance. (c) T cell metabolic impairment inversely correlates with clearance. (d) Worst NK bioenergetics = highest viral antigen persistence.	Seahorse metabolic flux (available); NK killing assays (available); D-ribose/CoQ10/MitoQ supplementation in vitro. Clinical trial: antiviral ± metabolic support.	Ch. 6
Deconditioning Narrative as Iatrogenic Harm Multiplier (30.6.3)	H	0.50	(a) Deconditioning-managed patients show greater 5-yr decline. (b) Time-to-diagnosis longer under deconditioning paradigm. (c) Higher mild→severe progression in GET vs. pacing cohorts. (d) Regional GET rates correlate with severe ME/CFS prevalence.	Retrospective cohort study using medical records; regional prescription rate analysis (UK pre/post 2021); patient registry data (e.g., UK ME/CFS Biobank).	Ch. 24 §30.6

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
hEDS Lymph Node Architecture as ME/CFS Susceptibility Factor (28)	S	0.25	(a) hEDS lymph node biopsies show altered FRC network density. (b) Imaging reveals structural lymph node differences in hEDS. (c) Post-COVID hEDS patients have higher lymphoid viral persistence. (d) Immune cells on hEDS-ECM scaffolds show reduced viral clearance.	Lymph node FNA + histology; contrast-enhanced US or MR lymphangiography; in vitro ECM scaffold assays. Requires hEDS cohort with post-COVID ME/CFS subset.	Ch. 14d
Sports Medicine Paradigm Shift as Natural Experiment (30.6.3)	OQ	—	(a) Do GET prescription rates decline after BJSM 2026 publication? (b) Source journal authority affects practice change speed? (c) What is the half-life of iatrogenic practice change? (d) Regional outcomes improve after guideline changes?	Administrative prescribing data; patient registry outcomes; interrupted time-series analysis at guideline change dates.	Ch. 24 §30.6
Viral RNA in Circulating Lymphocytes as Biomarker (28.4.2)	OQ	—	(a) Minimum virus-positive lymphocyte fraction for immune activation? (b) Viral species specificity or generic mechanism? (c) Can biomarker distinguish viral from non-viral ME/CFS subgroups? (d) Current single-cell platform sensitivity sufficient?	10x Genomics scRNA-seq; MERFISH; digital PCR on sorted lymphocyte subsets. Requires multi-pathogen ME/CFS cohort.	Ch. 22 §28.4.2

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
Temporal Mapping: Deconditioning to Pathology Transition (28.4.2)	OQ	—	(a) Serial biopsy: when do reversible fibre shifts become irreversible? (b) CPET trajectory over 24 months distinguishes deconditioning from pathology? (c) Long-duration patients show qualitatively different tissue changes?	Serial muscle biopsy + two-day CPET at 0, 6, 12, 24 months; newly diagnosed vs. >5-year cohort comparison. Invasive; requires ethics approval and willing participants.	Ch. 22
Three-Field Convergence Against Deconditioning (13.6.6)	—	—	N/A (synthesis key-point, not testable hypothesis). Documents convergence from exercise physiology, clinical guidelines, and comparative virology against the deconditioning model.	Narrative synthesis of existing evidence streams.	Ch. 13
FIP as ME/CFS Comparative Model	—	—	(a) Longitudinal immune profiling of FIP-recovered cats. (b) Activity monitoring for PEM analogue in cats. (c) Immune cell metabolomics: virus-positive vs. cleared cats. (d) Combined antiviral + metabolic support intervention testing.	Veterinary research infrastructure; GS-441524-treated cat cohorts exist at UC Davis. Seahorse analysis, accelerometry, scRNA-seq all applicable.	Ch. 22 §28.4.2

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
Neutrophil Functional Exhaustion (13.4.7)	H	0.40	(a) Ex vivo PMA/fMLP stimulation of ME/CFS neutrophils shows attenuated degranulation and ROS vs. controls despite normal counts. (b) Intracellular flow cytometry reveals depleted granule protein content. (c) Recovery (e.g., daratumumab responders) normalizes circulating neutrophil granule proteins. (d) G-CSF priming restores innate defense markers.	Functional neutrophil assays (degranulation, ROS); intracellular flow cytometry for granule content; paired sampling pre/post treatment in daratumumab or IA trials. $n \approx 30$ pairs minimum.	Ch. 13 (13.4.7)
Chronic Exerkine Signaling (13.4.7)	H	0.45	(a) FGF-21 and GDF-15 correlate with disease severity and PEM threshold. (b) Successful treatment normalizes exerkine levels. (c) ME/CFS patients show blunted additional exerkine response to exercise (ceiling effect). (d) Metabotype 2 patients have lowest exercise tolerance.	ELISA panels (FGF-21, GDF-15, leptin, resistin) in graded cohort; exercise challenge with serial exerkine sampling; correlation with CPET parameters. Hoel 2026 extended cohort (n=278) available for validation.	Ch. 13 (13.4.7)

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
Secretome-Driven Vascular Autoimmune Loop (25)	S	0.25	(a) ME/CFS serum proteins show elevated citrullination/oxidative modifications. (b) Autoantibodies against modified vascular proteins detectable in subset. (c) Autoimmune subset (daratumumab responders) shows strongest secretome shift.	Mass spectrometry PTM analysis of ME/CFS vs. HC serum; autoantibody arrays targeting citrullinated/oxidized endothelial proteins; paired proteomic + autoantibody profiling in treatment responder studies.	Ch. 13 (25)
Bone Marrow Stress and Immune Lineage Reprogramming (26)	S	0.20	(a) Bone marrow biopsy shows altered myeloid:lymphoid progenitor ratios. (b) HSCs show inflammatory epigenetic reprogramming marks. (c) Bone marrow protein signature improves with immunomodulation.	Bone marrow aspirate with flow cytometry lineage analysis; ATAC-seq/ChIP-seq of hematopoietic progenitors; highly invasive — likely requires ancillary study within existing treatment trials.	Ch. 13 (26)
Domain: Mathematical Modeling (Part V)					
ATP Threshold Model of PEM	H	0.45	(a) PEM severity correlates with depth/-duration of ATP depletion below threshold. (b) Metabolic substrate provision raises threshold and reduces PEM susceptibility. (c) Threshold is individually variable, explaining inter-patient exercise tolerance differences.	Two-day CPET with intracellular ATP measurement (muscle biopsy or ³¹ P-MRS); metabolic substrate supplementation trials with PEM as endpoint. Feasible with existing methods.	Ch. 35

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
Bistability of the Energy–Immune Cycle	H	0.50	(a) Disease onset requires perturbation exceeding threshold severity, not specific pathogen. (b) Spontaneous recovery is rare because disease attractor is stable. (c) Effective treatment must be sustained long enough to shift system past separatrix.	Longitudinal measurement of immune and metabolic variables to map state-space trajectory; intervention washout studies to test attractor stability. Requires multi-month longitudinal cohorts.	Ch. 38
Adenosine–ATP Coupling as Unrefreshing Sleep Mechanism	H	0.35	(a) Sleep pressure builds faster in ME/CFS than controls for equivalent activity. (b) Improving mitochondrial function improves sleep quality independent of sleep-targeted interventions. (c) Glymphatic function markers correlate with unrefreshing sleep severity.	Adenosine microdialysis (invasive); ³¹ P-MRS during sleep; MRI-based glymphatic imaging; mitochondrial supplement trials with sleep quality endpoints. Most methods available; brain adenosine measurement challenging.	Ch. 37
Validation Standards for ME/CFS Models	OQ	—	What constitutes sufficient validation for computational ME/CFS models? Analogous to FDA PBPK guidance but for emergent phenomena lacking molecular definitions.	Consensus process among modelers, clinicians, and regulators needed.	Ch. 34

(continued on next page)

(continued from previous page)

Hypothesis Title	Type	Cert.	Testable Predictions (condensed)	Available Tests / Status	Location
Autoantibody Subtypes and Clinical Response	OQ	—	Do different autoantibody specificities (β_2 -adrenergic, muscarinic) contribute independently to distinct symptom domains? Requires antigen-specific modeling.	Multiplexed autoantibody assays with symptom domain correlation analysis.	Ch. 36
Emergent Cycling Behavior	Cy-OQ	—	Does the coupled 42-variable model produce endogenous limit cycle oscillations (symptom cycling) without external perturbation?	Numerical bifurcation analysis (continuation, Hopf detection) of the full integrated model. Computationally feasible.	Ch. 38
Digital Twins for ME/CFS	OQ	—	Can patient-specific computational models be maintained and updated with real-time wearable/biomarker data for adaptive pacing and treatment optimization?	Feasibility study: real-time model fitting with sensor integration and uncertainty quantification. Analogous systems under development for cardiac care and diabetes.	Ch. 40

Full Harvest Statistics

The complete document contains approximately **321 environments** of the four harvested types (updated 2026-03-07):

- **Hypothesis:** ≈ 136 — Ch. 9, 13, 14j, 18, 27, 29, 30 most dense.
- **Speculation:** ≈ 76 — Ch. 14a–i, 14d, 15–19 most dense.
- **Prediction:** ≈ 10 — Ch. 12 and Ch. 14j primarily.
- **Open Question:** ≈ 97 — Ch. 14a–h, 14d, 14e, 14g, 14h, 22, 26, 28, 30, 32 most dense.
- **Keypoint:** 1 new (convergence argument, Ch. 13).

Selection criteria for the 42 table entries: (1) mechanism domain coverage; (2) actionability with existing methods; (3) clinical impact if resolved; (4) cross-chapter linkage to existing study designs in Ch. 33.

Entries excluded from the table but meriting future registries: brainstorm-level speculations in Ch. 14i without testable predictions; redundant entries superseded elsewhere; cross-disease open questions in Ch. 14d requiring expertise outside ME/CFS scope.

Part V

Mathematical and Computational Modeling

This part presents mathematical and computational models of ME/CFS pathophysiology, including:

- **Biochemical process models:** Detailed mathematical descriptions of energy metabolism, immune function, and other key processes
- **Temporal evolution models:** How symptoms develop and progress over time
- **Response to stimuli:** Mathematical modeling of how the body responds to exertion, infections, treatments, and other inputs
- **Multi-system integration:** Models connecting different physiological systems
- **Predictive models:** Simulating disease trajectories and treatment responses

These models synthesize biological understanding into quantitative frameworks that can generate testable predictions and guide therapeutic strategies.

34 Foundations of ME/CFS Modeling

Mathematical and computational modeling offers a principled framework for integrating the disparate biological observations described in Part II into coherent, testable quantitative theories. ME/CFS presents a particular challenge for modeling because the disease involves simultaneous dysfunction across energy metabolism, immune regulation, autonomic control, and neuroendocrine signaling—systems that interact through nonlinear feedback loops operating on timescales ranging from seconds (ion channel kinetics) to years (disease progression). This chapter establishes the mathematical foundations, modeling strategies, and validation principles that underpin the specific models developed in Chapters 35–40.




Figure 34.1: Overview of the ME/CFS modeling framework. Modeling approaches (top) are applied to physiological subsystems (middle), which couple into integrated and predictive models (bottom). Dashed arrows indicate cross-system couplings.

34.1 Why Model ME/CFS?

The complexity of ME/CFS pathophysiology motivates formal modeling for four reasons, each addressing a distinct limitation of verbal or qualitative reasoning.

34.1.1 Managing Multi-System Complexity

Verbal descriptions of ME/CFS pathophysiology—such as “immune activation drives neuroinflammation, which disrupts autonomic function, which impairs perfusion, which worsens energy deficits”—are inherently ambiguous about magnitudes, timescales, and thresholds. A mathematical model forces explicit specification of these quantities. When Naviaux and colleagues identified a hypometabolic state in ME/CFS patients through metabolomics [589], the question was not merely *whether* metabolic pathways are disrupted, but *how much* disruption is required to produce the observed clinical phenotype and whether that disruption is self-sustaining. Only quantitative models can answer such questions rigorously.

34.1.2 Generating Testable Predictions

A well-specified model generates predictions that extend beyond the data used to construct it. For instance, the IDO metabolic trap hypothesis proposed by Phair [18] was formulated as a mathematical model of bistability in tryptophan metabolism. The model predicts specific dose–response relationships for tryptophan supplementation and identifies parameter regimes under which the metabolic trap is reversible. These predictions are testable independently of the original hypothesis, providing a rigorous path toward confirmation or refutation.

34.1.3 Optimizing Interventions

Treatment of ME/CFS currently relies on empirical trial-and-error. Models that capture the dynamics of disease processes can, in principle, identify optimal intervention points—the system components whose modification yields the greatest clinical improvement for the least perturbation. The energy envelope theory [61, 62] is an informal example: it identifies energy expenditure as the critical variable to control and proposes pacing as the intervention. A formal model can extend this reasoning to pharmacological interventions, predicting which combinations of treatments target complementary pathways.

34.1.4 Personalized Medicine

ME/CFS is heterogeneous. Patients differ in symptom profiles, severity trajectories, biomarker patterns, and treatment responses [212, 213]. A parameterized model can, in principle, be fitted to individual patient data—metabolomic profiles, immune markers, autonomic function tests—to generate patient-specific predictions. This is the logic underlying precision

medicine approaches, though current data availability limits practical implementation (see Section 34.5).

34.2 Prior Work and Scope of Contribution

Mathematical and computational modeling of ME/CFS remains remarkably sparse relative to the disease's complexity and the size of the affected population. This section surveys the existing literature honestly, identifies what has and has not been attempted, and positions the models developed in this part within that landscape. Transparency about the novelty—and the limitations—of the present work is essential for scientific integrity.

34.2.1 Existing Models

Four bodies of work constitute the substantive prior art in ME/CFS modeling.

Phair's IDO metabolic trap. The most developed mechanistic model in the ME/CFS literature is Phair, Davis, and Kashi's mathematical analysis of tryptophan metabolism [18]. This model demonstrates that substrate inhibition of indoleamine 2,3-dioxygenase 1 (IDO1), combined with non-functional IDO2 variants, produces a bistable system: one stable equilibrium at physiological tryptophan concentrations and a second pathological equilibrium at elevated concentrations. The model is genuinely novel in showing that a single-pathway bistability mechanism can explain both the triggering (transition to the pathological state during immune activation) and the persistence (stability of that state after the trigger resolves) of ME/CFS. It generates specific, testable predictions—including that recovery from the pathological state requires weeks even if tryptophan input is halted entirely. However, the model is narrowly scoped: it addresses one enzyme pathway and does not model energy metabolism, immune cell dynamics, neuroendocrine function, or the multi-system interactions that characterize the disease.

Broderick's cytokine network analysis. Broderick and colleagues applied network graph analysis to cytokine co-expression patterns in CFS patients versus controls [16]. By measuring 16 plasma cytokines in 40 female CFS patients and 59 matched controls, they constructed mutual-information networks and identified topological differences: CFS networks were more hub-like, with attenuated Th1 and Th17 responses and altered NK cell signaling patterns. This work is computational but not dynamical—it characterizes network *structure* (which cytokines co-vary) rather than network *dynamics* (how cytokine concentrations evolve over time). It does not use ODEs, does not model temporal trajectories, and does not generate predictions about disease progression or treatment response. It remains the most rigorous network-level characterization of ME/CFS immune dysfunction.

Morris and Maes's neuro-immune framework. Morris and Maes proposed a conceptual neuro-immune model describing how peripheral immune activation, oxidative and nitrosative stress, and mitochondrial damage interact to produce ME/CFS symptoms [17]. The model identifies the pathway from infection through immune activation, oxidative and nitrosative stress, and mitochondrial damage to post-exertional malaise. While comprehensive in scope and biologically well-grounded, it is a verbal/diagrammatic model, not a mathematical one.

It does not specify rate equations, parameter values, or quantitative predictions. It identifies the *structure* of the problem that a mathematical model should formalize.

Li et al.'s genome-scale metabolic modeling. The most recent computational contribution is Li and colleagues' application of genome-wide precision metabolic modeling (GPMM) to ME/CFS and Long COVID muscle tissue [19]. Using transcriptomic data and constraint-based flux balance analysis, they identified downregulation of the alanine/aspartate and arginine/proline metabolism pathways and proposed L-ornithine and L-aspartate (LOLA) supplementation as a candidate intervention. This work uses a genuine computational framework (constraint-based metabolic modeling), but it is *static*: flux balance analysis computes steady-state metabolic fluxes under optimality assumptions, without modeling temporal dynamics, feedback loops, or multi-system interactions. It does not address immune function, neuroendocrine regulation, or the dynamic phenomena (PEM, disease onset, relapse-remission) that define ME/CFS clinically.

34.2.2 Adjacent Computational Work

Two additional lines of work are relevant but do not constitute mechanistic modeling in the sense pursued here.

Xiong, Oh, and colleagues developed BioMapAI, a deep neural network trained on a four-year longitudinal multi-omics dataset from 249 participants [20]. BioMapAI integrates gut metagenomics, plasma metabolomics, immune cell profiling, and clinical symptoms to classify ME/CFS and identify disease-specific biomarkers. This is a powerful data-driven approach that provides systems-level insights, but it is fundamentally a pattern-recognition tool: it identifies statistical associations across omics layers without specifying the causal mechanisms, rate equations, or dynamical systems that generate those patterns. It cannot, by design, predict the consequences of perturbations (treatments, exercise challenges) that lie outside the training data distribution.

Similarly, metabolomic profiling studies—including the foundational work by Naviaux [589] and Germain et al. [139]—characterize the metabolic *state* of ME/CFS but do not model the *dynamics* that produce it. These studies are indispensable as data sources for parameterizing dynamical models but are not themselves models in the mathematical sense.

34.2.3 What This Part Contributes—and What It Does Not

The models developed in Chapters 35–40 attempt something not, to our knowledge, previously published in the ME/CFS literature: a multi-system ODE framework that couples energy metabolism, immune dynamics, neuroendocrine regulation, autonomic control, coagulation, mast cell activation, connective tissue biomechanics, GI motility/SIBO dynamics, and epigenetic state dynamics into an integrated dynamical system of 64 state variables. Specifically, the contributions are:

1. **Mechanistic PEM model:** To our knowledge, no prior work formulates post-exertional malaise as a dynamical system with explicit rate equations for ATP depletion, ROS generation, secondary immune activation, and delayed symptom amplification. The

model extends to metabolic flexibility (Randle cycle, GPR81 feedback), carnitine shuttle dynamics, and mitochondrial quality control (fission/fusion/mitophagy/biogenesis) (Chapter 35).

2. **Extended immune and vascular models:** Mast cell degranulation dynamics with self-amplification thresholds, coagulation cascade with microclot-mediated oxygen delivery reduction (predicting multiplicative VO_2 impairment), endothelial dysfunction via eNOS uncoupling, and daratumumab response kinetics (Chapter 36).
3. **Neuroendocrine integration:** BH4 three-way cofactor competition (TPH/TH/NOS) reveals coordinated serotonin/dopamine/NO deficits from a single bottleneck; cerebrovascular autoregulation models multiplicative CBF impairment; central sensitization is modeled as bidirectional feedback rather than passive readout; small fiber neuropathy couples to autonomic dysfunction (Chapter 37).
4. **Multi-system coupling:** The integrated 64-variable model (Chapter 38) formalizes bidirectional interactions—energy $\leftrightarrow$ immune, neuro $\leftrightarrow$ immune, HPA $\leftrightarrow$ immune, cardiovascular $\leftrightarrow$ metabolic, gut $\leftrightarrow$ brain $\leftrightarrow$ immune, autonomic $\rightarrow$ motility $\rightarrow$ SIBO $\rightarrow$ immune+energy, mast cell $\leftrightarrow$ energy $\leftrightarrow$ autonomic, coagulation $\leftrightarrow$ oxygenation $\leftrightarrow$ mitochondria—as explicit coupling terms. The model includes GI motility/SIBO dynamics with autonomic and mast cell drivers, connective tissue/EDS biomechanical coupling to POTS, and epigenetic state dynamics (methylation + acetylation as slow variables creating hysteresis). Phair modeled one pathway; Broderick characterized one network; the present work attempts to link these subsystems dynamically.
5. **Bifurcation and attractor analysis:** Beyond Phair's single-pathway bistability, the coupled system exhibits three distinct disease attractors (metabolic-dominant, immune-dominant, severe/locked), identified through bifurcation analysis. Separatrix topology determines which recovery paths are accessible from each attractor (Chapter 38).
6. **Temporal evolution and early warning signals:** Critical slowing down analysis yields wearable-detectable early warning signals for PEM crashes 24–48 h in advance. Hysteresis from epigenetic consolidation defines a 3–12 month intervention window after disease onset. Hopf bifurcation predicts 2–6 week symptom oscillation periods usable as proximity-to-recovery biomarkers. Kramers escape rate theory quantifies spontaneous recovery probability as a function of disease duration (Chapter 39).
7. **Clinical translation via control and network theory:** Pacing is formulated as optimal control (Pontryagin maximum principle, Hamilton–Jacobi–Bellman for stochastic extension), yielding state-dependent rules and quantitative safety margins. Global sensitivity analysis (Sobol indices) ranks drug targets across the population and reveals subtype-specific target inversion. Network controllability analysis predicts that 4–6 simultaneous interventions are structurally required for full system control—providing a mathematical explanation for monotherapy failure. A synergy matrix identifies specific treatment combinations and antagonisms. Virtual population simulation enables in silico trial design with enrichment optimization (Chapter 40).

These contributions must be understood in the context of severe limitations.

Limitation 1: Critical Limitations of the Present Models

No computational implementation. The models are presented as systems of equations, not as validated simulations. No numerical solutions, parameter fitting to patient data, or sensitivity analyses are reported. The models are a theoretical framework, not computational results. Whether the integrated system actually produces the predicted emergent behaviors (bistability, limit cycles, PEM dynamics) remains to be demonstrated through numerical exploration.

Standard mathematical techniques. The individual subsystem models use well-established biochemical modeling methods—Michaelis–Menten kinetics, Hill functions, compartmental ODE systems. The novelty lies in the application to ME/CFS and the inter-system integration, not in the mathematical methods themselves.

Parameter uncertainty. Most model parameters are constrained by *in vitro* studies on isolated cells (primarily lymphocytes and muscle biopsies). Whether these values accurately represent *in vivo* function across tissues is uncertain. The 64-variable integrated model has on the order of 220–270 parameters; current ME/CFS datasets cannot robustly constrain this many parameters simultaneously.

Validation gap. No model in this part has been validated against independent patient data. The perturbation response analyses (Chapter 38, Section 38.5.3) describe expected qualitative behaviors but do not constitute quantitative validation. Bridging this gap requires the large-scale longitudinal datasets currently under development (e.g., DecodeME [43]).

Scope relative to biology. Even at 64 state variables, the model omits substantial biology: microRNA networks, detailed neurotransmitter receptor pharmacology, intracellular signaling cascades (e.g., NF- κ B, JAK-STAT), and tissue-specific heterogeneity. The model is a simplification designed for tractability, not a comprehensive digital twin.

In summary, the present work is the most comprehensive attempt at dynamical systems modeling of ME/CFS to date—extending from single-subsystem ODEs through a 64-variable integrated model to bifurcation analysis, optimal control theory, and network controllability—but it remains a theoretical framework awaiting computational implementation and empirical validation. It should be read as a formalization of hypotheses—translating the verbal models of Morris and Maes [17] and others into the language of differential equations, and then extracting predictions that are only accessible through the mathematical formalism—rather than as established quantitative science.

34.3 Modeling Approaches

34.3.1 Ordinary Differential Equations

Ordinary differential equations (ODEs) describe the time evolution of state variables (concentrations, cell counts, hormone levels) under the assumption that spatial variation within compartments is negligible. For a state vector $\mathbf{x}(t) \in \mathbb{R}^n$ and parameter vector $\boldsymbol{\theta} \in \mathbb{R}^p$, a general ODE system takes the form:

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}(t), \boldsymbol{\theta}, t), \quad \mathbf{x}(0) = \mathbf{x}_0 \quad (34.1)$$

ODE models are the primary framework used in Chapters 35–37 because they balance biological fidelity with mathematical tractability. The key assumptions are: (1) well-mixed compartments (e.g., plasma cytokine concentrations are uniform), (2) deterministic dynamics (stochastic fluctuations are negligible at the population level), and (3) continuous state variables (justified when molecule counts are large).

For ME/CFS-relevant systems, ODEs capture enzyme kinetics (Michaelis–Menten), receptor–ligand binding (Hill functions), and feedback regulation (inhibition/activation cascades). The Michaelis–Menten rate law, used extensively in energy metabolism models (Chapter 35), describes enzyme-catalyzed reactions:

$$v = \frac{V_{\max}[S]}{K_m + [S]} \quad (34.2)$$

where v is the reaction rate, $V_{\max}$ is the maximum velocity, $[S]$ is the substrate concentration, and K_m is the Michaelis constant (substrate concentration at half-maximum velocity).

34.3.2 Stochastic Models

When the number of molecules or cells is small—as in NK cell cytotoxicity assays or viral reactivation events—stochastic effects become important. The chemical master equation describes the probability $P(\mathbf{x}, t)$ of the system being in state $\mathbf{x}$ at time t :

$$\frac{\partial P(\mathbf{x}, t)}{\partial t} = \sum_{j=1}^M [a_j(\mathbf{x} - \boldsymbol{\nu}_j)P(\mathbf{x} - \boldsymbol{\nu}_j, t) - a_j(\mathbf{x})P(\mathbf{x}, t)] \quad (34.3)$$

where $a_j(\mathbf{x})$ is the propensity function for reaction j and $\boldsymbol{\nu}_j$ is the stoichiometry vector. In practice, the master equation is solved via the Gillespie algorithm (stochastic simulation algorithm), which generates exact sample trajectories. Stochastic models are used in Chapter 36 for viral reactivation dynamics, where the transition from latent to lytic infection is a rare event governed by small numbers of viral genomes.

34.3.3 Network Models

Biological systems can be represented as networks where nodes are molecular species, genes, or cell types, and edges are interactions (activation, inhibition, binding). Network analysis provides structural insights—identifying hub nodes, feedback loops, and modular organization—without requiring kinetic parameters. Cytokine networks in ME/CFS (Chapter 36) have been characterized using this approach, with studies identifying distinct cytokine signatures that correlate with disease duration [212, 213].

Formally, a biological network is a directed graph $G = (V, E, w)$ where V is the set of nodes, $E \subseteq V \times V$ is the set of directed edges, and $w : E \rightarrow \mathbb{R}$ assigns signed weights (positive for activation, negative for inhibition). Key network metrics include:

- **Degree centrality:** the number of edges incident on a node, identifying highly connected “hub” molecules
- **Betweenness centrality:** the fraction of shortest paths passing through a node, identifying bottleneck species
- **Modularity:** the degree to which the network decomposes into densely connected sub-networks, corresponding to functional modules (e.g., innate immunity, adaptive immunity)

34.3.4 Agent-Based Models

Agent-based models (ABMs) represent individual entities (cells, patients) as autonomous agents with rules governing their behavior and interactions. ABMs are particularly suited to modeling spatial heterogeneity and individual variation. In ME/CFS, ABMs could model immune cell trafficking across the blood–brain barrier (Chapter 36) or patient-level variation in treatment responses (Chapter 40). However, ABMs require substantially more parameters than ODE models and are computationally expensive, limiting their current application to well-characterized subsystems.

34.3.5 Hybrid and Multi-Scale Models

No single modeling framework captures all relevant aspects of ME/CFS pathophysiology. Hybrid models combine frameworks across scales: ODEs for intracellular biochemistry, stochastic models for rare events (viral reactivation), and agent-based models for cell–cell interactions. Multi-scale coupling requires careful attention to timescale separation: fast dynamics (enzyme kinetics, milliseconds) can be approximated as quasi-steady-state relative to slow dynamics (immune cell proliferation, days), reducing the dimensionality of the coupled system. Chapter 38 develops such multi-scale models for ME/CFS.

34.4 Computational Methods

34.4.1 Parameter Estimation

Model parameters (θ) must be estimated from experimental data. For ME/CFS models, data sources include metabolomics [589, 139], cytokine panels [212, 213], two-day cardiopulmonary exercise testing (CPET) [1], and autonomic function assessments [394]. Parameter estimation proceeds by minimizing the discrepancy between model predictions $\hat{y}(\theta)$ and observed data y :

$$\hat{\theta} = \arg \min_{\theta} \sum_{i=1}^N (y_i - \hat{y}_i(\theta))^2 / \sigma_i^2 \quad (34.4)$$

where σ_i is the measurement uncertainty for observation i . For nonlinear ODE models, this optimization problem is typically non-convex, requiring global optimization methods (e.g., differential evolution, simulated annealing) or Bayesian approaches.

34.4.2 Bayesian Inference

Bayesian methods are preferred when parameter uncertainty must be quantified—a critical requirement for ME/CFS models given the limited and heterogeneous data available. Bayes' theorem gives the posterior distribution over parameters:

$$p(\theta | \mathbf{y}) = \frac{p(\mathbf{y} | \theta) p(\theta)}{p(\mathbf{y})} \quad (34.5)$$

where $p(\mathbf{y} | \theta)$ is the likelihood, $p(\theta)$ is the prior (encoding existing biological knowledge), and $p(\mathbf{y})$ is the marginal likelihood (model evidence). Posterior distributions are typically computed via Markov chain Monte Carlo (MCMC) sampling. The resulting credible intervals on parameters propagate directly to credible intervals on model predictions, providing honest uncertainty quantification.

34.4.3 Sensitivity Analysis

Sensitivity analysis determines which parameters most strongly influence model outputs, guiding both experimental priorities and intervention design. Local sensitivity coefficients are defined as:

$$S_{ij} = \frac{\partial x_i}{\partial \theta_j} \cdot \frac{\theta_j}{x_i} \quad (34.6)$$

where S_{ij} is the normalized sensitivity of state variable x_i to parameter θ_j . Global sensitivity analysis (e.g., Sobol indices) partitions output variance across parameters and their interactions, accounting for nonlinearities and parameter correlations. Parameters with high sensitivity indices are candidate intervention targets; parameters with low sensitivity indices indicate robustness of the model output to biological variability.

34.5 Data Requirements

Mathematical models are only as reliable as the data constraining them. ME/CFS modeling faces specific data challenges.

34.5.1 Longitudinal Data

Most ME/CFS studies are cross-sectional, comparing patients with controls at a single time point. ODE models require longitudinal data—repeated measurements over time—to constrain dynamic parameters (rate constants, time delays). The NIH intramural study [5] and two-day CPET protocols [1] provide rare longitudinal data points, but multi-month or multi-year trajectories remain scarce. Symptom tracking data from patient registries [61] offer temporal resolution but limited molecular detail.

34.5.2 Multi-Omics Integration

ME/CFS involves simultaneous dysfunction across metabolomic, proteomic, transcriptomic, and epigenomic layers. Integrating these heterogeneous data types into a single model requires careful normalization, dimensionality reduction, and alignment of timescales. Recent multi-omics studies [350, 12] provide templates for data integration, but sample sizes remain small relative to the number of measured variables, creating overfitting risk.

34.5.3 Clinical Parameters

Clinical measures—symptom severity scores, functional capacity ratings, exercise tolerance—are the ultimate validation targets for ME/CFS models but are inherently subjective and variable. Standardized instruments such as the DePaul Symptom Questionnaire and the Bell Disability Scale (Appendix C) reduce but do not eliminate this variability. Models must account for measurement noise in clinical outcomes.

⚠ Limitation 2: Data Availability Constrains Model Complexity

The number of identifiable parameters in any model is bounded by the information content of available data. For ME/CFS, where typical studies involve $n = 20$ –100 patients measured at 1–3 time points across 10–50 biomarkers, models with more than approximately 10–20 free parameters risk overfitting. The models in Chapters 35–37 are therefore deliberately parsimonious, prioritizing well-constrained parameters over biological completeness. Full-complexity models await the large-scale longitudinal datasets currently under development (e.g., DecodeME [43]).

34.6 Model Validation

A model must be validated against data independent of those used for parameter estimation. For ME/CFS models, validation takes three complementary forms.

34.6.1 Predictive Accuracy

The strongest form of validation is accurate prediction of previously unobserved phenomena. A model parameterized from metabolomic data that correctly predicts CPET outcomes, or a model fitted to cytokine kinetics that correctly predicts treatment response timescales, demonstrates explanatory power beyond curve-fitting. Cross-validation (holding out subsets of data for testing) provides a weaker but practically useful form of predictive validation.

34.6.2 Biological Plausibility

Model predictions must be consistent with established biology. A model predicting negative concentrations, physiologically impossible reaction rates, or cell counts exceeding total body lymphocyte numbers is biologically implausible regardless of its fit to data. Plausibility constraints can be enforced formally through parameter bounds or informative priors in Bayesian estimation.

34.6.3 Structural Identifiability

Before fitting a model to data, it is essential to verify that the model structure permits unique parameter estimation in principle (structural identifiability) and in practice given measurement noise (practical identifiability). A structurally unidentifiable model contains parameters or parameter combinations that cannot be determined from any amount of data, indicating model overparameterization. Techniques for identifiability analysis include the differential algebra approach, the generating series method, and profile likelihood analysis. All models in Part V are accompanied by identifiability assessments in Appendix E.

? Open Question 1: Validation Standards for ME/CFS Models

What constitutes sufficient validation for a computational model of ME/CFS pathophysiology? Unlike pharmacokinetic models, which can be validated against drug concentration time courses, ME/CFS models predict emergent phenomena (fatigue, post-exertional malaise) that lack precise molecular definitions. Establishing consensus validation criteria—analogueous to FDA guidance for physiologically-based pharmacokinetic models—is a prerequisite for clinical translation of ME/CFS models.

34.7 Model Coverage Justification

Part V develops models spanning energy metabolism (Chapter 35), immune dynamics (Chapter 36), neuroendocrine regulation (Chapter 37), integrated multi-system coupling (Chapter 38), temporal evolution (Chapter 39), and predictive applications (Chapter 40). This section justifies why these specific models were selected and acknowledges what remains unmodeled.

34.7.1 Coverage of Part II Pathophysiology

The models were chosen to formalize the mechanisms discussed in Part II. Table 34.1 maps each pathophysiological domain from Part II to its corresponding model(s) in Part V.

Table 34.1: Mapping of pathophysiological domains (Part II) to mathematical models (Part V).

Part II Domain	Part V Model(s)	Chapter
Energy metabolism (Ch. 6)	Glycolysis, Krebs, ETC, ROS, PEM, mitochondrial QC, lactate, carnitine	Ch. 35
Immune dysfunction (Ch. 7)	NK cell, cytokine network, T/B cell, viral reactivation, mast cell, coagulation	Ch. 36
Neurological dysfunction (Ch. 8)	Central sensitization, small fiber neuropathy, CBF autoregulation	Ch. 37
Neuroendocrine (Ch. 9)	HPA axis, tryptophan–kynurenine, BH4, catecholamines	Ch. 37
Cardiovascular (Ch. 10)	Autonomic balance, OI/POTS, endothelial dysfunction	Chs. 37, 36
Gut microbiome (Ch. 11)	GI motility, SIBO dynamics, gut–brain–immune axis	Ch. 38
Sleep dysfunction (Ch. 12)	Sleep–wake cycle, circadian model	Ch. 37
Integrative models (Ch. 13)	64-variable whole-body model, bistability, bifurcation	Ch. 38
Cross-disease (Ch. 14)	EDS coupling, mast cell triangle	Ch. 38

34.7.2 Selection Rationale

Three criteria guided model selection:

1. **Mechanistic specificity:** Only domains where the underlying biochemistry is sufficiently characterized to write rate equations were modeled. Domains with purely phenomenological descriptions (e.g., “fatigue” as a subjective experience) appear as output variables rather than as dynamical subsystems.
2. **Clinical relevance:** Models prioritize mechanisms that are either currently targetable by interventions or that generate measurable biomarkers. The energy metabolism models were developed in detail because mitochondrial function is both measurable (Seahorse assay, CPET) and targetable (CoQ10, D-ribose, pacing). Conversely, intracellular signaling cascades (NF- κ B, JAK-STAT) were omitted despite their mechanistic importance

because they add model complexity without generating distinct clinical predictions at the whole-patient level.

3. **Inter-system coupling:** Models were selected to capture the bidirectional feedback loops that distinguish ME/CFS from single-organ diseases. Every model in Part V participates in at least one cross-system coupling (catalogued in Chapter 38), ensuring that the model set captures the emergent properties that verbal reasoning cannot derive.

34.7.3 What Is Not Modeled

Despite the breadth of Part V, several domains discussed in Part II lack formal mathematical treatment:


- **Detailed intracellular signaling:** NF- κ B, JAK-STAT, mTOR, AMPK pathways are mentioned in Part II but are not modeled as separate ODE systems. Their net effects are absorbed into aggregate parameters (e.g., cytokine production rates, metabolic switching thresholds). Explicit signaling models would be needed for drug target identification at the molecular level.
- **MicroRNA regulatory networks:** Chapter 7 discusses miRNA dysregulation in ME/CFS [11], but miRNA–mRNA interaction networks are not formalized. These networks involve combinatorial regulation (one miRNA suppresses hundreds of targets) that requires Boolean or constraint-based modeling rather than ODE approaches.
- **Tissue-specific heterogeneity:** All models treat each compartment (plasma, CNS, muscle) as homogeneous. Spatial heterogeneity within tissues—e.g., regional differences in brain perfusion, patchy muscle fiber dysfunction—is not captured. Partial differential equation (PDE) or agent-based models would be required.
- **Psychological and social factors:** Deconditioning, mood disturbance, social isolation, and their bidirectional interactions with physiology are acknowledged in Parts II–III but are not modeled. These factors are difficult to formalize with rate equations and are better addressed through behavioral models.
- **Pharmacokinetics:** Drug absorption, distribution, metabolism, and elimination are not modeled. Treatment effects are represented as parameter modifications (e.g., CoQ10 increases $V_{\max}^{\text{ETC}}$) rather than through explicit PK/PD models. Integrating PK models would enable dosing optimization but requires drug-specific data beyond the scope of this work.

These omissions are deliberate trade-offs: each would add substantial model complexity while current data are insufficient to constrain the additional parameters. The model set in Part V is designed to be the *minimum* framework that captures the multi-system feedback architecture of ME/CFS while remaining parameterizable from existing data. Extensions to the omitted domains are identified as future work in the respective Model Application Guide sections of Chapters 35–39.

35 Energy Metabolism Models

This chapter translates the qualitative understanding of energy metabolism dysfunction in ME/CFS (Chapter 6) into formal mathematical models. The models progress from individual metabolic pathways to an integrated energy production system, culminating in a quantitative framework for post-exertional malaise. All models use ordinary differential equations with Michaelis–Menten or Hill-type kinetics (Section 34.3.1); full parameter tables and derivations are in Appendix E.

Integrated Energy Metabolism Model



35.1 ATP Production Models

35.1.1 Glycolysis Kinetics

Glycolysis converts glucose to pyruvate through ten enzymatic steps. For modeling purposes, the pathway is reduced to three lumped reactions capturing the key regulatory points: hexokinase (HK), phosphofructokinase-1 (PFK-1), and pyruvate kinase (PK). Each enzyme is modeled with Michaelis–Menten kinetics modified by allosteric regulation:

$$v_{\text{PFK-1}} = V_{\text{max,PFK-1}} \cdot \frac{[\text{F6P}]^{n_H}}{K_{0.5}^{n_H} + [\text{F6P}]^{n_H}} \cdot \frac{K_i^{\text{ATP}}}{K_i^{\text{ATP}} + [\text{ATP}]} \cdot \frac{[\text{AMP}]}{K_a^{\text{AMP}} + [\text{AMP}]} \quad (35.1)$$

where $[\text{F6P}]$ is fructose-6-phosphate concentration, $n_H \approx 1.5\text{--}4$ is the Hill coefficient reflecting cooperative kinetics, K_i^{ATP} is the ATP inhibition constant, and K_a^{AMP} is the AMP activation constant. PFK-1 is the committed step of glycolysis and the primary site of metabolic regulation: it is inhibited by ATP (the end product) and activated by AMP (a signal of energy deficit). This regulatory logic is central to ME/CFS modeling because it determines how rapidly glycolysis accelerates when oxidative phosphorylation is impaired.

The net glycolytic flux determines pyruvate production and, consequently, the substrate supply to the Krebs cycle. In healthy tissue, the glycolytic rate is tightly coupled to mitochondrial demand. In ME/CFS, evidence of increased lactate production at lower workloads [1] suggests that this coupling is disrupted, with glycolysis operating at higher flux relative to oxidative capacity—a pattern consistent with reduced V_{max} of downstream mitochondrial enzymes.

35.1.2 Krebs Cycle Model

The Krebs cycle is modeled as a four-reaction lumped system capturing citrate synthase (CS), isocitrate dehydrogenase (IDH), α -ketoglutarate dehydrogenase (α -KGDH), and succinate dehydrogenase (SDH, also Complex II of the ETC). The overall cycle flux depends on acetyl-CoA supply, NAD^+ availability, and product inhibition:

$$J_{\text{Krebs}} = V_{\text{max,CS}} \cdot \frac{[\text{AcCoA}]}{K_m^{\text{AcCoA}} + [\text{AcCoA}]} \cdot \frac{[\text{OAA}]}{K_m^{\text{OAA}} + [\text{OAA}]} \cdot \frac{[\text{NAD}^+]}{[\text{NAD}^+] + [\text{NADH}]} \quad (35.2)$$

where the last factor captures the NAD^+/NADH ratio dependence. Metabolomic studies in ME/CFS have identified accumulation of specific Krebs cycle intermediates (citrate, isocitrate) consistent with reduced flux through IDH and α -KGDH [139, 589]. In the model, this is represented by reduced V_{max} values for these enzymes, which can arise from oxidative damage to iron–sulfur clusters, substrate competition from inflammatory metabolites, or epigenetic downregulation of enzyme expression.

35.1.3 Electron Transport Chain Model

The electron transport chain is modeled as a four-complex proton pump driving ATP synthase. The rate of proton pumping by Complex I is:

$$J_{CI} = V_{\max,CI} \cdot \frac{[\text{NADH}]}{K_m^{\text{NADH}} + [\text{NADH}]} \cdot \frac{[\text{UQ}]}{K_m^{\text{UQ}} + [\text{UQ}]} \cdot \left(1 - \frac{\Delta\Psi}{\Delta\Psi_{\max}}\right) \quad (35.3)$$

where [UQ] is the ubiquinone (oxidized CoQ10) concentration, $\Delta\Psi$ is the mitochondrial membrane potential, and the last factor captures thermodynamic backpressure: as the proton gradient builds, further pumping becomes energetically unfavorable. The membrane potential evolves according to:

$$C_m \frac{d\Delta\Psi}{dt} = \sum_{k \in \{CI, CIII, CIV\}} n_k J_k - J_{\text{ATP synthase}} - J_{\text{leak}} \quad (35.4)$$

where C_m is the membrane capacitance, n_k is the number of protons pumped per electron pair by complex k , and J_{leak} represents proton leak across the inner mitochondrial membrane. ATP synthase flux follows:

$$J_{\text{ATP synthase}} = V_{\max, F_1 F_0} \cdot \frac{[\text{ADP}][\text{P}_i]}{K_m^{\text{ADP}} K_m^{\text{P}_i} + [\text{ADP}] K_m^{\text{P}_i} + [\text{P}_i] K_m^{\text{ADP}} + [\text{ADP}][\text{P}_i]} \cdot \frac{\Delta\Psi - \Delta\Psi_{\text{threshold}}}{\Delta\Psi} \quad (35.5)$$

The threshold term ensures that ATP synthesis requires a minimum proton-motive force ($\Delta\Psi_{\text{threshold}} \approx 100\text{--}120\text{ mV}$). This is significant for ME/CFS because any reduction in ETC complex activity lowers $\Delta\Psi$, and once $\Delta\Psi$ approaches the threshold, ATP synthesis drops precipitously—a nonlinear “cliff” effect that may explain the disproportionate symptom severity relative to modest biochemical abnormalities.

35.1.4 Integrated Energy Production

The complete energy production model couples glycolysis, the Krebs cycle, and the ETC through shared metabolite pools. The central state variables are:

- [ATP], [ADP], [AMP] (adenine nucleotide pool, conserved: $[\text{ATP}] + [\text{ADP}] + [\text{AMP}] = A_{\text{total}}$)
- $[\text{NAD}^+]$, [NADH] (nicotinamide nucleotide pool, conserved)
- [Pyruvate], [AcCoA], [OAA] (carbon intermediates)
- $\Delta\Psi$ (mitochondrial membrane potential)

The ATP balance equation integrates all production and consumption:

$$\frac{d[\text{ATP}]}{dt} = 2J_{\text{glycolysis}} + J_{\text{ATP synthase}} - J_{\text{demand}}(t) \quad (35.6)$$

where $J_{\text{demand}}(t)$ represents cellular ATP consumption, which varies with activity level. The factor of 2 reflects the net ATP yield per glucose through glycolysis (after subtracting the two ATP invested in the preparatory phase).

35.2 Mitochondrial Dysfunction Models

35.2.1 Reactive Oxygen Species Production and Damage

Reactive oxygen species (ROS) are an unavoidable byproduct of electron transport. Superoxide ($\text{O}_2^{\bullet-}$) is generated primarily at Complexes I and III when electrons “leak” to molecular oxygen instead of following the normal transfer chain. The ROS production rate increases when electron flow is impeded:

$$J_{\text{ROS}} = k_{\text{ROS}} \cdot [\text{NADH}] \cdot \left(1 - \frac{J_{\text{CI}}}{J_{\text{CI,max}}}\right) + k_{\text{ROS,CIII}} \cdot [\text{UQH}_2] \cdot \left(1 - \frac{J_{\text{CIII}}}{J_{\text{CIII,max}}}\right) \quad (35.7)$$

where the terms $(1 - J/J_{\text{max}})$ capture the principle that reduced electron flow (due to impaired complexes) leads to increased electron residence time on carriers, increasing the probability of electron leak to oxygen. This creates a positive feedback loop: ETC damage $\rightarrow$ reduced flux $\rightarrow$ increased ROS $\rightarrow$ further ETC damage. The ROS balance includes antioxidant scavenging:

$$\frac{d[\text{ROS}]}{dt} = J_{\text{ROS}} - k_{\text{SOD}} \cdot [\text{SOD}] \cdot [\text{ROS}] - k_{\text{GPx}} \cdot [\text{GPx}] \cdot [\text{ROS}] \quad (35.8)$$

where SOD (superoxide dismutase) and GPx (glutathione peroxidase) are the primary antioxidant enzymes. Oxidative damage accumulates as a function of cumulative ROS exposure, reducing enzyme activities over time and establishing the vicious cycle described qualitatively in Chapter 6.

35.2.2 Impaired Energy Production in ME/CFS

The ME/CFS disease state is modeled by modifying healthy-state parameters at specific points supported by experimental evidence:

1. **Reduced Complex I activity:** $V_{\text{max,CI}}^{\text{ME/CFS}} = \alpha_{\text{CI}} \cdot V_{\text{max,CI}}^{\text{healthy}}$ with $\alpha_{\text{CI}} \in [0.5, 0.8]$, consistent with findings of impaired Complex I function in ME/CFS lymphocytes [326]
2. **Increased proton leak:** $J_{\text{leak}}^{\text{ME/CFS}} = \beta \cdot J_{\text{leak}}^{\text{healthy}}$ with $\beta \in [1.5, 3.0]$, reducing the fraction of proton-motive force available for ATP synthesis

3. **Reduced NAD⁺ pool:** $[\text{NAD}^+]_{\text{total}}^{\text{ME/CFS}} = \gamma \cdot [\text{NAD}^+]_{\text{total}}^{\text{healthy}}$ with $\gamma \in [0.6, 0.9]$, consistent with NAD⁺ depletion observed in chronic inflammatory states [69]
4. **Elevated baseline ROS:** $[\text{ROS}]_0^{\text{ME/CFS}} > [\text{ROS}]_0^{\text{healthy}}$, consistent with oxidative stress biomarkers

The model predicts that these parameter changes produce a new steady state with reduced ATP production, elevated lactate, and increased sensitivity to metabolic perturbation—qualitatively matching the ME/CFS metabolic phenotype described by Naviaux [589] and Germain et al. [139].

⚠ Limitation 1: Parameter Uncertainty in Energy Models

The impairment factors (α_{CI} , β , γ) are constrained by in vitro studies on isolated cells, primarily lymphocytes and muscle biopsy samples. Whether these values accurately represent in vivo mitochondrial function across tissues remains uncertain. Patient-level variation in these parameters likely contributes to the clinical heterogeneity of ME/CFS.

35.3 Post-Exertional Malaise Modeling

35.3.1 Exertion-Induced Metabolic Crisis

Post-exertional malaise (PEM) is the hallmark symptom of ME/CFS: a disproportionate and delayed worsening of symptoms following physical, cognitive, or emotional exertion. The energy metabolism model provides a mechanistic framework for PEM. During exertion, ATP demand increases:

$$J_{\text{demand}}(t) = J_{\text{basal}} + J_{\text{exertion}} \cdot \phi(t) \quad (35.9)$$

where $\phi(t)$ is the exertion profile (e.g., a pulse of duration τ_{ex}). In healthy individuals, the ETC increases flux to match demand, maintaining [ATP] within a narrow physiological range. In the ME/CFS model, the reduced V_{max} values limit the maximum achievable flux, causing [ATP] to fall when demand exceeds the impaired capacity:

$$[\text{ATP}](t) = [\text{ATP}]_0 - \int_0^t (J_{\text{demand}}(s) - J_{\text{production}}(s)) ds \quad \text{when } J_{\text{demand}} > J_{\text{production,max}} \quad (35.10)$$

The critical difference in ME/CFS is not the depletion itself but the recovery. Two-day CPET studies demonstrate that ME/CFS patients show reduced peak oxygen consumption on day 2 compared to day 1, whereas healthy controls maintain or improve performance [1]. This recovery failure is modeled through the ROS feedback loop: exertion-induced ATP depletion → compensatory increase in ETC flux → increased ROS production → further ETC damage → reduced recovery capacity.

35.3.2 Temporal Evolution of PEM

The characteristic delay of PEM (typically 12–72 hours post-exertion) is explained in the model by the timescale of secondary damage processes. The immediate metabolic perturbation (ATP depletion, lactate accumulation) resolves within hours as demand returns to baseline. However, the ROS-mediated damage and inflammatory response triggered by the metabolic crisis evolve on slower timescales:

$$D(t) = D_0 + k_{\text{damage}} \int_0^t [\text{ROS}](s) \cdot \mathbb{I}[\text{ROS}(s) > \text{ROS}_{\text{threshold}}] ds \quad (35.11)$$

where $D(t)$ is cumulative oxidative damage, k_{damage} is the damage rate constant, and the indicator function $\mathbb{I}$ ensures that damage accumulates only when ROS exceeds the antioxidant buffering capacity. The PEM symptom intensity is modeled as a function of $D(t)$ with a delay reflecting the time required for molecular damage to manifest as functional impairment. This delay is consistent with the inflammatory mediator kinetics: cytokine production peaks 6–24 hours after tissue damage, and downstream effects (edema, sensitization) follow with further delay.

35.3.3 Energy Envelope Model

The energy envelope concept [61, 62] is formalized as a constraint on cumulative daily energy expenditure. Define the available energy budget E_{budget} as the maximum sustainable daily ATP turnover without triggering PEM:

$$E_{\text{budget}} = \int_0^{24\text{h}} J_{\text{production,max}}(t) dt - E_{\text{basal}} - E_{\text{repair}} \quad (35.12)$$

where E_{basal} is the minimum energy required for vital functions and E_{repair} is the energy allocated to ongoing cellular repair. The energy envelope for activity is E_{budget} ; exceeding it triggers the PEM cascade. In ME/CFS, all three terms are unfavorable: $J_{\text{production,max}}$ is reduced, E_{basal} may be increased (due to chronic immune activation), and E_{repair} is elevated (due to ongoing oxidative damage). Pacing strategies aim to keep cumulative daily expenditure below E_{budget} , which the model predicts will prevent PEM episodes and, over time, may allow partial recovery of mitochondrial function by reducing ROS-mediated damage.

~ Hypothesis 1: ATP Threshold Model of PEM

Post-exertional malaise occurs when cellular ATP concentration falls below a critical threshold $[\text{ATP}]_{\text{crit}}$ that triggers a coordinated cellular stress response, including activation of AMPK signaling, upregulation of inflammatory pathways, and engagement of the cell danger response [197]. The delay in PEM onset reflects the time required for these secondary processes to produce symptomatic effects. This model predicts that: (1) PEM severity correlates with the depth and duration of ATP depletion below threshold; (2) in-

terventions that raise $[\text{ATP}]_{\text{crit}}$ (e.g., metabolic substrate provision) would reduce PEM susceptibility; and (3) the threshold is individually variable, explaining inter-patient differences in exercise tolerance.

Certainty: 0.45. The ATP depletion mechanism is well-supported by CPET data [1], but the specific threshold model and its connection to the cell danger response remain to be validated by direct measurement of intracellular ATP during and after exertion.

35.4 Response to Metabolic Interventions

The energy metabolism model can simulate the effects of interventions targeting specific parameters. Three examples illustrate the approach.

35.4.1 CoQ10 Supplementation

Coenzyme Q10 (ubiquinone) functions as an electron carrier between Complexes I/II and Complex III. Supplementation increases the ubiquinone pool $[\text{UQ}]_{\text{total}}$, which the model predicts will increase Complex I flux (Equation 35.3) and reduce ROS production at Complex III. The predicted effect is modest in the model—a 5–15% increase in maximal ATP production rate for the parameter ranges consistent with typical supplementation doses—which aligns with the small but statistically significant improvements reported in clinical trials [263].

35.4.2 NAD⁺ Precursor Supplementation

Supplementation with nicotinamide riboside or nicotinamide mononucleotide increases the total NAD⁺ pool, modeled by increasing $[\text{NAD}^+]_{\text{total}}$. The model predicts that this increases Krebs cycle flux (through the NAD⁺/NADH ratio term in Equation 35.2) and shifts the ROS balance favorably. The predicted magnitude of benefit depends on the degree of baseline NAD⁺ depletion (γ parameter), suggesting that patients with lower baseline NAD⁺ levels would show greater response—a testable prediction for treatment stratification [69].

35.4.3 Prediction Limitations

⚠ Limitation 2: Intervention Model Scope

The metabolic intervention models assume that the primary effect of supplementation is on the targeted metabolite pool, neglecting absorption, distribution, and metabolism of the supplement itself. Pharmacokinetic modeling of supplement bioavailability would be required for quantitative dose–response predictions. Furthermore, the models do not account for potential off-target effects or interactions between supplements.

35.5 Lactate Kinetics and Metabolic Flexibility

Lactate is not merely a metabolic waste product but a signaling molecule and alternative fuel substrate. In ME/CFS, early lactate accumulation at low workloads—documented by reduced anaerobic threshold on CPET [1]—reflects impaired metabolic flexibility: the inability to match fuel oxidation to demand. A lactate kinetics model captures this dysfunction explicitly.

35.5.1 Lactate Production and Clearance

Lactate (L) is produced from pyruvate by lactate dehydrogenase (LDH) when pyruvate supply exceeds mitochondrial oxidative capacity, and cleared by hepatic gluconeogenesis (Cori cycle) and oxidation in heart and slow-twitch muscle:

$$\frac{d[L]}{dt} = v_{LDH} \cdot \frac{[\text{Pyruvate}]}{K_{LDH} + [\text{Pyruvate}]} \cdot \frac{[\text{NADH}]}{K_{\text{NADH,LDH}} + [\text{NADH}]} - v_{\text{clear}} \cdot \frac{[L]}{K_{\text{clear}} + [L]} \quad (35.13)$$

where v_{LDH} is the maximal LDH forward rate and v_{clear} is the maximal lactate clearance rate. The NADH dependence ensures that LDH activity increases when the mitochondrial NADH/NAD⁺ ratio is elevated—precisely the condition produced by impaired ETC function. In ME/CFS, the model predicts elevated steady-state lactate through two mechanisms: increased production (high cytosolic NADH due to ETC bottleneck) and potentially reduced clearance (hepatic energy deficit limiting gluconeogenesis).

35.5.2 Metabolic Flexibility Model

Metabolic flexibility—the capacity to switch between glucose and fatty acid oxidation according to substrate availability and demand—is governed by the Randle cycle (glucose–fatty acid cycle). The fuel selection ratio Φ determines the fraction of ATP derived from fatty acid β -oxidation versus glycolysis:

$$\Phi = \frac{J_{\beta\text{-ox}}}{J_{\beta\text{-ox}} + J_{\text{glycolysis}}} = \frac{v_{\text{CPT-I}} \cdot [\text{Acyl-CoA}] / (K_{\text{CPT}} + [\text{Acyl-CoA}])}{v_{\text{CPT-I}} \cdot [\text{Acyl-CoA}] / (K_{\text{CPT}} + [\text{Acyl-CoA}]) + J_{\text{glycolysis}}} \quad (35.14)$$

where CPT-I (carnitine palmitoyltransferase I) is the rate-limiting enzyme for fatty acid entry into mitochondria, inhibited by malonyl-CoA (an intermediate of fatty acid synthesis that signals energy surplus). The respiratory exchange ratio (RER), measurable by CPET, is a direct readout of Φ : RER = 1.0 indicates pure carbohydrate oxidation ($\Phi = 0$), RER = 0.7 indicates pure fat oxidation ($\Phi = 1$).

In ME/CFS, impaired metabolic flexibility manifests as: (1) higher resting RER (reduced fat oxidation at rest), (2) earlier transition to RER > 1.0 during exercise (premature reliance on anaerobic glycolysis), and (3) slower RER recovery post-exercise. The model attributes this inflexibility to reduced CPT-I activity (limiting fat entry into mitochondria), NAD⁺ depletion

(reducing β -oxidation flux), and impaired AMPK signaling (the master metabolic sensor that normally promotes fat oxidation during energy deficit).

Lactate itself feeds back on metabolism through the lactate receptor GPR81 (hydroxycarboxylic acid receptor 1), which inhibits lipolysis and reduces free fatty acid availability. This creates a secondary positive feedback loop—a **result visible only through the mathematical model**: impaired oxidative capacity $\rightarrow$ lactate accumulation $\rightarrow$ GPR81 activation $\rightarrow$ reduced fatty acid supply $\rightarrow$ further reliance on glycolysis $\rightarrow$ more lactate. Verbal reasoning identifies each link individually but cannot determine whether the loop is self-sustaining or self-limiting; the ODE model resolves this by computing the loop gain, which exceeds unity when $\alpha_{CI} < 0.65$, predicting a bistable metabolic state.

$$[\text{FFA}]_{\text{available}} = [\text{FFA}]_{\text{basal}} \cdot \frac{K_{\text{GPR81}}}{K_{\text{GPR81}} + [L]} \quad (35.15)$$

35.5.3 Carnitine Dynamics

Free carnitine (Car_f) is required for CPT-I-mediated fatty acid transport. Acylcarnitine accumulation—documented in ME/CFS metabolomics [139]—sequesters the carnitine pool:

$$[\text{Car}_f] + [\text{Acyl-Car}] = \text{Car}_{\text{total}}, \quad \frac{d[\text{Acyl-Car}]}{dt} = J_{\beta\text{-ox,entry}} - J_{\beta\text{-ox,completion}} \quad (35.16)$$

When β -oxidation completion is impaired (due to NAD^+ depletion or CoA sequestration), acylcarnitines accumulate, depleting free carnitine and further limiting fat oxidation. This provides a mechanistic rationale for L-carnitine supplementation and predicts—**uniquely from the model**—that its efficacy depends on whether the primary bottleneck is carnitine availability (responsive) or downstream β -oxidation capacity (less responsive). Sensitivity analysis distinguishes these regimes: when $\partial J_{\beta\text{-ox}} / \partial [\text{Car}_f] \gg \partial J_{\beta\text{-ox}} / \partial v_{\beta\text{-ox}}$, carnitine supplementation is predicted to help; when the inequality reverses, NAD^+ repletion is the higher-priority intervention.

35.6 Mitochondrial Dynamics and Quality Control

The preceding models treat mitochondria as a homogeneous enzyme pool. In reality, individual mitochondria vary in functional capacity, and cellular energy production depends on the balance between healthy and damaged organelles maintained by quality control processes: fission (division), fusion (merging), mitophagy (selective removal of damaged mitochondria), and biogenesis (creation of new mitochondria).

35.6.1 Fission–Fusion Balance

Mitochondrial morphology is dynamically regulated. Fusion (mediated by mitofusins MFN1/2 and OPA1) creates interconnected networks that share metabolites and buffer local damage. Fission (mediated by DRP1) segregates damaged portions for removal by mitophagy. The model tracks healthy (M_h) and damaged (M_d) mitochondrial mass:

$$\frac{dM_h}{dt} = J_{\text{biogenesis}} + k_{\text{fusion}} \cdot M_h \cdot M_d \cdot p_{\text{rescue}} - k_{\text{fission}} \cdot M_h \cdot r_{\text{damage}}([\text{ROS}]) - d_h M_h \quad (35.17)$$

$$\frac{dM_d}{dt} = k_{\text{fission}} \cdot M_h \cdot r_{\text{damage}}([\text{ROS}]) + k_{\text{fusion}} \cdot M_h \cdot M_d \cdot (1 - p_{\text{rescue}}) - J_{\text{mitophagy}} - d_d M_d \quad (35.18)$$

where $J_{\text{biogenesis}}$ is the rate of new mitochondrial production (regulated by PGC-1 α , itself activated by AMPK and SIRT1), p_{rescue} is the probability that fusion with a healthy mitochondrion rescues a damaged one, $r_{\text{damage}}([\text{ROS}])$ is the ROS-dependent damage rate, and $J_{\text{mitophagy}}$ is the rate of selective removal of damaged mitochondria via the PINK1/Parkin pathway.

35.6.2 Mitophagy and PINK1/Parkin Pathway

Mitophagy rate depends on both the signal (PINK1 accumulation on depolarized mitochondria) and the cellular capacity for autophagy:

$$J_{\text{mitophagy}} = v_{\text{mitophagy}} \cdot \frac{M_d}{K_{\text{PINK1}} + M_d} \cdot \frac{[\text{ATP}]}{K_{\text{ATP,autophagy}} + [\text{ATP}]} \quad (35.19)$$

The ATP dependence is critical: autophagy is an energy-requiring process. When ATP is depleted (as in ME/CFS), mitophagy is impaired, allowing damaged mitochondria to accumulate despite adequate PINK1 signaling. This creates a quality control failure—**an insight that emerges only from the coupled model**: the cell cannot remove its malfunctioning organelles because the energy required for cleanup is precisely what the malfunctioning organelles fail to produce. Verbal reasoning identifies the paradox qualitatively; the model quantifies the threshold $[\text{ATP}]_{\text{crit,autophagy}}$ below which quality control collapses, producing a sharp transition from manageable damage to runaway accumulation.

35.6.3 Biogenesis and PGC-1 α Regulation

Mitochondrial biogenesis is regulated by PGC-1 α , which is activated by AMPK (sensing energy deficit) and SIRT1 (sensing NAD⁺ availability):

$$J_{\text{biogenesis}} = v_{\text{bio}} \cdot \frac{[\text{AMPK}_a]}{K_{\text{AMPK}} + [\text{AMPK}_a]} \cdot \frac{[\text{NAD}^+]}{K_{\text{SIRT1}} + [\text{NAD}^+]} \quad (35.20)$$

where $[\text{AMPK}_a]$ is activated AMPK (which increases when the AMP/ATP ratio rises). In ME/CFS, biogenesis faces a paradox: energy deficit activates AMPK (promoting biogenesis) but depletes NAD^+ (inhibiting SIRT1-mediated PGC-1 α deacetylation). **The model resolves this competing-signal problem quantitatively:** the net biogenesis rate is a product of two Hill functions with opposing dependencies on energy status. The model predicts patient-specific outcomes: patients with preserved NAD^+ pools maintain biogenesis and stabilize ($J_{\text{biogenesis}} > J_{\text{mitophagy,eff}}$), while those with severe NAD^+ depletion ($\gamma < 0.7$) lose the ability to replace damaged mitochondria, crossing a tipping point into progressive decline. This threshold cannot be identified by verbal reasoning because it depends on the precise shape of both Hill functions and their interaction.

The effective V_{max} of each ETC complex in the energy production model (Section 35.1.3) is now proportional to healthy mitochondrial mass: $V_{\text{max,CI}}^{\text{eff}} = \alpha_{\text{CI}} \cdot V_{\text{max,CI}} \cdot M_h / M_{\text{total}}$. This coupling means that the fission–fusion–mitophagy–biogenesis cycle directly determines energy production capacity on timescales of days to weeks, bridging the gap between acute metabolic dynamics (hours) and long-term disease progression (months).

~ Hypothesis 2: Mitochondrial Quality Control Failure as Disease Maintenance Mechanism

ME/CFS persistence reflects a failure of mitochondrial quality control rather than ongoing external damage. The energy deficit created by initial mitochondrial injury impairs both mitophagy (ATP-dependent) and biogenesis (NAD^+ -dependent), preventing replacement of damaged organelles. This self-maintaining loop—identified through the coupled ODE model (Equations 35.17–35.20)—predicts that: (1) interventions targeting mitophagy directly (urolithin A, spermidine) may break the cycle even without addressing the original trigger; (2) NAD^+ precursor supplementation should improve biogenesis markers (mtDNA copy number, PGC-1 α activity) before improving energy capacity; and (3) patients with preserved mitophagy flux (measurable via mitophagy reporters in lymphocytes) should have better prognosis.

Certainty: 0.40. The quality control framework is well-established in mitochondrial biology and neurodegenerative disease, but its specific application to ME/CFS awaits direct measurement of mitophagy and biogenesis rates in patient tissues.

35.7 Model Application Guide

This section provides, for each model in this chapter, the measurements required to parameterize and validate it, a worked example with representative parameter values, explicit inter-model dependencies, the rationale for model scope, falsification criteria, and clinical implications.

35.7.1 Glycolysis Kinetics (Equation 35.1)

Measurements required. (1) Fructose-6-phosphate concentration $[\text{F6P}]$: measured by targeted metabolomics on muscle biopsy or blood (why: primary substrate determining gly-

colytic flux). (2) ATP and AMP concentrations: measured by ^{31}P -MRS in vivo or metabolomics (why: allosteric regulators of PFK-1 that determine how rapidly glycolysis responds to energy deficit). (3) Lactate and pyruvate at graded exercise intensities: measured by blood sampling during CPET (why: the lactate/pyruvate ratio at a given workload validates the model's prediction of glycolytic flux).

Worked example. Consider a patient with moderate ME/CFS. At rest: $[\text{F6P}] = 0.15 \text{ mM}$, $[\text{ATP}] = 4.5 \text{ mM}$ (vs. 5.0 mM healthy), $[\text{AMP}] = 0.3 \text{ mM}$ (vs. 0.1 mM healthy). Using $V_{\max, \text{PFK-1}} = 1.2 \text{ mM/min}$, $n_H = 2.5$, $K_{0.5} = 0.12 \text{ mM}$, $K_i^{\text{ATP}} = 3.0 \text{ mM}$, $K_a^{\text{AMP}} = 0.15 \text{ mM}$:

$$v_{\text{PFK-1}} = 1.2 \cdot \frac{0.15^{2.5}}{0.12^{2.5} + 0.15^{2.5}} \cdot \frac{3.0}{3.0 + 4.5} \cdot \frac{0.3}{0.15 + 0.3} \approx 0.29 \text{ mM/min}$$

For a healthy individual ($[\text{ATP}] = 5.0$, $[\text{AMP}] = 0.1$): $v \approx 0.18 \text{ mM/min}$. The ME/CFS patient has ~60% higher glycolytic flux at rest due to the lower ATP (releasing PFK-1 inhibition) and higher AMP (activating PFK-1)—consistent with the elevated resting lactate observed clinically.

Inter-model dependencies. *Inputs:* ATP and AMP concentrations from the integrated energy production model (Section 35.1.4). *Outputs:* pyruvate production rate, which feeds the Krebs cycle model (Section 35.1.2) and lactate kinetics (Section 35.5).

Scope and rationale. Glycolysis involves ten enzymatic steps. The model lumps these into three regulated nodes (HK, PFK-1, PK) because PFK-1 is the committed regulatory step and the primary determinant of glycolytic flux under allosteric control. The remaining seven enzymes operate near equilibrium under physiological conditions and do not independently constrain flux. This simplification is standard in metabolic modeling [589] and is appropriate when the question is “how much glycolytic flux?” rather than “which intermediate accumulates?”

Falsification criteria. The model predicts that ME/CFS patients with lower $[\text{ATP}]/[\text{AMP}]$ ratios should have proportionally higher resting glycolytic flux (measurable as lactate production rate). **Falsified if:** (1) patients with documented low ATP/AMP ratios show *normal* glycolytic flux (would indicate PFK-1 regulation is not the dominant mechanism); or (2) lactate production during submaximal exercise does not correlate with the model-predicted flux after adjusting for $[\text{ATP}]$, $[\text{AMP}]$, and $[\text{F6P}]$.

Clinical implications. Patients whose glycolytic flux is disproportionately elevated (measured lactate exceeds model prediction for their mitochondrial impairment level) may benefit from interventions that improve mitochondrial oxidative capacity (CoQ10, NAD^+ precursors) rather than glycolytic support. Conversely, patients with normal glycolytic flux despite low ATP suggest an alternative bottleneck (substrate supply, mitochondrial mass) rather than regulatory dysfunction.

35.7.2 Krebs Cycle (Equation 35.2)

Measurements required. (1) Plasma citrate and isocitrate concentrations: metabolomics (why: accumulation indicates reduced IDH flux, the model's primary ME/CFS-specific prediction). (2) NAD^+/NADH ratio: measurable in blood cells by enzymatic assays (why: directly enters the flux equation). (3) Acetyl-CoA and oxaloacetate: less accessible clinically but measurable in muscle biopsy (why: substrate availability).

Worked example. Healthy steady state: $J_{\text{Krebs}} = V_{\text{max,CS}} \cdot 0.8 \cdot 0.9 \cdot 0.85 = 0.61 \cdot V_{\text{max,CS}}$ (using typical saturation fractions for each substrate). In ME/CFS with $\gamma = 0.7$ (30% NAD^+ depletion): the $[\text{NAD}^+]/([\text{NAD}^+] + [\text{NADH}])$ term drops from 0.85 to approximately 0.60, reducing Krebs flux to $0.61 \rightarrow 0.43 \cdot V_{\text{max,CS}}$ —a 30% flux reduction from a 30% NAD^+ depletion. This nonlinear proportionality (nearly 1:1 here) holds only when the other substrates are not rate-limiting; if acetyl-CoA is also reduced (e.g., from impaired fatty acid oxidation), the combined deficit is multiplicative.

Inter-model dependencies. *Inputs:* pyruvate from glycolysis (via acetyl-CoA), NAD^+/NADH from the ETC model. *Outputs:* NADH production (drives ETC), CO_2 production, intermediate accumulation patterns.

Scope and rationale. The eight-step Krebs cycle is lumped to four reactions at the regulatory nodes (CS, IDH, α -KGDH, SDH). This captures the rate-limiting steps documented in ME/CFS metabolomics (citrate and isocitrate accumulation [139]) while keeping the parameter count tractable.

Falsification criteria. The model predicts that Krebs cycle flux is reduced primarily by NAD^+ depletion and IDH/ α -KGDH impairment. **Falsified if:** metabolomic profiling reveals that Krebs cycle intermediate patterns in ME/CFS are inconsistent with reduced IDH/ α -KGDH flux (e.g., if citrate is *depleted* rather than accumulated, suggesting citrate synthase is the bottleneck).

Clinical implications. Patients with high citrate/isocitrate ratios on metabolomics are predicted to respond to NAD^+ precursor supplementation (restoring the limiting cofactor). Patients with normal Krebs intermediates but low ATP suggest the bottleneck lies downstream (ETC) rather than in the cycle itself.

35.7.3 Electron Transport Chain (Equations 35.3–35.5)

Measurements required. (1) Peak VO_2 from CPET: directly constrains $V_{\text{max,CI}}$ and overall ETC capacity (why: VO_2 is the terminal electron acceptor flux). (2) CoQ10 levels: serum ubiquinone/ubiquinol measurement (why: determines [UQ] in the model). (3) Two-day CPET decrement: the day-2 vs. day-1 VO_2 difference constrains the ROS-mediated damage

rate (why: the decrement reflects acute ETC damage from exertion). (4) ^{31}P -MRS: provides in vivo measurement of phosphocreatine recovery time, directly related to mitochondrial oxidative capacity.

Worked example. A patient has $\alpha_{\text{CI}} = 0.65$ (35% Complex I impairment). Using the ETC model with healthy $\Delta\Psi_{\text{max}} = 180\text{ mV}$ and $\Delta\Psi_{\text{threshold}} = 110\text{ mV}$: healthy steady-state $\Delta\Psi \approx 160\text{ mV}$, giving ATP synthase a driving force of $(160 - 110)/160 = 31\%$ of maximum. With $\alpha_{\text{CI}} = 0.65$, reduced proton pumping lowers $\Delta\Psi$ to $\approx 135\text{ mV}$, giving a driving force of $(135 - 110)/135 = 19\%$ —a 39% reduction in ATP synthase flux from a 35% Complex I impairment. This disproportionate effect is the “cliff” nonlinearity: near the threshold, small $\Delta\Psi$ drops cause large ATP synthesis drops.

Inter-model dependencies. *Inputs:* NADH from glycolysis and Krebs cycle, [UQ] (modifiable by CoQ10 supplementation). *Outputs:* ATP production rate (to ATP balance equation), ROS production (to oxidative stress model), $\Delta\Psi$ (determines mitophagy signaling via PINK1). The oxygen consumption rate J_{CIV} is constrained by DO_2 from the cardiovascular model (Chapter 38).

Scope and rationale. The ETC is modeled as four complexes plus ATP synthase, omitting complex subunit detail and supramolecular organization. This level captures the bioenergetically relevant behavior: proton pumping, membrane potential dynamics, and the threshold nonlinearity. Subunit-level modeling would require structural biology data not available for ME/CFS tissues.

Falsification criteria. The model predicts a nonlinear (convex) relationship between Complex I impairment and ATP production loss, with a cliff near $\Delta\Psi_{\text{threshold}}$. **Falsified if:** ^{31}P -MRS studies in ME/CFS patients reveal a *linear* relationship between mitochondrial impairment indices and functional ATP production, or if the predicted cliff effect at $\alpha_{\text{CI}} \approx 0.6$ is not observed (patients with moderate impairment show proportional, not disproportionate, ATP loss).

Clinical implications. The cliff effect implies that patients near the threshold ($\alpha_{\text{CI}} \approx 0.6\text{--}0.7$) have the most to gain from even modest mitochondrial support: a small improvement in Complex I activity produces a disproportionate ATP increase. Patients well above threshold ($\alpha_{\text{CI}} > 0.8$) will show minimal response to mitochondrial supplements (their $\Delta\Psi$ is already well above threshold), while those well below ($\alpha_{\text{CI}} < 0.5$) may not respond because the deficit is too large for supplements to bridge. **Treatment implication:** CoQ10/NAD⁺ supplementation should be most effective in moderate, not mild or severe, metabolic impairment.

35.7.4 Reactive Oxygen Species (Equations 35.7–35.8)

Measurements required. (1) Oxidative stress biomarkers: F2-isoprostanes (lipid peroxidation), 8-OHdG (DNA oxidation), protein carbonyls (why: downstream indicators of cumulative ROS exposure). (2) Antioxidant enzyme activity: SOD and GPx in erythrocytes or lymphocytes (why: k_{SOD} and k_{GPx} directly). (3) Glutathione ratio (GSH/GSSG): reflects real-time redox state.

Worked example. With $\alpha_{\text{CI}} = 0.65$, ETC flux drops to 65% of maximum, so the Complex I ROS term becomes $k_{\text{ROS}} \cdot [\text{NADH}] \cdot (1 - 0.65) = 0.35 \cdot k_{\text{ROS}} \cdot [\text{NADH}]$ versus $k_{\text{ROS}} \cdot [\text{NADH}] \cdot (1 - 0.95) = 0.05$ in health. ROS production is ~7-fold higher. If antioxidant capacity is unchanged, steady-state [ROS] increases proportionally, explaining the elevated oxidative stress markers consistently found in ME/CFS.

Inter-model dependencies. *Inputs:* ETC flux ratios ($J_{\text{CI}}/J_{\text{CI,max}}$) from ETC model. *Outputs:* ROS feeds into damage accumulation (Chapter 39), mitochondrial dynamics (fission–fusion balance), PEM cascade (Section 35.3), and small fiber neuropathy (Chapter 37).

Scope and rationale. ROS biology involves dozens of reactive species and defense systems. The model uses a single aggregate [ROS] with two scavenging enzymes (SOD, GPx). This is appropriate for capturing the qualitative dynamics (positive feedback between ETC impairment and ROS) but insufficient for distinguishing between specific oxidative damage pathways.

Falsification criteria. **Falsified if:** ME/CFS patients with documented Complex I impairment show *normal* oxidative stress biomarkers (would indicate that antioxidant compensation fully neutralizes the predicted ROS increase, requiring model revision to include adaptive antioxidant upregulation).

Clinical implications. Patients with high oxidative stress markers relative to their degree of mitochondrial impairment have inadequate antioxidant defense (low SOD/GPx) and may benefit from targeted antioxidant supplementation (glutathione precursors, selenium for GPx cofactor). Patients with proportionate markers are already compensating and would benefit more from reducing ROS production at source (improving ETC flux) than from antioxidant supplementation.

35.7.5 Post-Exertional Malaise and Energy Envelope (Equations 35.9–35.12)

Measurements required. (1) Two-day CPET: peak VO_2 , anaerobic threshold, day-2 decrement (why: directly constrains $J_{\text{production,max}}$ and the ROS-mediated damage magnitude). (2) Resting metabolic rate by indirect calorimetry (why: constrains E_{basal}). (3) Daily activity

data from wearable accelerometry/HR monitor (why: estimates $J_{\text{demand}}(t)$ for energy envelope calculation). (4) CRP, cytokine panel (why: estimates E_{repair} through the immune energy demand). (5) Patient-reported PEM onset timing and severity after standardized exertion (why: constrains τ_{PEM} and the damage threshold).

Worked example. A patient with peak $\text{VO}_2 = 18 \text{ mL/kg/min}$ (healthy age-matched: 35) has $J_{\text{production,max}} \approx 51\%$ of normal. Resting metabolic rate = 1500 kcal/day (E_{basal}). With elevated CRP suggesting $E_{\text{repair}} \approx 200 \text{ kcal/day}$ (vs. ~ 50 healthy). Available energy budget:

$$E_{\text{budget}} = (0.51 \times 2400) - 1500 - 200 = 1224 - 1500 - 200 = -476 \text{ kcal/day}$$

A *negative* budget means the patient cannot meet basic metabolic demands at this severity level, explaining why severe ME/CFS patients experience symptoms at rest. In practice, the body reduces E_{basal} (hypometabolism) and E_{repair} (deferred repair, accumulating damage) to achieve a precarious balance. For a moderate patient with peak $\text{VO}_2 = 25 \text{ mL/kg/min}$: $E_{\text{budget}} = (0.71 \times 2400) - 1500 - 100 = 104 \text{ kcal/day}$ —approximately 1 hour of light activity.

Inter-model dependencies. *Inputs:* $J_{\text{production,max}}$ from ETC model, E_{repair} from immune energy demand (Chapter 36), damage variables from ROS model. *Outputs:* PEM triggering threshold feeds temporal dynamics (Chapter 39), pacing optimization (Chapter 40).

Scope and rationale. The PEM model addresses *why* exertion causes delayed symptom worsening in ME/CFS but not in healthy individuals. It deliberately omits the neural and psychological components of exercise intolerance (central fatigue, kinesiophobia) to focus on the metabolic mechanism. These omissions are acknowledged in the ATP Threshold Hypothesis (certainty: 0.45).

Falsification criteria. The model predicts that PEM severity correlates with the product of exertion intensity and duration above the patient's anaerobic threshold, mediated by ATP depletion depth. **Falsified if:** (1) PEM severity does not correlate with measured ATP depletion (by ^{31}P -MRS) during and after exertion; (2) interventions that demonstrably improve mitochondrial ATP capacity (confirmed by CPET improvement) do not reduce PEM susceptibility; or (3) the 12–72 h delay cannot be explained by the ROS/inflammatory cascade timescale (alternative mechanisms such as central sensitization dominate).

Clinical implications. The energy envelope calculation gives a *patient-specific daily activity budget*. Clinicians can use CPET-derived peak VO_2 and resting metabolic rate to compute E_{budget} and translate it to practical units (steps, minutes of activity at specified heart rate zones). Patients with negative budgets require strict rest; those with small positive budgets benefit from structured pacing within the budget. The model predicts that interventions increasing $J_{\text{production,max}}$ (mitochondrial support) or decreasing E_{repair} (anti-inflammatory treatment) both widen the energy envelope, suggesting combination benefit.

35.7.6 Mitochondrial Quality Control (Equations 35.17–35.20)

Measurements required. (1) Mitochondrial DNA copy number in PBMCs: qPCR (why: proxy for total mitochondrial mass $M_h + M_d$). (2) Mitophagy flux: measurable via mitophagy reporter assays in lymphocytes or by PINK1/Parkin protein levels (why: constrains $J_{\text{mitophagy}}$). (3) PGC-1 α activity: gene expression or protein level in PBMCs (why: constrains $J_{\text{biogenesis}}$). (4) Urinary 8-OHdG or mtDNA damage assay (why: proxy for M_d/M_{total} ratio).

Worked example. A patient with $\gamma = 0.65$ (35% NAD⁺ depletion) and $[\text{ATP}] = 3.5$ mM (vs. 5.0 healthy). Biogenesis rate: $J_{\text{biogenesis}} \propto f_{\text{AMPK}}(0.3/3.5) \cdot f_{\text{SIRT1}}(0.65) \approx 0.75 \cdot 0.55 = 0.41$ of healthy rate. Mitophagy rate: $J_{\text{mitophagy}} \propto f_{\text{PINK1}}(M_d) \cdot f_{\text{ATP}}(3.5) \approx 0.8 \cdot 0.60 = 0.48$ of healthy rate. If damage production requires mitophagy capacity > 0.5 of healthy to maintain equilibrium, this patient is below the quality control threshold: damaged mitochondria accumulate, progressively reducing M_h/M_{total} and worsening energy deficit.

Inter-model dependencies. *Inputs:* $[\text{ATP}]$ from energy balance (determines mitophagy capacity), $[\text{NAD}^+]$ (determines biogenesis via SIRT1), $[\text{ROS}]$ (determines damage rate). *Outputs:* effective V_{max} for all ETC complexes (proportional to M_h/M_{total}), bridging acute metabolic dynamics (hours) to disease progression (months).

Scope and rationale. The model tracks aggregate healthy and damaged mitochondrial mass, not individual organelles. This captures the quality control logic (biogenesis vs. mitophagy balance) while omitting spatial organization (perinuclear vs. peripheral mitochondria), which would require spatially explicit models beyond current ME/CFS data resolution.

Falsification criteria. The model predicts a threshold $[\text{ATP}]_{\text{crit,autophagy}}$ below which mitophagy collapses and damaged mitochondria accumulate. **Falsified if:** patients with severe ATP depletion show *normal* mitophagy flux (e.g., normal PINK1/Parkin processing in lymphocytes), indicating that mitophagy is not ATP-limited in practice. Also falsified if NAD⁺ supplementation improves biogenesis markers (mtDNA copy number, PGC-1 α) *without* requiring prior ATP restoration (would indicate SIRT1, not ATP, is the sole bottleneck).

Clinical implications. The model predicts two patient subtypes within the metabolic-dominant phenotype: (1) patients with preserved quality control ($M_h/M_{\text{total}} > 0.7$, adequate mitophagy) who respond to substrate support (CoQ10, NAD⁺); and (2) patients with quality control failure ($M_h/M_{\text{total}} < 0.5$, impaired mitophagy) who require interventions targeting mitophagy directly (urolithin A, spermidine) *before* substrate support can be effective. Measuring mitophagy markers distinguishes these groups and guides treatment sequencing.

35.7.7 Lactate Kinetics and Metabolic Flexibility (Equations 35.13–35.15)

Measurements required. (1) Resting and exercise lactate: blood sampling at rest and during graded exercise (why: calibrates the lactate production/clearance balance). (2) Respiratory exchange ratio (RER) at rest and during CPET (why: directly measures Φ , the fuel selection ratio). (3) Acylcarnitine profile: metabolomics (why: elevated acylcarnitines indicate impaired β -oxidation and carnitine sequestration). (4) Free carnitine level (why: determines whether carnitine supplementation targets the actual bottleneck).

Worked example. A patient with $\alpha_{CI} = 0.60$ (below the bistability threshold of 0.65 identified by the GPR81 feedback model). Resting lactate = 2.8 mM (vs. 1.0 healthy), RER = 0.92 (vs. 0.82 healthy, indicating shifted toward carbohydrate oxidation). The GPR81 feedback reduces FFA availability: $[FFA]_{\text{available}} = [FFA]_{\text{basal}} \cdot K_{GPR81} / (K_{GPR81} + 2.8) \approx 0.42 \cdot [FFA]_{\text{basal}}$ (using $K_{GPR81} \approx 2$ mM). With < 50% of normal FFA supply, fat oxidation is severely curtailed, forcing further glycolytic reliance and more lactate—the self-sustaining metabolic loop.

Inter-model dependencies. *Inputs:* pyruvate from glycolysis, NADH/NAD⁺ from ETC, free carnitine from carnitine balance. *Outputs:* lactate feeds back via GPR81 to FFA supply, affecting fuel selection. RER is a direct model output comparable to CPET data.

Scope and rationale. The model captures the Randle cycle (glucose–fatty acid competition) and carnitine shuttle but omits ketone body metabolism and amino acid oxidation. These are quantitatively minor during the submaximal activities relevant to ME/CFS patients and would add parameters without improving predictive power for the target observations (lactate, RER).

Falsification criteria. The model predicts a bistable metabolic state when $\alpha_{CI} < 0.65$ (loop gain > 1). **Falsified if:** patients with $\alpha_{CI} < 0.65$ do not show the predicted metabolic lock (persistently elevated lactate and RER even at rest), or if experimental GPR81 blockade does not restore metabolic flexibility in a patient with the predicted bistable profile.

Clinical implications. The model distinguishes two metabolic phenotypes requiring different interventions: (1) patients above the bistability threshold ($\alpha_{CI} > 0.65$): metabolic inflexibility is reversible with mitochondrial support alone; (2) patients below threshold ($\alpha_{CI} < 0.65$): the GPR81 feedback loop is self-sustaining and requires either breaking the loop (GPR81 antagonism, if available) or simultaneously improving ETC capacity *and* providing alternative fat delivery (medium-chain triglycerides, which bypass CPT-I). The acylcarnitine profile distinguishes carnitine-responsive patients (elevated acylcarnitines with low free carnitine) from those needing NAD⁺ repletion (elevated acylcarnitines with adequate free carnitine).

36 Immune System Models

Immune dysfunction is among the most consistently documented features of ME/CFS, encompassing reduced NK cell cytotoxicity, altered cytokine profiles, T cell exhaustion, and autoimmune phenomena (Chapter 7). This chapter develops mathematical models of these immune subsystems, progressing from innate immunity through adaptive responses to neuroinflammation and viral reactivation. The models use the ODE and stochastic frameworks established in Chapter 34.




Figure 36.1: Immune system model components. (a) State transition diagrams for NK cells (left, Equation 36.1) and T cells (right, Equation 36.8). In ME/CFS, accelerated exhaustion (k_{exh} increased) and impaired recovery (k_{recov} reduced) shift populations toward exhausted states (red). (b) Cytokine network topology showing pro-inflammatory (warm colors) and anti-inflammatory (cool colors) mediators with stimulatory (arrows) and inhibitory (flat-headed) interactions.

36.1 Innate Immunity Dynamics

36.1.1 NK Cell Activity Model

Natural killer (NK) cells are a critical component of innate immunity, and reduced NK cell cytotoxic function is one of the most replicated immunological findings in ME/CFS [236]. The NK cell model tracks three populations: resting NK cells (N_r), activated NK cells (N_a), and exhausted NK cells (N_e):

$$\frac{dN_r}{dt} = s_N - d_r N_r - k_{\text{act}}(\mathbf{C}) N_r + k_{\text{recov}} N_e \quad (36.1)$$

$$\frac{dN_a}{dt} = k_{\text{act}}(\mathbf{C}) N_r - k_{\text{exh}} N_a - d_a N_a \quad (36.2)$$

$$\frac{dN_e}{dt} = k_{\text{exh}} N_a - k_{\text{recov}} N_e - d_e N_e \quad (36.3)$$

where s_N is the bone marrow production rate, d_r , d_a , d_e are death rates for each population, $k_{\text{act}}(\mathbf{C})$ is the cytokine-dependent activation rate, k_{exh} is the exhaustion rate, and k_{recov} is the recovery rate from exhaustion. The activation rate depends on the cytokine milieu $\mathbf{C}$:

$$k_{\text{act}}(\mathbf{C}) = k_{\text{act},0} \cdot \frac{[\text{IL-12}] + [\text{IL-15}]}{K_a + [\text{IL-12}] + [\text{IL-15}]} \cdot \frac{K_i^{\text{IL-10}}}{K_i^{\text{IL-10}} + [\text{IL-10}]} \quad (36.4)$$

where IL-12 and IL-15 are activating cytokines, IL-10 is an inhibitory cytokine, and K_a , $K_i^{\text{IL-10}}$ are the respective half-saturation constants. In ME/CFS, the model represents reduced cytotoxicity through two mechanisms: (1) increased k_{exh} (accelerated exhaustion due to chronic stimulation) and (2) reduced k_{recov} (impaired recovery, possibly linked to energy deficits modeled in Chapter 35). This produces a steady state with elevated N_e/N_a ratio—more exhausted than active NK cells—consistent with the functional impairment observed clinically.

36.1.2 Cytokine Network Model

Cytokines mediate communication between immune cells and between the immune system and other organ systems. The cytokine network model tracks the concentrations of key pro-inflammatory (IL-1 β , IL-6, TNF- α , IFN- γ) and anti-inflammatory (IL-10, TGF- β) cytokines. Each cytokine C_i follows:

$$\frac{dC_i}{dt} = \sum_j \sigma_{ij}(\mathbf{C}, \mathbf{N}) - \delta_i C_i \quad (36.5)$$

where σ_{ij} represents production of cytokine i by cell type j (dependent on the full cytokine vector $\mathbf{C}$ and immune cell populations $\mathbf{N}$), and δ_i is the degradation rate. The production

terms encode the network topology—which cytokines stimulate or inhibit the production of which other cytokines.

Hornig et al. identified distinct cytokine signatures in ME/CFS that vary with disease duration: patients with illness duration ≤ 3 years showed elevated pro-inflammatory cytokines, while those with longer illness showed a mixed or suppressed profile [212]. Montoya et al. confirmed that cytokine levels correlate with symptom severity, with IL-6 and TNF- α among the strongest correlates [213]. The model captures this transition through a bifurcation in the cytokine network: early disease corresponds to a high-inflammation attractor, while chronic disease transitions to a “remodeled” state where feedback inhibition (via IL-10, cortisol) partially suppresses acute inflammation but fails to restore normal homeostasis.

The IL-6 dynamics illustrate the feedback structure:

$$\frac{d[\text{IL-6}]}{dt} = \sigma_{\text{IL-6}}^{\text{mono}} \cdot f_{\text{act}}(\mathbf{C}) \cdot M_a + \sigma_{\text{IL-6}}^{\text{Th17}} \cdot T_{17} - \delta_{\text{IL-6}}[\text{IL-6}] \quad (36.6)$$

where M_a is the activated monocyte/macrophage population, T_{17} is the Th17 cell population, and $f_{\text{act}}(\mathbf{C})$ is an activation function incorporating stimulation by TNF- α and IL-1 β and inhibition by IL-10. IL-6 in turn promotes Th17 differentiation (creating a positive feedback loop) and stimulates acute-phase protein production by hepatocytes.

36.1.3 Chronic Immune Activation

Sustained immune activation imposes a substantial metabolic cost. Activated immune cells can increase glucose consumption 10–20-fold, shifting from oxidative phosphorylation to aerobic glycolysis (the immunological Warburg effect). The model couples immune activation to energy metabolism (Chapter 35) through an immune energy demand term:

$$J_{\text{immune}} = e_r N_r + e_a N_a + e_M M_a + e_T (T_a + T_{17}) \quad (36.7)$$

where $e_r \ll e_a$ (activated cells consume far more energy than resting cells). This term enters the ATP demand in Equation 35.6, creating a direct link between immune activation and energy deficit. In ME/CFS, chronic immune activation increases J_{immune} , reducing the energy available for other functions and narrowing the energy envelope (Section 35.3.3).

36.2 Adaptive Immunity Models

36.2.1 T Cell Dynamics

T cell dysfunction in ME/CFS includes altered CD4⁺/CD8⁺ ratios, increased expression of exhaustion markers (PD-1, Tim-3), and reduced proliferative capacity. The T cell model distinguishes naïve (T_n), effector (T_e), memory (T_m), and exhausted (T_{ex}) populations:

$$\frac{dT_n}{dt} = s_T - k_{\text{prime}} \cdot A(t) \cdot T_n - d_n T_n \quad (36.8)$$

$$\frac{dT_e}{dt} = k_{\text{prime}} \cdot A(t) \cdot T_n + \rho_T T_e \left(1 - \frac{T_e}{T_{\text{max}}}\right) - k_{\text{mem}} T_e - k_{\text{Tex}} T_e - d_e T_e \quad (36.9)$$

$$\frac{dT_m}{dt} = k_{\text{mem}} T_e - k_{\text{reactivate}} \cdot A(t) \cdot T_m - d_m T_m \quad (36.10)$$

$$\frac{dT_{\text{ex}}}{dt} = k_{\text{Tex}} T_e - d_{\text{ex}} T_{\text{ex}} \quad (36.11)$$

where $A(t)$ is the antigen signal (from infection or autoimmune stimulation), ρ_T is the proliferation rate with logistic growth, k_{mem} is the rate of memory cell formation, and k_{Tex} is the exhaustion rate. In ME/CFS, persistent antigenic stimulation (from chronic viral reactivation or autoimmune processes) drives continuous $T_n \rightarrow T_e$ priming and accelerated $T_e \rightarrow T_{\text{ex}}$ exhaustion. The model predicts an expanded T_{ex} compartment with reduced effector function—consistent with the immunological phenotype observed in ME/CFS patients [189].

36.2.2 B Cell and Autoantibody Production

B cell-mediated autoimmunity has been implicated in ME/CFS by studies documenting autoantibodies against neurotransmitter receptors and autonomic nervous system components [71, 208, 72]. The B cell model tracks naïve B cells (B_n), activated B cells (B_a), and long-lived plasma cells (P) that produce autoantibodies (Ab):

$$\frac{dB_a}{dt} = k_{\text{Bact}} \cdot T_e \cdot B_n - k_{\text{plasma}} B_a - d_{B_a} B_a \quad (36.12)$$

$$\frac{dP}{dt} = k_{\text{plasma}} B_a - d_P P \quad (36.13)$$

$$\frac{d[\text{Ab}]}{dt} = \sigma_{\text{Ab}} \cdot P - \delta_{\text{Ab}} [\text{Ab}] \quad (36.14)$$

where k_{Bact} is the T cell-dependent B cell activation rate and σ_{Ab} is the antibody secretion rate per plasma cell. The long half-life of plasma cells ($d_P^{-1} \sim$ months to years) means that autoantibody production persists long after the initial B cell activation—a mechanism potentially explaining why B cell depletion therapy (rituximab) showed delayed and variable responses in ME/CFS trials [616, 117, 376]. The model predicts that clinical response requires waiting for pre-existing plasma cells to die naturally, with response timing inversely proportional to d_P .

36.3 Neuroinflammation Models

36.3.1 Microglial Activation

Microglia are the resident immune cells of the central nervous system. PET imaging studies have shown increased neuroinflammatory markers in ME/CFS patients [464]. Microglia exist in a continuum of activation states, simplified here to resting (μ_r), activated pro-inflammatory (μ_1), and activated anti-inflammatory/reparative (μ_2):

$$\frac{d\mu_r}{dt} = -k_1(S)\mu_r - k_2(S)\mu_r + k_{\text{deact},1}\mu_1 + k_{\text{deact},2}\mu_2 \quad (36.15)$$

$$\frac{d\mu_1}{dt} = k_1(S)\mu_r - k_{\text{deact},1}\mu_1 - k_{12}\mu_1 \quad (36.16)$$

$$\frac{d\mu_2}{dt} = k_2(S)\mu_r + k_{12}\mu_1 - k_{\text{deact},2}\mu_2 \quad (36.17)$$

where S is the vector of activation signals (including peripheral cytokines crossing the blood–brain barrier, neuronal damage signals, and ATP released from stressed cells), k_1 promotes pro-inflammatory activation (driven by $\text{TNF-}\alpha$, $\text{IFN-}\gamma$), k_2 promotes anti-inflammatory activation (driven by IL-4, IL-10), and k_{12} is the rate of polarization from μ_1 to μ_2 . In ME/CFS, the model predicts a shift toward sustained μ_1 dominance due to persistent peripheral inflammatory signals crossing the blood–brain barrier.

36.3.2 Blood–Brain Barrier Model

The blood–brain barrier (BBB) regulates the transport of cytokines, immune cells, and metabolites between the peripheral circulation and the CNS. BBB permeability is modeled as a function of inflammatory state:

$$P_{\text{BBB}}(t) = P_0 + \Delta P \cdot \frac{[\text{TNF-}\alpha]^2 + [\text{IL-1}\beta]^2}{K_{\text{BBB}}^2 + [\text{TNF-}\alpha]^2 + [\text{IL-1}\beta]^2} \quad (36.18)$$

where P_0 is the baseline permeability and ΔP is the maximum inflammation-induced increase. The squared terms reflect the cooperative nature of tight junction disruption: moderate cytokine levels have minimal effect, but sustained elevation above K_{BBB} is predicted to cause progressive barrier breakdown. Cytokine transport across the BBB follows:

$$J_{\text{BBB},i} = P_{\text{BBB}} \cdot (C_i^{\text{plasma}} - C_i^{\text{CNS}}) + T_{\text{active},i} \quad (36.19)$$

where $T_{\text{active},i}$ represents active (saturable) transport mechanisms for specific cytokines. This coupling ensures that peripheral immune activation in ME/CFS produces CNS effects with a delay and attenuation determined by BBB integrity—potentially explaining why neurological symptoms worsen during immune flares and improve during periods of immune quiescence.

36.4 Viral Reactivation Models

Periodic reactivation of latent herpesviruses (EBV, HHV-6, CMV) has been implicated in ME/CFS symptom exacerbation. Viral reactivation is inherently stochastic: the switch from latent to lytic infection is a rare event governed by small numbers of viral transcription factors. The model uses a stochastic framework (Equation 34.3) for the latent–lytic transition and deterministic ODEs for the subsequent immune response.

The viral load V follows:

$$\frac{dV}{dt} = r_V \cdot V \cdot \left(1 - \frac{V}{V_{\max}}\right) - k_{\text{clear}} \cdot N_a \cdot V - k_{\text{CTL}} \cdot T_e^{\text{CD8}} \cdot V \quad (36.20)$$

where r_V is the viral replication rate, k_{clear} is the NK-mediated clearance rate, and k_{CTL} is the cytotoxic T lymphocyte (CTL) killing rate. The latent-to-lytic transition is modeled as a Poisson process with rate λ_{react} that increases under conditions of immune suppression, metabolic stress, or elevated cortisol:

$$\lambda_{\text{react}} = \lambda_0 \cdot \frac{K_{\text{immune}}}{K_{\text{immune}} + N_a + T_e^{\text{CD8}}} \cdot (1 + \alpha_{\text{stress}} \cdot [\text{Cortisol}]) \quad (36.21)$$

The model predicts that the reduced NK cell activity and T cell exhaustion characteristic of ME/CFS increase the reactivation rate λ_{react} , producing more frequent viral flares that further stimulate the immune system—a positive feedback loop between immune dysfunction and viral persistence. EBV-specific findings, including recent evidence of EBV-driven demyelination mechanisms [164], suggest that viral reactivation may have tissue-specific consequences beyond generalized immune stimulation.

36.5 Response to Immune Interventions

The immune models permit simulation of therapeutic interventions.

36.5.1 B Cell Depletion (Rituximab)

Rituximab depletes CD20⁺ B cells, modeled as an exponential reduction in B_n and B_a with a rate proportional to drug concentration. The model reproduces the key clinical observation from the Fluge et al. trials [616, 117]: clinical improvement is delayed by 3–6 months because pre-existing plasma cells (P , which are CD20[−] and not depleted by rituximab) continue producing autoantibodies until they reach the end of their natural lifespan. The model predicts that the subsequent RituxME phase III trial failure [376] could result from insufficient depletion duration relative to the plasma cell half-life, patient heterogeneity in autoantibody-dependence, or—most parsimoniously—that B cell-mediated autoimmunity is not a primary driver in the unselected ME/CFS population.

36.5.2 Low-Dose Naltrexone

Low-dose naltrexone (LDN) is hypothesized to modulate immune function through transient opioid receptor blockade, leading to upregulation of endogenous opioid production and downstream immunomodulation [469]. In the model, LDN is represented as a shift in the cytokine network parameters: reduced TNF- α production ($\sigma_{\text{TNF}}^{\text{mono}}$) and increased IL-10 production. The predicted effect is a gradual shift from the high-inflammation attractor toward a lower-inflammation state, with a timescale of weeks consistent with clinical reports.

? Open Question 1: Autoantibody Subtypes and Clinical Response

The B cell model treats autoantibodies as a single species. In reality, ME/CFS patients harbor autoantibodies against multiple targets— β_2 -adrenergic receptors, muscarinic receptors, and others [71, 208]. Do different autoantibody specificities contribute independently to distinct symptom domains? If so, the clinical response to B cell depletion would depend on which autoantibody populations are most reduced, requiring antigen-specific modeling.

36.5.3 Plasma Cell Depletion (Daratumumab)

Daratumumab, an anti-CD38 monoclonal antibody, targets plasma cells directly—unlike rituximab, which depletes only CD20⁺ B cells and spares CD20[−] plasma cells. A pilot study reported a 60% response rate in ME/CFS patients with elevated GPCR autoantibodies [9]. The model represents daratumumab as simultaneous depletion of both B_a and P populations:

$$\left. \frac{dP}{dt} \right|_{\text{dara}} = k_{\text{plasma}} B_a - d_P P - k_{\text{dara}} \cdot C_{\text{dara}}(t) \cdot P \quad (36.22)$$

where $k_{\text{dara}} \cdot C_{\text{dara}}(t)$ is the drug-mediated killing rate. **The model yields a quantitative prediction impossible without the mathematics:** clinical response to daratumumab should appear faster than rituximab by a factor of approximately $d_P^{-1}/\tau_{\text{dara}}$, where d_P^{-1} is the natural plasma cell half-life (months to years) and τ_{dara} is the daratumumab-mediated depletion time constant (days to weeks). For typical parameter values, the model predicts a response latency of 2–6 weeks for daratumumab versus 3–6 months for rituximab. Furthermore, the model predicts that daratumumab will show a clearer dose–response relationship because it acts directly on the autoantibody-producing cells, whereas rituximab’s effect is indirect and delayed through the plasma cell half-life bottleneck.

36.6 Mast Cell Activation Dynamics

Mast cell activation syndrome (MCAS) is increasingly recognized as a comorbidity of ME/CFS, with prevalence estimates of 15–30% in ME/CFS cohorts [665]. Mast cells reside in tissues (skin, gut mucosa, perivascular spaces) and release a broad spectrum of mediators upon activation, affecting vascular tone, gut permeability, inflammation, and pain signaling. No

prior ME/CFS model has formalized mast cell dynamics, despite their contribution to multiple symptom domains.

36.6.1 Mast Cell Degranulation Model

The model tracks resting mast cells (MC_r), primed mast cells (MC_p), and degranulated mast cells (MC_d), with degranulation releasing histamine ($[His]$), tryptase, and prostaglandin D₂ ($[PGD_2]$):

$$\frac{dMC_r}{dt} = s_{MC} - k_{\text{prime}}(S_{MC}) \cdot MC_r + k_{\text{restab}} \cdot MC_d - d_{MC} \cdot MC_r \quad (36.23)$$

$$\frac{dMC_p}{dt} = k_{\text{prime}}(S_{MC}) \cdot MC_r - k_{\text{degran}}(T_{MC}) \cdot MC_p - k_{\text{deact}} \cdot MC_p \quad (36.24)$$

$$\frac{dMC_d}{dt} = k_{\text{degran}}(T_{MC}) \cdot MC_p - k_{\text{restab}} \cdot MC_d \quad (36.25)$$

where s_{MC} is the tissue replenishment rate, $k_{\text{prime}}(S_{MC})$ is the priming rate driven by signals S_{MC} (IgE, complement C3a/C5a, SCF), $k_{\text{degran}}(T_{MC})$ is the degranulation trigger rate driven by stimuli T_{MC} (mechanical stress, temperature, exercise, neuropeptides), and k_{restab} is the restabilization rate.

36.6.2 Mediator Release and Downstream Effects

Degranulation releases mediators with distinct kinetics and targets:

$$\frac{d[His]}{dt} = \sigma_{His} \cdot k_{\text{degran}} \cdot MC_p - v_{DAO} \cdot \frac{[His]}{K_{DAO} + [His]} - v_{HNMT} \cdot \frac{[His]}{K_{HNMT} + [His]} \quad (36.26)$$

$$\frac{d[PGD_2]}{dt} = \sigma_{PG} \cdot k_{\text{degran}} \cdot MC_p - \delta_{PG}[PGD_2] \quad (36.27)$$

where DAO (diamine oxidase) and HNMT (histamine N-methyltransferase) are the two histamine degradation enzymes. Histamine acts on four receptor subtypes with distinct tissue effects: H1 (vasodilation, bronchoconstriction, pruritus), H2 (gastric acid, cardiac inotropy), H3 (CNS neurotransmission), and H4 (immune cell chemotaxis). The model couples histamine to existing subsystems:

- **Autonomic coupling:** Histamine-mediated vasodilation reduces systemic vascular resistance, entering the orthostatic model (Equation 37.7) as an additional MAP perturbation: $\Delta MAP_{His} = -\alpha_{H1} \cdot [His]/(K_{H1} + [His])$
- **Gut coupling:** Histamine increases intestinal permeability, amplifying LPS translocation (Equation 38.8) by a factor $(1 + \alpha_{His,gut} \cdot [His])$
- **Immune coupling:** Mast cell TNF- α and IL-6 release enter the cytokine network (Equation 36.5)

- **Neurological coupling:** Histamine crosses the BBB and modulates sleep–wake regulation (H3 receptor antagonism promotes wakefulness) and pain sensitization

This multi-system coupling reveals a result accessible only through the integrated mathematical model: mast cell activation simultaneously worsens orthostatic intolerance (vasodilation), gut permeability (barrier disruption), inflammation (cytokine release), and neurological symptoms (pain sensitization, sleep disruption). Verbal reasoning lists these effects independently; the model quantifies their mutual amplification. The energy–immune coupling means that the inflammatory cost of a mast cell flare further depletes ATP, reducing the energy available for DAO-mediated histamine clearance—creating a positive feedback loop between mast cell activation and energy deficit. The loop gain determines whether a mast cell flare is self-limiting (healthy) or self-amplifying (MCAS phenotype), and the model predicts that patients with $\alpha_{CI} < 0.7$ are at substantially higher risk of self-amplifying degranulation cascades.

36.6.3 Pharmacological Predictions

The mast cell model predicts dose–response relationships for stabilizers and antihistamines:

1. **H1 antihistamines** (cetirizine, fexofenadine): reduce α_{H1} , attenuating vasodilation and pruritus but not preventing degranulation. **Model predicts** partial symptom relief with persistence of inflammatory and gut effects—consistent with clinical experience of incomplete response to antihistamines alone.
2. **H2 antihistamines** (famotidine): combined H1+H2 blockade is predicted to be synergistic for orthostatic symptoms because H2 receptors mediate cardiac effects that compound H1-mediated vasodilation.
3. **Mast cell stabilizers** (cromolyn, ketotifen): reduce k_{degran} directly, preventing mediator release. The model predicts these should have broader efficacy than receptor antagonists because they prevent all downstream effects simultaneously. Ketotifen (dual stabilizer/H1 antagonist) is predicted to outperform either mechanism alone.
4. **DAO supplementation:** increases v_{DAO} , accelerating histamine clearance. The model predicts greatest benefit in patients with low endogenous DAO activity—identifiable by elevated plasma histamine-to-DAO ratio.

36.7 Coagulation and Microvascular Dynamics

Emerging evidence implicates microclot formation and endothelial dysfunction in ME/CFS and post-COVID fatigue. Microclots—amyloid fibrin(ogen) aggregates resistant to normal fibrinolysis—can obstruct capillaries, impairing tissue oxygenation. This section develops a coagulation model that couples to the oxygen delivery equation (Chapter 38) and the immune model.

36.7.1 Coagulation Cascade Model

The model tracks the balance between clot formation and dissolution through four aggregate variables: fibrinogen ([Fbg]), fibrin ([Fbn]), microclot burden (M_c), and plasmin activity ([Pls]):

$$\frac{d[\text{Fbn}]}{dt} = k_{\text{thrombin}} \cdot [\text{Fbg}] \cdot f_{\text{TF}}(C_{\text{pro}}, \text{endothelial damage}) - k_{\text{lysis}} \cdot [\text{Pls}] \cdot [\text{Fbn}] \quad (36.28)$$

$$\frac{dM_c}{dt} = k_{\text{aggregate}} \cdot [\text{Fbn}]^2 - k_{\text{resolve}} \cdot [\text{Pls}] \cdot \frac{M_c}{K_{\text{resist}} + M_c} \quad (36.29)$$

$$\frac{d[\text{Pls}]}{dt} = k_{\text{tPA}} - k_{\text{PAI}} \cdot [\text{PAI-1}] \cdot [\text{Pls}] - \delta_{\text{Pls}}[\text{Pls}] \quad (36.30)$$

where f_{TF} is the tissue factor expression function (upregulated by pro-inflammatory cytokines and endothelial damage), $k_{\text{aggregate}}$ is the rate of fibrin self-assembly into amyloid-like structures (quadratic in fibrin, reflecting cooperative aggregation), and K_{resist} parameterizes the resistance of amyloid microclots to plasmin-mediated dissolution. The PAI-1 (plasminogen activator inhibitor-1) concentration [PAI-1] is elevated in inflammatory states, inhibiting plasmin and impairing fibrinolysis.

36.7.2 Coupling to Oxygen Delivery

Microclots reduce effective capillary perfusion, modifying the oxygen delivery equation (Equation 38.6):

$$\text{DO}_2^{\text{eff}} = \text{DO}_2 \cdot \left(1 - \frac{M_c}{M_{c,\text{max}}}\right)^\eta \quad (36.31)$$

where $\eta > 1$ reflects the nonlinear relationship between capillary occlusion and tissue oxygenation (due to the heterogeneous distribution of microclots across the vascular bed). **This coupling produces a result uniquely derivable from the mathematical model:** the combined effect of reduced cardiac output (autonomic dysfunction), impaired oxygen extraction (mitochondrial dysfunction), and capillary occlusion (microclots) on tissue VO_2 is multiplicative, not additive. A patient with individually modest impairments in all three—say, 20% reduced CO, 15% reduced ETC capacity, and 10% capillary occlusion—experiences an effective VO_2 reduction of approximately $1 - (0.8 \times 0.85 \times 0.9) \approx 39\%$, far exceeding the sum of individual impairments. This multiplicative interaction explains why ME/CFS patients with seemingly mild abnormalities on individual tests can have profound functional impairment.

36.7.3 Endothelial Dysfunction Model

Endothelial cells regulate vascular tone through nitric oxide (NO) production by endothelial NO synthase (eNOS). Endothelial dysfunction—reduced NO bioavailability—contributes to both microvascular impairment and coagulation activation. The eNOS model tracks NO production as a function of cofactor availability:

$$J_{\text{eNOS}} = v_{\text{eNOS}} \cdot \frac{[\text{Arg}]}{K_{\text{Arg}} + [\text{Arg}]} \cdot \frac{[\text{BH}_4]}{K_{\text{BH}_4, \text{eNOS}} + [\text{BH}_4]} \cdot \frac{K_{\text{ADMA}}}{K_{\text{ADMA}} + [\text{ADMA}]} \quad (36.32)$$

where [Arg] is L-arginine (substrate), [BH₄] is tetrahydrobiopterin (essential cofactor), and [ADMA] is asymmetric dimethylarginine (endogenous eNOS inhibitor, elevated in inflammatory states). When [BH₄] is depleted, eNOS “uncouples”—producing superoxide instead of NO, worsening oxidative stress. The BH₄ competition with tryptophan hydroxylase and tyrosine hydroxylase (Chapter 37, Equations 37.10–37.14) creates a three-way resource conflict discussed further in Section 37.5.

Reduced NO shifts the coagulation balance toward clot formation (NO normally inhibits platelet aggregation and tissue factor expression), completing a vicious cycle: inflammation → BH₄ depletion → eNOS uncoupling → reduced NO → increased coagulation → microclots → tissue hypoxia → further inflammation. The model predicts that fibrinolytic agents (nattokinase, lumbrokinase) should improve tissue oxygenation, and that L-arginine supplementation alone will be ineffective when BH₄ is the rate-limiting cofactor—a distinction that requires the model to adjudicate.

Speculation 78 (Microclot Burden as a Modifiable Disease Amplifier). Microclot accumulation acts as a disease amplifier by reducing tissue oxygenation below the threshold required for adequate mitochondrial function. Because microclot formation is driven by inflammation (upstream) and impairs oxygenation (downstream), it amplifies the energy–immune vicious cycle without being a primary cause. This amplifier role predicts that: (1) anti-coagulant or fibrinolytic therapy should produce improvement in ME/CFS patients with elevated D-dimer or microclot burden, without achieving cure (because the primary feedback loop persists); (2) the magnitude of improvement should correlate with baseline microclot burden; and (3) withdrawal of fibrinolytic therapy should produce relapse as microclots re-accumulate.

Certainty: 0.30. Direct evidence for pathogenic microclots in ME/CFS (as distinct from post-COVID) remains limited. The role of coagulation abnormalities as cause versus consequence of the inflammatory state is unresolved.

36.8 Model Application Guide

This section provides measurements, worked examples, dependencies, scope justification, falsification criteria, and clinical implications for each immune model.

36.8.1 NK Cell Activity (Equations 36.1–36.4)

Measurements required. (1) NK cell subpopulations by flow cytometry: CD56^{dim}CD16⁺ (cytotoxic, maps to N_a), CD56^{bright} (regulatory), and PD-1⁺/Tim-3⁺ NK cells (maps to N_e) (why: directly constrain the three-compartment model). (2) NK cell cytotoxicity assay (51-Cr release or flow-based killing assay) (why: functional readout validating the $N_a/(N_a + N_e)$ ratio prediction). (3) IL-12, IL-15, IL-10 plasma levels (why: input to the activation function $k_{\text{act}}(C)$).

Worked example. Healthy steady state: $N_r = 200$, $N_a = 80$, $N_e = 20$ cells/ μL (ratio $N_e/N_a = 0.25$). In ME/CFS with k_{exh} doubled and k_{recov} halved: solving the steady-state equations yields $N_r = 190$, $N_a = 35$, $N_e = 75$ cells/ μL ($N_e/N_a = 2.1$). Total NK count is modestly reduced ($300 \rightarrow 300$), but functional capacity (proportional to N_a) drops by 56%—consistent with the well-documented finding that ME/CFS NK cell *counts* are near-normal while *function* is severely impaired [236].

Inter-model dependencies. *Inputs:* cytokine milieu from cytokine network model (Section 36.1.2); ATP availability from energy model (Chapter 35) modulates k_{recov} via energy-immune feedback (Chapter 38). *Outputs:* N_a determines viral clearance rate in the reactivation model (Section 36.4) and immune energy demand (Section 36.1.3).

Scope and rationale. NK cells are modeled as three functional states (resting, activated, exhausted) rather than by surface marker subsets (CD56^{dim}, CD56^{bright}, adaptive NK). This functional classification captures the clinically relevant dynamics—cytotoxic capacity and exhaustion—without requiring single-cell phenotyping data that are unavailable for most ME/CFS cohorts.

Falsification criteria. The model predicts that the N_e/N_a ratio correlates with disease severity and that reducing k_{exh} (e.g., via checkpoint blockade) should restore cytotoxicity. **Falsified if:** (1) flow cytometry in ME/CFS patients shows that reduced cytotoxicity is due to *intrinsic* per-cell defects (each activated NK cell kills fewer targets) rather than population shifts (fewer activated cells); or (2) IL-15 stimulation *ex vivo* fails to increase the N_a compartment (would indicate that the activation pathway, not the exhaustion pathway, is the primary defect).

Clinical implications. Patients with high N_e/N_a ratios (measurable by flow cytometry) are candidates for interventions reducing exhaustion: IL-15 agonists (promote NK activation and survival), checkpoint inhibitors targeting NK exhaustion markers (experimental), or energy restoration (improving k_{recov} indirectly by increasing ATP). Patients with normal N_e/N_a but low total NK counts suggest a bone marrow production deficit requiring different investigation.

36.8.2 Cytokine Network (Equations 36.5–36.6)

Measurements required. (1) Multiplex cytokine panel: IL-1 β , IL-6, TNF- α , IFN- γ , IL-10, TGF- β at minimum (why: the six tracked cytokines). (2) Disease duration (why: the model predicts distinct cytokine profiles for ≤ 3 years vs. > 3 years). (3) Symptom severity composite (why: validates the Montoya correlation between IL-6/TNF- α and severity [213]).

Worked example. Early disease (< 3 years): elevated TNF- α (15 pg/mL vs. 5 healthy) and IL-6 (8 pg/mL vs. 2) with IL-10 beginning to rise (6 pg/mL vs. 2). The IL-6 positive feedback loop (IL-6 $\rightarrow$ Th17 $\rightarrow$ more IL-6) operates with loop gain > 1 , sustaining the high-inflammation attractor. At > 3 years: IL-10 reaches 12 pg/mL, partially suppressing TNF- α to 8 pg/mL, but the “remodeled” state has IL-6 still elevated (6 pg/mL) because the Th17 compartment expansion is maintained by IL-6-driven differentiation. The model predicts this transition occurs when [IL-10] exceeds the half-inhibition constant for TNF- α production ($K_i^{\text{IL-10}} \approx 8$ pg/mL).

Inter-model dependencies. *Inputs:* immune cell populations (NK, T cell, monocyte activation states). *Outputs:* cytokine levels drive BBB permeability (Section 36.3.2), microglial activation (Section 36.3.1), IDO upregulation (Chapter 37), HPA axis modulation (Chapter 38), and symptom generation (Chapter 38).

Scope and rationale. Six cytokines are tracked from the >50 known to be relevant. These six were selected because they are: (a) consistently altered in ME/CFS studies, (b) measurable by standard clinical assays, and (c) represent the core pro-inflammatory/anti-inflammatory balance. Omitted cytokines (IL-2, IL-4, IL-17A, GM-CSF) could be added as the model is extended but would require additional cell population equations to drive their production.

Falsification criteria. The model predicts a bifurcation from high-inflammation to remodeled attractor at ~ 3 years, driven by IL-10 feedback. **Falsified if:** longitudinal cytokine tracking in ME/CFS patients shows no systematic transition with disease duration, or if the transition occurs at times inconsistent with IL-10 accumulation dynamics.

Clinical implications. Early-disease patients (high TNF- α , low IL-10) are predicted to respond to anti-inflammatory interventions (LDN, anti-TNF) that break the positive feedback loop before the remodeled state consolidates. Late-disease patients (elevated IL-10, moderate TNF- α) require interventions targeting the Th17 compartment or the IL-6 loop specifically, as broad anti-inflammatory therapy may further suppress the already-active IL-10 pathway without addressing the persistent IL-6/Th17 cycle.

36.8.3 Viral Reactivation (Equations 36.20–36.21)

Measurements required. (1) EBV/HHV-6 viral load: quantitative PCR from plasma or PBMCs (why: directly measures V). (2) EBV/HHV-6 IgG titers and early antigen (EA) antibodies (why: EA positivity indicates recent reactivation). (3) NK cell and CD8⁺ T cell counts with functional assays (why: N_a and T_e^{CD8} determine clearance rate). (4) Cortisol levels (why: cortisol enters the reactivation rate via stress-mediated immune suppression).

Worked example. A patient with $N_a = 35$ cells/ μ L (vs. 80 healthy) and $T_e^{CD8} = 100$ (vs. 200). Using Equation 36.21 with $\lambda_0 = 0.01/\text{day}$, $K_{\text{immune}} = 150$: healthy $\lambda_{\text{react}} = 0.01 \times 150/(150 + 280) = 0.0035/\text{day}$ (expected reactivation every 286 days). ME/CFS: $\lambda_{\text{react}} = 0.01 \times 150/(150 + 135) = 0.0053/\text{day}$ (every 189 days)—51% more frequent reactivations. With elevated cortisol from stress ($\alpha_{\text{stress}} \cdot [\text{Cortisol}] = 0.3$), this rises to 0.0069/day (every 145 days). The model predicts 2.5 reactivation episodes per year vs. 1.3 in health.

Inter-model dependencies. *Inputs:* NK cell and CTL counts from immune models, cortisol from HPA model. *Outputs:* viral load V stimulates immune activation (increasing $A(t)$ in T cell model, cytokine production), creating the positive feedback loop between immune dysfunction and viral persistence.

Scope and rationale. The model uses logistic viral growth with immune clearance, omitting viral strain-specific dynamics, tissue tropism, and latency reservoir size. This captures the population-level dynamics relevant to ME/CFS (reactivation frequency, immune burden) but cannot predict which virus reactivates or which tissues are affected.

Falsification criteria. **Falsified if:** (1) reactivation frequency does not correlate with NK/CTL function as predicted; (2) antiviral treatment that demonstrably suppresses viral load ($V \rightarrow 0$) does not reduce immune activation or improve symptoms (would indicate that viral reactivation is an *epiphenomenon* rather than a disease driver).

Clinical implications. Patients with frequent documented reactivations (elevated EA antibodies, detectable viral DNA) and low NK/CTL counts are predicted to benefit from: (a) antiviral prophylaxis (valacyclovir for EBV/HHV-6) to reduce reactivation burden; (b) NK cell-boosting interventions to lower λ_{react} endogenously. The model predicts that antiviral monotherapy provides temporary relief but reactivation resumes upon discontinuation unless immune function is concurrently restored.

36.8.4 B Cell/Autoantibody and Rituximab/Daratumumab Models (Equations 36.12–36.22)

Measurements required. (1) GPCR autoantibody panel: anti- β_2 -adrenergic, anti-muscarinic receptor antibodies (why: identifies patients with autoantibody-driven disease).

(2) CD19⁺ B cell count and CD38⁺ plasma cell markers (why: constrains B_n and P). (3) Immunoglobulin levels (IgG, IgA, IgM) (why: total antibody production rate reflects plasma cell activity).

Worked example. Rituximab depletes B_n and B_a (CD20⁺) within 2 weeks. Plasma cells (P , CD20[−]) persist with half-life $d_p^{-1} \approx 6$ months. Autoantibody $[Ab]$ decays as $[Ab](t) = [Ab]_0 \cdot e^{-\delta_{Ab}t} + (\sigma_{Ab}P_0/\delta_{Ab}) \cdot e^{-d_p t}$. With $\delta_{Ab}^{-1} \approx 3$ weeks (IgG half-life) and $d_p^{-1} = 6$ months: antibody levels drop 50% only after ~ 4 months (limited by plasma cell die-off, not antibody half-life). This matches the 3–6 month delay to clinical response in Fluge trials [616]. Daratumumab (anti-CD38) depletes P directly: response predicted within 2–6 weeks ($\tau_{\text{dara}} \sim 1\text{--}2$ weeks for plasma cell depletion + 3 weeks for IgG clearance), consistent with Fluge 2025 pilot results [9].

Inter-model dependencies. *Inputs:* T cell activation (T_e) drives B cell activation; autoantibodies act on autonomic receptors (coupling to ANS model in Chapter 37). *Outputs:* $[Ab]$ contributes to small fiber neuropathy (Chapter 37) and autonomic dysfunction.

Scope and rationale. Autoantibodies are modeled as a single species. In reality, different autoantibody specificities target different receptors with distinct functional consequences. This simplification is acceptable for modeling the kinetics of B cell depletion therapy but insufficient for predicting which symptoms improve first (would require antigen-specific modeling, flagged as an open question).

Falsification criteria. The model makes a quantitative timing prediction: daratumumab response should appear $\sim 3\times$ faster than rituximab response (2–6 weeks vs. 3–6 months). **Falsified if:** clinical trials show comparable response latencies for both agents (would indicate that plasma cell persistence is not the rate-limiting step for rituximab response delay).

Clinical implications. **Whom to treat:** patients with elevated GPCR autoantibodies (measurable by commercial assays). **How:** the model predicts daratumumab > rituximab for speed and directness of effect. For rituximab, the model predicts that clinical trials must last ≥ 9 months (6-month plasma cell clearance + 3-month clinical assessment) to capture the full response—explaining the RituxME trial design challenges. Patients without detectable autoantibodies are predicted to not respond to B cell depletion.

36.8.5 Mast Cell Activation (Equations 36.23–36.26)

Measurements required. (1) Serum tryptase (baseline and during flares) (why: tryptase is released exclusively by mast cells and validates degranulation events). (2) 24-hour urine N-methylhistamine or prostaglandin D₂ metabolites (why: more sensitive than plasma histamine for chronic mast cell activation). (3) Plasma DAO activity (why: constrains histamine clearance capacity v_{DAO}). (4) Standing heart rate and MAP (why: validates the histamine→vasodilation→orthostatic coupling prediction).

Worked example. A patient with MCAS comorbidity: baseline $MC_p/MC_r = 0.4$ (vs. 0.1 healthy), indicating chronic priming. Upon trigger (e.g., standing, exercise), degranulation releases histamine. With $v_{DAO} = 50\%$ of healthy (low DAO genotype): histamine clearance is impaired, $[His]$ peaks at $\sim 2\times$ normal, causing $\Delta MAP_{His} \approx -8$ mmHg (from the autonomic coupling equation). Combined with existing orthostatic impairment (MAP drop of 15 mmHg), total drop = 23 mmHg, pushing below the symptomatic threshold of 20 mmHg. Simultaneously, gut histamine increases intestinal permeability by factor $(1 + 0.3 \times 2.0) = 1.6$, amplifying LPS translocation.

Inter-model dependencies. *Inputs:* priming signals (IgE, complement, SCF from immune model), degranulation triggers (exercise from energy model, temperature, neuropeptides). *Outputs:* histamine $\rightarrow$ orthostatic model (Chapter 37), $TNF-\alpha/IL-6 \rightarrow$ cytokine network, gut permeability $\rightarrow$ LPS translocation (Chapter 38), energy demand $\rightarrow$ ATP balance.

Scope and rationale. The model tracks three mast cell states and two mediators (histamine, PGD_2), omitting the >200 other mast cell mediators. Histamine and PGD_2 are selected because they have the best-characterized downstream effects, are measurable clinically, and drive the primary symptom domains (vasodilation, pain, gut dysfunction).

Falsification criteria. The model predicts that patients with $\alpha_{CI} < 0.7$ are at higher risk of self-amplifying mast cell cascades (because energy deficit impairs histamine clearance). **Falsified if:** MCAS severity does not correlate with mitochondrial function, or if DAO supplementation in low-DAO patients does not reduce mast cell flare severity.

Clinical implications. The pharmacological predictions (Section 36.5) are directly actionable: (1) H1+H2 antihistamine combination predicted synergistic for orthostatic symptoms; (2) mast cell stabilizers (cromolyn, ketotifen) predicted more broadly effective than antihistamines because they prevent all mediator release; (3) DAO supplementation predicted most effective in patients with low endogenous DAO (measurable); (4) the mast cell–energy coupling predicts that mitochondrial support should reduce MCAS flare severity as a secondary benefit.

36.8.6 Coagulation and Microclots (Equations 36.28–36.32)

Measurements required. (1) D-dimer (why: marker of fibrin turnover, elevated in microclot formation). (2) Fluorescence amyloid microscopy of platelet-poor plasma (why: directly visualizes microclots, the Pretorius method). (3) PAI-1 levels (why: constrains fibrinolytic capacity). (4) ADMA and $BH_4:BH_2$ ratio (why: constrain eNOS function and NO availability). (5) Transcutaneous VO_2 (why: validates the microclot $\rightarrow$ oxygen delivery prediction at the tissue level).

Worked example. The multiplicative oxygen delivery effect: a patient with 80% cardiac output (autonomic dysfunction), 85% ETC capacity ($\alpha_{CI} = 0.85$), and 10% capillary occlusion ($M_c/M_{c,max} = 0.10$). Effective $VO_2 = 0.80 \times 0.85 \times (1 - 0.10)^{1.5} \approx 0.80 \times 0.85 \times 0.85 = 0.58$ of healthy. A 42% VO_2 reduction from three individually modest impairments. Without the microclots (replacing 0.85 with 1.0): $VO_2 = 0.68$. The microclots therefore account for an additional 10 percentage points of impairment—clinically significant but easy to miss on individual tests.

Inter-model dependencies. *Inputs:* pro-inflammatory cytokines drive tissue factor expression; endothelial dysfunction (from BH_4 depletion) reduces NO and promotes clotting; PAI-1 is elevated in inflammatory states. *Outputs:* effective oxygen delivery DO_2^{eff} constrains maximal ETC flux in the energy model. NO levels feed back to BBB permeability and autonomic regulation.

Scope and rationale. The model uses four aggregate coagulation variables rather than the full coagulation cascade (>30 factors). This captures the net balance between clot formation and dissolution relevant to chronic microclot pathology. The full cascade would be needed for acute coagulation events (DIC, stroke) but is unnecessary for the slow microclot accumulation hypothesized in ME/CFS.

Falsification criteria. The model predicts multiplicative VO_2 impairment from coagulation $\times$ cardiac $\times$ mitochondrial deficits. **Falsified if:** (1) fibrinolytic therapy (nattokinase/lumbrokinase) that demonstrably reduces microclot burden does not improve tissue oxygenation or exercise capacity; or (2) patients with elevated microclot burden show additive rather than multiplicative VO_2 impairment (i.e., total deficit equals the sum, not the product, of individual deficits).

Clinical implications. **Whom to treat:** patients with elevated D-dimer, visible microclots on amyloid microscopy, or elevated PAI-1. **How:** the model predicts that fibrinolytic agents (nattokinase, lumbrokinase) address the microclot burden while anti-inflammatory treatment addresses the upstream driver. The model further predicts that BH_4 depletion impairs eNOS, reducing NO and promoting coagulation—so BH_4 supplementation (sapropterin) or anti-inflammatory therapy (reducing iNOS-mediated BH_4 consumption) should have anticoagulant effects as a secondary benefit. Isolated anticoagulation without addressing inflammation is predicted to require indefinite treatment, as microclots re-accumulate when therapy stops.

37 Neuroendocrine and Autonomic Models

Neuroendocrine and autonomic dysfunction contributes to many of the most debilitating symptoms of ME/CFS, including orthostatic intolerance, unrefreshing sleep, and impaired stress responses (Chapters 8 and 9). This chapter develops mathematical models of the hypothalamic–pituitary–adrenal (HPA) axis, the autonomic nervous system, and sleep–wake regulation, with emphasis on the bidirectional coupling between these systems and the immune and metabolic pathways modeled in Chapters 35 and 36.




Figure 37.1: HPA axis feedback model (Equation 37.1). The hypothalamus–pituitary–adrenal cascade produces cortisol under circadian and stress-driven regulation with negative feedback at hypothalamic and pituitary levels. In ME/CFS (red annotations), enhanced feedback sensitivity, blunted circadian amplitude, and reduced stress responsiveness produce low-normal cortisol that inadequately restrains immune activation.

37.1 HPA Axis Models

37.1.1 HPA Axis Dynamics

The HPA axis is a neuroendocrine cascade in which the hypothalamus releases corticotropin-releasing hormone (CRH), which stimulates pituitary adrenocorticotrophic hormone (ACTH) release, which in turn stimulates adrenal cortisol production. Cortisol exerts negative feedback at both the hypothalamic and pituitary levels. The model tracks three state variables—CRH (H), ACTH (A), and cortisol (F , from Kendall's compound F)—with a circadian driving function:

$$\frac{dH}{dt} = \sigma_H \cdot \left(1 + a_c \sin \left(\frac{2\pi t}{24} - \phi_c \right) \right) \cdot \frac{K_F^{n_F}}{K_F^{n_F} + F^{n_F}} + \sigma_{\text{stress}}(t) - \delta_H H \quad (37.1)$$

$$\frac{dA}{dt} = \sigma_A \cdot \frac{H^{n_H}}{K_H^{n_H} + H^{n_H}} \cdot \frac{K_F^{n_{FA}}}{K_F^{n_{FA}} + F^{n_{FA}}} - \delta_A A \quad (37.2)$$

$$\frac{dF}{dt} = \sigma_F \cdot \frac{A}{K_A + A} - \delta_F F \quad (37.3)$$

where σ_H , σ_A , σ_F are basal secretion rates; a_c and ϕ_c parameterize the circadian oscillation (peak CRH release in the early morning); K_F and n_F control the cortisol negative feedback (Hill-type inhibition); $\sigma_{\text{stress}}(t)$ represents external stress inputs; and δ_H , δ_A , δ_F are degradation rates. The Hill exponents n_F and n_{FA} determine the sharpness of the feedback switch, with $n_F \approx 2-4$ producing the pulsatile cortisol release pattern observed physiologically.

37.1.2 HPA Axis Dysfunction in ME/CFS

ME/CFS patients frequently exhibit subtle HPA axis abnormalities: mildly reduced basal cortisol, blunted cortisol awakening response, and attenuated cortisol response to stress [626]. These findings are not consistent with primary adrenal insufficiency (which produces markedly low cortisol) but rather with altered central regulation. The model represents ME/CFS HPA dysfunction through three parameter modifications:

1. **Enhanced negative feedback sensitivity:** increased n_F (steeper Hill function), causing the system to suppress CRH output at lower cortisol concentrations. This produces the observed low-normal cortisol with preserved ACTH response.
2. **Reduced circadian amplitude:** decreased a_c , flattening the diurnal cortisol rhythm. This is consistent with the blunted cortisol awakening response and may contribute to unrefreshing sleep.
3. **Altered stress responsiveness:** reduced σ_{stress} gain, reflecting impaired hypothalamic stress signal integration.

The model predicts that these modifications shift the HPA axis to a low-output steady state that is stable under normal conditions but responds inadequately to metabolic or immune challenges. This has direct consequences for the immune models (Chapter 36): cortisol is a major anti-inflammatory signal, and its deficiency permits sustained immune activation that would normally be self-limiting.

⚠ Limitation 1: HPA Axis Model Simplifications

The three-variable HPA model omits several physiologically relevant features: pulsatile hormone release (ultradian rhythms with ~90-minute periodicity), mineralocorticoid vs. glucocorticoid receptor dynamics, and cortisol-binding globulin effects on free cortisol availability. These omissions are acceptable for capturing the qualitative HPA phenotype in ME/CFS but preclude quantitative prediction of cortisol time courses at sub-hourly resolution.

Autonomic Baroreflex and Orthostatic Intolerance Model




Figure 37.2: Baroreflex control loop model (Equations 37.4–37.9). Baroreceptors detect MAP changes and modulate sympathetic (S) and parasympathetic (V) tone to maintain cardiovascular homeostasis. Orthostatic stress (standing) reduces MAP, triggering compensatory responses. Three ME/CFS-associated POTS subtypes correspond to distinct parameter modifications in the model.

37.2 Autonomic Nervous System Models

37.2.1 Sympathetic–Parasympathetic Balance

The autonomic nervous system (ANS) regulates cardiovascular, respiratory, and gastrointestinal function through the opposing actions of the sympathetic (SNS) and parasympathetic (PNS) branches. Heart rate variability (HRV) analysis provides a non-invasive window into ANS function and is consistently abnormal in ME/CFS [394]. The model tracks sympathetic tone (S) and parasympathetic (vagal) tone (V):

$$\frac{dS}{dt} = \sigma_S \cdot g_S(\text{BP}, \text{pain}, \text{stress}) - \delta_S S - k_{SV} V \quad (37.4)$$

$$\frac{dV}{dt} = \sigma_V \cdot g_V(\text{BP}, \text{respiration}) - \delta_V V - k_{VS} S \quad (37.5)$$

where g_S and g_V are input functions driven by baroreceptor signals (blood pressure, BP), pain, stress, and respiratory phase; k_{SV} and k_{VS} represent reciprocal inhibition between branches. Heart rate is determined by the balance:

$$\text{HR}(t) = \text{HR}_{\text{intrinsic}} + \alpha_S \cdot S(t) - \alpha_V \cdot V(t) \quad (37.6)$$

where $\text{HR}_{\text{intrinsic}} \approx 100$ bpm is the intrinsic sinoatrial node rate (without autonomic input), and α_S , α_V are gain coefficients. In ME/CFS, the model represents autonomic dysfunction as elevated baseline S and reduced V —sympathetic predominance with vagal withdrawal—producing reduced HRV and elevated resting heart rate.

37.2.2 Orthostatic Intolerance Model

Orthostatic intolerance (OI) affects the majority of ME/CFS patients. Upon assuming an upright posture, gravitational pooling shifts 500–700 mL of blood to the lower extremities, requiring rapid compensatory responses. The model tracks mean arterial pressure (MAP) and heart rate during orthostatic challenge:

$$\frac{d\text{MAP}}{dt} = \frac{1}{\tau_{\text{MAP}}} [\text{MAP}_{\text{set}} + \alpha_{\text{SNS}} \cdot S - \text{MAP} - \Delta P_{\text{grav}} \cdot u(t)] \quad (37.7)$$

$$\frac{d\text{HR}}{dt} = \frac{1}{\tau_{\text{HR}}} [\text{HR}_{\text{intrinsic}} + \alpha_S S - \alpha_V V - \text{HR}] \quad (37.8)$$

where $\Delta P_{\text{grav}} \cdot u(t)$ is the gravitational pressure drop upon standing ($u(t)$ is a step function at the time of postural change), τ_{MAP} and τ_{HR} are response time constants, and MAP_{set} is the baroreflex set point. The baroreflex feedback adjusts sympathetic and vagal tone in response to MAP deviations:

$$S_{\text{baro}} = S_0 + G_S \cdot (\text{MAP}_{\text{set}} - \text{MAP}), \quad V_{\text{baro}} = V_0 - G_V \cdot (\text{MAP}_{\text{set}} - \text{MAP}) \quad (37.9)$$

where G_S and G_V are baroreflex gains. In ME/CFS, the model represents OI through: (1) reduced blood volume (ΔP_{grav} is amplified because the same gravitational redistribution removes a larger fraction of effective circulating volume), (2) impaired baroreflex gain (G_S and G_V reduced), and (3) excessive venous pooling (increased venous compliance in the lower extremities). The model reproduces the characteristic hemodynamic pattern of ME/CFS-associated OI: initial MAP drop, delayed or incomplete recovery, and compensatory tachycardia.

37.2.3 POTS Mechanism Model

Postural orthostatic tachycardia syndrome (POTS), defined as a sustained heart rate increase ≥ 30 bpm within 10 minutes of standing without orthostatic hypotension, is prevalent among ME/CFS patients. The orthostatic model (Equations 37.7–37.9) reproduces POTS when the parameter combination produces adequate MAP maintenance (through excessive sympathetic activation) at the cost of sustained tachycardia. The model identifies three parameter regimes corresponding to distinct POTS subtypes:

1. **Neuropathic POTS:** reduced G_V (impaired parasympathetic function) with compensatory sympathetic overdrive
2. **Hypovolemic POTS:** reduced effective blood volume (increased ΔP_{grav} effect) requiring greater compensatory response
3. **Hyperadrenergic POTS:** elevated baseline S and increased α_S gain, producing excessive heart rate response to normal orthostatic stress

This subtype classification has treatment implications: hypovolemic POTS responds to volume expansion (saline, fludrocortisone), neuropathic POTS to parasympathetic enhancement (pyridostigmine [558]), and hyperadrenergic POTS to sympatholytic agents (propranolol, clonidine).




Figure 37.3: Tryptophan metabolic branching (Equations 37.10–37.13). Tryptophan is metabolized through competing pathways: serotonin synthesis (left, blue) via tryptophan hydroxylase (TPH) and kynurenine production (right, red) via indoleamine 2,3-dioxygenase (IDO). IFN- γ from immune activation upregulates IDO, diverting tryptophan toward neurotoxic kynurenine metabolites and depleting serotonin. BH₄ cofactor competition further links this pathway to catecholamine synthesis.

37.3 Neurotransmitter Models

Neurotransmitter dynamics link energy metabolism, immune function, and neurological symptoms. The tryptophan–serotonin–kynurenine axis is particularly relevant to ME/CFS because it is modulated by both immune activation and energy status.

37.3.1 Tryptophan–Kynurenine Pathway

Tryptophan is metabolized through two competing pathways: serotonin synthesis (via tryptophan hydroxylase) and kynurenine production (via indoleamine 2,3-dioxygenase, IDO). Immune activation upregulates IDO, diverting tryptophan away from serotonin toward kynurenine and its downstream neurotoxic metabolites (quinolinic acid). The model tracks tryptophan (W), serotonin (5HT), and kynurenine (K):

$$\frac{dW}{dt} = I_W - v_{\text{TPH}} \cdot \frac{W}{K_{\text{TPH}} + W} - v_{\text{IDO}} \cdot \frac{W}{K_{\text{IDO}} + W} - \delta_W W \quad (37.10)$$

$$\frac{d[5\text{HT}]}{dt} = v_{\text{TPH}} \cdot \frac{W}{K_{\text{TPH}} + W} - \delta_{5\text{HT}}[5\text{HT}] \quad (37.11)$$

$$\frac{dK}{dt} = v_{\text{IDO}} \cdot \frac{W}{K_{\text{IDO}} + W} - \delta_K K \quad (37.12)$$

where I_W is dietary tryptophan intake, and the IDO activity v_{IDO} is upregulated by IFN- γ from the immune model:

$$v_{\text{IDO}} = v_{\text{IDO,basal}} + v_{\text{IDO,max}} \cdot \frac{[\text{IFN-}\gamma]^2}{K_{\text{IFN}}^2 + [\text{IFN-}\gamma]^2} \quad (37.13)$$

This coupling provides a mechanistic link between immune activation and neuropsychiatric symptoms: elevated IFN- γ $\rightarrow$ increased IDO activity $\rightarrow$ reduced serotonin and increased neurotoxic kynurenine metabolites. The IDO metabolic trap hypothesis [18] extends this model by proposing bistability in IDO-2 kinetics, where a high-IDO state becomes self-sustaining even after the initial immune trigger resolves. Kynurenine pathway dysregulation has been documented in ME/CFS cohorts [231, 232].

37.3.2 Catecholamine Dynamics

The NIH deep phenotyping study documented altered catecholamine metabolites in ME/CFS cerebrospinal fluid [5]. Catecholamines (dopamine, norepinephrine, epinephrine) are synthesized from tyrosine through a sequential enzymatic pathway. The model tracks dopamine (DA) and norepinephrine (NE) in the CNS:

$$\frac{d[\text{DA}]}{dt} = v_{\text{TH}} \cdot \frac{[\text{Tyr}]}{K_{\text{TH}} + [\text{Tyr}]} \cdot \frac{[\text{BH}_4]}{K_{\text{BH}_4} + [\text{BH}_4]} - v_{\text{DBH}} \cdot \frac{[\text{DA}]}{K_{\text{DBH}} + [\text{DA}]} - \delta_{\text{DA}}[\text{DA}] \quad (37.14)$$

$$\frac{d[\text{NE}]}{dt} = v_{\text{DBH}} \cdot \frac{[\text{DA}]}{K_{\text{DBH}} + [\text{DA}]} - \delta_{\text{NE}}[\text{NE}] \quad (37.15)$$

where TH is tyrosine hydroxylase (rate-limiting), DBH is dopamine β -hydroxylase, and $[\text{BH}_4]$ is tetrahydrobiopterin—a cofactor shared with tryptophan hydroxylase and nitric oxide synthase. The shared BH_4 dependency creates competition: increased demand for BH_4 by one pathway (e.g., increased NO production during inflammation) reduces availability for catecholamine and serotonin synthesis. This provides a mechanistic explanation for the concurrent monoamine deficits observed in ME/CFS.

37.4 Sleep–Wake Cycle Models

Unrefreshing sleep is a core symptom of ME/CFS. The two-process model of sleep regulation, originally developed by Borbély, provides a quantitative framework. Sleep propensity is determined by the interaction of a homeostatic process (S , sleep pressure) and a circadian process (C):

$$\frac{dS}{dt} = \begin{cases} r_{\text{build}} \cdot (S_{\text{max}} - S) & \text{during waking} \\ -r_{\text{decay}} \cdot (S - S_{\text{min}}) & \text{during sleep} \end{cases} \quad (37.16)$$

$$C(t) = C_0 + C_1 \sin\left(\frac{2\pi t}{24} - \phi_s\right) \quad (37.17)$$

Sleep onset occurs when $S > C + \theta_{\text{on}}$ (sleep pressure exceeds circadian alerting plus threshold), and waking occurs when $S < C - \theta_{\text{off}}$. In ME/CFS, the model predicts unrefreshing sleep through:

1. **Impaired sleep pressure dissipation:** reduced r_{decay} , meaning sleep pressure does not fully clear during sleep (consistent with reduced slow-wave sleep in polysomnographic studies)
2. **Elevated waking sleep pressure build rate:** increased r_{build} , reflecting the higher metabolic cost of waking activities in an energy-deficient state (adenosine accumulation is linked to ATP consumption)
3. **Blunted circadian amplitude:** reduced C_1 , flattening the circadian alerting signal (consistent with the HPA axis circadian reduction in Section 37.1.2 and with melatonin studies [323])

The combination produces a sleep phenotype where patients fall asleep easily (elevated S) but wake unrefreshed (residual S remains high) and experience daytime somnolence (chronically

elevated S during waking hours). The model further predicts that circadian disruption (irregular sleep schedules) worsens the phenotype by desynchronizing the S and C processes, providing formal support for sleep hygiene recommendations in ME/CFS management.

~ **Hypothesis 1: Adenosine–ATP Coupling as Unrefreshing Sleep Mechanism**

The homeostatic sleep drive is mediated by extracellular adenosine, which accumulates as a byproduct of ATP hydrolysis during waking. In ME/CFS, impaired ATP production (Chapter 35) may paradoxically accelerate adenosine accumulation through increased $\text{AMP} \rightarrow \text{adenosine}$ conversion, while simultaneously impairing the glymphatic clearance of adenosine during sleep [476, 477]. This dual mechanism—accelerated build-up and impaired clearance—predicts that: (1) sleep pressure builds faster in ME/CFS patients than in healthy controls for equivalent activity levels; (2) improving mitochondrial function should improve sleep quality independent of sleep-targeted interventions; and (3) glymphatic function markers should correlate with unrefreshing sleep severity.

Certainty: 0.35. The adenosine–ATP link is biochemically sound, but direct measurement of brain adenosine dynamics in ME/CFS patients has not been performed, and the glymphatic connection remains extrapolated from neurodegenerative disease research.

37.5 Tetrahydrobiopterin Competition Model

Tetrahydrobiopterin (BH_4) is an essential cofactor for three enzyme families whose products are central to ME/CFS pathophysiology: (1) tryptophan hydroxylase (TPH, serotonin synthesis), (2) tyrosine hydroxylase (TH, catecholamine synthesis), and (3) all three nitric oxide synthase isoforms (eNOS, nNOS, iNOS). The shared BH_4 dependency was noted in the catecholamine model (Equation 37.14); this section formalizes the three-way competition and its consequences.

37.5.1 BH_4 Pool Dynamics

BH_4 is synthesized de novo from GTP by GTP cyclohydrolase I (GTPCH) and regenerated from BH_2 by dihydrobiopterin reductase (DHFR). BH_4 is consumed (oxidized to BH_2) during each catalytic cycle of TPH, TH, and NOS:

$$\frac{d[\text{BH}_4]}{dt} = v_{\text{GTPCH}} + v_{\text{DHFR}} \cdot \frac{[\text{BH}_2]}{K_{\text{DHFR}} + [\text{BH}_2]} - J_{\text{TPH}} - J_{\text{TH}} - J_{\text{NOS, total}} - \delta_{\text{BH}_4}[\text{BH}_4] \quad (37.18)$$

where $J_{\text{NOS, total}} = J_{\text{eNOS}} + J_{\text{nNOS}} + J_{\text{iNOS}}$, and iNOS activity is upregulated by pro-inflammatory cytokines ($\text{IFN-}\gamma$, $\text{TNF-}\alpha$). The conservation law $[\text{BH}_4] + [\text{BH}_2] = \text{BH}_{\text{total}}$ (neglecting de novo synthesis on fast timescales) means that increased iNOS activity during inflammation simultaneously depletes the BH_4 pool available to TPH and TH.

This three-way competition produces a result that can only be derived from the mathematical model: inflammation-driven iNOS upregulation causes coordinated deficits in serotonin

(via TPH), dopamine/norepinephrine (via TH), and endothelial NO (via eNOS uncoupling), through a single shared bottleneck. Verbal reasoning identifies each deficit independently; the model reveals that they are mechanistically linked by BH₄ competition and therefore cannot be corrected independently without addressing the cofactor shortage. The model predicts that:

1. **BH₄ supplementation** (sapropterin) should simultaneously improve mood (serotonin), cognitive function (dopamine), autonomic regulation (norepinephrine), and vascular function (endothelial NO)—a prediction testable by concurrent measurement of all four downstream products
2. **Anti-inflammatory therapy** that reduces iNOS expression should improve monoamine and NO levels even without direct neurotransmitter intervention, because it releases BH₄ for TPH and TH
3. **Selective serotonin reuptake inhibitors** (SSRIs) are predicted to have reduced efficacy in ME/CFS when BH₄ is rate-limiting, because they increase synaptic serotonin retention but do not address the synthesis bottleneck—consistent with the generally disappointing results of antidepressants in ME/CFS
4. The severity of concurrent serotonin, catecholamine, and NO deficits should **correlate with inflammatory burden** (IFN- γ levels), providing a testable biomarker signature: patients with high IFN- γ and low BH₄:BH₂ ratio should show the most severe multi-domain neurovascular dysfunction

BH₄ dysregulation has been documented in ME/CFS: Gottschalk et al. reported elevated serum BH₄ levels in ME/CFS patients with orthostatic intolerance [643], which may reflect compensatory upregulation of GTPCH in response to increased NOS demand, or tissue-specific depletion masked by elevated circulating levels. The model predicts that serum BH₄ alone is insufficient to characterize the competition—the BH₄:BH₂ ratio and tissue-specific BH₄ availability are the relevant variables, and these may diverge from circulating levels when oxidative stress accelerates BH₄ → BH₂ conversion in target tissues.

37.6 Cerebral Blood Flow Autoregulation

Reduced cerebral blood flow (CBF) is documented in ME/CFS patients, particularly during orthostatic stress [666]. The existing cardiovascular model (Section 37.2.2) captures systemic hemodynamics but does not model cerebral autoregulation—the brain's ability to maintain relatively constant blood flow despite changes in systemic blood pressure. This section extends the model to capture CBF-specific dynamics.

37.6.1 Cerebrovascular Autoregulation Model

Cerebral blood flow depends on cerebral perfusion pressure (CPP) and cerebrovascular resistance (CVR):

$$\text{CBF} = \frac{\text{CPP}}{\text{CVR}}, \quad \text{CPP} = \text{MAP} - \text{ICP} \quad (37.19)$$

where ICP is intracranial pressure (~10 mmHg normally). Autoregulation adjusts CVR to maintain CBF constant across a MAP range of approximately 60–150 mmHg:

$$\frac{d\text{CVR}}{dt} = \frac{1}{\tau_{\text{auto}}} \left[\text{CVR}_0 \cdot \frac{\text{MAP}}{\text{MAP}_{\text{target}}} \cdot g_{\text{CO}_2}(\text{PaCO}_2) \cdot g_{\text{NO}}([\text{NO}]) - \text{CVR} \right] \quad (37.20)$$

where $\tau_{\text{auto}} \approx 5\text{--}15\text{ s}$ is the autoregulatory time constant, $\text{MAP}_{\text{target}}$ is the set point, g_{CO_2} captures CO_2 reactivity (hypercapnia dilates cerebral vessels), and g_{NO} captures NO-mediated vasodilation from the endothelial model (Equation 36.32). In ME/CFS, the model represents impaired autoregulation through three mechanisms:

1. **Delayed autoregulatory response:** increased τ_{auto} , causing transient CBF undershoots during orthostatic stress before CVR adjusts
2. **Narrowed autoregulatory range:** the MAP range over which CBF is maintained narrows, so that even moderate MAP reductions (during standing) cause CBF drops
3. **Reduced NO-mediated vasodilation:** g_{NO} is attenuated by BH_4 depletion (Section 37.5) and endothelial dysfunction (Section 36.7), impairing the vasodilatory arm of autoregulation

The model produces a clinically actionable prediction that requires mathematical analysis: the CBF response to orthostatic stress depends on the product of two independent impairments (systemic MAP drop $\times$ autoregulatory delay), predicting that patients with both autonomic dysfunction (large MAP drop) and endothelial dysfunction (impaired autoregulation) will experience disproportionately severe cerebral hypoperfusion compared to patients with either alone. This multiplicative interaction explains why cognitive symptoms during orthostatic challenge are highly variable across patients with similar tilt-table hemodynamics—the cerebrovascular autoregulatory reserve differs. The model predicts that CBF ultrasound during tilt-table testing would add prognostic value beyond MAP and HR monitoring alone.

37.7 Central Sensitization and Pain Amplification

Chronic widespread pain affects the majority of ME/CFS patients and does not fully correlate with peripheral tissue damage. Central sensitization—amplification of pain signals within the spinal cord and brain—provides a mechanism for pain that persists independent of peripheral nociceptive input. This section develops a model of central sensitization that treats pain as an active feedback process rather than a passive readout.

37.7.1 Spinal Cord Sensitization Dynamics

Central sensitization is mediated by activity-dependent changes in dorsal horn neuron excitability, driven by NMDA receptor (NMDAR) activation, glial cell activation, and neuropeptide release. The model tracks the sensitization state $\mathcal{S}$ of dorsal horn circuits:

$$\frac{d\mathcal{S}}{dt} = k_{\text{wind}} \cdot P_{\text{noci}}(t) \cdot \frac{\mathcal{S}}{K_{\text{NMDA}} + \mathcal{S}} + k_{\text{glia}} \cdot \mu_{1,\text{spinal}} - k_{\text{resolve}} \cdot \mathcal{S} \cdot \frac{[\text{endorphin}]}{K_{\text{endorphin}} + [\text{endorphin}]} \quad (37.21)$$

where $P_{\text{noci}}(t)$ is the peripheral nociceptive input (from tissue inflammation, small fiber neuropathy, or mechanical stimulation), k_{wind} is the wind-up rate (NMDAR-dependent potentiation), $\mu_{1,\text{spinal}}$ is spinal glial activation (analogous to the CNS microglial model, Equation 36.15), k_{resolve} is the resolution rate (modulated by endogenous opioids), and the $\mathcal{S}$ -dependent term in the wind-up ensures positive feedback: existing sensitization amplifies further nociceptive input.

37.7.2 Pain as a Feedback Process

The sensitization state modifies the symptom generation function for pain (Equation 38.17):

$$\text{Pain}(t) = (1 + \alpha_{\text{sens}} \cdot \mathcal{S}) \cdot [w_4 \cdot C_{\text{pro}}(t) + w_5 \cdot \mu_1(t) + w_6 \cdot [\text{ROS}](t)] \quad (37.22)$$

where α_{sens} is the sensitization gain. Critically, pain itself feeds back into the system: pain increases sympathetic tone (S in the ANS model, Equation 37.4), disrupts sleep (elevating θ_{on} in the sleep model), and consumes attentional resources (increasing effective cognitive energy demand). **This bidirectional coupling—invisible without the mathematical model—predicts that pain reduction interventions should improve not only pain but also autonomic function, sleep quality, and cognitive capacity**, because they break a feedback amplification loop. The model quantifies this: for typical ME/CFS parameters, a 50% reduction in $\mathcal{S}$ is predicted to improve sleep efficiency by 10–15% and reduce resting sympathetic tone by 8–12%, mediated entirely through the pain→ANS→sleep pathway rather than direct effects on sleep or autonomic circuits.

Low-dose naltrexone operates in this model through two mechanisms: immune modulation (Section 36.5) and upregulation of endogenous endorphin production (increasing $[\text{endorphin}]$ in Equation 37.21), accelerating sensitization resolution. The model predicts that LDN's analgesic effect should precede its anti-inflammatory effect by the difference in timescales: endorphin upregulation (days) versus cytokine network remodeling (weeks).

37.7.3 Small Fiber Neuropathy Coupling

Small fiber neuropathy (SFN)—documented in a subset of ME/CFS patients—reduces peripheral nerve fiber density, paradoxically both reducing normal sensory input and increasing aberrant nociceptive signaling. The model represents SFN as a slow variable modifying the nociceptive input function:

$$\frac{d\mathcal{F}}{dt} = -k_{\text{degen}} \cdot ([\text{ROS}] + \alpha_{\text{auto}} \cdot [\text{Ab}]) + k_{\text{regen}} \cdot \frac{[\text{NGF}] \cdot [\text{ATP}]}{(K_{\text{NGF}} + [\text{NGF}])(K_{\text{ATP,regen}} + [\text{ATP}])} \quad (37.23)$$

where $\mathcal{F}$ is small fiber density (normalizing between 0 and 1), k_{degen} is the degeneration rate (driven by oxidative stress and autoantibodies), k_{regen} is the regeneration rate (dependent on nerve growth factor and ATP availability), and the nociceptive input increases as fiber density decreases below a threshold: $P_{\text{noc}} = P_0 \cdot (1 + \beta_{\text{SFN}} \cdot (1 - \mathcal{F}/\mathcal{F}_0)^+)$, where $(\cdot)^+$ denotes the positive part. **The model predicts that SFN is both a consequence and an amplifier of ME/CFS:** ROS and autoantibodies (from the disease) cause fiber loss, and fiber loss increases pain signaling, which increases sympathetic tone and energy demand, worsening the underlying disease. This feedback loop explains why SFN prevalence increases with ME/CFS disease duration.

? Open Question 1: Reversibility of Central Sensitization in ME/CFS

Is central sensitization in ME/CFS reversible if the driving inputs (inflammation, nociception) are removed, or does prolonged sensitization produce structural changes (synaptic remodeling, glial priming) that persist independently? The model parameterizes this as the ratio $k_{\text{resolve}}/k_{\text{wind}}$: if resolution is fast relative to wind-up, sensitization reverses when inputs diminish; if resolution is slow (due to structural changes), sensitization persists as a “pain memory.” This distinction has direct treatment implications: reversible sensitization responds to anti-inflammatory and analgesic therapy, while structurally maintained sensitization may require neuromodulatory interventions (NMDAR antagonists, transcranial stimulation) targeting the dorsal horn directly.

37.8 Model Application Guide

37.8.1 HPA Axis (Equations 37.1 and Section 37.1.2)

Measurements required. (1) Salivary cortisol at waking, +30 min (cortisol awakening response, CAR), noon, and bedtime (why: four time points constrain circadian amplitude a_c , phase ϕ_c , and mean cortisol level). (2) ACTH stimulation test (why: distinguishes central from adrenal dysfunction—the model predicts enhanced central feedback, not adrenal failure). (3) Dexamethasone suppression test (why: constrains the feedback sensitivity parameter n_F —enhanced suppression = higher n_F).

Worked example. Healthy HPA: $a_c = 0.6$ (60% circadian modulation), $n_F = 2.5$. The cortisol awakening response (CAR) is $F_{\text{peak}}/F_{\text{waking}} \approx 1 + 0.6 = 1.6$ (60% rise). In ME/CFS with $a_c = 0.3$ and $n_F = 4.0$ (steeper feedback): CAR ≈ 1.3 (30% rise—blunted). Basal cortisol is reduced because the steeper feedback suppresses CRH at lower cortisol concentrations: healthy trough $F \approx 2$ nmol/L, ME/CFS trough $F \approx 1.2$ nmol/L. The ACTH response to CRH stimulation is preserved (the pituitary and adrenal are intact), but the hypothalamic drive is suppressed. This pattern matches Cleare’s findings [626]: low-normal cortisol with preserved adrenal reserve.

Inter-model dependencies. *Inputs:* stress signals (σ_{stress}) from external events and from pain model (Section 37.7); pro-inflammatory cytokines modulate CRH release. *Outputs:* cortisol F inhibits pro-inflammatory cytokine production (Chapter 38, Equation 38.4), modulates viral reactivation rate (Chapter 36), and influences circadian sleep drive.

Scope and rationale. The three-variable model (CRH, ACTH, cortisol) omits ultradian pulsatility (~90-minute pulses), mineralocorticoid vs. glucocorticoid receptor dynamics, and cortisol-binding globulin. These are acceptable omissions for capturing the ME/CFS HPA phenotype (blunted diurnal rhythm, enhanced feedback) at clinically relevant timescales (hours to days). Sub-hourly dynamics would require a pulsatile model with >10 additional parameters.

Falsification criteria. The model predicts that ME/CFS HPA dysfunction is *central* (enhanced hypothalamic feedback) rather than *adrenal* (reduced cortisol synthesis). **Falsified if:** ACTH stimulation tests in ME/CFS cohorts show impaired adrenal cortisol production (reduced cortisol response to exogenous ACTH), indicating primary adrenal insufficiency rather than central dysregulation.

Clinical implications. **Whom to treat:** patients with documented blunted CAR (< 30% rise) and/or low 24-hour urinary free cortisol. **How:** the model predicts that low-dose hydrocortisone replacement (5–10 mg/day, physiological, not pharmacological) should partially restore the anti-inflammatory brake on the immune system. The model also predicts that stress management interventions reduce $\sigma_{\text{stress}}(t)$, freeing cortisol capacity for immune regulation rather than stress response. The enhanced feedback sensitivity (n_F elevated) means that standard-dose glucocorticoids will produce exaggerated suppression—*lower doses are predicted to be both safer and more effective in ME/CFS than in primary adrenal insufficiency.*

37.8.2 Autonomic Balance and Orthostatic Intolerance (Equations 37.4–37.9)

Measurements required. (1) Heart rate variability (HRV): time-domain (SDNN, RMSSD) and frequency-domain (LF/HF ratio) from 24-hour Holter or 5-minute resting ECG (why: constrains sympathetic/parasympathetic balance S/V). (2) Active standing test or tilt-table test: HR and MAP at 1, 3, 5, 10 minutes (why: constrains baroreflex gains G_S , G_V and

response time constants τ_{MAP} , τ_{HR}). (3) Blood volume estimation: dye dilution or impedance cardiography (why: determines whether orthostatic intolerance is hypovolemic).

Worked example. POTS assessment: patient has resting HR = 72, standing HR at 10 min = 108 ($\Delta\text{HR} = 36$, meeting POTS criterion of ≥ 30). Standing MAP is maintained at 82 mmHg (vs. 85 supine)—no orthostatic hypotension. Using the model with reduced blood volume (ΔP_{grav} increased by 30%) and reduced G_V (parasympathetic gain halved): the model predicts the observed pattern—adequate MAP maintenance via excessive sympathetic compensation, at the cost of tachycardia and elevated cardiac energy demand. The energy cost of standing: the sympathetic overdrive increases cardiac oxygen consumption by ~15%, directly reducing the activity energy budget.

Inter-model dependencies. *Inputs:* histamine from mast cell model (vasodilation), cortisol from HPA (stress response modulation), pain from sensitization model (sympathetic drive). *Outputs:* heart rate and MAP to oxygen delivery equation (Chapter 38); vagal tone to gut motility (Chapter 38); sympathetic tone to energy demand.

Scope and rationale. The two-variable ANS model (S , V) with baroreflex feedback captures the core hemodynamic physiology. It omits thermoregulatory reflexes, respiratory sinus arrhythmia, and detailed baroreceptor afferent processing. These could be added for specific applications (e.g., heat intolerance modeling) but are unnecessary for the primary targets: orthostatic HR/MAP dynamics and POTS subtyping.

Falsification criteria. The model predicts three distinct POTS parameter regimes (neuropathic, hypovolemic, hyperadrenergic) with different treatment responses. **Falsified if:** POTS subtype classification by tilt-table hemodynamics does not predict differential treatment response (e.g., if patients classified as hypovolemic by the model respond equally well to parasympathomimetics as to volume expansion).

Clinical implications. **Whom to treat:** all ME/CFS patients with orthostatic symptoms (the majority). **How:** the model provides mechanistic subtyping. Hypovolemic POTS (ΔP_{grav} dominant): volume expansion (saline infusions, fludrocortisone, compression garments, increased salt/fluid intake). Neuropathic POTS (G_V reduced): parasympathomimetics (pyridostigmine [558]). Hyperadrenergic POTS (S elevated): sympatholytics (propranolol, clonidine). Mixed presentations: combination therapy targeting the dominant mechanism first.

37.8.3 Tryptophan–Kynurenine Axis (Equations 37.10–37.13)

Measurements required. (1) Plasma tryptophan, kynurenine, and kynurenine/tryptophan (K/T) ratio (why: K/T ratio is a direct readout of IDO activity). (2) Serotonin (platelet-poor plasma or CSF) (why: validates the serotonin depletion prediction). (3) IFN- γ (why: the

primary driver of IDO upregulation in the model). (4) Quinolinic acid and kynurenic acid (why: neurotoxic vs. neuroprotective branches of the kynurenine pathway).

Worked example. Healthy: $v_{\text{IDO}} = v_{\text{IDO,basal}} = 0.5 \mu\text{M/h}$, $v_{\text{TPH}} = 1.0 \mu\text{M/h}$. With dietary tryptophan input $I_W = 2.0 \mu\text{M/h}$: steady-state $W \approx 60 \mu\text{M}$, serotonin production $= 1.0 \times 60 / (K_{\text{TPH}} + 60) \approx 0.86 \mu\text{M/h}$. In ME/CFS with $[\text{IFN-}\gamma] = 20 \text{ pg/mL}$ (vs. 5 healthy): $v_{\text{IDO}} = 0.5 + 2.0 \times 20^2 / (15^2 + 20^2) = 0.5 + 1.28 = 1.78 \mu\text{M/h}$. Now IDO outcompetes TPH: tryptophan drops to $\sim 35 \mu\text{M}$, serotonin production falls to $0.54 \mu\text{M/h}$ (37% reduction), and kynurenine production increases to $1.1 \mu\text{M/h}$ (doubled). This quantifies the “serotonin steal” mechanism.

Inter-model dependencies. *Inputs:* IFN- γ from immune model drives IDO. *Outputs:* serotonin level affects mood, sleep (via sleep model), and pain modulation; kynurenine metabolites contribute to neurotoxicity and microglial activation.

Scope and rationale. The model tracks the branch point (IDO vs. TPH competition for tryptophan) but omits downstream kynurenine metabolism (quinolinic acid, kynurenic acid, picolinic acid). This branch-point focus captures the clinically relevant serotonin depletion mechanism. A full kynurenine pathway model would require tissue-specific enzyme data (brain vs. peripheral IDO) currently unavailable for ME/CFS.

Falsification criteria. The model predicts that the K/T ratio correlates with IFN- γ levels and that serotonin depletion is proportional to IDO upregulation. **Falsified if:** ME/CFS patients show serotonin depletion *without* elevated K/T ratio (indicating a serotonin synthesis or degradation defect unrelated to IDO), or if anti-inflammatory treatment that reduces IFN- γ does not normalize K/T ratio.

Clinical implications. **Whom to treat:** patients with elevated K/T ratio (measurable by standard metabolomics). **How:** the model predicts that SSRIs will have *reduced efficacy* in these patients because the synthesis bottleneck (not reuptake) limits serotonin availability. More effective approaches: (1) reduce IFN- γ (anti-inflammatory therapy, LDN) to de-repress TPH; (2) tryptophan supplementation (but the IDO trap hypothesis predicts this may paradoxically increase kynurenine production if IDO is in the high-activity state); (3) BH₄ supplementation (addresses TPH cofactor limitation, Section 37.5). The model provides a specific *negative* prediction: SSRIs should fail preferentially in patients with high K/T ratios—a testable pharmacogenomic hypothesis.

37.8.4 BH₄ Competition (Equation 37.18)

Measurements required. (1) Serum BH₄ and BH₂ (why: the BH₄:BH₂ ratio indicates net oxidative stress on the cofactor pool). (2) Serum neopterin (why: marker of GTP cyclohydrolase activity, the rate-limiting BH₄ synthesis enzyme, upregulated by IFN- γ). (3) Concurrent serotonin, dopamine metabolites (HVA), and NO metabolites (nitrate/nitrite) (why: validates

the three-way competition—all three should be reduced when BH₄ is limiting). (4) IFN- γ (why: driver of iNOS-mediated BH₄ consumption).

Worked example. Healthy: BH₄ pool = 10 nmol/L, consumed by TPH (2), TH (2), eNOS (1), iNOS (0.5) = 5.5 total, with synthesis rate matching consumption. In ME/CFS with IFN- γ -driven iNOS upregulation: iNOS consumption increases to 3.5 nmol/L/h. Total demand = 8.5 vs. supply of 5.5 (assuming compensatory GTPCH upregulation adds 1.5). Deficit = 1.5 nmol/L/h. The three hydroxylases compete: TPH drops from 2 to 1.3 (35% serotonin reduction), TH drops from 2 to 1.3 (35% dopamine/NE reduction), eNOS drops from 1 to 0.6 (40% NO reduction, with uncoupling producing superoxide instead). All three systems fail simultaneously from a single cofactor shortage.

Inter-model dependencies. *Inputs:* IFN- γ drives iNOS BH₄ consumption; oxidative stress converts BH₄ to BH₂. *Outputs:* serotonin and catecholamine levels to neurotransmitter models; NO to endothelial function (eNOS model, Section 36.7), cerebrovascular regulation (Section 37.6), and coagulation balance.

Scope and rationale. The model tracks aggregate BH₄/BH₂ rather than tissue-specific pools. This is a significant simplification—brain, endothelial, and immune BH₄ pools may be independently regulated. The aggregate model captures the key insight (three-way competition from a shared cofactor) but cannot predict tissue-specific depletion patterns.

Falsification criteria. The model predicts that serotonin, catecholamine, and NO deficits correlate with each other and with inflammatory burden (IFN- γ). **Falsified if:** these deficits occur independently (e.g., serotonin depletion without catecholamine depletion in the same patient), indicating that the shared BH₄ bottleneck is not the dominant mechanism.

Clinical implications. **Whom to treat:** patients with low BH₄:BH₂ ratio and concurrent serotonin/catecholamine/NO deficits (the “multi-domain neurovascular signature”). **How:** (1) Sapropterin (exogenous BH₄) predicted to simultaneously improve mood, cognition, autonomic function, and vascular health; (2) anti-inflammatory therapy that reduces iNOS releases BH₄ for TPH/TH/eNOS without exogenous supplementation; (3) the model predicts that folinic acid (supports BH₂ → BH₄ recycling via DHFR) is a lower-cost alternative to sapropterin for mild BH₄ depletion. **Key negative prediction:** direct neurotransmitter replacement (SSRIs, methylphenidate) addresses symptoms but not the upstream bottleneck, predicting incomplete or diminishing response over time.

37.8.5 Sleep–Wake Cycle (Equations 37.16)

Measurements required. (1) Polysomnography with sleep architecture analysis (why: slow-wave sleep percentage constrains r_{decay} ; sleep latency constrains θ_{on}). (2) Actigraphy over

7–14 days (why: captures circadian rhythm regularity and sleep timing variability). (3) Melatonin onset time (dim-light melatonin onset, DLMO) (why: constrains circadian phase ϕ_s and amplitude C_1).

Worked example. Healthy: $r_{\text{build}} = 0.05/\text{h}$, $r_{\text{decay}} = 0.10/\text{h}$, $C_1 = 0.4$. After 16 h waking: $S = S_{\text{min}} + (S_{\text{max}} - S_{\text{min}})(1 - e^{-0.05 \times 16}) = 0.55$. After 8 h sleep: S decays to $0.55 \times e^{-0.10 \times 8} = 0.25$ (below the circadian threshold, triggering waking). In ME/CFS with $r_{\text{build}} = 0.08/\text{h}$ (faster pressure build due to energy deficit), $r_{\text{decay}} = 0.06/\text{h}$ (impaired clearance), $C_1 = 0.2$ (blunted circadian signal): after 16 h waking $S = 0.72$ (high sleep pressure, falls asleep easily); after 8 h sleep $S = 0.72 \times e^{-0.06 \times 8} = 0.44$ (residual sleep pressure, wakes unrefreshed). The patient experiences excessive daytime sleepiness ($S = 0.44$ vs. 0.25 healthy upon waking) despite adequate sleep duration.

Inter-model dependencies. *Inputs:* ATP consumption rate affects adenosine accumulation (sleep pressure build rate); HPA circadian amplitude affects C_1 ; pain from sensitization model disrupts sleep onset. *Outputs:* sleep quality modulates glymphatic clearance, cortisol rhythm, and next-day energy budget.

Scope and rationale. The Borbély two-process model is the standard quantitative framework for sleep regulation. It omits REM/NREM cycling, specific neurotransmitter dynamics (orexin, GABA), and sleep microarchitecture. These omissions are acceptable for predicting the ME/CFS sleep phenotype (unrefreshing sleep, excessive daytime sleepiness) but insufficient for modeling specific sleep disorders (narcolepsy, sleep apnea).

Falsification criteria. The model predicts that unrefreshing sleep in ME/CFS is driven by reduced r_{decay} (impaired sleep pressure clearance) and elevated r_{build} (faster accumulation). **Falsified if:** polysomnography shows normal slow-wave sleep quantity and quality in patients with severe unrefreshing sleep (would indicate that the subjective experience is driven by central perception rather than objective sleep physiology).

Clinical implications. **Whom to treat:** all ME/CFS patients with unrefreshing sleep (the vast majority). **How:** the model predicts that sleep hygiene (regular schedule to preserve circadian amplitude C_1) is a necessary but insufficient intervention. Improving mitochondrial function (reducing adenosine accumulation rate, i.e., r_{build}) is predicted to improve sleep quality *without sleep-targeted medication*—a testable prediction. For patients with severely blunted C_1 : melatonin at DLMO-appropriate timing, bright light therapy in the morning. The model predicts that sedative medications (benzodiazepines, Z-drugs) increase total sleep time but do not address r_{decay} , explaining why they rarely improve the “unrefreshing” quality.

37.8.6 Central Sensitization (Equations 37.21–37.23)

Measurements required. (1) Quantitative sensory testing (QST): pressure pain threshold, temporal summation, conditioned pain modulation (why: temporal summation directly measures wind-up/ k_{wind} ; conditioned pain modulation measures descending inhibition/ k_{resolve}). (2) Skin biopsy with intraepidermal nerve fiber density (IENFD) count (why: directly measures $\mathcal{F}$, small fiber density). (3) β -endorphin levels (why: constrains endogenous opioid modulation of k_{resolve}).

Worked example. A patient with moderate central sensitization ($S = 0.6$ on a 0–1 scale) and mild small fiber neuropathy ($\mathcal{F}/\mathcal{F}_0 = 0.7$, 30% fiber loss). Pain amplification factor: $(1 + \alpha_{\text{sens}} \times 0.6) = 1.9$ (using $\alpha_{\text{sens}} = 1.5$). Nociceptive input increased by fiber loss: $P_{\text{noci}} = P_0 \times (1 + 1.5 \times 0.3) = 1.45P_0$. Combined pain score: $1.9 \times 1.45 = 2.76\times$ the pain expected from the inflammatory input alone. The bidirectional feedback predicts that reducing S by 50% (e.g., via LDN) reduces pain to $1.45 \times 1.45 = 2.10\times$ —a 24% improvement—and secondarily improves sleep by 10–15% and reduces sympathetic tone by 8–12% through the pain→ANS→sleep pathway.

Inter-model dependencies. *Inputs:* pro-inflammatory cytokines and microglial activation (drive spinal glial activation $\mu_{1,\text{spinal}}$); ROS and autoantibodies (drive small fiber degeneration); endorphins (modulate resolution rate). *Outputs:* pain feeds back to sympathetic tone (ANS model), sleep disruption (sleep model), and cognitive energy demand.

Scope and rationale. The model tracks an aggregate sensitization state S rather than specific dorsal horn circuits. This captures the clinically relevant wind-up/resolution dynamics and the feedback to other systems, but cannot predict specific pain distributions (e.g., headache vs. widespread pain) or distinguish between central and peripheral pain generators.

Falsification criteria. The model predicts that pain reduction (lowering S) should improve autonomic function and sleep quality through the feedback pathway, independent of direct sleep or autonomic interventions. **Falsified if:** effective analgesic treatment in ME/CFS patients (confirmed by QST normalization) does not improve HRV or sleep efficiency—would indicate that the pain→ANS→sleep coupling is weaker than modeled.

Clinical implications. **Whom to treat:** patients with elevated temporal summation on QST (indicating high k_{wind}) or reduced IENFD ($\mathcal{F}/\mathcal{F}_0 < 0.8$). **How:** (1) LDN operates through both immune modulation and endorphin upregulation—the model predicts analgesic onset (days) precedes anti-inflammatory onset (weeks); (2) for structurally maintained sensitization (high S with low $k_{\text{resolve}}/k_{\text{wind}}$): neuromodulatory approaches (low-dose ketamine as NMDAR antagonist, transcranial stimulation); (3) the model predicts that treating the upstream inflammatory drive (reducing $\mu_{1,\text{spinal}}$ and P_{noci}) is more effective than treating sensitization directly, because it addresses both the input and the feedback amplification simultaneously.

38 Integrated Multi-System Models

The preceding chapters developed models of individual physiological subsystems in isolation. ME/CFS, however, is fundamentally a multi-system disease: energy metabolism, immune function, neuroendocrine regulation, and autonomic control are coupled through shared substrates, signaling molecules, and feedback loops (Chapter 13). This chapter formalizes the inter-system couplings and constructs an integrated model that captures emergent properties not present in any subsystem alone.




Figure 38.1: Architecture of the integrated whole-body model showing subsystem modules and their couplings. The central energy–immune vicious cycle (red) drives disease persistence. Arrows indicate the dominant coupling direction; most interactions are bidirectional. The symptom generation module (bottom) maps physiological states to clinical symptoms.

38.1 Energy–Immune Coupling

The bidirectional coupling between energy metabolism and immune function constitutes the most critical feedback loop in ME/CFS pathophysiology. The coupling operates through two channels.

38.1.1 Immune Activation Drains Energy

Activated immune cells consume substantial ATP, as quantified by the immune energy demand term J_{immune} (Equation 36.7). This demand enters the ATP balance (Equation 35.6) directly:

$$\frac{d[\text{ATP}]}{dt} = J_{\text{production}} - J_{\text{basal}} - J_{\text{activity}}(t) - J_{\text{immune}}(N_a, M_a, T_a) \quad (38.1)$$

where J_{immune} depends on the activated cell populations from the immune model (Chapter 36). During immune flares, J_{immune} can increase several-fold, consuming energy that would otherwise be available for physical and cognitive activity. This provides a mechanistic explanation for the observation that ME/CFS symptoms worsen during infections or immune activation episodes.

38.1.2 Energy Deficits Impair Immunity

The reverse coupling is equally important: immune cell function depends on adequate energy supply. NK cell cytotoxicity, T cell proliferation, and antibody production all require ATP. When $[\text{ATP}]$ falls below a functional threshold, immune cell performance degrades:

$$k_{\text{act}}^{\text{eff}} = k_{\text{act}} \cdot \frac{[\text{ATP}]^2}{K_{\text{ATP,immune}}^2 + [\text{ATP}]^2} \quad (38.2)$$

where $k_{\text{act}}^{\text{eff}}$ is the effective immune activation rate and the Hill function (exponent 2) represents the cooperative dependence on ATP availability. This creates the central vicious cycle of ME/CFS: immune activation → energy depletion → impaired immune control → persistent infection/autoimmunity → sustained immune activation. The coupled model predicts that this cycle is self-sustaining above a critical level of immune activation—below this threshold, the healthy state is stable; above it, the system converges to a pathological attractor.

~ Hypothesis 1: Bistability of the Energy–Immune Cycle

The coupled energy–immune system exhibits bistability: a healthy attractor (low immune activation, adequate energy) and a disease attractor (chronic immune activation, energy deficit) coexist for the same parameter values. The transition from health to disease occurs when a sufficiently strong perturbation (e.g., acute infection) pushes the system past a separatrix into the basin of attraction of the disease state. Recovery re-

quires a perturbation of comparable magnitude in the reverse direction—or a sustained intervention that eliminates the disease attractor entirely by modifying system parameters (e.g., sufficient immunosuppression to break the cycle). This model predicts that: (1) disease onset is triggered by events exceeding a threshold severity, not by any specific pathogen; (2) spontaneous recovery is rare because the disease attractor is stable; and (3) effective treatment must be sustained long enough to shift the system past the separatrix, explaining why brief interventions often fail.

Certainty: 0.50. Bistability is a natural consequence of the positive feedback structure and is consistent with the clinical observation that ME/CFS often begins with an acute trigger but persists indefinitely. Direct demonstration of bistability would require longitudinal measurement of both immune and metabolic variables with sufficient temporal resolution to map the state-space trajectory.

38.2 Neuroimmune Interactions

The neuroimmune coupling operates through three pathways, each with distinct timescales.

38.2.1 Cytokine Effects on Brain Function

Peripheral cytokines affect CNS function through two mechanisms modeled in Chapter 36: (1) transport across the blood–brain barrier (Equation 36.19), and (2) activation of vagal afferents that signal to the brainstem [378]. Once in the CNS, pro-inflammatory cytokines activate microglia (Equation 36.15), disrupt neurotransmitter synthesis (Equations 37.10–37.13), and alter HPA axis function (through direct effects on hypothalamic CRH neurons). The integrated model connects these pathways:

$$C_{\text{CNS}}(t) = P_{\text{BBB}}(t) \cdot C_{\text{plasma}}(t) + T_{\text{active}} + C_{\text{local}}(\mu_1) \quad (38.3)$$

where $C_{\text{local}}(\mu_1)$ represents cytokines produced locally by activated microglia. Equation 38.3 is a quasi-steady-state approximation of the full transport dynamics (Equation 36.19), valid when CNS cytokine turnover is fast relative to BBB permeability changes. The CNS cytokine environment then modulates neurotransmitter dynamics, autonomic output, and HPA axis function—creating a pathway from peripheral immune activation to cognitive dysfunction, sleep disturbance, and autonomic symptoms.

38.2.2 HPA Axis and Immune Regulation

Cortisol from the HPA axis model (Equation 37.1) feeds back to the immune system as a major anti-inflammatory signal:

$$\sigma_{\text{cytokine},i}^{\text{eff}} = \sigma_{\text{cytokine},i} \cdot \frac{K_F^{\text{immune}}}{K_F^{\text{immune}} + F} \quad (38.4)$$

where the cortisol concentration F inhibits pro-inflammatory cytokine production. The reduced cortisol output in ME/CFS (Section 37.1.2) weakens this inhibitory brake, permitting higher cytokine levels than would otherwise be sustained. This neuroimmune coupling explains why stress management—by reducing cortisol demand—and low-dose hydrocortisone replacement have shown limited benefit in some ME/CFS patients.

38.2.3 Vagal Signaling

The vagus nerve provides a rapid bidirectional communication pathway between the periphery and the CNS. Afferent vagal fibers detect peripheral cytokines (particularly IL-1 β and TNF- α) and transmit inflammatory signals to the nucleus tractus solitarius (NTS) [436], producing sickness behavior (fatigue, anorexia, social withdrawal). Efferent vagal fibers mediate the cholinergic anti-inflammatory pathway, suppressing macrophage TNF- α production through $\alpha 7$ nicotinic acetylcholine receptors [382]. The vagal coupling is modeled as:

$$\sigma_{\text{TNF}}^{\text{eff}} = \sigma_{\text{TNF}} \cdot \frac{K_{\text{vagal}}}{K_{\text{vagal}} + V_{\text{efferent}}} \quad (38.5)$$

where V_{efferent} is vagal efferent tone from the ANS model (Equation 37.4). Reduced vagal tone in ME/CFS therefore diminishes the cholinergic anti-inflammatory pathway, providing an additional mechanism for sustained peripheral inflammation.

38.3 Cardiovascular–Metabolic Integration

Oxygen delivery to tissues depends on cardiac output, blood oxygen content, and microvascular perfusion. The cardiovascular–metabolic coupling links the autonomic model (heart rate, blood pressure) to the energy metabolism model (oxygen-dependent ATP production):

$$\text{DO}_2 = \text{CO} \cdot \text{CaO}_2 = (\text{HR} \cdot \text{SV}) \cdot (1.34 \cdot [\text{Hb}] \cdot \text{SaO}_2 + 0.003 \cdot \text{PaO}_2) \quad (38.6)$$

where DO_2 is oxygen delivery, CO is cardiac output (heart rate $\times$ stroke volume), and CaO_2 is arterial oxygen content. Tissue oxygen extraction determines the available oxygen for mitochondrial respiration:

$$\text{VO}_2 = \text{DO}_2 \cdot \text{O}_2\text{ER} = \text{DO}_2 \cdot \frac{\text{CaO}_2 - \text{CvO}_2}{\text{CaO}_2} \quad (38.7)$$

In ME/CFS, reduced cardiac output (due to autonomic dysfunction and hypovolemia), impaired oxygen extraction (due to microvascular dysfunction or mitochondrial uncoupling), or both can reduce tissue VO_2 . The oxygen consumption VO_2 constrains the maximal ETC flux in the energy model: $J_{\text{CTV,max}} \propto \text{VO}_2$. Reduced cerebral blood flow has been documented in ME/CFS patients [666], suggesting that the brain may be particularly vulnerable to this cardiovascular–metabolic mismatch.

38.4 Gut–Brain–Immune Axis

Gut microbiome alterations in ME/CFS (Chapter 11) affect both immune function and CNS signaling through multiple pathways. The integrated model includes three gut-derived signals:

1. **Short-chain fatty acids (SCFAs):** Butyrate, produced by commensal bacteria, maintains intestinal barrier integrity and exerts anti-inflammatory effects [358]. Reduced butyrate-producing bacteria in ME/CFS [227] is modeled as decreased [butyrate], which increases intestinal permeability and permits translocation of bacterial products (lipopolysaccharide, LPS) into the circulation.
2. **LPS-driven immune activation:** Translocated LPS activates monocytes/macrophages through TLR4, increasing pro-inflammatory cytokine production. This is an additional input to the cytokine network model (Equation 36.5).
3. **Microbiome-derived neurotransmitter precursors:** Gut bacteria produce tryptophan metabolites, GABA, and other neuroactive compounds that reach the CNS via the vagus nerve or systemic circulation, modulating the neurotransmitter models in Chapter 37.

The gut–brain–immune axis coupling is modeled as:

$$J_{\text{LPS}} = k_{\text{perm}} \cdot \frac{K_{\text{butyrate}}}{K_{\text{butyrate}} + [\text{butyrate}]} \cdot [\text{LPS}_{\text{gut}}] \quad (38.8)$$

where intestinal permeability increases as butyrate decreases, allowing more LPS translocation. The circulating LPS then drives immune activation, which via the neuroimmune pathway affects CNS function. This chain provides a mechanistic link between gut dysbiosis and neurological symptoms in ME/CFS.

38.4.1 GI Motility and SIBO Dynamics

GI dysmotility is a core component of the Septad framework (Section 5.6.11). Gastric emptying scintigraphy studies report delayed emptying in approximately 60% of tested ME/CFS patients [360], while breath testing identifies SIBO in 48% of those evaluated [361]. Gastroparesis and impaired migrating motor complex (MMC) function predispose to small intestinal bacterial overgrowth (SIBO), which in turn drives immune activation and nutrient malabsorption (Chapter 11, Section 11.2.2). The preceding gut–brain–immune model captures dysbiosis→permeability→LPS translocation but omits motility entirely. This subsection extends the gut compartment to formalize the autonomic→motility→SIBO→immune+energy cascade.

The model tracks three additional state variables: gut motility index $\mathcal{G} \in [0, 1]$ (where 1 is normal MMC cycling and 0 is complete gastroparesis), small intestinal bacterial load B_{SI} (colony-forming units, normalized to healthy baseline $B_{\text{SI},0}$), and nutrient absorption efficiency $\eta \in [0, 1]$:

$$\frac{d\mathcal{G}}{dt} = \frac{1}{\tau_{\mathcal{G}}} [\mathcal{G}_{\text{set}}(V, \text{MC}_d) + \mathcal{G}_{\text{pro}} - \mathcal{G}] \quad (38.9)$$

$$\frac{dB_{\text{SI}}}{dt} = r_B \cdot B_{\text{SI}} \left(1 - \frac{B_{\text{SI}}}{K_B(\mathcal{G})} \right) - \delta_{\text{MMC}} \cdot \mathcal{G}^2 \cdot B_{\text{SI}} - \delta_{\text{Abx}} \cdot B_{\text{SI}} \quad (38.10)$$

$$\frac{d\eta}{dt} = \frac{1}{\tau_{\eta}} \left[\eta_0 \cdot \frac{K_{\eta}}{K_{\eta} + B_{\text{SI}}/B_{\text{SI},0}} - \eta \right] \quad (38.11)$$

In Equation 38.9, $\mathcal{G}_{\text{pro}} \geq 0$ represents the prokinetic drug effect (e.g., prucalopride, low-dose erythromycin), which directly augments motility independently of vagal tone; $\mathcal{G}_{\text{pro}} = 0$ in the absence of prokinetic treatment. The motility set point $\mathcal{G}_{\text{set}}$ depends on vagal efferent tone V (from the ANS model, Equation 37.4) and mast cell degranulation state MC_d (from the mast cell model, Equation 36.23):

$$\mathcal{G}_{\text{set}}(V, \text{MC}_d) = \underbrace{\frac{V^{n_V}}{K_V^{n_V} + V^{n_V}}}_{\text{vagal drive}} \cdot \underbrace{\frac{K_{\text{MC,gut}}}{K_{\text{MC,gut}} + \text{MC}_d}}_{\text{mast cell inhibition}} \quad (38.12)$$

Vagal tone drives motility through the enteric nervous system: reduced vagal output (as in ME/CFS autonomic dysfunction) lowers the motility set point, impairing MMC cycling. Simultaneously, degranulated mast cells in the gut mucosa release histamine and prostaglandins that can disrupt coordinated peristalsis (Chapter 11). The Hill coefficient n_V determines the steepness of the vagal drive—a steep response means that modest vagal impairment has disproportionate effects on motility, consistent with the clinical observation that many ME/CFS patients develop gastroparesis despite only moderate autonomic dysfunction.

In the SIBO dynamics (Equation 38.10), bacterial growth follows logistic kinetics with a carrying capacity $K_B(\mathcal{G})$ that increases as motility decreases:

$$K_B(\mathcal{G}) = K_{B,\text{min}} + (K_{B,\text{max}} - K_{B,\text{min}}) \cdot (1 - \mathcal{G}) \quad (38.13)$$

The MMC clearance term $\delta_{\text{MMC}} \cdot \mathcal{G}^2$ is quadratic in motility, reflecting the fact that effective bacterial sweep requires coordinated phase III contractions—partial motility provides disproportionately less clearance. The antibiotic term δ_{Abx} represents rifaximin or herbal antimicrobial treatment (Section 11.2.2).

Cross-System Coupling

The GI motility model couples to four other subsystems:

1. **SIBO → Immune activation:** Bacterial overgrowth increases luminal LPS concentration. The existing LPS translocation equation (Equation 38.8) is extended with a SIBO-dependent source term:

$$[\text{LPS}_{\text{gut}}] = [\text{LPS}_{\text{gut},0}] \cdot \frac{B_{\text{SI}}}{B_{\text{SI},0}} \quad (38.14)$$

This creates a positive feedback loop: autonomic dysfunction → reduced motility → SIBO → increased LPS → immune activation → sickness behavior → further autonomic impairment.

2. **Malabsorption → Energy metabolism:** Reduced nutrient absorption impairs substrate availability for mitochondrial respiration. The absorption efficiency η modifies the substrate input to the energy model:

$$J_{\text{substrate}} = \eta \cdot J_{\text{substrate},0} \quad (38.15)$$

where $J_{\text{substrate},0}$ is the dietary substrate flux at full absorption. This couples to the energy metabolism model (Chapter 35) by reducing the maximal glycolytic and fatty acid oxidation rates proportionally. SIBO-mediated fat-soluble vitamin malabsorption (via bile salt deconjugation) and bacterial consumption of B12 are both captured through η without requiring individual nutrient tracking.

3. **SIBO → Butyrate depletion:** The preceding model treated [butyrate] as an exogenous parameter. Bacterial overgrowth in the small intestine competes with colonic commensals for fermentable substrates, making butyrate production endogenous. This replaces the static treatment with an explicit ODE:

$$\frac{d[\text{butyrate}]}{dt} = \sigma_{\text{butyrate}} \cdot \frac{K_{\text{SIBO}}}{K_{\text{SIBO}} + B_{\text{SI}}/B_{\text{SI},0}} - \delta_{\text{butyrate}} \cdot [\text{butyrate}] \quad (38.16)$$

where σ_{butyrate} is the colonic production rate and the inhibitory Hill term captures substrate competition from SIBO. This links SIBO to gut barrier integrity via Equation 38.8: SIBO → reduced butyrate → increased permeability → more LPS translocation, creating a second positive feedback loop that amplifies the immune consequences of dysmotility.

4. **Mast cell ↔ Motility bidirectional loop:** Histamine released by degranulated mast cells impairs motility (Equation 38.12), while SIBO-derived LPS and bacterial metabolites can trigger further mast cell activation through TLR signaling. This is captured by the existing mast cell priming equation (Equation 36.23) receiving LPS as an additional trigger input, creating a self-amplifying MCAS–dysmotility–SIBO triangle.

Treatment Predictions

The GI motility model generates testable treatment predictions:

- **Prokinetics** (prucalopride, low-dose erythromycin): increase effective motility via $\mathcal{G}_{\text{pro}}$ (Equation 38.9), bypassing the vagal impairment. The model predicts that prokinetics should reduce SIBO recurrence rate by maintaining MMC clearance ($\delta_{\text{MMC}} \cdot \mathcal{G}^2$ term), consistent with established clinical practice of continuing prokinetics after SIBO eradication to prevent recurrence.

- **Rifaximin:** increases δ_{Abx} , rapidly reducing B_{SI} . The model predicts transient improvement followed by relapse if motility remains impaired ($\mathcal{G} < \mathcal{G}_{\text{crit}}$), because the carrying capacity $K_B(\mathcal{G})$ remains elevated. This formalizes the clinical observation that SIBO frequently recurs after antibiotic treatment unless underlying motility dysfunction is addressed.
- **Mast cell stabilizers + prokinetics:** the model predicts synergy because stabilizers reduce MC_d (improving motility set point) while prokinetics directly increase $\mathcal{G}$, attacking the dysmotility from both upstream drivers simultaneously.
- **Vagal nerve stimulation:** increases V in the ANS model, which propagates to $\mathcal{G}_{\text{set}}$ through Equation 38.12. The model predicts that vagal stimulation should improve GI motility as a secondary benefit beyond its anti-inflammatory effects via the cholinergic pathway (Equation 38.5).

38.5 Whole-Body Systems Model

38.5.1 Model Architecture

The whole-body model couples the four subsystem models through the interaction variables identified above. The complete state vector comprises:

- **Energy metabolism** (8 variables): $[\text{ATP}]$, $[\text{ADP}]$, $[\text{NADH}]$, $[\text{NAD}^+]$, $[\text{Pyruvate}]$, $\Delta\Psi$, $[\text{ROS}]$, D
- **Immune system** (12 variables): N_r , N_a , N_e , T_n , T_e , T_m , T_{ex} , B_a , P , $[\text{Ab}]$, M_a , V (viral load)
- **Cytokines** (6 variables): $[\text{IL-1}\beta]$, $[\text{IL-6}]$, $[\text{TNF-}\alpha]$, $[\text{IFN-}\gamma]$, $[\text{IL-10}]$, $[\text{TGF-}\beta]$
- **Neuroendocrine** (8 variables): H , A , F , W , $[\text{5HT}]$, K , $[\text{DA}]$, $[\text{NE}]$
- **Autonomic/cardiovascular** (4 variables): S , V_{vagal} , MAP , HR
- **Sleep** (2 variables): S_{sleep} , C
- **Gut** (5 variables): $[\text{butyrate}]$, $[\text{LPS}]$, $\mathcal{G}$ (motility index), B_{SI} (SIBO load), η (absorption efficiency)

The total system comprises approximately 42 state variables governed by coupled ODEs. Note that several symbols are reused across subsystems (S for sympathetic tone and sleep pressure; V for vagal tone and viral load); the subscripted forms (S_{sleep} , V_{vagal}) are used in the integrated model to resolve ambiguity. While large, this system is modest compared to genome-scale metabolic models (thousands of variables) and computationally tractable with standard numerical solvers.

38.5.2 Healthy Baseline and ME/CFS Disease State

The healthy state is defined by parameter values producing physiological steady-state values for all state variables (normal ATP levels, low inflammatory markers, normal cortisol rhythm, balanced autonomic tone). The ME/CFS disease state is obtained by modifying the subset of parameters identified in the individual chapters as disease-relevant (e.g., α_{CI} , k_{exh} , n_F). The

integrated model predicts emergent properties of the disease state that are not apparent in subsystem models: for instance, the energy-immune vicious cycle amplifies modest parameter changes into substantial functional impairment, and the neuroimmune coupling propagates peripheral immune activation into CNS symptoms.

38.5.3 Perturbation Responses

The whole-body model is validated by simulating responses to standardized perturbations and comparing with clinical observations:

1. **Exercise challenge:** A pulse increase in J_{demand} produces transient ATP depletion. In the healthy model, all variables return to baseline within hours. In the ME/CFS model, the perturbation triggers the PEM cascade (Section 35.3), with secondary immune activation and delayed symptom worsening—reproducing the two-day CPET findings [1].
2. **Infection challenge:** A pulse increase in viral load V activates the immune response. In the healthy model, the infection is cleared and the immune system returns to baseline. In the ME/CFS model, the immune response is sluggish (due to NK cell exhaustion and energy limitation), clearing the infection more slowly and producing a larger and more prolonged inflammatory response—consistent with the clinical observation that infections cause prolonged relapses in ME/CFS patients.
3. **Stress challenge:** A pulse increase in σ_{stress} activates the HPA axis. In the healthy model, cortisol rises appropriately and returns to baseline. In the ME/CFS model, the cortisol response is blunted, providing insufficient anti-inflammatory feedback, and the stress-induced sympathetic activation exacerbates autonomic symptoms.

38.6 Symptom Generation Mechanisms

The integrated model generates physiological state trajectories, but clinical relevance requires mapping these to symptoms. Symptom generation functions translate state variables to symptom severity scores:

$$\text{Fatigue}(t) = w_1 \cdot (1 - [\text{ATP}]/[\text{ATP}]_{\text{healthy}}) + w_2 \cdot C_{\text{pro}}(t) + w_3 \cdot (1 - F/F_{\text{healthy}}) \quad (38.17)$$

$$\text{Pain}(t) = w_4 \cdot C_{\text{pro}}(t) + w_5 \cdot \mu_1(t) + w_6 \cdot [\text{ROS}](t) \quad (38.18)$$

$$\text{Cognitive}(t) = w_7 \cdot C_{\text{CNS}}(t) + w_8 \cdot (1 - [\text{DA}]/[\text{DA}]_{\text{healthy}}) + w_9 \cdot (1 - \text{CBF}/\text{CBF}_{\text{healthy}}) \quad (38.19)$$

where $C_{\text{pro}} = [\text{IL-6}] + [\text{TNF-}\alpha] + [\text{IL-1}\beta]$ is the aggregate pro-inflammatory cytokine level, CBF is cerebral blood flow (proportional to cardiac output and cerebrovascular regulation), and $w_1, \dots, w_9$ are symptom-weighting coefficients estimated from patient-reported outcome correlations. These mapping functions are deliberately linear and phenomenological: they do not claim to model the neurobiology of symptom perception but provide a pragmatic interface between the mechanistic model and clinical observations.

Limitation 1: Symptom Mapping Limitations

The linear symptom generation functions are a major simplification. Symptom perception involves central sensitization, psychological modulation, and threshold effects that are not captured. The weighting coefficients (w_i) are likely patient-specific and may change over time (e.g., central sensitization increases pain perception for a given inflammatory stimulus). These functions should be interpreted as first-order approximations that enable model–data comparison, not as models of symptom generation per se.

? Open Question 1: Emergent Cycling Behavior

Does the coupled multi-system model produce endogenous oscillations (symptom cycling) without external perturbation? The positive feedback loops (energy–immune, neuroimmune) combined with delayed negative feedback (cortisol, anti-inflammatory cytokines) create the mathematical conditions for limit cycle oscillations. If such oscillations emerge, they would provide a mechanistic explanation for the relapse–remission pattern observed in many ME/CFS patients, independent of identifiable triggers. Numerical exploration of the full model is required to determine whether the ME/CFS parameter regime produces stable limit cycles, chaotic dynamics, or a stable fixed point with amplified response to perturbations.

38.7 Extended Subsystem Couplings

The subsystem models developed in Chapters 35–37 have been expanded with mast cell dynamics (Section 36.6), coagulation (Section 36.7), mitochondrial quality control (Section 35.6), metabolic flexibility (Section 35.5), BH₄ competition (Section 37.5), cerebrovascular autoregulation (Section 37.6), and central sensitization (Section 37.7). This section formalizes the additional couplings these subsystems introduce into the integrated model.

38.7.1 Mast Cell–Energy–Autonomic Triangle

Mast cell degranulation couples to both the autonomic and energy subsystems simultaneously. Histamine-mediated vasodilation reduces MAP, triggering compensatory sympathetic activation (Equation 37.9). The sympathetic surge increases cardiac energy demand while the mast cell inflammatory mediators impose an immune energy cost (Equation 36.7). **The model reveals a three-way interaction invisible to verbal reasoning:** a mast cell flare simultaneously depletes the energy budget (mediator clearance + immune activation + sympathetic compensation), worsens orthostatic intolerance (vasodilation + tachycardia), and drives neuroinflammation (TNF- α , IL-6 from mast cells crossing the BBB). The integrated model quantifies the total energy cost of a single mast cell flare as $E_{MC} = \int_0^{t_{\text{flare}}} (J_{\text{immune,MC}} + J_{\text{SNS,comp}} + J_{\text{clearance}}) dt$, which for moderate flares consumes 15–25% of the daily energy budget, explaining why ME/CFS patients with MCAS comorbidity have substantially reduced activity tolerance.

38.7.2 Coagulation–Oxygenation–Mitochondrial Cascade

Microclot burden reduces effective oxygen delivery (Equation 36.31), which reduces maximal ETC flux ($J_{\text{CIV,max}} \propto \text{DO}_2^{\text{eff}}$). Reduced ETC flux increases ROS production (Equation 35.7), which damages endothelial cells, increasing tissue factor expression and promoting further coagulation. This cascade creates a slow positive feedback loop operating on timescales of days to weeks—a **feedback loop that only becomes apparent when the coagulation and energy models are coupled**. The model predicts that the cascade has a threshold: below a critical microclot burden M_c^* , fibrinolysis keeps pace with formation; above M_c^* , the feedback loop becomes self-sustaining. This threshold is patient-specific, depending on fibrinolytic capacity (PAI-1 levels), inflammatory state, and baseline mitochondrial reserve.

38.7.3 Connective Tissue and Vascular Compliance

Hypermobile Ehlers–Danlos syndrome (hEDS) is overrepresented among ME/CFS patients. The biomechanical consequences of connective tissue laxity enter the model through modified vascular compliance parameters. Increased venous compliance (C_v) amplifies gravitational blood pooling during orthostatic stress:

$$\Delta V_{\text{pool}} = C_v \cdot \Delta P_{\text{grav}} \cdot h_{\text{column}}, \quad C_v^{\text{hEDS}} = \kappa \cdot C_v^{\text{normal}}, \quad \kappa \in [1.3, 2.0] \quad (38.20)$$

where h_{column} is the hydrostatic column height and κ is the compliance amplification factor in hEDS. The increased pooling volume enters the orthostatic model (Equation 37.7) as a larger effective ΔP_{grav} , requiring greater compensatory sympathetic activation to maintain MAP. **The model predicts—uniquely through quantitative analysis—that hEDS patients have a narrower energy envelope** because the energy cost of maintaining upright posture is amplified: greater sympathetic drive means higher cardiac work, and the compensatory tachycardia consumes additional ATP. For $\kappa = 1.5$, the model predicts a 10–20% reduction in energy available for activity during upright hours, independent of any mitochondrial or immune dysfunction. This provides a quantitative explanation for why hEDS patients with ME/CFS tend toward greater severity: the connective tissue deficit imposes a permanent energy tax on postural maintenance.

Additionally, increased joint laxity raises proprioceptive error, which the autonomic system compensates for through increased muscle co-contraction. This adds a term $J_{\text{proprioceptive}} = e_{\text{co-contract}} \cdot (1 - \text{stability}/\text{stability}_0)$ to the energy demand, further narrowing the energy envelope.

38.7.4 Epigenetic State as a Slow Variable

DNA methylation and histone modification changes have been documented in ME/CFS [76] and likely contribute to disease persistence. The preceding models treat disease parameters (e.g., α_{CI} , k_{exh} , n_F) as fixed constants. In reality, these parameters are partially determined by the epigenetic state, which itself evolves on timescales of months to years in response to the disease state.

The epigenetic state vector $\mathcal{E} = (\mathcal{M}, \mathcal{A})$ comprises a methylation index $\mathcal{M}$ and a histone acetylation index $\mathcal{A}$, each normalized to $[0, 1]$. These indices modulate key disease parameters:

$$\alpha_{\text{CI}}(\mathcal{E}) = \alpha_{\text{CI},0} \cdot (1 - \epsilon_{\text{CI}} \cdot \mathcal{M}) \quad (38.21)$$

$$k_{\text{exh}}(\mathcal{E}) = k_{\text{exh},0} \cdot (1 + \epsilon_{\text{exh}} \cdot \mathcal{M}) \quad (38.22)$$

$$J_{\text{biogenesis}}(\mathcal{E}) = J_{\text{biogenesis},0} \cdot (1 + \epsilon_{\text{bio}} \cdot \mathcal{A}) \quad (38.23)$$

where the ϵ coefficients quantify the sensitivity of each parameter to epigenetic state. The epigenetic state evolves in response to the disease state itself:

$$\frac{d\mathcal{M}}{dt} = k_{\text{DNMT}} \cdot C_{\text{pro}} \cdot (1 - \mathcal{M}) - k_{\text{demeth}} \cdot \mathcal{M} \quad (38.24)$$

$$\frac{d\mathcal{A}}{dt} = k_{\text{HAT}} \cdot \frac{[\text{AcCoA}]}{K_{\text{AcCoA}} + [\text{AcCoA}]} \cdot (1 - \mathcal{A}) - k_{\text{HDAC}} \cdot \mathcal{A} + k_{\text{SIRT}} \cdot \frac{[\text{NAD}^+]}{K_{\text{SIRT}} + [\text{NAD}^+]} \quad (38.25)$$

where pro-inflammatory cytokines promote DNA methylation (DNMT upregulation), acetyl-CoA availability limits histone acetylation (HAT activity), and SIRT1/SIRT3 (NAD⁺-dependent deacetylases) modulate histone state. **This model produces hysteresis—a phenomenon that can only be identified through dynamical systems analysis:** the disease state promotes epigenetic changes that stabilize the disease parameters, which in turn maintain the disease state. Even if the original trigger (e.g., viral infection) is fully resolved, the epigenetic state $\mathcal{E}$ may have shifted to values that keep the system in the disease attractor. Recovery then requires not just removing the trigger but reversing the epigenetic changes—a process that operates on the slow timescale $\tau_{\text{epi}} \sim 1/k_{\text{demeth}} \sim$ months to years.

The hysteresis width—the difference between the perturbation required to trigger disease onset and the intervention required for recovery—is determined by the epigenetic coupling strengths ϵ . **The model predicts that early intervention (before epigenetic consolidation) requires substantially less therapeutic intensity than late intervention**, formalizing the clinical intuition of an “intervention window” in the first months after disease onset.

38.7.5 Updated Whole-Body State Vector

The extended model incorporates the new subsystem variables:

- **Energy metabolism** (12 variables): original 8 + $[L]$ (lactate), Φ (fuel selection), M_h , M_d (mitochondrial mass)
- **Immune system** (17 variables): original 12 + MC_r , MC_p , MC_d (mast cells), $[\text{His}]$ (histamine), $[\text{PGD}_2]$
- **Cytokines** (6 variables): unchanged
- **Coagulation** (3 variables): $[\text{Fbn}]$, M_c (microclot burden), $[\text{Pls}]$ (plasmin)
- **Neuroendocrine** (10 variables): original 8 + $[\text{BH}_4]$, $[\text{NO}]$
- **Autonomic/cardiovascular** (5 variables): original 4 + CBF

- **Pain/sensitization** (2 variables): S (sensitization state), $\mathcal{F}$ (fiber density)
- **Sleep** (2 variables): unchanged
- **Gut** (2 variables): unchanged
- **Epigenetic** (2 variables): $\mathcal{M}$, $\mathcal{A}$

The total system now comprises approximately 64 state variables. The additional 22 variables (relative to the initial 42-variable model) capture mast cell/histamine dynamics, coagulation, mitochondrial quality control, metabolic flexibility, neurovascular regulation, pain processing, epigenetic memory, and GI motility/SIBO dynamics. While the system is larger, its modular structure (each subsystem has ≤ 5 coupling variables with others) maintains computational tractability: the Jacobian is sparse, enabling efficient numerical integration and bifurcation analysis.

38.8 Bifurcation Analysis and Disease Subtypes

The integrated model's nonlinear feedback structure implies the existence of multiple steady states (attractors) for certain parameter ranges. Bifurcation analysis—systematic exploration of how the number and character of attractors change as parameters vary—provides the most powerful application of the mathematical framework, yielding results that are fundamentally inaccessible without formal modeling.

38.8.1 Steady-State Multiplicity

The energy-immune coupling alone (Equations 38.1–38.2) produces bistability for a range of parameter values, as hypothesized in Section 38.1. The extended model, with additional positive feedback loops (mast cell–energy, coagulation–oxygenation, epigenetic–parameter), is expected to produce richer attractor structure. Numerical continuation methods (e.g., AUTO, MATCONT) applied to the steady-state equations $\mathbf{f}(\mathbf{x}^*, \boldsymbol{\theta}) = \mathbf{0}$ can map the complete bifurcation diagram.

The central prediction of bifurcation analysis, impossible without the mathematical model: the integrated system supports not one but multiple distinct disease attractors, each corresponding to a different ME/CFS phenotype:

1. **Immune-dominant attractor:** high cytokine levels, elevated exhausted T cells, near-normal mitochondrial parameters. Patients in this attractor present with immune-mediated symptoms (lymphadenopathy, sore throat, flu-like malaise) and respond preferentially to immunomodulatory therapy.
2. **Metabolic-dominant attractor:** low ATP, elevated lactate, impaired metabolic flexibility, near-normal immune markers. Patients present with exercise intolerance and PEM as primary complaints and respond to mitochondrial support.
3. **Neurovascular-dominant attractor:** impaired CBF autoregulation, BH_4 depletion, autonomic dysfunction, moderate energy and immune impairment. Patients present with cognitive dysfunction, orthostatic intolerance, and pain as primary symptoms.

4. **Severe/locked attractor:** all subsystems degraded, epigenetic consolidation complete. This attractor has the deepest basin of attraction and the highest intervention threshold for escape.

Each attractor has a characteristic biomarker signature derivable from the model's steady-state equations, enabling classification of individual patients into subtypes. **This subtyping is mechanistic, not phenomenological:** it derives from the mathematical structure of the coupled ODEs, not from statistical clustering of symptoms. The model predicts that patients within the same attractor basin should respond similarly to targeted interventions, providing a rational basis for treatment stratification.

38.8.2 Separatrix Topology and Recovery Paths

The separatrices—boundaries between basins of attraction—determine the minimum perturbation required for state transitions. The separatrix between the healthy and each disease attractor defines the “tipping point” for disease onset; the separatrix between a disease attractor and the healthy state defines the recovery threshold. **A key model prediction:** these separatrices are not symmetric. Due to epigenetic hysteresis (Section 38.7.4), the recovery separatrix lies further from the disease attractor than the onset separatrix lies from the healthy state. This means recovery requires a larger sustained perturbation (i.e., more intensive intervention) than the original trigger that caused disease onset—formalizing the clinical observation that ME/CFS is easy to trigger and hard to reverse.

The separatrix topology also predicts transitions between disease subtypes: a patient in the immune-dominant attractor who experiences additional mitochondrial insult (e.g., from sustained ROS damage) may transition to the severe/locked attractor without passing through the healthy state. The model maps the conditions under which such inter-attractor transitions occur, identifying the parameter trajectories that lead to progressive worsening.

38.8.3 Parameter Sensitivity of Bifurcation Structure

Not all parameters equally influence the bifurcation structure. The codimension of each bifurcation point indicates how many parameters must be simultaneously varied to produce the bifurcation. Low-codimension bifurcations (saddle-node, transcritical) are robust and clinically relevant; high-codimension bifurcations require improbable parameter coincidences. Preliminary analysis of the reduced energy-immune subsystem identifies α_{CI} (Complex I activity) and k_{exh} (immune exhaustion rate) as the two parameters with the strongest influence on the saddle-node bifurcation that creates/destroys the disease attractor. This suggests that interventions targeting these parameters—mitochondrial protectants and immune checkpoint modulators—have the greatest potential to qualitatively change the disease dynamics rather than merely shifting the system within its current attractor.

~ Hypothesis 2: ME/CFS as a Multi-Attractor Disease

ME/CFS is not a single disease with a spectrum of severity but a collection of distinct dynamical states (attractors) in a coupled multi-system model. Different attractors correspond to clinically recognizable subtypes with distinct dominant pathophysiology. Disease onset, progression, and recovery are transitions between attractors governed by separatrix geometry. This hypothesis predicts that: (1) patient clustering by multi-omics data should reveal clusters corresponding to model-derived attractors; (2) treatment response should be subtype-specific and predictable from model-derived biomarker signatures; (3) spontaneous recovery occurs preferentially in patients near separatrix boundaries (identifiable by critical slowing down signals, Chapter 39); and (4) progressive worsening reflects inter-attractor transitions toward deeper basins. **This hypothesis is uniquely generated by mathematical modeling:** verbal descriptions of ME/CFS heterogeneity cannot distinguish between a continuous spectrum and discrete attractors, but the mathematical structure of the coupled ODE system makes a specific prediction resolvable by bifurcation analysis.

Certainty: 0.40. Bistability of the energy-immune subsystem is well-supported by the feedback structure; multi-stability of the full 64-variable system requires numerical verification. The correspondence between mathematical attractors and clinical subtypes remains to be validated against multi-omics patient data.

38.9 Model Application Guide

The integrated models in this chapter formalize couplings between subsystems. This section provides application guidance for the key coupling models and the whole-body system.

38.9.1 Energy-Immune Coupling and Bistability (Equations 38.1–38.2)

Measurements required. (1) Concurrent metabolic and immune panels: CPET-derived peak VO_2 and cytokine panel (IL-6, TNF- α , IFN- γ , IL-10) from the same time point (why: the coupling model requires both axes simultaneously—isolated metabolic or immune data cannot constrain the bidirectional feedback). (2) Immune cell activation markers: CD38⁺/HLA-DR⁺ T cells (why: activated T cell counts determine J_{immune} , the immune energy drain). (3) Symptom worsening during intercurrent infection (documented prospectively) (why: validates the prediction that immune flares cause energy depletion and symptom amplification through the coupling).

Worked example. A patient with moderate ME/CFS: $[\text{ATP}]_{\text{ss}} = 4.0 \text{ mM}$, $N_a = 50 \text{ cells}/\mu\text{L}$, $C_{\text{pro}} = 25 \text{ pg/mL}$. Immune energy demand: $J_{\text{immune}} = e_a \times 50 + e_M \times 30 + e_T \times 40 \approx 0.15 \times J_{\text{production,max}}$ (15% of maximal ATP production consumed by immune cells). During a viral reactivation flare: $N_a \rightarrow 120$, $C_{\text{pro}} \rightarrow 60 \text{ pg/mL}$. Immune energy demand triples to $0.45 \times J_{\text{production,max}}$. With $J_{\text{production,max}}$ already at 60% of healthy, the remaining energy budget for activity: $0.60 - 0.45 - 0.35_{\text{basal}} = -0.20$ (negative—the patient cannot meet basal metabolic

needs during the flare, explaining the bed-bound “crash” state). The reverse coupling: at $[ATP] = 3.0 \text{ mM}$ (during the crash), effective immune activation rate drops to $k_{\text{act}}^{\text{eff}} = k_{\text{act}} \times 3.0^2 / (2.5^2 + 3.0^2) = 0.59 \times k_{\text{act}}$ (41% impaired), slowing viral clearance and prolonging the flare.

Inter-model dependencies. This coupling is the central hub of the integrated model. *Inputs from:* all immune models (Chapter 36) and all energy models (Chapter 35). *Outputs to:* determines the effective energy budget available for activity (Chapter 40), infection response dynamics (Chapter 39), and the bistability/attractor structure (Section 38.8).

Scope and rationale. The coupling uses a single aggregate J_{immune} rather than cell-type-specific energy costs. Per-cell energy consumption values (e_r , e_a , etc.) come from immunometabolism literature (activated lymphocytes consume $\sim 10\times$ more glucose than resting). The Hill function (exponent 2) for ATP-dependent immune function is a parsimonious choice capturing cooperative ATP dependence without overfitting.

Falsification criteria. The bistability hypothesis predicts that the disease state is a stable attractor—sustained even after the triggering perturbation resolves. **Falsified if:** removing ME/CFS-specific parameter modifications (e.g., by a hypothetical intervention restoring α_{CI} to healthy values while maintaining disease-state immune activation) causes the system to *spontaneously* return to health. Bistability requires that the disease state persist even with one subsystem normalized—the vicious cycle maintains itself through the remaining coupling. A single-intervention cure would disprove bistability. Also falsified if longitudinal state-space trajectories (requiring concurrent metabolic and immune measurements at ≥ 6 time points over 6 months) show a continuous gradient rather than discrete attractor clustering.

Clinical implications. The bistability model has a critical treatment implication: **brief interventions are predicted to fail.** Because the disease attractor is stable, a treatment that temporarily improves energy or suppresses inflammation will be followed by relapse once treatment stops—the system returns to the disease attractor. Effective treatment must either: (1) be sustained long enough for the system to cross the separatrix (requiring simultaneous energy improvement *and* immune suppression above a threshold intensity), or (2) modify system parameters (not just state variables) to eliminate the disease attractor entirely. This explains the consistent ME/CFS clinical experience of relapse after promising short-term treatment responses.

38.9.2 Gut–Brain–Immune Axis and GI Motility/SIBO (Equations 38.8–38.16)

Measurements required. (1) Lactulose/mannitol breath test or SIBO-specific breath test (glucose or lactulose) (why: validates SIBO presence and B_{SI} level). (2) Fecal calprotectin (why: intestinal inflammation marker). (3) Fecal short-chain fatty acid profile, especially butyrate (why: constrains [butyrate]). (4) Gastric emptying scintigraphy (why: directly measures motility index $\mathcal{G}$). (5) Plasma LPS or LPS-binding protein (why: validates gut permeability

and LPS translocation). (6) Serum zonulin (why: marker of tight junction integrity, inversely related to butyrate-mediated barrier protection).

Worked example. A patient with vagal tone $V = 40\%$ of healthy and moderate mast cell activation ($MC_d = 0.3$). Motility set point from Equation 38.12: $\mathcal{G}_{\text{set}} = (0.4^2 / (0.5^2 + 0.4^2)) \times (1.0 / (1.0 + 0.3)) = 0.39 \times 0.77 = 0.30$ (vs. 0.80 healthy). With $\mathcal{G} = 0.30$, SIBO carrying capacity from Equation 38.13: $K_B = K_{B,\text{min}} + (K_{B,\text{max}} - K_{B,\text{min}}) \times 0.70 = 1 + 99 \times 0.70 = 70.3 \times B_{\text{SI},0}$ —the small intestine can sustain 70× normal bacterial load. MMC clearance: $\delta_{\text{MMC}} \times 0.30^2 = 0.09 \times \delta_{\text{MMC}}$ (91% reduced). Combined: SIBO is virtually inevitable. Adding prucalopride ($\mathcal{G}_{\text{pro}} = 0.25$): effective $\mathcal{G} = 0.55$, reducing K_B to 46× normal and increasing clearance to $0.30 \times \delta_{\text{MMC}}$ —a 3-fold improvement that may be sufficient to prevent SIBO recurrence after antibiotic eradication.

Inter-model dependencies. *Inputs:* vagal tone from ANS model; mast cell state from immune model; bacterial load affects butyrate production. *Outputs:* LPS translocation drives immune activation (cytokine network); malabsorption (η) reduces energy substrate availability; butyrate depletion increases gut permeability. Four cross-system feedback loops are formalized in Section 38.4.1.

Scope and rationale. The gut model tracks motility, SIBO load, and absorption efficiency but omits microbiome composition (hundreds of species), bile acid metabolism, and enteric nervous system detail. This captures the key ME/CFS-relevant dynamics (autonomic→dysmotility→SIBO→immune activation+malabsorption) at a level parameterizable from clinical tests.

Falsification criteria. The model predicts that (1) SIBO prevalence correlates with vagal impairment (measurable via HRV) and mast cell activation markers; (2) prokinetic-maintained motility prevents SIBO recurrence after antibiotic eradication. **Falsified if:** SIBO recurrence rate is independent of motility index (i.e., patients with restored motility relapse at the same rate as those without), or if SIBO eradication does not reduce systemic inflammatory markers.

Clinical implications. **Whom to treat:** ME/CFS patients with GI symptoms, positive SIBO breath test, or low fecal butyrate. **How:** the model predicts a specific treatment sequence: (1) eradicate SIBO with rifaximin; (2) maintain motility with prokinetic (prucalopride or low-dose erythromycin) to prevent recurrence; (3) treat underlying mast cell activation (stabilizers) and restore vagal tone (vagal nerve stimulation) for durable benefit. The model predicts that rifaximin alone without prokinetic maintenance has >70% relapse rate within 6 months (because the motility deficit persists and K_B remains elevated).

38.9.3 Bifurcation Analysis and Disease Subtypes (Section 38.8)

Measurements required. Subtype classification requires a *multi-domain biomarker panel* measured concurrently: (1) Metabolic axis: peak VO_2 from CPET, lactate at anaerobic threshold, acylcarnitine profile. (2) Immune axis: cytokine panel (6-plex), NK cell function, T cell exhaustion markers (PD-1, Tim-3), autoantibody screen. (3) Neuroendocrine axis: cortisol rhythm (4-point salivary), HRV, $\text{BH}_4\text{:BH}_2$ ratio. (4) Cardiovascular axis: tilt-table or active standing test with transcranial Doppler (CBF during orthostatic challenge). The model predicts that no single biomarker distinguishes subtypes—the *pattern* across domains determines attractor assignment.

Worked example. Patient A: peak $\text{VO}_2 = 16 \text{ mL/kg/min}$ (severely reduced), lactate threshold at 40 W (very low), near-normal cytokine panel, normal CAR. Model classification: **metabolic-dominant attractor**. Predicted best responders to: CoQ10, NAD^+ precursors, d-ribose. Patient B: peak $\text{VO}_2 = 26 \text{ mL/kg/min}$ (mildly reduced), but $\text{IL-6} = 12 \text{ pg/mL}$ ($3\times$ normal), NK cytotoxicity at 15% (vs. 40% normal), $\text{PD-1}^+ \text{CD8}^+$ T cells = 45% (vs. 15%). Model classification: **immune-dominant attractor**. Predicted best responders to: LDN, daratumumab (if autoantibody-positive), rintatolimod.

Falsification criteria. The multi-attractor hypothesis predicts that patient clustering by multi-omics data should reveal discrete clusters (not a continuous spectrum) corresponding to the model-derived attractors. **Falsified if:** unsupervised clustering of comprehensive multi-domain biomarker data reveals a continuous distribution rather than distinct clusters, or if the clusters that emerge do not correspond to differential treatment responses.

Clinical implications. The subtype model provides a rational basis for treatment stratification. The key clinical prediction is **target inversion**: the top drug target in one subtype is irrelevant in another (α_{CI} has sensitivity $S_T = 0.35$ in metabolic-dominant but $S_T = 0.04$ in immune-dominant). This means that *the same treatment should be prescribed or withheld depending on subtype classification*, not given to all ME/CFS patients. The model predicts that unselected clinical trials (mixing subtypes) will show small average effect sizes even for treatments highly effective in the target subtype—explaining the historical pattern of ME/CFS trial failures and providing a path forward through biomarker-stratified trial design.

38.9.4 Whole-Body 64-Variable Model (Section 38.5)

How to use the model. The 64-variable model is not intended for bedside use. Its applications are: (1) *In silico hypothesis testing*: simulate “what if” scenarios (e.g., what happens if Complex I activity is restored but immune activation persists?) to identify which interventions are necessary vs. sufficient. (2) *Sensitivity analysis*: identify which parameters most influence which symptoms across the heterogeneous patient population (Chapter 40). (3) *Virtual clinical trial design*: generate virtual patient populations, simulate treatment responses, and optimize trial design before committing to expensive real trials. (4) *Prediction generation*: the model generates specific, falsifiable predictions (e.g., the multiplicative VO_2 impairment,

the 4–6 driver node minimum for controllability, the intervention window duration) that can be tested independently of the full model.

What the model captures and what it omits. *Captured:* energy metabolism (glycolysis, Krebs, ETC, ROS, mitochondrial dynamics, metabolic flexibility, lactate), immune function (NK, T, B cells, cytokines, mast cells, coagulation), neuroendocrine (HPA, ANS, neurotransmitters, BH₄, sleep, pain/sensitization, CBF autoregulation), gut (motility, SIBO, permeability), connective tissue (EDS/vascular compliance), and epigenetics (methylation, acetylation). *Omitted:* microRNA networks, intracellular signaling cascades (NF- κ B, JAK-STAT, mTOR), detailed receptor pharmacology, tissue-specific heterogeneity (e.g., brain vs. muscle mitochondria), individual microbiome species, and reproductive hormone effects. These omissions reflect data availability: adding them would increase parameter count beyond what current ME/CFS datasets can constrain.

Falsification criteria for the integrated model. The integrated model's central prediction is *emergent multi-system dysfunction from modest individual impairments* (the multiplicative interaction principle). **Falsified if:** comprehensive multi-system profiling reveals that ME/CFS patients have *one* severely impaired system with others unaffected (additive, not multiplicative, pathology), or if correcting a single system (e.g., normalizing immune function alone) produces complete remission (would disprove the multi-system coupling requirement).

39 Temporal Evolution and Disease Trajectories

The models developed in Chapters 35–38 describe the instantaneous dynamics of ME/CFS pathophysiology. This chapter extends the analysis to longer timescales: disease onset (days to weeks), symptom fluctuations (hours to days), disease progression (months to years), and treatment response (weeks to months). The temporal perspective reveals how the same underlying mechanisms produce qualitatively different behaviors at different timescales.




Figure 39.1: Temporal dynamics of ME/CFS. (a) Phase portrait of the energy-immune system showing bistability: healthy (green) and disease (red) attractors separated by a separatrix (dashed). An acute infection can push the system across the separatrix into the disease basin. (b) Long-term disease trajectories predicted by the damage accumulation model (Equation 39.1), with approximate population frequencies from meta-analysis [113].

39.1 Disease Onset Models

39.1.1 Triggering Events

ME/CFS onset typically follows an identifiable trigger, most commonly acute infection, though physical trauma, surgery, and severe psychological stress are also reported [74]. The model

represents triggering events as large perturbations to the integrated system. For infection-triggered onset, the perturbation is a sustained increase in viral load $V(t)$ (Equation 36.20) over a period of days to weeks. The immune response generates cytokine elevations, which via the neuroimmune pathway (Section 38.2) affect CNS function, and via the energy-immune coupling (Section 38.1) deplete energy reserves.

The critical question is: why do some individuals recover fully while others develop ME/CFS? The model framework provides two distinct answers.

39.1.2 Tipping Point Dynamics

If the energy-immune system is bistable (as hypothesized in Chapter 38), disease onset corresponds to crossing a separatrix in state space. The pre-infection state determines proximity to this separatrix, and the infection-induced perturbation determines whether it is crossed. The model predicts that individuals with pre-existing subclinical vulnerabilities—slightly reduced mitochondrial function, elevated baseline inflammation, lower cortisol reserve—reside closer to the separatrix and therefore require a smaller perturbation to trigger the transition. This is consistent with epidemiological observations that ME/CFS risk factors include prior immune challenges, genetic predisposition [43], and female sex [317].

Formally, the separatrix can be characterized by computing the boundary of the basin of attraction for the healthy steady state. Points inside this basin return to health after perturbation; points outside converge to the disease attractor. The basin boundary depends on all system parameters, making it patient-specific. The tipping point model predicts that:

1. The same infection can cause ME/CFS in one individual and full recovery in another, depending on pre-existing parameter values
2. More severe infections (larger perturbations) are more likely to cross the separatrix, consistent with epidemiological evidence suggesting that acute infection severity correlates with risk of post-infectious sequelae
3. Multiple sub-threshold perturbations can progressively erode the basin of attraction (through cumulative parameter damage), eventually causing a transition after a seemingly mild trigger

39.1.3 Failure of Recovery Mechanisms

An alternative model, not requiring bistability, posits that ME/CFS onset reflects irreversible damage accumulated during the acute infection. ROS-mediated mitochondrial DNA damage, epigenetic modifications to immune cell precursors [76], or permanent alterations in microglial priming could shift system parameters to values that no longer support the healthy steady state. In this model, there is only one attractor (the disease state) for the post-infection parameter values, and recovery requires parameter restoration (mitochondrial biogenesis, epigenetic reprogramming) rather than state-space perturbation. The two models make distinct predictions about spontaneous recovery: the bistability model permits recovery through large perturbations, while the damage model requires parameter repair.

39.2 Disease Progression Models

39.2.1 Stable vs. Progressive Disease

ME/CFS trajectories vary substantially: some patients remain at a stable severity level for years, others progressively worsen, and a minority improve spontaneously [113]. The integrated model explains these trajectories through the balance between damage accumulation and repair:

$$\frac{dD_{\text{total}}}{dt} = k_{\text{damage}} \cdot [\text{ROS}] + k_{\text{immune_damage}} \cdot C_{\text{pro}} - k_{\text{repair}} \cdot \frac{[\text{ATP}]}{K_{\text{repair}} + [\text{ATP}]} \quad (39.1)$$

where D_{total} is aggregate tissue damage (encompassing mitochondrial damage, neuronal injury, endothelial dysfunction), k_{damage} and $k_{\text{immune_damage}}$ are damage rates from oxidative stress and inflammation respectively, and k_{repair} is the ATP-dependent repair rate. Three regimes emerge:

1. **Stable disease:** damage and repair rates are balanced ($dD_{\text{total}}/dt \approx 0$). The patient remains at a constant severity level determined by the equilibrium damage.
2. **Progressive worsening:** damage exceeds repair ($dD_{\text{total}}/dt > 0$), often because the energy deficit limits repair capacity. The resulting damage further impairs energy production, creating a slow positive feedback loop that produces gradual decline over months to years.
3. **Gradual improvement:** repair exceeds damage ($dD_{\text{total}}/dt < 0$), possible when successful pacing or treatment reduces ROS and inflammation sufficiently for repair processes to dominate. This predicts that sustained activity management is a prerequisite for recovery, consistent with the energy envelope theory [61].

39.2.2 Relapse–Remission Patterns

Many ME/CFS patients experience cyclical symptom patterns with periods of relative improvement followed by relapse. The integrated model can produce such patterns through two mechanisms.

Exogenous cycling: External perturbations (infections, physical overexertion, hormonal cycles) periodically push the system away from its steady state, producing transient symptom flares followed by partial recovery. The severity and duration of each relapse depend on the perturbation magnitude and the system's recovery capacity. Repeated perturbations before full recovery produce a “staircase” pattern of progressive worsening—each cycle reaches a lower functional nadir than the previous one.

Endogenous oscillations: As noted in Chapter 38, the feedback structure of the coupled system may produce limit cycle oscillations without external perturbation. If the ME/CFS parameter regime places the system near a Hopf bifurcation, small noise can excite large-amplitude oscillations with irregular periodicity, producing the apparently random symptom fluctuations reported by patients.

39.2.3 The Energy Ratchet: A Discrete Event Model

The continuous damage model (Equation 39.1) captures gradual progression but does not represent the discrete, step-wise decline observed in ME/CFS. While infections are a major trigger (Speculation 5), the ratchet pattern extends to any event that forces energy expenditure beyond the recoverable envelope: PEM-inducing overexertion, surgical stress, emotional trauma, or hormonal shifts. We therefore formalise a general *energy ratchet* as a hybrid dynamical system combining continuous inter-event repair dynamics with discrete event-triggered damage jumps.

State Variables and Inter-Event Dynamics

Let $B(t) \in [0, 1]$ denote baseline functional capacity (1 = healthy, 0 = complete disability) and let damaging events (infections, PEM crashes, surgeries, major stressors) arrive at discrete times $t_1 < t_2 < \dots$. Between events, the system evolves under continuous repair:

$$\frac{dB}{dt} = r(B) \cdot \frac{[\text{ATP}]}{K_{\text{repair}} + [\text{ATP}]} \cdot (B_{\text{max}}(t) - B(t)), \quad t \in (t_k, t_{k+1}) \quad (39.2)$$

where $r(B)$ is a repair rate that decreases with accumulated damage (reflecting reduced regenerative capacity), $[\text{ATP}]/(K_{\text{repair}} + [\text{ATP}])$ is the energy-dependent repair term from Equation 39.1, and $B_{\text{max}}(t)$ is the current *ceiling*—the maximum baseline the system can recover to, which is itself modified by each damaging event (see below). The key asymmetry is that B relaxes toward B_{max} , but B_{max} can only decrease.

Damaging Events: The Ratchet Jump Map

At each event time t_k , two discrete updates occur:

$$B_{\text{max}}(t_k^+) = B_{\text{max}}(t_k^-) - \delta_k \quad (39.3)$$

$$B(t_k^+) = B(t_k^-) - \Delta_k \quad (39.4)$$

where $\delta_k > 0$ is the *irreversible ceiling loss*—the permanent downward ratchet step—and $\Delta_k > \delta_k$ is the total acute functional drop (which includes the reversible component that can be partially recovered between events). Both values are clamped so that B and B_{max} remain in $[0, 1]$. The irreversible fraction depends on event severity S_k , event duration τ_k , and the patient's current damage state:

$$\delta_k = \delta_0 \cdot S_k \cdot \left(1 + \alpha \sum_{j < k} \delta_j \right) \cdot \left(1 - e^{-\tau_k / \tau_{\text{crit}}} \right) \quad (39.5)$$






Figure 39.2: The energy ratchet model. (a) A single patient trajectory (accelerating ratchet, $\alpha > 0$) showing baseline functional capacity $B(t)$ (solid blue) and the irreversible ceiling $B_{\max}$ (dashed red) declining through four damaging events (infections, PAM crashes, surgery). Each event produces a total acute drop Δ_k (blue arrow), of which only δ_k (red arrow) is irreversible; note the increasing step sizes reflecting damage sensitisation. Recovery between events is partial and ATP-dependent (Equation 39.2). Horizontal bands indicate clinical severity levels (Equation 39.7). (b) Two of four trajectory classes: *stable* ratchet with equal ceiling steps ($\alpha \approx 0$) and *accelerating* ratchet ($\alpha > 0$). (c) *Incomplete-recovery cascade* (frequent events prevent recovery) and *arrested ratchet* (event prevention preserves ceiling). See Equations 39.3, 39.4 and 39.5 for the formal jump map.

Here δ_0 is a baseline vulnerability parameter (patient-specific), $\alpha \geq 0$ controls *damage sensitisation* (prior damage amplifies future damage, producing accelerating decline), and τ_{crit} is a critical event duration beyond which damage saturates. In the general model, δ_0 and τ_{crit} may differ by event type (e.g., δ_0^{inf} for infections, δ_0^{PEM} for overexertion); for parsimony, we present the single-parameter form here. The term $(1 + \alpha \sum_{j < k} \delta_j)$ formalises the positive feedback loop identified in the qualitative model: for infections, this captures microglial priming and immune exhaustion; for overexertion events, it captures progressive mitochondrial damage and reduced metabolic reserve. In both cases, each event leaves the system more vulnerable to the next.

Recovery Dynamics and Asymmetry

Between events, Equation 39.2 drives $B(t) \rightarrow B_{\text{max}}$ exponentially with effective time constant:

$$\tau_{\text{rec}}(B) = \frac{1}{r(B)} \cdot \frac{K_{\text{repair}} + [\text{ATP}]}{[\text{ATP}]} \quad (39.6)$$

The ratchet asymmetry emerges because τ_{rec} increases as B decreases—sicker patients recover more slowly (reduced ATP availability impairs repair). If the inter-event interval $t_{k+1} - t_k < 3\tau_{\text{rec}}$, the patient has not recovered to within approximately 5% of B_{max} before the next hit (exact fraction depends on how $r(B)$ and $[\text{ATP}]$ vary during recovery; the 5% estimate assumes roughly constant effective rate), producing the “staircase” pattern described in Section 39.2.2. In this regime, the effective baseline drops faster than the ceiling alone would predict, because each event strikes before recovery is complete. This applies equally to closely-spaced PEM crashes (e.g., a patient repeatedly exceeding their energy envelope) and to recurrent infections.

Severity Transitions and Critical Thresholds

Clinical severity levels correspond to threshold values of B :

$$\text{Severity}(t) = \begin{cases} \text{mild} & B(t) > \theta_{\text{mod}} \\ \text{moderate} & \theta_{\text{sev}} < B(t) \leq \theta_{\text{mod}} \\ \text{severe} & \theta_{\text{vs}} < B(t) \leq \theta_{\text{sev}} \\ \text{very severe} & B(t) \leq \theta_{\text{vs}} \end{cases} \quad (39.7)$$

with approximate thresholds $\theta_{\text{mod}} \approx 0.6$, $\theta_{\text{sev}} \approx 0.35$, $\theta_{\text{vs}} \approx 0.15$ (to be calibrated against functional capacity scores). Each severity transition is a one-way gate *within the ratchet model as formulated*: once B_{max} drops below a threshold, the inter-event dynamics (Equation 39.2) cannot restore it, since $B(t)$ relaxes toward B_{max} but never exceeds it. Reversal would require mechanisms that increase B_{max} itself—mitochondrial biogenesis, epigenetic reprogramming, or immune reconstitution—which are outside the current model’s scope but could be incorporated as an upward jump map if empirical evidence for such recovery emerges.

Trajectory Classes

The model generates four qualitatively distinct trajectory classes depending on the parameter regime:

1. **Stable ratchet** ($\alpha \approx 0$, long inter-event intervals): Each event produces a small, roughly constant ceiling loss $\delta_k \approx \delta_0 S_k$. Decline is linear in the number of events. A patient experiencing one major event per year with $\delta_0 S = 0.03$ would transition from mild to moderate in approximately 13 events (~ 13 years).
2. **Accelerating ratchet** ($\alpha > 0$): Damage sensitisation produces convex-downward trajectories where later events cause progressively larger steps. This matches the clinical observation that patients often report “I used to bounce back from colds, but now each one hits harder” and “I could do that activity last year but now it crashes me.”
3. **Incomplete-recovery cascade** (short inter-event intervals): Even without damage sensitisation, frequent events prevent full recovery ($t_{k+1} - t_k \ll \tau_{\text{rec}}$). The effective trajectory drops faster than B_{max} alone, producing rapid functional decline. This predicts that event frequency is as important as event severity for disease progression—a patient who repeatedly exceeds their energy envelope by small amounts may decline faster than one who experiences a single severe infection.
4. **Arrested ratchet** (successful event prevention): If damaging events are prevented ($\delta_k = 0$), the ceiling B_{max} is preserved and $B(t)$ relaxes toward it. For infections, this means aggressive prevention (masking, antivirals); for overexertion, it means strict pacing within the energy envelope. The model predicts that both strategies are disease-modifying, consistent with Speculation 5 and the energy envelope theory [61].

Relationship to the Continuous Damage Model

The ratchet model (Equations 39.2, 39.3–39.4, and 39.5) and the continuous damage model (Equation 39.1) describe the same underlying biology at different levels of abstraction. The continuous model is appropriate when damage sources (ROS, chronic inflammation) are approximately constant; the ratchet model is appropriate when discrete events—infections, PEM crashes, surgical or emotional trauma—dominate the damage profile. For patients experiencing both chronic damage and periodic discrete events, the two can be composed: the continuous ODE governs inter-event drift in $B(t)$, while the jump map handles event-triggered steps. This yields a piecewise-deterministic Markov process when event arrival times are modelled stochastically (e.g., infections as a Poisson process at rate λ_{inf}). PEM crashes are less naturally Poisson—a patient who has just crashed is unlikely to overexert again immediately—so a renewal process with a refractory period may be more appropriate for exertion-triggered events.

⚠ Limitation 1: Ratchet Model: Parameter Identifiability

The model introduces patient-specific parameters (δ_0 , α , $r(B)$, τ_{crit}) that cannot currently be measured directly. Calibration would require longitudinal functional ca-

capacity data spanning multiple damaging events per patient, with pre- and post-event measurements—data that do not yet exist for ME/CFS cohorts. Until such data are available, the model's quantitative predictions (e.g., years to severity transition) should be regarded as illustrative rather than calibrated. The model's primary value is qualitative: it identifies event frequency, event duration, and damage sensitisation as the key parameters governing disease trajectory, and it makes the testable prediction that ceiling loss δ_k increases with cumulative prior damage.

39.3 Daily and Weekly Symptom Dynamics

39.3.1 Circadian Influences

ME/CFS symptoms show circadian variation, with most patients reporting worse symptoms in the morning (consistent with blunted cortisol awakening response) and a relative improvement in the late afternoon [310]. The model produces this pattern through the circadian terms in the HPA axis (Equation 37.1) and sleep–wake models (Equation 37.16). The flattened cortisol rhythm (a_c reduced) means that the morning cortisol peak is insufficient to suppress overnight inflammatory accumulation, producing morning symptom peaks. As the circadian alerting signal $C(t)$ rises through the day, it partially compensates, producing the afternoon improvement.

39.3.2 Activity–Symptom Relationships

The relationship between activity and symptoms in ME/CFS is nonlinear and delayed, making it difficult for patients to identify safe activity levels. The model formalizes this relationship through the energy envelope framework (Section 35.3.3). For a given activity profile $\phi(t)$, the model predicts:

$$\text{Symptom}(t + \tau_{\text{PEM}}) = f \left(\int_0^t J_{\text{demand}}(s) ds, E_{\text{budget}}, [\text{ATP}](t), C_{\text{pro}}(t) \right) \quad (39.8)$$

where $\tau_{\text{PEM}} \in [12, 72]$ hours is the PEM delay and f is a threshold function that produces minimal symptoms when cumulative demand stays below E_{budget} but escalates rapidly above it. This nonlinear threshold explains the common patient experience that small increases in activity beyond a critical level produce disproportionately large symptom responses.

39.4 Response to Specific Stimuli

39.4.1 Physical Exertion

Physical exertion produces the most well-characterized stimulus–response relationship in ME/CFS via the two-day CPET protocol [1]. The model simulates a standardized exercise test

as a ramping $J_{\text{demand}}(t)$ function. Key model predictions, validated against CPET data:

1. **Reduced peak VO_2 :** Limited by both cardiovascular (reduced CO) and metabolic (reduced ETC capacity) constraints
2. **Early anaerobic threshold:** Transition to glycolytic dominance at lower workloads because oxidative capacity ceiling is reduced
3. **Day-2 decrement:** The model predicts that ROS-mediated damage from day-1 exertion reduces ETC capacity by 5–15% on day 2, lowering peak VO_2 and anaerobic threshold
4. **Prolonged recovery:** The model predicts that post-exercise ATP and cytokine levels require 48–96 hours to return to pre-exercise baseline, versus < 24 hours in healthy controls

39.4.2 Cognitive Exertion

Cognitive exertion increases brain ATP demand by 10–20% above baseline (versus 5–10-fold increases for intense physical exercise in skeletal muscle). Despite the smaller absolute demand, the brain is particularly vulnerable because it has minimal energy reserves and depends on continuous aerobic metabolism. The model predicts that cognitive PEM is mediated by the same energy depletion mechanism as physical PEM but manifests at lower total energy expenditure because brain tissue is operating closer to its metabolic ceiling.

39.4.3 Infections

Intercurrent infections are the most common trigger for ME/CFS relapses. The model simulates infection as a combined perturbation: increased $V(t)$, increased immune activation (N_a , T_e , cytokines), and increased energy demand (J_{immune}). In the ME/CFS model, the immune response is both delayed (slower NK cell and T cell activation due to exhaustion and energy limitation) and prolonged (impaired viral clearance extends the infectious period), resulting in greater cumulative damage. The model predicts that the severity of relapse depends on the product of infection intensity and duration, suggesting that early antiviral intervention could reduce relapse severity.

39.4.4 Environmental Factors

Temperature extremes, chemical exposures, and sensory overload are reported ME/CFS triggers. The model represents these as perturbations to specific subsystems: temperature extremes alter metabolic rate and autonomic demand; chemical exposures (volatile organic compounds, fragrances) can trigger mast cell activation and neuroinflammation [665]; sensory overload increases CNS energy demand and sympathetic activation. The common pathway is an increase in total energy demand J_{demand} above the reduced energy envelope, triggering the PEM cascade.

39.5 Treatment Response Modeling

39.5.1 Time Course of Improvement

Treatment responses in ME/CFS are characteristically slow, with meaningful improvement often requiring weeks to months. The model explains this through the timescales of parameter recovery:

1. **Fast** (hours to days): Direct pharmacological effects—e.g., symptom relief from analgesics, blood pressure support from fludrocortisone, heart rate reduction from beta-blockers. These modify J_{demand} or compensatory variables without addressing underlying dysfunction.
2. **Medium** (weeks to months): Repair of accumulated damage—mitochondrial biogenesis ($\tau \sim 2\text{--}4$ weeks), immune system remodeling ($\tau \sim 4\text{--}8$ weeks), neuroplastic changes ($\tau \sim 6\text{--}12$ weeks). These modify the damage variable D_{total} , gradually improving system capacity.
3. **Slow** (months to years): Epigenetic reprogramming, immune cell lineage reconstitution (after B cell depletion, new naïve B cells repopulate over 6–12 months [616]), and neuroendocrine axis resetting.

The model predicts that treatments targeting fast timescale effects (symptom management) provide immediate but limited benefit, while treatments targeting medium and slow timescale processes (disease modification) show delayed onset but potentially greater magnitude of improvement.

39.5.2 Combination Therapy Effects

The integrated model enables simulation of multi-target interventions. Combination therapy can produce synergistic effects when interventions target complementary pathways. For example, the model predicts that combining CoQ10 (increasing ETC capacity) with an anti-inflammatory agent (reducing J_{immune}) produces greater ATP surplus than either alone, because the energy freed from reduced immune demand is processed more efficiently through the enhanced ETC. Formally, synergy occurs when:

$$\Delta[\text{ATP}]_{\text{A+B}} > \Delta[\text{ATP}]_{\text{A}} + \Delta[\text{ATP}]_{\text{B}} \quad (39.9)$$

The model identifies the parameter combinations most likely to produce synergy, providing a rational basis for combination therapy design rather than empirical trial-and-error.

39.6 Long-Term Trajectories

Long-term outcomes in ME/CFS are variable. Meta-analyses suggest that full recovery occurs in approximately 5% of patients, while 20–30% show some improvement, and the remainder remain stable or worsen [113]. The model reproduces these statistics through patient-to-patient variation in parameters. Monte Carlo simulation with parameter distributions derived from population data generates a distribution of trajectories:

1. **Recovery** (~5%): Patients whose parameters place them near the separatrix boundary, where spontaneous fluctuations or modest interventions can push the system back to the healthy attractor
2. **Improvement** (~25%): Patients where damage accumulation is slow (k_{damage} low) and repair is sufficient to gradually reduce D_{total} , especially with effective pacing
3. **Stable** (~50%): Patients at approximate damage–repair equilibrium, whose severity depends on management quality
4. **Progressive** (~20%): Patients where damage outpaces repair, often due to severe energy deficit limiting repair capacity or repeated perturbations (infections, overexertion) preventing equilibrium

⚠ Limitation 2: Long-Term Prediction Uncertainty

Long-term trajectory predictions are highly sensitive to parameter values and perturbation history, both of which are poorly characterized for individual patients. The trajectory distribution above should be interpreted as a population-level statistical prediction, not as a prognosis for any individual patient. Personalized trajectory prediction requires longitudinal data collection over months to years with sufficient biomarker resolution to constrain patient-specific parameters—a capability not yet available in clinical practice.

39.7 Critical Slowing Down and Early Warning Signals

Near bifurcation points—the tipping points where the system transitions between attractors—dynamical systems exhibit a universal phenomenon called *critical slowing down*: the rate at which the system returns to equilibrium after small perturbations decreases as the bifurcation is approached. This phenomenon, well-established in ecology, climate science, and financial systems, has direct clinical applications for ME/CFS.

39.7.1 Mathematical Basis

The recovery rate from perturbation is determined by the dominant eigenvalue λ_1 of the Jacobian matrix $J = \partial f / \partial x$ evaluated at the current steady state. At a saddle-node bifurcation (where the disease attractor is created or destroyed), $\lambda_1 \rightarrow 0$. As the system approaches the bifurcation—whether through parameter drift (disease progression) or state-space proximity

to the separatrix (accumulation of perturbations)—the recovery time $\tau_{\text{recovery}} = -1/\text{Re}(\lambda_1)$ diverges.

This produces measurable early warning signals that are a direct and unique consequence of the mathematical model:

1. **Increased autocorrelation:** As $\lambda_1 \rightarrow 0$, the autocorrelation of physiological time series (HRV, activity level, symptom scores) at lag Δt increases: $\rho(\Delta t) = e^{\lambda_1 \Delta t} \rightarrow 1$. The system “remembers” perturbations longer because it returns to baseline more slowly.
2. **Increased variance:** Fluctuations grow because the restoring force weakens: $\text{Var}(x) \propto \sigma^2/(2|\lambda_1|) \rightarrow \infty$, where σ^2 is the noise intensity.
3. **Flickering:** Near the bifurcation, noise-driven excursions occasionally reach the alternative attractor, producing brief “flickers” of the alternative state before the system returns. In ME/CFS, this manifests as intermittent symptom spikes of unusual character or duration.
4. **Asymmetric recovery:** Recovery from positive perturbations (toward the separatrix) slows more than recovery from negative perturbations (away from the separatrix), producing a skewed perturbation response.

39.7.2 Clinical Application: Crash Prediction

The early warning signals above are, in principle, detectable from wearable sensor data (continuous heart rate, accelerometry, skin conductance) without requiring blood draws or clinic visits. The model predicts that **24–48 hours before a PEM crash, patients near the energy envelope threshold should show:**

- Rising autocorrelation in resting heart rate (HR recovery after standing or minor exertion slows)
- Increasing variance in step count or activity level (more erratic activity patterns)
- Reduced HRV (reflecting the narrowing margin between current state and the PEM threshold)
- Elevated resting HR trend (sympathetic compensation for shrinking energy reserve)

This prediction is uniquely generated by the mathematical model: without the bifurcation framework, there is no theoretical basis for expecting these specific statistical signatures to precede crashes, nor for specifying the 24–48 hour timescale (which derives from the eigenvalue dynamics near the energy envelope threshold). A smartphone application implementing these detectors could warn patients to reduce activity before the crash threshold is crossed, potentially preventing PEM episodes. Retrospective validation against existing wearable datasets with crash annotations would constitute a strong test of the model.

39.7.3 Recovery Proximity Detection

The same early warning signals operate in reverse near recovery transitions. A patient approaching the separatrix from the disease side (improving toward recovery) should show

increasing autocorrelation and variance in the *disease-state* variables—paradoxically appearing less stable just before recovery. The model predicts that this “recovery instability” is a positive prognostic sign: the system is losing commitment to the disease attractor. Clinicians and patients aware of this prediction would interpret transient symptom instability during recovery differently—as a sign of approaching transition rather than deterioration.

39.8 Hysteresis and the Intervention Window

Hysteresis occurs when the path from health to disease differs from the path from disease to health. In the ME/CFS model, hysteresis arises from two sources: (1) the saddle-node bifurcation structure of the energy-immune system (structural hysteresis), and (2) the epigenetic consolidation of disease parameters (parametric hysteresis, Section 38.7.4).

39.8.1 Structural Hysteresis

In a bistable system, the disease attractor appears at a critical parameter value θ_{onset} (saddle-node bifurcation) but does not disappear until a different value $\theta_{\text{recovery}} \neq \theta_{\text{onset}}$. The hysteresis width $\Delta\theta = |\theta_{\text{recovery}} - \theta_{\text{onset}}|$ quantifies how much further the system must be pushed for recovery compared to onset. For the energy-immune subsystem, the model predicts:

$$\Delta\alpha_{\text{CI}} = \alpha_{\text{CI}}^{\text{recovery}} - \alpha_{\text{CI}}^{\text{onset}} > 0 \quad (39.10)$$

meaning that Complex I activity must be restored to a level *above* the value at which disease was triggered, not merely to the trigger value. **This asymmetry is invisible without the bifurcation diagram:** verbal reasoning about positive feedback loops identifies that the disease is self-sustaining but cannot quantify how much intervention overshoot is required for recovery. The model computes $\Delta\alpha_{\text{CI}}$ as a function of the feedback strengths, providing a quantitative recovery target.

39.8.2 Parametric Hysteresis and the Intervention Window

Epigenetic consolidation (Section 38.7.4) adds a second layer of hysteresis that increases over time. The methylation index $\mathcal{M}$ evolves toward values that stabilize the disease attractor, progressively widening the hysteresis loop. This predicts an **intervention window**: the period after disease onset during which the epigenetic contribution to hysteresis is still small, and recovery requires only modest intervention above the structural hysteresis threshold.

The model estimates the intervention window duration as the time required for $\mathcal{M}$ to reach half its equilibrium disease value:

$$\tau_{\text{window}} \approx \frac{\ln 2}{k_{\text{DNMT}} \cdot \bar{C}_{\text{pro}}} \quad (39.11)$$

where $\bar{C}_{\text{pro}}$ is the mean pro-inflammatory cytokine level in the disease state. For typical parameter values, $\tau_{\text{window}} \sim 3\text{--}12$ months, consistent with clinical observations that early intervention (within the first year of ME/CFS) is associated with better outcomes. After the intervention window closes, recovery requires both sufficient perturbation to cross the structural separatrix *and* sustained intervention to reverse epigenetic changes—a process operating on the slow timescale τ_{epi} .

This intervention window concept is a unique product of the coupled dynamical model: it emerges from the interaction of two timescales (fast state dynamics and slow epigenetic dynamics) that cannot be analyzed without formal mathematical treatment. It provides a quantitative basis for the clinical urgency of early diagnosis and treatment in ME/CFS.

39.9 Endogenous Oscillations and Hopf Bifurcation

The open question posed in Chapter 38—whether the coupled model produces endogenous oscillations—can be addressed through Hopf bifurcation analysis. A Hopf bifurcation occurs when a pair of complex conjugate eigenvalues of the Jacobian crosses the imaginary axis, destabilizing a fixed point and creating a limit cycle (periodic orbit).

39.9.1 Conditions for Oscillation

The ME/CFS model contains the structural requirements for Hopf bifurcation: positive feedback loops (energy–immune, mast cell–energy, coagulation–oxygenation, SIBO–immune–motility) combined with delayed negative feedback (cortisol suppression of inflammation, IL-10 anti-inflammatory signaling, sleep-mediated repair). The imaginary part of the critical eigenvalue pair determines the oscillation period:

$$T_{\text{osc}} = \frac{2\pi}{|\text{Im}(\lambda_{\text{crit}})|} = \frac{2\pi}{\omega_0} \quad (39.12)$$

Preliminary analysis of the reduced energy–immune–HPA subsystem (6 variables: [ATP], N_a , C_{pro} , F , [ROS], D) identifies a Hopf bifurcation when the cortisol feedback gain n_F exceeds a critical value while the energy–immune coupling strength is in the intermediate range. The predicted oscillation period is:

$$T_{\text{osc}} \approx 2\pi \sqrt{\frac{\tau_{\text{cortisol}} \cdot \tau_{\text{immune}}}{\text{gain}_{\text{loop}}}} \sim 2\text{--}6 \text{ weeks} \quad (39.13)$$

where $\tau_{\text{cortisol}} \sim 1$ day (HPA axis response), $\tau_{\text{immune}} \sim 1\text{--}2$ weeks (cytokine network re-modeling), and $\text{gain}_{\text{loop}}$ is the total loop gain. This 2–6 week period is consistent with the relapse–remission cycling reported by many ME/CFS patients who describe “good weeks and bad weeks” without identifiable external triggers.

39.9.2 Oscillation as a Biomarker

If endogenous oscillations exist, the oscillation period itself becomes a model-derived biomarker:

- **Shorter period** indicates higher loop gain (more active disease dynamics)—symptom fluctuations are rapid and intense
- **Longer period** indicates lower loop gain (attenuated feedback)—slow, gradual waxing and waning
- **Period changes over time** signal parameter drift: shortening period indicates approaching instability; lengthening period indicates the system is moving away from the Hopf bifurcation (toward stable disease or toward recovery)
- **Loss of periodicity** can signal either recovery (the oscillation amplitude shrinks to zero as the system stabilizes in the healthy attractor) or transition to chaos (if the system crosses a period-doubling cascade)

Detecting and characterizing these oscillations requires longitudinal symptom tracking with sufficient temporal resolution (daily or finer) over multiple oscillation periods (months). Wearable devices and electronic symptom diaries provide the data streams; spectral analysis (Fourier transform, wavelet analysis) provides the detection methods. The model predicts specific spectral signatures distinguishable from random fluctuation and from exogenous cycling (which would correlate with identifiable triggers).

39.9.3 Stochastic Resonance and Noise-Assisted Recovery

In bistable systems driven by noise, a phenomenon called *stochastic resonance* occurs: moderate noise can facilitate transitions between attractors that would be impossible in a deterministic system. Applied to ME/CFS, this predicts that random physiological fluctuations (minor infections, hormonal cycles, sleep variations) can occasionally push a patient from the disease attractor into the basin of attraction of the healthy state—but only if the patient is sufficiently close to the separatrix.

The Kramers escape rate from the disease attractor is:

$$r_{\text{escape}} = \frac{\omega_0 \omega_b}{2\pi} \exp\left(-\frac{2\Delta U}{\sigma^2}\right) \quad (39.14)$$

where ω_0 is the oscillation frequency at the attractor, ω_b is the curvature at the saddle point (separatrix), ΔU is the effective potential barrier height (distance from attractor to separatrix), and σ^2 is the noise intensity. **This equation produces three predictions unique to the mathematical framework:**

1. **Spontaneous recovery rate depends exponentially on disease severity:** patients deep in the disease attractor (high ΔU) have astronomically low escape probability, while those near the separatrix can recover spontaneously—consistent with the ~5% spontaneous recovery rate

2. **Optimal noise intensity exists:** too little noise → no escape; too much noise → the system is destabilized before reaching the healthy basin and falls back. This predicts that there is an optimal level of mild physiological perturbation that maximizes recovery probability—neither complete rest nor aggressive stimulation
3. **Hormonal cycling may assist recovery:** for female patients, the menstrual cycle provides periodic perturbations of fixed amplitude and frequency. If this perturbation amplitude is near optimal for stochastic resonance, it could explain potential sex differences in spontaneous recovery rates—a testable prediction

Speculation 79 (Therapeutic Perturbation Protocols). If stochastic resonance operates in ME/CFS, carefully calibrated periodic perturbations might facilitate recovery in patients near the separatrix. A “perturbation protocol” would involve controlled, mild physiological challenges (brief cold exposure, short exercise below anaerobic threshold, intermittent fasting) at frequencies tuned to the system’s natural oscillation period. The model predicts that the optimal perturbation amplitude is proportional to $\sqrt{\sigma_{\text{optimal}}^2} \propto \sqrt{\Delta U / \text{gain}_{\text{loop}}}$ —patient-specific and calculable from model parameters. This hypothesis predicts that: (1) patients near the separatrix (identified by critical slowing down signals) would benefit most; (2) patients deep in the disease attractor would not respond; and (3) perturbation frequency matching the endogenous oscillation period T_{osc} would be most effective.

Certainty: 0.25. Stochastic resonance is well-established in physics and neuroscience but has not been demonstrated in disease dynamics. The practical challenges of estimating separatrix proximity and optimal perturbation parameters in individual patients are substantial. This concept should be considered a research direction, not a treatment recommendation.

39.10 Model Application Guide

This section consolidates practical guidance for each temporal model: what to measure, how to use the model, worked examples, inter-model dependencies, falsification criteria, and clinical implications.

39.10.1 Tipping Point and Disease Onset Models

Measurements required. Pre-infection baseline: mitochondrial respiratory capacity (Seahorse assay or muscle biopsy oximetry), hs-CRP, IL-6, morning cortisol. During acute infection: serial viral load (PCR), daily cytokine panel (TNF- α , IL-6, IL-10), daily fatigue VAS score. Post-infection: repeat baseline panel at 4, 8, and 12 weeks to determine whether the system has returned to the pre-infection attractor or settled at a new steady state.

How to use. Compute the basin of attraction boundary (separatrix) from the energy-immune bistability model (Chapter 38) using the patient’s pre-infection parameter values. Overlay the infection-induced state-space trajectory. If the trajectory crosses the separatrix, the model predicts ME/CFS onset; if it remains within the healthy basin, full recovery is expected. The

separatrix distance d_{sep} —computed as the Euclidean distance from the patient’s pre-infection state to the nearest separatrix point—serves as a risk score.

Worked example. A 35-year-old female presents with baseline Complex I activity $\alpha_{\text{CI}} = 0.75$ (mildly reduced), hs-CRP = 1.8 mg/L (low-grade inflammation), and morning cortisol = 8 $\mu\text{g}/\text{dL}$ (low-normal). The model places her separatrix distance at $d_{\text{sep}} = 0.12$ (low margin). An acute COVID-19 infection produces a state-space displacement of magnitude 0.18 along the energy–immune axes. Since $0.18 > 0.12$, the trajectory crosses the separatrix, and the model predicts transition to the disease attractor. Post-infection monitoring at week 8 confirms persistent fatigue (VAS > 6), elevated IL-6 (> 4 pg/mL), and reduced ATP synthesis (< 70% of pre-infection baseline), consistent with the disease steady state.

Inter-model dependencies. The tipping point model depends on the bistability analysis from Chapter 38 for the separatrix computation, on the viral dynamics model (Chapter 36) for infection trajectory magnitude, and on the energy–immune coupling model for the state-space geometry. It feeds into the damage accumulation model (Section 39.2.1) by setting initial conditions for the post-onset trajectory, and into the energy ratchet model (Section 39.2.3) by establishing the initial ceiling $B_{\text{max}}(t_0)$.

Falsification criteria. The tipping point model is falsified if: (1) patients with large separatrix distance ($d_{\text{sep}} > 0.3$) develop ME/CFS after mild infections at rates comparable to high-risk individuals; (2) patients with identical pre-infection parameters show no dose–response relationship between infection severity and ME/CFS onset probability; (3) the bistability model fails validation (Chapter 38), eliminating the separatrix entirely.

Clinical implications. The separatrix distance d_{sep} identifies individuals at elevated risk for post-infectious ME/CFS *before* infection occurs. High-risk individuals ($d_{\text{sep}} < 0.15$) could be prioritized for aggressive infection prevention (vaccination, antivirals), early intervention during acute illness (rest, anti-inflammatory support), and close post-infection monitoring. The model does not prescribe a specific treatment but identifies the *window* (during and immediately after acute infection) and the *target* (preventing the state-space trajectory from crossing the separatrix) for preventive intervention.

39.10.2 Damage Accumulation and Energy Ratchet Models

Measurements required. Longitudinal functional capacity: 6-minute walk test or step count (monthly), self-reported activity level (daily via wearable). Biomarkers of damage: 8-OHdG (oxidative DNA damage), cell-free mitochondrial DNA, neurofilament light chain (neuronal damage). Energy status: resting metabolic rate, lactate at standardized workload. Event log: dates and severity of infections, PEM crashes, surgeries, and major stressors (patient diary).

How to use. Calibrate the ratchet model (Equations 39.2–39.5) from the patient’s event history. Fit δ_0 , α , and $r(B)$ to the observed trajectory of functional capacity $B(t)$ over at least 3 damaging events. Once calibrated, simulate forward: given the patient’s current B and $B_{\max}$, predict how many additional events of a given severity can occur before the next severity threshold is crossed. This provides a quantitative basis for event prevention urgency.

Worked example. A patient diagnosed 3 years ago has experienced 5 documented infections and an estimated 20 PEM crashes. Functional capacity has declined from $B_0 = 0.72$ (mild) to $B = 0.48$ (approaching moderate threshold at $\theta_{\text{mod}} = 0.6$; current ceiling $B_{\max} = 0.55$). Fitting the ratchet model yields $\delta_0 = 0.025$, $\alpha = 0.8$ (moderate damage sensitisation). Forward simulation predicts that 3 more infections of similar severity will drop $B_{\max}$ below 0.35 (severe threshold) within 2 years, whereas strict event prevention (reducing infection rate by 80% through masking and antivirals) extends the time to severe threshold to > 8 years. The model quantifies the stakes of infection prevention for this specific patient.

Inter-model dependencies. The ratchet model takes initial conditions from the tipping point model (Section 39.1.2). The repair rate $r(B)$ depends on ATP availability from the energy metabolism models (Chapter 35). Damage sensitisation parameter α connects to immune exhaustion (Chapter 36) and microglial priming (Chapter 37). The continuous damage model (Equation 39.1) provides the inter-event drift dynamics when chronic ROS/inflammation is significant.

Falsification criteria. The ratchet model is falsified if: (1) patients show no correlation between event frequency and rate of functional decline over 2+ years of prospective tracking; (2) damage sensitisation ($\alpha > 0$) is not observed—i.e., later events produce the same or smaller ceiling losses as earlier ones; (3) functional capacity spontaneously exceeds $B_{\max}$ without identifiable disease-modifying intervention (this would violate the one-way ceiling constraint).

Clinical implications. The ratchet model provides the quantitative case for aggressive event prevention as disease-modifying therapy. For each patient, the model computes: (a) current ceiling $B_{\max}$ and remaining margin to the next severity threshold, (b) expected events per year given current lifestyle and infection exposure, (c) projected time to severity transition under current conditions vs. with enhanced prevention. This transforms the abstract advice “avoid crashes” into a concrete risk quantification. Patients with high α (strong damage sensitisation) and low remaining margin benefit most from aggressive prevention strategies.

39.10.3 Critical Slowing Down and Early Warning Signals

Measurements required. Continuous physiological time series from wearables: heart rate (beat-to-beat for HRV), accelerometry (step count, activity patterns), skin temperature. Daily

symptom diary: fatigue, cognitive function, pain (0–10 scales). Minimum monitoring duration: 4 weeks for autocorrelation estimation, 3+ months for oscillation period detection. Sampling: heart rate at ≥ 1 Hz; activity at 1-minute epochs; symptoms at least daily.

How to use. Compute rolling-window statistics from the wearable time series: (1) lag-1 autocorrelation ρ_1 of detrended resting heart rate (window = 7 days, step = 1 day); (2) variance of daily step count (window = 14 days); (3) spectral power ratio (low-frequency to high-frequency HRV). Rising ρ_1 and variance signal approach to a bifurcation (crash or severity transition). Declining ρ_1 and variance during recovery signal retreat from the bifurcation. Threshold alerts: if ρ_1 exceeds 0.85 (up from a patient-specific baseline of ~ 0.5 – 0.7), issue a crash warning.

Worked example. A patient wears a heart rate monitor and logs daily symptoms for 8 weeks. During weeks 1–5, resting HR autocorrelation is $\rho_1 = 0.62 \pm 0.05$ and step-count variance is $1200 \pm 200 \text{ steps}^2/\text{day}^2$. During week 6, ρ_1 rises to 0.78 and variance increases to $2100 \text{ steps}^2/\text{day}^2$. The model flags this as an early warning signal. The patient reduces activity by 30%. By week 7, ρ_1 returns to 0.65 and no PEM crash occurs. Without the warning, the patient’s planned weekend activity would likely have exceeded the energy envelope, triggering a multi-day crash.

Inter-model dependencies. Early warning signal detection requires the bifurcation structure computed in Chapter 38 (which eigenvalues approach zero, and at what parameter values). The autocorrelation predictions derive from the Jacobian eigenvalues of the full integrated model. Crash prediction feeds into the energy ratchet model: each prevented crash is a prevented ratchet step, directly slowing disease progression.

Falsification criteria. The early warning signal model is falsified if: (1) prospective monitoring shows no increase in autocorrelation or variance in the 24–72 hours before documented PEM crashes (tested in ≥ 20 patients over ≥ 50 crash events); (2) the statistical signatures are indistinguishable from random fluctuation (receiver operating characteristic AUC < 0.6 for crash prediction); (3) the predicted 24–48 hour warning timescale is wrong by more than an order of magnitude.

Clinical implications. Early warning signals enable *preventive pacing*: reducing activity before rather than after a crash. The model predicts that this is disease-modifying (each prevented crash preserves the ratchet ceiling B_{max}). Implementation requires: (a) a wearable device with continuous HR monitoring, (b) software computing rolling autocorrelation and variance, (c) patient-facing alerts calibrated to individual baselines. The clinical trial design is straightforward: randomize crash-prone patients to alert vs. no-alert groups and compare PEM frequency, severity, and functional trajectory over 6–12 months.

39.10.4 Hysteresis and Intervention Window

Measurements required. Disease duration (time since onset). Epigenetic markers: global DNA methylation (LINE-1 assay), gene-specific methylation at ME/CFS-associated loci (e.g., glucocorticoid receptor NR3C1, immune regulatory genes [76]). Cytokine levels for estimating methylation rate parameter $k_{\text{DNMT}} \cdot \bar{C}_{\text{pro}}$. Functional capacity trajectory (is the patient stable, improving, or worsening?).

How to use. Estimate the intervention window τ_{window} from Equation 39.11 using the patient's mean pro-inflammatory cytokine level. Compare disease duration to τ_{window} : if duration $< \tau_{\text{window}}$, the patient is within the intervention window and recovery requires only crossing the structural hysteresis gap $\Delta\alpha_{\text{CI}}$. If duration $\gg \tau_{\text{window}}$, epigenetic consolidation has widened the hysteresis loop, and treatment must address both state-space displacement and epigenetic reprogramming.

Worked example. A patient develops ME/CFS after EBV reactivation. At month 4 post-onset, mean IL-6 is 6 pg/mL. Using $k_{\text{DNMT}} = 0.015 \text{ month}^{-1}$ per pg/mL, the model estimates $\tau_{\text{window}} = \ln 2 / (0.015 \times 6) \approx 7.7$ months. The patient is within the intervention window. The structural hysteresis width $\Delta\alpha_{\text{CI}} = 0.08$ means Complex I activity must be restored 8% above onset threshold. A mitochondrial support protocol (CoQ10 300 mg/day, D-ribose, pacing) is initiated. At month 3 of treatment (month 7 post-onset, still within the window), the patient shows meaningful improvement with ATP synthesis rising 12% from nadir. Had treatment been delayed to month 18 post-onset ($> 2\tau_{\text{window}}$), the model predicts that epigenetic methylation would have approximately doubled the effective hysteresis width, requiring more aggressive and prolonged intervention for equivalent improvement.

Inter-model dependencies. The hysteresis model requires the bifurcation diagram from Chapter 38 for structural hysteresis width, the epigenetic dynamics model (Section 38.7.4 in Chapter 38) for parametric hysteresis evolution, and cytokine data from the immune models (Chapter 36) for methylation rate estimation. It informs treatment response modeling (Section 39.5) by predicting treatment magnitude requirements as a function of disease duration.

Falsification criteria. The intervention window model is falsified if: (1) treatment response does not correlate with disease duration—i.e., patients treated at 5 years respond as well as those treated at 5 months, controlling for severity; (2) ME/CFS-associated epigenetic changes do not show progressive consolidation over the first 1–2 years (measured in a longitudinal methylation study); (3) the predicted window duration ($\tau_{\text{window}} \sim 3\text{--}12$ months) is inconsistent with clinical data on early vs. late treatment efficacy.

Clinical implications. The intervention window model provides the strongest model-derived argument for early diagnosis and treatment. It predicts that each month of delay narrows the treatment response, not because the patient is “getting used to being sick” but because epigenetic consolidation physically widens the barrier to recovery. For clinicians:

prioritize early, aggressive treatment within the first year post-onset. For health systems: reduce diagnostic delay [403] as a disease-modifying intervention at the population level. For patients: the model quantifies why “wait and see” is not a neutral strategy.

39.10.5 Oscillation Analysis and Stochastic Resonance

Measurements required. Daily symptom scores (fatigue, pain, cognitive function) for ≥ 3 months (to capture multiple oscillation periods). Weekly cytokine panel (TNF- α , IL-6, IL-10) and morning cortisol for at least 8 weeks. Continuous wearable data (HR, activity) for spectral analysis. Menstrual cycle tracking for female patients (to separate endogenous oscillations from hormonal cycling).

How to use. Apply spectral analysis (Lomb–Scargle periodogram for unevenly sampled data, wavelet transform for non-stationary signals) to the symptom and biomarker time series. Identify dominant spectral peaks and compare their periods to the model prediction of $T_{\text{osc}} \approx 2\text{--}6$ weeks. If a peak is present, estimate loop gain from the period (Equation 39.13). Track period changes over time to assess disease trajectory: shortening period signals increasing instability; lengthening period signals stabilisation.

Worked example. A patient records daily fatigue scores for 6 months. Lomb–Scargle analysis reveals a spectral peak at period $T = 3.2$ weeks ($p < 0.01$ vs. red noise null). Using Equation 39.13 with $\tau_{\text{cortisol}} = 1$ day and $\tau_{\text{immune}} = 10$ days, the estimated loop gain is $\text{gain}_{\text{loop}} \approx 4\pi^2 \tau_{\text{cortisol}} \tau_{\text{immune}} / T^2 \approx 0.78$. After 3 months of LDN therapy (which modulates immune feedback gain), repeat analysis shows the period has lengthened to $T = 4.8$ weeks, corresponding to reduced loop gain of 0.35—a quantitative measure of treatment effect on the underlying dynamical structure.

Inter-model dependencies. The oscillation analysis requires the Jacobian eigenvalues from the full integrated model (Chapter 38), cortisol dynamics from the HPA axis model (Chapter 37), and immune feedback timescales from the cytokine network model (Chapter 36). The stochastic resonance analysis additionally requires the bistability model for attractor depth ΔU and the noise characterisation from daily physiological variability.

Falsification criteria. The Hopf bifurcation oscillation model is falsified if: (1) longitudinal symptom tracking in ≥ 50 ME/CFS patients (daily data, ≥ 6 months) reveals no spectral peaks in the 2–6 week range above red noise; (2) observed oscillations correlate entirely with identifiable exogenous triggers (infections, menstrual cycle, activity patterns) with no residual endogenous component; (3) treatments that reduce immune feedback gain (as measured by cytokine response to standardised challenge) do not alter oscillation period. The stochastic resonance prediction is falsified if spontaneous recovery rates show no relationship to disease severity (attractor depth) after controlling for disease duration.

Clinical implications. If validated, oscillation period becomes a non-invasive biomarker of disease dynamics obtainable from symptom diaries and wearables alone. Clinicians could track treatment efficacy through period changes rather than waiting months for functional capacity improvement. The stochastic resonance concept, while speculative, suggests that some degree of controlled physiological variability (not strict rest, not aggressive exercise) may be optimal for patients near the recovery threshold—providing theoretical support for graded activity approaches *only for patients near the separatrix*, while predicting harm for patients deep in the disease attractor. This patient-specific prediction distinguishes the model-based approach from one-size-fits-all exercise recommendations.

39.10.6 Scope and Rationale

The temporal models in this chapter were selected to capture phenomena that the instantaneous models of Chapters 35–38 cannot explain: why some individuals develop ME/CFS after infection while others recover (tipping point dynamics), why disease severity changes over months to years (damage accumulation and ratchet), why symptoms fluctuate on multi-week timescales (endogenous oscillations), why early treatment produces better outcomes (intervention window and hysteresis), and why sudden transitions occur between severity levels (bifurcation analysis). These phenomena are reported consistently in patient surveys and clinical studies but have lacked formal mechanistic explanations.

The chapter does not model *every* temporal phenomenon in ME/CFS. Notably absent are: intra-day energy fluctuations beyond the circadian model (which would require coupling to meal timing, thermoregulation, and postural changes at minute-to-hour timescales), the specific dynamics of immune reconstitution after B-cell depletion therapy (which would require a dedicated lymphocyte population model at finer resolution than Chapter 36 provides), and the multi-generational epigenetic transmission of ME/CFS risk (which would require a population genetics framework). These are identified as directions for future model development where clinical data become available to constrain them.

40 Predictive Applications and Clinical Translation

The mathematical models developed in this part are not ends in themselves but tools for improving clinical care. This chapter examines how the models can be translated into practical applications: personalized pacing optimization, treatment selection, prognosis prediction, clinical decision support, and drug development. Each application is assessed for its current feasibility given available data and technology, with explicit identification of remaining barriers to clinical implementation.




Figure 40.1: Clinical translation pipeline from patient data to treatment recommendations. Patient biomarkers (Stage 1) are used to estimate individual model parameters (Stage 2), which enable simulation of candidate interventions (Stage 3) and generation of personalized treatment recommendations (Stage 4). The implementation roadmap (bottom) shows the staged path from current research tools to validated clinical instruments.

40.1 Personalized Pacing Optimization

Pacing—the deliberate management of activity to remain within the energy envelope—is the most widely recommended management strategy for ME/CFS [61, 62]. Current pacing advice is qualitative: “listen to your body,” “rest before you feel tired.” The energy envelope model (Section 35.3.3) enables a quantitative approach.

40.1.1 Individual Energy Budget Calculation

The energy budget E_{budget} (Equation 35.12) is patient-specific, depending on mitochondrial capacity ($J_{\text{production,max}}$), baseline metabolic demands (E_{basal}), and repair costs (E_{repair}). In principle, these can be estimated from clinical data:

- $J_{\text{production,max}}$: estimated from peak VO_2 on CPET, which directly measures maximal oxidative capacity [1]
- E_{basal} : estimated from resting metabolic rate (indirect calorimetry)
- E_{repair} : estimated indirectly from inflammatory biomarker levels (higher inflammation implies greater repair costs)

The resulting E_{budget} defines the maximum daily activity expenditure that avoids PEM. Converting this to practical units (e.g., steps, minutes of activity at specified intensity) requires calibration against wearable device data. The model predicts that E_{budget} is not constant but varies with disease state, sleep quality, immune status, and ambient conditions—explaining why a “safe” activity level on one day can trigger PEM on another.

40.1.2 Activity Planning Algorithms

Given $E_{\text{budget}}(t)$ and a set of planned activities with estimated energy costs, an optimization algorithm can schedule activities to maximize function while respecting the energy constraint:

$$\max_{\phi(t)} \int_0^T U(\phi(t)) dt \quad \text{subject to} \quad \int_0^T J_{\text{demand}}(\phi(t)) dt \leq \int_0^T E_{\text{budget}}(t) dt \quad (40.1)$$

where $U(\phi(t))$ is a utility function reflecting the value of activities to the patient and T is the planning horizon (typically one day). The constraint is the energy envelope. Additional constraints can encode PEM dynamics: not only must total daily expenditure stay below E_{budget} , but peak instantaneous demand must stay below $J_{\text{production,max}}$ to avoid acute ATP depletion, and recovery periods of minimum duration must separate activity bouts. This optimization problem is a constrained scheduling problem solvable with standard methods (linear programming for simplified versions, dynamic programming for time-dependent constraints).

40.1.3 Real-Time Monitoring Integration

Wearable devices (heart rate monitors, accelerometers, continuous glucose monitors) can provide real-time estimates of energy expenditure, enabling dynamic updating of the remaining daily energy budget. The model-based approach adds value beyond simple step counting by accounting for the nonlinear relationship between activity intensity and energy cost: moderate activity for extended duration may cost less total energy than brief intense exertion, because the latter triggers disproportionate ROS production and immune activation.

Limitation 1: Pacing Model Calibration

Personalized pacing optimization requires individual calibration of E_{budget} and its time-varying components. Current clinical practice lacks the routine measurements needed for this calibration (CPET is not widely available, resting metabolic rate is rarely measured). Until these measurements become routine, the pacing model serves as a conceptual framework rather than a clinical tool.

40.2 Treatment Selection and Optimization

40.2.1 Biomarker-Based Treatment Prediction

The integrated model (Chapter 38) maps from biological parameters to treatment response. If a patient's parameter values can be estimated from biomarker measurements, the model predicts which treatments will be most effective for that individual. The logic proceeds in three steps:

1. **Parameter estimation:** Fit the model to the patient's available data (metabolomics, cytokine panel, CPET, HRV, cortisol rhythm) to estimate the patient-specific parameter vector $\hat{\theta}_{\text{patient}}$
2. **Intervention simulation:** For each candidate treatment, modify the appropriate parameters (e.g., CoQ10 increases $[\text{UQ}]_{\text{total}}$, LDN modifies cytokine production rates) and simulate the model forward to predict symptom trajectories
3. **Treatment ranking:** Rank treatments by predicted symptom improvement, accounting for parameter uncertainty through Bayesian credible intervals

This approach would enable rational treatment selection based on the patient's dominant pathophysiological mechanism. A patient with primarily metabolic dysfunction (low α_{CI} , near-normal immune markers) would be directed toward mitochondrial support, while a patient with dominant immune activation (elevated cytokines, near-normal CPET) would be directed toward immunomodulation.

40.2.2 Dosing Optimization

For a given treatment, the model can predict the optimal dose by simulating dose–response relationships. The pharmacokinetic–pharmacodynamic (PK–PD) interface links drug plasma concentration $C_{\text{drug}}(t)$ (from standard PK models) to parameter modifications in the pathophysiology model. For example, CoQ10 supplementation modifies $[\text{UQ}]_{\text{total}}$ as:

$$[\text{UQ}]_{\text{total}}(t) = [\text{UQ}]_{\text{baseline}} + \Delta[\text{UQ}]_{\text{max}} \cdot \frac{C_{\text{CoQ10}}(t)}{K_{\text{CoQ10}} + C_{\text{CoQ10}}(t)} \quad (40.2)$$

where $C_{\text{CoQ10}}(t)$ follows oral absorption and distribution kinetics. The model predicts the steady-state ATP improvement as a function of daily dose, identifying the dose above which

additional supplementation provides negligible benefit (saturation) and below which the effect is indistinguishable from placebo.

40.2.3 Combination Therapy Design

The synergy analysis from Chapter 39 (Equation 39.9) extends to systematic screening of treatment combinations. For n candidate treatments, the model can simulate all $\binom{n}{2}$ pairwise combinations and rank them by predicted synergy. Combinations targeting independent rate-limiting steps (e.g., ETC capacity + substrate supply + immune demand reduction) are predicted to show the greatest synergy, while combinations targeting the same bottleneck show diminishing returns.

40.3 Prognosis Prediction

Predicting disease trajectory is among the most clinically valuable applications of the model. Patients and clinicians need to know whether the disease is likely to improve, remain stable, or worsen. The damage accumulation model (Equation 39.1) provides a framework: the sign and magnitude of dD_{total}/dt at presentation, estimated from serial biomarker measurements over 3–6 months, predicts the trajectory.

Risk stratification assigns patients to prognostic categories based on model-derived indices:

1. **Repair-dominant** ($dD/dt < 0$): favorable prognosis; prioritize activity management to sustain the repair advantage
2. **Equilibrium** ($dD/dt \approx 0$): stable prognosis; focus on preventing perturbations (infections, overexertion) that could tip the balance toward damage dominance
3. **Damage-dominant** ($dD/dt > 0$): unfavorable prognosis; aggressive intervention warranted to reduce damage rate (anti-inflammatory, antioxidant) and increase repair capacity (metabolic support)

△ Warning 1: Prognosis Prediction Is Not Prognosis Determination

Model-based prognosis predictions are probabilistic estimates, not deterministic outcomes. Individual patient trajectories are influenced by unpredictable events (infections, life stressors), treatment decisions, and biological variability not captured in the model. Prognosis predictions should be communicated to patients as ranges of possibility, not as fixed outcomes, and should always be accompanied by the caveat that effective management can improve prognosis relative to the model's prediction.

40.4 Clinical Decision Support

Translation of models into clinical practice requires decision support tools that present model outputs in clinically actionable formats.

40.4.1 Dashboard Concept

A model-based clinical dashboard would display:

1. **Current state assessment:** estimated positions along the energy, immune, neuroendocrine, and autonomic axes, derived from the most recent biomarker panel
2. **Trajectory projection:** predicted 3–6 month trajectory under current management, with uncertainty bounds
3. **Treatment comparison:** side-by-side predicted outcomes for candidate interventions
4. **Alert indicators:** flags for approaching thresholds (e.g., damage rate accelerating, energy budget shrinking, immune markers trending upward)

Such a dashboard requires automated model fitting, fast simulation, and validated uncertainty quantification—capabilities that exist in related fields (pharmacometrics, systems biology) but have not yet been adapted for ME/CFS.

40.4.2 Treatment Algorithms

Model-derived treatment algorithms formalize clinical reasoning as decision trees parameterized by biomarker values. A simplified example:

1. If $\alpha_{CI} < 0.6$ (severe metabolic impairment): prioritize mitochondrial support (CoQ10, NAD⁺ precursors, d-ribose)
2. If $C_{pro} > 2 \times$ healthy reference: prioritize immunomodulation (LDN, anti-inflammatory agents)
3. If $HR_{standing} - HR_{supine} > 30$ bpm: prioritize autonomic support (volume expansion, compression, pyridostigmine [558])
4. If multiple criteria met: combination therapy targeting dominant pathway first, adding sequential interventions with 4–6 week monitoring intervals

These algorithms are model-informed but require clinical judgment for implementation. They do not replace physician decision-making but provide a structured framework for the complex multi-system reasoning that ME/CFS management demands.

40.5 Drug Development Applications

40.5.1 Target Identification

Sensitivity analysis (Equation 34.6) identifies the model parameters whose modification produces the greatest improvement in the symptom generation functions (Equation 38.17). Parameters with high sensitivity indices are candidate drug targets. The integrated model predicts that the highest-sensitivity parameters are:

1. α_{CI} (Complex I activity): directly limits ATP production capacity

2. k_{exh} (immune cell exhaustion rate): controls the size of the exhausted immune cell compartment
3. n_F (cortisol feedback sensitivity): determines HPA axis set point
4. P_0 (baseline BBB permeability): gates CNS exposure to peripheral inflammation

These targets correspond to distinct pharmacological classes: mitochondrial protectants, immune checkpoint modulators, glucocorticoid receptor modulators, and BBB-stabilizing agents. Some of these are already under investigation in ME/CFS (e.g., mitochondrial support supplements), while others (checkpoint modulators, BBB agents) represent novel directions suggested by the model.

40.5.2 Virtual Clinical Trials

The model enables *in silico* clinical trials: simulating the response of a virtual patient population to a candidate treatment. A virtual population is generated by sampling parameter vectors from distributions representing the known heterogeneity of ME/CFS. Each virtual patient is simulated under treatment and control conditions, generating predicted effect sizes, responder rates, and optimal trial designs (duration, endpoints, enrichment criteria).

Virtual trials are particularly valuable for ME/CFS because:

- Real clinical trials are expensive and slow to recruit
- The heterogeneity of ME/CFS means that average treatment effects may be small even when a subgroup responds dramatically—virtual trials can identify the responding subgroup and design enrichment strategies
- Ethical constraints limit the types of perturbation studies possible in patients—the model permits simulation of interventions that would be impractical to test in humans

? Open Question 1: Digital Twins for ME/CFS

Could a patient-specific computational model—a “digital twin”—be maintained and updated throughout the course of illness, receiving real-time data from wearable sensors and periodic biomarker panels? Such a system would continuously refine its parameter estimates and predictions, enabling adaptive pacing, early detection of relapses, and prospective treatment optimization. The technical requirements are substantial (real-time model fitting, sensor integration, uncertainty quantification), but analogous approaches are under development for cardiac care and diabetes management. The feasibility for ME/CFS depends on whether the biological signals accessible through non-invasive monitoring contain sufficient information to constrain the model’s critical parameters.

40.6 Optimal Control Theory for Pacing

The pacing optimization in Equation 40.1 treats energy expenditure as a static budget constraint. Optimal control theory provides a fundamentally more powerful formulation by

treating the patient's physiological state as a dynamical system with activity as the control input, enabling optimization over trajectories rather than budgets.

40.6.1 Pontryagin Maximum Principle Formulation

Define the state vector $\mathbf{x}(t) = (E(t), D(t), I(t), F(t))^T$ representing energy reserves, cumulative damage, immune activation, and fatigue perception respectively, with activity intensity $u(t) \in [0, u_{\max}]$ as the control variable. The system dynamics follow from the models in Chapters 35–38:

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, u) = \begin{pmatrix} J_{\text{production}}(\mathbf{x}) - J_{\text{demand}}(u) - E_{\text{repair}}(\mathbf{x}) \\ k_{\text{damage}} \cdot \text{ROS}(u, \mathbf{x}) - k_{\text{repair}} \cdot R(\mathbf{x}) \\ \sigma(\mathbf{x}, u) - \delta_I \cdot I \\ \alpha_F \cdot (D + \beta_F \cdot I) - \gamma_F \cdot F \end{pmatrix} \quad (40.3)$$

The patient seeks to maximize total functional capacity over a planning horizon $[0, T]$ while penalizing damage accumulation:

$$J[u] = \int_0^T \left[U(u(t)) - \lambda_D \cdot D(t) - \lambda_F \cdot F(t) \right] dt + \Phi(\mathbf{x}(T)) \quad (40.4)$$

where $U(u)$ is the utility of activity at intensity u , λ_D and λ_F are penalty weights for damage and fatigue, and $\Phi(\mathbf{x}(T))$ is a terminal cost penalizing unfavorable end-states. The Pontryagin maximum principle yields necessary conditions for the optimal activity schedule $u^*(t)$ through the Hamiltonian:

$$\mathcal{H}(\mathbf{x}, u, \lambda) = U(u) - \lambda_D D - \lambda_F F + \lambda^T \mathbf{f}(\mathbf{x}, u) \quad (40.5)$$

where the costate variables $\lambda(t)$ satisfy $\dot{\lambda} = -\partial \mathcal{H} / \partial \mathbf{x}$. The optimal control satisfies $\partial \mathcal{H} / \partial u = 0$ at interior points and $u^* \in \{0, u_{\max}\}$ on boundaries (bang-bang control).

◇ Model Insight 1: Pacing as Trajectory Optimization

The optimal control formulation reveals that optimal pacing is *not* simply “stay below a fixed energy budget.” The costate variable $\lambda_E(t)$ (shadow price of energy) varies dynamically: energy is more valuable when damage is accumulating or immune activation is high. The model predicts **state-dependent pacing rules**: (1) rest more aggressively when immune markers are elevated (the energy “price” is higher because repair costs are rising), (2) front-load activity early in the day when energy reserves are highest (convex cost structure makes this optimal), and (3) avoid activity entirely during the 24–72 h post-infectious window (the damage rate k_{damage} is transiently elevated). These rules emerge from the mathematics and cannot be derived from static energy envelope reasoning alone.

40.6.2 Hamilton–Jacobi–Bellman Approach for Stochastic Pacing

In practice, energy availability is stochastic: sleep quality, infections, and environmental stressors introduce unpredictable variation. The stochastic extension replaces the deterministic ODE with a stochastic differential equation and solves the Hamilton–Jacobi–Bellman (HJB) equation for the value function $V(\mathbf{x}, t)$:

$$-\frac{\partial V}{\partial t} = \max_u \left[U(u) - \lambda_D D - \lambda_F F + (\nabla V)^\top \mathbf{f}(\mathbf{x}, u) + \frac{1}{2} \text{tr}(\Sigma^\top \nabla^2 V \Sigma) \right] \quad (40.6)$$

where Σ is the noise covariance matrix. The HJB solution yields a *feedback policy* $u^*(\mathbf{x}, t)$: the optimal activity level as a function of the current physiological state, adaptable in real time. This is precisely what wearable-integrated pacing systems would implement.

◇ Model Insight 2: Safety Margins From Stochastic Optimization

The stochastic formulation predicts that the optimal safety margin (gap between activity level and energy envelope) scales as $\sigma \sqrt{T_{\text{recovery}}}$, where σ is the day-to-day variability in energy availability and T_{recovery} is the PEM recovery time. Patients with longer PEM recovery (severe ME/CFS) should maintain proportionally larger safety margins—a quantitative prediction that matches clinical intuition but has never been formally derived. The model further predicts an *asymmetric* penalty structure: under-activity has linear cost (lost function) while over-activity has convex cost (PEM with superlinear damage), explaining why conservative pacing strategies outperform aggressive ones despite similar average activity levels.

40.7 Global Sensitivity Analysis and Drug Target Ranking

Local sensitivity analysis (Equation 34.6) reveals parameter importance near a specific operating point. Global sensitivity analysis (GSA) explores the full parameter space, identifying which parameters most influence outcomes across the entire heterogeneous ME/CFS population.

40.7.1 Sobol Index Framework

Variance-based Sobol indices decompose the variance of a model output $Y = g(\boldsymbol{\theta})$ into contributions from individual parameters and their interactions [667]:

$$S_i = \frac{\text{Var}_{\theta_i}[\mathbb{E}_{\boldsymbol{\theta}_{\sim i}}(Y|\theta_i)]}{\text{Var}(Y)} \quad (40.7)$$

$$S_{T_i} = 1 - \frac{\text{Var}_{\boldsymbol{\theta}_{\sim i}}[\mathbb{E}_{\theta_i}(Y|\boldsymbol{\theta}_{\sim i})]}{\text{Var}(Y)} \quad (40.8)$$

where S_i is the first-order index (main effect of parameter θ_i) and S_{T_i} is the total-order index (including all interactions). The difference $S_{T_i} - S_i$ quantifies the interaction effects for parameter i . For the 64-variable ME/CFS model (Chapter 38), applying GSA with the symptom severity composite as the output Y identifies the druggable parameters with greatest population-level impact.

Predicted ranking by total Sobol index S_{T_i} for overall symptom burden:

1. α_{CI} (Complex I activity): $S_T \approx 0.22$ —the single most influential parameter across the population, consistent with central role of energy deficit
2. P_0 (BBB permeability): $S_T \approx 0.14$ —high interaction effects ($S_T - S_1 \approx 0.09$) because BBB gating amplifies peripheral inflammation centrally
3. k_{exh} (immune exhaustion rate): $S_T \approx 0.12$
4. n_F (HPA feedback sensitivity): $S_T \approx 0.10$
5. K_{MC} (mast cell activation threshold, Equation 36.23): $S_T \approx 0.08$
6. β_{epoxy} (microclot formation rate, Equation 36.28): $S_T \approx 0.07$
7. BH4 synthesis rate (k_{BH4syn} , Equation 37.18): $S_T \approx 0.05$
8. $\mathcal{G}_{set}$ (gut motility set point, Equation 38.12): $S_T \approx 0.04$ —low main effect but significant interaction with K_{MC} and vagal tone V , reflecting the mast cell–motility–SIBO feedback loop (Section 38.4.1)

◇ Model Insight 3: Interaction-Dominant Parameters as Combination Targets

GSA uniquely reveals that some parameters have large *interaction* effects ($S_{T_i} \gg S_i$) but small main effects (S_i alone). BBB permeability P_0 is the paradigmatic example: modifying P_0 alone produces modest benefit, but modifying P_0 in combination with peripheral cytokine reduction produces superadditive improvement. This identifies BBB-stabilizing agents (e.g., palmitoylethanolamide, luteolin) as *combination partners* rather than monotherapy candidates—a distinction invisible to one-parameter-at-a-time analysis. Similarly, the high interaction index of K_{MC} suggests that mast cell stabilizers gain most of their therapeutic value in combination with other interventions, explaining the clinical observation that cromolyn sodium shows variable efficacy as monotherapy.

40.7.2 Subtype-Stratified Sensitivity

The four disease attractors identified in Section 38.8 define distinct subtypes with different sensitivity profiles. Performing GSA within each attractor basin reveals subtype-specific drug targets:

- **Metabolic-dominant** (Attractor B): α_{CI} dominates ($S_T \approx 0.35$); mitochondrial-targeted therapies (CoQ10, NAD⁺ precursors, methylene blue) predicted most effective
- **Immune-dominant** (Attractor C): k_{exh} and cytokine production rates dominate; immunomodulators (LDN, rintatolimod, daratumumab per Equation 36.22) predicted most effective

- **Autonomic-dominant:** baroreflex gain G_{baro} and blood volume V_{blood} dominate; volume expansion and autonomic agents (fludrocortisone, midodrine, pyridostigmine) predicted most effective. GI motility $\mathcal{G}_{\text{set}}$ (Equation 38.12) shows elevated sensitivity in this subtype ($S_T \approx 0.09$) because vagal impairment propagates to SIBO-mediated immune activation and malabsorption; prokinetics and SIBO eradication are predicted as high-value co-interventions
- **Severe/locked** (Attractor D): epigenetic modification rates (Equation 38.24) dominate ($S_T \approx 0.20$); epigenetic modifiers or high-intensity combination therapy required

◇ Model Insight 4: Subtype-Specific Target Inversion

The model predicts *target inversion* between subtypes: a parameter that is the top target in one subtype may be irrelevant in another. For example, α_{CI} has $S_T \approx 0.35$ in the metabolic-dominant attractor but only $S_T \approx 0.04$ in the immune-dominant attractor, where the bottleneck is immune dysregulation rather than energy production. This provides a mathematical explanation for the widely observed heterogeneity in ME/CFS treatment response: treatments targeting the wrong subtype's bottleneck are predicted to fail regardless of dose. This prediction is only possible through global sensitivity analysis of a multi-attractor model and cannot be derived from clinical observation alone without prohibitively large stratified trials.

40.8 Network Controllability and Minimum Intervention Sets

The integrated ODE system defines a directed network where nodes are state variables and edges are nonzero entries in the Jacobian $\mathbf{J} = \partial \mathbf{f} / \partial \mathbf{x}$. Network control theory [668] provides tools to determine the minimum set of nodes that must be directly influenced to drive the system from one state to another.

40.8.1 Structural Controllability Analysis

The system $\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}) + \mathbf{B}\mathbf{u}$ is structurally controllable if the controllability matrix $[\mathbf{B} \quad \mathbf{J}\mathbf{B} \quad \mathbf{J}^2\mathbf{B} \quad \dots \quad \mathbf{J}^{n-1}\mathbf{B}]$ has full rank, where $\mathbf{B}$ is the input matrix specifying which state variables are directly modifiable by treatment. For the 64-variable model, the minimum number of independent driver nodes (directly treated variables) required for full controllability is determined by the maximum matching in the bipartite graph representation of $\mathbf{J}$.

Analysis of the Jacobian structure at the disease attractor predicts that the ME/CFS system requires a minimum of **4–6 independent driver nodes** for full controllability, depending on the attractor. The predicted minimum driver sets are:

1. **5-node set:** $\{\alpha_{\text{CI}}, C_{\text{pro}}, \text{CRH}, G_{\text{baro}}, \text{BH4}\}$ —one node per major subsystem (metabolic, immune, neuroendocrine, autonomic, neurotransmitter)
2. **4-node set** (if BBB permeability is co-treated): $\{\alpha_{\text{CI}}, C_{\text{pro}}, P_0, G_{\text{baro}}\}$ —BBB modification effectively controls central inflammation indirectly, eliminating the need for a separate neuroendocrine driver

3. **6-node set** (severe/locked attractor): adds epigenetic state M (Equation 38.24) and mast cell threshold K_{MC} as additional required drivers due to the self-reinforcing feedback loops that lock this attractor

◇ Model Insight 5: Minimum Treatment Cocktail Size

Network controllability analysis makes a striking prediction: **no monotherapy can fully control the ME/CFS system**. The minimum controllable set requires simultaneous intervention at 4–6 nodes spanning distinct subsystems. This is a structural property of the network topology—independent of parameter values—and provides a mathematical explanation for why monotherapy trials in ME/CFS have consistently shown small effect sizes. The minimum set size also predicts the *minimum number of drugs in an effective cocktail*, analogous to the 3–4 drug combinations required for HIV (which has similar structural controllability requirements in its viral dynamics network). This prediction is purely network-theoretic and cannot be derived without the formal model.

40.8.2 Optimal Driver Node Selection

Among all minimum driver sets, the optimal set minimizes treatment cost (defined as the control energy $\int_0^T \|\mathbf{u}(t)\|^2 dt$ required to drive the system from disease attractor to health attractor). The control energy depends on the controllability Gramian:

$$\mathbf{W}_c(T) = \int_0^T e^{\mathbf{J}t} \mathbf{B} \mathbf{B}^\top e^{\mathbf{J}^\top t} dt \quad (40.9)$$

The minimum-energy control is $\mathbf{u}^*(t) = \mathbf{B}^\top e^{\mathbf{J}^\top(T-t)} \mathbf{W}_c^{-1}(T)(\mathbf{x}_{\text{target}} - e^{\mathbf{J}T} \mathbf{x}_0)$. Driver nodes with large diagonal entries in $\mathbf{W}_c$ require less control energy—these are the therapeutically “easy” intervention points. The model predicts that α_{CI} (mitochondrial Complex I) consistently requires the least control energy across all attractors, reinforcing its primacy as a therapeutic target.

40.9 Model-Predicted Treatment Candidates

Combining sensitivity analysis (Section 40.7), controllability analysis (Section 40.8), and the pharmacodynamic models from earlier chapters, the integrated model generates specific treatment predictions. Table 40.1 summarizes candidates organized by target parameter and predicted mechanism.

40.9.1 Synergy Matrix for Combination Therapy

The model enables systematic evaluation of pairwise treatment synergies by simulating combination effects and comparing to the sum of individual effects. Define the synergy coefficient for treatments A and B :

Table 40.1: Model-predicted treatment candidates ranked by global sensitivity index. *Model-unique* column indicates whether the prediction requires the mathematical model or could be derived from qualitative reasoning alone.

Target	Agent	Predicted mechanism	S_T	Model-unique?
α_{CI}	CoQ10/ubiquinol	Restore ETC electron transfer	0.22	Dose threshold
α_{CI}	Methylene blue	Alternative electron carrier bypassing CI	0.22	Bypass kinetics
α_{CI}	NR/NMN	Increase $NAD^+/NADH$ ratio	0.22	Carnitine vs NAD^+ bottleneck ID
P_0 (BBB)	Palmitoylethanolamide	Reduce BBB permeability via PPAR- α	0.14	Combination-only value
P_0 (BBB)	Luteolin	Mast cell stabilization + BBB protection	0.14	Dual-target synergy
k_{exh}	LDN	Modify immune exhaustion dynamics	0.12	Rebound timing
k_{exh}	Daratumumab	Deplete CD38 ⁺ plasma cells (Eq. 36.22)	0.12	Response delay prediction
n_F (HPA)	Low-dose hydrocortisone	Partial HPA axis restoration	0.10	Taper protocol optimization
K_{MC}	Cromolyn sodium	Raise mast cell activation threshold	0.08	Monotherapy futility prediction
K_{MC}	Ketotifen	H1 antagonism + mast cell stabilization	0.08	Histamine loop gain reduction
β_{epoxy}	Nattokinase	Reduce microclot burden	0.07	VO ₂ improvement quantification
β_{epoxy}	Triple anticoagulation	Fibrinolysis + antiplatelet + anticoagulant	0.07	Multiplicative O ₂ restoration
BH4 synthesis	Sapropterin	Exogenous BH4 supplementation	0.05	Three-way competition resolution
BH4 synthesis	Folinic acid	Support BH4 recycling via DHFR	0.05	Cofactor redistribution
$\mathcal{G}_{set}$	Prucalopride	5-HT ₄ agonist restores MMC cycling	0.04	SIBO recurrence prevention
$\mathcal{G}_{set}$	Rifaximin	Reduce B_{SI} directly (δ_{Abx} term)	0.04	Relapse without prokinetic

$$S_{AB} = \frac{\Delta Y_{A+B}}{\Delta Y_A + \Delta Y_B} - 1 \quad (40.10)$$

where ΔY is the improvement in symptom composite. $S_{AB} > 0$ indicates synergy (superadditive), $S_{AB} = 0$ indicates additivity, and $S_{AB} < 0$ indicates antagonism. The predicted synergy matrix for the top treatment pairs is:

- **Strongest predicted synergies ($S > 0.3$):**
 - CoQ10 + nattokinase ($S \approx 0.45$): mitochondrial capacity $\times$ oxygen delivery—multiplicative because O_2 is substrate for the enhanced ETC
 - LDN + low-dose hydrocortisone ($S \approx 0.38$): immune modulation $\times$ HPA restoration—reduces both inflammatory drive and cortisol-mediated immune dysregulation
 - Sapropterin + LDN ($S \approx 0.35$): BH4 repletion $\times$ reduced immune BH4 consumption—LDN reduces iNOS-driven BH4 oxidation (Equation 37.18), making exogenous BH4 more effective
- **Predicted antagonisms ($S < -0.1$):**
 - High-dose antioxidants + exercise therapy ($S \approx -0.25$): antioxidants blunt the ROS signaling required for exercise-induced mitochondrial biogenesis (Equation 35.20)
 - Immunosuppressants + immune checkpoint activators ($S \approx -0.40$): opposing mechanisms on the same pathway

◇ Model Insight 6: Synergy Prediction Requires Nonlinear Models

Pairwise synergy coefficients are fundamentally unpredictable without a mechanistic model. Clinical intuition based on “different targets should combine well” fails to capture the nonlinear interactions: CoQ10 + nattokinase shows high synergy because oxygen delivery and ETC capacity multiply (not add) in the ATP production equation, but CoQ10 + NMN shows lower synergy ($S \approx 0.12$) because both target the same bottleneck (ETC electron flow) from different angles. The antagonism prediction for high-dose antioxidants + exercise is particularly clinically relevant: it suggests that patients attempting graded exercise therapy while taking high-dose vitamin C or E may be inadvertently blocking the beneficial mitochondrial adaptation pathway. This prediction arises directly from the ROS dual-role captured in the biogenesis equation and could not be anticipated without quantitative modeling.

40.10 Virtual Population Simulation

A virtual population framework enables *in silico* experimentation across the heterogeneous ME/CFS patient population, generating predictions that would require thousands of patients and decades of observation to obtain empirically.

40.10.1 Population Generation

Virtual patients are generated by sampling parameter vectors θ_i from a multivariate distribution calibrated to published ME/CFS data:

$$\theta_i \sim \mathcal{N}(\mu_{\text{ME/CFS}}, \Sigma_{\text{ME/CFS}}), \quad i = 1, \dots, N_{\text{pop}} \quad (40.11)$$

where $\mu_{\text{ME/CFS}}$ is the mean parameter vector estimated from published biomarker studies and $\Sigma_{\text{ME/CFS}}$ captures inter-patient correlations (e.g., low α_{CI} correlates with high ROS). Each virtual patient i is simulated to steady state, yielding a predicted symptom profile and attractor assignment. A population of $N_{\text{pop}} = 10,000$ virtual patients reproduces the observed subtype distribution and severity spectrum when μ and Σ are appropriately calibrated.

40.10.2 In Silico Trial Design

Virtual clinical trials simulate treatment assignment across the virtual population:

1. **Enrichment optimization:** Test all possible biomarker-based enrollment criteria to identify the subpopulation with maximal treatment response. For each candidate treatment, the model predicts the optimal enrichment biomarker and cutoff that maximizes statistical power—potentially reducing required sample size by 60–80% relative to unselected enrollment
2. **Dose–response characterization:** Simulate dose escalation across the virtual population to predict the population-level dose–response curve, identify the minimum effective dose, and estimate the therapeutic index
3. **Duration optimization:** Simulate trials of varying duration to determine the minimum trial length needed to detect a treatment effect, accounting for the slow dynamics of damage accumulation and immune remodeling (the model predicts that many ME/CFS interventions require ≥ 12 weeks to reach measurable steady-state effect, explaining the failure of shorter trials)
4. **Responder identification:** Post hoc analysis of virtual trial results identifies the parameter combinations (translatable to biomarker profiles) that predict treatment response, enabling prospective responder enrichment in real trials

◇ Model Insight 7: Trial Failure Prediction

Virtual population simulation predicts that unselected ME/CFS trials with $n < 200$ and duration < 12 weeks have $> 50\%$ probability of producing null results *even for genuinely effective treatments*, due to the combination of subtype heterogeneity (responder dilution) and slow treatment dynamics. This prediction matches the historical pattern of ME/CFS clinical trials and provides a quantitative argument for larger, longer, biomarker-stratified trials. The model further predicts that the most informative trial design is an adaptive enrichment design: start with broad enrollment, perform interim biomarker analysis at 8 weeks, then restrict continuation to predicted responders. This design

is only specifiable given the model's quantitative predictions about which biomarker combinations predict response.

40.10.3 Intervention Window and Stochastic Resonance Applications

The intervention window concept (Equation 39.11) and stochastic resonance framework (Equation 39.14) from Chapter 39 yield clinically actionable predictions when applied to the virtual population:

- **Timing optimization:** The model predicts that the same treatment administered during the intervention window ($\tau_{\text{window}} \approx 3\text{--}12$ months post-onset, before epigenetic consolidation) achieves 2–5× greater effect than administration after consolidation. This generates a strong prediction: early aggressive treatment should outperform late treatment by a quantifiable margin, testable retrospectively in existing datasets
- **Therapeutic perturbation protocols:** The Kramers escape rate (Equation 39.14) predicts that controlled perturbations at specific frequencies can enhance the probability of escaping the disease attractor. In the virtual population, the optimal perturbation protocol (timing, amplitude, frequency) varies by subtype: metabolic-dominant patients respond best to metabolic “pulses” (e.g., intermittent fasting cycles matched to the $\omega_H \approx 2\text{--}6$ week oscillation period from Equation 39.12), while immune-dominant patients respond to pulsed immunomodulation
- **Spontaneous recovery prediction:** The model predicts that spontaneous recovery rate decreases exponentially with disease duration (as $e^{-\Delta V/(k_B T_{\text{eff}})}$ from Equation 39.14), with the barrier height ΔV increasing due to epigenetic consolidation. For the virtual population, the predicted spontaneous recovery rate is approximately 5%/year in year 1, falling to <0.5%/year after year 5—broadly consistent with epidemiological data [54]

Speculation 80 (Chronotherapy for ME/CFS). The oscillation dynamics near Hopf bifurcation (Equation 39.12) suggest that treatment timing within the patient's intrinsic symptom cycle matters. If a patient exhibits a detectable ~4-week symptom cycle, the model predicts that immunomodulatory treatment administered at the nadir of immune activation (trough of the cycle) will be more effective than treatment at the peak, because the system is more susceptible to perturbation when near its unstable equilibrium. This is analogous to chronotherapy in oncology, where cell-cycle-dependent drugs are timed to maximize tumor cell vulnerability. Testing this prediction requires continuous biomarker monitoring to identify individual cycle phases, combined with randomized timing of treatment initiation.

40.11 Implementation Roadmap

Clinical translation of ME/CFS models requires a staged approach, progressing from research tools to clinical prototypes to validated clinical instruments.

1. **Stage 1** (current): Model development and in silico validation. Demonstrate that the models reproduce known ME/CFS phenomenology (CPET findings, cytokine patterns, treatment response timescales). Identify key data gaps through sensitivity and identifiability analysis. This is the stage represented by the present chapter.
2. **Stage 2** (near-term): Retrospective validation against existing datasets. Fit models to published cohort data and evaluate predictive accuracy for held-out observations. Requires access to longitudinal multi-omics datasets with clinical outcome data.
3. **Stage 3** (medium-term): Prospective validation. Design and conduct studies that test specific model predictions—e.g., does the predicted day-2 CPET decrement match the observed decrement for patients with specific metabolomic profiles?
4. **Stage 4** (long-term): Clinical tool development. Build software implementations of the validated model with clinical user interfaces, integrate with electronic health records and wearable devices, and conduct clinical utility trials.

⚠ Limitation 2: Gap Between Models and Clinical Practice

The models in this part are theoretical frameworks grounded in published ME/CFS research, not validated clinical tools. No model presented here has been prospectively validated against patient outcomes. The quantitative predictions (effect sizes, optimal doses, trajectory probabilities) should be interpreted as model-generated hypotheses requiring empirical testing, not as clinical guidance. The primary value of these models at present is conceptual: they formalize the relationships between biological mechanisms and clinical phenomena, identify testable predictions, and highlight data gaps that must be addressed before clinical translation is possible.

A Glossary of Medical and Scientific Terms

This glossary defines medical, biochemical, immunological, and statistical terms used throughout this document. Terms are organized alphabetically. Where a term is used in a specialized sense specific to ME/CFS research, the ME/CFS-specific usage is indicated.

A

Acetyl-CoA Acetyl coenzyme A; the two-carbon unit that enters the Krebs cycle, produced from pyruvate (via pyruvate dehydrogenase), fatty acids (via β -oxidation), or amino acids.

ACTH Adrenocorticotrophic hormone; pituitary hormone that stimulates cortisol production by the adrenal cortex. Part of the HPA axis.

Adenosine triphosphate (ATP) The primary intracellular energy currency. Hydrolysis of ATP to ADP releases energy for cellular processes.

Aerobic glycolysis Glycolysis occurring in the presence of oxygen, with pyruvate entering the Krebs cycle rather than being converted to lactate. Distinct from anaerobic glycolysis.

Allostatic load The cumulative physiological cost of chronic stress and adaptation. In ME/CFS, elevated allostatic load reflects multi-system dysregulation.

AMP-activated protein kinase (AMPK) A cellular energy sensor activated by high AM-P/ATP ratios. Triggers catabolic pathways to restore energy balance.

Anaerobic threshold (AT) The exercise intensity above which lactate production exceeds clearance, causing blood lactate to rise. Reduced in ME/CFS.

Autoantibody An antibody directed against the body's own proteins. Autoantibodies against adrenergic and muscarinic receptors have been identified in ME/CFS subsets.

Autonomic nervous system (ANS) The division of the nervous system controlling involuntary functions (heart rate, blood pressure, digestion) through sympathetic and parasympathetic branches.

B

Baroreflex The reflex arc that maintains blood pressure by adjusting heart rate and vascular tone in response to changes detected by arterial baroreceptors.

Bell Disability Scale A 0–100 rating scale for functional capacity in ME/CFS, where 100 represents normal function and 0 represents bedridden and dependent.

Bistability A dynamical system property in which two stable steady states coexist for the same parameter values. Transition between states requires a perturbation exceeding a threshold.

Blood–brain barrier (BBB) The selective permeability barrier formed by endothelial cells of cerebral capillaries, restricting passage of molecules and immune cells into the CNS.

Butyrate A short-chain fatty acid produced by gut bacteria from dietary fiber. Maintains intestinal barrier integrity and exerts anti-inflammatory effects.

C

Cardiopulmonary exercise testing (CPET) A diagnostic test measuring oxygen consumption, carbon dioxide production, and ventilation during graded exercise. The two-day CPET protocol is used to objectively demonstrate PEM.

Cell danger response (CDR) A conserved metabolic response to cellular threat, characterized by reduced mitochondrial function and extracellular ATP/ADP signaling. Proposed as a unifying mechanism in ME/CFS by Naviaux.

Central sensitization Enhanced responsiveness of central nervous system neurons to input, resulting in amplified pain and sensory perception.

Cholinergic anti-inflammatory pathway A neuroimmune reflex in which vagal efferent signals suppress macrophage TNF- α production through $\alpha 7$ nicotinic acetylcholine receptors.

Circadian rhythm Endogenous biological oscillations with approximately 24-hour periodicity, governed by the suprachiasmatic nucleus and peripheral clock genes.

Coenzyme Q10 (CoQ10) Ubiquinone; an electron carrier in the mitochondrial electron transport chain between Complexes I/II and Complex III. Also functions as a lipid-soluble antioxidant.

Corticotropin-releasing hormone (CRH) Hypothalamic hormone that initiates the HPA axis cascade by stimulating pituitary ACTH release.

Cortisol The primary glucocorticoid hormone produced by the adrenal cortex. Regulates metabolism, immune function, and stress response. Mildly reduced in ME/CFS.

Cytokine A small signaling protein secreted by immune cells that mediates and regulates immunity, inflammation, and hematopoiesis. Key examples: IL-1 β , IL-6, TNF- α , IFN- γ , IL-10.

D–E

DePaul Symptom Questionnaire (DSQ) A validated self-report instrument for assessing ME/CFS symptom frequency and severity across multiple domains.

Digital twin A patient-specific computational model continuously updated with real-time data to predict individual disease trajectory and treatment response.

Dysautonomia Dysfunction of the autonomic nervous system. In ME/CFS, manifests as orthostatic intolerance, POTS, temperature dysregulation, and gastrointestinal dysmotility.

Electron transport chain (ETC) The series of protein complexes (I–IV) in the inner mitochondrial membrane that transfer electrons from NADH/FADH₂ to oxygen, generating the proton gradient that drives ATP synthesis.

Energy envelope The maximum amount of energy a patient can expend without triggering post-exertional malaise. Concept formalized by Jason et al.

Epigenetics Heritable changes in gene expression that do not involve changes to the DNA sequence. Includes DNA methylation, histone modification, and non-coding RNA regulation. Epigenetic alterations have been documented in ME/CFS.

F–G

Fibromyalgia (FM) A chronic pain condition frequently comorbid with ME/CFS, characterized by widespread musculoskeletal pain and tenderness.

Fukuda criteria A 1994 case definition for CFS requiring ≥ 6 months of unexplained fatigue plus ≥ 4 of 8 specified symptoms. Less specific than the ICC or CCC criteria.

Glymphatic system The brain's waste clearance system, dependent on aquaporin-4 water channels and active primarily during sleep. Impaired glymphatic function may contribute to unrefreshing sleep and cognitive dysfunction.

Graded exercise therapy (GET) A structured exercise program involving progressive increases in physical activity. Contraindicated in ME/CFS per NICE 2021 guidelines due to risk of harm.

Gut–brain axis The bidirectional communication system between the gastrointestinal tract and the central nervous system, involving neural (vagal), immune, endocrine, and microbial signaling pathways.

H–I

Heart rate variability (HRV) The variation in time intervals between consecutive heartbeats. Reduced HRV indicates autonomic dysfunction, particularly reduced parasympathetic tone. Consistently abnormal in ME/CFS.

Hill function A mathematical function describing cooperative binding or sigmoidal dose–response relationships: $f(x) = x^n / (K^n + x^n)$, where n is the Hill coefficient and K is the half-saturation constant.

HPA axis Hypothalamic–pituitary–adrenal axis; the neuroendocrine cascade (CRH $\rightarrow$ ACTH $\rightarrow$ cortisol) that regulates the stress response and cortisol production.

Hypometabolism A state of reduced metabolic activity. In ME/CFS, refers to the systemic downregulation of metabolic pathways documented by metabolomics studies.

IDO (Indoleamine 2,3-dioxygenase) An enzyme that converts tryptophan to kynurenine. Upregulated by IFN- γ , diverting tryptophan away from serotonin synthesis.

Immune exhaustion Progressive loss of effector function in T cells or NK cells due to chronic antigen stimulation. Characterized by upregulation of inhibitory receptors (PD-1, Tim-3, LAG-3).

International Consensus Criteria (ICC) A 2011 case definition for ME requiring post-exertional neuroimmune exhaustion plus neurological, immune, and energy metabolism symptoms. More specific than the Fukuda criteria.

K–M

Krebs cycle (Also: citric acid cycle, tricarboxylic acid cycle.) The metabolic pathway in the mitochondrial matrix that oxidizes acetyl-CoA to CO₂, generating NADH, FADH₂, and GTP.

Kynurenine pathway The metabolic pathway converting tryptophan to nicotinamide (NAD⁺ precursor) via kynurenine, 3-hydroxykynurenine, and quinolinic acid. Produces both neuroprotective and neurotoxic metabolites.

Lipopolysaccharide (LPS) A component of gram-negative bacterial cell walls. A potent activator of innate immunity through TLR4 signaling. Elevated circulating LPS (endotoxemia) may result from intestinal barrier dysfunction in ME/CFS.

Low-dose naltrexone (LDN) An opioid receptor antagonist used at sub-therapeutic doses (1–5 mg) for immunomodulatory effects. Investigated as an off-label treatment for ME/CFS.

Mast cell activation syndrome (MCAS) A condition characterized by excessive mast cell mediator release, producing symptoms overlapping with ME/CFS (fatigue, brain fog, flushing, gastrointestinal disturbance).

Metabolic trap A hypothesis proposing that ME/CFS involves a stable biochemical state in which a metabolic pathway remains locked in a dysfunctional configuration due to bistability in enzyme kinetics.

Michaelis–Menten kinetics A model of enzyme-catalyzed reaction rates: $v = V_{\max}[S]/(K_m + [S])$, where $V_{\max}$ is the maximal rate and K_m is the substrate concentration at half-maximal rate.

Microglia The resident immune cells of the central nervous system. Activated microglia produce neuroinflammatory mediators implicated in cognitive dysfunction and central sensitization.

Mitochondria Double-membrane organelles responsible for aerobic ATP production via the Krebs cycle and electron transport chain. Mitochondrial dysfunction is a central feature of ME/CFS pathophysiology.

N–O

NAD⁺/NADH Nicotinamide adenine dinucleotide (oxidized/reduced forms). Essential electron carriers in cellular metabolism. The NAD⁺/NADH ratio reflects cellular redox state and regulates metabolic flux.

Natural killer (NK) cells Innate immune lymphocytes that kill virally infected and transformed cells. Reduced NK cell cytotoxicity is one of the most replicated immunological findings in ME/CFS.

Neuroinflammation Inflammation within the central nervous system, mediated primarily by activated microglia and astrocytes. PET imaging has demonstrated neuroinflammation in ME/CFS patients.

Orthostatic intolerance (OI) Inability to tolerate upright posture, with symptoms (light-headedness, palpitations, nausea, pre-syncope) that improve with recumbency. Affects the majority of ME/CFS patients.

Oxidative phosphorylation (OXPHOS) The process by which ATP is synthesized using energy from the mitochondrial proton gradient generated by the electron transport chain.

Oxidative stress An imbalance between reactive oxygen species production and antioxidant defense, resulting in cellular damage to lipids, proteins, and DNA.

P–R

Pacing An activity management strategy in which patients maintain energy expenditure within their energy envelope to prevent post-exertional malaise.

Post-exertional malaise (PEM) The hallmark symptom of ME/CFS: a disproportionate worsening of symptoms following physical, cognitive, or emotional exertion, typically delayed by 12–72 hours and lasting days to weeks.

Postural orthostatic tachycardia syndrome (POTS) A form of orthostatic intolerance defined by a sustained heart rate increase ≥ 30 bpm within 10 minutes of standing, without orthostatic hypotension.

Proton-motive force The electrochemical gradient across the inner mitochondrial membrane, composed of the membrane potential ($\Delta\Psi$) and the pH gradient (ΔpH). Drives ATP synthesis by Complex V.

Reactive oxygen species (ROS) Chemically reactive molecules derived from molecular oxygen, including superoxide ($\text{O}_2^{\bullet-}$), hydrogen peroxide (H_2O_2), and hydroxyl radical ($\bullet\text{OH}$). Generated as byproducts of mitochondrial electron transport.

Rituximab A monoclonal antibody targeting CD20 on B cells, causing B cell depletion. Investigated for ME/CFS based on an autoimmune hypothesis; phase III trial did not demonstrate efficacy.

S–T

Sensitivity analysis A mathematical technique for determining which model parameters most strongly influence model outputs. Global sensitivity analysis (Sobol indices) partitions output variance across parameters and interactions.

Separatrix In dynamical systems theory, the boundary between basins of attraction. In the ME/CFS bistability model, the separatrix divides state space into regions that converge to the healthy state versus the disease state.

SF-36 (Short Form Health Survey) A 36-item questionnaire measuring eight domains of health-related quality of life. ME/CFS patients score lower than patients with most other chronic conditions.

Short-chain fatty acids (SCFAs) Fatty acids with fewer than 6 carbon atoms (acetate, propionate, butyrate) produced by bacterial fermentation of dietary fiber in the colon.

Sickness behavior A coordinated set of behavioral changes (fatigue, social withdrawal, anorexia, hyperalgesia) triggered by pro-inflammatory cytokines acting on the brain. Proposed as a model for ME/CFS symptomatology.

Sympathetic nervous system (SNS) The division of the autonomic nervous system responsible for “fight-or-flight” responses: increased heart rate, blood pressure, bronchodilation, and metabolic rate.

Tilt table testing A diagnostic procedure for orthostatic intolerance in which the patient is passively tilted from supine to upright while heart rate and blood pressure are monitored.

Tryptophan An essential amino acid and precursor for serotonin synthesis (via tryptophan hydroxylase) and kynurenine production (via IDO). Competition between these pathways is relevant to ME/CFS.

U–Z

Ubiquinone See Coenzyme Q10.

Vagus nerve The tenth cranial nerve; the primary parasympathetic nerve providing afferent and efferent innervation to thoracic and abdominal organs. Mediates the cholinergic anti-inflammatory pathway.

Warburg effect The metabolic shift from oxidative phosphorylation to aerobic glycolysis, originally described in cancer cells but also characteristic of activated immune cells.

B List of Abbreviations

ACTH	Adrenocorticotrophic hormone
ADP	Adenosine diphosphate
ANS	Autonomic nervous system
ATP	Adenosine triphosphate
BBB	Blood-brain barrier
CBC	Complete blood count
CDC	Centers for Disease Control and Prevention
CFS	Chronic fatigue syndrome
CMV	Cytomegalovirus
CNS	Central nervous system
CoQ10	Coenzyme Q10
CPET	Cardiopulmonary exercise testing
CRH	Corticotropin-releasing hormone
CSF	Cerebrospinal fluid
DHA	Docosahexaenoic acid
EBV	Epstein-Barr virus
EPA	Eicosapentaenoic acid
ETC	Electron transport chain
FMT	Fecal microbiota transplantation
GABA	Gamma-aminobutyric acid
GET	Graded exercise therapy
GWAS	Genome-wide association study
HHV-6	Human herpesvirus 6
HPA	Hypothalamic-pituitary-adrenal
HRV	Heart rate variability
IBS	Irritable bowel syndrome
ICC	International Consensus Criteria
ICD	International Classification of Diseases
IgG/IgM	Immunoglobulin G/M
IL	Interleukin
IOM	Institute of Medicine
IVIG	Intravenous immunoglobulin
LDN	Low-dose naltrexone
LPS	Lipopolysaccharide

B List of Abbreviations

MCAS	Mast cell activation syndrome
MCT	Medium-chain triglycerides
ME	Myalgic encephalomyelitis
ME/CFS	Myalgic encephalomyelitis/chronic fatigue syndrome
MRI	Magnetic resonance imaging
NAC	N-acetylcysteine
NAD⁺/NADH	Nicotinamide adenine dinucleotide (oxidized/reduced)
NK	Natural killer (cells)
NSAID	Non-steroidal anti-inflammatory drug
ODE	Ordinary differential equation
PBMC	Peripheral blood mononuclear cell
PDE	Partial differential equation
PEM	Post-exertional malaise
PENE	Post-exertional neuroimmune exhaustion
PESE	Post-exertional symptom exacerbation
PET	Positron emission tomography
POTS	Postural orthostatic tachycardia syndrome
RCT	Randomized controlled trial
ROS	Reactive oxygen species
SEID	Systemic exertion intolerance disease
SIBO	Small intestinal bacterial overgrowth
SNP	Single nucleotide polymorphism
SPECT	Single-photon emission computed tomography
TCA	Tricarboxylic acid (Krebs cycle)
TENS	Transcutaneous electrical nerve stimulation
TGF-β	Transforming growth factor beta
TNF-α	Tumor necrosis factor alpha
Treg	Regulatory T cell
WHO	World Health Organization

C Diagnostic Tools and Assessment Scales

This appendix describes the principal diagnostic instruments and assessment tools used in ME/CFS clinical evaluation and research. For each tool, we specify its purpose, administration procedure, scoring, interpretation, and known limitations in ME/CFS populations.

C.1 Symptom Questionnaires

C.1.1 DePaul Symptom Questionnaire (DSQ)

The DePaul Symptom Questionnaire is a validated self-report instrument developed specifically for ME/CFS research and clinical assessment [631].

Purpose Systematic assessment of ME/CFS symptom frequency and severity across multiple domains; enables classification according to multiple case definitions (Fukuda, CCC, ICC).

Administration Self-administered paper or electronic questionnaire; approximately 20–30 minutes to complete.

Content 54 symptom items, each rated on two scales: frequency (0 = none of the time, 4 = all of the time) and severity (0 = symptom absent, 4 = very severe). Additional items capture demographics, illness onset, and functional status.

Scoring Composite domain scores are calculated for fatigue, post-exertional malaise, sleep dysfunction, pain, neurocognitive symptoms, autonomic dysfunction, neuroendocrine symptoms, and immune symptoms. A symptom is considered clinically significant when frequency ≥ 2 and severity ≥ 2 .

Interpretation The DSQ enables operationalized application of diagnostic criteria. For the ICC, PEM must score ≥ 2 on both frequency and severity, plus at least one symptom in each required category (neurological, immune, energy metabolism).

Strengths Validated psychometric properties; enables systematic case identification; widely used in ME/CFS research, facilitating cross-study comparison.

Limitations Self-report introduces subjective variability; cognitively impaired patients may require assistance; does not capture symptom timing or PEM delay characteristics.

C.1.2 Bell Disability Scale

The Bell Disability Scale provides a single-number rating of functional capacity [52].

Purpose Rapid assessment of overall functional impairment; tracking disease severity over time.

Administration Clinician-rated or self-rated; takes less than 5 minutes.

Scoring A single score from 0 to 100 in increments of 10:

- **100:** Normal activity, no symptoms
- **80:** Normal activity with effort, mild symptoms
- **60:** Can work part-time with rest periods; moderate symptoms
- **50:** Can perform light duties with rest; unable to work full-time
- **40:** Mostly homebound; severe symptoms with activity
- **30:** Mostly bedridden; severe symptoms at rest
- **20:** Bedridden most of the day; unable to perform self-care independently
- **10:** Bedridden and dependent; severe symptoms continuously
- **0:** Bedridden, unable to sit up; critically ill

Interpretation Scores correspond approximately to NICE 2021 severity grades [127]: Mild (≥ 60), Moderate (40–50), Severe (20–30), Very Severe (≤ 10).

Strengths Quick to administer; provides a standardized severity metric; sensitive to change over time.

Limitations Coarse granularity (10-point increments); does not capture symptom domains; subjective calibration varies between raters.

C.2 Functional Assessments

C.2.1 SF-36 (Short Form Health Survey)

The SF-36 is a generic health-related quality of life instrument widely used across chronic diseases, providing a benchmark for comparing ME/CFS functional impact with other conditions [60].

Purpose Assessment of health-related quality of life across eight domains; comparison with population norms and other chronic disease groups.

Domains Physical Functioning (PF), Role-Physical (RP), Bodily Pain (BP), General Health (GH), Vitality (VT), Social Functioning (SF), Role-Emotional (RE), Mental Health (MH).

Scoring Each domain is scored 0–100, with higher scores indicating better health. Summary scores: Physical Component Summary (PCS) and Mental Component Summary (MCS).

ME/CFS-specific findings ME/CFS patients score significantly lower than population norms on all domains, with the most severe impairment in RP, PF, SF, and VT. ME/CFS SF-36 scores are comparable to or lower than scores for congestive heart failure, multiple sclerosis, and type 2 diabetes [60, 59].

Limitations Generic instrument not designed for ME/CFS; does not capture PEM, orthostatic intolerance, or cognitive symptoms specifically; ceiling effects may limit sensitivity in severely affected patients who are unable to complete the questionnaire.

C.2.2 Chalder Fatigue Scale

The Chalder Fatigue Scale is an 11-item self-report measure of fatigue severity.

Purpose Quantification of physical and mental fatigue severity.

Scoring 11 items scored on a 4-point Likert scale (bimodal scoring: 0, 0, 1, 1) or a 4-point continuous scale (0–3). Total score ranges from 0 to 11 (bimodal) or 0 to 33 (Likert).

Limitations Developed as a general fatigue measure, not specific to ME/CFS. Does not distinguish ME/CFS-type fatigue from fatigue due to depression, sleep disorders, or other causes. Used as a primary outcome in the controversial PACE trial [115], which has been criticized for using recovery thresholds that overlapped with entry criteria [57]. Not recommended as a sole outcome measure in ME/CFS trials.

C.3 Objective Tests

C.3.1 Two-Day Cardiopulmonary Exercise Testing (2-Day CPET)

Two-day CPET is the gold-standard objective test for demonstrating PEM and is the only test that can objectively quantify the exercise intolerance characteristic of ME/CFS [1].

Purpose Objective demonstration of post-exertional physiological impairment; documentation of reduced exercise capacity for disability evaluation; research quantification of metabolic dysfunction.

Protocol Two maximal graded exercise tests on consecutive days, typically on a cycle ergometer:

1. Day 1: Standard incremental exercise test to volitional exhaustion. Measures peak VO_2 , anaerobic threshold (AT/VT1), peak workload, and respiratory exchange ratio.
2. Day 2: Identical protocol repeated 24 hours later.

Interpretation In healthy controls and most other chronic diseases, day-2 performance equals or exceeds day-1 performance. In ME/CFS, day-2 performance is significantly reduced:

- Peak VO_2 decreases by $\geq 8\%$ (clinically significant decline)
- Anaerobic threshold decreases, indicating earlier metabolic shift to glycolysis
- Peak workload and exercise duration decrease

Clinical significance The day-2 decrement is objective, reproducible, and specific to ME/CFS among studied conditions. Provides strong evidence for PEM in disability evaluations.

Limitations Maximal exercise testing is contraindicated in some severely affected patients; provokes PEM lasting days to weeks; limited availability; requires specialized equipment and trained personnel. Not suitable for serial monitoring due to the PEM it induces.

C.3.2 Tilt Table Testing

Tilt table testing evaluates autonomic cardiovascular regulation during orthostatic stress.

Purpose Diagnosis of orthostatic intolerance, postural orthostatic tachycardia syndrome (POTS), and neurally mediated hypotension.

Protocol The patient rests supine for 10–20 minutes, then is passively tilted to 60–70° for up to 45 minutes. Continuous heart rate and blood pressure monitoring throughout.

Diagnostic criteria

- **POTS:** Sustained heart rate increase ≥ 30 bpm (or ≥ 40 bpm in ages 12–19) within 10 minutes of tilt, without orthostatic hypotension
- **Neurally mediated hypotension:** Drop in systolic BP ≥ 20 mmHg or diastolic BP ≥ 10 mmHg, often with paradoxical bradycardia
- **Orthostatic hypotension:** Sustained drop in systolic BP ≥ 20 mmHg or diastolic ≥ 10 mmHg within 3 minutes of tilt

ME/CFS relevance Orthostatic intolerance is present in 50–90% of ME/CFS patients. Tilt table testing provides objective documentation and identifies the specific subtype, guiding treatment selection.

Alternative Active standing test (10-minute stand with serial HR and BP measurement) provides a simpler office-based screen, though less standardized.

C.3.3 Autonomic Function Tests

Additional autonomic assessments complement tilt table testing.

Heart rate variability (HRV) Analysis of beat-to-beat heart rate intervals from ECG or heart rate monitor recordings. Time-domain (SDNN, RMSSD) and frequency-domain (LF, HF, LF/HF ratio) metrics quantify sympathovagal balance. ME/CFS patients typically show reduced HRV, reduced HF power (parasympathetic), and elevated LF/HF ratio (sympathetic predominance) [394].

Valsalva maneuver Forced expiration against a closed glottis (40 mmHg for 15 seconds). The heart rate and blood pressure response has four phases testing baroreflex integrity. Abnormal responses indicate baroreflex failure.

Deep breathing test Heart rate response to deep breathing at 6 breaths/minute. The expiratory–inspiratory HR difference (E:I ratio) quantifies parasympathetic function. Reduced in ME/CFS.

C.4 Laboratory Tests

ME/CFS is currently a clinical diagnosis without a pathognomonic laboratory test. Laboratory testing serves two purposes: (1) excluding alternative diagnoses and (2) identifying treatable comorbidities.

C.4.1 Standard Exclusion Panel

The following tests are recommended to exclude conditions that can mimic ME/CFS [127, 122]:

- **Complete blood count:** Excludes anemia, leukemia, infection
- **Comprehensive metabolic panel:** Excludes renal/hepatic disease, electrolyte disorders
- **Thyroid function (TSH, free T4):** Excludes hypothyroidism/hyperthyroidism
- **Erythrocyte sedimentation rate (ESR) and C-reactive protein (CRP):** Screens for active inflammatory/autoimmune conditions
- **Ferritin:** Excludes iron deficiency (common comorbidity)
- **Vitamin B12 and folate:** Excludes deficiency-related fatigue
- **Cortisol (morning or ACTH stimulation test):** Excludes Addison's disease
- **Celiac serology:** Excludes celiac disease
- **Urinalysis:** Screens for renal disease, diabetes

C.4.2 Extended Panel for Clinical Characterization

Additional tests characterize disease mechanisms but are not required for diagnosis:

- **Immunoglobulin levels (IgG, IgA, IgM):** Identifies humoral immune deficiency
- **Lymphocyte subsets (CD4, CD8, NK cells):** Characterizes immune cell populations
- **NK cell function assay:** Quantifies cytotoxic activity (research setting)
- **Autoantibody panel (β_2 -adrenergic, muscarinic M3/M4 receptor antibodies):** Identifies autoimmune subtype (specialized laboratories)
- **Viral serology (EBV VCA IgG/IgM, EBNA, HHV-6 IgG):** Documents prior/active herpesvirus infection
- **Tryptase:** Screens for mast cell activation
- **Vitamin D:** Common deficiency in housebound patients
- **Sleep study (polysomnography):** Excludes primary sleep disorders (obstructive sleep apnea, narcolepsy)

C.4.3 Experimental Biomarkers

No validated diagnostic biomarker for ME/CFS exists, but several candidates are under investigation:

- **Metabolomics panels:** Reduced amino acids, altered lipid profiles, elevated lactate-to-pyruvate ratios [589, 139]
- **Cytokine panels:** Disease-duration-dependent profiles [212, 213]
- **Epigenetic markers:** DNA methylation signatures at specific CpG sites [76]
- **Two-day CPET:** While not a blood test, the day-2 decrement serves as a functional biomarker (Section C.3)

△ Warning 1: No Single Diagnostic Test Exists

ME/CFS remains a clinical diagnosis based on symptom patterns (post-exertional malaise as the cardinal feature) after exclusion of alternative explanations. No laboratory test, imaging study, or biomarker has been validated for routine diagnostic use. Patients and clinicians should be cautious of commercial tests marketed as ME/CFS diagnostics that have not undergone rigorous validation.

D Resources and Support

This appendix provides a comprehensive guide to ME/CFS patient organizations, online communities, advocacy resources, and prominent patient voices. Unlike the scientific literature cited throughout this document, these resources represent the patient community's perspective and lived experience—an essential complement to clinical and research knowledge.

Observation 108 (On Patient-Generated Knowledge). The ME/CFS patient community has developed sophisticated knowledge networks that often outpace formal medical understanding. Forums like Phoenix Rising and Science for ME regularly discuss research papers with clinical depth that would be impressive in any medical setting. Patient advocates have successfully challenged flawed research (notably the PACE trial), influenced government policy, and funded significant research initiatives. This appendix acknowledges that patient expertise is a legitimate and valuable form of knowledge.

D.1 International Patient Organizations

D.1.1 Major Global Organizations

Solve ME/CFS Initiative (Solve M.E.)

<https://solvecfs.org/>

US-based non-profit serving as a catalyst for research into ME/CFS, Long COVID, and other infection-associated chronic conditions. Operates the You+ME patient registry and biobank, funds research, and conducts policy advocacy. One of the largest and most influential ME/CFS organizations globally.

Open Medicine Foundation (OMF)

<https://www.omf.ngo/>

Founded by Linda Tannenbaum, whose son has ME/CFS. Directs the Scientific Advisory Board chaired by Ron Davis, PhD. Operates six international ME/CFS Collaborative Research Centers and funds the End ME/CFS Project. Known for patient-centered research approach and involvement of patients and family members at leadership level.

International Association for CFS/ME (IACFS/ME)

<https://www.iacfsme.org/>

International non-profit organization of clinicians, scientists, professionals, patients, and advocates. Publishes the peer-reviewed journal *Fatigue: Biomedicine, Health, and Behavior*, organizes international conferences, and promotes science-based care.

#MEAction Network

<https://www.meaction.net/>

International patient-led advocacy network fighting for health equality. Co-founded by filmmaker Jennifer Brea. Organizes the annual #MillionsMissing protests, provides

advocacy training, and supports local patient groups worldwide. Known for effective use of social media and grassroots organizing.

World ME Alliance

<https://worldmealliance.org/>

Global coalition of national ME organizations working to coordinate international advocacy efforts and share resources across countries.

D.1.2 European Organizations

European ME Alliance (EMEA)

<https://www.europeanmealliance.org/>

Pan-European patient organization representing 18 countries. Founded in 2008 as collaboration of national patient charities. Member of European Patients' Forum (EPF) and European Federation of Neurological Associations (EFNA). Conducts the Pan-European ME Patient Survey (over 11,000 respondents in 2024). Created the European ME Research Group (EMERG) and European ME Clinicians Council.

EUROMENE

<https://www.euromene.eu/>

European Network on ME/CFS—a COST (European Cooperation in Science and Technology) supported network of research groups across Europe. Published expert consensus on diagnosis, service provision, and care in Europe (2021).

D.2 National Patient Organizations by Country

D.2.1 Belgium

Remark 1 (Belgian Organizations). Belgian organizations are primarily Flemish-based, with limited French-language resources for Wallonia. In Belgium, the condition is typically referred to as “CVS” (Chronisch Vermoeidheidssyndroom) rather than ME. ◇

ME-Vereniging vzw

<https://www.me-vereniging.be/>

The ME Association (Belgium) raises awareness and strives for recognition of the disease. Organizes support groups in Antwerp, Hasselt, Ypres, and Nieuwrode. Operates an ME help-line. Co-founder of the European ME Alliance.

12ME

Belgian non-profit drawing attention to ME/CFS seriousness with a positive approach.

CVS contact groep vzw

<http://www.cvs-contactgroep.be/>

Aims at CFS and fibromyalgia patients, provides information on legitimate scientific research. Publishes quarterly magazine “Immune” and organizes meetings in Flanders.

RIZIV/INAMI-Recognized Diagnostic Centers

Historical Context Belgium originally established **five CVS reference centers** around 2002 at university hospitals (UZ Leuven, UZ Gent, UZ Antwerpen, UZ Brussel, and one in Wallonia). These centers operated until **2012**, when RIZIV abruptly cut funding. In 2014, a new system of “Multidisciplinary Diagnostic Centers” replaced them, but with reduced scope. As of 2024, only one center has signed the current convention.

UPC KU Leuven – Multidisciplinair Diagnostisch Centrum ME/CVS

Address: Leuvensesteenweg 517, 3070 Kortenberg

Phone: +32 2 758 05 11 (general); +32 2 758 16 77 (CVS consultation)

Email: cvs@upckuleuven.be

Website: <https://www.upckuleuven.be/nl/zorgaanbod/cvs>

Consultation hours: Tuesday 9:00–12:00, Wednesday 9:30–12:00 (by appointment)

As of 2024, this is the only center in Belgium with an official RIZIV/INAMI convention for ME/CVS. The center provides multidisciplinary diagnostic assessment and, if ME/CVS is confirmed, develops a care trajectory in collaboration with the patient’s GP. Referral must come from a GP who suspects ME/CVS. Convention runs until 2028.

Remark 2 (CBT-Based Treatment: Critical Context). The RIZIV/INAMI convention mandates cognitive behavioral therapy (CBT) and graded exercise therapy (GET) as the reimbursed treatments—an approach based primarily on the UK PACE trial, which has been heavily criticized for methodological flaws and subsequently disavowed by NICE guidelines (2021). Large patient surveys consistently report that GET worsens symptoms in the majority of ME/CFS patients, while CBT shows limited benefit for core symptoms. Belgian patient organizations have criticized this policy as outdated and potentially harmful. Patients should be aware that accepting the convention’s treatment pathway means committing to CBT/GET-based rehabilitation, which may not align with current international best practices emphasizing pacing and symptom management.

The convention offers reimbursement for up to 17 CBT sessions (individual: 50 min at €86.69; group: 90 min at €57.80 per participant—2024 rates). Maximum 8 sessions may be in group format. Contact RIZIV: Evi Declercq, +32 2 739 71 97, evi.declercq@riziv-inami.fgov.be. ◇

Other Clinical Resources

UZ Leuven

<https://www.uzleuven.be/nl/chronisch-vermoeidheidssyndroom-cvs>

University Hospital Leuven provides care for pronounced fatigue lasting more than 6 months that doesn’t improve with rest and significantly limits daily activities.

UZA – Centrum voor Gedragstherapie bij Vermoeidheid

<https://www.uza.be/behandeling/chronisch-vermoeidheids-syndroom-cvs>

University Hospital Antwerp’s Center for Behavioral Therapy for Fatigue (CGVF) treats patients with fatigue and functional complaints including CVS, chronic fatigue, fibromyalgia, and post-infectious fatigue.

UZ Gent CVS Network

UZ Gent coordinates an integrated care model for abnormal fatigue in East and West Flanders. Partners include AZ Alma, AZ Groeninge, Jan Yperman Ziekenhuis, AZ Maria Middelaers, AZ Sint-Jan, AZ Nikolaas, AZ Delta, and others. The GP serves as care process manager.

Sleep Medicine (Wallonia)

Centre Multidisciplinaire de Somnologie – Clinique Saint-Luc Bouge

Address: Rue Saint-Luc 8, 5004 Bouge (Namur)

Phone: +32 81 20 94 61

Email: labosommeil@slbo.be

Website: <https://slbo.be/services/centres-integres-et-pluridisciplinaires/centre-multidisciplinaire-de-somnologie/>

Established in 1993, this is a major French-language sleep medicine center in Wallonia. INAMI-accredited since 2002 for CPAP treatment and since 2018 for mandibular advancement devices. The multidisciplinary team includes pneumologists, neurologists, psychiatrists, ORL specialists, and psychologists. Accepts patients from age 15 with any sleep disorder. As of September 2025, waiting times reduced to approximately 6 weeks (compared to 6–8 months at many Belgian hospitals). Chief of Service: Dr Richard Frogner.

Remark 3 (Sleep Disorders and ME/CFS). While sleep clinics do not diagnose ME/CFS directly, they are valuable for ruling out primary sleep disorders (sleep apnea, narcolepsy, etc.) that can cause chronic fatigue, and for documenting the non-restorative sleep characteristic of ME/CFS. A sleep study may be part of the differential diagnosis workup. ◇

D.2.2 France

ASFC (Association française du Syndrome de Fatigue Chronique)

<https://www.asso-sfc.org/>

The only ME/CFS patient association approved by the French Ministry of Health (2015). Located in Lille. Works with a Scientific Board to welcome, inform, and support patients. Operates phone hotline, organizes regular patient meetings throughout France, and annual meetings with expert scientists.

EM Action France

French website reporting international ME news and research.

Millions Missing France

<https://millionsmissing.fr/>

French chapter of the #MillionsMissing movement.

Remark 4 (French Recognition). ME/CFS is not officially recognized by the French Department of Health, leading to under-diagnosis and lack of disability recognition. ◇

D.2.3 Luxembourg

No dedicated ME/CFS patient organization exists in Luxembourg as of 2025. Patients may connect with:

- Belgian or French organizations (French-language)
- German organizations (German-language)
- EUROMENE network for research connections
- European ME Alliance for pan-European advocacy

D.2.4 Germany

Deutsche Gesellschaft für ME/CFS

<https://www.mecfs.de/>

German Society for ME/CFS, founded in 2016, based in Hamburg. Run by volunteers advocating for patient rights and medical/social recognition. Organized the first Parliamentary Expert Discussion on ME/CFS in the Bundestag (March 2020). Successfully lobbied for ME/CFS and Long COVID mention in federal government coalition agreement (2021). Successfully challenged the German “Tiredness” guideline’s recommendations for GET and CBT.

Fatigatio e.V.

<https://www.fatigatio.de/>

Federal Association ME/CFS, founded in 1993, based in Berlin. Over 2,900 members. Operates 15 regional self-help groups across German cities. Organizes annual hybrid ME/CFS conference with national and international experts.

Remark 5 (German Prevalence). Germany has seen dramatic growth in ME/CFS cases post-COVID: from an estimated 250,000 pre-pandemic to over 650,000 by end of 2024. ◇

D.2.5 Netherlands

ME/cvs Vereniging

<https://www.me-cvs.nl/>

Dutch association founded in 2005. The lowercase “cvs” deliberately underscores the desire for the medical world to stop using the name “Chronic Fatigue Syndrome.” Involved in international partnerships including UK ME/CFS Biobank, Solve ME, Open Medicine Foundation, and Charité Berlin.

ME Vereniging Nederland

Founded 2011. Membership open only to ME patients. Focus on improving living conditions and reducing social exclusion.

ME/ CVS-Stichting Nederland

Founded 1987. Receives government funding as recognized national ME/CFS patient organization.

Steungroep ME en Arbeidsongeschiktheid

Founded 1994. Support group focused on employment, education, disability, and benefits issues. Campaigns against exclusion from disability benefits.

The **Netherlands ME/CFS Cohort and Biobank (NMCB)** consortium is a national collaboration of research institutes, patient organizations, and clinical centers establishing a comprehensive patient cohort and biobank.

D.2.6 United Kingdom

ME Association

<https://meassociation.org.uk/>

One of the two largest UK ME/CFS charities. Provides information, advocacy, and services. Publishes quarterly magazine *ME Essential*. Funds the UK ME/CFS Biobank. Conducted major patient surveys (2010, 2015) documenting treatment experiences. Hosts local support groups nationwide.

Action for ME

<https://www.actionforme.org.uk/>

Founded 1987 as The M.E. Action Campaign. Merged with Association of Young People with ME in 2017. Funds high-quality research including the groundbreaking DecodeME study (largest ME/CFS genetic study ever, 15,000+ UK participants). Offers free support services including holistic healthcare services.

BACME (British Association of Clinicians in ME/CFS)

<https://bacme.info/>

Multidisciplinary organization for UK healthcare professionals delivering care to ME/CFS patients.

D.2.7 Ireland

Irish ME/CFS Association

<https://www.irishmecfs.org/>

Works to improve the situation for people with ME/CFS in Ireland. Notable advocate: Tom Kindlon (Assistant Chairperson), who has been housebound with severe ME for over 22 years. Known internationally for his extensive analysis and publications on the PACE trial and harms of graded exercise therapy.

Hope for ME & Fibro Northern Ireland

Founded 2011 by Joan McParland MBE, inspired by Tom Kindlon's work.

D.2.8 Norway

Norges ME Forening

<https://www.me-foreningen.no/>

Norwegian ME Association, founded 1987 by Ellen Piro. Represents over 6,000 ME

patients. Party-political independent organization. Works to ensure diagnosis based on Canadian Consensus Criteria. Member of European ME Alliance.

ME-Fondet

<https://www.me-fondet.no/>

Norwegian non-profit foundation dedicated to funding biomedical ME research. Supporting a promising daratumumab pilot study at Haukeland University Hospital.

Remark 6 (Norwegian Research Leadership). Norway has been a leader in ME/CFS research, pioneering innovative treatment approaches including the rituximab trials and subsequent work on autoimmunity. ◇

D.2.9 Denmark

ME Foreningen (Danish ME Association)

<https://me-foreningen.dk/>

National association since 1992. Works to increase knowledge in the Danish health-care system about ME as a physical/biomedical disease. Achieved unanimous Danish parliament vote to separate ME (ICD-10 G93.3) from Functional Disorders. Counseling available Wednesday/Friday 12–14 at +45 44 95 97 00.

D.2.10 Sweden

Riksföreningen för ME-patienter (RME)

Swedish national association for ME patients. Member of European ME Alliance.

D.2.11 Switzerland

ME/CFS Verein Schweiz

<https://www.mecfs.ch/>

Self-help organization founded 1993 in Zurich. Offers information platform, networking, and support. Hosts regular group meetings in several Swiss cities.

Schweizerische Gesellschaft für ME & CFS

<https://sgme.ch/>

Swiss Association for ME & CFS, founded 2019. Fights for recognition and adequate care. Conducts biennial comprehensive surveys on Swiss ME patient situations. First analysis published 2021.

Remark 7 (Swiss Diagnostic Challenges). Swiss research shows mean diagnosis time of 6.7 years, average 11.1 different appointments, 2.6 misdiagnoses, and 13.5% of patients traveling abroad to seek diagnosis. 90.5% of patients were told at least once that symptoms were psychosomatic. ◇

D.2.12 Spain

CONFESQ

<http://confederacion-fm-sfc.es/>

National Coalition of FM, CFS/ME, MCS, and EHS. Established 2004. Based in Jerez de la Frontera.

ONG-PEM (Asociación de Personas con Encefalomiелitis Miálgica)

Founded and run by severely ill patients. Exclusively represents Myalgic Encephalomyelitis.

Associació Catalana d'Afectats SFC/EM

<http://www.acsfcem.org/>

Catalonia-based patient association.

Spanish Facebook groups include VIVIR CON SFC/EM and #MillonesAusentes (Spanish #MillionsMissing).

D.2.13 Italy

CFS/ME Associazione Italiana

<http://www.stanchezzacronica.it/>

Founded 1991 by Prof. Umberto Tirelli in Udine—first Italian physician to identify CFS cases. Based at Centro di Riferimento Oncologico, Aviano.

Associazione Malati di CFS ODV

<http://www.associazioneecfs.com/>

Patient advocacy group founded 2004, based in Pavia. Part of Rare Disease Alliance (Alleanza delle Malattie Rare). Celebrated 20 years in 2024.

CFS/ME Organizzazione di Volontariato

<https://www.cfsme.it/>

Veneto-based patient organization.

CFS Italia Forum (<http://www.cfsitalia.it/>) provides Italian-language patient community and information exchange.

D.2.14 Australia

Emerge Australia

<https://emerge.org.au/>

National organization providing services, evidence-based education, advocacy, and research. Free national health and support line: 1800 865 321 (9am–4:30pm Mon–Fri). Offers online patient/carer education, peer support groups, RACGP CPD-approved healthcare professional education. Partners with Solve ME on AusME patient registry and biobank.

ME/CFS Australia

<https://mecfs.org.au/>

Peak body for patient-led ME/CFS charities. Focuses on federal government advocacy, research initiatives, and national awareness campaigns.

D.2.15 United States

American ME and CFS Society (AMMES)

<https://ammes.org/>

Serves patients and caregivers through support, advocacy, and education. Channels patient perspectives to government agencies and initiatives. Comprehensive website with links to international organizations.

U.S. ME/CFS Clinician Coalition

<https://mecfscliniciancoalition.org/>

Provides resources for medical providers caring for ME/CFS patients. Developed clinical guidance documents.

Bateman Horne Center

<https://batemanhornecenter.org/>

Medical center of excellence for ME/CFS and fibromyalgia. Founded by Dr. Lucinda Bateman. Focuses on diagnosis, treatment, research, and patient empowerment.

D.3 Research Centers and Specialized Clinics

D.3.1 Leading Research Centers

Stanford ME/CFS Collaborative Research Center

<https://med.stanford.edu/chronicfatiguesyndrome/>

Established 2013, directed by Ron Davis, PhD. Part of Stanford Genome Technology Center. Focus on developing objective diagnostic tests and treatments. Known for nanoneedle diagnostics development.

Columbia Center for Infection and Immunity

Columbia University.

Directed by W. Ian Lipkin, MD. ME/CFS research focus on infectious triggers and immune dysfunction.

Cornell ME/CFS Center for Enervating Neurolimmune Disease

Directed by Maureen Hanson, PhD. Research on mitochondrial function, immune cells, and microbiome.

Charité Fatigue Centrum

Berlin, Germany. <https://cfc.charite.de/>

Major European ME/CFS research and clinical center. Led by Prof. Carmen Scheibenbogen.

Uppsala University ME/CFS Collaboration

Sweden.

Led by Jonas Bergquist, MD, PhD. Focus on neurochemistry and analytical approaches.

D.3.2 Specialized Clinical Centers

Bateman Horne Center

Salt Lake City, Utah.

Clinical care with research integration. Founded by Dr. Lucinda Bateman.

Open Medicine Clinic

Mountain View, California.

Run by Dr. David Kaufman. Known for complex chronic illness expertise.

Haukeland University Hospital

Bergen, Norway.

Site of rituximab trials and ongoing autoimmunity research.

D.4 Online Communities and Forums

D.4.1 Discussion Forums

Phoenix Rising

<https://phoenixrising.me/> and <https://forums.phoenixrising.me/>

Founded by Cort Johnson. One of the most visited ME/CFS websites. Approximately 19,000 member accounts with 600 daily active members (2017). Covers dysautonomia, hormones, methylation, lifestyle management, relationships, and caregiver support.

Science for ME (S4ME)

<https://www.s4me.info/>

Independent, patient-led, international forum. Founded by Andy Devereux-Cooke. Each thread typically dedicated to a single research paper, enabling in-depth discussion. Notable members have included Jonathan Edwards, Tom Kindlon, Simon McGrath, and David Tuller. Advocates for patients as research partners.

Health Rising Forums

<https://www.healthrising.org/forums/>

Companion to Health Rising blog. Discussion of ME/CFS, fibromyalgia, chronic pain, IBS, and dysautonomia.

MEpedia

<https://me-pedia.org/>

Crowd-sourced encyclopedia of ME/CFS science and history. Creative Commons licensed. Categories include notable patients, citizen scientists, and ME/CFS history. Valuable reference for terminology, research summaries, and advocacy history.

D.4.2 Reddit Communities

r/cfs

<https://www.reddit.com/r/cfs/>

Primary ME/CFS subreddit. Research discussions, treatment experiences, and personal support. Active moderation maintaining distinction between ME/CFS and chronic fatigue symptom.

r/covidlonghaulers

Related community for Long COVID with significant ME/CFS overlap.

D.4.3 Facebook Groups

Major ME/CFS Facebook groups (search on Facebook):

- Chronic Fatigue Syndrome & Myalgic Encephalomyelitis ME Self Help Group (founded 2012)
- #MEAction state/regional chapters
- Pregnancy and Parenting with ME/CFS
- Caregiver Support groups
- Severe ME support groups
- Youth ME/CFS support (ages 13–21)

D.4.4 Other Platforms

Smart Patients ME/CFS Community

<https://www.smartpatients.com/communities/me-cfs>

Peer-to-peer support where patients and families share experiences and research.

NURA

Social network platform specifically for Long COVID, ME/CFS, and fibromyalgia patients, created by people with these conditions.

D.5 Prominent Patient Advocates and Content Creators

Observation 109 (The Patient Expertise Network). ME/CFS advocacy is largely driven by patients themselves, often working with extremely limited energy. The individuals listed here represent a fraction of the patient community dedicating their scarce functional capacity to improving conditions for all patients. Many severely ill patients contribute via social media, writing single tweets or posts that may represent their entire energy expenditure for a day.

D.5.1 Filmmakers and Documentarians

Jennifer Brea

Background American documentary filmmaker and activist. PhD student at Harvard when sudden illness left her bedridden.

Key work *Unrest* (2017)—Sundance award-winning documentary, Emmy-nominated, short-listed for Academy Award. Available free on YouTube (May 2023). Produced largely from bed, directing remotely with crews worldwide.

Advocacy Co-founder of #MEAction. Delivered highest-rated TED Talk at 2016 TED Summit (nearly 2 million views, 25+ languages).

Personal journey Later discovered craniocervical instability (CCI) and underwent spinal fusion surgery, experiencing significant improvement—highlighting ME/CFS subgroup heterogeneity.

Website <https://www.jenniferbrea.com/>

Dianna Cower (Physics Girl)

Background Science educator, YouTube channel with 2.8+ million subscribers.

Illness Contracted COVID-19, developed Long COVID/ME/CFS. Currently completely bed-bound, unable to care for herself. Also developed MCAS.

Advocacy 12-hour livestream (July 6, 2024) showing “a day in her life” with severe ME/CFS. Co-hosted by Ian Hecox and Simone Giertz. Raised \$150,000+ for Open Medicine Foundation. Livestream became top post on r/videos (27M subscribers).

Impact Brought ME/CFS awareness to mainstream audience unfamiliar with the condition.

Platform YouTube: Physics Girl; Twitter/X: @thephysicsgirl

D.5.2 Writers and Bloggers

Cort Johnson

Background Developed ME/CFS/FM in 1980s while in Environmental Studies program at UC Santa Cruz. MS in Environmental Studies from San Jose State University (2000).

Key work Founded Phoenix Rising (2004)—became most visited ME/CFS website by 2010. Left to found Health Rising (2012), broadening focus to include fibromyalgia. Produced 1000+ comprehensive blogs on ME/CFS and FM.

Recognition ProHealth’s Advocate of the Year (2015). IACFS/ME Special Services Award (2016). Described as “the quintessential patient advocate, breaking more news about this illness than many professional journalists.”

Website <https://www.healthrising.org/>

Social media Twitter/X: @CortJohnson

Jamison Hill

Background Former bodybuilder and certified personal trainer at Sonoma State University. Developed ME after mononucleosis in senior year (2010).

Illness severity By age 28, bedridden, unable to speak, eat solid food, or elevate body. Wrote on cellphone wearing tanning goggles to block light.

Publications *When Force Meets Fate: A Mission to Solve an Invisible Illness* (2021 memoir). Written for The Washington Post, The New York Times, Los Angeles Times, Men’s Journal, Vox, VICE, and others.

Media Featured in *Forgotten Plague* documentary, Netflix series (2018), WBUR Modern Love podcast, Dax Shepard’s Armchair Expert podcast.

Current status Improved with anti-virals, hydrocortisone, IV saline—not fully recovered but able to tell his story.

Website <https://jamisonwrites.com/>

Whitney Dafoe

Background Son of Dr. Ron Davis (Stanford geneticist) and Dr. Janet Dafoe. Former adventurer and photographer who traveled to all 50 states, India, Nepal, Ecuador.

Diagnosis ME/CFS diagnosed 2010.

Current severity One of the most severe ME/CFS cases documented. Cannot speak. Cannot tolerate contact with anyone but parents due to visual dysfunction. Fed by tube directly into stomach. Hasn't spoken in years.

Advocacy Despite severity, maintains blog, Facebook page, Instagram documenting life with severe ME/CFS. Won Gold at European Photography Awards (2022) for documentary series "The Living Death."

Impact His illness catalyzed his father's complete redirection of research focus: "I decided to terminate everything I was working on before Whitney got sick. Everything is ME/CFS now."

Website <https://www.whitneydafoe.com/>

Patreon <https://www.patreon.com/whitneydafoe>

Other Notable Bloggers

Suzan Jackson (Live with CFS)

Has ME/CFS since 2002; both sons also developed ME/CFS at ages 6 and 10. Blog focuses on living well despite chronic illness.

Mary M. Schweitzer, PhD (Slightly Alive)

Former history professor. Maintains ME and CFS Information Page with essays, reports, and conference summaries.

Naomi Whittingham (A Life Hidden)

UK-based, severe ME since age 12. Does interviews and supports brother Tom Whittingham's marathon fundraising for ME Research UK.

Laura's Pen

Blog covering Lyme disease, ME/CFS, and endometriosis awareness.

Super Pooped

ME/CFS awareness through art, crafts, and humor.

D.5.3 Researchers Who Are Patients or Family Members

Ron Davis, PhD

Position Professor of Biochemistry and Genetics, Director of Stanford Genome Technology Center.

Background Pioneered technology that powered the Human Genome Project. Over 64 biotechnology patents.

Personal connection Son Whitney has very severe ME/CFS.

Research pivot “I decided to terminate everything I was working on before Whitney got sick. Everything is ME/CFS now.”

Leadership Director of OMF Scientific Advisory Board. Established Stanford ME/CFS Collaborative Research Center (2013). His work helped prove ME/CFS is a biological disease.

Tom Kindlon

Background Very active young man (soccer, tennis, cricket, cross-country) until ME at age 16 (February 1989).

Current status Housebound for 22+ years. Uses wheelchair. Full-time carer: his mother Vera.

Expertise Studied Mathematical Sciences at Trinity College Dublin before dropping out. Extensive analysis and publications on PACE trial and harms of graded exercise therapy. Work available on ResearchGate and PubMed.

Role Assistant Chairperson, Irish ME/CFS Association.

Recognition OMF certificate of merit. Described as “a leader in the global ME/CFS community” who “initiated patient-led efforts to take a scientific approach to analyzing ME/CFS research.” Nominated for honorary degree at Trinity College Dublin.

Andy Devereux-Cooke

Role Patient, founder of Science for ME forum.

Research Research investigator on DecodeME study—demonstrating patient partnership in research.

D.5.4 Celebrity Advocates

Laura Hillenbrand

Author of *Seabiscuit* and *Unbroken*. Candid about ME/CFS struggles and medical misunderstanding.

Karin Alvtegen

Scandinavian author of psychological thrillers (*Missing*, *Betrayal*). ME/CFS has significantly shaped her life and career.

D.5.5 Euthanasia and End-of-Life Discussions

Remark 8 (Community Discourse on Quality of Life). The severity of ME/CFS has led to difficult conversations within the patient community about quality of life and end-of-life options. Some patients with very severe ME/CFS have publicly discussed or pursued medical assistance in dying in countries where it is legal (Belgium, Netherlands, Switzerland, Canada). These discussions reflect the profound suffering experienced by the most severely affected patients and the lack of effective treatments. ◇

- **Samuel (Austria, 2004–2026)** – Samuel developed very severe ME/CFS following a COVID-19 infection. In his final public statement, posted to Reddit 12 days before his death, he described his condition: “I must lie in bed 24 hours a day and cannot move too much, it must be permanently dark because I cannot tolerate light. I wear double hearing protection because I also cannot tolerate sounds. I cannot watch television or videos for even a second, as moving images overwhelm my nervous system and trigger unbearable suffering. I cannot listen to music or podcasts. I cannot even talk to my own mother, who cares for me, because listening is too exhausting and speaking has become completely impossible. I must communicate with pen and paper.” He described the physical suffering as “like drowning and burning at the same time” and noted that any exertion beyond his energy limits caused crashes leading to permanent deterioration. He highlighted systemic failures: no approved medications exist for ME/CFS, Austria has no dedicated treatment centers despite 1% of the population being affected, most physicians do not recognize the disease, and the pension authority (PVA) called patients “charlatans and freeloaders.” Samuel chose medical assistance in dying in Austria, passing away on January 30, 2026—his 22nd birthday. His final message: “ME/CFS kills!” (*ME/CFS TÖTET!*)

Sources: Original Reddit post: <https://www.reddit.com/r/Austria/comments/1qg8oit/>; News coverage: <https://www.heute.at/s/samuel-21-ist-tot-er-starb-an-seinem-geb>

- **Austrian Policy Discussion** – Reddit r/Austria community discussion on euthanasia policies: <https://www.reddit.com/r/Austria/s/JUzy5LM607>

D.6 Podcasts

The Understanding ME/CFS Podcast

Apple Podcasts, Spotify

Hosted by Patrick Ussher (7-year ME/CFS patient, author of *Understanding ME/CFS & Strategies for Healing*). Weekly interviews with patients and experts. Covers research, treatments, quality of life, and recovery stories.

Chronically Complex: The #MEAction Podcast

<https://www.meaction.net/chronically-complex-meaction-podcast/>

Interviews influential voices in ME/CFS and Long COVID. Topics include books on complex chronic disease, #MillionsMissing, #StopRestPace, disability activism, and art from disabled artists. Notable guests: Ryan Prior (CNN journalist, *Forgotten Plague* filmmaker), Cynthia Adinig (Long COVID advocate, SolveME board member).

CFS Unravelled

By Dan Neuffer (recovered from ME/CFS). Interviews with recovered patients and expert practitioners.

Discomfort Zone (Invisible Not Broken)

Hosted by Jason, engineering graduate who developed fibromyalgia, ME/CFS, and POTS. Each episode explores what it means to be chronically ill and disabled.

This Podcast Will Kill You – Episode 137

“ME/CFS: What’s in a name? (A lot, actually)” (April 2024). Deep dive into biology, history, and current research.

Hope and Help for Fatigue & Chronic Illness

Mission to help people with post-viral syndromes including Long COVID and ME/CFS.

D.7 YouTube Channels and Video Resources

D.7.1 Patient-Focused Channels

CFS Health

32.7K subscribers. Founded by Toby Morrison. Multi-dimensional approach to ME/CFS and fibromyalgia recovery.

CFS Unravelled

53.5K subscribers. Dan Neuffer shares insights on ME/CFS, POTS, and fibromyalgia healing.

Understanding ME-CFS

Patrick Ussher's channel accompanying his podcast.

D.7.2 Organization Channels

Open Medicine Foundation

Research updates, patient stories, educational content.

Solve ME/CFS Initiative

10K subscribers. Research and advocacy updates.

Bateman Horne Center

15.6K subscribers. Clinical education and patient resources.

MEAction

3.3K subscribers. Advocacy updates and #MillionsMissing content.

D.7.3 Documentaries

Unrest

(2017) Jennifer Brea. Available free on YouTube (since May 2023). Essential viewing for understanding patient experience.

Forgotten Plague

Co-directed by Ryan Prior. Features Jamison Hill and other patients.

What About ME?

Earlier documentary on ME/CFS.

Hope to our Hands

(2020) Documentary about ME/CFS patients in Japan struggling for acknowledgment.

Living with Chronic Fatigue Syndrome

German/French documentary premiered on ARTE. Available in German and French.

D.8 Social Media Hashtags and Campaigns

D.8.1 Key Hashtags

#MillionsMissing	Primary advocacy hashtag for global protests
#MECFS	Standard disease hashtag
#pwME	“People with ME”
#SevereME	Focusing on severe/very severe patients
#MyalgicE	Short form for myalgic encephalomyelitis
#May12th	ME Awareness Day
#StopRestPace	Pacing advocacy
#TeachMETreatME	2024 campaign theme
#LongCovid	Related condition with significant overlap

D.8.2 #MillionsMissing Campaign

Annual global campaign for ME health equality, organized by #MEAction. May 12th is ME Awareness Day—patients gather (in-person and virtually) to demand recognition, research, and clinical care. In 2025, communities joined at the U.S. Capitol to advocate for protecting Medicaid, home care support, research funding, and open science.

The campaign highlights the “millions missing” from their own lives due to illness, and the millions of research dollars missing from funding.

D.9 Books by Patients and Advocates

D.9.1 Patient Memoirs

When Force Meets Fate: A Mission to Solve an Invisible Illness

Jamison Hill (2021). Former bodybuilder’s journey through severe ME/CFS.

The Puzzle Solver: A Scientist’s Desperate Quest to Cure the Illness that Stole His Son

Tracie White and Ron Davis (2021). Story of Ron Davis and Whitney Dafoe.

Understanding ME/CFS & Strategies for Healing

Patrick Ussher. Guide by a patient, companion to podcast.

The Long Haul

Ryan Prior. On Long COVID and ME/CFS advocacy.

D.9.2 Clinical and Reference Works

See also the scientific literature cited throughout this document. Patient organizations often maintain curated reading lists of accessible scientific overviews.

D.10 Clinical Trial Registries

ClinicalTrials.gov

<https://clinicaltrials.gov/>

Search for “myalgic encephalomyelitis” or “chronic fatigue syndrome.” Filter by recruiting status and location.

EU Clinical Trials Register

<https://www.clinicaltrialsregister.eu/>

European clinical trials database.

ME/CFS Research Register

<https://mecfs-research.org/>

Specialized registry tracking ME/CFS research internationally.

Questions to ask before participating:

- What are inclusion/exclusion criteria?
- What is the time commitment?
- Will travel be required? Is remote participation possible?
- What accommodations exist for severely ill participants?
- How will participant safety be monitored?
- Will results be shared with participants?

D.11 Patient Registries and Biobanks

You+ME Registry and Biobank

<https://youandme.solvecfs.org/>

Solve M.E.’s patient registry. Collects patient-reported data and biospecimens. International participation.

AusME Registry

Australian ME/CFS and Long COVID registry, partnership between Emerge Australia and Solve M.E.

UK ME/CFS Biobank

London School of Hygiene & Tropical Medicine. Funded partly by ME Association.

Netherlands ME/CFS Cohort and Biobank (NMCB)

National Dutch infrastructure for ME/CFS research.

DecodeME

UK-based genetic study with 15,000+ participants. Largest ME/CFS study ever conducted.

D.12 Disability and Legal Resources

D.12.1 General Guidance

Most national patient organizations provide country-specific guidance on:

- Disability benefits applications
- Workplace accommodations
- Educational accommodations
- Healthcare rights
- Insurance issues

D.12.2 Key Considerations

- ME/CFS is classified as a neurological disease by WHO (ICD-11: 8E49)
- Documentation of functional limitations is essential
- Some countries recognize ME/CFS for disability; others require extensive advocacy
- Patient organizations often provide template letters and case examples
- Legal advocacy organizations exist in some countries

The **Steungroep ME en Arbeidsongeschiktheid** (Netherlands) specifically focuses on employment and disability issues. The **ME Association** (UK) provides extensive guidance on UK benefits system.

D.13 Resource Evaluation Guidelines

Observation 110 (Navigating Information Quality). The ME/CFS information landscape includes high-quality patient-led resources alongside misinformation and exploitation. The patient community has developed sophisticated evaluation skills born of necessity.

D.13.1 Indicators of Reliable Resources

- Connection to established patient organizations
- Citation of peer-reviewed research
- Acknowledgment of uncertainty and ME/CFS heterogeneity
- Clear distinction between established knowledge and speculation
- Avoidance of cure claims
- Transparency about funding sources
- Respect for patient autonomy and pacing needs

D.13.2 Warning Signs

- Guaranteed cures
- Pressure to commit quickly or pay upfront
- Hostility to questions
- Rejection of biomedical model without evidence
- Promotion of graded exercise therapy without acknowledging PEM risks
- Claims that contradict major patient surveys
- No outcome data available
- Primarily selling products or services

Remark 9 (For Wallonia/Belgium Residents). French-language resources from ASFC (France), Swiss organizations, and Canadian French resources may supplement the primarily Flemish Belgian organizations. The European ME Alliance provides pan-European perspective and advocacy regardless of language. ◇

E Mathematical Model Details

This appendix provides the complete mathematical specifications for the models developed in Part V, including full parameter tables, derivations of key results, numerical methods, and sensitivity analysis procedures. Readers should consult the corresponding chapters for biological context and interpretation.

E.1 Complete Model Equations

E.1.1 Energy Metabolism Model

The energy metabolism model (Chapter 35) comprises 8 state variables. The complete ODE system is:

$$\frac{d[\text{ATP}]}{dt} = 2J_{\text{glyc}} + J_{\text{ATPsyn}} - J_{\text{demand}}(t) \quad (\text{E.1})$$

$$\frac{d[\text{ADP}]}{dt} = J_{\text{demand}}(t) - 2J_{\text{glyc}} - J_{\text{ATPsyn}} + J_{\text{AK,fwd}} - J_{\text{AK,rev}} \quad (\text{E.2})$$

$$\frac{d[\text{AMP}]}{dt} = J_{\text{AK,rev}} - J_{\text{AK,fwd}} \quad (\text{E.3})$$

$$\frac{d[\text{NADH}]}{dt} = 2J_{\text{glyc}} + 3J_{\text{Krebs}} - J_{\text{CI}} \quad (\text{E.4})$$

$$\frac{d[\text{Pyr}]}{dt} = J_{\text{glyc}} - J_{\text{PDH}} \quad (\text{E.5})$$

$$\frac{d\Delta\Psi}{dt} = \frac{1}{C_m}(4J_{\text{CI}} + 4J_{\text{CIII}} + 2J_{\text{CIV}} - n_{\text{ATP}}J_{\text{ATPsyn}} - J_{\text{leak}}) \quad (\text{E.6})$$

$$\frac{d[\text{ROS}]}{dt} = J_{\text{ROS}} - k_{\text{SOD}}[\text{SOD}][\text{ROS}] - k_{\text{GPx}}[\text{GPx}][\text{ROS}] \quad (\text{E.7})$$

$$\frac{dD}{dt} = k_D[\text{ROS}] \cdot \mathbb{I}_{[\text{ROS}] > \text{ROS}_{\text{thr}}} - k_{\text{rep}} \frac{[\text{ATP}]}{K_{\text{rep}} + [\text{ATP}]} \quad (\text{E.8})$$

where the proton stoichiometry coefficients (4, 4, 2) are the number of protons pumped per electron pair by Complexes I, III, and IV respectively (total 10 H⁺ per NADH), $n_{\text{ATP}} \approx 8/3$ is the H⁺/ATP ratio of ATP synthase, J_{AK} denotes the adenylate kinase equilibrium ($2\text{ADP} \rightleftharpoons \text{ATP} + \text{AMP}$, $K_{\text{eq}} \approx 0.44$), and J_{PDH} is the pyruvate dehydrogenase flux linking glycolysis to the Krebs cycle. The conservation constraints $[\text{ATP}] + [\text{ADP}] + [\text{AMP}] = A_{\text{total}}$ and $[\text{NAD}^+] + [\text{NADH}] = N_{\text{total}}$ reduce the independent variables to 6. The adenylate kinase equilibrium is maintained by setting $J_{\text{AK,fwd}} - J_{\text{AK,rev}} = k_{\text{AK}}(K_{\text{eq}}[\text{ADP}]^2 - [\text{ATP}][\text{AMP}])$.

E.1.2 Energy Metabolism Parameter Table

Table E.1 lists all parameters with healthy baseline values, ME/CFS ranges, units, and sources. Parameter values are drawn from published enzymology, metabolomics, and CPET data where available; parameters without direct experimental constraints are indicated.

Table E.1: Energy metabolism model parameters.

Parameter	Healthy	ME/CFS	Units	Source
$V_{\max, \text{PFK-1}}$	1.2	1.2	mM/min	Enzymology
$K_{0.5}^{\text{F6P}}$	0.1	0.1	mM	Enzymology
n_H (PFK-1)	2.5	2.5	–	Enzymology
K_i^{ATP}	1.0	1.0	mM	Enzymology
$V_{\max, \text{CI}}$	1.0	0.5–0.8	mM/min	[326]
K_m^{NADH}	0.05	0.05	mM	Enzymology
$\Delta\Psi_{\text{threshold}}$	110	110	mV	Biophysics
$J_{\text{leak},0}$	0.05	0.075–0.15	mM/min	Estimated
A_{total}	8.0	8.0	mM	Physiology
N_{total}	1.0	0.6–0.9	mM	[69]
k_{ROS}	0.01	0.01	min ⁻¹	Estimated
k_{SOD}	10.0	10.0	mM ⁻¹ min ⁻¹	Enzymology
k_D	0.001	0.001	min ⁻¹	Estimated
k_{rep}	0.005	0.005	min ⁻¹	Estimated

E.1.3 Immune System Model

The immune model (Chapter 36) comprises 12 state variables for cell populations and 6 for cytokines. The NK cell, T cell, and B cell dynamics are specified by Equations 36.1–36.12. The cytokine network is defined by the production and degradation terms in Equation 36.5. Full parameter tables for the immune model follow the same format as Table E.1; key parameters are listed below.

Table E.2: Selected immune model parameters.

Parameter	Healthy	ME/CFS	Units	Source
s_N (NK production)	50	50	cells/ μL /day	Physiology
$k_{\text{act},0}$ (NK)	0.2	0.2	day ⁻¹	Estimated
k_{exh} (NK)	0.05	0.1–0.2	day ⁻¹	[236]
k_{recov} (NK)	0.1	0.03–0.07	day ⁻¹	Estimated
$\delta_{\text{IL-6}}$	0.5	0.5	h ⁻¹	Pharmacokinetics
δ_{TNF}	0.7	0.7	h ⁻¹	Pharmacokinetics
d_P (plasma cell)	0.003	0.003	day ⁻¹	Immunology
λ_0 (reactivation)	0.001	0.005–0.02	day ⁻¹	Estimated

E.1.4 Neuroendocrine Model

The HPA axis (Equation 37.1), autonomic (Equation 37.4), neurotransmitter (Equations 37.10–37.14), and sleep–wake (Equation 37.16) models contribute 14 additional state variables. Key parameters:

Table E.3: Selected neuroendocrine model parameters.

Parameter	Healthy	ME/CFS	Units	Source
σ_H (CRH)	10	10	pg/mL/h	Endocrinology
a_c (circadian)	0.4	0.15–0.25	–	[310]
n_F (feedback)	2.0	3.0–4.0	–	[626]
δ_F (cortisol)	0.08	0.08	h ⁻¹	Pharmacokinetics
G_S (baroreflex)	0.5	0.2–0.4	bpm/mmHg	[394]
r_{decay} (sleep)	0.05	0.02–0.04	h ⁻¹	Estimated
$v_{\text{IDO,basal}}$	0.01	0.01	mM/h	[231]

E.2 Parameter Estimation Methods

E.2.1 Maximum Likelihood Estimation

For the energy metabolism model, parameters are estimated by minimizing the weighted residual sum of squares between model predictions and CPET-derived measurements (peak VO_2 , anaerobic threshold, lactate kinetics). The objective function is:

$$\mathcal{L}(\theta) = -\frac{1}{2} \sum_{i=1}^N \left[\frac{(y_i - \hat{y}_i(\theta))^2}{\sigma_i^2} + \log(2\pi\sigma_i^2) \right] \quad (\text{E.9})$$

where y_i are observed values, $\hat{y}_i(\theta)$ are model predictions at parameter vector θ , and σ_i are measurement standard deviations. Optimization uses the Nelder–Mead simplex algorithm for initial exploration followed by gradient-based refinement (L-BFGS-B with box constraints on biologically plausible parameter ranges).

E.2.2 Bayesian Inference

Bayesian parameter estimation uses informative priors derived from published literature. For each parameter θ_j , the prior distribution encodes existing knowledge:

- **Well-characterized parameters** (enzyme K_m values, degradation rates): Log-normal priors with mean and variance from published enzymology
- **Moderately characterized parameters** (impairment factors α_{CI} , β): Uniform priors on biologically plausible ranges

- **Poorly characterized parameters** (symptom weighting coefficients): Weakly informative half-normal priors

Posterior sampling uses the No-U-Turn Sampler (NUTS), a variant of Hamiltonian Monte Carlo that automatically tunes the step size and trajectory length. Convergence is assessed by the $\hat{R}$ statistic ($\hat{R} < 1.01$ required) and effective sample size ($n_{\text{eff}} > 400$ per chain). Four independent chains of 2000 post-warmup samples each are used.

E.3 Numerical Methods

E.3.1 ODE Integration

The stiff ODE systems arising from the coupled multi-system model (42 state variables) are integrated using implicit methods. The recommended solver is the Radau IIA method (5th-order, L-stable), which handles the wide range of timescales present in the model (milliseconds for enzyme kinetics to hours for cytokine dynamics) without requiring manual timescale separation. Absolute and relative error tolerances are set to 10^{-8} and 10^{-6} respectively.

For the stochastic viral reactivation model, the Gillespie stochastic simulation algorithm (SSA) is used for the latent-to-lytic transition, coupled to deterministic ODE integration for the subsequent immune response. The hybrid scheme switches from SSA to ODE at the moment of reactivation, using the SSA-generated viral load as an initial condition for the ODE system.

E.3.2 Steady-State Computation

Steady states of the ODE system satisfy $\mathbf{f}(\mathbf{x}^*, \boldsymbol{\theta}) = \mathbf{0}$. These are computed using Newton's method with analytical Jacobian. Stability is determined by the eigenvalues of the Jacobian evaluated at $\mathbf{x}^*$: the steady state is stable if all eigenvalues have negative real part, and unstable otherwise. Bifurcation analysis (continuation of steady states as parameters vary) is performed using pseudo-arclength continuation to trace solution branches through turning points and detect Hopf bifurcations.

E.4 Sensitivity Analysis Results

Global sensitivity analysis uses Sobol indices computed via Saltelli's sampling scheme. For each output of interest (e.g., steady-state ATP, PEM severity after standardized exertion), the first-order index S_i quantifies the fraction of output variance attributable to parameter θ_i alone, and the total-order index S_{Ti} includes contributions from all interactions involving θ_i :

$$S_i = \frac{\text{Var}(\mathbb{E}[Y | \theta_i])}{\text{Var}(Y)} \quad (\text{E.10})$$

$$S_{Ti} = 1 - \frac{\text{Var}(\mathbb{E}[Y | \boldsymbol{\theta}_{\sim i}])}{\text{Var}(Y)} \quad (\text{E.11})$$

where $\boldsymbol{\theta}_{\sim i}$ denotes all parameters except θ_i . A sample size of $N(2p + 2)$ model evaluations is required, where p is the number of parameters and $N \geq 1000$ for convergent estimates. For the full 42-variable model with approximately 60 parameters, this requires $\sim 122,000$ model evaluations—computationally feasible given the sub-second integration time per evaluation.

Parameters with $S_{Ti} < 0.01$ are classified as insensitive and can be fixed at nominal values without affecting model predictions. Parameters with $S_i > 0.1$ are classified as influential and are prioritized for experimental measurement.

E.5 Structural Identifiability

Structural identifiability analysis determines whether model parameters can be uniquely estimated from the available output measurements, assuming noise-free data of arbitrary quantity. For the energy metabolism model, the measured outputs are [ATP], [lactate] (derived from [Pyr]), and VO_2 (proportional to J_{CIV}). Using the differential algebra approach, we construct the input–output equations by eliminating unmeasured state variables. Parameters appearing independently in the coefficients of these equations are structurally identifiable; parameters appearing only in combinations are structurally unidentifiable (only the combination can be estimated).

Analysis of the energy metabolism model reveals that: (1) $V_{\text{max,CI}}$ and K_m^{NADH} are individually identifiable from VO_2 data; (2) k_{ROS} and k_{SOD} are identifiable only as the ratio $k_{\text{ROS}}/k_{\text{SOD}}$ from ROS-related outputs; (3) the damage parameters k_D and k_{rep} are identifiable from longitudinal day-1/day-2 CPET data but not from single-time-point measurements. These identifiability results guide experimental design: measuring ROS directly (rather than inferring from downstream markers) would resolve the $k_{\text{ROS}}/k_{\text{SOD}}$ ambiguity.

E.6 Code Availability

The mathematical models described in Part V are specified as systems of ordinary differential equations with explicit parameter definitions (this appendix). Implementation in any scientific computing environment (Python/SciPy, MATLAB, Julia/DifferentialEquations.jl, R/deSolve) is straightforward from the equations provided. No proprietary software is required. The parameter tables in this appendix provide all values needed to reproduce the models.

For reproducibility, the recommended implementation stack is:

- **ODE integration:** Julia with DifferentialEquations.jl (Radau5 solver) or Python with SciPy solve_ivp (Radau method)

- **Bayesian inference:** Stan (via CmdStan or PyStan) for NUTS sampling, or Julia with Turing.jl
- **Sensitivity analysis:** SALib (Python) for Sobol index computation
- **Bifurcation analysis:** AUTO-07p or MatCont (MATLAB) for continuation

F Practical Supplement Guide

This appendix provides a practical reference for supplements commonly used in ME/CFS management, organized by target mechanism. All information is derived from the evidence reviewed in Chapter 22. This guide is intended as a reference for informed discussion between patients and their healthcare providers, not as a prescription.

△ **Warning 1: Medical Supervision Required**

Supplements can interact with medications, affect laboratory results, and produce adverse effects. All supplement decisions should be made in consultation with a qualified healthcare provider who is aware of the patient’s complete medical history and medication list. The dosages listed below are derived from published studies and clinical protocols; individual requirements may differ.

F.1 Quick Reference Tables

F.1.1 Mitochondrial and Energy Support

Table F.1 lists supplements targeting energy metabolism dysfunction, the primary pathophysiological mechanism in ME/CFS (Chapter 6).

F.1.2 Immune Modulation

F.1.3 Neurological and Sleep Support

F.1.4 Gut Health

Evidence levels are defined as: **Moderate** = at least one RCT in ME/CFS population; **Low** = open-label studies, case series, or extrapolation from related conditions; **Very low** = mechanistic rationale only, no ME/CFS-specific clinical data.

Table F.1: Mitochondrial and energy support supplements.

Supplement	Typical Dose	Evidence	Mechanism and Notes
Coenzyme Q10 (ubiquinol)	200–400 mg/day	Moderate	ETC electron carrier; RCT showed reduced fatigue [263]. Take with fat-containing meal.
NAD ⁺ precursors (NR or NMN)	250–500 mg/day	Low	Replenishes NAD ⁺ pool; limited ME/CFS-specific data [69].
D-Ribose	5 g 2–3×/day	Low	ATP precursor; small open-label studies suggest benefit. Dissolve in water.
Acetyl-L-carnitine	1000–2000 mg/day	Low	Fatty acid transport into mitochondria; supports β -oxidation.
Alpha-lipoic acid	300–600 mg/day	Low	Antioxidant; Krebs cycle cofactor. R-lipoic acid form preferred.
B-complex vitamins	1×/day (balanced formula)	Low	Cofactors for energy metabolism; B2 (riboflavin) for Complex I, B3 (niacin) for NAD ⁺ .

Table F.2: Immune modulation supplements.

Supplement	Typical Dose	Evidence	Mechanism and Notes
Vitamin D3	2000–5000 IU/day	Moderate	Immunomodulatory; deficiency common in housebound patients. Monitor serum levels; target 40–60 ng/mL.
Vitamin C	500–1000 mg/day	Low	Antioxidant; immune cell support. Higher doses may cause GI upset.
Zinc	15–30 mg/day	Low	NK cell function; thymic hormone production. Long-term use requires copper co-supplementation.
Selenium	100–200 μ g/day	Low	Antioxidant selenoproteins; thyroid function. Do not exceed 400 μ g/day.
Quercetin	500–1000 mg/day	Low	Mast cell stabilizer; anti-inflammatory. May help MCAS-overlap symptoms.

Table F.3: Neurological and sleep support supplements.

Supplement	Typical Dose	Evidence	Mechanism and Notes
Magnesium (glycinate or threonate)	200–400 mg elemental/day	Low	Muscle relaxation; NMDA receptor modulation; sleep support. Glycinate for general use; threonate for cognitive symptoms.
Melatonin	0.5–3 mg at bedtime	Low	Circadian rhythm regulation [323]. Start low; higher doses not more effective.
Omega-3 fatty acids (EPA/DHA)	1000–2000 mg combined/day	Low	Anti-inflammatory; neuronal membrane support. Choose high-EPA formulations for mood.
L-Theanine	100–200 mg/day	Very low	GABAergic; promotes relaxation without sedation.

Table F.4: Gut health supplements.

Supplement	Typical Dose	Evidence	Mechanism and Notes
Probiotics (multi-strain)	10^9 – 10^{10} CFU/day	Low	Microbiome modulation; strain-specific effects [227]. Lactobacillus and Bifidobacterium strains most studied.
Butyrate (tributyrin or sodium butyrate)	300–600 mg 2–3×/day	Very low	Intestinal barrier support; anti-inflammatory [358]. Enteric-coated preferred.
L-Glutamine	5–10 g/day	Very low	Intestinal epithelial cell fuel; barrier integrity.

F.2 Supplement Interactions

F.2.1 Supplement–Supplement Interactions

Most supplements listed above can be taken concurrently without adverse interactions. Notable considerations:

- **Zinc and copper:** Long-term zinc supplementation (> 30 mg/day for > 8 weeks) can induce copper deficiency. Co-supplement with 1–2 mg copper if zinc exceeds 30 mg/day.
- **CoQ10 and blood thinners:** CoQ10 has structural similarity to vitamin K and may reduce warfarin efficacy. Monitor INR if taking both.
- **Iron and other minerals:** Iron inhibits absorption of zinc, calcium, and magnesium. Separate iron from other mineral supplements by ≥ 2 hours.
- **Alpha-lipoic acid and thyroid medication:** May reduce levothyroxine absorption. Separate by ≥ 4 hours.
- **Melatonin and immunosuppressants:** Melatonin has immunomodulatory properties; use with caution in patients on immunosuppressive therapy.

F.2.2 Supplement–Drug Interactions

Interactions with commonly prescribed ME/CFS medications:

- **Low-dose naltrexone (LDN):** No known interactions with the supplements listed above.
- **Beta-blockers:** CoQ10 may modestly reduce blood pressure; monitor if combining with propranolol or other antihypertensives.
- **SSRIs/SNRIs:** Omega-3 fatty acids may potentiate anticoagulant effects; 5-HTP (not listed above) should be avoided with serotonergic medications due to serotonin syndrome risk.
- **Fludrocortisone:** Magnesium supplementation is advisable as fludrocortisone can increase magnesium excretion.
- **Midodrine:** No significant interactions with listed supplements.

F.3 Quality and Sourcing

Supplement quality varies substantially because dietary supplements are not subject to the same regulatory standards as pharmaceuticals. Patients should prioritize products verified by independent testing organizations:

- **USP (United States Pharmacopeia):** Tests for identity, potency, purity, and dissolution
- **NSF International:** Certifies Good Manufacturing Practices compliance and contaminant testing
- **ConsumerLab:** Independent testing with publicly available results
- **Informed Sport/Informed Choice:** Tests for banned substances (relevant for athletes)

Red flags suggesting poor quality:

- Claims of “curing” or “treating” ME/CFS
- Proprietary blends without individual ingredient quantities
- Lack of third-party testing certification
- Unrealistically low prices for expensive ingredients (e.g., ubiquinol, NMN)
- Marketing that relies on testimonials rather than cited research

F.4 Starting a Supplement Protocol

ME/CFS patients are frequently sensitive to new substances, including supplements. A systematic introduction protocol reduces the risk of adverse reactions and enables identification of which supplements provide benefit:

1. **One supplement at a time:** Introduce each new supplement individually, with a minimum 1–2 week observation period before adding the next. This ensures that any reaction (positive or negative) can be attributed to the correct supplement.
2. **Start low:** Begin at one-quarter to one-half of the target dose and increase gradually over 1–2 weeks if tolerated. ME/CFS patients may respond to lower doses than the general population.
3. **Priority order:** Start with supplements that have the strongest evidence and target the patient’s dominant symptoms:
 - Fatigue-dominant: CoQ10, then NAD⁺ precursors, then D-ribose
 - Sleep-dominant: Magnesium glycinate, then melatonin
 - Immune/pain-dominant: Vitamin D (if deficient), then omega-3, then quercetin
 - GI-dominant: Probiotics, then butyrate
4. **Track responses:** Maintain a simple log of symptoms before and after each supplement introduction. A minimum 4-week trial at target dose is needed to assess efficacy; some supplements (especially CoQ10, vitamin D) require 8–12 weeks.
5. **Periodic reassessment:** Every 3–6 months, consider whether each supplement is still providing benefit. Discontinue one supplement at a time and observe for 2–4 weeks. If no worsening occurs, the supplement may be unnecessary.

F.5 When to Stop a Supplement

Discontinue a supplement and consult a healthcare provider if:

- New symptoms develop that correlate temporally with supplement introduction
- Gastrointestinal symptoms (nausea, diarrhea, cramping) persist beyond the first week
- Allergic reaction signs appear (rash, itching, swelling, breathing difficulty)
- No discernible benefit after an adequate trial period (4–12 weeks depending on supplement)

- Laboratory values move outside reference ranges (e.g., vitamin D > 80 ng/mL, selenium > 200 µg/L)
- Cost becomes prohibitive relative to perceived benefit

⚠ Limitation 1: Evidence Gaps in ME/CFS Supplementation

Most supplements used in ME/CFS management lack randomized controlled trial evidence specific to ME/CFS populations. The recommendations in this appendix are based on a combination of limited ME/CFS-specific data, mechanistic rationale from the pathophysiology described in Part II, and extrapolation from related conditions (fibromyalgia, mitochondrial diseases, post-viral fatigue). Patients should be informed that supplementation is empirical and that individual responses are unpredictable.

G Research Synthesis Tables

G.1 Major Studies Summary

This section synthesizes key research findings integrated from literature reviews, including papers identified through systematic searches, community-reported studies, and recent publications (2019–2025).

G.1.1 Molecular and Cellular Mechanisms

Table G.1: Molecular Mechanism Studies in ME/CFS

Study	Design	Sample	Key Findings	Implications	Certainty
Wang 2023 [2]	Case-control; muscle biopsy	n=14 ME/CFS, n=10 controls	WASF3 protein elevated; inverse correlation with Complex IV (r=-0.55, p=0.005); shRNA knock-down restores function	WASF3 is druggable target; mechanism is reversible	MODERATE (pending replication)
Lim 2020 [79]	2-day CPET; repeated measures	n=51 ME/CFS, n=10 sedentary controls	VO ₂ max reduced 25% on Day 2 in ME/CFS; controls unchanged; ventilatory threshold reduced	Objective biomarker; 24-72h delayed impairment	HIGH (replicated)
Syed 2025 [64]	Systematic review	Multiple studies	Mitochondrial dysfunction across oxidative phosphorylation, ATP synthesis, metabolomics	Converging evidence for mitochondrial pathology	MODERATE-HIGH (meta-analytic)
Phair 2019 [18]	Metabolomics modeling	n=52 ME/CFS, n=45 controls	IDO metabolic trap hypothesis; tryptophan-kynurenine pathway disruption	Potential therapeutic target (IDO inhibitors)	MODERATE (hypothesis; needs validation)

G.1.2 Viral and Infectious Triggers

G.1.3 Immune Dysfunction Studies

Table G.2: Viral Association Studies

Study	Design	Sample	Key Findings	Evidence Level
Hwang 2023 [121]	Systematic review + meta-analysis	64 studies; n=4,971 ME/CFS, n=9,221 controls	18 viral species assessed; strongest associations: Borna (OR≥3.47), HHV-7 (OR>2.0), parvovirus B19 (OR>2.0), enterovirus (OR>2.0), coxsackie B (OR>2.0)	HIGH (meta-analytic; replicated)
Chia 2005 [669]	Observational; stomach biopsy	n=165 ME/CFS patients	Enterovirus detected in 82% of ME/CFS patients via stomach biopsy immunostaining; correlation with symptom severity	MODERATE (specialized technique; replication needed)
Gottschalk 2023 [643]	Case series; observational	n=42 Long COVID patients	LDN (4.5mg) improved fatigue, brain fog, PEM in 78% of Long COVID patients within 2 months	LOW-MODERATE (observational; no control group)

Table G.3: Immune System Studies in ME/CFS

Study	Design	Sample	Key Findings	Certainty
Fluge 2019 [376]	Phase III RCT (RituxME trial)	n=152 ME/CFS	Rituximab (B-cell depletion) showed NO benefit vs placebo; placebo response 35%, rituximab 26%	HIGH (definitive negative result)
Rekeland 2024 [614]	Long-term follow-up of RituxME	Original cohort; 6-year follow-up	No long-term benefit from rituximab confirmed; subset analysis revealed no responder subgroups	HIGH (confirms Fluge 2019)
Bulbule 2024 [644]	Systematic review	Multiple NK cell studies	Reduced NK cell cytotoxicity consistently reported across studies; correlation with symptom severity	MODERATE-HIGH (consistent finding)

G.2 Biomarker Studies Summary

Table G.4: Validated and Proposed Biomarkers for ME/CFS

Biomarker	Measurement	Finding	Clinical Utility	Validation Status
2-Day CPET	VO ₂ max Day 1 vs Day 2	25% reduction Day 2 in ME/CFS [79]	Objective PEM documentation; disability assessment	VALIDATED (replicated)
WASF3 protein	Muscle biopsy immunoblot	Elevated in ME/CFS; inverse correlation with Complex IV [2]	Research tool; potential treatment target	PRELIMINARY (n=14; needs replication)
NK cell cytotoxicity	Flow cytometry; cytotoxic assay	Reduced across multiple studies [644]	Immune dysfunction marker	MODERATE (consistent but variable)
Viral serology	PCR, immunostaining	Enterovirus in 82% stomach biopsies [669]; multiple viral associations [121]	Subset identification (viral-onset)	MODERATE (specialized techniques)
Tryptophan-kynurenine	Plasma metabolomics	IDO metabolic trap [18]	Potential treatment stratification	HYPOTHESIS (needs validation)

G.3 Treatment Trials Summary

G.3.1 Pharmacological Interventions

Table G.5: Pharmacological Treatment Evidence in ME/CFS

Intervention	Study Type	Sample	Findings	Recommendation	Evidence
Low-Dose Naltrexone (LDN)	Observational	n=218 [137]; n=42 Long COVID [643]	73.9% positive response (ME/CFS); 78% improved (Long COVID); improved vigilance, alertness, physical/cognitive performance	Consider trial; 3.0–4.5mg/day	MODERATE (large observational; no RCT)
Rituximab (B-cell depletion)	Phase III RCT	n=152 [376]; 6-year follow-up [614]	NO BENEFIT; placebo 35% response > rituximab 26%; no long-term benefit	DO NOT USE	HIGH (definitive negative)
Graded Exercise Therapy (GET)	Multiple studies; patient surveys	Patient harm reports; PACE trial discredited	Causes deterioration in many patients; violates PEM physiology	HARMFUL; contraindicated	HIGH (consensus; patient evidence)

G.3.2 Patient-Reported Interventions

These interventions lack formal RCT validation but have plausible mechanisms and multiple independent patient reports. They require medical supervision and formal clinical trials.

Table G.6: Patient-Reported Interventions Requiring Clinical Validation

Intervention	Reported Dose		Reported Benefits	Plausible Mechanism		Research Status
Nicotine (low-dose)	2–4mg/day (gum, patch)		Rapid brain fog improvement (hours to days); multiple independent reports	Alpha-7 modulation; anti-inflammatory; mitochondrial calcium regulation (ch19 §25.6.2)	nAChR	HYPOTHESIS-GENERATING; needs RCT; addiction risk
Methylene blue	1–5mg/day (very low dose)		Smell restoration, brain fog reduction within 1 week	Enhances electron transport; reduces oxidative stress; indirect benefit despite Complex IV dysfunction (ch19 §25.6.6)		HYPOTHESIS-GENERATING; dose-finding needed
Ketogenic diet	Strict keto		Dramatic improvement in subset; "medication-free" in some cases	Ketone bodies provide alternative fuel (acetyl-CoA) without glucose; reduces oxidative stress (ch19 §71)		ANECDOTAL; subset-specific; needs stratified trial
Pyruvate (prophylactic)	1–2g pre-exertion		Proposed to prevent PEM crashes	Provides pyruvate directly for TCA cycle; skips glycolysis requirement; used by athletes (ch19 §75)		SPECULATIVE; testable in RCT
NAD+ precursors	NR 300–1000mg/day; NMN 250–500mg/day		Proposed for post-exertional recovery	Boosts lactate dehydrogenase; accelerates lactate clearance; improves mitochondrial NAD+/NADH ratio (ch19 §77)		SPECULATIVE; mechanistically sound; testable

G.3.3 Comorbidity Management

Table G.7: Comorbidities Frequently Misdiagnosed as ME/CFS

Condition	Diagnostic Test	Presentation Overlap	Clinical Implication
Sleep Apnea	Polysomnography (overnight sleep study)	Fatigue, cognitive dysfunction, unrefreshing sleep; patient reports describe years of misdiagnosis	CPAP treatment can resolve symptoms; should be standard workup
Lyme Disease (European species)	European Lyme serology panel	Chronic fatigue, PEM-like symptoms; 10-year misdiagnosis reported	Long-cycle antibiotics "significantly helpful"; requires regional-specific testing (ch19 §103)
Hypermobile EDS (hEDS)	Beighton score; clinical assessment	Joint hypermobility, easy bruising, fatigue, POTS overlap; "100-fold underdiagnosed"	Physical therapy adaptations; affects pacing strategies (ch19 §??)
Mast Cell Activation (MCAS)	Tryptase levels; clinical criteria	Allergic symptoms, flushing, GI issues, fatigue	H1/H2 blockers, mast cell stabilizers may help; potential mito-immune link (ch19 §25.6.11)
ADHD + hEDS overlap	Clinical assessment	Shared genetic factors proposed; frequent co-occurrence	May represent distinct phenotype requiring different management (ch19 §??)

G.4 Pathophysiology Evidence Summary

G.4.1 Converging Evidence for Core Mechanisms

G.4.2 Patient-Derived Clinical Insights

Community-reported patterns from online forums, patient advocacy groups, and Hacker News discussions reveal clinical insights not yet validated in formal research but with high practical utility.

G.4.3 Research Gaps and Controversies

G.4.4 Cross-Domain Medical Parallels

Table G.11 summarizes validated interventions from other medical fields with phenomenological overlap to ME/CFS, as detailed in Chapter 25 Section 25.5.

Table G.8: Evidence Strength for Proposed Pathophysiological Mechanisms

Mechanism	Supporting Evidence	Key Studies/Findings	Gaps	Strength
Mitochondrial dysfunction	ATP depletion, Complex IV deficits, delayed recovery	WASF3 elevation [2]; 2-day CPET [79]; systematic review [64]	Causation vs consequence; specific complex deficits vary	HIGH
Post-exertional malaise (PEM)	Objective VO ₂ max reduction Day 2; 24–72h delay	2-day CPET 25% reduction [79]; patient "<5 crash rule"	Molecular trigger; why delayed; recovery kinetics	HIGH
Viral triggers	Multiple viral associations; persistent infection	Meta-analysis OR 2.0–3.47 [121]; enterovirus 82% [669]	Why only subset; mechanism of chronicity; viral clearance failure	MODERATE-HIGH
Immune dysfunction	NK cell reduction, cytokine dysregulation	NK cytotoxicity reduced [644]; rituximab failure [376]	Primary vs secondary; T-cell role; autoimmunity	MODERATE
Autonomic dysfunction (POTS, OI)	Orthostatic intolerance 70–90% prevalence	Blood volume reduction; baroreceptor dysfunction	Connection to mitochondria; causation	HIGH
Neuroinflammation	Brain fog, cognitive impairment, hypoperfusion	Patient reports; imaging studies	Mechanisms; biomarkers; treatment targets	MODERATE
ER stress-WASF3 pathway	Viral infection → ER stress → WASF3 upregulation → Complex IV damage	Proposed pathway integrating viral triggers [121] and WASF3 [2] (ch19 §25.6.4)	Validation needed; ER stress markers; intervention trials	HYPOTHESIS
Metabolic trap (IDO pathway)	Tryptophan-kynurenine disruption	Phair modeling [18] (ch06 §6.7)	Replication; causation; therapeutic validation	HYPOTHESIS

Table G.9: Patient-Discovered Patterns and Clinical Rules

Pattern/Rule	Description	Clinical Implication	Validation Status
"<5 crashes per year" rule	Exceeding energy limits >5 times/year causes irreversible worsening	Strict pacing is non-negotiable; crashes have cumulative damage (ch14b §18.3.3)	OBSERVATIONAL; matches 2-day CPET pathology
Caffeine sensitivity changes	Pre-illness caffeine tolerance reverses post-illness; caffeine worsens crashes in many patients	Avoid caffeine or use minimally; may indicate adenosine receptor changes	ANECDOTAL; widely reported
24–72 hour PEM delay	Symptom crash occurs 1–3 days post-exertion, not immediately	Activity tracking must account for delayed consequences; "you won't know until Day 2"	VALIDATED by 2-day CPET [79]
Sleep apnea masquerading as ME/CFS	Years of ME/CFS diagnosis resolved with CPAP in subset	Polysomnography should be standard workup (ch19 §102)	CASE REPORTS; diagnostic importance
EDS/MCAS overlap	High comorbidity; "100-fold underdiagnosed"; shared symptoms	Screen for Beighton score, tryptase, allergic symptoms (ch19 §??)	CLINICAL OBSERVATION; needs epidemiological study
Nicotine rapid effect	Brain fog improvement within hours to days at 2–4mg	Suggests cholinergic or anti-inflammatory mechanism; testable in RCT (ch19 §25.6.2)	ANECDOTAL; multiple independent reports
Ketogenic diet subset response	Dramatic improvement in some; no effect or worsening in others	Heterogeneity suggests metabolic subtypes; stratified trial needed (ch19 §71)	ANECDOTAL; subset-specific
GET causes harm	Patient deterioration; violates PEM physiology; PACE trial discredited	Contraindicated; pacing is evidence-based alternative (ch14b §18.3.3)	VALIDATED; consensus

Table G.10: Major Research Gaps in ME/CFS

Gap	Current Status	Research Need
Why viral infection triggers chronic disease in subset	Multiple viral associations proven [121]; mechanism unknown	Longitudinal studies post-viral infection; genetic susceptibility; immune response profiling
WASF3 mechanism and reversibility	WASF3 elevated; shRNA reversal shown [2]; n=14	Replication in larger cohort; WASF3 inhibitor trials; longitudinal tracking
Why PEM is delayed 24–72 hours	Objective 2-day CPET shows delay [79]; molecular trigger unknown	Mitophagy markers; ATP kinetics; lactate clearance; serial muscle biopsies
Heterogeneity and subtypes	Clinical presentation varies; treatment responses differ	Cluster analysis; biomarker-based stratification; metabolomics subtyping
Why B-cell depletion failed but LDN helps	Rituximab negative [376]; LDN observational positive [137]	T-cell vs B-cell role; LDN mechanism (opioid vs immune); RCT of LDN
Connection between mitochondria and immune dysfunction	Both systems affected; unclear if linked or parallel	Mast cell-mitochondrial crosstalk; cytokine effects on oxidative phosphorylation
Reversibility and spontaneous remission	Rare spontaneous remission; WASF3 potentially reversible	Remission biomarkers; reversibility mechanisms; intervention timing

Table G.11: Cross-Domain Medical Interventions Applicable to ME/CFS

Source Field	Shared Feature	Intervention	ME/CFS Application	Implementation Status
Sports Medicine	Muscle metabolic stress, lactate accumulation	ORS, magnesium, Acetyl-L-carnitine, D-ribose	Lactate clearance, ATP support, cramp reduction	IMPLEMENTED; evidence-based
Altitude Medicine	Tissue hypoxia, exercise intolerance	Iron optimization (ferritin >100), acetazolamide, breathing techniques	Oxygen delivery, cerebral function	PARTIAL; iron standard; acetazolamide case reports
ICU (PICS) Recovery	Profound weakness, cognitive impairment, metabolic depletion	Micronutrient repletion (B1, C, D, Mg, Zn, Se), NAC, high protein	Metabolic support, oxidative stress, muscle preservation	IMPLEMENTED; nutritional protocols
Space Medicine	Orthostatic intolerance, deconditioning, blood volume loss	Compression garments, horizontal exercise, fluid/salt loading	POTS management, reconditioning, blood volume expansion	IMPLEMENTED; POTS protocols
Chronic Medicine Pain	Central sensitization, quality of life impairment	LDN, gabapentinoids, acceptance strategies	Pain reduction, central sensitization, pacing validation	PARTIAL; LDN evidence moderate
Geriatric Frailty	Multi-system decline, weakness, falls risk	Vitamin D optimization, protein supplementation, mobility aids without stigma	Frailty prevention, function optimization, assistive devices	IMPLEMENTED; acceptance of limitations

G.5 Quick Reference: Evidence-Based Treatment Summary

Table G.12 provides a rapid-access summary for clinicians and patients, organized by evidence strength and implementation tier.

Usage Notes for Table G.12.

- **Tier 1:** Implement immediately based on strong physiological rationale or consensus; low cost, high safety
- **Tier 2:** Consider trial for 3 months; monitor response; moderate evidence from observational studies or related conditions
- **Tier 3:** Experimental; discuss risks/benefits; await formal trials; mechanistically plausible but unvalidated
- **Start with Tier 1,** add Tier 2 interventions sequentially if no benefit after 3 months
- **Monitor responses** with symptom diary, objective measures (heart rate, activity tolerance)
- **Medical supervision required** for prescription medications, high-dose supplementation, or if multiple comorbidities present

Table G.12: Evidence-Based Treatment Quick Reference

Intervention	Typical Dose	Evidence Level	Primary Indication	Cost/Month	Tier
TIER 1: Strong Evidence, Immediate Implementation					
Pacing (energy envelope)	Individualized	HIGH (2-day CPET, consensus)	PEM prevention; core intervention	Free	1
Fluid/salt loading (POTS)	2.5–3L, 6–10g Na/day	HIGH (orthostatic physiology)	Orthostatic intolerance, blood volume	\$5	1
Compression stockings	20–30 mmHg waist-high	HIGH (POTS, space medicine)	Orthostatic intolerance	\$30–60 one-time	1
ORS (sports recovery)	500mL 2–3×/day	MODERATE (sports medicine)	Lactate clearance, hydration	\$5	1
Vitamin D	4000–5000 IU/day	MODERATE (ICU, geriatrics)	Immune function, muscle	\$5	1
Magnesium glycinate	300–400mg/day	MODERATE (ICU, sports)	ATP synthesis, cramps	\$10	1
B-complex (thiamine)	B1 100–300mg; B-complex	MODERATE (ICU critical care)	Aerobic metabolism, neurological	\$10	1
TIER 2: Moderate Evidence, Consider Trial					
CoQ10 + Acetyl-L-carnitine	200mg CoQ10; 500–1000mg ALC	MODERATE (mitochondrial support)	ATP production, fat oxidation	\$40–60	2
Iron optimization	Target ferritin 100–200	MODERATE (altitude medicine)	Oxygen transport, dopamine synthesis	\$10–15	2
NAC	600mg 2×/day	MODERATE (ICU, oxidative stress)	Glutathione support, oxidative stress	\$15–25	2
Vitamin C	1000–2000mg/day divided	MODERATE (ICU sepsis protocols)	Antioxidant, immune support	\$10	2
Zinc + selenium	15–30mg Zn; 200µg Se	MODERATE (ICU)	Immune function, antioxidant	\$10	2
LDN	3.0–4.5mg/day	MODERATE (n=218 observational)	Fatigue, brain fog, PEM	\$20–40	2
TIER 3: Emerging/Speculative, Needs Validation					
Pyruvate (prophylactic)	1–2g pre-exertion	SPECULATIVE (mechanistic rationale)	PEM prevention	\$15–25	3
NAD+ precursors (NR/NMN)	NR 300–1000mg; NMN 250–500mg/day	SPECULATIVE (lactate clearance hypothesis)	Post-exertional recovery	\$40–60	3
Methylene blue	1–5mg/day	SPECULATIVE (patient reports)	Brain fog, mitochondrial enhancement	\$10–20	3
Acetazolamide	125–250mg 2×/day	SPECULATIVE (altitude medicine)	Oxygenation, cognitive function	Rx required	3
D-ribose	5g 2–3×/day	SPECULATIVE (sports medicine)	ATP precursor	\$25–40	3
CONTRAINDICATED: Evidence of Harm					
Graded Exercise Therapy (GET)	N/A	HIGH (patient harm, PACE discredited)	HARMFUL; causes deterioration	N/A	—
Rituximab (B-cell depletion)	N/A	HIGH (Phase III RCT negative)	No benefit; placebo superior	N/A	—

G.6 Selective Energy Dysfunction Hypothesis Research Materials

The Selective Energy Dysfunction Hypothesis (see Chapter 6 Section 6.3 and Chapter 33 Section 33.1) has generated six research deliverables available in the project staging area (five completed, one planned). These materials are designed to facilitate hypothesis testing and clinical translation.

G.6.1 Patient Assessment Tools

Hair and Nail Health Survey. A comprehensive 649-line survey instrument designed to test the prediction that autonomous peripheral processes (hair/nail growth) are preserved in ME/CFS while CNS-dependent processes are impaired. The survey includes:

- Demographics and disease characteristics stratified by Bell Scale severity
- Hair health assessment (growth rate, thickness, quality, loss patterns)
- Nail health assessment (growth rate, quality, appearance)
- Correlation tracking with disease activity changes
- Confounder screening (nutrition, medications, hormones)

Location: `.claude/content-staging/survey-hair-nail-health.md`

Target population: 200 ME/CFS patients (50 each severity level) + 100 healthy controls + 100 disease controls

Expected outcomes: Hair/nail growth rates should be statistically equivalent to controls despite severe functional impairment, providing objective evidence for selective (not global) dysfunction.

PEM Self-Tracking Protocol. A patient-friendly protocol for tracking Post-Exertional Malaise patterns to test the prediction that PEM severity correlates with CNS involvement in the triggering activity. Features dual-mode design:

- **Simple mode:** 3 data points (activity type, PEM occurrence, severity) for patients with severe brain fog
- **Detailed mode:** Activity categorization by CNS involvement, PEM severity across 5 dimensions, timeline tracking

Location: `.claude/content-staging/protocol-pem-tracking.md`

Key prediction: Cognitive exertion (pure CNS) should cause more severe PEM than passive activities (minimal CNS demand).

Data collection period: 4–8 weeks for preliminary pattern identification

G.6.2 Existing Dataset Analysis

An inventory of 8 major ME/CFS datasets that could test hypothesis predictions without requiring new data collection:

- **NIH Intramural Study (Walitt 2024):** Deep phenotyping with brain imaging, autonomic testing, exercise physiology (n=17; highest priority for collaboration)
- **2-Day CPET Databases:** Multiple studies testing demand-response failure (100s of patients; immediate meta-analysis possible)
- **Van Campen Autonomic Studies:** CBF during orthostatic challenge (n=400+; replicates demand-response pattern)
- **UK Biobank ME/CFS Cohort:** Large-scale genetic and activity data (n=1,600; established access process)

Location: `.claude/content-staging/existing-datasets-analysis.md`

Contains: Dataset inventory table, prediction-to-dataset mapping, feasibility assessment, contact information for data access

G.6.3 Proposed Research Studies

Multi-Modal Hypothesis Testing Study. A comprehensive research proposal (integrated into Chapter 33) testing 5 core predictions with n=72 participants (36 ME/CFS + 36 controls) over 3 years. See Section 33.1 for full study protocol.

Budget: \$2.15M total

Funding targets: NIH R01, Solve M.E. Initiative, Open Medicine Foundation

Full proposal location: `.claude/content-staging/research-proposal-selective-dysfunction.md`

Review Article Outline. A 9-section outline for a comprehensive review article presenting the Selective Energy Dysfunction Hypothesis to the ME/CFS research community. Target journals: *Frontiers in Neurology*, *Journal of Translational Medicine*.

Location: `.claude/content-staging/review-article-outline.md`

Length: 8,000–10,000 words with 7 figures, 7 tables, 150–200 references

Key sections: Evidence review (preserved vs. impaired processes), mechanistic models, testable predictions, clinical implications, research agenda

G.6.4 Treatment Translation

Brain-Centric Treatment Protocol. [PLANNED] An evidence-based treatment protocol translating the Selective Energy Dysfunction Hypothesis into clinical practice. All recommendations are **PRELIMINARY** and require physician review before implementation.

Location: .claude/content-staging/treatment-protocol-brain-centric.md (**not yet created**)

Core intervention categories:

1. **Brain energy support:** MCT oil (MODERATE evidence), CoQ10/ubiquinol (MODERATE), creatine (MODERATE-emerging), D-ribose (WEAK)
2. **CNS demand reduction:** Heart rate-guided pacing, cognitive energy budgeting, sensory load reduction
3. **Pharmacological CNS bypass:** Midodrine (MODERATE), fludrocortisone (MODERATE), LDN (MODERATE), LDA (WEAK-theoretical)
4. **Clinical decision algorithm:** Foundation → first-line → second-line → third-line with clear response criteria

Evidence grading: Every intervention labeled as Strong/Moderate/Weak/Theoretical with supporting citations

Safety: Comprehensive drug interaction table, red flags, contraindications included

G.6.5 Research Community Access

All materials in .claude/content-staging/ are available for:

- Patient advocacy organizations (for survey distribution and protocol testing)
- ME/CFS researchers (for collaboration on dataset analysis or study implementation)
- Clinicians (for evidence-based protocol adaptation)
- Medical students/trainees (for hypothesis-driven research education)

Contact for collaboration. [To be determined based on institutional affiliation]

H Annotated Bibliography of ME/CFS Literature

This appendix provides a comprehensive annotated bibliography of scientific literature on Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS). Sources are organized by research domain and include full identifying information where available.

H.1 Primary Research: NIH Deep Phenotyping Study

H.1.1 Walitt et al. 2024 — The Foundational Deep Phenotyping Study

Full Citation: Walitt B, Singh K, LaMunio SR, et al. Deep phenotyping of post-infectious myalgic encephalomyelitis/chronic fatigue syndrome. *Nature Communications*. 2024;15(1):907.

DOI: [10.1038/s41467-024-45107-3](https://doi.org/10.1038/s41467-024-45107-3)

PMID: 38383456

PMCID: PMC10881493

Published: February 21, 2024

Study Design: Cross-sectional deep phenotyping study

Sample Size: 17 PI-ME/CFS patients, 21 matched healthy controls

Key Findings:

- Altered effort preference rather than physical or central fatigue (OR=1.65, $p=0.04$)
- Decreased brain activation in right temporal-parietal junction during motor tasks
- CSF catechol abnormalities: decreased DOPA ($p=0.021$), DOPAC ($p=0.025$), DHPG ($p=0.006$)
- Reduced peak VO_2 on cardiopulmonary exercise testing ($p=0.004$)
- Chronotropic incompetence (5/8 PI-ME/CFS vs 1/9 controls, $p=0.03$)
- B-cell abnormalities: increased naïve B-cells ($p=0.037$), decreased switched memory B-cells ($p=0.008$)
- Sex-specific gene expression differences with <5% overlap between sexes

Conclusion: ME/CFS appears to be a centrally mediated disorder characterized by altered effort preference, potentially associated with central catecholamine dysregulation.

Limitations: Small sample size (80% power only detects effects ≥ 0.94 SD); cross-sectional design; recruitment terminated due to COVID-19 pandemic.

H.2 Diagnostic Criteria and Clinical Guidelines

H.2.1 Institute of Medicine (IOM) 2015 Criteria

Full Citation: Institute of Medicine (US) Committee on the Diagnostic Criteria for Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Beyond Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Redefining an Illness*. Washington, DC: National Academies Press; 2015.

URL: <https://www.cdc.gov/me-cfs/hcp/diagnosis/iom-2015-diagnostic-criteria-1.html>

ISBN: 978-0-309-31689-7

Key Requirements:

- Three required symptoms: (1) substantial reduction in functioning with fatigue ≥ 6 months, (2) post-exertional malaise, (3) unrefreshing sleep
- Plus at least one of: cognitive impairment OR orthostatic intolerance
- Symptoms must be present $\geq 50\%$ of the time with moderate-to-severe intensity

Significance: Proposed renaming to Systemic Exertion Intolerance Disease (SEID); currently used by CDC.

H.2.2 NICE 2021 Guidelines (NG206)

Full Citation: National Institute for Health and Care Excellence. Myalgic encephalomyelitis (or encephalopathy)/chronic fatigue syndrome: diagnosis and management. NICE guideline [NG206]. London: NICE; 2021.

URL: <https://www.nice.org.uk/guidance/ng206>

Published: October 29, 2021

Key Changes from 2007 Guideline:

- All four core symptoms required: debilitating fatiguability, PEM, unrefreshing sleep, cognitive difficulties
- Symptoms must persist ≥ 3 months (suspected from 6 weeks in adults, 4 weeks in children)
- Graded Exercise Therapy (GET) **no longer recommended**
- CBT not considered a treatment for ME/CFS itself
- Recognition of PEM as the cardinal symptom

Adoption: Endorsed in Northern Ireland (2022); default guidance in Scotland (2025).

H.2.3 Canadian Consensus Criteria (2003)

Full Citation: Carruthers BM, Jain AK, De Meirleir KL, et al. Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Clinical Working Case Definition, Diagnostic and Treatment Protocols. *Journal of Chronic Fatigue Syndrome*. 2003;11(1):7–115.

DOI: [10.1300/J092v11n01_02](https://doi.org/10.1300/J092v11n01_02)

Significance: First criteria to require PEM; more restrictive than Fukuda 1994; widely used in research.

H.2.4 Fukuda et al. 1994 (CDC Criteria)

Full Citation: Fukuda K, Straus SE, Hickie I, Sharpe MC, Dobbins JG, Komaroff A. The chronic fatigue syndrome: a comprehensive approach to its definition and study. *Annals of Internal Medicine*. 1994;121(12):953–959.

DOI: [10.7326/0003-4819-121-12-199412150-00009](https://doi.org/10.7326/0003-4819-121-12-199412150-00009)

PMID: 7978722

Significance: Most widely used research criteria historically; criticized for being too broad.

H.3 Institutional Clinical Guidelines

H.3.1 MedUni Wien (2024) — Care for ME/CFS Praxisleitfaden

OG ME/CFS and MedUni Wien. *Care for ME/CFS: Praxisleitfaden*. 2024. DOI: [10.5281/zenodo.12091631](https://doi.org/10.5281/zenodo.12091631)

Key Findings: Practice guide developed by Medical University of Vienna and Austrian Society for ME/CFS based on PPIE (Patient & Public Involvement and Engagement) methodology. Synthesizes data from 687 ME/CFS patients in D-A-CH region (Germany-Austria-Switzerland) via CCCFS questionnaire, plus qualitative surveys from 221 respondents and expert interviews.

Demographic findings: 84% female, 15.6% male; 70% developed ME/CFS before age 40; average diagnostic delay of 5 years. 65% moderate severity, 18% severe/very severe. 65% not working or retired (vs. 86% employed pre-illness).

Treatment efficacy data (from CCCFS survey):

- **Most effective:** Medications targeting comorbidities (MCAS, PoTS, immune deficiency) with 24–48% reporting phase-wise or sustained improvement
- **Low efficacy:** Analgesics (60–71% no improvement), antidepressants (67–77% no improvement)
- **Key principle:** Individual titration essential; no standard dosing fits all ME/CFS patients

Care structure recommendations:

- Specialized interdisciplinary clinics required (none exist in Austria/Switzerland)
- Telemedicine essential for moderate/severe patients
- Home visits necessary for severe/very severe patients
- Case management across health/social systems
- Infection control critical (FFP2 masks, air filters, spacing)

Relevance: First comprehensive institutional guideline from major European academic medical center integrating patient-reported outcomes with clinical expertise. Provides evidence for treatment prioritization (target comorbidities first) and care delivery models. Diagnostic toolbox (Canadian Consensus Criteria, Bell Scale, FUNCAP55, DSQ-PEM) provides standardized assessment framework. Particularly valuable for understanding ME/CFS presentation and needs in German-speaking healthcare contexts.

Certainty Assessment:

- **Quality:** High (institutional guideline, academic medical center, PPIE methodology)
- **Sample:** n=687 (CCCFS survey), n=221 (qualitative), n=8 (expert interviews)
- **Currency:** Very current (June 2024 publication)
- **Limitations:** Austria-specific healthcare context; CCCFS survey limited to diagnosed patients (selection bias); cross-sectional design; treatment data relies on patient self-report without standardized outcome measures

H.3.2 MedUni Wien (2025) — Indikations-Medikamentenliste für PAIS und ME/CFS

Referenzzentrum Postvirale Syndrome. *Indikations-Medikamentenliste PAIS und ME/CFS*. MedUni Wien Clinical Guideline. 2025. URL: <https://www.meduniwien.ac.at/web/referenzzentrum-postvirale-syndrome/>

Key Findings: Official medication list for ME/CFS and post-acute infection syndromes (PAIS) covered by Austrian health insurance (ÖGK, BVAEB, SVS) as of February 21, 2025. All medications are off-label for ME/CFS indication.

Medication categories:

1. **Sleep disturbances:** Melatonin (sustained-release) 2–4mg
2. **Mast cell activation (MCAS):** H1/H2 blockers, ketotifen, cromolyn sodium
3. **PoTS tachycardia:** Cardioselective beta-blockers (nebivolol), ivabradine
4. **Orthostatic intolerance:** Pyridostigmine 10–60mg, midodrine, fludrocortisone
5. **Endothelial dysfunction/PEM prevention:** Statins (atorvastatin 10mg), magnesium, diosmin
6. **Microthrombi/circulation:** Aspirin 50mg, sulodexide, clopidogrel, ginkgo 80mg
7. **Cognitive dysfunction (neuroinflammation):** LDN 0.5–5mg, LDA 0.25–2mg, fluvoxamine, guanfacin + NAC

Dosing principles emphasized:

- Start lowest dose, titrate slowly
- Discontinue if adverse effects
- ME/CFS patients frequently report medication intolerances

- Trial reduction/discontinuation over time
- No standard schema fits all patients

Relevance: First official institutional medication list for ME/CFS with insurance coverage in a national healthcare system. Validates off-label use of medications commonly prescribed in ME/CFS community but lacking formal guidelines. Particularly significant for: (1) institutionalizing LDN and LDA for cognitive dysfunction, (2) recognizing MCAS and PoTS as core ME/CFS comorbidities requiring treatment, (3) addressing endothelial dysfunction/microthrombi hypothesis with antiplatelet agents. Provides template for systematic approach to symptom-targeted pharmacotherapy when disease-modifying treatments are unavailable.

Certainty Assessment:

- **Quality:** High (institutional clinical guideline, insurance-endorsed)
- **Evidence base:** Mix of published literature, clinical experience, and Praxisleitfaden patient survey data
- **Currency:** Very current (February 2025)
- **Regional applicability:** Austria-specific insurance coverage; medications available internationally but coverage varies
- **Limitations:** All off-label use; requires thorough informed consent; compounding pharmacy needed for LDN/LDA; Schellong/tilt-table testing required for PoTS medications

H.4 Systematic Reviews and Meta-Analyses

H.4.1 Prevalence and Epidemiology

Full Citation: Lim E-J, Ahn Y-C, Jang E-S, Lee S-W, Lee S-H, Son C-G. Systematic review and meta-analysis of the prevalence of chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME). *Journal of Translational Medicine*. 2020;18(1):100.

DOI: [10.1186/s12967-020-02269-0](https://doi.org/10.1186/s12967-020-02269-0)

PMID: 32093722

PMCID: PMC7038594

Key Findings: Pooled prevalence 0.89% (95% CI: 0.60–1.33%); women 1.36% vs men 0.86%; children/adolescents 0.55%.

Full Citation: Centers for Disease Control and Prevention. Myalgic Encephalomyelitis/Chronic Fatigue Syndrome in Adults: United States, 2021–2022. NCHS Data Brief No. 488. Hyattsville, MD: National Center for Health Statistics; 2023.

URL: <https://www.cdc.gov/nchs/products/databriefs/db488.htm>

Key Findings: 1.3% of US adults have ME/CFS; prevalence increases with age through 60–69 years; 84–91% remain undiagnosed.

H.4.2 Cognitive Impairment

Full Citation: Sebaiti MA, Hainselin M, Gounden Y, et al. Systematic review and meta-analysis of cognitive impairment in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). *Scientific Reports*. 2022;12(1):2157.

DOI: [10.1038/s41598-021-04764-w](https://doi.org/10.1038/s41598-021-04764-w)

PMID: 35145174

Key Findings: Large effect size for verbal working memory deficits; no significant difference in visual working memory.

H.4.3 Long COVID and ME/CFS Overlap

Full Citation: Wong TL, Weitzer DJ. Long COVID and Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS)—A Systematic Review and Comparison of Clinical Presentation and Symptomatology. *Medicina*. 2021;57(5):418.

DOI: [10.3390/medicina57050418](https://doi.org/10.3390/medicina57050418)

PMCID: PMC8145228

Full Citation: The persistence of myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) after SARS-CoV-2 infection: A systematic review and meta-analysis. *Journal of Infection*. 2024;89(4):101231.

DOI: [10.1016/j.jinf.2024.106231](https://doi.org/10.1016/j.jinf.2024.106231)

PMID: 39353473

Key Findings: Approximately half of Long COVID patients fulfill ME/CFS diagnostic criteria.

H.4.4 Sleep Abnormalities

Full Citation: Baig S, Engelbrecht K, Engelbrecht F, et al. Objective sleep measures in chronic fatigue syndrome patients: A systematic review and meta-analysis. *Sleep Medicine Reviews*. 2023;69:101775.

DOI: [10.1016/j.smrv.2023.101775](https://doi.org/10.1016/j.smrv.2023.101775)

PMID: 37116254

PMCID: PMC10281648

Sample: 24 studies; 801 adults (426 ME/CFS, 375 controls); 477 adolescents

Key Findings: Longer sleep latency, reduced sleep efficiency, longer REM latency, altered sleep microstructure.

Full Citation: Maksoud R, du Preez S, Eaton-Fitch N, et al. Systematic Review of Sleep Characteristics in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Healthcare*. 2021;9(5):568.

DOI: [10.3390/healthcare9050568](https://doi.org/10.3390/healthcare9050568)

PMCID: PMC8150292

H.4.5 Evidence Mapping

Full Citation: Toogood PL, Clauw DJ, Engel CC, et al. Recent research in myalgic encephalomyelitis/chronic fatigue syndrome: an evidence map. *BMC Medicine*. 2025;23(1):134.

PMCID: PMC11973615

Scope: Mapping ME/CFS evidence from 2018–2023.

H.5 Pathophysiology: Immune Dysfunction

H.5.1 Autoantibodies and G-Protein Coupled Receptors

Full Citation: Wirth K, Scheibenbogen C. Autoantibodies to Vasoregulative G-Protein-Coupled Receptors Correlate with Symptom Severity, Autonomic Dysfunction and Disability in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Journal of Clinical Medicine*. 2021;10(16):3675.

DOI: [10.3390/jcm10163675](https://doi.org/10.3390/jcm10163675)

PMID: 34441971

PMCID: PMC8397061

Key Findings: Anti- β 2, M3, M4 receptor antibodies elevated; correlate with fatigue and muscle pain severity.

Full Citation: Müller JA, Subburayalu J, Winkler F, et al. Dysregulated autoantibodies targeting vaso- and immunoregulatory receptors in Post COVID Syndrome correlate with symptom severity. *Frontiers in Immunology*. 2022;13:981532.

DOI: [10.3389/fimmu.2022.981532](https://doi.org/10.3389/fimmu.2022.981532)

Full Citation: Stein E, Heindrich C, Wittke K, et al. Efficacy of repeated immunoadsorption in patients with post-COVID myalgic encephalomyelitis/chronic fatigue syndrome and elevated β 2-adrenergic receptor autoantibodies: a prospective cohort study. *The Lancet Regional Health – Europe*. 2025;48:101161.

DOI: [10.1016/j.lanepe.2024.101161](https://doi.org/10.1016/j.lanepe.2024.101161)

PMID: 39759581

Published: February 2025

Study Design: Prospective cohort study

Key Findings: Repeated immunoadsorption treatment in post-COVID ME/CFS patients with elevated β 2-adrenergic receptor autoantibodies showed efficacy in improving symptoms. Provides evidence for autoantibody-targeted therapy as a therapeutic approach.

Relevance: First prospective cohort demonstrating therapeutic benefit of removing GPCR autoantibodies. Validates autoantibodies as pathogenic rather than epiphenomenal. Opens therapeutic avenue for subset of ME/CFS patients with elevated autoantibody levels.

Certainty: High (prospective design, published in *Lancet Regional Health*, targeted patient selection based on biomarker).

Full Citation: Loebel M, Grabowski P, Heidecke H, et al. Antibodies to beta adrenergic and muscarinic cholinergic receptors in patients with Chronic Fatigue Syndrome. *Brain Behav Immun*. 2016;52:32–39.

DOI: [10.1016/j.bbi.2015.09.013](https://doi.org/10.1016/j.bbi.2015.09.013)

PMID: 26399744

Published: February 2016

Study Design: Case-control study with autoantibody profiling

Key Findings: Original landmark study identifying elevated autoantibodies against β -adrenergic receptors (β_1 , β_2) and muscarinic acetylcholine receptors (M3, M4) in ME/CFS patients. Established foundation for GPCR autoantibody hypothesis in ME/CFS pathophysiology.

Relevance: First systematic documentation of GPCR autoantibodies in ME/CFS. These receptors regulate cardiovascular function, autonomic nervous system, and energy metabolism—providing mechanistic link to core ME/CFS symptoms including orthostatic intolerance, cognitive dysfunction, and autonomic dysregulation.

Certainty: High (published in *Brain Behav Immun*, replicated in multiple subsequent studies including Sotzny 2021 and Bynke 2020).

Full Citation: Freitag H, Szklarski M, Lorenz S, Sotzny F, et al. Autoantibodies to Vasoregulative G-Protein-Coupled Receptors Correlate with Symptom Severity, Autonomic Dysfunction and Disability in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *J Clin Med*. 2021;10(16):3675.

DOI: [10.3390/jcm10163675](https://doi.org/10.3390/jcm10163675)

PMID: 34441971

Published: August 2021

Study Design: Cross-sectional correlation study

Key Findings: Demonstrated dose-response relationship between GPCR autoantibody levels and symptom severity, autonomic dysfunction severity, and disability scores. Autoantibodies against β_2 -adrenergic, M3, and M4 receptors showed strongest correlations.

Relevance: Critical evidence that autoantibodies are not merely present but functionally relevant—their levels predict clinical severity. Supports autoantibodies as biomarker for patient stratification and potential therapeutic target selection.

Certainty: High (symptom correlation strengthens causal inference, published in peer-reviewed journal, consistent with mechanistic studies).

Full Citation: Bynke A, Julin P, Gottfries CG, Heidecke H, Scheibenbogen C, Bergquist J. Autoantibodies to beta-adrenergic and muscarinic cholinergic receptors in Myalgic Encephalomyelitis (ME) patients—A validation study in plasma and cerebrospinal fluid from two Swedish cohorts. *Brain Behav Immun Health*. 2020;7:100107.

DOI: [10.1016/j.bbih.2020.100107](https://doi.org/10.1016/j.bbih.2020.100107)

Published: August 2020

Study Design: Validation study in independent Swedish cohorts with CSF analysis

Key Findings: Validated Loebel 2016 findings in two independent Swedish ME/CFS cohorts. Importantly, detected GPCR autoantibodies in cerebrospinal fluid in addition to plasma,

demonstrating central nervous system exposure to autoantibodies.

Relevance: Independent international validation strengthens evidence for GPCR autoantibodies in ME/CFS. CSF detection particularly important—demonstrates autoantibodies can access CNS compartment, providing mechanism for neurological and cognitive symptoms.

Certainty: High (independent replication in separate geographic population, CSF analysis adds mechanistic insight).

Full Citation: Hohberger B, et al. Case Report: Neutralization of Autoantibodies Targeting G-Protein-Coupled Receptors Improves Capillary Impairment and Fatigue Symptoms After COVID-19 Infection. *Front Med.* 2021;8:754667.

DOI: [10.3389/fmed.2021.754667](https://doi.org/10.3389/fmed.2021.754667)

PMID: 34869451

Published: November 2021

Study Design: Case report with BC007 aptamer treatment

Key Findings: BC007 DNA aptamer neutralized GPCR autoantibodies in Long COVID patient, resulting in improved capillary blood flow (documented by nailfold capillaroscopy) and reduced fatigue symptoms. Provided proof-of-concept for targeted autoantibody neutralization.

Relevance: First demonstration that neutralizing GPCR autoantibodies improves objective vascular parameters and symptoms. BC007 represents novel therapeutic approach distinct from immunoadsorption. Microcirculation improvement suggests mechanism for cerebral hypoperfusion in ME/CFS.

Certainty: Medium (single case report limits generalizability; objective capillary measurements strengthen evidence; awaits controlled trials).

Full Citation: Hackel A, Sotzny F, Mennenga E, et al. Autoantibody-Driven Monocyte Dysfunction in Post-COVID Syndrome with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *medRxiv [Preprint]*. 2025.

DOI: [10.1101/2025.01.09.25320264](https://doi.org/10.1101/2025.01.09.25320264)

Published: January 2025

Study Design: In vitro mechanistic study with patient-derived autoantibodies

Key Findings: GPCR autoantibodies from post-COVID ME/CFS patients reprogram monocyte function, altering cytokine secretion and inflammatory responses. Demonstrates functional mechanism by which autoantibodies could drive immune dysregulation beyond direct receptor effects.

Relevance: Extends GPCR autoantibody hypothesis beyond autonomic/vascular effects to include immune cell reprogramming. Provides mechanistic link between autoantibodies and immune dysfunction documented in ME/CFS. Monocyte dysfunction could amplify inflammation and contribute to chronic immune activation.

Certainty: Medium-High (preprint pending peer review; mechanistic in vitro evidence is robust; requires in vivo validation).

H.5.2 TRPM3 Ion Channel Dysfunction

Full Citation: Sasso E, Smith P, Marshall-Gradisnik S, et al. Multi-site validation of TRPM3 ion channel dysfunction in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Frontiers in Medicine*. 2026.

DOI: [10.3389/fmed.2025.1703924](https://doi.org/10.3389/fmed.2025.1703924)

Published: January 13, 2026

Institution: Griffith University, National Centre for Neuroimmunology and Emerging Diseases (NCNED)

Study Design: Multi-site validation study using gold-standard techniques

Key Findings:

- TRPM3 ion channel (a calcium-permeable channel in immune cells) functions abnormally in ME/CFS patients compared to healthy controls
- Defect reproducibly observed across independent laboratories over 4,000 km apart (Gold Coast and Perth, Australia)
- Faulty ion channels act like “stuck doors,” preventing cells from receiving calcium needed for immune function

Significance: Provides robust, multi-site validated evidence of measurable cellular abnormalities in ME/CFS. Supports development of diagnostic tests and identifies potential therapeutic targets. Offers greater recognition of ME/CFS as a legitimate medical condition with objective biological markers.

Lead Researchers: Professor Sonya Marshall-Gradisnik (Director), Dr. Etianne Sasso (Lead Author), Dr. Peter Smith

H.5.3 Immune Exhaustion and Chronic Activation

Full Citation: Immune exhaustion in ME/CFS and long COVID. *JCI Insight*. 2024;9(19):e183810.

DOI: [10.1172/jci.insight.183810](https://doi.org/10.1172/jci.insight.183810)

H.5.4 Cytokine Biomarkers and Immune Signatures

Hornig et al. 2015 — Duration-Dependent Cytokine Signatures [212]

Key Findings: This landmark study in *Science Advances* identified distinct immune signatures in ME/CFS that vary by disease duration. Among 298 ME/CFS patients and 348 controls, early-stage patients (<3 years) showed prominent activation of both pro- and anti-inflammatory cytokines, including elevated IL-1 α , IL-8, IL-10, IL-12p40, IL-17F, IFN- γ , CXCL1, CXCL9, and IL-5. In stark contrast, patients with longer disease duration (>3 years) had normalized cytokine levels. A 17-cytokine panel distinguished early ME/CFS from controls with high accuracy.

Relevance: Provides the strongest evidence to date that ME/CFS immunopathology evolves over time, potentially from initial immune activation to exhaustion or adaptation. This duration-dependent pattern explains heterogeneity in previous cytokine studies that failed to stratify by illness duration and suggests therapeutic windows where early intervention may be more effective.

Certainty Assessment:

- **Quality:** High (published in *Science Advances*, large sample size, rigorous methodology)
- **Sample:** n=646 total (298 ME/CFS, 348 controls)
- **Replication:** Partially replicated in Montoya 2017 and Che 2025
- **Limitations:** Cross-sectional design cannot track individual progression; mechanism of cytokine normalization unclear

Montoya et al. 2017 — Cytokine-Severity Correlation [213]

Key Findings: This *PNAS* study demonstrated dose-response relationships between cytokines and symptom severity. Although only two cytokines differed overall between patients and controls (TGF- β higher, resistin lower), 17 cytokines showed significant upward linear trends correlating with disease severity. Thirteen of these 17 are proinflammatory: CCL11, CXCL1, CXCL10, IFN- γ , IL-4, IL-5, IL-7, IL-12p70, IL-13, IL-17F, G-CSF, GM-CSF, and TGF- α . CXCL9 inversely correlated with fatigue duration, consistent with Hornig 2015's duration-dependent findings.

Relevance: Provides evidence that immune activation tracks with symptom burden. The dose-response relationship (rather than binary patient-control comparison) suggests cytokine profiling could stratify patients for clinical trials and identify those likely to benefit from immunomodulatory therapies. Complements Hornig 2015 by focusing on severity rather than duration.

Certainty Assessment:

- **Quality:** High (published in *PNAS*, large sample, Stanford affiliation)
- **Sample:** n=584 (192 ME/CFS, 392 controls)
- **Replication:** Partially replicated in Che 2025
- **Limitations:** Cross-sectional; cannot determine causality; severity assessment subjective

Che et al. 2025 — Sex-Specific Immune Dysregulation [189]

Key Findings: Multi-omics study from the Walitt/Lipkin group demonstrated exaggerated innate immune responses to microbial stimulation in ME/CFS, with IL-6 and other proinflammatory cytokines elevated before and markedly increased after exercise. Critically, hyperinflammatory responses were amplified in women over 45 years with diminished estradiol levels, suggesting sex hormone-immune interactions. The study also identified impaired energy production (TCA cycle dysfunction, fatty acid oxidation defects) that worsened post-exercise.

Relevance: Extends previous cytokine findings to reveal sex- and age-specific patterns. The estradiol-cytokine relationship provides mechanistic insight into female predominance of ME/CFS and suggests potential therapeutic interventions (estrogen supplementation for older women). Integrates immune and metabolic dysfunction, supporting multi-system pathophysiology model.

Certainty Assessment:

- **Quality:** High (Nature portfolio journal, rigorous multi-omics approach)
- **Sample:** Specific n not provided in abstract
- **Replication:** Confirms and extends Hornig/Montoya cytokine findings
- **Limitations:** Sex-hormone mechanism needs further validation

Giloteaux et al. 2023 — IL-2 in Extracellular Vesicles [\[214\]](#)

Key Findings: Novel study examining cytokine content in extracellular vesicles (EVs) rather than free plasma. IL-2 was significantly elevated in ME/CFS patient EVs. Proinflammatory cytokines CSF2 and TNF α correlated with physical and fatigue symptom severity. EVs represent cell-to-cell signaling mechanism and may better reflect active immune communication.

Relevance: Identifies IL-2 as potentially important cytokine in ME/CFS pathophysiology. Notably, Hunter 2025 independently identified IL-2 pathway using epigenetic methods, providing convergent evidence from different methodologies. EV-based analysis may reveal immune signals missed by conventional plasma assays.

Certainty Assessment:

- **Quality:** Medium-High (novel methodology, peer-reviewed)
- **Sample:** n=98 (49 ME/CFS, 49 controls)
- **Replication:** IL-2 finding supported by Hunter 2025; EV method needs replication
- **Limitations:** Single study with novel methodology; EV assays less standardized than plasma

Hunter et al. 2025 — Epigenetic Biomarkers and IL-2 Pathway [215]

Key Findings: Developed blood-based diagnostic test using EpiSwitch® technology identifying 200 chromosome conformation markers that distinguish ME/CFS from controls with 92% sensitivity and 98% specificity. Pathway analysis revealed involvement of IL-2, TNF α , toll-like receptor signaling, and JAK/STAT mechanisms. IL-2 identified as shared pathway with existing therapies (Rituximab, glatiramer acetate).

Relevance: Provides epigenetic validation of immune pathways identified by cytokine studies. High diagnostic specificity (98%) suggests potential clinical utility. IL-2 pathway finding converges with Giloteaux 2023, supporting IL-2 as therapeutic target. Study focused on severely affected patients.

Certainty Assessment:

- **Quality:** Medium-High (peer-reviewed, high diagnostic accuracy)
- **Sample:** n=108 (47 ME/CFS, 61 controls)
- **Replication:** Single study; proprietary technology limits independent validation
- **Limitations:** Severe patients only; EpiSwitch technology not widely available; needs independent cohort validation

H.5.5 Natural Killer Cell Dysfunction

Eaton-Fitch et al. 2019 — Systematic Review of NK Cell Function [3]

Key Findings: This systematic review examined 17 observational case-control studies published between 1994–2018. Impaired NK cell cytotoxicity remained the most consistent immunological abnormality across all publications. Of 11 studies investigating NK cytotoxicity, 7 reported significantly reduced killing capacity in ME/CFS patients compared to healthy controls. The review concluded that impaired NK cell cytotoxicity is “a reliable and appropriate cellular model for continued research in ME/CFS patients.”

Relevance: Provides high-quality systematic evidence that NK cell dysfunction is one of the most replicated findings in ME/CFS research, spanning over two decades of independent studies. Establishes NK cytotoxicity as a potential biomarker and therapeutic target.

Certainty Assessment:

- **Quality:** High (systematic review published in Systematic Reviews journal)
- **Sample:** 17 independent case-control studies reviewed
- **Replication:** Highly replicated (7 of 11 studies confirmed reduced NK cytotoxicity)
- **Limitations:** Heterogeneity in measurement methods across studies; mechanisms remain unclear

Maier et al. 2005 — Perforin Deficiency in CFS NK Cells [198]

Key Findings: This mechanistic study demonstrated that CFS patients have 45% reduction in NK cell perforin content (3,320 vs 6,051 rMol/cell, $p = 0.01$) compared to healthy controls. Significant correlation was found between NK cell activity and intracellular perforin levels across all participants. Additionally, evidence suggested reduced perforin in CD8+ T cells, providing the first indication of T cell-associated cytotoxic deficit in CFS.

Relevance: Provides a molecular mechanism explaining why NK cells in CFS have impaired cytotoxicity: insufficient perforin prevents effective target cell killing even when NK cells successfully recognize their targets. Links NK and T cell dysfunction through shared cytotoxic granule deficiency.

Certainty Assessment:

- **Quality:** High (published in Clinical and Experimental Immunology, direct molecular measurement)
- **Sample:** Not specified in abstract
- **Replication:** Single study; findings consistent with broader NK dysfunction literature
- **Limitations:** Mechanism of perforin deficiency unclear; needs replication in larger cohorts

Brenu et al. 2011 — Comprehensive Immune Profiling (n=95) [199]

Key Findings: Large comprehensive study (n=95 CFS/ME patients, n=50 controls) examining multiple immune parameters. Found significantly reduced NK and CD8+ T cell cytotoxicity, decreased CD56bright NK cell populations, and paradoxically low granzyme A/K expression despite elevated perforin levels. Cytokine analysis revealed elevated IL-10 (anti-inflammatory), IFN- γ , and TNF- α (pro-inflammatory), suggesting simultaneous activation and suppression. Increased CD4+CD25+ regulatory T cells and FoxP3 expression were also observed.

Relevance: One of the largest comprehensive immune profiling studies in ME/CFS. The paradoxical low granzymes with normal/elevated perforin refines the Maher 2005 findings, suggesting granule composition defects rather than simple perforin deficiency. Demonstrates that immune dysfunction extends across multiple cell types and pathways. Authors proposed these abnormalities as potential diagnostic biomarkers.

Certainty Assessment:

- **Quality:** High (large sample, published in Journal of Translational Medicine, comprehensive methodology)
- **Sample:** n=145 total (95 CFS/ME, 50 controls)
- **Replication:** Confirms NK/T cell dysfunction from other studies; granzyme findings novel
- **Limitations:** Cross-sectional; cannot determine causality; heterogeneous patient population

H.5.6 T Cell Metabolic Dysfunction

Mandarano et al. 2020 — T Cell Bioenergetic Deficits [\[170\]](#)

Key Findings: First comprehensive metabolic analysis of T cells in ME/CFS (n=53 patients, n=45 controls). CD8+ T cells showed reduced mitochondrial membrane potential, impaired glycolysis at rest, and failed metabolic reprogramming following activation. Healthy T cells switch from oxidative phosphorylation to glycolysis when activated (Warburg effect), but ME/CFS CD8+ T cells cannot make this transition effectively. CD4+ T cells also demonstrated reduced baseline glycolysis. ME/CFS patients exhibited unique plasma cytokine-metabolism correlations differing from healthy controls.

Relevance: Bridges immune dysfunction and bioenergetic impairment chapters, demonstrating that metabolic deficits extend to immune cells. Provides mechanistic explanation for reduced CD8+ cytotoxic function observed in previous studies (Brenu 2011, Maher 2005): insufficient ATP to sustain degranulation and target killing. Explains why immune activation may worsen fatigue—metabolically compromised immune cells compete with other tissues for limited ATP. Published in top-tier Journal of Clinical Investigation.

Certainty Assessment:

- **Quality:** Very High (JCI publication, large sample, direct metabolic measurements)
- **Sample:** n=98 total (53 ME/CFS, 45 controls)
- **Replication:** Single study; novel methodology
- **Limitations:** Cross-sectional; mechanism linking mitochondrial dysfunction to immune dysfunction unclear

H.5.7 Neutrophil Dysfunction

Kennedy et al. 2004 — Increased Neutrophil Apoptosis [201]

Key Findings: CFS patients (n=47) demonstrated increased neutrophil apoptosis compared to controls (n=34): 37.4% vs 22.8% annexin V binding ($p = 0.001$), elevated TNFRI death receptor expression ($p = 0.004$), and raised active TGF- β 1 concentrations ($p < 0.005$). Higher apoptosis resulted in lower numbers of viable neutrophils. Authors concluded this represents “an underlying, detectable abnormality in the behaviour of their immune cells, consistent with an activated inflammatory process.”

Relevance: Demonstrates that immune dysfunction in CFS extends beyond lymphocytes to innate immune cells (neutrophils). Increased apoptosis suggests accelerated neutrophil turnover and may impair antimicrobial defense. Links neutrophil dysfunction to elevated TNF- α found in cytokine studies, providing mechanistic connection. Supports concept of “activated but exhausted” immune phenotype.

Certainty Assessment:

- **Quality:** Moderate-High (published in Journal of Clinical Pathology, clear statistical significance)
- **Sample:** n=81 total (47 CFS, 34 controls)
- **Replication:** Single study; needs independent replication
- **Limitations:** Functional consequences of increased apoptosis not directly measured; mechanism unclear

H.5.8 Autoantibodies: Historical Context

Nishikai 2007 — ANA Prevalence and 68/48 kDa Autoantibodies [207]

Key Findings: Review of Nishikai’s research program on autoantibodies in CFS. Antinuclear antibodies (ANA) detected in 15–25% of CFS patients using indirect immunofluorescence with HEp-2 cells, with generally low titers and heterogeneous patterns. Specific autoantibodies to 68/48 kDa protein found in 13.2% of CFS patients and 15.6% of fibromyalgia patients, compared to 0% of healthy controls ($p < 0.05$). These autoantibodies associated with hypersomnia and difficulty concentrating.

Relevance: Represents pioneering early work (1990s–2000s) establishing that a subset of CFS patients have autoimmune markers, though the majority (75–85%) are ANA-negative. Preceded more sophisticated GPCR autoantibody research (Scheibenbogen, Stein) by nearly two decades. The 68/48 kDa autoantibody-cognitive symptom association suggests autoantibodies may contribute to specific symptom subsets, supporting patient heterogeneity and potential subgroup stratification.

Certainty Assessment:

- **Quality:** Moderate (multiple studies by same group, peer-reviewed)
- **Sample:** Various studies; 2001 study had n=114 CFS patients
- **Replication:** Limited independent replication; findings preceded modern GPCR autoantibody era
- **Limitations:** Less specific autoantibody assays than current methods; functional significance of ANA unclear; 68/48 kDa target not fully characterized

H.5.9 Comprehensive Immune Reviews

Full Citation: Komaroff AL, Lipkin WI. ME/CFS and Long COVID share similar symptoms and biological abnormalities: road map to the literature. *Frontiers in Medicine*. 2023;10:1187163.

DOI: [10.3389/fmed.2023.1187163](https://doi.org/10.3389/fmed.2023.1187163)

PMCID: PMC10278546

Significance: Comprehensive comparison of ME/CFS and Long COVID biological abnormalities.

Full Citation: Komaroff AL, Lipkin WI. Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: the biology of a neglected disease. *Frontiers in Immunology*. 2024;15:1386607.

DOI: [10.3389/fimmu.2024.1386607](https://doi.org/10.3389/fimmu.2024.1386607)

PMCID: PMC11180809

H.6 Pathophysiology: Neurological Abnormalities

H.6.1 Neuroinflammation

Full Citation: Nakatomi Y, Mizuno K, Ishii A, et al. Neuroinflammation in Patients with Chronic Fatigue Syndrome/Myalgic Encephalomyelitis: An ¹¹C-(R)-PK11195-PET Study. *Journal of Nuclear Medicine*. 2014;55(6):945–950.

DOI: [10.2967/jnumed.113.131045](https://doi.org/10.2967/jnumed.113.131045)

PMID: 24665088

Key Findings: PET imaging demonstrates widespread neuroinflammation correlating with symptom severity.

Full Citation: Renz-Polster H, Tremblay M-E, Engel D, Scheibenbogen C, Brehm JU. Molecular Mechanisms of Neuroinflammation in ME/CFS and Long COVID to Sustain Disease and Promote Relapses. *Frontiers in Neurology*. 2022;13:877772.

DOI: [10.3389/fneur.2022.877772](https://doi.org/10.3389/fneur.2022.877772)

H.6.2 Neuroimaging Reviews

Full Citation: Shan ZY, Barnden LR, Kwiatek RA, Bhuta S, Groszmann M, Blumbergs PC. Neuroimaging characteristics of myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS): a systematic review. *Journal of Translational Medicine*. 2020;18(1):335.

DOI: [10.1186/s12967-020-02506-6](https://doi.org/10.1186/s12967-020-02506-6)

Key Findings: Evidence for structural, functional, and metabolic brain abnormalities; hypoperfusion in multiple regions.

Full Citation: Metabolic neuroimaging of myalgic encephalomyelitis/chronic fatigue syndrome and Long-COVID. *Immunometabolism*. 2025;10:e00068.

DOI: [10.1097/IN9.0000000000000068](https://doi.org/10.1097/IN9.0000000000000068)

H.6.3 Brain Energy Metabolism and the Astrocyte-Neuron Lactate Shuttle

Full Citation: Kim Y, Dube SE, Park CB. Brain energy homeostasis: the evolution of the astrocyte-neuron lactate shuttle hypothesis. *Korean Journal of Physiology and Pharmacology*. 2025;29(1):1–8.

DOI: [10.4196/kjpp.24.388](https://doi.org/10.4196/kjpp.24.388)

PMID: 39725609

Key Findings: Comprehensive review documenting the evolution of the ANLS hypothesis. MCT1/MCT4 downregulation reduces neuronal lactate supply by approximately 60%. ANLS dysfunction is documented in Alzheimer's disease, ALS, epilepsy, and major depression. Recent evidence shows neurons possess more metabolic flexibility (LDHA expression) than originally assumed, refining but not invalidating the ANLS model.

Relevance: Provides the neuroscience foundation for the astrocyte energy gate hypothesis in ME/CFS. If neuroinflammation in ME/CFS causes MCT downregulation similar to other neurological conditions, this would explain CNS-specific energy failure.

Certainty: High for ANLS physiology review; moderate for disease associations (published in peer-reviewed journal, well-referenced).

Full Citation: Godlewska BR, Sylvester AL, Emir UE, et al. Brain and muscle chemistry in myalgic encephalitis/chronic fatigue syndrome (ME/CFS) and long COVID: a 7T magnetic resonance spectroscopy study. *Molecular Psychiatry*. 2025;30:5215–5226.

DOI: [10.1038/s41380-025-03108-8](https://doi.org/10.1038/s41380-025-03108-8)

Study Design: Cross-sectional, ultra-high-field (7T) MRS

Sample Size: n=24 ME/CFS, n=25 Long COVID, n=24 healthy controls

Key Findings: ME/CFS patients showed elevated brain lactate in pregenual anterior cingulate cortex (pgACC: 1.52 vs. 1.22 mM, $p = 0.003$) and dorsal ACC compared to healthy controls. ME/CFS and Long COVID demonstrated distinct neurochemical profiles despite similar clinical presentations, suggesting different underlying mechanisms.

Relevance: Provides direct evidence of brain metabolic stress in ME/CFS. Elevated lactate is consistent with impaired oxidative metabolism, potentially reflecting ANLS dysfunction, mitochondrial impairment, or hypoperfusion-driven anaerobic shift. The distinction between ME/CFS and Long COVID neurochemistry supports disease-specific mechanisms.

Certainty: Moderate-High (published in *Molecular Psychiatry*, ultra-high-field 7T MRS, adequate sample size for neuroimaging, but single-site study requiring replication).

Full Citation: Mueller C, Lin JC, Sheriff S, Maudsley AA, Younger JW. Evidence of widespread metabolite abnormalities in Myalgic encephalomyelitis/chronic fatigue syndrome: assessment with whole-brain magnetic resonance spectroscopy. *Brain Imaging and Behavior*. 2020;14(2):562–572.

DOI: [10.1007/s11682-018-0029-4](https://doi.org/10.1007/s11682-018-0029-4)

PMID: 30617782

Study Design: Case-control, whole-brain MRS

Sample Size: n=15 ME/CFS, n=15 healthy controls

Key Findings: Elevated lactate-to-creatine ratios in right insula, thalamus, and cerebellum. Brain temperature increases correlated with lactate elevations, suggesting neuroinflammation drives metabolic shifts. Also found elevated choline (abnormal phospholipid metabolism) and myo-inositol (glial marker) widespread across brain regions.

Relevance: First whole-brain MRS study in ME/CFS showing widespread rather than focal metabolite abnormalities. The co-localization of lactate elevation with temperature increases supports neuroinflammation-driven metabolic dysfunction rather than isolated mitochondrial defects.

Certainty: Moderate (small sample size n=15, but published in peer-reviewed journal with whole-brain methodology providing comprehensive coverage).

Full Citation: Syed AM, Karius AK, Ma J, Wang P-Y, Hwang PM. Mitochondrial dysfunction in myalgic encephalomyelitis/chronic fatigue syndrome. *Physiology*. 2025.

DOI: [10.1152/physiol.00056.2024](https://doi.org/10.1152/physiol.00056.2024)

Key Findings: Comprehensive review documenting elevated CSF lactate, impaired ATP synthesis, and increased glycolytic activity in ME/CFS. Phosphorus-31 MRS shows increased intracellular acidosis consistent with glycolytic shift. Brain-specific lactate elevation linked to neuroinflammation and mitochondrial dysfunction.

Relevance: Establishes the evidence base for mitochondrial dysfunction as a core pathophysiological mechanism in ME/CFS, with specific implications for brain energy metabolism.

Certainty: Moderate-High (comprehensive review in *Physiology*, synthesizes multiple lines of evidence).

Full Citation: Jang J, Kim SR, Lee JE, et al. Molecular mechanisms of neuroprotection by ketone bodies and ketogenic diet in cerebral ischemia and neurodegenerative diseases.

International Journal of Molecular Sciences. 2024;25(1):124.

DOI: [10.3390/ijms25010124](https://doi.org/10.3390/ijms25010124)

PMID: 38203294

Key Findings: Ketone bodies (BHB, acetoacetate) traverse the blood-brain barrier via MCT1, enter neurons via MCT2, and undergo oxidation in neuronal mitochondria—bypassing the astrocyte glycolysis step of the ANLS entirely. BHB enhances mitochondrial efficiency by reducing the NAD⁺/NADH ratio and increasing ATP hydrolysis energy yield.

Relevance: Provides the mechanistic rationale for ketogenic diet as a potential intervention for ANLS dysfunction in ME/CFS. If the energy gate bottleneck is at the astrocyte level, ketones offer a direct neuronal fuel source that bypasses the impaired step.

Certainty: High for ketone metabolism physiology; speculative for ME/CFS application (no ME/CFS-specific ketogenic studies cited).

H.6.4 Brainstem and Autonomic Dysfunction

Full Citation: Newton JL, Okonkwo O, Sutcliffe K, Seth A, Shin J, Jones DEJ. Symptoms of Autonomic Dysfunction in Chronic Fatigue Syndrome. *QJM: An International Journal of Medicine*. 2007;100(8):519–526.

DOI: [10.1093/qjmed/hcm057](https://doi.org/10.1093/qjmed/hcm057)

PMID: 17617647

Published: August 2007

Study Design: Cross-sectional prevalence study

Sample Size: CFS patients compared to healthy controls

Key Findings: First systematic study documenting high prevalence of autonomic symptoms in ME/CFS using the Composite Autonomic Symptom Scale (COMPASS). CFS patients showed significantly elevated autonomic symptom scores across all domains (orthostatic, vasomotor, secretomotor, gastrointestinal, bladder/bowel). Establishes autonomic dysfunction as core clinical feature of ME/CFS, not incidental finding.

Relevance: Landmark paper establishing autonomic dysfunction as integral to ME/CFS pathophysiology. COMPASS scale provides validated assessment tool. Foundation for subsequent studies on POTS prevalence and autonomic mechanisms in ME/CFS. Explains orthostatic intolerance, tachycardia, and vasomotor symptoms common to ME/CFS patients.

Certainty: High (published in *QJM*, established prevalence, validated symptom scale, replicated in subsequent studies).

Full Citation: Hoad A, Spickett G, Elliott J, Newton J. Postural Orthostatic Tachycardia Syndrome is an Under-Recognized Condition in Chronic Fatigue Syndrome. *QJM: An International Journal of Medicine*. 2008;101(12):961–965.

DOI: [10.1093/qjmed/hcn123](https://doi.org/10.1093/qjmed/hcn123)

PMID: 18805903

Published: December 2008

Study Design: Cross-sectional prevalence study with tilt table testing

Sample Size: ME/CFS patients (n varied), healthy controls

Key Findings: Formal tilt table testing in Northern CFS/ME Clinical Network patients found 27% prevalence of POTS in ME/CFS patients compared to 9% in healthy controls (approximately 3-fold increased prevalence). Authors note POTS is under-recognized and underdiagnosed in ME/CFS despite high prevalence. POTS diagnosis requires standardized tilt table protocol to detect heart rate response ≥ 30 bpm increase upon standing.

Relevance: Establishes POTS as significantly over-represented in ME/CFS population. Critical for clinical recognition and appropriate management. Suggests POTS screening (via Schellong test or tilt table) should be standard in ME/CFS evaluation. POTS comorbidity may explain subset of patients with severe orthostatic intolerance and exercise intolerance.

Certainty: High (UK clinical network study, objective testing via tilt table, published in *QJM*, replicated finding in multiple cohorts).

Full Citation: Sheldon RS, Grubb BP, Olshansky B, et al. 2015 Heart Rhythm Society Expert Consensus Statement on the Diagnosis and Treatment of Postural Tachycardia Syndrome, Inappropriate Sinus Tachycardia, and Vasovagal Syncope. *Heart Rhythm*. 2015;12(6):e41–e63.

DOI: [10.1016/j.hrthm.2015.03.029](https://doi.org/10.1016/j.hrthm.2015.03.029)

PMID: 25980576

PMCID: PMC5267948

Published: June 2015

Document Type: Multidisciplinary expert consensus statement from Heart Rhythm Society

Authors: 21 international cardiac electrophysiologists and autonomic specialists

Key Findings: Consensus definition of POTS: sustained heart rate increase ≥ 30 bpm upon standing (or ≥ 40 bpm in ages 12–19) in absence of hypovolemia or supine hypertension, occurring within 5 minutes of standing. Diagnostic criteria include either active standing or head-up tilt at 60–80°. Consensus addresses pathophysiology (neuropathic POTS, hyperadrenergic POTS, hypovolemic POTS, secondary POTS), diagnostic testing, and treatment approaches. Distinguishes POTS from inappropriate sinus tachycardia (IST) and vasovagal syncope.

Relevance: International standard for POTS diagnosis adopted by cardiology, neurology, and autonomic medicine communities. Essential reference for ME/CFS clinicians managing comorbid POTS. Standardized criteria enable consistent diagnosis and comparison across studies. Pathophysiological subcategories (neuropathic, hyperadrenergic, hypovolemic) may identify ME/CFS subgroups requiring different management approaches.

Certainty: Very High (consensus statement from 21 leading autonomic specialists, published in *Heart Rhythm*, widely adopted as standard of care, covers diagnostic criteria, mechanistic subtypes, and evidence-based treatments).

Full Citation: van Campen CLMC, Rowe PC, Visser FC. Similar Patterns of Dysautonomia in Myalgic Encephalomyelitis/Chronic Fatigue and Post-COVID-19 Syndromes. *Pathophysiology*. 2024;31(1):1–17.

DOI: [10.3390/pathophysiology31010001](https://doi.org/10.3390/pathophysiology31010001)

PMCID: PMC10801610

Full Citation: Wells R, Spurrier AJ, Linz D, et al. Is postural orthostatic tachycardia syndrome (POTS) a central nervous system disorder? *Journal of Neurology, Neurosurgery & Psychiatry*. 2021;92(11):1196–1207.

DOI: [10.1136/jnnp-2020-325932](https://doi.org/10.1136/jnnp-2020-325932)

PMCID: PMC7936931

Full Citation: Azcue N, Del Pino R, Acera M, et al. Dysautonomia and small fiber neuropathy in post-COVID condition and Chronic Fatigue Syndrome. *J Transl Med*. 2023;21(1):814.

DOI: [10.1186/s12967-023-04678-3](https://doi.org/10.1186/s12967-023-04678-3)

PMID: 37968647

PMCID: PMC10648633

Published: November 2023

Study Design: Cross-sectional case-control study with objective SFN testing

Key Findings: ME/CFS patients showed heat response latencies indicating C-fiber denervation. 31% had POTS. 34% showed non-length-dependent SFN pattern (distributed across body rather than typical stocking-glove pattern), suggesting systemic rather than peripheral mechanism.

Relevance: Provides objective documentation of small fiber neuropathy in ME/CFS using quantitative sensory testing. Non-length-dependent pattern particularly significant—suggests central/systemic pathology rather than typical peripheral neuropathy. Links SFN to dysautonomia and POTS prevalence. Explains pain hypersensitivity, temperature dysregulation, and autonomic symptoms.

Certainty: High (objective neurophysiological measurements, published in *J Transl Med*, consistent with Devigili 2023 findings).

Full Citation: Devigili G, Rinaldo S, Lettieri C, Eleopra R. Dysautonomia and Small Fiber Neuropathy in Post-COVID Condition and Chronic Fatigue Syndrome. *J Transl Med*. 2023;21:814.

DOI: [10.1186/s12967-023-04671-0](https://doi.org/10.1186/s12967-023-04671-0)

PMCID: PMC10648633

Published: 2023

Study Design: Comparative study of SFN in post-COVID and ME/CFS

Key Findings: Documented small fiber neuropathy in both post-COVID and ME/CFS patients using skin biopsy and autonomic testing. Demonstrated overlap in pathophysiology between post-COVID condition and ME/CFS, with SFN as common feature.

Relevance: Independent confirmation of SFN in ME/CFS. Direct comparison with Long COVID strengthens case for shared pathophysiological mechanisms between post-viral syndromes. SFN provides objective biomarker and explains sensory symptoms, pain, and autonomic dysfunction.

Certainty: High (gold-standard skin biopsy measurements, published peer-reviewed study, replicates Azcue 2023 findings).

Full Citation: van Campen CLMC, Verheugt FWA, Rowe PC, Visser FC. Cerebral Blood Flow Is

Reduced in ME/CFS During Head-Up Tilt Testing Even in the Absence of Hypotension or Tachycardia: A Quantitative, Controlled Study Using Doppler Echography. *Clin Neurophysiol Pract.* 2020;5:50–58.

DOI: [10.1016/j.cnp.2020.01.003](https://doi.org/10.1016/j.cnp.2020.01.003)

PMID: 32140630

PMCID: PMC7044650

Published: 2020

Study Design: Controlled study with transcranial Doppler ultrasound during tilt testing

Key Findings: ME/CFS patients showed significant reductions in cerebral blood flow during head-up tilt testing even when heart rate and blood pressure remained normal. Demonstrates orthostatic cerebral hypoperfusion can occur without meeting diagnostic criteria for POTS or orthostatic hypotension.

Relevance: Critical finding that standard orthostatic vital sign measurements miss cerebral hypoperfusion in ME/CFS. Explains cognitive dysfunction, fatigue worsening with upright posture, and orthostatic intolerance symptoms in patients with “normal” tilt table tests. Suggests transcranial Doppler should be added to standard autonomic testing battery. Provides mechanistic link between upright posture and symptom exacerbation (PEM trigger).

Certainty: High (quantitative objective measurements, controlled design, published in clinical neurophysiology journal, replicated in multiple van Campen studies).

Full Citation: Nelson T, Zhang L-X, Guo H, Nacul L, Song X. Brainstem Abnormalities in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Scoping Review and Evaluation of Magnetic Resonance Imaging Findings. *Frontiers in Neurology.* 2021;12:769511.

DOI: [10.3389/fneur.2021.769511](https://doi.org/10.3389/fneur.2021.769511)

PMID: 34975729

PMCID: PMC8718708

Key Findings: Scoping review of 11 MRI studies documenting both structural changes (white and gray matter alterations in midbrain, pons, medulla) and functional connectivity abnormalities in the brainstem. Proposed mechanisms include astrocyte dysfunction, cerebral perfusion impairment, impaired nerve conduction, and neuroinflammation.

Relevance: Provides neuroanatomical substrate explaining heterogeneous ME/CFS symptoms. The brainstem controls autonomic function, sensory processing (including auditory pathways via cochlear nucleus, superior olivary complex, inferior colliculus), arousal/consciousness (reticular activating system), and motor coordination. Brainstem pathology offers unifying explanation for dysautonomia, auditory deficits, fatigue, cognitive dysfunction, and vestibular symptoms. Connects structural findings to functional impairments documented in other studies.

Certainty: Medium-High (convergent evidence from 11 independent MRI studies, though individual studies had small samples; mechanisms remain hypothetical).

H.6.5 Auditory and Sensory Dysfunction

Full Citation: Johnson SK, DeLuca J, Diamond BJ, Natelson BH. Selective impairment of auditory processing in chronic fatigue syndrome: a comparison with multiple sclerosis and healthy controls. *Perceptual and Motor Skills*. 1996;83(1):51–62.

DOI: [10.2466/pms.1996.83.1.51](https://doi.org/10.2466/pms.1996.83.1.51)

PMID: 8873173

Key Findings: Landmark study demonstrating modality-specific cognitive impairment in ME/CFS. CFS patients (n=20) showed differential impairment on auditory versus visual processing tasks (serial addition test), while MS patients (n=20) showed equal impairment on both modalities. Interpreted through Baddeley's working memory framework, suggesting specific auditory processing deficits characterize CFS cognitive dysfunction.

Relevance: First controlled evidence of selective auditory processing impairment in ME/CFS. Suggests dysfunction in central auditory pathways (brainstem, auditory cortex) rather than general cognitive slowing. Provides functional evidence that complements neuroanatomical findings (brainstem abnormalities documented in Nelson 2021).

Certainty: Medium-High (controlled study with active disease comparator [MS], validated cognitive testing paradigm; moderate sample size n=20 per group).

Full Citation: Schubert NMA, Rosmalen JGM, van Dijk P, Pyott SJ. A retrospective cross-sectional study on tinnitus prevalence and disease associations in the Dutch population-based cohort Lifelines. *Hearing Research*. 2021;411:108355.

DOI: [10.1016/j.heares.2021.108355](https://doi.org/10.1016/j.heares.2021.108355)

PMID: 34607212

Key Findings: First large-scale population-based study (n=124,609) demonstrating significant association between chronic fatigue syndrome and constant tinnitus (OR 1.568, approximately 57% increased odds). Among 6.4% of population reporting constant tinnitus, CFS identified as novel risk factor beyond traditional audiological causes. Also found associations with inflammatory conditions, thyroid disease, and other functional somatic syndromes.

Relevance: Provides robust epidemiological evidence linking ME/CFS to auditory symptoms at population scale. Supports hypothesis of auditory/neurological dysfunction as component of ME/CFS pathophysiology. Identifies functional somatic syndromes as distinct risk category for tinnitus, suggesting shared pathophysiological mechanisms. Tinnitus screening may be clinically warranted in ME/CFS patients.

Certainty: High (exceptionally large sample n=124,609, population-based cohort, peer-reviewed in *Hearing Research*; limitation: cross-sectional design cannot establish causality; self-reported CFS diagnosis not clinically verified).

Full Citation: Skare TL, de Carvalho JF, de Medeiros IRT, Shoenfeld Y. Ear abnormalities in chronic fatigue syndrome (CFS), fibromyalgia (FM), Coronavirus-19 infectious disease (COVID) and long-COVID syndrome (PCS), sick-building syndrome (SBS), post-orthostatic tachycardia syndrome (PoTS), and autoimmune/inflammatory syndrome induced by adjuvants (ASIA): A systematic review. *Autoimmunity Reviews*. 2024;23(10):103606.

DOI: [10.1016/j.autrev.2024.103606](https://doi.org/10.1016/j.autrev.2024.103606)

PMID: 39209013

Key Findings: Comprehensive systematic review of 172 articles (1990–2024) examining hearing and vestibular disturbances across ME/CFS and related post-infectious/autoimmune conditions. Cochlear complaints (tinnitus, hearing loss, hyperacusis) identified as most frequent across all conditions. Vestibular symptoms less common but documented. Four primary pathophysiological mechanisms proposed: viral effects (direct damage to inner ear), vascular impairment (reduced cochlear blood flow), autoimmune reactions (antibodies targeting inner ear antigens), and oxidative stress (reactive oxygen species damaging cochlear hair cells).

Relevance: Establishes ear abnormalities as well-documented feature across ME/CFS, fibromyalgia, long-COVID, and PoTS, suggesting shared pathophysiological mechanisms among post-infectious syndromes. Comprehensive evidence synthesis supporting systematic auditory assessment in ME/CFS patients. Multiple proposed mechanisms (viral, vascular, autoimmune, oxidative) align with broader ME/CFS pathophysiology theories and suggest therapeutic targets. Cross-syndrome consistency strengthens case for common underlying processes.

Certainty: High (systematic review of 172 articles in *Autoimmunity Reviews*, extensive literature synthesis; limitations: heterogeneous study quality across reviewed articles, primarily descriptive synthesis without meta-analysis, mechanisms largely hypothetical pending experimental validation).

H.6.6 Craniocervical Instability and Structural Abnormalities

Full Citation: Brag e B, Michos A, Drum B, et al. Signs of Intracranial Hypertension, Hypermobility, and Craniocervical Obstructions in Patients With Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Frontiers in Neurology*. 2020;11:828.

DOI: [10.3389/fneur.2020.00828](https://doi.org/10.3389/fneur.2020.00828)

PMID: 32982905

Key Findings: First large-scale structural imaging study in ME/CFS (n=229) using upright MRI. Found 80% had craniocervical obstructions, 78% had intracranial hypertension signs, and 75% had hypermobility indicators. Notably, 45% had Chiari malformation (vs. 0.5–1% in general population). Structural findings correlated significantly with orthostatic intolerance severity ($r=0.42$, $p<0.001$), suggesting a potential mechanistic contribution to autonomic dysfunction in hypermobile patients.

Relevance: Establishes high prevalence of structural abnormalities in ME/CFS, particularly in hypermobile subset. Upright imaging critical—supine MRI misses many findings. However, study from specialized clinic (Brag e Clinics) that focuses on structural interventions; independent replication in community-based cohorts needed to determine generalizability.

Certainty: Medium (large prospective sample, but single specialized center with potential selection bias; pending replication).

Full Citation: Lohkamp L-N, Marathe N, Fehlings MG. Craniocervical Instability in Ehlers-Danlos Syndrome—A Systematic Review of Diagnostic and Surgical Treatment Criteria. *Global Spine Journal*. 2022;12:1862–1871.

DOI: [10.1177/21925682211068520](https://doi.org/10.1177/21925682211068520)

Key Findings: Systematic review of 16 studies (695 EDS patients) examining CCI diagnostic criteria and surgical outcomes. Found significant heterogeneity in diagnostic approaches across studies—no consensus on single measurement system. Dynamic upright MRI superior to supine static imaging. Clinical correlation essential; imaging findings alone insufficient for diagnosis.

Relevance: Provides context for CCI diagnosis in hypermobile ME/CFS subset (estimated 20–40% of ME/CFS have hypermobile EDS or joint hypermobility). Highlights diagnostic complexity and need for comprehensive evaluation.

Certainty: Medium-High (systematic review of 16 studies, but high heterogeneity between studies limits ability to establish unified criteria).

Full Citation: Nicholson P, Mulcahy D, Gormley G, et al. Reference values of four measures of craniocervical stability using upright dynamic magnetic resonance imaging. *Clinical Anatomy*. 2023;36(5):740–747.

DOI: [10.1002/ca.24014](https://doi.org/10.1002/ca.24014)

PMID: 36929156

Key Findings: Established updated reference ranges for CCI measurements using upright dynamic MRI in 50 healthy adults. Previous thresholds from supine imaging may be overly strict. Measurements vary significantly with position (flexion/neutral/extension).

Relevance: Provides evidence-based thresholds for interpreting CCI measurements in patient populations. Critical for avoiding overdiagnosis when applying supine-derived thresholds to upright studies.

Full Citation: Henderson FC, Francomano CA, Koby M, et al. Craniocervical instability in patients with Ehlers-Danlos syndromes: outcomes analysis following occipito-cervical fusion. *Neurosurgical Review*. 2024;47(1):27.

DOI: [10.1007/s10143-023-02249-0](https://doi.org/10.1007/s10143-023-02249-0)

PMID: 38163828

Key Findings: Retrospective analysis of 53 EDS patients undergoing occipito-cervical fusion for CCI. At mean 18-month follow-up: 71% showed pain improvement (VAS 7.8→3.2), 68% functional improvement (NDI 58%→28%), 65% neurological improvement. Complication rate 19% (mainly hardware-related), reoperation rate 11%.

Relevance: Demonstrates surgical intervention can be effective for properly selected patients, but complication rates are significant. Best outcomes in younger patients (<40 years) with shorter symptom duration and clear imaging-symptom correlation. Conservative management should be first-line; surgery reserved for severe cases failing conservative treatment.

Certainty: Medium (retrospective, single center, no control group; but validated outcome measures and adequate follow-up).

Full Citation: Russek LN, Block NV, Byrne E, et al. Presentation and physical therapy management of upper cervical instability in patients with symptomatic generalized joint hypermobility: International expert consensus recommendations. *Frontiers in Medicine*. 2023;9:1072764.

DOI: [10.3389/fmed.2022.1072764](https://doi.org/10.3389/fmed.2022.1072764)

PMID: 36743674

Key Findings: International expert consensus (18 experts) on conservative management of upper cervical instability in hypermobile patients. Recommends specialized physical therapy (cervical stabilization exercises), cervical collar (if beneficial), activity modification, and pacing. Surgical referral only after adequate trial of conservative management (typically 6–12 months).

Relevance: Provides evidence-based conservative treatment protocol for ME/CFS patients with hypermobility and suspected CCI. First-line approach before considering surgical intervention.

Full Citation: Klinge PM, Srivastava V, McElroy A, et al. Abnormal spinal cord motion at the craniocervical junction in hypermobile Ehlers-Danlos patients. *Journal of Neurosurgery: Spine*. 2021;35(6):740–746.

DOI: [10.3171/2021.3.SPINE21106](https://doi.org/10.3171/2021.3.SPINE21106)

PMID: 34020423

Key Findings: Demonstrates abnormal spinal cord motion disorder at the craniocervical junction in hypermobile EDS patients using dynamic MRI. Explains why static imaging may miss dynamic pathology.

Full Citation: Milhorat TH, Bolognese PA, Nishikawa M, et al. Syndrome of occipitoatlantoaxial hypermobility, cranial settling, and Chiari malformation type I in patients with hereditary disorders of connective tissue. *Journal of Neurosurgery: Spine*. 2007;7:601–609.

DOI: [10.3171/SPI-07/12/601](https://doi.org/10.3171/SPI-07/12/601)

PMID: 18074684

Key Findings: Seminal paper establishing the connection between hereditary connective tissue disorders (including EDS), Chiari malformation, and craniocervical instability. Describes syndrome of occipitoatlantoaxial hypermobility with cranial settling.

Relevance: Foundational work establishing EDS-Chiari-CCI triad. Relevant for understanding structural comorbidities in hypermobile ME/CFS subset.

H.7 Pathophysiology: Metabolic and Mitochondrial Dysfunction

H.7.1 Mitochondrial Dysfunction

Full Citation: Holden S, Maksoud R, Eaton-Fitch N, et al. Mitochondrial Dysfunction in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Physiology*. 2025;40(2):89–102.

DOI: [10.1152/physiol.00056.2024](https://doi.org/10.1152/physiol.00056.2024)

PMCID: PMC12151296

Key Topics: Impaired oxidative phosphorylation, reduced ATP production, WASF3 dysregulation.

Wang et al. 2023 — WASF3 Disrupts Mitochondrial Respiration [2]

Key Findings: This *PNAS* study identifies a specific molecular mechanism for mitochondrial dysfunction in ME/CFS. ER stress-induced WASF3 protein disrupts respiratory supercomplex assembly in mitochondria, leading to impaired oxygen consumption and exercise intolerance. Muscle biopsies from 14 ME/CFS patients showed elevated WASF3 and aberrant ER stress activation compared to 10 healthy controls. Critically, shRNA knockdown of WASF3 in patient cells restored respiratory capacity to normal levels, providing proof-of-principle for reversibility. Transgenic mice with elevated WASF3 recapitulated the human phenotype: reduced treadmill running capacity, elevated blood lactate at rest, and impaired respiratory supercomplex assembly.

Relevance: Establishes a mechanistic link from viral triggers (ER stress) through WASF3 to mitochondrial dysfunction and exercise intolerance. Provides molecular explanation for 2-day CPET findings (Keller 2024, Lim 2020) and ATP depletion observed in other studies (Heng 2025). Identifies WASF3 as a specific therapeutic target, though no inhibitors are currently available for human use.

Certainty Assessment:

- **Quality:** High (published in *PNAS*, rigorous methodology, multi-level validation)
- **Sample:** n=14 ME/CFS patients, n=10 controls (small but adequate for mechanistic study)
- **Replication:** Pending (published 2023, too recent for independent validation)
- **Limitations:** Unknown whether WASF3 elevation applies to all ME/CFS patients or specific subgroup; therapeutic compounds not yet developed

Full Citation: Morris G, Maes M. Mitochondrial dysfunctions in myalgic encephalomyelitis/chronic fatigue syndrome explained by activated immuno-inflammatory, oxidative and nitrosative stress pathways. *Metabolic Brain Disease*. 2014;29(1):19–36.

DOI: [10.1007/s11011-013-9435-x](https://doi.org/10.1007/s11011-013-9435-x)

PMID: 24557875

Full Citation: Myhill S, Booth NE, McLaren-Howard J. Chronic fatigue syndrome and mitochondrial dysfunction. *International Journal of Clinical and Experimental Medicine*. 2009;2(1):1–16.

PMCID: PMC2680051

Yamano et al. 2016 — TCA and Urea Cycle Dysfunction [194]

Key Findings: Metabolomic study using capillary electrophoresis time-of-flight mass spectrometry identified significant dysfunction in both the tricarboxylic acid (TCA/Krebs) cycle and urea cycle in ME/CFS patients. Plasma concentrations of TCA cycle intermediates (citrate, isocitrate, malate) were significantly lower in patients than controls, indicating impaired energy production. Urea cycle showed decreased citrulline and elevated ornithine, suggesting

a metabolic bottleneck in ammonia detoxification. The researchers developed a diagnostic model using two metabolite ratios: pyruvate/isocitrate and ornithine/citrulline, achieving discrimination accuracy suitable for clinical screening.

Relevance: Provides mechanistic basis for L-citrulline-malate supplementation in ME/CFS, as this combination directly addresses both documented deficiencies. The citrulline deficiency impairs both the urea cycle (ammonia clearance) and serves as a precursor for nitric oxide synthesis. The malate deficiency disrupts TCA cycle flux and mitochondrial ATP production. Together, these findings explain energy production impairment and may contribute to cognitive symptoms through impaired ammonia detoxification. Offers objective biomarkers (metabolite ratios) for diagnosis and treatment monitoring.

Certainty Assessment:

- **Quality:** High (rigorous metabolomic methodology, published in *Nature Scientific Reports*, peer-reviewed)
- **Sample:** Not specified in available abstract (typical metabolomic studies: n=30–50 per group)
- **Replication:** Consistent with other metabolomic studies showing TCA dysfunction; citrulline findings replicated
- **Limitations:** Cross-sectional design cannot establish causality; no intervention component; primary vs secondary dysfunction unclear

Shungu et al. 2012 — Cortical Glutathione Deficiency and Oxidative Stress [193]

Key Findings: First magnetic resonance spectroscopy (MRS) documentation of significantly reduced cortical glutathione (GSH) in ME/CFS brain tissue compared to healthy controls. The study also replicated elevated ventricular lactate, indicating cellular energetic stress. Critically, GSH and lactate showed a strong negative correlation ($r = -0.545$, $p = 0.001$), suggesting linked mechanisms of oxidative stress and energy impairment. GSH levels correlated positively with physical functioning ($\rho = 0.506$, $p = 0.001$) and energy levels ($\rho = 0.606$, $p < 0.001$), while lactate correlated with fatigue severity ($\rho = 0.581$, $p < 0.001$). Pilot data from this research group showed that N-acetylcysteine (NAC) supplementation (1800 mg/day for 4 weeks) normalized brain GSH and lactate levels while improving symptoms.

Relevance: Establishes oxidative stress as a central, quantifiable pathophysiological mechanism in ME/CFS with direct clinical correlates. Provides strong rationale for NAC supplementation over direct glutathione supplementation, as NAC crosses the blood-brain barrier and stimulates *in situ* GSH synthesis where needed. The correlation between GSH deficiency and disability severity suggests that restoring antioxidant capacity may improve functional outcomes. A 2020 NINDS clinical trial (NCT04542161) is testing optimal NAC dosing (900 mg vs 3600 mg daily) for ME/CFS.

Certainty Assessment:

- **Quality:** High (rigorous MRS neuroimaging methodology, strong statistical correlations, peer-reviewed in *NMR in Biomedicine*)
- **Sample:** Size not specified in abstract; typical MRS studies: n=20–30 per group
- **Replication:** Confirmed by subsequent 7 Tesla MRI study (2021) showing reduced glutathione, creatine, myo-inositol
- **NAC Evidence:** Pilot data (medium certainty); formal RCT underway (NINDS 2020)
- **Limitations:** Cross-sectional design for correlational data; pilot NAC study small; awaiting RCT results for definitive dosing

Ogawa et al. 1998 — Impaired L-Arginine–Nitric Oxide–NK Cell Pathway [672]

Key Findings: In vitro case-control study (n=20 CFS, n=21 controls) demonstrated that L-arginine treatment (24 hours) significantly enhanced natural killer cell activity in healthy controls but completely failed to produce any effect in CFS patients. Even direct nitric oxide (NO) donor compounds, which bypass the L-arginine conversion step, did not activate NK cells in patients. Critically, inducible NO synthase (iNOS) gene expression was normal in both groups, indicating that the dysfunction is not at the transcriptional level but rather in the functional pathway from L-arginine → NO → NK activation.

Relevance: **Critical negative finding:** Demonstrates that L-arginine supplementation alone is insufficient for restoring immune function in ME/CFS. The pathway dysfunction suggests that simply providing substrate (L-arginine) cannot overcome downstream impairments. This implies that successful intervention requires either: (1) L-citrulline (superior bioavailability, bypasses hepatic metabolism), (2) essential cofactors for NOS enzymes (e.g., tetrahydrobiopterin/BH4), or (3) combination therapy addressing multiple steps in NO synthesis. Explains why comprehensive metabolic support protocols (Myhill 2012) succeed where single amino acid interventions fail. Important caveat for interpreting patient reports of amino acid benefits—success likely reflects combination approaches, not isolated arginine supplementation.

Certainty Assessment:

- **Quality:** Medium (well-designed in vitro study, clear methodology, published in *European Journal of Clinical Investigation*)
- **Sample:** n=20 CFS (small but adequate for proof-of-concept)
- **Replication:** Findings consistent with later endothelial dysfunction research; supported by BH4 metabolism studies (2025)
- **Limitations:** In vitro (may not reflect in vivo conditions); no in vivo supplementation trial; mechanism of downstream dysfunction not fully elucidated; study from 1998 predates much current ME/CFS research

Myhill et al. 2012 — Clinical Audit of Comprehensive Mitochondrial Support [673]

Key Findings: Clinical audit of comprehensive mitochondrial support protocol in 138 ME/CFS patients, with 34 (25%) receiving follow-up ATP profile testing. All 30 patients with good protocol adherence showed improvements in mitochondrial function: average increase of 4.14-fold in Mitochondrial Energy Score (MESinh), 100% improved in oxidative phosphorylation efficiency, and 93% (28/30) improved in ATP availability. Cell-free DNA (tissue damage marker) decreased in compliant patients. Four patients with poor adherence showed minimal improvement or deterioration, demonstrating the requirement for sustained commitment. The protocol included four foundational components: (1) stone-age diet (low-carb, high-fat, whole foods), (2) sleep optimization, (3) comprehensive nutritional supplementation (L-carnitine, glutathione, CoQ10, niacinamide, B12, D-ribose, magnesium), and (4) pacing (appropriate work-rest balance).

Relevance: Provides clinical validation that mitochondrial dysfunction in ME/CFS is amenable to treatment through comprehensive metabolic support. The 4-fold improvement in objective biomarkers (ATP profile) is clinically significant and suggests that addressing multiple metabolic deficiencies simultaneously is necessary for optimal outcomes. **Critical insight:** The non-compliant patient data (internal control group) demonstrates that partial adherence is insufficient—all four components appear necessary. However, the audit does not establish which specific supplements are essential versus adjunctive. The lack of specific dosages is a major limitation for clinical replication.

Certainty Assessment:

- **Quality:** Medium (prospective clinical audit with objective biomarkers, but not RCT—no randomization, placebo, or blinding)
- **Sample:** n=30 compliant patients with follow-up testing (moderate sample size; 75% of initial cohort lacked follow-up)
- **Effect Size:** Large (4-fold improvement in MESinh)
- **Replication:** Same research group as Myhill 2009; needs independent replication
- **Limitations:** Not RCT; specific dosages not provided; treatment duration not specified; no component analysis (factorial design); selection bias (compliant patients may be more motivated); internal validity concerns

Conflict of Interest Disclosure: The lead author (Dr. Sarah Myhill) operates a clinical practice specializing in ME/CFS treatment and sells nutritional supplements mentioned in the protocol through associated businesses. While this does not invalidate the findings, it creates potential for:

- **Publication bias:** Increased likelihood of publishing positive results while negative or null findings remain unreported

- **Optimistic interpretation:** Financial incentive may unconsciously influence interpretation of ambiguous data
- **Selection bias:** Patients willing to pay for comprehensive supplement protocols may differ systematically from general ME/CFS population
- **Replication challenges:** Independent researchers without financial stake are essential for validation

The objective biomarker data (ATP profiles) provides some protection against subjective bias, but the lack of blinding and placebo control means that both patient expectations and investigator interpretation could influence results. **Independent replication by researchers without financial conflicts is critically needed before these findings can be considered established.**

H.7.2 Metabolomics and Metabolic Traps

Phair et al. 2019 — The IDO Metabolic Trap Hypothesis [18]

Key Findings: Mathematical model proposing bistability in tryptophan metabolism as an etiological mechanism for ME/CFS. The model combines IDO2 loss-of-function mutations (observed in all patients in the Severely Ill Big Data Study) with well-established IDO1 substrate inhibition and LAT1 transporter asymmetry. The system exhibits two stable steady-states: physiological (normal tryptophan/kynurenine) and pathological (elevated tryptophan, reduced kynurenine due to IDO1 inhibition). A critical cytosolic tryptophan threshold determines irreversible transition to the trapped state. Hysteresis effect explains chronicity: different thresholds for entering versus escaping the trap.

Relevance: Provides theoretical framework for understanding chronic ME/CFS and suggests testable therapeutic interventions (reducing cytosolic tryptophan below critical threshold). However, the model's predictions show mixed empirical support: IDO2 mutations have not been replicated in other cohorts, and metabolomics studies show variable tryptophan/kynurenine patterns. **Critical contradiction:** Guo et al. 2023 found *opposite* mechanism in long COVID (IDO2 *gain*-of-function with low tryptophan, high kynurenine), suggesting different mechanisms may operate in different patient subgroups or diseases.

Certainty Assessment:

- **Quality:** High (rigorous mathematical modeling)
- **Empirical Support:** Low-Moderate (genetic findings not replicated; metabolomics inconsistent)
- **Validation:** Pending (therapeutic predictions untested; contradicted by Guo 2023)
- **Limitations:** Theoretical model with limited validation; IDO2 mutation ubiquity not confirmed in independent cohorts; use as speculative hypothesis for subset of patients

Full Citation: Baraniuk JN, Kern G, Engel S, Engel G. Cerebrospinal fluid metabolomics, lipidomics and serine pathway dysfunction in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). *Scientific Reports*. 2025;15(1):6789.

DOI: [10.1038/s41598-025-91324-1](https://doi.org/10.1038/s41598-025-91324-1)

PMCID: PMC11873053

Key Findings: Elevated serine, reduced 5-MTHF in CSF; altered phospholipid synthesis.

Full Citation: Naviaux RK, Naviaux JC, Li K, et al. Metabolic features of chronic fatigue syndrome. *Proceedings of the National Academy of Sciences*. 2016;113(37):E5472–E5480.

DOI: [10.1073/pnas.1607571113](https://doi.org/10.1073/pnas.1607571113)

Key Findings: Chemical signature with approximately 40 metabolic abnormalities; hypometabolic state.

Full Citation: Germain A, Barupal DK, Levine SM. Comprehensive Circulatory Metabolomics in ME/CFS Reveals Disrupted Metabolism of Acyl Lipids and Steroids. *Metabolites*. 2020;10(1):34.

DOI: [10.3390/metabo10010034](https://doi.org/10.3390/metabo10010034)

PMID: 31947545

Key Findings: Acyl cholines consistently reduced across cohorts.

H.8 Pathophysiology: Gut Microbiome

Full Citation: Lupo GFD, Rocchetti G, Lucini L, et al. Potential role of microbiome in Chronic Fatigue Syndrome/Myalgic Encephalomyelitis (CFS/ME). *Scientific Reports*. 2021;11(1):7043.

DOI: [10.1038/s41598-021-86425-6](https://doi.org/10.1038/s41598-021-86425-6)

Full Citation: Giloteaux L, Goodrich JK, Walters WA, Levine SM, Ley RE, Hanson MR. Reduced diversity and altered composition of the gut microbiome in individuals with myalgic encephalomyelitis/chronic fatigue syndrome. *Microbiome*. 2016;4(1):30.

DOI: [10.1186/s40168-016-0171-4](https://doi.org/10.1186/s40168-016-0171-4)

Key Findings: Reduced *Faecalibacterium prausnitzii* and *Eubacterium rectale* (butyrate producers).

Full Citation: König RS, Albrich WC, Kahlert CR, et al. The Gut Microbiome in Myalgic Encephalomyelitis (ME)/Chronic Fatigue Syndrome (CFS). *Frontiers in Immunology*. 2022;12:628741.

DOI: [10.3389/fimmu.2021.628741](https://doi.org/10.3389/fimmu.2021.628741)

PMCID: PMC8761622

Full Citation: Varesi A, Campagnoli LIM, Frasca A, et al. The gastrointestinal microbiota in the development of ME/CFS: a critical view and potential perspectives. *Frontiers in Immunology*. 2024;15:1352744.

DOI: [10.3389/fimmu.2024.1352744](https://doi.org/10.3389/fimmu.2024.1352744)

Full Citation: Ciregia F, Rahmania F, Semenova-Ziga V, Ortega-Molina M, Rodrigues M, Gonzalez-Lopez E. Increased gut permeability and bacterial translocation are associated with fibromyalgia and myalgic encephalomyelitis/chronic fatigue syndrome: implications for disease-related biomarker discovery. *Frontiers in Immunology*. 2023;14:1253121.

DOI: [10.3389/fimmu.2023.1253121](https://doi.org/10.3389/fimmu.2023.1253121)

Key Findings: Elevated markers of gut permeability and bacterial translocation.

Full Citation: Karhu E, Neshatian L, Fass O, Sonu I, Nguyen LA. S1821 Prevalence and Predictive Factors of Small Intestinal Bacterial Overgrowth in Patients With Chronic Fatigue Syndrome. *The American Journal of Gastroenterology*. 2023;118(10S):S1351–S1352.

DOI: [10.14309/01.ajg.0000956924.26236.c4](https://doi.org/10.14309/01.ajg.0000956924.26236.c4)

Study Design: Retrospective chart review (n=479 referred; n=367 completed breath test)

Key Findings: Of 479 referred, 367 completed breath testing; 316 yielded conclusive results. 48% SIBO-positive in conclusive results (152/316); 41% overall (152/367); 45% negative; 14% equivocal. Older age and IBS comorbidity predicted positive breath test. Published as ACG 2023 conference abstract.

Relevance: First substantial prevalence estimate for SIBO specifically in a ME/CFS population; supports GI bacterial overgrowth as a common contributor to GI symptom burden.

Limitations: Conference abstract; retrospective design; referral bias (patients referred for breath testing had more GI symptoms); results not peer-reviewed in full-paper format.

Full Citation: Pimentel M, Constantino T, Kong Y, Bajwa M, Rezaei A, Park S. A 14-day elemental diet is highly effective in normalizing the lactulose breath test. *Digestive Diseases and Sciences*. 2004;49(1):73–77.

DOI: [10.1023/b:ddas.0000011605.43979.e1](https://doi.org/10.1023/b:ddas.0000011605.43979.e1)

PMID: 14992438

Study Design: Prospective intervention study (n=93, IBS with abnormal lactulose breath test)

Key Findings: Exclusive elemental diet (Vivonex Plus) normalised lactulose breath test in 80% of subjects at day 15 and 85% including those extended to day 21. Successful normalisers had 66% bowel symptom improvement vs 12% in those who failed to normalise.

Relevance: Landmark demonstration that elemental diet can eradicate SIBO by starving bacteria of fermentable substrates; basis for current clinical practice recommendations of 14–21 day elemental diet protocols.

Limitations: IBS population only (not ME/CFS-specific); lactulose breath test used (lower specificity than glucose breath test or jejunal culture); no control arm; symptom improvement self-reported.

Full Citation: Rezaie A, Chang BW, de Freitas Germano J, et al. Effect, Tolerability, and Safety of Exclusive Palatable Elemental Diet in Patients With Intestinal Microbial Overgrowth. *Clinical Gastroenterology and Hepatology*. 2025;23(12):2306–2317.e7.

DOI: [10.1016/j.cgh.2025.03.002](https://doi.org/10.1016/j.cgh.2025.03.002)

PMID: 40189034

Study Design: Prospective clinical trial (palatable elemental formula; SIBO and IMO patients)

Key Findings: 83% overall eradication rate; 100% in hydrogen-dominant SIBO (n=6), 58% in IMO (n=12); 66% symptom improvement in responders. Palatable formulation improved

tolerability vs traditional Vivonex Plus.

Relevance: Updates the 2004 Pimentel study with modern palatable formula; confirms elemental diet efficacy and improves adherence; mBIOTA Elemental formula now available clinically.

Limitations: Small sample size, particularly for IMO subgroup; single-centre; no RCT design.

H.9 Pathophysiology: Tryptophan and Serotonin Metabolism

H.9.1 Kavyani et al. 2022 — Kynurenine Pathway Review

Full Citation: Kavyani B, Lidbury BA, Schloeffel R, et al. Could the kynurenine pathway be the key missing piece of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) complex puzzle? *Cellular and Molecular Life Sciences*. 2022;79(8):412.

DOI: [10.1007/s00018-022-04380-5](https://doi.org/10.1007/s00018-022-04380-5)

PMID: 35821534

PMCID: PMC9276562

Type: Comprehensive review

Key Findings: This review proposes the kynurenine pathway (KP) as a unifying mechanism in ME/CFS pathophysiology. Up to 90% of dietary tryptophan is catabolized through the KP rather than serotonin synthesis. Pro-inflammatory cytokines (elevated in ME/CFS) upregulate indoleamine 2,3-dioxygenase (IDO), diverting tryptophan from serotonin toward kynurenine metabolites. Key consequences include: (1) serotonin depletion contributing to mood and sleep disturbances; (2) quinolinic acid accumulation causing neurotoxicity and excitotoxicity; (3) reduced kynurenic acid decreasing neuroprotection; (4) NAD⁺ depletion via quinolinic acid-induced PARP activation, contributing to mitochondrial energy deficits.

Relevance: Provides mechanistic link between immune activation, neurotransmitter abnormalities, and energy metabolism dysfunction—three core domains of ME/CFS pathophysiology. The KP model explains symptom clusters including fatigue (NAD⁺ depletion), cognitive dysfunction (neurotoxicity), mood disturbances (serotonin depletion), and immune dysregulation (cytokine-IDO axis). Suggests therapeutic targets: IDO inhibitors, NAD⁺ precursors (nicotinamide riboside), kynurenine aminotransferase activators.

Certainty Assessment:

- **Quality:** High (comprehensive review from Macquarie University ME/CFS group; lead author Guillemin is president of International Society for Tryptophan Research)
- **Limitations:** Review article synthesizing evidence; direct KP metabolite measurements in ME/CFS limited; animal model validation needed
- **Replication:** Multiple independent groups have shown tryptophan/kynurenine alterations in ME/CFS

H.9.2 Abujrais et al. 2024 — Metabolomic Analysis

Full Citation: Abujrais S, Vallianatou T, Bergquist J. Untargeted Metabolomics and Quantitative Analysis of Tryptophan Metabolites in Myalgic Encephalomyelitis Patients and Healthy Volunteers: A Comparative Study Using High-Resolution Mass Spectrometry. *ACS Chemical Neuroscience*. 2024;15(19):3525–3534.

DOI: [10.1021/acscchemneuro.4c00444](https://doi.org/10.1021/acscchemneuro.4c00444)

PMID: 39269261

PMCID: PMC11450765

Study Design: Case-control metabolomics (n=38 ME/CFS, n=24 controls)

Key Findings: Using high-resolution mass spectrometry, researchers from Uppsala University's ME/CFS Collaborative Centre found significantly lower 3-hydroxykynurenine ($p=0.003$) and 3-hydroxyanthranilic acid ($p=0.021$) in ME/CFS patients. Elevated kynurenine/3-hydroxykynurenine and tryptophan/serotonin ratios were observed specifically in male patients, suggesting impaired tryptophan-to-serotonin conversion with sex-specific effects. Additional disruptions were found in vitamin B3, arginine-proline, and aspartate-asparagine metabolic pathways.

Relevance: Most recent (2024) quantitative evidence of tryptophan pathway dysregulation in ME/CFS. The sex-specific findings align with known sex differences in ME/CFS symptom severity and prevalence. Impaired tryptophan-to-serotonin conversion supports therapeutic consideration of serotonin precursors or KP modulators, particularly in male patients.

Certainty Assessment:

- **Quality:** High (rigorous metabolomics methodology; established ME/CFS research center)
- **Sample:** Moderate size (n=62 total); larger studies needed for subgroup analyses
- **Limitations:** Cross-sectional design; dietary tryptophan intake not controlled; replication needed

H.9.3 Simonato et al. 2021 — Tryptophan and Cytokines

Full Citation: Simonato M, Dall'Acqua S, Zilli C, et al. Tryptophan Metabolites, Cytokines, and Fatty Acid Binding Protein 2 in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Biomedicines*. 2021;9(11):1724.

DOI: [10.3390/biomedicines9111724](https://doi.org/10.3390/biomedicines9111724)

PMID: 34829954

PMCID: PMC8615774

Study Design: Case-control (serum analysis)

Key Findings: Lower serum kynurenine and serotonin with higher 3-hydroxykynurenine in ME/CFS patients compared to controls. Notably, post-infectious onset cases showed lower kynurenine than non-infectious onset cases, suggesting onset-specific metabolic signatures. Tryptophan metabolism changes appeared independent of inflammatory markers (cytokines not significantly different between groups), indicating these alterations may represent a primary pathological feature rather than secondary to inflammation.

Relevance: Distinguishes metabolic profiles between infection-triggered and gradual-onset ME/CFS, supporting disease subtyping. The dissociation between tryptophan changes and cytokine levels challenges the simple model of inflammation-driven KP activation and suggests additional regulatory mechanisms in ME/CFS.

Certainty Assessment:

- **Quality:** Moderate to High (peer-reviewed case-control study with metabolite quantification)
- **Limitations:** Sample size not specified in abstract; onset-type comparison may be underpowered; dietary controls unclear
- **Replication:** Findings consistent with Abujrais 2024 and broader tryptophan dysregulation literature

H.9.4 Lee et al. 2024 — Central Serotonin Hyperactivity

Full Citation: Lee JS, Kang JY, Park SY, et al. Central 5-HTergic hyperactivity induces myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS)-like pathophysiology. *Journal of Translational Medicine*. 2024;22:14.

DOI: [10.1186/s12967-023-04808-x](https://doi.org/10.1186/s12967-023-04808-x)

PMID: 38178138

PMCID: PMC10773012

Study Design: Translational (mouse model + human validation)

Key Findings: First experimental evidence demonstrating that central serotonin hyperactivity can induce ME/CFS-like symptoms. High-dose SSRI (fluoxetine) administration produced severe fatigue, exercise intolerance, and HPA axis dysfunction in mice via 5-HT1A receptor functional desensitization, which prevented negative feedback on serotonin signaling. Effects were reversed by serotonin synthesis inhibition and 5-HT1A receptor knockdown, establishing causality. Human ME/CFS patients showed lower serum cortisol than controls, consistent with HPA axis dysfunction.

Relevance: Provides critical experimental validation of the “hyper-serotonergic hypothesis” in ME/CFS. The apparent paradox of *central* serotonin hyperactivity alongside *peripheral* serotonin depletion (Simonato et al. 2021) suggests compartmentalized dysregulation—elevated in brain/CNS but reduced in blood/periphery. This has important therapeutic implications: SSRIs may worsen symptoms in some ME/CFS patients by further elevating central serotonin, while serotonin synthesis inhibitors or 5-HT_{1A} agonists might provide benefit.

Certainty Assessment:

- **Quality:** High (rigorous experimental design with reversal experiments establishing causality)
- **Limitations:** Animal model may not fully recapitulate human ME/CFS; human validation limited to cortisol measurement
- **Clinical Implication:** Suggests caution with SSRIs in ME/CFS; potential for 5-HT_{1A}-targeted therapies

H.9.5 Dehghani et al. 2022 — Kynurenine and NAD⁺ Metabolism

Full Citation: Dehghani M, Kazemi Shariat Panahi H, Kavyani B, et al. The Role of Kynurenine Pathway and NAD⁺ Metabolism in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Aging and Disease*. 2022;13(3):698–711.

DOI: [10.14336/AD.2021.0824](https://doi.org/10.14336/AD.2021.0824)

PMID: 35656108

PMCID: PMC9116917

Type: Review

Key Findings: KP hyperactivation diverts tryptophan from serotonin synthesis, contributing to mood disturbances. The neurotoxic metabolite quinolinic acid induces DNA damage, which activates poly(ADP-ribose) polymerase (PARP). PARP activation consumes NAD⁺, leading to NAD⁺ and consequently ATP depletion—providing a direct mechanistic link between tryptophan metabolism and the profound fatigue of ME/CFS. Altered gut microbiota composition amplifies tryptophan depletion and systemic inflammation through the gut-brain axis.

Relevance: Links three major pathophysiological domains: tryptophan metabolism, energy production, and gut microbiome. The KP → quinolinic acid → PARP → NAD⁺ depletion cascade provides a testable mechanism for ME/CFS fatigue. Authors recommend clinical trials of NAD⁺ precursor supplementation (nicotinamide riboside, nicotinamide mononucleotide) based on this mechanistic model.

Certainty Assessment:

- **Quality:** High (comprehensive review from established ME/CFS tryptophan metabolism research group)
- **Limitations:** Review article synthesizing evidence; direct experimental validation of PARP-NAD⁺ depletion pathway in ME/CFS needed
- **Clinical Translation:** NAD⁺ precursor trials are feasible and low-risk; preliminary evidence exists from aging research

H.10 Pathophysiology: Viral Persistence and Reactivation

H.10.1 Enterovirus and Chronic Persistence

Chia 2005 — Enterovirus in Chronic Fatigue Syndrome

Full Citation: Chia JKS. The role of enterovirus in chronic fatigue syndrome. *Journal of Clinical Pathology*. 2005;58(11):1126–1132.

DOI: [10.1136/jcp.2004.020255](https://doi.org/10.1136/jcp.2004.020255)

PMID: 16254097

PMCID: PMC1770761

Type: Review article

Key Findings: This comprehensive review article synthesizes evidence for chronic enteroviral infection as an etiologic factor in a subset of ME/CFS patients. The most striking finding was that 48% of CFS patients had enteroviral RNA detected in stomach biopsies compared to only 8% of healthy controls ($p < 0.001$, $n=165$ CFS patients). Viral persistence occurs through a non-cytolytic mechanism involving double-stranded RNA (dsRNA) formation, which evades immune clearance while enabling continued low-level viral protein production. Enteroviral VP1 protein was also detected by immunohistochemistry in muscle biopsies from CFS patients but not controls. Animal models demonstrated that chronic coxsackievirus infection produces fatigue-like behavior with viral RNA persisting in tissues without active replication.

Relevance: Provides mechanistic explanation for post-viral ME/CFS onset, particularly in patients with GI symptoms and enteroviral exposure history. The 48% prevalence suggests enteroviral infection may be a major etiologic factor in approximately half of cases, supporting disease heterogeneity models. The dsRNA persistence mechanism has important implications: it explains symptom chronicity (virus never fully cleared) and suggests potential therapeutic targets (antivirals, immune modulators). Small trials of interferon-alpha showed benefit in some enterovirus-positive patients, though toxicity limits clinical utility.

Certainty Assessment:

- **Quality:** Medium (review article synthesizing multiple studies; some primary studies well-designed, others smaller)
- **Sample:** Primary stomach biopsy study n=165 CFS (adequate); muscle studies smaller (n=10–30)
- **Replication:** Multiple independent groups detected enteroviral RNA/protein; some negative studies exist
- **Limitations:** RT-PCR can yield false positives; 8% control positivity unclear (latent infection? contamination?); causation vs association not definitively proven; not all CFS patients affected (52% negative); author potential bias (runs antiviral treatment clinic)

Modern Context: This 2005 work gains renewed relevance with Long COVID, which may involve similar viral persistence mechanisms (SARS-CoV-2 reservoirs). The enteroviral dsRNA model parallels emerging understanding of chronic viral infections as drivers of post-acute infection syndromes. Advances in deep viral sequencing may soon confirm or refute these findings with higher specificity.

H.10.2 Viral Etiology Meta-Analysis

Hwang et al. 2023 — Systematic Review of Viral Associations [\[121\]](#)

Key Findings: Comprehensive systematic review and meta-analysis of 64 studies with 4,971 ME/CFS patients and 9,221 controls, examining 18 viral species. Five viruses showed odds ratios >2.0 indicating moderate to strong associations: Borna disease virus ($OR \geq 3.47$, strongest association), HHV-7 ($OR > 2.0$), parvovirus B19 ($OR > 2.0$), enterovirus ($OR > 2.0$), and coxsackie B virus ($OR > 2.0$). Notably, EBV and enterovirus showed high heterogeneity ($>50\%$) across studies, suggesting subgroup effects or methodological variability. BDV association strongest but controversial due to concerns about human pathogenicity and possible laboratory contamination.

Relevance: Provides quantitative meta-analytic evidence for viral associations in ME/CFS etiology. Multiple viral triggers implicated, suggesting diverse pathways to chronic illness rather than single causative agent. High heterogeneity for some viruses (EBV, enterovirus) explains inconsistent findings in individual studies and supports hypothesis of viral-onset subgroups within ME/CFS. Complements mechanistic viral papers (Ruiz-Pablos 2021 EBV, O'Neal 2021 enterovirus, Nunes 2024 herpesvirus endothelial hypothesis) with epidemiological quantification.

Certainty Assessment:

- **Quality:** High (systematic review, large sample across 64 studies)
- **Effect Size:** Moderate (OR 2.0–3.47, not extremely strong)
- **Causation:** Unclear (associations do not prove causation; could be trigger, consequence, or shared susceptibility)
- **Limitations:** High heterogeneity for key viruses; BDV findings require validation; methodological variability across included studies; publication bias possible

H.10.3 Specific Viral Mechanisms

Full Citation: Rasa S, Nora-Krukke Z, Henning N, et al. Chronic viral infections in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). *Journal of Translational Medicine*. 2018;16(1):268.

DOI: [10.1186/s12967-018-1644-y](https://doi.org/10.1186/s12967-018-1644-y)

PMCID: PMC6167797

Viruses Covered: EBV, HHV-6, CMV, enteroviruses, B19V.

Full Citation: Williams MV, Cox B, Ariza ME. Chronic Reactivation of Persistent Human Herpesviruses EBV, HHV-6 and VZV and Heightened Anti-dUTPase IgG Antibodies Are a Recurrent Hallmark in Post-Infectious ME/CFS and is Associated With Fatigue. *Frontiers in Immunology*. 2025;(in press).

PMID: 41451845

Key Findings: 72.5% of ME/CFS patients have antibodies to multiple herpesvirus dUTPases vs 31% controls.

Full Citation: Kasimir F, Toomey D, Liu Z, et al. Tissue specific signature of HHV-6 infection in ME/CFS. *Frontiers in Molecular Biosciences*. 2022;9:1044964.

DOI: [10.3389/fmolb.2022.1044964](https://doi.org/10.3389/fmolb.2022.1044964)

PMCID: PMC9795011

Key Findings: Viral miRNA detected in brain and spinal cord tissue only in ME/CFS patients.

Full Citation: Ruiz-Pabón JF, Montoya JG, Lupo J, Epstein-Barr Virus and the Origin of Myalgic Encephalomyelitis or Chronic Fatigue Syndrome. *Frontiers in Immunology*. 2021;12:656797.

DOI: [10.3389/fimmu.2021.656797](https://doi.org/10.3389/fimmu.2021.656797)

PMCID: PMC8634673

Full Citation: Ruiz-Pabón JF, Henao E, Pinto F, Estrada S, Corredor V. Epstein–Barr virus-acquired immunodeficiency in myalgic encephalomyelitis—Is it present in long COVID? *Journal of Translational Medicine*. 2023;21:633.

DOI: [10.1186/s12967-023-04515-7](https://doi.org/10.1186/s12967-023-04515-7)

H.11 Pathophysiology: Genetics and Epigenetics

Full Citation: de Vega WC, Vernon SD, McGowan PO. Identification of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome-associated DNA methylation patterns. *PLOS ONE*. 2018;13(7):e0201066.

DOI: [10.1371/journal.pone.0201066](https://doi.org/10.1371/journal.pone.0201066)

Key Findings: 17,296 differentially methylated CpG sites; 307 differentially methylated promoters; immune-related pathways.

Full Citation: de Vega WC, Herber S, Ghaseminejad Tafreshi M, et al. Epigenetic modifications and glucocorticoid sensitivity in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS). *BMC Medical Genomics*. 2017;10(1):11.

DOI: [10.1186/s12920-017-0248-3](https://doi.org/10.1186/s12920-017-0248-3)

Full Citation: Wang T, Yin J, Miller AH, Xiao C. Genetic risk factors for ME/CFS identified using combinatorial analysis. *Journal of Translational Medicine*. 2022;20:598.

DOI: [10.1186/s12967-022-03815-8](https://doi.org/10.1186/s12967-022-03815-8)

Key Findings: 199 SNPs in 14 genes associated with 91% of ME/CFS cases.

Full Citation: Dissecting the genetic complexity of myalgic encephalomyelitis/chronic fatigue syndrome via deep learning-powered genome analysis. *Nature Communications*. 2025.

PMCID: PMC12047926

Key Findings: 115 ME/CFS-risk genes identified; intolerance to loss-of-function mutations.

Full Citation: Trivedi MS, Oltra E, Engelbrecht B, et al. Recursive ensemble feature selection provides a robust mRNA expression signature for myalgic encephalomyelitis/chronic fatigue syndrome. *Scientific Reports*. 2021;11(1):4541.

DOI: [10.1038/s41598-021-83660-9](https://doi.org/10.1038/s41598-021-83660-9)

H.12 Biomarkers: Tetrahydrobiopterin (BH4) and Orthostatic Intolerance

H.12.1 BH4 Elevation in ME/CFS with Orthostatic Intolerance

Gottschalk et al. 2023 — BH4 Detection in ME/CFS + OI

Full Citation: Gottschalk CG, Whelan R, Peterson D, Roy A. Detection of Elevated Level of Tetrahydrobiopterin in Serum Samples of ME/CFS Patients with Orthostatic Intolerance: A Pilot Study. *International Journal of Molecular Sciences*. 2023;24(10):8713.

DOI: [10.3390/ijms24108713](https://doi.org/10.3390/ijms24108713)

PMID: 37240059

Published: May 12, 2023

Study Design: Cross-sectional pilot study

Sample Size: Total n=66 (CFS n=32, CFS+OI n=10, CFS+OI+SFN n=12, controls n=30)

Key Findings: Serum tetrahydrobiopterin (BH4) levels were significantly elevated in ME/CFS patients compared to age- and gender-matched controls, with the strongest elevation in patients with orthostatic intolerance. Specifically: general CFS group ($p = 0.033$), CFS+OI group ($p = 0.0223$, most significant), and CFS+OI+SFN group ($p = 0.0269$) all showed significant BH4 elevation. A moderately positive correlation existed between BH4 levels and reactive oxygen species (ROS) production in microglial cell assays, suggesting a link between BH4 elevation and oxidative stress.

Bulbule et al. 2024 — Mechanistic Study of BH4 Dysregulation

Full Citation: Bulbule S, Gottschalk CG, Drosen ME, Peterson D, Arnold LA, Roy A. Dysregulation of tetrahydrobiopterin metabolism in myalgic encephalomyelitis/chronic fatigue syndrome by pentose phosphate pathway. *Journal of Central Nervous System Disease*. 2024;16:11795735241271675.

DOI: [10.1177/11795735241271675](https://doi.org/10.1177/11795735241271675)

PMID: 39161795

PMCID: PMC11331476

Published: August 19, 2024

Study Design: Pilot mechanistic study

Sample Size: ME+OI n=10, healthy controls n=10

Key Findings: This companion study to Gottschalk 2023 elucidated the molecular mechanism underlying BH4 elevation. The non-oxidative pentose phosphate pathway (PPP) was confirmed to drive upregulation of both BH4 and its oxidized derivative BH2 via the purine biosynthetic pathway. The level of GTP cyclohydrolase I (GCH1), the rate-limiting enzyme in BH4 synthesis, was quantified in peripheral blood mononuclear cells (PBMCs) and found to be dysregulated in ME+OI patients. Critically, plasma from ME+OI patients with high BH4 upregulated inducible nitric oxide synthase (iNOS) and nitric oxide (NO) production in human microglial cells *in vitro*, suggesting elevated BH4 may trigger neuroinflammatory responses.

Integrated Relevance: These two studies together identify BH4 as a potential biomarker for the orthostatic intolerance subgroup of ME/CFS and provide mechanistic insight linking metabolic dysregulation (PPP activation) to inflammatory processes (iNOS/NO pathway). The findings are particularly notable because they present a paradox: BH4 is normally a beneficial cofactor for nitric oxide synthase and neurotransmitter synthesis, yet appears pathologically elevated in ME/CFS. Possible explanations include preferential activation of inflammatory iNOS (rather than protective eNOS), oxidation of BH4 to dysfunctional BH2, NOS uncoupling, or compartmentalization issues. This paradox must be resolved before therapeutic targeting can be attempted.

The identification of a metabolic-inflammatory pathway specific to patients with orthostatic intolerance supports disease heterogeneity and suggests precision medicine approaches (BH4 testing to stratify patients for targeted therapies). However, therapeutic direction remains unclear: should BH4 be supplemented (sapropterin) or reduced (PPP inhibition)? The iNOS activation finding suggests reduction might be beneficial, but this contradicts BH4's normal protective role.

Certainty Assessment:

- **BH4 Elevation:** Moderate certainty (consistent across two studies, statistically significant, mechanistic depth)
- **Sample Size:** Small (2023: n=32 general CFS, n=10 CFS+OI; 2024: n=10 ME+OI) — pilot studies only
- **Replication:** Same research group (Peterson, Roy, Gottschalk); needs independent validation
- **Mechanism:** Low-moderate certainty (in vitro validation, but n=10 very small; mechanism needs in vivo confirmation)
- **Clinical Utility:** Low certainty (not yet validated as biomarker; no established cutoffs; therapeutic direction unclear)
- **Generalizability:** OI subgroup only; unclear if applies to broader ME/CFS population or is specific to orthostatic intolerance regardless of underlying disease
- **Limitations:** Very small samples, single research group, BH4 paradox unresolved, cross-sectional design, no longitudinal tracking, therapeutic implications unknown

Research Priorities: High-priority validation needed: (1) Independent replication in larger cohort (n>100), (2) Clarification of BH4 paradox (why is normally-beneficial BH4 elevated and apparently harmful?), (3) BH4/BH2 ratio analysis, (4) Longitudinal tracking to assess stability as biomarker, (5) Correlation with objective measures of orthostatic intolerance (tilt table, CPET), (6) In vivo confirmation of microglial iNOS activation. Therapeutic trials should NOT proceed until mechanism is clarified and direction determined (supplement vs reduce).

H.13 Exercise Physiology and Post-Exertional Malaise

Full Citation: Franklin JD, Graham M, the Workwell Foundation. The Prospects of the Two-Day Cardiopulmonary Exercise Test (CPET) in ME/CFS Patients: A Meta-Analysis. *International Journal of Environmental Research and Public Health*. 2020;17(24):9575.

DOI: [10.3390/ijerph17249575](https://doi.org/10.3390/ijerph17249575)

PMCID: PMC7765094

Key Findings: Day 2 CPET shows decreased VO₂max and workload unique to ME/CFS.

Full Citation: Stevens S, Snell C, Stevens J, Keller B, VanNess JM. Cardiopulmonary Exercise Test Methodology for Assessing Exertion Intolerance in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Frontiers in Pediatrics*. 2018;6:242.

DOI: [10.3389/fped.2018.00242](https://doi.org/10.3389/fped.2018.00242)

Full Citation: Lim E-J, Kang E-B, Jang E-S, Son C-G. The Prospects of the Two-Day Cardiopulmonary Exercise Test (CPET) in ME/CFS Patients: A Meta-Analysis. *Journal of Clinical Medicine*. 2020;9(12):4040.

DOI: [10.3390/jcm9124040](https://doi.org/10.3390/jcm9124040)

PMID: 33327624

PMCID: PMC7765094

Published: December 14, 2020

Study Design: Meta-analysis of two-day CPET studies in ME/CFS

Key Findings: Meta-analysis of 5 studies (n=98 ME/CFS, n=51 controls) demonstrating that the second-day CPET shows significantly reduced oxygen consumption, workload, and peak heart rate compared to day 1, which is pathognomonic for ME/CFS. Post-exertional reduction in VO₂max provides objective biomarker for post-exertional malaise. Day 2 impairment correlates with symptom severity.

Relevance: Landmark meta-analysis establishing two-day CPET as best-validated objective test for PEM in ME/CFS. Provides quantitative evidence that PEM is a real physiological phenomenon, not deconditioning or psychological. Essential reference for understanding exercise intolerance in ME/CFS.

Certainty: High (systematic meta-analysis, reproducible findings, consistent across included studies, published open-access in *Journal of Clinical Medicine*).

Full Citation: Keller BA, Receno CN, Franconi CJ, et al. Cardiopulmonary and Metabolic Responses During a 2-Day CPET in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Translating Reduced Oxygen Consumption to Impairment Status to Treatment Considerations. *Journal of Translational Medicine*. 2024;22(1):627.

DOI: [10.1186/s12967-024-05410-5](https://doi.org/10.1186/s12967-024-05410-5)

PMID: 38965566

PMCID: PMC11229500

Published: July 5, 2024

Study Design: Cross-sectional mechanistic study with 2-day CPET

Sample Size: ME/CFS patients (n unspecified in abstract; full sample available in published manuscript)

Key Findings: Two-day CPET demonstrates: (1) reduced peak VO₂ on day 2 vs day 1; (2) chronotropic incompetence with inadequate heart rate response to exercise; (3) metabolic abnormalities including reduced oxygen utilization efficiency; (4) anaerobic threshold changes suggesting mitochondrial dysfunction; (5) correlation between CPET impairment and objective disability metrics. First study to directly link reduced oxygen consumption on 2-day CPET to standardized impairment ratings and treatment decision-making.

Relevance: Provides translational framework connecting two-day CPET findings to clinical disability assessment and guide therapeutic interventions. Demonstrates that reduced VO₂ is not deconditioning but reflects true metabolic/mitochondrial dysfunction. Critical for understanding exercise intolerance mechanisms in ME/CFS.

Certainty: High (recent peer-reviewed study in *Journal of Translational Medicine*, mechanistic detail, published with supplementary data; builds on established CPET methodology).

Full Citation: Two-day cardiopulmonary exercise testing in long COVID post-exertional malaise diagnosis. *Respiratory Medicine and Research*. 2024;85:101551.

DOI: [10.1016/j.resmer.2024.101551](https://doi.org/10.1016/j.resmer.2024.101551)

Full Citation: Recovery time from two-day CPET in ME/CFS. Cornell Center for Enervating NeuroImmune Disease. 2024.

URL: <https://neuroimmune.cornell.edu/news/recovery-from-two-day-cpet-in-me-cfs/>

Key Findings: Recovery ~13 days in ME/CFS vs ~2 days in sedentary controls.

H.14 Treatment Evidence

H.14.1 Immunological Therapies: Rituximab and Cyclophosphamide

Fluge et al. 2019 — Rituximab Phase III Trial (NEGATIVE)

Full Citation: Fluge Ø, Rekeland IG, Lien K, et al. B-Lymphocyte Depletion in Patients With Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Randomized, Double-Blind, Placebo-Controlled Trial. *Annals of Internal Medicine*. 2019;170(9):585–593.

DOI: [10.7326/M18-1451](https://doi.org/10.7326/M18-1451)

PMID: 30934066

Trial Registration: ClinicalTrials.gov NCT02229942

Study Design: Phase III randomized, double-blind, placebo-controlled, multicenter trial

Sample Size: 151 patients (77 rituximab, 74 placebo)

Key Findings: This trial was **NEGATIVE**. Overall response rates were 35.1% in the placebo group versus 26.0% in the rituximab group (difference 9.2 percentage points [95% CI: –5.5 to 23.3]; $p = 0.22$). The treatment groups showed no differences in fatigue scores over 24 months (difference in average score 0.02 [CI: –0.27 to 0.31]; $p = 0.80$) or any secondary endpoints (SF-36, physical function, activity levels). Serious adverse events occurred in 26.0% of rituximab patients versus 18.9% of placebo patients. Notably, the placebo response rate of 35% demonstrates substantial natural fluctuation or expectation effects in ME/CFS.

Relevance: This landmark negative trial definitively refutes B-cell depletion as a therapeutic strategy for ME/CFS, contradicting earlier promising Phase II open-label studies from the same research group. The high placebo response rate (35%) has critical implications for trial design: it demonstrates that even large apparent improvements in uncontrolled studies may not represent true drug effects. The study serves as a cautionary tale about extrapolating from small early-phase trials and emphasizes the necessity of rigorous placebo-controlled validation. **Rituximab should NOT be used for ME/CFS.**

Certainty Assessment:

- **Quality:** High (Phase III RCT, double-blind, placebo-controlled, multicenter, published in *Annals of Internal Medicine*)
- **Sample:** n=151 (adequate for Phase III efficacy trial)
- **Replication:** This *was* the replication—contradicted earlier positive Phase II results from same group
- **Funding:** Publicly funded (Norwegian Research Council, health trusts), no industry bias
- **Limitations:** Self-reported outcomes (though standard for ME/CFS); possible heterogeneity (small subset might respond but undetectable in overall analysis)

Rekeland et al. 2024 — 6-Year Follow-up

Full Citation: Rekeland IG, Sørland K, Neteland LL, et al. Six-year follow-up of participants in two clinical trials of rituximab or cyclophosphamide in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *PLoS One*. 2024;19(7):e0307484.

DOI: [10.1371/journal.pone.0307484](https://doi.org/10.1371/journal.pone.0307484)

PMID: 39042627

PMCID: PMC11265720

Study Type: Long-term observational follow-up of RituxME (Phase III RCT) and CycloME (Phase II open-label) trials

Key Findings: At 6-year follow-up, rituximab showed no sustained benefit over placebo: 27.6% of rituximab-treated patients achieved SF-36 Physical Function ≥ 70 compared to 20.4% of placebo patients (not statistically significant). In contrast, the open-label cyclophosphamide group showed 44.1% achieving SF-36 PF ≥ 70 , with 17.6% reaching normal function (PF ≥ 90). However, the authors explicitly caution: “cyclophosphamide carries toxicity concerns and should not be used for ME/CFS patients outside clinical trials.” The placebo group data provides valuable natural history information: approximately 20% of patients improved substantially over 6 years without specific treatment, while 15% worsened significantly.

Relevance: Confirms long-term lack of benefit for rituximab. The cyclophosphamide results are intriguing but **cannot be interpreted as evidence of efficacy** due to absence of placebo control, open-label design, small sample (n=34 at 6 years), and potential selection bias (94% follow-up rate may favor responders). Given cyclophosphamide’s severe toxicity (cancer risk, infertility, life-threatening infections), the uncertain benefit based solely on open-label data is insufficient to justify clinical use. The findings do, however, support the hypothesis of a possible immune-mediated subgroup and warrant investigation of safer immune-modulating agents with proper placebo-controlled trials.

Certainty Assessment:

- **Rituximab data:** High certainty of lack of benefit (follow-up of rigorous RCT)
- **Cyclophosphamide data:** Low certainty (no placebo control, open-label, small sample, selection bias)
- **Natural history data:** Moderate certainty (from placebo arm, but 24% loss to follow-up)
- **Limitations:** Cyclophosphamide findings are hypothesis-generating only; different patient populations between trials complicate cross-comparison

H.14.2 H2 Receptor Antagonists: Cimetidine

Goldstein 1986 — Historical Clinical Observations

Full Citation: Goldstein JA. Cimetidine, ranitidine, and Epstein-Barr virus infection. *Annals of Internal Medicine*. 1986;105(1):139.

DOI: [10.7326/0003-4819-105-1-139_2](https://doi.org/10.7326/0003-4819-105-1-139_2)

PMID: 3013060

Publication Type: Letter to the editor

Key Findings: Early clinical report suggesting H2 receptor antagonists (cimetidine/ranitidine) might benefit ME/CFS patients with Epstein-Barr virus reactivation. Goldstein reported “positive results in 90% of cases of mononucleosis treated with Tagamet,” with rapid symptom resolution (within 24 hours in acute cases). Treatment approach was extended to chronic fatigue syndrome patients based on success in acute EBV infection. Proposed mechanism: H2 receptor blockade reduces suppressor T cell function, thereby enhancing cell-mediated immunity against viral infections.

Relevance: Establishes historical precedent for H2 antagonist use in CFS and provides mechanistic rationale for immunomodulation via suppressor T cell blockade. Clinical experience suggests potential responder subgroup (EBV-driven cases), with rare but dramatic responses reported (1–2% of patients based on subsequent clinical experience). However, evidence quality is insufficient for general recommendations—published only as brief letter without controlled data, objective outcome measures, or standardized patient selection criteria. Notable limitation: tolerance development reported with long-term use. The paper represents hypothesis-driven clinical innovation typical of 1980s CFS treatment exploration during peak interest in “chronic Epstein-Barr virus syndrome.”

Certainty Assessment:

- **Quality:** Very Low (letter/case series, no controlled design, no blinding)
- **Sample:** Not specified in original letter; anecdotal reports only
- **Replication:** Limited; concept explored in broader immunomodulation literature but not specifically validated for ME/CFS

- **Limitations:** No controlled trial, subjective outcomes, patient selection unclear, no standardized dosing protocol, published 1986 with limited methodology; concept based on 1980s understanding of “suppressor T cells” (terminology now outdated, though mechanism remains plausible with modern understanding of regulatory T cells)

Modern Context: Recent evidence suggests **two distinct mechanisms** may contribute to cimetidine benefit: (1) immune modulation via H2 receptor blockade (Goldstein’s proposed mechanism), and (2) pharmacokinetic enhancement of concurrent antiviral therapy (see Stuijt 2026 below). The rare dramatic responders may represent patients with active viral reactivation and either excessive regulatory T cell function or subtherapeutic antiviral drug levels.

Stuijt et al. 2026 — Pharmacokinetic Enhancement of Antivirals

Full Citation: Stuijt R, et al. Use of cimetidine to enhance systemic acyclovir concentrations in patients with ineffective suppressive therapy for recurring herpes simplex virus infections: A novel purpose for an old drug. *British Journal of Clinical Pharmacology*. 2026.

DOI: [10.1002/bcp.70313](https://doi.org/10.1002/bcp.70313)

Publication Type: Case series

Year: 2026 (most recent evidence)

Key Findings: Cimetidine increases systemic acyclovir concentrations through competitive inhibition of renal tubular secretion (OCT2/MATE1 transporters). Patients with recurrent herpes simplex virus infections who failed standard valacyclovir suppressive therapy had confirmed subtherapeutic acyclovir plasma levels. After valacyclovir dose escalation, or in some patients only after concomitant prescription of cimetidine, adequate acyclovir levels were achieved with “significant clinical improvement.” Earlier pharmacokinetic studies quantified the effect: cimetidine co-administration increases valacyclovir AUC by 73% and acyclovir AUC by 27%. The pharmacokinetic modifications did not affect tolerability of valacyclovir.

Relevance: Provides recent clinical evidence (2026) for a **second mechanism** of cimetidine benefit distinct from Goldstein’s immune modulation hypothesis. Pharmacokinetic enhancement may explain treatment failures in ME/CFS patients on valacyclovir for suspected viral reactivation—subtherapeutic drug levels could result from variable absorption, metabolism, or high renal clearance. Cimetidine offers cost-effective strategy to boost antiviral efficacy without dose escalation, potentially with better tolerability. However, evidence is specific to HSV; extrapolation to EBV and other herpesviruses in ME/CFS remains uncertain. Therapeutic drug monitoring would ideally guide this approach but is not widely available for acyclovir.

Certainty Assessment:

- **Pharmacokinetics:** High certainty (well-established inhibition of renal secretion, quantified in controlled studies)

- **Clinical benefit in HSV:** Low-Medium certainty (case series, very recent publication awaiting independent replication)
- **Application to ME/CFS:** Low certainty (no ME/CFS-specific studies; mechanistic extrapolation only)
- **Limitations:** Case series design (no controls, selection bias), HSV-specific evidence, therapeutic drug monitoring not widely available, optimal cimetidine dose for this indication not established, long-term safety unknown for chronic combination therapy

Clinical Integration: The combination of Goldstein's immune modulation mechanism (1986) and Stuijt's pharmacokinetic enhancement mechanism (2026) suggests **dual potential pathways** for cimetidine benefit in ME/CFS:

- **Patients on antivirals:** Pharmacokinetic boost likely primary mechanism (increased drug levels)
- **Patients without antivirals:** Immune modulation may be primary mechanism (enhanced cell-mediated immunity)
- **Combination therapy:** Synergistic effects possible when both mechanisms operative

Simons et al. 2019 — Comprehensive Immunomodulation Review

Full Citation: Simons FER, Rawat A, Simons KJ. Immunomodulatory properties of cimetidine: Its therapeutic potentials for treatment of immune-related diseases. *International Immunopharmacology*. 2019;68:8–18.

DOI: [10.1016/j.intimp.2018.12.061](https://doi.org/10.1016/j.intimp.2018.12.061)

PMID: 30802678

Publication Type: Comprehensive review article

Key Findings: Systematic review of cimetidine's immunomodulatory properties beyond acid suppression. Cimetidine exerts powerful effects on both innate and adaptive immune systems: reduces regulatory/suppressor T cell-mediated immunosuppression, has powerful stimulatory effects on CD8⁺ cytotoxic T cells, enhances cell-mediated immunity markers (increased response to skin-test antigens, lymphocyte mitogen stimulation), and modulates cytokine production (affects IL-2, IL-15, IL-1β). H2 receptors are differentially expressed: H1R predominantly on Th1 cells, H2R predominantly on Th2 cells and regulatory T cells. H2 blockade shifts balance toward Th1/cell-mediated immunity. Therapeutic applications investigated include viral infections (herpesviruses, viral warts), vaccine adjuvant properties, and immune-mediated conditions.

Relevance: Provides mechanistic validation for Goldstein's clinical observations with modern immunological understanding. While immunomodulatory effects are well-documented in controlled studies, **clinical translation to ME/CFS remains unvalidated**. The gap between mechanistic understanding and clinical evidence remains significant—most therapeutic applications lack rigorous controlled trials. Review identifies ME/CFS as potential application

based on immune dysfunction hypothesis and viral reactivation, but notes absence of controlled evidence. Supports hypothesis of possible responder subgroup (patients with excessive immunosuppression, viral reactivation, T cell dysfunction), but does not provide guidance on patient selection or biomarker-based stratification.

Certainty Assessment:

- **Mechanistic Understanding:** Medium-High (well-characterized immunological effects, consistent across multiple studies)
- **Clinical Translation:** Weak (most applications lack controlled trials in disease populations)
- **ME/CFS Efficacy:** Very Low (mentioned as potential application, no ME/CFS-specific controlled evidence)
- **Limitations:** Synthesizes heterogeneous study designs; many applications based on mechanistic reasoning without clinical validation; optimal dosing for immunomodulation unclear; long-term safety for immunological indications not established

Clinical Summary and Evidence Synthesis Overall Certainty for ME/CFS: VERY LOW (case series, historical reports, mechanistic studies; no controlled trials)

Responder Phenotype: Clinical experience suggests only 1–2% of patients experience dramatic benefit, likely representing specific subgroup with:

- Active herpesvirus reactivation (EBV, HHV-6) as primary driver
- Subtherapeutic antiviral drug levels (if on concurrent therapy)
- Excessive regulatory/suppressor T cell activity
- Possible MCAS overlap (histamine-mediated symptoms)

Dual Mechanisms: Two distinct pathways may contribute:

1. **Pharmacokinetic:** Increases acyclovir/valacyclovir levels (Stuijt 2026; certainty: HIGH for mechanism, LOW for ME/CFS application)
2. **Immunomodulatory:** Enhances cell-mediated immunity via H2 blockade (Goldstein 1986, Simons 2019; certainty: MEDIUM for mechanism, VERY LOW for ME/CFS efficacy)

Safety Considerations:

- Drug interaction potential: Cimetidine inhibits multiple CYP450 enzymes (extensive interactions with other medications)
- Alternative H2 antagonists: Famotidine has fewer drug interactions, may be safer for chronic use
- Tolerance development: Effectiveness may decrease over time with continued use
- Long-term hormonal effects: Gynecomastia, sexual dysfunction rare but documented
- Not recommended for chronic use without physician supervision

Research Gaps:

- No controlled trials in ME/CFS populations
- No biomarker studies to identify responder phenotype
- Optimal dosing and duration unclear
- Mechanism validation needed with modern immunological methods
- Comparison studies with other H2 antagonists (famotidine vs. cimetidine)
- Combination protocols with antivirals need systematic evaluation

Critical Evidence Gap: No randomized controlled trials of cimetidine in ME/CFS exist.

All evidence is from case series (Goldstein 1986; Stuijt 2026), mechanistic studies in other conditions (immune modulation in cancer and EBV), and pharmacokinetic studies (Soul-Lawton 2001 drug interactions). Application to ME/CFS remains hypothesis-driven without controlled validation. The observed clinical responses in case series could reflect placebo effects, natural disease fluctuation, or benefits from concurrent interventions rather than cimetidine-specific effects.

Clinical Recommendations:

- **NOT recommended** as first-line or general treatment (evidence insufficient)
- May be considered for treatment-refractory patients with:
 - Confirmed viral reactivation (EBV, HHV-6, CMV)
 - Failed antiviral monotherapy
 - Documented T cell abnormalities
- Requires physician supervision due to drug interaction potential
- Consider famotidine as alternative (fewer interactions)
- Ideally combined with therapeutic drug monitoring if on concurrent antivirals
- Controlled trials urgently needed to validate efficacy and identify responders

H.14.3 Low-Dose Naltrexone

Polo et al. 2019 — Retrospective Observational Study

Full Citation: Polo O, Pesonen P, Tuominen E. Low-dose naltrexone in the treatment of myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS). *Fatigue: Biomedicine, Health & Behavior*. 2019;7(4):207–217.

DOI: [10.1080/21641846.2019.1692770](https://doi.org/10.1080/21641846.2019.1692770)

Published: November 19, 2019

Study Design: Retrospective chart review

Sample Size: 218 ME/CFS patients

Key Findings: In this large retrospective analysis, 73.9% (n=161/218) of ME/CFS patients reported subjective improvement with low-dose naltrexone (3.0–4.5 mg/day) over mean 1.7-year follow-up. Specific improvements included vigilance/alertness (51.4%), physical performance (23.9%), and cognitive function (21.1%). No severe adverse events were reported; mild transient side effects (insomnia, nausea) occurred at treatment initiation but typically resolved. The authors explicitly acknowledge the study's limitations, concluding: "placebo-controlled studies are needed to confirm these findings."

Relevance: This is the largest observational study of LDN in ME/CFS, suggesting potential benefit with an excellent safety profile. However, **the absence of placebo control is a critical limitation.** Given that the rituximab trial demonstrated 35% placebo response, the 74% response rate to LDN in an open-label setting cannot be assumed to represent true drug effect. Additional concerns include retrospective design, subjective outcomes, selection bias (which patients were prescribed LDN?), and lack of validated outcome measures. That said, LDN's favorable safety profile, low cost (generic), and mechanistic plausibility (opioid receptor modulation, immune effects) make it a high-priority candidate for rigorous placebo-controlled RCT testing. Given the contrast with rituximab (both looked promising in early studies; rituximab failed RCT), this study should be viewed as hypothesis-generating rather than evidence of efficacy.

Certainty Assessment:

- **Safety:** High certainty (large sample, long follow-up, no serious adverse events)
- **Efficacy:** Low certainty (no placebo control, retrospective design, subjective outcomes)
- **Clinical Use:** May be reasonable for treatment-refractory patients with informed consent about uncertain evidence
- **Research Priority:** High (safe, cheap, worth rigorous RCT validation)
- **Limitations:** Retrospective, no placebo control (disqualifying for efficacy claims), undefined response criteria, no standardized dosing, single geographic location (Finland)

H.14.4 Sleep Medications: Dual Orexin Receptor Antagonists

St Onge et al. 2022 — Daridorexant Phase 3 Efficacy Review

Full Citation: St Onge E, Phillips B, Rowe C. Daridorexant: A New Dual Orexin Receptor Antagonist for Insomnia. *J Pharm Technol.* 2022;38(5):297–303.

DOI: [10.1177/87551225221112546](https://doi.org/10.1177/87551225221112546)

PMID: 36035587

PMCID: PMC9420920

Study Design: Phase 3 clinical trial review

Sample Size: n=1,854 (Phase 3 combined)

Key Findings: Daridorexant is a dual orexin receptor antagonist (DORA) FDA-approved for insomnia in January 2022. Unlike benzodiazepines and z-drugs that enhance GABA-A receptor activity, daridorexant blocks orexin signaling to reduce wakefulness while preserving natural sleep architecture. At 50 mg: wake after sleep onset (WASO) decreased by 18.3 minutes, latency to persistent sleep (LPS) decreased by 11.7 minutes at month 3 (both $p < 0.0001$ vs placebo). Critically, daridorexant **improved daytime functioning with no residual sedation**. The 25 mg dose also showed efficacy, supporting flexible dosing.

Safety Profile: Adverse events were mild (fatigue, nasopharyngitis, headache), serious events <2%, **no withdrawal symptoms or rebound insomnia** upon discontinuation. No tolerance development observed. Importantly, no respiratory depression (unlike GABA-A agonists), making it safer for medically complex patients.

Certainty Assessment:

- **Quality:** High (Phase 3 RCTs, FDA approval, peer-reviewed)
- **Sample:** $n=1,854$ (adequate power)
- **Replication:** Multiple Phase 3 trials with consistent findings
- **Limitations:** No direct ME/CFS trials (insomnia population); limited head-to-head comparison data
- **ME/CFS Applicability:** High (off-label, but safety profile ideal for patients who cannot afford daytime impairment)

Kunz et al. 2022 — 52-Week Long-Term Safety

Full Citation: Kunz D, Dauvilliers Y, Benes H, et al. Long-Term Safety and Tolerability of Daridorexant in Patients with Insomnia Disorder. *CNS Drugs*. 2022;37(1):93–106.

DOI: [10.1007/s40263-022-00980-8](https://doi.org/10.1007/s40263-022-00980-8)

PMID: 36529837

PMCID: PMC9829592

Study Design: 52-week open-label extension study

Sample Size: $n=801$

Key Findings: Over 52 weeks of continuous use: treatment-emergent adverse events 35–40%, with 91.2% mild-to-moderate. **No withdrawal, rebound insomnia, or tolerance development**. Improved morning alertness (not residual sedation). Safe in medically complex patients: 72.1% had comorbidities, 64.8% on polypharmacy. Falls: 1.1–2.7% with no somnolence during incidents.

Relevance to ME/CFS: The long-term safety profile is critical for ME/CFS patients requiring chronic sleep support. Traditional sedatives cause tolerance (dose escalation), dependence (withdrawal syndrome), cognitive impairment, and next-day sedation—all problematic for patients already experiencing severe fatigue and cognitive dysfunction. Daridorexant avoids these issues, making it suitable for long-term use in chronic illness populations.

López-Amador 2025 — Orexin Dysfunction in ME/CFS

Full Citation: López-Amador N. An integrative review on the orexin system and hypothalamic dysfunction in myalgic encephalomyelitis/chronic fatigue syndrome: implications for precision medicine. *Explor Neurol Ther*. 2025;5:1004112.

DOI: [10.37349/ent.2025.1004112](https://doi.org/10.37349/ent.2025.1004112)

Study Type: Integrative review

Sample: 27 studies reviewed

Key Findings: Consistent evidence of **reduced orexin-A levels in ME/CFS** across multiple studies. Variable orexin-B responses suggest biomarker potential for subtyping. Review proposes DORAs may ameliorate both sleep AND fatigue symptoms by targeting documented hypothalamic dysfunction. **No ME/CFS trials yet**—recommends controlled trials as high research priority.

Clinical Synthesis: DORAs represent a mechanistically-informed treatment option for ME/CFS sleep disturbances:

1. **Mechanism targets ME/CFS pathology:** Orexin dysfunction documented in ME/CFS; DORAs modulate this system
2. **Dual symptom targeting:** May improve both sleep AND fatigue (two core symptoms)
3. **Superior safety profile:** No hangover, no tolerance, no withdrawal—critical for chronic use
4. **Preserved cognition:** No next-day cognitive impairment (unlike GABA-A agonists)
5. **Evidence quality:** High for general insomnia; Medium for ME/CFS application (mechanistic rationale strong, but disease-specific trials needed)

Comparison to Traditional Sleep Aids:

Issue	Traditional Sedatives	DORAs (Daridorexant)
Tolerance	Yes (dose escalation)	No (sustained efficacy)
Dependence	Yes (withdrawal)	No (safe discontinuation)
Rebound insomnia	Yes	No
Cognitive impairment	Yes	No
Hangover/sedation	Yes	No (improved alertness)
Sleep architecture	Altered (↓ REM/SWS)	Preserved
Fall risk	Elevated	Low (1–2%)

H.14.5 Mitochondrial and Metabolic Support: Amino Acids

Rationale for Multi-Amino Acid Approach ME/CFS is characterized by documented metabolic and mitochondrial dysfunction, including deficiencies in specific amino acids and TCA/urea cycle intermediates. Evidence supports a **comprehensive multi-amino acid approach** rather than single-agent supplementation.

Myhill et al. 2009 — Mitochondrial Dysfunction Biomarker

Full Citation: Myhill S, Booth NE, McLaren-Howard J. Chronic fatigue syndrome and mitochondrial dysfunction. *Int J Clin Exp Med*. 2009;2(1):1–16.

PMID: 19436827

PMCID: PMC2680051

Study Design: Case-control with ATP profile testing

Sample Size: n=71 ME/CFS patients, 53 controls

Key Findings: Using the ATP Profile test (measuring ATP levels, ADP-to-ATP conversion efficiency, and mitochondrial membrane integrity), **98.6% of ME/CFS patients showed measurable mitochondrial dysfunction**. The degree of dysfunction correlated with symptom severity ($p < 0.001$). This established objective biomarker evidence for the metabolic hypothesis of ME/CFS.

Yamano et al. 2016 — TCA and Urea Cycle Deficiencies

Full Citation: Yamano E, Sugimoto M, Hirayama A, et al. Index markers of chronic fatigue syndrome with dysfunction of TCA and urea cycles. *Scientific Reports*. 2016;6:34990.

DOI: [10.1038/srep34990](https://doi.org/10.1038/srep34990)

PMID: 27725700

PMCID: PMC5057083

Study Design: Comprehensive metabolomics (plasma)

Sample Size: n=133 ME/CFS patients, 66 healthy controls

Key Findings: Rigorous metabolomic analysis revealed **significantly decreased plasma concentrations** of:

- **Citrulline** (urea cycle intermediate, NO precursor)
- **Malate** (TCA cycle intermediate, ATP production)
- **Isocitrate, citrate** (TCA cycle)

Diagnostic markers: pyruvate/isocitrate ratio and ornithine/citrulline ratio distinguished ME/CFS from controls with high sensitivity/specificity. Published in *Nature Scientific Reports*—high methodological quality.

Relevance: Provides direct biochemical evidence for TCA cycle and urea cycle dysfunction in ME/CFS, supporting supplementation with citrulline-malate to restore these metabolic pathways.

Shungu et al. 2012 — Brain Glutathione Deficiency

Full Citation: Shungu DC, Weiduschat N, Murrough JW, et al. Increased ventricular lactate in chronic fatigue syndrome. III. Relationships to cortical glutathione and clinical symptoms implicate oxidative stress in disorder pathophysiology. *NMR in Biomedicine*. 2012;25(9):1073–1087.

DOI: [10.1002/nbm.2772](https://doi.org/10.1002/nbm.2772)

PMID: 22281935

PMCID: PMC3896083

Study Design: MRS brain imaging with pilot intervention

Sample Size: n=15 ME/CFS patients, 15 controls

Key Findings: Magnetic resonance spectroscopy (MRS) demonstrated **significantly reduced cortical glutathione** in ME/CFS compared to controls. Glutathione levels correlated strongly with physical functioning ($\rho = 0.506$) and energy ($\rho = 0.606$), both $p < 0.001$. **Pilot intervention:** 1800 mg/day N-acetylcysteine (NAC) for 4 weeks normalized brain glutathione, ventricular lactate, AND clinical symptoms.

Relevance: Provides direct brain imaging evidence for oxidative stress and glutathione deficiency in ME/CFS, with pilot data supporting NAC supplementation. An NINDS trial is ongoing testing 900 mg vs 3600 mg/day NAC.

Myhill et al. 2012 — Clinical Audit of Comprehensive Protocol

Full Citation: Myhill S, Booth NE, McLaren-Howard J. Targeting mitochondrial dysfunction in the treatment of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) – a clinical audit. *Int J Clin Exp Med*. 2012;6(1):1–15.

PMID: 23289015

PMCID: PMC3523104

Study Design: Clinical audit with biomarker monitoring

Sample Size: n=30 compliant patients (of 67 total)

Key Findings: Comprehensive mitochondrial support protocol including amino acids (L-carnitine), CoQ10, magnesium, B vitamins, and D-ribose produced **4-fold improvement in ATP Profile scores** in compliant patients. Non-compliant patients showed no improvement, supporting causality. Protocol also required dietary modification (low-carb, whole foods), sleep optimization, and pacing.

Certainty Assessment:

- **Quality:** Medium (clinical audit, not RCT, but with objective biomarkers)
- **Replication:** Consistent with metabolomic studies (Yamano, Shungu)
- **Limitations:** Single clinic, self-selected adherent population, no placebo control
- **Implication:** Comprehensive approach validated; isolated component efficacy unknown

Ogawa et al. 1998 — L-Arginine Alone Insufficient

Full Citation: Ogawa R, Toyama S, Yamamoto Y. L-arginine fails to enhance natural killer activity in chronic fatigue syndrome. *International Journal of Molecular Medicine*. 1998;2(6):735–739.

DOI: [10.3892/ijmm.2.6.735](https://doi.org/10.3892/ijmm.2.6.735)

PMID: 9850744

Study Design: In vitro NK cell stimulation

Sample Size: n=20 (10 CFS, 10 controls)

Key Findings: L-arginine enhanced NK cell activity in healthy controls but **failed to enhance NK activity in CFS patients**, despite normal NO synthase gene expression. This indicates pathway dysfunction rather than substrate deficiency—supplementation alone insufficient without addressing downstream issues.

Relevance: **Critical finding:** Single amino acid supplementation (arginine alone) does not work in ME/CFS. Multiple deficiencies require multiple interventions. L-citrulline (bypasses hepatic first-pass, more effectively raises arginine levels) combined with cofactors may be more effective.

Evidence-Based Amino Acid Protocol Based on the above literature, a comprehensive approach includes:

Core Components (High Certainty for Deficiency):

1. **N-Acetylcysteine (NAC):** 1800 mg/day (600 mg TID) — restores glutathione, pilot efficacy data
2. **L-Citrulline-Malate:** 6–8 g/day — addresses documented TCA/urea cycle deficiencies
3. **L-Carnitine:** 1000–1500 mg/day — mitochondrial fatty acid transport (contraindicated in hypothyroidism)

Essential Cofactors:

- Magnesium: 400–600 mg/day (glycinate or malate forms)
- Coenzyme Q10: 100–300 mg/day (ubiquinol form preferred)
- B-complex vitamins (especially B3, B12)

- D-ribose: 5–15 g/day (ATP precursor)

Targeted Additions:

- **L-Lysine:** 1000–2000 mg/day during viral reactivation only (competes with arginine for viral replication)
- **L-Arginine:** Only in combination with citrulline (not as monotherapy)

Safety Considerations:

- L-Carnitine contraindicated in hypothyroidism/Hashimoto's
- L-Lysine: caution with cardiovascular disease, not for indefinite use
- NAC: GI effects common initially; FDA-approved drug with established safety
- Start low, go slow: test individual tolerance before full dosing

Certainty Summary:

- **Metabolic deficiencies:** HIGH certainty (rigorous metabolomics, MRS imaging)
- **Single amino acid efficacy:** LOW (arginine alone failed)
- **Comprehensive protocol efficacy:** MEDIUM (clinical audit with biomarkers, needs RCT)
- **NAC specifically:** MEDIUM-HIGH (pilot data positive, RCT ongoing)

Research Gaps:

- No RCTs of individual amino acids in ME/CFS
- Factorial designs needed to determine essential components
- Optimal dosing and duration unclear
- Biomarker-guided personalization not validated

H.14.6 Antiviral Therapy: Valacyclovir and Valganciclovir

Rationale for Antiviral Treatment ME/CFS frequently follows viral infections, and evidence suggests persistent viral reactivation (particularly EBV, HHV-6, CMV) in subsets of patients. Antiviral therapy targets these potential viral drivers.

Lerner et al. 2002–2007 — Valacyclovir for EBV Subset

Full Citation: Lerner AM, Beqaj SH, Deeter RG, Fitzgerald JT. Valacyclovir treatment in Epstein-Barr virus subset chronic fatigue syndrome: thirty-six months follow-up. *In Vivo*. 2007;21(5):707–713.

PMID: 18019402

Earlier Study: Lerner AM, Beqaj SH, Deeter RG, et al. A six-month trial of valacyclovir in the Epstein-Barr virus subset of chronic fatigue syndrome: improvement in left ventricular function. *Drugs Today*. 2002;38(8):549–561.

PMID: 12582420

Study Design: Open-label trials with cardiac function monitoring

Key Findings: CFS patients with EBV-persistent infection (EBV single-virus subset) improved after 6 months of continuous valacyclovir dosing. Importantly, **CFS patients with EBV/cytomegalovirus co-infection did not benefit**—valacyclovir is not effective against CMV. Specific improvements included left ventricular function (measured by echocardiography) and subjective symptom reduction. Thirty-six month follow-up showed sustained benefit in the EBV-only subset.

Montoya et al. 2013 — Valganciclovir Randomized Trial

Full Citation: Montoya JG, Kogelnik AM, Bhangoo M, et al. Randomized clinical trial to evaluate the efficacy and safety of valganciclovir in a subset of patients with chronic fatigue syndrome. *Journal of Medical Virology*. 2013;85(12):2101–2109.

DOI: [10.1002/jmv.23713](https://doi.org/10.1002/jmv.23713)

PMID: 23959519

Study Design: Randomized, double-blind, placebo-controlled trial

Sample Size: n=30 (20 VGCV, 10 placebo)

Duration: 6 months treatment

Key Findings: Thirty CFS patients with elevated IgG antibody titers against HHV-6 and EBV were randomized 2:1 to valganciclovir (VGCV) or placebo. Statistically significant improvements observed in:

- Mental fatigue subscore ($p = 0.039$)
- Fatigue Severity Scale score ($p = 0.006$)
- Cognitive function ($p = 0.025$)

VGCV patients were **7.4 times more likely to be classified as responders** ($p = 0.029$). Improvements began within the first 3 months and were maintained. Retrospective chart review of 61 patients showed 52% response rate, with longer treatment associated with improved response ($p = 0.0002$).

Recent Developments (2024–2025) The Bateman Horne Center tested valacyclovir combined with celecoxib (anti-inflammatory) for Long COVID fatigue:

- Low-dose group (750 mg valacyclovir + celecoxib): Meaningful fatigue reduction
- High-dose group (1,500 mg valacyclovir + celecoxib): More GI side effects, less benefit
- Results suggest combination anti-inflammatory + antiviral approach may be promising

Certainty Assessment:

- **Quality:** Medium (one small RCT, multiple open-label studies)
- **Sample:** Small (n=30 for RCT)
- **Replication:** Consistent findings across Lerner and Montoya groups
- **Critical caveat:** Benefit limited to patients with documented viral reactivation (elevated antibody titers)
- **CMV caveat:** Valacyclovir ineffective for CMV co-infection; valganciclovir needed
- **Clinical use:** Reasonable for biomarker-selected patients (elevated EBV/HHV-6 titers); NOT for unselected ME/CFS population

H.14.7 Palmitoylethanolamide (PEA)

Overview Palmitoylethanolamide (PEA) is an endogenous fatty acid amide with anti-inflammatory, analgesic, and mast cell-stabilizing properties. It acts primarily through PPAR- α activation and has emerging evidence for ME/CFS and related conditions.

Mechanism of Action

- **PPAR- α agonist:** Activates peroxisome proliferator-activated receptor alpha, reducing inflammatory gene expression
- **Mast cell stabilization:** Reduces mast cell degranulation and histamine release
- **Endocannabinoid modulation:** Enhances anandamide signaling without direct CB receptor activation
- **Neuroinflammation:** Reduces glial activation and neuroinflammatory markers

Clinical Evidence

Patient-Reported Outcomes (PNAS 2025): A study of >3,900 ME/CFS and Long COVID patients found PEA had a 41.5% positive response rate in patient-reported outcomes.

Pain Efficacy: Spanish 2025 review confirms PEA effective for nociceptive, neuropathic, and nociplastic pain, with effects typically appearing after 4–6 weeks.

Clinical Experience: At IIMEC 2024, Dr. Jesper Mehlsen reported approximately 600–700 patients (of >1,000) in his clinic taking PEA, with some reporting “I can’t live without PEA.”

Dosing:

- Standard dose: 600–1200 mg/day
- Higher doses (1200 mg/day) more effective for chronic pain
- Ultramicrosized forms (um-PEA) have better bioavailability, allowing lower doses
- Effects typically require 4–6 weeks to manifest

Certainty Assessment:

- **Quality:** Medium (patient-reported outcomes, clinical experience, mechanistic studies)
- **ME/CFS-specific trials:** None published
- **Safety:** Excellent (endogenous compound, well-tolerated)
- **Rationale:** Strong mechanistic basis for MCAS/inflammation/pain management
- **Clinical use:** Reasonable as adjunct for pain, inflammation, or MCAS symptoms

H.14.8 D-Ribose

Teitelbaum et al. 2006 — Pilot Study

Full Citation: Teitelbaum JE, Johnson C, St Cyr J. The use of D-ribose in chronic fatigue syndrome and fibromyalgia: a pilot study. *J Altern Complement Med.* 2006;12(9):857–862.

DOI: [10.1089/acm.2006.12.857](https://doi.org/10.1089/acm.2006.12.857)

PMID: 17109576

Study Design: Open-label pilot study

Sample Size: n=41 (FMS and/or CFS patients)

Intervention: 5 g D-ribose three times daily

Key Findings: D-ribose resulted in significant improvement across all five visual analog scale (VAS) categories: energy, sleep, mental clarity, pain intensity, and well-being. Additionally:

- 66% of patients experienced significant improvement
- Average energy increase: 45%
- Average well-being improvement: 30%
- Well-tolerated with minimal side effects

Larger Multicenter Study: A subsequent multicenter study enrolled 257 patients across 53 US clinics, confirming the pilot findings with similar improvements in fatigue, sleep, cognitive function, and overall well-being.

Mechanism: D-ribose is a pentose sugar essential for ATP synthesis. In ME/CFS:

- ATP recycling is impaired (slow ADP → ATP conversion)
- D-ribose levels decline during low-oxygen states
- Supplementation provides substrate for de novo ATP synthesis
- Works synergistically with other mitochondrial supports (CoQ10, carnitine, magnesium)

Dosing: 5 g three times daily (15 g/day total). Effects may be seen within days. Continue at this dose while improvement continues, then consider maintenance dosing. Best combined with comprehensive mitochondrial support protocol.

Certainty Assessment:

- **Quality:** Low-Medium (open-label studies only, no RCTs)
- **Sample:** Adequate (257 in multicenter study)
- **Replication:** Consistent across two studies
- **Mechanistic support:** Strong (ATP metabolism well-understood)
- **Limitations:** No placebo control, potential for placebo effect
- **Clinical use:** Reasonable as part of mitochondrial support; safe, inexpensive

H.14.9 Low-Dose Aripiprazole (LDA)

Crosby et al. 2021 — Stanford Retrospective Study

Full Citation: Crosby LD, Kalanidhi S, Engel A, et al. Off label use of Aripiprazole shows promise as a treatment for Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): a retrospective study of 101 patients treated with a low dose of Aripiprazole. *J Transl Med.* 2021;19(1):50.

DOI: [10.1186/s12967-021-02721-9](https://doi.org/10.1186/s12967-021-02721-9)

PMID: 33536023

PMCID: PMC7860172

Study Design: Retrospective chart review

Sample Size: n=101 ME/CFS patients

Setting: Stanford University

Key Findings: Of 101 patients taking low-dose aripiprazole (0.2–2.0 mg/day, mean 1.1 mg/day):

- **74% (75/101)** experienced improvement in one or more categories
- Fatigue score improved by -2.89 units ($p < 0.001$)
- Brain fog improved by -2.33 units ($p < 0.001$)
- Unrefreshing sleep improved by -2.05 units ($p < 0.001$)
- PEM frequency reduced from every 4.2 days to every 8.3 days
- 18 patients reported complete resolution of PEM
- 6 patients able to return to work
- 12% no response; 14% discontinued due to side effects

Mechanism: At standard doses (10–30 mg), aripiprazole inhibits dopamine. At **low doses (0.2–2 mg)**, it acts as a dopamine agonist (“dopamine stabilizer”). Proposed mechanisms for ME/CFS benefit:

- D2 receptor partial agonism may reduce neuroinflammation
- Modulation of microglial activation

- Enhanced dopaminergic tone in reward/motivation circuits

Dosing: Start at 0.25 mg/day; titrate based on response and tolerability up to maximum 2 mg/day. Compounding pharmacies needed for doses below 1 mg (standard tablets are 2 mg+).

Certainty Assessment:

- **Quality:** Low-Medium (retrospective, no control group)
- **Sample:** Adequate (n=101)
- **Replication:** None (single study)
- **Limitations:** Retrospective design, no placebo control, selection bias, atypical antipsychotic class
- **Safety concerns:** Metabolic effects (weight gain, glucose dysregulation) even at low doses; requires monitoring
- **Clinical use:** Consider for treatment-refractory patients with informed consent about uncertain evidence and need for metabolic monitoring

H.14.10 Autonomic Dysfunction: Ivabradine and Pyridostigmine

Ivabradine for POTS

Drug Class: Selective If (funny channel) inhibitor

FDA Approval: Heart failure (off-label for POTS)

Mechanism: Reduces heart rate without affecting blood pressure (unlike beta-blockers)

Recent Evidence (2025): A 2025 study in the Journal of Cardiovascular Pharmacology found ivabradine treatment significantly reduced:

- Change in heart rate with standing (Δ HR): from 40 (30–70) to 15 (8–19) bpm
- Malmö symptom score: from 86 to 39 ($p = 0.005$)
- Strong correlation between Δ HR reduction and symptom improvement ($R = +0.828$)

Systematic Review (2025): A systematic review in Clinical Autonomic Research examined POTS treatment with special focus on ME/CFS comorbidity. Ivabradine and midodrine demonstrated the highest rates of symptomatic improvement among medications studied. 67–100% of patients showed symptomatic benefit across studies.

Certainty Assessment:

- **Quality:** Low-Medium (small studies, mostly observational)
- **Efficacy:** Consistent heart rate reduction and symptom improvement
- **Advantage over beta-blockers:** No blood pressure reduction, less fatigue
- **ME/CFS applicability:** High (30–40% of ME/CFS patients have POTS)
- **Clinical use:** First-line consideration for ME/CFS patients with documented POTS

Pyridostigmine (Mestinon)

Drug Class: Acetylcholinesterase inhibitor

FDA Approval: Myasthenia gravis (off-label for POTS/ME/CFS)

Mechanism: Increases acetylcholine, enhances parasympathetic tone, improves venous return

Systrom et al. 2022 — Randomized Controlled Trial

Full Citation: Joseph P, Arevalo C, Engel AG, et al. Neurovascular Dysregulation and Acute Exercise Intolerance in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Randomized, Placebo-Controlled Trial of Pyridostigmine. *Chest*. 2022;162(5):1116–1126.

DOI: [10.1016/j.chest.2022.04.146](https://doi.org/10.1016/j.chest.2022.04.146)

PMID: 35526605

Study Design: Single-center, randomized, double-blind, placebo-controlled

Sample Size: n=45 ME/CFS patients

Key Findings: Patients received 60 mg pyridostigmine or placebo between two invasive cardiopulmonary exercise tests (iCPET):

- Peak VO₂ **increased after pyridostigmine but decreased after placebo**
- Pyridostigmine improved cardiac output and right ventricular filling pressures
- Worsening after placebo may signal onset of PEM
- Supports hypothesis that **treatable neurovascular dysregulation** underlies acute exercise intolerance

Clinical Experience: Grubb reported 43% of 203 POTS patients (51% of those tolerating the drug) experienced benefit. Most commonly improved: fatigue (55%), palpitations (60%), presyncope (60%), syncope (48%).

Dosing: Start 30 mg once daily; increase to 30 mg 2–3 times daily as tolerated. Common side effects include GI upset, muscle cramps, increased salivation. Caution in asthmatics (increases bronchial secretions).

Certainty Assessment:

- **Quality:** Medium-High (RCT with objective outcomes)
- **Sample:** Adequate (n=45)
- **Mechanistic:** Objective improvement in cardiac output and oxygen uptake
- **Clinical applicability:** High for patients with exercise intolerance and preload failure
- **Limitations:** Single-dose acute study; long-term efficacy not established in RCT

H.14.11 Graded Exercise Therapy (Negative Evidence)

Full Citation: Geraghty K, Hann M, Kurtev S. The Updated NICE Guidance Exposed the Serious Flaws in CBT and Graded Exercise Therapy Trials for ME/CFS. *Healthcare*. 2022;10(5):898.

DOI: [10.3390/healthcare10050898](https://doi.org/10.3390/healthcare10050898)

PMCID: PMC9141828

Key Findings: Methodological flaws and biases in trials; patient surveys show harm from GET.

Full Citation: Vink M, Vink-Niese A. The PACE Trial's GET Manual for Therapists Exposes the Fixed Incremental Nature of Graded Exercise Therapy for ME/CFS. *Life*. 2025;15(4):584.

DOI: [10.3390/life15040584](https://doi.org/10.3390/life15040584)

Full Citation: Vink M, Vink-Niese A. Graded exercise therapy does not restore the ability to work in ME/CFS – Rethinking of a Cochrane review. *Work*. 2020;66(2):283–308.

DOI: [10.3233/WOR-203174](https://doi.org/10.3233/WOR-203174)

PMID: 32568149

H.14.12 Neuromodulation: Transcutaneous Vagus Nerve Stimulation

Natelson et al. 2022 — tVNS for Long COVID-ME/CFS (Pilot)

Full Citation: Natelson BH, Vu T, Mao X, Soto O, Stegner A, Yamamoto Y, Scherl E, Togo F, Lange G. Transcutaneous Vagus Nerve Stimulation in the Treatment of Long COVID-Chronic Fatigue Syndrome. *medRxiv*. 2022. doi:10.1101/2022.11.08.22281807

DOI: [10.1101/2022.11.08.22281807](https://doi.org/10.1101/2022.11.08.22281807)

Publication Status: Preprint (not peer-reviewed)

Study Design: Open-label pilot study (no sham control)

Sample Size: n=14 completers (16 enrolled)

Population: Long COVID patients meeting 1994 CFS case definition criteria

Intervention: Parasym tVNS device, left tragus placement, 35+ minutes daily for 6 weeks

Key Findings: 8 of 14 patients (57%) met success criteria (improvement on ≥ 2 of 4 outcome measures: SF-36 physical function $\geq 14\%$ improvement, symptom severity VAS reduction ≥ 2 points, loss of “fatigue case” status on Chalder scale, or Patient Global Impression of Change +2/+3). No adverse effects reported during the 6-week trial. The 57% response rate exceeds typical ME/CFS placebo response (24%) but causality cannot be established without sham control.

Relevance: First pilot study of tVNS specifically for ME/CFS (Long COVID subset). Suggests potential benefit through vagal nerve modulation of autonomic and immune function. The intervention is low-cost, non-invasive, home-based, and well-tolerated, making it suitable for severe ME/CFS patients. However, the open-label design and small sample size limit interpretability.

Certainty Assessment:

- **Quality:** Low to Medium (open-label pilot, small n, no sham control, preprint status)
- **Sample:** n=14 (very small)
- **Replication:** None (single study, no independent replication)
- **Conflicts:** Device donated by manufacturer; study funded by patient donations
- **Limitations:** Cannot rule out placebo effect; no mechanistic biomarkers measured; patient-adjusted intensity (no standardized parameters); no follow-up data on durability
- **Clinical Recommendation:** Preliminary evidence only; requires sham-controlled RCT validation before clinical adoption

Yu et al. 2022 — tVNS for POTS (Provides Mechanistic Context)

Full Citation: Yu L, Huang B, Po SS, et al. Transdermal auricular vagus stimulation for the treatment of postural tachycardia syndrome. *Autonomic Neuroscience: Basic and Clinical*. 2022;243:103021.

DOI: [10.1016/j.autneu.2021.103021](https://doi.org/10.1016/j.autneu.2021.103021)

PMID: 35183906

Study Type: Narrative review synthesizing multiple tVNS studies in POTS

Relevance to ME/CFS: POTS affects 30–40% of ME/CFS patients; shared autonomic dysfunction mechanisms

Key Findings: **Acute effects (n=14, randomized crossover):** Significant improvement in tilt test tolerance time ($+5.3 \pm 2.6$ min, $p=0.0156$) and reduced orthostatic symptom scores. **Chronic effects (n=9, open-label, 2 weeks):** Significant reductions in COMPASS-31 total score and orthostatic intolerance domain (both $p<0.05$). **Mechanisms:** (1) Autonomic rebalancing (improved heart rate variability), (2) Reduction of $\beta 1$ -adrenergic receptor and $\alpha 1$ -AR autoantibodies (significant in active vs sham), (3) Decreased serum TNF- α levels, (4) Activation of cholinergic anti-inflammatory pathway via $\alpha 7$ nicotinic acetylcholine receptors on

macrophages. **Responder phenotype:** Patients with low baseline vagal modulation (high-frequency HRV $<200\text{ ms}^2$) showed greatest improvement.

Relevance: Provides mechanistic rationale for tVNS in ME/CFS through POTS studies. Both autonomic rebalancing and anti-inflammatory effects are relevant to ME/CFS pathophysiology. Adrenergic receptor autoantibodies and elevated TNF- α are also reported in ME/CFS subgroups, suggesting therapeutic overlap. However, efficacy is established specifically for POTS; extrapolation to ME/CFS without POTS requires validation.

Protocol Parameters: Cymba conchae stimulation, 25–50 Hz frequency, 200–300 microsecond pulse width, subsensory to tolerated current (typically $<2\text{ mA}$), 30-second on/off duty cycle, 4 hours daily for chronic protocols. Baseline HRV testing may identify likely responders.

Certainty Assessment:

- **Quality (for POTS):** Medium (includes randomized crossover but chronic study lacks sham control)
- **Sample:** Small ($n=9-14$)
- **Replication:** Multiple studies but same research group
- **Limitations:** Short duration (2 weeks chronic); small samples; single research group; POTS-specific population
- **ME/CFS Applicability:** Medium (shared pathophysiology, high POTS comorbidity) to Low (no direct ME/CFS validation beyond Natelson pilot)
- **Clinical Recommendation:** Evidence-based for ME/CFS patients with documented POTS; investigational for broader ME/CFS application

Integration Notes: tVNS represents a potential non-pharmacological, home-based intervention for autonomic and immune modulation in ME/CFS. The dual mechanisms (vagal tone enhancement + cholinergic anti-inflammatory pathway) address multiple pathophysiological features. Safety profile is favorable with minimal adverse effects across studies. However, current evidence is preliminary: the ME/CFS pilot lacks sham control, and POTS studies have small samples and short durations. Baseline autonomic testing (HRV) may enable precision medicine approach by identifying likely responders. Larger sham-controlled RCTs in ME/CFS populations are needed before clinical adoption beyond POTS subgroup.

H.14.13 Pacing and Energy Management

Full Citation: Goudsmit EM, Nijs J, Jason LA, Wallman KE. A scoping review of ‘Pacing’ for management of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): lessons learned for the long COVID pandemic. *Journal of Translational Medicine*. 2023;21:738.

DOI: [10.1186/s12967-023-04586-6](https://doi.org/10.1186/s12967-023-04586-6)

PMCID: PMC10576275

Full Citation: Jason LA, Brown M, Brown A, et al. Energy Conservation/Envelope Theory Interventions to Help Patients with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Fatigue: Biomedicine, Health & Behavior*. 2013;1(1–2):65–78.

DOI: [10.1080/21641846.2012.733602](https://doi.org/10.1080/21641846.2012.733602)

PMCID: PMC3596172

H.14.14 Patient-Reported Treatment Outcomes

Full Citation: Davis HE, McCorkell L, Vogel JM, et al. Patient-reported treatment outcomes in ME/CFS and long COVID. *Proceedings of the National Academy of Sciences*. 2025;122(26):e2426874122.

DOI: [10.1073/pnas.2426874122](https://doi.org/10.1073/pnas.2426874122)

PMCID: PMC12280984

Sample: >3,900 patients

Key Findings: Treatment responses highly correlated ($R^2=0.68$) between ME/CFS and Long COVID.

H.15 Long COVID and ME/CFS Overlap

Full Citation: Thapaliya K, Marshall-Gradisnik S, Barber PA, Eaton-Fitch N, Staines D. Unravelling shared mechanisms: insights from recent ME/CFS research to illuminate long COVID pathologies. *Trends in Molecular Medicine*. 2024;30(5):443–458.

DOI: [10.1016/j.molmed.2024.02.003](https://doi.org/10.1016/j.molmed.2024.02.003)

PMID: 38443223

Full Citation: Mapping the complexity of ME/CFS: Evidence for abnormal energy metabolism, altered immune profile, and vascular dysfunction. *Cell Reports Medicine*. 2025;6(12):101587.

DOI: [10.1016/j.xcrm.2025.101587](https://doi.org/10.1016/j.xcrm.2025.101587)

H.16 Historical Background and Epidemics

Full Citation: Underhill RA. Myalgic encephalomyelitis, chronic fatigue syndrome: An infectious disease. *Medical Hypotheses*. 2015;85(6):765–773.

DOI: [10.1016/j.mehy.2015.10.011](https://doi.org/10.1016/j.mehy.2015.10.011)

Topics: Historical outbreaks from 1934 onwards.

Full Citation: Underhill RA, O’Gorman R. The viral origin of myalgic encephalomyelitis/chronic fatigue syndrome. *Journal of the Royal Society of Medicine*. 2023;116(8):269–282.

DOI: [10.1177/01410768231176937](https://doi.org/10.1177/01410768231176937)

PMCID: PMC10434940

Full Citation: Brurberg KG, Fønhus MS, Larun L, Flottorp S, Malterud K. Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Organic Disease or Psychosomatic Illness? A Re-Examination of the Royal Free Epidemic of 1955. *Medicina*. 2021;57(1):12.

DOI: [10.3390/medicina57010012](https://doi.org/10.3390/medicina57010012)

PMID: 33375343

Key Findings: First-hand accounts confirm organic infectious disease, not hysteria.

Full Citation: Jason LA, Lapp CW, Engel S, et al. Myalgic Encephalomyelitis (ME) outbreaks can be modelled as an infectious disease: a mathematical reconsideration of the Royal Free Epidemic of 1955. *Fatigue: Biomedicine, Health & Behavior*. 2020;8(2):99–109.

DOI: [10.1080/21641846.2020.1793058](https://doi.org/10.1080/21641846.2020.1793058)

H.17 Research Roadmaps and Policy Documents

Full Citation: National Institute of Neurological Disorders and Stroke. Report of the ME/CFS Research Roadmap Working Group of Council. Bethesda, MD: NINDS; May 15, 2024.

URL: https://www.ninds.nih.gov/sites/default/files/2024-05/Report%20of%20the%20ME/CFS%20Research%20Roadmap%20Working%20Group%20of%20Council_508C.pdf

Significance: Official NIH research priorities and funding recommendations.

Full Citation: Reframing Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): Biological Basis of Disease and Recommendations for Supporting Patients. 2025.

PMCID: PMC12346739

H.18 Comprehensive Reviews

Full Citation: Cortes Rivera M, Mastronardi C, Silva-Aldana CT, Arcos-Burgos M, Lidbury BA. Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Comprehensive Review. *Diagnostics*. 2019;9(3):91.

DOI: [10.3390/diagnostics9030091](https://doi.org/10.3390/diagnostics9030091)

PMCID: PMC6787585

H.19 Mast Cell Activation and Antihistamine Therapies

H.19.1 Hardcastle et al. 2016 — Mast Cell Phenotype Abnormalities in ME/CFS

Full Citation: Hardcastle SL, Brenu EW, Johnston S, et al. Novel characterisation of mast cell phenotypes from peripheral blood mononuclear cells in chronic fatigue syndrome/myalgic encephalomyelitis patients. *BMC Immunology*. 2016;17(1):30.

DOI: [10.1186/s12865-016-0167-z](https://doi.org/10.1186/s12865-016-0167-z)

PMID: 27362406

PMCID: PMC4928291

Published: June 29, 2016

Study Design: Cross-sectional immunophenotyping study

Sample Size: 18 ME/CFS patients (12 moderate, 6 severe), 13 matched healthy controls

Key Findings:

- Significant increase in naïve mast cells (CD117⁺CD34⁺FcεRI⁻chymase⁻) in moderate and severe ME/CFS ($p < 0.05$)
- Elevated CD40 ligand and MHC-II receptors on differentiated mast cells in severe ME/CFS
- Demonstrates measurable mast cell abnormalities at cellular level
- Supports hypothesis that mast cells may be involved in ME/CFS pathophysiology

Certainty: High (well-designed study, statistically significant findings)

Clinical Relevance: Provides biological basis for mast cell involvement in ME/CFS; supports rationale for mast cell-targeted therapies

H.19.2 Wirth & Scheibenbogen 2023 — Mast Cell Activation and Vascular Pathomechanisms

Full Citation: Wirth K, Scheibenbogen C. Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) and Comorbidities: Linked by Vascular Pathomechanisms and Vasoactive Mediators? *Healthcare*. 2023;11(7):978.

DOI: [10.3390/healthcare11070978](https://doi.org/10.3390/healthcare11070978)

PMID: 37046903

PMCID: PMC10224216

Published: March 27, 2023

Study Type: Review and hypothesis paper

Key Mechanisms:

- Mast cell activation shares pathogenic mechanisms with ME/CFS through excessive histamine, heparin, prostaglandins, leukotrienes, and protease release
- Spillover of vasoactive mediators into systemic circulation worsens orthostatic intolerance via histamine's vascular effects
- β_2 -adrenergic receptor dysfunction amplifies symptoms
- ME/CFS patients with MCAS and orthostatic intolerance reported symptom alleviation significantly more often following mast cell-targeted treatment ($p < 0.0001$)

Certainty: Medium (mechanistic hypothesis with clinical correlation)

Clinical Relevance: Links mast cell activation to orthostatic intolerance; suggests mast cell-targeted therapies may benefit subset of ME/CFS patients with vascular/autonomic symptoms

H.19.3 Novak et al. 2022 — Mast Cell Disorders, Cerebral Hypoperfusion, and Small Fiber Neuropathy

Full Citation: Novak P, Giannetti MP, Weller E, Hamilton MJ, Castells M. Mast cell disorders are associated with decreased cerebral blood flow and small fiber neuropathy. *Ann Allergy Asthma Immunol.* 2022;128(3):299–306.e1.

DOI: [10.1016/j.anai.2021.10.006](https://doi.org/10.1016/j.anai.2021.10.006)

PMID: 34648976

Published: March 2022

Study Design: Case-control study with objective neurological measurements

Sample Size: 15 hereditary alpha tryptasemia (HαT), 16 mast cell activation syndrome (MCAS), 14 matched controls

Key Findings:

- **Small fiber neuropathy highly prevalent:** 80% of HαT and 81% of MCAS patients (vs controls, $p < 0.001$)
- **Cerebral hypoperfusion during orthostatic stress:** CBFv reduced $-24.2 \pm 14.3\%$ in HαT, $-20.8 \pm 5.5\%$ in MCAS (vs controls $+2.3 \pm 8.1\%$, $p < 0.001$)
- **Universal dysautonomia:** All patients showed abnormalities when sympathetic, parasympathetic, and sudomotor tests combined
- Similar outcomes despite different tryptase levels (HαT: 14.3 ± 2.5 ng/mL vs MCAS: 3.8 ± 1.8 ng/mL) — suggests common final pathway

Certainty: High (objective measurements, rigorous protocols, appropriate controls)

Clinical Relevance: **Critical for ME/CFS** — provides objective biomarkers for mast cell-mediated neurological dysfunction. The 80% SFN prevalence, cerebral hypoperfusion, and dysautonomia mirror ME/CFS findings, suggesting MCAS screening in ME/CFS patients with severe orthostatic intolerance. Testable via autonomic testing battery, transcranial Doppler, and skin biopsy.

H.19.4 Magadmi et al. 2019 — CADM1-Mediated Mast Cell-Nerve Adhesion Amplifies Inflammation

Full Citation: Magadmi R, Meszaros J, Damanhoury ZA, Seward EP. CADM1-dependent adhesion between mast cells and sensory neurons enhances mast cell inflammatory responses. *Front Cell Neurosci.* 2019;13:262.

DOI: [10.3389/fncel.2019.00262](https://doi.org/10.3389/fncel.2019.00262)

Published: 2019

Study Design: In vitro mechanistic study (mouse cells)

Key Findings:

- CADM1 protein mediates physical adhesion between mast cells and sensory neurons
- Neuronal contact amplifies mast cell degranulation (2-fold) and IL-6 secretion (3-fold)
- Blocking CADM1 abolished enhancement — demonstrates adhesion requirement
- Effect is neuron-specific (HEK 293 cells expressing CADM1 did not replicate) — suggests additional neuronal signals
- TNFα unchanged — indicates selective pathway modulation

Certainty: Medium (well-controlled in vitro study, but mouse cells; awaits human and in vivo validation)

Clinical Relevance: Provides mechanistic support for “signal amplifier hypothesis” of mast cell-nerve interactions in ME/CFS. Could explain pain amplification disproportionate to tissue damage and the connection between mast cell activation and small fiber neuropathy documented in Novak et al. 2022.

H.19.5 Nakamura et al. 2014 — Circadian Clock Regulates Mast Cell Function

Full Citation: Nakamura Y, Nakano N, Ishimaru K, et al. Circadian regulation of allergic reactions by the mast cell clock in mice. *J Allergy Clin Immunol.* 2014;133(2):568–575.

DOI: [10.1016/j.jaci.2013.07.040](https://doi.org/10.1016/j.jaci.2013.07.040)

PMID: 24060274

Published: February 2014

Study Design: Genetic mouse model with circadian clock manipulation

Key Findings:

- Mast cells possess intrinsic circadian clocks that drive time-of-day variation in degranulation
- *Clock* gene mutation abolished temporal variation in IgE-mediated responses (both in vivo and in vitro)
- FcεRI receptor expression and signaling show circadian rhythms in wild-type cells, lost in *Clock*-mutant
- Adrenalectomy disrupts mast cell clock rhythms — demonstrates adrenal hormones entrain mast cell clocks

Certainty: High for mechanism (elegant genetics, replicated); Low-Medium for ME/CFS relevance (extrapolation from mouse allergy model)

Clinical Relevance: Provides basis for “temporal priming hypothesis” — HPA axis dysfunction in ME/CFS could disrupt mast cell circadian regulation, leading to inappropriate activation timing. May explain symptom timing patterns, sleep-symptom interactions, and potential for chronotherapy (timing mast cell treatments to peak reactivity periods).

H.19.6 Steinberg et al. 1996 — Terfenadine Trial (Negative)

Full Citation: Steinberg P, McNutt BE, Marshall P, et al. A double-blind placebo-controlled study of the efficacy of oral terfenadine in the chronic fatigue syndrome. *J Allergy Clin Immunol.* 1996;97(1 Pt 1):119–126.

DOI: [10.1016/S0091-6749\(96\)80212-6](https://doi.org/10.1016/S0091-6749(96)80212-6)

PMID: 8568124

Published: January 1996

Study Design: Double-blind, placebo-controlled RCT

Sample Size: 30 CFS patients enrolled, 28 completed

Intervention: Terfenadine 60 mg twice daily for 8 weeks (H1 antihistamine only)

Results:

- **NO therapeutic benefit detected**
- No improvement in symptom amelioration

- No improvement in physical or social functioning
- No improvement in health perceptions or mental health
- Additional finding: 73% had atopy, 53% had positive immediate skin test results

Conclusion: “Terfenadine is unlikely to be of clinical benefit in treating CFS symptoms”

Certainty: High (well-designed RCT with negative results)

Clinical Implications: H1 antihistamine alone insufficient; suggests combination therapy (H1+H2 or H1+mast cell stabilizer) may be necessary

H.19.7 Davis et al. 2023 — Long COVID Case with H1/H2 Combination Success

Full Citation: Davis HE, McCorkell L, Vogel JM, Topol EJ. Case Study of ME/CFS Care Applied to Long COVID: Hypothesis Regarding Exercise Intolerance, Orthostatic Intolerance, Mast Cell Activation, Sleep Dysfunction, Neuropathy, and Viral Persistence. *Healthcare*. 2023;11(6):896.

DOI: [10.3390/healthcare11060896](https://doi.org/10.3390/healthcare11060896)

PMID: 36981567

PMCID: PMC10048325

Published: March 21, 2023

Study Type: Single case report (n=1)

Patient: Long COVID patient meeting ME/CFS criteria

Interventions and Outcomes: • **H1 blockers** (loratadine 10 mg OR fexofenadine 180 mg): “helpful with energy and cognitive dysfunction”

- **H2 blocker** (famotidine 40 mg BID): “helpful with energy and cognitive dysfunction”

- **Discontinuation test:** Stopping fexofenadine and famotidine → “increased fatigue and increased cognitive dysfunction, both of which improved rapidly upon resumption”

- **Cromolyn** (400 mg QID): Peak heart rate during walking fell from 130–140 bpm to 100–105 bpm

- **Quercetin** (1000 mg BID): “Improvement in fatigue and allergic symptoms”

Certainty: Low (n=1 case report, but dramatic response with discontinuation-rechallenge confirmation)

Clinical Relevance: Demonstrates potential for H1+H2 combination therapy; suggests mast cell-targeted approach may benefit post-viral fatigue syndromes

H.19.8 Theoharides et al. 2012 — Quercetin Superior to Cromolyn

Full Citation: Theoharides TC, Asadi S, Panagiotidou S. Quercetin in combination with IL-6 inhibits histamine and TNF release from mast cells through interaction with the IL-6 receptor. *PLOS ONE*. 2012;7(3):e33805.

DOI: [10.1371/journal.pone.0033805](https://doi.org/10.1371/journal.pone.0033805)

PMID: 22470478

PMCID: PMC3314669

Published: March 29, 2012

Study Design: In vitro comparison + clinical pilot trials

Concentration: Quercetin 100 μ M (approximated by 2 g/day oral dosing)

Key Findings:

- **IgE/Anti-IgE stimulation:** Quercetin inhibited histamine (82% vs 67%), PGD₂ (77% vs 75%), leukotrienes (99% vs 88%) comparably to cromolyn
- **Substance P stimulation:** Quercetin dramatically outperformed cromolyn — IL-8 reduced from 437.2 to 115.4 pg/mL (quercetin) vs 362.9 pg/mL (cromolyn)
- **Mechanism:** Quercetin worked prophylactically (30 min pre-stimulus); cromolyn required simultaneous addition
- **Clinical trial — Contact dermatitis:** Quercetin 2 g/day for 3 days reduced nickel patch reactions >50% in 8 of 10 patients; pruritus eliminated completely
- **Clinical trial — Photosensitivity:** Quercetin 1 g increased minimal erythema dose in all patients ($p=0.002$)

Certainty: Medium-High (strong in vitro data, pilot clinical success)

Clinical Relevance: Quercetin may be superior to prescription cromolyn for mast cell stabilization; available over-the-counter; well-tolerated

H.19.9 Clemons et al. 2011 — Amitriptyline Mast Cell Inhibition

Full Citation: Clemons A, Vasiadi M, Kempuraj D, et al. Amitriptyline and prochlorperazine inhibit proinflammatory mediator release from human mast cells: possible relevance to chronic fatigue syndrome. *J Clin Psychopharmacol.* 2011;31(3):385–387.

DOI: [10.1097/JCP.0b013e3182196e50](https://doi.org/10.1097/JCP.0b013e3182196e50)

PMID: 21532369

PMCID: PMC3498825

Published: June 2011

Study Design: In vitro study on human mast cells

Key Findings:

- Amitriptyline (AMI) and prochlorperazine (PRO) at 25 μ M significantly reduced IL-8, VEGF, and IL-6 release from stimulated human mast cells
- Bupropion, citalopram, and atomoxetine did NOT inhibit mast cells
- Mechanism involves modulation of intracellular calcium (FURA2 AM calcium indicator assays)
- AMI inhibits histamine release while permitting serotonin release

Conclusion: “The ability of amitriptyline, but not other antidepressants, to inhibit human mast cell release of pro-inflammatory cytokines may be relevant to their apparent benefit in CFS”

Certainty: Medium (mechanistic in vitro study, explains clinical observations)

Clinical Relevance: Amitriptyline’s benefit in ME/CFS may involve mast cell inhibition beyond pain/sleep effects; specific pharmacological mechanism

H.19.10 Rupatadine — Dual H1/PAF Antagonist with Mast Cell Stabilization

Full Citations: • Piñero-González J, et al. Rupatadine inhibits proinflammatory mediator secretion from human mast cells triggered by different stimuli. *J Investig Allergol Clin Immunol*. 2017;27(3):161–168. PMID: 19672095; PMCID: PMC7065400.

- Mullol J, Bousquet J, Bachert C, et al. Rupatadine in allergic rhinitis and chronic urticaria. *Allergy*. 2008;63(Suppl 87):5–28. PMID: 18339040.

Mechanism: Triple action — (1) H1 receptor antagonist, (2) PAF (platelet-activating factor) antagonist, (3) Direct mast cell stabilizer

Mast Cell Effects: • Rupatadine (10–50 μ M) inhibited IL-8 (80%), VEGF (73%), histamine (88%) release from LAD2 mast cell line

- Also inhibited IL-6, IL-8, IL-10, IL-13, and TNF release from human cord blood-derived cultured mast cells
- More effective than levocetirizine and desloratadine at PAF-induced mast cell inhibition

PAF Antagonism Potency: • Rupatadine IC₅₀ = 4.6 μ M (most potent)

- Loratadine IC₅₀ = 142 μ M (~31× less potent)
- Cetirizine IC₅₀ >200 μ M (>43× less potent)
- Fexofenadine IC₅₀ >200 μ M (>43× less potent)

Efficacy Ranking: Network meta-analysis for allergic rhinitis (SUCRA scores):

- Rupatadine 20 mg: 99.7% (highest rank)
- Rupatadine 10 mg: 76.3%
- Fexofenadine, cetirizine: moderate
- Loratadine 10 mg: lowest (inferior to all others)

Certainty: High (multiple RCTs, network meta-analysis, in vitro mechanistic data)

Clinical Relevance: Superior to standard H1 antihistamines; unique PAF antagonism may benefit ME/CFS patients with mast cell activation and vascular/orthostatic symptoms

Note: PAF is a key inflammatory mediator in ME/CFS contributing to vascular leakage, brain fog, and orthostatic issues

H.19.11 Moldofsky et al. 2015 — Ketotifen in Fibromyalgia (Negative)

Full Citation: Moldofsky H, Harris HW, Archambault WT, Kwong T, Lederman S. A randomized, double-blind, placebo-controlled Phase 1 trial of ketotifen in fibromyalgia. *J Rheumatol*. 2015;42(12):2505–2513.

DOI: [10.3899/jrheum.150460](https://doi.org/10.3899/jrheum.150460)

PMID: 26472411

PMCID: PMC4417653

Published: December 2015

Study Design: Phase 1 RCT, double-blind, placebo-controlled

Sample Size: 51 fibromyalgia patients (24 ketotifen, 27 placebo)

Intervention: Ketotifen 2 mg BID for 8 weeks (after 1-week titration)

Results: NO significant differences in primary outcomes:

- Pain intensity: ketotifen −1.3 vs placebo −1.5 ($p=0.7$)
- FIQR scores: −12.1 vs −12.2 ($p=0.9$)
- Side effect: Transient sedation 28.6% vs 4%

Certainty: High (well-designed RCT showing no benefit)

Clinical Relevance: Mast cell stabilization alone may not address core pathophysiology in central pain syndromes like fibromyalgia; relevance to ME/CFS unclear

Note: Despite this negative finding, retrospective ME/CFS study (not included here) showed 77% of continuers had significant PEM reduction with ketotifen

H.20 Blood Volume and Cardiovascular Dysfunction

H.20.1 Hypovolemia and RAAS Dysfunction

Chronic hypovolemia (reduced blood volume) is a well-documented feature of ME/CFS, with direct consequences for oxygen delivery, exercise capacity, and orthostatic symptoms. Paradoxically, the renin-angiotensin-aldosterone system (RAAS) and antidiuretic hormone (ADH) — which normally activate in response to low blood volume — show suppression in ME/CFS patients.

Miwa & Fujita 2017 [674]: This study identified paradoxical down-regulation of volume-regulatory hormones in ME/CFS. Despite documented hypovolemia and reduced cardiac output, plasma aldosterone was 33% lower (104 ± 37 vs 157 ± 67 pg/ml, $p=0.004$) and ADH was 33% lower (2.2 ± 1.0 vs 3.3 ± 1.5 pg/ml, $p=0.02$) compared to healthy controls ($n=14$ patients, $n=13$ controls). Treatment trial: desmopressin (ADH analog) improved orthostatic symptoms in 50% of patients. **Certainty:** Medium (peer-reviewed, significant findings, but small sample awaiting replication). **Implication:** Hypovolemia results from central dysregulation of volume-regulatory systems, not excessive fluid loss.

Raj et al. 2005 [325]: Landmark study demonstrating the “renin-aldosterone paradox” in postural tachycardia syndrome (POTS), a condition overlapping with ME/CFS. Blood volume was markedly reduced (3583 ± 579 vs 4319 ± 578 mL, $p<0.0001$), with plasma volume 21% lower (2172 ± 429 vs 2763 ± 437 mL, $p<0.0001$). Despite this, plasma renin activity was unchanged and aldosterone was frankly low (7.0 ± 5.3 vs 12.3 ± 6.4 ng/dL, $p=0.01$). Strong positive correlation between blood volume and aldosterone ($r=0.56$, $p=0.001$). $n=33$ POTS patients, $n=13$ controls. **Certainty:** High (large sample, replicated, published in *Circulation*).

Mustafa et al. 2011 [675]: Identified abnormalities in angiotensin II regulation in POTS. Plasma Ang II was significantly elevated (43 ± 3 vs 28 ± 3 pg/mL, $p=0.006$), while estimated ACE2 activity was reduced (0.25 ± 0.02 vs 0.33 ± 0.03 , $p=0.038$). Elevated Ang II may contribute to peripheral vasoconstriction and reduced NO bioavailability. **Certainty:** Medium-High (peer-reviewed, mechanistic insight into RAAS dysfunction).

Stewart et al. 2006 [333]: Increased plasma angiotensin II in low-flow POTS patients related to reduced blood flow and blood volume. Suggests Ang II elevation is compensatory attempt to maintain blood pressure despite hypovolemia, but may contribute to local blood flow dysregulation. **Certainty:** Medium (mechanistic study linking Ang II to blood volume deficit).

Farquhar et al. 2002 [676]: Early study demonstrating relationship between blood volume and exercise capacity in CFS. Patients had significantly lower peak VO₂ consumption with trend toward lower blood volume. Strong correlation between blood volume and peak oxygen consumption, suggesting hypovolemia as physiological contributor to exercise intolerance. **Certainty:** Medium (established blood volume-exercise link).

van Campen et al. 2018 [340]: Dual-isotope blood volume measurement in ME/CFS adults. Mean absolute blood volume was 59(8) ml/kg, representing -11(7) ml/kg deficit below reference values. Blood volume reduction correlated with presence of orthostatic intolerance symptoms (n=20). **Certainty:** High (precise measurement technique, clear clinical correlation).

H.20.2 Cardiac Dysfunction and Natriuretic Peptides

Newton et al. 2016 [324]: CFS patients had significantly reduced cardiac volumes (both end-systolic and end-diastolic) with reduced end-diastolic wall masses. Strong positive correlations between total blood volume, red cell volume, plasma volume and cardiac end-diastolic wall mass. Critically, no relationship between disease length and cardiac/plasma volumes, ruling out deconditioning as sole cause. **Certainty:** High (cardiac MRI, objective measures, n=42 CFS patients). **Implication:** Reduced cardiac volumes are primary feature, not secondary to inactivity.

Tomas et al. 2017 [326]: Brain natriuretic peptide (BNP) levels significantly elevated in CFS cohort ($p=0.013$). Patients with high BNP (>400 pg/mL) had significantly lower cardiac volumes in both end-systolic and end-diastolic measurements ($p=0.05$). BNP elevation associated with cardiac dysfunction, not just volume overload. **Certainty:** Medium-High (established biomarker, cardiac imaging correlation).

H.20.3 Endothelial Dysfunction and Vascular Pathology

Scherbakov et al. 2020 [177]: Peripheral endothelial dysfunction found in 51% of ME/CFS patients vs 20% of healthy controls ($p<0.05$). Endothelial dysfunction assessed via flow-mediated dilation. Associated with disease severity and severity of immune symptoms (n=35 patients). **Certainty:** High (objective vascular measurement, peer-reviewed, ESC Heart Failure). **Implication:** Vascular pathology contributes to reduced blood flow and tissue perfusion.

Appel et al. 2024 [328]: Comprehensive review of endothelial dysfunction in ME/CFS. Elevated adhesion molecules (ICAM-1, VCAM-1), impaired flow-mediated dilation, chronic inflammatory state contributing to vascular pathology. Links endothelial dysfunction to exercise intolerance and post-exertional symptoms. **Certainty:** High (systematic review, multiple lines of evidence).

Miller et al. 2020 [334]: Arterial elasticity significantly increased in Ehlers-Danlos syndrome patients. Central pulse wave velocity significantly lower in EDS (4.73 m/s vs controls), indicating increased arterial elasticity that impairs baroreceptor-mediated blood pressure control. Explains orthostatic intolerance mechanism in EDS and related hypermobility conditions. n=46 EDS patients across multiple subtypes (primarily hEDS). **Certainty:** High (objective arterial measurements, published in *Genes*, significant finding). **Relevance to ME/CFS:** High comorbidity between ME/CFS and hEDS/hypermobility spectrum disorders. Increased arterial compliance reduces effectiveness of baroreceptor responses, contributing to orthostatic intolerance and POTS. Provides vascular mechanism linking connective tissue disorders to autonomic symptoms common in ME/CFS population.

H.20.4 Erythropoiesis and Red Blood Cell Function

Streiten DHP, Bell DS. 1998 [280]: Landmark early study measuring circulating blood volume in CFS patients using radiolabeled RBC and plasma volume techniques. Found: red blood cell mass reduced in 93.8% of female and 50% of male ME/CFS patients; plasma volume subnormal in 52.6%. Documented that blood volume deficits were consistent and substantial, with clear correlation to orthostatic intolerance symptoms. Provided first objective evidence that hypovolemia (not just deconditioning) contributes to exercise intolerance and symptom severity in ME/CFS. **Certainty:** High (gold-standard blood volume measurement technique, clear patient selection, objective methodology). **Clinical Implication:** Hypovolemia is a primary physiological feature of ME/CFS, not secondary consequence.

Saha et al. 2019 [180]: Red blood cell deformability significantly reduced in CFS patients using microfluidic measurements. Impaired RBC deformability can impair oxygen delivery to tissues and contribute to exercise intolerance and fatigue. n=20 CFS, n=20 controls. **Certainty:** Medium-High (novel methodology, replicated findings, published in clinical journal). **Implication:** Even with adequate RBC count, oxygen delivery may be compromised.

Winkler et al. 2004 [677]: Evaluation of serum erythropoietin levels and autonomic function in CFS. Examined potential relationships with anemia and fatigue severity. **Certainty:** Medium (exploratory study).

Świątczak et al. 2022 [678]: CFS patients show deteriorated iron metabolism: low serum iron, elevated ferritin, reduced transferrin saturation — pattern consistent with inflammatory anemia. Not true iron deficiency but iron sequestration due to inflammation. **Certainty:**

Medium-High (clear pattern, n=multiple cohorts). **Link to cytokines:** IL-6 and hepcidin drive iron restriction.

Morceau et al. 2009 [679]: Mechanistic review: pro-inflammatory cytokines (IL-1, IL-6, TNF- α , IFN- γ) suppress erythropoiesis via multiple pathways including hepcidin induction, direct EPO suppression, and shortened RBC lifespan. **Relevance to ME/CFS:** Elevated cytokines documented in ME/CFS may contribute to functional anemia.

McCranor et al. 2014 [680]: IL-6 directly impairs erythroid differentiation in vitro, providing mechanistic link between cytokine elevation and anemia of chronic disease. **Certainty:** High (mechanistic study, controlled conditions).

Fraenkel 2017 [681]: Comprehensive review of anemia of inflammation: cytokine-mediated suppression of erythropoiesis, hepcidin-induced iron restriction, shortened RBC survival. **Application to ME/CFS:** Framework for understanding functional anemia despite normal hemoglobin in some patients.

H.20.5 Integrated Mechanisms: The Hypovolemia Cascade

The blood volume deficit in ME/CFS results from convergent mechanisms:

1. **RAAS/ADH suppression:** Paradoxical down-regulation prevents compensatory volume retention (Miwa 2017, Raj 2005)
2. **Plasma volume reduction:** Primary deficit in fluid compartment (Raj 2005: 21% reduction; van Campen 2018: -11 ml/kg)
3. **Cardiac consequences:** Reduced preload → reduced cardiac output → exercise intolerance (Newton 2016)
4. **Endothelial dysfunction:** Impaired vascular regulation → tissue hypoperfusion (Scherbakov 2020)
5. **RBC dysfunction:** Reduced deformability + inflammatory anemia → impaired oxygen delivery (Saha 2019, Świątczak 2022)
6. **Cytokine-mediated effects:** IL-6 and other cytokines suppress erythropoiesis and sequester iron (Morceau 2009, McCranor 2014)

This multi-hit model explains why simple volume expansion (saline infusion) provides only temporary benefit: underlying regulatory systems remain dysfunctional.

Clinical Implications:

- **Diagnostics:** Dual-isotope blood volume measurement may identify hypovolemic subgroup

- **Treatment targets:** Desmopressin for ADH-deficient patients (Miwa 2017); fludrocortisone for aldosterone supplementation; management of endothelial dysfunction; optimization of iron availability despite inflammation
- **Subtype identification:** Not all ME/CFS patients show same degree of hypovolemia; responders to volume-expanding interventions may represent distinct subgroup

Research Gaps:

- Mechanism of RAAS/ADH suppression (central dysregulation? autoimmune?)
- Predictors of desmopressin response
- Longitudinal blood volume changes over disease course
- Relationship between blood volume deficit and PEM severity
- Role of capillary permeability in plasma volume loss

H.21 Additional Key Resources

H.21.1 Patient Advocacy and Information

MEpedia: <https://me-pedia.org/> — Comprehensive patient-edited wiki on ME/CFS.

ME Association (UK): <https://meassociation.org.uk/> — Patient support and research summaries.

Bateman Horne Center: <https://batemanhornecenter.org/> — Clinical and educational resources.

Open Medicine Foundation: <https://www.openmedicinefoundation.ngo/> — Research funding and updates.

Solve ME/CFS Initiative: <https://solvecfs.org/> — US-based research and advocacy.

H.21.2 Research Centers

Cornell Center for Enervating NeuroImmune Disease: <https://neuroimmune.cornell.edu/>

Griffith University National Centre for Neuroimmunology and Emerging Diseases:
Queensland, Australia

Charité Fatigue Center: Berlin, Germany

Stanford ME/CFS Initiative: Stanford University, California

H.22 Neuroimaging and Structural Brain Findings

H.22.1 Shan et al. 2020 — Neuroimaging Systematic Review (JTM)

Full Citation: Shan ZY, Barnden LR, Kwiatek RA, Bhuta S, Hermens DF, Lagopoulos J. Neuroimaging characteristics of myalgic encephalomyelitis/chronic fatigue syndrome

(ME/CFS): a systematic review. *Journal of Translational Medicine*. 2020;18(1):335.

DOI: [10.1186/s12967-020-02506-6](https://doi.org/10.1186/s12967-020-02506-6)

PMID: 32887623

Study Design: Systematic review of structural and functional neuroimaging studies in ME/CFS

Key Findings:

- Inconsistent but recurring structural brain abnormalities across cohorts
- Prefrontal myelination upregulation reported in some studies
- Basal ganglia metabolite differences (elevated choline) and functional changes
- Insula gray matter volume increases in some studies
- Widespread functional connectivity disruptions across networks

Limitations: High heterogeneity across studies; inconsistent findings prevent definitive regional conclusions.

H.22.2 Finkelmeyer et al. 2017 — VBM Gray and White Matter in CFS

Full Citation: Finkelmeyer A, He J, Maclachlan L, Watson S, Gallagher P, Newton JL, Blamire AM. Grey and white matter differences in Chronic Fatigue Syndrome — A voxel-based morphometry study. *NeuroImage: Clinical*. 2017;17:24–30.

DOI: [10.1016/j.nicl.2017.09.024](https://doi.org/10.1016/j.nicl.2017.09.024)

PMID: 29021956

Sample Size: 42 CFS patients, 30 healthy controls; 3T MRI

Key Findings:

- Increased gray matter volume in the amygdala and insula (not reductions as initially hypothesized)
- Reduced white matter volume in midbrain, pons, and right temporal lobe
- Insula changes consistent with altered interoceptive processing
- Brainstem white matter reductions suggest pathway disruption

Note: Direction of regional changes contrasts with some earlier studies; illustrates the heterogeneity of neuroimaging findings in ME/CFS.

H.22.3 Puri et al. 2012 — VBM Regional Gray/White Matter Changes

Full Citation: Puri BK, Jakeman PM, Agour M, et al. Regional grey and white matter volumetric changes in myalgic encephalomyelitis (chronic fatigue syndrome): a voxel-based morphometry 3 T MRI study. *British Journal of Radiology*. 2012;85(1015):e270–e273.

DOI: [10.1259/bjr/93889091](https://doi.org/10.1259/bjr/93889091)

PMID: 22128128

Key Findings:

- Reduced gray matter volume in bilateral occipital poles, lateral occipital cortex, right angular gyrus, and posterior parahippocampal gyrus
- White matter reductions in the left occipital lobe
- Parahippocampal changes consistent with the memory impairment characteristic of ME/CFS

H.22.4 de Lange et al. 2004 — fMRI Neural Correlates of CFS

Full Citation: de Lange FP, Kalkman JS, Bleijenberg G, Hagoort P, van der Werf SP, van der Meer JWM, Toni I. Neural correlates of the chronic fatigue syndrome—an fMRI study. *Brain*. 2004;127(9):1948–1957.

DOI: [10.1093/brain/awh225](https://doi.org/10.1093/brain/awh225)

PMID: 15240435

Sample Size: 16 CFS patients, 16 healthy controls

Key Findings:

- CFS patients recruited additional or atypical brain regions during motor imagery tasks
- Stronger responses in visually related structures during motor tasks
- Error-related brain activity differed between groups, suggesting motor planning dysfunction
- Pattern consistent with compensatory over-recruitment and inefficient neural processing

H.23 Neurotransmitter and Neurochemical Findings

H.23.1 Yamamoto et al. 2004 — Serotonin Transporter Reduction in CFS

Full Citation: Yamamoto S, Ouchi Y, Onoe H, et al. Reduction of serotonin transporters of patients with chronic fatigue syndrome. *NeuroReport*. 2004;15(17):2571–2574.

DOI: [10.1097/00001756-200412030-00002](https://doi.org/10.1097/00001756-200412030-00002)

PMID: 15570154

Key Findings:

- PET imaging (DASB ligand) showed significantly reduced serotonin transporter density in CFS
- Reduction was most pronounced in the rostral subdivision of the anterior cingulate cortex (ACC)
- Demonstrates in vivo serotonergic dysfunction in ME/CFS using direct neuroimaging
- Provides neurobiological basis for sleep, mood, and cognitive symptoms in ME/CFS

H.23.2 Miller et al. 2014 — Basal Ganglia Activation and Dopamine in CFS

Full Citation: Miller AH, Jones JF, Drake DF, Tian H, Unger ER, Pagnoni G. Decreased Basal Ganglia Activation in Subjects with Chronic Fatigue Syndrome: Association with Symptoms of Fatigue. *PLoS One*. 2014;9(5):e98156.

DOI: [10.1371/journal.pone.0098156](https://doi.org/10.1371/journal.pone.0098156)

PMID: 24858857

Sample Size: 18 CFS patients, 41 healthy controls

Key Findings:

- Significantly reduced right caudate nucleus and globus pallidus activation during reward tasks

- Diminished globus pallidus responsivity correlated with mental fatigue ($r^2 = 0.49$, $p = 0.001$)
- Proposes inflammatory cytokine-mediated disruption of basal ganglia dopamine availability
- Supports dopaminergic dysfunction as a contributor to ME/CFS fatigue

H.23.3 Dehhaghi et al. 2022 — Kynurenine Pathway and NAD⁺ in ME/CFS

Full Citation: Dehhaghi M, Panahi HKS, Kavyani B, et al. The Role of Kynurenine Pathway and NAD⁺ Metabolism in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Aging and Disease*. 2022;13(3):698–711.

DOI: [10.14336/AD.2021.0824](https://doi.org/10.14336/AD.2021.0824)

PMID: 35656104

Key Findings:

- Under inflammatory/IDO-1 activation, up to 90% of tryptophan is catabolized through the kynurenine pathway
- Pro-inflammatory cytokines (IFN- γ , TNF- α) documented in ME/CFS drive IDO-1 activity
- Kynurenine pathway overactivation depletes NAD⁺ and produces neurotoxic quinolinic acid accumulation
- Mechanistic explanation for reduced serotonin precursor availability and neurotransmitter imbalances in ME/CFS

H.24 Prognosis and Outcomes

H.24.1 Joyce, Hotopf and Wessely 1997 — Prognosis of CFS: Systematic Review

Full Citation: Joyce J, Hotopf M, Wessely S. The prognosis of chronic fatigue and chronic fatigue syndrome: a systematic review. *QJM: An International Journal of Medicine*. 1997;90(3):223–233.

DOI: [10.1093/qjmed/90.3.223](https://doi.org/10.1093/qjmed/90.3.223)

PMID: 9093600

Key Findings:

- **Pediatric:** 54–94% of children made good or complete recovery at 13–72 months (4 studies, $n = 15$ –31)
- **Adults:** Recovery rates substantially lower; improvement common but full recovery rare
- Recovery definition heterogeneity limits cross-study comparison
- Dramatically better pediatric outcomes support early and aggressive intervention

Significance: The 54–94% pediatric recovery range from this 1997 review remains the most cited figure in guidelines on pediatric ME/CFS prognosis.

H.24.2 Cairns and Hotopf 2005 — Adult CFS Prognosis: Systematic Review

Full Citation: Cairns R, Hotopf M. A systematic review describing the prognosis of chronic fatigue syndrome. *Occupational Medicine*. 2005;55(1):20–31.

DOI: [10.1093/occmed/kqi013](https://doi.org/10.1093/occmed/kqi013)

PMID: 15699087

Study Design: Systematic review of 14 studies meeting operational CFS criteria

Key Findings:

- Median full recovery rate: 5% (range 0–31%)
- Median proportion improving during follow-up: 39.5% (range 8–63%)
- Full recovery is rare in adult ME/CFS; sustained improvement without full recovery is more common
- Prognostic factors: illness beliefs, coping style, psychiatric comorbidity

Significance: Demonstrates the striking contrast with pediatric outcomes (54–94% recovery vs. 5% median in adults), supporting the urgency of early treatment in newly diagnosed patients.

H.25 Orthostatic Intolerance

H.25.1 Schondorf and Freeman 1999 — Importance of OI in CFS

Full Citation: Schondorf R, Freeman R. The importance of orthostatic intolerance in the chronic fatigue syndrome. *American Journal of Medical Sciences*. 1999;317(2):117–123.

DOI: [10.1097/00000441-199902000-00006](https://doi.org/10.1097/00000441-199902000-00006)

PMID: 10037115

Key Findings:

- Established orthostatic intolerance as a frequent and clinically important feature of CFS
- Demonstrates treatable autonomic dysfunction underlying many CFS symptoms
- Prevalence estimates across studies range from 50–97% depending on testing method and population
- Supports universal OI screening as a standard component of ME/CFS clinical evaluation

H.26 Treatment Evidence

H.26.1 Li et al. 2022 — tDCS of DLPFC for Neuropsychiatric Disorders

Full Citation: Li Q, Fu Y, Liu C, Meng Z. Transcranial Direct Current Stimulation of the Dorsolateral Prefrontal Cortex for Treatment of Neuropsychiatric Disorders. *Frontiers in Behavioral Neuroscience*. 2022;16:893955.

DOI: [10.3389/fnbeh.2022.893955](https://doi.org/10.3389/fnbeh.2022.893955)

Key Findings:

- DLPFC is central to attention, decision-making, working memory, and executive function

- Anodal tDCS over left DLPFC improves working memory and attentional control in multiple studies
- Mechanism: modulates neuronal excitability of primary executive control regions
- Provides rationale for tDCS as a cognitive enhancement strategy in conditions of executive dysfunction

Note: Evidence base is primarily from depression and healthy volunteer studies; direct ME/CFS tDCS trials are limited.

H.26.2 Lang-Illievich et al. 2023 — PEA for Chronic Pain: Systematic Review and Meta-Analysis

Full Citation: Lang-Illievich K, Klivinyi C, Lasser C, et al. Palmitoylethanolamide in the Treatment of Chronic Pain: A Systematic Review and Meta-Analysis of Double-Blind Randomized Controlled Trials. *Nutrients*. 2023;15(6):1350.

DOI: [10.3390/nu15061350](https://doi.org/10.3390/nu15061350)

PMID: 36986081

Study Design: Systematic review and meta-analysis of 13 double-blind RCTs

Key Findings:

- Pooled effect favors PEA with mean pain reduction of 1.68 points on 11-point scale
- Pain reductions documented at 6 weeks, 8 weeks, and 24–26 weeks, supporting extended treatment
- Safety profile excellent; minimal side effects across all trials
- Superiority of micronized vs. standard PEA on clinical outcomes unclear; enhanced solubility may affect absorption

Significance: Provides the strongest meta-analytic evidence for PEA in chronic pain, supporting its use in ME/CFS patients with a pain component.

H.27 Coagulation and Vascular Pathology

H.27.1 Nunes et al. 2022 — Hyperactivated Platelets and Fibrinoid Microclots in ME/CFS

Full Citation: Nunes JM, Kruger A, Proal A, Kell DB, Pretorius E. The Occurrence of Hyperactivated Platelets and Fibrinoid Microclots in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS). *Pharmaceuticals (Basel)*. 2022;15(8):931.

DOI: [10.3390/ph15080931](https://doi.org/10.3390/ph15080931)

PMID: 36015078

Study Design: Case-control study; ME/CFS patients vs. healthy controls; thromboelastography + fluorescent platelet microscopy

Key Findings:

- Fibrinoid microclot burden >10-fold greater in ME/CFS plasma than healthy controls
- ~80% of ME/CFS participants showed platelet hyperactivation (mean spreading score 2.72 vs. 1.00)

- Thromboelastography revealed hypercoagulable state in ~50% of participants
- Amyloid-resistant microclots may occlude microcapillaries, impairing tissue perfusion

Significance: First direct demonstration of ME/CFS-specific coagulation pathology paralleling Long COVID findings; opens potential therapeutic avenue via anticoagulation.

H.27.2 Suárez et al. 2010 — Nitric Oxide Metabolites During Exercise in CFS

Full Citation: Suárez A, Guillamó E, Roig T, et al. Nitric oxide metabolite production during exercise in chronic fatigue syndrome: a case-control study. *Journal of Women's Health*. 2010;19(6):1073–1077.

DOI: [10.1089/jwh.2008.1255](https://doi.org/10.1089/jwh.2008.1255)

PMID: 20469961

Study Design: Case-control; 44 female CFS patients vs. 25 healthy controls; maximal exercise testing

Key Findings:

- Plasma nitrate concentrations significantly higher in CFS patients during exercise
- Maximum difference ~295% above controls at maximal intensity
- Elevated NO metabolite response may indicate abnormal vascular regulation, not simple deficiency
- Authors suggest exercise-induced NO measurement as potential CFS biomarker

Significance: Demonstrates abnormal NO metabolism during exertion; challenges simple “endothelial NO deficiency” narrative; consistent with dysregulated vasoregulation rather than reduced production.

H.28 Lifestyle Intervention Safety

H.28.1 Kindlon 2011 — Reporting of Harms from GET and CBT in ME/CFS

Full Citation: Kindlon T. Reporting of harms associated with graded exercise therapy and cognitive behavioural therapy in myalgic encephalomyelitis/chronic fatigue syndrome. *Bulletin of the IACFS/ME*. 2011;19(2):59–111.

Key Findings:

- Systematic review of 10 patient surveys from four countries
- 51% of respondents reported GET worsened their health
- 20% reported similar worsening from CBT
- Critical methodological failures in RCTs: adverse events rarely tracked, patient compliance issues, subjective outcome bias

Significance: Foundational document establishing the evidence base for GET’s harm profile in ME/CFS; directly influenced subsequent NICE guideline revision in 2021 removing GET as a recommended intervention.

H.29 Adenosine, Sleep Pressure, and Neuroinflammation

H.29.1 Huang et al. 2024 — Adenosine Receptors in Sleep Regulation

Full Citation: Huang L, Zhu W, Li N, Zhang B, Dai W, Li S, Xu H. Functions and mechanisms of adenosine and its receptors in sleep regulation. *Sleep Medicine*. 2024;115:210–217. DOI: 10.1016/j.sleep.2024.02.012.

Key Findings:

- Adenosine promotes sleep by inhibiting arousal systems and activating sleep-promoting systems via A1 and A2A receptor subtypes
- Astrocyte-derived adenosine is the primary source of homeostatic sleep pressure signal
- Caffeine's wake-promoting effect is mediated entirely through adenosine receptor blockade
- A1 receptors inhibit wake-active neurons; A2A receptors in the nucleus accumbens core drive sleep pressure

Relevance: Establishes the mechanistic framework for adenosine as homeostatic sleep signal. Forms the foundational reference for Section 15.3 (*Adenosine Accumulation and Pathological Sleep Pressure*).

Certainty Assessment:

- **Quality:** High (peer-reviewed review, *Sleep Medicine*)
- **Study type:** Narrative/mechanistic review
- **ME/CFS specificity:** None — establishes normal physiology
- **Limitation:** No primary data; no ME/CFS cohort

H.29.2 Rétey et al. 2007 — ADORA2A Genotype and Caffeine Sensitivity

Full Citation: Rétey JV, Adam M, Khatami R, Luhmann UFO, Jung HH, Berger W, Landolt HP. A genetic variation in the adenosine A2A receptor gene (ADORA2A) contributes to individual sensitivity to caffeine effects on sleep. *Clinical Pharmacology & Therapeutics*. 2007;81(5):692–698. DOI: 10.1038/sj.clpt.6100102.

Key Findings:

- ADORA2A c.1083T>C genotype distribution differs significantly between caffeine-sensitive and caffeine-insensitive adults
- Specific ADORA2A genotypes determine how closely caffeine-induced EEG changes resemble insomnia patterns
- Caffeine sensitivity is a functional readout of A2A receptor sensitivity at the individual level

Relevance: Provides the pharmacogenetic basis for understanding ME/CFS patients' variable caffeine responses as a potential marker of A2A receptor function. Supports the speculation that ME/CFS-associated A2A upregulation could alter caffeine pharmacodynamics.

- Certainty Assessment:**
- **Quality:** High (Clinical Pharmacology & Therapeutics, primary human study)
 - **ME/CFS specificity:** None — general population caffeine-sleep study
 - **Limitation:** No ME/CFS cohort; ME/CFS-specific caffeine-ADORA2A study remains unpublished

H.29.3 Orr et al. 2009 — A2A Receptor Mediates Microglial Process Retraction

Full Citation: Orr AG, Orr AL, Li X-J, Gross RE, Traynelis SF. Adenosine A2A receptor mediates microglial process retraction. *Nature Neuroscience*. 2009;12(7):872–878. DOI: 10.1038/nn.2341.

- Key Findings:**
- A2A adenosine receptors are upregulated on microglia coincident with P2Y12 downregulation during brain inflammation
 - This receptor shift produces chemotactic reversal: activated microglia retract processes in response to adenosine (the ATP breakdown product) rather than extending toward it
 - A2A signaling operates through Gs-adenylate cyclase-PKA pathway to drive amoeboid microglial morphology
 - Human microglia show the same receptor expression pattern as mouse models

Relevance: Seminal mechanistic paper establishing the A2A upregulation-neuroinflammation link. Directly supports the hypothesis in Section 15.3 that ME/CFS neuroinflammation maintains elevated A2A receptor density in sleep-regulatory brain regions.

- Certainty Assessment:**
- **Quality:** High (Nature Neuroscience, top-tier journal)
 - **Study type:** Primary mechanistic (mouse + human cell data)
 - **ME/CFS specificity:** None — mechanism study
 - **Limitation:** In vitro/animal model; extrapolation to ME/CFS requires supporting in vivo evidence

H.29.4 Navia et al. 2020 — Adenosine Receptors as Neuroinflammation Modulators

Full Citation: Navia M, et al. Adenosine Receptors as Neuroinflammation Modulators: Role of A1 Agonists and A2A Antagonists. *Cells*. 2020;9(7):1739. DOI: 10.3390/cells9071739.

- Key Findings:**
- A1 receptor agonism prevents cytokine (TNF- α , IL-1 β , IFN- γ) induced inflammation in mixed glial cells
 - A2A receptor antagonism shows superior anti-inflammatory and antioxidant properties vs. A1 agonism
 - A2A antagonist is effective in both in vitro glial models and in vivo neuroinflammation animal models
 - Authors conclude A2A antagonism is a candidate strategy for chronic neuroinflammation therapy

Relevance: Provides mechanistic rationale for A2A receptor-targeted interventions in conditions with chronic glial activation, such as ME/CFS.

- Certainty Assessment:**
- **Quality:** Medium (Cells journal, in vitro + animal; no human trial)
 - **ME/CFS specificity:** None
 - **Limitation:** No clinical data; compounds tested not yet in ME/CFS trials

H.29.5 Chang et al. 2021 — Dysregulated Adenosine Homeostasis in Brain Disorders

Full Citation: Chang C-P, Wu K-C, Lin C-Y, Chern Y. Emerging roles of dysregulated adenosine homeostasis in brain disorders with a specific focus on neurodegenerative diseases. *Journal of Biomedical Science*. 2021;28(1):70. DOI: 10.1186/s12929-021-00766-y.

- Key Findings:**
- Adenosine homeostasis is regulated by ectonucleotidases (CD39, CD73), adenosine deaminase (ADA), and equilibrative nucleoside transporters (ENT1/ENT2)
 - Dysregulation of any component leads to aberrant extracellular adenosine accumulation
 - In neuroinflammatory and neurodegenerative conditions, impaired clearance through glial transporters is a consistent finding
 - Adenosine functions as an endogenous anti-inflammatory agent; its dysregulation sustains microglial activation in a positive feedback loop

Relevance: Establishes the molecular machinery of adenosine clearance relevant to Section 15.3. ME/CFS reactive astrogliosis would be predicted to impair ENT1/ENT2-mediated clearance, raising basal adenosine.

- Certainty Assessment:**
- **Quality:** High (Journal of Biomedical Science, comprehensive review)
 - **ME/CFS specificity:** None — focused on neurodegeneration
 - **Limitation:** Mechanistic extrapolation to ME/CFS; no direct measurement of ENT expression in ME/CFS glia

H.29.6 Rábago-Monzón et al. 2025 — Astrocytes, Microglia, and Sleep Dysregulation

Full Citation: Rábago-Monzón ÁR, Osuna-Ramos JF, Armienta-Rojas DA, et al. Stress-Induced Sleep Dysregulation: The Roles of Astrocytes and Microglia in Neurodegenerative and Psychiatric Disorders. *Biomedicines*. 2025;13(5):1121. DOI: 10.3390/biomedicines13051121.

- Key Findings:**
- Astrocytes regulate sleep architecture via two primary mechanisms: adenosine signaling and glymphatic clearance
 - Chronic stress disrupts both pathways, reducing restorative sleep quality
 - Microglial neuroinflammation further degrades sleep quality through synaptic dysfunction
 - Convergence of adenosine dysregulation and glymphatic failure on synaptic pruning and plasticity is documented

Relevance: Directly links the glial biology of ME/CFS (neuroinflammation, reactive astrogliosis) to the sleep complaints that are central to the disease. Supports the integration of Section 15.3 with Section 15.10 (glymphatic failure).

Certainty Assessment: • **Quality:** Medium (Biomedicines, mechanistic synthesis review; 2025)

• **ME/CFS specificity:** None

• **Limitation:** Synthesis paper; primary data for ME/CFS application is indirect

H.29.7 Nakatomi et al. 2014 — TSPO-PET Neuroinflammation in ME/CFS

Full Citation: Nakatomi Y, Mizuno K, Ishii A, et al. Neuroinflammation in Patients with Chronic Fatigue Syndrome/Myalgic Encephalomyelitis: An ¹¹C-(R)-PK11195 PET Study. *Journal of Nuclear Medicine*. 2014;55(6):945–950. DOI: 10.2967/jnumed.113.131045.

Key Findings: • TSPO binding (microglial activation marker) elevated 45–199% in ME/CFS patients vs. healthy controls

- Affected regions: cingulate cortex, hippocampus, amygdala, thalamus, midbrain, pons
- Regional neuroinflammation correlated with cognitive impairment, pain, and depression severity
- First direct in vivo neuroinflammation evidence in ME/CFS via PET imaging

Relevance: Landmark study providing the neuroinflammatory substrate that mechanistically supports A2A upregulation and adenosine dysregulation in ME/CFS. The affected brain regions overlap substantially with sleep-regulatory circuitry (cingulate, thalamus, brainstem).

Certainty Assessment: • **Quality:** High (Journal of Nuclear Medicine; in vivo patient data)

• **Sample:** n=9 ME/CFS patients, n=10 healthy controls

• **Replication:** Partially replicated by subsequent PET and MRS studies (Younger group)

• **Limitation:** Small sample; single centre; TSPO binding is a marker of glial activation, not exclusive to microglia

H.29.8 Maksoud et al. 2021 — Systematic Review of Sleep in ME/CFS

Full Citation: Maksoud R, Eaton-Fitch N, Matula M, Cabanas H, Staines D, Marshall-Gradisnik S. Systematic Review of Sleep Characteristics in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Healthcare*. 2021;9(5):568. DOI: 10.3390/healthcare9050568.

Key Findings: • Non-restorative sleep reported by approximately 91% of ME/CFS patients

- Increased microarousal index (MAI) is the single most consistent polysomnographic abnormality: elevated in all five studies that measured it
- Sleep onset latency is uniformly normal across 13 studies — sleep initiation is intact
- Major discrepancy between subjective sleep complaints and aggregate polysomnographic measures

- Pattern suggests pathology in sleep maintenance and quality, not in sleep initiation

Relevance: Provides the clinical sleep phenotype that the adenosine hypothesis in Section 15.3 must explain. The preserved sleep onset latency with elevated microarousal index is mechanistically consistent with adenosine-mediated sleep maintenance disruption rather than a simple sleep pressure deficit.

Certainty Assessment: • **Quality:** High (systematic review, Healthcare)

- **Study type:** Systematic review of polysomnographic studies
- **Limitation:** Heterogeneous populations and methods across included studies; no biomarker data linking sleep findings to adenosine specifically

Section 15.6 — Melatonin Dysfunction and Circadian Disruption

McCarthy 2022 — Circadian Rhythm Disruption in ME/CFS and Long COVID

M. J. McCarthy. “Circadian rhythm disruption in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Implications for the post-acute sequelae of COVID-19”. In: *Brain, Behavior, & Immunity – Health* 20 (2022). Narrative review linking circadian disruption to ME/CFS and long COVID pathophysiology. Key finding: ME/CFS patients lack the midday temperature increase seen in controls and show distinct evening temperature drop, indicating multi-system circadian decoupling. DLMO timing is dissociated from temperature rhythm (intact in controls, absent in ME/CFS). TGFB dysregulation proposed as mechanistic driver of peripheral clock disruption. Melatonin effective only in phase-delayed subset., p. 100412. DOI: [10.1016/j.bbih.2022.100412](https://doi.org/10.1016/j.bbih.2022.100412)

Key Findings: • ME/CFS patients lack the midday temperature increase seen in healthy controls and show a distinct evening temperature drop, indicating multi-system circadian decoupling beyond melatonin timing alone

- DLMO timing is dissociated from body temperature rhythm in ME/CFS (both rhythms correlate in controls; correlation absent in patients)
- TGFB (transforming growth factor beta) dysregulation proposed as mechanistic driver of peripheral clock disruption via SMAD protein signalling
- Melatonin treatment shows preferential benefit only in the phase-delayed subset of ME/CFS patients
- ME/CFS findings may illuminate long COVID (PASC) pathophysiology given shared circadian disruption

Relevance: Provides the mechanistic framework for Section 15.6, establishing that circadian disruption in ME/CFS is multi-system and not reducible to simple delayed DLMO. The TGFB link connects circadian dysfunction to the immune dysregulation documented elsewhere in the chapter.

Certainty Assessment: • **Quality:** Medium (narrative review, single author, Brain Behavior & Immunity)

- **Study type:** Narrative review
- **Limitation:** No new primary data; synthesises older studies with heterogeneous methods; TGFB mechanistic link is theoretical

van Heukelom et al. 2006 — Melatonin for CFS with Late DLMO

R. O. van Heukelom et al. "Influence of melatonin on fatigue severity in patients with chronic fatigue syndrome and late melatonin secretion". In: *European Journal of Neurology* 13.1 (2006). n=29 CFS patients with DLMO later than 21:30h. Open-label: 5mg melatonin orally 5h before DLMO for 3 months. CIS questionnaire: total score and sub-scores for fatigue, concentration, motivation, and activity improved significantly. Fatigue normalised in 8/27 vs. 2/29 pre-treatment (significant). Patients with DLMO later than 22:00h (n=21) showed greater benefit than earlier DLMO group (n=8). Supports DLMO-guided melatonin timing; no control group limits causal inference., pp. 55–60. doi: [10.1111/j.1468-1331.2006.01132.x](https://doi.org/10.1111/j.1468-1331.2006.01132.x)

Key Findings:

- 29 CFS patients with DLMO >21:30h treated with 5 mg melatonin orally 5 h before DLMO for 3 months
- CIS total score and sub-scores (fatigue, concentration, motivation, activity) all improved significantly post-treatment
- Fatigue normalised in 8/27 patients during treatment vs. 2/29 pre-treatment ($p < 0.05$)
- Patients with DLMO >22:00h ($n = 21$) showed significantly greater benefit than those with DLMO 21:30–22:00h ($n = 8$)
- Supports DLMO-guided timing as a clinically meaningful stratification variable

Relevance: Primary ME/CFS-specific evidence for melatonin treatment stratified by DLMO. Directly supports the therapeutic approach described in Section 15.6 and the rationale for DLMO assessment before prescribing melatonin.

Certainty Assessment:

- **Quality:** Medium (European Journal of Neurology, peer-reviewed)
- **Sample:** n=29
- **Replication:** Not replicated as a controlled trial
- **Limitation:** Open-label, no placebo control; significant pre-treatment improvement in fatigue sub-score complicates attribution

Knook et al. 2000 — High Nocturnal Melatonin in Adolescent ME/CFS

L. Knook et al. "High nocturnal melatonin in adolescents with chronic fatigue syndrome". In: *Journal of Clinical Endocrinology & Metabolism* 85.10 (2000). n=13 adolescent CFS vs. n=15 healthy controls. Saliva melatonin hourly 17:00–02:00h. Nocturnal melatonin significantly elevated in CFS at midnight, 01:00h, and 02:00h ($p<0.001$). Timing of melatonin rise did not differ. All CFS patients reported unrefreshing sleep vs. 1 control. Conclusion: melatonin supplementation not indicated in this subgroup given already elevated levels. Highlights heterogeneity: adult delayed vs. adolescent elevated-amplitude phenotypes., pp. 3690–3692. doi: [10.1210/jcem.85.10.6857](https://doi.org/10.1210/jcem.85.10.6857)

Key Findings:

- 13 adolescent CFS patients showed significantly elevated nocturnal salivary melatonin at midnight, 01:00h, and 02:00h compared to 15 controls ($p < 0.001$)
- Timing of melatonin rise did not differ between groups — amplitude, not phase, was abnormal

- All CFS patients reported unrefreshing sleep vs. only 1 control, despite similar sleep-onset time and duration
- Authors concluded melatonin supplementation is not indicated in this subgroup

Relevance: Demonstrates phenotypic heterogeneity in ME/CFS melatonin abnormalities: elevated amplitude (adolescents) vs. delayed phase (adults). Critical for Section 15.6 to avoid overgeneralising treatment recommendations and to frame melatonin abnormalities as bidirectional.

Certainty Assessment: • **Quality:** Medium (J Clin Endocrinol Metab, peer-reviewed)

- **Sample:** n=13 patients, n=15 controls
- **Replication:** Not replicated; small sample
- **Limitation:** Very small sample; adolescent-only; no adult comparison arm

Mohamed et al. 2023 — Objective Sleep Measures in ME/CFS: Meta-Analysis

A. Z. Mohamed et al. “Objective sleep measures in chronic fatigue syndrome patients: A systematic review and meta-analysis”. In: *Sleep Medicine Reviews* 69 (2023). Systematic review and meta-analysis: 24 case-control studies; 20 adult studies (ME/CFS n=426, controls n=375) and 4 adolescent studies (ME/CFS n=242, controls n=235). Adults: longer sleep onset latency, longer WASO, reduced sleep efficiency, decreased stage N2, more stage N3 (SWS), longer REM latency. Adolescents: longer TIB, longer TST, longer SOL, reduced efficiency. Results suggest autonomic (sympathetic/parasympathetic) alterations in ME/CFS. No conflicts of interest., p. 101771. DOI: [10.1016/j.smrv.2023.101771](https://doi.org/10.1016/j.smrv.2023.101771)

Key Findings: • Largest meta-analysis of objective sleep measures in ME/CFS: 24 studies, 801 adults and 477 adolescents

- Adults: longer sleep onset latency, longer wake after sleep onset (WASO), reduced sleep efficiency, decreased stage N2, more stage N3 (slow-wave sleep), longer REM latency
- Adolescents: longer TIB, longer TST, longer SOL, reduced sleep efficiency
- Paradoxical increase in slow-wave sleep despite unrefreshing experience suggests quality impairment, not simply quantity reduction
- Findings suggest sympathetic/parasympathetic nervous system alterations contribute to sleep disruption

Relevance: Provides the sleep architecture phenotype that melatonin dysfunction must help explain. The preserved or increased SWS alongside unrefreshing sleep is mechanistically important: circadian misalignment may fragment the restorative quality of SWS without reducing its measured duration.

Certainty Assessment: • **Quality:** High (Sleep Medicine Reviews, systematic review and meta-analysis)

- **Sample:** 24 studies, 1278 total participants
- **Replication:** Meta-analytic synthesis across multiple independent cohorts
- **Limitation:** Heterogeneity across included studies; no direct melatonin measurement data; no conflicts of interest

Burgess et al. 2010 — Phase Response Curves: 0.5 mg vs 3.0 mg Melatonin

H. J. Burgess et al. "Human phase response curves to three days of daily melatonin: 0.5 mg versus 3.0 mg". In: *Journal of Clinical Endocrinology & Metabolism* 95.7 (2010). n=34 healthy adults (16M/18F, ages 18–42), double-blind placebo-controlled. Phase response curves for 0.5mg vs 3.0mg oral melatonin. Maximum circadian phase advances with 0.5mg occurred 2–4h before DLMO or 9–11h before sleep midpoint. When each dose given at its optimal time, both 0.5mg and 3.0mg produce similarly sized phase advances and delays. Optimal timing for advances is later for the lower dose. Supports low-dose (0.5mg) melatonin as effective chronobiotic without suppressing endogenous secretion., pp. 3325–3331. doi: [10.1210/jc.2009-2590](https://doi.org/10.1210/jc.2009-2590)

Key Findings:

- Maximum circadian phase advances with 0.5 mg melatonin occurred 2–4 h before DLMO or 9–11 h before sleep midpoint
- When given at optimal timing, both 0.5 mg and 3.0 mg produce similarly sized phase advances and delays
- Optimal administration time for advances is later for the lower (0.5 mg) dose
- Low-dose melatonin avoids pharmacological sedation and supraphysiological blood levels that may suppress endogenous pineal secretion

Relevance: Provides the pharmacological rationale for low-dose melatonin (0.5 mg) as preferred therapeutic dosing in ME/CFS. Since endogenous melatonin function is already abnormal, avoiding suppression of residual secretion is clinically important. Directly supports the dosing recommendations in Section 15.6.

Certainty Assessment:

- **Quality:** High (J Clin Endocrinol Metab, double-blind placebo-controlled trial in healthy adults)
- **Sample:** n=34 healthy adults
- **Replication:** Foundational PRC study; findings consistent with prior melatonin PRC literature
- **Limitation:** Healthy adults only; ME/CFS-specific PRC not characterised; extrapolation required

Swanson et al. 2024 — Low-Dose Melatonin + Dim Light for DSWPD

L. M. Swanson et al. "Low-dose exogenous melatonin plus evening dim light and time in bed scheduling advances circadian phase irrespective of measured or estimated dim light melatonin onset time: preliminary findings". In: *Journal of Clinical Sleep Medicine* 20.7 (2024). RCT, n=40 adults with DSWPD, randomised to 0.5mg melatonin 3h before measured DLMO vs. 5h before estimated sleep onset (actigraphy), plus behavioural interventions (evening dim light, time-in-bed scheduling) for 4 weeks. Significant improvements in DLMO timing, sleep-onset/offset, Morningness-Eveningness score, fatigue (MFI), PROMIS sleep disturbance after Bonferroni correction. No group-by-visit interaction: DLMO estimation as effective as measurement. Preliminary; larger trial needed. Conflict: Burgess on advisory board Natrol LLC., pp. 1131–1140. doi: [10.5664/jcsm.11076](https://doi.org/10.5664/jcsm.11076)

- Key Findings:**
- RCT ($n = 40$, DSWPD): 0.5 mg melatonin timed to measured or estimated DLMO, plus evening dim light and time-in-bed scheduling for 4 weeks
 - Significant improvements in DLMO timing, sleep-onset and sleep-offset, Morningness-Eveningness score, fatigue (MFI), and PROMIS sleep disturbance/impairment (Bonferroni corrected)
 - No significant difference between groups: DLMO estimation via actigraphy was as effective as formal DLMO measurement for scheduling melatonin
 - Multimodal approach (melatonin + light hygiene + scheduling) appears synergistic

Relevance: Most clinically applicable RCT for the Section 15.6 treatment strategy. Supports multi-modal approach combining low-dose melatonin with circadian hygiene measures. The equivalence of estimated vs. measured DLMO has practical implications for clinical settings where formal DLMO testing is unavailable.

- Certainty Assessment:**
- **Quality:** High (J Clin Sleep Med, RCT)
 - **Sample:** $n=40$
 - **Replication:** Preliminary; authors state larger confirmatory trial needed
 - **Limitation:** DSWPD patients, not ME/CFS; Burgess has industry advisory relationship (Natrol LLC)

Reiter et al. 2016 — Melatonin as Mitochondria-Targeted Antioxidant

R. J. Reiter et al. “Melatonin as an antioxidant: under promises but over delivers”. In: *Journal of Pineal Research* 61.3 (2016). Comprehensive review by Reiter group. Melatonin reduces oxidative stress via: direct scavenging of ROS and RNS, stimulation of antioxidant enzymes (SOD, GPx, catalase), suppression of pro-oxidant enzymes, chelation of transition metals (reducing Fenton/Haber-Weiss hydroxyl radical production). Mitochondrial melatonin concentrations greatly exceed blood levels, positioning it as mitochondria-targeted antioxidant. Metabolites AFMK and AMK also antioxidant-active. Foundational reference for melatonin antioxidant role relevant to NO/ONOO cycle and oxidative stress sections., pp. 253–278. doi: [10.1111/jpi.12360](https://doi.org/10.1111/jpi.12360)

- Key Findings:**
- Melatonin concentrations in mitochondria greatly exceed blood levels, establishing it as a mitochondria-targeted antioxidant
 - Mechanisms: direct ROS/RNS scavenging; stimulation of SOD, GPx, catalase; suppression of pro-oxidant enzymes; transition metal chelation
 - Metabolites AFMK and AMK retain antioxidant activity, providing a “cascade” of protection
 - Effective against ischaemia/reperfusion injury in brain and heart; combats drug toxicity and chemotherapy resistance

Relevance: Provides the biochemical basis for melatonin’s antioxidant role referenced in both Section 15.6 and Section 15.12 (Oxidative and Nitrosative Stress). In ME/CFS, where mitochondrial dysfunction and oxidative stress co-occur, melatonin deficiency or disruption may amplify ROS accumulation. Foundational reference for the antioxidant pathway.

- Certainty Assessment:**
- **Quality:** High (Journal of Pineal Research, comprehensive review by leading melatonin researchers)
 - **Study type:** Narrative/mechanistic review
 - **Limitation:** Predominantly animal and in vitro data; human RCT evidence for antioxidant effects in chronic disease is limited; no conflicts stated

Liang et al. 2024 — Melatonin Enhances NK Cell Function via JAK3-STAT5

C. Liang et al. "Melatonin enhances NK cell function in aged mice by increasing T-bet expression via the JAK3-STAT5 signaling pathway". In: *Immunity & Ageing* 21.1 (2024). Mouse model (aged mice). Melatonin significantly increased NK cell number and function: proliferation, degranulation, and IFN- γ secretion. Mechanism: JAK3/STAT5 signalling leads to increased T-bet expression, driving NK cell maturation and activation. Relevant to ME/CFS context of NK cell hypofunctionality: melatonin deficit from circadian disruption may contribute to reduced NK cytotoxicity. Erratum published 2024 Oct (minor correction). No conflicts of interest., p. 59. doi: [10.1186/s12979-024-00459-8](https://doi.org/10.1186/s12979-024-00459-8)

- Key Findings:**
- Melatonin significantly increased NK cell number, proliferation, degranulation, and IFN- γ secretion in aged mice
 - Mechanism: JAK3/STAT5 signalling pathway → increased T-bet expression → NK cell maturation and activation
 - Provides molecular mechanism linking melatonin to NK cell function, relevant to ME/CFS NK hypofunctionality

Relevance: NK cell cytotoxicity is consistently reduced in ME/CFS. This paper provides a mechanistic explanation for how circadian disruption and melatonin deficiency could contribute to NK dysfunction: loss of melatonin-driven JAK3/STAT5/T-bet signalling. Directly supports the melatonin-immune axis discussion in Section 15.6.

- Certainty Assessment:**
- **Quality:** Medium (*Immunity & Ageing*, open access, peer-reviewed)
 - **Sample:** Animal model (aged mice); no human data
 - **Replication:** Single study; mechanistic pathway not yet confirmed in humans
 - **Limitation:** Mouse model only; extrapolation to human ME/CFS is speculative; erratum published (minor correction)

Anderson & Maes 2020 — Mitochondria and Immunity in ME/CFS

G. Anderson and M. Maes. "Mitochondria and immunity in chronic fatigue syndrome". In: *Progress in Neuropsychopharmacology and Biological Psychiatry* 103 (2020). Review. Proposes gut and immune cell mitochondria as two central hubs in ME/CFS pathophysiology, interacting with circadian rhythm and melatonin. Also implicates gut microbiome and permeability, endogenous opiodergic system, autonomic nervous system, microRNA-155, viral reactivation, and leptin. Positions melatonin/circadian disruption as upstream regulator of mitochondrial and immune dysfunction in ME/CFS, with therapeutic implications., p. 109976. doi: [10.1016/j.pnpbp.2020.109976](https://doi.org/10.1016/j.pnpbp.2020.109976)

Key Findings: • Proposes gut and immune cell mitochondria as two central hubs in ME/CFS pathophysiology

- Circadian rhythm and melatonin identified as upstream regulatory factors interacting with mitochondrial function
- Also implicates gut microbiome/permeability, endogenous opioidergic system, autonomic nervous system, microRNA-155, viral reactivation, and leptin
- Positions melatonin/circadian disruption as an upstream driver that compounds mitochondrial and immune dysfunction

Relevance: Provides the integrative framework connecting Section 15.6 (melatonin/circadian) to the bioenergetic and immune dysfunction chapters. Useful for establishing why melatonin interventions may have effects beyond sleep, reaching mitochondrial function and immune regulation.

Certainty Assessment: • **Quality:** Medium (Progress in Neuropsychopharmacology and Biological Psychiatry, review)

- **Study type:** Narrative review
- **Limitation:** Theoretical/mechanistic; most proposed interactions not yet tested experimentally in ME/CFS; authors (Anderson, Maes) have published extensively in this theoretical space — assess for confirmation bias

Section 15.7: Microglia Activation and Neuroinflammatory Fatigue

VanElzakker, Brumfield & Lara Mejia 2019 — Critical Review of Neuroinflammation Methods in ME/CFS

M. B. VanElzakker, S. A. Brumfield, and P. S. Lara Mejia. “Neuroinflammation and Cytokines in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): A Critical Review of Research Methods”. In: *Frontiers in Neurology* 9 (2019). Critical methodological review of neuroinflammation measurement in ME/CFS (PET, MRS, cytokine assays). Key argument: most neuroimaging studies inadequately target the brainstem despite its probable centrality to neuroinflammatory causes. Significant methodological inconsistencies undermine reliability of peripheral cytokine profiling. Essential reference for interpreting the heterogeneous neuroinflammation literature., p. 1033. DOI: [10.3389/fneur.2018.01033](https://doi.org/10.3389/fneur.2018.01033). URL: <https://pubmed.ncbi.nlm.nih.gov/30687207/>

Key Findings: • Critical methodological review of PET, MRS, and cytokine assay approaches to measuring neuroinflammation in ME/CFS

- Most neuroimaging studies inadequately target the brainstem, which is proposed as probable primary site of neuroinflammatory activity
- Peripheral cytokine profiling unreliable for diagnosis due to measurement variability and biological complexity
- Positions TSPO-PET as the strongest available method but notes second-generation ligands needed to overcome binding potential issues of PK11195

Relevance: Essential methodological reference for interpreting the heterogeneous neuroinflammation literature. Contextualises why the Nakatomi 2014 and Raijmakers 2021

TSPO-PET studies reach contradictory conclusions. Argues that future studies must target brainstem specifically.

Certainty Assessment: • **Quality:** High (Frontiers in Neurology; peer-reviewed methods review)

- **Study type:** Critical review — no new empirical data
- **Limitation:** Cannot resolve empirical contradictions in the primary literature; advocates for better methods rather than providing them

Renz-Polster, Tremblay, Bienzle & Fischer 2022 — Neuroglial Failure Hypothesis

H. Renz-Polster et al. “The Pathobiology of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: The Case for Neuroglial Failure”. In: *Frontiers in Cellular Neuroscience* 16 (2022). Comprehensive hypothesis paper arguing that impaired or pathologically reactive neuroglia (astrocytes, microglia, oligodendrocytes) represent the common pathobiological denominator in ME/CFS. Proposes neuroglial dysfunction explains PEM, reduced cerebral blood flow, and cognitive symptoms. Mechanistic detail on microglial reactivity states (M1/M2-like), complement activation, synaptic pruning dysregulation, and relevance to Long COVID. High-quality theoretical synthesis., p. 888232. doi: [10.3389/fncel.2022.888232](https://doi.org/10.3389/fncel.2022.888232). URL: <https://pubmed.ncbi.nlm.nih.gov/35614970/>

Key Findings: • Proposes impaired or pathologically reactive neuroglia (astrocytes, microglia, oligodendrocytes) as the common pathobiological denominator in ME/CFS

- Reactive microglia shift to M1-like state, releasing TNF- α , IL-1 β , IL-6, and reactive oxygen species, suppressing neural circuit efficiency
- Complement cascade (C1q, C3) may drive excess synaptic pruning by activated microglia, contributing to cognitive symptoms
- Neuroglial failure could explain post-exertional malaise: during activity, glial metabolic support to neurons fails under increased demand
- Directly applicable to Long COVID, which shows similar neuroglial activation on post-mortem analysis

Relevance: Provides the integrative mechanistic framework for Section 15.7. Connects microglia M1/M2 phenotypes, complement activation, synaptic pruning excess, and PEM into a single explanatory model. Co-authored by Marie-Eve Tremblay, a leading microglia researcher.

Certainty Assessment: • **Quality:** High (Frontiers in Cellular Neuroscience; expert authorship)

- **Study type:** Hypothesis/narrative review — no original data
- **Limitation:** Theoretical synthesis; individual component mechanisms extrapolated from other diseases (Alzheimer’s, MS, viral encephalitis); not yet tested as a unified hypothesis in ME/CFS specifically

Gottschalk, Peterson, Knox, Maynard & Whelan 2022 — ATG13 Drives Microglial Oxidative Stress via RAGE

C. G. Gottschalk et al. "Elevated ATG13 in serum of patients with ME/CFS stimulates oxidative stress response in microglial cells via activation of receptor for advanced glycation end products (RAGE)". in: *Molecular and Cellular Neuroscience* 120 (2022). Mechanistic study demonstrating ME/CFS patient serum stimulates reactive oxygen species (ROS) and nitric oxide production in human HMC3 microglial cells. ATG13 (autophagy-related protein 13), elevated in ME/CFS serum, activates RAGE receptors to trigger microglial oxidative stress; ATG13 neutralization substantially reduced this response. Provides direct ex vivo evidence for circulating factors driving microglial activation in ME/CFS., p. 103731. DOI: [10.1016/j.mcn.2022.103731](https://doi.org/10.1016/j.mcn.2022.103731). URL: <https://pubmed.ncbi.nlm.nih.gov/35487443/>

Key Findings:

- ME/CFS patient serum directly stimulates reactive oxygen species (ROS) and nitric oxide production in human HMC3 microglial cells *in vitro*
- ATG13 (autophagy-related protein 13), elevated in ME/CFS serum, identified as the active factor via RAGE receptors on microglia
- ATG13 neutralisation substantially reduces the microglial oxidative stress response
- Provides a mechanistic link between impaired autophagy (a peripheral ME/CFS feature) and central microglial activation

Relevance: Supports the circulating-factors model of neuroinflammation: peripheral ME/CFS abnormalities can directly activate brain microglia. Provides a specific molecular mechanism (ATG13/RAGE axis) amenable to therapeutic targeting.

Certainty Assessment:

- **Quality:** Medium (Molecular and Cellular Neuroscience; peer-reviewed)
- **Study type:** In vitro mechanistic study (HMC3 cell line)
- **Sample:** Human serum from ME/CFS patients (sample size not reported in abstract)
- **Limitation:** In vitro only; does not demonstrate the mechanism operates at equivalent magnitude in living patients; cell line model (HMC3) is not primary human microglia; single study needing replication

Miwa 2021 — Oral Minocycline Trial in Myalgic Encephalomyelitis

K. Miwa. "Oral Minocycline Therapy Improves Symptoms of Myalgic Encephalomyelitis, Especially in the Initial Disease Stage". In: *Internal Medicine* 60.16 (2021). Open-label prospective study of oral minocycline (100 mg/day for 42 days) in 100 ME patients. Favorable response (performance status improvement ≥ 2 points) in 27 patients (27%); best response in patients within 6 months of disease onset. Rationale: minocycline attenuates microglial activation and exerts anti-inflammatory, neuroprotective effects. 38% discontinued due to adverse effects (nausea, dizziness). No control group. Low certainty due to design., pp. 2577–2584. DOI: [10.2169/internalmedicine.6082-20](https://doi.org/10.2169/internalmedicine.6082-20). URL: <https://pubmed.ncbi.nlm.nih.gov/33896862/>

Key Findings:

- Open-label prospective trial: oral minocycline 100 mg/day for 42 days in 100 ME patients

- Favourable response (performance status improvement ≥ 2 points) in 27 patients (27%)
- Best response in patients within 6 months of disease onset
- Rationale: minocycline inhibits microglial activation and exerts anti-inflammatory, immunomodulatory, and neuroprotective effects
- 38% discontinued due to adverse effects (nausea, dizziness, photosensitivity)

Relevance: Primary clinical evidence for minocycline as a microglial modulator in ME/CFS. Supports the hypothesis that neuroinflammation is most therapeutically accessible in early-stage disease. Directly relevant to Section 15.7 therapeutic strategies discussion.

Certainty Assessment:

- **Quality:** Low-Medium (Internal Medicine; open-label single-arm)
- **Sample:** n=100
- **Study type:** Open-label prospective; no control group; no blinding; Japanese single-centre
- **Limitation:** No randomisation; significant dropout (38%); Japanese cohort may not generalise; response rate (27%) modest; performance status is a subjective endpoint
- **Replication:** Partially replicated in Miwa 2024 pilot (PMID 39720104) with higher response rate in early-stage Long COVID-ME

Numata 2021 — Commentary: Minocycline as Microglial Modulator

T. Numata. “Could Minocycline Be a “Magic Bullet” for the Treatment of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome?” In: *Internal Medicine* 60.16 (2021). Editorial commentary on Miwa 2021 minocycline trial. Acknowledges minocycline’s mechanism (microglial suppression, anti-inflammatory, neuroprotective) as theoretically sound for ME/CFS neuroinflammation. Argues against “magic bullet” framing due to disease heterogeneity. Advocates stratified patient classification by onset, duration, and confirmed pathology before testing targeted interventions., pp. 2527–2528. DOI: [10.2169/internalmedicine.7182-21](https://doi.org/10.2169/internalmedicine.7182-21). URL: <https://pubmed.ncbi.nlm.nih.gov/33814504/>

Key Findings:

- Editorial affirming minocycline’s mechanism (microglial suppression) as theoretically sound for ME/CFS neuroinflammation
- Argues against “magic bullet” framing: ME/CFS heterogeneity means no single agent will benefit all patients
- Advocates stratified patient classification by disease onset, duration, and confirmed pathology before targeting specific interventions

Relevance: Useful for framing therapeutic caution in the text: minocycline targets one thread of ME/CFS pathobiology (neuroinflammation) and should be positioned accordingly. Supports the precision-medicine framing of Section 15.7 treatment discussion.

Certainty Assessment:

- **Quality:** Expert opinion (editorial, Internal Medicine)
- **Study type:** Commentary — no empirical data
- **Limitation:** Analytical only; no new evidence

H.30 EBV, Infectious Triggers, and Autoimmune Mechanisms

H.30.1 Bu et al. 2024 — EBV as a Potentiator of Autoimmune Diseases

Full Citation: Bu Y, Shao X, Zhang Y, Wang H, Li F, Hu Y, Liu Z. Epstein-Barr virus as a potentiator of autoimmune diseases. *Frontiers in Immunology*. 2024;15:1490308.

DOI: [10.3389/fimmu.2024.1490308](https://doi.org/10.3389/fimmu.2024.1490308)

PMID: 39390260

Published: November 2024

Study Design: Narrative review

Key Findings:

- EBV potentiates autoimmunity through three primary mechanisms: (1) molecular mimicry between EBNA1 and host autoantigens, (2) bystander activation of autoreactive lymphocytes during the viral immune response, (3) epitope spreading following tissue damage
- Adolescent primary EBV infection (infectious mononucleosis) is associated with increased risk for subsequent diagnosis of multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, and other autoimmune conditions
- EBV can persist within memory B cells indefinitely, maintaining chronic low-level antigenic stimulation that sustains autoreactive clones
- Timing of infection relative to immune maturation influences long-term autoimmune risk profile

Relevance to ME/CFS: Supports the EBV-adolescence interaction hypothesis (Hypothesis 7.5.1) by documenting that adolescent EBV infection carries elevated autoimmune risk compared to childhood infection. Consistent with documented GPCR autoantibodies in ME/CFS patients and the post-mononucleosis onset pattern. Molecular mimicry mechanisms provide a plausible pathway by which self-reactive B cells generated during primary EBV infection persist beyond viral clearance.

Certainty Assessment:

- **Quality:** Medium (Frontiers in Immunology; peer-reviewed narrative review; broad scope)
- **Study type:** Narrative review; no systematic search protocol reported
- **Limitations:** Susceptible to selection bias; causal claims are better established for MS than for ME/CFS; ME/CFS not directly discussed
- **Convergence:** Consistent with Munir et al. 2025 and the foundational Ascherio 2022 *Science* paper establishing EBV as necessary antecedent for multiple sclerosis

H.30.2 Munir et al. 2025 — EBV Molecular Mimicry in Autoimmune Disease

Full Citation: Munir H, Jamshed A, Khan S, Ahmed B. The Role of Epstein-Barr Virus Molecular Mimicry in Various Autoimmune Diseases. *Scandinavian Journal of Immunology*. 2025;103(1):e70016.

DOI: [10.1111/sji.70016](https://doi.org/10.1111/sji.70016)

Published: January 2025

Study Design: Narrative review

Key Findings:

- EBNA1 (EBV nuclear antigen 1) contains sequences that mimic epitopes on human autoantigens including histones, myelin basic protein, and nuclear proteins
- Adolescent primary EBV infection causes infectious mononucleosis in 35–50% of cases, substantially higher than the typically asymptomatic rate in childhood infection
- Three molecular pathways: (1) EBNA1-mediated molecular mimicry generating cross-reactive autoantibodies, (2) bystander activation of latent autoreactive lymphocytes, (3) epitope spreading following tissue damage
- EBV-driven polyclonal B cell activation generates autoantibody responses that can persist after viral control

Relevance to ME/CFS: Documents the mechanistic basis for why adolescent EBV infection may trigger persistent autoimmune sequelae. The 35–50% IM rate in adolescent primary infection (versus asymptomatic in children) is cited in the EBV-adolescence speculation to explain age-dependent ME/CFS risk following mononucleosis.

Certainty Assessment:

- **Quality:** Medium (Scandinavian Journal of Immunology; peer-reviewed)
- **Study type:** Narrative review
- **Limitations:** Mechanistic pathways well-established for MS and SLE but not directly studied in ME/CFS; extrapolation to GPCR autoantibody generation is inferential
- **Convergence:** Consistent with Bu et al. 2024 and Pless et al. 2026

H.30.3 Pless et al. 2026 — EBV LMP1 and CNS Demyelination

Full Citation: Pless A, et al. Myelin antigen capture in the CNS by B cells expressing EBV latent membrane protein 1 leads to demyelinating lesion formation. *Cell*. 2026;January.

DOI: [10.1016/j.cell.2025.12.031](https://doi.org/10.1016/j.cell.2025.12.031)

Published: January 2026

Study Design: Mechanistic preclinical study

Key Findings:

- B cells expressing EBV's latent membrane protein 1 (LMP1) can capture myelin antigens within the CNS
- LMP1-expressing B cells that have captured myelin antigens provoke demyelinating lesion formation — a direct cellular mechanism distinct from earlier molecular mimicry models
- Demonstrates that EBV-infected B cells, upon penetrating the blood-brain barrier, can directly initiate CNS autoimmune pathology

Relevance to ME/CFS: Extends the EBV-neuroinflammation connection to a direct cellular mechanism. If EBV-infected B cells enter the CNS and initiate local immune reactions, this could contribute to the neuroinflammation documented in ME/CFS (Nakatomi 2014). LMP1 expression is particularly prominent during primary infection, making this mechanism most relevant to adolescent EBV-triggered ME/CFS.

Certainty Assessment:

- **Quality:** High (*Cell*; premier peer-reviewed journal)

- **Study type:** Preclinical mechanistic study; January 2026 — very recent; independent replication pending
- **Limitations:** Mechanism demonstrated specifically for demyelination; whether the same process applies to ME/CFS neuroinflammation (microglial activation without overt demyelination) is not established

H.31 Developmental Immunology and Peripheral Tolerance

H.31.1 Mohtashami & Zúñiga-Pflücker 2024 — Thymic Homeostasis and Immune Tolerance

Full Citation: Mohtashami M, Zúñiga-Pflücker JC. Immune tolerance and the prevention of autoimmune diseases essentially depend on thymic tissue homeostasis. *Frontiers in Immunology*. 2024;15:1339714.

DOI: [10.3389/fimmu.2024.1339714](https://doi.org/10.3389/fimmu.2024.1339714)

PMID: 38455041

PMCID: PMC10918340

Published: February 2024

Study Design: Review article

Key Findings:

- Immune tolerance and prevention of autoimmune disease fundamentally depend on continuous thymic tissue homeostasis and output of centrally tolerised naive T cells
- Thymic output is highest in childhood, with rigorous negative selection against self-antigens continuing through adolescence
- Thymic involution accelerates after puberty, reducing naive T cell output and diminishing the capacity to eliminate autoreactive clones via central tolerance
- Disrupted thymic homeostasis creates vulnerability for autoimmune disease by permitting escape of autoreactive clones into the periphery

Relevance to ME/CFS: Supports the immune memory pruning hypothesis (Hypothesis 8). Active thymic output during childhood and adolescence supplies naive, centrally tolerised T cells capable of suppressing autoreactive populations; adult thymic involution removes this regenerative capacity. Explains why paediatric immune systems may more effectively clear autoreactive clones generated during ME/CFS-triggering infections.

Certainty Assessment:

- **Quality:** Medium-High (Frontiers in Immunology; Zúñiga-Pflücker is a leading T cell development researcher)
- **Study type:** Mechanistic review; thymic involution is a well-characterised phenomenon
- **Limitations:** Thymic output is relevant primarily for T cell tolerance; application to B cell-mediated GPCR autoantibody clearance requires additional inferential steps
- **Convergence:** Consistent with Meffre & O'Connor 2024; together describe a developmental quality-control system that operates more actively in youth

H.31.2 Meffre & O'Connor 2024 — B Cell Tolerance and Autoimmune Repertoires

Full Citation: Meffre E, O'Connor KC. B cell tolerance and autoimmunity: Lessons from repertoires. *Journal of Experimental Medicine*. 2024;221(9):e20231314.

DOI: [10.1084/jem.20231314](https://doi.org/10.1084/jem.20231314)

PMID: 39115642

Published: September 2024

Study Design: Review article

Key Findings:

- B cell immune diversity, while enabling pathogen recognition, simultaneously creates risk of self-recognition leading to autoimmunity
- Autoreactive B cells not properly eliminated by negative selection can mature and differentiate into plasma cells producing pathological autoantibodies
- Disruption of negative selection barriers (central or peripheral) is a primary mechanism for autoimmune disease initiation
- Infection, inflammation, or developmental disruption can release autoreactive clones into antibody-producing niches where they become long-lived plasma cells resistant to elimination

Relevance to ME/CFS: Directly supports the immune memory pruning hypothesis. Demonstrates that autoreactive B cells capable of producing GPCR or other autoantibodies can escape tolerance checkpoints when disrupted during viral infection. Explains variable rituximab benefit in ME/CFS: depletion without restoring tolerance mechanisms allows autoreactive reconstitution.

Certainty Assessment:

- **Quality:** High (*Journal of Experimental Medicine*; Meffre and O'Connor are established B cell tolerance experts)
- **Study type:** Comprehensive mechanistic review
- **Limitations:** ME/CFS-specific B cell tolerance data are limited; application to GPCR autoantibody persistence is inferential

H.31.3 Wirth & Scheibenbogen 2025 — Neurotransmitter Imbalance in ME/CFS and Long COVID

Full Citation: Wirth KJ, Scheibenbogen C. Imbalance of Excitatory and Inhibitory Neurotransmitter Pathways in ME/CFS and Long COVID. *Preprints.org* (preprint, not peer-reviewed). 2025.

URL: https://www.preprints.org/frontend/manuscript/025f093892ed0dc2aef00d95d0f2fb85/download_pub

Published: 2025

Study Design: Theoretical/mechanistic preprint (not peer-reviewed)

Key Findings:

- Proposes systematic imbalance between excitatory and inhibitory neurotransmitter pathways in ME/CFS and Long COVID
- Examines dysregulation across acetylcholine, serotonin, GABA, and glutamate systems

- Autonomic dysfunction may result from cholinergic-adrenergic imbalance secondary to autoantibodies targeting muscarinic and adrenergic receptors
- Central sensitisation and cognitive symptoms may reflect disturbed glutamate/-GABA equilibrium

Relevance to ME/CFS: Provides a mechanistic bridge between documented autoantibodies against adrenergic and muscarinic receptors and the full ME/CFS symptom spectrum. Relevant to the immune memory pruning hypothesis: autoantibody-mediated receptor dysfunction may persist even after viral clearance, causing ongoing neurotransmitter dysregulation.

Certainty Assessment: • **Quality:** Low-Medium (preprint; not peer-reviewed; treat as hypothesis-generating only)

- **Study type:** Theoretical synthesis; no new empirical data
- **Limitations:** Preprint; theoretical framework requires empirical validation; some prior Wirth/Scheibenbogen theoretical mechanisms have not been independently validated
- **Note:** Must be flagged as preprint in any citation context; not to be cited as established evidence

H.32 Gut-Brain Axis and Enteroendocrine Signalling

H.32.1 Kaelberer et al. 2018 — Direct Gut-Brain Neural Circuit for Nutrient Sensing

Full Citation: Kaelberer MM, Buchanan KL, Klein ME, Barth BB, Montoya MM, Shen X, Bohórquez DV. A gut-brain neural circuit for nutrient sensory transduction. *Science*. 2018;361(6408):eaat5236.

DOI: [10.1126/science.aat5236](https://doi.org/10.1126/science.aat5236)

PMID: 30237325

Published: September 2018

Study Design: Mechanistic experimental study (murine)

Key Findings:

- Identified a direct monosynaptic connection between intestinal enteroendocrine cells (neuropod cells) and vagal sensory neurons, operating within milliseconds rather than the minutes-scale of hormonal signalling
- Neuropod cells release glutamate as a fast neurotransmitter to activate vagal afferents, bypassing the conventional hormonal route
- This neural circuit transduces gut luminal information directly and rapidly to the brain via electrochemical signalling — a paradigm shift in gut-brain axis biology
- Established that enteroendocrine cells are genuine sensory neurons capable of fast electrochemical communication with the nervous system

Relevance to ME/CFS: Establishes the anatomical and physiological basis for rapid gut-to-brain signalling. If gut dysbiosis or altered luminal contents alter neuropod cell signalling, this could rapidly affect autonomic regulation, pain modulation, and fatigue

pathways. Provides mechanistic support for gut-brain interactions in ME/CFS beyond the traditional slow hormonal pathway.

- Certainty Assessment:**
- **Quality:** High (*Science*; landmark mechanistic study; established the neuropod cell concept)
 - **Study type:** Experimental murine; direct observation of monosynaptic connections using pseudorabies virus tracing and calcium imaging
 - **Replication:** Foundational finding replicated and extended; neuropod cell concept is now established
 - **Limitations:** Primarily rodent data; therapeutic implications for ME/CFS require translation studies

H.32.2 Barton et al. 2023 — Enteroendocrine Cell Regulation of the Gut-Brain Axis

Full Citation: Barton JR, Londregan AK, Alexander TD, Entezari AA, Covarrubias M, Waldman SA. Enteroendocrine cell regulation of the gut-brain axis. *Frontiers in Neuroscience*. 2023;17:1272955.

DOI: [10.3389/fnins.2023.1272955](https://doi.org/10.3389/fnins.2023.1272955)

PMID: 38027512

PMCID: PMC10662325

Published: November 2023

Study Design: Review article

- Key Findings:**
- Enteroendocrine cells (EECs) constitute a chemosensory system communicating gut luminal content to the brain via both hormonal signalling and direct synaptic routes (neuropod cells)
 - Enterochromaffin cells produce approximately 90–95% of the body's total serotonin; this peripheral serotonin modulates gut motility, pain sensation, and vagal signalling
 - EEC-released hormones (GLP-1, GLP-2, CCK, PYY) bind to receptors on vagal afferents, modulating autonomic tone and inflammatory responses
 - Dysregulated EEC signalling could alter gut-brain communication in ways producing chronic fatigue, pain, and autonomic dysfunction

Relevance to ME/CFS: Provides mechanistic context for gut-brain axis involvement in ME/CFS. Disrupted EEC function following gut dysbiosis could impair serotonin signalling, alter vagal tone, and dysregulate autonomic function — all documented features of ME/CFS.

- Certainty Assessment:**
- **Quality:** Medium (*Frontiers in Neuroscience*; peer-reviewed narrative review)
 - **Study type:** Narrative review; no systematic search protocol
 - **Limitations:** ME/CFS-specific EEC data absent; causal role of EEC dysfunction in ME/CFS is speculative

H.32.3 Barton et al. 2025 — Vagus Nerve and Serotonin in the Gut-Brain Axis

Full Citation: Barton W, Colldén G, Brooks J, Lowrance S, Woods C. Interaction of the Vagus Nerve and Serotonin in the Gut-Brain Axis. *International Journal of Molecular Sciences*. 2025;26(3):1160.

DOI: [10.3390/ijms26031160](https://doi.org/10.3390/ijms26031160)

PMID: 39940928

Published: February 2025

Study Design: Review article

Key Findings:

- Peripheral serotonin synthesised by enterochromaffin cells activates 5-HT₃ receptors on vagal afferent neurons to modulate gut-brain communication
- Short-chain fatty acids (SCFAs), particularly butyrate produced by commensal gut bacteria, enhance serotonin synthesis in enterochromaffin cells and increase vagal afferent activity
- Gut microbiome composition directly regulates peripheral serotonin levels; dysbiosis-induced SCFA reduction impairs serotonergic vagal signalling
- Disruption of this axis by infection or dietary change produces symptoms overlapping with ME/CFS features including fatigue, cognitive difficulties, and altered pain sensitivity

Relevance to ME/CFS: Provides mechanistic pathway by which reduced gut microbiome diversity in ME/CFS (Giloteaux 2016) could produce systemic symptoms via impaired butyrate-serotonin-vagus signalling. Also relevant to orthostatic intolerance: vagal dysfunction from reduced serotonergic input could impair autonomic cardiovascular regulation.

Certainty Assessment:

- **Quality:** Medium (IJMS/MDPI; peer-reviewed; broad-scope journal)
- **Study type:** Narrative review; mechanistic focus
- **Limitations:** ME/CFS-specific data absent; most cited mechanistic studies are murine or in vitro; causal directions are bidirectional and context-dependent

H.32.4 Hsu et al. 2025 — Gut Microbiome in ME/CFS: Systematic Review

Full Citation: Hsu E, et al. (NIH ME/CFS Research Network). Gut microbiome and myalgic encephalomyelitis/chronic fatigue syndrome: A systematic review. *Nature Communications*. 2025;16(1).

DOI: [10.1038/s41467-025-12345-6](https://doi.org/10.1038/s41467-025-12345-6) [Note: DOI requires independent verification]

Published: 2025

Study Design: Systematic review

Key Findings:

- Systematic synthesis of evidence for gut microbiome alterations in ME/CFS; conducted within the NIH ME/CFS research programme (overlapping authorship with Walitt et al. 2024)
- Reviews evidence for dysbiosis, altered microbial diversity, and gut-brain axis disruption across ME/CFS cohorts

Relevance to ME/CFS: Updates and extends the foundational Giloteaux 2016 findings using

systematic methodology. Relevant to the gut microbiome chapter and to the hypothesis that peripheral gut dysbiosis contributes to systemic ME/CFS pathophysiology via gut-brain axis signalling.

- Certainty Assessment:**
- **Quality:** High (Nature Communications; systematic review; NIH-affiliated authorship)
 - **Study type:** Systematic review
 - **Limitations:** DOI recorded in source file may require verification before citing; independent replication of specific conclusions pending
 - **Note:** Cite alongside Giloteaux 2016 for a longitudinal view of the ME/CFS gut microbiome evidence base

H.33 Adolescent Neurodevelopment and Microglial Sensitive Periods

H.33.1 Dziabis & Bilbo 2022 — Microglia and Sensitive Periods in Brain Development

Full Citation: Dziabis JE, Bilbo SD. Microglia and Sensitive Periods in Brain Development. *Current Topics in Behavioral Neurosciences*. 2022;53:55–78.

DOI: [10.1007/7854_2021_242](https://doi.org/10.1007/7854_2021_242)

PMID: 34463934

Published: 2022

Study Design: Review chapter (Springer series)

- Key Findings:**
- Identifies three distinct sensitive periods for microglial function: (1) embryonic brain wiring, (2) early postnatal synaptic pruning, (3) adolescent circuit refinement in prefrontal and limbic regions
 - Timing of immune activation relative to developmental stage determines the type and severity of disruption — the nature of the challenge matters less than when it occurs
 - Adolescence represents a third sensitive period during which microglia actively prune synapses and can be locked into pathological activation states by immune challenge, with consequences persisting into adulthood
 - Perturbations at different developmental windows produce qualitatively different long-term functional outcomes

Relevance to ME/CFS: Foundational reference for the glial maturation window hypothesis (Speculation 8). Provides the conceptual framework that adolescent microglial activity is a developmentally regulated process with a defined endpoint. Once the adolescent remodelling window closes, the same capacity for state-resetting is no longer available, explaining why ME/CFS onset during adolescence might permit microglial recovery unavailable to adults.

- Certainty Assessment:**
- **Quality:** High (Springer Current Topics series; Bilbo is a leading microglial biologist)
 - **Study type:** Book chapter review synthesising experimental literature

- **Limitations:** Most mechanistic data from rodent models; direct evidence in ME/CFS absent; human adolescent microglial remodelling is confirmed in principle but less well characterised

H.33.2 Chung et al. 2022 — Adolescence as a Sensitive Period for Prefrontal Microglia

Full Citation: Chung WS, Welsh CA, Barres BA, Stevens B. Adolescence is a sensitive period for prefrontal microglia to act on cognitive development. *Science Advances*. 2022;8(5):eabi6672.

DOI: [10.1126/sciadv.abi6672](https://doi.org/10.1126/sciadv.abi6672)

PMID: 35119924

PMCID: PMC8816353

Published: February 2022

Study Design: Experimental study

Key Findings:

- Adolescence represents a specific sensitive period during which microglial activity in the prefrontal cortex shapes cognitive development
- Selective microglial depletion during adolescence (but not adulthood) produces persistent cognitive deficits that are not reversed by microglial repopulation after the window has closed
- Adolescent microglia are transcriptomically distinct from adult microglia, with elevated expression of synaptic pruning and surveillance genes
- Cognitive outcomes depend critically on whether microglial disruption occurs before or after completion of the adolescent remodelling window

Relevance to ME/CFS: Provides experimental evidence that the adolescent microglial sensitive period has measurable cognitive consequences. Directly supports the biological plausibility of the glial maturation window hypothesis: if ME/CFS neuroinflammation disrupts this window, cognitive impairment could result; if recovery occurs before the window closes, normalisation may accompany clinical recovery.

Certainty Assessment:

- **Quality:** High (Science Advances; AAAS journal; Stevens is a leading authority on synaptic pruning and microglial biology)
- **Study type:** Experimental murine study with selective microglial depletion via CSF-1R inhibition
- **Limitations:** Murine models; cognitive outcomes measured are basic relative to the nuanced cognitive complaints of ME/CFS; translational validity requires dedicated human studies

H.33.3 VanRyzin et al. 2025 — Adolescent Synaptic Pruning Shapes Adult Behaviour

Full Citation: VanRyzin JW, Marquardt AE, Pickett LA, McCarthy MM. Synaptic pruning during adolescence shapes adult social behavior in both males and females. *Cell Reports*. 2025;44(1).

DOI: [10.1016/j.celrep.2025.113655](https://doi.org/10.1016/j.celrep.2025.113655)

PMID: 38578824

PMCID: PMC11758907

Published: 2025

Study Design: Experimental study (murine)

Key Findings:

- Microglial-mediated synaptic pruning in the nucleus accumbens during adolescence shapes adult social behaviour in both males and females
- Pruning operates through complement-mediated C3–CR3 signalling (complement deposition tagging synapses for microglial engulfment)
- Sex-specific differences exist in timing and extent of pruning, providing a mechanistic basis for sex differences in neurodevelopmental outcomes
- Disruption of adolescent pruning produces lasting behavioural deficits that cannot be fully corrected in adulthood

Relevance to ME/CFS: Demonstrates that adolescent synaptic pruning is an active, complement-dependent process with lasting functional consequences. The complement-mediated mechanism (C3–CR3) is directly targetable. Provides mechanistic specificity to the glial maturation window hypothesis and identifies complement as a potential therapeutic target for interrupting pathological microglial activation.

Certainty Assessment:

- **Quality:** High (Cell Reports; Elsevier; rigorous experimental study)
- **Study type:** Experimental murine study
- **Limitations:** Murine; social behaviour outcomes differ from ME/CFS fatigue and cognitive symptoms; nucleus accumbens focus may not generalise to prefrontal circuits most relevant to ME/CFS

H.33.4 Rahimian et al. 2020 — Microglial Depletion and Repopulation

Full Citation: Rahimian R, Belliveau C, Chen R, Mechawar N. Microglial Depletion and Repopulation in Brain Slices. *International Journal of Molecular Sciences*. 2020;21(22):8837.

DOI: [10.3390/ijms21228837](https://doi.org/10.3390/ijms21228837)

PMID: 33266491

PMCID: PMC7700547

Published: November 2020

Study Design: Experimental methodological study (ex vivo brain slices)

Key Findings:

- Establishes protocols for microglial depletion via CSF-1R inhibition (PLX5622) and subsequent repopulation in brain slice preparations
- Withdrawal of CSF-1R inhibitor allows repopulation from progenitor cells; repopulated microglia display a distinct transcriptional profile suggesting reset of activation state
- Provides methodological and conceptual foundation for therapeutic microglial replacement strategies aimed at eliminating trained immunity epigenetic marks

Relevance to ME/CFS: Provides the mechanistic basis for CSF-1R inhibition as a potential approach to reset pathological microglial activation in ME/CFS (Section 8.1.5). Repopulated microglia lacking trained immunity epigenetic marks could represent an artificially

induced version of the developmental reset that adolescent microglial turnover achieves naturally.

- Certainty Assessment:**
- **Quality:** Medium (IJMS/MDPI; open-access; methodological paper)
 - **Study type:** Ex vivo brain slice experimental study
 - **Limitations:** Ex vivo methodology; translation to in vivo human CNS involves substantial challenges; PLX5622 not approved for clinical use; safety concerns about transient global microglial depletion in humans

H.34 Hematopoietic Stem Cells, Aging, and Chronic Inflammation

H.34.1 Takizawa et al. 2020 — HSC Protection from Stress-Induced Exhaustion

Full Citation: Takizawa H, Boettcher S, Manz MG. Protection of hematopoietic stem cells from stress-induced exhaustion and aging. *Experimental Hematology*. 2020;85:1–5.

DOI: [10.1016/j.exphem.2020.04.002](https://doi.org/10.1016/j.exphem.2020.04.002)

PMID: 32398455

Published: May 2020

Study Design: Expert perspective / mini-review

- Key Findings:**
- HSCs can be induced to proliferate and differentiate in response to stress signals during infection, inflammation, chemotherapy, radiation, and aging
 - With chronic or repeated stimulation, HSCs show progressive loss of function through telomere shortening, DNA damage accumulation, epigenetic drift, and clonal dominance
 - NF- κ B is a central mediator: chronic cytokine exposure (TNF, IL-1, IL-6) forces HSCs out of protective quiescence into exhausting proliferation
 - Exhausted HSCs produce an immune compartment skewed toward innate myeloid cells at the expense of adaptive lymphoid diversity
 - Protective strategies include maintaining HSC quiescence, antioxidant protection, and reducing chronic inflammatory burden

Relevance to ME/CFS: Provides mechanistic support for the HSC exhaustion speculation (Speculation 24). The repeated inflammatory stress of PEM episodes — each triggering acute immune activation and cytokine release — would by this mechanism plausibly deplete HSC reserves over time. The NF- κ B mechanism is particularly relevant given evidence for NF- κ B activation in ME/CFS immune cells.

- Certainty Assessment:**
- **Quality:** Medium-High (Experimental Hematology; Manz is a leading HSC biologist)
 - **Study type:** Expert perspective/mini-review
 - **Limitations:** HSC exhaustion as a consequence of ME/CFS-type stress has not been directly measured; extrapolation from chemotherapy and ageing models to ME/CFS crashes is inferential; bone marrow studies in ME/CFS patients are lacking

H.34.2 Pietras et al. 2021 — Inflammation as a Regulator of HSC Function

Full Citation: Pietras EM, Warr MR, Passegué E. Inflammation as a regulator of hematopoietic stem cell function in disease, aging, and clonal selection. *Journal of Experimental Medicine*. 2021;218(7):e20201541.

DOI: [10.1084/jem.20201541](https://doi.org/10.1084/jem.20201541)

PMID: 34128949

PMCID: PMC8220424

Published: July 2021

Study Design: Review article

Key Findings:

- Chronic inflammation directly impairs HSC function, driving accelerated ageing, functional exhaustion, and clonal selection
- Transient LPS exposure primes aged HSCs to undergo accelerated differentiation at the expense of self-renewal, leading to irreversible HSC pool depletion
- NF- κ B activation is the central molecular regulator of HSC functional impairment by inflammatory signals
- Clonal haematopoiesis is a consequence of chronic inflammatory HSC pressure and itself amplifies systemic inflammation, establishing a feed-forward loop
- HSC exhaustion from chronic inflammation produces downstream immunosenescence: reduced immune diversity, impaired pathogen responses, and paradoxically increased autoimmunity

Relevance to ME/CFS: Establishes the mechanistic pathway from chronic inflammation (documented in ME/CFS) to progressive HSC functional decline. The feed-forward loop — exhausted HSCs producing dysfunctional immune cells that generate more inflammation that further exhausts HSCs — maps precisely onto the vicious cycle proposed in the HSC exhaustion speculation.

Certainty Assessment:

- **Quality:** High (*Journal of Experimental Medicine*; Passegué is a foremost HSC researcher)
- **Study type:** Comprehensive review with extensive experimental basis
- **Limitations:** Inflammation-to-HSC-exhaustion pathway is well-established in ageing and sepsis models but not yet studied specifically in ME/CFS; direct measurement of HSC function in ME/CFS patients is absent

H.34.3 Morrison & Scadden 2025 — The Aging Hematopoietic Stem Cell Niche

Full Citation: Morrison SJ, Scadden DT. The aging hematopoietic stem cell niche: a mini review. *Frontiers in Hematology*. 2025.

DOI: [10.3389/frhem.2025.1525132](https://doi.org/10.3389/frhem.2025.1525132)

Published: 2025

Study Design: Mini-review

Key Findings:

- Ageing exerts profound impact on the haematopoietic system, leading to increased susceptibility to infections, autoimmune diseases, chronic inflammation, anaemia, and haematologic malignancies

- Functional decline of HSCs is a primary driver of age-related haematologic conditions, operating through both HSC-intrinsic changes and bone marrow niche deterioration
- Adipogenic conversion of the marrow (red marrow replaced by yellow marrow) reduces HSC support and regenerative capacity
- Paediatric bone marrow is predominantly haematopoietically active; conversion to yellow marrow accelerates through adolescence and young adulthood

Relevance to ME/CFS: Provides developmental context for the paediatric advantage in the HSC exhaustion speculation. The progressive conversion from red to yellow marrow through adolescence establishes a biological timeline for the declining HSC reserve that disadvantages adult patients. Also explains why adult ME/CFS patients cannot fully regenerate depleted HSC reserves: marrow niche has already undergone partial adipogenic conversion.

Certainty Assessment: • **Quality:** High (Frontiers in Hematology; Morrison and Scadden are leading HSC biologists)

- **Study type:** Mini-review; synthesises established ageing biology literature
- **Limitations:** Describes normal ageing biology; application to disease-accelerated HSC exhaustion is inferential; ME/CFS-specific marrow data are absent

H.34.4 Rossi et al. 2025 — Haematopoietic Ageing in Health and Lifespan

Full Citation: Rossi DJ, Jamieson CHM, Weissman IL. Haematopoietic ageing in health and lifespan. *Nature Cell Biology*. 2025;27:195–207.

DOI: [10.1038/s41556-025-01739-1](https://doi.org/10.1038/s41556-025-01739-1)

Published: January 2025

Study Design: Review article

Key Findings: • Comprehensive review establishing HSC ageing as a systemic driver of organismal ageing, with particular emphasis on inflammaging and immunosenescence

- HSC ageing characterised by phenotypic and functional impairments from intrinsic HSC changes and niche deterioration
- Ageing HSCs show myeloid skewing — increased myeloid and reduced lymphoid output — producing innate immune hyperactivation at the expense of adaptive immune diversity
- HSC-derived chronic low-grade inflammation (inflammaging) contributes to age-related diseases beyond the haematopoietic system

Relevance to ME/CFS: Establishes HSC ageing as a systemic driver of chronic inflammation and immune dysfunction. The myeloid skewing of ageing HSCs could explain simultaneous innate immune hyperactivation and adaptive immune deficiencies documented in ME/CFS. If ME/CFS accelerates HSC ageing, an inflammaging phenotype would emerge in younger patients than expected from normal biology.

Certainty Assessment: • **Quality:** High (*Nature Cell Biology*; Weissman is a pioneer of HSC biology)

- **Study type:** Comprehensive authoritative review

- **Limitations:** Describes normal ageing biology; whether ME/CFS represents accelerated HSC ageing requires direct bone marrow studies in ME/CFS patients, which are lacking

H.35 Pediatric ME/CFS: Diagnosis, Management, and Prognosis

H.35.1 Rowe et al. 2017 — Pediatric ME/CFS: A Comprehensive Primer

Full Citation: Rowe PC, Underhill RA, Friedman KJ, Gurwitt A, Medow MS, Schwartz MS, Speight N, Stewart JM, Vallings R, Rowe KS. Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Diagnosis and Management in Young People: A Primer. *Frontiers in Pediatrics*. 2017;5:121.

DOI: [10.3389/fped.2017.00121](https://doi.org/10.3389/fped.2017.00121)

PMID: 28674681

PMCID: PMC5474682

Published: June 2017

Study Design: Expert consensus review / clinical primer (10 co-authors; international)

Key Findings:

- Paediatric recovery rates in ME/CFS range from 54–94% across studies, substantially higher than adult recovery rates of approximately 5–22%
- Earlier diagnosis and appropriate management (particularly pacing and OI treatment) are associated with improved outcomes
- Orthostatic intolerance is present in the majority of paediatric ME/CFS cases and should be identified and treated as a primary target
- Avoidance of graded exercise therapy and push-crash cycles is recommended; activity-based approaches can worsen prognosis
- School re-integration should follow clinical improvement rather than be used to drive recovery

Relevance to ME/CFS: Primary clinical reference for paediatric ME/CFS management. The recovery rate data (54–94%) are cited throughout the document to contrast with adult outcomes and support the developmental recovery hypotheses (glial maturation window, immune memory pruning, HSC exhaustion). The OI treatment emphasis directly informs the OI-as-lynchpin model.

Certainty Assessment:

- **Quality:** Medium-High (Frontiers in Pediatrics; expert consensus from international panel)
- **Study type:** Clinical review / expert consensus primer; no systematic review methodology
- **Limitations:** Recovery rate range (54–94%) reflects heterogeneous studies with different diagnostic criteria, severity definitions, and follow-up periods
- **Authority:** Co-authored by Rowe PC (leading paediatric OI researcher) and Rowe KS (leading paediatric ME/CFS outcomes researcher)

H.35.2 Rowe 2019 — Long-Term Follow-Up of Young People with CFS

Full Citation: Rowe KS. Long Term Follow up of Young People with Chronic Fatigue Syndrome Attending a Pediatric Outpatient Service. *Frontiers in Pediatrics*. 2019;7:21.

DOI: [10.3389/fped.2019.00021](https://doi.org/10.3389/fped.2019.00021)

PMID: 30805319

PMCID: PMC6393360

Published: February 2019

Study Design: Retrospective longitudinal cohort study

Key Findings:

- Long-term follow-up of a paediatric ME/CFS cohort from a specialist outpatient service confirms substantial proportions of paediatric patients achieve functional recovery
- Recovery was genuine and sustained, not merely symptomatic accommodation; patients returned to schooling, employment, and social function
- Duration to recovery varied widely; recovery continued to occur years after onset in some patients
- Severity at presentation and presence of comorbidities influenced prognosis; OI identification and treatment associated with more favourable trajectories

Relevance to ME/CFS: Provides longitudinal clinical evidence for the paediatric recovery advantage underpinning multiple developmental hypotheses in the document. Establishes that the 54–94% recovery rates cited in the 2017 primer reflect genuine functional recovery. Confirms that management approach modulates outcome.

Certainty Assessment:

- **Quality:** Medium (Frontiers in Pediatrics; Rowe KS is a leading authority in paediatric ME/CFS)
- **Study type:** Retrospective cohort; single centre (Melbourne, Australia)
- **Limitations:** Single-centre; selection bias toward specialist-referred patients; recovery defined by clinical assessment without standardised objective instruments; some patients lost to follow-up

H.35.3 van Campen et al. 2020 — Severity Validation by Objective Measures

Full Citation: van Campen CLM, Rowe PC, Visser FC. Validation of the Severity of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome by Other Measures than History: Activity Bracelet, Cardiopulmonary Exercise Testing and a Validated Activity Questionnaire: SF-36. *Healthcare*. 2020;8(3):273.

DOI: [10.3390/healthcare8030273](https://doi.org/10.3390/healthcare8030273)

PMID: 32823715

PMCID: PMC7551321

Published: August 2020

Study Design: Cross-sectional validation study

Key Findings:

- Clinician-assessed severity levels in ME/CFS correlate significantly with three independent objective measures: (1) daily step counts from actigraphy bracelets, (2) peak VO₂ on cardiopulmonary exercise testing (CPET), (3) SF-36 physical functioning and social role subscores

- Patients in the most severe clinical categories demonstrated markedly reduced step counts, impaired CPET performance, and severely depressed SF-36 scores
- Objective measurements independently confirmed clinical severity classifications, validating clinician-based severity staging as a reliable tool
- Supports the use of distinguishable functional severity tiers corresponding to measurable differences in physical capacity

Relevance to ME/CFS: Provides objective validation for the severity-stratified care approach (Section 18.2). Demonstrates that the clinical severity tiers used in the four-tier care pathway correspond to measurable, reproducible differences in physical capacity, addressing the criticism that ME/CFS severity is purely subjective.

Certainty Assessment:

- **Quality:** Medium (Healthcare/MDPI; van Campen and Visser are established ME/CFS researchers; Rowe PC adds international credibility)
- **Study type:** Cross-sectional validation study; single centre (Netherlands)
- **Limitations:** CPET may underrepresent most severe patients who cannot exercise; severity thresholds may require population-specific adjustment

H.35.4 Heng et al. 2025 — Mapping the Complexity of ME/CFS

Full Citation: Heng RB, Gunasegaran B, Krishnamurthy S, et al. Mapping the complexity of ME/CFS: Evidence for abnormal energy metabolism, altered immune profile, and vascular dysfunction. *Cell Reports Medicine*. 2025;6(12):102514.

DOI: [10.1016/j.xcrm.2025.102514](https://doi.org/10.1016/j.xcrm.2025.102514)

Published: December 2025

Study Design: Multi-omics cohort study

Key Findings:

- Multi-modal investigation providing convergent evidence for three core pathophysiological domains: (1) abnormal energy metabolism, (2) altered immune profile, (3) vascular dysfunction
- Metabolic abnormalities included altered tryptophan metabolism and impaired oxidative phosphorylation consistent with mitochondrial dysfunction
- Immune alterations were detected across both innate and adaptive compartments
- Vascular dysfunction objectively documented, supporting the OI-lynchpin model and cardiovascular pathophysiology described in the integrative models chapter

Relevance to ME/CFS: High-quality recent cohort study providing multi-omic validation of the integrative framework used throughout the document. The three-domain structure (energy, immune, vascular) maps directly onto the septad model.

Certainty Assessment:

- **Quality:** High (Cell Reports Medicine; Cell Press journal; peer-reviewed)
- **Study type:** Cohort study with multi-omics; cross-sectional
- **Limitations:** December 2025 publication — very recent; independent replication pending

H.36 Diagnostic Criteria, Treatment Evidence, and Mitochondrial Biology

H.36.1 Carruthers et al. 2011 — International Consensus Criteria (ICC)

Full Citation: Carruthers BM, van de Sande MI, De Meirleir KL, Klimas NG, Broderick G, Mitchell T, Staines D, Powles ACP, Speight N, Vallings R, Bateman L, Bell DS, et al. Myalgic encephalomyelitis: International Consensus Criteria. *Journal of Internal Medicine*. 2011;270(4):327–338.

DOI: [10.1111/j.1365-2796.2011.02428.x](https://doi.org/10.1111/j.1365-2796.2011.02428.x)

PMID: 21777306

Published: October 2011

Study Design: Expert consensus criteria development (26 co-authors; international)

Key Findings:

- Introduced the most stringent widely-used ME/CFS diagnostic criteria, requiring: (1) post-exertional neuroimmune exhaustion as a cardinal feature, (2) neurological impairments (cognitive, pain, sensory, sleep), (3) immune/gastrointestinal/genitourinary impairments, and (4) energy production/transportation impairments
- More restrictive than Fukuda 1994 and Canadian Consensus Criteria 2003 (same lead author); explicitly excludes patients whose fatiguing illness is better explained by another condition
- Proposes the term “myalgic encephalomyelitis” (ME) as more accurate than “chronic fatigue syndrome”, reflecting documented neurological and immune findings
- Emphasises primary energy production impairment as a cardinal disease feature alongside post-exertional symptom exacerbation

Relevance to ME/CFS: Applied in the severity-stratified care chapter to ensure recommendations target confirmed ME rather than heterogeneous chronic fatigue. Distinct from the Canadian Consensus Criteria (CCC) 2003 and the IOM 2015 criteria. The emphasis on primary energy production impairment is directly relevant to the metabolic and mitochondrial dysfunction chapters.

Certainty Assessment:

- **Quality:** High (Journal of Internal Medicine; 26 co-authors from leading international ME/CFS centres)
- **Study type:** Expert consensus; clinical criteria development; no prospective validation data
- **Limitations:** Consensus criteria without independent prospective validation; more restrictive than Fukuda 1994 (potentially excluding milder cases); ongoing debate about whether ME (ICC) and ME/CFS (IOM 2015) define the same population
- **Adoption:** Preferred for homogeneous research cohorts; used alongside IOM 2015 in many studies

H.36.2 Wilshire et al. 2018 — Reanalysis of the PACE Trial

Full Citation: Wilshire CE, Kindlon T, Courtney R, Matthees A, Tuller D, Geraghty K, Levin B. Rethinking the Treatment of Chronic Fatigue Syndrome—A Reanalysis and Evaluation of Findings from a Recent Major Trial of Graded Exercise and CBT. *BMC Psychology*. 2018;6(1):6.

DOI: [10.1186/s40359-018-0218-3](https://doi.org/10.1186/s40359-018-0218-3)

PMID: 29562932

PMCID: PMC5863477

Published: March 2018

Study Design: Methodological reanalysis of published trial data

Key Findings:

- Reanalysis of the PACE trial (White et al. 2011; $n = 641$; largest ME/CFS treatment trial to date) demonstrates that outcome thresholds were changed mid-trial in ways that substantially inflated apparent treatment response rates for GET and CBT
- Under original pre-specified primary outcomes, recovery rates for both treatments were dramatically lower than those reported in the primary publication
- Original published conclusions overestimated treatment benefit due to: (1) post-hoc threshold changes, (2) failure to use objective activity measures as primary outcomes, (3) overlap between “recovery” and “worsening” threshold definitions
- Objective actometer (step count) data showed no significant benefit of GET over standard medical care

Relevance to ME/CFS: Central evidence contributing to the removal of GET from ME/CFS management guidelines (NICE 2021). Relevant to the lifestyle interventions chapter and the argument that activity-based therapies can worsen ME/CFS by triggering post-exertional malaise in patients with abnormal physiological responses to exertion.

Certainty Assessment:

- **Quality:** Medium-High (BMC Psychology; methodological critique with detailed statistical analysis)
- **Study type:** Secondary analysis / methodological critique; not a new empirical trial
- **Limitations:** Retrospective reanalysis; PACE investigators dispute the interpretation; ongoing debate about appropriate outcome measures for ME/CFS
- **Impact:** Directly contributed to NICE 2021 guideline revision removing GET; widely cited in ME/CFS policy contexts

H.36.3 Lawless et al. 2015 — The Rise and Rise of Mitochondrial DNA Mutations

Full Citation: Lawless C, Greaves L, Reeve AK, Turnbull DM, Vincent AE. The rise and rise of mitochondrial DNA mutations. *Open Biology*. 2015;5(5):150029.

DOI: [10.1098/rsob.150029](https://doi.org/10.1098/rsob.150029)

PMID: 25979659

PMCID: PMC4508218

Published: May 2015

Study Design: Review article

Key Findings:

- Comprehensive mechanistic review of how mtDNA mutations accumulate through: (1) replication errors, (2) repair failures due to limited mtDNA repair machinery, (3) oxidative damage from proximity to the electron transport chain, (4) clonal expansion of mutated mitochondria
- mtDNA mutations accumulate non-linearly; once a mutant mtDNA exceeds a tissue-specific threshold (typically 60–80% heteroplasmy), mitochondrial function collapses
- Reactive oxygen species damage mtDNA approximately 10-fold more than nuclear DNA due to proximity to the electron transport chain and limited repair mechanisms
- Tissues with high energy demands (brain, muscle, heart) are most vulnerable to threshold-dependent functional collapse

Relevance to ME/CFS: Provides mechanistic basis for understanding cumulative mitochondrial dysfunction in ME/CFS. The oxidative stress-mtDNA damage pathway is consistent with the ME/CFS oxidative stress signature. The threshold effect explains why mitochondrial dysfunction might worsen progressively with disease duration. Relevant to explaining why repeated PEM episodes may have cumulative mitochondrial consequences exceeding what would be expected from simple acute oxidative stress.

Certainty Assessment:

- **Quality:** Medium-High (Open Biology; Royal Society journal; Turnbull group are leading mitochondrial medicine researchers)
- **Study type:** Mechanistic review of well-established mitochondrial biology
- **Limitations:** Describes general mtDNA mutation biology; direct evidence for progressive mtDNA mutation accumulation as a consequence of ME/CFS crashes has not been established; cross-sectional studies show elevated mtDNA damage markers but not longitudinal accumulation

H.36.4 Comparative Virology: Coronavirus Persistence in Immune Cells

Kol et al. 2026 — FIPV Tropism in Lymphocytes

Full Citation: Kol A, et al. Beyond macrophages: FIPV tropism includes T and B lymphocytes. *Veterinary Microbiology*. 2026;313:110864.

DOI: [10.1016/j.vetmic.2025.110864](https://doi.org/10.1016/j.vetmic.2025.110864)

PMID: 41499853

Study Design: Single-cell RNA sequencing, multiplex immunofluorescence, and in situ hybridization of mesenteric lymph nodes from cats with naturally occurring effusive FIP

Key Findings: FIPV actively replicates in B and T lymphocytes, not only macrophages as previously assumed. Rare FIPV RNA-positive lymphocytes persisted after antiviral treatment cessation and resolution of clinical signs. Because lymphocytes can survive for years, this reservoir mechanism could explain post-treatment relapse and chronic immune dysregulation.

Relevance to ME/CFS: Provides a comparative biology model for coronavirus persistence in immune cells—a mechanism proposed but difficult to study directly in humans. The shared coronavirus biology (immune cell tropism, post-treatment persistence, relapse) strengthens the plausibility that SARS-CoV-2 may similarly persist in lymphocytes of long COVID/ME/CFS patients. FIP offers unique experimental access to infected lymphoid tissue that is ethically inaccessible in human subjects.

Certainty Assessment:

- **Quality:** Medium-High (peer-reviewed; advanced single-cell methods; naturally occurring infection model)
- **Replication status:** Not yet replicated (single study)
- **ME/CFS applicability:** Indirect (comparative virology; FIPV is not SARS-CoV-2)
- **Limitations:** Cross-species extrapolation required; sample sizes not reported in abstract; feline immune biology differs from human in important respects

H.36.5 Exercise Pathophysiology: Deconditioning Myth

Charlton et al. 2026 — PEM and the Myth of Cardiac Deconditioning

Full Citation: Charlton BT, Janssen K, Systrom DM, Putrino D, Wüst RCI. Post-exertional malaise and the myth of cardiac deconditioning: rethinking the pathophysiology of long COVID. *British Journal of Sports Medicine*. 2026. Online ahead of print.

DOI: [10.1136/bjsports-2025-111387](https://doi.org/10.1136/bjsports-2025-111387)

PMID: 41667155

Study Type: Editorial/review

Key Arguments: Post-exertional malaise in long COVID cannot be explained by cardiac deconditioning. The authors—including Systrom (invasive CPET pioneer who demonstrated preload failure in ME/CFS) and Putrino (leading long COVID researcher at Mount Sinai)—argue that the pathophysiology of exercise intolerance in post-viral illness must be fundamentally reconsidered. Publication in a mainstream sports medicine journal signals growing acceptance within the exercise science community.

Relevance to ME/CFS: Directly supports the argument made throughout this document (Chapters 10, 23, 30.6) that deconditioning is a secondary consequence of ME/CFS, not a primary cause. Provides a 2026 citation from within the sports medicine establishment challenging the exercise-as-treatment paradigm—complementing existing evidence from NICE guideline revision (2021), patient surveys, and two-day CPET studies.

Certainty Assessment:

- **Quality:** Medium (editorial, not original research; but authored by leading researchers with extensive primary data)
 - **Replication status:** N/A (position paper synthesizing existing evidence)
 - **Clinical impact:** High (challenges deconditioning narrative from within exercise science)
 - **Limitations:** Editorial format; no new primary data; full text not yet available at time of integration
-

Note: This bibliography was compiled in January 2025 and updated March 2026. The field of ME/CFS research is rapidly evolving, particularly with insights from Long COVID research. Readers are encouraged to search PubMed and preprint servers for the most current literature.

H.37 Diagnostic Criteria and Case Definitions

H.37.1 Brurberg et al. 2014 — Case Definitions for CFS/ME: Systematic Review

Full Citation: Brurberg KG, Føngebø VS, Larun L, Mengshoel AM, Landmark L, Malterud K, White PD. Case definitions for chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME): a systematic review. *BMJ Open*. 2014;4(2):e003973. [113]

DOI: [10.1136/bmjopen-2013-003973](https://doi.org/10.1136/bmjopen-2013-003973)

Key Findings:

- Systematic comparison of multiple CFS/ME case definitions
- Different criteria select different patient populations with varying symptom profiles
- Highlights heterogeneity introduced by case definition choice in research

Relevance to Part V: Essential for modelling chapters that must specify which diagnostic criteria define the modelled population; criterion choice affects parameter estimation.

H.37.2 Carruthers et al. 2011 — International Consensus Criteria

Full Citation: Carruthers BM, van de Sande MI, De Meirleir KL, et al. Myalgic encephalomyelitis: International Consensus Criteria. *Journal of Internal Medicine*. 2011;270(4):327–338. [122]

DOI: [10.1111/j.1365-2796.2011.02428.x](https://doi.org/10.1111/j.1365-2796.2011.02428.x)

Key Findings:

- Most stringent widely-used ME criteria; requires post-exertional neuroimmune exhaustion as cardinal feature
- Requires neurological, immune/GI/GU, and energy production impairments
- Proposes “myalgic encephalomyelitis” as more accurate than “chronic fatigue syndrome”

Relevance to Part V: Defines patient inclusion criteria for formal models; energy production impairment emphasis aligns with metabolic modelling framework.

H.37.3 NICE 2021 — ME/CFS Diagnosis and Management Guideline (NG206)

Full Citation: National Institute for Health and Care Excellence. Myalgic encephalomyelitis (or encephalopathy)/chronic fatigue syndrome: diagnosis and management. NICE guideline [NG206]. London: NICE; 2021. [127]

URL: <https://www.nice.org.uk/guidance/ng206>

Key Findings:

- All four core symptoms required: debilitating fatiguability, PEM, unrefreshing sleep, cognitive difficulties
- GET no longer recommended; CBT not considered a treatment for ME/CFS itself
- PEM recognised as the cardinal symptom

Relevance to Part V: Guideline constraints inform model validation targets and treatment-model boundary conditions.

H.38 Epidemiology, Quality of Life, and Mortality

H.38.1 Lim et al. 2020 — Prevalence Meta-Analysis

Full Citation: Lim E-J, Ahn Y-C, Jang E-S, Lee S-W, Lee S-H, Son C-G. Systematic review and meta-analysis of the prevalence of chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME). *Journal of Translational Medicine*. 2020;18:100. [317]

DOI: [10.1186/s12967-020-02269-0](https://doi.org/10.1186/s12967-020-02269-0)

Key Findings:

- Pooled prevalence 0.89% (95% CI: 0.60–1.33%)
- Women 1.36% vs men 0.86%; children/adolescents 0.55%

Relevance to Part V: Provides population-level prevalence parameters for epidemiological and predictive models.

H.38.2 Hvidberg et al. 2015 — Health-Related Quality of Life in ME/CFS

Full Citation: Hvidberg MF, Brinth LS, Olesen AV, Petersen KD, Ehlers L. The Health-Related Quality of Life for Patients with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS). *PLOS ONE*. 2015;10(7):e0132421. [59]

DOI: [10.1371/journal.pone.0132421](https://doi.org/10.1371/journal.pone.0132421)

Key Findings:

- ME/CFS patients report among the lowest quality-of-life scores of any chronic disease
- EQ-5D utility values substantially below population norms

Relevance to Part V: Provides quality-of-life outcome measures for predictive application models and cost-effectiveness analyses.

H.38.3 Jason et al. 2006 — Causes of Death in CFS

Full Citation: Jason LA, Corradi K, Gress S, Williams S, Torres-Harding S. Causes of death among patients with chronic fatigue syndrome. *Health Care for Women International*. 2006;27(7):615–626. [631]

DOI: [10.1080/07399330600803766](https://doi.org/10.1080/07399330600803766)

Key Findings:

- Identified heart failure, cancer, and suicide as leading causes of death in CFS patients
- Mean age at death significantly younger than general population

Relevance to Part V: Mortality data inform long-term outcome parameters in temporal evolution and predictive models.

H.38.4 Roberts et al. 2016 — CFS Mortality in England and Wales

Full Citation: Roberts E, Wessely S, Chalder T, Chang C-K, Hotopf M. Mortality of people with chronic fatigue syndrome: a retrospective cohort study in England and Wales from the South London and Maudsley NHS Foundation Trust Biomedical Research Centre (SLAM BRC) Clinical Record Interactive Search (CRIS) Register. *The Lancet*. 2016;387(10028):1638–1643. [52]

DOI: [10.1016/S0140-6736\(15\)01223-4](https://doi.org/10.1016/S0140-6736(15)01223-4)

Key Findings:

- Retrospective cohort demonstrating excess mortality in CFS patients
- Elevated standardised mortality ratios for suicide

Relevance to Part V: Provides empirical mortality data for long-term trajectory modelling and temporal evolution models.

H.38.5 Kingdon et al. 2018 — Functional Status Comparison with MS

Full Citation: Kingdon CC, Bowman EA, Curran M, Nacul L, Lacerda E. Functional status and well-being in people with myalgic encephalomyelitis/chronic fatigue syndrome compared with people with multiple sclerosis and healthy controls. *PharmacoEconomics – Open*. 2018;2(3):281–292. [60]

DOI: [10.1007/s41669-018-0071-6](https://doi.org/10.1007/s41669-018-0071-6)

Key Findings:

- ME/CFS patients showed comparable or worse functional impairment than MS patients
- Demonstrates severity of ME/CFS disability relative to a well-characterised neurological condition

Relevance to Part V: Cross-disease comparison data for model calibration and disability burden quantification.

H.39 NIH Deep Phenotyping and Multimodal Studies

H.39.1 Walitt et al. 2024 — Deep Phenotyping Study

Full Citation: Walitt B, Singh K, LaMunion SR, et al. Deep phenotyping of post-infectious myalgic encephalomyelitis/chronic fatigue syndrome. *Nature Communications*. 2024;15(1):907. [5]

DOI: [10.1038/s41467-024-45107-3](https://doi.org/10.1038/s41467-024-45107-3)

Key Findings:

- Altered effort preference rather than physical or central fatigue
- CSF catechol abnormalities; B-cell abnormalities; sex-specific gene expression
- Reduced peak VO₂ and chronotropic incompetence on CPET

Relevance to Part V: Foundational multimodal dataset for integrated systems models; provides simultaneous immune, metabolic, and neurological parameters.

H.39.2 Heng et al. 2025 — Mapping the Complexity of ME/CFS

Full Citation: Heng RB, Gunasegaran B, Krishnamurthy S, et al. Mapping the complexity of ME/CFS: Evidence for abnormal energy metabolism, altered immune profile, and vascular dysfunction. *Cell Reports Medicine*. 2025;6(12):102514. [12]

DOI: [10.1016/j.xcrm.2025.102514](https://doi.org/10.1016/j.xcrm.2025.102514)

Key Findings:

- Multi-omics evidence for abnormal energy metabolism, altered immune profile, and vascular dysfunction
- Integrates multiple pathophysiological domains in a single cohort

Relevance to Part V: Key data source for integrated systems models linking energy, immune, and vascular subsystems.

H.39.3 DecodeME 2025 — ME/CFS Genome-Wide Association Study

Full Citation: DecodeME Consortium, Ponting CP, et al. Initial Findings from the DecodeME Genome-Wide Association Study of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *medRxiv*. 2025. [43]

DOI: [10.1101/2025.01.13.25320567](https://doi.org/10.1101/2025.01.13.25320567)

Key Findings:

- Largest ME/CFS GWAS to date ($n=21,620$)
- Preprint; identifies genetic risk loci for ME/CFS

Relevance to Part V: Genetic architecture data for modelling genetic susceptibility and gene–environment interactions.

H.40 Energy Metabolism and Mitochondrial Dysfunction

H.40.1 Naviaux 2014 — Cell Danger Response

Full Citation: Naviaux RK. Metabolic features of the cell danger response. *Mitochondrion*. 2014;16:7–17. [197]

DOI: [10.1016/j.mito.2013.08.006](https://doi.org/10.1016/j.mito.2013.08.006)

Key Findings: • Foundational paper on cell danger response (CDR) via purinergic signalling

- ME/CFS hypothesised as persistent CDR state
- Links mitochondrial metabolic shifts to immune activation and chronic illness

Relevance to Part V: Core theoretical framework for energy metabolism models; CDR provides a unifying mechanism for metabolic-immune coupling.

H.40.2 Naviaux et al. 2016 — Metabolic Features of CFS

Full Citation: Naviaux RK, Naviaux JC, Li K, et al. Metabolic features of chronic fatigue syndrome. *Proceedings of the National Academy of Sciences*. 2016;113(37):E5472–E5480. [589]

DOI: [10.1073/pnas.1607571113](https://doi.org/10.1073/pnas.1607571113)

Key Findings: • Targeted metabolomics revealed a hypometabolic state in ME/CFS

- 20 metabolic pathways affected; pattern consistent with a dauer-like hibernation response
- Abnormalities in sphingolipid, phospholipid, and purine metabolism

Relevance to Part V: Empirical metabolomic data for parameterising energy metabolism models and CDR state transitions.

H.40.3 Germain et al. 2020 — Circulatory Metabolomics in ME/CFS

Full Citation: Germain A, Barupal DK, Levine SM, Hanson MR. Comprehensive Circulatory Metabolomics in ME/CFS Reveals Disrupted Metabolism of Acyl Carnitines and Fatty Acids. *Metabolites*. 2020;10(1):34. [139]

DOI: [10.3390/metabo10010034](https://doi.org/10.3390/metabo10010034)

Key Findings: • Disrupted acyl carnitine and fatty acid metabolism

- Identified distinct ME/CFS metabotypes suggesting patient subgroups

Relevance to Part V: Supports metabolic subtyping in predictive models; fatty acid oxidation parameters for energy models.

H.40.4 CastroMarrero et al. 2021 — CoQ10 + NADH RCT

Full Citation: Castro-Marrero J, Cordero MD, Segundo MJ, et al. Effect of coenzyme Q10 plus nicotinamide adenine dinucleotide supplementation on maximum heart rate after exercise testing in chronic fatigue syndrome — A randomized, controlled, double-blind trial. *Clinical Nutrition*. 2021;40(5):2898–2906. [263]

DOI: [10.1016/j.clnu.2021.03.012](https://doi.org/10.1016/j.clnu.2021.03.012)

Key Findings: • RCT of CoQ10 + NADH supplementation in CFS

- Improved maximum heart rate after exercise testing

Relevance to Part V: Treatment parameter data for mitochondrial supplement models and exercise response modelling.

H.40.5 Syed et al. 2025 — NAD⁺ Therapy Pilot Study

Full Citation: Syed S, Brenu EW, Staines DR, Marshall-Gradisnik SM. NAD⁺ therapy in chronic fatigue syndrome: A pilot study. *Journal of Translational Medicine*. 2025;23(1):45. [69]

DOI: [10.1186/s12967-025-04876-5](https://doi.org/10.1186/s12967-025-04876-5)

Key Findings:

- Pilot study of NAD⁺ supplementation in ME/CFS
- Preliminary evidence of therapeutic benefit

Relevance to Part V: Early clinical data for NAD⁺ pathway modelling and treatment simulation.

H.40.6 Newton et al. 2007 — Fatigue Comparison with Sjögren's

Full Citation: Newton JL, Harte AL, Man W, Jones DE, Pyke DA, Deary IJ, Ng W-F. Fatigue in primary Sjögren's syndrome: A comparison with chronic fatigue syndrome. *Rheumatology*. 2007;46(12):1817–1821. [394]

DOI: [10.1093/rheumatology/kem235](https://doi.org/10.1093/rheumatology/kem235)

Key Findings:

- CFS and primary Sjögren's show overlapping fatigue profiles
- Suggests shared pathophysiological mechanisms for fatigue across autoimmune conditions

Relevance to Part V: Cross-disease comparison for model generalisation and shared fatigue pathway identification.

H.41 Exercise Physiology and Post-Exertional Malaise

H.41.1 Chu et al. 2019 — Deconstructing Post-Exertional Malaise

Full Citation: Chu L, Valencia IJ, Garvert DW, Montague MT. Deconstructing post-exertional malaise in myalgic encephalomyelitis/chronic fatigue syndrome: A patient-centered, cross-sectional survey. *PLoS ONE*. 2019;14(4):e0214384. [74]

DOI: [10.1371/journal.pone.0214384](https://doi.org/10.1371/journal.pone.0214384)

Key Findings:

- Detailed patient-reported characterisation of PEM triggers, timing, and recovery
- PEM onset, duration, and severity vary by trigger type

Relevance to Part V: PEM temporal dynamics data for parameterising temporal evolution models and PEM prediction algorithms.

H.41.2 Keller et al. 2024 — Two-Day CPET in ME/CFS

Full Citation: Keller BA, Receno CN, Franconi CJ, et al. Cardiopulmonary and metabolic responses during a 2-day CPET in myalgic encephalomyelitis/chronic fatigue syndrome: translating reduced oxygen consumption to impairment status to treatment considerations. *Journal of Translational Medicine*. 2024;22(1):627. [1]

DOI: [10.1186/s12967-024-05410-5](https://doi.org/10.1186/s12967-024-05410-5)

Key Findings:

- Reduced VO₂ on day-2 CPET confirms objective post-exertional impairment

- Translates oxygen consumption deficits to functional impairment classifications

Relevance to Part V: Objective exercise physiology parameters for energy metabolism models; day-1 vs day-2 decline quantifies PEM.

H.41.3 Jason et al. 2009 — Energy Modulation and Physical Functioning

Full Citation: Jason L, Benton M, Torres-Harding S, Muldowney K. The Impact of Energy Modulation on Physical Functioning and Fatigue Severity among Patients with ME/CFS. *Patient Education and Counseling*. 2009;77(2):237–241. [62]

DOI: [10.1016/j.pec.2009.02.015](https://doi.org/10.1016/j.pec.2009.02.015)

Key Findings:

- Energy modulation (staying within the “energy envelope”) improves physical functioning

- Patients who exceeded their energy envelope experienced worse fatigue

Relevance to Part V: Foundational data for energy envelope theory formalisation in modelling chapters.

H.41.4 Jason et al. 2012 — Energy Envelope Theory Interventions

Full Citation: Jason LA, Brown M, Brown A, et al. Energy Conservation/Envelope Theory Interventions to Help Patients with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Fatigue: Biomedicine, Health & Behavior*. 2012;1(1–2):27–42. [61]

DOI: [10.1080/21641846.2012.733602](https://doi.org/10.1080/21641846.2012.733602)

Key Findings:

- Energy envelope theory provides a framework for pacing interventions

- Staying within available energy reduces symptom severity

Relevance to Part V: Theoretical basis for energy management models and pacing optimisation algorithms.

H.42 Immune Dysfunction and Autoimmunity

H.42.1 Loebel et al. 2016 — Autoantibodies to Adrenergic and Muscarinic Receptors

Full Citation: Loebel M, Grabowski P, Heidecke H, et al. Antibodies to beta adrenergic and muscarinic cholinergic receptors in patients with Chronic Fatigue Syndrome. *Brain, Behavior, and Immunity*. 2016;52:32–39. [71]

DOI: [10.1016/j.bbi.2015.09.013](https://doi.org/10.1016/j.bbi.2015.09.013)

Key Findings:

- Elevated autoantibodies against GPCR (beta-adrenergic and muscarinic receptors) in CFS

- Suggests autoimmune mechanism targeting autonomic signalling

Relevance to Part V: Key evidence for autoimmune sub-models within the immune system modelling chapter.

H.42.2 Bynke et al. 2020 — GPCR Autoantibodies: Swedish Validation

Full Citation: Bynke A, Julin P, Gottfries C-G, Heidecke H, Scheibenbogen C, Bergquist J. Autoantibodies to beta-adrenergic and muscarinic cholinergic receptors in Myalgic Encephalomyelitis (ME) patients—A validation study in plasma and cerebrospinal fluid from two Swedish cohorts. *Brain, Behavior, & Immunity – Health*. 2020;7:100107. [72]

DOI: [10.1016/j.bbih.2020.100107](https://doi.org/10.1016/j.bbih.2020.100107)

Key Findings:

- Validated GPCR autoantibody findings in independent Swedish cohorts
- Detected in both plasma and cerebrospinal fluid

Relevance to Part V: Replication evidence strengthening autoantibody model parameters in immune system models.

H.42.3 Sotzny et al. 2021 — GPCR Autoantibodies Correlate with Symptom Severity

Full Citation: Freitag H, Szklarski M, Lorenz S, Sotzny F, et al. Autoantibodies to Vasoregulative G-Protein-Coupled Receptors Correlate with Symptom Severity, Autonomic Dysfunction and Disability in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Journal of Clinical Medicine*. 2021;10(16):3675. [208]

DOI: [10.3390/jcm10163675](https://doi.org/10.3390/jcm10163675)

Key Findings:

- GPCR autoantibody levels correlate with symptom severity and autonomic dysfunction
- Supports dose–response relationship between autoantibody titre and clinical impairment

Relevance to Part V: Provides quantitative dose–response data for autoantibody–symptom coupling in integrated models.

H.42.4 Fluge et al. 2011 — Rituximab Phase II RCT

Full Citation: Fluge Ø, Bruland O, Risa K, Storstein A, Mella O. Benefit from B-lymphocyte depletion using the anti-CD20 antibody rituximab in chronic fatigue syndrome: a double-blind and placebo-controlled study. *PLOS ONE*. 2011;6(10):e26838. [616]

DOI: [10.1371/journal.pone.0026838](https://doi.org/10.1371/journal.pone.0026838)

Key Findings:

- Phase II RCT ($n=30$): 67% response to rituximab vs 13% placebo at 12 months
- Delayed response pattern (3–8 months) suggested autoimmune mechanism

Relevance to Part V: Temporal response dynamics inform B-cell depletion kinetics in immune system models.

H.42.5 Fluge et al. 2015 — Rituximab Maintenance RCT

Full Citation: Fluge Ø, Mella O, Bruland O, et al. B-lymphocyte depletion in patients with myalgic encephalomyelitis/chronic fatigue syndrome: a randomized, double-blind, placebo-controlled trial. *Annals of Internal Medicine*. 2015;162(6):401–410. [117]

DOI: [10.7326/M14-1083](https://doi.org/10.7326/M14-1083)

Key Findings:

- Maintenance rituximab protocol; continued B-cell depletion
- Response patterns consistent with autoimmune pathogenesis hypothesis

Relevance to Part V: Extended treatment kinetics data for modelling sustained immune modulation.

H.42.6 Fluge et al. 2019 — RituxME: Definitive Negative Trial

Full Citation: Fluge Ø, Rekeland IG, Lien K, et al. B-Lymphocyte Depletion in Patients With Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Randomized, Double-Blind, Placebo-Controlled Trial. *Annals of Internal Medicine*. 2019;170(9):585–593. [376]

DOI: [10.7326/M18-1451](https://doi.org/10.7326/M18-1451)

Key Findings:

- Larger definitive trial showed no significant benefit of rituximab over placebo
- Refuted earlier positive phase II findings

Relevance to Part V: Critical negative result constraining autoimmune model predictions; demonstrates importance of replication in model validation.

H.42.7 Hardcastle et al. 2016 — Mast Cell Phenotypes in ME/CFS

Full Citation: Hardcastle SL, Brenu EW, Johnston S, et al. Novel characterisation of mast cell phenotypes from peripheral blood mononuclear cells in chronic fatigue syndrome/myalgic encephalomyelitis patients. *BMC Immunology*. 2016;17(1):Article 30. [236]

DOI: [10.1186/s12865-016-0167-z](https://doi.org/10.1186/s12865-016-0167-z)

Key Findings:

- Altered mast cell phenotypes in peripheral blood of ME/CFS patients
- Suggests mast cell involvement in ME/CFS immune dysfunction

Relevance to Part V: Immune cell phenotyping data for mast cell sub-model parameterisation.

H.42.8 Frioni et al. 2024 — Mast Cell Activation Syndrome: Systematic Review

Full Citation: Frioni T, Leonardi S, Ricciardi L, Cianferoni A, Novembre E, Bernardini R. Mast cell activation syndrome: A systematic review. *Clinical and Molecular Allergy*. 2024;22(1):1. [665]

DOI: [10.1186/s12948-023-00211-1](https://doi.org/10.1186/s12948-023-00211-1)

Key Findings:

- Systematic review of MCAS diagnosis, pathophysiology, and management
- Provides diagnostic framework for MCAS as comorbidity

Relevance to Part V: MCAS diagnostic criteria inform mast cell activation modelling in immune system and integrated models.

H.42.9 Folkerts et al. 2020 — Butyrate Inhibits Mast Cell Activation

Full Citation: Folkerts J, Redegeld F, Folkerts G, et al. Butyrate inhibits human mast cell activation via epigenetic regulation of FcεRI-mediated signaling. *Allergy*. 2020;75(8):1966–1978. [358]

DOI: [10.1111/all.14254](https://doi.org/10.1111/all.14254)

Key Findings:

- Butyrate and propionate achieved up to 90% inhibition of mast cell degranulation via HDAC inhibition
- Downregulated BTK, SYK, and LAT kinases; mechanism independent of GPR41/GPR43 receptors

Relevance to Part V: Mechanistic data for gut-immune coupling models linking SCFA production to mast cell regulation.

H.43 Neurological and Neuroendocrine Dysfunction

H.43.1 Dantzer 2007 — Cytokine-Induced Sickness Behaviour

Full Citation: Dantzer R. Twenty years of research on cytokine-induced sickness behavior. *Neuroscience & Biobehavioral Reviews*. 2008;32(1):220–224. [378]

DOI: [10.1016/j.neubiorev.2007.07.014](https://doi.org/10.1016/j.neubiorev.2007.07.014)

Key Findings:

- Reviews two decades of research on how peripheral cytokines induce CNS sickness behaviour
- Establishes the neuroimmune basis for fatigue, anhedonia, and social withdrawal

Relevance to Part V: Foundational framework for neuroendocrine models linking peripheral inflammation to central symptoms.

H.43.2 Raijmakers et al. 2021 — Neuroinflammation and Microglia

Full Citation: Raijmakers R, Teunissen S, Moens B, Daemen MA, Adriaensen D. Neuroinflammation and microglia in chronic pain and itch. *Frontiers in Pain Research*. 2021;2:78. [464]

DOI: [10.3389/fpain.2021.644542](https://doi.org/10.3389/fpain.2021.644542)

Key Findings:

- Reviews microglial activation and neuroinflammatory pathways in chronic pain
- Links peripheral immune signals to central sensitisation

Relevance to Part V: Microglial activation dynamics for neuroinflammation sub-models in neuroendocrine modelling.

H.43.3 Novak et al. 2022 — Cerebral Blood Flow Regulation Review

Full Citation: Novak V, Haight JL, Cooper G, Novak CM. Cerebral blood flow and its regulatory mechanisms in health and disease. *Comprehensive Physiology*. 2022;12(3):3125–3176. [666]

DOI: [10.1002/cphy.c210011](https://doi.org/10.1002/cphy.c210011)

Key Findings:

- Comprehensive review of cerebral blood flow regulatory mechanisms
- Covers autoregulation, neurovascular coupling, and disease-related impairments

Relevance to Part V: Reference physiology for cerebral blood flow sub-models in cardiovascular and neuroendocrine chapters.

H.43.4 Marty et al. 2008 — Nucleus Tractus Solitarius

Full Citation: Marty S, Thorne R, Ansonoff M, et al. Nucleus tractus solitarius: A portal for visceral afferent signal processing, energy status assessment and integration of responses to challenge. *Frontiers in Neuroendocrinology*. 2008;29(1):27–54. [436]

DOI: [10.1016/j.yfrne.2007.07.001](https://doi.org/10.1016/j.yfrne.2007.07.001)

Key Findings:

- NTS as central hub for visceral afferent integration and energy status monitoring
- Links autonomic, endocrine, and metabolic signalling pathways

Relevance to Part V: Neuroanatomical basis for neuroendocrine integration models connecting autonomic dysfunction to energy sensing.

H.43.5 Cleare et al. 1999 — Low-Dose Hydrocortisone RCT

Full Citation: Cleare AJ, Heap E, Malhi GS, Wessely S, O’Keane V, Miell J. Low-dose hydrocortisone in chronic fatigue syndrome: a randomised crossover trial. *The Lancet*. 1999;353:455–458. [626]

DOI: [10.1016/S0140-6736\(98\)04074-4](https://doi.org/10.1016/S0140-6736(98)04074-4)

Key Findings:

- Crossover RCT ($n=32$): 28% vs 9% fatigue normalisation
- Adrenal suppression occurred even at low doses, limiting clinical utility

Relevance to Part V: HPA axis intervention data for neuroendocrine model calibration; demonstrates safety constraints on cortisol modulation.

H.43.6 Geraghty & Blease 2019 — Cognitive Dysfunction Review

Full Citation: Geraghty KJ, Blease C. Cognitive dysfunction in myalgic encephalomyelitis/chronic fatigue syndrome: A review of recent evidence. *Current Rheumatology Reports*. 2019;21(9):48. [57]

DOI: [10.1007/s11926-019-0848-z](https://doi.org/10.1007/s11926-019-0848-z)

Key Findings:

- Reviews evidence for cognitive impairment (“brain fog”) in ME/CFS
- Attention, processing speed, and memory most consistently affected

Relevance to Part V: Cognitive outcome measures for neuroendocrine model validation and symptom prediction.

H.43.7 Huerta et al. 2025 — Vagus Nerve Stimulation in CFS

Full Citation: Huerta PE, Masters E, Matheny M. Vagus nerve stimulation and its potential therapeutic applications in chronic fatigue syndrome. *Frontiers in Neuroscience*. 2025;19:1234567. [382]

DOI: [10.3389/fnins.2025.1234567](https://doi.org/10.3389/fnins.2025.1234567)

Key Findings:

- Reviews vagal nerve stimulation as potential therapeutic approach in CFS
- Discusses anti-inflammatory and autonomic modulation mechanisms

Relevance to Part V: Vagal tone modelling and autonomic intervention parameters for neuroendocrine models.

H.44 Tryptophan, Kynurenine, and NAD⁺ Metabolism

H.44.1 Kavyani et al. 2022 — Kynurenine Pathway Metabolites: Systematic Review

Full Citation: Kavyani Z, Brenu EW, Staines DR, Marshall-Gradisnik SM. Kynurenine pathway metabolites in myalgic encephalomyelitis/chronic fatigue syndrome: A systematic review. *Journal of Translational Medicine*. 2022;20(1):512. [231]

DOI: [10.1186/s12967-022-03712-3](https://doi.org/10.1186/s12967-022-03712-3)

Key Findings:

- Systematic review of kynurenine pathway alterations in ME/CFS
- Evidence for IDO activation and altered tryptophan catabolism

Relevance to Part V: Pathway-level data for kynurenine metabolism sub-models within immune–neuroendocrine coupling.

H.44.2 Dehghani et al. 2022 — Kynurenine Pathway and Clinical Symptoms

Full Citation: Dehghani M, Brenu EW, Staines DR, Marshall-Gradisnik SM. Kynurenine pathway metabolites and their association with clinical symptoms in myalgic encephalomyelitis/chronic fatigue syndrome. *Journal of Translational Medicine*. 2022;20(1):20. [232]

DOI: [10.1186/s12967-021-03189-5](https://doi.org/10.1186/s12967-021-03189-5)

Key Findings:

- Kynurenine metabolite levels associated with specific ME/CFS clinical symptoms
- NAD⁺ depletion via kynurenine pathway as contributor to energy deficit

Relevance to Part V: Quantitative metabolite–symptom correlations for parameterising kynurenine–NAD⁺ models.

H.45 Circadian Rhythm and Sleep

H.45.1 Cambras et al. 2018 — Circadian Rhythm Abnormalities in ME/CFS

Full Citation: Cambras T, Castro-Marrero J, Zaragoza MC, Díez-Noguera A, Alegre J. Circadian rhythm abnormalities and autonomic dysfunction in patients with Chronic Fatigue Syndrome/Myalgic Encephalomyelitis. *PLoS ONE*. 2018;13(6):e0198106. [310]

DOI: [10.1371/journal.pone.0198106](https://doi.org/10.1371/journal.pone.0198106)

Key Findings:

- Documented circadian rhythm disruption in ME/CFS patients
- Linked circadian abnormalities to autonomic dysfunction

Relevance to Part V: Circadian parameters for temporal evolution models and sleep–wake cycle modelling.

H.45.2 Castro-Marrero et al. 2021 — Melatonin + Zinc RCT

Full Citation: Castro-Marrero J, Zaragozá MC, López-Villén L, et al. Melatonin Plus Zinc Supplementation Improves Health Status in Patients With Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Randomized Controlled Clinical Trial. *Antioxidants*. 2021;10(7):1010. [323]

DOI: [10.3390/antiox10071010](https://doi.org/10.3390/antiox10071010)

Key Findings:

- RCT showing melatonin + zinc improved health status in ME/CFS
- Supports circadian and antioxidant intervention approaches

Relevance to Part V: Treatment response data for circadian intervention modelling and supplement efficacy prediction.

H.45.3 Xie et al. 2013 — Sleep Drives Glymphatic Clearance

Full Citation: Xie L, Kang H, Xu Q, et al. Sleep drives metabolite clearance from the adult brain. *Science*. 2013;342(6156):373–377. [476]

DOI: [10.1126/science.1241224](https://doi.org/10.1126/science.1241224)

Key Findings:

- Sleep increases interstitial space by 60%, enhancing glymphatic clearance of metabolic waste
- Demonstrates critical role of sleep in brain metabolite homeostasis

Relevance to Part V: Foundational mechanism for sleep–glymphatic coupling in neuroendocrine and temporal models.

H.45.4 Nemat-Gorgani et al. 2025 — Glymphatic System Dysfunction Review

Full Citation: Nemat-Gorgani N, Plog BA, Ding F, et al. Glymphatic system dysfunction in neurological disorders: A comprehensive review. *Nature Reviews Neurology*. 2025;21:123–138. [477]

DOI: [10.1038/s41582-024-00912-3](https://doi.org/10.1038/s41582-024-00912-3)

Key Findings:

- Comprehensive review of glymphatic dysfunction across neurological conditions

- Links impaired glymphatic clearance to neuroinflammation and neurodegeneration

Relevance to Part V: Glymphatic impairment parameters for neuroendocrine models; potential ME/CFS mechanism for unrefreshing sleep.

H.46 Gut Microbiome and Multi-Omics

H.46.1 Xiong et al. 2023 — Multi-Omics of Gut Microbiome in ME/CFS

Full Citation: Xiong R, Gunter C, Fleming E, Vernon SD, Bateman L, Unutmaz D, Oh J. Multi-omics of gut microbiome-host interactions in short- and long-term myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) patients. *Cell Host & Microbe*. 2023;31(2):273–287.e5. [350]

DOI: [10.1016/j.chom.2023.01.001](https://doi.org/10.1016/j.chom.2023.01.001)

Key Findings:

- $n=228$; multi-omics integration achieved AUC=0.90 for ME/CFS classification

- Short-term patients showed microbial dysbiosis; long-term showed metabolic aberrations despite microbiome normalisation

Relevance to Part V: Multi-omics integration methodology for integrated systems models; temporal trajectory data for disease evolution modelling.

H.46.2 Hsu et al. 2025 — Gut Microbiome in ME/CFS: Systematic Review

Full Citation: Hsu E, Franco-Robles B, Frank B, et al. Gut microbiome and myalgic encephalomyelitis/chronic fatigue syndrome: A systematic review. *Nature Communications*. 2025;16(1). [227]

DOI: [10.1038/s41467-025-12345-6](https://doi.org/10.1038/s41467-025-12345-6)

Key Findings:

- Systematic review of gut microbiome alterations in ME/CFS
- Synthesises evidence across multiple cohort studies

Relevance to Part V: Consolidated microbiome evidence base for gut-immune axis modelling parameters.

H.47 Epigenetics and DNA Methylation

H.47.1 de Vega et al. 2021 — DNA Methylation Meta-Analysis

Full Citation: de Vega WC, Vernon SD, McGowan PO. DNA methylation meta-analysis reveals cellular alterations in myalgic encephalomyelitis/chronic fatigue syndrome. *Journal of Translational Medicine*. 2021;19(1):113. [76]

DOI: [10.1186/s12967-021-02780-9](https://doi.org/10.1186/s12967-021-02780-9)

Key Findings:

- Meta-analysis of DNA methylation changes in ME/CFS
- Identified epigenetic alterations affecting immune and metabolic gene networks

Relevance to Part V: Epigenetic data for modelling gene regulation layers in integrated systems models.

H.48 Viral Persistence and EBV Mechanisms

H.48.1 Pless et al. 2026 — EBV LMP1 and CNS Demyelination

Full Citation: Pless A, et al. Myelin antigen capture in the CNS by B cells expressing EBV latent membrane protein 1 leads to demyelinating lesion formation. *Cell*. 2026. [164]

DOI: [10.1016/j.cell.2025.12.031](https://doi.org/10.1016/j.cell.2025.12.031)

Key Findings:

- EBV LMP1 expression in B cells enables myelin antigen capture leading to CNS demyelination
- Demonstrates direct mechanism linking EBV to autoimmune neurological damage

Relevance to Part V: Mechanistic basis for viral-trigger models; EBV–autoimmunity pathway for immune–neuroendocrine coupling.

H.49 Cardiovascular and Autonomic Dysfunction

H.49.1 Raj et al. 2005 — Pyridostigmine RCT in CFS

Full Citation: Raj SR, Fennell E, Natelson BH, Frangos S, Shungu DC. Pyridostigmine in the treatment of chronic fatigue syndrome: A randomized, double-blind, placebo-controlled trial. *Journal of Clinical Psychiatry*. 2005;66(4):508–512. [558]

DOI: [10.4088/JCP.v66n0414](https://doi.org/10.4088/JCP.v66n0414)

Key Findings:

- RCT of pyridostigmine (acetylcholinesterase inhibitor) for CFS
- Tested autonomic modulation as therapeutic approach

Relevance to Part V: Autonomic intervention data for cardiovascular model calibration and cholinergic pathway modelling.

H.50 Treatment: Pharmacological Interventions

H.50.1 Polo et al. 2019 — Low Dose Naltrexone for Fibromyalgia

Full Citation: Polo O, Smith S, Jones DE, Newton JL. Low dose naltrexone for the treatment of fibromyalgia: Findings of a small, randomized, double-blind, placebo-controlled, counterbalanced, crossover trial assessing daily pain levels. *Arthritis & Rheumatology*. 2019;71(10):1691–1699. [469]

DOI: [10.1002/art.40900](https://doi.org/10.1002/art.40900)

Key Findings:

- Crossover RCT of low-dose naltrexone (LDN) in fibromyalgia
- Assessed daily pain levels as primary outcome

Relevance to Part V: LDN efficacy data for neuroinflammation treatment modelling; relevant to ME/CFS–fibromyalgia overlap models.

H.50.2 White et al. 2011 — PACE Trial

Full Citation: White PD, Goldsmith KA, Johnson AL, et al. Comparison of adaptive pacing therapy, cognitive behaviour therapy, graded exercise therapy, and specialist medical care for chronic fatigue syndrome (PACE): a randomised trial. *The Lancet*. 2011;377(9768):823–836. [115]

DOI: [10.1016/S0140-6736\(11\)60096-2](https://doi.org/10.1016/S0140-6736(11)60096-2)

Key Findings:

- Largest ME/CFS treatment trial ($n=641$) comparing CBT, GET, APT, and specialist medical care
- Originally reported benefit for CBT and GET; subsequently criticised for methodological flaws
- NICE 2021 reversed GET recommendation based on reanalysis

Relevance to Part V: Historical benchmark trial; methodological lessons for model validation and outcome measure selection.

I Ongoing and Planned ME/CFS Research Studies

This appendix catalogues major ongoing and planned research studies in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS). Each entry provides structured metadata—principal investigator, institution, registry identifiers, cohort size, and study focus—followed by a brief discussion of the study’s potential to advance ME/CFS understanding.

Data sources. Entries were compiled from ClinicalTrials.gov, the ISRCTN registry, institutional announcements, and the Solve ME/CFS Initiative trial tracker. Registry identifiers (NCT numbers, ISRCTN numbers) are included where available.

Currency. This registry was last updated on 2026-03-07. Study status information may have changed since that date; readers should verify current status via the URLs provided.

Scope. The registry includes studies that are actively recruiting, enrolled, or in analysis as of the currency date, plus selected planned studies with confirmed funding. Completed studies with published results are covered in the main text and Appendix H.

⚠ Limitation 1: Registry Completeness

This registry is necessarily incomplete. Smaller institutional studies, industry-sponsored trials in early phases, and studies registered outside ClinicalTrials.gov or ISRCTN may not be captured. Inclusion does not imply endorsement of study design or methodology; exclusion does not imply irrelevance.

I.1 Multi-Omics and Systems Biology

I.1.1 Rosetta Stone Study

Principal Investigator: Danny Altmann, Professor of Immunology

Institution: Imperial College London, United Kingdom

Contact/URL: <https://meassociation.org.uk/2025/12/driving-discovery-the-me-association-inv>

Funder: ME Association, Ramsay Research Fund

Status: Active (recruitment began late 2025)

Phase: Observational, multi-omics

Cohort: 250 ME/CFS + 250 Long Covid + matched healthy controls

Mechanism/Focus: Side-by-side cellular and molecular analyses of ME/CFS and Long Covid. Stool, blood, and saliva samples analysed across multiple omics platforms, with machine learning applied to discover shared molecular pathways between the two conditions. Partnership with ELAROS for digital phenotyping.

Primary Outcomes: Shared and distinguishing molecular signatures between ME/CFS and Long Covid; candidate biomarkers

Estimated Completion: 2028 (3-year study)

Timeline: Recruitment started Q4 2025; 3-month progress update published March 2026; interim analysis expected mid-study

Publication Medium: Not yet announced; likely high-impact immunology or multidisciplinary journal; open access expected given charity funding

Document Relevance: Ch. 7 (immune dysfunction), Ch. 11 (gut microbiome), Ch. 13 (integrative models), Ch. 14 §14.17 (cross-disease comparison)

The Rosetta Stone Study represents the largest single investment in ME/CFS biomedical research by a UK charity (£1.1M). Its defining strength is the parallel recruitment of ME/CFS and Long Covid cohorts with identical multi-omics profiling, enabling direct molecular comparison of two conditions widely hypothesised to share pathophysiological mechanisms. If the study identifies shared immune or metabolic signatures, it could accelerate biomarker development by leveraging the larger Long Covid research infrastructure. The inclusion of stool samples positions the study to contribute microbiome data that remains scarce in well-phenotyped ME/CFS cohorts.

I.1.2 BioQuest Study

Principal Investigator: Jonas Bergquist (proteomics lead); coordinated by Open Medicine Foundation (OMF) research team

Institution: Uppsala University (Sweden) and Open Medicine Foundation collaborative centres

Contact/URL: <https://ns1.omf.ngo/bioquest-update/>

Funder: Open Medicine Foundation

Status: Active (sample collection complete; analysis phase)

Phase: Observational, multi-omics biomarker discovery

Cohort: 400 ME/CFS + 400 healthy controls + 200 disease controls (50 each: multiple sclerosis, burnout syndrome, depression, Long Covid). Combines samples from the Uppsala Collaborative Centre and the NIH Chronic Fatigue Initiative biobank.

Mechanism/Focus: Large-scale metabolomics, lipidomics, proteomics, and cytokine profiling (>10,000 analytes) with AI-driven subgroup identification. Goal: identify a 5–20 protein/metabolite biomarker panel unique to ME/CFS.

Primary Outcomes: Validated ME/CFS biomarker panel; biologically defined patient subgroups

Estimated Completion: Results expected 2026–2027

Timeline: Sample integration approved 2025; testing began late 2025; bioinformatic analysis ongoing

Publication Medium: Expected in high-impact journal; OMF typically publishes open access

Document Relevance: Ch. 6 (energy metabolism), Ch. 7 (immune dysfunction), Ch. 13 (integrative models), Ch. 20 (biomarker research)

BioQuest is the largest ME/CFS biomarker study to date by sample size ($n = 1,000$). Its inclusion of disease controls (MS, depression, burnout, Long Covid) directly addresses a longstanding criticism that ME/CFS biomarker candidates lack disease specificity. The multi-omics approach measuring over 10,000 analytes, combined with AI-driven clustering, has realistic potential to identify biologically defined subgroups—a prerequisite for precision medicine approaches. If successful, BioQuest could provide the first blood-based diagnostic panel capable of distinguishing ME/CFS from conditions with overlapping symptom profiles.

I.2 Genomics and Epigenetics

I.2.1 DecodeME

Principal Investigator: Chris Ponting, Professor of Medical Bioinformatics

Institution: University of Edinburgh, United Kingdom

Contact/URL: <https://www.decode.me.org.uk/>

Registry ID: ISRCTN15960771

Funder: MRC/NIHR (UK Research and Innovation)

Status: Active (genotyping and analysis phase)

Phase: Observational, genome-wide association study (GWAS)

Cohort: >20,000 ME/CFS patients (DNA samples); UK-based, self-reported diagnosis with GP confirmation

Mechanism/Focus: First large-scale GWAS for ME/CFS. Aims to identify genetic variants associated with susceptibility, severity, and symptom subtypes. DNA collected via saliva kits mailed to participants.

Primary Outcomes: Genome-wide significant loci for ME/CFS; genetic correlation with related conditions; polygenic risk scores

Estimated Completion: First results expected 2026

Timeline: Recruitment launched 2022; sample collection completed 2024; genotyping in progress; first GWAS results anticipated 2026

Publication Medium: Expected in genetics/genomics journal (e.g., *Nature Genetics*); open access

Document Relevance: Ch. 12 (genetic and epigenetic factors), Ch. 20 (biomarker research), Ch. 23 (epidemiology)

DecodeME is unprecedented in ME/CFS genetics: with over 20,000 participants, it has the statistical power to detect common variants of modest effect size that previous underpowered GWAS could not identify. Genetic findings could reveal causal pathways (e.g., immune regulation, ion channel function, autonomic control) and enable Mendelian randomisation analyses to test whether associations observed in smaller studies reflect causal relationships. The study's scale also allows subgroup analyses by onset trigger, severity, and symptom profile.

I.2.2 EpiSwitch CFS Epigenetic Diagnostic

Principal Investigator: Ewan Hunter (lead author, validation study)

Institution: Oxford BioDynamics, United Kingdom

Contact/URL: <https://doi.org/10.1186/s12967-025-07203-w>

Registry ID: N/A (proof-of-concept; clinical validation pending)

Funder: Oxford BioDynamics (industry)

Status: Proof-of-concept published October 2025; clinical validation studies planned

Phase: Diagnostic development (pre-clinical validation)

Cohort: Initial study: 47 severe ME/CFS patients + 61 healthy controls. Larger validation cohorts needed.

Mechanism/Focus: EpiSwitch 3D genomic profiling of chromosome conformation captures (CCs) in blood. Identified a 200-marker epigenetic model. Pathways implicated: interleukins, TNF α , neuroinflammation, toll-like receptor signalling, JAK/STAT.

Primary Outcomes: Sensitivity 92%, specificity 98%, overall accuracy 96% in proof-of-concept validation cohort

Estimated Completion: Clinical validation timeline not yet announced

Timeline: Proof-of-concept published October 2025; independent replication and prospective validation required before clinical use

Publication Medium: *Journal of Translational Medicine* (published); future validation studies TBD

Document Relevance: Ch. 12 (genetic and epigenetic factors), Ch. 20 (biomarker research), Ch. 25 (translational findings)

The EpiSwitch proof-of-concept achieved striking diagnostic accuracy (96%), but the small sample size ($n = 108$) and retrospective design warrant caution. If validated in larger, prospective, multi-site cohorts with disease controls, this approach could provide the first epigenetic blood test for ME/CFS. The implicated pathways (JAK/STAT, TLR signalling, neuroinflammation) align with immune findings reported in Chapters 7 and 8, providing convergent evidence from an independent methodological angle. The critical next step is independent replication with blinded samples.

I.2.3 PrecisionLife Combinatorial Genomics

Principal Investigator: Steve Gardner (CEO, PrecisionLife)

Institution: PrecisionLife Ltd, Oxford, United Kingdom

Contact/URL: <https://precisionlife.com/news-and-events/me-genetics-study>

Registry ID: N/A (secondary analysis of DecodeME and UK Biobank data)

Funder: PrecisionLife (industry) with access to DecodeME consortium data

Status: Published December 2025

Phase: Computational genomics (AI-led combinatorial analytics)

Cohort: Two DecodeME cohorts + UK Biobank; results confirmed across three independent datasets

Mechanism/Focus: Applied combinatorial analytics platform to identify gene–gene interaction networks in ME/CFS. Unlike standard GWAS (which tests single variants), this approach identifies combinations of variants that jointly confer risk.

Primary Outcomes: Identified 250+ core genes associated with ME/CFS, of which 76 genes overlap with long COVID. Networks cluster around immune regulation, metabolic function, and neurological pathways.

Estimated Completion: Published

Publication Medium: PrecisionLife press release and preprint; peer-reviewed publication expected

Document Relevance: Ch. 12 (genetic factors), Ch. 13 (integrative models), Ch. 14 §14.17 (long COVID overlap)

The PrecisionLife analysis complements the DecodeME GWAS (Section 1.2.1) by detecting higher-order gene–gene interactions that conventional single-variant approaches miss. The 76-gene overlap with long COVID provides the strongest genetic evidence yet for shared biological mechanisms between the two conditions, supporting the hypothesis explored in Ch. 14 §14.17. However, combinatorial approaches can generate false positives when the number of tested combinations is large; independent replication (ideally in non-European ancestries) is essential before clinical or mechanistic conclusions are drawn. [682]

1.3 Biomarker Discovery and Diagnostics

1.3.1 BioMapAI (Gut Microbiome and Metabolomics)

Principal Investigator: Derya Unutmaz (Jackson Laboratory); AI platform developed with Duke University

Institution: Jackson Laboratory (JAX) and Duke University School of Medicine, United States

Contact/URL: <https://medschool.duke.edu/news/ai-thats-finally-making-sense-chronic-fatigue>

Funder: NIH and institutional funding

Status: Active (longitudinal follow-up ongoing)

Phase: Observational, AI-integrated multi-omics

Cohort: 153 ME/CFS patients + 96 healthy controls, followed longitudinally over 4 years

Mechanism/Focus: AI platform integrating gut metagenomics, plasma metabolomics, immune cell profiles, standard blood tests, and clinical symptoms. Achieved 90% accuracy in distinguishing ME/CFS from controls. Identified gut microbiome signatures as strong predictors.

Primary Outcomes: Multi-modal diagnostic classifier; longitudinal trajectory modelling; microbiome-metabolome interaction networks

Estimated Completion: Ongoing longitudinal follow-up; initial results published 2025

Timeline: Recruitment 2020–2023; initial AI classifier published July 2025; longitudinal analysis ongoing

Publication Medium: Initial results in high-impact journal (2025); longitudinal follow-up expected 2026–2027

Document Relevance: Ch. 11 (gut microbiome), Ch. 6 (energy metabolism), Ch. 13 (integrative models), Ch. 20 (biomarker research)

BioMapAI is notable for its longitudinal design and AI-first analytical approach. Unlike cross-sectional biomarker studies, the 4-year follow-up can capture disease trajectory and distinguish state markers (fluctuating with symptoms) from trait markers (stable features of the disease). The gut microbiome component strengthens the microbiome–immune–metabolism axis described in Chapter 11. The 90% classification accuracy, while promising, requires external validation in independent cohorts.

I.3.2 ME Association PhD Project (Metabolite and Infection Markers)

Principal Investigator: Aleya Lumsden (PhD candidate); supervised across multiple institutions

Institution: Rosalind Franklin Institute, Imperial College London, and National Phenome Centre, United Kingdom

Contact/URL: <https://meassociation.org.uk/2025/11/research-new-funding-awarded-to-phd-project>

Funder: ME Association Ramsay Research Fund and UKRI (joint)

Status: Active (PhD project started 2025)

Phase: Observational, metabolomics and infection biomarkers

Cohort: To be determined (likely leveraging existing biobank samples)

Mechanism/Focus: Advanced mass spectrometry and informatics to identify unknown metabolites and uncover evidence of infection in ME/CFS. Uses facilities across three major UK research centres.

Primary Outcomes: Novel metabolite identification; infection marker discovery; candidate biomarkers for further validation

Estimated Completion: 2028–2029 (typical PhD duration)

Timeline: Funding awarded November 2025; project started 2025–2026

Publication Medium: PhD thesis and associated journal publications

Document Relevance: Ch. 6 (energy metabolism), Ch. 7 (immune dysfunction), Ch. 20 (biomarker research)

This PhD project complements the larger Rosetta Stone Study by focusing specifically on unknown metabolites—compounds not yet characterised in standard metabolomics panels. The infection marker component addresses the long-standing hypothesis that persistent or reactivated infections drive ME/CFS pathology (Chapter 7). Access to the Rosalind Franklin Institute’s advanced mass spectrometry facilities gives this project analytical capabilities beyond typical academic metabolomics studies.

I.3.3 MEA–UCL Blood Biomarker Pilot

Principal Investigator: Sophie Hicks (PhD student); supervisors Dr. Amanda Heslegrave, Dr. Michael Zandi, Prof. Henrik Zetterberg

Institution: UK DRI Fluid Biomarker Laboratory, UCL Institute of Neurology, London

Contact/URL: ME Association announcement, February 2026

Registry ID: Not yet registered (pilot/observational)

Funder: ME Association (UK)

Status: Active (12-month project, started February 2026)

Phase: Observational pilot

Cohort: Blood samples from ME/CFS patients (UK ME/CFS Biobank), Long Covid patients (UCLH STIMULATE-ICP study), and healthy volunteers

Mechanism/Focus: Searches for blood-based biomarkers linking specific symptom clusters (brain fog, headaches, muscle pain) to measurable biological changes. Applies neuroimmunology expertise from the Zetterberg group (world-leading in fluid biomarkers for neurodegeneration, e.g., neurofilament light chain).

Primary Outcomes: Candidate blood biomarkers correlated with symptom clusters

Estimated Completion: ~February 2027

Timeline: 12-month pilot; may lead to larger validation study

Publication Medium: Not yet announced

Document Relevance: Ch. 7 (immune dysfunction), Ch. 22 (mechanistic studies)

The Zetterberg laboratory's involvement is notable: their deep expertise in neurofilament light chain and related neurodegeneration biomarkers has not previously been applied to ME/CFS. Cross-disease biomarker candidates (shared between ME/CFS and neurodegenerative conditions) could emerge from this work, informing the diagnostic overlap discussion in Chapter 14.17.

I.3.4 Cornell Cell-Free RNA Liquid Biopsy

Principal Investigator: Iwijn De Vlaminck (Biomedical Engineering) and Maureen Hanson (Molecular Biology and Genetics)

Institution: Cornell University, Ithaca, NY

Contact/URL: <https://doi.org/10.1073/pnas.2507345122>

Funder: NIH / institutional

Status: Published August 2025

Phase: Biomarker discovery (cross-sectional)

Cohort: 93 ME/CFS cases + 75 healthy sedentary controls

Mechanism/Focus: Profiled circulating cell-free RNA (cfRNA) in plasma by RNA sequencing. Identified over 700 differentially expressed transcripts. Immune cfRNA deconvolution revealed elevated plasmacytoid dendritic cell, monocyte, and T cell-derived cfRNA in ME/CFS, with signatures of cytokine signalling and T cell exhaustion.

Primary Outcomes: GLM-LASSO classifier AUC 0.81, accuracy 77%. Not yet diagnostic-grade but substantial advance over prior attempts

Estimated Completion: Published

Publication Medium: *Proceedings of the National Academy of Sciences*

Document Relevance: Ch. 7 (immune dysfunction), Ch. 20 (biomarker research), Ch. 25 (translational findings)

Cell-free RNA offers a minimally invasive “liquid biopsy” approach that captures tissue-level biology without biopsies. The 77% classification accuracy is insufficient for a standalone diagnostic but demonstrates that cfRNA carries ME/CFS-relevant biological signal [205]. The T cell exhaustion and platelet activation signatures align with independent findings from single-cell transcriptomics (Section I.9.1) and proteomics (Section I.3.5). Prospective validation in a larger, multi-site cohort with disease controls (fibromyalgia, depression, long COVID) is needed to assess clinical utility.

I.3.5 Cell Reports Medicine Multi-Modal Study

Principal Investigator: Multiple collaborating groups

Institution: Multi-centre (USA)

Contact/URL: <https://doi.org/10.1016/j.xcrm.2025.102012>

Funder: NIH / institutional

Status: Published 2025

Phase: Observational, multi-modal

Cohort: ME/CFS patients + healthy controls

Mechanism/Focus: Integrated energy metabolism assessment, immune profiling, and plasma proteomics. Immune cells from ME/CFS patients showed elevated AMP and ADP with reduced ATP/ADP ratio. NK cell and dendritic cell subset abnormalities. Vascular dysfunction markers including soluble THBS1, VWF, and VASN.

Primary Outcomes: Converging evidence across energy, immune, and vascular axes; reduced CD56^{low}CD16⁺ NK cells (antiviral), increased plasmacytoid DCs

Estimated Completion: Published

Publication Medium: *Cell Reports Medicine*

Document Relevance: Ch. 6 (energy metabolism), Ch. 7 (immune dysfunction), Ch. 10 (cardiovascular), Ch. 13 (integrative models)

This study is noteworthy for its simultaneous assessment of three pathophysiological axes—energy, immune, and vascular—in the same cohort, enabling correlation analyses that single-axis studies cannot perform [169]. The reduced ATP/ADP ratio in immune cells provides a mechanistic link between energy metabolism (Ch. 6) and immune dysfunction (Ch. 7). The vascular findings (elevated thrombospondin-1, von Willebrand factor) converge with coagulation abnormalities reported in the microclot literature [166].

I.4 Autoimmunity and Immunomodulation

I.4.1 Charité Immunoadsorption Trial (IA-PACS-CFS)

Principal Investigator: Carmen Scheibenbogen, MD, Professor of Immunology

Institution: Charité – Universitätsmedizin Berlin, Charité Fatigue Centrum, Germany

Contact/URL: https://cfc.charite.de/en/clinical_research/nksg/trial_ia_pacs_cfs

Registry ID: NCT05710770

Funder: Institutional and ME/CFS Research Foundation

Status: Active (double-blind, sham-controlled)

Phase: Exploratory interventional trial

Cohort: ME/CFS patients (post-COVID and other post-infectious) with elevated β_2 -adrenergic receptor autoantibodies

Mechanism/Focus: Five immunoadsorption sessions over 10 days to remove pathogenic autoantibodies targeting G-protein coupled receptors (GPCRs), compared with sham treatment. Prior open-label cohort ($n = 20$) showed clinical improvement.

Primary Outcomes: Clinical improvement at 3 months (Chalder Fatigue Scale); autoantibody levels; 6-month follow-up

Estimated Completion: Not yet announced

Timeline: Open-label cohort 2022–2023 (published January 2025); double-blind RCT ongoing

Publication Medium: Open-label results in peer-reviewed journal (2025); RCT results expected in high-impact immunology journal

Document Relevance: Ch. 7 (immune dysfunction—autoantibodies), Ch. 10 (autonomic dysfunction), Ch. 19 (integrative approaches)

The Charité immunoadsorption programme is the most advanced clinical investigation of the autoantibody hypothesis in ME/CFS. Scheibenbogen's group has systematically built from case reports through an open-label cohort to a sham-controlled RCT—the methodological progression needed to establish causality. The requirement for elevated β_2 -adrenergic receptor autoantibodies as an inclusion criterion represents a biomarker-stratified trial design, testing both the treatment and the underlying autoimmune mechanism simultaneously. A new 2026-funded neuroimaging substudy will assess whether thalamocortical hyperconnectivity normalises after immunoadsorption, linking peripheral autoimmunity to central nervous system changes.

I.4.2 BC007 (Rovunaptabin) Phase II Trials

Principal Investigator: Multiple (industry-sponsored)

Institution: Berlin Cures GmbH, Germany (multi-site)

Contact/URL: <https://clinicaltrials.gov/study/NCT05911009>

Registry ID: NCT05911009 (Phase II, $n = 114$); earlier Phase IIa ($n = 29$)

Funder: Berlin Cures GmbH (industry)

Status: Phase II completed/in analysis

Phase: Phase IIa (crossover, $n = 29$) and Phase II ($n = 114$)

Cohort: Post-COVID syndrome patients with functional GPCR autoantibodies and predominant fatigue

Mechanism/Focus: Rovunaptabin (BC007) is an aptamer that neutralises functional autoantibodies against GPCRs. Unlike immunoadsorption (which removes all immunoglobulins non-selectively), BC007 targets pathogenic autoantibodies specifically.

Primary Outcomes: Fatigue severity; autoantibody neutralisation; safety

Estimated Completion: Results expected 2026

Timeline: Phase IIa enrolled 29 patients; Phase II enrolled 114 patients; analysis ongoing

Publication Medium: Not yet announced; industry-sponsored trials typically target high-impact journals

Document Relevance: Ch. 7 (immune dysfunction—autoantibodies), Ch. 25 (translational findings)

BC007 and immunoadsorption represent complementary approaches to the same autoantibody hypothesis: immunoadsorption removes antibodies broadly, while BC007 neutralises specific pathogenic autoantibodies in situ. The Phase II trial ($n = 114$) is the largest RCT testing autoantibody-targeted therapy in post-infectious fatigue conditions. Positive results would validate the functional GPCR autoantibody hypothesis and open a pharmacological treatment pathway. However, the trials enrolled post-COVID patients specifically; extension to non-COVID ME/CFS cohorts will be needed.

I.4.3 ResetME Trial (Daratumumab)

Principal Investigator: Øystein Flüge and Olav Mella

Institution: Haukeland University Hospital, Bergen, Norway (with Oslo University Hospital)

Contact/URL: EudraCT / EU Clinical Trials Register

Registry ID: European registry (no NCT number)

Funder: ME Fund Norway; institutional

Status: Recruiting (opened June 2025)

Phase: Phase II, randomised, double-blind, placebo-controlled (2:1 ratio)

Cohort: 66 participants; moderate-to-severe ME/CFS (CCC criteria), minimum 2-year disease duration

Mechanism/Focus: Daratumumab is an anti-CD38 monoclonal antibody that depletes plasma cells, causing transient reduction of pathogenic autoantibodies (including anti-adrenergic and anti-cholinergic receptor antibodies). Follow-on from the open-label KTS9 pilot ($n = 10$, EudraCT 2022-000281-18), in which 6/10 patients showed marked improvement and mean SF-36 Physical Function rose from 25.9 to 55.0.

Primary Outcomes: Symptom reduction (SF-36 Physical Function); safety

Estimated Completion: ~2027 (72-week total participation)

Timeline: 26-week treatment period, 72-week follow-up from June 2025 start

Publication Medium: Not yet announced; pilot published in *Frontiers in Medicine*, June 2025

Document Relevance: Ch. 7 (immune dysfunction—autoantibodies), Ch. 10 (adrenergic receptor autoantibodies), Ch. 19 (integrative approaches)

ResetME is among the highest-priority ongoing ME/CFS trials. Flüge and Mella are the same

group that conducted the rituximab trials; daratumumab targets plasma cells more selectively via CD38. Unlike rituximab (which depletes CD20+ B cells), daratumumab directly reduces the antibody-secreting plasma cells hypothesised to produce pathogenic autoantibodies. The pilot response rate (60%, with near-doubling of physical function scores) is among the strongest signals in ME/CFS treatment research. If the RCT replicates these findings, it would establish the first disease-modifying treatment for an autoimmune subset of ME/CFS. Combined with BC007 (Section I.4.2) and immunoadsorption (Section I.4.1), this trial represents the third independent approach to the autoantibody hypothesis.

I.5 Neuroimaging and Ion Channel Research

I.5.1 NCNED TRPM3 Ion Channel Programme

Principal Investigator: Sonya Marshall-Gradisnik, Professor of Immunology

Institution: National Centre for Neuroimmunology and Emerging Diseases (NCNED), Griffith University, Australia

Contact/URL: <https://www.griffith.edu.au/research/health/national-centre-neuroimmunology-em>

Funder: Institutional and NHMRC (Australia)

Status: Active (multiple ongoing studies)

Phase: Observational and interventional (LDN trials planned)

Cohort: ME/CFS and Long Covid patients (multiple cohorts)

Mechanism/Focus: Systematic investigation of TRPM3 ion channel dysfunction in ME/CFS. January 2026 publication confirmed consistent TRPM3 malfunction using gold-standard patch-clamp techniques. Also conducting 7T MRI studies revealing hippocampal volume alterations. Preparing LDN clinical trials for both ME/CFS and Long Covid.

Primary Outcomes: TRPM3 channel function; hippocampal volume changes; LDN trial endpoints TBD

Estimated Completion: Ongoing programme (multiple studies)

Timeline: TRPM3 confirmation published January 2026; 7T MRI study published February 2025; LDN trials in preparation; \$438,000 grant for longitudinal brain imaging (progression study)

Publication Medium: Published in *PLOS ONE* and specialty journals; future publications expected similarly

Document Relevance: Ch. 8 (neurological dysfunction), Ch. 7 (immune dysfunction—ion channels), Ch. 20 (biomarker research)

The NCNED programme is distinctive for its sustained, systematic focus on a single molecular target (TRPM3) across multiple methodologies. The January 2026 patch-clamp confirmation elevates TRPM3 dysfunction from an association to a consistently reproducible cellular phenotype—a rare achievement in ME/CFS research. The planned LDN trials will test whether modulating neuroinflammation (a proposed upstream driver of ion channel dysfunction) improves TRPM3 function and clinical outcomes, creating a mechanistic link between

treatment and pathophysiology. The 7T MRI hippocampal findings add a neuroanatomical dimension that connects to cognitive complaints reported by patients.

I.5.2 Charité Neuroimaging Biomarker Study

Principal Investigator: Carmen Scheibenbogen, MD (PI); neuroimaging team at Charité

Institution: Charité – Universitätsmedizin Berlin, Germany

Contact/URL: <https://mecfs-research.org/en/researchfunding-charite-biomarker2/>

Funder: ME/CFS Research Foundation (Germany)

Status: Funded; starting January 2026

Phase: Observational (neuroimaging biomarker development)

Cohort: ME/CFS patients undergoing immunoadsorption treatment

Mechanism/Focus: Investigates whether functional brain changes— particularly thalamo-cortical hyperconnectivity—can be modulated by immunoadsorption. Tests whether CNS abnormalities normalise after peripheral autoantibody removal, linking the autoimmune and neurological hypotheses.

Primary Outcomes: Pre/post-treatment fMRI connectivity changes; correlation with clinical improvement and autoantibody levels

Estimated Completion: Not yet announced (recently funded)

Timeline: Funding confirmed late 2025; project starting January 2026

Publication Medium: Not yet announced

Document Relevance: Ch. 8 (neurological dysfunction), Ch. 7 (immune dysfunction—autoantibodies), Ch. 13 (integrative models)

This study bridges two major ME/CFS hypotheses: autoimmunity and central nervous system dysfunction. By imaging the brain before and after immunoadsorption, it can test whether peripheral autoantibody removal reverses central connectivity abnormalities—a question that goes beyond symptom improvement to mechanistic understanding. If thalamocortical hyperconnectivity normalises with autoantibody reduction, it would provide the strongest evidence yet that peripheral immune dysregulation drives CNS pathology in ME/CFS rather than the reverse.

I.5.3 Hyperbaric Oxygen Therapy for ME/CFS

Principal Investigator: Multiple PIs

Institution: Multi-site (USA / Germany)

Contact/URL: <https://doi.org/10.1101/2025.10.29.25339096>

Registry ID: NCT06118138 (post-COVID/ME/CFS HBOT study)

Funder: ME/CFS Research Foundation / institutional

Status: Pilot results published (preprint October 2025); larger RCT in planning

Phase: Phase 1–2 (open-label pilot)

Cohort: $n = 30$ ME/CFS patients, 40 HBOT sessions each

Mechanism/Focus: Hyperbaric oxygen therapy (2.4 ATA, 100% O₂) to address tissue hypoxia, neuroinflammation, and endothelial dysfunction. Pre- and post-treatment fMRI assessed thalamic connectivity.

Primary Outcomes: SF-36 physical functioning significantly improved. Exercise capacity, muscle strength, and information processing speed all improved. Thalamic hyperconnectivity normalised post-HBOT

Estimated Completion: Pilot published; RCT timeline TBD

Publication Medium: medRxiv preprint; peer review pending

Document Relevance: Ch. 8 (neurological dysfunction), Ch. 10 (cardiovascular), Ch. 19 (integrative approaches)

The normalisation of thalamic hyperconnectivity after HBOT provides the most direct interventional evidence to date linking brain connectivity abnormalities to a treatable target in ME/CFS [683]. However, the open-label design and small sample size preclude definitive conclusions; placebo effects are substantial for fatigue interventions, and a sham-controlled design (e.g., 1.3 ATA room air) is essential. The overlap with brainstem hypoperfusion research strengthens the rationale for oxygen-targeting approaches.

I.6 Large-Scale Federally Funded Programmes

I.6.1 RECOVER-TLC Clinical Trials Programme

Principal Investigator: Multiple PIs across sites

Institution: NIH-funded, multi-site across the United States

Contact/URL: <https://recovercovid.org/>

Registry ID: Multiple NCT numbers (programme-level)

Funder: NIH (National Institutes of Health)

Status: Active; multiple trials at different stages

Phase: Multiple interventional trials (Phase II–III)

Cohort: Long Covid patients (with substantial ME/CFS overlap)

Mechanism/Focus: The RECOVER-Treating Long COVID (TLC) programme includes multiple treatment trials:

- **REVERSE-LC:** Baricitinib (JAK inhibitor) for neurocognitive and cardiopulmonary symptoms; recruiting
- **RECOVER Energize:** Structured pacing for PEM ($n = 600$); completing December 2025
- **LDN trial:** Low-dose naltrexone for fatigue in ages 6–25; enrolment estimated summer 2026
- **GLP-1 agonist trial:** Targeting immune dysregulation; enrolment estimated late summer 2026

Primary Outcomes: Vary by trial; include symptom severity, exercise tolerance, immune markers, quality of life

Estimated Completion: Results from first-round trials expected by end of 2026

Timeline: RECOVER programme initiated 2021; TLC trials began 2024–2025; second-round trials planning 2026

Publication Medium: High-impact journals; all results expected to be open access (NIH-funded)

Document Relevance: Ch. 7 (immune dysfunction), Ch. 10 (cardiovascular), Ch. 17 (pacing), Ch. 14 §14.17 (cross-disease)

RECOVER-TLC is the largest federally funded treatment trial programme relevant to ME/CFS. While focused on Long Covid, the substantial clinical overlap means positive results are directly translatable. The baricitinib (JAK inhibitor) trial is particularly noteworthy given the JAK/STAT pathway abnormalities identified in ME/CFS (Chapter 7) and the EpiSwitch epigenetic findings. The structured pacing trial (RECOVER Energize, $n = 600$) will provide the first large-scale evidence for pacing—the most widely recommended non-pharmacological intervention in ME/CFS—in a rigorous clinical trial framework. The paediatric LDN trial fills a critical gap, as most ME/CFS treatment evidence excludes children and adolescents.

I.7 Clinical Trials (Interventional)

I.7.1 LIFT Trial (Low-Dose Naltrexone and Pyridostigmine)

Principal Investigator: David Systrom, MD

Institution: Brigham and Women's Hospital, Harvard Medical School, United States

Contact/URL: <https://solvecfs.org/exploring-current-me-cfs-and-long-covid-clinical-trials->

Funder: Solve ME/CFS Initiative (Ramsay Research Grant)

Status: Recruiting

Phase: Clinical trial (interventional)

Cohort: ME/CFS patients (specific n to be confirmed)

Mechanism/Focus: Tests whether low-dose naltrexone (LDN) and pyridostigmine reduce symptoms and improve quality of life. LDN targets neuroinflammation via glial cell modulation; pyridostigmine is an acetylcholinesterase inhibitor used for autonomic dysfunction.

Primary Outcomes: Symptom severity; quality of life measures

Estimated Completion: Not yet announced

Timeline: Recruitment ongoing as of 2026

Publication Medium: Not yet announced

Document Relevance: Ch. 16 (supplements and nutraceuticals), Ch. 10 (cardiovascular/autonomic), Ch. 17 (lifestyle interventions)

The LIFT trial addresses two of the most widely used off-label treatments in ME/CFS clinical practice. LDN has accumulated substantial anecdotal and preliminary evidence (Chapter 16) but lacks rigorous RCT data in ME/CFS specifically. Pyridostigmine targets the preload failure mechanism identified in invasive cardiopulmonary exercise testing by Systrom's own

group (Chapter 10). A positive result could provide the first high-quality evidence base for two treatments already in widespread empirical use.

I.7.2 UAB LDN Dose-Finding Study

Principal Investigator: Jarred Younger, PhD

Institution: University of Alabama at Birmingham, United States

Contact/URL: <https://clinicaltrials.gov/study/NCT07285473>

Registry ID: NCT07285473

Funder: Institutional

Status: Recruiting (remote/decentralised, US-wide)

Phase: Dose-finding study

Cohort: Adults with ME/CFS (specific *n* TBD)

Mechanism/Focus: Dose-finding study for low-dose naltrexone in ME/CFS. Distinct from the LIFT trial (which combines LDN with pyridostigmine, Section I.7.1) and the RECOVER paediatric LDN trial (Section I.6.1): this study focuses on determining optimal adult LDN dosing specifically for ME/CFS. Remote design enables broad US enrolment.

Primary Outcomes: Dose-response for symptom reduction; safety and tolerability

Estimated Completion: Not yet announced

Timeline: Recruitment ongoing as of 2026

Publication Medium: Not yet announced

Document Relevance: Ch. 16 (supplements—LDN), Ch. 7 (immune dysfunction—microglial hypothesis)

Younger's laboratory has prior immunology work on LDN in ME/CFS (NCT02965768). While the LIFT trial tests LDN in combination, this study isolates LDN's independent effect and seeks to establish the optimal dose—a fundamental gap in the current evidence base. Most clinicians prescribe 1.5–4.5 mg empirically; systematic dose-finding data could substantially improve clinical practice.

I.7.3 Low-Dose Rapamycin in ME/CFS and Long Covid

Principal Investigator: Multi-site collaboration

Institution: Simmaron Research Institute, Bateman Horne Center, CDC, and Mayo Clinic, United States

Contact/URL: <https://batemanhornecenter.org/promising-clinical-trials/>

Funder: Multi-institutional

Status: Active

Phase: Clinical trial (interventional)

Cohort: ME/CFS and Long Covid patients (specific *n* TBD)

Mechanism/Focus: Rapamycin (sirolimus) is an mTOR inhibitor with immunomodulatory and anti-inflammatory properties. Targets mTOR pathway dysregulation implicated in ME/CFS immune and metabolic abnormalities.

Primary Outcomes: Symptom improvement; immune marker changes; safety and tolerability

Estimated Completion: Not yet announced

Timeline: Study active as of 2025–2026

Publication Medium: Not yet announced

Document Relevance: Ch. 7 (immune dysfunction), Ch. 6 (energy metabolism), Ch. 19 (integrative approaches)

The rapamycin trial is notable for its multi-institutional design (Simmaron, Bateman Horne, CDC, Mayo Clinic), which lends credibility and ensures phenotyping rigour. mTOR pathway dysregulation has been reported in ME/CFS (Chapter 7) and connects immune activation to metabolic dysfunction (Chapter 6). This trial directly tests a mechanistic hypothesis rather than treating symptoms empirically, making it particularly informative regardless of outcome.

I.7.4 Lumbrokinase for ME/CFS and Long Covid

Principal Investigator: David Putrino, PhD

Institution: Abilities Research Center, Mount Sinai Icahn School of Medicine, United States

Contact/URL: <https://solvecfs.org/exploring-current-me-cfs-and-long-covid-clinical-trials->

Funder: Institutional

Status: Recruiting

Phase: Clinical trial (interventional)

Cohort: Adults with Long Covid, post-treatment Lyme disease syndrome, and ME/CFS

Mechanism/Focus: Lumbrokinase is a fibrinolytic enzyme targeting microclot pathology. The microclot hypothesis proposes that persistent fibrin amyloid microclots impair microcirculation, contributing to tissue hypoxia and fatigue.

Primary Outcomes: Fibrinolytic markers; symptom scores; exercise tolerance

Estimated Completion: Not yet announced

Timeline: Recruitment ongoing as of 2026

Publication Medium: Not yet announced

Document Relevance: Ch. 10 (cardiovascular dysfunction), Ch. 19 (integrative approaches)

The microclot hypothesis has generated significant interest since 2021 but lacks interventional trial data. This study directly tests whether fibrinolytic treatment improves outcomes, providing a causal test of the hypothesis. The inclusion of three post-infectious conditions (Long Covid, post-Lyme, ME/CFS) mirrors the Rosetta Stone Study's cross-disease approach, potentially revealing whether microclot pathology is a shared mechanism.

I.7.5 Hydrogen Water for ME/CFS

Principal Investigator: Fred Friedberg, PhD

Institution: Stony Brook University School of Medicine, United States

Contact/URL: <https://solvecfs.org/exploring-current-me-cfs-and-long-covid-clinical-trials->

Funder: Solve ME/CFS Initiative

Status: Recruiting

Phase: Clinical trial, dosing study

Cohort: ME/CFS patients (specific *n* TBD)

Mechanism/Focus: Hydrogen-rich water as an antioxidant intervention. Molecular hydrogen (H₂) selectively scavenges hydroxyl radicals and may reduce oxidative stress and neuroinflammation.

Primary Outcomes: Fatigue severity; stress markers; physical function

Estimated Completion: Not yet announced

Timeline: Recruitment ongoing as of 2026

Publication Medium: Not yet announced

Document Relevance: Ch. 6 (energy metabolism—oxidative stress), Ch. 16 (supplements and nutraceuticals)

This dosing study addresses a practical gap: even if hydrogen water has antioxidant benefits, optimal dosing for ME/CFS has not been established. The oxidative stress hypothesis in ME/CFS (Chapter 6) provides a mechanistic rationale, though the evidence base for molecular hydrogen in any chronic condition remains preliminary. Results will be most informative if they include objective oxidative stress biomarkers alongside subjective symptom measures.

I.7.6 RESTORE-ME: Oxaloacetate for ME/CFS Fatigue

Principal Investigator: Alan Cash, PhD

Institution: Terra Biological LLC, United States (multi-site RCT)

Contact/URL: <https://clinicaltrials.gov/study/NCT05273372>

Registry ID: NCT05273372

Funder: Terra Biological LLC (industry)

Status: Results published (November 2024)

Phase: Phase II, randomised, double-blind, placebo-controlled

Cohort: 82 ME/CFS patients (2,000 mg oxaloacetate vs control daily for 3 months)

Mechanism/Focus: Oxaloacetate (OAA) is a mitochondrial metabolite that supports the citric acid cycle and NAD⁺/NADH ratio. Targets metabolic dysfunction by supplementing a key TCA cycle intermediate.

Primary Outcomes: Fatigue reduced >25% from baseline in OAA group (*p* = 0.0039 vs control); cognitive improvement at day 60 (*p* = 0.034); modest increase in uptime at day 30

Estimated Completion: Completed

Timeline: RCT completed 2024; results published November 2024 in *Frontiers in Neurology*; cognitive follow-up published 2025

Publication Medium: *Frontiers in Neurology* (open access)

Document Relevance: Ch. 6 (energy metabolism—TCA cycle), Ch. 16 (supplements and nutraceuticals)

RESTORE-ME is one of very few completed RCTs showing statistically significant fatigue reduction in ME/CFS. The mechanistic rationale is direct: oxaloacetate is an obligate TCA cycle intermediate, and ME/CFS metabolomic studies have consistently identified TCA cycle disruption (Chapter 6). The 25% fatigue reduction is clinically meaningful, though the modest sample size ($n = 82$) and industry sponsorship warrant independent replication. The cognitive improvement findings strengthen the case that metabolic support can address both physical and cognitive symptoms.

I.7.7 Stellate Ganglion Block for ME/CFS

Principal Investigator: Deborah Duricka, PhD; Lipeng Liu, MD

Institution: Multi-site, United States

Contact/URL: <https://doi.org/10.1080/21641846.2025.2455876>

Funder: Solve ME/CFS Initiative (Ramsay Research Grant)

Status: Pilot study published (February 2025); larger trials needed

Phase: Prospective cohort pilot ($n = 10$)

Cohort: 10 patients meeting both WHO Long Covid and IOM ME/CFS criteria

Mechanism/Focus: Sequential bilateral stellate ganglion blocks (local anaesthetic injection near the cervical sympathetic chain) administered over 3 consecutive weeks. Hypothesised to “reset” sympathetic/parasympathetic balance of the autonomic nervous system, increasing regional blood flow and reducing sympathetic hyperactivation.

Primary Outcomes: Significant improvements in cognition, physical and social functioning, vitality; reduced orthostatic intolerance and POTS symptoms. Cortisol levels did not change.

Estimated Completion: Pilot completed; larger RCT recommended

Timeline: Pilot published February 2025; larger trial planning stage

Publication Medium: *Fatigue: Biomedicine, Health & Behavior* (2025)

Document Relevance: Ch. 10 (cardiovascular/autonomic dysfunction), Ch. 8 (neurological dysfunction), Ch. 19 (integrative approaches)

The stellate ganglion block is a well-established procedure in pain medicine repurposed for autonomic dysfunction. The pilot results are notable for improvement across both autonomic (OI, POTS) and cognitive domains, suggesting a shared upstream mechanism. The “autonomic reset” hypothesis aligns with evidence of sympathetic hyperactivation in ME/CFS (Chapter 10). The very small sample ($n = 10$) limits conclusions, but the effect sizes and the procedure’s established safety profile justify a larger sham-controlled RCT.

I.7.8 Transcutaneous Vagus Nerve Stimulation (tVNS) for Long Covid/ME/CFS

Principal Investigator: Multiple investigators (several parallel studies)

Institution: Multi-site (US and European studies)

Contact/URL: <https://clinicaltrials.gov/study/NCT06585254>

Registry ID: NCT06585254 (US dosing study); additional European trials ongoing

Funder: Multiple (Solve ME/CFS Initiative, institutional)

Status: Recruiting (multiple trials)

Phase: Interventional (parameter optimisation and efficacy)

Cohort: Long Covid and ME/CFS patients fulfilling CFS criteria; prior studies $n = 10\text{--}25$ per trial

Mechanism/Focus: Non-invasive auricular stimulation of the vagus nerve to activate the cholinergic anti-inflammatory pathway (CAP). Targets neuroinflammation, autonomic imbalance, and systemic inflammation. Prior pilot studies showed improvements in cognition, anxiety, depression, sleep, and autonomic scores (COMPASS-31).

Primary Outcomes: Optimal stimulation parameters; health-related quality of life; fatigue severity; autonomic function

Estimated Completion: Varies by trial; results expected 2026–2027

Timeline: Multiple pilot studies published 2024–2025; current trials optimising parameters and assessing efficacy in larger cohorts

Publication Medium: *Frontiers in Neurology*, *PLOS ONE*, and others (prior studies published open access)

Document Relevance: Ch. 8 (neurological dysfunction), Ch. 10 (autonomic dysfunction), Ch. 7 (immune dysfunction—inflammation)

Vagus nerve stimulation represents a compelling intersection of neurology and immunology: the cholinergic anti-inflammatory pathway provides a mechanism through which neural stimulation can modulate systemic inflammation. The consistency of improvements across multiple small pilot studies (cognition, autonomic function, sleep) is encouraging, though no large RCT has yet been completed. The non-invasive nature, low side-effect profile, and consumer device availability (e.g., Nurosym) make tVNS a practical intervention if efficacy is confirmed. The current parameter-optimisation trials are a necessary step before definitive efficacy studies.

I.7.9 Masitinib for Severe Mast Cell Activation Syndrome

Principal Investigator: Multiple investigators

Institution: AB Science (sponsor), multi-site

Contact/URL: <https://clinicaltrials.gov/study/NCT05449444>

Registry ID: NCT05449444

Funder: AB Science (industry)

Status: Phase 2 (active)

Phase: Phase 2 clinical trial

Cohort: Patients with severe mast cell activation syndrome (MCAS)

Mechanism/Focus: Masitinib is a tyrosine kinase inhibitor that targets mast cell activity directly, unlike antihistamines that only address downstream mediator release. MCAS is identified in an estimated 50% of ME/CFS patients by some specialists.

Primary Outcomes: MCAS symptom severity; safety and tolerability

Estimated Completion: Not yet announced

Timeline: Trial active as of 2025–2026

Publication Medium: Not yet announced

Document Relevance: Ch. 7 (immune dysfunction—mast cells), Ch. 19 (integrative approaches)

Mast cell activation has emerged as a significant comorbidity in ME/CFS subsets, with some clinicians reporting MCAS in up to half of their ME/CFS patients. A 2026 cromolyn sodium case series from Dutch and Johns Hopkins investigators showed improvement in ME/CFS-related symptoms alongside MCAS management. Masitinib's upstream mechanism (direct tyrosine kinase inhibition rather than antihistamine blockade) could provide more complete mast cell stabilisation. If effective, this trial could open a treatment pathway for the MCAS subset of ME/CFS patients who respond poorly to conventional antihistamine regimens.

I.7.10 Cornell NAC–Glutathione Brain Study

Principal Investigator: Dikoma Shungu, PhD

Institution: Weill Cornell Medical College, United States

Contact/URL: <https://clinicaltrials.gov/study/NCT04542161>

Registry ID: NCT04542161

Funder: NINDS (National Institute of Neurological Disorders and Stroke)

Status: Recruiting (Phase 2)

Phase: Phase 2, double-blind, placebo-controlled, randomised

Cohort: 60 participants (20 per arm: 0 mg / 900 mg / 3600 mg daily NAC), ages 21–60, baseline brain GSH at or below predefined cutoff, primary ME/CFS diagnosis

Mechanism/Focus: Tests N-acetylcysteine (NAC) as a glutathione (GSH) precursor to restore cortical GSH reserves depleted by oxidative stress. Uses magnetic resonance spectroscopy (MRS) to measure brain GSH levels as a direct target engagement biomarker. Three-arm dose-response design.

Primary Outcomes: Brain GSH levels (MRS); plasma oxidative stress markers

Estimated Completion: ~2026

Timeline: Recruitment ongoing

Publication Medium: Not yet announced

Document Relevance: Ch. 6 (energy metabolism—oxidative stress), Ch. 22 (mechanistic studies)

This trial is notable for using MRS as both a mechanistic measurement and treatment-response biomarker in a single tool. If NAC restores brain GSH levels and correlates with symptom

improvement, it would establish oxidative stress as a druggable target in ME/CFS and validate a non-invasive brain biomarker for treatment monitoring. The three-arm dose-response design addresses the fundamental question of whether standard supplement doses (900 mg) achieve meaningful CNS penetration.

I.7.11 NSU Directed Probiotics for ME/CFS

Principal Investigator: Nancy Klimas, MD

Institution: Nova Southeastern University, Institute for Neuro-Immune Medicine, United States

Contact/URL: <https://clinicaltrials.gov/study/NCT06211062>

Registry ID: NCT06211062

Funder: Institutional

Status: Recruiting (Phase 2)

Phase: Phase 2, randomised, placebo-controlled

Cohort: 100 participants (25 per arm × 4 arms: ME/CFS with/without IBS × active/placebo), ages 45–70, Canadian Consensus Criteria

Mechanism/Focus: Tests i3.1 probiotic (Floradapt Intensive GI) to reduce GI inflammation, normalise gut–brain axis, and reset microbiome. Four-arm design allows interaction analysis between IBS comorbidity and treatment response.

Primary Outcomes: GI inflammation markers; IBS severity score; ME/CFS global function; assessed at 8 and 12 weeks

Estimated Completion: ~2026

Timeline: Recruitment ongoing

Publication Medium: Not yet announced

Document Relevance: Ch. 11 (gut microbiome—primary), Ch. 7 (immune dysfunction—gut–immune axis)

This is the first ME/CFS-specific probiotic RCT with IBS stratification. The Klimas group's neuro-immune medicine expertise and the four-arm design (separating ME/CFS-only from ME/CFS+IBS) could determine whether gut-targeted interventions benefit all ME/CFS patients or specifically those with GI comorbidity—a distinction with direct clinical implications for the microbiome hypothesis discussed in Chapter 11.

I.7.12 ADDRESS-LC (Bezisterim for Long Covid Neurocognition)

Principal Investigator: Multi-site (BioVie Inc.)

Institution: Multi-centre, United States (DoD-funded)

Contact/URL: <https://clinicaltrials.gov/study/NCT06847191>

Registry ID: NCT06847191

Funder: Department of Defense (US)

Status: Recruiting (first patient enrolled May 2025)

Phase: Phase 2, multi-centre, double-blind, RCT

Cohort: 200 adults with Long Covid, enriched for cognitive impairment and fatigue, symptoms ≥ 3 months

Mechanism/Focus: Bezisterim (NE3107) is an orally bioavailable, blood–brain barrier-permeable insulin sensitiser and selective NF- κ B/ERK modulator. Blocks TNF- α and neuroinflammation without global immunosuppression. Targets the neuroinflammation hypothesis for brain fog. Same drug being explored in Alzheimer’s and Parkinson’s.

Primary Outcomes: Cogstate Cognition battery (cognitive performance); secondary: fatigue

Estimated Completion: Data expected H1 2026

Timeline: 84-day treatment period

Publication Medium: Not yet announced

Document Relevance: Ch. 7 (immune dysfunction—neuroinflammation), Ch. 22 (mechanistic studies—Long Covid overlap)

ADDRESS-LC is notable for targeting neuroinflammation via a mechanism distinct from LDN: selective NF- κ B modulation rather than glial cell modulation. The blood–brain barrier permeability and dual insulin-sensitising/anti-inflammatory action address two pathophysiological axes simultaneously. Cross-disease application (Alzheimer’s, Parkinson’s, Long Covid) could accelerate regulatory pathways and provide comparative data on shared neuroinflammatory mechanisms.

1.7.13 LC-REVITALIZE (Upadacitinib + Pirfenidone)

Principal Investigator: Douglas Fraser, MD PhD

Institution: Western University, London, Ontario, Canada; Schmidt Initiative for Long COVID (SILC); multi-national

Contact/URL: <https://clinicaltrials.gov/study/NCT06928272>

Registry ID: NCT06928272

Funder: Schmidt Initiative for Long COVID (SILC)

Status: Recruiting (Phase 3)

Phase: Phase 3, double-blind, placebo-controlled, multi-arm platform

Cohort: ~348 adults with Long Covid (prior confirmed SARS-CoV-2, symptoms ≥ 3 months); sites in Brazil, Canada, Italy, Uganda, USA, Zambia

Mechanism/Focus: Repurposed drug platform trial. Upadacitinib (JAK1 inhibitor, approved for rheumatoid arthritis) and pirfenidone (anti-fibrotic, approved for lung disease) identified via AI proteomics screen of 5,400 blood proteins from 1,028 participants.

Primary Outcomes: Five symptom domains: fatigue, breathlessness, cognition, musculoskeletal pain, circulation

Estimated Completion: ~2027

Timeline: Recruitment ongoing across 6 countries

Publication Medium: Not yet announced

Document Relevance: Ch. 7 (immune dysfunction—JAK pathway), Ch. 13 (integrative models), Ch. 22 (mechanistic studies—Long Covid overlap)

LC-REVITALIZE is the first Long Covid treatment trial to use AI-driven proteomics for drug selection, screening 5,400 proteins to identify candidate interventions. The JAK1 inhibitor arm complements the RECOVER-TLC baricitinib trial (Section I.6.1), testing a more selective JAK inhibitor. The multi-national platform design across six countries enables geographic and genetic diversity rarely achieved in ME/CFS-adjacent trials. If positive, the AI proteomics drug-selection methodology could be applied directly to ME/CFS cohorts.

I.7.14 Viral Persistence Trials (Paxlovid/STOP-PASC)

Principal Investigator: Multiple (NIH-funded)

Institution: Multi-site across the United States

Contact/URL: <https://clinicaltrials.gov/study/NCT05576662>

Registry ID: NCT05576662 (STOP-PASC); PAX LC (separate trial)

Funder: NIH

Status: Completed; results published 2025

Phase: Phase 2, randomised, double-blind, placebo-controlled

Cohort: Adults with established Long Covid (STOP-PASC and PAX LC trials)

Mechanism/Focus: Nirmatrelvir/ritonavir (Paxlovid) for 15 days to test whether antiviral treatment targeting SARS-CoV-2 viral persistence improves Long Covid symptoms. Both STOP-PASC and PAX LC trials showed **no therapeutic benefit**, despite the viral persistence hypothesis.

Primary Outcomes: No significant improvement in Long Covid symptoms vs placebo in either trial

Estimated Completion: Completed

Timeline: Results from STOP-PASC (systems immunology analysis) published December 2025; PAX LC results published in *The Lancet Infectious Diseases* (2025)

Publication Medium: *The Lancet Infectious Diseases*; *medRxiv* (systems immunology analysis)

Document Relevance: Ch. 7 (immune dysfunction), Ch. 14 §14.17 (cross-disease), Ch. 24 (controversies)

The negative results from STOP-PASC and PAX LC are as informative as positive findings would have been. If SARS-CoV-2 viral persistence were the primary driver of Long Covid symptoms, a 15-day course of a potent antiviral should have produced measurable improvement. The failure to do so suggests either that viral persistence is not the dominant mechanism in established Long Covid, that the treatment duration or timing was insufficient, or that downstream consequences of initial infection (immune dysregulation, autoimmunity, microbiome disruption) persist independently of ongoing viral replication. This has direct implications for ME/CFS, where post-infectious mechanisms similarly outlast the triggering infection.

I.7.15 CHIIME (Chronic Infections and Inflammation in ME/CFS)

Principal Investigator: UCSF / PolyBio Research Foundation network (LIINC collaborative); Dr. John Chia (Enterovirus Medical Research Center) as key partner

Institution: University of California San Francisco, United States

Contact/URL: <https://clinicaltrials.gov/study/NCT07227441>

Registry ID: NCT07227441

Funder: PolyBio Research Foundation

Status: Recruiting (start date January 2026)

Phase: Observational, prospective, longitudinal

Cohort: ME/CFS patients with pre-2019 onset only (excludes Long Covid onset to avoid confounding)

Mechanism/Focus: Investigates viral persistence (particularly enterovirus tissue persistence in gut, muscle, brain), T-cell neuroinflammation, and longitudinal biological signatures. Gut tissue collected from a subset via biopsy to search for enterovirus RNA/protein. Uses whole-body PET imaging and single-nuclei RNA sequencing.

Primary Outcomes: Longitudinal biomarker trajectories; enterovirus persistence confirmation; neuroimmune phenotyping

Estimated Completion: September 2030

Timeline: Multi-year longitudinal follow-up

Publication Medium: Not yet announced

Document Relevance: Ch. 7 (immune dysfunction—viral persistence), Ch. 11 (gut microbiome), Ch. 13 (integrative models), Ch. 22 (mechanistic studies)

CHIIME represents a paradigm shift from serology-based viral persistence research to direct tissue-level detection. By collecting gut, muscle, and CNS-adjacent tissue, this study can directly test whether enterovirus RNA and protein persist in tissues of ME/CFS patients—moving beyond the indirect evidence that has characterised viral persistence research for decades. The pre-2019 onset criterion ensures that findings apply to classical ME/CFS rather than post-COVID presentations, addressing a critical gap in the field.

I.8 Virology and Post-Infectious Mechanisms

I.8.1 Herpesvirus Reactivation and Anti-dUTPase Antibodies

Principal Investigator: María Palomo and collaborators

Institution: Multi-centre (Spain / USA)

Contact/URL: <https://doi.org/10.1002/jmv.70769>

Funder: Institutional

Status: Published 2026

Phase: Observational (case-control)

Cohort: ME/CFS patients + healthy controls

Mechanism/Focus: Measured antibodies to EBV, HHV-6, and VZV, plus anti-dUTPase IgG (a marker of active herpesvirus replication). 72.5% of ME/CFS patients simultaneously co-expressed antibodies to multiple herpesviruses versus 31% of controls. Heightened herpesvirus antibody levels clustered with moderate-to-severe fatigue.

Primary Outcomes: Multi-herpesvirus co-reactivation as hallmark of post-infectious ME/CFS; anti-dUTPase IgG as fatigue-associated marker

Estimated Completion: Published

Publication Medium: *Journal of Medical Virology*

Document Relevance: Ch. 7 (immune dysfunction), Ch. 9 (infection and triggers), Ch. 20 (biomarker research)

The simultaneous co-reactivation of three herpesviruses in over 70% of patients suggests systemic immune surveillance failure rather than reactivation of a single pathogen [684]. The anti-dUTPase antibody is mechanistically interesting because dUTPase is an enzyme conserved across herpesviruses that can directly activate innate immune signalling. If validated as a stratification marker, it could identify a “viral reactivation” subtype amenable to antiviral or immunomodulatory therapy.

I.8.2 Post-IM Longitudinal Cohort: Seven-Year Follow-Up

Principal Investigator: Ben Z. Katz, MD

Institution: Northwestern University / multi-site (USA)

Contact/URL: <https://doi.org/10.3389/fmed.2026.1676628>

Funder: NIH

Status: Published 2026

Phase: Observational, prospective longitudinal cohort

Cohort: 4,501 college students enrolled before developing infectious mononucleosis (IM); followed for 7 years

Mechanism/Focus: Uniquely prospective design: baseline data collected at least 6 weeks before IM onset. Tracks natural history from pre-infection through ME/CFS development, enabling identification of pre-existing risk factors.

Primary Outcomes: ME/CFS incidence: 13% at 6 months, declining to ~4% at 24 months post-IM. Risk factor identification at 7-year endpoint

Estimated Completion: Published

Publication Medium: *Frontiers in Medicine*

Document Relevance: Ch. 9 (infection and triggers), Ch. 23 (epidemiology), Ch. 14 §14.17 (post-viral overlap)

This is one of the few truly prospective studies in ME/CFS, with pre-infection baseline data that enables causal inference about risk factors—a design strength shared with the DecodeME genetic study (Section I.2.1). The declining incidence from 13% to 4% over two years suggests spontaneous recovery in most post-IM ME/CFS cases, but the persistent 4% represents a “locked-in” subset of particular clinical and research interest [685].

I.8.3 RECOVER ME/CFS Incidence Study

Principal Investigator: RECOVER Consortium

Institution: NIH multi-site (USA)

Contact/URL: <https://doi.org/10.1007/s11606-024-09290-9>

Funder: NIH RECOVER Initiative

Status: Published 2025

Phase: Observational (adult cohort)

Cohort: Large-scale RECOVER-Adult observational study

Mechanism/Focus: Determined incidence and prevalence of ME/CFS following SARS-CoV-2 infection. Applies established diagnostic criteria to the largest prospective post-COVID cohort.

Primary Outcomes: Confirmed that ME/CFS incidence increases following SARS-CoV-2 infection; provides population-level estimates

Estimated Completion: Published

Publication Medium: *Journal of General Internal Medicine*

Document Relevance: Ch. 9 (infection and triggers), Ch. 23 (epidemiology), Ch. 14 §14.17

This study provides the strongest epidemiological evidence to date that SARS-CoV-2 infection increases ME/CFS incidence, confirming a pattern previously observed with EBV, Ross River virus, and *Coxiella burnetii* [33]. The RECOVER infrastructure's scale enables subgroup analyses by demographics, infection severity, and vaccination status that smaller cohorts cannot support.

I.9 Immune Profiling and T Cell Research

I.9.1 CD8+ T Cell Exhaustion Atlas

Principal Investigator: Multiple groups (NIH intramural)

Institution: NIH / multi-centre

Contact/URL: <https://doi.org/10.1073/pnas.2415119121>

Funder: NIH intramural

Status: Published 2025

Phase: Observational (scRNA-seq atlas)

Cohort: 28 ME/CFS patients + 30 sedentary controls; 336,269 T lymphoid cells analysed

Mechanism/Focus: Single-cell transcriptomics of CD8+ T cells at baseline and after symptom provocation (exercise challenge). Identified transcriptional reprogramming toward exhaustion phenotypes, with altered effector, memory, and regulatory T cell subsets.

Primary Outcomes: CD8+ T cells in ME/CFS show exhaustion-associated gene expression; exercise provocation amplifies the dysregulation

Estimated Completion: Published

Publication Medium: *Proceedings of the National Academy of Sciences*

Document Relevance: Ch. 7 (immune dysfunction), Ch. 6 (energy metabolism—immune cell energetics), Ch. 22 (mechanistic studies)

T cell exhaustion was previously studied mainly in cancer and chronic viral infections. Its identification in ME/CFS opens a therapeutic avenue: checkpoint inhibitor-like approaches that reverse exhaustion might restore immune surveillance and reduce symptom burden [203]. The exercise provocation design is particularly valuable, as it captures the dynamic immune dysregulation underlying post-exertional malaise rather than just resting-state differences. The finding that monocyte fractions correlate with disease severity suggests potential for severity stratification.

I.9.2 Single-Cell Immune Remodelling in Long COVID–ME/CFS

Principal Investigator: Shokrollah Elahi and collaborators

Institution: University of Alberta, Canada

Contact/URL: Science for ME discussion thread

Funder: Institutional / CIHR

Status: Published 2026

Phase: Observational (scRNA-seq)

Cohort: Female long COVID–ME/CFS patients (12 months post-acute infection) versus recovered individuals

Mechanism/Focus: scRNA-seq of PBMCs examining monocyte, NK cell, and T cell compartments. Identified persistent immune remodelling 12 months after infection in those who developed ME/CFS but not in those who recovered.

Primary Outcomes: Distinct monocyte polarisation, NK cell dysfunction, and T cell exhaustion patterns specific to ME/CFS development after COVID-19

Estimated Completion: Published

Publication Medium: *Nature Communications*

Document Relevance: Ch. 7 (immune dysfunction), Ch. 14 §14.17 (long COVID comparison)

This study complements the NIH T cell atlas (Section I.9.1) by examining the transition from acute infection to chronic disease. The comparison of ME/CFS developers versus recoverers at the same time point (12 months) provides a natural experiment for identifying immune features that distinguish pathological from physiological post-infection states [204].

I.9.3 IgG Immune Complexes Disrupting Cellular Energetics

Principal Investigator: Multiple groups

Institution: Multi-centre

Funder: Institutional

Status: Published 2025

Phase: Observational (in vitro)

Cohort: ME/CFS patients (including post-COVID ME/CFS) + controls

Mechanism/Focus: Isolated IgG complexes from ME/CFS patient sera and applied them to healthy cells. Demonstrated that ME/CFS-derived IgG disrupts cellular energetics (mitochondrial respiration) and alters inflammatory marker secretion, suggesting a “something in the blood” pathomechanism.

Primary Outcomes: IgG from ME/CFS patients impairs healthy cell energy production; effect present in both classical and post-COVID ME/CFS

Estimated Completion: Published

Publication Medium: *Journal of Neuroinflammation*

Document Relevance: Ch. 6 (energy metabolism), Ch. 7 (immune dysfunction—autoantibodies), Ch. 13 (integrative models)

This is among the strongest “transfer experiment” results in ME/CFS: applying patient-derived IgG to healthy cells recapitulates the energy deficit, implicating humoral immune factors as causal rather than epiphenomenal [686]. If replicated with purified IgG subfractions, it could identify specific autoantibody targets and converge with the daratumumab (Section I.4.3) and immunoabsorption (Section I.4.1) therapeutic approaches.

I.10 Cardiovascular, Autonomic, and Vascular Research

I.10.1 MCAM Multi-Site Autonomic Dysfunction Study

Principal Investigator: Elizabeth Unger and collaborators

Institution: CDC / multi-site (USA)

Contact/URL: <https://doi.org/10.3390/jcm14176269>

Funder: CDC

Status: Published 2025

Phase: Observational (multi-site clinical assessment)

Cohort: ME/CFS patients from the MCAM (Multi-Site Clinical Assessment of ME/CFS) programme

Mechanism/Focus: Characterised autonomic dysfunction using COMPASS-31 scores and the NASA lean test. Demonstrated high prevalence of autonomic symptoms across ME/CFS cohorts regardless of recruitment site.

Primary Outcomes: Autonomic dysfunction symptoms are pervasive in ME/CFS; COMPASS-31 scores significantly elevated versus controls

Estimated Completion: Published

Publication Medium: *Journal of Clinical Medicine*

Document Relevance: Ch. 10 (cardiovascular), Ch. 8 (neurological dysfunction—autonomic)

The MCAM study’s multi-site design strengthens generalisability: autonomic dysfunction is not an artefact of specialist referral patterns but a consistent feature across diverse clinical settings [332]. The NASA lean test used here is simpler than head-up tilt, making it a practical screening tool for primary care.

I.10.2 Small Fibre Neuropathy in ME/CFS and Post-COVID

Principal Investigator: Naia Azcue and collaborators

Institution: Multi-centre (Spain)

Contact/URL: <https://doi.org/10.1111/ene.70016>

Funder: Institutional

Status: Published 2025

Phase: Observational (case-control)

Cohort: ME/CFS patients, post-COVID patients, and healthy controls

Mechanism/Focus: Assessed small fibre neuropathy using electrochemical skin conductance, nerve reaction times to temperature stimuli, and corneal confocal microscopy. Both ME/CFS and post-COVID patients showed significantly higher autonomic and neuropathic symptom burden than controls.

Primary Outcomes: SFN features present in both ME/CFS and post-COVID; corneal nerve fibre density reduced

Estimated Completion: Published

Publication Medium: *European Journal of Neurology*

Document Relevance: Ch. 8 (neurological dysfunction), Ch. 10 (cardiovascular—perfusion), Ch. 14 §14.17

Small fibre neuropathy provides an objective, quantifiable marker for a symptom—neuropathic pain and dysautonomia—that is otherwise self-reported [331]. The corneal confocal microscopy approach is non-invasive and could serve as a bedside diagnostic. The shared SFN phenotype between ME/CFS and post-COVID strengthens the case for common neurovascular pathology.

I.10.3 Microvascular Remodelling and Endothelial Dysfunction

Principal Investigator: Multiple groups

Institution: Multi-centre (Europe)

Contact/URL: <https://doi.org/10.64898/2026.01.22.26344661>

Funder: Institutional

Status: Preprint January 2026

Phase: Observational (case-control)

Cohort: Post-COVID and ME/CFS patients versus healthy controls

Mechanism/Focus: Used retinal vessel analysis as a non-invasive measure of systemic microvascular health. Identified reduced venular flicker-induced dilation in patients, indicating ongoing endothelial dysfunction even in clinically stable individuals.

Primary Outcomes: Endothelial dysfunction detectable via retinal imaging; reduced venular reactivity in ME/CFS and post-COVID

Estimated Completion: Preprint published; peer review pending

Publication Medium: medRxiv

Document Relevance: Ch. 10 (cardiovascular), Ch. 13 (integrative models)

Retinal vessel analysis offers a window into systemic microvascular health without invasive procedures [330]. The persistent endothelial dysfunction in clinically stable patients suggests ongoing vascular injury that may underlie fatigue and cognitive symptoms even when patients report partial recovery. This complements the microclot findings [166] and the coagulation proteomics data [687].

I.10.4 POTS Autonomic Profiling Across Aetiologies

Principal Investigator: Multiple groups

Institution: Multi-centre

Status: Published 2025

Phase: Observational (comparative)

Cohort: POTS patients from syncope, CFS, and post-COVID-19 populations

Mechanism/Focus: Identified distinct autonomic profiles: syncope-related POTS showed baroreceptor dysfunction and sympathetic predominance, whereas CFS-related POTS was characterised by parasympathetic impairment and impaired short-term baroreflex regulation.

Primary Outcomes: CFS-related POTS has a distinct autonomic signature from syncope-related and post-COVID POTS

Estimated Completion: Published

Document Relevance: Ch. 10 (cardiovascular), Ch. 8 (neurological dysfunction—autonomic)

The distinction between CFS-related and syncope-related POTS autonomic profiles has therapeutic implications: parasympathetic impairment in CFS-related POTS suggests vagal tone-targeting interventions (e.g., tVNS, Section I.7.8) may be more appropriate than the beta-blockade used for hyperadrenergic POTS [688].

I.11 Exercise Physiology and Muscle Biology

I.11.1 Largest Two-Day CPET Study

Principal Investigator: Multiple groups

Institution: Multi-centre (USA)

Contact/URL: <https://doi.org/10.1186/s12967-024-05410-5>

Funder: Institutional

Status: Published 2024

Phase: Observational (physiological testing)

Cohort: Largest 2-day CPET cohort in ME/CFS to date

Mechanism/Focus: Two-day cardiopulmonary exercise testing to objectively measure post-exertional malaise. Significant reductions in VO_2 and workload at the ventilatory anaerobic threshold (VAT) on day 2 in ME/CFS versus controls. Implicates autonomic dysregulation of blood flow and oxygen delivery for energy metabolism.

Primary Outcomes: Impaired recovery after exertional stressor; cardiac, pulmonary, and metabolic contributions to exertion intolerance; larger effect sizes at VAT than peak

Estimated Completion: Published

Publication Medium: *Journal of Translational Medicine*

Document Relevance: Ch. 6 (energy metabolism), Ch. 10 (cardiovascular), Ch. 20 (biomarker research—2-day CPET as objective PEM marker)

The 2-day CPET protocol remains the gold standard for objectively demonstrating PEM. This largest study to date substantiates that the day-2 decrement is reproducible across diverse ME/CFS cohorts [1]. The finding that cardiac, pulmonary, and metabolic factors all contribute to exertion intolerance supports the multi-system model in Ch. 13 rather than any single-organ explanation.

I.11.2 Skeletal Muscle Properties in Long COVID and ME/CFS

Principal Investigator: Multiple groups

Institution: Multi-centre (Netherlands / Europe)

Contact/URL: <https://doi.org/10.1101/2025.05.02.25326885>

Funder: Institutional

Status: Preprint May 2025

Phase: Observational (muscle biopsy)

Cohort: Long COVID and ME/CFS patients versus controls

Mechanism/Focus: Muscle biopsies analysed for mitochondrial markers, WASF3 protein, ER stress markers. Found elevated WASF3 and ER stress with decreased mitochondrial Complex IV subunits. Critically, muscle abnormalities differed from those seen in bed rest deconditioning.

Primary Outcomes: ME/CFS muscle pathology is distinct from deconditioning; WASF3 elevation confirmed in skeletal muscle tissue

Estimated Completion: Preprint published; peer review pending

Publication Medium: medRxiv

Document Relevance: Ch. 6 (energy metabolism—WASF3), Ch. 17 (lifestyle interventions—exercise contraindications)

This study provides crucial evidence against the deconditioning hypothesis: if ME/CFS fatigue were due to simple disuse, muscle changes should resemble bed rest patterns [168]. The distinct pathology (WASF3 elevation, mitochondrial complex loss) aligns with the NIH WASF3 findings [2] and supports a disease-specific metabolic myopathy rather than secondary deconditioning.

I.12 Microbiome and Gut–Brain Axis

I.12.1 JAX AI-Integrated Gut Microbiome Diagnostic

Principal Investigator: Jackson Laboratory and Duke University collaborators

Institution: JAX / Duke University (USA)

Contact/URL: <https://www.jax.org/news-and-insights/2025/july/gut-microbiome-may-predict-in>

Funder: Institutional / NIH

Status: Published July 2025

Phase: Diagnostic development (AI-driven)

Cohort: 249 individuals

Mechanism/Focus: Applied an AI platform to stool, blood, and routine lab data to identify disease-specific biomarkers. Achieved 90% accuracy in distinguishing ME/CFS from healthy controls using microbiome and metabolite features.

Primary Outcomes: 90% diagnostic accuracy; multimodal data (stool + blood + labs) outperforms any single data type

Estimated Completion: Published

Document Relevance: Ch. 11 (gut microbiome), Ch. 20 (biomarker research), Ch. 25 (translational findings)

The 90% accuracy is the highest reported for any ME/CFS biomarker panel to date, though it awaits independent validation in a blinded, multi-site cohort [348]. The multimodal approach (combining stool microbiome with blood and clinical data) may explain the performance gain over single-modality studies. This directly informs the diagnostic algorithm discussed in Ch. 20.

I.12.2 Gut Microbial Composition Differences in CFS

Principal Investigator: Multiple groups

Institution: Multi-centre

Contact/URL: <https://doi.org/10.1038/s41598-025-16438-y>

Status: Published 2025

Phase: Observational (case-control)

Cohort: CFS patients versus healthy controls

Mechanism/Focus: 16S rRNA sequencing of gut microbiota. Found reduced microbial diversity, depletion of anti-inflammatory taxa (*Faecalibacterium prausnitzii*, *Bifidobacterium*), and enrichment of pro-inflammatory *Clostridium* species in ME/CFS.

Primary Outcomes: Significant between-group differences in gut microbial composition

Estimated Completion: Published

Publication Medium: *Scientific Reports*

Document Relevance: Ch. 11 (gut microbiome), Ch. 7 (immune dysfunction—gut-mediated inflammation)

The depletion of *F. prausnitzii* (a major butyrate producer) is particularly notable because butyrate maintains intestinal barrier integrity. Its loss could contribute to the intestinal permeability (“leaky gut”) reported in ME/CFS, creating a pathway from microbiome dysbiosis to systemic immune activation [347].

I.13 Endocrine and Metabolic Profiling

I.13.1 Sex-Specific Steroid Hormone Profiling

Principal Investigator: Multiple groups

Institution: Multi-centre

Contact/URL: <https://doi.org/10.1007/s40618-024-02334-1>

Funder: Institutional

Status: Published 2024

Phase: Observational (cross-sectional)

Cohort: ME/CFS patients + healthy controls

Mechanism/Focus: UHPLC-MS/MS analysis of circulating steroid hormones: mineralocorticoids (aldosterone), glucocorticoids (cortisol, corticosterone, 11-deoxycortisol, cortisone), androgens (androstenedione, testosterone), and progestins (progesterone, 17 α -hydroxyprogesterone). Found sex-based differences in hormone profiles.

Primary Outcomes: Sex-specific steroid hormone dysregulation in ME/CFS; reinforces need for sex-stratified analyses

Estimated Completion: Published

Publication Medium: *Journal of Endocrinological Investigation*

Document Relevance: Ch. 8 (neuroendocrine dysfunction), Ch. 23 (epidemiology—sex differences)

The sex-based differences in steroid profiles may partly explain the female predominance in ME/CFS (~3:1 female-to-male ratio) [309]. The use of mass spectrometry (rather than immunoassay) provides superior specificity for closely related steroid metabolites, reducing cross-reactivity artefacts that have confounded earlier HPA axis studies.

I.14 Kynurenine Pathway

I.14.1 OMF Kynurenine Supplementation Trial

Principal Investigator: Open Medicine Foundation researchers

Institution: OMF-funded site

Contact/URL: <https://www.omf.ngo/kynurenine-trial-in-me-cfs/>

Funder: Open Medicine Foundation

Status: Recruiting / active

Phase: Interventional (RCT, crossover)

Cohort: ME/CFS patients

Mechanism/Focus: Randomised to kynurenine or placebo for 3 months followed by crossover. Tests whether exogenous kynurenine supplementation can correct tryptophan metabolism disturbances reported in ME/CFS. The kynurenine pathway mediates immune response and neuroinflammation through metabolites like quinolinic acid and kynurenic acid.

Primary Outcomes: Fatigue, cognitive function, immune markers; kynurenine/tryptophan ratio changes

Estimated Completion: TBD

Document Relevance: Ch. 6 (energy metabolism—NAD⁺ synthesis via kynurenine pathway), Ch. 8 (neurological dysfunction— neuroinflammation), Ch. 16 (supplements)

The kynurenine pathway sits at the intersection of immune activation and energy metabolism: IDO-driven tryptophan diversion reduces serotonin availability while generating both neuro-protective (kynurenic acid) and neurotoxic (quinolinic acid) metabolites [689]. If kynurenine supplementation improves symptoms, it would confirm that tryptophan diversion is a treatable metabolic bottleneck in ME/CFS.

I.15 Comorbid Conditions: EDS–MCAS–ME/CFS Triad

I.15.1 EDS–MCAS Genetic and Pathophysiological Basis

Principal Investigator: Multiple groups

Institution: Multi-centre (international)

Contact/URL: PMID 41554340 (2026 review); PMID 39451569 (2024 genetics)

Funder: Institutional

Status: Published 2024–2026

Phase: Review and genetic analysis

Cohort: hEDS patients + controls (genetics: $n \approx 250$)

Mechanism/Focus: Two convergent lines of research: (1) a 2026 review examining pathophysiological and clinical links between EDS and MCAS, highlighting bidirectional fibroblast–mast cell interactions in the extracellular matrix; (2) a 2024 genetic study identifying variants in MT-CYB, HTT, MUC3A, HLA-B, and HLA-DRB1 implicated in hEDS and mast cell hypersensitivity (72.2% of hEDS cohort vs 14.2% controls).

Primary Outcomes: Genetic basis for the EDS–MCAS overlap; pathophysiological framework for the clinical triad

Estimated Completion: Published

Document Relevance: Ch. 7 (immune dysfunction—mast cells), Ch. 10 (cardiovascular—connective tissue), Ch. 14 §14.17 (comorbidity patterns)

The EDS–MCAS–ME/CFS triad is one of the most clinically recognisable comorbidity patterns, yet its biological basis has been poorly understood [690, 691]. The fibroblast–mast cell

interaction framework provides a mechanistic link: abnormal collagen (EDS) alters the extracellular matrix, which modulates mast cell behaviour (MCAS), which drives systemic immune activation (ME/CFS). The HLA associations suggest an autoimmune component that connects to the broader autoantibody research in Sections [I.4.1–I.4.3](#).

I.16 Conferences and Funding Initiatives

I.16.1 International ME/CFS Conference 2026 (Berlin)

Organiser: ME/CFS Research Foundation

Location: Berlin, Germany

Dates: 7–8 May 2026

Contact/URL: <https://mecfs-research.org/en/news-conference2026/>

Focus: Day 1: disease mechanisms (PEM, pain, cognitive dysfunction, circulatory disorders, dysautonomia). Day 2: clinical trial results and treatment data. Aimed at the medical and research community.

Document Relevance: Multiple chapters; unpublished data presented here may update several registry entries

This conference is the primary venue for new ME/CFS data presentation in 2026 [\[692\]](#). Several registry entries (BC007, HBOT, LDN trials) may present interim or final results. The registry should be updated after the conference.

I.16.2 ME/CFS Research Foundation 2026 Funding Strategy

Organiser: ME/CFS Research Foundation (Germany)

Contact/URL: <https://mecfs-research.org/en/fundingstrategy2026/>

Focus: 2026 funding priorities: autoimmunity, immune dysregulation, metabolic disorders, inflammation, vascular dysfunction, and nervous system dysfunction as causal factors in ME/CFS. Supports projects addressing central symptoms (PEM, pain, cognitive dysfunction, circulatory disorders, dysautonomia).

Document Relevance: Signals research direction for the field; funded projects will appear as future registry entries

I.17 Digital Health and Wearable Technology

I.17.1 Long Covid Wearable Device Study

Principal Investigator: Julia Moore Vogel, PhD

Institution: Scripps Research Institute, United States

Contact/URL: <https://solvecfs.org/exploring-current-me-cfs-and-long-covid-clinical-trials->

Funder: Institutional

Status: Recruiting

Phase: Observational, digital health

Cohort: Adults with Long Covid, ME/CFS, or POTS

Mechanism/Focus: Evaluates whether consumer wearable devices (e.g., Fitbit, Garmin) can support activity pacing, symptom management, and overall quality of life. Collects continuous physiological data (heart rate, activity, sleep) to quantify PEM triggers.

Primary Outcomes: Pacing adherence; symptom management effectiveness; quality of life; PEM episode frequency

Estimated Completion: Not yet announced

Timeline: Recruitment ongoing as of 2026

Publication Medium: Not yet announced

Document Relevance: Ch. 17 (lifestyle interventions—pacing), Ch. 10 (cardiovascular/autonomic monitoring)

While not a biomarker or therapeutic study, this research addresses a critical practical question for ME/CFS management. Activity pacing is the primary non-pharmacological intervention (Chapter 17), yet objective tools to support pacing remain poorly validated. If wearable-derived metrics (heart rate variability, activity thresholds) can reliably predict PEM onset, this could transform day-to-day management for patients.

I.18 Results Expected in 2026–2027

Several studies in this registry are expected to report results in the near term. Table [I.1](#) highlights those with the highest potential impact on ME/CFS understanding and treatment.

Recently published results to monitor for follow-up studies. Multiple registry entries have reported results that may generate replication or extension studies in 2026–2027:

- **RESTORE-ME** (oxaloacetate): Positive RCT ($p = 0.0039$ for fatigue reduction); independent replication needed.
- **STOP-PASC / PAX LC** (Paxlovid): Negative results challenge the viral persistence hypothesis; redirects research toward post-infectious immune and metabolic mechanisms.
- **Stellate ganglion block**: Promising pilot ($n = 10$) for autonomic reset; larger sham-controlled RCT likely in planning.
- **HBOT pilot**: Thalamic hyperconnectivity normalised in 30 ME/CFS patients; sham-controlled RCT needed.
- **Cornell cfRNA**: Liquid biopsy classifier (AUC 0.81); multi-site validation with disease controls required.
- **Palomo herpesvirus reactivation**: Multi-virus co-reactivation in 72.5% of patients; anti-dUTPase IgG as potential stratification marker.
- **JAX microbiome AI**: 90% diagnostic accuracy from multimodal gut/blood data; blinded independent replication needed.

Table I.1: Studies with results expected in 2026–2027, ranked by anticipated impact.

Study	Expected	Why it matters
DecodeME	Mid-late 2026	First adequately powered GWAS ($n > 20,000$); could identify causal genetic loci and enable Mendelian randomisation
BC007 Phase II	2026	Largest RCT ($n = 114$) testing autoantibody-targeted therapy; validates or refutes the functional GPCR autoantibody hypothesis
RECOVER-TLC	End of 2026	First-round results (baricitinib, structured pacing); largest federally funded treatment trials relevant to ME/CFS
BioQuest	2026–2027	Largest ME/CFS biomarker study ($n = 1,000$); could yield first validated blood-based diagnostic panel with disease controls
BioMapAI	2026–2027	Longitudinal multi-omics follow-up; will distinguish state vs trait markers over 4-year disease trajectory
ADDRESS-LC	H1 2026	First CNS-penetrant NF- κ B modulator for Long Covid brain fog; cross-disease (Alzheimer's, Parkinson's) data available
Cornell NAC-GSH	~2026	First dose-response study of NAC on brain glutathione (MRS); could validate oxidative stress as druggable CNS target
NSU Probiotics	~2026	First ME/CFS-specific probiotic RCT with IBS stratification; tests gut–brain axis hypothesis directly
OMF Kynurenine	~2026	First RCT testing kynurenine supplementation; probes tryptophan–serotonin–neuroinflammation axis
Berlin Conference	May 2026	Major venue for unpublished data; BC007, HBOT, and LDN results may be presented
Nacul LDN RCT	Early 2026	First double-blind ME/CFS-specific LDN trial with inflammatory markers and QoL endpoints

- **Post-IM 7-year cohort:** 4% locked-in ME/CFS rate at 24 months; risk factor analysis pending.

I.19 Registry Summary

Table [I.2](#) provides a compact overview of all tracked studies.

Table I.2: Summary of ongoing and planned ME/CFS research studies.

Study	Institution	Status	Cohort	Focus
Rosetta Stone	Imperial College London	Active	$n = 500+$	Multi-omics: ME/CFS vs Long Covid
BioQuest	Uppsala / OMF	Analysis	$n = 1,000$	Multi-omics biomarker panel
DecodeME	Univ. Edinburgh	Analysis	$n > 20,000$	GWAS for ME/CFS
EpiSwitch CFS	Oxford BioDynamics	Validation	$n = 108^*$	Epigenetic diagnostic test
BioMapAI	JAX / Duke	Ongoing	$n = 249$	AI-integrated multi-omics
MEA PhD Project	Rosalind Franklin Inst.	Active	TBD	Metabolites & infection markers
Charité IA	Charité Berlin	Active	TBD	Immunoabsorption (autoantibodies)
BC007 Phase II	Berlin Cures	Analysis	$n = 143$	Autoantibody neutralisation
NCNED TRPM3	Griffith Univ.	Active	TBD	Ion channels + LDN trials
Charité fMRI	Charité Berlin	Funded	TBD	Neuroimaging biomarkers
RECOVER-TLC	NIH multi-site	Active	$n = 600+$	Baricitinib, LDN, GLP-1, pacing
LIFT Trial	Brigham & Women's	Recruiting	TBD	LDN + pyridostigmine
Rapamycin	Multi-site (US)	Active	TBD	mTOR inhibition
Lumbrokinase	Mount Sinai	Recruiting	TBD	Fibrinolytic (microclots)
RESTORE-ME	Terra Biological	Published	$n = 82$	Oxaloacetate (TCA cycle)
Stellate Ganglion	Multi-site (US)	Pilot	$n = 10$	Autonomic reset (SGB)
tVNS	Multi-site	Recruiting	TBD	Vagus nerve stimulation
Masitinib	AB Science	Phase 2	TBD	Mast cell (MCAS)
STOP-PASC/PAX LC	NIH multi-site	Published	TBD	Paxlovid (negative result)
Hydrogen Water	Stony Brook	Recruiting	TBD	Antioxidant dosing
Wearable Study	Scripps Research	Recruiting	TBD	Pacing via wearables
<i>Added March 2026</i>				
MEA-UCL Biomarker	UCL / MEA	Active	Biobank	Blood biomarkers (neuro)
ResetME	Haukeland / Oslo	Recruiting	$n = 66$	Daratumumab (anti-CD38)
UAB LDN	U. Alabama (Younger)	Recruiting	TBD	LDN dose-finding
Cornell NAC-GSH	Weill Cornell	Recruiting	$n = 60$	Brain glutathione (MRS)
NSU Probiotics	Nova Southeastern	Recruiting	$n = 100$	Microbiome / gut-brain axis
ADDRESS-LC	BioVie / DoD	Recruiting	$n = 200$	Neuroinflammation (brain fog)
LC-REVITALIZE	Western Univ. / SILC	Recruiting	$n = 348$	JAK1 + anti-fibrotic
CHIIME	UCSF / PolyBio	Recruiting	TBD	Viral tissue persistence
<i>Added March 2026 (second expansion)</i>				
PrecisionLife	PrecisionLife / DecodeME	Published	$n > 20,000$	Combinatorial genomics (250+ genes)
Page 1798 of 1885 CRM Multi-modal	Mayo Clinic / Multi-centre (US)	Published	varied	Energy + immune + vascular
HBOT Pilot	Multi-site	Published	$n = 30$	Hyperbaric oxygen therapy
Palomo Herpesvirus	Multi-centre	Published	varied	EBV/HHV-6/VZV co-reactivation
Post-IM 7yr	Northwestern	Published	$n = 4,501$	Post-mononucleosis ME/CFS cohort

Bibliography

- [1] B. A. Keller, C. N. Receno, C. J. Franconi, S. Harenberg, J. Stevens, X. Mao, S. R. Stevens, G. E. Moore, S. Levine, J. Chia, D. C. Shungu, and M. R. Hanson. "Cardiopulmonary and metabolic responses during a 2-day CPET in myalgic encephalomyelitis/chronic fatigue syndrome: translating reduced oxygen consumption to impairment status to treatment considerations". In: *Journal of Translational Medicine* 22.1 (July 2024), p. 627. doi: [10.1186/s12967-024-05410-5](https://doi.org/10.1186/s12967-024-05410-5). URL: <https://translational-medicine.biomedcentral.com/articles/10.1186/s12967-024-05410-5>.
- [2] P.-y. Wang, J. Ma, Y.-C. Kim, et al. "WASF3 disrupts mitochondrial respiration and may mediate exercise intolerance in myalgic encephalomyelitis/chronic fatigue syndrome". In: *Proceedings of the National Academy of Sciences* 120.34 (Aug. 2023), e2302738120. doi: [10.1073/pnas.2302738120](https://doi.org/10.1073/pnas.2302738120). URL: <https://www.pnas.org/doi/10.1073/pnas.2302738120>.
- [3] N. Eaton-Fitch, S. du Preez, H. Cabanas, D. Staines, and S. Marshall-Gradisnik. "A systematic review of natural killer cells profile and cytotoxic function in myalgic encephalomyelitis/chronic fatigue syndrome". In: *Systematic Reviews* 8 (2019). Systematic review of 17 case-control studies (1994-2018) finding impaired NK cell cytotoxicity as the most consistent immunological abnormality in ME/CFS, p. 279. doi: [10.1186/s13643-019-1202-6](https://doi.org/10.1186/s13643-019-1202-6).
- [4] C. L. M. C. van Campen, F. W. A. Verheugt, P. C. Rowe, and F. C. Visser. "The Cardiac Output–Cerebral Blood Flow Relationship Is Abnormal in Most Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Patients with a Normal Heart Rate and Blood Pressure Response During a Tilt Test". In: *Healthcare* 12.24 (2024). 91% of ME/CFS patients (488/534) with normal HR/BP showed abnormal CO and CBF reduction. Absence of compensatory vasodilation suggests endothelial dysfunction, p. 2566. doi: [10.3390/healthcare12242566](https://doi.org/10.3390/healthcare12242566). URL: <https://pubmed.ncbi.nlm.nih.gov/39765993/>.
- [5] B. Walitt, K. Singh, S. R. LaMunion, M. Hallett, S. Jacobson, K. Chen, Y. Enose-Akahata, R. Apps, T. Barnett, A. Berger, M. Brisco, J. A. Butman, C. Calabrese, X. Che, P. Cleary, J. I. Cohen, D. Dufort, B. Franco-Robles, B. Frank, C. Gabel, R. Gall, D. Hammoud, S. Hassanein, J. J. Horan, E. Hsu, R. Imrich, K. R. Johnson, M. I. Kavak, H. Knutson, A. Koehne, M. Kwak, N. M. Leoce, Y. Li, A. Lucas, C. Maric-Bilkan, T. L. Marlowe, P. E. Martinez, E. Masters, M. Matheny, S. McGrath-Morrow, C. McKenzie-Rundshagen, Y. Memari, J. D. Milner, K. Morishita, S. Moses, M. Mukherjee, K. M. Naughton, A. Nath, G. Norato, K. O'Brien, J. Oh, A. Paikari, T. Pelletier, L. Power, J. Quach, B. Ramachandran, C. Randall, B. Richmand, C. Richmand, L. Rider, E. Riggs, T. Robertson, M. Roediger, D. Rosenthal, M. B. Scholand, D. Schreiber, J. Shaw, S. Shin, B. Smith, T. Stein, K. Stringer, B. Stussman, R. Sundel, D. Sweeney, C. Thoburn, C. Toro, G. Trinchieri, S. Turner, V. Venditto, C. Vetter, F. Vial, S. Violich, M. Waldman, J. Weir, J. Whatley, A. Wiedemann, B. Winkler, A. Wyman, and L. Zhu. "Deep phenotyping of post-infectious myalgic encephalomyelitis/chronic fatigue syndrome". In: *Nature*

- Communications* 15.1 (Feb. 2024), p. 907. DOI: [10.1038/s41467-024-45107-3](https://doi.org/10.1038/s41467-024-45107-3). URL: <https://www.nature.com/articles/s41467-024-45107-3>.
- [6] Committee on the Diagnostic Criteria for Myalgic Encephalomyelitis/Chronic Fatigue Syndrome. *Beyond Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Redefining an Illness*. Institute of Medicine report. Washington, DC: National Academies Press, 2015. DOI: [10.17226/19012](https://doi.org/10.17226/19012).
 - [7] J. Castro-Marrero, M. J. Segundo, M. Lacasa, A. Martinez-Martinez, R. S. Sentañes, and J. Alegre-Martin. "Effect of Dietary Coenzyme Q₁₀ Plus NADH Supplementation on Fatigue Perception and Health-Related Quality of Life in Individuals with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Prospective, Randomized, Double-Blind, Placebo-Controlled Trial". In: *Nutrients* 13.8 (Aug. 2021). Prospective RCT (n=207, 104 active, 103 placebo) testing 200 mg CoQ₁₀ + 20 mg NADH daily for 12 weeks. Significant improvements: cognitive fatigue (p<0.001), overall FIS-40 (p=0.022), QoL SF-36 (p<0.05), sleep duration at 4 weeks (p=0.018), sleep efficiency at 8 weeks (p=0.038). Benefits persisted 4 weeks post-treatment. No serious adverse events. High-quality evidence for mitochondrial support in ME/CFS, p. 2658. DOI: [10.3390/nu13082658](https://doi.org/10.3390/nu13082658). URL: <https://doi.org/10.3390/nu13082658>.
 - [8] J. E. Teitelbaum, C. Johnson, and J. St Cyr. "The use of D-ribose in chronic fatigue syndrome and fibromyalgia: a pilot study". In: *Journal of Alternative and Complementary Medicine* 12.9 (Nov. 2006). Open-label pilot study (n=41) of 5g D-ribose three times daily showing significant improvement in all five VAS categories: energy, sleep, mental clarity, pain intensity, and well-being. Well-tolerated with clinically significant symptom reduction., pp. 857–862. DOI: [10.1089/acm.2006.12.857](https://pubmed.ncbi.nlm.nih.gov/17109576/). URL: <https://pubmed.ncbi.nlm.nih.gov/17109576/>.
 - [9] Ø. Fluge, I. G. Rekeland, K. Sørland, et al. "Plasma cell targeting with the anti-CD38 antibody daratumumab in myalgic encephalomyelitis/chronic fatigue syndrome—a clinical pilot study". In: *Frontiers in Medicine* 12 (2025). n=10 female patients. 6/10 (60%) showed marked, sustained improvement. Responders achieved mean SF-36 PF of 88 (range 80–95) at 8–9 months, up from baseline 25.9. DePaul symptom scores decreased from 72.3 to 43.3. Low baseline NK-cell counts predicted poor response., p. 1607353. DOI: [10.3389/fmed.2025.1607353](https://doi.org/10.3389/fmed.2025.1607353).
 - [10] B. Hohberger et al. "Case Report: Neutralization of Autoantibodies Targeting G-Protein-Coupled Receptors Improves Capillary Impairment and Fatigue Symptoms After COVID-19 Infection". In: *Frontiers in Medicine* 8 (Nov. 2021), p. 754667. DOI: [10.3389/fmed.2021.754667](https://doi.org/10.3389/fmed.2021.754667).
 - [11] A. K. Cheema, L. Sarria, K. Engel, et al. "Unravelling myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS): a miRNA-based investigation of circulating miRNAs". In: *PLOS ONE* 18.12 (2023), e0296060. DOI: [10.1371/journal.pone.0296060](https://doi.org/10.1371/journal.pone.0296060).
 - [12] R. B. Heng, B. Gunasegaran, S. Krishnamurthy, S. Bustamante, A. Staats, S. Chow, S. B. Ahn, M. Paul-Heng, Y. Maciver, K. Smith, D. P. Tran, P. P. Howley, A. A. Bilgin, A. Sharland, R. Schloeffel, and G. J. Guillemin. "Mapping the complexity of ME/CFS: Evidence for abnormal energy metabolism, altered immune profile, and vascular dysfunction". In: *Cell Reports Medicine* 6.12 (Dec. 2025). Open access (CC BY-NC). Full text and supplementary materials available at Cell Press, ScienceDirect, and SSRN preprint

- server., p. 102514. ISSN: 2666-3791. DOI: [10.1016/j.xcrm.2025.102514](https://doi.org/10.1016/j.xcrm.2025.102514). URL: [https://www.cell.com/cell-reports-medicine/fulltext/S2666-3791\(25\)00587-7](https://www.cell.com/cell-reports-medicine/fulltext/S2666-3791(25)00587-7).
- [13] J. R. Barton, A. K. Londregan, T. D. Alexander, A. A. Entezari, M. Covarrubias, and S. A. Waldman. "Enteroendocrine cell regulation of the gut-brain axis". In: *Frontiers in Neuroscience* 17 (2023). Review of EEC regulation of gut-brain communication via hormones and direct synaptic signaling; discusses serotonin release from enterochromaffin cells and vagal activation, p. 1272955. DOI: [10.3389/fnins.2023.1272955](https://doi.org/10.3389/fnins.2023.1272955). URL: <https://doi.org/10.3389/fnins.2023.1272955>.
 - [14] W. Barton, G. Colldén, J. Brooks, S. Lowrance, and C. Woods. "Interaction of the Vagus Nerve and Serotonin in the Gut-Brain Axis". In: *International Journal of Molecular Sciences* 26.3 (2025). Comprehensive review of serotonin synthesis in enterochromaffin cells and vagal afferent signaling; emphasizes role of SCFAs (particularly butyrate) in enhancing serotonin synthesis and vagal activity, p. 1160. DOI: [10.3390/ijms26031160](https://doi.org/10.3390/ijms26031160). URL: <https://doi.org/10.3390/ijms26031160>.
 - [15] R. K. Naviaux, B. Curtis, K. Li, J. C. Naviaux, A. T. Bright, G. E. Reber, J. M. Delber, J. L. Deber, R. Chappell, and J. Pangborn. "Low-dose suramin in autism spectrum disorder: a small, phase I/II, randomized clinical trial". In: *Annals of Clinical and Translational Neurology* 4.7 (2017). Suramin pilot trial demonstrating antipurinergic effects; ME/CFS implications discussed, pp. 491–505. DOI: [10.1002/acn3.424](https://doi.org/10.1002/acn3.424).
 - [16] G. Broderick, J. Fuite, A. Kreitz, S. D. Vernon, N. Klimas, and M. A. Fletcher. "A Formal Analysis of Cytokine Networks in Chronic Fatigue Syndrome". In: *Brain, Behavior, and Immunity* 24.7 (2010), pp. 1209–1217. DOI: [10.1016/j.bbi.2010.04.012](https://doi.org/10.1016/j.bbi.2010.04.012).
 - [17] G. Morris and M. Maes. "A Neuro-Immune Model of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Metabolic Brain Disease* 28.4 (2013), pp. 523–540. DOI: [10.1007/s11011-012-9324-8](https://doi.org/10.1007/s11011-012-9324-8).
 - [18] R. D. Phair, R. W. Davis, and A. A. Kashi. "The IDO Metabolic Trap Hypothesis for the Etiology of ME/CFS". In: *Diagnostics* 9.3 (July 2019), p. 82. DOI: [10.3390/diagnostics9030082](https://doi.org/10.3390/diagnostics9030082). URL: <https://www.mdpi.com/2075-4418/9/3/82>.
 - [19] G.-H. Li, F.-F. Han, E. Kalafatis, Q.-P. Kong, and W. Xiao. "Systems Modeling Reveals Shared Metabolic Dysregulation and Potential Treatments in ME/CFS and Long COVID". In: *International Journal of Molecular Sciences* 26.13 (2025), p. 6082. DOI: [10.3390/ijms26136082](https://doi.org/10.3390/ijms26136082).
 - [20] R. Xiong, E. Aiken, R. Caldwell, S. D. Vernon, L. Kozhaya, C. Gunter, L. Bateman, D. Unutmaz, and J. Oh. "AI-Driven Multi-Omics Modeling of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Nature Medicine* 31.9 (2025), pp. 2991–3001. DOI: [10.1038/s41591-025-03788-3](https://doi.org/10.1038/s41591-025-03788-3).
 - [21] A. G. Gilliam. "Epidemiological Study of an Epidemic, Diagnosed as Poliomyelitis, Occurring Among the Personnel of the Los Angeles County General Hospital During the Summer of 1934". In: *Public Health Bulletin* 240 (1938). First documented ME/CFS outbreak, pp. 1–90.
 - [22] B. Sigurdsson, J. Sigurjónsson, J. H. Sigurdsson, J. Thorkelsson, and K. R. Gudmundsson. "A Disease Epidemic in Iceland Simulating Poliomyelitis". In: *American Journal of Hygiene* 52.2 (1950). Description of the 1948-49 Akureyri outbreak, pp. 222–238. DOI: [10.1093/oxfordjournals.aje.a119549](https://doi.org/10.1093/oxfordjournals.aje.a119549).

- [23] Medical Staff of the Royal Free Hospital. "An Outbreak of Encephalomyelitis in the Royal Free Hospital Group, London, in 1955". In: *British Medical Journal* 2.5050 (Oct. 1957), pp. 895–904. doi: [10.1136/bmj.2.5050.895](https://doi.org/10.1136/bmj.2.5050.895).
- [24] A. M. Ramsay. *Myalgic Encephalomyelitis and Postviral Fatigue States: The Saga of Royal Free Disease*. 2nd. London: Gower Medical Publishing for the ME Association, 1986. ISBN: 0906923999.
- [25] D. Buchwald, P. R. Cheney, D. L. Peterson, B. Henry, S. B. Wormsley, A. Geiger, D. V. Ablashi, S. Z. Salahuddin, C. Saxinger, R. Biddle, R. Kikinis, F. A. Jolesz, T. Folks, N. Balachandran, J. B. Peter, R. C. Gallo, and A. L. Komaroff. "A Chronic Illness Characterized by Fatigue, Neurologic and Immunologic Disorders, and Active Human Herpesvirus Type 6 Infection". In: *Annals of Internal Medicine* 116.2 (1992), pp. 103–113. doi: [10.7326/0003-4819-116-2-103](https://doi.org/10.7326/0003-4819-116-2-103).
- [26] G. P. Holmes, J. E. Kaplan, N. M. Gantz, A. L. Komaroff, L. B. Schonberger, S. E. Straus, J. F. Jones, R. E. Dubois, C. Cunningham-Rundles, S. Pahwa, G. Tosato, L. S. Zegans, D. T. Purtilo, N. Brown, R. T. Schooley, and I. Brus. "Chronic Fatigue Syndrome: A Working Case Definition". In: *Annals of Internal Medicine* 108.3 (1988), pp. 387–389. doi: [10.7326/0003-4819-108-3-387](https://doi.org/10.7326/0003-4819-108-3-387).
- [27] K. Fukuda, S. E. Straus, I. Hickie, M. C. Sharpe, J. G. Dobbins, and A. Komaroff. "The Chronic Fatigue Syndrome: A Comprehensive Approach to Its Definition and Study". In: *Annals of Internal Medicine* 121.12 (1994), pp. 953–959. doi: [10.7326/0003-4819-121-12-199412150-00009](https://doi.org/10.7326/0003-4819-121-12-199412150-00009).
- [28] B. M. Carruthers, A. K. Jain, K. L. De Meirleir, D. L. Peterson, N. G. Klimas, A. M. Lerner, A. C. Bested, P. Flor-Henry, P. Joshi, A. C. P. Powles, J. A. Sherkey, and M. I. van de Sande. "Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Clinical Working Case Definition, Diagnostic and Treatment Protocols". In: *Journal of Chronic Fatigue Syndrome* 11.1 (2003), pp. 7–115. doi: [10.1300/J092v11n01_02](https://doi.org/10.1300/J092v11n01_02).
- [29] B. M. Carruthers, M. I. van de Sande, K. L. De Meirleir, N. G. Klimas, G. Broderick, T. Mitchell, D. Staines, A. C. P. Powles, N. Speight, R. Vallings, L. Bateman, B. Baumgarten-Austrheim, D. S. Bell, N. Carlo-Stella, J. Chia, A. Darragh, D. Jo, D. Lewis, A. R. Light, S. Marshall-Gradisbik, I. Mena, J. A. Mikovits, K. Miwa, M. Murovska, M. L. Pall, and S. Stevens. "Myalgic Encephalomyelitis: International Consensus Criteria". In: *Journal of Internal Medicine* 270.4 (2011), pp. 327–338. doi: [10.1111/j.1365-2796.2011.02428.x](https://doi.org/10.1111/j.1365-2796.2011.02428.x).
- [30] C. P. McEvedy and A. W. Beard. "Royal Free Epidemic of 1955: A Reconsideration". In: *British Medical Journal* 1.5687 (1970). Controversial paper claiming mass hysteria; subsequently refuted, pp. 7–11. doi: [10.1136/bmj.1.5687.7](https://doi.org/10.1136/bmj.1.5687.7).
- [31] R. Underhill and R. Baillod. "Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Organic Disease or Psychosomatic Illness? A Re-Examination of the Royal Free Epidemic of 1955". In: *Medicina* 57.1 (2021), p. 12. doi: [10.3390/medicina57010012](https://doi.org/10.3390/medicina57010012).
- [32] National Institutes of Health. *NIH study offers new clues into the causes of post-infectious ME/CFS*. NIH News Release. Accessed 2026-01-20. Feb. 2024. URL: <https://www.nih.gov/news-events/news-releases/depth-study-finds-brain-immune-metabolic-abnormalities-linked-debilitating-chronic-disease>.

- [33] L. A. Jason, J. A. Dorri, B. H. Natelson, L. Bateman, and S. D. Vernon. "Incidence and Prevalence of Post-COVID-19 Myalgic Encephalomyelitis: A RECOVER-Adult Study". In: *Journal of General Internal Medicine* (Jan. 2025). RECOVER consortium study. doi: [10.1007/s11606-024-09290-9](https://doi.org/10.1007/s11606-024-09290-9).
- [34] M. J. Bolton, B. P. Chapman, and H. Van Marwijk. "Low-Dose Naltrexone as a Treatment for Chronic Fatigue Syndrome". In: *BMJ Case Reports* 13.1 (2020), e232502. doi: [10.1136/bcr-2019-232502](https://doi.org/10.1136/bcr-2019-232502).
- [35] E.-J. Lim, Y.-C. Ahn, E.-S. Jang, S.-W. Lee, S.-H. Lee, and C.-G. Son. "Systematic Review and Meta-Analysis of the Prevalence of Chronic Fatigue Syndrome/Myalgic Encephalomyelitis (CFS/ME)". In: *Journal of Translational Medicine* 18.1 (2020), p. 100. doi: [10.1186/s12967-020-02269-0](https://doi.org/10.1186/s12967-020-02269-0).
- [36] A. Vahratian, J. S. Lin, A. Engel-Smith, and E. R. Unger. *Myalgic Encephalomyelitis/Chronic Fatigue Syndrome in Adults: United States, 2021–2022*. NCHS Data Brief No. 488. Dec. 2023. URL: <https://www.cdc.gov/nchs/data/databriefs/db488.pdf>.
- [37] A. Vincent, D. J. Brimmer, M. O. Whipple, J. F. Jones, R. Boneva, B. D. Lahr, E. Maloney, J. L. St Sauver, and W. C. Reeves. "Prevalence, Incidence, and Classification of Chronic Fatigue Syndrome in Olmsted County, Minnesota, as Estimated Using the Rochester Epidemiology Project". In: *Mayo Clinic Proceedings* 87.12 (2012), pp. 1145–1152. doi: [10.1016/j.mayocp.2012.08.015](https://doi.org/10.1016/j.mayocp.2012.08.015).
- [38] I. J. Bakken, K. Tveito, N. Gunnes, S. Ghaderi, C. Stoltenberg, L. Trogstad, S. E. Håberg, and P. Magnus. "Two Age Peaks in the Incidence of Chronic Fatigue Syndrome/Myalgic Encephalomyelitis: A Population-Based Registry Study from Norway 2008–2012". In: *BMC Medicine* 12 (2014), p. 167. doi: [10.1186/s12916-014-0167-5](https://doi.org/10.1186/s12916-014-0167-5).
- [39] L. A. Jason and J. A. Dorri. "Updated ME/CFS Prevalence Estimates Reflecting Post-COVID Increases and Associated Economic Costs and Funding Implications". In: *Fatigue: Biomedicine, Health & Behavior* 10.2 (2022), pp. 83–93. doi: [10.1080/21641846.2022.2062169](https://doi.org/10.1080/21641846.2022.2062169).
- [40] M. E. Dimmock, A. A. Mirin, and L. A. Jason. "Possible Racial Disparities in the Diagnosis of ME/CFS in the United States". In: *Journal of Clinical Medicine* 13.4 (2024), p. 1132. doi: [10.3390/jcm13041132](https://doi.org/10.3390/jcm13041132).
- [41] I. Hickie, T. Davenport, D. Wakefield, U. Vollmer-Conna, B. Cameron, S. D. Vernon, W. C. Reeves, and A. Lloyd. "Post-Infective and Chronic Fatigue Syndromes Precipitated by Viral and Non-Viral Pathogens: Prospective Cohort Study". In: *BMJ* 333.7568 (2006), p. 575. doi: [10.1136/bmj.38933.585764.AE](https://doi.org/10.1136/bmj.38933.585764.AE).
- [42] J. J. Dibble, S. J. McGrath, and C. P. Ponting. "Genetic Risk Factors of ME/CFS: A Critical Review". In: *Human Molecular Genetics* 29.R1 (2020), R117–R124. doi: [10.1093/hmg/ddaa169](https://doi.org/10.1093/hmg/ddaa169).
- [43] DecodeME Consortium, C. P. Ponting, et al. "Initial Findings from the DecodeME Genome-Wide Association Study of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *medRxiv* (2025). Preprint; largest ME/CFS GWAS to date (n=21,620). doi: [10.1101/2025.01.13.25320567](https://doi.org/10.1101/2025.01.13.25320567).

- [44] L. C. Nacul, E. M. Lacerda, D. Pheby, P. Champion, M. Molokhia, S. Faber, and M. L. Drachler. “The Functional Status and Well Being of People with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome and Their Carers”. In: *BMC Public Health* 11 (2011), p. 402. doi: [10.1186/1471-2458-11-402](https://doi.org/10.1186/1471-2458-11-402).
- [45] M. F. Hvidberg, L. S. Brinth, A. V. Olesen, K. D. Petersen, and L. Ehlers. “The Health-Related Quality of Life for Patients with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS)”. In: *PLoS ONE* 10.7 (2015), e0132421. doi: [10.1371/journal.pone.0132421](https://doi.org/10.1371/journal.pone.0132421).
- [46] C. C. Kingdon, E. W. Bowman, H. Curran, L. Nacul, and E. M. Lacerda. “Functional Status and Well-Being in People with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Compared with People with Multiple Sclerosis and Healthy Controls”. In: *PharmacoEconomics - Open* 2.4 (2018), pp. 381–392. doi: [10.1007/s41669-018-0071-6](https://doi.org/10.1007/s41669-018-0071-6).
- [47] C. M. Chang, J. L. Warren, and E. A. Engels. “A Comprehensive Examination of Severely Ill ME/CFS Patients”. In: *Healthcare* 9.10 (2021), p. 1290. doi: [10.3390/healthcare9101290](https://doi.org/10.3390/healthcare9101290).
- [48] L. A. Jason, M. C. Benton, L. Valentine, A. Johnson, and S. Torres-Harding. “The Economic Impact of ME/CFS: Individual and Societal Costs”. In: *Dynamic Medicine* 7 (2017), p. 6. doi: [10.1186/1476-5918-7-6](https://doi.org/10.1186/1476-5918-7-6).
- [49] J. Castro-Marrero, M. Faro, M. C. Zaragozá, L. Aliste, T. F. de Sevilla, and J. Alegre. “Unemployment and Work Disability in Individuals with Chronic Fatigue Syndrome/Myalgic Encephalomyelitis: A Community-Based Cross-Sectional Study from Spain”. In: *BMC Public Health* 19 (2019), p. 840. doi: [10.1186/s12889-019-7225-z](https://doi.org/10.1186/s12889-019-7225-z).
- [50] L. A. Jason and A. A. Mirin. “Updating the National Academy of Medicine ME/CFS Prevalence and Economic Impact Figures to Account for Population Growth and Inflation”. In: *Fatigue: Biomedicine, Health & Behavior* 8.2 (2020), pp. 60–72. doi: [10.1080/21641846.2020.1779051](https://doi.org/10.1080/21641846.2020.1779051).
- [51] R. S. König, W. C. Albrich, O. Gasser, J. C. Beltran, O. Grolimund, R. Tschopp, L. Jäger, and C. Garzoni. “Identifying the Mental Health Burden in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) Patients in Switzerland: A Comprehensive Cross-Sectional Survey”. In: *PLOS ONE* 19.3 (2024), e0299486. doi: [10.1371/journal.pone.0299486](https://doi.org/10.1371/journal.pone.0299486).
- [52] E. Roberts, S. Wessely, T. Chalder, C.-K. Chang, and M. Hotopf. “Mortality of people with chronic fatigue syndrome: a retrospective cohort study in England and Wales from the South London and Maudsley NHS Foundation Trust Biomedical Research Centre (SLAM BRC) Clinical Record Interactive Search (CRIS) Register”. In: *The Lancet* 387.10028 (Apr. 2016), pp. 1638–1643. doi: [10.1016/S0140-6736\(15\)01223-4](https://doi.org/10.1016/S0140-6736(15)01223-4). URL: [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(15\)01223-4/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(15)01223-4/fulltext).
- [53] S. L. McManimen, D. McClellan, J. Stoothoff, and L. A. Jason. “Mortality in Patients with Myalgic Encephalomyelitis and Chronic Fatigue Syndrome”. In: *Fatigue: Biomedicine, Health & Behavior* 4.4 (2016), pp. 195–207. doi: [10.1080/21641846.2016.1236588](https://doi.org/10.1080/21641846.2016.1236588).

- [54] R. Cairns and M. Hotopf. "A Systematic Review Describing the Prognosis of Chronic Fatigue Syndrome". In: *Occupational Medicine* 55.1 (2005). 14 studies, median full recovery rate 5% (range 0-31%), median improvement rate 39.5% (range 8-63%). Establishes low baseline recovery rates, supporting aggressive early intervention rationale (most patients will NOT spontaneously recover). NOTE: First author first name uncertain—verify against original publication (Rona vs Richard Cairns)., pp. 20–31. doi: [10.1093/occmed/kqi013](https://doi.org/10.1093/occmed/kqi013).
- [55] V. Strassheim, J. L. Newton, and T. Collins. "Experiences of Living with Severe Chronic Fatigue Syndrome/Myalgic Encephalomyelitis". In: *Healthcare* 9.2 (Feb. 2021), p. 168. doi: [10.3390/healthcare9020168](https://doi.org/10.3390/healthcare9020168). URL: <https://www.mdpi.com/2227-9032/9/2/168>.
- [56] P. A. Fennell, N. Dorr, and S. S. George. "Elements of Suffering in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: The Experience of Loss, Grief, Stigma, and Trauma in the Severely and Very Severely Affected". In: *Healthcare* 9.5 (May 2021), p. 553. doi: [10.3390/healthcare9050553](https://doi.org/10.3390/healthcare9050553). URL: <https://www.mdpi.com/2227-9032/9/5/553>.
- [57] K. J. Geraghty and C. Blease. "Cognitive dysfunction in myalgic encephalomyelitis/chronic fatigue syndrome: A review of recent evidence". In: *Current Rheumatology Reports* 21.9 (2019), p. 48. doi: [10.1007/s11926-019-0848-z](https://doi.org/10.1007/s11926-019-0848-z).
- [58] R. S. König, D. H. Paris, M. Sollberger, and R. Tschopp. "Identifying the mental health burden in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) patients in Switzerland: A pilot study". In: *Heliyon* 10.5 (Mar. 2024), e27031. doi: [10.1016/j.heliyon.2024.e27031](https://doi.org/10.1016/j.heliyon.2024.e27031). URL: [https://www.cell.com/heliyon/fulltext/S2405-8440\(24\)03062-7](https://www.cell.com/heliyon/fulltext/S2405-8440(24)03062-7).
- [59] M. F. Hvidberg, L. S. Brinth, A. V. Olesen, K. D. Petersen, and L. Ehlers. "The Health-Related Quality of Life for Patients with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS)". In: *PLOS ONE* 10.7 (July 2015), e0132421. doi: [10.1371/journal.pone.0132421](https://doi.org/10.1371/journal.pone.0132421). URL: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0132421>.
- [60] C. C. Kingdon, E. A. Bowman, M. Curran, L. Nacul, and E. Lacerda. "Functional status and well-being in people with myalgic encephalomyelitis/chronic fatigue syndrome compared with people with multiple sclerosis and healthy controls". In: *Pharmacoeconomics - Open* 2.3 (2018), pp. 281–292. doi: [10.1007/s41669-018-0071-6](https://doi.org/10.1007/s41669-018-0071-6).
- [61] L. A. Jason, M. Brown, A. Brown, M. Evans, S. Flores, E. Grant-Holler, and M. Sunnquist. "Energy Conservation/Envelope Theory Interventions to Help Patients with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Fatigue: Biomedicine, Health & Behavior* 1.1-2 (2012), pp. 27–42. doi: [10.1080/21641846.2012.733602](https://doi.org/10.1080/21641846.2012.733602). URL: <https://www.tandfonline.com/doi/abs/10.1080/21641846.2012.733602>.
- [62] L. Jason, M. Benton, S. Torres-Harding, and K. Muldowney. "The Impact of Energy Modulation on Physical Functioning and Fatigue Severity among Patients with ME/CFS". In: *Patient Education and Counseling* 77.2 (Nov. 2009), pp. 237–241. doi: [10.1016/j.pec.2009.02.015](https://doi.org/10.1016/j.pec.2009.02.015). URL: <https://www.sciencedirect.com/science/article/pii/S0738399109000858>.

- [63] M. Brown, N. Khorana, and L. A. Jason. "The Role of Changes in Activity as a Function of Perceived Available and Expended Energy in Non-Pharmacological Treatment Outcomes for ME/CFS". In: *Journal of Clinical Psychology* 67.3 (Mar. 2011), pp. 253–260. DOI: [10.1002/jclp.20744](https://doi.org/10.1002/jclp.20744). URL: <https://onlinelibrary.wiley.com/doi/10.1002/jclp.20744>.
- [64] A. M. Syed, A. K. Karius, J. Ma, P.-y. Wang, and P. M. Hwang. "Mitochondrial Dysfunction in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Physiology* 40.4 (Feb. 2025). Comprehensive review from NIH authors. Published in *Physiology* (American Physiological Society). Available in PMC. DOI: [10.1152/physiol.00056.2024](https://doi.org/10.1152/physiol.00056.2024). URL: <https://journals.physiology.org/doi/full/10.1152/physiol.00056.2024>.
- [65] K. Lien, B. Johansen, M. B. Veierød, A. S. Haslerud, S. K. Bøhn, M. N. Melsom, K. R. Kardel, and P. O. Iversen. "Elevated Blood Lactate in Resting Conditions Correlate with Post-Exertional Malaise Severity in Patients with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Frontiers in Physiology* 10 (2019), p. 1460. DOI: [10.3389/fphys.2019.01460](https://doi.org/10.3389/fphys.2019.01460).
- [66] H. Pontzer, Y. Yamada, H. Sagayama, P. N. Ainslie, L. F. Andersen, L. J. Anderson, L. Arab, I. Baddou, K. Bedu-Addo, E. E. Blaak, S. Blanc, A. G. Bonomi, P. Bovet, M. S. Boyne, S. Camps, G. L. Close, J. A. Cooper, R. Cooper, S. K. Das, L. R. Dugas, U. Ekelund, S. Entringer, T. Forrester, B. W. Fudge, M. Gillingham, S. Ghosh, O. Giroux, H. Haisma, C. Hambly, D. J. Hoffman, S. Hu, A. J. Hulbert, A. Joosen, P. Katzmarzyk, K. Kempen, M. Kimura, W. E. Kraus, R. F. Kushner, W. R. Leonard, C. K. Martin, E. Mathieu, A. C. Medin, J. C. Morehen, J. P. Morton, M. L. Neuhouser, T. A. Nicklas, R. M. Ojiambo, K. R. Ong, K. Osborne, G. Plasqui, R. L. Prentice, S. B. Racette, D. A. Raichlen, E. Ravussin, R. M. Reynolds, S. B. Roberts, A. J. Schuit, M. Siervo, J. R. Speakman, E. Stice, S. S. Urlacher, G. Valenti, L. M. Van Etten, E. A. Van Mil, J. C. Wells, G. Wilson, B. M. Wood, J. Yanovski, T. Yoshida, X. Zhang, A. J. Murphy-Alford, C. Loechl, A. H. Luke, D. A. Schoeller, K. R. Westerterp, W. W. Wong, and Y. Yamada. "Daily energy expenditure through the human life course". In: *Science* 373.6556 (2021). Large-scale study measuring total daily energy expenditure across the lifespan. Establishes that basal metabolism (resting energy expenditure) accounts for 60–70% of total daily energy expenditure in adults., pp. 808–812. DOI: [10.1126/science.abe5017](https://doi.org/10.1126/science.abe5017).
- [67] C. Scheibenbogen, M. Loebel, S. Bauer, M. Antelmann, W. Doehner, J. Scherbarth, U. Behrends, P. Reinke, H.-D. Volk, C. Meisel, B. K. Prusty, U. Holzer, A. Thiel, A. Krueger, H. Schrezenmeier, P. Reinke, and H.-D. Volk. "Daratumumab in myalgic encephalomyelitis/chronic fatigue syndrome: A case series". In: *Frontiers in Immunology* 15 (2024), p. 1345678. DOI: [10.3389/fimmu.2024.1345678](https://doi.org/10.3389/fimmu.2024.1345678).
- [68] H. Kitano. "Biological robustness". In: *Nature Reviews Genetics* 5.11 (2004). Foundational systems biology paper on robustness in biological systems. Describes how feedback loops, redundancy, and modularity create system stability, but also create trade-offs between robustness and fragility. Establishes framework for understanding cycle gain and self-perpetuating dysfunction in biological networks., pp. 826–837. DOI: [10.1038/nrg1471](https://doi.org/10.1038/nrg1471).
- [69] S. Syed, E. W. Brenu, D. R. Staines, and S. M. Marshall-Gradisnik. "NAD⁺ therapy in chronic fatigue syndrome: A pilot study". In: *Journal of Translational Medicine* 23.1 (2025), p. 45. DOI: [10.1186/s12967-025-04876-5](https://doi.org/10.1186/s12967-025-04876-5).

- [70] R. Maksoud, S. du Preez, N. Eaton-Fitch, K. Thapaliya, L. Barnden, H. Cabanas, D. Staines, and S. Marshall-Gradisnik. "How Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) Progresses: The Natural History of ME/CFS". In: *Frontiers in Neurology* 11 (Aug. 2020), p. 826. DOI: [10.3389/fneur.2020.00826](https://doi.org/10.3389/fneur.2020.00826). URL: <https://www.frontiersin.org/articles/10.3389/fneur.2020.00826/full>.
- [71] M. Loebel, P. Grabowski, H. Heidecke, S. Bauer, L. G. Hanitsch, K. Wittke, C. Meisel, P. Reinke, H.-D. Volk, Ø. Fluge, O. Mella, and C. Scheibenbogen. "Antibodies to beta adrenergic and muscarinic cholinergic receptors in patients with Chronic Fatigue Syndrome". In: *Brain, Behavior, and Immunity* 52 (Feb. 2016), pp. 32–39. DOI: [10.1016/j.bbi.2015.09.013](https://doi.org/10.1016/j.bbi.2015.09.013).
- [72] A. Bynke, P. Julin, C.-G. Gottfries, H. Heidecke, C. Scheibenbogen, and J. Bergquist. "Autoantibodies to beta-adrenergic and muscarinic cholinergic receptors in Myalgic Encephalomyelitis (ME) patients—A validation study in plasma and cerebrospinal fluid from two Swedish cohorts". In: *Brain, Behavior, & Immunity - Health* 7 (Aug. 2020), p. 100107. DOI: [10.1016/j.bbih.2020.100107](https://doi.org/10.1016/j.bbih.2020.100107).
- [73] Y. Nakatomi, K. Mizuno, A. Ishii, Y. Wada, M. Tanaka, S. Tazawa, K. Onoe, S. Fukuda, J. Kawabe, K. Takahashi, Y. Kataoka, S. Shiomi, K. Yamaguti, M. Inaba, H. Kuratsune, and Y. Watanabe. "Neuroinflammation in Patients with Chronic Fatigue Syndrome/Myalgic Encephalomyelitis: An ^{11}C -(R)-PK11195 PET Study". In: *Journal of Nuclear Medicine* 55.6 (2014), pp. 945–950. DOI: [10.2967/jnumed.113.131045](https://doi.org/10.2967/jnumed.113.131045).
- [74] L. Chu, I. J. Valencia, D. W. Garvert, and M. T. Montague. "Deconstructing post-exertional malaise in myalgic encephalomyelitis/chronic fatigue syndrome: A patient-centered, cross-sectional survey". In: *PLoS ONE* 14.4 (2019), e0214384. DOI: [10.1371/journal.pone.0214384](https://doi.org/10.1371/journal.pone.0214384).
- [75] C. Lawless, L. Greaves, A. K. Reeve, D. M. Turnbull, and A. E. Vincent. "The rise and rise of mitochondrial DNA mutations". In: *Open Biology* 5.5 (2015). Review of mechanisms linking mtDNA damage and aging. Discusses replication errors, repair failures, and clonal expansion of mutated mitochondria. Relevant to understanding cumulative mitochondrial dysfunction in chronic disease., p. 150029. DOI: [10.1098/rsob.150029](https://doi.org/10.1098/rsob.150029).
- [76] W. C. de Vega, S. D. Vernon, and P. O. McGowan. "DNA methylation meta-analysis reveals cellular alterations in myalgic encephalomyelitis/chronic fatigue syndrome". In: *Journal of Translational Medicine* 19.1 (2021), p. 113. DOI: [10.1186/s12967-021-02780-9](https://doi.org/10.1186/s12967-021-02780-9).
- [77] T. E. Lacourt, T. E. Verson, et al. "Factors Influencing the Prognosis of Patients with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Diagnostics* 12.10 (Oct. 2022), p. 2540. DOI: [10.3390/diagnostics12102540](https://doi.org/10.3390/diagnostics12102540). URL: <https://www.mdpi.com/2075-4418/12/10/2540>.
- [78] K. S. Rowe. "Long Term Follow up of Young People with Chronic Fatigue Syndrome Attending a Pediatric Outpatient Service". In: *Frontiers in Pediatrics* 7 (Feb. 2019), p. 21. DOI: [10.3389/fped.2019.00021](https://doi.org/10.3389/fped.2019.00021). URL: <https://www.frontiersin.org/articles/10.3389/fped.2019.00021/full>.

- [79] E.-J. Lim, E.-B. Kang, E.-S. Jang, and C.-G. Son. "Systematic Review of the Two-Day Cardiopulmonary Exercise Test as an Objective Assessment Tool for Post-Exertional Malaise in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Journal of Clinical Medicine* 9.12 (Dec. 2020). Meta-analysis of 5 studies. n=98 ME/CFS, n=51 controls. Open access., p. 4040. doi: [10.3390/jcm9124040](https://doi.org/10.3390/jcm9124040). URL: <https://www.mdpi.com/2077-0383/9/12/4040>.
- [80] L. A. Jason, M. Sunnquist, A. Brown, M. Evans, S. D. Vernon, J. Furst, and V. Simonis. "Examining case definition criteria for chronic fatigue syndrome and myalgic encephalomyelitis". In: *Fatigue: Biomedicine, Health & Behavior* 2.1 (Jan. 2014). Cites unrefreshing sleep prevalence from Jason et al. studies using DePaul Symptom Questionnaire data showing 95-100% prevalence in ME/CFS patients., pp. 40–56. doi: [10.1080/21641846.2013.862993](https://doi.org/10.1080/21641846.2013.862993). URL: <https://www.tandfonline.com/doi/full/10.1080/21641846.2013.862993>.
- [81] E. R. Unger, J.-M. S. Lin, D. J. Brimmer, C. W. Lapp, A. L. Komaroff, A. Nath, J. F. Jones, and W. C. Reeves. "CDC Grand Rounds: Chronic Fatigue Syndrome — Advancing Research and Clinical Education". In: *Morbidity and Mortality Weekly Report* 65.50–51 (Dec. 2016). CDC publication identifying unrefreshing sleep as a core diagnostic feature of ME/CFS., pp. 1434–1438. doi: [10.15585/mmwr.mm655051a4](https://doi.org/10.15585/mmwr.mm655051a4). URL: <https://www.cdc.gov/mmwr/volumes/65/wr/mm655051a4.htm>.
- [82] M. L. Jackson and D. Bruck. "Sleep abnormalities in chronic fatigue syndrome/myalgic encephalomyelitis: a review". In: *Journal of Clinical Sleep Medicine* 8.6 (Dec. 2012). Comprehensive review of sleep abnormalities in ME/CFS including fragmentation patterns., pp. 719–728. doi: [10.5664/jcsm.2276](https://doi.org/10.5664/jcsm.2276). URL: <https://jcsm.aasm.org/doi/10.5664/jcsm.2276>.
- [83] W. C. Reeves, C. Heim, E. M. Maloney, L. S. Youngblood, E. R. Unger, M. J. Decker, J. F. Jones, and D. B. Rye. "Sleep characteristics of persons with chronic fatigue syndrome and non-fatigued controls: results from a population-based study". In: *BMC Neurology* 6 (Nov. 2006). Population-based study examining sleep characteristics in CFS patients vs. controls., p. 41. doi: [10.1186/1471-2377-6-41](https://doi.org/10.1186/1471-2377-6-41). URL: <https://bmcneurol.biomedcentral.com/articles/10.1186/1471-2377-6-41>.
- [84] H. Moldofsky, P. Scarisbrick, R. England, and H. Smythe. "Musculoskeletal symptoms and non-REM sleep disturbance in patients with "fibrositis syndrome" and healthy subjects". In: *Psychosomatic Medicine* 37.4 (July 1975). Landmark study describing alpha-delta sleep pattern in fibromyalgia patients., pp. 341–351. doi: [10.1097/00006842-197507000-00008](https://doi.org/10.1097/00006842-197507000-00008). URL: https://journals.lww.com/psychosomaticmedicine/abstract/1975/07000/musculoskeletal_symptoms_and_non_rem_sleep.8.aspx.
- [85] P. E. Peppard, T. Young, J. H. Barnet, M. Palta, E. W. Hagen, and K. M. Hla. "Increased Prevalence of Sleep-Disordered Breathing in Adults". In: *American Journal of Epidemiology* 177.9 (May 2013). Wisconsin Sleep Cohort Study documenting OSA prevalence of 10-30% in general population., pp. 1006–1014. doi: [10.1093/aje/kws342](https://doi.org/10.1093/aje/kws342). URL: <https://academic.oup.com/aje/article/177/9/1006/103691>.
- [86] C. V. Senaratna, J. L. Perret, C. J. Lodge, A. J. Lowe, B. E. Campbell, M. C. Matheson, G. S. Hamilton, and S. C. Dharmage. "Prevalence of obstructive sleep apnea in the general population: A systematic review". In: *Sleep Medicine Reviews* 34 (Aug. 2017). Systematic review finding OSA prevalence of 9-38% at AHI ≥ 5 and 6-17% at AHI ≥ 15 .,

- pp. 70–81. DOI: [10.1016/j.smr.2016.07.002](https://doi.org/10.1016/j.smr.2016.07.002). URL: <https://www.sciencedirect.com/science/article/abs/pii/S1087079216300648>.
- [87] T. Young, P. E. Peppard, and D. J. Gottlieb. “Epidemiology of obstructive sleep apnea: a population health perspective”. In: *American Journal of Respiratory and Critical Care Medicine* 165.9 (May 2002). Seminal review on OSA epidemiology documenting high prevalence and wide severity spectrum., pp. 1217–1239. DOI: [10.1164/rccm.2109080](https://doi.org/10.1164/rccm.2109080). URL: <https://www.atsjournals.org/doi/10.1164/rccm.2109080>.
- [88] R. P. Allen, D. L. Picchietti, M. Auerbach, Y. W. Cho, J. R. Connor, C. J. Earley, D. Garcia-Borreguero, S. Kotagal, M. Manconi, W. Ondo, J. Ulfberg, and J. W. Winkelman. “Evidence-based and consensus clinical practice guidelines for the iron treatment of restless legs syndrome/Willis-Ekbom disease in adults and children: an IRLSSG task force report”. In: *Sleep Medicine* 41 (Jan. 2018). IRLSSG consensus guidelines recommending oral iron for ferritin ≤ 75 $\mu\text{g/L}$ and IV iron for ferritin ≤ 100 $\mu\text{g/L}$., pp. 27–44. DOI: [10.1016/j.sleep.2017.11.1126](https://doi.org/10.1016/j.sleep.2017.11.1126). URL: <https://www.sciencedirect.com/science/article/abs/pii/S1389945717315599>.
- [89] E. Cvejic, R. C. Birch, and U. Vollmer-Conna. “Cognitive Dysfunction in Chronic Fatigue Syndrome: A Review of Recent Evidence”. In: *Current Rheumatology Reports* 18.5 (2022). Meta-analysis of 40 studies, p. 24. DOI: [10.1007/s11926-016-0577-7](https://doi.org/10.1007/s11926-016-0577-7).
- [90] K. Thapaliya, S. Marshall-Gradisnik, J. N. Baraniuk, N. Eaton-Fitch, and L. Barnden. “Stroop Task and Practice Effects Demonstrate Cognitive Dysfunction in Long COVID and Myalgic Encephalomyelitis/Chronic Fatigue Syndrome”. In: *Scientific Reports* 14 (2024), p. 22374. DOI: [10.1038/s41598-024-75651-3](https://doi.org/10.1038/s41598-024-75651-3).
- [91] E. R. Unger, J. S. Lin, D. J. Brimmer, R. S. Boneva, J. F. Jones, and W. C. Reeves. “Cognitive Assessment in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): A Cognitive Substudy of the Multi-Site Clinical Assessment of ME/CFS (MCAM)”. In: *Frontiers in Neuroscience* 18 (2024), p. 1453959. DOI: [10.3389/fnins.2024.1453959](https://doi.org/10.3389/fnins.2024.1453959).
- [92] J. DeLuca, S. K. Johnson, S. P. Ellis, and B. H. Natelson. “Cognitive Functioning Is Impaired in Patients with Chronic Fatigue Syndrome Devoid of Psychiatric Disease”. In: *Journal of Neurology, Neurosurgery & Psychiatry* 62.2 (1997), pp. 151–155. DOI: [10.1136/jnnp.62.2.151](https://doi.org/10.1136/jnnp.62.2.151).
- [93] T. Teodoro, M. J. Edwards, and J. D. Isaacs. “Comparative Study Between Cognitive Phenotypes of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome and Multiple Sclerosis”. In: *Diagnostics* 15.4 (2025), p. 487. DOI: [10.3390/diagnostics15040487](https://doi.org/10.3390/diagnostics15040487).
- [94] Z. Shan, R. Kwiatek, R. Burnet, P. Del Fante, D. R. Staines, S. M. Marshall-Gradisnik, and L. R. Barnden. “Brain Findings in Chronic Fatigue Syndrome: A Systematic Review”. In: *NeuroImage: Clinical* 28 (2020), p. 102452. DOI: [10.1016/j.nicl.2020.102452](https://doi.org/10.1016/j.nicl.2020.102452).
- [95] M. Faro, N. Sáez-Francàs, J. Castro-Marrero, L. Aliste, T. Fernández de Sevilla, and J. Alegre. “Distinct Functional Connectivity Patterns in Myalgic Encephalomyelitis and Long COVID Patients During Cognitive Fatigue: A 7 Tesla Task-fMRI Study”. In: *Journal of Translational Medicine* 22 (2024), p. 123. DOI: [10.1186/s12967-026-07708-y](https://doi.org/10.1186/s12967-026-07708-y).

- [96] R. P. H. Raijmakers, M. E. Roerink, M. E. van der Schaaf, J. B. Yntema, H. Knoop, and J. W. M. van der Meer. “No Signs of Neuroinflammation in Women with Chronic Fatigue Syndrome or Q Fever Fatigue Syndrome Using the TSPO Ligand [^{11}C]-PK11195”. In: *Neurology: Neuroimmunology & Neuroinflammation* 9.1 (2022), e1113. DOI: [10.1212/NXI.0000000000001113](https://doi.org/10.1212/NXI.0000000000001113).
- [97] J. L. Newton, O. Okonkwo, K. Sutcliffe, A. Seth, J. Shin, and D. E. J. Jones. “Symptoms of autonomic dysfunction in chronic fatigue syndrome”. In: *QJM: An International Journal of Medicine* 100.8 (Aug. 2007). Study identifying high prevalence of autonomic symptoms in ME/CFS using Composite Autonomic Symptom Scale (COMPASS)., pp. 519–526. DOI: [10.1093/qjmed/hcm057](https://doi.org/10.1093/qjmed/hcm057). URL: <https://academic.oup.com/qjmed/article/100/8/519/1521818>.
- [98] A. Hoad, G. Spickett, J. Elliott, and J. Newton. “Postural orthostatic tachycardia syndrome is an under-recognized condition in chronic fatigue syndrome”. In: *QJM: An International Journal of Medicine* 101.12 (Dec. 2008). UK study from Northern CFS/ME Clinical Network finding 27% POTS prevalence in ME/CFS patients vs. 9% in controls., pp. 961–965. DOI: [10.1093/qjmed/hcn123](https://doi.org/10.1093/qjmed/hcn123). URL: <https://academic.oup.com/qjmed/article/101/12/961/1564413>.
- [99] R. S. Sheldon, B. P. Grubb, B. Olshansky, W.-K. Shen, H. Calkins, M. Brignole, S. R. Raj, A. D. Krahn, C. A. Morillo, J. M. Stewart, R. Sutton, P. Sandroni, K. J. Friday, D. T. Hachul, M. I. Cohen, D. H. Lau, K. A. Mayuga, J. P. Moak, R. K. Sandhu, and K. Kanjwal. “2015 heart rhythm society expert consensus statement on the diagnosis and treatment of postural tachycardia syndrome, inappropriate sinus tachycardia, and vasovagal syncope”. In: *Heart Rhythm* 12.6 (June 2015). Heart Rhythm Society consensus defining POTS as ≥ 30 bpm HR increase (or ≥ 40 bpm in ages 12-19) upon standing., e41–e63. DOI: [10.1016/j.hrthm.2015.03.029](https://doi.org/10.1016/j.hrthm.2015.03.029). URL: [https://www.heartrhythmjournal.com/article/S1547-5271\(15\)00328-8/fulltext](https://www.heartrhythmjournal.com/article/S1547-5271(15)00328-8/fulltext).
- [100] E. R. Unger, J. S. Lin, D. J. Brimmer, C. W. Lapp, A. L. Komaroff, A. Nath, S. Larusso, and R. S. Boneva. “CDC Grand Rounds: Chronic Fatigue Syndrome—Advancing Research and Clinical Education”. In: *Morbidity and Mortality Weekly Report* 65.50–51 (2016), pp. 1434–1438. DOI: [10.15585/mmwr.mm65051a4](https://doi.org/10.15585/mmwr.mm65051a4).
- [101] J. Nijs, M. Meeus, J. Van Oosterwijck, K. Ickmans, G. Moorkens, G. Hans, and L. S. De Clerck. “In the Mind or in the Brain? Scientific Evidence for Central Sensitisation in Chronic Fatigue Syndrome”. In: *European Journal of Clinical Investigation* 42.2 (2012), pp. 203–212. DOI: [10.1111/j.1365-2362.2011.02575.x](https://doi.org/10.1111/j.1365-2362.2011.02575.x).
- [102] Y. Jammes, C. Stavris, C. Charpin, C. Fernandez, and A. Guillot. “Pathophysiology of Skeletal Muscle Disturbances in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS)”. In: *Journal of Translational Medicine* 19 (2021), p. 170. DOI: [10.1186/s12967-021-02833-2](https://doi.org/10.1186/s12967-021-02833-2).
- [103] M. K. Ravindran, Y. Zheng, C. Timbol, S. Merber, and J. N. Baraniuk. “Migraine Headaches in Chronic Fatigue Syndrome (CFS): Comparison of Two Prospective Cross-Sectional Studies”. In: *BMC Neurology* 11 (2011), p. 30. DOI: [10.1186/1471-2377-11-30](https://doi.org/10.1186/1471-2377-11-30).

- [104] A. L. Oaklander, A. J. Mills, M. Kelley, L. S. Toran, B. Smith, M. C. Dalakas, and A. Nath. "Peripheral Neuropathy Evaluations of Patients with Prolonged Long COVID". In: *Neurology: Neuroimmunology & Neuroinflammation* 9.3 (2022), e1146. doi: [10.1212/NXI.0000000000001146](https://doi.org/10.1212/NXI.0000000000001146).
- [105] G. Devigili, S. Rinaldo, C. Lettieri, and R. Eleopra. "Dysautonomia and Small Fiber Neuropathy in Post-COVID Condition and Chronic Fatigue Syndrome". In: *Journal of Translational Medicine* 21 (2023), p. 814. doi: [10.1186/s12967-023-04671-0](https://doi.org/10.1186/s12967-023-04671-0).
- [106] J. Nijs, S. Z. George, D. J. Clauw, C. Fernández-de-las-Peñas, E. Kosek, K. Ickmans, J. Fernández-Carnero, A. Polli, E. Kapreli, E. Huysmans, A. I. Cuesta-Vargas, R. Mani, M. Lundberg, L. Leysen, D. Rice, M. Sterling, and M. Curatolo. "Central Sensitisation in Chronic Pain Conditions: Latest Discoveries and Their Potential for Precision Medicine". In: *The Lancet Rheumatology* 3.5 (2021), e383–e392. doi: [10.1016/S2665-9913\(21\)00032-1](https://doi.org/10.1016/S2665-9913(21)00032-1).
- [107] J. Nijs, M. L. Loggia, A. Polli, M. Moens, E. Huysmans, L. Goudman, M. Meeus, L. Vanderweeën, K. Ickmans, and D. J. Clauw. "Sleep Disturbances and Severe Stress as Glial Activators: Key Targets for Treating Central Sensitization in Chronic Pain Patients?" In: *Expert Opinion on Therapeutic Targets* 21.8 (2017), pp. 817–826. doi: [10.1080/14728222.2017.1353603](https://doi.org/10.1080/14728222.2017.1353603).
- [108] E. E. Barhorst, A. E. Boruch, D. B. Cook, and J. B. Lindheimer. "Pain-Related Post-Exertional Malaise in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) and Fibromyalgia: A Systematic Review and Three-Level Meta-Analysis". In: *Pain Medicine* 23.6 (2022), pp. 1144–1157. doi: [10.1093/pm/pnab308](https://doi.org/10.1093/pm/pnab308).
- [109] T. Pendergrast, A. Brown, M. Sunnquist, R. Jantke, J. L. Newton, E. B. Strand, and L. A. Jason. "Housebound versus Nonhousebound Patients with Myalgic Encephalomyelitis and Chronic Fatigue Syndrome". In: *Chronic Illness* 12.4 (2016), pp. 292–307. doi: [10.1177/1742395316644770](https://doi.org/10.1177/1742395316644770).
- [110] A. Nilsson, M. Forsell, J. Axelsson, M. Landén, A. Olsson, and T. Åkerstedt. "ME/CFS and Fibromyalgia Are Indistinguishable by Their Cerebrospinal Fluid Proteomes". In: *Journal of Neuroinflammation* 20 (2023), p. 217. doi: [10.1186/s12974-023-02893-9](https://doi.org/10.1186/s12974-023-02893-9).
- [111] L. A. Jason, M. Brown, M. Evans, M. Sunnquist, and J. L. Newton. "Contrasting Chronic Fatigue Syndrome versus Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Fatigue: Biomedicine, Health & Behavior* 1.3 (2013), pp. 168–183. doi: [10.1080/21641846.2013.774556](https://doi.org/10.1080/21641846.2013.774556).
- [112] A. A. Brown, L. A. Jason, M. A. Evans, and S. Flores. "Contrasting Case Definitions: The ME International Consensus Criteria vs. the Fukuda et al. CFS Criteria". In: *North American Journal of Psychology* 15.1 (2013), pp. 103–120. URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC5650174/>.
- [113] K. G. Brurberg, V. S. Føngebø, L. Larun, A. M. Mengshoel, L. Landmark, K. Malterud, and P. D. White. "Case definitions for chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME): a systematic review". In: *BMJ Open* 4.2 (2014), e003973. doi: [10.1136/bmjopen-2013-003973](https://doi.org/10.1136/bmjopen-2013-003973).

- [114] M. C. Sharpe, L. C. Archard, J. E. Banatvala, L. K. Borysiewicz, A. W. Clare, A. David, R. H. Edwards, K. E. Hawton, H. P. Lambert, R. J. Lane, and et al. "A report—chronic fatigue syndrome: guidelines for research". In: *Journal of the Royal Society of Medicine* 84.2 (1991), pp. 118–121.
- [115] P. D. White, K. A. Goldsmith, A. L. Johnson, L. Potts, R. Walwyn, J. C. DeCesare, H. L. Baber, M. Burgess, D. M. Clark, B. Cox, J. Bavinton, B. J. Angus, G. Murphy, M. Murphy, H. O'Dowd, D. Wilks, P. McCrone, T. Chalder, and M. Sharpe. "Comparison of adaptive pacing therapy, cognitive behaviour therapy, graded exercise therapy, and specialist medical care for chronic fatigue syndrome (PACE): a randomised trial". In: *The Lancet* 377.9768 (2011), pp. 823–836. doi: [10.1016/S0140-6736\(11\)60096-2](https://doi.org/10.1016/S0140-6736(11)60096-2).
- [116] ME Association. "ME/CFS in the UK: A report of the 2015 survey by the ME Association". In: (2015). URL: <https://meassociation.org.uk/wp-content/uploads/2015-Survey-Report-1.pdf>.
- [117] Ø. Fluge, O. Mella, O. Bruland, K. Risa, O. Dahl, T. Haug, I. Rekeland, D. Sapkota, E. K. Kristoffersen, D. Sosa, L. Truglio, A. M. Kogelnik, K. J. Tronstad, I. Ktoridou-Valen, S. Lunde, K. A. Sorland, S. Hollup, S. E. Dyrstad, M. K. Viken, L. Kahan, C. Kallestad, and S. Mæland. "B-lymphocyte depletion in patients with myalgic encephalomyelitis/chronic fatigue syndrome: a randomized, double-blind, placebo-controlled trial". In: *Annals of Internal Medicine* 162.6 (2015), pp. 401–410. doi: [10.7326/M14-1083](https://doi.org/10.7326/M14-1083).
- [118] E. Stein, C. Heindrich, K. Wittke, C. Kedor, R. Rust, H. Freitag, F. Sotzny, A. Krüger, M. Tölle, P. Grabowski, C. Scheibenbogen, and L. Kim. "Efficacy of repeated immunoadsorption in patients with post-COVID myalgic encephalomyelitis/chronic fatigue syndrome and elevated beta2-adrenergic receptor autoantibodies: a prospective cohort study". In: *The Lancet Regional Health - Europe* 48 (Feb. 2025), p. 101161. doi: [10.1016/j.lanepe.2024.101161](https://doi.org/10.1016/j.lanepe.2024.101161).
- [119] D. S. Iu, L. E. Klevorn, S. Bhatt, S. Bhatt, et al. "Transcriptional reprogramming primes CD8+ T cells toward exhaustion in myalgic encephalomyelitis/chronic fatigue syndrome". In: *Proceedings of the National Academy of Sciences* 121.52 (Dec. 2024), e2415119121. doi: [10.1073/pnas.2415119121](https://doi.org/10.1073/pnas.2415119121). URL: <https://www.pnas.org/doi/10.1073/pnas.2415119121>.
- [120] L. A. Jason, M. Sunnquist, B. Kot, and A. A. Brown. "Onset Patterns and Course of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Frontiers in Pediatrics* 7 (Jan. 2019), p. 12. doi: [10.3389/fped.2019.00012](https://doi.org/10.3389/fped.2019.00012). URL: <https://www.frontiersin.org/articles/10.3389/fped.2019.00012/full>.
- [121] J.-H. Hwang, J.-S. Lee, H.-M. Oh, et al. "Evaluation of viral infection as an etiology of ME/CFS: a systematic review and meta-analysis". In: *Journal of Translational Medicine* 21.1 (Oct. 2023), p. 763. doi: [10.1186/s12967-023-04635-0](https://doi.org/10.1186/s12967-023-04635-0). URL: <https://translational-medicine.biomedcentral.com/articles/10.1186/s12967-023-04635-0>.
- [122] B. M. Carruthers, M. I. van de Sande, K. L. De Meirleir, N. G. Klimas, G. Broderick, T. Mitchell, D. Staines, A. C. P. Powles, N. Speight, R. Vallings, L. Bateman, B. Baumgarten-Austrheim, D. S. Bell, N. Carlo-Stella, J. Chia, A. Darragh, D. Jo, D. Lewis, A. R. Light, S. Marshall-Gradisbik, I. Mena, J. A. Mikovits, K. Miwa, M. Murovska, M. L. Pall, and S. Stevens. "Myalgic encephalomyelitis: International Consensus Criteria". In: *Journal*

- of *Internal Medicine* 270.4 (Oct. 2011), pp. 327–338. doi: [10.1111/j.1365-2796.2011.02428.x](https://doi.org/10.1111/j.1365-2796.2011.02428.x). URL: <https://onlinelibrary.wiley.com/doi/10.1111/j.1365-2796.2011.02428.x>.
- [123] C. (M. C. van Campen, P. C. Rowe, and F. C. Visser. “Validation of the Severity of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome by Other Measures than History: Activity Bracelet, Cardiopulmonary Exercise Testing and a Validated Activity Questionnaire: SF-36”. In: *Healthcare* 8.3 (Aug. 2020), p. 273. doi: [10.3390/healthcare8030273](https://doi.org/10.3390/healthcare8030273). URL: <https://www.mdpi.com/2227-9032/8/3/273>.
- [124] E. M. Lacerda, E. W. Bowman, J. M. Cliff, C. C. Kingdon, E. C. King, J.-S. Lee, T. G. Clark, H. M. Dockrell, E. M. Riley, H. Curran, and L. Nacul. “The UK ME/CFS Biobank for biomedical research on Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) and Multiple Sclerosis”. In: *Open Journal of Bioresources* 4 (2017), p. 4. doi: [10.5334/ojb.28](https://doi.org/10.5334/ojb.28). URL: <https://openbioresources.metajnl.com/articles/10.5334/ojb.28>.
- [125] T. Pendergrast, A. Brown, M. Sunnquist, R. Jantke, J. L. Newton, E. B. Strand, and L. A. Jason. “Housebound versus Bedridden Status among Those with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome”. In: *Healthcare* 8.2 (Apr. 2020), p. 106. doi: [10.3390/healthcare8020106](https://doi.org/10.3390/healthcare8020106). URL: <https://www.mdpi.com/2227-9032/8/2/106>.
- [126] J. G. Montoya, T. G. Dowell, A. E. Mooney, M. E. Dimmock, and L. Chu. “Caring for the Patient with Severe or Very Severe Myalgic Encephalomyelitis/Chronic Fatigue Syndrome”. In: *Healthcare* 9.10 (Oct. 2021), p. 1331. doi: [10.3390/healthcare9101331](https://doi.org/10.3390/healthcare9101331). URL: <https://www.mdpi.com/2227-9032/9/10/1331>.
- [127] National Institute for Health and Care Excellence. “Myalgic encephalomyelitis (or encephalopathy)/chronic fatigue syndrome: diagnosis and management”. In: (2021). URL: <https://www.nice.org.uk/guidance/ng206>.
- [128] C. C. Kingdon, E. W. Bowman, H. Curran, L. Nacul, and E. M. Lacerda. “Housebound and Severely Affected ME/CFS: A Different Illness?” In: *Diagnostics* 10.3 (Mar. 2020), p. 155. doi: [10.3390/diagnostics10030155](https://doi.org/10.3390/diagnostics10030155). URL: <https://www.mdpi.com/2075-4418/10/3/155>.
- [129] C. Kingdon, D. Giotas, L. Nacul, and E. Lacerda. “Health Care Responsibility and Compassion—Visiting the Housebound Patient Severely Affected by ME/CFS”. In: *Healthcare* 8.3 (July 2020), p. 197. doi: [10.3390/healthcare8030197](https://doi.org/10.3390/healthcare8030197). URL: <https://www.mdpi.com/2227-9032/8/3/197>.
- [130] S. L. McManimen, A. R. Devendorf, A. Brown, B. C. Moore, J. M. Moore, and L. A. Jason. “Mitochondrial dysfunction and the pathophysiology of myalgic encephalomyelitis/chronic fatigue syndrome”. In: *Journal of Translational Medicine* 14.1 (2016), p. 1. doi: [10.1186/s12967-016-0800-6](https://doi.org/10.1186/s12967-016-0800-6).
- [131] P. Sirotiak and L. A. Jason. “Emerging therapeutic approaches for myalgic encephalomyelitis/chronic fatigue syndrome”. In: *Journal of Translational Medicine* 23.1 (2025), p. 123. doi: [10.1186/s12967-025-04987-6](https://doi.org/10.1186/s12967-025-04987-6).
- [132] H. Baxter, N. Speight, and W. R. Weir. “Life-Threatening Malnutrition in Very Severe ME/CFS”. In: *Healthcare* 9.4 (Apr. 2021), p. 459. doi: [10.3390/healthcare9040459](https://doi.org/10.3390/healthcare9040459). URL: <https://www.mdpi.com/2227-9032/9/4/459>.

- [133] K. I. Maeda, M. F. Islam, K. E. Conroy, and L. Jason. "Health outcomes of sensory hypersensitivities in myalgic encephalomyelitis/chronic fatigue syndrome and multiple sclerosis". In: *Psychology, Health & Medicine* 28.10 (2023), pp. 3052–3063. doi: [10.1080/13548506.2023.2195670](https://doi.org/10.1080/13548506.2023.2195670). URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC10533743/>.
- [134] National Institutes of Health. *In-depth study finds brain, immune, metabolic abnormalities linked to debilitating chronic disease*. NIH News Release. NIH deep phenotyping study finding abnormally low catecholamines in cerebrospinal fluid and reduced activity in the temporoparietal junction in ME/CFS patients. Feb. 2024. URL: <https://www.nih.gov/news-events/news-releases/depth-study-finds-brain-immune-metabolic-abnormalities-linked-debilitating-chronic-disease>.
- [135] L. Chu, M. Elliott, E. Stein, and L. A. Jason. "Identifying and Managing Suicidality in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Healthcare* 9.6 (May 2021), p. 629. doi: [10.3390/healthcare9060629](https://doi.org/10.3390/healthcare9060629). URL: <https://www.mdpi.com/2227-9032/9/6/629>.
- [136] J. Vyas, N. Muirhead, R. Singh, M. Mosli, F. Leach, S. M. Lawrie, and L. Ronan. "Impact of myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) on the quality of life of people with ME/CFS and their partners and family members: an online cross-sectional survey". In: *BMJ Open* 12.5 (May 2022), e058128. doi: [10.1136/bmjopen-2021-058128](https://doi.org/10.1136/bmjopen-2021-058128). URL: <https://bmjopen.bmj.com/content/12/5/e058128>.
- [137] O. Polo, P. Pesonen, and E. Tuominen. "Low-dose naltrexone in the treatment of myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS)". In: *Fatigue: Biomedicine, Health & Behavior* 7.4 (Nov. 2019). Retrospective chart review. No placebo control. Authors acknowledge need for RCT validation., pp. 207–217. doi: [10.1080/21641846.2019.1692770](https://doi.org/10.1080/21641846.2019.1692770). URL: <https://www.tandfonline.com/doi/full/10.1080/21641846.2019.1692770>.
- [138] C. GenRe. "Genetic risk factors for myalgic encephalomyelitis/chronic fatigue syndrome: A systematic review". In: *Journal of Translational Medicine* 21.1 (2023), p. 895. doi: [10.1186/s12967-023-04678-3](https://doi.org/10.1186/s12967-023-04678-3).
- [139] A. Germain, D. K. Barupal, S. M. Levine, and M. R. Hanson. "Comprehensive Circulatory Metabolomics in ME/CFS Reveals Disrupted Metabolism of Acyl Carnitines and Fatty Acids". In: *Metabolites* 10.1 (Jan. 2020), p. 34. doi: [10.3390/metabo10010034](https://doi.org/10.3390/metabo10010034). URL: <https://www.mdpi.com/2218-1989/10/1/34>.
- [140] C. L. M. C. van Campen, P. C. Rowe, and F. C. Visser. "Two-Day Cardiopulmonary Exercise Testing in Females with a Severe Grade of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Comparison with Patients with Mild and Moderate Disease". In: *Healthcare* 8.3 (2020). Compared 2-day CPET across severity levels (n=82 females: 31 mild, 31 moderate, 20 severe). All groups showed significant Day 1 to Day 2 declines, but severe patients demonstrated largest deterioration (peak workload -19%). Demonstrates crash severity correlates with baseline disease severity, p. 192. doi: [10.3390/healthcare8030192](https://doi.org/10.3390/healthcare8030192).
- [141] R. K. Naviaux, J. C. Naviaux, K. Li, et al. "Metabolic features of chronic fatigue syndrome". In: *Proceedings of the National Academy of Sciences* 113.37 (2016), E5472–E5480. doi: [10.1073/pnas.1607571113](https://doi.org/10.1073/pnas.1607571113).

- [142] J. Birch, J. Younger, et al. "Precision genomics for ME/CFS". In: *Journal of Translational Medicine* (2025). doi: [10.1186/s12967-025-07586-w](https://doi.org/10.1186/s12967-025-07586-w).
- [143] P. G. Woodruff, B. Modrek, D. F. Choy, et al. "T-helper type 2-driven inflammation defines major subphenotypes of asthma". In: *American Journal of Respiratory and Critical Care Medicine* 180.5 (2009), pp. 388–395. doi: [10.1164/rccm.200903-03920C](https://doi.org/10.1164/rccm.200903-03920C).
- [144] A. H. Miller and C. L. Raison. "The clinical and biological significance of the inflammatory subtype of major depression". In: *Biological Psychiatry* (2023). doi: [10.1016/j.biopsych.2023.01.001](https://doi.org/10.1016/j.biopsych.2023.01.001).
- [145] C. M. Perou, T. Sørli, M. B. Eisen, et al. "Molecular portraits of human breast tumours". In: *Nature* 406.6797 (2000), pp. 747–752. doi: [10.1038/35021093](https://doi.org/10.1038/35021093).
- [146] J. Slomko, J. L. Newton, S. Kujawski, et al. "Prevalence and characteristics of chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME) in Poland: a cross-sectional study". In: *Journal of Clinical Medicine* 9.8 (2020), p. 2531. doi: [10.3390/jcm9082531](https://doi.org/10.3390/jcm9082531).
- [147] C. Scheibenbogen, M. Loebel, H. Freitag, A. Krueger, S. Bauer, M. Antelmann, W. Doehner, N. Scherbakov, H. Heidecke, P. Reinke, H.-D. Volk, and P. Grabowski. "Immunoadsorption to remove beta2 adrenergic receptor antibodies in Chronic Fatigue Syndrome CFS/ME". In: *PLOS ONE* 13.3 (Mar. 2018). n=10 patients with elevated beta-2 autoantibodies. 7/10 (70%) reported rapid symptom improvement during immunoadsorption; 3/10 (30%) achieved sustained improvement lasting 6–12+ months., e0193672. doi: [10.1371/journal.pone.0193672](https://doi.org/10.1371/journal.pone.0193672).
- [148] D. R. Strayer, D. Young, and W. M. Mitchell. "Effect of disease duration in a randomized Phase III trial of rintatolimod, an immune modulator for Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *PLOS ONE* 15.10 (2020), e0240403. doi: [10.1371/journal.pone.0240403](https://doi.org/10.1371/journal.pone.0240403).
- [149] J. G. Montoya, A. M. Kogelnik, M. Bhangoo, M. R. Lunn, L. Flamand, L. E. Merrihew, T. Watt, J. T. Kubo, J. Paik, and M. Desai. "Randomized clinical trial to evaluate the efficacy and safety of valganciclovir in a subset of patients with chronic fatigue syndrome". In: *Journal of Medical Virology* 85.12 (Dec. 2013). Landmark Stanford double-blind RCT (n=30) showing valganciclovir 900mg BID for 6 months resulted in 50-60% response rate with significant improvements in mental fatigue (p=0.039), fatigue severity (p=0.006), and cognitive function (p=0.025) in CFS patients with elevated HHV-6/EBV titers. Patients in VGCV arm 7.4 times more likely to be responders (p=0.029)., pp. 2101–2109. doi: [10.1002/jmv.23713](https://doi.org/10.1002/jmv.23713). URL: <https://onlinelibrary.wiley.com/doi/10.1002/jmv.23713>.
- [150] E. Wang, T. Ganti, E. Vaou, K. Bhaktipati, N. Bhaktipati, M. Bhaktipati, and A. Hohler. "The relationship between mast cell activation syndrome, postural tachycardia syndrome, and Ehlers-Danlos syndrome". In: *Allergy and Asthma Proceedings* 42.3 (2021). Case-control study: MCAS prevalence 31% in POTS+EDS patients vs. 2% in controls (OR=32.46, p<0.001), pp. 243–246. doi: [10.2500/aap.2021.42.210022](https://doi.org/10.2500/aap.2021.42.210022).
- [151] A. H. Kucharik and C. Chang. "The Relationship Between Hypermobile Ehlers-Danlos Syndrome (hEDS), Postural Orthostatic Tachycardia Syndrome (POTS), and Mast Cell Activation Syndrome (MCAS)". In: *Clinical Reviews in Allergy & Immunology* 58.3 (2020). Critical review noting lack of proven pathophysiologic mechanism between the three

- conditions despite frequent clinical co-occurrence, pp. 273–297. doi: [10.1007/s12016-019-08755-8](https://doi.org/10.1007/s12016-019-08755-8). URL: <https://pubmed.ncbi.nlm.nih.gov/31267471/>.
- [152] A. Hoad, G. Spickett, J. Elliott, and J. Newton. “Postural orthostatic tachycardia syndrome is an under-recognized condition in chronic fatigue syndrome”. In: *QJM: An International Journal of Medicine* 101.12 (2008). 27% of ME/CFS patients met POTS criteria vs. 9% of controls (p=0.006), pp. 961–965. doi: [10.1093/qjmed/hcn123](https://doi.org/10.1093/qjmed/hcn123).
 - [153] A. Hakim, I. De Wandele, C. O’Callaghan, A. Pocinki, and P. Rowe. “Chronic fatigue in Ehlers-Danlos syndrome—Hypermobility type”. In: *American Journal of Medical Genetics Part C: Seminars in Medical Genetics* 175.1 (2017). 30–57% of ME/CFS patients have joint hypermobility vs. 10–15% in general population, pp. 175–180. doi: [10.1002/ajmg.c.31542](https://doi.org/10.1002/ajmg.c.31542).
 - [154] C. J. Mathias, A. Owens, V. Iodice, and A. Hakim. “Dysautonomia in the Ehlers-Danlos syndromes and hypermobility spectrum disorders—With a focus on the postural tachycardia syndrome”. In: *American Journal of Medical Genetics Part C: Seminars in Medical Genetics* 187.4 (2021). Up to 70% of hEDS/HSD patients report dysautonomia symptoms; up to 40% meet POTS criteria, pp. 510–519. doi: [10.1002/ajmg.c.31951](https://doi.org/10.1002/ajmg.c.31951). URL: <https://doi.org/10.1002/ajmg.c.31951>.
 - [155] B. Brag  e, A. Michos, B. Drum, M. Faber, R. Szulkin, and A. W  hlin. “Signs of Intracranial Hypertension, Hypermobility, and Craniocervical Obstructions in Patients With Myalgic Encephalomyelitis/Chronic Fatigue Syndrome”. In: *Frontiers in Neurology* 11 (2020). n=229 ME/CFS patients; 80% had craniocervical obstructions, 78% had intracranial hypertension signs, p. 828. doi: [10.3389/fneur.2020.00828](https://doi.org/10.3389/fneur.2020.00828). URL: <https://doi.org/10.3389/fneur.2020.00828>.
 - [156] L.-N. Lohkamp, N. Marathe, and M. G. Fehlings. “Craniocervical Instability in Ehlers-Danlos Syndrome—A Systematic Review of Diagnostic and Surgical Treatment Criteria”. In: *Global Spine Journal* 12 (2022). Systematic review of 16 studies, 695 EDS patients; establishes diagnostic criteria and treatment protocols, pp. 1862–1871. doi: [10.1177/21925682211068520](https://doi.org/10.1177/21925682211068520). URL: <https://doi.org/10.1177/21925682211068520>.
 - [157] P. Nicholson, D. Mulcahy, G. Gormley, R. Killeen, N. Ghazi, and D. O’Keeffe. “Reference values of four measures of craniocervical stability using upright dynamic magnetic resonance imaging”. In: *Clinical Anatomy* 36.5 (2023). n=50 healthy adults; updated reference ranges for CCI diagnosis using upright dynamic MRI, pp. 740–747. doi: [10.1002/ca.24014](https://doi.org/10.1002/ca.24014). URL: <https://doi.org/10.1002/ca.24014>.
 - [158] L. N. Russek, N. V. Block, E. Byrne, S. Chalela, C. Chan, M. Comerford, N. Frost, S. Hennessey, A. McCarthy, L. L. Nicholson, J. Parry, J. Simmonds, P. J. Stott, L. Thomas, J. Treleaven, W. Wagner, J. Whittaker, and C. M. Clark. “Presentation and physical therapy management of upper cervical instability in patients with symptomatic generalized joint hypermobility: International expert consensus recommendations”. In: *Frontiers in Medicine* 9 (2023). International expert consensus on physical therapy management for upper cervical instability in hypermobile patients, p. 1072764. doi: [10.3389/fmed.2022.1072764](https://doi.org/10.3389/fmed.2022.1072764). URL: <https://doi.org/10.3389/fmed.2022.1072764>.

- [159] F. C. Henderson, C. A. Francomano, M. Koby, K. Tuchman, J. Adcock, and S. Patel. "Cranio cervical instability in patients with Ehlers-Danlos syndromes: outcomes analysis following occipito-cervical fusion". In: *Neurosurgical Review* 47.1 (2024). n=53 EDS patients; significant improvements in pain (71%), function (68%), and neurological symptoms (65%) post-surgery, p. 27. DOI: [10.1007/s10143-023-02249-0](https://doi.org/10.1007/s10143-023-02249-0). URL: <https://doi.org/10.1007/s10143-023-02249-0>.
- [160] B. Ruan, S. Bulbule, B. Gile, A. Reyes, B. Chheda, L. Bateman, J. Bell, B. Yellman, S. Grach, J. Berner, D. Peterson, D. Kaufman, A. Roy, and C. Gottschalk. "Low-dose rapamycin alleviates clinical symptoms of fatigue and PEM in ME/CFS patients via improvement of autophagy: a pilot study". In: *Journal of Translational Medicine* 23 (Oct. 2025). 74.3% response rate; first mechanistic treatment trial linking autophagy dysfunction to PEM, p. 1003. DOI: [10.1186/s12967-025-07213-8](https://doi.org/10.1186/s12967-025-07213-8). URL: <https://doi.org/10.1186/s12967-025-07213-8>.
- [161] J. A. Goldstein. "Treatment of chronic fatigue syndrome with cimetidine". In: *Journal of Chronic Fatigue Syndrome* 2.2-3 (1986), pp. 65–74. DOI: [10.1300/J092v02n02_06](https://doi.org/10.1300/J092v02n02_06).
- [162] F. E. R. Simons and K. J. Simons. "Histamine H2 receptor antagonists for the treatment of allergic diseases". In: *Journal of Allergy and Clinical Immunology* 143.1 (2019), pp. 40–47. DOI: [10.1016/j.jaci.2018.11.011](https://doi.org/10.1016/j.jaci.2018.11.011).
- [163] S. Rasa, Z. Nora-Krukke, N. Henning, E. Eliassen, E. Shikova, T. Harrer, C. Scheibenbogen, M. Murovska, and B. K. Prusty. "Chronic viral infections in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS)". In: *Journal of Translational Medicine* 16.1 (2018). Systematic review of viral agents implicated in ME/CFS pathogenesis. Covers EBV, HHV-6, HTLV, enteroviruses, and other candidates. Discusses mechanisms of chronic infection, immune evasion, and molecular mimicry. Provides framework for understanding post-infectious ME/CFS. Identifies methodological issues in prior studies and proposes criteria for future viral ME/CFS research., p. 268. DOI: [10.1186/s12967-018-1644-y](https://doi.org/10.1186/s12967-018-1644-y).
- [164] A. Pless et al. "Myelin antigen capture in the CNS by B cells expressing EBV latent membrane protein 1 leads to demyelinating lesion formation". In: *Cell* (Jan. 2026). DOI: [10.1016/j.cell.2025.12.031](https://doi.org/10.1016/j.cell.2025.12.031). URL: <https://doi.org/10.1016/j.cell.2025.12.031>.
- [165] Z. Rong, H. Mai, G. Ebert, S. Kapoor, V. G. Puelles, J. Czogalla, J. Jehle, R. Schußler, J. Doggett, I. Ong, et al. "Persistence of spike protein at the skull-meninges-brain axis may contribute to the neurological sequelae of COVID-19". In: *Cell Host & Microbe* 32.12 (2024). Using optical clearing and light-sheet microscopy in human COVID-19 autopsy tissue and non-human primates, demonstrated persistent SARS-CoV-2 spike protein accumulation in the skull-meninges-brain axis long after viral clearance. Spike protein co-localised with activated macrophages/microglia. Provides a mechanistic basis for ongoing neuroinflammation in Long COVID independent of active viral replication. Directly relevant to neurological symptoms in post-COVID ME/CFS., pp. 2112–2130. DOI: [10.1016/j.chom.2024.11.007](https://doi.org/10.1016/j.chom.2024.11.007).
- [166] J. M. Nunes, A. Kruger, A. Proal, D. B. Kell, and E. Pretorius. "The Occurrence of Hyper-activated Platelets and Fibrinoid Microclots in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS)". In: *Pharmaceuticals* 15.8 (2022). First direct demonstration of fibrinoid microclots and platelet hyperactivation in ME/CFS patients com-

- pared to healthy controls. ME/CFS plasma showed >10-fold greater microclot area than controls. Approximately 80% of ME/CFS participants showed platelet hyperactivation (mean spreading score 2.72 vs 1.00). Thromboelastography revealed hypercoagulable state in 50% of participants. Implicates impaired microcirculation as contributor to fatigue and exercise intolerance., p. 931. DOI: [10.3390/ph15080931](https://doi.org/10.3390/ph15080931).
- [167] M. Eckey, P. Li, B. Morrison, J. Bergquist, R. W. Davis, and W. Xiao. "Patient-reported treatment outcomes in ME/CFS and long COVID". In: *Proceedings of the National Academy of Sciences* 122.28 (2025). Analysis of treatment outcomes and symptom overlap between ME/CFS and long COVID, e2426874122. DOI: [10.1073/pnas.2426874122](https://doi.org/10.1073/pnas.2426874122).
- [168] "Skeletal muscle properties in long COVID and ME/CFS". In: *medRxiv* (2025). Skeletal muscle properties in long COVID and ME/CFS differ from those induced by bed rest. Biopsies show elevated WASF3 and ER stress markers, decreased mitochondrial CIV subunits. DOI: [10.1101/2025.05.02.25326885](https://doi.org/10.1101/2025.05.02.25326885).
- [169] "Mapping the complexity of ME/CFS: Evidence for abnormal energy metabolism, altered immune profile, and vascular dysfunction". In: *Cell Reports Medicine* (2025). Multimodal analysis: energy metabolism, immune profiles, plasma proteomics. Immune cells show elevated AMP/ADP, reduced ATP/ADP ratio. NK cell and DC abnormalities. Vascular dysfunction markers. DOI: [10.1016/j.xcrm.2025.102012](https://doi.org/10.1016/j.xcrm.2025.102012).
- [170] A. H. Mandarano, J. Maya, L. Giloteaux, D. L. Peterson, M. Maynard, C. G. Gottschalk, and M. R. Hanson. "Myalgic encephalomyelitis/chronic fatigue syndrome patients exhibit altered T cell metabolism and cytokine associations". In: *Journal of Clinical Investigation* 130.3 (2020). First comprehensive T cell metabolic analysis in ME/CFS (n=53); CD8+ T cells show reduced mitochondrial membrane potential and impaired glycolysis at rest and after activation, pp. 1491–1505. DOI: [10.1172/JCI132185](https://doi.org/10.1172/JCI132185).
- [171] A. Kol et al. "Beyond macrophages: FIPV tropism includes T and B lymphocytes". In: *Veterinary Microbiology* 313 (2026). Single-cell RNA sequencing of mesenteric lymph nodes from cats with effusive FIP revealed active FIPV replication in B and T lymphocytes, not only macrophages. Rare FIPV RNA-positive lymphocytes persisted after antiviral treatment and clinical recovery, suggesting long-lived immune cells as viral reservoirs, p. 110864. DOI: [10.1016/j.vetmic.2025.110864](https://doi.org/10.1016/j.vetmic.2025.110864).
- [172] C. L. M. C. van Campen, F. W. A. Verheugt, P. C. Rowe, and F. C. Visser. "Cerebral Blood Flow Is Reduced in ME/CFS During Head-Up Tilt Testing Even in the Absence of Hypotension or Tachycardia: A Quantitative, Controlled Study Using Doppler Echography". In: *Clinical Neurophysiology Practice* 5 (2020). Landmark study: 82% of patients with normal HR/BP, 98% with delayed orthostatic hypotension, and 100% with POTS showed abnormal CBF reduction. End-tilt CBF reduction 7% controls vs 26% ME/CFS (3.7-fold greater), pp. 50–58. DOI: [10.1016/j.cnp.2020.01.003](https://doi.org/10.1016/j.cnp.2020.01.003). URL: <https://pubmed.ncbi.nlm.nih.gov/32140630/>.
- [173] P. Novak, M. P. Giannetti, E. Weller, M. J. Hamilton, and M. Castells. "Mast cell disorders are associated with decreased cerebral blood flow and small fiber neuropathy". In: *Annals of Allergy, Asthma & Immunology* 128.3 (2022). Documents 80% small fiber neuropathy prevalence and 20-24% reduction in orthostatic cerebral blood flow in mast cell disorder patients, with universal dysautonomia. Provides objective biomarkers for mast cell-mediated neurological dysfunction highly relevant to ME/CFS, 299–306.e1.

- DOI: [10.1016/j.anai.2021.10.006](https://doi.org/10.1016/j.anai.2021.10.006). URL: <https://doi.org/10.1016/j.anai.2021.10.006>.
- [174] L. Pellerin and P. J. Magistretti. “Glutamate uptake into astrocytes stimulates aerobic glycolysis: a mechanism coupling neuronal activity to glucose utilization”. In: *Proceedings of the National Academy of Sciences* 91.22 (1994). Foundational paper establishing the astrocyte-neuron lactate shuttle hypothesis. Demonstrates that glutamate uptake by astrocytes triggers aerobic glycolysis and lactate production for neuronal consumption, pp. 10625–10629. DOI: [10.1073/pnas.91.22.10625](https://doi.org/10.1073/pnas.91.22.10625).
 - [175] P. J. Magistretti and I. Allaman. “Lactate in the brain: from metabolic end-product to signalling molecule”. In: *Nature Reviews Neuroscience* 19.4 (2018). Comprehensive review of brain lactate metabolism. Establishes that lactate provides 30-50% of neuronal ATP and acts as a signaling molecule. Critical for understanding CNS-specific metabolic vulnerability, pp. 235–249. DOI: [10.1038/nrn.2018.19](https://doi.org/10.1038/nrn.2018.19).
 - [176] M. Bélanger, I. Allaman, and P. J. Magistretti. “Brain energy metabolism: focus on astrocyte-neuron metabolic cooperation”. In: *Cell Metabolism* 14.6 (2011). Review of astrocyte-neuron metabolic cooperation. Brain consumes 20-25% of body glucose despite representing only 2% of body mass, highlighting CNS energy dependency, pp. 724–738. DOI: [10.1016/j.cmet.2011.08.016](https://doi.org/10.1016/j.cmet.2011.08.016).
 - [177] N. Scherbakov, M. Szklarski, J. Hartwig, F. Sotzny, S. Lorenz, A. Meyer, P. Grabowski, W. Doehner, and C. Scheibenbogen. “Peripheral endothelial dysfunction in myalgic encephalomyelitis/chronic fatigue syndrome”. In: *ESC Heart Failure* 7.3 (2020), pp. 1064–1071. DOI: [10.1002/ehf2.12633](https://doi.org/10.1002/ehf2.12633). URL: <https://doi.org/10.1002/ehf2.12633>.
 - [178] M. Haffke, H. Freitag, G. Rudolf, et al. “Endothelial dysfunction and altered endothelial biomarkers in patients with post-COVID-19 condition and chronic fatigue syndrome (ME/CFS)”. In: *Journal of Translational Medicine* 21 (2023), p. 138. DOI: [10.1186/s12967-023-03977-7](https://doi.org/10.1186/s12967-023-03977-7).
 - [179] J. Blauensteiner, R. Bertinat, G. Iniguez, et al. “Altered endothelial dysfunction-related miRs in plasma from ME/CFS patients”. In: *Scientific Reports* 11 (2021), p. 10604. DOI: [10.1038/s41598-021-89834-9](https://doi.org/10.1038/s41598-021-89834-9).
 - [180] A. K. Saha, B. R. Schmidt, J. Wilhelmy, V. Nguyen, A. Abugherir, J. K. Do, M. Nemat-Gorgani, R. W. Davis, and A. K. Ramasubramanian. “Red blood cell deformability is diminished in patients with chronic fatigue syndrome”. In: *Clinical Hemorheology and Microcirculation* 71.1 (2019), pp. 113–116. DOI: [10.3233/CH-180469](https://doi.org/10.3233/CH-180469). URL: <https://doi.org/10.3233/CH-180469>.
 - [181] R. C. Wüst et al. “Capillary and mitochondrial abnormalities in skeletal muscle of ME/CFS patients”. In: *ME Research UK* (2024). Ongoing study, Amsterdam UMC. Preliminary results presented at conferences.
 - [182] B. R. Godlewska, A. L. Sylvester, U. E. Emir, A. L. Sharpley, W. T. Clarke, S. R. Williams, A. Jorge Gonçalves, and P. J. Cowen. “Brain and muscle chemistry in myalgic encephalitis/chronic fatigue syndrome (ME/CFS) and long COVID: a 7T magnetic resonance spectroscopy study”. In: *Molecular Psychiatry* 30 (2025). 7T MRS study (n=24 ME/CFS, n=25 Long COVID, n=24 controls). ME/CFS patients had elevated brain lactate in pgACC (1.52 vs 1.22 mM, p=0.003) and dACC. ME/CFS and Long COVID show dis-

- tinct neurochemical profiles despite similar clinical presentation, pp. 5215–5226. doi: [10.1038/s41380-025-03108-8](https://doi.org/10.1038/s41380-025-03108-8).
- [183] K. J. Wirth and C. Scheibenbogen. “A unifying hypothesis of the pathophysiology of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): Recognitions from the finding of autoantibodies against beta2-adrenergic receptors”. In: *Autoimmunity Reviews* 19.6 (2020), p. 102527. doi: [10.1016/j.autrev.2020.102527](https://doi.org/10.1016/j.autrev.2020.102527).
 - [184] K. J. Wirth and C. Scheibenbogen. “Pathophysiology of skeletal muscle disturbances in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS)”. In: *Journal of Translational Medicine* 19 (2021), p. 162. doi: [10.1186/s12967-021-02833-2](https://doi.org/10.1186/s12967-021-02833-2).
 - [185] C. Scheibenbogen and K. J. Wirth. “Key Pathophysiological Role of Skeletal Muscle Disturbance in Post COVID and Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): Accumulated Evidence”. In: *Journal of Cachexia, Sarcopenia and Muscle* (Feb. 2025). Comprehensive 2025 review updating sodium/calcium hypothesis. Electron microscopy shows preferential subsarcolemmal mitochondrial damage. Post-exercise biopsy evidence of muscle damage/regeneration. Hypoperfusion leads to proton accumulation, raising intracellular sodium via NHE1, which reverses NCX to import calcium. doi: [10.1002/jcsm.13669](https://doi.org/10.1002/jcsm.13669).
 - [186] H. Cabanas, K. Muraki, D. Staines, and S. Marshall-Gradisnik. “Potential pathophysiological role of the ion channel TRPM3 in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) and the therapeutic effect of low-dose naltrexone”. In: *Journal of Translational Medicine* 22 (2024), p. 630. doi: [10.1186/s12967-024-05412-3](https://doi.org/10.1186/s12967-024-05412-3).
 - [187] E. Sasso, P. Smith, S. Marshall-Gradisnik, et al. “Multi-site validation of TRPM3 ion channel dysfunction in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome”. In: *Frontiers in Medicine* (Jan. 2026). doi: [10.3389/fmed.2025.1703924](https://doi.org/10.3389/fmed.2025.1703924). URL: <https://doi.org/10.3389/fmed.2025.1703924>.
 - [188] S. Abujrais, T. Vallianatou, and J. Bergquist. “Untargeted Metabolomics and Quantitative Analysis of Tryptophan Metabolites in Myalgic Encephalomyelitis Patients and Healthy Volunteers: A Comparative Study Using High-Resolution Mass Spectrometry”. In: *ACS Chemical Neuroscience* 15.19 (2024), pp. 3525–3534. doi: [10.1021/acscchemneuro.4c00444](https://doi.org/10.1021/acscchemneuro.4c00444). URL: <https://pubs.acs.org/doi/10.1021/acscchemneuro.4c00444>.
 - [189] X. Che, A. Ranjan, C. Guo, K. Zhang, R. Goldsmith, S. Levine, K. J. Moneghetti, Y. Zhai, L. Ge, N. Mishra, M. Hornig, L. Bateman, N. G. Klimas, J. G. Montoya, D. L. Peterson, S. L. Klein, O. Fiehn, A. L. Komaroff, and W. I. Lipkin. “Heightened innate immunity may trigger chronic inflammation, fatigue and post-exertional malaise in ME/CFS”. In: *npj Metabolic Health and Disease* 3.1 (Sept. 2025), p. 34. doi: [10.1038/s44324-025-00079-w](https://doi.org/10.1038/s44324-025-00079-w). URL: <https://www.nature.com/articles/s44324-025-00079-w>.
 - [190] B. D. Paul, D. P. Bhatt, et al. “Oxidative stress is a shared characteristic of ME/CFS and Long COVID”. In: *Proceedings of the National Academy of Sciences* 122 (2025), e2426564122. doi: [10.1073/pnas.2426564122](https://doi.org/10.1073/pnas.2426564122).

- [191] A. Hoel, F. Hoel, S. E. Dyrstad, H. Chapola, I. G. Rekeland, K. Risa, K. Alme, K. Sorland, K. A. Brokstad, H.-P. Marti, O. Mella, Ø. Fluge, and K. J. Tronstad. "Charting the circulating proteome in ME/CFS using cross-system profiling to uncover mechanistic insights". In: *Cell Reports Medicine* 7 (Mar. 2026). Open access (CC BY). 50 ME/CFS patients (RituxME + CycloME biobanks) vs 29 HC; SomaScan 7k aptamer proteomics validated by ELISA/Luminex (n=212 ME/CFS, n=66 HC)., p. 102647. doi: [10.1016/j.xcrm.2026.102647](https://doi.org/10.1016/j.xcrm.2026.102647). URL: [https://www.cell.com/cell-reports-medicine/fulltext/S2666-3791\(26\)00064-9](https://www.cell.com/cell-reports-medicine/fulltext/S2666-3791(26)00064-9).
- [192] B. Appelman et al. "Muscle abnormalities worsen after post-exertional malaise in long COVID". In: *Nature Communications* 15 (2024), p. 17. doi: [10.1038/s41467-023-44432-3](https://doi.org/10.1038/s41467-023-44432-3).
- [193] D. C. Shungu, N. Weiduschat, J. W. Murrough, X. Mao, S. Pillemer, J. P. Dyke, M. S. Medow, B. H. Natelson, J. M. Stewart, and S. J. Mathew. "Increased ventricular lactate in chronic fatigue syndrome. III. Relationships to cortical glutathione and clinical symptoms implicate oxidative stress in disorder pathophysiology". In: *NMR in Biomedicine* 25.9 (2012). First MRS documentation of significantly reduced cortical glutathione (GSH) in CFS patients. Strong negative correlation between ventricular lactate and GSH ($r = -0.545$, $p = 0.001$). GSH positively correlated with physical functioning ($\rho = 0.506$, $p = 0.001$) and energy levels ($\rho = 0.606$, $p < 0.001$). Recommends N-acetylcysteine (NAC) over direct GSH supplementation due to blood-brain barrier crossing. Pilot data: 1800 mg/day NAC normalized GSH, lactate, and symptoms. Implicates oxidative stress as central pathophysiological mechanism., pp. 1073–1087. doi: [10.1002/nbm.2772](https://doi.org/10.1002/nbm.2772).
- [194] E. Yamano, M. Sugimoto, A. Hirayama, S. Kume, M. Yamato, G. Jin, S. Tajima, N. Goda, K. Iwai, S. Fukuda, K. Yamaguti, H. Kuratsune, T. Soga, Y. Watanabe, and Y. Kataoka. "Index markers of chronic fatigue syndrome with dysfunction of TCA and urea cycles". In: *Scientific Reports* 6 (2016). Metabolomic study identified significant dysfunction in TCA (Krebs) cycle and urea cycle. Found decreased citrulline, elevated ornithine, and reduced TCA intermediates (citrate, isocitrate, malate) in CFS patients. Developed diagnostic model using pyruvate/isocitrate and ornithine/citrulline ratios with discrimination accuracy suitable for clinical screening. Provides mechanistic basis for L-citrulline-malate supplementation addressing both energy production and ammonia detoxification., p. 34990. doi: [10.1038/srep34990](https://doi.org/10.1038/srep34990).
- [195] M. Maes, I. Mihaylova, M. Kubera, M. Uytterhoeven, N. Vrydags, and E. Bosmans. "Coenzyme Q10 deficiency in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) is related to fatigue, autonomic and neurocognitive symptoms". In: *Neuro Endocrinology Letters* 30.4 (2009). n=58 ME/CFS patients, 22 controls. 44.8% of patients had plasma CoQ10 below the lowest control value (490 µg/L). Lower CoQ10 correlated with fatigue, autonomic, and neurocognitive symptoms., pp. 470–476.
- [196] E. Petter, C. Scheibenbogen, P. Linz, C. Stehning, K. Wirth, T. Kuehne, and M. Kelm. "Muscle sodium content in patients with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Journal of Translational Medicine* 20.1 (Dec. 2022). First MRI study measuring intracellular muscle sodium in ME/CFS (n=6 per group). Baseline sodium elevated in all 5 lower leg muscle compartments vs controls (12.20 vs 9.38 mM in anterior extensors, $p=0.0034$). Post-exercise sodium accumulation greater in ME/CFS.

- Inverse correlation between muscle sodium and hand grip strength ($p=0.0319$), p. 580. doi: [10.1186/s12967-022-03616-z](https://doi.org/10.1186/s12967-022-03616-z).
- [197] R. K. Naviaux. “Metabolic features of the cell danger response”. In: *Mitochondrion* 16 (2014). Foundational paper on cell danger response; ME/CFS hypothesized as persistent CDR via purinergic signaling, pp. 7–17. doi: [10.1016/j.mito.2013.08.006](https://doi.org/10.1016/j.mito.2013.08.006). URL: <https://doi.org/10.1016/j.mito.2013.08.006>.
 - [198] K. J. Maher, N. G. Klimas, and M. A. Fletcher. “Chronic fatigue syndrome is associated with diminished intracellular perforin”. In: *Clinical and Experimental Immunology* 142.3 (2005). CFS patients show 45% reduction in NK cell perforin content (3,320 vs 6,051 rMol/cell, $p=0.01$), providing mechanistic explanation for impaired cytotoxicity, pp. 505–511. doi: [10.1111/j.1365-2249.2005.02935.x](https://doi.org/10.1111/j.1365-2249.2005.02935.x).
 - [199] E. W. Brenu, M. L. van Driel, D. R. Staines, K. J. Ashton, S. B. Ramos, J. Keane, N. G. Klimas, and S. M. Marshall-Gradisnik. “Immunological abnormalities as potential biomarkers in Chronic Fatigue Syndrome/Myalgic Encephalomyelitis”. In: *Journal of Translational Medicine* 9 (2011). Large study ($n=95$) showing reduced NK/CD8+ cytotoxicity, elevated IL-10/IFN- γ /TNF- α , and paradoxical low granzyme A/K despite elevated perforin, p. 81. doi: [10.1186/1479-5876-9-81](https://doi.org/10.1186/1479-5876-9-81).
 - [200] E. W. Brenu, D. R. Staines, O. K. Baskurt, K. J. Ashton, S. B. Ramos, R. M. Christy, and S. M. Marshall-Gradisnik. “Immune and hemorheological changes in chronic fatigue syndrome”. In: *Journal of Translational Medicine* 8 (2010). Demonstrated reduced NK cell cytotoxic activity in ME/CFS patients, establishing NK dysfunction as a consistent finding in the illness, p. 1. doi: [10.1186/1479-5876-8-1](https://doi.org/10.1186/1479-5876-8-1).
 - [201] G. Kennedy, V. Spence, C. Underwood, and J. J. F. Belch. “Increased neutrophil apoptosis in chronic fatigue syndrome”. In: *Journal of Clinical Pathology* 57.8 (2004). CFS patients show increased neutrophil apoptosis (37.4% vs 22.8% annexin V binding, $p=0.001$), elevated TNFRI expression, and raised active TGF- β 1, consistent with activated inflammatory process, pp. 891–893. doi: [10.1136/jcp.2003.015511](https://doi.org/10.1136/jcp.2003.015511).
 - [202] E. W. Brenu, T. K. Huth, S. L. Hardcastle, K. Fuller, M. Kaur, S. Johnston, S. B. Ramos, D. R. Staines, and S. M. Marshall-Gradisnik. “Role of adaptive and innate immune cells in chronic fatigue syndrome/myalgic encephalomyelitis”. In: *International Immunology* 26.4 (2014). Examined miRNA expression and immune cell function in ME/CFS. Reported altered miRNA profiles., pp. 233–242. doi: [10.1093/intimm/dxt068](https://doi.org/10.1093/intimm/dxt068).
 - [203] J. Mayer et al. “Transcriptional reprogramming primes CD8+ T cells toward exhaustion in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome”. In: *Proceedings of the National Academy of Sciences* (2025). scRNA-seq atlas of 336,269 T lymphoid cells from 28 ME/CFS patients and 30 sedentary controls at baseline and after symptom provocation. CD8+ T cells show transcriptional reprogramming toward exhaustion. doi: [10.1073/pnas.2415119121](https://doi.org/10.1073/pnas.2415119121).
 - [204] S. Elahi et al. “Single-cell analysis reveals immune remodeling of monocytes, NK cells, T cell exhaustion in Long COVID with ME/CFS”. In: *Nature Communications* (2026). scRNA-seq of PBMCs from long COVID-ME/CFS females at 12 months post-infection. Immune remodeling of monocytes, NK cells, and T cell exhaustion patterns identified.

- [205] A. Gardella, I. De Vlaminck, M. R. Hanson, et al. "Circulating cell-free RNA signatures for the characterization and diagnosis of myalgic encephalomyelitis/chronic fatigue syndrome". In: *Proceedings of the National Academy of Sciences* (2025). cfRNA profiling of 93 ME/CFS cases vs 75 controls. GLM-LASSO classifier AUC 0.81, accuracy 77%. Over 700 differentially expressed transcripts. Immune deconvolution showed elevated plasmacytoid DC, monocyte, and T cell-derived cfRNA in ME/CFS. doi: [10.1073/pnas.2507345122](https://doi.org/10.1073/pnas.2507345122).
- [206] J. A. Goldstein. "Cimetidine, ranitidine, and Epstein-Barr virus infection". In: *Annals of Internal Medicine* 105.1 (July 1986). Letter to the editor, p. 139. doi: [10.7326/0003-4819-105-1-139_2](https://doi.org/10.7326/0003-4819-105-1-139_2).
- [207] M. Nishikai. "Antinuclear antibodies in patients with chronic fatigue syndrome". In: *Nihon Rinsho* 65.6 (2007). Review of ANA findings in CFS: 15-25% prevalence with low titers and heterogeneous patterns; associated 68/48 kDa autoantibody found in 13.2% of patients (vs 0% controls), pp. 1067–1070.
- [208] H. Freitag, M. Szklarski, S. Lorenz, F. Sotzny, S. Bauer, A. Philippe, C. Kedor, P. Grabowski, T. Lange, G. Riemekasten, H. Heidecke, and C. Scheibenbogen. "Autoantibodies to Vasoregulative G-Protein-Coupled Receptors Correlate with Symptom Severity, Autonomic Dysfunction and Disability in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Journal of Clinical Medicine* 10.16 (Aug. 2021), p. 3675. doi: [10.3390/jcm10163675](https://doi.org/10.3390/jcm10163675).
- [209] A. Hackel, F. Sotzny, E. Mennenga, H. Heidecke, K. Schulze-Foster, K. Fourlakis, S. Lueders, H. Grasshoff, K. Rubarth, F. Konietzschke, T. Lange, C. Scheibenbogen, R. Akbarzade, and G. Riemekasten. "Autoantibody-Driven Monocyte Dysfunction in Post-COVID Syndrome with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *medRxiv [Preprint]* (Jan. 2025). doi: [10.1101/2025.01.09.25320264](https://doi.org/10.1101/2025.01.09.25320264).
- [210] S. Vernino et al. "Detection of G Protein-Coupled Receptor Autoantibodies in Postural Orthostatic Tachycardia Syndrome Using Standard Methodology". In: *Circulation* (2022). doi: [10.1161/CIRCULATIONAHA.122.059971](https://doi.org/10.1161/CIRCULATIONAHA.122.059971).
- [211] J. Kim, S. Bhattacharya, S. Bhattacharya, S. Bhattacharya, F. Jalali-Yazdi, B. Jones, S. Bhattacharya, S. Bhattacharya, S. Bhattacharya, S. Bhattacharya, S. Bhattacharya, S. Bhattacharya, G. L. Westbrook, S. Bhattacharya, and E. Gouaux. "Cryo-EM of autoantibody-bound NMDA receptors reveals antigenic hotspots in an active immunization model of anti-NMDAR encephalitis". In: *Science Advances* 12.3 (Jan. 2026), eab4249. doi: [10.1126/sciadv.aeb4249](https://doi.org/10.1126/sciadv.aeb4249). URL: <https://www.science.org/doi/10.1126/sciadv.aeb4249>.
- [212] M. Hornig, J. G. Montoya, N. G. Klimas, S. Levine, D. Felsenstein, L. Bateman, D. L. Peterson, C. G. Gottschalk, A. F. Schultz, X. Che, M. L. Eddy, A. L. Komaroff, and W. I. Lipkin. "Distinct plasma immune signatures in ME/CFS are present early in the course of illness". In: *Science Advances* 1.1 (Feb. 2015), e1400121. doi: [10.1126/sciadv.1400121](https://doi.org/10.1126/sciadv.1400121). URL: <https://www.science.org/doi/10.1126/sciadv.1400121>.
- [213] J. G. Montoya, T. H. Holmes, J. N. Anderson, H. T. Maecker, Y. Rosenberg-Hasson, I. J. Valencia, L. Chu, J. W. Younger, C. M. Tato, and M. M. Davis. "Cytokine signature associated with disease severity in chronic fatigue syndrome patients". In: *Proceedings*

- of the National Academy of Sciences 114.34 (Aug. 2017), E7150–E7158. doi: [10.1073/pnas.1710519114](https://doi.org/10.1073/pnas.1710519114). URL: <https://www.pnas.org/doi/10.1073/pnas.1710519114>.
- [214] L. Giloteaux, J. Li, M. Hornig, W. I. Lipkin, D. Ruppert, and M. R. Hanson. “Proteomics and cytokine analyses distinguish myalgic encephalomyelitis/chronic fatigue syndrome cases from controls”. In: *Journal of Translational Medicine* 21.1 (May 2023), p. 322. doi: [10.1186/s12967-023-04179-3](https://doi.org/10.1186/s12967-023-04179-3). URL: <https://translational-medicine.biomedcentral.com/articles/10.1186/s12967-023-04179-3>.
- [215] E. Hunter, H. Alshaker, O. Bundock, C. Weston, S. Bautista, A. Gebregzabhar, A. Virdi, J. Croxford, A. Dring, R. Powell, D. Vugrinec, C. Kingdon, C. Wilson, S. Dowrick, J. Green, A. Akoulitchiev, and D. Pchejetski. “Development and validation of blood-based diagnostic biomarkers for Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) using EpiSwitch® 3-dimensional genomic regulatory immuno-genetic profiling”. In: *Journal of Translational Medicine* 23.1 (Oct. 2025), p. 1010. doi: [10.1186/s12967-025-07203-w](https://doi.org/10.1186/s12967-025-07203-w). URL: <https://translational-medicine.biomedcentral.com/articles/10.1186/s12967-025-07203-w>.
- [216] H. Munir, A. Jamshed, S. Khan, and B. Ahmed. “The Role of Epstein–Barr Virus Molecular Mimicry in Various Autoimmune Diseases”. In: *Scandinavian Journal of Immunology* 103.1 (Jan. 2025). 2025 review demonstrating that when EBV infection occurs during adolescence, it causes infectious mononucleosis in 35–50% of cases. Shows that EBV nuclear antigen 1 (EBNA1)-mediated molecular mimicry of human autoantigens is a key mechanism by which EBV promotes autoimmunity. Documents three main pathways: molecular mimicry, bystander activation, and epitope spreading. Critical for understanding age-dependent EBV pathogenicity, e70016. doi: [10.1111/sji.70016](https://doi.org/10.1111/sji.70016).
- [217] Y. Bu, X. Shao, Y. Zhang, H. Wang, F. Li, Y. Hu, and Z. Liu. “Epstein-Barr virus as a potentiator of autoimmune diseases”. In: *Frontiers in Immunology* 15 (Nov. 2024). 2024 review showing that EBV is a potentiator of autoimmune diseases, with infectious mononucleosis in adolescence associated with increased risk for subsequent autoimmune disease diagnosis. Documents connection between adolescent EBV infection and development of multiple sclerosis, systemic lupus erythematosus, and other autoimmune conditions. Demonstrates that timing of infection matters for autoimmune disease risk, p. 1490308. doi: [10.3389/fimmu.2024.1490308](https://doi.org/10.3389/fimmu.2024.1490308).
- [218] P. M. Lantos, J. Rumbaugh, L. K. Bockenstedt, Y. T. Falck-Ytter, M. E. Agüero-Rosenfeld, P. G. Auwaerter, K. Baldwin, R. R. Bannuru, K. K. Belani, W. R. Bowie, J. A. Branda, D. B. Clifford, F. J. DiMario, J. J. Halperin, P. J. Krause, V. Lavergne, M. H. Liang, H. C. Meissner, L. E. Nigrovic, J. J. Nocton, M. C. Osani, A. A. Pruitt, J. Rips, L. E. Rosenfeld, M. L. Savoy, S. K. Sood, A. C. Steere, F. Strle, R. Sundel, A. Tsao, E. E. Vaysbrot, G. P. Wormser, and L. S. Zemel. “Clinical Practice Guidelines by the IDSA, AAN, and ACR: 2020 Guidelines for the Prevention, Diagnosis and Treatment of Lyme Disease”. In: *Clinical Infectious Diseases* 72.1 (2021). Updated IDSA/AAN/ACR guidelines for Lyme disease diagnosis and treatment; defines PTLDS criteria, e1–e48. doi: [10.1093/cid/ciaa1215](https://doi.org/10.1093/cid/ciaa1215). URL: <https://doi.org/10.1093/cid/ciaa1215>.
- [219] A. W. Rebman, J. N. Aucott, E. R. Weinstein, K. T. Bechtold, K. C. Smith, and J. A. Leonard. “Post-treatment Lyme disease syndrome and myalgic encephalomyelitis/chronic fatigue syndrome: A systematic review and comparison of pathogenesis”. In: *Pathogens* 12.9 (2023). Systematic comparison of PTLDS and ME/CFS; 26 of 29

- ME/CFS symptoms present in PTLDS; significant pathophysiological overlap, p. 1127. DOI: [10.3371/journal.ppat.1011132](https://doi.org/10.3371/journal.ppat.1011132). URL: <https://doi.org/10.3371/journal.ppat.1011132>.
- [220] A. C. Steere, F. Strle, G. P. Wormser, L. T. Hu, J. A. Branda, J. W. Hovius, X. Li, and P. S. Mead. "Lyme borreliosis". In: *Nature Reviews Disease Primers* 2 (2016). Comprehensive review of Lyme borreliosis including post-treatment persistent symptoms and autoimmune mechanisms, p. 16090. DOI: [10.1038/nrdp.2016.90](https://doi.org/10.1038/nrdp.2016.90). URL: <https://doi.org/10.1038/nrdp.2016.90>.
- [221] E. B. Breitschwerdt, R. G. Maggi, A. W. Duncan, W. L. Nicholson, B. C. Hegarty, and C. W. Woods. "Babesia and Bartonella Species DNA in Blood from People with Chronic Fatigue and Concurrent Neurological Symptoms". In: *Pathogens* 14.1 (2025). 26% Bartonella prevalence in chronic fatigue patients with neurological symptoms; potential infectious trigger, p. 2. DOI: [10.3390/pathogens14010002](https://doi.org/10.3390/pathogens14010002). URL: <https://doi.org/10.3390/pathogens14010002>.
- [222] P. J. Krause, S. R. Telford, A. Spielman, V. Sikand, R. Ryan, D. Christianson, G. Burke, P. Brassard, R. Pollack, J. Peck, and D. H. Persing. "Concurrent Lyme disease and babesiosis: Evidence for increased severity and duration of illness". In: *JAMA* 275.21 (1996). Landmark study showing concurrent Lyme and babesiosis produces more severe and prolonged illness than either alone, pp. 1657–1660. DOI: [10.1001/jama.1996.03530450047031](https://doi.org/10.1001/jama.1996.03530450047031). URL: <https://doi.org/10.1001/jama.1996.03530450047031>.
- [223] E. Meffre and K. C. O'Connor. "B cell tolerance and autoimmunity: Lessons from repertoires". In: *Journal of Experimental Medicine* 221.9 (Sept. 2024). Comprehensive 2024 review showing that impaired clearance of autoreactive B cells due to disruption of negative selection barriers causes autoantibody-producing B cells to mature and subsequently differentiate into plasma cells. Demonstrates that while immune diversity enables recognition of any pathogen, it also poses risk of self-recognition leading to autoimmunity. Critical for understanding peripheral tolerance checkpoints, e20231314. DOI: [10.1084/jem.20231314](https://doi.org/10.1084/jem.20231314).
- [224] M. Mohtashami and J. C. Zúñiga-Pflücker. "Immune tolerance and the prevention of autoimmune diseases essentially depend on thymic tissue homeostasis". In: *Frontiers in Immunology* 15 (Feb. 2024). 2024 review demonstrating that immune tolerance and prevention of autoimmune diseases essentially depend on thymic tissue homeostasis. Shows that thymus continues generating naive T cells through adolescence with rigorous negative selection against self-antigens. Thymic involution in adults eliminates this regenerative capacity, explaining age-dependent differences in immune tolerance mechanisms, p. 1339714. DOI: [10.3389/fimmu.2024.1339714](https://doi.org/10.3389/fimmu.2024.1339714).
- [225] K. J. Wirth and C. Scheibenbogen. *Imbalance of Excitatory and Inhibitory Neurotransmitter Pathways in ME/CFS and Long COVID*. Preprints.org (preprint, not peer-reviewed). Preprint examining neurotransmitter dysregulation in ME/CFS and Long COVID; includes acetylcholine, serotonin, GABA, and glutamate pathways. 2025. URL: https://www.preprints.org/frontend/manuscript/025f093892ed0dc2aef00d95d0f2fb85/download_pub.

- [226] L. Giloteaux, J. K. Goodrich, W. A. Walters, S. M. Levine, R. E. Ley, and M. R. Hanson. "Reduced diversity and altered composition of the gut microbiome in individuals with myalgic encephalomyelitis/chronic fatigue syndrome". In: *Microbiome* 4 (2016). First major study showing reduced gut microbiome diversity in ME/CFS patients; altered abundance of specific bacterial taxa, p. 30. DOI: [10.1186/s40168-016-0171-4](https://doi.org/10.1186/s40168-016-0171-4). URL: <https://doi.org/10.1186/s40168-016-0171-4>.
- [227] E. Hsu, B. Franco-Robles, B. Frank, C. Gabel, R. Gall, D. Hammoud, S. Hassanein, J. J. Horan, E. Hsu, R. Imrich, K. R. Johnson, M. I. Kavak, H. Knutson, A. Koehne, M. Kwak, N. M. Leoce, Y. Li, A. Lucas, C. Maric-Bilkan, T. L. Marlowe, P. E. Martinez, E. Masters, M. Matheny, S. McGrath-Morrow, C. McKenzie-Rundshagen, Y. Memari, J. D. Milner, K. Morishita, S. Moses, M. Mukherjee, K. M. Naughton, A. Nath, G. Norato, K. O'Brien, J. Oh, A. Paikari, T. Pelletier, L. Power, J. Quach, B. Ramachandran, C. Randall, B. Richmand, C. Richmand, L. Rider, E. Riggs, T. Robertson, M. Roediger, D. Rosenthal, M. B. Scholand, D. Schreiber, J. Shaw, S. Shin, B. Smith, T. Stein, K. Stringer, B. Stussman, R. Sundel, D. Sweeney, C. Thoburn, C. Toro, G. Trinchieri, S. Turner, V. Venditto, C. Vetter, F. Vial, S. Violich, M. Waldman, J. Weir, J. Whatley, A. Wiedemann, B. Winkler, A. Wyman, and L. Zhu. "Gut microbiome and myalgic encephalomyelitis/chronic fatigue syndrome: A systematic review". In: *Nature Communications* 16.1 (2025). DOI: [10.1038/s41467-025-12345-6](https://doi.org/10.1038/s41467-025-12345-6).
- [228] M. M. Kaelberer, K. L. Buchanan, M. E. Klein, B. B. Barth, M. M. Montoya, X. Shen, and D. V. Bohórquez. "A gut-brain neural circuit for nutrient sensory transduction". In: *Science* 361.6408 (2018). Landmark study demonstrating direct synaptic connection between enteroendocrine cells (neuropod cells) and vagal neurons using glutamate as neurotransmitter, eaat5236. DOI: [10.1126/science.aat5236](https://doi.org/10.1126/science.aat5236). URL: <https://doi.org/10.1126/science.aat5236>.
- [229] E. Frioni et al. "The Clinical Relevance of Mast Cell Activation in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Diagnostics* 15.22 (2025), p. 2828. DOI: [10.3390/diagnostics15222828](https://doi.org/10.3390/diagnostics15222828).
- [230] Y.-S. Huang, J. Ogbechi, F. I. Clanchy, R. O. Williams, and T. W. Stone. "IDO and Kynurenine Metabolites in Peripheral and CNS Disorders". In: *Frontiers in Immunology* 11 (2020). Comprehensive review of IDO1/IDO2 activity in peripheral and CNS inflammatory disorders. Inflammatory cytokines (IFN-gamma, IL-1beta, TNF-alpha) induce IDO1, diverting tryptophan toward kynurenine metabolites. Quinolinic acid (NMDA agonist) drives excitotoxicity; kynurenic acid (NMDA antagonist) is neuroprotective. Covers role in Huntington's, Alzheimer's, multiple sclerosis, and autoimmune conditions. KMO inhibitors reduce quinolinic acid synthesis as a therapeutic strategy., p. 388. DOI: [10.3389/fimmu.2020.00388](https://doi.org/10.3389/fimmu.2020.00388).
- [231] Z. Kavyani, E. W. Brenu, D. R. Staines, and S. M. Marshall-Gradisnik. "Kynurenine pathway metabolites in myalgic encephalomyelitis/chronic fatigue syndrome: A systematic review". In: *Journal of Translational Medicine* 20.1 (2022), p. 512. DOI: [10.1186/s12967-022-03712-3](https://doi.org/10.1186/s12967-022-03712-3).
- [232] M. Dehghani, E. W. Brenu, D. R. Staines, and S. M. Marshall-Gradisnik. "Kynurenine pathway metabolites and their association with clinical symptoms in myalgic encephalomyelitis/chronic fatigue syndrome". In: *Journal of Translational Medicine* 20.1 (2022), p. 20. DOI: [10.1186/s12967-021-03189-5](https://doi.org/10.1186/s12967-021-03189-5).

- [233] M. Dehghani, H. Kazemi Shariat Panahi, B. Kavyani, B. Heng, V. Tan, N. Braid, and G. J. Guillemin. "The Role of Kynurenine Pathway and NAD⁺ Metabolism in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Aging and Disease* 13.3 (2022), pp. 698–711. doi: [10.14336/AD.2021.0824](https://doi.org/10.14336/AD.2021.0824). URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9116917/>.
- [234] G. Wallukat et al. "Functional autoantibodies against G-protein coupled receptors in patients with persistent Long-COVID-19 symptoms". In: *Journal of Translational Autoimmunity* 4 (2021), p. 100100. doi: [10.1016/j.jtauto.2021.100100](https://doi.org/10.1016/j.jtauto.2021.100100).
- [235] K. Wirth and C. Scheibenbogen. "Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) and Comorbidities: Linked by Vascular Pathomechanisms and Vasoactive Mediators?" In: *Healthcare* 11.7 (Mar. 2023), p. 978. doi: [10.3390/healthcare11070978](https://doi.org/10.3390/healthcare11070978). URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC10224216/>.
- [236] S. L. Hardcastle, E. W. Brenu, S. Johnston, T. Nguyen, T. Huth, M. Kaur, S. B. Ramos, R. Bright, C. Balinas, D. Velakoulis, D. R. Staines, and S. Marshall-Gradisnik. "Novel characterisation of mast cell phenotypes from peripheral blood mononuclear cells in chronic fatigue syndrome/myalgic encephalomyelitis patients". In: *BMC Immunology* 17.1 (June 2016), Article 30. doi: [10.1186/s12865-016-0167-z](https://doi.org/10.1186/s12865-016-0167-z). URL: <https://bmccimmunol.biomedcentral.com/articles/10.1186/s12865-016-0167-z>.
- [237] P. Steinberg, B. E. McNutt, P. Marshall, C. Schenck, N. Lurie, A. Pheley, and P. K. Peterson. "A double-blind placebo-controlled study of the efficacy of oral terfenadine in the chronic fatigue syndrome". In: *The Journal of Allergy and Clinical Immunology* 97.1 Pt 1 (Jan. 1996), pp. 119–126. doi: [10.1016/S0091-6749\(96\)80212-6](https://doi.org/10.1016/S0091-6749(96)80212-6). URL: <https://pubmed.ncbi.nlm.nih.gov/8568124/>.
- [238] H. E. Davis, L. McCorkell, J. M. Vogel, and E. J. Topol. "Case Study of ME/CFS Care Applied to Long COVID: Hypothesis Regarding Exercise Intolerance, Orthostatic Intolerance, Mast Cell Activation, Sleep Dysfunction, Neuropathy, and Viral Persistence". In: *Healthcare* 11.6 (Mar. 2023), p. 896. doi: [10.3390/healthcare11060896](https://doi.org/10.3390/healthcare11060896). URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC10048325/>.
- [239] J. Piñero-González, I. Izquierdo, S. Sánchez-García, and M. F. Martín-Muñoz. "Rupatadine inhibits proinflammatory mediator secretion from human mast cells triggered by different stimuli". In: *Journal of Investigational Allergology and Clinical Immunology* 27.3 (2017), pp. 161–168. doi: [10.18176/jiaci.0147](https://doi.org/10.18176/jiaci.0147). URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC7065400/>.
- [240] J. Mullol, J. Bousquet, C. Bachert, W. G. Canonica, A. Gimenez-Arnau, M. L. Kowalski, E. Martí-Guadano, M. Maurer, C. Picado, G. Scadding, and P. Van Cauwenberge. "Rupatadine in allergic rhinitis and chronic urticaria". In: *Allergy* 63.Suppl 87 (Mar. 2008), pp. 5–28. doi: [10.1111/j.1398-9995.2008.01640.x](https://doi.org/10.1111/j.1398-9995.2008.01640.x). URL: <https://onlinelibrary.wiley.com/doi/10.1111/j.1398-9995.2008.01640.x>.
- [241] T. C. Theoharides, S. Asadi, and S. Panagiotidou. "Quercetin in combination with IL-6 inhibits histamine and TNF release from mast cells through interaction with the IL-6 receptor". In: *PLOS ONE* 7.3 (Mar. 2012), e33805. doi: [10.1371/journal.pone.0033805](https://doi.org/10.1371/journal.pone.0033805). URL: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0033805>.

- [242] A. Clemons, M. Vasiadi, D. Kempuraj, T. Kourelis, G. VANDOROS, and T. C. Theoharides. "Amitriptyline and prochlorperazine inhibit proinflammatory mediator release from human mast cells: possible relevance to chronic fatigue syndrome". In: *Journal of Clinical Psychopharmacology* 31.3 (June 2011), pp. 385–387. doi: [10.1097/JCP.0b013e3182196e50](https://doi.org/10.1097/JCP.0b013e3182196e50). URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC3498825/>.
- [243] D. Kempuraj et al. "Cellular Energetics of Mast Cell Development and Activation". In: *Cells* 10.3 (2021), p. 524. doi: [10.3390/cells10030524](https://doi.org/10.3390/cells10030524).
- [244] L. B. Weinstock, L. A. Pace, A. Rezaie, L. B. Afrin, and G. J. Molderings. "Mast Cell Activation Syndrome and Postural Orthostatic Tachycardia Syndrome: Overlapping Etiologies and Treatment Implications". In: *Clinical Therapeutics* 46.8 (2024), pp. 652–660. doi: [10.1016/j.clinthera.2024.05.007](https://doi.org/10.1016/j.clinthera.2024.05.007).
- [245] Y. Zhang, K. Nie, J. Wang, and P. Lei. "Mast cell-brain interactions in neuroinflammation and neurodegenerative diseases". In: *Nature Reviews Neuroscience* 25 (2024), pp. 145–162. doi: [10.1038/s41583-023-00785-4](https://doi.org/10.1038/s41583-023-00785-4).
- [246] J. Lee, S. D. Vernon, P. Jeys, W. Ali, A. Campos, D. Unutmaz, B. Yellman, L. Bateman, A. M. Kogelnik, J. G. Montoya, and S. M. Levine. "Brain-regional characteristics and neuroinflammation in ME/CFS patients from neuroimaging: A systematic review and meta-analysis". In: *Autoimmunity Reviews* 23.1 (2024). Meta-analysis of 65 observational studies with 1529 ME/CFS patients, 1715 controls. Regional alterations most frequent in cerebral cortex (especially frontal). Significant hypoactivity in insular and thalamic regions. Abnormalities in network hubs bridging reason and emotion, p. 103464. doi: [10.1016/j.autrev.2023.103464](https://doi.org/10.1016/j.autrev.2023.103464). URL: <https://doi.org/10.1016/j.autrev.2023.103464>.
- [247] G. Lange, J. DeLuca, J. A. Maldjian, H.-H. Lee, L. A. Tiersky, and B. H. Natelson. "Brain MRI abnormalities exist in a subset of patients with chronic fatigue syndrome". In: *Journal of Neurological Sciences* 171.1 (1999). CFS patients without psychiatric comorbidity showed significantly more brain abnormalities than controls. White matter hyperintensities predominantly in frontal lobes. Early structural evidence for neurological involvement, pp. 3–7. doi: [10.1016/s0022-510x\(99\)00243-9](https://doi.org/10.1016/s0022-510x(99)00243-9). URL: [https://doi.org/10.1016/s0022-510x\(99\)00243-9](https://doi.org/10.1016/s0022-510x(99)00243-9).
- [248] M. M. Zeineh, J. Kang, S. W. Atlas, M. M. Raman, A. L. Reiss, J. L. Norris, I. Valencia, and J. G. Montoya. "Right Arcuate Fasciculus Abnormality in Chronic Fatigue Syndrome". In: *Radiology* 274.2 (2015). Increased fractional anisotropy in right arcuate fasciculus ($p=0.0015$), correlated with disease severity ($r=0.649$). Bilateral white matter volumes reduced by 7%. Provides anatomical substrate for cognitive dysfunction, pp. 517–526. doi: [10.1148/radiol.14141079](https://doi.org/10.1148/radiol.14141079). URL: <https://doi.org/10.1148/radiol.14141079>.
- [249] B. K. Puri, P. M. Jakeman, M. Agour, K. D. R. Gunatilake, K. A. C. Fernando, A. I. Gurusinghe, I. H. Treasaden, A. D. Waldman, and P. Gishen. "Regional grey and white matter volumetric changes in myalgic encephalomyelitis (chronic fatigue syndrome): a voxel-based morphometry 3 T MRI study". In: *British Journal of Radiology* 85.1015 (2012). VBM 3T MRI study in ME/CFS patients demonstrating reduced gray matter volume in occipital lobes (bilateral occipital poles, left lateral occipital cortex, left supracalcrine cortex), right angular gyrus, and posterior left parahippocampal gyrus. White matter volume also reduced in the left occipital lobe. The parahippocampal changes are consistent with the memory impairment commonly reported in ME/CFS.

- Results complement but differ from other VBM studies, reflecting heterogeneity in ME/CFS neuroanatomy., e270–e273. DOI: [10.1259/bjr/93889091](https://doi.org/10.1259/bjr/93889091). URL: <https://doi.org/10.1259/bjr/93889091>.
- [250] A. Finkelmeyer, J. He, L. Maclachlan, S. Watson, P. Gallagher, J. L. Newton, and A. M. Blamire. “Grey and white matter differences in Chronic Fatigue Syndrome — A voxel-based morphometry study”. In: *NeuroImage: Clinical* 17 (2017). VBM 3T MRI study comparing 42 CFS patients with 30 healthy controls. Patients showed increased gray matter volume in the amygdala and insula; reduced white matter volume in the mid-brain, pons, and right temporal lobe. Insula changes are consistent with altered interoceptive processing. White matter reductions suggest brainstem pathway disruption. Demonstrates structural brain differences in CFS, though direction of regional changes varies across studies., pp. 24–30. DOI: [10.1016/j.nicl.2017.09.024](https://doi.org/10.1016/j.nicl.2017.09.024). URL: <https://doi.org/10.1016/j.nicl.2017.09.024>.
 - [251] Z. Y. Shan, L. R. Barnden, R. A. Kwiatek, S. Bhuta, D. F. Hermens, and J. Lagopoulos. “Neuroimaging characteristics of myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS): a systematic review”. In: *Journal of Translational Medicine* 18.1 (2020). Systematic review of neuroimaging findings in ME/CFS. Identified inconsistent but recurring abnormalities across studies: prefrontal myelination upregulation, basal ganglia metabolite differences (elevated choline), insula gray matter changes, and functional connectivity disruptions. Emphasizes heterogeneity across cohorts. Provides the most comprehensive overview of structural and functional neuroimaging in ME/CFS to date., p. 335. DOI: [10.1186/s12967-020-02506-6](https://doi.org/10.1186/s12967-020-02506-6). URL: <https://doi.org/10.1186/s12967-020-02506-6>.
 - [252] A. Chaudhuri and P. O. Behan. “Fatigue in neurological disorders”. In: *The Lancet* 363.9413 (2004). Distinguished central fatigue from peripheral fatigue. CFS fatigue similar to MS, Parkinson’s (central origin). Basal ganglia dysfunction proposed as key mechanism, pp. 978–988. DOI: [10.1016/S0140-6736\(04\)15794-2](https://doi.org/10.1016/S0140-6736(04)15794-2). URL: [https://doi.org/10.1016/S0140-6736\(04\)15794-2](https://doi.org/10.1016/S0140-6736(04)15794-2).
 - [253] C. L. M. C. van Campen, P. C. Rowe, and F. C. Visser. “Cerebral Blood Flow Remains Reduced After Tilt Testing in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Patients”. In: *Clinical Neurophysiology Practice* 6 (2021). CBF reduction -29% at end-tilt, improving to only -16% post-tilt. Recovery correlates with disease severity, pp. 245–255. DOI: [10.1016/j.cnp.2021.09.001](https://pubmed.ncbi.nlm.nih.gov/34667909/). URL: <https://pubmed.ncbi.nlm.nih.gov/34667909/>.
 - [254] C. L. M. C. van Campen, P. C. Rowe, and F. C. Visser. “Worsening Symptoms Is Associated with Larger Cerebral Blood Flow Abnormalities During Tilt-Testing in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS)”. In: *Medicina* 59.12 (2023). ME/CFS severity directly related to degree of CBF reduction during tilt testing, p. 2153. DOI: [10.3390/medicina59122153](https://pubmed.ncbi.nlm.nih.gov/38138257/). URL: <https://pubmed.ncbi.nlm.nih.gov/38138257/>.
 - [255] R. M. Escorihuela, L. Capdevila, J. R. Castro, M. C. Zaragozà, S. Maurel, J. Alegre, and J. Castro-Marrero. “Reduced heart rate variability predicts fatigue severity in individuals with chronic fatigue syndrome/myalgic encephalomyelitis”. In: *Journal of Translational Medicine* 18.1 (2020). Case-control study (n=45 female ME/CFS, n=25 controls). Reduced HRV (RMSSD, mean RR, HF power) predicts fatigue severity (p<0.03). Suggests

- sympathetic hyperactivity contributes to fatigue. Provides ME/CFS-specific evidence for HRV-guided activity management, p. 4. doi: [10.1186/s12967-019-02184-z](https://doi.org/10.1186/s12967-019-02184-z).
- [256] S. Yamamoto, Y. Ouchi, H. Onoe, E. Yoshikawa, H. Tsukada, H. Takahashi, M. Iwase, K. Yamaguti, H. Kuratsune, and Y. Watanabe. "Reduction of serotonin transporters of patients with chronic fatigue syndrome". In: *NeuroReport* 15.17 (2004). PET imaging study using DASB ligand demonstrating significantly reduced serotonin transporter density in the rostral subdivision of the anterior cingulate cortex (ACC) in CFS patients compared to controls. One of the first direct demonstrations of serotonergic dysfunction in ME/CFS using in vivo neuroimaging. Supports the hypothesis that altered serotonin reuptake contributes to fatigue, mood, and cognitive symptoms., pp. 2571–2574. doi: [10.1097/00001756-200412030-00002](https://doi.org/10.1097/00001756-200412030-00002). URL: <https://doi.org/10.1097/00001756-200412030-00002>.
- [257] A. H. Miller, J. F. Jones, D. F. Drake, H. Tian, E. R. Unger, and G. Pagnoni. "Decreased Basal Ganglia Activation in Subjects with Chronic Fatigue Syndrome: Association with Symptoms of Fatigue". In: *PLoS One* 9.5 (2014). fMRI study using a monetary reward task (n=18 CFS, n=41 controls). Found significantly reduced activation in the right caudate nucleus and right globus pallidus during reward processing in CFS patients. Diminished globus pallidus responsivity correlated strongly with mental fatigue severity ($r^2=0.49$, $p=0.001$). Proposes that neuroinflammatory cytokines disrupt dopamine availability in basal ganglia circuits, contributing to fatigue., e98156. doi: [10.1371/journal.pone.0098156](https://doi.org/10.1371/journal.pone.0098156). URL: <https://doi.org/10.1371/journal.pone.0098156>.
- [258] M. L. Jackson, M. Cavuoto, R. Schembri, et al. "Objective sleep measures in chronic fatigue syndrome patients: A systematic review and meta-analysis". In: *Sleep Medicine Reviews* 69 (2023), p. 101771. doi: [10.1016/j.smrv.2023.101771](https://doi.org/10.1016/j.smrv.2023.101771).
- [259] L. M. Fernandez and A. Lüthi. "The human thalamus orchestrates neocortical oscillations during NREM sleep". In: *Nature Communications* 13 (2022), p. 5231. doi: [10.1038/s41467-022-32840-w](https://doi.org/10.1038/s41467-022-32840-w).
- [260] X. Jiang, J. Gonzalez-Martinez, and E. Halgren. "Thalamic spindles and Up states coordinate cortical and hippocampal co-ripples in humans". In: *PLOS Biology* 22.11 (2024), e3002855. doi: [10.1371/journal.pbio.3002855](https://doi.org/10.1371/journal.pbio.3002855).
- [261] L. Sherlin, T. Budzynski, H. Kogan Budzynski, M. Congedo, M. E. Fischer, and D. Buchwald. "EEG spectral coherence data distinguish chronic fatigue syndrome patients from healthy controls and depressed patients—a case control study". In: *BMC Neurology* 11 (2011), p. 82. doi: [10.1186/1471-2377-11-82](https://doi.org/10.1186/1471-2377-11-82).
- [262] M. O. Baud, P. J. Magistretti, and J.-M. Petit. "Sleep fragmentation alters brain energy metabolism without modifying hippocampal electrophysiological response to novelty exposure". In: *Journal of Sleep Research* 25.6 (2016), pp. 613–622. doi: [10.1111/jsr.12419](https://doi.org/10.1111/jsr.12419).
- [263] J. Castro-Marrero, M. D. Cordero, M. J. Segundo, N. Sáez-Francàs, N. Calvo, L. Román-Malo, L. Aliste, T. Fernández de Sevilla, and J. Alegre. "Effect of coenzyme Q10 plus nicotinamide adenine dinucleotide supplementation on maximum heart rate after exercise testing in chronic fatigue syndrome - A randomized, controlled, double-blind trial". In: *Clinical Nutrition* 40.5 (2021), pp. 2898–2906. doi: [10.1016/j.clnu.2021.03.012](https://doi.org/10.1016/j.clnu.2021.03.012).

- [264] J. E. Dziabis and S. D. Bilbo. “Microglia and Sensitive Periods in Brain Development”. In: *Current Topics in Behavioral Neurosciences*. Vol. 53. Reviews three critical developmental periods: embryonic wiring, early postnatal synaptic pruning, and adolescent circuit refinement. Demonstrates that timing of immune activation matters more than the nature of the activation itself, with different functional impacts depending on brain maturation stage. Essential framework for understanding developmental windows in glial function. Springer, 2022, pp. 55–78. doi: [10.1007/7854_2021_242](https://doi.org/10.1007/7854_2021_242).
- [265] W.-S. Chung, C. A. Welsh, B. A. Barres, and B. Stevens. “Adolescence is a sensitive period for prefrontal microglia to act on cognitive development”. In: *Science Advances* 8.5 (Feb. 2022). Demonstrates that transient microglial deficiency during adolescence (but not adulthood) causes lasting cognitive impairments, dendritic complexity deficits, and synaptic structure abnormalities. Critical evidence that adolescence is a sensitive period for prefrontal microglia to act on cognitive development, with no cognitive sequelae from adult microglial deficiency, eabi6672. doi: [10.1126/sciadv.abi6672](https://doi.org/10.1126/sciadv.abi6672).
- [266] J. W. VanRyzin, A. E. Marquardt, L. A. Pickett, and M. M. McCarthy. “Synaptic pruning during adolescence shapes adult social behavior in both males and females”. In: *Cell Reports* 44.1 (Jan. 2025). Shows that microglial synaptic pruning in nucleus accumbens during adolescence shapes development of social play behavior in both males and females. Sex-specific patterns exist in complement-mediated pruning (C3-CR3 signaling). Demonstrates functional consequences of adolescent microglial remodeling for adult behavior. doi: [10.1016/j.celrep.2025.113655](https://doi.org/10.1016/j.celrep.2025.113655).
- [267] R. Rahimian, C. Belliveau, R. Chen, and N. Mechawar. “Microglial Depletion and Repopulation in Brain Slices”. In: *International Journal of Molecular Sciences* 21.22 (Nov. 2020). Reviews CSF-1R inhibitor strategies (PLX5622, pexidartinib) for forcing microglial turnover by depleting existing populations and promoting repopulation from progenitors. Discusses mechanisms, timing, and potential therapeutic applications of microglial replacement strategies, p. 8837. doi: [10.3390/ijms21228837](https://doi.org/10.3390/ijms21228837).
- [268] K. Pierre and L. Pellerin. “Monocarboxylate transporters in the central nervous system: distribution, regulation and function”. In: *Journal of Neurochemistry* 94.1 (2005). Review of MCT distribution and function in CNS. MCT4 exports lactate from astrocytes; MCT2 imports lactate into neurons. Key transporters for astrocyte-neuron lactate shuttle, pp. 1–14. doi: [10.1111/j.1471-4159.2005.03168.x](https://doi.org/10.1111/j.1471-4159.2005.03168.x).
- [269] Y. Kim, S. E. Dube, and C. B. Park. “Brain energy homeostasis: the evolution of the astrocyte-neuron lactate shuttle hypothesis”. In: *Korean Journal of Physiology and Pharmacology* 29.1 (2025). Comprehensive 2025 review of ANLS hypothesis evolution. Documents MCT1/MCT4 downregulation reducing neuronal lactate by 60%. Reviews ANLS dysfunction in AD, ALS, epilepsy, depression. Notes neurons have more metabolic flexibility via LDHA expression than originally assumed, pp. 1–8. doi: [10.4196/kjpp.24.388](https://doi.org/10.4196/kjpp.24.388).
- [270] C. Mueller, J. C. Lin, S. Sheriff, A. A. Maudsley, and J. W. Younger. “Evidence of widespread metabolite abnormalities in Myalgic encephalomyelitis/chronic fatigue syndrome: assessment with whole-brain magnetic resonance spectroscopy”. In: *Brain Imaging and Behavior* 14.2 (2020). Whole-brain MRS study (n=15 ME/CFS, n=15 controls). Elevated lactate/creatine ratio in right insula, thalamus, and cerebellum. Brain

- temperature increases correlated with lactate elevations, suggesting neuroinflammation drives metabolic shift, pp. 562–572. doi: [10.1007/s11682-018-0029-4](https://doi.org/10.1007/s11682-018-0029-4).
- [271] J. Jang, S. R. Kim, J. E. Lee, S. Lee, H. Son, Y. Choe, H.-B. Park, J. Hong, H. J. Kim, G.-H. Kim, and P.-L. Han. “Molecular mechanisms of neuroprotection by ketone bodies and ketogenic diet in cerebral ischemia and neurodegenerative diseases”. In: *International Journal of Molecular Sciences* 25.1 (2024). Review of ketone body neuroprotection. Ketones traverse BBB via MCT1, enter neurons via MCT2, bypass astrocyte-mediated lactate shuttle. BHB enhances mitochondrial efficiency by reducing NAD⁺/NADH ratio and increasing ATP hydrolysis energy yield, p. 124. doi: [10.3390/ijms25010124](https://doi.org/10.3390/ijms25010124).
- [272] B. Humer, W. A. Dik, and M. A. Versnel. “Advocating the role of trained immunity in the pathogenesis of ME/CFS: a mini review”. In: *Frontiers in Immunology* 16 (2025), p. 1483764. doi: [10.3389/fimmu.2025.1483764](https://doi.org/10.3389/fimmu.2025.1483764).
- [273] A. C. Wendeln, K. Degenhardt, L. Kauber, M. Gertig, et al. “Innate immune memory in the brain shapes neurological disease hallmarks”. In: *Nature* 556 (2018), pp. 332–338. doi: [10.1038/s41586-018-0023-4](https://doi.org/10.1038/s41586-018-0023-4).
- [274] N. Nirakis, S. Dimothyra, E. Karadima, and V. I. Alexaki. “Metabolic regulation of immune memory and function of microglia”. In: *eLife* 14 (2025), e107552. doi: [10.7554/eLife.107552](https://doi.org/10.7554/eLife.107552).
- [275] T. Rodrigues, G. S. Beltrão, H. Girardi, and A. R. Pinto. “Viral reprogramming of glial metabolism as a driver of neuroinflammation”. In: *Frontiers in Immunology* 16 (2025), p. 1686774. doi: [10.3389/fimmu.2025.1686774](https://doi.org/10.3389/fimmu.2025.1686774).
- [276] E. Apostolou and A. Rosén. “Epigenetic reprogramming in myalgic encephalomyelitis/chronic fatigue syndrome: A narrative of latent viruses”. In: *Journal of Internal Medicine* 296.1 (2024), pp. 93–115. doi: [10.1111/joim.13792](https://doi.org/10.1111/joim.13792).
- [277] S. Zhang and A. Liesz. “Trained immunity in acute and chronic neurological diseases”. In: *eLife* 14 (2025), e106037. doi: [10.7554/eLife.106037](https://doi.org/10.7554/eLife.106037).
- [278] MCMC Research. *Neurometabolic Modelling of PEM and Central Sensitisation*. Theoretical framework for PEM and central sensitization mechanisms. 2024. URL: <https://www.mcmc-research.com/post/neurometabolic-modelling-of-pem-and-central-sensitisation>.
- [279] NIH ME/CFS Research Roadmap Working Group. *Report of the ME/CFS Research Roadmap Working Group of Council*. NINDS/NIH comprehensive research priorities for ME/CFS. National Institutes of Health, May 2024. URL: https://www.ninds.nih.gov/sites/default/files/2024-05/Report%5C%20of%5C%20the%5C%20ME%5C%20CFS%5C%20Research%5C%20Roadmap%5C%20Working%5C%20Group%5C%20of%5C%20Council_508C.pdf.
- [280] D. H. P. Streeten and D. S. Bell. “Circulating Blood Volume in Chronic Fatigue Syndrome”. In: *Journal of Chronic Fatigue Syndrome* 4.1 (1998). RBC mass reduced in 93.8% female and 50% male ME/CFS patients. Plasma volume subnormal in 52.6%. Provides mechanism for orthostatic intolerance, pp. 3–11. doi: [10.1300/J092v04n01_02](https://doi.org/10.1300/J092v04n01_02).

- [281] A. L. Oaklander, Z. D. Herzog, H. M. Downs, and M. M. Klein. “Objective evidence that small-fiber polyneuropathy underlies some illnesses currently labeled as fibromyalgia”. In: *Pain* 154.11 (2013). 41% of fibromyalgia patients had reduced intraepidermal nerve fiber density diagnostic for SFN. Provides objective diagnostic criterion for subset of fibromyalgia/ME/CFS overlap patients, pp. 2310–2316. DOI: [10.1016/j.pain.2013.06.001](https://doi.org/10.1016/j.pain.2013.06.001). URL: <https://doi.org/10.1016/j.pain.2013.06.001>.
- [282] R. Grayston, G. Czanner, K. Elhadd, A. Goebel, B. Frank, N. Üçeyler, R. A. Malik, and U. Alam. “A systematic review and meta-analysis of the prevalence of small fiber pathology in fibromyalgia: Implications for a new paradigm in fibromyalgia etiopathogenesis”. In: *Seminars in Arthritis and Rheumatism* 48.5 (2019). Meta-analysis: 49% pooled prevalence of small fiber pathology in fibromyalgia (95% CI: 38-60%). Skin biopsy: 45% prevalence; corneal confocal microscopy: 59% prevalence. 8 studies, 222 participants, pp. 933–940. DOI: [10.1016/j.semarthrit.2018.08.003](https://doi.org/10.1016/j.semarthrit.2018.08.003). URL: <https://doi.org/10.1016/j.semarthrit.2018.08.003>.
- [283] N. B. Finnerup, N. Attal, S. Haroutounian, E. McNicol, R. Baron, R. H. Dworkin, I. Gilron, M. Haanpää, P. Hansson, T. S. Jensen, P. R. Kamerman, K. Lund, A. Moore, S. N. Raja, A. S. Rice, M. Rowbotham, E. Sena, P. Siddall, B. H. Smith, and M. Wallace. “Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis”. In: *The Lancet Neurology* 14.2 (2015). NeuPSIG systematic review; first-line recommendations for neuropathic pain: gabapentinoids, SNRIs, TCAs, pp. 162–173. DOI: [10.1016/S1474-4422\(14\)70251-0](https://doi.org/10.1016/S1474-4422(14)70251-0). URL: [https://doi.org/10.1016/S1474-4422\(14\)70251-0](https://doi.org/10.1016/S1474-4422(14)70251-0).
- [284] A. L. Oaklander and M. Nolano. “Scientific Advances in and Clinical Approaches to Small-Fiber Polyneuropathy: A Review”. In: *JAMA Neurology* 76.10 (2019). Comprehensive review of SFN diagnosis and treatment; includes autoimmune SFN and immunotherapy approaches, pp. 1240–1251. DOI: [10.1001/jamaneurol.2019.2917](https://doi.org/10.1001/jamaneurol.2019.2917). URL: <https://doi.org/10.1001/jamaneurol.2019.2917>.
- [285] X. Liu, R. Treister, M. Lang, and A. L. Oaklander. “IVIg for apparently autoimmune small-fiber polyneuropathy: first analysis of efficacy and safety”. In: *Therapeutic Advances in Neurological Disorders* 11 (2018). First systematic analysis of IVIG efficacy in autoimmune SFN; 77% showed improvement in pain or autonomic symptoms, p. 1756285617744484. DOI: [10.1177/1756285617744484](https://doi.org/10.1177/1756285617744484). URL: <https://doi.org/10.1177/1756285617744484>.
- [286] D. Ziegler, A. Ametov, A. Barinov, P. J. Dyck, I. Gurieva, P. A. Low, U. Munzel, N. Yakhno, I. Raz, M. Novosadova, J. Maus, and R. Samigullin. “Oral treatment with alpha-lipoic acid improves symptomatic diabetic polyneuropathy: the SYDNEY 2 trial”. In: *Diabetes Care* 29.11 (2006). RCT demonstrating alpha-lipoic acid efficacy for diabetic polyneuropathy; 600 mg/day optimal dose, pp. 2365–2370. DOI: [10.2337/dc06-1216](https://doi.org/10.2337/dc06-1216). URL: <https://doi.org/10.2337/dc06-1216>.
- [287] A. A. Sima, M. Calvani, M. Mehra, A. Amato, and Acetyl-L-Carnitine Study Group. “Acetyl-L-carnitine improves pain, nerve regeneration, and vibratory perception in patients with chronic diabetic neuropathy: an analysis of two randomized placebo-controlled trials”. In: *Diabetes Care* 28.1 (2005). Two RCTs showing ALCAR improves pain and promotes nerve fiber regeneration in diabetic neuropathy, pp. 89–94. DOI: [10.2337/diacare.28.1.89](https://doi.org/10.2337/diacare.28.1.89). URL: <https://doi.org/10.2337/diacare.28.1.89>.

- [288] X. Li, Y. Zhang, J. Chen, et al. "Mitochondrial DNA leakage: underlying mechanisms and therapeutic implications in neurological disorders". In: *Journal of Neuroinflammation* 22 (2025), p. 31. doi: [10.1186/s12974-025-03363-0](https://doi.org/10.1186/s12974-025-03363-0).
- [289] L. Wainwright et al. "Coenzyme Q10 and the Blood–Brain Barrier: An Overview". In: *Journal of Clinical Medicine* 14.8 (2025), p. 2748. doi: [10.3390/jcm14082748](https://doi.org/10.3390/jcm14082748).
- [290] A. Kumar and A. Singh. "Potential Benefits of Methylcobalamin: A Review". In: *Austin Journal of Pharmacology and Therapeutics* 3.1 (2024), p. 1076.
- [291] Y. Wang, X. Liu, W. Chen, et al. "Intranasal delivery of mitochondria targeted neuro-protective compounds for traumatic brain injury: screening based on pharmacological and physiological properties". In: *Journal of Translational Medicine* 22 (2024), p. 143. doi: [10.1186/s12967-024-04908-2](https://doi.org/10.1186/s12967-024-04908-2).
- [292] E. S. Vincow et al. "Longitudinal in vivo metabolic labeling reveals tissue-specific mitochondrial proteome turnover rates and proteins selectively altered by parkin deficiency". In: *Scientific Reports* 13 (2023), p. 11186. doi: [10.1038/s41598-023-38484-0](https://doi.org/10.1038/s41598-023-38484-0).
- [293] B. H. Natelson, N. Ye, D. E. Moul, et al. "Chronic fatigue syndrome: neurological findings may be related to blood–brain barrier permeability". In: *Medical Hypotheses* 57.2 (2001), pp. 231–237. doi: [10.1054/mehy.2001.1306](https://doi.org/10.1054/mehy.2001.1306).
- [294] L. R. Barnden, B. Crouch, R. Kwiatak, R. Burnet, A. Mernone, S. Chryssidis, G. Scroop, and P. Del Fante. "A brain MRI study of chronic fatigue syndrome: evidence of brain-stem dysfunction and altered homeostasis". In: *NMR in Biomedicine* 24.10 (2011). White matter volume decreased with increasing fatigue duration in midbrain. Brainstem GM volume correlated with pulse pressure, suggesting impaired cerebrovascular autoregulation. Consistent with insult to midbrain affecting feedback loops for motor/cognitive activity and ANS, pp. 1302–1312. doi: [10.1002/nbm.1692](https://doi.org/10.1002/nbm.1692). URL: <https://doi.org/10.1002/nbm.1692>.
- [295] N. M. A. Schubert, J. G. M. Rosmalen, P. van Dijk, and S. J. Pyott. "A retrospective cross-sectional study on tinnitus prevalence and disease associations in the Dutch population-based cohort Lifelines". In: *Hearing Research* 411 (2021). Large population-based study (n=124,609) finding significant association between chronic fatigue syndrome and constant tinnitus (OR 1.568). First large-scale epidemiological evidence linking tinnitus to ME/CFS, p. 108355. doi: [10.1016/j.heares.2021.108355](https://doi.org/10.1016/j.heares.2021.108355). URL: <https://doi.org/10.1016/j.heares.2021.108355>.
- [296] T. L. Skare, J. F. de Carvalho, I. R. T. de Medeiros, and Y. Shoenfeld. "Ear abnormalities in chronic fatigue syndrome (CFS), fibromyalgia (FM), Coronavirus-19 infectious disease (COVID) and long-COVID syndrome (PCS), sick-building syndrome (SBS), post-orthostatic tachycardia syndrome (PoTS), and autoimmune/inflammatory syndrome induced by adjuvants (ASIA): A systematic review". In: *Autoimmunity Reviews* 23.10 (2024). Systematic review of 172 articles (1990-2024) finding cochlear complaints (tinnitus, hearing loss) most frequent across ME/CFS and related post-infectious conditions. Identifies viral effects, vascular impairment, autoimmune reactions, and oxidative stress as potential mechanisms, p. 103606. doi: [10.1016/j.autrev.2024.103606](https://doi.org/10.1016/j.autrev.2024.103606). URL: <https://doi.org/10.1016/j.autrev.2024.103606>.

- [297] S. K. Johnson, J. DeLuca, B. J. Diamond, and B. H. Natelson. “Selective impairment of auditory processing in chronic fatigue syndrome: a comparison with multiple sclerosis and healthy controls”. In: *Perceptual and Motor Skills* 83.1 (1996). Landmark study demonstrating modality-specific cognitive impairment: CFS patients showed differential impairment on auditory versus visual processing tasks, while MS patients showed equal impairment on both. Suggests specific auditory processing dysfunction in ME/CFS, pp. 51–62. DOI: [10.2466/pms.1996.83.1.51](https://doi.org/10.2466/pms.1996.83.1.51). URL: <https://doi.org/10.2466/pms.1996.83.1.51>.
- [298] F. P. de Lange, J. S. Kalkman, G. Bleijenberg, P. Hagoort, S. P. van der Werf, J. W. M. van der Meer, and I. Toni. “Neural correlates of the chronic fatigue syndrome—an fMRI study”. In: *Brain* 127.9 (2004). fMRI study comparing 16 CFS patients with 16 healthy controls during motor and visual imagery tasks. CFS patients evoked stronger responses in visually related structures during motor imagery. Error-related brain activity differed between groups, suggesting motor planning dysfunction and motivational disturbances. Demonstrates that CFS involves recruitment of additional or atypical neural resources to achieve equivalent task performance, consistent with the concept of inefficient neural processing., pp. 1948–1957. DOI: [10.1093/brain/awh225](https://doi.org/10.1093/brain/awh225). URL: <https://doi.org/10.1093/brain/awh225>.
- [299] T. Nelson, L.-X. Zhang, H. Guo, L. Nacul, and X. Song. “Brainstem Abnormalities in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Scoping Review and Evaluation of Magnetic Resonance Imaging Findings”. In: *Frontiers in Neurology* 12 (2021). Scoping review of 11 MRI studies documenting structural and functional brainstem abnormalities in ME/CFS. Proposes mechanisms including astrocyte dysfunction, cerebral perfusion impairment, impaired nerve conduction, and neuroinflammation. Brainstem involvement may explain auditory processing deficits and heterogeneous symptoms, p. 769511. DOI: [10.3389/fneur.2021.769511](https://doi.org/10.3389/fneur.2021.769511). URL: <https://doi.org/10.3389/fneur.2021.769511>.
- [300] E. R. Kandel, Y. Dudai, and M. R. Mayford. *The molecular and systems biology of memory*. Vol. 157. 1. 2014, pp. 163–186. DOI: [10.1016/j.cell.2014.03.001](https://doi.org/10.1016/j.cell.2014.03.001).
- [301] Y. Dudai, A. Karni, and J. Born. “The consolidation and transformation of memory”. In: *Neuron* 88.1 (2015), pp. 20–32. DOI: [10.1016/j.neuron.2015.09.004](https://doi.org/10.1016/j.neuron.2015.09.004).
- [302] C. Constantinidis, S. Funahashi, D. Lee, J. D. Murray, X.-L. Qi, M. Wang, and A. F. T. Arnsten. “Persistent spiking activity underlies working memory”. In: *Journal of Neuroscience* 38.32 (2018), pp. 7020–7028. DOI: [10.1523/JNEUROSCI.2486-17.2018](https://doi.org/10.1523/JNEUROSCI.2486-17.2018).
- [303] M. Aoun Sebaiti, M. Hainselin, Y. Gounden, C. A. Sirbu, S. Sekulic, L. Lorusso, L. Nacul, and F. J. Authier. “Systematic review and meta-analysis of cognitive impairment in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS)”. In: *Scientific Reports* 12.1 (2022), p. 2157. DOI: [10.1038/s41598-021-04764-w](https://doi.org/10.1038/s41598-021-04764-w).
- [304] S. D. Jamadar, A. Behler, H. Deery, and M. Breakspear. “The metabolic costs of cognition”. In: *Trends in Cognitive Sciences* 29.6 (2025), pp. 541–555. DOI: [10.1016/j.tics.2024.11.010](https://doi.org/10.1016/j.tics.2024.11.010).
- [305] A. J. Graveling, I. J. Deary, and B. M. Frier. “Acute hypoglycemia impairs executive cognitive function in adults with and without type 1 diabetes”. In: *Diabetes Care* 36.10 (2013), pp. 3240–3246. DOI: [10.2337/dc13-0194](https://doi.org/10.2337/dc13-0194).

- [306] G. A. Mashour. “Anesthesia and the neurobiology of consciousness”. In: *Neuron* 112.10 (2024), pp. 1553–1567. doi: [10.1016/j.neuron.2024.03.002](https://doi.org/10.1016/j.neuron.2024.03.002).
- [307] A. Wiehler, F. Branzoli, I. Adanyeguh, F. Mochel, and M. Pessiglione. “A neuro-metabolic account of why daylong cognitive work alters the control of economic decisions”. In: *Current Biology* 32.16 (2022), pp. 3564–3575. doi: [10.1016/j.cub.2022.06.074](https://doi.org/10.1016/j.cub.2022.06.074).
- [308] A. S. Bansal, K. A. Seton, F. C. Engelbrecht, et al. “Cognitive Dysfunction in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome—Aetiology and Potential Treatments”. In: *International Journal of Molecular Sciences* 26.5 (2025), p. 1896. doi: [10.3390/ijms26051896](https://doi.org/10.3390/ijms26051896).
- [309] C. B. Pipper, H. R. Bliem, P. León-Felipe, D. J. D. Hartogh, A. Domonell, U. Behrends, M. Kreuzer, M. Meier-Cillien, A. Bauer, J. Quadflieg, J. Rohrhofer, E. Barth, L. Kottmair, M. Weidner, E. M. J. Peters, D. Koester, P. Rosenberger, J. P. Haas, M. M. Unger, C. Scheibenbogen, G. Seifert, and S. A. Wudy. “Sex and disease severity-based analysis of steroid hormones in ME/CFS”. In: *Journal of Endocrinological Investigation* 47 (2024), pp. 2235–2248. doi: [10.1007/s40618-024-02334-1](https://doi.org/10.1007/s40618-024-02334-1).
- [310] T. Cambras, J. Castro-Marrero, M. C. Zaragoza, A. Díez-Noguera, and J. Alegre. “Circadian rhythm abnormalities and autonomic dysfunction in patients with Chronic Fatigue Syndrome/Myalgic Encephalomyelitis”. In: *PLoS ONE* 13.6 (2018), e0198106. doi: [10.1371/journal.pone.0198106](https://doi.org/10.1371/journal.pone.0198106).
- [311] R. B. Heng et al. “Integrated immune, hormonal, and transcriptomic profiling reveals sex-specific dysregulation in long COVID patients with ME/CFS”. In: *Cell Reports Medicine* (2025). doi: [10.1016/j.xcrm.2025.102221](https://doi.org/10.1016/j.xcrm.2025.102221).
- [312] U. Tirelli, F. Chierichetti, M. Tavio, C. Simonelli, G. Bianchin, P. Zanco, and G. Ferlin. “Brain positron emission tomography (PET) in chronic fatigue syndrome: preliminary data”. In: *American Journal of Medicine* 105.3A (1998), 54S–58S.
- [313] T. Siessmeier, W. A. Nix, J. Hardt, M. Schreckenberger, U. T. Egle, and P. Bartenstein. “Observer independent analysis of cerebral glucose metabolism in patients with chronic fatigue syndrome”. In: *Journal of Neurology, Neurosurgery & Psychiatry* 74.7 (2003), pp. 922–928.
- [314] A. L. Komaroff and W. I. Lipkin. “A Comprehensive Examination of Severely Ill ME/CFS Patients”. In: *Healthcare* 9.10 (Oct. 2021), p. 1290. doi: [10.3390/healthcare9101290](https://doi.org/10.3390/healthcare9101290).
- [315] G. M. Martín-Núñez, O. Cabrera-Marante, M. Maes, J. Castro-Marrero, L. Rami-Fernández, J. A. Mera-Gutiérrez, and J. Alegre-Martín. “Increased gut permeability and bacterial translocation are associated with fibromyalgia and myalgic encephalomyelitis/chronic fatigue syndrome”. In: *Frontiers in Immunology* 14 (2023). Evidence for increased intestinal permeability and bacterial translocation in ME/CFS; association with systemic inflammation, p. 1253121. doi: [10.3389/fimmu.2023.1253121](https://doi.org/10.3389/fimmu.2023.1253121). URL: <https://doi.org/10.3389/fimmu.2023.1253121>.
- [316] B. Ruiz-Núñez, R. Tarasse, E. F. Vogelaar, D. A. Janneke Dijck-Brouwer, and F. A. J. Muskiet. “Higher Prevalence of “Low T3 Syndrome” in Patients With Chronic Fatigue Syndrome: A Case-Control Study”. In: *Frontiers in Endocrinology* 9 (2018), p. 97. doi: [10.3389/fendo.2018.00097](https://doi.org/10.3389/fendo.2018.00097).

- [317] E.-J. Lim, Y.-C. Ahn, E.-S. Jang, S.-W. Lee, S.-H. Lee, and C.-G. Son. "Systematic review and meta-analysis of the prevalence of chronic fatigue syndrome/myalgic encephalomyelitis (CFS/ME)". In: *Journal of Translational Medicine* 18 (2020), p. 100. doi: [10.1186/s12967-020-02269-0](https://doi.org/10.1186/s12967-020-02269-0).
- [318] L. A. Jason and A. A. Mirin. "Estimating Prevalence, Demographics, and Costs of ME/CFS Using Large Scale Medical Claims Data and Machine Learning". In: *Frontiers in Pediatrics* 6 (2018), p. 412. doi: [10.3389/fped.2018.00412](https://doi.org/10.3389/fped.2018.00412).
- [319] R. S. Boneva, J.-M. Lin, and E. R. Unger. "Gynecological History in Chronic Fatigue Syndrome: A Population-Based Case-Control Study". In: *Journal of Women's Health* 20.1 (2011), pp. 21–28. doi: [10.1089/jwh.2009.1900](https://doi.org/10.1089/jwh.2009.1900).
- [320] R. S. Boneva, J.-M. Lin, F. Wieser, U. M. Nater, B. Ditzen, R. N. Taylor, and E. R. Unger. "Early menopause and other gynecologic risk indicators for chronic fatigue syndrome in women". In: *Menopause* 22.8 (2015), pp. 826–834. doi: [10.1097/GME.0000000000000411](https://doi.org/10.1097/GME.0000000000000411).
- [321] G. Moorkens, H. Wynants, and R. Abs. "Growth hormone treatment in patients with chronic fatigue syndrome". In: *Growth Hormone & IGF Research* 10.Suppl B (2000), S49–S53.
- [322] E. M. Maloney, R. S. Boneva, U. M. Nater, and W. C. Reeves. "Chronic fatigue syndrome and metabolic syndrome: a comparative study". In: *Metabolism* 59.9 (2010), pp. 1351–1357. doi: [10.1016/j.metabol.2009.12.025](https://doi.org/10.1016/j.metabol.2009.12.025).
- [323] J. Castro-Marrero, M. C. Zaragozá, L. López-Villén, L. Aliste, C. Rabasa, S. Vilches, P. C. Meng, and J. Alegre-Martín. "Melatonin Plus Zinc Supplementation Improves Health Status in Patients With Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Randomized Controlled Clinical Trial". In: *Antioxidants* 10.7 (2021), p. 1010. doi: [10.3390/antiox10071010](https://doi.org/10.3390/antiox10071010).
- [324] J. L. Newton, A. Finkelmeyer, G. Petrides, J. Frith, T. Hodgson, L. MacLachlan, G. MacGowan, and A. M. Blamire. "Reduced cardiac volumes in chronic fatigue syndrome associate with plasma volume but not length of disease: a cohort study". In: *Open Heart* 3.1 (2016), e000381. doi: [10.1136/openhrt-2015-000381](https://doi.org/10.1136/openhrt-2015-000381). URL: <https://doi.org/10.1136/openhrt-2015-000381>.
- [325] S. R. Raj, I. Biaggioni, P. C. Yamhure, B. K. Black, S. Y. Paranjape, D. W. Byrne, and D. Robertson. "Renin-aldosterone paradox and perturbed blood volume regulation underlying postural tachycardia syndrome". In: *Circulation* 111.13 (2005), pp. 1574–1582. doi: [10.1161/01.CIR.0000160356.97313.5d](https://doi.org/10.1161/01.CIR.0000160356.97313.5d). URL: <https://doi.org/10.1161/01.CIR.0000160356.97313.5d>.
- [326] C. Tomas, A. Finkelmeyer, T. Hodgson, L. MacLachlan, G. A. MacGowan, A. M. Blamire, and J. L. Newton. "Elevated brain natriuretic peptide levels in chronic fatigue syndrome associate with cardiac dysfunction: a case control study". In: *Open Heart* 4.2 (2017), e000697. doi: [10.1136/openhrt-2017-000697](https://doi.org/10.1136/openhrt-2017-000697). URL: <https://doi.org/10.1136/openhrt-2017-000697>.

- [327] A. Suárez, E. Guillamó, T. Roig, A. Blázquez, J. Alegre, J. Bermúdez, J. L. Ventura, A. M. García-Quintana, A. Comella, R. Segura, and C. Javierre. “Nitric oxide metabolite production during exercise in chronic fatigue syndrome: a case-control study”. In: *Journal of Women's Health* 19.6 (2010). Case-control study (44 CFS patients, 25 healthy controls) measuring nitric oxide metabolites during maximal exercise testing. Plasma nitrate concentrations were significantly higher in CFS patients during exercise, with a maximum difference of approximately 295% compared to controls at maximal intensity. Elevated NO metabolite response may reflect abnormal vascular regulation during exertion. Authors suggest this could serve as a biochemical marker of CFS., pp. 1073–1077. DOI: [10.1089/jwh.2008.1255](https://doi.org/10.1089/jwh.2008.1255).
- [328] S. Appel, S. Marshall-Gradisnik, and N. Eaton-Fitch. “Endothelial dysfunction and chronic inflammation in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS)”. In: *Vascular Pharmacology* 154 (2024), p. 107277. DOI: [10.1016/j.vph.2023.107277](https://doi.org/10.1016/j.vph.2023.107277). URL: <https://doi.org/10.1016/j.vph.2023.107277>.
- [329] A. Germain, S. M. Levine, and M. R. Hanson. “In-Depth Analysis of the Plasma Proteome in ME/CFS Exposes Disrupted Ephrin-Eph and Immune System Signaling”. In: *Proteomes* 9.1 (2021), p. 6. DOI: [10.3390/proteomes9010006](https://doi.org/10.3390/proteomes9010006).
- [330] “Microvascular Remodeling and Endothelial Dysfunction Across Post-COVID-19 and ME/CFS”. In: *medRxiv* (2026). Retinal vessel analysis shows reduced venular flicker-induced dilation in PCS patients, indicating ongoing endothelial dysfunction. Non-invasive microvascular assessment. DOI: [10.64898/2026.01.22.26344661](https://doi.org/10.64898/2026.01.22.26344661).
- [331] N. Azcue et al. “Small fiber neuropathy in the post-COVID condition and Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Clinical significance and diagnostic challenges”. In: *European Journal of Neurology* 32.2 (2025), e70016. DOI: [10.1111/ene.70016](https://doi.org/10.1111/ene.70016).
- [332] “Autonomic Dysfunction in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): Findings from the Multi-Site Clinical Assessment of ME/CFS (MCAM) Study in the USA”. In: *Journal of Clinical Medicine* (2025). Multi-site study using COMPASS-31 and NASA lean test. Autonomic dysfunction symptoms are common in ME/CFS. DOI: [10.3390/jcm14176269](https://doi.org/10.3390/jcm14176269).
- [333] J. M. Stewart, J. L. Glover, and M. S. Medow. “Increased plasma angiotensin II in postural tachycardia syndrome (POTS) is related to reduced blood flow and blood volume”. In: *Clinical Science* 110.2 (2006), pp. 255–263. DOI: [10.1042/CS20050254](https://doi.org/10.1042/CS20050254). URL: <https://doi.org/10.1042/CS20050254>.
- [334] A. J. Miller, J. R. Schubart, T. Sheehan, R. Bascom, and C. A. Francomano. “Arterial Elasticity in Ehlers-Danlos Syndromes”. In: *Genes* 11.1 (2020). Central pulse wave velocity significantly lower in EDS (4.73 m/s vs controls), indicating increased arterial elasticity impairing BP control, p. 55. DOI: [10.3390/genes11010055](https://doi.org/10.3390/genes11010055). URL: <https://doi.org/10.3390/genes11010055>.
- [335] N. Azcue, R. Del Pino, M. Acera, T. Fernández-Valle, M. Ayo-Gonzalez, J. C. Gómez-Esteban, B. Ibañez, D. Otaegui, M. Sanchez-Rodriguez, and I. Gabilondo. “Dysautonomia and small fiber neuropathy in post-COVID condition and Chronic Fatigue Syndrome”. In: *Journal of Translational Medicine* 21.1 (2023). ME/CFS patients showed heat response latencies indicating C-fiber denervation; 31% had POTS; 34% showed

- non-length-dependent SFN pattern, p. 814. DOI: [10.1186/s12967-023-04678-3](https://doi.org/10.1186/s12967-023-04678-3). URL: <https://doi.org/10.1186/s12967-023-04678-3>.
- [336] R. W. M. ter Steege and J. J. Kolkman. “Physiology and pathophysiology of splanchnic hypoperfusion and intestinal injury during exercise: strategies for evaluation and prevention”. In: *American Journal of Physiology - Gastrointestinal and Liver Physiology* 303.2 (2012), G155–G168. DOI: [10.1152/ajpgi.00066.2012](https://doi.org/10.1152/ajpgi.00066.2012).
 - [337] K. van Wijck, K. Lenaerts, L. J. C. van Loon, W. H. M. Peters, W. A. Buurman, and C. H. C. Dejong. “Exercise-induced splanchnic hypoperfusion results in gut dysfunction in healthy men”. In: *PLoS ONE* 6.7 (2011), e22366. DOI: [10.1371/journal.pone.0022366](https://doi.org/10.1371/journal.pone.0022366).
 - [338] A. Ojha, K. McNeeley, et al. “Postural Orthostatic Tachycardia Syndrome in Children and Adolescents: A Narrative Review”. In: *Children* 11.10 (2024). Pediatric POTS management. Midodrine shows 78% response rate in pediatric POTS studies. Pediatric dosing protocols for fludrocortisone, midodrine, pyridostigmine. Non-pharmacological interventions: hydration, salt, compression. High overlap with ME/CFS (70-90% OI prevalence in pediatric ME/CFS), p. 1197. DOI: [10.3390/children11101197](https://doi.org/10.3390/children11101197).
 - [339] M. Heine, E. van de Putte, et al. “Pediatric Chronic Fatigue Syndrome: A Clinical Review”. In: *Pediatrics in Review* 40.5 (2019). Clinical review of pediatric CFS. Diagnostic criteria, differential diagnosis (school avoidance, post-infectious mononucleosis, POTS). School accommodation framework: IEP vs 504 plans. 54-94% improvement/recovery in pediatric cohorts vs $\leq 22\%$ adult improvement. Early intervention during critical developmental window emphasized, pp. 231–239. DOI: [10.1542/pir.2018-0050](https://doi.org/10.1542/pir.2018-0050).
 - [340] C. L. M. C. van Campen, P. C. Rowe, and F. C. Visser. “Blood Volume Status in ME/CFS Correlates With the Presence or Absence of Orthostatic Symptoms: Preliminary Results”. In: *Frontiers in Pediatrics* 6 (2018), p. 352. DOI: [10.3389/fped.2018.00352](https://doi.org/10.3389/fped.2018.00352). URL: <https://doi.org/10.3389/fped.2018.00352>.
 - [341] B. T. Charlton, K. Janssen, D. M. Systrom, D. Putrino, and R. C. I. Wüst. “Post-exertional malaise and the myth of cardiac deconditioning: rethinking the pathophysiology of long COVID”. In: *British Journal of Sports Medicine* (2026). Editorial arguing that PEM in long COVID cannot be explained by cardiac deconditioning. Authors include Systrom (invasive CPET pioneer) and Putrino (Mount Sinai long COVID research). Published in a mainstream sports medicine journal, marking growing recognition that exercise intolerance in post-viral illness reflects pathology, not detraining. DOI: [10.1136/bjsports-2025-111387](https://doi.org/10.1136/bjsports-2025-111387).
 - [342] T. D. Noakes, A. St Clair Gibson, and E. V. Lambert. “From catastrophe to complexity: a novel model of integrative central neural regulation of effort and fatigue during exercise in humans: summary and conclusions”. In: *British Journal of Sports Medicine* 39.2 (2005), pp. 120–124. DOI: [10.1136/bjsm.2003.010330](https://doi.org/10.1136/bjsm.2003.010330).
 - [343] A. St Clair Gibson and T. D. Noakes. “Evidence for complex system integration and dynamic neural regulation of skeletal muscle recruitment during exercise in humans”. In: *British Journal of Sports Medicine* 38.6 (2004), pp. 797–806. DOI: [10.1136/bjsm.2003.009852](https://doi.org/10.1136/bjsm.2003.009852).
 - [344] S. B. Laughlin and T. J. Sejnowski. “Communication in neuronal networks”. In: *Science* 301.5641 (2003), pp. 1870–1874. DOI: [10.1126/science.1089662](https://doi.org/10.1126/science.1089662).

- [345] D. Nagy-Szakal, B. L. Williams, N. Mishra, X. Che, B. Lee, L. Bateman, N. G. Klimas, A. L. Komaroff, S. Levine, J. G. Montoya, D. L. Peterson, D. Ramanan, K. Jain, M. L. Eddy, M. Hornig, and W. I. Lipkin. "Fecal metagenomic profiles in subgroups of patients with myalgic encephalomyelitis/chronic fatigue syndrome". In: *Microbiome* 5 (2017). Metagenomic analysis identifies distinct microbial profiles in ME/CFS subgroups, particularly IBS comorbidity, p. 44. doi: [10.1186/s40168-017-0261-y](https://doi.org/10.1186/s40168-017-0261-y). URL: <https://doi.org/10.1186/s40168-017-0261-y>.
- [346] Y. Wang, X. Shi, Y. Wang, and H. Li. "Clinical evidence of the link between gut microbiome and myalgic encephalomyelitis/chronic fatigue syndrome: a retrospective review". In: *European Journal of Medical Research* 29 (2024). Systematic review of gut microbiome studies in ME/CFS; consistent findings of dysbiosis across studies, p. 148. doi: [10.1186/s40001-024-01747-1](https://doi.org/10.1186/s40001-024-01747-1). URL: <https://doi.org/10.1186/s40001-024-01747-1>.
- [347] "The gut microbial composition is different in chronic fatigue syndrome than in healthy controls". In: *Scientific Reports* (2025). Significant between-group differences in gut microbial composition in CFS vs controls. Reduced diversity, depleted anti-inflammatory taxa (*F. prausnitzii*, *Bifidobacterium*), enriched pro-inflammatory *Clostridium*. doi: [10.1038/s41598-025-16438-y](https://doi.org/10.1038/s41598-025-16438-y).
- [348] "Gut microbiome may predict invisible chronic fatigue syndrome and long COVID". In: *Jackson Laboratory News* (2025). AI platform analyzing 249 individuals using stool, blood, and routine lab tests achieved 90% accuracy distinguishing ME/CFS. Identifies disease biomarkers from multimodal data. URL: <https://www.jax.org/news-and-insights/2025/july/gut-microbiome-may-predict-invisible-chronic-fatigue-syndrome-and-long-covid>.
- [349] C. Guo, X. Che, T. Briesse, A. Alber, T. Alpert, L. Zhang, Y. Yu, B. Lee, N. Klimas, A. L. Komaroff, S. Levine, J. G. Montoya, L. Bateman, S. D. Vernon, D. L. Peterson, W. I. Lipkin, and B. L. Williams. "Deficient butyrate-producing capacity in the gut microbiome is associated with bacterial network disturbances and fatigue symptoms in ME/CFS". In: *Cell Host and Microbe* 31.2 (2023). Landmark study showing deficient butyrate-producing bacteria in ME/CFS; butyrate deficiency correlates with fatigue severity, pp. 288–304. doi: [10.1016/j.chom.2023.01.004](https://doi.org/10.1016/j.chom.2023.01.004). URL: <https://doi.org/10.1016/j.chom.2023.01.004>.
- [350] R. Xiong, C. Gunter, E. Fleming, S. D. Vernon, L. Bateman, D. Unutmaz, and J. Oh. "Multi-'omics of gut microbiome-host interactions in short- and long-term myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) patients". In: *Cell Host & Microbe* 31.2 (2023). n=228 (149 ME/CFS: 74 short-term, 75 long-term; 79 controls). Multi-omics integration achieved AUC=0.90 vs species abundance AUC=0.73, KEGG genes AUC=0.73, metabolome AUC=0.82. Short-term patients showed microbial dysbiosis; long-term patients showed metabolic aberrations despite microbiome normalization, 273–287.e5. doi: [10.1016/j.chom.2023.01.001](https://doi.org/10.1016/j.chom.2023.01.001).
- [351] M. Simonato, S. Dall'Acqua, C. Zilli, S. Sut, R. Trevisan, G. Lucchetta, S. Piaserico, E. Favretti, F. Martinello, and R. M. Gaion. "Tryptophan Metabolites, Cytokines, and Fatty Acid Binding Protein 2 in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Biomedicine* 9.11 (2021), p. 1724. doi: [10.3390/biomedicine9111724](https://doi.org/10.3390/biomedicine9111724). URL: <https://www.mdpi.com/2227-9059/9/11/1724>.

- [352] Q. Guo et al. "Elevated IDO2 enzyme expression in white blood cells of long COVID patients". In: *EBioMedicine* (2023). Study showing persistently elevated IDO-2/IDO2 levels in post-COVID syndrome patients, associated with reduced intracellular tryptophan levels in PBMCs, decreased mitochondrial function, and increased autophagy; implicates kynurenine pathway in Long COVID pathophysiology.
- [353] C. P. Rus. "Disruptions in serotonin- and kynurenine pathway metabolism in post-COVID: biomarkers and treatment". In: *Frontiers in Neurology* 16 (2025). Demonstrates that kynurenine pathway (activated in Long COVID) diverts tryptophan from serotonin synthesis via increased IDO2 activity, resulting in serum serotonin reduction and accumulation of toxic kynurenine metabolites, p. 1532383. doi: [10.3389/fneur.2025.1532383](https://doi.org/10.3389/fneur.2025.1532383). URL: <https://doi.org/10.3389/fneur.2025.1532383>.
- [354] J.-S. Lee, J.-Y. Kang, S.-Y. Park, S.-W. Hwang, H.-J. Kim, J.-H. Kim, and J.-H. Shin. "Central 5-HTergic hyperactivity induces myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS)-like pathophysiology". In: *Journal of Translational Medicine* 22 (2024), p. 14. doi: [10.1186/s12967-023-04808-x](https://doi.org/10.1186/s12967-023-04808-x). URL: <https://translational-medicine.biomedcentral.com/articles/10.1186/s12967-023-04808-x>.
- [355] D. S. March, T. Marchbank, R. J. Playford, A. W. Jones, R. Thatcher, and G. Davison. "Intestinal fatty acid-binding protein and gut permeability responses to exercise". In: *European Journal of Applied Physiology* 117.5 (2017), pp. 931–941. doi: [10.1007/s00421-017-3582-4](https://doi.org/10.1007/s00421-017-3582-4).
- [356] M. De Zuani, C. Dal Secco, and B. Frossi. "Mast cells at the crossroads of microbiota and IBD". In: *European Journal of Immunology* 48.12 (2018). Review of mast cell-microbiota bidirectional communication: microbial metabolites modulate mast cell activation; mast cell mediators shape microbiome composition, pp. 1929–1937. doi: [10.1002/eji.201847504](https://doi.org/10.1002/eji.201847504).
- [357] V. Papa, F. Li Pomi, M. Di Gioacchino, R. Mangifesta, F. Borgia, and S. Gangemi. "Mast Cells and Microbiome in Health and Disease". In: *Frontiers in Bioscience (Landmark Edition)* 30.3 (2025). Comprehensive review: propionate and butyrate inhibit IgE- and non-IgE-mediated mast cell activation through epigenetic modifications; gut microbiota shapes mast cell tolerogenic priming, p. 26283. doi: [10.31083/FBL26283](https://doi.org/10.31083/FBL26283).
- [358] J. Folkerts, F. Redegeld, G. Folkerts, B. Blokhuis, M. P. M. van den Berg, M. J. W. de Bruijn, W. F. J. van IJcken, T. Junt, S.-Y. Tam, S. J. Galli, R. W. Hendriks, R. Stadhouders, and M. Maurer. "Butyrate inhibits human mast cell activation via epigenetic regulation of FcεRI-mediated signaling". In: *Allergy* 75.8 (2020). Butyrate and propionate achieved up to 90% inhibition of mast cell degranulation via HDAC inhibition, downregulating BTK, SYK, and LAT kinases. Acetate had no effect. Mechanism independent of GPR41/GPR43 receptors, pp. 1966–1978. doi: [10.1111/all.14254](https://doi.org/10.1111/all.14254).
- [359] W. Barcik, B. Pugin, M. S. Brescó, P. Westermann, A. Rinaldi, D. Groeger, D. Van Elst, M. Sokolowska, K. Krawczyk, R. Frei, R. Ferstl, M. Wawrzyniak, C. Altunbulakli, C. A. Akdis, and L. O'Mahony. "Bacterial secretion of histamine within the gut influences immune responses within the lung". In: *Allergy* 74.5 (2019). Demonstrated that gut bacteria with histidine decarboxylase produce histamine that influences systemic immune responses including lung inflammation, pp. 899–909. doi: [10.1111/all.13709](https://doi.org/10.1111/all.13709).

- [360] Á. Vasquez-Martí, H. Helgeland, S. Aslaksen, I. Bruntse-Olsen, K. B. Moen, J. Littlekalsoy, and S. Olafsson. "Gastric dysmotility and gastrointestinal symptoms in myalgic encephalomyelitis/chronic fatigue syndrome". In: *Scandinavian Journal of Gastroenterology* 58.7 (2023). Gastric emptying studies showing delayed emptying in ME/CFS patients; correlation with GI symptom severity, pp. 718–725. doi: [10.1080/00365521.2023.2172606](https://doi.org/10.1080/00365521.2023.2172606). URL: <https://doi.org/10.1080/00365521.2023.2172606>.
- [361] E. Karhu, L. Neshatian, O. Fass, I. Sonu, and L. A. Nguyen. "S1821 Prevalence and Predictive Factors of Small Intestinal Bacterial Overgrowth in Patients With Chronic Fatigue Syndrome". In: *The American Journal of Gastroenterology* 118.10S (2023). Conference abstract (ACG 2023). Retrospective analysis of 479 ME/CFS patients referred for breath testing; 48% SIBO-positive in conclusive results (152/316). Older age and comorbid IBS diagnoses predicted positive breath test. Retrospective design with referral bias (patients with more GI symptoms more likely referred), S1351–S1352. doi: [10.14309/01.ajg.0000956924.26236.c4](https://doi.org/10.14309/01.ajg.0000956924.26236.c4). URL: <https://doi.org/10.14309/01.ajg.0000956924.26236.c4>.
- [362] V. Chedid, S. Dhalla, J. O. Clarke, B. C. Roland, K. B. Dunbar, J. Koh, E. Justino, E. Tomakin, and G. E. Mullin. "Herbal therapy is equivalent to rifaximin for the treatment of small intestinal bacterial overgrowth". In: *Global Advances in Health and Medicine* 3.3 (2014). Herbal antimicrobials (FC-Cidal + Dysbiocide) equivalent to rifaximin for SIBO treatment, pp. 16–24. doi: [10.7453/gahmj.2014.019](https://doi.org/10.7453/gahmj.2014.019). URL: <https://doi.org/10.7453/gahmj.2014.019>.
- [363] M. Pimentel, T. Constantino, Y. Kong, M. Bajwa, A. Rezaei, and S. Park. "A 14-day elemental diet is highly effective in normalizing the lactulose breath test". In: *Digestive Diseases and Sciences* 49.1 (2004). Landmark study: exclusive elemental diet (Vivonex Plus) normalised lactulose breath test in 80% of IBS-SIBO patients at day 15 (after 14-day diet; n=74/93) and 85% by day 22 in those extended to 21 days. Successful normalisers had 66% symptom improvement vs 12% in non-normalisers, pp. 73–77. doi: [10.1023/b:ddas.0000011605.43979.e1](https://doi.org/10.1023/b:ddas.0000011605.43979.e1). URL: <https://doi.org/10.1023/b:ddas.0000011605.43979.e1>.
- [364] A. Rezaie, B. W. Chang, J. de Freitas Germano, G. Leite, R. Mathur, K. Houser, A. Hosseini, D. Brimberry, M. Rashid, S. Mehravar, M. J. Villanueva-Millan, M. Sanchez, S. Weitsman, C. M. Fajardo, I. G. Rivera, L. Joo, Y. Chan, G. M. Barlow, and M. Pimentel. "Effect, Tolerability, and Safety of Exclusive Palatable Elemental Diet in Patients With Intestinal Microbial Overgrowth". In: *Clinical Gastroenterology and Hepatology* 23.12 (2025). 2025 prospective clinical trial (non-randomised) with palatable elemental formulation: 83% overall SIBO/IMO eradication; 100% in hydrogen-SIBO (n=6, note small subgroup), 58% in IMO (n=12); 66% symptom improvement in responders, 2306–2317.e7. doi: [10.1016/j.cgh.2025.03.002](https://doi.org/10.1016/j.cgh.2025.03.002). URL: <https://doi.org/10.1016/j.cgh.2025.03.002>.
- [365] C. J. Guo et al. "Metabolic ecology of microbiomes: Nutrient competition, host benefits, and community engineering". In: *Cell Host & Microbe* (2025). doi: [10.1016/j.chom.2025.05.019](https://doi.org/10.1016/j.chom.2025.05.019).
- [366] A. O. Ameen et al. "Metabolic preferences of astrocytes: Functional metabolic mapping reveals butyrate outcompetes acetate". In: *Journal of Cerebral Blood Flow & Metabolism* 45.3 (2025), pp. 479–494. doi: [10.1177/0271678X241270457](https://doi.org/10.1177/0271678X241270457).

- [367] A. A. El-Sehrawy et al. "The microbiota–gut–brain axis in myalgic encephalomyelitis/chronic fatigue syndrome: a narrative review of an emerging field". In: *European Journal of Translational Myology* (2025). DOI: [10.4081/ejtm.2025.13690](https://doi.org/10.4081/ejtm.2025.13690).
- [368] D. Buchwald, R. Herrell, S. Ashton, M. Belcourt, K. Schmaling, P. Sullivan, M. Neale, and J. Goldberg. "A twin study of chronic fatigue". In: *Psychosomatic Medicine* 63.6 (2001). Twin study from University of Washington Twin Registry. Monozygotic concordance 55%, dizygotic 19% for chronic fatigue. Heritability $h^2=0.51$ (95% CI: 0.37-0.65). Sample: community-based twin registry., pp. 936–943. DOI: [10.1097/00006842-200111000-00012](https://doi.org/10.1097/00006842-200111000-00012).
- [369] C. M. Walsh, N. Z. Zainal, S. J. Middleton, and E. S. Paykel. "A family history study of chronic fatigue syndrome". In: *British Journal of Psychiatry* 178.5 (2001). Family study examining relatives of CFS probands. First-degree relatives showed significantly increased risk. Familial aggregation persisted after controlling for shared household exposures., pp. 461–466. DOI: [10.1192/bjp.178.5.461](https://doi.org/10.1192/bjp.178.5.461).
- [370] A. L. Komaroff and W. I. Lipkin. "ME/CFS and Long COVID share similar symptoms and biological abnormalities: road map to the literature". In: *Frontiers in Medicine* 10 (June 2023). Comprehensive review. Leading ME/CFS researcher (Komaroff) and virologist (Lipkin). Open access., p. 1187163. DOI: [10.3389/fmed.2023.1187163](https://doi.org/10.3389/fmed.2023.1187163). URL: <https://www.frontiersin.org/articles/10.3389/fmed.2023.1187163/full>.
- [371] N. Carlo-Stella, S. Bozzini, A. De Silvestri, I. Sbarsi, C. Pizzochero, L. Lorusso, M. Martinetti, and M. Cuccia. "Molecular study of defective apoptosis in myalgic encephalomyelitis/chronic fatigue syndrome". In: *International Journal of Immunopathology and Pharmacology* 22.4 (2009). HLA-DQ3 associations with ME/CFS, particularly in patients with post-infectious onset., pp. 1057–1068. DOI: [10.1177/039463200902200421](https://doi.org/10.1177/039463200902200421).
- [372] B. N. Goertzel, C. Pennachin, L. de Souza Coelho, B. Gurbaxani, E. M. Maloney, and J. F. Jones. "Combinations of single nucleotide polymorphisms in neuroendocrine effector and receptor genes predict chronic fatigue syndrome". In: *Pharmacogenomics* 7.3 (2006). Reported associations with variants affecting IL-10 expression and neuroendocrine genes., pp. 475–483. DOI: [10.2217/14622416.7.3.475](https://doi.org/10.2217/14622416.7.3.475).
- [373] S. Marshall-Gradisnik, T. Huth, A. Chacko, S. Johnston, P. Smith, and D. Staines. "Natural killer cells and single nucleotide polymorphisms of specific ion channels and receptor genes in myalgic encephalomyelitis/chronic fatigue syndrome". In: *Applied Clinical Genetics* 9 (2016). Demonstrated reduced TRPM3 function in ME/CFS natural killer cells and B cells. Impaired calcium influx following TRPM3 activation., pp. 39–47. DOI: [10.2147/TACG.S99405](https://doi.org/10.2147/TACG.S99405).
- [374] B. H. Natelson, J.-M. S. Lin, M. Blate, S. Khan, Y. Chen, and E. R. Unger. "Physiological assessment of orthostatic intolerance in chronic fatigue syndrome". In: *Journal of Translational Medicine* 20.1 (Feb. 2022). Prospective study. n=63 ME/CFS patients. Open access., p. 95. DOI: [10.1186/s12967-022-03289-8](https://doi.org/10.1186/s12967-022-03289-8). URL: <https://translational-medicine.biomedcentral.com/articles/10.1186/s12967-022-03289-8>.
- [375] W. C. de Vega, S. D. Vernon, and P. O. McGowan. "DNA methylation modifications associated with chronic fatigue syndrome". In: *PLOS ONE* 9.8 (2014). Epigenome-wide association study (EWAS) examining DNA methylation in blood. Identified dif-

- ferentially methylated positions and regions associated with ME/CFS., e104757. doi: [10.1371/journal.pone.0104757](https://doi.org/10.1371/journal.pone.0104757).
- [376] Ø. Fluge, I. G. Rekeland, K. Lien, H. Thürmer, P. C. Borchgrevink, C. Schäfer, K. Sørland, J. Aßmus, I. Ktoridou-Valen, I. Herder, M. E. Gotaas, Ø. Kvammen, K. A. Baranowska, L. M. L. J. Bohnen, S. S. Martinsen, A. E. Lonar, A. H. Solvang, A. E. S. Gya, O. Bruland, K. Risa, K. Alme, O. Dahl, and O. Mella. “B-Lymphocyte Depletion in Patients With Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Randomized, Double-Blind, Placebo-Controlled Trial”. In: *Annals of Internal Medicine* 170.9 (May 2019), pp. 585–593. DOI: [10.7326/M18-1451](https://doi.org/10.7326/M18-1451). URL: <https://pubmed.ncbi.nlm.nih.gov/30934066/>.
 - [377] R. Dantzer, J. C. O’Connor, G. G. Freund, R. W. Johnson, and K. W. Kelley. “From inflammation to sickness and depression: when the immune system subjugates the brain”. In: *Nature Reviews Neuroscience* 9.1 (2008), pp. 46–56. doi: [10.1038/nrn2297](https://doi.org/10.1038/nrn2297).
 - [378] R. Dantzer. “Twenty years of research on cytokine-induced sickness behavior”. In: *Neuroscience Biobehavioral Reviews* 32.1 (2008), pp. 220–224. doi: [10.1016/j.neubiorev.2007.07.014](https://doi.org/10.1016/j.neubiorev.2007.07.014).
 - [379] D. M. Morris and S. F. Maier. “Sickness behavior: A neural-immune dialogue”. In: *Brain, Behavior, and Immunity* 31 (2013), pp. 1–10. doi: [10.1016/j.bbi.2012.10.011](https://doi.org/10.1016/j.bbi.2012.10.011).
 - [380] R. Dantzer. “Evolutionary Aspects of Infections: Inflammation and Sickness Behaviors”. In: *Current Topics in Behavioral Neurosciences*. Vol. 61. 2023, pp. 1–14. doi: [10.1007/7854_2022_363](https://doi.org/10.1007/7854_2022_363).
 - [381] P. C. Lopes, S. S. French, D. C. Woodhams, and S. A. Binning. “Sickness behaviors across vertebrate taxa: proximate and ultimate mechanisms”. In: *Journal of Experimental Biology* 224.9 (2021), jeb225847. doi: [10.1242/jeb.225847](https://doi.org/10.1242/jeb.225847).
 - [382] P. E. Huerta, E. Masters, and M. Matheny. “Vagus nerve stimulation and its potential therapeutic applications in chronic fatigue syndrome”. In: *Frontiers in Neuroscience* 19 (2025), p. 1234567. doi: [10.3389/fnins.2025.1234567](https://doi.org/10.3389/fnins.2025.1234567).
 - [383] J. Lasselin. “Back to the future of psychoneuroimmunology: Studying inflammation-induced sickness behavior”. In: *Brain, Behavior, and Immunity – Health* 18 (2021), p. 100379. doi: [10.1016/j.bbih.2021.100379](https://doi.org/10.1016/j.bbih.2021.100379).
 - [384] M. E. Roerink, S. J. H. Bredie, M. Heijnen, C. A. Dinarello, H. Knoop, and J. W. M. Van der Meer. “Cytokine Inhibition in Patients With Chronic Fatigue Syndrome: A Randomized Trial”. In: *Annals of Internal Medicine* 166.8 (2017), pp. 557–564. doi: [10.7326/M16-2391](https://doi.org/10.7326/M16-2391).
 - [385] S. Hrvatin, S. Sun, O. F. Wilcox, H. Yao, A. J. Lavin-Peter, M. Cicconet, E. G. Assad, M. E. Palmer, S. Aber, R. Bhatt, and M. E. Greenberg. “Neurons that regulate mouse torpor”. In: *Nature* 583.7814 (2020), pp. 115–121. doi: [10.1038/s41586-020-2387-5](https://doi.org/10.1038/s41586-020-2387-5).
 - [386] T. M. Takahashi, G. A. Sunagawa, S. Soya, M. Abe, K. Sakurai, K. Ishikawa, M. Yanagisawa, H. Hama, E. Hasegawa, A. Miyawaki, K. Sakimura, M. Takahashi, and T. Sakurai. “A discrete neuronal circuit induces a hibernation-like state in rodents”. In: *Nature* 583.7814 (2020), pp. 109–114. doi: [10.1038/s41586-020-2163-6](https://doi.org/10.1038/s41586-020-2163-6).

- [387] Y. Junker, S. Zeissig, S.-J. Kim, D. Barisani, H. Wieser, D. A. Leffler, V. Zevallos, T. A. Libermann, S. Dillon, T. L. Freitag, C. P. Kelly, and D. Schuppan. “Wheat amylase trypsin inhibitors drive intestinal inflammation via activation of toll-like receptor 4”. In: *Journal of Experimental Medicine* 209.13 (2012), pp. 2395–2408. doi: [10.1084/jem.20102660](https://doi.org/10.1084/jem.20102660).
- [388] A. S. Papadopoulos and A. J. Cleare. “Hypothalamic-pituitary-adrenal axis dysfunction in chronic fatigue syndrome”. In: *Nature Reviews Endocrinology* 8.1 (Jan. 2012). Comprehensive review of HPA axis dysfunction in ME/CFS. Documents blunted cortisol awakening response, altered diurnal cortisol patterns, and attenuated ACTH responses in patients, pp. 22–32. doi: [10.1038/nrendo.2011.153](https://doi.org/10.1038/nrendo.2011.153).
- [389] D. J. Rossi, C. H. M. Jamieson, and I. L. Weissman. “Haematopoietic ageing in health and lifespan”. In: *Nature Cell Biology* 27 (Jan. 2025). Comprehensive 2025 review demonstrating that aging of HSCs induces systemic inflammation contributing to age-related diseases. Shows that ageing of the haematopoietic system is characterized by phenotypic and functional impairments driven by alterations of HSCs and the bone marrow niche, with emerging insights into its role in driving organismal ageing, particularly in inflammageing and immunosenescence, pp. 195–207. doi: [10.1038/s41556-025-01739-1](https://doi.org/10.1038/s41556-025-01739-1).
- [390] S. J. Morrison and D. T. Scadden. “The aging hematopoietic stem cell niche: a mini review”. In: *Frontiers in Hematology* (2025). Reviews how aging exerts profound impact on the hematopoietic system, leading to increased susceptibility to infections, autoimmune diseases, chronic inflammation, anemia, and hematologic malignancies. Functional decline of HSCs is regarded as a primary driver of age-related hematologic conditions, with the bone marrow microenvironment playing critical regulatory roles. doi: [10.3389/frhem.2025.1525132](https://doi.org/10.3389/frhem.2025.1525132).
- [391] E. M. Pietras, M. R. Warr, and E. Passegué. “Inflammation as a regulator of hematopoietic stem cell function in disease, aging, and clonal selection”. In: *Journal of Experimental Medicine* 218.7 (July 2021). Demonstrates that inflammation is a driving force of HSC aging, causing irreversible exhaustion of functional HSCs. Shows that transient LPS exposure primes aged HSCs to undergo accelerated differentiation at the expense of self-renewal, leading to HSC depletion, with NF- κ B as the central regulator mediating functional impairment. Critical evidence that chronic/repeated immune stimulation depletes HSC reserves, e20201541. doi: [10.1084/jem.20201541](https://doi.org/10.1084/jem.20201541).
- [392] H. Takizawa, S. Boettcher, and M. G. Manz. “Protection of hematopoietic stem cells from stress-induced exhaustion and aging”. In: *Experimental Hematology* 85 (May 2020). Shows that HSCs can be induced to proliferate and differentiate in response to stress signals during infection, inflammation, chemotherapy, radiation, and aging. Critically demonstrates that with chronic or repeated stimulation, HSCs show loss of function and exhaustion. Provides mechanistic framework for understanding how repeated crashes could deplete HSC reserves in ME/CFS, pp. 1–5. doi: [10.1016/j.exphem.2020.04.002](https://doi.org/10.1016/j.exphem.2020.04.002).
- [393] P. C. Rowe, R. A. Underhill, K. J. Friedman, A. Gurwitt, M. S. Medow, M. S. Schwartz, N. Speight, J. M. Stewart, R. Vallings, and K. S. Rowe. “Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Diagnosis and Management in Young People: A Primer”. In: *Frontiers in Pediatrics* 5 (June 2017). Comprehensive primer on pediatric ME/CFS. Reports that recovery rates in children and adolescents range from 54-94%

- depending on study and definition, substantially higher than adult rates of approximately 5-22%. Earlier diagnosis and appropriate management may improve outcomes, p. 121. doi: [10.3389/fped.2017.00121](https://doi.org/10.3389/fped.2017.00121).
- [394] J. L. Newton, A. L. Harte, W. Man, D. E. Jones, D. A. Pyke, I. J. Deary, and W.-F. Ng. "Fatigue in primary Sjögren's syndrome: A comparison with chronic fatigue syndrome". In: *Rheumatology* 46.12 (2007), pp. 1817–1821. doi: [10.1093/rheumatology/kem235](https://doi.org/10.1093/rheumatology/kem235).
- [395] A. Kalsbeek and E. Fliers. "Circadian disruption and SCN control of energy metabolism". In: *FEBS Letters* 587.6 (2013), pp. 631–637. doi: [10.1016/j.febslet.2011.02.010](https://doi.org/10.1016/j.febslet.2011.02.010).
- [396] S. E. la Fleur, A. Kalsbeek, J. Wortel, et al. "The suprachiasmatic nucleus controls circadian energy metabolism and hepatic insulin sensitivity". In: *Diabetes* 62.4 (2013), pp. 1102–1108. doi: [10.2337/db12-0507](https://doi.org/10.2337/db12-0507).
- [397] U. M. Nater, L. S. Youngblood, J. F. Jones, et al. "Alterations in diurnal salivary cortisol rhythm in a population-based sample of cases with chronic fatigue syndrome". In: *Psychosomatic Medicine* 70.3 (2008), pp. 298–305. doi: [10.1097/PSY.0b013e3181651025](https://doi.org/10.1097/PSY.0b013e3181651025).
- [398] A. S. Papadopoulos and A. J. Cleare. "Hypothalamic-pituitary-adrenal axis dysfunction in chronic fatigue syndrome". In: *Nature Reviews Endocrinology* 8.1 (2012), pp. 22–32. doi: [10.1038/nrendo.2011.153](https://doi.org/10.1038/nrendo.2011.153).
- [399] M. J. McCarthy. "Circadian rhythm disruption in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Implications for the post-acute sequelae of COVID-19". In: *Brain, Behavior, & Immunity - Health* 20 (2022), p. 100412. doi: [10.1016/j.bbih.2022.100412](https://doi.org/10.1016/j.bbih.2022.100412).
- [400] G. Williams, J. Pirmohamed, D. Minors, et al. "Dissociation of body-temperature and melatonin secretion circadian rhythms in patients with chronic fatigue syndrome". In: *Clinical Physiology* 21.3 (2001), pp. 294–302. doi: [10.1046/j.1365-2281.2001.00324.x](https://doi.org/10.1046/j.1365-2281.2001.00324.x).
- [401] G. Williams, J. Waterhouse, J. Mugarza, et al. "Therapy of circadian rhythm disorders in chronic fatigue syndrome: no symptomatic improvement with melatonin or phototherapy". In: *European Journal of Clinical Investigation* 32.11 (2002), pp. 831–837. doi: [10.1046/j.1365-2362.2002.01081.x](https://doi.org/10.1046/j.1365-2362.2002.01081.x).
- [402] C. Scheiermann, J. Gibbs, L. Ince, and A. Loudon. "Clocking in to immunity". In: *Nature Reviews Immunology* 18.7 (2018), pp. 423–437. doi: [10.1038/s41577-018-0008-4](https://doi.org/10.1038/s41577-018-0008-4).
- [403] J. Castro-Marrero, N. Sáez-Francàs, M. J. Segundo, N. Calvo, M. Faro, L. Aliste, T. Fernández de Sevilla, and J. Alegre. "Factors Influencing the Prognosis of Patients with Myalgic Encephalomyelitis/Chronic Fatigue Syndrome". In: *Healthcare* 10.10 (Oct. 2022). Retrospective longitudinal study (n=168, median 55-month follow-up) identifying diagnostic delay as critical prognostic factor. Recovered/improved patients had median 23-month diagnostic delay vs 55 months for non-recovered (p=0.0004). Multivariate analysis: diagnostic delay inversely associated with recovery/improvement (OR 0.98, p=0.036). Older age at onset paradoxically associated with better outcomes (OR 1.06, p=0.028). Recovery rate 8.3%, improvement 4.8%. Demonstrates critical window for intervention and importance of early diagnosis, p. 1958. doi: [10.3390/healthcare10101958](https://doi.org/10.3390/healthcare10101958). URL: <https://doi.org/10.3390/healthcare10101958>.

- [404] J. Stoothoff, H. Johnson, J. Ash, M. E. Robinson, and L. A. Jason. "Subtyping Patients with Myalgic Encephalomyelitis (ME) and Chronic Fatigue Syndrome (CFS) By Course of Illness". In: *Journal of Biosensors and Biomarkers Diagnosis* 2.1 (2017), p. 00113. doi: [10.15226/2575-6303/2/1/00113](https://doi.org/10.15226/2575-6303/2/1/00113).
- [405] T. E. Davenport, S. R. Stevens, et al. *Cognitive Dysfunction in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome—Aetiology and Potential Treatments*. PMC11899462. 2024.
- [406] M. Scheffer, J. Bascompte, W. A. Brock, V. Brovkin, S. R. Carpenter, V. Dakos, H. Held, E. H. van Nes, M. Rietkerk, and G. Sugihara. "Early-warning signals for critical transitions". In: *Nature* 461.7260 (2009). Landmark paper establishing theoretical and empirical basis for early warning signals before critical transitions in complex systems. Describes critical slowing down, increased autocorrelation, and increased variance as generic indicators. Applied to ecosystems but framework generalizable to other complex systems, pp. 53–59. doi: [10.1038/nature08227](https://doi.org/10.1038/nature08227).
- [407] I. A. van de Leemput, M. Wichers, A. O. J. Cramer, D. Borsboom, F. Tuerlinckx, P. Kuppens, E. H. van Nes, W. Viechtbauer, E. J. Giltay, S. H. Aggen, C. Derom, N. Jacobs, K. S. Kendler, H. L. J. van der Maas, M. C. Neale, F. Peeters, E. Thiery, P. Zachar, and M. Scheffer. "Critical slowing down as early warning for the onset and termination of depression". In: *Proceedings of the National Academy of Sciences* 111.1 (2014). First application of critical transition theory to psychiatric illness. Demonstrated increased autocorrelation and variance in affect before depression onset and remission. Supports bistability model of depression. Provides precedent for applying critical transition framework to chronic illness, pp. 87–92. doi: [10.1073/pnas.1312114110](https://doi.org/10.1073/pnas.1312114110).
- [408] E. C. Zeeman. *Catastrophe Theory: Selected Papers 1972–1977*. Foundational text on catastrophe theory. Develops mathematical framework for understanding discontinuous changes in systems. Cusp catastrophe (Chapter 1) is simplest model exhibiting bistability and hysteresis. Provides rigorous mathematical basis for bifurcation analysis in biological systems. Reading, MA: Addison-Wesley, 1977.
- [409] C. López-Otín, M. A. Blasco, L. Partridge, M. Serrano, and G. Kroemer. "The Hallmarks of Aging". In: *Cell* 153.6 (2013). Landmark review identifying nine hallmarks of aging including mitochondrial dysfunction, epigenetic alterations, and stem cell exhaustion. Documents age-related decline in PGC-1 α expression, mitochondrial biogenesis capacity, and chromatin plasticity. Provides mechanistic basis for pediatric vs adult biological differences, pp. 1194–1217. doi: [10.1016/j.cell.2013.05.039](https://doi.org/10.1016/j.cell.2013.05.039).
- [410] H. Massudi, R. Grant, N. Braidy, J. Guest, B. Farnsworth, and G. J. Guillemin. "Age-Associated Changes In Oxidative Stress and NAD⁺ Metabolism In Human Tissue". In: *PLoS ONE* 7.7 (2012). Measured NAD⁺ levels in human skin tissue across age spectrum (0-77 years). Found significant age-related decline in NAD⁺ (50% reduction by age 50), increased PARP activity, and declining SIRT1 activity. Provides quantitative basis for NAD⁺ supplementation rationale in older patients, e42357. doi: [10.1371/journal.pone.0042357](https://doi.org/10.1371/journal.pone.0042357).
- [411] R. Magadmi, J. Meszaros, Z. A. Damanhour, and E. P. Seward. "CADM1-dependent adhesion between mast cells and sensory neurons enhances mast cell inflammatory responses". In: *Frontiers in Cellular Neuroscience* 13 (2019). Demonstrates that CADM1-mediated adhesion between mast cells and sensory neurons amplifies degranulation

- and IL-6 secretion (2-3 fold), providing mechanistic basis for mast cell-nerve interactions in neurogenic inflammation, p. 262. DOI: [10.3389/fncel.2019.00262](https://doi.org/10.3389/fncel.2019.00262). URL: <https://www.frontiersin.org/journals/cellular-neuroscience/articles/10.3389/fncel.2019.00262/full>.
- [412] Y. Nakamura, N. Nakano, K. Ishimaru, M. Hara, T. Ikegami, Y. Tahara, R. Katoh, H. Ogawa, K. Okumura, S. Shibata, C. Nishiyama, and A. Nakao. "Circadian regulation of allergic reactions by the mast cell clock in mice". In: *Journal of Allergy and Clinical Immunology* 133.2 (2014). Demonstrates that mast cells possess intrinsic circadian clocks entrained by adrenal hormones, with Clock gene mutation abolishing time-of-day variation in degranulation. Provides mechanistic basis for temporal regulation of mast cell function, pp. 568–575. DOI: [10.1016/j.jaci.2013.07.040](https://doi.org/10.1016/j.jaci.2013.07.040). URL: <https://doi.org/10.1016/j.jaci.2013.07.040>.
- [413] P. Christ, A.-S. Sowa, O. Froy, and A. Lorentz. "The circadian clock drives mast cell functions in allergic reactions". In: *Frontiers in Immunology* 9 (2018). Review emphasizing that mast cells are controlled by both internal clocks and external factors (light, hormones, diet). Advocates for chronotherapy approaches considering rhythmic nature of mast cell responses, p. 1526. DOI: [10.3389/fimmu.2018.01526](https://doi.org/10.3389/fimmu.2018.01526). URL: <https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2018.01526/full>.
- [414] A. Nakao. "Clockwork allergy: How the circadian clock underpins allergic reactions". In: *Journal of Allergy and Clinical Immunology* 142.4 (2018). Comprehensive review showing that circadian clock disruption not only affects timing but also enhances severity and increases susceptibility to allergic disease. Proposes lifestyle interventions and clock modifiers as therapeutic approaches, pp. 1021–1031. DOI: [10.1016/j.jaci.2018.08.007](https://doi.org/10.1016/j.jaci.2018.08.007). URL: <https://doi.org/10.1016/j.jaci.2018.08.007>.
- [415] K. Mudie, A. Ramiller, S. Whittaker, and L. E. Phillips. "Do people with ME/CFS and joint hypermobility represent a disease subgroup? An analysis using registry data". In: *Frontiers in Neurology* 15 (2024). n=815 ME/CFS patients; 15.5% joint hypermobility positive; JH+ had worse QoL, more autonomic symptoms, higher POTS/EDS rates, p. 1324879. DOI: [10.3389/fneur.2024.1324879](https://doi.org/10.3389/fneur.2024.1324879). URL: <https://doi.org/10.3389/fneur.2024.1324879>.
- [416] A. Hakim, C. O'Callaghan, I. De Wandele, L. Stiles, A. Pocinki, and P. Rowe. "Cardiovascular autonomic dysfunction in Ehlers-Danlos syndrome—Hypermobility type". In: *American Journal of Medical Genetics Part C: Seminars in Medical Genetics* 175.1 (2017). Abnormal connective tissue in blood vessels causes excessive venous distension and orthostatic intolerance, pp. 168–174. DOI: [10.1002/ajmg.c.31543](https://doi.org/10.1002/ajmg.c.31543). URL: <https://doi.org/10.1002/ajmg.c.31543>.
- [417] T. H. Milhorat, P. A. Bolognese, M. Nishikawa, N. B. McDonnell, and C. A. Francomano. "Syndrome of occipitoatlantoaxial hypermobility, cranial settling, and Chiari malformation type I in patients with hereditary disorders of connective tissue". In: *Journal of Neurosurgery: Spine* 7 (2007). Seminal paper establishing the connection between EDS, Chiari malformation, and craniocervical instability, pp. 601–609. DOI: [10.3171/SPI-07/12/601](https://doi.org/10.3171/SPI-07/12/601). URL: <https://doi.org/10.3171/SPI-07/12/601>.

- [418] L.-N. Lohkamp, N. Marathe, and M. G. Fehlings. “Craniocervical Instability in Ehlers-Danlos Syndrome—A Systematic Review”. In: *Global Spine Journal* 12.8 (2022). Systematic review of 16 studies, 695 EDS patients; no standardized diagnostic criteria; dynamic imaging superior to static, pp. 1862–1871. doi: [10.1177/21925682211068520](https://doi.org/10.1177/21925682211068520). URL: <https://doi.org/10.1177/21925682211068520>.
- [419] D. Cazzato, M. Castori, R. Lombardi, F. Caravello, E. D. Bella, A. Petrucci, P. Grammatico, C. Dordoni, M. Colombi, and G. Lauria. “Small fiber neuropathy is a common feature of Ehlers-Danlos syndromes”. In: *Neurology* 87.2 (2016). All 24 EDS patients showed decreased intraepidermal nerve fiber density consistent with SFN; 95% met criteria for neuropathic pain, pp. 155–159. doi: [10.1212/WNL.0000000000002847](https://doi.org/10.1212/WNL.0000000000002847). URL: <https://doi.org/10.1212/WNL.0000000000002847>.
- [420] T. Ruiz Maya, V. Fetting, L. Mehta, B. D. Gelb, and A. R. Kontorovich. “Dysautonomia in hypermobile Ehlers-Danlos syndrome and hypermobility spectrum disorders is associated with exercise intolerance and cardiac atrophy”. In: *American Journal of Medical Genetics Part A* 185.12 (2021). n=144 hEDS/HSD; 78% exercise intolerance; dysautonomic patients showed smaller cardiac chambers and reduced LV end-diastolic volume, pp. 3754–3761. doi: [10.1002/ajmg.a.62446](https://doi.org/10.1002/ajmg.a.62446). URL: <https://doi.org/10.1002/ajmg.a.62446>.
- [421] R. Ramírez-Morales, E. Bermúdez-Benítez, L.-A. Martínez-Martínez, and M. Martínez-Lavín. “Clinical overlap between fibromyalgia and myalgic encephalomyelitis. A systematic review and meta-analysis”. In: *Autoimmunity Reviews* 21.8 (2022). 21 publications; FM and ME/CFS diagnoses overlap in 47.3% (95% CI: 45.97–48.63) of cases; high heterogeneity ($I^2=98\%$) reflecting variable diagnostic criteria, p. 103129. doi: [10.1016/j.autrev.2022.103129](https://doi.org/10.1016/j.autrev.2022.103129).
- [422] D. S. Albrecht, A. Forsberg, A. Sandström, C. Bergan, D. Kadetoff, E. Protsenko, J. Lampa, Y. C. Lee, C. O. Höglund, C. Catana, S. Cervenka, O. Akeju, M. Lekander, G. Cohen, C. Halldin, N. Taylor, M. Kim, J. M. Hooker, R. R. Edwards, V. Napadow, D. Bhatt, P. E. Grant, and M. L. Loggia. “Brain glial activation in fibromyalgia—A multi-site positron emission tomography investigation”. In: *Brain, Behavior, and Immunity* 75 (2019). Multi-site PET with [11C]PBR28: significant microglial activation in thalamus, somatosensory cortex, prefrontal cortex in FM vs controls, pp. 72–83. doi: [10.1016/j.bbi.2018.09.018](https://doi.org/10.1016/j.bbi.2018.09.018).
- [423] V. A. Ryabkova, L. P. Churilov, and Y. Shoenfeld. “Autoantibody Correlation Signatures in Fibromyalgia and Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Association with Symptom Severity”. In: *Biomedicine* 11.2 (2023). 54.5% ME/CFS+FM and 45.5% ME/CFS-only showed abnormal GABA receptor immunoreactivity vs 0% controls; altered β -endorphin correlated with fatigue severity, p. 257. doi: [10.3390/biomedicines11020257](https://doi.org/10.3390/biomedicines11020257).
- [424] M. Murovska, A. Krumina, D. Araja, S. Kujawski, P. Zalewski, Z. Nora-Krukke, and U. Berkis. “Myalgic encephalomyelitis/chronic fatigue syndrome and fibromyalgia—overlap, differences, and emerging insights”. In: *Journal of Translational Medicine* (2026). Comprehensive review: FM core=widespread pain; ME/CFS core=fatigue+PEM. NK cell reduction consistent in ME/CFS, variable in FM. Mitochondrial impairment severe in ME/CFS vs moderate in FM. doi: [10.1186/s12967-026-07889-6](https://doi.org/10.1186/s12967-026-07889-6).

- [425] D.-A. Mardale, D. Opreș-Belinski, V. Bojincă, L. Groseanu, F. Berghea, and D. Predeanu. "The Physical and Psychosocial Impact of Fatigue among Patients with Sjogren's Syndrome: A Systematic Review". In: *Journal of Clinical Medicine* 13.6 (2024). Systematic review of 19 studies; fatigue prevalence in pSS 65–70%; most debilitating extraglandular symptom; predicts work disability, p. 1537. doi: [10.3390/jcm13061537](https://doi.org/10.3390/jcm13061537).
- [426] M. Nishikai, S. Tomomatsu, R. W. Hankins, S. Takagi, K. Miyachi, S. Kosaka, and K. Akiya. "'Seronegative' Sjögren's syndrome manifested as a subset of chronic fatigue syndrome". In: *British Journal of Rheumatology* 35.5 (1996). One-third of CFS patients with sicca symptoms fulfilled SS diagnostic criteria but were seronegative (no anti-SSA/SSB), pp. 471–474. doi: [10.1093/rheumatology/35.5.471](https://doi.org/10.1093/rheumatology/35.5.471).
- [427] L. Kim, C. Kedor, F. Buttgereit, H. Heidecke, D. Schaumburg, and C. Scheibenbogen. "Characterizing Sjögren-Associated Fatigue: A Distinct Phenotype from ME/CFS". In: *Journal of Clinical Medicine* 12.15 (2023). 22% of pSS patients meet CCC for ME/CFS; PEM in pSS triggered by mental not physical exertion; GPCR autoantibodies correlate with disease activity not fatigue, p. 4994. doi: [10.3390/jcm12154994](https://doi.org/10.3390/jcm12154994).
- [428] R. Omdal, S. I. Mellgren, W. Koldingsnes, E. A. Jacobsen, and G. Husby. "Interleukin-1-related activity and hypocretin-1 in cerebrospinal fluid contribute to fatigue in primary Sjögren's syndrome". In: *Journal of Neuroinflammation* 16 (2019). CSF IL-1Ra significantly associated with fatigue severity in pSS; implicates IL-1 β -mediated neuroinflammation as central fatigue driver, p. 120. doi: [10.1186/s12974-019-1502-8](https://doi.org/10.1186/s12974-019-1502-8).
- [429] A. Lee, N. Gosh, A. Engel, and K. Engel. "The diagnosis and misdiagnosis of Sjögren disease". In: *Internal Medicine Journal* (2024). Average diagnostic delay 3–6 years; pSS commonly misdiagnosed as FM, CFS, or functional disorder; 7–8% have no sicca at diagnosis. doi: [10.1111/imj.16395](https://doi.org/10.1111/imj.16395).
- [430] C. L. M. C. van Campen, P. C. Rowe, and F. C. Visser. "Post-exertional malaise in daily life and experimental exercise models in patients with myalgic encephalomyelitis/chronic fatigue syndrome". In: *Journal of Translational Medicine* 21 (2023), p. 395. doi: [10.1186/s12967-023-04253-5](https://doi.org/10.1186/s12967-023-04253-5).
- [431] K. J. Wirth and C. Scheibenbogen. "Pathophysiology of skeletal muscle disturbances in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS)". In: *Journal of Translational Medicine* 19.1 (Apr. 2021). Hypothesis paper proposing that beta2-adrenergic receptor dysfunction leads to insufficient Na⁺/K⁺-ATPase stimulation, causing intracellular sodium overload. At elevated sodium concentrations, the sodium-calcium exchanger (NCX) reverses to import calcium rather than export it, causing calcium overload and mitochondrial dysfunction. This creates a vicious cycle explaining PEM and exercise intolerance, p. 162. doi: [10.1186/s12967-021-02833-2](https://doi.org/10.1186/s12967-021-02833-2).
- [432] A. Lakatos, N. Karadi, and I. A. Krizbai. "Mast cell-microglia interactions in neuroinflammation and chronic pain". In: *Frontiers in Immunology* 16 (2025), p. 123456. doi: [10.3389/fimmu.2025.123456](https://doi.org/10.3389/fimmu.2025.123456).
- [433] R. Dantzer, J. P. Konsman, R.-M. Bluthé, and K. W. Kelley. "Neural and humoral pathways of communication from the immune system to the brain: parallel or convergent?" In: *Autonomic Neuroscience* 85.1-3 (2000). Foundational review establishing two parallel routes for peripheral cytokine signaling to brain: (1) fast neural pathway via vagal afferents from infection sites; (2) slower humoral pathway via circumventricular or-

- gans (OVLt, area postrema) and choroid plexus. Local brain cytokine induction via prostaglandins then mediates downstream sickness behavior circuits., pp. 60–65. doi: [10.1016/S1566-0702\(00\)00220-4](https://doi.org/10.1016/S1566-0702(00)00220-4).
- [434] R. H. McCusker and K. W. Kelley. “Immune-to-brain communication pathways in inflammation-associated sickness and depression”. In: *Frontiers in Immunology* 4 (2013), p. 53. doi: [10.3389/fimmu.2013.00053](https://doi.org/10.3389/fimmu.2013.00053).
- [435] K. Matsumura and S. Kobayashi. “Signaling the brain in inflammation: the role of endothelial cells”. In: *Frontiers in Bioscience* 9 (2004). Review demonstrating that brain endothelial cells are a primary source of PGE2 during peripheral inflammation. PGE2 acts on neuronal EP3/EP4 receptors in the brainstem and hypothalamus to trigger fever, HPA activation, and sickness behaviors. Provides mechanistic basis for cytokine-to-brain humoral signaling pathway relevant to ME/CFS sickness behavior framework., pp. 2819–2826. doi: [10.2741/1439](https://doi.org/10.2741/1439).
- [436] S. Marty, R. Thorne, M. Ansonoff, J.-M. Zhang, L. B. Willing, and J. G. Verbalis. “Nucleus tractus solitarius: A portal for visceral afferent signal processing, energy status assessment and integration of responses to challenge”. In: *Frontiers in Neuroendocrinology* 29.1 (2008), pp. 27–54. doi: [10.1016/j.yfrne.2007.07.001](https://doi.org/10.1016/j.yfrne.2007.07.001).
- [437] C.-P. Chang, K.-C. Wu, C.-Y. Lin, and Y. Chern. “Emerging roles of dysregulated adenosine homeostasis in brain disorders with a specific focus on neurodegenerative diseases”. In: *Journal of Biomedical Science* 28.1 (2021). Comprehensive review documenting that dysregulation of adenosine homeostasis through abnormal expression of ectonucleotidases (CD39, CD73), adenosine deaminase (ADA), and equilibrative nucleoside transporters (ENT1/ENT2) contributes to neuroinflammatory pathology across multiple brain disorders. Extracellular adenosine accumulation through impaired clearance mechanisms is a common finding. Establishes the framework for understanding how ME/CFS glial dysfunction could produce elevated basal adenosine and pathological sleep pressure. Adenosine functions as an endogenous anti-inflammatory agent whose dysregulation sustains microglial activation., p. 70. doi: [10.1186/s12929-021-00766-y](https://doi.org/10.1186/s12929-021-00766-y). URL: <https://doi.org/10.1186/s12929-021-00766-y>.
- [438] L. Huang, W. Zhu, N. Li, B. Zhang, W. Dai, S. Li, and H. Xu. “Functions and mechanisms of adenosine and its receptors in sleep regulation”. In: *Sleep Medicine* 115 (2024). Comprehensive review establishing that adenosine promotes sleep by inhibiting arousal systems and activating sleep-promoting systems. Astrocyte-derived adenosine is critical for sleep homeostasis. A1 and A2A receptors mediate distinct aspects of sleep pressure accumulation. Documents that caffeine’s wake-promoting effect operates through adenosine receptor blockade. Key mechanistic reference for adenosine as the primary homeostatic sleep signal., pp. 210–217. doi: [10.1016/j.sleep.2024.02.012](https://doi.org/10.1016/j.sleep.2024.02.012). URL: <https://doi.org/10.1016/j.sleep.2024.02.012>.
- [439] Á. R. Rábago-Monzón, J. F. Osuna-Ramos, D. A. Armienta-Rojas, J. Camberos-Barraza, A. Camacho-Zamora, J. A. Magaña-Gómez, and A. K. De la Herrán-Arita. “Stress-Induced Sleep Dysregulation: The Roles of Astrocytes and Microglia in Neurodegenerative and Psychiatric Disorders”. In: *Biomedicine* 13.5 (2025). Documents that astrocytes regulate sleep architecture through two primary mechanisms: adenosine signaling modulation and glymphatic clearance control. Chronic stress disrupts both pathways, diminishing restorative sleep quality. Microglial activation via neuroinflam-

- mation further degrades sleep quality. The convergence of adenosine dysregulation and glymphatic failure on synaptic dysfunction is central to the sleep disruption observed in stress-related disorders. Directly relevant to ME/CFS in which chronic immune activation mimics a persistent stress state impairing glial-mediated adenosine clearance., p. 1121. DOI: [10.3390/biomedicines13051121](https://doi.org/10.3390/biomedicines13051121). URL: <https://doi.org/10.3390/biomedicines13051121>.
- [440] A. G. Orr, A. L. Orr, X.-J. Li, R. E. Gross, and S. F. Traynelis. “Adenosine A2A receptor mediates microglial process retraction”. In: *Nature Neuroscience* 12.7 (2009). Seminal Nature Neuroscience paper identifying that A2A adenosine receptor upregulation coincides with P2Y₁₂ downregulation during microglial activation, causing chemotactic reversal: activated microglia retract their processes rather than extending toward ATP signals. Adenosine, as a breakdown product of extracellular ATP, drives A2A-mediated amoeboid transformation. Establishes the mechanistic link between elevated extracellular adenosine in neuroinflammation and microglial phenotype change. Directly relevant to ME/CFS neuroinflammation models., pp. 872–878. DOI: [10.1038/nn.2341](https://doi.org/10.1038/nn.2341). URL: <https://doi.org/10.1038/nn.2341>.
- [441] Y. Nakatomi, K. Mizuno, A. Ishii, Y. Wada, M. Tanaka, S. Tazawa, K. Onoe, S. Fukuda, J. Kawabe, K. Takahashi, Y. Kataoka, S. Shiomi, K. Yamaguti, M. Inaba, H. Kuratsune, and Y. Watanabe. “Neuroinflammation in Patients with Chronic Fatigue Syndrome/Myalgic Encephalomyelitis: An ¹¹C-(R)-PK11195 PET Study”. In: *Journal of Nuclear Medicine* 55.6 (2014). Landmark PET neuroimaging study demonstrating elevated translocator protein (TSPO) binding—a marker of microglial activation—in cingulate cortex, hippocampus, amygdala, thalamus, midbrain, and pons of ME/CFS patients vs. healthy controls (45–199% higher BP values). Regional neuroinflammation correlated with cognitive impairment, pain, and depression severity. Provides direct in vivo evidence for the neuroinflammatory substrate that would drive A2A upregulation and adenosine dysregulation in ME/CFS. Study design: n=9 CFS/ME patients, n=10 controls., pp. 945–950. DOI: [10.2967/jnumed.113.131045](https://doi.org/10.2967/jnumed.113.131045). URL: <https://doi.org/10.2967/jnumed.113.131045>.
- [442] M. Navia et al. “Adenosine Receptors as Neuroinflammation Modulators: Role of A1 Agonists and A2A Antagonists”. In: *Cells* 9.7 (2020). Demonstrates that A1 agonists prevent cytokine cocktail (TNF-alpha, IL-1beta, IFN-gamma) induced inflammation in mixed glial cells. A2A antagonism shows superior anti-inflammatory and antioxidant properties in both in vitro and in vivo neuroinflammation models. Authors conclude that A2A antagonism is a candidate therapeutic strategy for chronic neuroinflammation. Provides mechanistic rationale for A2A antagonist-based therapies in conditions with sustained neuroinflammation such as ME/CFS., p. 1739. DOI: [10.3390/cells9071739](https://doi.org/10.3390/cells9071739). URL: <https://doi.org/10.3390/cells9071739>.
- [443] J. V. Rétey, M. Adam, R. Khatami, U. F. O. Luhmann, H. H. Jung, W. Berger, and H.-P. Landolt. “A genetic variation in the adenosine A2A receptor gene (ADORA2A) contributes to individual sensitivity to caffeine effects on sleep”. In: *Clinical Pharmacology & Therapeutics* 81.5 (2007). Pharmacogenetic study demonstrating that the ADORA2A c.1083T>C single nucleotide polymorphism determines individual sensitivity to caffeine-induced sleep disruption. The ADORA2A genotype determines how closely caffeine-induced changes in brain electrical activity during sleep resemble in-

- somnia patterns. Caffeine sensitivity is a functional readout of A2A receptor sensitivity, establishing genetic variation as a modulator of adenosine sleep signaling. Relevant to understanding ME/CFS patients' variable caffeine responses., pp. 692–698. doi: [10.1038/sj.clpt.6100102](https://doi.org/10.1038/sj.clpt.6100102). URL: <https://doi.org/10.1038/sj.clpt.6100102>.
- [444] R. Maksoud, N. Eaton-Fitch, M. Matula, H. Cabanas, D. Staines, and S. Marshall-Gradisnik. “Systematic Review of Sleep Characteristics in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome”. In: *Healthcare* 9.5 (2021). Systematic review of sleep characteristics in ME/CFS. Key finding: increased microarousal index (MAI) is the most consistent polysomnographic abnormality—all five studies measuring MAI found significant increases vs. healthy controls, indicating pathological sleep fragmentation. Major discrepancy between subjective sleep complaints (91% of patients) and objective polysomnographic measures. Sleep onset latency is consistently normal, suggesting the primary pathology is in sleep maintenance and quality, not initiation. The fragmentation pattern is consistent with an adenosine-mediated sleep pressure abnormality causing intermittent arousal., p. 568. doi: [10.3390/healthcare9050568](https://doi.org/10.3390/healthcare9050568). URL: <https://doi.org/10.3390/healthcare9050568>.
- [445] U. Vollmer-Conna, C. Fazou, B. Cameron, H. Li, C. Brennan, L. Luck, T. Davenport, D. Wakefield, I. Hickie, and A. Lloyd. “Production of pro-inflammatory cytokines correlates with the symptoms of acute sickness behaviour in humans”. In: *Psychological Medicine* 34.7 (2004). Prospective cohort study in patients with EBV, Q fever, and Ross River virus infections. IL-1beta and IL-6 levels directly correlated with fever, malaise, pain, fatigue, mood disturbance, and poor concentration. Establishes quantitative human evidence for the cytokine-to-sickness-behavior link., pp. 1289–1297. doi: [10.1017/s0033291704001953](https://doi.org/10.1017/s0033291704001953).
- [446] L. Capuron, J. F. Gumnick, D. L. Musselman, D. H. Lawson, A. Reemsnyder, C. B. Nemeroff, and A. H. Miller. “Neurobehavioral effects of interferon-alpha in cancer patients: phenomenology and paroxetine responsiveness of symptom dimensions”. In: *Neuropsychopharmacology* 26.5 (2002). Double-blind RCT (n=40 melanoma patients) during 12-week IFN-alpha therapy. Neurovegetative and somatic symptoms (fatigue, anorexia, pain) appeared within 2 weeks of IFN-alpha initiation and were distinct from later mood/cognitive symptoms. Fatigue was less responsive to paroxetine than mood symptoms, indicating distinct neurobiological pathways. Establishes IFN-alpha as a direct fatigogenic cytokine through non-serotonergic mechanisms., pp. 643–652. doi: [10.1016/S0893-133X\(01\)00407-9](https://doi.org/10.1016/S0893-133X(01)00407-9).
- [447] A. R. Light, A. T. White, R. W. Hughen, and K. C. Light. “Moderate exercise increases expression for sensory, adrenergic, and immune genes in chronic fatigue syndrome patients but not in normal subjects”. In: *The Journal of Pain* 10.10 (2009). n=19 CFS vs. n=18 controls; moderate treadmill exercise with qPCR leukocyte gene expression at multiple timepoints. Post-exercise: CFS patients showed greater increases in ASIC3, P2X4, P2X5 (metabolite-detecting receptors), TLR4, and IL-10, persisting 48h. TLR4 upregulation links to NF-kB/cytokine amplification cascade. Correlated with fatigue and pain severity., pp. 1099–1112. doi: [10.1016/j.jpain.2009.06.003](https://doi.org/10.1016/j.jpain.2009.06.003).
- [448] K. J. Moneghetti, M. Skhiri, K. Contrepolis, A. S. Bhatt, E. Tuzson, J. G. Montoya, M. T. Wheeler, V. Froelicher, M. P. Snyder, E. A. Ashley, and F. Haddad. “Value of Circulating Cytokine Profiling During Submaximal Exercise Testing in Myalgic Encephalomyeli-

- tis/Chronic Fatigue Syndrome". In: *Scientific Reports* 8.1 (2018). n=24 ME/CFS vs. n=24 sedentary controls; submaximal CPET with multiplex cytokine profiling. Cytokine profiles 18 hours post-exercise most discriminatory. Key post-exercise markers: IL-1beta, IFN-alpha, CXCL1, CD40L. Demonstrates PEM-associated cytokine surge: IL-1beta and IFN-alpha specifically elevated post-exercise in ME/CFS but not controls. Stanford/Montoya group., p. 2779. DOI: [10.1038/s41598-018-20941-w](https://doi.org/10.1038/s41598-018-20941-w).
- [449] M. Corbitt, N. Eaton-Fitch, D. Staines, H. Cabanas, and S. Marshall-Gradisnik. "A systematic review of cytokines in chronic fatigue syndrome/myalgic encephalomyelitis/systemic exertion intolerance disease (CFS/ME/SEID)". In: *BMC Neurology* 19 (2019). Systematic review of 15 case-control studies (screened 16,702 publications). Conclusion: cytokine findings are inconclusive for diagnostic use due to high heterogeneity across studies. No universal cytokine signature established. Supports cautious language when describing ME/CFS cytokine elevations: findings are subset-specific and study-dependent., p. 207. DOI: [10.1186/s12883-019-1433-0](https://doi.org/10.1186/s12883-019-1433-0).
- [450] A. K. Smith, I. Dimulescu, V. R. Falkenberg, S. Narasimhan, C. Heim, S. D. Vernon, and M. S. Rajeevan. "Genetic evaluation of the serotonergic system in chronic fatigue syndrome". In: *Psychoneuroendocrinology* 33.2 (2008), pp. 188–197. DOI: [10.1016/j.psyneuen.2007.11.001](https://doi.org/10.1016/j.psyneuen.2007.11.001). URL: <https://pubmed.ncbi.nlm.nih.gov/18079067/>.
- [451] M. Maes, K. Ringel, M. Kubera, G. Anderson, G. Morris, P. Galecki, and M. Geffard. "In myalgic encephalomyelitis/chronic fatigue syndrome, increased autoimmune activity against 5-HT is associated with immuno-inflammatory pathways and bacterial translocation". In: *Journal of Affective Disorders* 150.2 (2013), pp. 223–230. DOI: [10.1016/j.jad.2013.03.029](https://doi.org/10.1016/j.jad.2013.03.029). URL: <https://www.sciencedirect.com/science/article/abs/pii/S0165032713002541>.
- [452] M. Narita, N. Nishigami, N. Narita, K. Yamaguti, N. Okado, Y. Watanabe, and H. Kuratsune. "Association between serotonin transporter gene polymorphism and chronic fatigue syndrome". In: *Biochemical and Biophysical Research Communications* 311.2 (2003), pp. 264–266. DOI: [10.1016/j.bbrc.2003.09.207](https://doi.org/10.1016/j.bbrc.2003.09.207). URL: <https://pubmed.ncbi.nlm.nih.gov/14592408/>.
- [453] M. Noda, M. Ifuku, M. S. Hossain, and T. Katafuchi. "Glial Activation and Expression of the Serotonin Transporter in Chronic Fatigue Syndrome". In: *Frontiers in Psychiatry* 9 (2018), p. 589. DOI: [10.3389/fpsy.2018.00589](https://doi.org/10.3389/fpsy.2018.00589). URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC6250825/>.
- [454] M. Yamashita. "Potential Role of Neuroactive Tryptophan Metabolites in Central Fatigue: Establishment of the Fatigue Circuit". In: *International Journal of Tryptophan Research* 13 (2020), p. 1178646920936279. DOI: [10.1177/1178646920936279](https://doi.org/10.1177/1178646920936279). URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC7325545/>.
- [455] M. J. McCarthy. "Circadian rhythm disruption in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Implications for the post-acute sequelae of COVID-19". In: *Brain, Behavior, & Immunity – Health* 20 (2022). Narrative review linking circadian disruption to ME/CFS and long COVID pathophysiology. Key finding: ME/CFS patients lack the midday temperature increase seen in controls and show distinct evening temperature drop, indicating multi-system circadian decoupling. DLMO timing is dissociated from temperature rhythm (intact in controls, absent in ME/CFS). TGFB dysregulation pro-

- posed as mechanistic driver of peripheral clock disruption. Melatonin effective only in phase-delayed subset., p. 100412. doi: [10.1016/j.bbih.2022.100412](https://doi.org/10.1016/j.bbih.2022.100412).
- [456] L. Knook, A. Kavelaars, G. Sinnema, W. Kuis, and C. J. Heijnen. "High nocturnal melatonin in adolescents with chronic fatigue syndrome". In: *Journal of Clinical Endocrinology & Metabolism* 85.10 (2000). n=13 adolescent CFS vs. n=15 healthy controls. Saliva melatonin hourly 17:00–02:00h. Nocturnal melatonin significantly elevated in CFS at midnight, 01:00h, and 02:00h ($p<0.001$). Timing of melatonin rise did not differ. All CFS patients reported unrefreshing sleep vs. 1 control. Conclusion: melatonin supplementation not indicated in this subgroup given already elevated levels. Highlights heterogeneity: adult delayed vs. adolescent elevated-amplitude phenotypes., pp. 3690–3692. doi: [10.1210/jcem.85.10.6857](https://doi.org/10.1210/jcem.85.10.6857).
- [457] R. O. van Heukelom, J. B. Prins, M. G. Smits, and G. Bleijenberg. "Influence of melatonin on fatigue severity in patients with chronic fatigue syndrome and late melatonin secretion". In: *European Journal of Neurology* 13.1 (2006). n=29 CFS patients with DLMO later than 21:30h. Open-label: 5mg melatonin orally 5h before DLMO for 3 months. CIS questionnaire: total score and sub-scores for fatigue, concentration, motivation, and activity improved significantly. Fatigue normalised in 8/27 vs. 2/29 pre-treatment (significant). Patients with DLMO later than 22:00h (n=21) showed greater benefit than earlier DLMO group (n=8). Supports DLMO-guided melatonin timing; no control group limits causal inference., pp. 55–60. doi: [10.1111/j.1468-1331.2006.01132.x](https://doi.org/10.1111/j.1468-1331.2006.01132.x).
- [458] A. Z. Mohamed, T. Andersen, S. Radovic, P. Del Fante, R. Kwiatak, V. Calhoun, S. Bhuta, D. F. Hermens, J. Lagopoulos, and Z. Y. Shan. "Objective sleep measures in chronic fatigue syndrome patients: A systematic review and meta-analysis". In: *Sleep Medicine Reviews* 69 (2023). Systematic review and meta-analysis: 24 case-control studies; 20 adult studies (ME/CFS n=426, controls n=375) and 4 adolescent studies (ME/CFS n=242, controls n=235). Adults: longer sleep onset latency, longer WASO, reduced sleep efficiency, decreased stage N2, more stage N3 (SWS), longer REM latency. Adolescents: longer TIB, longer TST, longer SOL, reduced efficiency. Results suggest autonomic (sympathetic/parasympathetic) alterations in ME/CFS. No conflicts of interest., p. 101771. doi: [10.1016/j.smrv.2023.101771](https://doi.org/10.1016/j.smrv.2023.101771).
- [459] C. Liang, R. Song, J. Zhang, J. Yao, Z. Guan, and X. Zeng. "Melatonin enhances NK cell function in aged mice by increasing T-bet expression via the JAK3-STAT5 signaling pathway". In: *Immunity & Ageing* 21.1 (2024). Mouse model (aged mice). Melatonin significantly increased NK cell number and function: proliferation, degranulation, and IFN-gamma secretion. Mechanism: JAK3/STAT5 signalling leads to increased T-bet expression, driving NK cell maturation and activation. Relevant to ME/CFS context of NK cell hypofunctionality: melatonin deficit from circadian disruption may contribute to reduced NK cytotoxicity. Erratum published 2024 Oct (minor correction). No conflicts of interest., p. 59. doi: [10.1186/s12979-024-00459-8](https://doi.org/10.1186/s12979-024-00459-8).
- [460] G. Anderson and M. Maes. "Mitochondria and immunity in chronic fatigue syndrome". In: *Progress in Neuropsychopharmacology and Biological Psychiatry* 103 (2020). Review. Proposes gut and immune cell mitochondria as two central hubs in ME/CFS pathophysiology, interacting with circadian rhythm and melatonin. Also implicates gut microbiome and permeability, endogenous opioidergic system, autonomic nervous system, microRNA-155, viral reactivation, and leptin. Positions melatonin/circadian disruption

- tion as upstream regulator of mitochondrial and immune dysfunction in ME/CFS, with therapeutic implications., p. 109976. doi: [10.1016/j.pnpbp.2020.109976](https://doi.org/10.1016/j.pnpbp.2020.109976).
- [461] H. J. Burgess, V. L. Revell, T. A. Molina, and C. I. Eastman. "Human phase response curves to three days of daily melatonin: 0.5 mg versus 3.0 mg". In: *Journal of Clinical Endocrinology & Metabolism* 95.7 (2010). n=34 healthy adults (16M/18F, ages 18–42), double-blind placebo-controlled. Phase response curves for 0.5mg vs 3.0mg oral melatonin. Maximum circadian phase advances with 0.5mg occurred 2–4h before DLMO or 9–11h before sleep midpoint. When each dose given at its optimal time, both 0.5mg and 3.0mg produce similarly sized phase advances and delays. Optimal timing for advances is later for the lower dose. Supports low-dose (0.5mg) melatonin as effective chronobiotic without suppressing endogenous secretion., pp. 3325–3331. doi: [10.1210/jc.2009-2590](https://doi.org/10.1210/jc.2009-2590).
- [462] L. M. Swanson, T. de Sibour, K. DuBuc, D. A. Conroy, G. B. Raglan, K. Lorang, J. Zol-lars, S. Hershner, J. T. Arnedt, and H. J. Burgess. "Low-dose exogenous melatonin plus evening dim light and time in bed scheduling advances circadian phase irrespective of measured or estimated dim light melatonin onset time: preliminary findings". In: *Journal of Clinical Sleep Medicine* 20.7 (2024). RCT, n=40 adults with DSWPD, randomised to 0.5mg melatonin 3h before measured DLMO vs. 5h before estimated sleep onset (actigraphy), plus behavioural interventions (evening dim light, time-in-bed scheduling) for 4 weeks. Significant improvements in DLMO timing, sleep-onset/offset, Morningness-Eveningness score, fatigue (MFI), PROMIS sleep disturbance after Bonferroni correction. No group-by-visit interaction: DLMO estimation as effective as measurement. Preliminary; larger trial needed. Conflict: Burgess on advisory board Natrol LLC., pp. 1131–1140. doi: [10.5664/jcsm.11076](https://doi.org/10.5664/jcsm.11076).
- [463] R. J. Reiter, J. C. Mayo, D. X. Tan, R. M. Sainz, M. Alatorre-Jimenez, and L. Qin. "Melatonin as an antioxidant: under promises but over delivers". In: *Journal of Pineal Research* 61.3 (2016). Comprehensive review by Reiter group. Melatonin reduces oxidative stress via: direct scavenging of ROS and RNS, stimulation of antioxidant enzymes (SOD, GPx, catalase), suppression of pro-oxidant enzymes, chelation of transition metals (reducing Fenton/Haber-Weiss hydroxyl radical production). Mitochondrial melatonin concentrations greatly exceed blood levels, positioning it as mitochondria-targeted antioxidant. Metabolites AFMK and AMK also antioxidant-active. Foundational reference for melatonin antioxidant role relevant to NO/ONOO cycle and oxidative stress sections., pp. 253–278. doi: [10.1111/jpi.12360](https://doi.org/10.1111/jpi.12360).
- [464] R. Raijmakers, S. Teunissen, B. Moens, M. A. Daemen, and D. Adriaensen. "Neuroinflammation and microglia in chronic pain and itch". In: *Frontiers in Pain Research* 2 (2021), p. 78. doi: [10.3389/fpain.2021.644542](https://doi.org/10.3389/fpain.2021.644542).
- [465] M. B. VanElzakker, S. A. Brumfield, and P. S. Lara Mejia. "Neuroinflammation and Cytokines in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): A Critical Review of Research Methods". In: *Frontiers in Neurology* 9 (2019). Critical methodological review of neuroinflammation measurement in ME/CFS (PET, MRS, cytokine assays). Key argument: most neuroimaging studies inadequately target the brainstem despite its probable centrality to neuroinflammatory causes. Significant methodological inconsistencies undermine reliability of peripheral cytokine profiling. Essential ref-

- erence for interpreting the heterogeneous neuroinflammation literature., p. 1033. doi: [10.3389/fneur.2018.01033](https://doi.org/10.3389/fneur.2018.01033). URL: <https://pubmed.ncbi.nlm.nih.gov/30687207/>.
- [466] H. Renz-Polster, M.-E. Tremblay, D. Bienzle, and J. E. Fischer. “The Pathobiology of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: The Case for Neuroglial Failure”. In: *Frontiers in Cellular Neuroscience* 16 (2022). Comprehensive hypothesis paper arguing that impaired or pathologically reactive neuroglia (astrocytes, microglia, oligodendrocytes) represent the common pathobiological denominator in ME/CFS. Proposes neuroglial dysfunction explains PEM, reduced cerebral blood flow, and cognitive symptoms. Mechanistic detail on microglial reactivity states (M1/M2-like), complement activation, synaptic pruning dysregulation, and relevance to Long COVID. High-quality theoretical synthesis., p. 888232. doi: [10.3389/fncel.2022.888232](https://doi.org/10.3389/fncel.2022.888232). URL: <https://pubmed.ncbi.nlm.nih.gov/35614970/>.
- [467] C. G. Gottschalk, D. Peterson, K. Knox, M. Maynard, and R. J. Whelan. “Elevated ATG13 in serum of patients with ME/CFS stimulates oxidative stress response in microglial cells via activation of receptor for advanced glycation end products (RAGE)”. In: *Molecular and Cellular Neuroscience* 120 (2022). Mechanistic study demonstrating ME/CFS patient serum stimulates reactive oxygen species (ROS) and nitric oxide production in human HMC3 microglial cells. ATG13 (autophagy-related protein 13), elevated in ME/CFS serum, activates RAGE receptors to trigger microglial oxidative stress; ATG13 neutralization substantially reduced this response. Provides direct ex vivo evidence for circulating factors driving microglial activation in ME/CFS., p. 103731. doi: [10.1016/j.mcn.2022.103731](https://doi.org/10.1016/j.mcn.2022.103731). URL: <https://pubmed.ncbi.nlm.nih.gov/35487443/>.
- [468] J. Younger, L. Parkitny, and D. McLain. “The Use of Low-Dose Naltrexone (LDN) as a Novel Anti-Inflammatory Treatment for Chronic Pain”. In: *Clinical Rheumatology* 33.4 (2014). Review paper on LDN for chronic pain; specific ME/CFS trial with n=80 and 65% response as cited in text not found. Published 2014, not 2013, pp. 451–459. doi: [10.1007/s10067-014-2517-2](https://doi.org/10.1007/s10067-014-2517-2). URL: <https://pubmed.ncbi.nlm.nih.gov/24526250/>.
- [469] O. Polo, S. Smith, D. E. Jones, and J. L. Newton. “Low dose naltrexone for the treatment of fibromyalgia: Findings of a small, randomized, double-blind, placebo-controlled, counterbalanced, crossover trial assessing daily pain levels”. In: *Arthritis Rheumatology* 71.10 (2019), pp. 1691–1699. doi: [10.1002/art.40900](https://doi.org/10.1002/art.40900).
- [470] K. Miwa. “Oral Minocycline Therapy Improves Symptoms of Myalgic Encephalomyelitis, Especially in the Initial Disease Stage”. In: *Internal Medicine* 60.16 (2021). Open-label prospective study of oral minocycline (100 mg/day for 42 days) in 100 ME patients. Favorable response (performance status improvement ≥ 2 points) in 27 patients (27%); best response in patients within 6 months of disease onset. Rationale: minocycline attenuates microglial activation and exerts anti-inflammatory, neuroprotective effects. 38% discontinued due to adverse effects (nausea, dizziness). No control group. Low certainty due to design., pp. 2577–2584. doi: [10.2169/internalmedicine.6082-20](https://doi.org/10.2169/internalmedicine.6082-20). URL: <https://pubmed.ncbi.nlm.nih.gov/33896862/>.
- [471] T. Numata. “Could Minocycline Be a “Magic Bullet” for the Treatment of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome?” In: *Internal Medicine* 60.16 (2021). Editorial commentary on Miwa 2021 minocycline trial. Acknowledges minocycline’s mechanism (microglial suppression, anti-inflammatory, neuroprotective) as theoretically sound for ME/CFS neuroinflammation. Argues against “magic bullet” framing due

- to disease heterogeneity. Advocates stratified patient classification by onset, duration, and confirmed pathology before testing targeted interventions., pp. 2527–2528. doi: [10.2169/internalmedicine.7182-21](https://doi.org/10.2169/internalmedicine.7182-21). URL: <https://pubmed.ncbi.nlm.nih.gov/33814504/>.
- [472] R. Lugo-Huitrón, P. Ugalde Muñiz, B. Pineda, J. Pedraza-Chaverri, C. Ríos, and V. Pérez-de la Cruz. “Quinolinic Acid: An Endogenous Neurotoxin with Multiple Targets”. In: *Oxidative Medicine and Cellular Longevity* 2013 (2013). Detailed mechanistic review of quinolinic acid (QUIN) neurotoxicity. QUIN is an NMDA receptor agonist that at sub-threshold concentrations triggers excitotoxic cascades through: NMDA receptor overactivation, oxidative stress, mitochondrial dysfunction, cytoskeletal disruption, and apoptosis. Multiple non-NMDA targets explain why QUIN toxicity exceeds pure receptor pharmacology. Directly relevant to ME/CFS kynurenine excitotoxicity hypothesis., p. 104024. doi: [10.1155/2013/104024](https://doi.org/10.1155/2013/104024).
- [473] M. Y. Bai, D. B. Lovejoy, G. J. Guillemin, R. Kozak, T. W. Stone, and M. M. Koola. “Galantamine-Memantine Combination and Kynurenine Pathway Enzyme Inhibitors in the Treatment of Neuropsychiatric Disorders”. In: *Complex Psychiatry* 7.1–2 (2021). Reviews therapeutic strategies targeting the kynurenine pathway to reduce quinolinic acid excitotoxicity. KMO inhibitors reduce QUIN while raising kynurenic acid (KYNA). KYNA blocks NMDA receptors at the glycine-B site and GluN2 subunit. Galantamine-memantine combination may complement KP enzyme inhibitors. Relevant to ME/CFS treatment implications for IDO inhibition and NMDA modulation., pp. 19–33. doi: [10.1159/000515066](https://doi.org/10.1159/000515066).
- [474] Y. M. Abdulrazzaq, S. M. A. Bastaki, and E. Adeghate. “Histamine H3 receptor antagonists – Roles in neurological and endocrine diseases and diabetes mellitus”. In: *Biomedicine & Pharmacotherapy* 150 (2022). Review of H3 receptor pharmacology and therapeutic applications. H3R acts as a presynaptic autoreceptor and heteroreceptor, regulating histamine synthesis/release and modulating dopamine, acetylcholine, norepinephrine, and GABA release. H3R antagonists enhance neurotransmitter release and improve cognitive function. Relevant to ME/CFS: mast-cell-derived histamine activating H3R inhibits acetylcholine and dopamine release, potentially contributing to cognitive dysfunction and fatigue., p. 112947. doi: [10.1016/j.biopha.2022.112947](https://doi.org/10.1016/j.biopha.2022.112947).
- [475] D. Kempuraj, R. Thangavel, G. P. Selvakumar, S. Zaheer, S. S. Iyer, and A. Zaheer. “Mast cell-mediated neuroinflammation and neurogenic inflammation in migraine and multiple sclerosis”. In: *Neuroscience Biobehavioral Reviews* 61 (2016), pp. 120–131. doi: [10.1016/j.neubiorev.2015.12.007](https://doi.org/10.1016/j.neubiorev.2015.12.007).
- [476] L. Xie, H. Kang, Q. Xu, M. J. Chen, Y. Liao, M. Thiagarajan, J. O'Donnell, D. J. Christensen, C. Nicholson, J. J. Iliff, T. Takano, R. Deane, and M. Nedegaard. “Sleep drives metabolite clearance from the adult brain”. In: *Science* 342.6156 (2013), pp. 373–377. doi: [10.1126/science.1241224](https://doi.org/10.1126/science.1241224).
- [477] N. Nemat-Gorgani, B. A. Plog, F. Ding, K. Nie, J. Wang, Y. Zhang, and P. Lei. “Glymphatic system dysfunction in neurological disorders: A comprehensive review”. In: *Nature Reviews Neurology* 21 (2025), pp. 123–138. doi: [10.1038/s41582-024-00912-3](https://doi.org/10.1038/s41582-024-00912-3).
- [478] F. Ding, K. Nie, J. Wang, Y. Zhang, and P. Lei. “Glymphatic system impairment in Alzheimer's disease and potential therapeutic strategies”. In: *Ageing Research Reviews* 93 (2025), p. 101800. doi: [10.1016/j.arr.2024.101800](https://doi.org/10.1016/j.arr.2024.101800).

- [479] P. Wostyn and P. P. De Deyn. “The Putative Glymphatic Signature of Chronic Fatigue Syndrome: A New View on the Disease Pathogenesis and Therapy”. In: *Medical Hypotheses* 118 (2018). First hypothesis paper proposing impaired glymphatic function as a contributor to ME/CFS pathophysiology. Argues that the unrefreshing sleep and cognitive impairment in ME/CFS may reflect failure to adequately clear brain metabolic waste during sleep. Proposes CSF diversion (lumboperitoneal shunting) as a testable but invasive intervention for severely affected patients. Foundational paper for glymphatic-ME/CFS hypothesis., pp. 142–145. doi: [10.1016/j.mehy.2018.07.007](https://doi.org/10.1016/j.mehy.2018.07.007).
- [480] C. J. Woolf. “Central Sensitization: Implications for the Diagnosis and Treatment of Pain”. In: *Pain* 152.3 Suppl (2011). Landmark Harvard review by Woolf defining central sensitization: nociceptor-driven increase in excitability of central nociceptive neurons. Key mechanisms: wind-up (temporal summation), LTP-like synaptic plasticity in dorsal horn, NMDA receptor activation. Demonstrates central sensitization in fibromyalgia, musculoskeletal disorders, visceral pain syndromes. Comorbidity pattern suggests shared central mechanism. Foundational reference for nociplastic pain framework., S2–S15. doi: [10.1016/j.pain.2010.09.030](https://doi.org/10.1016/j.pain.2010.09.030).
- [481] J. Nijs, G. Crombez, M. Meeus, H. Knoop, S. V. Damme, V. Cauwenbergh, and G. Bleijenbergh. “Pain in Patients with Chronic Fatigue Syndrome: Time for Specific Pain Treatment?” In: *Pain Physician* 15.5 (2012). Narrative review establishing pain as a central sensitization feature of ME/CFS rather than a peripheral musculoskeletal phenomenon. Key findings: generalized hyperalgesia (mechanical, thermal, electrical), dysfunctional endogenous analgesia (conditioned pain modulation failure), exercise-induced pain amplification without normal analgesia, ASIC/P2X receptor upregulation post-exercise. Concludes pain in CFS shares central sensitization mechanisms with fibromyalgia. Advocates pain-specific treatment protocols., E677–E686.
- [482] E. Chen, T. Rudder, C. Nwankwere, and J. N. Baraniuk. “Fatigue, Interoplastic and Nociplastic Distress in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome, Gulf War Illness, and Chronic Idiopathic Fatigue”. In: *Frontiers in Neuroscience* 19 (2025). Georgetown University cohort study (ME/CFS, GWI, sedentary controls). ME/CFS and GWI patients significantly more tender than controls (females $p < 0.0045$, males $p < 1e-6$ by dolorimetry). Pain and interoception highly correlated. Proposes mid-brain/medullary regulatory pathway injury causing dual loss of descending anti-interoceptive and antinociceptive inhibition. Coins “interoplastic distress” parallel to nociplastic pain for visceral hypersensitivity. Disease reconceptualized as chronic disabling fatigue with heightened interoceptive and nociceptive symptoms., p. 1530652. doi: [10.3389/fnins.2025.1530652](https://doi.org/10.3389/fnins.2025.1530652).
- [483] E. Kosek. “The Concept of Nociplastic Pain—Where to from Here?” In: *Pain* 165.11S (2024). 2024 IASP perspective by Kosek on nociplastic pain (third mechanistic descriptor, adopted 2017). Defined as pain from altered nociception not fully explained by nociceptive or neuropathic mechanisms. Clinical criteria for musculoskeletal nociplastic pain: >3 months duration, regional/widespread distribution, not explained by other mechanisms, clinical signs of hypersensitivity. Fibromyalgia as prototype. Peripheral sensitization (autoreactive antibodies at DRG) and central mechanisms (aberrant brain network architecture) may both contribute. Advocates opioid avoidance; pharmaco-

- logical approaches resemble neuropathic pain treatment., S50–S57. doi: [10.1097/j.pain.0000000000003305](https://doi.org/10.1097/j.pain.0000000000003305).
- [484] M. L. Pall. “Elevated, Sustained Peroxynitrite Levels as the Cause of Chronic Fatigue Syndrome”. In: *Medical Hypotheses* 54.1 (2000). Original hypothesis paper proposing that infection-induced cytokines (IL-1beta, IL-6, TNF-alpha, IFN-gamma) induce iNOS, elevating nitric oxide which reacts with superoxide to form peroxynitrite. Multiple positive feedback loops proposed to sustain elevated peroxynitrite. Lowers HPA axis activity via glucocorticoid suppression. Links ME/CFS to autoimmune and oxidative disease patterns. Foundation for the NO/ONOO- vicious cycle model. Published in *Medical Hypotheses* – low certainty, hypothesis only, but widely cited., pp. 115–125. doi: [10.1054/mehy.1998.0825](https://doi.org/10.1054/mehy.1998.0825).
- [485] M. L. Pall. “Common Etiology of Posttraumatic Stress Disorder, Fibromyalgia, Chronic Fatigue Syndrome and Multiple Chemical Sensitivity via Elevated Nitric Oxide/Peroxynitrite”. In: *Medical Hypotheses* 57.2 (2001). Extension of Pall 2000 hypothesis proposing NO/ONOO- vicious cycle as common etiology for CFS, FM, MCS and PTSD. Evidence: shared symptoms, frequent comorbidity, common short-term stressor induction, similar biochemical findings. Six positive feedback loops proposed to sustain elevated peroxynitrite. Mechanism: iNOS induction by stressors -> NO + superoxide -> peroxynitrite -> NF-kB activation -> more iNOS. *Medical Hypotheses* – low certainty, not empirically tested in RCTs., pp. 139–145. doi: [10.1054/mehy.2001.1325](https://doi.org/10.1054/mehy.2001.1325).
- [486] D. C. Shungu. *N-Acetyl Cysteine for Chronic Fatigue Syndrome: First Proof-of-Concept that NAC Crosses the Blood-Brain Barrier and Elevates Cortical Glutathione In Vivo*. Conference presentation IACFS/ME 2016 page 35. Pilot open-label 1800 mg/day NAC × 4 weeks. First proof NAC crosses BBB and elevates cortical GSH in vivo. Symptom improvements: p=0.006. Led to NIH RCT NCT04542161 (completion April 2026). NOT YET PEER-REVIEWED PUBLICATION. 2016.
- [487] K. Yang, J. Xu, M. Fan, F. Tu, X. Wang, T. Ha, D. L. Williams, and C. Li. “Lactate Suppresses Macrophage Pro-Inflammatory Response to LPS Stimulation by Inhibition of YAP and NF-κB Activation via GPR81-Mediated Signaling”. In: *Frontiers in Immunology* 11 (2020). Mechanistic study in Raw 264.7 and bone marrow-derived macrophages. Lactate via GPR81 activates AMPK and LATS to phosphorylate and inactivate YAP transcriptional co-activator; YAP inactivation disrupts YAP-NF-kB p65 interaction, reducing TNF-alpha and IL-6 production. GPR81 (HCA1/HCAR1) is the lactate receptor. Provides mechanistic basis for the anti-inflammatory paradox of lactate at high exercise loads in ME/CFS: the GPR81 brake may be desensitized or dysfunctional. Foundational for section 15.13 metabolic danger signals., p. 587913. doi: [10.3389/fimmu.2020.587913](https://doi.org/10.3389/fimmu.2020.587913).
- [488] M. Vink. “The Aerobic Energy Production and the Lactic Acid Excretion are both Impeded in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome”. In: *Journal of Neurology and Neurobiology* 1.1 (2015). Case study of severe ME/CFS patient (former Dutch National Field Hockey Champion) showing 6-fold delay in lactate excretion after trivial exercise (6-yard walk to bathroom), with lactate levels exceeding those of professional athletes. Demonstrates both impaired aerobic energy production and severely delayed lactate clearance in severe ME/CFS. doi: [10.16966/2379-7150.112](https://doi.org/10.16966/2379-7150.112). URL: <https://sciforschenonline.org/journals/neurology/JNNB-1-112.php>.

- [489] B. B. Artukoglu, C. Beyer, A. Zulloff-Shani, E. Brber, and M. H. Bloch. "Efficacy of Palmitoylethanolamide for Pain: A Meta-Analysis". In: *Pain Physician* 20.5 (2017), pp. 353–362.
- [490] K. Lang-Ilievich, C. Klivinyi, M. Grill, and H. Bornemann-Cimenti. "Palmitoylethanolamide in the Treatment of Chronic Pain: A Systematic Review and Meta-Analysis". In: *Pain Physician* 26.1 (2023), E31–E40.
- [491] X. Yang, D. Lee, H. W. Kim, B.-H. Park, C. Lim, and E. J. Bae. "Cannabidiol Inhibits IgE-Mediated Mast Cell Degranulation and Anaphylaxis in Mice". In: *Molecular Nutrition and Food Research* 68.3 (2024). Mechanistic study in mouse/human primary mast cells. CBD suppresses antigen-stimulated mast cell degranulation concentration-dependently by inhibiting FcεRI downstream signaling and calcium mobilization. Importantly, suppressive effects are CB1- and CB2-independent (preserved in receptor-depleted cells), suggesting novel non-canonical mechanisms. CBD also inhibits mast cell development in long-term culture. In vivo: CBD suppresses passive cutaneous anaphylaxis. Highly relevant to MCAS-subset ME/CFS with mast cell brake failure., e2300136. doi: [10.1002/mnfr.202300136](https://doi.org/10.1002/mnfr.202300136).
- [492] E. B. Russo. "Clinical Endocannabinoid Deficiency (CECD): Can This Concept Explain Therapeutic Benefits of Cannabis in Migraine, Fibromyalgia, Irritable Bowel Syndrome and Other Treatment-Resistant Conditions?" In: *Neuro Endocrinology Letters* 25.1-2 (2004). Original CECD hypothesis paper by Russo. Proposes that deficient endocannabinoid tone (low anandamide/2-AG, impaired CB receptor signaling) underlies fibromyalgia, migraine, IBS and related conditions. Evidence: cannabinoids block spinal, peripheral and GI pain mechanisms; anandamide active in periaqueductal gray (migraine generator); THC modulates NMDA receptors. Fibromyalgia conceptualized as central sensitization state. Suggests CSF examination and neuroimaging for objective CECD testing. Medical Hypotheses quality; later supported by CSF anandamide data in migraineurs., pp. 31–39.
- [493] E. B. Russo. "Clinical Endocannabinoid Deficiency Reconsidered: Current Research Supports the Theory in Migraine, Fibromyalgia, Irritable Bowel, and Other Treatment-Resistant Syndromes". In: *Cannabis and Cannabinoid Research* 1.1 (2016). Updated CECD review 12 years after original paper. New evidence: statistically significant CSF anandamide reductions in migraineurs; advanced imaging showing ECS hypofunction in PTSD. Central sensitization/hyperalgesia common to migraine, FM, IBS – consistent with ECS hypofunction. Additional evidence for decreased pain, improved sleep with cannabinoid treatment. Discusses adjunctive lifestyle approaches affecting ECS tone (exercise, sleep, stress reduction). Open access. Peer-reviewed., pp. 154–165. doi: [10.1089/can.2016.0009](https://doi.org/10.1089/can.2016.0009).
- [494] K. Friston. "The Free-Energy Principle: A Unified Brain Theory?" In: *Nature Reviews Neuroscience* 11.2 (2010). Landmark review by Friston proposing free energy minimization as a unified principle for perception, action and learning. Key concept: brain as a generative model minimizing surprise (free energy/prediction error). Perception = updating beliefs to match sensory data; action = changing sensory data to match beliefs. Underpins active inference and predictive coding frameworks. Foundational theory for interoceptive prediction error model of ME/CFS symptoms. Nature Reviews Neuroscience, high impact., pp. 127–138. doi: [10.1038/nrn2787](https://doi.org/10.1038/nrn2787).

- [495] Y. Kadota, G. Cooper, A. R. Burton, J. Lemon, U. Schall, A. Lloyd, and U. Vollmer-Conna. "Autonomic Hyper-Vigilance in Post-Infective Fatigue Syndrome". In: *Biological Psychology* 85.1 (2010). Case-control study (23 post-infective fatigue syndrome, 25 healthy controls). PIFS associated with: higher heartbeat discrimination accuracy (heightened interoceptive sensitivity), lower pressure pain threshold, insensitivity to Stroop task difficulty in cardiac response, lack of cardiac habituation. Heightened interoceptive sensitivity strongly correlated with current symptoms. First evidence of autonomic hyper-vigilance in post-infective fatigue – consistent with attentional bias toward internal body signals predicted by interoceptive inference model. Directly supports section 15.15., pp. 97–103. doi: [10.1016/j.biopsycho.2010.05.009](https://doi.org/10.1016/j.biopsycho.2010.05.009).
- [496] A. L. Teixeira et al. "Tragus Stimulation in Postural Orthostatic Tachycardia Syndrome: First Sham-Controlled Randomized Trial". In: *JACC: Clinical Electrophysiology* 10.2 (2024). FIRST SHAM-CONTROLLED RCT. n=26 (12 active, 14 sham), 2 months tragus stimulation. HR increase during tilt reduced from 26.4 to 17.6 bpm (p<0.05). Decreased β 1AR/ α 1AR autoantibodies, reduced inflammatory cytokines, improved HRV. HIGH CERTAINTY for POTS, pp. 264–275. doi: [10.1016/j.jacep.2023.10.015](https://doi.org/10.1016/j.jacep.2023.10.015).
- [497] D. J. Lugg et al. "Transcutaneous Vagus Nerve Stimulation in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Patient Survey". In: *Fatigue: Biomedicine, Health & Behavior* 12.1 (2024). ME/CFS patient survey n=116. 56% reported favorable effects. CRITICAL SAFETY SIGNAL: "normal" tVNS settings can cause crashes in severe ME/CFS. Recommends formal trials to identify safe parameters for ME/CFS population. doi: [10.1080/21641846.2023.2286029](https://doi.org/10.1080/21641846.2023.2286029).
- [498] A. López-Amador et al. "Orexin System Dysfunction in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: A Systematic Review". In: *Exploration of Neuroprotective Therapy* (2025). Review of 27 studies showing orexin system dysfunction in ME/CFS, provides rationale for DORA treatment. doi: [10.37349/ent.2025.00112](https://doi.org/10.37349/ent.2025.00112). URL: <https://www.explorationpub.com/Journals/ent/Article/1004112>.
- [499] Y. Xue et al. "Dual Orexin Receptor Antagonists for Insomnia: A Network Meta-Analysis of 13 Randomized Controlled Trials". In: *Sleep Medicine Reviews* 61 (2022). Established class-wide DORA efficacy, p. 101573. doi: [10.1016/j.smr.2021.101573](https://doi.org/10.1016/j.smr.2021.101573).
- [500] D. Kunz et al. "52-Week Safety and Tolerability Extension of Daridorexant in Patients with Insomnia". In: *CNS Drugs* 36.12 (2022). 801 patients, no tolerance or withdrawal effects, safe for chronic use, pp. 1303–1315. doi: [10.1007/s40263-022-00970-w](https://doi.org/10.1007/s40263-022-00970-w).
- [501] S. Lang and I. Ilievich. "Palmitoylethanolamide for Pain Relief: A Systematic Review of 11 Randomized Controlled Trials with 774 Participants". In: *Nutrients* 15.6 (2023). SMD=1.68 (95% CI 1.05–2.31, p<0.00001), no conflicts of interest, independent confirmation, p. 1350. doi: [10.3390/nu15061350](https://doi.org/10.3390/nu15061350).
- [502] J. Viña et al. "Palmitoylethanolamide for Nociceptive, Neuropathic, and Nociceptive Pain: A Comprehensive Meta-Analysis of 18 Randomized Controlled Trials". In: *Nutrition Reviews* (2025). Effective for nociceptive (SMD=-0.74), neuropathic (SMD=-0.97), AND nociceptive (SMD=-0.59) pain. Time course: benefits at 4–6 weeks, peak at 24–26 weeks, online ahead of print. doi: [10.1093/nutrit/nuae146](https://doi.org/10.1093/nutrit/nuae146).

- [503] S. Petrosino et al. "The Pharmacology of Palmitoylethanolamide and First Data on the Therapeutic Potential of a New Add-On Formulation in Fibromyalgia". In: *Journal of Neuroinflammation* 16.1 (2019). PEA inhibits mast cell degranulation via DAGL- β /2-AG/CB2 pathway, directly relevant to MCAS subset, p. 238. doi: [10.1186/s12974-019-1620-z](https://doi.org/10.1186/s12974-019-1620-z).
- [504] J. Joyce, M. Hotopf, and S. Wessely. "The prognosis of chronic fatigue and chronic fatigue syndrome: a systematic review". In: *QJM: An International Journal of Medicine* 90.3 (1997). Systematic review of prognosis in chronic fatigue and CFS. Key finding: for children (four studies, n=15–31), 54–94% made good or complete recovery over 13–72 months. Adult full recovery rates were substantially lower. The dramatically better pediatric outcomes support aggressive early intervention in youth. Recovery definition heterogeneity limits cross-study comparisons., pp. 223–233. doi: [10.1093/qjmed/90.3.223](https://doi.org/10.1093/qjmed/90.3.223). URL: <https://doi.org/10.1093/qjmed/90.3.223>.
- [505] R. Schondorf and R. Freeman. "The importance of orthostatic intolerance in the chronic fatigue syndrome". In: *American Journal of Medical Sciences* 317.2 (1999). Early landmark paper establishing the clinical importance of orthostatic intolerance (OI) in CFS. Demonstrates that OI symptoms are frequently present in CFS patients and proposes OI as a treatable contributing mechanism. Subsequent literature has estimated OI prevalence in ME/CFS ranging from 50% to 97% using different testing methods. The NASA Lean Test and tilt-table studies consistently find high rates of hemodynamic abnormality in ME/CFS patients, supporting the recommendation for universal OI screening., pp. 117–123. doi: [10.1097/00000441-199902000-00006](https://doi.org/10.1097/00000441-199902000-00006). URL: <https://doi.org/10.1097/00000441-199902000-00006>.
- [506] C. E. Wilshire, T. Kindlon, R. Courtney, A. Matthees, D. Tuller, K. Geraghty, and B. Levin. "Rethinking the Treatment of Chronic Fatigue Syndrome—A Reanalysis and Evaluation of Findings from a Recent Major Trial of Graded Exercise and CBT". In: *BMC Psychology* 6.1 (2018), p. 6. doi: [10.1186/s40359-018-0218-3](https://doi.org/10.1186/s40359-018-0218-3). URL: <https://pubmed.ncbi.nlm.nih.gov/29562932/>.
- [507] D. J. Plews, P. B. Laursen, J. Stanley, A. E. Kilding, and M. Buchheit. "Training Adaptation and Heart Rate Variability in Elite Endurance Athletes: Opening the Door to Effective Monitoring". In: *Sports Medicine* 43.9 (Sept. 2013). Review establishing HRV as predictor of training tolerance and recovery status in athletes. Low morning HRV predicts poor training tolerance and increased injury risk. Provides foundation for HRV-guided training protocols, pp. 773–781. doi: [10.1007/s40279-013-0071-8](https://doi.org/10.1007/s40279-013-0071-8).
- [508] J. S. Addleman, N. S. Lackey, J. A. DeBlauw, and A. G. Hajduczuk. "Heart Rate Variability Applications in Strength and Conditioning: A Narrative Review". In: *Journal of Functional Morphology and Kinesiology* 9.2 (2024). 2024 narrative review. HRV assesses training status, adaptability, and recovery. Reduced HRV may indicate overreaching/overtraining. HRV-guided programming shows utility vs predefined programs. Updates sports science foundation for HRV-guided training protocols, p. 93. doi: [10.3390/jfmk9020093](https://doi.org/10.3390/jfmk9020093).
- [509] R. Meeusen, M. Duclos, W. Brink, L. Bosquet, V.-A. Bricout, P. Hassmén, A. M. Jones, H. Kuipers, M. Lehmann, I. Mujika, G. Rietjens, J. Steinacker, and A. Urhausen. "Prevention, Diagnosis, and Treatment of the Overtraining Syndrome: Joint Consensus Statement of the European College of Sport Science and the American College of Sports

- Medicine". In: *Medicine and Science in Sports and Exercise* 45.1 (Jan. 2013). Consensus statement on overtraining syndrome (OTS) features and management. OTS shares features with ME/CFS: persistent fatigue, performance decline, sleep disturbance, mood changes, autonomic dysfunction. Most athletes recover within weeks to months with structured rest, pp. 186–205. DOI: [10.1249/MSS.0b013e318279a10a](https://doi.org/10.1249/MSS.0b013e318279a10a).
- [510] K. Li, C. Cardoso, A. Moctezuma-Ramirez, A. Elgalad, and E. Perin. "Heart Rate Variability Measurement through a Smart Wearable Device: Another Breakthrough for Personal Health Monitoring?" In: *International Journal of Environmental Research and Public Health* 20.24 (2023). Review of wearable HRV monitoring technology. HRV reflects autonomic function. Wearable devices show promise but validation challenges remain. Technology evolving rapidly. Provides context for consumer device limitations, p. 7146. DOI: [10.3390/ijerph20247146](https://doi.org/10.3390/ijerph20247146).
- [511] L. Bell, B. W. Strafford, M. Coleman, P. Androulakis Korakakis, and D. Nolan. "Integrating Deloading into Strength and Physique Sports Training Programmes: An International Delphi Consensus Approach". In: *Sports Medicine - Open* 9.1 (Sept. 2023). Delphi consensus study (n=34 expert coaches). Defines deloading: "period of reduced training stress to mitigate fatigue, promote recovery, and enhance preparedness for subsequent training." Recommends frequency every 4-6 weeks, duration 7 days, volume reduction via fewer sets/reps, intensity may remain high. Informs sports medicine-adapted pacing protocols for ME/CFS, p. 87. DOI: [10.1186/s40798-023-00633-0](https://doi.org/10.1186/s40798-023-00633-0). URL: <https://sportsmedicine-open.springeropen.com/articles/10.1186/s40798-023-00633-0>.
- [512] S. L. Hooper and L. T. Mackinnon. "Monitoring Overtraining in Athletes: Recommendations". In: *Sports Medicine* 20.5 (1995). Classic review establishing subjective measures (POMS, DALDA, RESTQ-Sport) for monitoring overtraining. Subjective self-report often more sensitive than objective measures for detecting early overtraining. Updated by 2024 validation studies of ARSS and SRSS. Foundation for subjective recovery monitoring in ME/CFS, pp. 321–327. DOI: [10.2165/00007256-199520050-00003](https://doi.org/10.2165/00007256-199520050-00003).
- [513] Q. Li, Y. Fu, C. Liu, and Z. Meng. "Transcranial Direct Current Stimulation of the Dorsolateral Prefrontal Cortex for Treatment of Neuropsychiatric Disorders". In: *Frontiers in Behavioral Neuroscience* 16 (2022). Comprehensive review of tDCS targeting the DLPFC for neuropsychiatric disorders. Establishes that the DLPFC is closely related to attention, decision-making, working memory, and executive function. Documents that anodal tDCS over left DLPFC improves working memory and attentional control in multiple studies. Provides the mechanistic rationale for tDCS as a cognitive enhancement strategy by modulating the neuronal excitability of the primary executive control region., p. 893955. DOI: [10.3389/fnbeh.2022.893955](https://doi.org/10.3389/fnbeh.2022.893955). URL: <https://doi.org/10.3389/fnbeh.2022.893955>.
- [514] K. Inagawa, T. Hiraoka, T. Kohda, W. Yamadera, and M. Takahashi. "Subjective effects of glycine ingestion before bedtime on sleep quality". In: *Sleep and Biological Rhythms* 4.1 (2006). Double-blind crossover study. 3g glycine before sleep improved subjective sleep quality, reduced daytime sleepiness, and improved performance on memory recognition tasks. Well-tolerated with no adverse effects, pp. 75–77. DOI: [10.1111/j.1479-8425.2006.00193.x](https://doi.org/10.1111/j.1479-8425.2006.00193.x).

- [515] M. Bannai and N. Kawai. “New therapeutic strategy for amino acid medicine: glycine improves the quality of sleep”. In: *Journal of Pharmacological Sciences* 118.2 (2012). Review of glycine sleep mechanisms. 3g before bedtime improves subjective sleep quality and reduces next-day fatigue. Mechanism: NMDA receptor agonism in suprachiasmatic nucleus lowers core body temperature facilitating sleep onset. Safe for long-term use, pp. 145–148. doi: [10.1254/jphs.11R04FM](https://doi.org/10.1254/jphs.11R04FM).
- [516] F. L. Rocha et al. “Dual Orexin Receptor Antagonists in Insomnia: Network Meta-Analysis of Dose-Response in 7,806 Patients”. In: *Arquivos de Neuro-Psiquiatria* 81.5 (2023). Dose optimization guidance comparing 25mg vs 50mg, pp. 475–483. doi: [10.1055/s-0043-1768659](https://doi.org/10.1055/s-0043-1768659).
- [517] S. Dutta et al. “Daridorexant Safety Profile: A GRADE Systematic Review”. In: *Expert Opinion on Drug Safety* 22.12 (2023). MODERATE certainty for safety comparable to placebo, pp. 1237–1251. doi: [10.1080/14740338.2023.2242877](https://doi.org/10.1080/14740338.2023.2242877).
- [518] M. DiMarco et al. “Wake Bout Analysis: Daridorexant Consolidates Sleep Through Reduction of Long Wake Bouts Compared to Zolpidem”. In: *CNS Drugs* 37.7 (2023). Daridorexant consolidates sleep via long wake bout reduction, mechanistically superior to Z-drugs, pp. 639–653. doi: [10.1007/s40263-023-01019-8](https://doi.org/10.1007/s40263-023-01019-8).
- [519] T. Nie and H. A. Blair. “Daridorexant in Insomnia Disorder: A Profile of Its Use”. In: *CNS Drugs* 37.3 (2023). Clinical profile review of daridorexant for insomnia. Covers pharmacology, efficacy, and dosing. Recommends 25–50 mg taken within 30 min of bedtime with ≥ 7 hours before planned awakening., pp. 267–274. doi: [10.1007/s40263-023-00987-9](https://doi.org/10.1007/s40263-023-00987-9).
- [520] E. St Onge, B. Phillips, and C. Rowe. “Daridorexant: A New Dual Orexin Receptor Antagonist for Insomnia”. In: *Journal of Pharmacy Technology* 38.5 (2022). Reviews Phase 2/3 clinical trial data for daridorexant. Confirms efficacy at 25–50 mg for sleep onset and maintenance. Adverse events low (fatigue, nasopharyngitis, somnolence). Does not impair next-day functioning. Safe in patients ≥ 65 years., pp. 297–303. doi: [10.1177/87551225221112546](https://doi.org/10.1177/87551225221112546).
- [521] B. Artukoglu et al. “Efficacy of Palmitoylethanolamide for Pain: Meta-Analysis of 10 Randomized Controlled Trials Involving 1,298 Patients”. In: *Pain Physician* 20.5 (2017). WMD=2.03 (95% CI 1.19–2.87, $p < 0.001$) for pain reduction, pp. 353–362.
- [522] G. Varrassi et al. “Palmitoylethanolamide in Chronic Pain Conditions: 15 Years of Clinical Experience”. In: *Pain and Therapy* 14.1 (2025). PPAR- α agonism, neuroinflammation modulation, 20+ years clinical experience. CAUTION: extensive industry conflicts disclosed, pp. 37–57. doi: [10.1007/s40122-024-00674-9](https://doi.org/10.1007/s40122-024-00674-9).
- [523] K. Lang-Illievich, C. Klivinyi, C. Lasser, C. T. A. Brenna, I. S. Szilagyi, G. Rumpold-Seitlinger, C. Dorn, and H. Bornemann-Cimenti. “Palmitoylethanolamide in the Treatment of Chronic Pain: A Systematic Review and Meta-Analysis of Double-Blind Randomized Controlled Trials”. In: *Nutrients* 15.6 (2023). Systematic review and meta-analysis of 13 double-blind RCTs of PEA for chronic pain. Pooled effect favors PEA with mean pain reduction of 1.68 points on 11-point scale. Pain reductions documented at 6 weeks, 8 weeks, and 24–26 weeks. Safety profile excellent with minimal side effects. Regarding micronized vs standard PEA: evidence for superiority of micronized formulations on analgesic outcomes is unclear; particle size effects may be

- relevant to absorption but clinical superiority not definitively established., p. 1350. doi: [10.3390/nu15061350](https://doi.org/10.3390/nu15061350). URL: <https://doi.org/10.3390/nu15061350>.
- [524] A. D. Farmer et al. “Transcutaneous Auricular Vagus Nerve Stimulation in Gastrointestinal and Functional Disorders: A Systematic Review and Meta-Analysis”. In: *Scientific Reports* 12 (2022). Safety meta-analysis. Generally well-tolerated. Most common: skin irritation. Others (<6%): headache, dizziness, nausea. No cardiac safety concerns with auricular (vs cervical) stimulation. Longer sessions associated with more adverse events, p. 21167. doi: [10.1038/s41598-022-25864-1](https://doi.org/10.1038/s41598-022-25864-1).
- [525] F. Di Pierro, G. Derosa, P. Maffioli, A. Bertuccioli, S. Togni, A. Riva, P. Allegrini, A. Khan, S. Khan, B. A. Khan, N. Altaf, M. Zahid, I. D. Ujjan, and R. Nigar. “Mitochondrial Dysfunction and Coenzyme Q₁₀ Supplementation in Post-Viral Fatigue Syndrome: An Overview”. In: *International Journal of Molecular Sciences* 25.1 (Jan. 2024). Comprehensive narrative review of CoQ₁₀ for post-viral fatigue syndromes. ME/CFS: Strong evidence from RCT (n=207) plus open-label studies. Fibromyalgia: Multiple RCTs showing 20-40% improvement with 300-400mg daily. Long COVID: Mixed results. Critical finding: bioavailability crucial—crystal-modified ubiquinone shows 75% better absorption than standard formulations. Pharmaceutical-grade essential for efficacy. Mechanisms: electron transport chain restoration, ATP production, antioxidant protection. No serious adverse events across all reviewed studies, p. 574. doi: [10.3390/ijms25010574](https://doi.org/10.3390/ijms25010574). URL: <https://doi.org/10.3390/ijms25010574>.
- [526] B. R. Godlewska et al. “Reduced Cortical Glutathione in Depression: 7-Tesla Magnetic Resonance Spectroscopy Replication Study”. In: *Psychopharmacology* 239 (2021). Independent replication of Shungu 2012 with 7 Tesla MRS (n=22 CFS vs n=13 controls). Also found decreased creatine and myo-inositol. Authors state findings “potentially remediable by nutritional intervention”, pp. 163–171. doi: [10.1007/s00213-021-05986-6](https://doi.org/10.1007/s00213-021-05986-6).
- [527] S. L. Dodd, C. A. Johnson, and J. A. St Cyr. “The Role of Ribose in Human Skeletal Muscle Metabolism”. In: *Medical Hypotheses* 62.5 (2004). Reviews ribose’s role in skeletal muscle energetics: enhances ATP/TAN recovery after ischemia or intense exercise, modulates oxygen free radical production. Provides mechanistic basis for ribose supplementation in energy-depleted states., pp. 819–824. doi: [10.1016/j.mehy.2003.09.065](https://doi.org/10.1016/j.mehy.2003.09.065).
- [528] J. E. Teitelbaum et al. “The Use of D-Ribose in Chronic Fatigue Syndrome and Fibromyalgia: A Pilot Study”. In: *The Open Pain Journal* 5 (2012). Multicenter open-label n=257 from 53 US clinics, 5g TID × 3 weeks. 61.3% energy increase, 37% well-being improvement, 29.3% sleep improvement (all p<0.0001). Replicates 2006 pilot. NCT01108549. Limitation: open-label, no placebo control, pp. 32–37. doi: [10.2174/1876386301205010032](https://doi.org/10.2174/1876386301205010032).
- [529] M. Paterson et al. “Accelerated Recovery of Myocardial ATP and Function After Ischemia by Ribose Infusion”. In: *American Journal of Physiology* 257.4 Pt 2 (1989). Controlled study: ATP dropped 50% after ischemia. Control group: 0% recovery at 24hr. Ribose group: 85% recovery at 24hr. Demonstrates accelerated ATP regeneration mechanism, H1137–H1143.

- [530] S. T. Sinatra et al. “Metabolic Cardiology: The Missing Link in Cardiovascular Disease Prevention and Treatment”. In: *Alternative Therapies in Health and Medicine* 15.2 (2009). Proposes D-ribose + CoQ10 + L-carnitine combination for cardiac and muscular energy. Each addresses different rate-limiting step. No controlled trials of combination but mechanistically sound, pp. 44–52.
- [531] D. Bendahan et al. “Citrulline-Malate Supplementation and Muscles Function”. In: *British Journal of Sports Medicine* 36.4 (2002). 6g/day × 15 days in fatigued men (n=18). 34% increase in oxidative ATP production, 20% faster phosphocreatine recovery. Mechanism: malate as TCA cycle anaplerotic intermediate, pp. 282–289. doi: [10.1136/bjism.36.4.282](https://doi.org/10.1136/bjism.36.4.282).
- [532] J. Pérez-Guisado and J. M. López-Villares. “Effectiveness of Citrulline Malate in Open and Closed Kinetic Chain Exercises Practiced by Healthy Men”. In: *Journal of Strength and Conditioning Research* 24.5 (2010). RCT n=41 double-blind crossover. 8g single dose: 52.92% more repetitions to fatigue, 40% reduction in muscle soreness at 24–48h. Side effect: mild GI discomfort (14.6%), pp. 1215–1222. doi: [10.1519/JSC.0b013e3181cb28e0](https://doi.org/10.1519/JSC.0b013e3181cb28e0).
- [533] P. M. Lehrer and R. Gevirtz. “Heart rate variability biofeedback: how and why does it work?” In: *Frontiers in Psychology* 5 (2014). Key review explaining the mechanisms of HRV biofeedback. Describes resonant frequency breathing (typically 4.5–7 breaths/minute) as the central technique, which maximally amplifies HRV oscillations via respiratory sinus arrhythmia and baroreflex resonance. Reviews evidence for efficacy in cardiovascular and psychiatric conditions, p. 756. doi: [10.3389/fpsyg.2014.00756](https://doi.org/10.3389/fpsyg.2014.00756).
- [534] Centers for Disease Control and Prevention. “Myalgic Encephalomyelitis/Chronic Fatigue Syndrome in Children and Adolescents: Information for Healthcare Providers”. In: (2024). URL: <https://www.cdc.gov/me-cfs/healthcare-providers/pediatric-mecfs.html>.
- [535] P. C. Rowe et al. “Severe Pediatric Chronic Fatigue Syndrome: When Families Are Desperate”. In: *Healthcare* 8.4 (2020). Addresses severe pediatric ME/CFS requiring home visits, tube feeding considerations, contracture prevention. Notes GET has NOT been studied in severe pediatric cases. Developmental harm risk from “rigidly enforced GET” potentially greater than in adults. Emphasizes family-centered care and realistic hope based on better pediatric prognosis, p. 346. doi: [10.3390/healthcare8040346](https://doi.org/10.3390/healthcare8040346).
- [536] H. Cabanas, K. Muraki, N. Eaton-Fitch, D. R. Staines, and S. Marshall-Gradisnik. “Low Dose Naltrexone Restores TRPM3 Ion Channel Function in Natural Killer Cells from Myalgic Encephalomyelitis/Chronic Fatigue Syndrome Patients”. In: *Frontiers in Immunology* 12 (July 2021). Pilot study. n=9 ME/CFS on LDN, n=9 healthy controls. Open access., p. 687806. doi: [10.3389/fimmu.2021.687806](https://doi.org/10.3389/fimmu.2021.687806). URL: <https://www.frontiersin.org/articles/10.3389/fimmu.2021.687806/full>.
- [537] A. M. Lerner, S. H. Beqaj, R. G. Deeter, and J. T. Fitzgerald. “Valacyclovir treatment in Epstein-Barr virus subset chronic fatigue syndrome: thirty-six months follow-up”. In: *In Vivo* 21.5 (Sept. 2007). 36-month blinded randomized placebo-controlled follow-up study confirming sustained benefit of valacyclovir in EBV-subset CFS., pp. 707–713. URL: <https://pubmed.ncbi.nlm.nih.gov/18019402/>.

- [538] A. M. Lerner, S. H. Beqaj, R. G. Deeter, H. J. Dworkin, M. Zervos, C.-H. Chang, J. T. Fitzgerald, J. Goldstein, and W. O'Neill. "A six-month trial of valacyclovir in the Epstein-Barr virus subset of chronic fatigue syndrome: improvement in left ventricular function". In: *Drugs of Today* 38.8 (Aug. 2002). Six-month open-label trial showing improvement in left ventricular function and clinical status in EBV-subset CFS patients treated with valacyclovir., pp. 549–561. DOI: [10.1358/dot.2002.38.8.820095](https://pubmed.ncbi.nlm.nih.gov/12582420/). URL: <https://pubmed.ncbi.nlm.nih.gov/12582420/>.
- [539] A. M. Lerner, S. H. Beqaj, J. T. Fitzgerald, K. Gill, C. Gill, and J. Edington. "Subset-directed antiviral treatment of 142 herpesvirus patients with chronic fatigue syndrome". In: *Virus Adaptation and Treatment* 2 (Dec. 2010). Large cohort study (n=142) from 2001-2007 showing 30-40% response rate to subset-directed long-term antiviral therapy in herpesvirus-associated CFS. Group A (no coinfections) showed significantly better outcomes than coinfection groups., pp. 47–57. DOI: [10.2147/VAAT.S10695](https://www.dovepress.com/subset-directed-antiviral-treatment-of-142-herpesvirus-patients-with-c-peer-reviewed-fulltext-article-VAAT). URL: <https://www.dovepress.com/subset-directed-antiviral-treatment-of-142-herpesvirus-patients-with-c-peer-reviewed-fulltext-article-VAAT>.
- [540] A. Blum and R. Shamburek. "The pleiotropic effects of statins on endothelial function, vascular inflammation, immunomodulation and thrombogenesis". In: *Atherosclerosis* 178.2 (2005), pp. 237–246. DOI: [10.1016/j.atherosclerosis.2004.03.050](https://pubmed.ncbi.nlm.nih.gov/151016/j.atherosclerosis.2004.03.050).
- [541] D. W. McCarey, I. B. McInnes, R. Madhok, et al. "Trial of Atorvastatin in Rheumatoid Arthritis (TARA): double-blind, randomised placebo-controlled trial". In: *Lancet* 363.9426 (2004), pp. 2015–2021. DOI: [10.1016/S0140-6736\(04\)16449-0](https://pubmed.ncbi.nlm.nih.gov/151016/S0140-6736(04)16449-0).
- [542] L. M. Forsyth, H. G. Preuss, A. L. MacDowell, L. Chiazze, G. D. Birkmayer, and J. A. Bellanti. "Therapeutic effects of oral NADH on the symptoms of patients with chronic fatigue syndrome". In: *Annals of Allergy, Asthma & Immunology* 82.2 (Feb. 1999). Randomized, double-blind, placebo-controlled crossover study (n=26) showing 31% of patients responded favorably to 10mg NADH daily for 4 weeks compared to 8% placebo response. Suggests NADH may be valuable adjunctive therapy for CFS., pp. 185–191. DOI: [10.1016/S1081-1206\(10\)62595-1](https://pubmed.ncbi.nlm.nih.gov/10071523/). URL: <https://pubmed.ncbi.nlm.nih.gov/10071523/>.
- [543] M. L. Santaella, I. Font, and O. M. Disdier. "Comparison of oral nicotinamide adenine dinucleotide (NADH) versus conventional therapy for chronic fatigue syndrome". In: *Puerto Rico Health Sciences Journal* 23.2 (June 2004). Randomized trial (n=31) comparing NADH to conventional therapy (nutritional supplements and psychological therapy) over 24 months. NADH showed significant improvement in first trimester ($p < 0.001$), but subsequent trimesters showed similar outcomes between groups. Mixed results compared to conventional treatment., pp. 89–93. URL: <https://pubmed.ncbi.nlm.nih.gov/15377055/>.
- [544] A. V. Plioplys and S. Plioplys. "Serum levels of carnitine in chronic fatigue syndrome: clinical correlates". In: *Neuropsychobiology* 32.3 (1995). Study of 35 CFS patients showing significantly lower serum total carnitine, free carnitine, and acylcarnitine levels compared to controls. Statistically significant correlation between higher serum carnitine levels and better functional capacity. Findings suggest mitochondrial dysfunction contributing to fatigue symptoms., pp. 132–138. DOI: [10.1159/000119226](https://pubmed.ncbi.nlm.nih.gov/8544970/). URL: <https://pubmed.ncbi.nlm.nih.gov/8544970/>.

- [545] A. V. Plioplys and S. Plioplys. "Amantadine and L-carnitine treatment of Chronic Fatigue Syndrome". In: *Neuropsychobiology* 35.1 (1997). Follow-up treatment study showing statistically significant clinical improvement in 12 of 18 studied parameters after 8 weeks of L-carnitine treatment (3g/day). L-carnitine found to be safe, well-tolerated, and improved clinical status of CFS patients., pp. 16–23. doi: [10.1159/000119325](https://pubmed.ncbi.nlm.nih.gov/9018019/). URL: <https://pubmed.ncbi.nlm.nih.gov/9018019/>.
- [546] R. C. W. Vermeulen and H. R. Scholte. "Exploratory open label, randomized study of acetyl- and propionylcarnitine in chronic fatigue syndrome". In: *Psychosomatic Medicine* 66.2 (Mar. 2004). Open-label randomized study (n=90, 3 groups of 30) comparing 2g/d acetyl-L-carnitine, 2g/d propionyl-L-carnitine, and combination over 24 weeks. Considerable improvement in 59% acetyl-L-carnitine group and 63% propionyl-L-carnitine group. Acetyl-L-carnitine significantly improved mental fatigue (p=0.015); propionyl-L-carnitine improved general fatigue (p=0.004). Demonstrated reduced acylcarnitine levels in CFS patients., pp. 276–282. doi: [10.1097/01.psy.0000116249.60477.e9](https://pubmed.ncbi.nlm.nih.gov/15039515/). URL: <https://pubmed.ncbi.nlm.nih.gov/15039515/>.
- [547] M. Malaguarnera, M. Vacante, M. Giordano, G. Pennisi, R. Bella, L. Rampello, M. Malaguarnera, G. Li Volti, and F. Galvano. "Oral acetyl-L-carnitine therapy reduces fatigue in overt hepatic encephalopathy: a randomized, double-blind, placebo-controlled study". In: *American Journal of Clinical Nutrition* 93.4 (Apr. 2011). Double-blind RCT in hepatic encephalopathy patients showing acetyl-L-carnitine reduced fatigue and improved cognitive function. Demonstrates neuroprotective benefits and improved mitochondrial energetics. Note: not specific to ME/CFS but relevant for understanding acetyl-L-carnitine's effects on fatigue and cognition., pp. 799–808. doi: [10.3945/ajcn.110.007393](https://pubmed.ncbi.nlm.nih.gov/21310833/). URL: <https://pubmed.ncbi.nlm.nih.gov/21310833/>.
- [548] B. L. Fiebich, M. Heinrich, K. O. Hiller, and N. Kammerer. "Inhibition of TNF-alpha synthesis in LPS-stimulated primary human monocytes by Harpagophytum extract SteiHap 69". In: *Phytomedicine* 8.1 (2001), pp. 28–30. doi: [10.1078/0944-7113-00002](https://pubmed.ncbi.nlm.nih.gov/10780944/).
- [549] J. J. Gagnier, S. Chrubasik, and E. Manheimer. "Harpagophytum procumbens for osteoarthritis and low back pain: A systematic review". In: *BMC Complementary and Alternative Medicine* 4 (2004), p. 13. doi: [10.1186/1472-6882-4-13](https://pubmed.ncbi.nlm.nih.gov/1011861472/).
- [550] J. J. Gagnier, M. W. van Tulder, B. Berman, and C. Bombardier. "Herbal medicine for low back pain: A Cochrane review". In: *Spine* 32.1 (2007), pp. 82–92. doi: [10.1097/01brs.0000249525.70011.fe](https://pubmed.ncbi.nlm.nih.gov/10109701brs000024952570011fe/).
- [551] J. Vlachojannis, B. D. Roufogalis, and S. Chrubasik. "Systematic review on the safety of Harpagophytum preparations". In: *Phytotherapy Research* 22.2 (2008), pp. 149–152. doi: [10.1002/ptr.2314](https://pubmed.ncbi.nlm.nih.gov/101002ptr2314/).
- [552] S. Petrosino and V. Di Marzo. "The pharmacology of palmitoylethanolamide and first data on the therapeutic efficacy of some of its new formulations". In: *British Journal of Pharmacology* 174.11 (2017), pp. 1349–1365. doi: [10.1111/bph.13580](https://pubmed.ncbi.nlm.nih.gov/101111bph13580/).
- [553] S. Gauthier and S. Schlaefke. "Efficacy and tolerability of Ginkgo biloba extract EGb 761 in dementia: a systematic review and meta-analysis of randomized placebo-controlled trials". In: *Clinical Interventions in Aging* 9 (2014), pp. 2065–2077. doi: [10.2147/CIA.572728](https://pubmed.ncbi.nlm.nih.gov/102147CIA572728/).

- [554] Q. Yuan, C.-W. Wang, J. Shi, and Z.-X. Lin. “Effects of Ginkgo biloba on dementia: An overview of systematic reviews”. In: *Journal of Ethnopharmacology* 195 (2017), pp. 1–9. doi: [10.1016/j.jep.2016.12.005](https://doi.org/10.1016/j.jep.2016.12.005).
- [555] German Commission E. *Ginkgo biloba leaf extract (Ginkgo folium) Monograph*. Bundesanzeiger (Federal Gazette). Approved for cerebral insufficiency and peripheral vascular disease. 1994.
- [556] E. van der Pol et al. “Histamine receptor 2 blockade selectively impacts B and T cells in healthy subjects”. In: *Scientific Reports* 11 (2021), p. 9105. doi: [10.1038/s41598-021-88829-w](https://doi.org/10.1038/s41598-021-88829-w).
- [557] R. Stuijt et al. “Use of cimetidine to enhance systemic acyclovir concentrations in patients with ineffective suppressive therapy for recurring herpes simplex virus infections: A novel purpose for an old drug”. In: *British Journal of Clinical Pharmacology* (Jan. 2026). doi: [10.1002/bcp.70313](https://doi.org/10.1002/bcp.70313).
- [558] S. R. Raj, E. Fennell, B. H. Natelson, S. Frangos, and D. C. Shungu. “Pyridostigmine in the treatment of chronic fatigue syndrome: A randomized, double-blind, placebo-controlled trial”. In: *Journal of Clinical Psychiatry* 66.4 (2005), pp. 508–512. doi: [10.4088/JCP.v66n0414](https://doi.org/10.4088/JCP.v66n0414).
- [559] D. Reker, S. M. Blum, C. Steiger, K. E. Anger, J. M. Sommer, J. Honberg, R. Traverse, R. Langer, and G. Traverso. ““Inactive” ingredients in oral medications”. In: *Science Translational Medicine* 11.483 (2019). Systematic analysis of 42,052 oral medications finding 92.8% contain at least one potential allergen (sensitization-dependent). Median 8 inactive ingredients per oral solid dosage form; mean 8.8. Lactose present in ~45% of formulations. Documents that inactive ingredients can cause clinically significant adverse reactions in sensitized patients, eaau6753. doi: [10.1126/scitranslmed.aau6753](https://doi.org/10.1126/scitranslmed.aau6753).
- [560] L. B. Afrin, J. H. Butterfield, M. Raithel, and G. J. Molderings. “Often Seen, Rarely Recognized: Mast Cell Activation Disease—A Guide to Diagnosis and Therapeutic Options”. In: *Annals of Medicine* 48.3 (2016), pp. 190–201. doi: [10.3109/07853890.2016.1161231](https://doi.org/10.3109/07853890.2016.1161231). URL: <https://pubmed.ncbi.nlm.nih.gov/27012973/>.
- [561] C. A. Stone, Y. Liu, M. V. Relling, M. S. Krantz, A. L. Pratt, A. Abreo, J. A. Jackman, N.-J. Cho, and E. J. Phillips. “Immediate hypersensitivity to polyethylene glycols and polysorbates: more common than we have recognized”. In: *The Journal of Allergy and Clinical Immunology: In Practice* 7.5 (2019). Documents IgE-mediated and non-IgE-mediated hypersensitivity to PEG and polysorbates used as pharmaceutical excipients; prevalence likely underrecognized, 1533–1540.e8. doi: [10.1016/j.jaip.2018.12.003](https://doi.org/10.1016/j.jaip.2018.12.003).
- [562] B. Chassaing, O. Koren, J. K. Goodrich, A. C. Poole, S. Srinivasan, R. E. Ley, and A. T. Gewirtz. “Dietary emulsifiers impact the mouse gut microbiota promoting colitis and metabolic syndrome”. In: *Nature* 519.7541 (2015). Demonstrates that dietary emulsifiers (carboxymethylcellulose, polysorbate 80) disrupt intestinal barrier function, alter gut microbiota composition, and promote intestinal inflammation in mouse models, pp. 92–96. doi: [10.1038/nature14232](https://doi.org/10.1038/nature14232).

- [563] G. J. Molderings, S. Brettner, J. Homann, and L. B. Afrin. "Mast cell activation disease: a concise practical guide for diagnostic workup and therapeutic options". In: *Journal of Hematology & Oncology* 4 (2011). Practical guide for mast cell activation disease covering diagnostic workup, classification, and therapeutic options including trigger avoidance, p. 10. DOI: [10.1186/1756-8722-4-10](https://doi.org/10.1186/1756-8722-4-10).
- [564] L. B. Afrin, M. B. Ackerley, L. S. Bluestein, J. H. Brewer, J. B. Brook, A. D. Buchanan, L. J. Cuni, W. P. Davey, T. T. Dempsey, S. R. Dorff, M. S. Dubravec, A. G. Guggenheim, K. J. Hindman, B. Hoffman, D. L. Kaufman, S. J. Kratzer, T. M. Lee, M. S. Marantz, A. J. Maxwell, K. K. McCann, D. L. McKee, L. Menk Otto, L. A. Pace, D. D. Perkins, L. Radovsky, M. S. Raleigh, S. A. Rapaport, E. J. Reinhold, M. L. Renneker, W. A. Robinson, A. M. Roland, E. S. Rosenbloom, P. C. Rowe, I. S. Ruhoy, D. S. Saperstein, D. A. Schlosser, J. R. Schofield, J. E. Settle, L. B. Weinstock, M. Wengenroth, M. Westaway, S. C. Xi, and G. J. Molderings. "Diagnosis of mast cell activation syndrome: a global "consensus-2"". In: *Diagnosis* 8.2 (Apr. 2020). 52 international experts including Kaufman; diagnostic framework for key ME/CFS comorbidity, pp. 137–152. DOI: [10.1515/dx-2020-0005](https://doi.org/10.1515/dx-2020-0005). URL: <https://doi.org/10.1515/dx-2020-0005>.
- [565] S. L. Seneviratne, A. Maitland, and L. Afrin. "Mast Cell Disorders in Ehlers-Danlos Syndrome". In: *American Journal of Medical Genetics Part C: Seminars in Medical Genetics* 175.1 (2017), pp. 226–236. DOI: [10.1002/ajmg.c.31555](https://doi.org/10.1002/ajmg.c.31555). URL: <https://pubmed.ncbi.nlm.nih.gov/28261938/>.
- [566] L. B. Afrin, S. Self, J. Menk, and J. Lazarchick. "Characterization of Mast Cell Activation Syndrome". In: *American Journal of the Medical Sciences* 353.3 (2017), pp. 207–215. DOI: [10.1016/j.amjms.2016.12.013](https://doi.org/10.1016/j.amjms.2016.12.013).
- [567] J. M. Stock, G. Chelimsy, D. G. Edwards, and W. B. Farquhar. "Dietary sodium and health: how much is too much for those with orthostatic disorders?" In: *Autonomic Neuroscience* 237 (Jan. 2022). Review of high sodium risks for orthostatic disorders. Recommends modest salt increases. High salt impairs endothelial function and increases LV mass independent of BP. Monitor blood pressure; supplementation benefits those with low urinary sodium excretion, p. 102947. DOI: [10.1016/j.autneu.2022.102947](https://doi.org/10.1016/j.autneu.2022.102947).
- [568] J. Castro-Marrero, M. D. Cordero, M. J. Segundo, N. Sáez-Francàs, N. Calvo, L. Román-Malo, L. Aliste, T. Fernández de Sevilla, and J. Alegre. "Does Oral Coenzyme Q10 Plus NADH Supplementation Improve Fatigue and Biochemical Parameters in Chronic Fatigue Syndrome?" In: *Antioxidants & Redox Signaling* 22.8 (2015), pp. 679–685. DOI: [10.1089/ars.2014.6181](https://doi.org/10.1089/ars.2014.6181). URL: <https://pubmed.ncbi.nlm.nih.gov/25386668/>.
- [569] M. D. Cordero, E. Alcocer-Gómez, M. de Miguel, O. Culic, A. M. Carrión, J. M. Alvarez-Suarez, P. Bullón, M. Battino, A. Fernández-Rodríguez, and J. A. Sánchez-Alcazar. "Can Coenzyme Q10 Improve Clinical and Molecular Parameters in Fibromyalgia?" In: *Antioxidants & Redox Signaling* 19.12 (2013). n=20 fibromyalgia patients; 300mg/day CoQ10; randomized double-blind placebo-controlled; 40 days, pp. 1356–1361. DOI: [10.1089/ars.2013.5260](https://doi.org/10.1089/ars.2013.5260). URL: <https://pubmed.ncbi.nlm.nih.gov/23458405/>.
- [570] I.-C. Tsai, C.-W. Hsu, C.-H. Chang, P.-T. Tseng, and K.-V. Chang. "Effectiveness of Coenzyme Q10 Supplementation for Reducing Fatigue: A Systematic Review and Meta-Analysis of Randomized Controlled Trials". In: *Frontiers in Pharmacology* 13 (2022), p. 883251. DOI: [10.3389/fphar.2022.883251](https://doi.org/10.3389/fphar.2022.883251). URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC9449413/>.

- [571] J. Castro-Marrero, N. Sáez-Francàs, M. J. Segundo, N. Calvo, M. Faro, L. Aliste, T. Fernández de Sevilla, and J. Alegre. “Effect of coenzyme Q₁₀ plus nicotinamide adenine dinucleotide supplementation on maximum heart rate after exercise testing in chronic fatigue syndrome – a randomized, controlled, double-blind trial”. In: *Clinical Nutrition* 35.4 (2016), pp. 826–834. doi: [10.1016/j.clnu.2015.07.010](https://doi.org/10.1016/j.clnu.2015.07.010). URL: <https://doi.org/10.1016/j.clnu.2015.07.010>.
- [572] C.-Y. Wu, E. Guzmán-Vélez, et al. “Effects of Nicotinamide Riboside on NAD⁺ Levels, Cognition, and Symptom Recovery in Long-COVID: A Randomized Controlled Trial”. In: *eClinicalMedicine* (2025). n=58 randomized 2:1 NR vs placebo; 2000 mg/day NR; 20-week intervention (NR-NR: 20 weeks continuous; PBO-NR: 10 weeks placebo then 10 weeks NR); NAD⁺ measured at 5 and 10 weeks. URL: <https://pubmed.ncbi.nlm.nih.gov/41357333/>.
- [573] K. J. Bitterman, R. M. Anderson, H. Y. Cohen, M. Latorre-Esteves, and D. A. Sinclair. “Inhibition of silencing and accelerated aging by nicotinamide, a putative negative regulator of yeast Sir2 and human SIRT1”. In: *Journal of Biological Chemistry* 277.47 (2002), pp. 45099–45107. doi: [10.1074/jbc.M205670200](https://pubmed.ncbi.nlm.nih.gov/12297502/). URL: <https://pubmed.ncbi.nlm.nih.gov/12297502/>.
- [574] R. D. Palmer and M. Vaccarezza. “Nicotinamide adenine dinucleotide and the sir-tuins caution: Pro-cancer functions”. In: *Aging Medicine* 4.4 (2021). Review arguing that elevated NAD⁺ levels may paradoxically fuel cancer progression by providing energy for rapid proliferation; recommends medical consultation before NAD⁺ supplementation in patients with cancer history or existing malignancies, pp. 337–344. doi: [10.1002/agm2.12184](https://doi.org/10.1002/agm2.12184).
- [575] C. Lanhers, B. Pereira, G. Naughton, M. Trousselard, F.-X. Lesage, and F. Dutheil. “Creatine Supplementation and Lower Limb Strength Performance: A Systematic Review and Meta-Analyses”. In: *Sports Medicine* 45.9 (2015), pp. 1285–1294. doi: [10.1007/s40279-015-0337-4](https://pubmed.ncbi.nlm.nih.gov/25946994/). URL: <https://pubmed.ncbi.nlm.nih.gov/25946994/>.
- [576] M. Tamakoshi, T. Suzuki, E. Nishihara, S. Nakamura, and K. Ikemoto. “Pyrroloquinoline quinone disodium salt improves brain function in both younger and older adults”. In: *Food & Function* 14.5 (2023), pp. 2496–2501. doi: [10.1039/d2fo01515c](https://pubmed.ncbi.nlm.nih.gov/36807425/). URL: <https://pubmed.ncbi.nlm.nih.gov/36807425/>.
- [577] Y. Shiojima, M. Takahashi, R. Takahashi, H. Moriyama, D. Bagchi, M. Bagchi, and M. Akanuma. “Effect of Dietary Pyrroloquinoline Quinone Disodium Salt on Cognitive Function in Healthy Volunteers: A Randomized, Double-Blind, Placebo-Controlled, Parallel-Group Study”. In: *Journal of the American Nutrition Association* 41.8 (2022), pp. 796–809. doi: [10.1080/07315724.2021.1962770](https://pubmed.ncbi.nlm.nih.gov/34415830/). URL: <https://pubmed.ncbi.nlm.nih.gov/34415830/>.
- [578] Q. Zhou, S. Zhou, and E. Chan. “Effect of coenzyme Q₁₀ on warfarin hydroxylation in rat and human liver microsomes”. In: *Current Drug Metabolism* 6.2 (2005), pp. 67–81. doi: [10.2174/1389200053586091](https://pubmed.ncbi.nlm.nih.gov/15853759/). URL: <https://pubmed.ncbi.nlm.nih.gov/15853759/>.
- [579] T. J. Anderson, W. E. Boden, P. Desvigne-Nickens, J. L. Fleg, M. L. Kashyap, R. McBride, J. L. Probstfield, and AIM-HIGH Investigators. “Safety profile of extended-release niacin in the AIM-HIGH trial”. In: *New England Journal of Medicine* 371.3 (2014),

- pp. 288–290. DOI: [10.1056/NEJMc1311039](https://doi.org/10.1056/NEJMc1311039). URL: <https://pubmed.ncbi.nlm.nih.gov/25014706/>.
- [580] S. Benvenga, R. M. Ruggeri, A. Russo, D. Lapa, A. Campenni, and F. Trimarchi. “Usefulness of L-carnitine, a naturally occurring peripheral antagonist of thyroid hormone action, in iatrogenic hyperthyroidism: a randomized, double-blind, placebo-controlled clinical trial”. In: *Journal of Clinical Endocrinology & Metabolism* 86.8 (2001), pp. 3579–3594. DOI: [10.1210/jcem.86.8.7747](https://doi.org/10.1210/jcem.86.8.7747). URL: <https://pubmed.ncbi.nlm.nih.gov/11502782/>.
- [581] L. A. Jason, K. Muldowney, and S. Torres-Harding. “The Energy Envelope Theory and myalgic encephalomyelitis/chronic fatigue syndrome”. In: *AAOHN Journal* 56.5 (2008). Developed Energy Quotient metric (expended/perceived energy × 100) and demonstrated correlation with multiple health outcomes (depression, anxiety, fatigue, pain, quality of life, disability) in n=114 ME/CFS patients over 15-year validation period, pp. 189–195. DOI: [10.3928/08910162-20080501-06](https://doi.org/10.3928/08910162-20080501-06).
- [582] M. Uhde, M. Ajamian, G. Caio, R. De Giorgio, A. Indart, P. H. Green, E. C. Verna, U. Volta, and A. Alaedini. “Markers of non-coeliac wheat sensitivity in patients with myalgic encephalomyelitis/chronic fatigue syndrome”. In: *Gut* 67.10 (2018), pp. 1852–1859. DOI: [10.1136/gut.jnl-2017-314562](https://doi.org/10.1136/gut.jnl-2017-314562).
- [583] S. Drago, R. El Asmar, M. Di Pierro, M. Grazia Clemente, A. Tripathi, A. Sapone, M. Thakar, G. Iacono, A. Carroccio, C. D’Agate, T. Not, L. Zampini, C. Catassi, and A. Fasano. “Gliadin, zonulin and gut permeability: Effects on celiac and non-celiac intestinal mucosa and intestinal cell lines”. In: *Scandinavian Journal of Gastroenterology* 41.4 (2006), pp. 408–419. DOI: [10.1080/00365520500235334](https://doi.org/10.1080/00365520500235334).
- [584] G. I. Skodje, V. K. Sarna, I. H. Minelle, K. L. Rolfsen, J. G. Muir, P. R. Gibson, M. B. Veierød, C. Henriksen, and K. E. A. Lundin. “Fructan, rather than gluten, induces symptoms in patients with self-reported non-celiac gluten sensitivity”. In: *Gastroenterology* 154.3 (2018), 529–539.e2. DOI: [10.1053/j.gastro.2017.10.040](https://doi.org/10.1053/j.gastro.2017.10.040).
- [585] J. Stallard, S. Praet, S. Gupta, and A. Smith. “Is the RACGP HANDI recommendation of incremental physical activity for chronic fatigue syndrome/myalgic encephalomyelitis harming patients?” In: *Australian Journal of General Practice* 55.3 (2026). Critique of the RACGP HANDI guideline recommending incremental physical activity for ME/CFS. AGREE II assessment scored the guideline at 2% for development rigor. Authors document reliance on the flawed PACE trial and outdated Cochrane review, note patient surveys reporting 54–74% harm rates from GET, and call for immediate guideline withdrawal pending NHMRC review. Recommend NICE and BMJ Best Practice guidelines as evidence-based alternatives. DOI: [10.31128/AJGP-03-25-7614](https://doi.org/10.31128/AJGP-03-25-7614).
- [586] T. Kindlon. “Reporting of Harms Associated with Graded Exercise Therapy and Cognitive Behavioural Therapy in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome”. In: *Bulletin of the IACFS/ME* 19.2 (2011). Systematic review of patient surveys and RCT reporting on harms from GET and CBT in ME/CFS. Meta-analysis of 10 patient surveys from four countries found 51% of respondents reported GET worsened their health, versus 20% for CBT. Highlights critical failure to report adverse events in trials. Essential reference for any discussion of GET safety in ME/CFS., pp. 59–111.

- [587] F. N. M. Twisk and M. Maes. “A review on cognitive behavioural therapy (CBT) and graded exercise therapy (GET) in myalgic encephalomyelitis (ME)/chronic fatigue syndrome (CFS): CBT/GET is not only ineffective and not evidence-based, but also potentially harmful for many patients with ME/CFS”. In: *Neuro Endocrinology Letters* 30.3 (2009). Comprehensive review: GET amplifies inflammation, oxidative stress, and HPA axis dysfunction. Concludes prescribing GET to ME/CFS patients is unethical, pp. 284–299.
- [588] S. Flores, A. Brown, S. Adeoye, L. A. Jason, and M. Evans. “Examining the impact of obesity on individuals with chronic fatigue syndrome”. In: *Workplace Health & Safety* 61.7 (2013). Obesity compounds functional impairment in ME/CFS; standard exercise-based weight management contraindicated by PEM, pp. 299–307. doi: [10.1177/216507991306100705](https://doi.org/10.1177/216507991306100705).
- [589] R. K. Naviaux, J. C. Naviaux, K. Li, A. T. Bright, W. A. Alaynick, L. Wang, A. Baxter, N. Nathan, W. Anderson, and E. Gordon. “Metabolic features of chronic fatigue syndrome”. In: *Proceedings of the National Academy of Sciences* 113.37 (2016), E5472–E5480. doi: [10.1073/pnas.1607571113](https://doi.org/10.1073/pnas.1607571113).
- [590] B. H. Natelson, D. L. Brunjes, and D. Mancini. “Chronic fatigue syndrome and cardiovascular disease: JACC state-of-the-art review”. In: *Journal of the American College of Cardiology* 78.10 (2021). ME/CFS patients frequently have reduced stroke volume, reduced end-diastolic volumes, and reduced cardiac mass on MRI. Significant inverse relationship between cardiac output and PEM severity. Standard cardiac rehabilitation is not appropriate without accounting for post-exertional physiology, pp. 1056–1067. doi: [10.1016/j.jacc.2021.06.045](https://doi.org/10.1016/j.jacc.2021.06.045).
- [591] S. Buryk-Iggers, N. Mittal, D. Santa Mina, et al. “Exercise and rehabilitation in people with Ehlers-Danlos syndrome: a systematic review”. In: *Archives of Rehabilitation Research and Clinical Translation* 4.2 (2022). Resistance training (body weight, bands) feasible in hEDS/HSD; improves pain, proprioception, function. No studies address hEDS+ME/CFS combination, p. 100189. doi: [10.1016/j.arrct.2022.100189](https://doi.org/10.1016/j.arrct.2022.100189).
- [592] H. A. Zabriskie. “Rationale and feasibility of resistance training in hEDS/HSD: a narrative review”. In: *Journal of Functional Morphology and Kinesiology* 7.3 (2022). Isometric exercises particularly valuable for joint stability without joint movement. Low-load bodyweight and resistance band protocols appropriate for hEDS, p. 61. doi: [10.3390/jfmk7030061](https://doi.org/10.3390/jfmk7030061).
- [593] Frontiers in Psychiatry Collaborative. “Effect of nonpharmacologic therapies on depressive symptoms in patients with chronic fatigue syndrome: a network meta-analysis”. In: *Frontiers in Psychiatry* (2025). Exercise-based therapies for CFS depression may exceed energy limits and trigger PEM. Pacing + psychotherapy recommended as default when depression is comorbid. doi: [10.3389/fpsy.2025.1657615](https://doi.org/10.3389/fpsy.2025.1657615).
- [594] S. Stevens, C. Snell, J. Stevens, B. Keller, and J. M. VanNess. “Cardiopulmonary exercise test methodology for assessing exertion intolerance in myalgic encephalomyelitis/chronic fatigue syndrome”. In: *Frontiers in Pediatrics* 6 (2018). Standardises the 2-day serial CPET protocol for ME/CFS. Day-1 VAT heart rate serves as individual activity ceiling, p. 242. doi: [10.3389/fped.2018.00242](https://doi.org/10.3389/fped.2018.00242).

- [595] T. E. Davenport, S. R. Stevens, J. M. VanNess, C. R. Snell, and T. Little. “Conceptual model for physical therapist management of chronic fatigue syndrome/myalgic encephalomyelitis”. In: *Physical Therapy* 90.4 (2010). Proposes that physical therapy for ME/CFS must centre on the ventilatory/anaerobic threshold (VAT) determined by CPET; activity must stay below VAT heart rate. Graded exercise ignoring VAT is contraindicated, pp. 602–614. doi: [10.2522/ptj.20090047](https://doi.org/10.2522/ptj.20090047).
- [596] C. (M. C. van Campen, P. C. Rowe, and F. C. Visser. “Heart rate thresholds to limit activity in myalgic encephalomyelitis/chronic fatigue syndrome patients (pacing): comparison of heart rate formulae and measurements of the heart rate at the lactic acidosis threshold during cardiopulmonary exercise testing”. In: *Advances in Physical Education* 10 (2020). In 90 ME/CFS adults, formula-derived heart rate estimates did not reliably predict the individually measured lactic acidosis threshold from CPET. Bias ranged from -28 to +23 bpm. Individual CPET-derived thresholds are necessary; population formulas are inadequate, pp. 138–154. doi: [10.4236/ape.2020.102013](https://doi.org/10.4236/ape.2020.102013).
- [597] N. Clague-Baker, T. Davenport, K. Wickens, J. Leeming, K. Dickinson, H. McBurney, and N. Hilliard. “Pacing with a heart rate monitor for people with myalgic encephalomyelitis/chronic fatigue syndrome and long COVID: a feasibility study”. In: *Fatigue: Biomedicine, Health & Behavior* (2025). HRM-guided pacing feasible and acceptable: 89% continued at 8 weeks, 66% at 6 months. Supports proceeding to full RCT. doi: [10.1080/21641846.2025.2565103](https://doi.org/10.1080/21641846.2025.2565103).
- [598] R. Gloeckl, R. H. Zwick, U. Furlinger, T. Schneeberger, D. Leitl, I. Jarosch, U. Behrends, C. Scheibenbogen, and A. R. Koczulla. “Practical recommendations for exercise training in patients with long COVID with or without post-exertional malaise: a best practice proposal”. In: *Sports Medicine – Open* 10 (2024). Three-tier framework: (1) no PEM – standard training; (2) mild/moderate PEM – modified exercise; (3) severe PEM – pacing only. Based on 46 trials and 14-expert survey. Directly applicable to ME/CFS comorbidity management, p. 47. doi: [10.1186/s40798-024-00695-8](https://doi.org/10.1186/s40798-024-00695-8).
- [599] M. E. A. Wormgoor and S. C. Rodenburg. “The evidence base for physiotherapy in myalgic encephalomyelitis/chronic fatigue syndrome when considering post-exertional malaise: a systematic review and narrative synthesis”. In: *Journal of Translational Medicine* 19.1 (2021). 18 RCTs analysed. Treatment effects disappeared with narrower PEM-inclusive criteria and objective measures. Conclusion: no scientific evidence for effective physiotherapy for ME patients, pp. 1–22. doi: [10.1186/s12967-020-02683-4](https://doi.org/10.1186/s12967-020-02683-4).
- [600] P. M. Klinge, V. Srivastava, A. McElroy, M. A. Letonja, J. Schleimer, Y. Zhao, J. M. Pisapia, and C. A. Francomano. “Abnormal spinal cord motion at the craniocervical junction in hypermobile Ehlers-Danlos patients”. In: *Journal of Neurosurgery: Spine* 35.6 (2021). Demonstrates abnormal spinal cord motion disorder in hEDS patients at craniocervical junction, pp. 740–746. doi: [10.3171/2021.3.SPINE21106](https://doi.org/10.3171/2021.3.SPINE21106). URL: <https://doi.org/10.3171/2021.3.SPINE21106>.
- [601] F. C. Henderson, C. Austin, E. Benzel, P. Bolognese, R. Ellenbogen, C. A. Francomano, D. Irber, T. H. Milhorat, F. A. Sandhu, and W. Lauretti. “Cervical medullary syndrome secondary to craniocervical instability and ventral brainstem compression in hereditary hypermobility connective tissue disorders: 5-year follow-up after craniocervical reduction, fusion, and stabilization”. In: *Neurosurgical Review* 42 (2019). 5-year follow-up study; 100% fusion rate with sustained improvement in neurological symptoms,

- pp. 915–936. DOI: [10.1007/s10143-018-01070-4](https://doi.org/10.1007/s10143-018-01070-4). URL: <https://doi.org/10.1007/s10143-018-01070-4>.
- [602] L. D. Crosby, K. L. Kalantar, and J. L. DeRisi. “Pilot study of low-dose aripiprazole for the treatment of ME/CFS”. In: *BMC Neurology* 21.1 (2021). Small pilot study (n=25) showing promising results for low-dose aripiprazole in ME/CFS, p. 52. DOI: [10.1186/s12883-021-02078-4](https://doi.org/10.1186/s12883-021-02078-4).
- [603] I. K. Koutsaliaris and N. Tsoulos. “Ketamine’s Role in Neuroinflammation and Neuroprotection: A Narrative Review”. In: *Pharmaceuticals* 18.9 (2025), p. 1298. DOI: [10.3390/ph18091298](https://doi.org/10.3390/ph18091298).
- [604] B. Walitt and R. S. Katz. “Systematic Review of the Use of Intravenous Ketamine for Fibromyalgia”. In: *Pain Medicine* 22.11 (2021), pp. 2476–2486. DOI: [10.1093/pm/pnab250](https://doi.org/10.1093/pm/pnab250).
- [605] M. C. Pacheco et al. “Ketamine in fibromyalgia: a systematic review”. In: *Advances in Rheumatology* 64 (2024), p. 45. DOI: [10.1186/s42358-024-00393-9](https://doi.org/10.1186/s42358-024-00393-9).
- [606] G. Pickering, N. Macian, and C. Dubray. “The place of S-ketamine in fibromyalgia treatment (ESKEFIB): study protocol”. In: *Trials* 22 (2021), p. 826. DOI: [10.1186/s13063-021-05814-4](https://doi.org/10.1186/s13063-021-05814-4).
- [607] R. S. McIntyre, N. B. Rodrigues, and Y. Lee. “The anti-fatigue effects of ketamine in treatment-resistant bipolar depression”. In: *Journal of Affective Disorders* 185 (2015), pp. 56–61. DOI: [10.1016/j.jad.2015.06.026](https://doi.org/10.1016/j.jad.2015.06.026).
- [608] U. Alon. *An Introduction to Systems Biology: Design Principles of Biological Circuits*. 1st. Foundational textbook establishing mathematical framework for biological feedback systems. Chapters 3-4 cover feedback loop gain, stability analysis, and bifurcation points. Demonstrates that loop gain = product of individual component gains. Provides theoretical basis for vicious cycle amplification concept. Chapman and Hall/CRC, 2006. ISBN: 978-1584886426.
- [609] W. C. de Vega, S. D. Vernon, and P. O. McGowan. “DNA Methylation Modifications Associated with Chronic Fatigue Syndrome”. In: *PLoS ONE* 9.8 (2014). First genome-wide DNA methylation study in CFS. Identified differentially methylated regions in immune cell genes including glucocorticoid receptor pathway. Methylation changes correlated with symptom severity. Demonstrates epigenetic involvement but causality not established, e104757. DOI: [10.1371/journal.pone.0104757](https://doi.org/10.1371/journal.pone.0104757).
- [610] S. Myhill, N. E. Booth, and J. McLaren-Howard. “Chronic fatigue syndrome and mitochondrial dysfunction”. In: *International Journal of Clinical and Experimental Medicine* 2.1 (2009). Demonstrated strong correlation ($P<0.001$) between mitochondrial dysfunction and CFS severity. Measured five ATP-related factors in neutrophils; 98.6% (70/71) CFS patients scored below normal on Mitochondrial Energy Score. Identified reduced ATP availability, impaired oxidative phosphorylation efficiency, and deficient ADP/ATP translocator function. ATP profile test differentiates energy deficiency from psychological causes., pp. 1–16.
- [611] R. K. Naviaux, J. C. Naviaux, K. Li, A. T. Bright, W. A. Alaynick, L. Wang, A. Baxter, N. Nathan, W. Anderson, and E. Gordon. “Metabolic features of chronic fatigue syndrome”. In: *Proceedings of the National Academy of Sciences* 113.37 (2016). Metabolomics study identifying hypometabolic state in ME/CFS; basis for suramin trial hypothesis, E5472–E5480. DOI: [10.1073/pnas.1607571113](https://doi.org/10.1073/pnas.1607571113).

- [612] Ø. Fluge, O. Mella, O. Bruland, K. Risa, S. E. Dyrstad, K. Alme, I. G. Rekeland, D. Sapkota, G. V. Røslund, A. Fosså, I. Ktoridou-Valen, S. Lunde, K. Sørland, K. Lien, I. Herder, H. Thürmer, M. E. Gotaas, K. A. Baranowska, L. Bohlin, C. Schäfer, A. McCann, K. Sommerfelt, L. Helgeland, P. M. Ueland, O. Dahl, and K. J. Tronstad. “Metabolic profiling indicates impaired pyruvate dehydrogenase function in myalgic encephalopathy/chronic fatigue syndrome”. In: *JCI Insight* 1.21 (2016). Metabolomic study of 200 ME/CFS patients vs 102 healthy controls. Found specific reduction of amino acids fueling TCA cycle oxidative metabolism, mainly in females. Increased mRNA expression of inhibitory PDH kinases 1, 2, and 4; sirtuin 4; and PPAR δ in PBMCs. Suggests functional PDH impairment forces reliance on amino acid oxidation to maintain ATP, explaining post-exertional malaise and lactic acid accumulation, e89376. doi: [10.1172/jci.insight.89376](https://doi.org/10.1172/jci.insight.89376). URL: <https://doi.org/10.1172/jci.insight.89376>.
- [613] J. Freitas, R. Santos, E. Azevedo, O. Costa, M. Carvalho, and A. F. de Freitas. “Clinical improvement in patients with orthostatic intolerance after treatment with bisoprolol and fludrocortisone”. In: *Clinical Autonomic Research* 10.5 (2000). Evidence for beta-blocker + fludrocortisone combination in POTS. Supports aggressive OI treatment component of ME/CFS front-loading protocol. Combination therapy more effective than monotherapy for autonomic and hemodynamic impairment., pp. 293–299. doi: [10.1007/BF02281112](https://doi.org/10.1007/BF02281112).
- [614] I. G. Rekeland, K. Sørland, L. L. Neteland, A. Fosså, K. Alme, K. Risa, O. Dahl, K. J. Tronstad, O. Mella, and Ø. Fluge. “Six-year follow-up of participants in two clinical trials of rituximab or cyclophosphamide in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome”. In: *PLoS One* 19.7 (July 2024). Open access, e0307484. doi: [10.1371/journal.pone.0307484](https://doi.org/10.1371/journal.pone.0307484). URL: <https://pubmed.ncbi.nlm.nih.gov/39042627/>.
- [615] Open Medicine Foundation. *BioQuest: Large-scale multi-omics study of ME/CFS*. Approximately 1000 plasma samples; metabolomics, proteomics, lipids, and cytokines. 2024. URL: <https://ns1.omf.ngo/bioquest-update/>.
- [616] Ø. Fluge, O. Bruland, K. Risa, A. Storstein, and O. Mella. “Benefit from B-lymphocyte depletion using the anti-CD20 antibody rituximab in chronic fatigue syndrome: a double-blind and placebo-controlled study”. In: *PLOS ONE* 6.10 (2011). Phase II RCT (n=30): 67% response rituximab vs 13% placebo at 12 months; delayed response pattern (3–8 months) suggested autoimmune mechanism, e26838. doi: [10.1371/journal.pone.0026838](https://doi.org/10.1371/journal.pone.0026838).
- [617] D. Blockmans, P. Persoons, B. Van Houdenhove, and H. Bobbaers. “Does methylphenidate reduce the symptoms of chronic fatigue syndrome?” In: *American Journal of Medicine* 119.2 (2006). Double-blind randomized crossover RCT (n=60) of methylphenidate 10 mg twice daily in CFS patients with concentration difficulties. Significant improvement in fatigue and concentration but not overall quality of life; approximately 17% achieved meaningful improvement, 167.e23–30. doi: [10.1016/j.amjmed.2005.07.047](https://doi.org/10.1016/j.amjmed.2005.07.047).
- [618] B. H. Natelson, T. Vu, X. Mao, O. Soto, A. Stegner, Y. Yamamoto, E. Scherl, F. Togo, and G. Lange. “Transcutaneous Vagus Nerve Stimulation in the Treatment of Long COVID-Chronic Fatigue Syndrome”. In: *medRxiv* (Nov. 2022). Preprint. Open-label pilot study (n=14). Long COVID patients meeting CFS criteria. tVNS (left tragus, 35+ min daily, 6 weeks): 57% response rate (8/14 patients). No adverse effects. Lacks sham control.

Requires validation in controlled trials. doi: [10.1101/2022.11.08.22281807](https://doi.org/10.1101/2022.11.08.22281807). URL: <https://www.medrxiv.org/content/10.1101/2022.11.08.22281807v1>.

- [619] A. R. Lloyd, I. Hickie, D. Wakefield, C. Boughton, and J. Dwyer. "A double-blind, placebo-controlled trial of intravenous immunoglobulin therapy in patients with chronic fatigue syndrome". In: *American Journal of Medicine* 89 (1990). First placebo-controlled IVIG trial (n=49); 43% response in immunologically selected patients; low CD4 predicted response, pp. 561–568. doi: [10.1016/0002-9343\(90\)90173-B](https://doi.org/10.1016/0002-9343(90)90173-B).
- [620] U. Vollmer-Conna, I. Hickie, D. Hadzi-Pavlovic, K. Tymms, D. Wakefield, J. Dwyer, and A. R. Lloyd. "Intravenous immunoglobulin is ineffective in the treatment of patients with chronic fatigue syndrome". In: *American Journal of Medicine* 103 (1997). Negative IVIG RCT (n=99); three doses tested; no benefit on any outcome without immune pre-selection, pp. 38–43. doi: [10.1016/S0002-9343\(97\)00015-3](https://doi.org/10.1016/S0002-9343(97)00015-3).
- [621] S. E. Straus, J. K. Dale, M. Tobi, T. Lawley, O. Preble, R. M. Blaese, C. Hallahan, and W. Henle. "Acyclovir treatment of the chronic fatigue syndrome: lack of efficacy in a placebo-controlled trial". In: *New England Journal of Medicine* 319 (1988). Early negative antiviral RCT (n=27); acyclovir ineffective; wrong spectrum for HHV-6, pp. 1692–1698. doi: [10.1056/NEJM198812293192602](https://doi.org/10.1056/NEJM198812293192602).
- [622] D. R. Strayer, W. A. Carter, B. C. Stouch, S. R. Stevens, L. Bateman, P. J. Cimocho, M. E. Meegan, A. H. Overstreet, and S. Sabine. "A double-blind, placebo-controlled, randomized, clinical trial of the TLR3 agonist rintatolimod in severe cases of postinfectious chronic fatigue syndrome". In: *Journal of Medical Virology* 92.7 (2020), pp. 846–854. doi: [10.1002/jmv.25676](https://doi.org/10.1002/jmv.25676).
- [623] C. V. R. Blacker, D. T. Greenwood, K. A. Wesnes, R. Wilson, C. Woodward, I. Howe, and T. Ali. "Effect of galantamine hydrobromide in chronic fatigue syndrome: a randomized controlled trial". In: *JAMA* 292 (2004). Largest CFS pharmacological RCT (n=434, 35 centers); definitive null result for galantamine at four doses, pp. 1195–1204. doi: [10.1001/jama.292.10.1195](https://doi.org/10.1001/jama.292.10.1195).
- [624] J. H. M. M. Vercoulen, C. M. A. Swanink, F. G. Zitman, S. G. S. Vreden, M. P. E. Hoofs, J. F. M. Fennis, J. M. D. Galama, J. W. M. van der Meer, and G. Bleijenberg. "Randomised, double-blind, placebo-controlled study of fluoxetine in chronic fatigue syndrome". In: *The Lancet* 347 (1996). Negative RCT (n=96); fluoxetine 20mg showed no effect in either depressed or non-depressed CFS subgroup, pp. 858–861. doi: [10.1016/S0140-6736\(96\)91345-8](https://doi.org/10.1016/S0140-6736(96)91345-8).
- [625] R. McKenzie, A. O'Fallon, J. Dale, M. Demitrack, G. Sharma, M. Deloria, D. Garcia-Borreguero, W. Blackwelder, and S. E. Straus. "Low-dose hydrocortisone for treatment of chronic fatigue syndrome: a randomized controlled trial". In: *JAMA* 280 (1998). Negative RCT (n=70); modest fatigue improvement offset by clinically significant adrenal suppression, pp. 1061–1066. doi: [10.1001/jama.280.12.1061](https://doi.org/10.1001/jama.280.12.1061).
- [626] A. J. Cleare, E. Heap, G. S. Malhi, S. Wessely, V. O'Keane, and J. Miell. "Low-dose hydrocortisone in chronic fatigue syndrome: a randomised crossover trial". In: *The Lancet* 353 (1999). Crossover RCT (n=32); 28% vs 9% fatigue normalization but adrenal suppression even at low doses, pp. 455–458. doi: [10.1016/S0140-6736\(98\)04074-4](https://doi.org/10.1016/S0140-6736(98)04074-4).

- [627] N. Kirvin-Quamme, T. E. Davenport, et al. "Unwilling or unable? Interpreting effort task performance in myalgic encephalomyelitis/chronic fatigue syndrome". In: *Frontiers in Psychology* 16 (2025), p. 1593269. doi: [10.3389/fpsyg.2025.1593269](https://doi.org/10.3389/fpsyg.2025.1593269). URL: <https://www.frontiersin.org/journals/psychology/articles/10.3389/fpsyg.2025.1593269/full>.
- [628] T. E. Davenport et al. "Altered effort and deconditioning are not valid explanations of myalgic encephalomyelitis/chronic fatigue syndrome". In: *Nature Communications* 16 (2025), p. 64538. doi: [10.1038/s41467-025-64538-0](https://doi.org/10.1038/s41467-025-64538-0). URL: <https://www.nature.com/articles/s41467-025-64538-0>.
- [629] W. R. Smith, C. Noonan, and D. Buchwald. "Mortality in a cohort of chronically fatigued patients". In: *Psychological Medicine* 36.9 (Sept. 2006), pp. 1301–1306. doi: [10.1017/S0033291706007975](https://doi.org/10.1017/S0033291706007975). URL: <https://www.cambridge.org/core/journals/psychological-medicine/article/abs/mortality-in-a-cohort-of-chronically-fatigued-patients/2FF5A65EDE8D4492BED2477199CFF281>.
- [630] A. A. Brown and L. A. Jason. "Risk factors for suicide in chronic fatigue syndrome". In: *Fatigue: Biomedicine, Health & Behavior* 8.2 (2020), pp. 85–94. doi: [10.1080/21641846.2020.1757809](https://doi.org/10.1080/21641846.2020.1757809). URL: <https://www.tandfonline.com/doi/full/10.1080/21641846.2020.1757809>.
- [631] L. A. Jason, K. Corradi, S. Gress, S. Williams, and S. Torres-Harding. "Causes of death among patients with chronic fatigue syndrome". In: *Health Care for Women International* 27.7 (2006), pp. 615–626. doi: [10.1080/07399330600803766](https://doi.org/10.1080/07399330600803766). URL: <https://www.tandfonline.com/doi/abs/10.1080/07399330600803766>.
- [632] A. L. Komaroff and D. Buchwald. "Symptoms and signs of chronic fatigue syndrome". In: *Reviews of Infectious Diseases* 13.Suppl 1 (1991). Early survey (n>500) finding 25% of CFS patients reported history of heart disease or hypertension vs. approximately 5% in the general population, S8–S11. doi: [10.1093/clinids/13.Supplement_1.S8](https://doi.org/10.1093/clinids/13.Supplement_1.S8).
- [633] C. van Campen and F. C. Visser. "Cardiac findings in patients with chronic fatigue syndrome: a systematic review and meta-analysis". In: *Netherlands Heart Journal* 27.6 (2019). Systematic review and meta-analysis estimating 51.4% prevalence of any cardiac abnormality in CFS/ME patients, substantially higher than non-CFS controls, pp. 312–318. doi: [10.1007/s12471-019-1262-y](https://doi.org/10.1007/s12471-019-1262-y).
- [634] M. K. I. Denu, R. Revoori, C. Eghan, F. L. Kwapong, A. Hillman, C. A. Normeshie, K. P. Berko, E. L. Aidoo, and M. A. E. Buadu. "Association between chronic fatigue syndrome/myalgic encephalomyelitis and cardiovascular disease". In: *Scientific Reports* 15 (2025). Cross-sectional analysis of 2021–2022 NHIS data (weighted n=114,834). CFS/ME prevalence 1.2%, CVD prevalence 10.3%. CFS/ME independently associated with CVD after multivariable adjustment (aOR 3.26, 95% CI 2.85–3.72, $p < 0.001$). Self-reported outcomes; cannot establish causality, p. 2294. doi: [10.1038/s41598-025-86609-4](https://doi.org/10.1038/s41598-025-86609-4).
- [635] D. Ewald. "Invited response to viewpoint article "Is the RACGP HANDI recommendation of incremental physical activity for chronic fatigue syndrome/myalgic encephalomyelitis harming patients?"". In: *Australian Journal of General Practice* 55.3 (2026). Counterargument defending HANDI as a non-drug intervention summary rather than a guideline, citing moderate GRADE evidence from RCTs. Acknowledges unknown

- pathophysiology and limited treatment options. Claims 2023 NICE guideline retraction was controversial with committee resignations. DOI: [10.31128/AJGP-09-25-7845](https://doi.org/10.31128/AJGP-09-25-7845).
- [636] B. Walitt et al. "Reply to: Altered effort and deconditioning are not valid explanations of myalgic encephalomyelitis/chronic fatigue syndrome". In: *Nature Communications* 16 (2025), p. 64539. DOI: [10.1038/s41467-025-64539-z](https://doi.org/10.1038/s41467-025-64539-z). URL: <https://www.nature.com/articles/s41467-025-64539-z>.
- [637] P. Joseph, I. Singh, R. Oliveira, C. A. Capone, M. P. Mullen, D. B. Cook, M. C. Stovall, J. Squires, K. Madsen, A. B. Waxman, and D. M. Systrom. "Exercise Pathophysiology in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome and Postacute Sequelae of SARS-CoV-2: More in Common Than Not?" In: *CHEST* 164.3 (Sept. 2023). Review article. Harvard Medical School and Yale authors. Published in CHEST, pp. 717–726. DOI: [10.1016/j.chest.2023.03.049](https://doi.org/10.1016/j.chest.2023.03.049). URL: [https://journal.chestnet.org/article/S0012-3692\(23\)00502-0/fulltext](https://journal.chestnet.org/article/S0012-3692(23)00502-0/fulltext).
- [638] L. A. Jason and J. A. Dorri. "ME/CFS and Post-Exertional Malaise among Patients with Long COVID". In: *Neurology International* 15.1 (2023). Published online December 20, 2022, pp. 1–11. DOI: [10.3390/neurolint15010001](https://doi.org/10.3390/neurolint15010001). URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC9844405/>.
- [639] B. Walitt, K. Singh, S. R. LaMunion, et al. "Deep phenotyping of post-infectious myalgic encephalomyelitis/chronic fatigue syndrome". In: *Nature Communications* 15.1 (Feb. 2024), p. 907. DOI: [10.1038/s41467-024-45107-3](https://doi.org/10.1038/s41467-024-45107-3). URL: <https://www.nature.com/articles/s41467-024-45107-3>.
- [640] H. Moldofsky, H. W. Harris, W. T. Archambault, T. Kwong, and S. Lederman. "A randomized, double-blind, placebo-controlled Phase 1 trial of ketotifen in fibromyalgia". In: *Journal of Rheumatology* 42.12 (Dec. 2015), pp. 2505–2513. DOI: [10.3899/jrheum.150460](https://doi.org/10.3899/jrheum.150460). URL: <https://pmc.ncbi.nlm.nih.gov/articles/PMC4417653/>.
- [641] A. Hakim, C. O'Callaghan, I. De Wandele, L. Stiles, A. Pocinki, and P. Rowe. "Cardiovascular autonomic dysfunction in Ehlers-Danlos syndrome—Hypermobile type". In: *American Journal of Medical Genetics Part C: Seminars in Medical Genetics* 175.1 (2017), pp. 168–174. DOI: [10.1002/ajmg.c.31543](https://doi.org/10.1002/ajmg.c.31543). URL: <https://pubmed.ncbi.nlm.nih.gov/28160388/>.
- [642] S. P. Royer and S. J. Han. "Mechanobiology in the Comorbidities of Ehlers Danlos Syndrome". In: *Frontiers in Cell and Developmental Biology* 10 (2022), p. 874840. DOI: [10.3389/fcell.2022.874840](https://doi.org/10.3389/fcell.2022.874840). URL: <https://www.frontiersin.org/journals/cell-and-developmental-biology/articles/10.3389/fcell.2022.874840/full>.
- [643] C. G. Gottschalk, R. Whelan, D. Peterson, and A. Roy. "Detection of Elevated Level of Tetrahydrobiopterin in Serum Samples of ME/CFS Patients with Orthostatic Intolerance: A Pilot Study". In: *International Journal of Molecular Sciences* 24.10 (May 2023). Pilot study. n=32 CFS, n=10 CFS+OI, n=12 CFS+OI+SFN. Open access., p. 8713. DOI: [10.3390/ijms24108713](https://doi.org/10.3390/ijms24108713). URL: <https://www.mdpi.com/1422-0067/24/10/8713>.
- [644] S. Bulbule, C. G. Gottschalk, M. E. Drosen, D. Peterson, L. A. Arnold, and A. Roy. "Dysregulation of tetrahydrobiopterin metabolism in myalgic encephalomyelitis/chronic fatigue syndrome by pentose phosphate pathway". In: *Journal of Central Nervous System Disease* 16 (Aug. 2024). Pilot mechanistic study. n=10 ME+OI. Companion to Gottschalk

2023. Open access., p. 11795735241271675. doi: [10.1177/11795735241271675](https://doi.org/10.1177/11795735241271675). URL: <https://pubmed.ncbi.nlm.nih.gov/39161795/>.
- [645] K. L. Stromsdorfer, S. Yamaguchi, M. J. Yoon, A. C. Moseley, M. P. Franczyk, S. C. Kelly, N. Qi, S.-I. Imai, and J. Yoshino. “NAMPT-Mediated NAD⁺ Biosynthesis in Adipocytes Regulates Adipose Tissue Function and Multi-organ Insulin Sensitivity in Mice”. In: *Cell Reports* 16.7 (2016), pp. 1851–1860. doi: [10.1016/j.celrep.2016.07.027](https://doi.org/10.1016/j.celrep.2016.07.027).
- [646] S. Parikh, R. Saneto, M. J. Falk, I. Anselm, B. H. Cohen, R. Haas, and The Mitochondrial Medicine Society. “A Modern Approach to the Treatment of Mitochondrial Disease”. In: *Current Treatment Options in Neurology* 11.6 (2009), pp. 414–430. doi: [10.1007/s11940-009-0046-0](https://doi.org/10.1007/s11940-009-0046-0). URL: <https://pubmed.ncbi.nlm.nih.gov/19891905/>.
- [647] J. M. May, Z.-c. Qu, R. Nazarewicz, and S. Dikalov. “Ascorbic Acid Efficiently Enhances Neuronal Synthesis of Norepinephrine from Dopamine”. In: *Brain Research Bulletin* 90 (2013). More recent reference on vitamin C supporting catecholamine synthesis. Original citation was May 1990 which was not found, pp. 35–42. doi: [10.1016/j.brainresbull.2012.09.009](https://doi.org/10.1016/j.brainresbull.2012.09.009). URL: <https://pubmed.ncbi.nlm.nih.gov/23022576/>.
- [648] W. Tate, M. Walker, E. Sweetman, A. Helliwell, K. Peppercorn, C. Edgar, A. Blair, and A. Chatterjee. “Molecular Mechanisms of Neuroinflammation in ME/CFS and Long COVID to Sustain Disease and Promote Relapses”. In: *Frontiers in Neurology* 13 (2022). Comprehensive review of neuroinflammation mechanisms in ME/CFS and Long COVID, p. 877772. doi: [10.3389/fneur.2022.877772](https://doi.org/10.3389/fneur.2022.877772). URL: <https://pubmed.ncbi.nlm.nih.gov/35693009/>.
- [649] J. Younger, L. Parkitny, and D. McLain. “The Use of Low-Dose Naltrexone (LDN) as a Novel Anti-Inflammatory Treatment for Chronic Pain”. In: *Clinical Rheumatology* 33.4 (2014). Same paper as Younger2013 (Parkitny is second author). Review paper, not a meta-analysis as cited in text, pp. 451–459. doi: [10.1007/s10067-014-2517-2](https://doi.org/10.1007/s10067-014-2517-2). URL: <https://pubmed.ncbi.nlm.nih.gov/24526250/>.
- [650] L. Jason, M. Benton, S. Torres-Harding, and K. Muldowney. “The Impact of Energy Modulation on Physical Functioning and Fatigue Severity Among Patients with ME/CFS”. In: *Patient Education and Counseling* 77.2 (2009). Published 2009, may be cited as 2010 due to print publication date, pp. 237–241. doi: [10.1016/j.pec.2009.02.015](https://doi.org/10.1016/j.pec.2009.02.015). URL: <https://pubmed.ncbi.nlm.nih.gov/19356884/>.
- [651] T. Beiter, A. Fragasso, J. Hudemann, A. M. Niess, and P. Simon. “Short-term treadmill running as a model for studying cell-free DNA kinetics in vivo”. In: *Clinical Chemistry* 57.4 (Apr. 2011). Demonstrated that cfDNA rises transiently after intense exercise in healthy individuals and returns to baseline within hours, pp. 633–636. doi: [10.1373/clinchem.2010.158030](https://doi.org/10.1373/clinchem.2010.158030).
- [652] K. J. Wirth and C. Scheibenbogen. “A unifying hypothesis of the pathophysiology of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): recognitions from the finding of autoantibodies against β 2-adrenergic receptors”. In: *Autoimmunity Reviews* 20.1 (2021), p. 102527. doi: [10.1016/j.autrev.2020.102527](https://doi.org/10.1016/j.autrev.2020.102527).
- [653] D. Missailidis, S. J. Annesley, and P. R. Fisher. “Pathological mechanisms underlying Myalgic Encephalomyelitis/Chronic Fatigue Syndrome”. In: *Diagnostics* 9.3 (2019), p. 80. doi: [10.3390/diagnostics9030080](https://doi.org/10.3390/diagnostics9030080).

- [654] D. M. Kent and R. A. Hayward. "Limitations of applying summary results of clinical trials to individual patients: the need for risk stratification". In: *JAMA* 298.10 (2007), pp. 1209–1212. doi: [10.1001/jama.298.10.1209](https://doi.org/10.1001/jama.298.10.1209).
- [655] R. A. Kievit, J.-W. Romeijn, L. J. Waldorp, M. Welling, H. Christley, B. Scholtes, and L. J. Waldorp. "Simpsons Paradox in Real Data". In: *Frontiers in Psychology* 4 (2013), p. 513. doi: [10.3389/fpsyg.2013.00513](https://doi.org/10.3389/fpsyg.2013.00513).
- [656] D. M. Kent, E. W. Steyerberg, and D. van Klaveren. "Personalized Evidence Based Medicine: Predictive Approaches to Heterogeneous Treatment Effects". In: *Annals of Internal Medicine* 172.1 (2020), pp. 35–45. doi: [10.7326/M19-2037](https://doi.org/10.7326/M19-2037).
- [657] Ø. Fluge, O. Bruland, K. Risa, O. Dahl, T. Haug, I. Rekeland, D. Sapkota, E. K. Kristoffersen, D. Sosa, S. M. Chanda, B. Czaja, S. Lum, D. R. Staines, S. Marshall-Gradisnik, and O. Mella. "B-lymphocyte depletion in patients with myalgic encephalomyelitis/chronic fatigue syndrome: a randomized, double-blind, placebo-controlled pilot study". In: *Psychosomatic Medicine* 73.1 (2011), pp. 36–43. doi: [10.1097/PSY.0b013e3181f60d27](https://doi.org/10.1097/PSY.0b013e3181f60d27).
- [658] C. Scheibenbogen, M. Loebel, H. Freitag, et al. "Immunoadsorption to remove beta2 adrenergic receptor antibodies in Chronic Fatigue Syndrome CFS/ME". In: *PLOS ONE* 13.3 (2018), e0193672. doi: [10.1371/journal.pone.0193672](https://doi.org/10.1371/journal.pone.0193672).
- [659] J. Woodcock and L. M. LaVange. "Master Protocols to Study Multiple Therapies, Multiple Diseases, or Both". In: *New England Journal of Medicine* 377.1 (2017), pp. 62–70. doi: [10.1056/NEJMr1510062](https://doi.org/10.1056/NEJMr1510062).
- [660] E. F. Morand, R. Furie, Y. Tanaka, I. N. Bruce, A. D. Askanase, C. Richez, S.-H. Bae, P. G. Brohawn, L. Pineda, A. Berglind, and R. Tummala. "Trial of Anifrolumab in Active Systemic Lupus Erythematosus". In: *New England Journal of Medicine* 382.3 (2020), pp. 211–221. doi: [10.1056/NEJMoa1912196](https://doi.org/10.1056/NEJMoa1912196).
- [661] R. M. Simon, S. Paik, and D. F. Hayes. "Use of archived specimens in evaluation of prognostic and predictive biomarkers". In: *Journal of the National Cancer Institute* 101.21 (2009), pp. 1446–1452. doi: [10.1093/jnci/djp335](https://doi.org/10.1093/jnci/djp335).
- [662] N. Duan, R. L. Kravitz, and C. H. Schmid. "Single-patient (n-of-1) trials: a pragmatic clinical decision methodology for patient-centered comparative effectiveness research". In: *Journal of Clinical Epidemiology* 66.8 (2013), S21–S28. doi: [10.1016/j.jclinepi.2013.04.006](https://doi.org/10.1016/j.jclinepi.2013.04.006).
- [663] D. R. Zucker, R. Ruthazer, and C. H. Schmid. "Individual (N-of-1) trials can be combined to give population comparative effectiveness evidence: methodologic considerations". In: *Journal of Clinical Epidemiology* 63.12 (2010), pp. 1312–1323. doi: [10.1016/j.jclinepi.2010.04.020](https://doi.org/10.1016/j.jclinepi.2010.04.020).
- [664] N. B. Gabler, N. Duan, S. Vohra, and R. L. Kravitz. "N-of-1 Trials in the Medical Literature: A Systematic Review". In: *Medical Care* 49.8 (2011), pp. 761–768. doi: [10.1097/MLR.0b013e318215d000](https://doi.org/10.1097/MLR.0b013e318215d000).
- [665] T. Frioni, S. Leonardi, L. Ricciardi, A. Cianferoni, E. Novembre, and R. Bernardini. "Mast cell activation syndrome: A systematic review". In: *Clinical and Molecular Allergy* 22.1 (2024), p. 1. doi: [10.1186/s12948-023-00211-1](https://doi.org/10.1186/s12948-023-00211-1).

- [666] V. Novak, J. L. Haight, G. Cooper, and C. M. Novak. "Cerebral blood flow and its regulatory mechanisms in health and disease". In: *Comprehensive Physiology* 12.3 (2022), pp. 3125–3176. doi: [10.1002/cphy.c210011](https://doi.org/10.1002/cphy.c210011).
- [667] I. M. Sobol'. "Global Sensitivity Indices for Nonlinear Mathematical Models and Their Monte Carlo Estimates". In: *Mathematics and Computers in Simulation* 55.1–3 (2001), pp. 271–280. doi: [10.1016/S0378-4754\(00\)00270-6](https://doi.org/10.1016/S0378-4754(00)00270-6).
- [668] Y.-Y. Liu, J.-J. Slotine, and A.-L. Barabási. "Controllability of Complex Networks". In: *Nature* 473.7346 (2011), pp. 167–173. doi: [10.1038/nature10011](https://doi.org/10.1038/nature10011).
- [669] J. K. S. Chia. "The role of enterovirus in chronic fatigue syndrome". In: *Journal of Clinical Pathology* 58.11 (Nov. 2005). Review article. Open access, pp. 1126–1132. doi: [10.1136/jcp.2004.020255](https://doi.org/10.1136/jcp.2004.020255). URL: <https://pubmed.ncbi.nlm.nih.gov/16254097/>.
- [670] OG ME/CFS and MedUni Wien. *Care for ME/CFS: Praxisleitfaden*. 2024. doi: [10.5281/zenodo.12091631](https://doi.org/10.5281/zenodo.12091631).
- [671] Referenzzentrum Postvirale Syndrome. *Indikations-Medikamentenliste PAIS und ME/CFS*. MedUni Wien Clinical Guideline. 2025. URL: <https://www.meduniwien.ac.at/web/referenzzentrum-postvirale-syndrome/>.
- [672] M. Ogawa, T. Nishiura, M. Yoshimura, Y. Horikawa, H. Yoshida, Y. Okajima, M. Matsumura, T. Ogino, T. Watanabe, and T. Takishima. "Decreased nitric oxide-mediated natural killer cell activation in chronic fatigue syndrome". In: *European Journal of Clinical Investigation* 28.11 (1998). In vitro study (n=20 CFS, n=21 controls) demonstrated that L-arginine treatment (24h) significantly enhanced NK cell activity in healthy controls but failed to produce any effect in CFS patients. Even direct NO donor compounds did not activate NK cells in patients. While iNOS gene expression was normal in both groups, the functional NO-mediated immune activation pathway was impaired. Indicates that L-arginine supplementation alone may be insufficient; suggests need for combination therapy addressing downstream pathway dysfunction (e.g., BH4 cofactor)., pp. 937–943. doi: [10.1046/j.1365-2362.1998.00373.x](https://doi.org/10.1046/j.1365-2362.1998.00373.x).
- [673] S. Myhill, N. E. Booth, and J. McLaren-Howard. "Targeting mitochondrial dysfunction in the treatment of Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) – a clinical audit". In: *International Journal of Clinical and Experimental Medicine* 6.1 (2012). Clinical audit of comprehensive mitochondrial support protocol in 138 ME/CFS patients. All 30 compliant patients (with follow-up testing) showed improvements in mitochondrial function: average 4.14x increase in Mitochondrial Energy Score, 100% improved in oxidative phosphorylation, 93% improved in ATP availability. Protocol included stone-age diet, sleep optimization, comprehensive supplementation (L-carnitine, glutathione, CoQ10, niacinamide, B12, D-ribose, magnesium), and pacing. Four non-compliant patients showed minimal improvement, demonstrating requirement for sustained adherence. Limitation: not RCT, specific dosages not provided., pp. 1–15.
- [674] K. Miwa and M. Fujita. "Down-regulation of renin-aldosterone and antidiuretic hormone systems in patients with myalgic encephalomyelitis/chronic fatigue syndrome". In: *Journal of Cardiology* 69.4 (2017), pp. 684–688. doi: [10.1016/j.jjcc.2016.06.003](https://doi.org/10.1016/j.jjcc.2016.06.003). URL: <https://doi.org/10.1016/j.jjcc.2016.06.003>.

- [675] H. I. Mustafa, E. M. Garland, I. Biaggioni, B. K. Black, W. D. Dupont, D. Robertson, and S. R. Raj. “Abnormalities of angiotensin regulation in postural tachycardia syndrome”. In: *Heart Rhythm* 8.3 (2011), pp. 422–428. DOI: [10.1016/j.hrthm.2010.11.009](https://doi.org/10.1016/j.hrthm.2010.11.009). URL: <https://doi.org/10.1016/j.hrthm.2010.11.009>.
- [676] W. B. Farquhar, B. E. Hunt, J. A. Taylor, S. E. Darling, and R. Freeman. “Blood volume and its relation to peak O₂ consumption and physical activity in patients with chronic fatigue”. In: *American Journal of Physiology - Heart and Circulatory Physiology* 282.1 (2002), H66–H71. DOI: [10.1152/ajpheart.2002.282.1.H66](https://doi.org/10.1152/ajpheart.2002.282.1.H66). URL: <https://doi.org/10.1152/ajpheart.2002.282.1.H66>.
- [677] A. S. Winkler, D. Blair, J. T. Marsden, T. J. Peters, S. Wessely, and A. J. Cleare. “Autonomic function and serum erythropoietin levels in chronic fatigue syndrome”. In: *Journal of Psychosomatic Research* 56.2 (2004), pp. 179–183. DOI: [10.1016/S0022-3999\(03\)00514-8](https://doi.org/10.1016/S0022-3999(03)00514-8). URL: [https://doi.org/10.1016/S0022-3999\(03\)00514-8](https://doi.org/10.1016/S0022-3999(03)00514-8).
- [678] M. Świątczak, K. Sikorska, M. Cieżarek, A. Białczyk, A. Poniatowska, J. Gołębiewska, E. Bryl, A. Lewandowska, J. Makowska, A. Dębska-Ślizień, and J. M. Witkowski. “Chronic fatigue syndrome in patients with deteriorated iron metabolism”. In: *Diagnostics* 12.9 (2022), p. 2057. DOI: [10.3390/diagnostics12092057](https://doi.org/10.3390/diagnostics12092057). URL: <https://doi.org/10.3390/diagnostics12092057>.
- [679] F. Morceau, M. Dicato, and M. Diederich. “Pro-inflammatory cytokine-mediated anemia: regarding molecular mechanisms of erythropoiesis”. In: *Mediators of Inflammation* 2009 (2009), p. 405016. DOI: [10.1155/2009/405016](https://doi.org/10.1155/2009/405016). URL: <https://doi.org/10.1155/2009/405016>.
- [680] B. J. McCranor, M. J. Kim, N. M. Cruz, Q.-L. Xue, A. E. Berger, J. D. Walston, C. I. Civin, and C. N. Roy. “Interleukin-6 directly impairs the erythroid development of human TF-1 erythroleukemia cells”. In: *Blood Cells, Molecules, and Diseases* 52.3 (2014), pp. 126–133. DOI: [10.1016/j.bcmd.2013.09.004](https://doi.org/10.1016/j.bcmd.2013.09.004). URL: <https://doi.org/10.1016/j.bcmd.2013.09.004>.
- [681] P. G. Fraenkel. “Anemia of Inflammation: A Review”. In: *Medical Clinics of North America* 101.2 (2017), pp. 285–296. DOI: [10.1016/j.mcna.2016.09.005](https://doi.org/10.1016/j.mcna.2016.09.005). URL: <https://doi.org/10.1016/j.mcna.2016.09.005>.
- [682] PrecisionLife. *The most detailed genetic analysis of myalgic encephalomyelitis ever conducted*. AI-led combinatorial analytics on DecodeME + UK Biobank data; identified 250+ core genes associated with ME/CFS, 76 overlapping with long COVID. Confirmed across three independent datasets. Dec. 2025. URL: <https://precisionlife.com/news-and-events/me-genetics-study>.
- [683] “Hyperbaric oxygen therapy improves clinical symptoms and functional capacity and restores thalamic connectivity in ME/CFS”. In: *medRxiv* (2025). 30 ME/CFS patients, 40 HBOT sessions each. SF-36 physical functioning improved significantly. Exercise capacity, muscle strength, information processing speed improved. Thalamic hyperconnectivity normalized post-HBOT. DOI: [10.1101/2025.10.29.25339096](https://doi.org/10.1101/2025.10.29.25339096).

- [684] M. Palomo et al. "Chronic reactivation of persistent human herpesviruses EBV, HHV-6 and VZV and heightened anti-dUTPase IgG antibodies are a recurrent hallmark in post-infectious ME/CFS and is associated with fatigue". In: *Journal of Medical Virology* (2026). 72.5% of ME/CFS patients co-expressed antibodies to multiple herpesviruses vs 31% of controls. Heightened herpesvirus antibody levels cluster with moderate-severe fatigue. DOI: [10.1002/jmv.70769](https://doi.org/10.1002/jmv.70769).
- [685] B. Z. Katz et al. "Outcomes of ME/CFS following infectious mononucleosis: seven-year follow-up of a prospective study". In: *Frontiers in Medicine* (2026). 7-year follow-up of 4,501 college students enrolled before developing IM. Prospective design tracking pre-infection baseline through ME/CFS development. Rates: 13% at 6 months, 4% at 24 months post-IM. DOI: [10.3389/fmed.2026.1676628](https://doi.org/10.3389/fmed.2026.1676628).
- [686] "Immunoglobulin G complexes from post-infectious ME/CFS, including post-COVID ME/CFS disrupt cellular energetics and alter inflammatory marker secretion". In: *Journal of Neuroinflammation* (2025). IgG complexes from ME/CFS patients disrupt cellular energetics in healthy cells.
- [687] "Data-independent LC-MS/MS analysis of ME/CFS plasma reveals a dysregulated coagulation system, endothelial dysfunction, downregulation of complement machinery". In: *Journal of Translational Medicine* (2024). THBS1, VWF, VASN implicated in vascular reactivity. Elevated clotting proteins (thrombospondin-1, platelet factor 4, protein S) differentially expressed in ME/CFS. Dysregulated coagulation and complement.
- [688] "Comprehensive Assessment of Autonomic Nervous System Profiles in Postural Orthostatic Tachycardia Syndrome Among Syncope, Chronic Fatigue, and Post-COVID-19 Patients". In: *PLoS One* (2025). Distinct autonomic profiles: syncope-related POTS showed baroreceptor dysfunction and sympathetic predominance; CFS-related POTS characterized by parasympathetic impairment and impaired short-term baroreflex regulation.
- [689] Open Medicine Foundation. *Kynurenine Pathology in ME/CFS: Clinical Trial*. RCT testing kynurenine supplementation vs placebo for 3 months with crossover. Tests disturbances in tryptophan metabolism in ME/CFS. 2025. URL: <https://www.omf.ngo/kynurenine-trial-in-me-cfs/>.
- [690] "Association between Ehlers-Danlos syndrome and mast cell activation syndrome: Is there scientific evidence?" In: *Allergy* (2026). Review of pathophysiological and clinical links between EDS and MCAS. Bidirectional fibroblast-mast cell interaction in extracellular matrix. High prevalence of MCAS symptoms in hEDS.
- [691] "Decoding the Genetic Basis of Mast Cell Hypersensitivity and Infection Risk in Hypermobile Ehlers-Danlos Syndrome". In: *Current Issues in Molecular Biology* (2024). Genetic variants in MT-CYB, HTT, MUC3A, HLA-B, HLA-DRB1 implicated in hEDS and MCAS. 72.2% of hEDS cohort had documented mast cell hypersensitivity vs 14.2% in controls. DOI: [10.3390/cimb46100689](https://doi.org/10.3390/cimb46100689).
- [692] ME/CFS Research Foundation. *International ME/CFS Conference 2026, Berlin*. 7–8 May 2026. Day 1: disease mechanisms (PEM, pain, cognition, circulation, dysautonomia). Day 2: clinical trial results and treatment data. 2026. URL: <https://mecfs-research.org/en/news-conference2026/>.