

Research Question

- Mesoscale eddies detected by sea surface height (SSH) fields have been observed to display coherent temperature (SST) structures, which lead to the term of **SSH-SST coherent eddies**. For example,
- AEs** ---> **SSHA+** ---> **SSTA+**
CEs ---> **SSHA-** ---> **SSTA-**
- However, recent statistics have shown that ~20% eddies are unconventionally **warm CEs and cold AEs** (e.g., Moschos et al., 2022), which refer to the definition of **SSH-SST incoherent eddies**. For example,
- AEs** ---> **SSHA+** ---> **SSTA-**
CEs ---> **SSHA-** ---> **SSTA+**
- In this study, we focus on proportions and mechanisms of both eddy types in the **North Indian Ocean and Global Ocean** to understand air-sea coupling induced by eddies, and related heat transport.




Mesoscale meridional heat transport inferred from sea surface observations

Yanxu Chen

Woods Hole Oceanographic Institution

Contact: yanxu.chen@whoi.edu




Data and methods

Data:


- The air-sea flux datasets are sourced from **OAFlux2**, the second generation of OAFlux sponsored by NASA's MEaSUREs program (Yu, 2023). These datasets consist of 1/4 degree gridded satellite-derived products, including turbulent heat fluxes, wind and wind stress etc.
- The Mesoscale Eddy Trajectories Atlas (**META3.2 DT**) is derived from the altimetric absolute dynamic topography (ADT) (Pegliasco et al., 2022).

Procedure:

- For time: apply a band-pass Butterworth window to preserve **7-90 days**;
- For space: employ a moving average Hann window to filter spatial scales larger than **600 km**;
- Subtract the large scale from time-filtered maps to obtain anomaly signals of the mesoscale.





SSH-SST coherent and incoherent eddies



Mesoscale meridional heat transport carried by geostrophic components is **10 times** larger than (and opposite in direction to) that of the wind-driven Ekman components.

An offset between SSH-SST coherent and incoherent eddies in the Ekman heat transport is apparent, whereas the geostrophic heat transport is contained within coherent eddies.



Conclusions

In the North Indian Ocean, semi-annual reversal of monsoon winds influences the proportion of coherent and incoherent eddies.

The combination of coherent and incoherent eddies leads to:

- monopolar structure similar to coherent eddies;
- compensation resulting in null net flux;
- dipolar pattern known as eddy-stirring effect.

Mechanisms of incoherent eddies might include:

- continuous air-sea heat exchange along eddy pathways;
- subsurface diffusion below the mixed layer;
- interior mode water resulted from water mass subduction.

The geostrophic component of meridional heat transport (MHT) at the mesoscale is 10 times larger than the Ekman component.

SSH-SST coherent eddies dominate the spatial patterns of MHT at the mesoscale.

References:

- Chen and Yu (2024). Mesoscale Meridional Heat Transport Inferred From Sea Surface Observations. *Geophysical Research Letters*.
- Chen and Yu (2024). Signature of Mesoscale Eddies on Air-Sea Heat Fluxes in the North Indian Ocean. *JGR: Oceans*.