
VE320 – Summer 2024

Introduction to Semiconductor Devices

Instructor: Yaping Dan (但亚平)
yaping.dan@sjtu.edu.cn

Chapter 2 Introduction to Quantum Mechanics

Outline

2.1 2nd order differential equations and waves

2.2 Historic events in developing quantum mechanics

2.3 A case study

2.4 Electrons in infinite quantum well

2.5 Electrons in finite quantum well

2.6 Electrons in an atom

Outline

2.1 2nd order differential equations and waves

2.2 Historic events in developing quantum mechanics

2.3 A case study

2.4 Electrons in infinite quantum well

2.5 Electrons in finite quantum well

2.6 Electrons in an atom

2.1 2nd order differential equations and waves

$$\frac{\partial^2 y}{\partial x^2} = k^2 y$$

General solution: $y = Ae^{bx}$

Plug into the equation: $b^2 Ae^{bx} = k^2 Ae^{bx}$

$$\Rightarrow b = \pm k$$

$$\Rightarrow y = A_1 e^{kx} + A_2 e^{-kx}$$

2.1 2nd order differential equations and waves

$$\frac{\partial^2 y}{\partial x^2} = k^2 y$$

General solution: $y = Ae^{bx}$

Plug into the equation: $b^2 Ae^{bx} = k^2 Ae^{bx}$

$$\Rightarrow b = \pm k$$

$$\Rightarrow y = A_1 e^{kx} + A_2 e^{-kx}$$

$$\frac{\partial^2 y}{\partial x^2} = -k^2 y$$

General solution: $y = Ae^{bx}$

Plug into the equation: $b^2 Ae^{bx} = -k^2 Ae^{bx}$

$$\Rightarrow b = \pm ki$$

$$\Rightarrow y = A_1 e^{ikx} + A_2 e^{-ikx}$$

2.1 2nd order differential equations and waves



$$\frac{\partial^2 y}{\partial x^2} = k^2 y$$

General solution: $y = Ae^{bx}$



Plug into the equation: $b^2 Ae^{bx} = k^2 Ae^{bx}$

$$\Rightarrow b = \pm k$$

$$\Rightarrow y = A_1 e^{kx} + A_2 e^{-kx}$$



2.1 2nd order differential equations and waves



$$\frac{\partial^2 y}{\partial x^2} = -k^2 y$$



General solution: $y = Ae^{bx}$

Plug into the equation: $b^2 Ae^{bx} = -k^2 Ae^{bx}$

$$\Rightarrow b = \pm ki$$

$$\Rightarrow y = A_1 e^{ikx} + A_2 e^{-ikx}$$

2.1 2nd order differential equations and waves



1. Give a wave propagating along x with a wavelength λ_0 , please write the static 2nd order differential equation that governs the behavior of this wave.

2.1 2nd order differential equations and waves

- Electromagnetic (EM) wave

$$\left\{ \begin{array}{l} \nabla \cdot E = 4\pi\rho \\ \nabla \times E = -\frac{1}{\sqrt{\mu\varepsilon}} \frac{\partial B}{\partial t} \\ \nabla \cdot B = 0 \\ \nabla \times B = \frac{4\pi}{c} J + \frac{1}{\sqrt{\mu\varepsilon}} \frac{\partial E}{\partial t} \end{array} \right.$$




James Maxwell

Static differential equation describing the electromagnetic wave

$$\frac{\partial^2 E_x}{\partial x^2} = -\omega^2 \mu \varepsilon E_x = -(\omega \sqrt{\mu \varepsilon})^2 E_x \quad E_x = E_{x0} e^{-i\omega \sqrt{\mu \varepsilon} x}$$

2.1 2nd order differential equations and waves

- Electromagnetic (EM) wave



Outline

2.1 2nd order differential equations and waves

2.2 Historic events in developing quantum mechanics

2.3 A case study


2.4 Electrons in infinite quantum well

2.5 Electrons in finite quantum well

2.6 Electrons in an atom

2.2 Historic events in developing quantum mechanics

① Speed of light in 1862 $v = 2.98 \times 10^8 \text{ m/s}$



Leon Foucault

Rotation mirror

2.2 Historic events in developing quantum mechanics

② Maxwell Equations in the year 1865

$$\left\{ \begin{array}{l} \nabla \cdot E = 4\pi\rho \\ \nabla \times E = -\frac{1}{\sqrt{\mu\varepsilon}} \frac{\partial B}{\partial t} \\ \nabla \cdot B = 0 \\ \nabla \times B = \frac{4\pi}{c} J + \frac{1}{\sqrt{\mu\varepsilon}} \frac{\partial E}{\partial t} \end{array} \right.$$



James Maxwell

Static differential equation describing the electromagnetic wave

$$\frac{\partial^2 E_x}{\partial x^2} = -\omega^2 \mu \varepsilon E_x$$

$$E_x = E_{x0} e^{-i\omega \sqrt{\mu\varepsilon} x}$$

Light is an electromagnetic wave!

$$v = \frac{1}{\sqrt{\mu\varepsilon}} = 2.99 \times 10^8 \text{ m/s}$$

2.2 Historic events in developing quantum mechanics

③ Light wave-particle duality in 1905





Photo-electric experiment




Albert Einstein
Nobel Prize 1921

- ☐ Light frequency higher than a certain frequency \rightarrow electron ejection
- ☐ Not a function of light intensity

2.2 Historic events in developing quantum mechanics

③ Light wave-particle duality in 1905



□ Light wave is a particle: $h\nu = K_{\max} + W_c$

2.2 Historic events in developing quantum mechanics

③ Light wave-particle duality in 1905





Photo-electric experiment



Albert Einstein
Nobel Prize 1921

$$E = h\nu = \hbar\omega$$

$$E = mc^2$$

$$p = mc = \frac{E}{c} = \frac{h\nu}{c} = \frac{h}{\lambda} = \hbar k$$

Light is a particle!

$$k = \frac{2\pi}{\lambda}$$

2.2 Historic events in developing quantum mechanics

④ Matter wave hypothesis in 1924

$$E = \frac{1}{2}mv^2$$

$$p = mv$$

Matter particle

$$E = h\nu = \hbar\omega$$

$$p = \frac{h}{\lambda} = \hbar k$$



Wave- particle



Louis Victor de Broglie
Nobel Prize 1929

2.2 Historic events in developing quantum mechanics

④ Matter wave hypothesis in 1924



Outline

2.1 2nd order differential equations and waves

2.2 Historic events in developing quantum mechanics

2.3 A case study

2.4 Electrons in infinite quantum well

2.5 Electrons in finite quantum well

2.6 Electrons in an atom

2.3 A case study


- Maxwell Equation in the year 1865
- Light wave-particle duality in 1905
- Matter wave hypothesis in 1924

$$\frac{\partial^2 y}{\partial x^2} = -k^2 y$$

$$E = \frac{1}{2}mv^2 \quad E = h\nu = \hbar\omega$$

$$p = mv \quad p = \frac{h}{\lambda} = \hbar k$$

Quiz #1:



Can you find a differential equation that governs the wave behavior of electrons?

2.3 A case study

$$E = qV_0 = \frac{1}{2}mv^2$$

$$p = mv$$

$$E = h\nu = \hbar\omega$$

$$p = \frac{h}{\lambda} = \hbar k$$

$$v = \sqrt{\frac{2qV_0}{m}}$$


$$k = \frac{m}{\hbar} v = \sqrt{\frac{2mqV_0}{\hbar^2}}$$

$$\frac{\partial^2 \Psi}{\partial x^2} = -k^2 \Psi = -\frac{2mqV_0}{\hbar^2} \Psi$$

$$\Rightarrow -\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} = E\Psi$$

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V(x)\Psi = E\Psi$$

Schrodinger Equation !



Can you find a differential equation that governs the wave behavior of electrons?

Outline

2.1 2nd order differential equations and waves

2.2 Historic events in developing quantum mechanics

2.3 A case study

2.4 Electrons in infinite quantum well

2.5 Electrons in finite quantum well

2.6 Electrons in an atom

2.4 Electrons in Infinite Quantum Well

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V(x)\Psi = E\Psi$$


Conditions:

for $x \leq 0, x \geq a$

$$V(x) = +\infty;$$

for $0 < x < a$

$$V(x) = 0$$



2.4 Electrons in Infinite Quantum Well

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V(x)\Psi = E\Psi$$


Conditions:

for $x \leq 0, x \geq a$

$$V(x) = +\infty; \Rightarrow \Psi(x) = 0$$

for $0 < x < a$

$$V(x) = 0$$



$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} = E\Psi$$



$$\frac{\partial^2 \Psi}{\partial x^2} = -\frac{2mE}{\hbar^2} \Psi$$



$$\frac{\partial^2 \Psi}{\partial x^2} = -k^2 \Psi$$

$$k = \sqrt{\frac{2mE}{\hbar^2}}$$

2.4 Electrons in Infinite Quantum Well

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} = E\Psi$$



$$\frac{\partial^2 \Psi}{\partial x^2} = -\frac{2mE}{\hbar^2} \Psi$$



$$\frac{\partial^2 \Psi}{\partial x^2} = -k^2 \Psi$$

$$k = \sqrt{\frac{2mE}{\hbar^2}}$$

The general solution

$$\Psi(x) = Ae^{-ikx} + Be^{ikx}$$

Boundary conditions:

$$\begin{cases} \Psi(x)|_{x=a,0} = 0 \\ \int_0^a \Psi(x)\Psi^*(x)dx = 1 \end{cases}$$

2.4 Electrons in Infinite Quantum Well

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} = E\Psi$$



$$\frac{\partial^2 \Psi}{\partial x^2} = -\frac{2mE}{\hbar^2} \Psi$$



$$\frac{\partial^2 \Psi}{\partial x^2} = -k^2 \Psi$$

$$k = \sqrt{\frac{2mE}{\hbar^2}}$$

The general solution

$$\Psi(x) = Ae^{-ikx} + Be^{ikx}$$

Boundary conditions:

$$\begin{cases} \Psi(x)|_{x=0} = 0 \\ \int_0^a \Psi(x)\Psi^*(x)dx = 1 \end{cases}$$

$$\Psi(x) = Ae^{-ik0} + Be^{ik0} = 0 \Rightarrow A = -B$$

$$\Psi(x) = Ae^{-ika} + Be^{ika} = 0 \Rightarrow \sin(ka) = 0$$

$$ka = n\pi \quad \text{where } n = 0, \pm 1, \pm 2, \dots$$

2.4 Electrons in Infinite Quantum Well

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} = E\Psi$$



$$\frac{\partial^2 \Psi}{\partial x^2} = -\frac{2mE}{\hbar^2} \Psi$$



$$\frac{\partial^2 \Psi}{\partial x^2} = -k^2 \Psi$$

$$k = \sqrt{\frac{2mE}{\hbar^2}}$$

The general solution

$$\Psi(x) = Ae^{-ikx} + Be^{ikx}$$

Boundary conditions:

$$\begin{cases} \Psi(x)|_{x=a,0} = 0 \\ \int_0^a \Psi(x)\Psi^*(x)dx = 1 \end{cases} \Rightarrow \begin{aligned} k &= \frac{n\pi}{a} \quad n = 0, \pm 1, \pm 2, \dots \\ E &= \frac{k^2 \hbar^2}{2m} = \frac{n^2 \pi^2 \hbar^2}{2ma^2} \end{aligned}$$

2.4 Electrons in Infinite Quantum Well

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V(x)\Psi = E\Psi$$


Conditions:

$$\text{for } x \leq 0, x \geq a$$

$$V(x) = +\infty; \Psi(x) = 0$$

$$\text{for } 0 < x < a$$

$$V(x) = 0$$





$$E = \frac{k^2 \hbar^2}{2m} = \frac{n^2 \pi^2 \hbar^2}{2ma^2}$$

2.4 Electrons in Infinite Quantum Well

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V(x)\Psi = E\Psi$$




$$E = \frac{k^2 \hbar^2}{2m}$$



$$E = \frac{k^2 \hbar^2}{2m} = \frac{n^2 \pi^2 \hbar^2}{2ma^2}$$


$$k = \frac{n\pi}{a} \quad n = 0, \pm 1, \pm 2, \dots$$

2.4 Electrons in Infinite Quantum Well




2.4 Electrons in Infinite Quantum Well

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V(x)\Psi = E\Psi$$



2-dimentional Quantum well



http://en.wikipedia.org/wiki/Particle_in_a_box

$$E = \frac{k^2 \hbar^2}{2m} = \frac{n^2 \pi^2 \hbar^2}{2ma^2}$$

Outline

2.1 2nd order differential equations and waves

2.2 Historic events in developing quantum mechanics

2.3 A case study

2.4 Electrons in infinite quantum well

2.5 Electrons in finite quantum well

2.6 Electrons in an atom

2.5 Electrons in Finite Quantum Well

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V(x)\Psi = E\Psi$$


Conditions:

for $x \leq 0, x \geq a$

$$V(x) = V_0;$$

for $0 < x < a$

$$V(x) = 0$$



2.5 Electrons in Finite Quantum Well

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V(x)\Psi = E\Psi$$


Conditions:

for $x \leq 0, x \geq a$

$$V(x) = V_0;$$

for $0 < x < a$

$$V(x) = 0$$



$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} = (E - V_0)\Psi$$

$$\Psi(r) = Ae^{-ik_1x} + Be^{ik_1x}$$

$$k_1 = \sqrt{\frac{2m(E - V_0)}{\hbar^2}}$$

$$-\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} = E\Psi$$

$$\Psi(r) = Ce^{-ik_2x} + De^{ik_2x}$$

$$k_2 = \sqrt{\frac{2mE}{\hbar^2}}$$

2.5 Electrons in Finite Quantum Well

Boundary Conditions:

$$\Psi(x)|_{x=a,0} \text{ continuous}$$

$$\Psi'(x)|_{x=a,0} \text{ continuous}$$

$$\int_{-\infty}^{\infty} \Psi(x)\Psi^*(x)dx = 1$$

for $x < 0, x > a$


$$\Psi(x) = Ae^{-ik_1x} + Be^{ik_1x}$$

$$k_1 = \sqrt{\frac{2m(E - V_0)}{\hbar^2}}$$

for $0 \leq x \leq a$

$$\Psi(x) = Ce^{-ik_2x} + De^{ik_2x}$$

$$k_2 = \sqrt{\frac{2mE}{\hbar^2}}$$



2.5 Electrons in Finite Quantum Well

$$\text{If } E < V_0 \quad k_1 = i \sqrt{\frac{2m(V_0 - E)}{\hbar^2}}$$

Boundary Conditions:

$$\Psi(x)|_{x=a,0} \text{ continuous}$$

$$\Psi'(x)|_{x=a,0} \text{ continuous}$$

$$\int_{-\infty}^{\infty} \Psi(x) \Psi^*(x) dx = 1$$

for $x < 0, x > a$


$$\Psi(x) = Ae^{-ik_1x} + Be^{ik_1x}$$

$$k_1 = \sqrt{\frac{2m(E - V_0)}{\hbar^2}}$$

for $0 \leq x \leq a$

$$\Psi(x) = Ce^{-ik_2x} + De^{ik_2x}$$

$$k_2 = \sqrt{\frac{2mE}{\hbar^2}}$$



2.5 Electrons in Finite Quantum Well

Boundary Conditions:

$$\Psi(x)|_{x=a,0} \text{ continuous}$$

$$\Psi'(x)|_{x=a,0} \text{ continuous}$$


$$\int_{-\infty}^{\infty} \Psi(x)\Psi^*(x)dx = 1$$

for $x < 0, x > a$

$$\Psi(r) = Ae^{-ik_1x} + Be^{ik_1x}$$

for $0 \leq x \leq a$

$$\Psi(r) = Ce^{-ik_2x} + De^{ik_2x}$$



Outline

2.1 2nd order differential equations and waves

2.2 Historic events in developing quantum mechanics

2.3 A case study


2.4 Electrons in infinite quantum well

2.5 Electrons in finite quantum well


2.6 Electrons in an atom

2.6 Electrons in an Atom

- 2D




Periodic boundary conditions




2.6 Electrons in an Atom





- 2D



Periodic boundary conditions




$f(r, \phi)$



2.6 Electrons in an Atom

- 3D



$$\Psi_{r,\theta,\phi} = R_n^l(r) Y_l^m(\phi, \theta)$$


2.6 Electrons in an Atom

- 3D

$3s^2 3p^6 3d^6$

$2s^2 2p^6$

$1s^2$



2.6 Electrons in an Atom

- 3D

Table 2.1 | Initial portion of the periodic table (shape: s,p)

Element	Notation	$n(\text{orbital})$	l	m	s
Hydrogen	$1s^1$	1	0	0	$+\frac{1}{2}$ or $-\frac{1}{2}$
Helium	$1s^2$	1	0	0	$+\frac{1}{2}$ and $-\frac{1}{2}$
Lithium	$1s^2 2s^1$	2	0	0	$+\frac{1}{2}$ or $-\frac{1}{2}$
Beryllium	$1s^2 2s^2$	2	0	0	$+\frac{1}{2}$ and $-\frac{1}{2}$
Boron	$1s^2 2s^2 2p^1$	2	1	<div style="display: flex; align-items: center; justify-content: center;"> <div style="margin-right: 10px;"> $m=0, -1, 1$ </div> <div style="border: 1px solid black; padding: 5px; text-align: center;"> <div style="display: flex; justify-content: space-around;"> <div style="border: 1px solid black; padding: 2px;">$\uparrow\downarrow$</div> <div style="border: 1px solid black; padding: 2px;">$\uparrow\downarrow$</div> <div style="border: 1px solid black; padding: 2px;">\uparrow</div> <div style="border: 1px solid black; padding: 2px; width: 20px;"></div> <div style="border: 1px solid black; padding: 2px; width: 20px;"></div> </div> </div> </div>	$m = 0, -1, +1$ $s = +\frac{1}{2}, -\frac{1}{2}$
Carbon	$1s^2 2s^2 2p^2$	2	1		
Nitrogen	$1s^2 2s^2 2p^3$	2	1		
Oxygen	$1s^2 2s^2 2p^4$	2	1		
Fluorine	$1s^2 2s^2 2p^5$	2	1		
Neon	$1s^2 2s^2 2p^6$	2	1		