PtoP Supplementary MATERIAL

1 User Study Interface



Figure 1: User Study Interface

2 Weighted Fleiss' Kappa Calculation

In our rating scheme, we acknowledge that larger rating discrepancies should correspond to greater disagreement. To account for this, we introduce the Weighted Fleiss' Kappa calculation. We first transfer our Likert scale rating category from text value to numerical value:

- *"Very unrealistic"* is assigned the value 1.
- *"Slightly unrealistic"* is assigned the value 2.
- *"Neutral"* is assigned the value 3.
- *"Slightly realistic"* is assigned the value 4.
- *"Very realistic"* is assigned the value 5.

Definition of the Weight Matrix

Since ratings are ordinal, we apply a quadratic weight function to penalize large discrepancies more than small ones. The weight matrix w_{ij} is defined as:

$$w_{ij} = 1 - \frac{(i-j)^2}{(k-1)^2} \tag{1}$$

where:

- i, j are rating categories $(1, 2, \ldots, k)$.
- k is the total number of rating categories.
- The further apart the categories, the smaller the weight.

In our case, the $k = 5$, an example of the weight matrix is:

$$W = \begin{bmatrix} 1.00 & 0.75 & 0.44 & 0.19 & 0.00 \\ 0.75 & 1.00 & 0.75 & 0.44 & 0.19 \\ 0.44 & 0.75 & 1.00 & 0.75 & 0.44 \\ 0.19 & 0.44 & 0.75 & 1.00 & 0.75 \\ 0.00 & 0.19 & 0.44 & 0.75 & 1.00 \end{bmatrix} \tag{2}$$

Observed Agreement

For each subject (e.g., a video being rated), let f_{ij} be the number of raters assigning a rating of j. The total number of ratings for a given subject i is:

$$p_i = \sum_{j=1}^{k} f_{ij} \tag{3}$$

The observed agreement for subject i is calculated as:

$$P_i = \frac{\sum_{j=1}^{k} \sum_{l=1}^{k} w_{jl} \cdot f_{ij} \cdot f_{il}}{p_i(p_i - 1)} \tag{4}$$

where:

- f_{ij} is the count of raters giving category j to subject i.
- f_{il} represents the number of raters who assigned rating l to subject i.
- w_{jl} is the weight between rating category j and l.
- $p_i(p_i - 1)$ is the total number of rating pairs.
- If $p_i = 1$, then P_i is set to 0 (as no agreement is possible).

The overall observed agreement across all subjects is:

$$\bar{P} = \frac{1}{N} \sum_{i=1}^{N} P_i \tag{5}$$

where N is the total number of subjects.

Expected Agreement

The proportion of ratings assigned to each category across all subjects is:

$$p_j = \frac{\sum_{i=1}^{N} f_{ij}}{\sum_{i=1}^{N} p_i} \tag{6}$$

The expected agreement, assuming ratings were randomly assigned, is:

$$P_e = \sum_{j=1}^{k} \sum_{l=1}^{k} w_{jl} \cdot p_j \cdot p_l \tag{7}$$

Computation of Weighted Fleiss' Kappa

The final formula for the weighted Fleiss' Kappa is:

$$\kappa_w = \frac{\bar{P} - P_e}{1 - P_e} \tag{8}$$

where:

- $\bar{P}$ is the observed agreement.
- P_e is the expected agreement.
- If $\bar{P} = P_e$, then $\kappa_w = 0$, indicating no agreement beyond chance.

Interpretation of Weighted Fleiss' Kappa

The interpretation of κ_w is as follows:

- $0.00 - 0.20$: Slight agreement
- $0.21 - 0.40$: Fair agreement
- $0.41 - 0.60$: Moderate agreement
- $0.61 - 0.80$: Substantial agreement
- $0.81 - 1.00$: Almost perfect agreement

For example, if we compute:

$$\kappa_w \approx 0.714 \tag{9}$$

this suggests **substantial agreement** among raters.

3 Gradient to Motion