

Calculus

Yaohui Wu

May 22, 2024

For all lovers of mathematics.

Introduction

Calculus is the study of continuous change established by **Issac Newton** (1643–1727) and **Gottfried Wilhelm Leibniz** (1646–1716) in the 17th century. **Single variable calculus** studies **derivatives** and **integrals** of functions of one variable and their relationship stated by the **fundamental theorem of calculus**.

$$\int_a^b f(x) dx = F(b) - F(a)$$

Contents

1	Functions and Limits	3
1.1	The Limit of a Function	3
1.1.1	Functions	3
1.1.2	Intuitive Definition of a Limit	3
1.2	The Precise Definition of a Limit	4
1.2.1	Epsilon-Delta Definition of a Limit	4
1.3	Computing Limits	5
1.4	Continuity	5
1.5	Limits and Infinity	5
2	Derivatives	5
2.1	Derivatives	5
2.2	Differentiation Formulas	5
2.3	Implicit Differentiation	5
2.4	Derivatives of Inverse Functions	5
2.5	Indeterminate Forms and l'Hospital's Rule	5

3	Applications of Differentiation	5
3.1	Maximum and Minimum Values	5
3.2	The Mean Value Theorem	5
3.3	Derivatives and Graphs	5
3.4	Antiderivatives	5
4	Integrals	5
4.1	Definite Integrals	5
4.2	Evaluating Definite Integrals	5
4.3	The Fundamental Theorem of Calculus	5
4.4	The Substitution Rule	5
5	Techniques of Integration	6
5.1	Integration by Parts	6
5.2	Trigonometric Integrals and Substitutions	6
5.3	Partial Fractions	6
5.4	Improper Integrals	6
6	Applications of Integration	6
6.1	Areas	6
6.2	Volumes	6
6.3	Arc Length	6
7	Sequences and Series	6
7.1	Sequences	6
7.2	Series	6
7.3	Convergence Tests	6
7.4	Power Series	6
7.5	Taylor Series	6
8	Parametric Equations and Polar Coordinates	6
8.1	Calculus of Parametric Equations	6
8.2	Calculus in Polar Coordinates	6
9	Differential Equations	6
9.1	Ordinary Differential Equations	6
9.2	Direction Fields and Euler's Method	6
9.3	Separable Equations	6

1 Functions and Limits

1.1 The Limit of a Function

1.1.1 Functions

A function $f : X \mapsto Y$ is a rule that assigns each element x in set X to exactly one element y in set Y . We have a formal definition of a function.

Definition 1.1. A **function** f is a binary relation R between domain X and codomain Y that satisfies:

- R is a subset of the Cartesian product of X and Y .

$$R \subset \{(x, y) \mid x \in X, y \in Y\}$$

- For every x in X , there exists a y in Y such that (x, y) is in R .

$$\forall x \in X, \exists y \in Y, (x, y) \in R$$

- If (x, y) and (x, z) are in R , then $y = z$.

$$(x, y) \in R \wedge (x, z) \in R \implies y = z$$

1.1.2 Intuitive Definition of a Limit

Newton and Leibniz introduced a working definition of a limit. Let $f(x)$ be a function defined on some open interval that contains the number a , except possibly at a itself.

Definition 1.2. The **limit** of $f(x)$ as x approaches a equals L if we can make $f(x)$ arbitrarily close to L by taking x sufficiently close to a from the left and the right but $x \neq a$.

$$\lim_{x \rightarrow a} f(x) = L$$

Definition 1.3. The **left-hand limit** of $f(x)$ as x approaches a from the left equals L if we can make $f(x)$ arbitrarily close to L by taking x sufficiently close to a where $x < a$.

$$\lim_{x \rightarrow a^-} f(x) = L$$

Definition 1.4. The **right-hand limit** of $f(x)$ as x approaches a from the right equals L if we can make $f(x)$ arbitrarily close to L by taking x sufficiently close to a where $x > a$.

$$\lim_{x \rightarrow a^+} f(x) = L$$

The limit **exists** if the left-hand limit and the right-hand limit of $f(x)$ as x approaches a equal L , otherwise the limit **does not exist**.

$$\lim_{x \rightarrow a} f(x) = L \iff \lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = L$$

1.2 The Precise Definition of a Limit

1.2.1 Epsilon-Delta Definition of a Limit

Augustin-Louis Cauchy (1789–1857) and **Karl Weierstrass** (1815–1897) developed a rigorous definition of a limit.

Definition 1.5.

$$\lim_{x \rightarrow a} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |f(x) - L| < \varepsilon$$

Definition 1.6.

$$\lim_{x \rightarrow a^-} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$a - \delta < x < a \implies |f(x) - L| < \varepsilon$$

Definition 1.7.

$$\lim_{x \rightarrow a^+} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$a < x < a + \delta \implies |f(x) - L| < \varepsilon$$

Problem 1.1. Prove that

$$\lim_{x \rightarrow 3} (4x - 5) = 7$$

Solution. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x - 3| < \delta \implies |(4x - 5) - 7| < \varepsilon$$

We simplify to get $|(4x - 5) - 7| = |4x - 12| = 4|x - 3|$ so we have

$$4|x - 3| < \varepsilon \iff |x - 3| < \frac{\varepsilon}{4}$$

Let $\delta = \varepsilon/4$, we have

$$0 < |x - 3| < \delta = \frac{\varepsilon}{4} \iff 0 < 4|x - 3| < \varepsilon \iff 0 < |(4x - 5) - 7| < \varepsilon$$

Therefore, by the definition of a limit,

$$\lim_{x \rightarrow 3} (4x - 5) = 7$$

■

1.3 Computing Limits

1.4 Continuity

1.5 Limits and Infinity

2 Derivatives

2.1 Derivatives

2.2 Differentiation Formulas

2.3 Implicit Differentiation

2.4 Derivatives of Inverse Functions

2.5 Indeterminate Forms and l'Hospital's Rule

3 Applications of Differentiation

3.1 Maximum and Minimum Values

3.2 The Mean Value Theorem

3.3 Derivatives and Graphs

3.4 Antiderivatives

4 Integrals

4.1 Definite Integrals

4.2 Evaluating Definite Integrals

4.3 The Fundamental Theorem of Calculus

4.4 The Substitution Rule

5 Techniques of Integration

5.1 Integration by Parts

5.2 Trigonometric Integrals and Substitutions

5.3 Partial Fractions

5.4 Improper Integrals

6 Applications of Integration

6.1 Areas

6.2 Volumes

6.3 Arc Length

7 Sequences and Series

7.1 Sequences

7.2 Series

7.3 Convergence Tests

7.4 Power Series

7.5 Taylor Series

8 Parametric Equations and Polar Coordinates

8.1 Calculus of Parametric Equations

8.2 Calculus in Polar Coordinates

9 Differential Equations

9.1 Ordinary Differential Equations

9.2 Direction Fields and Euler's Method

9.3 Separable Equations