

Calculus

Yaohui Wu

May 21, 2024

For all lovers of mathematics.

Introduction

Calculus is the study of continuous change established by **Issac Newton** (1643–1727) and **Gottfried Wilhelm Leibniz** (1646–1716) in the 17th century. **Single variable calculus** studies **derivatives** and **integrals** of functions of one variable and their relationship stated by the **fundamental theorem of calculus**.

$$\int_a^b f(x) dx = F(b) - F(a)$$

Contents

1	Functions and Limits	3
1.1	The Limit of a Function	3
1.2	The Precise Definition of a Limit	4
1.3	Computing Limits	4
1.4	Continuity	4
1.5	Limits and Infinity	4
2	Derivatives	4
2.1	Derivatives	4
2.2	Differentiation Formulas	4
2.3	Implicit Differentiation	4
2.4	Derivatives of Inverse Functions	4
2.5	Indeterminate Forms and l'Hospital's Rule	4
3	Applications of Differentiation	4
3.1	Maximum and Minimum Values	4
3.2	The Mean Value Theorem	4
3.3	Derivatives and Graphs	4
3.4	Antiderivatives	4

4	Integrals	4
4.1	Definite Integrals	4
4.2	Evaluating Definite Integrals	4
4.3	The Fundamental Theorem of Calculus	4
4.4	The Substitution Rule	4
5	Techniques of Integration	5
5.1	Integration by Parts	5
5.2	Trigonometric Integrals and Substitutions	5
5.3	Partial Fractions	5
5.4	Improper Integrals	5
6	Applications of Integration	5
6.1	Areas	5
6.2	Volumes	5
6.3	Arc Length	5
7	Sequences and Series	5
7.1	Sequences	5
7.2	Series	5
7.3	Convergence Tests	5
7.4	Power Series	5
7.5	Taylor Series	5
8	Parametric Equations and Polar Coordinates	5
8.1	Calculus of Parametric Equations	5
8.2	Calculus in Polar Coordinates	5
9	Differential Equations	5
9.1	Ordinary Differential Equations	5
9.2	Direction Fields and Euler's Method	5
9.3	Separable Equations	5

1 Functions and Limits

1.1 The Limit of a Function

A function $f : X \mapsto Y$ is a rule that assigns each element in set X to exactly one element $y = f(x)$ in set Y .

Definition 1.1. A **function** f is a binary relation R with domain X and codomain Y that satisfies:

- R is a subset of the Cartesian product of X and Y .

$$R \subset \{(x, y) \mid x \in X, y \in Y\}$$

- For every x in X , there exists a $y = f(x)$ such that (x, y) is in R .

$$\forall x \in X, \exists y = f(x) \in Y, (x, y) \in R$$

- If (x, y) and (x, z) are in R , then $y = z$.

$$(x, y) \in R \wedge (x, z) \in R \implies y = z$$

1.2 The Precise Definition of a Limit

1.3 Computing Limits

1.4 Continuity

1.5 Limits and Infinity

2 Derivatives

2.1 Derivatives

2.2 Differentiation Formulas

2.3 Implicit Differentiation

2.4 Derivatives of Inverse Functions

2.5 Indeterminate Forms and l'Hospital's Rule

3 Applications of Differentiation

3.1 Maximum and Minimum Values

3.2 The Mean Value Theorem

3.3 Derivatives and Graphs

3.4 Antiderivatives

4 Integrals

4.1 Definite Integrals

4.2 Evaluating Definite Integrals

4.3 The Fundamental Theorem of Calculus

4.4 The Substitution Rule

5 Techniques of Integration

5.1 Integration by Parts

5.2 Trigonometric Integrals and Substitutions

5.3 Partial Fractions

5.4 Improper Integrals

6 Applications of Integration

6.1 Areas

6.2 Volumes

6.3 Arc Length

7 Sequences and Series

7.1 Sequences

7.2 Series

7.3 Convergence Tests

7.4 Power Series

7.5 Taylor Series

8 Parametric Equations and Polar Coordinates

8.1 Calculus of Parametric Equations

8.2 Calculus in Polar Coordinates

9 Differential Equations

9.1 Ordinary Differential Equations

9.2 Direction Fields and Euler's Method

9.3 Separable Equations