

Calculus

Yaohui Wu

May 24, 2024

For all lovers of mathematics.

Introduction

Calculus is the study of continuous change established by **Issac Newton** (1643–1727) and **Gottfried Wilhelm Leibniz** (1646–1716) in the 17th century. **Single variable calculus** studies **derivatives** and **integrals** of functions of one variable and their relationship stated by the **fundamental theorem of calculus**.

$$\int_a^b f(x) dx = F(b) - F(a)$$

Contents

1	Functions and Limits	3
1.1	The Limit of a Function	3
1.1.1	Functions	3
1.1.2	Intuitive Definition of a Limit	3
1.2	The Precise Definition of a Limit	4
1.2.1	Epsilon-Delta Definition of a Limit	4
1.3	Computing Limits	5
1.3.1	Limit Laws	5
1.4	Continuity	12
1.5	Limits and Infinity	13
2	Derivatives	13
2.1	Derivatives	13
2.2	Differentiation Formulas	13
2.3	Implicit Differentiation	13
2.4	Derivatives of Inverse Functions	13
2.5	Indeterminate Forms and l'Hospital's Rule	13

3	Applications of Differentiation	13
3.1	Maximum and Minimum Values	13
3.2	The Mean Value Theorem	13
3.3	Derivatives and Graphs	13
3.4	Antiderivatives	13
4	Integrals	13
4.1	Definite Integrals	13
4.2	Evaluating Definite Integrals	13
4.3	The Fundamental Theorem of Calculus	13
4.4	The Substitution Rule	13
5	Techniques of Integration	14
5.1	Integration by Parts	14
5.2	Trigonometric Integrals and Substitutions	14
5.3	Partial Fractions	14
5.4	Improper Integrals	14
6	Applications of Integration	14
6.1	Areas	14
6.2	Volumes	14
6.3	Arc Length	14
7	Sequences and Series	14
7.1	Sequences	14
7.2	Series	14
7.3	Convergence Tests	14
7.4	Power Series	14
7.5	Taylor Series	14
8	Parametric Equations and Polar Coordinates	14
8.1	Calculus of Parametric Equations	14
8.2	Calculus in Polar Coordinates	14
9	Differential Equations	14
9.1	Ordinary Differential Equations	14
9.2	Direction Fields and Euler's Method	14
9.3	Separable Equations	14

1 Functions and Limits

1.1 The Limit of a Function

1.1.1 Functions

A function $f : X \mapsto Y$ is a rule that assigns each element x in set X to exactly one element y in set Y . We have a formal definition of a function.

Definition 1.1. A **function** f is a binary relation R between domain X and codomain Y that satisfies:

- R is a subset of the **Cartesian product** of X and Y .

$$R \subset \{(x, y) \mid x \in X, y \in Y\}$$

- For every x in X , there exists a y in Y such that (x, y) is in R .

$$\forall x \in X, \exists y \in Y, (x, y) \in R$$

- If (x, y) and (x, z) are in R , then $y = z$.

$$(x, y) \in R \wedge (x, z) \in R \implies y = z$$

The real line is the 1-dimensional **Euclidean space** defined as the set of real numbers \mathbb{R} . The xy -plane in the **Cartesian coordinate system** by **René Descartes** (1596-1650) is the 2-dimensional Euclidean space defined as the set of all ordered pairs of real numbers $(x, y) \in \mathbb{R}^2$. A **function of a real variable** is a function whose domain is the set of real numbers \mathbb{R} . A real function is a real-valued function of a real variable whose domain and codomain is \mathbb{R} .

1.1.2 Intuitive Definition of a Limit

Newton and Leibniz introduced a working definition of a limit. Let $f(x)$ be a function defined on some open interval that contains the number a , except possibly at a itself.

Definition 1.2. The **limit** of $f(x)$ as x approaches a equals L if we can make $f(x)$ arbitrarily close to L by taking x sufficiently close to a from the left and the right but $x \neq a$.

$$\lim_{x \rightarrow a} f(x) = L$$

Definition 1.3. The **left-hand limit** of $f(x)$ as x approaches a from the left equals L if we can make $f(x)$ arbitrarily close to L by taking x sufficiently close to a where $x < a$.

$$\lim_{x \rightarrow a^-} f(x) = L$$

Definition 1.4. The **right-hand limit** of $f(x)$ as x approaches a from the right equals L if we can make $f(x)$ arbitrarily close to L by taking x sufficiently close to a where $x > a$.

$$\lim_{x \rightarrow a^+} f(x) = L$$

The limit **exists** if the left-hand limit and the right-hand limit of $f(x)$ as x approaches a equal L , otherwise the limit **does not exist**.

$$\lim_{x \rightarrow a} f(x) = L \iff \lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = L$$

1.2 The Precise Definition of a Limit

1.2.1 Epsilon-Delta Definition of a Limit

Augustin-Louis Cauchy (1789–1857) and **Karl Weierstrass** (1815–1897) formalized a rigorous definition of a limit.

Definition 1.5.

$$\lim_{x \rightarrow a} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |f(x) - L| < \varepsilon$$

Definition 1.6.

$$\lim_{x \rightarrow a^-} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$a - \delta < x < a \implies |f(x) - L| < \varepsilon$$

Definition 1.7.

$$\lim_{x \rightarrow a^+} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$a < x < a + \delta \implies |f(x) - L| < \varepsilon$$

Problem 1.1. Prove that

$$\lim_{x \rightarrow 3} (4x - 5) = 7$$

Solution. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x - 3| < \delta \implies |(4x - 5) - 7| < \varepsilon$$

We simplify to get $|(4x - 5) - 7| = |4x - 12| = 4|x - 3|$ so we have

$$4|x - 3| < \varepsilon \iff |x - 3| < \frac{\varepsilon}{4}$$

Let $\delta = \varepsilon/4$, we have

$$0 < |x - 3| < \frac{\varepsilon}{4} \implies 4|x - 3| < \varepsilon \implies |(4x - 5) - 7| < \varepsilon$$

Therefore, by the definition of a limit, it is proved that

$$\lim_{x \rightarrow 3} (4x - 5) = 7$$

■

Problem 1.2. Prove that

$$\lim_{x \rightarrow 3} x^2 = 9$$

Solution. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x - 3| < \delta \implies |x^2 - 9| < \varepsilon$$

We simplify to get

$$|x^2 - 9| = |x + 3| |x - 3| < \varepsilon$$

Let C be a positive constant such that

$$|x + 3| |x - 3| < C |x - 3| < \varepsilon \iff |x - 3| < \frac{\varepsilon}{C}$$

Since we are interested only in values of x that are close to 3, it is reasonable to assume that $|x - 3| < 1$ such that $|x + 3| < 7$ so $C = 7$. Let $\delta = \min\{1, \varepsilon/7\}$, we have

$$0 < |x - 3| < 1 \iff |x + 3| < 7$$

$$0 < |x - 3| < \frac{\varepsilon}{7} \iff 7|x - 3| < \varepsilon$$

$$|x + 3| |x - 3| < 7|x - 3| < \varepsilon \implies |x^2 - 9| < \varepsilon$$

Therefore, it is proved that

$$\lim_{x \rightarrow 3} x^2 = 9$$

■

Problem 1.3. Prove that

$$\lim_{x \rightarrow 0^+} \sqrt{x} = 0$$

Solution. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < x < \delta \implies |\sqrt{x} - 0| < \varepsilon$$

We simplify to get $\sqrt{x} < \varepsilon \iff x < \varepsilon^2$. Let $\delta = \varepsilon^2$, we have

$$0 < x < \varepsilon^2 \implies |\sqrt{x} - 0| < \varepsilon$$

Therefore, it is proved that

$$\lim_{x \rightarrow 0^+} \sqrt{x} = 0$$

■

1.3 Computing Limits

1.3.1 Limit Laws

Suppose that c is a constant and the limits

$$\lim_{x \rightarrow a} f(x) = L$$

$$\lim_{x \rightarrow a} g(x) = M$$

exist. We have the following properties of limits called the **limit laws** to compute limits.

Theorem 1.1.

$$\lim_{x \rightarrow a} c = c$$

Proof. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |c - c| < \varepsilon$$

We have $|c - c| = 0 < \varepsilon$ so the trivial inequality is always true for any number $\delta > 0$. Therefore, by the definition of the limit, it is proved that

$$\lim_{x \rightarrow a} c = c \quad \blacksquare$$

Theorem 1.2.

$$\lim_{x \rightarrow a} x = a$$

Proof. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |x - a| < \varepsilon$$

Let $\delta = \varepsilon$, we have

$$0 < |x - a| < \delta = \varepsilon \implies |x - a| < \varepsilon$$

Therefore, by the definition of the limit, it is proved that

$$\lim_{x \rightarrow a} x = a \quad \blacksquare$$

Theorem 1.3 Constant Multiple Law. The limit of a constant times a function is the constant times the limit of the function.

$$\lim_{x \rightarrow a} [c f(x)] = c \lim_{x \rightarrow a} f(x) = cL$$

Proof. Note that if $c = 0$, then $cf(x) = 0$ and we have

$$\lim_{x \rightarrow a} [0 \cdot f(x)] = \lim_{x \rightarrow a} 0 = 0 = 0 \cdot \lim_{x \rightarrow a} f(x)$$

Let $\varepsilon > 0$ and $c \neq 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |cf(x) - c \lim_{x \rightarrow a} f(x)| < \varepsilon$$

We simplify to get

$$|f(x) - L| < \frac{\varepsilon}{|c|}$$

By the definition of the limit, there is a number $\delta_1 > 0$ such that

$$0 < |x - a| < \delta_1 \implies |f(x) - L| < \frac{\varepsilon}{|c|}$$

Let $\delta = \delta_1$, we have

$$0 < |x - a| < \delta \implies |cf(x) - c \lim_{x \rightarrow a} f(x)| < \varepsilon$$

Therefore, by the definition of the limit, it is proved that

$$\lim_{x \rightarrow a} [c f(x)] = c \lim_{x \rightarrow a} f(x) = cL \quad \blacksquare$$

Theorem 1.4 Sum and Difference Law. The limit of a sum or difference is the sum or difference of the limits.

$$\lim_{x \rightarrow a} [f(x) \pm g(x)] = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x) = L \pm M$$

Proof. First we prove the sum law. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |f(x) + g(x) - (L + M)| < \varepsilon$$

By the **triangle inequality** $|a + b| \leq |a| + |b|$, we have

$$|f(x) + g(x) - (L + M)| = |f(x) - L + g(x) - M| \leq |f(x) - L| + |g(x) - M|$$

Since $\lim_{x \rightarrow a} f(x) = L$, there is a number δ_1 such that

$$0 < |x - a| < \delta_1 \implies |f(x) - L| < \frac{\varepsilon}{2}$$

Similarly, there exists $\delta_2 > 0$, there is a number δ_2 such that

$$0 < |x - a| < \delta_2 \implies |g(x) - M| < \frac{\varepsilon}{2}$$

$$0 < |x - a| < \delta_2 \implies |g(x) - M| < \frac{\varepsilon}{2}$$

Let $\delta = \min\{\delta_1, \delta_2\}$ such that

$$0 < |x - a| < \delta \implies 0 < |x - a| < \delta_1, \delta_2$$

$$|f(x) - L| + |g(x) - M| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

Then we have

$$0 < |x - a| < \delta \implies |f(x) + g(x) - (L + M)| < \varepsilon$$

Therefore, by the definition of the limit, it is proved that

$$\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) = L + M$$

We prove the difference law using the sum law and the constant multiple law with $c = -1$.

$$\begin{aligned} \lim_{x \rightarrow a} [f(x) - g(x)] &= \lim_{x \rightarrow a} [f(x) + (-1)g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} (-1)g(x) \\ &= \lim_{x \rightarrow a} f(x) + (-1) \lim_{x \rightarrow a} g(x) \\ &= \lim_{x \rightarrow a} f(x) - \lim_{x \rightarrow a} g(x) = L - M \end{aligned}$$

Therefore, it is proved that

$$\lim_{x \rightarrow a} [f(x) \pm g(x)] = \lim_{x \rightarrow a} f(x) \pm \lim_{x \rightarrow a} g(x) = L \pm M \quad \blacksquare$$

Theorem 1.5 Product Law. The limit of a product is the product of the limits.

$$\lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x) = L \cdot M$$

Proof. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |f(x)g(x) - LM| < \varepsilon$$

By the triangle inequality, we have

$$\begin{aligned} |f(x)g(x) - LM| &= |f(x)g(x) - Lg(x) + Lg(x) - LM| = |[f(x) - L]g(x) + L[g(x) - M]| \\ &\leq |[f(x) - L]g(x)| + |L[g(x) - M]| = |f(x) - L| |g(x)| + |L| |g(x) - M| \end{aligned}$$

We want to make both of the terms less than $\varepsilon/2$. Since $\lim_{x \rightarrow a} f(x) = L$, there is a number $\delta_1 > 0$ such that

$$0 < |x - a| < \delta_1 \implies |f(x) - L| < \frac{\varepsilon}{2(1 + |M|)}$$

Since $\lim_{x \rightarrow a} g(x) = M$, there is a number $\delta_2 > 0$ such that

$$0 < |x - a| < \delta_2 \implies |g(x) - M| < \frac{\varepsilon}{2(1 + |L|)}$$

Also, there is a number $\delta_3 > 0$ such that

$$0 < |x - a| < \delta_3 \implies |g(x) - M| < 1$$

and therefore

$$|g(x)| = |g(x) - M + M| \leq |g(x) - M| + |M| < 1 + |M|$$

Let $\delta = \min\{\delta_1, \delta_2, \delta_3\}$ such that

$$0 < |x - a| < \delta \implies 0 < |x - a| < \delta_1, \delta_2, \delta_3$$

so we can combine the inequalities to get

$$\begin{aligned} |f(x)g(x) - LM| &\leq |f(x) - L| |g(x)| + |L| |g(x) - M| \\ &< \frac{\varepsilon}{2(1 + |M|)} (1 + |M|) + (1 + |L|) \frac{\varepsilon}{2(1 + |L|)} \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon \end{aligned}$$

Therefore, it is proved that

$$\lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x) = L \cdot M \quad \blacksquare$$

Theorem 1.6 Quotient Law. The limit of a quotient is the quotient of the limits (if that the limit of the denominator is not 0).

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)} = \frac{L}{M} \iff \lim_{x \rightarrow a} g(x) = M \neq 0$$

Proof. First we prove that

$$\lim_{x \rightarrow a} \frac{1}{g(x)} = \frac{1}{M}$$

Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x - a| < \delta \implies \left| \frac{1}{g(x)} - \frac{1}{M} \right| < \varepsilon$$

Notice that

$$\left| \frac{1}{g(x)} - \frac{1}{M} \right| = \frac{|M - g(x)|}{|Mg(x)|} = \frac{|g(x) - M|}{|Mg(x)|}$$

Since $\lim_{x \rightarrow a} g(x) = M$, there is a number δ_1 such that

$$0 < |x - a| < \delta_1 \implies |g(x) - M| < \frac{|M|}{2}$$

and therefore

$$|M| = |M - g(x) + g(x)| \leq |M - g(x)| + |g(x)| = |g(x) - M| + |g(x)| < \frac{|M|}{2} + |g(x)|$$

It is shown that

$$0 < |x - a| < \delta_1 \implies \frac{|M|}{2} < |g(x)| \implies \frac{1}{|g(x)|} < \frac{2}{|M|}$$

It follows that for these values of x ,

$$\frac{1}{|Mg(x)|} = \frac{1}{|M| |g(x)|} < \frac{1}{|M|} \frac{2}{|M|} = \frac{2}{M^2}$$

Also, there is a number $\delta_2 > 0$ such that

$$0 < |x - a| < \delta_2 \implies |g(x) - M| < \frac{M^2}{2} \varepsilon$$

Let $\delta = \min\{\delta_1, \delta_2\}$, if $0 < |x - a| < \delta$, then

$$\left| \frac{1}{g(x)} - \frac{1}{M} \right| = \frac{|M - g(x)|}{|Mg(x)|} = \frac{1}{|Mg(x)|} |g(x) - M| < \frac{2}{M^2} \frac{M^2}{2} \varepsilon = \varepsilon$$

which is what we want to show. We apply the product law to prove the quotient law

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} f(x) \left(\frac{1}{g(x)} \right) = \lim_{x \rightarrow a} f(x) \lim_{x \rightarrow a} \frac{1}{g(x)} = L \cdot \frac{1}{M} = \frac{L}{M} \quad \blacksquare$$

Theorem 1.7 Power Law.

$$\lim_{x \rightarrow a} [f(x)]^n = [\lim_{x \rightarrow a} f(x)]^n, n \in \mathbb{R}$$

Theorem 1.8 Root Law.

$$\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{\lim_{x \rightarrow a} f(x)}, n \in \mathbb{R}$$

Theorem 1.9 Direct Substitution Property. If f is a polynomial function, rational function, or trigonometric function and a is in the domain of f , then

$$\lim_{x \rightarrow a} f(x) = f(a)$$

Thus, we have the following limits

$$\lim_{\theta \rightarrow 0} \sin \theta = 0$$

$$\lim_{\theta \rightarrow 0} \cos \theta = 1$$

If $f(x) = g(x)$ when $x \neq a$, then $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x)$ if the limits exist.

Problem 1.4. Show that

$$\lim_{x \rightarrow 0} |x| = 0$$

Solution. Since $|x| = x$ for $x > 0$, we have

$$\lim_{x \rightarrow 0^+} |x| = \lim_{x \rightarrow 0^+} x = 0$$

For $x < 0$ we have $|x| = -x$ so

$$\lim_{x \rightarrow 0^-} |x| = \lim_{x \rightarrow 0^-} (-x) = 0$$

Therefore, it is shown that

$$\lim_{x \rightarrow 0} |x| = 0$$

■

Theorem 1.10. If $f(x) \leq g(x)$ for all x in an open interval that contains a , except possibly at a , and

$$\lim_{x \rightarrow a} f(x) = L$$

$$\lim_{x \rightarrow a} g(x) = M$$

then

$$\lim_{x \rightarrow a} f(x) \leq \lim_{x \rightarrow a} g(x) \iff L \leq M$$

Proof. We use the method of proof by contradiction. Suppose that $L > M$, then we have

$$\lim_{x \rightarrow a} [g(x) - f(x)] = M - L$$

Therefore, for any number $\varepsilon > 0$, there exists $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |g(x) - f(x) - (M - L)| < \varepsilon$$

Note that $L - M > 0$ by the hypothesis. Let $\varepsilon = L - M$, there exists $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |g(x) - f(x) - (M - L)| < L - M$$

Since $a \leq |a|$ for any number a , we have

$$0 < |x - a| < \delta \implies g(x) - f(x) - (M - L) < L - M$$

which simplifies to

$$0 < |x - a| < \delta \implies g(x) < f(x)$$

but this is a contradiction since given $f(x) \leq g(x)$. Then the inequality $L > M$ must be false so $L \leq M$ must be true. Therefore, it is proved that

$$\lim_{x \rightarrow a} f(x) \leq \lim_{x \rightarrow a} g(x) \quad \blacksquare$$

Theorem 1.11 Squeeze Theorem. If $f(x) \leq g(x) \leq h(x)$ for all x in an open interval that contains a , except possibly at a , and

$$\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L$$

then

$$\lim_{x \rightarrow a} g(x) = L$$

Proof. Let $\varepsilon > 0$ be given. Since $\lim_{x \rightarrow a} f(x) = L$, there exists $\delta_1 > 0$ such that

$$0 < |x - a| < \delta_1 \implies |f(x) - L| < \varepsilon \implies L - \varepsilon < f(x) < L + \varepsilon$$

Since $\lim_{x \rightarrow a} h(x) = L$, there exists $\delta_2 > 0$ such that

$$0 < |x - a| < \delta_2 \implies |h(x) - L| < \varepsilon \implies L - \varepsilon < h(x) < L + \varepsilon$$

Let $\delta = \min\{\delta_1, \delta_2\}$. If $0 < |x - a| < \delta$, then we have

$$L - \varepsilon < f(x) \leq g(x) \leq h(x) < L + \varepsilon \implies L - \varepsilon < g(x) < L + \varepsilon \implies |g(x) - L| < \varepsilon$$

which is what we want to prove. Therefore, it is proved that

$$\lim_{x \rightarrow a} g(x) = L \quad \blacksquare$$

By **algebra**, **geometry**, and **trigonometry**, we can get the following result by the **Pythagorean theorem** $a^2 + b^2 = c^2$. If $0 < \theta < \pi/2$, then

$$\sin \theta < \theta \implies \frac{\sin \theta}{\theta} < 1 \qquad \theta < \tan \theta = \frac{\sin \theta}{\cos \theta} \implies \cos \theta < \frac{\sin \theta}{\theta}$$

so we have the following inequality

$$\cos \theta < \frac{\sin \theta}{\theta} < 1$$

Since $(\sin \theta)/\theta$ is an even function, its left and right limits must be equal. Therefore, we have the following limit by the squeeze theorem.

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$$

Problem 1.5. Evaluate

$$\lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{\theta}$$

Solution. By the **Pythagorean identity** $\sin^2 \theta + \cos^2 \theta = 1$, we have

$$\frac{\cos \theta - 1}{\theta} = \frac{\cos \theta - 1}{\theta} \left(\frac{\cos \theta + 1}{\cos \theta + 1} \right) = \frac{\cos^2 \theta - 1}{\theta(\cos \theta + 1)} = \frac{-\sin^2 \theta}{\theta(\cos \theta + 1)} = \frac{\sin \theta}{\theta} \left(\frac{-\sin \theta}{\cos \theta + 1} \right)$$

We take the limit and we have

$$\lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{\theta} = \lim_{\theta \rightarrow 0} \left[\frac{\sin \theta}{\theta} \left(\frac{-\sin \theta}{\cos \theta + 1} \right) \right] = \left(\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} \right) \left(\lim_{\theta \rightarrow 0} \frac{-\sin \theta}{\cos \theta + 1} \right) = 0 \left(\frac{-1 \cdot 0}{1 + 1} \right) = 0$$

Therefore, it is shown that

$$\lim_{\theta \rightarrow 0} \frac{\cos \theta - 1}{\theta} = 0 \quad \blacksquare$$

1.4 Continuity

Let $f(x)$ be a function and the number a is in the domain of f so $f(a)$ is defined. If the limit exists, then we have the following definition.

Definition 1.8. A function f is **continuous** at the number a if

$$\lim_{x \rightarrow a} f(x) = f(a)$$

A function f is continuous from the left at a if the left-hand limit equals $f(a)$ and it is continuous from the right at a if the right-hand limit equals $f(a)$. A function f is continuous on an interval if it is continuous at every number in the interval. If f is not continuous at a , then it is a discontinuous function at a .

Theorem 1.12. If f and g are continuous functions at a and c is a constant, then the following functions are also continuous at a .

$$\bullet f + g \quad \bullet f - g \quad \bullet cf \quad \bullet f \cdot g \quad \bullet \frac{f}{g} \iff g(x) \neq 0$$

Theorem 1.13. Let $P(x)$ be any polynomial, then $P(x)$ is continuous on $\mathbb{R} = (-\infty, \infty)$.

Proof. A polynomial $P(x)$ is a function of the form

$$a_n x^n + a_{n-1} x^{n-1} + \cdots + a_2 x^2 + a_1 x + a_0$$

where the coefficients a_i are constants. $P(x)$ is the sum of power functions with a constant multiple and therefore it is continuous. \blacksquare

Theorem 1.14. Let f be any rational function, then f is continuous on its domain.

Proof. A rational function f is a function of the form

$$f(x) = \frac{P(x)}{Q(x)}$$

where P and Q are polynomials. We know that polynomials are continuous so a rational function is continuous on its domain. \blacksquare

Theorem 1.15. Polynomials, rational functions, root functions, and trigonometric functions are continuous on their domain.

1.5 Limits and Infinity

2 Derivatives

2.1 Derivatives

2.2 Differentiation Formulas

2.3 Implicit Differentiation

2.4 Derivatives of Inverse Functions

2.5 Indeterminate Forms and l'Hospital's Rule

3 Applications of Differentiation

3.1 Maximum and Minimum Values

3.2 The Mean Value Theorem

3.3 Derivatives and Graphs

3.4 Antiderivatives

4 Integrals

4.1 Definite Integrals

4.2 Evaluating Definite Integrals

4.3 The Fundamental Theorem of Calculus

4.4 The Substitution Rule

5 Techniques of Integration

5.1 Integration by Parts

5.2 Trigonometric Integrals and Substitutions

5.3 Partial Fractions

5.4 Improper Integrals

6 Applications of Integration

6.1 Areas

6.2 Volumes

6.3 Arc Length

7 Sequences and Series

7.1 Sequences

7.2 Series

7.3 Convergence Tests

7.4 Power Series

7.5 Taylor Series

8 Parametric Equations and Polar Coordinates

8.1 Calculus of Parametric Equations

8.2 Calculus in Polar Coordinates

9 Differential Equations

9.1 Ordinary Differential Equations

9.2 Direction Fields and Euler's Method

9.3 Separable Equations