Calculus

Yaohui Wu

May 22, 2024

For all lovers of mathematics.

Introduction

Calculus is the study of continuous change established by Issac Newton (1643–1727) and Gottfried Wilhelm Leibniz (1646–1716) in the 17th century. Single variable calculus studies derivatives and integrals of functions of one variable and their relationship stated by the fundamental theorem of calculus.

$$\int_{a}^{b} f(x) dx = F(b) - F(a)$$

Contents

1	Fun	ections and Limits				
	1.1	The Limit of a Function				
		1.1.1 Functions				
		1.1.2 Intuitive Definition of a Limit				
	1.2	The Precise Definition of a Limit				
		1.2.1 Epsilon-Delta Definition of a Limit				
	1.3	Computing Limits				
	1.4	Continuity				
	1.5	Limits and Infinity				
2	Derivatives					
	2.1	Derivatives				
	2.2	Differentiation Formulas				
	2.3	Implicit Differentiation				
	2.4	Derivatives of Inverse Functions				
	2.5	Indeterminate Forms and l'Hospital's Rule				

3	App	olications of Differentiation	5			
	3.1	Maximum and Minimum Values	5			
	3.2	The Mean Value Theorem	5			
	3.3	Derivatives and Graphs	5			
	3.4	Antiderivatives	5			
4	Integrals					
	4.1	Definite Integrals	5			
	4.2	Evaluating Definite Integrals	5			
	4.3	The Fundamental Theorem of Calculus	5			
	4.4	The Substitution Rule	5			
5	Techniques of Integration					
	5.1	Integration by Parts	6			
	5.2	Trigonometric Integrals and Substitutions	6			
	5.3	Partial Fractions	6			
	5.4	Improper Integrals	6			
6	Applications of Integration					
	6.1	Areas	6			
	6.2	Volumes	6			
	6.3	Arc Length	6			
7	Sequences and Series					
	7.1	Sequences	6			
	7.2	Series	6			
	7.3	Convergence Tests	6			
	7.4	Power Series	6			
	7.5	Taylor Series	6			
8	Parametric Equations and Polar Coordinates					
	8.1	Calculus of Parametric Equations	6			
	8.2	Calculus in Polar Coordinates	6			
9	Differential Equations					
	9.1	Ordinary Differential Equations	6			
	9.2	Direction Fields and Euler's Method	6			
	9.3	Separable Equations	6			

1 Functions and Limits

1.1 The Limit of a Function

1.1.1 Functions

A function $f: X \mapsto Y$ is a rule that assigns each element x in set X to exactly one element y in set Y. We have a formal definition of a function.

Definition 1.1. A function f is a binary relation R between domain X and codomain Y that satisfies:

• R is a subset of the Cartesian product of X and Y.

$$R \subset \{(x,y) \mid x \in X, y \in Y\}$$

• For every x in X, there exists a y in Y such that (x, y) is in R.

$$\forall x \in X, \exists y \in Y, (x, y) \in R$$

• If (x, y) and (x, z) are in R, then y = z.

$$(x,y) \in R \land (x,z) \in R \implies y = z$$

1.1.2 Intuitive Definition of a Limit

Newton and Leibniz introduced a working definition of a limit. Let f(x) be a function defined on some open interval that contains the number a, except possibly at a itself.

Definition 1.2. The **limit** of f(x) as x approaches a equals L if we can make f(x) arbitrarily close to L by taking x sufficiently close to a from the left and the right but $x \neq a$.

$$\lim_{x \to a} f(x) = L$$

Definition 1.3. The **left-hand limit** of f(x) as x approaches a from the left equals L if we can make f(x) arbitrarily close to L by taking x sufficiently close to a where x < a.

$$\lim_{x \to a^{-}} f(x) = L$$

Definition 1.4. The **right-hand limit** of f(x) as x approaches a from the right equals L if we can make f(x) arbitrarily close to L by taking x sufficiently close to a where x > a.

$$\lim_{x \to a^+} f(x) = L$$

The limit **exists** if the left-hand limit and the right-hand limit of f(x) as x approaches a equal L, otherwise the limit **does not exist**.

$$\lim_{x \to a} f(x) = L \iff \lim_{x \to a^{-}} f(x) = \lim_{x \to a^{+}} f(x) = L$$

1.2 The Precise Definition of a Limit

1.2.1 Epsilon-Delta Definition of a Limit

Augustin-Louis Cauchy (1789–1857) and **Karl Weierstrass** (1815–1897) developed a rigorous definition of a limit.

Definition 1.5.

$$\lim_{x \to a} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |f(x) - L| < \varepsilon$$

Definition 1.6.

$$\lim_{x \to a^{-}} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$a - \delta < x < a \implies |f(x) - L| < \varepsilon$$

Definition 1.7.

$$\lim_{x \to a^+} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$a < x < a + \delta \implies |f(x) - L| < \varepsilon$$

Problem 1.1. Prove that

$$\lim_{x \to 3} (4x - 5) = 7$$

Solution. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x-3| < \delta \implies |(4x-5)-7| < \varepsilon$$

- 1.3 Computing Limits
- 1.4 Continuity
- 1.5 Limits and Infinity
- 2 Derivatives
- 2.1 Derivatives
- 2.2 Differentiation Formulas
- 2.3 Implicit Differentiation
- 2.4 Derivatives of Inverse Functions
- 2.5 Indeterminate Forms and l'Hospital's Rule
- 3 Applications of Differentiation
- 3.1 Maximum and Minimum Values
- 3.2 The Mean Value Theorem
- 3.3 Derivatives and Graphs
- 3.4 Antiderivatives
- 4 Integrals
- 4.1 Definite Integrals
- 4.2 Evaluating Definite Integrals
- 4.3 The Fundamental Theorem of Calculus
- 4.4 The Substitution Rule

5 Techniques of Integration

- 5.1 Integration by Parts
- 5.2 Trigonometric Integrals and Substitutions
- 5.3 Partial Fractions
- 5.4 Improper Integrals
- 6 Applications of Integration
- 6.1 Areas
- 6.2 Volumes
- 6.3 Arc Length
- 7 Sequences and Series
- 7.1 Sequences
- 7.2 Series
- 7.3 Convergence Tests
- 7.4 Power Series
- 7.5 Taylor Series
- 8 Parametric Equations and Polar Coordinates
- 8.1 Calculus of Parametric Equations
- 8.2 Calculus in Polar Coordinates
- 9 Differential Equations
- 9.1 Ordinary Differential Equations
- 9.2 Direction Fields and Euler's Method
- 9.3 Separable Equations