

Calculus

Yaohui Wu

May 22, 2024

For all lovers of mathematics.

Introduction

Calculus is the study of continuous change established by **Issac Newton** (1643–1727) and **Gottfried Wilhelm Leibniz** (1646–1716) in the 17th century. **Single variable calculus** studies **derivatives** and **integrals** of functions of one variable and their relationship stated by the **fundamental theorem of calculus**.

$$\int_a^b f(x) dx = F(b) - F(a)$$

Contents

1	Functions and Limits	3
1.1	The Limit of a Function	3
1.1.1	Functions	3
1.1.2	Intuitive Definition of a Limit	3
1.2	The Precise Definition of a Limit	4
1.2.1	Epsilon-Delta Definition of a Limit	4
1.3	Computing Limits	6
1.3.1	Limit Laws	6
1.4	Continuity	7
1.5	Limits and Infinity	7
2	Derivatives	7
2.1	Derivatives	7
2.2	Differentiation Formulas	7
2.3	Implicit Differentiation	7
2.4	Derivatives of Inverse Functions	7
2.5	Indeterminate Forms and l'Hospital's Rule	7

3	Applications of Differentiation	7
3.1	Maximum and Minimum Values	7
3.2	The Mean Value Theorem	7
3.3	Derivatives and Graphs	7
3.4	Antiderivatives	7
4	Integrals	7
4.1	Definite Integrals	7
4.2	Evaluating Definite Integrals	7
4.3	The Fundamental Theorem of Calculus	7
4.4	The Substitution Rule	7
5	Techniques of Integration	8
5.1	Integration by Parts	8
5.2	Trigonometric Integrals and Substitutions	8
5.3	Partial Fractions	8
5.4	Improper Integrals	8
6	Applications of Integration	8
6.1	Areas	8
6.2	Volumes	8
6.3	Arc Length	8
7	Sequences and Series	8
7.1	Sequences	8
7.2	Series	8
7.3	Convergence Tests	8
7.4	Power Series	8
7.5	Taylor Series	8
8	Parametric Equations and Polar Coordinates	8
8.1	Calculus of Parametric Equations	8
8.2	Calculus in Polar Coordinates	8
9	Differential Equations	8
9.1	Ordinary Differential Equations	8
9.2	Direction Fields and Euler's Method	8
9.3	Separable Equations	8

1 Functions and Limits

1.1 The Limit of a Function

1.1.1 Functions

A function $f : X \mapsto Y$ is a rule that assigns each element x in set X to exactly one element y in set Y . We have a formal definition of a function.

Definition 1.1. A **function** f is a binary relation R between domain X and codomain Y that satisfies:

- R is a subset of the Cartesian product of X and Y .

$$R \subset \{(x, y) \mid x \in X, y \in Y\}$$

- For every x in X , there exists a y in Y such that (x, y) is in R .

$$\forall x \in X, \exists y \in Y, (x, y) \in R$$

- If (x, y) and (x, z) are in R , then $y = z$.

$$(x, y) \in R \wedge (x, z) \in R \implies y = z$$

1.1.2 Intuitive Definition of a Limit

Newton and Leibniz introduced a working definition of a limit. Let $f(x)$ be a function defined on some open interval that contains the number a , except possibly at a itself.

Definition 1.2. The **limit** of $f(x)$ as x approaches a equals L if we can make $f(x)$ arbitrarily close to L by taking x sufficiently close to a from the left and the right but $x \neq a$.

$$\lim_{x \rightarrow a} f(x) = L$$

Definition 1.3. The **left-hand limit** of $f(x)$ as x approaches a from the left equals L if we can make $f(x)$ arbitrarily close to L by taking x sufficiently close to a where $x < a$.

$$\lim_{x \rightarrow a^-} f(x) = L$$

Definition 1.4. The **right-hand limit** of $f(x)$ as x approaches a from the right equals L if we can make $f(x)$ arbitrarily close to L by taking x sufficiently close to a where $x > a$.

$$\lim_{x \rightarrow a^+} f(x) = L$$

The limit **exists** if the left-hand limit and the right-hand limit of $f(x)$ as x approaches a equal L , otherwise the limit **does not exist**.

$$\lim_{x \rightarrow a} f(x) = L \iff \lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = L$$

1.2 The Precise Definition of a Limit

1.2.1 Epsilon-Delta Definition of a Limit

Augustin-Louis Cauchy (1789–1857) and **Karl Weierstrass** (1815–1897) developed a rigorous definition of a limit.

Definition 1.5.

$$\lim_{x \rightarrow a} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$0 < |x - a| < \delta \implies |f(x) - L| < \varepsilon$$

Definition 1.6.

$$\lim_{x \rightarrow a^-} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$a - \delta < x < a \implies |f(x) - L| < \varepsilon$$

Definition 1.7.

$$\lim_{x \rightarrow a^+} f(x) = L$$

if for every number $\varepsilon > 0$, there is a number $\delta > 0$ such that

$$a < x < a + \delta \implies |f(x) - L| < \varepsilon$$

Problem 1.1. Prove that

$$\lim_{x \rightarrow 3} (4x - 5) = 7$$

Solution. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x - 3| < \delta \implies |(4x - 5) - 7| < \varepsilon$$

We simplify to get $|(4x - 5) - 7| = |4x - 12| = 4|x - 3|$ so we have

$$4|x - 3| < \varepsilon \iff |x - 3| < \frac{\varepsilon}{4}$$

Let $\delta = \varepsilon/4$, we have

$$0 < |x - 3| < \frac{\varepsilon}{4} \implies 4|x - 3| < \varepsilon \implies |(4x - 5) - 7| < \varepsilon$$

Therefore, by the definition of a limit,

$$\lim_{x \rightarrow 3} (4x - 5) = 7$$

■

Problem 1.2. Prove that

$$\lim_{x \rightarrow 3} x^2 = 9$$

Solution. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < |x - 3| < \delta \implies |x^2 - 9| < \varepsilon$$

We simplify to get

$$|x^2 - 9| = |x + 3| |x - 3| < \varepsilon$$

Let C be a positive constant such that

$$|x + 3| |x - 3| < C |x - 3| < \varepsilon \iff |x - 3| < \frac{\varepsilon}{C}$$

Since we are interested only in values of x that are close to 3, it is reasonable to assume that $|x - 3| < 1$ such that $|x + 3| < 7$ so $C = 7$. Let $\delta = \min\{1, \varepsilon/7\}$, we have

$$0 < |x - 3| < 1 \iff |x + 3| < 7$$

$$0 < |x - 3| < \frac{\varepsilon}{7} \iff 7|x - 3| < \varepsilon$$

$$|x + 3| |x - 3| < 7|x - 3| < \varepsilon \implies |x^2 - 9| < \varepsilon$$

Therefore, it is proved that

$$\lim_{x \rightarrow 3} x^2 = 9$$

■

Problem 1.3. Prove that

$$\lim_{x \rightarrow 0^+} \sqrt{x} = 0$$

Solution. Let $\varepsilon > 0$ be given, we want to find a number $\delta > 0$ such that

$$0 < x < \delta \implies |\sqrt{x} - 0| < \varepsilon$$

We simplify to get $\sqrt{x} < \varepsilon \iff x < \varepsilon^2$. Let $\delta = \varepsilon^2$, so we have

$$0 < x < \varepsilon^2 \implies |\sqrt{x} - 0| < \varepsilon$$

Therefore, it is proved that

$$\lim_{x \rightarrow 0^+} \sqrt{x} = 0$$

■

1.3 Computing Limits

1.3.1 Limit Laws

Suppose that c is a constant and the limits

$$\lim_{x \rightarrow a} f(x) = L \qquad \lim_{x \rightarrow a} g(x) = M$$

exists. We have the following properties of limits called the **limit laws** to compute limits.

Theorem 1.1.

$$\lim_{x \rightarrow a} c = c$$

Theorem 1.2.

$$\lim_{x \rightarrow a} x = a$$

Theorem 1.3 (Sum and Difference Law).

$$\lim_{x \rightarrow a} [f(x) \pm g(x)] = L \pm M$$

Theorem 1.4 (Constant Multiple Law).

$$\lim_{x \rightarrow a} [c f(x)] = c \lim_{x \rightarrow a} f(x)$$

Theorem 1.5 (Product Law).

$$\lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$$

Theorem 1.6 (Quotient Law).

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}$$

1.4 Continuity

1.5 Limits and Infinity

2 Derivatives

2.1 Derivatives

2.2 Differentiation Formulas

2.3 Implicit Differentiation

2.4 Derivatives of Inverse Functions

2.5 Indeterminate Forms and l'Hospital's Rule

3 Applications of Differentiation

3.1 Maximum and Minimum Values

3.2 The Mean Value Theorem

3.3 Derivatives and Graphs

3.4 Antiderivatives

4 Integrals

4.1 Definite Integrals

4.2 Evaluating Definite Integrals

4.3 The Fundamental Theorem of Calculus

4.4 The Substitution Rule

5 Techniques of Integration

5.1 Integration by Parts

5.2 Trigonometric Integrals and Substitutions

5.3 Partial Fractions

5.4 Improper Integrals

6 Applications of Integration

6.1 Areas

6.2 Volumes

6.3 Arc Length

7 Sequences and Series

7.1 Sequences

7.2 Series

7.3 Convergence Tests

7.4 Power Series

7.5 Taylor Series

8 Parametric Equations and Polar Coordinates

8.1 Calculus of Parametric Equations

8.2 Calculus in Polar Coordinates

9 Differential Equations

9.1 Ordinary Differential Equations

9.2 Direction Fields and Euler's Method

9.3 Separable Equations