

REAL ANALYSIS

Yaohui Wu

October 20, 2024

1 The Real Number System

1.1 The Set of Real Numbers

We define

1. The set of natural numbers is $\mathbb{N} = \{1, 2, 3, \dots\}$.
2. The set of integers is $\mathbb{Z} = \{0, 1, -1, 2, -2, \dots\}$.
3. The set of rational numbers is $\mathbb{Q} = \left\{\frac{m}{n} \mid m, n \in \mathbb{Z}, n \neq 0\right\}$.
4. The set of real numbers is \mathbb{R} .

Note that $\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}$.

Definition 1.1.1 (Well Ordering Property of \mathbb{N}). Every nonempty subset of \mathbb{N} has a least element.

Definition 1.1.2. An *ordered set* is a set S with a relation $<$ such that

1. (*Trichotomy*) If $x, y \in S$, then one and only one of the statements $x < y, x = y, y < x$ is true.
2. (*Transitivity*) If $x, y, z \in S$, $x < y$, and $y < z$, then $x < z$.

Definition 1.1.3. Suppose S is an ordered set and $E \subset S$. If there exists a $b \in S$ such that $x \leq b$ for all $x \in E$, then E is *bounded above* and b is an *upper bound* of E . If there exists an upper bound a such that $a \leq b$ for all upper bounds b of E , then a is the *least upper bound* or the *supremum* of E denoted by $\sup E$.

Similar definition applies for a set that is *bounded below* and *lower bound*. The *greatest lower bound* or the *infimum* of E is denoted by $\inf E$. A set is *bounded* if it is bounded above and below.

Definition 1.1.4 (Dedekind Completeness). An ordered set S has the *least-upper-bound property* if every nonempty subset $E \subset S$ that is bounded above has a least upper bound, that is, $\sup E$ exists in S .

Theorem 1.1.1. *There exists a unique ordered field \mathbb{R} with the least-upper-bound property such that $\mathbb{Q} \subset \mathbb{R}$.*

1.2 Archimedean Property

Theorem 1.2.1. *If $x, \varepsilon \in \mathbb{R}$ and $x \leq \varepsilon$ for all $\varepsilon > 0$, then $x \leq 0$.*

Theorem 1.2.2. *Suppose $x, y \in \mathbb{R}$.*

1. *(Archimedean property) If $x > 0$, then there exists an $n \in \mathbb{N}$ such that $nx > y$.*
2. *(\mathbb{Q} is dense in \mathbb{R}) If $x < y$, then there exists an $r \in \mathbb{Q}$ such that $x < r < y$.*

If $A \subset \mathbb{R}$ and $\sup A \in A$, then the supremum is the *maximum* of A denoted by $\max A$. Similarly, the infimum is the *minimum* of A denoted by $\min A$.

Theorem 1.2.3 (Triangle Inequality). *If $x, y \in \mathbb{R}$, then $|x + y| \leq |x| + |y|$.*

2 Sequences and Series

2.1 Sequences

Definition 2.1.1. A *sequence* $\{x_n\}$ is a function with domain \mathbb{N} and range \mathbb{R} .

Definition 2.1.2. A sequence $\{x_n\}$ is *bounded* if there exists a $B \in \mathbb{R}$ such that $|x_n| \leq B$ for all $n \in \mathbb{N}$.

Definition 2.1.3. A sequence is *convergent* and has the unique *limit* L and we write

$$\lim_{n \rightarrow \infty} \{x_n\} = L$$

if for every $\varepsilon > 0$, there exists an $N \in \mathbb{N}$ such that $|x_n - L| < \varepsilon$ for all $n \geq N$.

If a sequence is not convergent, then it is *divergent*.

Theorem 2.1.1. *If a sequence is convergent, then it is bounded.*

Theorem 2.1.2 (Squeeze Theorem). *If $a_n \leq b_n \leq c_n$ for all $n \geq N$, and $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = L$, then*

$$\lim_{n \rightarrow \infty} b_n = L$$

2.2 Monotonic Sequences

Definition 2.2.1. A sequence $\{x_n\}$ is *increasing* if $x_n \leq x_{n+1}$ or *decreasing* if $x_n \geq x_{n+1}$ for all $n \in \mathbb{N}$. A sequence is *monotonic* if it is increasing or decreasing.

Theorem 2.2.1 (Monotone Convergence Theorem). *A monotonic sequence is convergent if and only if it is bounded.*

If $\{x_n\}$ is increasing and bounded above, then

$$\lim_{n \rightarrow \infty} x_n = \sup x_n$$

If $\{x_n\}$ is decreasing and bounded below, then

$$\lim_{n \rightarrow \infty} x_n = \inf x_n$$

2.3 Subsequences

Definition 2.3.1. Let n_k be a sequence with $n_k < n_{k+1}$ for all $k \in \mathbb{N}$. The sequence $\{x_{n_k}\}$ is a *subsequence* of $\{x_n\}$.

Theorem 2.3.1. *If $\{x_n\}$ is convergent, then every subsequence $\{x_{n_k}\}$ is convergent, and*

$$\lim_{n \rightarrow \infty} x_n = \lim_{i \rightarrow \infty} x_{n_k}$$

2.4 Bolzano-Weierstrass Theorem

Theorem 2.4.1 (Bolzano-Weierstrass Theorem). *Every bounded sequence has a convergent subsequence.*

2.5 Cauchy Sequences

Definition 2.5.1. A sequence $\{x_n\}$ is a *Cauchy sequence* if for every $\varepsilon > 0$ there exists an $N \in \mathbb{N}$ such that $|x_m - x_n| < \varepsilon$ for all $m, n \geq N$.

Theorem 2.5.1. *If a sequence is a Cauchy sequence, then it is bounded.*

Theorem 2.5.2 (Cauchy Completeness). *A sequence is convergent if and only if it is a Cauchy sequence.*