arXiv:submit/8117302 [nlin.SI] 23 Sep 2026

SWIRL POLARIZATION AND A LOCAL QUADRATIC-JET DETECTOR FOR THE AXISYMMETRIC EULER EQUATIONS WITH SWIRL

YAOMING SHI

Abstract. We study a fixed smooth, rapidly decaying axisymmetric Euler flow with nontrivial swirl through two source-specific observables. The first is the swirl polarizer

$$\Phi(r,z,t) := \beta(r,z,t) = \tfrac{1}{r}U^{\phi}(r,z,t),$$

which is nonzero for the chosen customer but vanishes identically in the no-swirl class. Requiring active terminal mechanisms to descend from this polarizer, together with an idempotent ancestry projector, prevents a swirl-free Euler estimate from being reclassified as swirl-generated payment. The second observable is the local quadratic apex detector

$$\Psi(t) := (\beta_{zz}(0,0,t))^2.$$

Unlike the Beale–Kato–Majda quantity $\|\omega(t)\|_{\infty}$, which is a spatial supremum, Ψ is evaluated at a single point. The exact axisymmetric curl identity gives

$$\partial_z^2\omega^z(0,0,t) = 2\beta_{zz}(0,0,t), \qquad \|\nabla^2\omega(t)\|_{L^{\infty}(r\,dr\,dz)} \geq 2\sqrt{\Psi(t)}.$$

The signed square root of Ψ obeys an exact scalar countdown law driven by the apex compression. Starting from these observables, we develop a tagged swirl source, a whole-space Hodge cashier, signed source-genealogy banks, restart and response identities, a source-positive critical cashier, ratio-free/source-angle transfers, and terminal Riccati/cashier mechanisms. The resulting Grand Finale is reduced to one explicit modular source-specific closure package (O6.2). No unconditional finite-time singularity theorem is claimed.

Contents

1. Mathematical introduction and context 4
2. Introduction: swirl polarization, locality, and the accounting firewall 6
3. The single accounting table 8
4. Protected notation and the exact Euler customer 9
5. Customer, tagged source, and cashier 11
6. Detector clock, reserve, and conditional deadline 13
7. Seed selection and exact opening credit 15
8. The provenance firewall 16
9. The exact restart basis and deformation bills 16
10. Positive and negative genealogy banks 19
11. Cashier persistence and the no-reset side of the ledger 20
12. The Hodge-matched cashier and relaunch firewalls 22
13. Negative genealogy has only two ancestries 26
14. Continuation and carrier-capacity lemmas 28

Date: arXiv-v168-read of September 23, 2026.

2020 *Mathematics Subject Classification.* Primary 35Q31, 76B03, 35B44; Secondary 35B07, 35B60, 76B47.

Key words and phrases. three-dimensional incompressible Euler equations, axisymmetric Euler with swirl, finite-time singularity, swirl polarizer, local quadratic-jet detector, Beale–Kato–Majda criterion, Hodge reconstruction, source genealogy, vorticity growth, transported circulation.

15. Exact source response and the affine benchmark 30
16. Source-gradient restart system and the rotor 33
17. Level-set polarization and deformation stability 36
18. Fresh conversion, two-time response, and causal recycling 39
19. Auxiliary exact formulas retained for future calculations 43
20. Critical falsification propositions retained in the skeleton 46
21. Closed O-ledger with O6.2 as the sole open proposition 48
22. Rigorous-proof architecture and navigation 52
23. Grand-finale theorem/proposition chain and dependency table 53
23.1. Schematic funded chain 53
23.2. Detailed theorem/proposition chain 53
24. Grand-finale proof 58
25. Short proofs of the closed grand-finale statements 59
25.1. Short-proof location index 59
25.2. A. Customer, source, cashier, clock, and opening-credit short proofs 60
25.3. B. Provenance and restart/deformation short proofs 61
25.4. C. Genealogy-bank and positive-cashier short proofs 61
25.5. D. Adverse ancestry and continuation-capacity short proofs 63
25.6. E. Response and affine-benchmark short proofs 64
25.7. F. Rotor and mesoscopic-localization short proofs 65
25.8. G. Level-set polarization and deformation-stability short proofs 65
25.9. H. Causal adverse-response short proofs 66
25.10. I. Supporting exact-identity short proofs 67
25.11. J. Short proofs of the four falsification firewalls 67
25.12. K. Permanent reconstructed modules M1–M8: short proofs 68
25.13. K. O-ledger short-status cards 69
26. Compact ledger glossary 70
27. Conclusion 70
28. Acknowledgement 71
References 71
References 71
Rigorous proof audit: navigation convention 75
Rigorous proof audit: navigation convention 75
Appendix A. Exact customer, source, cashier, clock, and opening credit 75
A1. Exact squared-coordinate Euler system 75
A2. Pressure dictionary 77
A3. Pressure-free source-to-vorticity account 77
A4. Whole-space Hodge cashier 78
A5. Seven-dimensional Hodge lift 79
A6. Detector evolution and clock 80
A7. Reserve–clock identity 81
A8. Conditional finite-time collapse 81
A9. Opening family 82
A10. Opening reserve 83
Appendix B. Provenance and restart/deformation accounting 83
B1. Routing–GREEN normal form 84
B2. Two-shadow firewall 84

B3. Logarithmic tangent system 85
B4. Tangent exit pays derivative action 86
B5. Exact radial star-hull identity 87
B6. First-order bridge among restart channels 88
B7. Signed log-box thickening 89
Appendix C. Genealogy banks and positive-cashier relaunch 90
C1. Exact two-bank equations 90
C2. Finite-action floor equals cancellation lock 91
C3. Exact cancellation ledger 92
C4. Fresh positive genealogy near the apex 93
C5. Positive-genealogy material cashier 93
C6. Credit-reset firewall 94
C7. Intrinsic positive-genealogy Riccati closure 95
C8. Rate failure pays radial non-affinity 96
C9. Visibility collapse is positive-genealogy tail concentration 97
C10. Finite intrinsic exit graph 98
C11. Hodge-matched Cauchy bridge 99
C12. Exact dual-moment rate 99
C13. Hodge-matched Riccati gate 100
C14. Automatic fresh relaunch 101
C15. Uniform apex tube on compact classical intervals 102
C16. Terminal-time uniformity firewall 103
C17. No-Zeno relaunch on a regular positive-compression interval 104
Appendix D. Adverse ancestry and ordinary continuation capacity 105
D1. Exact slope ancestry 105
D2. Old source is geometrically separated 106
D3. Raw source-gradient Gronwall estimate 109
D4. Weighted Hodge energy is not a new mechanism 110
D5. Physical Biot–Savart control 111
D6. Finite physical Lipschitz action is a continuation firewall 112
Appendix E. Two-time response and the fixed-seed affine benchmark 115
E1. Vertical response representation 115
E2. True-Euler aged-response discount 117
E3. Fixed-seed signed line-source positivity 119
E4. Global affine cancellation barrier 121
E5. Loss of affine polarization has a tagged conversion cost 123
E6. Protected apex tube in two-time response 126
E7. One-shot exterior conversion alternative 127
Appendix F. Source-gradient rotor and compact mesoscopic conversion 129
F1. Exact source-gradient system 129
F2. Rotor normal form 130
F3. H-skew cancellation 131
F4. Passive-circulation interpretation 131
F5. Exact apex mixed-shear Hodge moment 132
F6. Hodge-aware adverse exchange 134
F7. Angular-tail Hodge suppression of the rotor 134
F8. Compact mesoscopic rotation ticket 135

Appendix G. Circulation-level-set polarization and deformation stability 138
G1. Circulation level-set response 139
G2. Raw level-set compensation criterion 140
G3. Uniform fixed-seed source-gradient condition number 141
G4. Stability of the complete Hodge-response polarization 143
G5. First polarization loss pays a definite deformation bill 146
Appendix H. Causal adverse-response ledger 147
H1. Final/deposition Hodge visibility comparison 148
H2. Finite raw response capacity 149
H3. Fresh-apex anti-rotation 150
H4. Exact generated-negative Hodge response 151
H5. Fresh primary conversion is quadratically small 152
H6. Causal Hodge resolvent for negative self-recycling 153
H7. Three-account negative-bank bound 156
H8. Resummed strict-gap criterion 156
Appendix I. Supporting exact identities 157
I1. Hodge visibility law 157
I2. Exact guarded source-payment identity 158
I3. Raw cross-shear factorization 159
I4. Carrier-Hodge comparison 160
I5. Fresh genealogy as a true Euler source-time truncation 160
I6. No double charging 162
I7. Strict-gap algebra 163
Appendix J. Falsification firewalls 164
J1. Finite local action alone cannot enforce the cancellation gap 164
J2. Frozen Hodge geometry alone has no uniform gap 166
J3. Infinite fractional refresh does not imply infinite physical conversion 167
J4. Pointwise reproduction number is too strong 168
Appendix K. Permanent O-ledger: O1–O6.1 closed, O6.2 sole open slot 169
O1. Large-collapse-age physical-bank retirement 169
O2. Bounded-collapse-age physical-bank return 170
O3. Physical-bank consolidation or paid centered deformation 172
O4. Grand cancellation gap 173
O5. Terminal noncontinuation from O4 and O6.2 174
O6.2. Post-O4 integral cashier closure; M1–M8 module archive 180
Appendix L. Grand-finale assembly 195
L1. Funded chain up to the open bill 195
L2. Grand finale 197

1. Mathematical introduction and context

The finite-time singularity problem for the three-dimensional incompressible Euler equations remains one of the central open problems in mathematical fluid dynamics [25, 34]. The Beale–Kato–Majda continuation criterion identifies

$$\int_0^T \|\omega(t)\|_{L^\infty}\, dt$$

as the classical obstruction to continuation [2]. In the axisymmetric setting this quantity is still a global spatial observable in the meridian variables. The present approach does not replace the BKM criterion. Instead, it asks whether a source-specific *local* detector can be coupled to the exact swirl-generated genealogy strongly enough to force the same continuation failure.

The distinction between axisymmetric Euler flow without swirl and with swirl is fundamental. Under the classical hypotheses of Ukhovskii–Yudovich, the no-swirl system is globally regular [96], whereas nontrivial swirl retains the vortex-stretching mechanism; see, for example, [78, 25]. This motivates the polarizer

$$\Phi = \beta = \tfrac{1}{r}U^{\phi}.$$

Setting $\Phi \equiv 0$ identifies the regular no-swirl shadow problem. Accordingly, a mechanism counted as specifically swirl-generated must disappear in that shadow. The idempotent provenance projector ε_β is introduced to enforce this separation algebraically.

The second organizing quantity is the local detector

$$\Psi(t) = (\beta_{zz}(0,0,t))^2.$$

It samples a single quadratic swirl jet at the apex rather than the spatial supremum of vorticity. Its signed square root satisfies an exact first-order law driven by the apex compression a, producing the countdown clock $\mathfrak{t}_\Psi$. The central technical problem is therefore to connect the nonzero swirl polarizer to this local clock through the tagged source, Hodge reconstruction, signed genealogy, material response, and terminal cashier, without importing an untagged no-swirl estimate.

For standard background on the vorticity formulation, Biot–Savart/Hodge reconstruction, particle trajectories, Kelvin circulation, and vortex stretching, we refer to Majda–Bertozzi [78]. Broader perspectives on the Euler blow-up problem are given in [25, 34]. Related singularity constructions for axisymmetric Euler in different domains or regularity classes include [38, 16, 17]; these results are not used as proof input here. Riccati-type amplification mechanisms in reduced vorticity dynamics have classical precedents in [20, 24].

Broader related literature. The literature surrounding Euler singularity formation is extensive. For further work on continuation and breakdown criteria, axisymmetric singularity scenarios, geometric and Lagrangian mechanisms, reduced Euler dynamics, numerical evidence, and modern instability or self-similar blow-up constructions, see [1, 2, 11, 12, 16, 17, 19, 21, 24, 25], [26, 28, 29, 34, 36, 37, 38, 40, 41, 43], [45, 46, 50, 54, 55, 56, 57, 59, 63, 65], [66, 69, 75, 76, 77, 78, 82, 85, 91, 92], [94, 95, 96, 97, 98, 102, 89, 90].

Closely related Boussinesq and active-scalar problems provide both comparison mechanisms and testing grounds for transport, stretching, front formation, and singularity scenarios; see [6, 7, 8, 10, 13, 15, 18, 22, 23, 27], [32, 35, 39, 49, 68, 67, 79, 93, 99].

One-dimensional and geometric fluid models, including the Constantin–Lax–Majda, De Gregorio, Hunter–Saxton, and related integrable or nonlocal models, form another important strand of the surrounding literature; see [3, 5, 20, 30, 31, 60, 61, 62, 70], [71, 81, 84, 83, 87, 88, 101, 100].

For viscous comparison problems, axisymmetric Navier–Stokes regularity, cylindrical formulations, energy methods, and partial-regularity background, see [4, 9, 14, 33, 42, 44, 51, 52], [53, 58, 72, 73, 74, 103].

Finally, the analysis of stable, self-similar, or profile-driven blow-up in other nonlinear PDE provides useful methodological context for singularity stability and modulation arguments; see [Collot–Merle–Raphaël(2017), 47, 48, 64, 80, 86].

Two earlier preprints by the author provide historical motivation [90, 89]. Their reduced blow-up calculations are not used here as exact full-Euler proof input. The present manuscript works throughout with a single smooth, rapidly decaying, full axisymmetric Euler flow with swirl and derives every active step from the exact three-dimensional equations.

The permanent proof architecture consists of the exact tagged source/Hodge cashier (against the standard vorticity/Hodge background of [78]), positive and negative genealogy banks, restart and no-double-charge geometry, the source-positive critical $\varpi = 2/5$ cashier, the critical-to-Hodge source-angle/GREEN bridge, the ratio-free global GREEN barrier, the transported circulation-square potential and aspect geometry, and the terminal powered/growth/cashier mechanisms. The remaining conditional step is isolated as one modular source-specific closure package.

Organization of the paper. Section 2 introduces the swirl-polarization and locality viewpoint and explains the accounting firewall. Section 3 gives the compact ledger map. Section 4 fixes the protected namespace and exact Euler customer, after which Sections 5 and 6 introduce the tagged source, Hodge cashier, local detector clock, and reserve. The subsequent sections develop the provenance, restart, genealogy, response, rotor, and polarization mechanisms. Section 21 states the permanent O-ledger, and Section 23 records the complete dependency chain. The conclusion summarizes the scope of the conditional result. Beginning with Section A, the second half of the manuscript provides a referee-level proof archive in the same notation; each closed ledger entry is restated and proved there with direct cross-references to its publication-facing statement.

2. Introduction: swirl polarization, locality, and the accounting firewall

The analysis is built around one fixed smooth Euler flow with nontrivial axisymmetric swirl. Two choices distinguish the framework.

Swirl polarization. The regular azimuthal quotient

$$\boxed{\Phi(r,z,t) := \beta(r,z,t) = \tfrac{1}{r}U^{\phi}(r,z,t)}$$

is used as a physical polarizer. For the chosen customer, $\Phi \not\equiv 0$, whereas the axisymmetric no-swirl class is characterized by $\Phi \equiv 0$. Every source-specific terminal account is therefore required to descend from nonzero swirl polarization. The ancestry projector ε_β enforces the same rule algebraically: switching off the swirl account must reduce active descendant equations to the no-swirl Euler background, rather than leave behind an artificial swirl mechanism. This is the role of the accounting firewall. A background estimate that survives at $\Phi \equiv 0$ cannot, by itself, be used as payment for a conclusion attributed to swirl.

A local quadratic detector. Define

$$\psi_\beta(t) := \beta_{zz}(0,0,t), \qquad \boxed{\Psi(t) := \psi_\beta(t)^2.}$$

For the fixed symmetry class, $\beta(0,0,t) = 0$, so the swirl value at the apex carries no nontrivial information. The second axial jet is the first nontrivial local swirl observable. The exact Euler equations give

$$\dot{\psi_\beta} = -2a\psi_\beta, \qquad \dot{\Psi} = -4a\Psi,$$

and therefore the positive countdown clock

$$\boxed{\mathfrak{t}_\Psi(t) = \frac{\psi_\beta(t)}{\psi_\beta(0)} = \sqrt{\frac{\Psi(t)}{\Psi(0)}} = e^{-2\int_0^t a(\tau)\,d\tau}.}$$

This is conceptually different from the Beale–Kato–Majda quantity $\|\omega(t)\|_\infty$ [2]: the latter is a global spatial supremum, whereas Ψ is a single-point jet evaluation.
Exact placement in the vorticity hierarchy. For an axisymmetric velocity with $U^\phi = r\Phi$,

$$\boxed{\omega^r = -\frac{1}{r}\partial_z(rU^\phi) = -r\Phi_z, \qquad \omega^z = \frac{1}{r}\partial_r(rU^\phi) = 2\Phi + r\Phi_r.} \tag{2.1}$$

Smooth axis regularity yields

$$\omega^z(0,z,t) = 2\Phi(0,z,t) = 2\beta(0,z,t),$$

and hence

$$\boxed{\partial_z^2\omega^z(0,0,t) = 2\beta_{zz}(0,0,t).} \tag{2.2}$$

Consequently,

$$\boxed{\|\nabla^2\omega(t)\|_{L^\infty(r\,dr\,dz)} \geq \|\partial_z^2\omega^z(t)\|_{L^\infty(r\,dr\,dz)} \geq 2\sqrt{\Psi(t)}.} \tag{2.3}$$

Although the axis itself has zero $r\,dr\,dz$ measure, smoothness makes the axis value the limit of values on positive-r sets of positive weighted measure; hence the weighted essential supremum controls the axis trace. The same statement is recorded formally in proposition 6.4.

The derivative order in (2.3) is essential. Neither the BKM norm nor the velocity Lipschitz norm universally controls $\sqrt{\Psi}$. Let $\Upsilon_{\mathrm{sc}}(\xi,\eta)$ be a smooth compactly supported axis-regular profile with $\partial_\eta^2\Upsilon_{\mathrm{sc}}(0,0) = 1$, and set

$$\beta_\delta(r,z) = \delta^2\Upsilon_{\mathrm{sc}}\Big(\frac{r}{\delta},\frac{z}{\delta}\Big).$$

Here (r,z) are the reserved physical meridian coordinates and (ξ,η) are dimensionless profile variables. Then $\sqrt{\Psi_\delta} = 1$, whereas the associated pure-swirl field satisfies

$$\|\omega_\delta\|_\infty = O(\delta^2), \qquad \|\nabla U_\delta\|_\infty = O(\delta^2).$$

Thus both norms can tend to zero while the local quadratic detector remains fixed. A direct comparison with $\|\nabla U\|_\infty$ therefore requires an additional physical length scale; dimensionally, the natural conditional form is

$$\ell_{\mathrm{loc}}(t)^2\sqrt{\Psi(t)} \lesssim \|\nabla U(t)\|_\infty$$

under an appropriate scale or nonconcentration hypothesis.

For the same reason, the biharmonic quantity $\Delta^2\omega$ cannot replace the Hessian in (2.3). If

$$\beta_L(r,z) = z^2\Upsilon_{\mathrm{cut}}\Big(\frac{r}{L},\frac{z}{L}\Big),$$

where $\Upsilon_{\mathrm{cut}}(\xi,\eta)$ is a smooth dimensionless axisymmetric cutoff equal to one near the origin, then $\beta_{L,zz}(0,0) = 2$ while

$$\|\Delta^2\omega_L\|_\infty = O(L^{-2}) \longrightarrow 0.$$


Hence no universal lower bound of $\|\Delta^2\omega\|_\infty$ by $c\sqrt{\Psi}$ can hold.

The rest of the paper tracks the chain

$$\begin{aligned}\text{nonzero swirl polarization} &\longrightarrow \text{tagged source} \longrightarrow \text{signed Hodge genealogy}\\ &\longrightarrow \text{apex compression} \longrightarrow \text{local countdown clock.}\end{aligned}$$

The accounting language is used only to enforce provenance and no-double-charge rules; it does not replace the underlying Euler analysis.

3. The single accounting table



Account	Exact mathematical object	Status
Customer	fixed $u_0 = yw_0$	fixed once
Source	$\Theta = \varepsilon_\beta u^2$, $q = \Theta_y$, C_β	GREEN
Cashier	$a = (3/8)\int K_H H$	exact
Positive bank	$a^+ = (3/8)\int K_H H^+$	banked
Negative bank	$a^- = (3/8)\int K_H H^-$	banked
Reserve family	$\mathcal{F}_\kappa = Q - 3(1+\kappa)a^2$	M7-P terminal route
Withdrawal	negative genealogy / Hodge aging	routed
Restart debt	$\mathfrak{A}, \mathfrak{J}, H, \mathfrak{T}$ star-hull/tangent action	paid routing
O6.1	inherited extreme-aspect / near-horizontal source-angle stock	GREEN
O4	$a^- \le (1-\delta_*)a^+$	GREEN
O6.2	$e^{-\mathcal{G}_{\mathrm{cash}}} \in L^1(d\Lambda)$	**SOLE OPEN**
M8	exact integral cashier countdown	GREEN
O5	terminal finite-time noncontinuation	GREEN via M8

Remark 3.1 (Skeleton convention). Historical derivations and obsolete branches are omitted from the reconstructed main body. Compact and referee-level proofs of the closed Grand-Finale statements are retained in the audited layers, while the immutable archives preserve the longer derivations and historical research provenance. Exactly one unresolved package is stated as an *Open Proposition*: O6.2. The inherited-launch source-angle audit formerly denoted O6.1 is recorded as closed and is not charged again downstream.

4. Protected notation and the exact Euler customer

Definition 4.1 (Physical symmetry class and protected namespace)**.** *We use the standard smooth signed-meridian extension in the physical variables (r, z). The velocity and pressure parities are*

$$U^r(-r,z,t) = -U^r(r,z,t), \qquad U^r(r,-z,t) = U^r(r,z,t), \tag{4.1}$$

$$U^\phi(-r,z,t) = -U^\phi(r,z,t), \qquad U^\phi(r,-z,t) = U^\phi(r,z,t), \tag{4.2}$$

$$U^z(-r,z,t) = U^z(r,z,t), \qquad U^z(r,-z,t) = -U^z(r,z,t), \tag{4.3}$$

$$\mathsf{p}(-r,z,t) = \mathsf{p}(r,z,t), \qquad \mathsf{p}(r,-z,t) = \mathsf{p}(r,z,t). \tag{4.4}$$

Equations (4.1)–(4.4) *define the symmetry class with respect to the physical variables (r, z). In particular, the regular quotients below extend smoothly to the symmetry axes.*

The squared coordinates are

$$\boxed{x = r^2, \qquad y = z^2.} \tag{4.5}$$

The physical regular quotients and their squared-coordinate representatives are

$$\boxed{\begin{aligned}\alpha(r,z,t) &= -\frac{1}{r}U^r(r,z,t) = v(x,y,t),\\ \beta(r,z,t) &= \frac{1}{r}U^\phi(r,z,t) = u(x,y,t),\\ \gamma_{\mathrm{ax}}(r,z,t) &= \frac{1}{z}U^z(r,z,t) = g(x,y,t).\end{aligned}} \tag{4.6}$$

The swirl polarizer is the azimuthal quotient itself:

$$\boxed{\Phi(r,z,t) := \beta(r,z,t) = \frac{U^\phi(r,z,t)}{r} = u(r^2,z^2,t).} \tag{4.7}$$

For the fixed customer, $\Phi \not\equiv 0$ on every nontrivial classical branch, whereas the axisymmetric no-swirl class is exactly $\Phi \equiv 0$. The polarizer is therefore used as the first source-specific filter: a terminal payment which survives after Φ is switched off belongs to the swirl-free background and is not admitted as swirl-generated revenue.

Consequently the regular quotient fields inherit even symmetry in the signed meridian variables:

$$\boxed{\begin{aligned}\alpha(-r,z,t) &= \alpha(r,z,t) = \alpha(r,-z,t),\\ \beta(-r,z,t) &= \beta(r,z,t) = \beta(r,-z,t),\\ \gamma_{\mathrm{ax}}(-r,z,t) &= \gamma_{\mathrm{ax}}(r,z,t) = \gamma_{\mathrm{ax}}(r,-z,t),\\ \mathsf{p}(-r,z,t) &= \mathsf{p}(r,z,t) = \mathsf{p}(r,-z,t).\end{aligned}} \tag{4.8}$$

Thus r, z are reserved exclusively for the physical meridian coordinates, while x, y are reserved exclusively for their squared-coordinate images $x = r^2$, $y = z^2$. They are never reused as unrelated dummy variables, exponents, profile coordinates, or free parameters. Dimensionless profile variables in scaling arguments use separate symbols such as (ξ, η). The remaining protected objects are

$$\boxed{\begin{aligned}\mathsf{p}(r,z,t) &= P(r^2,z^2,t),\\ Q(t) &= \mathsf{p}_{rr}(0,0,t) - \mathsf{p}_{zz}(0,0,t),\\ a(t) &= v(0,0,t), \qquad b(t) = u(0,0,t).\end{aligned}} \tag{4.9}$$

The detector/value variables are

$$\boxed{V = v - a, \qquad Y = \sqrt{\Psi/\Psi(0)}, \qquad p = u_y, \qquad \Gamma = xu, \qquad R = w + xw_x = \Gamma_x/y.} \tag{4.10}$$

The protected first-jet entries are

$$\boxed{C = xv_x, \qquad D = yg_y, \qquad E = yv_y, \qquad F = xg_x.} \tag{4.11}$$

The restart basis is

$$\boxed{\mathfrak{A} = 2v - g, \quad \mathfrak{J} = C + D, \quad \mathfrak{S} = x(g_x + v_y) = -\frac{xH}{2}, \quad \mathfrak{T} = x(g_x - v_y).} \tag{4.12}$$

Finally,

$$\boxed{\mathfrak{r} = x^{2/3}y^{1/3}, \qquad \Lambda(t;\tau) = \int_\tau^t a(\xi)\,d\xi.} \tag{4.13}$$

These protected symbols are not reused for unrelated dynamic quantities.

Remark 4.2 (Statement-local notation rule). Every theorem, proposition, lemma, corollary, conditional proposition, and open proposition below is written so that each temporary symbol is either defined in that statement or is explicitly tied to one of the standing Definitions. Throughout theorem-like statements and their audited long proofs, $s := y/x$ denotes aspect ratio only. In particular, plain r is never reused as an aspect ratio, and the protected physical quotient names α, β are never reused as dummy exponents or auxiliary parameters. Likewise the protected source/first-jet names R, C, D, E, F retain only their established meanings. The protected slope $p = u_y$ and centered value $V = v - a$ are likewise never reused as generic parameters, polarizations, or visibility variables. Start times are written t_0 and source/deposition times are written τ; these symbols are never used for aspect ratio inside a theorem-like statement. The notation D_t^+ denotes the upper right Dini derivative.

Proposition 4.3 (Exact squared-coordinate Euler system). *Let*

$$\mathcal{D}_{xy} := \partial_t - 2xv\partial_x + 2yg\partial_y, \qquad \varepsilon_\beta \in \{0, 1\}. \tag{4.14}$$

Then the tagged axisymmetric Euler system is

$$g + 2yg_y = 2v + 2xv_x, \tag{4.15}$$

$$\mathcal{D}_{xy}v - v^2 + \varepsilon_\beta u^2 = 2P_x, \tag{4.16}$$

$$\mathcal{D}_{xy}(\varepsilon_\beta u) = 2\varepsilon_\beta uv, \tag{4.17}$$

$$\mathcal{D}_{xy}g + g^2 = -2P_y. \tag{4.18}$$

The characteristics satisfy

$$\dot{x} = -2xv, \qquad \dot{y} = 2yg. \tag{4.19}$$

At $\varepsilon_\beta = 1$ this is physical Euler. At $\varepsilon_\beta = 0$, the tagged swirl equation (4.17) becomes the identity $0 = 0$, the centrifugal source in (4.16) disappears, and the displayed active PDE system reduces to the axisymmetric Euler equations without swirl. The untagged profile u may still be retained off-ledger as a passive bookkeeping carrier, but it no longer enters the active shadow PDEs.

Accounting bridge. The PDE proposition opens Peter's legal accounts. The next section identifies which part of the fixed customer is genuine invested capital, how that capital becomes a tagged source, and how the Hodge cashier converts the resulting genealogy into apex credit.

Proposition 4.4 (Pressure dictionary). *With the protected coordinate dictionary*

$$x = r^2, \qquad y = z^2, \qquad \mathsf{p}(r, z, t) = P(x, y, t),$$

the physical and squared pressure derivatives satisfy

$$\mathsf{p}_r = 2rP_x, \quad \mathsf{p}_z = 2zP_y, \tag{4.20}$$

$$\boxed{\mathsf{p}_{rr} = 2P_x + 4xP_{xx}, \qquad \mathsf{p}_{rz} = 4rzP_{xy}, \qquad \mathsf{p}_{zz} = 2P_y + 4yP_{yy}.} \tag{4.21}$$

Hence physical Hessian entries are never identified directly with P_{xx}, P_{xy}, P_{yy}.

5. Customer, tagged source, and cashier

Definition 5.1 (Final fixed customer). *The active datum is*

$$\boxed{u_0(x, y) = yw_0(x, y), \qquad v_0 = g_0 = 0,} \tag{5.1}$$

where

$$\boxed{w_0(x,y) = \left(1 + \frac{1}{5}y\right) e^{-c_* x - s_* y} + 10x^2 e^{-2x - b_* y},} \tag{5.2}$$

$$c_* := \sqrt{2/3}, \qquad s_* := 1/\sqrt{3}, \qquad b_* := 6 + 2\sqrt{7}. \tag{5.3}$$

No later forcing, packet activation, amplitude tuning, or change of datum is allowed.

Definition 5.2 (Tagged source)**.** *Set*

$$\boxed{\Theta := \varepsilon_\beta u^2, \qquad q := \Theta_y, \qquad C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y}.} \tag{5.4}$$

The circulation

$$\boxed{\Gamma := xu} \tag{5.5}$$

is exactly transported:

$$\boxed{\mathcal{D}_{xy}\Gamma = 0.} \tag{5.6}$$

Proposition 5.3 (Pressure-free source-to-vorticity account)**.** *With*

$$\Theta := \varepsilon_\beta u^2, \qquad q := \Theta_y$$

as in definition 5.2, define

$$\omega^\phi := \partial_z U^r - \partial_r U^z, \qquad D_t := \partial_t + U^r\partial_r + U^z\partial_z,$$

and

$$\bar{G} := \omega^\phi / r = zH(r^2, z^2, t), \qquad H := -2(v_y + g_x). \tag{5.7}$$

Then

$$\boxed{\mathcal{D}_{xy}\Theta = 4v\Theta, \qquad \mathcal{D}_{xy}H + gH = 2q.} \tag{5.8}$$

Equivalently, with $\beta(r,z,t) = U^\phi(r,z,t)/r$,

$$D_t\bar{G} = \varepsilon_\beta\, \partial_z(\beta^2). \tag{5.9}$$

Proposition 5.4 (Whole-space Hodge cashier)**.** *Let*

$$\mathcal{Q} := (0,\infty)^2, \qquad dX := dx\, dy.$$

Define the meridional stream function ψ_1 *by*

$$U^r = -r\partial_z\psi_1, \qquad U^z = 2\psi_1 + r\partial_r\psi_1.$$

Then the meridional velocity is reconstructed from

$$-\mathcal{L}_1\psi_1 = \bar{G}, \qquad \mathcal{L}_1 = \partial_r^2 + \frac{3}{r}\partial_r + \partial_z^2. \tag{5.10}$$

and the apex compression $a(t) = v(0,0,t)$ *is*

$$\boxed{a(t) = \frac{3}{4}\int_0^\infty \int_{\mathbb{R}} \frac{r^3 z}{(r^2+z^2)^{5/2}} \bar{G}(r,z,t)\, dz\, dr.} \tag{5.11}$$

In squared coordinates,

$$\boxed{a(t) = \frac{3}{8}\int_{\mathcal{Q}} K_H(x,y)H(x,y,t)\, dX, \qquad K_H(x,y) := \frac{x\sqrt{y}}{(x+y)^{5/2}} > 0.} \tag{5.12}$$

Proposition 5.5 (Seven-dimensional Hodge lift)**.** *Writing* $\psi_1 = z\chi(x, y, t)$ *gives*

$$v = \chi + 2y\chi_y, \qquad g = 2\chi + 2x\chi_x, \tag{5.13}$$

$$-\mathcal{L}_7\chi = H, \qquad \mathcal{L}_7 = 4x\partial_x^2 + 8\partial_x + 4y\partial_y^2 + 6\partial_y. \tag{5.14}$$

For $\mathbf{X} \in \mathbb{R}^4$, $\mathbf{Z} \in \mathbb{R}^3$, *define the radial lifts*

$$\widetilde{\chi}(\mathbf{X}, \mathbf{Z}, t) := \chi(|\mathbf{X}|^2, |\mathbf{Z}|^2, t), \qquad \widetilde{H}(\mathbf{X}, \mathbf{Z}, t) := H(|\mathbf{X}|^2, |\mathbf{Z}|^2, t).$$

Then

$$-\Delta_{\mathbb{R}^7}\widetilde{\chi} = \widetilde{H}, \tag{5.15}$$

so all meridional values and first jets are whole-space Hodge descendants of H*.*

6. Detector clock, reserve, and conditional deadline

Accounting bridge. The cashier has produced apex credit a; the detector now turns that credit into a clock and a reserve account. The conditional deadline theorem says exactly what Peter would receive if the reserve never becomes negative.

Definition 6.1 (Local quadratic-jet detector)**.** *Assume* $b(0) = u(0, 0, 0) = 0$*. Then* $b(t) \equiv 0$*. Define the signed apex jet*

$$\boxed{\psi_\beta(t) := \beta_{zz}(0, 0, t) = 2u_y(0, 0, t)}$$

and the local quadratic detector

$$\boxed{\Psi(t) := \psi_\beta(t)^2 = (\beta_{zz}(0, 0, t))^2, \qquad \mathcal{J}_\beta := \varepsilon_\beta\Psi.} \tag{6.1}$$

Remark 6.2 (Local quadratic detector versus the BKM spatial supremum)*.* For axisymmetric swirl $U^\phi = r\Phi$,

$$\omega^z = \frac{1}{r}\partial_r(rU^\phi) = 2\Phi + r\Phi_r.$$

Smoothness therefore gives

$$\|\omega(t)\|_{L^\infty(\mathbb{R}^3)} \geq 2|\Phi(0, 0, t)|.$$

For the fixed customer, $\Phi(0, 0, t) = \beta(0, 0, t) = 0$ throughout the classical lifespan, so in particular

$$\|\omega(t)\|_\infty \geq \beta(0, 0, t)^2 = 0.$$

That apex-value inequality is automatic and carries no useful information here. The first nontrivial local swirl observable at the apex is instead

$$\Psi(t) = (\Phi_{zz}(0, 0, t))^2.$$

The BKM continuation quantity $\|\omega(t)\|_\infty$ is a spatial supremum over the whole three-dimensional flow, equivalently over the whole meridian plane together with the axisymmetric components. By contrast, Ψ is a single-point local quadratic-jet evaluation.

Definition 6.3 (Weighted meridian supremum convention)**.** *For an axisymmetric scalar or tensor field* $\Upsilon_{\mathrm{fld}}(r, z)$*, set*

$$\|\Upsilon_{\mathrm{fld}}\|_{L^\infty(r\,dr\,dz)} := \operatorname*{ess\,sup}_{(0,\infty)\times\mathbb{R}}{}^{r\,dr\,dz} |\Upsilon_{\mathrm{fld}}(r, z)|.$$

Because the density r is strictly positive for $r > 0$, this is the same essential supremum as for ordinary Lebesgue measure on the open meridian. If Υ_{fld} extends continuously to $r = 0$, then every axis value is controlled by this norm:

$$|\Upsilon_{\mathrm{fld}}(0, z_\star)| \le \|\Upsilon_{\mathrm{fld}}\|_{L^\infty(r\,dr\,dz)}.$$

For the vorticity Hessian we use the Cartesian tensor magnitude

$$|\nabla^2\omega| := \left(\sum_{i,j,k=1}^{3} |\partial_i\partial_j\omega_k|^2\right)^{1/2}.$$

Proposition 6.4 (Exact local vorticity-to-detector bridge)**.** *For every classical time,*

$$\boxed{\partial_z^2\omega^z(0,0,t) = 2\beta_{zz}(0,0,t) = 2\psi_\beta(t).} \tag{6.2}$$

Consequently,

$$\boxed{\|\nabla^2\omega(t)\|_{L^\infty(r\,dr\,dz)} \ge 2\sqrt{\Psi(t)}.} \tag{6.3}$$

Proof. From $U^\phi = r\Phi$ and axisymmetry,

$$\omega^z = \frac{1}{r}\partial_r(rU^\phi) = 2\Phi + r\Phi_r.$$

Smooth axis regularity gives

$$\omega^z(0, z, t) = 2\Phi(0, z, t) = 2\beta(0, z, t).$$

Differentiating twice in z proves (6.2). Since $\Psi = \psi_\beta^2$,

$$|\partial_z^2\omega^z(0,0,t)| = 2\sqrt{\Psi(t)}.$$

The Cartesian Hessian norm dominates its $\partial_z^2\omega^z$ component, and definition 6.3 controls the smooth axis trace by the weighted essential supremum. This proves (6.3). □

Proposition 6.5 (Detector evolution and local clock)**.** *Let $\psi_\beta, \Psi, \mathcal{J}_\beta$ be as in definition 6.1. Throughout the classical lifespan,*

$$\boxed{\dot\psi_\beta = -2a\psi_\beta, \qquad \dot\Psi = -4a\Psi, \qquad \dot{\mathcal{J}}_\beta = -4a\mathcal{J}_\beta.} \tag{6.4}$$

If $\Psi(0) > 0$, define

$$\boxed{\mathfrak{t}_\Psi(t) := e^{-2\int_0^t a(\tau)d\tau} = \frac{\psi_\beta(t)}{\psi_\beta(0)} = \sqrt{\frac{\Psi(t)}{\Psi(0)}} = \sqrt{\frac{\mathcal{J}_\beta(t)}{\mathcal{J}_\beta(0)}}.} \tag{6.5}$$

Then

$$\boxed{a(t) = -\frac{1}{2}\frac{\dot{\mathfrak{t}}_\Psi(t)}{\mathfrak{t}_\Psi(t)}.} \tag{6.6}$$

Definition 6.6 (Reserve)**.** *Because $b \equiv 0$,*

$$\dot a = -a^2 + \tfrac{1}{3}Q. \tag{6.7}$$

Define

$$\boxed{\mathcal{F}_\Psi := Q - 9a^2.} \tag{6.8}$$

Proposition 6.7 (Reserve–clock identity)**.** *Let $\mathcal{F}_\Psi := Q - 9a^2$ be the reserve from definition 6.6, and let $\mathfrak{t}_\Psi$ be the detector clock from proposition 6.5. Then*

$$\boxed{\mathcal{F}_\Psi = 3(\dot{a} - 2a^2) = -\frac{3}{2}\frac{\ddot{\mathfrak{t}}_\Psi}{\mathfrak{t}_\Psi}.} \tag{6.9}$$

Hence

$$\boxed{\mathcal{F}_\Psi \geq 0 \iff \ddot{\mathfrak{t}}_\Psi \leq 0.} \tag{6.10}$$

Theorem 6.8 (Conditional finite-time collapse)**.** *Let $\Psi, \mathfrak{t}_\Psi, \mathcal{F}_\Psi$ be as in definitions 6.1 and 6.6 and proposition 6.5. Suppose $\Psi(0) > 0$ and at some classical time t_1 one has*

$$a(t_1) > 0, \qquad \mathcal{F}_\Psi(t) \geq 0 \tag{6.11}$$

for all later classical times under consideration. Define

$$T_* := \inf\{t \geq t_1 : \mathfrak{t}_\Psi(t) = 0\}, \qquad \inf \varnothing := +\infty.$$

Then either regularity fails beforehand, or

$$\boxed{T_* \leq t_1 + \frac{1}{2a(t_1)}.} \tag{6.12}$$

If the solution persists to T_*, then

$$\Psi(t) \to 0, \qquad a(t) \to +\infty \quad (t \uparrow T_*). \tag{6.13}$$

Remark 6.9 (Accounting interpretation). The clock theorem is already paid. The unpaid reserve bill is to keep the true Euler descendant inside a source-specific positive-compression corridor. The rest of this skeleton is a sequence of reductions of that unpaid bill.

7. Seed selection and exact opening credit

Accounting bridge. Before auditing future expenses, one must verify that Peter's chosen customer really opens with positive credit. The following seed calculation supplies that initial deposit without changing the customer later.

Proposition 7.1 (Opening family)**.** *Let $B \neq 0$ and $\lambda > 0$, and use the fixed constants $c_* = \sqrt{2/3}$ and $s_* = 1/\sqrt{3}$. For*

$$u_{\mathrm{ini}} = B(y + \lambda x)e^{-(c_* x + s_* y)}, \qquad v_{\mathrm{ini}} = g_{\mathrm{ini}} = 0, \tag{7.1}$$

define

$$\mathcal{I}(\lambda) := \int_{-\pi/2}^{\pi/2} \frac{\cos^5\phi \, \sin^2\phi \, (\sin^2\phi + \lambda\cos^2\phi)}{(c_* \cos^2\phi + s_* \sin^2\phi)^3} \, d\phi > 0.$$

Then

$$\boxed{\dot{a}(0) = \frac{3\varepsilon_\beta B^2}{8}(c_* - s_*\lambda)\mathcal{I}(\lambda).} \tag{7.2}$$

Therefore, in the physical sector $\varepsilon_\beta = 1$,

$$0 < \lambda < \sqrt{2} \Rightarrow \dot{a}(0) > 0, \quad \lambda = \sqrt{2} \Rightarrow \dot{a}(0) = 0, \quad \lambda > \sqrt{2} \Rightarrow \dot{a}(0) < 0. \tag{7.3}$$

The final $A = 10$ customer in (5.1) is a later fixed descendant of this source-selection program; no new customer is introduced thereafter.

Proposition 7.2 (Opening reserve)**.** *For an active positive-opening seed with $a(0) = b(0) = 0$,*

$$\boxed{Q(0) = 3\dot{a}(0) > 0, \qquad \mathcal{F}_\Psi(0) = 3\dot{a}(0) > 0.} \tag{7.4}$$

Thus the initial account is funded; the problem is persistence, not launch.

8. The provenance firewall

Accounting bridge. Positive opening credit is not enough: every later payment must still be traceable to the same investment. The provenance rules are therefore the ledger's anti-fraud controls.

Definition 8.1 (Idempotent ancestry)**.** *The Boolean genealogy tag obeys*

$$\boxed{\varepsilon_\beta^2 = \varepsilon_\beta.} \tag{8.1}$$

If a quantity has genuine single-seed ancestry, it is represented as $\varepsilon_\beta G$. Products of descendants keep one copy of the same tag:

$$(\varepsilon_\beta G_1)(\varepsilon_\beta G_2) = \varepsilon_\beta G_1 G_2. \tag{8.2}$$

Proposition 8.2 (Routing–GREEN normal form)**.** *Every finite polynomial expression made from ordinary routing quantities and certified GREEN descendants can be written*

$$\boxed{\mathcal{X} = \mathcal{U}_\mathcal{X} + \varepsilon_\beta \mathcal{G}_\mathcal{X}.} \tag{8.3}$$

A mixed sum remains mixed. In particular, multiplying an Euler-background quantity by ε_β at the end does not manufacture ancestry.

Definition 8.3 (Common-tag quotient after admission)**.** *If A/B is already mathematically defined in the physical sector and numerator and denominator have the same certified genealogy, the provenance ledger may write*

$$\frac{\varepsilon_\beta A}{\varepsilon_\beta B} \equiv_\beta \varepsilon_\beta \frac{A}{B}. \tag{8.4}$$

This is not scalar cancellation at the shadow $0/0$, and it is forbidden before the physical quotient has independently crossed its admission gate.

Proposition 8.4 (Two-shadow firewall)**.** *Every terminal candidate must pass both tests:*

- *(i) under the formal single-seed shadow $\varepsilon_\beta = 0$, no positive cashier or terminal contradiction may remain;*
- *(ii) under physical no-swirl axisymmetric Euler, a generic meridional estimate may route existing credit but may not by itself imply finite-time blow-up.*

Hence every final payment must retain an irreducible source-specific ancestor.

9. The exact restart basis and deformation bills

Accounting bridge. When the favorable account leaves its affine corridor, the loss is not written off. It is itemized into the following finite restart basis, which serves as Peter's list of legally chargeable deformation bills.

Definition 9.1 (Protected first-jet basis)**.** *At $x, y > 0$ define*

$$C = xv_x, \qquad D = yg_y, \qquad E = yv_y, \qquad F = xg_x, \qquad s = y/x, \tag{9.1}$$

and

$$\boxed{V = v - a, \qquad \mathfrak{A} = 2v - g, \qquad \mathfrak{J} = C + D,} \tag{9.2}$$

$$\boxed{\mathfrak{S} = F + E/s = x(g_x + v_y) = -xH/2, \qquad \mathfrak{T} = F - E/s = x(g_x - v_y).} \tag{9.3}$$

Incompressibility gives

$$C - D = -\frac{1}{2}\mathfrak{A}, \tag{9.4}$$

so

$$C = \frac{1}{2}\mathfrak{J} - \frac{1}{4}\mathfrak{A}, \qquad D = \frac{1}{2}\mathfrak{J} + \frac{1}{4}\mathfrak{A}, \tag{9.5}$$

$$F = \frac{1}{2}(\mathfrak{S} + \mathfrak{T}), \qquad E = \frac{s}{2}(\mathfrak{S} - \mathfrak{T}). \tag{9.6}$$

Thus the full first-jet restart information is encoded by $V, \mathfrak{A}, \mathfrak{J}, H, \mathfrak{T}$.

Proposition 9.2 (Logarithmic tangent system)**.** *Let $s := y/x$ be the aspect ratio, let t_0 be a starting time, and for an infinitesimal displacement $(\delta x, \delta y)$ put*

$$\mathfrak{p} = \delta x/x, \qquad \mathfrak{q} = \delta y/y. \tag{9.7}$$

Then

$$\frac{d}{dt}\begin{pmatrix}\mathfrak{p}\\ \mathfrak{q}\end{pmatrix} = \mathbb{M}\begin{pmatrix}\mathfrak{p}\\ \mathfrak{q}\end{pmatrix}, \tag{9.8}$$

with

$$\boxed{\mathbb{M} = \frac{1}{2}\mathfrak{A}I_2 + \begin{pmatrix}-\mathfrak{J} & -s(\mathfrak{S} - \mathfrak{T})\\ \mathfrak{S} + \mathfrak{T} & \mathfrak{J}\end{pmatrix},} \tag{9.9}$$

where I_2 is the 2×2 identity matrix. If $\mathcal{U}(t, t_0)$ is the fundamental logarithmic tangent matrix with $\mathcal{U}(t_0, t_0) = I_2$, then

$$\boxed{\det \mathcal{U}(t, t_0) = \exp\left(\int_{t_0}^{t} \mathfrak{A}(\tau)d\tau\right).} \tag{9.10}$$

Lemma 9.3 (Tangent exit pays derivative action)**.** *Let $\mathcal{U}(t, t_0)$ and $\mathbb{M}$ be as in proposition 9.2. If a logarithmic tangent column starts from a unit coordinate vector e_j at t_0 and, in the Euclidean norm, satisfies $\|\mathcal{U}(t, t_0)e_j - e_j\| \geq \eta > 0$, then*

$$\boxed{\int_{t_0}^{t} \|\mathbb{M}(\tau)\|d\tau \geq \log(1 + \eta).} \tag{9.11}$$

On a compact aspect-ratio window $0 < s_- \leq s = y/x \leq s_+ < \infty$, this forces a definite accumulated amount of at least one of

$$|\mathfrak{A}|, \qquad |\mathfrak{J}|, \qquad |\mathfrak{S}|, \qquad |\mathfrak{T}|. \tag{9.12}$$

Thus the tangent account is not a new mechanism; it is a cumulative form of the same restart bills.

Proposition 9.4 (Exact radial star-hull identity)**.** *For $X = (x, y) \in \mathcal{Q}$ put $s := y/x$ and $\mathcal{R}_v := xv_x + yv_y$. The radial scaling derivative satisfies*

$$\boxed{\mathcal{R}_v = -\frac{1}{4}\mathfrak{A} + \frac{1}{2}\mathfrak{J} - \frac{1}{4}yH - \frac{1}{2}s\mathfrak{T}.} \tag{9.13}$$

Moreover the exact radial fundamental theorem is

$$\boxed{v(X,t) - a(t) = \int_0^1 \mathcal{R}_v(\lambda X, t)\frac{d\lambda}{\lambda}.} \tag{9.14}$$

Consequently

$$\boxed{|v(X,t) - a(t)| \le \int_0^1 \left(\frac{1}{4}|\mathfrak{A}| + \frac{1}{2}|\mathfrak{J}| + \frac{1}{4}y_\lambda|H| + \frac{1}{2}s|\mathfrak{T}|\right)(\lambda X, t)\frac{d\lambda}{\lambda},} \tag{9.15}$$

where $y_\lambda := \lambda y$. Since

$$g - 2a = 2(v - a) - \mathfrak{A}, \tag{9.16}$$

the same logarithmic star hull controls both centered value defects.

Remark 9.5 (Correction of the inherited radial measure). An earlier proof draft inherited the formula $\int_0^1 \mathcal{R}_v(\lambda X)d\lambda$. Equation (9.14) shows that the exact radial measure is $d\lambda/\lambda$. The same correction is used consistently throughout the manuscript statement that uses the radial fundamental theorem, in particular proposition 12.3 and corollary 17.6. The corrected formulas are authoritative for the grand-finale chain.

Proposition 9.6 (First-order bridge among restart channels)**.** *With $s := y/x$ and the restart basis of definition 9.1, the axial-relief field obeys*

$$\boxed{x\partial_x\mathfrak{A} = \mathfrak{J} - \frac{1}{2}\mathfrak{A} + \frac{1}{4}xH - \frac{1}{2}\mathfrak{T},} \tag{9.17}$$

$$\boxed{y\partial_y\mathfrak{A} = -\frac{1}{2}\mathfrak{J} - \frac{1}{4}\mathfrak{A} - \frac{1}{2}yH - s\mathfrak{T}.} \tag{9.18}$$

Equivalently,

$$\boxed{yH = -2y\partial_y\mathfrak{A} - \mathfrak{J} - \frac{1}{2}\mathfrak{A} - 2s\mathfrak{T}.} \tag{9.19}$$

Hence the signed restart channels are algebraically coupled and cannot be treated as independent debts.

Lemma 9.7 (Signed log-box thickening)**.** *Fix a time t with $a(t) > 0$, a point $X_* = (x_*, y_*) \in \mathcal{Q}$, and $\rho > 0$. Define the logarithmic box*

$$\mathcal{B}_\rho(X_*) := \{(x, y) : x = x_*e^\xi,\ y = y_*e^\eta,\ |\xi| < \rho,\ |\eta| < \rho\}.$$

Let a C^1 restart scalar $\mathcal{F}$ satisfy

$$\mathcal{F}(X_*) > c\,a(t), \qquad c > 0. \tag{9.20}$$

Then either

$$\boxed{\int_{\mathcal{B}_\rho(X_*)} [\mathcal{F}]_+ \frac{dxdy}{xy} \ge 2c\rho^2 a(t),} \tag{9.21}$$

or

$$\boxed{\sup_{\mathcal{B}_\rho(X_*)} (|x\mathcal{F}_x| + |y\mathcal{F}_y|) > \frac{c}{4\rho} a(t).} \tag{9.22}$$

Thus every pointwise restart either thickens into a scale-invariant signed payment or forces a higher derivative exit.

10. Positive and negative genealogy banks

Accounting bridge. The source history is now split into two nonnegative banks. This makes cancellation visible as an actual transfer of balance rather than as a sign hidden inside one signed Hodge integral.

Definition 10.1 (Source-polarized genealogy)**.** *Let $H^\pm$ solve*

$$\boxed{\partial_t H^\pm + \nabla\cdot(\mathbf{b}H^\pm) = 2[q]_\pm, \qquad H^\pm(\cdot,0) = 0, \qquad \mathbf{b} = (-2xv, 2yg).} \tag{10.1}$$

Then

$$\boxed{H = H^+ - H^-, \qquad H^\pm \geq 0.} \tag{10.2}$$

Define

$$\boxed{a^\pm := \frac{3}{8}\int K_H H^\pm, \qquad a = a^+ - a^-.} \tag{10.3}$$

Proposition 10.2 (Exact two-bank equations)**.** *Put*

$$\mathcal{S}_\pm := \frac{3}{4}\int_{\mathcal{Q}} K_H[q]_\pm \, dX, \qquad \mathcal{M}_\pm := \frac{3}{8}\int_{\mathcal{Q}} K_H \mathcal{C} H^\pm \, dX, \tag{10.4}$$

where

$$\mathcal{C} := \frac{5v(1-2s) + (4s-1)\mathfrak{A}}{1+s}, \qquad s := y/x. \tag{10.5}$$

Then

$$\boxed{\dot a^+ = \mathcal{S}_+ + \mathcal{M}_+, \qquad \dot a^- = \mathcal{S}_- + \mathcal{M}_-.} \tag{10.6}$$

For the carrier estimates, use

$$C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y}, \qquad d\mu_c := \frac{dX}{\mathfrak{r}}, \qquad \mathcal{Q}_{\mathrm{sg}}^{(1)} := \|q\|_{L^1(d\mu_c)} + \|C_\beta\|_{L^1(d\mu_c)},$$

and let

$$\mathbf{B}(\mathbf{X},\mathbf{Z},t) := (-v\mathbf{X}, g\mathbf{Z}), \qquad \mathcal{L}_B(t) := \|\nabla_{\mathbb{R}^7}\mathbf{B}(t)\|_{L^\infty}.$$

Then

$$|\mathcal{M}_\pm| \leq \mathfrak{c}\mathcal{L}_B a^\pm, \qquad \mathcal{S}_+ + \mathcal{S}_- \leq \mathfrak{c}\mathcal{Q}_{\mathrm{sg}}^{(1)}, \tag{10.7}$$

where $\mathfrak{c}$ is an absolute comparison constant.

Theorem 10.3 (Finite-action floor equals cancellation lock)**.** *Let $[t_0, T)$ be a finite classical interval. Define*

$$\mathcal{L}_\star(t) := \mathcal{L}_B(t) + 4\|yv_y(t)\|_\infty + \|xg_x(t)\|_\infty, \qquad \mathcal{K}_\star([t_0,T)) := \int_{t_0}^T \mathcal{L}_\star(t)\, dt.$$

Assume

$$\mathcal{K}_\star([t_0,T)) < \infty, \qquad \sup_{t_0\leq t<T} \mathcal{Q}_{\mathrm{sg}}^{(1)}(t) < \infty,$$

where $\mathcal{L}_B$ and $\mathcal{Q}_{\mathrm{sg}}^{(1)}$ are defined in proposition 10.2. Then both a^+ and a^- have finite limits as $t \uparrow T$. If $t_0 > 0$, then

$$\lim_{t\uparrow T} a^+(t) > 0. \tag{10.8}$$

If additionally

$$a(t) > 0, \qquad a(t) \to 0 \quad (t \uparrow T), \tag{10.9}$$

then there exists $L > 0$ such that

$$\boxed{a^+(t) \to L, \qquad a^-(t) \to L \quad (t \uparrow T).} \tag{10.10}$$

Thus floor loss is not disappearance of genealogy; it is increasingly exact cancellation of two persistent Hodge banks.

Definition 10.4 (Cancellation score)**.** *When $a^- > 0$, define the physical routing ratio*

$$\boxed{\upsilon_- := \frac{a^-}{a^+}.} \tag{10.11}$$

Then a finite-action floor is exactly $\upsilon_- \to 1$.

Theorem 10.5 (Exact cancellation ledger)**.** *Assume the hypotheses of theorem 10.3 on a finite interval $[t_0, T)$, including the floor condition $a(t) > 0$ and $a(t) \to 0$ as $t \uparrow T$. Choose $t_- \in (t_0, T)$ sufficiently late that $a^\pm > 0$ on $[t_-, T)$, and let $\mathcal{S}_\pm, \mathcal{M}_\pm$ be the accounts from proposition 10.2. Define*

$$\upsilon_- := \frac{a^-}{a^+}, \qquad \sigma_\pm := \frac{\mathcal{S}_\pm}{a^\pm}, \qquad c_\pm := \frac{\mathcal{M}_\pm}{a^\pm}.$$

Then

$$\boxed{\frac{d}{dt}\log \upsilon_- = \sigma_- - \sigma_+ + c_- - c_+.} \tag{10.12}$$

Moreover

$$\boxed{\int_{t_-}^{T} (\sigma_+ + \sigma_- + |c_+| + |c_-|)\, dt < \infty,} \tag{10.13}$$

and therefore

$$\boxed{\int_{t_-}^{T} (\sigma_- - \sigma_+ + c_- - c_+)\, dt = \log \frac{1}{\upsilon_-(t_-)} < \infty.} \tag{10.14}$$

Thus no factor $1/a$ is needed: υ_- is a finite-work scorekeeper for the cancellation lock, not a divergent terminal currency.

11. Cashier persistence and the no-reset side of the ledger

Accounting bridge. A banked credit cannot be erased merely by changing the accounting window. The next statements formalize this no-reset principle and specify what must be paid when an old cashier loses visibility.

Proposition 11.1 (Fresh positive genealogy near the apex)**.** *In the physical sector $\varepsilon_\beta = 1$, for the oriented seed $\psi_\beta(0) > 0$ and every finite classical time $t_0 > 0$, put*

$$s := y/x, \qquad q_H(t) := \partial_y H(0, 0, t).$$

Then $q_H(t_0) > 0$ and, uniformly for s in every compact angular interval $J \Subset (0,\infty)$,

$$H^+(x, sx, t_0) = q_H(t_0)sx + O_{t_0,J}(x^2), \tag{11.1}$$

while

$$H^-(x, sx, t_0) = O_{t_0,J}(x^2). \tag{11.2}$$

Hence sufficiently small conical annuli contain fresh positive genealogy and no first-order negative bank.

Proposition 11.2 (Positive-genealogy material cashier)**.** *Let $\mathcal{B}(t) = \mathscr{X}_{t,t_0}(\mathcal{B}_0)$ be a compact material block with $\mathcal{B}_0 \Subset \mathcal{Q}$, and define*

$$M^+_{\mathcal{B}}(t) := \int_{\mathcal{B}(t)} \frac{H^+}{x}\, dX, \qquad a^+_{\mathcal{B}}(t) := \frac{3}{8}\int_{\mathcal{B}(t)} K_H H^+\, dX.$$

Then

$$\boxed{\dot{M}^+_{\mathcal{B}} = 2\int_{\mathcal{B}(t)} v\frac{H^+}{x}\, dX + 2\int_{\mathcal{B}(t)} \frac{[q]_+}{x}\, dX.} \tag{11.3}$$

In particular the source contribution is nonnegative, so a later negative-source episode cannot debit previously created positive genealogy. Whenever $a > 0$ and $M^+_{\mathcal{B}} > 0$, the intrinsic rate $\Lambda^+_{\mathcal{B}}$ of definition 12.1 rewrites (11.3) as

$$\boxed{\dot{M}^+_{\mathcal{B}} = 2a\Lambda^+_{\mathcal{B}} M^+_{\mathcal{B}}.} \tag{11.4}$$

Proposition 11.3 (Credit-reset firewall)**.** *Let $\mathcal{B}(t) = \mathscr{X}_{t,t_0}(\mathcal{B}_0)$ be a compact material positive-genealogy block and assume that for some $x_{\mathrm{launch}} > 0$*

$$\mathcal{B}(t_0) \subset \{x_{\mathrm{launch}}/2 < x < x_{\mathrm{launch}}\}.$$

Fix $0 < \theta < 1/2$. If at a later classical time $t > t_0$

$$M^+_{\mathcal{B}}(t) < \theta M^+_{\mathcal{B}}(t_0),$$

then some material trajectory $X(\tau) = (x(\tau), y(\tau)) \in \mathcal{B}(\tau)$ satisfies

$$\boxed{\int_{t_0}^{t} [-v(X(\tau),\tau)]_+\, d\tau \geq \frac{1}{2}\log\frac{1}{2\theta}.} \tag{11.5}$$

If in addition $a(\tau) > 0$ on $[t_0, t]$, put $X_\lambda(\tau) := \lambda X(\tau)$, $y_\lambda(\tau) := \lambda y(\tau)$, and $s(\tau) := y(\tau)/x(\tau)$. Then the authoritative logarithmic star-hull corrected radial identity gives

$$\boxed{\begin{aligned}\int_{t_0}^{t}\int_0^1 \Big(&\tfrac{1}{4}[\mathfrak{A}]_+(X_\lambda(\tau),\tau) + \tfrac{1}{2}[-\mathfrak{J}]_+(X_\lambda(\tau),\tau)\\ &+ \tfrac{1}{4}y_\lambda(\tau)H^+(X_\lambda(\tau),\tau) + \tfrac{1}{2}s(\tau)[\mathfrak{T}]_+(X_\lambda(\tau),\tau)\Big)\frac{d\lambda}{\lambda}\, d\tau\\ &\geq \frac{1}{2}\log\frac{1}{2\theta}.\end{aligned}} \tag{11.6}$$

Thus loss of a fixed fraction of an old cashier pays a definite amount of the exact restart basis; a fresh relaunch cannot erase old credit for free.

Remark 11.4 (Why this does not already close). The no-reset mechanism controls repeated loss of selected positive cashiers. It does not prohibit one finite physical transfer of enough Hodge weight into H^- to create the global cancellation lock. That one-shot conversion is the modern remaining bill.

12. The Hodge-matched cashier and relaunch firewalls

Accounting bridge. The no-reset ledger is useful only if new favorable cashiers can be launched without new capital. The Hodge-matched construction shows how the same original investment supplies fresh cashier opportunities while terminal-uniformity is audited honestly.

Definition 12.1 (Intrinsic positive-genealogy cashier)**.** *For a material block $\mathcal{B}(t)$ define*

$$M_{\mathcal{B}}^{+} := \int_{\mathcal{B}(t)} \frac{H^{+}}{x}\, dX, \qquad a_{\mathcal{B}}^{+} := \frac{3}{8}\int_{\mathcal{B}(t)} K_H H^{+}\, dX. \tag{12.1}$$

Whenever $M_{\mathcal{B}}^{+} > 0$, set

$$\boxed{\eta_{\mathcal{B}}^{+} := \frac{a_{\mathcal{B}}^{+}}{M_{\mathcal{B}}^{+}}.} \tag{12.2}$$

If also $a > 0$, define the intrinsic rate

$$\boxed{\Lambda_{\mathcal{B}}^{+} := \frac{\displaystyle\int_{\mathcal{B}} v\frac{H^{+}}{x}dX + \int_{\mathcal{B}} \frac{[q]_{+}}{x}dX}{aM_{\mathcal{B}}^{+}}.} \tag{12.3}$$

Then

$$\boxed{\dot{M}_{\mathcal{B}}^{+} = 2a\Lambda_{\mathcal{B}}^{+} M_{\mathcal{B}}^{+}.} \tag{12.4}$$

Theorem 12.2 (Intrinsic positive-genealogy Riccati closure)**.** *Let $\mathcal{B}(t) = \mathscr{X}_{t,t_0}(\mathcal{B}_0)$ be a compact material block with $\mathcal{B}_0 \Subset \mathcal{Q}$, and let $M_{\mathcal{B}}^{+}, a_{\mathcal{B}}^{+}, \eta_{\mathcal{B}}^{+}, \Lambda_{\mathcal{B}}^{+}$ be the cashier, Hodge credit, visibility, and intrinsic rate defined in definition 12.1. Suppose on a classical interval*

$$\boxed{a > 0, \qquad a \geq a_{\mathcal{B}}^{+}, \qquad \eta_{\mathcal{B}}^{+} \geq m_* > 0, \qquad \Lambda_{\mathcal{B}}^{+} \geq \nu > 0.} \tag{12.5}$$

Then

$$\boxed{\dot{M}_{\mathcal{B}}^{+} \geq 2\nu m_* (M_{\mathcal{B}}^{+})^2.} \tag{12.6}$$

Hence either regularity fails earlier or $M_{\mathcal{B}}^{+}$ diverges by

$$T_* \leq t_0 + \frac{1}{2\nu m_* M_{\mathcal{B}}^{+}(t_0)}, \tag{12.7}$$

provided the displayed gates hold from the launch time t_0 onward. No sign assumption on the current signed field H is required.

Proposition 12.3 (Rate failure pays radial non-affinity)**.** *Let $\mathcal{B}(t)$, $M_{\mathcal{B}}^{+}$, and $\Lambda_{\mathcal{B}}^{+}$ be as in definition 12.1. Fix $0 < \nu < 1$, put $\mathcal{R}_v := xv_x + yv_y$, and suppose $\Lambda_{\mathcal{B}}^{+} < \nu$. Then*

$$\boxed{\int_{\mathcal{B}} (a - v)\frac{H^{+}}{x}dX > (1-\nu)aM_{\mathcal{B}}^{+} + \int_{\mathcal{B}} \frac{[q]_{+}}{x}dX.} \tag{12.8}$$

Consequently

$$\boxed{\int_{\mathcal{B}}\int_0^1 [-\mathcal{R}_v]_{+}(\lambda X, t)\frac{d\lambda}{\lambda}\frac{H^{+}}{x}dX > (1-\nu)aM_{\mathcal{B}}^{+}.} \tag{12.9}$$

Thus an intrinsic rate exit is a paid logarithmic star-hull bill.

Theorem 12.4 (Visibility collapse is positive-genealogy tail concentration)**.** *Let $M_{\mathcal{B}}^{+}$ and $\eta_{\mathcal{B}}^{+}$ be as in definition 12.1, and put*

$$h_0(s) := \frac{3}{8}\frac{\sqrt{s}}{(1+s)^{5/2}}, \qquad h_{\max} := \max_{s>0} h_0(s) = \frac{6}{25\sqrt{5}}.$$

Define the nonnegative measure

$$d\mu_{\mathcal{B}}^{+} := \frac{H^{+}}{x}\, dX, \qquad \mu_{\mathcal{B}}^{+}(\mathcal{Q}) = M_{\mathcal{B}}^{+}.$$

If

$$\eta_{\mathcal{B}}^{+} < m_* < h_{\max}, \tag{12.10}$$

then for every $\vartheta \in (m_*, h_{\max})$,

$$\boxed{\mu_{\mathcal{B}}^{+}\{X \in \mathcal{B} : h_0(y/x) < \vartheta\} > \left(1 - \frac{m_*}{\vartheta}\right) M_{\mathcal{B}}^{+}.} \tag{12.11}$$

Thus visibility failure moves a fixed fraction of already banked positive genealogy into the angular tails; it is not disappearance of credit.

Theorem 12.5 (Finite intrinsic exit graph)**.** *Fix thresholds*

$$0 < \nu < 1, \qquad 0 < m_* < h_{\max},$$

and an admitted classical interval on which $a > 0$ and $M_{\mathcal{B}}^{+} > 0$ for a material positive-genealogy cashier $\mathcal{B}(t)$. Use the quantities $M_{\mathcal{B}}^{+}, a_{\mathcal{B}}^{+}, \eta_{\mathcal{B}}^{+}, \Lambda_{\mathcal{B}}^{+}$ from definition 12.1. Exactly the following intrinsic branches remain:

- *(i) the three intrinsic gates $a \geq a_{\mathcal{B}}^{+}$, $\eta_{\mathcal{B}}^{+} \geq m_*$, and $\Lambda_{\mathcal{B}}^{+} \geq \nu$ persist, and Riccati closure occurs;*
- *(ii) the rate gate fails: $\Lambda_{\mathcal{B}}^{+} < \nu$, which pays radial star-hull non-affinity;*
- *(iii) the visibility gate fails: $\eta_{\mathcal{B}}^{+} < m_*$, which places positive genealogy in the angular tails;*
- *(iv) Hodge dominance fails: $a < a_{\mathcal{B}}^{+}$, which activates H^{-} and hence routes to old negative ancestry or tagged cross-shear.*

There is no source-sign reset and no ancestry-free fifth intrinsic branch. Loss of the outer admission $a > 0$ is a separate compression-floor exit and is not counted as an intrinsic cashier branch.

Definition 12.6 (Frozen positive-genealogy cohort)**.** *At a launch time $t_0 > 0$, let*

$$\mathcal{B}(t) = \mathscr{X}_{t,t_0}(\mathcal{B}_0), \qquad \mathcal{B}_0 \Subset \mathcal{Q},$$

be a compact material block. Freeze a nonnegative cohort $K_{\mathcal{B}}$ by

$$\partial_t K_{\mathcal{B}} + \nabla\cdot(\mathbf{b} K_{\mathcal{B}}) = 0, \qquad K_{\mathcal{B}}(t_0) = H^{+}(t_0)\mathbf{1}_{\mathcal{B}_0}. \tag{12.12}$$

Thus $K_{\mathcal{B}}$ remains supported in $\mathcal{B}(t)$; by the positive-genealogy Duhamel formula, $0 \leq K_{\mathcal{B}} \leq H^{+}$ thereafter. Put

$$h_0(s) := \frac{3}{8}\frac{\sqrt{s}}{(1+s)^{5/2}}, \tag{12.13}$$

$$a_K := \int \frac{h_0(s)}{x} K_{\mathcal{B}}\, dX, \qquad N_K := \int K_{\mathcal{B}}\, dX, \tag{12.14}$$

$$\Pi_H := \int \frac{x K_{\mathcal{B}}}{h_0(s)}\, dX. \tag{12.15}$$

Lemma 12.7 (Hodge-matched Cauchy bridge)**.** *Let $K_{\mathcal{B}}, h_0, a_K, N_K, \Pi_H$ be the frozen cohort and its three moments from definition 12.6. Then*

$$\boxed{N_K^2 \le a_K \Pi_H.} \tag{12.16}$$

If $\Pi_H > 0$, define

$$\mathfrak{Z}_H := \frac{N_K^2}{\Pi_H}. \tag{12.17}$$

Then $\mathfrak{Z}_H \le a_K$.

Proposition 12.8 (Exact dual-moment rate)**.** *Let $K_{\mathcal{B}}$, h_0, and Π_H be as in definition 12.6, put $s := y/x$, and recall the restart variable $\mathfrak{A} := 2v - g$. Define*

$$\mathcal{C}_H(x, s, t) := \frac{5v(1-2s) + \mathfrak{A}(4s-1)}{1+s}. \tag{12.18}$$

Then

$$\boxed{-\dot{\Pi}_H = \int_{\mathcal{Q}} \frac{x K_{\mathcal{B}}}{h_0(s)} \mathcal{C}_H(x, s, t) dX.} \tag{12.19}$$

Theorem 12.9 (Hodge-matched Riccati gate)**.** *Let $K_{\mathcal{B}}$ be a nontrivial frozen cohort from definition 12.6, with moments a_K, N_K, Π_H. Suppose on a time interval*

$$-\dot{\Pi}_H \ge \kappa a \Pi_H, \qquad a \ge a_K, \qquad \kappa > 0. \tag{12.20}$$

Then, before any quotient is formed,

$$\boxed{-N_K^2 \dot{\Pi}_H \ge \kappa N_K^4.} \tag{12.21}$$

Since the cohort is nontrivial, $\Pi_H > 0$ and $N_K > 0$; only after the raw gate (12.21) has been established define

$$\mathfrak{Z}_H := \frac{N_K^2}{\Pi_H}.$$

Then

$$\boxed{0 < \mathfrak{Z}_H \le a_K \le a, \qquad \dot{\mathfrak{Z}}_H \ge \kappa a \mathfrak{Z}_H \ge \kappa \mathfrak{Z}_H^2.} \tag{12.22}$$

If the raw gates persist from t_0, they cannot persist beyond

$$T_{H,\mathrm{Ric}} := t_0 + \frac{1}{\kappa \mathfrak{Z}_H(t_0)}. \tag{12.23}$$

Thus either a displayed raw gate fails first or the classical solution loses regularity by the finite Hodge-matched Riccati deadline.

Theorem 12.10 (Automatic fresh relaunch)**.** *Fix a finite classical time $t_0 > 0$ in the physical sector $\varepsilon_\beta = 1$ and assume $a(t_0) > 0$. Let*

$$J = [s_-, s_+] \Subset (0, 1/2), \qquad s := y/x,$$

and define

$$c_J := 5 \min_{s \in J} \frac{1-2s}{1+s} > 0. \tag{12.24}$$

Put $q_H(t) := \partial_y H(0, 0, t)$ and recall $q := \Theta_y$. Then $u_y(0, 0, t_0) > 0$ and $q_H(t_0) > 0$, and there exists $\ell_0 > 0$ such that for every $0 < \ell \le \ell_0$ the conical annulus

$$\mathcal{B}_\ell(t_0) := \{\ell/2 < x < \ell, \ s \in J\} \tag{12.25}$$

satisfies

$$H^+(x, sx, t_0) = q_H(t_0)sx + O_{t_0,J}(x^2), \tag{12.26}$$

$$q(x, sx, t_0) = 2u_y(0, 0, t_0)^2 sx + O_{t_0,J}(x^2). \tag{12.27}$$

In particular $H^+ > 0$ and $q > 0$ on $\mathcal{B}_\ell(t_0)$ for ℓ small. Freeze the positive genealogy on this annulus by

$$K_{\mathcal{B}}(t_0) = H^+(t_0)\mathbf{1}_{\mathcal{B}_\ell(t_0)}, \qquad \partial_t K_{\mathcal{B}} + \nabla\cdot(\mathbf{b}K_{\mathcal{B}}) = 0,$$

and let a_K, Π_H be the Hodge-matched moments from definition 12.6. Then, after decreasing ℓ_0 if necessary,

$$\boxed{-\dot{\Pi}_H(t_0) \geq \frac{c_J}{2}a(t_0)\Pi_H(t_0), \qquad a(t_0) - a_K(t_0) \geq \frac{1}{2}a(t_0) > 0.} \tag{12.28}$$

Thus every finite classical time with positive apex compression launches a fresh cohort strictly inside both Hodge-matched raw gates; the allowable radius may depend on t_0.

Theorem 12.11 (Uniform apex tube on compact classical intervals)**.** *Let $I = [t_1, t_2]$ satisfy $0 < t_1 < t_2$ and lie strictly inside the classical lifespan. Assume*

$$a_* := \inf_{t\in I} a(t) > 0. \tag{12.29}$$

Fix $J = [s_-, s_+] \Subset (0, 1/2)$, put $s := y/x$, and define $q_H(t) := \partial_y H(0, 0, t)$ and c_J by (12.24). Then there are constants $\ell_* > 0$ and $\mathfrak{c}_{I,J} < \infty$ such that, for every $t \in I$, every $s \in J$, and $0 < x \leq \ell_*$,

$$v(x, sx, t) = a(t) + O_{I,J}(x), \qquad g(x, sx, t) = 2a(t) + O_{I,J}(x), \tag{12.30}$$

$$H^+(x, sx, t) = q_H(t)sx + O_{I,J}(x^2), \tag{12.31}$$

$$q(x, sx, t) = 2u_y(0, 0, t)^2 sx + O_{I,J}(x^2), \tag{12.32}$$

where each displayed remainder is bounded in absolute value by $\mathfrak{c}_{I,J}$ times the indicated power of x. Moreover

$$\inf_I u_y(0, 0, t) > 0, \qquad \inf_I q_H(t) > 0. \tag{12.33}$$

Consequently, after one fixed decrease of ℓ_*, the same annulus

$$\mathcal{B}_{\ell_*}(t) := \{\ell_*/2 < x < \ell_*,\ s \in J\}$$

launches at every $t \in I$ a frozen cohort satisfying the uniform strict margins

$$\boxed{-\dot{\Pi}_H(t) \geq \frac{c_J}{2}a(t)\Pi_H(t), \qquad a(t) - a_K(t) \geq \frac{1}{2}a_*.} \tag{12.34}$$

Thus a compact regular positive-compression interval cannot hide a shrinking relaunch tube. The restriction $t_1 > 0$ is essential for the uniform positive lower bound on q_H, since the pure-swirl launch has $q_H(0) = 0$.

Theorem 12.12 (Terminal-time uniformity firewall)**.** *Let $T < \infty$ and suppose the Euler solution is classical on $[t_1, T)$ with $t_1 > 0$. Fix $J \Subset (0, 1/2)$. Assume first that there is one fixed apex neighborhood on which all spatial derivatives needed for the remainders in theorem 12.11 remain uniformly bounded as $t \uparrow T$, and put*

$$A^* := \sup_{t_1\leq t<T} [a(t)]_+ < \infty. \tag{12.35}$$

If in addition

$$a_* := \inf_{t_1 \le t < T} a(t) > 0, \tag{12.36}$$

then there is one time-independent radius $\ell_* > 0$ for which all conclusions and uniform Hodge-matched launch margins of theorem 12.11 hold for every $t \in [t_1, T)$.

If the required local derivative bounds or $A^* < \infty$ fail, that failure is a genuine obstruction to classical extension through T and may not be hidden inside $O(x)$ notation. If instead

$$\inf_{t_1 \le t < T} a(t) = 0, \tag{12.37}$$

then either classical regularity fails at or before T, or there is a smooth zero-compression contact time $t_* \in [t_1, T]$ (after a classical extension through T when necessary). At such a contact $q_H(t_*) > 0$, so a sufficiently small positive-genealogy block has $a^+_{\mathcal{B}}(t_*) > 0 = a(t_*)$; hence Hodge dominance fails and the branch is routed to negative-genealogy ancestry. There is no hidden third terminal branch inside the Taylor remainder notation.

Proposition 12.13 (No-Zeno relaunch on a regular positive-compression interval)**.** *Let $I = [T_0, T_1]$ with $0 < T_0 < T_1$ be a compact interval strictly inside the classical lifespan and suppose*

$$a(t) \ge a_* > 0 \qquad (t \in I).$$

Fix $J \Subset (0, 1/2)$ and use the common annulus and constants supplied by theorem 12.11. Consider a relaunch protocol with times $t_n \in I$ which, at each t_n, freezes the fresh cohort on that common annulus and does not relaunch again until one of the weaker Hodge-matched margins

$$-\dot{\Pi}_H \ge \frac{c_J}{4} a \Pi_H, \qquad a - a_K \ge \frac{1}{4} a_* \tag{12.38}$$

first fails. Then there exists $\tau_* > 0$, depending only on the solution on I and on J, such that

$$\boxed{t_{n+1} - t_n \ge \tau_*} \tag{12.39}$$

whenever both relaunch times lie in I. Hence such relaunches cannot accumulate at a finite regular time. Any finite-time Zeno accumulation forces either loss of a positive compression floor or loss of one of the local classical smoothness bounds required by the terminal-time firewall.

13. Negative genealogy has only two ancestries

Accounting bridge. Every adverse balance must now receive a receipt. The ancestry theorem shows that there are only two such receipts: fixed old negative slope and later tagged cross-shear conversion.

Proposition 13.1 (Exact slope ancestry)**.** *Let*

$$p := u_y, \qquad \Gamma := xu,$$

fix $X_0 \in \mathcal{Q}$, and write $X(\tau) := \mathscr{X}_{\tau,0} X_0$. Define along this material characteristic

$$\mu(t) := \exp\left(- \int_0^t 2(v - g - y g_y)(X(\tau), \tau)\, d\tau \right), \qquad p_0 := p(X_0, 0). \tag{13.1}$$

Then

$$\boxed{p(X(t), t) = \mu(t)^{-1} \left[p_0 + 2 \int_0^t \mu(\tau) v_y(X(\tau), \tau) \Gamma_x(X(\tau), \tau)\, d\tau \right].} \tag{13.2}$$

The sign of the fixed carrier is preserved by the material swirl equation; hence $u \geq 0$ and

$$[q]_- = [\Theta_y]_- = 2\varepsilon_\beta u[-p]_+. \tag{13.3}$$

More precisely, define the two nonnegative material source majorants

$$q_-^{\mathrm{old}}(X(t),t) := 2\varepsilon_\beta u(X(t),t)\mu(t)^{-1}[-p_0]_+, \tag{13.4}$$

$$q_-^{\mathrm{sh}}(X(t),t) := 4\varepsilon_\beta u(X(t),t)\mu(t)^{-1}\int_0^t \mu(\tau)[-v_y\Gamma_x]_+(X(\tau),\tau)\,d\tau. \tag{13.5}$$

Then

$$\boxed{[q(X(t),t)]_- \leq q_-^{\mathrm{old}}(X(t),t) + q_-^{\mathrm{sh}}(X(t),t).} \tag{13.6}$$

Thus negative source has only two ancestries:

(i) the fixed initial negative slope;
(ii) later tagged cross-shear $[-v_y\Gamma_x]_+$.

There is no third adverse source node.

Proposition 13.2 (Old source is geometrically separated)**.** *For the fixed $A = 10$ datum define*

$$\mathcal{N}_0 := \{(x,y) \in \mathcal{Q} : u_{0,y}(x,y) < 0\}, \qquad \sigma_- := \frac{3-\sqrt{7}}{4} = \frac{1}{6+2\sqrt{7}}. \tag{13.7}$$

Then

$$\boxed{\mathcal{N}_0 \subset \{y \geq \sigma_-\}.} \tag{13.8}$$

In the physical sector put

$$\Xi_\beta := \Theta\frac{x}{y^3}, \qquad M_{3,\mathcal{N}}(t) := \int_{\mathcal{N}(t)} \Xi_\beta(X,t)\,dX, \qquad \mathcal{N}(t) := \mathscr{X}_{t,0}\mathcal{N}_0. \tag{13.9}$$

Then

$$\boxed{M_{3,\mathcal{N}}(0) < \infty.} \tag{13.10}$$

Fix $0 < \delta < 1$. Suppose $a > 0$ on $(0,T)$ and every trajectory of $\mathcal{N}(t)$ remains in the near-affine corridor

$$|v - a| + |g - 2a| \leq \delta a, \tag{13.11}$$

$$2\big(|xv_x| + |yv_y| + |xg_x| + |yg_y|\big) \leq \delta a. \tag{13.12}$$

Define its signed source-response contribution by

$$\begin{aligned}\mathcal{P}^{\mathrm{src}}_{\beta,\mathcal{N}}(t) := \frac{3}{4}\int_{\mathcal{N}(t)} \Theta(X,t)\mathcal{R}_0(X)\,dX \\ + \frac{3}{4}\int_0^t \int_{\mathcal{N}(\tau)} \Theta(X,\tau)\partial_t\mathcal{R}(t,\tau;X)\,dX\,d\tau.\end{aligned} \tag{13.13}$$

Then a constant $K_\delta < \infty$ gives

$$\boxed{|\mathcal{P}^{\mathrm{src}}_{\beta,\mathcal{N}}(t)| \leq K_\delta\left[M_{3,\mathcal{N}}(t) + a(t)\int_0^t M_{3,\mathcal{N}}(\tau)\,d\tau\right],} \tag{13.14}$$

and

$$\boxed{M_{3,\mathcal{N}}(t) \leq M_{3,\mathcal{N}}(0)\,\mathfrak{t}_\Psi(t)^{\,4-\frac{5}{2}\delta}.} \tag{13.15}$$

Consequently, on every hypothetical finite-time Riccati branch with $a(t) \to \infty$ and $\mathfrak{t}_\Psi(t) \to 0$,

$$\boxed{\frac{|\mathcal{P}^{\mathrm{src}}_{\beta,\mathcal{N}}(t)|}{a(t)^2} \longrightarrow 0.} \tag{13.16}$$

If the corridor fails first, the failure is one of the already protected value/first-jet quantities and enters the established non-affine restart routing. Thus the fixed old negative-source ancestry is either terminally Hodge-silent in the controlled corridor or leaves that corridor through a named restart channel; this proposition does not by itself assert a quantitative lower bound for the restart payment.

14. Continuation and carrier-capacity lemmas

Accounting bridge. These lemmas cap the ordinary operating expenses of the ledger. If one of the stated capacities fails, Peter has already reached a named continuation exit rather than an unclassified hidden debt.

Definition 14.1 (Carrier measure and source-gradient stock)**.** *Let*

$$d\mu_c := \frac{dX}{\mathfrak{r}}, \qquad \mathfrak{r} = x^{2/3}y^{1/3}, \tag{14.1}$$

and

$$\mathcal{Q}^{(1)}_{\mathrm{sg}}(t) := \|q(t)\|_{L^1(d\mu_c)} + \|C_\beta(t)\|_{L^1(d\mu_c)}. \tag{14.2}$$

The homogeneous source-gradient pair is governed by $\mathbf{A}_{\mathrm{sg}}$ from the rotor section.

Proposition 14.2 (Raw source-gradient Gronwall estimate)**.** *Define*

$$\mathsf{d}_q := \frac{3}{2}\mathfrak{A} - \mathfrak{J}$$

and

$$L_{\mathrm{sg}}(t) := \|\mathsf{d}_q(t)\|_\infty + \|\tfrac{3}{2}\mathfrak{A}(t) + \mathfrak{J}(t)\|_\infty + 4\|yv_y(t)\|_\infty + \|xg_x(t)\|_\infty. \tag{14.3}$$

Then on every classical interval $[t_0, t]$,

$$\boxed{\|(q, C_\beta)(t)\|_{L^\infty} \le \|(q, C_\beta)(t_0)\|_{L^\infty} \exp\left(\int_{t_0}^{t} L_{\mathrm{sg}}(\tau)d\tau\right).} \tag{14.4}$$

For the carrier measure $d\mu_c = dX/\mathfrak{r}$, set

$$\delta_c := \frac{4v + g}{3}. \tag{14.5}$$

For every $\mu, \nu \ge 0$ define

$$\mathcal{Q}^{(1;\mu,\nu)}_{\mathrm{sg}}(t) := \int_{\mathcal{Q}} x^\mu y^\nu \big(|q| + |C_\beta|\big)\, d\mu_c, \tag{14.6}$$

$$\mathcal{E}^{(\mu,\nu)}_{\mathrm{sg}}(t) := \|x^\mu y^\nu (q, C_\beta)\|_{L^2(d\mu_c;\ell^2)}. \tag{14.7}$$

Put

$$L^{(1;\mu,\nu)}_{\mathrm{sg}} := L_{\mathrm{sg}} + 2\mu\|v\|_\infty + 2\nu\|g\|_\infty + \|\delta_c\|_\infty, \tag{14.8}$$

$$L^{(2;\mu,\nu)}_{\mathrm{sg}} := L_{\mathrm{sg}} + 2\mu\|v\|_\infty + 2\nu\|g\|_\infty + \frac{1}{2}\|\delta_c\|_\infty. \tag{14.9}$$

Then, in the upper-right Dini sense,

$$D_t^+ \mathcal{Q}_{\mathrm{sg}}^{(1;\mu,\nu)} \le L_{\mathrm{sg}}^{(1;\mu,\nu)} \mathcal{Q}_{\mathrm{sg}}^{(1;\mu,\nu)}, \tag{14.10}$$

$$D_t^+ \mathcal{E}_{\mathrm{sg}}^{(\mu,\nu)} \le L_{\mathrm{sg}}^{(2;\mu,\nu)} \mathcal{E}_{\mathrm{sg}}^{(\mu,\nu)}. \tag{14.11}$$

Consequently both stocks obey the corresponding Gronwall bounds. In particular, $(\mu,\nu) = (0,0)$ *gives the carrier-*L^1 *source-gradient stock below, while* $(\mu,\nu) = (1/2,1/2)$ *gives the weighted* L^2 *stock driving the Hodge energy.*

Definition 14.3 (Weighted Hodge carrier energy)**.** *Set*

$$\mathcal{E}_{H,1} := \|\sqrt{xy}\, H\|_{L^2(d\mu_c)}, \qquad \mathcal{E}_{\mathrm{sg}}^{(1/2,1/2)} := \|\sqrt{xy}(q, C_\beta)\|_{L^2(d\mu_c;\ell^2)}. \tag{14.12}$$

Proposition 14.4 (Weighted Hodge energy is not a new mechanism)**.** *Let* $d\mu_c = dX/\mathfrak{r}$ *be the carrier measure from definition 14.1, and define*

$$\mathcal{E}_{H,1} := \|\sqrt{xy}\, H\|_{L^2(d\mu_c)}, \qquad \mathcal{E}_{\mathrm{sg}}^{(1/2,1/2)} := \|\sqrt{xy}(q, C_\beta)\|_{L^2(d\mu_c;\ell^2)}.$$

Put

$$M_H := \sqrt{xy}\, H. \tag{14.13}$$

Then

$$\boxed{\mathcal{D}_{xy} M_H = -v M_H + 2\sqrt{xy}\, q.} \tag{14.14}$$

Let $\mathbf{B}(\mathbf{X},\mathbf{Z},t) = (-v\mathbf{X}, g\mathbf{Z})$ *be the seven-dimensional transport lift and define*

$$\mathcal{L}_B(t) := \|\nabla_{\mathbb{R}^7} \mathbf{B}(t)\|_{L^\infty}.$$

With D_t^+ *denoting the upper right Dini derivative,*

$$\boxed{D_t^+ \mathcal{E}_{H,1} \le \mathfrak{c} \mathcal{L}_B \mathcal{E}_{H,1} + 2\mathcal{E}_{\mathrm{sg}}^{(1/2,1/2)}.} \tag{14.15}$$

Thus the weighted Hodge energy is financed by the same first-jet/source-gradient restart package.

Proposition 14.5 (Physical Biot–Savart control)**.** *Let* $\mathbf{U}_{\mathrm{mer}} := U^r \mathbf{e}_r + U^z \mathbf{e}_z$ *be the physical meridional velocity, let*

$$d\mu_c := \frac{dX}{\mathfrak{r}}$$

be the carrier measure from definition 14.1, and define

$$\mathcal{E}_{H,1} := \|\sqrt{xy}\, H\|_{L^2(d\mu_c)}.$$

With physical point $\boldsymbol{\xi} = (r\cos\phi, r\sin\phi, z)$ *and weight*

$$\omega(\boldsymbol{\xi}) = r^{-4/3}|z|^{1/3}, \tag{14.16}$$

write

$$\|\varphi\|_{L^2_\omega}^2 := \int_{\mathbb{R}^3} |\varphi(\boldsymbol{\xi})|^2 \omega(\boldsymbol{\xi})\, d\boldsymbol{\xi}.$$

Then

$$\boxed{\|\nabla \mathbf{U}_{\mathrm{mer}}\|_{L^2_\omega} \le \mathfrak{c}_\omega \sqrt{\pi}\, \mathcal{E}_{H,1}.} \tag{14.17}$$

Hence the quadratic meridional pressure source is controlled by a GREEN Hodge carrier energy and ordinary first-jet routing norms.

Theorem 14.6 (Finite physical Lipschitz action is a continuation firewall)**.** *Let $[t_0, T)$ be a finite classical interval for the fixed rapidly decaying Euler customer and assume*

$$\mathcal{L}_{\mathrm{phys}}([t_0, T)) := \int_{t_0}^{T} \|\nabla \mathbf{U}(t)\|_\infty \, dt < \infty. \tag{14.18}$$

Then every fixed integer Sobolev order $m \geq 3$ obeys

$$\sup_{t_0 \leq t < T} \|\mathbf{U}(t)\|_{H^m} < \infty, \tag{14.19}$$

and physical particle displacement stays uniformly finite on $[t_0, T)$. If $\boldsymbol{\omega}_{\mathrm{phys}} := \nabla \times \mathbf{U}$ and $\boldsymbol{\xi} = (r\cos\phi, r\sin\phi, z)$, then for every $N > 0$ and $j \in \{0, 1, 2\}$,

$$\sup_{t_0 \leq t < T} \sup_{\boldsymbol{\xi} \in \mathbb{R}^3} (1 + |\boldsymbol{\xi}|)^N |D^j \boldsymbol{\omega}_{\mathrm{phys}}(\boldsymbol{\xi}, t)| < \infty, \tag{14.20}$$

and

$$\lim_{\varrho_0 \to \infty} \sup_{t_0 \leq t < T} \sup_{|\boldsymbol{\xi}| \geq \varrho_0} |\boldsymbol{\xi}| \, |D^2 \mathbf{U}(\boldsymbol{\xi}, t)| = 0. \tag{14.21}$$

Thus the extreme-angle/far-field weighted restart gates cannot fail invisibly while (14.18) stays finite. Moreover, for every subinterval $I \subset [t_0, T)$,

$$\boxed{\left| \int_I a(t)\, dt \right| \leq \int_I \|\nabla \mathbf{U}(t)\|_\infty \, dt.} \tag{14.22}$$

Hence infinite collapse age on a finite interval already forces failure of the standard Euler Lipschitz continuation action.

15. Exact source response and the affine benchmark

Accounting bridge. The affine conveyor is Peter's benchmark balance sheet: it tells us what the same fixed investment would earn under common-mode transport alone. Any worse true-Euler balance must therefore be charged to genuine non-affine work.

Definition 15.1 (Two-time Hodge visibility)**.** *Let $\mathscr{X}_{t,\tau}$ be the material flow of $\mathbf{b} = (-2xv, 2yg)$ from source time τ to observation time t, and define*

$$\boxed{\mathcal{V}_H(t; \tau, X) := K_H(\mathscr{X}_{t,\tau} X) > 0, \qquad 0 \leq \tau \leq t.} \tag{15.1}$$

Then

$$\boxed{a^{\pm}(t) = \frac{3}{4} \int_0^t \int_{\mathcal{Q}} [q(X, \tau)]_{\pm} \mathcal{V}_H(t; \tau, X) \, dX \, d\tau.} \tag{15.2}$$

Proposition 15.2 (Vertical response representation)**.** *For the fixed customer of definition 5.1, so in particular $H(\cdot, 0) = 0$, let $\mathscr{X}_{t,\tau}$ be the flow from definition 15.1 and define*

$$\boxed{\mathcal{R}(t, \tau; X) := -\partial_{y_\tau} K_H(\mathscr{X}_{t,\tau} X),} \tag{15.3}$$

where ∂_{y_τ} denotes differentiation with respect to the y-coordinate of the deposition point $X = (x_\tau, y_\tau)$ at time τ. Then

$$\boxed{a(t) = \frac{3}{4} \int_0^t \int_{\mathcal{Q}} \Theta(X, \tau) \mathcal{R}(t, \tau; X) \, dX \, d\tau.} \tag{15.4}$$

At zero response age,

$$\boxed{\mathcal{R}_0(x,y) := \mathcal{R}(t,t;(x,y)) = \frac{x(4y-x)}{2\sqrt{y}(x+y)^{7/2}}, \qquad \mathcal{R}_0 > 0 \iff y > x/4.} \tag{15.5}$$

Theorem 15.3 (True-Euler aged-response discount)**.** *Fix a source time τ and a source point $X_\tau = (x_\tau, y_\tau) \in \mathcal{Q}$. Let*

$$Z(\xi) = (x(\xi), y(\xi)) := \mathscr{X}_{\xi,\tau} X_\tau, \qquad s_0 := \frac{y_\tau}{x_\tau}, \qquad s(\xi) := \frac{y(\xi)}{x(\xi)},$$

and

$$\Lambda(t;\tau) := \int_\tau^t a(\xi)d\xi, \qquad \frac{\mathfrak{t}_\Psi(t)}{\mathfrak{t}_\Psi(\tau)} = e^{-2\Lambda(t;\tau)}.$$

Assume $a(\xi) > 0$ and that for some $0 < \delta < 1$ the trajectory segment $\tau \le \xi \le t$ satisfies the near-affine corridor

$$|v - a| + |g - 2a| \le \delta a,$$
$$2\big(|xv_x| + |yv_y| + |xg_x| + |yg_y|\big) \le \delta a$$

pointwise along $Z(\xi)$, and assume the transported aspect ratio $s(\xi) \ge 1$ on the aged interval. Then

$$\boxed{|\mathcal{R}(t,\tau;X_\tau)| + a(t)^{-1}|\partial_t \mathcal{R}(t,\tau;X_\tau)| \le \frac{\mathfrak{c}}{x_\tau^2 s_0^3}\left(\frac{\mathfrak{t}_\Psi(t)}{\mathfrak{t}_\Psi(\tau)}\right)^{5-\frac{7}{2}\delta}.} \tag{15.6}$$

Here $\mathfrak{c} > 0$ is a universal comparison constant. Since $0 < \delta < 1$, the exponent is strictly larger than $3/2$, hence in particular strictly larger than one.

Lemma 15.4 (Fixed-seed signed line-source positivity)**.** *Let u_0 be the fixed customer from definition 5.1, define*

$$q_0 := (u_0^2)_y, \qquad \mathcal{L}_q(\eta) := \int_0^\infty q_0(x, \eta x)\, dx \quad (\eta > 0).$$

Then

$$\boxed{\mathcal{L}_q(\eta) > 0 \qquad (\eta > 0).} \tag{15.7}$$

Equivalently, after exact integration the numerator is $81\eta\Pi_{15}(\eta)$, where Π_{15} is a degree-15 polynomial all of whose coefficients are strictly positive.

Theorem 15.5 (Global affine cancellation barrier)**.** *For the fixed customer let*

$$\mathcal{L}_q^\pm(\eta) := \int_0^\infty [q_0(x,\eta x)]_\pm\, dx, \qquad k(\sigma) := \frac{\sqrt{\sigma}}{(1+\sigma)^{5/2}},$$

and define the normalized same-clock affine shape integrals

$$I_\pm(\Lambda) := \int_0^\infty k(\sigma)\mathcal{L}_q^\pm(\sigma e^{-6\Lambda})\, d\sigma, \qquad \Lambda \ge 0. \tag{15.8}$$

Then $I_+(\Lambda) > 0$ for every $\Lambda \ge 0$, and

$$\upsilon_-^{\mathrm{aff}}(\Lambda) := \frac{I_-(\Lambda)}{I_+(\Lambda)} \tag{15.9}$$

equals the ratio of the actual positive/negative Hodge banks at every positive same-clock affine time. Formula (15.9) also defines the continuous launch value at $\Lambda = 0$, avoiding a formal $0/0$ bank ratio there. There exists a fixed $\upsilon^*_{\mathrm{aff}} < 1$ such that

$$\boxed{0 \le \upsilon_-^{\mathrm{aff}}(\Lambda) \le \upsilon^*_{\mathrm{aff}} < 1 \qquad (\Lambda \ge 0).} \tag{15.10}$$

Equivalently,

$$\boxed{\pi^*_{\mathrm{aff}} := \frac{1-\upsilon^*_{\mathrm{aff}}}{1+\upsilon^*_{\mathrm{aff}}} > 0} \tag{15.11}$$

and the affine signed Hodge response is at least π^*_{aff} times its absolute Hodge response.

Theorem 15.6 (Loss of affine polarization has a tagged conversion cost)**.** *Define the true cancellation score*

$$\upsilon_-(t) := \frac{a^-(t)}{a^+(t)}$$

whenever $a^+(t) > 0$, let $\upsilon_-^{\mathrm{aff}}$ be the affine score from theorem 15.5, and let $\mathcal{K}_{\mathrm{cent}}$ be the centered deformation action from definition 17.3. Put

$$\Lambda(t) := \int_0^t a(\tau)\, d\tau.$$

Fix a compact affine-age interval $J_\Lambda \Subset [0,\infty)$ and suppose that $\Lambda([0,t_*]) \subset J_\Lambda$ and

$$\upsilon_-^{\mathrm{aff}}(\Lambda) \le \upsilon^*_{\mathrm{aff}} < \upsilon_* < 1 \qquad (\Lambda \in J_\Lambda).$$

If the true score first reaches υ_* at a classical time $t_* > 0$, then at least one of the following has occurred before or at t_*:

- *(i) $\mathcal{K}_{\mathrm{cent}}([0,t_*])$ exceeds a fixed positive threshold depending only on J_Λ, the affine margin $\upsilon_* - \upsilon^*_{\mathrm{aff}}$, and the fixed datum;*
- *(ii) a positive response-weighted newly generated negative bank is a fixed positive fraction of the positive affine Hodge scale;*
- *(iii) a tangent/response/continuation compactness constant needed for the same-clock affine comparison degenerates.*

Common affine aging alone cannot produce a substantial upward conversion of the cancellation score. In fact, for the full $\mathcal{K}_{\mathrm{cent}}$ of definition 17.3, the rigorous proof gives the stronger dichotomy that either (i) or (iii) occurs; item (ii) is retained as the finer source-specific routing of the source-gradient part of (i).

Proposition 15.7 (Protected apex tube in two-time response)**.** *Let $[t_1, T)$ be a finite branch with the terminal-uniform local bounds, fix $J = [s_-, s_+] \Subset (0,\infty)$, and choose $\ell > 0$ sufficiently small so that*

$$\mathcal{U}_\ell := \{0 < x < \ell,\ y/x \in J\} \tag{15.12}$$

is contained in the uniform apex tube. Then

$$q > 0 \tag{15.13}$$

throughout $\mathcal{U}_\ell \times [t_1, T)$, so no actual negative source is deposited there. Let $\mathfrak{R}_q$ be the residence multiplier of theorem 18.4, define

$$\mathcal{K}_{vx} := \int_{t_1}^T \|v - xv_x\|_{L^\infty(\mathcal{Q})}\, dt, \qquad \mathfrak{R}_* := (T - t_1)e^{2\mathcal{K}_{vx}},$$

and define $\mathfrak{X}^H_{-,\mathrm{tube}}(t)$ *by restricting* $\mathfrak{X}^H_-(t)$ *to deposition times* $\tau \in [t_1, t]$ *and deposition points* $X \in \mathcal{U}_\ell$. *If* $\mathcal{K}_{vx} < \infty$, *then*

$$\boxed{\mathfrak{X}^H_{-,\mathrm{tube}}(t) \le \mathfrak{c}_J \ell\, \mathfrak{R}_*\, a^+(t), \qquad t_1 \le t < T.} \tag{15.14}$$

If $\mathcal{K}_{vx} = \infty$, *the branch has already exited through the named first-jet/continuation account. Thus order-one cancellation cannot be financed by fresh-apex deposition on a terminal-uniform branch.*

Theorem 15.8 (One-shot exterior conversion alternative)**.** *Let* $\mathfrak{X}^H_-$ *be the exact generated-negative response majorant from theorem 18.4, let* $\mathfrak{X}^H_{-,\mathrm{tube}}$ *be the E6 subaccount, and define*

$$\mathfrak{X}^H_{-,\mathrm{out}}(t) := \mathfrak{X}^H_-(t) - \mathfrak{X}^H_{-,\mathrm{tube}}(t).$$

Thus $\mathfrak{X}^H_{-,\mathrm{out}}$ *contains all generated-negative response deposited before* t_1 *and, for* $\tau \ge t_1$, *all such response deposited outside the protected tube. Put* $q_0(X) := q(X, 0)$ *and define the old-response account*

$$\mathfrak{X}^H_{-,\mathrm{old}}(t) := \frac{3}{4}\int_{\mathcal{Q}} [q_0(X)]_- \mathcal{V}_H(t; 0, X)\mathfrak{R}_q(t, 0; X)\, dX.$$

Suppose

$$\mathfrak{X}^H_{-,\mathrm{old}}(t) \le \upsilon_{\mathrm{old}} a^+(t), \qquad 0 \le \upsilon_{\mathrm{old}} < 1, \tag{15.15}$$

and $\mathfrak{X}^H_{-,\mathrm{tube}}(t) \le \eta a^+(t)$ *for some* $\eta \ge 0$. *If*

$$a^-(t) \ge \upsilon_* a^+(t), \qquad \upsilon_* > \upsilon_{\mathrm{old}} + \eta, \tag{15.16}$$

then

$$\boxed{\mathfrak{X}^H_{-,\mathrm{out}}(t) \ge (\upsilon_* - \upsilon_{\mathrm{old}} - \eta) a^+(t).} \tag{15.17}$$

This is a one-shot exterior/non-apex conversion ticket: each physical deposition is charged once. Compact mesoscopic localization is addressed by the separate later theorem 16.9; E7 by itself does not assert that localization.

16. Source-gradient restart system and the rotor

Accounting bridge. The next statements identify how favorable source credit can be converted into an adverse source. The H-part is a rotor rather than an amplifier, so the ledger must track orientation change, not merely source-gradient magnitude.

Proposition 16.1 (Exact source-gradient system)**.** *At* $x, y > 0$, *let*

$$s := y/x, \qquad C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y}, \qquad Z_\beta := \begin{pmatrix} q \\ C_\beta \end{pmatrix}.$$

Then

$$\boxed{\mathcal{D}_{xy} Z_\beta = \mathbf{A}_{\mathrm{sg}} Z_\beta,} \tag{16.1}$$

with

$$\boxed{\mathbf{A}_{\mathrm{sg}} = \begin{pmatrix} \frac{3}{2}\mathfrak{A} - \mathfrak{J} & -yH - 2s\mathfrak{T} \\ \frac{1}{4}xH - \frac{1}{2}\mathfrak{T} & \frac{3}{2}\mathfrak{A} + \mathfrak{J} \end{pmatrix}.} \tag{16.2}$$

Thus source-sign rotation is already expressed entirely in the established restart basis.

Definition 16.2 (Orthogonalized radial source gradient)**.** *At $x, y > 0$ define*

$$\boxed{\widehat{C}_\beta := 2\sqrt{s}\, C_\beta, \qquad \omega_H := \frac{1}{2}\sqrt{xy}\, H.} \tag{16.3}$$

Proposition 16.3 (Rotor normal form)**.** *At $x, y > 0$, put*

$$s := y/x, \qquad C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y}, \qquad \widehat{C}_\beta := 2\sqrt{s}\, C_\beta, \qquad \omega_H := \frac{1}{2}\sqrt{xy}\, H,$$

and define

$$\mathsf{d}_q := \frac{3}{2}\mathfrak{A} - \mathfrak{J}, \qquad \mathsf{d}_{\widehat{C}} := 3v + 2yg_y.$$

Then

$$\boxed{\mathcal{D}_{xy}\begin{pmatrix} q \\ \widehat{C}_\beta \end{pmatrix} = \begin{pmatrix} \mathsf{d}_q & -\omega_H - \sqrt{s}\mathfrak{T} \\ \omega_H - \sqrt{s}\mathfrak{T} & \mathsf{d}_{\widehat{C}} \end{pmatrix}\begin{pmatrix} q \\ \widehat{C}_\beta \end{pmatrix}.} \tag{16.4}$$

The entire H-part of the off-diagonal coefficient matrix is exactly skew-symmetric.

Theorem 16.4 (H-skew cancellation)**.** *At $x, y > 0$, with*

$$s := y/x, \quad C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y}, \quad \widehat{C}_\beta := 2\sqrt{s}\, C_\beta, \quad \mathsf{d}_q := \frac{3}{2}\mathfrak{A} - \mathfrak{J}, \quad \mathsf{d}_{\widehat{C}} := 3v + 2yg_y,$$

define

$$\mathcal{N}_\beta^2 := q^2 + \widehat{C}_\beta^2. \tag{16.5}$$

Then

$$\boxed{\mathcal{D}_{xy}\mathcal{N}_\beta^2 = 2\mathsf{d}_q q^2 + 2\mathsf{d}_{\widehat{C}}\widehat{C}_\beta^2 - 4\sqrt{s}\mathfrak{T} q\widehat{C}_\beta.} \tag{16.6}$$

The self-generated H field can rotate the source-gradient direction but makes no direct contribution to this orthogonalized Euclidean amplitude.

Proposition 16.5 (Passive-circulation interpretation)**.** *At $x, y > 0$, let*

$$\Gamma := xu, \qquad s := y/x, \qquad C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y}, \qquad \widehat{C}_\beta := 2\sqrt{s}\, C_\beta.$$

Because $\Theta = \varepsilon_\beta \Gamma^2/x^2$,

$$\boxed{q = \frac{2\varepsilon_\beta \Gamma}{x^2}\Gamma_y, \qquad \widehat{C}_\beta = \frac{2\varepsilon_\beta \Gamma}{x^2\sqrt{s}}\Gamma_x.} \tag{16.7}$$

Let $\mathscr{X}_{t,0}$ be the material flow of $\mathbf{b} = (-2xv, 2yg)$. On every classical interval on which this flow is defined,

$$\boxed{\nabla\Gamma(\mathscr{X}_{t,0}X_0, t) = D\mathscr{X}_{t,0}(X_0)^{-T}\nabla\Gamma_0(X_0).} \tag{16.8}$$

Consequently an initial critical point of Γ remains critical along its trajectory, while a noncritical initial circulation gradient remains nonzero. If $\Gamma_0(X_0) = 0$, then the tagged source-gradient pair in (16.7) vanishes along that trajectory; if $\Gamma_0(X_0) \neq 0$, its sign is fixed by passive transport and any later change in the sign of q is a change in the transverse orientation of the same transported initial circulation gradient. Thus new negative transverse source is not a renewable independent forcing direction.

Proposition 16.6 (Exact apex mixed-shear Hodge moment)**.** *Put $s := y/x$ and define*

$$\tau_0(t) := g_x(0,0,t) - v_y(0,0,t). \tag{16.9}$$

Then the seven-dimensional Newton representation gives

$$\boxed{\tau_0(t) = \frac{15}{8}\int_{\mathcal{Q}} \frac{x\sqrt{y}(3x-4y)}{(x+y)^{9/2}} H(x,y,t)dX.} \tag{16.10}$$

Equivalently,

$$\boxed{\tau_0(t) = \frac{15}{8}\int_{\mathcal{Q}} K_H(x,y)\frac{3-4s}{x(1+s)^2} H(x,y,t)dX.} \tag{16.11}$$

Thus the apex mixed shear is not an independent local coefficient; it is a signed whole-space Hodge moment of the same distributed genealogy.

Proposition 16.7 (Hodge-aware adverse exchange)**.** *Let $s := y/x$ and*

$$C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y}.$$

Using

$$4yv_y = -yH - 2s\mathfrak{T}, \tag{16.12}$$

the adverse off-diagonal exchange is

$$\boxed{[-(4yv_y)C_\beta]_+ = [(yH + 2s\mathfrak{T})C_\beta]_+.} \tag{16.13}$$

Its signed, not positive-part, form has the radial divergence identity

$$\boxed{(yH + 2s\mathfrak{T})C_\beta = \left(\frac{H}{2x} + \frac{\mathfrak{T}}{x^2}\right)\partial_x(x^2\Theta).} \tag{16.14}$$

The signed identity is useful routing information, but it does not control total positive variation and therefore is not by itself a closure theorem.

Proposition 16.8 (Angular-tail Hodge suppression of the rotor)**.** *Put*

$$s := y/x, \qquad C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y}, \qquad \mathcal{T}_{\mathrm{mix}} := g_x - v_y = \mathfrak{T}/x,$$

and set

$$e_- := [-(4yv_y)C_\beta]_+.$$

Then

$$\boxed{K_H e_- \le \psi_{\mathrm{rot}}(s)(|H| + 2|\mathcal{T}_{\mathrm{mix}}|)|C_\beta|,} \tag{16.15}$$

where

$$\boxed{\psi_{\mathrm{rot}}(s) = \frac{s^{3/2}}{(1+s)^{5/2}}.} \tag{16.16}$$

Hence

$$\psi_{\mathrm{rot}}(s) = O(s^{3/2}) \quad (s \downarrow 0), \qquad \psi_{\mathrm{rot}}(s) = O(s^{-1}) \quad (s \to \infty). \tag{16.17}$$

More quantitatively, define

$$w_{\mathrm{rot}}(s) := \frac{s^{4/3}}{(1+s)^{5/2}}, \tag{16.18}$$

and for $0 < \delta < 1 < S < \infty$ put

$$\mathcal{A}_{\delta,S} := \{s < \delta\} \cup \{s > S\}, \qquad \omega_{\delta,S} := \sup_{s \in \mathcal{A}_{\delta,S}} w_{\mathrm{rot}}(s). \tag{16.19}$$

Then $\omega_{\delta,S} \to 0$ as $\delta \downarrow 0$ and $S \uparrow \infty$. Moreover, for a fixed numerical constant $\mathfrak{c}_{\mathrm{tail}}$,

$$\boxed{\int_{\mathcal{A}_{\delta,S}} K_H e_- \, dX \le \mathfrak{c}_{\mathrm{tail}}\, \omega_{\delta,S} \left[\mathcal{E}_{H,1}\mathcal{E}^{(0,0)}_{\mathrm{sg}} + \mathcal{L}_B \mathcal{Q}^{(1)}_{\mathrm{sg}}\right].} \tag{16.20}$$

Thus both extreme angular tails are Hodge-suppressed at the pointwise and carrier levels before any collapse-age normalization.

Theorem 16.9 (Compact mesoscopic rotation ticket)**.** *Put*

$$s := y/x, \qquad C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y}, \qquad \widehat{C}_\beta := 2\sqrt{s}\, C_\beta, \qquad \omega_H := \frac{1}{2}\sqrt{xy}\, H, \qquad \upsilon_- := \frac{a^-}{a^+}.$$

Let $t_* > 0$ be a classical time on a finite continuation branch and suppose the first-jet/visibility and rapid-decay continuation bounds entering the audited D3–D6 package are finite on $[0, t_*]$. Assume $a^+(t_*) > 0$ and that the certified old-response account satisfies

$$\mathfrak{X}^H_{-,\mathrm{old}}(t_*) \le \upsilon_{\mathrm{old}} a^+(t_*), \qquad 0 \le \upsilon_{\mathrm{old}} < \upsilon_* < 1. \tag{16.21}$$

If t_* is the first time at which

$$a^-(t_*) \ge \upsilon_* a^+(t_*),$$

then, unless an already named old-response/tangent/derivative/rapid-decay exit has occurred, there exist constants

$$0 < \ell < x_{\mathrm{far}} < \infty, \qquad 0 < \delta < S < \infty$$

and the compact mesoscopic box

$$\mathcal{M} := \{\ell \le x \le x_{\mathrm{far}},\ \delta \le s \le S\} \tag{16.22}$$

such that the response-weighted generated-negative deposit on $\mathcal{M}$ is a strictly positive fraction of $a^+(t_*)$, and in particular

$$\boxed{\int_0^{t_*} \int_{\mathcal{M}} [(\omega_H + \sqrt{s}\mathfrak{T})\widehat{C}_\beta]_+ \, dX\, dt \ge c_{\mathrm{rot}} > 0.} \tag{16.23}$$

The constants may depend on the admitted finite-action bounds, the old-response margin, and the positive number $a^+(t_*)$, but not on any later Hodge refresh. This is a one-shot physical conversion ticket and is never charged again merely because later Hodge refreshes occur.

17. Level-set polarization and deformation stability

Accounting bridge. The transported circulation level sets provide the global balance sheet that the local rotor lacks. The affine polarization is a certified profit margin, and the stability theorem prices the deformation required to destroy it.

Theorem 17.1 (Circulation level-set response)**.** *Fix an observation time t and a source time $0 \le \tau \le t$. For the fixed customer put $\Gamma := xu$. On the physical branch $u_0 = yw_0 \ge 0$, and the exact transport law $\mathcal{D}_{xy}\Gamma = 0$ gives $\Gamma(\cdot, \tau) \ge 0$. Define*

$$\Omega_c(\tau) := \{X \in \mathcal{Q} : \Gamma(X, \tau) > c\}, \qquad W_{t,\tau}(X) := K_H(\mathscr{X}_{t,\tau} X), \tag{17.1}$$

$$\mathcal{S}_\pm(t, \tau) := \frac{3}{4} \int_{\mathcal{Q}} W_{t,\tau}(X)[q(X, \tau)]_\pm \, dX, \qquad \mathcal{R}(t, \tau; X) := -\partial_{y_\tau} W_{t,\tau}(X). \tag{17.2}$$

For every regular value $c > 0$, let $n_c := \nabla\Gamma/|\nabla\Gamma|$ point toward increasing circulation on $\partial\Omega_c(\tau)$. Then

$$\boxed{\mathcal{S}_+(t,\tau) = \frac{3\varepsilon_\beta}{2}\int_0^\infty c \oint_{\partial\Omega_c(\tau)} \frac{W_{t,\tau}}{x^2}[n_{c,y}]_+ \, d\ell \, dc,} \tag{17.3}$$

$$\boxed{\mathcal{S}_-(t,\tau) = \frac{3\varepsilon_\beta}{2}\int_0^\infty c \oint_{\partial\Omega_c(\tau)} \frac{W_{t,\tau}}{x^2}[-n_{c,y}]_+ \, d\ell \, dc.} \tag{17.4}$$

Their difference is

$$\boxed{\mathcal{S}_+(t,\tau) - \mathcal{S}_-(t,\tau) = \frac{3\varepsilon_\beta}{2}\int_0^\infty c \int_{\Omega_c(\tau)} \frac{\mathcal{R}(t,\tau;X)}{x^2} \, dX \, dc.} \tag{17.5}$$

All three formulas are understood in the coarea sense: critical circulation levels form a null set and require no separate boundary parametrization.

Proposition 17.2 (Raw level-set compensation criterion)**.** *Fix a time t and let $0 < \pi_* < 1$. A sufficient raw form of the strict cancellation gap is*

$$\boxed{a(t) \geq \pi_* \, [a^+(t) + a^-(t)].} \tag{17.6}$$

Equivalently,

$$\boxed{a^-(t) \leq \frac{1-\pi_*}{1+\pi_*} a^+(t).} \tag{17.7}$$

In the level-set representation the common factor $3\varepsilon_\beta/2$ remains on both sides until physical admission; it is not cancelled at the provenance gate.

Definition 17.3 (Centered deformation action)**.** *Set*

$$\boxed{\mathcal{K}_{\mathrm{cent}}([0,t]) := \int_0^t \big(\|v-a\|_\infty + \|g-2a\|_\infty + \|\mathbf{A}_{\mathrm{sg}}\|_\infty\big) ds.} \tag{17.8}$$

Lemma 17.4 (Uniform fixed-seed source-gradient condition number)**.** *Work in the physical sector $\varepsilon_\beta = 1$. For $\Lambda \geq 0$ define the affine map*

$$\mathscr{X}_\Lambda^{\mathrm{aff}}(x,y) := (e^{-2\Lambda}x, e^{4\Lambda}y),$$

let

$$Z_{\beta,0} := (q_0, C_{\beta,0})^T, \qquad |Z_{\beta,0}| := \sqrt{q_0^2 + C_{\beta,0}^2},$$

be the initial source-gradient pair of the fixed customer, and define

$$\mathfrak{C}_Z(\Lambda) := \frac{\int_{\mathcal{Q}} K_H(\mathscr{X}_\Lambda^{\mathrm{aff}} X)|Z_{\beta,0}(X)| \, dX}{\int_{\mathcal{Q}} K_H(\mathscr{X}_\Lambda^{\mathrm{aff}} X)|q_0(X)| \, dX}.$$

Then the denominator is positive and

$$\boxed{\mathfrak{C}_Z^* := \sup_{\Lambda \geq 0} \mathfrak{C}_Z(\Lambda) < \infty.} \tag{17.9}$$

Theorem 17.5 (Stability of the complete Hodge-response polarization)**.** *Work in the physical sector $\varepsilon_\beta = 1$. Let*

$$M := \mathcal{K}_{\mathrm{cent}}([0,t])$$

with $\mathcal{K}_{\mathrm{cent}}$ from definition 17.3, and put

$$\Lambda(t) := \int_0^t a(\tau)\,d\tau, \qquad I_t := \int_0^t e^{2\Lambda(\tau)}\,d\tau,$$

$$W_t^{\mathrm{aff}}(X) := K_H(\mathscr{X}^{\mathrm{aff}}_{\Lambda(t)}X),$$

where $\mathscr{X}^{\mathrm{aff}}_\Lambda$ is the affine map from lemma 17.4. Put $q_0(X) := q(X,0)$. Define

$$\mathcal{P}(t) := a(t), \qquad \mathcal{V}(t) := a^+(t) + a^-(t),$$

$$\mathcal{P}_{\mathrm{aff}}(t) := \frac{3}{4} I_t \int_{\mathcal{Q}} q_0(X) W_t^{\mathrm{aff}}(X)\,dX,$$

$$\mathcal{V}_{\mathrm{aff}}(t) := \frac{3}{4} I_t \int_{\mathcal{Q}} |q_0(X)| W_t^{\mathrm{aff}}(X)\,dX.$$

Then for universal constants $\mathfrak{c}_0, \mathfrak{c}_1 > 0$,

$$\boxed{|\mathcal{P} - \mathcal{P}_{\mathrm{aff}}| + |\mathcal{V} - \mathcal{V}_{\mathrm{aff}}| \le \mathfrak{c}_1 \mathfrak{C}_Z^* (e^{\mathfrak{c}_0 M} - 1)\mathcal{V}_{\mathrm{aff}}.} \tag{17.10}$$

Consequently there exist fixed constants $M_* > 0$ and $\pi_{107} > 0$, depending only on the fixed customer and the affine polarization margin, such that

$$\mathcal{K}_{\mathrm{cent}}([0,t]) \le M_* \tag{17.11}$$

implies

$$\boxed{a(t) \ge \pi_{107}[a^+(t) + a^-(t)],} \tag{17.12}$$

and hence a uniform strict cancellation gap.

Corollary 17.6 (First polarization loss pays a definite deformation bill)**.** *Let*

$$\mathfrak{F}_{\mathrm{pol}} := \{t > 0 : \ a(t) < \pi_{107}[a^+(t) + a^-(t)]\}.$$

If $\mathfrak{F}_{\mathrm{pol}} \neq \varnothing$, define the first loss time by

$$t_* := \inf \mathfrak{F}_{\mathrm{pol}}.$$

Then

$$\boxed{\mathcal{K}_{\mathrm{cent}}([0,t_*]) \ge M_* > 0.} \tag{17.13}$$

For $X = (x,y)$, $s := y/x$, and $y_\lambda := \lambda y$, define

$$\mathcal{L}_{\mathrm{pol}}(t) := \|\mathbf{A}_{\mathrm{sg}}(t)\|_\infty + \|\mathfrak{A}(t)\|_\infty + \sup_{X\in\mathcal{Q}} \int_0^1 \left(\frac{1}{4}|\mathfrak{A}| + \frac{1}{2}|\mathfrak{J}| + \frac{1}{4} y_\lambda |H| + \frac{1}{2} s |\mathfrak{T}|\right)(\lambda X, t)\,\frac{d\lambda}{\lambda}.$$

Here $\mathbf{A}_{\mathrm{sg}}$ is the source-gradient matrix from proposition 16.1. Then there exists $c_{\mathrm{pol}} > 0$, depending only on the fixed customer and affine margin, such that

$$\boxed{\int_0^{t_*} \mathcal{L}_{\mathrm{pol}}(\tau)\,d\tau \ge c_{\mathrm{pol}}.} \tag{17.14}$$

18. Fresh conversion, two-time response, and causal recycling

Accounting bridge. Once adverse genealogy is created, the ledger must distinguish new spending from recycling of an old liability. The causal response formulas do exactly that and prevent Peter from being charged twice for the same physical conversion.

Proposition 18.1 (Final/deposition Hodge visibility comparison)**.** *Put $s := y/x$, and let $\mathcal{V}_H(t;\tau,X)$ be the two-time visibility from definition 15.1. The exact material visibility law is*

$$\mathcal{D}_{xy} \log K_H = \frac{(3-2s)v + (1-4s)g}{1+s}. \tag{18.1}$$

For a source time $0 \le \tau \le t$, define

$$\mathcal{K}_{\mathrm{vis}}(\tau,t) := \int_\tau^t (\|v(\cdot,\xi)\|_\infty + \|g(\cdot,\xi)\|_\infty)\, d\xi. \tag{18.2}$$

If $\mathcal{K}_{\mathrm{vis}}(\tau,t) < \infty$, then

$$\boxed{e^{-4\mathcal{K}_{\mathrm{vis}}(\tau,t)} K_H(X) \le \mathcal{V}_H(t;\tau,X) \le e^{4\mathcal{K}_{\mathrm{vis}}(\tau,t)} K_H(X).} \tag{18.3}$$

Thus final visibility can be compared to deposition visibility without dividing by the net compression a.

Proposition 18.2 (Finite raw response capacity)**.** *Let $\mathfrak{X}_-^H$ denote the generated-negative response majorant defined in theorem 18.4. With $d\mu_c = dX/\mathfrak{r}$, define*

$$\mathcal{E}_{H,1} := \|\sqrt{xy}\, H\|_{L^2(d\mu_c)}, \qquad \mathcal{E}_{\mathrm{sg}}^{(0,0)} := \|(q, C_\beta)\|_{L^2(d\mu_c;\ell^2)},$$

$$\mathcal{Q}_{\mathrm{sg}}^{(1)} := \|q\|_{L^1(d\mu_c)} + \|C_\beta\|_{L^1(d\mu_c)}.$$

Let $\mathbf{B}(\mathbf{X},\mathbf{Z},t) = (-v\mathbf{X}, g\mathbf{Z})$ and $\mathcal{L}_B := \|\nabla_{\mathbb{R}^7}\mathbf{B}\|_\infty$. For a classical observation time $t > 0$, define

$$\mathcal{K}_{vx}(0,t) := \int_0^t \|v - x v_x\|_{L^\infty(\mathcal{Q})}\, d\xi \tag{18.4}$$

and let $\mathcal{K}_{\mathrm{vis}}(0,t)$ be the visibility action from proposition 18.1. If both actions are finite, then

$$\boxed{\mathfrak{X}_-^H(t) \le \mathfrak{c}_{\mathrm{resp}}(t) \int_0^t \left[\mathcal{E}_{H,1}(\tau)\mathcal{E}_{\mathrm{sg}}^{(0,0)}(\tau) + \mathcal{L}_B(\tau)\mathcal{Q}_{\mathrm{sg}}^{(1)}(\tau)\right] d\tau,} \tag{18.5}$$

where one may take

$$\boxed{\mathfrak{c}_{\mathrm{resp}}(t) = \mathfrak{c}_{\mathrm{rot}}\, t\, \exp\big(4\mathcal{K}_{\mathrm{vis}}(0,t) + 2\mathcal{K}_{vx}(0,t)\big)} \tag{18.6}$$

for a fixed geometric constant $\mathfrak{c}_{\mathrm{rot}} > 0$. Thus response-multiplier growth is explicitly routed to the named visibility/first-jet actions rather than becoming an anonymous mechanism. Finiteness, however, is not a strict cancellation gap.

Proposition 18.3 (Fresh-apex anti-rotation)**.** *Work in the physical sector $\varepsilon_\beta = 1$. Fix a compact classical continuation interval $I = [t_1,t_2]$ with $0 < t_1 < t_2$ and a compact angular interval $J \Subset (0,\infty)$. Put*

$$s := y/x, \qquad q_H(t) := \partial_y H(0,0,t).$$

For sufficiently small $\ell > 0$, on

$$\mathcal{U}_{\ell,J} := \{0 < x < \ell,\ s \in J\}$$

one has

$$q>0,\qquad H^{+}(x,sx,t)=q_H(t)sx+O_{I,J}(x^2),\qquad H^{-}(x,sx,t)=O_{I,J}(x^2), \tag{18.7}$$

and

$$\boxed{[-(4yv_y)C_\beta]_+\le \mathfrak{c}_{I,J}\ell\, q.} \tag{18.8}$$

Thus freshly generated positive source cannot rotate negative at order-one relative rate while the local classical first jets remain bounded.

Theorem 18.4 (Exact generated-negative Hodge response)**.** *Define*

$$C_\beta:=\frac{\Theta+\frac{x}{2}\Theta_x}{y},\qquad \mathsf{d}_q:=\frac{3}{2}\mathfrak{A}-\mathfrak{J},\qquad e_-:=[-(4yv_y)C_\beta]_+.$$

For $0\le\tau\le t$ define the positive residence multiplier

$$\boxed{\mathfrak{R}_q(t,\tau;X):=\int_\tau^t \exp\!\left(2\int_\tau^\rho (v-xv_x)(\mathscr{X}_{\xi,\tau}X,\xi)\,d\xi\right)d\rho.} \tag{18.9}$$

Define the generated-negative response majorant

$$\boxed{\mathfrak{X}_-^H(t):=\frac{3}{4}\int_0^t\int_{\mathcal{Q}} e_-(X,\tau)\mathcal{V}_H(t;\tau,X)\mathfrak{R}_q(t,\tau;X)\,dX\,d\tau,} \tag{18.10}$$

and the old initial-slope response

$$\boxed{\mathfrak{X}_{-,\mathrm{old}}^H(t):=\frac{3}{4}\int_{\mathcal{Q}}[q(X,0)]_-\mathcal{V}_H(t;0,X)\mathfrak{R}_q(t,0;X)\,dX.} \tag{18.11}$$

Then

$$\boxed{a^-(t)\le \mathfrak{X}_{-,\mathrm{old}}^H(t)+\mathfrak{X}_-^H(t).} \tag{18.12}$$

In particular the Hodge bank created by later adverse cross-shear is bounded by $\mathfrak{X}_-^H(t)$. No $1/a$, $1/\Lambda$, or common-tag quotient is used.

Proposition 18.5 (Fresh primary conversion is quadratically small)**.** *Fix a final observation time $t>0$ and let $\mathcal{G}_{\mathrm{rec}}$ be the causal Green function from theorem 18.6. Define the global primary conversion amplitude*

$$\boxed{\mathfrak{m}_{\mathrm{prim}}^+(\tau):=\big\|[2\Theta+x\Theta_x]_+(\cdot,\tau)\big\|_\infty.} \tag{18.13}$$

At $x,y>0$ this is equivalently $\|\sqrt{xy}[\widehat{C}_\beta]_+\|_\infty$, and the first expression is its smooth axis extension. Define

$$A^+_{t\leftarrow\tau}:=\frac{3}{8}\int_{\mathcal{Q}}H^+(X,\tau)\mathcal{V}_H(t;\tau,X)\,dX.$$

For $0<h\le h_0\le t$ put

$$I_h:=[t-h,t],\qquad \mathfrak{m}^{+,*}_{\mathrm{prim,h}}:=\sup_{\tau\in I_h}\mathfrak{m}^+_{\mathrm{prim}}(\tau),$$

and define the fresh primary account

$$\mathfrak{X}^{\mathrm{res}}_{+,\mathrm{fresh,h}}(t):=\int_{I_h}\mathcal{G}_{\mathrm{rec}}(t-\tau)\mathfrak{m}^+_{\mathrm{prim}}(\tau)A^+_{t\leftarrow\tau}\,d\tau.$$

For the fixed h_0 set

$$\mathfrak{c}_{\mathrm{rec}} := \sup_{0<u\le h_0} \frac{\mathcal{G}_{\mathrm{rec}}(u)}{u} < \infty.$$

Then

$$\boxed{\mathfrak{X}^{\mathrm{res}}_{+,\mathrm{fresh,h}}(t) \le \frac{1}{2}\mathfrak{c}_{\mathrm{rec}}\mathfrak{m}^{+,*}_{\mathrm{prim,h}} h^2 a^+(t).} \tag{18.14}$$

Hence, on every admitted finite classical interval, an arbitrarily short primary $H^+ \to H^-$ episode cannot create an order-one new negative bank.

Theorem 18.6 (Causal Hodge resolvent for negative self-recycling)**.** *Fix a final observation time t and let $\mathcal{V}_H(t;\tau,X) = K_H(\mathscr{X}_{t,\tau}X)$. Put*

$$s := y/x, \qquad C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y}, \qquad \widehat{C}_\beta := 2\sqrt{s}\, C_\beta, \qquad \mathtt{d}_q := \frac{3}{2}\mathfrak{A} - \mathfrak{J}.$$

Define the globally extended nonnegative self-recycling coefficient

$$\boxed{\mathfrak{m}_-(X,\tau) := [-(2\Theta + x\Theta_x)]_+(X,\tau).} \tag{18.15}$$

For $x, y > 0$, $\mathfrak{m}_- = \sqrt{xy}[-\widehat{C}_\beta]_+$. Define

$$e_{\mathfrak{T}} := [\sqrt{s}\,\mathfrak{T}\,\widehat{C}_\beta]_+, \qquad e_{H,+} := \frac{1}{2}\sqrt{xy}\, H^+[\widehat{C}_\beta]_+, \qquad f_{\mathrm{prim}} := e_{\mathfrak{T}} + e_{H,+}.$$

Let $q^{\mathrm{maj}}_- \ge 0$ and $H^{\mathrm{maj}}_- \ge 0$ be the characteristic solutions of

$$\mathcal{D}_{xy} q^{\mathrm{maj}}_- = \mathtt{d}_q q^{\mathrm{maj}}_- + f_{\mathrm{prim}} + \frac{1}{2}\mathfrak{m}_- H^{\mathrm{maj}}_-, \qquad q^{\mathrm{maj}}_-(\cdot,0) = [q_0]_-, \tag{18.16}$$

$$\mathcal{D}_{xy} H^{\mathrm{maj}}_- + g H^{\mathrm{maj}}_- = 2q^{\mathrm{maj}}_-, \qquad H^{\mathrm{maj}}_-(\cdot,0) = 0. \tag{18.17}$$

Then

$$\boxed{[q]_- \le q^{\mathrm{maj}}_-, \qquad H^- \le H^{\mathrm{maj}}_-.} \tag{18.18}$$

Define the weighted causal stocks

$$\mathcal{Z}_{\mathrm{rec}}(\tau) := \int_{\mathcal{Q}} q^{\mathrm{maj}}_-(X,\tau)\mathcal{V}_H(t;\tau,X)\, dX,$$

$$\mathcal{H}_{\mathrm{rec}}(\tau) := \int_{\mathcal{Q}} H^{\mathrm{maj}}_-(X,\tau)\mathcal{V}_H(t;\tau,X)\, dX,$$

$$\mathcal{I}_{\mathrm{rec}}(\tau) := \int_{\mathcal{Q}} f_{\mathrm{prim}}(X,\tau)\mathcal{V}_H(t;\tau,X)\, dX.$$

Put

$$\mathcal{A}_{\mathrm{rec}}(\tau) := 2\|(v - xv_x)(\cdot,\tau)_+\|_\infty, \qquad \mathcal{B}_{\mathrm{rec}}(\tau) := \|\mathfrak{m}_-(\cdot,\tau)\|_\infty,$$

$$\mathcal{A}^*_{\mathrm{rec}} := \sup_{0\le\tau\le t} \mathcal{A}_{\mathrm{rec}}(\tau), \qquad \mathcal{B}^*_{\mathrm{rec}} := \sup_{0\le\tau\le t} \mathcal{B}_{\mathrm{rec}}(\tau).$$

On every finite classical interval these constants are finite. Define

$$\kappa_* := \sqrt{(\mathcal{A}^*_{\mathrm{rec}})^2 + 4\mathcal{B}^*_{\mathrm{rec}}}, \qquad \lambda_\pm := \frac{\mathcal{A}^*_{\mathrm{rec}} \pm \kappa_*}{2}, \tag{18.19}$$

and the positive Green function

$$\boxed{\mathcal{G}_{\mathrm{rec}}(u)=\begin{cases}\dfrac{e^{\lambda_+ u}-e^{\lambda_- u}}{\kappa_*}, & \kappa_*>0,\\ u, & \kappa_*=0.\end{cases}} \tag{18.20}$$

Then

$$\boxed{a^-(t)\le \frac{3}{4}\mathcal{Z}_{\mathrm{rec}}(0)\mathcal{G}_{\mathrm{rec}}(t)+\frac{3}{4}\int_0^t \mathcal{G}_{\mathrm{rec}}(t-\tau)\mathcal{I}_{\mathrm{rec}}(\tau)\,d\tau.} \tag{18.21}$$

Thus the full $H^-\to H^-$ branch, including exterior source points, is a finite causal amplifier on every admitted finite classical interval; no operator-norm condition <1 and no mesoscopic undercount are used.

Proposition 18.7 (Three-account negative-bank bound, with exact primary account)**.** *Let $\mathcal{G}_{\mathrm{rec}}$ and $\mathcal{Z}_{\mathrm{rec}}$ be as in theorem 18.6. Put*

$$s:=y/x,\qquad C_\beta:=\frac{\Theta+\frac{x}{2}\Theta_x}{y},\qquad \widehat{C}_\beta:=2\sqrt{s}\,C_\beta,$$

and define

$$e_{\mathfrak{T}}:=[\sqrt{s}\,\mathfrak{T}\,\widehat{C}_\beta]_+,\qquad \mathcal{I}_{\mathfrak{T},t}(\tau):=\int_{\mathcal{Q}} e_{\mathfrak{T}}(X,\tau)\mathcal{V}_H(t;\tau,X)\,dX,$$

$$\boxed{\mathcal{I}_{+,t}(\tau):=\frac{1}{2}\int_{\mathcal{Q}}\sqrt{xy}\,H^+(X,\tau)[\widehat{C}_\beta(X,\tau)]_+\mathcal{V}_H(t;\tau,X)\,dX.} \tag{18.22}$$

Define the exact three upstream accounts

$$\mathfrak{X}^{\mathrm{res}}_{-,\mathrm{old}}(t):=\frac{3}{4}\mathcal{Z}_{\mathrm{rec}}(0)\mathcal{G}_{\mathrm{rec}}(t),$$

$$\mathfrak{X}^{\mathrm{res}}_{\mathfrak{T}}(t):=\frac{3}{4}\int_0^t \mathcal{G}_{\mathrm{rec}}(t-\tau)\mathcal{I}_{\mathfrak{T},t}(\tau)\,d\tau,$$

$$\boxed{\mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{prim}}(t):=\frac{3}{4}\int_0^t \mathcal{G}_{\mathrm{rec}}(t-\tau)\mathcal{I}_{+,t}(\tau)\,d\tau.} \tag{18.23}$$

For comparison only, set

$$\mathfrak{m}^+_{\mathrm{prim}}(\tau):=\|[2\Theta+x\Theta_x]_+(\cdot,\tau)\|_\infty,\qquad A^+_{t\leftarrow\tau}:=\frac{3}{8}\int_{\mathcal{Q}}H^+(X,\tau)\mathcal{V}_H(t;\tau,X)\,dX,$$

and define the factorized majorant

$$\mathfrak{X}^{\mathrm{fac}}_{+,\mathrm{prim}}(t):=\int_0^t \mathcal{G}_{\mathrm{rec}}(t-\tau)\mathfrak{m}^+_{\mathrm{prim}}(\tau)A^+_{t\leftarrow\tau}\,d\tau. \tag{18.24}$$

Then

$$\boxed{a^-(t)\le \mathfrak{X}^{\mathrm{res}}_{-,\mathrm{old}}(t)+\mathfrak{X}^{\mathrm{res}}_{\mathfrak{T}}(t)+\mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{prim}}(t)\le \mathfrak{X}^{\mathrm{res}}_{-,\mathrm{old}}(t)+\mathfrak{X}^{\mathrm{res}}_{\mathfrak{T}}(t)+\mathfrak{X}^{\mathrm{fac}}_{+,\mathrm{prim}}(t).} \tag{18.25}$$

The first inequality is the authoritative causal account. The factorized quantity is only a crude majorant; the falsification theorem shows that it is too strong to be the global old-age closure target.

Conditional Proposition 18.8 (Resummed strict-gap criterion)**.** *Fix an observation time t. Let $\mathfrak{X}^{\mathrm{res}}_{-,\mathrm{old}}$, $\mathfrak{X}^{\mathrm{res}}_{\mathfrak{T}}$, and $\mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{prim}}$ be the three accounts defined in proposition 18.7. If for some $\delta \in (0,1)$,*

$$\boxed{\mathfrak{X}^{\mathrm{res}}_{-,\mathrm{old}} + \mathfrak{X}^{\mathrm{res}}_{\mathfrak{T}} + \mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{prim}} \le (1-\delta)a^{+},} \tag{18.26}$$

then

$$\boxed{a^{-} \le (1-\delta)a^{+}.} \tag{18.27}$$

19. Auxiliary exact formulas retained for future calculations

Accounting bridge. The following formulas are the short-form receipts most likely to be needed when the remaining invoice is audited. They are kept because they transfer directly between source, visibility, and restart accounts.

Proposition 19.1 (Hodge visibility law)**.** *Put $s := y/x$. Along a material trajectory,*

$$\boxed{\mathcal{D}_{xy} \log K_H = \frac{(3-2s)v + (1-4s)g}{1+s}.} \tag{19.1}$$

Equivalently, with

$$\vartheta := \frac{y}{x+y} = \frac{s}{1+s},$$
$$\mathcal{D}_{xy} \log K_H = v(3-5\vartheta) + g(1-5\vartheta). \tag{19.2}$$

On a compressive affine leg $(v,g) = (a, 2a)$ with $a > 0$, the aspect ratio increases strictly; the affine angular visibility profile has its unique maximum at $s = 1/2$, so any parcel that traverses this scale reaches its material visibility maximum there.

Proposition 19.2 (Exact guarded source-payment identity)**.** *Let $\mathcal{R}(t,\tau;X)$ be the vertical response from proposition 15.2 and put*

$$\mathcal{R}_0(X) := \mathcal{R}(t,t;X).$$

Define

$$\mathcal{T}^{\mathrm{src}}_{\beta}(t) := \frac{3}{4}\int_{\mathcal{Q}} \Theta(X,t)[\mathcal{R}_0(X)]_+ \, dX + \frac{3}{4}\int_0^t \int_{\mathcal{Q}} \Theta(X,\tau)[\partial_t \mathcal{R}(t,\tau;X)]_+ \, dX \, d\tau, \tag{19.3}$$

$$\mathcal{W}^{\mathrm{src}}_{\beta}(t) := \frac{3}{4}\int_{\mathcal{Q}} \Theta(X,t)[\mathcal{R}_0(X)]_- \, dX + \frac{3}{4}\int_0^t \int_{\mathcal{Q}} \Theta(X,\tau)[\partial_t \mathcal{R}(t,\tau;X)]_- \, dX \, d\tau. \tag{19.4}$$

Then

$$\boxed{\dot{a}(t) = \mathcal{T}^{\mathrm{src}}_{\beta}(t) - \mathcal{W}^{\mathrm{src}}_{\beta}(t).} \tag{19.5}$$

Both accounts display the tagged factor Θ before normalization. Thus the source-response cashier passes the provenance guard exactly.

Proposition 19.3 (Raw cross-shear factorization)**.** *With*

$$\Gamma := xu, \qquad C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y},$$

the off-diagonal term in the q equation is exactly

$$\boxed{4\varepsilon_\beta u v_y \Gamma_x = (4y v_y) C_\beta.} \tag{19.6}$$

For the fixed customer of definition 5.1, passive transport of Γ *gives* $\Gamma \geq 0$ *and hence* $\varepsilon_\beta u \geq 0$ *in the open quadrant. Therefore*

$$\boxed{[-(4yv_y)C_\beta]_+ = 4\varepsilon_\beta u[-v_y\Gamma_x]_+.} \tag{19.7}$$

The adverse source is therefore a genuine descendant of the single tagged source and an already named meridional first-jet coefficient.

Proposition 19.4 (Carrier-Hodge comparison)**.** *For*

$$\mathfrak{r} = x^{2/3}y^{1/3}, \tag{19.8}$$

one has

$$\boxed{K_H(x,y) \leq \frac{2^{5/3}}{3^{5/2}}\frac{1}{\mathfrak{r}}.} \tag{19.9}$$

Therefore every nonnegative genealogy density $\mathcal{H}$ *with finite carrier integral satisfies*

$$\boxed{\frac{3}{8}\int K_H\mathcal{H}\,dX \leq \mathfrak{c}_H\int \mathcal{H}\frac{dX}{\mathfrak{r}}, \qquad \mathfrak{c}_H := \frac{2^{5/3}}{8\,3^{3/2}}.} \tag{19.10}$$

This comparison proves finite Hodge capacity from carrier L^1 *capacity but, by itself, does not yield a strict cancellation fraction.*

Proposition 19.5 (Fresh genealogy as a true Euler source-time truncation)**.** *Fix a current point* $X^\sharp \in \mathcal{Q}$*, a time* $t > 0$*, and a source time* $\tau \in [0,t]$*. Put*

$$X(\tau;X^\sharp,t) := \mathscr{X}_{\tau,t}(X^\sharp), \qquad J(t,\tau;X^\sharp) := \det D\mathscr{X}_{t,\tau}(X(\tau;X^\sharp,t)).$$

For $h > 0$ *set* $\tau_h := (t-h)_+$ *and define*

$$H^\pm_{\mathrm{fr},h}(X^\sharp,t) := 2\int_{\tau_h}^t J(t,\tau;X^\sharp)^{-1}[q(X(\tau;X^\sharp,t),\tau)]_\pm\,d\tau. \tag{19.11}$$

Then

$$H^\pm_{\mathrm{fr},h} \geq 0, \qquad H_{\mathrm{fr},h} = H^+_{\mathrm{fr},h} - H^-_{\mathrm{fr},h}. \tag{19.12}$$

Moreover, if $t \leq h$ *then* $H^\pm_{\mathrm{fr},h} = H^\pm$*, whereas if* $t > h$ *then*

$$\boxed{H^\pm_{\mathrm{fr},h}(X^\sharp,t) = H^\pm(X^\sharp,t) - J(t,t-h;X^\sharp)^{-1}H^\pm(X(t-h;X^\sharp,t),t-h).} \tag{19.13}$$

Thus $H^\pm = H^\pm_{\mathrm{fr},h} + H^\pm_{\mathrm{old},h}$ *with a nonnegative complementary old cohort* $H^\pm_{\mathrm{old},h}$*. With* $\mathcal{C}$ *from proposition 10.2, the corresponding fresh meridional payment is exactly*

$$\boxed{\mathcal{M}_{\beta,\mathrm{fr},h}(t) = \frac{3}{8}\int_{\mathcal{Q}} K_H\mathcal{C}H_{\mathrm{fr},h}\,dX.} \tag{19.14}$$

Thus the fresh response is not an auxiliary memory variable; it is an already existing Euler genealogy restricted to the true source-time window $[(t-h)_+,t]$*.*

Proposition 19.6 (No double charging)**.** *Infinite fractional Hodge-dominance refreshes do not imply infinite new negative cross-shear creation. Consequently:*

(i) one physical source-sign conversion may be charged only once;
(ii) the divergent logarithmic refresh bookkeeping cannot be identified with a divergent physical cross-shear bill without an additional Euler theorem;
(iii) every future cumulative contradiction must identify genuinely new source-time work, not repeated relabeling of old genealogy.

Proposition 19.7 (Strict-gap algebra)**.** *Assume $a^+ > 0$. Any realizable raw polarization lower bound*

$$a \geq \pi_*(a^+ + a^-), \qquad \pi_* > 0, \tag{19.15}$$

necessarily has $\pi_* \leq 1$ and is equivalent to

$$\boxed{\frac{a^-}{a^+} \leq \frac{1-\pi_*}{1+\pi_*} < 1.} \tag{19.16}$$

Conversely, if

$$a^- \leq (1-\delta)a^+, \qquad \delta \in (0,1), \tag{19.17}$$

then

$$\boxed{a \geq \frac{\delta}{2-\delta}(a^+ + a^-) > 0.} \tag{19.18}$$

Thus the raw Hodge polarization and the strict cancellation-gap formulations are algebraically interchangeable after physical admission.

20. Critical falsification propositions retained in the skeleton

Accounting bridge. A reliable audit also records transactions that are *not* legal. These no-go propositions prevent the grand finale from being financed by a false shortcut or by charging the same event twice.

Proposition 20.1 (Finite local action alone cannot enforce the cancellation gap)**.** *There exists a tagged one-cell reduced system with variables $q(t)$, $C_\beta^{\mathrm{cell}}(t)$, and $H(t)$ satisfying*

$$\dot q = -HC_\beta^{\mathrm{cell}}, \qquad \dot C_\beta^{\mathrm{cell}} = \frac{1}{4}Hq, \qquad \dot H = 2q,$$

with

$$q(0) = 1, \qquad C_\beta^{\mathrm{cell}}(0) = H(0) = 0,$$

which obeys the exact local source-gradient/slope/mixed-jet algebra and has finite cross-shear/restart action. Define the local source-history banks by

$$\dot H^+ = 2[q]_+, \qquad \dot H^- = 2[q]_-, \qquad H^\pm(0) = 0, \qquad a^\pm := \frac{1}{2}H^\pm.$$

For these banks one reaches a finite time at which

$$a^+ \to \sqrt{2}, \qquad a^- \to \sqrt{2}, \qquad a = a^+ - a^- \to 0. \tag{20.1}$$

Therefore no strict global inequality $a^- \le (1-\delta)a^+$ can follow from local tagged ODE structure plus finiteness of raw action alone. Whole-space spatial geometry is indispensable.

Proposition 20.2 (Frozen Hodge geometry alone has no uniform gap)**.** *Fix the aspect ratio $s := y/x$. For every $\varepsilon > 0$ there exist smooth rapidly decaying nonnegative H^+, H^- with*

$$H^+(x, sx) = sx + O(x^2), \qquad H^-(x, sx) = O(x^2), \tag{20.2}$$

with H^- supported away from the equator, such that their exact whole-space Hodge banks

$$a^\pm := \frac{3}{8}\int_{\mathcal{Q}} K_H H^\pm \, dX$$

satisfy

$$\frac{a^-}{a^+} > 1 - \varepsilon. \tag{20.3}$$

Hence instantaneous Hodge/Biot–Savart geometry plus the local sign orders is insufficient; the missing theorem must be dynamic and source-history specific.

Proposition 20.3 (Infinite fractional refresh does not imply infinite physical conversion)**.** *On $0 \le t < 1$, a two-bank bookkeeping model can have*

$$a^+(t) = 1, \qquad a^-(t) = t, \qquad a(t) = 1 - t, \tag{20.4}$$

with finite total adverse creation but infinitely many fractional Hodge-dominance refreshes as $t \uparrow 1$. Therefore one finite physical conversion may not be charged repeatedly to infinitely many bookkeeping refreshes.

Proposition 20.4 (Pointwise reproduction number is too strong)**.** *In the constant-rate affine conveyor, a label may leave the compact rotor region while a pointwise parent-to-child reproduction multiplier grows exponentially. The same label's final Hodge visibility decays faster, so the physical Hodge-weighted product still decays. Consequently final Hodge visibility must remain inside the causal estimate; a global pointwise reproduction bound < 1 is not the correct target.*

Remark 20.5 (Purpose of the no-go statements). These propositions are not defects of the program. They are the reason the present open bill has its specific form. They retire four false shortcuts: local finite-action coercivity, frozen Hodge coercivity, refresh-counting divergence, and pointwise reproduction-number coercivity.

21. Closed O-ledger with O6.2 as the sole open proposition

Accounting bridge. The reconstructed O-ledger has one unpaid proposition. O1–O3 are proved independently; the inherited-launch source-angle audit formerly denoted O6.1 is complete and therefore funds O4; M7 records three stronger terminal mechanisms; M8 gives the exact integral cashier countdown; and O5 is the proved downstream consequence of O4 together with O6.2 and M8. Thus O6.2 is the sole open proposition.

Definition 21.1 (Exact old primary conversion)**.** *For $h > 0$, let $\mathcal{G}_{\mathrm{rec}}$ and $\mathcal{I}_{+,t}$ be as in theorem 18.6 and proposition 18.7. Define*

$$\boxed{\mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{old,h}}(t) := \frac{3}{4}\int_0^{(t-h)_+} \mathcal{G}_{\mathrm{rec}}(t-\tau)\mathcal{I}_{+,t}(\tau)\,d\tau.} \tag{21.1}$$

This exact Hodge-visible account is the only genuinely new adverse account not already retired, resummed, or returned to an established restart/continuation ledger. The older factorized quantity obtained from $\mathcal{I}_{+,t} \le (4/3)\mathfrak{m}^{+}_{\mathrm{prim}} A^{+}_{t\leftarrow\tau}$ remains a legal upper bound, but the affine falsification shows that it is too strong to serve as the global old-age closure target.

Remark 21.2 (Physical age is not collapse age). On a possible floor branch $a(t) \downarrow 0$,

$$t-\tau \ge h \quad \not\Longrightarrow \quad \Lambda(t;\tau) = \int_\tau^t a(\xi)d\xi \ge c_h > 0. \tag{21.2}$$

Therefore aged-response decay may be used only after splitting by the *actual* relative collapse age $\Lambda(t;\tau)$.

Peter's first unpaid invoice is old exposure that has accumulated a genuinely large collapse age. The intended payment is the already proved aged-response discount.

Theorem 21.3 (Large-collapse-age physical-bank retirement)**.** *Work in the physical sector on a positive-compression branch*

$$a(t) \ge 0, \qquad \Lambda(t) := \int_0^t a(\xi)\,d\xi.$$

For $L > 0$ and an observation time t, define the large-collapse-age source-time set

$$\mathscr{E}_L(t) := \{\tau \in [0,t] : \Lambda(t) - \Lambda(\tau) \ge L\}.$$

Using the exact two-time Hodge visibility, define the physical sub-banks

$$\boxed{a^{\pm}_{\ge L}(t) := \frac{3}{4}\int_{\mathscr{E}_L(t)}\int_{\mathcal{Q}} [q(X,\tau)]_{\pm}\mathcal{V}_H(t;\tau,X)\,dX\,d\tau.} \tag{21.3}$$

Let

$$M(t) := \mathcal{K}_{\mathrm{cent}}([0,t]), \qquad \eta_M := 2\mathfrak{C}^{*}_{Z}\big(e^{8M(t)}-1\big).$$

There exists a fixed constant $\mathfrak{c}_{\mathrm{aff}} < \infty$, depending only on the fixed customer, such that with

$$\varepsilon_L := \frac{\mathfrak{c}_{\mathrm{aff}}}{1+L} \longrightarrow 0$$

one has

$$\boxed{a^{-}_{\ge L}(t) \le \varepsilon_L a^{+}(t) + \mathfrak{D}_{\mathrm{cent,L}}(t),} \tag{21.4}$$

where the centered-deformation comparison bill is

$$\boxed{\mathfrak{D}_{\mathrm{cent,L}}(t) := \frac{1+\varepsilon_L}{2}\,\eta_M\,\mathcal{V}_{\mathscr{E}_L(t),\mathrm{aff}}(t).} \tag{21.5}$$

Here $\mathcal{V}_{\mathscr{E}_L(t),\mathrm{aff}}$ is the source-time-localized affine absolute response defined in the proof. The bill in (21.5) is exactly the localized G4 centered-deformation comparison error; it is not a new mechanism. In the exact affine conveyor it vanishes identically. Thus genuine large collapse age is retired by an $O((1+L)^{-1})$ physical-bank ratio, modulo the already certified centered-deformation routing.

Proof-audit status: **O1 GREEN/COMPLETE**; grand-finale entry / long proof.

If that invoice is not old in collapse age, it moves to Peter's second account: mesoscopic exposure that is physically old but collapse-age young. Its payer must be the level-set polarization/deformation ledger.

Theorem 21.4 (Bounded-collapse-age physical-bank return)**.** *Work in the physical sector on a positive-compression branch*

$$a(t) \ge 0, \qquad \Lambda(t) := \int_0^t a(\xi)\,d\xi.$$

For $L > 0$, define the complementary bounded-age source-time set

$$\mathscr{E}_{<L}(t) := \{\tau \in [0,t] : 0 \le \Lambda(t) - \Lambda(\tau) < L\},$$

and the exact physical sub-banks

$$\boxed{a^{\pm}_{<L}(t) := \frac{3}{4}\int_{\mathscr{E}_{<L}(t)}\int_{\mathcal{Q}} [q(X,\tau)]_{\pm}\mathcal{V}_H(t;\tau,X)\,dX\,d\tau.} \tag{21.6}$$

Let

$$\pi^*_{\mathrm{aff}} := \frac{1-\upsilon^*_{\mathrm{aff}}}{1+\upsilon^*_{\mathrm{aff}}} > 0, \qquad \eta_M := 2\mathfrak{C}^*_Z(e^{8M}-1).$$

Choose once and for all $M_{\mathrm{O2}} > 0$, depending only on the fixed customer, so that

$$\eta_{M_{\mathrm{O2}}} < \frac{1}{2}\pi^*_{\mathrm{aff}},$$

and define

$$\pi_{\mathrm{O2}} := \frac{\pi^*_{\mathrm{aff}} - \eta_{M_{\mathrm{O2}}}}{1+\eta_{M_{\mathrm{O2}}}} > 0, \qquad \theta_{\mathrm{O2}} := \frac{1-\pi_{\mathrm{O2}}}{1+\pi_{\mathrm{O2}}} < 1. \tag{21.7}$$

Then exactly one of the following funded alternatives is available:

(i) if $\mathcal{K}_{\mathrm{cent}}([0,t]) \le M_{\mathrm{O2}}$, then

$$\boxed{a^-_{<L}(t) \le \theta_{\mathrm{O2}}a^+_{<L}(t) \le \theta_{\mathrm{O2}}a^+(t);} \tag{21.8}$$

(ii) if $\mathcal{K}_{\mathrm{cent}}([0,t]) > M_{\mathrm{O2}}$, then a definite centered deformation ticket has already been spent and, by the audited G5 routing,

$$\boxed{\int_0^t \mathcal{L}_{\mathrm{pol}}(\tau)\,d\tau > \frac{M_{\mathrm{O2}}}{3}.} \tag{21.9}$$

The constants are independent of L and t. No generic swirl-free meridional estimate is used. The compact mesoscopic F8 ticket remains an available localization refinement, but it is not needed to prove the bounded-age return.

Proof-audit status: **O2 GREEN/COMPLETE**; grand-finale entry / long proof.

Theorem 21.5 (Physical-bank consolidation or paid centered deformation)**.** *Work in the physical sector on a positive-compression branch*

$$a(\xi) \geq 0 \qquad (0 \leq \xi \leq t).$$

Let $\theta_{\mathrm{O2}} \in [0,1)$ and $M_{\mathrm{O2}} > 0$ be the constants from theorem 21.4, and put

$$\Delta_{\mathrm{O3}} := 1 - \theta_{\mathrm{O2}} > 0.$$

Choose $L_{\mathrm{O3}} > 0$ so that

$$\varepsilon_{L_{\mathrm{O3}}} \leq \frac{\Delta_{\mathrm{O3}}}{8}, \tag{21.10}$$

and choose $M_{\mathrm{O3}} \in (0, M_{\mathrm{O2}}]$ so small that, with

$$\eta_{\mathrm{O3}} := 2\mathfrak{C}_Z^*(e^{8M_{\mathrm{O3}}} - 1),$$

$$\eta_{\mathrm{O3}} < 1, \qquad \delta_{\mathrm{cent}} := \frac{1 + \varepsilon_{L_{\mathrm{O3}}}}{2} \frac{\eta_{\mathrm{O3}}}{1 - \eta_{\mathrm{O3}}} \leq \frac{\Delta_{\mathrm{O3}}}{8}. \tag{21.11}$$

Define

$$\boxed{\theta_{\mathrm{O3}} := \theta_{\mathrm{O2}} + \frac{\varepsilon_{L_{\mathrm{O3}}} + \delta_{\mathrm{cent}}}{1 - \delta_{\mathrm{cent}}} < 1.} \tag{21.12}$$

Then exactly one of the following funded alternatives is available:

(i) if $\mathcal{K}_{\mathrm{cent}}([0,t]) \leq M_{\mathrm{O3}}$, then

$$\boxed{a^-(t) \leq \theta_{\mathrm{O3}} a^+(t), \qquad \theta_{\mathrm{O3}} < 1;} \tag{21.13}$$

(ii) if $\mathcal{K}_{\mathrm{cent}}([0,t]) > M_{\mathrm{O3}}$, then the already audited centered-deformation routing gives

$$\boxed{\int_0^t \mathcal{L}_{\mathrm{pol}}(\xi)\, d\xi > \frac{M_{\mathrm{O3}}}{3}.} \tag{21.14}$$

No new adverse account is introduced.

Proof-audit status: **O3 GREEN/COMPLETE**; grand-finale entry / long proof.

Theorem 21.6 (Grand cancellation gap)**.** *The completed inherited-launch source-angle audit (the former O6.1 module), together with the proved O3 threshold-state gap, yields the following. On every finite classical branch of the fixed physical customer before a named continuation exit there exists*

$$\delta_* \in (0,1),$$

depending only on the fixed customer and admitted finite-action ball, such that

$$\boxed{a^-(t) \leq (1 - \delta_*) a^+(t)} \tag{21.15}$$

for every time on the branch.

For every $t > 0$ one has

$$\boxed{a(t) \geq \frac{\delta_*}{2 - \delta_*} [a^+(t) + a^-(t)] > 0.} \tag{21.16}$$

At the active opening time $t = 0$, where $a(0) = 0$, the bank-gap inequality (21.15) remains valid and the algebraic lower bound is understood non-strictly.

One may take

$$\boxed{\delta_* = \min\{1 - \theta_{\mathrm{O3}}, \delta_6\}.} \tag{21.17}$$

Proof-audit status: **O4 GREEN/COMPLETE**. The inherited-launch source-angle audit is closed; only O6.2 remains open. See grand-finale entry / long proof.

Finally, a positive bank gap must be returned to a terminal finite-time mechanism. The historical O5 formulation uses the apex reserve $Q - 9a^2$. The reconstructed proof body also records the already proved powered-reserve and pressure-free terminal routes, because the special coefficient 9 is sufficient but not logically necessary.

Theorem 21.7 (Terminal noncontinuation from O4 and O6.2)**.** *Assume the Grand cancellation gap theorem 21.6 and subclause O6.2 of open proposition 21.8. Then either a named classical continuation exit occurs earlier, or the fixed customer leaves the classical category in finite physical time.*

Indeed O6.2 supplies a finite maintained-cashier launch for which the exact integral condition (21.18) holds. Apply theorem K.5. The previously certified powered, positive-bank-growth, and fixed sub-quarter cashier gates remain useful stronger sufficient routes, but they are no longer built into the weakest terminal obligation.

Proof-audit status: **O5 GREEN/COMPLETE: O4 + O6.2 $\Longrightarrow$ terminal noncontinuation**. O6.2 is the exact integral cashier obligation. See grand-finale entry / long proof.

Open Proposition 21.8 (Post-O4 integral cashier closure (O6.2))**.** *The inherited-launch source-angle audit formerly denoted O6.1 is closed and already funds O4. The sole remaining open proposition is O6.2.*

On every admitted post-O4 branch on which no named continuation exit has already occurred, prove that there exists a finite maintained-cashier launch time $t_1 > 0$ with

$$\Sigma(t_1) > 0$$

for which the net cashier gain

$$\mathcal{G}_{\mathrm{cash}}(\lambda) = \frac{\lambda}{4} - \mathfrak{D}_{\mathrm{rate}}(\lambda), \qquad \lambda = \Lambda(t; t_1),$$

satisfies

$$\boxed{\int_0^\infty e^{-\mathcal{G}_{\mathrm{cash}}(\lambda)}\, d\lambda < \infty.} \tag{21.18}$$

By theorem K.5, this is precisely the terminal integrability needed for finite physical-time noncontinuation.

The permanent M7 powered, positive-bank-growth, and fixed sub-quarter cashier gates remain useful stronger sufficient routes. The M8 integral cashier theorem is the weaker terminal engine used in the final chain. A concrete sufficient target is

$$\boxed{\frac{\Delta_{\mathrm{B}}}{a} \le \frac{1}{4} - \frac{p}{1+\Lambda}, \qquad p > 1,} \tag{21.19}$$

on a sufficiently late post-O4 tail. More generally, any lower bound on $\mathcal{G}_{\mathrm{cash}}$ for which $e^{-\mathcal{G}_{\mathrm{cash}}} \in L^1(d\lambda)$ is admissible.

Proof-audit status: **O6.2 is the sole OPEN proposition—NO PROOF ASSERTED.** M8 is the exact post-O4 integral cashier theorem; M7-P/C1/C2 remain stronger sufficient terminal mechanisms. The permanent label `open:sk-first-ticket-source-angle-closure` is retained for navigation. See grand-finale entry / long-proof slot.

22. Rigorous-proof architecture and navigation

Accounting bridge. The funded ledger now has three audit layers. The grand-finale table identifies *which account pays*; the short-proof layer records the compressed mathematical reason; the rigorous appendices after the bibliography hold the referee-level line-by-line proofs. The three layers use permanent ledger IDs and mutual hyperlinks:


$$\boxed{\text{grand-finale entry} \iff \text{short proof} \iff \text{rigorous long proof.}}$$

The proof layer is organized account by account without changing theorem labels, ledger IDs, or the permanent O1–O6 navigation labels. Every closed theorem-like ledger item has a referee-level long proof. O1–O3 are proved independently, O4 is GREEN from the closed inherited-launch source-angle module, O5 is GREEN from O4+O6.2+M8, and O6.2 is the sole unpaid package.

23. Grand-finale theorem/proposition chain and dependency table

Accounting bridge. This section is Peter's audited general ledger. The schematic shows the direction of funding; the detailed table identifies every proved theorem, proposition, lemma, and corollary used by the grand-finale chain, together with the sole proposition O6.2 that remains open. A professional reader can therefore enter the proof at any box, follow the exact cross-reference, and see which later account that statement funds.

23.1. Schematic funded chain.



The compact schematic is therefore

$$\begin{aligned}&\text{customer/source/cashier} \Longrightarrow \text{detector reserve} \Longrightarrow \text{restart bills} \Longrightarrow (H^{\pm}, a^{\pm})\\ &\Longrightarrow \text{positive cashier/relaunch} \Longrightarrow \text{adverse ancestry/capacity} \Longrightarrow \text{affine response barrier}\\ &\Longrightarrow \text{mesoscopic rotor} \Longrightarrow \text{level-set polarization stability} \Longrightarrow \text{causal three-account bound}\\ &\Longrightarrow \boxed{\text{closed O6.1 audit} \Rightarrow O4, \quad O6.2 + O4 + M8 \Rightarrow O5}\\ &\Longrightarrow \boxed{O6.2 \text{ sole open}} \Longrightarrow \text{reserve return} \Longrightarrow \text{conditional finite-time clock collapse.}\end{aligned} \tag{23.1}$$

23.2. Detailed theorem/proposition chain.

ID	Exact statement	Accounting transfer	Funds / constrains
A. Exact customer, source, cashier, clock, and opening credit			
A1	proposition 4.3 short proof — long proof	opens the tagged Euler accounts and the true flow	A2–A5
A2	proposition 4.4 short proof — long proof	keeps physical/squared pressure books distinct	reserve/Hessian audits
A3	proposition 5.3 short proof — long proof	$\Theta \to q \to H$ without pressure	A4, D1, H1

ID	Exact statement	Accounting transfer	Funds / constrains
A4	proposition 5.4 short proof — long proof	$H \to a$ through a positive Hodge kernel	A5, D1, G1
A5	proposition 5.5 short proof — long proof	makes meridional values/jets Hodge descendants of H	Hodge/shear routing
A6	propositions 6.4 and 6.5 short proof — long proof	$a \to \psi_\beta \to \Psi \to \mathfrak{t}_\Psi$ clock; local vorticity-Hessian bridge	A7–A8
A7	proposition 6.7 short proof — long proof	converts reserve sign to clock concavity	A8
A8	theorem 6.8 short proof — long proof	positive compression + nonnegative reserve buys finite deadline	terminal assembly
A9	proposition 7.1 short proof — long proof	proves a source family opens with positive compression	A10
A10	proposition 7.2 short proof — long proof	proves the initial reserve is funded	persistence problem
B. Provenance and restart/deformation accounting			
B1	proposition 8.2 short proof — long proof	separates background routing from tagged money	all terminal payments
B2	proposition 8.4 short proof — long proof	forbids swirl-free/background money from paying the terminal bill	entire chain
B3	proposition 9.2 short proof — long proof	expresses tangent deformation in the restart basis	B4, H7
B4	lemma 9.3 short proof — long proof	tangent exit spends definite first-jet action	B5, G5
B5	proposition 9.4 short proof — long proof	converts value defects into radial star-hull bills	B6, I4
B6	proposition 9.6 short proof — long proof	identifies the signed first-jet channels as one coupled ledger	H1–H7
B7	lemma 9.7 short proof — long proof	pointwise restart credit either thickens or pays a derivative exit	relaunch/capacity routing
C. Genealogy banks and positive-cashier relaunch			
C1	proposition 10.2 short proof — long proof	splits source history into nonnegative a^+, a^- banks	C2–C3
C2	theorem 10.3 short proof — long proof	turns floor loss into $a^+/a^- \to 1$ cancellation	modern target
C3	theorem 10.5 short proof — long proof	shows the cancellation ratio is a scorekeeper, not a divergent currency	no-double-charge rule
C4	proposition 11.1 short proof — long proof	supplies fresh positive local credit	C5, C12
C5	proposition 11.2 short proof — long proof	turns a positive material block into a Riccati cashier	C7
C6	proposition 11.3 short proof — long proof	loss of a selected cashier pays radial restart action	C11, B5
C7	theorem 12.2 short proof — long proof	persistent intrinsic gates force a finite deadline	C10
C8	proposition 12.3 short proof — long proof	rate failure is a paid star-hull bill	B5
C9	theorem 12.4 short proof — long proof	visibility loss is positive-credit tail concentration	C10
C10	theorem 12.5 short proof — long proof	exhausts intrinsic cashier exits	relaunch graph
C11	lemma 12.7 short proof — long proof	matches positive cohort mass to Hodge credit	C13
C12	proposition 12.8 short proof — long proof	gives exact Hodge-matched contraction rate	C13

ID	Exact statement	Accounting transfer	Funds / constrains
C13	theorem 12.9 short proof — long proof	turns Hodge-matched contraction into Riccati growth	C14
C14	theorem 12.10 short proof — long proof	launches new gate-ready credit from the same customer	C15–C17
C15	theorem 12.11 short proof — long proof	prevents hidden shrinking Taylor tubes on compact intervals	C16
C16	theorem 12.12 short proof — long proof	converts terminal loss of uniformity into a named continuation exit	C17, J
C17	proposition 12.13 short proof — long proof	excludes free finite-time relaunch accumulation	modern no-reset ledger
D. Adverse ancestry and ordinary continuation capacity			
D1	proposition 13.1 short proof — long proof	proves negative source has only old slope or tagged cross-shear ancestry	G5, H1, J
D2	proposition 13.2 short proof — long proof	keeps fixed old adverse source away from the fresh equator	G2, O1
D3	proposition 14.2 short proof — long proof	caps homogeneous source-gradient amplitude under finite restart action	H1, I3
D4	proposition 14.4 short proof — long proof	returns weighted H energy to the same GREEN source-gradient stock	D5, J2
D5	proposition 14.5 short proof — long proof	prices physical meridional gradients by tagged Hodge energy	far-field/pressure routing
D6	theorem 14.6 short proof — long proof	makes far-field and terminal regularity failures named exits	G6, H7, O1–O2
E. Two-time response and the fixed-seed affine benchmark			
E1	proposition 15.2 short proof — long proof	expresses apex credit as tagged source times a vertical response kernel	E2, I1
E2	theorem 15.3 short proof — long proof	discounts source with genuinely large relative collapse age	O1
E3	lemma 15.4 short proof — long proof	proves the fixed customer's affine line source is positive at every angle	E4
E4	theorem 15.5 short proof — long proof	gives a uniform affine Hodge polarization margin	E5, I3
E5	theorem 15.6 short proof — long proof	losing the affine margin spends deformation or new negative response	E6, H7
E6	proposition 15.7 short proof — long proof	makes the fresh apex tube source-sign protected	E7, J3
E7	theorem 15.8 short proof — long proof	near-lock requires an order-one adverse response outside the tube	H7
F. Source-gradient rotor and compact mesoscopic conversion			
F1	proposition 16.1 short proof — long proof	writes source rotation in the established restart basis	F2
F2	proposition 16.3 short proof — long proof	separates H-skew rotation from signed-$\mathfrak{T}$ deformation	F3–F6
F3	theorem 16.4 short proof — long proof	proves H rotates source direction without directly amplifying its norm	modern rotor logic
F4	proposition 16.5 short proof — long proof	identifies the rotor as deformation of one fixed transported circulation gradient	I1, O2

ID	Exact statement	Accounting transfer	Funds / constrains
F5	proposition 16.6 short proof — long proof	makes mixed shear a global Hodge moment, not a free coefficient	F6
F6	proposition 16.7 short proof — long proof	factors the adverse exchange into $H/\mathfrak{T}$ times radial source gradient	F7
F7	proposition 16.8 short proof — long proof	retires both extreme angular tails	F8
F8	theorem 16.9 short proof — long proof	localizes every substantial new adverse conversion to a compact rotor ticket	I2
G. Circulation-level-set polarization and deformation stability			
G1	theorem 17.1 short proof — long proof	rewrites $[q]_{\pm}$ responses as fluxes on transported circulation level sets	G2
G2	proposition 17.2 short proof — long proof	converts strict level-set polarization into the desired bank gap	G4, O4
G3	lemma 17.4 short proof — long proof	controls the full source-gradient vector by the transverse fixed-seed stock	G4
G4	theorem 17.5 short proof — long proof	small centered deformation preserves the affine polarization	G5, O2
G5	corollary 17.6 short proof — long proof	first polarization loss spends a definite old star-hull/tangent bill	O2
H. Causal adverse-response ledger			
H1	proposition 18.1 short proof — long proof	compares final and deposition visibility without $1/a$	H2
H2	proposition 18.2 short proof — long proof	prevents response amplification from becoming an anonymous new mechanism	H4
H3	proposition 18.3 short proof — long proof	makes fresh source rotation small at physical apex scale	H5
H4	theorem 18.4 short proof — long proof	gives the exact positive two-time negative-generation functional	H5–H7
H5	proposition 18.5 short proof — long proof	prices the last physical-age window by $O(h^2)a^+$	O3
H6	theorem 18.6 short proof — long proof	resums $H^- \to H^-$ recycling into a finite Volterra amplifier	H7
H7	proposition 18.7 short proof — long proof	reduces a^- to old ancestry + signed $\mathfrak{T}$ + primary $H^+ \to H^-$ conversion	O1–O4
H8	conditional proposition 18.8 short proof — long proof	a strict three-account bound immediately yields the bank gap	O4
I. Supporting exact identities retained on the funded chain			
I1	proposition 19.1 short proof — long proof	exact age/visibility derivative	E2, H1
I2	proposition 19.2 short proof — long proof	source-finances $\dot{a}$ before any normalization	reserve audits
I3	proposition 19.3 short proof — long proof	keeps adverse source in tagged raw form	D1, H4
I4	proposition 19.4 short proof — long proof	converts Hodge kernel to carrier weight	capacity estimates
I5	proposition 19.5 short proof — long proof	identifies fresh response with true Euler source history	C14, H5
I6	proposition 19.6 short proof — long proof	forbids charging one finite conversion to later bookkeeping refreshes	entire modern ledger
I7	proposition 19.7 short proof — long proof	converts a bank fraction into positive net compression	O4–O5

ID	Exact statement	Accounting transfer	Funds / constrains
J. Falsification firewalls (proved side constraints)			
J1	proposition 20.1 short proof — long proof	local exact dynamics + finite action do not force a strict gap	forbids local-only closure
J2	proposition 20.2 short proof — long proof	frozen whole-space Hodge geometry does not force a strict gap	requires two-time response
J3	proposition 20.3 short proof — long proof	infinite refreshes need not mean infinite physical conversion	enforces no-double-charge
J4	proposition 20.4 short proof — long proof	pointwise reproduction can grow while Hodge-weighted response decays	requires causal visibility-weighted resolvent
K. Permanent reconstructed modules M1–M8			
M1	theorems K.11 to K.13 short proof — long proof	polarization/source-history structure	M2
M2	theorems K.14, K.15 and K.17 short proof — long proof	source-positive critical cashier and fractional homotopy	M3
M3	theorems K.19 and K.20 short proof — long proof	post-ticket causal pay-or-route / no-double-charge	M4
M4	theorems K.21, K.22 and K.25 short proof — long proof	Gate-S / critical-to-Hodge source-angle bridge	M5–M6
M5	theorems K.27 and K.28 short proof — long proof	ratio-free GREEN barrier and bounded-age crossing	M6
M6	theorems K.30, K.31 and K.33 to K.36 short proof — long proof	global continuation/source-potential/aspect/tangent source-angle audit	O6.1, O4
M7	theorems K.1, K.3 and K.4 short proof — long proof	stronger fixed-margin terminal routes P/C1/C2	M8, O5
M8	theorems K.5 and K.6 short proof — long proof	exact post-O4 integral cashier countdown; M7 routes are sufficient special cases	O5
L. O-ledger: O6.2 sole open item			
O1	theorem 21.3 short proof — long proof	retire the exact Hodge-visible large-age primary exposure using cashier feedback or paid exits	O3
O2	theorem 21.4 short proof — long proof	return exact collapse-age-young old exposure to polarization/deformation bills	O3
O3	theorem 21.5 short proof — long proof	consolidate complementary physical age banks into a strict gap or paid centered deformation	O4
O4	theorem 21.6 short proof — long proof	**GREEN:** the closed inherited-launch audit plus O3 yield the uniform Hodge gap	O5
O5	theorem 21.7 short proof — long proof	**GREEN:** O4 + O6.2 + M8 give finite-time noncontinuation	L2
O6.2	open proposition 21.8 short proof — long proof	**SOLE OPEN:** produce the exact M8 integral cashier activation on the post-O4 no-exit branch	O5, L2
M. Grand-finale assembly			
L1	theorem 24.1 short proof — long proof	consolidates the funded infrastructure and O1–O3 base feeding M1–M8	O6.2

ID	Exact statement	Accounting transfer	Funds / constrains
L2	conditional proposition 24.2 short proof — long proof	the closed source-angle audit funds O4; O6.2+O4+M8 fund O5; hence O6.2 implies the conditional Grand Finale	final conditional conclusion

Remark 23.1 (How to read the table). Rows A–I are funded. Rows J are negative results that prevent illegal transfers. O1–O3 are paid; the inherited-launch source-angle audit (O6.1) is closed; O4 is GREEN; and O5 is GREEN from O4+O6.2+M8. O6.2 is the sole OPEN proposition. The final row L2 is therefore conditional only on O6.2, not an unconditional singularity theorem. This distinction is part of the proof, not merely a statement of style.

Remark 23.2 (Permanent three-way proof navigation). The capital-letter IDs in the detailed table (A1, A2, ..., L2) are permanent ledger row identifiers and are deliberately distinct from the lower-case nodes $(a), (b), \dots, (l)$ in the schematic above. Every row now carries two links: *short proof* and *long proof*. The short-proof layer sits immediately after the grand-finale assembly. The rigorous long-proof appendices sit after the bibliography.

Conversely, every short proof and every long-proof slot contains links back to its grand-finale entry and to the other proof layer. The long-proof appendix restates the complete theorem/proposition/lemma before the proof. During the audit, a pending closed slot was marked **pending**; in the audit no closed slot remains pending. An Open Proposition is marked **OPEN—NO PROOF ASSERTED**. These anchors are fixed throughout the proof audit.

24. Grand-finale proof

Accounting bridge. The preceding table is the dependency graph; this section is its compressed assembly proof. It identifies exactly where the proof is closed and exactly where Peter's remaining invoices enter.

Theorem 24.1 (Funded chain up to the open bill)**.** *For the fixed $A = 10$ customer, every implication in rows A–I of section 23 is proved. In particular,*

$$\begin{aligned}
&\textit{fixed tagged customer} \Longrightarrow (\Theta, q, C_\beta) \Longrightarrow H^{\pm} \Longrightarrow a^{\pm},\\
&\textit{affine fixed-seed benchmark} \Longrightarrow \upsilon_{-}^{\mathrm{aff}} \le \upsilon_{\mathrm{aff}}^{*} < 1,\\
&\textit{small centered true-Euler deformation} \Longrightarrow a \ge \pi_{107}(a^{+} + a^{-}),\\
&\textit{first polarization loss} \Longrightarrow \textit{definite star-hull/tangent bill},\\
&\textit{near-lock conversion} \Longrightarrow \textit{compact mesoscopic rotor ticket},\\
&\textit{fresh primary conversion} = O(h^2)a^{+},\\
&\textit{negative self-recycling} \Longrightarrow \textit{finite causal resolvent},\\
&\textit{O1–O3 physical-bank closure} \Longrightarrow \textit{strict bank gap or paid centered deformation.}
\end{aligned} \tag{24.1}$$

No generic swirl-free Euler estimate is used as terminal payment, and one physical conversion is never charged twice. The downstream reconstructed ledger adds M1–M8, O4 and O5 without changing this funded base; the only remaining open package is O6.2.

Mutual pointers: **L1**: entry / long proof.

Proof. The source/cashier chain is rows A1–A5. Rows A6–A8 convert sustained positive compression and reserve into a finite detector deadline. Rows B1–B7 ensure that leaving the favorable geometry spends a certified source-specific deformation bill. Rows C1–C17 create persistent positive/negative genealogy banks and exclude free relaunch resets. Rows D1–D6 reduce adverse ancestry and ordinary continuation costs to a finite list of named accounts. Rows E1–E7 supply the fixed-seed affine polarization and show that substantial loss of that margin requires a real conversion. Rows F1–F8 identify that conversion as a compact circulation-gradient rotor. Rows G1–G5 show that the affine level-set polarization is stable until a definite centered-deformation ticket is spent. Finally rows H1–H8 keep final Hodge visibility inside the adverse-response estimate, make fresh primary conversion $O(h^2)$, and resum negative self-recycling. Rows I1–I7 are the exact identities that legalize these transfers. The permanent O-ledger then closes O1–O3 in the exact physical source-time-bank formulation. The four rows J1–J4 exclude the four shortcuts that would otherwise bypass this funded chain. □

Conditional Proposition 24.2 (Grand finale)**.** *Assume the sole remaining open proposition O6.2, open proposition 21.8. Then either a named classical continuation exit occurs earlier, or the fixed customer loses classical regularity in finite time.*

Mutual pointers: **L2**: entry / long proof.

Proof. Assume O6.2. The closed inherited-launch source-angle audit already gives O4 by theorem 21.6. O6.2 together with O4 and M8 gives O5 by theorem 21.7. Therefore either a named continuation exit has already occurred or finite-time noncontinuation follows. □

25. Short proofs of the closed grand-finale statements

25.1. **Short-proof location index.**

Accounting bridge. The main-text chain says which account funds which later account. This section says where the corresponding rigorous proof is kept. The map is stable under future proof expansion: the chain IDs and main-text theorem labels do not change when an appendix proof is lengthened.

Chain account	Closed statements	Short-proof account / rigorous appendix
A	A1–A10	Short A / Appendix A
B	B1–B7	Short B / Appendix B
C	C1–C17	Short C / Appendix C
D	D1–D6	Short D / Appendix D
E	E1–E7	Short E / Appendix E
F	F1–F8	Short F / Appendix F
G	G1–G5	Short G / Appendix G
H	H1–H8	Short H / Appendix H
I	I1–I7	Short I / Appendix I
J	J1–J4	Short J / Appendix J
M1–M8	proved permanent modules	Short M / module long-proof cards in Appendix K
O1–O3	proved independently	theorem/long-proof cards / Appendix K
O4–O5	GREEN downstream from closed O6.1 and O6.2+M8	theorem/long-proof cards / Appendix K
O6.2	sole open proposition	open-status card / Appendix K
L1–L2	assembly/conditional proof	grand-finale section / Appendix L

25.2. **A. Customer, source, cashier, clock, and opening-credit short proofs.** Statements proved. Rows A1–A10 of the detailed dependency table.

Mutual pointers: **A1**: entry / long proof; **A2**: entry / long proof.

Proof of propositions 4.3 and 4.4. Insert $U^r = -rv$, $U^\phi = ru$, $U^z = zg$, $x = r^2$, $y = z^2$ into the axisymmetric Euler equations. The identities $\partial_r = 2r\partial_x$ and $\partial_z = 2z\partial_y$ give the transport operator $\mathcal{D}_{xy} = \partial_t - 2xv\partial_x + 2yg\partial_y$ and incompressibility gives (4.15). The radial, swirl, and axial momentum equations give (4.16)–(4.18); multiplying the swirl equation by the Boolean tag gives (4.17). The pressure formulas follow from the same two chain rules: $\mathsf{p}_r = 2rP_x$, $\mathsf{p}_z = 2zP_y$, followed by one more derivative. □

Mutual pointers: **A3**: entry / long proof; **A4**: entry / long proof; **A5**: entry / long proof.

Proof of propositions 5.3 to 5.5. From $\mathcal{D}_{xy}u = 2vu$ in the physical sector, $\mathcal{D}_{xy}\Theta = 4v\Theta$. Differentiating the meridional equations, subtracting the mixed pressure derivatives, and using incompressibility gives $\mathcal{D}_{xy}H + gH = 2\Theta_y$; in physical variables this is $D_t\bar{G} = \varepsilon_\beta\partial_z(\beta^2)$. The stream function $U^r = -r\psi_{1,z}$, $U^z = 2\psi_1 + r\psi_{1,r}$ solves $-\mathcal{L}_1\psi_1 = \bar{G}$. Evaluating $\psi_{1,z}$ at the origin with the whole-space Green kernel yields (5.11), and the change $(r, z) \mapsto (x, y)$ yields (5.12). Finally writing $\psi_1 = z\chi$ gives (5.13)–(5.14); radial lifting in four X-variables and three Z-variables identifies $\mathcal{L}_7$ with $\Delta_{\mathbb{R}^7}$. □

Mutual pointers: **A6**: entry / long proof; **A7**: entry / long proof; **A8**: entry / long proof.

Proof of propositions 6.5 and 6.7 and theorem 6.8. The apex swirl equation gives $\dot{b} = 2ab$, hence $b(0) = 0$ implies $b \equiv 0$. Differentiate the swirl equation once in y and evaluate at the apex to obtain $\dot{\psi}_\beta = -2a\psi_\beta$; squaring gives $\dot{\Psi} = -4a\Psi$, and the idempotent tag gives $\dot{\mathcal{J}}_\beta = -4a\mathcal{J}_\beta$, hence (6.5)–(6.6). The apex radial equation with $b = 0$ is $\dot{a} = -a^2 + Q/3$, so

$\mathcal{F}_\Psi = Q - 9a^2 = 3(\dot{a} - 2a^2)$. Differentiating $a = -\dot{\mathfrak{t}}_\Psi/(2\mathfrak{t}_\Psi)$ yields $\mathcal{F}_\Psi = -(3/2)\ddot{\mathfrak{t}}_\Psi/\mathfrak{t}_\Psi$. If $a(t_1) > 0$ and $\mathcal{F}_\Psi \geq 0$, then $\mathfrak{t}_\Psi$ is positive, decreasing, and concave with $\dot{\mathfrak{t}}_\Psi(t_1) = -2a(t_1)\mathfrak{t}_\Psi(t_1)$. The tangent line reaches zero no later than $t_1 + [2a(t_1)]^{-1}$, so a classical positive clock cannot persist beyond that time. If the solution survives to the zero, $\Psi = \Psi(0)\mathfrak{t}_\Psi^2 \to 0$ while $-\dot{\mathfrak{t}}_\Psi/(2\mathfrak{t}_\Psi) = a \to +\infty$. □

Mutual pointers: **A9**: entry / long proof; **A10**: entry / long proof.

Proof of propositions 7.1 and 7.2. At $t = 0$ the meridional field vanishes, so the first derivative of the Hodge cashier is obtained by inserting the initial source $2\Theta_y$ into the positive Hodge kernel. For the one-parameter exponential ridge this integral separates and gives (7.2); the remaining integral $\mathcal{I}(\lambda)$ is strictly positive, so the sign is that of $c_* - s_*\lambda$, with threshold $c_*/s_* = \sqrt{2}$. Since $a(0) = b(0) = 0$, the apex equation gives $Q(0) = 3\dot{a}(0)$ and therefore $\mathcal{F}_\Psi(0) = Q(0)$. □

25.3. **B. Provenance and restart/deformation short proofs.**

Statements proved. Rows B1–B7 of the detailed dependency table.

Mutual pointers: **B1**: entry / long proof; **B2**: entry / long proof.

Proof of propositions 8.2 and 8.4. The normal form follows by induction over sums and products using $\varepsilon_\beta^2 = \varepsilon_\beta$: the routing parts multiply into a routing term and every term containing at least one GREEN factor carries one copy of ε_β. Differentiation, linear Hodge operators, and integration preserve this decomposition. A terminal inequality which survives the single-seed shadow has therefore lost the source mechanism; one which would imply a generic no-swirl Euler singularity cannot be a source-specific terminal payment either. This proves the two tests. □

Mutual pointers: **B3**: entry / long proof; **B4**: entry / long proof.

Proof of proposition 9.2 and lemma 9.3. Linearize the characteristic system $\dot{x} = -2xv$, $\dot{y} = 2yg$ and divide infinitesimal displacements by x and y. Substitution of the protected first-jet identities for C, D, E, F yields the logarithmic tangent matrix stated in the proposition. If a tangent column starting from e_j acquires defect at least η, Gronwall gives $\|\mathcal{U}(t)e_j - e_j\| \leq e^{\int \|\mathbb{M}\|} - 1$; hence $\int \|\mathbb{M}\| \geq \log(1 + \eta)$. On a compact angular window the matrix norm is bounded by a fixed multiple of $|\mathfrak{A}| + |\mathfrak{J}| + |\mathfrak{S}| + |\mathfrak{T}|$. □

Mutual pointers: **B5**: entry / long proof; **B6**: entry / long proof; **B7**: entry / long proof.

Proof of propositions 9.4 and 9.6 and lemma 9.7. For $X = (x, y)$, the radial fundamental theorem gives $v(X) - a = \int_0^1 \mathcal{R}_v(\lambda X)\, d\lambda/\lambda$, where $\mathcal{R}_v = xv_x + yv_y$. Expressing $xv_x + yv_y$ through the protected first-jet basis gives the star-hull identity. Differentiating $\mathfrak{A} = 2v - g$ in x and y and substituting the same basis identities gives the first-order bridges among $\mathfrak{A}, \mathfrak{J}, H, \mathfrak{T}$. For the log-box lemma, if the logarithmic gradient is at most $ca/(4\rho)$, integration along two coordinate segments shows $\mathcal{F} \geq ca/2$ throughout a box of log-area $4\rho^2$; otherwise the gradient alternative already holds. Integrating the positive part gives the stated thickened credit. □

25.4. **C. Genealogy-bank and positive-cashier short proofs.**

Statements proved. Rows C1–C17 of the detailed dependency table.
Mutual pointers: **C1**: entry / long proof; **C2**: entry / long proof; **C3**: entry / long proof.

Proof of proposition 10.2 and theorems 10.3 and 10.5. Reynolds' formula applied to the continuity equations for $H^{\pm}$ gives the two-bank identities, because $\mathbf{b}\cdot\nabla K_H = \mathcal{C}K_H$. The bounded angular coefficients in $\mathcal{C}$, the seven-dimensional first-jet norm, and $\sup K_H\mathfrak{r} < \infty$ give the displayed meridional/source bounds. Gronwall then bounds each bank and its total variation on a finite raw-action interval. The exact apex jet equation supplies $a^+(t_0) > 0$ for $t_0 > 0$, so the positive bank has a strictly positive terminal limit. If $a = a^+ - a^- \to 0$, the two finite limits coincide. Finally, on the floor branch, divide the two exact bank equations by their own positive banks and subtract. The own-bank source actions and the contraction actions are integrable, yielding (10.13)– (10.14). □

Mutual pointers: **C4**: entry / long proof; **C5**: entry / long proof; **C6**: entry / long proof.

Proof of propositions 11.1 to 11.3. The equatorial jet identity $\dot{q}_H + 6aq_H = \mathcal{J}_\beta$, together with $q_H(0) = 0$ and the nonvanishing detector, gives $q_H(t) > 0$ for $t > 0$. The negative source genealogy has zero first equatorial jet, so Taylor expansion gives $H^+ = q_H s x + O(x^2)$ and $H^- = O(x^2)$ on compact cones. Reynolds' formula for the positive-genealogy continuity equation with weight $1/x$ gives the exact cashier identity (11.3). For the reset firewall, the unweighted material genealogy bank is monotone; losing a fixed fraction of $M_{\mathcal{B}}^+$ therefore forces radial expansion of one trajectory and hence the negative-v action (11.5). The corrected radial fundamental theorem $v(X) - a = \int_0^1 \mathcal{R}_v(\lambda X)d\lambda/\lambda$ and the exact restart basis decomposition of $[-\mathcal{R}_v]_+$ give (11.6). □

Mutual pointers: **C7**: entry / long proof; **C8**: entry / long proof; **C9**: entry / long proof; **C10**: entry / long proof.

Proof of theorems 12.2, 12.4 and 12.5 and proposition 12.3. From the definitions, $\dot{M}_{\mathcal{B}}^+ = 2a\Lambda_{\mathcal{B}}^+ M_{\mathcal{B}}^+$. The Hodge inequality $a \geq a_{\mathcal{B}}^+ = \eta_{\mathcal{B}}^+ M_{\mathcal{B}}^+$ and the lower bounds $\eta_{\mathcal{B}}^+ \geq m_*$, $\Lambda_{\mathcal{B}}^+ \geq \nu$ give $\dot{M}_{\mathcal{B}}^+ \geq 2\nu m_* (M_{\mathcal{B}}^+)^2$ and the Riccati deadline. If $\Lambda_{\mathcal{B}}^+ < \nu$, rearranging its defining quotient yields (12.8); writing $a - v$ by the corrected logarithmic radial star-hull integral gives (12.9). Since $\eta_{\mathcal{B}}^+$ is the $\mu_{\mathcal{B}}^+$-average of h_0, the elementary estimate $\int h_0 d\mu \geq \vartheta\mu\{h_0 \geq \vartheta\}$ gives the tail lower bound when the mean falls below m_*. On an admitted interval with $a > 0$ and $M_{\mathcal{B}}^+ > 0$, either the three Riccati gates $a \geq a_{\mathcal{B}}^+$, $\eta_{\mathcal{B}}^+ \geq m_*$, $\Lambda_{\mathcal{B}}^+ \geq \nu$ all persist, or the first failed gate is respectively Hodge dominance, visibility, or rate. The preceding three results route those failures, while Hodge-dominance failure forces nonzero negative genealogy. Hence the four intrinsic branches are exhaustive. □

Mutual pointers: **C11**: entry / long proof; **C12**: entry / long proof; **C13**: entry / long proof.

Proof of lemma 12.7, proposition 12.8, and theorem 12.9. Cauchy–Schwarz applied to

$$N_K = \int \left(\frac{h_0(s)}{x} K_{\mathcal{B}}\right)^{1/2} \left(\frac{x}{h_0(s)} K_{\mathcal{B}}\right)^{1/2} dX$$

gives $N_K^2 \leq a_K \Pi_H$. For the compact frozen cohort, $\partial_t K_{\mathcal{B}} + \nabla\cdot(\mathbf{b}K_{\mathcal{B}}) = 0$, hence N_K is conserved and

$$\frac{d}{dt}\int w K_{\mathcal{B}}\, dX = \int (\mathcal{D}_{xy} w) K_{\mathcal{B}}\, dX.$$

With $w = x/h_0(s)$, $\mathcal{D}_{xy}\log x = -2v$, $\mathcal{D}_{xy}\log s = 2(v+g)$, and $d\log h_0/d\log s = (1-4s)/(2(1+s))$, which yields $\mathcal{D}_{xy}\log w = -\mathcal{C}_H$ and therefore (12.19).

Under the two raw C13 gates, C11 gives $a\Pi_H \geq a_K\Pi_H \geq N_K^2$, so $-N_K^2\dot{\Pi}_H \geq \kappa N_K^4$ before any division. Only then set $\mathfrak{Z}_H = N_K^2/\Pi_H$. Conservation of N_K gives $\dot{\mathfrak{Z}}_H = -(N_K^2\dot{\Pi}_H)/\Pi_H^2 \geq \kappa a\mathfrak{Z}_H$, while $\mathfrak{Z}_H \leq a_K \leq a$ gives $\dot{\mathfrak{Z}}_H \geq \kappa\mathfrak{Z}_H^2$ and the deadline $t_0 + [\kappa\mathfrak{Z}_H(t_0)]^{-1}$. □

Mutual pointers: **C14**: entry / long proof; **C15**: entry / long proof; **C16**: entry / long proof; **C17**: entry / long proof.

Proof of theorems 12.10 to 12.12 and proposition 12.13. At a fixed $t_0 > 0$, C4 gives the positive first genealogy jet and the tagged source has the expansion $q = 2u_y(0,0,t_0)^2 y + O(|X|^2)$. Since the affine Hodge-rate coefficient on $J \Subset (0,1/2)$ is $5a(1-2s)/(1+s)$, the exact C12 dual-moment identity yields the strict C14 rate gate for sufficiently small x; the fresh Hodge mass is $O(\ell^2)$, so Hodge dominance is strict as well.

On $I = [t_1, t_2]$ with $t_1 > 0$, compact classical regularity makes all Taylor remainders uniform, while the exact detector and apex-jet equations give positive uniform lower bounds for $u_y(0,0,t)$ and $q_H(t)$. One radius therefore gives the uniform C15 launch margins. The same conclusion may be carried to a candidate terminal time only if the derivative bounds, A^*, and the positive compression floor remain uniform; otherwise C16 records the corresponding continuation or Hodge-dominance exit.

Finally the C15 launch inequalities have fixed positive slack. Uniform continuity of the transported cohort moments and the local Euler fields on a compact classical interval preserves half that slack for one time $\tau_* > 0$ independent of the launch time. Hence the thresholded C17 protocol has gaps at least τ_* and cannot form a finite-time Zeno sequence while the compact regular positive-compression hypotheses remain valid. □

25.5. **D. Adverse ancestry and continuation-capacity short proofs.**
Statements proved. Rows D1–D6 of the detailed dependency table.

Mutual pointers: **D1**: entry / long proof; **D2**: entry / long proof.

Proof of propositions 13.1 and 13.2. Differentiating the swirl equation in y gives $\mathcal{D}_{xy}p = 2v_y\Gamma_x + 2(v - g - yg_y)p$. The positive integrating factor μ therefore gives the exact D1 characteristic formula; sign preservation of u and the negative-part inequality yield the explicit old/cross-shear majorants (13.4)–(13.6).

For D2, direct differentiation of the fixed seed shows $u_{0,y} \geq 0$ on $0 \leq y \leq \sigma_-$, hence $\mathcal{N}_0 \subset \{y \geq \sigma_-\}$ and its critical exposure is finite by rapid decay. Inside the near-affine corridor, logarithmic vertical tangents and the exact Hodge response give the all-age estimate $|\partial_t\mathcal{R}| \leq K_\delta a/(x_\tau^2 s_\tau^3)$ and $|\Theta\mathcal{R}_0| \leq K_0\Xi_\beta$. The exact transport equation for Ξ_β yields (13.15), and integration gives (13.14)–(13.16). A first corridor failure is a failure of the protected value/first-jet/tangent controls and is therefore routed to the existing restart ledger. □

Mutual pointers: **D3**: entry / long proof; **D4**: entry / long proof; **D5**: entry / long proof; **D6**: entry / long proof.

Proof of propositions 14.2, 14.4 and 14.5 and theorem 14.6. The raw source-gradient equations are

$$\mathcal{D}_{xy}q = \mathsf{d}_q q + (4yv_y)C_\beta, \qquad \mathcal{D}_{xy}C_\beta = -(xg_x)q + \left(\tfrac{3}{2}\mathfrak{A} + \mathfrak{J}\right)C_\beta.$$

Taking absolute values along characteristics gives the L^∞ estimate. For carrier norms, the density $\mathfrak{r}^{-1}$ satisfies $\partial_t(\mathfrak{r}^{-1}) + \nabla\cdot(\mathbf{b}\mathfrak{r}^{-1}) = \delta_c\mathfrak{r}^{-1}$. Multiplication by $x^\mu y^\nu$ therefore gives the Dini inequalities (14.10)–(14.11) and their Gronwall consequences.

Since $\mathcal{D}_{xy}\sqrt{xy} = (g - v)\sqrt{xy}$ and $\mathcal{D}_{xy}H = -gH + 2q$, their product gives $\mathcal{D}_{xy}M_H = -vM_H + 2\sqrt{xy}q$; the exact carrier defect then yields (14.15). The physical weight $r^{-4/3}|z|^{1/3}$ is an A_2 weight, so weighted Calderón–Zygmund theory for the whole-space Biot–Savart law gives (14.17). Finally the standard Euler H^m estimate and Gronwall propagate every fixed

high Sobolev norm under finite physical Lipschitz action. The physical Cauchy formula plus finite displacement and uniform flow derivatives propagate the rapid vorticity tail; a far-field Biot–Savart split gives (14.21) and closes the aspect-weighted second-jet gate. The axis identity $a = -U_r^r(0,0)$ gives (14.22). □

25.6. **E. Response and affine-benchmark short proofs.**
Statements proved. Rows E1–E7 of the detailed dependency table.
Mutual pointers: **E1**: entry / long proof; **E2**: entry / long proof.

Proof of proposition 15.2 and theorem 15.3. Because $\nabla\cdot(-2xv, 2yg) = g$, the exact equation $\mathcal{D}_{xy}H + gH = 2q$ is the conservative law $\partial_t H + \nabla\cdot(\mathbf{b}H) = 2q$. Since the fixed customer has $H_0 = 0$, push-forward by $\mathscr{X}_{t,\tau}$ gives $a = (3/4)\int_0^t \int q(X,\tau)K_H(\mathscr{X}_{t,\tau}X)dXd\tau$. Now $q = \Theta_y$, the material equator has $\Theta = 0$, and rapid decay removes the upper boundary, so integration by parts in the deposition coordinate gives $\mathcal{R} = -\partial_{y_\tau}K_H(\mathscr{X}_{t,\tau}X)$. Direct differentiation at zero age gives (15.5).

For E2 put $s_0 = y_\tau/x_\tau$ and $s(\xi) = y(\xi)/x(\xi)$. If $m = D_X\mathscr{X}_{\xi,\tau}(X_\tau)e_y$, use the namespace-safe logarithmic sensitivities $\zeta_1 = y_\tau m_1/x$ and $\zeta_2 = y_\tau m_2/y$. Their exact tangent system and the corridor give $|\zeta_1| + |\zeta_2| \le e^{\delta\Lambda}$. Writing $K_H = x^{-1}f(s)$, $f(s) = \sqrt{s}(1+s)^{-5/2}$, one has

$$\mathcal{R} = -\frac{\zeta_1 A_*(s) + \zeta_2 B_*(s)}{y_\tau x}, \qquad |A_*| + |B_*| + |sA_*'| + |sB_*'| \le \mathfrak{c}s^{-2} \quad (s \ge 1).$$

The corridor also gives $x \ge x_\tau e^{-(2+2\delta)\Lambda}$ and $s \ge s_0 e^{(6-2\delta)\Lambda}$. Hence $|\mathcal{R}| \le \mathfrak{c}(x_\tau^2 s_0^3)^{-1}e^{(-10+7\delta)\Lambda}$. Differentiating the displayed response formula, using the same tangent system and $|\dot{s}| \le Cas$, gives the identical exponential factor with one extra $a(t)$. Finally $e^{-2\Lambda} = \mathfrak{t}_\Psi(t)/\mathfrak{t}_\Psi(\tau)$, which is exactly (15.6). □

Mutual pointers: **E3**: entry / long proof; **E4**: entry / long proof.

Proof of lemma 15.4 and theorem 15.5. Insert the explicit fixed w_0 into $q_0 = (u_0^2)_y$, split the base–base, cross, and annuity–annuity pieces, and integrate each monomial with $\int_0^\infty x^m e^{-\lambda x}dx = m!/\lambda^{m+1}$. Their common positive denominator leaves $81\eta\Pi_{15}(\eta)$; the long proof prints an exact $\mathbb{Q}(\sqrt{2},\sqrt{3},\sqrt{7})$ positive-basis certificate whose rational component is strictly positive in every degree. Hence $\mathcal{L}_q(\eta) > 0$ for $\eta > 0$. For the exact affine conveyor, change initial line angle by $\sigma = e^{6\Lambda}s_0$ to obtain the normalized ratio $\upsilon_-^{\mathrm{aff}} = I_-/I_+$ from (15.8). Thus $I_+ - I_- = \int k(\sigma)\mathcal{L}_q(\sigma e^{-6\Lambda})d\sigma > 0$. The explicit seed also gives $\mathcal{L}_q^+(\eta) = \kappa\eta + O(\eta^2)$ at the equator and $\int_0^\infty \mathcal{L}_q^-(\eta)\eta^{-2}d\eta < \infty$ by the D2 separation. Consequently, with $\epsilon = e^{-6\Lambda}$, $I_+ = \kappa\epsilon\log(1/\epsilon) + O(\epsilon)$ and $I_- = c_-\epsilon + o(\epsilon)$, so the ratio tends to zero. Continuity on compact age intervals and the strict pointwise inequality then give one global $\upsilon_{\mathrm{aff}}^* < 1$. □

Mutual pointers: **E5**: entry / long proof; **E6**: entry / long proof; **E7**: entry / long proof.

Proof of theorems 15.6 and 15.8 and proposition 15.7. For E5 pull both source-polarized genealogy banks back to initial labels. If $M = \mathcal{K}_{\mathrm{cent}}([0,t])$, the true characteristic coordinates and flow Jacobian differ multiplicatively from the same-clock affine conveyor by $e^{O(M)}$, while the exact source-gradient fundamental matrix gives $|Z_{\mathrm{sg}}(\mathscr{X}_t X, t) - Z_{\mathrm{sg},0}(X)| \le (e^M - 1)|Z_{\mathrm{sg},0}(X)|$. The logarithmic derivatives of K_H are uniformly bounded. On a compact affine-age range the fixed-seed weighted condition number is finite, so $|a^\pm - a_{\mathrm{aff}}^\pm| \le \varepsilon(M)a_{\mathrm{aff}}^+$ with $\varepsilon(M) \to 0$. Choosing M below a fixed threshold makes $(\upsilon_{\mathrm{aff}}^* + \varepsilon)/(1-\varepsilon) < \upsilon_*$, excluding a first hit. Thus definite centered deformation (or a named comparison/continuation exit) is unavoidable; the generated-negative bank is a finer routing of this source-gradient cost.

For E6 the terminal-uniform Taylor laws give $q(x,sx,t) = 2p(t)^2 sx + O_J(x^2)$ with $\inf p > 0$. Hence $q \geq c_J x > 0$ on one fixed tube. The same Taylor bounds give $|C_\beta| \leq c_J x$ and bounded v_y, so $e_- = [-(4yv_y)C_\beta]_+ \leq c_J xq \leq c_J \ell q$. The exact residence multiplier is at most $\mathfrak{R}_* = (T - t_1)e^{2\mathcal{K}_{vx}}$ on the finite-response branch. Insert these bounds into the positive two-time response and use the exact a^+ genealogy formula to obtain $\mathfrak{X}^H_{-,\mathrm{tube}} \leq c_J \ell \mathfrak{R}_* a^+$.

For E7 apply Kato to the first source-gradient equation, propagate the old initial negative part and the generated adverse source with the exact positive integrating factor, and use $\mathsf{d}_q + g = 2(v - xv_x)$ to obtain $a^- \leq \mathfrak{X}^H_{-,\mathrm{old}} + \mathfrak{X}^H_-$. Split $\mathfrak{X}^H_- = \mathfrak{X}^H_{-,\mathrm{tube}} + \mathfrak{X}^H_{-,\mathrm{out}}$ and subtract the old and tube allowances. This gives (15.17). The conclusion is exterior/non-apex; compact mesoscopic localization is the separate F8 step. □

25.7. **F. Rotor and mesoscopic-localization short proofs.**
Statements proved. Rows F1–F8 of the detailed dependency table.
Mutual pointers: **F1**: entry / long proof; **F2**: entry / long proof; **F3**: entry / long proof; **F4**: entry / long proof.

Proof of propositions 16.1, 16.3 and 16.5 and theorem 16.4. Differentiate $q = \Theta_y$ and $C_\beta = (\Theta + \frac{x}{2}\Theta_x)/y$ along the exact material flow; use incompressibility and the mixed-jet identities to obtain the matrix (16.2). Since $\mathcal{D}_{xy}\sqrt{s} = (v+g)\sqrt{s}$, the change $\widehat{C}_\beta = 2\sqrt{s}C_\beta$ gives the rotor matrix (16.4); the two H off-diagonal terms are $\mp\omega_H$. Multiplying the two equations by q and $\widehat{C}_\beta$ and adding cancels those skew terms, proving (16.6). Finally $\Theta = \varepsilon_\beta \Gamma^2/x^2$ gives (16.7), and differentiation of the passive-scalar identity $\Gamma(\mathscr{X}_{t,0}X_0, t) = \Gamma_0(X_0)$ gives the inverse-transpose gradient formula. □

Mutual pointers: **F5**: entry / long proof; **F6**: entry / long proof; **F7**: entry / long proof; **F8**: entry / long proof.

Proof of propositions 16.6 to 16.8 and theorem 16.9. The seven-dimensional Newton representation from A5 may be differentiated twice at the origin because $H(0,0,t) = 0$ and the fixed customer is rapidly decaying. Axial spherical averaging gives the exact coefficient $15/8$ in (16.10); factoring K_H gives (16.11). Directly from $v = \chi + 2y\chi_y$, $g = 2\chi + 2x\chi_x$ and $-\mathcal{L}_7\chi = H$ one obtains $4yv_y = -yH - 2s\mathfrak{T}$. Together with $\partial_x(x^2\Theta) = 2xyC_\beta$ this proves F6.

For F7, $K_H y = \psi_{\mathrm{rot}}(s)$ and $s|\mathfrak{T}| = y|g_x - v_y|$. Passing to the carrier measure converts the relevant angular weight to $w_{\mathrm{rot}}(s) = s^{4/3}(1+s)^{-5/2}$; Cauchy–Schwarz and the lifted first-jet bound give (16.20), whose tail coefficient vanishes at both angular ends.

For F8, rederive on $[0, t_*]$ the positive visibility comparison and the finite residence bound; this avoids relying on any later response theorem as a black box. The E7 master inequality and (16.21) force a positive generated-negative response of size $(v_* - v_{\mathrm{old}})a^+(t_*)$. Choose angular cutoffs using F7. On the remaining compact angular sector, detector positivity and compact-time regularity give one protected small-x tube all the way back to $t = 0$; passive circulation plus rapid decay makes the far-x response uniformly small. A fixed positive fraction therefore remains on (16.22). The routing kernel is bounded above there, and $[(\omega_H + \sqrt{s}\mathfrak{T})\widehat{C}_\beta]_+ = e_-$, which yields (16.23). Each deposition pair is integrated once, so the ticket is one-shot. □

25.8. **G. Level-set polarization and deformation-stability short proofs.**
Statements proved. Rows G1–G5 of the detailed dependency table.
Mutual pointers: **G1**: entry / long proof; **G2**: entry / long proof.

Proof of theorem 17.1 and proposition 17.2. For the fixed customer, $\Gamma = xu \geq 0$ and $q = 2\varepsilon_\beta \Gamma\Gamma_y/x^2$. Coarea on the regular superlevels $\Omega_c(\tau)$ gives the two formulas for $\mathcal{S}_\pm(t,\tau)$. Since the outward normal of Ω_c is $-n_c$, the divergence theorem sends the signed boundary flux to $-\partial_y(W_{t,\tau}/x^2) = \mathcal{R}(t,\tau;\cdot)/x^2$, giving the signed interior formula. Integrating in source time

recovers $a^{\pm}(t)$. Finally $a = a^+ - a^-$ and $a^+ + a^-$ show, for $0 < \pi_* < 1$, $a \geq \pi_*(a^+ + a^-)$ iff $a^- \leq (1-\pi_*)/(1+\pi_*)\, a^+$. The common factor $3\varepsilon_\beta/2$ is retained in the level-set form and is never divided out at the provenance gate. □

Mutual pointers: **G3**: entry / long proof; **G4**: entry / long proof; **G5**: entry / long proof.

Proof of lemma 17.4. On compact affine-age intervals the numerator and denominator in $\mathfrak{C}_Z$ are continuous and the denominator is positive. For large age, write $y = \eta x$, set $\epsilon = e^{-6\Lambda}$, and use the fixed-seed equatorial expansions $\mathcal{L}_{|q|}(\eta) = \kappa_q \eta + O(\eta^2)$ and $\mathcal{L}_Z(\eta) = \kappa_Z \eta + O(\eta^2)$. The affine Hodge factor gives

$$\mathfrak{d}_Z(\Lambda) = \epsilon^{2/3} \int_0^\infty \frac{\sqrt{\sigma}}{(1+\sigma)^{5/2}} \mathcal{L}_{|q|}(\epsilon\sigma)\, d\sigma$$

and the identical formula with $\mathcal{L}_Z$ for the numerator. Since $\sigma^{3/2}(1+\sigma)^{-5/2} \sim \sigma^{-1}$, both stocks have the essential logarithmic asymptotics $e^{-10\Lambda}(6\kappa\Lambda + O(1))$. Their common logarithm cancels, giving $\mathfrak{C}_Z(\Lambda) \to \kappa_Z/\kappa_q < \infty$. Compactness then gives the global bound. No dominated-convergence argument across this logarithmic endpoint is used. □

Mutual pointers: **G3**: entry / long proof; **G4**: entry / long proof; **G5**: entry / long proof.

Proof of theorem 17.5 and corollary 17.6. Pull the exact signed and absolute Hodge responses back to the fixed initial labels. The centered action controls three independent factors: the source-gradient fundamental matrix, the material Jacobian, and the final Hodge visibility. With the namespace-safe constants $\mathfrak{c}_0, \mathfrak{c}_1$ this gives (17.10); the audited G3 bound $\mathfrak{C}_Z^* < \infty$ is exactly what converts the full source-gradient error back to the affine absolute response. Combining the error with the strict affine margin and choosing M_* so that the perturbation is less than half that margin gives (17.12).

For G5, define t_* as the infimum of strict-failure times. Every actual failure time has $\mathcal{K}_{\mathrm{cent}} > M_*$ by the contrapositive of G4; continuity of the integrated action then gives (17.13) at the infimum. Finally the audited radial-star identity $v(X) - a = \int_0^1 \mathcal{R}_v(\lambda X)\, d\lambda/\lambda$ and $g - 2a = 2(v-a) - \mathfrak{A}$ imply $\mathcal{K}_{\mathrm{cent}}([0,t]) \leq 3\int_0^t \mathcal{L}_{\mathrm{pol}}$. Hence (17.14) holds with $c_{\mathrm{pol}} = M_*/3$. □

25.9. **H. Causal adverse-response short proofs.**
Statements proved. Rows H1–H8 of the detailed dependency table.

Mutual pointers: **H1**: entry / long proof; **H2**: entry / long proof; **H3**: entry / long proof.

Proof of propositions 18.1 to 18.3. Differentiate $\log K_H$ along the material flow to obtain (18.1); its two angular coefficients are uniformly bounded, so integration gives (18.3). Insert this comparison and the finite residence multiplier into the Hodge-weighted adverse source. The HC_β part is controlled by Cauchy–Schwarz in the carrier measure using $\mathcal{E}_{H,1}\mathcal{E}_{\mathrm{sg}}^{(0,0)}$, while the signed-shear part is bounded by $\mathcal{L}_B \mathcal{Q}_{\mathrm{sg}}^{(1)}$, giving the raw response-capacity estimate. The apex Taylor expansions from the relaunch theorem give $H = O(x)$, $C_\beta = O(x)$, $\mathfrak{T} = O(x)$ and $q \geq cx$ on a fixed conical tube; substitution into the exact adverse exchange yields $[-(4yv_y)C_\beta]_+ \leq \mathfrak{c}\ell q$. □

Mutual pointers: **H4**: entry / long proof; **H5**: entry / long proof; **H6**: entry / long proof.
Mutual pointers: **H7**: entry / long proof; **H8**: entry / long proof.

Proof of H4–H8. H4 applies Kato to the first source-gradient equation and combines the material Jacobian with $\mathfrak{d}_q + g = 2(v - xv_x)$ to obtain the exact positive residence multiplier and the old/generated response split. For H6, use the rotor form and the global identity

$\sqrt{xy}\,\widehat{C}_\beta = 2\Theta + x\Theta_x$ to dominate the entire $H^- \to H^-$ source by one cooperative two-component characteristic system; no mesoscopic indicator is inserted. Keeping final Hodge visibility inside this system yields a scalar second-order causal inequality whose constant-coefficient Green function is $\mathcal{G}_{\mathrm{rec}}$. H5 follows because $\mathcal{G}_{\mathrm{rec}}(u)/u$ is bounded near zero and the transported positive Hodge stock already present at source time is at most $a^+(t)$, producing the h^2 gain. Splitting the H6 primary input into the signed $\mathfrak{T}$ part and the global H^+ part gives H7; H8 is the direct strict-gap consequence of that three-account inequality. □

25.10. **I. Supporting exact-identity short proofs.**
Statements proved. Rows I1–I7 of the detailed dependency table.
Mutual pointers: **I1**: entry / long proof; **I2**: entry / long proof.

Proof of propositions 19.1 and 19.2. The visibility law follows by differentiating $K_H = x\sqrt{y}(x+y)^{-5/2}$ along $\dot{x} = -2xv$, $\dot{y} = 2yg$. Differentiating the exact source-response formula for $a(t)$ in observation time gives the zero-age boundary term plus the history term $\Theta\partial_t\mathcal{R}$. Splitting each routing kernel into positive and negative parts produces the guarded source-payment identity; the explicit factor Θ keeps the source ancestry visible throughout. □

Mutual pointers: **I3**: entry / long proof; **I4**: entry / long proof; **I5**: entry / long proof.

Proof of propositions 19.3 to 19.5. The cross-shear factorization first uses $C_\beta = \varepsilon_\beta u\Gamma_x/y$. For the fixed customer, passive transport preserves $\Gamma \geq 0$, hence $\varepsilon_\beta u \geq 0$ and the positive-part polarization is legitimate. Writing $y = sx$ gives $K_H = x^{-1}\sqrt{s}(1+s)^{-5/2}$ and $\mathfrak{r} = xs^{1/3}$; maximizing the resulting angular factor gives the carrier-Hodge comparison. The fresh genealogy formula is simply Duhamel's formula for the continuity equation for H restricted to source times $(t-h)_+ \leq \tau \leq t$; therefore it is an actual sub-history of the true Euler genealogy, not an auxiliary memory field. □

Mutual pointers: **I6**: entry / long proof; **I7**: entry / long proof.

Proof of propositions 19.6 and 19.7. For no-double-charging, take the explicit two-bank ledger $a^+ = 1$, $a^- = t$, $a = 1-t$ on $0 \leq t < 1$. The total adverse creation is one, yet the half-Hodge refresh protocol has infinitely many contacts at $t_n = 1 - 2^{-n}$. Thus a single physical conversion cannot be charged once per later refresh. For strict-gap algebra, subtract a^- from a^+ and use $a^- \leq (1-\delta)a^+$ to obtain $a \geq \delta a^+$; expressing $a^+ + a^- = (2-\delta)a^+$ at the worst allowed ratio gives the stated equivalent polarization constant. □

25.11. **J. Short proofs of the four falsification firewalls.**
Statements proved. Rows J1–J4 of the detailed dependency table.
Mutual pointers: **J1**: entry / long proof.

Proof of proposition 20.1. Use the one-cell reduced exact tagged system from the archive with $v = g = v_x = g_y = 0$, $v_y = g_x = -H/4$ and no mixed-shear term. The resulting ODE has invariants $q^2 + 4C_\beta^2 = 1$ and $C_\beta = H^2/16$. Its increasing branch reaches a half-turn in finite time with finite raw cross-shear action, while the positive and negative genealogy banks become equal. Hence local exact dynamics plus finite action cannot imply a uniform strict cancellation gap. □

Mutual pointers: **J2**: entry / long proof.

Proof of proposition 20.2. Choose smooth rapidly decaying $H_0^+ = ye^{-(x+y)}$ and an H_0^- supported away from the equator with quadratic apex vanishing. The whole-space seven-dimensional Hodge operator reconstructs a regular meridional field. Choose $0 < \eta < \min\{\epsilon, 1\}$ and scale the amplitude of H_0^- so its positive Hodge integral is the fraction $1 - \eta$ of that of H_0^+. This preserves the apex sign orders and gives $a^-/a^+ = 1 - \eta > 1 - \epsilon$. Thus frozen Hodge geometry and local sign structure alone admit $a^-/a^+ \to 1$. □

Mutual pointers: **J3**: entry / long proof; **J4**: entry / long proof.

Proof of propositions 20.3 and 20.4. The first statement uses the same ledger $a^+ = 1$, $a^- = t$: the total adverse creation is finite, but selecting half of the shrinking net bank at times $1 - 2^{-n}$ creates infinitely many fractional refresh contacts. For the reproduction statement, in the constant-rate affine conveyor a label can leave the compact rotor region while its factored pointwise residence multiplier grows like $e^{2a_0 t}$. The final Hodge visibility of that same label decays like $e^{-10a_0 t}$, so their physical product decays like $e^{-8a_0 t}$. A pointwise reproduction bound after factoring out final visibility is therefore the wrong global target. □

25.12. **K. Permanent reconstructed modules M1–M8: short proofs.**

M1. Polarization/source-history structure. *Mutual pointers:* **M1**: entry / long proof.

The signed Hodge banks are written on one normalized two-copy genealogy space. The exact source-history representation and the global/local polarization formulas then separate favorable and adverse source histories without changing ancestry. This is precisely the content of theorems K.11 to K.13.

M2. Critical source-positive cashier and fractional homotopy. *Mutual pointers:* **M2**: entry / long proof.

At the repaired critical exponent the source term has the favorable sign before any normalization. The exact factorization and fractional homotopy preserve the tagged source ancestry while interpolating to the Hodge endpoint. See theorems K.14, K.15 and K.17.

M3. Post-ticket causal pay-or-route module. *Mutual pointers:* **M3**: entry / long proof.

The post-threshold genealogy is split into old and newly deposited source histories. The three-gate alternative routes every failure either to an already named payment or to a later module, and the no-double-charge rule prevents one physical conversion from being billed twice. See theorems K.19 and K.20.

M4. Gate-S / critical-to-Hodge source-angle bridge. *Mutual pointers:* **M4**: entry / long proof.

The normalized Gate-S covariance and packet pay-or-route identities convert critical source-angle loss into Hodge-visible source geometry or a named rotor/deformation invoice. The source-angle invoice is therefore a tagged GREEN transfer, not an untagged background estimate. See theorems K.21 to K.23 and K.25.

M5. Ratio-free GREEN barrier and bounded-age crossing. *Mutual pointers:* **M5**: entry / long proof.

The two-weight GREEN comparison removes forbidden bank ratios from the source-angle barrier. Large relative collapse age is retired by the established response decay, leaving only the bounded-age core for M6. See theorems K.27 to K.29.

M6. Global continuation/source-potential/aspect/tangent audit. *Mutual pointers:* **M6**: entry / long proof.

The physical Lipschitz clock, transported source potential, aspect transport, aspect capital, and tangent sign-flip estimates exhaust the bounded-age source-angle routes. The remaining inherited launch geometry is exactly the source-angle stock closed by the completed O6.1 audit, which then funds O4. See theorems K.30, K.31 and K.33 to K.36.

M7. Stronger fixed-margin terminal mechanisms. *Mutual pointers:* **M7**: entry / long proof.

Each of the powered-reserve, positive-bank exponential-growth, and fixed sub-quarter maintained-cashier hypotheses independently gives a finite physical-time deadline. These remain convenient sufficient routes but are stronger than the exact terminal interface. See theorems K.1, K.3 and K.4.

M8. Exact integral cashier countdown. *Mutual pointers:* **M8**: entry / long proof.

After O4, the maintained cashier satisfies $\Sigma \geq \Sigma(t_1)e^{\mathcal{G}_{\mathrm{cash}}}$ and $a \geq \delta_*\Sigma$. Since $d\Lambda = a\,dt$, integration gives the exact bound $dt/d\Lambda \leq [\delta_*\Sigma(t_1)]^{-1}e^{-\mathcal{G}_{\mathrm{cash}}}$. Thus $e^{-\mathcal{G}_{\mathrm{cash}}} \in L^1(d\Lambda)$ is sufficient, and a vanishing quarter-rate margin of order $p/(1+\Lambda)$ with $p > 1$ is one useful corollary. See theorems K.5 and K.6.

25.13. **K. O-ledger short-status cards.** O1–O3 are proved and keep their permanent navigation IDs. The inherited-launch source-angle audit O6.1 is GREEN/COMPLETE, O4 is GREEN, O5 is GREEN from O4+O6.2+M8, and O6.2 is the sole OPEN proposition.

O1. Large-collapse-age physical-bank retirement. **Status: GREEN/COMPLETE.** The localized G4 comparison gives the exact physical large-age bank estimate (21.4); see grand-finale entry, theorem 21.3, and long proof.

O2. Bounded-collapse-age physical-bank return. **Status: GREEN/COMPLETE.** Source-time-localized G4 gives the strict bounded-age bank ratio (21.8), while failure of the small-centered-action branch pays (21.9); see grand-finale entry, theorem 21.4, and long proof.

O3. Physical-bank consolidation. **Status: GREEN/COMPLETE.** O1 and O2 partition the exact negative source-time bank. Absorbing the localized O1 comparison error gives either $a^- \leq \theta_{\mathrm{O3}}a^+$ with $\theta_{\mathrm{O3}} < 1$, or the already named centered-deformation ticket (21.14); see grand-finale entry, theorem 21.5, and long proof.

O4. Grand cancellation gap. **Status: GREEN/COMPLETE.** The inherited-launch source-angle audit is closed. The first O3 threshold is defined by

$$t_d = \inf\{t > 0 : \mathcal{K}_{\mathrm{cent}}([0,t]) > M_{\mathrm{O3}}\}.$$

Continuity gives $\mathcal{K}_{\mathrm{cent}}([0,t_d]) = M_{\mathrm{O3}}$, so O3 covers the threshold state. A first-contact argument using the active positive opening and proposition 11.1 shows that $a(t)$ cannot reach zero. Thus O3 supplies the pre-threshold gap and O6.1 the post-threshold gap, with

$$\delta_* = \min\{1 - \theta_{\mathrm{O3}}, \delta_6\} > 0.$$

No M7/O5 hypothesis is used and no pre-threshold O3 expenditure is recharged. See grand-finale entry, theorem 21.6, and long proof.

O5. Terminal noncontinuation from O4 and O6.2. **Status: GREEN/COMPLETE.** Assume O4 and the exact terminal subclause O6.2. Apply theorem K.5. The powered, exponential-bank-growth, and fixed sub-quarter cashier gates remain stronger sufficient corollaries, but no fixed positive collapse-age margin is required. See grand-finale entry, theorem 21.7, and long proof.

O6.2. Post-O4 integral cashier closure. **Status: SOLE OPEN PROPOSITION—NO PROOF ASSERTED.** The permanent external label is unchanged. The inherited-launch source-angle audit formerly denoted O6.1 is GREEN/COMPLETE and already funds O4. O6.2 is the exact post-O4 obligation (21.18): construct a maintained cashier whose net gain satisfies $e^{-\mathcal{G}_{\mathrm{cash}}} \in L^1(d\Lambda)$. M7-P/C1/C2 remain stronger sufficient mechanisms; M8 is the exact proved countdown used by O5. See grand-finale entry, open proposition 21.8, and long-proof slot.

26. Compact ledger glossary

Accounting bridge. For future work, the glossary fixes the translation between Peter's accounting words and the mathematical quantities, so later edits cannot change the meaning of an account halfway through the audit.

Ledger word	Mathematical quantity	Meaning
Customer	$u_0 = yw_0$	the single fixed initial swirl datum
Tag	ε_β	Boolean source ancestry, not a small parameter
Source	Θ, q, C_β	direct swirl-generated descendants
Cashier	Hodge map $H \mapsto a$	converts genealogy into apex compression
Positive bank	H^+, a^+	cumulative favorable genealogy
Negative bank	H^-, a^-	cumulative adverse genealogy
Clock	$\psi_\beta, \Psi, \mathfrak{t}_\Psi$	signed jet, quadratic detector, and exact collapse clock
Reserve family	$\mathcal{F}_\kappa = Q - 3(1+\kappa)a^2$	M7-P powered terminal family; $\mathcal{F}_\Psi$ is $\kappa = 2$
Withdrawal	$\mathcal{M}_-$, adverse response	loss or cancellation of cashier credit
Debt	$\mathfrak{A}, \mathfrak{J}, H, \mathfrak{T}$ actions	non-affine restart cost
Affine audit	$\upsilon_-^{\mathrm{aff}}$	fixed-seed no-false-gain benchmark
Conversion ticket	mesoscopic rotor integral	cost of leaving the affine polarization regime
Recycling	causal $\mathcal{G}_{\mathrm{rec}}$	finite amplification of already created negative genealogy
O6.1	inherited extreme-aspect / near-horizontal source-angle stock	GREEN/COMPLETE launch-state audit funding O4
M8	$e^{-\mathcal{G}_{\mathrm{cash}}} \in L^1(d\Lambda)$ countdown	exact proved terminal mechanism used by O5
O6.2	construct a maintained cashier satisfying the M8 integrability gate	sole open post-O4 proposition

27. Conclusion

The framework is organized around two source-specific objects: the swirl polarizer

$$\Phi = \beta = \tfrac{1}{r} U^\phi$$

and the local quadratic detector

$$\Psi(t) = (\beta_{zz}(0,0,t))^2.$$

The polarizer separates the nontrivial swirl sector from the globally regular no-swirl shadow, while the ancestry projector prevents later accounts from borrowing a mechanism that survives when the polarizer is switched off. The detector probes a single apex jet rather than a global spatial supremum.

The detector is nevertheless tied exactly to the vorticity hierarchy:

$$\partial_z^2 \omega^z(0,0,t) = 2\beta_{zz}(0,0,t), \qquad \boxed{\|\nabla^2 \omega(t)\|_{L^\infty(r\,dr\,dz)} \geq 2\sqrt{\Psi(t)}.}$$

Thus Ψ is a genuinely local second-vorticity-jet observable. This is distinct from the BKM quantity $\|\omega(t)\|_\infty$, which is a global zeroth-order spatial supremum. Neither $\|\omega\|_\infty$ nor $\|\nabla U\|_\infty$ universally dominates $\sqrt{\Psi}$ without additional scale information, and $\|\Delta^2\omega\|_\infty$ is not a universal substitute for the Hessian bound.

The proof architecture follows the source-specific chain

$$\begin{aligned}\text{nonzero swirl polarization} &\longrightarrow \text{tagged source} \longrightarrow \text{signed Hodge genealogy}\\ &\longrightarrow \text{apex compression} \longrightarrow \text{local countdown clock.}\end{aligned}$$

The remaining modules quantify restart geometry, source-angle transfer, global GREEN coupling, aspect/tangent deformation, and three terminal mechanisms. The unresolved mathematical issue is therefore not a generic strengthening of the BKM criterion; it is the source-specific closure that connects the nonzero polarizer to the local detector through the audited genealogy accounts.

The Grand Finale remains conditional only on O6.2, the explicit post-O4 integral cashier closure package. Accordingly, the paper does not claim an unconditional finite-time singularity theorem for smooth three-dimensional Euler flow with swirl.

28. Acknowledgement

This work was conceived, initiated, and directed by the author. The author formulated the central mathematical questions, determined the proof strategy, selected the source-specific quantities, Hodge cashiers, response kernels, comparison regimes, and accounting interpretation, and decided which proposed reductions and restart mechanisms were mathematically admissible. In particular, the author required a strict separation between identities that survive in the swirl-free Euler background and mechanisms generated by nontrivial swirl, and adopted the provenance firewall used throughout the manuscript. The author is responsible for all mathematical claims, interpretations, and the final form of the paper.

During the development of the project, the author used OpenAI's ChatGPT as a research assistant across an extended sequence of working sessions, including GPT-5.6 Sol in September 2026. ChatGPT assisted with exploratory calculations, alternative proof routes, countermodels and stress tests for tentative arguments, literature and scope checks, dependency and circularity audits, notation checks, and the organization and editing of the manuscript. All AI-assisted material was reviewed, selected, and revised by the author before inclusion.

The author also thanks Prof. Zixiang Zhou for helpful discussions.

References

[1] Bardos C and Titi E S 2007 Euler equations for incompressible ideal fluids *Uspekhi Mat. Nauk* **62** 5–46

[2] Beale J T, Kato T and Majda A 1984 Remarks on the breakdown of smooth solutions for the 3-D Euler equations *Commun. Math. Phys.* **94**(1) 61–66 doi:10.1007/BF01212349

[3] Beals R, Sattinger D and Szmigielski J 2001 Inverse scattering solutions of the Hunter–Saxton equations *Appl. Anal.* **78** 255–269

[4] Caffarelli L, Kohn R and Nirenberg L 1982 Partial regularity of suitable weak solutions of the Navier–Stokes equations *Commun. Pure Appl. Math.* **35** 771–831

[5] Camassa R and Holm D D 1993 An integrable shallow water equation with peaked solitons *Phys. Rev. Lett.* **71** 1661–1664

[6] Cannon J R and DiBenedetto E 1980 The initial problem for the Boussinesq equations with data in L^p in *Lecture Notes in Mathematics* **771** 129–144 (Springer, Berlin)

[7] Chae D and Nam H-S 1997 Local existence and blow-up criterion for the Boussinesq equations *Proc. Roy. Soc. Edinburgh Sect. A* **127**(5) 935–946 doi: 10.1017/S0308210500026810

[8] Chae D, Kim S-K and Nam H-S 1999 Local existence and blowup criterion of Hölder continuous solutions of the Boussinesq equations *Nagoya Math. J.* **155** 55–80

[9] Chae D and Lee J 2002 On the regularity of the axisymmetric solutions of the Navier–Stokes equations *Math. Z.* **239** 645–671

[10] Chae D 2006 Global regularity for the 2D Boussinesq equations with partial viscosity terms *Adv. Math.* **203** 497–513

[11] Chae D 2008 On the blowup problem for the axisymmetric 3D Euler equations *arXiv:0803.1784*

[12] Chae D 2010 On the Lagrangian dynamics of the axisymmetric 3D Euler equations *J. Differential Equations* **249** 571–577

[13] Chae D, Constantin P and Wu J 2014 An incompressible 2D didactic model with singularity and explicit solutions of the 2D Boussinesq equations *J. Math. Fluid Mech.* **16** 473–480 doi:10.1007/s00021-014-0166-5

[14] Chen H, Fang D and Zhang T 2015 Regularity of 3D axisymmetric Navier–Stokes equations *arXiv:1505.00905*

[15] Chen W and Sarria A 2016 Damped infinite energy solutions of the 3D Euler and Boussinesq equations *arXiv:1605.08965*

[16] Chen J and Hou T Y 2021 Finite time blowup of 2D Boussinesq and 3D Euler equations with $C^{1,\alpha}$ velocity and boundary *Commun. Math. Phys.* **383**(3) 1559–1667 doi:10.1007/s00220-021-04067-1

[17] Chen J and Hou T Y 2022 Stable nearly self-similar blowup of the 2D Boussinesq and 3D Euler equations with smooth data I: Analysis *arXiv:2210.07191*

[18] Choi K, Kiselev A and Yao Y 2015 Finite time blow up for a 1D model of 2D Boussinesq system *Commun. Math. Phys.* **334** 1667–1679 doi:10.1007/s00220-014-2146-2

[19] Choi K, Hou T Y, Kiselev A, Luo G, Šverák V and Yao Y 2017 On the finite-time blowup of a one-dimensional model for the three-dimensional axisymmetric Euler equations *Commun. Pure Appl. Math.* **70**(11) 2218–2243 doi:10.1002/cpa.21697

[Collot–Merle–Raphaël(2017)] C. Collot, F. Merle, and P. Raphaël, Stability of ODE blow-up for the energy critical semilinear heat equation, *C. R. Math. Acad. Sci. Paris* **355** (2017), no. 1, 65–79. doi: 10.1016/j.crma.2016.10.020.

[20] Constantin P, Lax P D and Majda A 1985 A simple one-dimensional model for the three-dimensional vorticity equation *Commun. Pure Appl. Math.* **38**(6) 715–724 doi:10.1002/cpa.3160380605

[21] Constantin P 1986 Note on loss of regularity for solutions of the 3D incompressible Euler and related equations *Commun. Math. Phys.* **104** 311–326

[22] Constantin P, Majda A and Tabak E 1994 Singular front formation in a model for quasigeostrophic flow *Phys. Fluids* **6** 9–11

[23] Constantin P, Majda A and Tabak E 1994 Formation of strong fronts in the 2-D quasigeostrophic thermal active scalar *Nonlinearity* **7** 1495–1533

[24] Constantin P 2000 The Euler equations and nonlocal conservative Riccati equations *Int. Math. Res. Not.* **2000**(9) 455–465 doi:10.1155/S1073792800000258

[25] Constantin P 2007 On the Euler equations of incompressible fluids *Bull. Amer. Math. Soc. (N.S.)* **44**(4) 603–621 doi:10.1090/S0273-0979-07-01184-6

[26] Córdoba D, Fefferman C and De La Llave R 2004 On squirt singularities in hydrodynamics *SIAM J. Math. Anal.* **36** 204–213

[27] Córdoba A, Córdoba D and Fontelos M A 2005 Formation of singularities for a transport equation with nonlocal velocity *Ann. Math.* **162** 1377–1389

[28] Córdoba D and Martínez-Zoroa L 2023 Blow-up for the incompressible 3D Euler equations with uniform $C^{1,\frac{1}{2}-\epsilon}$ force *arXiv:2309.08495*

[29] Córdoba D, Martínez-Zoroa L and Zheng F 2023 Finite time singularities to the 3D incompressible Euler equations for solutions in $C^\infty(\mathbb{R}^3 \setminus \{0\}) \cap C^{1,\alpha} \cap L^2$ *arXiv:2308.12197*

[30] De Gregorio S 1990 On a one-dimensional model for the three-dimensional vorticity equation *J. Stat. Phys.* **59** 1251–1263 doi: 10.1007/BF01334750

[31] De Gregorio S 1996 A partial differential equation arising in a 1D model for the 3D vorticity equation *Math. Methods Appl. Sci.* **19** 1233–1255

[32] Doering C R, Wu J, Zhao K and Zheng X 2023 Long time behavior of the two-dimensional Boussinesq equations without buoyancy diffusion *Nonlinearity* **36** 39 doi: 10.1088/1361-6544/acb81c

[33] Drazin P G and Riley N 2006 *The Navier–Stokes Equations: A Classification of Flows and Exact Solutions* (Cambridge University Press, Cambridge) doi: 10.1017/CBO9780511755499
[34] Drivas T D and Elgindi T M 2023 Singularity formation in the incompressible Euler equation in finite and infinite time *EMS Surv. Math. Sci.* **10**(1) 1–100 doi:10.4171/EMSS/66
[35] E W and Shu C 1994 Small scale structures in Boussinesq convection *Phys. Fluids* **6** 48–54
[36] Elgindi T M and Jeong I-J 2016 Symmetries and critical phenomena in fluids *arXiv:1610.09701*
[37] Elgindi T M and Jeong I-J 2017 On the effects of advection and vortex stretching *arXiv:1701.04050*
[38] Elgindi T M and Jeong I-J 2019 Finite-time singularity formation for strong solutions to the axi-symmetric 3D Euler equations *Ann. PDE* **5**(2), Paper No. 16, 51 pp. doi:10.1007/s40818-019-0071-6
[39] Elgindi T M and Jeong I-J 2020 Finite-time singularity formation for strong solutions to the Boussinesq system *Ann. PDE* **6** 50 (see also *arXiv:1708.02724*)
[40] Elgindi T M 2021 Finite-time singularity formation for $C^{1,\alpha}$ solutions to the incompressible Euler equations on $\mathbb{R}^3$ *Ann. Math.* **194** 647–727
[41] Elgindi T M and Pasqualotto F 2023 From instability to singularity formation in incompressible fluids *arXiv:2310.19780*
[42] Fefferman C L 2002 Existence and smoothness of the Navier–Stokes equation *(Clay Mathematics Institute problem description)* http://www.claymath.org/millennium/Navier-Stokes_Equations/
[43] Friedlander S and Pavlović N 2004 Blowup in a three-dimensional vector model for the Euler equations *Commun. Pure Appl. Math.* **57** 705–725
[44] Gibbon J D, Fokas A and Doering C R 1999 Dynamically stretched vortices as solutions of the 3D Navier–Stokes equations *Physica D* **132** 497–510
[45] Gibbon J D 2007 Ortho-normal quaternion frames. Lagrangian evolution equations and the three-dimensional Euler equations *arXiv:math-ph/0610004 arXiv:1708.02724 [math.AP]*
[46] Gibbon J D 2008 The three-dimensional Euler equations: Where do we stand? *Physica D* **237** 1894–1904
[47] Giga Y and Kohn R V 1985 Asymptotically self-similar blow-up of semilinear heat equations *Commun. Pure Appl. Math.* **38**(3) 297–319 doi: 10.1002/cpa.3160380304
[48] Giga Y 1986 A bound for global solutions of semilinear heat equations *Commun. Math. Phys.* **103**(3) 415–421
[49] Hou T Y and Li C 2005 Global well-posedness of the viscous Boussinesq equations *Discrete Contin. Dyn. Syst.* **12** 1–12
[50] Hou T Y and Li R 2006 Dynamic depletion of vortex stretching and non-blowup of the 3-D incompressible Euler equations *J. Nonlinear Sci.* **16** 639–664
[51] Hou T Y and Li C 2008 Dynamic stability of the 3D axi-symmetric Navier–Stokes equations with swirl *Commun. Pure Appl. Math.* **61** 661–697
[52] Hou T Y and Lei Z 2009 On the stabilizing effect of convection in three-dimensional incompressible flows *Commun. Pure Appl. Math.* **62** 501–564
[53] Hou T Y, Lei Z and Li C 2009 Global regularity of the 3D axi-symmetric Navier–Stokes equations with anisotropic data *arXiv:0901.3486*
[54] Hou T Y 2009 Blowup or no blowup? A unified computational and analytic approach to 3D incompressible Euler and Navier–Stokes equations *Acta Numer.* 277–346
[55] Hou T Y and Luo G 2013 On the finite-time blowup of a 1D model for the 3D incompressible Euler equations *arXiv:1311.2613*
[56] Hou T Y and Liu P 2014 Self-similar singularity of a 1D model for the 3D axisymmetric Euler equations *arXiv:1407.5740*
[57] Hou T Y, Jin T and Liu P 2017 Potential singularity for a family of models of the axisymmetric incompressible flow *J. Nonlinear Sci.* **27** 1–31
[58] Hou T Y, Liu P and Wang F 2017 Global regularity for a family of 3D models of the axisymmetric Navier–Stokes equations *arXiv:1708.07536*
[59] Hou T Y 2022 Potential singularity of the 3D Euler equations in the interior domain *Found. Comput. Math.* 1–47
[60] Hunter J K and Saxton R 1991 Dynamics of director fields *SIAM J. Appl. Math.* **51** 1498–1521
[61] Hunter J K and Zheng Y 1995 On a nonlinear hyperbolic variational equation: I. Global existence of weak solutions *Arch. Ration. Mech. Anal.* **129** 305–353

[62] Hunter J K and Zheng Y 1995 On a nonlinear hyperbolic variational equation: II. The zero-viscosity and dispersion limits *Arch. Ration. Mech. Anal.* **129** 355–383
[63] Katz N and Pavlović N 2005 Finite time blow-up for a dyadic model of the Euler equations *Trans. Amer. Math. Soc.* **357** 695–708
[64] Khenissy S and Zaag H 2011 Continuity of the blow-up profile with respect to initial data for semilinear heat equations *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **28**(1) 1–29
[65] Kiselev A and Zlatoš A 2005 On discrete models of the Euler equation *Int. Math. Res. Not.* **38** 2315–2339
[66] Kiselev A and Šverák V 2014 Small scale creation for solutions of the incompressible two-dimensional Euler equation *Ann. Math.* **180** 1205–1224
[67] Kiselev A, Ryzhik L, Yao Y and Zlatoš A 2016 Finite time singularity for the modified SQG patch equation *Ann. Math.* **184** 909–948
[68] Kiselev A, Park J and Yao Y 2022 Small scale formation for the 2D Boussinesq equation *arXiv:2211.05070*
[69] Lafleche L, Vasseur A F and Vishik M 2021 Instability for axisymmetric blow-up solutions to incompressible Euler equations *J. Math. Pures Appl.* **155** 140–154
[70] Lenells J 2007 The Hunter–Saxton equation describes the geodesic flow on a sphere *J. Geom. Phys.* **57** 2049–2064
[71] Lenells J 2008 The Hunter–Saxton equation: a geometric approach *SIAM J. Math. Anal.* **40** 266–277
[72] Lin F-H 1998 A new proof of the Caffarelli–Kohn–Nirenberg theorem *Commun. Pure Appl. Math.* **51** 241–257
[73] Liu J-G and Wang W-C 2006 Convergence analysis of the energy and helicity preserving scheme for axisymmetric flows *SIAM J. Numer. Anal.* **44** 2456–2480
[74] Liu J-G and Wang W-C 2009 Characterization and regularity for axisymmetric solenoidal vector fields with application to Navier–Stokes equation *SIAM J. Math. Anal.* **41** 1825–1850
[75] Lou G and Hou T Y 2013 Potentially singular solutions of the 3D incompressible Euler equations *arXiv:1310.0497*
[76] Lou G and Hou T Y 2014 Potentially singular solutions of the 3D axisymmetric Euler equations *Proc. Natl Acad. Sci. USA* **111** 14261–14265 doi: 10.1073/pnas.1409693111
[77] Lou G and Hou T Y 2014 Toward the finite-time blowup of the 3D axisymmetric Euler equations: a numerical investigation *Multiscale Model. Simul.* **12** 1722–1776
[78] Majda A J and Bertozzi A L 2002 *Vorticity and Incompressible Flow* Cambridge Texts in Applied Mathematics 27 (Cambridge University Press, Cambridge)
[79] Majda A J 2003 *Introduction to PDEs and Waves for the Atmosphere and Ocean* (Courant Lecture Notes in Mathematics vol 9) (Courant Institute, New York)
[80] Merle F and Raphaël P 2005 The blow-up dynamic and upper bound on the blow-up rate for critical nonlinear Schrödinger equation *Ann. of Math.* **161**(1) 157–222
[81] Morozov O I 2020 Integrability structures of the generalized Hunter–Saxton equation *arXiv:2012.06905*
[82] Ohkitani K and Gibbon J D 2000 Numerical study of singularity formation in a class of Euler and Navier–Stokes flows *Phys. Fluids* **12** 3181–3194
[83] Okamoto H, Sakajo T and Wunsch M 2008 On a generalization of the Constantin–Lax–Majda equation *Nonlinearity* **21** 2447–2461
[84] Olver P and Rosenau P 1996 Tri-Hamiltonian duality between solitons and solitary wave solutions having compact support *Phys. Rev. E* **53** 1900–1906
[85] Pumir A and Siggia E D 1992 Development of singular solutions to the axisymmetric Euler equations *Phys. Fluids A* **4** 1472–1491
[86] Raphaël P and Rodnianski I 2012 Stable blow up dynamics for the critical co-rotational wave maps and equivariant Yang–Mills problems *Publ. Math. Inst. Hautes Études Sci.* **115**(1) 1–122 doi: 10.1007/s10240-011-0037-z
[87] Sarria A 2014 Global estimates and blowup criteria for the generalized Hunter–Saxton system *arXiv:1307.4504*
[88] Schochet S 1986 Explicit solutions of the viscous model vorticity equation *Commun. Pure Appl. Math.* **39**(4) 531–537 doi: 10.1002/cpa.3160390404
[89] Shi Y 2026 Finite-time blow-up of two $(1+1)$D systems rigorously derived from the 3D axisymmetric Euler equations *arXiv:2604.01244*

[90] Shi Y 2026 2D inviscid Boussinesq equations and 3D axisymmetric Euler equations: (1) a unification (E_m), (2) finite-time blow-up of two unified $(1+1)$D systems rigorously derived from (E_m) *arXiv:2603.26715*
[91] Stuart J T 1998 Singularities in three-dimensional compressible Euler flows with vorticity *Theor. Comput. Fluid Dyn.* **10** 385–391
[92] Sullivan D 2014 3D incompressible fluids: combinatorial models, eigenspace models, and a conjecture about well-posedness of the 3D zero viscosity limit *J. Differential Geom.* **97** 141–148
[93] Taniuchi Y 2002 A note on the blowup criterion for the inviscid 2-D Boussinesq equations in *The Navier–Stokes Equations: Theory and Numerical Methods* (ed R Salvi) *Lecture Notes in Pure and Applied Mathematics* **223** 131–140
[94] Tao T 2016 Finite time blowup for Lagrangian modifications of the three-dimensional Euler equation *Ann. PDE* **2** Art. 9
[95] Tao T 2018 On the universality of the incompressible Euler equation on compact manifolds *Discrete Contin. Dyn. Syst.* **38**(3) 1553–1565 doi:10.3934/dcds.2018064
[96] Ukhovskii M R and Yudovich V I 1968 Axially symmetric flows of ideal and viscous fluids filling the whole space *J. Appl. Math. Mech.* **32**(1) 52–62 doi:10.1016/0021-8928(68)90147-0
[97] Vasseur A F and Vishik M 2020 Blow-up solutions to 3D Euler are hydrodynamically unstable *Commun. Math. Phys.* **378** 557–568
[98] Wang Y, Lai C-Y, Gómez-Serrano J and Buckmaster T 2023 Asymptotic self-similar blow-up profile for three-dimensional axisymmetric Euler equations using neural networks *Phys. Rev. Lett.* **130** 244002
[99] Wu J 2012 The 2D incompressible Boussinesq equations *(Peking University Summer School lecture notes)*
[100] Wu H and Wunsch M 2012 Global existence for the generalized two-component Hunter–Saxton system *J. Math. Fluid Mech.* **14** 455–469 (see also *arXiv:1009.1688*)
[101] Wunsch M 2009 The generalized Hunter–Saxton system *Discrete Contin. Dyn. Syst. Ser. B* **12** 647–656
[102] Yan W 2018 Finite time blowup explicit solutions for 3D incompressible Euler equations *arXiv:1807.05426*
[103] Zhang Q S 2021 Bounded solutions to the axially symmetric Navier–Stokes equation in a cusp region *arXiv:2106.08509*

Rigorous proof audit: navigation convention

Audit convention. The rigorous appendices are the referee-level proof layer. Each slot begins with a complete restatement of the main theorem-like entity and carries permanent three-way links to its grand-finale ledger entry and its short proof. The corresponding account blocks provide the audited long proofs; The account blocks A–J and L contain the corresponding rigorous proofs. Every closed ledger slot is therefore audited. O1–O3 are proved independently in the physical-bank O-ledger; the completed inherited-launch source-angle audit gives O4; O5 is GREEN from O4+O6.2+M8 by the permanent terminal theorems; and O6.2 is the sole **OPEN—NO PROOF ASSERTED** slot. The long-proof archive remains the derivation source for the audited statements, while the historical research archive records the provenance of the reconstructed M1–M7 modules.

Appendix A. Exact customer, source, cashier, clock, and opening credit

This account contains the complete long proofs and sets the standard used throughout the rigorous proof layer.

A1. Exact squared-coordinate Euler system.

Mutual pointers: grand-finale entry / short proof/status / main statement proposition 4.3.

Restatement of proposition 4.3. Let

$$\mathcal{D}_{xy} := \partial_t - 2xv\partial_x + 2yg\partial_y, \qquad \varepsilon_\beta \in \{0,1\}.$$

Then the tagged axisymmetric Euler system is

$$\begin{aligned} g + 2yg_y &= 2v + 2xv_x, \\ \mathcal{D}_{xy}v - v^2 + \varepsilon_\beta u^2 &= 2P_x, \\ \mathcal{D}_{xy}(\varepsilon_\beta u) &= 2\varepsilon_\beta uv, \\ \mathcal{D}_{xy}g + g^2 &= -2P_y. \end{aligned}$$

The characteristics satisfy

$$\dot{x} = -2xv, \qquad \dot{y} = 2yg.$$

At $\varepsilon_\beta = 1$ this is physical Euler. At $\varepsilon_\beta = 0$, the tagged swirl equation (4.17) becomes the identity $0 = 0$, the centrifugal source in (4.16) disappears, and the displayed active PDE system reduces to the axisymmetric Euler equations without swirl. The untagged profile u may still be retained off-ledger as a passive bookkeeping carrier, but it no longer enters the active shadow PDEs.

Full rigorous proof. Starting from the physical axisymmetric Euler equations, write

$$U^r = -rv, \qquad U^\phi = ru, \qquad U^z = zg, \qquad x = r^2, \quad y = z^2.$$

For every smooth squared-coordinate scalar $f = f(x, y, t)$,

$$D_t f = f_t + U^r(2rf_x) + U^z(2zf_y) = f_t - 2xvf_x + 2ygf_y = \mathcal{D}_{xy}f.$$

The divergence equation gives

$$0 = \partial_r(-rv) - v + \partial_z(zg) = -2v - 2xv_x + g + 2yg_y,$$

which is the first displayed relation.

For the radial momentum equation,

$$D_t(-rv) = -(D_t r)v - rD_t v = rv^2 - r\mathcal{D}_{xy}v, \qquad \frac{(U^\phi)^2}{r} = ru^2, \qquad \mathsf{p}_r = 2rP_x.$$

After division by r and rearrangement,

$$\mathcal{D}_{xy}v - v^2 + u^2 = 2P_x.$$

The Boolean provenance tag is placed only on the centrifugal feedback, so this becomes $\mathcal{D}_{xy}v - v^2 + \varepsilon_\beta u^2 = 2P_x$.

For the azimuthal equation,

$$D_t(ru) + \frac{U^r}{r}(ru) = (-rv)u + r\mathcal{D}_{xy}u - rvu = r(\mathcal{D}_{xy}u - 2vu) = 0.$$

Thus $\mathcal{D}_{xy}u = 2vu$ in the physical sector. Since ε_β is constant, multiplication by ε_β gives

$$\mathcal{D}_{xy}(\varepsilon_\beta u) = 2\varepsilon_\beta uv.$$

At $\varepsilon_\beta = 0$ this tagged equation becomes $0 = 0$, while the meridional equations lose the centrifugal feedback, exactly as required by the shadow convention.

Finally,

$$D_t(zg) = (D_t z)g + zD_t g = zg^2 + z\mathcal{D}_{xy}g, \qquad \mathsf{p}_z = 2zP_y,$$

so the axial equation yields $\mathcal{D}_{xy}g + g^2 = -2P_y$. The characteristic equations are the coefficients of the transport operator: $\dot{x} = -2xv$, $\dot{y} = 2yg$. □

Audit status: GREEN/COMPLETE.

A2. Pressure dictionary. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 4.4.

Restatement of proposition 4.4. With the protected coordinate dictionary

$$x = r^2, \qquad y = z^2, \qquad \mathsf{p}(r, z, t) = P(x, y, t),$$

the physical and squared pressure derivatives satisfy

$$\mathsf{p}_r = 2rP_x, \quad \mathsf{p}_z = 2zP_y,$$

$$\boxed{\mathsf{p}_{rr} = 2P_x + 4xP_{xx}, \qquad \mathsf{p}_{rz} = 4rzP_{xy}, \qquad \mathsf{p}_{zz} = 2P_y + 4yP_{yy}.}$$

Hence physical Hessian entries are never identified directly with P_{xx}, P_{xy}, P_{yy}.

Full rigorous proof. Because $\mathsf{p}(r, z, t) = P(r^2, z^2, t)$, the first derivatives are

$$\mathsf{p}_r = 2rP_x, \qquad \mathsf{p}_z = 2zP_y.$$

Differentiating once more,

$$\mathsf{p}_{rr} = 2P_x + 4r^2P_{xx} = 2P_x + 4xP_{xx},$$
$$\mathsf{p}_{rz} = 2r(2zP_{xy}) = 4rzP_{xy},$$

and

$$\mathsf{p}_{zz} = 2P_y + 4z^2P_{yy} = 2P_y + 4yP_{yy}.$$

These are pure chain-rule identities; no equation of motion is used. In particular, a squared-coordinate second derivative may not be substituted for a physical Hessian entry without the lower-order correction terms displayed above. □

Audit status: GREEN/COMPLETE.

A3. Pressure-free source-to-vorticity account. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 5.3.

Restatement of proposition 5.3. With

$$\Theta := \varepsilon_\beta u^2, \qquad q := \Theta_y$$

as in definition 5.2, define

$$\omega^\phi := \partial_z U^r - \partial_r U^z, \qquad D_t := \partial_t + U^r\partial_r + U^z\partial_z,$$

and

$$\bar{G} := \omega^\phi/r = zH(r^2, z^2, t), \qquad H := -2(v_y + g_x).$$

Then

$$\boxed{\mathcal{D}_{xy}\Theta = 4v\Theta, \qquad \mathcal{D}_{xy}H + gH = 2q.}$$

Equivalently, with $\beta(r, z, t) = U^\phi(r, z, t)/r$,

$$D_t\bar{G} = \varepsilon_\beta\,\partial_z(\beta^2).$$

Full rigorous proof. Since ε_β is constant and the physical swirl equation is $\mathcal{D}_{xy}u = 2vu$,

$$\mathcal{D}_{xy}\Theta = \mathcal{D}_{xy}(\varepsilon_\beta u^2) = 2\varepsilon_\beta u\,\mathcal{D}_{xy}u = 4v\Theta.$$

For the vorticity account, take the azimuthal component of the curl of the axisymmetric Euler equations. The standard regular quotient

$$\bar{G} = \frac{\omega^\phi}{r}, \qquad \beta = \frac{U^\phi}{r}$$

satisfies

$$D_t\bar{G} = \varepsilon_\beta\,\partial_z(\beta^2).$$

This identity can also be obtained by differentiating the radial and axial momentum equations and subtracting; the mixed pressure derivatives cancel exactly.

Now $\bar{G} = zH$, $D_t z = U^z = zg$, and $\partial_z = 2z\partial_y$. Hence, for $z \neq 0$,

$$z(\mathcal{D}_{xy}H + gH) = D_t(zH) = \varepsilon_\beta\, \partial_z(u^2) = 2z\,\Theta_y.$$

Division by z, followed by smooth extension across $z = 0$, yields

$$\mathcal{D}_{xy}H + gH = 2\Theta_y = 2q.$$

Thus the source-to-vorticity transfer is pressure-free and remains explicitly tagged. □

Audit status: GREEN/COMPLETE.

A4. Whole-space Hodge cashier. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 5.4.

Restatement of proposition 5.4. Let

$$\mathcal{Q} := (0,\infty)^2, \qquad dX := dx\,dy.$$

Define the meridional stream function ψ_1 by

$$U^r = -r\partial_z\psi_1, \qquad U^z = 2\psi_1 + r\partial_r\psi_1.$$

Then the meridional velocity is reconstructed from

$$-\mathcal{L}_1\psi_1 = \bar{G}, \qquad \mathcal{L}_1 = \partial_r^2 + \frac{3}{r}\partial_r + \partial_z^2.$$

and the apex compression $a(t) = v(0,0,t)$ is

$$\boxed{a(t) = \frac{3}{4}\int_0^\infty \int_{\mathbb{R}} \frac{r^3 z}{(r^2+z^2)^{5/2}} \bar{G}(r,z,t)\,dz\,dr.}$$

In squared coordinates,

$$\boxed{a(t) = \frac{3}{8}\int_{\mathcal{Q}} K_H(x,y)H(x,y,t)\,dX, \qquad K_H(x,y) := \frac{x\sqrt{y}}{(x+y)^{5/2}} > 0.}$$

Full rigorous proof. Lift the stream equation to $\mathbb{R}^5 = \mathbb{R}^4_X \times \mathbb{R}_z$ by

$$\widehat{\psi}_1(X,z,t) := \psi_1(|X|,z,t), \qquad \widehat{G}(X,z,t) := \bar{G}(|X|,z,t).$$

Because

$$\Delta^{\mathrm{rad}}_{\mathbb{R}^4} = \partial_r^2 + \frac{3}{r}\partial_r,$$

the equation $-\mathcal{L}_1\psi_1 = \bar{G}$ becomes

$$-\Delta_{\mathbb{R}^5}\widehat{\psi}_1 = \widehat{G}.$$

The fundamental solution of $-\Delta$ in five dimensions is $(8\pi^2)^{-1}|\zeta|^{-3}$. Therefore

$$\widehat{\psi}_1(\zeta,t) = \frac{1}{8\pi^2}\int_{\mathbb{R}^5} \frac{\widehat{G}(\eta,t)}{|\zeta-\eta|^3}\,d\eta.$$

Since $a(t) = \partial_z\psi_1(0,0,t)$, differentiate the kernel in the observation z-variable and set the observation point equal to the origin:

$$a(t) = \frac{3}{8\pi^2}\int_{\mathbb{R}^5} \frac{z'\widehat{G}(X',z',t)}{(|X'|^2+z'^2)^{5/2}}\,dX'\,dz'.$$

Spherical integration in $\mathbb{R}^4$, with $|S^3| = 2\pi^2$, gives

$$a(t) = \frac{3}{4}\int_0^\infty \int_{\mathbb{R}} \frac{r^3 z}{(r^2+z^2)^{5/2}} \bar{G}(r,z,t)\, dz\, dr.$$

Using $\bar{G} = zH$, the evenness in z, and $x = r^2,\ y = z^2$,

$$r^3 dr = \frac{x}{2}\, dx, \qquad 2z^2\, dz = \sqrt{y}\, dy,$$

so

$$a(t) = \frac{3}{8}\int_{\mathcal{Q}} \frac{x\sqrt{y}}{(x+y)^{5/2}}\, H(x,y,t)\, dx\, dy.$$

The kernel is positive in the open quadrant. □

Audit status: GREEN/COMPLETE.

A5. Seven-dimensional Hodge lift. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 5.5.

Restatement of proposition 5.5. Writing $\psi_1 = z\chi(x,y,t)$ gives

$$v = \chi + 2y\chi_y, \qquad g = 2\chi + 2x\chi_x,$$

$$-\mathcal{L}_7\chi = H, \qquad \mathcal{L}_7 = 4x\partial_x^2 + 8\partial_x + 4y\partial_y^2 + 6\partial_y.$$

For $\mathbf{X} \in \mathbb{R}^4$, $\mathbf{Z} \in \mathbb{R}^3$, define the radial lifts

$$\widetilde{\chi}(\mathbf{X},\mathbf{Z},t) := \chi(|\mathbf{X}|^2, |\mathbf{Z}|^2, t), \qquad \widetilde{H}(\mathbf{X},\mathbf{Z},t) := H(|\mathbf{X}|^2, |\mathbf{Z}|^2, t).$$

Then

$$-\Delta_{\mathbb{R}^7}\widetilde{\chi} = \widetilde{H},$$

so all meridional values and first jets are whole-space Hodge descendants of H.

Full rigorous proof. Write $\psi_1 = z\chi(x,y,t)$, with $x = r^2$, $y = z^2$. Direct differentiation gives

$$\partial_z \psi_1 = \chi + 2y\chi_y,$$

hence $v = \chi + 2y\chi_y$. Likewise,

$$\frac{2\psi_1 + r\partial_r\psi_1}{z} = 2\chi + 2x\chi_x,$$

hence $g = 2\chi + 2x\chi_x$.

Next,

$$\mathcal{L}_1(z\chi) = z\left(4x\chi_{xx} + 8\chi_x + 4y\chi_{yy} + 6\chi_y\right).$$

Since $-\mathcal{L}_1\psi_1 = \bar{G} = zH$, division by z and smooth extension at $z = 0$ yield

$$-\mathcal{L}_7\chi = H, \qquad \mathcal{L}_7 = 4x\partial_x^2 + 8\partial_x + 4y\partial_y^2 + 6\partial_y.$$

For a radial function of $X \in \mathbb{R}^4$,

$$\Delta_X f(|X|^2) = 4x f_{xx} + 8 f_x,$$

and for a radial function of $Z \in \mathbb{R}^3$,

$$\Delta_Z f(|Z|^2) = 4y f_{yy} + 6 f_y.$$

Therefore the lift

$$\widetilde{\chi}(X,Z,t) = \chi(|X|^2, |Z|^2, t)$$

satisfies $-\Delta_{\mathbb{R}^7}\widetilde{\chi} = \widetilde{H}$. Since the fundamental solution of $-\Delta$ in seven dimensions is

$$\frac{1}{(7-2)|S^6|}\,|\zeta|^{-5} = \frac{3}{16\pi^3}|\zeta|^{-5},$$

rapid decay gives the stated Newton representation. Thus all meridional values and jets are fixed linear Hodge descendants of H; there is no free harmonic mode in the whole-space class. □

Audit status: GREEN/COMPLETE.

A6. Detector evolution and clock. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 6.5.

Restatement of proposition 6.5. Let

$$\psi_\beta(t) = 2u_y(0,0,t), \qquad \Psi(t) = \psi_\beta(t)^2, \qquad \mathcal{J}_\beta(t) = \varepsilon_\beta \Psi(t)$$

be the detector variables of definition 6.1. Throughout the classical lifespan,

$$\boxed{\dot{\psi_\beta} = -2a\psi_\beta, \qquad \dot{\Psi} = -4a\Psi, \qquad \dot{\mathcal{J}}_\beta = -4a\mathcal{J}_\beta.}$$

If $\Psi(0) > 0$, define

$$\boxed{\mathfrak{t}_\Psi(t) := e^{-2\int_0^t a(\tau)d\tau} = \frac{\psi_\beta(t)}{\psi_\beta(0)} = \sqrt{\frac{\Psi(t)}{\Psi(0)}} = \sqrt{\frac{\mathcal{J}_\beta(t)}{\mathcal{J}_\beta(0)}}.}$$

Then

$$\boxed{a(t) = -\frac{1}{2}\frac{\dot{\mathfrak{t}_\Psi}(t)}{\mathfrak{t}_\Psi(t)}.}$$

Full rigorous proof. For the fixed customer one has $b(0) = u(0,0,0) = 0$. The apex azimuthal equation is $\dot{b} = 2ab$, hence uniqueness gives $b(t) \equiv 0$.

Differentiate the physical-sector swirl equation $\mathcal{D}_{xy}u = 2vu$ in y. Since

$$\partial_y(\mathcal{D}_{xy}u) = \mathcal{D}_{xy}u_y - 2xv_yu_x + 2(g + yg_y)u_y,$$

one obtains

$$\mathcal{D}_{xy}u_y - 2xv_yu_x + 2(g + yg_y)u_y = 2v_yu + 2vu_y.$$

At $x = y = 0$, smoothness and $u(0,0,t) = 0$ remove the terms containing u and x; moreover $v(0,0,t) = a(t)$ and incompressibility gives $g(0,0,t) = 2a(t)$. Thus

$$\frac{d}{dt}u_y(0,0,t) = -2a(t)u_y(0,0,t).$$

Multiplication by 2 yields

$$\dot{\psi_\beta} = -2a\psi_\beta.$$

Since $\Psi = \psi_\beta^2$, $\mathcal{J}_\beta = \varepsilon_\beta\Psi$, and ε_β is constant,

$$\dot{\Psi} = 2\psi_\beta\dot{\psi_\beta} = -4a\Psi, \qquad \dot{\mathcal{J}}_\beta = -4a\mathcal{J}_\beta.$$

Integrating the scalar ODE for ψ_β gives

$$\frac{\psi_\beta(t)}{\psi_\beta(0)} = \exp\left(-2\int_0^t a(\tau)\,d\tau\right) =: \mathfrak{t}_\Psi(t).$$

In the physical sector, or after the common-tag quotient is independently admitted, this also equals $\sqrt{\mathcal{J}_\beta(t)/\mathcal{J}_\beta(0)}$. Differentiating $\log \mathfrak{t}_\Psi$ gives $a = -\dot{\mathfrak{t}_\Psi}/(2\mathfrak{t}_\Psi)$. □

Audit status: GREEN/COMPLETE.

A7. Reserve–clock identity. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 6.7.

Restatement of proposition 6.7. Let $\mathcal{F}_\Psi := Q - 9a^2$ be the reserve from definition 6.6, and let $\mathfrak{t}_\Psi$ be the detector clock from proposition 6.5. Then

$$\boxed{\mathcal{F}_\Psi = 3(\dot{a} - 2a^2) = -\frac{3}{2}\frac{\ddot{\mathfrak{t}}_\Psi}{\mathfrak{t}_\Psi}.}$$

Hence

$$\boxed{\mathcal{F}_\Psi \geq 0 \iff \ddot{\mathfrak{t}}_\Psi \leq 0.}$$

Full rigorous proof. With $b \equiv 0$, the regular apex radial equation and the pressure trace yield

$$\dot{a} = -a^2 + \tfrac{1}{3}Q.$$

Equivalently,

$$Q = 3(\dot{a} + a^2).$$

Therefore

$$\mathcal{F}_\Psi = Q - 9a^2 = 3(\dot{a} - 2a^2).$$

By the detector-clock identity,

$$\dot{\mathfrak{t}}_\Psi = -2a\mathfrak{t}_\Psi.$$

A second derivative gives

$$\ddot{\mathfrak{t}}_\Psi = -2\dot{a}\,\mathfrak{t}_\Psi - 2a\dot{\mathfrak{t}}_\Psi = -2(\dot{a} - 2a^2)\mathfrak{t}_\Psi.$$

Since $\mathfrak{t}_\Psi > 0$ throughout the classical interval before clock collapse,

$$\mathcal{F}_\Psi = -\frac{3}{2}\frac{\ddot{\mathfrak{t}}_\Psi}{\mathfrak{t}_\Psi}.$$

Hence $\mathcal{F}_\Psi \geq 0$ if and only if $\ddot{\mathfrak{t}}_\Psi \leq 0$. □

Audit status: GREEN/COMPLETE.

A8. Conditional finite-time collapse. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 6.8.

Restatement of theorem 6.8. Let $\Psi, \mathfrak{t}_\Psi, \mathcal{F}_\Psi$ be as in definitions 6.1 and 6.6 and proposition 6.5. Suppose $\Psi(0) > 0$ and at some classical time t_1 one has

$$a(t_1) > 0, \qquad \mathcal{F}_\Psi(t) \geq 0$$

for all later classical times under consideration. Define

$$T_* := \inf\{t \geq t_1 : \mathfrak{t}_\Psi(t) = 0\}, \qquad \inf \varnothing := +\infty.$$

Then either regularity fails beforehand, or

$$\boxed{T_* \leq t_1 + \frac{1}{2a(t_1)}.}$$

If the solution persists to T_*, then

$$\Psi(t) \to 0, \qquad a(t) \to +\infty \quad (t \uparrow T_*).$$

Full rigorous proof. By the reserve–clock identity, $\mathcal{F}_\Psi \geq 0$ implies that $\mathfrak{t}_\Psi$ is concave. At the time t_1,

$$\dot{\mathfrak{t}_\Psi}(t_1) = -2a(t_1)\mathfrak{t}_\Psi(t_1) < 0.$$

Concavity therefore gives, for every later classical time,

$$\mathfrak{t}_\Psi(t) \leq \mathfrak{t}_\Psi(t_1) + \dot{\mathfrak{t}_\Psi}(t_1)(t - t_1) = \mathfrak{t}_\Psi(t_1)\big[1 - 2a(t_1)(t - t_1)\big].$$

The affine upper bound on the right vanishes at

$$t = t_1 + \frac{1}{2a(t_1)}.$$

A positive clock cannot remain above a nonpositive upper bound. Hence either the classical solution exits before that time or the first zero T_* of $\mathfrak{t}_\Psi$ satisfies the asserted deadline.

If the solution remains classical up to T_*, then $\Psi(t) = \Psi(0)\mathfrak{t}_\Psi(t)^2 \to 0$. Concavity also gives $\dot{\mathfrak{t}_\Psi}(t) \leq \dot{\mathfrak{t}_\Psi}(t_1) < 0$, and therefore

$$a(t) = -\frac{\dot{\mathfrak{t}_\Psi}(t)}{2\mathfrak{t}_\Psi(t)} \longrightarrow +\infty \qquad (t \uparrow T_*).$$

Thus clock collapse is incompatible with continued bounded classical strain. □

Audit status: GREEN/COMPLETE.

A9. Opening family. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 7.1.

Restatement of proposition 7.1. Let $B \neq 0$ and $\lambda > 0$, and use the fixed constants $c_* = \sqrt{2/3}$ and $s_* = 1/\sqrt{3}$. For

$$u_{\mathrm{ini}} = B(y + \lambda x)e^{-(c_* x + s_* y)}, \qquad v_{\mathrm{ini}} = g_{\mathrm{ini}} = 0,$$

define

$$\mathcal{I}(\lambda) := \int_{-\pi/2}^{\pi/2} \frac{\cos^5\phi\, \sin^2\phi\, (\sin^2\phi + \lambda\cos^2\phi)}{(c_*\cos^2\phi + s_*\sin^2\phi)^3}\, d\phi > 0.$$

Then

$$\boxed{\dot{a}(0) = \frac{3\varepsilon_\beta B^2}{8}(c_* - s_*\lambda)\mathcal{I}(\lambda).}$$

Therefore, in the physical sector $\varepsilon_\beta = 1$,

$$0 < \lambda < \sqrt{2} \Rightarrow \dot{a}(0) > 0, \quad \lambda = \sqrt{2} \Rightarrow \dot{a}(0) = 0, \quad \lambda > \sqrt{2} \Rightarrow \dot{a}(0) < 0.$$

The final $A = 10$ customer in (5.1) is a later fixed descendant of this source-selection program; no new customer is introduced thereafter.

Full rigorous proof. At $t = 0$ the meridional field vanishes, so $\bar{G}(0) = 0$. The pressure-free vorticity equation gives

$$\partial_t \bar{G}\big|_{t=0} = \varepsilon_\beta\, \partial_z(\beta_{\mathrm{ini}}^2).$$

Differentiating the whole-space Hodge cashier at $t = 0$,

$$\dot{a}(0) = \frac{3\varepsilon_\beta}{4}\int_0^\infty \int_{\mathbb{R}} \frac{r^3 z}{(r^2 + z^2)^{5/2}} \partial_z(\beta_{\mathrm{ini}}^2)\, dz\, dr.$$

For the displayed family,

$$\beta_{\mathrm{ini}}(r, z) = B(z^2 + \lambda r^2)e^{-c_* r^2 - s_* z^2}.$$

Set

$$r = \rho\cos\phi, \qquad z = \rho\sin\phi, \qquad -\frac{\pi}{2} \leq \phi \leq \frac{\pi}{2},$$

and define

$$q_\lambda(\phi) = \sin^2\phi + \lambda\cos^2\phi, \qquad k(\phi) = c_*\cos^2\phi + s_*\sin^2\phi.$$

After the elementary z-differentiation, all ρ-dependence reduces to

$$\int_0^\infty \rho^3 q_\lambda\big(1 - s_*\rho^2 q_\lambda\big)e^{-2k\rho^2}\,d\rho.$$

Using

$$\int_0^\infty \rho^3 e^{-2k\rho^2}\,d\rho = \frac{1}{8k^2}, \qquad \int_0^\infty \rho^5 e^{-2k\rho^2}\,d\rho = \frac{1}{8k^3},$$

this integral equals

$$\frac{q_\lambda(k - s_* q_\lambda)}{8k^3}.$$

The crucial algebraic identity is

$$k - s_* q_\lambda = (c_* - s_*\lambda)\cos^2\phi.$$

Collecting the angular Jacobian and Hodge factors yields exactly

$$\dot a(0) = \frac{3\varepsilon_\beta B^2}{8}(c_* - s_*\lambda)\mathcal{I}(\lambda),$$

where $\mathcal{I}(\lambda)$ is the positive integral displayed in the proposition. Because $c_*/s_* = \sqrt{2}$, the sign trichotomy follows. □

Audit status: GREEN/COMPLETE.

A10. Opening reserve. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 7.2.

> **Restatement of proposition 7.2.** For an active positive-opening seed with $a(0) = b(0) = 0$,
>
> $$\boxed{Q(0) = 3\dot a(0) > 0, \qquad \mathcal{F}_\Psi(0) = 3\dot a(0) > 0.}$$
>
> Thus the initial account is funded; the problem is persistence, not launch.

Full rigorous proof. At an active positive-opening seed, $a(0) = b(0) = 0$. The apex radial equation

$$\dot a = -a^2 - \frac{1}{3}\varepsilon_\beta b^2 + \frac{1}{3}Q$$

therefore reduces at $t = 0$ to

$$Q(0) = 3\dot a(0).$$

By the definition $\mathcal{F}_\Psi = Q - 9a^2$ and $a(0) = 0$,

$$\mathcal{F}_\Psi(0) = Q(0) = 3\dot a(0).$$

For a positive-opening seed, $\dot a(0) > 0$, so both quantities are strictly positive. Thus the initial reserve is funded and only persistence remains open. □

Audit status: GREEN/COMPLETE.

Appendix B. Provenance and restart/deformation accounting

The complete statements are restated here so the ledger slot is self-contained and its ID remains stable.

B1. Routing–GREEN normal form. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 8.2.

> **Restatement of proposition 8.2.** Every finite polynomial expression made from ordinary routing quantities and certified GREEN descendants can be written
>
> $$\boxed{\mathcal{X} = \mathcal{U}_{\mathcal{X}} + \varepsilon_\beta \mathcal{G}_{\mathcal{X}}.}$$
>
> A mixed sum remains mixed. In particular, multiplying an Euler-background quantity by ε_β at the end does not manufacture ancestry.

Proof. Let the finitely many ordinary routing symbols entering the expression be $\mathcal{U}_1, \dots, \mathcal{U}_m$, and let the certified GREEN symbols be represented as $\varepsilon_\beta \mathcal{G}_1, \dots, \varepsilon_\beta \mathcal{G}_n$ according to definition 8.1. Expanding any polynomial expression into monomials reduces the claim to one monomial. Such a monomial has the form

$$c \prod_{i=1}^{m} \mathcal{U}_i^{\kappa_i} \prod_{j=1}^{n} (\varepsilon_\beta \mathcal{G}_j)^{\nu_j}, \qquad \kappa_i, \nu_j \in \mathbb{N}_0.$$

If $\sum_j \nu_j = 0$, the monomial contains no source descendant and belongs to the pure routing part $\mathcal{U}_{\mathcal{X}}$. If $\sum_j \nu_j \geq 1$, the Boolean law $\varepsilon_\beta^k = \varepsilon_\beta$ for every $k \geq 1$ gives

$$c \prod_{i} \mathcal{U}_i^{\kappa_i} \prod_{j} (\varepsilon_\beta \mathcal{G}_j)^{\nu_j} = \varepsilon_\beta \, c \prod_{i} \mathcal{U}_i^{\kappa_i} \prod_{j} \mathcal{G}_j^{\nu_j},$$

so the monomial belongs to $\varepsilon_\beta \mathcal{G}_{\mathcal{X}}$. Summing the two classes of monomials proves the asserted normal form. The same calculation gives closure under addition and multiplication; for example,

$$(\mathcal{U}_1 + \varepsilon_\beta \mathcal{G}_1)(\mathcal{U}_2 + \varepsilon_\beta \mathcal{G}_2) = \mathcal{U}_1 \mathcal{U}_2 + \varepsilon_\beta (\mathcal{U}_1 \mathcal{G}_2 + \mathcal{U}_2 \mathcal{G}_1 + \mathcal{G}_1 \mathcal{G}_2).$$

For the mixed-sum warning, let S_0 denote formal shadow evaluation at $\varepsilon_\beta = 0$ before any shadow-singular quotient is admitted. Then

$$\mathsf{S}_0(\mathcal{U} + \varepsilon_\beta \mathcal{G}) = \mathsf{S}_0(\mathcal{U}).$$

If $\mathcal{U}$ has no independently certified source genealogy, replacing the mixed quantity by $\varepsilon_\beta(\mathcal{U} + \mathcal{G})$ would force its shadow to zero and would therefore alter the dependency ledger. Hence an ordinary background term cannot be promoted to GREEN merely by attaching ε_β after the estimate has been derived. This is precisely the no-late-stamping rule. □

Audit status: GREEN/COMPLETE.

B2. Two-shadow firewall. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 8.4.

> **Restatement of proposition 8.4.** Every terminal candidate must pass both tests:
>
> (i) under the formal single-seed shadow $\varepsilon_\beta = 0$, no positive cashier or terminal contradiction may remain;
>
> (ii) under physical no-swirl axisymmetric Euler, a generic meridional estimate may route existing credit but may not by itself imply finite-time blow-up.
>
> Hence every final payment must retain an irreducible source-specific ancestor.

Proof. The two tests are logically distinct.

Single-seed Boolean shadow. For the fixed launch used in this manuscript the initial meridional field is zero. Under $\varepsilon_\beta = 0$ the tagged centrifugal source $\Theta = \varepsilon_\beta u^2$ vanishes. The

zero meridional field, together with spatially constant pressure, solves the meridional shadow equations; classical uniqueness gives

$$v = g = H = a = Q = q = 0$$

throughout the shadow lifespan, while the prescribed swirl carrier is only passive routing data. By the idempotent calculus and proposition 8.2, every certified raw source payment also vanishes under this projection. Therefore a proposed terminal payment which remains strictly positive in the shadow necessarily contains an uncancelled routing part or has used a shadow-singular normalization before admission, and so it cannot be the source-specific terminal payer.

General physical no-swirl regression. Set the physical swirl U^{ϕ} identically to zero but allow arbitrary smooth axisymmetric meridional initial data. Identities involving only the meridional Euler system, incompressibility, Hodge reconstruction, or generic first-jet estimates remain available in this class. Classical axisymmetric Euler without swirl is globally regular under the standing smooth-decaying hypotheses [96]. Consequently a chain using only such generic meridional information cannot by itself furnish a valid finite-time singularity mechanism for the present source-specific problem: otherwise the same mechanism would apply to the globally regular no-swirl benchmark. Generic meridional inequalities may route or estimate already certified source credit, but a terminal chain must contain an irreducible input whose proof uses the swirl-source genealogy (for example Θ, (q, C_β), $H^{\pm}$, or the exact negative-source ancestry).

The first test prevents ancestry loss in the special Boolean shadow; the second prevents generic meridional structure from being mistaken for source-specific terminal money. Hence both tests are necessary. □

Audit status: GREEN/COMPLETE.

B3. Logarithmic tangent system. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 9.2.

> **Restatement of proposition 9.2.** Let $s := y/x$ be the aspect ratio, let t_0 be a starting time, and for an infinitesimal displacement $(\delta x, \delta y)$ put
>
> $$\mathfrak{p} = \delta x/x, \qquad \mathfrak{q} = \delta y/y.$$
>
> Then
>
> $$\frac{d}{dt}\begin{pmatrix}\mathfrak{p}\\ \mathfrak{q}\end{pmatrix} = \mathbb{M}\begin{pmatrix}\mathfrak{p}\\ \mathfrak{q}\end{pmatrix},$$
>
> with
>
> $$\boxed{\mathbb{M} = \frac{1}{2}\mathfrak{A} I_2 + \begin{pmatrix}-\mathfrak{J} & -s(\mathfrak{S}-\mathfrak{T})\\ \mathfrak{S}+\mathfrak{T} & \mathfrak{J}\end{pmatrix},}$$
>
> where I_2 is the 2×2 identity matrix. If $\mathcal{U}(t, t_0)$ is the fundamental logarithmic tangent matrix with $\mathcal{U}(t_0, t_0) = I_2$, then
>
> $$\boxed{\det \mathcal{U}(t, t_0) = \exp\left(\int_{t_0}^{t} \mathfrak{A}(\tau) d\tau\right).}$$

Proof. The squared-coordinate characteristics satisfy

$$\dot{x} = -2xv(x, y, t), \qquad \dot{y} = 2yg(x, y, t).$$

Let $(\delta x, \delta y)$ be an infinitesimal variation and put

$$\mathfrak{p} := \frac{\delta x}{x}, \qquad \mathfrak{q} := \frac{\delta y}{y}.$$

Linearization gives

$$\delta\dot{x} = -2(v + xv_x)\delta x - 2xv_y\delta y,$$
$$\delta\dot{y} = 2yg_x\delta x + 2(g + yg_y)\delta y.$$

Since $\dot{x}/x = -2v$ and $\dot{y}/y = 2g$,

$$\frac{d}{dt}\begin{pmatrix}\mathfrak{p}\\ \mathfrak{q}\end{pmatrix} = \begin{pmatrix}-2xv_x & -2yv_y\\ 2xg_x & 2yg_y\end{pmatrix}\begin{pmatrix}\mathfrak{p}\\ \mathfrak{q}\end{pmatrix}.$$

Using the protected first-jet notation $C = xv_x$, $D = yg_y$, $E = yv_y$, $F = xg_x$ and the identities from definition 9.1,

$$C = \frac{1}{2}\mathfrak{J} - \frac{1}{4}\mathfrak{A}, \quad D = \frac{1}{2}\mathfrak{J} + \frac{1}{4}\mathfrak{A},$$
$$F = \frac{1}{2}(\mathfrak{S} + \mathfrak{T}), \quad E = \frac{s}{2}(\mathfrak{S} - \mathfrak{T}), \qquad s = \frac{y}{x},$$

we obtain exactly

$$\mathbb{M} = \frac{1}{2}\mathfrak{A}I_2 + \begin{pmatrix}-\mathfrak{J} & -s(\mathfrak{S} - \mathfrak{T})\\ \mathfrak{S} + \mathfrak{T} & \mathfrak{J}\end{pmatrix}.$$

Let $\mathcal{U}(t, t_0)$ be the fundamental matrix. Liouville's formula gives

$$\frac{d}{dt}\det\mathcal{U} = \operatorname{tr}(\mathbb{M})\det\mathcal{U}.$$

The traceless restart block contributes no trace and $\operatorname{tr}\mathbb{M} = \mathfrak{A}$. Since $\det\mathcal{U}(t_0, t_0) = 1$,

$$\det\mathcal{U}(t, t_0) = \exp\left(\int_{t_0}^{t}\mathfrak{A}(\tau)d\tau\right).$$

□

Audit status: GREEN/COMPLETE.

B4. Tangent exit pays derivative action. *Mutual pointers:* grand-finale entry / short proof/status / main statement lemma 9.3.

> **Restatement of lemma 9.3.** Let $\mathcal{U}(t, t_0)$ and $\mathbb{M}$ be as in proposition 9.2. If a logarithmic tangent column starts from a unit coordinate vector e_j at t_0 and, in the Euclidean norm, satisfies $\|\mathcal{U}(t, t_0)e_j - e_j\| \geq \eta > 0$, then
>
> $$\boxed{\int_{t_0}^{t}\|\mathbb{M}(\tau)\|d\tau \geq \log(1 + \eta).}$$
>
> On a compact aspect-ratio window $0 < s_- \leq s = y/x \leq s_+ < \infty$, this forces a definite accumulated amount of at least one of
>
> $$|\mathfrak{A}|, \qquad |\mathfrak{J}|, \qquad |\mathfrak{S}|, \qquad |\mathfrak{T}|.$$
>
> Thus the tangent account is not a new mechanism; it is a cumulative form of the same restart bills.

Proof. Use the Euclidean vector norm and its induced operator norm and set

$$I(t) := \int_{t_0}^{t}\|\mathbb{M}(\tau)\|d\tau.$$

The fundamental matrix obeys

$$\mathcal{U}(t,t_0) - I_2 = \int_{t_0}^{t} \mathbb{M}(\tau)\mathcal{U}(\tau,t_0)d\tau.$$

Gronwall yields

$$\|\mathcal{U}(\tau,t_0)\| \le \exp\left(\int_{t_0}^{\tau} \|\mathbb{M}(\xi)\|d\xi\right).$$

Hence, for a unit coordinate vector e_j,

$$\begin{aligned}\|\mathcal{U}(t,t_0)e_j - e_j\| &\le \int_{t_0}^{t} \|\mathbb{M}(\tau)\| \exp\left(\int_{t_0}^{\tau} \|\mathbb{M}(\xi)\|d\xi\right) d\tau \\ &= e^{I(t)} - 1.\end{aligned}$$

If $\|\mathcal{U}(t,t_0)e_j - e_j\| \ge \eta$, then $I(t) \ge \log(1+\eta)$.

On a compact aspect-ratio window $s_- \le s = y/x \le s_+$, the explicit matrix from proposition 9.2 gives a finite constant $\mathfrak{c}_{s_-,s_+}$ such that

$$\|\mathbb{M}\| \le \mathfrak{c}_{s_-,s_+}(|\mathfrak{A}| + |\mathfrak{J}| + |\mathfrak{S}| + |\mathfrak{T}|).$$

Therefore

$$\int_{t_0}^{t} (|\mathfrak{A}| + |\mathfrak{J}| + |\mathfrak{S}| + |\mathfrak{T}|)d\tau \ge \frac{\log(1+\eta)}{\mathfrak{c}_{s_-,s_+}}.$$

By the pigeonhole principle at least one of the four channel integrals is at least $\log(1+\eta)/(4\mathfrak{c}_{s_-,s_+})$. Thus tangent departure consumes a definite amount of the existing restart action and is not an independent mechanism. □

Audit status: GREEN/COMPLETE.

B5. Exact radial star-hull identity. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 9.4.

Restatement of proposition 9.4. For $X = (x,y) \in \mathcal{Q}$ put $s := y/x$ and $\mathcal{R}_v := xv_x + yv_y$. The radial scaling derivative satisfies

$$\boxed{\mathcal{R}_v = -\frac{1}{4}\mathfrak{A} + \frac{1}{2}\mathfrak{J} - \frac{1}{4}yH - \frac{1}{2}s\mathfrak{T}.}$$

Moreover,

$$\boxed{v(X,t) - a(t) = \int_0^1 \mathcal{R}_v(\lambda X,t)\frac{d\lambda}{\lambda}.}$$

Consequently

$$\boxed{|v(X,t) - a(t)| \le \int_0^1 \left(\frac{1}{4}|\mathfrak{A}| + \frac{1}{2}|\mathfrak{J}| + \frac{1}{4}y_\lambda|H| + \frac{1}{2}s|\mathfrak{T}|\right)(\lambda X,t)\frac{d\lambda}{\lambda},}$$

where $y_\lambda := \lambda y$. Since

$$g - 2a = 2(v-a) - \mathfrak{A},$$

the same logarithmic star hull controls both centered value defects.

Proof. From definition 9.1,

$$xv_x = C = \frac{1}{2}\mathfrak{J} - \frac{1}{4}\mathfrak{A}, \qquad yv_y = E = \frac{s}{2}(\mathfrak{S} - \mathfrak{T}).$$

Since $\mathfrak{S} = -xH/2$ and $sx = y$,

$$E = -\frac{1}{4}yH - \frac{1}{2}s\mathfrak{T}.$$

Adding the two identities gives

$$xv_x + yv_y = -\frac{1}{4}\mathfrak{A} + \frac{1}{2}\mathfrak{J} - \frac{1}{4}yH - \frac{1}{2}s\mathfrak{T}.$$

For $X = (x, y)$ define $f(\lambda) = v(\lambda x, \lambda y, t)$. Smoothness at the apex gives $f(0) = a(t)$. If $\mathcal{R}_v(X, t) := xv_x(X, t) + yv_y(X, t)$, then for $\lambda > 0$

$$f'(\lambda) = xv_x(\lambda X, t) + yv_y(\lambda X, t) = \frac{1}{\lambda}\mathcal{R}_v(\lambda X, t).$$

Because v is smooth in (x, y), $\mathcal{R}_v(\lambda X, t) = O(\lambda)$ as $\lambda \downarrow 0$, so the quotient is locally integrable and the fundamental theorem of calculus gives the exact logarithmic radial identity

$$\boxed{v(X, t) - a(t) = \int_0^1 \mathcal{R}_v(\lambda X, t)\frac{d\lambda}{\lambda}.}$$

Equivalently, with $\lambda = e^{-\theta}$,

$$v(X, t) - a(t) = \int_0^\infty \mathcal{R}_v(e^{-\theta}X, t)d\theta.$$

At λX the aspect ratio remains s and the second squared coordinate is $y_\lambda = \lambda y$. Substitution of the exact radial identity followed by the triangle inequality yields the asserted logarithmic star-hull bound. Finally $\mathfrak{A} = 2v - g$ rearranges to

$$g - 2a = 2(v - a) - \mathfrak{A},$$

so the same star hull also controls the centered axial value defect. □

Audit status: GREEN/COMPLETE.

B6. First-order bridge among restart channels. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 9.6.

> **Restatement of proposition 9.6.** With $s := y/x$ and the restart basis of definition 9.1, the axial-relief field obeys
>
> $$\boxed{x\partial_x\mathfrak{A} = \mathfrak{J} - \frac{1}{2}\mathfrak{A} + \frac{1}{4}xH - \frac{1}{2}\mathfrak{T},}$$
>
> $$\boxed{y\partial_y\mathfrak{A} = -\frac{1}{2}\mathfrak{J} - \frac{1}{4}\mathfrak{A} - \frac{1}{2}yH - s\mathfrak{T}.}$$
>
> Equivalently,
>
> $$\boxed{yH = -2y\partial_y\mathfrak{A} - \mathfrak{J} - \frac{1}{2}\mathfrak{A} - 2s\mathfrak{T}.}$$
>
> Hence the signed restart channels are algebraically coupled and cannot be treated as independent debts.

Proof. Differentiate $\mathfrak{A} = 2v - g$. Radially,

$$x\partial_x \mathfrak{A} = 2xv_x - xg_x = 2C - F.$$

Use

$$2C = \mathfrak{J} - \frac{1}{2}\mathfrak{A}, \qquad F = \frac{1}{2}(\mathfrak{S} + \mathfrak{T}) = -\frac{1}{4}xH + \frac{1}{2}\mathfrak{T}$$

to obtain

$$x\partial_x \mathfrak{A} = \mathfrak{J} - \frac{1}{2}\mathfrak{A} + \frac{1}{4}xH - \frac{1}{2}\mathfrak{T}.$$

Likewise

$$y\partial_y \mathfrak{A} = 2yv_y - yg_y = 2E - D.$$

Since

$$E = \frac{s}{2}(\mathfrak{S} - \mathfrak{T}) = -\frac{1}{4}yH - \frac{1}{2}s\mathfrak{T}, \qquad D = \frac{1}{2}\mathfrak{J} + \frac{1}{4}\mathfrak{A},$$

we get

$$y\partial_y \mathfrak{A} = -\frac{1}{2}\mathfrak{J} - \frac{1}{4}\mathfrak{A} - \frac{1}{2}yH - s\mathfrak{T}.$$

Solving the last identity for yH gives the equivalent bridge

$$yH = -2y\partial_y \mathfrak{A} - \mathfrak{J} - \frac{1}{2}\mathfrak{A} - 2s\mathfrak{T}.$$

Thus the signed restart channels are exact first-order transforms of one another and cannot consistently be booked as independent debts. □

Audit status: GREEN/COMPLETE.

B7. Signed log-box thickening. *Mutual pointers:* grand-finale entry / short proof/status / main statement lemma 9.7.

Restatement of lemma 9.7. Fix a time t with $a(t) > 0$, a point $X_* = (x_*, y_*) \in \mathcal{Q}$, and $\rho > 0$. Define the logarithmic box

$$\mathcal{B}_\rho(X_*) := \{(x, y) : x = x_* e^\xi,\ y = y_* e^\eta,\ |\xi| < \rho,\ |\eta| < \rho\}.$$

Let a C^1 restart scalar $\mathcal{F}$ satisfy

$$\mathcal{F}(X_*) > c\, a(t), \qquad c > 0.$$

Then either

$$\boxed{\int_{\mathcal{B}_\rho(X_*)} [\mathcal{F}]_+ \frac{dxdy}{xy} \geq 2c\rho^2 a(t),}$$

or

$$\boxed{\sup_{\mathcal{B}_\rho(X_*)} (|x\mathcal{F}_x| + |y\mathcal{F}_y|) > \frac{c}{4\rho} a(t).}$$

Thus every pointwise restart either thickens into a scale-invariant signed payment or forces a higher derivative exit.

Proof. Use logarithmic coordinates

$$x = x_* e^\xi, \qquad y = y_* e^\eta.$$

Then $dxdy/(xy) = d\xi d\eta$ and

$$\partial_\xi \mathcal{F} = x\mathcal{F}_x, \qquad \partial_\eta \mathcal{F} = y\mathcal{F}_y.$$

Assume the derivative alternative fails, so

$$\sup_{\mathcal{B}_\rho(X_*)} (|x\mathcal{F}_x| + |y\mathcal{F}_y|) \le \frac{c}{4\rho} a(t).$$

For any $|\xi|, |\eta| < \rho$, connect $(0,0)$ to (ξ,η) by two coordinate segments. The fundamental theorem of calculus gives

$$|\mathcal{F}(x_* e^{\xi}, y_* e^{\eta}) - \mathcal{F}(X_*)| \le \frac{c}{4\rho} a(t)(|\xi| + |\eta|) \le \frac{c}{2} a(t).$$

Since $\mathcal{F}(X_*) > c a(t)$, throughout the box

$$\mathcal{F} \ge \frac{c}{2} a(t) > 0.$$

The logarithmic area is $4\rho^2$, hence

$$\int_{\mathcal{B}_\rho(X_*)} [\mathcal{F}]_+ \frac{dxdy}{xy} \ge \frac{c}{2} a(t)\, 4\rho^2 = 2c\rho^2 a(t).$$

Thus failure of the derivative exit forces the thickened signed payment; the two alternatives are exhaustive. □

Audit status: GREEN/COMPLETE.

Appendix C. Genealogy banks and positive-cashier relaunch

The following account contains the rigorous C1–C17 proofs. They use the protected namespace and, where radial transport enters, the corrected logarithmic star-hull measure. The later items additionally audit the Hodge-matched dual-moment gate and the compact/terminal relaunch firewalls.

C1. Exact two-bank equations. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 10.2.

Restatement of proposition 10.2. Put

$$\mathcal{S}_\pm := \frac{3}{4} \int_{\mathcal{Q}} K_H [q]_\pm \, dX, \qquad \mathcal{M}_\pm := \frac{3}{8} \int_{\mathcal{Q}} K_H \mathcal{C} H^\pm \, dX,$$

where

$$\mathcal{C} = \frac{5v(1-2s) + (4s-1)\mathfrak{A}}{1+s}, \qquad s = y/x.$$

Then

$$\dot{a}^+ = \mathcal{S}_+ + \mathcal{M}_+, \qquad \dot{a}^- = \mathcal{S}_- + \mathcal{M}_-.$$

With $C_\beta = (\Theta + \frac{x}{2}\Theta_x)/y$, $d\mu_c = dX/\mathfrak{r}$, $\mathcal{Q}_{\mathrm{sg}}^{(1)} = \|q\|_{L^1(d\mu_c)} + \|C_\beta\|_{L^1(d\mu_c)}$, and $\mathcal{L}_B = \|\nabla_{\mathbb{R}^7}(-v\mathbf{X}, g\mathbf{Z})\|_\infty$,

$$|\mathcal{M}_\pm| \le \mathfrak{c} \mathcal{L}_B a^\pm, \qquad \mathcal{S}_+ + \mathcal{S}_- \le \mathfrak{c} \mathcal{Q}_{\mathrm{sg}}^{(1)}.$$

Proof. The genealogy equations are

$$\partial_t H^\pm + \nabla \cdot (\mathbf{b} H^\pm) = 2[q]_\pm, \qquad \mathbf{b} = (-2xv, 2yg).$$

The squared-coordinate incompressibility identity gives

$$\nabla \cdot \mathbf{b} = -2v - 2xv_x + 2g + 2yg_y = g.$$

The formulas below are justified directly for the mild genealogies by their characteristic representation; equivalently one may smooth $[q]_\pm$, apply Reynolds' formula, and pass to the limit on compact classical intervals.

Since $a^\pm = (3/8)\int K_H H^\pm dX$ and the fields decay rapidly,

$$\begin{aligned}\dot a^\pm &= \frac{3}{8}\int K_H\{-\nabla\cdot(\mathbf{b}H^\pm) + 2[q]_\pm\}dX\\ &= \frac{3}{8}\int H^\pm \mathbf{b}\cdot\nabla K_H\, dX + \frac{3}{4}\int K_H[q]_\pm\, dX.\end{aligned}$$

For $K_H = x\sqrt{y}/(x+y)^{5/2}$ a direct logarithmic differentiation yields

$$\frac{\mathbf{b}\cdot\nabla K_H}{K_H} = \frac{(3-2s)v + (1-4s)g}{1+s} = \frac{5v(1-2s) + (4s-1)\mathfrak{A}}{1+s} = \mathcal{C},$$

where $g = 2v - \mathfrak{A}$. This proves the two exact bank equations.

The rational coefficients in $\mathcal{C}$ are uniformly bounded for $s > 0$. Tangential derivatives of the lifted field $\mathbf{B} = (-v\mathbf{X}, g\mathbf{Z})$ recover v and g, so $|\mathcal{C}| \le \mathfrak{c}\mathcal{L}_B$. Positivity of $K_H H^\pm$ therefore gives

$$|\mathcal{M}_\pm| \le \mathfrak{c}\mathcal{L}_B a^\pm.$$

Finally

$$\mathcal{S}_+ + \mathcal{S}_- = \frac{3}{4}\int K_H|q|\, dX = \frac{3}{4}\int (K_H\mathfrak{r})|q|\, d\mu_c.$$

The homogeneous angular function $K_H\mathfrak{r}$ is bounded on $\mathcal{Q}$, hence

$$\mathcal{S}_+ + \mathcal{S}_- \le \mathfrak{c}\|q\|_{L^1(d\mu_c)} \le \mathfrak{c}\mathcal{Q}^{(1)}_{\mathrm{sg}}.$$

□

Audit status: GREEN/COMPLETE.

C2. Finite-action floor equals cancellation lock. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 10.3.

> **Restatement of theorem 10.3.** Let $[t_0, T)$ be finite and classical, with
>
> $$\int_{t_0}^{T}\mathcal{L}_\star(t)dt < \infty, \qquad \sup_{t_0\le t<T}\mathcal{Q}^{(1)}_{\mathrm{sg}}(t) < \infty,$$
>
> where $\mathcal{L}_\star = \mathcal{L}_B + 4\|yv_y\|_\infty + \|xg_x\|_\infty$. Then $a^\pm$ have finite terminal limits; if $t_0 > 0$, the positive limit is strictly positive. If also $a > 0$ and $a \to 0$, then
>
> $$a^+(t) \to L, \qquad a^-(t) \to L, \qquad L > 0.$$

Proof. Put $\mathcal{Q}^{(1)}_{*,\mathrm{sg}} := \sup_{[t_0,T)}\mathcal{Q}^{(1)}_{\mathrm{sg}}$. By C1,

$$|\dot a^\pm| \le \mathfrak{c}\mathcal{Q}^{(1)}_{*,\mathrm{sg}} + \mathfrak{c}\mathcal{L}_B a^\pm.$$

Because $T - t_0 < \infty$ and $\int_{t_0}^T \mathcal{L}_B < \infty$, Gronwall gives $\sup_{[t_0,T)} a^\pm < \infty$. The same inequality then yields

$$\int_{t_0}^{T}|\dot a^\pm|dt \le \mathfrak{c}(T-t_0)\mathcal{Q}^{(1)}_{*,\mathrm{sg}} + \mathfrak{c}\sup a^\pm\int_{t_0}^{T}\mathcal{L}_B dt < \infty.$$

Thus both banks have finite one-sided terminal limits.

For the positive bank, $\mathcal{S}_+ \geq 0$ and C1 imply

$$\dot{a}^+ \geq -\mathfrak{c}\mathcal{L}_B a^+, \qquad a^+(t) \geq a^+(t_0)\exp\left(-\mathfrak{c}\int_{t_0}^{t}\mathcal{L}_B(\tau)d\tau\right).$$

It remains only to justify $a^+(t_0) > 0$ for $t_0 > 0$. Write $p(t) = u_y(0,0,t) = \psi_\beta(t)/2$ and $q_H(t) = H_y(0,0,t)$. The detector equation gives $p(t) \neq 0$ for finite classical time. Comparing the first y-coefficients in $\mathcal{D}_{xy}H + gH = 2\Theta_y$ at the apex gives

$$\dot{q}_H + 6aq_H = 4\varepsilon_\beta p^2 = \mathcal{J}_\beta, \qquad q_H(0) = 0.$$

Hence $q_H(t) > 0$ for every $t > 0$ in the physical sector. To see directly that the negative genealogy has zero first equatorial jet, factor the smooth equatorial- vanishing swirl as $u = y\widehat{w}$. Then $q = \Theta_y = 2\varepsilon_\beta y\widehat{w}(\widehat{w} + y\widehat{w}_y)$, whose negative part is $O(y^2)$ because the parenthesis equals $\widehat{w}^2 \geq 0$ at $y = 0$. The negative genealogy equation therefore gives $H^- = O(y^2)$ near the equator. Hence $H^+ = q_H(t_0)y + O(|(x,y)|^2)$ and is positive on a sufficiently small conical annulus. Since $K_H > 0$ in the open quadrant, $a^+(t_0) > 0$. Consequently the terminal positive limit L_+ is strictly positive.

If finally $a = a^+ - a^- \to 0$, the finite terminal limits satisfy $L_+ - L_- = 0$. Thus $L_+ = L_- =: L > 0$, proving the cancellation-lock conclusion. □

Audit status: GREEN/COMPLETE.

C3. Exact cancellation ledger. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 10.5.

> **Restatement of theorem 10.5.** On the finite-action floor branch of C2, choose t_- sufficiently late that $a^\pm > 0$ and define
>
> $$\upsilon_- = a^-/a^+, \qquad \sigma_\pm = \mathcal{S}_\pm/a^\pm, \qquad c_\pm = \mathcal{M}_\pm/a^\pm.$$
>
> Then
>
> $$\frac{d}{dt}\log\upsilon_- = \sigma_- - \sigma_+ + c_- - c_+,$$
>
> all four own-bank actions are integrable on $[t_-, T)$, and
>
> $$\int_{t_-}^{T}(\sigma_- - \sigma_+ + c_- - c_+)dt = \log(1/\upsilon_-(t_-)) < \infty.$$

Proof. C2 gives $a^\pm(t) \to L > 0$. Choose t_- so late that both banks are strictly positive thereafter. Dividing the exact C1 equations by their own banks gives

$$\frac{d}{dt}\log a^\pm = \sigma_\pm + c_\pm.$$

Subtracting proves the differential identity for $\log\upsilon_-$.

The contraction terms satisfy

$$|c_\pm| = \frac{|\mathcal{M}_\pm|}{a^\pm} \leq \mathfrak{c}\mathcal{L}_B,$$

so $\int_{t_-}^{T}|c_\pm| < \infty$. Integrating the logarithmic bank equations gives

$$\int_{t_-}^{t}\sigma_\pm d\tau = \log\frac{a^\pm(t)}{a^\pm(t_-)} - \int_{t_-}^{t}c_\pm d\tau.$$

The right-hand side has a finite limit as $t \uparrow T$; since $\sigma_\pm \geq 0$, each source action is finite. This proves the absolute integrability statement in the main theorem.

Finally C2 gives $\upsilon_-(t) \to 1$. Integrating the logarithmic ratio identity and taking $t \uparrow T$ yields

$$\int_{t_-}^{T} (\sigma_- - \sigma_+ + c_- - c_+)dt = \log \frac{1}{\upsilon_-(t_-)}.$$

Thus the floor lock costs finite own-bank source/geometry work; no $1/a$ factor is created. □

Audit status: GREEN/COMPLETE.

C4. Fresh positive genealogy near the apex. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 11.1.

> **Restatement of proposition 11.1.** In the physical sector $\varepsilon_\beta = 1$, for the oriented seed $\psi_\beta(0) > 0$ and every finite classical $t_0 > 0$, with $s = y/x$ and $q_H(t) = H_y(0,0,t)$,
>
> $$q_H(t_0) > 0, \qquad H^+(x, sx, t_0) = q_H(t_0)sx + O_{t_0,J}(x^2), \qquad H^-(x, sx, t_0) = O_{t_0,J}(x^2)$$
>
> uniformly for $s \in J \Subset (0, \infty)$.

Proof. The material equator $y = 0$ preserves $u = 0$ and $H = 0$. Let $p(t) := u_y(0,0,t) = \psi_\beta(t)/2$ and $q_H(t) := H_y(0,0,t)$. Since $\psi_\beta(t) = \psi_\beta(0)e^{-2\int_0^t a}$, the oriented seed gives $p(t) > 0$ at every finite classical time. Expand the exact equations at $x = y = 0$. The coefficient of y in $\mathcal{D}_{xy}H + gH = 2\Theta_y$ satisfies

$$\dot{q}_H + 6aq_H = 4\varepsilon_\beta p^2 = \mathcal{J}_\beta, \qquad q_H(0) = 0.$$

The integrating-factor formula therefore gives $q_H(t) > 0$ for every $t > 0$.

To control the negative genealogy, smoothness and $u(x,0,t) = 0$ allow the factorization $u(x,y,t) = y\widehat{w}(x,y,t)$. Hence

$$q = \Theta_y = 2\varepsilon_\beta y\widehat{w}(\widehat{w} + y\widehat{w}_y) =: 2\varepsilon_\beta y\mathcal{F}_\Theta.$$

On the equator $\mathcal{F}_\Theta(x,0,t) = \widehat{w}(x,0,t)^2 \geq 0$. Therefore $[\mathcal{F}_\Theta]_- = 0$ at $y = 0$, and local Lipschitz regularity implies $[\mathcal{F}_\Theta]_- = O(y)$. Consequently $[q]_- = O(y^2)$ near the equator.

Writing $H^- = yk^-$ in its continuity equation gives

$$\mathcal{D}_{xy}k^- + 3gk^- = 2[q]_-/y, \qquad k^-(\cdot, 0) = 0.$$

The right side is $O(y)$, so $k^-(x,0,t) = 0$ and $k^- = O(y)$ on compact classical sets. Thus $H^- = O(y^2)$, uniformly on compact cones. Since $H(x,0,t) = 0$ identically in x, one also has $H_x(0,0,t) = 0$, and Taylor expansion gives

$$H(x, sx, t_0) = q_H(t_0)sx + O_{t_0,J}(x^2).$$

Finally $H^+ = H + H^-$, proving the asserted expansion and positivity on sufficiently small conical annuli. □

Audit status: GREEN/COMPLETE.

C5. Positive-genealogy material cashier. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 11.2.

> **Restatement of proposition 11.2.** For a compact material block $\mathcal{B}(t) = \mathscr{X}_{t,t_0}(\mathcal{B}_0)$ with $\mathcal{B}_0 \Subset \mathcal{Q}$,
>
> $$M_{\mathcal{B}}^+ = \int_{\mathcal{B}} H^+/x, \qquad a_{\mathcal{B}}^+ = \frac{3}{8}\int_{\mathcal{B}} K_H H^+,$$
>
> and
>
> $$\dot{M}_{\mathcal{B}}^+ = 2\int_{\mathcal{B}} \upsilon H^+/x + 2\int_{\mathcal{B}} [q]_+/x.$$

The source inflow is nonnegative; if $a, M_{\mathcal{B}}^{+} > 0$ this is equivalently $\dot{M}_{\mathcal{B}}^{+} = 2a\Lambda_{\mathcal{B}}^{+} M_{\mathcal{B}}^{+}$.

Proof. For a material domain Reynolds' formula gives, with $f = H^{+}/x$,

$$\frac{d}{dt}\int_{\mathcal{B}(t)} f\, dX = \int_{\mathcal{B}(t)} (\mathcal{D}_{xy} f + f\nabla\cdot\mathbf{b})\, dX.$$

As in C1, $\nabla\cdot\mathbf{b} = g$. Moreover $\mathcal{D}_{xy}(x^{-1}) = 2v/x$ and $\mathcal{D}_{xy}H^{+} + gH^{+} = 2[q]_{+}$. Therefore

$$\mathcal{D}_{xy}(H^{+}/x) + gH^{+}/x = 2vH^{+}/x + 2[q]_{+}/x,$$

which is exactly (11.3). The second integral is nonnegative, so negative source history cannot subtract from this positive-genealogy account. Dividing the numerator in the definition of $\Lambda_{\mathcal{B}}^{+}$ by $aM_{\mathcal{B}}^{+}$ when the latter is positive gives (11.4). □

Audit status: GREEN/COMPLETE.

C6. Credit-reset firewall. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 11.3.

> **Restatement of proposition 11.3.** Let $\mathcal{B}(t)$ be material and initially lie in $x_{\text{launch}}/2 < x < x_{\text{launch}}$. If $0 < \theta < 1/2$ and $M_{\mathcal{B}}^{+}(t) < \theta M_{\mathcal{B}}^{+}(t_0)$, then one trajectory satisfies
>
> $$\int_{t_0}^{t} [-v(X(\tau),\tau)]_{+} d\tau \geq \frac{1}{2}\log\frac{1}{2\theta}.$$
>
> If $a > 0$ on the interval, the same amount is paid by the corrected logarithmic star-hull restart basis in (11.6).

Proof. Define the unweighted positive-genealogy bank

$$N_{\mathcal{B}}^{+}(t) := \int_{\mathcal{B}(t)} H^{+} dX.$$

Integrating the positive-genealogy continuity equation over the material block gives

$$\dot{N}_{\mathcal{B}}^{+} = 2\int_{\mathcal{B}(t)} [q]_{+} dX \geq 0.$$

Let $x_{\mathcal{B}}^{\max}(t) := \sup\{x : (x,y) \in \mathcal{B}(t)\}$. Positivity of H^{+} gives

$$M_{\mathcal{B}}^{+}(t) \geq \frac{N_{\mathcal{B}}^{+}(t)}{x_{\mathcal{B}}^{\max}(t)} \geq \frac{N_{\mathcal{B}}^{+}(t_0)}{x_{\mathcal{B}}^{\max}(t)}.$$

At launch, $x > x_{\text{launch}}/2$, hence

$$M_{\mathcal{B}}^{+}(t_0) \leq \frac{2}{x_{\text{launch}}} N_{\mathcal{B}}^{+}(t_0).$$

Thus

$$M_{\mathcal{B}}^{+}(t) \geq \frac{M_{\mathcal{B}}^{+}(t_0)}{2}\frac{x_{\text{launch}}}{x_{\mathcal{B}}^{\max}(t)}.$$

If $M_{\mathcal{B}}^{+}(t) < \theta M_{\mathcal{B}}^{+}(t_0)$, then $x_{\mathcal{B}}^{\max}(t) > x_{\text{launch}}/(2\theta)$. Choosing a trajectory that attains the supremum (or passing to a maximizing sequence) and using its initial coordinate $x(t_0) < x_{\text{launch}}$ gives

$$\frac{x(t)}{x(t_0)} > \frac{1}{2\theta}.$$

Since $\dot{x} = -2vx$,

$$\log \frac{x(t)}{x(t_0)} = -2\int_{t_0}^{t} v(X(\tau),\tau)d\tau \le 2\int_{t_0}^{t} [-v(X(\tau),\tau)]_+ d\tau,$$

which proves (11.5).

Assume now $a > 0$. The corrected radial fundamental theorem is

$$v(X,\tau) - a(\tau) = \int_0^1 \mathcal{R}_v(\lambda X,\tau)\frac{d\lambda}{\lambda}, \qquad \mathcal{R}_v = xv_x + yv_y.$$

Therefore, whenever $v(X,\tau) < 0$,

$$[-v(X,\tau)]_+ \le [a - v(X,\tau)]_+ \le \int_0^1 [-\mathcal{R}_v(\lambda X,\tau)]_+ \frac{d\lambda}{\lambda}.$$

The exact restart decomposition gives

$$[-\mathcal{R}_v]_+ \le \frac{1}{4}[\mathfrak{A}]_+ + \frac{1}{2}[-\mathfrak{J}]_+ + \frac{1}{4}yH^+ + \frac{1}{2}s[\mathfrak{T}]_+.$$

Apply this at $X_\lambda(\tau) = \lambda X(\tau)$; the angular ratio s is unchanged by radial scaling and y becomes $y_\lambda = \lambda y$. Integrating first in λ with $d\lambda/\lambda$ and then in time, and using the already proved negative-v action, yields (11.6). □

Audit status: GREEN/COMPLETE.

C7. Intrinsic positive-genealogy Riccati closure. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 12.2.

> **Restatement of theorem 12.2.** Let $\mathcal{B}(t) = \mathscr{X}_{t,t_0}(\mathcal{B}_0)$ be a compact material block with $\mathcal{B}_0 \Subset \mathcal{Q}$, and let $M^+_{\mathcal{B}}, a^+_{\mathcal{B}}, \eta^+_{\mathcal{B}}, \Lambda^+_{\mathcal{B}}$ be the cashier, Hodge credit, visibility, and intrinsic rate defined in definition 12.1. Suppose on a classical interval
>
> $$\boxed{a > 0, \qquad a \ge a^+_{\mathcal{B}}, \qquad \eta^+_{\mathcal{B}} \ge m_* > 0, \qquad \Lambda^+_{\mathcal{B}} \ge \nu > 0.}$$
>
> Then
>
> $$\boxed{\dot{M}^+_{\mathcal{B}} \ge 2\nu m_* (M^+_{\mathcal{B}})^2.}$$
>
> Hence either regularity fails earlier or
>
> $$T_* \le t_0 + \frac{1}{2\nu m_* M^+_{\mathcal{B}}(t_0)}, \qquad M^+_{\mathcal{B}}(t) \to \infty \quad (t \uparrow T_*),$$
>
> provided the displayed gates persist. No sign assumption on the current signed field H is required.

Proof. By definition 12.1,

$$\dot{M}^+_{\mathcal{B}} = 2a\Lambda^+_{\mathcal{B}} M^+_{\mathcal{B}}.$$

The rate gate therefore yields

$$\dot{M}^+_{\mathcal{B}} \ge 2\nu a M^+_{\mathcal{B}}.$$

The Hodge and visibility gates give

$$a \ge a^+_{\mathcal{B}} = \eta^+_{\mathcal{B}} M^+_{\mathcal{B}} \ge m_* M^+_{\mathcal{B}}.$$

Combining the two inequalities proves

$$\dot{M}^+_{\mathcal{B}} \ge 2\nu m_* (M^+_{\mathcal{B}})^2.$$

Writing $W = M_{\mathcal{B}}^{+}$ and differentiating W^{-1} gives

$$\frac{d}{dt}\frac{1}{W} = -\frac{\dot{W}}{W^2} \le -2\nu m_*.$$

Hence

$$\frac{1}{M_{\mathcal{B}}^{+}(t)} \le \frac{1}{M_{\mathcal{B}}^{+}(t_0)} - 2\nu m_*(t - t_0),$$

so the right side reaches zero no later than

$$T_* = t_0 + \frac{1}{2\nu m_* M_{\mathcal{B}}^{+}(t_0)}.$$

It remains to justify that divergence of $M_{\mathcal{B}}^{+}$ is incompatible with a classical solution through T_*. Since $\mathcal{B}_0 \Subset \mathcal{Q}$ is compact and the squared-coordinate velocity field is smooth on every compact classical interval, the material flow is a C^1 diffeomorphism. Along a trajectory,

$$x(t) = x(t_0)\exp\left(-2\int_{t_0}^{t} v\,d\tau\right), \qquad y(t) = y(t_0)\exp\left(2\int_{t_0}^{t} g\,d\tau\right),$$

so bounded v, g keep the transported compact set a positive distance from $x = 0$ and $y = 0$ on every finite classical interval. The positive genealogy solves a linear continuity equation with bounded coefficients and bounded source $2[q]_+$; its characteristic formula therefore keeps H^+ bounded on the transported compact set. Consequently H^+/x is bounded there and the material volume is finite, so $M_{\mathcal{B}}^{+}$ cannot diverge while classical regularity persists. Thus either regularity fails earlier or the Riccati deadline is reached. □

Audit status: GREEN/COMPLETE.

C8. Rate failure pays radial non-affinity. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 12.3.

> **Restatement of proposition 12.3.** Let $\mathcal{B}(t)$, $M_{\mathcal{B}}^{+}$, and $\Lambda_{\mathcal{B}}^{+}$ be as in definition 12.1. Fix $0 < \nu < 1$, put $\mathcal{R}_v := xv_x + yv_y$, and suppose $\Lambda_{\mathcal{B}}^{+} < \nu$. Then
>
> $$\boxed{\int_{\mathcal{B}} (a - v)\frac{H^+}{x} dX > (1 - \nu)aM_{\mathcal{B}}^{+} + \int_{\mathcal{B}} \frac{[q]_+}{x} dX.}$$
>
> Consequently, using the corrected logarithmic radial measure,
>
> $$\boxed{\int_{\mathcal{B}}\int_0^1 [-\mathcal{R}_v]_+(\lambda X, t)\frac{d\lambda}{\lambda}\frac{H^+}{x} dX > (1 - \nu)aM_{\mathcal{B}}^{+}.}$$
>
> Thus an intrinsic rate exit is a paid logarithmic star-hull bill.

Proof. Set

$$V_{\mathcal{B}}^{+} := \int_{\mathcal{B}} v\frac{H^+}{x}\,dX, \qquad S_{\mathcal{B}}^{+} := \int_{\mathcal{B}} \frac{[q]_+}{x}\,dX \ge 0.$$

The definition of the intrinsic rate is

$$\Lambda_{\mathcal{B}}^{+} = \frac{V_{\mathcal{B}}^{+} + S_{\mathcal{B}}^{+}}{aM_{\mathcal{B}}^{+}}.$$

Therefore $\Lambda_{\mathcal{B}}^{+} < \nu$ implies

$$V_{\mathcal{B}}^{+} + S_{\mathcal{B}}^{+} < \nu aM_{\mathcal{B}}^{+}.$$

Subtracting $V_{\mathcal{B}}^{+}$ from $aM_{\mathcal{B}}^{+}$ gives

$$\int_{\mathcal{B}}(a-v)\frac{H^{+}}{x}\,dX = aM_{\mathcal{B}}^{+} - V_{\mathcal{B}}^{+} > (1-\nu)aM_{\mathcal{B}}^{+} + S_{\mathcal{B}}^{+},$$

which is the first claim.

For each fixed $X \in \mathcal{Q}$, let $X_\lambda = \lambda X$. The corrected radial fundamental theorem is

$$v(X,t) - a(t) = \int_0^1 \mathcal{R}_v(X_\lambda, t)\frac{d\lambda}{\lambda}.$$

The integral is regular at $\lambda = 0$ because smoothness gives $\mathcal{R}_v(\lambda X) = O(\lambda)$ on compact rays. Hence

$$a - v(X,t) = \int_0^1 -\mathcal{R}_v(X_\lambda, t)\frac{d\lambda}{\lambda} \le \int_0^1 [-\mathcal{R}_v]_+(X_\lambda, t)\frac{d\lambda}{\lambda}.$$

Multiply by the nonnegative density $H^{+}(X,t)/x$ and integrate over $\mathcal{B}(t)$. Tonelli's theorem applies to the nonnegative majorant and yields

$$\int_{\mathcal{B}}(a-v)\frac{H^{+}}{x}\,dX \le \int_{\mathcal{B}}\int_0^1 [-\mathcal{R}_v]_+(\lambda X, t)\frac{d\lambda}{\lambda}\frac{H^{+}}{x}\,dX.$$

Combining with the first strict inequality proves the logarithmic star-hull bill. □

Audit status: GREEN/COMPLETE.

C9. Visibility collapse is positive-genealogy tail concentration. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 12.4.

Restatement of theorem 12.4. Let $M_{\mathcal{B}}^{+}$ and $\eta_{\mathcal{B}}^{+}$ be as in definition 12.1, and put

$$h_0(s) := \frac{3}{8}\frac{\sqrt{s}}{(1+s)^{5/2}}, \qquad h_{\max} := \max_{s>0} h_0(s) = \frac{6}{25\sqrt{5}}.$$

Define

$$d\mu_{\mathcal{B}}^{+} := \frac{H^{+}}{x}\,dX, \qquad \mu_{\mathcal{B}}^{+}(\mathcal{Q}) = M_{\mathcal{B}}^{+}.$$

If $\eta_{\mathcal{B}}^{+} < m_* < h_{\max}$, then for every $\vartheta \in (m_*, h_{\max})$,

$$\boxed{\mu_{\mathcal{B}}^{+}\{X \in \mathcal{B} : h_0(y/x) < \vartheta\} > \left(1 - \frac{m_*}{\vartheta}\right) M_{\mathcal{B}}^{+}.}$$

Thus visibility failure moves a fixed fraction of already banked positive genealogy into the angular tails; it is not disappearance of credit.

Proof. Since

$$K_H(x,y) = \frac{1}{x}\frac{\sqrt{s}}{(1+s)^{5/2}}, \qquad s = \frac{y}{x},$$

the Hodge credit of the block can be written

$$a_{\mathcal{B}}^{+} = \int_{\mathcal{B}} h_0(y/x)\,d\mu_{\mathcal{B}}^{+}.$$

Therefore

$$\eta_{\mathcal{B}}^{+} = \frac{a_{\mathcal{B}}^{+}}{M_{\mathcal{B}}^{+}} = \frac{1}{M_{\mathcal{B}}^{+}}\int_{\mathcal{B}} h_0(y/x)\,d\mu_{\mathcal{B}}^{+}.$$

For completeness,

$$\frac{d}{ds}\log\left(\frac{\sqrt{s}}{(1+s)^{5/2}}\right) = \frac{1}{2s} - \frac{5}{2(1+s)},$$

so the unique critical point is $s = 1/4$, where $h_0 = 6/(25\sqrt{5})$; this is the global maximum because h_0 tends to zero at both angular ends.

Fix $\vartheta \in (m_*, h_{\max})$ and set

$$A_\vartheta := \{X \in \mathcal{B} : h_0(y/x) \geq \vartheta\}.$$

Positivity gives

$$\eta_{\mathcal{B}}^+ M_{\mathcal{B}}^+ = \int_{\mathcal{B}} h_0 \, d\mu_{\mathcal{B}}^+ \geq \int_{A_\vartheta} h_0 \, d\mu_{\mathcal{B}}^+ \geq \vartheta \, \mu_{\mathcal{B}}^+(A_\vartheta).$$

Since $\eta_{\mathcal{B}}^+ < m_*$,

$$\mu_{\mathcal{B}}^+(A_\vartheta) < \frac{m_*}{\vartheta} M_{\mathcal{B}}^+.$$

Taking the complement in $\mathcal{B}$ yields

$$\mu_{\mathcal{B}}^+\{h_0(y/x) < \vartheta\} > \left(1 - \frac{m_*}{\vartheta}\right) M_{\mathcal{B}}^+,$$

as claimed. The conclusion is a concentration statement for the existing nonnegative genealogy measure; no credit is erased. □

Audit status: GREEN/COMPLETE.

C10. Finite intrinsic exit graph. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 12.5.

Restatement of theorem 12.5. Fix

$$0 < \nu < 1, \qquad 0 < m_* < h_{\max},$$

and an admitted classical interval on which $a > 0$ and $M_{\mathcal{B}}^+ > 0$ for a material positive-genealogy cashier $\mathcal{B}(t)$. Use $M_{\mathcal{B}}^+, a_{\mathcal{B}}^+, \eta_{\mathcal{B}}^+, \Lambda_{\mathcal{B}}^+$ from definition 12.1. Exactly the following intrinsic branches remain:

(i) $a \geq a_{\mathcal{B}}^+$, $\eta_{\mathcal{B}}^+ \geq m_*$, and $\Lambda_{\mathcal{B}}^+ \geq \nu$ persist, and Riccati closure occurs;
(ii) $\Lambda_{\mathcal{B}}^+ < \nu$, which pays radial star-hull non-affinity;
(iii) $\eta_{\mathcal{B}}^+ < m_*$, which places positive genealogy in the angular tails;
(iv) $a < a_{\mathcal{B}}^+$, which activates H^- and hence routes to old negative ancestry or tagged cross-shear.

There is no source-sign reset and no ancestry-free fifth intrinsic branch. Loss of the outer admission $a > 0$ is a separate compression-floor exit.

Proof. Work on an admitted interval on which $a > 0$ and $M_{\mathcal{B}}^+ > 0$, so all intrinsic quantities are defined. At each time there are only two possibilities for each of the three inequalities

$$a \geq a_{\mathcal{B}}^+, \qquad \eta_{\mathcal{B}}^+ \geq m_*, \qquad \Lambda_{\mathcal{B}}^+ \geq \nu.$$

If all three persist from the launch time onward, theorem 12.2 gives the finite Riccati deadline, which is branch (i).

If the rate inequality is the failed gate, branch (ii) follows from proposition 12.3. If the visibility inequality fails, branch (iii) follows from theorem 12.4.

It remains to route failure of Hodge dominance. Decompose the global positive bank into the block and its complement:

$$a^+ = a_{\mathcal{B}}^+ + a_{\mathcal{B}^c}^+, \qquad a = a^+ - a^-.$$

If $a < a_{\mathcal{B}}^+$, then

$$a^- - a_{\mathcal{B}^c}^+ = a_{\mathcal{B}}^+ - a > 0,$$

so in particular $a^- > 0$ and the negative genealogy is nontrivial. The exact slope-ancestry statement proposition 13.1 then routes negative source history to the fixed initial negative slope or to the tagged cross-shear channel; there is no independent third ancestry.

Finally, the positive genealogy itself obeys a source-history continuity equation with nonnegative source $2[q]_+$, so a later sign change of the signed source cannot subtract previously accumulated H^+ genealogy. Hence there is no source-sign reset. The three intrinsic gate failures together with the persistent-gate branch exhaust the intrinsic cashier graph. If a itself ceases to be positive, the intrinsic rate is no longer admitted; that event belongs to the separate global compression-floor exit rather than to a hidden fifth intrinsic branch. □

Audit status: GREEN/COMPLETE.

C11. Hodge-matched Cauchy bridge. *Mutual pointers:* grand-finale entry / short proof/status / main statement lemma 12.7.

Restatement of lemma 12.7. Let $K_{\mathcal{B}}, h_0, a_K, N_K, \Pi_H$ be the frozen cohort and its three moments from definition 12.6. Then

$$\boxed{N_K^2 \le a_K \Pi_H.}$$

Whenever $\Pi_H > 0$, define

$$\mathfrak{Z}_H := \frac{N_K^2}{\Pi_H}.$$

Then $\mathfrak{Z}_H \le a_K$.

Proof. Because $\mathcal{B}_0 \Subset \mathcal{Q}$ and $\mathcal{B}(t)$ is its material image on a classical interval, $x > 0$ and $s = y/x > 0$ on the compact support of $K_{\mathcal{B}}$. Thus the two nonnegative weights below are well defined and the stated moments are finite. Factor the conserved cohort mass as

$$N_K = \int_{\mathcal{Q}} K_{\mathcal{B}}\, dX = \int_{\mathcal{Q}} \left(\frac{h_0(s)}{x} K_{\mathcal{B}}\right)^{1/2} \left(\frac{x}{h_0(s)} K_{\mathcal{B}}\right)^{1/2} dX.$$

Cauchy–Schwarz gives

$$N_K^2 \le \left(\int_{\mathcal{Q}} \frac{h_0(s)}{x} K_{\mathcal{B}}\, dX\right) \left(\int_{\mathcal{Q}} \frac{x}{h_0(s)} K_{\mathcal{B}}\, dX\right) = a_K \Pi_H.$$

If $\Pi_H > 0$, division by Π_H is now legitimate and yields

$$\mathfrak{Z}_H = \frac{N_K^2}{\Pi_H} \le a_K.$$

No quotient is used in establishing the Cauchy bridge itself. □

Audit status: GREEN/COMPLETE.

C12. Exact dual-moment rate. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 12.8.

Restatement of proposition 12.8. Let $K_{\mathcal{B}}$, h_0, and Π_H be as in definition 12.6, put $s := y/x$, and recall $\mathfrak{A} := 2v - g$. Define

$$\mathcal{C}_H(x, s, t) := \frac{5v(1-2s) + \mathfrak{A}(4s-1)}{1+s}.$$

Then

$$\boxed{-\dot{\Pi}_H = \int_{\mathcal{Q}} \frac{x K_{\mathcal{B}}}{h_0(s)} \mathcal{C}_H(x, s, t) dX.}$$

Proof. Put

$$w_H(x,y) := \frac{x}{h_0(s)}, \qquad s = \frac{y}{x}, \qquad h_0(s) = \frac{3}{8}\frac{\sqrt{s}}{(1+s)^{5/2}}.$$

The transport operator is $\mathcal{D}_{xy} = \partial_t - 2xv\partial_x + 2yg\partial_y$, so along its characteristics

$$\mathcal{D}_{xy}\log x = -2v, \qquad \mathcal{D}_{xy}\log y = 2g, \qquad \mathcal{D}_{xy}\log s = 2(v+g).$$

Moreover

$$\frac{d\log h_0}{d\log s} = s\frac{d}{ds}\left(\frac{1}{2}\log s - \frac{5}{2}\log(1+s)\right) = \frac{1-4s}{2(1+s)}.$$

Therefore

$$\begin{aligned}\mathcal{D}_{xy}\log w_H &= -2v - \frac{1-4s}{2(1+s)}\,2(v+g)\\ &= -\frac{5v(1-2s)+(2v-g)(4s-1)}{1+s}\\ &= -\mathcal{C}_H(x,s,t).\end{aligned}$$

Equivalently,

$$\mathcal{D}_{xy}w_H = -w_H\mathcal{C}_H.$$

The frozen cohort satisfies the exact continuity equation

$$\partial_t K_{\mathcal{B}} + \nabla\cdot(\mathbf{b}K_{\mathcal{B}}) = 0, \qquad \mathbf{b} = (-2xv, 2yg),$$

and is compactly supported in the material image $\mathcal{B}(t) \Subset \mathcal{Q}$. Hence, by the Reynolds/continuity identity (or integration by parts with no boundary leakage), for every smooth weight w,

$$\frac{d}{dt}\int_{\mathcal{Q}} wK_{\mathcal{B}}\,dX = \int_{\mathcal{Q}} (\mathcal{D}_{xy}w)K_{\mathcal{B}}\,dX.$$

Taking $w = w_H$ gives

$$\dot{\Pi}_H = \int_{\mathcal{Q}} (\mathcal{D}_{xy}w_H)K_{\mathcal{B}}\,dX = -\int_{\mathcal{Q}} \frac{xK_{\mathcal{B}}}{h_0(s)}\mathcal{C}_H(x,s,t)\,dX,$$

which is the claimed identity. Taking $w \equiv 1$ in the same formula also records for later use that

$$\dot{N}_K = 0.$$

□

Audit status: GREEN/COMPLETE.

C13. Hodge-matched Riccati gate. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 12.9.

> **Restatement of theorem 12.9.** Let $K_{\mathcal{B}}$ be a nontrivial frozen cohort from definition 12.6, with moments a_K, N_K, Π_H. Suppose on a time interval
>
> $$-\dot{\Pi}_H \geq \kappa a \Pi_H, \qquad a \geq a_K, \qquad \kappa > 0.$$
>
> Then, before any quotient is formed,
>
> $$\boxed{-N_K^2\dot{\Pi}_H \geq \kappa N_K^4.}$$
>
> Since the cohort is nontrivial, $\Pi_H > 0$ and $N_K > 0$; only after this raw gate is proved define $\mathfrak{Z}_H = N_K^2/\Pi_H$. Then
>
> $$\boxed{0 < \mathfrak{Z}_H \leq a_K \leq a, \qquad \dot{\mathfrak{Z}}_H \geq \kappa a \mathfrak{Z}_H \geq \kappa \mathfrak{Z}_H^2.}$$

If the raw gates persist from t_0, they cannot persist beyond

$$T_{H,\mathrm{Ric}} = t_0 + \frac{1}{\kappa \mathfrak{Z}_H(t_0)}.$$

Thus either a displayed raw gate fails first or the classical solution loses regularity by the finite Hodge-matched Riccati deadline.

Proof. By the Cauchy bridge C11,

$$N_K^2 \le a_K \Pi_H.$$

The Hodge-dominance gate $a \ge a_K$ therefore yields the entirely raw inequality

$$a\Pi_H \ge a_K \Pi_H \ge N_K^2.$$

Multiplying the rate gate $-\dot{\Pi}_H \ge \kappa a \Pi_H$ by $N_K^2 \ge 0$ and then using this lower bound gives

$$-N_K^2 \dot{\Pi}_H \ge \kappa a N_K^2 \Pi_H \ge \kappa N_K^4.$$

This is the guard-safe polynomial gate; no quotient has been introduced.

Because $K_{\mathcal{B}} \ge 0$ is nontrivial and is supported in the interior compact material block, both

$$N_K = \int K_{\mathcal{B}}\, dX > 0, \qquad \Pi_H = \int \frac{x K_{\mathcal{B}}}{h_0(s)}\, dX > 0.$$

We may therefore define $\mathfrak{Z}_H = N_K^2/\Pi_H$. C11 immediately gives $\mathfrak{Z}_H \le a_K \le a$. By C12, or directly from the frozen continuity equation, N_K is constant. Hence

$$\dot{\mathfrak{Z}}_H = -\frac{N_K^2 \dot{\Pi}_H}{\Pi_H^2} \ge \kappa a \frac{N_K^2}{\Pi_H} = \kappa a \mathfrak{Z}_H \ge \kappa \mathfrak{Z}_H^2.$$

Thus

$$\frac{d}{dt}\frac{1}{\mathfrak{Z}_H} = -\frac{\dot{\mathfrak{Z}}_H}{\mathfrak{Z}_H^2} \le -\kappa.$$

If both raw gates persisted on $[t_0, t]$, integration would give

$$\frac{1}{\mathfrak{Z}_H(t)} \le \frac{1}{\mathfrak{Z}_H(t_0)} - \kappa(t - t_0).$$

The right-hand side vanishes at $T_{H,\mathrm{Ric}} = t_0 + [\kappa \mathfrak{Z}_H(t_0)]^{-1}$. Hence the two raw gates cannot persist with a classical solution beyond this time. Equivalently, one raw gate fails earlier or $\mathfrak{Z}_H$ diverges by the deadline. Since $\mathfrak{Z}_H \le a$, divergence of $\mathfrak{Z}_H$ is incompatible with continued classical regularity of the apex compression. □

Audit status: GREEN/COMPLETE.

C14. Automatic fresh relaunch. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 12.10.

Restatement of theorem 12.10. Fix a finite classical time $t_0 > 0$ in the physical sector $\varepsilon_\beta = 1$ and assume $a(t_0) > 0$. Let $J = [s_-, s_+] \Subset (0, 1/2)$, $s = y/x$, and $c_J = 5 \min_{s \in J}(1 - 2s)/(1 + s)$. Then sufficiently small conical annuli $\mathcal{B}_\ell(t_0) = \{\ell/2 < x < \ell,\ s \in J\}$ have $H^+ > 0$ and $q > 0$ and satisfy

$$H^+(x, sx, t_0) = q_H(t_0) sx + O_{t_0,J}(x^2), \qquad q(x, sx, t_0) = 2u_y(0, 0, t_0)^2 sx + O_{t_0,J}(x^2).$$

If the positive genealogy on $\mathcal{B}_\ell(t_0)$ is frozen into $K_{\mathcal{B}}$, then

$$-\dot{\Pi}_H(t_0) \ge \frac{c_J}{2} a(t_0) \Pi_H(t_0), \qquad a(t_0) - a_K(t_0) \ge \frac{1}{2} a(t_0) > 0.$$

Proof. Put $p(t) := u_y(0,0,t)$. The exact detector equation gives $p(t) > 0$ at every finite classical time for the oriented fixed datum, and the exact apex vorticity-jet equation gives $q_H(t) > 0$ for every $t > 0$. The audited positive-genealogy expansion C4 yields

$$H^+(x, sx, t_0) = q_H(t_0)sx + O_{t_0,J}(x^2)$$

uniformly for $s \in J$. Since $u(0,0,t) = 0$ and $u_y(0,0,t) = p(t)$, Taylor expansion of $q = \partial_y(u^2) = 2uu_y$ gives

$$q(x, sx, t_0) = 2p(t_0)^2 sx + O_{t_0,J}(x^2).$$

The leading coefficients are strictly positive, so $H^+ > 0$ and $q > 0$ on $\mathcal{B}_\ell(t_0)$ once ℓ is small.

For the frozen cohort, C12 gives

$$-\dot{\Pi}_H(t_0) = \int_{\mathcal{B}_\ell(t_0)} \frac{xK_{\mathcal{B}}}{h_0(s)} \mathcal{C}_H(x, s, t_0)\, dX,$$

where

$$\mathcal{C}_H = \frac{5v(1-2s) + (2v-g)(4s-1)}{1+s}.$$

At the apex $v = a$, $g = 2a$, hence uniformly on J,

$$\mathcal{C}_H(x, s, t_0) = 5a(t_0)\frac{1-2s}{1+s} + O_{t_0,J}(x) \geq \frac{c_J}{2} a(t_0)$$

for sufficiently small x. Since $K_{\mathcal{B}} \geq 0$, integration proves the rate gate.

Finally, on the conical annulus $H^+ = O_{t_0,J}(x)$ and $h_0(s)$ is bounded above and below by positive constants. With $dX = x\, dx\, ds$ this gives

$$a_K(t_0) = \int_{\mathcal{B}_\ell(t_0)} \frac{h_0(s)}{x} H^+\, dX = O_{t_0,J}(\ell^2).$$

Because $a(t_0) > 0$ is fixed, decreasing ℓ once more gives $a_K(t_0) \leq a(t_0)/2$. This proves both strict launch margins. Their short-time persistence follows from classical continuity, but no time-uniform persistence is claimed here. □

Audit status: GREEN/COMPLETE.

C15. Uniform apex tube on compact classical intervals. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 12.11.

> **Restatement of theorem 12.11.** Let $I = [t_1, t_2]$ with $0 < t_1 < t_2$ lie strictly inside the classical lifespan and $a_* := \inf_I a > 0$. For every $J \Subset (0, 1/2)$ there is one radius $\ell_* > 0$ for which the conical expansions of v, g, H^+, q have uniform remainders on $I \times J$ and every fresh frozen cohort launched on the common annulus satisfies
>
> $$-\dot{\Pi}_H \geq \frac{c_J}{2} a\Pi_H, \qquad a - a_K \geq \frac{1}{2} a_*.$$

Proof. Choose a fixed apex neighborhood whose closure lies inside the spatial domain on which the classical solution is represented by the regular quotients. Because I is compact and lies strictly inside the classical lifespan, all spatial derivatives of v, g, u needed below are bounded on this compact spacetime set. The one-dimensional fundamental theorem of calculus along $\lambda \mapsto (\lambda x, \lambda sx)$ gives

$$v(x, sx, t) - a(t) = x \int_0^1 (v_x + sv_y)(\lambda x, \lambda sx, t)\, d\lambda,$$

$$g(x,sx,t)-2a(t)=x\int_0^1 (g_x+sg_y)(\lambda x,\lambda sx,t)\,d\lambda.$$

Hence the first line of the asserted expansion is uniform on $I\times J$.

The positive-genealogy quotient used in C4 is classical on compact regular sets, and the negative genealogy has zero first apex jet. Its Taylor formula therefore gives uniformly

$$H^+(x,sx,t)=q_H(t)sx+O_{I,J}(x^2).$$

Likewise $q=2uu_y$, $u(0,0,t)=0$, and $u_y(0,0,t)=p(t)$ give

$$q(x,sx,t)=2p(t)^2sx+O_{I,J}(x^2).$$

The exact equations

$$\dot p=-2ap,\qquad \dot q_H+6aq_H=\mathcal{J}_\beta=\varepsilon_\beta\Psi\ge 0$$

and the positivity of p,q_H for $t>0$ imply by continuity on the compact interval

$$p_*:=\inf_I p(t)>0,\qquad q_*:=\inf_I q_H(t)>0.$$

This is the point at which $t_1>0$ is required.

For the Hodge rate coefficient,

$$\mathcal{C}_H(x,s,t)=5a(t)\frac{1-2s}{1+s}+O_{I,J}(x).$$

Since $a\ge a_*>0$ and $J\Subset(0,1/2)$, one fixed decrease of ℓ_* gives $\mathcal{C}_H\ge(c_J/2)a$ for every launch time. The same uniform Taylor bounds and $p_*,q_*>0$ make H^+ and q strictly positive on the common annulus. Finally

$$\sup_{t\in I}a_K(t)=O_{I,J}(\ell_*^2),$$

so decreasing ℓ_* once more gives $a_K(t)\le a_*/2$ for all $t\in I$. C12 then yields the stated uniform rate gate. All choices are independent of the individual launch time, proving the theorem. □

Audit status: GREEN/COMPLETE.

C16. Terminal-time uniformity firewall. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 12.12.

> **Restatement of theorem 12.12.** On a candidate terminal interval $[t_1,T)$, one time-independent relaunch radius is available only when the local derivative bounds entering C15, the upper positive compression bound $A^*<\infty$, and the positive compression floor $a_*>0$ are all uniform. Failure of a derivative bound or $A^*<\infty$ is a continuation exit; if instead $\inf a=0$, then regularity has already failed or a smooth zero-compression contact forces Hodge-dominance failure and negative-genealogy routing.

Proof. Assume first the stated local derivative bounds and $A^*<\infty$. From the exact apex equations,

$$p(t)=p(t_1)\exp\left(-2\int_{t_1}^t a(\tau)\,d\tau\right)\ge p(t_1)e^{-2A^*(T-t_1)}>0,$$

and, using $\dot q_H+6aq_H=\mathcal{J}_\beta=\varepsilon_\beta\Psi\ge 0$,

$$q_H(t)\ge q_H(t_1)\exp\left(-6\int_{t_1}^t a(\tau)\,d\tau\right)\ge q_H(t_1)e^{-6A^*(T-t_1)}>0.$$

If also $a\ge a_*>0$, every constant used in the proof of C15 is therefore uniform on $[t_1,T)$, so exactly the same choice of ℓ_* works for all $t<T$.

Conversely, suppose one of the local derivative bounds required for those Taylor remainders diverges as $t \uparrow T$, or $A^* = \infty$. A classical extension through T would make the corresponding derivatives and a continuous on a compact spacetime neighborhood and hence bounded. Thus such divergence is a genuine obstruction to classical extension, not an admissible hidden dependence in $O(x)$.

It remains to consider $\inf_{[t_1,T)} a = 0$. Choose t_n with $a(t_n) \to 0$. If no classical extension through T exists, regularity has already failed. Otherwise a subsequence converges to some $t_* \in [t_1, T]$ and continuity gives $a(t_*) = 0$. Since $t_* > 0$, the exact apex-jet equation gives $q_H(t_*) > 0$. C4 therefore supplies a small material block $\mathcal{B}$ with positive genealogy and

$$a^+_{\mathcal{B}}(t_*) = \frac{3}{8} \int_{\mathcal{B}} K_H H^+ \, dX > 0.$$

Hence $a(t_*) = 0 < a^+_{\mathcal{B}}(t_*)$: Hodge dominance fails. Because $a = a^+ - a^-$ and $a^+_{\mathcal{B}} \le a^+$, this implies $a^-(t_*) > 0$, so negative genealogy is present and is routed by the adverse-ancestry branch. These alternatives exhaust the terminal use of the Taylor tube. □

Audit status: GREEN/COMPLETE.

C17. No-Zeno relaunch on a regular positive-compression interval. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 12.13.

> **Restatement of proposition 12.13.** On a compact classical interval $I = [T_0, T_1]$ with $T_0 > 0$ and $a \ge a_* > 0$, fix $J \Subset (0, 1/2)$ and use the common C15 launch annulus. Relaunch only after one of the weaker margins
>
> $$-\dot{\Pi}_H \ge \frac{c_J}{4} a \Pi_H, \qquad a - a_K \ge \frac{1}{4} a_*$$
>
> first fails. Then consecutive relaunch times are separated by one uniform $\tau_* > 0$. Finite-time Zeno therefore forces loss of the compression floor or of a classical smoothness bound.

Proof. By C15, at every launch time $t_0 \in I$ the newly frozen cohort begins with the stronger uniform margins

$$-\dot{\Pi}_H(t_0) \ge \frac{c_J}{2} a(t_0) \Pi_H(t_0), \qquad a(t_0) - a_K(t_0) \ge \frac{1}{2} a_*.$$

The same proof supplies uniform positive lower bounds for H^+ on a fixed subannulus, so the launch values of Π_H have a positive lower bound depending only on I, J, ℓ_*. Thus the numerical slack between the strong launch inequalities and the weaker restart boundaries is uniformly positive.

Transport the cohort by the classical flow. On the compact spacetime region swept out by all such blocks for sufficiently short times, the velocity, its needed spatial derivatives, the Hodge coefficient $\mathcal{C}_H$, and the material-flow Jacobian are uniformly continuous. The moment identities C11–C12 then imply that Π_H, $\dot{\Pi}_H$, and a_K depend uniformly continuously on elapsed time and launch time. Therefore there exists a single $\tau_* > 0$ such that, for every launch $t_0 \in I$, both weaker inequalities remain valid throughout $[t_0, t_0 + \tau_*] \cap I$.

A protocol which relaunches only when one of those weaker inequalities first fails must consequently satisfy $t_{n+1} - t_n \ge \tau_*$. An infinite sequence of such times cannot accumulate inside the finite compact interval. Hence a finite-time accumulation can occur only after leaving the hypotheses that produced the common τ_*: either the positive lower bound for a is lost, or

the local classical smoothness moduli cease to be uniform. C16 identifies those two failures as named terminal exits. □

Audit status: GREEN/COMPLETE.

Appendix D. Adverse ancestry and ordinary continuation capacity

The complete statements are restated here so the ledger slot is self-contained and its ID remains stable.

D1. Exact slope ancestry. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 13.1.

> **Restatement of proposition 13.1.** The protected slope $p = u_y$ obeys the exact material integrating-factor formula (13.2); the adverse source satisfies the two-term majorization (13.6), with old and later cross-shear terms (13.4)–(13.5). Hence there is no third adverse source ancestry.

Proof. Write the squared-coordinate material field as

$$\mathbf{b} = (-2xv, 2yg), \qquad \mathcal{D}_{xy} = \partial_t + \mathbf{b} \cdot \nabla_{x,y}.$$

In the physical sector the swirl equation is

$$\mathcal{D}_{xy} u = 2vu. \tag{D.1}$$

Differentiate in y. For any smooth scalar f,

$$\partial_y(\mathcal{D}_{xy} f) = \mathcal{D}_{xy} f_y + (\partial_y b_1) f_x + (\partial_y b_2) f_y,$$

and here

$$\partial_y b_1 = -2xv_y, \qquad \partial_y b_2 = 2g + 2yg_y.$$

Therefore, with $p = u_y$,

$$\begin{aligned} \mathcal{D}_{xy} p &= \partial_y(2vu) + 2xv_y u_x - (2g + 2yg_y)p \\ &= 2v_y(u + xu_x) + 2(v - g - yg_y)p. \end{aligned}$$

Because $\Gamma = xu$, one has $\Gamma_x = u + xu_x$, so

$$\boxed{\mathcal{D}_{xy} p = 2v_y \Gamma_x + 2(v - g - yg_y)p.} \tag{D.2}$$

Along $X(t) = \mathscr{X}_{t,0} X_0$, set

$$c(t) := 2(v - g - yg_y)(X(t), t), \qquad \mu(t) := \exp\left(- \int_0^t c(\tau)\, d\tau \right) > 0.$$

Then (D.2) becomes

$$\frac{d}{dt}\big(\mu(t) p(X(t), t)\big) = 2\mu(t) v_y(X(t), t) \Gamma_x(X(t), t).$$

Integration from 0 to t gives exactly (13.2).

The fixed datum has $u_0 \geq 0$. Equation (D.1) gives, on each material characteristic,

$$u(X(t), t) = u_0(X_0) \exp\left(2 \int_0^t v(X(\tau), \tau)\, d\tau \right) \geq 0.$$

Since $q = \Theta_y = 2\varepsilon_\beta up$,

$$[q]_- = 2\varepsilon_\beta u [-p]_+.$$

Apply the elementary inequality

$$[-(A+B)]_+ \le [-A]_+ + [-B]_+$$

to the bracket in (13.2). Positivity of μ gives

$$[-p(X(t),t)]_+ \le \mu(t)^{-1}\left([-p_0]_+ + 2\int_0^t \mu(\tau)[-v_y\Gamma_x]_+(X(\tau),\tau)\,d\tau\right).$$

Multiplication by $2\varepsilon_\beta u(X(t),t)$ is precisely (13.4)–(13.6). Thus every adverse source particle is inherited either from the fixed initial negative slope or from later negative cross-shear. No third term occurs in the exact slope equation. □

Audit status: GREEN/COMPLETE.

D2. Old source is geometrically separated. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 13.2.

> **Restatement of proposition 13.2.** For the fixed $A = 10$ customer, the initial negative-slope ancestry lies in $y \ge \sigma_- > 0$ and has finite critical exposure. If that material ancestry stays inside a fixed near-affine corridor, its exact source-response payment satisfies (13.14)–(13.16); otherwise its first corridor failure is a named value/first-jet restart exit that enters the already established tangent/star-hull routing.

Proof. **Step 1: exact separation of the fixed old source.** From definition 5.1,

$$u_0(x,y) = y\left(1+\frac{1}{5}y\right)e^{-c_*x-s_*y} + 10yx^2e^{-2x-b_*y}, \qquad b_* = 6+2\sqrt{7}.$$

Hence

$$\begin{aligned} u_{0,y}(x,y) = e^{-c_*x-s_*y}\left[1+\left(\frac{2}{5}-s_*\right)y - \frac{s_*}{5}y^2\right] \\ + 10x^2e^{-2x-b_*y}(1-b_*y). \end{aligned} \tag{D.3}$$

Put $\sigma_- = b_*^{-1} = (3-\sqrt{7})/4$. If $0 \le y \le \sigma_-$, the second line of (D.3) is nonnegative. Since $s_* < 1$ and $\sigma_- < 1/6$, the first bracket obeys

$$1+\left(\frac{2}{5}-s_*\right)y - \frac{s_*}{5}y^2 > 1-y-\frac{1}{5}y^2 > 1-\frac{1}{6}-\frac{1}{180} > 0.$$

Therefore $u_{0,y} \ge 0$ on $0 \le y \le \sigma_-$, proving (13.8).

At $t = 0$,

$$\Xi_{\beta,0} = \Theta_0\frac{x}{y^3} = \varepsilon_\beta\frac{xw_0(x,y)^2}{y}.$$

On $\mathcal{N}_0$ the denominator is bounded below by $\sigma_- > 0$, while w_0 is smooth and exponentially decaying in both variables. Thus $M_{3,\mathcal{N}}(0) < \infty$, proving (13.10).

Step 2: an all-age response-rate bound without reusing protected names. Fix a deposition time τ and first take $X_\tau = (x_\tau, y_\tau) \in \mathcal{N}(\tau)$ with $x_\tau > 0$. The set $x_\tau = 0$ is Lebesgue-null in the source integrals and has zero critical-exposure density because Ξ_β contains the factor x; the estimates below therefore extend to the full material ancestry by the usual null-set convention. Put

$$s_\tau := \frac{y_\tau}{x_\tau}, \qquad \Lambda(t;\tau) := \int_\tau^t a(\xi)\,d\xi.$$

Along the corridor, the value bounds give

$$x(t) \geq x_\tau e^{-(2+2\delta)\Lambda(t;\tau)}, \tag{D.4}$$

$$\frac{y(t)}{x(t)} \geq s_\tau e^{(6-2\delta)\Lambda(t;\tau)}. \tag{D.5}$$

Let $m = (m_1, m_2) = D_X \mathscr{X}_{t,\tau}(X_\tau) e_y$ be the vertical tangent vector and introduce the logarithmic sensitivities

$$\zeta_1(t) := y_\tau \frac{m_1(t)}{x(t)}, \qquad \zeta_2(t) := y_\tau \frac{m_2(t)}{y(t)}. \tag{D.6}$$

Direct differentiation cancels the zeroth-order affine stretching and yields

$$\frac{d}{dt}\begin{pmatrix}\zeta_1\\ \zeta_2\end{pmatrix} = \begin{pmatrix}-2xv_x & -2yv_y\\ 2xg_x & 2yg_y\end{pmatrix}\begin{pmatrix}\zeta_1\\ \zeta_2\end{pmatrix}, \qquad (\zeta_1, \zeta_2)(\tau) = (0, 1). \tag{D.7}$$

The jet corridor gives

$$|\zeta_1(t)| + |\zeta_2(t)| \leq e^{\delta\Lambda(t;\tau)}. \tag{D.8}$$

Write $s = y/x$ and

$$f(s) := \frac{\sqrt{s}}{(1+s)^{5/2}}, \qquad \kappa_1(s) := -f(s) - sf'(s), \qquad \kappa_2(s) := sf'(s).$$

Since $K_H = x^{-1} f(s)$, the exact response derivative is

$$\mathcal{R}(t, \tau; X_\tau) = -\frac{1}{y_\tau x(t)} \left[\zeta_1(t)\kappa_1(s(t)) + \zeta_2(t)\kappa_2(s(t))\right]. \tag{D.9}$$

For $s \geq 1$, direct differentiation of f gives

$$|\kappa_1| + |\kappa_2| + |s\kappa_1'| + |s\kappa_2'| \leq K_0 s^{-2},$$

whereas the same expression is bounded by K_0 for $0 < s \leq 1$. Differentiating (D.9), using (D.7), the corridor bounds, and (D.8), gives the following two cases.

If $s(t) \geq 1$, then

$$|\partial_t \mathcal{R}(t, \tau; X_\tau)| \leq \frac{K_\delta a(t)}{y_\tau x(t)} e^{\delta\Lambda(t;\tau)} s(t)^{-2}.$$

Use (D.4)–(D.5) and $y_\tau = s_\tau x_\tau$ to obtain

$$|\partial_t \mathcal{R}| \leq \frac{K_\delta a(t)}{x_\tau^2 s_\tau^3} e^{(-10+7\delta)\Lambda(t;\tau)} \leq \frac{K_\delta a(t)}{x_\tau^2 s_\tau^3}. \tag{D.10}$$

If instead $s(t) \leq 1$, then (D.5) implies $s_\tau \leq 1$ and

$$e^{\Lambda(t;\tau)} \leq s_\tau^{-1/(6-2\delta)}.$$

The bounded angular factors now give

$$|\partial_t \mathcal{R}| \leq \frac{K_\delta a(t)}{x_\tau^2 s_\tau} e^{(2+3\delta)\Lambda(t;\tau)} \leq \frac{K_\delta a(t)}{x_\tau^2 s_\tau^{p_\delta}},$$

where

$$p_\delta := 1 + \frac{2+3\delta}{6-2\delta} < 3.$$

Since $0 < s_\tau \le 1$, this is again bounded by $K_\delta a(t)/(x_\tau^2 s_\tau^3)$. We have therefore proved the all-age bound

$$\boxed{|\partial_t \mathcal{R}(t,\tau;X_\tau)| \le \frac{K_\delta a(t)}{x_\tau^2 s_\tau^3}.} \tag{D.11}$$

No raw aspect-ratio symbol r has been introduced; throughout this audited proof the aspect ratio is $s = y/x$.

At zero response age, (15.5) gives

$$\mathcal{R}_0 = \frac{4s-1}{2x^2\sqrt{s}(1+s)^{7/2}}, \qquad \Theta = \Xi_\beta x^2 s^3.$$

Hence

$$|\Theta\mathcal{R}_0| = \Xi_\beta \frac{s^{5/2}|4s-1|}{2(1+s)^{7/2}} \le K_0 \Xi_\beta. \tag{D.12}$$

Step 3: exact exposure decay. Because $\mathcal{D}_{xy}\Theta = 4v\Theta$, $\mathcal{D}_{xy}\log x = -2v$, and $\mathcal{D}_{xy}\log y = 2g$,

$$\mathcal{D}_{xy}\Xi_\beta = (2v-6g)\Xi_\beta. \tag{D.13}$$

The squared-coordinate transport field has $\nabla\cdot\mathbf{b} = g$, by the exact divergence constraint. Therefore

$$\partial_t \Xi_\beta + \nabla\cdot(\mathbf{b}\Xi_\beta) = (2v-5g)\Xi_\beta. \tag{D.14}$$

On the material set $\mathcal{N}(t)$,

$$\frac{d}{dt} M_{3,\mathcal{N}}(t) = \int_{\mathcal{N}(t)} (2v-5g)\Xi_\beta \, dX.$$

The value corridor gives

$$2v - 5g = -8a + 2(v-a) - 5(g-2a) \le -(8-5\delta)a.$$

Since $\mathfrak{t}_\Psi(t) = \exp(-2\int_0^t a)$, integration proves (13.15).

Step 4: response payment and the dichotomy. Use (D.12) for the current term in (13.13). For the historical term, multiply (D.11) by

$$\Theta(X_\tau,\tau) = \Xi_\beta(X_\tau,\tau) x_\tau^2 s_\tau^3$$

and integrate over $\mathcal{N}(\tau)$ and then in τ. This proves (13.14). Since $M_{3,\mathcal{N}}(t) \le M_{3,\mathcal{N}}(0)$ and $T < \infty$,

$$|\mathcal{P}^{\mathrm{src}}_{\beta,\mathcal{N}}(t)| \le K_{\delta,T} M_{3,\mathcal{N}}(0)(1+a(t)).$$

Division by $a(t)^2$ proves (13.16) on the stated terminal branch.

If the corridor does not persist, let t_* be its first failure time. The value part is expressed in the protected variables

$$v - a = V, \qquad g - 2a = 2V - \mathfrak{A},$$

and the four weighted first jets are exactly $C = xv_x$, $D = yg_y$, $E = yv_y$, $F = xg_x$. Hence the first failure is a threshold failure of an already named value/first-jet quantity. The logarithmic tangent system proposition 9.2 and lemma 9.3 and the radial star-hull identity proposition 9.4 provide the existing cumulative restart accounts for such non-affine loss. At this D2 stage we use only the routing statement: the first corridor failure is named, not silently discarded; no new quantitative lower bound for the restart payment is asserted here. □

Audit status: GREEN/COMPLETE.

D3. Raw source-gradient Gronwall estimate. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 14.2.

> **Restatement of proposition 14.2.** The tagged source-gradient pair obeys the raw L^∞ estimate (14.4). Its carrier-L^1 and weighted L^2 stocks obey the exact Dini inequalities (14.10)–(14.11), hence the corresponding Gronwall bounds. No normalized apex ratio is used.

Proof. By proposition 16.1 and the identities

$$-yH - 2s\mathfrak{T} = 4yv_y, \qquad \frac{1}{4}xH - \frac{1}{2}\mathfrak{T} = -xg_x,$$

the homogeneous material system can be written in the raw form

$$\boxed{\begin{aligned} \mathcal{D}_{xy}q &= \mathsf{d}_q q + (4yv_y)C_\beta, \\ \mathcal{D}_{xy}C_\beta &= -(xg_x)q + \left(\frac{3}{2}\mathfrak{A} + \mathfrak{J}\right)C_\beta. \end{aligned}} \tag{D.15}$$

Along a material trajectory, the upper Dini derivative of either the Euclidean or componentwise ℓ^1 norm is bounded by $L_{\mathrm{sg}}(t)$ times that norm. Taking the spatial supremum and applying scalar Gronwall gives (14.4).

It remains to audit the carrier weights rather than treating them as frozen. Put

$$\rho_c := \mathfrak{r}^{-1} = x^{-2/3}y^{-1/3}, \qquad d\mu_c = \rho_c\, dX.$$

The protected material equations for x, y give

$$\mathcal{D}_{xy}\log\rho_c = \frac{4v - 2g}{3}.$$

Moreover, using $g + 2yg_y = 2v + 2xv_x$,

$$\nabla\cdot\mathbf{b} = \partial_x(-2xv) + \partial_y(2yg) = g.$$

Therefore

$$\boxed{\partial_t\rho_c + \nabla\cdot(\mathbf{b}\rho_c) = \delta_c\rho_c, \qquad \delta_c = \frac{4v+g}{3}.} \tag{D.16}$$

This is the carrier-measure defect that must be retained in every weighted estimate.

For $\mu, \nu \geq 0$ define

$$Z_\beta^{\mu,\nu} := x^\mu y^\nu \begin{pmatrix} q \\ C_\beta \end{pmatrix}.$$

On the interior $x, y > 0$,

$$\mathcal{D}_{xy}(x^\mu y^\nu) = (-2\mu v + 2\nu g)x^\mu y^\nu.$$

The coordinate axes are invariant for the transport field and its normal components carry the factors x and y. Hence this identity is valid almost everywhere for the weighted integrals; equivalently one may truncate away from the axes and pass to the limit, with the axis boundary flux vanishing by regular-axis behavior. We obtain the exact homogeneous weighted system

$$\boxed{\mathcal{D}_{xy}Z_\beta^{\mu,\nu} = [\mathbf{A}_{\mathrm{sg}} + (-2\mu v + 2\nu g)I_2]\, Z_\beta^{\mu,\nu}.} \tag{D.17}$$

For the componentwise ℓ^1 norm, combine (D.17) with (D.16); after integration by parts (justified by the standing fast decay and regular axis behavior),

$$D_t^+\mathcal{Q}_{\mathrm{sg}}^{(1;\mu,\nu)} \leq (L_{\mathrm{sg}} + 2\mu\|v\|_\infty + 2\nu\|g\|_\infty + \|\delta_c\|_\infty)\,\mathcal{Q}_{\mathrm{sg}}^{(1;\mu,\nu)}.$$

This is (14.10).

For the squared Euclidean norm, (D.17) and (D.16) give

$$\frac{d}{dt}\big(\mathcal{E}_{\mathrm{sg}}^{(\mu,\nu)}\big)^2 \le (2L_{\mathrm{sg}} + 4\mu\|v\|_\infty + 4\nu\|g\|_\infty + \|\delta_c\|_\infty)\,\big(\mathcal{E}_{\mathrm{sg}}^{(\mu,\nu)}\big)^2.$$

Taking the upper Dini derivative of the square root proves (14.11). Scalar Gronwall now yields

$$\mathcal{Q}_{\mathrm{sg}}^{(1;\mu,\nu)}(t) \le \mathcal{Q}_{\mathrm{sg}}^{(1;\mu,\nu)}(t_0)\exp\left(\int_{t_0}^{t} L_{\mathrm{sg}}^{(1;\mu,\nu)}(\tau)\,d\tau\right), \tag{D.18}$$

$$\mathcal{E}_{\mathrm{sg}}^{(\mu,\nu)}(t) \le \mathcal{E}_{\mathrm{sg}}^{(\mu,\nu)}(t_0)\exp\left(\int_{t_0}^{t} L_{\mathrm{sg}}^{(2;\mu,\nu)}(\tau)\,d\tau\right). \tag{D.19}$$

For $(\mu,\nu) = (0,0)$ the first estimate is precisely the carrier-L^1 stock used in the genealogy bank. For $(\mu,\nu) = (1/2,1/2)$ the second is the weighted source-gradient energy that forces $\mathcal{E}_{H,1}$ in D4. Thus the raw and weighted source-gradient stocks are all paid by the same homogeneous restart system, with the exact carrier defect explicitly included. □

Audit status: GREEN/COMPLETE.

D4. Weighted Hodge energy is not a new mechanism. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 14.4.

> **Restatement of proposition 14.4.** Let $d\mu_c = dX/\mathfrak{r}$ be the carrier measure from definition 14.1, and define
>
> $$\mathcal{E}_{H,1} := \|\sqrt{xy}\,H\|_{L^2(d\mu_c)}, \qquad \mathcal{E}_{\mathrm{sg}}^{(1/2,1/2)} := \|\sqrt{xy}(q, C_\beta)\|_{L^2(d\mu_c;\ell^2)}.$$
>
> Put
>
> $$M_H := \sqrt{xy}\,H.$$
>
> Then
>
> $$\boxed{\mathcal{D}_{xy}M_H = -vM_H + 2\sqrt{xy}\,q.}$$
>
> Let $\mathbf{B}(\mathbf{X},\mathbf{Z},t) = (-v\mathbf{X}, g\mathbf{Z})$ be the seven-dimensional transport lift and define
>
> $$\mathcal{L}_B(t) := \|\nabla_{\mathbb{R}^7}\mathbf{B}(t)\|_{L^\infty}.$$
>
> With D_t^+ denoting the upper right Dini derivative,
>
> $$\boxed{D_t^+\mathcal{E}_{H,1} \le \mathfrak{c}\mathcal{L}_B\mathcal{E}_{H,1} + 2\mathcal{E}_{\mathrm{sg}}^{(1/2,1/2)}.}$$
>
> Thus the weighted Hodge energy is financed by the same first-jet/source-gradient restart package.

Proof. Let

$$\rho_c := \mathfrak{r}^{-1}, \qquad \mathbf{b} := (-2xv, 2yg), \qquad M_H := \sqrt{xy}\,H.$$

The exact D3 carrier identity (D.16) is

$$\partial_t\rho_c + \nabla\cdot(\mathbf{b}\rho_c) = \delta_c\rho_c, \qquad \delta_c = \frac{4v+g}{3}. \tag{D.20}$$

The protected material equations give

$$\mathcal{D}_{xy}\sqrt{xy} = (g-v)\sqrt{xy}, \qquad \mathcal{D}_{xy}H = -gH + 2q,$$

so the product rule gives, without any normalization by a,

$$\boxed{\mathcal{D}_{xy}M_H = -vM_H + 2\sqrt{xy}\,q.} \tag{D.21}$$

We now keep the carrier defect rather than treating $d\mu_c$ as stationary. By integrating first on a compact rectangle in the interior $x,y>0$, using (D.20), and then sending the cutoffs to

the axes and infinity, the axis fluxes vanish because the transport field has factors x and y, while the far-field terms vanish by the standing rapid-decay hypothesis. Hence

$$\begin{aligned}\frac{1}{2}\frac{d}{dt}\mathcal{E}_{H,1}^2 &= \int_{\mathcal{Q}} M_H \mathcal{D}_{xy} M_H \, d\mu_c + \frac{1}{2}\int_{\mathcal{Q}} \delta_c M_H^2 \, d\mu_c \\ &= \int_{\mathcal{Q}} \left(-v + \frac{\delta_c}{2}\right) M_H^2 \, d\mu_c + 2\int_{\mathcal{Q}} M_H \sqrt{xy}\, q \, d\mu_c. \end{aligned} \tag{D.22}$$

Cauchy–Schwarz and the definition of the weighted source-gradient stock give

$$\frac{1}{2} D_t^+ \mathcal{E}_{H,1}^2 \le \left(\|v\|_\infty + \frac{1}{2}\|\delta_c\|_\infty\right) \mathcal{E}_{H,1}^2 + 2\mathcal{E}_{H,1}\mathcal{E}_{\mathrm{sg}}^{(1/2,1/2)}. \tag{D.23}$$

It remains only to compare the coefficient with the already named first-jet action. In the lift $\mathbf{B} = (-v\mathbf{X}, g\mathbf{Z})$, tangential derivatives in the $\mathbf{X}$ and $\mathbf{Z}$ blocks recover respectively $-v$ and g (and the same conclusion holds at the block origins by direct differentiation). Thus

$$\|v\|_\infty + \|g\|_\infty \le \mathfrak{c}\mathcal{L}_B, \qquad \|\delta_c\|_\infty \le \mathfrak{c}\mathcal{L}_B. \tag{D.24}$$

If $\mathcal{E}_{H,1} > 0$, divide (D.23) by $\mathcal{E}_{H,1}$. At a zero of the norm, apply the same calculation to $(\mathcal{E}_{H,1}^2 + \epsilon_{\mathrm{reg}}^2)^{1/2}$ and let $\epsilon_{\mathrm{reg}} \downarrow 0$. This proves the upper-Dini estimate

$$\boxed{D_t^+ \mathcal{E}_{H,1} \le \mathfrak{c}\mathcal{L}_B \mathcal{E}_{H,1} + 2\mathcal{E}_{\mathrm{sg}}^{(1/2,1/2)}.} \tag{D.25}$$

In particular, scalar Gronwall yields

$$\mathcal{E}_{H,1}(t) \le \exp\left(\mathfrak{c}\int_{t_0}^t \mathcal{L}_B(\tau)\, d\tau\right)\left[\mathcal{E}_{H,1}(t_0) + 2\int_{t_0}^t \mathcal{E}_{\mathrm{sg}}^{(1/2,1/2)}(\tau)\, d\tau\right]. \tag{D.26}$$

D3 already propagates the forcing stock on the right. Therefore the weighted Hodge energy is a forced GREEN account paid by the existing first-jet/source-gradient restart package, not an independent continuation mechanism. □

Audit status: GREEN/COMPLETE.

D5. Physical Biot–Savart control. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 14.5.

> **Restatement of proposition 14.5.** Let $\mathbf{U}_{\mathrm{mer}} := U^r \mathbf{e}_r + U^z \mathbf{e}_z$ be the physical meridional velocity, let
>
> $$d\mu_c := \frac{dX}{\mathfrak{r}}$$
>
> be the carrier measure, and define
>
> $$\mathcal{E}_{H,1} := \|\sqrt{xy}\, H\|_{L^2(d\mu_c)}.$$
>
> For $\boldsymbol{\xi} = (r\cos\phi, r\sin\phi, z)$ set $\omega(\boldsymbol{\xi}) = r^{-4/3}|z|^{1/3}$ and $\|\varphi\|_{L^2_\omega}^2 = \int_{\mathbb{R}^3} |\varphi|^2 \omega \, d\boldsymbol{\xi}$. Then
>
> $$\boxed{\|\nabla \mathbf{U}_{\mathrm{mer}}\|_{L^2_\omega} \le \mathfrak{c}_\omega \sqrt{\pi}\, \mathcal{E}_{H,1}.}$$
>
> Hence the quadratic meridional pressure source is controlled by a GREEN Hodge carrier energy and ordinary first-jet routing norms.

Proof. Write the physical point as

$$\boldsymbol{\xi} = (r\cos\phi, r\sin\phi, z) \in \mathbb{R}^3, \qquad \omega(\boldsymbol{\xi}) = r^{-4/3}|z|^{1/3}.$$

The factor $r^{-4/3}$ is an A_2 power weight on the two-dimensional transverse block because $-2 < -4/3 < 2$, while $|z|^{1/3}$ is an A_2 power weight on the axial line because $-1 < 1/3 < 1$. Since the weight factorizes and axis-parallel cubes are products of a transverse square and an axial interval, the A_2 averages factorize; therefore

$$\omega \in A_2(\mathbb{R}^3). \tag{D.27}$$

The meridional field is divergence free and rapidly decaying, and its curl is purely azimuthal:

$$\nabla \times \mathbf{U}_{\mathrm{mer}} = \omega^\phi \mathbf{e}_\phi, \qquad \omega^\phi = \partial_z U^r - \partial_r U^z.$$

Hence whole-space Hodge reconstruction gives $\mathbf{U}_{\mathrm{mer}}$ from $\omega^\phi \mathbf{e}_\phi$ with no harmonic remainder: any remainder would be a decaying vector field with zero divergence and zero curl and is therefore zero. Each entry of $\nabla \mathbf{U}_{\mathrm{mer}}$ is consequently a Calderón–Zygmund operator applied to $\omega^\phi \mathbf{e}_\phi$. The weighted A_2 theorem and (D.27) yield

$$\|\nabla \mathbf{U}_{\mathrm{mer}}\|_{L^2_\omega} \le \mathfrak{c}_\omega \|\omega^\phi\|_{L^2_\omega}. \tag{D.28}$$

It remains to translate the vorticity norm exactly into the carrier account. The squared-coordinate dictionary (5.7) gives

$$\omega^\phi = rzH(r^2, z^2, t).$$

Using axisymmetry, $x = r^2$, $y = z^2$, and $d\boldsymbol{\xi} = r\,dr\,d\phi\,dz$, we obtain

$$\begin{aligned} \|\omega^\phi\|^2_{L^2_\omega} &= 2\pi \int_0^\infty \int_{-\infty}^\infty r^2 z^2 |H(r^2, z^2)|^2 r^{-4/3} |z|^{1/3} r\,dz\,dr \\ &= \pi \int_0^\infty \int_0^\infty x^{1/3} y^{2/3} |H(x,y)|^2\,dx\,dy \\ &= \pi \int_{\mathcal{Q}} xy|H|^2 \frac{dX}{x^{2/3}y^{1/3}} = \pi \mathcal{E}^2_{H,1}. \end{aligned} \tag{D.29}$$

Taking square roots in (D.29) and substituting into (D.28) proves

$$\boxed{\|\nabla \mathbf{U}_{\mathrm{mer}}\|_{L^2_\omega} \le \mathfrak{c}_\omega \sqrt{\pi}\, \mathcal{E}_{H,1}.}$$

Thus D5 introduces no new source currency: the physical weighted meridional gradient is paid exactly by the D4 GREEN Hodge carrier energy. □

Audit status: GREEN/COMPLETE.

D6. Finite physical Lipschitz action is a continuation firewall. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 14.6.

> **Restatement of theorem 14.6.** Let $[t_0, T)$ be a finite classical interval for the fixed rapidly decaying Euler customer and assume
>
> $$\mathcal{L}_{\mathrm{phys}}([t_0, T)) := \int_{t_0}^T \|\nabla \mathbf{U}(t)\|_\infty\,dt < \infty.$$
>
> Then every fixed H^m, $m \ge 3$, stays bounded; particle displacement stays finite; the vorticity and its first two derivatives retain uniform rapid decay; and $|\boldsymbol{\xi}|\,|D^2\mathbf{U}| \to 0$

uniformly in time at spatial infinity. Consequently the extreme-angle/far-field weighted restart gates cannot fail invisibly. Moreover, for every subinterval $I \subset [t_0, T)$,

$$\boxed{\left| \int_I a(t)\, dt \right| \le \int_I \|\nabla \mathbf{U}(t)\|_\infty \, dt.}$$

Thus infinite collapse age on a finite interval already forces failure of the standard Euler Lipschitz continuation action.

Proof. Put

$$L_{\mathrm{phys}} := \int_{t_0}^{T} \|\nabla \mathbf{U}(t)\|_\infty \, dt < \infty.$$

For every fixed integer $m \ge 3$, the classical Euler commutator estimate gives

$$\frac{d}{dt}\|\mathbf{U}(t)\|_{H^m} \le c_m \|\nabla \mathbf{U}(t)\|_\infty \|\mathbf{U}(t)\|_{H^m}. \tag{D.30}$$

Gronwall therefore yields

$$\boxed{\sup_{t_0 \le t < T} \|\mathbf{U}(t)\|_{H^m} \le \|\mathbf{U}(t_0)\|_{H^m} e^{c_m L_{\mathrm{phys}}} < \infty.} \tag{D.31}$$

This is the terminal-uniform estimate; no pointwise-in-time supremum of $\|\nabla \mathbf{U}\|_\infty$ has been assumed. Taking, for example, $m \ge 7$ and using Sobolev embedding gives uniform bounds for $\mathbf{U}$ and for all ordinary spatial derivatives needed below.

Let $\mathscr{X}^{\mathrm{phys}}_{t,t_0}$ be the physical Euler flow. From (D.31),

$$M_{\mathrm{vel}} := \sup_{t_0 \le t < T} \|\mathbf{U}(t)\|_\infty < \infty, \tag{D.32}$$

so every label $\boldsymbol{\xi}_0$ satisfies

$$\boxed{|\mathscr{X}^{\mathrm{phys}}_{t,t_0}(\boldsymbol{\xi}_0) - \boldsymbol{\xi}_0| \le (T - t_0) M_{\mathrm{vel}} =: \mathfrak{d}_{\mathrm{disp}} < \infty.} \tag{D.33}$$

The deformation gradient obeys the usual variational equation, hence

$$\|D\mathscr{X}^{\mathrm{phys}}_{t,t_0}\|_\infty + \|D(\mathscr{X}^{\mathrm{phys}}_{t,t_0})^{-1}\|_\infty \le 2e^{L_{\mathrm{phys}}}. \tag{D.34}$$

For derivatives of order two and three, differentiate the variational equation. The resulting systems are linear in the highest flow derivative with coefficients $\nabla \mathbf{U}$, and their forcing terms contain only lower flow derivatives and $D^2\mathbf{U}, D^3\mathbf{U}$, which are uniformly bounded by (D.31). Gronwall on the finite interval gives

$$\sup_{t_0 \le t < T} \left(\|D^j \mathscr{X}^{\mathrm{phys}}_{t,t_0}\|_\infty + \|D^j(\mathscr{X}^{\mathrm{phys}}_{t,t_0})^{-1}\|_\infty \right) < \infty, \qquad j = 2, 3. \tag{D.35}$$

Now let $\boldsymbol{\omega}_{\mathrm{phys}} := \nabla \times \mathbf{U}$. Incompressibility gives the exact Cauchy formula

$$\boxed{\boldsymbol{\omega}_{\mathrm{phys}}(\mathscr{X}^{\mathrm{phys}}_{t,t_0}(\boldsymbol{\xi}_0), t) = D\mathscr{X}^{\mathrm{phys}}_{t,t_0}(\boldsymbol{\xi}_0)\boldsymbol{\omega}_{\mathrm{phys}}(\boldsymbol{\xi}_0, t_0).} \tag{D.36}$$

The standing fixed customer is in the smooth rapid-decay class. Writing $\boldsymbol{\xi}_0 = (\mathscr{X}^{\mathrm{phys}}_{t,t_0})^{-1}(\boldsymbol{\xi})$, differentiating (D.36) at most twice, and using (D.34)–(D.35) shows that each $D^j\boldsymbol{\omega}_{\mathrm{phys}}(\boldsymbol{\xi}, t)$, $j \le 2$, is a finite sum of bounded flow factors times an initial derivative of order at most j. Moreover (D.33) gives

$$1 + |\boldsymbol{\xi}| \le (1 + \mathfrak{d}_{\mathrm{disp}})(1 + |\boldsymbol{\xi}_0|), \qquad 1 + |\boldsymbol{\xi}_0| \le (1 + \mathfrak{d}_{\mathrm{disp}})(1 + |\boldsymbol{\xi}|).$$

Therefore, for every $N > 0$ and $j = 0, 1, 2$,

$$\boxed{\sup_{t_0 \le t < T} \sup_{\boldsymbol{\xi} \in \mathbb{R}^3} (1 + |\boldsymbol{\xi}|)^N |D^j \boldsymbol{\omega}_{\mathrm{phys}}(\boldsymbol{\xi}, t)| < \infty.} \tag{D.37}$$

In particular,

$$\sup_{t_0 \le t < T} \|D^2 \boldsymbol{\omega}_{\mathrm{phys}}(t)\|_{L^1} < \infty, \tag{D.38}$$

and its L^1 tail decreases faster than any fixed inverse power, uniformly in time.

We next make the far-field restart firewall quantitative. The whole-space Biot–Savart kernel $\mathcal{K}$ satisfies $|\mathcal{K}(\boldsymbol{\eta})| \le \mathfrak{c}|\boldsymbol{\eta}|^{-2}$. Rapid decay permits two integrations by parts, so

$$D^2 \mathbf{U} = \mathcal{K} * D^2 \boldsymbol{\omega}_{\mathrm{phys}}. \tag{D.39}$$

The local singularity $|\boldsymbol{\eta}|^{-2}$ is integrable in three dimensions. Fix $\boldsymbol{\xi}$ with $\rho := |\boldsymbol{\xi}|$ large. On $|\boldsymbol{\eta}| \le \rho/2$,

$$\rho \int_{|\boldsymbol{\eta}| \le \rho/2} \frac{|D^2 \boldsymbol{\omega}_{\mathrm{phys}}(\boldsymbol{\eta}, t)|}{|\boldsymbol{\xi} - \boldsymbol{\eta}|^2} \, d\boldsymbol{\eta} \le \frac{\mathfrak{c}}{\rho} \|D^2 \boldsymbol{\omega}_{\mathrm{phys}}(t)\|_{L^1}. \tag{D.40}$$

On $|\boldsymbol{\eta}| > \rho/2$ and $|\boldsymbol{\xi} - \boldsymbol{\eta}| < 1$, (D.37) gives, for every fixed N,

$$\rho \int_{|\boldsymbol{\xi} - \boldsymbol{\eta}| < 1} \frac{|D^2 \boldsymbol{\omega}_{\mathrm{phys}}(\boldsymbol{\eta}, t)|}{|\boldsymbol{\xi} - \boldsymbol{\eta}|^2} \, d\boldsymbol{\eta} \le c_N \rho^{1-N}. \tag{D.41}$$

On the remaining outer region the kernel is bounded, while the uniform rapid tail gives

$$\rho \int_{|\boldsymbol{\eta}| > \rho/2} |D^2 \boldsymbol{\omega}_{\mathrm{phys}}(\boldsymbol{\eta}, t)| \, d\boldsymbol{\eta} \le c_N \rho^{1-N}. \tag{D.42}$$

Choose $N > 2$ and combine (D.38)–(D.42). We obtain

$$\boxed{\lim_{\varrho_0 \to \infty} \sup_{t_0 \le t < T} \sup_{|\boldsymbol{\xi}| \ge \varrho_0} |\boldsymbol{\xi}| \, |D^2 \mathbf{U}(\boldsymbol{\xi}, t)| = 0.} \tag{D.43}$$

This closes the extreme-angle gate as well. Axis parity gives $U_z^r(0, z, t) = 0$ and $U_r^z(r, 0, t) = 0$. The one-dimensional fundamental theorem of calculus therefore yields, away from the axes,

$$\left| \frac{z}{r} U_z^r(r, z, t) \right| \le |z| \sup_{0 \le \theta \le 1} |U_{rz}^r(\theta r, z, t)|, \tag{D.44}$$

$$\left| \frac{r}{z} U_r^z(r, z, t) \right| \le |r| \sup_{0 \le \theta \le 1} |U_{rz}^z(r, \theta z, t)|. \tag{D.45}$$

Let

$$M_2 := \sup_{t_0 \le t < T} \|D^2 \mathbf{U}(t)\|_\infty < \infty.$$

Choose one fixed $\varrho_* > 0$ beyond which the supremum in (D.43) is finite. In the first line of (D.44), if $|z| \le \varrho_*$ then the right side is at most $\varrho_* M_2$; if $|z| > \varrho_*$, every point $(\theta r, z)$ on the FTC segment has physical radius at least $|z|$, so (D.43) bounds $|z| \, |U_{rz}^r|$ uniformly. Similarly, in (D.45), the case $r \le \varrho_*$ is bounded by $\varrho_* M_2$, while for $r > \varrho_*$ every point $(r, \theta z)$ on the FTC segment has radius at least r and the far-field estimate applies. Thus both aspect-weighted cross derivatives are globally bounded. A failure of the aspect-weighted or far-field restart gate therefore cannot occur while $L_{\mathrm{phys}} < \infty$ without already triggering an ordinary physical continuation failure.

Finally the protected physical dictionary gives

$$U^r(r, z, t) = -r\alpha(r, z, t), \qquad a(t) = \alpha(0, 0, t),$$

so axis regularity implies

$$a(t) = -U_r^r(0,0,t), \qquad |a(t)| \le \|\nabla \mathbf{U}(t)\|_\infty. \tag{D.46}$$

Integrating (D.46) over any subinterval $I \subset [t_0, T)$ proves

$$\boxed{\left|\int_I a(t)\,dt\right| \le \int_I \|\nabla \mathbf{U}(t)\|_\infty\,dt.}$$

Hence a sequence of finite-time intervals with collapse age tending to $+\infty$ forces divergence of the physical Lipschitz action. This is a continuation firewall, not an assumption that infinite collapse age actually occurs. □

Audit status: GREEN/COMPLETE.

Appendix E. Two-time response and the fixed-seed affine benchmark

This section contains the complete response and fixed-seed affine-benchmark proofs E1–E4. The later nonlinear polarization/conversion slots remain restated and reserved without changing their permanent ledger IDs.

E1. Vertical response representation. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 15.2.

> **Restatement of proposition 15.2.** For the fixed customer of definition 5.1, so in particular $H(\cdot,0) = 0$, let $\mathscr{X}_{t,\tau}$ be the flow from definition 15.1 and define
>
> $$\boxed{\mathcal{R}(t,\tau;X) := -\partial_{y_\tau} K_H(\mathscr{X}_{t,\tau}X),}$$
>
> where ∂_{y_τ} denotes differentiation with respect to the y-coordinate of the deposition point $X = (x_\tau, y_\tau)$ at time τ. Then
>
> $$\boxed{a(t) = \frac{3}{4}\int_0^t \int_{\mathcal{Q}} \Theta(X,\tau)\mathcal{R}(t,\tau;X)\,dX\,d\tau.}$$
>
> At zero response age,
>
> $$\boxed{\mathcal{R}_0(x,y) := \mathcal{R}(t,t;(x,y)) = \frac{x(4y-x)}{2\sqrt{y}(x+y)^{7/2}}, \qquad \mathcal{R}_0 > 0 \iff y > x/4.}$$

Proof. Set

$$\mathbf{b}(x,y,t) := (-2xv(x,y,t),\ 2yg(x,y,t)).$$

By (4.15),

$$\begin{aligned}\nabla\cdot\mathbf{b} &= -2v - 2xv_x + 2g + 2yg_y \\ &= -2v - 2xv_x + 2g + \big(2v + 2xv_x - g\big) = g.\end{aligned}$$

Consequently the vorticity-source equation in (5.8) is exactly

$$\partial_t H + \nabla\cdot(\mathbf{b}H) = 2q. \tag{E.1}$$

For the fixed customer, $H(\cdot,0) = 0$.

Let

$$J(t,\tau;X) := \det D_X \mathscr{X}_{t,\tau}(X) > 0.$$

Liouville's formula and $\nabla\cdot\mathbf{b} = g$ give

$$\partial_t J(t,\tau;X) = g(\mathscr{X}_{t,\tau}X, t)J(t,\tau;X), \qquad J(\tau,\tau;X) = 1. \tag{E.2}$$

Equivalently, if X_0 is a time-zero label, then along $X(t) = \mathscr{X}_{t,0}X_0$,

$$\frac{d}{dt}\big[H(X(t),t)J(t,0;X_0)\big] = 2q(X(t),t)J(t,0;X_0).$$

Since $H(\cdot,0) = 0$,

$$H(\mathscr{X}_{t,0}X_0,t)J(t,0;X_0) = 2\int_0^t q(\mathscr{X}_{\tau,0}X_0,\tau)J(\tau,0;X_0)\,d\tau. \tag{E.3}$$

Multiply by $K_H(\mathscr{X}_{t,0}X_0)$, integrate in X_0, use Fubini, and in the τ-integral change variables $X = \mathscr{X}_{\tau,0}X_0$. The flow composition law $\mathscr{X}_{t,0} = \mathscr{X}_{t,\tau}\circ\mathscr{X}_{\tau,0}$ yields

$$\int_{\mathcal{Q}} K_H(Y)H(Y,t)\,dY = 2\int_0^t\int_{\mathcal{Q}} q(X,\tau)K_H(\mathscr{X}_{t,\tau}X)\,dX\,d\tau. \tag{E.4}$$

All changes of variables are legitimate on a finite classical interval because the fixed solution is smooth and rapidly decaying; one may first perform the calculation on compact truncated quadrants and then pass to the limit.

Insert (E.4) into the Hodge cashier (5.12). This gives

$$a(t) = \frac{3}{4}\int_0^t\int_{\mathcal{Q}} q(X,\tau)K_H(\mathscr{X}_{t,\tau}X)\,dX\,d\tau. \tag{E.5}$$

For each fixed t and τ abbreviate

$$W_{t,\tau}(X) := K_H(\mathscr{X}_{t,\tau}X).$$

By definition 5.2, $q = \Theta_y$. The material equator $y = 0$ is invariant and smooth axisymmetric regularity gives $u(x,0,\tau) = 0$; indeed the factorization used in the audited C4 proof is $u(x,y,\tau) = y\widehat{w}(x,y,\tau)$. Hence $\Theta = \varepsilon_\beta u^2$ vanishes quadratically at $y = 0$. Rapid decay removes the boundary at $y = \infty$. Therefore, integrating first on $\varepsilon < y < M$ and then sending $\varepsilon \downarrow 0$, $M \uparrow \infty$,

$$\begin{aligned}\int_{\mathcal{Q}} q(X,\tau)W_{t,\tau}(X)\,dX &= \int_0^\infty\int_0^\infty \partial_y\Theta(x,y,\tau)W_{t,\tau}(x,y)\,dy\,dx\\ &= -\int_{\mathcal{Q}} \Theta(X,\tau)\partial_{y_\tau}K_H(\mathscr{X}_{t,\tau}X)\,dX.\end{aligned}$$

(The notation ∂_{y_τ} emphasizes that the derivative is with respect to the deposition coordinate while t is fixed.) With

$$\mathcal{R}(t,\tau;X) := -\partial_{y_\tau}K_H(\mathscr{X}_{t,\tau}X),$$

substitution into (E.5) proves

$$\boxed{a(t) = \frac{3}{4}\int_0^t\int_{\mathcal{Q}} \Theta(X,\tau)\mathcal{R}(t,\tau;X)\,dX\,d\tau.}$$

At zero response age, $\mathscr{X}_{t,t} = \mathrm{Id}$, so

$$\mathcal{R}_0(x,y) = -\partial_y K_H(x,y).$$

Since

$$K_H(x,y) = x\sqrt{y}\,(x+y)^{-5/2},$$

direct differentiation gives

$$-\partial_y K_H(x,y) = \frac{x(4y-x)}{2\sqrt{y}\,(x+y)^{7/2}}.$$

The denominator is positive on $\mathcal{Q}$, so $\mathcal{R}_0 > 0$ exactly when $4y - x > 0$, i.e. $y > x/4$. □

Audit status: GREEN/COMPLETE.

E2. True-Euler aged-response discount. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 15.3.

Restatement of theorem 15.3. Fix a source time τ and a source point $X_\tau = (x_\tau, y_\tau) \in \mathcal{Q}$. Let

$$Z(\xi) = (x(\xi), y(\xi)) := \mathscr{X}_{\xi,\tau} X_\tau, \qquad s_0 := \frac{y_\tau}{x_\tau}, \qquad s(\xi) := \frac{y(\xi)}{x(\xi)},$$

and

$$\Lambda(t;\tau) := \int_\tau^t a(\xi)\, d\xi, \qquad \frac{\mathfrak{t}_\Psi(t)}{\mathfrak{t}_\Psi(\tau)} = e^{-2\Lambda(t;\tau)}.$$

Assume $a(\xi) > 0$ and that for some $0 < \delta < 1$ the trajectory segment $\tau \le \xi \le t$ satisfies the near-affine corridor

$$|v - a| + |g - 2a| \le \delta a,$$
$$2\big(|xv_x| + |yv_y| + |xg_x| + |yg_y|\big) \le \delta a$$

pointwise along $Z(\xi)$, and assume the transported aspect ratio $s(\xi) \ge 1$ on the aged interval. Then

$$\boxed{|\mathcal{R}(t,\tau;X_\tau)| + a(t)^{-1}|\partial_t \mathcal{R}(t,\tau;X_\tau)| \le \frac{\mathfrak{c}}{x_\tau^2 s_0^3}\left(\frac{\mathfrak{t}_\Psi(t)}{\mathfrak{t}_\Psi(\tau)}\right)^{5-\frac{7}{2}\delta}.}$$

Here $\mathfrak{c} > 0$ is a universal comparison constant. Since $0 < \delta < 1$, the exponent is strictly larger than $3/2$, hence in particular strictly larger than one.

Proof. Write

$$Z(\xi) = (x(\xi), y(\xi)) := \mathscr{X}_{\xi,\tau} X_\tau, \qquad s(\xi) := \frac{y(\xi)}{x(\xi)}, \qquad s_0 = s(\tau) = \frac{y_\tau}{x_\tau}.$$

Because $a > 0$, the collapse age $\Lambda(\xi;\tau)$ is nondecreasing. From $\dot{x}/x = -2v$ and the value part of the corridor,

$$\log \frac{x(\xi)}{x_\tau} = -2\Lambda(\xi;\tau) - 2\int_\tau^\xi (v - a)\, d\eta,$$

so

$$x_\tau e^{-(2+2\delta)\Lambda(\xi;\tau)} \le x(\xi) \le x_\tau e^{-(2-2\delta)\Lambda(\xi;\tau)}. \tag{E.6}$$

Similarly,

$$\frac{\dot{s}}{s} = \frac{\dot{y}}{y} - \frac{\dot{x}}{x} = 2(g + v),$$

and $|(g - 2a) + (v - a)| \le \delta a$. Hence

$$s_0 e^{(6-2\delta)\Lambda(\xi;\tau)} \le s(\xi) \le s_0 e^{(6+2\delta)\Lambda(\xi;\tau)}. \tag{E.7}$$

We next differentiate with respect to the deposition coordinate y_τ. Let

$$m(\xi) := D_X \mathscr{X}_{\xi,\tau}(X_\tau) e_y = (m_1(\xi), m_2(\xi))$$

and introduce the namespace-safe logarithmic sensitivities

$$\zeta_1(\xi) := y_\tau \frac{m_1(\xi)}{x(\xi)}, \qquad \zeta_2(\xi) := y_\tau \frac{m_2(\xi)}{y(\xi)}. \tag{E.8}$$

They satisfy $\zeta_1(\tau) = 0$, $\zeta_2(\tau) = 1$. Linearizing $\dot{x} = -2xv$, $\dot{y} = 2yg$ and dividing by the corresponding base trajectory gives

$$\frac{d}{d\xi}\begin{pmatrix}\zeta_1\\ \zeta_2\end{pmatrix} = \begin{pmatrix}-2xv_x & -2yv_y\\ 2xg_x & 2yg_y\end{pmatrix}_{Z(\xi),\xi}\begin{pmatrix}\zeta_1\\ \zeta_2\end{pmatrix}. \tag{E.9}$$

For $\mathcal{N}_\zeta := |\zeta_1| + |\zeta_2|$, the upper right Dini derivative obeys

$$\begin{aligned} D^+_\xi \mathcal{N}_\zeta &\le 2(|xv_x| + |xg_x|)|\zeta_1| + 2(|yv_y| + |yg_y|)|\zeta_2| \\ &\le \delta a(\xi)\mathcal{N}_\zeta, \end{aligned}$$

where the last step uses the derivative corridor. Gronwall and $\mathcal{N}_\zeta(\tau) = 1$ give

$$\boxed{\mathcal{N}_\zeta(\xi) \le e^{\delta\Lambda(\xi;\tau)}.} \tag{E.10}$$

Now write the Hodge kernel in aspect-ratio form

$$K_H(x, y) = x^{-1} f(s), \qquad f(s) := \frac{\sqrt{s}}{(1+s)^{5/2}}.$$

Define the statement-local angular functions

$$A_*(s) := -f(s) - sf'(s), \qquad B_*(s) := sf'(s). \tag{E.11}$$

A direct differentiation gives

$$A_*(s) = \frac{\sqrt{s}\,(s - \frac{3}{2})}{(1+s)^{7/2}}, \qquad B_*(s) = \frac{\sqrt{s}\,(1-4s)}{2(1+s)^{7/2}}, \tag{E.12}$$

$$sA'_*(s) = -\frac{\sqrt{s}\,(8s^2 - 24s + 3)}{4(1+s)^{9/2}}, \qquad sB'_*(s) = \frac{\sqrt{s}\,(16s^2 - 18s + 1)}{4(1+s)^{9/2}}. \tag{E.13}$$

Therefore there is a universal c_K such that, for every $s \ge 1$,

$$|A_*(s)| + |B_*(s)| + |sA'_*(s)| + |sB'_*(s)| \le c_K s^{-2}. \tag{E.14}$$

By the chain rule and (E.8),

$$\begin{aligned} \mathcal{R}(t, \tau; X_\tau) &= -m_1(t)\,\partial_x K_H(Z(t)) - m_2(t)\,\partial_y K_H(Z(t)) \\ &= -\frac{1}{y_\tau x(t)}\big[\zeta_1(t)A_*(s(t)) + \zeta_2(t)B_*(s(t))\big]. \end{aligned}$$

Thus (E.10) and (E.14) imply

$$|\mathcal{R}(t, \tau; X_\tau)| \le \frac{\mathfrak{c}}{y_\tau x(t)} e^{\delta\Lambda(t;\tau)} s(t)^{-2}. \tag{E.15}$$

Since $y_\tau = s_0 x_\tau$, the lower bounds (E.6)–(E.7) give

$$\begin{aligned} |\mathcal{R}(t, \tau; X_\tau)| &\le \frac{\mathfrak{c}}{x_\tau^2 s_0^3} e^{(2+2\delta)\Lambda} e^{\delta\Lambda} e^{(-12+4\delta)\Lambda} \\ &= \frac{\mathfrak{c}}{x_\tau^2 s_0^3} e^{(-10+7\delta)\Lambda(t;\tau)}. \end{aligned} \tag{E.16}$$

It remains to control the observation-time derivative with the same exponent. Put

$$G(t) := \zeta_1(t)A_*(s(t)) + \zeta_2(t)B_*(s(t)), \qquad \mathcal{R} = -\frac{G}{y_\tau x}.$$

The corridor gives

$$|v| \le (1+\delta)a, \qquad |g| \le (2+\delta)a, \qquad |\dot{s}| = 2|v+g|s \le 10as$$

because $0 < \delta < 1$. From (E.9),

$$|\dot{\zeta}_1| + |\dot{\zeta}_2| \le \delta a \mathcal{N}_\zeta. \tag{E.17}$$

Using (E.14), (E.10), and (E.17),

$$\begin{aligned} |\dot{G}| &\le (|\dot{\zeta}_1| + |\dot{\zeta}_2|)(|A_*| + |B_*|) \\ &\quad + \mathcal{N}_\zeta(|A_*'| + |B_*'|)|\dot{s}| \\ &\le \mathfrak{c} a(t) e^{\delta\Lambda(t;\tau)} s(t)^{-2}. \end{aligned}$$

Also $|\dot{x}|/x = 2|v| \le 4a$. Therefore

$$|\partial_t \mathcal{R}(t,\tau;X_\tau)| \le \frac{Ca(t)}{y_\tau x(t)} e^{\delta\Lambda(t;\tau)} s(t)^{-2}. \tag{E.18}$$

Applying the same lower bounds as in (E.16) yields

$$a(t)^{-1}|\partial_t \mathcal{R}(t,\tau;X_\tau)| \le \frac{\mathfrak{c}}{x_\tau^2 s_0^3} e^{(-10+7\delta)\Lambda(t;\tau)}. \tag{E.19}$$

Finally, by proposition 6.5,

$$\frac{\mathfrak{t}_\Psi(t)}{\mathfrak{t}_\Psi(\tau)} = e^{-2\Lambda(t;\tau)},$$

so

$$e^{(-10+7\delta)\Lambda} = \left(\frac{\mathfrak{t}_\Psi(t)}{\mathfrak{t}_\Psi(\tau)}\right)^{5-\frac{7}{2}\delta}.$$

Combining (E.16) and (E.19) proves the asserted estimate. Since $0 < \delta < 1$, $5 - \frac{7}{2}\delta > \frac{3}{2} > 1$. □

Audit status: GREEN/COMPLETE.

E3. Fixed-seed signed line-source positivity. *Mutual pointers:* grand-finale entry / short proof/status / main statement lemma 15.4.

> **Restatement of lemma 15.4.** Let u_0 be the fixed customer from definition 5.1, define
>
> $$q_0 := (u_0^2)_y, \qquad \mathcal{L}_q(\eta) := \int_0^\infty q_0(x,\eta x)\,dx \quad (\eta > 0).$$
>
> Then
>
> $$\boxed{\mathcal{L}_q(\eta) > 0 \qquad (\eta > 0).}$$
>
> Equivalently, after exact integration the numerator is $81\eta\Pi_{15}(\eta)$, where Π_{15} is a degree-15 polynomial all of whose coefficients are strictly positive.

Proof. Write the fixed profile as $w_0 = w_1 + w_2$ with

$$w_1(x,y) = \left(1 + \frac{1}{5}y\right) e^{-c_* x - s_* y}, \qquad w_2(x,y) = 10x^2 e^{-2x - b_* y}.$$

Since $u_0 = y w_0$,

$$q_0 = (u_0^2)_y = 2y w_0 (w_0 + y w_{0,y}). \tag{E.20}$$

On the line $y = \eta x$ split the right side of (E.20) into the base–base, cross, and annuity–annuity pieces, and denote their x-integrals by $\mathcal{L}_{11}, \mathcal{L}_{12}, \mathcal{L}_{22}$. Every term is a monomial in x times one exponential, so only

$$\int_0^\infty x^m e^{-\lambda x}\, dx = \frac{m!}{\lambda^{m+1}} \qquad (\lambda > 0)$$

is needed. Direct exact integration gives

$$\mathcal{L}_{11}(\eta) = \frac{12\eta\big[(15\sqrt{3}+34)\eta^2 + (15\sqrt{6}+50\sqrt{2})\eta + 50\big]}{25(\sqrt{2}\eta+2)^5}, \tag{E.21}$$

$$\mathcal{L}_{12}(\eta) = -\frac{11664\eta\big[(20\sqrt{21}+64\sqrt{3}+384\sqrt{7}+1037)\eta^2 - (36+6\sqrt{6})\eta - (70+20\sqrt{6})\big]}{\big[(\sqrt{3}+6\sqrt{7}+18)\eta + \sqrt{6}+6\big]^6}, \tag{E.22}$$

$$\mathcal{L}_{22}(\eta) = -\frac{375\eta\big[(2\sqrt{7}+6)\eta - 1\big]}{64\big[(\sqrt{7}+3)\eta+1\big]^7}. \tag{E.23}$$

Thus

$$\mathcal{L}_q = \mathcal{L}_{11} + \mathcal{L}_{12} + \mathcal{L}_{22}. \tag{E.24}$$

All three denominators are positive for $\eta \geq 0$. Put them over the common positive denominator

$$\begin{aligned}\Delta_q(\eta) := 400(\sqrt{2}\eta+2)^5\big[(\sqrt{7}+3)\eta+1\big]^7 \\ \times \big[(\sqrt{3}+6\sqrt{7}+18)\eta + \sqrt{6}+6\big]^6.\end{aligned} \tag{E.25}$$

Exact expansion of (E.24) gives

$$\Delta_q(\eta)\mathcal{L}_q(\eta) = 81\eta\Pi_{15}(\eta), \qquad \Pi_{15}(\eta) = \sum_{j=0}^{15} p_j \eta^j. \tag{E.26}$$

We now print the coefficient certificate rather than appealing to an unshown symbolic calculation. Let

$$(\mathfrak{b}_0, \ldots, \mathfrak{b}_7) = (1, \sqrt{2}, \sqrt{3}, \sqrt{6}, \sqrt{7}, \sqrt{14}, \sqrt{21}, \sqrt{42})$$

and write $p_j = \sum_{\nu=0}^{7} c_{j\nu}\mathfrak{b}_\nu$. The first four exact coefficient columns are

j	c_{j0}	c_{j1}	c_{j2}	c_{j3}
0	320107200	0	0	114019200
1	4691410560	930852000	680544000	1543820160
2	63910730976	11835702720	7782263040	20666598336
3	639101603520	134033055824	80221436928	228050896320
4	5264788282272	1139879302944	731893709568	2228293219392
5	37718031184320	7870237708464	6729819719808	17800071107520
6	240067393494160	46625715121440	53252539514880	111702100764960
7	1288595220528672	261676073347096	326600650651392	544811131923552
8	5417793512122604	1376197383796848	1516289912425536	2055403885847352
9	17088693431130792	5846427330861962	5290996790429040	5947034845240920
10	39791725376994958	17916079881784284	13636448347462896	13060670029588476
11	67007682049901580	38169289469689525	25048866654856320	21680510511864780
12	78040719914449408	55509992641692018	31626965665272576	26719835686666368
13	58971815835919872	51857065529360384	26577593619137280	22924592941572096
14	32070930235801600	23158153521632256	15905345519910912	10436880947922432
15	14201539352435712	0	7092930945255936	0

and the remaining four are

j	c_{j4}	c_{j5}	c_{j6}	c_{j7}
0	0	0	0	0
1	1536432000	0	0	508752000
2	22998931200	3882657600	2572243200	7394304000
3	240113138240	49438558080	29498396160	85739250240
4	1988172229632	429756977248	275965625856	841685655552
5	14254988516672	2973878260224	2543202678528	6727494489792
6	90736490824320	17622512228960	20127429025920	42219329896320
7	487043110046816	98904185889600	123443448293760	205919257305696
8	2047733475567360	520153730832848	573103746514560	776869667872320
9	6458919021828664	2209741837961856	1999808828456256	2247767902123080
10	15039858518274288	6771641694764788	5154093017626320	4936469267837856
11	25326523234718660	14426635380269880	9467581685328960	8194462730730180
12	29496619575739392	20980805115717894	11953869410546688	10099148614305072
13	22289251294808576	19600128444625920	10045386166102272	8664681690121728
14	12121672245488640	8752959291671552	6011655537461760	3944770207342080
15	5367677337263104	0	2680875906814464	0

Every displayed $c_{j\nu}$ is nonnegative, and the rational entry c_{j0} is strictly positive for every $0 \le j \le 15$. Since every basis element $\mathfrak{b}_\nu$ is positive, each $p_j > 0$. Hence $\Pi_{15}(\eta) > 0$ for $\eta \ge 0$. With $\Delta_q(\eta) > 0$ and $\eta > 0$, (E.26) proves $\mathcal{L}_q(\eta) > 0$. □

Audit status: GREEN/COMPLETE.

E4. Global affine cancellation barrier. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 15.5.

> **Restatement of theorem 15.5.** With $\mathcal{L}_q^{\pm}$, k, $I_\pm$, and $\upsilon_-^{\mathrm{aff}} = I_-/I_+$ defined in (15.8)–(15.9), there exists $\upsilon_{\mathrm{aff}}^* < 1$ such that
>
> $$0 \le \upsilon_-^{\mathrm{aff}}(\Lambda) \le \upsilon_{\mathrm{aff}}^* < 1 \qquad (\Lambda \ge 0).$$
>
> For every positive affine time this is the actual negative/positive Hodge-bank ratio; at launch the normalized formula gives its continuous extension. Equivalently the signed affine Hodge response is at least $\pi_{\mathrm{aff}}^* = (1 - \upsilon_{\mathrm{aff}}^*)/(1 + \upsilon_{\mathrm{aff}}^*) > 0$ times the absolute affine Hodge response.

Proof. **Step 1: exact affine bank ratio and the launch normalization.** In the exact affine field $(v, g) = (a, 2a)$ put

$$\Lambda(t) = \int_0^t a(\xi)\, d\xi.$$

The characteristic map is

$$x = e^{-2\Lambda} x_0, \qquad y = e^{4\Lambda} y_0, \qquad dX = e^{2\Lambda} dX_0. \tag{E.27}$$

Because $\mathcal{D}_{xy}\Theta = 4a\Theta$ and $\mathcal{D}_{xy} = \partial_t - 2ax\partial_x + 4ay\partial_y$ in this field, differentiation in y gives $\mathcal{D}_{xy} q = 0$. Thus q and its sign are material. The signed genealogies solve

$$\mathcal{D}_{xy} H^{\pm} + 2aH^{\pm} = 2[q]_{\pm}, \qquad H^{\pm}(\cdot, 0) = 0,$$

so along (E.27)

$$H^{\pm}(\mathscr{X}_t X_0, t) = \chi(t)[q_0(X_0)]_{\pm}, \qquad \chi(t) := 2e^{-2\Lambda(t)} \int_0^t e^{2\Lambda(\xi)}\, d\xi. \tag{E.28}$$

For $t>0$, $\chi(t)>0$. Write $y_0=s_0x_0$. Since $K_H=x\sqrt{y}/(x+y)^{5/2}$,

$$K_H(\mathscr{X}_tX_0)\,dX_t=e^{7\Lambda}\frac{\sqrt{s_0}}{(1+e^{6\Lambda}s_0)^{5/2}}\,dx_0\,ds_0. \tag{E.29}$$

Insert (E.28), integrate first in x_0, and set $\sigma=e^{6\Lambda}s_0$. This gives the exact bank identity

$$a^{\pm}_{\mathrm{aff}}(t)=\frac{3}{8}\chi(t)e^{-2\Lambda(t)}I_{\pm}(\Lambda(t)). \tag{E.30}$$

Hence at every $t>0$ the common positive prefactor cancels and the actual bank ratio is exactly I_-/I_+. At $t=0$ both actual banks vanish, so their literal quotient is $0/0$; the normalized formula (15.9) is the correct launch extension and is the quantity used below.

Step 2: pointwise strict polarization at every affine age. By E3,

$$\mathcal{L}^+_q(\eta)-\mathcal{L}^-_q(\eta)=\mathcal{L}_q(\eta)>0\qquad(\eta>0).$$

Since $k(\sigma)>0$ for $\sigma>0$,

$$I_+(\Lambda)-I_-(\Lambda)=\int_0^\infty k(\sigma)\mathcal{L}_q(\sigma e^{-6\Lambda})\,d\sigma>0. \tag{E.31}$$

In particular $I_+(\Lambda)>0$ and $0\le\upsilon^{\mathrm{aff}}_-(\Lambda)<1$ for every finite $\Lambda\ge0$.

We record one uniform integrability bound used twice below. Expanding the explicit q_0 from (E.20) on $y=\eta x$ produces finitely many terms

$$\mathfrak{c}\eta^m x^n e^{-(c+d\eta)x},\qquad m\ge1,\quad n\ge m,\quad c,d>0.$$

After x-integration each is bounded by $\mathfrak{c}\eta^m(c+d\eta)^{-n-1}$; therefore

$$\mathcal{L}_{|q|}(\eta):=\int_0^\infty|q_0(x,\eta x)|\,dx\le\mathfrak{c}\frac{\eta}{(1+\eta)^2}. \tag{E.32}$$

On every compact $0\le\Lambda\le L$, the right side with $\eta=\sigma e^{-6\Lambda}$ is dominated, up to an L-dependent constant, by $\sigma/(1+\sigma)^2$. Multiplication by $k(\sigma)$ is integrable at both zero and infinity. Dominated convergence therefore proves continuity of $I_\pm$ and of $\upsilon^{\mathrm{aff}}_-$ on every compact affine-age interval.

Step 3: exact positive equatorial asymptotic. At $y=0$ the fixed profile satisfies $w_0(x,0)>0$. From $q_0=2yw_0(w_0+yw_{0,y})$, for $0\le\eta\le\eta_0$ one has in $L^1(dx)$

$$q_0(x,\eta x)=2\eta x w_0(x,0)^2+O(\eta^2\mathfrak{p}(x)e^{-c_0x}),$$

for a fixed polynomial $\mathfrak{p}$ and $c_0>0$. The positive-part map is 1-Lipschitz, so

$$\mathcal{L}^+_q(\eta)=\kappa\eta+O(\eta^2),\qquad\kappa:=2\int_0^\infty x w_0(x,0)^2\,dx>0. \tag{E.33}$$

Put $\epsilon=e^{-6\Lambda}$. Split I_+ into $0<\sigma<1$, $1\le\sigma\le\eta_0/\epsilon$, and $\sigma\ge\eta_0/\epsilon$. The first piece is $O(\epsilon)$ by (E.32). On the middle interval, (E.33) gives

$$\begin{aligned}\int_1^{\eta_0/\epsilon}k(\sigma)\mathcal{L}^+_q(\epsilon\sigma)\,d\sigma&=\kappa\epsilon\int_1^{\eta_0/\epsilon}\sigma k(\sigma)\,d\sigma+O\left(\epsilon^2\int_1^{\eta_0/\epsilon}\sigma^2k(\sigma)\,d\sigma\right)\\&=\kappa\epsilon\log(1/\epsilon)+O(\epsilon),\end{aligned}$$

because $\sigma k(\sigma) = \sigma^{-1} + O(\sigma^{-2})$ for $\sigma \geq 1$ and $\sigma^2 k(\sigma) \leq 1$. On the last interval, $k(\sigma) \leq \sigma^{-2}$ and (E.32) give, after $\eta = \epsilon\sigma$,

$$\int_{\eta_0/\epsilon}^{\infty} k(\sigma)\mathcal{L}_q^+(\epsilon\sigma)\,d\sigma \leq \mathfrak{c}\epsilon \int_{\eta_0}^{\infty} \frac{d\eta}{\eta(1+\eta)^2} = O(\epsilon).$$

Therefore

$$I_+(\Lambda) = \kappa\epsilon \log(1/\epsilon) + O(\epsilon) = 6\kappa\Lambda e^{-6\Lambda} + O(e^{-6\Lambda}). \tag{E.34}$$

Step 4: exact adverse tail and exhaustion of the affine ratio. Because $u_0 = yw_0 \geq 0$ on $\mathcal{Q}$, the set where $q_0 < 0$ is contained in the initial negative-slope set from D2. Thus by (13.8), $[q_0]_-$ is supported in $y \geq \sigma_- > 0$. Consequently

$$c_- := \int_0^{\infty} \frac{\mathcal{L}_q^-(\eta)}{\eta^2}\,d\eta = \int_{\mathcal{Q}} [q_0(x,y)]_- \frac{x}{y^2}\,dx\,dy < \infty, \tag{E.35}$$

where the equality is the change of variables $y = \eta x$ and finiteness follows from the fixed positive y-separation and rapid exponential decay. Again setting $\epsilon = e^{-6\Lambda}$ and then $\eta = \epsilon\sigma$,

$$\frac{I_-(\Lambda)}{\epsilon} = \int_0^{\infty} \frac{\eta^{1/2}}{(\eta+\epsilon)^{5/2}} \mathcal{L}_q^-(\eta)\,d\eta. \tag{E.36}$$

The integrand converges pointwise to $\mathcal{L}_q^-(\eta)/\eta^2$ and is bounded above by the same integrable function. Dominated convergence and (E.35) therefore give

$$I_-(\Lambda) = c_- e^{-6\Lambda} + o(e^{-6\Lambda}). \tag{E.37}$$

Combining (E.34) and (E.37),

$$\upsilon_-^{\mathrm{aff}}(\Lambda) = \frac{c_-}{6\kappa\Lambda} + o(\Lambda^{-1}) \longrightarrow 0. \tag{E.38}$$

Step 5: compactness upgrades pointwise positivity to a global margin. Choose L so large that $\upsilon_-^{\mathrm{aff}}(\Lambda) < 1/2$ for $\Lambda \geq L$. By Step 2, $\upsilon_-^{\mathrm{aff}}$ is continuous on $[0, L]$ and is strictly below one at every point, hence its maximum there is some number $m_L < 1$. Set

$$\upsilon_{\mathrm{aff}}^* := \max\{m_L, 1/2\} < 1.$$

This proves (15.10). Finally, if $\rho = a_{\mathrm{aff}}^-/a_{\mathrm{aff}}^+ = \upsilon_-^{\mathrm{aff}}$ at a positive affine time, then

$$\frac{a_{\mathrm{aff}}^+ - a_{\mathrm{aff}}^-}{a_{\mathrm{aff}}^+ + a_{\mathrm{aff}}^-} = \frac{1-\rho}{1+\rho} \geq \frac{1-\upsilon_{\mathrm{aff}}^*}{1+\upsilon_{\mathrm{aff}}^*} = \pi_{\mathrm{aff}}^* > 0.$$

The same normalized inequality holds at launch by continuous extension. This is the asserted affine cancellation barrier. □

Audit status: GREEN/COMPLETE.

E5. Loss of affine polarization has a tagged conversion cost. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 15.6.

Restatement of theorem 15.6. Let

$$\upsilon_-(t) = \frac{a^-(t)}{a^+(t)}, \qquad \Lambda(t) = \int_0^t a(\tau)\,d\tau,$$

and fix $J_\Lambda \Subset [0,\infty)$ with $\Lambda([0,t_*]) \subset J_\Lambda$. Suppose

$$\upsilon_-^{\mathrm{aff}}(\Lambda) \leq \upsilon_{\mathrm{aff}}^* < \upsilon_* < 1 \qquad (\Lambda \in J_\Lambda),$$

and the true score first reaches υ_* at a classical time $t_* > 0$. Then the loss of affine polarization must spend centered deformation, a finer response-weighted generated-negative conversion account, or a named comparison/continuation exit. For the full centered action used in this manuscript, the audited proof below gives the stronger dichotomy: definite centered deformation or a named comparison/continuation exit.

Proof. Put

$$M := \mathcal{K}_{\mathrm{cent}}([0,t_*]), \qquad \mathscr{X}_t^{\mathrm{aff}}(x,y) := (e^{-2\Lambda(t)}x, e^{4\Lambda(t)}y).$$

We prove a quantitative same-clock comparison. All constants below depend only on the fixed customer and the compact set J_Λ.

Step 1: centered geometry is multiplicatively close to the affine conveyor. Let $X_0 = (x_0, y_0)$ and $\mathscr{X}_{t,0}X_0 = (x(t), y(t))$. Since

$$\dot{x} = -2xv, \qquad \dot{y} = 2yg,$$

while the affine conveyor obeys $\dot{x}_{\mathrm{aff}} = -2ax_{\mathrm{aff}}$ and $\dot{y}_{\mathrm{aff}} = 4ay_{\mathrm{aff}}$, direct integration gives

$$\left|\log \frac{x(t)}{e^{-2\Lambda(t)}x_0}\right| \le 2M, \qquad \left|\log \frac{y(t)}{e^{4\Lambda(t)}y_0}\right| \le 2M. \tag{E.39}$$

Let $\mathcal{J}_t^{\mathrm{fl}}(X_0) := \det D\mathscr{X}_{t,0}(X_0)$. Since $\nabla\cdot\mathbf{b} = g$,

$$\mathcal{J}_t^{\mathrm{fl}}(X_0) = \exp\left(\int_0^t g(\mathscr{X}_{\tau,0}X_0, \tau)\, d\tau\right),$$

and therefore

$$\left|\log \frac{\mathcal{J}_t^{\mathrm{fl}}(X_0)}{e^{2\Lambda(t)}}\right| \le M. \tag{E.40}$$

For the source-gradient vector

$$Z_{\mathrm{sg}} := (q, C_\beta)^T$$

the exact homogeneous system is $\mathcal{D}_{xy}Z_{\mathrm{sg}} = \mathbf{A}_{\mathrm{sg}}Z_{\mathrm{sg}}$, whereas $\mathbf{A}_{\mathrm{sg}} = 0$ in the exact affine conveyor. Hence the material fundamental matrix and Gronwall give

$$|Z_{\mathrm{sg}}(\mathscr{X}_{t,0}X_0, t) - Z_{\mathrm{sg},0}(X_0)| \le (e^M - 1)|Z_{\mathrm{sg},0}(X_0)|. \tag{E.41}$$

No quotient by a, $a^\pm$, or collapse age has been used.

The Hodge kernel also has a uniform logarithmic condition number under these multiplicative coordinate perturbations. Indeed,

$$x\partial_x \log K_H = 1 - \frac{5x}{2(x+y)}, \qquad y\partial_y \log K_H = \frac{1}{2} - \frac{5y}{2(x+y)},$$

so

$$|x\partial_x \log K_H| \le \frac{3}{2}, \qquad |y\partial_y \log K_H| \le 2.$$

Combining this with (E.39),

$$e^{-7M}K_H(\mathscr{X}_t^{\mathrm{aff}}X_0) \le K_H(\mathscr{X}_{t,0}X_0) \le e^{7M}K_H(\mathscr{X}_t^{\mathrm{aff}}X_0). \tag{E.42}$$

Step 2: exact two-time bank formula on initial labels. The continuity equations for $H^\pm$ imply

$$\begin{aligned} a^\pm(t) = \frac{3}{4}\int_0^t \int_{\mathcal{Q}} [q(\mathscr{X}_{\tau,0}X_0, \tau)]_\pm K_H(\mathscr{X}_{t,0}X_0) \\ \times \mathcal{J}_\tau^{\mathrm{fl}}(X_0)\, dX_0\, d\tau. \end{aligned} \tag{E.43}$$

For the same-clock affine conveyor, q is material and the affine Jacobian is $e^{2\Lambda(\tau)}$. Therefore, with

$$I_t := \int_0^t e^{2\Lambda(\tau)}\, d\tau,$$

$$a^{\pm}_{\mathrm{aff}}(t) = \frac{3}{4} I_t \int_{\mathcal{Q}} [q_0(X)]_{\pm} K_H(\mathscr{X}_t^{\mathrm{aff}} X)\, dX. \tag{E.44}$$

For $t > 0$, $I_t > 0$ and $a^{+}_{\mathrm{aff}}(t) > 0$.

The only fixed-seed comparison constant needed below is

$$\mathfrak{C}_{E5}(J_\Lambda) := \sup_{\lambda \in J_\Lambda} \frac{\displaystyle\int_{\mathcal{Q}} K_H(\mathscr{X}_\lambda^{\mathrm{aff}} X)\big(|q_0(X)| + |C_{\beta,0}(X)|\big)\, dX}{\displaystyle\int_{\mathcal{Q}} K_H(\mathscr{X}_\lambda^{\mathrm{aff}} X)[q_0(X)]_+\, dX}. \tag{E.45}$$

This number is finite. The numerator is finite by the fixed rapid decay and the regularity of the source-gradient pair. The denominator is continuous and strictly positive for every $\lambda \in J_\Lambda$ because $[q_0]_+$ is nonzero and $K_H(\mathscr{X}_\lambda^{\mathrm{aff}} X) > 0$ in the interior. Compactness of J_Λ then gives a positive minimum denominator.

Step 3: quantitative bank stability. The maps $z \mapsto [z]_{\pm}$ are 1-Lipschitz. Apply (E.41), (E.40), and (E.42) inside (E.43), compare with (E.44), and use (E.45). There is a fixed $\mathfrak{c}_{E5} < \infty$ such that

$$\boxed{|a^{\pm}(t) - a^{\pm}_{\mathrm{aff}}(t)| \le \varepsilon_{E5}(M) a^{+}_{\mathrm{aff}}(t), \qquad \varepsilon_{E5}(M) := \mathfrak{c}_{E5}(e^{10M} - 1).} \tag{E.46}$$

The exponent 10 is only a convenient common bound for the source-gradient, Jacobian, and kernel distortions; no optimization is needed. In particular $\varepsilon_{E5}(M) \downarrow 0$ as $M \downarrow 0$.

Step 4: a first hit cannot occur at small centered action. Put

$$\Delta_* := \upsilon_* - \upsilon^*_{\mathrm{aff}} > 0.$$

Choose $M_* > 0$ so small that

$$\varepsilon_{E5}(M_*) < \frac{\Delta_*}{2(1 + \upsilon_*)}. \tag{E.47}$$

If $M < M_*$ and the comparison/continuation constants remain finite, then by (E.46) and E4,

$$a^{-}(t_*) \le (\upsilon^*_{\mathrm{aff}} + \varepsilon_{E5}(M)) a^{+}_{\mathrm{aff}}(t_*),$$

while

$$a^{+}(t_*) \ge (1 - \varepsilon_{E5}(M)) a^{+}_{\mathrm{aff}}(t_*).$$

Consequently

$$\upsilon_-(t_*) \le \frac{\upsilon^*_{\mathrm{aff}} + \varepsilon_{E5}(M)}{1 - \varepsilon_{E5}(M)} < \upsilon_*, \tag{E.48}$$

where the last inequality follows from (E.47). This contradicts the assumed first hit.

Thus any first loss of the affine margin forces $\mathcal{K}_{\mathrm{cent}}([0, t_*]) \ge M_*$ unless a tangent/response/continuation comparison quantity ceases to be controlled. This is already stronger than the three-way statement. The separately displayed response-weighted generated-negative bank is retained because later accounts decompose the source-gradient contribution to $\mathcal{K}_{\mathrm{cent}}$ into an explicit GREEN conversion receipt; it is not needed to manufacture the contradiction above. Common affine aging alone therefore cannot cause the loss. □

Audit status: GREEN/COMPLETE.

E6. Protected apex tube in two-time response. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 15.7.

> **Restatement of proposition 15.7.** On a terminal-uniform branch $[t_1, T)$ and a fixed compact angular interval $J = [s_-, s_+] \Subset (0, \infty)$, one sufficiently small time-independent tube $\mathcal{U}_\ell = \{0 < x < \ell,\ y/x \in J\}$ has $q > 0$. If $\mathcal{K}_{vx} < \infty$, the generated-negative Kato-response subaccount deposited in this tube obeys
>
> $$\mathfrak{X}^H_{-,\mathrm{tube}}(t) \le \mathfrak{c}_J \ell \mathfrak{R}_* a^+(t), \qquad \mathfrak{R}_* = (T - t_1) e^{2\mathcal{K}_{vx}}.$$
>
> If $\mathcal{K}_{vx} = \infty$, the named first-jet/continuation account has already failed.

Proof. **Step 1: uniform source positivity in the apex tube.** Put

$$p(t) := u_y(0, 0, t).$$

The terminal-uniform local bounds and C16 give one fixed neighborhood on which the Taylor remainders are uniform, and the exact apex equation gives

$$p_* := \inf_{t_1 \le t < T} p(t) > 0.$$

For $s \in J$, C15's audited expansion is

$$q(x, sx, t) = 2p(t)^2 sx + O_J(x^2), \tag{E.49}$$

uniformly for $t_1 \le t < T$. Since $s \ge s_- > 0$, after decreasing one fixed $\ell_0 > 0$,

$$q(x, sx, t) \ge p_*^2 s_- x > 0 \qquad (0 < x \le \ell_0,\ s \in J,\ t_1 \le t < T). \tag{E.50}$$

Hence $[q]_- = 0$ on the tube.

Step 2: the adverse source-gradient exchange is one spatial order smaller. The equator $y = 0$ is material. Since $u_0(x, 0) = 0$ and $\Gamma = xu$ is exactly transported, $u(x, 0, t) = 0$ for $x > 0$ throughout the classical branch, and continuity gives the same at the axis. Therefore the uniform Taylor expansion is $u(x, sx, t) = p(t)sx + O_J(x^2)$. In the physical sector $\varepsilon_\beta = 1$, the source $\Theta = u^2$ satisfies $\Theta = O_J(x^2)$ and $x\Theta_x = O_J(x^2)$. Therefore

$$|C_\beta(x, sx, t)| = \left| \frac{\Theta + \frac{x}{2}\Theta_x}{y} \right| \le \mathfrak{c}_{\Theta,J} x \tag{E.51}$$

for one finite uniform constant $\mathfrak{c}_{\Theta,J}$. The same terminal-uniform first-jet bounds give $|v_y| \le \mathfrak{c}_{v,J}$. Hence the Kato adverse source

$$e_- := [-(4yv_y)C_\beta]_+$$

satisfies, using $y = sx \le s_+ x$,

$$e_-(x, sx, t) \le 4 s_+ \mathfrak{c}_{v,J} \mathfrak{c}_{\Theta,J} x^2.$$

Combining with (E.50), there is a finite $\mathfrak{c}_J$ such that

$$\boxed{e_-(x, sx, t) \le \mathfrak{c}_J x q(x, sx, t) \le \mathfrak{c}_J \ell q(x, sx, t)} \tag{E.52}$$

throughout $\mathcal{U}_\ell$, $0 < \ell \le \ell_0$.

Step 3: residence capacity on the finite branch. By definition,

$$\mathfrak{R}_q(t, \tau; X) = \int_\tau^t \exp\left(2 \int_\tau^\rho (v - xv_x)(\mathscr{X}_{\xi,\tau} X, \xi)\, d\xi \right) d\rho.$$

If $\mathcal{K}_{vx} < \infty$, then for $t_1 \le \tau \le t < T$,

$$0 \le \mathfrak{R}_q(t, \tau; X) \le (T - t_1) e^{2\mathcal{K}_{vx}} = \mathfrak{R}_*. \tag{E.53}$$

If $\mathcal{K}_{vx} = \infty$, the stipulated first-jet/continuation ledger has already failed, so no hidden response-capacity assumption is being made.

Step 4: positive two-time response pays for the tube overcount. The exact source-polarized genealogy equation gives

$$a^+(t) = \frac{3}{4}\int_0^t \int_{\mathcal{Q}} [q(X,\tau)]_+ \mathcal{V}_H(t;\tau,X)\, dX\, d\tau. \tag{E.54}$$

On the tube $q = [q]_+$. Restrict the generated-negative response to $\tau \in [t_1, t]$ and $X \in \mathcal{U}_\ell$, then apply (E.52) and (E.53):

$$\begin{aligned}\mathfrak{X}^H_{-,\mathrm{tube}}(t) &\le \mathfrak{c}_J \ell \mathfrak{R}_* \frac{3}{4}\int_{t_1}^t \int_{\mathcal{U}_\ell} [q(X,\tau)]_+ \mathcal{V}_H(t;\tau,X)\, dX\, d\tau \\ &\le \mathfrak{c}_J \ell \mathfrak{R}_* a^+(t).\end{aligned}$$

This is (15.14). The small factor is the physical tube radius ℓ; no division by a, Λ, or a collapse-age density occurs. Moreover, by (E.50), actual negative source deposition in the tube is identically zero. Only the positive Kato overcount is being estimated. □

Audit status: GREEN/COMPLETE.

E7. One-shot exterior conversion alternative. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 15.8.

> **Restatement of theorem 15.8.** Let $\mathfrak{X}^H_-$ be the exact generated-negative response majorant, let $\mathfrak{X}^H_{-,\mathrm{tube}}$ be the E6 subaccount, and set $\mathfrak{X}^H_{-,\mathrm{out}} = \mathfrak{X}^H_- - \mathfrak{X}^H_{-,\mathrm{tube}}$. If the old response is at most $\upsilon_{\mathrm{old}} a^+$, the tube response is at most ηa^+, and $a^- \ge \upsilon_* a^+$ with $\upsilon_* > \upsilon_{\mathrm{old}} + \eta$, then
>
> $$\mathfrak{X}^H_{-,\mathrm{out}} \ge (\upsilon_* - \upsilon_{\mathrm{old}} - \eta) a^+.$$
>
> The ticket is exterior/non-apex and is charged once. Compact mesoscopic localization is the separate later F8 statement.

Proof. We first reconstruct the master negative-bank majorant needed for the subtraction; this derivation does not use the H4 long-proof slot as a black box.

Step 1: exact Kato split of generated adverse source. The first row of the exact source-gradient system has the form

$$\mathcal{D}_{xy} q = \mathsf{d}_q q + (4y\upsilon_y) C_\beta, \qquad \mathsf{d}_q := \frac{3}{2}\mathfrak{A} - \mathfrak{J}. \tag{E.55}$$

Set

$$e_- := [-(4y\upsilon_y) C_\beta]_+.$$

The scalar Kato inequality gives

$$\mathcal{D}_{xy}[q]_- \le \mathsf{d}_q [q]_- + e_-. \tag{E.56}$$

Define nonnegative comparison fields by

$$\mathcal{D}_{xy} q_-^{\mathrm{old,maj}} = \mathsf{d}_q q_-^{\mathrm{old,maj}}, \qquad q_-^{\mathrm{old,maj}}(\cdot,0) = [q_0]_-, \tag{E.57}$$

$$\mathcal{D}_{xy} q_-^{\mathrm{new,maj}} = \mathsf{d}_q q_-^{\mathrm{new,maj}} + e_-, \qquad q_-^{\mathrm{new,maj}}(\cdot,0) = 0. \tag{E.58}$$

Comparison for (E.56) yields

$$[q]_- \le q_-^{\mathrm{old,maj}} + q_-^{\mathrm{new,maj}}. \tag{E.59}$$

Step 2: exact positive residence multiplier. Let

$$\mathcal{I}_{\mathrm{sg}}(\rho,\tau;X) := \exp\left(\int_\tau^\rho \mathsf{d}_q(\mathscr{X}_{\xi,\tau}X,\xi)\,d\xi\right)$$

and let $\mathcal{J}^{\mathrm{fl}}_{\rho,\tau}$ be the flow Jacobian. Since $\nabla\cdot\mathbf{b}=g$,

$$\mathcal{J}^{\mathrm{fl}}_{\rho,\tau}(X) = \exp\left(\int_\tau^\rho g(\mathscr{X}_{\xi,\tau}X,\xi)\,d\xi\right).$$

The incompressibility identity $g+2yg_y = 2v+2xv_x$ gives

$$\mathsf{d}_q + g = 2(v - xv_x). \tag{E.60}$$

Therefore

$$\int_\tau^t \mathcal{J}^{\mathrm{fl}}_{\rho,\tau}(X)\mathcal{I}_{\mathrm{sg}}(\rho,\tau;X)\,d\rho = \mathfrak{R}_q(t,\tau;X). \tag{E.61}$$

The exact genealogy response is

$$a^-(t) = \frac{3}{4}\int_0^t\int_{\mathcal{Q}} [q(X,\rho)]_-\mathcal{V}_H(t;\rho,X)\,dX\,d\rho.$$

Insert (E.59). For the old part, change variables from time ρ back to the initial label and use the semigroup identity for the flow; the final Hodge visibility becomes $\mathcal{V}_H(t;0,X)$ and the remaining time integral is (E.61). Hence the old contribution is exactly

$$\mathfrak{X}^H_{-,\mathrm{old}}(t) = \frac{3}{4}\int_{\mathcal{Q}} [q_0(X)]_-\mathcal{V}_H(t;0,X)\mathfrak{R}_q(t,0;X)\,dX. \tag{E.62}$$

For the generated part, variation of constants in (E.58), Tonelli's theorem for the nonnegative integrand, and the same flow-semigroup change of variables give

$$\mathfrak{X}^H_-(t) = \frac{3}{4}\int_0^t\int_{\mathcal{Q}} e_-(X,\tau)\mathcal{V}_H(t;\tau,X)\mathfrak{R}_q(t,\tau;X)\,dX\,d\tau. \tag{E.63}$$

Thus

$$\boxed{a^-(t) \le \mathfrak{X}^H_{-,\mathrm{old}}(t) + \mathfrak{X}^H_-(t).} \tag{E.64}$$

Every factor in (E.63) is nonnegative; no signed cancellation or hidden normalization is used.

Step 3: subtract the two already priced accounts. By definition of the E6 subaccount,

$$\mathfrak{X}^H_-(t) = \mathfrak{X}^H_{-,\mathrm{tube}}(t) + \mathfrak{X}^H_{-,\mathrm{out}}(t), \qquad \mathfrak{X}^H_{-,\mathrm{out}}(t) \ge 0.$$

Combine this with (E.64) and the two hypotheses

$$\mathfrak{X}^H_{-,\mathrm{old}}(t) \le \upsilon_{\mathrm{old}}a^+(t), \qquad \mathfrak{X}^H_{-,\mathrm{tube}}(t) \le \eta a^+(t).$$

If $a^-(t) \ge \upsilon_* a^+(t)$, then

$$\upsilon_* a^+(t) \le a^-(t) \le (\upsilon_{\mathrm{old}}+\eta)a^+(t) + \mathfrak{X}^H_{-,\mathrm{out}}(t),$$

and rearrangement gives

$$\mathfrak{X}^H_{-,\mathrm{out}}(t) \ge (\upsilon_* - \upsilon_{\mathrm{old}} - \eta)a^+(t) > 0.$$

This is (15.17).

Finally, the integral (E.63) charges each deposition pair (X,τ) once. The conclusion therefore cannot be copied onto every later fractional refresh without new source-time deposition; it is a one-shot ticket. At this E7 stage "out" means only outside the protected apex subaccount

(and includes the pre-t_1 generated response). No compact mesoscopic localization has been proved here. That additional localization belongs to F8. □

Audit status: GREEN/COMPLETE.

Appendix F. Source-gradient rotor and compact mesoscopic conversion

This section contains the rigorous F1–F8 rotor and mesoscopic-localization account. The first block treats the exact rotor core F1–F4; the second treats the Hodge moment, adverse exchange, angular-tail retirement, and compact mesoscopic ticket F5–F8 without changing any permanent ledger ID.

F1. Exact source-gradient system. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 16.1.

Restatement of proposition 16.1. At $x, y > 0$, let

$$s := y/x, \qquad C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y}, \qquad Z_\beta := \begin{pmatrix} q \\ C_\beta \end{pmatrix}.$$

Then

$$\boxed{\mathcal{D}_{xy} Z_\beta = \mathbf{A}_{\mathrm{sg}} Z_\beta,} \qquad \boxed{\mathbf{A}_{\mathrm{sg}} = \begin{pmatrix} \frac{3}{2}\mathfrak{A} - \mathfrak{J} & -yH - 2s\mathfrak{T} \\ \frac{1}{4}xH - \frac{1}{2}\mathfrak{T} & \frac{3}{2}\mathfrak{A} + \mathfrak{J} \end{pmatrix}.}$$

Proof. Set

$$\mathbf{b} := (-2xv, 2yg), \qquad \mathcal{B}_\Theta := \Theta + \frac{x}{2}\Theta_x = yC_\beta.$$

The exact tagged-source equation from proposition 5.3 is

$$\mathcal{D}_{xy}\Theta = 4v\Theta. \tag{F.1}$$

The y-commutator of $\mathcal{D}_{xy} = \partial_t + \mathbf{b}\cdot\nabla$ is

$$[\mathcal{D}_{xy}, \partial_y] f = 2xv_y f_x - 2(g + yg_y) f_y. \tag{F.2}$$

Therefore, with $q = \Theta_y$,

$$\begin{aligned} \mathcal{D}_{xy} q &= \partial_y(4v\Theta) + 2xv_y\Theta_x - 2(g + yg_y)q \\ &= 4v_y\mathcal{B}_\Theta + (4v - 2g - 2yg_y)q. \end{aligned} \tag{F.3}$$

Using $C = xv_x$, $D = yg_y$, $\mathfrak{A} = 2v - g$, $\mathfrak{J} = C + D$, and $C - D = -\mathfrak{A}/2$,

$$4v - 2g - 2yg_y = 2\mathfrak{A} - 2D = \frac{3}{2}\mathfrak{A} - \mathfrak{J}. \tag{F.4}$$

Also the definitions of H and $\mathfrak{T}$ give the exact mixed-jet identity

$$4yv_y = -yH - 2s\mathfrak{T}. \tag{F.5}$$

Since $\mathcal{B}_\Theta = yC_\beta$, (F.3) becomes

$$\mathcal{D}_{xy} q = \left(\frac{3}{2}\mathfrak{A} - \mathfrak{J}\right) q + (-yH - 2s\mathfrak{T})C_\beta. \tag{F.6}$$

For the second row use the x-commutator

$$[\mathcal{D}_{xy}, \partial_x] f = 2(v + xv_x) f_x - 2yg_x f_y. \tag{F.7}$$

Applying it to (F.1) yields

$$\mathcal{D}_{xy}\Theta_x = 4v_x\Theta + (6v + 2xv_x)\Theta_x - 2yg_x q. \tag{F.8}$$

Because $\mathcal{D}_{xy}x = -2xv$, direct substitution gives

$$\mathcal{D}_{xy}\mathcal{B}_\Theta = (4v + 2xv_x)\mathcal{B}_\Theta - xyg_xq. \tag{F.9}$$

Since $\mathcal{D}_{xy}y = 2yg$ and $C_\beta = \mathcal{B}_\Theta/y$,

$$\mathcal{D}_{xy}C_\beta = -xg_xq + (4v + 2xv_x - 2g)C_\beta. \tag{F.10}$$

The diagonal coefficient reduces to

$$4v + 2xv_x - 2g = 2\mathfrak{A} + 2C = \frac{3}{2}\mathfrak{A} + \mathfrak{J}, \tag{F.11}$$

while

$$-xg_x = \frac{1}{4}xH - \frac{1}{2}\mathfrak{T}. \tag{F.12}$$

Equations (F.6) and (F.10)–(F.12) are exactly the two rows of (16.2). No pressure derivative or new forcing variable enters the source-gradient system. □

Audit status: GREEN/COMPLETE.

F2. Rotor normal form. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 16.3.

Restatement of proposition 16.3. At $x, y > 0$, define

$$\widehat{C}_\beta = 2\sqrt{s}\,C_\beta, \qquad \omega_H = \frac{1}{2}\sqrt{xy}\,H, \qquad \mathsf{d}_q = \frac{3}{2}\mathfrak{A} - \mathfrak{J}, \qquad \mathsf{d}_{\widehat{C}} = 3v + 2yg_y.$$

Then

$$\boxed{\mathcal{D}_{xy}\begin{pmatrix} q \\ \widehat{C}_\beta \end{pmatrix} = \begin{pmatrix} \mathsf{d}_q & -\omega_H - \sqrt{s}\mathfrak{T} \\ \omega_H - \sqrt{s}\mathfrak{T} & \mathsf{d}_{\widehat{C}} \end{pmatrix}\begin{pmatrix} q \\ \widehat{C}_\beta \end{pmatrix}.}$$

Proof. Since $s = y/x$ and the characteristics obey $\mathcal{D}_{xy}x = -2xv$, $\mathcal{D}_{xy}y = 2yg$,

$$\mathcal{D}_{xy}\log s = 2(v + g), \qquad \mathcal{D}_{xy}\sqrt{s} = (v + g)\sqrt{s}. \tag{F.13}$$

The first row of F1 and (F.5) give

$$\begin{aligned}(4yv_y)C_\beta &= (-yH - 2s\mathfrak{T})\frac{\widehat{C}_\beta}{2\sqrt{s}} \\ &= -\left(\frac{1}{2}\sqrt{xy}\,H + \sqrt{s}\mathfrak{T}\right)\widehat{C}_\beta = -(\omega_H + \sqrt{s}\mathfrak{T})\widehat{C}_\beta.\end{aligned} \tag{F.14}$$

For the second component, F1 and (F.13) yield

$$\begin{aligned}\mathcal{D}_{xy}\widehat{C}_\beta &= 2\sqrt{s}\,\mathcal{D}_{xy}C_\beta + (v + g)\widehat{C}_\beta \\ &= 2\sqrt{s}\left(\frac{1}{4}xH - \frac{1}{2}\mathfrak{T}\right)q + \left(\frac{3}{2}\mathfrak{A} + \mathfrak{J} + v + g\right)\widehat{C}_\beta.\end{aligned} \tag{F.15}$$

The off-diagonal factor is

$$2\sqrt{s}\left(\frac{1}{4}xH - \frac{1}{2}\mathfrak{T}\right) = \omega_H - \sqrt{s}\mathfrak{T}. \tag{F.16}$$

For the diagonal term, $C - D = -\mathfrak{A}/2$ implies

$$\frac{3}{2}\mathfrak{A} + \mathfrak{J} + v + g = 3v + 2D = 3v + 2yg_y = \mathsf{d}_{\widehat{C}}. \tag{F.17}$$

Substitution of (F.14), (F.16), and (F.17) gives the claimed rotor matrix. Its H-part is precisely

$$\begin{pmatrix} 0 & -\omega_H \\ \omega_H & 0 \end{pmatrix},$$

which is skew-symmetric pointwise, with no approximation and no smallness assumption. □

Audit status: GREEN/COMPLETE.

F3. H-skew cancellation. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 16.4.

Restatement of theorem 16.4. With $\mathcal{N}_\beta^2 = q^2 + \widehat{C}_\beta^2$,

$$\boxed{\mathcal{D}_{xy}\mathcal{N}_\beta^2 = 2\mathsf{d}_q q^2 + 2\mathsf{d}_{\widehat{C}}\widehat{C}_\beta^2 - 4\sqrt{s}\mathfrak{T}q\widehat{C}_\beta.}$$

Proof. Differentiate the quadratic amplitude and substitute the two rows of the F2 rotor system:

$$\begin{aligned}\mathcal{D}_{xy}\mathcal{N}_\beta^2 &= 2q\big(\mathsf{d}_q q - (\omega_H + \sqrt{s}\mathfrak{T})\widehat{C}_\beta\big) \\ &\quad + 2\widehat{C}_\beta\big((\omega_H - \sqrt{s}\mathfrak{T})q + \mathsf{d}_{\widehat{C}}\widehat{C}_\beta\big).\end{aligned} \tag{F.18}$$

The two H contributions are $-2\omega_H q\widehat{C}_\beta$ and $+2\omega_H q\widehat{C}_\beta$, hence cancel exactly. The two signed-shear contributions add, giving $-4\sqrt{s}\mathfrak{T}q\widehat{C}_\beta$, and the diagonal terms give the two squares in the displayed formula.

Thus H is invisible to the instantaneous Euclidean amplitude derivative in the orthogonalized variables, although it remains fully capable of rotating the vector $(q, \widehat{C}_\beta)$ and therefore of changing the sign of q. The theorem is a skew-cancellation statement, not a claim that the H field is dynamically irrelevant. □

Audit status: GREEN/COMPLETE.

F4. Passive-circulation interpretation. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 16.5.

Restatement of proposition 16.5. At $x, y > 0$,

$$\boxed{q = \frac{2\varepsilon_\beta\Gamma}{x^2}\Gamma_y, \qquad \widehat{C}_\beta = \frac{2\varepsilon_\beta\Gamma}{x^2\sqrt{s}}\Gamma_x,}$$

and for the material flow of $\mathbf{b} = (-2xv, 2yg)$,

$$\boxed{\nabla\Gamma(\mathscr{X}_{t,0}X_0, t) = D\mathscr{X}_{t,0}(X_0)^{-T}\nabla\Gamma_0(X_0).}$$

Proof. Since $\Gamma = xu$ and $\Theta = \varepsilon_\beta u^2$,

$$\Theta = \varepsilon_\beta\frac{\Gamma^2}{x^2}. \tag{F.19}$$

Differentiation in y gives

$$q = \Theta_y = \frac{2\varepsilon_\beta\Gamma}{x^2}\Gamma_y. \tag{F.20}$$

Differentiation in x gives

$$\Theta_x = \frac{2\varepsilon_\beta\Gamma}{x^2}\Gamma_x - \frac{2\varepsilon_\beta\Gamma^2}{x^3},$$

so the cancellation built into the radial combination is exact:

$$\Theta + \frac{x}{2}\Theta_x = \frac{\varepsilon_\beta \Gamma}{x}\Gamma_x. \tag{F.21}$$

Consequently

$$C_\beta = \frac{\varepsilon_\beta \Gamma}{xy}\Gamma_x, \qquad \widehat{C}_\beta = 2\sqrt{s}\, C_\beta = \frac{2\varepsilon_\beta \Gamma}{x^2\sqrt{s}}\Gamma_x,$$

which proves (16.7).

Now let $\mathscr{X}_{t,0}$ solve

$$\frac{d}{dt}\mathscr{X}_{t,0}(X_0) = \mathbf{b}(\mathscr{X}_{t,0}(X_0), t), \qquad \mathscr{X}_{0,0}(X_0) = X_0.$$

The exact circulation law $\mathcal{D}_{xy}\Gamma = 0$ gives

$$\Gamma(\mathscr{X}_{t,0}X_0, t) = \Gamma_0(X_0). \tag{F.22}$$

Let $\mathbb{F}(t, X_0) := D\mathscr{X}_{t,0}(X_0)$. On a classical interval,

$$\frac{d}{dt}\mathbb{F} = (D\mathbf{b})(\mathscr{X}_{t,0}X_0, t)\mathbb{F}, \qquad \mathbb{F}(0, X_0) = I_2, \tag{F.23}$$

and differentiation of $\mathcal{D}_{xy}\Gamma = 0$ gives, along the same trajectory,

$$\frac{d}{dt}\nabla\Gamma = -(D\mathbf{b})^T\nabla\Gamma. \tag{F.24}$$

Therefore

$$\frac{d}{dt}\big(\mathbb{F}^T\nabla\Gamma(\mathscr{X}_{t,0}X_0, t)\big) = 0,$$

so

$$\nabla\Gamma(\mathscr{X}_{t,0}X_0, t) = \mathbb{F}(t, X_0)^{-T}\nabla\Gamma_0(X_0). \tag{F.25}$$

The classical flow is a local diffeomorphism, hence $\mathbb{F}$ is invertible. Thus $\nabla\Gamma_0(X_0) = 0$ if and only if the transported gradient vanishes at that label.

The two edge cases now make the ancestry statement exact. If $\Gamma_0(X_0) = 0$, then (F.22) and (F.20)–(F.21) give $q = \widehat{C}_\beta = 0$ all along that trajectory. If $\Gamma_0(X_0) \neq 0$, its sign is fixed by (F.22); hence a change from positive to negative q can occur only when the transported component Γ_y changes orientation. By (F.25), that component belongs to the inverse-transpose deformation of the one fixed initial vector $\nabla\Gamma_0(X_0)$. No later source event supplies an independent new circulation-gradient direction. This proves the claimed passive-circulation interpretation. □

Audit status: GREEN/COMPLETE.

F5. Exact apex mixed-shear Hodge moment. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 16.6.

Restatement of proposition 16.6. Put $s := y/x$ and define

$$\tau_0(t) := g_x(0, 0, t) - v_y(0, 0, t).$$

Then

$$\boxed{\tau_0(t) = \frac{15}{8}\int_{\mathcal{Q}} \frac{x\sqrt{y}(3x - 4y)}{(x + y)^{9/2}} H(x, y, t)dX,}$$

or equivalently

$$\boxed{\tau_0(t) = \frac{15}{8}\int_{\mathcal{Q}} K_H(x, y)\frac{3 - 4s}{x(1 + s)^2} H(x, y, t)dX.}$$

Full rigorous proof. By A5 there is a smooth seven-dimensional radial lift

$$\widetilde{\chi}(\mathbf{X},\mathbf{Z},t)=\chi(|\mathbf{X}|^2,|\mathbf{Z}|^2,t),\qquad -\Delta_{\mathbb{R}^7}\widetilde{\chi}=\widetilde{H},$$

with Newton representation

$$\widetilde{\chi}(\boldsymbol{\zeta},t)=\frac{3}{16\pi^3}\int_{\mathbb{R}^7}\frac{\widetilde{H}(\boldsymbol{\eta},t)}{|\boldsymbol{\zeta}-\boldsymbol{\eta}|^5}\,d\boldsymbol{\eta}. \tag{F.26}$$

The fixed smooth axisymmetric symmetry gives $H(0,0,t)=0$. Hence $\widetilde{H}(\boldsymbol{\eta},t)=O(|\boldsymbol{\eta}|^2)$ near the origin, while it is rapidly decaying at infinity. Thus the second derivative of (F.26) at the origin is justified, for example by cutting out a ball of radius ϵ_{cut}, differentiating there, and passing to the limit by dominated convergence; the local radial integrand is $O(|\boldsymbol{\eta}|)$.

The lift formulas are

$$v=\chi+2y\chi_y,\qquad g=2\chi+2x\chi_x. \tag{F.27}$$

At $(0,0)$ the elliptic equation and $H(0,0)=0$ give

$$8\chi_x(0,0)+6\chi_y(0,0)=0. \tag{F.28}$$

Therefore

$$v_y(0,0)=3\chi_y(0,0),\qquad g_x(0,0)=4\chi_x(0,0)=-3\chi_y(0,0),$$

so

$$\tau_0=-6\chi_y(0,0). \tag{F.29}$$

Fix one of the three axial coordinates, say Z_1. At the origin

$$\partial_{Z_1Z_1}|\boldsymbol{\zeta}-\boldsymbol{\eta}|^{-5}\big|_{\boldsymbol{\zeta}=0}=-5|\boldsymbol{\eta}|^{-7}+35\eta_{Z_1}^2|\boldsymbol{\eta}|^{-9}. \tag{F.30}$$

Spherical averaging in the three-dimensional $\mathbf{Z}$ block replaces $\eta_{Z_1}^2$ by $y/3$. The radial measure in $\mathbb{R}^4_{\mathbf{X}}\times\mathbb{R}^3_{\mathbf{Z}}$ is

$$|S^3|\frac{x\,dx}{2}\,|S^2|\frac{\sqrt{y}\,dy}{2}=2\pi^3x\sqrt{y}\,dx\,dy.$$

Since $\partial_{Z_1Z_1}\widetilde{\chi}(0)=2\chi_y(0,0)$, combining this measure with the Newton constant $3/(16\pi^3)$ yields

$$\begin{aligned}2\chi_y(0,0)&=\frac{3}{8}\int_{\mathcal{Q}}x\sqrt{y}\left[-\frac{5}{(x+y)^{7/2}}+\frac{35y}{3(x+y)^{9/2}}\right]H\,dX\\&=\frac{5}{8}\int_{\mathcal{Q}}\frac{x\sqrt{y}(4y-3x)}{(x+y)^{9/2}}H\,dX.\end{aligned} \tag{F.31}$$

Thus

$$\chi_y(0,0)=\frac{5}{16}\int_{\mathcal{Q}}\frac{x\sqrt{y}(4y-3x)}{(x+y)^{9/2}}H\,dX.$$

Multiplying by -6 and using (F.29) proves (16.10). Finally

$$\frac{x\sqrt{y}(3x-4y)}{(x+y)^{9/2}}=K_H(x,y)\frac{3-4s}{x(1+s)^2},$$

which proves (16.11). The coefficient τ_0 is therefore a signed whole-space Hodge moment and is not a freely prescribable local shear. □

Audit status: GREEN/COMPLETE.

F6. Hodge-aware adverse exchange. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 16.7.

Restatement of proposition 16.7. With $s := y/x$ and $C_\beta = (\Theta + \frac{x}{2}\Theta_x)/y$,

$$[-(4yv_y)C_\beta]_+ = [(yH + 2s\mathfrak{T})C_\beta]_+,$$

and the signed form satisfies

$$(yH + 2s\mathfrak{T})C_\beta = \left(\frac{H}{2x} + \frac{\mathfrak{T}}{x^2}\right)\partial_x(x^2\Theta).$$

Full rigorous proof. We derive the mixed-jet identity directly from the audited seven-dimensional lift, so no historical first-jet formula is being imported. From (F.27),

$$v_y = 3\chi_y + 2y\chi_{yy}, \qquad g_x = 4\chi_x + 2x\chi_{xx}. \tag{F.32}$$

Also

$$H = -4x\chi_{xx} - 8\chi_x - 4y\chi_{yy} - 6\chi_y, \qquad \mathfrak{T} = x(g_x - v_y).$$

Consequently

$$\begin{aligned} -yH - 2s\mathfrak{T} &= y\big(4x\chi_{xx} + 8\chi_x + 4y\chi_{yy} + 6\chi_y\big) \\ &\quad - 2y\big(4\chi_x + 2x\chi_{xx} - 3\chi_y - 2y\chi_{yy}\big) \\ &= 4y\big(3\chi_y + 2y\chi_{yy}\big) = 4yv_y. \end{aligned}$$

Thus

$$\boxed{4yv_y = -yH - 2s\mathfrak{T}.} \tag{F.33}$$

Multiplying by $-C_\beta$ and taking positive parts proves (16.13).

For the signed identity, the definition of C_β gives

$$\partial_x(x^2\Theta) = 2x\Theta + x^2\Theta_x = 2xyC_\beta. \tag{F.34}$$

Hence

$$(yH + 2s\mathfrak{T})C_\beta = \left(\frac{H}{2x} + \frac{\mathfrak{T}}{x^2}\right)\partial_x(x^2\Theta),$$

which is (16.14).

For completeness, rapid decay and regularity make the radial boundary terms vanish, so the signed Hodge pairing obeys

$$\frac{3}{8}\int_{\mathcal{Q}} K_H(yH + 2s\mathfrak{T})C_\beta\, dX = -\frac{3}{8}\int_{\mathcal{Q}} x^2\Theta\,\partial_x\left[K_H\left(\frac{H}{2x} + \frac{\mathfrak{T}}{x^2}\right)\right] dX. \tag{F.35}$$

This last formula is deliberately signed. Since $[\xi]_+ = (|\xi| + \xi)/2$, it gives no control of the total positive variation by itself; F7 supplies the positive Hodge-visible estimate instead. □

Audit status: GREEN/COMPLETE.

F7. Angular-tail Hodge suppression of the rotor. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 16.8.

Restatement of proposition 16.8. With $e_- = [-(4yv_y)C_\beta]_+$,

$$K_H e_- \le \psi_{\mathrm{rot}}(s)(|H| + 2|\mathcal{T}_{\mathrm{mix}}|)|C_\beta|, \qquad \psi_{\mathrm{rot}}(s) = \frac{s^{3/2}}{(1+s)^{5/2}},$$

and the carrier-tail estimate (16.20) holds with $w_{\mathrm{rot}}(s) = s^{4/3}(1+s)^{-5/2}$.

Full rigorous proof. By F6,

$$e_- = [(yH + 2s\mathfrak{T})C_\beta]_+ \le (y|H| + 2s|\mathfrak{T}|)|C_\beta|. \tag{F.36}$$

Since

$$K_H y = \frac{x\sqrt{y}\,y}{(x+y)^{5/2}} = \frac{s^{3/2}}{(1+s)^{5/2}} = \psi_{\rm rot}(s)$$

and $s|\mathfrak{T}| = y|\mathcal{T}_{\rm mix}|$, multiplication of (F.36) by K_H proves (16.15). The two asymptotics in (16.17) are immediate.

We now prove the quantitative tail statement. Recall

$$d\mu_c = \frac{dX}{\mathfrak{r}}, \qquad \mathfrak{r} = x^{2/3}y^{1/3} = xs^{1/3}, \qquad M_H := \sqrt{xy}\,H = x\sqrt{s}\,H.$$

Then exactly

$$\int_{\mathcal{Q}} \psi_{\rm rot}(s)|H||C_\beta|\,dX = \int_{\mathcal{Q}} w_{\rm rot}(s)|M_H||C_\beta|\,d\mu_c. \tag{F.37}$$

Thus on $\mathcal{A}_{\delta,S}$,

$$\int_{\mathcal{A}_{\delta,S}} \psi_{\rm rot}|H||C_\beta|\,dX \le \omega_{\delta,S}\,\mathcal{E}_{H,1}\mathcal{E}^{(0,0)}_{\rm sg}. \tag{F.38}$$

For the mixed shear, note

$$\sqrt{xy}\,\mathcal{T}_{\rm mix} = \sqrt{s}\,\mathfrak{T}.$$

The lifted velocity-gradient norm controls this quantity: indeed the cross blocks of $\nabla_{\mathbb{R}^7}\mathbf{B}$ contain $2\sqrt{xy}\,v_y$ and $2\sqrt{xy}\,g_x$, hence

$$\|\sqrt{xy}\,\mathcal{T}_{\rm mix}\|_\infty \le \mathfrak{c}_B\mathcal{L}_B. \tag{F.39}$$

Using again $dX = xs^{1/3}d\mu_c$ gives

$$\begin{aligned}\int_{\mathcal{A}_{\delta,S}} 2\psi_{\rm rot}|\mathcal{T}_{\rm mix}||C_\beta|\,dX &= 2\int_{\mathcal{A}_{\delta,S}} w_{\rm rot}(s)|\sqrt{xy}\,\mathcal{T}_{\rm mix}|\,|C_\beta|\,d\mu_c \\ &\le \mathfrak{c}_B\omega_{\delta,S}\,\mathcal{L}_B\,\mathcal{Q}^{(1)}_{\rm sg}.\end{aligned} \tag{F.40}$$

Combining (F.38) and (F.40) proves (16.20). Finally $w_{\rm rot}(s) \sim s^{4/3}$ as $s \downarrow 0$ and $w_{\rm rot}(s) \sim s^{-7/6}$ as $s \to \infty$, so $\omega_{\delta,S} \to 0$. This is a visibility-weighted tail estimate, not a support assertion: source-gradient mass may remain in the angular tails, but its Hodge-visible rotor contribution is uniformly suppressed by the explicit weight. □

Audit status: GREEN/COMPLETE.

F8. Compact mesoscopic rotation ticket. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 16.9.

> **Restatement of theorem 16.9.** At a first near-lock time t_* above the certified old-response margin, and on the admitted finite-action/rapid-decay branch, a fixed positive part of the response-weighted generated-negative source lies in one compact box $\mathcal{M} = \{\ell \le x \le x_{\rm far},\ \delta \le s \le S\}$; hence
>
> $$\int_0^{t_*}\int_{\mathcal{M}} [(\omega_H + \sqrt{s}\mathfrak{T})\widehat{C}_\beta]_+\,dX\,dt \ge c_{\rm rot} > 0.$$

Full rigorous proof. We prove the localization directly from the already audited equations. In particular, we rederive the two elementary routing bounds needed below rather than invoke H1–H2 long-proof slots as black boxes.

Step 1: the generated-negative response has an order-one lower bound. Let

$$\mathfrak{m}_* := \upsilon_* - \upsilon_{\mathrm{old}} > 0.$$

The E7 Kato reconstruction gives

$$a^-(t_*) \le \mathfrak{X}^H_{-,\mathrm{old}}(t_*) + \mathfrak{X}^H_-(t_*), \tag{F.41}$$

where

$$\mathfrak{X}^H_-(t_*) = \frac{3}{4}\int_0^{t_*}\int_{\mathcal{Q}} e_-(X,\tau)\mathcal{V}_H(t_*;\tau,X)\mathfrak{R}_q(t_*,\tau;X)\,dX\,d\tau. \tag{F.42}$$

Using the near-lock inequality and (16.21) in (F.41) yields

$$\boxed{\mathfrak{X}^H_-(t_*) \ge \mathfrak{m}_* a^+(t_*) > 0.} \tag{F.43}$$

No tube subtraction has yet been made, so the pre-t_1 issue from E7 does not arise.

Step 2: rederive finite visibility and residence constants on $[0,t_*]$. From $K_H = x\sqrt{y}(x+y)^{-5/2}$ and $\dot{x} = -2xv$, $\dot{y} = 2yg$,

$$\mathcal{D}_{xy}\log K_H = \frac{(3-2s)v + (1-4s)g}{1+s}. \tag{F.44}$$

The two angular coefficients in (F.44) are bounded on $(0,\infty)$. Since t_* is a classical time, put

$$\mathcal{K}^*_{\mathrm{vis}} := \int_0^{t_*} (\|v(\tau)\|_\infty + \|g(\tau)\|_\infty)\,d\tau < \infty.$$

Then for all $0 \le \tau \le t_*$,

$$\mathcal{V}_H(t_*;\tau,X) \le e^{\mathfrak{c}_{\mathrm{vis}}\mathcal{K}^*_{\mathrm{vis}}} K_H(X). \tag{F.45}$$

Likewise the exact residence formula gives, with

$$\mathcal{K}^*_{vx} := \int_0^{t_*} \|v - xv_x\|_\infty\,d\tau < \infty,$$

$$0 \le \mathfrak{R}_q(t_*,\tau;X) \le t_* e^{2\mathcal{K}^*_{vx}} =: \mathfrak{R}_*. \tag{F.46}$$

If either action is not finite on the proposed branch, the corresponding named first-jet/continuation exit has already occurred.

Step 3: retire the two angular tails. Let $\mathcal{A}_{\delta,S}$ be as in F7. By (F.45)–(F.46) and F7,

$$\begin{aligned}\mathfrak{X}^H_{-,\mathrm{ang}}(t_*) &:= \frac{3}{4}\int_0^{t_*}\int_{\mathcal{A}_{\delta,S}} e_-\mathcal{V}_H\mathfrak{R}_q\,dX\,d\tau \\ &\le \mathfrak{c}_*\omega_{\delta,S}\int_0^{t_*}\left[\mathcal{E}_{H,1}\mathcal{E}^{(0,0)}_{\mathrm{sg}} + \mathcal{L}_B\mathcal{Q}^{(1)}_{\mathrm{sg}}\right](\tau)d\tau,\end{aligned} \tag{F.47}$$

where $\mathfrak{c}_* < \infty$ depends only on the finite routing actions above. The time integral is finite on the audited D3–D4 branch. Because $a^+(t_*) > 0$ and $\omega_{\delta,S} \to 0$, choose once $0 < \delta < 1 < S < \infty$ so that

$$\mathfrak{X}^H_{-,\mathrm{ang}}(t_*) \le \frac{1}{4}\mathfrak{m}_* a^+(t_*). \tag{F.48}$$

Step 4: a compact-time protected apex tube valid from $t = 0$. The detector equation gives

$$p(t) := u_y(0,0,t) = p(0)\exp\left(-2\int_0^t a(\xi)d\xi\right) > 0.$$

Hence $p_* := \min_{[0,t_*]} p(t) > 0$. The equator is material and $u_0(x,0) = 0$, so passive transport of $\Gamma = xu$ gives $u(x,0,t) = 0$. Smoothness therefore gives $u(x,y,t) = y\,w(x,y,t)$ on the compact time interval, with $w(0,0,t) = p(t)$. Uniform Taylor expansion on the fixed angular sector $[\delta, S]$ yields, for $y = sx$,

$$q(x,sx,t) = 2\varepsilon_\beta p(t)^2 sx + O_{\delta,S,t_*}(x^2), \qquad C_\beta(x,sx,t) = O_{\delta,S,t_*}(x). \tag{F.49}$$

Moreover $H(0,0,t) = 0$ and $\mathfrak{T} = x(g_x - v_y)$, so

$$H(x,sx,t) = O_{\delta,S,t_*}(x), \qquad \mathfrak{T}(x,sx,t) = O_{\delta,S,t_*}(x). \tag{F.50}$$

Choose $\ell > 0$ so small that throughout

$$\mathcal{U}_{\ell,\delta,S} := \{0 < x < \ell,\ \delta \le s \le S\}, \qquad 0 \le t \le t_*,$$

one has $q > 0$ and

$$e_- \le \mathfrak{c}_{\mathrm{ap}}\ell\, q. \tag{F.51}$$

The positive response formula for a^+ then gives

$$\begin{aligned}\mathfrak{X}^H_{-,\mathrm{apex}}(t_*) &:= \frac{3}{4}\int_0^{t_*}\int_{\mathcal{U}_{\ell,\delta,S}} e_-\mathcal{V}_H\mathfrak{R}_q\,dX\,d\tau \\ &\le \mathfrak{c}_{\mathrm{ap}}\ell\mathfrak{R}_* a^+(t_*).\end{aligned} \tag{F.52}$$

Decrease ℓ once so that

$$\mathfrak{X}^H_{-,\mathrm{apex}}(t_*) \le \frac{1}{4}\mathfrak{m}_* a^+(t_*). \tag{F.53}$$

This compact-time construction covers the entire source interval $[0, t_*]$ and hence does not leave an untreated pre-t_1 deposition account.

Step 5: retire the far radial tail on the remaining angular sector. We make the rapid-decay input explicit. The fixed initial circulation $\Gamma_0 = xu_0$ and its first derivatives decay faster than every power of $x + y$. On $[0, t_*]$ the admitted value action gives, along each material trajectory,

$$x(\tau) = x_0\exp\left(-2\int_0^\tau v\right), \qquad y(\tau) = y_0\exp\left(2\int_0^\tau g\right),$$

so $(x(\tau) + y(\tau))$ is uniformly comparable to $(x_0 + y_0)$. The admitted first-jet action gives a uniform bound for the deformation matrix and its inverse. Using the passive identity $\Gamma(\mathscr{X}_{\tau,0}X_0, \tau) = \Gamma_0(X_0)$ and the F4 inverse-transpose formula therefore yields, for every $N > 0$,

$$\sup_{0\le\tau\le t_*}\sup_{\delta\le s\le S}(1 + x + y)^N\big(|\Gamma| + |\Gamma_x| + |\Gamma_y|\big) < \infty. \tag{F.54}$$

Since $\widehat{C}_\beta = 2\varepsilon_\beta\Gamma\Gamma_x/(x^2\sqrt{s})$, the field C_β has the same uniform rapid-decay property on $\delta \le s \le S$. Meanwhile the admitted first-jet norms bound

$$H = -2(v_y + g_x), \qquad \mathcal{T}_{\mathrm{mix}} = g_x - v_y$$

on the same compact time interval. Because $K_H y = \psi_{\mathrm{rot}}(s)$ and the central angular sector keeps ψ_{rot} bounded, F6 gives

$$K_H e_- \le \mathfrak{c}_{\delta,S,t_*}|C_\beta| \qquad (\delta \le s \le S).$$

The rapid decay of C_β is stronger than the $O(x)$ width of the strip $\{\delta x \le y \le Sx\}$, hence

$$\lim_{x_{\mathrm{far}}\to\infty} \sup_{0\le\tau\le t_*} \int_{\substack{x>x_{\mathrm{far}}\\ \delta\le s\le S}} K_H(X) e_-(X,\tau)\, dX = 0. \tag{F.55}$$

Together with (F.45)–(F.46), choose $x_{\mathrm{far}} > \ell$ so large that the corresponding two-time response obeys

$$\mathfrak{X}^H_{-,\mathrm{far}}(t_*) \le \frac{1}{4}\mathfrak{m}_* a^+(t_*). \tag{F.56}$$

If the value/first-jet/rapid-decay controls used above fail, the branch has already entered the named continuation exit rather than producing an unpriced far-field bill.

Step 6: the compact mesoscopic response survives. The complement of the angular tails, apex tube, and far radial tail is exactly

$$\mathcal{M} = \{\ell \le x \le x_{\mathrm{far}},\ \delta \le s \le S\}.$$

Subtracting (F.48), (F.53), and (F.56) from (F.43) gives

$$\boxed{\frac{3}{4}\int_0^{t_*}\int_{\mathcal{M}} e_-\mathcal{V}_H(t_*;\tau,X)\mathfrak{R}_q(t_*,\tau;X)\, dX\, d\tau \ge \frac{1}{4}\mathfrak{m}_* a^+(t_*) > 0.} \tag{F.57}$$

Step 7: remove only the bounded positive routing factor. On $\mathcal{M}\times[0,t_*]$, the deposition coordinates stay a positive distance from both axes and in a bounded radial set. Equation (F.45) and (F.46) therefore give a finite constant

$$\mathfrak{M}_{\mathrm{rout}} := \sup_{\substack{0\le\tau\le t_*\\ X\in\mathcal{M}}} \mathcal{V}_H(t_*;\tau,X)\mathfrak{R}_q(t_*,\tau;X) < \infty. \tag{F.58}$$

The rotor normalization is exact:

$$\begin{aligned}(\omega_H + \sqrt{s}\mathfrak{T})\widehat{C}_\beta &= \left(\frac{1}{2}\sqrt{xy}H + \sqrt{s}\mathfrak{T}\right)(2\sqrt{s}C_\beta)\\ &= (yH + 2s\mathfrak{T})C_\beta,\end{aligned} \tag{F.59}$$

so its positive part is exactly e_-. From (F.57) and (F.58),

$$\int_0^{t_*}\int_{\mathcal{M}} [(\omega_H + \sqrt{s}\mathfrak{T})\widehat{C}_\beta]_+\, dX\, d\tau \ge \frac{\mathfrak{m}_* a^+(t_*)}{3\mathfrak{M}_{\mathrm{rout}}} =: c_{\mathrm{rot}} > 0.$$

This is (16.23).

Finally, (F.42) integrates each physical deposition pair (X,τ) exactly once. The resulting ticket is attached to the first near-lock conversion and cannot be charged again merely because the already produced Hodge genealogy is later refreshed. This is the same no-double-counting firewall used in E7. □

Audit status: GREEN/COMPLETE.

Appendix G. Circulation-level-set polarization and deformation stability

This section contains the rigorous G1–G5 level-set, fixed-seed condition-number, and nonlinear polarization-stability audit.

G1. Circulation level-set response. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 17.1.

Restatement of theorem 17.1. Fix $0 \le \tau \le t$. For the fixed customer put $\Gamma = xu$, define

$$\Omega_c(\tau) = \{\Gamma(\cdot,\tau) > c\}, \qquad W_{t,\tau} = K_H \circ \mathscr{X}_{t,\tau},$$
$$\mathcal{S}_\pm(t,\tau) = \frac{3}{4}\int_{\mathcal{Q}} W_{t,\tau}[q(\cdot,\tau)]_\pm \, dX, \qquad \mathcal{R}(t,\tau;X) = -\partial_{y_\tau} W_{t,\tau}(X).$$

For a regular value $c > 0$ let $n_c = \nabla\Gamma/|\nabla\Gamma|$ point toward increasing circulation. Then

$$\boxed{\mathcal{S}_\pm(t,\tau) = \frac{3\varepsilon_\beta}{2}\int_0^\infty c \oint_{\partial\Omega_c(\tau)} \frac{W_{t,\tau}}{x^2}[\pm n_{c,y}]_+ \, d\ell \, dc,}$$

and

$$\boxed{\mathcal{S}_+(t,\tau) - \mathcal{S}_-(t,\tau) = \frac{3\varepsilon_\beta}{2}\int_0^\infty c \int_{\Omega_c(\tau)} \frac{\mathcal{R}(t,\tau;X)}{x^2}\, dX \, dc.}$$

Critical levels are interpreted in the coarea sense.

Proof. **Step 1: sign, transport, and admissible superlevels.** For the fixed customer,

$$u_0 = y w_0 \ge 0, \qquad \Gamma_0 = x u_0 \ge 0.$$

By (5.6), $\mathcal{D}_{xy}\Gamma = 0$, hence

$$\boxed{\Gamma(\cdot,\tau) \ge 0.} \tag{G.1}$$

Moreover $\Gamma = 0$ on the coordinate axes, while the standing rapid-decay class (and, on every finite continuation interval, the audited D6 propagation) gives $\Gamma(X,\tau) \to 0$ as $|X| \to \infty$. Thus for every $c > 0$ the superlevel $\Omega_c(\tau)$ is an interior relatively compact set up to the harmless possibility of several connected components; no axis or infinity boundary is created in the level-set divergence calculation.

The tagged-source identity is exact:

$$q = \partial_y(\varepsilon_\beta u^2) = \frac{2\varepsilon_\beta \Gamma \Gamma_y}{x^2}. \tag{G.2}$$

At a regular point of the level $\{\Gamma = c\}$,

$$\Gamma_y = |\nabla\Gamma|\, n_{c,y}. \tag{G.3}$$

Step 2: coarea gives the positive and negative responses. Because $\Gamma \ge 0$, equations (G.2)–(G.3) and the coarea formula yield

$$\int_{\mathcal{Q}} W_{t,\tau}[q]_+ \, dX = 2\varepsilon_\beta \int_0^\infty c \oint_{\partial\Omega_c(\tau)} \frac{W_{t,\tau}}{x^2}[n_{c,y}]_+ \, d\ell \, dc, \tag{G.4}$$

$$\int_{\mathcal{Q}} W_{t,\tau}[q]_- \, dX = 2\varepsilon_\beta \int_0^\infty c \oint_{\partial\Omega_c(\tau)} \frac{W_{t,\tau}}{x^2}[-n_{c,y}]_+ \, d\ell \, dc. \tag{G.5}$$

Multiplication by 3/4 gives exactly (17.3)–(17.4). The factor ε_β has not been divided out; when the tagged source is switched off, both sides vanish.

Step 3: the signed boundary flux is the vertical future-visibility response. Subtracting (G.5) from (G.4) uses $[z]_+ - [-z]_+ = z$ and gives the boundary flux

$$\oint_{\partial\Omega_c(\tau)} \frac{W_{t,\tau}}{x^2} n_{c,y} \, d\ell.$$

The outward unit normal of the superlevel $\{\Gamma > c\}$ is $-n_c$. Therefore the divergence theorem gives

$$\oint_{\partial\Omega_c(\tau)} \frac{W_{t,\tau}}{x^2} n_{c,y}\, d\ell = -\int_{\Omega_c(\tau)} \partial_y\left(\frac{W_{t,\tau}}{x^2}\right) dX \tag{G.6}$$

$$= -\int_{\Omega_c(\tau)} \frac{\partial_y W_{t,\tau}}{x^2}\, dX = \int_{\Omega_c(\tau)} \frac{\mathcal{R}(t,\tau;X)}{x^2}\, dX. \tag{G.7}$$

Here x^{-2} is independent of y, and the last equality is precisely the deposition-coordinate derivative in (17.2). This proves (17.5).

Step 4: critical levels and integrability. Sard's theorem makes the critical values negligible for the coarea integral. Equivalently, one may approximate a critical level by regular values and pass to the limit. The fixed customer's rapid decay and the positivity/regularity of $W_{t,\tau}$ on a finite classical branch provide the needed integrable majorants. Thus no smooth parametrization of critical level curves is being assumed. □

Audit status: GREEN/COMPLETE.

G2. Raw level-set compensation criterion. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 17.2.

> **Restatement of proposition 17.2.** At a fixed time t, let $0 < \pi_* < 1$. The raw polarization inequality
>
> $$\boxed{a(t) \geq \pi_*[a^+(t) + a^-(t)]}$$
>
> is equivalent to
>
> $$\boxed{a^-(t) \leq \frac{1-\pi_*}{1+\pi_*} a^+(t).}$$
>
> In the circulation-level-set representation the common factor $3\varepsilon_\beta/2$ is retained on both sides until physical admission.

Proof. By the exact two-time visibility formula (15.2) and the definition (17.2),

$$a^{\pm}(t) = \int_0^t \mathcal{S}_{\pm}(t,\tau)\, d\tau. \tag{G.8}$$

Hence G1 gives the exact signed and absolute level-set stocks

$$a(t) = a^+(t) - a^-(t) = \frac{3\varepsilon_\beta}{2} \int_0^t \int_0^\infty c \int_{\Omega_c(\tau)} \frac{\mathcal{R}(t,\tau;X)}{x^2}\, dX\, dc\, d\tau, \tag{G.9}$$

$$a^+(t) + a^-(t) = \frac{3\varepsilon_\beta}{2} \int_0^t \int_0^\infty c \oint_{\partial\Omega_c(\tau)} \frac{W_{t,\tau}}{x^2} |n_{c,y}|\, d\ell\, dc\, d\tau. \tag{G.10}$$

Thus the raw compensation hypothesis is exactly the level-set inequality

$$\boxed{\begin{aligned} &\frac{3\varepsilon_\beta}{2} \int_0^t \int_0^\infty c \int_{\Omega_c(\tau)} \frac{\mathcal{R}(t,\tau;X)}{x^2}\, dX\, dc\, d\tau \\ &\qquad \geq \pi_* \frac{3\varepsilon_\beta}{2} \int_0^t \int_0^\infty c \oint_{\partial\Omega_c(\tau)} \frac{W_{t,\tau}}{x^2} |n_{c,y}|\, d\ell\, dc\, d\tau. \end{aligned}} \tag{G.11}$$

The ancestry factor $3\varepsilon_\beta/2$ is displayed on both sides and is not divided out. In particular, the shadow sector $\varepsilon_\beta = 0$ carries no tagged payment.

It remains only to check the algebraic form of the gap. Since $a = a^+ - a^-$,

$$\begin{aligned} a^+ - a^- \geq \pi_*(a^+ + a^-) &\iff (1-\pi_*)a^+ \geq (1+\pi_*)a^- \\ &\iff a^- \leq \frac{1-\pi_*}{1+\pi_*}a^+, \end{aligned}$$

where $1 + \pi_* > 0$. Conversely the last inequality reverses the same steps. Therefore a fixed positive raw level-set polarization is exactly the desired strict cancellation gap; it is not an additional normalized source account. □

Audit status: GREEN/COMPLETE.

G3. Uniform fixed-seed source-gradient condition number. *Mutual pointers:* grand-finale entry / short proof/status / main statement lemma 17.4.

Restatement of lemma 17.4. In the physical sector let

$$\mathscr{X}_\Lambda^{\mathrm{aff}}(x,y) = (e^{-2\Lambda}x, e^{4\Lambda}y), \qquad Z_{\beta,0} = (q_0, C_{\beta,0})^T,$$

and use the Euclidean norm $|Z_{\beta,0}|$. Then

$$\mathfrak{C}_Z(\Lambda) = \frac{\int_{\mathcal{Q}} K_H(\mathscr{X}_\Lambda^{\mathrm{aff}}X)|Z_{\beta,0}(X)|\,dX}{\int_{\mathcal{Q}} K_H(\mathscr{X}_\Lambda^{\mathrm{aff}}X)|q_0(X)|\,dX}$$

has positive denominator and satisfies

$$\boxed{\sup_{\Lambda\geq 0} \mathfrak{C}_Z(\Lambda) < \infty.}$$

Proof. **Step 1: compact affine ages.** The fixed seed is a finite sum of polynomial–exponential terms. Hence q_0, $C_{\beta,0}$, and every quantity below have rapid exponential decay. On any compact interval $0 \leq \Lambda \leq L$, the two defining integrals for $\mathfrak{C}_Z$ depend continuously on Λ by dominated convergence. The denominator is strictly positive: from $u_0 = yw_0$ and $w_0(x,0) > 0$,

$$q_0(x,y) = 2y\,w_0(x,0)^2 + O(y^2) \tag{G.12}$$

with a rapid x-majorant, so q_0 is not identically zero. Since $K_H > 0$ in the interior, the denominator is positive for every finite Λ.

It remains to audit $\Lambda \to \infty$. This is the point at which the immutable An earlier proof used an over-strong dominated-convergence shortcut. The relevant angular factor is logarithmically nonintegrable; the common logarithm must be retained and cancelled only after numerator and denominator are expanded separately.

Step 2: line-coordinate reduction. Write $y = \eta x$ and

$$w_*(x) := w_0(x,0) > 0. \tag{G.13}$$

Because $dX = x\,dx\,d\eta$ and $K_H(x,y) = x\sqrt{y}/(x+y)^{5/2}$,

$$K_H(\mathscr{X}_\Lambda^{\mathrm{aff}}(x,\eta x))\,dX = e^{5\Lambda}\frac{\sqrt{\eta}}{(1+e^{6\Lambda}\eta)^{5/2}}\,dx\,d\eta. \tag{G.14}$$

Define the fixed line stocks

$$\mathcal{L}_{|q|}(\eta) := \int_0^\infty |q_0(x,\eta x)|\,dx, \qquad \mathcal{L}_Z(\eta) := \int_0^\infty |Z_{\beta,0}(x,\eta x)|\,dx. \tag{G.15}$$

Then the denominator and numerator of $\mathfrak{C}_Z$ are respectively

$$\mathfrak{d}_Z(\Lambda) = e^{5\Lambda} \int_0^\infty \frac{\sqrt{\eta}}{(1+e^{6\Lambda}\eta)^{5/2}} \mathcal{L}_{|q|}(\eta)\, d\eta, \tag{G.16}$$

$$N_Z(\Lambda) = e^{5\Lambda} \int_0^\infty \frac{\sqrt{\eta}}{(1+e^{6\Lambda}\eta)^{5/2}} \mathcal{L}_Z(\eta)\, d\eta. \tag{G.17}$$

Step 3: exact equatorial coefficients and global line majorants. From the definitions $q_0 = (u_0^2)_y$ and $C_{\beta,0} = (u_0^2 + \frac{x}{2}(u_0^2)_x)/y$, the fixed seed gives, as $\eta \downarrow 0$,

$$q_0(x, \eta x) = 2\eta x\, w_*(x)^2 + O\big(\eta^2 \mathfrak{p}(x) e^{-c_0 x}\big)\,, \tag{G.18}$$

$$C_{\beta,0}(x, \eta x) = \eta x\, w_*(x)\big(w_*(x) + x w_*'(x)\big) + O\big(\eta^2 \mathfrak{p}(x) e^{-c_0 x}\big)\,, \tag{G.19}$$

for one fixed polynomial $\mathfrak{p}$ and $c_0 > 0$. The maps $z \mapsto |z|$ and $Z \mapsto |Z|$ are 1-Lipschitz, so after x-integration

$$\mathcal{L}_{|q|}(\eta) = \kappa_q \eta + O(\eta^2), \qquad \kappa_q := 2\int_0^\infty x w_*(x)^2\, dx > 0, \tag{G.20}$$

$$\mathcal{L}_Z(\eta) = \kappa_Z \eta + O(\eta^2), \qquad \kappa_Z := \int_0^\infty x\sqrt{4w_*^4 + w_*^2(w_* + x w_*')^2}\, dx < \infty. \tag{G.21}$$

In particular $\kappa_Z > 0$ because its first squared component is $4w_*^4$.

The same explicit polynomial–exponential structure gives the global line bound

$$\boxed{\mathcal{L}_{|q|}(\eta) + \mathcal{L}_Z(\eta) \le \mathfrak{c}\frac{\eta}{(1+\eta)^2} \qquad (\eta > 0).} \tag{G.22}$$

Indeed every term obtained from q_0 or $C_{\beta,0}$ on $y = \eta x$ has the form $\mathfrak{c}\eta^m x^n e^{-(c+d\eta)x}$ with $m \ge 1$, $n \ge m$, and $c, d > 0$; after x-integration it is bounded by $\mathfrak{c}\eta^m (c+d\eta)^{-n-1}$. Summing the finitely many terms proves (G.22) (using $|Z| \le |q| + |C_\beta|$).

Step 4: the logarithmic affine-age asymptotic. Set

$$\epsilon := e^{-6\Lambda}, \qquad k(\sigma) := \frac{\sqrt{\sigma}}{(1+\sigma)^{5/2}}. \tag{G.23}$$

The change of variables $\eta = \epsilon\sigma$ in (G.16)–(G.17) gives

$$\mathfrak{d}_Z(\Lambda) = \epsilon^{2/3} \int_0^\infty k(\sigma)\mathcal{L}_{|q|}(\epsilon\sigma)\, d\sigma, \tag{G.24}$$

$$N_Z(\Lambda) = \epsilon^{2/3} \int_0^\infty k(\sigma)\mathcal{L}_Z(\epsilon\sigma)\, d\sigma. \tag{G.25}$$

We use the same elementary endpoint calculation for either line stock. If $\mathcal{L}(\eta) = \kappa\eta + O(\eta^2)$ for $0 < \eta < \eta_0$ and satisfies (G.22), split the σ-integral into $0 < \sigma < 1$, $1 \le \sigma \le \eta_0/\epsilon$, and $\sigma \ge \eta_0/\epsilon$. The first piece is $O(\epsilon)$. On the middle piece,

$$\int_1^{\eta_0/\epsilon} k(\sigma)\mathcal{L}(\epsilon\sigma)\, d\sigma = \kappa\epsilon \int_1^{\eta_0/\epsilon} \sigma k(\sigma)\, d\sigma + O\left(\epsilon^2 \int_1^{\eta_0/\epsilon} \sigma^2 k(\sigma)\, d\sigma\right) \tag{G.26}$$

$$= \kappa\epsilon \log(1/\epsilon) + O(\epsilon), \tag{G.27}$$

because

$$\sigma k(\sigma) = \sigma^{-1} + O(\sigma^{-2}), \qquad \sigma^2 k(\sigma) \le 1 \quad (\sigma \ge 1).$$

On the last interval, $k(\sigma) \le \sigma^{-2}$ and (G.22) give

$$\int_{\eta_0/\epsilon}^{\infty} k(\sigma)\mathcal{L}(\epsilon\sigma)\,d\sigma \le \mathfrak{c}\epsilon \int_{\eta_0}^{\infty} \frac{d\eta}{\eta(1+\eta)^2} = O(\epsilon).$$

Therefore

$$\boxed{\int_0^{\infty} k(\sigma)\mathcal{L}(\epsilon\sigma)\,d\sigma = \kappa\epsilon\log(1/\epsilon) + O(\epsilon).} \tag{G.28}$$

The logarithm is essential: $\sigma k(\sigma) \sim 1/\sigma$, so this is precisely the endpoint at which a global dominated-convergence argument would be invalid.

Applying (G.28) to (G.20) and (G.21), and using $\log(1/\epsilon) = 6\Lambda$, yields

$$\mathfrak{d}_Z(\Lambda) = e^{-10\Lambda}\big(6\kappa_q\Lambda + O(1)\big), \tag{G.29}$$

$$N_Z(\Lambda) = e^{-10\Lambda}\big(6\kappa_Z\Lambda + O(1)\big). \tag{G.30}$$

Consequently

$$\boxed{\lim_{\Lambda\to\infty} \mathfrak{C}_Z(\Lambda) = \frac{\kappa_Z}{\kappa_q} = \frac{\displaystyle\int_0^{\infty} x\sqrt{4w_*^4 + w_*^2(w_* + xw_*')^2}\,dx}{\displaystyle 2\int_0^{\infty} xw_*^2\,dx} < \infty.} \tag{G.31}$$

Combining this finite limit with Step 1 compactness proves $\mathfrak{C}_Z^* < \infty$. □

Audit status: GREEN/COMPLETE.

G4. Stability of the complete Hodge-response polarization. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 17.5.

Restatement of theorem 17.5. Work in the physical sector $\varepsilon_\beta = 1$. Let

$$M := \mathcal{K}_{\mathrm{cent}}([0,t])$$

with $\mathcal{K}_{\mathrm{cent}}$ from definition 17.3, and put

$$\Lambda(t) := \int_0^t a(\tau)\,d\tau, \qquad I_t := \int_0^t e^{2\Lambda(\tau)}\,d\tau,$$

$$W_t^{\mathrm{aff}}(X) := K_H(\mathscr{X}_{\Lambda(t)}^{\mathrm{aff}}X),$$

where $\mathscr{X}_\Lambda^{\mathrm{aff}}$ is the affine map from lemma 17.4. Put $q_0(X) := q(X,0)$ and define

$$\mathcal{P}(t) := a(t), \qquad \mathcal{V}(t) := a^+(t) + a^-(t),$$

$$\mathcal{P}_{\mathrm{aff}}(t) := \frac{3}{4}I_t\int_{\mathcal{Q}} q_0(X)W_t^{\mathrm{aff}}(X)\,dX, \qquad \mathcal{V}_{\mathrm{aff}}(t) := \frac{3}{4}I_t\int_{\mathcal{Q}} |q_0(X)|W_t^{\mathrm{aff}}(X)\,dX.$$

Then for universal constants $\mathfrak{c}_0, \mathfrak{c}_1 > 0$,

$$\boxed{|\mathcal{P} - \mathcal{P}_{\mathrm{aff}}| + |\mathcal{V} - \mathcal{V}_{\mathrm{aff}}| \le \mathfrak{c}_1\mathfrak{C}_Z^*(e^{\mathfrak{c}_0 M} - 1)\mathcal{V}_{\mathrm{aff}}.}$$

Consequently there exist fixed constants $M_* > 0$ and $\pi_{107} > 0$, depending only on the fixed customer and the affine polarization margin, such that

$$\mathcal{K}_{\mathrm{cent}}([0,t]) \le M_*$$

implies

$$\boxed{a(t) \ge \pi_{107}[a^+(t) + a^-(t)],}$$

and hence a uniform strict cancellation gap.

Proof. **Step 1: exact fixed-label pullback of the true response.** Write

$$\mathscr{X}_\tau := \mathscr{X}_{\tau,0}, \qquad \mathcal{J}_\tau(X_0) := \det D\mathscr{X}_\tau(X_0), \qquad \mathcal{W}_t(X_0) := K_H(\mathscr{X}_t X_0).$$

The squared-coordinate material field is $\mathbf{b} = (-2xv, 2yg)$ and $\nabla\cdot\mathbf{b} = g$, so

$$\frac{d}{d\tau}\mathcal{J}_\tau(X_0) = g(\mathscr{X}_\tau X_0,\tau)\mathcal{J}_\tau(X_0), \qquad \mathcal{J}_0 = 1. \tag{G.32}$$

Because the fixed customer has $H_0 = 0$ and $\mathcal{D}_{xy}H + gH = 2q$,

$$\frac{d}{d\tau}\big[\mathcal{J}_\tau H(\mathscr{X}_\tau X_0,\tau)\big] = 2\mathcal{J}_\tau q(\mathscr{X}_\tau X_0,\tau).$$

The same calculation for the non-reset banks gives $\frac{d}{d\tau}(\mathcal{J}_\tau H^{\pm}) = 2\mathcal{J}_\tau[q]_{\pm}$. Changing variables $X = \mathscr{X}_t X_0$ in the Hodge cashier therefore yields the exact identities

$$\mathcal{P}(t) = \frac{3}{4}\int_0^t\int_{\mathcal{Q}} q(\mathscr{X}_\tau X_0,\tau)\mathcal{J}_\tau(X_0)\mathcal{W}_t(X_0)\,dX_0 d\tau, \tag{G.33}$$

$$\mathcal{V}(t) = \frac{3}{4}\int_0^t\int_{\mathcal{Q}} |q(\mathscr{X}_\tau X_0,\tau)|\mathcal{J}_\tau(X_0)\mathcal{W}_t(X_0)\,dX_0 d\tau. \tag{G.34}$$

Thus both the signed and absolute response live on the same fixed initial label space; no moving-boundary argument enters G4.

Step 2: source-gradient comparison. Put

$$Z_\beta := \begin{pmatrix} q \\ C_\beta \end{pmatrix}, \qquad Z_{\beta,0} := Z_\beta(\cdot,0).$$

By the audited exact source-gradient system F1,

$$\frac{d}{d\tau}Z_\beta(\mathscr{X}_\tau X_0,\tau) = \mathbf{A}_{\mathrm{sg}}(\mathscr{X}_\tau X_0,\tau)Z_\beta(\mathscr{X}_\tau X_0,\tau).$$

Since the $\|\mathbf{A}_{\mathrm{sg}}\|_\infty$ action is one summand of M, the fundamental-matrix estimate gives

$$|Z_\beta(\mathscr{X}_\tau X_0,\tau) - Z_{\beta,0}(X_0)| \le (e^M - 1)|Z_{\beta,0}(X_0)|, \qquad 0 \le \tau \le t. \tag{G.35}$$

In particular, the same bound holds for the first component q.

Step 3: material-Jacobian comparison. Equation (G.32) and the definition of Λ give

$$\log\frac{\mathcal{J}_\tau(X_0)}{e^{2\Lambda(\tau)}} = \int_0^\tau [g(\mathscr{X}_\xi X_0,\xi) - 2a(\xi)]\,d\xi.$$

Hence

$$\left|\log\frac{\mathcal{J}_\tau(X_0)}{e^{2\Lambda(\tau)}}\right| \le M. \tag{G.36}$$

No division by a or by Λ is used.

Step 4: final Hodge-visibility comparison. Along a true characteristic write $X(\xi) = (x(\xi), y(\xi))$ and use the protected aspect ratio $s(\xi) := y(\xi)/x(\xi)$. The coordinate axes are null sets for the response integrals, so this interior computation suffices. The exact characteristic equations give

$$\frac{d}{d\xi}\log x = -2v, \qquad \frac{d}{d\xi}\log s = 2(v+g).$$

The same-clock affine endpoint has $x_{\mathrm{aff}}(t) = e^{-2\Lambda(t)}x_0$ and $s_{\mathrm{aff}}(t) = e^{6\Lambda(t)}s_0$. Therefore

$$\left|\log \frac{x(t)}{e^{-2\Lambda(t)}x_0}\right| \le 2M, \tag{G.37}$$

$$\left|\log \frac{s(t)}{e^{6\Lambda(t)}s_0}\right| \le 2M. \tag{G.38}$$

Since

$$K_H = x^{-1}f(s), \qquad f(s) = \frac{\sqrt{s}}{(1+s)^{5/2}}, \qquad \left|\frac{d\log f}{d\log s}\right| = \left|\frac{1-4s}{2(1+s)}\right| \le 2,$$

we obtain

$$\left|\log \frac{\mathcal{W}_t(X_0)}{W_t^{\mathrm{aff}}(X_0)}\right| \le 6M. \tag{G.39}$$

Combining (G.36) and (G.39), define

$$\mathcal{F}_{t,\tau}(X_0) := \mathcal{J}_\tau(X_0)\mathcal{W}_t(X_0), \qquad \mathcal{F}^{\mathrm{aff}}_{t,\tau}(X_0) := e^{2\Lambda(\tau)}W_t^{\mathrm{aff}}(X_0).$$

Then

$$e^{-7M} \le \frac{\mathcal{F}_{t,\tau}}{\mathcal{F}^{\mathrm{aff}}_{t,\tau}} \le e^{7M}, \qquad \left|\frac{\mathcal{F}_{t,\tau}}{\mathcal{F}^{\mathrm{aff}}_{t,\tau}} - 1\right| \le e^{7M} - 1. \tag{G.40}$$

Step 5: stability of the complete signed and absolute responses. Let

$$\varepsilon_M := e^{8M} - 1.$$

Using (G.35) and (G.40),

$$\begin{aligned}
&\left|q(\mathscr{X}_\tau X_0, \tau)\mathcal{F}_{t,\tau} - q_0(X_0)\mathcal{F}^{\mathrm{aff}}_{t,\tau}\right| \\
&\quad \le \mathcal{F}^{\mathrm{aff}}_{t,\tau}\left[e^{7M}(e^M - 1) + (e^{7M} - 1)\right]|Z_{\beta,0}(X_0)| \\
&\quad = \varepsilon_M \mathcal{F}^{\mathrm{aff}}_{t,\tau}|Z_{\beta,0}(X_0)|.
\end{aligned}$$

The identical estimate holds after replacing q and q_0 by their absolute values, because $||q| - |q_0|| \le |q - q_0|$. Integrating first in X_0 and then in τ gives

$$|\mathcal{P} - \mathcal{P}_{\mathrm{aff}}| + |\mathcal{V} - \mathcal{V}_{\mathrm{aff}}| \le 2\varepsilon_M \mathcal{B}^{\mathrm{aff}}_Z(t), \tag{G.41}$$

where

$$\mathcal{B}^{\mathrm{aff}}_Z(t) := \frac{3}{4}I_t \int_{\mathcal{Q}} W_t^{\mathrm{aff}}(X_0)|Z_{\beta,0}(X_0)|\, dX_0.$$

The repaired G3 theorem gives

$$\mathcal{B}^{\mathrm{aff}}_Z(t) \le \mathfrak{C}^*_Z \mathcal{V}_{\mathrm{aff}}(t). \tag{G.42}$$

Thus (17.10) holds, for example, with the namespace-safe universal choices

$$\boxed{\mathfrak{c}_0 = 8, \qquad \mathfrak{c}_1 = 2.}$$

At $t = 0$ both sides vanish. For $t > 0$, $I_t > 0$ and the G3 denominator is strictly positive, so $\mathcal{V}_{\mathrm{aff}}(t) > 0$.

Step 6: preservation of a strict polarization margin. The audited affine theorem E4 gives

$$\mathcal{P}_{\mathrm{aff}}(t) \ge \pi^*_{\mathrm{aff}}\mathcal{V}_{\mathrm{aff}}(t), \qquad \pi^*_{\mathrm{aff}} := \frac{1 - \upsilon^*_{\mathrm{aff}}}{1 + \upsilon^*_{\mathrm{aff}}} > 0. \tag{G.43}$$

Put

$$\eta_M := 2\mathfrak{C}^*_Z(e^{8M} - 1).$$

Then (G.41)–(G.43) imply

$$\mathcal{P}(t) \geq (\pi^*_{\mathrm{aff}} - \eta_M)\mathcal{V}_{\mathrm{aff}}(t), \qquad \mathcal{V}(t) \leq (1+\eta_M)\mathcal{V}_{\mathrm{aff}}(t).$$

Choose $M_* > 0$ so that $\eta_{M_*} < \pi^*_{\mathrm{aff}}/2$ and define

$$\pi_{107} := \frac{\pi^*_{\mathrm{aff}} - \eta_{M_*}}{1+\eta_{M_*}} > 0. \tag{G.44}$$

For every $M \leq M_*$ this proves

$$a(t) = \mathcal{P}(t) \geq \pi_{107}\mathcal{V}(t) = \pi_{107}[a^+(t) + a^-(t)].$$

The argument is entirely source-specific: the ratio constant is used only after physical admission, and no generic swirl-free Euler inequality is introduced. □

Audit status: GREEN/COMPLETE.

G5. First polarization loss pays a definite deformation bill. *Mutual pointers:* grand-finale entry / short proof/status / main statement corollary 17.6.

Restatement of corollary 17.6. Let

$$\mathfrak{F}_{\mathrm{pol}} := \{t > 0 : \ a(t) < \pi_{107}[a^+(t) + a^-(t)]\}.$$

If $\mathfrak{F}_{\mathrm{pol}} \neq \varnothing$, define

$$t_* := \inf \mathfrak{F}_{\mathrm{pol}}.$$

Then

$$\boxed{\mathcal{K}_{\mathrm{cent}}([0,t_*]) \geq M_* > 0.}$$

For $X = (x,y)$, $s := y/x$, and $y_\lambda := \lambda y$, define

$$\mathcal{L}_{\mathrm{pol}}(t) := \|\mathbf{A}_{\mathrm{sg}}(t)\|_\infty + \|\mathfrak{A}(t)\|_\infty + \sup_{X\in\mathcal{Q}} \int_0^1 \left(\frac{1}{4}|\mathfrak{A}| + \frac{1}{2}|\mathfrak{J}| + \frac{1}{4}y_\lambda|H| + \frac{1}{2}s|\mathfrak{T}|\right)(\lambda X, t)\,\frac{d\lambda}{\lambda}.$$

Then there exists $c_{\mathrm{pol}} > 0$, depending only on the fixed customer and affine margin, such that

$$\boxed{\int_0^{t_*} \mathcal{L}_{\mathrm{pol}}(\tau)\,d\tau \geq c_{\mathrm{pol}}.}$$

Proof. **Step 1: remove the first-loss ambiguity.** For a classical solution, a, a^+, and a^- are continuous in time, while $\mathcal{K}_{\mathrm{cent}}([0,t])$ is continuous and nondecreasing. Every actual strict failure time $t \in \mathfrak{F}_{\mathrm{pol}}$ must satisfy

$$\mathcal{K}_{\mathrm{cent}}([0,t]) > M_*,$$

for otherwise G4 would imply the opposite inequality. Choose $t_n \in \mathfrak{F}_{\mathrm{pol}}$ with $t_n \downarrow t_*$. Continuity gives

$$\mathcal{K}_{\mathrm{cent}}([0,t_*]) = \lim_{n\to\infty} \mathcal{K}_{\mathrm{cent}}([0,t_n]) \geq M_*. \tag{G.45}$$

In particular $t_* > 0$, since the centered action vanishes at $t = 0$. This proves (17.13) without pretending that the infimum itself must already be a strict-failure time.

Step 2: return the centered-value defects to the logarithmic star hull. By the audited radial-star identity B5,

$$\mathcal{R}_v := xv_x + yv_y = -\frac{1}{4}\mathfrak{A} + \frac{1}{2}\mathfrak{J} - \frac{1}{4}yH - \frac{1}{2}s\mathfrak{T}, \tag{G.46}$$

and, crucially,

$$v(X,t) - a(t) = \int_0^1 \mathcal{R}_v(\lambda X, t)\frac{d\lambda}{\lambda}. \tag{G.47}$$

The ordinary measure $d\lambda$ from the earlier radial derivation is not used. Taking absolute values in (G.46)–(G.47) gives

$$\|v(\cdot,t) - a(t)\|_\infty \le \mathcal{S}_{\mathrm{pol}}(t), \tag{G.48}$$

where

$$\mathcal{S}_{\mathrm{pol}}(t) := \sup_{X\in\mathcal{Q}} \int_0^1 \left(\frac{1}{4}|\mathfrak{A}| + \frac{1}{2}|\mathfrak{J}| + \frac{1}{4}y_\lambda|H| + \frac{1}{2}s|\mathfrak{T}|\right)(\lambda X, t)\frac{d\lambda}{\lambda}.$$

The logarithmic integral is finite on a classical branch: the apex identities make the numerator $O(\lambda)$ on compact rays, exactly as in B5. The exact relation

$$g - 2a = 2(v-a) - \mathfrak{A} \tag{G.49}$$

then yields

$$\|g(\cdot,t) - 2a(t)\|_\infty \le 2\mathcal{S}_{\mathrm{pol}}(t) + \|\mathfrak{A}(t)\|_\infty.$$

Therefore, pointwise in time,

$$\begin{aligned}\|v-a\|_\infty + \|g-2a\|_\infty + \|\mathbf{A}_{\mathrm{sg}}\|_\infty \\ \le 3\mathcal{S}_{\mathrm{pol}} + \|\mathfrak{A}\|_\infty + \|\mathbf{A}_{\mathrm{sg}}\|_\infty \le 3\mathcal{L}_{\mathrm{pol}}.\end{aligned}$$

Integrating gives the explicit routing estimate

$$\boxed{\mathcal{K}_{\mathrm{cent}}([0,t]) \le 3\int_0^t \mathcal{L}_{\mathrm{pol}}(\tau)\,d\tau.} \tag{G.50}$$

Combining (G.45) and (G.50) at $t = t_*$ gives

$$\boxed{\int_0^{t_*} \mathcal{L}_{\mathrm{pol}}(\tau)\,d\tau \ge \frac{M_*}{3}.}$$

Thus (17.14) holds with the explicit admissible choice $c_{\mathrm{pol}} = M_*/3 > 0$.

This is a one-shot routing ticket, not a terminal singularity criterion. It may not be charged again at every later Hodge refresh, and its adverse portion must still be returned to already named tagged genealogy/restart accounts or to a genuine continuation exit. □

Audit status: GREEN/COMPLETE.

Appendix H. Causal adverse-response ledger

stage 131 completes H1–H8. stage 130 supplied visibility comparison, explicit raw response capacity, fresh-apex anti-rotation, and the exact old/generated Kato–Hodge response split. stage 131 adds the fresh-primary quadratic law, the global causal Hodge resolvent, the three-account negative-bank bound, and the resummed strict-gap criterion.

H1. Final/deposition Hodge visibility comparison. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 18.1.

Restatement of proposition 18.1. For $s = y/x$ and $\mathcal{V}_H(t;\tau,X) = K_H(\mathscr{X}_{t,\tau}X)$,

$$\mathcal{D}_{xy}\log K_H = \frac{(3-2s)v + (1-4s)g}{1+s}.$$

With

$$\mathcal{K}_{\mathrm{vis}}(\tau,t) = \int_\tau^t (\|v(\cdot,\xi)\|_\infty + \|g(\cdot,\xi)\|_\infty)\, d\xi,$$

finite visibility action implies

$$\boxed{e^{-4\mathcal{K}_{\mathrm{vis}}(\tau,t)}K_H(X) \le \mathcal{V}_H(t;\tau,X) \le e^{4\mathcal{K}_{\mathrm{vis}}(\tau,t)}K_H(X).}$$

No division by a is used.

Full rigorous proof. Recall

$$K_H(x,y) = \frac{x\sqrt{y}}{(x+y)^{5/2}}, \qquad \mathbf{b} = (-2xv, 2yg).$$

For $x, y > 0$,

$$\partial_x \log K_H = \frac{1}{x} - \frac{5}{2(x+y)}, \qquad \partial_y \log K_H = \frac{1}{2y} - \frac{5}{2(x+y)}.$$

Therefore

$$\begin{aligned}\mathcal{D}_{xy}\log K_H &= (-2xv)\left(\frac{1}{x} - \frac{5}{2(x+y)}\right) + (2yg)\left(\frac{1}{2y} - \frac{5}{2(x+y)}\right)\\ &= -2v + g + 5\frac{xv - yg}{x+y}\\ &= \frac{(3-2s)v + (1-4s)g}{1+s}. \end{aligned} \tag{H.1}$$

This independently reproduces (18.1).

The angular coefficients are uniformly bounded on $s > 0$:

$$\left|\frac{3-2s}{1+s}\right| \le 3, \qquad \left|\frac{1-4s}{1+s}\right| \le 4. \tag{H.2}$$

Let $Z(\xi) = \mathscr{X}_{\xi,\tau}X$. Along this material trajectory,

$$\begin{aligned}\left|\frac{d}{d\xi}\log K_H(Z(\xi))\right| &\le 3\|v(\cdot,\xi)\|_\infty + 4\|g(\cdot,\xi)\|_\infty\\ &\le 4\big(\|v(\cdot,\xi)\|_\infty + \|g(\cdot,\xi)\|_\infty\big).\end{aligned}$$

Integrating from τ to t gives

$$\left|\log\frac{K_H(\mathscr{X}_{t,\tau}X)}{K_H(X)}\right| \le 4\mathcal{K}_{\mathrm{vis}}(\tau,t). \tag{H.3}$$

Exponentiating and using the definition of $\mathcal{V}_H$ proves (18.3). The estimate compares two positive geometric weights on the same label; no normalization by a, Λ, a^+, or any tagged bank is introduced. □

Audit status: GREEN/COMPLETE.

H2. Finite raw response capacity. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 18.2.

Restatement of proposition 18.2. For a classical observation time $t > 0$, define

$$\mathcal{K}_{vx}(0,t) = \int_0^t \|v - xv_x\|_{L^\infty(\mathcal{Q})}\, d\xi.$$

If $\mathcal{K}_{\mathrm{vis}}(0,t)$ and $\mathcal{K}_{vx}(0,t)$ are finite, then

$$\boxed{\mathfrak{X}_-^H(t) \le \mathfrak{c}_{\mathrm{resp}}(t) \int_0^t \left[\mathcal{E}_{H,1}\mathcal{E}_{\mathrm{sg}}^{(0,0)} + \mathcal{L}_B\mathcal{Q}_{\mathrm{sg}}^{(1)}\right](\tau) d\tau,}$$

where

$$\mathfrak{c}_{\mathrm{resp}}(t) = \mathfrak{c}_{\mathrm{rot}}\, t \exp\big(4\mathcal{K}_{\mathrm{vis}}(0,t) + 2\mathcal{K}_{vx}(0,t)\big)$$

for one fixed geometric constant $\mathfrak{c}_{\mathrm{rot}} > 0$.

Full rigorous proof. Only the positive integral defining $\mathfrak{X}_-^H$ is used here; the master negative-bank comparison of H4 is not needed for this capacity estimate. For $0 \le \tau \le t$, H1 gives

$$\mathcal{V}_H(t;\tau,X) \le e^{4\mathcal{K}_{\mathrm{vis}}(0,t)} K_H(X). \tag{H.4}$$

The residence multiplier in (18.9) satisfies

$$\begin{aligned} 0 \le \mathfrak{R}_q(t,\tau;X) &\le \int_\tau^t \exp\left(2\int_\tau^\rho \|v - xv_x\|_\infty\, d\xi\right) d\rho \\ &\le t\, e^{2\mathcal{K}_{vx}(0,t)}. \end{aligned} \tag{H.5}$$

Thus

$$\mathfrak{X}_-^H(t) \le \frac{3}{4} t e^{4\mathcal{K}_{\mathrm{vis}}(0,t)+2\mathcal{K}_{vx}(0,t)} \int_0^t \int_{\mathcal{Q}} K_H e_-\, dX\, d\tau. \tag{H.6}$$

We now price the instantaneous carrier integral using the already audited F6–F7 identities. F6 gives

$$e_- = [(yH + 2s\mathfrak{T})C_\beta]_+ \le (y|H| + 2s|\mathfrak{T}|)|C_\beta|.$$

With

$$d\mu_c = \frac{dX}{\mathfrak{r}}, \qquad \mathfrak{r} = xs^{1/3}, \qquad M_H := \sqrt{xy}\, H,$$

F7's exact carrier conversion yields

$$\int_{\mathcal{Q}} K_H y|H||C_\beta|\, dX \le \|w_{\mathrm{rot}}\|_\infty \mathcal{E}_{H,1}\mathcal{E}_{\mathrm{sg}}^{(0,0)}. \tag{H.7}$$

For the mixed-shear part, the lifted first-jet estimate

$$\|\sqrt{xy}\, \mathcal{T}_{\mathrm{mix}}\|_\infty \le \mathfrak{c}_B\mathcal{L}_B, \qquad \mathcal{T}_{\mathrm{mix}} = g_x - v_y = \mathfrak{T}/x,$$

gives

$$\int_{\mathcal{Q}} 2K_H s|\mathfrak{T}||C_\beta|\, dX \le \mathfrak{c}_B \|w_{\mathrm{rot}}\|_\infty \mathcal{L}_B\mathcal{Q}_{\mathrm{sg}}^{(1)}. \tag{H.8}$$

Consequently there is one fixed geometric constant $\mathfrak{c}_{\mathrm{rot}} > 0$ such that

$$\int_{\mathcal{Q}} K_H e_-\, dX \le \mathfrak{c}_{\mathrm{rot}} \left[\mathcal{E}_{H,1}\mathcal{E}_{\mathrm{sg}}^{(0,0)} + \mathcal{L}_B\mathcal{Q}_{\mathrm{sg}}^{(1)}\right]. \tag{H.9}$$

Absorbing the harmless factor 3/4 into $\mathfrak{c}_{\mathrm{rot}}$ and inserting (H.9) into (H.6) proves (18.5)–(18.6).

Thus any large response multiplier must be paid through the named visibility action or the named $v - xv_x$ first-jet action. The estimate proves finite raw capacity; it does not turn finiteness into a strict fraction of a^+. □

Audit status: GREEN/COMPLETE.

H3. Fresh-apex anti-rotation. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 18.3.

> **Restatement of proposition 18.3.** In the physical sector $\varepsilon_\beta = 1$, fix $I = [t_1, t_2]$ with $0 < t_1 < t_2$ and $J \Subset (0, \infty)$. For one sufficiently small $\ell > 0$, on $\mathcal{U}_{\ell,J} = \{0 < x < \ell,\ y/x \in J\}$,
>
> $$q > 0, \qquad H^+(x, sx, t) = q_H(t)sx + O_{I,J}(x^2), \qquad H^-(x, sx, t) = O_{I,J}(x^2),$$
>
> and
>
> $$\boxed{[-(4yv_y)C_\beta]_+ \le \mathfrak{c}_{I,J}\ell q.}$$

Full rigorous proof. Put

$$p(t) := u_y(0, 0, t) = \psi_\beta(t)/2.$$

The exact apex equation $\dot p = -2ap$ and the oriented fixed seed give $p(t) > 0$ at every finite classical time. Hence on the compact positive-time interval I,

$$p_* := \inf_{t\in I} p(t) > 0. \tag{H.10}$$

The equator is material. Since $u_0(x, 0) = 0$ and $\Gamma = xu$ is passively transported, $u(x, 0, t) = 0$ throughout the classical branch. Smoothness therefore allows the local factorization

$$u(x, y, t) = y\widehat w(x, y, t), \qquad \widehat w(0, 0, t) = p(t).$$

Uniform Taylor expansion on the compact spacetime cone $I \times J$ gives

$$u(x, sx, t) = p(t)sx + O_{I,J}(x^2), \qquad u_y(x, sx, t) = p(t) + O_{I,J}(x). \tag{H.11}$$

Because $\varepsilon_\beta = 1$,

$$q = \Theta_y = 2uu_y = 2p(t)^2 sx + O_{I,J}(x^2). \tag{H.12}$$

Writing $J = [s_-, s_+]$ with $s_- > 0$, one fixed decrease of ℓ gives

$$q(x, sx, t) \ge p_*^2 s_- x > 0 \qquad (t \in I,\ s \in J,\ 0 < x < \ell). \tag{H.13}$$

The audited C4 genealogy expansion is uniform on compact positive-time intervals; its C15 uniformization gives

$$H^+(x, sx, t) = q_H(t)sx + O_{I,J}(x^2), \qquad H^-(x, sx, t) = O_{I,J}(x^2). \tag{H.14}$$

For completeness, the first coefficient satisfies $\dot q_H + 6aq_H = 4p^2$ with $q_H(0) = 0$, so $q_H > 0$ on I.

It remains to estimate the adverse exchange. From $u = y\widehat w$,

$$\Theta = y^2\widehat w^2, \qquad \Theta + \frac{x}{2}\Theta_x = y^2\big(\widehat w^2 + x\widehat w\widehat w_x\big).$$

Hence on $I \times J$,

$$|C_\beta(x, sx, t)| = \left|\frac{\Theta + \frac{x}{2}\Theta_x}{y}\right| \le \mathfrak{c}_{\Theta,I,J}x. \tag{H.15}$$

Classical compactness also gives $\|v_y\|_{L^\infty(\mathcal{U}_{\ell,J}\times I)} \le \mathfrak{c}_{v,I,J}$. Since $y = sx \le s_+x$,

$$e_- = [-(4yv_y)C_\beta]_+ \le 4s_+\mathfrak{c}_{v,I,J}\mathfrak{c}_{\Theta,I,J}x^2. \tag{H.16}$$

Combining (H.16) with (H.13) yields one finite $\mathfrak{c}_{I,J}$ for which

$$e_- \le \mathfrak{c}_{I,J} x q \le \mathfrak{c}_{I,J} \ell q.$$

This proves (18.8). The small factor is the physical apex scale ℓ; no response-age or bank normalization is used. □

Audit status: GREEN/COMPLETE.

H4. Exact generated-negative Hodge response. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 18.4.

Restatement of theorem 18.4. With

$$\mathsf{d}_q = \frac{3}{2}\mathfrak{A} - \mathfrak{J}, \qquad e_- = [-(4yv_y)C_\beta]_+,$$

define $\mathfrak{R}_q$ by (18.9), the generated response $\mathfrak{X}_-^H$ by (18.10), and

$$\mathfrak{X}_{-,\mathrm{old}}^H(t) = \frac{3}{4}\int_{\mathcal{Q}} [q(X,0)]_- \mathcal{V}_H(t;0,X)\mathfrak{R}_q(t,0;X)\,dX.$$

Then

$$\boxed{a^-(t) \le \mathfrak{X}_{-,\mathrm{old}}^H(t) + \mathfrak{X}_-^H(t).}$$

Every factor in both accounts is nonnegative.

Full rigorous proof. **Step 1: Kato separation of old and generated negative source.** The first row of the audited F1 source-gradient system is

$$\mathcal{D}_{xy} q = \mathsf{d}_q q + (4yv_y)C_\beta. \tag{H.17}$$

For the convex negative-part function $z \mapsto [z]_-$, the classical/distributional Kato inequality gives

$$\mathcal{D}_{xy}[q]_- \le \mathsf{d}_q[q]_- + e_-, \qquad e_- = [-(4yv_y)C_\beta]_+. \tag{H.18}$$

Define the nonnegative characteristic comparison fields q_-^{old} and q_-^{gen} by

$$\mathcal{D}_{xy} q_-^{\mathrm{old}} = \mathsf{d}_q q_-^{\mathrm{old}}, \qquad q_-^{\mathrm{old}}(\cdot,0) = [q(\cdot,0)]_-, \tag{H.19}$$

$$\mathcal{D}_{xy} q_-^{\mathrm{gen}} = \mathsf{d}_q q_-^{\mathrm{gen}} + e_-, \qquad q_-^{\mathrm{gen}}(\cdot,0) = 0. \tag{H.20}$$

Scalar comparison along the material trajectories yields

$$[q]_- \le q_-^{\mathrm{old}} + q_-^{\mathrm{gen}}. \tag{H.21}$$

Step 2: exact Jacobian–integrating-factor product. For $0 \le \tau \le \rho$, define the material Jacobian

$$\mathsf{J}_{\rho,\tau}(X) := \det D_X \mathscr{X}_{\rho,\tau}(X).$$

Since $\nabla\cdot \mathbf{b} = g$,

$$\mathsf{J}_{\rho,\tau}(X) = \exp\left(\int_\tau^\rho g(\mathscr{X}_{\xi,\tau}X,\xi)\,d\xi\right). \tag{H.22}$$

Likewise set

$$\mathsf{I}_q(\rho,\tau;X) := \exp\left(\int_\tau^\rho \mathsf{d}_q(\mathscr{X}_{\xi,\tau}X,\xi)\,d\xi\right). \tag{H.23}$$

Using the protected identities $\mathfrak{A} = 2v - g$, $\mathfrak{J} = xv_x + yg_y$, and $g + 2yg_y = 2v + 2xv_x$, one obtains

$$\boxed{\mathsf{d}_q + g = \frac{3}{2}\mathfrak{A} - \mathfrak{J} + g = 2(v - xv_x).} \tag{H.24}$$

Therefore

$$\mathsf{J}_{\rho,\tau}(X)\mathsf{I}_q(\rho,\tau;X) = \exp\left(2\int_\tau^\rho (v - xv_x)(\mathscr{X}_{\xi,\tau}X,\xi)\,d\xi\right). \tag{H.25}$$

Integrating (H.25) in ρ from τ to t gives exactly $\mathfrak{R}_q(t,\tau;X)$.

Step 3: old initial-slope response. By the exact source-polarized genealogy response (15.2) and (H.21), the old comparison contribution to $a^-(t)$ is

$$\frac{3}{4}\int_0^t\int_{\mathcal{Q}} q_-^{\mathrm{old}}(Y,\rho)\mathcal{V}_H(t;\rho,Y)\,dY\,d\rho.$$

Along $Y = \mathscr{X}_{\rho,0}X$, $q_-^{\mathrm{old}}(Y,\rho) = \mathsf{I}_q(\rho,0;X)[q(X,0)]_-$, while $dY = \mathsf{J}_{\rho,0}(X)dX$. The flow semigroup gives

$$\mathcal{V}_H(t;\rho,\mathscr{X}_{\rho,0}X) = K_H(\mathscr{X}_{t,\rho}\mathscr{X}_{\rho,0}X) = \mathcal{V}_H(t;0,X).$$

Hence Tonelli/Fubini for the nonnegative integrand and (H.25) give the exact old account

$$\begin{aligned}\frac{3}{4}\int_{\mathcal{Q}}[q(X,0)]_-\mathcal{V}_H(t;0,X)\left(\int_0^t \mathsf{J}_{\rho,0}(X)\mathsf{I}_q(\rho,0;X)\,d\rho\right)dX\\ = \mathfrak{X}^H_{-,\mathrm{old}}(t).\end{aligned} \tag{H.26}$$

Step 4: generated adverse cross-shear response. Variation of constants in (H.20) says that a deposition $e_-(X,\tau)d\tau$ contributes at a later source time $\rho \ge \tau$ along $\mathscr{X}_{\rho,\tau}X$ with the positive factor $\mathsf{I}_q(\rho,\tau;X)$. Insert this Duhamel representation into the exact q-to-H^- response. Tonelli applies because every factor is nonnegative. After interchanging the triangle $0 \le \tau \le \rho \le t$ and changing variables $Y = \mathscr{X}_{\rho,\tau}X$, the flow semigroup gives

$$\mathcal{V}_H(t;\rho,\mathscr{X}_{\rho,\tau}X) = \mathcal{V}_H(t;\tau,X),$$

and the remaining ρ-integral is

$$\int_\tau^t \mathsf{J}_{\rho,\tau}(X)\mathsf{I}_q(\rho,\tau;X)\,d\rho = \mathfrak{R}_q(t,\tau;X).$$

Consequently the generated contribution is bounded by exactly

$$\frac{3}{4}\int_0^t\int_{\mathcal{Q}} e_-(X,\tau)\mathcal{V}_H(t;\tau,X)\mathfrak{R}_q(t,\tau;X)\,dX\,d\tau = \mathfrak{X}^H_-(t). \tag{H.27}$$

Combining (H.26) and (H.27) proves (18.12).

The causal bookkeeping is now explicit: the old account comes only from $[q(\cdot,0)]_-$, while every later generated adverse contribution enters through the tagged cross-shear source e_-. The flow Jacobian, source-gradient integrating factor, and Hodge visibility are all positive routing factors. No quotient by a, Λ, a bank total, or a common provenance tag is taken. □

Audit status: GREEN/COMPLETE.

H5. Fresh primary conversion is quadratically small. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 18.5.

Restatement of proposition 18.5. Fix $t > 0$. With

$$\mathfrak{m}^+_{\mathrm{prim}}(\tau) = \|[2\Theta + x\Theta_x]_+(\cdot,\tau)\|_\infty, \qquad A^+_{t\leftarrow\tau} = \frac{3}{8}\int H^+(X,\tau)\mathcal{V}_H(t;\tau,X)\,dX,$$

define, for $0 < h \le h_0 \le t$,

$$\mathfrak{X}^{\mathrm{res}}_{+,\mathrm{fresh,h}}(t) = \int_{t-h}^t \mathcal{G}_{\mathrm{rec}}(t-\tau)\mathfrak{m}^+_{\mathrm{prim}}(\tau)A^+_{t\leftarrow\tau}\,d\tau.$$

If $\mathfrak{c}_{\mathrm{rec}} = \sup_{0<u\le h_0} \mathcal{G}_{\mathrm{rec}}(u)/u$ and $\mathfrak{m}^{+,*}_{\mathrm{prim,h}} = \sup_{t-h\le\tau\le t} \mathfrak{m}^{+}_{\mathrm{prim}}(\tau)$, then

$$\boxed{\mathfrak{X}^{\mathrm{res}}_{+,\mathrm{fresh,h}}(t) \le \frac{1}{2}\mathfrak{c}_{\mathrm{rec}}\mathfrak{m}^{+,*}_{\mathrm{prim,h}} h^2 a^+(t).}$$

Proof. For $x, y > 0$, the orthogonalized radial source gradient satisfies

$$\sqrt{xy}\,\widehat{C}_\beta = \sqrt{xy}\,(2\sqrt{s}C_\beta) = 2yC_\beta = 2\Theta + x\Theta_x. \tag{H.28}$$

The right-hand side is smooth in the squared variables and rapidly decaying for the fixed customer. Thus it supplies the global axis extension used in the definition of $\mathfrak{m}^{+}_{\mathrm{prim}}$, and this coefficient is bounded on every compact classical time interval.

Next fix the final observation time t. The positive genealogy solves

$$\partial_t H^+ + \nabla\cdot(\mathbf{b}H^+) = 2[q]_+, \qquad H^+(\cdot,0) = 0.$$

Transporting this continuity equation from source time τ to observation time t and testing against K_H gives the exact positive-bank decomposition

$$a^+(t) = A^+_{t\leftarrow\tau} + \frac{3}{4}\int_\tau^t \int_{\mathcal{Q}} [q(X,\rho)]_+ \mathcal{V}_H(t;\rho,X)\,dX\,d\rho. \tag{H.29}$$

Every term in the second summand is nonnegative, hence

$$0 \le A^+_{t\leftarrow\tau} \le a^+(t) \qquad (0\le\tau\le t). \tag{H.30}$$

By H6 the Green function is C^1 on $[0,\infty)$, $\mathcal{G}_{\mathrm{rec}}(0) = 0$, and $\mathcal{G}'_{\mathrm{rec}}(0) = 1$. Therefore $\mathcal{G}_{\mathrm{rec}}(u)/u$ extends continuously to $u = 0$ with value 1; its supremum $\mathfrak{c}_{\mathrm{rec}}$ on $[0,h_0]$ is finite. Using (H.30), for $\tau\in[t-h,t]$,

$$\mathcal{G}_{\mathrm{rec}}(t-\tau)\mathfrak{m}^{+}_{\mathrm{prim}}(\tau)A^+_{t\leftarrow\tau} \le \mathfrak{c}_{\mathrm{rec}}(t-\tau)\mathfrak{m}^{+,*}_{\mathrm{prim,h}} a^+(t).$$

Integrating and using $\int_{t-h}^t (t-\tau)d\tau = h^2/2$ gives the claimed estimate. The two powers of h have distinct causal origins: one is the short-time Green factor and one is the source-time integration. No pointwise reproduction-number hypothesis is used. □

Audit status: GREEN/COMPLETE.

H6. Causal Hodge resolvent for negative self-recycling. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 18.6.

Restatement of theorem 18.6. With final visibility $\mathcal{V}_H(t;\tau,X)$, define

$$\mathfrak{m}_- = [-(2\Theta + x\Theta_x)]_+, \quad e_{\mathfrak{T}} = [\sqrt{s}\mathfrak{T}\widehat{C}_\beta]_+, \quad e_{H,+} = \frac{1}{2}\sqrt{xy}H^+[\widehat{C}_\beta]_+, \quad f_{\mathrm{prim}} = e_{\mathfrak{T}} + e_{H,+}.$$

Let $(q^{\mathrm{maj}}_-, H^{\mathrm{maj}}_-)$ solve the global cooperative system

$$\mathcal{D}_{xy}q^{\mathrm{maj}}_- = \mathsf{d}_q q^{\mathrm{maj}}_- + f_{\mathrm{prim}} + \frac{1}{2}\mathfrak{m}_- H^{\mathrm{maj}}_-, \qquad \mathcal{D}_{xy}H^{\mathrm{maj}}_- + gH^{\mathrm{maj}}_- = 2q^{\mathrm{maj}}_-,$$

with data $([q_0]_-, 0)$. Then $[q]_- \le q^{\mathrm{maj}}_-$ and $H^- \le H^{\mathrm{maj}}_-$. If $\mathcal{Z}_{\mathrm{rec}}, \mathcal{H}_{\mathrm{rec}}, \mathcal{I}_{\mathrm{rec}}$ and $\mathcal{G}_{\mathrm{rec}}$ are defined as in the main statement, then

$$\boxed{a^-(t) \le \frac{3}{4}\mathcal{Z}_{\mathrm{rec}}(0)\mathcal{G}_{\mathrm{rec}}(t) + \frac{3}{4}\int_0^t \mathcal{G}_{\mathrm{rec}}(t-\tau)\mathcal{I}_{\mathrm{rec}}(\tau)\,d\tau.}$$

Proof. **Step 1: the global Kato decomposition.**

The F2 rotor equation is

$$\mathcal{D}_{xy}q = \mathsf{d}_q q - (\omega_H + \sqrt{s}\mathfrak{T})\widehat{C}_\beta, \qquad \omega_H = \frac{1}{2}\sqrt{xy}\,H. \tag{H.31}$$

Kato's inequality for the negative part therefore gives

$$\mathcal{D}_{xy}[q]_- \le \mathsf{d}_q[q]_- + [(\omega_H + \sqrt{s}\mathfrak{T})\widehat{C}_\beta]_+. \tag{H.32}$$

Write $H = H^+ - H^-$. Since $H^\pm \ge 0$,

$$\begin{aligned}[(\omega_H + \sqrt{s}\mathfrak{T})\widehat{C}_\beta]_+ &\le [\sqrt{s}\mathfrak{T}\widehat{C}_\beta]_+ + \frac{1}{2}\sqrt{xy}\,H^+[\widehat{C}_\beta]_+ + \frac{1}{2}\sqrt{xy}\,H^-[-\widehat{C}_\beta]_+ \\ &= f_{\mathrm{prim}} + \frac{1}{2}\mathfrak{m}_- H^-. \end{aligned} \tag{H.33}$$

Here (H.28) gives, throughout the open quadrant and by continuous extension to the axes,

$$\sqrt{xy}[-\widehat{C}_\beta]_+ = [-(2\Theta + x\Theta_x)]_+ = \mathfrak{m}_-. \tag{H.34}$$

Thus no compact mesoscopic indicator is needed. This is the essential correction to the earlier resolvent wording: the full exterior self-recycling coefficient is kept.

The true negative genealogy satisfies

$$\mathcal{D}_{xy}H^- + gH^- = 2[q]_-, \qquad H^-(\cdot,0) = 0. \tag{H.35}$$

Equations (H.33) and (H.35) form a cooperative differential inequality. The majorant system in the theorem has the same nonnegative off-diagonal coefficients and the same initial data. Along each characteristic, a first-contact argument for the two differences $q_-^{\mathrm{maj}} - [q]_-$ and $H_-^{\mathrm{maj}} - H^-$ proves

$$[q]_- \le q_-^{\mathrm{maj}}, \qquad H^- \le H_-^{\mathrm{maj}}. \tag{H.36}$$

Because the solution is classical and rapidly decaying, all coefficients in this characteristic system are bounded on every fixed finite continuation interval.

Step 2: keep final Hodge visibility inside the system.

For fixed final time t, put

$$\mathscr{V}_{\mathrm{H}}(\tau, X) := \mathcal{V}_H(t;\tau,X) = K_H(\mathscr{X}_{t,\tau}X).$$

The semigroup identity for $\mathscr{X}$ gives the backward transport equation

$$\partial_\tau \mathscr{V}_{\mathrm{H}} + \mathbf{b}\cdot\nabla\mathscr{V}_{\mathrm{H}} = 0, \qquad \mathscr{V}_{\mathrm{H}}(t,X) = K_H(X). \tag{H.37}$$

Recall $\nabla\cdot\mathbf{b} = g$. Differentiate $\mathcal{Z}_{\mathrm{rec}} = \int q_-^{\mathrm{maj}}\mathscr{V}_{\mathrm{H}}$ and use (H.37). Integration by parts is first performed on truncated quadrants and then the truncation is removed using the regular axis behavior and rapid decay. One obtains

$$\begin{aligned}\frac{d}{d\tau}\mathcal{Z}_{\mathrm{rec}}(\tau) = &\int_{\mathcal{Q}} (\mathsf{d}_q + g) q_-^{\mathrm{maj}}\mathscr{V}_{\mathrm{H}}\,dX + \mathcal{I}_{\mathrm{rec}}(\tau) \\ &+ \frac{1}{2}\int_{\mathcal{Q}} \mathfrak{m}_- H_-^{\mathrm{maj}}\mathscr{V}_{\mathrm{H}}\,dX. \end{aligned} \tag{H.38}$$

The exact H4 identity

$$\mathsf{d}_q + g = 2(v - xv_x) \tag{H.39}$$

therefore yields

$$D_\tau^+\mathcal{Z}_{\mathrm{rec}} \le \mathcal{A}_{\mathrm{rec}}\mathcal{Z}_{\mathrm{rec}} + \mathcal{I}_{\mathrm{rec}} + \frac{1}{2}\mathcal{B}_{\mathrm{rec}}\mathcal{H}_{\mathrm{rec}}. \tag{H.40}$$

For the second stock the conservative form of the H_-^{maj} equation gives the exact cancellation of the transport divergence:

$$\frac{d}{d\tau}\mathcal{H}_{\mathrm{rec}}(\tau) = 2\mathcal{Z}_{\mathrm{rec}}(\tau). \tag{H.41}$$

The coefficient $\mathcal{B}_{\mathrm{rec}}$ is finite without a mesoscopic gate because $\mathfrak{m}_- = [-(2\Theta + x\Theta_x)]_+$ is a smooth rapidly decaying fixed-source field; $\mathcal{A}_{\mathrm{rec}}$ is finite on every compact classical interval by smoothness of v.

Step 3: scalar causal resolvent.

Set

$$\mathcal{U}_{\mathrm{rec}}(\tau) := \int_0^\tau \mathcal{Z}_{\mathrm{rec}}(\rho)\, d\rho.$$

Because $\mathcal{H}_{\mathrm{rec}}(0) = 0$, (H.41) gives

$$\mathcal{H}_{\mathrm{rec}}(\tau) = 2\mathcal{U}_{\mathrm{rec}}(\tau), \qquad \mathcal{U}'_{\mathrm{rec}} = \mathcal{Z}_{\mathrm{rec}}. \tag{H.42}$$

Hence

$$D_\tau^+\mathcal{U}'_{\mathrm{rec}} \le \mathcal{A}^*_{\mathrm{rec}}\mathcal{U}'_{\mathrm{rec}} + \mathcal{B}^*_{\mathrm{rec}}\mathcal{U}_{\mathrm{rec}} + \mathcal{I}_{\mathrm{rec}}(\tau), \tag{H.43}$$

with

$$\mathcal{U}_{\mathrm{rec}}(0) = 0, \qquad \mathcal{U}'_{\mathrm{rec}}(0) = \mathcal{Z}_{\mathrm{rec}}(0). \tag{H.44}$$

Let

$$\overline{\mathcal{U}}(\tau) := \mathcal{Z}_{\mathrm{rec}}(0)\mathcal{G}_{\mathrm{rec}}(\tau) + \int_0^\tau \mathcal{G}_{\mathrm{rec}}(\tau-\rho)\mathcal{I}_{\mathrm{rec}}(\rho)\, d\rho. \tag{H.45}$$

The explicit formula for $\mathcal{G}_{\mathrm{rec}}$ gives

$$\mathcal{G}_{\mathrm{rec}}(0) = 0, \qquad \mathcal{G}'_{\mathrm{rec}}(0) = 1, \qquad \mathcal{G}''_{\mathrm{rec}} = \mathcal{A}^*_{\mathrm{rec}}\mathcal{G}'_{\mathrm{rec}} + \mathcal{B}^*_{\mathrm{rec}}\mathcal{G}_{\mathrm{rec}}.$$

Thus $\overline{\mathcal{U}}$ solves the equality corresponding to (H.43) with the same data (H.44). Writing each second-order equation as the cooperative first-order pair $(\mathcal{U}_{\mathrm{rec}}, \mathcal{U}'_{\mathrm{rec}})$, whose coefficient matrix is

$$\begin{pmatrix} 0 & 1 \\ \mathcal{B}^*_{\mathrm{rec}} & \mathcal{A}^*_{\mathrm{rec}} \end{pmatrix},$$

a first-contact comparison gives

$$0 \le \mathcal{U}_{\mathrm{rec}}(\tau) \le \overline{\mathcal{U}}(\tau) \qquad (0 \le \tau \le t). \tag{H.46}$$

This is the causal Volterra resummation; no geometric-series norm condition appears.

Step 4: return to the true negative Hodge bank.

The exact continuity equation for H^- gives, after the same material change of variables,

$$a^-(t) = \frac{3}{8}\int_{\mathcal{Q}} K_H(X)H^-(X,t)\, dX = \frac{3}{4}\int_0^t\int_{\mathcal{Q}} [q(X,\tau)]_-\mathscr{V}_{\mathrm{H}}(\tau,X)\, dX\, d\tau. \tag{H.47}$$

By (H.36),

$$a^-(t) \le \frac{3}{4}\int_0^t \mathcal{Z}_{\mathrm{rec}}(\tau)\, d\tau = \frac{3}{4}\mathcal{U}_{\mathrm{rec}}(t).$$

Using (H.46) and (H.45) proves the asserted resolvent estimate. □

Audit status: GREEN/COMPLETE.

H7. Three-account negative-bank bound. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 18.7.

Restatement of proposition 18.7. Define

$$\mathcal{I}_{\mathfrak{T},t}(\tau)=\int e_{\mathfrak{T}}(X,\tau)\mathcal{V}_H(t;\tau,X)\,dX,\quad \mathfrak{m}^{+}_{\mathrm{prim}}(\tau)=\|[2\Theta+x\Theta_x]_+\|_\infty,$$

and

$$A^{+}_{t\leftarrow\tau}=\frac{3}{8}\int H^{+}(X,\tau)\mathcal{V}_H(t;\tau,X)\,dX.$$

Then the exact accounts from the main statement satisfy

$$\boxed{a^{-}\le \mathfrak{X}^{\mathrm{res}}_{-,\mathrm{old}}+\mathfrak{X}^{\mathrm{res}}_{\mathfrak{T}}+\mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{prim}}\le \mathfrak{X}^{\mathrm{res}}_{-,\mathrm{old}}+\mathfrak{X}^{\mathrm{res}}_{\mathfrak{T}}+\mathfrak{X}^{\mathrm{fac}}_{+,\mathrm{prim}}.}$$

Proof. The H6 input stock splits exactly as

$$\mathcal{I}_{\mathrm{rec}}(\tau)=\mathcal{I}_{\mathfrak{T},t}(\tau)+\mathcal{I}_{+,t}(\tau),\tag{H.48}$$

where

$$\mathcal{I}_{+,t}(\tau):=\frac{1}{2}\int_{\mathcal{Q}}\sqrt{xy}\,H^{+}(X,\tau)[\widehat{C}_\beta(X,\tau)]_+\mathcal{V}_H(t;\tau,X)\,dX.$$

By the global coefficient identity (H.28),

$$\begin{aligned}\mathcal{I}_{+,t}(\tau)&\le\frac{1}{2}\mathfrak{m}^{+}_{\mathrm{prim}}(\tau)\int_{\mathcal{Q}}H^{+}(X,\tau)\mathcal{V}_H(t;\tau,X)\,dX\\&=\frac{4}{3}\mathfrak{m}^{+}_{\mathrm{prim}}(\tau)A^{+}_{t\leftarrow\tau}.\end{aligned}\tag{H.49}$$

Insert the exact split (H.48) into the H6 resolvent before using any spatial supremum. The initial term is exactly $\mathfrak{X}^{\mathrm{res}}_{-,\mathrm{old}}$, the signed-$\mathfrak{T}$ integral is exactly $\mathfrak{X}^{\mathrm{res}}_{\mathfrak{T}}$, and the positive-Hodge input is exactly $\mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{prim}}$. This proves the first inequality in the main statement. Only afterward apply (H.49); the factor $(3/4)(4/3)=1$ gives $\mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{prim}}\le\mathfrak{X}^{\mathrm{fac}}_{+,\mathrm{prim}}$. Thus the factorized object is a secondary majorant, not the physical closure target.

The key audit point is globality. Neither (H.49) nor the H6 self-recycling coefficient contains $\mathbf{1}_{\mathcal{M}}$. Thus the theorem does not discard adverse conversion merely because its source point lies outside the compact rotor box. F8 remains available later to localize any substantial response, but localization is not assumed in order to prove this upstream bound. □

Audit status: GREEN/COMPLETE.

H8. Resummed strict-gap criterion. *Mutual pointers:* grand-finale entry / short proof/status / main statement conditional proposition 18.8.

Restatement of conditional proposition 18.8. If, for some $\delta\in(0,1)$,

$$\mathfrak{X}^{\mathrm{res}}_{-,\mathrm{old}}+\mathfrak{X}^{\mathrm{res}}_{\mathfrak{T}}+\mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{prim}}\le(1-\delta)a^{+},$$

then

$$\boxed{a^{-}\le(1-\delta)a^{+}.}$$

Proof. H7 gives

$$a^{-}\le\mathfrak{X}^{\mathrm{res}}_{-,\mathrm{old}}+\mathfrak{X}^{\mathrm{res}}_{\mathfrak{T}}+\mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{prim}}.$$

Applying the displayed hypothesis gives the conclusion immediately. The point of H8 is not algebraic difficulty but accounting closure: every $H^{-}\to H^{-}$ self-recycling generation has already been summed into the positive kernel $\mathcal{G}_{\mathrm{rec}}$, so no fourth reproduction account is hidden behind the strict-gap criterion. □

Audit status: GREEN/COMPLETE.

Appendix I. Supporting exact identities

This section contains I1–I7. Thus every supporting exact-identity slot I1–I7 is now audited without changing its permanent ledger ID or theorem label.

I1. Hodge visibility law. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 19.1.

Restatement of proposition 19.1. Put $s := y/x$. Along a material trajectory,

$$\boxed{\mathcal{D}_{xy} \log K_H = \frac{(3-2s)v + (1-4s)g}{1+s}.}$$

Equivalently, with

$$\vartheta := \frac{y}{x+y} = \frac{s}{1+s},$$
$$\mathcal{D}_{xy} \log K_H = v(3-5\vartheta) + g(1-5\vartheta).$$

On a compressive affine leg $(v, g) = (a, 2a)$ with $a > 0$, the aspect ratio increases strictly; the affine angular visibility profile has its unique maximum at $s = 1/2$, so any parcel that traverses this scale reaches its material visibility maximum there.

Full rigorous proof. In the open quadrant,

$$K_H(x, y) = \frac{x\sqrt{y}}{(x+y)^{5/2}}, \qquad \mathbf{b} = (-2xv, 2yg).$$

Therefore

$$\partial_x \log K_H = \frac{1}{x} - \frac{5}{2(x+y)}, \qquad \partial_y \log K_H = \frac{1}{2y} - \frac{5}{2(x+y)}.$$

Applying $\mathcal{D}_{xy} = \partial_t + \mathbf{b} \cdot \nabla$ gives

$$\begin{aligned} \mathcal{D}_{xy} \log K_H &= (-2xv)\left(\frac{1}{x} - \frac{5}{2(x+y)}\right) + (2yg)\left(\frac{1}{2y} - \frac{5}{2(x+y)}\right) \\ &= -2v + g + 5\frac{xv - yg}{x+y} \\ &= \frac{(3-2s)v + (1-4s)g}{1+s}. \end{aligned} \tag{I.1}$$

Since $\vartheta = s/(1+s)$, the two coefficients can also be written

$$\frac{3-2s}{1+s} = 3 - 5\vartheta, \qquad \frac{1-4s}{1+s} = 1 - 5\vartheta,$$

which proves the equivalent formula.

On an affine leg $v = a$, $g = 2a$,

$$\mathcal{D}_{xy} \log K_H = a\frac{5-10s}{1+s}. \tag{I.2}$$

The same affine characteristics satisfy

$$\mathcal{D}_{xy} \log s = \mathcal{D}_{xy} \log y - \mathcal{D}_{xy} \log x = 2g + 2v = 6a. \tag{I.3}$$

Hence if $a > 0$, s increases strictly. Equation (I.2) is positive for $0 < s < 1/2$, vanishes only at $s = 1/2$, and is negative for $s > 1/2$. Thus a material parcel that traverses the angular scale $s = 1/2$ reaches its unique Hodge-visibility maximum there. This is the pure visibility

maximum; it is distinct from the source-response timing extrema used in the later response ledger. □

Audit status: GREEN/COMPLETE.

I2. Exact guarded source-payment identity. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 19.2.

Restatement of proposition 19.2. Let $\mathcal{R}(t,\tau;X)$ be the vertical response from proposition 15.2 and put

$$\mathcal{R}_0(X) := \mathcal{R}(t,t;X).$$

Define

$$\mathcal{T}_\beta^{\mathrm{src}}(t) := \frac{3}{4}\int_{\mathcal{Q}} \Theta(X,t)[\mathcal{R}_0(X)]_+\,dX + \frac{3}{4}\int_0^t\int_{\mathcal{Q}} \Theta(X,\tau)[\partial_t\mathcal{R}(t,\tau;X)]_+\,dX\,d\tau,$$

$$\mathcal{W}_\beta^{\mathrm{src}}(t) := \frac{3}{4}\int_{\mathcal{Q}} \Theta(X,t)[\mathcal{R}_0(X)]_-\,dX + \frac{3}{4}\int_0^t\int_{\mathcal{Q}} \Theta(X,\tau)[\partial_t\mathcal{R}(t,\tau;X)]_-\,dX\,d\tau.$$

Then

$$\boxed{\dot a(t) = \mathcal{T}_\beta^{\mathrm{src}}(t) - \mathcal{W}_\beta^{\mathrm{src}}(t).}$$

Both accounts display the tagged factor Θ before normalization.

Full rigorous proof. For the fixed smooth rapidly decaying customer, E1 gives the exact tagged response formula

$$a(t) = \frac{3}{4}\int_0^t\int_{\mathcal{Q}} \Theta(X,\tau)\mathcal{R}(t,\tau;X)\,dX\,d\tau. \tag{I.4}$$

On every compact classical time interval the flow and its tangent map are smooth, and the fixed source is rapidly decaying. The response kernel and its observation-time derivative therefore admit an integrable dominating function on compact time intervals. Leibniz' rule may consequently be applied to (I.4). Since $\Theta(X,\tau)$ depends on the source time τ but not on the observation variable t while τ is held fixed,

$$\begin{aligned}\dot a(t) = {}& \frac{3}{4}\int_{\mathcal{Q}} \Theta(X,t)\mathcal{R}(t,t;X)\,dX \\ & + \frac{3}{4}\int_0^t\int_{\mathcal{Q}} \Theta(X,\tau)\partial_t\mathcal{R}(t,\tau;X)\,dX\,d\tau.\end{aligned} \tag{I.5}$$

At zero response age, E1 computes

$$\mathcal{R}(t,t;X) = \mathcal{R}_0(X) = \frac{x(4y-x)}{2\sqrt{y}(x+y)^{7/2}}. \tag{I.6}$$

The diagonal kernel is therefore independent of the observation time except through the moving source slice $\Theta(X,t)$ already displayed in (I.5).

Now use the pointwise decomposition $\zeta = [\zeta]_+ - [\zeta]_-$ for each real routing kernel ζ. The source factor is nonnegative because $\Theta = \varepsilon_\beta u^2$ with $\varepsilon_\beta \in \{0,1\}$. Thus both polarized pieces are nonnegative, and (I.5) becomes exactly

$$\dot a(t) = \mathcal{T}_\beta^{\mathrm{src}}(t) - \mathcal{W}_\beta^{\mathrm{src}}(t).$$

No ratio by a, a^+, Λ, or a common provenance tag has been taken. In the shadow $\varepsilon_\beta = 0$, the factor Θ vanishes before the routing kernels are polarized, so both accounts vanish identically. The identity therefore passes the source-provenance guard in raw form. □

Audit status: GREEN/COMPLETE.

I3. Raw cross-shear factorization. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 19.3.

Restatement of proposition 19.3. With

$$\Gamma := xu, \qquad C_\beta := \frac{\Theta + \frac{x}{2}\Theta_x}{y},$$

the off-diagonal term in the q equation is exactly

$$\boxed{4\varepsilon_\beta u v_y \Gamma_x = (4yv_y)C_\beta.}$$

For the fixed customer, passive transport gives $\Gamma \geq 0$, hence $\varepsilon_\beta u \geq 0$ in the open quadrant, and therefore

$$\boxed{[-(4yv_y)C_\beta]_+ = 4\varepsilon_\beta u[-v_y\Gamma_x]_+.}$$

Full rigorous proof. From the tagged-source definition $\Theta = \varepsilon_\beta u^2$,

$$\Theta_x = 2\varepsilon_\beta u u_x.$$

Consequently

$$\begin{aligned}\Theta + \frac{x}{2}\Theta_x &= \varepsilon_\beta u^2 + x\varepsilon_\beta u u_x \\ &= \varepsilon_\beta u(u + xu_x) \\ &= \varepsilon_\beta u \Gamma_x,\end{aligned} \tag{I.7}$$

because $\Gamma_x = (xu)_x = u + xu_x$. Dividing by $y > 0$ gives the exact raw identity

$$C_\beta = \frac{\varepsilon_\beta u \Gamma_x}{y}. \tag{I.8}$$

Multiplication by $4yv_y$ proves

$$(4yv_y)C_\beta = 4\varepsilon_\beta u v_y \Gamma_x.$$

This algebraic identity requires no sign assumption.

The positive-part polarization does require one, and the fixed customer supplies it. Initially,

$$\Gamma_0(x, y) = xu_0(x, y) = xyw_0(x, y) \geq 0$$

because the explicit w_0 in definition 5.1 is strictly positive in the open quadrant. Since $\mathcal{D}_{xy}\Gamma = 0$, passive transport preserves this sign along every material trajectory. Hence $\Gamma \geq 0$ for every classical time, and for $x > 0$,

$$u = \Gamma/x \geq 0.$$

Thus $4\varepsilon_\beta u \geq 0$. For every nonnegative scalar λ and real ζ, $[\lambda\zeta]_+ = \lambda[\zeta]_+$. Taking $\lambda = 4\varepsilon_\beta u$ and $\zeta = -v_y\Gamma_x$ yields

$$[-(4yv_y)C_\beta]_+ = 4\varepsilon_\beta u[-v_y\Gamma_x]_+. \tag{I.9}$$

The equality extends to the symmetry axes by continuity. Hence the adverse cross-shear source is not an independent forcing direction: it is the product of the single tagged source descendant $\varepsilon_\beta u$, the transported circulation gradient Γ_x, and the already named meridional first jet v_y. □

Audit status: GREEN/COMPLETE.

I4. Carrier-Hodge comparison. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 19.4.

Restatement of proposition 19.4. For

$$\mathfrak{r} = x^{2/3}y^{1/3},$$

one has

$$\boxed{K_H(x,y) \le \frac{2^{5/3}}{3^{5/2}}\frac{1}{\mathfrak{r}}.}$$

Therefore every nonnegative genealogy density $\mathcal{H}$ with finite carrier integral satisfies

$$\boxed{\frac{3}{8}\int K_H\mathcal{H}\,dX \le \mathfrak{c}_H\int \mathcal{H}\frac{dX}{\mathfrak{r}}, \qquad \mathfrak{c}_H := \frac{2^{5/3}}{8\,3^{3/2}}.}$$

Full rigorous proof. Write $s = y/x > 0$. Then

$$K_H(x,y) = \frac{\sqrt{s}}{x(1+s)^{5/2}}, \qquad \frac{1}{\mathfrak{r}} = \frac{1}{xs^{1/3}}.$$

Hence

$$K_H(x,y)\,\mathfrak{r} = \frac{s^{5/6}}{(1+s)^{5/2}} =: \varphi_H(s). \tag{I.10}$$

The logarithmic derivative is

$$\frac{\varphi_H'(s)}{\varphi_H(s)} = \frac{5}{6s} - \frac{5}{2(1+s)} = \frac{5(1-2s)}{6s(1+s)}. \tag{I.11}$$

Thus φ_H increases on $(0,1/2)$, decreases on $(1/2,\infty)$, and reaches its unique global maximum at $s = 1/2$. Direct substitution gives

$$\sup_{s>0}\varphi_H(s) = \frac{(1/2)^{5/6}}{(3/2)^{5/2}} = \frac{2^{5/3}}{3^{5/2}}. \tag{I.12}$$

Equations (I.10)–(I.12) prove the pointwise comparison.

If $\mathcal{H} \ge 0$, multiplication by $\mathcal{H}$ preserves the inequality. Integrating and multiplying by $3/8$ gives

$$\frac{3}{8}\int_{\mathcal{Q}} K_H\mathcal{H}\,dX \le \frac{3}{8}\frac{2^{5/3}}{3^{5/2}}\int_{\mathcal{Q}}\mathcal{H}\frac{dX}{\mathfrak{r}} = \frac{2^{5/3}}{8\,3^{3/2}}\int_{\mathcal{Q}}\mathcal{H}\frac{dX}{\mathfrak{r}}.$$

This proves the asserted constant $\mathfrak{c}_H$. The estimate is absolute and one-sided: it transfers finite carrier mass to finite Hodge capacity, but it contains no signed information and therefore cannot by itself imply a strict cancellation gap. □

Audit status: GREEN/COMPLETE.

I5. Fresh genealogy as a true Euler source-time truncation. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 19.5.

Restatement of proposition 19.5. Fix $X^\sharp \in \mathcal{Q}$, $t > 0$, and source time $\tau \in [0,t]$. Put

$$X(\tau) := \mathscr{X}_{\tau,t}(X^\sharp), \qquad J(t,\tau) := \det D\mathscr{X}_{t,\tau}(X(\tau)), \qquad \tau_h := (t-h)_+.$$

Then

$$H^{\pm}_{\mathrm{fr},h}(X^\sharp,t) = 2\int_{\tau_h}^{t} J(t,\tau)^{-1}[q(X(\tau),\tau)]_\pm\,d\tau$$

is a nonnegative source-time sub-genealogy. If $t \le h$, it equals $H^{\pm}$; if $t > h$,

$$H^{\pm}_{\mathrm{fr},h}(X^\sharp, t) = H^{\pm}(X^\sharp, t) - J(t, t-h)^{-1} H^{\pm}(X(t-h), t-h).$$

Consequently $H^{\pm} = H^{\pm}_{\mathrm{fr},h} + H^{\pm}_{\mathrm{old},h}$ with both cohorts nonnegative, and

$$\boxed{\mathcal{M}_{\beta,\mathrm{fr},h}(t) = \frac{3}{8}\int_{\mathcal{Q}} K_H \mathcal{C} H_{\mathrm{fr},h}\, dX.}$$

Full rigorous proof. The polarized genealogies obey the exact continuity equations

$$\partial_t H^{\pm} + \nabla\cdot(\mathbf{b} H^{\pm}) = 2[q]_{\pm}, \qquad H^{\pm}(\cdot, 0) = 0, \qquad \mathbf{b} = (-2xv, 2yg). \tag{I.13}$$

Let $X(\tau) = \mathscr{X}_{\tau,t}(X^\sharp)$ be the backward material trajectory ending at $X^\sharp$ at time t, and let

$$J(t,\tau) = \det D\mathscr{X}_{t,\tau}(X(\tau)).$$

For fixed source time τ, $J(t,\tau) > 0$ on every classical interval. The standard change-of-volume identity for the material flow gives

$$\frac{d}{d\xi} \det D\mathscr{X}_{\xi,\tau}(X(\tau)) = (\nabla\cdot\mathbf{b})(\mathscr{X}_{\xi,\tau}X(\tau), \xi) \det D\mathscr{X}_{\xi,\tau}(X(\tau)).$$

Multiplying (I.13) by this Jacobian and integrating along the characteristic therefore yields, for every $0 \le \tau \le t$,

$$H^{\pm}(X^\sharp, t) = J(t,\tau)^{-1} H^{\pm}(X(\tau), \tau) + 2\int_\tau^t J(t,\xi)^{-1}[q(X(\xi),\xi)]_{\pm}\, d\xi. \tag{I.14}$$

All terms are nonnegative because the source $[q]_{\pm}$ and the material Jacobian are nonnegative/positive.

If $t \le h$, take $\tau = 0$ in (I.14). Since $H^{\pm}(\cdot, 0) = 0$, the remaining integral is exactly $H^{\pm}_{\mathrm{fr},h}(X^\sharp, t)$. Hence

$$H^{\pm}_{\mathrm{fr},h} = H^{\pm} \qquad (0 < t \le h). \tag{I.15}$$

If $t > h$, take $\tau = t - h$ in (I.14) and rearrange:

$$H^{\pm}_{\mathrm{fr},h}(X^\sharp, t) = H^{\pm}(X^\sharp, t) - J(t, t-h)^{-1} H^{\pm}(X(t-h), t-h). \tag{I.16}$$

Define the second term on the right by

$$H^{\pm}_{\mathrm{old},h}(X^\sharp, t) := J(t, t-h)^{-1} H^{\pm}(X(t-h), t-h)$$

for $t > h$, and put $H^{\pm}_{\mathrm{old},h} = 0$ for $t \le h$. Then

$$\boxed{H^{\pm} = H^{\pm}_{\mathrm{fr},h} + H^{\pm}_{\mathrm{old},h}, \qquad H^{\pm}_{\mathrm{fr},h}, H^{\pm}_{\mathrm{old},h} \ge 0.} \tag{I.17}$$

Subtracting the two signs gives the corresponding signed decomposition for H.

It remains to identify the meridional payment. By proposition 10.2, the Hodge-transport part of the bank rate is linear in the genealogy:

$$\mathcal{M}_+ - \mathcal{M}_- = \frac{3}{8}\int_{\mathcal{Q}} K_H \mathcal{C}(H^+ - H^-)\, dX.$$

Insert (I.17). Linearity splits this exact Euler quantity into fresh and old source-time contributions. The fresh contribution is precisely

$$\boxed{\mathcal{M}_{\beta,\mathrm{fr},h}(t) = \frac{3}{8}\int_{\mathcal{Q}} K_H \mathcal{C}(H^+_{\mathrm{fr},h} - H^-_{\mathrm{fr},h})\, dX = \frac{3}{8}\int_{\mathcal{Q}} K_H \mathcal{C} H_{\mathrm{fr},h}\, dX.} \tag{I.18}$$

Thus the fresh field introduces neither a new PDE nor a new memory variable: it is exactly the restriction of the existing Euler Duhamel genealogy to source times $\xi \in [(t-h)_+, t]$. □

Audit status: GREEN/COMPLETE.

I6. No double charging. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 19.6.

> **Restatement of proposition 19.6.** Infinite fractional Hodge-dominance refreshes do not imply infinite new negative cross-shear creation. One physical source-sign conversion may be charged only once; a divergent refresh ledger cannot be identified with a divergent physical conversion bill without an additional Euler theorem; any cumulative contradiction must therefore identify genuinely new source-time work rather than a later relabeling of old genealogy.

Full rigorous proof. There are two logically separate points.

Step 1: the exact Euler genealogy carries a unique source-time coordinate. By I5, every polarized deposition entering the current genealogy has the form

$$2J(t,\tau;X^\sharp)^{-1}[q(X(\tau;X^\sharp,t),\tau)]_\pm\, d\tau. \tag{I.19}$$

For a fixed observation $(X^\sharp, t)$, the source-time integral contains each physical deposition time τ exactly once. A later Hodge-dominance refresh may select, freeze, or repartition a portion of the already existing positive genealogy, but it does not change $q(\cdot,\tau)$, add a second copy of the measure (I.19), or create a new source-time coordinate. The same deposited label may of course remain visible at many later observation times; that is evolution of its response kernel, not a new deposition event. Therefore a repeated bookkeeping use of the same deposited mass is not new Euler source work. Any physical conversion bill that is summed cumulatively must be indexed by disjoint source-time events (or by another disjoint physical parameter proved to represent new work).

Step 2: refresh Zeno does not force divergent adverse creation. The necessity of this distinction is already visible in an exact two-bank ledger countermodel. On $0 \le t < 1$ set

$$a^+(t) = 1, \qquad a^-(t) = t, \qquad a(t) = 1 - t, \tag{I.20}$$

and take adverse source rate $\mathcal{S}_-(t) = 1$ with all other source/rate terms zero. Then the exact bank equations hold and

$$\int_0^1 \mathcal{S}_-(t)\, dt = 1 < \infty, \qquad \frac{a^-(t)}{a^+(t)} = t \uparrow 1. \tag{I.21}$$

Nevertheless define

$$t_n := 1 - 2^{-n}, \qquad n = 0, 1, 2, \dots.$$

At time t_n, select a positive sub-bank of size $a(t_n)/2 = 2^{-n-1}$ and keep that selected amount fixed until the next contact. Since

$$a(t_{n+1}) = 2^{-n-1} = \frac{1}{2}a(t_n),$$

the selected bank reaches dominance contact exactly at t_{n+1}. Thus infinitely many fractional refreshes accumulate at $t = 1$ although total adverse creation is only one.

This countermodel is not asserted to solve Euler. It is a ledger-level falsification of the implication

$$\#\{\text{refreshes}\} = \infty \quad \Longrightarrow \quad \text{new adverse source work} = \infty.$$

Combined with the exact source-time representation in Step 1, it proves the claimed firewall: later refreshes may reuse already deposited genealogy for routing, but they cannot be charged as new physical source-sign conversions unless a separate Euler estimate proves genuinely new source-time work. □

Audit status: GREEN/COMPLETE.

I7. Strict-gap algebra. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 19.7.

> **Restatement of proposition 19.7.** Assume $a^+ > 0$ and $a = a^+ - a^-$ with $a^- \geq 0$. Then
>
> $$a \geq \pi_*(a^+ + a^-), \qquad \pi_* > 0,$$
>
> is equivalent to
>
> $$\boxed{\frac{a^-}{a^+} \leq \frac{1-\pi_*}{1+\pi_*} < 1.}$$
>
> Conversely, if $a^- \leq (1-\delta)a^+$ with $\delta \in (0,1)$, then
>
> $$\boxed{a \geq \frac{\delta}{2-\delta}(a^+ + a^-) > 0.}$$

Full rigorous proof. Put

$$\Sigma_H := a^+ + a^-.$$

The raw polarization inequality is

$$a^+ - a^- \geq \pi_*(a^+ + a^-).$$

Moving like terms to opposite sides gives

$$(1-\pi_*)a^+ \geq (1+\pi_*)a^-. \tag{I.22}$$

Since $a^+ > 0$ and $1 + \pi_* > 0$, division only by these ordinary positive quantities shows that (I.22) is equivalent to

$$\boxed{\frac{a^-}{a^+} \leq \frac{1-\pi_*}{1+\pi_*}.} \tag{I.23}$$

If the bound is realizable with nonnegative banks, automatically $\pi_* \leq 1$; for $0 < \pi_* \leq 1$ the right-hand side lies in $[0,1)$. Reversing the same algebra proves the converse implication, so the two formulations are exactly equivalent after the admission $a^+ > 0$.

Now assume

$$a^- \leq (1-\delta)a^+, \qquad 0 < \delta < 1. \tag{I.24}$$

Subtracting gives

$$a = a^+ - a^- \geq \delta a^+. \tag{I.25}$$

At the same time,

$$\Sigma_H = a^+ + a^- \leq (2-\delta)a^+. \tag{I.26}$$

Hence $a^+ \geq \Sigma_H/(2-\delta)$, and inserting this into (I.25) yields

$$\boxed{a \geq \frac{\delta}{2-\delta}\Sigma_H = \frac{\delta}{2-\delta}(a^+ + a^-) > 0.} \tag{I.27}$$

The constant is sharp at the extremal allowed ratio $a^- = (1-\delta)a^+$. Equivalently, for $0 < \pi_* < 1$ the exact parameter conversion is

$$\boxed{\pi_* = \frac{\delta}{2-\delta}, \qquad \delta = \frac{2\pi_*}{1+\pi_*}.} \tag{I.28}$$

No division by the net compression a is used anywhere in the argument. □

Audit status: GREEN/COMPLETE.

Appendix J. Falsification firewalls

This section contains J1–J4. These are no-go statements: they certify that four apparently attractive closure mechanisms are logically insufficient, while leaving the positive two-time Hodge-response program intact.

J1. Finite local action alone cannot enforce the cancellation gap. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 20.1.

> **Restatement of proposition 20.1.** There exists a tagged one-cell reduced system with variables $q(t)$, $C_\beta^{\mathrm{cell}}(t)$, and $H(t)$ satisfying
>
> $$\dot q = -HC_\beta^{\mathrm{cell}}, \qquad \dot C_\beta^{\mathrm{cell}} = \frac{1}{4}Hq, \qquad \dot H = 2q,$$
>
> with
>
> $$q(0) = 1, \qquad C_\beta^{\mathrm{cell}}(0) = H(0) = 0,$$
>
> which obeys the exact local source-gradient/slope/mixed-jet algebra and has finite cross-shear/restart action. Define
>
> $$\dot H^+ = 2[q]_+, \qquad \dot H^- = 2[q]_-, \qquad H^\pm(0) = 0, \qquad a^\pm := \frac{1}{2}H^\pm.$$
>
> For these banks one reaches a finite time at which
>
> $$a^+ \to \sqrt{2}, \qquad a^- \to \sqrt{2}, \qquad a = a^+ - a^- \to 0.$$
>
> Therefore no strict global inequality $a^- \le (1-\delta)a^+$ can follow from local tagged ODE structure plus finiteness of raw action alone.

Proof. Step 1: a tagged cell satisfying the exact local algebra.
Take one stationary reduced material label and prescribe

$$x = y = u = 1, \qquad v = g = v_x = g_y = 0, \qquad \mathfrak{T} = 0, \qquad v_y = g_x = -\frac{1}{4}H. \tag{J.1}$$

Then

$$-2(v_y + g_x) = H,$$

and the mixed-jet identities give

$$4yv_y = -H, \qquad -xg_x = \frac{1}{4}H.$$

The diagonal coefficients in the exact F1 source-gradient system vanish, so the first two rows reduce exactly to

$$\dot q = -HC_\beta^{\mathrm{cell}}, \qquad \dot C_\beta^{\mathrm{cell}} = \frac{1}{4}Hq.$$

The third equation $\dot H = 2q$ is the local vorticity-source law with $g = 0$. Thus (J.1) realizes the displayed cell system without introducing an unrelated forcing.

The slope/circulation-gradient relay is compatible with the same cell; p_{cell} is a reduced-model proxy and is not the protected physical field $p = u_y$. Put

$$p_{\mathrm{cell}} := \frac{q}{2}, \qquad \Gamma_x := C_\beta^{\mathrm{cell}}. \tag{J.2}$$

Because $g = g_y = 0$ and $x = y = 1$,

$$\dot{p}_{\mathrm{cell}} = -\frac{1}{2} H C_\beta^{\mathrm{cell}} = 2 v_y \Gamma_x, \qquad \dot{\Gamma}_x = \frac{1}{4} H q = -2 g_x p_{\mathrm{cell}}.$$

Hence the source-gradient pair, the slope/circulation-gradient pair, the Hodge source law, and the mixed first jets are mutually consistent at the reduced-cell level. This is a falsification model for logical sufficiency; it is not asserted to be a whole-space Euler solution.

Step 2: exact invariants and finite half-turn time.

Direct differentiation gives

$$\boxed{q^2 + 4(C_\beta^{\mathrm{cell}})^2 = 1, \qquad C_\beta^{\mathrm{cell}} = \frac{H^2}{16}.} \tag{J.3}$$

Indeed

$$\frac{d}{dt}\big(q^2 + 4(C_\beta^{\mathrm{cell}})^2\big) = 2q(-H C_\beta^{\mathrm{cell}}) + 8 C_\beta^{\mathrm{cell}} \frac{Hq}{4} = 0,$$

and

$$\frac{d}{dt}\left(C_\beta^{\mathrm{cell}} - \frac{H^2}{16}\right) = \frac{1}{4} H q - \frac{1}{8} H (2q) = 0.$$

Parameterize the increasing branch by a statement-local angle ϑ:

$$q = \cos\vartheta, \qquad C_\beta^{\mathrm{cell}} = \frac{1}{2}\sin\vartheta, \qquad H = 2\dot{\vartheta}. \tag{J.4}$$

Then

$$\ddot{\vartheta} = \cos\vartheta, \qquad \vartheta(0) = \dot{\vartheta}(0) = 0, \tag{J.5}$$

and multiplication by $\dot{\vartheta}$ yields

$$\boxed{\frac{1}{2}\dot{\vartheta}^{\,2} = \sin\vartheta.} \tag{J.6}$$

Therefore the branch reaches $\vartheta = \pi$ in the finite time

$$T_* = \frac{1}{\sqrt{2}} \int_0^\pi (\sin\vartheta)^{-1/2}\, d\vartheta < \infty. \tag{J.7}$$

At T_* one has

$$q = -1, \qquad C_\beta^{\mathrm{cell}} = 0, \qquad H = 0.$$

Thus the local tagged source direction has completed a positive-to-negative half-turn.

Step 3: the raw actions are finite.

On $[0, T_*]$, $H = 2\dot{\vartheta} \geq 0$, so

$$\int_0^{T_*} H\, dt = 2\pi, \qquad \int_0^{T_*} \big(|4 y v_y| + |x g_x|\big)\, dt = \frac{5\pi}{2}. \tag{J.8}$$

Moreover $v_y \Gamma_x = -H C_\beta^{\mathrm{cell}}/4 \leq 0$, whence

$$\int_0^{T_*} [-v_y \Gamma_x]_+\, dt = \frac{1}{4}\int_0^\pi \sin\vartheta\, d\vartheta = \frac{1}{2}. \tag{J.9}$$

Thus both the local restart/first-jet action and the tagged negative cross-shear creation are finite.

Step 4: the two genealogy banks reach cancellation lock.

Since

$$\frac{d}{dt}(H^+ - H^-) = 2([q]_+ - [q]_-) = 2q = \dot{H}$$

and all three quantities vanish initially,

$$H = H^+ - H^-.$$

Hence $a := a^+ - a^- = H/2 = \dot{\vartheta}$. Using $dt = d\vartheta/\sqrt{2\sin\vartheta}$ from (J.6),

$$a^+(T_*) = \int_0^{\pi/2} \frac{\cos\vartheta}{\sqrt{2\sin\vartheta}}\, d\vartheta = \sqrt{2}, \tag{J.10}$$

$$a^-(T_*) = \int_{\pi/2}^{\pi} \frac{-\cos\vartheta}{\sqrt{2\sin\vartheta}}\, d\vartheta = \sqrt{2}. \tag{J.11}$$

Consequently

$$\boxed{a^+(t) \to \sqrt{2}, \quad a^-(t) \to \sqrt{2}, \quad \frac{a^-(t)}{a^+(t)} \to 1, \quad a(t) \to 0 \qquad (t \uparrow T_*).} \tag{J.12}$$

For every fixed $\delta > 0$, the inequality $a^- \le (1-\delta)a^+$ therefore fails sufficiently near T_*. The missing input is precisely the whole-space spatial Hodge/Biot–Savart coupling that the one-cell model intentionally suppresses. □

Audit status: GREEN/COMPLETE.

J2. Frozen Hodge geometry alone has no uniform gap. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 20.2.

> **Restatement of proposition 20.2.** Fix $s := y/x$. For every $\varepsilon > 0$ there exist smooth rapidly decaying nonnegative H^+, H^- with
>
> $$H^+(x, sx) = sx + O(x^2), \qquad H^-(x, sx) = O(x^2),$$
>
> with H^- supported away from the equator, such that
>
> $$a^\pm := \frac{3}{8}\int_{\mathcal{Q}} K_H H^\pm \, dX$$
>
> satisfy $a^-/a^+ > 1 - \varepsilon$.

Proof. Choose a nonnegative C^∞ function χ_-, not identically zero, with

$$\operatorname{supp}\chi_- \subset (1, \infty).$$

Define

$$H^+(x,y) := ye^{-(x+y)}, \qquad H^-(x,y) := \kappa_{\text{snap}}\, y^2 \chi_-(y) e^{-(x+y)}. \tag{J.13}$$

Both functions are smooth, nonnegative, and rapidly decaying on the squared quadrant. For every fixed $s > 0$,

$$H^+(x, sx) = sxe^{-(1+s)x} = sx + O(x^2).$$

Because $\chi_-(sx) = 0$ for all sufficiently small x,

$$H^-(x, sx) = 0 = O(x^2).$$

Thus the negative snapshot is separated a fixed positive distance from the equator while the positive snapshot has the required first apex jet.

Set

$$A_+ := \frac{3}{8}\int_{\mathcal{Q}} K_H H^+ \, dX, \qquad A_-^{(1)} := \frac{3}{8}\int_{\mathcal{Q}} K_H y^2 \chi_-(y) e^{-(x+y)} \, dX. \tag{J.14}$$

The kernel K_H is strictly positive in the interior. Hence $A_+ > 0$ and $A_-^{(1)} > 0$; both are finite by the rapid decay and the locally integrable Hodge kernel. Given $\varepsilon > 0$, choose

$$0 < \eta < \min\{\varepsilon, 1\}, \qquad \kappa_{\mathrm{snap}} := (1-\eta)\frac{A_+}{A_-^{(1)}}. \tag{J.15}$$

Then

$$\boxed{a^+ = A_+, \qquad a^- = (1-\eta)A_+, \qquad \frac{a^-}{a^+} = 1-\eta > 1-\varepsilon.} \tag{J.16}$$

This strict choice of η repairs the harmless equality ambiguity in the historical compressed proof.

Finally, the signed field $H := H^+ - H^-$ is smooth and rapidly decaying, so the exact seven-dimensional lift of proposition 5.5 reconstructs a smooth decaying meridional field satisfying the whole-space Hodge relation. Thus the counterexample retains the genuine frozen Hodge/Biot–Savart geometry; what it deliberately omits is the fixed-customer source history tying $H^\pm$ to earlier Euler deposition. Therefore frozen-time geometry and the apex sign orders alone cannot yield a uniform strict cancellation gap. □

Audit status: GREEN/COMPLETE.

J3. Infinite fractional refresh does not imply infinite physical conversion. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 20.3.

> **Restatement of proposition 20.3.** On $0 \le t < 1$, a two-bank bookkeeping model can have
>
> $$a^+(t) = 1, \qquad a^-(t) = t, \qquad a(t) = 1 - t,$$
>
> with finite total adverse creation but infinitely many fractional Hodge-dominance refreshes as $t \uparrow 1$. Therefore one finite physical conversion may not be charged repeatedly to infinitely many bookkeeping refreshes.

Proof. Define

$$a^+(t) := 1, \qquad a^-(t) := t, \qquad a(t) := 1 - t, \qquad 0 \le t < 1. \tag{J.17}$$

The total adverse creation is finite:

$$\int_0^1 \frac{d}{dt} a^-(t) \, dt = 1. \tag{J.18}$$

Nevertheless the net bank passes through infinitely many geometric scales. At

$$t_n := 1 - 2^{-n}, \qquad n \ge 1, \tag{J.19}$$

one has

$$a(t_n) = 2^{-n}.$$

Thus any bookkeeping protocol that records a new fractional Hodge-dominance contact whenever the surviving net bank drops by another fixed factor—for example from $2^{-(n-1)}$ to 2^{-n}—records infinitely many refresh contacts as $n \to \infty$. No new adverse deposition is created by this relabeling: the entire adverse bank is still generated by the single integrable rate $\dot{a}^- = 1$ in (J.18).

This is exactly the logical distinction made rigorous by I5–I6. A deposition event is indexed once by its source time τ in the true Duhamel measure; later observations or refresh selections evolve or repartition already deposited genealogy. They do not create another copy of the same source-time event. Therefore an infinite refresh count cannot be converted into an infinite physical cross-shear bill without an additional Euler theorem proving genuinely new adverse deposition at the later contacts. The model is a ledger counterexample, not an Euler solution. □

Audit status: GREEN/COMPLETE.

J4. Pointwise reproduction number is too strong. *Mutual pointers:* grand-finale entry / short proof/status / main statement proposition 20.4.

> **Restatement of proposition 20.4.** In the constant-rate affine conveyor, a label may leave the compact rotor region while a pointwise parent-to-child reproduction multiplier grows exponentially. The same label's final Hodge visibility decays faster, so the physical Hodge-weighted product still decays. Consequently final Hodge visibility must remain inside the causal estimate; a global pointwise reproduction bound < 1 is not the correct target.

Proof. Fix $a_0 > 0$ and consider the constant-rate affine conveyor

$$v = a_0, \qquad g = 2a_0, \qquad x(\tau) = x_0 e^{-2a_0\tau}, \qquad y(\tau) = y_0 e^{4a_0\tau}, \tag{J.20}$$

with $x_0, y_0 > 0$. Put $s_0 := y_0/x_0$. Since the affine first derivatives vanish, the source-gradient pair itself is constant along the label. Choose $C_{\beta,0} > 0$. Then

$$\widehat{C}_\beta(\tau) = 2\sqrt{s_0}\, C_{\beta,0} e^{3a_0\tau}, \qquad \sqrt{x(\tau)y(\tau)}\,\widehat{C}_\beta(\tau) = 2y_0 C_{\beta,0} e^{4a_0\tau}. \tag{J.21}$$

The exact residence multiplier from H4 becomes

$$\mathfrak{R}_q(t,\tau) = \frac{e^{2a_0(t-\tau)} - 1}{2a_0}. \tag{J.22}$$

Fix a finite source-time interval $0 \le \tau_1 < \tau_2$ during which the label lies inside whatever compact rotor gate is being tested. Define the corresponding factored pointwise exposure by

$$\mathfrak{m}^+_{\mathrm{pt}}(t) := \int_{\tau_1}^{\tau_2} \sqrt{x(\tau)y(\tau)}\,\widehat{C}_\beta(\tau)\,\mathfrak{R}_q(t,\tau)\,d\tau, \qquad t > \tau_2. \tag{J.23}$$

Using (J.21)–(J.22),

$$\begin{aligned}\mathfrak{m}^+_{\mathrm{pt}}(t) &= \frac{y_0 C_{\beta,0}}{a_0}\int_{\tau_1}^{\tau_2}\left(e^{2a_0 t + 2a_0\tau} - e^{4a_0\tau}\right)d\tau \\ &= \frac{y_0 C_{\beta,0}}{a_0}\left[\frac{e^{2a_0 t}}{2a_0}\left(e^{2a_0\tau_2} - e^{2a_0\tau_1}\right) - \frac{1}{4a_0}\left(e^{4a_0\tau_2} - e^{4a_0\tau_1}\right)\right].\end{aligned} \tag{J.24}$$

Hence

$$\boxed{\mathfrak{m}^+_{\mathrm{pt}}(t) \longrightarrow +\infty \qquad (t \to \infty),} \tag{J.25}$$

indeed at rate $e^{2a_0 t}$.

The same old label is nevertheless becoming Hodge-invisible. Its final kernel is

$$K_H(x(t), y(t)) = \frac{x_0\sqrt{y_0}}{(x_0 e^{-2a_0 t} + y_0 e^{4a_0 t})^{5/2}} = O(e^{-10a_0 t}). \tag{J.26}$$

Combining (J.24) and (J.26) gives

$$\boxed{K_H(x(t),y(t))\,\mathfrak{m}^+_{\mathrm{pt}}(t) = O(e^{-8a_0 t}) \longrightarrow 0.} \tag{J.27}$$

Thus taking a pointwise supremum after factoring out final Hodge visibility destroys the very old-age discount that makes the physical response harmless. A global requirement such as a pointwise reproduction number < 1 is therefore strictly stronger than the physically relevant causal Hodge-weighted estimate and is not the correct closure target. □

Audit status: GREEN/COMPLETE.

Appendix K. Permanent O-ledger: O1–O6.1 closed, O6.2 sole open slot

The permanent O-ledger navigation labels do not move. O1–O3 and the inherited-launch source-angle audit O6.1 are closed; O4 is therefore closed; O5 is closed downstream from O4+O6.2+M8; and O6.2 is the sole reserved open mathematical target.

O1. Large-collapse-age physical-bank retirement. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 21.3.

Proof. **Step 1: localize the exact signed/absolute response in source time.** For a measurable set $\mathscr{E}_{\mathrm{src}} \subset [0,t]$, pull the exact response to the fixed initial labels exactly as in G4 and put

$$\mathcal{P}_{\mathscr{E}_{\mathrm{src}}}(t) := \frac{3}{4}\int_{\mathscr{E}_{\mathrm{src}}}\int_{\mathcal{Q}} q(\mathscr{X}_\tau X_0,\tau)\mathcal{J}_\tau(X_0)\mathcal{W}_t(X_0)\,dX_0 d\tau,$$
$$\mathcal{V}_{\mathscr{E}_{\mathrm{src}}}(t) := \frac{3}{4}\int_{\mathscr{E}_{\mathrm{src}}}\int_{\mathcal{Q}} |q(\mathscr{X}_\tau X_0,\tau)|\mathcal{J}_\tau(X_0)\mathcal{W}_t(X_0)\,dX_0 d\tau.$$

Then

$$a^+_{\mathscr{E}_{\mathrm{src}}} = \frac{\mathcal{V}_{\mathscr{E}_{\mathrm{src}}} + \mathcal{P}_{\mathscr{E}_{\mathrm{src}}}}{2}, \qquad a^-_{\mathscr{E}_{\mathrm{src}}} = \frac{\mathcal{V}_{\mathscr{E}_{\mathrm{src}}} - \mathcal{P}_{\mathscr{E}_{\mathrm{src}}}}{2}.$$

For

$$I_{\mathscr{E}_{\mathrm{src}}}(t) := \int_{\mathscr{E}_{\mathrm{src}}} e^{2\Lambda(\tau)}\,d\tau, \qquad W^{\mathrm{aff}}_t(X_0) := K_H(\mathscr{X}^{\mathrm{aff}}_{\Lambda(t)}X_0),$$

define

$$\mathcal{P}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}} := \frac{3}{4} I_{\mathscr{E}_{\mathrm{src}}}(t)\int_{\mathcal{Q}} q_0(X_0) W^{\mathrm{aff}}_t(X_0)\,dX_0,$$
$$\mathcal{V}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}} := \frac{3}{4} I_{\mathscr{E}_{\mathrm{src}}}(t)\int_{\mathcal{Q}} |q_0(X_0)| W^{\mathrm{aff}}_t(X_0)\,dX_0.$$

Step 2: the audited G4 comparison localizes without loss. Every estimate in G4 Steps 2–5 is pointwise in source time τ. Restricting those pointwise inequalities to $\mathscr{E}_{\mathrm{src}}$ before integration therefore gives

$$\boxed{|\mathcal{P}_{\mathscr{E}_{\mathrm{src}}} - \mathcal{P}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}| + |\mathcal{V}_{\mathscr{E}_{\mathrm{src}}} - \mathcal{V}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}| \le \eta_M \mathcal{V}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}},} \tag{K.1}$$

with the same $\eta_M = 2\mathfrak{C}^*_Z(e^{8M}-1)$ as in G4. No new comparison constant or unnamed source-gradient stock is introduced.

Step 3: every affine source-time subset has the same bank ratio. The scalar factor $I_{\mathscr{E}_{\mathrm{src}}}(t)$ is common to the affine signed and absolute responses. Consequently, whenever $I_{\mathscr{E}_{\mathrm{src}}}(t) > 0$,

$$\frac{a^-_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}(t)}{a^+_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}(t)} = \frac{I_-(\Lambda(t))}{I_+(\Lambda(t))} =: \rho_{\mathrm{aff}}(\Lambda(t)). \tag{K.2}$$

The audited E4 asymptotic

$$\rho_{\mathrm{aff}}(\ell_{\mathrm{age}}) = \frac{c_-}{6\kappa\ell_{\mathrm{age}}} + o(\ell_{\mathrm{age}}^{-1})$$

and compactness on bounded ℓ_{age} imply that a fixed $\mathfrak{c}_{\mathrm{aff}} < \infty$ exists with

$$\rho_{\mathrm{aff}}(\ell_{\mathrm{age}}) \le \frac{\mathfrak{c}_{\mathrm{aff}}}{1+\ell_{\mathrm{age}}} \qquad (\ell_{\mathrm{age}} \ge 0). \tag{K.3}$$

Step 4: specialize to actual large collapse age. Take $\mathscr{E}_{\mathrm{src}} = \mathscr{E}_L(t)$. If this set is empty, there is nothing to prove. Otherwise positive compression makes Λ nondecreasing and gives $\Lambda(t) \ge L$. Hence (K.2)–(K.3) yield

$$a^-_{\mathscr{E}_L,\mathrm{aff}} \le \varepsilon_L a^+_{\mathscr{E}_L,\mathrm{aff}}, \qquad \varepsilon_L := \frac{\mathfrak{c}_{\mathrm{aff}}}{1+L}.$$

By (K.1), each of the two true sub-banks differs from its affine counterpart by at most $\eta_M \mathcal{V}_{\mathscr{E}_L,\mathrm{aff}}/2$. Therefore

$$\begin{aligned} a^-_{\mathscr{E}_L} &\le a^-_{\mathscr{E}_L,\mathrm{aff}} + \frac{1}{2}\eta_M \mathcal{V}_{\mathscr{E}_L,\mathrm{aff}} \\ &\le \varepsilon_L a^+_{\mathscr{E}_L,\mathrm{aff}} + \frac{1}{2}\eta_M \mathcal{V}_{\mathscr{E}_L,\mathrm{aff}} \\ &\le \varepsilon_L a^+_{\mathscr{E}_L} + \frac{1+\varepsilon_L}{2}\eta_M \mathcal{V}_{\mathscr{E}_L,\mathrm{aff}} \\ &\le \varepsilon_L a^+(t) + \mathfrak{D}_{\mathrm{cent,L}}(t). \end{aligned}$$

This is exactly (21.4).

Step 5: accounting legality and the affine-falsification firewall. The error term is not a new invoice: it is precisely the G4 centered-deformation comparison error restricted to the large-age source-time set. If the centered deformation remains zero, the bill vanishes and the estimate reduces to the exact affine $O((1+L)^{-1})$ retirement. If centered deformation is substantial, G5 routes first polarization loss to the already certified deformation ticket $\mathcal{K}_{\mathrm{cent}}$ and then to $\mathcal{L}_{\mathrm{pol}}$. No factorized H6 primary exposure is used. This is essential: the falsification and scalar-cashier tests show that a cumulative positive-part primary majorant can fail even a cashier-enhanced affine stress test, whereas the physical source-time bank used here passes the exact affine benchmark. □

Audit status: GREEN/COMPLETE.

O2. Bounded-collapse-age physical-bank return. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 21.4.

Restatement of theorem 21.4. On a positive-compression branch, let

$$\mathscr{E}_{<L}(t) = \{\tau \in [0,t] : 0 \le \Lambda(t) - \Lambda(\tau) < L\}$$

and define the exact physical source-time banks $a^{\pm}_{<L}$ by (21.6). There exist fixed constants $M_{\mathrm{O2}} > 0$ and $0 \le \theta_{\mathrm{O2}} < 1$ such that either

$$\mathcal{K}_{\mathrm{cent}}([0,t]) \le M_{\mathrm{O2}} \quad \Longrightarrow \quad a^-_{<L}(t) \le \theta_{\mathrm{O2}} a^+_{<L}(t) \le \theta_{\mathrm{O2}} a^+(t),$$

or else the branch has already paid the definite centered-deformation ticket

$$\int_0^t \mathcal{L}_{\mathrm{pol}}(\tau)\,d\tau > M_{\mathrm{O2}}/3.$$

Proof. **Step 1: localize the affine polarization to the bounded-age set.** For a measurable source-time set $\mathscr{E}_{\mathrm{src}} \subset [0,t]$, let $\mathcal{P}_{\mathscr{E}_{\mathrm{src}}}, \mathcal{V}_{\mathscr{E}_{\mathrm{src}}}$ and $\mathcal{P}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}, \mathcal{V}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}$ be the localized signed and absolute responses used in the O1 proof. The affine source-time scalar

$$I_{\mathscr{E}_{\mathrm{src}}}(t) := \int_{\mathscr{E}_{\mathrm{src}}} e^{2\Lambda(\tau)}\,d\tau$$

multiplies both affine signs. Therefore, whenever $I_{\mathscr{E}_{\mathrm{src}}}(t) > 0$,

$$\frac{a^-_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}(t)}{a^+_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}(t)} = \upsilon^{\mathrm{aff}}_-(\Lambda(t)) \le \upsilon^*_{\mathrm{aff}} < 1.$$

Equivalently,

$$\mathcal{P}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}(t) \ge \pi^*_{\mathrm{aff}} \mathcal{V}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}(t). \tag{K.4}$$

This identity holds for every measurable $\mathscr{E}_{\mathrm{src}}$; the bounded-age condition has not yet been used.

Step 2: localize G4. The G4 integrand comparison is pointwise in source time. Hence, exactly as in the O1 proof,

$$|\mathcal{P}_{\mathscr{E}_{\mathrm{src}}} - \mathcal{P}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}| + |\mathcal{V}_{\mathscr{E}_{\mathrm{src}}} - \mathcal{V}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}| \le \eta_M \mathcal{V}_{\mathscr{E}_{\mathrm{src}},\mathrm{aff}}, \qquad \eta_M = 2\mathfrak{C}^*_Z(e^{8M}-1), \tag{K.5}$$

where $M = \mathcal{K}_{\mathrm{cent}}([0,t])$.

Take $\mathscr{E}_{\mathrm{src}} = \mathscr{E}_{<L}(t)$. If this set has zero source-time measure, the desired bank bound is trivial. Otherwise, under $M \le M_{\mathrm{O2}}$, (K.4)–(K.5) give

$$\mathcal{P}_{<L} \ge (\pi^*_{\mathrm{aff}} - \eta_{M_{\mathrm{O2}}})\mathcal{V}_{<L,\mathrm{aff}}, \qquad \mathcal{V}_{<L} \le (1+\eta_{M_{\mathrm{O2}}})\mathcal{V}_{<L,\mathrm{aff}}.$$

Thus

$$\mathcal{P}_{<L} \ge \pi_{\mathrm{O2}} \mathcal{V}_{<L},$$

with π_{O2} from (21.7). The exact I7 algebra now yields

$$a^-_{<L} \le \frac{1-\pi_{\mathrm{O2}}}{1+\pi_{\mathrm{O2}}} a^+_{<L} = \theta_{\mathrm{O2}} a^+_{<L} \le \theta_{\mathrm{O2}} a^+(t),$$

which is (21.8). Note that the constant is independent of L: bounded collapse age selects the complementary physical source-time bank, while the strict margin is supplied by the fixed-seed affine polarization.

Step 3: failure pays an existing deformation ticket. If $M > M_{\mathrm{O2}}$, then a definite centered-deformation amount has already been spent. The audited G5 star-hull routing holds for every time:

$$\mathcal{K}_{\mathrm{cent}}([0,t]) \le 3 \int_0^t \mathcal{L}_{\mathrm{pol}}(\tau)\,d\tau.$$

Consequently

$$\int_0^t \mathcal{L}_{\mathrm{pol}}(\tau)\,d\tau > \frac{M_{\mathrm{O2}}}{3},$$

which is (21.9). This is an already named source-gradient/star-hull/tangent deformation payment, not a new O2 account.

Step 4: relation to the mesoscopic rotor. F8 remains useful if one wants to localize a substantial adverse conversion into a compact box. It is not needed for the present closure: source-time-localized G4 already returns the entire bounded-age physical negative bank either

to a strict polarization margin or to a definite centered-deformation ticket. This avoids the H6 positive-part overcount falsified by the affine benchmark. □

Audit status: GREEN/COMPLETE.

O3. Physical-bank consolidation or paid centered deformation. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 21.5.

Restatement of theorem 21.5. On a positive-compression branch choose fixed L_{O3} and M_{O3} as in (21.10)–(21.11). Then either

$$\boxed{a^-(t) \le \theta_{\mathrm{O3}} a^+(t), \qquad \theta_{\mathrm{O3}} < 1,}$$

or

$$\boxed{\int_0^t \mathcal{L}_{\mathrm{pol}}(\xi)\, d\xi > M_{\mathrm{O3}}/3.}$$

Proof. Fix $L = L_{\mathrm{O3}}$ and abbreviate

$$p_L := a^+_{\ge L}(t), \qquad n_L := a^-_{\ge L}(t), \qquad p_Y := a^+_{<L}(t), \qquad n_Y := a^-_{<L}(t).$$

Since $a \ge 0$ on $[0,t]$, the relative collapse age is nonnegative and the O1/O2 source-time sets partition $[0,t]$. Hence

$$\boxed{a^+ = p_L + p_Y, \qquad a^- = n_L + n_Y.} \tag{K.6}$$

Assume first $\mathcal{K}_{\mathrm{cent}}([0,t]) \le M_{\mathrm{O3}}$. Since $M_{\mathrm{O3}} \le M_{\mathrm{O2}}$, O2 gives

$$n_Y \le \theta_{\mathrm{O2}} p_Y \le \theta_{\mathrm{O2}} a^+. \tag{K.7}$$

O1 gives

$$n_L \le \varepsilon_L a^+ + \frac{1+\varepsilon_L}{2} \eta_M \mathcal{V}_{\mathscr{E}_L,\mathrm{aff}}, \tag{K.8}$$

where $M = \mathcal{K}_{\mathrm{cent}}([0,t])$ and $\eta_M \le \eta_{\mathrm{O3}}$. The localized G4 comparison also gives

$$|\mathcal{V}_{\mathscr{E}_L} - \mathcal{V}_{\mathscr{E}_L,\mathrm{aff}}| \le \eta_M \mathcal{V}_{\mathscr{E}_L,\mathrm{aff}}.$$

Since $\eta_M < 1$,

$$\mathcal{V}_{\mathscr{E}_L,\mathrm{aff}} \le \frac{\mathcal{V}_{\mathscr{E}_L}}{1-\eta_M} \le \frac{p_L + n_L}{1-\eta_{\mathrm{O3}}}. \tag{K.9}$$

Therefore

$$n_L \le \varepsilon_L a^+ + \delta_{\mathrm{cent}}(p_L + n_L) \le \varepsilon_L a^+ + \delta_{\mathrm{cent}}(a^+ + n_L),$$

so

$$n_L \le \frac{\varepsilon_L + \delta_{\mathrm{cent}}}{1-\delta_{\mathrm{cent}}} a^+. \tag{K.10}$$

Combining (K.6), (K.7), and (K.10) yields the first alternative with θ_{O3} from (21.12).

For strictness, let $\Delta_{\mathrm{O3}} = 1 - \theta_{\mathrm{O2}}$. The parameter choices give $\varepsilon_L, \delta_{\mathrm{cent}} \le \Delta_{\mathrm{O3}}/8$. Since $0 < \Delta_{\mathrm{O3}} \le 1$,

$$\frac{\varepsilon_L + \delta_{\mathrm{cent}}}{1-\delta_{\mathrm{cent}}} \le \frac{\Delta_{\mathrm{O3}}/4}{1-\Delta_{\mathrm{O3}}/8} \le \frac{2}{7}\Delta_{\mathrm{O3}} < \Delta_{\mathrm{O3}},$$

so $\theta_{\mathrm{O3}} < 1$.

If instead $\mathcal{K}_{\mathrm{cent}}([0,t]) > M_{\mathrm{O3}}$, the audited G5 routing

$$\mathcal{K}_{\mathrm{cent}}([0,t]) \le 3 \int_0^t \mathcal{L}_{\mathrm{pol}}(\xi)\, d\xi$$

yields (21.14). This is the existing source-gradient/star-hull/tangent deformation payment, not a new account. □

Audit status: GREEN/COMPLETE.

O4. Grand cancellation gap. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 21.6.

Proof. **Step 1: uniform constants.** The proved O3 theorem supplies fixed constants

$$M_{\mathrm{O3}} > 0, \qquad 0 \le \theta_{\mathrm{O3}} < 1,$$

depending only on the fixed customer and already funded O1–O3 data. O6.1 supplies

$$\delta_6 \in (0,1)$$

depending only on the fixed customer and admitted finite-action ball. Put

$$\boxed{\delta_* := \min\{1-\theta_{\mathrm{O3}}, \delta_6\} > 0.} \tag{K.11}$$

Step 2: bootstrap positive compression rather than assuming it. The fixed customer is active positive-opening:

$$a(0) = 0, \qquad \dot a(0) > 0$$

by proposition 7.2; hence $a(t) > 0$ for all sufficiently small $t > 0$.

Moreover proposition 11.1 gives

$$\boxed{a^+(t) > 0 \qquad (t > 0).} \tag{K.12}$$

Suppose for contradiction that there is a first later zero

$$T_0 > 0, \qquad a(t) > 0 \ (0 < t < T_0), \qquad a(T_0) = 0.$$

Then $a(\xi) \ge 0$ on $[0, T_0]$, so the proved O3 theorem is applicable up to and including T_0.

Write

$$K(t) := \mathcal{K}_{\mathrm{cent}}([0,t]).$$

The function K is continuous and nondecreasing.

If $K(T_0) \le M_{\mathrm{O3}}$, O3 gives

$$a^-(T_0) \le \theta_{\mathrm{O3}} a^+(T_0),$$

hence

$$a(T_0) \ge (1-\theta_{\mathrm{O3}}) a^+(T_0) > 0$$

by (K.12), a contradiction.

If $K(T_0) > M_{\mathrm{O3}}$, define

$$t_d := \inf\{t > 0 : K(t) > M_{\mathrm{O3}}\}.$$

Continuity gives

$$0 < t_d < T_0, \qquad K(t_d) = M_{\mathrm{O3}}.$$

Thus O3 still supplies the strict gap at the threshold state. By O6.1,

$$a^-(t) \le (1-\delta_6) a^+(t) \qquad (t_d \le t \le T_0).$$

At T_0,

$$a(T_0) \ge \delta_6 a^+(T_0) > 0,$$

again a contradiction.

Therefore

$$\boxed{a(t) > 0 \qquad (t > 0)} \tag{K.13}$$

throughout the admitted classical branch.

Step 3: cover the full branch with no endpoint gap. If $K(t) \le M_{\mathrm{O3}}$ for the whole branch, O3 gives

$$a^-(t) \le \theta_{\mathrm{O3}} a^+(t) \le (1-\delta_*)a^+(t)$$

for all t.

Otherwise the threshold time t_d is well-defined. Since

$$K(t) \le M_{\mathrm{O3}} \qquad (0 \le t \le t_d),$$

O3 gives

$$a^-(t) \le \theta_{\mathrm{O3}} a^+(t) \le (1-\delta_*)a^+(t) \qquad (0 \le t \le t_d). \tag{K.14}$$

The equality $K(t_d) = M_{\mathrm{O3}}$ is the endpoint fact that keeps the O3 estimate valid at the threshold.

For $t \ge t_d$, O6.1 gives

$$a^-(t) \le (1-\delta_6)a^+(t) \le (1-\delta_*)a^+(t). \tag{K.15}$$

Thus (21.15) holds on the entire branch.

Step 4: convert the bank gap to positive compression. For $t > 0$, (K.12) and (21.15) give

$$a = a^+ - a^- \ge \delta_* a^+ > 0.$$

Also

$$a^+ + a^- \le (2-\delta_*)a^+,$$

hence

$$a \ge \frac{\delta_*}{2-\delta_*}(a^+ + a^-) > 0.$$

Step 5: dependency/no-double-charge audit. The interval $[0, t_d]$ is financed solely by the proved O1–O3 package. The threshold state is launch data for the completed inherited-launch source-angle audit. That audit charges only post-t_d response/geometry or inherited stock explicitly present at launch; no pre-threshold centered action is reused.

No O5 or M7 theorem enters the proof. In particular, no powered reserve, quarter-rate debt, cap-work, or bank/deformation-overlap hypothesis is used.

Therefore O6.1 implies O4 with the uniform constant (K.11). □

Audit status: GREEN/COMPLETE. The inherited-launch source-angle audit O6.1 is closed. Only O6.2 remains OPEN—NO PROOF ASSERTED.

O5. Terminal noncontinuation from O4 and O6.2. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 21.7.

Standing O4 solvency. Assume that from some finite classical time t_0 onward,

$$\boxed{a^-(t) \le (1-\delta_*)a^+(t), \qquad 0 < \delta_* < 1.} \tag{K.16}$$

Then

$$\boxed{a(t) = a^+(t) - a^-(t) \ge \delta_* a^+(t) > 0.} \tag{K.17}$$

The permanent terminal module M7 consists of the following already proved routes.

Mutual pointers: **M7**: grand-finale entry / short proof.

Theorem K.1 (M7-P: powered-reserve terminal closure)**.** *Fix $\kappa > 0$ and define*

$$\boxed{\mathcal{F}_\kappa := Q - 3(1+\kappa)a^2.} \tag{K.18}$$

Suppose that at some classical time $t_1 \geq t_0$,

$$a(t_1) > 0,$$

and that

$$\mathcal{F}_\kappa(t) \geq 0$$

for every later classical time under consideration. Then either regularity fails earlier or

$$\boxed{T_* \leq t_1 + \frac{1}{\kappa a(t_1)}.} \tag{K.19}$$

The historical reserve $\mathcal{F}_\Psi = Q - 9a^2$ is the special case $\kappa = 2$.

Proof. The exact apex equation is

$$\dot{a} = -a^2 + \tfrac{1}{3}Q.$$

Hence

$$\boxed{\mathcal{F}_\kappa = 3(\dot{a} - \kappa a^2).} \tag{K.20}$$

The hypothesis therefore gives

$$\dot{a} \geq \kappa a^2.$$

Since $a(t_1) > 0$,

$$\frac{d}{dt}\frac{1}{a} = -\frac{\dot{a}}{a^2} \leq -\kappa.$$

Thus

$$\frac{1}{a(t)} \leq \frac{1}{a(t_1)} - \kappa(t - t_1),$$

and a positive classical a cannot persist beyond $t_1 + [\kappa a(t_1)]^{-1}$. □

Remark K.2 (Powered detector interpretation). If $\mathfrak{t}_\Psi = \sqrt{\Psi/\Psi(0)} > 0$ and $\mathcal{Z}_\kappa := \mathfrak{t}_\Psi^{\kappa/2}$, then

$$\mathcal{F}_\kappa = -\frac{3}{\kappa}\frac{\ddot{\mathcal{Z}}_\kappa}{\mathcal{Z}_\kappa}.$$

Thus the powered-reserve condition is exactly concavity of a positive detector clock. The permanent O5 reserve $Q - 9a^2$ is therefore one member of a family of sufficient terminal clocks, not a logically necessary coefficient.

Theorem K.3 (M7-C1: positive-bank collapse-age growth closure)**.** *Assume* (K.16)*. Suppose there are constants $\kappa_* > 0$ and $C_0 < \infty$ such that*

$$\boxed{\log \frac{a^+(t)}{a^+(t_0)} \geq \kappa_* \Lambda(t; t_0) - C_0} \tag{K.21}$$

throughout the later classical interval. Then either regularity fails earlier or

$$\boxed{T_* \leq t_0 + \frac{e^{C_0}}{\kappa_* \delta_* a^+(t_0)}.} \tag{K.22}$$

Proof. From (K.21),

$$a^+(t) \geq a^+(t_0)e^{-C_0}e^{\kappa_*\Lambda(t;t_0)}.$$

Using (K.17),

$$\dot{\Lambda} = a \geq \delta_* a^+(t_0)e^{-C_0}e^{\kappa_*\Lambda}.$$

Therefore

$$\frac{d}{dt}e^{-\kappa_*\Lambda} \leq -\kappa_*\delta_* a^+(t_0)e^{-C_0}.$$

The positive left side cannot persist beyond the deadline (K.22). □

Maintained positive cashier. Let $\Sigma(t) > 0$ denote the dominance-free maintained positive cashier on a post-O4 interval and let $\Delta_{\mathrm{B}}(t) \geq 0$ denote its instantaneous quarter-rate shortage. Define

$$\boxed{\mathfrak{D}_{\mathrm{rate}}(t) := \int_{t_0}^{t} \Delta_{\mathrm{B}}(\tau)\, d\tau.} \tag{K.23}$$

The already audited cashier inequality is

$$\boxed{\dot{\Sigma} \geq \left(\frac{1}{4}a - \Delta_{\mathrm{B}}\right)\Sigma} \tag{K.24}$$

between maintenance operations. Residual positive-genealogy top-ups and same-genealogy refinancings have nonnegative jumps in Σ, and after O4 the historical Hodge-dominance refresh is never performed.

Theorem K.4 (M7-C2: dominance-free sub-quarter cashier closure)**.** *Assume O4 and the dominance-free maintained cashier above. Suppose*

$$\boxed{\mathfrak{D}_{\mathrm{rate}}(t) \leq \delta_{\mathrm{rate}}\Lambda(t;t_0) + C_{\mathrm{rate}}, \qquad 0 \leq \delta_{\mathrm{rate}} < \frac{1}{4}.} \tag{K.25}$$

Then either regularity fails earlier or

$$\boxed{T_* - t_0 \leq \frac{e^{C_{\mathrm{rate}}}}{(\frac{1}{4} - \delta_{\mathrm{rate}})\delta_*\Sigma(t_0)}.} \tag{K.26}$$

Proof. Integrating (K.24) across the continuous pieces and using the nonnegative maintenance jumps gives

$$\log\frac{\Sigma(t)}{\Sigma(t_0)} \geq \frac{1}{4}\Lambda(t;t_0) - \mathfrak{D}_{\mathrm{rate}}(t). \tag{K.27}$$

Moreover O4 gives the permanent bank comparison

$$a \geq \delta_*\Sigma.$$

With

$$\kappa_{\mathrm{rate}} := \frac{1}{4} - \delta_{\mathrm{rate}} > 0,$$

equation (K.25) yields

$$\dot{\Lambda} = a \geq \delta_*\Sigma(t_0)e^{-C_{\mathrm{rate}}}e^{\kappa_{\mathrm{rate}}\Lambda}.$$

The same scalar countdown as in theorem K.3 gives (K.26). □

Mutual pointers: **M8**: grand-finale entry / short proof.

Theorem K.5 (Exact post-O4 integral cashier countdown)**.** *Assume O4 and a dominance-free maintained positive cashier launched at a classical time $t_1 > 0$ with $\Sigma(t_1) > 0$. Put*

$$\lambda := \Lambda(t; t_1), \qquad \mathfrak{D}_{\mathrm{rate}}(\lambda) := \int_{t_1}^{t(\lambda)} \Delta_{\mathrm{B}}(\tau)\, d\tau$$

and define the net cashier gain

$$\boxed{\mathcal{G}_{\mathrm{cash}}(\lambda) := \frac{\lambda}{4} - \mathfrak{D}_{\mathrm{rate}}(\lambda).} \tag{K.28}$$

If

$$\boxed{\int_0^\infty e^{-\mathcal{G}_{\mathrm{cash}}(\lambda)}\, d\lambda < \infty,} \tag{K.29}$$

then either a named classical continuation exit occurs earlier or the classical branch cannot persist for infinite collapse age. More precisely,

$$\boxed{T_* - t_1 \le \frac{1}{\delta_* \Sigma(t_1)} \int_0^\infty e^{-\mathcal{G}_{\mathrm{cash}}(\lambda)}\, d\lambda.} \tag{K.30}$$

Proof. Integrating the maintained-cashier inequality

$$\dot{\Sigma} \ge \left(\frac{1}{4} a - \Delta_{\mathrm{B}}\right) \Sigma$$

across continuous pieces and using the nonnegative maintenance jumps gives

$$\Sigma(t) \ge \Sigma(t_1) e^{\mathcal{G}_{\mathrm{cash}}(\lambda)}.$$

O4 yields

$$a(t) \ge \delta_* \Sigma(t).$$

Since $d\lambda = a(t)\, dt$,

$$\frac{dt}{d\lambda} = \frac{1}{a(t)} \le \frac{1}{\delta_* \Sigma(t_1)} e^{-\mathcal{G}_{\mathrm{cash}}(\lambda)}.$$

Integration proves (K.30). □

Corollary K.6 (Vanishing quarter-rate margin)**.** *Under the hypotheses of theorem K.5, suppose that for some $p > 1$ and all sufficiently large collapse ages,*

$$\boxed{\frac{\Delta_{\mathrm{B}}}{a} \le \frac{1}{4} - \frac{p}{1+\lambda}.} \tag{K.31}$$

Then finite physical-time noncontinuation follows unless a named continuation exit occurs earlier.

Proof. At regular collapse-age points,

$$\mathcal{G}'_{\mathrm{cash}}(\lambda) = \frac{1}{4} - \frac{\Delta_{\mathrm{B}}}{a} \ge \frac{p}{1+\lambda}.$$

Hence

$$\mathcal{G}_{\mathrm{cash}}(\lambda) \ge p \log(1+\lambda) - C,$$

so

$$e^{-\mathcal{G}_{\mathrm{cash}}(\lambda)} \le C'(1+\lambda)^{-p}.$$

The latter is integrable because $p > 1$. □

Remark K.7 (Exact scalar falsification of fixed-margin minimality). The fixed-margin terminal gates P, C1 and C2 are useful sufficient conditions, but they are not necessary even for the exact scalar cashier countdown. On the tail $\lambda \geq 8$, set

$$a(\lambda) = a^{+}(\lambda) = \Sigma(\lambda) = (1+\lambda)^2,$$

and prescribe

$$\frac{\Delta_{\mathrm{B}}}{a} = \frac{1}{4} - \frac{2}{1+\lambda}.$$

The shortage is nonnegative because

$$\frac{1}{4} - \frac{2}{1+\lambda} \geq \frac{1}{4} - \frac{2}{9} = \frac{1}{36} > 0.$$

The cashier law is then an equality:

$$\frac{d}{d\lambda} \log \Sigma = \frac{2}{1+\lambda} = \frac{1}{4} - \frac{\Delta_{\mathrm{B}}}{a}.$$

Moreover

$$\int_8^{\infty} \frac{d\lambda}{a(\lambda)} = \int_8^{\infty} \frac{d\lambda}{(1+\lambda)^2} = \frac{1}{9} < \infty.$$

Nevertheless,

$$\frac{d}{d\lambda} \log a = \frac{2}{1+\lambda} \longrightarrow 0,$$

so no P gate with a fixed $\kappa > 0$ holds, and

$$\frac{\log(a^{+}(\lambda)/a^{+}(8))}{\lambda - 8} = \frac{2\log((1+\lambda)/9)}{\lambda - 8} \longrightarrow 0,$$

so no fixed positive C1 exponent holds. Finally,

$$\mathfrak{D}_{\mathrm{rate}}(\lambda) = \frac{\lambda - 8}{4} - 2\log\frac{1+\lambda}{9},$$

and therefore

$$\frac{\mathfrak{D}_{\mathrm{rate}}(\lambda)}{\lambda - 8} \longrightarrow \frac{1}{4},$$

so no C2 condition with a fixed $\delta_{\mathrm{rate}} < 1/4$ holds.

For direct verification, the exact values at four sample ages are

λ	Δ_{B}/a	$\mathcal{G}_{\mathrm{cash}}$	$e^{-\mathcal{G}_{\mathrm{cash}}}$	$dt/d\lambda$
8	1/36	0	1	1/81
17	5/36	$2\log 2$	1/4	1/324
26	19/108	$2\log 3$	1/9	1/729
35	7/36	$2\log 4$	1/16	1/1296

where in this scalar sample $dt/d\lambda = (1+\lambda)^{-2}$ exactly. Thus the verification may be performed with exact rational arithmetic and elementary logarithms; no floating-point constant is needed.

This is an exact scalar falsification sample, not a numerical approximation and not an Euler solution. Its role is only to show that the old fixed-margin interface is stronger than the exact terminal requirement. A single exact counterexample is sufficient for that falsification claim, but it does not prove the open Euler closure.

Remark K.8 (No refresh-count payment). The theorem above contains no Hodge-dominance refresh debit. After O4, the global comparison $a \geq \delta_* \Sigma$ makes that refresh unnecessary. Thus the terminal countdown charges only the continuous shortage $\mathfrak{D}_{\mathrm{rate}}$ and cannot mistake an intervention count for physical work.

Count-free physical debt. On a maintained interval $I = [t_0, t]$, put

$$d\lambda := a(t)\, dt, \qquad \Lambda_I := \int_I d\lambda.$$

Let $\mathcal{J}_I^{\mathrm{car}}$ denote the count-free carrier-forcing ledger, $\mathcal{V}_{I,\lambda}$ the maintained-bank radial/value-deformation ledger, and $\mathcal{A}_{I,\lambda}$ the signed axial-deformation ledger. Under a fixed carrier-ratio cap, the audited O4-transplanted estimate has the form

$$\boxed{\mathfrak{D}_{\mathrm{rate}}(I) \leq C_{\mathrm{init}} + C_J \mathcal{J}_I^{\mathrm{car}} + C_V \mathcal{V}_{I,\lambda} + C_A \mathcal{A}_{I,\lambda},} \tag{K.32}$$

where the constants are finite and depend only on the fixed cap, fine-cell parameters, and δ_*. No top-up count, refinancing count, or dominance-refresh count appears.

Corollary K.9 (M7-C3: zero-density pressure-free closure)**.** *Assume O4 on an unbounded-collapse-age branch. Suppose a fixed carrier-ratio cap is available on a terminal tail and*

$$\boxed{\frac{\mathcal{J}^{\mathrm{car}}_{[t_0,t]}}{\Lambda(t;t_0)} \to 0, \qquad \frac{\mathcal{V}_{[t_0,t],\lambda}}{\Lambda(t;t_0)} \to 0, \qquad \frac{\mathcal{A}_{[t_0,t],\lambda}}{\Lambda(t;t_0)} \to 0.} \tag{K.33}$$

Then finite-time noncontinuation follows. If no fixed carrier-ratio cap is maintainable, the alternative is divergent carrier cap-work, which is already a named physical failure account.

Proof. Divide (K.32) by $\Lambda(t;t_0)$. The finite initial term and the three physical ledgers have zero density, so

$$\frac{\mathfrak{D}_{\mathrm{rate}}}{\Lambda} \to 0.$$

Hence for all sufficiently large collapse age,

$$\mathfrak{D}_{\mathrm{rate}} \leq \delta_{\mathrm{rate}} \Lambda + C_{\mathrm{rate}}$$

with some $\delta_{\mathrm{rate}} < 1/4$. Apply theorem K.4 from a sufficiently late maintained-bank launch. □

Remark K.10 (M7-F: bank/deformation segregation firewall). Let

$$\mathcal{V}_B(t) := \langle |v - a| \rangle_{\mu_B(t)}, \qquad \mathcal{V}_{I,\lambda} = \int_I \mathcal{V}_B(t)\, dt.$$

Although the global energy/continuation audit yields

$$\int_{t_0}^{t} \|v - a\|_\infty \, ds \geq \Lambda(t;t_0) - C_{\mathrm{sw}}(t_0),$$

one only has

$$0 \leq \mathcal{V}_B(t) \leq \|v - a\|_\infty.$$

Thus raw global deformation cannot be substituted for the maintained-bank debt.

Whenever $\|v - a\|_\infty > 0$, define

$$\kappa_B(t) := \frac{\mathcal{V}_B(t)}{\|v - a\|_\infty} \in [0, 1]$$

and, when the denominator is nonzero,

$$\overline{\kappa}_B(I) := \frac{\mathcal{V}_{I,\lambda}}{\int_I \|v-a\|_\infty \, dt}.$$

Then exactly

$$\mathcal{V}_{I,\lambda} = \overline{\kappa}_B(I) \int_I \|v-a\|_\infty \, dt.$$

Consequently a small maintained-bank value-debt density on an unbounded-age branch is possible only through quantitatively small bank/deformation overlap. No positive lower bound for that overlap is asserted here.

Continuation firewall. On a finite physical-time classical interval, the global energy audit also gives

$$\Lambda(t;t_0) \to \infty \quad \Longrightarrow \quad \int_{t_0}^{T} \|\nabla U(t)\|_\infty \, dt = \infty.$$

Thus unbounded collapse age is already a named physical Lipschitz continuation exit when a finite endpoint has been fixed. The pressure-free M7-C route remains useful because it constructs a finite endpoint from the maintained positive cashier itself.
Terminal-activation audit. The permanent M7 terminal engines do not follow from O4 alone. In particular, theorem K.10 forbids replacing the maintained-bank value debt by the raw global deformation forced by energy, and positivity of a does not imply an eventual powered reserve.

Accordingly the exact missing activation is stated openly as O6.2 rather than being hidden in O5.

Proof of O5 from O4 and O6.2. Assume O4, O6.2, and no earlier named continuation exit. By O6.2 there is a finite maintained-cashier launch time t_1 with $\Sigma(t_1) > 0$ such that, writing $\lambda = \Lambda(t;t_1)$ and $\mathcal{G}_{\mathrm{cash}}(\lambda) = \lambda/4 - \mathfrak{D}_{\mathrm{rate}}(\lambda)$,

$$\int_0^\infty e^{-\mathcal{G}_{\mathrm{cash}}(\lambda)} \, d\lambda < \infty.$$

Apply the exact M8 countdown theorem K.5. It gives

$$T_* - t_1 \le \frac{1}{\delta_* \Sigma(t_1)} \int_0^\infty e^{-\mathcal{G}_{\mathrm{cash}}(\lambda)} \, d\lambda < \infty.$$

Thus the fixed customer leaves the classical category in finite physical time unless a named continuation exit occurred earlier. The three M7 gates remain stronger sufficient special cases, but O5 needs only the M8 integral criterion. □

Audit status: GREEN/COMPLETE as O4 + O6.2 + M8 $\Longrightarrow$ O5. The completion is conditional only on the sole open proposition O6.2.

O6.2. Post-O4 integral cashier closure; M1–M8 module archive. *Mutual pointers:* grand-finale entry / short proof/status / main statement open proposition 21.8.
Permanent funded upstream modules M1–M2. The exact polarization/source-history structure and the repaired critical fractional cashier are permanent proved modules. Together with the subsequent source-angle modules they form the closed inherited-launch audit feeding O4.

M1: polarization/source-history structure. *Mutual pointers:* **M1**: grand-finale entry / short proof.

Put

$$A_{\mathrm{abs}} := a^+ + a^-, \qquad \Pi := \frac{a^+ - a^-}{A_{\mathrm{abs}}} \tag{K.34}$$

whenever $A_{\mathrm{abs}} > 0$. On the two-copy genealogy space $\mathcal{Q} \times \{+,-\}$ define

$$d\nu_{\mathrm{abs}}(X,\varsigma) := \frac{\frac{3}{8} K_H(X) H^{\varsigma}(X,t)\, dX}{A_{\mathrm{abs}}(t)}, \qquad \sigma_H(X,\pm) := \pm 1. \tag{K.35}$$

Let

$$\mathcal{S}_{\mathrm{abs}} := \mathcal{S}_+ + \mathcal{S}_-, \qquad q_{\mathrm{abs}} := \frac{\mathcal{S}_{\mathrm{abs}}}{A_{\mathrm{abs}}},$$

and, when $\mathcal{S}_{\mathrm{abs}} > 0$,

$$\Pi_{\mathrm{src}} := \frac{\mathcal{S}_+ - \mathcal{S}_-}{\mathcal{S}_{\mathrm{abs}}}.$$

Theorem K.11 (M1-A: global Hodge-polarization balance)**.** *Whenever $A_{\mathrm{abs}} > 0$,*

$$\boxed{\dot{\Pi} = \mathrm{Cov}_{\nu_{\mathrm{abs}}}(\mathcal{C}, \sigma_H) + q_{\mathrm{abs}}(\Pi_{\mathrm{src}} - \Pi).} \tag{K.36}$$

Hence global cancellation has exactly two mechanisms: differential Hodge selection of already banked genealogy and fresh-source polarization mismatch.

Proof. The exact bank equations are

$$\dot{a}^{\pm} = \mathcal{S}_{\pm} + \mathcal{M}_{\pm}, \qquad \mathcal{M}_{\pm} = \frac{3}{8} \int K_H \mathcal{C} H^{\pm}\, dX.$$

Differentiate $\Pi = (a^+ - a^-)/(a^+ + a^-)$. The geometry numerator is

$$(\mathcal{M}_+ - \mathcal{M}_-) - \Pi(\mathcal{M}_+ + \mathcal{M}_-) = A_{\mathrm{abs}}\, \mathrm{Cov}_{\nu_{\mathrm{abs}}}(\mathcal{C}, \sigma_H),$$

while the source numerator is

$$(\mathcal{S}_+ - \mathcal{S}_-) - \Pi \mathcal{S}_{\mathrm{abs}} = \mathcal{S}_{\mathrm{abs}}(\Pi_{\mathrm{src}} - \Pi).$$

Divide by A_{abs}. □

Let

$$H_{\mathrm{abs}} := H^+ + H^-, \qquad \pi_H := \frac{H}{H_{\mathrm{abs}}}$$

on $\{H_{\mathrm{abs}} > 0\}$, and let $\sigma_q := q/|q|$ on $\{q \neq 0\}$.

Theorem K.12 (M1-B: local genealogy source-relaxation law)**.** *On $\{H_{\mathrm{abs}} > 0\}$,*

$$\boxed{\mathcal{D}_{xy} \pi_H = \frac{2|q|}{H_{\mathrm{abs}}} (\sigma_q - \pi_H).} \tag{K.37}$$

In particular, on a source-free material interval, π_H is constant along the trajectory.

Proof. The signed and absolute genealogy equations are

$$\mathcal{D}_{xy} H = 2q - gH, \qquad \mathcal{D}_{xy} H_{\mathrm{abs}} = 2|q| - g H_{\mathrm{abs}}.$$

Differentiate $\pi_H = H/H_{\mathrm{abs}}$. The common geometry terms cancel identically, leaving (K.37). □

Let μ_t be the spatial absolute-Hodge probability

$$d\mu_t := \frac{\frac{3}{8}K_H H_{\mathrm{abs}}\, dX}{A_{\mathrm{abs}}}.$$

For each current point X, let $\widetilde{\nu}^{\mathrm{hist}}_{X,t}$ be the signed source-history probability on source-time/sign space whose sign mean equals $\pi_H(X,t)$. Define

$$\mathcal{H}_{\mathrm{hist}}(t) := \iint \mathrm{JS}\big(\widetilde{\nu}^{\mathrm{hist}}_{X,t}, \widetilde{\nu}^{\mathrm{hist}}_{Y,t}\big)\, d\mu_t(X) d\mu_t(Y). \tag{K.38}$$

Theorem K.13 (M1-C: source-history selection bound)**.** *One has*

$$\boxed{|\mathrm{Cov}_{\mu_t}(\mathcal{C}, \pi_H)| \le 2\sqrt{\mathrm{Var}_{\mu_t}(\mathcal{C})\mathcal{H}_{\mathrm{hist}}(t)}.} \tag{K.39}$$

Thus adverse Hodge selection requires both nontrivial contraction-rate variance and nontrivial signed source-history separation.

Proof. The signed source-history representation gives

$$|\pi_{H,X} - \pi_{H,Y}| \le 2\,\mathrm{TV}\left(\widetilde{\nu}^{\mathrm{hist}}_{X,t}, \widetilde{\nu}^{\mathrm{hist}}_{Y,t}\right).$$

Pinsker applied through Jensen–Shannon divergence yields

$$|\pi_{H,X} - \pi_{H,Y}|^2 \le 8\,\mathrm{JS}\left(\widetilde{\nu}^{\mathrm{hist}}_{X,t}, \widetilde{\nu}^{\mathrm{hist}}_{Y,t}\right),$$

hence $\mathrm{Var}_{\mu_t}(\pi_H) \le 4\mathcal{H}_{\mathrm{hist}}$. Cauchy–Schwarz for covariance then gives (K.39). □

M2: source-positive critical fractional cashier. *Mutual pointers:* **M2**: grand-finale entry / short proof.

Recall

$$\phi(s) = \frac{3}{8}\frac{s^{5/6}}{(1+s)^{5/2}}, \qquad \psi_\varpi(s) := \phi_*^{1-\varpi}\phi(s)^\varpi, \qquad W_\varpi(x,y) := \frac{\psi_\varpi(y/x)}{\mathfrak{r}(x,y)}.$$

Theorem K.14 (M2-A: sharp source-positive critical cashier)**.** *Universal signed-source nonnegativity*

$$2\int W_\varpi q\, dX \ge 0$$

for every nonnegative tagged swirl source holds precisely for

$$0 \le \varpi \le \frac{2}{5}.$$

At the maximal exponent $\varpi = 2/5$,

$$\boxed{W_c(x,y) := W_{2/5}(x,y) = \frac{\sqrt{3}}{12(x+y)},} \tag{K.40}$$

and

$$\boxed{2\int W_c q\, dX = \frac{\sqrt{3}}{6}\int \frac{\Theta}{(x+y)^2}\, dX \ge 0.} \tag{K.41}$$

The right side is strictly positive for nontrivial tagged swirl.

Proof. At fixed x, signed-source positivity after integration by parts is equivalent to $-\partial_y W_\varpi \geq 0$, or

$$\frac{d}{d\log s}\log(\psi_\varpi(s)s^{-1/3}) \leq 0.$$

Since

$$\frac{d}{d\log s}\log\phi(s) = \frac{5(1-2s)}{6(1+s)},$$

whose supremum is $5/6$, the sharp condition is $\varpi(5/6) \leq 1/3$, i.e. $\varpi \leq 2/5$. At $\varpi = 2/5$,

$$\psi_{2/5}(s) = \frac{\sqrt{3}}{12}\frac{s^{1/3}}{1+s},$$

so (K.40) follows from $\mathfrak{r} = xs^{1/3}$. Integration by parts then gives (K.41). □

Define the critical sign banks

$$A_c^\pm := \int_{\mathcal{Q}} W_c H^\pm \, dX = \frac{\sqrt{3}}{12}\int_{\mathcal{Q}} \frac{H^\pm}{x+y}\, dX. \tag{K.42}$$

Put

$$\vartheta := \frac{y}{x+y}, \qquad \omega(\vartheta) := \frac{3\sqrt{3}}{2}(1-\vartheta)\sqrt{\vartheta}, \qquad 0 \leq \omega \leq 1.$$

Whenever $A_c^\pm > 0$, define

$$d\nu_{c,\pm} := \frac{W_c H^\pm \, dX}{A_c^\pm}, \qquad \zeta_\pm := \langle\omega\rangle_{\nu_{c,\pm}}.$$

Theorem K.15 (M2-B: critical-cashier/Hodge factorization)**.** *For each sign,*

$$\boxed{a^\pm = A_c^\pm \zeta_\pm, \qquad 0 < \zeta_\pm \leq 1.} \tag{K.43}$$

Hence

$$\boxed{\frac{a^-}{a^+} = \frac{A_c^-}{A_c^+}\frac{\zeta_-}{\zeta_+}.} \tag{K.44}$$

Proof. The true Hodge kernel factors exactly as

$$\frac{3}{8}K_H = W_c\,\omega.$$

Integrating against $H^\pm$ gives $a^\pm = A_c^\pm\langle\omega\rangle_{\nu_{c,\pm}}$. □

Define the residual relative efficiency

$$\Delta_c := \log\frac{\zeta_+}{\zeta_-}.$$

Theorem K.16 (M2-C: common-affine cancellation and bounded source relaxation)**.** *The geometry part of Δ_c obeys*

$$\boxed{\dot{\Delta}_c\Big|_{\text{geom}} = \mathfrak{R}_+^{\text{vis}} - \mathfrak{R}_-^{\text{vis}},} \tag{K.45}$$

so the common affine rate cancels exactly. Moreover each sign efficiency satisfies

$$\boxed{\dot{\zeta}_\pm = \zeta_\pm D_\pm + s_{c,\pm}(\omega_{\text{src},\pm} - \zeta_\pm),} \tag{K.46}$$

where $0 \leq \omega_{\text{src},\pm} \leq 1$.

Proof. The exact difference between the Hodge and critical geometry rates is a common affine contribution plus a centered remainder. Subtracting the two sign equations removes the common affine term, yielding (K.45). Differentiating $\zeta_\pm = a^\pm/A_c^\pm$ and using the corresponding Hodge and critical source rates gives (K.46). □

For $\varpi \in [0,1]$ define

$$\ell_\phi(s) := \log \frac{\phi_*}{\phi(s)} \geq 0, \qquad A_\varpi^\pm := \int \psi_\varpi(s) \frac{H^\pm}{\mathfrak{r}}\, dX,$$

and, whenever $A_\varpi^\pm > 0$,

$$d\nu_{\varpi,\pm} := \frac{\psi_\varpi H^\pm\, dX/\mathfrak{r}}{A_\varpi^\pm}, \qquad m_{\varpi,\pm} := \langle \ell_\phi \rangle_{\nu_{\varpi,\pm}}.$$

Let

$$h_\varpi := \frac{1}{2} \log \frac{A_\varpi^+}{A_\varpi^-}.$$

Theorem K.17 (M2-D: critical-to-Hodge fractional homotopy)**.** *For each sign,*

$$\partial_\varpi \log A_\varpi^\pm = -m_{\varpi,\pm}. \tag{K.47}$$

Consequently

$$\boxed{h_1 = h_{2/5} - \frac{1}{2} \int_{2/5}^1 (m_{\varpi,+} - m_{\varpi,-})\, d\varpi.} \tag{K.48}$$

Equivalently, with

$$\Pi_\varpi := \frac{A_\varpi^+ - A_\varpi^-}{A_\varpi^+ + A_\varpi^-},$$

one has

$$\boxed{\partial_\varpi \Pi_\varpi = -\operatorname{Cov}_{\nu_{\varpi,\mathrm{abs}}}(\ell_\phi, \sigma_H).} \tag{K.49}$$

Thus the transfer from the source-positive critical cashier to the true Hodge cashier is pure sign-dependent visibility-age selection.

Proof. Since $\partial_\varpi \psi_\varpi = -\ell_\phi \psi_\varpi$, differentiate $A_\varpi^\pm$ to obtain (K.47). Subtract the two sign equations and integrate from $2/5$ to 1 to obtain (K.48). The two-copy normalized derivative gives (K.49). □

Remark K.18 (M2-F: affine-benchmark firewall). The one-sided negative-carrier/Hodge envelope is not an authoritative closure route. On the exact affine conveyor,

$$\frac{\phi_* B_-^{\mathrm{aff}}}{a_+^{\mathrm{aff}}} \to +\infty$$

even though the exact affine Hodge ratio retains a uniform strict gap. The repaired M2 route avoids this false obstruction by starting at the source-positive critical exponent, using relative rather than one-sided Hodge efficiency, and cancelling the common affine visibility rate exactly.

M1–M2 funded interface. The permanent post-ticket starting chain is now

$$\boxed{\begin{aligned}\text{O3 ticket state} &\Longrightarrow \text{M1 source/history decomposition}\\ &\Longrightarrow \text{M2 source-positive critical cashier}\\ &\Longrightarrow \text{relative critical-to-Hodge age/sign debit.}\end{aligned}} \tag{K.50}$$

No arrow in (K.50) by itself claims the inherited source-angle conclusion. The later M3–M6 modules must route the remaining relative debit to named post-ticket accounts or leave precisely the young inherited source-angle branch.

M3: post-ticket causal pay-or-route module. *Mutual pointers:* **M3**: grand-finale entry / short proof.

Let t_d be the first O3 ticket time. Freeze the genealogy present there:

$$\boxed{\begin{aligned} H_O(\cdot,t_d) &:= H(\cdot,t_d), \quad \partial_t H_O + \nabla\cdot(\mathbf{b}H_O) = 0,\\ H_N &:= H - H_O, \quad \partial_t H_N + \nabla\cdot(\mathbf{b}H_N) = 2q, \qquad H_N(\cdot,t_d) = 0.\end{aligned}} \tag{K.51}$$

For $J \in \{O, N\}$ write

$$H_J = H_J^+ - H_J^-, \qquad a_J^\pm := \frac{3}{8}\int_{\mathcal{Q}} K_H H_J^\pm \, dX, \qquad P_J := a_J^+ - a_J^-.$$

Then

$$a = P_O + P_N, \qquad P_O(t_d) = a(t_d) > 0, \qquad P_N(t_d) = 0.$$

Theorem K.19 (M3-A: ticket-time old/new Hodge-lock dichotomy)**.** *Suppose $T_* > t_d$ is a terminal Hodge lock,*

$$a(T_*) = 0, \qquad a^+(T_*) = a^-(T_*) > 0.$$

Then exactly one of the following occurs.

(1) ***OLD reversal:*** *$P_O(T_*) \le 0$, in which case*

$$\boxed{15\Lambda(T_*;t_d) + 20\int_{t_d}^{T_*} \|v-a\|_\infty \, dt + 8\int_{t_d}^{T_*} \|\mathfrak{A}\|_\infty \, dt \ge \log\frac{1}{\theta_{\mathrm{O3}}}.} \tag{K.52}$$

(2) ***NEW adverse genealogy:*** *$P_O(T_*) > 0$ and $P_N(T_*) < 0$, where H_N has zero initial genealogy at t_d and is generated entirely from post-ticket source times.*

Proof. At the terminal lock,

$$P_O(T_*) + P_N(T_*) = 0.$$

If $P_O(T_*) \le 0$, the source-free old bank has crossed from the strict O3 ticket gap to nonpositive sign. Its negative-to-positive ratio therefore reaches one. The source-free Hodge sign-bank rate is the difference of two averages of the same contraction field $\mathcal{C}$, whose post-ticket oscillation satisfies

$$\operatorname{osc}\mathcal{C} \le 15a + 20\|v-a\|_\infty + 8\|\mathfrak{A}\|_\infty.$$

Integration from t_d gives (K.52). If instead $P_O(T_*) > 0$, the terminal balance forces $P_N(T_*) = -P_O(T_*) < 0$. Since $H_N(t_d) = 0$, this genealogy is genuinely post-ticket. □

The theorem is the exact no-double-charge cut: Alternative OLD charges only post-ticket physical-age/geometry work; Alternative NEW contains no inherited O3 deformation capital.

Define the old-reversal and critical-geometry exposures

$$\mathfrak{G}_O(T) := 15\Lambda(T;t_d) + 20\int_{t_d}^{T} \|v-a\|_\infty\,dt + 8\int_{t_d}^{T} \|\mathfrak{A}\|_\infty\,dt, \tag{K.53}$$

$$\mathfrak{G}_c(T) := 3\Lambda(T;t_d) + 4\int_{t_d}^{T} \|v-a\|_\infty\,dt + 2\int_{t_d}^{T} \|\mathfrak{A}\|_\infty\,dt. \tag{K.54}$$

Let $\pi_c^{\mathrm{src}} > 0$ denote the restarted critical source-polarization floor on the admitted no-exit branch. When

$$\mathfrak{G}_c(T_*) < \frac{1}{2}\pi_c^{\mathrm{src}},$$

the restarted critical comparison gives the setwise packet floor

$$\boxed{m_c^T(\tau) \geq \frac{1}{2}\pi_c^{\mathrm{src}} \quad \text{for } \rho_{1,N}\text{-a.e. } \tau.} \tag{K.55}$$

Define the same-source-time critical-to-Hodge sign drop

$$\boxed{\mathfrak{D}_{\mathrm{same}}(T) := \int_{t_d}^{T} [m_c^T(\tau) - m_1^T(\tau)]_+\,d\rho_{1,N}(\tau).} \tag{K.56}$$

Theorem K.20 (M3-B: closure-grade three-gate alternative)**.** *Every post-O3 terminal Hodge lock satisfies at least one of*

$$\boxed{\mathfrak{G}_O \geq \log(1/\theta_{\mathrm{O3}}), \qquad \mathfrak{G}_c \geq \frac{1}{2}\pi_c^{\mathrm{src}}, \qquad \mathfrak{D}_{\mathrm{same}} > \frac{1}{2}\pi_c^{\mathrm{src}}.} \tag{K.57}$$

The three alternatives are called Gate O, Gate C, and Gate S, respectively.

Proof. If Gate O fails, theorem K.19 forces the NEW branch $P_N(T_*) < 0$. If Gate C also fails, the setwise critical packet floor (K.55) holds. Since the final Hodge mean of the NEW genealogy is negative while its critical packet means are uniformly positive, averaging over the Hodge source-time law forces the same-packet sign-drop debit $\mathfrak{D}_{\mathrm{same}} > \pi_c^{\mathrm{src}}/2$. □

Thus Gates O and C are already explicit post-ticket physical-age/geometry invoices. Gate S is the only qualitatively different branch.

M4: normalized Gate-S / source-angle GREEN bridge. *Mutual pointers:* **M4**: grand-finale entry / short proof.

Fix a post-ticket source time $\tau \in [t_d, T]$ and let $\widehat{\sigma}_{c,\tau}^T$ be the final critical conditional law of that packet. Let

$$\Sigma := \operatorname{sgn} q(X,\tau) \in \{-1,+1\}, \qquad \omega := \frac{W_1}{W_c} \in [0,1],$$

and define

$$m_c^T(\tau) := \langle \Sigma \rangle_c, \qquad \zeta(\tau) := \langle \omega \rangle_c.$$

The final Hodge conditional law is the ω-tilt

$$\boxed{d\widehat{\sigma}_{1,\tau}^T = \frac{\omega}{\zeta(\tau)} d\widehat{\sigma}_{c,\tau}^T.} \tag{K.58}$$

Theorem K.21 (M4-A: sign-conditioned Gate-S identity)**.** *For every packet with* $\zeta(\tau) > 0$,

$$\boxed{m_c^T(\tau) - m_1^T(\tau) = -\frac{\mathrm{Cov}_c(\omega, \Sigma)}{\zeta(\tau)}.} \tag{K.59}$$

If ρ_c *is the final critical source-time law and* $\bar{\zeta} := \langle \zeta \rangle_{\rho_c}$*, then*

$$\boxed{\bar{\zeta}\, \mathfrak{D}_{\mathrm{same}} = \int [-\mathrm{Cov}_c(\omega, \Sigma)]_+ \, d\rho_c.} \tag{K.60}$$

Moreover, writing

$$\omega_\pm(\tau) := \langle \omega \mid \Sigma = \pm 1 \rangle_c,$$

one has

$$\boxed{[-\mathrm{Cov}_c(\omega, \Sigma)]_+ = \frac{1 - (m_c^T)^2}{2} [\omega_- - \omega_+]_+.} \tag{K.61}$$

Proof. Equation (K.58) gives

$$m_1^T = \frac{\langle \Sigma \omega \rangle_c}{\zeta},$$

which yields (K.59). The Hodge source-time law is the ζ-tilt of ρ_c, so the local denominator cancels after integration, giving (K.60). Finally, for a two-sign law,

$$\mathrm{Cov}_c(\omega, \Sigma) = 2p_+p_-(\omega_+ - \omega_-) = \frac{1 - (m_c^T)^2}{2}(\omega_+ - \omega_-).$$

□

Thus Gate S is not raw visibility loss. It requires a negative-sign residual Hodge visibility advantage inside sign-mixed critical packets.

For the same source time, let

$$L(\tau) := \Lambda(T; \tau), \qquad \omega_L(X, \tau) := \omega(e^{6L(\tau)} y/x),$$

and let $\sigma_{c,\tau}^{\pm}$ be the critical source laws conditioned on source sign. Put

$$\omega_{\mathrm{src},\pm}^0 := \langle \omega(y/x) \rangle_{\sigma_{c,\tau}^{\pm}}.$$

Define the packetwise centered angular action

$$D_{\mathrm{ang}}^*(\tau) := \operatorname*{ess\,sup}_{|q(X,\tau)|>0} \int_\tau^T \big(|v - a| + |g - 2a|\big)(\mathscr{X}_{s,\tau} X, s)\, ds,$$

and the critical response-selection oscillation

$$\delta_c(\tau) := \mathrm{osc}_{|q(X,\tau)|>0} \log G_{c,T;\tau}(X).$$

Theorem K.22 (M4-B: Gate-S packet pay-or-route)**.** *For every post-ticket source time,*

$$\boxed{\begin{aligned}[\omega_-^T - \omega_+^T]_+ \le\ & [\omega_{\mathrm{src},-}^0 - \omega_{\mathrm{src},+}^0]_+ \\ & + 12\Lambda(T; \tau) \\ & + 2\left(e^{\delta_c(\tau) + 2D_{\mathrm{ang}}^*(\tau)} - 1\right).\end{aligned}} \tag{K.62}$$

Proof. The exact aspect-ratio law compares true final visibility with the same-clock affine visibility by the multiplicative factor $e^{\pm 2D^*_{\mathrm{ang}}}$. The critical response law differs from the deposition law by a likelihood ratio with oscillation δ_c. This gives

$$[\omega^T_- - \omega^T_+]_+ \le [\omega^L_{\mathrm{src},-} - \omega^L_{\mathrm{src},+}]_+ + 2(e^{\delta_c + 2D^*_{\mathrm{ang}}} - 1).$$

Since $|d\omega/d\log s| \le 1$, the affine physical-age change satisfies

$$[\omega^L_{\mathrm{src},-} - \omega^L_{\mathrm{src},+}]_+ \le [\omega^0_{\mathrm{src},-} - \omega^0_{\mathrm{src},+}]_+ + 12L.$$

Combine the two estimates. □

The three causal payer classes are therefore deposition source-angle selection, physical relative age, and post-deposition response deformation.

To identify the deposition payer, define the orthogonalized source-gradient rotor

$$\varrho_Z := \sqrt{q^2 + \widehat{C}^2_\beta}, \qquad q = \varrho_Z \cos\theta_Z,$$

and put

$$c_Z := |\cos\theta_Z|, \qquad h_Z := \cos\theta_Z.$$

Let $\sigma_{\rho,c}$ be the orientation-free critical rotor-amplitude law and

$$\bar{c}_Z := \langle c_Z \rangle_{\sigma_{\rho,c}}.$$

Then the physical critical source law is

$$d\sigma_c = \frac{c_Z}{\bar{c}_Z}\, d\sigma_{\rho,c}.$$

With

$$\Pi_{\mathrm{src},c} := \langle \Sigma \rangle_{\sigma_c}, \qquad R_Z := h_Z - \Pi_{\mathrm{src},c} c_Z,$$

one has $\langle R_Z \rangle_{\sigma_{\rho,c}} = 0$.

Theorem K.23 (M4-C: centered rotor representation of deposition source angle)**.**

$$\boxed{\mathrm{Cov}_{\sigma_c}(\omega, \Sigma) = \frac{1}{\bar{c}_Z} \mathrm{Cov}_{\sigma_{\rho,c}}(\omega, R_Z).} \tag{K.63}$$

Proof. Using $h_Z = c_Z \Sigma$ and the c_Z-tilt,

$$\langle \omega\Sigma \rangle_{\sigma_c} = \frac{\langle \omega h_Z \rangle_{\sigma_{\rho,c}}}{\bar{c}_Z}, \qquad \langle \omega \rangle_{\sigma_c} = \frac{\langle \omega c_Z \rangle_{\sigma_{\rho,c}}}{\bar{c}_Z}.$$

Subtract $\Pi_{\mathrm{src},c} \langle \omega \rangle_{\sigma_c}$ and use the definition of R_Z. □

This replaces absolute negative-source fraction by a centered orientation covariance.

Ticket source-angle reserve. Define the additive critical-to-Hodge source-angle reserve

$$\boxed{\Delta_{\mathrm{src}} := \Pi_{\mathrm{src},1} - \Pi_{\mathrm{src},c}.} \tag{K.64}$$

Theorem K.24 (M4-D: ticket source-angle reserve certificate)**.** *For the fixed customer, the O3 small-action threshold may be chosen compatibly with the audited affine source-angle certificate so that at the first ticket time*

$$\boxed{\Delta_{\mathrm{src}}(t_d) \ge \delta_{\mathrm{src}} > 0.} \tag{K.65}$$

Proof. The exact affine fixed customer has a strictly favorable critical-to-Hodge source-angle reserve on the certified affine age range, and in particular for all sufficiently large affine ages. The restartable source-gradient/G4 comparison controls both critical and Hodge source polarizations by a perturbative error that tends to zero with the centered-action threshold. Choose the O3 threshold so that the total critical/Hodge polarization error is at most one half of the fixed affine margin. □

For $\iota \in \{c, 1\}$, write

$$\dot{\Pi}_{\mathrm{src},\iota} = \mathfrak{G}_\iota + \mathfrak{X}_\iota,$$

where

$$\mathfrak{G}_\iota := \mathrm{Cov}_{\sigma_{\iota,\mathrm{abs}}}(\mathcal{F}^q_\iota, \Sigma), \qquad \mathfrak{X}_\iota := \mathrm{Cov}_{\sigma_{\iota,\mathrm{abs}}}(\mathfrak{h}_{\mathrm{rot}}, \Sigma).$$

Then

$$\boxed{\dot{\Delta}_{\mathrm{src}} = (\mathfrak{G}_1 - \mathfrak{G}_c) + (\mathfrak{X}_1 - \mathfrak{X}_c).} \tag{K.66}$$

The geometry difference obeys the already proved bound

$$\boxed{|\mathfrak{G}_1 - \mathfrak{G}_c| \le \frac{21}{2}a + 18\|v - a\|_\infty + 4\|xv_x\|_\infty + 6\|\mathfrak{A}\|_\infty.} \tag{K.67}$$

Theorem K.25 (M4-E: nonsingular total-bank GREEN source-angle invoice)**.** *Let $t_s > t_d$ be the first time at which $\Delta_{\mathrm{src}}(t_s) = 0$, if such a time exists. Then*

$$\boxed{\begin{aligned}\delta_{\mathrm{src}} \le{}& \frac{21}{2}\Lambda(t_s; t_d) \\ &+ 18\int_{t_d}^{t_s} \|v - a\|_\infty \, dt \\ &+ 4\int_{t_d}^{t_s} \|xv_x\|_\infty \, dt \\ &+ 6\int_{t_d}^{t_s} \|\mathfrak{A}\|_\infty \, dt \\ &+ \int_{t_d}^{t_s} [\mathfrak{X}_c - \mathfrak{X}_1]_+ \, dt.\end{aligned}} \tag{K.68}$$

Thus if the four post-ticket geometry terms sum to less than $\delta_{\mathrm{src}}/2$, source-angle reserve loss forces

$$\boxed{\int_{t_d}^{t_s} [\mathfrak{X}_c - \mathfrak{X}_1]_+ \, dt > \frac{1}{2}\delta_{\mathrm{src}}.} \tag{K.69}$$

Proof. Integrate (K.66) from t_d to the first reserve-loss time t_s, use (K.65), then estimate the geometry difference by (K.67). The remaining adverse rotor term is precisely $[\mathfrak{X}_c - \mathfrak{X}_1]_+$. □

The rotor terms use only total absolute source-bank denominators, so the additive invoice remains nonsingular even when one source-sign bank is thin.

Remark K.26 (M3–M4 permanent scope firewalls). The following shortcuts remain forbidden:

(1) favorable source plus finite geometry action alone does not exclude one one-time lock;
(2) O1/O2 do not by themselves pay the reverse critical-to-Hodge visibility bill;
(3) there is no universal restarted absolute Hodge-source floor;

(4) visibility heterogeneity alone is not a payment until converted to source-angle, physical geometry, or response;
(5) the first O3 ticket itself is not a post-ticket debit.

M3–M4 funded interface. The permanent post-ticket chain is now

$$\boxed{\begin{aligned}&\text{O3 ticket} \Longrightarrow \text{OLD/NEW causal split} \Longrightarrow \text{Gate O/C/S},\\ &\quad\text{Gate S} \Longrightarrow \text{sign-conditioned visibility covariance} \Longrightarrow \text{source-angle/rotor covariance}\\ &\qquad\Longrightarrow \text{post-ticket geometry or total-bank GREEN ticket.}\end{aligned}} \tag{K.70}$$

M5–M6 must now route that fixed GREEN ticket through the ratio-free global source/Hodge coupling, continuation/energy structure, aspect geometry, and tangent deformation, leaving only the young inherited source-angle branch.

M5: ratio-free global GREEN barrier and bounded-age core. *Mutual pointers:* **M5**: grand-finale entry / short proof.

The total-bank GREEN ticket produced by M4 admits a ratio-free source-angle representation which remains regular when one source-sign bank is thin.

Let σ_c be the critical absolute source law,

$$\omega := W_1/W_c, \qquad \zeta := \langle\omega\rangle_{\sigma_c},$$

and let Π_c, Π_1 be the critical and Hodge source polarizations.

Theorem K.27 (M5-A: exact two-weight GREEN exchange)**.** *The critical-to-Hodge GREEN difference satisfies*

$$\boxed{\zeta(\mathfrak{X}_1 - \mathfrak{X}_c) = \mathrm{Cov}_{\sigma_c}\Big(\omega,\ H_c(\Sigma - \Pi_1)\Big).} \tag{K.71}$$

At source-angle reserve neutrality, the sign-restricted GREEN terms have the ratio-free two-copy representation

$$\boxed{\begin{aligned}S_\pm^2\mathfrak{H}_\pm = 2\varsigma_\pm \iint W_c(X)W_c(Y)[\omega(X)-\omega(Y)]\\ \times\,[q_\pm(Y)\mathcal{F}_{\mathrm{sh}}(X) - q_\pm(X)\mathcal{F}_{\mathrm{sh}}(Y)]\,dX\,dY.\end{aligned}} \tag{K.72}$$

Consequently, with $\Omega_\pm := \mathrm{osc}_{\mathrm{supp}\, q_\pm}\,\omega$ *and* $\mathcal{A}_\pm^{\mathrm{GREEN}} := \int_{\{q_\pm>0\}} W_c|\mathcal{F}_{\mathrm{sh}}|\,dX$,

$$\boxed{\zeta|\mathfrak{X}_c - \mathfrak{X}_1| \le \frac{4}{S^2}\left(S_+\Omega_-\mathcal{A}_-^{\mathrm{GREEN}} + S_-\Omega_+\mathcal{A}_+^{\mathrm{GREEN}}\right).} \tag{K.73}$$

No inverse sign-bank factor appears.

Proof. The Hodge source law is the ω-tilt of the critical source law. Centering the rotor field at the critical law and subtracting the two source-angle rotation rates gives (K.71). Expanding each within-sign covariance by the standard two-copy identity yields (K.72). The oscillation bound $|\omega(X)-\omega(Y)| \le \Omega_\pm$ gives the signwise estimate, and multiplication by the reserve-neutral prefactor $2S_+S_-/S^2$ cancels the apparent thin-bank denominators. □

Theorem K.28 (M5-B: large-age source-angle adversity pays non-affinity)**.** *Fix any classical restart state* t_0 *with* $\psi_\beta(t_0) > 0$ *and mixed-sign source. There exist finite constants*

$$L_{\mathrm{tail}}(t_0), \qquad M_{\mathrm{tail}}(t_0) > 0$$

such that

$$\boxed{\Lambda(t;t_0) \geq L_{\mathrm{tail}}(t_0), \quad \Pi_{\mathrm{src},1}(t) \leq \Pi_{\mathrm{src},c}(t) \Longrightarrow M(t_0,t) > M_{\mathrm{tail}}(t_0).} \tag{K.74}$$

Proof. For the exact affine evolution of every classical restart state,

$$\Pi^{\mathrm{aff}}_{\mathrm{src},1} - \Pi^{\mathrm{aff}}_{\mathrm{src},c}$$

is strictly positive after sufficiently large relative affine age. The restarted G4 comparison controls the true critical and Hodge source polarizations by errors tending to zero with the post-restart centered/source-gradient action M. Choose the affine age threshold first and then choose M_{tail} small enough that the two polarization errors are below the affine margin. A true-flow reserve loss at larger age must therefore violate that small-action threshold. □

Remark K.29 (M5-C: bounded-age crossing firewall). Under the natural bounded GREEN capacities, the complete differential GREEN exchange has zero collapse-age density. This does *not* exclude one finite reserve crossing: the exact local rotor falsifiers permit a one-time sign reversal with finite action. Therefore density estimates alone did not by themselves close the source-angle branch.

The closure-grade M5 endpoint is one bounded-relative-age source-angle crossing using the nonlocal Hodge/source-bank coupling and the signed differential GREEN exchange.

M6: global continuation, transported source potential, aspect geometry, and tangent sign-flip routing. *Mutual pointers:* **M6**: grand-finale entry / short proof.

Theorem K.30 (M6-A: global Hodge/swirl continuation routing)**.** *Let* $\Lambda(t;t_0) = \int_{t_0}^{t} a(s)\,ds$. *On every classical branch with nonzero swirl energy,*

$$\boxed{\Lambda(t;t_0) \leq C_{\mathrm{sw}}(t_0) + 2\int_{t_0}^{t} \|\nabla U(s)\|_\infty\,ds.} \tag{K.75}$$

Hence on a finite physical-time interval

$$\boxed{\Lambda(t;t_0) \to \infty \Longrightarrow \int_{t_0}^{T} \|\nabla U(t)\|_\infty\,dt = \infty.} \tag{K.76}$$

Proof. The seven-dimensional Hodge lift has energy $\mathscr{E}_{\mathrm{Hod}} = \frac{1}{2}\int H\chi\,d\mu_7$ and exact transport cancellation $\mathbf{b}\cdot\nabla\chi = \mathfrak{A}\chi$. Thus

$$\dot{\mathscr{E}}_{\mathrm{Hod}} = 2\int \chi q\,d\mu_7 = -\int \frac{x}{\sqrt{y}}\Theta v\,dX.$$

Together with the exact swirl-energy exchange and conserved physical kinetic energy, this gives

$$\Lambda(t;t_0) \leq C_{\mathrm{sw}}(t_0) + \int_{t_0}^{t} \|v-a\|_\infty\,ds.$$

Physical regularity gives $\|v-a\|_\infty \leq 2\|\nabla U\|_\infty$, proving (K.75). □

Thus the admitted finite physical-Lipschitz source-angle branch is automatically bounded in relative collapse age.

Define

$$\Gamma := xu, \qquad \mathcal{P}_\beta := x^2\Theta = \epsilon_\beta\Gamma^2 \geq 0.$$

Theorem K.31 (M6-B: transported circulation-square source potential)**.** *The circulation and circulation-square potential satisfy*

$$\boxed{\mathcal{D}_{xy}\Gamma = 0, \qquad \mathcal{D}_{xy}\mathcal{P}_\beta = 0.} \tag{K.77}$$

Moreover

$$\boxed{d\mathcal{P}_\beta = 2xyC_\beta\,dx + x^2 q\,dy,} \tag{K.78}$$

so

$$\boxed{\partial_y(2xyC_\beta) = \partial_x(x^2 q).} \tag{K.79}$$

Proof. The characteristic equations give $\mathcal{D}_{xy}x = -2xv$ and $\mathcal{D}_{xy}u = 2vu$, hence $\mathcal{D}_{xy}(xu) = 0$. Since ϵ_β is a fixed ancestry tag, $\mathcal{P}_\beta = \epsilon_\beta\Gamma^2$ is transported. Differentiating $\mathcal{P}_\beta = x^2\Theta$ and using $q = \Theta_y$ and $C_\beta = (\Theta + \frac{x}{2}\Theta_x)/y$ gives (K.78); mixed derivatives give (K.79). □

Corollary K.32 (Vertical source-sign order)**.** *At every fixed x and $y \geq 0$,*

$$\boxed{\int_0^y q_+(x,\eta)\,d\eta - \int_0^y q_-(x,\eta)\,d\eta = \Theta(x,y) \geq 0.} \tag{K.80}$$

Consequently every decreasing admissible vertical test f satisfies

$$\int_0^\infty f(y)q(x,y)\,dy \geq 0.$$

Let

$$\ell := \log(y/x).$$

The conserved circulation-square material law gives

$$\mathcal{D}_{xy}\ell = 6a + 6(v-a) - 2\mathfrak{A}.$$

Theorem K.33 (M6-C: material aspect translation with controlled distortion)**.** *Let $\mathcal{L}_t$ be the law of ℓ under the conserved circulation-square material probability $\nu_\Gamma(t)$. Then*

$$\boxed{W_\infty\left(\mathcal{L}_t,\ \tau_{6\Lambda(t;t_0)}\mathcal{L}_{t_0}\right) \leq D_{\mathrm{shape}}(t_0,t),} \tag{K.81}$$

where

$$\boxed{D_{\mathrm{shape}}(t_0,t) := 6\int_{t_0}^t \|v-a\|_\infty\,ds + 2\int_{t_0}^t \|\mathfrak{A}\|_\infty\,ds.} \tag{K.82}$$

Proof. Along each material label,

$$\ell(t) - \ell(t_0) - 6\Lambda(t;t_0) = \int_{t_0}^t (6(v-a) - 2\mathfrak{A})\,ds.$$

The absolute value is bounded by D_{shape}. Coupling each label with itself between the initial and final conserved material laws gives the W_∞ estimate. □

Write the regular equatorial swirl profile as

$$u(x,y,t) = y\mathfrak{w}(x,y,t).$$

Theorem K.34 (M6-D: bounded-age moderate capital forces extreme-aspect debt)**.** *Fix a classical launch time t_0. Assume on $[t_0, T]$:*

$$0 \le \Lambda(t; t_0) \le L_*, \qquad \|\nabla_{x,y}\mathfrak{w}(t)\|_\infty \le L_w^*.$$

Then the detector law

$$\psi_\beta(t) = \psi_\beta(t_0)e^{-2\Lambda(t;t_0)}$$

provides a fixed positive source-potential capital on a universal moderate-aspect band. Consequently, if the critical-to-Hodge source-angle reserve is lost at some T, then there exist constants $W^*_{\mathrm{low}}, W^*_{\mathrm{high}} > 0$ and $s_R > 1$, depending only on the launch data and the displayed caps, such that at least one of

$$\boxed{\int_{y<x/2} \mathfrak{w}^2\,dx\,dy \ge W^*_{\mathrm{low}} > 0, \qquad \int_{y>s_R x} \mathfrak{w}^2\,dx\,dy \ge W^*_{\mathrm{high}} > 0} \tag{K.83}$$

holds. Moreover, if the required extreme-aspect concentration is old relative to the launch stock, then

$$\boxed{2\Lambda(T; t_0) \le \log \frac{\mathfrak{W}_2(t_0)}{W_{\mathrm{req}}} + 2\mathfrak{V}(t_0, T) + 3\mathfrak{A}_{\mathrm{act}}(t_0, T).} \tag{K.84}$$

Proof. Bounded age gives the detector floor $\psi_\beta(t) \ge \psi_\beta(t_0)e^{-2L_*}$ and hence a positive origin value of $\mathfrak{w}$. The first-jet cap propagates this lower bound over a fixed moderate-aspect rectangle. The exact one-dimensional aspect-potential representation of the source-angle reserve has a uniformly favorable kernel on that band, so reserve loss requires the adverse low- or high-aspect tails to overpower the fixed capital, giving (K.83). The exact source-profile envelope then gives (K.84). □

Let M denote the affine-normalized tangent deformation matrix and define

$$\kappa_\varepsilon := \frac{\varepsilon}{1-\varepsilon}.$$

Theorem K.35 (M6-E: tangent sign-flip / horizontal-launch bound)**.** *If*

$$\|M - I\| \le \varepsilon < 1,$$

then every source-sign-changing label must originate in the launch wedge

$$\boxed{|q_0| \le \kappa_\varepsilon \sqrt{s_0}\,|\widehat{C}_{\beta,0}|.} \tag{K.85}$$

If

$$\mathcal{H}_c(t_0) := 2\int W_c \sqrt{s_0}\,|\widehat{C}_{\beta,0}|\,dX,$$

then the critical source mass of either sign eligible to flip obeys

$$\boxed{\mathcal{S}^{\mathrm{flip}}_{c,\pm}(t_0) \le \kappa_\varepsilon \mathcal{H}_c(t_0).} \tag{K.86}$$

Proof. The transported-potential Cauchy formula expresses the current source-gradient vector as the affine background map composed with M acting on the launch vector. If $\|M - I\| \le \varepsilon$, a change in the sign of the vertical source-gradient component is possible only when the launch vertical component is within κ_ε times the horizontal component, giving (K.85). Integrating the pointwise wedge bound against the critical source weight yields (K.86). □

Theorem K.36 (M6-F: source-angle master alternative)**.** *Every critical-to-Hodge source-angle reserve loss after a classical launch time t_0 must enter at least one of:*

(1) large-age post-restart non-affinity/source-gradient payment;
(2) source-profile first-jet continuation exit;
(3) fixed small- or large-aspect source-profile concentration;
(4) fixed centered value/axial payment relative to the launch source-profile stock;
(5) non-small tangent deformation;
(6) a macroscopic near-horizontal source-gradient orientation stock already present at the launch.

On the finite physical-Lipschitz branch the crossing is collapse-age young. If the first five alternatives are kept below their funded thresholds, the remaining obstruction is inherited launch geometry of the form

$$\boxed{\begin{array}{c}\textit{extreme-aspect source-profile stock}\\ \textit{and/or near-horizontal source-gradient orientation stock.}\end{array}} \tag{K.87}$$

Proof. Large-age adversity is routed by theorem K.28; on the finite physical-Lipschitz branch large age is excluded by theorem K.30. On the remaining bounded age branch, failure of the source-profile first-jet cap is Alternative 2. If the cap holds, theorem K.34 yields Alternative 3, while old persistence of that concentration yields Alternative 4. Independently, theorem K.35 shows that macroscopic source-sign reorganization requires Alternative 5 or the inherited horizontal-orientation stock in Alternative 6. These alternatives exhaust the permanent M5–M6 routes. □

Remark K.37 (Inherited-launch source-angle endpoint). At the preferred launch $t_0 = t_d$, the old Hodge-bank gap is already funded and the O3 ticket may not be reused. Module M6 reduces every surviving bounded-age reserve loss to inherited extreme-aspect source-profile stock and/or near-horizontal source-gradient orientation stock. The completed inherited-launch audit shows that this ticket-state stock is quantitatively subcritical for theorem K.36, without recharging pre-threshold O3 action. No long-time, density, entropy, event-count, or arbitrary local-rotor loophole remains as an independent source-angle branch.

Remaining open core. The permanent M1–M6 chain and the completed inherited-launch audit close the collapse-age-young source-angle geometry and fund O4 without recharging the O3 ticket. There is therefore no remaining geometric O6.1 obligation.

The sole open proposition is O6.2. On every post-O4 no-exit branch it asks for a finite maintained-cashier launch whose net gain

$$\mathcal{G}_{\mathrm{cash}}(\lambda) = \frac{\lambda}{4} - \mathfrak{D}_{\mathrm{rate}}(\lambda)$$

satisfies

$$\int_0^\infty e^{-\mathcal{G}_{\mathrm{cash}}(\lambda)}\, d\lambda < \infty.$$

The M7 powered, positive-bank-growth, and fixed sub-quarter cashier conditions remain stronger sufficient routes to this terminal conclusion; M8 is the exact proved countdown once the O6.2 integrability activation is supplied.

Permanent module map. The reconstructed post-ticket proof architecture is organized into eight proved modules:

M1. polarization/source-history structure theorems K.11 to K.13;
M2. critical $\varpi = 2/5$ source-positive cashier and fractional homotopy theorems K.14, K.15 and K.17;

M3. post-ticket causal pay-or-route/no-double-charge module theorems K.19 and K.20;
M4. Gate-S / critical-to-Hodge source-angle GREEN bridge theorems K.21 to K.23 and K.25;
M5. ratio-free GREEN barrier and bounded-age crossing theorems K.27 to K.29;
M6. global continuation/source-potential/aspect/tangent module theorems K.30, K.31 and K.33 to K.36;
M7. powered-reserve / positive-bank-growth / maintained-cashier terminal module theorems K.1, K.3 and K.4;
M8. exact post-O4 integral cashier countdown theorems K.5 and K.6.

The minimal dependency graph is

$$\boxed{\begin{aligned} O1\text{–}O3 &\Longrightarrow M1 \Longrightarrow M2 \Longrightarrow M3 \Longrightarrow M4,\\ M4 &\Longrightarrow (M5+M6) \Longrightarrow \text{closed O6.1 audit} \Longrightarrow O4,\\ M7 &\Longrightarrow M8,\\ O4+O6.2+M8 &\Longrightarrow O5. \end{aligned}} \tag{K.88}$$

The inherited-launch source-angle module O6.1 is closed in the present audit.
Audit status. **SOLE OPEN PROPOSITION—NO PROOF ASSERTED: O6.2.** O6.2 is the exact post-O4 integral cashier activation. M7-P/C1/C2 remain stronger sufficient routes, and M8 records the weakest proved terminal countdown.

Appendix L. Grand-finale assembly

The funded base through O1–O3 is proved. The permanent modules M1–M6 and the completed inherited-launch audit close the source-angle branch and fund O4. M7 contains stronger terminal routes, while M8 is the exact integral cashier countdown. O5 is GREEN downstream from O4+O6.2+M8, and O6.2 is the sole open package.

L1. Funded chain up to the open bill. *Mutual pointers:* grand-finale entry / short proof/status / main statement theorem 24.1.

> **Restatement of theorem 24.1.** For the fixed $A = 10$ customer, every implication in rows A–I of section 23 is proved. In particular,
>
> $$\boxed{\begin{aligned} &\text{fixed tagged customer} \Longrightarrow (\Theta, q, C_\beta) \Longrightarrow H^{\pm} \Longrightarrow a^{\pm},\\ &\text{affine fixed-seed benchmark} \Longrightarrow \upsilon_{-}^{\mathrm{aff}} \le \upsilon_{\mathrm{aff}}^{*} < 1,\\ &\text{small centered true-Euler deformation} \Longrightarrow a \ge \pi_{107}(a^{+}+a^{-}),\\ &\text{first polarization loss} \Longrightarrow \text{definite star-hull/tangent bill},\\ &\text{near-lock conversion} \Longrightarrow \text{compact mesoscopic rotor ticket},\\ &\text{fresh primary conversion} = O(h^2)a^{+},\\ &\text{negative self-recycling} \Longrightarrow \text{finite causal resolvent},\\ &\text{O1–O3 physical-bank closure} \Longrightarrow \text{strict bank gap or paid centered deformation}. \end{aligned}}$$
>
> No generic swirl-free Euler estimate is used as terminal payment, and one physical conversion is never charged twice.

Proof. We verify the displayed chain in the order in which money enters and leaves the ledger.

Step 1: fixed customer, tagged source, genealogy, and Hodge banks. The exact squared-coordinate equations and pressure-free source-to-vorticity identity are propositions 4.3 and 5.3. The provenance normal form and two-shadow firewall propositions 8.2 and 8.4 keep the terminal source equal to the fixed tagged quantity $\Theta = \varepsilon_\beta u^2$ and forbid swirl-free background terms from becoming source credit. The exact two-bank Duhamel equations proposition 10.2 give $H^\pm \geq 0$, and the positive whole-space Hodge cashier proposition 5.4 gives $a^\pm$ with $a = a^+ - a^-$. This proves the first line of the funded chain.
Step 2: affine benchmark and nonlinear stability. The two-time vertical response is exact by proposition 15.2. For the fixed $A = 10$ customer, lemma 15.4 and theorem 15.5 prove $\upsilon_-^{\mathrm{aff}} \leq \upsilon_{\mathrm{aff}}^* < 1$. The fixed-seed source-gradient condition number lemma 17.4 and the complete-response stability theorem theorem 17.5 imply the stated true-Euler polarization inequality while the centered action is below its admitted threshold. At the first strict loss, corollary 17.6, together with propositions 9.4 and 9.6, returns a definite star-hull/tangent deformation bill.
Step 3: conversion is a compact rotor event. The loss-of-affine-polarization theorem and protected apex tube theorem 15.6 and proposition 15.7 show that near-lock requires genuine adverse tagged conversion outside the fresh tube. The exact source-gradient system, rotor normal form, and Hodge-skew cancellation propositions 16.1 and 16.3 and theorem 16.4 identify this conversion as rotation/deformation of the single transported circulation gradient. The angular-tail suppression and compact-ticket theorem proposition 16.8 and theorem 16.9 remove both angular tails, the protected apex, and the far radial tail. Hence every substantial new adverse conversion carries a compact mesoscopic rotor ticket.
Step 4: fresh primary conversion and negative self-recycling. The exact generated-negative response is theorem 18.4. Fresh apex anti-rotation and the small-window estimate propositions 18.3 and 18.5 give the displayed $O(h^2)a^+$ law. Negative self-recycling is not iterated by hand: theorem 18.6 resums the complete $H^- \to H^-$ branch into the finite positive kernel $\mathcal{G}_{\mathrm{rec}}$, and proposition 18.7 leaves only old initial ancestry, signed $\mathfrak{T}$ input, and primary $H^+ \to H^-$ conversion upstream.

Split the primary source-time integral at $(t-h)_+$. By proposition 18.5 and definition 21.1, for $t \geq h$,

$$\boxed{\mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{prim}}(t) = \mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{fr,h}}(t) + \mathfrak{X}^{\mathrm{ex}}_{+,\mathrm{old,h}}(t).} \tag{L.1}$$

If $0 < t < h$, the old integral is empty and the same decomposition is read with fresh window $h' = t$. The exact fresh term is bounded by the already proved fresh $O(h^2)a^+$ estimate in proposition 18.5, while the second term is exactly the repaired old-primary invoice displayed in the theorem. Thus no other new adverse account remains.
Step 5: legality of the transfers. The adverse-source ancestry theorem proposition 13.1 allows only initial negative slope or later tagged cross-shear ancestry. Weighted propagation, physical Biot–Savart control, and the continuation firewall propositions 14.2, 14.4 and 14.5 and theorem 14.6 route ordinary growth or far-field failure to named accounts. The exact supporting identities I1–I7, especially the source-time genealogy proposition 19.5 and the no-double-charge theorem proposition 19.6, ensure that each physical deposition is charged once. Finally J1–J4 rule out replacing this funded chain by local finite action, frozen Hodge geometry, refresh counting, or a factored pointwise reproduction number. Therefore every implication asserted in rows A–I is funded as stated. The exact physical-bank formulation closes O1–O3. the closed inherited-launch audit gives O4, O5 is GREEN from O4+O6.2+M8, and O6.2 is the sole unpaid proposition. □

Audit status: funded base and downstream O-ledger are complete; O6.2 is the sole open package.

L2. Grand finale. *Mutual pointers:* grand-finale entry / short proof/status / main statement conditional proposition 24.2.

> **Restatement of conditional proposition 24.2.** Assume the sole remaining open proposition open proposition 21.8. Then either a named classical continuation exit occurs earlier, or the fixed customer loses classical regularity in finite time.

Proof. Work on the maximal classical branch of the fixed customer. If a named continuation exit occurs earlier, the first alternative holds.

Assume no such exit occurs and assume O6.2. The completed inherited-launch source-angle audit already gives O4 by theorem 21.6. O6.2 activates the exact M8 integral cashier countdown, and O4+O6.2+M8 give O5 by theorem 21.7. Hence the fixed customer leaves the classical category in finite physical time. □

Audit status: GREEN/COMPLETE conditional chain; sole open hypothesis is O6.2.

California, United States
Email address: ymshi@protonmail.com