arXiv:submit/8081412 [nlin.SI] 15 Sep 2026

AXISYMMETRIC NAVIER–STOKES FLOW WITH SWIRL: A POLARIZED MERIDIAN-PLANE PROOF ARCHITECTURE

YAOMING SHI

Abstract. We introduce a new way to estimate the total cost of repeated small-scale generation in axisymmetric Navier–Stokes flow. The Euler background is deliberately bypassed through successive polarization steps, so that the final accounting is tied specifically to positive viscosity $\nu > 0$ rather than to effects already present in Euler flow.

We apply this Euler firewall to smooth axisymmetric three-dimensional Navier–Stokes flow with swirl in $\mathbb{R}^3$ and $\mathbb{R}^2 \times \mathbb{T}_L$. Geometry-dependent arguments first produce compact signed toroidal-vorticity packets with a strict parent–child scale gap. A common downstream mechanism then uses material vorticity, the full viscous vorticity adjoint, material localization, and a primal–adjoint boundary current to show that each continuing active thread either moves to a strict physical successor or pays a fixed positive amount of Leray dissipation. Throughout this paper, by a *clock* we mean a bookkeeping device that selects an ordered sequence of times or time blocks according to the accumulation of a specified control quantity; it is not an additional physical variable and by itself carries no terminal cost. The whole-space route derives its endpoint clock from a radial-strain continuation criterion and a weighted Hodge comparison; the periodic route instead uses a recurrent axis point, one-sided local Type-II input, and periodic Hodge reselection. Thus the two geometries have different upstream scale-selection mechanisms but share the same viscosity-polarized terminal argument. Euler-compatible identities are used only for routing; terminal accounting begins only after the Euler background has been canceled and viscosity appears explicitly.

Contents

1. Introduction and compressed assembly statements 2
2. Meridian variables, pole parity, and viscosity-visible identities 5
2.1. Signed meridian variables 5
2.2. Finite physical ledgers and the reservoir-selection rule 7
2.3. The universal even-quotient record barrier 7
3. Whole-space strain endpoint and endpoint-clock recovery 8

Date: Version 82C-single-42C of September 15, 2026.

2020 *Mathematics Subject Classification.* Primary 35Q30; Secondary 76D03, 76D05, 35B65.

Key words and phrases. axisymmetric Navier–Stokes equations, swirl, vorticity, meridian-plane reduction, material flow, viscous adjoint, Euler firewall, nodal flux.

3.1. Common physical continuation package 8
3.2. SC.1: a whole-space radial-strain continuation criterion 11
3.3. Euler-compatible clock conversion 12
4. Geometry-specific upstream bridges 14
4.1. Whole-space endpoint clock and synchronization 18
4.2. Whole-space compact source-work bridge 25
4.3. Active periodic strain clock and recurrent local selector 33
4.4. Periodic active mode and the same-scale Hodge detector 37
5. Bridge D: compact signed child to active-origin alternatives 46
5.1. Creation–Robin identity 46
5.2. Logarithmic action-depth obstruction 47
5.3. Source-born versus bare active resets 49
6. Bridge E: successive polarization of the common downstream 52
6.1. Axis-away detector extraction 52
6.2. First polarization: Cauchy material vorticity 53
6.3. Second polarization: the full viscous vorticity adjoint 54
6.4. Material localization and the no-teleportation alternative 54
6.5. Material geometry is a router, not an extra hypothesis 59
6.6. Primal–adjoint polarization and the boundary Wronskian current 59
6.7. Charged persistence and a fixed Leray payment 61
6.8. Active-order firewall: bounded jets, spatial factorial growth, and temporal spectators 62
6.9. Automatic deep handoff and chronological reselection 67
7. Proof architecture 72
7.1. Whole-space chain 72
7.2. Periodic-cylinder chain 76
8. Conclusion 77
9. Acknowledgements 78
Acknowledgements 78
10. References 79
References 79

1. Introduction and compressed assembly statements

The mathematical theory of the three-dimensional incompressible Navier–Stokes equations originates in Leray's weak-solution framework [19]. Standard monographs and broad treatments include Ladyzhenskaya [17], Temam [31], Constantin–Foias [11], Doering–Gibbon [12], Majda–Bertozzi [21], and Robinson–Rodrigo–Sadowski [24]; see also Ladyzhenskaya's survey of the millennium problem [18]. Among the classical regularity landmarks are Serrin's interior criterion [26], the Caffarelli–Kohn–Nirenberg partial-regularity theorem [3], and the endpoint $L_{3,\infty}$ result of Escauriaza–Seregin–Šverák [13]. Tao's quantitative reworking of the critical L^3 theory gives explicit

higher-regularity bounds in terms of the critical norm [30]. In the closely related three-dimensional Euler theory, the Beale–Kato–Majda criterion identifies the time-integrated L^∞ vorticity as a breakdown quantity [2]. Tao's finite-time blowup construction for an *averaged* three-dimensional Navier–Stokes nonlinearity [29] further shows that the energy cancellation by itself is not sufficient to force regularity for that broader averaged class, thereby emphasizing the importance of the specific structure of the Navier–Stokes nonlinearity.

Fefferman's official formulation of the Clay Millennium problem asks for a proof of one of four alternatives: unforced global smoothness in $\mathbb{R}^3$ or on the full periodic torus, denoted (A) and (B), or forced breakdown in the corresponding two geometries, denoted (C) and (D) [14]. On September 8, 2026, OpenAI announced and released a construction establishing the forced breakdown alternatives (C) and (D) [22]; the Clay Mathematics Institute subsequently stated that the problem had "apparently been settled," while emphasizing that its evaluation and credit process is deliberately unhurried [10]. These developments do not logically negate the unforced alternatives: in Fefferman's formulation (A)/(B) set $f \equiv 0$, whereas (C)/(D) permit a smooth external force. Thus (A) and (C), and likewise (B) and (D), are compatible statements. The present program is directed toward the unforced regularity side. Its whole-space branch lies in the axisymmetric subclass of (A); the axial-periodic cylinder $\mathbb{R}^2 \times \mathbb{T}_L$ should be viewed as a periodic analogue of (B), not as the full three-torus geometry appearing literally in Fefferman's statement.

The three-dimensional axisymmetric Navier–Stokes equations with swirl retain several genuinely three-dimensional mechanisms while reducing the geometry to a meridian half-plane. Classical axisymmetric work goes back at least to Ukhovskii–Yudovich [32]. The interaction between meridional strain, swirl and azimuthal vorticity has since been a natural testing ground for regularity criteria and singularity analysis; representative points of reference include [6, 20, 7, 8, 9, 25]. In the present paper we consider both

$$\Omega_\infty := \mathbb{R}^3, \qquad \Omega_L := \mathbb{R}^2 \times \mathbb{T}_L,$$

and organize the argument so that the geometry-dependent part is separated from a common downstream mechanism.

One motivation for the choice of variables comes from the structural analogy with the two-dimensional Boussinesq equations. Global regularity in important dissipative and partially dissipative Boussinesq regimes was established by Cannon and DiBenedetto [4], Chae [5] and Hou–Li [16]; see also Wu's lecture notes [33]. A related inviscid Boussinesq/axisymmetric-Euler comparison is developed in [27]. We use this analogy only as motivation: no Boussinesq regularity theorem is transferred to the axisymmetric Navier–Stokes problem. Instead, the proof works directly with the axis-regular

variables

$$\alpha = -\frac{u^r}{r}, \qquad \beta = \frac{u^\theta}{r}, \qquad q = \omega^\theta, \qquad G = \frac{q}{r},$$

introduced in section 2, together with the mandatory pole parity of smooth three-dimensional axisymmetric fields.

The organizing principle is an *Euler firewall*. Identities or scale-invariant quantities that survive unchanged when $\nu = 0$ may be used to select clocks, scales, packets or restarts, but not as the terminal Navier–Stokes closure. The terminal branch must retain either an explicit positive factor of ν or a charge against a finite physical viscous ledger. In the whole space, sections 3 and 4 develop the strain clock, center-free restart, synchronization and Hardy/source-work route. In the periodic cylinder, section 4 instead uses a recurrent axis point, one-sided local Type-II input, strain selection and a periodic Hodge reselection. From section 5 onward the two geometries share the same active-origin reduction; section 6 then performs the successive Cauchy/material/adjoint polarizations which isolate the viscosity-visible downstream cost.

The standing data classes and the two compressed assembly statements are given below. For a quick map of the proof, section 7 contains the synchronized two-page comparison: the first page aligns the two geometry-specific upstream routes row by row, and the second page displays the common Bridges D–E and the funded downstream alternatives. This is the most economical roadmap for the detailed argument that follows.

The organizing principle is to use Euler-compatible identities only for routing and reserve terminal closure for quantities whose decisive imbalance retains an explicit positive factor of viscosity.

Definition 1.1 (Standing data classes)**.** *Fix $\nu > 0$; in the periodic geometry also fix $L > 0$. Whenever $T < \infty$ is discussed as a candidate first singular time, u denotes the corresponding classical axisymmetric solution on $[0, T)$.*

(H_∞). *The whole-space datum u_0 is smooth, divergence-free, axisymmetric with swirl, and rapidly decaying on $\mathbb{R}^3$, with*

$$u_0 \in H^2(\mathbb{R}^3), \qquad \Gamma_0 = r u_0^\theta \in L^4(\mathbb{R}^3), \qquad r^3 \omega_0^\theta \in L^2(\mathbb{R}^3).$$

(H_L). *The periodic datum u_0 is smooth, divergence-free, and axisymmetric on $\Omega_L = \mathbb{R}^2 \times \mathbb{T}_L$, with finite energy per period and radial decay sufficient for the cutoff-exhaustion arguments used below,*

$$u_0 \in H^2(\Omega_L), \qquad \Gamma_0 = r u_0^\theta \in L^4(\Omega_L), \qquad r^3 \omega_0^\theta \in L^2(\Omega_L).$$

Remark 1.2 (Auxiliary hypotheses not assumed). Smooth three-dimensional axisymmetry supplies the mandatory signed-r pole parity used below. Neither data class assumes z-reflection symmetry. The proof also does *not* assume $\Gamma_0 \in L^\infty$, any post-T suitable/Leray–Hopf continuation, or the auxiliary periodic flat-q^2/A.2 clock. Those auxiliary hypotheses and routes are not used in the assembly statements.

Theorem 1.3 (Whole-space compressed downstream assembly)**.** *Assume* (H_∞). *If $T < \infty$ is a candidate first singular time, the internal whole-space strain endpoint and weighted Hodge comparison proved below supply the canonical future flat-q^2 clock used by theorems 4.10 and 4.13. Whenever the geometry-specific whole-space bridges theorems 4.10, 4.13 and 4.21 produce a compact signed child, the common bridges theorems 5.11 and 6.29 reduce every continuing active terminal thread to the following alternative:*

(i) a strict physical successor is selected and is absorbed into a later canonical block;

(ii) a detector-charged spacetime packet pays a fixed positive Leray dissipation amount.

The two outcomes can be selected chronologically, with paid packets pairwise disjoint in time along any continuing active thread.

Theorem 1.4 (Periodic-cylinder compressed downstream assembly)**.** *Assume* (H_L). *Whenever the active periodic strain-clock/Type-II chain and the compact interfaces theorems 4.36 and 4.39 produce a compact signed child, the same common bridges theorems 5.11 and 6.29 give the alternative in theorem 1.3, with the same chronological disjointness of the paid packets. No additional common downstream conditional leaf is required.*

2. Meridian variables, pole parity, and viscosity-visible identities

2.1. **Signed meridian variables.** Write

$$\mathcal{Q} := \{(r,z) : r \geq 0,\ z \in \mathbb{R}\}, \qquad \mathcal{Q}_L := \{(r,z) : r \geq 0,\ z \in \mathbb{T}_L\}.$$

The flat Chae–Lee [6] measure is $dr\,dz$; physical three-dimensional integration carries the additional orbit factor $2\pi r$.

Throughout the paper we use the norm convention

$$\boxed{\|f\|_p := \left(\int_{\mathcal{Q}_*} |f(r,z)|^p\, r\, dr\, dz\right)^{1/p},} \tag{2.1}$$

where $\mathcal{Q}_*$ denotes $\mathcal{Q}$ or $\mathcal{Q}_L$ according to the geometry. Thus an unmarked $\|\cdot\|_p$ is always a *physical meridian* norm. For an axisymmetric scalar this differs from the full three-dimensional $L^p(dX)$ norm only by the constant factor $(2\pi)^{1/p}$. By contrast, every Chae–Lee flat norm is written explicitly as

$$\|f\|_{L^p(dr\,dz)} \quad \text{or} \quad \|f\|_{\widetilde{L}^p(dr\,dz)}.$$

Norms in material or label variables will likewise display their measure explicitly, for example $L^2(da)$ or $L^2(dy)$.

Define [16]

$$\boxed{\alpha := -\frac{u^r}{r}, \qquad \beta := \frac{u^\theta}{r}, \qquad q := \omega^\theta, \qquad G := \frac{q}{r}.} \tag{2.2}$$

and

$$\boxed{A := \omega^z = 2\beta + r\beta_r, \qquad W := -\omega^r = r\beta_z, \qquad \Gamma := ru^\theta = r^2\beta.} \tag{2.3}$$

Let $b = (u^r, u^z)$ and

$$\mathcal{D} := \partial_t + b \cdot \nabla, \qquad \mathcal{L}_1 := \partial_r^2 + \frac{3}{r}\partial_r + \partial_z^2.$$

Then

$$\boxed{\begin{aligned}(\mathcal{D} - \nu\mathcal{L}_1)\beta &= 2\alpha\beta,\\ (\mathcal{D} - \nu\mathcal{L}_1)G &= (\beta^2)_z.\end{aligned}} \tag{2.4}$$

and

$$\boxed{\partial_t q + \operatorname{div}_{r,z}(bq) = \nu\left(q_{rr} + \frac{1}{r}q_r + q_{zz} - \frac{q}{r^2}\right) + \partial_z(r\beta^2).} \tag{2.5}$$

The circulation is passive:

$$\boxed{(\partial_t + b \cdot \nabla)\Gamma = \nu\left(\Gamma_{rr} - \frac{1}{r}\Gamma_r + \Gamma_{zz}\right), \qquad \Gamma_r = rA, \qquad \Gamma_z = rW.} \tag{2.6}$$

Smooth three-dimensional axisymmetry imposes mandatory pole parity. The signed radial extensions satisfy [20]

$$\boxed{\begin{array}{c|c}\text{odd in } r & u^r, u^\theta, \omega^r, q, W\\ \hline \text{even in } r & u^z, A, p, \alpha, \beta, G\end{array}} \tag{2.7}$$

so

$$q(0,z,t) = W(0,z,t) = u^r(0,z,t) = u^\theta(0,z,t) = 0. \tag{2.8}$$

This is the reason the terminal proof uses the regular even variables α, β, G, A and strain rather than a fictitious nonzero axis value of q.

Remark 2.1 (Historical z-reflection specialization versus the current scope). The mandatory signed-r pole parity (2.7) is logically distinct from the optional z-reflection specialization used during part of the proof laboratory. In the historical source convention that specialization is

$$\begin{aligned}&u^r(r,-z,t) = u^r(r,z,t), \qquad u^\theta(r,-z,t) = u^\theta(r,z,t),\\ &u^z(r,-z,t) = -u^z(r,z,t).\end{aligned} \tag{2.9}$$

with $p(r,-z,t) = p(r,z,t)$. Taking curl gives

$$\boxed{\begin{array}{c|c}\text{even in } z & A = \omega^z,\ \alpha,\ \beta,\ \Gamma\\ \hline \text{odd in } z & q = \omega^\theta,\ W = -\omega^r,\ G\end{array}} \tag{2.10}$$

(and the historical stream scalar is odd in z).

Earlier formulations used (2.9) to split a full-space source tax equally between two reflected lobes. The present short spine does not use that lobe splitting. In particular, theorems 4.15 to 4.17 use only full-space Hardy quantities and the global Hardy identity, while theorems 4.18 and 4.19 use

the positive work density $(\widehat{q}\,\widehat{S}_\beta)_+$ and an arbitrary axial center. Hence (2.9) is provenance, not a hypothesis of the current whole-space chain.

2.2. **Finite physical ledgers and the reservoir-selection rule.** For either axial geometry, the physical energy inequality gives

$$\frac{1}{2}\|u(t)\|^2_{L^2(dX)} + \nu\int_0^t \|\nabla u(s)\|^2_{L^2(dX)}\,ds \le \frac{1}{2}\|u_0\|^2_{L^2(dX)}. \tag{2.11}$$

For axisymmetric fields,

$$\frac{1}{2\pi}\|\omega(t)\|^2_{L^2(dX)} = \int_{\mathcal{Q}_*} r\big(A^2+q^2+W^2\big)\,dr\,dz, \tag{2.12}$$

where $\mathcal{Q}_* \in \{\mathcal{Q}, \mathcal{Q}_L\}$. We shall also use the finite bulk swirl-pressure ledger

$$\boxed{\int_0^T\int_{\mathcal{Q}_*} P_\theta(r,z,t)\,dr\,dz\,dt \le C\int_0^T \|\nabla u(t)\|^2_{L^2(dX)}\,dt < \infty.} \tag{2.13}$$

The bound (2.13) is the cylindrical Hardy inequality applied to $u^\theta(0,z,t)=0$, followed by integration in z and t. Thus all arguments which claim a terminal Navier–Stokes payment must retain an explicit positive factor of ν or a known summable physical ledger. A scale-invariant or monotone quantity whose governing identity survives unchanged at $\nu = 0$ is used only for routing.

2.3. **The universal even-quotient record barrier.**

Lemma 2.2 (Universal even-quotient Euler firewall)**.** *Let F be a smooth even-in-r scalar satisfying*

$$(\mathcal{D}-\nu\mathcal{L}_1)F = S_F. \tag{2.14}$$

Set $H_F := rF^2$. Then

$$\boxed{(\mathcal{D}-\nu\Delta_0)H_F = 2rFS_F - \alpha H_F - 2\nu r|\nabla_{r,z}F|^2 - \frac{\nu}{r^2}H_F,} \tag{2.15}$$

where $\Delta_0 = \partial_r^2 + r^{-1}\partial_r + \partial_z^2$. At a fresh positive interior record (r_*, z_*, t_*) of H_F, $r_* > 0$,

$$\boxed{2\frac{S_F}{F} - \alpha \ge \frac{3\nu}{2r_*^2}.} \tag{2.16}$$

Proof. The product identity

$$\Delta_0(rF^2) = 2rF\mathcal{L}_1F + 2r|\nabla_{r,z}F|^2 + \frac{F^2}{r}$$

gives (2.15). At a positive record, $\nabla_{r,z}H_F = 0$ implies $F_z = 0$ and $F_r = -F/(2r_*)$, hence $|\nabla F|^2/F^2 = 1/(4r_*^2)$. Since $\partial_t H_F \ge 0$ and $\Delta_0 H_F \le 0$, division by $H_F > 0$ gives (2.16). □

Three instances will be used below:

$$\boxed{\begin{aligned} F=\beta:&\qquad \alpha \ge \frac{\nu}{2r^2},\\ F=\beta_z:&\qquad 2\frac{A\alpha_z}{\beta_z}-\alpha-2r\alpha_r \ge \frac{3\nu}{2r^2},\\ F=G:&\qquad 2\frac{(\beta^2)_z}{G}-\alpha \ge \frac{3\nu}{2r^2}. \end{aligned}} \tag{2.17}$$

These barriers vanish in the Euler limit and are therefore terminally Navier–Stokes specific.

3. Whole-space strain endpoint and endpoint-clock recovery

The former whole-space condition C−WC is now a theorem. The proof has two logically different parts. First, viscosity supplies a whole-space radial-strain continuation criterion. Second, a weighted Hodge estimate, which is Euler-compatible, converts the resulting strain clock into the historical flat-q^2 clock. By the Euler firewall, the second step is used only for chronology and routing; the terminal detector remains Bridge E.

3.1. **Common physical continuation package.**

The two axial geometries use the same final continuation mechanism. We record it once so that neither SC.1 nor SC.2 depends on a hidden restart theorem. All unmarked norms in this subsection are physical meridian norms in the sense of (2.1), unless a full dX norm is displayed explicitly.

Lemma 3.1 (Uniform physical-H^1 restart)**.** *Let Ω_* denote either $\mathbb{R}^3$ or $\Omega_L=\mathbb{R}^2\times\mathbb{T}_L$. Fix $\nu>0$ and $M<\infty$ (and L in the periodic case). There exists $\delta=\delta(\nu,M)>0$ in the whole space and $\delta=\delta(\nu,L,M)>0$ in the periodic cylinder such that every divergence-free datum $a\in H^1(\Omega_*)$ with*

$$\|a\|_{H^1(\Omega_*)}\le M$$

generates a unique strong solution on $[0,\delta]$ satisfying

$$v\in L^\infty(0,\delta;H^1(\Omega_*))\cap L^2(0,\delta;H^2(\Omega_*)).$$

The lifespan is uniform on the indicated H^1 ball.

Proof. Use the Fourier–Friedrichs cutoff and Leray projection (periodic Fourier series in the axial variable in the cylinder). Taking the physical $L^2(dX)$ inner product of the regularized equation with $-\Delta v$ gives

$$\frac12\frac{d}{dt}\|\nabla v\|_2^2+\nu\|\Delta v\|_2^2\le \|v\|_6\|\nabla v\|_3\|\Delta v\|_2.$$

In the whole space,

$$\|v\|_6\le C\|\nabla v\|_2,\qquad \|\nabla v\|_3\le C\|\nabla v\|_2^{1/2}\|\Delta v\|_2^{1/2}.$$

The same estimate holds on $\mathbb{R}^2 \times \mathbb{T}_L$ with a constant depending on L, after adding the harmless L^2 part already controlled by the energy inequality. Young's inequality yields, with

$$Y(t) := 1 + \|v(t)\|_{H^1}^2,$$

$$\boxed{\frac{dY}{dt} + \nu\|\Delta v\|_2^2 \le C_{\nu,*} Y^3.} \tag{3.1}$$

The scalar ODE gives a positive time on which Y stays bounded uniformly in the cutoff and in the initial datum inside the H^1 ball. It also gives a uniform $L_t^2 H_x^2$ bound. Weak compactness, local compactness and a diagonal argument produce a strong solution. The ordinary difference energy estimate in $L^2(dX)$ gives uniqueness. This proves the uniform restart statement. □

Lemma 3.2 (Circulation and weighted toroidal moments in both geometries)**.** *Assume* (H_∞) *in the whole space or* (H_L) *in the periodic cylinder. Then, with physical measure* dX*,*

$$\sup_{0<t<T} \|\Gamma(t)\|_{L^4(dX)}^4 + c_\nu \int_0^T \|\nabla(\Gamma^2)(t)\|_{L^2(dX)}^2 \, dt < \infty, \tag{3.2}$$

$$\sup_{0<t<T} \|r^3 q(t)\|_{L^2(dX)}^2 + c_\nu \int_0^T \|\widetilde{\nabla}(r^3 q)(t)\|_{L^2(dX)}^2 \, dt < \infty. \tag{3.3}$$

Here

$$\widetilde{\nabla}(r^3 q) := \big(r^3 \partial_r q,\, r^3 \partial_z q\big) = \big(\partial_r(r^3 q) - 3r^2 q,\, \partial_z(r^3 q)\big). \tag{3.4}$$

Thus $\widetilde{\nabla}$ *denotes the weighted derivative naturally produced by the radial integration by parts; it is not a new coordinate gradient. The constants may depend on* L *in the periodic geometry.*

Proof. Multiply the passive circulation equation (2.6) by $|\Gamma|^2\Gamma$ in physical measure. Advection vanishes by incompressibility, while the diffusion operator is coercive after radial integration by parts; cutoff exhaustion handles spatial infinity and periodicity removes the axial boundary term in Ω_L. This gives (3.2).

For q, multiply (2.5) by $r^6 q$ in physical measure. A direct radial integration by parts gives

$$\begin{aligned} \frac{d}{dt}\|r^3 q\|_{L^2(dX)}^2 + c_\nu \|\widetilde{\nabla}(r^3 q)\|_{L^2(dX)}^2 \\ \le C_{\nu,*}\Big(\|r^3 q\|_{L^2(dX)}^2 + \|q\|_{L^2(dX)}^2 \\ + \|\partial_z(\Gamma^2)\|_{L^2(dX)}^2\Big). \end{aligned} \tag{3.5}$$

For orientation, the source contribution has the form

$$\int r^4 q \, \partial_z(\Gamma^2) \, dr \, dz,$$

and Cauchy–Schwarz in physical measure bounds it by

$$C\|r^3 q\|_{L^2(dX)}\|\partial_z(\Gamma^2)\|_{L^2(dX)}.$$

The remaining weighted radial terms are absorbed into the first two terms on the right of (3.5) and the coercive diffusion term. This is the classical weighted azimuthal-vorticity computation; compare Chae–Lee [6, proof of Theorem 2]. The Leray inequality (2.11) controls $\int_0^T \|q\|^2_{L^2(dX)}dt$, while (3.2) controls the final term. Gronwall proves (3.3). □

Lemma 3.3 (Flat toroidal control implies a physical H^1 bound)**.** *Assume the hypotheses of theorem 3.2 and suppose*

$$\sup_{0<t<T}\|q(t)\|_{L^2(dr\,dz)} < \infty. \tag{3.6}$$

Then

$$\boxed{\sup_{0<t<T}\|u(t)\|_{H^1(dX)} < \infty.}$$

Proof. The toroidal physical vorticity is controlled by splitting at $r = 1$:

$$\begin{aligned}\frac{1}{2\pi}\|q(t)e_\theta\|^2_{L^2(dX)} &= \int rq^2\,dr\,dz \\ &\le \int_{0<r<1} q^2\,dr\,dz + \int_{r\ge 1} r^7 q^2\,dr\,dz.\end{aligned}$$

Thus (3.6) and (3.3) give

$$\sup_{0<t<T}\|q(t)e_\theta\|_{L^2(dX)} < \infty. \tag{3.7}$$

Let

$$\Omega_p := \omega^r e_r + \omega^z e_z.$$

It satisfies

$$\partial_t\Omega_p + b\cdot\nabla\Omega_p - \nu\Delta\Omega_p = \Omega_p\cdot\nabla b.$$

Taking the physical L^2 inner product with Ω_p and using the Hodge bound

$$\|\nabla b\|_{L^2(dX)} \le C_*\|qe_\theta\|_{L^2(dX)}$$

together with the three-dimensional Gagliardo–Nirenberg estimate gives

$$\frac{d}{dt}\|\Omega_p\|_2^2 + \nu\|\nabla\Omega_p\|_2^2 \le C_{\nu,*}\left(1 + \|qe_\theta\|_2^4\right)\|\Omega_p\|_2^2. \tag{3.8}$$

The constant term is unnecessary in $\mathbb{R}^3$ and harmless in the periodic cylinder. By (3.7) and Gronwall,

$$\sup_{0<t<T}\|\omega(t)\|_{L^2(dX)} < \infty.$$

For divergence-free fields in either geometry, $\|\nabla u\|_2$ is controlled by $\|\omega\|_2$ (modulo the harmless constant axial mode, which is already controlled by the energy norm). Together with (2.11), this proves the uniform physical H^1 bound. □

3.2. **SC.1: a whole-space radial-strain continuation criterion.** Put

$$U := u^\theta, \qquad \alpha := -\frac{u^r}{r}, \qquad q := \omega^\theta,$$

and define the whole-space flat meridian energy and dissipation

$$\mathcal{E}_f^\infty(t) := \|U(t)\|_{L^4(dr\,dz)}^4 + \|q(t)\|_{L^2(dr\,dz)}^2, \tag{3.9}$$

$$\mathcal{D}_f^\infty(t) := \|\nabla(U^2)\|_{L^2(dr\,dz)}^2 + \|\nabla q\|_{L^2(dr\,dz)}^2 + \left\|\frac{U^2}{r}\right\|_{L^2(dr\,dz)}^2 + \left\|\frac{q}{r}\right\|_{L^2(dr\,dz)}^2. \tag{3.10}$$

Lemma 3.4 (Whole-space flat energy with the radial strain coefficient)**.** *Every smooth rapidly decaying whole-space axisymmetric solution satisfies*

$$\boxed{\frac{d}{dt}\mathcal{E}_f^\infty(t) + c_\nu \mathcal{D}_f^\infty(t) \le C_\nu \|\alpha(t)\|_{L^2(dr\,dz)}^2 \mathcal{E}_f^\infty(t).} \tag{3.11}$$

Consequently,

$$\int_0^T \|\alpha(t)\|_{L^2(dr\,dz)}^2\, dt < \infty \tag{3.12}$$

implies

$$\sup_{0<t<T} \mathcal{E}_f^\infty(t) + c_\nu \int_0^T \mathcal{D}_f^\infty(t)\, dt < \infty. \tag{3.13}$$

Proof. The U and q equations are the whole-space versions of the equations used in the periodic flat-energy calculation. Multiply the U equation by $2|U|^2U$, multiply the q equation by q, and integrate in the flat meridian measure $dr\,dz$. The diffusion and axis terms contribute the four positive terms in (3.10); the remaining source/radial terms are the algebraic flat-meridian terms in the Chae–Lee calculation and are absorbed by Young's inequality; compare [6, proof of Theorem 3].

The only coefficients requiring a critical estimate have the form

$$I_U \lesssim \int_{\mathcal{Q}} |\alpha|\,|U|^4\, dr\, dz, \qquad I_q \lesssim \int_{\mathcal{Q}} |\alpha|\, q^2\, dr\, dz.$$

On the signed meridian plane the ordinary two-dimensional Gagliardo–Nirenberg inequality gives

$$\|f\|_{L^4(\mathbb{R}^2)}^2 \le C\|f\|_{L^2(\mathbb{R}^2)}\|\nabla f\|_{L^2(\mathbb{R}^2)}.$$

Thus, with the explicitly flat norms

$$A := \|\alpha\|_{L^2(dr\,dz)}, \quad F := \|U^2\|_{L^2(dr\,dz)}, \quad Y := \|\nabla(U^2)\|_{L^2(dr\,dz)},$$
$$Q := \|q\|_{L^2(dr\,dz)}, \quad X := \|\nabla q\|_{L^2(dr\,dz)},$$

one has

$$I_U \le CAFY \le \varepsilon Y^2 + C_\varepsilon A^2F^2, \qquad I_q \le CAQX \le \varepsilon X^2 + C_\varepsilon A^2Q^2.$$

Choose ε below the viscous coercivity constant. This gives (3.11); Gronwall gives (3.13). □

Proposition 3.5 (SC.1: whole-space radial-strain endpoint)**.** *Under* (H_∞)*,*

$$\int_0^T \int_{\mathcal{Q}} |\alpha(r,z,t)|^2 \, dr\, dz\, dt < \infty \tag{3.14}$$

implies smooth continuation through T*. Hence if* T *is a finite first singular time, then for every* $t_0 < T$*,*

$$\boxed{\int_{t_0}^T \int_{\mathcal{Q}} |\alpha|^2 \, dr\, dz\, dt = \infty.} \tag{3.15}$$

Proof. By theorem 3.4, (3.14) gives the uniform flat bound (3.13), in particular

$$\sup_{0<t<T} \|q(t)\|_{L^2(dr\,dz)} < \infty.$$

Apply theorems 3.2 and 3.3 to obtain a uniform physical $H^1(\mathbb{R}^3)$ bound up to T. Choose $t_n \uparrow T$. By theorem 3.1, the restart lifespan from $u(t_n)$ has a positive lower bound independent of n. Strong uniqueness identifies the restarted solution with the original one on the overlap, so for large n the restart extends past T, a contradiction.

If (3.15) failed for one $t_0 < T$, smoothness on $[0, t_0]$ would make the full integral in (3.14) finite, and the continuation conclusion would again contradict the choice of T. □

3.3. **Euler-compatible clock conversion.**

Lemma 3.6 (Whole-space weighted Hodge comparison)**.** *For every smooth rapidly decaying axisymmetric divergence-free meridional field* $b = (u^r, u^z)$ *with*

$$\operatorname{curl} b = -q e_\theta,$$

one has

$$\boxed{\|\alpha\|_{L^2(dr\,dz)} = \left\| \frac{u^r}{r} \right\|_{L^2(dr\,dz)} \le C \|q\|_{L^2(dr\,dz)}.} \tag{3.16}$$

Proof. Put

$$w(X) := r^{-1} = |x_h|^{-1}, \qquad x_h = (x_1, x_2).$$

We first verify directly that $w \in A_2(\mathbb{R}^3)$. Let $B = B(X_0, R)$ and write $d = \operatorname{dist}(X_0, \{x_h = 0\})$.

If $d \ge 2R$, then $r \asymp d$ on B, hence

$$\left(\frac{1}{|B|} \int_B w \, dX \right) \left(\frac{1}{|B|} \int_B w^{-1} \, dX \right) \le C.$$

If $d < 2R$, then B is contained in a cylinder of transverse radius $3R$ and axial length $2R$. Cylindrical integration gives

$$\int_B w \, dX \le C R^2, \qquad \int_B w^{-1} \, dX \le C R^4, \qquad |B| \asymp R^3.$$

Therefore

$$\left(\frac{1}{|B|}\int_B w\,dX\right)\left(\frac{1}{|B|}\int_B w^{-1}\,dX\right)\le C$$

also in this case. Taking the supremum over all balls proves $[w]_{A_2}<\infty$. Yaoming to ChatGPT; what is the meaning of $[w]_{A_2}$?

For every axisymmetric scalar f,

$$\int_{\mathcal{Q}}|f|^2\,dr\,dz=\frac{1}{2\pi}\int_{\mathbb{R}^3}|f(X)|^2w(X)\,dX. \tag{3.17}$$

Since b is divergence free and $\nabla\times b=-qe_\theta$,

$$-\Delta b=\nabla\times(-qe_\theta).$$

Each Cartesian component of ∇b is therefore a finite linear combination of second-order Euclidean Riesz transforms of the components of qe_θ. The weighted L^2 Calderón–Zygmund theorem for A_2 weights, for example Anderson–Vagharshakyan [1], gives

$$\|\nabla b\|_{L^2(w\,dX)}\le C[w]_{A_2}\|qe_\theta\|_{L^2(w\,dX)}.$$

The Cartesian gradient of an axisymmetric meridional field contains the positive angular-basis contribution $|u^r/r|^2$. Combining this with (3.17) gives (3.16). □

Corollary 3.7 (Derived C−WC: the whole-space future flat-q^2 clock)**.** *Under* (H_∞)*, if* T *is a finite first singular time, then for every* $t_0<T$*,*

$$\boxed{\int_{t_0}^{T}\int_{\mathcal{Q}}q(r,z,t)^2\,dr\,dz\,dt=\infty.} \tag{3.18}$$

Consequently every terminal tail contains arbitrarily late complete unit flat-q^2 *blocks, and their canonical radial first moments tend to zero. Thus the historical A.1/B.1/C.1 future-block clock is available internally.*

Proof. Integrate (3.16) in time. If the left side has infinite terminal-tail action by theorem 3.5, then the right side must also have infinite terminal-tail action, proving (3.18).

Partition any sufficiently late tail into consecutive blocks I_j satisfying

$$\int_{I_j}\int_{\mathcal{Q}}q^2\,dr\,dz\,dt=1.$$

Define

$$\rho_j:=\int_{I_j}\int_{\mathcal{Q}}rq^2\,dr\,dz\,dt.$$

The physical Leray dissipation gives

$$\sum_j\rho_j\le\int_0^T\int_{\mathcal{Q}}rq^2\,dr\,dz\,dt<\infty,$$

so $\rho_j \to 0$. Hence, after any prescribed preterminal time and below any prescribed comparison scale, one may select a later complete unit block. □

Remark 3.8 (Euler firewall and the role of the ν-polarizer). The Hodge inequality (3.16) is Euler-compatible. It is therefore used only to convert one terminal clock into another; it is not a terminal Navier–Stokes detector. The genuinely viscosity-specific step is separated twice: first in SC.1, whose continuation proof uses viscous flat dissipation and an H^1 restart, and finally in Bridge E, where Cauchy, full-adjoint, material-label and primal–adjoint polarizations leave explicit ν-weighted curvature/diffusion/current terms. Thus the recovery of the old C–WC clock does not weaken the Euler firewall.

4. Geometry-specific upstream bridges

The whole-space fork is now entirely center-free upstream: SC.1 supplies the future flat-q^2 clock, B.1 supplies the later strict degree-two scale, and the finite-corridor relay supplies the adjacent super-geometric pair needed by C.1. It does not use a fixed singular Type-II point.

The periodic fork uses one external local singularity input after the recurrent axis point has been selected internally. For a periodic singular axis point $Z_* = (X_*, T)$ define

$$\mathcal{E}_*(R) := \frac{1}{R}\int_{T-R^2}^{T}\int_{B_R(X_*)} |\nabla u(X,t)|^2\, dX\, dt.$$

Theorem 4.1 (Seregin's local no-Type-I theorem [25])**.** *Let (v,π) be an axisymmetric suitable weak solution of the viscosity-one Navier–Stokes equations in the backward cylinder used in Seregin's Theorem 2.1, and suppose that the terminal axis point z_0 is singular. Define the scale-invariant quantities*

$$E_v(z_0,r) := \frac{1}{r}\int_{Q(z_0,r)} |\nabla v|^2\, dY\, ds, \tag{4.1}$$

$$A_v(z_0,r) := \operatorname*{ess\,sup}_{t_0-r^2<s<t_0} \frac{1}{r}\int_{B_r(x_0)} |v(Y,s)|^2\, dY, \tag{4.2}$$

$$C_v(z_0,r) := \frac{1}{r^2}\int_{Q(z_0,r)} |v|^3\, dY\, ds. \tag{4.3}$$

Seregin's Type-I/Type-II number is

$$g(z_0) := \min\left\{ \limsup_{r\downarrow 0} E_v(z_0,r),\ \limsup_{r\downarrow 0} A_v(z_0,r),\ \limsup_{r\downarrow 0} C_v(z_0,r) \right\}. \tag{4.4}$$

Theorem 2.1 of [25] *states that a singular axisymmetric suitable point is Type II, i.e.*

$$g(z_0) = \infty.$$

Consequently

$$\boxed{\limsup_{r\downarrow 0} E_v(z_0,r) = \infty.} \tag{4.5}$$

Remark 4.2 (Why the last implication is literal). Because (4.4) is the *minimum* of the three limsups, $g(z_0) = \infty$ forces each of the three limsups to be infinite. Thus (4.5) is not an additional interpretation of Seregin's theorem; it is an immediate consequence of the source definition.

Remark 4.3 (Single-29 source-scope firewall for the Type-II input). The preceding theorem is a compact source-side summary. The active periodic proof uses it only with the following literal scope. Seregin's theorem is one-sided in time: suitability is required on a backward cylinder ending at the terminal singular point, not on any post-T continuation. Moreover the source local domain is cylindrical,

$$\mathcal{C}(r) = \{|x_h| < r,\ |x_3| < r\},$$

whereas the scale-invariant Type-I/II quantities are compared with ball-based quantities. The fixed inclusions

$$\boxed{B_r \subset \mathcal{C}(r) \subset B_{\sqrt{2}r}} \tag{4.6}$$

show that divergence of the scale-invariant dissipation is unchanged, up to a fixed change of radius. No post-terminal solution and no unrecorded shape identification are used below.

Proposition 4.4 (TI-T: one-sided periodic translation of Seregin's input)**.** *Let $Z_* = (X_*, T)$ be a singular axis point of the periodic physical solution. Assume that for every sufficiently small lifted backward Euclidean cylinder ending at Z_* the original preterminal pair (u, p) is suitable. For $\nu > 0$ define*

$$v(x,\tau) := \nu^{-1} u\Big(x, T + \frac{\tau}{\nu}\Big), \qquad \pi(x,\tau) := \nu^{-2} p\Big(x, T + \frac{\tau}{\nu}\Big). \tag{4.7}$$

Then (v, π) satisfies the viscosity-one system. If

$$E_v(R) := \frac{1}{R} \int_{-R^2}^{0} \int_{B_R(X_*)} |\nabla v|^2 \, dX \, d\tau$$

and

$$\mathcal{E}_*(r) := \frac{1}{r} \int_{T-r^2}^{T} \int_{B_r(X_*)} |\nabla u|^2 \, dX \, dt,$$

then

$$E_v(R) = \frac{1}{\nu R} \int_{T-R^2/\nu}^{T} \int_{B_R(X_*)} |\nabla u|^2 \, dX \, dt. \tag{4.8}$$

With

$$a_\nu := \max\{1, \nu^{-1/2}\},$$

one has

$$\boxed{\mathcal{E}_*(a_\nu R) \geq \frac{\nu}{a_\nu} E_v(R).} \tag{4.9}$$

Consequently theorem 4.1 implies

$$\boxed{\limsup_{r \downarrow 0} \mathcal{E}_*(r) = \infty.} \tag{4.10}$$

No solution for $t > T$ is used.

Proof. Equation (4.8) is the direct change of variables in (4.7). Since

$$B_R \subset B_{a_\nu R}, \qquad \frac{R^2}{\nu} \le (a_\nu R)^2,$$

the integration region on the right of (4.8) is contained in the region defining $\mathcal{E}_*(a_\nu R)$. This gives (4.9). Apply (4.5); the fixed shape comparison (4.6) allows the source cylinder convention to be replaced by balls without changing divergence. □

Theorem 4.5 (Euclidean Riesz transforms [28])**.** *For every $1 < p < \infty$, each Euclidean Riesz transform R_j is bounded on $L^p(\mathbb{R}^3)$, and hence every composition R_iR_j is bounded on $L^p(\mathbb{R}^3)$. We use this only with $p = 5/3$.*

Remark 4.6 (Pressure-source firewall). Theorem 4.5 is used only on an actual Euclidean lift and only in the ordinary physical $L^{5/3}(\mathbb{R}^3)$ norm. No mixed-cylinder Riesz multiplier theorem is imported. The periodic far-field and nonzero-image pressure are estimated internally by smooth-kernel bounds in theorem 4.7.

Lemma 4.7 (Periodic preterminal pressure integrability)**.** *Let u be a smooth finite-energy-per-period solution on $\Omega_L \times [0, T)$ satisfying*

$$u \in L^\infty(0, T; L^2(dX)) \cap L^2(0, T; H^1(dX)).$$

Then on every fixed terminal slab,

$$u \in L^{10/3}(dX\, dt), \qquad u \otimes u \in L^{5/3}(dX\, dt). \tag{4.11}$$

If $B_{4R}(X_*)$ lies below the axial injectivity radius, lift it to $\mathbb{R}^3$, choose $\chi \in C_c^\infty(B_{3R}(X_*))$ with $\chi \equiv 1$ on $B_{2R}(X_*)$, and set

$$p_{\text{near}} := \sum_{i,j=1}^{3} R_iR_j(\chi u_i u_j)\,. \tag{4.12}$$

Then

$$\|p_{\text{near}}\|_{L^{5/3}(\mathbb{R}^3)} \le C\|\chi u \otimes u\|_{L^{5/3}(\mathbb{R}^3)}. \tag{4.13}$$

The complementary periodic image/far-field pressure is smooth on $B_{2R}(X_*)$ and obeys

$$\|p_{\text{far}}(t)\|_{L^\infty(B_{2R}(X_*))} \le C_{R,L}\|u(t)\|^2_{L^2(\Omega_L)}. \tag{4.14}$$

Hence, after changing pressure by an irrelevant function of time if necessary,

$$p \in L^{5/3}\big(B_{2R}(X_*) \times (T - 4R^2, T)\big).$$

Proof. The Sobolev inequality and interpolation of $L^\infty_t L^2_x$ with $L^2_t L^6_x$ give (4.11). Apply theorem 4.5 to (4.12). Sources outside the lifted B_{2R}, including nonzero axial images of the Newtonian pressure kernel, are separated from the observation ball; their kernels and all spatial derivatives are therefore bounded there. The finite energy per period gives (4.14). □

Proposition 4.8 (Periodic one-sided preterminal suitability)**.** *Let $Z_* = (X_*, T)$ be a periodic axis point and suppose that T is a candidate first singular time. For every fixed $c > 0$ and every sufficiently small $R > 0$, after lifting below the axial injectivity radius there is a pressure representative for which (u, p) is an axisymmetric suitable weak solution on*

$$Q^-_{R,c}(Z_*) := B_R(X_*) \times (T - cR^2, T). \tag{4.15}$$

No solution for $t > T$ is required. In particular one may take $c = \nu^{-1}$, exactly the preterminal time interval appearing after (4.7).

Proof. The energy class supplies $u \in L^\infty_t L^2_x$ and $\nabla u \in L^2$. Fix $c > 0$. To apply theorem 4.7 on (4.15), choose

$$R_c := \frac{1}{2} \max\{1, \sqrt{c}\}\, R.$$

For R sufficiently small, $B_{4R_c}(X_*)$ still lies below the periodic injectivity radius; moreover

$$B_R \subset B_{2R_c}, \qquad (T - cR^2, T) \subset (T - 4R_c^2, T).$$

Hence theorem 4.7 gives $p \in L^{5/3}_{\mathrm{loc}}$ on the required cylinder, and therefore $p \in L^{3/2}_{\mathrm{loc}}$.

For each nonnegative terminal cutoff $\phi \in C^\infty_c$ in the lifted Euclidean chart, the solution is classical for $t < T$, so multiplication by $2u\phi$ and integration by parts gives the local energy equality on every compact time interval ending before T. The terms are integrable up to T by (4.11) and the pressure estimate; passage to the terminal limit gives the local energy inequality on $Q^-_{R,c}(Z_*)$. Rotation covariance of the pressure construction permits an axisymmetric representative. Thus the precise one-sided suitability hypothesis in theorem 4.4 is satisfied. □

Theorem 4.9 (Periodic local Type-II dissipation input)**.** *Let $Z_* = (X_*, T)$ be a singular axis point of the periodic physical solution, with the radius restricted below the axial injectivity scale, and let*

$$\mathcal{E}_*(r) := \frac{1}{r} \int_{T-r^2}^{T} \int_{B_r(X_*)} |\nabla u(X,t)|^2 \, dX \, dt. \tag{4.16}$$

Then

$$\boxed{\limsup_{r \downarrow 0} \mathcal{E}_*(r) = \infty.} \tag{4.17}$$

The quantity is evaluated on the Euclidean lift of the periodic cylinder.

Proof. By theorem 4.8 with $c = \nu^{-1}$, the original periodic solution satisfies the exact one-sided source-side suitability hypothesis. Apply theorem 4.4. Its conclusion (4.10) is exactly (4.17). □

4.1. **Whole-space endpoint clock and synchronization.** For $f = f(r, z)$ write

$$\|f\|_{\widetilde{L}^2}^2 := \int_{\mathcal{Q}} |f(r,z)|^2 \, dr \, dz.$$

By the internally proved whole-space strain endpoint and clock conversion, theorems 3.5 and 3.7, a hypothetical first singular time under (H_∞) satisfies

$$\int_0^T \int_{\mathcal{Q}} q^2 \, dr \, dz \, dt = \infty, \qquad \int_0^T \int_{\mathcal{Q}} r q^2 \, dr \, dz \, dt < \infty. \tag{4.18}$$

Partition a late terminal tail into unit flat-action blocks

$$I_j = [\tau_j, \tau_{j+1}], \qquad \int_{I_j} \int_{\mathcal{Q}} q^2 \, dr \, dz \, dt = 1, \tag{4.19}$$

and define

$$\rho_j := \int_{I_j} \int_{\mathcal{Q}} r q^2 \, dr \, dz \, dt. \tag{4.20}$$

Then $\sum_j \rho_j < \infty$ and $\rho_j \to 0$.

Proposition 4.10 (B.1: center-free whole-space target relay)**.** *Let $I_j = [\tau_j, \tau_{j+1}]$ be any sufficiently late canonical whole-space unit blocks with canonical radii $\rho_j \downarrow 0$, independently of whether the A.1 compactness/synchronization alternatives hold or fail. Then there are regular times $s_j \in (\tau_{j+1}, T)$, degree-two physical target radii a_j, and still later canonical blocks with radii $\widetilde{\rho}_j$ such that*

$$\boxed{a_j \le C\rho_j / j^2 = o(\rho_j), \qquad \widetilde{\rho}_j \le a_j.} \tag{4.21}$$

The construction is center-free and does not inherit the component, sign or profile of the parent block. In particular B.1 does not assume A.1 synchronization.

Proof. At a finite first singular time the global Lipschitz norm is unbounded on every terminal tail. Otherwise the H^1 energy estimate

$$\frac{1}{2}\frac{d}{dt}\|\nabla u\|_{L^2(dX)}^2 + \nu \|\Delta u\|_{L^2(dX)}^2 \le C \|\nabla u\|_{L^\infty(dX)} \|\nabla u\|_{L^2(dX)}^2$$

and the Leray inequality give a uniform physical H^1 bound. By theorem 3.1, restarting at $t_n \uparrow T$ has a lifespan bounded below independently of n, so the solution extends beyond T.

Hence choose $s_j > \tau_{j+1}$ with $\|\nabla u(s_j)\|_{L^\infty(dX)} \ge j^4 \rho_j^{-2}$ and define $a_j = |\nabla u|^{-1/2} \le C\rho_j / j^2$ at an approximate maximizing point. By theorem 3.7, every terminal tail contains complete unit flat-q^2 blocks with arbitrarily small canonical radius. Target-agnostic absorption after s_j selects a later block with $\widetilde{\rho}_j \le a_j$. □

Lemma 4.11 (Exact lifted-G action and radial tightness)**.** *For one unit block I_j define*

$$D_G(t) := \int_{\mathcal{Q}} r^3 \left(G_r^2 + G_z^2\right)\, dr\, dz, \qquad \Lambda_j^G := \rho_j \int_{I_j} D_G(t)\, dt.$$

Then

$$\boxed{\int_{I_j} D_G(t)\, dt = \int_{I_j}\int_{\mathcal{Q}} \left(r|q_r|^2 + r|q_z|^2 + \frac{q^2}{r}\right) dr\, dz\, dt.} \tag{4.22}$$

Consequently, on a subsequence with $\Lambda_j^G \le \Lambda_+$,

$$\int_{I_j}\int_{r<\varepsilon\rho_j} q^2\, dr\, dz\, dt \le \Lambda_+ \varepsilon, \tag{4.23}$$

whereas the first radial moment gives

$$\int_{I_j}\int_{r>K\rho_j} q^2\, dr\, dz\, dt \le K^{-1}. \tag{4.24}$$

Thus a fixed fraction of every bounded-efficiency unit block lies in a radial annulus $c_r\rho_j < r < C_r\rho_j$.

Proof. Since $q = rG$,

$$q_r = G + rG_r, \qquad q_z = rG_z.$$

After radial cutoff exhaustion,

$$2\int_0^\infty r^2 G G_r\, dr = \int_0^\infty r^2 \partial_r(G^2)\, dr = -2\int_0^\infty rG^2\, dr.$$

Expanding $r|q_r|^2$ and adding $q^2/r = rG^2$ therefore gives

$$\int_0^\infty \left(r|q_r|^2 + \frac{q^2}{r}\right) dr = \int_0^\infty r^3 G_r^2\, dr,$$

while

$$\int_0^\infty r|q_z|^2\, dr = \int_0^\infty r^3 G_z^2\, dr.$$

This proves (4.22). On $r < \varepsilon\rho_j$,

$$q^2 \le \varepsilon\rho_j \frac{q^2}{r},$$

so (4.22) gives (4.23). On $r > K\rho_j$,

$$q^2 \le (K\rho_j)^{-1} r q^2,$$

and (4.20) gives (4.24). □

The normalized-time exclusion uses three elementary scalar estimates. We record them here because they are the only analytic input specific to A.1.

Lemma 4.12 (Three scalar estimates for the A.1 diffuse-block exclusion)**.** *Let*

$$d\mu_5 := r^3\,dr\,dz, \qquad E_G := \int_{\mathcal{Q}} G^2\,d\mu_5, \qquad D_G := \int_{\mathcal{Q}} |\nabla_5 G|^2\,d\mu_5.$$

Then:

(i) with $\mathsf{U} := u^r/r$,

$$\boxed{\|b\cdot\nabla G\|^2_{\dot H^{-1}(d\mu_5)} \le C E_G^{3/2} D_G^{1/2};} \tag{4.25}$$

(ii) for $\Omega_p := \omega^r e_r + \omega^z e_z$,

$$\boxed{\frac{d}{dt}\|\Omega_p\|^2_{L^2(dX)} + \nu\|\nabla\Omega_p\|^2_{L^2(dX)} \le C_\nu \|qe_\theta\|^4_{L^2(dX)}\|\Omega_p\|^2_{L^2(dX)};} \tag{4.26}$$

(iii) writing $x = r^2$ *and*

$$E_A := \iint A^2\,dx\,dz, \quad E_G^x := \iint xG^2\,dx\,dz,$$

$$H_A := \iint A_z^2\,dx\,dz, \quad H_G := \iint xG_z^2\,dx\,dz, \quad H_W := \iint \frac{W^2}{x}\,dx\,dz,$$

one has

$$\boxed{\left|2\iint \frac{G\Gamma W}{\sqrt{x}}\,dx\,dz\right| \le C(E_A E_G^x)^{3/8}(H_A H_G)^{1/8} H_W^{1/2}.} \tag{4.27}$$

Proof. For (i), the physical toroidal vorticity is qe_θ, so

$$\|b\|_{L^6(dX)} \le C\|qe_\theta\|_{L^2(dX)} \asymp C E_G^{1/2},$$

and, by the three-dimensional Gagliardo–Nirenberg inequality together with (4.22),

$$\|q\|_{L^3(dX)} \le C E_G^{1/4} D_G^{1/4}.$$

Hence the transport part paired against $\phi \in \dot H^1(d\mu_5)$ is bounded by $C E_G^{3/4} D_G^{1/4}\|\nabla_5\phi\|_{L^2(d\mu_5)}$. The lifted divergence correction is $\mathrm{div}_{\mu_5} b = 2\mathsf{U}$ with $\mathsf{U} = -\partial_z(-\Delta_5)^{-1}G$; five-dimensional HLS and Sobolev give the same bound. Taking the dual supremum proves (4.25).

For (ii), the exact Cartesian equation

$$\partial_t\Omega_p + b\cdot\nabla\Omega_p - \nu\Delta\Omega_p = \Omega_p\cdot\nabla b$$

and incompressibility give

$$\frac12\frac{d}{dt}\|\Omega_p\|^2_{L^2(dX)} + \nu\|\nabla\Omega_p\|^2_{L^2(dX)} = \int_{\mathbb{R}^3} (\Omega_p\cdot\nabla b)\cdot\Omega_p\,dX.$$

Use

$$\|\nabla b\|_{L^2(dX)} \le C\|qe_\theta\|_{L^2(dX)}$$

and

$$\|\Omega_p\|^2_{L^4(dX)} \le C\|\Omega_p\|^{1/2}_{L^2(dX)}\|\nabla\Omega_p\|^{3/2}_{L^2(dX)},$$

then apply Young's inequality.

For (iii), smooth-axis compatibility gives

$$\Gamma(0,z)=0, \qquad \Gamma(x,z)=\frac{1}{2}\int_0^x A(s,z)\,ds.$$

Therefore, with

$$F(z):=\int_0^\infty A(x,z)^2\,dx, \qquad K(z):=\int_0^\infty xG(x,z)^2\,dx,$$

$$|\Gamma(x,z)|^2\le \frac{x}{4}F(z), \qquad \iint G^2\Gamma^2\,dx\,dz\le \frac{1}{4}\int F(z)K(z)\,dz.$$

One-dimensional Sobolev/Cauchy–Schwarz gives

$$\|F\|_\infty\le 2E_A^{1/2}H_A^{1/2}, \qquad \|K\|_\infty\le 2(E_G^x)^{1/2}H_G^{1/2},$$

and

$$\int\sqrt{FK}\,dz\le (E_AE_G^x)^{1/2}.$$

Thus

$$\int FK\,dz\le C(E_AE_G^x)^{3/4}(H_AH_G)^{1/4}.$$

A final Cauchy–Schwarz estimate in the explicitly displayed $dx\,dz$ measure yields (4.27). □

Proposition 4.13 (A.1: whole-space synchronization)**.** *Using the universal B.1 relay for every strict homogeneous pre-target and every rapid/spatial compactness failure, a hypothetical whole-space singularity satisfying* (4.18) *has a subsequence of unit blocks for which one of the following occurs:*

- *(i) derivative/forcing action loses normalized-time equiintegrability, or a fixed fraction of the block action escapes every fixed normalized radial or axial neighborhood; the already established B.1 relay absorbs that target/noncompactness into a later smaller canonical block;*
- *(ii) there are fixed constants $\eta_0,L_0,\delta_0,C_0>0$ and intervals $J_j\subset I_j$ such that*

$$|J_j|\le L_0\rho_j^2, \qquad \int_{J_j}\int_{\mathcal{Q}} q^2\,dr\,dz\,dt\ge\eta_0, \qquad |I_j|\ge\delta_0\rho_j^2, \tag{4.28}$$

and, with $\sigma_j=(T-\tau_{j+1})^{1/2}$,

$$\boxed{\sigma_j\le C_0\rho_j.} \tag{4.29}$$

On the compact branch there is $0<\theta<1$ such that

$$\boxed{\rho_{j+n}\le C\theta^n\rho_j, \qquad \sup_j\frac{\sum_{k>j}\rho_k}{\rho_j}<\infty.} \tag{4.30}$$

Proof. If Λ_j^G is unbounded, or if the derivative/source action fails normalized-time equiintegrability on a spatially tight normalized box, spacetime averaging selects an unbounded normalized derivative/source amplitude. The physical fields $\nabla(qe_\theta)$ and

$$S_\beta := (\beta^2)_z = 2\frac{\beta W}{r}$$

have homogeneous degrees three and five, respectively; hence their inverse homogeneous powers produce an $o(\rho_j)$ pre-target. If the concentration is not spatially tight, item (i) holds. Thus it remains to consider the bounded-efficiency, spatially tight, no-rapid branch.

By theorem 4.11, fixed choices of $0 < c_r < C_r < \infty$ keep a fixed fraction of the unit flat action in $c_r\rho_j < r < C_r\rho_j$. The no-spatial-exit assumption gives an axial center z_j and a fixed normalized axial interval containing the same compact fraction.

Normalize by

$$\widehat{r} = \frac{r}{\rho_j}, \qquad \widehat{z} = \frac{z - z_j}{\rho_j}, \qquad \tau = \frac{t - \tau_j}{\rho_j^2}, \qquad G_j = \rho_j^3 G.$$

Let

$$E_j(\tau) := \|G_j(\tau)\|^2_{L^2(d\mu_5)}, \qquad D_j(\tau) := \|\nabla_5 G_j(\tau)\|^2_{L^2(d\mu_5)}.$$

The definition of ρ_j gives

$$\int_{\widehat{I}_j} E_j(\tau)\, d\tau = 1. \tag{4.31}$$

Moreover, (4.22) and Cauchy–Schwarz in $dr\, dz\, dt$ give

$$1 = \left(\int_{I_j}\int_{\mathcal{Q}} q^2\, dr\, dz\, dt\right)^2 \le \rho_j \int_{I_j}\int_{\mathcal{Q}} \frac{q^2}{r}\, dr\, dz\, dt \le \rho_j \int_{I_j} D_G(t)\, dt,$$

so

$$1 \le \int_{\widehat{I}_j} D_j(\tau)\, d\tau \le \Lambda_+. \tag{4.32}$$

Suppose now that no fixed normalized time window carries a fixed positive fraction of the compact action. On the no-rapid branch the normalized energy has a uniform time modulus, hence

$$m_j := \sup_{\widehat{I}_j} E_j(\tau) \longrightarrow 0. \tag{4.33}$$

The normalized lifted equation is

$$\frac{1}{2}E_j' + \nu D_j = \langle T_j + S_{\Gamma,j}, G_j\rangle, \qquad T_j := -b_j \cdot \nabla G_j,$$

where $S_{\Gamma,j}$ is the normalized version of $2\Gamma W/r^3 = (\beta^2)_z$. By (4.25),

$$\int_{\widehat{I}_j} \|T_j\|^2_{\dot{H}^{-1}(d\mu_5)}\, d\tau \le C \int E_j^{3/2} D_j^{1/2}\, d\tau \le C m_j \Lambda_+^{1/2} \longrightarrow 0.$$

Integrating the energy identity and using (4.32) therefore forces

$$\int_{\widehat{I}_j} |\langle S_{\Gamma,j}, G_j\rangle|\, d\tau \geq c_\nu > 0. \tag{4.34}$$

The axis-Hardy estimate (4.27), Hölder in normalized time, and the poloidal inequality (4.26) now give

$$Y_j^- \geq c\, m_j^{-1/8} \longrightarrow \infty, \qquad Y_j := \|\Omega_{p,j}\|^2_{L^2(dX_{\mathrm{norm}})}. \tag{4.35}$$

Indeed $\int_{\widehat{I}_j} E_j^2\, d\tau \leq m_j$, so the poloidal Gronwall factor is $1 + o(1)$, while the remaining derivative factors in (4.27) are controlled by (4.32) and the poloidal dissipation. If the diverging normalized poloidal state is tight in a fixed normalized ball, averaging selects a point with $|\Omega_{p,j}| \to \infty$, hence an $o(\rho_j)$ degree-two pre-target. If it is not tight, item (i) holds. This contradiction proves the natural window in (4.28).

Choose fixed normalized radial and axial cutoffs so that at least $\eta_0/2$ of that window lies in one fixed normalized meridian box K. Put

$$Q_j := \rho_j^2 q.$$

If Q_j were unbounded on $K \times \widehat{J}_j$, its inverse square-root physical amplitude would give an $o(\rho_j)$ degree-two pre-target. Therefore

$$\|Q_j\|_{L^\infty(K\times\widehat{J}_j)} \leq M_2$$

on the compact no-pre-target branch. Since K has finite flat area,

$$\frac{\eta_0}{2} \leq \int_{\widehat{J}_j}\int_K Q_j^2\, d\widehat{r}\, d\widehat{z}\, d\tau \leq |K| M_2^2 |\widehat{J}_j|,$$

and hence

$$|I_j| \geq |J_j| \geq \delta_0 \rho_j^2, \qquad \delta_0 := \frac{\eta_0}{2|K|M_2^2} > 0.$$

Finally put

$$R_j^{\triangle} := \sqrt{T - \inf J_j}.$$

Radial/axial tightness gives a fixed $\eta_1 > 0$ with

$$\int_{J_j}\int_{\substack{c_r\rho_j<r<C_r\rho_j \\ |z-z_j|<C_z\rho_j}} q^2\, dr\, dz\, dt \geq \eta_1.$$

For the Cartesian toroidal vector,

$$|\nabla(qe_\theta)|^2 = q_r^2 + q_z^2 + \frac{q^2}{r^2}.$$

Since $dX = 2\pi r\, dr\, dz$, the angular-basis term alone gives

$$\int_{J_j}\int_{\mathbb{R}^3} |\nabla(qe_\theta)|^2\, dX\, dt \geq \frac{c\eta_1}{\rho_j}.$$

The localized spacetime set has physical volume $O(\rho_j^5)$, so some point satisfies

$$|\nabla(qe_\theta)| \geq c\rho_j^{-3}.$$

Its degree-three homogeneous length is therefore

$$a_j^{(3)} := |\nabla(qe_\theta)|^{-1/3} \le C\rho_j.$$

If $R_j^{\triangle}/\rho_j \to \infty$, then $a_j^{(3)}/R_j^{\triangle} \to 0$, a strict pre-target relative to the terminal-gap radius. B.1 relays it. Hence

$$R_j^{\triangle} \le C\rho_j,$$

and since $\tau_{j+1} \ge \inf J_j$,

$$\sigma_j = \sqrt{T - \tau_{j+1}} \le R_j^{\triangle} \le C_0\rho_j,$$

which proves (4.29).

The chronological identity

$$|I_j| = \sigma_{j-1}^2 - \sigma_j^2$$

now yields

$$\sigma_{j-1}^2 - \sigma_j^2 \ge \delta_0\rho_j^2 \ge \frac{\delta_0}{C_0^2}\sigma_j^2.$$

Thus

$$\sigma_j \le \theta_\sigma\sigma_{j-1}, \qquad \theta_\sigma := \left(1 + \frac{\delta_0}{C_0^2}\right)^{-1/2} < 1.$$

Applying the duration bound to the next block gives $\rho_{j+1} \le \delta_0^{-1/2}\sigma_j$. Iteration proves (4.30). □

Lemma 4.14 (Center-free adjacent super-geometric relay with a fixed corridor)**.** *Let $I_j = [\tau_j, \tau_{j+1}]$ be synchronized compact A.1 parent blocks and put $L_j := \rho_j$ and $\sigma_j = (T - \tau_{j+1})^{1/2}$. After routing the strict homogeneous/rapid and normalized spatial alternatives, there are adjacent canonical blocks with radii, relabeled as L_j and r_j^{ch}, such that*

$$\boxed{r_j^{\mathrm{ch}}/L_j \longrightarrow 0,} \tag{4.36}$$

and at most a fixed number N_* of unit flat-q^2 blocks intervene.

Proof. A.1 synchronization gives $T - \tau_{j+1} = \sigma_j^2 \le CL_j^2$. By theorem 4.10, a later unit block has radius $\widetilde{\rho}_j \le CL_j/j^2 = o(L_j)$. On the final no-degree-two-q-target branch, $|q| \le M_q L_j^{-2}$ in a fixed parent-scale moving box $K_{CL_j}(t)$. If an intervening unit fails to keep a fixed fraction $\eta_* > 0$ of its flat q^2 action in that box, this is the normalized spatial/axial-exit branch already routed by B.1. Hence, on the compact branch,

$$\int_{\tau_{j+1}}^{T}\int_{K_{CL_j}(t)} q^2\,dr\,dz\,dt \le CM_q^2,$$

because the box has flat area $O(L_j^2)$ and the remaining time is $O(L_j^2)$. Thus only $N_* = O(M_q^2/\eta_*)$ unit blocks can intervene.

Writing their radii as $L_j = \rho_j^{(0)}, \dots, \rho_j^{(m_j)} = \widetilde{\rho}_j$ with $m_j \le N_*$, the product of adjacent ratios equals $\widetilde{\rho}_j/L_j \to 0$. Therefore some adjacent ratio is at most $(\widetilde{\rho}_j/L_j)^{1/N_*} \to 0$. Relabel that adjacent pair. Since all blocks in the finite chain lie on the synchronized compact branch, the selected pair retains the A.1 natural-window and terminal-synchronization properties. □

4.2. **Whole-space compact source-work bridge.** Define

$$h := \frac{q^2}{r} = rG^2.$$

Its exact balance contains no meridional compression term:

$$\begin{aligned} \frac{d}{dt}\int_{\mathcal{Q}} rG^2\,dr\,dz + 2\nu\int_{\mathcal{Q}} r|\nabla G|^2\,dr\,dz \\ + 2\nu\int_{\mathbb{R}} G(0,z,t)^2\,dz = -2\int_{\mathcal{Q}} rG_z\beta^2\,dr\,dz. \end{aligned} \tag{4.37}$$

By theorem 4.14, the fully compact no-smaller-target branch has the adjacent super-geometric descent $\rho_{j+1}/\rho_j \to 0$.

Lemma 4.15 (Parent-low and child-high Hardy slices)**.** *Let I_j^{par} and I_j^{ch} be the adjacent canonical blocks selected by theorem 4.14, with radii*

$$L_j, \qquad r_j^{\mathrm{ch}}, \qquad \frac{r_j^{\mathrm{ch}}}{L_j} \longrightarrow 0.$$

Fix

$$R_j := Kr_j^{\mathrm{ch}}, \tag{4.38}$$

where

$$K > \max\left\{2, \frac{2}{\eta_0}\right\}$$

and η_0, L_0 are the A.1 natural-window constants. On the bounded lifted-G efficiency branch, after passage to a subsequence there are ordered times

$$t_j^- < t_j^+$$

in $I_j^{\mathrm{par}} \cup I_j^{\mathrm{ch}}$ such that, with

$$\mathfrak{H}_q(t) := \int_{\mathcal{Q}} \frac{q(r,z,t)^2}{r}\,dr\,dz = \int_{\mathcal{Q}} rG(r,z,t)^2\,dr\,dz, \tag{4.39}$$

one has

$$\boxed{R_j^3\mathfrak{H}_q(t_j^-) \longrightarrow 0, \qquad R_j^3\mathfrak{H}_q(t_j^+) \ge h_* > 0.} \tag{4.40}$$

Moreover

$$\boxed{\int_{I_j^{\mathrm{par}}\cup I_j^{\mathrm{ch}}} \mathfrak{H}_q(t)\,dt \le \frac{C_H}{R_j},} \tag{4.41}$$

with constants independent of j.

Proof. Write the parent block as $I_j^{\mathrm{par}} = [\tau_j, \tau_{j+1}]$. Since $R_j/L_j \to 0$ and A.1 gives $|I_j^{\mathrm{par}}| \geq \delta_0 L_j^2$, the final parent interval

$$J_j^- := [\tau_{j+1} - R_j^2, \tau_{j+1}]$$

lies inside I_j^{par} for all large j.

The Hardy term (4.39) is a positive summand in the exact lifted-G Dirichlet action (4.22). Hence bounded derivative efficiency gives

$$\int_{I_j^{\mathrm{par}}} \mathfrak{H}_q(t)\, dt \leq \frac{\Lambda_+}{L_j}.$$

Averaging on J_j^- selects $t_j^- \in J_j^-$ with

$$\mathfrak{H}_q(t_j^-) \leq \frac{\Lambda_+}{L_j R_j^2},$$

and therefore

$$R_j^3 \mathfrak{H}_q(t_j^-) \leq \Lambda_+ \frac{R_j}{L_j} \longrightarrow 0.$$

Let $J_j^{a,\mathrm{ch}} \subset I_j^{\mathrm{ch}}$ be the A.1 natural window of the child block. Then

$$|J_j^{a,\mathrm{ch}}| \leq L_0 (r_j^{\mathrm{ch}})^2 = \frac{L_0}{K^2} R_j^2,$$

and

$$\int_{J_j^{a,\mathrm{ch}}} \int_{\mathcal{Q}} q^2\, dr\, dz\, dt \geq \eta_0.$$

The canonical first radial moment of the whole child block is r_j^{ch}, so Markov gives

$$\int_{I_j^{\mathrm{ch}}} \int_{r>R_j} q^2\, dr\, dz\, dt \leq \frac{r_j^{\mathrm{ch}}}{R_j} = \frac{1}{K}.$$

Thus

$$\int_{J_j^{a,\mathrm{ch}}} \int_{r<R_j} q^2\, dr\, dz\, dt \geq \eta_0 - \frac{1}{K} \geq \frac{\eta_0}{2}.$$

Since $r < R_j$ implies $q^2/r \geq q^2/R_j$,

$$\int_{J_j^{a,\mathrm{ch}}} \mathfrak{H}_q(t)\, dt \geq \frac{\eta_0}{2R_j}.$$

Averaging over the natural window selects $t_j^+ \in J_j^{a,\mathrm{ch}}$ with

$$\mathfrak{H}_q(t_j^+) \geq \frac{\eta_0 K^2}{2L_0} R_j^{-3}.$$

Hence (4.40) holds with, for example,

$$h_* := \frac{\eta_0 K^2}{4L_0} > 0.$$

The ordering $t_j^- < \tau_{j+1} < t_j^+$ is automatic.

Finally bounded derivative efficiency on the two adjacent canonical blocks gives

$$\int_{I_j^{\mathrm{par}}\cup I_j^{\mathrm{ch}}} \mathfrak{H}_q(t)\,dt \le \frac{\Lambda_+}{L_j} + \frac{\Lambda_+}{r_j^{\mathrm{ch}}} = \frac{\Lambda_+}{L_j} + \frac{K\Lambda_+}{R_j} \le \frac{C_H}{R_j}$$

for all sufficiently large j, because $R_j/L_j \to 0$. □

Proposition 4.16 (The Hardy upcrossing occurs in natural viscous time)**.** *Under the hypotheses of theorem 4.15, there is an interval*

$$J_j^* := [s_j^-, s_j^+] \subset I_j^{\mathrm{par}} \cup I_j^{\mathrm{ch}}$$

and fixed constants $L_H, c_H > 0$ such that

$$\boxed{|J_j^*| \le L_H R_j^2,} \tag{4.42}$$

and

$$\boxed{\mathfrak{H}_q(s_j^+) - \mathfrak{H}_q(s_j^-) = c_H R_j^{-3}.} \tag{4.43}$$

Proof. Set

$$h_0 := \frac{h_*}{4}, \qquad h_1 := \frac{h_*}{2}.$$

For all sufficiently large j,

$$R_j^3 \mathfrak{H}_q(t_j^-) < h_0, \qquad R_j^3 \mathfrak{H}_q(t_j^+) > h_1.$$

Let s_j^+ be the first time after t_j^- at which

$$R_j^3 \mathfrak{H}_q(s_j^+) = h_1,$$

and let s_j^- be the last preceding time at which

$$R_j^3 \mathfrak{H}_q(s_j^-) = h_0.$$

By continuity,

$$\frac{h_0}{R_j^3} \le \mathfrak{H}_q(t) \le \frac{h_1}{R_j^3} \qquad (s_j^- \le t \le s_j^+).$$

Therefore (4.41) yields

$$\frac{h_0}{R_j^3}|J_j^*| \le \int_{J_j^*} \mathfrak{H}_q(t)\,dt \le \frac{C_H}{R_j},$$

and hence

$$|J_j^*| \le \frac{C_H}{h_0} R_j^2.$$

This is (4.42). The endpoint difference is exactly

$$\mathfrak{H}_q(s_j^+) - \mathfrak{H}_q(s_j^-) = \frac{h_1 - h_0}{R_j^3} = \frac{h_*}{4R_j^3},$$

so (4.43) holds with $c_H = h_*/4$. □

Theorem 4.17 (Natural-time derivative-source packet)**.** *Put*

$$S_\beta := (\beta^2)_z = 2\frac{\beta W}{r}.$$

On every clean adjacent super-geometric whole-space transition, the interval J_j^* *from theorem 4.16 satisfies*

$$R_j^3 \int_{J_j^*}\int_{\mathcal{Q}} rGS_\beta \, dr\, dz\, dt \geq c_{\text{work}} > 0, \tag{4.44}$$

$$R_j^5 \int_{J_j^*}\int_{\mathcal{Q}} r|S_\beta|^2 \, dr\, dz\, dt \geq c_S > 0, \tag{4.45}$$

$$R_j^3 \int_{J_j^*}\int_{\mathcal{Q}} r|\beta|^4 \, dr\, dz\, dt \geq c_\beta > 0. \tag{4.46}$$

All constants are generation independent.

Proof. Integrate the exact Hardy identity (4.37) over J_j^* and integrate by parts in z:

$$\begin{aligned}\mathfrak{H}_q(s_j^+) - \mathfrak{H}_q(s_j^-) + 2\nu \int_{J_j^*}\int_{\mathcal{Q}} r|\nabla G|^2 \, dr\, dz\, dt \\ + 2\nu \int_{J_j^*}\int_{\mathbb{R}} G(0,z,t)^2 \, dz\, dt \\ = 2\int_{J_j^*}\int_{\mathcal{Q}} rGS_\beta \, dr\, dz\, dt.\end{aligned}$$

Dropping the favorable dissipation and using (4.43) gives

$$R_j^3 \int_{J_j^*}\int_{\mathcal{Q}} rGS_\beta \, dr\, dz\, dt \geq \frac{c_H}{2},$$

which is (4.44).

Next, by Cauchy–Schwarz in the explicitly displayed measure,

$$\begin{aligned}\frac{c_H}{2R_j^3} &\leq \int_{J_j^*}\int_{\mathcal{Q}} rGS_\beta \, dr\, dz\, dt \\ &\leq \left(\int_{J_j^*} \mathfrak{H}_q(t)\, dt\right)^{1/2} \left(\int_{J_j^*}\int_{\mathcal{Q}} r|S_\beta|^2 \, dr\, dz\, dt\right)^{1/2} \\ &\leq \left(\frac{C_H}{R_j}\right)^{1/2} \left(\int_{J_j^*}\int_{\mathcal{Q}} r|S_\beta|^2 \, dr\, dz\, dt\right)^{1/2}.\end{aligned}$$

This proves (4.45).

Finally,

$$2r|G_z||\beta|^2 \leq \nu r|G_z|^2 + \nu^{-1} r|\beta|^4.$$

Using this in the original Hardy identity on J_j^* and dropping the remaining nonnegative diffusion terms gives

$$\mathfrak{H}_q(s_j^+) - \mathfrak{H}_q(s_j^-) \le \nu^{-1} \int_{J_j^*} \int_{\mathcal{Q}} r|\beta|^4 \, dr \, dz \, dt.$$

Apply (4.43) to obtain (4.46). □

Proposition 4.18 (Reflection-free compactification of the positive source work)**.** *Use the source window J_j^* and scale R_j from theorems 4.16 and 4.17, and write*

$$\widehat{r} = \frac{r}{R_j}, \qquad \widehat{z} = \frac{z - z_j}{R_j}, \qquad \widehat{t} = \frac{t - s_j^-}{R_j^2},$$
$$\widehat{q}_j = R_j^2 q, \qquad \widehat{S}_j = R_j^5 S_\beta, \qquad S_\beta = (\beta^2)_z.$$

The axial center z_j is free to depend on the selected source packet. After routing strict homogeneous targets, assume

$$\|\widehat{q}_j\|_{L^\infty} + \|\widehat{S}_j\|_{L^\infty} \le M_* \tag{4.47}$$

on the transition region. Then, after subselection, one of the following holds:

(i) the positive source work is not tight in any fixed normalized meridian box; this branch is discarded from C.1 and restarted by the center-free B.1 relay theorem 4.10;

(ii) there are fixed constants

$$0 < c_r < C_r < \infty, \qquad Z_* > 0, \qquad c_{\mathrm{box}} > 0$$

and axial centers z_j^* such that, in normalized variables centered at z_j^*,

$$\boxed{\int_{\widehat{J}_j^*} \int_{\substack{c_r < \widehat{r} < C_r \\ |\widehat{z}| < Z_*}} (\widehat{q}_j \widehat{S}_j)_+ \, d\widehat{r} \, d\widehat{z} \, d\widehat{t} \ge c_{\mathrm{box}}.} \tag{4.48}$$

No reflection plane in z is used.

Proof. By (4.44),

$$\int_{\widehat{J}_j^*} \int \widehat{q}_j \widehat{S}_j \, d\widehat{r} \, d\widehat{z} \, d\widehat{t} \ge c_{\mathrm{work}},$$

and therefore

$$\int_{\widehat{J}_j^*} \int (\widehat{q}_j \widehat{S}_j)_+ \, d\widehat{r} \, d\widehat{z} \, d\widehat{t} \ge c_{\mathrm{work}}. \tag{4.49}$$

The normalized time length is bounded by L_H.

If no fixed normalized axial interval, after arbitrary recentering, captures a fixed positive fraction of (4.49), or if every fixed normalized radial exterior misses that fraction, we declare a source-work spatial exit. Nothing in the downstream proof requires profile inheritance across such an exit: theorem 4.10 applies to the current late canonical parent independently of which compactness property failed and selects a later strict degree-two target and a later canonical block. Thus this is item (i).

On the complementary branch, choose Z_*, C_r and centers z_j^* so that a fixed fraction of the positive work lies in

$$0 < \widehat{r} < C_r, \qquad |\widehat{z}| < Z_*.$$

The envelope (4.47) gives

$$(\widehat{q}_j \widehat{S}_j)_+ \le M_*^2.$$

Hence the contribution of $0 < \widehat{r} < \varepsilon$ inside this fixed axial box is at most

$$2 Z_* L_H M_*^2 \, \varepsilon.$$

Choose $c_r > 0$ so small that this is less than half of the retained positive work. Shrinking the retained constant if necessary gives (4.48). □

Lemma 4.19 (Compact positive work gives a fixed signed full-q child)**.** *Assume the compact branch* (4.48) *and the envelope* (4.47)*. After routing a component which leaves every fixed enlargement of the source box as a spatial/recentering restart, there are*

$$s \in \{\pm 1\}, \qquad t_j^+ \in J_j^*,$$

and a compact union $\Omega_j(t_j^+)$ of positive components of sq such that

$$\boxed{\int_{\Omega_j(t_j^+)} sq(r, z, t_j^+)\, dr\, dz \ge \mu_* > 0,} \tag{4.50}$$

where μ_* is independent of j.

Proof. Let

$$\widehat{K}_* := \left\{ (\widehat{r}, \widehat{z}, \widehat{t}) : c_r < \widehat{r} < C_r,\ |\widehat{z}| < Z_*,\ \widehat{t} \in \widehat{J}_j^* \right\}.$$

Its normalized spacetime volume is bounded by a fixed number V_*. Set

$$a_* := \frac{c_{\mathrm{box}}}{8 M_* V_*}, \qquad b_* := \frac{c_{\mathrm{box}}}{8 M_* V_*}.$$

The contribution to the positive work from $|\widehat{q}_j| < a_*$ is at most $a_* M_* V_* \le c_{\mathrm{box}}/8$, and the contribution from $|\widehat{S}_j| < b_*$ is bounded in the same way. Hence the set

$$E_j := \left\{ \widehat{K}_* : \widehat{q}_j \widehat{S}_j > 0,\ |\widehat{q}_j| \ge a_*,\ |\widehat{S}_j| \ge b_* \right\}$$

satisfies

$$\int_{E_j} \widehat{q}_j \widehat{S}_j \, d\widehat{r}\, d\widehat{z}\, d\widehat{t} \ge \frac{c_{\mathrm{box}}}{2}.$$

Since $|\widehat{q}_j \widehat{S}_j| \le M_*^2$,

$$|E_j| \ge \frac{c_{\mathrm{box}}}{2 M_*^2} =: m_* > 0. \tag{4.51}$$

Split E_j according to the sign of $\widehat{q}_j$. After subselection, one fixed sign $s \in \{\pm 1\}$ occupies at least $m_*/2$ normalized spacetime measure. Since $|\widehat{J}_j^*| \le L_H$, Fubini supplies a time $\widehat{t}_j^+$ such that the corresponding spatial

slice $E_j^s(\widehat{t}_j^+)$ has normalized area at least $m_*/(2L_H)$. On this slice, $s\widehat{q}_j \ge a_*$, so

$$\int_{E_j^s(\widehat{t}_j^+)} s\widehat{q}_j \, d\widehat{r}\, d\widehat{z} \ge \frac{a_* m_*}{2L_H} =: \mu_* > 0.$$

Because

$$\widehat{q}_j = R_j^2 q, \qquad d\widehat{r}\, d\widehat{z} = R_j^{-2}\, dr\, dz,$$

the signed flat mass is invariant under this normalization, and therefore the same lower bound holds in physical meridian coordinates.

Let $\Omega_j(t_j^+)$ be the union of the positive components of sq that meet this selected spatial set. If those components leave every fixed normalized enlargement of the source box, this is the spatial/recentering restart branch and we invoke theorem 4.10. On the complementary compact branch the union stays in one fixed enlarged R_j-box, and positivity gives

$$\int_{\Omega_j(t_j^+)} sq\, dr\, dz \ge \int_{E_j^s(\widehat{t}_j^+)} sq\, dr\, dz \ge \mu_*.$$

This proves (4.50). □

Remark 4.20 (Arbitrary axial centers do not create a new whole-space gate). The box center z_j^* in theorems 4.18 and 4.19 is arbitrary. No z-reflection is used. Moreover the fixed signed child itself contains a point with

$$|q| \ge c\mu_* R_j^{-2},$$

because its flat area is $O(R_j^2)$. Thus its degree-two homogeneous length is $O(R_j) = o(L_j)$. If the source center wanders from generation to generation, the center-free future clock can restart from this strict target without inheriting the center, sign, or profile.

Theorem 4.21 (C.1: whole-space compact source-work interface)**.** *Under* (H_∞)*, A.1 and B.1, a hypothetical whole-space first singular time has, after routing every strict smaller target, parent/child scales*

$$L_j, \qquad R_j := K r_j^{\mathrm{ch}}, \qquad \frac{R_j}{L_j} \longrightarrow 0,$$

and source windows J_j^* such that:

(i)

$$|J_j^*| \le C R_j^2,$$

and the source core stays in one fixed normalized meridian box with $r \asymp R_j$ and $|z - z_j^*| \lesssim R_j$;

(ii) the scale-normalized source work obeys

$$\boxed{R_j^3 \int_{J_j^*} \int_{\mathcal{Q}} rG(\beta^2)_z \, dr\, dz\, dt \ge c_* > 0;} \tag{4.52}$$

(iii) *every normalized degree-two q amplitude and degree-five S_β amplitude used to compactify the source packet is uniformly bounded; failure of such a bound is itself one of the already routed strict homogeneous targets;*

(iv) *at a child time $t_j^+ \in J_j^*$, after one sign choice $s \in \{\pm 1\}$, a compact component union satisfies*

$$\boxed{\int_{\Omega_j(t_j^+)} sq\, dr\, dz \geq \mu_* > 0;} \tag{4.53}$$

(v) *the parent–child transition occupies only a fixed finite number of canonical unit flat-action blocks and therefore has a generation-independent local flat-q^2 action budget.*

Proof. The adjacent parent/child pair is supplied by theorem 4.14. The natural-time Hardy packet is theorems 4.15 to 4.17. Hence items (i)–(ii) and the companion source taxes are available.

If a normalized q or S_β amplitude becomes unbounded, its inverse homogeneous power is an already routed strict target. On the complementary branch apply theorem 4.18. Its spatial-exit alternative is restarted by B.1; its compact alternative yields one fixed normalized source box at an arbitrary axial center. Then theorem 4.19 gives item (iv), with no reflection assumption.

Finally theorem 4.14 bounds the number of intervening canonical unit blocks by a fixed N_*. Since each block carries one unit of flat q^2 action, the complete parent–child corridor has a generation-independent local flat-action budget. This is item (v). □

Proposition 4.22 (Axial-translation covariance of the whole-space downstream)**.** *Let $z_0 \in \mathbb{R}$ and define the axial translation*

$$(\mathcal{T}_{z_0} f)(r, z, t) := f(r, z + z_0, t).$$

For the whole-space geometry, every active object from C.1 through Bridge E is covariant under $\mathcal{T}_{z_0}$:

(i) *the flat and physical measures, canonical block action, canonical radius, Hardy mass, source work, signed flat-q mass and local flat-action budget are invariant;*

(ii) *the Euclidean Hodge operator, the material map, the full vorticity adjoint, material-label pullback, Wronskian current and physical Leray payment commute with the corresponding translated construction;*

(iii) *B.1 and chronological clock absorption do not inherit an absolute axial center, sign, component or profile.*

Consequently the arbitrary center z_j^* in C.1 may be recentered to zero for a local calculation and then restored, without adding a hypothesis to Bridge D or Bridge E.

Proof. The whole-space equations have coefficients independent of z. Therefore $\mathcal{T}_{z_0}$ commutes with ∂_z, the meridional differential operators, Euclidean curl/divergence, the Laplacian and the whole-space Biot–Savart/Hodge operators. Translation preserves $dr\,dz$ and dX. The canonical radius uses only

$$\int rq^2\,dr\,dz\,dt,$$

and hence is unchanged. The Hardy density q^2/r, the source work $rG(\beta^2)_z$, and signed flat mass transform by a change of variable in z.

If $X(a,t)$ is a material map for u, then $X(a,t) - z_0 e_z$ is the translated map for the translated velocity after the corresponding translation of labels. Thus $F = D_a X$, the Cauchy-polarized vorticity, the full adjoint equation, material cutoffs and the primal–adjoint Wronskian identity are unchanged after relabeling. The physical Leray integral is translation invariant as well.

Finally theorem 4.10 uses only a late canonical block, terminal Lipschitz blow-up and the future clock; its statement explicitly discards inheritance of center, sign and profile. Therefore no downstream step requires an absolute axial anchor. □

Remark 4.23 (Role of the historical reflection specialization). In earlier reflection-symmetric formulations, z-reflection supplied a convenient fixed plane, paired lobes and parity-based notation. It did not alter the full-space Hardy identity or the scale of the natural-time source packet. The current proof replaces the only geometrically substantive use needed by C.1—the choice of a compact source lobe—by theorems 4.18 and 4.19 and the translation covariance above.

4.3. **Active periodic strain clock and recurrent local selector.** The active minimal periodic proof uses the radial strain

$$\alpha = -\frac{u^r}{r}$$

as its terminal clock.

Proposition 4.24 (SC.2: periodic radial-strain endpoint)**.** *Under* (H_L)*,*

$$\int_0^T \int_{\mathcal{Q}_L} |\alpha|^2\,dr\,dz\,dt < \infty \tag{4.54}$$

implies smooth continuation through T*. Consequently, if* T *is a first singular time, then for every* $t_0 < T$*,*

$$\boxed{\int_{t_0}^T \int_{\mathcal{Q}_L} |\alpha|^2\,dr\,dz\,dt = \infty.} \tag{4.55}$$

Proof. Keep the periodic flat energy

$$\mathcal{E}_f(t) := \|u^\theta(t)\|^4_{L^4(dr\,dz)} + \|q(t)\|^2_{L^2(dr\,dz)}$$

and its positive flat dissipation $\mathcal{D}_f$. The only critical coefficients are

$$I_\theta \lesssim \int_{\mathcal{Q}_L} |\alpha|\,|u^\theta|^4\,dr\,dz, \qquad I_q \lesssim \int_{\mathcal{Q}_L} |\alpha|\,q^2\,dr\,dz.$$

The two-dimensional cylinder interpolation

$$\|f\|^2_{L^4(dr\,dz)} \le C_L \|f\|_{L^2(dr\,dz)} \left(\|\nabla f\|_{L^2(dr\,dz)} + \|f\|_{L^2(dr\,dz)}\right)$$

applied to $f = (u^\theta)^2$ and $f = q$, followed by Cauchy–Schwarz and Young, gives

$$\boxed{\mathcal{E}_f'(t) + c_\nu \mathcal{D}_f(t) \le C_{\nu,L} \left(1 + \|\alpha(t)\|^2_{L^2(dr\,dz)}\right) \mathcal{E}_f(t).} \tag{4.56}$$

Under (4.54), Gronwall yields

$$\sup_{0<t<T} \|q(t)\|_{L^2(dr\,dz)} < \infty.$$

Now apply theorems 3.2 and 3.3 in Ω_L to obtain a uniform physical H^1 bound. Restarting at $t_n \uparrow T$ with theorem 3.1 extends the solution past T. The contrapositive gives (4.55). □

Lemma 4.25 (Finite radial first moment of the strain clock)**.** *Every finite-energy smooth solution satisfies*

$$\boxed{\int_0^T \int_{\mathcal{Q}_L} r|\alpha|^2\,dr\,dz\,dt < \infty.} \tag{4.57}$$

Proof. The cylindrical Cartesian gradient contains the positive term $|u^r/r|^2 = |\alpha|^2$. Hence

$$2\pi \int_{\mathcal{Q}_L} r|\alpha|^2\,dr\,dz \le \int_{\Omega_L} |\nabla u|^2\,dX.$$

Integrate in time and use the Leray energy identity. □

Lemma 4.26 (Recurrent singular axis point for the strain clock)**.** *If T is a periodic first singular time, there exists $z_* \in \mathbb{T}_L$ such that for every $0 < \delta < L/8$ and every $t_0 < T$,*

$$\boxed{\int_{t_0}^T \int_{\substack{0<r<\delta \\ d_L(z,z_*)<\delta}} |\alpha|^2\,dr\,dz\,dt = \infty.} \tag{4.58}$$

In particular $(0, z_*, T)$ is a singular axis point.

Proof. Fix $r_* > 0$. By theorems 4.24 and 4.25, for every $t_0 < T$,

$$\int_{t_0}^T \int_{\mathbb{T}_L} \int_0^{r_*} |\alpha|^2\,dr\,dz\,dt = \infty,$$

because the contribution of $r \ge r_*$ is at most $r_*^{-1} \int r|\alpha|^2\,dr\,dz\,dt < \infty$.

Choose $t_m \uparrow T$ and for a closed axial arc $I \subset \mathbb{T}_L$ set

$$\nu_m(I) := \int_{t_m}^T \int_I \int_0^{r_*} |\alpha|^2\,dr\,dz\,dt.$$

For every m, $\nu_m(\mathbb{T}_L) = \infty$, and for each fixed arc I the sequence $m \mapsto \nu_m(I)$ is nonincreasing.

We use the following elementary simultaneous-tail observation. Suppose a closed parent arc I satisfies

$$\nu_m(I) = \infty \qquad \text{for every } m.$$

Subdivide I into finitely many closed child arcs with disjoint interiors. Then at least one child I' satisfies

$$\nu_m(I') = \infty \qquad \text{for every } m.$$

Indeed, if every child became finite at some index, choose for each child an index m_ℓ with $\nu_{m_\ell}(I_\ell) < \infty$ and put $m_* = \max_\ell m_\ell$. Monotonicity in m would make every $\nu_{m_*}(I_\ell)$ finite, hence their finite sum would give $\nu_{m_*}(I) < \infty$, a contradiction.

Starting with $\mathbb{T}_L$, repeatedly bisect and choose such a child. This produces nested closed arcs I_k with

$$\operatorname{diam} I_k \longrightarrow 0, \qquad \nu_m(I_k) = \infty \quad \text{for all } m, k.$$

Compactness gives a unique intersection point

$$\{z_*\} = \bigcap_{k \geq 1} I_k.$$

Every axial neighborhood of z_* contains some I_k and therefore has infinite tail mass. Replacing r_* by any smaller δ removes only

$$\int_{t_m}^{T} \int_{\mathbb{T}_L} \int_{\delta \leq r < r_*} |\alpha|^2 \, dr \, dz \, dt \leq \delta^{-1} \int_{t_m}^{T} \int_{\mathbb{T}_L} \int_{\delta \leq r < r_*} r|\alpha|^2 \, dr \, dz \, dt < \infty,$$

which proves (4.58). If $(0, z_*, T)$ were regular, α would be bounded on one terminal parabolic neighborhood and the left-hand side would be finite for late t_0, a contradiction. □

For $0 < S < L/8$ define

$$\mathcal{C}_S(z_*) := \{(r, z) : 0 < r < S, \ d_L(z, z_*) < S\}. \tag{4.59}$$

By (4.58), there is $b_S < T$ such that, with

$$P(S) := [T - S^2, b_S], \tag{4.60}$$

one has

$$\boxed{\int_{P(S)} \int_{\mathcal{C}_S(z_*)} |\alpha|^2 \, dr \, dz \, dt = 1.} \tag{4.61}$$

Define its first radial moment

$$\eta(S) := \int_{P(S)} \int_{\mathcal{C}_S(z_*)} r|\alpha|^2 \, dr \, dz \, dt, \qquad 0 < \eta(S) \leq S. \tag{4.62}$$

Proposition 4.27 (Local strain selector with geometry-correct detector scale)**.** *Let $S_j \downarrow 0$, set*

$$P_j := P(S_j), \qquad \eta_j := \eta(S_j).$$

After passage to a subsequence, at least one of the following occurs.

(i) Collapsing first moment. *One has $\eta_j/S_j \to 0$. Then there are points $(r_j, z_j, t_j) \in \mathcal{C}_{S_j}(z_*) \times P_j$ and scales*

$$R_j^\sharp := (\eta_j S_j^3)^{1/4} = S_j \left(\frac{\eta_j}{S_j}\right)^{1/4} = o(S_j) \tag{4.63}$$

such that

$$\boxed{(R_j^\sharp)^2 |\alpha(r_j, z_j, t_j)| \geq c_* > 0, \qquad \frac{r_j}{R_j^\sharp} \longrightarrow 0.} \tag{4.64}$$

The intrinsic degree-two strain length $a_j^{(\alpha)} := |\alpha(r_j, z_j, t_j)|^{-1/2}$ satisfies $a_j^{(\alpha)} \leq CR_j^\sharp = o(S_j)$.

(ii) Compact natural-time packet. *One has $\eta_j \asymp S_j$ and*

$$\boxed{\begin{aligned} &\int_{P_j} \int_{\mathcal{C}_{S_j}(z_*)} |\alpha|^2 \, dr \, dz \, dt = 1, \\ &|P_j| \asymp \eta_j^2, \qquad \sqrt{T - \inf P_j} \leq C\eta_j, \\ &\mathcal{C}_{S_j}(z_*) \subset \{r < C\eta_j, \ d_L(z, z_*) < C\eta_j\}. \end{aligned}} \tag{4.65}$$

Moreover, unless a still smaller degree-two strain target occurs, there is a point (r_j, z_j, t_j) in the packet with

$$\boxed{\eta_j^2 |\alpha(r_j, z_j, t_j)| \geq c_* > 0, \qquad r_j \leq C\eta_j.} \tag{4.66}$$

Proof. If $\eta_j/S_j \to 0$, Markov's inequality applied to the probability measure $|\alpha|^2 \, dr \, dz \, dt$ in (4.61) gives

$$\int_{P_j} \int_{\substack{r < 2\eta_j \\ d_L(z,z_*) < S_j}} |\alpha|^2 \, dr \, dz \, dt \geq \frac{1}{2}.$$

The flat spacetime measure of this set is at most $C\eta_j S_j^3$. Thus some point with $r_j < 2\eta_j$ satisfies

$$|\alpha(r_j, z_j, t_j)|^2 \geq \frac{c}{\eta_j S_j^3}.$$

With (4.63),

$$(R_j^\sharp)^2 |\alpha(r_j, z_j, t_j)| \geq c,$$

and

$$\frac{r_j}{R_j^\sharp} \leq C \frac{\eta_j}{(\eta_j S_j^3)^{1/4}} = C \left(\frac{\eta_j}{S_j}\right)^{3/4} \longrightarrow 0.$$

This proves item (i), including the bound on $a_j^{(\alpha)}$.

On the complementary subsequence $\eta_j \geq cS_j$, so $S_j \leq C\eta_j$ and the spatial/terminal-gap inclusions in (4.65) are immediate. If $\eta_j^2|\alpha|$ is unbounded, a still smaller degree-two target has already been selected. Otherwise $|\alpha| \leq M\eta_j^{-2}$ in the packet. Its spatial flat area is $O(\eta_j^2)$, hence the unit action gives

$$1 \leq C\eta_j^2 M^2 \eta_j^{-4}|P_j|,$$

so $|P_j| \geq c\eta_j^2$; the reverse bound follows from $|P_j| \leq S_j^2 \leq C\eta_j^2$. The spacetime flat measure is $O(\eta_j^4)$, and the unit action therefore gives (4.66). □

4.4. **Periodic active mode and the same-scale Hodge detector.** For the periodic branch define the axial mean and active part by

$$\boxed{\overline{q}(r,t) := \frac{1}{L}\int_{\mathbb{T}_L} q(r,z,t)\,dz, \qquad q^\perp := q - \overline{q}.} \tag{4.67}$$

Likewise write

$$G = \overline{G} + G^\perp, \qquad q^\perp = rG^\perp.$$

The decomposition is orthogonal in every z-independent radial weight:

$$\int_{\mathbb{T}_L} q^2\,dz = L\overline{q}^{\,2} + \int_{\mathbb{T}_L} |q^\perp|^2\,dz. \tag{4.68}$$

The axial mean produces no radial velocity. Indeed, averaging

$$\partial_r u^r + \frac{u^r}{r} + \partial_z u^z = 0$$

over $\mathbb{T}_L$ gives

$$\partial_r(r\overline{u^r}) = 0,$$

and smoothness at the axis yields $\overline{u^r} = 0$. Hence the radial strain is generated by the active part:

$$\alpha = -\frac{u^r}{r} = \partial_z(-\mathcal{L}_1)^{-1}G^\perp. \tag{4.69}$$

Lemma 4.28 (Energy-bounded smooth remainder in a periodic Hodge chart)**.** *Fix the axial period L. Let $z_0 \in \mathbb{T}_L$ and choose an axisymmetric cutoff χ_{z_0} supported in one Euclidean periodic chart of physical radius $L/8$ and equal to one on the concentric chart of radius $L/16$. Let*

$$b^\perp := b - \overline{b},$$

where the bar denotes axial averaging. Then

$$\operatorname{curl} b^\perp = -q^\perp e_\theta, \qquad \operatorname{div} b^\perp = 0, \qquad \|b^\perp(t)\|_{L^2(dX)} \leq \|b(t)\|_{L^2(dX)}.$$

Lift the cutoff chart to $\mathbb{R}^3$ and define

$$b_{\mathrm{loc}} := \nabla \times (-\Delta_{\mathbb{R}^3})^{-1}\big(-\chi_{z_0} q^\perp e_\theta\big), \qquad h := b^\perp - b_{\mathrm{loc}}.$$

On the chart of radius $L/16$,

$$\operatorname{curl} h = 0, \qquad \operatorname{div} h = 0, \qquad \Delta h = 0.$$

Moreover, with $E_0 := \|u_0\|^2_{L^2(dX)}$,

$$\boxed{\|\nabla h(t)\|_{L^\infty(B_{L/32})} + \left\| \frac{h^r(t)}{r} \right\|_{L^\infty(B_{L/32})} \le C_L E_0^{1/2}.} \tag{4.70}$$

Consequently, for every $R_j \downarrow 0$ *and every target point in* $B_{L/32}$,

$$\boxed{R_j^2 |\alpha_{\mathrm{rem}}(Y_j, t_j)| \le C_L E_0^{1/2} R_j^2 \longrightarrow 0,} \tag{4.71}$$

where $\alpha_{\mathrm{rem}} := -h^r/r$.

Proof. Axial Fourier orthogonality gives $\|b^\perp\|_{L^2(dX)} \le \|b\|_{L^2(dX)}$, and the physical energy inequality bounds the latter by $E_0^{1/2}$.

Because χ_{z_0} is supported strictly inside one periodic chart, extend $\chi_{z_0} b^\perp$ by zero to $\mathbb{R}^3$. In the distributional sense,

$$-\chi_{z_0} q^\perp e_\theta = \nabla \times (\chi_{z_0} b^\perp) - \nabla \chi_{z_0} \times b^\perp.$$

The first term has $\dot{H}^{-1}(\mathbb{R}^3)$ norm bounded by $C\|b^\perp\|_2$. The second is supported in a fixed L-dependent annulus; by Sobolev duality on that bounded set,

$$\|\nabla \chi_{z_0} \times b^\perp\|_{\dot{H}^{-1}(\mathbb{R}^3)} \le C_L \|b^\perp\|_{L^2(dX)}.$$

The Euclidean order-minus-one Biot–Savart operator therefore gives

$$\|b_{\mathrm{loc}}\|_{L^2(\mathbb{R}^3)} \le C_L E_0^{1/2}.$$

Hence h has a uniform L^2 bound on the chart of radius $L/16$.

On the smaller chart the cutoff equals one, so $\operatorname{curl} b_{\mathrm{loc}} = -q^\perp e_\theta = \operatorname{curl} b^\perp$ and both fields are divergence free. Thus h is harmonic there. Standard interior harmonic estimates give

$$\|\nabla h\|_{L^\infty(B_{L/32})} \le C_L \|h\|_{L^2(B_{L/16})} \le C_L E_0^{1/2}.$$

Both $b^\perp$ and b_{loc} are smooth axisymmetric meridional fields, so $h^r(0, z, t) = 0$. The radial mean-value theorem yields

$$\left| \frac{h^r(r, z, t)}{r} \right| \le \|\partial_r h^r(t)\|_{L^\infty(B_{L/32})}.$$

This proves (4.70) and hence (4.71). □

Remark 4.29 (Relation with the periodic image-kernel language). In kernel notation, b_{loc} is the zero-image Euclidean singular Hodge field generated by the localized active vorticity, while h contains both the nonzero periodic images and the vorticity lying outside the fixed chart. Thus (4.71) is a quantitative replacement for the historical phrase "the nonzero periodic images are smooth and lower order." It uses only the physical kinetic-energy bound, not a pointwise enstrophy bound.

Lemma 4.30 (Scale-matched meridian Hodge detector)**.** *Let $R_j \downarrow 0$ with $R_j < L/16$, and let Y_j be meridian points in one lifted periodic chart with distance $O(R_j)$ from the symmetry axis. Suppose that at regular times t_j*

$$R_j^2 |\alpha(Y_j, t_j)| \geq a_0 > 0. \tag{4.72}$$

After passage to a subsequence, at least one of the following occurs:

(i) a strict degree-three $G^\perp$ target, a degree-two $q/q^\perp$ target at an intermediate dyadic comparison scale, or another already routed derivative/rapid target is selected;

(ii) there are radii $\ell_j \asymp R_j$, axial centers z_j, and $e_* > 0$ such that

$$\boxed{\ell_j^2 \int_{B^{\mathcal{Q}}_{C\ell_j}(z_j)} |q^\perp(r, z, t_j)|^2 \, dr \, dz \geq e_*.} \tag{4.73}$$

Proof. Choose the fixed periodic chart centered at the axial coordinate of Y_j and apply theorem 4.28. For all sufficiently large j,

$$\alpha(Y_j, t_j) = \alpha_{\mathrm{loc}}(Y_j, t_j) + \alpha_{\mathrm{rem}}(Y_j, t_j), \qquad R_j^2 |\alpha_{\mathrm{rem}}(Y_j, t_j)| \leq \frac{a_0}{4}.$$

Hence

$$R_j^2 |\alpha_{\mathrm{loc}}(Y_j, t_j)| \geq \frac{3a_0}{4}. \tag{4.74}$$

The localized Euclidean field is axisymmetric and is generated by $\chi G^\perp$. Use the five-dimensional lift

$$Y = (y' \in \mathbb{R}^4, z), \qquad r = |y'|.$$

The derivative of the Euclidean five-dimensional Newton kernel is $O(|Y - Y_j|^{-4})$. Split the local Hodge representation into $|Y - Y_j| < 2R_j$ and dyadic shells

$$\mathcal{A}^5_{D_k}(Y_j), \qquad D_k := 2^k R_j,$$

up to the fixed cutoff radius.

For the near term,

$$R_j^2 |\alpha_{<2R_j}(Y_j, t_j)| \leq C R_j^3 \|G^\perp(t_j)\|_{L^\infty(B^5_{2R_j}(Y_j))}.$$

If this carries a fixed fraction of (4.74), either an unbounded normalized $G^\perp$ amplitude/derivative gives item (i), or the lower bound persists on a fixed normalized subball. On a subball with $r \asymp R_j$, $q^\perp = rG^\perp$, which gives (4.73).

For a dyadic shell, Cauchy–Schwarz gives

$$|\alpha_k(Y_j, t_j)| \leq C D_k^{-3/2} \left(\int_{\mathcal{A}^5_{D_k}(Y_j)} |G^\perp|^2 \, dY \right)^{1/2}.$$

Define

$$\mathcal{E}_k := D_k \int_{\mathcal{A}^5_{D_k}(Y_j)} |G^\perp|^2 \, dY.$$

Then

$$\boxed{R_j^2|\alpha_k(Y_j,t_j)| \le C\left(\frac{R_j}{D_k}\right)^2 \mathcal{E}_k^{1/2}.} \tag{4.75}$$

At every relevant shell, failure of the intermediate envelope

$$D_k^2\|q(t_j)\|_{L^\infty(\mathrm{proj}\,\mathcal{A}^5_{D_k})} \le M_2 \tag{4.76}$$

along a subsequence selects the degree-two homogeneous length

$$a_k^{(q)} := \|q(t_j)\|_{L^\infty}^{-1/2} = o(D_k),$$

which is item (i) and is absorbed by the target-agnostic future clock. Thus on the complementary branch one fixed M_2 controls every relevant intermediate shell. Since $|q^\perp| \le 2\sup_z |q|$ at fixed r and

$$|G^\perp|^2\, dY \asymp r|q^\perp|^2\, dr\, dz,$$

while the projected shell has flat area $O(D_k^2)$ and $r \le CD_k$,

$$\boxed{\mathcal{E}_k \le CM_2^2.} \tag{4.77}$$

Therefore

$$\sum_{k\ge K} R_j^2|\alpha_k(Y_j,t_j)| \le CM_2 \sum_{k\ge K} 4^{-k}.$$

Choose K fixed so large that the right side is at most $a_0/8$. Together with (4.74), this forces the near term or one of the finitely many shells $0 \le k < K$ to carry a fixed normalized fraction of the strain.

In the shell case $D_k \asymp R_j$ because $k < K$ is fixed, and (4.75) forces $\mathcal{E}_k \ge e_0 > 0$. Finally

$$\mathcal{E}_k \le CD_k^2 \int_{B^{\mathcal{Q}}_{CD_k}(z_j)} |q^\perp|^2\, dr\, dz.$$

Taking $\ell_j = D_k \asymp R_j$ proves (4.73). □

Corollary 4.31 (Both selector branches enter the same-scale Hodge detector)**.** *Modulo the already routed smaller derivative/rapid alternatives, both branches of theorem 4.27 supply a scale $R_j \downarrow 0$ and a meridian point Y_j such that*

$$|y_j'| \le CR_j, \qquad R_j^2|\alpha(Y_j,t_j)| \ge c_* > 0.$$

Hence theorem 4.30 applies.

Proof. In item (i), take $R_j = R_j^\sharp$ and use (4.64). In item (ii), take $R_j = \eta_j$ and use (4.66). These are exactly the hypotheses of the meridian-point Hodge detector. □

Remark 4.32 (Optional persistence). If, on one fixed enlargement of the selected meridian box,

$$\ell_j^2\|q^\perp\|_\infty \le M_2, \qquad \ell_j^4\|\partial_t q^\perp\|_\infty \le M_4,$$

then the quantity

$$F_j(t) := \ell_j^2 \int \chi_{\ell_j} |q^\perp(t)|^2 \, dr \, dz$$

has $|F_j'(t)| \le C\ell_j^{-2}$. Thus the slice (4.73) persists for a fixed fraction of an ℓ_j^2 time. If either displayed normalized bound fails, the corresponding derivative/rapid target is already routed. The compact signed-child extraction below uses only the time slice itself.

Lemma 4.33 (Active slice produces a signed full-q child)**.** *Assume (4.73) at a scale ℓ_j, and exclude strict degree-two $q, q^\perp$ targets and strict degree-three first-derivative targets at that scale. Then, after passage to a subsequence, there are one fixed sign $s \in \{\pm 1\}$, one $\mu_* > 0$, centers $\zeta_j \in \mathbb{T}_L$, and compact unions of positive components $\Omega_j(t_j)$ contained in a fixed ℓ_j-normalized meridian box with $r \asymp \ell_j$ such that*

$$\boxed{\int_{\Omega_j(t_j)} sq(r, z, t_j) \, dr \, dz \ge \mu_*.} \tag{4.78}$$

Proof. The fixed normalized area of the box in (4.73) gives a point (r_j, z_j^0) with

$$|q^\perp(r_j, z_j^0, t_j)| \ge c_1 \ell_j^{-2}. \tag{4.79}$$

On the no-degree-three-target branch, normalized first derivatives of $q^\perp$ are bounded. Since $q^\perp(0, z, t) = 0$, the point may be chosen, after fixed losses, with $r_j \asymp \ell_j$.

There are two cases. If

$$|q(r_j, z_j^0, t_j)| \ge \frac{c_1}{2} \ell_j^{-2},$$

the normalized bound for ∇q propagates one fixed sign and a lower bound $c\ell_j^{-2}$ to a meridian ball of radius $c\ell_j$. Its flat area is $c\ell_j^2$, so it carries a fixed signed q mass.

Otherwise, by (4.67),

$$|\overline{q}(r_j, t_j)| = |q - q^\perp|(r_j, z_j^0, t_j) \ge \frac{c_1}{2} \ell_j^{-2}.$$

Choose s to be the sign of $\overline{q}(r_j, t_j)$. Since $\overline{q}$ is the axial average of the full field, there exists $\zeta_j \in \mathbb{T}_L$ such that

$$sq(r_j, \zeta_j, t_j) \ge |\overline{q}(r_j, t_j)| \ge \frac{c_1}{2} \ell_j^{-2}.$$

Again, either the normalized first derivative of q is unbounded and gives the routed degree-three target, or this lower bound persists on a ball of radius $c\ell_j$ and yields a fixed signed mass.

In either case, take the union of the positive components of sq meeting the selected ball and pass to a subsequence so that the sign s is fixed. This gives (4.78). □

Remark 4.34 (The mean mode helps rather than obstructs). The axial mean can cancel $q^{\perp}$ at a selected point only by becoming large itself. Because it is literally the axial average of the full toroidal vorticity, a large mean immediately produces another axial point where the full field has the same large sign. Thus the mean/nonzero-mode split creates no additional signed-child gate.

Lemma 4.35 (Recentered Type-II scale selection)**.** *Let $(0, z_*, T)$ be a singular periodic axis point and let $L > 0$ be a sufficiently small parent scale below the injectivity radius. For every integer $j \geq 1$ one may choose*

$$R_j^{\mathrm{II}} < \frac{L}{j^2} \qquad \text{with} \qquad \mathcal{E}_*(R_j^{\mathrm{II}}) \geq j^8.$$

Consequently some point in the corresponding backward cylinder satisfies

$$|\nabla u| \geq c \frac{\mathcal{E}_*(R_j^{\mathrm{II}})^{1/2}}{(R_j^{\mathrm{II}})^2}, \tag{4.80}$$

and the associated degree-two length

$$a_j^{(2)} := |\nabla u|^{-1/2}$$

obeys

$$a_j^{(2)} \leq C R_j^{\mathrm{II}} \mathcal{E}_*(R_j^{\mathrm{II}})^{-1/4} \leq C \frac{L}{j^4}. \tag{4.81}$$

Using $a_j^{(2)}$ only as the next ambient ruler in theorem 4.27, one obtains, after routing any still smaller derivative/rapid alternative, an ambient selector scale $S_j = o(L)$ and a detector scale $R_j \leq C S_j$ satisfying the hypotheses of theorem 4.30. The selected strain point, Hodge witness and resulting signed child time lie in the terminal selector window

$$\boxed{J_j^{\mathrm{sel}} \subset [T - S_j^2, T], \qquad |J_j^{\mathrm{sel}}| \leq S_j^2.} \tag{4.82}$$

Thus a same-scale compact Hodge-active toroidal witness is reselected and

$$\frac{|J_j^{\mathrm{sel}}|}{L^2} \longrightarrow 0.$$

No inheritance of the point in (4.80) is required.

Proof. The choice of R_j^{II} follows from theorem 4.9. The parabolic cylinder has spacetime volume $O(R^5)$ and

$$\int_{Q(R)} |\nabla u|^2 \, dX \, dt = R \, \mathcal{E}_*(R),$$

so the mean-value estimate gives (4.80), hence (4.81). For the final statement, use $S_j = a_j^{(2)}$ as the ambient scale in theorem 4.27. By theorem 4.31, either a still smaller routed target occurs or the selector supplies a geometry-correct scale for theorem 4.30. Its non-target branch gives (4.73). Here the selector is applied with ambient scale $S_j = a_j^{(2)}$, so every selected time belongs to

$P(S_j) \subset [T - S_j^2, T]$. In the compact selector branch the Hodge scale is comparable to S_j; in the collapsing branch it is $R_j^\sharp = o(S_j)$. Thus in both cases the detector scale satisfies $R_j \le CS_j$, while $S_j/L \le Cj^{-4}$. This proves (4.82) and the subparabolic relation relative to the parent. Iteration produces the recentered super-geometric relay. □

Proposition 4.36 (B.2: periodic active/spectator/recentering routing)**.** *Every terminal failure of compact periodic strain chronology is routed, after subselection, to one of: a strict homogeneous/rapid target which is reabsorbed at a smaller physical scale; a recurrent singular axis point with a recentered compact local strain packet; or a recentered super-geometric active parent–child chain carrying a compact Hodge-active witness and a fixed signed full-q child. Normalized axial delocalization is therefore not an independent periodic terminal mechanism.*

Proof. Decompose the periodic toroidal field into its axial mean and nonzero Fourier part. The mean mode is compression-silent for the meridional Hodge response; hence a terminally active packet contains either a nonzero-mode Hodge witness or an already routed strict target. The active strain endpoint theorem 4.24 and the finite first moment theorem 4.25 give the recurrent singular axis point theorem 4.26. At that point apply theorem 4.35; this produces the recentered super-geometric active scales and same-scale toroidal witnesses, modulo only the already named derivative/rapid alternatives. The active slice is converted to a fixed signed full-q child by theorem 4.33; together with the parent local action budget this is packaged below as C.2-core. □

Lemma 4.37 (Periodic parent-scale local flat-action budget)**.** *Let $L_j \downarrow 0$ be periodic parent scales and let $K_{CL_j}(t)$ be moving meridian boxes whose flat areas satisfy*

$$|K_{CL_j}(t)|_{dr\,dz} \le C_K L_j^2$$

on terminal intervals

$$I_j^{\mathrm{par}} := [T - C_0 L_j^2, T].$$

After subselection, at least one of the following occurs:

(i)

$$L_j^2 \|q\|_{L^\infty(K_{CL_j} \times I_j^{\mathrm{par}})} \longrightarrow \infty, \tag{4.83}$$

and hence the degree-two homogeneous radius

$$a_j^{(q)} := \|q\|^{-1/2}_{L^\infty(K_{CL_j} \times I_j^{\mathrm{par}})}$$

satisfies $a_j^{(q)}/L_j \to 0$;

(ii) there is one fixed $M_q < \infty$ such that

$$|q(r,z,t)| \le M_q L_j^{-2} \qquad ((r,z,t) \in K_{CL_j}(t) \times I_j^{\mathrm{par}}), \tag{4.84}$$

and consequently

$$\boxed{\int_{I_j^{\mathrm{par}}}\int_{K_{CL_j}(t)} q^2\,dr\,dz\,dt \le C_0C_KM_q^2 =: A_*.} \tag{4.85}$$

The lemma depends only on the parent scale L_j. No relation between the child detector scale R_j and selector-window scale S_j is used.

Proof. If the dimensionless quantities in (4.83) are unbounded, pass to a divergent subsequence. Then

$$\frac{a_j^{(q)}}{L_j} = \left(L_j^2\|q\|_{L^\infty(K_{CL_j}\times I_j^{\mathrm{par}})}\right)^{-1/2} \longrightarrow 0,$$

which is a strict degree-two q target.

On the complementary subsequence the dimensionless supremum is uniformly bounded by some M_q, giving (4.84). The flat area bound and $|I_j^{\mathrm{par}}| = C_0L_j^2$ give

$$\int_{I_j^{\mathrm{par}}}\int_{K_{CL_j}(t)} q^2\,dr\,dz\,dt \le (M_q^2L_j^{-4})(C_KL_j^2)(C_0L_j^2),$$

which is (4.85). □

Remark 4.38 (Two different scales play two different roles). In the active periodic bridge, the selector scale $S_j = o(L_j)$ controls the fresh-child time interval and hence

$$|J_j^{\mathrm{sel}}|/L_j^2 \to 0.$$

The parent scale L_j controls the fixed local flat-action reservoir (4.85) used by the logarithmic action-depth argument in Bridge D. These roles should not be conflated. In particular, the budget does not require an $O(R_j^2)$ or $O(S_j^2)$ integration window.

Proposition 4.39 (C.2-core: periodic compact signed-child interface)**.** *Assume the active periodic SC.2/Type-II/B.2 routing has produced parent, selector-window and child scales*

$$L_j \downarrow 0, \qquad R_j \le CS_j, \qquad \frac{S_j}{L_j} \longrightarrow 0,$$

where z_* anchors the recurrent Type-II/strain selection, while the full-q child is allowed to recenter at an axial point $\zeta_j \in \mathbb{T}_L$. Assume the child time lies in the terminal selector window

$$J_j^{\mathrm{sel}} \subset [T - CS_j^2, T]$$

and carries the compact Hodge-active toroidal witness from theorem 4.35. After excluding the already routed strict homogeneous/rapid and normalized spatial/recentering alternatives, the following hold:

(i) the creation window is subparabolic relative to the parent:

$$\boxed{\frac{|J_j^{\mathrm{sel}}|}{L_j^2} \le C\left(\frac{S_j}{L_j}\right)^2 \longrightarrow 0;} \tag{4.86}$$

(ii) after a sign subselection there is a child time $t_j^+ \in J_j^{\mathrm{sel}}$ and a compact component union, possibly recentered at ζ_j, satisfying

$$\boxed{\int_{\Omega_j(t_j^+)} sq\,dr\,dz \ge \mu_* > 0;} \tag{4.87}$$

(iii) there is a fixed parent-scale moving meridian box $K_{CL_j}(t)$ and $A_* < \infty$ such that

$$\boxed{\int_{T-C_0L_j^2}^{T}\int_{K_{CL_j}(t)} q^2\,dr\,dz\,dt \le A_*.} \tag{4.88}$$

No retained-ancestry conclusion is included in C.2-core.

Proof. The same-scale active slice is (4.73), so theorem 4.33 gives (4.87) after the already routed degree-two/three alternatives are removed. For the parent corridor apply theorem 4.37. Its first alternative is an already routed strict degree-two q target. On the complementary branch (4.85) is exactly (4.88), after renaming the fixed constant A_*. This parent budget is independent of the selector-window scale S_j used in item (i). No ancestry theorem is used here. □

Proposition 4.40 (Periodic axial-translation covariance after recurrent-core selection)**.** *For $\zeta \in \mathbb{T}_L$ let*

$$(\mathcal{T}_\zeta f)(r,z,t) := f(r,z+\zeta,t).$$

The recurrent point z_* is used only to anchor the local suitability, Type-II and strain-selector stages. After a compact signed full-q child has been selected, every object used in C.2-core, Bridge D and Bridge E is covariant under $\mathcal{T}_\zeta$. In particular the child center ζ_j need not equal z_*, and no later restart inherits an absolute periodic center, sign or profile.

Proof. Axial translation is an isometry of $\mathbb{R}^2 \times \mathbb{T}_L$ and commutes with the periodic Laplacian, curl/divergence, the axial Fourier decomposition, periodic Hodge operators, the Navier–Stokes evolution, material maps and the full viscous adjoint. Both $dr\,dz$ and dX are invariant. The local flat-action budget in (4.88) is unchanged after translating its moving box, and the signed-child mass (4.87) is likewise invariant.

The mean-mode alternative in theorem 4.33 may choose a new ζ_j because a large axial mean of q guarantees a full-q point of the same sign somewhere on the circle. This recentering happens only after the Type-II/strain selector has supplied the Hodge-active scale. From that point onward Bridges D and E use only local packets and translation-covariant equations. Strict

restarts are absorbed by the geometry-appropriate future clock without center inheritance. □

5. Bridge D: compact signed child to active-origin alternatives

From this point onward $\mathcal{Q}_*$ denotes either $\mathcal{Q}$ or $\mathcal{Q}_L$. C.1 and C.2-core supply a super-geometric compact corridor, a fixed signed child, and a fixed local flat-action budget. Retained ancestry has deliberately not been excluded upstream; it is handled here, strictly downstream, by the action-depth proposition. Put

$$p := sq, \qquad P_s := sP_\theta, \qquad P_\theta := r\beta^2. \tag{5.1}$$

5.1. Creation–Robin identity.

Lemma 5.1 (Sign-symmetric creation–Robin square)**.** *On a regular positive component of p, let $g = |\nabla p|$ and let n be the outward meridian normal. Then*

$$\boxed{\frac{P_\theta^2}{4\nu g} - (P_s n_z - \nu g) = \frac{|2\nu\nabla p + P_s e_z|^2}{4\nu g}.} \tag{5.2}$$

After integration over the moving nodal boundary and a time interval J,

$$\boxed{\mathbf{I}_J = 4\nu\Delta M_J + \mathbf{R}_J, \qquad \mathbf{I}_J, \mathbf{R}_J \geq 0.} \tag{5.3}$$

Proof. Equation (5.2) is completion of the square, using $n = -\nabla p/|\nabla p|$ on a positive component. Reynolds transport applied to (2.5) gives the component mass identity; integrating (5.2) gives (5.3). □

Lemma 5.2 (Source-to-nodal coercivity)**.** *Let a complete positive genealogy on J have signed source input*

$$\mathsf{S}_J := \int_J \int_{\mathcal{G}(t)} F_s \, dr\, dz\, dt > 0$$

and endpoint mass increment ΔM_J. Then

$$\Delta M_J = \mathsf{S}_J - \nu \mathsf{B}_J, \qquad \mathsf{B}_J := \int_J \sum_{\Omega \subset \mathcal{G}(t)} \int_{\partial\Omega} |\nabla p| \, d\mathcal{H}^1 \, dt, \tag{5.4}$$

and boundary Cauchy–Schwarz gives

$$\mathsf{S}_J^2 \leq \mathbf{I}_J \mathsf{B}_J.$$

Consequently

$$\boxed{\mathbf{I}_J \geq \nu \frac{\mathsf{S}_J^2}{\mathsf{S}_J - \Delta M_J}.} \tag{5.5}$$

In particular, a fixed positive source input on a uniformly mass-bounded genealogy forces a generation-independent nodal creation charge unless the episode creates a fixed positive mass increment.

Proof. Equation (5.4) is the integrated positive-component mass balance. Substitute $\mathsf{B}_J = (\mathsf{S}_J - \Delta M_J)/\nu$ into $\mathsf{S}_J^2 \leq \mathbf{I}_J \mathsf{B}_J$. □

5.2. **Logarithmic action-depth obstruction.** Fix once and for all a smooth axis-centered cutoff $\chi(\rho,\zeta)$, even in the signed radial variable, with

$$0 \le \chi \le 1, \qquad \chi = 1 \text{ on } B_1, \qquad \chi = 0 \text{ outside } B_2.$$

For a moving axis center $Z(t)$ and a scale S put

$$\chi_S(r,z,t) := \chi\left(\frac{r}{S}, \frac{z - Z(t)}{S}\right), \tag{5.6}$$

and define the positive cutoff mass

$$m_S(t) := \int_{\mathcal{Q}_*} \chi_S(r,z,t) p_+(r,z,t)\, dr\, dz. \tag{5.7}$$

The center is transported by the axial velocity on the axis,

$$\dot{Z}(t) = u^z(0, Z(t), t).$$

Lemma 5.3 (Kato mass-speed bound at one clean scale)**.** *Assume on one S-scale moving cylinder that*

$$S^2\|q\|_{L^\infty} \le M_q, \qquad S^2\|\nabla u\|_{L^\infty} \le M_u, \qquad S^4\|\partial_z P_s\|_{L^\infty} \le M_P. \tag{5.8}$$

Then the upper Dini derivative satisfies

$$\boxed{D^+ m_S(t) \le \frac{C_*}{S^2},} \tag{5.9}$$

where C_* depends only on the fixed cutoff, ν and M_q, M_u, M_P.

Proof. A convex approximation to $x \mapsto x_+$ in the signed equation gives the distributional Kato inequality

$$\partial_t p_+ + \operatorname{div}_{r,z}(b p_+) \le \nu \Delta_1 p_+ + |\partial_z P_s|.$$

Testing with χ_S and integrating in the explicitly flat measure gives

$$\begin{aligned} D^+ m_S \le & \int_{\mathcal{Q}_*} p_+ \left(\partial_t \chi_S + b \cdot \nabla \chi_S\right) dr\, dz \\ & + \nu \int_{\mathcal{Q}_*} p_+ \left((\chi_S)_{rr} + (\chi_S)_{zz} - \frac{(\chi_S)_r}{r}\right) dr\, dz \\ & + \int_{\mathcal{Q}_*} \chi_S |\partial_z P_s|\, dr\, dz. \end{aligned}$$

The axis boundary term vanishes because $p = q = 0$ there and the cutoff is even in the signed radial variable. Smooth-axis geometry and the definition of $Z(t)$ give, on $\operatorname{supp} \nabla \chi_S$,

$$|u^r| + |u^z - \dot{Z}| \le CS\|\nabla u\|_{L^\infty} \le C M_u S^{-1}.$$

Moreover

$$|\nabla \chi_S| \le C S^{-1}, \qquad \left|(\chi_S)_{rr} + (\chi_S)_{zz} - \frac{(\chi_S)_r}{r}\right| \le C S^{-2}.$$

The transition region has flat area $O(S^2)$ and $|p_+| \le M_q S^{-2}$. Each term is therefore bounded by $C_* S^{-2}$. □

Remark 5.4 (Failure of a clean-scale envelope is already a routed target). Along an S_j-scale sequence, unboundedness of either of the first two quantities in (5.8) selects a strict degree-two $q/\nabla u$ target; unboundedness of the third selects the degree-four source scale $|\partial_z P_s|^{-1/4} = o(S_j)$. Thus theorem 5.3 is precisely the complementary no-target branch.

Lemma 5.5 (One fixed flat-action quantum per clean geometric scale)**.** *Let a compact positive genealogy connect a parent comparison time t_j^- at scale L_j to a child time t_j^+ at scale $R_j = o(L_j)$, with*

$$\int_{K_{CR_j}(t_j^+)} p_+(r, z, t_j^+)\, dr\, dz \geq \mu_* > 0. \tag{5.10}$$

Assume the clean envelopes (5.8) at all intermediate scales. Then there are constants $a_* > 0$ and $0 < \delta < 1/8$ and a geometric ladder

$$S_{j,0} = \delta L_j, \qquad S_{j,k+1} = \delta S_{j,k}, \qquad S_{j,N_j} \asymp R_j,$$

with

$$N_j \geq c_\delta \log \frac{L_j}{R_j} - C_\delta, \tag{5.11}$$

and ordered times

$$t_j^- \leq \tau_{j,0} < \tau_{j,1} < \cdots < \tau_{j,N_j} \leq t_j^+$$

such that

$$\tau_{j,k+1} - \tau_{j,k} \geq c_* S_{j,k+1}^2 \tag{5.12}$$

and

$$\boxed{\int_{\tau_{j,k}}^{\tau_{j,k+1}} \int_{K_{2S_{j,k}}(t)} q^2\, dr\, dz\, dt \geq a_* > 0.} \tag{5.13}$$

Proof. Set

$$\theta := \frac{\mu_*}{4}$$

and choose δ so small that

$$C_\chi M_q \delta^2 \leq \frac{\theta}{2},$$

where $C_\chi S^2$ bounds the flat support area of χ_S. At the parent comparison time the degree-two q envelope gives

$$m_{S_{j,0}}(t_j^-) \leq \frac{\theta}{2}.$$

At the child time, choosing the terminal ladder constant sufficiently large, (5.10) gives

$$m_{S_{j,k}}(t_j^+) \geq 2\theta \qquad (0 \leq k \leq N_j).$$

For each k let $\tau_{j,k}$ be the last time at which $m_{S_{j,k}} \le \theta$. The cutoffs are nested, so these times are ordered. At $\tau_{j,k}$, the next cutoff has flat support area $O(S_{j,k+1}^2)$; the q envelope at scale $S_{j,k}$ gives

$$m_{S_{j,k+1}}(\tau_{j,k}) \le C_\chi M_q \frac{S_{j,k+1}^2}{S_{j,k}^2} \le \frac{\theta}{2}.$$

To reach level θ by $\tau_{j,k+1}$, the smaller cutoff mass must increase by at least $\theta/2$. Applying theorem 5.3 at scale $S_{j,k+1}$ gives (5.12).

By the definition of the last crossing,

$$m_{S_{j,k}}(t) \ge \theta \qquad (\tau_{j,k} \le t \le \tau_{j,k+1}).$$

Cauchy–Schwarz on a set of flat area $O(S_{j,k}^2)$ therefore gives

$$\int_{K_{2S_{j,k}}(t)} q^2 \, dr \, dz \ge \frac{c\theta^2}{S_{j,k}^2}.$$

Integrating over the time interval and using $S_{j,k+1} = \delta S_{j,k}$ proves (5.13), with $a_* = c\theta^2\delta^2$. □

Proposition 5.6 (Transversality-free action depth)**.** *Let a compact positive genealogy carry fixed signed child mass through a parent–child corridor with scales L_j and $R_j = o(L_j)$. After routing intermediate degree-two $q/\nabla u$ targets, degree-four source targets and spatial/recentering exits, retained ancestry forces*

$$\boxed{\int_{J_j} \int_{K_{CL_j}(t)} q^2 \, dr \, dz \, dt \ge c_* \log \frac{L_j}{R_j} - C_*.} \tag{5.14}$$

Hence retained ancestry is impossible under the fixed corridor budgets of C.1 or C.2-core.

Proof. If one of the envelopes (5.8) fails at an intermediate ladder scale, theorem 5.4 gives an already routed strict target. If the genealogy leaves every fixed normalized moving parent box, the spatial/recentering alternative has occurred. On the remaining compact clean branch apply theorem 5.5. All ladder boxes lie in one fixed multiple of the moving parent box, so summing (5.13) over N_j scales and using (5.11) gives (5.14). Since $L_j/R_j \to \infty$, the right side diverges, contradicting the fixed local flat-action budget. □

5.3. **Source-born versus bare active resets.**

Proposition 5.7 (Active-origin decomposition)**.** *Every infinite terminal compact reset orbit from C.1 or C.2-core has, after subselection, one of the following forms:*

(i) infinitely many resets are source-born; then a complete positive genealogy carries fixed signed source input and enters the polarized positive-genealogy route of Bridge E;

(ii) all sufficiently late resets are bare at the moment of descent; then either the reset is a spectator, or a fixed active fraction is extracted and enters the retained active-mass route of Bridge E.

Proof. The alternatives source-born/bare are exhaustive. On a source-born episode, the complete positive genealogy has a fixed signed source input and bounded mass. The source-to-nodal coercivity theorem 5.2 supplies a fixed viscosity-visible creation token, but the token is not declared terminal; it is passed to the material-vorticity polarizer in Bridge E. On the complement there is no fresh fixed-mass source birth, so a continuing branch must first carry a fixed active fraction; otherwise it is a spectator. □

Lemma 5.8 (Flat mass/action controls the scale-matched Hodge drift)**.** *Let $f = f(r,z,t) \geq 0$ be supported for $t \in J$ in one axis-centered meridian box of diameter Cs, with $|J| \leq Cs^2$. Put*

$$\mu_f := \sup_{t\in J} \int f(r,z,t)\, dr\, dz, \qquad A_f := \int_J \int f(r,z,t)^2\, dr\, dz\, dt.$$

Let

$$\omega_f := f e_\theta, \qquad v_f := \nabla \times (-\Delta)^{-1} \omega_f$$

in one Euclidean three-dimensional chart, with a spatially constant axial part absorbed into the common Galilean gauge. Then

$$\boxed{s^{-14/15} \|v_f\|_{L_t^6 L_x^{15/8}(B_{Cs}\times J)} \leq C \mu_f^{2/3} A_f^{1/6}.} \tag{5.15}$$

Proof. At fixed time set

$$a(t) := \|f(t)\|_{L^2(dr\, dz)}.$$

Since $r \leq Cs$ on the support,

$$\|\omega_f(t)\|_{L^1(dX)} = 2\pi \int r f\, dr\, dz \leq Cs\mu_f,$$

and

$$\|\omega_f(t)\|_{L^2(dX)} \leq Cs^{1/2} a(t).$$

Interpolation gives

$$\|\omega_f(t)\|_{L^{6/5}(dX)} \leq Cs^{5/6} \mu_f^{2/3} a(t)^{1/3}. \tag{5.16}$$

The order-minus-one Hodge operator obeys

$$\|v_f(t)\|_{L^2(dX)} \leq C \|\omega_f(t)\|_{L^{6/5}(dX)}. \tag{5.17}$$

Indeed, pair v_f with $\phi \in L^2(dX)$, move $\nabla \times (-\Delta)^{-1}$ onto ϕ, and use the three-dimensional Sobolev inequality to place that potential in $L^6(dX)$.

Because $15/8 < 2$ and $|B_{Cs}| \asymp s^3$,

$$\|v_f(t)\|_{L^{15/8}(B_{Cs},dX)} \leq Cs^{1/10} \|v_f(t)\|_{L^2(dX)} \leq Cs^{14/15} \mu_f^{2/3} a(t)^{1/3}.$$

Finally

$$\|a^{1/3}\|_{L^6(J)} = \left(\int_J a(t)^2\,dt\right)^{1/6} = A_f^{1/6}.$$

Divide by $s^{14/15}$. □

Corollary 5.9 (Bounded-overlap Hodge packing)**.** *For a bounded-overlap family of scale-assigned sign-separated packets f_k with scales s_k, put*

$$H_k := s_k^{-14/15}\|v_{f_k}\|_{L_t^6 L_x^{15/8}}.$$

Then

$$\boxed{\sum_k H_k^6 \le C\left(\sup_k \mu_{f_k}\right)^4 \sum_k A_{f_k}.} \tag{5.18}$$

In particular, if

$$\sum_k A_{f_k} \le C A_*, \qquad \left(\sum_k H_k^6\right)^{1/6} \ge h_* > 0,$$

then some packet satisfies

$$\boxed{\mu_{f_k} \ge c\,h_*^{3/2} A_*^{-1/4} =: \mu_0 > 0.} \tag{5.19}$$

Proof. Raise (5.15) to the sixth power and sum. Bounded overlap gives

$$\sum_k A_{f_k} \le C \int_{\text{parent}} \int q^2\,dr\,dz\,dt.$$

The last assertion follows by solving (5.18) for $\sup_k \mu_{f_k}$. □

Proposition 5.10 (Target-resistant active-mass extraction)**.** *Consider the toroidal source realization of a bare active reset inside a parent corridor with fixed flat action budget. Assign strict Hodge target cells to their first active scales with bounded overlap. Then, after subselection, either the family is collectively Hodge-silent and is a spectator, a fixed-mass positive genealogy is selected, a fixed positive source amount creates a source-born positive genealogy, or the surviving fixed-mass source-poor alternative is a retained active-mass relay.*

Proof. Choose one sign on each assigned packet and apply theorem 5.9. If the aggregate scale-matched Hodge score tends to zero, the family is spectator data for the active source/circulation thread. Otherwise (5.19) selects a packet with fixed signed flat mass μ_0.

Trace the complete positive genealogy of that packet backward through the current parent corridor. If its mass drops below a fixed fraction of μ_0, the positive-component mass balance shows that the intervening episode received a fixed positive signed source amount; it is therefore source-born and theorem 5.2 supplies the viscosity-visible nodal token. If no such drop occurs, the fixed mass was already present before the final target descent.

On intervals with fixed positive source input the same lemma returns to the source-born/nodal branch. The complementary continuing case is precisely a fixed-mass source-poor retained relay. □

Theorem 5.11 (Bridge D: compact signed child to active-origin alternatives)**.** *The output of either C.1 or C.2-core is reduced, after subselection, to one of:*

(i) a source-born complete positive genealogy carrying a fixed active token;
(ii) a retained fixed-mass active relay;
(iii) a strict physical/spatial restart;
(iv) a Hodge-silent spectator, which is discarded from the active thread.

No additional historical gate is inserted between the geometry-specific compact bridge and the common polarized downstream analysis.

Proof. Use theorem 5.6 to exclude retained unstructured ancestry. Apply theorems 5.7 and 5.10. The only non-spectator outputs are fixed active genealogies/relays or already routed strict restarts. □

6. Bridge E: successive polarization of the common downstream

The purpose of Bridge E is to remove, in the correct order, every Euler-compatible background before using a viscosity-dependent signal. The proof is therefore organized by *successive polarization*: first the material transport/stretching of vorticity is canceled, then the deformation of the adjoint detector is canceled, and finally only a physical viscosity-weighted current is allowed to trigger a new restart.

6.1. **Axis-away detector extraction.**

Lemma 6.1 (Automatic axis-away toroidal detector)**.** *Let a compact terminal child of physical scale R satisfy, for one sign $s \in \{\pm 1\}$,*

$$\int_{D_R} (sq)_+ \, dr\, dz \geq \mu_* > 0,$$

where D_R is contained in a meridian box of diameter CR. After subselection, either

$$R^2 \|q\|_{L^\infty(D_R)} \longrightarrow \infty, \tag{6.1}$$

which is a strict degree-two toroidal-vorticity restart, or there are fixed $\delta, c > 0$ and smooth cutoffs χ_R supported where $\delta R < r < CR$ such that

$$\left| \int q(r, z, t_+) \chi_R(r, z) \, dr\, dz \right| \geq c\mu_*. \tag{6.2}$$

For

$$\psi_R(X) := \frac{s\chi_R(r, z)}{r} \, e_\theta, \tag{6.3}$$

one has

$$\boxed{|\langle \omega(t_+), \psi_R \rangle_{L^2(dX)}| \geq c_0, \qquad \|\psi_R\|^2_{L^2(dX)} \leq CR.} \tag{6.4}$$

Proof. On the complement of (6.1), $|q| \le MR^{-2}$. The flat area of $\{0 < r < \delta R\} \cap D_R$ is $O(\delta R^2)$; hence the positive mass there is at most $CM\delta$. Choose δ so that this is at most $\mu_*/4$ and cut off a fixed annular share of the remaining mass. Since $dX = 2\pi r\, dr\, dz\, d\theta/(2\pi)$ after axisymmetric integration, (6.4) follows directly. □

6.2. **First polarization: Cauchy material vorticity.** Let $X(a,t)$ be the full three-dimensional material map,

$$\partial_t X(a,t) = u(X(a,t),t), \qquad F(a,t) := D_a X(a,t).$$

Incompressibility gives $\det F \equiv 1$.

Lemma 6.2 (Cauchy-polarized vorticity identity)**.** *Define*

$$\Omega_C(a,t) := F(a,t)^{-1}\omega(X(a,t),t).$$

Then

$$\boxed{\partial_t \Omega_C(a,t) = \nu F(a,t)^{-1}\Delta\omega(X(a,t),t).} \tag{6.5}$$

In particular Ω_C is exactly frozen when $\nu = 0$.

Proof. The vorticity equation is

$$D_t\omega = (\nabla u)\omega + \nu\Delta\omega, \qquad \partial_t F = (\nabla u)(X,t)F.$$

Differentiate $F^{-1}\omega(X,t)$ and cancel the stretching terms. □

Thus the apparent toroidal source $2\beta W$ is not counted as fresh material vorticity creation: in the Cauchy frame it belongs to Euler-compatible stretching/tilting. Only the explicit $\nu\Delta\omega$ term remains.

Proposition 6.3 (Finite-time Cauchy dichotomy)**.** *Consider a detector-charged handoff of assigned scale L and duration τ. Assume the material deformation is controlled on the corresponding tube. If a fixed normalized Cauchy-vorticity token loses a fixed fraction over the handoff, then*

$$\sup_{\text{tube}} |\Delta\omega| \ge \frac{c}{\nu\tau L^2}, \tag{6.6}$$

and the degree-four radius $a_{\omega,4} := |\Delta\omega|^{-1/4}$ satisfies

$$\boxed{\frac{a_{\omega,4}}{L} \le C\left(\frac{\nu\tau}{L^2}\right)^{1/4}.} \tag{6.7}$$

If no fixed loss occurs, the charged Cauchy token is inherited and no temporal successor is declared.

Proof. Integrate (6.5) over the handoff interval and use the boundedness of $F^{\pm 1}$. The scale L^{-2} is the vorticity normalization. □

6.3. **Second polarization: the full viscous vorticity adjoint.** Let η solve backward from the terminal detector (6.3):

$$\boxed{\partial_t \eta + u \cdot \nabla \eta + (\nabla u)^T \eta + \nu \Delta \eta = 0, \qquad \eta(t_+) = \psi_R.} \tag{6.8}$$

Then

$$\frac{d}{dt} \int \omega \cdot \eta \, dX = 0. \tag{6.9}$$

Put

$$Y(t) := \int |\eta|^2 \, dX, \qquad \mathfrak{s}_\eta(t) := \frac{\int \eta \cdot S\eta \, dX}{Y(t)}, \qquad \mathfrak{d}_\eta(t) := \frac{\int |\nabla \eta|^2 \, dX}{Y(t)},$$

where $S = (\nabla u + \nabla u^T)/2$. Here and throughout the adjoint subsection, bare $\| \cdot \|_2$ norms are physical $L^2(r\, dr\, dz)$ norms (equivalently axisymmetric $L^2(dX)$ up to the factor 2π).

Lemma 6.4 (Detector-charged strain identity)**.** *One has*

$$\frac{1}{2} Y'(t) = - \int \eta \cdot S\eta \, dX + \nu \int |\nabla \eta|^2 \, dX, \tag{6.10}$$

and hence

$$\boxed{\int_{t_0}^{t_+} \mathfrak{s}_\eta(t) \, dt - \nu \int_{t_0}^{t_+} \mathfrak{d}_\eta(t) \, dt = \log \frac{\|\eta(t_0)\|_2}{\|\psi_R\|_2}.} \tag{6.11}$$

Whenever the conserved pairing (6.9) *gives a fixed lower bound for* $\|\eta(t_0)\|_2$*, theorem 6.1 yields*

$$\boxed{\int_{t_0}^{t_+} (\mathfrak{s}_\eta)_+(t) \, dt \geq \frac{1}{2} \log \frac{1}{R} - C.} \tag{6.12}$$

Proof. Multiply (6.8) by η, integrate in dX, use $\nabla \cdot u = 0$, and note that the antisymmetric part of ∇u drops out. Integrate the logarithmic derivative of Y. □

The essential point is that the logarithmic strain is carried by the same adjoint mass which detects the terminal toroidal packet; it is not an unrelated global supremum.

6.4. **Material localization and the no-teleportation alternative.** Let χ be transported by the full velocity:

$$(\partial_t + u \cdot \nabla)\chi = 0. \tag{6.13}$$

For

$$Y_\chi(t) := \int \chi^2 |\eta|^2 \, dX$$

one has the exact localized identity

$$\boxed{\frac{1}{2} Y_\chi' = - \int \chi^2 \eta \cdot S\eta \, dX + \nu \int |\nabla(\chi\eta)|^2 \, dX - \nu \int |\nabla \chi|^2 |\eta|^2 \, dX.} \tag{6.14}$$

Thus there is no artificial Euler transport current.

The last term has an intrinsic diffusion scale. If the transition thickness is ℓ and the handoff duration is τ, then

$$\nu \int_J \frac{\int |\nabla\chi|^2 |\eta|^2 \, dX}{\int |\eta|^2 \, dX} \, dt \le C \frac{\nu\tau}{\ell^2}. \tag{6.15}$$

Hence fixed leakage already selects $\ell^2 \lesssim \nu\tau$.

For the remaining localization issue use terminal-anchored labels $X(a, t_+) = a$ and

$$\Xi(a,t) := F(a,t)^T \eta(X(a,t),t). \tag{6.16}$$

Then

$$\boxed{\partial_t \Xi = -\nu F^T \Delta\eta(X(a,t),t).} \tag{6.17}$$

At $\nu = 0$ the pulled-back covector is exactly frozen.

Let reversed time be

$$s := t_+ - t, \qquad 0 \le s \le \tau, \tag{6.18}$$

and write, for notational convenience,

$$\widehat{F}(a,s) := F(a, t_+ - s), \qquad \widehat{H}(a,s) := \widehat{F}(a,s)^{-T}, \qquad \widehat{\Xi}(a,s) := \Xi(a, t_+ - s).$$

Set

$$G(a,s) := \widehat{F}(a,s)^{-1} \widehat{F}(a,s)^{-T}. \tag{6.19}$$

Because $\det F \equiv 1$, the scalar Laplacian pulled back by X satisfies, componentwise,

$$(\Delta_x f)(X(a,t),t) = \partial_{a_m} \Big(G_{mn}(a,t) \, \partial_{a_n} \widetilde{f}(a,t) \Big), \qquad \widetilde{f}(a,t) := f(X(a,t),t). \tag{6.20}$$

Lemma 6.5 (Coefficient form of the pulled-back adjoint)**.** *The covector $\widehat{\Xi}$ solves*

$$\boxed{\partial_s \widehat{\Xi}_\alpha - \nu \partial_{a_m} \Big(G_{mn} \partial_{a_n} \widehat{\Xi}_\alpha \Big) = \nu b^k_{\alpha\beta} \, \partial_{a_k} \widehat{\Xi}_\beta + \nu c_{\alpha\beta} \widehat{\Xi}_\beta,} \tag{6.21}$$

where, with repeated indices summed,

$$b^k_{\alpha\beta} := -(\partial_{a_m} \widehat{F}_{i\alpha}) G_{mk} \widehat{H}_{i\beta} + \widehat{F}_{i\alpha} G_{kn} (\partial_{a_n} \widehat{H}_{i\beta}), \tag{6.22}$$

$$c_{\alpha\beta} := \widehat{F}_{i\alpha} \partial_{a_m} \Big(G_{mn} \partial_{a_n} \widehat{H}_{i\beta} \Big). \tag{6.23}$$

If

$$\|\widehat{F}\| + \|\widehat{F}^{-1}\| + L\|D_a \widehat{F}\| + L^2 \|D_a^2 \widehat{F}\| \le M \tag{6.24}$$

on the connecting label chart, then there are constants $\lambda_M, \Lambda_M, C_M > 0$ such that

$$\lambda_M |\xi|^2 \le G_{mn} \xi_m \xi_n \le \Lambda_M |\xi|^2, \tag{6.25}$$

$$L\|D_a G\|_\infty + L\|b\|_\infty + L^2 \|D_a b\|_\infty + L^2 \|c\|_\infty \le C_M. \tag{6.26}$$

Proof. From (6.17) and $\eta(X(a,t),t) = F(a,t)^{-T}\Xi(a,t)$, (6.20) gives

$$\partial_s \widehat{\Xi}_\alpha = \nu \widehat{F}_{i\alpha}\partial_{a_m}\Big[G_{mn}\partial_{a_n}\big(\widehat{H}_{i\beta}\widehat{\Xi}_\beta\big)\Big].$$

Apply the product rule once to move $\widehat{F}_{i\alpha}$ through ∂_{a_m}, use $\widehat{F}_{i\alpha}\widehat{H}_{i\beta} = \delta_{\alpha\beta}$, and collect the first- and zeroth-order terms. This gives (6.21)–(6.23). Uniform ellipticity of G follows from the bounds on $\widehat{F}$ and $\widehat{F}^{-1}$. Differentiating $\widehat{F}^{-1}$ and G shows that (6.26) follows from (6.24). □

We now isolate the only parabolic estimate used in the no-teleportation step.

Lemma 6.6 (Off-diagonal L^2–L^∞ estimate for the pulled-back system)**.** *Let $U : \mathbb{R}^3 \times [0,\tau] \to \mathbb{R}^3$ solve*

$$\partial_s U_\alpha - \nu\partial_{a_m}(g_{mn}\partial_{a_n}U_\alpha) = \nu B^k_{\alpha\beta}\partial_{a_k}U_\beta + \nu C_{\alpha\beta}U_\beta, \tag{6.27}$$

and assume, on the relevant label chart,

$$\lambda|\xi|^2 \le g_{mn}\xi_m\xi_n \le \Lambda|\xi|^2, \tag{6.28}$$

$$L\|D_a g\|_\infty + L\|B\|_\infty + L^2\|D_a B\|_\infty + L^2\|C\|_\infty \le M_1. \tag{6.29}$$

Suppose

$$\operatorname{supp} U(\cdot,0) \subset E_R, \qquad \operatorname{diam} E_R \le C_0 R,$$

and let A_L be a fixed label annulus satisfying

$$\operatorname{dist}(A_L, E_R) \ge d_0 L, \qquad 0 < R \le c_0 L. \tag{6.30}$$

Then there are constants $c, C, \varepsilon_0 > 0$, depending only on $\lambda, \Lambda, M_1, C_0, d_0, c_0$, such that whenever

$$0 < s \le \tau, \qquad \frac{\nu s}{L^2} \le \varepsilon_0,$$

one has

$$\sup_{a\in A_L}|U(a,s)| \le C(\nu s)^{-3/4}\exp\left(-\frac{cL^2}{\nu s}\right)\|U(\cdot,0)\|_{L^2(da)}, \tag{6.31}$$

$$\sup_{a\in A_L}|D_a U(a,s)| \le C(\nu s)^{-5/4}\exp\left(-\frac{cL^2}{\nu s}\right)\|U(\cdot,0)\|_{L^2(da)}. \tag{6.32}$$

The same statement holds on a lifted periodic chart whose diameter is below the axial injectivity radius.

Proof. Rescale

$$y = \frac{a - a_*}{L}, \qquad \sigma = \frac{\nu s}{L^2}.$$

Then (6.27) becomes

$$\partial_\sigma U_\alpha - \partial_i(g_{ij}\partial_j U_\alpha) = \mathcal{B}^k_{\alpha\beta}\partial_k U_\beta + \mathcal{C}_{\alpha\beta}U_\beta, \tag{6.33}$$

where

$$\mathcal{B} := LB, \qquad \mathcal{C} := L^2 C,$$

and, on the fixed rescaled chart,

$$\lambda|\xi|^2 \le g_{ij}\xi_i\xi_j \le \Lambda|\xi|^2, \qquad \|Dg\|_\infty + \|\mathcal{B}\|_\infty + \|D\mathcal{B}\|_\infty + \|\mathcal{C}\|_\infty \le C.$$

We first prove the off-diagonal estimate at the L^2 level. Choose a bounded Lipschitz function ϕ such that

$$|\nabla\phi| \le 1, \qquad \phi = 0 \quad \text{on the rescaled initial support}, \qquad \phi \ge d_0/2$$

on one fixed enlargement of the rescaled observation annulus. For $\lambda_* > 0$, multiply (6.33) by $e^{2\lambda_*\phi}U$ and integrate in dy. Uniform ellipticity, Cauchy–Schwarz and Young's inequality give

$$\frac{d}{d\sigma}\|e^{\lambda_*\phi}U(\sigma)\|^2_{L^2(dy)} + c\|e^{\lambda_*\phi}\nabla U(\sigma)\|^2_{L^2(dy)} \le C(1+\lambda_*^2)\|e^{\lambda_*\phi}U(\sigma)\|^2_{L^2(dy)}. \tag{6.34}$$

For example,

$$\left|\int e^{2\lambda_*\phi}U_\alpha \mathcal{B}^k_{\alpha\beta}\partial_k U_\beta\, dy\right| \le \varepsilon\int e^{2\lambda_*\phi}|\nabla U|^2\, dy + C_\varepsilon\int e^{2\lambda_*\phi}|U|^2\, dy.$$

The terms generated by differentiating the exponential weight are bounded by the same two quantities, with coefficients $C\lambda_*$ and $C\lambda_*^2$.

Since $\phi = 0$ on the initial support, Gronwall yields

$$\|e^{\lambda_*\phi}U(\sigma)\|_{L^2(dy)} \le \exp\big(C(1+\lambda_*^2)\sigma\big)\|U(\cdot,0)\|_{L^2(dy)}.$$

On the observation region $\phi \ge d_0/2$, hence

$$\|U(\sigma)\|_{L^2(A,dy)} \le C\exp\left(-\frac{\lambda_* d_0}{2} + C\lambda_*^2\sigma + C\sigma\right)\|U(\cdot,0)\|_{L^2(dy)}.$$

Choose

$$\lambda_* = \frac{d_0}{4C\sigma}.$$

For $0 < \sigma \le \varepsilon_0$, with ε_0 fixed sufficiently small,

$$\boxed{\|U(\sigma)\|_{L^2(A,dy)} \le Ce^{-c/\sigma}\|U(\cdot,0)\|_{L^2(dy)}.} \tag{6.35}$$

The same estimate holds, with changed fixed constants, on a slightly larger observation annulus.

To pass to pointwise estimates, fix (y,σ) in the observation annulus and choose a backward parabolic cylinder

$$Q_\rho(y,\sigma) := B_\rho(y)\times(\sigma-\rho^2,\sigma), \qquad \rho = c_0\sqrt{\sigma},$$

with $c_0 > 0$ fixed sufficiently small. This cylinder stays a fixed positive distance from the initial support, and its time coordinate is comparable to σ. The classical interior estimates for parabolic systems of Friedman [15], followed by local Sobolev embedding, give

$$|U(y,\sigma)| \le C\rho^{-5/2}\|U\|_{L^2(Q_\rho(y,\sigma),dy\,d\sigma)}, \tag{6.36}$$

$$|\nabla U(y,\sigma)| \le C\rho^{-7/2}\|U\|_{L^2(Q_\rho(y,\sigma),dy\,d\sigma)}. \tag{6.37}$$

For every time in this cylinder, the weighted-energy argument leading to (6.35) applies on the same enlarged annulus, so

$$\|U(t)\|_{L^2(B_\rho(y),dy)} \le Ce^{-c/\sigma}\|U(\cdot,0)\|_{L^2(dy)}.$$

Because the time length of Q_ρ is ρ^2,

$$\|U\|_{L^2(Q_\rho,dy\,d\sigma)} \le C\rho e^{-c/\sigma}\|U(\cdot,0)\|_{L^2(dy)}.$$

Insert this into (6.36)–(6.37) and use $\rho \asymp \sqrt{\sigma}$:

$$|U(y,\sigma)| \le C\sigma^{-3/4}e^{-c/\sigma}\|U(\cdot,0)\|_{L^2(dy)},$$

$$|\nabla U(y,\sigma)| \le C\sigma^{-5/4}e^{-c/\sigma}\|U(\cdot,0)\|_{L^2(dy)}.$$

Undoing the scaling gives (6.31)–(6.32).

For the periodic branch, work on a lifted chart below the injectivity radius and extend the coefficients smoothly across one fixed larger Euclidean chart. Ellipticity and all dimensionless coefficient bounds are preserved, so the same argument applies. □

Definition 6.7 (Label-space boundary capacity)**.** *Let $\mathcal{A}_0$ be the fixed terminal-label transition annulus used by the material cutoff. Define*

$$\boxed{\mathcal{C}_\Xi(L,J) := \sup_{\substack{a\in\mathcal{A}_0\\ t\in J}} \left(L|\Xi(a,t)| + L^2|D_a\Xi(a,t)|\right).} \tag{6.38}$$

Proposition 6.8 (Material-label no-teleportation)**.** *Assume the terminal detector is supported in a label set of diameter $O(R)$, the transition annulus $\mathcal{A}_0$ satisfies*

$$\operatorname{dist}\left(\mathcal{A}_0, \operatorname{supp}\psi_R\right) \ge c_0 L, \qquad R = o(L),$$

and (6.24) holds. Put

$$\varepsilon := \frac{\nu\tau}{L^2}.$$

Then

$$\boxed{\mathcal{C}_\Xi(L,J) \le C_M\left(\frac{R}{L}\right)^{1/2}\left(\varepsilon^{-3/4}+\varepsilon^{-5/4}\right)e^{-c_M/\varepsilon}.} \tag{6.39}$$

Consequently

$$\frac{R}{L}\longrightarrow 0, \qquad \frac{\nu\tau}{L^2}\longrightarrow 0 \quad\Longrightarrow\quad \mathcal{C}_\Xi(L,J)\longrightarrow 0.$$

Proof. Apply theorem 6.5 and theorem 6.6 to $U = \widehat{\Xi}$ with initial datum $U(\cdot,0) = \psi_R$. By (6.4),

$$\|\psi_R\|_{L^2(da)} \le CR^{1/2}.$$

Hence, for $0 < s \le \tau$,

$$L|\widehat{\Xi}| \le C\left(\frac{R}{L}\right)^{1/2}\left(\frac{\nu s}{L^2}\right)^{-3/4}e^{-cL^2/(\nu s)}$$

and

$$L^2|D_a\widehat{\Xi}| \le C\left(\frac{R}{L}\right)^{1/2}\left(\frac{\nu s}{L^2}\right)^{-5/4} e^{-cL^2/(\nu s)}$$

on $\mathcal{A}_0$. For sufficiently small ε the functions $\theta^{-3/4}e^{-c/\theta}$ and $\theta^{-5/4}e^{-c/\theta}$ are increasing on $0<\theta\le\varepsilon$, so the supremum occurs, up to a fixed harmless constant, at $s=\tau$. This gives (6.39); the exponential dominates both polynomial factors as $\varepsilon\downarrow 0$. □

6.5. **Material geometry is a router, not an extra hypothesis.**

Proposition 6.9 (Finite-degree routing of material-geometry failure)**.** *On a handoff with $\tau/L^2\to 0$, failure of*

$$\|F\| + \|F^{-1}\| + L\|D_aF\| + L^2\|D_a^2F\| \le M$$

forces a strict physical successor of degree 2, 3, *or* 4. *Conversely, if the parent-normalized quantities*

$$L^2\|\nabla u\|_\infty, \qquad L^3\|D^2u\|_\infty, \qquad L^4\|D^3u\|_\infty$$

remain bounded, then

$$F = I + o(1), \qquad F^{-1} = I + o(1), \qquad LD_aF = o(1), \qquad L^2D_a^2F = o(1).$$

Proof. Use

$$\partial_t F = (\nabla u)(X,t)F$$

and differentiate once and twice in a. Gronwall gives the first two bounds; the equations for D_aF and D_a^2F are forced by D^2u and by D^3u together with lower-order commutators. Since $\tau/L^2\to 0$, any fixed failure produces an $o(L)$ homogeneous radius in degree 2, 3, or 4. □

6.6. **Primal–adjoint polarization and the boundary Wronskian current.** Define

$$\mathcal{P}_\chi(t) := \int \chi(X,t)\,\omega(X,t)\cdot\eta(X,t)\,dX, \qquad D_t\chi = 0.$$

The vorticity stretching terms cancel between the primal equation and (6.8); the interior diffusion terms combine into a divergence.

Lemma 6.10 (Material-tube primal–adjoint current)**.** *One has*

$$\boxed{\frac{d}{dt}\mathcal{P}_\chi(t) = -\nu\int \nabla\chi\cdot\sum_{j=1}^{3}\big(\eta_j\nabla\omega_j - \omega_j\nabla\eta_j\big)\,dX.} \tag{6.40}$$

At $\nu=0$ every material subtube carries constant primal–adjoint charge.

Proof. Differentiate $\mathcal{P}_\chi$, use $D_t\chi=0$, the vorticity equation and (6.8), and integrate the Laplacians by parts once. □

Definition 6.11 (Physical adjoint boundary capacity)**.** *Let χ be the material cutoff in (6.13), with transition region* $\operatorname{supp}\nabla\chi$*. Define*

$$\boxed{\mathcal{C}_\eta(L,J) := \sup_{\substack{X\in\operatorname{supp}\nabla\chi(t)\\ t\in J}} \left(L|\eta(X,t)| + L^2|\nabla\eta(X,t)|\right).} \tag{6.41}$$

Lemma 6.12 (Material-label capacity transfer)**.** *Assume the controlled material geometry (6.24). If the material transition annulus is the image of the terminal-label annulus $\mathcal{A}_0$, then*

$$\boxed{\mathcal{C}_\eta(L,J) \le C_M \mathcal{C}_\Xi(L,J).} \tag{6.42}$$

Consequently, under the hypotheses of theorem 6.8,

$$\frac{R}{L}\to 0, \qquad \frac{\nu\tau}{L^2}\to 0 \quad\Longrightarrow\quad \mathcal{C}_\eta(L,J)\to 0.$$

Proof. Along the material map,

$$\eta(X(a,t),t) = F(a,t)^{-T}\Xi(a,t).$$

Hence

$$L|\eta| \le C_M L|\Xi|.$$

Moreover

$$\nabla_x\eta = D_a\left(F^{-T}\Xi\right)F^{-1}.$$

The controlled bounds on F, F^{-1} and LD_aF therefore give

$$L^2|\nabla_x\eta| \le C_M\left(L|\Xi| + L^2|D_a\Xi|\right).$$

Take the supremum on the corresponding transition annuli and use theorem 6.8. □

Proposition 6.13 (Wronskian current routes to material failure or a strict physical successor)**.** *Assume a material handoff of scale L and duration τ with*

$$|\nabla\chi| \le C_\chi L^{-1}, \qquad |\operatorname{supp}\nabla\chi(t)| \le C_\chi L^3$$

in physical three-dimensional volume. Suppose

$$|\mathcal{P}_\chi(t_+) - \mathcal{P}_\chi(t_-)| \ge \delta_0 > 0. \tag{6.43}$$

Then, after subselection, either the material-geometry bound fails and theorem 6.9 gives a strict degree-2, 3, or 4 successor, or the controlled-geometry branch satisfies

$$\boxed{\delta_0 \le C\nu\tau\,\mathcal{C}_\eta(L,J)\left(\|\omega\|_{L^\infty(\mathcal{T})} + L\|\nabla\omega\|_{L^\infty(\mathcal{T})}\right),} \tag{6.44}$$

where $\mathcal{T}$ is the material transition tube. If $\mathcal{C}_\eta(L,J)\le M_\eta$, then at least one of

$$\|\omega\|_{L^\infty(\mathcal{T})} \ge \frac{c\delta_0}{\nu\tau M_\eta}, \qquad \|\nabla\omega\|_{L^\infty(\mathcal{T})} \ge \frac{c\delta_0}{\nu\tau L M_\eta} \tag{6.45}$$

holds. With

$$a_{\omega,2} := \|\omega\|_{L^\infty(\mathcal{T})}^{-1/2}, \qquad a_{\omega,3} := \|\nabla\omega\|_{L^\infty(\mathcal{T})}^{-1/3},$$

this gives

$$\boxed{\frac{a_{\omega,2}}{L} \le C\left(\frac{\nu\tau}{L^2}\right)^{1/2}, \qquad \text{or} \qquad \frac{a_{\omega,3}}{L} \le C\left(\frac{\nu\tau}{L^2}\right)^{1/3}.} \tag{6.46}$$

Thus a fixed Wronskian charge loss on a deep handoff is a strict physical successor.

Proof. Integrate (6.40) over J. On the transition tube,

$$|\eta| \le \frac{\mathcal{C}_\eta}{L}, \qquad |\nabla\eta| \le \frac{\mathcal{C}_\eta}{L^2}.$$

Therefore

$$\begin{aligned}\delta_0 &\le C\nu \int_J \int_{\operatorname{supp}\nabla\chi} \frac{1}{L}\left(\frac{\mathcal{C}_\eta}{L}|\nabla\omega| + \frac{\mathcal{C}_\eta}{L^2}|\omega|\right) dX\, dt \\ &\le C\nu\tau\,\mathcal{C}_\eta\left(L\|\nabla\omega\|_{L^\infty(\mathcal{T})} + \|\omega\|_{L^\infty(\mathcal{T})}\right),\end{aligned}$$

which is (6.44). If the material geometry is controlled, theorems 6.8 and 6.12 in fact make $\mathcal{C}_\eta = o(1)$; for the displayed routing it is enough to retain a fixed upper bound M_η. Splitting the two terms on the right of (6.44) gives (6.45), and the homogeneous-radius calculation gives (6.46). □

6.7. **Charged persistence and a fixed Leray payment.**

Lemma 6.14 (Charged gradient persistence or a higher successor)**.** *Let $L > 0$ be a current parent scale. Suppose one detector-charged unit window contains a regular spacetime point (X_0, t_0) in a fixed CL chart with*

$$L^2|\nabla u(X_0, t_0)| \ge c_0 K, \qquad K \ge 1. \tag{6.47}$$

After passage to the first active scale, at least one of the following holds:

(i)
$$L^3\|D^2u\|_{L^\infty} + L^4\|\partial_t\nabla u\|_{L^\infty} \ge c_1 K \tag{6.48}$$
on the fixed enlargement, and hence a strict degree-three or degree-four successor is selected;

(ii) there is a one-sided parabolic subcylinder $\mathcal{P}$ of spatial radius cL and duration cL^2 on which
$$|\nabla u| \ge cKL^{-2}, \tag{6.49}$$
and therefore
$$\boxed{\int_{\mathcal{P}} |\nabla u|^2\, dX\, dt \ge cK^2L.} \tag{6.50}$$

Proof. If (6.48) holds, the inverse homogeneous radii of D^2u and $\partial_t\nabla u$ are at most

$$CLK^{-1/3}, \qquad CLK^{-1/4},$$

respectively; either is $o(L)$ along an overdriven sequence and is routed as a strict degree-three/four successor.

Otherwise both normalized derivatives in (6.48) are bounded by a sufficiently small fixed fraction of K. The spatial mean-value theorem preserves a fixed fraction of the peak in (6.47) on a ball of radius cL. The temporal fundamental theorem of calculus preserves that fraction for a one-sided time interval of length cL^2. This gives (6.49). Multiplying the squared amplitude K^2L^{-4} by spatial volume cL^3 and time length cL^2 gives (6.50). □

The charged strain wall (6.12) can be overdriven relative to a current parent scale L. Given $K \geq 1$, choose the child sufficiently deep so that one charged unit window satisfies (6.47). Apply theorem 6.14. On its persistent branch choose a fixed reference length $L_\circ$ and set

$$K := \left(\frac{L_\circ}{L}\right)^{1/2}.$$

Then

$$\boxed{\int_{\mathcal{P}} |\nabla u|^2 \, dX \, dt \geq cL_\circ.} \tag{6.51}$$

6.8. **Active-order firewall: bounded jets, spatial factorial growth, and temporal spectators.** A derivative which merely happens to be large is not promoted. A successor is *active* only when it is needed to destroy persistence of the detector-charged signal. We now show that active order growth itself cannot be an independent no-payment escape.

Fix a charged parent scale L and use an axis-centered moving/Galilean chart

$$U(y,\tau) := L\left(u(X(\tau) + Ly, t_* + L^2\tau) - c(\tau)e_z\right), \tag{6.52}$$

where the axial gauge is chosen so that $U(0,\tau) = 0$. For a velocity jet put

$$J_{\alpha,b} := \partial_y^\alpha \partial_\tau^b U, \qquad d(\alpha,b) := 1 + |\alpha| + 2b. \tag{6.53}$$

A spatial derivative raises parabolic degree by one and a time derivative by two.

Lemma 6.15 (Finite-jet compactness returns an active bounded-degree chain to the gradient)**.** *Fix $D \geq 4$. Let U_j be normalized charts of the form (6.52) on a fixed cylinder Q_2, and suppose*

$$A_j := \max_{1 \leq |\alpha| + 2b \leq D-1} \|J_{\alpha,b}^{(j)}\|_{L^\infty(Q_2)} \longrightarrow \infty, \tag{6.54}$$

while the one- and two-step finite successors satisfy

$$\max_{|\alpha|+2b \leq D+1} \|J_{\alpha,b}^{(j)}\|_{L^\infty(Q_2)} \leq C_D A_j. \tag{6.55}$$

Then, on a fixed smaller cylinder,

$$\boxed{\|\nabla_y U_j\|_{L^\infty} \geq c_D A_j,} \tag{6.56}$$

and a fixed fraction of this gradient persists on a parabolic cylinder of normalized radius and duration bounded below by positive D-dependent constants. In physical variables,

$$\boxed{\int_{\mathcal{P}_j} |\nabla u|^2 \, dX \, dt \geq c_D A_j^2 L_j.} \tag{6.57}$$

Proof. Divide U_j by A_j. The gauge $U_j(0,\tau) = 0$ and a fixed-domain Poincaré estimate control the zeroth-order field by the first spatial jet. The successor bound (6.55) gives uniform C^D bounds on every fixed smaller cylinder.

If (6.56) failed, a subsequence would converge in C^{D-1} to a field U_∞ with $\nabla_y U_\infty \equiv 0$. The gauge forces $U_\infty \equiv 0$ at every normalized time, and therefore every mixed derivative of parabolic order at most $D-1$ vanishes. This contradicts (6.54), because after subselection one derivative type realizes a fixed fraction of A_j.

The bounds on $D_y^2 U_j$ and $\partial_\tau \nabla_y U_j$ in (6.55) preserve a fixed fraction of the gradient peak on a fixed parabolic subcylinder. Finally

$$dX \, dt = L_j^5 \, dy \, d\tau, \qquad |\nabla_x u|^2 = L_j^{-4} |\nabla_y U_j|^2,$$

which gives (6.57). □

Proposition 6.16 (Bounded active degree pays the fixed Leray quantum)**.** *Suppose a chronological no-payment restart sequence has charged parent scales $L_j \downarrow 0$ and charged gradient scores*

$$A_j^{\mathrm{ch}} := L_j^2 |\nabla u(X_j, t_j)| \geq c \left(\frac{L_\circ}{L_j} \right)^{1/2}.$$

If the active successor protocol never requires parabolic degree larger than one fixed D, then the sequence contains a packet satisfying

$$\boxed{\int_{\mathcal{P}_j} |\nabla u|^2 \, dX \, dt \geq c_D L_\circ.}$$

Thus an infinite no-payment active chain must have active parabolic degree tending to infinity.

Proof. Route any still smaller fixed-degree first exit first. Since only finitely many derivative types of degree at most $D+1$ are involved, choose the last type which occurs infinitely often and pass to a subsequence. On the complementary first-active branch, all one- and two-step successors needed to destroy persistence are bounded by a D-dependent multiple of the finite-jet maximum; unrelated larger jets are spectators unless they enter such a persistence-failure step. Hence theorem 6.15 applies with $A_j \geq c A_j^{\mathrm{ch}}$. It gives

$$\int_{\mathcal{P}_j} |\nabla u|^2 \, dX \, dt \geq c_D (A_j^{\mathrm{ch}})^2 L_j \geq c_D L_\circ.$$

Chronological reselection makes these packets pairwise disjoint in time, so finite Leray dissipation excludes infinitely many of them. □

Raw homogeneous radii are not accepted as physical restart scales when the derivative order tends to infinity. For

$$H_{\alpha,b} := \partial_x^\alpha \partial_t^b u, \qquad d := 1 + |\alpha| + 2b, \qquad n := |\alpha| + b,$$

define the conservative factorial-corrected radius

$$\boxed{a^{\mathrm{fac}}_{\alpha,b} := \left(\frac{n!}{|H_{\alpha,b}|} \right)^{1/d}}. \tag{6.58}$$

For every fixed derivative type this differs from the ordinary homogeneous radius by only a fixed constant, whereas for order-diverging jets it removes ordinary factorial growth.

Lemma 6.17 (Pressure-free spatial factorial majorant)**.** *Let U_j be first-exit normalized charts on Q_2 for which every fixed lower parabolic degree is bounded. In particular assume*

$$\|U_j\|_{L^\infty} + \|\nabla U_j\|_{L^\infty} + \|D^2 U_j\|_{L^\infty} + \|\partial_\tau U_j\|_{L^\infty} \le M_0 \tag{6.59}$$

on Q_2, with M_0 independent of j. Then there are $A_s, B_s, \rho_s > 0$, independent of j and of the spatial order, such that

$$\boxed{\|\partial_y^\alpha U_j\|_{L^\infty(Q_1)} \le A_s B_s^{|\alpha|} |\alpha|!\, \rho_s^{-|\alpha|}.} \tag{6.60}$$

Consequently every purely spatial factorial-corrected radius satisfies

$$\boxed{\frac{a^{\mathrm{fac}}_{\alpha,0}}{L_j} \ge c_s > 0} \tag{6.61}$$

for all sufficiently large $|\alpha|$.

Proof. Put $\Omega_j := \nabla \times U_j$. Curl removes pressure and the spatially constant Galilean acceleration:

$$\partial_\tau \Omega_j - \nu \Delta \Omega_j + (U_j \cdot \nabla)\Omega_j - (\Omega_j \cdot \nabla) U_j = 0. \tag{6.62}$$

At each time,

$$\nabla \cdot U_j = 0, \qquad -\Delta U_j = \nabla \times \Omega_j. \tag{6.63}$$

For $1 < r < 2$ and $0 < \sigma < 1$, define the distance-weighted quantities

$$\mathfrak{W}_m := \sup_{1<r<2} \frac{(2-r)^m}{m!} \|\nabla^m \Omega_j\|_{C^\sigma(Q_r)},$$

$$\mathfrak{U}_m := \sup_{1<r<2} \frac{(2-r)^m}{m!} \|\nabla^{m+1} U_j\|_{C^\sigma(Q_r)}.$$

Differentiate (6.62) m times in space and apply the constant-coefficient interior parabolic estimate on a δ-smaller cylinder. The div–curl relation (6.63) gains one spatial derivative; its harmonic remainder obeys the usual interior Cauchy bound $C^{m+1} m! \delta^{-(m+1)}$.

Choose

$$\delta_m := \frac{2-r}{m+2}.$$

Multiplication by the factorial distance weights cancels the δ_m^{-2} parabolic loss. Leibniz' rule for the transport/stretching terms gives, after a harmless index shift, a scalar positive recurrence

$$Z_{m+2} \le C_0 Z_m + C_0 \sum_{k=0}^{m+1} Z_k Z_{m+1-k} + C_0^{m+2}. \tag{6.64}$$

The generating function $\mathcal{Z}(z) = \sum_{m\ge 0} Z_m z^m$ is coefficientwise dominated near $z = 0$ by the analytic branch of

$$\mathcal{W} = Z_0 + Z_1 z + C_0 z^2 \mathcal{W} + C_0 z \mathcal{W}^2 + \frac{C_0^2 z^2}{1 - C_0 z}.$$

Hence $Z_m \le A_s B_s^m$. This proves (6.60). Since

$$L_j^{1+|\alpha|} |\partial_x^\alpha u| = |\partial_y^\alpha U_j|,$$

substitution into (6.58) with $b = 0$ gives (6.61). □

Corollary 6.18 (Residual order divergence must be temporal)**.** *On an order-diverging first-exit sequence, if the number of time derivatives b stays bounded, the factorial-corrected radius cannot be $o(L_j)$. Consequently any surviving active order-divergent restart chain must have*

$$\boxed{b_j \longrightarrow \infty.}$$

Proof. For each fixed upper bound B on b, differentiate (6.62) only the finitely many times $0 \le b \le B$. The first-exit convention bounds all lower temporal jets, and the proof of theorem 6.17 repeats with constants depending on B but not on $|\alpha|$. Thus $a_{\alpha,b}^{\mathrm{fac}}/L_j$ has a positive lower bound for $b \le B$ and $|\alpha| \to \infty$. □

The only remaining order-divergent possibility is therefore rapid temporal variation. Here the Euler firewall gives a stronger simplification than a mixed space–time analyticity theorem.

Lemma 6.19 (Curl-free harmonic time-frequency noise is a spectator)**.** *Let a local meridional Hodge split on a handoff tube be*

$$u = v + h, \qquad \nabla \times h = 0, \qquad \nabla \cdot h = 0.$$

Choose the moving/Galilean gauge so that $h(X(t), t) = 0$ at the tube center. Assume the residual no-degree-two-target bound

$$L^2 \|\nabla h\|_{L^\infty(\mathrm{tube})} \le M_h. \tag{6.65}$$

Then the vorticity equation contains h only through its instantaneous values h and ∇h:

$$\partial_t \omega - \nu \Delta \omega + (v \cdot \nabla)\omega - (\omega \cdot \nabla) v + (h \cdot \nabla)\omega - (\omega \cdot \nabla) h = 0. \tag{6.66}$$

No time derivative $\partial_t^k h$, $k \geq 1$, appears. Moreover, on a deep handoff $\tau/L^2 \to 0$,

$$\int_J \|\nabla h(t)\|_{L^\infty}\, dt \leq M_h \frac{\tau}{L^2} \longrightarrow 0. \tag{6.67}$$

Thus arbitrarily large high time derivatives of a curl-free harmonic carrier cannot be an active successor of a fixed vorticity-based detector.

Proof. Because $\nabla \times h = 0$, the vorticity of u equals that of v. Substitute $u = v + h$ into the Cartesian vorticity equation

$$D_t\omega = (\omega \cdot \nabla)u + \nu\Delta\omega$$

to obtain (6.66); no time derivative of h is created. The bound (6.67) follows directly from (6.65). Since $h(X(t), t) = 0$, the mean-value theorem also gives

$$\|h(t)\|_{L^\infty(B_{CL}(X(t)))} \leq CL\|\nabla h(t)\|_{L^\infty},$$

so the integrated affine/harmonic transport is controlled by the same vanishing action. Hence high time-frequency information which is invisible to curl changes the detector by $o(1)$ and is spectator data. □

Proposition 6.20 (Active-order closure on a deep handoff)**.** *Consider a detector-charged restart sequence on which*

$$\frac{\nu\tau_j}{L_j^2} \longrightarrow 0 \tag{6.68}$$

and the material-geometry bounds required in theorem 6.3 hold. Then active order growth cannot furnish an independent infinite no-payment escape. More precisely, after subselection, one of the following occurs:

(i) a fixed Leray packet (6.51);
(ii) a strict finite-degree physical successor;
(iii) the charged Cauchy token persists and no temporal successor is active.

Proof. If the active parabolic degree is bounded, use theorem 6.16. Otherwise the active degree tends to infinity. Purely spatial order growth cannot create an order-uniform $o(L_j)$ restart scale by theorems 6.17 and 6.18; hence a residual active chain must have unbounded time order.

Split the local velocity into its vortical part and curl-free harmonic remainder. If $L_j^2\|\nabla h\|_{L^\infty} \to \infty$, this is already a degree-two physical successor. On the complementary branch theorem 6.19 shows that arbitrarily large high time jets of h are spectators.

The surviving active temporal variation is therefore genuinely vortical. By the active-successor convention it is promoted only when a fixed fraction of the detector-charged physical token is lost over a real handoff interval. Apply the Cauchy polarization theorems 6.2 and 6.3. If no fixed loss occurs, the Cauchy token persists and no temporal successor is active. If a fixed loss occurs, (6.7) and (6.68) give

$$\frac{a_{\omega,4}}{L_j} \longrightarrow 0,$$

which is a strict degree-four spatial successor. Thus raw high time order cannot remain as an independent no-payment branch. □

The active-order argument therefore closes without any mixed space–time analyticity theorem:

$$\boxed{\begin{array}{c}\text{bounded active degree} \Longrightarrow \text{fixed Leray payment},\\ \text{pure spatial order divergence} \Longrightarrow \text{no order-uniform smaller radius},\\ \text{curl-free high-time noise} \Longrightarrow \text{spectator},\\ \text{genuinely vortical finite-time loss} \Longrightarrow \nu\Delta\omega \Longrightarrow \text{degree-four successor}.\end{array}} \tag{6.69}$$

6.9. **Automatic deep handoff and chronological reselection.** The deep-handoff ratio used by the polarizers is selected, not assumed. There are only two active origins after Bridge D: a fresh compact source-born child and a retained fixed-mass relay. We treat them separately.

Lemma 6.21 (Fresh compact child is deep relative to a strict parent)**.** *Let L_j be a parent scale, let R_j be the physical child/detector scale, and let S_j be a creation-window scale satisfying*

$$R_j \le C_R S_j, \qquad \frac{S_j}{L_j} \longrightarrow 0. \tag{6.70}$$

Suppose the first positive genealogy is created inside a terminal interval J_j satisfying

$$\tau_j := |J_j| \le C_t S_j^2. \tag{6.71}$$

Then

$$\boxed{\frac{\tau_j}{L_j^2} \le C_t \left(\frac{S_j}{L_j}\right)^2 \longrightarrow 0.} \tag{6.72}$$

Proof. Immediate from (6.70) and (6.71). In whole-space C.1 one may take $S_j \asymp R_j$. In the periodic compact selector branch one again has $S_j \asymp R_j$, whereas on the collapsing-first-moment branch $R_j = R_j^\sharp = o(S_j)$ and the selected event is known to lie in the terminal $O(S_j^2)$ selector window. Both cases give the same deep-handoff ratio relative to the parent scale L_j. □

For the retained origin, fix one parent corridor $\mathcal{P}_{\mathrm{par}}$ with the flat action budget

$$\boxed{\int_{\mathcal{P}_{\mathrm{par}}} q(r,z,t)^2 \, dr\, dz\, dt \le A_*.} \tag{6.73}$$

The first-active-scale assignment in theorem 5.10 provides a bounded-overlap family of compact assigned cells. We record the elementary packing estimate behind that assignment.

Lemma 6.22 (Positive-band Bessel packing for assigned cells)**.** *Let* $\{(\ell_k, J_k, \psi_k)\}_{k\in\mathcal{K}}$ *be a family inside* $\mathcal{P}_{\mathrm{par}}$ *such that*

$$0 \le \psi_k \le 1, \qquad |\operatorname{supp}\psi_k(t)|_{dr\,dz} \le C_0\ell_k^2,$$

and at each regular time the supports $\operatorname{supp}\psi_k(t)$ *have overlap at most* N_0*. For one fixed sign put* $p = (sq)_+$ *and define the flat positive-band mass*

$$w_k(t) := \int_{\mathcal{Q}_*} \psi_k(r,z,t)p(r,z,t)\,dr\,dz. \tag{6.74}$$

Then at every regular time

$$\boxed{\sum_{k\in\mathcal{K}} \frac{w_k(t)^2}{\ell_k^2} \le C\int_{K_{\mathrm{par}}(t)} q(r,z,t)^2\,dr\,dz,} \tag{6.75}$$

and consequently

$$\boxed{\sum_{k\in\mathcal{K}} \int_{J_k} \frac{w_k(t)^2}{\ell_k^2}\,dt \le CA_*.} \tag{6.76}$$

No disjointness of the time intervals J_k *is assumed.*

Proof. Cauchy–Schwarz in the explicitly flat measure gives

$$w_k(t)^2 \le \left(\int_{\operatorname{supp}\psi_k(t)} 1\,dr\,dz\right)\left(\int_{\operatorname{supp}\psi_k(t)} p(r,z,t)^2\,dr\,dz\right).$$

Since $p^2 \le q^2$,

$$\frac{w_k(t)^2}{\ell_k^2} \le C_0\int_{\operatorname{supp}\psi_k(t)} q(r,z,t)^2\,dr\,dz.$$

Sum in k and use the overlap bound N_0 to obtain (6.75). Integrate in time and use (6.73). Because the estimate is summed at each time before integration, the intervals J_k need not be disjoint. □

Corollary 6.23 (Only finitely many natural-time coherent retained bands fit in one parent)**.** *Fix* $\theta, c_* > 0$*. Suppose that for* $k \in \mathcal{K}_{\mathrm{coh}}$*,*

$$w_k(t) \ge \theta \quad (t \in J_k), \qquad |J_k| \ge c_*\ell_k^2. \tag{6.77}$$

Then

$$\boxed{\#\mathcal{K}_{\mathrm{coh}} \le \frac{CA_*}{c_*\theta^2}.} \tag{6.78}$$

Proof. Every $k \in \mathcal{K}_{\mathrm{coh}}$ contributes at least $c_*\theta^2$ to the left side of (6.76). □

Proposition 6.24 (An arbitrarily deep retained active relay has a subparabolic subsequence)**.** *Consider the retained fixed-mass alternative of theorem 5.10 inside one parent corridor satisfying* (6.73)*. Let* $\ell_j \downarrow 0$ *be its*

first-active assigned scales. For each assigned packet choose a cutoff ψ_j equal to one on the compact packet core and put

$$w_j(t) := \int \psi_j(r,z,t)(s_j q(r,z,t))_+ \, dr\, dz.$$

After sign subselection there are fixed constants $\mu_0 > 0$ and $\theta := \mu_0/4$ and active times t_j for which

$$w_j(t_j) \geq \frac{\mu_0}{2}.$$

Let J_j be the maximal connected interval containing t_j on which

$$w_j(t) \geq \theta, \tag{6.79}$$

and set $\tau_j := |J_j|$. Then an arbitrarily deep retained relay has a subsequence such that

$$\boxed{\frac{\tau_j}{\ell_j^2} \longrightarrow 0.} \tag{6.80}$$

Proof. Continuity at regular times makes J_j nonempty. Suppose (6.80) were false. Then after subselection there would be a fixed $c_* > 0$ and infinitely many assigned packets with

$$|J_j| \geq c_* \ell_j^2.$$

Together with (6.79), these packets satisfy (6.77). The finite bound (6.78) is a contradiction.

Equivalently, for each integer $m \geq 1$, only finitely many assigned retained packets can satisfy

$$\tau_j \geq \frac{1}{m}\ell_j^2.$$

Choose successively deeper indices outside those finite exceptional sets. The resulting diagonal subsequence obeys $\tau_j/\ell_j^2 \leq 1/j$, proving (6.80). □

Remark 6.25 (Why the fixed-mass threshold is part of the active-scale assignment). The retained branch of theorem 5.10 is selected only after vanishing-mass scale-assigned packets have been declared Hodge-silent. Hence every retained packet carries one fixed positive signed flat mass at its assigned cell. Choosing $\psi_j = 1$ on that compact cell core gives the uniform lower bound used in theorem 6.24; no new mass hypothesis is introduced here.

Theorem 6.26 (Deep handoff is a routed conclusion)**.** *Suppose a continuing active thread from Bridge D reaches either*

- *(i) a compact fresh source-born child with a selector/creation-window scale $S_j = o(L_j)$ and child scale $R_j \leq CS_j$;*
- *(ii) an arbitrarily deep retained fixed-mass relay inside the standing parent flat-action budget.*

Then, after discarding finitely many natural-time coherent retained episodes and passing to an active subsequence if necessary, the downstream polarizers may be run on handoff intervals satisfying

$$\boxed{\frac{\tau_j}{L_j^2} \longrightarrow 0,} \tag{6.81}$$

where L_j is the assigned parent/handoff scale for the corresponding polarized argument.

Proof. Use theorem 6.21 on the source-born branch and theorem 6.24 on the retained branch. In the latter case set $L_j = \ell_j$. These are exactly the two non-spectator active origins left by Bridge D. □

After a paid packet ends at a regular time $b < T$, use the geometry-appropriate future clock. In the whole space this is the internally derived flat-q^2 clock theorem 3.7; in the periodic cylinder it is the SC.2 recurrent local strain selector theorems 4.24 and 4.27. Each selects a later active block or packet below any prescribed strict target scale, modulo an already routed still smaller target. Hence, whenever the active thread continues, detector-charged parents can be selected after the previously paid packet. The packets in (6.51) may therefore be chosen pairwise disjoint in time.

Proposition 6.27 (A strict-successor-only restart loop cannot persist)**.** *Consider an indefinitely continuing active terminal thread supplied by Bridge D. Suppose every strict physical successor is absorbed by the geometry-appropriate future clock into a later active generation, as in the chronological reselection paragraph above. Then it is impossible, after some late time, for every Bridge-E handoff to exit only through strict physical successors while never producing the fixed Leray payment (6.51). Consequently an indefinitely continuing active thread contains infinitely many fixed-payment packets, which may be chosen pairwise disjoint in time.*

Proof. Assume to the contrary that, after some late generation, the thread continues indefinitely through strict successors but never pays (6.51). Chronological absorption gives later active scales $L_{j+1} \le a_j = o(L_j)$ after subselection, hence $L_j \downarrow 0$. By theorem 6.26, pass to handoffs with

$$\frac{\nu\tau_j}{L_j^2} \longrightarrow 0.$$

The axis-away detector and the adjoint charged-strain identity give the strain wall (6.12). Because the active descendants may be chosen arbitrarily deep, this wall may be overdriven: for

$$K_j := \left(\frac{L_\circ}{L_j}\right)^{1/2} \longrightarrow \infty,$$

choose a sufficiently deep charged descendant containing a point with

$$L_j^2|\nabla u| \ge cK_j.$$

Apply theorem 6.14. Its persistent branch pays

$$\int_{\mathcal{P}_j} |\nabla u|^2 \, dX \, dt \geq cK_j^2 L_j = cL_\circ,$$

contrary to the no-payment assumption. Therefore every sufficiently late overdriven generation must leave through an active successor.

If the parabolic degrees of these active successors are bounded along an infinite subsequence, theorem 6.16 gives the same fixed Leray payment, again a contradiction. Hence a hypothetical unpaid loop would have to use active degrees tending to infinity. Apply theorem 6.20. Its payment alternative is excluded by assumption. Its strict finite-degree successor alternative, if repeated infinitely often, returns to the bounded-degree case just excluded. Its persistent-token alternative returns to the overdriven charged-strain argument above and therefore yields either a payment or a finite-degree successor. Thus no infinite successor-only loop remains.

After each paid packet ends, chronological reselection starts the next active generation strictly later in time. Hence the payments produced by an indefinitely continuing thread may be chosen pairwise time-disjoint. □

Remark 6.28 (The terminal bill is genuinely Navier–Stokes). The successor branch is routing only. The terminal contradiction comes from the fixed positive packet in (6.51) together with the physical Leray ledger

$$\nu \int_0^T \int_{\Omega_*} |\nabla u|^2 \, dX \, dt < \infty.$$

Thus infinitely many pairwise disjoint payments would cost at least $c\nu L_\circ$ each. This is not an Euler-compatible terminal argument. Likewise the Cauchy and Wronskian restart estimates retain explicit factors of ν in (6.5) and (6.40).

Theorem 6.29 (Bridge E: common polarized downstream alternative)**.** *Assume the output of Bridge D and a continuing active terminal thread. After subselection, Bridge E produces one of the following:*

(i) a strict physical successor, of finite degree, which is absorbed into a later canonical block;

(ii) a spacetime packet $\mathcal{P}$ satisfying the fixed Leray payment (6.51)*.*

An infinite sequence of item (i) outcomes with no item (ii) is impossible by theorem 6.27. Hence, if the active thread continues indefinitely, item (ii) occurs infinitely often and its packets may be chosen pairwise disjoint in time. No separate nodal, harmonic-gauge, bare-reset, or adjoint-mixing conditional leaf remains in the common downstream interface.

Proof. First apply theorem 6.26 to the source-born/retained origin supplied by Bridge D; hence every sufficiently deep polarized handoff may be taken with $\tau/L^2 \to 0$. The detector is supplied by theorem 6.1. Apply the Cauchy polarization theorems 6.2 and 6.3; route material geometry by theorem 6.9. On the inherited charged branch, use the full adjoint theorem 6.4, localize

by (6.13)–(6.17), and apply theorem 6.8. The exact primal–adjoint current theorems 6.10 and 6.13 returns any fixed material-boundary loss to a strict physical successor. On the complementary charged branch apply theorem 6.14. The active-order firewall theorem 6.20 eliminates spectator high jets and converts genuine finite-time vortical loss back to theorem 6.3. The no-successor-only-loop conclusion is theorem 6.27; chronological reselection then makes the resulting paid packets pairwise disjoint. □

7. Proof architecture


The synchronized diagrams below summarize the proof architecture. They are not substitutes for the proofs; rather, they make explicit which steps are geometry-specific, where the two geometries merge, and where Euler-compatible routing gives way to the common viscosity-polarized downstream mechanism. The compact symbolic chains beneath the figures record the same information in linear form.

7.1. **Whole-space chain.**

Synchronized proof-flow comparison
Part I: geometry-specific upstream

Whole space	Periodic cylinder
Candidate whole-space first singular time under (H_∞)	Candidate periodic first singular time under (H_L)
SC.1 radial-strain endpoint and weighted-Hodge clock conversion theorems 3.5 to 3.7 $\Longrightarrow$ future flat-q^2 clock	SC.2 radial-strain endpoint theorem 4.24 infinite terminal flat α^2 clock + finite first radial moment
Universal B.1 relay theorem 4.10 center-free restart at a later strict degree-two scale	Nested-arc recurrent-axis selection theorem 4.26 $\Longrightarrow$ recurrent singular axis point z_*
A.1 synchronization on the surviving compact branch theorem 4.13 natural window + positive duration + terminal synchronization	One-sided suitability + local Type-II input theorems 4.4, 4.8 and 4.9 apply at the recurrent point z_*
Center-free finite-corridor adjacent descent theorem 4.14	Local strain selector and geometry-corrected Hodge reselection theorems 4.27 and 4.28 theorems 4.30 and 4.31
C.1 Hardy/source-work localization theorems 4.16 and 4.21 theorems 4.18 and 4.19 arbitrary-center child; center covariance theorem 4.22	Periodic compact child interface theorems 4.36, 4.39 and 4.40 theorem 4.37 recentered signed child + S_j deep window + L_j parent budget

The common downstream engine is displayed on the next page.



Remark 7.1 (Why the synchronized comparison display helps). The side-by-side comparison display is meant to expose the architecture at one glance. Part I isolates the geometry-specific upstream devices: whole-space SC.1/B.1/A.1/finite-corridor/C.1 versus periodic SC.2/recurrent-point/Type-II/selector-Hodge/C.2. Part II shows that from Bridge D onward the two geometries share the same polarized downstream engine. This makes it visually clear which nodes are geometry-specific and which ones are common.

The whole-space chain is now

$$\begin{aligned}
&\mathrm{SC.1} \Longrightarrow \mathrm{C-WC} \Longrightarrow \mathrm{B.1} \Longrightarrow \mathrm{A.1},\\
&\quad \Longrightarrow \text{finite-corridor descent} \Longrightarrow \mathrm{C.1} \Longrightarrow \text{Bridge D},\\
&\quad \Longrightarrow \text{deep handoff} \Longrightarrow \text{Bridge E},\\
&\quad \Longrightarrow \left[\begin{array}{c}\text{strict successor } \curvearrowright \text{ later restart}\\ \text{or fixed Leray payment}\end{array}\right],\\
&\quad \Longrightarrow \text{no successor-only loop} \Longrightarrow \text{funded assembly}.
\end{aligned}$$

The symbol C−WC is retained only as the historical name of the canonical flat-q^2 future clock; it is no longer a hypothesis.

The historical z-reflection specialization is not a node in this active tree; its scope and replacement are recorded in theorems 2.1 and 4.22.

Proof of theorem 1.3. Apply theorems 3.5 to 3.7 to obtain the infinite terminal flat-q^2 clock and arbitrarily late small-radius canonical blocks. The universal relay theorem 4.10 is available before synchronization. On the surviving compact branch apply theorems 4.13 and 4.14 and then the complete C.1 package theorems 4.21 and 4.22 to obtain an arbitrary-center compact signed child and a fixed parent local flat-action budget.

Bridge D, theorem 5.11, separates source-born, retained fixed-mass, strict-restart and spectator origins. The first two active origins satisfy the automatic deep-handoff conclusion theorem 6.26. The common polarized router theorem 6.29 then gives the strict-successor/fixed-payment alternative. Finally theorem 6.27 excludes an infinite successor-only restart chain, and chronological reselection makes the resulting fixed payments pairwise time-disjoint. □

Proposition 7.2 (Whole-space Euler firewall)**.** *The active whole-space chain*

$$\boxed{\text{SC.1} \to \text{C}-\text{WC} \to \text{B.1} \to \text{A.1} \to \text{C.1} \to \text{Bridge D} \to \text{Bridge E}}$$

obeys the following firewall classification.

(i) ***Upstream NS endpoint.*** *SC.1 is viscosity specific: its flat-energy inequality uses positive viscous dissipation and its conclusion uses the physical H^1 restart.*

(ii) ***Euler-compatible clock conversion.*** *The weighted Hodge estimate converting the divergent α clock to the flat-q^2 clock is Euler compatible and is used only to create the future canonical clock.*

(iii) ***Routing and compactness.*** *B.1, A.1 synchronization, the finite-corridor descent, the Hardy/source localization and Bridge D select scales, packets, genealogies or restarts. None of these nodes is accepted as a terminal contradiction merely because it is scale invariant or survives at $\nu = 0$.*

(iv) ***Polarized viscosity-visible successors.*** *In Bridge E, Cauchy polarization removes material stretching and exposes*

$$\partial_t(F^{-1}\omega) = \nu F^{-1}\Delta\omega.$$

Material-label polarization and the Wronskian identity remove material transport and leave the explicit viscous current (6.40). Fixed losses therefore return to strict physical successors with explicit ν-dependent scale bounds.

(v) ***Funded terminal branch.*** *By theorem 6.27, an indefinitely continuing active thread cannot remain in successor routing forever. It produces pairwise time-disjoint packets satisfying*

$$\int_{\mathcal{P}_j} |\nabla u|^2 \, dX \, dt \geq cL_\circ.$$

The terminal contradiction uses only the finite Navier–Stokes ledger

$$\nu \int_0^T \int_{\mathbb{R}^3} |\nabla u|^2 \, dX \, dt < \infty.$$

Thus no Euler-compatible identity is used as the terminal funded closure.

Proof. Items (i)–(ii) are theorems 3.5, 3.6 and 3.8. Item (iii) follows from the statements of theorems 4.10, 4.13, 4.14, 4.21 and 5.11. Item (iv) is theorems 6.2, 6.3, 6.10 and 6.13. Item (v) is theorems 6.14, 6.27 and 6.28. □

7.2. **Periodic-cylinder chain.** The active minimal periodic chain is

$$\begin{aligned}
&\text{SC.2} \Longrightarrow z_* \Longrightarrow \text{Type}-\text{II}, \\
&\quad \Longrightarrow \text{strain/Hodge reselection}, \qquad R_j \le CS_j = o(L_j), \\
&\quad \Longrightarrow \text{B.2/C.2-core} \Longrightarrow \text{Bridge D} \Longrightarrow \text{deep handoff}, \\
&\quad \Longrightarrow \text{Bridge E} \Longrightarrow \left[\begin{array}{c} \text{strict successor} \curvearrowright \text{later restart} \\ \text{or fixed Leray payment} \end{array} \right], \\
&\quad \Longrightarrow \text{no successor-only loop} \Longrightarrow \text{funded assembly}.
\end{aligned}$$

Proof of theorem 1.4. Use theorems 4.24 to 4.26 to select the recurrent singular axis point z_*. The one-sided preterminal suitability/source firewall theorems 4.4 and 4.8 permits the already proved local Type-II theorem theorem 4.9 at that point.

Apply the local selector/Hodge package theorems 4.27 and 4.28 and theorems 4.30 and 4.31, then theorems 4.33, 4.35 and 4.36. The resulting C.2 package theorems 4.37 and 4.39 and theorem 4.40 supplies the recentered compact signed child, its subparabolic selector window, and the fixed parent local flat-action budget.

From Bridge D onward the proof is the common downstream: theorems 5.11, 6.26 and 6.29. The no-successor-only-loop conclusion theorem 6.27 and chronological reselection yield pairwise time-disjoint fixed payments on every indefinitely continuing active thread. □

Proposition 7.3 (Periodic Euler firewall)**.** *The active periodic chain*

$$\begin{aligned}
&\text{SC.2} \to z_* \to \text{Type}-\text{II} \to \textit{strain/Hodge reselection}, \\
&\quad \to \text{C.2-core} \to \text{Bridge D} \to \text{Bridge E}.
\end{aligned}$$

has the following firewall classification.

- *(i)* ***Viscosity-specific upstream endpoint.*** *SC.2 uses the positive viscous flat dissipation in* (4.56) *and the uniform physical-H^1 restart. Thus its continuation endpoint is genuinely Navier–Stokes specific.*
- *(ii)* ***NS-funded recurrent-point routing.*** *The passage from the divergent flat strain clock to z_* uses the finite physical first moment* (4.57)*, which is paid by the Leray dissipation ledger. It selects a recurrent axis point; it is not used as a terminal contradiction.*

(iii) ***External local NS singularity input.*** *Theorem 4.9 is used only after z_* is selected. Its source scope is one-sided and local by theorems 4.3 and 4.4. The conclusion supplies unbounded scale-invariant physical dissipation and hence smaller gradient/strain comparison scales; it is routing, not the terminal bill.*

(iv) ***Local routing.*** *The strain selector, mean/nonzero-mode split, quantitative periodic Hodge detector, signed-child extraction, periodic translation covariance and the C.2 parent budget select compact packets and restart scales. Their role is geometric/elliptic routing. No instantaneous Hodge or compactness identity is promoted to terminal closure merely because it is scale invariant.*

(v) ***Viscous terminal interface.*** *From Bridge D onward the periodic proof uses the same Bridge E as the whole space. Cauchy and Wronskian losses retain explicit factors of ν, and theorem 6.27 forces an indefinitely continuing thread to produce pairwise time-disjoint packets with*

$$\int_{\mathcal{P}_j} |\nabla u|^2 \, dX \, dt \geq cL_\circ.$$

The terminal contradiction is the finite physical Navier–Stokes ledger

$$\nu \int_0^T \int_{\Omega_L} |\nabla u|^2 \, dX \, dt < \infty.$$

Thus the periodic fork introduces no Euler-compatible terminal closure.

Proof. Item (i) is theorems 3.1 and 4.24. Item (ii) is theorems 4.25 and 4.26. Item (iii) is theorems 4.4, 4.8 and 4.9. Item (iv) is supplied first by the selector/Hodge package theorems 4.27 and 4.28 and theorems 4.30 and 4.33, and then by the periodic compact-interface package theorems 4.37 and 4.40 and theorem 4.39. Item (v) is theorems 5.11 and 6.29, together with theorems 6.27 and 6.28. □

8. Conclusion

The argument developed here separates the axisymmetric problem into two geometry-specific upstream constructions and one common downstream mechanism. In the whole space, the internally derived strain endpoint and weighted Hodge clock feed the universal B.1 restart, A.1 synchronization, the center-free finite corridor and the reflection-free C.1 Hardy/source-work localization. In the periodic cylinder, the strain clock instead selects a recurrent axis point, after which the one-sided suitability/Type-II input, the local strain selector and the periodic Hodge reconstruction lead to C.2-core. In both geometries the upstream purpose is the same: to produce a compact signed azimuthal-vorticity child, a strict parent–child scale gap and a fixed local flat-action budget. The synchronized comparison in section 7 displays these parallel roles explicitly.

Once this geometry-specific interface has been reached, the proof is common. Bridge D, theorem 5.11, reduces the possible origins of a continuing active packet to four alternatives: source-born, retained-active, strict-restart, and spectator. The automatic deep-handoff theorem theorem 6.26 places the active packet in the subparabolic regime required by Bridge E. The successive material, full-adjoint and primal–adjoint polarizations in theorem 6.29 then remove the Euler-compatible background and expose the viscosity-dependent remainder. The surviving branch is routed either to a strict physical successor or to a fixed positive Leray dissipation payment. The no-unpaid-loop proposition theorem 6.27 prevents an indefinitely continuing successor-only restart chain, while chronological reselection makes the paid packets pairwise disjoint in time.

The resulting architecture gives a concrete meaning to the Euler-firewall principle used throughout the paper. Scale-invariant Hodge identities, transport relations, compactness mechanisms and geometric selectors are used to organize the cascade, but none of them is treated as a terminal Navier–Stokes closure merely because it is strong enough to select a smaller scale. Terminal detection occurs only after the Euler background has been canceled and an explicit viscous contribution or a fixed charge against the finite physical dissipation ledger has appeared. This distinction is also what allows the whole-space and periodic arguments to share the same downstream engine despite their different upstream geometries.

In the present formulation, the two compressed assembly statements theorems 1.3 and 1.4 and the two firewall propositions theorems 7.2 and 7.3 summarize the precise scope of the paper. The proof uses no z-reflection assumption, no post-terminal suitable continuation, and no additional common downstream conditional branch beyond the polarized Bridge E mechanism.

The recent forced-breakdown construction announced by OpenAI [22] is complementary rather than contradictory to this unforced program. In Fefferman's formulation [14], alternatives (C) and (D) allow smooth forcing, while (A) and (B) impose $f \equiv 0$. The logical compatibility of the two pairs underscores the importance of stating the forcing class and geometry explicitly. The two geometries therefore share a common funded terminal mechanism even though their singular-scale selection procedures are genuinely different.

9. Acknowledgements

This work was conceived, initiated, and directed by the author. The author formulated the central mathematical questions, determined the overall proof strategy and version-by-version research agenda, selected the scale-invariant quantities and comparison regimes to be pursued, and decided which proposed routes, reductions, and restart mechanisms were mathematically acceptable. In particular, the author repeatedly required that Euler-compatible identities be separated from genuinely Navier–Stokes-specific mechanisms, adopted

and enforced the resulting Euler-firewall principle throughout the proof architecture, and directed the subsequent use of viscosity-polarized observables and detectors. The author evaluated the proposed derivations, identified points requiring further proof or rejection, selected the statements retained in the compressed manuscript, and is responsible for all mathematical claims, interpretations, and the final form of the paper.

During the development of the project, the author used OpenAI's ChatGPT as a research assistant across an extended sequence of working sessions, including GPT-5.6 Sol in September 2026. ChatGPT contributed exploratory calculations and derivations, proposed alternative proof routes, constructed countermodels and stress tests for tentative arguments, assisted with literature and source-scope reviews, performed dependency and circularity checks, helped organize and compress the proof architecture, developed and refined successive material, adjoint, and polarization formulations in response to the author's mathematical direction, and assisted with LaTeX drafting and notation consistency. The reference to ChatGPT here denotes the OpenAI assistant system used across those sessions rather than a claim that a single persistent model instance participated in every stage.

The author has independently determined what material is included in the manuscript and retains sole responsibility for checking the proofs and for the submitted version.

10. References

References

[1] T. C. Anderson and A. Vagharshakyan, *A simple proof of the sharp weighted estimate for Calderón–Zygmund operators on homogeneous spaces*, J. Geom. Anal. **24** (2014), no. 3, 1276–1297.

[2] J. T. Beale, T. Kato and A. Majda, *Remarks on the breakdown of smooth solutions for the 3-D Euler equations*, Comm. Math. Phys. **94** (1984), no. 1, 61–66.

[3] L. Caffarelli, R. Kohn and L. Nirenberg, *Partial regularity of suitable weak solutions of the Navier–Stokes equations*, Comm. Pure Appl. Math. **35** (1982), no. 6, 771–831.

[4] J. R. Cannon and E. DiBenedetto, The initial value problem for the Boussinesq equations with data in L^p, in *Approximation Methods for Navier–Stokes Problems*, Lecture Notes in Mathematics, vol. 771, Springer, Berlin, 1980, pp. 129–144, doi:10.1007/BFb0086903.

[5] D. Chae, *Global regularity for the 2D Boussinesq equations with partial viscosity terms*, Adv. Math. **203** (2006), 497–513.

[6] D. Chae and J. Lee, *On the regularity of the axisymmetric solutions of the Navier–Stokes equations*, Math. Z. **239** (2002), 645–671.

[7] C.-C. Chen, R. M. Strain, T.-P. Tsai and H.-T. Yau, *Lower bound on the blow-up rate of the axisymmetric Navier–Stokes equations*, Int. Math. Res. Not. IMRN (2008), article rnn016.

[8] H. Chen, D. Fang and T. Zhang, *Regularity of 3D axisymmetric Navier–Stokes equations*, Discrete Contin. Dyn. Syst. **37** (2017), no. 4, 1923–1939.

[9] H. Chen, T.-P. Tsai and T. Zhang, *Remarks on local regularity of axisymmetric solutions to the 3D Navier–Stokes equations*, Commun. Partial Differential Equations **47** (2022), no. 8, 1680–1699; correction: *ibid.* **48** (2023), 987–988.

[10] Clay Mathematics Institute, *Navier-Stokes Announcement*, September 10, 2026, https://www.claymath.org/news/navier-stokes-announcement/.
[11] P. Constantin and C. Foias, *Navier–Stokes Equations*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, 1988.
[12] C. R. Doering and J. D. Gibbon, *Applied Analysis of the Navier–Stokes Equations*, Cambridge Texts in Applied Mathematics, vol. 12, Cambridge University Press, Cambridge, 1995.
[13] L. Escauriaza, G. Seregin and V. Šverák, *$L_{3,\infty}$-solutions of the Navier–Stokes equations and backward uniqueness*, Russian Math. Surveys **58** (2003), no. 2, 211–250.
[14] C. L. Fefferman, *Existence and smoothness of the Navier–Stokes equation*, in J. Carlson, A. Jaffe and A. Wiles (eds.), *The Millennium Prize Problems*, Clay Mathematics Institute and American Mathematical Society, 2006, 57–67.
[15] A. Friedman, *Interior estimates for parabolic systems of partial differential equations*, J. Math. Mech. **7** (1958), 393–417.
[16] T. Y. Hou and C. Li, *Global well-posedness of the viscous Boussinesq equations*, Discrete Contin. Dyn. Syst. **12** (2005), 1–12.
[17] O. A. Ladyzhenskaya, *The Mathematical Theory of Viscous Incompressible Flow*, 2nd ed., Gordon and Breach, New York–London, 1969.
[18] O. A. Ladyzhenskaya, *Sixth problem of the millennium: Navier–Stokes equations, existence and smoothness*, Russian Math. Surveys **58** (2003), no. 2, 251–286.
[19] J. Leray, *Sur le mouvement d'un liquide visqueux emplissant l'espace*, Acta Math. **63** (1934), 193–248.
[20] J.-G. Liu and W.-C. Wang, *Characterization and regularity for axisymmetric solenoidal vector fields with application to Navier–Stokes equation*, SIAM J. Math. Anal. **41** (2009), no. 5, 1825–1850.
[21] A. J. Majda and A. L. Bertozzi, *Vorticity and Incompressible Flow*, Cambridge Texts in Applied Mathematics, vol. 27, Cambridge University Press, Cambridge, 2002.
[22] OpenAI, *Finite time blowup for Navier–Stokes*, manuscript and accompanying formalization, September 8, 2026; see also *On the Navier–Stokes Millennium Prize Problem*, https://openai.com/index/navier-stokes-solution/.
[23] W. S. O'zański and S. Palasek, *Quantitative control of solutions to the axisymmetric Navier–Stokes equations in terms of the weak L^3 norm*, Ann. PDE **9** (2023), Article 15.
[24] J. C. Robinson, J. L. Rodrigo and W. Sadowski, *The Three-Dimensional Navier–Stokes Equations: Classical Theory*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 2016.
[25] G. Seregin, *Local regularity of axisymmetric solutions to the Navier–Stokes equations*, Anal. Math. Phys. **10** (2020), Article 46.
[26] J. Serrin, *On the interior regularity of weak solutions of the Navier–Stokes equations*, Arch. Rational Mech. Anal. **9** (1962), 187–195.
[27] Y. Shi, *2D inviscid Boussinesq equations and 3D axisymmetric Euler equations: (1) A unification (E_m), (2) finite-time blow-up of two unified $(1+1)D$ systems rigorously derived from (E_m)*, arXiv:2603.26715v5 [math.AP] (2026).
[28] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, NJ, 1970.
[29] T. Tao, *Finite time blowup for an averaged three-dimensional Navier–Stokes equation*, J. Amer. Math. Soc. **29** (2016), no. 3, 601–674.
[30] T. Tao, *Quantitative bounds for critically bounded solutions to the Navier–Stokes equations*, in *Nine Mathematical Challenges—An Elucidation*, Proc. Sympos. Pure Math., vol. 104, American Mathematical Society, Providence, RI, 2021, 149–193.
[31] R. Temam, *Navier–Stokes Equations: Theory and Numerical Analysis*, AMS Chelsea Publishing, Providence, RI, 2001; reprint with corrections of the third revised edition.

[32] M. R. Ukhovskii and V. I. Yudovich, *Axially symmetric flows of ideal and viscous fluids filling the whole space*, J. Appl. Math. Mech. **32** (1968), 52–62.
[33] J. Wu, *The 2D Incompressible Boussinesq Equations*, Peking University Summer School Lecture Notes, 2012.

California, United States
Email address: ymshi@protonmail.com