

Equally-distributed-equivalent from an L -function viewpoint (Working paper)

Yao-Rui Yeo

New York University Grossman School of Medicine, New York, USA

yao-rui.yeo@nyulangone.org

June 15, 2026

Abstract

Atkinson's index is a measure of income inequality, first defined in 1970 by invoking the idea of Equally Distributed Equivalent, i.e. the level of income that, if equally distributed in a hypothetical scenario, gives the same level of welfare as the current income distribution. Originally a concept in economics, Atkinson's index is increasingly applied to studies in health inequality. Using functional analytic techniques with an L -function viewpoint, we give a new decomposition of Atkinson's index into a harmonic-like mean among subgroup welfare levels. Our decomposition clarifies how subgroup welfare levels are differentially weighted for Atkinson's index as inequality aversion varies, and gives rise to two applications. Firstly, we give a theoretical characterization on how inequality aversion influences resource allocation among different societal subgroups. Secondly, we obtain a duality principle for inequality aversion, extending applications of Atkinson's index to welfare metrics that vary inversely proportionately to income or health levels.

JEL classification: C65; D63; I14

1 Introduction

Many measures exist to estimate population inequality in both economics and health [3], with the Gini index being the most famous. However, a good measure for inequality analyses should satisfy three important properties [10]: subgroup decomposability, where total inequality is divided into its constituent components; the Pigou-Dalton Principle, where a transfer of a desirable variable (e.g. wealth) from the rich to the poor results in less inequality as long as it does not bring the rich to a worse situation than the poor; and avoids value judgement, for instance by including an explicit parameter that changes the weights placed on various percentiles of the distribution (to allow sensitivity analysis with respect to this parameter for policy decisions). Using this definition of goodness, the Gini index does not qualify as a good measure for two reasons: it does not avoid value judgement; and it is only subgroup decomposable if subgroups are explicitly ranked. A detailed analysis (by the same paper [10]) concluded that Atkinson's index [1] stands out as the best inequality measure in health. Additionally, a quantitative analysis of income inequality measures by Shorrocks imply the generalized entropy index [17, Equation 31] is the only one-parameter family that is subgroup decomposable under relatively weak restrictions on homogeneity, and the generalized entropy index can be viewed as a monotonic transformation of Atkinson's index. As such, we focus on Atkinson's index in this paper.

Atkinson defined his index in [1] by invoking the idea of Equally Distributed Equivalent (EDE), reflecting the willingness to trade off aggregate benefits for income to be more equally distributed. This index is defined on the real line as

$$EDE_H(\epsilon) := \begin{cases} \left(\sum_i H_i^{1-\epsilon} \mu_i \right)^{\frac{1}{1-\epsilon}} & \epsilon \neq 1 \\ \prod_i H_i^{\mu_i} & \epsilon = 1 \end{cases} \quad (1)$$

where H_i is the income level for subgroup i with each H_i distinct and nonzero, μ_i is a weight for subgroup i (with the sum of all μ_i equaling 1), and ϵ is Atkinson's inequality aversion parameter. The weights μ_i are chosen depending on the application. In Atkinson's original formulation, μ_i is simply the proportion of the total population at income level H_i , but μ_i can also be income-dependent, causality-dependent, and so on [15]. Applications of this index in health inequality studies, an emerging body

of research [7, 8, 14, 15, 19], replaces the variables H_i with subgroup health levels. For applicability of our results to both economics and health, we will henceforth call H_i the wealth level unless discussing specific examples.

Note that our definition of $EDE_H(\epsilon)$ is the non-normalized form of the original definition given by Atkinson. This is the perspective taken in health inequality studies, and we find that this form is most suitable for our arguments in this paper. Also, instead of restricting ϵ to be strictly non-negative, we allow use of the entire real line as it has important implications in our analysis (e.g. the duality statement stated at the end of this Section).

The main goal of this paper is to decompose Atkinson’s index $EDE_H(\epsilon)$ as weighted sums of the wealth levels H_i . To agree with the original work of Atkinson, such a weighted sum should give greater weight to subgroups with lower wealth at higher inequality aversion ϵ . Therefore, our decomposition should resemble a harmonic-like sum in order to emphasize sensitivity in lower wealth groups. Such a decomposition is also supported mathematically, for $EDE_H(\epsilon)$ is a generalized mean with negative exponents at $\epsilon > 1$.

Our harmonic-like sum decomposition of Atkinson’s index is obtained through functional analysis with an L -function viewpoint, and will be explained in Section 2. Following this, we demonstrate how our techniques give greater subgroup-level insights in EDE weighting of subgroups (Section 3). An example of such an insight is the following: subgroups that do not correspond to the highest or lowest wealth may not be monotonically weighted as ϵ varies. In other words, Atkinson’s index weighs subgroups in a highly interesting way, especially for subgroups with wealth levels sufficiently close to the one with the lowest wealth, and a precise formulation is given in the discussion leading up to Example 3.5.

After the above analyses, we explain how our decomposition naturally arises in resource allocation problems with health equity constraints (Section 4). In particular, our work provides a framework for Robson’s resource allocation experiments [15], which are used to elicit health-based Atkinson’s inequality aversion parameter ϵ in various settings.

Our decomposition can also be used to explain how Atkinson’s idea of EDE can be carried over to economics or health metrics associated with negative outcomes, such as poverty or mortality rate over a time horizon (Section 5). To our knowledge, this has not been done in the literature. Such an explanation is important: EDE for metrics

associated with negative outcomes should give greater weight to subgroups with higher negative outcomes at higher ϵ , but a straightforward application of Equation 1 will give lower negative outcomes as ϵ increases. In our argument to derive EDE for metrics associated with negative outcomes, we also obtain a surprising connection between opposing metrics in both economics (e.g. income and poverty) and health (e.g. life expectancy and mortality rate over a time horizon). We state this duality for income and poverty here, leaving a more general discussion to Section 5. Our duality principle vastly generalizes a recent observation of Sterck [23] that minimizing overall poverty level (at $\epsilon = 0$) is equivalent to maximizing EDE income at $\epsilon = 2$.

EDE Duality between Income and Poverty (c.f. Duality Principle 5.1). Minimizing EDE poverty at Atkinson’s inequality aversion parameter ϵ is equivalent to maximizing EDE income at $\epsilon + 2$.

In Section 6, we apply our mathematical methods to analyze potentially avoidable mortality in the setting of Ontario, Canada, before ending with some concluding remarks in Section 7.

2 Functional analysis on Atkinson’s index

Let us list some basic properties of Atkinson’s index $EDE_H(\epsilon)$ (Equation 1) that is almost immediate from definition. $EDE_H(\epsilon)$ gives increasingly greater weight to the subgroup with lowest wealth as ϵ increases, and is a strictly decreasing function in ϵ . In particular, $EDE_H(\epsilon)$ plateaus to the minimum wealth level among the subgroups as ϵ increases, and plateaus to the maximum wealth level among the subgroups as ϵ decreases. Further, by definition of the individual utility function for Atkinson’s index

$$U(x) = \frac{x^{1-\epsilon}}{1-\epsilon}$$

and its concavity, Atkinson’s index satisfies the main properties of social welfare measures, such as wealth homogeneity, population homogeneity, and the Pigou-Dalton Principle [1, 17].

The definition of Atkinson’s index is a little opaque for subgroup-level analyses at first glance. For instance, it does not inform us how the H_i ’s are differentially weighted as ϵ varies. While Shorrocks performed a subgroup decomposition on Atkinson’s index

into an arithmetic-like sum to obtain between-group and within-group factors [17, 18], we will decompose $EDE_H(\epsilon)$ in an alternative way as a harmonic-like sum

$$EDE_H(\epsilon) = \left(\sum_i \frac{1}{H_i} f_{i,\epsilon} \mu_i \right)^{-1}$$

for a family of functions $f_{i,\epsilon}$ in ϵ and depending on H_i and μ_i . Our decomposition does not have between-group and within-group factors, but a single “factor” $f_{i,\epsilon}$ per subgroup. The functions $f_{i,\epsilon}$ are not uniquely determined, but a “natural” expression can be derived by applying Euler’s Theorem for homogeneous functions. However, simply applying Euler’s Theorem does not give much economical or mathematical insights. As such, we do not apply Euler’s Theorem. Instead, we perform functional analysis on Atkinson’s index, guiding our work by analogy to the theory of L -functions (see Appendix A).

The purpose of this section is to state the following mathematical results, with proofs given in Appendix B. For convenience in the arguments, these results are formulated as the inverse of Atkinson’s index as stated in Equation 1.

Theorem 2.1. *Let $\epsilon > 2$, let $p = \epsilon - 1$, and let q be the number such that $1/p + 1/q = 1$. Then there exist a unique collection of non-negative values $f_{i,\epsilon}$ satisfying*

$$\sum_i f_{i,\epsilon}^q \mu_i = 1, \tag{2}$$

and such that these $f_{i,\epsilon}$ maximizes the expression

$$F(f_{i,\epsilon}) := \sum_i \frac{f_{i,\epsilon} \mu_i}{H_i}.$$

Furthermore, this maximum value equals $EDE_H(\epsilon)^{-1}$.

Equation 2 can be viewed as a q -analog of the usual weighted sum. Note that calling our Equation a q -analog is appropriate: As $\epsilon \rightarrow \infty$, necessarily $q \rightarrow 1$, and Equation 2 returns the usual weighted sum.

Corollary 2.2. *The values $f_{i,\epsilon}$ in Theorem 2.1 can be explicitly determined:*

$$f_{i,\epsilon} := \left(\sum_j \left(\frac{H_j}{H_i} \right)^{1-\epsilon} \mu_j \right)^{-(1+\frac{1}{1-\epsilon})} = \left(\frac{EDE_H(\epsilon)}{H_i} \right)^{\epsilon-2}.$$

The expressions above imply $f_{i,\epsilon}$ can be viewed as analytic functions at $\epsilon \neq 1$.

Theorem 2.3. *Let $\epsilon > 2$. Suppose $q > 1$ is a number satisfying the following two conditions.*

- *There exists unique values $f_i > 0$ such that*

$$\sum_i f_i^q \mu_i = 1$$

and f_i maximizes the expression

$$F(a_i) := \sum_i \frac{1}{H_i} a_i \mu_i.$$

- *$F(f_i)$ equals $EDE_H(\epsilon)^{-1}$, where f_i is specified in the above condition.*

Then q is unique, and equals the value in Theorem 2.1.

Theorem 2.4 (Analytic Continuation). *Let q be as defined in Theorem 2.1. Consider the real-valued function*

$$L(H_i, \epsilon) := \max_{\substack{\sum_i f_{i,\epsilon}^q \mu_i = 1 \\ f_i \geq 0}} \left\{ \sum_i \frac{f_{i,\epsilon} \mu_i}{H_i} \right\}$$

which is well-defined on $\epsilon > 2$ by Theorem 2.3. Then $L(H_i, \epsilon)$ can be analytically continued to the entire real line, except for a removable singularity at $\epsilon = 1$. Furthermore, $L(H_i, \epsilon)$ is a positive function with $L(H_i, \epsilon) = EDE_H(\epsilon)^{-1}$.

Theorem 2.5 (Functional Equation). *The function $L(H_i, \epsilon)$ defined in Theorem 2.4 satisfies*

$$L(H_i, -\epsilon)^{-1} = L(H_i^{-1}, 2 + \epsilon).$$

In summary, the five mathematical statements above narrate the following story. Theorem 2.1 gives a unique harmonic-like sum that is simultaneously equal to Atkinson's index at $\epsilon > 2$ and minimal under a q -analog weighted sum constraint, showing

a robust harmonic-like decomposition of Atkinson's index at $\epsilon > 2$. Corollary 2.2 explicitly defines the components $f_{i,\epsilon}$ of the harmonic-like decomposition. (Interestingly, the expression in this Corollary coincides with the “natural” definition of $f_{i,\epsilon}$ via Euler's Theorem for homogeneous functions.) Theorem 2.3 tells us that the technical condition imposed on $f_{i,\epsilon}$ allows us to deform Atkinson's weights in a well-defined manner as ϵ varies, and is the only sensible deformation that naturally extends Atkinson's weights as ϵ varies. In particular, this Theorem shows our definition of EDE-factors in the next Section makes sense (Definition 3.1). Theorem 2.4 mathematically justifies our harmonic-like decomposition agrees with Atkinson's index at all ϵ , not just $\epsilon > 2$. Finally, Theorem 2.5 exhibits a dual pairing on the harmonic-like decomposition, resulting in a duality principle that will be explained in Section 5 (Duality Principle 5.1).

3 EDE-factors

Due to Section 2, Atkinson's index can be decomposed as a harmonic-like sum

$$EDE_H(\epsilon) = \left(\sum_i \frac{1}{H_i} f_{i,\epsilon} \mu_i \right)^{-1}$$

where $f_{i,\epsilon}$ is defined as

$$f_{i,\epsilon} := \begin{cases} \left(\sum_j \left(\frac{H_j}{H_i} \right)^{1-\epsilon} \mu_j \right)^{-(1+\frac{1}{1-\epsilon})} & \epsilon \neq 1 \\ \prod_j \left(\frac{H_j}{H_i} \right)^{\mu_j} & \epsilon = 1. \end{cases}$$

These $f_{i,\epsilon}$ satisfy a technical weighted normality condition:

$$\sum_i f_{i,\epsilon}^q \mu_i = 1, \quad q = \frac{\epsilon - 1}{\epsilon - 2},$$

allowing us to extend Atkinson's weights μ_i through the expressions $f_{i,\epsilon}^q \mu_i$. In other words, we extend Atkinson's weights via a q -analog of the expression $f_{i,\epsilon} \mu_i$. Such an extension $w_i(\epsilon)$ requires agreement with Atkinson's weight without inequality aversion considerations, i.e. $w_i(0) = \mu_i$. As such, we need to shift the ϵ in $f_{i,\epsilon}$ by 1 for this

equality to hold.

Definition 3.1. The *EDE-factor* for subgroup i can be explicitly defined in two ways:

$$w_i(\epsilon) := \left(\sum_j \left(\frac{H_j}{H_i} \right)^{-\epsilon} \mu_j \right)^{-1} \mu_i = \left(\frac{EDE_H(\epsilon + 1)}{H_i} \right)^\epsilon \mu_i.$$

A reasonable question can be asked at this point: Why did we define the EDE-factor by (a shift of) $f_{i,\epsilon}^q \mu_i$ instead of a weighted proportion among $f_{i,\epsilon} \mu_i$? In fact, by Corollary 2.2, defining EDE-factors via weighted proportions will arrive at the same result as

$$\frac{f_{i,\epsilon} \mu_i}{\sum_k f_{k,\epsilon} \mu_k} = \left(\frac{EDE_H(\epsilon - 1)}{H_i} \right)^{\epsilon-2} \mu_i,$$


and a shift of ϵ by 2 is required to agree with Atkinson's weight without inequality aversion considerations.

Another reasonable question can also be asked: Why did we choose to shift ϵ when defining EDE-factors? Other than the logical agreement with Atkinson's weights at $\epsilon = 0$, we give two other benefits. Firstly, slightly shifting ϵ ahead (by 1 or 2) does not detract the value of comparing subgroup weights across ϵ for health policy making in real-world settings, as elicitation of ϵ in health generally gives rise to a distribution in a wide range between 0 and 20 [7, 14, 19]. Secondly, in Section 4, we will show that our definition of EDE-factors result in a more direct comparison for resource allocation problems.

Before discussing properties of EDE-factors, we give a simple example of EDE-factors on two hypothetical policy scenarios.

Example 3.2. Consider a population split equally into 2 subgroups ($\mu_1 = \mu_2 = 0.5$) with income levels $H_1 = \$70,000$ (lower income), $H_2 = \$100,000$ (higher income). Also consider two Scenarios: A, where individuals in the lower income subgroup H_1 are given additional \$5,000; and B, where individuals in the higher income subgroup H_2 are given additional \$20,000. Then the graphs of Atkinson's index (computing

EDE) and EDE-factors (computed using Definition 3.1) are graphed below.



If inequality aversion is not considered, Scenario B results in a higher average income as four-fold more resources are additionally given compared to Scenario A. However, a computation of EDE income shows a more rapid decrease as ϵ increases, with a tipping point at $\epsilon_{tip} \approx 3.37$ and $EDE_H(\epsilon_{tip}) \approx \$84,500$. This is also evident by the EDE-factors, as Scenario B puts a much higher weight on the lower income subgroup in EDE computations due to a larger income gap between subgroups.

Immediate properties from functional analysis

From our discussion in the previous Section, clearly the EDE-factors $w_i(\epsilon)$ are non-negative and sum to 1 for all ϵ :

$$\sum_i w_i(\epsilon) = 1.$$

At $\epsilon = 0$, this is simply the base case condition that the sum of all μ_i equal 1. These EDE-factors can be seen as a spiritual answer to questions raised in [15, 17] on a method to decompose measures, such as the Atkinson's index, as a simple weighted sum. We now discuss how EDE-factors imply most of the homogeneity and transfer properties in Atkinson's index. Proof of the statements are given in Appendix C.

Wealth homogeneity. By Definition 3.1, the EDE-factors $w_i(\epsilon)$ are not affected by a uniform scaling of wealth levels $H_i \mapsto kH_i$ for some positive constant k . Thus, by Theorem 2.4, this implies wealth homogeneity for Atkinson's index, i.e. $EDE_H(\epsilon)$ is multiplied by the same constant k under a uniform scaling of wealth levels.

Population homogeneity. Suppose each subgroup i is replicated n times with

welfare levels $H_{(i,1)} = \dots = H_{(i,n)}$ equal to H_i , and such that the replications are weighted $\mu_i\omega_1, \dots, \mu_i\omega_n$, with the ω_k 's summing to 1.

Proposition 3.3. *EDE_H(ϵ) is unchanged with respect to the subgroup replications described above.*

Pigou-Dalton Principle. The Pigou-Dalton Principle is a transfer principle that asserts any social welfare function must prefer allocations that are more equitable. Formally, if $H_i > H_j$, then a transfer of $\Delta > 0$ from H_i to H_j , in such a way that

$$H_i - \Delta \geq H_j + \Delta^*, \quad \Delta^* := \Delta \frac{\mu_i}{\mu_j}$$

must not decrease $EDE_H(\epsilon)$. The Pigou-Dalton Principle is easily seen to hold for Atkinson's index due to the concavity of the utility function. We offer another argument for $\epsilon > 2$ via the results in Section 2.

Proposition 3.4. *Theorem 2.1 implies the Pigou-Dalton Principle for Atkinson's index at $\epsilon > 2$.*

The above Proposition tells us that the Pigou-Dalton Principle can be inferred from Theorem 2.1 when $\epsilon > 2$, though the Theorem cannot be used to prove the Pigou-Dalton Principle at $0 < \epsilon < 2$. However, this is sufficient to show that Theorem 2.1 generalizes the Pigou-Dalton Principle for inequality studies using negative metrics (e.g. poverty level); see Section 5 for a discussion on this.

A non-monotonic property

As $EDE_H(\epsilon)$ tends to the minimum wealth level as ϵ increases, the EDE-factors obey the following asymptotic property:

$$\lim_{\epsilon \rightarrow \infty} w_i(\epsilon) = \begin{cases} 1 & \text{if } H_i = \min\{H_k\}_k; \\ 0 & \text{otherwise.} \end{cases}$$

However, EDE-factors demonstrate a very interesting property: subgroups that do not correspond to the highest or lowest wealth may not be monotonically weighted as ϵ increases. More precisely, let $w_i(\epsilon)$ correspond to the EDE-factor of such a subgroup.

By taking the derivative, one gets

$$\frac{d}{d\epsilon} w_i(\epsilon) = \left(\sum_j \left(\frac{H_j}{H_i} \right)^{-\epsilon} \mu_j \right)^{-2} \mu_i \left(\sum_j (\ln H_j - \ln H_i) \left(\frac{H_j}{H_i} \right)^{-\epsilon} \mu_j \right)$$

The above derivative is non-increasing exactly when the right-most sum is non-negative, or equivalently


$$\ln H_i \geq \frac{\sum_{j \neq i} H_j^{-\epsilon} \mu_j \ln H_j}{\sum_{j \neq i} H_j^{-\epsilon} \mu_j}.$$

As the right hand side is a decreasing function in ϵ (by applying the Cauchy-Schwarz inequality on its derivative), this implies

$$\ln H_i \geq \sum_{j \neq i} \ln H_j^{\mu_j}.$$

If we assume $H_i > 1$ (with no loss of generality by wealth homogeneity), the above inequality implies its EDE-factor will increase at lower levels of ϵ to a unique maxima before monotonically tending to 0 as long as H_i is less than the relative geometric mean of the other subgroups. In other words, Atkinson's index may increasingly weight subgroups close to the lowest wealth at lower levels of ϵ . This fact cannot be seen directly from the original definition of Atkinson's index (Equation 1).

Example 3.5. Consider a population split into 3 subgroups with Atkinson's weights $\mu_1 = 0.25$, $\mu_2 = 0.35$, $\mu_3 = 0.4$ and income levels $H_1 = \$70,000$ (lowest income), $H_2 = \$80,000$ (moderate income), $H_3 = \$100,000$ (highest income). Then the graphs of Atkinson's index and EDE-factors are graphed below.



In this example, the moderate income subgroup H_2 is increasingly weighted until $\epsilon_{2,peak} \approx 2.76$, with EDE-factor peaking at $w_2(\epsilon_{2,peak}) \approx 0.377$. This shows that at low levels of inequality aversion, the calculation of $EDE_H(\epsilon)$ puts emphasis in both H_1 and H_2 , and not just the lowest income subgroup H_1 .

4 Maximal EDE resource allocation

Consider the problem of reallocating current resources between subgroups to achieve maximal EDE wealth at inequality aversion ϵ . Let p_i be the (local) production factor of subgroup i , and let r_i be the amount of resources allocated to subgroup i . Then we can adopt a simplified Cobb-Douglas production function for the subgroup wealth levels: $H_i = p_i r_i^\alpha$, where $\alpha \in (0, 1]$ is the resource elasticity variable. Although proofs of everything in this Section can be done with $\alpha < 1$ (see Appendix D), we adopt a linear production function $\alpha = 1$ in the main paper for two reasons: simplicity in illustrations, and to directly compare our results with those in the literature.

Henceforth suppose $H_i = p_i r_i$. Assuming the sum of all current resources equals R , a resource reallocation problem that seeks to maximize EDE can be stated as follows: Maximize

$$EDE_H(\epsilon) = \left(\sum_i (p_i r_i)^{1-\epsilon} \mu_i \right)^{\frac{1}{1-\epsilon}}$$

subject to the condition

$$\sum_i r_i = R.$$

If inequality aversion is not a consideration ($\epsilon = 0$), this problem has a simple solution: Allocate all resources to the subgroup with the highest value of $p_i \mu_i$, i.e. best weighted production factor. However, if equity is a consideration ($\epsilon > 0$), we can solve this problem via Lagrange Multipliers, telling us that $EDE_H(\epsilon)$ is maximized if $H_i = \widetilde{H}_{i,\epsilon}$, where

$$\widetilde{H}_{i,\epsilon} := R \left(\sum_j \frac{1}{p_j} \left(\frac{p_j \mu_j}{p_i \mu_i} \right)^{\frac{1}{\epsilon}} \right)^{-1}. \quad (3)$$

Equation 3 can be compared to a similar setup described in [15].

EDE-factors offer a comparison of arriving at maximal EDE resource allocation

given current allocation. Writing $w_{i,\epsilon} = w_i(\epsilon)$, Definition 3.1 implies

$$H_i = d \left(w_{i,\epsilon}^{-1} \mu_i \right)^{\frac{1}{\epsilon}}$$

for an expression d that is constant across all subgroups. As $H_i = p_i r_i$, dividing by p_i and summing across i gives

$$R = d \sum_j \frac{1}{p_j} \left(w_{j,\epsilon}^{-1} \mu_j \right)^{\frac{1}{\epsilon}},$$

and a rearrangement gives

$$H_i = R \left(\sum_j \frac{1}{p_j} \left(\frac{w_{j,\epsilon}^{-1} \mu_j}{w_{i,\epsilon}^{-1} \mu_i} \right)^{\frac{1}{\epsilon}} \right)^{-1} \quad (4)$$

which is very similar to Equation 3 for $\widetilde{H_{i,\epsilon}}$. There exists a unique set of EDE-factors w_i achieving maximal $EDE_H(\epsilon)$, given by

$$w_i = \frac{1}{p_i} \left(\sum_j \frac{1}{p_j} \right)^{-1}. \quad (5)$$

In general, the EDE-factors achieving maximal $EDE_H(\epsilon)$ are not constant functions for non-linear production functions (see Equation 16 in Appendix D.)

Equations 3 and 4 are useful for policy making as it allows us to compare resource allocation as a ratio between subgroups without requiring explicit knowledge on total resources (the R 's cancel out under a ratio), allowing for scalability or if total resources are relatively unknown but with known effects. For instance, the expressions $p_i \mu_i$ and $w_{i,\epsilon}^{-1} \mu_i$, in the fractions of Equations 3 and 4 respectively, are related by a factor of $w_{i,\epsilon} p_i$:

$$p_i \mu_i = p_i w_{i,\epsilon} \cdot (w_{i,\epsilon}^{-1} \mu_i).$$

At current resource allocation, this means that, for each subgroup i , the deformation of the weighted production factor $p_i w_{i,\epsilon}$, consisting of production factor p_i and EDE-factor $w_{i,\epsilon}$, can serve as a measure of current resource allocation compared to the optimal allocation giving rise to the maximal EDE at a given $\epsilon > 0$.

5 EDE calculations for negative metrics

Let M be a metric that varies inversely proportional to wealth level H (e.g. poverty level, mortality rate across a specified time horizon). We call M a *negative metric*. Such metrics are important in applications for both economics and health [8]. Note that the EDE-adjusted M as ϵ varies cannot be calculated by directly substituting M into Atkinson's index $EDE_H(\epsilon)$ (Equation 1) as this would tend to the lowest level of M (i.e. highest level of wealth), contrary to what we expect.

With that said, calculations on negative metrics M can be done through our functional equation (Theorem 2.5). Let $EDE^\dagger(\epsilon)$ be the proposed EDE for M at inequality aversion parameter ϵ in the sense of Atkinson. For such an $EDE^\dagger(\epsilon)$ to be compatible with $EDE_H(\epsilon)$ for wealth, we require $EDE^\dagger(\epsilon)$ to satisfy the same fundamental properties as those outlined in [1, 17]. Therefore, using the fact that (non-normalized) Atkinson's index is a monotonic transformation of Shorrocks work in [17], $EDE^\dagger(\epsilon)$ has to equal $EDE_M(\eta)$ for some η . In particular, the results of Section 2 apply to $EDE^\dagger(\epsilon)$ through $EDE_M(\eta)$. Furthermore, η needs to be a non-positive function by fundamental results on constant relative risk aversion [12], and also needs to be chosen such that normalization holds (i.e. $EDE^\dagger(0)$ equals the usual arithmetic mean of M among subgroups).

By Theorem 2.5,

$$EDE_M(2 - \epsilon) = EDE_H(\epsilon)^{-1}. \quad (6)$$

In other words, EDE for M also varies inversely proportional to EDE for H , but at different inequality aversion parameters. As the parameters $2 - \epsilon$ and ϵ are negative linearly related, the most appropriate choice for η is $2 - \epsilon$, plus a (linear) translation of 2 to satisfy normalization. In summary, $-\epsilon$ is our choice of η , or equivalently

$$EDE^\dagger(\epsilon) = EDE_M(-\epsilon).$$

The above paragraph is a generalization of Sterck's observation [23] that minimizing overall poverty level is equivalent to maximizing EDE wealth at $\epsilon = 2$, for this observation is simply a consequence of substituting $\epsilon = 0$ into Equation 6. In other words, we extend Sterck's pointwise equivalence at $\epsilon = 0$ into a structural equivalence valid for all $\epsilon \geq 0$. This is reformulated as the Duality Principle below.

Duality Principle 5.1. Minimizing the EDE of a negative metric M at ϵ is equivalent

lent to maximizing the EDE of its inverse metric at $\epsilon + 2$.

Because of Equation 6, everything discussed in Sections 3 and 4 can be appropriately carried over to negative metrics by replacing ϵ with $-\epsilon$.

EDE-factors for negative metrics

For negative metrics, the EDE-factors are

$$w_i^\dagger(\epsilon) := \left(\sum_j \left(\frac{M_i}{M_j} \right)^{-\epsilon} \mu_j \right)^{-1} \mu_i = \left(\frac{M_i}{EDE_M(-\epsilon + 1)} \right)^\epsilon \mu_i.$$

The three main properties still hold (wealth homogeneity, population homogeneity, Pigou-Dalton Principle). In fact, our functional-analytic discussion in Section 2 actually implies the Pigou-Dalton Principle in this case.

Corollary 5.2. *Theorem 2.1 implies the Pigou-Dalton Principle for negative metrics at all $\epsilon \geq 0$.*

Proof. This is immediate by applying Proposition 3.4 to Equation 6. $\square$

The non-monotonic property of EDE-factors works the opposite way for negative metrics: $w_i^\dagger(\epsilon)$ is strictly non-increasing as ϵ increases precisely when

$$\ln M_i \leq \sum_{j \neq i} \ln M_j^{\mu_j}.$$

In other words, at lower levels of ϵ Atkinson's index may increasingly weigh a subgroup i with negative metric M_i if M_i greater than the relative geometric mean of (the negative metrics of) the other subgroups.

There also exists an arithmetic-like sum decomposition for $EDE^\dagger(\epsilon) = EDE_M(-\epsilon)$, given by

$$EDE^\dagger(\epsilon) = \sum_i M_i \left(w_i^\dagger(\epsilon + 1) \right)^{\frac{\epsilon}{1+\epsilon}} \mu_i.$$

This decomposition can be verified algebraically, and may be more suitable when working with EDE of negative metrics as larger values of subgroup-level M are emphasized (i.e. the most wealth-disadvantaged subgroups bear more weights when computing EDE of M).

Resource allocation for negative metrics

Let M be a negative metric. Typically, negative metrics are rates or probabilities (such as poverty level), and resource allocation problems seek to optimally allocate an amount of new resources in order to lower M . This is an important area of research in cost-effectiveness analysis, and while EDE-factors cannot immediately solve the issue of resource allocation to minimize M , it can offer a measurement on how far a proposed allocation strategy is from a hypothetical scenario where both M and resources can be traded to achieve the minimal EDE M as ϵ varies.

We outline the modifications required to apply techniques in Section 4, using linear production functions for the same reasons as stated in Section 4. Let M_i and p_i be the respective negative metric and production factor allocated to subgroup i . For the amount of resources allocated to subgroups i , we use the formula

$$r_i = \min\{m_i, s_i^2\}$$

where m_i is the maximum capacity of resource the subgroup can accept, and s_i^2 is the (non-negative) amount of resources allocated if below maximum capacity. Let R be the total amount of new resource to be allocated. If M_i is transformed to $M_i^o = M_i - p_i r_i$ after resource reallocation, and $w_{i,\epsilon}^o$ is the respective EDE-factor at inequality aversion parameter ϵ after resource allocation, then

$$M_i^o = \left(-R + \sum_j \frac{M_j}{p_j} \right) \left(\sum_j \frac{1}{p_j} \left(\frac{(w_{i,\epsilon}^o)^{-1} \mu_i}{(w_{j,\epsilon}^o)^{-1} \mu_j} \right)^{\frac{1}{\epsilon}} \right)^{-1}.$$

Consider a hypothetical scenario, where the minimal of

$$EDE^o(\epsilon) = \left(\sum_i (M_i - p_i r_i)^{1+\epsilon} \mu_i \right)^{\frac{1}{1+\epsilon}}$$

is attained subject to the condition

$$\sum_i r_i = R.$$

If $\epsilon = 0$, the solution is clear: allocate maximal resources to subgroups with highest weighted production factor $p_i \mu_i$. If $\epsilon > 0$, Lagrange Multipliers can be applied to

compute the associated resource allocation r_i . If the reallocation satisfies $0 < r_i < m_i$ for all i , then minimality can be achieved when $M_i = \widetilde{M}_{i,\epsilon}$, where

$$\widetilde{M}_{i,\epsilon} := \left(-R + \sum_j \frac{M_j}{p_j} \right) \left(\sum_j \frac{1}{p_j} \left(\frac{p_i \mu_i}{p_j \mu_j} \right)^{\frac{1}{\epsilon}} \right)^{-1}.$$

If $r_i < 0$ or $m_i > r_i$ for some i , then we will need to consider cases where $r_i = 0$ or $r_i = m_i$ and separately perform Lagrange Multipliers to obtain the resource allocation minimizing $EDE^o(\epsilon)$. This results in potentially 3^n different cases to consider, where n is the number of subgroups. However, many cases can be immediately discarded by considering the total number of available resources R (e.g. it is usually not feasible to have $r_i = m_i$ for all i unless resources are abundant). A computational example of resource allocation for negative metrics is presented in the following Section.

6 Example: EDE for potentially avoidable mortality

Ontario, Canada, has the only published province-level elicitation of health-based ϵ in North America, and around the world as far as we know, as of writing this paper. Results of the elicitation [7] show that health-based ϵ in Ontario is bimodally distributed, with two classes of nearly equal size (50.7% and 49.3%) having mean values $\epsilon = 0.339$ and $\epsilon = 7.304$ respectively. Additionally, there are no statistically significant socio-economic or demographic differences between these two classes.

We use our tools developed in Sections 2–5 on data from Public Health Ontario [13]. The dataset considered is stratified by 7 geographical regions, with the health outcome as age-standardized mortality rate from avoidable causes in 2023 (latest available year as of writing this paper). In the remainder of this section, “mortality rate” will abbreviate “age-standardized mortality rate from avoidable causes”.

As an illustration, consider a highly effective hypothetical intervention that reduces mortality rate over a one-year period by 75%. This percentage is informed by the lower values of medication non-adherence literature in Ontario [5, 24], as we assume such an intervention will be administered after a health consultation. As such, we also assume a scale-up of our hypothetical intervention to a coverage level of 87%, coinciding with the estimated percentage of adults in Ontario with access to a regular health provider in 2023 [21]. The resource allocation problem we consider is

the optimal strategy to administer the intervention among the 7 geographic regions such that EDE mortality rate is minimized. One unit of resource will be equal to giving the intervention to one person.

We consider three scenarios: "baseline", where mortality is as listed in [13]; "burden-preference", where our hypothetical intervention is rolled out instantly with a preference to regions with higher mortality rate; and "uniform", where our hypothetical intervention is rolled out instantly at 87% coverage to each of the 7 geographical regions. For simplicity, we assume the health production factor p_i is the product of intervention efficiency $e = 0.75$, region mortality rate M_i , and the inverse population $1/n_i$:


$$p_i = \frac{eM_i}{n_i}.$$

Writing total resources as $R = 0.87n$, where n is the total population of Ontario, this production factor allows us to capture the change in death rate by reallocating r_i resources to region i via $M_i - p_i r_i$, with $r_i \leq n_i$ as the upper bound. A summary of the three scenarios is listed in the following table.


Age-standardized mortality rate per 10,000 from avoidable causes, Ontario (2023)				
Region	Baseline scenario	Uniform scenario	Burden-preference scenario	Population
North West	331.1	115.1	82.8	224,481
North East	290.8	101.1	72.7	548,020
South West	222.1	77.2	55.5	1,696,228
East	194.3	67.5	48.6	1,879,714
Central West	185.6	64.5	46.4	2,841,315
Toronto	158.6	55.1	39.7	2,875,513
Central East	154.8	53.8	89.3	4,277,775

By Equation 6, the EDE mortality rate at ϵ can be calculated by substituting $-\epsilon$ into Atkinson's index (Equation 1). Also let μ_i be the original weights considered by Atkinson (i.e. μ_i is the proportion of population in subgroup i). The weighted mortality rate among the three scenarios are: 182.8 (baseline); 63.5 (uniform); 60.8 (burden-preference). A plot of EDE-factors for baseline and uniform scenarios (left), and burden-preference scenario (right), is given below. Note that the EDE-factors for

baseline and uniform scenarios are the same due to homogeneity (Definition 3.1).




One immediately notices two phenomena. Firstly, a rapidly increasing and disproportionate amount of weight is placed on the Central East region for the burden-preference scenario at $\epsilon < 3$. This implies a faster increase in EDE mortality rate for the burden-preference scenario compared to the uniform scenario at low ϵ , and a tipping point should exist where EDE mortality rate for the burden-preference scenario rises above that for the uniform scenario as ϵ increases. Secondly, EDE-factors for the burden-preference scenario is almost fully weighted by a single region (Central East) at $\epsilon > 6$, compared to two dominant regions (North West and North East) for the uniform scenario. This implies EDE mortality for the burden-preference scenario should have a flatter slope at $\epsilon > 6$ compared to the uniform scenario, suggesting another tipping point in this interval. Indeed, this is the case as the left graph below shows. The right graph below compares EDE mortality rate of all three scenarios as ϵ varies.



The first tipping point is at $\epsilon_1 \approx 1.398$ with $EDE^\dagger(\epsilon_1) \approx 65.3$, and the second tipping point is at $\epsilon_2 \approx 7.680$ with $EDE^\dagger(\epsilon_2) \approx 78.2$. Considering the bimodally distributed ϵ in Ontario, the population with mean $\epsilon = 0.339$ would prefer burden-preference to uniform scenarios (EDE mortality rate: 61.9 for burden-preference; 63.9 for uniform), while the population with mean $\epsilon = 7.304$ would prefer uniform to burden-preference scenarios (EDE mortality rate: 77.8 for burden-preference; 77.4 for uniform).

Following the discussion on resource allocation for negative metrics in Section 5, let us briefly compare the three scenarios considered above with “EDE-optimal”, where resources are allocated to maximize EDE mortality rate at ϵ , in such a way that resources are never allocated to subgroups beyond maximum capacity.



At $\epsilon = 0$, the burden-preference scenario achieves EDE-optimal. At $\epsilon = 0.339$, the burden-preference scenario is still closer to EDE-optimal than the uniform scenario, and to achieve EDE-optimal we should allocate maximal resources to all but Toronto and Central East, distribute remaining resources in the ratio of 45.629% to 54.371% among Central East and Toronto. At $\epsilon = 7.304$, the uniform scenario is closer to EDE-optimal than the burden-preference scenario, and to achieve EDE-optimal we should allocate maximal resources to all but Toronto and Central East, and distribute remaining resources in the ratio of 58.067% to 41.933% among Central East and Toronto, such that 16.5% of the Central East population and 23.4% of the Toronto population receives the hypothetical intervention.

7 Concluding remarks

This paper demonstrated a new decomposition of Atkinson’s index by way of EDE-factors (Definition 3.1). Although many kinds of wealth inequality measures exist [3], we focused on Atkinson’s index for applicability in both health and economics [1, 10].

The technical aspect of our paper contributes a novel way to decompose Atkinson’s index as we used an approach via a L -function viewpoint borrowed from number theory. The original motivation of our work was to seek a general principle behind Sterck’s observed duality between income at $\epsilon = 2$ and poverty at $\epsilon = 0$ [23]. An L -function viewpoint is essential to obtain our duality principle as it is mathematically expressed via a functional equation (Theorem 2.5). This duality is also hinted at in a recent working paper on decomposition of measures [11].

The empirical aspect of our paper applied EDE-factors to resource allocation problems, providing a framework for Robson’s experiments used to elicit health-based Atkinson’s inequality aversion parameter ϵ in various settings. We additionally considered a case study of age-adjusted mortality rate from avoidable causes in Ontario, Canada to illustrate our mathematical methods. This case study is interesting as elicitation of health-based ϵ in [4] tells us that the Ontario population is bimodally distributed, with one population class strongly inequality neutral while the other class has high inequality aversion. Our case study, on resource allocation of a hypothetical intervention, shows that the two different classes may support opposing public health policies.

We end by listing two limitations and directions for future work. Firstly, some important indicators of health do not vary proportionally or inversely to health, such as cholesterol level. While our functional analytic techniques allowed us to extend Atkinson’s index to negative metrics, more work is required to extend Atkinson’s index to other important health metrics. Secondly, there are many other inequality indices that are favored over Atkinson’s index in non-health applications. For instance, Kolm-Pollak’s index is used in environmental justice [16] and transportation problems [6]. As such, it is essential to extend our work to other classes of inequality indices, as our current methods rely on techniques that are specific to Atkinson’s index (such as the Riesz-Fréchet Representation Theorem). Kolm-Pollak’s index can be a promising start, as [16] had showed that this index satisfies an additive counterpart to our functional equation.

Acknowledgements

The author would like to thank R. Scott Braithwaite for his countless insights in inequality measures, and Nao Yamamoto for the numerous helpful criticisms in a preliminary draft of this work.

References

- [1] Atkinson, A. B. (1970). *On the measurement of inequality*. Journal of Economic Theory, 2(3), 244–263.
- [2] Bullen, P. S. (2003). *Handbook of means and their inequalities (2nd ed.)*. Springer.
- [3] De Maio, F. G. (2007). *Income inequality measures*. Journal of Epidemiology & Community Health, 61(10), 849–852.
- [4] Donnell, D., Gao, F., Hughes, J. P., Hanscom, B., Corey, L., Cohen, M. S., Sri-latha Edupuganti, Nyaradzo Mgodi, Rees, H., Baeten, J. M., Gray, G., Bekker, L., Hosseinipour, M. C., Sinead D.-M. (2023). Counterfactual estimation of efficacy against placebo for novel PrEP agents using external trial data: example of injectable cabotegravir and oral PrEP in women. Journal of the International AIDS Society, 26(6).
- [5] Holbrook A. M., Wang M., Lee M., Chen Z., Garcia M., Nguyen L., Ford A., Manji S., Law M. R. (2021). Cost-related medication nonadherence in Canada: a systematic review of prevalence, predictors, and clinical impact. Systematic Reviews, 10(1):11.
- [6] Drew, H., Joshua, M., Daphne, S., Emily, S., Tom, L. (2025). *A scalable optimization approach for equitable facility location: Methodology and transportation applications*. Transportation Research Part B: Methodological, 201, 103319.
- [7] Hurley, J., Mentzakis, E., Walli-Attai, M. (2020). *Inequality aversion in income, health, and income-related health*. Journal of Health Economics, 70, 102276.

- [8] Kim, H.-Y., Bershteyn, A., Russo, R., McGillen, J., Sisti, J., Ko, C., Shaff, J., Newton-Dame, R., Braithwaite, R. S. (2025). *Does prioritization of COVID vaccine distribution to communities with the highest COVID burden reduce health inequity?* Journal of Infection and Public Health, 18(11), 102904.
- [9] Krantz, S. G., Parks, H. R. (2002). *A primer of real analytic functions (2nd ed.)*. Birkhäuser.
- [10] Levy, J. I., Chemerynski, S. M., Tuchmann, J. L. (2006). *Incorporating concepts of inequality and inequity into health benefits analysis*. International Journal for Equity in Health, 5(1), 2.
- [11] Moramarco, D., Sterck, O. (2025). *Who drives inequality?* SSRN working paper, 5220560.
- [12] Pratt, J.W. (1964). *Risk aversion in the small and in the large*. Econometrica, 32(1/2). 122–136.
- [13] Public Health Ontario (2025). *Potentially Avoidable Mortality Snapshot (2014 to 2023)*. <https://www.publichealthontario.ca/en/Data-and-Analysis/Mortality-Overall-Health/Avoidable-Mortality>.
- [14] Robson, M., Asaria, M., Cookson, R., Tsuchiya, A., Ali, S. (2016). *Eliciting the level of health inequality aversion in England*. Health Economics, 26(10), 1328–1334.
- [15] Robson, M., O'Donnell, O., Tom Van Ourti. (2024). *Aversion to health inequality – Pure, income-related and income-caused*. Journal of Health Economics, 94, 102856.
- [16] Sheriff, G., Maguire, KB. (2020). *Health risk, inequality indexes, and environmental justice*. Risk Analysis, 40(12), 2661–2674.
- [17] Shorrocks, A. F. (1980). *The class of additively decomposable inequality measures*. Econometrica, 48, 613–625.
- [18] Shorrocks, A. F. (1984). *Inequality decomposition by population subgroups*. Econometrica, 52(6), 1369.

- [19] Slejko, J. F., Ricci, S., dosReis, S., Cookson, R., Kowal, S. (2025). *Health inequality aversion in the United States*. Value in Health, 29(1), 129–138.
- [20] Spinelli, M. A., Bisom-Rapp, E., Heise, M. J., Camp, C., Appa, A., Liu, A. Y., Sassaman, K., Shiels, M., Mayorga-Munoz, F., Chimezie, A., Nguyen, J., Oskarsson, J., Gandhi, M. (2024). High Retention and Adherence With Rapid Long-acting Injectable Preexposure Prophylaxis Implementation in an Urban Safety Net Clinic Population. Clinical Infectious Diseases.
- [21] Statistics Canada (2025). *Access to health care*. <https://www150.statcan.gc.ca/n1/pub/82-570-x/2024001/section4-eng.htm>.
- [22] Stein, E. M., Shakarchi, R. (2011). *Functional analysis: Introduction to further topics in analysis (Princeton Lectures in Analysis, Vol. 4)*. Princeton University Press.
- [23] Sterck, O. (2024). *Poverty without poverty line*. SSRN working paper, 4785458.
- [24] Stewart S. L., Baiden P. (2013). An exploratory study of the factors associated with medication nonadherence among youth in adult mental health facilities in Ontario, Canada. Psychiatry Research, May 30;207(3):212-7.

A Analogies of our analysis with L -functions

The following table summarizes how L -function analogies from number theory was used to guide our analysis in Section 2.

Atkinson’s index	L -functions analog
Sum decomposition (Theorem 2.1)	L -functions as sums at $\Re(s) > 1$
Values $f_{i,\epsilon}$ (Corollary 2.2; Theorem 2.3)	Twisting by Hecke characters
Analytic continuation at $\epsilon \neq 1$ (Theorem 2.4)	Analytic continuation at $s \neq 1$
Functional equation (Theorem 2.5)	Functional equation for L -functions

We believe such a framework to compare between mathematics and health economics can also be useful to analyze other indices of potential value. Although we do

not have an answer at present, and left for future work, we mention another index of potential value here. Kolm-Pollak's index is an absolute counterpart of Atkinson's index (the latter derived from relative constant risk aversion [1]). Recent work [16] argued that Kolm-Pollak's index may be more suitable for health studies where absolute differences are economically meaningful (e.g. financial risk protection), and additionally derived a Mirror Property for Kolm-Pollak's index. We hypothesize that this Mirror Property is an analog of the symmetric property of Kloosterman exponential sums, or certain trace formulas in automorphic functions.

B Proofs of statements in Section 2

Proof of Theorem 2.1. Let X be a countable measure space with discrete probability measure μ , and let F be an injective real-valued positive Lebesgue-measurable function on X . Then the p -norm of F is simply

$$\|F\|_p = \left(\sum_i F_i^p \mu_i \right)^{\frac{1}{p}}.$$

In our case where $F_i = 1/H_i$,

$$\|F\|_p = \left(\sum_i H_i^{-p} \mu_i \right)^{\frac{1}{p}}.$$

By the Riesz-Fréchet Representation Theorem for L^p -spaces [22, Chapter 1], there exist non-negative values f_i such that

$$\|F\|_p = \max_{\substack{\sum_i f_i^q \mu_i \leq 1 \\ f_i \geq 0}} \left\{ \sum_i \frac{1}{H_i} f_i \mu_i \right\},$$

where q is the number such that $1/p + 1/q = 1$. If we can find values f_i for this equality to hold, then we are done as the left-hand side $\|F\|_p$ equals $EDE_H(\epsilon)^{-1}$ after recalling $p = \epsilon - 1$.

Following this discussion, we need to solve an optimization problem: Find numbers

$f_i \geq 0$ that maximizes

$$\sum_i \frac{1}{H_i} f_i \mu_i \quad (7)$$

subject to the condition

$$\sum_i f_i^q \mu_i = l, \quad 0 \leq l \leq 1. \quad (8)$$

Clearly, f_i cannot be simultaneously zero for all i . By using Lagrange Multipliers, there exist a constant λ_l depending on l such that

$$\frac{1}{H_i} \mu_i = \lambda_l q f_i^{q-1} \mu_i.$$

Hence

$$f_i = \left(\frac{1}{\lambda_l q H_i} \right)^{\frac{1}{q-1}}.$$

Substituting f_i to Equations 7 and 8 gives

$$\sum_i \frac{1}{H_i} f_i \mu_i = \lambda_l^{-\frac{1}{q-1}} \sum_i \left(\frac{1}{q H_i^q} \right)^{\frac{1}{q-1}} \mu_i$$

and

$$\lambda_l^{-\frac{1}{q-1}} = l^{\frac{1}{q-1}} \left(\sum_i \left(\frac{1}{q H_i} \right)^{\frac{q}{q-1}} \mu_i \right)^{-\frac{1}{q}}$$

Since q, H_i, μ_i are all known constants, to maximize Equation 7, we will need to maximize $\lambda_l^{-\frac{1}{q-1}}$, which requires maximizing $l^{\frac{1}{q-1}}$, and this last expression is an increasing function on l as $q - 1 > 0$. Therefore, necessarily $l = 1$ for our optimization problem.

In summary, for each f_i ,

$$\begin{aligned} f_i &= \left(\frac{1}{\lambda_1 q H_i} \right)^{\frac{1}{q-1}} \\ &= \left(\frac{1}{q H_i} \right)^{\frac{1}{q-1}} \left(\sum_j \left(\frac{1}{q H_j} \right)^{\frac{q}{q-1}} \mu_j \right)^{-\frac{1}{q}} \\ &= \left(\sum_j \left(\frac{H_i}{H_j} \right)^{\frac{q}{q-1}} \mu_j \right)^{-\frac{1}{q}}. \end{aligned} \quad (9)$$

These f_i 's are exactly the $f_{i,\epsilon}$ we seek. $\square$

Proof of Corollary 2.2. This is an algebraic manipulation of Equation 9 to rewrite it into two different ways. $\square$

Proof of Theorem 2.3. By manipulating Equation 9,

$$f_i = \left(\frac{EDE_H(k)}{H_i} \right)^{-\frac{1}{q-1}}, \quad k = 2 + \frac{1}{q-1}.$$

Therefore, we can view both f_i and

$$H(q) := \sum_i \frac{1}{H_i} f_i \mu_i$$

as functions in q . To prove the Theorem, it suffices to show that $H(q)$ is monotone decreasing in q . A computation tells us that the derivative with respect to q is

$$\begin{aligned} \frac{d}{dq} H(q) &= \sum_i \frac{1}{H_i} \frac{df_i}{dq} \mu_i \\ &= \frac{1}{(q-1)^2} \sum_i \frac{f_i \mu_i}{H_i} \ln \left(\frac{EDE_H(k)}{H_i} \right) + \frac{1}{(q-1)^3} \sum_i \frac{f_i \mu_i}{H_i} \frac{EDE'_H(k)}{EDE_H(k)} \\ &= -\frac{1}{q-1} \sum_i \frac{f_i \mu_i}{H_i} \ln(f_i) + \frac{1}{(q-1)^3} \sum_i \frac{f_i \mu_i}{H_i} \frac{EDE'_H(k)}{EDE_H(k)}. \end{aligned} \tag{10}$$

Here, $EDE'_H(k)$ is the derivative of $EDE_H(k)$ with respect to k and not q .

We now need to show that the first and second term of Equation 10 are both negative. The second term is negative as $EDE'_H(k)$ is the only part of the term that is negative. For the first term, note that

$$\sum_i \frac{f_i \mu_i}{H_i} \ln(f_i) = \ln \left(\prod_i f_i^{\frac{f_i \mu_i}{H_i}} \right)$$

so we are reduced to showing that the product inside the logarithm is at least 1. Using the weighted power mean inequality, more specifically the weighted GM-HM

inequality [2, Chapter 3],

$$\prod_i f_i^{\frac{f_i \mu_i}{H_i}} \geq \left(\left(\frac{\sum_i \frac{f_i \mu_i}{H_i} f_i^{-1}}{\sum_i \frac{f_i \mu_i}{H_i}} \right)^{-1} \right)^{\sum_i \frac{f_i \mu_i}{H_i}}. \quad (11)$$

We now make the observation that

$$\sum_i \frac{f_i \mu_i}{H_i} f_i^{-1} = \sum_i \frac{1}{H_i} \cdot 1 \cdot \mu_i$$

so by the first condition in the statement of the Theorem,

$$\sum_i \frac{f_i \mu_i}{H_i} f_i^{-1} \leq \sum_i \frac{f_i \mu_i}{H_i}.$$

Hence the fraction in the right hand side of Equation 11 is at most 1, implying

$$\left(\left(\frac{\sum_i \frac{f_i \mu_i}{H_i} f_i^{-1}}{\sum_i \frac{f_i \mu_i}{H_i}} \right)^{-1} \right)^{\sum_i \frac{f_i \mu_i}{H_i}} \geq 1$$

and we are done. $\square$

Proof of Theorem 2.4. Note that both $L(H_i, \epsilon)$ and $EDE_H(\epsilon)^{-1}$ are analytic functions defined on the interval $(-\infty, 1) \cup (1, \infty)$. As $L(H_i, \epsilon) = EDE_H(\epsilon)^{-1}$ on $(2, \infty)$, the Identity Theorem [9, Chapter 1] implies they must also be equal on $(1, \infty)$.

We now apply the Identity Theorem on $(-\infty, 1)$ by showing that $L(H_i, \epsilon)$ and $EDE_H(\epsilon)^{-1}$ agree on a sequence converging to the limit point 0. In particular, we show that $L(H_i, 1/n)$ equals $EDE_H(1/n)^{-1}$ for all positive integers $n \geq 2$. By Corol-

lary 2.2,

$$\begin{aligned}
L\left(H_i, \frac{1}{n}\right) &= \sum_i \frac{1}{H_i} \left(\sum_j \left(\frac{H_j}{H_i} \right)^{1-\frac{1}{n}} \mu_j \right)^{-1+\frac{1}{1-\frac{1}{n}}} \mu_i \\
&= \left(\sum_j H_j^{1-\frac{1}{n}} \mu_j \right)^{-(1+\frac{n}{n-1})} \left(\sum_i H_i^{1-\frac{1}{n}} \mu_i \right) \\
&= EDE \left(\frac{1}{n} \right)^{-1},
\end{aligned}$$

as desired. □

Proof of Theorem 2.5. If $\epsilon > 0$, a calculation reveals

$$\begin{aligned}
L(H_i^{-1}, 2 + \epsilon) &= \sum_j H_j \left(\sum_i \left(\frac{H_j}{H_i} \right)^{-(1+\epsilon)} \mu_i \right)^{-(1-\frac{1}{1+\epsilon})} \mu_j \\
&= \left(\sum_j H_j^{1+\epsilon} \mu_j \right) \left(\sum_i H_i^{1+\epsilon} \mu_i \right)^{-(1-\frac{1}{1+\epsilon})} \\
&= \left(\sum_i H_i^{1+\epsilon} \mu_i \right)^{\frac{1}{1+\epsilon}} \\
&= EDE_H(-\epsilon) \\
&= L(H, -\epsilon)^{-1}
\end{aligned}$$

where the last equality is due to Theorem 2.1. Finally, the condition $\epsilon > 0$ can be dropped by the Identity Theorem. □

C Proofs of statements in Section 3

Proof of Proposition 3.3. To show that the EDE-factors satisfy population homogeneity, suppose each subgroup i is replicated n times $(i, 1), \dots, (i, n)$ and each replication is weighted $\omega_1, \dots, \omega_n$, with the ω_k 's summing to 1. Then $H_{(i,1)} = \dots = H_{(i,n)}$, and each subgroup (i, η) is weighted $\mu_i \omega_\eta$ in Atkinson's index. Therefore, its corresponding

EDE-factor is

$$\begin{aligned}
w_{(i,\eta)}(\epsilon) &= \left(\sum_{\alpha} \sum_j \left(\frac{H_{(j,\alpha)}}{H_{(i,\eta)}} \right)^{-\epsilon} \mu_j \omega_{\alpha} \right)^{-1} \mu_i \omega_{\eta} \\
&= \left(\sum_j \left(\frac{H_{(j,\alpha)}}{H_{(i,\eta)}} \right)^{-\epsilon} \mu_j \sum_{\alpha} \omega_{\alpha} \right)^{-1} \mu_i \omega_{\eta} \\
&= \omega_{\eta} \cdot w_i(\epsilon).
\end{aligned}$$

As the weights $w_i(\epsilon)$ are q -analogs of the expression $f_{i,\epsilon} \mu_i$,

$$\begin{aligned}
\left(\sum_{\alpha} \sum_i \frac{1}{H_{(i,\alpha)}} f_{(i,\alpha),\epsilon} \mu_i \omega_{\alpha} \right)^{-1} &= \left(\sum_i \frac{1}{H_i} f_{i,\epsilon} \mu_i \sum_{\alpha} \omega_{\alpha} \right)^{-1} \\
&= \left(\sum_i \frac{1}{H_i} f_{i,\epsilon} \mu_i \right)^{-1} \\
&= EDE_H(\epsilon)
\end{aligned}$$

which proves population homogeneity. $\square$

Proof of Proposition 3.4. Let $\{H_k^*\}_k$ be the wealth profile such that $H_k^* = H_k$ for $k \neq i, j$, with $H_i^* = H_i - \Delta$ and $H_j^* = H_j + \Delta^*$ where Δ, Δ^* are as defined above. Let $EDE^*(\epsilon)$ be Atkinson's index calculated with the wealth profile $\{H_k^*\}_k$. By Theorem 2.1, there exists $f_{i,\epsilon}, f_{i,\epsilon}^*$ satisfying the conditions of that Theorem such that

$$EDE_H(\epsilon) = \left(\sum_i \frac{1}{H_i} f_{i,\epsilon} \mu_i \right)^{-1}$$

and

$$EDE^*(\epsilon) = \left(\sum_i \frac{1}{H_i^*} f_{i,\epsilon}^* \mu_i \right)^{-1}.$$

We need to show that $EDE^*(\epsilon) \geq EDE_H(\epsilon)$. Note that

$$\sum_i \frac{1}{H_i^*} f_{i,\epsilon}^* \mu_i = \Delta \mu_i \left(\frac{f_i^*}{H_i H_i^*} - \frac{f_j^*}{H_j H_j^*} \right) + \sum_i \frac{1}{H_i} f_{i,\epsilon}^* \mu_i.$$

The summation on the right satisfies

$$\sum_i \frac{1}{H_i} f_{i,\epsilon}^* \mu_i \leq \sum_i \frac{1}{H_i} f_{i,\epsilon} \mu_i$$

by Theorem 2.1. The term on the left satisfies

$$\begin{aligned} \frac{f_i^*}{H_i H_i^*} - \frac{f_j^*}{H_j H_j^*} &= EDE^*(\epsilon)^{\epsilon-2} \left(\frac{1}{H_i (H_i^*)^{\epsilon-1}} - \frac{1}{H_j (H_j^*)^{\epsilon-1}} \right) \\ &< 0 \end{aligned}$$

where the equality is by definition of f_i^* (Corollary 2.1) and the inequality is because $H_i > H_j$ and $H_i^* > H_j^*$. Therefore

$$\sum_i \frac{1}{H_i^*} f_{i,\epsilon}^* \mu_i \leq \sum_i \frac{1}{H_i} f_{i,\epsilon} \mu_i$$

implying $EDE^*(\epsilon) \geq EDE_H(\epsilon)$. □

D Proofs of statements in Section 4

We prove the statements in Section 4 in more generality via the modified Cobb-Douglas production function $H_i = p_i r_i^\alpha$, where $\alpha \in (0, 1]$ is the resource elasticity variable. At present, our arguments to obtain explicit formulas do not apply if subgroups do not present similar diminishing marginal utilities (i.e. $H_i = p_i r_i^{\alpha_i}$ where α_i is not necessarily equal to α). This limitation of our work will be investigated in the future.

Assume the sum of all available resources equal R . Then we seek to maximize

$$EDE_H(\epsilon) = \left(\sum_i (p_i r_i^\alpha)^{1-\epsilon} \mu_i \right)^{\frac{1}{1-\epsilon}}$$

subject to the condition

$$\sum_i r_i = R.$$

By Lagrange Multipliers, there exists some $\lambda > 0$ such that

$$\partial_{r_i} EDE_H(\epsilon) = \lambda.$$

Taking the derivative on the left hand side, we get

$$d_{i,\epsilon} r_i^{\alpha(-\epsilon+1)-1} = \lambda, \quad \text{where} \quad d_{i,\epsilon} = \alpha p_i^{1-\epsilon} \mu_i \left(\sum_i H_i^{1-\epsilon} \mu_i \right)^{\frac{1}{1-\epsilon}-1}.$$

Therefore

$$r_i = \left(\frac{\lambda}{d_{i,\epsilon}} \right)^{\frac{1}{\alpha(-\epsilon+1)-1}}. \quad (12)$$

Summing across all r_i gives us

$$R = \sum_j \left(\frac{\lambda}{d_{j,\epsilon}} \right)^{\frac{1}{\alpha(-\epsilon+1)-1}},$$

implying

$$\lambda^{\frac{1}{\alpha(-\epsilon+1)-1}} = R \left(\sum_j \left(\frac{1}{d_{j,\epsilon}} \right)^{\frac{1}{\alpha(-\epsilon+1)-1}} \right)^{-1}.$$

Substituting the above equality into Equation 12,

$$r_i = R \left(\sum_j \left(\frac{d_{i,\epsilon}}{d_{j,\epsilon}} \right)^{\frac{1}{\alpha(-\epsilon+1)-1}} \right)^{-1}$$

implying

$$\begin{aligned} H_i &= p_i R^\alpha \left(\sum_j \left(\frac{d_{i,\epsilon}}{d_{j,\epsilon}} \right)^{\frac{1}{\alpha(-\epsilon+1)-1}} \right)^{-\alpha} \\ &= R^\alpha \left(\sum_j \frac{p_i^{-\frac{1}{\alpha} + \frac{\epsilon}{\alpha(\epsilon-1)+1}}}{p_j^{\frac{\epsilon}{\alpha(\epsilon-1)+1}}} \left(\frac{p_j \mu_j}{p_i \mu_i} \right)^{\frac{1}{\alpha(\epsilon-1)+1}} \right)^{-\alpha}. \end{aligned} \quad (13)$$

Note that substituting $\alpha = 1$ into Equation 13 gives Equation 3.

Now, let $w_i = w_i(\epsilon)$ be EDE-factors. Then we show that we can rewrite the

subgroup welfare levels H_i in terms of w_i . By Definition 3.1,

$$w_i = H_i^{-\epsilon} \mu_i e_\epsilon^{-1}, \quad \text{where} \quad e_\epsilon = \sum_j H_j^{-\epsilon} \mu_j.$$

Therefore

$$H_i = \left(\frac{e_\epsilon w_i}{\mu_i} \right)^{-\frac{1}{\epsilon}}. \quad (14)$$

Since $H_i = p_i r_i^\alpha$, this implies

$$r_i = \left(\frac{1}{p_i} \left(\frac{e_\epsilon w_i}{\mu_i} \right)^{-\frac{1}{\epsilon}} \right)^{\frac{1}{\alpha}}.$$

Summing across all r_i and separating out the e_ϵ term gives

$$e_\epsilon^{-\frac{1}{\epsilon\alpha}} = R \left(\sum_j \left(\frac{1}{p_j} \left(\frac{w_j}{\mu_j} \right)^{-\frac{1}{\epsilon}} \right)^{\frac{1}{\alpha}} \right)^{-1}.$$

Substituting the above equality into Equation 14,

$$H_i = R^\alpha \left(\sum_j \left(\frac{1}{p_j} \left(\frac{w_j^{-1} \mu_j}{w_i^{-1} \mu_i} \right)^{\frac{1}{\epsilon}} \right)^{\frac{1}{\alpha}} \right)^{-\alpha}. \quad (15)$$

Note that substituting $\alpha = 1$ into Equation 13 gives Equation 4.

Proposition D.1. *There exists a unique set of EDE-factors w_i that maximizes $EDE_H(\epsilon)$.*

Proof. To show uniqueness, suppose η_i is another set of EDE-factors maximizing $EDE_H(\epsilon)$. By Equation 13, both sets of EDE-factors admit the same subgroup welfare levels H_i for all i . In particular, using Equation 15,

$$R^\alpha \left(\sum_j \left(\frac{1}{p_j} \left(\frac{w_j^{-1} \mu_j}{w_i^{-1} \mu_i} \right)^{\frac{1}{\epsilon}} \right)^{\frac{1}{\alpha}} \right)^{-\alpha} = R^\alpha \left(\sum_j \left(\frac{1}{p_j} \left(\frac{\eta_j^{-1} \mu_j}{\eta_i^{-1} \mu_i} \right)^{\frac{1}{\epsilon}} \right)^{\frac{1}{\alpha}} \right)^{-\alpha},$$

implying the existence of some constant c such that

$$\frac{w_i}{\eta_i} = c$$

for all i . This implies $c = 1$ as w_i and η_i both sum to 1, showing uniqueness.

To show existence, we can equate Equations 13 and 15 term-wise:

$$\frac{p_i^{-\frac{1}{\alpha} + \frac{\epsilon}{\alpha(\epsilon-1)+1}}}{p_j^{\frac{\epsilon}{\alpha(\epsilon-1)+1}}} \left(\frac{p_j \mu_j}{p_i \mu_i} \right)^{\frac{1}{\alpha(\epsilon-1)+1}} = \left(\frac{1}{p_j} \left(\frac{w_j^{-1} \mu_j}{w_i^{-1} \mu_i} \right)^{\frac{1}{\epsilon}} \right)^{\frac{1}{\alpha}}.$$

Writing the left side of the above equation as $l_{i,j,\epsilon}$, a rearrangement gives

$$\frac{w_i}{w_j} = \frac{\mu_i}{\mu_j} p_j^{\epsilon} l_{i,j,\epsilon}^{\epsilon \alpha}.$$

Finally, summing across i , and taking the reciprocal, gives

$$w_j = \left(\sum_i \frac{\mu_i}{\mu_j} p_j^{\epsilon} l_{i,j,\epsilon}^{\epsilon \alpha} \right)^{-1} \quad (16)$$

as desired. $\square$

Note that substituting $\alpha = 1$ into Equation 16, and switching the indices from j to i , gives Equation 5.

We now make some comments if we consider the more general case of modified Cobb-Douglas production functions $H_i = p_i r_i^{\alpha_i}$, where $\alpha_i \in (0, 1]$ are resource elasticity variables (and not necessarily all equal to some α). If we seek to maximize

$$EDE_H(\epsilon) = \left(\sum_i (p_i r_i^{\alpha_i})^{1-\epsilon} \mu_i \right)^{\frac{1}{1-\epsilon}}$$

subject to the condition

$$\sum_i r_i = R,$$

a similar argument via Lagrange Multipliers gives the formulas

$$r_i = \left(\frac{\lambda}{d_{i,\epsilon}} \right)^{\frac{1}{\alpha_i(-\epsilon+1)-1}} \quad \text{and} \quad R = \sum_j \left(\frac{\lambda}{d_{j,\epsilon}} \right)^{\frac{1}{\alpha_j(-\epsilon+1)-1}}.$$

Since the right hand side is a monotonic (decreasing) function in λ , there exists a unique λ satisfying the optimization problem.

On the other hand, letting $w_i = w_i(\epsilon)$ be EDE-factors, a similar manipulation as before gives

$$r_i = \left(\frac{1}{p_i} \left(\frac{e_\epsilon w_i}{\mu_i} \right)^{-\frac{1}{\epsilon}} \right)^{\frac{1}{\alpha_i}} \quad \text{and} \quad R = \sum_j \left(\frac{1}{p_j} \left(\frac{e_\epsilon w_j}{\mu_j} \right)^{-\frac{1}{\epsilon}} \right)^{\frac{1}{\alpha_j}}$$

Since the right hand side is a monotonic (decreasing) function in λ , there exists unique w_i 's if we fix the r_i 's. This implies Proposition D.1 still holds even in the more general case where resource elasticity variables are different across subgroups, even though we cannot obtain explicit formulas for r_i .