


$$\frac{1}{4}(1-\alpha)S + \frac{\epsilon}{2}\sigma T_s^4 = \sigma T_s^4$$

$$\frac{1}{4}(1-\alpha)S = (1-\frac{\epsilon}{2})\sigma T_s^4$$

$$T_s = \left(\frac{(1-\alpha)S}{4(1-\frac{\epsilon}{2})\sigma}\right)^{\frac{1}{4}}$$




$$\frac{1}{4}(1-\alpha)S + \frac{\epsilon}{2}\sigma T_s^4 = \sigma T_s^4$$

$$\frac{1}{4}(1-\alpha)S=(1-\frac{\epsilon}{2})\sigma T_s^4$$

$$T_s = \left(\frac{(1-\alpha)S}{4(1-\frac{\varepsilon}{2})\sigma}\right)^{\frac{1}{4}}$$

1. Mars has an insolation of  $S = 586 \text{ W m}^{-2}$  and an albedo of  $\alpha = 0.25$ . It has a very thin atmosphere, so we can approximate  $\varepsilon = 0$ . Remember that  $\sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$ . Use the one-layer model to estimate the surface temperature of Mars, and compare to the actual value of 210 K.



$$\frac{1}{4}(1-\alpha)S + \frac{\epsilon}{2}\sigma T_s^4 = \sigma T_s^4$$

$$\frac{1}{4}(1-\alpha)S = (1-\frac{\epsilon}{2})\sigma T_s^4$$

$$T_s = \left(\frac{(1-\alpha)S}{4(1-\frac{\epsilon}{2})\sigma}\right)^{\frac{1}{4}}$$

1. Mars has an insolation of  $S = 586 \text{ W m}^{-2}$  and an albedo of  $\alpha = 0.25$ . It has a very thin atmosphere, so we can approximate  $\varepsilon = 0$ . Remember that  $\sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$ . Use the one-layer model to estimate the surface temperature of Mars, and compare to the actual value of 210 K.

$$T_s = \left(\frac{(1-0.25)586~W~m^{-2}}{4\times5.67\times10^{-8}~W~m^{-2}~K^{-4}}\right)^{\frac{1}{4}} = 210~K$$

This is actually a 0-layer model since we assume there is no atmosphere!