

MA 1521  
Tutorial 8 Solutions


1. The volume is given by the double integral

$$V = \iint_D f(x, y) dA$$

where  $D$  is the region bounded by the parabola  $y = 4 - x^2$  and straight line  $y = 3x$  and  $f(x, y)$  is the function whose graph is the plane  $x - z + 4 = 0$ .

Writing the equation of the plane as  $z = x + 4$ , we get the function  $f(x, y) = x + 4$ .

A rough sketch of the region  $D$  is shown below:



$D$  can be regarded as type A region

$$D : \quad 3x \leq y \leq 4 - x^2, \quad -4 \leq x \leq 1.$$

(The two limits  $-4$  and  $1$  of  $x$  are obtained by solving the two equation  $y = 3x$  and  $y = 4 - x^2$ .)

Hence

$$V = \int_{-4}^1 \int_{3x}^{4-x^2} (x+4) dy dx = \int_{-4}^1 (x+4)(4-x^2-3x) dx = \left[ 16x - 4x^2 - \frac{7}{3}x^3 - \frac{1}{4}x^4 \right]_{-4}^1 = \frac{625}{12}$$

2. Let  $z = \sqrt{2^2 - x^2 - y^2}$ . Then  
 $z_x = -x(4 - x^2 - y^2)^{-1/2}$  and  $z_y = -y(4 - x^2 - y^2)^{-1/2}$ .

Substitute  $z = 1$  into  $x^2 + y^2 + z^2 = 4$  gives

$$x^2 + y^2 + 1 = 4 \quad \Rightarrow \quad x^2 + y^2 = 3$$

which is the equation of a circle of radius  $\sqrt{3}$ .

This means the plane  $z = 1$  intersects the sphere at a circle of radius  $\sqrt{3}$ .

Hence the projected region  $R$  of the part of the sphere is a disk of radius  $\sqrt{3}$ .

In polar coordinates, this is given by

$$0 \leq r \leq \sqrt{3}, \quad 0 \leq \theta \leq 2\pi.$$

Thus,

$$\begin{aligned}
A(S) &= \iint_R \sqrt{\frac{x^2 + y^2}{4 - x^2 - y^2} + 1} dA = \int_0^{2\pi} \int_0^{\sqrt{3}} \left( \frac{r^2}{4 - r^2} + 1 \right)^{\frac{1}{2}} r dr d\theta \\
&= \int_0^{2\pi} \int_0^{\sqrt{3}} 2r(4 - r^2)^{-\frac{1}{2}} dr d\theta = \int_0^{2\pi} d\theta \left[ -2(4 - r^2)^{\frac{1}{2}} \right]_{r=0}^{r=\sqrt{3}} \\
&= (2\pi)[-2[(4 - 3)^{\frac{1}{2}} + 2[(4)^{\frac{1}{2}}] = 4\pi.
\end{aligned}$$

3.  $\frac{\partial z}{\partial x} = \frac{x}{\sqrt{x^2 + y^2}}$ ,  $\frac{\partial z}{\partial y} = \frac{y}{\sqrt{x^2 + y^2}}$ . Therefore

$$\sqrt{\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 + 1} = \sqrt{\frac{x^2}{x^2 + y^2} + \frac{y^2}{x^2 + y^2} + 1} = \sqrt{2}.$$

Note that  $D$  is given as a Type A domain. The surface area is then given by

$$\int \int_D \sqrt{2} dx dy = \int_{-1}^2 \left( \int_{x^2}^{x+2} \sqrt{2} dy \right) dx = \frac{9}{2} \sqrt{2}.$$

4. Write the equation of the saddle surface as  $z = \frac{1}{a}x^2 - \frac{1}{a}y^2$ , we have

$$z_x = \frac{2x}{a} \text{ and } z_y = \frac{-2y}{a}.$$

Let  $D$  denote the bounded circular region on the  $xy$ -plane bounded by the circle  $x^2 + y^2 = a^2$ .

Then the required surface area is given by

$$\begin{aligned}
S &= \int \int_D \sqrt{1 + z_x^2 + z_y^2} dx dy \\
&= \frac{1}{a} \int \int_D \sqrt{a^2 + 4x^2 + 4y^2} dx dy \\
&= \frac{1}{a} \int_0^{2\pi} \int_0^a \sqrt{a^2 + 4r^2} r dr d\theta \\
&= \frac{2\pi}{a} \int_0^a \sqrt{a^2 + 4r^2} d\left(\frac{a^2 + 4r^2}{8}\right) \\
&= \frac{\pi}{6a} \left[ (a^2 + 4r^2)^{\frac{3}{2}} \right]_0^a \\
&= \frac{\pi a^2}{6} \left( 5^{\frac{3}{2}} - 1 \right)
\end{aligned}$$