NM4QM 2026 · HANDS-ON SESSION · 45 MIN

Spin and anomalous Hall effects with WannierBerri and GPAW

Symmetry-adapted Wannier functions, spin–orbit coupling on demand, the spin Hall effect of graphene and the anomalous Hall effect of the altermagnet MnTe



Stepan Tsirkin

Yaroslav Zhumagulov
Chair of Computational Physics of Condensed Matter (C3MP), EPFL, Lausanne
DIPC, Donostia-San Sebastián · 13–16 October 2026

THE STORY

Two puzzles, two acts

0–8 min	8–20	20–37	37–45
Setup and ideas	Act 1: graphene	Act 2: MnTe	Your material, questions

TWO PUZZLES

Graphene: spin–orbit coupling opens a gap, a quantum spin Hall insulator

MnTe: no net magnetisation, yet an anomalous Hall effect

WHAT YOU WILL DO

Wannier functions from GPAW, SOC on demand, Hall conductivities

SETUP

Before we start: one check

```
cd wannier-berri-demo && conda activate wbdemo
python setup/check_setup.py        # should end with "Ready."
cd 01_graphene
python 1_dft.py                    # 10 s
jupyter lab graphene.ipynb
```


No environment: `bash setup/setup.sh`

DFT fails: `tar -xzf dft-files.tar.gz`

Fell behind: `bash run_all.sh`


Numbers on slides: checkpoints

One Python pipeline, GPAW to conductivities



Each box is one call; the shaded step needs no new DFT and no new Wannierisation

Hall, anomalous Hall and spin Hall effects



magnetic field B

Lorentz force

magnetisation M, no B

Berry curvature of the bands

no B, no M

spin current, no charge imbalance

Act 1: a quantised spin Hall effect in graphene · Act 2: an anomalous Hall effect in MnTe, without M

Berry curvature needs millions of k-points; DFT gives tens

Intrinsic anomalous Hall conductivity:

$$\sigma_{xy} = -\frac{e^2}{\hbar} \int_{\mathrm{BZ}} \frac{d^3k}{(2\pi)^3} \sum_n f(\varepsilon_{n\mathbf{k}})\, \Omega_n^z(\mathbf{k})$$

$$\Omega_n^z = -2\,\mathrm{Im} \sum_{m \neq n} \frac{v_{nm}^x\, v_{mn}^y}{(\varepsilon_n - \varepsilon_m)^2}$$

f: occupation, ε_n: band energy, $v_{nm}^a = \langle u_n | \partial_{k_a} H | u_m \rangle$. Ω_n^z is large and narrow wherever two bands come close.

Wannier functions give *H*(**k**) at any **k**:


$$H_{ij}(\mathbf{k}) = \sum_{\mathbf{R}} e^{i\mathbf{k}\cdot\mathbf{R}} \langle w_{i\mathbf{0}} | \hat{H} | w_{j\mathbf{R}} \rangle$$

$w_{j\mathbf{R}}$: Wannier function j in the cell at $\mathbf{R}$; a few hundred $\mathbf{R}$ are enough.

Symmetry-adapted Wannier functions: DFT on irreducible k only

Each Wannier function transforms like its orbital

Degeneracies come out exact



6 × 6 grid: 36 points, 7 irreducible (blue)

METHODS

Wannierisation: WannierBerri vs Wannier90

$$\Omega = \sum_n \left[\langle w_n | r^2 | w_n \rangle - \left| \langle w_n | \mathbf{r} | w_n \rangle \right|^2 \right] = \Omega_{\mathrm{I}} + \tilde{\Omega}$$

Ω: spread · Ω_{I}: fixed by the subspace (disentanglement) · $\tilde{\Omega}$: set by the gauge $U(\mathbf{k})$ (localisation)

	Wannier90	**WannierBerri, this demo**
Input	.amn, .mmn, .eig files from a DFT interface	computed in Python from the GPAW wave functions
k-points	full grid; irreducible ones with site_symmetry	irreducible ones
Algorithm	disentangle first, then minimise $\tilde{\Omega}$ by steepest descent or conjugate gradients	one loop: a disentanglement step and a local gauge update per iteration, no gradient descent
Symmetry	site_symmetry with a .dmn file from pw2wannier90, not with a frozen window	from irrep, for any code it reads; frozen window allowed
SOC	spinor DFT run with SOC	collinear run, SOC added afterwards; spin ↓ from symmetry in altermagnets

Marzari, Vanderbilt, PRB 56, 12847 (1997) · Souza, Marzari, Vanderbilt, PRB 65, 035109 (2001) · Sakuma, PRB 87, 235109 (2013) · Pizzi et al., JPCM 32, 165902 (2020)

Post-processing: WannierBerri vs postw90

$$H(\mathbf{K}+\boldsymbol{\kappa}) = \sum_{\mathbf{R}} e^{i\boldsymbol{\kappa}\cdot\mathbf{R}} \left[e^{i\mathbf{K}\cdot\mathbf{R}}\, H(\mathbf{R}) \right]$$

$\boldsymbol{\kappa}$: points of a coarse FFT grid · $\mathbf{K}$: shifts of that grid; one FFT per shift, the shifts together give the dense grid


	postw90	**WannierBerri**
$H(\mathbf{R}) \to H(\mathbf{k})$	sum over **R** at every k, cost $\propto N_{\mathbf{R}}$	FFT over sub-grids, cost $\propto \log N_{\mathrm{FFT}}$ per k
Symmetry	full grid	irreducible k-points only, result symmetrised
Hot spots	finer mesh where |Ω| exceeds a threshold	recursive adaptive refinement
Fermi levels	cost grows with their number	many at almost no extra cost
SOC axis	fixed by the DFT run	`set_soc_axis` : any axis and strength

Up to three orders of magnitude faster on dense grids

Wang, Yates, Souza, Vanderbilt, PRB 74, 195118 (2006) · Tsirkin, npj Comput. Mater. 7, 33 (2021) · Pizzi et al., JPCM 32, 165902 (2020)

METHODS

What WannierBerri computes



Blue: used in this demo


Does graphene have a gap?

Step	Run	Checkpoint
0 · DFT	`python 1_dft.py`	`gpaw/nscf.gpw`
1 · Symmetry	irrep	48 operations, 7 of 36 k-points
2 · Wannier functions	`wannierise`	two equal p_z-like functions
3 · Bands	interpolation	blue on grey in the window
4 · SOC on demand	`set_soc_axis`	gap ≈ 36 μeV (GPAW ≈ 39 μeV)
5 · Spin Hall	`wb.run`	plateau at e/2π

Kane, Mele, PRL 95, 226801 (2005)


Quantum spin Hall effect: the Kane–Mele model

second neighbours: $i\lambda\nu_{ij}\, s_z$



ν_{ij} = +1 counterclockwise · A filled, B open

helical edge states




each edge carries ↑ and ↓ in opposite directions

$$H_{\mathrm{SO}} = i\lambda \sum_{\langle\langle ij \rangle\rangle} \nu_{ij}\, c_i^\dagger s_z c_j, \qquad \Delta_{\mathrm{SO}} = 6\sqrt{3}\,\lambda$$

$$\sigma_{xy}^{s_z} = (e/2\pi)(C_\uparrow - C_\downarrow)/2 = e/2\pi$$


Graphene: two p_z Wannier functions from GPAW




One p_z Wannier function on each carbon, A and B

```
calc = GPAW(mode=PW(400), xc="PBE",
            kpts={"size": (6, 6, 1), "gamma": True})
atoms.calc = calc
atoms.get_potential_energy()
sg = SpaceGroup.from_gpaw(calc)                    # irrep
kpts = sg.get_irreducible_kpoints_grid((6, 6, 1))  # 7 of 36
calc.fixed_density(kpts=kpts, nbands=20).write(
    "gpaw/nscf.gpw", mode="all")

# graphene.ipynb
pz = Projection(position_num=[[1/3, 2/3, 0], [2/3, 1/3, 0]],
                orbital="pz", spacegroup=sg)
wandata = wb.WannierDataSOC.from_gpaw(
    calc, projections=ProjectionsSet([pz]))
wandata.wannierise(froz_min=EF - 2.9, froz_max=EF + 2.8,
                   frozen_states={0: [1]},  # pi at Gamma
                   num_iter=100, sitesym=True)
system = wb.SystemSOC.from_wannierdata(wandata, berry=True)
```



The Dirac cone of the Wannier model at K



SOC as a matrix you can rotate and scale

$$H(\mathbf{k}) = H_0(\mathbf{k}) \otimes \sigma_0 + \alpha \sum_c \Delta V^c(\mathbf{k}) \otimes \sigma_c(\hat{\mathbf{n}})$$

H_0: no SOC · $\sigma_c(\hat{\mathbf{n}})$: Pauli matrices along $\hat{\mathbf{n}}$ · α: SOC scale

ΔV: the PAW matrix elements of GPAW's SOC

New $\hat{\mathbf{n}}$ or α: no new DFT

```
K = [1/3, 1/3, 0]
energy = wb.calculators.tabulate.Energy(
    degen_thresh=1e-8)
for alpha in [0, 1]:
    system.set_soc_axis(alpha_soc=alpha)
    E = wb.evaluate_k(system, k=K,
            calculators={"E": energy})
```


```
              gap at K
alpha = 0      0
alpha = 1     36 ueV
GPAW SOC      39 ueV
```

Blöchl, PRB 50, 17953 (1994) · Olsen, PRB 94, 235106 (2016)

SOC gap at the Dirac point K

Wannier model, two p_z	**36 µeV**
GPAW, all 20 bands	**39 µeV**
Experiment (Sichau 2019)	**42 µeV**


From the d part of p_z (Gmitra 2009)




Kane, Mele, PRL 95, 226801 (2005) · Gmitra et al., PRB 80, 235431 (2009) · Sichau et al., PRL 122, 046403 (2019)

$$\sigma_{xy}^{s_z} = \frac{e}{\hbar}\int \frac{d^2k}{(2\pi)^2}\sum_n f_{n\mathbf{k}}\,\Omega_n^{s_z}, \qquad \Omega_n^{s_z} = -2\hbar^2\,\mathrm{Im}\sum_{m\neq n}\frac{(j_x^{s_z})_{nm}(v_y)_{mn}}{(\varepsilon_n-\varepsilon_m)^2}$$

$j_x^{s_z} = \frac{1}{2}\{s_z, v_x\}$: spin current, v: velocity. With s_z conserved, $\sigma_{xy}^{s_z} = (e/2\pi)(C_\uparrow - C_\downarrow)/2 = e/2\pi$



α = 1000 (gap 36 meV): plateau at e/2π



α = 1, the real 36 μeV gap: also e/2π

ACT 2 · MNTE.IPYNB

A Hall effect without magnetisation

Step	Run	Checkpoint
0 · DFT	`python 1_dft.py`	moments ≈ ±4.4 μ_B
1–2 · Spin ↑ Wannier	`altermagnetic=True`	24 of 144 k-points
3 · Splitting	no SOC	up to ≈ 1 eV at $k_z \neq 0$
4 · Turn $\hat{\mathbf{n}}$	`set_soc_axis`	SOC shift at A 4–5× smaller in plane
5 · Anomalous Hall	three $\hat{\mathbf{n}}$	0 for x and c, up to ≈ 80 S/cm for y
6 · Rotate in plane	0–30°	far from sin 3φ


Gonzalez Betancourt et al., PRL 130, 036702 (2023)

Altermagnets: zero net moment, spin-split bands without SOC

Spin ↑ and ↓ sublattices, related by a rotation


Zero net moment, yet an anomalous Hall effect

MnTe: altermagnetic up to $T_N \approx 310$ K




Te octahedra of Mn1 (↑) and Mn2 (↓), turned by 60°

Spin ↓ bands are spin ↑ bands at a rotated **k**: the splitting changes sign, g-wave





Wannierise spin ↑ only, spin ↓ follows by symmetry





```python
calc = GPAW("gpaw/nscf.gpw")   # PBE+U, 24 of 144 k
sg = SpaceGroup.from_gpaw(calc)
Te = [[1/3, 2/3, 1/4], [2/3, 1/3, 3/4]]
projections = ProjectionsSet([
    Projection(position_num=[0, 0, 0], orbital="d",
               spacegroup=sg),          # Mn1 only
    Projection(position_num=Te, orbital="sp2",
               spacegroup=sg, xaxis=[0, -1, 0],
               rotate_basis=True),
    Projection(position_num=Te, orbital="pz",
               spacegroup=sg)])
wandata = wb.WannierDataSOC.from_gpaw(
    calc, projections=projections,
    altermagnetic=True,   # spin down from symmetry
    IBstart=8)            # skip Mn 3s, 3p
wandata.wannierise(froz_min=-np.inf,
                   froz_max=EF + 0.1,
                   num_iter=200, sitesym=True)
system = wb.SystemSOC.from_wannierdata(wandata,
                                       berry=True)
```

Dudarev et al., PRB 57, 1505 (1998) · Kriegner et al., PRB 96, 214418 (2017) · Bossini et al., NJP 22, 083029 (2020)


$k_z = 0$
$k_z = 0.35$ Å^{-1}
0
−1
−2
−3
−4
−5
−6
E − E$_F$ (eV)
K
Γ
K|M̄
Γ̄
M̄
spin ↑
spin ↓






$\hat{\mathbf{n}} \parallel$ y, colour: spin along $\hat{\mathbf{n}}$


```
ahc = wb.calculators.static.AHC(Efermi=Efermi,
    degen_thresh=1e-4, smoother=smoother)
directions = {"x": (90, 0), "y": (90, 90),
              "z": (0, 0)}   # theta, phi
grids = {"x": (48, 48, 32), "y": (96, 96, 64),
         "z": (48, 48, 32)}
for name, (theta, phi) in directions.items():
    system.set_soc_axis(theta=theta, phi=phi,
                        units="degrees")
    result = wb.run(system,
        grid=wb.Grid(system, NK=grids[name]),
        calculators={"ahc": ahc})
```

Zero for $\hat{\mathbf{n}} \parallel x$ and $\hat{\mathbf{n}} \parallel c$, by symmetry




In-plane Néel vector at angle φ from a, amplitudes A_m in S/cm:

$$\sigma_{xy}(\varphi) = A_3 \sin 3\varphi + A_9 \sin 9\varphi + A_{15} \sin 15\varphi + \ldots$$



σ_{xy} = 0 along every a axis, sign flip every 60°



Filled: computed. Open: from symmetry

Your material next

```
calc = GPAW(...)                                  # SCF
kpts = SpaceGroup.from_gpaw(calc)\
           .get_irreducible_kpoints_grid(grid)
calc.fixed_density(kpts=kpts, nbands=...)  # NSCF
wandata = wb.WannierDataSOC.from_gpaw(
    calc, projections=...)
wandata.wannierise(froz_min=..., froz_max=...,
                   sitesym=True)
system = wb.SystemSOC.from_wannierdata(
    wandata, berry=True)
system.set_soc_axis(theta, phi)
wb.run(system, grid=..., calculators=...)
```

Project on the orbitals of your bands

Freeze every state you need

Check a few levels against GPAW first

SUMMARY

Take-aways, and how to run it

Symmetry: DFT only on irreducible k-points

SOC on demand: any spin axis, no new DFT

Graphene: spin Hall plateau e/2π

Altermagnets: Wannierise one spin, get both

RUN IT

github.com/yaroslav-zhumagulov/wannier-berri-demo

```
bash setup/setup.sh    # once, 5 min
conda activate wbdemo
bash run_all.sh        # 6 min, or the notebooks
```

Linux, macOS, WSL2 · GPAW 26.7 from conda-forge

BACKUP

Details

Kane–Mele form · nodal planes of MnTe · timings · references

The Wannier SOC matrix has the Kane–Mele form

$$H_{\mathrm{SO}} = i\lambda \sum_{\langle\langle ij \rangle\rangle} \nu_{ij}\, c_i^{\dagger} s_z c_j, \qquad \Delta_{\mathrm{SO}} = 6\sqrt{3}\,\lambda$$

- Second neighbours: **$\lambda \approx 3.4$ μeV**, purely imaginary, sign ±1 as in Kane–Mele
- $6\sqrt{3}\,\lambda \approx 36$ μeV: the whole gap of the full model
- $\Delta V^{x,y} = 0$: the mirror plane forbids Rashba terms, s_z is conserved
- No SOC between first neighbours: each bond lies in a vertical mirror

Shell	$\lvert d \rvert$ (Å)	$\lvert t \rvert$ (eV)	$\lvert \Delta V^{x,y} \rvert$ (eV)	$\lvert \Delta V^{z} \rvert$ (μeV)
1st, A–B	1.42	2.90	0	0
2nd, A–A	2.46	0.25	0	**3.4**
3rd, A–B	2.84	0.26	0	0
4th, A–B	3.76	0.02	0	0
5th, A–A	4.26	0.04	0	0
6th, A–A	4.92	0.03	0	0.3

Largest element per shell of $H(\mathbf{R})$ and $\Delta V(\mathbf{R})$; nothing fitted


Kane, Mele, PRL 95, 226801 (2005)

Four nodal planes make MnTe g-wave

$E_\uparrow(\mathbf{k}) = E_\downarrow(R_g\mathbf{k})$ for a spin-flipping g, so the splitting vanishes on planes that R_g leaves in place:

- m_z: $k_z = 0$ and $k_z = \pi/c$
- three c-glides: three vertical planes through Γ–K

In the model the splitting is exactly zero on these planes, and up to ≈ 1 eV along Γ–M at $k_z = 0.35$ Å^{-1}



$E_\uparrow - E_\downarrow$, top valence band, $k_z = 0.35$ Å^{-1}

BACKUP

Where the time goes

Step	Graphene	MnTe
GPAW SCF + NSCF, one core	10 s	45 s
`WannierDataSOC.from_gpaw`	3.3 s	9.7 s
`wannierise`	2.1 s	9.7 s
`SystemSOC.from_wannierdata`	1.0 s	6.6 s
Conductivities (step 5)	28 s	63 s
Néel scan (step 6)	–	66 s
Animation frames	1 s	22 s
Whole notebook	1.1 min	4.2 min

Apple M2, 8 cores, 8 Ray workers: 6 minutes for everything, try-its and bonus steps included

With the Wannier model in hand, a new spin axis, SOC strength or Fermi level costs seconds

References

CODES

- WannierBerri: S. S. Tsirkin, npj Comput. Mater. **7**, 33 (2021)
- GPAW: J. J. Mortensen et al., J. Chem. Phys. **160**, 092503 (2024)
- IrRep: M. Iraola et al., Comput. Phys. Commun. **272**, 108226 (2022)
- Wannier90: G. Pizzi et al., J. Phys.: Condens. Matter **32**, 165902 (2020)
- Wannier ecosystem: A. Marrazzo et al., Rev. Mod. Phys. **96**, 045008 (2024)

METHODS

- N. Marzari, D. Vanderbilt, PRB **56**, 12847 (1997); I. Souza et al., PRB **65**, 035109 (2001); N. Marzari et al., Rev. Mod. Phys. **84**, 1419 (2012)
- R. Sakuma, PRB **87**, 235109 (2013)
- D. Xiao, M.-C. Chang, Q. Niu, Rev. Mod. Phys. **82**, 1959 (2010)
- X. Wang et al., PRB **74**, 195118 (2006); J. Qiao et al., PRB **98**, 214402 (2018); J. H. Ryoo et al., PRB **99**, 235113 (2019)

PAW, SOC AND DFT+U

- P. E. Blöchl, PRB **50**, 17953 (1994); T. Olsen, PRB **94**, 235106 (2016); S. L. Dudarev et al., PRB **57**, 1505 (1998)

GRAPHENE

- C. L. Kane, E. J. Mele, PRL **95**, 226801 and 146802 (2005)
- M. Gmitra et al., PRB **80**, 235431 (2009); J. C. Boettger, S. B. Trickey, PRB **75**, 121402 (2007)
- J. Sichau et al., PRL **122**, 046403 (2019)

ALTERMAGNETS AND MNTE

- L. Šmejkal et al., Sci. Adv. **6**, eaaz8809 (2020); PRX **12**, 031042 and 040501 (2022)
- D. Kriegner et al., PRB **96**, 214418 (2017); D. Bossini et al., New J. Phys. **22**, 083029 (2020)
- R. D. Gonzalez Betancourt et al., PRL **130**, 036702 (2023)
- J. Krempaský et al., Nature **626**, 517 (2024)