1. The probability of a leap year selected at random contain 53 Sunday is:					
(a) 53/366 (b) 1/7 (c) 2/7 (d) 53/365					
2. A bag contains 3 red and 2 blue marbles. A marble is drawn at					
random. The probability of drawing a black ball is:					
(a) 3/5 (b) 2/5 (c) 0/5 (d) 1/5					
3. The probability that it will rain tomorrow is 0.85. What is the					
probability that it will not rain tomorrow					
(a) 0.25 (b) 0.145 (c) 3/20 (d) none of these					
4. What is the probability that a number selected from the numbers					
(1, 2, 3,,15) is a multiple of 4?					
(a) 1/5 (b) 4/5 (c) 2/15 (d) 1/3					
5. What are the total outcomes when we throw three coins?					
(a) 4 (b) 5 (c) 8 (d) 7					
6. The probability that a prime number selected at random from the					
numbers (1,2,3,35) is :					
(a) 12/35 (b) 11/35 (c) 13/35 (d) none of these					
7. The sum of the probability of an event and non event is:					
(a) 2 (b) 1 (c) 0 (d) none of these.					
8. The following probabilities are given; choose the correct answer					
for that which is not possible.					
(a) 0.15 (b) 2/7 (c) 7/5 (d) none of these.					
9. If three coins are tossed simultaneously, than the probability of					
getting at least two heads, is:					
(a) 1/4 (b) 3/8 (c) ½ (d) 1/8					
10. A letter is chosen at random from the letters of the word					
<b>♦</b> ASSASSINATION <b>♦</b> . The probability that the letter chosen has:					
(a) 6/13 (b) 7/13 (c) 1 (d) none of these.					
11. A dice is thrown. Find the probability of getting an even number.					
(A) 2/3 (B) 1 (C) 5/6 (D) 1/2					
12. Two coins are thrown at the same time. Find the probability of getting both heads.					
(A) 3/4 (B) 1/4 (C) 1/2 (D) 0					
13. Two dice are thrown simultaneously. The probability of getting a					

1

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D)	4/9	
	ds are number ime number.	ed from 1 to 1	00. Find th	ne probability of	f
(A) 3/4		(C) 1/4	1)	D) 29/100	
•	a blue ball is d			If the probabiles .If the probabiles	•
(A) 5	(B) 10	(C) 15	(D) 20		
			n the proba		
mixed thore the probabi	narked with nu oughly. One ca llity that the nu (B) 1/10	rd is drawn fro mber on card	om this box is a perfec	randomly, the t square.	n
<b>18. What is</b> (A) 1/7	s the probabilit (B) 53/3		•	in a leap year? 7/366	
probability	is drawn from of getting a kir (B) 3/26 (0	ng of red suit.		2 cards. Find th	1e
equally like 1,2,312	e of chance con ly to come to r ,then the proba (B) 1/12	est pointing to ability that it w	o one of the	e number an odd numbe	r is:
its outcome result i.e. the probability	e each time. Ar	yan wins if all hree tails and lose the game	the tosses loses othe e.	times and noting times the same the training the	

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
(A) 364/365	(B) 31/365	(C) 1/365	(D) 1/133225				
23. A number $x$ is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$							
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is:  (A) $10$ (B) $6$ (C) $8$ (D) $7$							
25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) $7/50$ (B) $4/25$ (C) $1/25$ (D) $2/25$							
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?							
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) 4/21				
	ed by batsman ir e standard devia	•	es are 50, 70, 82,				
		c) 25.29	d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
	b) 13, 18		d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) $^{1}/_{2}$	b) $^{1}/_{3}$	c) $\frac{1}{4}$					
		<b>d 3. The value of I</b> 2) 27 d)					
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?							

32.Out of t probability?	he following valu	ıes, which	one is not po	ssible in
a) $P(x) = 1$	b) ∑ x P(	(x) = 3		
	d) P(x)			
	2 and E(z) = 4, t	•	· •	
a) 2	b) 6	c) 0	d) I	nsufficient data
34.The cov	ariance of two in	dependen	t random vari	able is
a) 1	b) 0	c) - 1	d) (	Undefined
, ,	) = k <sup>2</sup> – 8 then, th			
a) 0	b) 1	c) 3	d)	Insufficient data
• •	<b>0.5 and x = 4, th</b> b) 0.5	<b>en E(x) = ?</b> c) 4	d) 2	2
is always?	erete probability			-
a) 0	b) Infinite	c) 1	d) U	Indefined
38.If the pr	obability of hitti	ng the targ	et is 0.4, find	mean and
	b) 0.6, 0.2	24	c) 0.4, 0.16	d) 0.6, 0.16
-	-	_	-	ace will strike the an and variance?
a) 0.6, 0.24		c) 0	· -	d) 4, 1.6
	e mean of tossing	~	Α	
a) 2 <b>41. What is</b>	,	8 <mark>ariance fo</mark>	d) 1 r <b>standard no</b> l	rmal distribution?

c) 5

d) 7

a) 3

b) 4

a) Mean is 0 and variance is 1 b) Mean is 1 and variance is 0 c) Mean is 0 and variance is $\infty$ d) Mean is $\infty$ and variance is 0								
<b>42.Variance of a random variable X is given by</b> a) E(X) b) E(X2) c) E(X2) - (E(X))2							. d) (E(X))2	
	B.Mean of E(X)	f <b>a random</b> b) E(X2		e X is giv c) E(X2	•		d) (E(X))2	
44.Mean of a constant 'a' is a) 0								
<b>45</b> .		e <b>of a cons</b> b) a		is c) a/2		d) 1		
46.	46.Find the mean and variance of X?							
	Х	0	1	2	3	4		
	f(x)	1/9	2/9	3/9	2/9	1/9		
a) :	2, 4/3	b) 3	3, 4/3		c) 2, 2/3	3	d) 3, 2/3	
47	47. Find the expectation of a random variable X?							

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) (	).5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

b) npq

c) np2q

d) npq2

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.** 

- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?** 

- a)  $\sqrt{np}$  b)  $\sqrt{pq}$  c) (np)2 d)  $\sqrt{npq}$