

# Finite element spaces for tensor fields with specified interelement continuity conditions

Yakov Berchenko-Kogan, joint with Evan Gawlik

Florida Institute of Technology  
Supported by NSF DMS-2411209

October 19, 2025

# Tangential and normal continuity of vector fields




Figure: Tangential continuity (left) vs. normal continuity (right)

## Tangential continuity

- Well-defined line integrals.
- In  $H(\text{curl})$ .



## Normal continuity

- Well-defined fluxes.
- In  $H(\text{div})$ .

# What's wrong with full continuity?


Finite element exterior calculus (FEEC) perspective: differential complexes

Gradients of scalar fields only have tangential continuity



Spurious eigenvalues of the  $\operatorname{curl curl}$  operator (AFW, 2010)

- Solve  $\operatorname{curl curl} \mathbf{u} = \lambda \mathbf{u}$ , where  $\mathbf{u}$  is a vector field on a square domain with appropriate boundary conditions.
- Using vector fields with **full continuity** yields **false** eigenvalue  $\lambda = 6$ .




# What's wrong with full continuity?

Geometric perspective

Extrinsic



Intrinsic




---

Four images from Wikipedia



# What's wrong with full continuity?

Geometric perspective

Extrinsic



Intrinsic



---

Four images from Wikipedia




# What's wrong with full continuity?

Geometric perspective

Extrinsic



Intrinsic



## Why compute intrinsically?

- Intrinsic problems, e.g. numerical relativity, Ricci flow.
- Structure preservation: independence of embedding.

Four images from Wikipedia

# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.

# What's wrong with full continuity?


Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?

# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?



# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?



# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?



# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?



# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?



# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?



# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?



# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?



# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?



# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?




# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?


continuous elements



# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?




# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?

continuous elements




# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?

continuous elements




# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?

continuous elements




# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?

continuous elements




continuous on each triangle

# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?

continuous elements




continuous on each triangle  
**discontinuous** across red edge

# What's wrong with full continuity?


Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?

continuous elements



blow-up elements




continuous on each triangle  
**discontinuous** across red edge

# What's wrong with full continuity?


Geometric perspective: Angle defect obstruction to continuous elements

- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?

continuous elements



blow-up elements




continuous on each triangle  
**discontinuous** across red edge

# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?

continuous elements



continuous on each triangle  
**discontinuous** across red edge

blow-up elements




continuous across all edges

# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?

continuous elements



continuous on each triangle  
**discontinuous** across red edge

blow-up elements




continuous across all edges  
discontinuous at vertices

# What's wrong with full continuity?

Geometric perspective: Angle defect obstruction to continuous elements


- Try to construct a tangent vector field on the icosahedron.
- What do we see when we zoom in on a vertex?

continuous elements



continuous on each triangle  
**discontinuous** across red edge

blow-up elements



continuous across all edges  
discontinuous at vertices

- See also later today: Alan Demlow (10am).

## One-forms $\Lambda^1$

- $M dx + N dy + P dz$
- Restricted to the  $xy$ -plane  $z = 0$ :
  - $M dx + N dy$ .
  - Tangential components.

## Two-forms $\Lambda^2$

- $M dy \wedge dz + N dz \wedge dx + P dx \wedge dy$ .
- Restricted to the  $xy$ -plane  $z = 0$ :
  - $P dx \wedge dy$ .
  - Normal component.

## Continuity conditions

- Vector fields with tangential continuity are one-forms.
- Vector fields with normal continuity are  $(n - 1)$ -forms.

## Continuity conditions for 2-tensors (matrix fields)

- tangential–tangential
- normal–normal
- normal–tangential

## Applications

- Strain/stress tensors
  - Elasticity (objects deforming under stress)
  - Fluid mechanics (Stokes equations)
- Numerical geometry/relativity
  - Riemannian/Minkowski metric
  - Curvature tensor
- See also later today: Francis Aznaran (9:30am), Alan Demlow (10am), Qingguo Hong (2pm), Bowen Shi (2:30pm).

# Double forms

## Vector fields ( $\mathbb{R}^3$ )

- Vector fields with tangential continuity are one-forms  $\Lambda^1$ .
- Vector fields with normal continuity are two-forms  $\Lambda^2$ .

## Matrix fields ( $\mathbb{R}^3 \otimes \mathbb{R}^3$ )

- Matrix fields with tangential–tangential continuity are  $(1, 1)$ -forms  
 $\Lambda^{1,1} := \Lambda^1 \otimes \Lambda^1$ .
- Matrix fields with normal–tangential continuity are  $(2, 1)$ -forms  
 $\Lambda^{2,1} := \Lambda^2 \otimes \Lambda^1$ .
- Matrix fields with normal–normal continuity are  $(2, 2)$ -forms  
 $\Lambda^{2,2} := \Lambda^2 \otimes \Lambda^2$ .

## More on double forms later today

- Evan Gawlik (8:30am).
- Anil Hirani (9am).

## Affine-invariant (metric-independent) finite element spaces

- FEEC differential forms  $\Lambda^k$  and their continuity conditions are defined without reference to a Riemannian metric.
- Same for double forms  $\Lambda^{p,q}$ .
- Angle defect cannot pose a problem since angle defect is not even defined without a Riemannian metric.
- In particular, for vector fields with tangential or normal continuity, FEEC works just as well on surface meshes as it does on the plane.

## Metric-dependent finite element spaces

- Defining finite element spaces of vector fields with full continuity requires a Riemannian metric (even via differential form proxies).
- Behavior depends on whether angle defect is zero or not.

## Theorem (Eigendecomposition of $s^*s$ )

$$\Lambda^{p,q} = \bigoplus_m \Lambda_m^{p,q}, \quad \max\{0, q-p\} \leq m \leq \min\{q, n-p\}.$$

### Example

- $\Lambda_0^{1,1}$ : Symmetric bilinear forms,  $\varphi(X; Y) = \varphi(Y; X)$ .
- $\Lambda_1^{1,1}$ :  $\Lambda^2$ , antisymmetric bilinear forms,  $\varphi(X; Y) = -\varphi(Y; X)$ .

---

- $\Lambda_0^{2,1}$ : spanned by  $\alpha \otimes \beta$  such that  $\alpha \wedge \beta = 0$ .
  - Matrix proxy in 3D: trace-free matrices.
- $\Lambda_1^{2,1}$ :  $\Lambda^3$ .
  - Matrix proxy in 3D: multiples of the identity matrix.

---

- $\Lambda_0^{2,2}$ : Symmetric, satisfying the algebraic Bianchi identity.
  - Riemann curvature tensor.
- $\Lambda_1^{2,2}$ : Antisymmetric,  $\varphi(X, Y; Z, W) = -\varphi(Z, W; X, Y)$ .
- $\Lambda_2^{2,2}$ :  $\Lambda^4$ .

## Theorem (—, Gawlik)

Apart from  $\Lambda_q^{p,q} \cong \Lambda^{p+q}$  with constant coefficients, there is a finite element space for every natural space of double forms  $\Lambda_m^{p,q}$  with polynomial coefficients of any degree (including zero).

## Example (Constant coefficient spaces)

- $\Lambda_0^{1,1}$ : symmetric matrices with tangential–tangential continuity (Regge, 1961).
  - Higher order: (Li, 2018).
- $\Lambda_0^{2,1}$  in 3D: trace-free matrices with normal–tangential continuity (Gopalakrishnan, Lederer, and Schöberl, 2019).
- $\Lambda_0^{2,2}$  in 3D: symmetric matrices with normal–normal continuity (Pechstein and Schöberl, 2011).
- $\Lambda_0^{2,2}$  (or  $\Lambda_0^{n-2,n-2}$ ) in any dimension: finite elements for the Riemann curvature tensor.

# Degrees of freedom for constant coefficient spaces


	$d$						
	0	1	2	3	4	5	6
$\Lambda_0^{1,1}$	0	<b>1</b>	0	0	0	0	0
$\Lambda_0^{2,1}$	0	0	<b>2</b>	0	0	0	0
$\Lambda_0^{2,2}$	0	0	<b>1</b>	<b>2</b>	0	0	0
$\Lambda_1^{2,2} \cong \Lambda_0^{3,1}$	0	0	0	<b>3</b>	0	0	0
$\Lambda_0^{3,2}$	0	0	0	<b>3</b>	<b>5</b>	0	0
$\Lambda_1^{3,2} \cong \Lambda_0^{4,1}$	0	0	0	0	<b>4</b>	0	0
$\Lambda_0^{3,3}$	0	0	0	<b>1</b>	<b>5</b>	<b>5</b>	0
$\Lambda_1^{3,3} \cong \Lambda_0^{4,2}$	0	0	0	0	<b>6</b>	<b>9</b>	0
$\Lambda_2^{3,3} \cong \Lambda_1^{4,2} \cong \Lambda_0^{5,1}$	0	0	0	0	0	<b>5</b>	0

**Table:** Number of degrees of freedom for  $\Lambda_m^{p,q}$  associated to a face of the triangulation of dimension  $d$  is  $\frac{p-q+2m+1}{p+m+1} \binom{d+1}{q-m} \binom{q-m-1}{d-p-m}$ .


## Motivating problem

- Goal: construct **intrinsic** discretizations of tangent vector fields on smooth surfaces that are **continuous across edges**.
- Obstruction to using classical Lagrange  $\mathcal{P}_1$  elements: **angle defect**.

continuous elements




blow-up elements



continuous on each triangle  
**discontinuous** across red edge


continuous across all edges  
**discontinuous at vertices**

# A simplicial analogue of the angular coordinate




$$\frac{\lambda_2}{\lambda_1 + \lambda_2}$$


# Lagrange $\mathcal{P}_1$ shape functions



$$\lambda_0$$





$$\lambda_1$$





$$\lambda_2$$



# Blow-up $b\mathcal{P}_1$ shape functions



$$\psi_{012} = \frac{\lambda_0 \lambda_1}{\lambda_1 + \lambda_2}, \quad \psi_{021} = \frac{\lambda_0 \lambda_2}{\lambda_2 + \lambda_1},$$



$$\psi_{102} = \frac{\lambda_1 \lambda_0}{\lambda_0 + \lambda_2}, \quad \psi_{120} = \frac{\lambda_1 \lambda_2}{\lambda_2 + \lambda_0},$$



$$\psi_{201} = \frac{\lambda_2 \lambda_0}{\lambda_0 + \lambda_1}, \quad \psi_{210} = \frac{\lambda_2 \lambda_1}{\lambda_1 + \lambda_0}.$$

## Shape function

$$\psi_{012} = \frac{\lambda_0 \lambda_1}{\lambda_1 + \lambda_2} = \frac{\lambda_0}{\lambda_0 + \lambda_1 + \lambda_2} \cdot \frac{\lambda_1}{\lambda_1 + \lambda_2} \cdot \frac{\lambda_2}{\lambda_2}.$$

## Shape function

$$\psi_{012} = \frac{\lambda_0 \lambda_1}{\lambda_1 + \lambda_2} = \frac{\lambda_0}{\lambda_0 + \lambda_1 + \lambda_2} \cdot \frac{\lambda_1}{\lambda_1 + \lambda_2} \cdot \frac{\lambda_2}{\lambda_2}.$$

## Earlier appearances

## Shape function

$$\psi_{012} = \frac{\lambda_0 \lambda_1}{\lambda_1 + \lambda_2} = \frac{\lambda_0}{\lambda_0 + \lambda_1 + \lambda_2} \cdot \frac{\lambda_1}{\lambda_1 + \lambda_2} \cdot \frac{\lambda_2}{\lambda_2}.$$

## Earlier appearances

- Geometric invariants (Chen, 1957).

## Shape function

$$\psi_{012} = \frac{\lambda_0 \lambda_1}{\lambda_1 + \lambda_2} = \frac{\lambda_0}{\lambda_0 + \lambda_1 + \lambda_2} \cdot \frac{\lambda_1}{\lambda_1 + \lambda_2} \cdot \frac{\lambda_2}{\lambda_2}.$$

## Earlier appearances

- Geometric invariants (Chen, 1957).
- Horse betting (Harville, 1973).


## Shape function

$$\psi_{012} = \frac{\lambda_0 \lambda_1}{\lambda_1 + \lambda_2} = \frac{\lambda_0}{\lambda_0 + \lambda_1 + \lambda_2} \cdot \frac{\lambda_1}{\lambda_1 + \lambda_2} \cdot \frac{\lambda_2}{\lambda_2}.$$

## Earlier appearances

- Geometric invariants (Chen, 1957).
- Horse betting (Harville, 1973).
- Intersection homology (Brasselet, Goresky, MacPherson, 1991; Bendiffalah, 1995).

## Classical Lagrange $\mathcal{P}_1$




Barycentric coordinates:  $\lambda_0 + \lambda_1 + \lambda_2 = 1$ .

- 0 :  $\lambda_0 = 1 \Leftrightarrow \lambda_1 = \lambda_2 = 0$
- 1 :  $\lambda_1 = 1 \Leftrightarrow \lambda_2 = \lambda_0 = 0$
- 2 :  $\lambda_2 = 1 \Leftrightarrow \lambda_0 = \lambda_1 = 0$

# Degrees of freedom


## Classical Lagrange $\mathcal{P}_1$



Barycentric coordinates:  $\lambda_0 + \lambda_1 + \lambda_2 = 1$ .


- 0 :  $\lambda_0 = 1 \Leftrightarrow \lambda_1 = \lambda_2 = 0$
- 1 :  $\lambda_1 = 1 \Leftrightarrow \lambda_2 = \lambda_0 = 0$
- 2 :  $\lambda_2 = 1 \Leftrightarrow \lambda_0 = \lambda_1 = 0$

## Blow-up $b\mathcal{P}_1$




- 012 :  $\lim_{\lambda_1 \rightarrow 0} \lim_{\lambda_2 \rightarrow 0}$
- 120 :  $\lim_{\lambda_2 \rightarrow 0} \lim_{\lambda_0 \rightarrow 0}$
- 201 :  $\lim_{\lambda_0 \rightarrow 0} \lim_{\lambda_1 \rightarrow 0}$
- 021 :  $\lim_{\lambda_2 \rightarrow 0} \lim_{\lambda_1 \rightarrow 0}$
- 102 :  $\lim_{\lambda_0 \rightarrow 0} \lim_{\lambda_2 \rightarrow 0}$
- 210 :  $\lim_{\lambda_1 \rightarrow 0} \lim_{\lambda_0 \rightarrow 0}$

- Scalar fields: we place a number at each dot.




Blow-up finite elements




Blow-up finite elements

- Scalar fields: we place a number at each dot.
- Vector fields: we place two numbers at each dot, for the tangential and normal components, respectively.




Blow-up finite elements

- Scalar fields: we place a number at each dot.
- Vector fields: we place two numbers at each dot, for the tangential and normal components, respectively.
  - Enforce continuity for **both** components, yielding **full continuity across edges**.




Blow-up finite elements

- Scalar fields: we place a number at each dot.
- Vector fields: we place two numbers at each dot, for the tangential and normal components, respectively.
  - Enforce continuity for **both** components, yielding **full continuity across edges**.
- Matrix fields: At each dot, we record the tangential–tangential component, the tangential–normal component, etc.




Blow-up finite elements

- Scalar fields: we place a number at each dot.
- Vector fields: we place two numbers at each dot, for the tangential and normal components, respectively.
  - Enforce continuity for **both** components, yielding **full continuity across edges**.
- Matrix fields: At each dot, we record the tangential–tangential component, the tangential–normal component, etc.
  - Can impose conditions on the components such as symmetry, trace-free, etc.



Blow-up finite elements

- Scalar fields: we place a number at each dot.
- Vector fields: we place two numbers at each dot, for the tangential and normal components, respectively.
  - Enforce continuity for **both** components, yielding **full continuity across edges**.
- Matrix fields: At each dot, we record the tangential–tangential component, the tangential–normal component, etc.
  - Can impose conditions on the components such as symmetry, trace-free, etc.
  - Can enforce continuity for all components or just some of them.



Blow-up finite elements

- Scalar fields: we place a number at each dot.
- Vector fields: we place two numbers at each dot, for the tangential and normal components, respectively.
  - Enforce continuity for **both** components, yielding **full continuity across edges**.
- Matrix fields: At each dot, we record the tangential–tangential component, the tangential–normal component, etc.
  - Can impose conditions on the components such as symmetry, trace-free, etc.
  - Can enforce continuity for all components or just some of them.
- General tensor fields are analogous.

## Hodge Laplacian

$$(dd^* + d^*d)v^\flat = \lambda v^\flat.$$


- Tangential continuity across edges suffices.
- Standard FEEC works.
- $L^2$  pairing suffices.

## Bochner Laplacian


$$\nabla^*\nabla v = \lambda v.$$

- Must have full continuity across edges.
- Can't use standard FEEC.
- Needs Riemannian metric.

## Bochner Laplacian on sphere using blow-up elements



Eigenvalue error



Eigenfield magnitude ( $\lambda = 11, 11, 19, 19$ )

# There's more

So far in this talk

## So far in this talk

- Lowest order blow-up elements in two dimensions,  $b\mathcal{P}_1(T^2)$ ,

## So far in this talk

- Lowest order blow-up elements in two dimensions,  $b\mathcal{P}_1(T^2)$ ,
  - including tensor fields with components in  $b\mathcal{P}_1(T^2)$ .

# There's more

## So far in this talk

- Lowest order blow-up elements in two dimensions,  $b\mathcal{P}_1(T^2)$ ,
  - including tensor fields with components in  $b\mathcal{P}_1(T^2)$ .

## Our paper

# There's more

## So far in this talk

- Lowest order blow-up elements in two dimensions,  $b\mathcal{P}_1(T^2)$ ,
  - including tensor fields with components in  $b\mathcal{P}_1(T^2)$ .

## Our paper

- Differential complex of blow-up Whitney forms in any dimension,  $b\mathcal{P}_1^-\Lambda^k(T^n)$ .

## There's more

### So far in this talk

- Lowest order blow-up elements in two dimensions,  $b\mathcal{P}_1(T^2)$ ,
  - including tensor fields with components in  $b\mathcal{P}_1(T^2)$ .

### Our paper

- Differential complex of blow-up Whitney forms in any dimension,  $b\mathcal{P}_1^-\Lambda^k(T^n)$ .
  - Shape functions previously studied in (Brasselet, Goresky, MacPherson, 1991), called shadow forms.

# There's more

## So far in this talk

- Lowest order blow-up elements in two dimensions,  $b\mathcal{P}_1(T^2)$ ,
  - including tensor fields with components in  $b\mathcal{P}_1(T^2)$ .

## Our paper

- Differential complex of blow-up Whitney forms in any dimension,  $b\mathcal{P}_1^-\Lambda^k(T^n)$ .
  - Shape functions previously studied in (Brasselet, Goresky, MacPherson, 1991), called shadow forms.
- Higher-order blow-up scalar fields  $b\mathcal{P}_r(T^n)$ .

# There's more

## So far in this talk

- Lowest order blow-up elements in two dimensions,  $b\mathcal{P}_1(T^2)$ ,
  - including tensor fields with components in  $b\mathcal{P}_1(T^2)$ .

## Our paper

- Differential complex of blow-up Whitney forms in any dimension,  $b\mathcal{P}_1^-\Lambda^k(T^n)$ .
  - Shape functions previously studied in (Brasselet, Goresky, MacPherson, 1991), called shadow forms.
- Higher-order blow-up scalar fields  $b\mathcal{P}_r(T^n)$ .
- A surprising connection to arrival times of Poisson processes, yielding simpler computations.

# There's more

## So far in this talk

- Lowest order blow-up elements in two dimensions,  $b\mathcal{P}_1(T^2)$ ,
  - including tensor fields with components in  $b\mathcal{P}_1(T^2)$ .

## Our paper

- Differential complex of blow-up Whitney forms in any dimension,  $b\mathcal{P}_1^-\Lambda^k(T^n)$ .
  - Shape functions previously studied in (Brasselet, Goresky, MacPherson, 1991), called shadow forms.
- Higher-order blow-up scalar fields  $b\mathcal{P}_r(T^n)$ .
- A surprising connection to arrival times of Poisson processes, yielding simpler computations.
  - Three radiation sources with rates  $\lambda_0$ ,  $\lambda_1$ , and  $\lambda_2$ , sum 1.

# There's more

## So far in this talk

- Lowest order blow-up elements in two dimensions,  $b\mathcal{P}_1(T^2)$ ,
  - including tensor fields with components in  $b\mathcal{P}_1(T^2)$ .

## Our paper

- Differential complex of blow-up Whitney forms in any dimension,  $b\mathcal{P}_1^-\Lambda^k(T^n)$ .
  - Shape functions previously studied in (Brasselet, Goresky, MacPherson, 1991), called shadow forms.
- Higher-order blow-up scalar fields  $b\mathcal{P}_r(T^n)$ .
- A surprising connection to arrival times of Poisson processes, yielding simpler computations.
  - Three radiation sources with rates  $\lambda_0$ ,  $\lambda_1$ , and  $\lambda_2$ , sum 1.
  - Let  $t_0$ ,  $t_1$ ,  $t_2$  be the times when the respective radiation sources produce their first particle.

# There's more

## So far in this talk

- Lowest order blow-up elements in two dimensions,  $b\mathcal{P}_1(T^2)$ ,
  - including tensor fields with components in  $b\mathcal{P}_1(T^2)$ .

## Our paper

- Differential complex of blow-up Whitney forms in any dimension,  $b\mathcal{P}_1^-\Lambda^k(T^n)$ .
  - Shape functions previously studied in (Brasselet, Goresky, MacPherson, 1991), called shadow forms.
- Higher-order blow-up scalar fields  $b\mathcal{P}_r(T^n)$ .
- A surprising connection to arrival times of Poisson processes, yielding simpler computations.
  - Three radiation sources with rates  $\lambda_0$ ,  $\lambda_1$ , and  $\lambda_2$ , sum 1.
  - Let  $t_0$ ,  $t_1$ ,  $t_2$  be the times when the respective radiation sources produce their first particle.
  - $\frac{\lambda_0\lambda_1}{\lambda_1+\lambda_2}$  is the probability that  $t_0 \leq t_1 \leq t_2$ .

# There's more


## So far in this talk

- Lowest order blow-up elements in two dimensions,  $b\mathcal{P}_1(T^2)$ ,
  - including tensor fields with components in  $b\mathcal{P}_1(T^2)$ .

## Our paper


- Differential complex of blow-up Whitney forms in any dimension,  $b\mathcal{P}_1^-\Lambda^k(T^n)$ .
  - Shape functions previously studied in (Brasselet, Goresky, MacPherson, 1991), called shadow forms.
- Higher-order blow-up scalar fields  $b\mathcal{P}_r(T^n)$ .
- A surprising connection to arrival times of Poisson processes, yielding simpler computations.
  - Three radiation sources with rates  $\lambda_0$ ,  $\lambda_1$ , and  $\lambda_2$ , sum 1.
  - Let  $t_0$ ,  $t_1$ ,  $t_2$  be the times when the respective radiation sources produce their first particle.
  - $\frac{\lambda_0\lambda_1}{\lambda_1+\lambda_2}$  is the probability that  $t_0 \leq t_1 \leq t_2$ .
- Degrees of freedom in terms of blow-up simplex.

# Blowing up



- Even on an individual triangle, the vector field is not continuous at the origin.
- But it is “continuous in polar coordinates,” i.e. in  $r$  and  $\theta$ .

# Blowing up



- Even on an individual triangle, the vector field is not continuous at the origin.
- But it is “continuous in polar coordinates,” i.e. in  $r$  and  $\theta$ .

## Blowing up manifolds with corners (Melrose, 1996)

- formalizes continuity/smoothness “in polar coordinates”



# Thank you

 Yakov Berchenko-Kogan and Evan S. Gawlik

Finite element spaces of double forms.

<https://arxiv.org/abs/2505.17243>

 Yakov Berchenko-Kogan

Duality in finite element exterior calculus and Hodge duality on the sphere.

*Found. Comput. Math.* 21(5):1153–1180, 2021.

 Evan S. Gawlik and Anil N. Hirani

Sequences from sequences, sans coordinates.

In preparation.

 Yakov Berchenko-Kogan and Evan S. Gawlik

Blow-up Whitney forms, shadow forms, and Poisson processes.

*Results in Applied Mathematics*, special issue on Hilbert complexes, Paper No. 100529, 2025.

 J. P. Brasselet, M. Goresky, and R. MacPherson.

Simplicial differential forms with poles.

*Amer. J. Math.*, 113(6):1019–1052, 1991.

Supported by NSF DMS-2411209.