

Discrete Random Variables 1

STAT-S520

Arturo Valdivia

01-26-23

These slides complement ISI Chapter 4

Discrete random variable

- ▶ Let X be a random variable. X is discrete if $X(S)$ is countable
 - ▶ A set is countable if it is finite or denumerable
 - ▶ A set is denumerable if there is a one-to-one correspondence with \mathbb{N}

Exercise 1

- ▶ Let $X(S) = \{0, 1, 2\}$, then X is ...
- ▶ Let $Y(S) = \mathbb{N}$ (naturals), then Y is ...
- ▶ Let $X_1(S) = \mathbb{R}$ (reals), then X_1 is ...
- ▶ Let $X_2(S) = \mathbb{Z}$ (integers), then X_2 is ...
- ▶ Let $X_3(S) = \mathbb{Q}$ (rationals), then X_3 is ...
- ▶ Let $Z(S) = [0, 1]$, then Z is ...

Probability Mass Function

Let X be a discrete random variable. The probability mass function (PMF) of X is a function f , with $f : \mathbb{R} \rightarrow [0, 1]$, such that, for any $y \in \mathbb{R}$:

$$f(y) = \begin{cases} P(X = y) & \text{if } y \in X(S) \\ 0 & \text{otherwise} \end{cases}$$

Exercise 2

We toss a fair coin twice and X is the number of heads. What is

▶ $f(1) =$

▶ $f(2) =$

▶ $f(\pi) =$

▶ $f(10) =$

CDF and PMF

If X is a discrete random variable with range $X(S)$ and for any given $y \in \mathbb{R}$ the relationship between the PMF and CDF is given by:

$$F(y) = \sum_{x: x \in L(y)} f(x)$$

where $L(y)$ is the set of numbers in the range of X that are less than or equal to y , or in math notation: $L(y) = X(S) \cap (-\infty, y]$,

Exercise 3

We toss a fair coin twice and X is the number of heads. Let's find $F(\sqrt{2})$ using the PMF

Bernoulli trials

- ▶ A Bernoulli trial is a random variable where $X(S) = \{0, 1\}$.
 - ▶ It is customary to call $X = 1$ a success and $X = 0$ a failure.
- ▶ There is a family of Bernoulli trials with parameter p where $p = P(X = 1)$ and we write

$$X \sim \text{Bernoulli}(p)$$

PMF of a Bernoulli trial

- ▶ If $X \sim \text{Bernoulli}(p)$, the PMF of X is given by

$$f(x) = \begin{cases} p & \text{if } x = 1 \\ 1 - p & x = 0 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ We can simplify this notation by writing

$$f(x) = p^x(1 - p)^{1-x}$$

for $x \in \{0, 1\}$ and $f(x) = 0$ otherwise.

Binomial distribution

- ▶ Let X_1, X_2, \dots, X_n independent Bernoulli trials with parameter p
- ▶ We construct a random variable as follows:

$$Y = \sum_{i=1}^n X_i = X_1 + \dots + X_n$$

We then say that Y follows a binomial distribution with parameters n and p and write

$$Y \sim \text{binomial}(n, p)$$

Exercise 4

- ▶ You invite 50 friends to your birthday party. We assume that
 - ▶ Friends don't influence each other
 - ▶ Each friend has the same chance to attend
- ▶ There is a 0.8 chance that any given friend attends the party.

PMF of a Binomial distribution

If $Y \sim \text{binomial}(n, p)$, the PMF of Y is given by

$$f(x) = P(Y = x) = \binom{n}{x} p^x (1 - p)^{n-x}$$

for $x \in Y(S)$ and $f(x) = 0$ otherwise.

Expected value

- ▶ Let X be a discrete random variable. The expected value of X is given by

$$EX = \sum_{x \in X(S)} x \cdot f(x)$$

- ▶ The expected value can be understood as the long-run average of values that X assigns when the experiment is performed many times

Variance

- ▶ Let X be a discrete random variable. The variance of X is given by

$$\text{Var}X = E\left((X - \mu)^2\right) = \sum_{x \in X(S)} (x - \mu)^2 \cdot f(x)$$

- ▶ The standard deviation of X is the square root of the variance

Properties of the expected value and variance

Let X, Y be random variables and a, b scalars (constant values).

- ▶ $E(a + X) = a + EX$
- ▶ $E(bX) = bEX$
- ▶ $E(X + Y) = EX + EY$

If in addition, X and Y are independent:

- ▶ $Var(a + X) = VarX$
- ▶ $Var(bX) = b^2 \cdot VarX$
- ▶ $Var(X + Y) = VarX + VarY$

Exercise 5

Let X, Y be independent random variables with $EX = 1$, $EY = 2$, $VarX = 4$, $VarY = 9$. Find the expected value and variance of $2X + 1$ and $X - Y$