

# GROMOV-WITTEN INVARIANTS OF RIEMANN-FINSLER MANIFOLDS

YASHA SAVELYEV

**ABSTRACT.** We define a  $\mathbb{Q}$ -valued deformation invariant of certain complete Riemann-Finsler manifolds. It is proved that every rational number is the value of this invariant for some compact Riemannian manifold. We use this to find the first generalizations, to non-compact fibrations, of Preissman's theorem on non-existence of negative sectional curvature metrics on compact products. Along the way, we also prove that sky catastrophes of smooth dynamical systems are not geodesible by a certain class of forward complete Riemann-Finsler metrics, in particular by Riemannian metrics with non-positive sectional curvature. This partially answers a question of Fuller and gives important examples for our theory here. In a sister paper [13], we study a direct generalization of this metric invariant, by lifting the count of geodesics to a Gromov-Witten count of elliptic curves in an associated locally conformally symplectic manifold.

## 1. INTRODUCTION

We will define certain rational number valued deformation invariants for certain complete Riemann-Finsler manifolds, in particular for complete Riemannian metrics with non-positive sectional curvature. These invariants can be directly interpreted as a part of certain elliptic Gromov-Witten invariants in an associated lcs manifold, [13]. However, in the more basic setting here, we can reduce the invariants to counts of closed geodesic strings (equivalence classes of closed unit speed geodesics up to reparametrization  $S^1$  action), via Fuller index [5]. And so this self contained more elementary story is developed separately here.

**Terminology 1.** All our metrics are Riemann-Finsler metrics unless specified to be Riemannian, and usually denoted by just  $g$ . Completeness, always means forward completeness. Curvature always means sectional curvature in the Riemannian case and flag curvature in the Finsler case. Thus we will usually just say complete metric  $g$ , for a forward complete Riemann-Finsler metric. A reader may certainly choose to interpret all metrics as Riemannian metrics, completeness as standard completeness, and curvature as sectional curvature.

In what follows  $\pi_1(X)$  denotes the set of free homotopy classes of continuous maps  $o : S^1 \rightarrow X$ .

**Definition 1.1.** Let  $X$  be a smooth manifold. Fix an exhaustion by nested compact sets  $\bigcup_{i \in \mathbb{N}} K_i = X$ ,  $K_i \supset K_{i-1}$  for all  $i \geq 1$ . We say that a class  $\beta \in \pi_1(X)$  is boundary compressible if  $\beta$  is in the image of

$$inc_* : \pi_1(X - K_i) \rightarrow \pi_1(X)$$

for all  $i$ , where  $inc : X - K_i \rightarrow X$  is the inclusion map. We say that  $\beta$  is **boundary incompressible** if it is not boundary compressible.

Let  $\pi_1^{inc}(X)$  denote the set of such boundary incompressible classes. When  $X$  is compact, we set  $\pi_1^{inc}(X) := \pi_1(X) - const$ , where  $const$  denotes the set of homotopy classes of constant loops.

It is easily seen that the above is well defined and moreover any homeomorphism  $X_1 \rightarrow X_2$  of a pair of manifolds induces a set isomorphism  $\pi_1^{inc}(X_1) \rightarrow \pi_1^{inc}(X_2)$ . Denote by  $L_\beta X$  the class  $\beta \in \pi_1^{inc}(X)$

---

*Key words and phrases.* locally conformally symplectic manifolds, Gromov-Witten theory, virtual fundamental class, Fuller index.

component of the free loop space of  $X$ , with its compact open topology. Let  $g$  be a complete metric on  $X$ , and let  $S(g, \beta) \subset L_\beta X$  denote the subspace of all unit speed parametrized, smooth, closed  $g$ -geodesics in class  $\beta$ .

**Definition 1.2.** We say that a metric  $g$  on  $X$  is  $\beta$ -**taut** if it is complete and  $S(g, \beta)$  is compact. We will say that  $g$  is **taut** if it is  $\beta$ -taut for each  $\beta \in \pi_1^{\text{inc}}(X)$ .

A complete metric all of whose boundary incompressible closed geodesics are minimizing in their homotopy class is taut, see Lemma 5.1. Thus, by the Cartan-Hadamard theorem [2], a basic example of a taut metric is a complete metric with non-positive curvature. It should be emphasized that taut metrics form a much larger class of metrics than just non-positive curvature metrics. One class of examples comes by way of Lemma 1.17 ahead, but this only scratches the surface.

**Definition 1.3.** Let  $\beta \in \pi_1^{\text{inc}}(X)$ , and let  $g_0, g_1$  be a pair of  $\beta$ -taut metrics on  $X$ . A  $\beta$ -**taut deformation** between  $g_0, g_1$ , is a continuous (in the topology of  $C^0$  convergence on compact sets) family  $\{g_t\}$ ,  $t \in [0, 1]$  of complete metrics on  $X$ , s.t.

$$S(\{g_t\}, \beta) := \{(o, t) \in L_\beta X \times [0, 1] \mid o \in S(g_t, \beta)\}$$

is compact. We say that  $\{g_t\}$  is a **taut deformation** if it is  $\beta$ -taut for each  $\beta \in \pi_1^{\text{inc}}(X)$ . The above definitions of tautness are extended naturally to the case of a smooth fibration  $X \hookrightarrow P \rightarrow [0, 1]$ , with a smooth fiber-wise family of metrics.

A useful criterion for  $\beta$ -tautness is the following.

**Theorem 1.4.** Let  $\{g_t\}_{t \in [0, 1]}$  be a continuous family of complete metrics on  $X$ . Suppose that:

$$\sup_t \left| \max_{o \in S(g_t, \beta)} l_{g_t}(o) - \min_{o \in S(g_t, \beta)} l_{g_t}(o) \right| < \infty,$$

where  $l_{g_t}$  is the length functional with respect to  $g_t$ , then  $\{g_t\}$  is  $\beta$ -taut. It follows that sky catastrophes of vector fields on closed manifolds are not geodesible by metrics all of whose geodesics are minimal, Appendix A.1.

For example, the hypothesis is trivially satisfied if  $g_t$  have the property that all their class  $\beta$  geodesics are minimal. In particular, if  $g_t$  have non-positive curvature then  $\{g_t\}$  is taut, again by the Cartan-Hadamard theorem.

Fuller at the end of [5] has asked for any metric conditions on vector fields to rule out sky catastrophes, see Appendix A.1. By the above, non-positivity of curvature is one such condition.

*Remark 1.5.* Note that if sky catastrophes were never geodesible then the geodesible Seifert conjecture would follow, by the main result of [12]. Hence, this is a subtle situation and it now seems to be plausible that sky catastrophes for geodesible and Reeb families of vector fields do exist. The qualitative structure of such potential geodesible or Reeb sky catastrophes is somewhat understood by results in [12], and [12, Theorem 1.10] in particular. But this does not greatly aid constructing potential examples, which must be topologically very complex, (there are necessarily infinitely many suitably synchronized bifurcation events). No results prior to the theorem above are known to me aside from those mentioned by Fuller himself in [5].

Let  $\mathcal{G}(X)$  be the set of equivalence classes of taut metrics  $g$ , where  $g_0$  is equivalent to  $g_1$  whenever there is a taut deformation between them. We may denote an equivalence class by its representative  $g$  by a slight abuse of notation.

**Theorem 1.6.** For each manifold  $X$  there is a natural functional

$$F : \mathcal{G}(X) \times \pi_1^{\text{inc}}(X) \rightarrow \mathbb{Q}.$$

The value  $F(g, \beta)$  can be interpreted as a count of the set of closed  $g$ -geodesic strings in class  $\beta$ . But one must take care of exactly how to count, as in general this set should be understood as an orbifold or rather a Kuranishi space of a certain kind, hence this is why  $F$  is  $\mathbb{Q}$  valued.

*Question 1.* Do there exist a pair of taut metrics  $g_1, g_2$  on a manifold  $X$  which are not taut homotopic?

Probably both possibilities are interesting. If the answer is ‘no’ then we may obtain much sharper applications, particularly in the setup of [13]. On the other hand, if the answer is ‘yes’, then the previous Theorem 1.6 becomes far more intriguing.

**Definition 1.7.** Let  $\beta \in \pi_1(X)$ . For any based point  $x_0 \in \text{image } \beta \subset X$  (for  $\text{image } \beta$  the image of some representative of  $\beta$ ) there is a naturally determined element  $\beta_{x_0} \in \pi_1(X, x_0)$  well defined up to an inner automorphism, (concatenate a representative of  $\beta$  with a path from  $x_0$  to a point in  $\text{image } \beta$ ). We say that a class  $\beta \in \pi_1(X, x_0)$  is at **most a  $k$ -power**, if whenever  $\beta = \alpha^n$ , with  $n > 0$  then  $n \leq k$ . Similarly,  $\beta \in \pi_1(X)$  is at most a  $k$ -power if for any  $x_0$  as above,  $\beta_{x_0}$  is at most a  $k$ -power. We say that  $\beta \in \pi_1(X)$  is **not a power** if it is at most a 1-power.

*Example 1.* Let  $g$  be a Riemannian metric with negative sectional curvature on a closed manifold  $X$  and  $\beta \in \pi_1(X)$  a class represented by a multiplicity  $n$  closed geodesic, then

$$(1.8) \quad F(g, \beta) = \frac{1}{n}.$$

In particular, if  $\beta$  is not a power then  $F(g, \beta) = 1$ . More generally, (1.8) holds whenever  $g$  has a unique and non-degenerate closed geodesic string in class  $\beta$ . Here and throughout the paper, a closed geodesic string is **non-degenerate** if the corresponding  $S^1$  family of closed geodesics is Morse-Bott non-degenerate.

**Theorem 1.9.** Every rational number has the form  $F(g, \beta)$  for some  $\beta$ -taut Riemannian  $g$  on some compact manifold  $X$  and for some  $\beta \in \pi_1^{\text{inc}}(X)$ .

If  $\beta \in \pi_1^{\text{inc}}(X)$  is not a power, then it is easy to see that the reparametrization  $S^1$  action on  $L_\beta X$  is free (see Appendix A), so that  $H_*^{S^1}(L_\beta X, \mathbb{Z}) \simeq H_*(L_\beta X/S^1, \mathbb{Z})$ , where  $H_*^{S^1}(L_\beta X, \mathbb{Z})$  denotes the  $S^1$ -equivariant homology. Moreover, we have:

**Theorem 1.10.** Suppose that  $\beta \in \pi_1^{\text{inc}}(X)$  is not a power, and  $X$  admits a  $\beta$ -taut metric, then  $H_*^{S^1}(L_\beta X, \mathbb{Z})$  is finite dimensional. Denote by  $\chi^{S^1}(L_\beta X)$  the Euler characteristic of this homology. Then for any  $\beta$ -taut metric  $g$  on  $X$ :

$$F(g, \beta) = \chi^{S^1}(L_\beta X).$$

Explicit examples for the theorem above can be found by the proof of Theorem 1.9. For these types of examples any negative integer may appear as the value of  $F(g, \beta)$ . We leave out the details.

*Remark 1.11.* If  $\beta$  is a power, the idea behind Theorem 1.10 breaks down, as the  $S^1$ -equivariant homology of  $L_\beta X$  may then be infinite dimensional even if  $X$  admits a  $\beta$ -taut  $g$ . As a trivial example this homology is already infinite dimensional when  $g$  is negatively curved, and the class  $\beta$  geodesic is  $k$ -covered, as then this homology is the group homology of  $\mathbb{Z}_k$ . In particular the connection with the Euler characteristic a priori breaks down. It is thus an interesting open problem if the functional  $F$  remains topological, this is related to question 1.

A celebrated theorem of Preissman [10] says that there are no negative sectional curvature metrics on compact products. Fibration counterexamples to Preissman’s product theorem certainly exist. In fact, every closed 3-manifold  $X^3$ , for which there is no injection  $\mathbb{Z}^2 \rightarrow \pi_1(X, x_0)$ , and which fibers over

a circle has a hyperbolic structure  $g_h$ , Thurston [14]. We are going to give a certain generalization of Preissman's theorem to fibrations, with possibly non-compact fibers, also replacing the negative sectional curvature condition by a significantly weaker condition.

**Definition 1.12.** Let  $Z \hookrightarrow X \xrightarrow{p} Y$  be a smooth fiber bundle with  $X$  having a  $\beta$ -taut Riemannian metric  $g$ , for  $\beta \in \pi_1^{inc}(X)$ , and let  $g_Y$  be a metric on  $Y$ . Suppose that

- (1) The fibers  $Z_y = p^{-1}(y)$  are totally  $g$ -geodesic, for closed geodesics in class  $\beta$ . We denote by  $g_y$  the metric  $g$  restricted to  $Z_y$ .
- (2) The fibers are parallel (the distribution  $T^{vert}X = \ker p_*$  is parallel along any smooth curve in  $X$  with respect to the Levi-Civita connection of  $g$ ).
- (3) For any pair of fibers  $(Z_{y_0}, g_{y_0})$ ,  $(Z_{y_1}, g_{y_1})$ , and a path  $\gamma : [0, 1] \rightarrow Y$  from  $y_0$  to  $y_1$  the fiber family  $\{(Z_{\gamma(t)}, g_{\gamma(t)})\}$  furnishes a taut deformation.
- (4)  $p$  projects  $g$ -geodesics to geodesics of  $Y, g_Y$ .

We then call  $p : X \rightarrow Y$  a  **$\beta$ -taut submersion**, with the metrics  $g, g_Y$  and  $g_Z$  all possibly implicit.

**Definition 1.13.** For  $Z \hookrightarrow X \rightarrow Y$  as above, we say that  $\beta \in \pi_1(X)$  is a **fiber class** if it is in the image of the inclusion  $i_Z : \pi_1(Z) \rightarrow \pi_1(X)$ .

In the above definition of a taut submersion and the following theorem we need the auxiliary metric  $g$  on  $X$  to be Riemannian, and there is no obvious extension of the theorem to the Riemann-Finsler case. However, the conclusions of the theorem are for Riemann-Finsler metrics.

**Theorem 1.14.** Let  $p : (X, g) \rightarrow (Y, g_Y)$  be a  $\beta$ -taut submersion, where  $\beta \in \pi_1^{inc}(X)$  is a fiber class. Suppose further that  $Y$  is connected and closed, and is such that all smooth closed contractible  $g_Y$  geodesics in  $Y$  are constant, then the following holds.

- If  $\chi(Y) \neq \pm 1$  and  $\beta$  is at most a  $k$ -power then  $X$  does not admit a complete Riemann-Finsler metric with a unique and non-degenerate class  $\beta$  geodesic string.
- If  $\chi(Y) \neq \pm 1$  then  $X$  does not admit a complete Riemann-Finsler metric with negative curvature.

Note that  $\chi(Y) \neq 1$  is of course essential, as the trivial fibration  $X \rightarrow \{pt\}$ , with  $X$  admitting a complete negatively curved metric, will satisfy the hypothesis. The condition that there is a fiber class  $\beta \in \pi_1^{inc}(X)$  is also essential, for any vector bundle over a manifold admitting a Riemannian metric of negative curvature admits a metric of negative curvature, Anderson [1].

**Corollary 1.15.** Let  $X = Z \times Y$  where  $Z, Y$  admit complete Riemannian metrics with non-positive sectional curvature,  $Y$  is closed and  $\beta \in \pi_1^{inc}(X)$  is in the image of the inclusion  $\pi_1(Z) \rightarrow \pi_1(X)$ .

- (1) If  $\chi(Y) \neq \pm 1$  and  $\beta$  is at most a  $k$ -power then  $X$  does not admit a complete (Riemann-Finsler) metric with a unique and non-degenerate class  $\beta$  geodesic string.
- (2) If  $\chi(Y) \neq \pm 1$  then  $X$  does not admit a complete metric of negative curvature.

A basic set of examples for the theorem is obtained by starting with any homomorphism

$$(1.16) \quad \phi : \pi_1(Y, y_0) \rightarrow \text{Isom}(Z, g_Z), \quad (\text{the group of all isometries}).$$

where  $g_Z$  is a taut Riemannian metric, and there is a class  $\beta_Z \in \pi_1^{inc}(Z)$ . Suppose further:

(1) The orbit

$$O := \bigcup_{\gamma \in \pi_1(Y, y_0)} \phi_*(\gamma)(\beta_Z)$$

is finite.

(2)  $Y$  is closed and connected.

(3) All contractible smooth closed  $g_Y$  geodesics in  $Y$  are constant.

We have the obvious induced diagonal action

$$\pi_1(Y, y_0) \rightarrow \text{Diff}(Z \times \tilde{Y}), \text{ (the group of all diffeomorphisms),}$$

$$\gamma \mapsto ((z, y) \mapsto (\phi(\gamma)(z), \gamma \cdot y)),$$

for  $\tilde{Y}$  the universal cover of  $Y$ . Taking the quotient of  $Z \times \tilde{Y}$  by this action, we get an associated “flat” bundle  $Z \hookrightarrow X_\phi \xrightarrow{p} Y$ , with a metric  $g_\phi$  induced from the product metric  $\tilde{g} = g_Z \oplus g_Y$ , on the covering space  $q : Z \times \tilde{Y} \rightarrow Z \times Y$ .

**Lemma 1.17.** *Let  $p : (X_\phi, g_\phi) \rightarrow (Y, g_Y)$  be as above, then this is a  $\beta$ -taut submersion, where  $\beta = i_*(\beta_Z)$ , for  $i_* : \pi_1^{inc}(Z) \rightarrow \pi_1^{inc}(X_\phi)$  induced by inclusion.*

By the lemma above,  $p : (X_\phi, g_\phi) \rightarrow (Y, g_Y)$  satisfies the hypothesis of the theorem above. Yet more concretely:

*Example 2.* Suppose we have  $\beta_Z \in \pi_1^{inc}(Z)$ , and let  $\phi : Z \rightarrow Z$  be an isometry of a taut metric  $g_Z$ . Then by the construction above, the mapping torus

$$(Z, g_Z) \hookrightarrow (X_\phi, g_\phi) \xrightarrow{\pi} S^1$$

has the structure of a  $\beta$ -taut submersion, satisfying the hypothesis of the theorem, for  $\beta = i_*(\beta_Z)$  as above.

The next corollary of Theorem 1.14 is immediate.

**Corollary 1.18.** *Let*

$$(Z_g, g_Z) \hookrightarrow (X_\phi, g_\phi) \rightarrow (Y, g_Y)$$

*be as in the construction above for  $Z, g_Z$  having non-positive curvature, and let  $\beta_Z \in \pi_1^{inc}(Z)$ . Then*

- (1) *If  $\chi(Y) \neq \pm 1$ ,  $X_\phi$  does not admit a complete Riemann-Finsler metric with negative curvature.*
- (2) *If  $\chi(Y) \neq \pm 1$ , and  $\beta_Z$  is at most a  $k$ -power, then  $X_\phi$  does not admit a Riemann-Finsler metric with a unique and non-degenerate class  $\beta$  geodesic string, for  $\beta = i_*(\beta_Z)$  as above.*

*As a special case, this applies to the mapping tori  $X_\phi$ , for  $\phi : Z \rightarrow Z$  an isometry of a complete Riemannian non-positively curved metric on  $Z$ , satisfying the finiteness condition 1. (The non-positive curvature hypothesis is for concreteness we may of course replace this condition by tautness.)*

In the special case when  $Z$  is compact, the first part of the above corollary readily follows by Preissman’s theorem (specifically, because of the condition 1), see also [3, Theorem 9.3.4] for a generalization which fits our Finsler setting. The second part is new even when  $Z$  is compact. In the non-compact case, as far as I know, the above corollary and the more basic Corollary 1.15, are the only presently known extensions of Preissman’s theorem.

## 2. PROOF OF THEOREM 1.4

The first part of the theorem clearly follows by the second part. So let  $\{g_t\}$ ,  $t \in [0, 1]$  be as in the hypothesis, with

$$\sup_t \left| \max_{o \in S(g_t, \beta)} l_{g_t}(o) - \min_{o \in S(g_t, \beta)} l_{g_t}(o) \right| < c,$$

and suppose that

$$\sup_{(o,t) \in \mathcal{O}(\{g_t\}, \beta)} l_{g_t}(o) = \infty.$$

Then we have a sequence  $\{o_k\}$ ,  $k \in \mathbb{N}$ , of closed class  $\beta$   $g_{t_k}$ -geodesics in  $X$ , satisfying:

- (1)  $\lim_{k \rightarrow \infty} t_k = t_\infty \in [0, 1]$ .
- (2)  $\lim_{k \rightarrow \infty} l_{g_{t_k}}(o_k) = \infty$ , where  $l_{g_{t_k}}(o_k)$  is the length with respect to  $g_{t_k}$ .

Let  $o_\infty$  be a minimal, class  $\beta$ ,  $g_\infty = g_{t_\infty}$  geodesic in  $X$ , with  $g_\infty$  length  $L$ . Let  $g_{aux}$  be a fixed auxiliary metric on  $X$ , and let  $L_{aux}$  be the  $g_{aux}$  length of  $o_\infty$ .

Define a pseudo-metric on the space of metrics on  $X$  as follows. Let  $K \subset X$  be a fixed compact set containing image  $o_\infty$ , and set

$$V \subset TX = \{v \in TX \mid \pi(v) \in K \text{ for } \pi : TX \rightarrow X \text{ the canonical projection, and } |v|_{aux} = 1\},$$

where  $|v|_{aux}$  is the norm taken with respect to  $g_{aux}$ .

Then define:

$$d_{C^0}(g_1, g_2) = \sup_{v \in V} ||v|_{g_1} - |v|_{g_2}|.$$

By properties 1 and 2 we may find a  $k > 0$  such that

$$(2.1) \quad d_{C^0}(g_{t_k}, g_{t_\infty}) < \epsilon$$

and

$$l_{g_{t_k}}(o_k) > c + L + L_{aux} \cdot \epsilon.$$

As  $L + L_{aux} \cdot \epsilon \geq l_{g_{t_k}}(o_\infty)$  by (2.1), we have:

$$l_{g_{t_k}}(o_k) > l_{g_{t_k}}(o_\infty) + c.$$

Since we may find a closed  $g_{t_k}$ -geodesic  $o'$  satisfying  $l_{g_{t_k}}(o') \leq l_{g_{t_k}}(o_\infty)$ , we get that

$$\left| \max_{o \in S(g_{t_k}, \beta)} l_{g_{t_k}}(o) - \min_{o \in S(g_{t_k}, \beta)} l_{g_{t_k}}(o) \right| > c,$$

and so we are in contradiction.

Thus,

$$\sup_{(o,t) \in \mathcal{O}(\{g_t\}, \beta)} l_{g_t}(o) < \infty.$$

It follows, by an analogue of Lemma 5.2, that the images of all elements  $o \in S(\{g_t\}, \beta)$  are contained in a fixed compact  $K \subset X$ . Compactness of  $S(\{g_t\}, \beta)$  then follows by the Arzella-Ascoli theorem.  $\square$

## 3. PROOF OF LEMMA 1.17

Let  $\phi_* : \pi_1(Y, y_0) \rightarrow \text{Aut}(\pi_1^{inc}(Z))$  be the natural induced action, where  $\text{Aut}(\pi_1^{inc}(Z))$  denotes the group of set isomorphisms of  $\pi_1^{inc}(Z)$ . And such that the orbit

$$O := \bigcup_{\gamma \in \pi_1(Y, y_0)} \phi_*(\gamma)(\beta_Z)$$

is finite.

As  $g_Z$  is taut,  $S(g_Z, \phi_*(\gamma)(\beta_Z))$  is compact for each  $\gamma$ , where  $S(g_Z, \phi_*(\gamma)(\beta_Z))$  is the space of geodesics as in Definition 1.2. By the condition on contractible geodesics of  $g_Y$ , we get:

$$\begin{aligned} S(g_\phi, \beta) &= q_*(S(g_Z \oplus g_Y, \beta)) \\ &= \bigcup_{\beta \in O} q_*(S(g_Z, \beta) \times \tilde{Y}), \end{aligned}$$

for  $q_* : L(Z \times \tilde{Y}) \rightarrow L(Z \times Y)$  induced by the quotient map  $q : Z \times \tilde{Y} \rightarrow Z \times Y$ , (as in the preamble to the statement of the lemma) and where  $S(g_Z, \gamma) \times \tilde{Y}$  is identified as a subset  $S(g_Z, \gamma) \times \tilde{Y} \subset L(Z) \times \tilde{Y} \subset L(Z \times \tilde{Y})$ . Given that  $O$  is finite, this then readily implies our claim.  $\square$

## 4. PRELIMINARIES ON REEB FLOW

Let  $(C^{2n+1}, \lambda)$  be a contact manifold with  $\lambda$  a contact form, that is a one form s.t.  $\lambda \wedge (d\lambda)^n \neq 0$ . Denote by  $R^\lambda$  the Reeb vector field satisfying:

$$d\lambda(R^\lambda, \cdot) = 0, \quad \lambda(R^\lambda) = 1.$$

Recall that a **closed  $\lambda$ -Reeb orbit** (or just Reeb orbit when  $\lambda$  is implicit) is a smooth map

$$o : (S^1 = \mathbb{R}/\mathbb{Z}) \rightarrow C$$

such that

$$\dot{o}(t) = cR^\lambda(o(t)),$$

with  $\dot{o}(t)$  denoting the time derivative, for some  $c > 0$  called period. Let  $S(R^\lambda, \beta)$  denote the space of all closed  $\lambda$ -Reeb orbits in free homotopy class  $\beta$ , with its compact open topology. And set

$$\mathcal{O}(R^\lambda, \beta) = S(R^\lambda, \beta)/S^1,$$

where  $S^1 = \mathbb{R}/\mathbb{Z}$  acts by reparametrization  $t \cdot o(\tau) = o(t + \tau)$ .

## 5. DEFINITION OF THE FUNCTIONAL F AND PROOFS OF AUXILIARY RESULTS

Let  $X$  be a manifold with a taut metric  $g$ . Let  $C$  be the unit cotangent bundle of  $X$ , with its Liouville contact 1-form  $\lambda_g$ . If  $o : S^1 = \mathbb{R}/\mathbb{Z} \rightarrow X$  is a unit speed closed geodesic, it has a canonical lift  $\tilde{o} : S^1 \rightarrow C$ . If  $\beta \in \pi_1^{inc}(X)$ , let  $\tilde{\beta} \in \pi_1(C)$  denote class  $[\tilde{o}] \in \pi_1(C)$ , where  $o$  is a unit speed closed geodesic representing  $\beta$ .

Let  $S(R^{\lambda_g}, \tilde{\beta})$  be the orbit space as in Section 4, for the Reeb flow of the contact form  $\lambda_g$ . And set

$$\mathcal{O}_{g, \beta} = \mathcal{O}(R^{\lambda_g}, \tilde{\beta}) := S(R^{\lambda_g}, \tilde{\beta})/S^1,$$

i.e. this can be identified with the space of class  $\beta$   $g$ -geodesic strings. By the tautness assumptions  $\mathcal{O}_{g, \beta}$  is compact.

We then define

$$F(g, \beta) = i(\mathcal{O}_{g, \beta}, R^{\lambda_g}, \tilde{\beta}) \in \mathbb{Q}$$

where the right hand side is the Fuller index of  $R^{\lambda_g}$  in class  $\tilde{\beta}$ . As a basic example we have:

**Lemma 5.1.** *Suppose that  $g$  is a complete metric on  $X$ , all of whose boundary incompressible geodesics are minimal, then  $g$  is taut.*

*Proof.* First we state a more basic lemma.

**Lemma 5.2.** *Suppose that  $g$  is a complete metric on  $X$ ,  $\beta \in \pi_1^{inc}(X)$  and let  $S \subset L_\beta X$  be a subset on which the  $g$ -length functional is bounded from above. Then the images in  $X$  of elements of  $S$  are contained in a fixed compact subset of  $X$ .*

*Proof.* Suppose otherwise. Fix an exhaustion by nested compact sets

$$\bigcup_{i \in \mathbb{N}} K_i = X, \quad K_i \supset K_{i-1}.$$

Then either there is sequence  $\{o_i\}_{i \in \mathbb{N}}$ ,  $o_i \in S$  s.t.  $o_i \in K_i^c$ , for  $K_i^c$  the complement of  $K_i$ , which contradicts the fact that  $\beta$  is incompressible. Or there is a sequence  $\{o_k\}_{k \in \mathbb{N}}$ ,  $o_k \in S$  s.t.:

- (1) Each  $o_k$  intersects  $K_{i_0}$  for some  $i_0$  fixed.
- (2) For each  $i \in \mathbb{N}$  there is a  $k_i > i$  s.t.  $o_{k_i}$  is not contained in  $K_i$ .

Now if  $\text{diam}(o_k)$  is bounded in  $k$ , then condition 1 implies that  $o_k$  are contained in a set of bounded diameter. (Here  $\text{diam}(o_k)$  denotes the diameter of image  $o_k$ .) Consequently, by Hopf-Rinow theorem [2],  $o_k$  are contained in a compact set. But this contradicts condition 2, and the fact that  $K_i$  form an exhaustion of  $X$ .

Thus, we conclude that  $\text{diam}(o_k)$  is unbounded, but this contradicts the hypothesis.  $\square$

Returning to the proof of the main lemma. By assumption, closed, class  $\beta \in \pi_1^{inc}(X)$  geodesics are  $g$ -minimizing in their homotopy class and in particular have fixed length. By the lemma above there is a fixed  $K \subset X$  s.t. every class  $\beta$  closed geodesic has image contained in  $K$ . Then compactness of  $S(g, \beta)$  follows by Arzella-Ascoli theorem.  $\square$

*Proof Theorem 1.6.* Let  $\beta \in \pi_1^{inc}(X)$ , be given and let  $g$  be  $\beta$ -taut. We just need to prove that  $F(g, \beta)$  is invariant under a  $\beta$ -taut deformation of  $g$ . So let  $\{g_t\}$ ,  $t \in [0, 1]$  be a  $\beta$ -taut deformation of metrics on a compact manifold  $X$ . Let  $R^{\lambda_{g_t}}$  be the geodesic flow on the  $g_t$  unit cotangent bundle  $C_t$ . Trivializing the family  $\{C_t\}$  we get a family  $\{R_t\}$  of flows on  $C \simeq C_t$ , with  $R_t$  conjugate to  $R^{\lambda_{g_t}}$ .

Let  $\mathcal{O}(\{R_t\}, \tilde{\beta})$  be the cobordism as in (A.2), where  $\tilde{\beta} \in \pi_1(C)$  is as above. Then  $\mathcal{O}(\{R_t\}, \tilde{\beta})$  is compact as  $S(\{g_t\}, \beta)$  is compact by assumption.

Basic invariance of the Fuller index, that is (A.3), immediately yields:  $F(g_0, \beta) = F(g_1, \beta)$ .  $\square$

*Proof of Theorem 1.10.* This is an application of Morse theory. As  $g$  is  $\beta$ -taut, and so  $S(g, \beta)$  is compact, we may find a  $C^\infty$ -nearby metric  $g'$ , s.t.  $g'$  has finitely many class  $\beta$  closed geodesic strings, all of which are non-degenerate. The notation  $L_\beta X$  now denotes the Hilbert manifold of  $H^1$  loops, as used for example in the classical work of Gromoll-Meyer [6]. This Hilbert manifold is well known to be homotopy equivalent to the standard free loop space with its compact open topology.

The energy function  $e_{g'} : L_\beta X \rightarrow S^1$ ,

$$(5.3) \quad e_{g'}(o) = \int_{S^1} \langle \dot{o}(t), \dot{o}(t) \rangle_{g'} dt$$

is smooth,  $S^1$  invariant and satisfies the Palais-Smale condition. The flow for its negative gradient vector field  $V$  is complete, and we can do Morse theory mostly as usual. This is understood starting with the work of Klingenberg [7], with the framework of Palais and Smale [9]. In our case,  $e_{g'}$  is moreover a Morse-Bott function with critical manifolds  $C_o$  corresponding to  $S^1$  families of geodesics, for each closed geodesic string  $o$ .



There is an induced Morse-Bott cell decomposition on  $L_\beta X$ , meaning a stratification formed by  $V$  unstable manifolds of the above mentioned critical manifolds  $C_o$ . This is Bott's extension of the fundamental Morse decomposition theorem. Now the  $S^1$  action on  $L_\beta X$  is free by the condition that  $\beta$  is not a power. This action is not smooth, but is continuous and so taking the topological  $S^1$  quotient we get a standard CW cell decomposition of  $L_\beta X/S^1$  with one  $k$ -cell for each closed  $g'$ -geodesic string  $o$  of index  $\text{morse}(o) = k$ . (The "index" means the Morse-Bott index of the critical manifold  $C_o$ .) All of the above is understood, see for instance [6].

From the above cell decomposition, we readily get that the homology  $H_*(L_\beta X/S^1, \mathbb{Z}) = H_*^{S^1}(L_\beta X, \mathbb{Z})$  is finite dimensional. We also get that:

$$\begin{aligned} \chi(L_\beta X/S^1) &= \sum_{o \in \mathcal{O}(R^{\lambda_{g'}}, \beta)} -1^{\text{morse}(o)} \quad (\text{immediate from the cell decomposition}) \\ &= F(g', \beta) \quad (\text{basic properties of the fixed point index, see for instance [12, Section 2]}) \\ &= F(g, \beta) \quad (\text{by the local invariance (A.4) of the Fuller index}). \end{aligned}$$

□

## 6. PROOF OF THEOREM 1.14 AND ITS COROLLARIES

We first prove:

**Theorem 6.1.** *Let  $p : X \rightarrow Y$  be a  $\beta$ -taut submersion as in the statement of Theorem 1.14 and  $\beta \in \pi_1^{\text{inc}}(X)$  a fiber class. Then*

$$(6.2) \quad F(g, \beta) = \text{card} \cdot \chi(Y) \cdot F(g_Z, \beta_Z),$$

where  $\text{card} \in \mathbb{N} - \{0\}$  is the cardinality of a certain orbit of the holonomy group (as explained in the proof), and where  $\beta_Z$  is as in Lemma 1.17.

*Proof.* We have a natural subset of  $\mathcal{O}' \subset \mathcal{O}g, \beta$ , consisting of all vertical geodesics, that is  $g$ -geodesics contained in fibers  $p^{-1}(y) = Z_y$ . In fact,  $\mathcal{O}' = \mathcal{O}g, \beta$ , for if  $o$  is any class  $\beta$  geodesic, the projection  $p(o)$  is a contractible  $g_Y$  geodesic, and by assumptions is constant.

In particular there a natural continuous projection  $\tilde{p} : \mathcal{O}g, \beta \rightarrow Y$ ,  $\tilde{p}(o) = y$  where  $y$  is determined by the condition that  $Z_y \supset \text{image } o$ . We will use this to construct a suitable (in a sense abstract i.e. not Reeb) perturbation of the vector field  $R^{\lambda_g}$ , using which we can calculate the invariant  $F(g, \beta)$ .

Fix a Morse function on  $f$  on  $Y$ , let  $C = S^*X$  denote the  $g$ -unit cotangent bundle of  $X$ . For  $v \in T_x X$  let  $\langle v|$  denote the functional  $T_x X \rightarrow \mathbb{R}$ ,  $w \mapsto \langle v, w \rangle_g$ . Define  $\tilde{f} : C \rightarrow \mathbb{R}$  by  $\tilde{f}(\langle v|) := f(p(v))$ , also define  $P : C \rightarrow \mathbb{R}$  by  $P(\langle v|) := |P^{\text{vert}}(v)|_g^2$ , where  $P^{\text{vert}}(v)$  denotes the  $g$ -orthogonal projection of  $v$  onto the  $T_x^{\text{vert}} X \subset T_x X$ , for  $T^{\text{vert}} X$  the vertical tangent bundle of  $X$ , i.e. the kernel of the map  $p_* : TX \rightarrow TY$ .

Next define  $F : C \rightarrow \mathbb{R}$  by:

$$F(\langle v|) := P(\langle v|) + \tilde{f}(\langle v|).$$

Set

$$V_t = R^{\lambda_g} - t \text{grad}_{g_S} F,$$

where the gradient is taken with respect to the Sasaki metric  $g_S$  on  $C$  [11] induced by  $g$ . The latter Sasaki metric is the natural metric for which we have an orthogonal splitting  $TC = T^{\text{vert}} C \oplus T^{\text{hor}} C$ , where  $T^{\text{vert}} C$  is the kernel of  $pr_* : TC \rightarrow TX$ , induced by the natural projection  $pr : C \rightarrow X$ , and where  $T^{\text{hor}} C$  is the  $g$  Levi-Civita horizontal sub-bundle.

Set  $\mathcal{O}_t = \mathcal{O}(V_t, \tilde{\beta})$ , where  $\tilde{\beta}$  is as in Section 5.

**Lemma 6.3.** (1) For all  $t \in [0, 1]$ ,  $N_t := \mathcal{O}_t \cap \mathcal{O}_{g,\beta}$  is open and closed in  $\mathcal{O}_t$ .

(2) For all  $t \in (0, 1]$ ,  $N_t = \cup_{y \in \text{crit}(f)} \tilde{p}^{-1}(y)$ , where  $\text{crit}(f)$  is the set of critical points of  $f$ .

*Proof.* It is easy to see that  $V_t$  is complete and without zeros. Suppose that  $t > 0$ . Let  $\langle v_\tau |$ ,  $\tau \in \mathbb{R}$  be the flow line of  $V_t$ , through  $\langle v_0 |$ , i.e.  $\langle v_\tau | = \phi_\tau(\langle v_0 |)$ , for  $\phi_\tau$  the time  $\tau$  flow map of  $V_t$ . By the fact that the fibers of  $p$  are assumed to be parallel, we have that

$$R^{\lambda_g}(P) = 0, \quad \text{using the derivation notation.}$$

Also,

$$\text{grad}_{g_S} \tilde{f}(P) = 0,$$

which readily follows by the conjunction of  $g_S$  being Sasaki and the fibers of  $p$  being parallel. Consequently, the function

$$\tau \mapsto P(\langle v_\tau |) = |P^{\text{vert}}(v_\tau)|_g^2$$

is monotonically decreasing unless either:

- (1)  $v_0$  is tangent to  $T^{\text{vert}}X$ , in which case for all  $\tau$ ,  $v_\tau$  are tangent to  $T^{\text{vert}}X$  and  $|P^{\text{vert}}(v_\tau)|_g^2 = 1$ .
- (2) For all  $\tau$ ,  $|P^{\text{vert}}(v_\tau)|_g^2 = 0$ .

In particular, the closed orbits of  $V_t$  split into two types.

- (1) Closed orbits  $o(\tau) = \langle v_\tau |$  with  $v_\tau$  always tangent to  $T^{\text{vert}}X$ . In this case we may immediately, conclude that  $o$  is a lift to  $C$  of a closed  $g$ -geodesic contained in the fiber over a critical point of  $f$ .
- (2) Closed orbits  $o(\tau) = \langle v_\tau |$  for which  $v_\tau$  is always  $g$ -orthogonal to  $T^{\text{vert}}X$ .

Clearly, the conclusion follows. □

*Remark 6.4.* It would be very fruitful to remove the condition on the fibers of  $p$  being parallel. But our argument would need to substantially change.

We return to the proof of the theorem. Set

$$\tilde{N} = \{(o, t) \in L_{\tilde{\beta}}C \times [0, \epsilon] \mid o \in N_t\},$$

where  $L_{\tilde{\beta}}C$  denotes the  $\tilde{\beta}$  component of the free loop space as previously. By part I of Lemma 6.3, this is an open compact subset of  $\mathcal{O}(\{V_t\}, \tilde{\beta})$  s.t.

$$\tilde{N} \cap (L_{\tilde{\beta}}C \times \{0\}) = \mathcal{O}(R^{\lambda_g}, \tilde{\beta}),$$

(equalities throughout are up to natural set theoretic identifications.)

By definitions:

$$N_t = \tilde{N} \cap (L_{\tilde{\beta}}C \times \{t\}).$$

Now the invariance of the Fuller index gives:

$$i(N_0, R^{\lambda_g}, \tilde{\beta}) = i(N_1, V_1, \tilde{\beta}).$$

We proceed to compute the right hand side. Fix any smooth Ehresmann connection  $\mathcal{A}$  on the fiber bundle  $p : X \rightarrow Y$ . This induces a holonomy homomorphism:

$$\text{hol}_y : \pi_1(Y, y) \rightarrow \text{Aut } \pi_1(Z_y) \quad (\text{the right-hand side is the group of set automorphisms}),$$

with image denoted  $\mathcal{H}_y \subset \text{Aut } \pi_1(Z_y)$ .

Let  $\beta_Z$  denote a class in  $\pi_1(Z_y)$  s.t.  $(i_{Z_y})_*(\beta_Z) = \beta$ , for  $i_{Z_y} : Z_y \rightarrow X$  the inclusion map. Set

$$S_y := \bigcup_{g \in \mathcal{H}_y} g(\beta_Z) \subset \pi_1(Z_y).$$

Then for another  $y' \in Y$ ,

$$(6.5) \quad h_* : S_{y'} \rightarrow S_y,$$

is an isomorphism, where  $h : Z_{y'} \rightarrow Z_y$  is a smooth map given by the  $\mathcal{A}$ -holonomy map determined by some path from  $y$  to  $y'$ , and where  $h_*$  is the naturally induced map.

Denoting by  $g_y$  the restriction of  $g$  to the fiber  $Z_y$ , let  $R^y$  denote the  $\lambda_{g_y}$  Reeb vector field on the  $g_{Z_y}$ -unit cotangent bundle  $C_y$  of  $Z_y$ . The cardinality  $\text{card}$  of  $S_y$  is finite, as otherwise we get a contradiction to the compactness of  $S(g, \beta)$ . Now

$$\tilde{p}^{-1}(y) = \bigcup_{\alpha \in S_y} \mathcal{O}(R^{\lambda_y}, \alpha).$$

From part 2 of Lemma 6.3 and by straightforward index computations we get:

$$i(N_1, V_1, \tilde{\beta}) = \sum_{y \in \text{crit}(f)} (-1)^{\text{morse}(y)} \cdot i(\tilde{p}^{-1}(y), R^{\lambda_y}, \tilde{\beta}),$$

where  $\text{morse}(y)$  denote the Morse index of  $y$ . Now

$$\begin{aligned} i(\tilde{p}^{-1}(y), R^{\lambda_y}, \tilde{\beta}) &= \sum_{\alpha \in S_y} i(\mathcal{O}(R^{\lambda_y}), R^{\lambda_y}, \tilde{\alpha}) \\ &= \sum_{\alpha \in S_y} F(g_y, \alpha). \\ &= \text{card} \cdot F(g_Z, \beta_Z), \end{aligned}$$

where the last equality follows by (6.5), and by the condition 3 in the Definition 1.2. And so the result follows.  $\square$

Now returning to the proof of the main theorem. As  $\beta$  is at most a  $k$ -power,  $\beta = \alpha^n$ , for some  $n$  where  $\alpha$  is not a power. By the assumption that all contractible  $g_Y$  geodesics are constant, classical Morse theory Milnor [8] tells us that  $Y$  has vanishing higher homotopy group  $\pi_k(Y, y_0)$ ,  $k \geq 2$ , and in particular  $i_{Z,*} : \pi_1(Z, p_0) \rightarrow \pi_1(X, p_0)$  is a group injection, by the long exact sequence of a fibration. It follows that  $\alpha \in \pi_1^{\text{inc}}(X)$  is also a fiber class.

Now, any  $\alpha$ -class closed  $g$ -geodesic must be tangent a fiber of  $p$ , as otherwise clearly  $p$  would not be  $\beta$ -taut. It follows that  $p : X \rightarrow Y$  is also  $\alpha$ -taut.

Now, to prove the first part of the conclusion, if  $\chi(Y) \neq \pm 1$  then by (6.2)  $F(g, \alpha) \neq \pm 1$ , since  $F(g_Z, \alpha_Z)$  is an integer by Theorem 1.10. And also by Theorem 1.10

$$F(g, \alpha) = \chi^{S^1}(L_\alpha X),$$

so that  $X$  cannot admit a complete metric with a unique and non-degenerate class  $\alpha$ -geodesic string as otherwise  $\chi^{S^1}(L_\alpha X) = 1$  (see Proof of Theorem 1.10). This also readily implies that  $X$  does not admit a metric with a unique and non-degenerate class  $\beta$ -geodesic string. So that the first conclusion follows.

To prove the second conclusion, it is enough to show that if  $X$  admits a complete metric  $g'$  of negative curvature then a class  $\beta \in \pi_1^{\text{inc}}(X)$  must be at most a  $k$ -power for some  $k$ . As then the needed result follows by the first conclusion.

Let then  $o$  be the unique, closed, class  $\beta$   $g'$ -geodesic string  $X$ . Then  $o$  covers a simple, closed, fiber class geodesic string  $\tilde{o}$  in class  $\tilde{\beta} \in \pi_1^{\text{inc}}(X)$ . We claim that  $\tilde{\beta}$  is not a power. Suppose otherwise, so

that  $\tilde{\beta}_{x_0} = \alpha^k$  for  $k > 1$  and  $\alpha \in \pi_1(X, x_0)$ , ( $\beta_{x_0}$  is as in Definition 1.7). Let  $u$  be the unique, class  $\alpha$ , closed  $g'$ -geodesic string (where  $\alpha$  also denotes the free homotopy class corresponding to the based class  $\alpha$ .) It is immediate that the  $k$  cover of  $u$ ,  $u^k$  represents  $\tilde{\beta}$  and is a closed  $g$ -geodesic string. As  $g'$  is negatively curved, we must have  $u^k = o$ , which gives a contradiction.

□

*Proof of Corollary 1.15.* Let  $g_Z, g_Y$  be complete Riemannian metrics on  $Z, Y$  with non-positive curvature. Take the product metric  $g = g_Z \times g_Y$  on  $X = Z \times Y$ . By Lemma 1.17 and the Cartan-Hadamard theorem,  $g$  is taut. Moreover, for a class  $\beta \in \pi_1(Z)$  in the image of the inclusion  $\pi_1(Z) \rightarrow \pi_1(X)$  the natural projection  $X \rightarrow S^1$  is automatically a  $\beta$ -taut submersion. Then the conclusion readily follows from Theorem 1.14.

□

*Proof of Theorem 1.9.* By Theorem 6.1 0 is certainly a value of the invariant  $F$ . We first prove that every negative rational number is the value of the invariant. Let  $p, q$  be positive integers. Let  $Y$  be a closed surface of genus  $(p+1) > 1$  with a hyperbolic metric  $g_Y$ , let  $Z$  be the genus 2 closed surface with a hyperbolic metric  $g_Z$  and let  $\beta_Z \in \pi_1^{inc}(Z)$  be the class represented by a  $2 \cdot q$ -fold covering of a simple loop representing a generator of the fundamental group of  $Z$ .

Let  $X = Y \times Z$  with the product metric  $g = g_Y \times g_Z$  and  $p : X \rightarrow Y$  the canonical projection. By Theorem 6.1

$$F(g, \beta) = \chi(Y) \cdot F(g_Z, \beta_Z) = (-2p) \cdot \frac{1}{2q} = -\frac{p}{q},$$

where  $\beta$  is as in Lemma 1.17. So we proved our first claim.

Let again  $p, q$  be positive integers. Let  $Y$  be closed surface of genus 2, with a hyperbolic metric  $g_Y$ . And let  $Z$  be a manifold satisfying  $F(g_Z, \beta_Z) = -\frac{p}{2q}$  for some  $\beta_Z$ -taut metric  $g_Z$  on  $Z$  and for some class  $\beta_Z \in \pi_1^{inc}(Z)$ . This exist by the discussion above. Let  $g = g_Y \times g_Z$  be the product metric on  $Y \times Z$ , and  $\beta$  as above. Analogously to the discussion above we get:

$$F(g, \beta) = \chi(Y) \cdot F(g_Z, \beta_Z) = (-2) \cdot \frac{-p}{2q} = \frac{p}{q}.$$

□

#### A. FULLER INDEX AND SKY CATASTROPHES

Let  $X$  be a complete vector field without zeros on a manifold  $M$ . Set

$$(A.1) \quad S(X, \beta) = \{o \in L_\beta M \mid \exists p \in (0, \infty), \ o : \mathbb{R}/\mathbb{Z} \rightarrow M \text{ is a periodic orbit of } pX\}.$$

The above  $p$  is uniquely determined and we denote it by  $p(o)$  called the period of  $o$ .

There is a natural  $S^1$  reparametrization action on  $S(X, \beta)$ :  $t \cdot o$  is the loop  $t \cdot o(\tau) = o(t + \tau)$ . The elements of  $\mathcal{O}(X, \beta) := S(X, \beta)/S^1$  will be called **orbit strings** or just closed orbits. Slightly abusing notation we just write  $o$  for the equivalence class of  $o$ .

The multiplicity  $m(o)$  of an orbit string is the ratio  $p(o)/l$  for  $l > 0$  the period of a simple orbit string covered by  $o$ .

We want a kind of fixed point index which counts orbit strings  $o$  with certain weights. Assume for simplicity that  $N \subset \mathcal{O}(X, \beta)$  is finite. (Otherwise, for a general open compact  $N \subset \mathcal{O}(X, \beta)$ , we need to perturb.) Then to such an  $(N, X, \beta)$  Fuller associates an index:

$$i(N, X, \beta) = \sum_{o \in N} \frac{1}{m(o)} i(o),$$

where  $i(o)$  is the fixed point index of the time  $p(o)$  return map of the flow of  $X$  with respect to a local surface of section in  $M$  transverse to the image of  $o$ .

Fuller then shows that  $i(N, X, \beta)$  has the following invariance property. For a continuous homotopy  $\{X_t\}$ ,  $t \in [0, 1]$  set

$$S(\{X_t\}, \beta) = \{(o, t) \in L_\beta M \times [0, 1] \mid o \in S(X_t)\}.$$

And given a continuous homotopy  $\{X_t\}$ ,  $X_0 = X$ ,  $t \in [0, 1]$ , suppose that  $\tilde{N}$  is an open compact subset of

$$(A.2) \quad \mathcal{O}(\{X_t\}, \beta) := S(\{X_t\}, \beta) / S^1,$$

such that

$$\tilde{N} \cap (L_\beta M \times \{0\}) / S^1 = N.$$

Then if

$$N_1 = \tilde{N} \cap (L_\beta M \times \{1\}) / S^1$$

we have

$$(A.3) \quad i(N, X, \beta) = i(N_1, X_1, \beta).$$

We call this **basic invariance**. In the case  $\mathcal{O}(X_0, \beta)$  is compact,  $\mathcal{O}(X_1, \beta)$  is compact for any sufficiently  $C^0$  nearby  $X_1$ , and in this case basic invariance implies (see for instance [12, Proof of Lemma 1.6]):

$$(A.4) \quad i(\mathcal{O}(X_0, \beta), X, \beta) = i(\mathcal{O}(X_1, \beta), X_1, \beta).$$

### A.1. Blue sky catastrophes.

**Definition A.5** (Preliminary). A **sky catastrophe** for a smooth family  $\{X_t\}$ ,  $t \in [0, 1]$ , of non-vanishing vector fields on a closed manifold  $M$  is a continuous family of closed orbit strings  $\tau \mapsto o_{t_\tau}$ ,  $o_{t_\tau}$  is an orbit string of  $X_{t_\tau}$ ,  $\tau \in [0, \infty)$ , such that the period of  $o_{t_\tau}$  is unbounded from above.

A sky catastrophe as above was initially constructed by Fuller [4]. Or rather his construction essentially contained this phenomenon. A more general definition appears in [12], we slightly extend it here to the case of non-compact manifolds. All these definitions become equivalent given certain regularity conditions on the family  $\{X_t\}$  and assuming  $M$  is compact.

**Definition A.6.** Let  $\{X_t\}$ ,  $t \in [0, 1]$  be a continuous family of non-zero, complete smooth vector fields on a manifold  $M$  and  $\beta \in \pi_1^{inc}(X)$ .

We say that  $\{X_t\}$  has a **catastrophe in class  $\beta$** , if there is an element

$$y \in \mathcal{O}(X_0, \beta) \sqcup \mathcal{O}(X_1, \beta) \subset \mathcal{O}(\{X_t\}, \beta)$$

such that there is no open compact subset of  $\mathcal{O}(\{X_t\}, \beta)$  containing  $y$ .

A vector field  $X$  on  $M$  is **geodesible** if there exists a metric  $g$  on  $M$  s.t. every flow line of  $X$  is a unit speed  $g$ -geodesic. A family  $\{X_t\}$  is **geodesible** if there is a continuous family  $\{g_t\}$  of metrics, with  $X_t$  geodesible with respect to  $g_t$  for each  $t$ . A family  $\{X_t\}$  is **geodesible** if there is a continuous family  $\{g_t\}$  of metrics with  $X_t$  geodesible with respect to  $g_t$  for each  $t$ . A **geodesible sky catastrophe** is a geodesible family  $\{X_t\}$  with a sky catastrophe. A **Reeb sky catastrophe** is a family of Reeb vector fields  $\{X_t\}$  with a sky catastrophe.

## 7. ACKNOWLEDGEMENTS

Larry Guth for related discussions. Dennis Sullivan and Misha Gromov for much welcome supportive comments.

## REFERENCES

- [1] M. T. ANDERSON, *Metrics of negative curvature on vector bundles*, Proc. Am. Math. Soc., 99 (1987), pp. 357–363.
- [2] D. BAO, S.-S. CHERN, AND Z. SHEN, *An introduction to Riemann-Finsler geometry*, vol. 200 of Grad. Texts Math., New York, NY: Springer, 2000.
- [3] D. BURAGO, Y. BURAGO, AND S. IVANOV, *A course in metric geometry*, vol. 33 of Grad. Stud. Math., Providence, RI: American Mathematical Society (AMS), 2001.
- [4] F. FULLER, *Note on trajectories in a solid torus.*, Ann. Math. (2), 56 (1952), pp. 438–439.
- [5] ———, *An index of fixed point type for periodic orbits.*, Am. J. Math., 89 (1967), pp. 133–145.
- [6] D. GROMOLL AND W. MEYER, *Periodic geodesics on compact Riemannian manifolds*, J. Differ. Geom., 3 (1969), pp. 493–510.
- [7] W. KLINGENBERG, *Closed geodesics*, Ann. Math. (2), 89 (1969), pp. 68–91.
- [8] J. MILNOR, *Morse theory*, Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, No. 51, Princeton University Press, Princeton, N.J., 1963.
- [9] R. S. PALAIS AND S. SMALE, *A generalized Morse theory*, Bull. Am. Math. Soc., 70 (1964), pp. 165–172.
- [10] A. PREISSMANN, *Quelques propriétés globales des espaces de Riemann*, Commentarii Mathematici Helvetici, (1942).
- [11] S. SASAKI, *On the differential geometry of tangent bundles of Riemannian manifolds*, Tôhoku Math. J. (2), 10 (1958), pp. 338–354.
- [12] Y. SAVELYEV, *Extended Fuller index, sky catastrophes and the Seifert conjecture*, International Journal of mathematics, 29 (2018).
- [13] ———, *Elliptic curves in lcs manifolds and metric invariants*, <http://yashamon.github.io/web2/papers/conformalsymplectic.pdf>, (2023).
- [14] W. P. THURSTON, *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Am. Math. Soc., New Ser., 19 (1988), pp. 417–431.

Email address: `yasha.savelyev@gmail.com`

FACULTY OF SCIENCE, UNIVERSITY OF COLIMA, MEXICO