

RELATING MEAN CURVATURE AND DIAMETER IN GENERAL RIEMANNIAN MANIFOLDS, AND ENERGY QUANTIZATION

YASHA SAVELYEV

ABSTRACT. Using a theorem of Topping we give a simple relation between mean curvature and intrinsic diameter for closed submanifolds of general compact Riemannian manifolds. We use to prove a quantization of energy result of holomorphic curves in a locally conformally symplectic manifold.

In [6] Topping gave via a concise but sophisticated argument based on ideas of Ricci flows a simple relation between intrinsic diameter and mean curvature for immersed submanifold of \mathbb{R}^n . Let us state it here:

Theorem 0.1 ([6]). *For Σ^m a smoothly immersed closed submanifold of \mathbb{R}^n we have:*

$$\text{diam}(\Sigma) \leq \text{Const}(m) \int_{\Sigma} |\mathbf{H}|^{m-1} d\text{vol},$$

for \mathbf{H} the mean curvature vector field along Σ , vol the volume measure induced by the standard ambient metric, and diam the intrinsic diameter: $\max_{x,y \in \Sigma} \text{dist}_{(\Sigma, g_{st})}(x, y)$.

Here we use the above and Nash embedding theorem to give via an otherwise elementary argument the following (mostly) simple relation between diameter and mean curvature of closed submanifolds of general compact Riemannian manifolds. We use this to give an application to energy quantization of holomorphic curves in locally conformally symplectic manifolds. In plain words the main statement is the following: if the volume of a given closed immersed submanifold of a compact Riemannian manifold is “small” but diameter “large” then the mean curvature must be somewhere large. Here is the more precise statement.

Theorem 0.2. *Consider the set $S = S(C)$ of immersed $\Sigma \subset X$, for Σ a closed smooth m -manifold and (X, g) a fixed compact Riemannian manifold, with the mean curvature of Σ bounded from above by $C > 0$. Let $\text{Vol}(\Sigma)$ denote the g -volume, and $\text{diam}(\Sigma)$ the (intrinsic) diameter in X, g . Then for all $\Sigma \in S$*

$$\text{diam}(\Sigma) \leq F(g, C, m) \text{vol}(\Sigma),$$

for some function F .

Proof. Pick an isometric Nash embedding N of (M, g) into \mathbb{R}^n , where n is large enough.

Lemma 0.3. *For all $\Sigma \in S(C)$ the magnitude of the mean curvature vector field along $N(\Sigma)$ in \mathbb{R}^n is bounded from above by some C' .*

Proof. In what follows we conflate the notation for Σ and its images $u : \Sigma \rightarrow M$, $N \circ f : \Sigma \rightarrow \mathbb{R}^n$. In other words we just think in terms of subspaces $\Sigma \subset M \subset \mathbb{R}^n$. Let h be the second fundamental form on $T_p M$:

$$h(v, w) = \tilde{\nabla}_v w - \nabla_v w,$$

where $\tilde{\nabla}$ is the Levi-Civita connection of (\mathbb{R}^n, g_{st}) , ∇ is the Levi-Civita connection of (Σ, g) and where we locally extend $v, w \in T_p \Sigma$ to vector fields tangent to Σ . If $\dim \Sigma = m$, the mean curvature vector

1991 *Mathematics Subject Classification.* 53C42.

Partially supported by PRODEP grant of SEP, Mexico.

of Σ in \mathbb{R}^n at p is given by:

$$\mathbf{H}(p) = \frac{1}{m} \sum_i h(e_i, e_i),$$

where $\{e_i\}$ is an orthonormal basis for $T_p\Sigma$. Likewise $\tilde{\nabla}$ will denote in what follows the Levi-Civita connection of (M, g) . So we have:

$$\begin{aligned} m|\mathbf{H}(p)| &= \left| \sum_i (\tilde{\nabla}_{e_i} e_i - \nabla_{e_i} e_i) \right| = \left| \sum_i (\tilde{\nabla}_{e_i} e_i - \tilde{\nabla}_{e_i} e_i + \tilde{\nabla}_{e_i} e_i - \nabla_{e_i} e_i) \right| \\ &\leq \left| \sum_i (\tilde{\nabla}_{e_i} e_i - \nabla_{e_i} e_i) \right| + mB \\ &\leq mC + mB, \end{aligned}$$

where $B = \sup_{e \in TM, |e|=1} |\tilde{\nabla}_e e - \nabla_e e|$. □

Since Σ is closed we get by Topping's theorem:

$$\text{diam}(N\Sigma) \leq \text{Const}(m) \int_{\Sigma} |\mathbf{H}_{\mathbb{R}^n}|^{m-1} d\text{Vol}.$$

By the lemma above the function $|\mathbf{H}_{\mathbb{R}^n}|$ on Σ is universally (independently of u) bounded from above by some C' . So we get:

$$\text{diam}(\Sigma) = \text{diam}(N\Sigma) \leq \text{Const}(m) \cdot (C')^{m-1} \cdot \text{vol}(\Sigma),$$

and so we get the required inequality. □

1. APPLICATION TO ENERGY QUANTIZATION

A locally conformally symplectic manifold or l.c.s. manifold, is a smooth $2n$ -fold M , with a non-degenerate 2-form ω , which is locally diffeomorphic to $e^f \omega_0$ for some functions f , and ω_0 the standard symplectic form on \mathbb{R}^{2n} . There has been some interest recently in developing Gromov-Witten theory for l.c.s. manifolds, [5], [1]. This is partly impeded by our lack of understanding of how badly can holomorphic curves behave in such a manifold. For instance, in the context of holomorphic curves in symplectic manifolds, we have the following energy quantization phenomenon:

Theorem 1.1. *Given a closed symplectic (M, ω) , J an ω -compatible almost complex structure there exists a constant $\hbar = \hbar(\omega, J)$, s.t. for any $u : (\Sigma, j) \rightarrow M$ a non-constant J -holomorphic map of a closed Riemann surface (Σ, j) , the energy of the map satisfies:*

$$e(u) = \int_{\Sigma} |du|^2 d\text{vol} = \int_{\Sigma} u^* \omega \geq \hbar,$$

for vol the measure induced by the ambient (ω, J) metric.

For $\Sigma = S^2$ this holds via a generalized mean value inequality, [4], [7] and the symplectic condition on M can be loosened to just almost complex. For more general Σ , but with M symplectic we can give a simple argument via geometric measure theory. We give this proof here for completeness, as I am not aware if this previously appeared and the proof is relatively elementary, given state of the art.

Proof. Suppose otherwise, then we have a sequence $\{u_i\}$ of J -holomorphic curves with $e(u_i) \rightarrow 0$ as $i \rightarrow \infty$. In particular, $|u_i| \rightarrow 0$, where $|\cdot|$ is the mass norm, and u_i are understood as integral 2-currents, Federer [2]. By the main compactness theorem for currents, $\{u_i\}$ has a convergent subsequence $\{u_{i_k}\}$ to an integral 2-current with mass necessarily 0, that is the 0 in the vector space of closed integral 2-currents $\mathcal{I}_2(M)$, since mass is a norm. Next it is proved in [2] that the space $\mathcal{H}_k(M)$ of closed integral k -currents modulo exact integral k -currents, is discrete with respect to induced mass norm, and isomorphic to singular integral k -homology. Moreover the natural map

$$q_k : \mathcal{I}_k(M) \rightarrow \mathcal{H}_k(M)$$

is continuous. Thus $\{q_2(u_{i_k})\}$ is eventually constant, which means that u_{i_k} are eventually in class 0, which is a contradiction, since all u_i have positive symplectic area. \square

We point out here that a partial extension of the above can be made to l.c.s. manifolds.

Theorem 1.2. *Given a closed l.c.s. (M, ω) , J an ω -compatible almost complex structure there exists a constant $\hbar = \hbar(\omega, J)$, s.t. for any $u : (\Sigma, j) \rightarrow M$ a J -holomorphic immersion of a closed Riemann surface (Σ, j) , the energy of the map satisfies:*

$$e(u) = \int_{\Sigma} |du|^2 d\text{vol} = \int_{\Sigma} u^* \omega \geq \hbar,$$

for vol the measure induced by the ambient (ω, J) metric.

Proof.

Theorem 1.3. *For $u : \Sigma \rightarrow M$ a J -holomorphic immersion, as above, the mean curvature $|\mathbf{H}|$ of the image of u is bounded from above by a universal constant $C(\omega, J) > 0$.*

Proof. We may fix a finite cover of M by charts $\phi_i : U_i \subset \mathbb{R}^{2n} \rightarrow M$, $\phi_i^* \omega = e^{f_i} \omega_0$ with ω_0 the standard symplectic form, and with U_i contractible. Then $\phi_i^{-1} \circ u$ is a $\phi_i^* J$ -holomorphic map defined on $u^{-1}(\phi_i(U_i))$ into \mathbb{R}^{2n} , and $\phi_i^* J$ is compatible with $f_i \omega_0$ and hence with ω_0 . Fix a cover of Σ by disk domains $\{V_{i_j}\}$, with each $V_{i_j} \subset u^{-1}(\phi_i(U_i))$ for some i . Then $\phi_i^{-1} \circ u|_{V_{i_j}}$ is an immersed $\phi_i^* J$ -holomorphic curve in \mathbb{R}^{2n} . So $(\omega_0, \phi_i^* J)$ is an almost Kahler manifold, and the image D_{i_j} of $\phi_i^{-1} \circ u|_{V_{i_j}}$ is hence a minimal surface, with respect to the metric $(\omega_0, \phi_i^* J)$, since ω is then a calibration, Harvey-Lawson [3]. Thus D_{i_j} has mean curvature 0. Since the cover $\{U_i\}$ is finite, the C^∞ norm of the functions f_i is universally bounded: $A < |f_i| < B$, for some A, B and all i . It follows that the magnitude of the mean curvature of the surface D_{i_j} with respect to the metric induced by $(e^{f_i} \omega_0, \phi_i^* J)$ is bounded from above by some universal constant $C(\omega, J)$. As the distortion of the mean curvature corresponding to the conformal distortion e^{f_i} can be readily bounded in terms of the functions f_i . Consequently the mean curvature of image $u|_{V_{i_j}}$ with respect to (ω, J) , is likewise bounded by $C(\omega, J)$, from which the result follows. \square

Now let ϵ be the Lebesgue covering number of $\{U_i\}$ with respect to the metric g . Combining Lemma 1.3 and Theorem 0.2 we get that for u non-constant as in the hypothesis if $\text{area}(u) < \hbar$ then $\text{diam}(u) < \epsilon$, for some \hbar independent of u . Consequently the image of u is contained in some U_i , and so $\phi_i^{-1} \circ u$ is a $\phi_i^* J$ -holomorphic map of a sphere into the almost Kahler contractible manifold $(U_i, \omega_0, \phi_i^* J)$ and so must be constant, which is a contradiction. \square

Question 1. Can the condition of u being an immersion in the statement of Theorem 1.2 be removed?

Acknowledgements. I am grateful to Peter Topping for comments on earlier ideas, Robert Kushner for noticing some typos.

REFERENCES

- [1] C. BAPTISTE AND A. MURPHY, *Conformal symplectic geometry of cotangent bundles*, arXiv, (2016).
- [2] H. FEDERER, *Geometric measure theory. Repr. of the 1969 ed.*, Berlin: Springer-Verlag, repr. of the 1969 ed. ed., 1996.
- [3] R. HARVEY AND H. B. LAWSON, *Calibrated geometries.*, Acta Math., 148 (1982), pp. 47–157.
- [4] D. MCDUFF AND D. SALAMON, *J-holomorphic curves and symplectic topology*, no. 52 in American Math. Society Colloquium Publ., Amer. Math. Soc., 2004.
- [5] Y. SAVELYEV, *Gromov Witten theory of a locally conformally symplectic manifold and the Fuller index*, arXiv, (2016).
- [6] P. TOPPING, *Relating diameter and mean curvature for submanifolds of Euclidean space.*, Comment. Math. Helv., 83 (2008), pp. 539–546.
- [7] K. WEHRHEIM, *Energy quantization and mean value inequalities for nonlinear boundary value problems.*, J. Eur. Math. Soc. (JEMS), 7 (2005), pp. 305–318.

E-mail address: yasha.savelyev@gmail.com

UNIVERSITY OF COLIMA, CUICBAS