

[Direct link to author's version](#)

GLOBAL FUKAYA CATEGORY I

YASHA SAVELYEV

ABSTRACT. Let $Ham(M, \omega)$ denote the Frechet Lie group of Hamiltonian symplectomorphisms of a monotone symplectic manifold (M, ω) . Let $NFuk(M, \omega)$ be the A_∞ -nerve of the Fukaya category $Fuk(M, \omega)$, and let $|(\mathbb{S}, NFuk(M, \omega))|$ denote the $NFuk(M, \omega)$ component of the “space” of ∞ -categories. Using Floer-Fukaya theory for a monotone (M, ω) we construct a natural up to homotopy classifying map

$$BHam(M, \omega) \rightarrow |(\mathbb{S}, NFuk(M, \omega))|.$$

This verifies one sense of a conjecture of Teleman on existence of action of $Ham(M, \omega)$ on the Fukaya category of (M, ω) . This construction is very closely related to the theory of the Seidel homomorphism and the quantum Chern classes of the author, and this map is intended to be the deepest expression of their underlying geometric theory. In part II the above map is shown to be non trivial by an explicit calculation. In particular we arrive at a new non-trivial “quantum” invariant of any smooth manifold, which motivates the statement of a kind of “quantum” Novikov conjecture.

CONTENTS

1. Introduction	2
1.1. A functor from the category of smooth simplices of X to A_∞ categories	2
1.2. Towards new invariants and quantum Novikov conjecture	5
1.3. Hochschild and geometric Hochschild cohomology and homotopy groups of $Ham(M, \omega)$	5
1.4. Organization	7
1.5. Acknowledgements	7
2. Notations and conventions and large categories	7
3. Preliminaries	8
3.1. The simplex category of a smooth manifold X	8
3.2. Preliminaries on Riemann surfaces	9
3.3. Hamiltonian fibrations	12
4. A system of natural maps from the universal curve to Δ^n	12
4.1. Outline of an explicit construction	21
5. Auxiliary data \mathcal{D}	24
5.1. Lagrangian labels and admissible connections	27
5.2. Gluing admissible connections.	28
5.3. Admissible fiber almost complex structures	29
5.4. Gluing admissible fiber almost complex structures	29
5.5. Combining admissible connections and fiber almost complex structures	30
5.6. The summary of the perturbation data \mathcal{D}	32
6. The functor F	32
6.1. F on a point	32
6.2. F on higher dimensional simplices	35

6.3. Unital replacement of F	38
6.4. Concordance classes of functors $F : \text{Simp}(X) \rightarrow A_\infty - \text{Cat}$	39
7. Global Fukaya category	39
7.1. Global Fukaya category as a colimit	40
7.2. Universal construction via diffeological spaces	41
7.3. Universal construction via smooth simplicial sets	42
7.4. Global Fukaya category and unital replacement, a remark	43
8. Extending F to degeneracies	43
9. Algebraic-topological considerations	45
9.1. Colimit of F	45
9.2. Naturality	48
Appendix A. ∞ -categories and Joyal model structure	49
A.1. Categorical equivalences, morphisms and equivalences	49
A.2. Inner fibrations	51
A.3. Categorical fibrations	51
A.4. A_∞ -nerve	52
References	53

1. INTRODUCTION

Smooth fibrations over a Lorentz 4-manifold with fiber a Calabi-Yau 6-fold are a model for the physical background in string theory. This suggests that there may be some string theory linked mathematical invariants of such a fibration. Indeed, when the structure group of $M \hookrightarrow P \rightarrow X$ can be reduced to the group of Hamiltonian symplectomorphisms of M , (with its C^∞ topology) in which case P is called a Hamiltonian fibration, there are a couple of basic invariants of such a fibration based on Floer-Gromov-Witten theory. One such example is the Seidel representation [23] and the related quantum characteristic classes of the author [22]. Related invariants are also proposed by Hutchings [8]. Even earlier there is work on parametric Gromov-Witten invariants of Hamiltonian fibrations by Le-Ono [9] and Olga Buse [2]. At the same time, Costello's theorem [3] on reconstruction of topological conformal field theories from Calabi-Yau A_∞ categories suggests that the above invariants must have a similar reconstruction principle.

For a given Hamiltonian fibration P as above, the A_∞ Fukaya categories of the fibers fit into a “family”, although exactly what this “family” should mean is a non-trivial problem by itself, since we must somehow remember the continuity of P . Then our basic idea is that associated to a Hamiltonian fibration there should be a classifying map from X into an appropriate “classifying” space of A_∞ categories, from which the other invariants can be reconstructed via a version of Toen's derived Morita theory, [26]. This can also be understood to say that $\text{Ham}(M, \omega)$ naturally (continuously) acts on $\text{Fuk}(M)$, verifying in one sense a conjecture of Teleman. We say more on this in Section 1.3.

This paper will be mostly self contained, as we will explain many (especially algebraic) concepts used.

1.1. A functor from the category of smooth simplices of X to A_∞ categories. The first basic ingredient for our construction is as follows. Given P as

above, and a choice of geometric-analysis theoretic perturbation data \mathcal{D} , to each smooth simplex

$$\Sigma : \Delta^n \rightarrow X,$$

we associate an A_∞ category $F(\Sigma)$. This data involves certain compatible choices of Hamiltonian connections and almost complex structures, similar to the kind in the construction of the Seidel morphism [23]. This will be done in Section 5.

A key geometric ingredient is the following. Let $\overline{\mathcal{R}}_d$, $d \geq 2$, denote the compactified moduli space of Riemann surfaces which are topologically disks with $d+1$ punctures on the boundary. We construct natural, axiomatically determined maps from the universal curves over $\overline{\mathcal{R}}_d$, for each d , into the standard topological simplices Δ^n . This topological-combinatorial connection of the universal curves with simplices is new, and is likely of independent interest.

Let X_\bullet denote the smooth singular set of X and let P be as above. Let

$$\Delta(X) = \Delta/X_\bullet$$

denote the category of simplices of X_\bullet , see Section 3.1.

The above data is then extended to a functor

$$F_P : \Delta(X) \rightarrow A_\infty - Cat^{unit}.$$

with $A_\infty - Cat^{unit}$ the category of small, unital, \mathbb{Z}_2 -graded A_∞ categories over \mathbb{Q} , with morphisms strict embeddings, which are moreover quasi-equivalences.

We had mentioned above the “space of A_∞ -categories”. However, technically it will be simpler to work with a related space (Kan complex) of ∞ -categories we denote by \mathbb{S} , discussed in Appendix A.1. A bit more completely, this is a certain Kan complex whose vertices are ∞ -categories, and whose edges are equivalences of ∞ -categories, called categorical equivalences.

The connection of the functor F_P with \mathbb{S} comes via the nerve functor

$$N : A_\infty - Cat^{unit} \rightarrow sSet,$$

with right hand side the category of simplicial sets. The functor N is an analogue for A_∞ categories of the classical nerve construction, which is due to Grothendieck. The A_∞ version, first suggested in Lurie [12], can be considered to be a special case of the more general nerve construction for simplicial categories, and was developed by Faonte [5]. See also, Tanaka [10]. What will be crucial for us is that N takes an A_∞ -category to a ∞ -category.

One basic reason that working with \mathbb{S} is useful, is that it will allow us to convert all the algebraic data of the functor F above, to the data of a single combinatorial-topological object, which we call the global Fukaya category $Fuk_\infty(P)$, as appearing in the title of the paper. More specifically, $Fuk_\infty(P)$ has the structure of ∞ -fibration $Fuk_\infty(P) \rightarrow X$, described in Section 7.1.3. (The latter is a special kind of simplicial fibration.) This will be crucial for computations in Part II [20].

We denote the connected component of an ∞ -category \mathcal{X} in \mathbb{S} by $(\mathbb{S}, \mathcal{X})$, cf. Definition A.2. The geometric realization $|\mathbb{S}|$ may thus be understood as the “space of infinity categories”, cf. Appendix A. In what follows \mathcal{X} will be $NFuk(M, \omega)$.

Together with suitable invariance, under deformation of the perturbation data \mathcal{D} , this yields the following.

Theorem 1.1. *For (M, ω) a monotone symplectic manifold, and $M \hookrightarrow P \rightarrow X$ a Hamiltonian fibration over a smooth manifold X , there is a natural up homotopy*

map

$$cl_P : X \rightarrow |(\mathbb{S}, NFuk(M, \omega))|.$$

Moreover this extends to the universal level, so that there is a natural up to homotopy map

$$cl : BHam(M, \omega) \rightarrow |(\mathbb{S}, NFuk(M, \omega))|,$$

corresponding to the universal Hamiltonian M -fibration $p : E_M \rightarrow BHam(M, \omega)$. This is natural so that

$$[cl_P] = [cl] \circ [\tilde{cl}_P],$$

for $\tilde{cl}_P : X \rightarrow BHam(M, \omega)$ the classifying map of the Hamiltonian fibration P , and $[\cdot]$ denoting the homotopy class.

Natural up to homotopy just means that the map is natural in the homotopy category, with morphisms homotopy classes of continuous maps. The name cl_P comes from the fact that in a certain sense cl_P is classifying. It classifies the fibration $Fuk_\infty(P) \rightarrow X_\bullet$. The proof of this theorem is in Section 7.2.

This theorem also be interpreted to say that $Ham(M, \omega)$ “continuously acts” on $NFuk(M)$.

Remark 1.2. For example, working in the category of simplicial sets, suppose we have a simplicial action of $Ham(M, \omega)_\bullet$ on $NFuk(M, \omega)$. Then we have an induced simplicial map

$$BHam(M, \omega)_\bullet \rightarrow BAut(NFuk(M, \omega)),$$

with $Aut(NFuk(M, \omega))$ interpreted as a simplicial group, and where BG denotes the simplicial nerve of a simplicial group G . And it’s easy to see that there is a natural map

$$BAut(NFuk(M, \omega)) \rightarrow (\mathbb{S}, NFuk(M, \omega)),$$

by the construction of \mathbb{S} . Starting with a simplicial map

$$(1.1) \quad BHam(M, \omega)_\bullet \rightarrow (\mathbb{S}, NFuk(M, \omega))$$

we do not generally get an induced homomorphism

$$Ham(M, \omega)_\bullet \rightarrow Aut(NFuk(M, \omega)),$$

in a natural way. But if we take the based loop space of both sides of (1.1) we get something approaching this homomorphism, since $\Omega BHam(M, \omega)_\bullet \simeq Ham(M, \omega)_\bullet$ (simplicial homotopy equivalence). And since the simplicial H -space

$$\Omega(\mathbb{S}, NFuk(M, \omega))$$

in a sense extends the group

$$Aut(NFuk(M, \omega)),$$

(loops in \mathbb{S} based at $NFuk(M, \omega)$ are in correspondence with categorical self-equivalences of $NFuk(M, \omega)$, which generalize simplicial automorphisms.)

The continuous action of $Ham(M, \omega)$ on $NFuk(M, \omega)$ is one interpretation of the existence of a “continuous action” of $Ham(M, \omega)$ on $Fuk(M, \omega)$. A discret version of such an action can be found in Seidel [24, Section 10c]. Thus, we verify one sense of a conjecture of Teleman ICM 2014 on the existence of such an action. Another interpretation of this action appears in the work of Oh-Tanaka [17]. There the functor of type F_P above is converted to a map of spaces by means of localizations. This provides an alternative algebraic topological perspective.

In part II [20] of this paper, this map is shown to be homotopically non-trivial in a specific example, and some possibly unexpected geometric applications of this are developed.

1.2. Towards new invariants and quantum Novikov conjecture. By the above discussion we automatically obtain a new invariant of a Hamiltonian fibration $M \hookrightarrow P \rightarrow X$ as the homotopy class of the classifying map $cl_P : X \rightarrow |\mathbb{S}|$.

It may be difficult to get intrinsic motivation for Hamiltonian fibrations for a reader outside of symplectic geometry, as a start one may read [7]. However as one particular case we can fiberwise projectivize the complexified tangent bundle:

$$P(X) = P(TX \otimes \mathbb{C}),$$

of a smooth manifold X . This $P(X)$ in particular has the structure of a smooth Hamiltonian fibration with fiber \mathbb{CP}^{r-1} for r the real dimension of X . In this way we also get a new invariant of a smooth r manifold X , given by the homotopy class of the classifying map

$$cl_{P(X)} : X \rightarrow (\mathbb{S}, NFuk(\mathbb{CP}^{r-1})),$$

induced by Theorem 1.1.

Recall that Pontryagin classes of a smooth manifold are defined as Chern classes of its complexified tangent bundle. Novikov has shown that rational Pontryagin classes are topologically invariant. It is then very natural to ask the following, “quantum” variant of the Novikov conjecture:

Question 1.3. *Suppose that $f : X \rightarrow Y$ is a homeomorphism of smooth manifolds. Is $cl_{P(X)}$ homotopic to $cl_{P(Y)} \circ f$?*

I suspect that the answer is yes, simply because the whole construction involves a kind of integration theory, not fantastically far removed from Chern-Pontryagin theory, (if we understand “Gromov-Witten counts” as integration). This would lead to further questions however, of how the resulting invariants are related to more classical topological invariants of smooth manifolds.

The answer of “no” is perhaps more interesting, since it means that our construction gives new smooth invariants of manifolds via holomorphic curves in symplectic geometry.

1.3. Hochschild and geometric Hochschild cohomology and homotopy groups of $Ham(M, \omega)$. This section is an excursion, meant to relate our geometric theory with the algebraic derived Morita theory of Toen. For an A_∞ category C we define

$$HH_{geom}^{2-i}(C) = \pi_i(\mathbb{S}, NC), \quad i > 2.$$

The left hand side is named geometric Hochschild cohomology, the name and notation will be justified shortly. By Theorem 1.1 above we then get:

Theorem 1.4. *For (M, ω) monotone, there is a natural group homomorphism*

$$(1.2) \quad \pi_{i-1}(Ham(M, \omega), id) \rightarrow HH_{geom}^{2-i}(Fuk(M, \omega)), \quad i > 2.$$

$HH^*(Fuk(M, \omega))$ is known to be isomorphic to $QH^*(M)$ in some cases, for example in the monotone setting, relevant to us here, this is due to Sheridan [25]. And so the above morphism, when $i > 2$ has the same formal form as (a special case of) the author’s quantum characteristic classes [22], provided there is a connection between $HH^*(Fuk(M))$ and $HH_{geom}^*(Fuk(M))$. Such a connection is described

further below. This would be the most basic form of the “reconstruction” that was mentioned in the first paragraph of the paper.

In Part II we calculate with Hamiltonian S^2 fibrations over S^4 to get:

Theorem 1.5. *The map*

$$(\pi_3(\text{Ham}(S^2, \omega), \text{id}) = \mathbb{Z}) \rightarrow (HH_{\text{geom}}^{-2}(\text{Fuk}(S^2)) = \pi_4(\mathbb{S}, N\text{Fuk}(S^2))),$$

determined by (1.2) is an injection.

This has some rather surprising consequences, particularly for the theory of singular connections.

1.3.1. Geometric Hochschild cohomology and Toen’s derived Morita theory. A small disclaimer. $HH_{\text{geom}}^*(C)$ is just a name for an object whose construction is immediate from work of Joyal and Lurie, and quite possibly appears elsewhere. We claim no originality for this construction. What may however be interesting is the connection to symplectic geometry that we discover in these papers, and perhaps $HH_{\text{geom}}^*(C)$ deserves a more careful study on its own.

Let us then very briefly indicate the connection of $HH_{\text{geom}}^*(C)$ with Hochschild cohomology via Toen’s derived Morita theory. Let $dg - \text{Cat}$ denote the category of differential graded categories, a.k.a. dg categories. Some terminology: a **component** of $C \in dg - \text{Cat}$ is the set ¹ of dg categories quasi-equivalent to C . We denote by the component of C by $(dg - \text{Cat}, C)$.

Theorem 1.6 (Corollary 8.4 [26]). *For a small dg category C , (with cohomological grading conventions) there are natural isomorphisms*

$$(1.3) \quad \pi_i|(dg - \text{Cat}, C)| \simeq HH^{2-i}(C), \text{ for } i > 2,$$

$$(1.4) \quad \pi_2|(dg - \text{Cat}, C)| \simeq HH^0(C)^\times,$$

with $HH^0(C)^\times$ denoting the multiplicative group of invertible elements, $(dg - \text{Cat}, C)$ denoting the C component of the subcategory of $dg - \text{Cat}$ with morphisms quasi-equivalences, and with $|\cdot|$ the geometric realization.

On the other hand the nerve functor N naturally induces a homomorphism,

$$N_* : \pi_i|(dg - \text{Cat}, C)| \rightarrow \pi_i|(\infty - \text{Cat}, NC)| \simeq \pi_i(\mathbb{S}, NC).$$

When C is \mathbb{Z} -graded, (pre)-triangulated dg category over \mathbb{Q} there are folklore theorems of Lurie (personal communication) to the effect that this is an isomorphism.

Thus, in this case for $i > 2$

$$HH^{2-i}(C) = \pi_i(\mathbb{S}, NC) = HH_{\text{geom}}^{2-i}(C),$$

by our definition. This should extend to \mathbb{Z} -graded rational (pre)-triangulated A_∞ categories, along the lines of Faonte [4].

Remark 1.7. *As I understand, but I am not an Fukaya category expert, these hypothesis apply to at least monotone symplectic manifolds, if we pre-triangulate the Fukaya categories. It is important to note however that we do not pre-triangulate the Fukaya categories in the main construction of the paper, it should be possible to do that, following the same ideas, but this possibly loses information, and it may*

¹It is a set if we are implicitly working with Grothendieck universes, otherwise a class. We ignore the set theoretic foundations here. Toen’s own paper makes use of universes, see also [21] which treats these issues explicitly in similar context.

make the computation in Part II [20] more difficult. Without pre-triangulating the connection of HH_{geom}^* and HH^* appears to be more complicated. It is also worth noting that even if we did identify HH_{geom}^* and HH^* then there is still a hard geometric problem of identifying the actual morphisms - the quantum characteristic classes/Seidel morphism and the morphisms from the data of F . So all in all the reconstruction is still an open problem. In [17] a different approach is taken, starting with the functor F , or a close cousin, the authors use categorical techniques of localization, which allows to more directly obtain a connection between $\pi_* B\text{Ham}(M, \omega)$ and HH^* , avoiding the space \mathbb{S} . This approach might however make computations more difficult.

Remark 1.8. In the case of $i = 2$, by Theorem 1.1, we have a homomorphism

$$\pi_1(\text{Ham}(M, \omega), id) \rightarrow \pi_2(\mathbb{S}, NFuk(M, \omega)).$$

So again, if we could again identify $\pi_2(\mathbb{S}, NFuk(M, \omega))$ with $(HH^0(Fuk(M, \omega)))^\times$ and the latter with $QH_{2n}(M)$, then we would, a priori only in form, recover the Seidel homomorphism [23].

1.4. Organization. Section 3 is concerned with preliminaries. The crucial construction of the system of maps from the universal curve to Δ^n is in Section 4. Perturbation data \mathcal{D} is constructed in Section 5. The main functor F is constructed in Section 6. Finally, the global Fukaya category is constructed in Sections 7, 8. With Section 7 containing the proofs of the main Theorems 1.1, 1.4.

1.5. Acknowledgements. I would like to thank Octav Cornea and Egor Shelukhin for discussions and support, Kevin Costello and Paul Seidel for interest. Hiro Lee Tanaka for enthusiasm, generously providing me with an early draft of his thesis and finding a number of misprints in a draft of the paper. Bertrand Toen for explaining to me an outline of the proof of some conjectures and for enthusiastic response. I also thank Jacob Lurie, for feedback on some questions. Special thanks to the referees for tremendous help in the shaping of this paper. The paper was primarily written while I was a CRM-ISM postdoctoral fellow, and I am grateful for the wonderful research atmosphere provided by CRM-Montreal. It was then substantially revised during my stay at RIMS at Kyoto university, and I also thank the staff and Kaoru Ono for great hospitality and discussions.

2. NOTATIONS AND CONVENTIONS AND LARGE CATEGORIES

We use diagrammatic order for composition of morphisms in the Fukaya category, and in ∞ -categories so $f \circ g$ means

$$\cdot \xrightarrow{f} \cdot \xrightarrow{g} \cdot,$$

as reversing order for composition in ∞ -categories is geometrically very confusing, since morphisms are identified with edges of simplices. Elsewhere we use the more common Leibnitz convention. Although this is somewhat contradictory in practice things should be clear from context. By simplex and notation Δ^n we will interchangeably mean the topological n -simplex and the standard representable n -simplex as a simplicial set, for the latter we may also write Δ_\bullet^n .

Given a category C the over-category of an object $c \in C$ is denoted by C/c . We say that a morphism in C is *over* c exactly if it is a morphism in the over-category of c .

Given an A_∞ category by the nerve we always mean the A_∞ nerve N , as previously described.

Some of our ∞ -categories are “large” with proper classes of simplices instead of sets. The standard formal treatment of this is to work with Grothendieck universes. This is a widely accepted extension of set theory. We shall not however need to make this explicit. For reference one paper that does make this kind of thing explicit is [26] also previously cited.

3. PRELIMINARIES

3.1. The simplex category of a smooth manifold X . Let Δ denote the category of combinatorial simplices, whose objects are totally ordered finite sets $[n] = \{0, \dots, n\}$, with $\text{hom}_\Delta([n], [m])$ the set of non-strictly increasing maps

$$\{0, \dots, n\} \rightarrow \{0, \dots, m\}.$$

A simplicial set S_\bullet is a functor $S_\bullet : \Delta^{op} \rightarrow \text{Set}$. We will usually write $S_\bullet(n)$ instead of $S_\bullet([n])$, and this is the set of n -simplices of S_\bullet .

A map of simplicial set $f : A_\bullet \rightarrow S_\bullet$ is a natural transformation of the corresponding functors.

If Δ_\bullet^n denotes the standard n -simplex, $\Delta_\bullet^n = \text{hom}_\Delta(\cdot, [n])$, then we have the category of simplices over S_\bullet , Δ/S_\bullet , whose set of objects is the set of natural transformations $\text{Nat}(\Delta_\bullet^n, S_\bullet)$ and morphisms commutative diagrams

$$\begin{array}{ccc} \Delta_\bullet^n & \longrightarrow & \Delta_\bullet^m \\ & \searrow & \downarrow \\ & & S_\bullet, \end{array}$$

with natural transformation $\Delta_\bullet^n \rightarrow \Delta_\bullet^m$ induced by maps $[n] \rightarrow [m]$. To simplify notation we rename:

$$\Delta(S_\bullet) := \Delta/S_\bullet.$$

Let Δ^n denote the standard topological n -simplex, i.e.

$$\Delta^n := \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_1 + \dots + x_n \leq 1, \text{ and } \forall i : x_i \geq 0\}.$$

The vertices of Δ^n are assumed ordered in the standard way $0, \dots, n$. Let X be a smooth manifold. We say that $\Sigma : \Delta^n \rightarrow X$ is a **smooth map** if it has an extension $V \subset \mathbb{R}^d \rightarrow X$, for $V \supset \Delta^n$ some open set.

Definition 3.1. We say that a smooth map $\Sigma : \Delta^n \rightarrow X$ is **collared** if there is a neighborhood $U \supset \partial\Delta^n$ in Δ^n , such that $\Sigma|_U = \Sigma \circ \text{ret}$ for $\text{ret} : U \rightarrow \partial\Delta^n$ some smooth retraction. Here smooth means that ret has an extension to a smooth map $V \subset \mathbb{R}^d \rightarrow \mathbb{R}^d$, with $V \supset \partial\Delta^n$ open in \mathbb{R}^d .

For X a smooth manifold, define a simplicial set X_\bullet by:

$$X_\bullet(n) := C_{col}^\infty(\Delta^n, X),$$

with the right hand side the set of all smooth collared maps $\Delta^n \rightarrow X$. It is easy to see that X_\bullet is a Kan complex. The same surely holds without the collared condition but the proof is more difficult. For simplicity, we will work with collared simplices and this will no longer be mentioned.

In this case the simplex category

$$\Delta(X) := \Delta(X_\bullet)$$

can be elaborated as follows. It is the category with objects smooth, collared maps $\Delta^n \rightarrow X$, and morphisms commutative diagrams

$$\begin{array}{ccc} \Delta^n & \longrightarrow & \Delta^m \\ & \searrow & \downarrow \\ & & X, \end{array}$$

and top horizontal arrow a **simplicial map**, that is an affine map taking vertices to vertices preserving the order. We say that $\Sigma : \Delta^n \rightarrow X$ is *non-degenerate* if it does not fit into a commutative diagram

$$\begin{array}{ccc} \Delta^n & \longrightarrow & \Delta^m \\ & \searrow \Sigma & \downarrow \\ & & X, \end{array}$$

with $m < n$.

We will denote by $\text{Simp}(X)$ the full subcategory of $\Delta(X)$, with objects non-degenerate singular simplices. The significance of $\text{Simp}(X)$ is that the perturbation data in the construction of F (as in Section 1.1) of the introduction, must first be constructed in the context of $\text{Simp}(X)$, and then formally extended to all simplices. This is necessary to insure functoriality of F on $\Delta(X)$.

3.2. Preliminaries on Riemann surfaces. Much of this material is adopted from the book of Seidel [24]. Although there are some notation changes, to fit better with our goals. Some other notions like the linear ordering, appearing further on, might be new, at least in present type of context.

Let S' be a nodal, connected, simply connected, Riemann surface, with each smooth component topologically a disk D^2 with some marked points on the boundary, indexed by a set I . Removing the marked points we obtain a surface S with ends. However, it is sometimes simpler to represent S as the original compact surface S' with marked points.

The ends are labeled by $\{e_i\}_{i \in I}$. These will also be called **punctures**, especially once we specify additional structure, which will be explained shortly. The nodal points are denoted by $\{n_j\}_{j \in J}$, again for some index set J , and these are distinct from the punctures. For each $j \in J$ we have a pair $S_{j,\pm}$ of smooth components, that are topologically disks with punctures

$$\{e_i\}_{i \in I_{j,\pm}} \subset \{e_i\}_{i \in I},$$

we explain the signs \pm shortly, for now they just distinguish the pair of components $S_{j,+}$ and $S_{j,-}$. More explicitly, $I_{j,+}$ respectively $I_{j,-}$ are just the subsets of I corresponding to the punctures on the components $S_{j,+}$ respectively $S_{j,-}$. If we remove the node n_j from S then

$$S_{j,\pm}^\circ := S_{j,\pm} - n_j$$

has an additional puncture $n_{j,\pm}$ called the **node end**.

We distinguish one puncture of S as the root, to be denoted as e_0 . Using the clockwise orientation of the boundary of S , and if $\text{Card}(I) = d$ we then have an induced ordering, e_0, \dots, e_d of the punctures.

It is sometimes convenient to depict such Riemann surfaces as stable, rooted semi-infinite trees, embedded in the plane, where stable means that the valency of each vertex is at least 3. We do this by assigning a vertex to each smooth component as above, a half infinite edge to each marked point, and an edge to each nodal point, as depicted in Figure 1.




FIGURE 1.

To make some arguments and notation cleaner we also introduce a linear ordering on the smooth components of S , or vertices, by “order of operation” defined as follows. The component with the root semi-infinite edge e_0 will be called the root vertex denoted by ω . In terms of the associated tree for the surface we have a pre-order on vertices given by the distance to the root vertex, (by giving each edge length 1). To get an actual order, first isometrically embed the tree in the plane, while preserving the clockwise ordering of each half-infinite edge, corresponding to the ordering of the punctures. Then clockwise order vertices equidistant to the root, as in Figure 1. We shall denote by α the furthestmost component from ω , i.e. it is the greatest element with respect to our order. Then β is the next furthestmost component, etc. (Pretending that we can’t run out of letters.) Note, that α may not be the leftmost component, in fact “leftmost” may be ambiguous (dependent on embedding) for vertices not equidistant to ω .

Remark 3.2. *This correspondence of letters to the order may seem counterintuitive, but this is motivated by idea that these trees are operadic trees determining composition. More explicitly, later on this is the composition in certain A_∞ Fukaya categories. Composition corresponding to furthestmost elements from ω is performed first. Hence α corresponds to the first operation we need to perform. Although the operations corresponding to components equidistant from ω can be performed in any order.*

As part of the data, at the i ’th puncture we ask for a holomorphic diffeomorphism having the name of the end:

$$e_i : [0, 1] \times [0, \infty) \rightarrow S,$$

$i \neq 0$, and called **strip end chart**. And at the 0’th puncture we ask for a holomorphic diffeomorphism

$$e_0 : [0, 1] \times (-\infty, 0] \rightarrow S.$$

Let $S_{j,\pm}^\circ, S_{j,\pm}$ be as above. We further specify the \pm distinction so that $S_{j,-} > S_{j,+}$ with respect to the linear order above. And we ask for a similar pair of holomorphic diffeomorphisms

$$(3.1) \quad e_{j,-} : [0, 1] \times (-\infty, 0] \rightarrow S_{j,-}^\circ,$$

$$(3.2) \quad e_{j,+} : [0, 1] \times [0, \infty) \rightarrow S_{j,+}^\circ$$

at the $n_{j,\pm}$ ends. The data of such diffeomorphisms, for a given possibly nodal surface, will be called a *strip end structure* and the particular diffeomorphisms *strip end charts*.

The moduli space of the Riemann surfaces as above, with $\text{Card } I = d$, will be denoted by $\overline{\mathcal{R}}_d$. (Note that Seidel [24] calls our $\overline{\mathcal{R}}_d$ by $\overline{\mathcal{R}}_{d+1}$.) $\overline{\mathcal{R}}_d$ is a real dimension $d-2$ manifold with corners. We will also denote by $\mathcal{R}_d \subset \overline{\mathcal{R}}_d$ the subspace corresponding to non-nodal surfaces.

For $d \geq 2$ let $\rho' : \mathcal{S}_d \rightarrow \overline{\mathcal{R}}_d$ denote the universal family of the Riemann surfaces S , as above. Denote by

$$\rho : \mathcal{S}_d^\circ \rightarrow \overline{\mathcal{R}}_d,$$

this universal family where the nodal points of the surface fibers have been removed.

Notation 3.3. We denote by $\mathcal{S}_{d,r}$ and sometimes just by \mathcal{S}_r the fiber $\rho^{-1}(r)$, for $r \in \overline{\mathcal{R}}_d$.

Choose r -smooth (varying smoothly with respect to r) families $\{e_{i,r}\}, \{e_{j,\pm,r}\}$ of strip end structures for the entire universal family $\mathcal{S}_d \rightarrow \overline{\mathcal{R}}_d$, (note that further on r is suppressed). These choices have to be consistent with gluing in the natural sense as explained in [24, Section 9g]. We will keep track of these systems of choices of strip end structures only implicitly.

3.2.1. Metric characterization of the moduli space. It will be helpful to recall the characterization of the moduli space \mathcal{R}_d and its compactification $\overline{\mathcal{R}}_d$, in terms of hyperbolic metrics on the punctured disks \mathcal{S}_r . The family $\{\mathcal{S}_{d,r}\}_{r \in \mathcal{R}_d}$ is in a bijective correspondence with a suitably universal family $\{\text{Met}_r\} = \{\text{Met}_{d,r}\}$ of constant curvature -1 metrics on the disk with $d+1$ punctures on the boundary. Under this correspondence the complex structure on \mathcal{S}_r is just the conformal structure induced by Met_r . This is of course classical, to see all this use Schwartz reflection to “double” each \mathcal{S}_r to a connected genus 0 Riemann surface \mathcal{D}_r , without boundary, and with $d+1$ punctures. This determines an embedding of \mathcal{R}_d into the Grothendieck-Knudsen moduli space $\overline{M}_{0,d+1}$ of genus 0, compact, connected, nodal Riemann surfaces with $d+1$ points removed. As $d > 2$, by the uniformization theorem, each \mathcal{D}_r is a quotient of the disk by a subgroup of $PSL(2, \mathbb{R})$, which must also preserve the hyperbolic metric. Therefore \mathcal{S}_r inherits a hyperbolic metric, that we call Met_r .

The metric point of view gives an illuminating description of the compactification $\overline{\mathcal{R}}_d$, for $d \geq 2$. Starting with some \mathcal{S}_r and taking r to a boundary stratum, corresponds to some fixed collection of embedded, disjoint geodesics on \mathcal{S}_r , with boundary in the boundary of \mathcal{S}_r , have their length shrunk to zero. Each boundary stratum of $\overline{\mathcal{R}}_d$ is completely determined by such a collection of geodesics.

3.2.2. Gluing. The gluing construction (see for example [24]) takes a surface \mathcal{S}_r , $r \in \partial \overline{\mathcal{R}}_d$ and produces a surface with one less node. This gluing is determined by gluing parameters which we parametrize by $[0, 1]$, assigned to each node. For us 0




FIGURE 2. This diagram is only schematic. The embedding into the plane is not meant to be holomorphic or isometric for the natural hyperbolic structure on the surface.

means don't glue, and 1 is meant to correspond to some small value of the gluing parameter used in actual gluing. We will write $d_{\alpha,\beta}$ for the parameter used in the gluing of components α, β , and likewise with other components.

The gluing construction for parameters in $[0, 1)$ determines an open neighborhood called **gluing neighborhood** of the boundary of $\overline{\mathcal{R}}_d$. We will call by N a **gluing normal neighborhood** of the boundary of $\overline{\mathcal{R}}_d$: an open neighborhood of the boundary, deformation retracting to the boundary, contained in the gluing neighborhood.

The gluing construction also induces a kind of thick-thin decomposition of the surface, with thin parts conformally identified with $[0, 1] \times [0, l]$ for $l > 0$ determined by the corresponding gluing parameter. This decomposition is not intrinsic, as it depends in particular on the choice of the family of strip end structures. However, instructively these gluing parameters can be thought of as lengths of geodesic segments, for example m_α, m_β in figure 2, and the thin parts are closely related to thin parts of thick-thin decomposition in hyperbolic geometry.

3.3. Hamiltonian fibrations. A **Hamiltonian fibration** is a smooth fiber bundle

$$M \hookrightarrow P \rightarrow X,$$

with structure group $\mathcal{G} = \text{Ham}(M, \omega)$ with its C^∞ Frechet topology. A **Hamiltonian connection**, in this paper usually denote by \mathcal{A} , is just an Ehresmann \mathcal{G} -invariant connection for a Hamiltonian fibration. When $P = M \times X$ is trivial, such \mathcal{A} may be specified as a one form valued in $C_{\text{norm}}^\infty(M)$ - the space of smooth functions on M with mean 0.

4. A SYSTEM OF NATURAL MAPS FROM THE UNIVERSAL CURVE TO Δ^n

We explain here a remarkable connection between the universal curve over $\overline{\mathcal{R}}_d$ and the standard topological simplices Δ^n . This will be used in our construction but may be of independent interest.

Let $\Pi(\Delta^n)$ be the small groupoid, whose objects set *obj* is the set of vertices of Δ^n . The morphisms set *hom* is the set of affine maps $m : [0, 1] \rightarrow \Delta^n$, (possibly

constant) sending end points to the vertices. The source map

$$s : \text{hom} \rightarrow \text{obj}$$

is defined by $s(m) = m(0)$ and the target map

$$t : \text{hom} \rightarrow \text{obj}$$

is defined by $t(m) = m(1)$. Thus the set $\text{hom}(v_1, v_2)$ of morphisms between $v_1, v_2 \in \Pi(\Delta^n)$ consists of a single morphism. It is the unique affine map $m : [0, 1] \rightarrow \Delta^n$, satisfying $m(0) = v_1$ and $m(1) = v_2$. Consequently the composition maps in $\Pi(\Delta^n)$ are forced, by the above uniqueness. And the identity at v is the constant $m : [0, 1] \rightarrow \Delta^n$, $m(0) = m(1) = v$.

Notation 4.1. We denote the composition of m_1, m_2 by $m_1 \cdot m_2$. The order is diagrammatic, so that this means the composition

$$\cdot \xrightarrow{m_1} \cdot \xrightarrow{m_2} \cdot$$

We say that (m_1, \dots, m_d) is a *composable chain* of morphisms $m_i \in \text{hom}(\Pi(\Delta^n))$ if $t(m_{i-1}) = s(m_i)$, for all $2 \leq i \leq d$. For future use let $m_{i-1,i}$ denote the unique morphism from the $i-1$ vertex to the i vertex in Δ^n .

The goal is to construct a “natural” system of maps

$$u(m_1, \dots, m_d, n) : \mathcal{S}_d^\circ \rightarrow \Delta^n,$$

$d \geq 2$, for each composable chain (m_1, \dots, m_d) , which in particular will satisfy the following four properties.

Definition 4.2. We say that $u(m_1, \dots, m_d, n)$ as above satisfy **partial naturality** if the following holds.

- (1) The maps $u(m_1, \dots, m_d, n)$ are continuous and their restrictions to

$$\rho^{-1}(\mathcal{R}_d \subset \overline{\mathcal{R}}_d)$$

are smooth. This condition is partly why we work with \mathcal{S}_d° instead of \mathcal{S}_d .

- (2) Let $u(m_1, \dots, m_d, n, r)$ denote the restriction of $u(m_1, \dots, m_d, n)$ to \mathcal{S}_r , and let

$$m_0 := m_1 \cdot \dots \cdot m_d$$

be the composition in $\Pi(\Delta^n)$. Then in the strip end coordinates $e_k : [0, 1] \times [0, \infty) \rightarrow \mathcal{S}_r$ at each e_k end, $1 \leq k \leq d$, $u(m_1, \dots, m_d, n, r)$ has the form of the projection to $[0, 1]$ composed with m_k .

- (3) Likewise, in the strip end coordinates $e_k : [0, 1] \times (-\infty, 0] \rightarrow \mathcal{S}_r$ at the end e_0 of \mathcal{S}_r , $u(m_1, \dots, m_d, n, r)$ has the form of the projection to $[0, 1]$ composed with m_0 .
- (4) For $1 \leq k \leq d$, the component of the boundary of \mathcal{S}_r between e_{k-1}, e_k ends is mapped to $s(m_k)$, and the component of the boundary between e_d and e_0 is mapped by $u(m_1, \dots, m_d, n, r)$ to $t(m_d)$, for each $r \in \overline{\mathcal{R}}_d$.

See Figure 3 for an example of a map satisfying naturality. After imposing certain additional to the properties above axioms, Theorem 4.8, which will follow, states that such a suitable system of maps exists and is unique up to suitable homotopy.

So far we haven’t put any conditions at the nodes. These conditions will be forced by certain additional naturality axioms, which arise from various gluing conditions. We start by explaining the natural gluing operations that appear.




FIGURE 3. Figure for a map $u(m_{0,1}, m_{1,2}, 2)$ satisfying partial naturality. The edges of the surface labeled 0, 1, 2 are mapped to the vertices 0, 1, 2 respectively. Colored ends are collapsed onto correspondingly colored edges.

First denote by $\mathcal{T}(m_1, \dots, m_d, n)$ the space of maps satisfying the partial naturality properties above. We have the natural gluing map

$$(4.1) \quad St_i : \overline{\mathcal{R}}_{s_1} \times \overline{\mathcal{R}}_{s_2} \times [0, 1) \rightarrow \overline{\mathcal{R}}_{s_1+s_2-1},$$

whose value on (r, r', τ) is given by gluing the surface \mathcal{S}_r at the root and $\mathcal{S}_{r'}$ at its i 'th marked point, with gluing parameter $\tau \in [0, 1)$, and then associating to this surface its isomorphism class in $\overline{\mathcal{R}}_{s_1+s_2-1}$. (When the value of the gluing parameter is 0, this is the composition map in the Stasheff topological A_∞ operad).

Given an element $u \in \mathcal{T}(m_1, \dots, m_{s_1}, n)$ and an element

$$u' \in \mathcal{T}(m'_1, \dots, m'_{i-1}, m_1 \cdot \dots \cdot m_{s_1} = m'_i, m'_{i+1}, \dots, m'_{s_2}, n),$$

we have a naturally induced map

$$(u \star_i u')_0 : \mathcal{S}_{s_1, s_2, 0}^\circ \rightarrow \Delta^n,$$

where

$$\mathcal{S}_{s_1, s_2, 0}^\circ \rightarrow \overline{\mathcal{R}}_{s_1} \times \overline{\mathcal{R}}_{s_2}$$

is the pullback of the family $\mathcal{S}_{d=s_1+s_2-1}^\circ$ by $St_i|_{\overline{\mathcal{R}}_{s_1} \times \overline{\mathcal{R}}_{s_2} \times \{0\}}$. More specifically, the fiber of $\mathcal{S}_{s_1, s_2, 0}^\circ$ over (r_1, r_2) is the disjoint union of a pair of distinguished (possibly disconnected) subspaces identified with \mathcal{S}_{s_1, r_1} , \mathcal{S}_{s_2, r_2} . So we may apply u to the first component and u' to the second, and this is the map $(u \star_i u')_0$, see Figure 4.

We can extend $(u \star_i u')_0$ to a map

$$(u \star_i u')_1 : \mathcal{S}_{s_1, s_2, 1}^\circ \rightarrow \Delta^n,$$

where

$$\mathcal{S}_{s_1, s_2, 1}^\circ \rightarrow \overline{\mathcal{R}}_{s_1} \times \overline{\mathcal{R}}_{s_2} \times [0, 1)$$

is the pullback of the universal family by $St_i|_{\overline{\mathcal{R}}_{s_1} \times \overline{\mathcal{R}}_{s_2} \times [0, 1)}$. To get this extension we specify $(u \star_i u')_1|_{\mathcal{S}_{r, r', \tau}}$, for $\mathcal{S}_{r, r', \tau}$ the fiber of $\mathcal{S}_{s_1, s_2, 1}^\circ$ over

$$(r, r', \tau) \in \overline{\mathcal{R}}_{s_1} \times \overline{\mathcal{R}}_{s_2} \times [0, 1), \quad \tau \in (0, 1).$$

Recall that the surface $\mathcal{S}_{r, r', \tau}$ glued from $\mathcal{S}_r, \mathcal{S}_{r'}$ has a sub-domain which we denote by $thin = thin_{\tau, i}$ that has a determined conformal identification with a strip of the form $[0, 1] \times [-\phi(\tau), \phi(\tau)]$, for some function (not explicitly needed)

$$(4.2) \quad \phi : [0, 1] \rightarrow (0, \infty),$$




FIGURE 4. The green shaded ends are collapsed onto the same edge of Δ^n .

determined by the particular parametrization of the gluing operation. Now $\mathcal{S}_{r,r'}$ – *thin* is the disjoint union of subspaces holomorphically identified with the regions

$$\text{Reg}_r \subset \mathcal{S}_r, \text{Reg}_{r'} \subset \mathcal{S}_{r'},$$

so that Reg_r is identified with the complement of $[0, 1] \times (-\infty, 0]$ in \mathcal{S}_r , for the strip coordinates on the root end. And likewise, so that $\text{Reg}_{r'}$ is identified with the complement of $[0, 1] \times [0, \infty)$ in $\mathcal{S}_{r'}$, for the strip end coordinates on the i 'th end. We then define $(u \star_i u')_1$ to coincide with u, u' on Reg_r respectively $\text{Reg}_{r'}$, while on *thin* in the coordinates

$$[0, 1] \times [-\phi(\tau), \phi(\tau)],$$

$(u \star u')_1$ is the map given by the projection

$$[0, 1] \times [-\phi(\tau), \phi(\tau)] \rightarrow [0, 1]$$

followed by the map m'_i . This operation is well defined for all $\tau \in (0, 1)$ and so determines the needed extension.

In order state the naturality axioms we need more geometry. Let N be a gluing normal neighborhood of the boundary $\partial\overline{\mathcal{R}}_d$, $d \geq 3$. Recall that \mathcal{R}_d has dimension $d - 2$. Let $S_0^{d-3} \subset N$ be an embedded sphere of dimension $d - 3$, not intersecting $\partial\overline{\mathcal{R}}_d$, homotopic to the inclusion $\partial\overline{\mathcal{R}}_d \simeq S^{d-3} \rightarrow N$, with \simeq homeomorphism. (A 0-dimensional sphere is understood as a pair of points.) Let R_0^{d-2} be the dimension $d - 2$ ball sub-domain of $\overline{\mathcal{R}}_d$ bounded by S_0^{d-3} . Finally, let \mathcal{S}_r – *ends* denote the compact Riemann surface with boundary obtained from \mathcal{S}_r by removing the ends. Specifically, for $1 \leq i \leq d$ we remove the images of the charts $e_i|_{[0,1] \times (0,\infty)} \rightarrow \mathcal{S}_r$, and the image of the chart $e_0 : [0, 1] \times (-\infty, 0) \rightarrow \mathcal{S}_r$.

For $r \notin \partial\overline{\mathcal{R}}_d$ set $D_r^2 := \mathcal{S}_r - \text{ends} \simeq D^2$. And let

$$G_r^2 := (D_r^2, \partial D_r^2) \simeq (D^2, \partial D^2)$$

denote the pair, where \simeq is homeomorphism.

So we get a fiber bundle of pairs

$$(D^2, \partial D^2) \hookrightarrow (\tilde{F}, A) \rightarrow R_0^{d-2},$$

with fiber over r : G_r^2 . And where A is just the corresponding sub-bundle with fiber over r : ∂D_r^2 .

We are almost ready to state our axioms. Let $C_s(\Delta^n)$ denote the set of composable chains (m_1, \dots, m_s) of length $s \geq 2$ in $\Pi(\Delta^n)$.

Definition 4.3. For $(m_1, \dots, m_s) \in C_s(\Delta^n)$ let $D(m_1, \dots, m_s)$ denote the minimal dimension of a subsimplex of Δ^n which contains the edges corresponding to the morphisms m_i .

A system of maps \mathcal{U} is an element of:

$$(4.3) \quad \prod_{n \in \mathbb{N}} \prod_{s \in \mathbb{N}_{\geq 2}} \prod_{(m_1, \dots, m_s) \in C_s(\Delta^n)} \mathcal{T}(m_1, \dots, m_s, n).$$

Given a system \mathcal{U} its projection onto $(n, s, (m_1, \dots, m_s))$ component will be denoted by $u(m_1, \dots, m_s, n)$. To put this another way, let

$$A = \{(m_1, \dots, m_s, n) | s \in \mathbb{N}_{\geq 2}, n \in \mathbb{N}, (m_1, \dots, m_s) \in C_s(\Delta^n)\},$$

then set theoretically the product (4.3) above is the set of certain set maps \mathcal{U} with domain A , (and co-domain a certain function space) then in this language $u(m_1, \dots, m_s, n) := \mathcal{U}(m_1, \dots, m_s, n)$.

Definition 4.4. We say that \mathcal{U} is **natural** if it satisfies the following axioms:

- (1) For all s_1, s_2 and for all i if $m'_i = m_1 \cdot \dots \cdot m_{s_1}$ then the map

$$(4.4) \quad (u(m_1, \dots, m_{s_1}, n) \star_i u(m'_1, \dots, m'_{s_2}, n))_1$$

coincides with the composition

$$(4.5) \quad \mathcal{S}_{s_1, s_2, 1}^\circ \xrightarrow{St_{i,*}} \mathcal{S}_{s_1+s_2-1}^\circ \xrightarrow{u(m'_1, \dots, m'_{i-1}, m_1, \dots, m_{s_1}, m_{i+1}, \dots, m'_{s_2}, n)} \Delta^n,$$

for $St_{i,*}$ the bundle map induced by St_i .

- (2) Let $f : \Delta^n \rightarrow \Delta^m$ be a simplicial map, which recall is an affine map preserving the orders of the vertices. Then there is an induced functor

$$f : \Pi(\Delta^n) \rightarrow \Pi(\Delta^m)$$

and we ask that

$$f \circ u(m_1, \dots, m_s, n) = u(f(m_1), \dots, f(m_s), m).$$

- (3) Let $(m_1, \dots, m_d) \in C_d(\Delta^d)$ for $d \geq 3$. Suppose that $D(m_1, \dots, m_d) = d$ then by the Lemma 4.6 below, $u(m_1, \dots, m_d, d)$ induces a map of pairs:

$$(4.6) \quad \tilde{u} : (\tilde{F}, A \sqcup \tilde{F}|_{S_0^{d-3}}) \rightarrow (\Delta^d, \partial \Delta^d),$$

and we ask that \tilde{u} be a homological degree 1 map.

Remark 4.5. Note that when $d = 2$ we may still state a natural version of Axiom 3, but it would be immediate from the partial naturality properties of Definition 4.2.

Lemma 4.6. Let \mathcal{U} satisfy the first pair of axioms in the Definition 4.4. For $(m_1, \dots, m_d) \in C_s(\Delta^d)$, suppose that $D(m_1, \dots, m_d) = d$. Then $\tilde{u} := u(m_1, \dots, m_d, d)$ is a map of pairs:

$$\tilde{u} : (\tilde{F}, A \sqcup \tilde{F}|_{S_0^{d-3}}) \rightarrow (\Delta^d, \partial \Delta^d).$$

Proof. The map \tilde{u} maps A to $\partial\Delta^d$ by the partial naturality properties 2,3 and 4 in Definition 4.2. In fact A is mapped to the edges of Δ^d .

For $u \in \mathcal{T}(m_1, \dots, m_s, n)$, with (m_1, \dots, m_s) and n general, set $u_r := u|_{S_r}$. Now since S_0^{d-3} is contained in N , by Axiom 1 for each $r \in S_0^{d-3}$, \tilde{u}_r has the form $u_1 \star_i u_2$ for some i , for some $u_1 \in \mathcal{T}(m_1, \dots, m_{s_1}, d)$ and for some

$$u_2 \in \mathcal{T}(m'_1, \dots, m'_{i-1}, m_1 \cdot \dots \cdot m_{s_1} = m'_i, m'_{i+1}, \dots, m'_{s_2}, d),$$

depending on r . In addition $s_1 < d$ and $s_2 < d$ since $s_1 + s_2 - 1 = d$, and since $s_1, s_2 \geq 2$. Now by the previous point and by Axiom 2 it readily follows that the image of u_1 and u_2 is contained in proper faces of Δ^d . Consequently, the image of $\tilde{u}|_{S_0^{d-3}} = u_1 \star_i u_2$ is contained in $\partial\Delta^d$ and so we are done. \square

Remark 4.7. Using an inductive procedure as in the proof of the following theorem, it should be possible to show that the final axiom 3 actually follows by the previous axioms 1,2.

Theorem 4.8. A natural system \mathcal{U} exists, and is unique up to homotopy (through natural systems).

We first give a not explicitly constructive proof in all generality, and afterwards describe a partial explicit construction.

Proof of 4.8. To construct our maps $u(m_1, \dots, m_s, n)$ we will proceed by induction. When $n = 0$ there is nothing to do, as we have unique maps for all $s \geq 2$, and they trivially satisfy Axioms 1,2,3. However, we have to be careful with the basis of the induction, as the induction step needs a sufficiently high n .

Given a composable sequence (m_1, \dots, m_s) of morphisms in $\Pi(\Delta^n)$, for some n , let $D(m_1, \dots, m_s)$ denote the least dimension of a non-degenerate subsimplex of Δ^n which contains the edges corresponding to $\{m_i\}$. Let $S(N)$ be the following statement: there are maps

$$(4.7) \quad u(m_1, \dots, m_s, n),$$

for all $s \geq 2$ and all $n \leq N$ and every composable chain (m_1, \dots, m_s) with $D(m_1, \dots, m_s) = s$ so that axioms 1,3 are satisfied and so that

$$(4.8) \quad \sigma \circ u(m_1, \dots, m_s, n) = u(\sigma(m_1), \dots, \sigma(m_s), n)$$

for

$$\sigma : \Delta^n \rightarrow \Delta^m$$

an injective simplicial map. That is we satisfy Axiom 2 only partially and only for restricted (m_1, \dots, m_s) at the moment. We will also denote by $S(N)$ the corresponding collection of maps (4.7). It can be seen fairly directly that $S(N)$ holds for $N = 0, 1, 2, 3$. $N = 0$ is the trivial case already considered above. Indeed, when $N = 4$ such a construction is given in the following Section 4.1, and this implies the cases of $N = 1, 2, 3$. We intend to prove:

$$(N \geq 3) \wedge S(N) \implies S(N+1),$$

moreover the collection of maps $S(N+1)$ can be chosen to extend the collection of maps $S(N)$. Hence, by Piano induction $\forall N : S(N)$, and we will construct our natural system from the corresponding data.

Note that the condition 4.8 and the maps (4.7) uniquely determine

$$(4.9) \quad u(m_1, \dots, m_s, N+1)$$

for all $\forall s \geq 2, \forall (m_1, \dots, m_s)$ with $D(m_1, \dots, m_s) = s \leq N$. We need an extension in the case $D(m_1, \dots, m_s) = s = N + 1$.

By assumption $N + 1 > 3$. Let then $(m_1^0, \dots, m_{N+1}^0)$, be a chosen composable sequence with $D(m_1^0, \dots, m_{N+1}^0) = N + 1$. Then gluing as in the Axiom 1 of naturality and the maps (4.9) naturally determine a map

$$(4.10) \quad u = u(m_1^0, \dots, m_{N+1}^0, N + 1) : Sub_{N+1} \rightarrow \Delta^{N+1},$$

where $Sub_{N+1} = \rho^{-1}(\partial \overline{\mathcal{R}}_{N+1})$, and where as before

$$\rho : \mathcal{S}_{N+1}^\circ \rightarrow \overline{\mathcal{R}}_{N+1}.$$

Let U be a gluing normal neighborhood of \mathcal{R}_{N+1} . (We previously used the letter N , but just for this proof we use the letter U .) Extend u in any way to $\rho^{-1}(U)$, so that Axiom 1 of naturality is satisfied. We then need to further extend u to \mathcal{S}_{N+1}° so that Axiom 3 of naturality is satisfied.

Let $S_0^{N-2} \subset U$ be an embedded sphere in U not intersecting $\partial \overline{\mathcal{R}}_{N+1}$, homotopic to the inclusion $\partial \overline{\mathcal{R}}_{N+1} \rightarrow U$. Then u induces a map of a pair

$$(4.11) \quad g : (\tilde{F}|_{S_0^{N-2}}, A|_{S_0^{N-2}}) \rightarrow (\partial \Delta^{N+1} \simeq S^N, loop),$$

where $loop$ is a topologically embedded S^1 in $\partial \Delta^{N+1}$ that is the image of the loop

$$\gamma = m_1^0 \cdot \dots \cdot m_{N+1}^0 \cdot m_{s(m_1^0), t(m_{N+1}^0)}^{-1},$$

where \cdot is concatenation of paths and the order of composition is diagrammatic. The map g is just the restriction of the map \tilde{u} , (4.6).

Lemma 4.9. *The map g is homological degree 1.*

Proof. As $N > 2$ $loop$ has codimension greater than 1, so that the meaning of homological degree is unambiguous, as the pair $(S^N, loop)$ has a well defined fundamental class by the homology long exact sequence for a pair. Moreover approximating g by a smooth map we may compute the homological degree via the smooth degree, (denote the approximation still by g). That is let f be a N -face of Δ^{N+1} and $p \in interior(f)$ a regular image point of g . The homological degree of g is then the count of elements of $g^{-1}(p)$ with signs given by whether dg_k , $k \in g^{-1}(p)$, is orientation preserving or reversing.

Suppose without loss of generality that the vertices of f are $0, \dots, N$. As the degree of g is clearly independent of the choice of S_0^{N-2} we may assume that S_0^{N-2} is chosen so that for some $\epsilon > 0$:

$$(4.12) \quad St_1 : (R_0^{N-2} \subset \overline{\mathcal{R}}_N) \times \overline{\mathcal{R}}_2 \times \{\epsilon\} \rightarrow \overline{\mathcal{R}}_{N+1}$$

is an embedding into S_0^{N-2} . We may in addition suppose that $S_0^{N-2} - T$ is covered by such embeddings corresponding to the various other faces of $\overline{\mathcal{R}}_N$, where the region $T \subset S_0^{N-2}$, is such that g maps $\rho^{-1}(T)$ into the union of $(N - 1)$ -faces of Δ^{N+1} . The image of the map (4.12) will be denote by V .

Then by the naturality Axiom 3 the face f is covered by the image of

$$\kappa = (u(m_1^0, \dots, m_N^0, N + 1) \star_1 u(m_1^0 \cdot \dots \cdot m_N^0, m_{N+1}^0, N + 1))|_{\tilde{V}},$$

where

$$\tilde{V} = \rho^{-1}(V).$$

By construction the smooth degree of $g|_{\tilde{V}}$ is the smooth degree of κ . But then, by the naturality Axiom 3, κ has smooth degree one. And again, by naturality and

the assumption on the form of S_0^{N-2} , as described above, no other point of $\tilde{F}|_{S_0^{N-2}}$ is in $g^{-1}(p)$. It follows that g is smooth degree one and so is homological degree one. \square

Lemma 4.10. *There is a degree one extension:*

$$\tilde{g} : (\tilde{F}, A \sqcup \tilde{F}|_{S_0^{N-2}}) \rightarrow (\Delta^{N+1}, \partial\Delta^{N+1}),$$

with $\tilde{g}(A) = \text{loop}$.

Proof. This is all very elementary topology so that we will not give exhaustive detail. Note that $\tilde{F} \simeq B^{N-1} \times D^2$, where B^{N-1} denotes the unit ball in \mathbb{R}^{N-1} . So to find the necessary degree one extension it is enough to show that given a degree one map $h : (S^{d-2} \times D^2, S^{d-2} \times \partial D^2) \rightarrow (S^d, \text{loop})$, there is a degree one extension

$$\tilde{h} : (B^{d-1} \times D^2, \partial B^{d-1} \times D^2) \rightarrow (B^{d+1}, \partial B^{d+1}).$$

First note that a degree one map

$$\tilde{t} : (B^{d-1} \times D^2, \partial B^{d-1} \times D^2) \rightarrow (B^{d+1}, \partial B^{d+1}),$$

exists by an elementary topology construction. (Start with the slicing diffeomorphism $[0, 1]^{d-1} \times [0, 1]^2 \rightarrow [0, 1]^{d+1}$.) We may in addition insure that its restriction to the boundary $S^{d-2} \times D^2$ is a map of a pair

$$t : (S^{d-2} \times D^2, S^{d-2} \times \partial D^2) \rightarrow (S^d, \text{loop}).$$

We may now apply an analogue of a theorem of Hopf, which says that homotopy classes of maps of closed d -manifolds to S^d are classified by degree. (We say analogue because we have a map of pairs.) In our case Hopf's theorem readily implies that t is homotopic to h through maps of pairs. Then use classical homotopy extension theorem, to get a homotopy from \tilde{t} to the needed map \tilde{h} , extending h . \square

Given this \tilde{g} we may readily construct our extension $u : \mathcal{S}_{N+1}^\circ \rightarrow \Delta^{N+1}$ so that

$$u \in \mathcal{T}(m_1^0, \dots, m_{N+1}^0, N+1)$$

and so that the last naturality axiom is satisfied.

Now given any other composable sequence (m_1, \dots, m_{N+1}) with

$$D(m_1, \dots, m_{N+1}) = N+1$$

let

$$\sigma : \Delta^{N+1} \rightarrow \Delta^{N+1}$$

be the unique simplicial bijective map taking (m_1, \dots, m_{N+1}) to $(m_1^0, \dots, m_{N+1}^0)$, (it is unique because the action is determined by the action on the vertices.) And we define

$$u(m_1, \dots, m_{N+1}, N+1) : \mathcal{S}_{N+1}^\circ \rightarrow \Delta^{N+1}$$

by

$$u(m_1, \dots, m_{N+1}, N+1) = \sigma^{-1} \circ u.$$

We then obtain a collection of maps,

$$\{u(m_1, \dots, m_{s'}, n)\}_{(m_1, \dots, m_{s'}, n) \leq N+1, D(m_1, \dots, m_{s'}) = s'},$$

and these satisfy maps satisfy the condition of the statement $S(N+1)$ by the construction and the inductive hypothesis. We thus complete the inductive step.

Moreover, the collection of maps $S(N+1)$ extends the collection $S(N)$. We then obtain the total collection:

$$\mathcal{U}' = \bigcup_N S(N) = \{u(m_1, \dots, m_{s'}, n) | n \in \mathbb{N}, (m_1, \dots, m_{s'}) \in C_{s'}(\Delta^n),$$

$$\text{such that } D(m_1, \dots, m_{s'}) = s'\},$$

It remains to extend the partial system \mathcal{U}' above to a full natural system \mathcal{U} , that is we need to remove restrictions on the D -number. Given (m_1, \dots, m_s) , a composable sequence in $\Pi(\Delta^n)$ with $s > n$, we may write $m_i = pr \circ \tilde{m}_i$ for $(\tilde{m}_1, \dots, \tilde{m}_s)$ a composable sequence in Δ^s s.t. $D(\tilde{m}_1, \dots, \tilde{m}_s) = s$, for $pr : \Delta^s \rightarrow \Delta^n$ surjective simplicial map. Moreover $(\tilde{m}_1, \dots, \tilde{m}_s)$ is clearly unique up to an action of a simplicial bijective map $\sigma : \Delta^s \rightarrow \Delta^s$ fixing the image $i(\Delta^n)$, for $i : \Delta^n \rightarrow \Delta^s$ inclusion of face s.t. $pr \circ i = id$. We then define

$$u(m_1, \dots, m_s, n) := pr \circ u(\tilde{m}_1, \dots, \tilde{m}_s, s).$$

Let $(\tilde{m}'_1, \dots, \tilde{m}'_s)$ be another choice of a composable sequence with $pr(\tilde{m}'_i) = m_i$, and let $\sigma : \Delta^s \rightarrow \Delta^s$ be a simplicial bijective map with $\sigma(\tilde{m}'_i) = \tilde{m}_i$ fixing the image of $i(\Delta^n)$ as above. Then we have

$$pr \circ u(\tilde{m}'_1, \dots, \tilde{m}'_s, s) = pr \circ \sigma \circ u(\tilde{m}_1, \dots, \tilde{m}_s, s).$$

But $pr \circ \sigma = pr$ since σ fixes $i(\Delta^n)$. So we obtain that

$$pr \circ u(\tilde{m}'_1, \dots, \tilde{m}'_s, s) = pr \circ u(\tilde{m}_1, \dots, \tilde{m}_s, s),$$

so that $u(m_1, \dots, m_s, n)$ is well defined. So we have constructed our system of maps satisfying all the axioms of naturality.

To prove uniqueness up to homotopy, note that by our axioms a system of natural maps is completely determined by all the maps $u(m_1, \dots, m_s, n)$ with

$$D(m_1, \dots, m_s) = s = n.$$

We then again proceed by induction. Suppose that we have a pair of natural systems $\mathcal{U}_1, \mathcal{U}_2$. Let $H(N)$ be the statement: there is continuous family of maps $u_t(m_1, \dots, m_s, n)$, $t \in [0, 1]$, for all $n \leq N$, $N \geq 0$, (m_1, \dots, m_s) , with

$$D(m_1, \dots, m_s) = s = n,$$

so that

$$u_{t=i}(m_1, \dots, m_s, n) = u_i(m_1, \dots, m_s, n)$$

for $i = 0, 1$, and so that for each t $u_t(m_1, \dots, m_s, n)$ satisfy the naturality axioms. As for $S(N)$ we also denote by $H(N)$ the corresponding collection of homotopies. The statement $H(N)$, for $N = 0, 1, 2$ holds trivially. We prove:

$$(N \geq 2) \wedge H(N) \implies H(N+1),$$

moreover the collection $H(N+1)$ can be chosen to extend $H(N)$. This then readily implies our claim.

So suppose $(N \geq 2) \wedge H(N)$. Given some composable sequence (m_1, \dots, m_s) in $\Pi(\Delta^{N+1})$, with

$$D(m_1, \dots, m_s) = s = N+1,$$

we have continuous in t families of induced (as before) maps:

$$u_t(m_1, \dots, m_s, N+1) : Sub_s \rightarrow \Delta^{N+1},$$

with Sub_s as before, and so that

$$u_{t=1}(m_1, \dots, m_s, N+1) : Sub_s \rightarrow \Delta^{N+1}$$

coincides with the map

$$u_1(m_1, \dots, m_s, N+1) : Sub_s \rightarrow \Delta^{N+1}.$$

Use the homotopy extension property to get a homotopy:

$$\tilde{u}_t(m_1, \dots, m_{N+1}, N+1) : \mathcal{S}_d^\circ \rightarrow \Delta^{N+1}$$

of the map $u_0(m_1, \dots, m_{N+1}, N+1)$. Now

$$\tilde{u}_1(m_1, \dots, m_{N+1}, N+1)$$

and

$$u_1(m_1, \dots, m_{N+1}, N+1)$$

do not necessarily coincide on \mathcal{S}_d° but they coincide on Sub_s since we used homotopy extension, and both these maps are “degree one” (that is they both satisfy the last naturality axiom). Then similarly to the proof of Lemma 4.10, we may then construct a homotopy relative to Sub_s , between the pair of maps $\tilde{u}_1(m_1, \dots, m_{N+1}, N+1)$, $u_1(m_1, \dots, m_{N+1}, N+1)$. Taking the composite homotopy we clearly obtain the needed homotopy. As the choice of (m_1, \dots, m_s) satisfying $D(m_1, \dots, m_s) = s = N+1$, was general, this concludes the proof of the claim. \square

4.1. Outline of an explicit construction. This section is not logically necessary, but in order to give the reader more intuition we now give a partial explicit construction of the maps that would satisfy the axioms. To be more formal, we give an explicit construction of the system of satisfying condition $S(4)$, as in the proof of Theorem 4.8. However, we will not formally check all the properties. (Although this would be straightforward.) This construction could be in principle extended to all generality but at the cost of much complexity.

Fix a geometric model for $\overline{\mathcal{R}}_d$, for example as the Stasheff associahedra. When $d = 4$ this is a pentagon. Recall that to each corner of $\overline{\mathcal{R}}_4$ we have a uniquely associated nodal Riemann surface with 3 components and 5 marked points, one of which is called the root. Recall that we label the root component by ω , the next component by β and the component furthest from root by α . (With respect to the linear ordering described earlier.) Denote by M_α the collection of marked points, different from the root e_0 , on α , likewise with β, ω . This determines a sub-composable sequence $mor(S_\alpha)$ of a composable sequence (m_1, \dots, m_4) , and likewise with β, ω , (note that M_ω, M_β could be empty).

Let r be in the gluing normal neighborhood of some corner, corresponding to non-zero gluing parameters $d_{\alpha, \beta}, d_{\beta, \omega}$. We now construct a map

$$f_r = f_r(m_1, \dots, m_4) : [0, 4] \times [0, 1] \rightarrow \Delta^4$$

In what follows by *concatenation* of a collection of paths we mean their product in the Moore path category of Δ^4 , the notation for composition will be assumed to be diagrammatic. This is the category with objects: points of Δ^4 , and morphisms from x_0 to x_1 : continuous paths $[0, T] \rightarrow \Delta^4$, $T > 0$, between x_0, x_1 , with composition the natural concatenation of paths. Note that this is quite different from our previously defined groupoid $\Pi(\Delta^4)$.

For a morphism m in $\Pi(\Delta^4)$ let $s(m)$ and $t(m)$ denote the source respectively target of m . Let $H^m : \Delta^4 \times [0, 1] \rightarrow \Delta^4$ denote the natural deformation retraction of Δ^4 onto the edge determined by $s(m), t(m)$, with time 1 map the orthogonal affine projection onto this edge (for the standard metric on Δ^n). Set $H_\tau^m = H^m|_{\Delta^n \times \{\tau\}}$. Next, for a general piece-wise affine path $p : [0, T] \rightarrow \Delta^4$, with end points $s(m), t(m)$, we have a homotopy $H_\tau^m \circ p$, $\tau \in [0, 1]$, from p to a path

$$\tilde{p} : [0, T] \rightarrow \Delta^4$$

with image in the edge determined by m . Let $D(p, \tau)$, $\tau \in [0, 1]$, be the concatenation of $H_\tau^m \circ p$ with the homotopy G_τ , $\tau \in [0, 1]$, of paths with fixed end points, from \tilde{p} to the map

$$\tilde{m} : [0, T] \rightarrow \Delta^4$$

linearly parametrizing the edge determined by m . This second homotopy G_τ can be chosen in a way that depends only on \tilde{p} . This can be done explicitly, using piece-wise linearity of p .

The map f_r^t from the $y = t$ slice $[0, 4] \times \{t\}$ is constructed as follows. Set

$$I_\alpha = (1 - d_{\alpha, \beta})/2,$$

set $f_{\alpha, r}$ to be the concatenation of the morphisms in $\text{mor}(M_\alpha)$. That is if

$$M_\alpha = (m_1^\alpha, \dots, m_k^\alpha)$$

then

$$f_{\alpha, r} = m_1^\alpha \cdot \dots \cdot m_k^\alpha.$$

Then for $t \in [0, I_\alpha]$ set $f_{\alpha, r}^t = D(f_{\alpha, r}, 2t)$. Then set f_r^t to be the concatenation of morphisms in $\text{mor}(M_\beta)$, $\text{mor}(M_\gamma)$ and of $f_{\alpha, r}^t$, in that order, although note that the order of the morphisms in the concatenation is uniquely determined by the end point conditions, this holds further on as well.

Next set

$$I_\beta = I_\alpha + (1 - I_\alpha)(1 - d_\beta)/2.$$

If α and β components have a nodal point in common we set

$$f_{\beta, r} : [0, 4] \times \{t\} \rightarrow \Delta^4$$

to be the concatenation of $f_{\alpha, r}^{I_\alpha}$ with morphisms in $\text{mor}(M_\beta)$, and for $t \in [I_\alpha, I_\beta]$ we set

$$f_{\beta, r}^t = D(f_{\beta, r}, \frac{2(t - I_\alpha)}{1 - I_\alpha}).$$

And then for $t \in [I_\alpha, I_\beta]$ set f_r^t to be the concatenation of morphisms in $\text{mor}(M_\omega)$ and of $f_{\beta, r}^t$.

Finally set $f_{\omega, r}$ to be the concatenation of $f_{\beta, r}^{I_\beta}$ with morphisms in $\text{mor}(M_\omega)$, and for $t \in [I_\beta, 1]$ set

$$f_r^t = D(f_{\omega, r}, \frac{2(t - I_\beta)}{1 - I_\beta}).$$

When α has a nodal point in common with the ω component set $f_{\beta, r}$ to be the concatenation of morphisms in $\text{mor}(M_\beta)$, and for $t \in [I_\alpha, I_\beta]$ set

$$f_{\beta, r}^t = D(f_{\beta, r}, \frac{2(t - I_\alpha)}{1 - I_\alpha}).$$

Then for $t \in [I_\alpha, I_\beta]$ set f_r^t to be the concatenation of morphisms $f_r^{I_\alpha}$ and $f_{\beta, r}^t$, and $\text{mor}(M_\omega)$ (although $\text{mor}(M_\omega)$ in this particular case is empty, we add this so




FIGURE 5. The uncolored enclosed regions labeled T_α , T_β surrounding segments m_α , m_β are “thin”.

that the degenerate case $M_\alpha = \emptyset$, $M_\beta = \emptyset$ makes sense, see the discussion below). Finally for $t \in [I_\beta, 1]$ set

$$f_r^t = D(f_r^{I_\beta}, \frac{2(t - I_\beta)}{1 - I_\beta}).$$

When $r \in \overline{\mathcal{R}}_4$ is in the gluing neighborhood of a face but not of a corner the construction of $f_r : [0, 4] \times [0, 1] \rightarrow \Delta^4$ is similar, in fact we can think of it as a special case of the above by setting $d_\beta = 1$, $M_\beta = \emptyset$. When $r \in \overline{\mathcal{R}}_4$ is not in the gluing neighborhood of the boundary we can also think of this as a special case of the above, with $M_\alpha = \emptyset$, $M_\beta = \emptyset$, $d_\alpha = 1$, $d_\beta = 1$ in the construction above.

4.1.1. *Retracting \mathcal{S}_r onto $[0, 4] \times [0, 1]$.* We now construct a smooth r -family of maps

$$ret_r : \mathcal{S}_r \rightarrow [0, 4] \times [0, 1],$$

$r \in \mathcal{R}_4$, suitably compatible with the maps

$$f_r : [0, 4] \times [0, 1] \rightarrow \Delta^4.$$

In figure 5 (a), (b), (c) represent cases where (c): r is not within gluing normal neighborhood of boundary, (b): r is in a gluing neighborhood of a side but not a corner and (a): r is within gluing neighborhood of a corner, (we picked a particular corner and side for these diagrams). The color shading will be explained in a moment. In each case (a), (b), (c) we first color shade $[0, 4] \times [0, 1]$ as in figure 6, the green region is the domain of $f_{\alpha,r}^t$ contained in $[0, 4] \times [0, I_\alpha]$, in the blue regions the map f_r is vertically constant, the red region is the domain of $f_{\beta,r}^t$ contained in $[0, 4] \times [I_\alpha, I_\beta]$ and yellow region is the rest of the domain of f_r . The maps ret_r are defined for each r by taking color shaded areas to color shaded areas, so that the following holds.

- (1) The ends e_0, e_1, \dots, e_4 of \mathcal{S}_r , colored in purple, are identified in strip coordinate charts as $[1, \infty) \times [0, 1]$ and in these coordinates ret_r is the composition of the projection $[1, \infty) \times [0, 1] \rightarrow [0, 1]$ with the map to the boundary




FIGURE 6. Diagram for S_d . Solid black border is boundary, while dashed red lines are open ends. The connection $\mathcal{A}(r, \{L_i\})$ preserves Lagrangians L_i over boundary components labeled L_i .

of $[0, 4] \times [0, 1]$, so that composition with f_r parametrizes the morphism $m_0 = m_1 \cdot \dots \cdot m_4, m_1, \dots, m_4$ in $\Pi(\Delta^4)$, respectively.

- (2) The boundary of \mathcal{S}_r goes either to the boundary of $[0, 4] \times [0, 1]$ or to the vertical boundary lines between colored regions.
- (3) The unshaded “thin” regions labeled T_α, T_β come from the gluing construction and are identified with $[0, 1] \times [-\phi(\tau_\alpha), \phi(\tau_\alpha)]$, respectively $[0, 1] \times [-\phi(\tau_\beta), \phi(\tau_\beta)]$. In these coordinates ret_r on T_α, T_β is the projection to $[0, 1]$ composed with a diffeomorphism onto the lower edge of green, respectively red region, (linear in respective natural coordinates).
- (4) The unshaded part of \mathcal{S}_r is collapsed onto the horizontal line bounding yellow region of $[0, 4] \times [0, 1]$.
- (5) Blue shaded regions are identified in strip coordinate charts $[0, \infty] \times [0, 1] \rightarrow \Sigma_r$, as $[0, 1] \times [0, 1]$, and are mapped to the corresponding blue regions in $[0, 4] \times [0, 1]$.

(The above prescription naturally extends to the boundary $\overline{\mathcal{R}}_4$.)

We then set

$$u(m_1, \dots, m_4, \Delta^4)|_{\mathcal{S}_r} = f_r \circ ret_r.$$

These are almost the maps we want but we need to collar them near the boundary of $\overline{\mathcal{R}}_4$ so that Axiom 1 of naturality is satisfied. We omit the details.

5. AUXILIARY DATA \mathcal{D}

Let

$$M \hookrightarrow P \xrightarrow{\pi} X$$

be a Hamiltonian fiber bundle, with model fiber (M^{2n}, ω) that we shall assume here to be a closed, monotone:

$$[\omega] = \text{const} \cdot 2c_1(TM),$$

$\text{const} > 0$, symplectic manifold. The constant const is called the monotonicity constant.

We now discuss geometric-analysis theoretic data needed for the construction of the functor $F_P : \Delta(X) \rightarrow A_\infty - Cat^{unit}$, as outlined in Section 1.1 of the introduction. Essentially, this data $\mathcal{D} = \mathcal{D}(P)$ specifies various choices of Hamiltonian connections, as well as certain choices of almost complex structures. These choices are to be made for each $\Sigma \in Simp(X)$, while being suitably compatible, so that we obtain our functor F_P .

Let (M, ω) be a monotone symplectic manifold with monotonicity constant $const$. We say that a Lagrangian submanifold $L \subset M$ is *monotone* if the homomorphisms given by symplectic area and Maslov class

$$[\omega] : H_2(M, L) \rightarrow \mathbb{R}, \quad \mu : H_2(M, L) \rightarrow \mathbb{Z}$$

are proportional:

$$[\omega] = const \cdot \mu.$$

For an $x : pt \rightarrow X$, define

$$(5.1) \quad F(x) = F_P(x),$$

to be the set of oriented, spin, monotone Lagrangian submanifold L in

$$P_x = x^*P \simeq M$$

with minimal (positive) Maslov number at least 2, and such that the inclusion map $\pi_1(L) \rightarrow \pi_1(M)$ vanishes. We call elements of $F_P(x)$ **objects**. These will in fact be objects of a certain category to be constructed.

Let $L \in F_P(x)$, and j be an almost complex structure on P_x compatible with ω_x the symplectic form on P_x . Let $\mathcal{M}(L, j)$ denote the moduli space of Maslov number 2 j -holomorphic discs in P_x , with one marked point on the boundary, with boundary of the disk going to L . It is well known, c.f. [25, Section 2.3] (which also contains a number of additional references) that for a generic such j , $\mathcal{M}(L, j)$ is regular, is a transversely cut out n -dimensional manifold and is compact. The compactness is due to the following fact. If $\mathcal{M}(L, j)$ were not compact, then by Gromov compactness there would be a sequence of curves in

$$\mathcal{M}(L, j)$$

degenerating to a nodal curve with at least a pair of components both of which must have Maslov number at least 2, (a spherical bubble would likewise contribute positively to the total Maslov number of the nodal curve) by our assumption on the minimal Maslov number. (The monotonicity, and energy positivity preclude negative Maslov number components.) This would clearly be a contradiction, by additivity of the Maslov number.

Then we have an evaluation map at the marked point:

$$ev : \mathcal{M}(L, j) \rightarrow L,$$

and we define $\omega(L) \in \mathbb{Z}$ as the degree of ev .

Given a smooth

$$\Sigma : \Delta^n \rightarrow X,$$

set $x_i := \Sigma(i) \in X$, $i \in \{0, \dots, n\}$ a corner of Δ^n . Also denote by x_i the corresponding inclusion map $x_i : pt \rightarrow X$. Set

$$(5.2) \quad F_P(\Sigma) := \bigsqcup_i F_P(x_i),$$

at the moment this is just a set of Lagrangians, but later on this will be the set of objects of a certain A_∞ category, (with the same name).

Given a pair of objects

$$L_0, L_1 \in F_P(\Sigma),$$

satisfying

$$\omega(L_0) = \omega(L_1),$$

and such that $L_0 \in F_P(x_i), L_1 \in F_P(x_j)$, let

$$m = m_{L_0, L_1} : [0, 1] \rightarrow \Delta^n$$

denote the edge between i, j corners of Δ^n , $i, j \in \{0, \dots, n\}$. We then set

$$\overline{m} := \Sigma \circ m.$$

For each such Σ , for each L_0, L_1 as above, the data \mathcal{D} prescribes a Hamiltonian connection

$$\mathcal{A}(L_0, L_1) = \mathcal{A}(L_0, L_1, \overline{m})$$

on \overline{m}^*P . (See Section 3.3 for definition of Hamiltonian connections.)

Denote by $\mathcal{A}(L_0, L_1)(L_0)$ the image in P_{x_j} of the $\mathcal{A}(L_0, L_1)$ -parallel transport of L_0 over $[0, 1]$. Then we require that $\mathcal{A}(L_0, L_1)(L_0)$ be transverse to L_1 .

Definition 5.1. *Let*

$$S(L_0, L_1) = S(L_0, L_1, \mathcal{A}(L_0, L_1))$$

*denote the space of $\mathcal{A}(L_0, L_1)$ -flat sections with boundary on L_0, L_1 , over 0 respectively over 1. In other words elements of $S(L_0, L_1)$ are sections $\gamma : [0, 1] \rightarrow \overline{m}^*P$, with vanishing $\mathcal{A}(L_0, L_1)$ -covariant derivative, and satisfying $\gamma(0) \in L_0$ and $\gamma(1) \in L_1$. By a **starting position** of an element $\gamma \in S(L_0, L_1)$ we mean $\gamma(0) \in L_0$. Likewise by an **ending position** of an element $\gamma \in S(L_0, L_1)$ we mean $\gamma(1) \in L_1$.*

Definition 5.2. *For each m as above, a family $\{j_t\} = \{j_t(L_0, L_1, \overline{m})\}$ of fiber-wise almost complex structures on \overline{m}^*P is said to be **admissible with respect to** $\mathcal{A}(L_0, L_1)$ if the following holds.*

- *Let $P_{\overline{m}(t)}$ denote the fiber of \overline{m}^*P over $\overline{m}(t)$. Then for each t , Chern number 1 j_t -holomorphic spheres in $P_{\overline{m}(t)} \subset \overline{m}^*P$ do not intersect any of the images of any elements of $S(L_0, L_1)$.*
- *The moduli spaces $\mathcal{M}(L_0, j_0), \mathcal{M}(L_1, j_1)$ are regular, and the evaluation map*

$$ev_0 : \mathcal{M}(L_0, j_0) \rightarrow L_0$$

does not intersect the set of starting positions of elements of $S(L_0, L_1)$. Likewise the evaluation map

$$ev_1 : \mathcal{M}(L_1, j_1) \rightarrow L_1$$

does not intersect the set of ending positions of elements of $S(L_0, L_1)$.

Such a family $\{j_t\}$ is easily seen to exist, cf. [25]. Our data \mathcal{D} then fixes a choice of such $\{j_t\}$ for each Σ, m as above. Next \mathcal{D} makes a choice of a natural system \mathcal{U} as previously described. Finally \mathcal{D} will specify a certain natural system of Hamiltonian connections and a system of complex structures that we now describe. This is to be done for all choices of certain Lagrangian labels. This is a bit notationally complicated but ultimately trivial, given the main geometric input of the system

\mathcal{U} . It is just a system of compatible perturbations in the sense of Seidel [24, Section 9i], but relative to \mathcal{U} . We now describe the ingredients for this.

Given a composable chain (m_1, \dots, m_d) and a map

$$u(m_1, \dots, m_d, n) : \mathcal{S}_d^\circ \rightarrow \Delta^n \text{ that is part of a natural system } \mathcal{U},$$

we have an induced fibration

$$M \hookrightarrow \tilde{S}(m_1, \dots, m_d, \Sigma^n) \rightarrow \mathcal{S}_d^\circ$$

by pulling back $M \hookrightarrow P \rightarrow X$ first by $\Sigma : \Delta^n \rightarrow X$ and then by $u(m_1, \dots, m_d, n)$. We have a natural projection

$$\tilde{S}(m_1, \dots, m_d, \Sigma_n) \rightarrow \overline{\mathcal{R}}_d$$

and we denote the fiber over $r \in \overline{\mathcal{R}}_d$ by $\tilde{S}(m_1, \dots, m_d, \Sigma, r)$, or simply by $\tilde{\mathcal{S}}_r$ where there can be no confusion. So $\tilde{\mathcal{S}}_r$ is naturally a Hamiltonian M -fibration over the surface \mathcal{S}_r , smooth over smooth components. To state this another way,

$$\tilde{\mathcal{S}}_r = (u(m_1, \dots, m_d, n, r) \circ \Sigma)^* P,$$

where

$$(5.3) \quad u(m_1, \dots, m_d, n, r) = u(m_1, \dots, m_d, n)|_{\mathcal{S}_r}.$$

By the partial naturality properties of the maps $u(m_1, \dots, m_d, n, r)$, at each e_i end, $1 \leq i \leq d$, of \mathcal{S}_r , we have natural trivializations

$$[0, \infty) \times \overline{m}_i^* P \rightarrow \tilde{\mathcal{S}}_r.$$

Similarly at the e_0 end. We shall call these *distinguished trivializations/coordinates*.

5.1. Lagrangian labels and admissible connections. Given $r \in \mathcal{R}_d$, and given choices

$$L_i \in F(\overline{m}_{i+1}(0)), \text{ for } 0 \leq i \leq d-1, \quad L_d \in F(\overline{m}_d(1)),$$

a *labeling* is the following construction. We assign L_i to the component of $\partial\mathcal{S}_r$ between the punctures $i, i+1$, $0 \leq i \leq d-1$ and we assign L_d to the component of $\partial\mathcal{S}_r$ between $d, 0$. We suppose in addition that $\omega(L_i) = \omega(L_d)$ for all i .

Extend the labeling strategy naturally to \mathcal{S}_r , with $r \in \partial\mathcal{R}_d$. In other words if we glue at some node of \mathcal{S}_r then each boundary component of the glued surface inherits a consistent label. See Figure 7 below. For the moment we do not specify any dependence of the labels on r .

Definition 5.3. We say that a Hamiltonian connection (cf. Section 3.3) \mathcal{A} on the Hamiltonian fibration $\tilde{S}(m_1, \dots, m_d, \Sigma, r) \rightarrow \mathcal{S}_r$, $r \in \overline{\mathcal{R}}_d$, is **admissible** with respect to a labeling L_0, \dots, L_d if:

- Parallel transport by \mathcal{A} over the boundary component(s) of $\partial\mathcal{S}_r$ labeled L_i preserves the Lagrangian L_i , $0 \leq i \leq d$. This condition is unambiguous, as by construction and by the partial naturality properties of Definition 4.2, over such a boundary component, $\tilde{\mathcal{S}}_r$ has a distinguished trivialization as $[0, 1] \times P_{s(m_{i+1})}$, $0 \leq i \leq d-1$, or as $[0, 1] \times P_{t(m_d)}$ in the case of L_d . Note that we have such trivializations, over the boundary components, also when $r \in \partial\mathcal{R}_d$. So that the above condition also makes sense for nodal \mathcal{S}_r .




FIGURE 7.

- For $1 \leq i \leq d$, in the distinguished coordinates $[0, \infty) \times \overline{m}_i^* P$, at each e_i end, $\mathcal{A} = pr^* \mathcal{A}(L_{i-1}, L_i, \overline{m}_i)$ for $pr : [0, 1] \times [1, \infty) \rightarrow [0, 1]$ the projection. Likewise, at the e_0 end in the distinguished coordinates $(-\infty, 0] \times \overline{m}_0^* P$, $\mathcal{A} = pr^* \mathcal{A}(L_{i-1}, L_i, \overline{m}_0)$ for $pr : [0, 1] \times (-\infty, 0] \rightarrow [0, 1]$ the projection.

We do not yet impose any conditions at the nodes, but certain conditions will be forced by the additional properties of the Definition 5.5 below. For later, we mention that \mathcal{D} will make a choice of such a connection for all possible L_0, \dots, L_d as above.

We also have a Lagrangian sub-fibration of

$$\tilde{\mathcal{S}}_r \rightarrow \mathcal{S}_r$$

over the boundary of \mathcal{S}_r , whose fiber over an element of the boundary component labeled L_i is L_i , with respect to the distinguished trivializations mentioned above. (This naturally extends to the case \mathcal{S}_r is nodal.)

We name this sub-fibration by

$$(5.4) \quad \mathcal{L}(\mathcal{U}, L_0, \dots, L_d, r),$$

so that if \mathcal{A} is admissible as above it preserves this sub-fibration. Denote by $\mathcal{T}(L_0, \dots, L_d, \Sigma, r)$ the space of Hamiltonian connections on

$$M \hookrightarrow \tilde{\mathcal{S}}(m_1, \dots, m_d, \Sigma, r) \rightarrow \mathcal{S}_r$$

admissible with respect to L_0, \dots, L_d .

5.2. Gluing admissible connections. Given an element \mathcal{A} in $\mathcal{T}(L_0, \dots, L_{s_1}, \Sigma, r)$ and an element

$$\mathcal{A}' \in \mathcal{T}(L'_0, \dots, L'_{i-1}, L'_i, L'_{i+1}, \dots, L'_{s_2}, \Sigma, r'),$$

with $L'_{i-1} = L_0, L'_i = L_{s_1}$, we have a connection naturally induced by the gluing maps St_i , denoted by

$$St_i(\mathcal{A}, \mathcal{A}', 0) \in \mathcal{T}(L'_0, \dots, L'_{i-2}, L_0, \dots, L_{s_1}, L'_{i+2}, \dots, L'_{s_2}, \Sigma, St_i(r, r', 0)).$$

Such a pair $\mathcal{A}, \mathcal{A}'$ will be called *composable*.

Using the second property in the Definition 5.3 we can naturally extend this to a family of connections $\{St_i(\mathcal{A}, \mathcal{A}', \tau)\}_\tau$, so that

$$St_i(\mathcal{A}, \mathcal{A}', \tau) \in \mathcal{T}(L'_0, \dots, L'_{i-2}, L_0, \dots, L_{s_1}, L'_{i+2}, \dots, L'_{s_2}, \Sigma, St_i(r, r', \tau)),$$

for $0 \leq \tau < 1$, and so that in addition we have the following. Outside of the thin region $thin_{\tau,i}$ of $\mathcal{S}_{St_i(r, r', \tau)}$ the connection $St_i(\mathcal{A}, \mathcal{A}', \tau)$ is naturally identified with $St_i(\mathcal{A}, \mathcal{A}', 0)$. On the other hand over $thin_{\tau,i}$, for $\tau > 0$, $\tilde{\mathcal{S}}_{St_i(r, r', \tau)}$ is naturally isomorphic to $(\overline{m}_i)^*P \times [-\phi(\tau), \phi(\tau)]$ by the Axiom 1 of the natural system \mathcal{U} , and where ϕ is as in (4.2). Over $thin_{\tau,i}$, in the above coordinate representation, $St_i(\mathcal{A}, \mathcal{A}', \tau)$ is the trivial in the second variable extension of the connection $\mathcal{A}(L_{i-1}, L_i, \overline{m}_i)$. (Analogously, to the second part of the Definition 5.3).

5.3. Admissible fiber almost complex structures.

Definition 5.4. We say that a family $\{j_z\}$ of fiber-wise, $\{\omega_z\}$ -compatible, almost complex structures on the Hamiltonian M -fibration $\tilde{\mathcal{S}}(m_1, \dots, m_d, \Sigma, r) \rightarrow \mathcal{S}_r$ is **admissible** with respect to L_0, \dots, L_d if:

- At the i 'th end of \mathcal{S}_r , $1 \leq i \leq d$ in the distinguished trivialization

$$[0, \infty) \times \overline{m}_i^*P \rightarrow \tilde{\mathcal{S}}(m_1, \dots, m_d, \Sigma, r),$$

the family $\{j_z\}$ coincides with the canonical, in the factor $[0, \infty)$, \mathbb{R} -translation invariant extension of the family $\{j_t(L_{i-1}, L_i, \overline{m}_i)\}$, with the latter as in Definition 5.2. More specifically, in the distinguished coordinates $[0, \infty) \times \overline{m}_i^*P$,

$$\forall (z = (t, s)) \in [0, 1] \times [0, \infty) : j_{(t,s)} = j_t(L_{i-1}, L_i, \overline{m}_i).$$

Similarly, at the e_0 end.

5.4. Gluing admissible fiber almost complex structures. Denote by

$$\mathcal{J}(L_0, \dots, L_s, \Sigma, r)$$

the space of families of fiberwise almost complex structures $\{j_z\}$ on

$$\tilde{\mathcal{S}}(m_1, \dots, m_s, \Sigma, r)$$

admissible with respect to L_0, \dots, L_s . Given an element $\{j_z\}$ in $\mathcal{J}(L_0, \dots, L_{s_1}, \Sigma, r)$ and an element

$$\{j'_z\} \in \mathcal{J}(L'_0, \dots, L'_{i-2}, L_0, L_{s_1}, L'_{i+1}, \dots, L'_{s_2}, \Sigma, r')$$

we have an induced element:

$$St_i(\{j_z\}, \{j'_z\}, \tau) \in \mathcal{J}(L'_0, \dots, L'_{i-2}, L_0, \dots, L_{s_1}, L'_{i+2}, \dots, L'_{s_2}, \Sigma, St_i(r, r', \tau)),$$

for each $0 \leq \tau < 1$. Such a pair $\{j_z\}, \{j'_z\}$ will be called *composable*.

5.5. Combining admissible connections and fiber almost complex structures.

Definition 5.5. Let $M \hookrightarrow P \rightarrow X$ be as above. A **system** $\mathcal{F} = \mathcal{F}(P)$ of connections, and almost complex structures compatible with a system of maps \mathcal{U} is an element of

$$\prod_{\Sigma \in \text{Simp}(X)} \prod_{s \geq 2} \prod_{\{(L_0, \dots, L_s) | L_i \in F(\Sigma)\}} \prod_{r \in \overline{\mathcal{R}}_s} \mathcal{T}(L_0, \dots, L_s, \Sigma, r) \times \mathcal{J}(L_0, \dots, L_s, \Sigma, r).$$

The projection of \mathcal{F} onto the $(\Sigma, s, (L_0, \dots, L_s), r)$ component will be denoted by $\mathcal{F}(L_0, \dots, L_s, \Sigma, r)$. Similarly as with \mathcal{U} , it might help to phrase this in functional language. Let

$$O = \{(L_0, \dots, L_s, \Sigma, r) | s \in \mathbb{N}_{\geq 2}, \Sigma \in \text{Simp}(X), \\ L_i \in F(\Sigma), r \in \overline{\mathcal{R}}_s\},$$

then set theoretically the product above is the set of certain maps \mathcal{F} with domain O . Then in this language $\mathcal{F}(L_0, \dots, L_s, n, r)$ is just the value $\mathcal{F}(L_0, \dots, L_s, n, r)$, of the map \mathcal{F} .

For shorthand we say that a Hamiltonian connection $\mathcal{A} \in \mathcal{F}$ if it is of the form $pr_1 \mathcal{F}(L_0, \dots, L_s, \Sigma, r)$, for some $(L_0, \dots, L_s, \Sigma, r)$ and for pr_i , $i = 1, 2$, the projections

$$pr_1 : \mathcal{T}(L_0, \dots, L_s, \Sigma, r) \times \mathcal{J}(L_0, \dots, L_s, \Sigma, r) \rightarrow \mathcal{T}(L_0, \dots, L_s, \Sigma, r) \\ pr_2 : \mathcal{T}(L_0, \dots, L_s, \Sigma, r) \times \mathcal{J}(L_0, \dots, L_s, \Sigma, r) \rightarrow \mathcal{J}(L_0, \dots, L_s, \Sigma, r).$$

Definition 5.6. We say that \mathcal{F} is **natural** if:

- (1) The families of connections/almost complex structures are smooth in the parameter r , over smooth components of the surfaces \mathcal{S}_r . (At the nodes the behavior will be characterized by the following properties.)
- (2) For a composable pair $\mathcal{A}, \mathcal{A}' \in \mathcal{F}$ as above the connection $St_i(\mathcal{A}, \mathcal{A}', 0)$ coincides with

$$pr_1 \mathcal{F}(L'_0, \dots, L'_{i-2}, L_0, \dots, L_{s_1}, L'_{i+1}, \dots, L'_{s_2}, \Sigma, St_i(r, r', 0)).$$

- (3) The pair of connections,

$$St_i(\mathcal{A}, \mathcal{A}', 0),$$

$$pr_1 \mathcal{F}(L'_0, \dots, L'_{i-2}, L_0, \dots, L_{s_1}, L'_{i+1}, \dots, L'_{s_2}, \Sigma, St_i(r, r', 0)),$$

also agree for a sufficiently small $\epsilon > 0$ on the “thin part” of $\mathcal{S}_{St_i(r, r', \epsilon)}$.

- (4) Given a face map $f : \Delta^{n-1} \rightarrow \Delta^n$, by the second naturality property of maps $u(m_1, \dots, m_d, n)$ determined by \mathcal{U} , there is bundle map

$$p : \tilde{\mathcal{S}}(m_1, \dots, m_d, \Sigma \circ f, r) \rightarrow \tilde{\mathcal{S}}(f(m_1), \dots, f(m_d), \Sigma, r),$$

and we ask that the pullback map takes $pr_1 \mathcal{F}(p(L_0), \dots, p(L_d), \Sigma, r)$ to

$$pr_1 \mathcal{F}(L_0, \dots, L_d, \Sigma \circ f, r).$$

- (5) There are analogous conditions on the families of almost complex structures $pr_2 \mathcal{F}(L_0, \dots, L_s, \Sigma^n, r)$ that we will not state.

Notation 5.7. We will sometimes write by abuse of notation $\mathcal{F}(\dots)$, for either the connection $pr_1\mathcal{F}(\dots)$, or the family of almost complex structures $pr_2\mathcal{F}(\dots)$, since there usually can be no confusion.

Lemma 5.8. A natural system \mathcal{F} compatible with a given \mathcal{U} exists and is unique up to homotopy.

Proof. When $n = 0$ this is the classical Fukaya category case, and the proof of existence of a natural system is given in Seidel [24, Section 9i] in the language of what Seidel calls compatible system of perturbations, which is completely analogous to our connection language. Although, in Seidel's book only the case of exact Lagrangians in exact symplectic manifolds is considered, this readily extends to our context, since we are not yet concerned with compactness or regularity properties.

In what follows we write Σ^n for an element of $\text{Simp}(X)$ of the form $\Sigma^n : \Delta^n \rightarrow X$. Now suppose that we have chosen an element

$$\mathcal{F} \in \prod_{r \in \overline{\mathcal{R}}_s} \prod_{\{(L_0, \dots, L_s) | L_i \in F(\Sigma^n)\}} \prod_{s \geq 2} \prod_{\{\Sigma^n | n \leq N\}} \mathcal{T}(L_0, \dots, L_s, \Sigma^n, r) \times \mathcal{J}(L_0, \dots, L_s, \Sigma^n, r)$$

satisfying naturality. We need to extend this to an element of

$$\prod_{r \in \overline{\mathcal{R}}_s} \prod_{\{(L_0, \dots, L_s) | L_i \in F(\Sigma^n)\}} \prod_{s \geq 2} \prod_{\{\Sigma^n | n \leq N+1\}} \mathcal{T}(L_0, \dots, L_s, \Sigma^n, r) \times \mathcal{J}(L_0, \dots, L_s, \Sigma^n, r),$$

also satisfying naturality.

Let $\Sigma^{N+1} : \Delta^{N+1} \rightarrow X$ be given. Let $L_0, \dots, L_s \in F(\Sigma^{N+1})$, so that each $L_i \in F(x_i)$ for $x_i = \Sigma^{N+1}(v_i)$ for some vertex $v_i \in \Delta^{N+1}$. In particular the set $\{L_0, \dots, L_s\}$ determines the set of vertices $\{v_0, \dots, v_s\}$ of Δ^{N+1} . Denote by $D(L_0, \dots, L_s)$ the least dimension of a subsimplex of Δ^{N+1} with vertices $\{v_0, \dots, v_s\}$. There is a unique extension of \mathcal{F} to an element

$$(5.5) \quad \mathcal{F} \in \prod_{r \in \overline{\mathcal{R}}_s} \prod_{\{(L_0, \dots, L_s) | N \geq D(L_0, \dots, L_s)\}} \prod_{s \geq 2} \prod_{\{\Sigma^n | n \leq N+1\}} \mathcal{T}(L_0, \dots, L_s, \Sigma^n, r) \times \mathcal{J}(L_0, \dots, L_s, \Sigma^n, r)$$

satisfying the naturality condition. We need to extend to the case $N+1 = D(L_0, \dots, L_s)$ and so that naturality is satisfied. For all (L_0, \dots, L_s) with

$$D(L_0, \dots, L_s) = N+1,$$

and given Σ^{N+1} , the naturality condition and \mathcal{F} from (5.5) determine

$$\mathcal{F}(L_0, \dots, L_s, \Sigma^{N+1}, r)$$

for r in the boundary of $\overline{\mathcal{R}}_s$, see the discussion following (4.10). Set

$$\mathcal{P} := \bigcup_{r \in \overline{\mathcal{R}}_s} \mathcal{T}(L_0, \dots, L_s, \Sigma^n, r) \times \mathcal{J}(L_0, \dots, L_s, \Sigma^n, r).$$

So we have a natural fibration $\mathcal{P} \rightarrow \overline{\mathcal{R}}_s$, with the fiber over $r \in \overline{\mathcal{R}}_s$ denoted by \mathcal{P}_r . The topology on \mathcal{P} is the natural metric ‘‘Gromov topology’’, constructed using gluing operations of Sections 5.2, 5.4. The fibers are non-empty, the corresponding statement for just connections follows by [1, Lemma 3.2]. The fibers are contractible, for the connection component this is just because the relevant space is naturally affine. For the almost complex structure component, this is basically classical by work of Gromov [6]. Moreover, this fibration is a Serre fibration, this

is only non-obvious at boundary points of $\overline{\mathcal{R}}_s$, but there the corresponding lifting property for cubes can be easily verified directly, again using the gluing operations of Sections 5.2, 5.4.

To summarize we have a Serre fibration $\mathcal{P} \rightarrow \overline{\mathcal{R}}_s$ with non-empty contractible fibers. We have a section of \mathcal{P} over $\partial\overline{\mathcal{R}}_s$ corresponding to the partially constructed family

$$\{\mathcal{F}(L_0, \dots, L_s, \Sigma^{N+1}, r)\}_{r \in \partial\overline{\mathcal{R}}_s}$$

above. By the classical obstruction theory, there is an extension to a section over $\overline{\mathcal{R}}_s$. The naturality conditions for the corresponding family then follow by construction. (We may need to homotopically adjust the section to satisfy the Property 2,3 of naturality, but this is straightforward.) Uniqueness up to homotopy is obvious from the argument above. \square

5.6. The summary of the perturbation data \mathcal{D} . Let $M \hookrightarrow P \rightarrow X$ be as above. To summarize, the perturbation data $\mathcal{D} = \mathcal{D}(P)$ consists of a choice of a natural system of maps \mathcal{U} , and a choice of a system $\mathcal{F} = \mathcal{F}(P)$ of connections/almost complex structures compatible with \mathcal{U} .

Proposition 5.9. *Any pair $\mathcal{D}_0(P), \mathcal{D}_1(P)$ of perturbation data for P are homotopic. That is there is a family $\{\tilde{\mathcal{D}}_t\}$, $t \in [0, 1]$, smooth in t , with $\tilde{\mathcal{D}}_i = \mathcal{D}_i$, $i = 0, 1$.*

Proof. This is just a direct consequence of Theorem 4.8 and Lemma 5.8. \square

6. THE FUNCTOR F

Let $A_\infty - Cat$ denote the category of small \mathbb{Z}_2 graded A_∞ categories over \mathbb{Q} , with morphisms fully-faithful embeddings, as defined below, that are in addition quasi-equivalences.

Definition 6.1. *We say that an A_∞ functor G is a **fully-faithful embedding**, if G has vanishing higher order components, is injective on objects and if the first component map on hom spaces is an isomorphism of chain complexes. In other words G above is just an identification map of a full A_∞ sub-category.*

We now describe the construction of the functor

$$F_{P, \mathcal{D}} : \text{Simp}(X) \rightarrow A_\infty - Cat$$

associated to a Hamiltonian fibration P and the chosen auxiliary data \mathcal{D} , described in the previous section. In what follows we usually drop \mathcal{D} and P from notation. For a point $x : pt \rightarrow X$ the associated category will be constructed following Sheridan [25]. In fact the analysis does not change for the case of higher dimensional simplices $\Delta^n \rightarrow X$, the geometry however needs to be substantially generalized.

6.1. F on a point. For $x : pt \rightarrow X$, $F(x)$ is defined to be the Fukaya A_∞ category $Fuk(x^*P)$ whose set of objects is the set $F_P(x)$ discussed in Section 5, cf. (5.1). For a pair $L_0, L_1 \in F(x)$, with $\omega(L_0) \neq \omega(L_1)$ we set $hom(L_0, L_1) = 0$, (to avoid dealing with curved A_∞ categories), otherwise we set

$$hom(L_0, L_1) = CF(L_0, L_1, \mathcal{D}),$$

where the latter is a \mathbb{Z}_2 graded Floer chain complex over \mathbb{Q} that is defined as follows.

Let $\mathcal{A}(L_0, L_1)$ be the Hamiltonian connection on $P_x \times [0, 1]$ determined by the chosen data \mathcal{D} , and likewise let $j(L_0, L_1)$ to be the family of almost complex structures determined by \mathcal{D} . Then $CF(L_0, L_1, \mathcal{D})$ is the vector space over \mathbb{Q} , freely

generated by elements of $S(L_0, L_1)$, where the latter is as in Definition 5.1. To quickly recall, $S(L_0, L_1)$ is the space of $\mathcal{A}(L_0, L_1)$ -flat sections γ of $P_x \times [0, 1]$, with boundary on the pair of Lagrangians

$$L_0 \subset P_x \times \{0\}, L_1 \subset P_x \times \{1\}.$$

These γ are called **geometric generators**. To relate this with more classical Floer homology generators, we point out that there is a natural set isomorphism:

$$\phi : S(L_0, L_1) \rightarrow (\mathcal{A}(L_0, L_1)L_0) \cap L_1$$

where $\mathcal{A}(L_0, L_1)L_0$ is as in the paragraph prior to the Definition 5.1. The map ϕ is given by

$$\phi(\gamma) = \gamma(1).$$

Then the \mathbb{Z}_2 grading of a generator $\gamma \in S(L_0, L_1)$ is given by the sign of the intersection point $\phi(\gamma)$.

6.1.1. *Differential on $CF(L_0, L_1, \mathcal{D})$.* For γ_0, γ_1 geometric generators of

$$CF(L_0, L_1, \mathcal{D}),$$

let $\mathcal{M}(\gamma_0; \gamma_1)$ denote the space of holomorphic (to be further explained) sections of

$$P_x \times ([0, 1] \times \mathbb{R}) \rightarrow [0, 1] \times \mathbb{R},$$

with boundary on the Lagrangian sub-bundles

$$L_0 \times \{0\} \times \mathbb{R} \rightarrow \mathbb{R}, L_1 \times \{1\} \times \mathbb{R} \rightarrow \mathbb{R},$$

and asymptotic to γ_i at the ∞ , respectively $-\infty$ ends. And let $\overline{\mathcal{M}}(\gamma_0; \gamma_1)$ denote the natural Floer compactification of the quotient $\mathcal{M}(\gamma_0; \gamma_1)/\mathbb{R}$, where \mathbb{R} acts by translation on the domain.

Terminology 6.2. *Here and elsewhere the term **holomorphic section** of various Hamiltonian fibrations over Riemann surfaces S will mean the following. Our Hamiltonian fibrations $\tilde{S} \rightarrow S$ always come with choices of a Hamiltonian connection \mathcal{A} , and a family of fiber-wise almost complex structures $\{j_z\}_{z \in S}$, determined by the perturbation data \mathcal{D} . This gives an induced almost complex structure $J(\mathcal{A}, \{j_z\})$ on \tilde{S} restricting to $\{j_z\}$ on the fibers, having a holomorphic projection map to the base, and preserving the horizontal distribution of \mathcal{A} . Holomorphic then means that the section has $J(\mathcal{A}, \{j_z\})$ -complex linear differential.*

In the above case, “holomorphic” is with respect to the almost complex structure induced by the flat, \mathbb{R} -translation invariant extension of $\mathcal{A}(L_0, L_1)$ to $P_x \times [0, 1] \times \mathbb{R}$, and likewise by the translation invariant extension of $j(L_0, L_1)$ to $P_x \times [0, 1] \times \mathbb{R}$.

For a generic pair $\mathcal{A}(L_0, L_1), j(L_0, L_1)$, all the moduli spaces $\overline{\mathcal{M}}(\gamma_0; \gamma_1)$ are transversely cut out for all γ_i , [25] but these kind of transversality results go much further back, see for example Oh [18].

The differential

$$\mu^1 : CF(L_0, L_1, \mathcal{D}) \rightarrow CF(L_0, L_1, \mathcal{D})$$

is defined as usual by

$$\mu^1(\gamma_i) = \sum_j \# \overline{\mathcal{M}}(\gamma_i; \gamma_j) \gamma_j,$$

for $\{\gamma_i\}$ a basis of geometric generators for $CF(L_0, L_1, \mathcal{D})$. Here $\# \overline{\mathcal{M}}(\gamma_i; \gamma_j)$ is defined to be zero, unless the virtual dimension of $\mathcal{M}(\gamma_i; \gamma_j)$ is zero in that case it is the signed count of points. The sum is finite by the monotonicity condition.

6.1.2. *Section classes.* Let $M \hookrightarrow \tilde{S} \rightarrow S$ be a Hamiltonian fibration over a Riemann surface with boundary and ends $\{e_i\}$, analogously to the setup in Section ???. Suppose we have fixed trivializations $\tilde{e}_i : [0, 1] \times (0, \infty) \times M \rightarrow \tilde{S}$, for each end. And suppose we have a Lagrangian sub-fibration \mathcal{L} over the components of the boundary ∂S , analogous to the sub-fibrations (5.4). Let σ be a section of \tilde{S} , so that $\sigma(\partial S) \in \mathcal{L}$. Suppose in addition that σ is continuous and is C^0 asymptotic at each end, to a section $\tilde{\sigma}$ which is translation invariant in the $[0, \infty)$ factor (in the distinguished trivialization). Here asymptotic just means C^0 convergence $\lim_{s \rightarrow \infty} \sigma|_{[0, 1] \times \{s\}} = \gamma$, for $\gamma = \tilde{\sigma}|_{[0, 1] \times s}$. ($\tilde{\sigma}$ is translation invariant so that this is well defined.) In this case, as may be apparent, we may define the homology class of σ , relative to the boundary and relative to the asymptotic constraints.

The above extends, to the case S is disconnected, of the form \mathcal{S}_r for $r \in \partial \mathcal{R}_s$. In this case we ask that our sections σ also have matching asymptotic constraints, at the corresponding nodal ends $n_{j, \pm}$, cf. Section 3.2. Given this we may again define a relative homology class of σ . We will not give exhaustive detail of this, as this is highly analogous to the definition of relative classes of curves in for example [?]. We just have a language change, instead of maps to a manifold M , we have sections of M -fibrations. Let us denote such homology classes by letters A .

6.1.3. *Higher multiplication maps.* The multiplication maps

$$(6.1) \quad \begin{aligned} \mu^d : \text{hom}(L_0, L_1) \otimes \text{hom}(L_1, L_2) \otimes \dots \\ \otimes \text{hom}(L_{d-1}, L_d) \rightarrow \text{hom}(L_0, L_d), \end{aligned}$$

$d > 1$ are defined as follows.

Notation 6.3. *In the rest of the paper we use the notation $\{\gamma_i^j\}_{i \in I^j} \in CF(L_{j-1}, L_j)$ for the basis of geometric generators. (The set I^j will usually be omitted from notation.) So the superscript in this notation refers to the vector space. Similarly, $\gamma_i^0 \in CF(L_0, L_d)$ will likewise denote the generators. If the subscript is not specified then we just mean general generator.*

Let \mathcal{U}, \mathcal{F} be the natural systems determined by \mathcal{D} . Then for generators $\gamma^j \in CF(L_{j-1}, L_j)$, $1 \leq j \leq d$, $\gamma^0 \in CF(L_0, L_d)$, we define the moduli space

$$(6.2) \quad \mathcal{M}(\{\gamma^j\}; \gamma^0, x, \mathcal{F}, A)$$

as follows. The elements are pairs (σ, r) , for σ a relative class A (to be further explained), $\mathcal{F}(\{L_i\}, x, r)$ -holomorphic section of

$$P_x \times \mathcal{S}_r \rightarrow \mathcal{S}_r,$$

and where $r \in \mathcal{R}_d$. And s.t. each pair (σ, r) satisfies:

- $\sigma(\partial \mathcal{S}_r) \subset \mathcal{L}(\mathcal{U}, L_0, \dots, L_d, r)$, see (5.4).
- By assumptions, at each e_i end of \mathcal{S}_r , $i \neq 0$, in the distinguished coordinates

$$(M \simeq P_x) \times [0, \infty) \rightarrow \tilde{\mathcal{S}}_r,$$

the data $\mathcal{F}(\{L_j\}, \Sigma, r)$ is \mathbb{R} -translation invariant, in the $[0, \infty)$ factor. Then we ask that σ be asymptotic to γ^j . Here, **asymptotic** means that

$$\lim_{s \rightarrow \infty} \sigma|_{[0, 1] \times \{s\}} = \gamma^i,$$

where the limit is C^∞ limit. Likewise, in the distinguished coordinates

$$M \times (-\infty, 0] \rightarrow \tilde{\mathcal{S}}_r,$$

at the e_0 end, we ask that σ be asymptotic to γ^0 .

- The pair of the conditions above mean that σ determines a relative homology class, as in Section 6.1.2, and we ask that all the σ are in the same class A .

Take a basis of geometric generators $\{\gamma_i^j\}_{i \in I^j}$ for $\text{hom}_{F(x)}(L_{j-1}, L_j)$, $1 \leq j \leq d$, $d \geq 2$, and a basis $\{\gamma_k^0\}$ for $\text{hom}_{F(x)}(L_0, L_d)$. Then assuming that $\mathcal{F}(\{L_j\}, x, r)$ is regular we define μ^d by:

$$(6.3) \quad \langle \mu^d(\gamma_{i_1}^1, \dots, \gamma_{i_d}^d), \gamma_{k_0}^0 \rangle = \sum_A \# \mathcal{M}(\gamma_{i_1}^1, \dots, \gamma_{i_d}^d; \gamma_{k_0}^0, x, \mathcal{F}, A),$$

when the above moduli spaces have dimension 0, for \langle, \rangle the natural inner product pairing induced by our choice of basis. The sum is finite by monotonicity.

6.1.4. *Compactification regularity, and associativity.* Given a certain dictionary, the moduli spaces

$$\mathcal{M}(\{\gamma^i\}; \gamma^0, x, \mathcal{F}, A)$$

are identical to the moduli spaces in Sheridan [25] with respect to a system of Hamiltonian perturbations, and a domain dependent family of almost complex structures on (in this case) trivial bundles, determined by \mathcal{F} .

To be more explicit, a Hamiltonian connection on a trivial M bundle over a surface S is the same as the data of a 1-form on S with values in $C_0^\infty(M)$: smooth functions with mean 0. This is the same as the data of a Hamiltonian perturbation. So in our case we just have a language change, the reason for which will be obvious when we shall construct the value of F on higher dimensional simplices of X . Consequently the compactification and regularity story is word for word identical to Sheridan [25]. (Again, given the right dictionary.)

Let us say a few more words on this. The compactification

$$\overline{\mathcal{M}}(\{\gamma^i\}; \gamma^0, x, \mathcal{F}, A),$$

is obtained by allowing $r \in \overline{\mathcal{R}}_d$ and allowing broken holomorphic sections over the disconnected surfaces \mathcal{S}_r , $r \in \partial \overline{\mathcal{R}}_d$. (This is in addition to the usual stable map compactification.) A broken holomorphic section of $\tilde{\mathcal{S}}_r \rightarrow \mathcal{S}_r$ is a holomorphic section over each smooth component of \mathcal{S}_r , so that at the node ends $n_{j,\pm}$, the corresponding sections are asymptotic to the same geometric generator γ . (In the natural bundle trivializations at the ends.)

6.1.5. *A_∞ associativity.* The maps μ^d satisfy the A_∞ -associativity equations (stated over \mathbb{F}_2 for simplicity)

$$(6.4) \quad \sum_{n,m} \mu^{d-m+1}(\gamma^1, \dots, \gamma^n, \mu^m(\gamma^{n+1}, \dots, \gamma^{m+n}), \gamma^{n+m+1}, \dots, \gamma^d) = 0,$$

This is shown as usual by considering the boundary of the one dimensional moduli spaces, of the form: $\overline{\mathcal{M}}(\{\gamma^i\}; \gamma^0, x, \mathcal{F}, A)$.

6.2. **F on higher dimensional simplices.** Let $\Sigma : \Delta^n \rightarrow X$ be smooth. The category $F(\Sigma)$ has objects $\bigsqcup_i \text{obj } F(x_i)$, where $x_i : pt \rightarrow X$ is the composition of the vertex inclusion $\Delta^0 \rightarrow \Delta^n$, corresponding to the i 'th corner, with the map Σ . Abusing notation we may also write x_i for $x_i(pt) \in X$.

Let

$$m : [0, 1] \rightarrow \Delta^n$$

be the edge between i, j corners of Δ^n and set

$$\overline{m} = \Sigma \circ m.$$

Given a pair of objects $L_0 \in F(x_i) \subset F(\Sigma)$, $L_1 \in F(x_j) \subset F(\Sigma)$, (including $i = j$) and given the Hamiltonian connection

$$\mathcal{A}(L_0, L_1) = \mathcal{A}(L_0, L_1, \overline{m})$$

on \overline{m}^*P , determined by \mathcal{D} , we define as before $\text{hom}_{F(\Sigma)}(L_0, L_1)$ to be the \mathbb{Z}_2 graded chain complex over \mathbb{Q} generated by the elements of $S(L_0, L_1, \mathcal{A}(L_0, L_1))$, cf. Definition 5.1. The grading is defined as before.

The differential μ^1 is defined identically to the differential on morphism spaces of categories $\text{Fuk}(P_x)$. The only difference is that \overline{m}^*P may no longer be naturally trivialized.

This completely describes all objects and morphisms of $F(\Sigma)$. We now need to describe the A_∞ structure. Given $\{L_{\rho(k)} \in F(x_{\rho(k)})\}_{k=0}^{k=d}$,

$$\rho : \{0, \dots, d\} \rightarrow \{0, \dots, n\},$$

with $\omega(L_{\rho(k)}) = \theta \in \mathbb{Z}$, we need to define the higher composition maps

$$(6.5) \quad \mu_\Sigma^d : \text{hom}(L_{\rho(0)}, L_{\rho(1)}) \otimes \dots \otimes \text{hom}(L_{\rho(d-1)}, L_{\rho(d)}) \rightarrow \text{hom}(L_{\rho(0)}, L_{\rho(d)}).$$

Note that by construction, to each morphism of $F(\Sigma)$ naturally corresponds either an edge or a vertex of Δ^n , in either case we may naturally associate to these a morphism in the groupoid $\Pi(\Delta^n)$. The collection $\{x_{\rho(k)}\}$ then clearly determines a composable chain (m_1, \dots, m_d) of morphisms in $\Pi(\Delta^n)$.

So let $\gamma^j \in CF(L_{\rho(j-1)}, L_{\rho(j)})$, $1 \leq j \leq d$, $\gamma^0 \in CF(L_{\rho(0)}, L_{\rho(d)})$, be geometric generators. Given the map

$$u(m_1, \dots, m_d, n) : \mathcal{S}_d^\circ \rightarrow \Delta^n \text{ that is part of a natural system } \mathcal{U},$$

and given the system \mathcal{F} determined by \mathcal{D} , we define the moduli space

$$\mathcal{M}(\{\gamma^j\}; \gamma^0, \Sigma, \mathcal{F}, A)$$

analogously to (6.2). The elements of our moduli space are pairs (σ, r) , $r \in \mathcal{R}_d$, and σ a class A (to be further explained), $\mathcal{F}(\{L_{\rho(j)}\}, \Sigma, r)$ -holomorphic section of

$$\tilde{\mathcal{S}}_r = \tilde{\mathcal{S}}(m_1, \dots, m_d, \Sigma, r) \rightarrow \mathcal{S}_r.$$

In addition each pair (σ, r) satisfies:

- $\sigma(\partial \mathcal{S}_r) \subset \mathcal{L}(\mathcal{U}, L_{\rho(0)}, \dots, L_{\rho(d)}, r)$, see (5.4). Recall that the right hand side is a sub-fibration of $\tilde{\mathcal{S}}_r$ over the boundary of \mathcal{S}_r .
- By assumptions, at the i 'th end of \mathcal{S}_r , $i \neq 0$, in the distinguished coordinates

$$\overline{m}_i^*P \times [0, \infty) \rightarrow \tilde{\mathcal{S}}_r,$$

the data $\mathcal{F}(\{L_{\rho(j)}\}, \Sigma, r)$ is \mathbb{R} -translation invariant, in the $[0, \infty)$ factor. Then we ask that σ be asymptotic to γ^j a geometric generator of

$$\text{hom}_{F(\Sigma)}(L_{\rho(j-1)}, L_{\rho(j)}),$$

where asymptotic is as in Section 6.1.3. Likewise, in the distinguished coordinates

$$\overline{m}_0^*P \times (-\infty, 0] \rightarrow \tilde{\mathcal{S}}_r,$$

we ask that σ be asymptotic to γ^0 a geometric generator of $\text{hom}_{F(\Sigma)}(L_{\rho(0)}, L_{\rho(d)})$.

- The pair of the conditions above mean that σ determines a relative homology class, as in Section 6.1.2, and we ask that all the σ are in the same class A .

6.2.1. *Compactness and regularity.* We do not need to reinvent the wheel proving compactness and regularity results for the above moduli spaces. (Although it obviously works the same way.) Instead pick a Hamiltonian trivialization of

$$M \times \Delta^n \xrightarrow{tr} \Sigma^* P,$$

then we may pullback the systems of connections $\mathcal{F}(\{L_i\}, \Sigma, r)$, and the systems of almost complex structures $\mathcal{J}(\{L_i\}, \Sigma)$ to $M \times \Delta^n$. Then in the coordinates of $M \times \Delta^n$, these systems of connections and complex structures are essentially equivalent to systems of compatible perturbations, in the sense of Seidel [24, Section 9i], and Sheridan [25] for the Lagrangian boundary conditions given by $\{\pi \circ tr^{-1}(L_i)\}$ for

$$\pi : M \times \Delta^n \rightarrow M$$

the projection. (Only “essentially” because we now allow extra copies of the same object.) Consequently compactness and regularity works the same way as described in Section 6.1.4, which is based on the work of Sheridan [25].

6.2.2. *Composition maps in the A_∞ category $F(\Sigma)$.* For $\{L_{\rho(j)}\}$, as above, given geometric generators $\gamma^j \in \text{hom}_{F(x)}(L_{\rho(j-1)}, L_{\rho(j)})$, $1 \leq j \leq d$, $d \geq 2$, and a geometric generator $\gamma^0 \in \text{hom}_{F(x)}(L_{\rho(0)}, L_{\rho(d)})$, assuming that $\mathcal{F}(\{L_{\rho(j)}\}, \Sigma, r)$ is regular we define $\mu_{F(\Sigma)}^d(\gamma^1, \dots, \gamma^d)$ by the pairing:

$$(6.6) \quad \langle \mu_{F(\Sigma)}^d(\gamma^1, \dots, \gamma^d), \gamma^0 \rangle = \sum_A \# \mathcal{M}(\gamma^1, \dots, \gamma^d; \gamma^0, \Sigma, \mathcal{F}, A),$$

when the above moduli spaces are of dimension 0, for \langle, \rangle as before the inner product pairing induced by our basis choice. Again the sum is finite by the monotonicity.

6.2.3. *Associativity.* This works as before.

Lemma 6.4. *The assignment $\Sigma \mapsto F(\Sigma)$ extends to a natural functor*

$$F : \text{Simp}(X) \rightarrow A_\infty - \text{Cat}.$$

Proof. Given a face map $f : \Delta^{n-1} \rightarrow \Delta^n$ and $\Sigma^n : \Delta^n \rightarrow X$, by the naturality property 4 of our connections there is a canonical functor $F(\Sigma^n \circ f) \rightarrow F(\Sigma^n)$ that is by construction a fully-faithful embedding. It follows via iteration that a morphism $\sigma : \Sigma^k \rightarrow \Sigma^l$, with $\Sigma^k, \Sigma^l \in \text{Simp}(X)$, $k < l$ induces a fully-faithful embedding:

$$F(\sigma) : F(\Sigma^k) \rightarrow F(\Sigma^l),$$

and this assignment is clearly functorial. Note that $F(\sigma)$ is essentially surjective on the cohomological level, which follows by a classical continuation argument, cf. [24, Section 10a], and so each $F(\sigma)$ is a quasi-equivalence. \square

Let us call the functor $F_{P, \mathcal{D}} : \text{Simp}(X) \rightarrow A_\infty - \text{Cat}$, as constructed geometrically in this section, a *geometric functor* to emphasize the origin.

6.3. Unital replacement of F . Let $A_\infty - Cat^{unit}$ denote the subcategory of $A_\infty - Cat$ consisting of strictly unital A_∞ categories and unital functors. By *unital replacement* for

$$F : Simp(X) \rightarrow A_\infty - Cat$$

we mean a functor

$$F^{unit} : Simp(X) \rightarrow A_\infty - Cat^{unit}$$

together with a natural transformation

$$N : F \rightarrow F^{unit},$$

which is object-wise quasi-equivalence.

Lemma 6.5. *Any functor $F : Simp(X) \rightarrow A_\infty - Cat$ has a unital replacement.*

Proof. To obtain this we proceed inductively: for each 0-simplex $x \in Simp(X)$, since each $F(x)$ is c-unital we may fix a formal diffeomorphism $\Phi_x : F(x) \rightarrow F(x)$, with first component maps Φ_x^1 the identity maps, such that the induced A_∞ -structure

$$F^{unit}(x) = (\Phi_x)_*(F(x))$$

is strictly unital, [24, Lemma 2.1]. Let

$$N_x : F(x) \rightarrow F^{unit}(x)$$

denote the induced A_∞ functor. Let F_k denote the restriction of F to $Simp^{\leq k}(X)$ with $Simp^{\leq k}(X)$ denoting the sub-category of $Simp(X)$, consisting of simplices whose degree is at most k . And suppose that the maps N_x can be extended to a natural transformation $N_k : F_k \rightarrow F_k^{unit}$ of functors

$$F_k : Simp^{\leq k}(X) \rightarrow A_\infty - Cat,$$

$$F_k^{unit} : Simp^{\leq k}(X) \rightarrow A_\infty - Cat^{unit},$$

$k > 0$, where for each Σ $N_k(\Sigma) : F(\Sigma) \rightarrow F^{unit}(\Sigma)$ is induced by a formal diffeomorphism $\Phi_\Sigma : F(\Sigma) \rightarrow F(\Sigma)$, whose first component maps are the identity maps.

We construct an extension N_{k+1} . For each given $\Sigma^{k+1} : \Delta^{k+1} \rightarrow X$ and $i : \Sigma^k \rightarrow \Sigma^{k+1}$, a morphism in $Simp(X)$, by assumption $F(i)$ is a fully-faithful embedding. Identifying $F(\Sigma^k)$ with a full subcategory of $F(\Sigma^{k+1})$, we may clearly construct, as in the proof of [24, Lemma 2.1], a formal diffeomorphism

$$\Phi_{\Sigma^{k+1}} : F(\Sigma^{k+1}) \rightarrow F(\Sigma^{k+1})$$

with $\Phi_{\Sigma^{k+1}}^*(\Sigma^{k+1})$ unital, and so that its restriction to $F(\Sigma^k)$ coincides with the formal diffeomorphisms $\{\Phi_{\Sigma^{k+1} \circ i}\}$, for each $i : \Sigma^k \rightarrow \Sigma^{k+1}$. The result then follows. \square

Let us write F^{unit} for the particular unital replacement of F as constructed in the proof of the lemma above.

6.4. Concordance classes of functors $F : \text{Simp}(X) \rightarrow A_\infty - \text{Cat}$. We say that a pair of functors $F_0, F_1 : \text{Simp}(X) \rightarrow A_\infty - \text{Cat}$ are *concordant* if there is a functor

$$T : \text{Simp}(X \times I) \rightarrow A_\infty - \text{Cat},$$

restricting to F_0, F_1 over $\text{Simp}(X \times \{0\})$, respectively over $\text{Simp}(X \times \{1\})$. Note that by the proof of Lemma 6.5 if F_1, F_2 are concordant then so are $F_1^{\text{unit}}, F_2^{\text{unit}}$.

Theorem 6.6. *Let $M \hookrightarrow P \rightarrow X$ be a smooth Hamiltonian fibration. For a given pair of data $\mathcal{D}_1, \mathcal{D}_2$ for P , the functors*

$$F_{P, \mathcal{D}_1} : \text{Simp}(X) \rightarrow A_\infty - \text{Cat},$$

$$F_{P, \mathcal{D}_2} : \text{Simp}(X) \rightarrow A_\infty - \text{Cat}$$

are concordant.

Proof. A given pair of choices $\mathcal{D}_1, \mathcal{D}_2$ of auxiliary data are homotopic, in the natural sense, through perturbation data $\{\mathcal{D}_t\} = (\{\mathcal{U}_t\}, \{\mathcal{F}_t\})$, $t \in I = [0, 1]$, by Theorem 4.8. Let $\tilde{P} \rightarrow X \times I$ be the Hamiltonian fibration given by pull-back of P by the projection map $X \times I \rightarrow X$.

We have partial perturbation data $\{\mathcal{D}_t\}$ for \tilde{P} , which is defined for any

$$\Sigma : \Delta^n \rightarrow X \times \{t\} \subset X \times I$$

by the data \mathcal{D}_t . Then extend the perturbation data $\{\mathcal{D}_t\}$ in any way to all $\text{Simp}(X \times I)$, call this data \mathcal{H} . The extension \mathcal{H} clearly exists once one has an extension for the natural system of maps $\{\mathcal{U}_t\}$, since there are no obstructions at all for extending $\{\mathcal{F}_t\}$. On the other hand the extension of $\{\mathcal{U}_t\}$ clearly exists by the same inductive argument as in the construction of a natural system \mathcal{U} .

Consequently we may define the functor T giving a concordance between

$$F_{P, \mathcal{D}_1}, F_{P, \mathcal{D}_2}$$

to be the geometric functor

$$F_{\tilde{P}, \mathcal{H}} : \text{Simp}(X \times I) \rightarrow A_\infty - \text{Cat}.$$

□

Remark 6.7. *Concordance relation is an equivalence relation (in the special case above). Although we will not show this here. The concordance class of the functor $F_{P, \mathcal{D}}$ is then the most fundamental invariant of the Hamiltonian fibration P that is constructed in this paper, however calculating with it may be very difficult.*

7. GLOBAL FUKAYA CATEGORY

Concordance “classes” of functors $F_{P, \mathcal{D}}$ constructed as above have a wealth of information but we will cut down on this information by assigning a more geometric object to it, via a certain colimit. We will assign to the concordance classes above equivalence classes of certain special simplicial fibrations over the singular set X_\bullet . These will be called ∞ -fibrations.

The first necessary ingredient for this story will be the notion of an ∞ -category, or a quasi-category in the specific model here. As this model is fixed in the paper we will no longer mention this. This is a simplicial set with an additional property, relaxing the notion of Kan complex. Whereas Kan complexes are fibrant objects in the Quillen model structure on the category $s\text{Set}$ of simplicial sets, ∞ -categories

are in turn the fibrant objects for a different non Quillen equivalent model structure on $sSet$ called the Joyal model structure.

For the reader's convenience we will review some of this theory of simplicial sets in the Appendix.

7.1. Global Fukaya category as a colimit. We will see in Section 8 how to enrich our construction so that our geometric functors $F_{P,\mathcal{D}}$ extend to functors

$$F_{P,\mathcal{D}} : \Delta(X) \rightarrow A_\infty - Cat^{unit},$$

in other words so that degeneracies are included. This is purely algebraic and we assume this for now.

Remark 7.1. *A naive idea for an invariant of the Hamiltonian fibration $M \hookrightarrow P \rightarrow X$ is to try to form the colimit directly:*

$$Fuk(P, \mathcal{D}) = \operatorname{colim}_{\Delta(X)} F_{P,\mathcal{D}},$$

which one may hope is an A_∞ -category. However, this has great technical difficulties. The colimit may not even exist, as in general colimits of diagrams of A_∞ categories may not exist. Our category $A_\infty - Cat^{unit}$ is a very special sub-category of all A_∞ categories, so that such co-limits may exist (this is perhaps open). But this is not good enough, as we need suitable invariance of $Fuk(P, \mathcal{D})$, say up to quasi-isomorphism, under change of \mathcal{D} , which means that in our case we need some kind of homotopy colimit, which means that our $A_\infty - Cat^{unit}$ needs to be some kind of model category. This is again a technical challenge particularly because $A_\infty - Cat^{unit}$ is so special. See however [11] where a kind of model structure is constructed on a more general category of A_∞ categories, (but with no co-limits!).

Because of technical issues as described in the above remark, we are going to use the nerve functor to land in a much more robust category of ∞ -categories, and then take the colimit. The use of the nerve functor has some perhaps unexpected benefits. We get a certain rich additional structure for our invariant object closely tied the geometry, (an ∞ -fibration structure) this is crucial for computations in Part II.

7.1.1. The A_∞ -nerve. The A_∞ -nerve is an analogue for A_∞ categories of the classical nerve functor from the category of small categories to the category of simplicial sets, (in-fact ∞ -categories). From now A_∞ -nerve will be just “nerve”: N , where there can be no confusion. This construction is due to Lurie [12, Construction 1.3.1.6], and the output is an ∞ -category. Thus, N is a functor from the category of all (strictly-unital) A_∞ categories, with morphisms A_∞ unital functors, to the category of ∞ -categories $\infty - Cat$. More precisely Lurie discusses the case of dg-categories and only indicates the case of A_∞ categories. A complete description of the nerve construction for A_∞ categories is contained in the thesis of Tanaka, [10], where it plays a central role, and is also carefully worked out in Faonte [5], where a number of properties are proved. We will reproduce it here for the reader's convenience.

It should be noted that in the Lagrangian cobordism approach to Fukaya category in Nadler-Tanaka [16] a stable ∞ -category \mathcal{Z} is constructed directly. The category \mathcal{Z} is expected to be closely related to the nerve of the triangulated envelope of the Fukaya category.

7.1.2. Outline of the (A_∞) -nerve construction. The first step in the construction of N is as follows. Let C be a strictly unital A_∞ category. The 2-skeleton of the nerve $N(C)$, has objects of C as 0-simplices, morphisms of C as 1-simplices and the 2-simplices consist of a triple of objects X, Y, Z , a triple of morphisms

$$f \in \text{hom}_C(X, Y), g \in \text{hom}_C(Y, Z), h \in \text{hom}_C(X, Z),$$

a morphism $e \in \text{hom}_C(X, Z)_1$, (subscript 1 corresponds to the degree) with $de = h - f \circ g$. We will describe the full nerve construction in the Appendix A following Tanaka [10].

7.1.3. The definition.

Definition 7.2. We define:

$$\text{Fuk}_\infty(P, \mathcal{D}) := \text{colim}_{\Delta(X)} N \circ F_{P, \mathcal{D}}^{\text{unit}} \in s\text{Set}.$$

An explicit construction of the colimit is given in Lemma 9.6. In principle the above definition could be very impractical since general objects in $s\text{Set}$ are difficult to deal with, while taking fibrant replacements for the Joyal model category structure could obfuscate all the original geometry contained in the Fukaya category. Thankfully none of this is necessary as we have a couple of miracles coming from the underlying geometry to save us. The content of this is the following theorem, to be proved in Section 9.

Theorem 7.3. As defined $\text{Fuk}_\infty(P, \mathcal{D}) \in \infty\text{-Cat}$, i.e. is a ∞ -category moreover there is a natural ∞ -fibration

$$N(\text{Fuk}(M, \omega)) \hookrightarrow \text{Fuk}_\infty(P, \mathcal{D}) \rightarrow X_\bullet,$$

whose concordance equivalence (Definition A.6) class is independent of the choice of \mathcal{D} .

7.2. Universal construction via diffeological spaces. Let $E_M \rightarrow B\text{Ham}(M, \omega)$ be the associated Hamiltonian M -bundle

$$E_M = E \times_{\text{Ham}(M, \omega)} M,$$

for E the universal principal $\text{Ham}(M, \omega)$ -bundle $E \rightarrow B\text{Ham}(M, \omega)$. $B\text{Ham}(M, \omega)$ or the classifying space of any smooth Lie group, based on Milnor's construction [15], admits a well defined notion of smooth maps into it from smooth manifolds. To be precise it has a natural diffeology, see Magnot-Watts [14] and likewise the universal M -bundle

$$E_M \rightarrow B\text{Ham}(M, \omega)$$

has a natural diffeology, so that for a diffeological map

$$f : B \rightarrow B\text{Ham}(M, \omega)$$

the pull-back bundle f^*E_M is naturally diffeological. If B is in addition a smooth dimension k manifold then f^*E_M is a diffeological space locally (diffeologically) diffeomorphic to $U \times M$, for $U \subset \mathbb{R}^k$. But the latter is clearly locally diffeomorphic to \mathbb{R}^{k+2n} , for $2n$ the dimension of M . Thus f^*E_M is a diffeological space locally diffeomorphic to \mathbb{R}^{k+2n} and hence is a smooth manifold. More formally, f^*E_M is contained in the full subcategory of the category of diffeological spaces corresponding to smooth manifolds.

In the case $B = \Delta^k$ is the k -simplex, and given an open U with $\Delta^k \subset U \subset \mathbb{R}^k$, by a smooth map $\Sigma : \Delta^k \rightarrow B\text{Ham}(M, \omega)$ we mean a map with a diffeological

extension $\tilde{\Sigma} : U \rightarrow BHam(M, \omega)$, with U given the canonical diffeology. Then we may conclude as above that $\tilde{\Sigma}^* E_M$ is naturally a smooth bundle.

So to each smooth in the sense above map $\Sigma : \Delta^k \rightarrow BHam(M, \omega)$ we have a naturally corresponding smooth bundle $\Sigma^* E_M$ over Δ^k . The construction of the previous section then works as before, associating to Σ an A_∞ category $F_{E_M, \mathcal{D}}(\Sigma)$. We may then define $BHam(M, \omega)_\bullet$ as the simplicial set, with $BHam(M, \omega)_\bullet(k)$ the set of diffeological, collared maps $\Sigma : \Delta^k \rightarrow BHam(M, \omega)$. $BHam(M, \omega)_\bullet$ is readily seen to be a Kan complex.

To summarize we have the following:

Proposition 7.4. *There is a natural functor*

$$F_{E_M, \mathcal{D}} : Simp(BHam(M, \omega)) \rightarrow A_\infty - Cat,$$

for $Simp(BHam(M, \omega))$ the category of smooth (diffeological simplices) as above.

The proof is omitted since this is just a minor extension of the construction of the previous section.

7.3. Universal construction via smooth simplicial sets. A more abstract and technically simpler approach to the above can be extracted from [21]. There an abstract simplicial set $BG_\bullet^\mathcal{U}$ ² is constructed, for any Frechet Lie group and each choice of a particular Grothendieck universe \mathcal{U} . The simplicial set $BG_\bullet^\mathcal{U}$ has the property that for every k -simplex $\Sigma \in BG_\bullet^\mathcal{U}(k)$, there is a canonically associated smooth G -fibration $P_\Sigma \rightarrow \Delta^k$. Using this, we immediately obtain a functor using our construction:

$$(7.1) \quad F_{E_M, \mathcal{D}} : Simp(BHam(M, \omega)_\bullet^\mathcal{U}) \rightarrow A_\infty - Cat,$$

where $Simp(BHam(M, \omega)_\bullet^\mathcal{U})$ denotes the simplex category of the simplicial set $BHam(M, \omega)_\bullet^\mathcal{U}$, cf. Section 3.1. Now, as follows by [21, Theorem 7.5],

$$|BHam(M, \omega)_\bullet^\mathcal{U}| \simeq BHam(M, \omega)$$

for $|\cdot|$ the geometric realization, and \simeq homotopy equivalence. In particular, to prove Theorems 1.1, 1.4 we may also start with (7.1). See the proof below. One advantage of this simplicial approach is that the connection with homotopy groups becomes elementary.

Proof of Theorems 1.1, 1.4. Given a smooth Hamiltonian fibration $M \hookrightarrow P \rightarrow X$, by Theorem 7.3 we obtain a well defined concordance class of an ∞ -fibration $Fuk_\infty(P) \rightarrow X_\bullet$. By Theorem A.9 this is classified by a homotopy class of a map:

$$cl_P : X_\bullet \rightarrow (\mathbb{S}, NFuk(M, \omega)).$$

Likewise, starting with

$$F_{P, \mathcal{D}} : Simp(BHam(M, \omega)) \rightarrow A_\infty - Cat,$$

by the previous constructions, we obtain an associated functor

$$F_{P, \mathcal{D}}^{unit} : \Delta(BHam(M, \omega)) \rightarrow A_\infty - Cat^{unit}.$$

Hence by Theorem 7.3 we obtain an ∞ -fibration

$$(7.2) \quad N(Fuk(M, \omega)) \hookrightarrow Fuk_\infty(E_M, \mathcal{D}) \rightarrow BHam(M, \omega)_\bullet.$$

²The notation in [21] omits \bullet subscript.

By Theorem A.9, there is then a uniquely determined (simplicial) homotopy class of the “classifying” simplicial map

$$cl = cl(Fuk_\infty(E_M)) : BHam(M, \omega)_\bullet \rightarrow (\mathbb{S}, NFuk(M)),$$

of the ∞ -fibration (7.2). Then we obtain a group homomorphism of simplicial homotopy groups

$$cl_* : \pi_i(BHam(M, \omega)_\bullet, x_0) \rightarrow \pi_i(\mathbb{S}, NFuk(M)).$$

If we knew that $BHam(M, \omega)_\bullet$ is weakly equivalent to the usual continuous singular set of $BHam(M, \omega)$, then we would obtain a group homomorphism $cl_* : \pi_i(BHam(M, \omega), *) \rightarrow \pi_i(|\mathbb{S}|, NFuk(M))$. This is probably true, but I don’t know if a ready reference exists. Alternatively, we can use the map

$$cl : BHam(M, \omega)_\bullet^{\mathcal{U}} \rightarrow (\mathbb{S}, NFuk(M)),$$

induced by (7.1). Since $|BHam(M, \omega)_\bullet^{\mathcal{U}}| \simeq BHam(M, \omega)$ we immediately obtain the homomorphism $cl_* : \pi_i(BHam(M, \omega), *) \rightarrow \pi_i(|\mathbb{S}|, NFuk(M))$. And this proves Theorem 1.4.

Now if $M \hookrightarrow P \rightarrow X$ is a smooth Hamiltonian fibration then $P \simeq f_P^* E_M$ for some diffeological smooth map $f_P : X \rightarrow BHam(M, \omega)$. Then by Theorem 9.9

$$Fuk_\infty(P) = f_{P,\bullet}^* Fuk_\infty(E_M),$$

with $f_{P,\bullet} : X_\bullet \rightarrow BHam(M, \omega)_\bullet$ denoting the induced simplicial map. In particular $Fuk_\infty(P)$ is classified as ∞ -fibration by the map $cl \circ f_P$. And so by Theorem A.9 $cl_P \simeq cl \circ f_P$. And so we have proved Theorem 1.1. \square

7.4. Global Fukaya category and unital replacement, a remark. In the construction of $Fuk_\infty(P)$ we had to take a unital replacement for the functor $F : \Delta/X_\bullet \rightarrow A_\infty - Cat$. One may worry then that this algebraic step will obfuscate the “geometry” of simplices of $Fuk_\infty(P)$. This is not really the case. First the A_∞ nerve NC of a non-unital A_∞ category C still exists as a semi-simplicial set, that is as a co-functor $\Delta^{inj} \rightarrow Set$, with Δ^{inj} the subcategory of Δ consisting of injective morphisms. For a unital replacement equivalence $C \rightarrow C^{unit}$ of C , constructed as in Section 6.3, we then have an induced morphism of semi-simplicial sets $NC \rightarrow NC^{unit}$, which by construction induces a bijection $NC([n]) \rightarrow NC^{unit}([n])$, for each $[n]$. So we may think without loss of geometric information, of simplices of NC^{unit} in terms of simplices of NC . (The former just have an extra formal algebraic structure.)

8. EXTENDING F TO DEGENERACIES

We have to construct our perturbation data \mathcal{D} for all simplexes in such a way that there is a natural functor:

$$(8.1) \quad F : \Delta(X) \rightarrow A_\infty - Cat^{unit},$$

extending the geometric functor

$$F_{\mathcal{D}} : Simp(X) \rightarrow A_\infty - Cat^{unit},$$

for this data \mathcal{D} as previously constructed. This perturbation data will be referred to as *extended perturbation data*.

Suppose that we are given a commutative diagram:

$$\begin{array}{ccccc} \Delta^0 & \xrightarrow{j+1} & \Delta^{n+1} & \xrightarrow{pr} & \Delta^n \\ & \searrow j_* & \downarrow \tilde{\Sigma} & \swarrow \Sigma & \\ & & X, & & \end{array}$$

where

$$pr : \Delta^{n+1} \rightarrow \Delta^n, \quad j \in [n]$$

is induced by the unique surjection $[n+1] \rightarrow [n]$ taking j and $j+1$ to j . Here $j_* = \Sigma \circ j$, and j also denotes the map $pt \rightarrow \Delta^n$ whose image is the vertex j .

Let \mathcal{D}_Σ be some chosen perturbation data for Σ . Then we construct certain induced perturbation data $\mathcal{D}_{\tilde{\Sigma}} = pr^* \mathcal{D}_\Sigma$. And using this we construct an A_∞ category $F(\tilde{\Sigma})$.

We may already define the objects of $F(\tilde{\Sigma})$ as before. So pr clearly induces a map of sets of objects

$$pr_* : \text{obj } F(\tilde{\Sigma}) \rightarrow \text{obj } F(\Sigma).$$

We need to specify our system of connections and almost complex structures. We say what to do with connections, the case of almost complex structures is analogous. Given vertices i, j of Δ^{n+1} and objects $L \in F(x_i), L' \in F(x_j)$ for $x_i = \tilde{\Sigma}(i), x_j = \tilde{\Sigma}(j)$ we set

$$\mathcal{A}(L, L') = \mathcal{A}(pr_* L, pr_* L'),$$

where the latter connection is determined by the given \mathcal{D}_Σ , and where the equality is with respect to the natural identification

$$(\Sigma \circ pr \circ m_{i,j})^* P = (\tilde{\Sigma} \circ m_{i,j})^* P.$$

Likewise given objects $L_0, \dots, L_s \in F(\tilde{\Sigma})$ we set

$$\mathcal{F}(L_0, \dots, L_s, \tilde{\Sigma}, r) = \mathcal{F}(pr_* L_0, \dots, pr_* L_s, \Sigma, r).$$

Where the equality is again with respect to the natural identification of the corresponding bundles. All together this determines a certain data $\mathcal{D}_{\tilde{\Sigma}}$.

Using this data $\mathcal{D}_{\tilde{\Sigma}}$ we then define an A_∞ category denoted by $F(\tilde{\Sigma})$ as previously. By construction, the natural map on objects:

$$pr_* : \text{obj } F(\tilde{\Sigma}) \rightarrow \text{obj } F(\Sigma)$$

extends to a strict A_∞ functor

$$F(pr) : F(\tilde{\Sigma}) \rightarrow F(\Sigma)$$

satisfying:

$$F(pr) \circ F(\sigma) = id,$$

where $\sigma : \Sigma \rightarrow \tilde{\Sigma}$ is induced by the map $d^{j+1} : [n] \rightarrow [n+1]$, which is the unique morphism in Δ whose image misses the vertex $j+1$. We are going to call the above the **extension construction for $\tilde{\Sigma}, \mathcal{D}_\Sigma$** with the corresponding extended data denoted by $\mathcal{D}_{\tilde{\Sigma}}$.

We are now going to proceed by induction. Suppose that we have constructed our data \mathcal{D} for all simplices up to degree N , and an extension of $F|_{\text{Simp}^N(X)}$ to a functor

$$F^N : \Delta(X)^N \rightarrow A_\infty - \text{Cat}^{unit},$$

where $\Delta(X)^N$ and $\text{Simp}^N(X)$ are the subcategories of simplices of degree at most N . Let Σ' be a general $(N+1)$ -simplex of X_\bullet . If Σ' is degenerate, so that $\Sigma' = \Sigma \circ pr$ for some degeneracy morphism pr , for some $\Sigma \in \text{Simp}^N(X)$, then define the data $\mathcal{D}_{\Sigma'}$ and $F^{N+1}(\Sigma')$ as in the extension construction above for $(\tilde{\Sigma} = \Sigma')$, \mathcal{D}_Σ .

Otherwise, if Σ is a non-degenerate $(N+1)$ -simplex then its faces are N -simplices for which we already have perturbation data \mathcal{D} , which is then extended arbitrarily to perturbation data \mathcal{D}_Σ for Σ . The extension is obtained as in the proof of Lemma 5.8. Using this data define $F^{N+1}(\Sigma)$ as normal. Then clearly F^{N+1} extends F^N , which completes the induction step. So that we obtain our extension (8.1).

9. ALGEBRAIC-TOPOLOGICAL CONSIDERATIONS

In this section by equivalence of ∞ -categories we always mean categorical equivalence. This and other categorical preliminaries needed for this section are discussed in the Appendix A. We will prove here Theorem 7.3.

9.1. Colimit of F .

Definition 9.1. A functor $F : \Delta(X) \rightarrow A_\infty - \text{Cat}^{\text{unit}}$ which is induced by a geometric functor $F_{\mathcal{D}} : \text{Simp}(X) \rightarrow A_\infty - \text{Cat}^{\text{unit}}$ as in Section ?? will be called **geometric**.

Remark 9.2. Instead of the above definition it would be more ideal to extract suitable algebraic axioms for our “geometric functors”. However, this may take us too far afield.

Given a geometric functor $F : \Delta(X) \rightarrow A_\infty - \text{Cat}^{\text{unit}}$, let

$$(9.1) \quad \text{Fuk}_\infty(F) := \text{colim}_{\Delta(X)} NF.$$

The category of simplicial sets is well known to be (co)-complete so that the limit certainly exists as a simplicial set. However, we shall show that this limit has additional structure, the content of which is the following proposition.

Remark 9.3. It is possible that the proposition below can be obtained as a consequence of more general principles, using general theory of colimits of ∞ -categories. However, I suspect that for a general functor of the form $G : \Delta(X) \rightarrow \infty - \text{Cat}$, we must first take a fibrant replacement of the functor G (for the Joyal model structure), if we want similar structure on $\text{colim}_{\Delta(X)} G$. (I am not however sure this is sufficient.)

Proposition 9.4. There is a natural projection of simplicial sets

$$p : \text{Fuk}_\infty(F) \rightarrow X_\bullet,$$

and this is an ∞ -fibration.

Proof. Recall that a given $\Sigma : \Delta^n \rightarrow X$ could equally be thought of as an element of $X_\bullet([n])$ or as a natural transformation $\Delta_\bullet^n \rightarrow X_\bullet$. Let us first give a more easily conceptualized presentation of $\text{Fuk}_\infty(F)$. We should say that we are just simplifying the standard, “level wise” construction, of colimits of simplicial sets, in our specific context, to keep the geometry as transparent as possible. This will be important for the computation in Part II [20].

Define a partial order $<$ on the set of pairs (f, Σ) , $f \in NF(\Sigma)(k)$ a k -simplex, $k \geq 0$, $\Sigma \in \Delta(X)$ as follows.

$$(f, \Sigma) < (f', \Sigma')$$

if there is a morphism

$$\sigma : \Sigma \rightarrow \Sigma'$$

in $\Delta(X)$ induced by injective $[n] \rightarrow [m]$ with $n \leq m$, i.e. a *face morphism*, s.t.

$$NF(\sigma)(f) = f'.$$

Clearly for every (f, Σ) there is a unique least pair

$$(f_{\min}, \Sigma_{\min}) < (f, \Sigma).$$

Note that if f is an k -simplex then Σ_{\min} is not necessarily a k -simplex. However, once we impose the following equivalence relation, we get something similar, see Lemma 9.5 below.

Let \tilde{C} be the set of minimal pairs. Define an equivalence relation on \tilde{C} first by defining

$$(f, \Sigma) \sim (f', \Sigma')$$

if there exists a degeneracy morphism $d : \Sigma \rightarrow \Sigma'$ induced by $[m] \rightarrow [n]$ with $m > n$, such that

$$NF(d)(f) = f'.$$

And then by imposing symmetry and transitivity. Denote the equivalence class of (f, Σ) by $[f, \Sigma]$.

The following is not formally necessary, but it might be helpful for visualization.

Lemma 9.5. *Each class $[f, \Sigma]$ has a unique natural representative (f_c, Σ_c) so that if f is a k -simplex then Σ_c is a k -simplex.*

Proof. Let (f, Σ) as above be given, with $\deg(f) = k$. And let $(f_{\min}, \Sigma_{\min}) \in \tilde{C}$ be as above. Then $\deg(\Sigma_{\min}) \leq (n = \deg(f_{\min}))$. Suppose that $\deg(\Sigma_{\min}) < n$. Then by the extension construction of Section 8 there there is a degeneracy

$$d_c : \Sigma_c \rightarrow \Sigma_{\min},$$

with $\deg(\Sigma_c) = k$ together with a k -simplex $f_c \in NF(\Sigma_c)$ so that $NF(d_c)(f_c) = f_{\min}$. Moreover (d_c, f_c) are uniquely determined, (by the extension construction). So that we set $\Sigma_c = \Sigma_{\min} \circ d_c$. Also note that Σ_c is just $p_{\Sigma}(f_{\min})$, with p_{Σ} as in (9.2) ahead. \square

Continuing with the proof of the proposition, we then define $C = \tilde{C} / \sim$. This is naturally a simplicial set, with

$$C([k]) = \{[f, \Sigma] \in C \mid f \in NF(\Sigma)([k])\}.$$

For example $C([0])$ is naturally isomorphic to

$$\sqcup_{x \in X} \text{Obj } F(x).$$

The following in particular will give a direct proof that the colimit (9.1) exists.

Lemma 9.6. *$C = \text{colim}_{\Delta(X)} NF$, with equality meaning natural isomorphism.*

Proof. Note first that C is a co-cone on the diagram NF . Indeed, for each Σ define $\phi_{\Sigma} : NF(\Sigma) \rightarrow C$ by

$$\phi_{\Sigma}(f) = [f_{\min}, \Sigma_{\min}].$$

It is easy to see that for a face morphism $i : \Sigma \rightarrow \Sigma'$ we have that the composition

$$NF(\Sigma) \xrightarrow{NF(i)} NF(\Sigma') \xrightarrow{\phi_{\Sigma'}} C,$$

coincides with ϕ_Σ . Likewise for a degeneracy morphism $d : \Sigma \rightarrow \Sigma'$ we have that the composition

$$NF(\Sigma) \xrightarrow{NF(d)} NF(\Sigma') \xrightarrow{\phi_{\Sigma'}} C,$$

coincides with ϕ_Σ , because of the equivalence relation \sim .

The universal property is also easy to verify, for given another co-cone C' with maps $\rho_\Sigma : NF(\Sigma) \rightarrow C'$, $\Sigma \in \Delta(X)$ we can naturally define $U : C \rightarrow C'$ by

$$U([f, \Sigma]) = \rho_\Sigma(f).$$

Then U is clearly well defined, by C' being a co-cone. And moreover, U is a map of co-cones (all the relevant diagrams commute). Since for a given $f \in NF(\Sigma)$, we have

$$U(\phi_\Sigma(f)) = \rho_{\Sigma_{\min}}(f_{\min}) = \rho_\Sigma(f),$$

where the last equality holds since $(C', \{\rho_\Sigma\})$ is a co-cone, and since by construction there is a morphism

$$i : NF(\Sigma_{\min}) \rightarrow NF(\Sigma),$$

with $F(i)f_{\min} = f$. □

Continuing with the proof of the proposition, for each n -simplex $\Sigma \in \Delta(X)$, we have a natural simplicial map

$$(9.2) \quad p_\Sigma : NF(\Sigma) \rightarrow \Sigma_*(\Delta_\bullet^n) \subset X_\bullet,$$

defined as follows. On the vertices of $NF(\Sigma)$, p_Σ is just the obvious projection. Now a k -simplex f in $NF(\Sigma)$ by definition determines a composable chain (f_1, \dots, f_k) in $NF(\Sigma)$, and hence determines a sequence of vertices $e_0, \dots, e_k \in NF(\Sigma)$ s.t. the source/target of f_i is e_{i-1} respectively e_i . This in turn determines a sequence of vertices $\{p_\Sigma(e_i)\} \subset \Sigma_*(\Delta_\bullet^n)$, and we set $p_\Sigma(f)$ to be the unique (possibly degenerate) k -simplex of $\Sigma_*(\Delta_\bullet^n)$ with these vertices. We will omit the verification that p_Σ is simplicial.

The simplicial projection

$$p : C \rightarrow X_\bullet$$

is then: send $[f, \Sigma]$ to $p_\Sigma(f)$, which is readily seen to be well defined.

It is immediate from the definition of inner fibration in Section A.2 that p is an inner-fibration if and only if the pre-image of every simplex $\Sigma : \Delta_\bullet^n \rightarrow X_\bullet$ by p is a ∞ -category, where “pre-image” $p^{-1}(\Sigma)$ is the pre-image by p of the simplicial subset $\Sigma(\Delta_\bullet^n)$. In our case this follows by construction as the preimage of Σ is clearly identified with $NF(\Sigma)$, which is an ∞ -category by properties of N .

We now verify that in addition p is a ∞ -fibration. By the Definition A.5 we need to show that for every equivalence $m : a \rightarrow b$ in $\mathcal{X} = X_\bullet$ and for every object $a' \in \text{Fuk}_\infty(P)$ with $p(a') = a$, there exists an equivalence $\widehat{m} : a' \rightarrow b'$ in $\mathcal{E} = \text{Fuk}_\infty(P)$ with $p(\widehat{m}) = m$.

Lemma 9.7. *The functor $N : A_\infty - \text{Cat}^{\text{unit}} \rightarrow \infty - \text{Cat}$, takes quasi-equivalences to weak equivalences in the Joyal model structure, i.e. categorical equivalences.*

Proof. The proof of this is contained in the proof of Proposition 1.3.1.20, Lurie [12]. We can also prove this directly by first recalling that quasi-equivalences of A_∞ -categories A, B are invertible (when working over a field with characteristic 0), up to homotopy, and then via the nerve construction translate this to a categorical equivalence of $N(A), N(B)$. □

Recall that the morphisms of $A_\infty - \text{Cat}$ are in particular quasi-equivalences, also recall Lemma 6.4. Since the inclusions $F(x_i) \rightarrow F(m)$ are quasi-equivalences, it follows by the lemma above, and by the construction of L that the inclusions of L_i into L_m are categorical equivalences of ∞ -categories, and so \tilde{m} as above must exist. \square

Proof of Theorem 7.3. By the discussion above we have an ∞ -fibration

$$\text{Fuk}_\infty(P, \mathcal{D}) \rightarrow X_\bullet.$$

The first part of the theorem follows by the following general fact: for an inner fibration of simplicial sets

$$p : P_\bullet \rightarrow X_\bullet,$$

if X_\bullet is a ∞ -category then P_\bullet is a ∞ -category. Let us prove this elementary point. Suppose we are given

$$\rho : \Delta_k^n \rightarrow P_\bullet$$

for $0 < k < n$. As X_\bullet is a ∞ -category there a simplex

$$\tilde{\rho} : \Delta_\bullet^n \rightarrow X_\bullet$$

extending $p \circ \rho$. But then ρ maps into the ∞ -category $p^{-1}(\tilde{\rho})$, and consequently there is an extension of ρ , c.f. Proposition A.4.

The final part of the theorem follows by the following.

Lemma 9.8. *For the geometric functor $F_{P, \mathcal{D}}$ the concordance class of the ∞ -fibration $p : \text{Fuk}_\infty(P, \mathcal{D}) \rightarrow X_\bullet$ is independent of the choice of \mathcal{D} .*

Proof. By Theorem 6.6 given a pair $\mathcal{D}_0, \mathcal{D}_1$ of perturbation data for P , there is a geometric functor:

$$\tilde{F} : \Delta(X \times I) \rightarrow A_\infty - \text{Cat}^{\text{unit}},$$

which gives a concordance of the functors

$$F_{P, \mathcal{D}_i} : \Delta(X) \rightarrow A_\infty - \text{Cat}^{\text{unit}}.$$

Then by the Proposition 9.4 there exists an ∞ -fibration:

$$\mathcal{T} \rightarrow X_\bullet \times I_\bullet,$$

whose restriction over $X_\bullet \times \partial I_\bullet$ coincides with

$$\text{Fuk}_\infty(P, \mathcal{D}_0) \sqcup \text{Fuk}_\infty(P, \mathcal{D}_1).$$

\square

\square

9.2. Naturality. We may expect if our constructions are really natural that

$$\text{Fuk}_\infty(P, \mathcal{D})$$

is functorial with respect to pull-back and this is indeed the case. Let $f : X \rightarrow Y$ be a smooth map, $P \rightarrow Y$ a smooth Hamiltonian fibration and $\mathcal{D} = \mathcal{D}(P)$ extended perturbation data. We may then naturally define pull-back extended perturbation data $f^*\mathcal{D}$ for $f^*P \rightarrow X$. Specifically, if $\Sigma \in X_\bullet$ then we set

$$\mathcal{F}(L_0, \dots, L_s, \Sigma, r) = \mathcal{F}(L_0, \dots, L_s, \tilde{\Sigma}, r),$$

for $\tilde{\Sigma} = f \circ \Sigma$.

Let $f_\bullet : X_\bullet \rightarrow Y_\bullet$ be the induced map of simplicial sets.

Theorem 9.9.

$$Fuk_{\infty}(f^*P, f^*\mathcal{D}) = f_{\bullet}^*Fuk_{\infty}(P, \mathcal{D}),$$

where $f^*\mathcal{D}$ is as above, and where $f_{\bullet}^*Fuk_{\infty}(P, \mathcal{D})$ denotes the standard pull-back of the simplicial fibration by f_{\bullet} .

Proof. The proof is immediate. \square

APPENDIX A. ∞ -CATEGORIES AND JOYAL MODEL STRUCTURE

We don't need absolutely everything in this section, particularly we can avoid ever mentioning model categories, but the latter helps with the narrative. A very good concise reference for much of this material is Riehl [19], which we will mostly follow. The material on co-Cartesian fibrations is taken from Lurie [13, Section 2.4]. First let us recall the notion of a Kan complex, which may be thought of as formalizing the property of a simplicial set to be like the singular set of a topological space, defined in Section 3.1.

Let Δ^n be the standard representable n -simplex: $\Delta([i]) = \Delta([i], [n])$. Previously we denoted this by Δ_{\bullet}^n , but as there are no topological simplices in this section we simplify the notation, which is also consistent with above references. Let $\Lambda_k^n \subset \Delta^n$ denote the sub-simplicial set corresponding to the “boundary” of Δ^n with the k 'th face removed, $0 \leq k \leq n$. By k 'th face we mean the face opposite to k 'th vertex. This is called a **horn**.

A simplicial set S_{\bullet} is said to be a *Kan complex* if for all n, k given a diagram with solid arrows

$$\begin{array}{ccc} \Lambda_k^n & \longrightarrow & S_{\bullet} \\ \downarrow & \nearrow & \\ \Delta^n & & \end{array},$$

there is a dotted arrow making the diagram commute.

An ∞ -category is a simplicial set S_{\bullet} for which the above extension property is only required to hold for inner horns Λ_k^n , i.e. those horns with $0 < k < n$. A Kan complex is a simplicial model of an ∞ -groupoid, as the Kan condition can be interpreted as giving the condition that morphisms are invertible up to (coherent) homotopy. Likewise the defining property of an ∞ -category tells us that while 1-morphisms/edges may not be invertible (up to homotopy), the higher morphisms are. It would perhaps take us too far afield too further motivate ∞ -categories here. However, the introductory sections of Lurie [13] should be highly accessible.

A morphism between ∞ -categories is just a simplicial map. We will denote ∞ -categories by calligraphic letters e.g. \mathcal{B} . In [13, Chapter 3] an ∞ -category of ∞ -categories is constructed, with 1-morphisms simplicial maps, and we call this Cat_{∞} . On the other hand the full-subcategory of the category of simplicial sets with objects ∞ -categories will be denoted by $\infty-Cat$.

Notation A.1. We denote the maximal Kan subcomplex of Cat_{∞} by S .

A.1. Categorical equivalences, morphisms and equivalences. We have a natural functor $\tau : sSet \rightarrow Cat$, defined as follows. $\tau(S_{\bullet})$ is the category with objects 0-simplices of S_{\bullet} , 1-simplices as morphisms, degenerate 1-simplices as identities and freely generated composition subject to the relation $g = f \circ h$ if there is a

2-simplex e with 0-face h , 2-face f and 1-face g . (Remembering our diagrammatic order for composition.) See the figure below.




FIGURE 8.

We have a functor $\tau_0 : sSet \rightarrow Set$ by sending A_\bullet to the set of isomorphism classes of objects in $\tau(A_\bullet)$.

If $S_\bullet = \mathcal{X}$ is a ∞ -category an edge $e \in \mathcal{X}$, i.e. a 1-simplex $e : \Delta^1 \rightarrow \mathcal{X}$, is said to be an *equivalence* if $\tau(e)$ is an isomorphism in $\tau(X)$. We may use morphism notation, so that $e : a \rightarrow b$ signifies that the edge e goes from the vertex a to b .

The *maximal Kan subcomplex* of a ∞ -category \mathcal{X} is the maximal sub-simplicial set $K(\mathcal{X}) \subset \mathcal{X}$ with all edges equivalences. (It can be constructed simply by removing edges that are not equivalences, and all simplices containing them.) The fact that $K(\mathcal{X})$ is forced to be a Kan complex can be readily verified using the ∞ -category structure of \mathcal{X} . (This is an instructive exercise.)

Definition A.2. Let \mathcal{X} be a Kan complex and $a \in \mathcal{X}(0)$. A **connected component** of a is the set of vertices sharing an edge with a . We may sometimes denote such a connected component by (\mathcal{X}, a) .

We define $sSet^{\tau_0}$ to be the category with the same objects as $sSet$ but with the morphisms given by $sSet^{\tau_0}(A_\bullet, B_\bullet) = \tau^0(B_\bullet^{A_\bullet})$. A map of simplicial sets

$$u : A_\bullet \rightarrow B_\bullet$$

is said to be a *categorical equivalence* if the induced map in $sSet^{\tau_0}$ is an equivalence. It is said to be a *weak categorical equivalence* if the pull-back map

$$sSet^{\tau_0}(B_\bullet, X_\bullet) \rightarrow sSet^{\tau_0}(A_\bullet, X_\bullet)$$

induced by u is an equivalence for all X_\bullet . A categorical equivalence is necessarily a weak categorical equivalence. Conversely a weak categorical equivalence between ∞ -categories is necessarily a categorical equivalence, this is a consequence of ∞ -categories being fibrant objects for the Joyal model structure, see Theorem A.3 below. (Technically, we will only need categorical equivalences.)

We will say that a pair of ∞ -categories are *categorically equivalent* if there is categorical equivalence between them.

As we are following Riehl [19], we refer the reader there for the following:

Theorem A.3. [Joyal, Lurie, Riehl] *There is a model structure on $sSet$, called the Joyal model structure, with weak equivalences weak categorical equivalences and cofibrations monomorphisms. Moreover the fibrant objects are ∞ -categories.*

We do not formally need the above theorem, but we hope it places things into some perspective, by making ∞ -categories a less mysterious object.

A.2. Inner fibrations. A map $p : \mathcal{A} \rightarrow \mathcal{B}$ of ∞ -categories is said to be an *inner fibration* if it has the lifting property with respect to all inner horn inclusions. More specifically if for $0 < k < n$ whenever we are given a commutative diagram with solid arrows:

$$(A.1) \quad \begin{array}{ccc} \Lambda_k^n & \longrightarrow & \mathcal{A} \\ \downarrow & \nearrow & \downarrow p \\ \Delta^n & \longrightarrow & \mathcal{B}, \end{array}$$

there exists a dashed arrow as indicated, making the whole diagram commutative. For reference p is said to be a *Kan fibration* if the above extension property holds for all horns. A Kan fibration is an analogue in the simplicial world of Serre fibrations of topological spaces. The following is immediate from definitions.

Proposition A.4. *A map $p : \mathcal{A} \rightarrow \mathcal{B}$ is an inner fibration, if and only if the pre-image of every simplex of \mathcal{B} is a ∞ -category.*

A.3. Categorical fibrations. These are the analogues of Serre fibrations for the Joyal model structure.

Definition A.5. *We say that $p : \mathcal{E} \rightarrow \mathcal{X}$ is a **categorical fibration** if:*

- (1) *The map p is an inner fibration.*
- (2) *For every equivalence $f : a \rightarrow b$ in \mathcal{X} and every object $a' \in \mathcal{E}$ with $p(a') = a$, there exists an equivalence $\tilde{f} : a' \rightarrow b'$ in \mathcal{E} with $p(\tilde{f}) = f$.*

A categorical fibration over a Kan complex will be called an ∞ -**fibration**.

Definition A.6. *We say that a pair of ∞ -fibrations $p_i : \mathcal{P}_i \rightarrow \mathcal{X}$, $i = 0, 1$ over a Kan complex \mathcal{X} are **concordant** if the following holds. There is a ∞ -fibration*

$$\mathcal{Y} \rightarrow \mathcal{X} \times \Delta_\bullet^1,$$

whose pull-back by $i_0 : \mathcal{X} \rightarrow \mathcal{X} \times \Delta_\bullet^1$ is identified with \mathcal{P}_0 and the pull-back by $i_1 : \mathcal{X} \rightarrow \mathcal{X} \times \Delta_\bullet^1$ is identified with \mathcal{P}_1 . Here the two maps i_0, i_1 correspond to the two vertex inclusions $\Delta_\bullet^0 \rightarrow \Delta_\bullet^1$.

Notation A.7. *We shall denote the set of concordance classes of ∞ -fibrations over a Kan complex \mathcal{X} by $\text{Fib}_\infty(\mathcal{X})$.*

For convenience we recall the basic definition.

Definition A.8. *We say that a pair of maps of simplicial sets $f, g : A_\bullet \rightarrow B_\bullet$ are **homotopic** if there is map of simplicial sets*

$$F : A_\bullet \times \Delta_\bullet^1 \rightarrow B_\bullet,$$

so that $F|_{A_\bullet \times \{0\}} = f$ and $F|_{A_\bullet \times \{1\}} = g$.

In [13, Section 3.3.2] Lurie constructs a universal ∞ -fibration over \mathbb{S} . (It is the restriction over \mathbb{S} of the universal Cartesian fibration over Cat_∞ , constructed in [13, Section 3.3.2].) As a direct consequence we have the following theorem. The statement should be interpreted with care however, since there are set theoretic issues. The most natural (and arguably standard) interpretation is via Grothendieck universes, as for example done explicitly in [21], in a very similar context.

Theorem A.9 (Lurie [13]). *For a Kan complex \mathcal{X} , there is a natural isomorphism*

$$\mathrm{Fib}_\infty(\mathcal{X}) \simeq [\mathcal{X}, \mathbb{S}],$$

with $[\mathcal{X}, \mathbb{S}]$ denoting the “set” of homotopy classes of maps $\mathcal{X} \rightarrow \mathbb{S}$.

A.4. A_∞ -nerve. This section mostly follows Tanaka [10, 2.3], except that for us everything will be ungraded, and for simplicity with \mathbb{F}_2 -coefficients.

For $[n] \in \Delta$, a *length s wedge decomposition* of $[n]$ is a collection of monomorphisms in Δ

$$j_i : [n_i] \rightarrow [n], \quad i = 1, \dots, s,$$

satisfying the following properties:

- $\forall i : n_i \geq 1$.
- $1 \in \mathrm{image}(j_1), n \in \mathrm{image}(j_s)$.
- $\forall 2 \leq i \leq n : \max_{[n_{i-1}]} j_{i-1} = \min_{[n_i]} j_i$

We denote the set of all decompositions of $[n]$ by $D[n]$. We may of course equally understand a length s decomposition as a finite set $\{J_1, \dots, J_s\}$ of subsets of $[n]$ decomposing $[n]$. However it is a bit simpler to formulate the following in terms of the maps j_i .

Definition A.10. *For A a small unital A_∞ category its nerve $N(A)$ is a simplicial set with the set of vertices the set of objects of A . A n -simplex f of $N(A)$ consists of the following data:*

- A map $[n] \rightarrow \mathrm{Objects}(A)$. We denote the corresponding objects X_0, \dots, X_n .
- For each mono-morphism $j : [n_j] \rightarrow [n]$ in Δ , with $n_j \geq 1$, an element

$$f_j \in \mathrm{hom}_A(X_{j(0)}, X_{j(n_j)}).$$

We may completely characterize each such j by its image set, and will sometimes write j for the corresponding set and vice versa, thus $f_{[n]}$ corresponds to the identity $j : [n] \rightarrow [n]$.

- For a given $j : [n_j] \rightarrow [n]$, and $i \in [n_j]$ denote by $j - j(i) : [n_j - 1] \rightarrow [n]$ the unique morphism in Δ with image set $j - j(i)$. Then the collection of these f_j is required to satisfy the following equation:

$$(A.2) \quad \mu^1(f_j) = \sum_{0 < i < n_j} f_{j-j(i)} + \sum_{s \geq 2} \sum_{\mathrm{decomp}_s \in D[n_j]} \mu^s(f_{j \circ j_1}, \dots, f_{j \circ j_s}),$$

with $\mathrm{decomp}_s \in D[n_j]$ denoting a length s decomposition and f_{j_i} , $1 \leq i \leq s$, its elements. This also corresponds to the discussion in Section 7.1.2 describing the case of 2-simplices of $N(A)$.

The simplicial maps are as follows. Given an injection $k : [m] \rightarrow [n]$ and an n simplex f , define an m -simplex f' by $\{f'_j = f_{k \circ j}\}$, where $j : [l] \rightarrow [m]$ is an injection.

On the other hand, given the unique surjection in Δ : $s_i : [n+1] \rightarrow [n]$, $s_i(i+1) = s_i(i)$, and given an n -simplex f , define an $(n+1)$ -simplex f' by setting

$$f'_j = \begin{cases} e_{X_i} & \text{if } j = \{i, i+1\}; \\ f_{s_i|_j} & \text{if } s_i|_j \text{ is injective.} \\ 0 & \text{otherwise,} \end{cases}$$

for $j : [l] \rightarrow [n+1]$ an injection. It is straightforward but tedious to verify that the latter is indeed a face and that simplicial relations are satisfied. On the other hand,

Faonte [5] given a conceptual construction of the above nerve so that the above is automatic.

Proposition A.11. [10, 2.3.2], [5] *For A a unital A_∞ category its nerve $\mathcal{A} = N(A)$ is a ∞ -category.*

For the reader's convenience we outline the proof here.

Proof. Suppose we have an inner horn $\rho_k : \Lambda_k^n \rightarrow NA$. In particular, by the construction of the simplices of NA , corresponding to the faces of the horn, there are determined $f_j \in \text{hom}(A)$, for all $j : [n_j] \rightarrow [n]$ except $j = [n] - \{k\}$ and $j = [n]$. Set $f_{[n]} = 0$, and set

$$f_{[n]-\{k\}} = \sum_{0 < i < n; i \neq k} f_{[n]-\{i\}} + \sum_{s \geq 2} \sum_{\text{decomp}_s \in D[n]} \mu^s(f_{j_1}, \dots, f_{j_s}),$$

Only thing left to check is that as defined the data $\{f_j\}$ determines a n -simplex.

This amounts to verifying a pair of identities:

$$\begin{aligned} 0 &= \sum_{0 < i < n} f_{[n]-i} + \sum_{s \geq 2} \sum_{\text{decomp}_s \in D[n]} \mu^s(f_{j_1}, \dots, f_{j_s}), \\ \mu^1(f_{n-\{k\}}) &= \sum_{0 < i < n-1; i \neq k} f_{[n]-k-i} + \sum_{s \geq 2} \sum_{\text{decomp}_s \in D([n-1])} \mu^s(f_{j_k \circ j_1}, \dots, f_{j_k \circ j_s}), \end{aligned}$$

where $j_k : [n-1] \rightarrow [n]$ is the inclusion with image $[n] - \{k\}$.

The first identity is immediate by the definition of $f_{[n]-\{k\}}$. For the second identity, a direct calculation is long but straightforward, using the A_∞ associativity equations. For $n = 2$ this is automatic and for $n = 3$ this can be checked in a few lines. However, for general n it is certainly better to prove this using conceptual methods as is done in Faonte [5]. \square

For $F : A \rightarrow B$ an A_∞ functor we define $NF : NA \rightarrow NB$ via the assignment:

$$f_j \mapsto \sum_{\text{decomp}_s \in D[n_j]} F^s(f_{j_1}, \dots, f_{j_s}).$$

Lemma A.12. [10], [5] *The assignment $A \mapsto NA$, and $F \mapsto NF$ as above, determines a functor*

$$N : A_\infty - \text{Cat}^{\text{unit}} \rightarrow \infty - \text{Cat}.$$

The details on why this constitutes a functor N are omitted.

REFERENCES

- [1] M. AKVELD AND D. SALAMON, *Loops of Lagrangian submanifolds and pseudoholomorphic discs.*, Geom. Funct. Anal., 11 (2001), pp. 609–650.
- [2] O. BUŞE, *Deformations of Whitehead products, symplectomorphism groups, and Gromov-Witten invariants.*, Int. Math. Res. Not., 2010 (2010), pp. 3303–3340.
- [3] K. COSTELLO, *Topological conformal field theories and Calabi-Yau categories.*, Adv. Math., 210 (2007), pp. 165–214.
- [4] G. FAONTE, *A_∞ -functors and homotopy theory of dg-categories*, J. Noncommut. Geom., 11 (2017), pp. 957–1000.
- [5] ———, *Simplicial nerve of an A_∞ -category.*, Theory Appl. Categ., 32 (2017), pp. 31–52.
- [6] M. GROMOV, *Pseudo holomorphic curves in symplectic manifolds.*, Invent. Math., 82 (1985), pp. 307–347.
- [7] V. GUILLEMIN, E. LERMAN, AND S. STERNBERG, *Symplectic fibrations and multiplicity diagrams*, Cambridge University Press, Cambridge, 1996.

- [8] M. HUTCHINGS, *Floer homology of families. I.*, Algebr. Geom. Topol., 8 (2008), pp. 435–492.
- [9] H.-V. LÊ AND K. ONO, *Parametrized Gromov-Witten invariants and topology of symplectomorphism groups.*, (2008), pp. 51–75.
- [10] H. LEE TANAKA, *A functor from Lagrangian cobordisms to the Fukaya category*, thesis, (2013).
- [11] K. LEFEVRE-HASEGAWA, *Sur les a_∞ -categories*, thesis, (2003).
- [12] J. LURIE, *Higher algebra*, <http://www.math.harvard.edu/~lurie/papers/HigherAlgebra.pdf>.
- [13] ———, *Higher topos theory.*, Annals of Mathematics Studies 170. Princeton, NJ: Princeton University Press. , 2009.
- [14] J.-P. MAGNOT AND J. WATTS, *The diffeology of Milnor’s classifying space*, Topology Appl., 232 (2017), pp. 189–213.
- [15] J. W. MILNOR, *Construction of universal bundles. II.*, Ann. Math. (2), 63 (1956), pp. 430–436.
- [16] D. NADLER AND H. L. TANAKA, *A stable infinity-category of lagrangian cobordisms*, <http://arXiv.org/abs/1109.4835>.
- [17] OH AND TANAKA, *Continuous and coherent actions on wrapped Fukaya categories*, <https://arxiv.org/abs/1911.00349>.
- [18] Y.-G. OH, *Floer cohomology of Lagrangian intersections and pseudo-holomorphic disks. I*, Commun. Pure Appl. Math., 46 (1993), pp. 949–994.
- [19] E. RIEHL, *A model structure for quasi-categories*, <http://www.math.harvard.edu/~eriehl/topic.pdf>.
- [20] Y. SAVELYEV, *Global Fukaya category II: singular connections, quantum obstruction theory and other applications*, <https://arxiv.org/abs/1408.3250>.
- [21] Y. SAVELYEV, *Smooth simplicial sets and universal Chern-Weil homomorphism*, <https://arxiv.org/abs/2112.13272>.
- [22] Y. SAVELYEV, *Quantum characteristic classes and the Hofer metric*, Geom. Topol., 12 (2008), pp. 2277–2326.
- [23] P. SEIDEL, π_1 of symplectic automorphism groups and invertibles in quantum homology rings, Geom. Funct. Anal., 7 (1997), pp. 1046–1095.
- [24] P. SEIDEL, *Fukaya categories and Picard-Lefschetz theory*, Zurich Lectures in Advanced Mathematics. Zürich: European Mathematical Society (EMS). vi, 326 p. EUR 46.00 , 2008.
- [25] N. SHERIDAN, *On the Fukaya category of a Fano hypersurface in projective space.*, Publ. Math., Inst. Hautes Étud. Sci., 124 (2016), pp. 165–317.
- [26] B. TOEN, *The homotopy theory of dg-categories and derived Morita theory.*, Invent. Math., 167 (2007), pp. 615–667.

UNIVERSITY OF COLIMA, CUICBAS

Email address: yasha.savelyev@gmail.com