

GEODESIC STRING COUNTING INVARIANTS OF MANIFOLDS

YASHA SAVELYEV

ABSTRACT. We study $\mathbb{Q}$ -valued metric/topological invariants of manifolds, by counting closed geodesics, and using the Fuller index. As one application, let g be a regular Finsler metric on T^n sufficiently nearby to the standard flat metric. Let o be a closed geodesic in a non-trivial class β , s.t. a given prime p divides multiplicity of o . Then there is at least one other geometrically distinct such o (with the same $p!$). This holds for any regular g with finitely many class β orbits, provided a certain conjectural topological invariance of our geodesic string count holds. The formulation extends from metrics to Reeb vector fields on the unit cotangent bundle of X . The conjecture is partially verified, and this plays a role for a number of other applications. Along the way, we also prove that sky catastrophes of smooth dynamical systems are not geodesible by a certain class of forward complete Riemann-Finsler metrics, in particular by complete Riemannian metrics with non-positive sectional curvature. This partially answers a question of Fuller and gives further examples for our theory here.

1. INTRODUCTION

Using counts of *geodesic strings* (equivalence classes of closed, constant speed geodesics up to reparametrization S^1 action), and the Fuller index [7], we will define certain rational number valued, deformation invariants for complete Riemann-Finsler manifolds. A basic case is a complete Riemannian manifold with non-positive sectional curvature, and in this case we get deformation invariants of the metric, provided the deformation is through non-positive sectional curvature metrics.

Studying connections of the Fuller index with Riemann-Finsler geometry was suggested by Fuller himself in the 1960's but was never developed. An outline of Fuller's theory and basic definitions are given here in the Appendix.

Suppose for the moment X is compact. For a non-constant, free homotopy class β of a loop in X , we say that a metric g on X is β -*regular* if all of its class β closed geodesics are non-degenerate the usual sense, or equivalently the closed orbits of the associated geodesic flow are dynamically non-degenerate. For a β -regular g , with finitely many class β geodesic strings, we have the following formula for the count:

$$(1.1) \quad F(g, \beta) = \sum_o \frac{(-1)^{\text{morse}(o)}}{\text{mult}(o)} \in \mathbb{Q},$$

where:

- The sum is over class β g -geodesic strings o .
- $\text{morse}(o)$ denotes the Morse index of o , (meaning the Morse-Bott index of the associated critical submanifold (diffeomorphic to S^1) of the loop space).
- $\text{mult}(o)$ is the geometric multiplicity, equivalently the order of the corresponding isotropy subgroup of S^1 , where S^1 is acting on the loop space by reparametrization.

Key words and phrases. locally conformally symplectic manifolds, Gromov-Witten theory, virtual fundamental class, Fuller index.

Supported by CONAHCYT research grant CF-2023-I.

Let $\pi_1(X)$ denote the set of free homotopy classes of loops in X , and let $L_\beta X$ denote the subset of the free loop space with elements in class $\beta \in \pi_1(X)$. The following is a direct corollary of Theorem 1.25, and exemplifies the basic topological invariance of the count:

Corollary 1.2. *Let X, g, β be as above and such that β is indivisible, (cannot be represented by a multiply covered loop). Then the S^1 equivariant homology $H_*^{S^1}(L_\beta X, \mathbb{Z})$ has finite total rank. Denote by $\chi^{S^1}(L_\beta X)$ the Euler characteristic of this homology. Then*

$$F(g, \beta) = \chi^{S^1}(L_\beta X).$$

Corollary 1.2 above, and its parent Theorem 1.25 lead us to conjecture full topological invariance, see Conjecture 1 in Section 1.1.1. The latter theorem is an important ingredient for some geometric applications concerning closed geodesic counts.

Assuming full topological invariance of our string counting invariant we get the following perhaps mysterious theorem.

Theorem 1.3 (Proof in Section 10). *Assume the Conjecture 1, let $\beta \in \pi_1(T^n)$ be a non-trivial class and let g be a β -regular Finsler metric on T^n having finitely many β class geodesic strings. Suppose that o is a g -geodesic string s.t.:*

- (1) o has class β .
- (2) $p \mid \text{mult}(o)$, where p is prime,

then there is at least one other such o (same p).

Remark 1.4. We can extend the above from metrics to Reeb vector fields on the unit cotangent bundle, representing the standard contact structure. But now counting closed Reeb orbits in classes $\tilde{\beta}$, lifting a class β as above. However, we need a stronger form of the Conjecture 1, see Remark 1.18. The following local result, Theorem 1.5, also holds in the Reeb context, with basically the same proof.

Without Conjecture 1 we have a local form of the above, which is one of our main concrete applications. Even this local form appears to be inaccessible to standard Morse theory techniques.

Theorem 1.5 (Proof in Section 8). *Let g_0 be the standard flat metric on T^n , and $\beta \in \pi_1(T^n)$ a non-trivial class. For all $\epsilon > 0$ sufficiently small, whenever g' is a Finsler metric C^2 ϵ close to g_0 and is β -regular, the following holds. Suppose that o is a g' -geodesic string s.t.:*

- (1) o has class β .
- (2) $p \mid \text{mult}(o)$, where p is prime,

then there is at least one other such o (same p). Moreover, any Finsler metric g_1 taut deformation equivalent to g_0 (see Definition 1.12) has the same property as g_0 , above. An example for such a metric is in Figure 1.

Recall that a now classical theorem of Preissmann [11] implies that there are no non-trivial compact products with negative sectional curvature. The following is one generalization. We denote by $\pi_1^{\text{inc}}(X)$ the set of free homotopy classes of loops incompressible to the ends, in the sense of Definition 1.9, (non-constant classes when X is closed).

Theorem 1.6 (Proof in Section 12). *Let $X = Z \times Y$ where Z, Y admit complete Riemannian metrics with non-positive sectional curvature, Y is closed, $\pi_1^{\text{inc}}(Z) \neq \emptyset$, $\chi(Y) \neq \pm 1$. Then:*

- X does not admit a complete metric of negative sectional curvature, or a forward complete Finsler metric with negative flag curvature.
- Moreover, X does not admit a forward complete Finsler metric with a unique and non-degenerate class β geodesic string, for any β in the image of the the set map

$$\pi_1^{\text{inc}}(Z) \longrightarrow \pi_1^{\text{inc}}(X),$$

induced by fiber inclusion $Z \rightarrow X$.

This theorem is very close to being sharp, for example the conclusion of the corollary is false if $Y = S^1$ and $Z = S^1 \times \mathbb{R}$. As $X = T^2 \times \mathbb{R}$ admits the warped product metric $g_{T^2} \times_{e^t} g_{\mathbb{R}}$ (with respect to the function $f = e^t$ on $\mathbb{R}$) where $g_{T^2}, g_{\mathbb{R}}$ are the flat metrics. This warped product has constant negative sectional curvature -1 . So it is essential that not only $\pi_1(Z)$ be non-trivial but also that there is a class incompressible to the ends. Of course, $\chi(Y) = 1$ is also obviously essential, otherwise we may take $Y = pt$. The condition $\chi(Y) \neq -1$ is however not obviously essential.

The following is a partial corollary of the theorem above.

Theorem 1.7 (Proof in Section 9). (1) *Let Σ be a connected, possibly infinite type oriented surface. There is a forward complete Finsler metric on $M = \Sigma \times T^n$ with negative flag curvature, if and only if Σ is the infinite cylinder or is $\mathbb{R}^2$.*

(2) *For any non-trivial class β , T^n does not admit a Finsler metric g with a unique and non-degenerate β -class g -geodesic string.*

When Σ is the infinite cylinder, a counterexample g to part one of the theorem can be given as the warped product Riemannian metric.

Remark 1.8. The Riemannian version of the first part of the theorem above can be proved via the remarkable flat strip theorem [5]. Furthermore, Theorems 1.7, 1.6 can be proved via S^1 equivariant Morse theory for the (Finsler) energy functional on the loop space of M , but the proofs are harder, and I believe the above is the first statement of the theorems.

We will also give various generalizations of the above results to fibrations. For fibrations, the most obvious analogue of Preissmann's theorem fails even assuming compactness. In fact, every closed 3-manifold X^3 , for which there is no injection $\mathbb{Z}^2 \rightarrow \pi_1(X, x_0)$, and which fibers over a circle has a hyperbolic structure g_h , Thurston [13].

1.1. Setup and more extensive statements. Proofs of many results here are postponed till subsequent sections.

Terminology 1. From now on, all our metrics are Riemann-Finsler (a.k.a. Finsler) metrics unless specified to be Riemannian, and usually denoted by just g . Completeness, always means forward completeness, and it is an assumption for all our metrics. Curvature always means sectional curvature in the Riemannian case and flag curvature in the Finsler case. Thus we will usually just say complete metric g , for a forward complete Riemann-Finsler metric. A reader may certainly choose to interpret all metrics as Riemannian metrics, completeness as standard completeness, and curvature as sectional curvature.

In what follows $\pi_1(X)$ denotes the set of free homotopy classes of continuous maps $o : S^1 \rightarrow X$.

Definition 1.9. *Let X be a smooth manifold. Fix an exhaustion by nested compact sets $\bigcup_{i \in \mathbb{N}} K_i = X$, $K_i \supset K_{i-1}^\circ$ for all $i \geq 1$. We say that a class $\beta \in \pi_1(X)$ is end compressible if β is in the image of*

$$\text{inc}_* : \pi_1(X - K_i) \rightarrow \pi_1(X)$$

for all i , where $\text{inc} : X - K_i \rightarrow X$ is the inclusion map. We say that β is **end incompressible** (or **incompressible to the ends**) if it is not end compressible.

Let $\pi_1^{\text{inc}}(X)$ denote the set of such end incompressible classes. When X is compact, we set $\pi_1^{\text{inc}}(X) := \pi_1(X) - \text{const}$, where const denotes the set of homotopy classes of constant loops.

It is easily seen that the above is well defined (independent of the choice of an exhaustion) and moreover any homeomorphism $X_1 \rightarrow X_2$ of a pair of manifolds induces a set isomorphism $\pi_1^{\text{inc}}(X_1) \rightarrow \pi_1^{\text{inc}}(X_2)$.

Denote by $L_\beta X$ the class $\beta \in \pi_1^{\text{inc}}(X)$ component of the free loop space of X , with its compact open topology. Let g be a complete metric on X , and let $S(g, \beta) \subset L_\beta X$ denote the subspace of all constant speed parametrized, smooth, closed g -geodesics in class β .

Definition 1.10. We say that a metric g on X is β -**taut** if it is complete and $S(g, \beta)$ is compact. We will say that g is **taut** if it is β -taut for each $\beta \in \pi_1^{\text{inc}}(X)$.

Lemma 1.11. A complete metric g with non-positive curvature satisfies:

- All of its closed geodesics are minimizing in their free homotopy class.
- It is taut.

Proof. The first part is a standard consequence of the Cartan-Hadamard theorem. The second part follows by the first part and Lemma 6.1. $\square$

It should be emphasized that taut metrics form a much larger class of metrics than just non-positive curvature metrics. For example any sufficiently C^1 small perturbation of a metric with non-positive curvature will be taut. (Indeed, this is crucial for the construction of our invariant.) Another class of examples comes by way of Lemma 1.33 ahead, these metrics may not be non-positively curved nor close to non-positively curved metrics. A simple concrete example very far away from being non-positively curved in Example (2).

Definition 1.12. Let $\beta \in \pi_1^{\text{inc}}(X)$, and let g_0, g_1 be a pair of β -taut metrics on X . A β -**taut deformation** between g_0, g_1 , is a continuous (in the topology of C^0 convergence on compact sets) family $\{g_t\}$, $t \in [0, 1]$ of complete metrics on X , s.t.

$$S(\{g_t\}, \beta) := \{(o, t) \in L_\beta X \times [0, 1] \mid o \in S(g_t, \beta)\}$$

is compact. We say that $\{g_t\}$ is a **taut deformation** if it is β -taut for each $\beta \in \pi_1^{\text{inc}}(X)$. The above definitions of tautness are extended naturally to the case of a smooth fibration $X \hookrightarrow P \rightarrow [0, 1]$, with a smooth fiber-wise family of metrics.

A useful criterion for β -tautness is the following.

Theorem 1.13 (Proof in Section 3). Let $\beta \in \pi_1^{\text{inc}}(X)$. Let $\{g_t\}_{t \in [0, 1]}$ be a continuous family of complete metrics on X . Suppose that:

$$\sup_t \left| \max_{o \in S(g_t, \beta)} l_{g_t}(o) - \min_{o \in S(g_t, \beta)} l_{g_t}(o) \right| < \infty,$$

where l_{g_t} is the length functional with respect to g_t , then $\{g_t\}$ is β -taut.

For example, the hypothesis is trivially satisfied if g_t have the property that all their class β closed geodesics are minimal in their homotopy class.

Corollary 1.14. If g_t , $t \in [0, 1]$ have non-positive curvature then $\{g_t\}$ is taut.

Proof. This follows by Theorem 1.13 and by Lemma 1.11. $\square$

1.1.1. *The geodesic string counting invariant F .* Let $\mathcal{G}(X)$ be the set of equivalence classes of taut metrics g , where g_0 is equivalent to g_1 whenever there is a taut deformation between them. We may denote an equivalence class by its representative g by a slight abuse of notation. The following is proved in Section 6.

Theorem 1.15. *For each manifold X there is a natural, non-trivial functional:*

$$F : \mathcal{G}(X) \times \pi_1^{\text{inc}}(X) \rightarrow \mathbb{Q},$$

extending the definition in (1.1).

For g β -regular we have already defined $F(g, \beta)$ at the beginning of the Introduction. For a general g we define $F(g, \beta) := F(g', \beta)$, where g' is a sufficiently small C^∞ -perturbation of g , is β -regular, and remains β -taut. The content of Theorem 1.15 is then that this is well defined. We will show that this is a consequence of general Fuller index theory.

Corollary 1.16. *Suppose for a pair g_0, g_1 of β -taut metrics on X :*

$$F(g_0, \beta) \neq F(g_1, \beta),$$

then any path $\{g_t\}$, connecting g_0, g_1 , is not β -taut.

Proof. The fact that any connecting $\{g_t\}$ is not β -taut is just a direct corollary of the theorem above. $\square$

I was unable to find such a pair g_0, g_1 , in fact there is strong evidence to believe the following:

Conjecture 1. *F does not depend on the choice of a smooth structure and taut metric. In other words we have the following. Let X be a topological manifold, define:*

$$\mathcal{S}(X) : \pi_1^{\text{inc}}(X) \rightarrow \mathbb{Q} \sqcup \{\infty\}$$

by:

$$\mathcal{S}(X)(\beta) = \begin{cases} \infty, & \text{if } X \text{ does not admit a smooth structure and a } \beta\text{-taut Finsler metric.} \\ F(g, \beta), & \text{if } g \text{ is a } \beta\text{-taut Finsler metric on } X. \end{cases},$$

then $\mathcal{S}$ is well defined and hence determines a topological invariant of topological manifolds. So if $f : X_1 \rightarrow X_2$ is a homeomorphism then $\mathcal{S}(X_1)(\beta) = \mathcal{S}(X_2)(f_(\beta))$.*

Theorem 1.25 in the following section partially proves this conjecture.

Remark 1.17. The smooth manifold version of this conjecture, i.e. that $\mathcal{S}(X)$ is a smooth manifold invariant, is implied by the Reeb sky catastrophe conjecture described in the next section. However, the above seems to be much more basic as will be apparent from the proof of Theorem 1.25.

Remark 1.18. The formulation of the conjecture naturally extends to Reeb vector fields. That is let λ be a contact form on the unit cotangent bundle C of a smooth manifold X (for simplicity closed), with λ representing the Liouville contact structure (i.e. the standard contact structure). Assume that the space of closed, class $\tilde{\beta}$ λ -Reeb orbits on C is compact. Then the Fuller index (See Section 6) of this compact set is a topological invariant of X .

1.1.2. A short digression on sky catastrophes. [A.1.](#)

Definition 1.19 (Preliminary). *A **sky catastrophe** for a smooth family $\{X_t\}$, $t \in [0, 1]$, of non-vanishing vector fields on a closed manifold M is a continuous family of closed orbit strings $\tau \mapsto o_{t_\tau}$, o_{t_τ} is an orbit string of X_{t_τ} , $\tau \in [0, \infty)$, such that the period of o_{t_τ} is unbounded from above.*

Sky catastrophes first appeared in the work of Fuller [6]. One general definition appears in [12], the point there being that there are no regularity assumptions on $\{X_t\}$. The preliminary definition becomes equivalent to the more general definition given certain regularity conditions on the family $\{X_t\}$.

The following corollary of Theorem [1.13](#) may be of independent interest.

Corollary 1.20. *Sky catastrophes of vector fields on closed manifolds are not geodesible by metrics all of whose geodesics are minimal. That is if a family $\{X_t\}$ has a sky catastrophe, there is no family $\{g_t\}$ of metrics such that the orbits of X_t are unit speed parametrized g_t -geodesics.*

Fuller at the end of [7] has asked for any metric conditions on vector fields to rule out sky catastrophes, see Appendix [A.1](#). By the above, non-positivity of curvature is one such condition. So this is a partial answer to his question.

The main conjecture in [12] is that geodesible, and more generally Reeb sky catastrophes ($\{X_t\}$ are Reeb) are unstable, in the sense that there is a C^0 nearby family $\{X'_t\}$ which does not have a sky catastrophe (at least in the fixed class β). Let's call this **Reeb sky catastrophe conjecture**. The reason to conjecture such a thing, is that a theorem in [12] says that if $\{o_{t_\tau}\}_\tau$ is a sky catastrophe for a family of Reeb vector fields $\{X_t\}$ then the length of the corresponding projection $\tau \mapsto t_\tau$ is infinite. (Assuming some minor regularity on $\{o_{t_\tau}\}$.) The latter suggests that the structure of such a sky catastrophe is very fragile.

Remark 1.21. The Reeb sky catastrophe conjecture implies the geodesible Seifert conjecture (apparently still an open problem), by the main result of [12]. Hence, this is a subtle question.

The following is a variant of Corollary [1.16](#).

Theorem 1.22. *Suppose for a pair g_0, g_1 of β -taut metrics on a closed manifold M :*

$$F(g_0, \beta) \neq F(g_1, \beta),$$

then any path $\{g_t\}$, connecting g_0, g_1 , has a sky catastrophe, meaning that the corresponding family $\{X_t\}$ of Reeb vector fields on the unit cotangent bundle (cf. Section [6](#)) has a sky catastrophe as in Definition [A.5](#).

Proof. This directly follows by [12, Theorem 3.2]. □

So that if such a pair g_0, g_1 exists the Reeb sky catastrophe conjecture is disproved, and in a rather extreme fashion.

1.1.3. Basic results on the invariant F .

Definition 1.23. *Let $\beta \in \pi_1(X)$. For any based point $x_0 \in \text{image } \beta \subset X$ (for $\text{image } \beta$ the image of some representative of β) there is a naturally determined element $\beta_{x_0} \in \pi_1(X, x_0)$ well defined up to an inner automorphism, (concatenate a representative of β with a path from x_0 to a point in $\text{image } \beta$).*

- We say that a class $\beta \in \pi_1(X, x_0)$ is **indivisible** if whenever $\beta = \alpha^k$ for some $\alpha, k > 0$ then $k = 1$.

- We say that a class $\beta \in \pi_1(X, x_0)$ is a **k-power** if $\beta = \alpha^k$, $k > 1$, for some α which is not a n -power for any n .
- We say that β is **atomic** if it is a k -power for some k .¹
- We say that $\beta \in \pi_1(X)$ is indivisible, respectively is a k -power, respectively is atomic if for any x_0 as above, β_{x_0} is indivisible, respectively is a k -power, respectively is atomic. This notion does not depend on the choice of β_{x_0} in its inner automorphism class. In fact, β is k -power just means that β has a smooth representative with multiplicity k .

Note that a class $\beta \in \pi_1^{inc}(X)$ is indivisible iff any representative of this class is not multiply covered.

Example 1. Let g be a Riemannian metric with negative sectional curvature on a closed manifold X and $\beta \in \pi_1(X)$ a class represented by a multiplicity n closed geodesic, then

$$(1.24) \quad F(g, \beta) = \frac{1}{n}.$$

In particular, if β is indivisible then $F(g, \beta) = 1$. More generally, (1.24) holds whenever g has a unique and non-degenerate geodesic string in class β , where non-degenerate is as in the Introduction.

If $\beta \in \pi_1^{inc}(X)$ is indivisible, then it is easy to see that the reparametrization S^1 action on $L_\beta X$ is free (see Appendix A), so that $H_*^{S^1}(L_\beta X, \mathbb{Z}) \simeq H_*(L_\beta X/S^1, \mathbb{Z})$, where $H_*^{S^1}(L_\beta X, \mathbb{Z})$ denotes the S^1 -equivariant homology. Moreover, we have the following result proved in Section 11.

Theorem 1.25. *Suppose that $\beta \in \pi_1^{inc}(X)$ is indivisible, and X admits a β -taut metric. Then $H_*^{S^1}(L_\beta X, \mathbb{Z})$ has finite total rank. Denote by $\chi^{S^1}(L_\beta X)$ the Euler characteristic of this homology. Then for any β -taut metric g on X :*

$$F(g, \beta) = \chi^{S^1}(L_\beta X).$$

Consequently, $F(g, \beta)$ is independent of the choice of β -taut metric g , for indivisible β . And Furthermore, Conjecture 1 holds on the subset of classes β which are indivisible.

Example 2. A basic example of a β -taut metric that is not nearby to a non-positive curvature metric is the following. Let g be a metric on the 2-torus with exactly two class β g -geodesic strings, for β one of the π_1 generators. We can take such a g_0 with arbitrarily large curvature, see Figure 1. The metric g_0 in this figure is moreover clearly taut deformation equivalent to the standard flat metric. Taking products of metrics of this sort gives higher dimensional examples.

More examples for the theorem above can be found by the proof of Theorem 1.27.

Remark 1.26. If β is not indivisible, the idea behind Theorem 1.25 breaks down, as the S^1 -equivariant homology of $L_\beta X$ may then be infinite dimensional even if X admits a β -taut g . As a trivial example, this homology is already infinite dimensional when g is negatively curved, and the class β geodesic is k -covered, as then this homology is the group homology of $\mathbb{Z}_k$. However for a general β we should still have that $F(g, \beta)$ is the orbifold Euler characteristic, of a suitable orbifold quotient $L_\beta X/S^1$ which in principle proves Conjecture 1. The difficulty in making this precise is that at the moment there is no such orbifold Euler characteristic theory as $L_\beta X/S^1$ is infinite dimensional (thinking of it as a stack or an orbifold).

We can use the above, and the product formula of Theorem 12.1 to get:

Theorem 1.27 (Proof in Section 12). *Every rational number has the form $F(g, \beta)$ for some β -taut Riemannian g on some compact manifold X and for some β .*

¹As we are working with non-compact manifold, we may in principle have infinitely divisible classes.




FIGURE 1. The class β is represented by the drawn curve.

1.1.4. *Applications to existence of negative curvature metrics.* A celebrated theorem of Preissmann [11] says that there are no negative sectional curvature metrics on compact products. Fibration counterexamples to Preissmann's product theorem certainly exist as mentioned in the Introduction. We are going to give a certain generalization of Preissmann's theorem to fibrations, with possibly non-compact fibers, also replacing the negative sectional curvature condition by a significantly weaker condition. Though the definitions may seem somewhat ad hoc, we just want enough generality to incorporate some interesting "flat bundle" examples. The latter are of interest in 3-manifold theory, for instance vis-à-vis Thurston's classification [13].

Definition 1.28. Let $Z \hookrightarrow X \xrightarrow{p} Y$ be a smooth fiber bundle, and let g be a Riemannian metric on X . Denote by

$$T^{\text{vert}} X = \ker(dp)$$

the vertical tangent bundle, and let

$$P^{\text{vert}} : TX \rightarrow T^{\text{vert}} X$$

be the g -orthogonal projection. We say that p is **parallel** if

$$\nabla P^{\text{vert}} = 0,$$

where ∇ is the Levi-Civita connection of g , and P^{vert} is viewed as a $(1,1)$ -tensor field on X . Equivalently, for all vector fields A, B on X ,

$$\nabla_A(P^{\text{vert}} B) = P^{\text{vert}}(\nabla_A B).$$

Definition 1.29. Let $Z \hookrightarrow X \xrightarrow{p} Y$ be a smooth fiber bundle with X having a β -taut Riemannian metric g , for $\beta \in \pi_1^{\text{inc}}(X)$, and let g_Y be a metric on Y . Suppose that:

- (1) The fibers $Z_y = p^{-1}(y)$ are totally g -geodesic, for closed geodesics in class β . We denote by g_y the metric g restricted to Z_y .
- (2) p is parallel.
- (3) For any pair of fibers (Z_{y_0}, g_{y_0}) , (Z_{y_1}, g_{y_1}) , and a path $\gamma : [0, 1] \rightarrow Y$ from y_0 to y_1 the fiber family $\{(Z_{\gamma(t)}, g_{\gamma(t)})\}$ furnishes a taut deformation.
- (4) p projects g -geodesics to geodesics of Y, g_Y .

We then call $p : X \rightarrow Y$ a **β -taut fibration**, with the metrics g, g_Y and g_Z all possibly implicit.

Definition 1.30. For $Z \hookrightarrow X \rightarrow Y$ as above, we say that $\beta \in \pi_1(X)$ is a **fiber class** if it is in the image of the set map $i_Z : \pi_1(Z) \rightarrow \pi_1(X)$, induced by inclusion.

In the above definition of a taut fibration and the following theorem we need the auxiliary metric g on X to be Riemannian, and there is no obvious extension of the theorem to the Riemann-Finsler case. However, the conclusions of the theorem are for Riemann-Finsler metrics. The following is perhaps the main technical result of the paper, proved in Section 12.

Theorem 1.31. *Let $p : (X, g) \rightarrow (Y, g_Y)$ be a β -taut fibration, where $\beta \in \pi_1^{inc}(X)$ is a fiber class. Suppose further that Y is connected, closed, $\chi(Y) \neq \pm 1$ and is such that all smooth closed contractible g_Y -geodesics in Y are constant. Then the following holds:*

- X does not admit a complete Riemann-Finsler metric with negative curvature.
- X does not admit a complete Riemann-Finsler metric with a unique and non-degenerate class β geodesic string.

Note that $\chi(Y) \neq 1$ is of course essential, as the trivial fibration $X \rightarrow \{pt\}$, with X admitting a complete negatively curved metric, will satisfy the hypothesis. The condition that there is a fiber class $\beta \in \pi_1^{inc}(X)$ is also essential, for any vector bundle over a manifold admitting a Riemannian metric of negative curvature admits a metric of negative curvature, Anderson [1].

Theorem 1.6 gives one set of examples. A further basic set of examples for the theorem is obtained by starting with any closed Y , and any homomorphism

$$(1.32) \quad \phi : \pi_1(Y, y_0) \rightarrow \text{Isom}(Z, g_Z), \quad (\text{the group of all isometries}).$$

where g_Z is a β_Z -taut Riemannian metric, for some $\beta_Z \in \pi_1^{inc}(Z)$ (for example (Z, g_Z) is a non-simply connected complete hyperbolic surface of genus at least one). Suppose further:

- (1) The orbit

$$\text{Orb} := \bigcup_{\gamma \in \pi_1(Y, y_0)} \phi_*(\gamma)(\beta_Z)$$

is finite.

- (2) Y is closed and connected.

- (3) All contractible smooth closed g_Y geodesics in Y are constant.

We have the obvious induced diagonal action

$$\begin{aligned} \pi_1(Y, y_0) &\rightarrow \text{Diff}(Z \times \tilde{Y}), \quad (\text{the group of all diffeomorphisms}), \\ \gamma &\mapsto ((z, y) \mapsto (\phi(\gamma)(z), \gamma \cdot y)), \end{aligned}$$

for $\tilde{Y}$ the universal cover of Y . Taking the quotient of $Z \times \tilde{Y}$ by this action, we get an associated “flat” bundle $Z \hookrightarrow X_\phi \xrightarrow{p} Y$, with a metric g_ϕ induced from the product metric $\tilde{g} = g_Z \oplus g_Y$, on the covering space $q : Z \times \tilde{Y} \rightarrow Z \times Y$.

Lemma 1.33 (Proof in Section 4). *Let $p : (X_\phi, g_\phi) \rightarrow (Y, g_Y)$ be as above, then this is a β -taut fibration, where $\beta = i_*(\beta_Z)$, for $i_* : \pi_1^{inc}(Z) \rightarrow \pi_1^{inc}(X_\phi)$ the set map induced by inclusion.*

By the lemma above, $p : (X_\phi, g_\phi) \rightarrow (Y, g_Y)$ satisfies the hypothesis of the theorem above. Yet more concretely:

Example 3. Suppose we have $\beta_Z \in \pi_1^{inc}(Z)$, and let $\phi : Z \rightarrow Z$ be an isometry of a taut metric g_Z . Then by the construction above, the mapping torus

$$(Z, g_Z) \hookrightarrow (X_\phi, g_\phi) \xrightarrow{\pi} S^1$$

has the structure of a β -taut fibration, satisfying the hypothesis of the theorem, for $\beta = i_*(\beta_Z)$ as above.

The next corollary of Theorem 1.31 is immediate.

Corollary 1.34. *Let*

$$(Z_{g,Z}) \hookrightarrow (X_\phi, g_\phi) \rightarrow (Y, g_Y)$$

be as in the construction above for Z, g_Z having non-positive curvature, and let $\beta_Z \in \pi_1^{\text{inc}}(Z)$. Then if $\chi(Y) \neq \pm 1$:

- (1) X_ϕ does not admit a complete Riemann-Finsler metric with negative curvature.
- (2) Moreover, X_ϕ does not admit a Riemann-Finsler metric with a unique and non-degenerate class β geodesic string, for $\beta = i_*(\beta_Z)$ as above.

As a special case, this applies to the mapping tori X_ϕ , for $\phi : Z \rightarrow Z$ an isometry of a complete Riemannian non-positively curved metric on Z , satisfying the finiteness condition 1. (The non-positive curvature hypothesis is for concreteness we may of course replace this condition by tautness.)

In the special case when Z is compact, the first part of the above corollary can be deduced, with some work, from Preissmann's theorem (specifically, because of the condition 1), see also [3, Theorem 9.3.4] for a generalization that fits our Finsler setting. The second part is new even when Z is compact.

1.2. Estimated counts of multiply covered geodesics. The following is a more extensive version of Theorem 1.5.

Theorem 1.35 (Proof in Section 7). *Suppose that g_1 is a taut metric on X , taut deformation equivalent to a complete metric of negative curvature (everything is Finsler). Suppose that $\beta \in \pi_1^{\text{inc}}(X)$ is a k -power. Let L_β be the length of a class β , g_1 -geodesic. For all $\epsilon > 0$ sufficiently small, whenever g' is C^2 ϵ close to g_1 , and is β -regular, we have:*

$$\sum_{o \in \mathcal{O}_{2L_\beta}(g', \beta)} \frac{(-1)^{\text{morse}(o)}}{\text{mult}(o)} = \frac{1}{k},$$

where $\mathcal{O}_{2L_\beta}(g', \beta)$ is the set of class β geodesic strings with g' -length less than $2L_\beta$.

2. COMPACTNESS PRELIMINARIES

Lemma 2.1. *Suppose that g is a complete metric on X , $\beta \in \pi_1^{\text{inc}}(X)$ and let $S \subset L_\beta X$ be a subset on which the g -length functional is bounded from above. Then the images in X of elements of S are contained in a fixed compact subset of X .*

Proof. Suppose otherwise. Fix an exhaustion by nested compact sets

$$\bigcup_{i \in \mathbb{N}} K_i = X, \quad K_i^\circ \supset K_{i-1}.$$

Then either there is sequence $\{o_i\}_{i \in \mathbb{N}}$, $o_i \in S$ s.t. $o_i \in K_i^c$, for K_i^c the complement of K_i , which contradicts the fact that β is end incompressible. Or there is a sequence $\{o_k\}_{k \in \mathbb{N}}$, $o_k \in S$ s.t.:

- (1) Each o_k intersects K_{i_0} for some i_0 fixed.
- (2) For each $i \in \mathbb{N}$ there is a $k_i > i$ s.t. o_{k_i} is not contained in K_i .

Now if $\text{diam}(o_k)$ is bounded in k , then condition 1 implies that o_k are contained in a set of bounded diameter. (Here $\text{diam}(o_k)$ denotes the diameter of image o_k .) Consequently, by Hopf-Rinow theorem [2], o_k are contained in a compact set. But this contradicts condition 2, and the fact that K_i form an exhaustion of X .

Thus, we conclude that $\text{diam}(o_k)$ is unbounded, but this contradicts the hypothesis. $\square$

Lemma 2.2. *Let g be a complete metric on X , and let $\beta \in \pi_1^{\text{inc}}(X)$. Then there exists a closed g -geodesic minimizing length among all loops in the free homotopy class β .*

Proof. Let

$$L_\beta = \inf_{\gamma \in L_\beta X} \ell_g(\gamma).$$

Choose a minimizing sequence $\{\gamma_j\} \subset L_\beta X$ such that

$$\ell_g(\gamma_j) \rightarrow L_\beta.$$

In particular, the lengths $\ell_g(\gamma_j)$ are uniformly bounded. By Lemma 2.1, the images of all γ_j lie in a fixed compact set $K \subset X$.

Parametrize γ_j by constant speed. Since the images lie in a fixed compact set and the lengths are uniformly bounded, the γ_j are equicontinuous. By Arzelà–Ascoli, a subsequence converges uniformly to a loop $\gamma_\infty \in L_\beta X$. Lower semicontinuity of length gives

$$\ell_g(\gamma_\infty) \leq \liminf_j \ell_g(\gamma_j) = L_\beta.$$

Thus γ_∞ is minimizing. □

3. PROOF OF THEOREM 1.13

Let $\{g_t\}$, $t \in [0, 1]$ be as in the hypothesis, with

$$(3.1) \quad \sup_t \left| \max_{o \in S(g_t, \beta)} l_{g_t}(o) - \min_{o \in S(g_t, \beta)} l_{g_t}(o) \right| < C,$$

and suppose that

$$\sup_{(o, t) \in \mathcal{O}(\{g_t\}, \beta)} l_{g_t}(o) = \infty.$$

Then we have a sequence $\{o_k\}$, $k \in \mathbb{N}$, of closed class β g_{t_k} -geodesics in X , satisfying:

- (1) $\lim_{k \rightarrow \infty} t_k = t_\infty \in [0, 1]$.
- (2) $\lim_{k \rightarrow \infty} l_{g_{t_k}}(o_k) = \infty$, where $l_{g_{t_k}}(o_k)$ is the length with respect to g_{t_k} .

By the preceding lemma, choose a length-minimizing closed g_{t_∞} -geodesic o_∞ in the class β . And let L denote its length g_∞ length. Let g_{aux} be a fixed auxiliary metric on X , and let L_{aux} be the g_{aux} length of o_∞ .

Define a pseudo-metric d_{C_0} on the space of metrics on X as follows. Set $K = \text{image } o_\infty$. And set

$$V \subset TX = \{v \in TX \mid \pi(v) \in K \text{ for } \pi : TX \rightarrow X \text{ the canonical projection, and } |v|_{aux} = 1\},$$

where $|v|_{aux}$ is the norm taken with respect to g_{aux} .

Then define:

$$d_{C^0}(g_1, g_2) = \sup_{v \in V} ||v|_{g_1} - |v|_{g_2}|.$$

By Properties 1 and 2 we may find a $k > 0$ such that:

$$(3.2) \quad d_{C^0}(g_{t_k}, g_{t_\infty}) < \epsilon$$

and

$$(3.3) \quad l_{g_{t_k}}(o_k) > C + L + L_{aux} \cdot \epsilon.$$

By (3.2), we have:

$$l_{g_{t_k}}(o_\infty) < l_{g_{t_\infty}}(o_\infty) + L_{aux} \cdot \epsilon = L + L_{aux} \cdot \epsilon.$$

Combining with (3.3) we get:

$$l_{g_{t_k}}(o_k) > l_{g_{t_k}}(o_\infty) + C.$$

Since we may find a closed g_{t_k} -geodesic o' satisfying $l_{g_{t_k}}(o') \leq l_{g_{t_k}}(o_\infty)$, we get that

$$\left| \max_{o \in S(g_{t_k}, \beta)} l_{g_{t_k}}(o) - \min_{o \in S(g_{t_k}, \beta)} l_{g_{t_k}}(o) \right| > C,$$

and so we are in contradiction.

Thus,

$$\sup_{(o, t) \in \mathcal{O}(\{g_t\}, \beta)} l_{g_t}(o) < \infty.$$

It follows, by an analogue of Lemma 2.1, that the images of all elements $o \in S(\{g_t\}, \beta)$ are contained in a fixed compact $T \subset X$. Compactness of $S(\{g_t\}, \beta)$ then readily follows by the Arzelà-Ascoli theorem. $\square$

4. PROOF OF LEMMA 1.33

Let

$$\phi_* : \pi_1(Y, y_0) \rightarrow \text{Bij}(\pi_1^{\text{inc}}(Z))$$

be the induced action on the set of end-incompressible free homotopy classes. Suppose that the orbit

$$\text{Orb} := \bigcup_{\gamma \in \pi_1(Y, y_0)} \phi_*(\gamma)(\beta_Z)$$

is finite.

As g_Z is taut, $S(g_Z, \phi_*(\gamma)(\beta_Z))$ is compact for each γ , where $S(g_Z, \phi_*(\gamma)(\beta_Z))$ is the space of geodesics as in Definition 1.10. By the condition on contractible geodesics of g_Y , we get:

$$\begin{aligned} S(g_\phi, \beta) &= q_*(S(g_Z \oplus g_Y, \beta)) \\ &= \bigcup_{\beta \in \text{Orb}} q_*(S(g_Z, \beta) \times \tilde{Y}), \end{aligned}$$

for $q_* : L(Z \times \tilde{Y}) \rightarrow L(Z \times Y)$ induced by the quotient map $q : Z \times \tilde{Y} \rightarrow Z \times Y$, (as in the preamble to the statement of the lemma) and where $S(g_Z, \gamma) \times \tilde{Y}$ is understood as a subset

$$S(g_Z, \gamma) \times \tilde{Y} \subset L(Z) \times \tilde{Y} \subset L(Z \times \tilde{Y}).$$

Given that Orb is finite, this then readily implies our claim. $\square$

5. PRELIMINARIES ON REEB FLOW

Let (C^{2n+1}, λ) be a contact manifold with λ a contact form, that is a one form s.t. $\lambda \wedge (d\lambda)^n \neq 0$. Denote by R^λ the Reeb vector field satisfying:

$$d\lambda(R^\lambda, \cdot) = 0, \quad \lambda(R^\lambda) = 1.$$

Recall that a **closed λ -Reeb orbit** (or just Reeb orbit when λ is implicit) is a smooth map

$$o : (S^1 = \mathbb{R}/\mathbb{Z}) \rightarrow C$$

such that

$$\dot{o}(t) = cR^\lambda(o(t)),$$

with $\dot{o}(t)$ denoting the time derivative, for some $c > 0$ called period. Let $S(R^\lambda, \beta)$ denote the space of all closed λ -Reeb orbits in free homotopy class β , with its compact open topology. And set

$$\mathcal{O}(R^\lambda, \beta) = S(R^\lambda, \beta)/S^1,$$

where $S^1 = \mathbb{R}/\mathbb{Z}$ acts by reparametrization $t \cdot o(\tau) = o(t + \tau)$.

6. DEFINITION OF THE FUNCTIONAL F AND PROOFS OF AUXILIARY RESULTS

Let X be a manifold with a taut metric g . Let C be the unit cotangent bundle of X , with its Liouville contact 1-form λ_g . If $o : S^1 = \mathbb{R}/\mathbb{Z} \rightarrow X$ is a constant speed closed geodesic, it has a canonical lift $\tilde{o} : S^1 \rightarrow C$, which is a closed flow line of $s \cdot R^{\lambda_g}$, where $s = |\frac{do}{dt}|$ is the speed of the geodesic, i.e. it is a Reeb orbit.

If $\beta \in \pi_1^{inc}(X)$, let $\tilde{\beta} \in \pi_1(C)$ denote class $[\tilde{o}] \in \pi_1(C)$, where o is a constant speed closed geodesic representing β .

Let $S(R^{\lambda_g}, \tilde{\beta})$ be the orbit space as in Section 5, for the Reeb flow of the contact form λ_g . And set

$$\mathcal{O}_{g,\beta} = \mathcal{O}(R^{\lambda_g}, \tilde{\beta}) := S(R^{\lambda_g}, \tilde{\beta})/S^1,$$

which by construction is identified with the space of class β g -geodesic strings. By the tautness assumptions $\mathcal{O}_{g,\beta}$ is compact.

We then define

$$F(g, \beta) = i(\mathcal{O}_{g,\beta}, R^{\lambda_g}, \tilde{\beta}) \in \mathbb{Q}$$

where the right hand side is the Fuller index as outlined in the Appendix A. As a basic example we have:

Lemma 6.1. *Suppose that g is a complete metric on X , all of whose class $\beta \in \pi_1^{inc}(X)$ geodesics are minimal, then g is β -taut.*

Proof. Returning to the proof of the main lemma. By assumption, closed, class $\beta \in \pi_1^{inc}(X)$ geodesics are g -minimizing in their homotopy class and in particular have fixed length. By the lemma above there is a fixed $K \subset X$ s.t. every class β closed geodesic has image contained in K . Then compactness of $S(g, \beta)$ follows by Arzelà-Ascoli theorem. $\square$

Proof Theorem 1.15. Let $\beta \in \pi_1^{inc}(X)$, be given and let g be β -taut. We just need to prove that $F(g, \beta)$ is invariant under a β -taut deformation of g . So let $\{g_t\}$, $t \in [0, 1]$ be a β -taut deformation of metrics on a compact manifold X . Let $R^{\lambda_{g_t}}$ be the geodesic flow on the g_t unit cotangent bundle C_t . Trivializing the family $\{C_t\}$ we get a family $\{R_t\}$ of flows on $C \simeq C_t$, with R_t conjugate to $R^{\lambda_{g_t}}$.

Let $\mathcal{O}(\{R_t\}, \tilde{\beta})$ be the cobordism as in (A.2), where $\tilde{\beta} \in \pi_1(C)$ is as above. Then $\mathcal{O}(\{R_t\}, \tilde{\beta})$ is compact as $S(\{g_t\}, \beta)$ is compact by assumption.

Basic invariance of the Fuller index, that is (A.3), immediately yields: $F(g_0, \beta) = F(g_1, \beta)$. $\square$

7. PROOF OF THEOREM 1.35

We already know by Example 1 that $F(g_0, \beta) = \frac{1}{k}$ and hence by Theorem 1.15 $F(g_1, \beta) = \frac{1}{k}$. Let U denote the open subset of $L_\beta X$ consisting of loops with g_1 -length less than $2L_\beta$. By [12, Lemma 4.1], for all $\epsilon > 0$ sufficiently small, for any g' , C^2 ϵ -close to g the following holds. Set $g'_t = (t-1) \cdot g_1 + t \cdot g'$, for $t \in [0, 1]$, then

$$\tilde{N} = \mathcal{O}(\{g'_t\}, \beta) \cap (U \times [0, 1])$$

is an open and compact subset of $\mathcal{O}(\{g'_t\}, \beta)$.

Now set

$$N_1 = \tilde{N} \cap (L_\beta X \times \{1\}),$$

and $N_0 = \mathcal{O}(g_1, \beta)$. By the invariance property (A.3) of the Fuller index, we then have that

$$\frac{1}{k} = i(N_0, R^{\lambda_{g_1}}) = i(N_1, R^{\lambda_{g'}}).$$

On the other hand, by construction and by index computations as in [12, Section 2]), we get:

$$i(N_1, R^{\lambda_{g'}}) = \sum_{o \in \mathcal{O}(g', \beta) \cap U} \frac{(-1)^{\text{morse}(o)}}{\text{mult}(o)}.$$

If ϵ is chosen to be sufficiently small then $\mathcal{O}(g', \beta) \cap U = \mathcal{O}_{2L_\beta}(g', \beta)$. So that we are done. $\square$

8. PROOF OF THEOREM 1.5

Let β be a non-zero class, as in the statement. Let $Y \subset T^n$ be a totally geodesic submanifold diffeomorphic to T^{n-1} , containing image β . Let $\alpha \in H^1(T^n, \mathbb{Z})$ be the Poincare dual of Y and let $p : T^n \rightarrow S^1$ be the classifying map of α . Then clearly p is a β -taut fibration with respect to g_0 .

By Theorem 1.25, $F(g_0, \beta) = 0$. Analogously to the proof of Theorem 1.35, we get that for all $\epsilon > 0$ sufficiently small, for any g' as in the statement of our theorem we have:

$$(8.1) \quad \sum_{o \in \mathcal{O}_{2L_\beta}(g', \beta)} \frac{(-1)^{\text{morse}(o)}}{\text{mult}(o)} = 0.$$

But if ϵ is sufficiently small then $\mathcal{O}_{2L_\beta}(g', \beta) = \mathcal{O}(g', \beta)$ by the following. A g' geodesic string o has g_0 geodesic curvature approximately 0, then o must be approximately g_0 minimizing, i.e. approximately of length L_β (this is specific to the Euclidean metric g_0). Hence, any g' geodesic string o can be assumed to have length less $2L_\beta$ by taking ϵ to be sufficiently small.

So we get:

$$\sum_{o \in \mathcal{O}(g', \beta)} \frac{(-1)^{\text{morse}(o)}}{\text{mult}(o)} = 0.$$

To finish the argument, enumerate elements of $\mathcal{O}(g', \beta)$ as $o_0, \dots, o_n$ we then get:

$$\sum_i (-1)^{\text{morse}(o_i)} \prod_{j \neq i} \text{mult}(o_j) = 0.$$

We show that if a prime p divides the multiplicity $\text{mult}(o_k)$ for some $k \in \{1, \dots, n\}$, then there exists at least one other index $m \neq k$ such that p divides $\text{mult}(o_m)$. This will complete the proof.

Here is the proof. Let $M_i = \prod_{j \neq i} \text{mult}(o_j)$ and let $\sigma_i = (-1)^{\text{morse}(o_i)}$. The original equation is:

$$\sum_{i=1}^n \sigma_i M_i = 0.$$

Suppose $p \mid \text{mult}(o_k)$. Then

$$(8.2) \quad \sum_{i=1}^n \sigma_i M_i \equiv 0 \pmod{p}.$$

For any index $i \neq k$, the product M_i contains the factor $\text{mult}(o_k)$:

$$M_i = \text{mult}(o_k) \cdot \prod_{j \neq i, k} \text{mult}(o_j).$$

Since $p \mid \text{mult}(o_k)$, it follows from (8.2) that:

$$\sigma_k M_k + 0 \equiv 0 \pmod{p}.$$

And so

$$M_k \equiv 0 \pmod{p}.$$

By definition, $M_k = \prod_{j \neq k} \text{mult}(o_j)$. Since p is prime and $p \mid \prod_{j \neq k} \text{mult}(o_j)$, by Euclid's Lemma,

$$p \mid \text{mult}(o_m) \text{ for some } m \neq k.$$

And this completes the proof of the claim and the main theorem. $\square$

9. PROOF THEOREM 1.7

The first part follows immediately by part one of Theorem 1.6, as the condition that the surface Σ is not $\mathbb{R}^2$ or $S^1 \times \mathbb{R}$ implies that $\pi_1^{inc}(\Sigma) \neq \emptyset$.

To prove the second part, suppose otherwise, and let g be such a metric. Let o be the unique and non-degenerate geodesic string in some class γ . If γ is indivisible, $F(g, \gamma)$ is g -independent by Theorem 1.25. So we get:

$$0 = F(g_0, \gamma) = F(g, \gamma) = 1,$$

where g_0 is as in the proof of Theorem 1.5 above. We have a contradiction, so that γ is a k -power, and so that $\gamma_{x_0} = \beta^k$ for $k \geq 1$ and $\beta \in \pi_1(X, x_0)$, (γ_{x_0} is as in Definition 1.23) and is indivisible.

Let o be a class β geodesic string, then its k -cover is a class γ geodesic string, and by the uniqueness assumption on the latter, o is likewise unique. Moreover, it is non-degenerate since its k -cover is non-degenerate. We may apply the argument above to get that β is a k' -power for some k' . But this is a contradiction. $\square$

10. PROOF OF THEOREM 1.3

Let $p : T^n \rightarrow S^1$ be the β -taut fibration with respect to g_0 , as in the proof of Theorem 1.5 above. By Theorem 1.25, $F(g_0, \beta) = 0$. Hence, as we assumed Conjecture 1, for any other β -regular g on M with finitely β -class geodesic strings we have:

$$\sum_{o \in \mathcal{O}(g, \beta)} \frac{(-1)^{\text{morse}(o)}}{\text{mult}(o)} = 0.$$

The conclusion then readily follows by basic arithmetic. $\square$

11. PROOF OF THEOREM 1.25

For simplicity, as the full argument is already lengthy, we prove the theorem under the assumption that g is Riemannian. In the Finsler case we must use a more modern variational setup for the Hilbert manifold of H^1 -loops. In the Finsler case the energy functional is generally only C^1 , rather than C^2 , on the Hilbert loop space. Nevertheless the required Palais–Smale compactness and critical point theory for closed Finsler geodesics are standard; see Caponio–Javaloyes–Masiello [4], Mercuri [9], and Lu [8]. With these replacements, the broken-geodesic approximation, the compactness argument for end-incompressible classes, and the Fuller-index invariance argument go through verbatim.

Let g be a β -taut Riemannian metric. By definition, the space $S(g, \beta)$ of constant-speed closed g -geodesics in the class β is compact.

We first record a consequence of the indivisibility of β . The circle S^1 acts on $L_\beta X$ by reparametrization:

$$(s \cdot \gamma)(t) = \gamma(t + s).$$

If some $\gamma \in L_\beta X$ had nontrivial stabilizer, then γ would factor through a nontrivial covering $S^1 \rightarrow S^1$. Equivalently, γ would be a multiple cover of another loop. This would make the free homotopy class β a nontrivial power, contrary to indivisibility. Therefore the S^1 -action on $L_\beta X$ is free. Hence

$$H_*^{S^1}(L_\beta X; \mathbb{Z}) \cong H_*(L_\beta X/S^1; \mathbb{Z}).$$

Let

$$E_g(\gamma) = \int_{S^1} |\dot{\gamma}(t)|_g^2 dt$$

be the energy functional. Since $S(g, \beta)$ is compact, E_g is bounded above on $S(g, \beta)$. Choose

$$E_0 > \sup_{o \in S(g, \beta)} E_g(o).$$

Set

$$U_{E_0} = \{\gamma \in L_\beta X \mid E_g(\gamma) < E_0\}.$$

We claim that the inclusion

$$U_{E_0} \hookrightarrow L_\beta X$$

is a homotopy equivalence.

Let

$$L_\beta^{\text{cs}} X \subset L_\beta X$$

denote the subspace of constant-speed loops in the class β , and set

$$U_{E_0}^{\text{cs}} = \{\gamma \in L_\beta^{\text{cs}} X \mid E_g(\gamma) < E_0\}.$$

Since β is nonconstant, every loop in $L_\beta X$ has positive length. Thus every loop in $L_\beta X$ has a well-defined constant-speed reparametrization. If γ_{cs} denotes the constant-speed reparametrization of γ , then

$$E_g(\gamma_{\text{cs}}) = \ell_g(\gamma)^2 \leq E_g(\gamma),$$

by Cauchy–Schwarz. The standard reparametrization homotopy gives a deformation equivalence

$$L_\beta X \simeq L_\beta^{\text{cs}} X,$$

and, since constant-speed reparametrization is energy nonincreasing, the same argument applies to energy sublevels. Thus it is enough to show that

$$U_{E_0}^{\text{cs}} \hookrightarrow L_\beta^{\text{cs}} X$$

is a homotopy equivalence.

We prove this using the finite-dimensional broken-geodesic approximation. Fix $A > E_0$, and set

$$\Lambda_A^{\text{cs}} = \{\gamma \in L_\beta^{\text{cs}} X \mid E_g(\gamma) \leq A\}.$$

For $\gamma \in \Lambda_A^{\text{cs}}$, constant speed gives

$$E_g(\gamma) = \ell_g(\gamma)^2.$$

Hence

$$\ell_g(\gamma) \leq A^{1/2}.$$

Since $\beta \in \pi_1^{\text{inc}}(X)$, bounded length in the class β prevents loops from escaping to infinity. By the compactness Lemma 2.1 for end-incompressible classes, there exists a compact set

$$K_A \subset X$$

such that

$$\text{image}(\gamma) \subset K_A$$

for every $\gamma \in \Lambda_A^{\text{cs}}$.

For each $x \in K_A$, choose a strongly convex normal ball B_x . Since K_A is compact, finitely many such balls cover K_A . Let $\delta_A > 0$ be a Lebesgue number for this finite cover, and choose $0 < \rho_A < \delta_A$ small enough so that whenever $p, q \in K_A$ satisfy

$$d_g(p, q) < \rho_A,$$

the points p and q lie in a common strongly convex normal ball. Consequently, p and q are joined by a unique short minimizing g -geodesic depending smoothly on the endpoints, as long as the endpoints remain in that common convex ball.

Choose N so large that

$$\frac{A^{1/2}}{N} < \rho_A.$$

Let

$$t_i = i/N, \quad i = 0, \dots, N.$$

For $\gamma \in \Lambda_A^{\text{cs}}$, we have

$$\begin{aligned} d_g(\gamma(t_i), \gamma(t_{i+1})) &\leq \int_{t_i}^{t_{i+1}} |\dot{\gamma}(t)|_g dt \\ &= \frac{1}{N} \ell_g(\gamma) \\ &\leq \frac{A^{1/2}}{N} \\ &< \rho_A. \end{aligned}$$

Therefore consecutive vertices $\gamma(t_i)$ and $\gamma(t_{i+1})$ lie in a common strongly convex normal ball. We may replace each segment

$$\gamma|_{[t_i, t_{i+1}]}$$

by the unique short minimizing geodesic joining its endpoints.

Let $B_{A,N}$ denote the resulting space of broken geodesics in the class β . Equivalently, $B_{A,N}$ is parametrized by tuples

$$(x_0, \dots, x_{N-1}) \in K_A^N$$

such that consecutive pairs x_i, x_{i+1} , with $x_N = x_0$, lie in a common strongly convex normal ball and the corresponding broken geodesic represents β . The connecting geodesic segments need not lie in K_A ; they lie in the corresponding strongly convex normal balls. This is harmless, since the finite-dimensional parameters are the vertices.

The replacement map

$$r_{A,N} : \Lambda_A^{\text{cs}} \rightarrow B_{A,N}$$

is continuous. The usual straightening homotopy, performed on each interval $[t_i, t_{i+1}]$, deforms γ to $r_{A,N}(\gamma)$. Since each segment is replaced by the minimizing geodesic with the same endpoints, this straightening does not increase energy. Hence the inclusion

$$B_{A,N} \hookrightarrow \Lambda_A^{\text{cs}}$$

is a homotopy equivalence, for N sufficiently large.

On $B_{A,N}$, the restriction of the energy is a finite-dimensional function. Its critical points are precisely the closed g -geodesics in the class β with energy at most A . By the choice of E_0 , all closed class- β g -geodesics have energy strictly smaller than E_0 . Therefore this finite-dimensional energy function has no critical points in the region

$$\{\gamma \in B_{A,N} \mid E_0 \leq E_g(\gamma) \leq A\}.$$

By the finite-dimensional deformation lemma, the inclusion

$$B_{E_0,N} \hookrightarrow B_{A,N}$$

is a homotopy equivalence. Transporting this through the broken-geodesic approximation gives a homotopy equivalence

$$U_{E_0}^{\text{cs}} \hookrightarrow \Lambda_A^{\text{cs}}.$$

Since $A > E_0$ was arbitrary and

$$L_\beta^{\text{cs}} X = \bigcup_{A > E_0} \Lambda_A^{\text{cs}},$$

we conclude that

$$U_{E_0}^{\text{cs}} \simeq L_\beta^{\text{cs}} X.$$

Using the deformation equivalence

$$L_\beta X \simeq L_\beta^{\text{cs}} X$$

and the corresponding equivalence on sublevels, we get

$$U_{E_0} \simeq L_\beta X.$$

Since the energy is S^1 -invariant, U_{E_0} is S^1 -invariant. Because the S^1 -action on $L_\beta X$ is free, it follows that

$$U_{E_0}/S^1 \simeq L_\beta X/S^1.$$

We now perturb g to a β -regular metric g' , chosen sufficiently close to g and within a β -taut deformation. By invariance of the Fuller index under β -taut deformations,

$$F(g, \beta) = F(g', \beta).$$

Since g' is β -regular and β -taut, the class- β closed geodesic strings of g' form a compact zero-dimensional space, hence there are finitely many of them.

Let

$$E_{g'}(\gamma) = \int_{S^1} |\dot{\gamma}(t)|_{g'}^2 dt$$

be the energy associated to g' . Choose $E_0' > 0$ larger than the energy of every class- β closed g' -geodesic, and set

$$V = \{\gamma \in \mathcal{L}_\beta X \mid E_{g'}(\gamma) < E_0'\},$$

where $\mathcal{L}_\beta X$ is the Hilbert manifold of H^1 -loops in the class β . By the same broken-geodesic approximation argument applied to g' , the inclusion

$$V \hookrightarrow \mathcal{L}_\beta X$$

is a homotopy equivalence. Since $\mathcal{L}_\beta X$ is homotopy equivalent to $L_\beta X$, and since the S^1 -action is free, we have

$$V/S^1 \simeq L_\beta X/S^1.$$

We next use Morse theory on the Hilbert loop manifold. The energy functional $E_{g'}$ is smooth, S^1 -invariant, satisfies the Palais–Smale condition, and its negative gradient flow has the usual compactness properties on V . Indeed, if $\{\gamma_j\} \subset V$ is a Palais–Smale sequence, then the energy bound gives a uniform length bound. Since $\beta \in \pi_1^{\text{inc}}(X)$, the compactness lemma for end-incompressible classes implies that all images $\gamma_j(S^1)$ lie in one compact subset of X . On that compact subset, the standard Palais–Smale compactness theorem for the geodesic energy functional gives a strongly convergent subsequence in H^1 .

Since g' is β -regular, the critical set of $E_{g'}$ in V is a finite union of nondegenerate critical circles

$$C_o \simeq S^1,$$

one for each class- β g' -geodesic string o . The nondegeneracy is transverse to the natural S^1 -direction.

The negative gradient flow gives the usual Morse–Bott decomposition of V . Namely, V is obtained by attaching the unstable disk bundles of the critical circles C_o . The rank of the unstable disk bundle over C_o is the Morse–Bott index

$$\text{morse}(o).$$

Because β is indivisible, the S^1 -action on V is free. After passing to the quotient by S^1 , each critical circle C_o becomes a point, and its unstable disk bundle becomes a cell of dimension

$$\text{morse}(o).$$

Therefore V/S^1 has a finite CW decomposition with one k -cell for each class- β geodesic string o satisfying

$$\text{morse}(o) = k.$$

Thus

$$\chi(V/S^1) = \sum_o (-1)^{\text{morse}(o)},$$

where the sum is over all class- β g' -geodesic strings.

On the other hand, since g' is β -regular, the Fuller index formula gives

$$F(g', \beta) = \sum_o \frac{(-1)^{\text{morse}(o)}}{\text{mult}(o)}.$$

Because β is indivisible, every class- β geodesic is geometrically simple. Hence

$$\text{mult}(o) = 1$$

for every class- β string o , and therefore

$$F(g', \beta) = \sum_o (-1)^{\text{morse}(o)} = \chi(V/S^1).$$

Since $F(g, \beta) = F(g', \beta)$ and

$$V/S^1 \simeq L_\beta X/S^1,$$

we obtain

$$F(g, \beta) = \chi(L_\beta X/S^1).$$

Finally, because the S^1 -action on $L_\beta X$ is free,

$$\chi(L_\beta X/S^1) = \chi^{S^1}(L_\beta X).$$

Thus

$$F(g, \beta) = \chi^{S^1}(L_\beta X).$$

The same finite CW decomposition of V/S^1 shows that $H_*(V/S^1; \mathbb{Z})$ has finite total rank. Since

$$V/S^1 \simeq L_\beta X/S^1$$

and

$$H_*^{S^1}(L_\beta X; \mathbb{Z}) \cong H_*(L_\beta X/S^1; \mathbb{Z}),$$

it follows that $H_*^{S^1}(L_\beta X; \mathbb{Z})$ has finite total rank. This completes the proof. $\square$

12. PROOF OF THEOREM 1.31 AND ITS COROLLARIES

We first prove the following.

Theorem 12.1 (Product formula). *Let*

$$Z \hookrightarrow X \xrightarrow{p} Y$$

be a β -taut fibration, where $\beta \in \pi_1^{\text{inc}}(X)$ is a fiber class. Suppose that Y is connected and closed, and that every smooth closed contractible g_Y -geodesic in Y is constant.

Fix $y_0 \in Y$, set $Z_0 = p^{-1}(y_0)$, and choose a fiber class

$$\beta_Z \in \pi_1(Z_0)$$

such that

$$(i_{y_0})_*(\beta_Z) = \beta,$$

where $i_{y_0} : Z_0 \hookrightarrow X$ is the inclusion. Let H_{y_0} be the holonomy group of the fibration acting on $\pi_1^{\text{inc}}(Z_0)$, and set

$$S_{y_0} = H_{y_0} \cdot \beta_Z.$$

Then S_{y_0} is finite, and

$$F(g, \beta) = \text{card}(S_{y_0}) \chi(Y) F(g_{y_0}, \beta_Z).$$

Proof. We first show that every class- β g -geodesic in X is vertical.

Let

$$o : S^1 \rightarrow X$$

be a closed g -geodesic representing the class β . Since β is a fiber class, the loop

$$p \circ o : S^1 \rightarrow Y$$

is contractible. By the definition of a β -taut fibration, p projects g -geodesics to g_Y -geodesics. Hence $p \circ o$ is a smooth closed contractible g_Y -geodesic. By hypothesis, every such geodesic is constant. Therefore $p \circ o$ is constant, and so o is contained in a single fiber $Z_y = p^{-1}(y)$.

Thus the class- β geodesic strings in X are precisely vertical geodesic strings.

Let

$$C = S^*X$$

be the unit cotangent bundle of (X, g) , with projection

$$\text{pr} : C \rightarrow X.$$

Let R^{λ_g} denote the Reeb vector field on C , equivalently the unit-speed geodesic flow. Let $\tilde{\beta} \in \pi_1(C)$ be the canonical lift of β . We want to compute:

$$F(g, \beta) = i(O(R^{\lambda_g}, \tilde{\beta}), R^{\lambda_g}, \tilde{\beta}).$$

The idea is to localize contributions over critical points of a Morse function on Y .

Let

$$P^{\text{vert}} : TX \rightarrow T^{\text{vert}}X = \ker(dp)$$

be the g -orthogonal projection. By assumptions on $p : X \rightarrow Y$,

$$\nabla P^{\text{vert}} = 0,$$

where P^{vert} is viewed as a $(1, 1)$ -tensor field.

For $\xi \in C$, let

$$v_\xi \in T_{\text{pr}(\xi)}X$$

be the g -dual unit vector to ξ . Define

$$P : C \rightarrow \mathbb{R}$$

by

$$P(\xi) = |P^{\text{vert}}v_\xi|_g^2.$$

Thus $P = 1$ on vertical unit covectors and $P = 0$ on horizontal unit covectors.

Let $f : Y \rightarrow \mathbb{R}$ be a Morse function. Define

$$\tilde{f} : C \rightarrow \mathbb{R}$$

by

$$\tilde{f} = f \circ p \circ \text{pr}.$$

Equip $C = S^*X$ with the Sasaki metric g_S induced by g . Thus

$$TC = T^{\text{vert}}C \oplus T^{\text{hor}}C$$

is an orthogonal splitting, where

$$T^{\text{vert}}C = \ker(\text{pr}_*)$$

and $T^{\text{hor}}C$ is the horizontal subbundle induced by the Levi-Civita connection of g .

We shall use the following two elementary identities.

First,

$$(12.2) \quad dP(R^{\lambda_g}) = 0.$$

Indeed, let $\xi(\tau)$ be a flow line of R^{λ_g} , and set

$$x(\tau) = \text{pr}(\xi(\tau)).$$

Then $x(\tau)$ is a unit-speed g -geodesic. Let

$$v(\tau) = \dot{x}(\tau) = v_{\xi(\tau)}$$

be its velocity vector field. Since $x(\tau)$ is a geodesic,

$$\nabla_\tau v = 0.$$

Since $\nabla P^{\text{vert}} = 0$, we have

$$\nabla_\tau(P^{\text{vert}}v) = P^{\text{vert}}(\nabla_\tau v) = 0.$$

Therefore

$$\begin{aligned} \frac{d}{d\tau}P(\xi(\tau)) &= \frac{d}{d\tau}|P^{\text{vert}}v(\tau)|_g^2 \\ &= 2\langle \nabla_\tau(P^{\text{vert}}v), P^{\text{vert}}v \rangle_g \\ &= 0. \end{aligned}$$

Hence $dP(R^{\lambda_g}) = 0$.

Second,

$$(12.3) \quad dP(\nabla_{g_S}\tilde{f}) = 0.$$

Since $\tilde{f} = f \circ p \circ \text{pr}$ depends only on the base point $\text{pr}(\xi)$:

$$\nabla_{g_S}\tilde{f} \in T^{\text{hor}}C.$$

More precisely, $\nabla_{g_S}\tilde{f}$ is the horizontal lift of

$$\nabla_g(f \circ p)$$

on X . Let $\xi(s)$ be a horizontal curve in C , and set

$$x(s) = \text{pr}(\xi(s)).$$

By definition of the Levi-Civita horizontal distribution on S^*X , the covectors $\xi(s)$ are parallel along $x(s)$. Equivalently, their g -dual vectors

$$v(s) = v_{\xi(s)}$$

are parallel along $x(s)$:

$$\nabla_s v(s) = 0.$$

Using again $\nabla P^{\text{vert}} = 0$, we get

$$\nabla_s(P^{\text{vert}}v(s)) = P^{\text{vert}}(\nabla_s v(s)) = 0.$$

Hence

$$\begin{aligned} \frac{d}{ds}P(\xi(s)) &= \frac{d}{ds}|P^{\text{vert}}v(s)|_g^2 \\ &= 2\langle \nabla_s(P^{\text{vert}}v(s)), P^{\text{vert}}v(s) \rangle_g \\ &= 0. \end{aligned}$$

Taking $\dot{\xi}(0) = \nabla_{g_S}\tilde{f}(\xi)$ gives

$$dP_\xi(\nabla_{g_S}\tilde{f}) = 0.$$

Since ξ was arbitrary,

$$dP(\nabla_{g_S}\tilde{f}) = 0.$$

Now set

$$V_\varepsilon = R^{\lambda_g} - \varepsilon \nabla_{g_S}(P + \tilde{f}),$$

for $\varepsilon > 0$ sufficiently small. Since the class- β orbit space of R^{λ_g} is compact by β -tautness, Fuller continuation gives a compact continued orbit set N_ε and

$$i(O(R^{\lambda_g}, \tilde{\beta}), R^{\lambda_g}, \tilde{\beta}) = i(N_\varepsilon, V_\varepsilon, \tilde{\beta}).$$

We now identify the closed orbits in N_ε . We show that P is a Lyapunov function for V_ε . Mainly, along a trajectory $\xi(\tau)$ of V_ε , we have

$$\begin{aligned} \frac{d}{d\tau} P(\xi(\tau)) &= dP(R^{\lambda_g}) - \varepsilon dP(\nabla_{g_S} P) - \varepsilon dP(\nabla_{g_S} \tilde{f}) \\ &= -\varepsilon |\nabla_{g_S} P|_{g_S}^2 \leq 0. \end{aligned}$$

Thus P is nonincreasing along the V_ε -flow.

If $\xi(\tau)$ is a closed orbit of V_ε , then $P(\xi(\tau))$ returns to its initial value after one period. Since P is nonincreasing, it must be constant along the orbit. Hence

$$\nabla_{g_S} P = 0$$

along the orbit.

On each unit cotangent sphere, P is the squared norm of the vertical component of the corresponding unit vector. Therefore the critical locus of P consists of the two branches

$$P = 1 \quad \text{and} \quad P = 0,$$

corresponding respectively to vertical and horizontal covectors. At $\varepsilon = 0$, the class- $\tilde{\beta}$ orbit set under consideration is vertical, hence lies in $P = 1$. For $\varepsilon > 0$ sufficiently small, the continued set N_ε remains in the continuation of this vertical component. Thus the horizontal branch $P = 0$ is excluded, and every closed orbit in N_ε lies in the vertical branch $P = 1$.

Consequently, every closed orbit in N_ε projects to a geodesic contained in a fiber Z_y .

It remains to see over which fibers these vertical orbits occur. On the vertical branch $P = 1$, the Reeb vector field restricts to the geodesic Reeb field of the fiber (Z_y, g_y) , while $\tilde{f}$ depends only on the base point y . The perturbation

$$-\varepsilon \nabla_{g_S} \tilde{f}$$

moves the base point in the negative gradient direction of f . Therefore a vertical closed orbit of V_ε can occur only over a critical point of f . Conversely, if $y \in \text{Crit}(f)$, then

$$\nabla_{g_S} \tilde{f} = 0$$

over Z_y , and the vertical part of V_ε is just the fiber geodesic flow, up to the harmless P -term which vanishes on the vertical critical branch. Thus the closed orbits of V_ε in N_ε are precisely the vertical closed geodesic orbits lying in fibers over critical points of f .

We now compute the index. For $y \in Y$, let

$$H_y \subset \text{Aut } \pi_1(Z_y)$$

be the holonomy group of the fibration at y . Choose

$$\beta_Z \in \pi_1(Z_y)$$

with

$$(i_y)_*(\beta_Z) = \beta.$$

Set

$$S_y = H_y \cdot \beta_Z.$$

This is the holonomy orbit of the fiber representative β_Z . Since representatives of distinct free homotopy classes lie in distinct connected components of the fiber loop space, and since the relevant vertical geodesic locus is compact, the set S_y is finite.

Moreover, $F(g_y, \cdot)$ is constant on S_y . Indeed, let

$$\gamma : [0, 1] \rightarrow Y$$

be a path based at y . Pulling back $p : X \rightarrow Y$ along γ gives a fibration over $[0, 1]$. By the definition of a β -taut fibration, this pullback is a taut deformation of the fiber metric from (Z_y, g_y) back to itself, with final fiber identified with the initial one by the holonomy map $h_\gamma : Z_y \rightarrow Z_y$. Fuller index invariance under taut deformation gives

$$F(g_y, \alpha) = F(g_y, h_{\gamma,*}\alpha).$$

Thus $F(g_y, \alpha)$ is constant on the holonomy orbit S_y . Hence

$$\sum_{\alpha \in S_y} F(g_y, \alpha) = \text{card}(S_y) F(g_y, \beta_Z).$$

If $y, y' \in Y$, holonomy along any path from y to y' identifies S_y with $S_{y'}$. The same path gives a taut deformation between the fiber metrics, so

$$\text{card}(S_y) F(g_y, \beta_Z)$$

is independent of y . We denote this common fiber contribution by

$$A = \text{card}(S_{y_0}) F(g_{y_0}, \beta_Z).$$

Near a critical point $y \in \text{Crit}(f)$, the linearized return map of the perturbed flow splits into a vertical part and a base Morse part. The vertical part is the Fuller index contribution of the fiber geodesic flow over the classes in S_y . The base part is the local index of the negative gradient vector field $-\nabla f$ at y , namely

$$(-1)^{\text{morse}(y)}.$$

Therefore

$$i(N_{\varepsilon,y}, V_\varepsilon, \tilde{\beta}) = (-1)^{\text{morse}(y)} A,$$

where $N_{\varepsilon,y}$ denotes the part of N_ε lying over the critical point y .

Summing over the critical points of f , we obtain

$$\begin{aligned} F(g, \beta) &= i(N_\varepsilon, V_\varepsilon, \tilde{\beta}) \\ &= \sum_{y \in \text{Crit}(f)} i(N_{\varepsilon,y}, V_\varepsilon, \tilde{\beta}) \\ &= \sum_{y \in \text{Crit}(f)} (-1)^{\text{morse}(y)} A \\ &= \chi(Y) A. \end{aligned}$$

Substituting the definition of A , we get

$$F(g, \beta) = \text{card}(S_{y_0}) \chi(Y) F(g_{y_0}, \beta_Z).$$

This proves the theorem. $\square$

We return to the proof of Theorem 1.31. The first part immediately follows from the second, as any class $\beta \in \pi_1^{\text{inc}}(X)$ geodesic strings of a complete negatively curved Riemannian manifold X are unique. We prove second part. Suppose first that β is an n -power, so that β has a representative with multiplicity n , and we let α denote the corresponding indivisible class. By the assumption that all contractible g_Y geodesics are constant, classical Morse theory Milnor [10] tells us that Y has vanishing higher homotopy groups $\pi_k(Y, y_0)$, $k \geq 2$. And in particular $i_{Z,*} : \pi_1(Z, p_0) \rightarrow \pi_1(X, p_0)$ is a group injection, by the long exact sequence of a fibration. It follows that $\alpha \in \pi_1^{\text{inc}}(X)$ is also a fiber class.

Now, any α -class g -geodesic string must be contained in a fiber of p . For otherwise we may find a β class g -geodesic string, which is not contained in a fiber of p , which would contradict verticality in the first part of the proof of the product formula. It readily follows that $p : X \rightarrow Y$ is also α -taut.

Now, if $\chi(Y) \neq \pm 1$ then by the product formula $F(g, \alpha) \neq 1$, since $F(g_{y_0}, \alpha_Z)$ is an integer by Theorem 1.25. By Theorem 1.25

$$F(g, \alpha) = \chi^{S^1}(L_\alpha X).$$

So if X admits a complete metric with a unique and non-degenerate class α g -geodesic string then we have:

$$F(g, \alpha) = \chi^{S^1}(L_\alpha X) = 1$$

(see Proof of Theorem 1.25), which is impossible. It follows that X does not admit a metric with a unique and non-degenerate class β g -geodesic string. For if o, o' are distinct, non-degenerate, class α g -geodesics strings, then the n -fold covers $o^n, (o')^n$ are class β , distinct non-degenerate g -geodesic strings.

We now prove the general case. Suppose by contradiction that X admits a metric with a unique and non-degenerate class β g -geodesic string o . By the above, β is not atomic. Now o covers a multiplicity one geodesic string $\tilde{o}$ in some class $\tilde{\beta} \in \pi_1^{inc}(X)$. Moreover, $\tilde{o}$ is the unique geodesic string in its class, otherwise o would not be unique in its class. We prove that $\tilde{\beta}$ is indivisible, which will be a contradiction to β not being atomic and will complete the proof.

Suppose otherwise, so that $\tilde{\beta}_{x_0} = \alpha^k$ for $k > 1$ and $\alpha \in \pi_1(X, x_0)$, (β_{x_0} is as in Definition 1.23). Let u be a class α , closed g -geodesic string (where α also denotes the class in $\pi_1^{inc}(X)$ corresponding to the based class α .) It is immediate that the k cover of u , u^k represents $\tilde{\beta}$ and is a g -geodesic string. By the uniqueness, $\tilde{o} = u^k$. But this contradicts simplicity of $\tilde{o}$. So $\tilde{\beta}$ is indivisible.

□

Proof of Theorem 1.6. Let g_Z, g_Y be complete Riemannian metrics on Z respectively Y with non-positive curvature. Take the product metric $g = g_Z \times g_Y$ on $X = Z \times Y$. For a class $\beta \in \pi_1^{inc}(X)$ in the image of the inclusion $\pi_1^{inc}(Z) \rightarrow \pi_1^{inc}(X)$, the natural projection $X \rightarrow Y$ is automatically a β -taut fibration. Then the conclusion readily follows from Theorem 1.31.

□

Proof of Theorem 1.27. By Theorem 12.1 0 is certainly a value of the invariant F . We first prove that every negative rational number is the value of the invariant. Let p, q be positive integers. Let Y be a closed surface of genus $(p+1) > 1$ with a hyperbolic metric g_Y , let Z be the genus 2 closed surface with a hyperbolic metric g_Z and let $\beta_Z \in \pi_1^{inc}(Z)$ be the class represented by a $2 \cdot q$ -fold covering of a simple closed loop representing a generator of the fundamental group of Z .

Let $X = Y \times Z$ with the product metric $g = g_Y \times g_Z$ and $p : X \rightarrow Y$ the canonical projection. By Theorem 12.1

$$F(g, \beta) = \chi(Y) \cdot F(g_Z, \beta_Z) = (-2p) \cdot \frac{1}{2q} = -\frac{p}{q},$$

where β is as in Lemma 1.33. So we proved our first claim.

Let again p, q be positive integers. Let Y be closed surface of genus 2, with a hyperbolic metric g_Y . And let Z be a manifold satisfying $F(g_Z, \beta_Z) = -\frac{p}{2q}$ for some β_Z -taut metric g_Z on Z and for some class $\beta_Z \in \pi_1^{inc}(Z)$. This exist by the discussion above. Let $g = g_Y \times g_Z$ be the product metric on $Y \times Z$, and β as above. Analogously to the discussion above we get:

$$F(g, \beta) = \chi(Y) \cdot F(g_Z, \beta_Z) = (-2) \cdot \frac{-p}{2q} = \frac{p}{q}.$$

□

A. FULLER INDEX AND SKY CATASTROPHES

Let X be a complete vector field without zeros on a manifold M . Set

$$(A.1) \quad S(X, \beta) = \{o \in L_\beta M \mid \exists p \in (0, \infty), \ o : \mathbb{R}/\mathbb{Z} \rightarrow M \text{ is a periodic orbit of } pX\}.$$

The above p is uniquely determined and we denote it by $p(o)$ called the period of o .

There is a natural S^1 reparametrization action on $S(X, \beta)$: $t \cdot o$ is the loop $t \cdot o(\tau) = o(t + \tau)$. The elements of $\mathcal{O}(X, \beta) := S(X, \beta)/S^1$ will be called **orbit strings**. Slightly abusing notation we just write o for the equivalence class of o .

The multiplicity $m(o)$ of an orbit string is the ratio $p(o)/l$ for $l > 0$ the period of a simple orbit string covered by o .

We want a kind of fixed point index which counts orbit strings o with certain weights. Assume for simplicity that $N \subset \mathcal{O}(X, \beta)$ is finite. (Otherwise, for a general open compact $N \subset \mathcal{O}(X, \beta)$, we need to perturb.) Then to such an (N, X, β) Fuller associates an index:

$$i(N, X, \beta) = \sum_{o \in N} \frac{1}{m(o)} i(o),$$

where $i(o)$ is the fixed point index of the time $p(o)$ return map of the flow of X with respect to a local surface of section in M transverse to the image of o .

Fuller then shows that $i(N, X, \beta)$ has the following invariance property. For a continuous homotopy $\{X_t\}$, $t \in [0, 1]$ set

$$S(\{X_t\}, \beta) = \{(o, t) \in L_\beta M \times [0, 1] \mid o \in S(X_t)\}.$$

And given a continuous homotopy $\{X_t\}$, $X_0 = X$, $t \in [0, 1]$, suppose that $\tilde{N}$ is an open compact subset of

$$(A.2) \quad \mathcal{O}(\{X_t\}, \beta) := S(\{X_t\}, \beta)/S^1,$$

such that

$$\tilde{N} \cap (L_\beta M \times \{0\})/S^1 = N.$$

Then if

$$N_1 = \tilde{N} \cap (L_\beta M \times \{1\})/S^1$$

we have

$$(A.3) \quad i(N, X, \beta) = i(N_1, X_1, \beta).$$

We call this **basic invariance**.

Given an open compact $N_0 \subset \mathcal{O}(X_0, \beta)$, and X_1 sufficiently C^0 nearby X_0 , there is a distinguished open compact $N_1 \subset \mathcal{O}(X_1, \beta)$ such that

$$(A.4) \quad i(N_0, X_0, \beta) = i(N_1, X_1, \beta).$$

See for instance [12, Proof of Lemma 1.6]). The set N_1 will be called the **continuation of N_0** .

A.1. Blue sky catastrophes. We gave a preliminary definition in Section 1.1.2. Here is a more encompassing definition.

Definition A.5. Let $\{X_t\}$, $t \in [0, 1]$ be a continuous family of non-zero, smooth vector fields on a closed manifold M . We say that $\{X_t\}$ has a **catastrophe in class β** , if there is an element

$$y \in \mathcal{O}(X_0, \beta) \sqcup \mathcal{O}(X_1, \beta) \subset \mathcal{O}(\{X_t\}, \beta)$$

such that there is no open compact subset of $\mathcal{O}(\{X_t\}, \beta)$ containing y .

13. ACKNOWLEDGEMENTS

I am grateful to the IAS for resources provided during my stay as a member, where some of this paper was written.

REFERENCES

- [1] M. T. ANDERSON, *Metrics of negative curvature on vector bundles*, Proc. Am. Math. Soc., 99 (1987), pp. 357–363.
- [2] D. BAO, S.-S. CHERN, AND Z. SHEN, *An introduction to Riemann-Finsler geometry*, vol. 200 of Grad. Texts Math., New York, NY: Springer, 2000.
- [3] D. BURAGO, Y. BURAGO, AND S. IVANOV, *A course in metric geometry*, vol. 33 of Grad. Stud. Math., Providence, RI: American Mathematical Society (AMS), 2001.
- [4] E. CAPONIO, M. A. JAVALOYES, AND A. MASIELLO, *On the energy functional on finsler manifolds and applications to stationary spacetimes*, Mathematische Annalen, 351 (2011), pp. 365–392.
- [5] P. EBERLEIN AND B. O’NEILL, *Visibility manifolds*, Pac. J. Math., 46 (1973), pp. 45–109.
- [6] F. FULLER, *Note on trajectories in a solid torus.*, Ann. Math. (2), 56 (1952), pp. 438–439.
- [7] ———, *An index of fixed point type for periodic orbits.*, Am. J. Math., 89 (1967), pp. 133–145.
- [8] G. LU, *Methods of infinite dimensional morse theory for geodesics on finsler manifolds*, 2012.
- [9] F. MERCURI, *The critical points theory for the closed geodesics problem*, Mathematische Zeitschrift, 156 (1977), pp. 231–246.
- [10] J. MILNOR, *Morse theory*, Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, No. 51, Princeton University Press, Princeton, N.J., 1963.
- [11] A. PREISSMANN, *Quelques propriétés globales des espaces de Riemann*, Commentarii Mathematici Helvetici, (1942).
- [12] Y. SAVELYEV, *Extended Fuller index, sky catastrophes and the Seifert conjecture*, International Journal of mathematics, 29 (2018).
- [13] W. P. THURSTON, *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Am. Math. Soc., New Ser., 19 (1988), pp. 417–431.

Email address: `yasha.savelyev@gmail.com`

FACULTY OF SCIENCE, UNIVERSITY OF COLIMA, MEXICO