

# GEODESIC STRING COUNTING INVARIANTS OF MANIFOLDS

YASHA SAVELYEV

**ABSTRACT.** We study rational valued geodesic-string (equivalence class of a closed geodesic) counts for complete Riemann-Finsler manifolds based on the Fuller index of the geodesic flow. This can be understood as a shadow of a conjectural orbifold Morse homology theory of the infinite dimensional quotient stack  $[LX/S^1]$ . One basic new phenomenon appears out of this. Let  $g$  be a regular Finsler metric on  $T^n$ ,  $n > 1$ , with finitely many class  $\beta$  geodesic strings. Suppose that  $g$  is  $\beta$ -taut homotopic to the flat metric (meaning that the  $g_t$  length of  $\beta$ -class  $g_t$  geodesics is uniformly bounded from above.) Fix a prime  $p$ , and suppose that there is one class- $\beta$   $g$ -geodesic string with multiplicity divisible by  $p$ , then there is another one. This theorem has interesting examples even on  $T^2$ . As the main technical ingredient we prove a fibration/product formula for the geodesic-string count. This also gives new arithmetic constraints on counts of closed geodesics in mapping tori and flat bundles, and constraints on existence of negative sectional curvature metrics.

## 1. INTRODUCTION

Using counts of *geodesic strings* (equivalence classes of closed, constant speed geodesics up to reparametrization  $S^1$  action), and the Fuller index [11], we will define certain rational number valued, deformation invariants for complete Riemann-Finsler manifolds. A basic case is a complete Riemannian manifold with nonpositive sectional curvature, and in this case we get deformation invariants of the metric, provided the deformation is through nonpositive sectional curvature metrics. A detailed outline of Fuller's theory and basic definitions are given here in the Appendix, so no prior knowledge is necessary.

Suppose for the moment  $X$  is compact. For a nonconstant, free homotopy class  $\beta$  of a loop in  $X$ , we say that a metric  $g$  on  $X$  is  $\beta$ -**regular** if all of its class  $\beta$  closed geodesics are nondegenerate in the usual sense, or equivalently the closed orbits of the associated geodesic flow are dynamically nondegenerate. For a  $\beta$ -regular  $g$ , with finitely many class  $\beta$  geodesic strings we have the following formula for the  $F$ -count:

$$(1.1) \quad F(g, \beta) = \sum_o \frac{(-1)^{\text{morse}(o)}}{\text{mult}(o)} \in \mathbb{Q},$$

where:

- The sum is over class  $\beta$   $g$ -geodesic strings  $o$ .
- $\text{morse}(o)$  denotes the Morse index of  $o$ , (meaning the Morse-Bott index of the associated critical submanifold (diffeomorphic to  $S^1$ ) of the loop space).
- $\text{mult}(o)$  is the geometric multiplicity, equivalently the order of the corresponding isotropy subgroup of  $S^1$ , where  $S^1$  is acting on the loop space by reparametrization.

For a free homotopy class  $\beta$  of loops in  $X$ , let  $L_\beta X$  denote the corresponding component of the free loop space. The following is a direct corollary of Theorem 1.23, and exemplifies the basic topological invariance of the  $F$ -count:

---

*Key words and phrases.* Nonpositive curvature, Fuller index, geodesic counts,  $S^1$ -equivariant homology of the loop space.

Supported by CONAHCYT research grant CF-2023-I.

**Corollary 1.2.** *Let  $X, g, \beta$  be as above and such that  $\beta$  is indivisible, (cannot be represented by a multiply covered loop). Then the  $S^1$ -equivariant homology  $H_*^{S^1}(L_\beta X, \mathbb{Z})$  has finite total rank. Denote by  $\chi^{S^1}(L_\beta X)$  the Euler characteristic of this homology. Then*

$$F(g, \beta) = \sum_o (-1)^{\text{morse}(o)} = \chi^{S^1}(L_\beta X),$$

(note  $\text{mult}(o) = 1$  by the indivisible condition.)

The proof of the above is based on  $S^1$ -equivariant Morse theory. For divisible classes, the homology theoretic interpretation of the  $F$ -count would require more than ordinary  $S^1$ -equivariant Morse homology for the  $g$ -energy functional. One would want a  $g$ -invariant up to quasi-isomorphism “orbifold Morse homology chain complex” for the quotient stack  $[LX/S^1]$ , together with (assuming certain finiteness) an algebraically derived  $\mathbb{Q}$ -valued characteristic coinciding with the  $F$ -count. This is likely related to Wall-Euler characteristic theory in group cohomology theory, which also appears in the more basic Conjecture 1. Existence of this orbifold Morse homology is at the moment conjectural.

We now present one new phenomenon, which uses a product formula for the  $F$ -count in the form of Theorem 8.1. For a compact  $X$ , we say a metric  $g$  is taut if the space of class  $\beta$  closed  $g$ -geodesics is compact for every  $\beta$ . (With the topology induced by the  $C^1$  topology on the loop space.) It is said to be  $\beta$ -taut if the same holds for just  $\beta$ -class geodesics. A homotopy  $\{g_t\}_{t \in [0,1]}$  is said to be  $\beta$ -taut if the space of pairs  $(o, t)$  with  $o$  class  $\beta$   $g_t$ -geodesic is compact (equivalent to the condition in the abstract). An example of a metric  $g$ ,  $\beta$ -taut homotopic to the flat metric is in Figure 1. We can construct such examples which are arbitrarily “far” from being flat. Figure 1 is also an example for the following.

**Theorem 1.3** (Proof in Section 5). *Let  $g$  be a  $\beta$ -regular Riemann-Finsler metric on  $T^n$ ,  $n > 1$ ,  $\beta$ -taut homotopic to the flat metric for  $\beta \in \pi_1(T^n)$  a nontrivial class. Fix prime  $p$ , if there is one  $g$ -geodesic string  $o$  s.t.:*

- (1)  $o$  has class  $\beta$ .
- (2)  $p \mid \text{mult}(o)$ .

*then there is another one.*

In Section 5 we will give a stronger form of this result which applies to general Riemann-Finsler metrics nearby to the flat metric.

It should be noted, there are no examples of  $\beta$ -taut metrics on  $T^n$  not  $\beta$ -taut homotopic to the flat metric.

*Remark 1.4.* In dimension 2, Hamilton’s normalized Ricci flow [12] converges smoothly to a flat metric, and a KAM confinement argument (which is far beyond present techniques) shows that every metric sufficiently  $C^k$ -close to a flat metric is  $\beta$ -taut homotopic to it; the question thus reduces to  $\beta$ -tautness of the flow path on a compact time interval, which we cannot presently verify.

This result is not accessible to  $S^1$ -equivariant Morse homology techniques, as the latter doesn’t suitably track isotropy information. Nevertheless, the arithmetic constraints are reminiscent of the resonance identities in closed-geodesic and closed-characteristic theory, where finite collections of periodic objects satisfy identities involving average Euler characteristics and mean indices. See, for example, [23], [14] and the later noncontractible resonance identities [16].

We will show that the count  $F(g, \beta) \in \mathbb{Q}$  can be defined for any  $\beta$ -taut  $g$  (Section 4). What follows is a special, more elaborate case of the Conjecture 2 (appearing further ahead) and can be understood as a divisible analogue of Corollary 1.2 in the aspherical case.




FIGURE 1. Schematic of a  $\beta$ -taut torus metric with mixed geodesic multiplicities in the class  $\beta = 4[x]$ . In coordinates

$$T^2 = \mathbb{R}^2 / \mathbb{Z}^2, \quad g_{\lambda, \epsilon} = dy^2 + \lambda^2 f_\epsilon(y)^2 dx^2, \quad f_\epsilon(y) = 1 + \epsilon \cos(4\pi y),$$

the  $x$ -direction is drawn as the meridian direction. The four critical levels of  $f_\epsilon$ ,

$$y = 0, \quad \frac{1}{4}, \quad \frac{1}{2}, \quad \frac{3}{4},$$

give four meridian geodesics; their 4-covers are class- $\beta$  geodesic strings of multiplicity 4. For a resonant choice of  $\lambda > (4\sqrt{\epsilon(1+\epsilon)})^{-1}$ , there is also a non-horizontal oscillating closed geodesic in the same class  $\beta$ , of multiplicity 2. This is in part classical Clairaut theory, see for instance [1], [8, §4-4, Example 5]. This latter geodesic comes in an  $S^1$ -family; after a small generic perturbation, this family breaks into cancelling (for the  $F$ -count) isolated strings. In particular, we can see how Theorem 1.3 is satisfied in this example taking  $p = 2$ . To see the  $\beta$ -taut homotopy to the flat metric take

$$g_s = dy^2 + \lambda^2 (1 + s\epsilon \cos(4\pi y))^2 dx^2, \quad s : 1 \rightarrow 0,$$

each  $g_s$  is Clairaut-integrable and tautness is then established by explicit tracking of geodesics.

**Conjecture 1** (Aspherical topological formula for  $F$ ). *Let  $X$  be a closed aspherical manifold, and let  $\beta \neq 1$  be a free homotopy class. Suppose that  $X$  admits a  $\beta$ -taut metric. Then the Borel construction*

$$(L_\beta X)_{hS^1} := L_\beta X \times_{S^1} ES^1$$

*is Wall-Euler finite, and for every  $\beta$ -taut metric  $g$ ,*

$$F(g, \beta) = \chi_{\text{Wall}}((L_\beta X)_{hS^1}).$$

*Equivalently, if  $G = \pi_1 X$ , if  $\beta = [\gamma_\beta]$  (the conjugacy class of  $\gamma_\beta \in G$ ), and if*

$$Q_\beta = C_G(\gamma_\beta) / \langle \gamma_\beta \rangle,$$

*then  $Q_\beta$  is Wall-Euler finite and*

$$F(g, \beta) = \chi_{\text{Wall}}(Q_\beta).$$

The conjecture implies that  $F(g, \beta)$  is independent of the  $\beta$ -taut metric  $g$ , for closed aspherical manifolds. For background on rational Euler characteristics of groups, we refer to Wall's original note [27]

and to Brown's treatment of Euler characteristics of groups in [6, Chapter IX]; see also Brown [5] for Euler characteristics of discrete groups and  $G$ -spaces.

Assuming the conjecture we get a generalization of Theorem 5.1.

**Theorem 1.5** (Proof in Section 6). *Assume the above conjecture, let  $\beta \in \pi_1(T^n)$ ,  $n > 1$ , be a nontrivial class and let  $g$  be a  $\beta$ -regular Finsler metric on  $T^n$  having finitely many  $\beta$  class geodesic strings. Suppose that  $o$  is a  $g$ -geodesic string s.t.:*

- (1)  $o$  has class  $\beta$ .
- (2)  $p \mid \text{mult}(o)$ , where  $p$  is prime,

*then there is at least one other such  $o$  (in particular same  $p$ ).*

The following is rather important open question in our context:

*Question 1.* Suppose that  $X$  is closed, does  $\beta$ -taut for some  $\beta$  imply aspherical?

*Remark 1.6.* A possible route to a counterexample is a mapping torus of a non-aspherical  $Z$  by a diffeomorphism with “hyperbolic behavior”. In the other direction, a closed aspherical manifold may admit no  $\beta$ -taut metric. By results of Mess [19] and Belegradek [4, Corollary 5.1], in every dimension  $n \geq 4$  there are closed aspherical  $n$ -manifolds whose fundamental groups contain nontrivial infinitely divisible elements. Then for such an infinitely divisible  $\beta$ , the manifold cannot have a  $\beta$ -taut metric. Here is the proof. Take any  $g$  and let  $l > 0$  be the systole of  $g$ , that is

$$l = \inf_{\{o \mid o \text{ is noncontractible}\}} \ell_g(o).$$

Then clearly if  $\beta$  is infinitely divisible, there are  $\beta$  class  $g$ -geodesics of length at least  $n \cdot l$  for arbitrarily large  $n \in \mathbb{N}$ .

**1.1. Geodesics in mapping tori.** A theorem of Preissmann [22] says that there are no negative sectional curvature metrics on compact products. For fibrations, the most obvious analogue of Preissmann's theorem fails even assuming compactness. By Thurston's fibered hyperbolization theorem, a closed surface bundle over  $S^1$  with fiber of negative Euler characteristic is hyperbolic if and only if its monodromy is pseudo-Anosov; see [26, 25]. What follows is a kind of partial, “only if” Finsler analogue in higher dimensions, noting that an isometry of a manifold has “anti-Anosov behavior” (for some reasonable generalization of pseudo-Anosov to higher dimensions).

We denote by  $\pi_1^{\text{inc}}(X)$  the set of free homotopy classes of loops incompressible to the ends, in the sense of Definition 1.8, (by convention nonconstant classes when  $X$  is closed).

**Corollary 1.7** (Corollary of Corollary 1.34). *Let  $X_\phi$  be the mapping torus of an isometry  $\phi : Z \rightarrow Z$  of a complete taut Riemannian metric on  $Z$ , with  $Z$  finite type, and where  $\beta_Z \in \pi_1^{\text{inc}}(Z)$  is indivisible. Let  $\beta \in \pi_1^{\text{inc}}(X_\phi)$  denote the image of  $\beta_Z$  under inclusion. Then:*

- (1)  $\chi^{S^1}(L_\beta X_\phi) = 0$ .
- (2)  $X_\phi$  does not admit a forward complete,  $\beta$ -regular Riemann–Finsler metric with an odd number of class  $\beta$  geodesic strings.
- (3)  $X_\phi$  does not admit a complete negative sectional curvature Riemannian metric.

*Example 1.* The corollary applies to the mapping torus of a finite type surface of genus at least one (the genus condition allows us to find a nontrivial class in  $\pi_1^{\text{inc}}(X_\phi)$ , we also need a suitable taut metric e.g. a nonpositive curvature metric, with an isometry). The conclusion that  $X_\phi$  does not admit a complete

negative sectional curvature metric, follows from Thurston's classification<sup>1</sup>. However the arithmetic constraints on geodesic counts seem interesting even for a surface map.

If we further assume that  $Z$  is aspherical, the first part of Corollary 1.7 can be proved via group cohomology theory, for which we already provided some references. But the main point is that in our setup the theorem has an elementary proof, once the general product formula of Theorem 8.1 for the  $F$ -count is established. The proof of the latter does involve a tricky perturbation, but Fuller index theory allows us to do this abstractly, perturbing a vector field rather than the metric/energy functional.

**1.2. Setup and more statements.** Proofs of many results here are postponed till subsequent sections.

**Terminology 1.** From now on, all our metrics are Riemann-Finsler (a.k.a. Finsler) metrics unless specified to be Riemannian, and usually denoted by just  $g$ . Completeness, always means forward completeness, and it is an assumption for all our metrics. Curvature always means sectional curvature in the Riemannian case and flag curvature in the Finsler case. Thus we will usually just say complete metric  $g$ , for a forward complete Riemann-Finsler metric. A reader may certainly choose to interpret all metrics as Riemannian metrics, completeness as standard completeness, and curvature as sectional curvature.

In what follows,  $\pi_k(X)$  denotes the usual based homotopy group, with basepoints suppressed when no confusion can arise. We do not use  $\pi_1(X)$  to denote free homotopy classes.

**Definition 1.8.** Let  $X$  be a smooth manifold. Fix an exhaustion by nested compact sets  $\bigcup_{i \in \mathbb{N}} K_i = X$ ,  $K_i \supset K_{i-1}^\circ$  for all  $i \geq 1$ . A free homotopy class  $\beta$  of loops in  $X$  is called **end compressible** if, for every  $i$ , it has a representative contained in  $X - K_i$ . We say that  $\beta$  is **end incompressible** (or incompressible to the ends) if it is not end compressible.

Let  $\pi_1^{inc}(X)$  denote the set of such end incompressible free homotopy classes. When  $X$  is compact, we set  $\pi_1^{inc}(X)$  to be the set of nonconstant free homotopy classes of loops in  $X$ .

It is easily seen that the above is well defined (independent of the choice of an exhaustion) and moreover any homeomorphism  $X_1 \rightarrow X_2$  of a pair of manifolds induces a set isomorphism  $\pi_1^{inc}(X_1) \rightarrow \pi_1^{inc}(X_2)$ .

Denote by  $L_\beta X$  the class  $\beta \in \pi_1^{inc}(X)$  component of the free loop space of  $X$ , with its compact open topology. Let  $g$  be a complete metric on  $X$ , and let  $S(g, \beta) \subset L_\beta X$  denote the subspace of all constant speed parametrized, smooth, closed  $g$ -geodesics in class  $\beta$ .

**Definition 1.9.** We say that a metric  $g$  on  $X$  is  **$\beta$ -taut** if it is complete and  $S(g, \beta)$  is compact. We will say that  $g$  is **taut** if it is  $\beta$ -taut for each  $\beta \in \pi_1^{inc}(X)$ .

**Lemma 1.10.** *A complete metric  $g$  with nonpositive curvature satisfies:*

- *All of its closed geodesics are minimizing in their free homotopy class.*
- *It is taut.*

*Proof.* The first part is a standard consequence of the Cartan-Hadamard theorem. The second part follows by the first part and Corollary 2.3.  $\square$

---

<sup>1</sup>Suitably adjusted, one may also need finite geometry (volume) assumptions.

**Definition 1.11.** Let  $\beta \in \pi_1^{inc}(X)$ , and let  $g_0, g_1$  be a pair of  $\beta$ -taut metrics on  $X$ . A  $\beta$ -**taut deformation** between  $g_0, g_1$ , is a continuous (in the topology of  $C^0$  convergence on compact sets) family  $\{g_t\}$ ,  $t \in [0, 1]$  of complete metrics on  $X$ , s.t.

$$S(\{g_t\}, \beta) := \{(o, t) \in L_\beta X \times [0, 1] \mid o \in S(g_t, \beta)\}$$

is compact. We say that  $\{g_t\}$  is a **taut deformation** if it is  $\beta$ -taut for each  $\beta \in \pi_1^{inc}(X)$ . The above definitions of tautness are extended naturally to the case of a smooth fibration  $X \hookrightarrow P \rightarrow [0, 1]$ , with a smooth fiber-wise family of metrics.

A useful criterion for  $\beta$ -tautness is the following.

**Theorem 1.12** (Proof in Section 2.1). *Let  $\beta \in \pi_1^{inc}(X)$ . Let  $\{g_t\}_{t \in [0, 1]}$  be a continuous family of complete metrics on  $X$ . Suppose that:*

$$\sup_t \left| \sup_{o \in S(g_t, \beta)} \ell_{g_t}(o) - \inf_{o \in S(g_t, \beta)} \ell_{g_t}(o) \right| < \infty,$$

where  $\ell_{g_t}$  is the length functional with respect to  $g_t$ , then  $\{g_t\}$  is  $\beta$ -taut.

For example, the hypothesis is trivially satisfied if  $g_t$  have the property that all their class  $\beta$  closed geodesics are minimizing in their homotopy class.

**Corollary 1.13.** *If  $g_t$ ,  $t \in [0, 1]$  have nonpositive curvature then  $\{g_t\}$  is taut.*

*Proof.* This follows by Theorem 1.12 and by Lemma 1.10. □

1.2.1. *The geodesic string counting invariant  $F$ .* Let  $\mathcal{G}(X)$  be the set of equivalence classes of taut metrics  $g$ , where  $g_0$  is equivalent to  $g_1$  whenever there is a taut deformation between them. We may denote an equivalence class by its representative  $g$  by a slight abuse of notation. The following is proved in Section 4.

**Theorem 1.14.** *For each manifold  $X$  there is a natural, nontrivial functional:*

$$F : \mathcal{G}(X) \times \pi_1^{inc}(X) \rightarrow \mathbb{Q},$$

extending the definition in (1.1).

For  $g$   $\beta$ -regular and  $\beta$ -taut we have already defined  $F(g, \beta)$  at the beginning of the Introduction. For a general  $\beta$ -taut  $g$  we define  $F(g, \beta)$  in terms of the Fuller index.

**Corollary 1.15.** *Suppose for a pair  $g_0, g_1$  of  $\beta$ -taut metrics on  $X$ :*

$$F(g_0, \beta) \neq F(g_1, \beta),$$

then any path  $\{g_t\}$ , connecting  $g_0, g_1$ , is not  $\beta$ -taut.

*Proof.* The fact that any connecting  $\{g_t\}$  is not  $\beta$ -taut is just a direct corollary of the theorem above. □

I was unable to find such a pair  $g_0, g_1$ , in fact there is strong evidence to believe the following:

**Conjecture 2.**  *$F$  does not depend on the choice of a smooth structure and taut metric. In other words we have the following. Let  $X$  be a topological manifold, define:*

$$\mathcal{S}(X) : \pi_1^{inc}(X) \rightarrow \mathbb{Q} \sqcup \{\infty\}$$

by:

$$\mathcal{S}(X)(\beta) = \begin{cases} \infty, & \text{if } X \text{ does not admit a smooth structure and a } \beta\text{-taut Finsler metric.} \\ F(g, \beta), & \text{if } g \text{ is a } \beta\text{-taut Finsler metric on } X. \end{cases},$$

then  $\mathcal{S}$  is well-defined and hence determines a topological invariant of topological manifolds. So if  $f : X_1 \rightarrow X_2$  is a homeomorphism then  $\mathcal{S}(X_1)(\beta) = \mathcal{S}(X_2)(f_*(\beta))$ .

Theorem 1.23 in the following section partially proves this conjecture.

*Remark 1.16.* The smooth manifold version of this conjecture, i.e. that  $\mathcal{S}(X)$  is a smooth manifold invariant, is implied by the Reeb sky catastrophe conjecture described in the next section. However, the above seems to be much more basic as will be apparent from the proof of Theorem 1.23.

### 1.2.2. A short digression on sky catastrophes.

**Definition 1.17** (Preliminary). A *sky catastrophe* for a smooth family  $\{X_t\}$ ,  $t \in [0, 1]$ , of nonvanishing vector fields on a closed manifold  $M$  is a continuous family of closed orbit strings  $\tau \mapsto o_{t_\tau}$ ,  $o_{t_\tau}$  is an orbit string of  $X_{t_\tau}$ ,  $\tau \in [0, \infty)$ , such that the period of  $o_{t_\tau}$  is unbounded from above.

Sky catastrophes first appeared in the work of Fuller [10]. One general definition appears in [24], which we review in the Appendix C, the main point of this definition is that there are no regularity assumptions on  $\{X_t\}$ . The preliminary definition becomes equivalent to the more general definition given certain regularity conditions on the family  $\{X_t\}$ .

The following corollary of Theorem 1.12 may be of independent interest.

**Corollary 1.18.** *Sky catastrophes of vector fields on closed manifolds are not geodesible by metrics all of whose geodesics are minimizing. That is, if a family  $\{X_t\}$  has a sky catastrophe, there is no family  $\{g_t\}$  of metrics (all of whose geodesics are minimizing in their homotopy class) such that the orbits of  $X_t$  are unit speed parametrized  $g_t$ -geodesics.*

Fuller at the end of [11] has asked for any metric conditions on vector fields to rule out sky catastrophes. By the above, nonpositivity of curvature is one such condition. So this is a partial answer to his question.

The main conjecture in [24] is that geodesible, and more generally Reeb sky catastrophes ( $\{X_t\}$  are Reeb) are unstable, in the sense that there is a  $C^0$  nearby family  $\{X'_t\}$  which does not have a sky catastrophe (at least in the fixed class  $\beta$ ). Let's call this **Reeb sky catastrophe conjecture**. The reason to conjecture such a thing, is that a theorem in [24] says that if  $\{o_{t_\tau}\}_\tau$  is a sky catastrophe for a family of Reeb vector fields  $\{X_t\}$  then the length of the corresponding projection  $\tau \mapsto t_\tau$  is infinite. (Assuming some minor regularity on  $\{o_{t_\tau}\}$ .) The latter suggests that the structure of such a sky catastrophe is very fragile.

*Remark 1.19.* The Reeb sky catastrophe conjecture implies the geodesible Seifert conjecture (apparently still an open problem), by the main result of [24]. Hence, this is a subtle question.

The following is a stronger form Corollary 1.15.

**Theorem 1.20.** *Suppose for a pair  $g_0, g_1$  of  $\beta$ -taut metrics on a closed manifold  $M$ :*

$$F(g_0, \beta) \neq F(g_1, \beta),$$

*then any path  $\{g_t\}$ , connecting  $g_0, g_1$ , has a sky catastrophe, meaning that the corresponding family  $\{X_t\}$  of Reeb vector fields on the unit cotangent bundle (cf. Section 4) has a sky catastrophe as in Definition C.1.*



*Proof.* This directly follows by [24, Theorem 3.2].  $\square$

So that if such a pair  $g_0, g_1$  exists, the Reeb sky catastrophe conjecture is disproved and in a rather extreme fashion.

### 1.2.3. Basic results on the invariant $F$ .

**Definition 1.21.** Let  $\beta$  be a free homotopy class of loops in  $X$ . Choose a representative of  $\beta$  and a base point  $x_0$  on its image. This choice determines an element

$$\beta_{x_0} \in \pi_1(X, x_0)$$

well-defined up to inner automorphism.

- An element  $\gamma \in \pi_1(X, x_0)$  is **indivisible** if whenever  $\gamma = \alpha^k$  for some  $\alpha \in \pi_1(X, x_0)$  and  $k > 0$ , then  $k = 1$ .
- An element  $\gamma \in \pi_1(X, x_0)$  is a  **$k$ -power** if  $\gamma = \alpha^k$ ,  $k > 1$ , for some  $\alpha$  which is not an  $n$ -power for any  $n > 1$ .
- An element  $\gamma \in \pi_1(X, x_0)$  is **atomic** if  $\gamma = \alpha^k$ , for some  $k \geq 1$ , with  $\alpha$  indivisible.

We say that the free homotopy class  $\beta$  is indivisible, respectively a  $k$ -power, respectively atomic, if  $\beta_{x_0}$  has the corresponding property for one, equivalently any, such choice of  $x_0$  and representative. This is well-defined because the corresponding notions are invariant under inner automorphism. Equivalently,  $\beta$  is a  $k$ -power precisely when it admits a smooth representative with multiplicity  $k$  whose underlying simple representative is indivisible.

Note that a class  $\beta \in \pi_1^{inc}(X)$  is indivisible iff any representative of this class is not multiply covered.

*Example 2.* Let  $g$  be a Riemannian metric with negative sectional curvature on a closed manifold  $X$  and  $\beta \in \pi_1^{inc}(X)$  a class represented by a multiplicity  $n$  closed geodesic, then

$$(1.22) \quad F(g, \beta) = \frac{1}{n}.$$

In particular, if  $\beta$  is indivisible then  $F(g, \beta) = 1$ . More generally, (1.22) holds whenever  $g$  has a unique and nondegenerate geodesic string in class  $\beta$ , where nondegenerate is as in the Introduction.

If  $\beta \in \pi_1^{inc}(X)$  is indivisible, then it is easy to see that the reparametrization  $S^1$  action on  $L_\beta X$  is free (Section 7), so that  $H_*^{S^1}(L_\beta X, \mathbb{Z}) \simeq H_*(L_\beta X/S^1, \mathbb{Z})$ , where  $H_*^{S^1}(L_\beta X, \mathbb{Z})$  denotes the  $S^1$ -equivariant homology. Moreover, we have the following result proved in Section 7.

**Theorem 1.23.** Suppose that  $\beta \in \pi_1^{inc}(X)$  is indivisible, and  $X$  admits a  $\beta$ -taut metric. Then  $H_*^{S^1}(L_\beta X, \mathbb{Z})$  has finite total rank. Denote by  $\chi^{S^1}(L_\beta X)$  the Euler characteristic of this homology. Then for any  $\beta$ -taut metric  $g$  on  $X$ :

$$F(g, \beta) = \chi^{S^1}(L_\beta X).$$

Consequently,  $F(g, \beta)$  is independent of the choice of  $\beta$ -taut metric  $g$ , for indivisible  $\beta$ . Furthermore, Conjecture 2 holds on the subset of classes  $\beta$  which are indivisible.

At least for surfaces  $\beta$ -taut metrics (distinct from nonpositive curvature metrics) can be constructed explicitly, recall the example of Figure 1.

We can use the above, and the product formula of Theorem 8.1 to get:

**Theorem 1.24** (Proof in Section 9). Every rational number has the form  $F(g, \beta)$  for some  $\beta$ -taut Riemannian  $g$  on some compact manifold  $X$  and for some  $\beta$ .



1.2.4. *Applications to existence of negative curvature metrics.* We are going to give a certain generalization of Preissmann's theorem to fibrations, with possibly noncompact fibers, also replacing the negative sectional curvature condition by a significantly weaker condition. Though the definitions may at first seem hard to understand, we just want enough generality to incorporate interesting “flat bundle” examples (with anti-Anosov or periodic holonomy behavior). The latter are of interest in 3-manifold theory, for instance vis-à-vis Thurston's classification [25].

**Definition 1.25.** For a fiber bundle  $Z \hookrightarrow X \rightarrow Y$ , we say that  $\beta \in \pi_1^{\text{inc}}(X)$  is a **fiber class** if it lies in the image of the set map

$$i_Z : \pi_1^{\text{inc}}(Z) \rightarrow \pi_1^{\text{inc}}(X)$$

induced by inclusion of a fiber.

**Lemma 1.26.** *Let*

$$Z \hookrightarrow X \xrightarrow{p} Y$$

*be a smooth fiber bundle with connected fiber and connected base. Assume*

$$\pi_2(Y) = 0.$$

*Fix  $y_0 \in Y$ , and let*

$$H = \pi_1(Y, y_0)$$

*act on  $\pi_1^{\text{inc}}(Z_{y_0})$  by holonomy. Let  $\beta \in \pi_1^{\text{inc}}(X)$  be a fiber class, and define*

$$\mathcal{R}_{y_0}(\beta) = \{ \alpha \in \pi_1^{\text{inc}}(Z_{y_0}) \mid (i_{y_0})_*(\alpha) = \beta \}.$$

*Then  $\mathcal{R}_{y_0}(\beta)$  is a single  $H$ -orbit. In particular, if*

$$\beta_Z, \beta'_Z \in \mathcal{R}_{y_0}(\beta),$$

*then*

$$H \cdot \beta_Z = H \cdot \beta'_Z.$$

*Thus the holonomy orbit associated to the fiber class  $\beta$  is independent of the choice of fiber representative  $\beta_Z$ .*

*Proof.* Choose basepoints  $z_0 \in Z_{y_0}$  and  $x_0 = i_{y_0}(z_0)$ . Since  $\pi_2(Y) = 0$ , the homotopy long exact sequence gives an exact sequence of based groups

$$1 \rightarrow \pi_1(Z_{y_0}, z_0) \xrightarrow{(i_{y_0})_*} \pi_1(X, x_0) \xrightarrow{p_*} \pi_1(Y, y_0) \rightarrow 1.$$

Thus we identify  $\pi_1(Z_{y_0}, z_0)$  with a normal subgroup of  $\pi_1(X, x_0)$ , and the quotient is  $\pi_1(Y, y_0)$ .

Let  $\alpha, \alpha' \in \mathcal{R}_{y_0}(\beta)$ . Choose based representatives of these fiber free homotopy classes. Since they determine the same total free homotopy class  $\beta$  in  $X$ , the corresponding elements of  $\pi_1(X, x_0)$  are conjugate. Thus, after conjugating inside  $\pi_1(Z_{y_0}, z_0)$  if necessary, there exists  $x \in \pi_1(X, x_0)$  such that

$$\alpha' = x\alpha x^{-1}.$$

Let

$$h = p_*(x) \in H.$$

Conjugation by  $x$  on the normal subgroup  $\pi_1(Z_{y_0}, z_0)$  induces the holonomy action of  $h$ , up to an inner automorphism of  $\pi_1(Z_{y_0}, z_0)$ . Passing to free homotopy classes in the fiber, we get

$$\alpha' = h \cdot \alpha.$$

Thus all fiber representatives of the total free homotopy class  $\beta$  belong to a single holonomy orbit.

Conversely, if  $\alpha' = h \cdot \alpha$ , choose  $x \in \pi_1(X, x_0)$  with  $p_*(x) = h$ . Then  $x\alpha x^{-1}$  represents the same total free homotopy class in  $X$  as  $\alpha$ , and its fiber free homotopy class is  $h \cdot \alpha$ . Hence the holonomy orbit is exactly the set of fiber representatives of  $\beta$ .  $\square$

We assume from now on that  $Z, Y$  are connected. For a fiber class  $\beta \in \pi_1^{\text{inc}}(X)$ , set

$$\mathcal{R}(\beta) = \{\alpha \in \pi_1^{\text{inc}}(Z) \mid (i_Z)_*(\alpha) = \beta\}.$$

Assume that  $\pi_2(Y) = 0$ , and put

$$H = \pi_1(Y, y_0).$$

By Lemma 1.26, the holonomy orbit

$$S = H \cdot \beta_Z$$

is independent of the choice of the fiber representative  $\beta_Z \in \mathcal{R}(\beta)$ . If another fiber inclusion is chosen, the corresponding holonomy actions are abstractly conjugate.

**Definition 1.27.** Let

$$Z \hookrightarrow X \xrightarrow{p} Y$$

be a smooth fiber bundle, with  $Y$  and  $\pi_2(Y) = 0$ , and let  $\beta \in \pi_1^{\text{inc}}(X)$  be a fiber class. We say that  $p$  is  **$\beta$ -holonomy-finite** if for one, equivalently any,  $\beta_Z \in \mathcal{R}(\beta)$ , the orbit

$$S = \pi_1(Y, y_0) \cdot \beta_Z$$

is finite.

**Definition 1.28.** Let

$$Z \hookrightarrow X \xrightarrow{p} Y$$

be a smooth fiber bundle, and let  $g$  be a Riemannian metric on  $X$ . Let

$$T^{\text{vert}}X = \ker(dp)$$

and let

$$\pi^{\text{vert}} : TX \rightarrow T^{\text{vert}}X$$

be the  $g$ -orthogonal projection. We say that  $p$  is  **$g$ -parallel** if

$$\nabla \pi^{\text{vert}} = 0.$$

Equivalently,

$$\nabla_A(\pi^{\text{vert}}B) = \pi^{\text{vert}}(\nabla_A B)$$

for all vector fields  $A, B$  on  $X$ .

**Definition 1.29.** Let  $\beta \in \pi_1^{\text{inc}}(X)$  be a fiber class, for a smooth fibration

$$p : Z \hookrightarrow (X, g) \rightarrow (Y, g_Y).$$

We say that this is a  **$\beta$ -periodic fibration** if:

- (1)  $g$  is a complete Riemannian metric.
- (2)  $Z, Y$  are connected and  $\pi_2(Y) = 0$ .
- (3)  $p$  is  $g$ -parallel and is a Riemannian submersion.
- (4)  $p$  is  $\beta$ -holonomy-finite.
- (5) the fiber metrics  $g_y$  on  $Z_y$  are taut.

The metrics  $g, g_Y$  may be kept implicit. In the above definition of a  $\beta$ -periodic fibration and the hypothesis of the following theorem we need the auxiliary metric  $g$  on  $X$  to be Riemannian, and there is no obvious extension of the theorem to the Riemann-Finsler case. However, the conclusions of the theorem are for Riemann-Finsler metrics. The following is perhaps the main technical result of the paper, proved in Section 9.

**Theorem 1.30.** *Let  $p : Z \hookrightarrow (X, g) \rightarrow (Y, g_Y)$  be a  $\beta$ -periodic fibration, where  $\beta \in \pi_1^{\text{inc}}(X)$  is a fiber class. Suppose further that  $Y$  is connected, closed,  $\chi(Y) \neq \pm 1$  and is such that all smooth closed contractible  $g_Y$ -geodesics in  $Y$  are constant. Then the following holds:*

- $X$  does not admit a complete Riemannian metric with negative curvature.
- $X$  does not admit a forward complete Riemann-Finsler metric with a unique nondegenerate class- $\beta$  geodesic string.

Note that  $\chi(Y) \neq 1$  is of course essential, as the trivial fibration  $X \rightarrow \{pt\}$ , with  $X$  admitting a complete negatively curved metric, will satisfy the hypothesis. The condition that there is a fiber class  $\beta \in \pi_1^{\text{inc}}(X)$  is also essential; as any vector bundle over a manifold admitting a Riemannian metric of negative curvature also admits a metric of negative curvature, Anderson [2].

**Corollary 1.31** (Proof in Section 9). *Let  $Y, Z$  be smooth manifolds, let  $\beta_Z \in \pi_1^{\text{inc}}(Z)$ , and assume that  $\chi(Y) \neq \pm 1$ . Suppose that  $Y$  is closed and admits a metric all of whose contractible geodesics are constant, for instance a nonpositive curvature Riemannian metric, and suppose that  $Z$  admits a taut metric. Let*

$$X = Y \times Z$$

with projection  $X \rightarrow Y$ , let  $i : Z \rightarrow X$  be a fiber inclusion, and set

$$\beta = i_*(\beta_Z).$$

Then:

- $X$  does not admit a complete Riemannian metric of negative sectional curvature.
- $X$  does not admit a forward complete Riemann-Finsler metric with a unique and nondegenerate class- $\beta$  geodesic string.

*Example 3.* Take  $Y$  any nonpositively curved closed manifold and  $Z$  a surface of genus at least one, possibly infinite type.

This theorem is very close to being sharp, for example the conclusion of the corollary is false if  $Y = S^1$  and  $Z = S^1 \times \mathbb{R}$ . As  $X = T^2 \times \mathbb{R}$  admits the warped product metric  $g_{\mathbb{R}} \times_{e^t} g_{T^2}$  (with respect to the function  $f = e^t$  on  $\mathbb{R}$ ) where  $g_{T^2}, g_{\mathbb{R}}$  are the flat metrics. This warped product has constant negative sectional curvature  $-1$ . So it is essential that not only  $\pi_1(Z)$  be nontrivial but also that there is a class incompressible to the ends. Of course,  $\chi(Y) \neq 1$  is also obviously essential, otherwise we may take  $Y = pt$ . The condition  $\chi(Y) \neq -1$  is however not obviously essential.

A further basic set of examples for the theorem is obtained by starting with a Riemannian manifold  $(Y, g_Y)$ , and any homomorphism

$$(1.32) \quad \phi : \pi_1(Y, y_0) \rightarrow \text{Isom}(Z, g_Z), \quad (\text{the group of all isometries}).$$

where  $g_Z$  is a taut Riemannian metric and  $\beta_Z \in \pi_1^{\text{inc}}(Z)$ . (For example  $(Z, g_Z)$  is a non-simply connected complete hyperbolic surface of genus at least one). Suppose further:

- (1) The orbit

$$\text{Orb}(\beta_Z) = \{\phi_*(\gamma)(\beta_Z) \mid \gamma \in \pi_1(Y, y_0)\}$$

is finite.

- (2)  $Y$  is closed and connected.

- (3) All contractible smooth closed  $g_Y$  geodesics in  $Y$  are constant.

We have the obvious induced diagonal action

$$\pi_1(Y, y_0) \rightarrow \text{Diff}(Z \times \tilde{Y}), \text{ (the group of all diffeomorphisms),}$$

$$\gamma \mapsto ((z, y) \mapsto (\phi(\gamma)(z), \gamma \cdot y)),$$

for  $\tilde{Y}$  the universal cover of  $Y$ . Taking the quotient of  $Z \times \tilde{Y}$  by this action, we get an associated “flat” bundle  $Z \hookrightarrow X_\phi \xrightarrow{p} Y$ , with a metric  $g_\phi$  induced from the product metric  $\tilde{g} = g_Z \oplus g_Y$ , on the covering space  $q : Z \times \tilde{Y} \rightarrow Z \times Y$ . The natural projection  $p : X_\phi \rightarrow Y$  is then a Riemannian submersion and is a  $g_\phi$ -parallel. The condition on the finiteness of  $\text{Orb}$  implies that  $p$  is  $\beta$ -holonomy finite where  $\beta$  is the image by  $(i_Z)_* : \pi_1^{\text{inc}}(Z) \rightarrow \pi_1^{\text{inc}}(X_\phi)$  of  $\beta_Z$ . It follows that  $p$  is a  $\beta$ -periodic fibration. Hence  $p : (X_\phi, g_\phi) \rightarrow (Y, g_Y)$  satisfies the hypothesis of Theorem 1.30.

*Example 4.* Suppose we have  $\beta_Z \in \pi_1^{\text{inc}}(Z)$ , and let  $\phi : Z \rightarrow Z$  be an isometry of a taut metric  $g_Z$ , such that  $\text{Orb}$  is finite. Then for  $\beta = (i_Z)_*(\beta_Z)$ , by the construction above, the mapping torus

$$(Z, g_Z) \hookrightarrow (X_\phi, g_\phi) \xrightarrow{\pi} S^1$$

has the structure of a  $\beta$ -periodic fibration, satisfying the hypothesis of Theorem 1.30.

The example can be made more concrete as follows.

**Lemma 1.33.** *Let  $Z$  be a complete Riemannian manifold of finite type. Then every isometry of  $Z$  has finite orbits on  $\pi_1^{\text{inc}}(Z)$ .*

*Proof.* We first prove that, for every  $L > 0$ , only finitely many classes in  $\pi_1^{\text{inc}}(Z)$  have representatives of length at most  $L$ . Choose a compact core

$$K_0 \subset Z$$

such that  $Z - K_0$  is a finite union of collar ends. Since  $Z$  is complete, the closed  $L$ -neighborhood

$$K_L = N_L(K_0)$$

is compact. If a representative of an end-incompressible class lies entirely in  $Z - K_0$ , then it lies in a collar end and can be pushed arbitrarily far out, contradicting end-incompressibility. Therefore every representative of an end-incompressible class must meet  $K_0$ . If such a representative has length at most  $L$ , then its image lies in  $K_L = N_L(K_0)$ .

It remains to use compactness. A compact Riemannian manifold, or compact domain after a harmless thickening of  $K_L$ , contains only finitely many free homotopy classes with representatives of length at most  $L$ . For instance, choose a finite good cover with Lebesgue number  $\delta > 0$ . A loop of length at most  $L$  is encoded by an edge loop in the nerve of the cover, of combinatorial length bounded by  $N(L, \delta)$ , and there are only finitely many such combinatorial loops. Hence there are only finitely many end-incompressible classes represented by loops of length at most  $L$ .

Now let  $\phi : Z \rightarrow Z$  be an isometry and let

$$\beta_Z \in \pi_1^{\text{inc}}(Z).$$

Choose a loop  $c$  representing  $\beta_Z$ , and set

$$L = \ell(c).$$

Then  $\phi^k(c)$  represents  $\phi_*^k(\beta_Z)$  and has the same length  $L$  for every  $k \in \mathbb{Z}$ . Since  $\phi$  preserves end-incompressibility, the orbit

$$\{\phi_*^k(\beta_Z) \mid k \in \mathbb{Z}\}$$

is contained in the finite set of end-incompressible classes represented by loops of length at most  $L$ . Therefore the orbit is finite.  $\square$

**Corollary 1.34.** *Let*

$$(Z, g_Z) \hookrightarrow (X_\phi, g_\phi) \rightarrow S^1$$

*be the mapping torus of an isometry of  $Z, g_Z$ , with  $Z$  finite type and  $g_Z$  taut. Let  $\beta \in \pi_1^{\text{inc}}(X_\phi)$  be the image of some  $\beta_Z \in \pi_1^{\text{inc}}(Z)$  under fiber inclusion, as previously.*

- (1)  $X_\phi$  does not admit a complete Riemannian metric with negative curvature.
- (2)  $X_\phi$  does not admit a forward complete Riemann-Finsler metric with a unique nondegenerate class- $\beta$  geodesic string.
- (3) If  $\beta$  is indivisible,  $\chi^{S^1}(L_\beta X_\phi) = 0$  and so  $X_\phi$  does not admit a  $\beta$ -regular forward complete Finsler metric with an odd number of  $\beta$ -class geodesic strings.

The following questions seem to be open.

*Question 2.* Is the finite type assumption necessary for Corollary 1.34 in the case of surfaces? The answer is likely no, as the intuition of Nielsen-Thurston classification (it cannot be directly applied in the infinite type setting) suggests that  $\phi$  should have pseudo-Anosov type behavior which an isometry even on an infinite type surface should not have.

## 2. COMPACTNESS PRELIMINARIES

**Lemma 2.1.** *Suppose that  $g$  is a complete metric on  $X$ ,  $\beta \in \pi_1^{\text{inc}}(X)$  and let  $S \subset L_\beta X$  be a subset on which the  $g$ -length functional is bounded from above. Then the images in  $X$  of elements of  $S$  are contained in a fixed compact subset of  $X$ .*

*Proof.* Suppose otherwise. Fix an exhaustion by nested compact sets

$$\bigcup_{i \in \mathbb{N}} K_i = X, \quad K_i^\circ \supset K_{i-1}.$$

Then either there is sequence  $\{o_i\}_{i \in \mathbb{N}}$ ,  $o_i \in S$  s.t.  $o_i \in K_i^c$ , for  $K_i^c$  the complement of  $K_i$ , which contradicts the fact that  $\beta$  is end incompressible. Or there is a sequence  $\{o_k\}_{k \in \mathbb{N}}$ ,  $o_k \in S$  s.t.:

- (1) Each  $o_k$  intersects  $K_{i_0}$  for some  $i_0$  fixed.
- (2) For each  $i \in \mathbb{N}$  there is a  $k_i > i$  s.t.  $o_{k_i}$  is not contained in  $K_i$ .

Now if  $\text{diam}(o_k)$  is bounded in  $k$ , then condition 1 implies that  $o_k$  are contained in a set of bounded diameter. (Here  $\text{diam}(o_k)$  denotes the diameter of image  $o_k$ .) Consequently, by Hopf-Rinow theorem [3],  $o_k$  are contained in a compact set. But this contradicts condition 2, and the fact that  $K_i$  form an exhaustion of  $X$ .

Thus, we conclude that  $\text{diam}(o_k)$  is unbounded, but this contradicts the hypothesis.  $\square$

**Lemma 2.2.** *Let  $g$  be a complete metric on  $X$ , and let  $\beta \in \pi_1^{\text{inc}}(X)$ . Denote by  $S_A(g, \beta) \subset S(g, \beta)$  the subset of elements with  $g$ -length at most  $A$ . Then  $S_A(g, \beta) \subset S(g, \beta)$  is compact.*

*Proof.* By Lemma 2.1, the images of elements of  $S_A(g, \beta) \subset S(g, \beta)$  are contained in compact  $K$ . Since elements of  $S_A(g, \beta) \subset S(g, \beta)$  are parametrized by constant speed, and since we have a length bound  $S_A(g, \beta)$  is an equicontinuous collection. The result then follows by the Arzelà-Ascoli theorem and standard arguments.  $\square$

**Corollary 2.3.** *Suppose that  $g$  is a complete metric on  $X$ , all of whose class  $\beta \in \pi_1^{\text{inc}}(X)$  geodesics are minimizing, then  $g$  is  $\beta$ -taut.*

**Lemma 2.4.** *Let  $g$  be a complete metric on  $X$ , and let  $\beta \in \pi_1^{\text{inc}}(X)$ . Then there exists a closed  $g$ -geodesic minimizing length among all loops in the free homotopy class  $\beta$ .*

*Proof.* Suppose otherwise and let

$$A = \inf_{\gamma \in L_\beta X} \ell_g(\gamma).$$

By the hypothesis that there is no geodesic with length  $A$ , and by Lemma 2.2, there is an  $\epsilon > 0$  such that there is no closed geodesic  $o$  with  $A \leq \ell_g(o) < A + \epsilon$ .

By Lemma 2.1 the subset  $M_{\beta,A} \subset L_\beta X$  consisting of elements with length less than  $A + \epsilon$  is contained in a compact  $K$ . Pick  $o \in M_{\beta,A}$  and apply the curve shortening process to it. Let  $o_k$ ,  $k \in \mathbb{N}$  be the stages of this process. Then  $\text{image } o_k \subset K$ . In particular  $\{o_k\}$  must converge to a geodesic, for instance [20], (we can also argue via Arzelà–Ascoli to prove convergence) but this contradicts the condition on  $\epsilon$ .  $\square$

**2.1. Proof of Theorem 1.12.** Let  $\{g_t\}$ ,  $t \in [0, 1]$  be as in the hypothesis, with

$$(2.5) \quad \sup_t \left| \sup_{o \in S(g_t, \beta)} \ell_{g_t}(o) - \inf_{o \in S(g_t, \beta)} \ell_{g_t}(o) \right| < C,$$

and suppose that

$$\sup_{(o,t) \in S(\{g_t\}, \beta)} \ell_{g_t}(o) = \infty.$$

Then we have a sequence  $\{o_k\}$ ,  $k \in \mathbb{N}$ , of closed class  $\beta$   $g_{t_k}$ -geodesics in  $X$ , satisfying:

- (1)  $\lim_{k \rightarrow \infty} t_k = t_\infty \in [0, 1]$ .
- (2)  $\lim_{k \rightarrow \infty} \ell_{g_{t_k}}(o_k) = \infty$ , where  $\ell_{g_{t_k}}(o_{t_k})$  is the length with respect to  $g_{t_k}$ .

Applying Lemma 2.4, we may choose a length-minimizing closed  $g_{t_\infty}$ -geodesic  $o_\infty$  in the class  $\beta$ . And let  $L$  denote its  $g_{t_\infty}$ -length. Let  $g_{aux}$  be a fixed auxiliary metric on  $X$ , and let  $L_{aux}$  be the  $g_{aux}$  length of  $o_\infty$ .

Define a pseudo-metric  $d_{C^0}$  on the space of metrics on  $X$  as follows. Set  $K = \text{image } o_\infty$ . And set

$$V = \{v \in TX \mid \pi(v) \in K \text{ for } \pi : TX \rightarrow X \text{ the canonical projection, and } |v|_{aux} = 1\},$$

where  $|v|_{aux}$  is the norm taken with respect to  $g_{aux}$ .

Then define:

$$d_{C^0}(g_1, g_2) = \sup_{v \in V} ||v|_{g_1} - |v|_{g_2}|.$$

By Properties 1 and 2, for every  $\epsilon > 0$  we may find a  $k > 0$  such that:

$$(2.6) \quad d_{C^0}(g_{t_k}, g_{t_\infty}) < \epsilon$$

and

$$(2.7) \quad \ell_{g_{t_k}}(o_k) > C + L + L_{aux} \cdot \epsilon.$$

By (2.6), we have:

$$\ell_{g_{t_k}}(o_\infty) < \ell_{g_{t_\infty}}(o_\infty) + L_{aux} \cdot \epsilon = L + L_{aux} \cdot \epsilon.$$

Combining with (2.7) we get:

$$\ell_{g_{t_k}}(o_k) > \ell_{g_{t_k}}(o_\infty) + C.$$

Since we may find a closed  $g_{t_k}$ -geodesic  $o'$  satisfying  $\ell_{g_{t_k}}(o') \leq \ell_{g_{t_k}}(o_\infty)$ , we get that

$$\left| \max_{o \in S(g_{t_k}, \beta)} \ell_{g_{t_k}}(o) - \min_{o \in S(g_{t_k}, \beta)} \ell_{g_{t_k}}(o) \right| > C,$$

and so we are in contradiction.

Thus,

$$\sup_{(o,t) \in \mathcal{O}(\{g_t\}, \beta)} \ell_{g_t}(o) < \infty.$$

Then compactness of  $\mathcal{S}(\{g_t\}, \beta)$  follows by a slight modification of the proof of Lemma 2.2.  $\square$

### 3. PRELIMINARIES ON REEB FLOW

Let  $(C^{2n+1}, \lambda)$  be a contact manifold with  $\lambda$  a contact form, that is a one form s.t.  $\lambda \wedge (d\lambda)^n \neq 0$ . Denote by  $R^\lambda$  the Reeb vector field satisfying:

$$d\lambda(R^\lambda, \cdot) = 0, \quad \lambda(R^\lambda) = 1.$$

Recall that a **closed  $\lambda$ -Reeb orbit** (or just Reeb orbit when  $\lambda$  is implicit) is a smooth map

$$o : (S^1 = \mathbb{R}/\mathbb{Z}) \rightarrow C$$

such that

$$\dot{o}(t) = cR^\lambda(o(t)),$$

with  $\dot{o}(t)$  denoting the time derivative, for some  $c > 0$  called period. Let  $S(R^\lambda, \beta)$  denote the space of all closed  $\lambda$ -Reeb orbits in free homotopy class  $\beta$ , with its compact open topology. And set

$$\mathcal{O}(R^\lambda, \beta) = S(R^\lambda, \beta)/S^1,$$

where  $S^1 = \mathbb{R}/\mathbb{Z}$  acts by reparametrization  $t \cdot o(\tau) = o(t + \tau)$ .

### 4. DEFINITION OF THE FUNCTIONAL $F$ AND PROOFS OF AUXILIARY RESULTS

Let  $X$  be a manifold. Fix once and for all an auxiliary Riemannian metric  $g_0$ , and set

$$C = S_{g_0}^* X.$$

For any Riemann–Finsler metric  $g$ , let  $F_g^*$  denote the dual norm, and let

$$\rho_g : C \rightarrow S_g^* X, \quad \rho_g(\xi) = \frac{\xi}{F_g^*(\xi)}$$

be the fiberwise radial projection. We define a contact form on the fixed manifold  $C$  by

$$\lambda_g = \rho_g^* \lambda_{\text{can}},$$

where  $\lambda_{\text{can}}$  is the Liouville one-form on  $T^*X$ . We write

$$R^{\lambda_g}$$

for the Reeb vector field of  $\lambda_g$  on this fixed manifold  $C$ . Under  $\rho_g$ , the flow of  $R^{\lambda_g}$  corresponds to the  $g$ -geodesic flow.

The same convention applies to the Sasaki metric. If  $g$  is Riemannian, let  $g_S^{\text{nat}}$  denote the usual Sasaki metric on  $S_g^* X$ , and set

$$g_S = \rho_g^* g_S^{\text{nat}}$$

on the fixed manifold  $C$ . Similarly, all metric-dependent functions on  $S_g^* X$ , such as verticality functions, are regarded as functions on  $C$  by pullback through  $\rho_g$ . This fixed- $C$  convention will be used throughout.



**4.1. Lifted geodesic strings.** We recall how closed geodesics in  $X$  are lifted to closed Reeb orbits in its cosphere bundle. Let  $C$  be the fixed cosphere model used in this paper, and let

$$q : C \rightarrow X$$

denote the projection. For a Riemann–Finsler metric  $g$ , the contact form  $\lambda_g$  on  $C$  has Reeb vector field  $R^{\lambda_g}$ . Its closed orbits project to oriented closed constant-speed  $g$ -geodesics in  $X$ .

Conversely, every oriented closed constant-speed  $g$ -geodesic has a unique Reeb lift to  $C$ . In the Riemannian case, if

$$o : S^1 \rightarrow X$$

is a nonconstant closed  $g$ -geodesic, its lift to the  $g$ -unit cotangent bundle is

$$\tilde{o}_g(t) = \frac{g_{o(t)}(\dot{o}(t), \cdot)}{|\dot{o}(t)|_g} \in S_g^*X.$$

Since  $o$  is a geodesic, this covector field is parallel along  $o$ . Thus, under the Levi–Civita splitting of  $T(T^*X)$ , the curve  $\tilde{o}_g$  is horizontal. Equivalently, after the constant reparametrization determined by the speed of  $o$ , it is a trajectory of the Reeb vector field. Passing to the fixed model  $C$  by radial projection gives the corresponding closed  $R^{\lambda_g}$ -orbit in  $C$ . The same statement holds in the Riemann–Finsler case using the Legendre transform.

The projection  $q$  induces a homeomorphism between closed  $R^{\lambda_g}$ -orbit strings in  $C$  and oriented closed constant-speed  $g$ -geodesic strings in  $X$ . Hence, for every class  $\beta \in \pi_1^{\text{inc}}(X)$ , if  $\mathcal{O}(g, \beta)$  denotes the class- $\beta$   $g$ -geodesic-string set as usual, then its Reeb lift

$$\tilde{\mathcal{O}}(g, \beta) = \{\tilde{o} \mid o \in \mathcal{O}(g, \beta)\} \subset \mathcal{O}(R^{\lambda_g})$$

is homeomorphic to  $\mathcal{O}(g, \beta)$ . In particular, if  $\mathcal{O}(g, \beta)$  is compact, then  $\tilde{\mathcal{O}}(g, \beta)$  is compact. Moreover,  $\tilde{\mathcal{O}}(g, \beta)$  is open in  $\mathcal{O}(R^{\lambda_g})$ , since the projected free homotopy class in  $X$  is locally constant on the loop space of  $C$ .

**4.2. Definition via the Fuller index.** Suppose then that  $g$  is  $\beta$ -taut. Define

$$F(g, \beta) = i(\tilde{\mathcal{O}}(g, \beta), R^{\lambda_g}) \in \mathbb{Q},$$

where the right hand side is the Fuller index, see Appendix A.

For a path of metrics  $\{g_t\}_{t \in [0,1]}$ , the lifted cobordism of orbit string spaces

$$\tilde{\mathcal{O}}(\{g_t\}, \beta) = \{(\tilde{o}, t) \mid o \in \mathcal{O}(g_t, \beta)\}$$

is likewise homeomorphic to the downstairs cobordism  $\mathcal{O}(\{g_t\}, \beta)$ .

*Proof of Theorem 1.14.* Let  $\beta \in \pi_1^{\text{inc}}(X)$  be given, and let  $\{g_t\}_{t \in [0,1]}$  be a  $\beta$ -taut deformation. As discussed above, the lifted cobordism  $\tilde{\mathcal{O}}(\{g_t\}, \beta)$  is compact by  $\beta$ -tautness. It is also open in the orbit cobordism, since the projected free homotopy class in  $X$  is locally constant. Basic invariance of the Fuller index, namely (A.3), then immediately gives

$$F(g_0, \beta) = F(g_1, \beta).$$

□

## 5. PROOF OF THEOREM 1.3

We prove a stronger result, partially removing the tautness condition.

**Theorem 5.1.** *Let  $g$  be the standard flat metric on  $T^n$ ,  $n > 1$ , and  $\beta \in \pi_1(T^n)$  a nontrivial class. For all  $L$  sufficiently large, there exists an  $\epsilon > 0$  such that whenever  $g'$  is a Finsler metric  $C^2$   $\epsilon$ -close to  $g$  and is  $\beta$ -regular, the following holds. Fix a prime  $p$ . Suppose that  $o$  is a  $g'$ -geodesic string s.t.:*

- (1) *The flat length satisfies  $\ell_g(o) < L$ .*
- (2)  *$o$  has class  $\beta$ .*
- (3)  *$p \mid \text{mult}(o)$ .*

*then there is at least one other such  $o$ .*

- *Any Finsler metric  $g_1$  taut homotopic to the flat metric has the same property as  $g$  above.*
- *In particular, if  $g_1$  is  $\beta$ -regular, is  $\beta$ -taut homotopic to the flat metric and if there is  $g_1$ -geodesic string  $o$  satisfying conditions (2), (3) then there is another one.*

*Proof.* We work on the fixed contact manifold

$$C = S_g^* T^n,$$

and let  $h$  be a fixed auxiliary metric on  $C$ , for instance the Sasaki metric associated to the flat metric  $g$ . The nearby unit cotangent bundles are identified with  $C$  by the fixed radial convention of Section 4; hence  $R^{\lambda_{g'}}$  denotes the corresponding pushed-forward Reeb vector field on  $C$ .

Let  $L_\beta$  denote the flat length of a class- $\beta$  geodesic, and let  $L > L_\beta$  be the length cutoff appearing in the statement. Since  $C$  is compact and  $R^{\lambda_{g'}} \rightarrow R^{\lambda_g}$  in  $C^0$  for  $g'$  sufficiently  $C^2$ -close to  $g$ , there is a preliminary  $C^2$ -neighborhood of  $g$  and constants

$$0 < a < A < \infty$$

such that, for every metric  $g''$  in this neighborhood,

$$|R^{\lambda_{g''}}|_h \leq A, \quad |\text{pr}_* R^{\lambda_{g''}}|_g \geq a$$

on  $C$ . Set

$$A_L = \left( \frac{A}{a} + 1 \right) L.$$

Define  $\widehat{U}_{L,A_L}$  to be the set of smooth loops  $\gamma : S^1 \rightarrow C$  such that

$$[\text{pr} \circ \gamma] = \beta, \quad \ell_g(\text{pr} \circ \gamma) < L, \quad \ell_h(\gamma) < A_L,$$

and let

$$U_{L,A_L} = \widehat{U}_{L,A_L} / S^1$$

be its image in the orbit-string quotient. This is a  $h$ -controlled neighborhood of the compact lifted flat orbit block

$$N_0 = \widetilde{\mathcal{O}}(g, \beta),$$

in the sense of Definition A.1. Indeed, the  $h$ -length upper bound is built into the definition of  $U_{L,A_L}$ , and the nonzero projected class  $\beta$  gives a positive lower  $h$ -length bound on  $\widehat{U}_{L,A_L}$ . The suitable compact set  $K$  comes from Lemma 2.1.

Since  $L > L_\beta$ , we have  $N_0 \subset U_{L,A_L}$ , and since the flat metric has no other class- $\beta$  geodesic strings,  $U_{L,A_L}$  is isolating for  $N_0$ .

Let  $W$  be as in Lemma A.4, and let  $\epsilon$  be smaller than both the corresponding Fuller constant for

$$(N_0, U_{L,A_L}, W)$$

and the size of the preliminary  $C^2$ -neighborhood above. If  $g'$  is sufficiently  $C^2$  close to  $g$ , such that the corresponding Reeb vector fields are  $C^0$   $\epsilon$ -close, Lemma A.6 gives

$$(5.2) \quad F(g, \beta) = i(N_1, R^{\lambda_{g'}}),$$

where

$$N_1 = U_{L, A_L} \cap \mathcal{O}(R^{\lambda_{g'}}).$$

We now identify the projection of  $N_1$  with the length-filtered class- $\beta$  geodesic strings of  $g'$ . Let  $\gamma$  be a closed  $R^{\lambda_{g'}}$ -orbit, parametrized by

$$\dot{\gamma} = TR^{\lambda_{g'}}(\gamma).$$

Then

$$\ell_h(\gamma) \leq AT,$$

while

$$\ell_g(\text{pr} \circ \gamma) = T \int_0^1 |\text{pr}_* R^{\lambda_{g'}}(\gamma(t))|_g dt \geq aT.$$

Thus

$$\ell_h(\gamma) \leq \frac{A}{a} \ell_g(\text{pr} \circ \gamma).$$

Consequently, for closed  $R^{\lambda_{g'}}$ -orbits,

$$\ell_g(\text{pr} \circ \gamma) < L \implies \ell_h(\gamma) < A_L.$$

Since the definition of  $U_{L, A_L}$  already includes the condition  $\ell_g(\text{pr} \circ \gamma) < L$ , the projection identifies  $N_1$  bijectively with the set of class- $\beta$   $g'$ -geodesic strings satisfying  $\ell_g(o) < L$ .

Let  $Z \subset T^n$  be a totally  $g$ -geodesic submanifold diffeomorphic to  $T^{n-1}$ , containing image  $\beta$ . Let  $\alpha \in H^1(T^n, \mathbb{Z})$  be the Poincaré dual of  $Z$  and let  $p : T^n \rightarrow S^1$  be the classifying map of  $\alpha$ . Then clearly  $p$  is a  $\beta$ -periodic fibration with respect to  $g$ . By Theorem 8.1,  $F(g, \beta) = 0$ , since  $\chi(S^1) = 0$ . And hence

$$(5.3) \quad i(N_1, R^{\lambda_{g'}}) = 0$$

By the previous paragraph, we may rewrite (5.3) as:

$$(5.4) \quad \sum_{o \in \mathcal{O}_L(g', \beta)} \frac{(-1)^{\text{morse}(o)}}{\text{mult}(o)} = 0,$$

Since  $g'$  is  $\beta$ -regular and  $N_1$  is compact,  $\mathcal{O}_L(g', \beta)$  is finite. Enumerate its elements as  $o_1, \dots, o_N$ . We then get:

$$\sum_i (-1)^{\text{morse}(o_i)} \prod_{j \neq i} \text{mult}(o_j) = 0.$$

We show that if a prime  $p$  divides the multiplicity  $\text{mult}(o_k)$  for some  $k \in \{1, \dots, N\}$ , then there exists at least one other index  $m \neq k$  such that  $p$  divides  $\text{mult}(o_m)$ . This will complete the proof of conclusion (1). The conclusions (2), (3) follow by the same argument using that  $F$  is a taut deformation invariant, Theorem 1.14.

Let  $M_i = \prod_{j \neq i} \text{mult}(o_j)$  and let  $\sigma_i = (-1)^{\text{morse}(o_i)}$ . The original equation is:

$$\sum_{i=1}^N \sigma_i M_i = 0.$$

Suppose  $p \mid \text{mult}(o_k)$ . Then

$$(5.5) \quad \sum_{i=1}^N \sigma_i M_i \equiv 0 \pmod{p}.$$

For any index  $i \neq k$ , the product  $M_i$  contains the factor  $\text{mult}(o_k)$ :

$$M_i = \text{mult}(o_k) \cdot \prod_{j \neq i, k} \text{mult}(o_j).$$

Since  $p \mid \text{mult}(o_k)$ , it follows from (5.5) that:

$$\sigma_k M_k + 0 \equiv 0 \pmod{p}.$$

And so

$$M_k \equiv 0 \pmod{p}.$$

By definition,  $M_k = \prod_{j \neq k} \text{mult}(o_j)$ . Since  $p$  is prime and  $p \mid \prod_{j \neq k} \text{mult}(o_j)$ , by Euclid's Lemma,

$$p \mid \text{mult}(o_j) \text{ for some } j \neq k.$$

And this completes the proof of the claim and the main theorem.  $\square$

## 6. PROOF OF THEOREM 1.5

Let  $p : T^n \rightarrow S^1$  be the  $\beta$ -periodic fibration with respect to  $g_0$ , as in the proof of Theorem 5.1 above. By Theorem 8.1,  $F(g_0, \beta) = 0$ . Hence, as we assumed Conjecture 2, for any other  $\beta$ -regular  $g$  on  $T^n$  with finitely many  $\beta$ -class geodesic strings we have:

$$\sum_{o \in \mathcal{O}(g, \beta)} \frac{(-1)^{\text{morse}(o)}}{\text{mult}(o)} = 0.$$

Then the conclusion of the Theorem follows by the same arithmetic argument as in the proof of Theorem 5.1 above.  $\square$

## 7. PROOF OF THEOREM 1.23

For simplicity, as the full argument is already lengthy, we prove the theorem under the assumption that  $g$  is Riemannian. In the Finsler case we must use a more modern variational setup for the Hilbert manifold of  $H^1$ -loops. In the Finsler case the energy functional is generally only  $C^1$ , rather than  $C^2$ , on the Hilbert loop space. Nevertheless the required Palais–Smale compactness and critical point theory for closed Finsler geodesics are standard; see Caponio–Javaloyes–Masiello [7], Mercuri [18], and Lu [17]. With these replacements, the compactness argument for end-incompressible classes, and the Fuller-index invariance argument go through without substantial change.

The circle  $S^1$  acts on  $L_\beta X$  by reparametrization:

$$(s \cdot \gamma)(t) = \gamma(t + s).$$

If some  $\gamma \in L_\beta X$  had nontrivial stabilizer, then  $\gamma$  would factor through a nontrivial covering  $S^1 \rightarrow S^1$ , contradicting indivisibility. Therefore the  $S^1$ -action on  $L_\beta X$  is free. Hence

$$H_*^{S^1}(L_\beta X, \mathbb{Z}) \cong H_*(L_\beta X/S^1, \mathbb{Z}),$$

and we will construct a finite total rank model for the latter.

Let

$$\mathcal{L}_\beta X = W^{1,2}(S^1, X)_\beta$$

be the Hilbert manifold of  $H^1$ -loops in the free homotopy class  $\beta$ . Consider the energy functional

$$\mathcal{E}_g : \mathcal{L}_\beta X \rightarrow \mathbb{R}, \quad \mathcal{E}_g(\gamma) = \int_{S^1} |\dot{\gamma}(t)|_g^2 dt.$$

Since  $S(g, \beta)$  is compact,  $\mathcal{E}_g$  is bounded above on the critical set of  $\mathcal{E}_g$  in class  $\beta$ . Choose

$$E_0 > \max_{\gamma \in S(g, \beta)} \mathcal{E}_g(\gamma) = M^2,$$

which exists by the  $\beta$ -tautness assumption. Set

$$\mathcal{U}_{E_0} = \{\gamma \in \mathcal{L}_\beta X \mid \mathcal{E}_g(\gamma) < E_0\}.$$

**Lemma 7.1.** *The inclusion*

$$\mathcal{U}_{E_0} \hookrightarrow \mathcal{L}_\beta X$$

*is an  $S^1$ -equivariant homotopy equivalence. Consequently,*

$$\mathcal{U}_{E_0}/S^1 \simeq \mathcal{L}_\beta X/S^1.$$

*Moreover, the inclusion of smooth loops into the Hilbert loop space is an  $S^1$ -equivariant homotopy equivalence, and hence*

$$\mathcal{L}_\beta X/S^1 \simeq L_\beta X/S^1.$$

*Therefore*

$$\mathcal{U}_{E_0}/S^1 \simeq L_\beta X/S^1.$$

*Proof.* The energy functional  $\mathcal{E}_g$  is smooth and  $S^1$ -invariant. We equip  $\mathcal{L}_\beta X$  with the standard  $S^1$ -invariant Hilbert metric induced by  $g$ . With respect to this metric, the negative gradient flow of  $\mathcal{E}_g$  is  $S^1$ -equivariant.

We first recall the Palais–Smale compactness needed for the deformation lemma. Let  $\{\gamma_j\} \subset \mathcal{L}_\beta X$  be a Palais–Smale sequence for  $\mathcal{E}_g$  with

$$\mathcal{E}_g(\gamma_j) \leq A.$$

Then

$$\ell_g(\gamma_j) \leq \mathcal{E}_g(\gamma_j)^{1/2} \leq A^{1/2}.$$

Since  $\beta \in \pi_1^{\text{inc}}(X)$ , Lemma 2.1 implies that the images  $\gamma_j(S^1)$  lie in a fixed compact subset of  $X$ . On this compact set, the usual Hilbert-loop compactness argument for the geodesic energy gives a strongly convergent subsequence in  $\mathcal{L}_\beta X$ . Thus  $\mathcal{E}_g$  satisfies the Palais–Smale condition on sublevels.

By the choice of  $E_0$ , the functional  $\mathcal{E}_g$  has no critical points in

$$\{\mathcal{E}_g \geq E_0\}.$$

Therefore, for every  $A > E_0$ , Palais’ deformation lemma gives an  $S^1$ -equivariant homotopy equivalence

$$\{\mathcal{E}_g < E_0\} \hookrightarrow \{\mathcal{E}_g < A\}.$$

Letting  $A \rightarrow \infty$ , or equivalently applying the negative gradient flow on the exhaustion by sublevels, we obtain an  $S^1$ -equivariant homotopy equivalence

$$\mathcal{U}_{E_0} \simeq_{S^1} \mathcal{L}_\beta X.$$

Passing to quotients gives

$$\mathcal{U}_{E_0}/S^1 \simeq \mathcal{L}_\beta X/S^1.$$

Finally, the inclusion

$$C^\infty(S^1, X)_\beta \hookrightarrow W^{1,2}(S^1, X)_\beta$$

is an  $S^1$ -equivariant homotopy equivalence by standard equivariant smoothing approximation for loops. Since  $L_\beta X$  denotes the same free homotopy component in the smooth or compact-open loop space, we get

$$\mathcal{L}_\beta X \simeq_{S^1} L_\beta X.$$

Combining the quotient equivalences gives

$$\mathcal{U}_{E_0}/S^1 \simeq L_\beta X/S^1.$$

□

We now compare the Fuller index with the equivariant Morse model. The only subtlety is that after perturbing  $g$  to a regular metric  $g'$ , the full class- $\beta$  geodesic set of  $g'$  need not be compact. We therefore work in an isolating neighborhood of the compact lifted orbit block coming from  $g$ .

Let  $g_S$  denote the Sasaki metric on the fixed contact manifold  $C$  induced by  $g$  under the convention of Section 4. Define the  $S^1$ -invariant loop set  $\widehat{U}_{\sqrt{E_0}}$  to consist of all loops  $\gamma : S^1 \rightarrow C$  such that

$$[q \circ \gamma] = \beta, \quad \ell_{g_S}(\gamma) < \sqrt{E_0}.$$

Let

$$U_{\sqrt{E_0}} = \widehat{U}_{\sqrt{E_0}} / S^1$$

be its image in the orbit-string quotient. By (A.7) and the choice of  $E_0$ , this is an isolating neighborhood of

$$N_0 = \widetilde{\mathcal{O}}(g, \beta)$$

in the sense of Lemma A.4. It is also  $g_S$ -controlled, in the sense of Definition A.1, with the lower length bound supplied by the nontrivial projected class  $\beta$  and  $K$  provided by Lemma 2.1.

Let  $W$  be as in Lemma A.4, and let  $\epsilon$  be the corresponding Fuller constant for the isolating data

$$(N_0, U_{\sqrt{E_0}}, W).$$

Choose a  $\beta$ -regular complete metric  $g'$  sufficiently  $C^2$ -close  $\epsilon'$ -close to  $g$  so that, on the fixed manifold  $C$ , the vector field  $R^{\lambda_{g'}}$  is  $C^0\epsilon$ -close to  $R^{\lambda_g}$ . Choose  $E'$  with

$$\max_{\gamma \in N_0} \ell_{g_S}(\gamma) < \sqrt{E'} < \sqrt{E_0}.$$

Set  $\widehat{U}$  to be the set of loops  $\gamma : S^1 \rightarrow C$  such that

$$[q \circ \gamma] = \beta, \quad \ell_{g'_S}(\gamma) < \sqrt{E'},$$

and set

$$U = \widehat{U} / S^1.$$

After shrinking the perturbation if necessary, we may assume

$$W \subset U \subset U_{\sqrt{E_0}}.$$

Thus Lemma A.6 gives

$$F(g, \beta) = i(N_1, R^{\lambda_{g'}}),$$

where

$$N_1 = U \cap \mathcal{O}(R^{\lambda_{g'}}).$$

It remains to compute the last Fuller index. Let

$$\mathcal{U}_{E'}^{g'} = \{\gamma \in \mathcal{L}_\beta X \mid \mathcal{E}_{g'}(\gamma) < E'\}.$$

For a constant-speed  $g'$ -geodesic, its canonical lift has  $g'_S$ -length equal to its  $g'$ -length. Hence the critical circles of  $\mathcal{E}_{g'}$  in  $\mathcal{U}_{E'}^{g'}$  are in one-to-one correspondence with the elements of  $N_1$ . Since  $g'$  is  $\beta$ -regular, this critical set is a finite union of nondegenerate critical circles

$$C_o \simeq S^1.$$

The Morse–Bott handlebody  $\mathcal{MB}$  obtained from the unstable disk bundles of these critical circles is  $S^1$ -invariant, and the standard Hilbert-manifold Morse–Bott deformation theorem gives an  $S^1$ -equivariant homotopy equivalence

$$\mathcal{MB} \simeq_{S^1} \mathcal{U}_{E'}^{g'}.$$

After passing to the quotient by  $S^1$ , the unstable disk bundle over  $C_o$  contributes one cell of dimension  $\text{morse}(o)$ . Therefore

$$\chi(\mathcal{U}_{E'}^{g'} / S^1) = \chi(\mathcal{MB} / S^1) = \sum_{o \in N_1} (-1)^{\text{morse}(o)}.$$

We now compare the  $g'$ -sublevel with the Hilbert sublevel for  $g$ . Choose numbers  $a, b$  and  $\delta > 0$  such that

$$M^2 < a < a + \delta < E' < b < E_0$$

and such that  $\mathcal{E}_g$  has no critical points on

$$\{a \leq \mathcal{E}_g \leq b\}.$$

Choosing a smaller  $\epsilon'$  above if necessary, the hypotheses of Lemma 7.2 hold for

$$A = U_M^g, \quad C = U_{E'}^{g'}, \quad B = U_{E_0}^g.$$

In particular,

$$U_M^g \subset U_{E'}^{g'} \subset U_{E_0}^g.$$

Lemma ?? then gives an  $S^1$ -equivariant homotopy equivalence

$$U_{E'}^{g'} \simeq U_{E_0}^g.$$

By Lemma 7.2, applied with  $a + \delta$  in place of  $a$ , the inclusion

$$\mathcal{U}_{E'}^{g'} \hookrightarrow \mathcal{U}_b^g$$

is an  $S^1$ -equivariant homotopy equivalence. Since all critical points of  $\mathcal{E}_g$  have energy at most  $M^2 < b$ , the inclusion

$$\mathcal{U}_b^g \hookrightarrow \mathcal{U}_{E_0}^g$$

is also an  $S^1$ -equivariant homotopy equivalence. Thus

$$\mathcal{U}_{E'}^{g'} \simeq_{S^1} \mathcal{U}_{E_0}^g.$$

By Lemma 7.1,

$$\mathcal{U}_{E_0}^g / S^1 \simeq L_\beta X / S^1.$$

Consequently

$$\chi(\mathcal{U}_{E'}^{g'} / S^1) = \chi(L_\beta X / S^1).$$

In particular  $H_*^{S^1}(L_\beta X, \mathbb{Z})$  has finite total rank.

On the other hand, since  $g'$  is  $\beta$ -regular, the Fuller index formula, summed over the finitely many lifted free homotopy classes met by  $N_1$ , gives

$$i(N_1, R^{\lambda_{g'}}) = \sum_{o \in N_1} \frac{(-1)^{\text{morse}(o)}}{\text{mult}(o)}.$$

Since  $\beta$  is indivisible, all multiplicities in this class are 1. Hence

$$i(N_1, R^{\lambda_{g'}}) = \chi(\mathcal{U}_{E'}^{g'} / S^1) = \chi(L_\beta X / S^1).$$

Therefore

$$F(g, \beta) = \chi(L_\beta X / S^1) = \chi^{S^1}(L_\beta X).$$

□

We state the following elementary lemma separately, since it may be of independent interest.

**Lemma 7.2.** *For a metric  $h$  and a number  $r$ , set*

$$\mathcal{U}_r^h = \{\gamma \in \mathcal{L}_\beta X \mid \mathcal{E}_h(\gamma) < r\}.$$

*Let  $a < b$ , and suppose that  $\mathcal{E}_g$  satisfies the Palais-Smale condition on the band*

$$\mathcal{B} = \{a \leq \mathcal{E}_g \leq b\}$$

*and has no critical points there. Let  $g'$  be sufficiently  $C^2$ -close to  $g$ , so that  $\nabla \mathcal{E}_{g'}$  is  $C^0$ -close to  $\nabla \mathcal{E}_g$  on  $\mathcal{B}$ . Suppose*

$$\mathcal{U}_a^g \subset \mathcal{C} \subset \mathcal{U}_b^g,$$

where

$$\mathcal{C} = \{\mathcal{E}_{g'} < c\}$$

is an  $S^1$ -invariant sublevel set of  $\mathcal{E}_{g'}$ . Then the inclusion

$$\mathcal{C} \hookrightarrow \mathcal{U}_b^g$$

is an  $S^1$ -equivariant homotopy equivalence.

*Proof.* By the Palais–Smale condition and the absence of critical points on  $\mathcal{B}$ , there exists  $\mu > 0$  such that

$$(7.3) \quad \|\nabla \mathcal{E}_g\| \geq \mu$$

on  $\mathcal{B}$ . For  $g'$  sufficiently close to  $g$ , we have

$$(7.4) \quad \|\nabla \mathcal{E}_{g'} - \nabla \mathcal{E}_g\| < \mu/2$$

on the same band. Therefore

$$\langle \nabla \mathcal{E}_g, \nabla \mathcal{E}_{g'} \rangle > 0$$

on  $\mathcal{B}$ .

Since

$$\mathcal{U}_a^g \subset \mathcal{C} \subset \mathcal{U}_b^g,$$

we have

$$\mathcal{U}_b^g - \mathcal{C} \subset \mathcal{U}_b^g - \mathcal{U}_a^g \subset \mathcal{B}.$$

Thus, as long as a negative  $\mathcal{E}_{g'}$ -gradient trajectory lies in  $\mathcal{U}_b^g - \mathcal{C}$ , we have

$$\frac{d}{dt} \mathcal{E}_g = -\langle \nabla \mathcal{E}_g, \nabla \mathcal{E}_{g'} \rangle < 0.$$

Consequently the flow cannot exit  $\mathcal{U}_b^g$  before entering  $\mathcal{C}$ . Moreover,  $\mathcal{C}$  is forward-invariant because it is a sublevel set of  $\mathcal{E}_{g'}$ .

The estimates above also imply that  $\mathcal{E}_{g'}$  has no critical points on  $\mathcal{U}_b^g - \mathcal{C}$ . The standard deformation argument of Palais [21, Section 10, Proposition (2)], applied to  $\mathcal{E}_{g'}$  on the region  $\mathcal{U}_b^g - \mathcal{C}$ , therefore gives a deformation retraction of  $\mathcal{U}_b^g$  onto  $\mathcal{C}$ . The energies and the Hilbert metrics are  $S^1$ -invariant, so the gradient flow is  $S^1$ -equivariant. Hence the deformation retraction is  $S^1$ -equivariant.  $\square$

## 8. PRODUCT FORMULA

In this section we prove a kind of product formula for the  $F$ -count. This is used in all our applications.

**Theorem 8.1** (Product formula). *Let*

$$Z \hookrightarrow (X, g) \xrightarrow{p} (Y, g_Y)$$

*be a  $\beta$ -periodic fibration, with  $Y$  closed, connected and such that all smooth closed contractible  $g_Y$ -geodesics in  $Y$  are constant. Then  $p$  is  $\beta$ -taut. Fix  $y_0 \in Y$ , set  $Z_0 = p^{-1}(y_0)$ , and choose*

$$\beta_Z \in \pi_1^{\text{inc}}(Z_0)$$

*such that*

$$(i_{y_0})_*(\beta_Z) = \beta,$$

*where  $i_{y_0} : Z_0 \hookrightarrow X$  is the inclusion. Denote by  $g_{y_0}$  the restriction of  $g$  to the fiber of  $y_0$ . Let*

$$H = \pi_1(Y, y_0)$$

*act on  $\pi_1^{\text{inc}}(Z_{y_0})$  by holonomy, and set*

$$S_{y_0} = H \cdot \beta_Z.$$



Then

$$F(g, \beta) = (-1)^{\dim Y} \cdot \text{card}(S_{y_0}) \cdot \chi(Y) \cdot F(g_{y_0}, \beta_Z).$$

*Proof.* We prove  $\beta$ -tautness. We first show that every class- $\beta$   $g$ -geodesic in  $X$  is vertical. Let

$$o : S^1 \rightarrow X$$

be a closed  $g$ -geodesic representing the class  $\beta$ . Since  $\beta$  is a fiber class, the loop

$$p \circ o : S^1 \rightarrow Y$$

is contractible. Since  $p$  is a parallel Riemannian submersion, the local de Rham product decomposition for  $p$  implies that  $p \circ o$  is a  $g_Y$ -geodesic, possibly constant. Hence  $p \circ o$  is a smooth closed contractible  $g_Y$ -geodesic. By hypothesis, every such geodesic is constant. Therefore  $p \circ o$  is constant, and so  $o$  is contained in a single fiber

$$Z_y = p^{-1}(y).$$

Thus the class- $\beta$  geodesic strings in  $X$  are precisely the vertical geodesic strings whose fiber class represents  $\beta$  in  $X$ .

It remains to prove compactness. Let  $o_j$  be a sequence of class- $\beta$   $g$ -geodesic strings in  $X$ . By the previous paragraph, each  $o_j$  is contained in a fiber  $Z_{y_j}$ . Since  $Y$  is compact, after passing to a subsequence we may assume

$$y_j \rightarrow y_\infty.$$

Choose a contractible coordinate ball

$$U \subset Y$$

around  $y_\infty$ . For all large  $j$ , we have  $y_j \in U$ . Since  $p$  is a parallel Riemannian submersion, the local de Rham product theorem gives a local Riemannian product decomposition

$$p^{-1}(U) \cong Z_{y_\infty} \times U.$$

Under this identification, vertical geodesics in the fibers  $Z_{y_j}$  are identified with geodesics of the fiber metric on  $Z_{y_\infty}$ , and their fiber free homotopy classes lie in the finite holonomy orbit

$$S_{y_\infty}.$$

By the fiber-tautness condition in the definition of a  $\beta$ -periodic fibration, the union of the geodesic-string sets

$$\bigcup_{\alpha \in S_{y_\infty}} S(g_{y_\infty}, \alpha)$$

is compact. Since  $S_{y_\infty}$  is finite, this is a finite union of compact sets. Therefore, after passing to a further subsequence, the vertical geodesic strings  $o_j$ , transported to the model fiber  $Z_{y_\infty}$ , converge to a fiber geodesic string. Transporting back through the local product chart gives convergence of the original  $o_j$  in the class- $\beta$  geodesic-string space of  $X$ .

Thus every sequence of class- $\beta$  geodesic strings has a convergent subsequence. Hence the class- $\beta$  geodesic-string set is compact, and  $g$  is  $\beta$ -taut.

Let

$$C = S^*X$$

be the unit cotangent bundle of  $(X, g)$ , with projection

$$\text{pr} : C \rightarrow X.$$

For this fixed metric calculation, this is equivalent to the fixed- $C$  convention of Section 4. Let  $R^{\lambda_g}$  denote the Reeb vector field on  $C$ . The  $F$ -invariant is computed from the compact lifted orbit set

$$N_0 = \tilde{\mathcal{O}}(g, \beta).$$

The idea is to localize contributions over critical points of a Morse function on  $Y$ .

Let

$$\pi^{\text{vert}} : TX \rightarrow T^{\text{vert}}X = \ker(dp)$$

be the  $g$ -orthogonal projection. By assumptions on  $p : X \rightarrow Y$ ,

$$\nabla \pi^{\text{vert}} = 0,$$

where  $\pi^{\text{vert}}$  is viewed as a  $(1, 1)$ -tensor field.

For  $\xi \in C$ , let

$$v_\xi \in T_{\text{pr}(\xi)}X$$

be the  $g$ -dual unit vector to  $\xi$ . Define

$$P : C \rightarrow \mathbb{R}$$

by

$$P(\xi) = |\pi^{\text{vert}}v_\xi|_g^2.$$

Thus  $P = 1$  on vertical unit covectors and  $P = 0$  on horizontal unit covectors.

Let  $f : Y \rightarrow \mathbb{R}$  be a Morse function. Define

$$\tilde{f} : C \rightarrow \mathbb{R}$$

by

$$\tilde{f} = f \circ p \circ \text{pr}.$$

Equip  $C = S^*X$  with the Sasaki metric  $g_S$  induced by  $g$ , as in the Proof of Theorem 1.23. We shall use the following two elementary identities.

First,

$$(8.2) \quad dP(R^{\lambda_g}) = 0.$$

Indeed, let  $\xi(\tau)$  be a flow line of  $R^{\lambda_g}$ , and set

$$x(\tau) = \text{pr}(\xi(\tau)).$$

Then  $x(\tau)$  is a unit-speed  $g$ -geodesic. Let

$$v(\tau) = \dot{x}(\tau) = v_{\xi(\tau)}$$

be its velocity vector field. Since  $x(\tau)$  is a geodesic,

$$\nabla_\tau v = 0.$$

Since  $\nabla \pi^{\text{vert}} = 0$ , we have

$$\nabla_\tau(\pi^{\text{vert}}v) = \pi^{\text{vert}}(\nabla_\tau v) = 0.$$

Therefore

$$\begin{aligned} \frac{d}{d\tau}P(\xi(\tau)) &= \frac{d}{d\tau}|\pi^{\text{vert}}v(\tau)|_g^2 \\ &= 2\langle \nabla_\tau(\pi^{\text{vert}}v), \pi^{\text{vert}}v \rangle_g \\ &= 0. \end{aligned}$$

Hence  $dP(R^{\lambda_g}) = 0$ .

Second,

$$(8.3) \quad dP(\nabla_{g_S}\tilde{f}) = 0.$$

Since  $\tilde{f} = f \circ p \circ \text{pr}$  depends only on the base point  $\text{pr}(\xi)$ :

$$\nabla_{g_S}\tilde{f} \in T^{\text{hor}}C.$$

More precisely,  $\nabla_{g_S}\tilde{f}$  is the horizontal lift of

$$\nabla_g(f \circ p)$$

on  $X$ . Let  $\xi(s)$  be a horizontal curve in  $C$ , and set

$$x(s) = \text{pr}(\xi(s)).$$

By definition of the Levi-Civita horizontal distribution on  $S^*X$ , the covectors  $\xi(s)$  are parallel along  $x(s)$ . Equivalently, their  $g$ -dual vectors

$$v(s) = v_{\xi(s)}$$

are parallel along  $x(s)$ :

$$\nabla_s v(s) = 0.$$

Using again  $\nabla \pi^{\text{vert}} = 0$ , we get

$$\nabla_s(\pi^{\text{vert}} v(s)) = \pi^{\text{vert}}(\nabla_s v(s)) = 0.$$

Hence

$$\begin{aligned} \frac{d}{ds} P(\xi(s)) &= \frac{d}{ds} |\pi^{\text{vert}} v(s)|_g^2 \\ &= 2 \langle \nabla_s(\pi^{\text{vert}} v(s)), \pi^{\text{vert}} v(s) \rangle_g \\ &= 0. \end{aligned}$$

Taking  $\dot{\xi}(0) = \nabla_{g_S} \tilde{f}(\xi)$  gives

$$dP_{\xi}(\nabla_{g_S} \tilde{f}) = 0.$$

Since  $\xi$  was arbitrary,

$$dP(\nabla_{g_S} \tilde{f}) = 0.$$

Now set

$$V_{\varepsilon} = R^{\lambda_g} - \varepsilon \nabla_{g_S}(P + \tilde{f}).$$

Recall first the discussion of Fuller continuation in the Appendix. The set  $N_0 = \tilde{\mathcal{O}}(g, \beta)$  is compact by  $\beta$ -tautness. Choose a controlled isolating neighborhood  $U$  of  $N_0$  and an open neighborhood  $W$  with  $\bar{W} \subset U$ , and let  $\epsilon'$  be the corresponding Fuller constant. Take  $\varepsilon > 0$  small enough so that  $V_{\varepsilon}$  is  $C^0$   $\epsilon'$ -close to  $R^{\lambda_g}$  on the compact region determined by  $U$ . Lemma A.6 gives a compact continued set  $N_{\varepsilon}$  and

$$F(g, \beta) = i(N_{\varepsilon}, V_{\varepsilon}).$$

Note that the above does not require any further properties of  $V_{\varepsilon}$ , such as completeness. We set up the Fuller index theory to sidestep such issues.

We now identify the closed orbits in  $N_{\varepsilon}$ . We show that  $P$  is a Lyapunov function for  $V_{\varepsilon}$ . Mainly, along a trajectory  $\xi(\tau)$  of  $V_{\varepsilon}$ , we have

$$\begin{aligned} \frac{d}{d\tau} P(\xi(\tau)) &= dP(R^{\lambda_g}) - \varepsilon dP(\nabla_{g_S} P) - \varepsilon dP(\nabla_{g_S} \tilde{f}) \\ &= -\varepsilon |\nabla_{g_S} P|_{g_S}^2 \leq 0. \end{aligned}$$

Thus  $P$  is nonincreasing along the  $V_{\varepsilon}$ -flow.

If  $\xi(\tau)$  is a closed orbit of  $V_{\varepsilon}$ , then  $P(\xi(\tau))$  returns to its initial value after one period. Since  $P$  is nonincreasing, it must be constant along the orbit. Hence

$$\nabla_{g_S} P = 0$$

along the orbit.

On each unit cotangent sphere,  $P$  is the squared norm of the vertical component of the corresponding unit vector. Therefore the critical locus of  $P$  consists of the two critical loci

$$P = 1 \quad \text{and} \quad P = 0,$$

corresponding respectively to vertical and horizontal covectors. At  $\varepsilon = 0$ , the lifted orbit set under consideration is vertical, hence lies in  $P = 1$ . Taking the controlled neighborhood  $U$  (above) so that all its elements are disjoint (as loops) from  $P = 0$ , we get that every closed orbit in  $N_{\varepsilon}$  lies in the

vertical critical locus  $P = 1$ . Consequently, every closed orbit in  $N_\varepsilon$  projects to a geodesic contained in a fiber  $Z_y$ .

It remains to see over which fibers these vertical orbits occur. On  $P = 1$ , the Reeb vector field restricts to the geodesic Reeb field of the fiber  $(Z_y, g_y)$ , while  $\tilde{f}$  depends only on the base point  $y$ . The perturbation

$$-\varepsilon \nabla_{g_S} \tilde{f}$$

moves the base point in the negative gradient direction of  $f$ . Therefore a vertical closed orbit of  $V_\varepsilon$  can occur only over a critical point of  $f$ . Conversely, if  $y \in \text{Crit}(f)$ , then

$$\nabla_{g_S} \tilde{f} = 0$$

over  $Z_y$ , and the vertical part of  $V_\varepsilon$  is just the fiber geodesic flow, up to the harmless  $P$ -term which vanishes on the vertical critical locus. Thus the closed orbits of  $V_\varepsilon$  in  $N_\varepsilon$  are precisely the vertical closed geodesic orbits lying in fibers over critical points of  $f$ .

We now compute the index. For  $y \in Y$ , put

$$H_y = \pi_1(Y, y),$$

acting on  $\pi_1^{\text{inc}}(Z_y)$  by holonomy. Choose

$$\beta_Z \in \pi_1^{\text{inc}}(Z_y)$$

with

$$(i_y)_*(\beta_Z) = \beta.$$

Set

$$S_y = H_y \cdot \beta_Z.$$

The Fuller index is additive over disjoint compact open subsets. Thus it remains to compute the local index contribution  $i(N_{\varepsilon, y}, V_\varepsilon)$  corresponding to the isolated subset of  $N_\varepsilon$  lying over a critical point  $y$  of  $f$ . We defer this computation to the Appendix as Lemma B.1; it yields that

$$(8.4) \quad i(N_{\varepsilon, y}, V_\varepsilon) = (-1)^{\dim Y + \text{morse}(y)} \sum_{\alpha \in S_y} F(g_y, \alpha),$$

Next, we show that  $F(g_y, \cdot)$  is constant on  $S_y$ . Indeed, let

$$\gamma : [0, 1] \rightarrow Y$$

be a path based at  $y$ . Pulling back  $p : X \rightarrow Y$  along  $\gamma$  gives a fibration over  $[0, 1]$ . By the same proof that  $p$  is  $\beta$ -taut, this pullback is a taut deformation of the fiber metric from  $(Z_y, g_y)$  back to itself, with final fiber identified with the initial one by the holonomy map  $h_\gamma : Z_y \rightarrow Z_y$ . Fuller index invariance under taut deformation gives

$$F(g_y, \alpha) = F(g_y, h_{\gamma, *} \alpha).$$

Thus  $F(g_y, \alpha)$  is constant on the holonomy orbit  $S_y$ . Hence

$$\sum_{\alpha \in S_y} F(g_y, \alpha) = \#S_y \cdot F(g_y, \beta_Z).$$

If  $y, y' \in Y$ , holonomy along any path from  $y$  to  $y'$  identifies  $S_y$  with  $S_{y'}$ . The same path gives a taut deformation between the fiber metrics, so

$$\sum_{\alpha \in S_y} F(g_y, \alpha)$$

is independent of  $y$ . So we get:

$$\begin{aligned}
F(g, \beta) &= \sum_{y \in \text{Crit}(f)} i(N_{\varepsilon, y}, V_{\varepsilon}) \\
&= \sum_{y \in \text{Crit}(f)} (-1)^{\dim Y + \text{morse}(y)} \sum_{\alpha \in S_y} F(g_y, \alpha), \\
&= \sum_{y \in \text{Crit}(f)} (-1)^{\dim Y + \text{morse}(y)} \cdot \#S_{y_0} \cdot F(g_{y_0}, \beta_Z) \\
&= (-1)^{\dim Y} \cdot \chi(Y) \cdot \#S_{y_0} \cdot F(g_{y_0}, \beta_Z).
\end{aligned}$$

□

**Corollary 8.5.** *Let  $X = Z \times Y$ , with  $Y$  closed, connected, such that all smooth closed contractible  $g_Y$ -geodesics in  $Y$  are constant and such that  $Z, g_Z$  is taut. Let  $\beta \in \pi_1^{\text{inc}}(X)$  be a fiber class. Then  $g = g_Z \times g_Y$  is  $\beta$ -taut. Furthermore,*

$$F(g, \beta) = (-1)^{\dim Y} \cdot \chi(Y) \cdot F(g_Z, \beta_Z).$$

*Proof.* This is immediate from the theorem. □

## 9. PROOF OF THEOREM 1.30 AND ITS COROLLARIES

We prove the first part. By Cartan–Hadamard, a complete negatively curved Riemannian manifold has at most one closed geodesic in each nontrivial free homotopy class; moreover the absence of nontrivial periodic Jacobi fields in negative curvature gives nondegeneracy. Existence of a geodesic in the end-incompressible class  $\beta$  follows from Lemma 2.4. Hence a complete negatively curved metric would give a unique nondegenerate class- $\beta$  geodesic string. It thus suffices to prove the second part.

We prove the second part. Suppose first that  $\beta$  is atomic, so that  $\beta$  has a representative that is a  $k$ -cover of a loop,  $k \geq 1$ , whose corresponding class  $\alpha$  is indivisible. By the assumption that all contractible  $g_Y$  geodesics are constant, classical Morse theory Milnor [20] tells us that  $Y$  has vanishing higher homotopy groups  $\pi_j(Y, y_0)$ ,  $j \geq 2$ . Thus by the long exact sequence of a fibration  $i_{Z,*} : \pi_1(Z, z_0) \rightarrow \pi_1(X, i_Z(z_0))$  is a group injection, and image  $i_{Z,*} = \ker p_*$ . Now  $\pi_1(Y, y_0)$  is torsion-free since it is a finite dimensional  $K(\pi_1(Y, y_0), 1)$  space, Hatcher [13, Prop. 2.45]. It follows that  $\alpha$  is also a fiber class since

$$p_*(\alpha^k) = 0 \implies p_*(\alpha) = 0.$$

Furthermore,  $\alpha \in \pi_1^{\text{inc}}(X)$  (otherwise  $\beta$  isn't).

**Lemma 9.1.** *Let*

$$Z \hookrightarrow (X, g) \xrightarrow{p} (Y, g_Y)$$

*be a  $g$ -parallel Riemannian submersion with complete total space, and assume that  $Y$  is connected with  $\pi_2(Y) = 0$ . Let*

$$\beta, \gamma \in \pi_1^{\text{inc}}(X)$$

*be fiber classes such that*

$$\beta = \gamma^m$$

*for some  $m \geq 1$ . If  $g$  is  $\beta$ -taut, then  $g$  is  $\gamma$ -taut and  $p$  is  $\gamma$ -holonomy-finite.*

*Proof.* We show  $\gamma$ -tautness. Every closed  $\gamma$ -geodesic  $o_\gamma$  has  $m$ -fold cover  $o_\gamma^m \in S(g, \beta)$ , so

$$\ell_g(o_\gamma) = \frac{1}{m} \ell_g(o_\gamma^m) \leq A/m, \quad A := \max_{o \in S(g, \beta)} \ell_g(o) < \infty,$$

where finiteness of  $A$  is  $\beta$ -tautness. Hence  $S(g, \gamma) = S_{A/m}(g, \gamma)$ , which is compact by Lemma 2.2.

We show  $\gamma$ -holonomy-finiteness. Choose

$$\gamma_Z \in \mathcal{R}(\gamma)$$

and set

$$S_{y_0} = \pi_1(Y, y_0) \cdot \gamma_Z.$$

Suppose, toward a contradiction, that  $S_{y_0}$  is infinite. For each

$$\gamma_i \in S_{y_0}$$

choose a closed  $g_{y_0}$ -geodesic

$$o_i \subset Z_{y_0}$$

representing  $\gamma_i$ . Such geodesics exist by Lemma 2.4. Since the classes  $\gamma_i$  are distinct in  $Z_{y_0}$ , the loops  $o_i$  are pairwise distinct as elements of the fiber loop space.

We first record a separation property for these loops in the ambient loop space  $LX$ . Choose a contractible trivializing neighborhood

$$U \subset Y$$

of  $y_0$ . For a loop  $o \subset Z_{y_0}$ , let

$$\mathcal{V}_{U,o} \subset LX$$

be the set of loops  $\eta : S^1 \rightarrow X$  such that

$$p(\eta(S^1)) \subset U$$

and such that  $\eta$ , regarded as a loop in  $p^{-1}(U)$ , is freely homotopic in  $p^{-1}(U)$  to  $o$ . The set  $\mathcal{V}_{U,o}$  is open in  $LX$ : the condition that the image lie in  $p^{-1}(U)$  is open, and free homotopy classes are open components of the loop space of  $p^{-1}(U)$ .

Moreover, if  $o_i$  and  $o_j$  represent distinct free homotopy classes in  $Z_{y_0}$ , then

$$\mathcal{V}_{U,o_i} \cap \mathcal{V}_{U,o_j} = \emptyset.$$

Indeed, an element of the intersection would give a free homotopy between  $o_i$  and  $o_j$  inside  $p^{-1}(U)$ . Since  $U$  is contractible, the inclusion

$$Z_{y_0} \hookrightarrow p^{-1}(U)$$

is a homotopy equivalence, and hence  $o_i$  and  $o_j$  would already be freely homotopic inside  $Z_{y_0}$ , contrary to construction.

On the other hand,  $S(g, \gamma)$  is compact. Thus the infinite family  $\{o_i\} \subset S(g, \gamma)$  has an accumulation point  $o_\infty \in S(g, \gamma)$ . Since every  $o_i$  lies in the fiber  $Z_{y_0}$ , the limit  $o_\infty$  also lies in  $Z_{y_0}$ . The open neighborhood  $\mathcal{V}_{U,o_\infty}$  of  $o_\infty$  therefore contains  $o_i$  for infinitely many  $i$ , forcing these  $o_i$ 's to lie in the same fiber free homotopy class as  $o_\infty$ . This contradicts the fact that the classes  $\gamma_i$  are pairwise distinct. Hence  $S_{y_0}$  is finite and  $p$  is  $\gamma$ -holonomy-finite.

□

By Theorem 8.1  $p$  is  $\beta$ -taut. By the lemma above  $p$  is  $\alpha$ -holonomy finite and hence is an  $\alpha$ -periodic fibration. Now, if  $\chi(Y) \neq \pm 1$  then by the product formula of Theorem 8.1  $F(g, \alpha) \neq \pm 1$ , since  $F(g_{y_0}, \alpha_Z)$  is an integer by Theorem 1.23.

Now, if  $X$  admits a complete metric  $h$  with a unique and nondegenerate class- $\beta$   $h$ -geodesic string  $o_\beta$ , then it clearly also has a unique class- $\alpha$   $h$ -geodesic string  $o_\alpha$ . Furthermore,  $o_\alpha$  must be nondegenerate as otherwise  $o_\beta$ , which is a  $k$ -fold cover of  $o_\alpha$  is degenerate. Then we have:

$$F(h, \alpha) = \pm 1.$$

But this is a contradiction, since  $F(h, \alpha) = F(g, \alpha)$  by Theorem 1.23.

We now prove the general case. Suppose by contradiction that  $X$  admits a complete metric  $h$  with a unique and nondegenerate class- $\beta$   $h$ -geodesic string  $o_\beta$ . By the preceding paragraph, the class  $\beta$  is not atomic. Now  $o_\beta$  covers a multiplicity one geodesic string  $o_\alpha$  in some class  $\alpha \in \pi_1^{inc}(X)$ . Moreover,  $o_\alpha$

is the unique geodesic string in its class, otherwise  $o_\beta$  would not be unique in its class. We prove that  $\alpha$  is indivisible, which will be a contradiction to  $\beta$  not being atomic and will complete the proof.

Suppose otherwise, so that  $\alpha_{x_0} = \rho^k$  for  $k > 1$  and  $\rho \in \pi_1(X, x_0)$ , (with the notation of Definition 1.21). Let  $o_\rho$  be a class  $\rho$ , closed  $h$ -geodesic string (where by slight abuse  $\rho$  also denotes the class in  $\pi_1^{inc}(X)$  corresponding to the based class  $\rho$ .) It is immediate that the  $k$  cover of  $o_\rho$ ,  $o_\rho^k$  represents  $\alpha$  and is a  $h$ -geodesic string. By the uniqueness,  $o_\alpha = o_\rho^k$ . But this contradicts multiplicity one of  $o_\alpha$ . So  $\alpha$  is indivisible.

□

*Proof of Corollary 1.31.* Let  $g_Z, g_Y$  be complete Riemannian metrics with  $g_Z$  taut, and  $g_Y$  not admitting contractible closed geodesics. Take the product metric  $g = g_Z \times g_Y$  on  $X = Z \times Y$ . For a class  $\beta \in \pi_1^{inc}(X)$  in the image of the inclusion  $\pi_1^{inc}(Z) \rightarrow \pi_1^{inc}(X)$ , the natural projection  $X \rightarrow Y$  is automatically a  $\beta$ -periodic fibration. Then the conclusion readily follows from Theorem 1.30. □

*Proof of Corollary 1.34.* The first two parts follow from Theorem 1.30, Lemma 1.33 and Example 4. The last part is deduced as follows. The product formula Theorem 8.1 gives us  $F(g_\phi, \beta) = 0$ . So for any  $\beta$ -regular  $g$  with finitely many class  $\beta$   $g$ -geodesic strings  $o$ , since  $\beta$  is indivisible, by Corollary 1.2 we have:

$$\chi^{S^1}(L_\beta X_\phi) = F(g, \beta) = \sum_o (-1)^{\text{morse } o} = 0.$$

Then the conclusion readily follows. □

*Proof of Theorem 1.24.* By Theorem 8.1 0 is certainly a value of the invariant  $F$ . We first prove that every negative rational number is the value of the invariant. Let  $p, q$  be positive integers. Let  $Y$  be a closed surface of genus  $(p+1) > 1$  with a hyperbolic metric  $g_Y$ , let  $Z$  be the genus 2 closed surface with a hyperbolic metric  $g_Z$  and let  $\beta_Z \in \pi_1^{inc}(Z)$  be the class represented by a  $2 \cdot q$ -fold covering of a simple closed loop representing a generator of the fundamental group of  $Z$ .

Let  $X = Y \times Z$  with the product metric  $g = g_Y \times g_Z$  and  $p : X \rightarrow Y$  the canonical projection. Taking  $\beta = (i_Z)_*(\beta_Z)$ , by Corollary 8.5 since  $(-1)^{\dim Y} = 1$ ,

$$F(g, \beta) = \chi(Y) \cdot F(g_Z, \beta_Z) = (-2p) \cdot \frac{1}{2q} = -\frac{p}{q}.$$

So we proved our first claim.

Let again  $p, q$  be positive integers. Let  $Y$  be closed surface of genus 2, with a hyperbolic metric  $g_Y$ . And let  $Z$  be a manifold satisfying  $F(g_Z, \beta_Z) = -\frac{p}{2q}$  for some  $\beta_Z$ -taut metric  $g_Z$  on  $Z$  and for some class  $\beta_Z \in \pi_1^{inc}(Z)$ . This exists by the discussion above. Let  $g = g_Y \times g_Z$  be the product metric on  $Y \times Z$ , and  $\beta$  as above. Analogously to the discussion above we get:

$$F(g, \beta) = \chi(Y) \cdot F(g_Z, \beta_Z) = (-2) \cdot \frac{-p}{2q} = \frac{p}{q}.$$

□

## APPENDIX A. FULLER INDEX AND SKY CATASTROPHES

In this appendix we recall the part of Fuller's periodic orbit index used in the paper. We also fix the notation for orbit strings and for continuation under small perturbations of vector fields.

Let  $X$  be a smooth manifold, and fix a complete auxiliary Riemannian metric  $h$  on  $X$ . Let

$$\mathcal{V}(X)$$

denote the space of smooth vector fields on  $X$ , with the  $C^0$ -topology on compact subsets.

Let

$$LX = C^\infty(S^1, X)$$

denote the smooth free loop space. For  $V \in \mathcal{V}(X)$ , define

$$S(V) = \{o \in LX \mid o \text{ is nonconstant, } \dot{o}(t) = T V(o(t)) \text{ for some } T > 0\}.$$

We call it the period of  $o$ , and write

$$T = T(o),$$

it is clearly uniquely determined.

The circle  $S^1 = \mathbb{R}/\mathbb{Z}$  acts on  $S(V)$  by reparametrization:

$$(s \cdot o)(t) = o(t + s).$$

We define the space of orbit strings by

$$\mathcal{O}(V) = S(V)/S^1.$$

If  $\beta$  is a free homotopy class in  $X$ , we write

$$S(V, \beta) \subset S(V), \quad \mathcal{O}(V, \beta) \subset \mathcal{O}(V)$$

for the corresponding subspaces. When no confusion is possible, we use the same letter  $o$  for a parametrized orbit and for its associated orbit string.

Let

$$q : LX \rightarrow LX/S^1$$

be the quotient map. If

$$U \subset LX/S^1,$$

we write

$$\widehat{U} = q^{-1}(U) \subset LX.$$

**Definition A.1.** We say that  $U$  is  *$h$ -controlled* if there are constants

$$0 < \ell_- < \ell_+ < \infty$$

and a compact set  $K \subset X$  such that every loop  $o \in \widehat{U}$  satisfies

$$\text{image}(o) \subset K, \quad \ell_- \leq \ell_h(o) \leq \ell_+.$$

The lower bound excludes collapse to constant loops; in the applications it is usually supplied by a nontrivial projected free homotopy class.

For an orbit string  $o \in \mathcal{O}(V)$ , its multiplicity

$$m(o) \in \mathbb{N}$$

is the unique positive integer such that  $o$  is the  $m(o)$ -fold cover of a primitive orbit string. Equivalently, if  $T(o)$  is the period of  $o$  and  $T_{\text{prim}}(o)$  is the period of the underlying primitive orbit string, then

$$m(o) = \frac{T(o)}{T_{\text{prim}}(o)}.$$



A.1. **Fuller index.** Suppose

$$N \subset \mathcal{O}(V)$$

is finite and consists of nondegenerate orbit strings. Fuller associates to  $(N, V)$  the rational number

$$i(N, V) = \sum_{o \in N} \frac{i(o)}{m(o)}.$$

Here  $i(o)$  is the fixed point index of the time- $T(o)$  return map of the flow of  $V$ , computed on a local hypersurface transverse to the image of  $o$ .

By a compact open subset  $N$  of an orbit-string space, we mean a subset which is compact and open in the relative topology. In this case Fuller again associates to this a rational index  $i(N, V)$ . The latter can be defined perturbatively, and using the formula as above.

In the special case of a nondegenerate closed geodesic orbit  $o$ , we have:

$$i(\tilde{o}) = (-1)^{\text{morse}(o)}.$$

Consequently, if  $N$  consists of finitely many nondegenerate closed orbits of the geodesic Reeb flow  $R^{\lambda_g}$  on  $C$ , we get:

$$i(N, V) = \sum_{o \in N} \frac{(-1)^{\text{morse}(o)}}{m(o)}.$$

A.2. **Continuation.** Let

$$\{V_t\}_{t \in [0,1]} \subset \mathcal{V}(X)$$

be a continuous family. Define

$$S(\{V_t\}) = \{(o, t) \in LX \times [0, 1] \mid o \in S(V_t)\},$$

and

$$(A.2) \quad \mathcal{O}(\{V_t\}) = S(\{V_t\})/S^1,$$

where  $S^1$  acts on the loop coordinate by reparametrization as before.

Let

$$\mathcal{N} \subset \mathcal{O}(\{V_t\})$$

be compact and open. Suppose that its endpoint slices are

$$N_0 = \mathcal{N} \cap \mathcal{O}(V_0), \quad N_1 = \mathcal{N} \cap \mathcal{O}(V_1).$$

Then

$$(A.3) \quad i(N_0, V_0) = i(N_1, V_1).$$

We refer to this as the **basic invariance** of the Fuller index.

For applications, we need a local version of this statement under small perturbations. This is a variation of [11, Lemma 4.1], with the added control needed for noncompact  $X$ .

**Lemma A.4** (Isolation lemma). *Let  $h$  be a complete auxiliary Riemannian metric on  $X$ . Let  $V_0 \in \mathcal{V}(X)$ , be a nonsingular vector field, and let*

$$N_0 \subset \mathcal{O}(V_0)$$

*be compact and open. Let*

$$U \subset LX/S^1$$

*be an  $h$ -controlled open neighborhood of  $N_0$  such that*

$$\overline{U} \cap \mathcal{O}(V_0) = N_0.$$

Let  $W$  be an open neighborhood of  $N_0$  satisfying

$$\overline{W} \subset U.$$

Then there exists  $\epsilon > 0$  with the following property. If  $V$  is a smooth vector field satisfying

$$\sup_{x \in K} |V(x) - V_0(x)|_h < \epsilon,$$

where  $K$  is the compact set from the  $h$ -control of  $U$ , then

$$\mathcal{O}(V) \cap U \subset W.$$

In particular,

$$\mathcal{O}(V) \cap U$$

is compact and open in  $\mathcal{O}(V)$ .

*Proof.* Let  $K$ ,  $\ell_-$ , and  $\ell_+$  be the compact set and length bounds from the  $h$ -control of  $U$ . Since  $V_0$  is nonsingular and  $K$  is compact, there are constants

$$0 < B < A < \infty$$

such that

$$B \leq |V_0(x)|_h \leq A$$

for all  $x \in K$ . We choose  $\epsilon < B/2$ . Then every vector field  $V$  satisfying

$$\sup_K |V - V_0|_h < \epsilon$$

obeys

$$|V(x)|_h \geq B/2, \quad |V(x)|_h \leq A + \epsilon$$

for all  $x \in K$ .

Suppose the conclusion failed for the chosen  $W$ . Then there would be vector fields

$$V_j \rightarrow V_0$$

uniformly on  $K$ , and orbit strings

$$o_j \in (\mathcal{O}(V_j) \cap U) - W.$$

Choose parametrized representatives, still denoted  $o_j$ , satisfying

$$(A.5) \quad \dot{o}_j(s) = T_j V_j(o_j(s)).$$

Since  $o_j \in \widehat{U}$ , controlledness gives

$$\text{image}(o_j) \subset K, \quad \ell_- \leq \ell_h(o_j) \leq \ell_+.$$

For  $j$  sufficiently large,

$$B/2 \leq |V_j|_h \leq A + \epsilon$$

on  $K$ . Hence

$$\frac{\ell_-}{A + \epsilon} \leq T_j \leq \frac{2\ell_+}{B}.$$

The velocities are also uniformly bounded:

$$|\dot{o}_j(s)|_h = T_j |V_j(o_j(s))|_h \leq \frac{2\ell_+(A + \epsilon)}{B}.$$

Thus the  $o_j$  are equicontinuous and have images in the fixed compact set  $K$ . By the Arzelà–Ascoli theorem, after passing to a subsequence,

$$o_j \rightarrow o_\infty$$

uniformly. After passing to a further subsequence,  $T_j \rightarrow T_\infty$ , with  $T_\infty > 0$ .

Hence  $o_j$  is a Cauchy sequence in the  $C^1$  space of maps, by the defining equation (A.5). Then  $C^1$  completeness gives that

$$\dot{o}_\infty = T_\infty V_0(o_\infty).$$

Thus  $o_\infty$  is a closed  $V_0$ -orbit string. Moreover

$$o_\infty \in \overline{U} - W.$$

This contradicts

$$\overline{U} \cap \mathcal{O}(V_0) = N_0 \subset W.$$

Therefore

$$\mathcal{O}(V) \cap U \subset W$$

for all  $V$  sufficiently close to  $V_0$  on  $K$ .

It remains to prove compactness. Let  $o_j$  be any sequence in  $\mathcal{O}(V) \cap U$ . The same estimates as above give uniformly bounded periods and velocities, with all images contained in  $K$ . Arzelà–Ascoli therefore gives a convergent subsequence. Its limit is again a closed  $V$ -orbit. Since the inclusion just proved gives

$$\mathcal{O}(V) \cap U = \mathcal{O}(V) \cap \overline{W}$$

and  $\overline{W} \subset U$ , the limit lies in  $\mathcal{O}(V) \cap U$ . Thus  $\mathcal{O}(V) \cap U$  is compact. It is open because  $U$  is open.  $\square$

We call such an  $\epsilon$  a **Fuller constant** for the data

$$(V_0, N_0, U, W).$$

It also depends on the auxiliary control data  $K, \ell_-, \ell_+$ , which will always be understood from the chosen  $U$ .

**Lemma A.6** (Fuller continuation). *Let  $V_0, N_0, U, W$ , and  $\epsilon$  be as in Lemma A.4. Let*

$$\{V_t\}_{t \in [0,1]}, \quad V_{t=0} = V_0$$

*be a continuous family such that for every  $t \in [0, 1]$ ,*

$$\sup_{x \in K} |V_t(x) - V_0(x)|_h < \epsilon,$$

*where  $K$  is the compact set from the  $h$ -control of  $U$ . Define*

$$\mathcal{N} = (U \times [0, 1]) \cap \mathcal{O}(\{V_t\}).$$

*Then  $\mathcal{N}$  is compact and open in the orbit cobordism  $\mathcal{O}(\{V_t\})$ . Its endpoint slices are*

$$N_0 = U \cap \mathcal{O}(V_0), \quad N_1 = U \cap \mathcal{O}(V_1).$$

*Consequently,*

$$i(N_0, V_0) = i(N_1, V_1).$$

*We call  $N_1$  the Fuller continuation of  $N_0$ .*

*Proof.* Since  $U$  is open,

$$\mathcal{N} = (U \times [0, 1]) \cap \mathcal{O}(\{V_t\})$$

is open in  $\mathcal{O}(\{V_t\})$ .

By Lemma A.4, applied to each  $V_t$ , we have

$$\mathcal{O}(V_t) \cap U \subset W$$

for every  $t \in [0, 1]$ . Since

$$\overline{W} \subset U,$$

this implies

$$\mathcal{N} = (\overline{W} \times [0, 1]) \cap \mathcal{O}(\{V_t\}).$$

The compactness then works as in Lemma A.4. The endpoint slices are

$$\mathcal{N} \cap \mathcal{O}(V_0) = U \cap \mathcal{O}(V_0) = N_0$$

and

$$\mathcal{N} \cap \mathcal{O}(V_1) = U \cap \mathcal{O}(V_1) = N_1.$$

Fuller's basic invariance applied to the compact open cobordism  $\mathcal{N}$  gives

$$i(N_0, V_0) = i(N_1, V_1).$$

□

*Example 5* (Geodesic flows and the Sasaki metric). Let  $(Y, g_Y)$  be a complete Riemannian manifold, and let

$$C_g = S^*Y$$

be its unit cotangent bundle with Liouville contact form  $\lambda_{g_Y}$ . The Reeb vector field  $R^{\lambda_{g_Y}}$  is the geodesic Reeb field.

Let  $g_S$  be the Sasaki metric on  $S^*Y$ . It is defined using the Levi-Civita connection of  $g_Y$ . Thus

$$TC = T^{\text{vert}}C \oplus T^{\text{hor}}C$$

is an orthogonal splitting, where

$$T^{\text{vert}}C = \ker(\text{pr}_*)$$

and  $T^{\text{hor}}C$  is the Levi-Civita horizontal subbundle. If  $\xi, \eta \in T(S^*Y)$  are horizontal, then

$$g_S(\xi, \eta) = g_Y(\text{pr}_* \xi, \text{pr}_* \eta),$$

and the vertical part is measured using the metric induced by  $g_Y$  on the cotangent fibers.

The basic property needed in the paper is the following. If  $o$  is a nonzero constant-speed  $g_Y$ -geodesic and  $\tilde{o}$  is its unit cotangent lift, then

$$(A.7) \quad \ell_{g_S}(\tilde{o}) = \ell_{g_Y}(o).$$

Indeed, the covector component of  $\tilde{o}$  is parallel along  $o$ , so the vertical part of  $\tilde{o}$  vanishes and the Sasaki speed equals the base speed.

If  $\beta \in \pi_1^{\text{inc}}(Y)$  and  $U \subset S(g_Y, \beta)$  is  $g_Y$ -length bounded, then the set of canonical lifts

$$\tilde{U} \subset S(R^{\lambda_{g_Y}})$$

is  $g_S$ -length bounded by (A.7).

## APPENDIX B. A LOCAL FIBERED CALCULATION OF $F$

We prove in this section the technical lemma computing the local index, used in Theorem 8.1.

Let

$$P : S^*X \rightarrow \mathbb{R}$$

be the verticality function as in the proof of Theorem 8.1,

$$P(\xi) = |\pi^{\text{vert}} v_\xi|_g^2.$$

Here  $v_\xi$  is the  $g$ -dual unit vector to  $\xi$ , and

$$\pi^{\text{vert}} : TX \rightarrow T^{\text{vert}}X$$

is the  $g$ -orthogonal projection. The notation  $\beta$  and  $\beta_Z$  is likewise as in Theorem 8.1.

**Lemma B.1** (Local product calculation). *Let*

$$Z \hookrightarrow X \xrightarrow{p} Y$$

*be a parallel Riemannian submersion. Let*

$$\tilde{f} = f \circ p \circ \text{pr}$$

*for a Morse function  $f : Y \rightarrow \mathbb{R}$ , and define the vector field on  $S^*X$*

$$V_\varepsilon = R^{\lambda_g} - \varepsilon \nabla_{g_S}(P + \tilde{f}).$$

*Fix*

$$y \in \text{Crit}(f),$$

*and let*

$$N_{\varepsilon, y}$$

*be the isolated set of closed  $V_\varepsilon$ -orbit strings lying over the fiber*

$$Z_y = p^{-1}(y),$$

*as in the proof of Theorem 8.1. Then*

$$i(N_{\varepsilon, y}, V_\varepsilon) = (-1)^{\dim Y + \text{morse}(y)} \sum_{\alpha \in S_y} F(g_y, \alpha),$$

*where  $S_y$  is the  $\pi_1(Y, y)$ -orbit of the corresponding fiber representative in  $\pi_1^{\text{inc}}(Z_y)$ . In particular, if the fibration is holonomy-finite in the relevant class, the sum is finite.*

*Proof.* Let

$$T^{\text{vert}}X = \ker(dp), \quad H = (T^{\text{vert}}X)^{\perp_g}.$$

Since the submersion is parallel, the two distributions

$$T^{\text{vert}}X \quad \text{and} \quad H$$

are preserved by the Levi-Civita connection. Since  $T^{\text{vert}}X$  and  $H = (T^{\text{vert}}X)^{\perp_g}$  are parallel complementary distributions, the local de Rham decomposition theorem gives a product decomposition

$$p^{-1}(D_y) \cong Z_y \times D_y, \quad g = g_y \oplus g_Y|_{D_y},$$

for a normal ball neighborhood  $D_y \subset Y$  centered at  $y$ . See Kobayashi–Nomizu [15, Chapter IV].

For this local calculation we suppress the fixed- $C$  identification and write  $S^*X$  for the  $g$ -unit cotangent bundle. Let

$$C^{\text{vert}} = \{P = 1\} \subset S^*X$$

be the vertical locus. Over  $Z_y$ , this locus is naturally identified with

$$S^*Z_y \subset S^*X.$$

Near

$$S^*Z_y \subset C^{\text{vert}},$$

the product decomposition gives the normal-bundle model

$$(B.2) \quad S^*Z_y \times D_y \times B_y^*,$$

where

$$B_y^* \subset T_y^*Y$$

is a small ball in the horizontal cotangent direction. In this model, the vertical locus is

$$S^*Z_y \times D_y \times \{0\}.$$

A covector near the vertical locus decomposes as

$$\xi = \xi_{\text{vert}} + \eta, \quad \eta \in B_y^*.$$

The unit condition is

$$|\xi_{\text{vert}}|_{g_y^*}^2 + |\eta|_{g_Y^*}^2 = 1.$$

Therefore

$$P(\xi) = |\xi_{\text{vert}}|_{g_y^*}^2 = 1 - |\eta|_{g_Y^*}^2.$$

Thus  $P$  has a Morse–Bott maximum along  $C^{\text{vert}}$ .

In the local model (B.2),

$$N_{\varepsilon, y} = N_y^{\text{fib}} \times \{y\} \times \{0\},$$

where  $N_y^{\text{fib}}$  is the relevant closed orbit set of the fiber geodesic Reeb flow on  $S^*Z_y$ .

We now describe the local return map at an element of  $N_{\varepsilon, y}$ . Along the vertical locus over  $Z_y$ , the fiber component of  $V_\varepsilon$  is the geodesic Reeb field  $R^{\lambda_{g_y}}$  on  $S^*Z_y$ . The base component is

$$-\varepsilon \nabla_{g_Y} f$$

on  $D_y$ . The normal covector component is the outward radial component coming from

$$-\varepsilon \nabla_{g_S} P$$

on  $B_y^*$ .

Let

$$o \in N_y^{\text{fib}},$$

and choose a product transverse section

$$\Sigma = \Sigma^{\text{fib}} \times D_y \times B_y^*,$$

where

$$\Sigma^{\text{fib}} \subset S^*Z_y$$

is transverse to the fiber Reeb flow near  $o$ . With respect to this product section, the local return map of  $V_\varepsilon$  is the product

$$\Phi_{\text{fib}} \times \psi_y \times \kappa.$$

Here:

- $\Phi_{\text{fib}}$  is the local return map of the fiber geodesic flow generated by  $R^{\lambda_{g_y}}$ ;
- $\psi_y$  is the small-time map of

$$-\varepsilon \nabla_{g_Y} f$$

near  $y$ ;

- $\kappa$  is the normal return map induced by

$$-\varepsilon \nabla_{g_S} P$$

near  $0 \in B_y^*$ .

The fixed point index is multiplicative under products, Dold [9, Chapter VII, Section 5]. Hence

$$i_{\text{fp}}(o; V_\varepsilon) = i_{\text{fp}}(o; R^{\lambda_{g_y}}) \cdot i_{\text{fp}}(\psi_y, y) \cdot i_{\text{fp}}(\kappa, 0).$$

Since  $\psi_y$  is the positive small-time map of  $-\varepsilon \nabla_{g_Y} f$ , the map  $\text{id} - \psi_y$  has the same local degree as  $\nabla_{g_Y} f$ . Thus, because  $y$  is a Morse critical point,

$$i_{\text{fp}}(\psi_y, y) = (-1)^{\text{morse}(y)}.$$

Since the normal component of

$$-\varepsilon \nabla_{g_S} P$$

has outward radial linearization on a vector space of dimension  $\dim Y$ , the map  $\text{id} - \kappa$  has local degree  $(-1)^{\dim Y}$ . Hence

$$i_{\text{fp}}(\kappa, 0) = (-1)^{\dim Y}.$$

Therefore

$$i_{\text{fp}}(o; V_\varepsilon) = (-1)^{\dim Y + \text{morse}(y)} i_{\text{fp}}(o; R^{\lambda_{g_y}}).$$

The multiplicity of  $o$  is unchanged by taking the product with the isolated base and normal fixed points. Thus

$$\frac{i_{\text{fp}}(o; V_\varepsilon)}{m(o)} = (-1)^{\dim Y + \text{morse}(y)} \frac{i_{\text{fp}}(o; R^{\lambda_{g_y}})}{m(o)}.$$

Summing over all fiber orbit strings in the local block gives

$$i(N_{\varepsilon, y}, V_\varepsilon) = (-1)^{\dim Y + \text{morse}(y)} \sum_{\alpha \in S_y} F(g_y, \alpha).$$

This proves the claim.  $\square$

### APPENDIX C. SKY CATASTROPHES

We now recall the topological formulation of sky catastrophes. This is the version used in Fuller's theory and in the applications above.

**Definition C.1.** Let

$$\{V_t\}_{t \in [0,1]} \subset \mathcal{V}(X)$$

be a continuous family. We say that the family has a sky catastrophe in class  $\beta$  if there is an endpoint orbit string

$$y \in \mathcal{O}(V_0, \beta) \cup \mathcal{O}(V_1, \beta) \subset \mathcal{O}(\{V_t\}, \beta)$$

such that no compact open subset of  $\mathcal{O}(\{V_t\}, \beta)$  contains  $y$ .

Equivalently, the endpoint orbit string  $y$  cannot be included in any compact continuation block through the family.

For suitably regular families, this agrees with the preliminary Definition 1.17.

### ACKNOWLEDGEMENTS

I am grateful to the IAS for resources provided during my stay as a member, where some of this paper was written. Thanks also to Mohammed Abouzaid, Semon Rezchikov and Viktor Ginzburg for interesting related discussions.

### REFERENCES

- [1] J. C. ALEXANDER, *Closed geodesics on certain surfaces of revolution*, J. Geom. Symmetry Phys., 8 (2006), pp. 1–16.
- [2] M. T. ANDERSON, *Metrics of negative curvature on vector bundles*, Proc. Am. Math. Soc., 99 (1987), pp. 357–363.
- [3] D. BAO, S.-S. CHERN, AND Z. SHEN, *An introduction to Riemann-Finsler geometry*, vol. 200 of Grad. Texts Math., New York, NY: Springer, 2000.
- [4] I. BELEGRADEK, *Aspherical manifolds, relative hyperbolicity, simplicial volume and assembly maps*, Algebr. Geom. Topol., 6 (2006), pp. 1341–1354.
- [5] K. S. BROWN, *Euler characteristics of discrete groups and  $g$ -spaces*, Inventiones Mathematicae, 27 (1974), pp. 229–264.
- [6] ———, *Cohomology of Groups*, vol. 87 of Graduate Texts in Mathematics, Springer, 1982.
- [7] E. CAPONIO, M. A. JAVALOYES, AND A. MASIELLO, *On the energy functional on finsler manifolds and applications to stationary spacetimes*, Mathematische Annalen, 351 (2011), pp. 365–392.
- [8] M. P. DO CARMO, *Differential Geometry of Curves and Surfaces*, Prentice-Hall, Englewood Cliffs, NJ, 1976.
- [9] A. DOLD, *Lectures on Algebraic Topology*, vol. 200 of Die Grundlehren der mathematischen Wissenschaften, Springer-Verlag, Berlin, 1972.
- [10] F. FULLER, *Note on trajectories in a solid torus.*, Ann. Math. (2), 56 (1952), pp. 438–439.
- [11] ———, *An index of fixed point type for periodic orbits.*, Am. J. Math., 89 (1967), pp. 133–145.
- [12] R. S. HAMILTON, *The Ricci flow on surfaces*, in Mathematics and General Relativity (Santa Cruz, CA, 1986), vol. 71 of Contemporary Mathematics, American Mathematical Society, Providence, RI, 1988, pp. 237–262.
- [13] A. HATCHER, *Algebraic topology*, Cambridge: Cambridge University Press, 2002.

- [14] N. HINGSTON AND H.-B. RADEMACHER, *Resonance for loop homology of spheres*, Journal of Differential Geometry, 93 (2013), pp. 133–174.
- [15] S. KOBAYASHI AND K. NOMIZU, *Foundations of Differential Geometry*, vol. 1 of Interscience Tracts in Pure and Applied Mathematics, Interscience Publishers, New York, 1963.
- [16] H. LIU AND Y. XIAO, *Resonance identity and multiplicity of non-contractible closed geodesics on Finsler  $\mathbb{R}^n$* , Advances in Mathematics, 318 (2017), pp. 158–190.
- [17] G. LU, *Methods of infinite dimensional morse theory for geodesics on finsler manifolds*, 2012.
- [18] F. MERCURI, *The critical points theory for the closed geodesics problem*, Mathematische Zeitschrift, 156 (1977), pp. 231–246.
- [19] G. MESS, *Examples of Poincaré duality groups*, Proceedings of the American Mathematical Society, 110 (1990), pp. 1145–1146.
- [20] J. MILNOR, *Morse theory*, Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, No. 51, Princeton University Press, Princeton, N.J., 1963.
- [21] R. S. PALAIS, *Morse theory on hilbert manifolds*, Topology, 2 (1963), pp. 299–340.
- [22] A. PREISSMANN, *Quelques propriétés globales des espaces de Riemann*, Commentarii Mathematici Helvetici, (1942).
- [23] H.-B. RADEMACHER, *On the average indices of closed geodesics*, Journal of Differential Geometry, 29 (1989), pp. 65–83.
- [24] Y. SAVELYEV, *Extended Fuller index, sky catastrophes and the Seifert conjecture*, International Journal of mathematics, 29 (2018).
- [25] W. P. THURSTON, *On the geometry and dynamics of diffeomorphisms of surfaces*, Bull. Am. Math. Soc., New Ser., 19 (1988), pp. 417–431.
- [26] ———, *Hyperbolic structures on 3-manifolds, II: Surface groups and 3-manifolds which fiber over the circle*, arXiv preprint, (1998). Based on a 1986 preprint.
- [27] C. T. C. WALL, *Rational euler characteristics*, Proceedings of the Cambridge Philosophical Society, 57 (1961), pp. 182–184.

Email address: yasha.savelyev@gmail.com

FACULTY OF SCIENCE, UNIVERSITY OF COLIMA, MEXICO