

GLOBAL FUKAYA CATEGORY II: APPLICATIONS

YASHA SAVELYEV

ABSTRACT. In part I, we constructed a natural homotopy coherent action of $\text{Ham}(M, \omega)$ on the Fukaya category of a closed monotone symplectic manifold. Here we show that this action is generally homotopically non-trivial by a sample computation. One principal application is differential geometric, and the other is about algebraic K -theory of the integers. We find new curvature constraint phenomena for smooth and singular $\mathcal{G}$ -connections on principal $\mathcal{G}$ -bundles over S^4 , where $\mathcal{G}$ is $\text{PU}(2)$ or $\text{Ham}(S^2)$. Even for the classical group $\text{PU}(2)$, these phenomena are inaccessible to known techniques like the Yang-Mills theory. The computation is the geometric component used to show that the categorified algebraic K -theory of the integers, defined in [6] following Toën, admits a $\mathbb{Z}$ injection in degree 4. In particular, the natural trace maps to the standard algebraic K -theory have a kernel in higher degrees.

1. INTRODUCTION

Given a smooth Hamiltonian fibration $M \hookrightarrow P \rightarrow X$, with M monotone, and X any smooth manifold, in Part I, that is in [8], we have constructed a functor which may be understood as giving a fibration of A_∞ categories over X , with fiber the Fukaya category of M . This construction is to the Fukaya category as vector bundles are to vector spaces.

More specifically, denote by $\Delta(X)$ the smooth simplex category of X , whose objects are smooth maps $\Sigma : \Delta^d \rightarrow X$. See Section 3 for more details. Denote by $A_\infty \text{Cat}_k^{\mathbb{Z}_2}$ the category of A_∞ categories over a commutative ring k , with $\mathbb{Z}_2$ graded hom complexes. ($\mathbb{Z}_2$ grading is part of the structure not a property.)

Then in Part I we construct a functor:

$$(1.1) \quad F_P : \Delta(B \text{Ham}(M, \omega)) \rightarrow A_\infty \text{Cat}_k^{\mathbb{Z}_2}.$$

Here we show, by a sample computation that this functor is (generally) “homotopically” non-trivial, and give some applications for this fact. ¹

We will show non-triviality of F_P , by showing homotopy non-triviality of the associated “classifying map”

$$(1.2) \quad f_P : B \text{Ham}(M, \omega) \rightarrow |\mathbb{S}|,$$

where $|\mathbb{S}|$ denotes the space of ∞ -categories, and where f_P maps into the component of $\text{NFuk}(M, \omega)$, with the latter denoting the Faonte-Lurie A_∞ nerve of $\text{Fuk}(M, \omega)$,

2000 *Mathematics Subject Classification.* 53D37, 55U35, 53C21.

Key words and phrases. Fukaya category, action of the group of Hamiltonian symplectomorphisms, infinity categories, singular connections, curvature constraints.

¹Homotopically non-trivial, more specifically means that the concordance class of the functor is non-trivial, where concordance of such functors is defined in Part I.

(see Appendix A.4 of Part I.) This classifying map is also constructed in Part I.

As explained in Part I, the homotopy non-triviality of F_P , may be understood to say that there is a non-trivial “homotopy coherent” action of $\text{Ham}(M, \omega)$ on $\text{Fuk}(M, \omega)$. Existence of such an action in a somewhat weaker form (on the level E_2 algebras rather than space level) has been independently conjectured by Teleman, ICM 2014, post-factum of first appearance of Part I.

The classifying map f_P , also induces a certain kind of simplicial fibration over the smooth singular set $X_\bullet$ of X , having the structure of a Cartesian fibration (in this case equivalently categorical fibration):

$$\text{NFuk}(M, \omega) \hookrightarrow \text{Fuk}_\infty(P) \xrightarrow{p_\bullet} X_\bullet.$$

This is called the global Fukaya category of P . We show that for P a non-trivial Hamiltonian S^2 fibration over S^4 , the maximal Kan sub-fibration of $\text{Fuk}_\infty(P)$ is non-trivial. In particular, $\text{Fuk}_\infty(P)$ is non-trivial as a Cartesian fibration and so f_P is homotopically non-trivial. In particular, this gives:

Theorem 1.1. *The homomorphism*

$$\mathbb{Z} = \pi_4(\text{BHam}(S^2, \omega)) \xrightarrow{h} \pi_4(\mathbb{S}, \text{NFuk}(S^2, \omega)),$$

induced by the map (1.2), is injective.

The theorem above is proved via geometric analysis as the main ingredient, and more specifically a certain computation from [7], using the notion of quantum Maslov classes. Note that $\pi_4(\text{BHam}(S^2, \omega)) \simeq \pi_3(SO(3)) = \mathbb{Z}$, by Smale’s theorem. Restricting to $k = \mathbb{Z}$, the theorem above, together with some additional, purely algebraic ideas (forthcoming in a future work), prove that there is a natural injection

$$\mathbb{Z} \rightarrow K_4^{\text{Cat}, \mathbb{Z}_2}(\mathbb{Z}),$$

where the right hand side is the categorified algebraic K -theory of the integers, introduced in [6], following ideas of Toën [10]. As explained in [6], this implies that the natural trace maps from $K_n^{\text{Cat}, \mathbb{Z}_2}(\mathbb{Z})$ to the classical algebraic K -theory of $\mathbb{Z}$ have a kernel. So we get a connection from Floer theory all the way to arithmetic.

1.1. An application to curvature bounds of singular and smooth connections.

1.1.1. *A non-metric measure of curvature.* Let G as above be a Frechet Lie group, we denote by $\text{lie } G$ its Lie algebra and let

$$\mathbf{n} : \text{lie } G \rightarrow \mathbb{R}$$

be an Ad invariant Finsler norm. For a principal G -bundle P over a Riemann surface (S, j) , and given a G connection $\mathcal{A}$ on P define a 2-form $\alpha_{\mathcal{A}}$ on S by:

$$\alpha_{\mathcal{A}}(v, jv) = \mathbf{n}(R_{\mathcal{A}}(v, jv)),$$

where $R_{\mathcal{A}}$ is the curvature 2-form of $\mathcal{A}$. More specifically, the latter form has the properties:

$$R_{\mathcal{A}}(v, w) \in \text{lie Aut } P_z,$$

for $z \in S$, $v, w \in T_z S$, P_z the fiber of P over z , $\text{Aut } P_z \simeq G$ the group of G -bundle automorphisms of P_z , and where $\simeq$ means non-canonical group isomorphism.

Define

$$(1.3) \quad \text{area}_n(\mathcal{A}) := \int_S \alpha_{\mathcal{A}}.$$

In the case $\mathcal{A}$ is singular with singular set C , $\alpha_{\mathcal{A}}$ is defined on $S - C$, so we define

$$\text{area}_n(\mathcal{A}) := \int_{S-C} \alpha_{\mathcal{A}},$$

with the right-hand side now being an extended integral. This area is a non-metric measurement meaning that no Riemannian metric on S is needed.

It is possible to extend the functional above to a functional on the space $\mathcal{C}$ of G -connections on principal G bundles $P \rightarrow \Delta^n$. It may seem that Δ^n has no connection to Riemann surfaces, but in fact there is an intriguing such connection. Let $\mathcal{S}_d$ denote the universal curve over $\overline{\mathcal{R}}_d$ - the moduli space of complex structures on the disk with $d + 1$ punctures on the boundary. And let $\mathcal{S}_d^\circ$ denote $\mathcal{S}_d$, with nodal points of the fibers removed. Then it is shown in Part I that there are certain axiomatized systems of maps:

$$u : \mathcal{S}_d^\circ \rightarrow \Delta^n, d, n \text{ varying.}$$

Such a system is uniquely determined up to suitable homotopy, and is referred to as $\mathcal{U}$.

There is then a natural functional:

$$(1.4) \quad \text{area}_{\mathcal{U}} : \mathcal{C} \rightarrow \mathbb{R}_{\geq 0},$$

defined with respect to a choice of $\mathcal{U}$, see Section 10. When $n = 2$ it is just the area functional as previously defined.

1.1.2. Curvature bounds. The following basic result can be formally understood as a corollary of Theorem 1.6. But it is more elementary to see it as a corollary of Theorem 10.4.

Corollary 1.2 (Of Theorem 1.6 and of Theorem 10.4). *Let P be a non-trivial $\text{PU}(2)$ (or $\text{Ham}(S^2)$) bundle $P \rightarrow S^4$, let $D_{\pm}^4$ be as above and let $\mathcal{A}$ be a smooth $\text{PU}(2)$ or $\text{Ham}(S^2, \omega)$ connection on P . For any $\mathcal{U}$, suppose that $\text{area}_{\mathcal{U}}(\mathcal{A})|_{D_-^4} < \frac{1}{2}$, then $\text{area}_{\mathcal{U}}(\mathcal{A})|_{D_+^4} \geq \frac{1}{2}$.*

Remark 1.3. *The proof of even the above corollary traverses the entirety of the theory here. As this is a very elementary result we may hope for a simpler argument. In the case of $\text{PU}(2)$, one idea might be to replace Floer theory, used here, by the technically simpler mathematical Yang-Mills theory over surfaces [1]. If we want to mimic the argument presented in this paper, then we should first extend Yang-Mills theory to work with G -bundles over surfaces with corners and holonomy constraints over boundary. This should be possible, but beyond this things are unclear, since, as we also use certain abstract algebraic topology to glue various analytic data, and it is not clear how this would work for Yang-Mills theory.*

1.1.3. *Abstract resolutions of singular connections.* We can use the computation of Theorem 1.1 to obtain lower bounds for the curvature of certain types of singular connections.

Definition 1.4. Let $G \hookrightarrow P \rightarrow X$ be a principal G bundle, where G is a Frechet Lie group. A **singular G -connection** on P is a closed subset $C \subset X$, and a smooth Ehresmann G -connection $\mathcal{A}$ on $P|_{X-C}$.

The above definition is basic, as one often puts additional conditions, see for instance [2], [9].

Avoiding generality, suppose that $\mathcal{A}$ is a singular G -connection on a principal G -bundle $P \rightarrow S^n$, with a single singularity x_0 . We will show that it is possible to control the curvature of the singular connection $\mathcal{A}$ if we impose a certain structure on the singularity of $\mathcal{A}$. The simplest way to do this is to ask for existence of a certain kind of abstract resolution.

First, a simplicial G -connection on $\mathcal{D}$ on P , as defined in Section 10.1, is basically a functorial assignment of a smooth G connection $\mathcal{D}_\Sigma$ on Σ^*P for each smooth

$$\Sigma : \Delta^n \rightarrow S^n.$$

Definition 1.5. For $\mathcal{A}, P$ as above a **simplicial resolution** of $\mathcal{A}$ is a simplicial G connection $\mathcal{A}^{res}$ on P , with the following property. Let $\Sigma_0 : \Delta^n \rightarrow S^n$ represent the generator of $\pi_n(S^n, x_0)$ (cf. Appendix A), then

$$(\Sigma_0|_{\text{interior } \Delta^n})^* \mathcal{A} = \mathcal{A}_{\Sigma_0}^{res}|_{\text{interior } \Delta^n}.$$

In the following theorem $G = PU(2)$, $n = 4$ and the norm $\mathfrak{n}$ on $\text{lie } PU(2)$ will be taken to be the operator norm, normalized so that the Finsler length of the shortest one parameter subgroup from id to $-id$ is $\frac{1}{2}$. We will omit $\mathfrak{n}$ in notation. We also impose an additional constraint on $\mathcal{A}^{res}$, so that the curvature at “ ∞ ” is bounded by a threshold, which means the following. Let

$$\Sigma_\infty : \Delta^4 \rightarrow x_0$$

be the constant map. Suppose that:

$$\text{area}_{\mathcal{U}} \mathcal{A}_{\Sigma_\infty}^{res} < 1/2,$$

and suppose for simplicity that $\mathcal{A}_{\Sigma_\infty}^{res}$ is trivial along the edges of Δ^4 , later on this condition is relaxed, see Proposition 10.4. (This condition can be completely removed but at the cost of significant additional complexity.)

We say in this case that $\mathcal{A}^{res}$ is a **sub-quantum resolution**. The following is proved in Section 10.

Theorem 1.6. Let $P \rightarrow S^4$ be a non-trivial principal $PU(2)$ bundle. Let $\mathcal{A}$ be a singular $PU(2)$ -connection on P with a single singularity at x_0 . Then for any sub-quantum resolution $\mathcal{A}^{res}$ of $\mathcal{A}$ and for any $\mathcal{U}$ as above

$$\text{area}_{\mathcal{U}}(\mathcal{A}_{\Sigma_0}^{res}) \geq 1/2.$$

The theorem has certain extensions to Hamiltonian singular connections $\mathcal{A}$, understanding P as a principal $\text{Ham}(S^2)$ bundle, see Section 10.

Example 1.7. Let P be as above, and $\mathcal{A}'$ be an ordinary smooth $PU(2)$ connection on P . Express S^4 as a union of sub-balls $D_{\pm}^4 \subset S^4$, intersecting only in the boundary. Suppose that we have the property that $\text{area}_{\mathcal{U}}(\mathcal{A}')|_{D_{-}^4} < \frac{1}{2}$. Let $\mathcal{A}$ be the singular connection on P obtained as the push-forward of $\mathcal{A}'$ by the bundle map $\tilde{q} : P \rightarrow P$ over the singular smooth map $q : S^4 \rightarrow S^4$ taking D_{-}^4 to a single point $\infty \in S^4$, with $q|_{\text{interior } D_{+}^4}$ an immersion. Then $\mathcal{A}$ has a sub-quantum resolution $\mathcal{A}^{res}$ by construction. In this case, the theorem above simply yields that $\text{area}_{\mathcal{U}}(\mathcal{A}'_{D_{+}^4}) \geq 1/2$.

Remark 1.8. There are possible physical interpretations for singular connections, as appearing in the context here. A $PU(2)$ connection $\mathcal{A}$ on P in physical terms represents a Yang-Mills field on the space-time S^4 . When the space-time has a black hole singularity, the fields solving the Einstein-Yang-Mills equations (mathematically connections as above) likewise develop singularities. There is a wealth of physics literature on this subject, here is one sample reference [4]. As quantum gravity is often related to simplicial ideas, it is not inconceivable that the mathematical sub-quantum resolution condition above also has a (quantum gravity theoretic) physical interpretation.

At this point the reader may be curious why Theorem 1.1 has something to do with Theorem 1.6. We cannot give the full story, but the idea is that the Cartesian fibration $\text{Fuk}_{\infty}(P)$ only sees the principal bundle P (and its curvature) by the behavior of certain holomorphic curves. When one has the sub-quantum condition on the curvature of $\mathcal{A}_{\Sigma_{\infty}}^{res}$, certain holomorphic curves are ruled out so that from the view point of $\text{Fuk}_{\infty}(P)$, $\mathcal{A}_{\Sigma_{\infty}}^{res}$ is the trivial connection, (its curvature is undetectable) but $\text{Fuk}_{\infty}(P)$ is non-trivial as a fibration so that the aforementioned holomorphic curves and consequently curvature must appear elsewhere.

1.2. First quantum obstruction and smooth invariants. We present here a construction of an integer valued smooth invariant which is based on our theory. This is probably just the beginning of the story for invariants of smooth manifolds based on Floer-Fukaya theory.

First we discuss a more general setup. Let $M \hookrightarrow P \xrightarrow{P} X$ be a Hamiltonian M -bundle, as previously. Let

$$\text{Fuk}_{\infty}(P) \xrightarrow{P_{\bullet}} X_{\bullet}$$

be the associated Cartesian fibration, and let

$$K(P) \rightarrow X_{\bullet}$$

be its maximal Kan sub-fibration as in Lemma 4.1. Then $|K(P)| \rightarrow X$ is a Serre fibration, where $|K(P)|$ is the geometric realization.

Define

$$\text{q-obs}(P) \in \mathbb{N} \sqcup \{\infty\},$$

to be the degree of the first obstruction to a section of $|K(P)|$. That is $\text{q-obs}(P)$ is the smallest integer n such that there is no section of $|K(P)|$ over the n skeleton of X , with respect to some chosen CW structure. This is independent of the choice of the CW structure, as any pair of CW structures on X are filtered (using cellular

filtration) homotopy equivalent up to a wedge sum with some collection of D^n , $n \in \mathbb{N}$ (with its canonical CW structure), see Faria [3, Theorem 2.4].

When no such n exists we set $\text{q-obs}(P) = \infty$.

Theorem 1.9. *Let $S^2 \hookrightarrow P \rightarrow S^4$ be a non-trivial Hamiltonian fibration then:*

$$\text{q-obs}(P) = 4.$$

Indeed, the proof of Theorem 1.1 can be understood as showing that the associated obstruction class in

$$H^4(S^4, \pi_3(NFuk(S^2)))$$

is non-trivial.

1.2.1. *First quantum obstruction as a manifold invariant.* Let X be a smooth manifold, and let $P(X)$ denote the fiber-wise projectivization of $TX \otimes \mathbb{C}$. We then define

$$\text{q-obs}(X) := \text{q-obs}(P(X)) \in \mathbb{N} \sqcup \{\infty\},$$

which is then an invariant of the smooth manifold X . It should be noted that this “first quantum obstruction” invariant is only sensitive to the tangent bundle, whereas for example Donaldson invariants can see finer aspects of the smooth structure. In fact the “quantum Novikov conjecture” of Part I would immediately imply that the first quantum obstruction is only a topological invariant of X , but in that case it is very likely a new topological invariant (not expressible in terms like Pontryagin numbers).

2. ACKNOWLEDGEMENTS

I am grateful to RIMS institute at Kyoto university and Kaoru Ono for the invitation, financial assistance and a number of discussions which took place there. Much thanks also ICMAT Madrid and Fran Presas for providing financial assistance, and a lovely research environment during my stay there. I have also benefited from conversations with (in no particular order) Hiro Lee Tanaka, Mohammed Abouzaid, Kevin Costello and Bertrand Toen, Nick Sheridan, and Paul Seidel.

CONTENTS

1. Introduction	1
1.1. An application to curvature bounds of singular and smooth connections	2
1.2. First quantum obstruction and smooth invariants	5
2. Acknowledgements	6
3. Preliminaries	7
4. Outline	8
5. Qualitative description of the perturbation data	9
5.1. Extending $\mathcal{D}_{pt}$ to higher dimensional simplices	10
5.2. The main lemma and immediate consequences	13
6. Towards the proof of Lemma 5.5	14

7. Construction of unexcited data	15
8. The product $\mu_{\Sigma_-}^4(\gamma_1, \dots, \gamma_4)$ and the quantum Maslov classes	17
8.1. Constructing suitable $\{\mathcal{A}_r\}$	17
8.2. Restructuring $\{\mathcal{A}_r\}$	20
8.3. Computing $[ev(\mathcal{H}^n, A_0)]$	22
9. Finishing up the proof of Lemma 5.5	23
10. Singular and simplicial connections and curvature bounds	23
10.1. Simplicial connections	24
Appendix A. Homotopy groups of Kan complexes	26
References	28

3. PRELIMINARIES

We quickly recall some notation from Part I and some other basics for convenience. We denote by Δ the simplex category:

- The set of objects of Δ is $\mathbb{N}$.
- $\text{hom}_{\Delta}(n, m)$ is the set of non-decreasing maps $[n] \rightarrow [m]$, where $[n] = \{0, 1, \dots, n\}$, with its natural order.

A simplicial set X is a functor

$$X : \Delta^{op} \rightarrow \text{Set}.$$

The set $X(n)$ is called the set of n -simplices of X . $\Delta_{\bullet}^d$ will denote a particular simplicial set: the standard representable d -simplex, with

$$\Delta_{\bullet}^d(n) = \text{hom}_{\Delta}(n, d).$$

We use notation Δ^n to denote the standard topological n -simplex.

Definition 3.1. *For Y smooth manifold, $\Delta(Y)$ will denote the **smooth simplex category of Y** . This is the category s.t.:*

- The set of objects $\text{obj } \Delta(Y)$ is the set of smooth maps:

$$\Sigma : \Delta^d \rightarrow X, \quad d \geq 0.$$

- Morphisms $f : \Sigma_1 \rightarrow \Sigma_2$ are commutative diagrams in $s\text{-Set}$:

$$(3.1) \quad \begin{array}{ccc} \Delta^d & \xrightarrow{\quad} & \Delta^n, \\ & \searrow \Sigma_1 & \downarrow \Sigma_2 \\ & & X. \end{array}$$

where $\Delta^n \rightarrow \Delta^m$ is a simplicial map, that is an affine map taking vertices to vertices, preserving the order.

When Y is a smooth manifold $Y_\bullet$ will denote the smooth singular set of X . That is $X_\bullet(n)$ is the set of smooth maps $\Delta^n \rightarrow X$.

If $p : X \rightarrow Y$ is a map of spaces, $p_\bullet : X_\bullet \rightarrow Y_\bullet$ will mean the induced simplicial map. We will denote abstract Kan complexes or quasi-categories by calligraphic letters e.g. $\mathcal{X}, \mathcal{Y}$.

4. OUTLINE

In what follows, when we say Part I we shall mean [8]. We will mostly follow the notation and setup of Part I, as well as some setup of [7]. The reader may review the basics of simplicial sets, in Section 3 of Part I. For a more detailed introduction, which also includes some theory of quasi-categories, we recommend Riehl [5]. Here are some specific summary points. Let us briefly review what we do in Part I. Let $M \hookrightarrow P \xrightarrow{p} X$ be a Hamiltonian fibration.

As in Part I, an auxiliary perturbation data $\mathcal{D}$ for P , (in particular) involves:

- A choice of a natural system $\mathcal{U}$, see Section 7, consisting of certain maps

$$u : \mathcal{S}_d^\circ \rightarrow \Delta^n, \text{ with } d, n \text{ varying,}$$

as already discussed in Section 1.1.1 of the Introduction.

- Choices of certain Hamiltonian connections, on Hamiltonian bundles associated to the maps u . (Oversimplified for this outline.)

Given such a $\mathcal{D}$, we construct in Part I a functor

$$F : \Delta(X) \rightarrow A_\infty - Cat,$$

where $A_\infty - Cat$ denotes the category of A_∞ categories. The properties of this functor are such that we may algebraically get an induced functor

$$F^{unit} : \Delta(X) \rightarrow A_\infty - Cat^{unit},$$

with $A_\infty - Cat^{unit}$ denoting the category of unital A_∞ categories, by taking unital replacements. In what follows we rename F by F^{raw} and F^{unit} by F , as F is the main object here.

We then define

$$\text{Fuk}_\infty(P) = \text{colim}_{\Delta(X)} NF,$$

which is shown to be an ∞ -category whose equivalence class (under concordance, see Definition 4.3) is independent of all choices. This also has the structure of a Cartesian fibration:

$$\text{NFuk}(M, \omega) \hookrightarrow \text{Fuk}_\infty(P) \xrightarrow{p_\bullet} X_\bullet,$$

where $\text{NFuk}(M, \omega)$ is the A_∞ nerve of the Fukaya category of M . We will extract from the above fibration a Kan fibration and work with that, since then we can just use standard tools of topology.

To this end we have the following elementary lemma.

Lemma 4.1. *Suppose we have a Cartesian fibration $p : \mathcal{Y} \rightarrow \mathcal{X}$, where $\mathcal{X}$ is a Kan complex. Let $K(\mathcal{Y})$ denote the maximal Kan sub-complex of $\mathcal{Y}$ then $p : K(\mathcal{Y}) \rightarrow \mathcal{X}$ is a Kan fibration.*

The proof is given in Appendix A. In particular by the above lemma

$$K(P) := K(\mathrm{Fuk}_\infty(P)) \xrightarrow{p_\bullet} X_\bullet$$

is a Kan fibration.

Notation 4.2. In what follows $p_\bullet$ will refer to this projection unless specified otherwise.

Definition 4.3. We say that a Kan fibration, or a Cartesian fibration $\mathcal{P}$ over a Kan complex $\mathcal{X}$ is **non-trivial** if it is not null-concordant. Here $\mathcal{P}$ is **null-concordant** means that there is a Kan respectively Cartesian fibration

$$\mathcal{Y} \rightarrow \mathcal{X} \times \Delta_\bullet^1,$$

whose pull-back by $i_0 : \mathcal{X} \rightarrow \mathcal{X} \times \Delta_\bullet^1$ is trivial and by $i_1 : \mathcal{X} \rightarrow \mathcal{X} \times \Delta_\bullet^1$ is $\mathcal{P}$. Here the two maps i_0, i_1 correspond to the two vertex inclusions $\Delta_\bullet^0 \rightarrow \Delta_\bullet^1$.

Theorem 4.4. Suppose that $p : P \rightarrow S^4$ is a non-trivial Hamiltonian S^2 fibration then $p_\bullet : K(P) \rightarrow S_\bullet^4$ does not admit a section. In particular $K(P)$ is a non-trivial Kan fibration over $S_\bullet^4$ and so $\mathrm{Fuk}_\infty(P)$ is a non-trivial Cartesian fibration over $S_\bullet^4$.

This is the main technical result of the paper. Although in a sense we just are just deducing existence of a certain holomorphic curve, for this deduction we need a global compatibility condition involving multiple moduli spaces, involved in multiple local datum's of Fukaya categories, so that this computation will not be straightforward.

The proof will be aided by constructing suitable perturbation data, and will be split into a number of sections.

5. QUALITATIVE DESCRIPTION OF THE PERTURBATION DATA

Let $\mathrm{Fuk}(S^2, \omega)$ denote the $\mathbb{Z}_2$ -graded A_∞ category over $\mathbb{Q}$, with objects oriented spin Lagrangian submanifolds Hamiltonian isotopic to the equator. Our particular construction of $\mathrm{Fuk}(M, \omega)$ is presented in Part I. In particular, we use the language of perturbation systems $\mathcal{D}$, see Section 5, and 6.1 Part I. The data $\mathcal{D}$ is generally associated to a Hamiltonian fibration, and uses the language of connections. As a symplectic manifold is a Hamiltonian fibration over a point, we write $\mathcal{D}_{pt}$ for this restricted data, needed for construction of $\mathrm{Fuk}(S^2, \omega)$.

Denote by $\mathrm{Fuk}^{\mathrm{eq}}(S^2, \omega) \subset \mathrm{Fuk}(S^2, \omega)$ the full sub-category obtained by restricting our objects to be equators in S^2 . We take our perturbation data $\mathcal{D}_{pt}$ so that the following is satisfied.

- All the connections $\mathcal{A}(L, L')$ for $L, L' \in \mathrm{Fuk}^{\mathrm{eq}}(S^2)$ are $PU(2)$ -connections.
- For L intersecting L' transversally, the $PU(2)$ connection $\mathcal{A}(L, L')$ is the trivial flat connection.
- For $L = L'$ the corresponding connection is generated by an autonomous Hamiltonian.

The associated cohomological Donaldson-Fukaya category $DF(S^2)$ is equivalent as a linear category over $\mathbb{Q}$ to $FH(L_0, L_0)$ (considered as a linear category with one object) for $L_0 \in \text{Fuk}(S^2, \omega)$.

It is easily verified that a morphism (1-edge) f is an isomorphism in the nerve $\text{NFuk}(S^2)$, if and only if it corresponds, under the nerve construction N , to a morphism in $\text{Fuk}(S^2, \omega)$ that induces an isomorphism in $DF(S^2)$. Such a morphism will be called a *c-isomorphism*.

Consequently the maximal Kan subcomplex $K(S^2)$ of $\text{NFuk}(S^2)$ is characterized as the maximal subcomplex with 1-simplices the images by N of *c-isomorphisms* in $\text{Fuk}(S^2)$.

5.1. Extending $\mathcal{D}_{pt}$ to higher dimensional simplices.

Terminology 5.1. *A bit of possibly non-standard terminology: we say that A is a model for B in some category, with weak equivalences, if there is a morphism $\text{mod} : A \rightarrow B$ which is a weak-equivalence. The map mod will be called a modelling map. In our context the modeling map mod always turns out to be a monomorphism.*

Let us model $D_\bullet^4$ and $S_\bullet^3$ as follows. Take the standard representable 3-simplex $\Delta_\bullet^3$, and the standard representable 0-simplex $\Delta_\bullet^0$. Then collapse all faces of $\Delta_\bullet^3$ to a point, that is take the colimit of the following diagram:

$$(5.1) \quad \begin{array}{ccccc} & & \Delta_\bullet^0 & & \\ & \nearrow & \uparrow & \nwarrow & \\ \Delta_\bullet^2 & & \Delta_\bullet^2 & & \Delta_\bullet^2 \\ & \searrow_{i_0} & \downarrow_{i_2} & \swarrow_{i_3} & \\ & & \Delta_\bullet^3 & & \end{array}$$

Here i_j are the inclusion maps of the non-degenerate 2-faces. This gives a simplicial set $S_\bullet^{3, \text{mod}}$ modeling the simplicial set $S_\bullet^3$; in other words there is a natural a weak-equivalence

$$S_\bullet^{3, \text{mod}} \rightarrow S_\bullet^3.$$

Now take the cone on $S_\bullet^{3, \text{mod}}$, denoted by $C(S_\bullet^{3, \text{mod}})$, and collapse the one non-degenerate 1-edge. The resulting simplicial set $D_\bullet^{4, \text{mod}}$ is our model for $D_\bullet^4$, it may be identified with a subcomplex of $D_\bullet^4$ so that the inclusion map $\text{mod} : D_\bullet^{4, \text{mod}} \rightarrow D_\bullet^4$ induces a weak homotopy equivalence of pairs

$$(5.2) \quad (D_\bullet^{4, \text{mod}}, S_\bullet^{3, \text{mod}}) \rightarrow (D_\bullet^4, S_\bullet^3).$$

We set $b_0 \in D_\bullet^4$ to be the vertex which is the image by mod of the unique 0-vertex in $D_\bullet^{4, \text{mod}}$.

Suppose we have a commutative diagram:

$$\begin{array}{ccccc} D^4 & \xrightarrow{h_+} & S^4 & \xleftarrow{h_-} & D^4 \\ & \nwarrow i & & \nearrow i & \\ & & S^3 & & \end{array}$$

where $i : S^3 \rightarrow D^4$ is the natural boundary inclusion, and s.t. the following is satisfied.

- $h_{\pm} : D^4 \rightarrow S^4$ are smooth, and their images cover S^4 .

•

$$h_+(D^4) \cap h_-(D^4)$$

is contained in the image E of

$$h_{\pm} \circ i : S^3 \rightarrow S^4.$$

- $h_{\pm}$ takes b_0 to x_0 .

For example, we may just let h_- represent the generator of $\pi_4(S^4, x_0)$ and h_+ to be the constant map to x_0 . We call such a pair $h_{\pm}$ a *complementary pair*.

We set

$$D_{\pm} := h_{\pm}(D_{\bullet}^{4,mod}) \subset S^4_{\bullet},$$

and we set $\Sigma_{\pm} \in S^4_{\bullet}$ to be the image by $h_{\pm}$ of the sole non-degenerate 4-simplex of $D_{\bullet}^{4,mod}$. Also set

$$\partial D_{\pm} := h_{\pm}(\partial D_{\bullet}^{4,mod}),$$

where $\partial D_{\bullet}^{4,mod}$ is the image of the natural inclusion $S_{\bullet}^{3,mod} \rightarrow D_{\bullet}^{4,mod}$.

Fix a Hamiltonian frame for the fiber P_{x_0} of P over x_0 , in other words a Hamiltonian bundle diffeomorphism

$$\begin{array}{ccc} S^2 & \longrightarrow & P \\ \downarrow & & \downarrow p \\ pt & \xrightarrow{x_0} & X. \end{array}$$

In particular, this allows us to identify $\text{Fuk}(S^2, \omega)$ with $F^{raw}(x_0)$, using the analytic perturbation data $\mathcal{D}_{pt}$ for both. Denote by $x_{0,\bullet}$ the image of the map

$$\Delta_{\bullet}^0 \rightarrow S_{\bullet}^{4,mod},$$

induced by the inclusion of the 0-simplex x_0 .

We continue with the description of the data $\mathcal{D} = \mathcal{D}(P)$. This must associate certain data $\mathcal{D}_{\Sigma}$ for each singular simplex $\Sigma : \Delta^n \rightarrow S^4$. Recall from Section 8 Part I, that given the data $\mathcal{D}_{\Sigma}$ for a non-degenerate simplex Σ , we assigned extended perturbation data $\mathcal{D}_{\tilde{\Sigma}}$ for all degeneracies $\tilde{\Sigma}$ of this simplex. So by this discussion, our chosen data $\mathcal{D}_{pt}$ induces perturbation data for all degeneracies of x_0 , that is for all simplices of $x_{0,\bullet}$, this data will again be denoted by $\mathcal{D}_{pt}$, for simplicity.

Fix an object $L_0 \in \text{Fuk}^{\text{eq}}(S^2) \subset F^{raw}(x_0)$. Denote by $\gamma \in \text{hom}_{F^{raw}(x_0)}(L_0, L_0)$ the generator of $FH_1(L_0, L_0)$, i.e. the fundamental chain, so that it corresponds to the identity in $DF(L_0, L_0)$. This γ is uniquely determined by our conditions and

corresponds to a single geometric section. Denote by L_0^i the image of L_0 by the embedding

$$F^{raw}(x_0) \rightarrow F^{raw}(\Sigma_+),$$

corresponding to the i 'th vertex inclusion into Δ^4 , $i = 0, \dots, 4$.

Let m_i be the edge between $i-1, i$ vertices and set

$$\overline{m}_i := \Sigma_+ \circ m_i.$$

Let Σ_i^0 denote the 0-simplex obtained by restriction of Σ^4 to the i 'th vertex. Note that each $\overline{m}_i$ is degenerate by construction, so we have an induced morphism

$$F^{raw}(pr) : F^{raw}(\overline{m}_i) \rightarrow F^{raw}(x_0),$$

for pr the degeneracy morphism in $\Delta(S^4)$:

$$pr : \overline{m}_i \rightarrow \Sigma_i^0.$$

Finally, for each L_0^{i-1}, L_0^i we have a c -isomorphism

$$\gamma_i : L_0^{i-1} \rightarrow L_0^i$$

in $F^{raw}(\overline{m}_i) \subset F^{raw}(\Sigma_+)$, which corresponds to γ , meaning that the fully-faithful projection $F^{raw}(pr)$ takes γ_i to γ . We will denote by $\gamma_{i,j}$ the analogous c -isomorphisms $L_0^i \rightarrow L_0^j$.

Notation 5.2. Let us denote from now on, the morphism spaces $\text{hom}_{F^{raw}(\Sigma_\pm)}(L_0, L_1)$ by $\text{hom}_{\Sigma_\pm}(L_0, L_1)$. And denote the A_∞ composition maps $\mu_{F^{raw}(\Sigma_\pm)}^d$, in the A_∞ category $F^{raw}(\Sigma_\pm)$, by $\mu_{\Sigma_\pm}^d$.

Definition 5.3. We call perturbation data $\mathcal{D}$ for P **unexcited** if it extends the data $\mathcal{D}_{pt}$ as above, and if with respect to $\mathcal{D}$

$$(5.3) \quad \mu_{\Sigma_+}^d(\gamma^1, \dots, \gamma^d) = 0, \text{ for } 2 < d < 4,$$

where $(\gamma^1, \dots, \gamma^d)$ is a composable chain, and each γ^k is of the form $\gamma_{i,j}$ as above.

We will see further on how to construct such unexcited data, assume for now that it is constructed.

Let $\{f_J\}$, corresponding to an n -simplex, be as in the definition of the A_∞ nerve in Appendix A.4 Part I, where J is a subset of $[n] = \{0, \dots, n\}$.

Lemma 5.4. Let $\mathcal{D}$ be unexcited as above, then there is a 4-simplex $\sigma \in NF^{raw}(\Sigma_+)$ with faces determined by the conditions:

- $f_J = 0$, for J any subset of $[4]$ with at least 3 elements.
- $f_{\{i-1, i\}} = \gamma_i$ for γ_i as before.

Proof. This follows by (5.3) and by the identity $\mu_{\Sigma_+}^2(\gamma, \gamma) = \gamma$. □

If we take our unital replacements so that γ corresponds to the unit, then σ induces (by the construction) a section of $K(P_+) \rightarrow D_+$, where $K(P_\pm)$ will be shorthand for $K(P)$ restricted over $D_\pm$.

Let

$$i : (K(P_+)|_{\partial D_+} := p_\bullet^{-1}(\partial D_+)) \rightarrow K(P_-),$$

be the natural inclusion map. Set

$$sec = i \circ \sigma \circ h_+|_{\partial D_\bullet^{4,mod}}.$$

5.2. The main lemma and immediate consequences.

Lemma 5.5. *Suppose that P is a non-trivial Hamiltonian fibration and $\mathcal{D}$ is unexcited data for P as above, then sec as above does not extend to a section of $K(P_-)$. Moreover, unexcited data $\mathcal{D}$ exists.*

This lemma involves all the ingredients of our theory, its proof that will be broken up in parts, and will follow shortly.

Proof of Theorem 1.9. Clearly $\text{q-obs}(P) \geq 4$, since the 3-skeleton of S^4 is trivial. By Lemma 5.5 above, $K(P)$ does not have a section over the 4-skeleton. $\square$

Remark 5.6. *When P is obtained by clutching with a generator of $\pi_3(PU(2))$, and when $h_\pm$ are embeddings, the class $[sec]$ in $\pi_3(K(P_-)) \simeq \pi_3(K(S^2))$ can be thought of as “quantum” analogue of the class of the classical Hopf map.*

Proof of Theorem 4.4. It might be helpful to first review Appendix A before reading the following. If we take any unexcited perturbation data $\mathcal{D}$ for P , then the first part follows immediately by Lemma 5.5. So $K(P)$ is non-trivial as a Kan fibration. This then implies that $\text{Fuk}_\infty(P)$ is non-trivial as a Cartesian fibration, which means in particular that its classifying map

$$f_P : S_\bullet^4 \rightarrow (\mathbb{S}, \text{NFuk}(S^2, \omega))$$

is not null-homotopic.

To see this, suppose otherwise that we have a Cartesian fibration

$$\tilde{\mathcal{P}} \rightarrow S_\bullet^4 \times I_\bullet,$$

restricting to $\text{Fuk}_\infty(P)$ over $S_\bullet^4 \times 0_\bullet$ and to $\text{NFuk}(S^2, \omega) \times S_\bullet^4$ over the other end $S_\bullet^4 \times 1_\bullet$. Here $0_\bullet$, respectively $1_\bullet$ are notation for the images of $i_{j,\bullet} : \Delta_\bullet^0 \rightarrow \Delta_\bullet^1$, $j = 0, 1$, where $i_{j,\bullet}$ are induced by the pair of boundary point inclusions.

Now take the maximal Kan sub-fibration of $\tilde{\mathcal{P}}$, then by Lemma 4.1 we obtain a trivialization of $K(P)$ which is a contradiction. $\square$

Proof of Theorem 1.1. Theorem 4.4 implies that the group homomorphism

$$\mathbb{Z} \simeq \pi_4(\text{BHam}(S^2, \omega) \rightarrow \pi_4(\mathbb{S}, \text{NFuk}(S^2, \omega))),$$

has vanishing kernel, so that the result follows. $\square$

6. TOWARDS THE PROOF OF LEMMA 5.5

We will denote by $L_{0,\bullet}$ the image of the map $\Delta_{\bullet}^0 \rightarrow K(P_-)$, induced by the inclusion of L_0 into $K(S^2)$ as a 0-simplex. Suppose that sec extends to a section of $K(P_-)$, so we have map

$$e : D_{\bullet}^{4,mod} \rightarrow K(P_-)$$

extending sec over $\partial D_{\bullet}^{4,mod}$. We may assume WLOG that e lies over h_- , meaning

$$p_{\bullet} \circ e = h_-.$$

since it can be homotoped to have this property. To see this, first take a relative homotopy of

$$p_{\bullet} \circ e : (D_{\bullet}^{4,mod}, \partial D_{\bullet}^{4,mod}) \rightarrow (D_-, \partial D_-)$$

to h_- , using that we have a homotopy equivalence of pairs (5.2), and then lift the homotopy to a relative homotopy upstairs using the defining lifting property of Kan fibrations.

And so we have a 4-simplex

$$T = e(\Sigma^4) \in K(P_-)$$

projecting to $\Sigma_- \in D_-$ by $p_{\bullet}$. Since T is in the image of e , all but one 3-faces of T are totally degenerate with image in $L_{0,\bullet}$. The exceptional 3-face is the sole non-degenerate 3-face of sec , (that is of $sec(\partial D_{\bullet}^{4,mod})$).

Let $m_{i,j}, \gamma_{i,j}$ be as in the previous section, but corresponding now to Σ_- rather than Σ_+ . Then by the boundary condition on e , the edges of T (which are all edges of sec) correspond, under the nerve construction, to the generators $\gamma_{i,j}$. As this is the condition for the edges of sec .

Lemma 6.1. *For $\mathcal{D}$ unexcited, and for the unital replacement F of F^{raw} as above, the simplex T exists if and only if $\mu_{\Sigma_-}^4(\gamma_1, \dots, \gamma_4)$ is exact.*

Proof. The following argument will be over $\mathbb{F}_2$ as opposed to $\mathbb{Q}$ as the signs will not matter. Recall that we take the unital replacement so that $\gamma \in hom_{F^{raw}(P_{x_0})}(L_0, L_0)$ corresponds to the unit in the unital replacement.

Now if $T \in K(P_-)$ as above exists, then it corresponds under unital replacement (see Remark 7.5 in Part I) to a 4-simplex $T' \in NF^{raw}(\Sigma_-)$ satisfying the following condition on its 4-face. Recalling the nerve construction, the morphism $f_{[4]} \in hom_{\Sigma_-}(L_0^0, L_0^4)$, figuring in the definition of the 4-face, satisfies:

$$(6.1) \quad \mu_{\Sigma_-}^1 f_{[4]} = \sum_{1 < i < 4} f_{[4]-i} + \sum_s \sum_{(J_1, \dots, J_s) \in decomp_s} \mu_{\Sigma_-}^s(f_{J_1}, \dots, f_{J_s}).$$

By our conditions on the boundary of T , by the condition on the unital replacement, and by the conditions in Lemma 5.4, we must have $f_J = 0$, for every proper subset $J \subset [4]$, in some length s decomposition of $[4]$, unless $J = \{i, j\}$ in which case $f_{i,j} = \gamma_{i,j}$. Given this (6.1) holds if and only if $\mu_{\Sigma_-}^4(\gamma_1, \dots, \gamma_4)$ is exact.

□

We are going to show that for some unexcited $\mathcal{D}$, $\mu_{\Sigma_-}^4(\gamma_1, \dots, \gamma_4)$ does not vanish in homology, which will finish the proof of the Lemma 5.5.

7. CONSTRUCTION OF UNEXCITED DATA

To forewarn, we use here notation and notions from Part I, as well some basic technical constructions and lemma's from [7].

Let $(m^1, \dots, m^d)$ be a composable chain of morphisms in $\Pi(\Delta^n)$, which we recall means that the target of m^{i-1} is the source of m^i for each i . The perturbation data $\mathcal{D}$, in particular specifies for each n and for each such composable chain, certain maps

$$u(m^1, \dots, m^d, n) : \mathcal{E}_d^\circ \rightarrow \Delta^n.$$

Here $\mathcal{E}_d$ is the universal curve over $\overline{\mathcal{R}}_d$, and $\mathcal{E}_d^\circ$ denotes $\mathcal{E}_d$ with nodal points of the fibers removed. The collection of these maps, satisfying certain axioms, is denoted by $\mathcal{U}$. We have already mentioned this in the introduction.

The restriction of $u(m^1, \dots, m^d, n)$ to the fiber $\mathcal{S}_r$ of $\mathcal{E}_r^\circ$ over r , is denoted by $u(m^1, \dots, m^d, n, r)$ which may also be abbreviated by u_r .

Let $\Sigma : \Delta^n \rightarrow X$ be smooth, denote $\overline{m}^i := \Sigma \circ m^i$, and denote $\tilde{\mathcal{S}}_r := (\Sigma \circ u_r)^* P$. Then $\mathcal{D}$ specifies for each: Σ , r , composable chain $(m^1, \dots, m^d)$ in $\Pi(\Delta^n)$, and for each chain of objects $L'_0, \dots, L'_d$ with $L'_i \subset P|_{\overline{m}_i(1)}$, $i \geq 1$, $L'_0 \subset P|_{\overline{m}_1(0)}$, a certain Hamiltonian connection $\mathcal{A}_r$ on

$$(7.1) \quad \tilde{\mathcal{S}}_r \rightarrow \mathcal{S}_r.$$

Using terminology of [7, Section 2.4], $\mathcal{D}$ induces a family Hamiltonian structure

$$\{\tilde{\mathcal{S}}_r, \mathcal{S}_r, \mathcal{L}_r, \mathcal{A}_r\}_{\overline{\mathcal{R}}_d},$$

where $\mathcal{L}_r$ is trivial over each boundary component, with fiber the corresponding objects L'_i .

Now, specialize to the case $n = 4$, $\Sigma = \Sigma_+$, $\forall i : L'_i = L_0$. In this case, we write $\mathcal{A}_r^+(m^1, \dots, m^d)$ for the corresponding connections, further abbreviated by $\mathcal{A}_r^+$ as $(m^1, \dots, m^d)$ will usually be implicit.

Suppose that $\mathcal{D}$ extends $\mathcal{D}_{pt}$ from before. If $\mathcal{L}_r \subset \tilde{\mathcal{S}}_r|_{\partial \mathcal{S}_r}$ denotes the trivial Lagrangian sub-bundle with fiber L_0 , then we obtain a Hamiltonian structure

$$\Theta^+ = \{\Theta_r^+\} = \{\tilde{\mathcal{S}}_r, \mathcal{S}_r, \mathcal{L}_r, \mathcal{A}_r^+\}_{\overline{\mathcal{R}}_d}.$$

By the properties necessitated by $\mathcal{D}$, at each end e_i of $\mathcal{S}_r$, $\mathcal{A}_r^+$ is compatible with the connection $\mathcal{A}_i = \mathcal{A}(L_0, L_0)$, where $\mathcal{A}(L_0, L_0)$ is the connection on $[0, 1] \times S^2$, also part of our data $\mathcal{D}$. Then Θ^+ is trivially taut since for each r $\mathcal{L}_r$ is naturally trivial and $\mathcal{A}_r$ is likewise trivial over $\partial \mathcal{S}_r$, for each r , again by the properties necessitated by $\mathcal{D}$.

Set

$$\hbar := \frac{1}{2} \text{area}(S^2, \omega).$$

Let κ denote the $L^\pm$ -length of the holonomy path in $\text{Ham}(S^2, \omega)$ of $\mathcal{A}_0 = \mathcal{A}(L_0, L_0)$. We may suppose that

$$(7.2) \quad \forall r : \text{area}(\mathcal{A}_r^+) < \hbar - 5\kappa,$$

is satisfied after taking κ to be sufficiently small. (There is no obstruction since the corresponding bundles are naturally trivializable, continuously in r .)

Fix a complex structure j_0 on M , and let $\{J_r = J(\mathcal{A}_r^+)\}$ be the corresponding, induced family of almost complex structures on $\{\tilde{\mathcal{S}}_r\}$ as defined in [7, Section 2.5].

Lemma 7.1. *As in Part I, let*

$$\overline{\mathcal{M}} = \overline{\mathcal{M}}(\gamma^1, \dots, \gamma^d; \gamma^0, \Sigma_+, \{J_r\}, A),$$

denote the set of elements of $\overline{\mathcal{M}}(\Theta^+, A)$ with asymptotic constraints γ^i at each e_i end. Here each γ^k , $k \neq 0$, is of the form $\gamma_{i,j}$ where this is as in Section 5.1. Then whenever the class A is such that $\mathcal{M}$ has virtual dimension 0, and d satisfies $2 < d \leq 4$, $\overline{\mathcal{M}}$ is empty.

Proof. Let

$$\Theta' := (\Theta^+)',$$

and A' be the capping off construction as in [7, Section 2.7]. For a fixed r , by the Riemann-Roch, [Appendix A][7], we get that the expected dimension of $\mathcal{M}(\Theta', A')$ is

$$1 + \text{Maslov}^{vert}(A').$$

Consequently, when $\gamma^0 = \gamma$, the expected dimension of $\mathcal{M}$ is:

$$(7.3) \quad 1 + \text{Maslov}^{vert}(A') - 1 + (\dim \mathcal{R}_d = d - 2).$$

We need the expected dimension of $\mathcal{M}$ to be 0, and $d \geq 3$, so $\text{Maslov}^{vert}(A') \leq -1$. But $\text{Maslov}^{vert}(A') = -1$ is impossible as the minimal positive Maslov number is 2.

Now, note that if $\text{Maslov}^{vert}(A') = -2$ then

$$-C \cdot \text{Maslov}^{vert}(A') = \hbar,$$

for C the monotonicity constant of (S^2, ω) and the equator L_0 . Consequently, the result follows by [7, Lemma 2.34] and by the property (7.2).

When γ^0 is the Poincare dual to γ , we would get $\text{Maslov}^{vert}(A') \leq -2$ so for the same reason the conclusion follows. $\square$

So if we choose our data $\mathcal{D}$ so that the hypothesis of the lemma above are satisfied, then with respect to this $\mathcal{D}$:

$$(7.4) \quad \mu_{\Sigma_{\pm}^4}^2(\gamma_{i,j}, \gamma_{j,k}) = \gamma_{i,k}$$

$$(7.5) \quad \mu_{\Sigma_{\pm}^4}^3(\gamma^1, \dots, \gamma^3) = 0, \text{ for } \gamma^i \text{ as above}$$

$$(7.6) \quad \mu_{\Sigma_{+}^4}^4(\gamma_1, \dots, \gamma_4) = 0.$$

In particular this $\mathcal{D}$ is unexcited.

8. THE PRODUCT $\mu_{\Sigma_-}^4(\gamma_1, \dots, \gamma_4)$ AND THE QUANTUM MASLOV CLASSES

The product

$$(8.1) \quad \mu_{\Sigma_-}^4(\gamma_1, \dots, \gamma_4),$$

a priori depends on various choices, like the choices of $h_{\pm}$, and then choice of data $\mathcal{D}_0$. For the purpose of computation we will take h_+ to be the constant map to x_0 and

$$h_- : (D^4, \partial D^4) \rightarrow (S^4, x_0)$$

to be the complementary map, that is representing the generator of $\pi_4(S^4, x_0) \simeq \mathbb{Z}$. We further suppose that h_- is an embedding in the interior of D^4 .

We are then going to reduce the computation of the product (8.1) to the computation of a certain quantum Maslov class, already done in [7]. This will require a number of geometric steps.

Let Σ_- be the 4-simplex of $S^4_{\bullet}$ corresponding to h_- as before in Section 5.1. We need to study the moduli spaces

$$(8.2) \quad \overline{\mathcal{M}}(\gamma_1, \dots, \gamma_4; \gamma^0, \Sigma_-, \{\mathcal{A}_r\}, A),$$

where $\mathcal{A}_r$ now denotes the connections on

$$(8.3) \quad \tilde{\mathcal{S}}_r := (\Sigma_- \circ u(m_1, \dots, m_4, 4, r))^* P \rightarrow \mathcal{S}_r,$$

part of some unexcited data $\mathcal{D}_0$ as above. We abbreviate $u(m_1, \dots, m_4, 4, r)$ by u_r in what follows.

By the dimension formula (7.3), since we need the expected dimension of (8.2) to be zero, the class $A^{\vee}$ satisfies:

$$Maslov^{vert}(A^{\vee}) = -2,$$

and we must have $\gamma^0 = \gamma_{0,4}$.

Notation 8.1. From now on, by slight abuse, A_0 refers to various section classes of various Hamiltonian structures such that the associated class $A_0^{\vee}$ satisfies:

$$Maslov^{vert}(A_0^{\vee}) = -2.$$

8.1. Constructing suitable $\{\mathcal{A}_r\}$. To get a handle on (8.2) we want to construct very special, unexcited data $\mathcal{D}_0$.

A Hamiltonian S^2 fibration over S^4 is classified by an element

$$[g] \in \pi_3(\text{Ham}(S^2, \omega), id) \simeq \pi_3(PU(2), id) \simeq \mathbb{Z}.$$

Such an element determines a fibration P_g over S^4 via the clutching construction:

$$P_g = D_-^4 \times S^2 \sqcup D_+^4 \times S^2 \sim,$$

with D_-^4, D_+^4 being 2 different names for the standard closed 4-ball D^4 , and where the equivalence relation $\sim$ is $(d, x) \sim \tilde{g}(d, x)$,

$$\tilde{g} : \partial D_-^4 \times S^2 \rightarrow \partial D_+^4 \times S^2, \quad \tilde{g}(d, x) = (d, g(d)^{-1}(x)).$$

We suppose that the that previously appeared point $x_0 = h_{\pm}(b_0)$, is in $D_+^4 \cap D_-^4 \subset S^4$.

From now on P_g will denote such a fibration for a non-trivial class $[g]$. Note that the fiber of P_g over the base point $x_0 \in S^3 \subset D_\pm^4$ (chosen for definition of the homotopy group $\pi_3(\text{Ham}(S^2, \omega), id)$) has a distinguished, by the construction, identification with S^2 . Take $\mathcal{A}$ to be a connection on $P \simeq P_g$ which is trivial in the distinguished trivialization over D_+^4 . This gives connections

$$\mathcal{A}'_r := (\tilde{u}_r)^* \mathcal{A}$$

on $\tilde{\mathcal{S}}_r$,

$$\tilde{u}_r = \Sigma_- \circ u_r.$$

By the last axiom for the system $\mathcal{U}$ introduced in Part I, we may choose $\{u_r\}$ so that the family $\{\tilde{u}_r(\mathcal{S}_r)\}$ induces a singular foliation of S^4 with the properties:

- The foliation is smooth outside x_0 . Note that x_0 is the image by $\tilde{u}_r$ of the ends (images of e_i), and the image of the boundary of each $\mathcal{S}_r$.
- Each $\tilde{u}_r$ is an embedding on the complement of $\tilde{u}_r^{-1}(x_0)$.

Denote by E the subset $S^3 \subset S^4$ bounding $D_\pm^4$. We may in addition suppose that each $\tilde{u}_r$ intersects E transversally, again on the complement of $\tilde{u}_r^{-1}(x_0)$.

By the above, the preimage by $\tilde{u}_r$ of E contains a smoothly embedded curve c_r as in Figure 1, and $\tilde{u}_r$ takes c_r into E . This c_r not uniquely determined, but we may fix a family $r \mapsto c_r$, with parametrizations

$$c_r : \mathbb{R} \rightarrow \mathcal{S}_r,$$

with the properties:

- c_r maps $(-\infty, 0)$ diffeomorphically onto $e_0(\{0\} \times (-\infty, 0))$.
- c_r maps $(1, \infty)$ diffeomorphically onto $e_0(\{1\} \times (-\infty, 0))$.
- $\{c_r\}$ is a C^0 continuous family in r .

We set:

$$\tilde{c}_r := \tilde{u}_r \circ c_r.$$

In Figure 1, the regions $R_\pm$ are the preimages by $\tilde{u}_r$ of $D_\pm^4 \subset S^4$, and c_r bounds R_- . It follows that $\{\tilde{c}_r\}$ likewise induces a singular foliation of the equator $E \simeq S^3$ that is smooth outside x_0 .

So each $\mathcal{A}'_r$ is flat in the region R_+ , in fact is trivial in the distinguished trivialization of $\tilde{\mathcal{S}}_r$ over R_+ , corresponding to the distinguished trivialization of P over D_+^4 . Likewise, we have a distinguished trivialization of $\tilde{\mathcal{S}}_r$ over R_- , corresponding to the distinguished trivialization of P over D_-^4 . In this latter trivialization let

$$\phi_r : \mathbb{R} \rightarrow \text{Ham}(S^2, \omega)$$

be the holonomy path of $\mathcal{A}'_r$ over c_r . Then by construction,

$$\phi_r|_{(-\infty, 0] \sqcup [1, \infty)} = id,$$

so that we may define

$$f(r) \in \Omega_{L_0} \text{Lag}(M),$$

by

$$f(r)(t) = \phi_r(t)(L_0), \quad t \in [0, 1]$$




FIGURE 1. The labels L_0 indicate that the Lagrangian subbundle is constant with corresponding fiber L_0 . The curve c_r bounds R_- .

where the right hand side means apply an element of $\text{Ham}(S^2, \omega)$ to L_0 to get a new Lagrangian. We will say that $f(r)$ is **generated by** $\mathcal{A}'_r$.

Note that by construction

$$(8.4) \quad \phi_r(t) = g(\tilde{c}_r(t)), \quad t \in [0, 1],$$

if we identify $\tilde{c}_r(t)$ with an element of S^3 .

Let $D_0^2 \subset \overline{\mathcal{R}}_4$ be an embedded closed disk, not intersecting the boundary $\partial \overline{\mathcal{R}}_4$, so that ∂D_0^2 is in the gluing normal neighborhood N of $\partial \overline{\mathcal{R}}_4$, as defined in Part I.

So we get a continuous map

$$f : D_0^2 \rightarrow \Omega_{L_0} Eq(S^2).$$

And $f(\partial D_0^2) = p_{L_0}$, with the right-hand side denoting the constant loop at L_0 . Then by construction, and (8.4) in particular, $f \simeq lag$, where $\simeq$ is a homotopy equivalence, and where

$$(8.5) \quad lag : S^2 \rightarrow \Omega_{L_0} Lag(S^2)$$

is the composition

$$S^2 \xrightarrow{g'} \Omega_{id} PU(2) \rightarrow \Omega_{L_0} Eq(S^2),$$

for g' naturally induced by g , and for the second map naturally induced by the map

$$PU(2) \rightarrow Eq(S^2), \quad \phi \mapsto \phi(L_0).$$

We then deform each $\mathcal{A}'_r$ to a connection $\mathcal{A}_r$, which is as follows. In the region R_+ $\mathcal{A}_r$ is still flat, but at each end e_i , $\mathcal{A}_r$ is compatible with $\mathcal{A}(L_0, L_0)$, where this is as in [7, Section 2.2], and so that $\mathcal{A}_r$ is still trivial over the boundary of $\mathcal{S}_r$.

Since $\tilde{\mathcal{S}}_r$ and $\mathcal{A}'_r$ are trivial for $r \in \overline{\mathcal{R}}_d - D_0^2$, with trivialization induced by the trivialization of P_+ , and since the condition (7.2) holds, we may ensure that

$$(8.6) \quad \text{area}(\mathcal{A}_r) < \hbar - 5\kappa,$$

for r in the complement of D_0^2 . In other words $\{\mathcal{A}_r\}$ extends to a system of connections corresponding to unexcited data $\mathcal{D}_0$ for P , as intended.

8.2. Restructuring $\{\mathcal{A}_r\}$. We quickly recall the following definition from [7].

Definition 8.2. *We say that a Hamiltonian structure $\{\Theta_r\}$ is **A-regular** if:*

- *The pairs $(\mathcal{A}_i, \{j_z\})$ are regular so that the Floer chain complexes $CF(\mathcal{A}_i)$ are defined.*
- *$\mathcal{M}(\{\Theta_r\}, A)$ is regular, (transversely cut out).*

*And we say that $\{\Theta_r\}$ is **regular** if it is A-regular for all A . We say that $\{\Theta_r\}$ is **A-admissible** if there are no elements*

$$(\sigma, r) \in \overline{\mathcal{M}}(\{\Theta_r\}, A),$$

for r in a neighborhood of the boundary of $\mathcal{K}$.

Applying [7, Lemma 2.34], and using (8.6) we get that the resulting Hamiltonian structure

$$\mathcal{H} := \{\mathcal{H}_r := (\tilde{\mathcal{S}}_r, \mathcal{S}_r, \mathcal{L}_r, \mathcal{A}_r)\}$$

is A_0 -admissible. We now further mold this data for the purposes of computation.

First cap off the ends e_i , $i \neq 0$, of each $\mathcal{H}_r$ as in the paragraph preceding [7, Lemma 2.34]. This gives a Hamiltonian structure

$$\mathcal{H}^\wedge := \{\tilde{\mathcal{S}}_r^\wedge, \mathcal{S}_r^\wedge, \mathcal{L}_r^\wedge, \mathcal{A}_r^\wedge\}_{\mathcal{K}=D_0^2},$$

satisfying

$$(8.7) \quad \text{area}(\mathcal{A}_r^\wedge) + \kappa < \hbar,$$

for each r . Again by [7, Lemma 2.34] $\mathcal{H}^\wedge$ is A_0 -admissible.

By the classical gluing of holomorphic curves it follows that

$$(8.8) \quad [\mu_{\Sigma^4}^4(\gamma_1, \dots, \gamma_4)] = [ev(\mathcal{H}^\wedge, A_0)].$$

It remains to compute the right-hand side, to this end we further restructure.

Let $p_1 : [0, 1] \rightarrow \text{Lag}(S^2, L_0)$ be the path generated by $\mathcal{A}(L_0, L_0)$, with p_1 starting at L_0 , and where generated has the same meaning as in the previous section. Suppose we have defined p_{i-1} , set $L_{i-1} := p_{i-1}(1)$ and define p_i to be the path in $\text{Lag}(S^2, L_0)$ starting at L_{i-1} , generated by $\mathcal{A}(L_0, L_0)$.

Now set $p_0 := p_1 \cdot \dots \cdot p_d$, where $\cdot$ is path concatenation in diagrammatic order. We may assume that L_0 is transverse to $L_4 = p_0(1)$ by adjusting the connection $\mathcal{A}(L_0, L_0)$ if necessary.

We construct a concordance of $\mathcal{H}^\wedge$ to another Hamiltonian structure:

$$\mathcal{H}^n := \{\tilde{\mathcal{S}}_r^\wedge, \mathcal{S}_r^\wedge, \mathcal{L}_r^n, \mathcal{A}_r^n\},$$




FIGURE 2. Over the boundary components with black labels L_i the Lagrangian subbundle $\mathcal{L}_r^n$ is constant with corresponding fiber L_i . Over the i 'th red boundary component the Lagrangian subbundle corresponds to the path of Lagrangians p_i . Likewise over the right boundary component of the blue region, the Lagrangian subbundle corresponds to the path of Lagrangians p_0^{-1} . In the red striped regions we have removed the curvature of the connection, in the blue striped region we have added it.

whose properties are illustrated in Figure 2.

The Hamiltonian connection $\mathcal{A}_r^n$ satisfies the following conditions, (referring to the Figure 2):

- $\mathcal{A}_r^n$ is flat in the entire region R_+ (which includes the red shaded finger regions).
- The blue region is contained in the strip end chart at the e_0 end, which is down in the figure.
- Along top boundary segment of the blue region, contained in the dashed line, $\mathcal{A}_r^n$ is the trivial connection in the distinguished trivialization at the e_0 end.
- At the e_0 end the connection is unchanged over $[0, 1] \times (s, \infty)$, for s large.

The connections $\mathcal{A}_r^n$ are forced by the conditions to gain curvature in some region, and we limit this region to the blue region of Figure 2.

We choose a concordance $\mathcal{T}$ from $\mathcal{H}^\wedge$ to $\mathcal{H}^n$ such that for the associated family of connections $\{\mathcal{A}_{r,t}\}$, $t \in [0, 1]$, the following is satisfied:

(1)

$$\mathcal{A}_{r,0} = \mathcal{A}_r^\wedge, \quad \mathcal{A}_{r,1} = \mathcal{A}_r^n.$$

(2)

$$\forall r \forall v \in T_z \mathcal{S}_r : \frac{d}{dt} |R_{\mathcal{A}_{r,t}}(v, jv)|_+ < 0,$$

for each $z \in \mathcal{S}_r$, except for z in the region which is blue shaded in Figure 2.

(3)

$$\forall t : |\text{area}(\mathcal{A}_{r,t}) - \text{area}(\mathcal{A}_r^\wedge)| = 0.$$

The last condition can be satisfied, since the gain of area in the blue region is exactly equal to the loss of area in the red regions.

It follows by the conditions above, and the usual use of [7, Lemma 2.34] that $\mathcal{T}$ is A_0 -admissible.

By [7, Lemma 2.13] we have:

$$[ev(\mathcal{H}^\wedge, A_0)] = [ev(\mathcal{H}^\mathfrak{n}, A_0)]$$

since the corresponding Hamiltonian structures are A_0 -admissibly concordant. This finishes our restructuring.

8.3. Computing $[ev(\mathcal{H}^\mathfrak{n}, A_0)]$. If we stretch the neck along the dashed line in Figure 2, the upper half of the resulting building gives us a new Hamiltonian structure:

$$\mathcal{H}^0 = \{\tilde{S}_r^0, S_r^0, \mathcal{L}_r^0, \mathcal{A}_r^0\}.$$

By the classical theory of continuation maps in Floer homology we clearly have that

$$[ev_\mathfrak{n}] := [ev(\mathcal{H}^\mathfrak{n}, A_0)] \in FH(L_0, L_0)$$

is non-zero iff

$$[ev_0] := [ev(\mathcal{H}^0, A_0)] \in FH(L_0, L_4)$$

is non-zero. Let $\mathcal{P}_{L_0, L_4} \text{Lag}(M)$ denote the space of smooth paths in $\text{Lag}(M)$ from L_0 to L_4 . Let

$$f' : D_0^2 \rightarrow \mathcal{P}_{L_0, L_4} \text{Lag}(S^2),$$

be like f but defined with respect to $\{\mathcal{A}_r^0\}$, so that

$$(8.9) \quad f'(t) = g(\tilde{c}_r(t))(p_0(t)),$$

if we suppose that the holonomy path of $\mathcal{A}_r^0$ over c_r , in the trivialization over R_+ , generates p_0 (which can be insured by adjusting $\{\mathcal{A}_r^0\}$ or the parametrizations $\{c_r\}$). Here the right-hand side of (8.9) means as before: apply an element of $\text{Ham}(S^2, \omega)$ to a Lagrangian to get a new Lagrangian. In this case

$$f'(\partial D_0^2) = p_0 \in \mathcal{P}_{L_0, L_4} \text{Lag}(M).$$

In particular, f' represents a class

$$(8.10) \quad a \in \pi_2(\mathcal{P}_{L_0, L_4} \text{Lag}(M), p_0)$$

We omit specifying the parameter space D_0^2 for r , since it will be the same everywhere. Let $\mathcal{L}_p$ be as in [7, Definition 2.27]

Lemma 8.3. *The A_0 -admissible Hamiltonian structure $\mathcal{H}_0 = \{\tilde{\mathcal{S}}_r^0, \mathcal{S}_r^0, \mathcal{L}_r^0, \mathcal{A}_r^0\}$ is A_0 -admissibly concordant to*

$$\Theta' = \{\mathcal{D} \times S^2, \mathcal{D}, \mathcal{L}_{f'}(r), \mathcal{B}_r\},$$

for certain Hamiltonian connections $\{\mathcal{B}_r\}$ (which are not explicitly relevant yet).

Proof. Let $R_\pm \subset \mathcal{S}_r^\wedge$ be as before. Fix a smooth deformation retraction

$$ret_r : \mathcal{S}_r^0 \times I \rightarrow \mathcal{S}_r^0,$$

of $\mathcal{S}_r^0$ onto R^- , smooth in r . Since $\mathcal{A}_r^0$ is flat over R^+ , the pull-back by ret_r of the data $\mathcal{H}_0$ then induces an A_0 -admissible concordance between $\mathcal{H}_0$ and

$$\{\mathcal{D} \times S^2, \mathcal{D}, \mathcal{L}_{f'}(r), \mathcal{B}_r = ret_r^* \mathcal{A}_r^0\},$$

once we use smooth Riemann mapping theorem to identify each $R^- \subset \mathcal{S}_r^0$ with its induced complex structure j_r with $(\mathcal{D}, j_{st})$, smoothly in r . $\square$

9. FINISHING UP THE PROOF OF LEMMA 5.5

The existence of unexcited data $\mathcal{D}$ is proved in Section 7. Given this existence, starting with (8.8) we showed that $[\mu_{\Sigma^4}^4(\gamma_1, \dots, \gamma_4)]$ is non-vanishing in Floer homology iff

$$[ev(\mathcal{H}_0, A_0)] \in HF(L_0, L_4),$$

is non-vanishing. We then use Lemma 8.3 to identify $[ev(\mathcal{H}_0, A_0)]$ with $[ev(\Theta', A_0)]$.

Now by definitions

$$[ev(\Theta', A_0)] = \Psi(a),$$

where the right hand side is the quantum Maslov class as defined in [7], and where a is a spherical class as in (8.10). Given that g is non-trivial the corresponding class a is non-trivial and so $\Psi(a)$ is non-trivial by Theorem [7, Theorem 4.13]. This together with Lemma 6.1 imply Lemma 5.5. $\square$

10. SINGULAR AND SIMPLICIAL CONNECTIONS AND CURVATURE BOUNDS

Let $\mathcal{A}$ be a G connection on a principal G bundle $P \rightarrow \Delta^n$, and the Finsler norm $\mathbf{n}$ on $\text{lie } G$ be as in Section 1.1.1 of the introduction. As previously discussed, a given system $\mathcal{U}$ in particular specifies maps:

$$u(m_1, \dots, m_n, r, n) : \mathcal{S}_r \rightarrow \Delta^n,$$

where $r \in \overline{\mathcal{R}}_n$, $\mathcal{S}_r$ is the fiber of $\mathcal{S}_n^\circ$ over r , and where $(m_1, \dots, m_n)$ is the composable chain of morphisms in $\Pi(\Delta^n)$, m_i being the edge morphism from the vertex $i-1$ to i . Then define

$$(10.1) \quad \text{area}_{\mathcal{U}}(\mathcal{A}) = \sup_r \text{area}_{\mathbf{n}}(u(m_1, \dots, m_n, r, n)^* \mathcal{A}),$$

where $\text{area}_{\mathbf{n}}$ on the right hand side is as defined in equation (1.3). In the case $G = \text{Ham}(M, \omega)$ we take

$$\mathbf{n} : \text{lie Ham}(M, \omega) \rightarrow \mathbb{R}$$

to be

$$\mathbf{n}(H) = |H|_+ = \max_M H.$$

Let ω be the area 1 Fubini-Study symplectic 2-form on $M = \mathbb{CP}^1$. Then the pull-back by the natural map

$$\mathrm{lie} h : \mathrm{lie} PU(2) \rightarrow \mathrm{lie} \mathrm{Ham}(\mathbb{CP}^1, \omega) \simeq C_0^\infty(\mathbb{CP}^1)$$

of the semi-norm: $|H|_+ = \max_M H$ is the operator norm on $PU(2)$, up to normalization. This will be used to get the specific form of Theorem 1.6, from the more general form here.

10.1. Simplicial connections. We now introduce a certain abstraction of simplicial connections, which can partly be understood as simplicial resolutions of singular connections. Let $G \hookrightarrow P \rightarrow X$ be a principal G bundle, where G is a Frechet Lie group. Denote by $X_\bullet$ the simplicial set whose set of n -simplices, $X_\bullet(n)$, consists of smooth maps $\Sigma : \Delta^n \rightarrow X$, with Δ^n standard topological n -simplex with vertices ordered $0, \dots, n$. And denote by $\mathrm{Simp}(X_\bullet)$ the category with objects $\cup_n X_\bullet(n)$ and with $\mathrm{hom}(\Sigma_0, \Sigma_1)$ commutative diagrams:

$$\begin{array}{ccc} \Delta^n & \xrightarrow{\mathrm{mor}} & \Delta^m \\ & \searrow \Sigma_0 & \downarrow \Sigma_1 \\ & & X, \end{array}$$

for mor a simplicial face map, that is an injective affine map preserving order of the vertices.

Definition 10.1. Define a **simplicial G -connection $\mathcal{A}$** on P to be the following data:

- For each $\Sigma : \Delta^n \rightarrow X$ in $X_\bullet(n)$ a smooth G -connection $\mathcal{A}_\Sigma$ on $\Sigma^* P \rightarrow \Delta^n$, (a usual Ehresmann G -connection.)
- For a morphism $\mathrm{mor} : \Sigma_0 \rightarrow \Sigma_1$ in $\mathrm{Simp}(X_\bullet)$, we ask that $\mathrm{mor}^* \mathcal{A}_{\Sigma_1} = \mathcal{A}_{\Sigma_0}$.

Example 10.2. If $\mathcal{A}$ is a smooth G -connection on P , define a simplicial connection by $\mathcal{A}_\Sigma = \Sigma^* \mathcal{A}$ for every simplex $\Sigma \in X_\bullet$. We call such a simplicial connection **induced**.

If we try to “push forward” a simplicial connection to get a “classical” connection on P over X , then we get a kind of multi-valued singular connection. Multi-valued because each $x \in X$ may be in the image of a number of $\Sigma : \Delta^n \rightarrow X$ and Σ itself may not be injective, and singular because each Σ is in general singular so that the naive push-forward may have blow up singularities. We will call the above the naive pushforward of a simplicial connection.

Proof of Theorem 1.6 and Corollary 1.2. We will prove this by way of a stronger result. Let P be a Hamiltonian fibration $S^2 \hookrightarrow P \rightarrow S^4$, and $\mathcal{A}$ a simplicial $\mathcal{G} = \mathrm{Ham}(S^2, \omega)$ connection on P . Denote by $\sigma_0^1 \in S_\bullet^4$ the degenerate 1-simplex at x_0 , in other words the constant map: $\sigma_0^1 : [0, 1] \rightarrow x_0$. Let κ be the $L^\pm$ -length of the holonomy path of $\mathcal{A}_{\sigma_0^1}$ over $[0, 1]$.

Finally, let $\Sigma_\pm \in S_\bullet^4(4)$ be a complementary pair as in Section 5.1. The connection $\mathcal{A}$ gives us a simplicial connection as in Example 10.2. By an inductive procedure as

in Part I, Lemma 5.6, we may find perturbation data $\mathcal{D}$ for P so that with respect to $\mathcal{D}$ the following is satisfied.

$$(10.2) \quad \forall r : pr_1 \mathcal{F}(L_0^0, \dots, L_0^n, \Sigma_\pm, r) \simeq_\delta u(m_1, \dots, m_s, r, n)^* \mathcal{A}_{\Sigma_\pm},$$

$$(10.3) \quad \mathcal{A}(L_0, L_0) \simeq_\delta \mathcal{A}_{\sigma_0^1},$$

where L_0^i are the objects as before, where $\simeq_\delta$ means δ -close in the metrized C^∞ topology, and δ is as small as we like. Here we are using notation of Part I as before. Set

$$\tilde{u}_r := \Sigma_- \circ u(m_1, \dots, m_s, r, 4),$$

so $\tilde{u}_r : \mathcal{S}_r \rightarrow S^4$. Set $\tilde{\mathcal{S}}_r := \tilde{u}_r^* P$, set $\mathcal{A}'_r := pr_1 \mathcal{F}(L_0^0, \dots, L_0^n, \Sigma_-, r)$ and set

$$\{\Theta_r\} := \{\tilde{\mathcal{S}}_r, \mathcal{S}_r, \mathcal{L}_r, \mathcal{A}'_r\}.$$

Definition 10.3. *We say that $\mathcal{A}$ is **perfect**, if for every arbitrarily small δ as above, $\mathcal{A}(L_0, L_0)$, as above, can be chosen so that the corresponding Floer chain complex $CF(L_0, L_1, \mathcal{A}(L_0, L_0))$ is perfect.*

Theorem 10.4. *Let $\mathcal{A}$ be a perfect simplicial Hamiltonian connections on P . If P is non-trivial as a Hamiltonian bundle then*

$$(\text{area}_{\mathcal{U}}(\mathcal{A}_{\Sigma_+}) \geq \hbar - 5\kappa) \vee (\text{area}_{\mathcal{U}}(\mathcal{A}_{\Sigma_-}) \geq \hbar - 5\kappa),$$

Proof. Suppose

$$(10.4) \quad \text{area}_{\mathcal{U}}(\mathcal{A}_{\Sigma_+}) < \hbar - 5\kappa.$$

Then by (10.2), (10.2) and by [7, Lemma 2.34] $\mathcal{D}$, as defined above, can be assumed to be unexcited provided δ is chosen to be sufficiently small. Take the unital replacement as in Lemma 6.1. Since we know that $K(P)$ does not admit a section by Theorem 4.4, the simplex T of the Lemma 6.1 does not exist. Hence, again by this lemma

$$ev(\{\Theta_r\}, A_0) = [\mu_{\Sigma_-}^4(\gamma_1, \dots, \gamma_4)] \neq 0.$$

In particular

$$\overline{\mathcal{M}}(\{\Theta_r\}, A_0) \neq \emptyset.$$

So by [7, Lemma 2.34] there exists an r_0 so that

$$(10.5) \quad \text{area}(\mathcal{A}'_{r_0}) \geq \hbar - 5\kappa'.$$

where κ' denotes the $L^\pm$ length of the holonomy path in $\text{Ham}(S^2, \omega)$ of $\mathcal{A}(L_0, L_0)$. By (10.3) $\kappa' \rightarrow \kappa$ as $\delta \rightarrow 0$. By (10.2), (10.5) passing to the limit as $\delta \rightarrow 0$ we get:

$$\text{area}_{\mathcal{U}}(\mathcal{A}_{\Sigma_-}) \geq \hbar - 5\kappa.$$

□

Corollary 10.5. *Let $\mathcal{A}$ be a $PU(2)$ connection on a non-trivial principal $PU(2)$ bundle $P \rightarrow S^4$. Then*

$$(\text{area}_{\mathcal{U}}(\mathcal{A}_{\Sigma_+}) \geq \hbar - 5\kappa) \vee (\text{area}_{\mathcal{U}}(\mathcal{A}_{\Sigma_-}) \geq \hbar - 5\kappa).$$

Proof. A simplicial $PU(2)$ connection $\mathcal{A}$ on a principal $PU(2)$ bundle $PU(2) \hookrightarrow P' \rightarrow S^4$ is automatically perfect, when understood as a Hamiltonian connection on the associated bundle $S^2 \hookrightarrow P \rightarrow S^4$. So that this is an immediate consequence of the theorem above. $\square$

To prove Corollary 1.2, we just note that for an induced simplicial Hamiltonian connection $\mathcal{A}$, as defined in Example 10.2, $\mathcal{A}_{\sigma_0^1}$ is trivial. And hence $\mathcal{A}$ is automatically perfect. So that this corollary follows by Theorem 10.4. $\square$

APPENDIX A. HOMOTOPY GROUPS OF KAN COMPLEXES

For convenience let us quickly review Kan complexes just to set notation. This notation is also used in Part I. Let

$$\Delta_{\bullet}^n(k) := \text{hom}_{\Delta}([k], [n]),$$

be the standard representable n -simplex, where Δ is the simplicial category with objects ordered finite sets $[n] = \{1, \dots, n\}$ and morphisms order preserving set maps.

Let $\Lambda_k^n \subset \Delta_{\bullet}^n$ denote the sub-simplicial set corresponding to the “boundary” of $\Delta_{\bullet}^n$ with the k 'th face removed, $0 \leq k \leq n$. By k 'th face we mean the face opposite to the k 'th vertex. Let $X_{\bullet}$ be an abstract simplicial set. A simplicial map

$$h : \Lambda_k^n \subset \Delta_{\bullet}^n \rightarrow X_{\bullet}$$

will be called a **horn**. A simplicial set $S_{\bullet}$ is said to be a **Kan complex** if for all $n, k \in \mathbb{N}$ given a diagram with solid arrows

$$\begin{array}{ccc} \Lambda_k^n & \xrightarrow{h} & S_{\bullet} \\ \downarrow i & \nearrow \tilde{h} & \\ \Delta_{\bullet}^n & & \end{array},$$

there is a dotted arrow making the diagram commute. The map $\tilde{h}$ will be called **the Kan filling** of the horn h . The k 'th face of $\tilde{h}$ will be called **Kan filled face along h** . As before we will denote Kan complexes and ∞ -categories by calligraphic letters.

Given a pointed Kan complex $(\mathcal{X}, x)$ and $n \geq 1$ the n 'th simplicial homotopy group of $(\mathcal{X}, x)$: $\pi_n(\mathcal{X}, x)$ is defined to be the set of equivalence classes of maps

$$\Sigma : \Delta_{\bullet}^n \rightarrow \mathcal{X},$$

such that Σ takes $\partial\Delta_{\bullet}^n$ to $x_{\bullet}$, with the latter denoting the image of $\Delta_{\bullet}^0 \rightarrow \mathcal{X}$, induced by the vertex inclusion $x \rightarrow \mathcal{X}$.

More precisely, we have a commutative diagram:

$$\begin{array}{ccc} \Delta_{\bullet}^n & \longrightarrow & \Delta_{\bullet}^0 \\ & \searrow \Sigma & \downarrow x \\ & & \mathcal{X}. \end{array}$$

Example A.1. When $\mathcal{X} = X_{\bullet}$ is the simplicial set of singular simplices of a topological space X , the maps above are in complete correspondence with maps:

$$\Sigma : \Delta^n \rightarrow X,$$

taking the topological boundary of Δ^n to x .

For $X_{\bullet}$ general simplicial set, a pair of maps $\Sigma_1 : \Delta_{\bullet}^n \rightarrow X_{\bullet}, \Sigma_2 : \Delta_{\bullet}^n \rightarrow X_{\bullet}$, are equivalent if there is a diagram, called simplicial homotopy:

$$\begin{array}{ccc} \Delta_{\bullet}^n & & \\ \downarrow i_0 & \searrow \Sigma_1 & \\ \Delta_{\bullet}^n \times I_{\bullet} & \longrightarrow & X_{\bullet} \\ \uparrow i_1 & \nearrow \Sigma_2 & \\ \Delta_{\bullet}^n & & \end{array}$$

such that $\partial\Delta_{\bullet}^n \times I_{\bullet}$ is taken by H to $x_{\bullet}$. The simplicial homotopy groups of a Kan complex $(\mathcal{X}, x)$ coincide with the classical homotopy groups of the geometric realization $(|\mathcal{X}|, x)$.

Proof of Lemma 4.1. We refer the reader to Part I, Appendix A.2, for more details on the notions here. We prove a stronger claim.

Lemma A.2. Let $p : \mathcal{Y} \rightarrow \mathcal{X}$ be an inner fibration of quasi-categories $\mathcal{Y}, \mathcal{X}$, with $\mathcal{X}$ a Kan complex. And let $K(\mathcal{Y}) \subset \mathcal{Y}$ denote the maximal Kan subcomplex. Then $p : K(\mathcal{Y}) \rightarrow \mathcal{X}$ is a Kan fibration.

The above is probably well known, but it is simple to just provide the proof for convenience.

Proof. By definition of an inner fibration, whenever we are given a commutative diagram with solid arrows and with $0 < k < n$:

$$(A.1) \quad \begin{array}{ccccc} \Lambda_k^n & \xrightarrow{\sigma} & K(\mathcal{Y}) & \hookrightarrow & \mathcal{Y} \\ \downarrow & & \searrow \Sigma & & \uparrow p \\ \Delta^n & \longrightarrow & \mathcal{X}, & & \end{array}$$

there exists a dashed arrow Σ as indicated, making the whole diagram commutative. When $n > 2$ the edges, i.e. 1-faces, of Σ are all automatically isomorphisms in $\mathcal{Y}$, as Σ extends σ , and all edges of σ are isomorphisms by definition. For $n = 2$ the edges

of Σ are either edges of σ , or are compositions of edges of σ in the quasi-category $\mathcal{Y}$, and hence again always invertible.

It follows that Σ maps into $K(\mathcal{Y}) \subset \mathcal{Y}$. Since the starting diagram was arbitrary, we just proved that $p : K(\mathcal{Y}) \rightarrow \mathcal{X}$ is an inner fibration. In particular the pre-images $p^{-1}(\Sigma(\Delta^n)) \subset K(\mathcal{Y})$ are quasi-categories, for all n , where $\Sigma : \Delta^n \rightarrow \mathcal{X}$ is any n -simplex, see Part I, Appendix A.2. But $K(\mathcal{Y})$ is a Kan complex, so that also the above pre-images $p^{-1}(\Sigma(\Delta^n))$ are Kan complexes. It readily follows from this that $p : K(\mathcal{Y}) \rightarrow \mathcal{X}$ is a Kan fibration. $\square$

The main lemma then follows, since if $p : \mathcal{Y} \rightarrow \mathcal{X}$ is a Cartesian fibration, it is in particular an inner fibration. $\square$

REFERENCES

- [1] M. F. ATIYAH AND R. BOTT, *The Yang-Mills equations over Riemann surfaces.*, Philos. Trans. R. Soc. Lond., A, 308 (1983), pp. 523–615.
- [2] R. HARVEY AND H. B. JUN. LAWSON, *A theory of characteristic currents associated with a singular connection*, Bull. Am. Math. Soc., New Ser., 31 (1994), pp. 54–63.
- [3] J. F. MARTINS, *On the homotopy type and the fundamental crossed complex of the skeletal filtration of a CW-complex*, Homology Homotopy Appl., 9 (2007), pp. 295–329.
- [4] F. NAEIMPOUR, B. MIRZA, AND F. JAHROMI, *Yang–mills black holes in quasitopological gravity*, The European Physical Journal C, 81 (2021).
- [5] E. RIEHL, *A model structure for quasi-categories*, <https://emilyriehl.github.io/files/topic.pdf>.
- [6] Y. SAVELYEV, *Hamiltonian elements in algebraic K-theory*, <http://yashamon.github.io/web2/papers/KtheoryCycles.pdf>.
- [7] Y. SAVELYEV, *Quantum Maslov classes*, <http://yashamon.github.io/web2/papers/quantumMaslov.pdf>.
- [8] ———, *Global Fukaya category I*, Int. Math. Res. Not., 2023 (2023), pp. 18302–18386.
- [9] L. M. SIBNER AND R. J. SIBNER, *Classification of singular Sobolev connections by their holonomy*, Commun. Math. Phys., 144 (1992), pp. 337–350.
- [10] B. TOËN, *The homotopy theory of dg-categories and derived Morita theory.*, Invent. Math., 167 (2007), pp. 615–667.

UNIVERSITY OF COLIMA, CUICBAS, BERNAL DÍAZ DEL CASTILLO 340, COL. VILLAS SAN SEBASTIAN, 28045, COLIMA, COLIMA, MEXICO

Email address: yasha.savelyev@gmail.com